

Properties of field functionals and characterization of local functionals

Christian Brouder

Sorbonne Universités, UPMC Univ. Paris 06, UMR CNRS 7590,
Institut de Minéralogie, de Physique des Matériaux et de Cosmochimie,
Muséum National d'Histoire Naturelle, IRD UMR 206,
4 place Jussieu, F-75005 Paris, France.

Nguyen Viet Dang

Institut de Mathématiques de Jussieu - Paris Rive Gauche,
Université Paris Diderot-Paris 7, CNRS UMR7586,
Case 7012, Bât. Sophie Germain, 75205 Paris cedex 13, France.

Camille Laurent-Gengoux

Institut Elie Cartan de Lorraine (IECL)
Université de Lorraine, Bât. A, Ile du Saulcy
F-57045 Metz cedex 1, France.

Kasia Rejzner

Department of Mathematics, University of York,
Heslington, York YO10 5DD, United Kingdom.

October 17, 2018

Abstract

Functionals (i.e. functions of functions) are widely used in quantum field theory and solid-state physics. In this paper, functionals are given a rigorous mathematical framework and their main properties are described. The choice of the proper space of test functions (smooth functions) and of the relevant concept of differential (Bastiani differential) are discussed.

The relation between the multiple derivatives of a functional and the corresponding distributions is described in detail. It is proved that, in a neighborhood of every test function, the support of a smooth functional is uniformly compactly supported and the order of the corresponding distribution is uniformly bounded. By leaning on a recent work by Yoann Dabrowski, several spaces of functionals are furnished with a complete and nuclear topology.

In view of physical applications, it is shown that most formal manipulations can be given a rigorous meaning. A new concept of local functionals is proposed and two characterizations of it are given: the first one uses the additivity (or Hammerstein) property, the second one is a variant of Peetre's theorem. Finally, the first step of a cohomological approach to quantum field theory is carried out by proving a global Poincaré lemma within our framework.

1 Motivation

Functionals (i.e. functions of functions) met with a tremendous success in many areas of physics. Since Schwinger's ground-breaking papers [1, 2], Green functions of quantum field theory are obtained as functional derivatives of the generating functional $Z(j)$ with respect to the functions

j (external sources). In solid-state and molecular physics, the exchange and correlation potential of density functional theory is computed from the functional derivative of the total energy $E(\rho)$ with respect to the electron density ρ [3, 4]. In perturbative algebraic quantum field theory, the observables are functionals $F(\varphi)$ of the classical field φ [5]. Functionals are also used in pure mathematics, for example loop space cohomology [6] and infinite dimensional integrable systems: the hierarchy of commuting Hamiltonians for the Korteweg de Vries is for instance all made of functionals [7].

In all these fields, the concept of locality is crucial: the Lagrangian of quantum field theory is local and the counterterms of the renormalization process have to be local, the approximations of $E(\rho)$ used in practice are local and it is an open question whether the true density functional $E(\rho)$ is local or not. Therefore, it is crucial to determine precisely what is meant by a *local functional*. According to the standard definition [8, 9, 10, 11], if φ is a classical field (i.e. a smooth section of a vector bundle over M and we momentarily consider $M = \mathbb{R}^d$ for notational convenience), then a functional $F(\varphi)$ is local if it is of the form

$$F(\varphi) = \int_{\mathbb{R}^d} dx f(x, \varphi(x), \partial_\mu \varphi(x), \dots, \partial_{\mu_1} \dots \partial_{\mu_k} \varphi(x)). \quad (1)$$

where f is a smooth compactly supported function with a finite number of arguments.

However, this definition of local functionals is not very handy in practice because it is global and sometimes too restrictive. For example, general relativity has no local gauge-invariant observables in the sense of Eq. (1), whereas it has local gauge-invariant observables when the concept of locality is slightly generalized, as discussed in [12] (see also the parallel work [13]). Note that the concept of locality presented in the present paper gives a proper topological framework for local functionals as understood by [14, 15]. Brunetti and Fredenhagen conjectured that F is local in the sense of Eq. (1) (without an explicit condition on the number of derivatives) if and only if the following conditions are satisfied: (i) F is smooth; (ii) F is additive (in the sense of section 5.3); (iii) F is compactly supported; (iv) the wave front set of all the derivatives $F_\varphi^{(k)}$ of F at φ is conormal to the small diagonal of M^k ([16, p. 139], [15, p. 1555], [17, p. 19], [14]). These concepts will be defined shortly.

The present paper puts forth the following formulation of the concept of locality:

Definition 1.1. *Let M be a manifold¹. Let U be an open subset of $C^\infty(M)$. A smooth functional $F : U \rightarrow \mathbb{K}$ is said to be local if, for every $\varphi \in U$, there is a neighborhood V of φ , an integer k , an open subset $\mathcal{V} \subset J^k M$ and a smooth function $f \in C^\infty(\mathcal{V})$ such that $x \in M \mapsto f(j_x^k \psi)$ is supported in a compact subset $K \subset M$ and*

$$F(\varphi + \psi) = F(\varphi) + \int_M f(j_x^k \psi) dx,$$

whenever $\varphi + \psi \in V$ and where $j_x^k \psi$ denotes the k -jet of ψ at x .

In other words, we require F to be local in the sense of Eq. (1), but only around each $\varphi \in U$ because the integer k and the function f can depend on the neighborhood V . In short, our local functionals are local in the “traditional sense”, but only locally in the configuration space (i.e. in a neighborhood of each φ). We do not need global locality to apply variational methods and derive Euler-Lagrange equations. We will show by exhibiting an example that this concept of locality is strictly more general than the traditional one. Our first main result is a simple characterization of local functionals in the sense of Def. 1.1:

Theorem 1.2. *Let U be an open subset of $C^\infty(M)$. A smooth functional $F : U \rightarrow \mathbb{K}$ (where $\mathbb{K} = \mathbb{R}$ or \mathbb{C}) is local if and only if*

1. F is additive (i.e. it satisfies $F(\varphi_1 + \varphi_2 + \varphi_3) = F(\varphi_1 + \varphi_2) + F(\varphi_2 + \varphi_3) - F(\varphi_2)$ whenever $\text{supp } \varphi_1 \cap \text{supp } \varphi_3 = \emptyset$)

¹In this article, all manifolds shall be assumed to be paracompact.

2. For every $\varphi \in U$, the differential DF_φ of F at φ is a distribution with empty wave front set. Thus, it can be represented by a function $\nabla F_\varphi \in \mathcal{D}(M)$ (with $\mathcal{D}(M)$ the space of compactly supported smooth functions on M , i.e. “test functions”).
3. The map $U \rightarrow \mathcal{D}(M)$ defined by $\varphi \mapsto \nabla F_\varphi$ is smooth (in the sense of Bastiani).

Let us stress that the notion of locality is quite subtle and depends strongly on the functional analytic setting. A functional characterization of a notion of local functionals on **measurable functions** might not be valid anymore when applied to **smooth functions** as is shown by the simple counterexample of section 5.2. We also make a conjecture as to how to generalize our main result to multi-vector fields and graded functionals, which is crucial for a rigorous version of the Batalin-Vilkovisky approach to gauge field theory and quantum gravity. The second main result is a proof of the global Poincaré lemma (in our context), which is crucial to set up the BRST and variational complexes. The last one is another characterization of local (and multilocal) functionals in the form of a Peetre’s theorem.

Along the way to these results, we prove interesting properties of general functionals that we briefly describe now. In section 2, we explain why we choose test functions that are only smooth instead of smooth and compactly supported, we describe the topology of the space and test functions and we present the concept of Bastiani differentiability and its main properties. In section 3, we show that a smooth functional is locally compactly supported (i.e. in a neighborhood of every test function), we prove that the k th derivative of a functional defines a continuous family of distributions whose order is locally bounded. Section 4, which relies heavily on Dabrowski’s work [18, 19], describes in detail a nuclear and complete topology on several spaces of functionals used in quantum field theory. Section 5 discusses the concept of additivity which characterizes local functionals. Sections 6 and 7 prove the main results discussed above. Note that the present paper has a somewhat foundational character, in as much as the choice of test-functions, additivity property and differential are carefully justified from the physical and mathematical points of view. It contributes to the formulation of a mathematically rigorous basis on which the quantum field theory of gauge fields and gravitation can be built.

Note also that this paper aims at both functional analysts and theoretical physicists. Because of this dual readership, the proofs are often more detailed than what would be required for experts in functional analysis.

2 Functionals and their derivatives

To set up a mathematical definition of functionals, we need to determine precisely which space of test functions (i.e. classical fields and sources) we consider and what we mean by a functional derivative.

2.1 The space of classical fields

Propagators and Green functions of quantum fields in flat spacetimes are tempered distributions [20, 21] and the corresponding test functions are rapidly decreasing. Tempered distributions are computationally convenient because they have Fourier transforms. However, tempered distributions cannot be canonically extended to curved spacetimes (i.e. Lorentzian smooth manifolds) because the rapid decrease of test functions at infinity is controlled by a Euclidian distance which is not canonically defined on general spacetime manifolds [22, p. 339].

The most natural spaces of test functions on a general spacetime M are the space $C^\infty(M)$ of real valued smooth functions on M and its subspace $\mathcal{D}(M)$ of compactly supported functions. These two spaces are identical when M is compact, but physically relevant spacetimes are not compact because they are globally hyperbolic, and a choice must be made.

In this paper, we choose $C^\infty(M)$ (or the set $\Gamma(M, B)$ of smooth sections of a vector bundle B). There is a strong physical reason for this [23]: in the quantization process we must be able to deal with on-shell fields φ , that are smooth solutions to normally hyperbolic equations and as such cannot be compactly supported. Therefore, the domain of the functionals can be $C^\infty(M)$ but

not $\mathcal{D}(M)$. There are also good mathematical arguments for this choice: In particular, $C^\infty(M)$ is a Fréchet space and its pointwise multiplication is continuous [24, p. 119]. Moreover, the Fréchet property of $C^\infty(M)$ saves us the trouble of distinguishing Bastiani from convenient differentiability which is treated in the monumental work [25].

The choice of $C^\infty(M)$ has, however, several drawbacks: (i) Since smooth functions are generally not integrable over M , the Lagrangian density $\mathcal{L}(\varphi)$ must be multiplied by a smooth compactly supported function g so that $\mathcal{L}(\varphi)g$ is integrable over M [26]. As a result, long-range interactions are suppressed and infrared convergence is enforced. This simplifies the problem but makes it difficult to deal with the physics of infrared divergence. (ii) The function g breaks the diffeomorphism invariance of the Einstein-Hilbert action. (iii) The effect of a perturbation $\varphi + \epsilon\psi$ is easier to deal with when ψ is compactly supported because it avoids the presence of boundary terms. This problem can be solved by considering $C^\infty(M)$ as a manifold modeled on $\mathcal{D}(M)$ [17, 5], but this is an additional complication.

2.2 Locally convex spaces

The spaces of test functions and of functionals considered in the paper are all locally convex. The most pedagogical introduction to locally convex spaces is probably Horvath's book [27], so we refer the reader to that publication for more details.

We describe now the topology of the spaces of test functions that we use. For the space of smooth test functions $C^\infty(\mathbb{R}^d)$, the topology is defined by the seminorms

$$\pi_{m,K}(f) = \sup_{x \in K} \sup_{|\alpha| \leq m} |\partial^\alpha f(x)|, \quad (2)$$

where $f \in C^\infty(\mathbb{R}^d)$, m is an integer, K is a compact subset of \mathbb{R}^d , $\alpha = (\alpha_1, \dots, \alpha_d)$ is a d -tuple of nonnegative integers, with $|\alpha| = \alpha_1 + \dots + \alpha_d$ and $\partial^\alpha = \partial_1^{\alpha_1} \dots \partial_d^{\alpha_d}$, with $\partial_i = \partial/\partial x^i$ the derivative with respect to the i -th coordinate of x [24, p. 88].

If U is open in \mathbb{R}^d , we denote by $C^\infty(U)$ the space of all functions defined on U which possess continuous partial derivatives of all orders. We equip $C^\infty(U)$ with the topology defined by the seminorms $\pi_{m,K}$ where K runs now over the compact subsets of U [27, p. 89]. For every open set $U \subset \mathbb{R}^d$, the space $C^\infty(U)$ is Fréchet, reflexive, Montel, barrelled [27, p. 239], bornological [27, p. 222] and nuclear [28, p. 530].

We define now $C^\infty(M)$, where M is a d -dimensional manifold (tacitly smooth, Hausdorff, paracompact and orientable) described by charts (U_α, ψ_α) . If for every $U_\alpha \subset M$ we are given a smooth function $g_\alpha \in C^\infty(\psi_\alpha(U_\alpha))$ such that $g_\beta = g_\alpha \circ \psi_\alpha \circ \psi_\beta^{-1}$ on $\psi_\beta(U_\alpha \cap U_\beta)$, we call the system g_α a smooth function g on M . The space of smooth functions on M is denoted by $C^\infty(M)$ [29, p. 143]. This definition is simple but to describe the topological properties of $C^\infty(M)$ the following more conceptual definition is useful.

Let M be a manifold and $B \rightarrow M$ a smooth vector bundle of rank r over M with projection π . Let $E = \Gamma(M, B)$ be the space of smooth sections of B furnished with the following topology [25].

Definition 2.1. *The topology on $\Gamma(M, B)$ is defined as follows. Choose a chart $(U_\alpha, \psi_\alpha)_\alpha$ and a trivialization map $\Phi_\alpha : \pi^{-1}(U_\alpha) \rightarrow \Omega \times \mathbb{R}^r$, where Ω is a fixed open set in \mathbb{R}^d . Then the map Φ_α allows to identify $\Gamma(U_\alpha, B)$ with $C^\infty(\Omega, \mathbb{R}^r)$ by*

$$\Phi_\alpha : (x, s(x)) \in \pi^{-1}U_\alpha \mapsto (\psi_\alpha(x), K_\alpha(s)(\psi_\alpha(x))) \in \Omega \times \mathbb{R}^d,$$

where

$$K_\alpha : s \in \Gamma(U_\alpha, B) \mapsto K_\alpha(s) \in C^\infty(\Omega, \mathbb{R}^r).$$

The topology on $\Gamma(M, B)$ is the weakest topology making all the maps K_α continuous.

This topology does not depend on the choice of charts or trivialization maps [25, p. 294]. To interpret this topology, denote by $\rho_\alpha : s \in \Gamma(M, B) \mapsto s|_{U_\alpha} \in \Gamma(U_\alpha, B)$ the restriction map of

sections on open sets of our open cover $(U_\alpha)_{\alpha \in I}$ of M . The space $\Gamma(M, B)$ fits into the following complex of vector spaces

$$0 \rightarrow \Gamma(M, B) \xrightarrow{(\rho_\alpha)_{\alpha \in I}} \prod_{\alpha \in I} \Gamma(U_\alpha, B) \simeq \prod_{\alpha \in I} C^\infty(\Omega, \mathbb{R}^r) \xrightarrow{(\rho_\alpha - \rho_\beta)_{\alpha \neq \beta}} \prod_{(\alpha \neq \beta) \in I^2} \Gamma(U_\alpha \cap U_\beta, B). \quad (3)$$

The topology on $\Gamma(U_\alpha, B)$ is given by the isomorphism $\Gamma(U_\alpha, B) \simeq C^\infty(\Omega, \mathbb{R}^r)$ hence it is nuclear Fréchet. The countable products $\prod_{\alpha \in I} \Gamma(U_\alpha, B)$ and $\prod_{(\alpha, \beta) \in I^2} \Gamma(U_\alpha \cap U_\beta, B)$ are therefore nuclear Fréchet. For every pair (α, β) of distinct elements of I , the difference of restriction maps $\rho_\alpha - \rho_\beta$ is continuous and the topology on $\Gamma(M, B)$ is the weakest topology which makes the above complex topological, which implies that it is nuclear Fréchet as the kernel of $\prod_{\alpha \neq \beta} \rho_\alpha - \rho_\beta$.

Locally convex spaces are very versatile and they are the proper framework to define spaces of smooth functionals, i.e. smooth functions on a space of functions (or sections of a bundle). The first step towards this goal is to provide a rigorous definition of functional derivatives.

2.3 Functional derivatives

To define the space of functionals, we consider the main examples $Z(j)$ and $F(\varphi)$. These two functionals send smooth classical fields to \mathbb{K} , where $\mathbb{K} = \mathbb{R}$ or $\mathbb{K} = \mathbb{C}$. Moreover, functional derivatives of Z and F of all orders are required to obtain the Green functions from $Z(j)$ and to quantize the product $F(\varphi)G(\varphi)$. Therefore, we must define the derivative of a function $f : E \rightarrow \mathbb{K}$, where E is the space of classical fields.

It will be useful to generalize the problem to functions f between arbitrary locally convex spaces E and F . To define such a derivative we start from

Definition 2.2. *Let U be an open subset of a Hausdorff locally convex space E and let f be a map from U to a Hausdorff locally convex space F . Then f is said to have a derivative at $x \in U$ in the direction of $v \in E$ if the following limit*

$$Df_x(v) := \lim_{t \rightarrow 0} \frac{f(x + tv) - f(x)}{t},$$

exists².

One can also consider the same definition restricted to $t > 0$ [30]. A function f is said to have a *Gâteaux differential* [31, 32] (or a *Gâteaux variation* [33]) at x if $Df_x(v)$ exists for every $v \in E$. However, this definition is far too weak for our purpose because $Df_x(v)$ is generally neither linear nor continuous in v and it can be linear without being continuous and continuous without being linear [34, p. 7]. Therefore, we will use a stronger definition, namely *Bastiani differentiability* [35], which is the fundamental concept of differentiability used throughout the paper:

Definition 2.3. *Let U be an open subset of a Hausdorff locally convex space E and let f be a map from U to a Hausdorff locally convex space F . Then f is Bastiani differentiable on U (denoted by $f \in C^1(U)$) if f has a Gâteaux differential at every $x \in U$ and the map $Df : U \times E \rightarrow F$ defined by $Df(x, v) = Df_x(v)$ is continuous on $U \times E$.*

With this definition, most of the properties used in physics textbooks (e.g. chain rule, Leibniz rule, linearity) are mathematically valid.

²Since an open set is absorbing [27, p. 80], for every $x \in U$ and every $v \in E$ there is an $\epsilon > 0$ such that $x + tv \in U$ if $|t| < \epsilon$. Thus, $f(x + tv)$ is well defined for every t such that $|t| < \epsilon$.

2.3.1 Examples

We shall consider several examples of functions from $C^\infty(M)$ to \mathbb{R} or \mathbb{C} , where $M = \mathbb{R}^d$:

$$\begin{aligned}
F(\varphi) &= \int_M f(x)\varphi^n(x)dx, \\
G(\varphi) &= \int_{M^n} g(x_1, \dots, x_n)\varphi(x_1)\dots\varphi(x_n)dx_1\dots dx_n, \\
H(\varphi) &= \sum_{\mu\nu} \int_M g^{\mu\nu}(x)h(x)\partial_\mu\varphi(x)\partial_\nu\varphi(x)dx, \\
I(\varphi) &= \int_M f(x)e^{\varphi(x)}dx, \\
J(\varphi) &= e^{\int_M f(x)\varphi(x)dx}, \\
K(\varphi) &= \int_M f(x)\sin(\varphi(x))dx,
\end{aligned}$$

where f , g , and h are smooth compactly supported functions and where g is a symmetric function of its arguments. Further examples can be found in [15, 36]. It is immediate to check that

$$\begin{aligned}
DF_\varphi(v) &= n \int_M g(x)\varphi^{n-1}(x)v(x)dx, \\
DG_\varphi(v) &= n \int_{M^n} g(x_1, \dots, x_n)\varphi(x_1)\dots\varphi(x_{n-1})v(x_n)dx_1\dots dx_n, \\
DH_\varphi(v) &= 2 \sum_{\mu\nu} \int_M g^{\mu\nu}(x)h(x)\partial_\mu\varphi(x)\partial_\nu v(x)dx, \\
DI_\varphi(v) &= \int_M f(x)e^{\varphi(x)}v(x)dx, \\
DJ_\varphi(v) &= e^{\int_M f(x)\varphi(x)dx} \int_M f(x)v(x)dx, \\
DK_\varphi(v) &= \int_M f(x)\cos(\varphi(x))v(x)dx.
\end{aligned}$$

2.3.2 Historical remarks

Definition 2.3 is due to Bastiani [37, 35] and looks quite natural. In fact, it is not so. For a long time, many different approaches were tried. For any reasonable definition of differentiability, the map $Df_x : E \rightarrow F$ is linear and continuous, so that $Df_x \in L(E, F)$. If E and F are Banach spaces, then a map $f : E \rightarrow F$ is defined to be continuously (Fréchet) differentiable if the map $x \rightarrow Df_x$ is continuous from U to $L_c(E, F)$, where $L_c(E, F)$ is the set of continuous maps from E to F equipped with the operator norm topology. But Fréchet differentiability is strictly stronger than Bastiani's differentiability specialized to Banach spaces [38]. This is why Bastiani's definition was often dismissed in the literature [39] and, for locally convex spaces that are not Banach, the map Df was generally required to be continuous from U to $L(E, F)$ equipped with some well-chosen topology. However, when E is not normable, no topology on $L(E, F)$ provides the nice properties of Bastiani's definition [40, p. 6] (Hamilton [36, p. 70] gives a simple example of a map which is continuous $U \times E \rightarrow E$ but such that the corresponding map $U \rightarrow L(E, E)$ is not continuous). Thus, $L(E, F)$ was equipped with various non-topological convergence structures [40, p. 23]. The result is an impressive zoology of differentiabilitys. Twenty-five of them were reviewed and classified by Averbukh and Smolyanov [41]. Still more can be found in the extensive lists given by Gähler [42] and Ver Eecke [43] covering the period up to 1983 (see also [34, 40, 44]).

Nowadays, essentially two concepts of differentiability survive, Bastiani's and the so-called *convenient approach* developed by Kriegl–Michor in the reference monograph [25], which is weaker than Bastiani's for general Hausdorff locally convex spaces. In particular, on any locally convex

space which is not bornological, there is a conveniently smooth map which is not continuous [37, p. 19]. However, a nice feature of both approaches is that for a Fréchet space E , a function $f : E \rightarrow \mathbb{K}$ is smooth in the sense of Bastiani iff it is smooth in the sense of the convenient calculus [45]. Bastiani differentiability became widespread after it was used by Michor [46], Hamilton [36] (for Fréchet spaces) and Milnor [47] and it is now vigorously developed by Glöckner and Neeb (see also [48, p. 8]).

To complete this section, we would like to mention that the Bastiani differential is sometimes called the Michal-Bastiani differential [49, 50, 51] (or even Michel-Bastiani differential [50]). This is not correct. The confusion comes from the fact that Bastiani defines her differentiability in several steps. She starts from the Gâteaux derivability, then she says that a map $f : U \rightarrow F$ is *differentiable at x* (see [37, p. 18] and [35, p. 18]) if: i) Df_x is linear and continuous from E to F and ii) the map $m_x : \mathbb{R} \times E \rightarrow F$ defined by

$$m_x(t, v) = \frac{f(x + tv) - f(x)}{t} - Df_x(v),$$

for $x + tv \in U$, is continuous at $(0, v)$ for all $v \in E$. This differentiability at x is indeed equivalent to the differentiability defined by Michal [52] in 1938, as proved in Refs. [41, 53], [40, p. 72] and [43, p. 202]. What we call Bastiani differentiability is called *differentiability on an open set* by Bastiani (see [37, p. 25] and [35, p. 44]) and is strictly stronger than Michal differentiability.

The same distinction between Michal-Bastiani differentiability and Bastiani differentiability is made by Keller [40, p. 72] in his thorough review. Bastiani's differentiability is denoted by C_c^1 by Keller [40], who also attributes the definition equivalent to C_c^1 to Bastiani alone [40, p. 11].

In her PhD thesis, Andrée Bastiani developed her concept of differentiability to define distributions on a locally convex space E with values in a locally convex space F . She started from Schwartz' remark that a distribution is, locally, the derivative of a continuous function. She used her differential D to defined F -valued distributions over E [54]. A drawback of Bastiani's framework with respect to the convenient framework is that her category is not Cartesian closed for locally convex spaces that are not Fréchet.

2.4 Properties of the differential

We review now some of the basic properties of functional derivatives which will be used in the sequel. We strongly recommend Hamilton's paper [36], adapted to locally convex spaces by Neeb [38].

2.4.1 Continuity

We characterize continuous (nonlinear) maps between two locally convex spaces.

Lemma 2.4. *Let E and F be locally convex spaces whose topology is defined by the families of seminorms $(p_i)_{i \in I}$ and $(q_j)_{j \in J}$, respectively. Then f is continuous at x iff, for every seminorm q_j of F and every $\epsilon > 0$, there is a finite number $\{p_{i_1}, \dots, p_{i_k}\}$ of seminorms of E and k strictly positive numbers η_1, \dots, η_k such that $p_{i_1}(x-y) < \eta_1, \dots, p_{i_k}(x-y) < \eta_k$ imply $q_j(f(y) - f(x)) < \epsilon$.*

Proof. This is just the translation in terms of seminorms of the fact that f is continuous at x if, for every open set V containing $f(x)$, there is an open set U containing x such that $f(U) \subset V$ [55, p. 86]. \square

When the seminorms of E are saturated [27, p. 96], as the seminorms $\pi_{m,K}$ of $C^\infty(\mathbb{R}^d)$, the condition becomes simpler: a map $f : C^\infty(\mathbb{R}^d) \rightarrow \mathbb{K}$ is continuous at x if and only if, for every $\epsilon > 0$, there is a seminorm $\pi_{m,K}$ and an $\eta > 0$ such that $\pi_{m,K}(x-y) < \eta$ implies $|f(y) - f(x)| < \epsilon$. Since Fréchet spaces are metrizable, we can also use the following characterization of continuity [56, p. 154]:

Proposition 2.5. *Let E be a metrizable topological space and F a topological space. Then, a map $f : E \rightarrow F$ is continuous at a point x iff, whenever a sequence $(x_n)_{n \in \mathbb{N}}$ converges to x in E , the sequence $f(x_n)_{n \in \mathbb{N}}$ converges to $f(x)$ in F .*

Another useful theorem is [57, p. III.30]:

Proposition 2.6. *Let E and F be two Fréchet spaces and G a locally convex space. Every separately continuous bilinear mapping from $E \times F$ to G is continuous.*

This result extends to multilinear mappings from a product $E_1 \times \cdots \times E_n$ of Fréchet spaces to a locally convex space [58, p. 178][59].

2.4.2 The fundamental theorem of calculus

The fundamental theorem of calculus for functionals reads

Theorem 2.7. *Let f be a Bastiani differentiable map between two Hausdorff locally convex spaces E and F . Let U be an open set in E , x in U and v in E such that $(x + tv) \in U$ for every t in an open neighborhood I of $[0, 1]$, so that $g : t \mapsto f(x + tv)$ is a map from I to F . Then,*

$$f(x + v) = f(x) + \int_0^1 g'(t)dt = f(x) + \int_0^1 Df_{x+tv}(v)dt. \quad (4)$$

To give a meaning to Eq. (4), we need to define an integral of a function taking its values in a locally convex space. To cut a long story short [60, p. 75] [61]:

Definition 2.8. *Let X be a locally compact space (for example \mathbb{R}^n or some finite dimensional manifold), μ a measure on X and F a Hausdorff locally convex space. Let f be a compactly supported continuous function from X to F . Let F' be the topological dual of F (i.e. the space of continuous linear maps from F to \mathbb{K}). If there is an element $y \in F$ such that*

$$\langle \alpha, y \rangle = \int_X \langle \alpha, f \rangle d\mu,$$

for every $\alpha \in F'$, where $\langle \cdot, \cdot \rangle$ denotes the duality pairing, then we say that f has a weak integral and we denote y by $\int_X f d\mu$.

The uniqueness of the weak integral follows from the fact that F is Hausdorff. In general, the existence of a weak integral requires some completeness property for F [60, p. 79]. However, this is not the case for the fundamental theorem of calculus [37, p. 27]. This point was stressed by Glöckner [62].

2.4.3 Additional properties

For maps between locally convex spaces, the linearity of the differential is not completely trivial [36].

Proposition 2.9. *Let E and F be locally convex spaces and f be a Bastiani differentiable map from an open subset U of E to F . Then, for every $x \in U$, the differential $Df_x : E \rightarrow F$ is a linear map.*

The chain rule for Bastiani-differentiable functions was first proved by Bastiani herself [37, p. 32] [35, p. 44] (see also [49]).

Proposition 2.10. *Assume that E, F, G are locally convex spaces, $U \subset E$ and $V \subset F$ are open subsets and $f : V \rightarrow G$ and $g : U \rightarrow V$ are two Bastiani-differentiable maps. Then, the composite map $f \circ g : U \rightarrow G$ is Bastiani differentiable and $D(f \circ g)_x = Df_{g(x)} \circ Dg_x$.*

By using these properties, the reader can prove that our examples are all Bastiani differentiable.

2.5 Smooth functionals

To define smooth functionals we first define multiple derivatives.

Definition 2.11. *Let U be an open subset of a locally convex space E and f a map from U to a locally convex space F . We say that f is k -times Bastiani differentiable on U if:*

- *The k th Gâteaux differential*

$$D^k f_x(v_1, \dots, v_k) = \left. \frac{\partial^k f(x + t_1 v_1 + \dots + t_k v_k)}{\partial t_1 \dots \partial t_k} \right|_{t_1 = \dots = t_k = 0},$$

exists for every $x \in U$ and every $v_1, \dots, v_k \in E$.

- *The map $D^k f : U \times E^k \rightarrow F$ is continuous.*

Notice that for a function f assumed to be k -times Bastiani differentiable, the restriction to any finite dimensional affine subspace is not only k -times differentiable (in the usual sense) but indeed of class C^k . The set of k -times Bastiani differentiable functions on U is denoted by $C^k(U)$, or $C^k(U, F)$ when the target space F has to be specified. Bastiani gives an equivalent definition, called k -times differentiability on U [37, p. 40] [35, p. 59], which is denoted by C_c^k by Keller [40].

Definition 2.12. *Let U be an open subset of a locally convex space E and f a map from U to a locally convex space F . We say that f is smooth on U if $f \in C^k(U, F)$ for every integer k .*

We now list a number of useful properties of the k -th Bastiani differential:

Proposition 2.13. *Let U be an open subset of a locally convex space E and $f \in C^k(U, F)$, where F is a locally convex space, then*

1. *$D^k f_x(v_1, \dots, v_k)$ is a k -linear symmetric function of v_1, \dots, v_k [36, p. 84].*
2. *The function f is of class C^m for all $0 \leq m \leq k$ [37, p. 40]. In particular, f is continuous.*
3. *The compositions of two functions in C^k is in C^k and the chain rule holds [37, p. 51] [35, p. 55] [36, p. 84].*
4. *The map $D^m f$ is in $C^{k-m}(U, L(E^m, F))$ [37, p. 40] where $L(E^m, F)$ is the space of jointly continuous m -linear maps from E to F , equipped with the locally convex topology of uniform convergence on the compact sets of E : i.e. the topology generated by the seminorms*

$$p_{C_1, \dots, C_m, j}(\alpha) = \sup_{(h_1, \dots, h_m) \in C_1 \times \dots \times C_m} q_j(\alpha(h_1, \dots, h_m)),$$

where C_i runs over the compact sets of E and $(q_j)_{j \in J}$ is a family of seminorms defining the topology of F .

5. *If E is metrizable, then $f \in C^k(U, F)$ if and only if f belongs to $C^{k-1}(U, F)$ and $D^{k-1} f : U \rightarrow L(E^{k-1}, F)$ is Bastiani differentiable [37, p. 43]. Here the metrizability hypothesis is used to obtain a canonical injection from $C(U \times E, L(E^{k-1}, F))$ to $C(U \times E^k, F)$.*

We refer the reader to [36, 38] and Bastiani's opus cited for the proofs. Other results on $C^k(U)$ functions can be found in Keller's book [40]. All the statements of Proposition 2.13 are valid for $k = \infty$, i.e. smooth functions. Bastiani also defines jets of smooth functions between locally convex spaces [37, p. 52] [35, p. 75].

Note that Neeb [63, p. 14] and Glöckner [64, p. 20] agree with Bastiani for the definition of the first derivative but they use an apparently simpler definition of higher derivatives by saying that f is C^k iff df is C^{k-1} iff $d^{k-1} f$ is C^1 . However, this definition is less natural because, for example, $f \in C^2$ if $df : U \times E \rightarrow F$ is C^1 . In the definition of the first derivative, U is now replaced by $U \times E$ and E by $E \times E$. In other words, d^2 is a continuous maps from $U \times E^3$ to F . More generally d^k is a continuous map from $U \times E^{2^k-1}$ to F [64, p. 20]. Moreover, according

to Proposition 1.3.13 [64, p. 23], a map f belongs to C^k if and only if it belongs to $C^k(U)$ in the sense of Bastiani, and Bastiani's $D^k f$ is denoted by $d^{(k)}f$ by Glöckner [64, p. 23] and called the k -th differential of f . The k -th derivatives $d^k f$ and $d^{(k)}f = D^k f$ are not trivially related. For example [64, p. 24]: $d^2 f(x, h_1, h_2, h_3) = D^2 f(x, h_1, h_2) + Df(x, h_3)$.

The Taylor formula with remainder for a function in $C^{n+1}(U)$ reads [37, p. 44]:

$$f(x + th) = f(x) + \sum_{k=1}^n \frac{t^k}{k!} D^k f_x(h^k) + \int_0^t \frac{(t-\tau)^n}{n!} D^{n+1} f_{x+\tau h}(h^{n+1}) d\tau, \quad (5)$$

$$= f(x) + \sum_{k=1}^n \frac{t^k}{k!} D^k f_x(h^k) + \int_0^t \frac{(t-\tau)^{n-1}}{(n-1)!} (D^n f_{x+\tau h}(h^n) - D^n f_x(h^n)) d\tau. \quad (6)$$

Taylor's formula with remainder is a very important tool to deal with smooth functions on locally convex spaces.

The reader can check that all our examples are smooth functionals in the sense of Bastiani.

$$D^k F_\varphi(v_1, \dots, v_k) = \frac{n!}{(n-k)!} \int_M f(x) \varphi^{n-k}(x) v_1(x) \dots v_k(x) dx,$$

for $k \leq n$ and $D^k F_\varphi = 0$ for $k > n$.

$$D^k G_\varphi(v_1, \dots, v_k) = \frac{n!}{(n-k)!} \int_{M^n} g(x_1, \dots, x_n) v_1(x_1) \dots v_k(x_k) \varphi(x_{k+1}) \dots \varphi(x_n) dx_1 \dots dx_n,$$

for $k \leq n$ and $D^k G_\varphi = 0$ for $k > n$. Recall that g is a symmetric, smooth compactly supported function of its arguments. The functional H has only two non-zero derivatives and

$$D^2 H_\varphi(v_1, v_2) = 2 \sum_{\mu\nu} \int_M g^{\mu\nu}(x) h(x) \partial_\mu v_1(x) \partial_\nu v_2(x).$$

The example I has an infinite number of nonzero derivatives:

$$D^k I_\varphi(v_1, \dots, v_k) = \int_M f(x) e^{\varphi(x)} v_1(x) \dots v_k(x).$$

Finally

$$D^k J_\varphi(v_1, \dots, v_k) = e^{\int_M f(x) \varphi(x) dx} \int_{M^k} f(x_1) \dots f(x_k) v_1(x_1) \dots v_k(x_k) dx_1 \dots dx_k.$$

The functionals F , G and H are polynomials in the sense of Bastiani [37, p. 53]:

Definition 2.14. *Let E and F be locally convex spaces. A polynomial of degree n on E is a smooth function $f : E \rightarrow F$ such that $D^k f = 0$ for all $k > n$.*

Let u be a distribution in $\mathcal{D}'(M^k)$, then the functional $f : \mathcal{D}(M) \rightarrow \mathbb{K}$ defined by $f(\varphi) = u(\varphi^{\otimes k})$ is polynomial in the sense of Bastiani and its k -derivative is:

$$D^k f_\varphi(v_1, \dots, v_k) = \sum_{\sigma} u(v_{\sigma(1)} \otimes \dots \otimes v_{\sigma(k)}),$$

where σ runs over the permutations of $\{1, \dots, k\}$ and the canonical inclusion $\mathcal{D}(M)^{\otimes k} \subset \mathcal{D}(M^k)$ was used.

If F and G are smooth maps from E to \mathbb{K} , we can compose the smooth map $\varphi \mapsto (F(\varphi), G(\varphi))$ and the multiplication in \mathbb{K} to show that:

Proposition 2.15. *Let E be a locally convex space, and U an open set in E . Then the space of smooth functionals from U to \mathbb{K} is a sub-algebra of the algebra of real valued functions.*

3 Properties of functionals

We now prove important properties of smooth functionals. We first investigate the support of a functional. The fact that DF_φ is continuous from $C^\infty(M)$ to \mathbb{K} exactly means that DF_φ is a compactly supported distribution for every φ . The support of F is then essentially the union over φ of the supports of DF_φ . We prove that, for any smooth functional and any $\varphi \in C^\infty(M)$, there is a neighborhood V of φ such that $F|_V$ is compactly supported.

The second property that we investigate is required to establish a link with quantum field theory. In this paper, we deal with functionals that are smooth functions F on an open subset U of $E = \Gamma(M, B)$, where $\Gamma(M, B)$ is the space of smooth sections of some finite rank vector bundle B on the manifold M . There is a discrepancy between $D^k F_\varphi$, which is a continuous *multilinear* map from E^k to \mathbb{K} , and the quantum field amplitudes (e.g. represented pictorially by Feynman diagrams) that are continuous *linear* maps from $E^{\hat{\otimes}_\pi k} = \Gamma(M^k, B^{\boxtimes k})$ to \mathbb{K} , i.e. elements of the space $\Gamma'(M^k, (B^*)^{\boxtimes k})$ of compactly supported distributions with values in the k -th external tensor power of the dual bundle B^* . It is easy to see that there is a canonical correspondence between $D^k F_\varphi$ and its associated distribution on $E^{\hat{\otimes}_\pi k}$, that we denote by $F_\varphi^{(k)}$. However, the equivalence between the continuity of $D^k F$ on $U \times E^k$ and the continuity of $F^{(k)}$ on $U \times E^{\hat{\otimes}_\pi k}$ requires a proof.

Finally, we show that the order of $F^{(k)}$ is locally bounded.

3.1 Support of a functional

Brunetti, Dütsch and Fredenhagen proposed to define the support of a functional F by the property that, if the support of the smooth function ψ does not meet the support of F , then $F(\varphi + \psi) = F(\varphi)$ for all φ . More precisely [15]:

Definition 3.1. *Let $F : U \rightarrow \mathbb{K}$ be a Bastiani smooth function, with U a subset of $C^\infty(M)$. The support of F is the set of points $x \in M$ such that, for every open set \mathcal{U}_x containing x , there is a $\varphi \in U$ and a ψ in $C^\infty(M)$ with $\varphi + \psi \in U$ such that $\text{supp } \psi \subset \mathcal{U}_x$ and $F(\varphi + \psi) \neq F(\varphi)$.*

We want to relate this definition of the support of F with the support of $D_\varphi F$, which is compactly supported as every distribution over $C^\infty(M)$ [29]. To do so, we need a technical lemma about connected open subsets in locally convex spaces.

Lemma 3.2. *Let U be a connected open set in a locally convex space E then any pair $(x, y) \in U^2$ can be connected by a piecewise affine path.*

Proof. Define the equivalence relation \sim in E as follows, two elements (x, y) are equivalent iff they are connected by a piecewise affine path. Let us prove that this equivalence relation is both open and closed hence any non empty equivalence class for \sim is both open and closed in U hence equal to U .

Let $x \in U$ then there exists a convex neighborhood V of x in U which means that every element in V lies in the class of x , the relation is open. Conversely let y be in the closure of the equivalence class of x , then any neighborhood V of y contains an element equivalent to x . Choose some convex neighborhood V then we find $z \in V$ s.t. $z \sim x$, but $z \sim y$ hence $x \sim z \sim y$ and we just proved that the equivalence class of x was closed. \square

We can now prove an alternative formula, due to Brunetti, Fredenhagen and Ribeiro [17].

Lemma 3.3. *For every Bastiani smooth function $F : U \rightarrow \mathbb{K}$, with U a connected open subset of $C^\infty(M)$:*

$$\text{supp}(F) = \overline{\bigcup_{\varphi \in U} \text{supp } F_\varphi^{(1)}}. \quad (7)$$

Proof. Using the result of Lemma 3.2, we may reduce to the case where U is an open convex set. We prove that both sets $\overline{\bigcup_{\varphi} \text{supp } F_\varphi^{(1)}}$ and $\text{supp } F$ as defined in Def. 3.1 have identical complements.

Indeed, for every point $x \in M$, $x \notin \text{supp } F$ means by definition of the support that there exists an open neighborhood Ω of x such that $\forall(\psi, \varphi) \in \mathcal{D}(\Omega) \times C^\infty(M)$, $F(\varphi + \psi) = F(\varphi)$. It follows that for all $\psi \in \mathcal{D}(\Omega)$, there exists $\varepsilon > 0$ such that $|t| \leq \varepsilon \implies \varphi + t\psi \in U$ and $t \in [-\varepsilon, \varepsilon] \mapsto F(\varphi + t\psi)$ is a constant function of t therefore $\frac{dF(\varphi+t\psi)}{dt}|_{t=0} = DF_\varphi(\psi) = 0$. This means that for all $\varphi \in U$, the support of $DF_\varphi \in \mathcal{E}'(M)$ does not meet Ω hence Ω lies in the complement of $\cup_{\varphi \in U} \text{supp } (DF_\varphi)$ and therefore $x \in \Omega$ does not meet the closure $\overline{\cup_{\varphi \in U} \text{supp } (DF_\varphi)}$.

Conversely if x does not meet the closure $\overline{\cup_{\varphi \in U} \text{supp } (DF_\varphi)}$, then there is some neighborhood Ω of x which does not meet $\cup_{\varphi \in U} \text{supp } (DF_\varphi)$ therefore for all $(\varphi, \psi) \in U \times \mathcal{D}(\Omega)$ s.t. $\varphi + \psi \in U$, the whole straight path $[\varphi, \varphi + \psi]$ lies in U (by convexity of U) hence

$$\forall t \in [0, 1], DF_{\varphi+t\psi}(\psi) = 0 \implies \int_0^1 dt DF_{\varphi+t\psi}(\psi) = 0,$$

and by the fundamental theorem of calculus $F(\varphi + \psi) = F(\varphi)$. \square

Now we show that any smooth functional is locally compactly supported:

Proposition 3.4. *Let $F : U \mapsto \mathbb{K}$ be a Bastiani smooth function, where U is an open connected subset of $E = C^\infty(M)$. For every $\varphi \in U$, there is a neighborhood V of φ in U and a compact subset $K \subset M$ such that the support of F restricted to V is contained in K . Moreover for all integers n and all $\varphi \in U$, the distributional support $\text{supp } (D^n F_\varphi)$ is contained in $K^n \subset M^n$.*

Proof. By definition of the support of a functional, it is enough to show that, for every $\varphi \in V$, $\text{supp } DF_\varphi \subset K$. If F is smooth, then $DF : U \times E \rightarrow \mathbb{K}$ is continuous. Thus, it is continuous in the neighborhood of $(\varphi_0, 0)$ for every $\varphi_0 \in U$. In other words, for every $\epsilon > 0$, there is a neighborhood V of φ_0 , a seminorm $\pi_{m,K}$ and an $\eta > 0$ such that $|DF_\varphi(\chi)| < \epsilon$ for every $\varphi \in V$ and every $\chi \in E$ such that $\pi_{m,K}(\chi) < \eta$. Now, for every $\psi \in E$ such that $\pi_{m,K}(\psi) \neq 0$, we see that $\chi = \psi\eta/(2\pi_{m,K}(\psi))$ satisfies $\pi_{m,K}(\chi) < \eta$. Thus, $|DF_\varphi(\chi)| < \epsilon$ for every $\varphi \in V$ and, by linearity, $|DF_\varphi(\psi)| < (2\epsilon/\eta)\pi_{m,K}(\psi)$. On the other hand, if $\pi_{m,K}(\psi) = 0$, then for any $\mu > 0$ $\pi_{m,K}(\mu\psi) = 0 < \eta$, so that $|DF_\varphi(\mu\psi)| < \epsilon$. By linearity, $|DF_\varphi(\psi)| < \epsilon/\mu$ for any $\mu > 0$ and we conclude that $|DF_\varphi(\psi)| = 0$. Thus, for every $\varphi \in V$ and every $\psi \in E$,

$$|DF_\varphi(\psi)| \leq 2\frac{\epsilon}{\eta}\pi_{m,K}(\psi).$$

Of course, this inequality implies that $DF_\varphi(\psi) = 0$ when ψ is identically zero on the compact subset K .

Let us show that this implies that the support of DF_φ is contained in K . To avoid possible problems at the boundary, take any compact neighborhood K' of K . Now, take an open set Ω in M such that $\Omega \cap K' = \emptyset$. Then, for every smooth function ψ supported in Ω , we have $\pi_{m,K}(\psi) = 0$ because the seminorm $\pi_{m,K}$ takes only into account the points of K . As a consequence, the restriction of DF_φ to Ω is zero, which means that Ω is outside the support of DF_φ for every $\varphi \in V$. Thus, $\text{supp } F|_V \subset K$ because, for every $\varphi \in V$, the support of DF_φ is included in every compact neighborhood of K . Finally we show that, if F is compactly supported, then all $D^n F_\varphi$ are compactly supported with $\text{supp } D^n F_\varphi \subset (\text{supp } F)^{\times n}$. This is easily seen by the following cutoff function argument: if F is compactly supported then for every cutoff function χ equal to 1 on an arbitrary compact neighborhood of $\text{supp } F$, we have: $F(\varphi) = F(\chi\varphi), \forall \varphi \in E$. Then it is immediate by definition of $D^n F_\varphi$ that

$$D^n F_\varphi(\psi_1, \dots, \psi_n) = \frac{d^n F(\chi(\varphi + t_1\psi_1 + \dots + t_n\psi_n))}{dt_1 \dots dt_n} = F_{\chi\varphi}^{(n)}(\chi\psi_1, \dots, \chi\psi_n)$$

thus $\text{supp } D^n F_\varphi \subset \text{supp } \chi^{\times n}$ for all test function χ s.t. $\chi|_{\text{supp } F} = 1$ and therefore $\text{supp } D^n F_\varphi \subset (\text{supp } F)^{\times n}$ since $\bigcap_{\chi|_{\text{supp } F}=1} \text{supp } \chi = \text{supp } F$. \square

3.2 A multilinear kernel theorem with parameters.

We work with M a smooth manifold and $B \rightarrow M$ a smooth vector bundle of finite rank over M . Let $E = \Gamma(M, B)$ be its space of smooth sections and U an open subset of E . Recall that we consider smooth maps $F : E \mapsto \mathbb{K}$ where \mathbb{K} is the field \mathbb{R} or \mathbb{C} . In this section we relate the Bastiani derivatives $D^k F$, which are k -linear on $\Gamma(M, \mathcal{E})$ to the distributions used in quantum field theory, which are linear on $\Gamma(M^k, \mathcal{E}^{\boxtimes k})$. Since the k -th derivative $D^k F$ of a smooth map is multilinear and continuous in the last k variables, we can use the following result [28, p. 438].

Lemma 3.5. *Let E be a Hausdorff locally convex space. There is a canonical isomorphism between any k -linear map $f : E^k \rightarrow \mathbb{K}$ and the map $\bar{f} : E^{\otimes k} \rightarrow \mathbb{K}$, where \otimes is the algebraic tensor product, which is linear and defined as follows: if $\chi = \sum_j \chi_1^j \otimes \cdots \otimes \chi_k^j$ is a finite sum of tensor products, then*

$$\bar{f}(\chi) = \sum_j f(\chi_1^j, \dots, \chi_k^j). \quad (8)$$

A fundamental property of the completed projective tensor product $\hat{\otimes}_\pi$ is that f is (jointly) continuous iff \bar{f} , still defined by Eq. (8) but extended by continuity to $E^{\hat{\otimes}_\pi k}$, is continuous [28, p. 441]. In particular, if $E = C^\infty(M)$, then $E^{\hat{\otimes}_\pi k} = C^\infty(M^k)$ [28, p. 530], and \bar{f} becomes a compactly supported distribution on M^k . More generally, if $E = \Gamma(M, B)$, then E is Fréchet nuclear and $E^{\hat{\otimes}_\pi k} = \Gamma(M^k, \mathcal{E}^{\boxtimes k})$ [65, p. 72]. Thus, \bar{f} becomes a compactly supported distributional section on M^k . If $f = D^k F_\varphi$ we denote \bar{f} by $F_\varphi^{(k)}$.

Recall that, if U is an open subset of a Hausdorff locally convex space E , a map $F : U \rightarrow \mathbb{K}$ is smooth iff every $D^k F$ is continuous from $U \times E^k$ to \mathbb{K} . According to the previous discussion, continuity on E^k is equivalent to continuity on $E^{\hat{\otimes}_\pi k}$. Therefore, it is natural to wonder when joint continuity on $U \times E^k$ is equivalent to joint continuity on $U \times E^{\hat{\otimes}_\pi k}$. This is the subject of the next sections.

3.2.1 An equicontinuity Lemma.

We recall the definition of a generating family of seminorms of the completed projective tensor product on Fréchet spaces following [28, p. 435]. The definition starts with the algebraic tensor product $E^{\otimes k}$, for any $\psi \in E^{\otimes k}$, for every seminorm $p_1 \times \cdots \times p_k$ on E^k , there exists a seminorm $p_1 \otimes \cdots \otimes p_k$ on $E^{\otimes k}$ such that

$$p_1 \otimes \cdots \otimes p_k(\psi) = \inf_n \sum_n p_1(e_{1,n}) \cdots p_k(e_{k,n}),$$

where the infimum is taken over representations of ψ as finite sums: $\psi = \sum_n e_{1,n} \otimes \cdots \otimes e_{k,n}$. The projective topology on $E^{\otimes k}$ is generated by the family of seminorms $p_1 \otimes \cdots \otimes p_k$ where p_1, \dots, p_k are a generating family of seminorms of E [28, p. 435]. Since E is Fréchet, its topology is defined by a countable family of seminorms and it follows that the family of seminorms $p_1 \otimes \cdots \otimes p_k$ on $E^{\otimes k}$ is countable. Hence they can be used to construct a metric on $E^{\otimes k}$ which defines the same topology as the projective topology and $E^{\hat{\otimes}_\pi k}$ is defined as the completion of $E^{\otimes k}$ relative to this metric or equivalently with respect to the projective topology.

Lemma 3.6. *Let E be a Fréchet space, U open in E , and $F : U \times E^k \mapsto \mathbb{K}$ a continuous map, multilinear in the last k variables. Then for every $\varphi_0 \in U$, there exist a neighborhood V of φ_0 , a seminorm q of $E^{\hat{\otimes}_\pi k}$ and a constant $C > 0$ such that*

$$\forall \varphi \in V, \forall \psi \in E^{\hat{\otimes}_\pi k}, \quad |\bar{F}(\varphi, \psi) - \bar{F}(\varphi_0, \psi)| \leq C q(\psi).$$

Proof. By continuity of $F : U \times E^k \mapsto \mathbb{K}$, for every $\epsilon > 0$, there exist a neighborhood V of φ_0 and neighborhoods U_1, \dots, U_k of zero such that $\varphi \in V$ and $e_i \in U_i$ for $i = 1, \dots, k$ imply $|F(\varphi, e_1, \dots, e_k)| \leq \epsilon$. Since E is locally convex, there are continuous seminorms p_1, \dots, p_k on E and strictly positive numbers η_1, \dots, η_k such that $e_i \in U_i$ if $p_i(e_i) \leq \eta_i$. Consider now arbitrary

elements e_1, \dots, e_k of E such that $p_i(e_i) \neq 0$. Then, if $f_i = e_i \eta_i / p_i(e_i)$ we have $p_i(f_i) \leq \eta_i$. Thus, $|F(\varphi, f_1, \dots, f_k)| < \epsilon$ and, by multilinearity, $|F(\varphi, e_1, \dots, e_k)| < M p_1(e_1) \dots p_k(e_k)$ where $M = \epsilon / (\eta_1, \dots, \eta_k)$. The same argument as in the proof of Proposition 3.4 shows that $|F(\varphi, e_1, \dots, e_k)| = 0$ if $p_i(e_i) = 0$. Therefore, for every $\varphi \in V$ and every e_1, \dots, e_k in E , $|F(\varphi, e_1, \dots, e_k)| \leq M p_1(e_1) \dots p_k(e_k)$. By defining $C = 2M$ we obtain

$$\forall \varphi \in V, \forall (e_1, \dots, e_k) \in E^k, |F(\varphi, e_1, \dots, e_k) - F(\varphi_0, e_1, \dots, e_k)| \leq C p_1(e_1) \dots p_k(e_k). \quad (9)$$

By definition of \overline{F} , for all $(\varphi, \psi) \in V \times E^{\otimes k}$:

$$\begin{aligned} |\overline{F}(\varphi, \psi) - \overline{F}(\varphi_0, \psi)| &\leq \sum_n |F(\varphi, e_{1,n}, \dots, e_{k,n}) - F(\varphi_0, e_{1,n}, \dots, e_{k,n})| \\ &\leq C \sum_n p_1(e_{1,n}) \dots p_k(e_{k,n}), \end{aligned}$$

for all representations of ψ as finite sum $\psi = \sum_n e_{1,n} \otimes \dots \otimes e_{k,n}$. Taking the infimum over such representations yields the estimate:

$$\forall (\varphi, \psi) \in V \times E^{\otimes k}, |\overline{F}(\varphi, \psi) - \overline{F}(\varphi_0, \psi)| \leq C q(\psi), \quad (10)$$

for the seminorm $q = p_1 \otimes \dots \otimes p_k$ on $E^{\otimes \pi^k}$ and the above inequality extends to any ψ of $E^{\hat{\otimes} \pi^k}$ since, in a Fréchet space, ψ can be approximated by a convergent sequence of elements in $E^{\otimes k}$ by density of $E^{\otimes k}$ in $E^{\hat{\otimes} \pi^k}$ and by continuity of the seminorm $p_1 \otimes \dots \otimes p_k$ for the topology of $E^{\hat{\otimes} \pi^k}$. \square

Another way to state the previous result is to say that the family of linear maps $\{F(\varphi, \cdot); \varphi \in V\}$ is equicontinuous [57, p. II.6].

3.2.2 Proof of the main result.

We are now ready to prove

Proposition 3.7. *Let E be a Fréchet space and $U \subset E$ an open subset. Then $F : U \times E^k \mapsto \mathbb{K}$, multilinear in the last k variables, is jointly continuous iff the corresponding map $\overline{F} : U \times E^{\hat{\otimes} \pi^k} \rightarrow \mathbb{K}$ is jointly continuous.*

Proof. One direction of this theorem is straightforward and holds if E is any locally convex space. Indeed, by definition of the projective tensor product, the canonical multilinear mapping $E^k \rightarrow E^{\hat{\otimes} k}$ is continuous [28, p. 434]. Therefore, if \overline{F} is continuous on $U \times E^{\hat{\otimes} k}$ then, by composition with the canonical multilinear mapping, F is continuous on $U \times E^k$.

Let us prove that the continuity of F implies the continuity of \overline{F} . According to Lemma 2.4, we have to show that, for every $\varphi_0 \in U$ and every $\epsilon > 0$, there exist a finite number of continuous seminorms q_i on $E^{\hat{\otimes} \pi^k}$, a neighborhood V of φ_0 and $\eta_i > 0$ such that, if φ belongs to V and $\psi \in E^{\hat{\otimes} \pi^k}$ satisfies $q_i(\psi - \psi_0) \leq \eta_i$, then $|\overline{F}(\varphi, \psi) - \overline{F}(\varphi_0, \psi_0)| \leq \epsilon$.

In order to bound $\overline{F}(\varphi, \psi) - \overline{F}(\varphi_0, \psi_0)$, we cut it into three parts

$$\begin{aligned} \overline{F}(\varphi, \psi) - \overline{F}(\varphi_0, \psi_0) &= \overline{F}(\varphi, \psi_k) - \overline{F}(\varphi_0, \psi_k) \\ &+ \overline{F}(\varphi, \psi - \psi_k) - \overline{F}(\varphi_0, \psi - \psi_k) \\ &+ \overline{F}(\varphi_0, \psi) - \overline{F}(\varphi_0, \psi_0), \end{aligned} \quad (11)$$

where ψ_k is some element of the algebraic tensor product $E^{\otimes k}$ close enough to ψ_0 that we choose now. The equicontinuity Lemma 3.6 yields a neighborhood V_2 of φ_0 , a constant $C > 0$ and a continuous seminorm q_2 on $E^{\hat{\otimes} \pi^k}$ so that

$$\forall \varphi \in V_2, \forall \psi \in E^{\hat{\otimes} \pi^k}, |\overline{F}(\varphi, \psi) - \overline{F}(\varphi_0, \psi)| \leq C q_2(\psi).$$

Now we use the fact that the algebraic tensor product $E^{\otimes k}$ is everywhere dense in $E^{\hat{\otimes} \pi^k}$ hence there is some element ψ_k in the algebraic tensor product $E^{\otimes k}$ such that $q_2(\psi_0 - \psi_k) \leq \eta_2$ with $\eta_2 := \frac{\epsilon}{6C}$.

Now that ψ_k is chosen, we can bound the second term of the sum (11), namely $\overline{F}(\varphi, \psi - \psi_k) - \overline{F}(\varphi_0, \psi - \psi_k)$. From the previous relation, for every $\varphi \in V_2$ and every ψ such that $q_2(\psi - \psi_0) \leq \eta_2$, the triangle inequality for $q_2(\psi - \psi_k)$ gives us

$$|\overline{F}(\varphi, \psi - \psi_k) - \overline{F}(\varphi_0, \psi - \psi_k)| \leq C(q_2(\psi - \psi_0) + q_2(\psi_0 - \psi_k)) \leq \frac{\varepsilon}{3}.$$

We continue by bounding the last term $\overline{F}(\varphi_0, \psi) - \overline{F}(\varphi_0, \psi_0)$ in the sum (11). Since φ_0 is fixed, the map $\psi \mapsto \overline{F}(\varphi_0, \psi)$ is continuous in ψ because, since $\overline{F}(\varphi_0, \cdot)$ is continuous on $E^{\otimes \pi k}$, its extension to the completion $E^{\hat{\otimes} \pi k}$, also denoted by $\overline{F}(\varphi_0, \cdot)$, is continuous. It follows that there is some seminorm q_1 of $E^{\hat{\otimes} \pi k}$ and a number $\eta_1 > 0$ such that if $\psi \in U$ satisfies $q_1(\psi - \psi_0) \leq \eta_1$ then

$$|\overline{F}(\varphi_0, \psi) - \overline{F}(\varphi_0, \psi_0)| \leq \frac{\varepsilon}{3}.$$

To bound the first term $\overline{F}(\varphi, \psi_k) - \overline{F}(\varphi_0, \psi_k)$ in the sum (11), we use the fact that $\psi_k \in E^{\otimes k}$. Thus, $\psi_k = \sum_{j=1}^p (e_{1,j} \otimes \cdots \otimes e_{k,j})$ for some $(e_{1,j}, \dots, e_{k,j}) \in E^k$. By definition of \overline{F} ,

$$\overline{F}(\varphi, \psi_k) - \overline{F}(\varphi_0, \psi_k) = \sum_{j=1}^p F(\varphi, e_{1,j}, \dots, e_{k,j}) - F(\varphi_0, e_{1,j}, \dots, e_{k,j}).$$

By continuity of F in the first factor, the finite sum $\sum_{j=1}^p F(\varphi, e_{1,j}, \dots, e_{k,j})$ is continuous in φ and there is some neighborhood V_3 of φ_0 such that for all $\varphi \in V_3$ the following bound

$$\left| \sum_{j=1}^p F(\varphi, e_{1,j}, \dots, e_{k,j}) - F(\varphi_0, e_{1,j}, \dots, e_{k,j}) \right| \leq \frac{\varepsilon}{3},$$

holds true.

Finally we found some neighborhood $V = V_2 \cap V_3$ of φ_0 , two seminorms q_1 and q_2 of $E^{\hat{\otimes} \pi k}$, and two numbers $\eta_1 > 0$ and $\eta_2 = \varepsilon/6C$ such that $q_1(\psi - \psi_0) < \eta_1$ and $q_2(\psi - \psi_0) < \eta_2$ imply

$$|\overline{F}(\varphi, \psi) - \overline{F}(\varphi_0, \psi_0)| \leq \varepsilon.$$

The proposition is proved. \square

Now we can specialize our result to the space of smooth sections of vector bundles. We recall a fundamental result on the projective tensor product of sections [65, p. 72]:

Proposition 3.8. *Let $\Gamma(M, B)$ be the space of smooth sections of some smooth finite rank vector bundle $B \rightarrow M$ on a manifold M . Then $\Gamma(M, B)^{\hat{\otimes} \pi k} = \Gamma(M^k, B^{\boxtimes k})$.*

Note that we could remove the index π in $\hat{\otimes}_\pi$ because we saw that $\Gamma(M, B)$ is nuclear. If we specialize Proposition 3.7 to sections of vector bundles (which is a Fréchet space) we obtain

Theorem 3.9. *Let $E = \Gamma(M, B)$ be the space of smooth sections of some smooth finite rank vector bundle $B \rightarrow M$. Then $F : U \times E^k \rightarrow \mathbb{K}$ multilinear in the last k variables is jointly continuous iff the corresponding map $\overline{F} : U \times \Gamma(M^k, B^{\boxtimes k}) \rightarrow \mathbb{K}$ is jointly continuous.*

Proof. The proof is an immediate consequence of the fact that $E^{\hat{\otimes} \pi k} = \Gamma(M^k, B^{\boxtimes k})$ and Proposition 3.7. \square

The definition of a Bastiani smooth functional implies the following corollary:

Theorem 3.10. *Let $E = \Gamma(M, B)$ be the space of smooth sections of some smooth finite rank vector bundle $B \rightarrow M$. A map $F : U \rightarrow \mathbb{K}$, where U is open in E , is Bastiani smooth iff the maps $F^{(k)} : U \times \Gamma(M^k, B^{\boxtimes k}) \rightarrow \mathbb{K}$ are (jointly) continuous for every $k \geq 1$.*

To interpret Theorem 3.7 in terms of distributional kernels, let $B \rightarrow M$ denote a smooth vector bundle of finite rank over a manifold M equipped with a fixed density $|dx|$ and $B^* \rightarrow M$ the corresponding dual bundle. Recall that $\Gamma(M, B)' \simeq \Gamma(M, B^*) \otimes_{C^\infty(M)} \mathcal{E}'(M)$ [66], where $\Gamma(M, B^*) \otimes_{C^\infty(M)} \mathcal{E}'(M)$ denotes the compactly supported distributional sections of the dual bundle B^* . In global analysis, to every continuous linear map $L : \Gamma(M, B) \rightarrow \Gamma(M, B)'$, we associate the continuous bilinear map $B : (\varphi_1, \varphi_2) \in \Gamma(M, B)^2 \mapsto \langle \varphi_1, L\varphi_2 \rangle \in \mathbb{K}$ where the pairing is understood as a pairing between a smooth function and a distribution once the smooth density on M is fixed.

The usual kernel theorem of the theory of distributions, states that a bilinear map can be represented by a distribution: $K_L \in \mathcal{E}'(M \times M) \otimes_{C^\infty(M^2)} \Gamma(M^2, B^* \boxtimes B^*)$ living on configuration space M^2 such that

$$\langle K_L, \varphi_1 \boxtimes \varphi_2 \rangle_{\Gamma(M^2, B \boxtimes B)', \Gamma(M^2, B \boxtimes B)} = \langle \varphi_1, L\varphi_2 \rangle_{\Gamma(M, B), \Gamma(M, B)'}$$

for every $(\varphi_1, \varphi_2) \in \Gamma(M, B)^2$. Theorem 3.7 generalizes the kernel theorem by using multilinear maps instead of bilinear ones and by considering that these multilinear maps depend continuously and non-linearly on a parameter φ .

3.3 Order of distributions

If F is a Bastiani smooth map from an open subset U of $E = C^\infty(M)$ to \mathbb{K} , then, for every $\varphi \in U$, $D^k F_\varphi$ is a compactly supported distribution. Therefore, the order of $F_\varphi^{(k)}$ is finite [24, p. 88]. For some applications, for example to local functionals, it is important to require the order of $F_\varphi^{(k)}$ to be *locally bounded*:

Proposition 3.11. *Let $E = C^\infty(M)$ and $F : E \rightarrow \mathbb{K}$ be a smooth functional on an open subset U of E . Then, for every $\varphi_0 \in U$ and every integer k , there is a neighborhood V of φ_0 , an integer m and a compact $K \subset M^k$ such that, for every $\varphi \in V$, the order of $F_\varphi^{(k)}$ is smaller than m and $F_\varphi^{(k)}$ is supported in K .*

Proof. According to Lemma 3.6, for every φ_0 in U , there is a neighborhood V of φ_0 , a constant C and a seminorm $\pi_{n,K}$ of $C^\infty(M)$ such that

$$|F_\varphi^{(k)}(\psi) - F_{\varphi_0}^{(k)}(\psi)| \leq C \pi_{n,K}(\psi).$$

This means that the order of $F_\varphi^{(k)} - F_{\varphi_0}^{(k)}$ is bounded by n [24, p. 64], and the order of $F_\varphi^{(k)}$ is bounded by n plus the order of $F_{\varphi_0}^{(k)}$. Moreover, if $\text{supp } \psi \cap K = \emptyset$, then $\pi_{n,K}(\psi) = 0$ and $F_\varphi^{(k)}(\psi) - F_{\varphi_0}^{(k)}(\psi) = 0$. This means that the support of $F_\varphi^{(k)} - F_{\varphi_0}^{(k)}$ is contained in K and the support of $F_\varphi^{(k)}(\psi)$ is contained in the compact $K \cup \text{supp } F_{\varphi_0}^{(k)}$. \square

Note also that, in general, the order of the distributions is not bounded on U :

Lemma 3.12. *Let $g \in \mathcal{D}(\mathbb{R})$ and $(\chi_n)_{n \in \mathbb{Z}}$ a sequence of functions such that $\chi_n \in \mathcal{D}([n-1, n+1])$ and $\sum_{n \in \mathbb{Z}} \chi_n = 1$. Then, the functional*

$$F(\varphi) = \sum_{n=-\infty}^{\infty} \int_{\mathbb{R}} \chi_n(\varphi(x)) \frac{d^{|n|} \varphi}{dx^{|n|}}(x) g(x) dx,$$

is Bastiani smooth on $C^\infty(\mathbb{R})$ but the order of $F^{(k)}$ is not bounded on $C^\infty(\mathbb{R})$.

Proof. The functional F is smooth because, for every $\varphi_0 \in C^\infty(\mathbb{R})$, we can define a neighborhood of φ_0 by $V = \{\varphi; \pi_{0,K}(\varphi - \varphi_0) < \epsilon\}$, where K is a compact neighborhood of the support of g . Let N be the smallest integer strictly greater than $\pi_{0,K}(\varphi_0) + \epsilon$. Then, $-N < \varphi(x) < N$ for every $\varphi \in V$ and every $x \in K$ and

$$F(\varphi) = \sum_{n=-N-1}^{N+1} \int \chi_n(\varphi(x)) \varphi^{(|n|)}(x) g(x) dx,$$

is a finite sum of smooth functionals.

However, the order of

$$F_\varphi^{(1)}(\psi) = \sum_{n=-\infty}^{\infty} \int \left(\chi_n(\varphi(x))\psi^{(|n|)}(x) + \chi'_n(\varphi(x))\psi^{(1)}(x)\varphi^{(|n|)}(x) \right) g(x)dx,$$

is not bounded on $C^\infty(\mathbb{R})$. Indeed, for any positive integer n , we can find a smooth function φ such that $\chi_n(\varphi(x))g(x) \neq 0$ for some $x \in \text{supp } g$. Since $F_\varphi^{(1)}(\psi)$ contains a factor $\psi^{(n)}(x)$ it is at least of order n . \square

3.4 Derivatives as smooth functionals

In the next section we equip several spaces of functionals with a topology. As a warm-up exercise, we show here that the maps $F^{(k)}$ are smooth functionals from $C^\infty(M)$ to $\mathcal{E}'(M^k)$.

We adapt to the case of functionals the general result given in item 4 of Prop. 2.13 stating that, if F is a smooth functional on U , then $D^k F$ is a Bastiani smooth map from U to $L(E^k, \mathbb{K})$. We need to identify the topology of $L(E^k, \mathbb{K})$ used by Bastiani. Let us start with $L(E, \mathbb{K})$. Bastiani furnishes E with the topology of convergence on all compact sets of E . In other words, the seminorms that define the topology of $L(E, \mathbb{K})$ are $p_C(u) = \sup_{f \in C} |\langle u, f \rangle|$, where C runs over the compact subsets of $C^\infty(M)$. Since $C^\infty(M)$ is a Montel space [27, p. 239], the topology of uniform convergence on compact sets is the same as the strong topology [27, p. 235]. This means that $L(E, \mathbb{R})$ is the space $\mathcal{E}'(M)$ of compactly supported distributions with its usual topology. Similarly, $L(E^k, \mathbb{R})$ can be identified to a subset of $\mathcal{E}'(M^k)$ with its usual topology. We just obtained the following result:

Proposition 3.13. *Let U be an open subset of $C^\infty(M)$ and $F : U \rightarrow \mathbb{K}$ a Bastiani smooth functional. Then, for every integer k , the map $F^{(k)} : U \rightarrow \mathcal{E}'(M^k)$ is smooth in the sense of Bastiani.*

4 Topologies on spaces of functionals

We need to define a topology on the various spaces of functionals used in quantum field theory. The generally idea is to define seminorms on F and its derivatives $F^{(k)}$. The topology proposed by Brunetti, Dütsch and Fredenhagen [15] is the initial topology of all the maps $F \rightarrow F_\varphi^{(k)}$, where each $F_\varphi^{(k)}$ belongs to a nuclear space determined by a wavefront set condition. This topology is nuclear, but the absence of a control of the dependence on φ makes it generally not complete. We then describe Bastiani's topology, which is complete but has two drawbacks: it does not take wavefront set conditions into account and it is generally not nuclear. Finally we shall describe the family of topologies proposed by Dabrowski [18] which are both nuclear and complete.

4.1 Bastiani's topology

Bastiani defines several topologies on the space of Bastiani smooth maps between two locally convex spaces [37, p. 65]. For the case of functionals, we consider the topology defined by the following seminorms:

$$\begin{aligned} p_{C_0}(F) &= \sup_{\varphi \in C_0} |F(\varphi)|, \\ p_{C_0, C_1, \dots, C_k}(F) &= \sup_{(\varphi, h_1, \dots, h_k) \in C_0 \times C_1 \times \dots \times C_k} |D^k F_\varphi(h_1, \dots, h_k)|, \end{aligned}$$

where C_i runs over the compact sets of $\Gamma(M, B)$. By using Bastiani's results [37, pp. 66] we obtain

Proposition 4.1. *Let $B \xrightarrow{\pi} M$ be a finite rank vector bundle over the manifold M and $\Gamma(M, B)$ be the space of smooth sections of B . Then, with the seminorms defined above, the space of smooth functionals on $\Gamma(M, B)$ is a complete locally convex space.*

A similar topology was used by Glöckner [67, p. 367] and Wockel [68, p. 29] [69, p. 12].

4.2 Nuclear and complete topologies

Quantum field theory uses different spaces of functionals defined by conditions on the wave front set of $F_\varphi^{(k)}$. Recall that the wave front set describes the points and the directions of singularity of a distribution [70]. Yoann Dabrowski [18] recently described nuclear and complete topologies for spaces of functionals with wave front set conditions. We present some of his topologies for several common spaces of functionals.

Dabrowski's definition differs from Bastiani in two respects. To describe the first difference, recall that, according to Proposition 3.13, if $F : U \rightarrow \mathbb{R}$ is a smooth functional, then the derivatives $F^{(k)} : U \rightarrow \mathcal{E}'(M^k)$ are smooth functionals. To add the wave front set conditions, Dabrowski requires $F^{(k)}$ to be smooth from U to $\mathcal{E}'_{\Gamma_k}(M^k)$, which is the space of compactly supported distributions whose wave front sets are included in Γ_k , a cone in T^*M^k . In fact, Dabrowski supplements this definition with a more refined wave front set (the dual wave front set) which enables him to equip $\mathcal{E}'_{\Gamma_k}(M^k)$ with a Montel, complete, ultrabornological and nuclear topology. He also considers support conditions which are different from compact.

To describe the second difference, recall that Bastiani's topology gives a locally convex space which is complete. However it is generally not nuclear. This is due to a theorem by Colombeau and Meise [71] which says, broadly speaking, that a function space over a Fréchet space cannot be nuclear for the topology of convergence over some balanced, convex, compact sets of infinite dimension. To avoid that problem, the variable φ is made to run over *finite dimensional* compact sets. More precisely, Dabrowski considers compact sets in \mathbb{R}^m for any finite value of m and smooth maps f from \mathbb{R}^m to an open subset of $C^\infty(M)$. He defines two families of seminorms:

$$p_{f,K}(F) = \sup_{\varphi \in f(K)} |F(\varphi)|, \quad (12)$$

$$p_{n,f,K,C}(F) = \sup_{\varphi \in f(K)} \sup_{v \in C} |\langle F_\varphi^{(n)}, v \rangle|, \quad (13)$$

where K is a compact subset of \mathbb{R}^m for some m and C is an equicontinuous subset of the dual of the space of distributions to which $F_\varphi^{(n)}$ belong. Dabrowski proved that, with this family of seminorms, the space of functionals F is a complete locally convex nuclear space [19].

We describe now several types of functionals that have been used in the literature and we specify more precisely their topologies.

4.3 The regular functionals

A polynomial functional of the form

$$F(\varphi) = \sum_n \int_{M^n} dx_1 \dots dx_n f_n(x_1, \dots, x_n) \varphi(x_1) \dots \varphi(x_n),$$

where the sum over n is finite and $f_n \in \mathcal{D}(M^n)$, is called a *regular functional* [72], because all its derivatives are smooth functions [73], i. e. the wave front set of $F_\varphi^{(k)}$ is empty. More generally, we define the space $F_{\text{reg}}(M)$ of regular functionals to be the set of Bastiani smooth functionals F such that $\text{WF}(F^{(n)}) = \emptyset$ for every $n > 0$. Thus, $F^{(n)} \in \mathcal{E}'_\emptyset(M^n) = \mathcal{D}(M^n)$ and the sets C in Equation (13) are the equicontinuous sets of $\mathcal{D}'(M^n)$. By a general theorem [27, p. 200], the topology of uniform convergence on the equicontinuous sets of $\mathcal{D}'(M^n)$ is equivalent to the topology given by the seminorms of its dual $\mathcal{D}(M^n)$. In other words, the topology of $F_{\text{reg}}(M)$ is defined by the seminorms [19]:

$$p_{f,K}(F) = \sup_{\varphi \in f(K)} |F(\varphi)|, \quad (14)$$

$$p_{n,f,K,\alpha}(F) = \sup_{\varphi \in f(K)} p_{\alpha,n}(F_\varphi^{(n)}), \quad (15)$$

where $p_{\alpha,n}$ runs over a defining family of seminorms of $\mathcal{D}(M^n)$ [74]. With this topology, $F_{\text{reg}}(M)$ is nuclear and complete.

Note that the tensor product of elements of $\mathcal{D}(M^m)$ with elements of $\mathcal{D}(M^n)$ is not continuous in $\mathcal{D}(M^{m+n})$ [75]. Thus, the product in $\mathbf{F}_{\text{reg}}(M)$ is hypocontinuous but not continuous.

4.4 The microcausal functionals

It is possible to describe quantum field theory (up to renormalization) as the deformation quantization of classical field theory [23]. For the deformation quantization of the product FG of two functionals to first order in \hbar , we need to evaluate $\langle DF_\varphi \otimes DG_\varphi, \Delta_+ \rangle$, where Δ_+ is a singular distribution (the Wightman propagator) and $\langle \cdot, \cdot \rangle$ is an extension of the duality pairing between distributions and test functions [76]. For this pairing to be meaningful to all orders in \hbar , the wave front set of Δ_+ imposes that the wave front set of $D_\varphi^{(k)}$ must not meet the cone Γ_k defined as follows [23].

Let M be a Lorentzian manifold with pseudo-metric g . Let V_x^+ (resp. V_x^-) be the set of $(x; \xi) \in T_x^*M$ such that $g^{\mu\nu}(x)\xi_\mu\xi_\nu \geq 0$ and $\xi_0 \geq 0$ (resp. $\xi_0 \leq 0$), where we assume that $g^{00} > 0$. We define the closed cone

$$\Gamma_k = \{(x_1, \dots, x_k; \xi_1, \dots, \xi_k) \in \dot{T}^*M^n; (\xi_1, \dots, \xi_n) \in (V_{x_1}^+ \times \dots \times V_{x_n}^+) \cup (V_{x_1}^- \times \dots \times V_{x_n}^-)\},$$

where \dot{T}^*M^n is the cotangent bundle T^*M^n without its zero section. The space \mathbf{F}_{mc} of microcausal functionals was originally defined as the set of Bastiani smooth functionals such that $F_\varphi^{(n)} \in \mathcal{E}'_{\Xi_n}(M^n)$ for every φ , where $\Xi_n = \dot{T}^*M^n \setminus \Gamma_n$ is an open cone [23, 77, 15, 78, 14, 17, 72].

However, the space $\mathcal{E}'_{\Xi_n}(M^n)$ being not even sequentially complete [76], it is not suitable to define a complete space of functionals. Therefore, Dabrowski defines the space \mathbf{F}_{mc} of microcausal functionals to be the set of Bastiani smooth functionals such that $F_\varphi^{(n)} \in \mathcal{E}'_{\overline{\Xi}_n, \Xi_n}(M^n)$, which is the completion of $\mathcal{E}'_{\Xi_n}(M^n)$. Dabrowski proved that $\mathcal{E}'_{\overline{\Xi}_n, \Xi_n}(M^n)$ is the set of compactly supported distributions $u \in \mathcal{E}'(M^n)$ such that the *dual wavefront set* of u is in Ξ_n and the wavefront set of u is in its closure $\overline{\Xi}_n$ (see [18] for a precise definition of these concepts and of the topology). This completion is not only complete, but even Montel and nuclear [18]. According to the general results of Ref. [18], the sets C are now equicontinuous sets of the bornologification of the normal topology of \mathcal{D}'_{Γ_n} . However, it was shown [18] that these equicontinuous sets are the same as the bounded sets of \mathcal{D}'_{Γ_n} with its normal topology. Therefore, the sets C are the well-known bounded sets of \mathcal{D}'_{Γ_n} [76].

With this topology, the space \mathbf{F}_{mc} is a complete nuclear algebra with hypocontinuous product.

4.5 Local functionals

As discussed in the introduction, local functionals are the basic building block (Lagrangian) of quantum field theory. We shall see that local functionals are a closed subset of the set of smooth functionals such that $F_\varphi^{(1)}$ can be identified with an element of $\mathcal{D}(M)$ that we denote by ∇F_φ and the wave front set of $F_\varphi^{(k)}$ is included in the conormal C_k of $D_k = \{(x_1, \dots, x_k) \in M^k; x_1 = \dots = x_k\}$. Recall that the conormal of D_k is the set of $(x_1, \dots, x_k; \xi_1, \dots, \xi_k) \in T^*M^k$ such that $x_1 = \dots = x_k$ and $\xi_1 + \dots + \xi_k = 0$.

Since the additivity property (defined in section 5.3) of local functionals complicates the matter, we follow Dabrowski [18] and, for any open set $\Omega \subset M$, we first define $\mathbf{F}_C(\Omega)$ to be the set of smooth maps such that $\varphi \mapsto \nabla F_\varphi$ is Bastiani smooth from $C^\infty(\Omega)$ to $\mathcal{D}(M)$ and, for every integer k , $\varphi \mapsto F_\varphi^{(k)}$ is Bastiani smooth from $C^\infty(\Omega)$ to $\mathcal{E}'_{C_k}(M^k)$ (we do not need to index $\mathcal{E}'(M^k)$ with two cones because C_k is closed [18]). The set $\mathbf{F}_{\text{loc}}(\Omega)$ of local functionals is then the subset of $\mathbf{F}_C(\Omega)$ satisfying the additivity condition.

The topology of \mathbf{F}_C is induced by the family of seminorms given by Eq. (15) that depend on the equicontinuous sets of the dual $\mathcal{D}'_{\Lambda_k, \overline{\Lambda}_k}(M^k)$ of $\mathcal{D}'_{C_k}(M^k)$, where $\Lambda_k = \dot{T}^*M^k \setminus C_k$. They were determined by Dabrowski [18, Lemma 28]:

Proposition 4.2. *A subset B of $\mathcal{D}'_{\Lambda_k}(M^k)$ is equicontinuous if and only if there is a closed cone $\Gamma \subset \Lambda_k$ such that $\text{WF}(u) \subset \Gamma$ for every $u \in B$ and B is bounded in $\mathcal{D}'_\Gamma(M^k)$.*

The bounded sets of $\mathcal{D}'_{\Gamma}(M^k)$ are characterized in detail in Ref. [76]. The topology of $\mathcal{D}'_{\Lambda_k}(M^k)$, where Λ_k is open, can be described as a non-countable inductive limit as follows. Write the complement $\Lambda_k^c = \cup \Gamma_n$, where each Γ_n is a compactly supported closed set. We write the open set Γ_n^c as a countable union of closed sets $\Gamma_n^c = \cup_m \Lambda_{n,m}$, so that $\Gamma_n = \cap_m \Lambda_{n,m}^c$ and $\Lambda_k^c = \cup_n \cap_m \Lambda_{n,m}^c$ is a countable union of countable intersections of open sets. We obtain $\Lambda_k = \cap_n \cup_m \Lambda_{n,m}$. We define for a sequence α the closed set $\Pi_\alpha = \cap_n \Lambda_{n,\alpha(n)}$, such that $\alpha \leq \beta$ implies $\Pi_\alpha \subset \Pi_\beta$. Then $\Lambda = \cup_\alpha \Pi_\alpha$ is a non-countable inductive limit of closed cones from which we can define the topology of $\mathcal{D}'_{\Lambda_k}(M^k)$ as a non-countable inductive limit of $\mathcal{D}'_{\Pi_\alpha}(M^k)$.

The space F_C furnished with the topology induced by the seminorms defined by Eqs. (12) and (13) is complete and nuclear. The space F_{loc} of local functionals is the closed subset of F_C defined by the additivity condition defined in the next section. As a closed subspace of a nuclear complete space, the space of local functionals is nuclear and complete.

Further examples of spaces of functionals are given by Dabrowski [19].

5 Additivity

The characterization of local functionals is a long-standing mathematical problem. According to Rao [79], the first criterium was proposed by Pinsker in 1938 and called *partial additivity* [80]. This criterium is also used in physics, but we shall see that it is not what we need by exhibiting a partially additive functional which is not local. Then, we shall discuss a more stringent criterium which is exactly what we need.

5.1 Partial additivity

When looking for an equation to characterize functionals having the form of Eq. (1), one can make the following observation. Let φ_1 and φ_2 be two smooth functions with disjoint support K_1 and K_2 and assume that $f(x, \varphi(x), \dots) = 0$ if $\varphi = 0$ on a neighborhood of x [80], so that $F(0) = 0$. Then, since the support of $\varphi_1 + \varphi_2$ is included in $K_1 \cup K_2$,

$$\begin{aligned} F(\varphi_1 + \varphi_2) &= \int_{K_1} dx f(x, \varphi_1(x) + \varphi_2(x), \dots) + \int_{K_2} dx f(x, \varphi_1(x) + \varphi_2(x), \dots) \\ &= \int_{K_1} dx f(x, \varphi_1(x), \dots) + \int_{K_2} dx f(x, \varphi_2(x), \dots) = F(\varphi_1) + F(\varphi_2). \end{aligned}$$

Therefore, it is tempting to use the condition of locality:

$$F(\varphi_1 + \varphi_2) = F(\varphi_1) + F(\varphi_2), \quad (16)$$

for φ_1 and φ_2 with disjoint support and functionals F such that $F(0) = 0$. And indeed, many authors since 1938, including Gelfand and Vilenkin [81, p. 275], used condition (16), but with disjoint support replaced by $\varphi_1 \varphi_2 = 0$ and smooth functions by measurable functions (see [79] for a review). In perturbative quantum field theory, partial additivity in our sense is also used when the function f in Eq. (1) is polynomial [82, 15, 83] because, in that case, partial additivity is equivalent to locality in the sense of Eq. (1) [15].

However, this definition of locality does not suit our purpose, essentially because the set of functions φ that can be written as $\varphi = \varphi_1 + \varphi_2$ (with $\text{supp } \varphi \cap \text{supp } \varphi_2 = \emptyset$) is not dense in the space of smooth functions. We show this now and we construct a partially additive functional which is not local.

5.2 A non-local partially additive functional

We work in the space $C^\infty(\mathbb{S}^1)$ of smooth functions on the unit circle. We denote by \mathcal{I} the subset of functions $f = \varphi_1 + \varphi_2$ which are sums of two elements of $C^\infty(\mathbb{S}^1)$ whose supports are disjoint. It is not a vector subspace of $C^\infty(\mathbb{S}^1)$.

The idea of the construction is the following. In the metric space $C^\infty(\mathbb{S}^1)$, we will show that the subset \mathcal{I} is bounded away from the constant function $f = 1$. This means that the functional equation (16) only concerns the restriction $F|_{\mathcal{I}}$ to a subset which is bounded away from 1. Therefore there is some open neighborhood of $f = 1$ which does not meet \mathcal{I} . Then we use Sobolev norms to build some cut-off function χ to glue a local functional near \mathcal{I} with a nonlocal functional near $f = 1$.

Lemma 5.1. *The constant function $f = 1$ is bounded away from \mathcal{I} in $C^\infty(\mathbb{S}^1)$: if $f \in \mathcal{I}$, then $\|f - 1\|_{C^0} = \sup_{x \in \mathbb{S}^1} |f(x) - 1| \geq 1$.*

Proof. Let us denote by $\|\cdot\|_{C^0}$ the norm $\|f\|_{C^0} = \sup_{x \in \mathbb{S}^1} |f(x)|$. It is a continuous norm for the Fréchet topology of $C^\infty(\mathbb{S}^1)$ because $\|f\|_{C^0} = \pi_{0, \mathbb{S}^1}(f)$. Then, if $\text{supp } \varphi_1 \cap \text{supp } \varphi_2 = \emptyset$ we have $\|\varphi_1 + \varphi_2 - 1\|_{C^0} \geq 1$. Indeed, the supports of φ_1 and φ_2 being compact, the fact that they do not meet implies that they are at a finite distance. Thus, there is a point $x \in \mathbb{S}^1$ such that $\varphi_1(x) = \varphi_2(x) = 0$. Hence, $|\varphi_1(x) + \varphi_2(x) - 1| = 1$ and $\sup_{x \in \mathbb{S}^1} |\varphi_1(x) + \varphi_2(x) - 1| \geq 1$. \square

The second step is to build a smooth function χ such that $\chi(1) = 1$ and $\chi|_{\mathcal{I}} = 0$.

Lemma 5.2. *There is a smooth function $\chi : C^\infty(\mathbb{S}^1) \rightarrow \mathbb{R}$ such that $\chi = 1$ on a neighborhood of $f = 1$ and $\chi(f) = 0$ if $\|f - 1\|_{C^0} \geq 1$. In particular, $\chi|_{\mathcal{I}} = 0$.*

Proof. First recall that the Sobolev norm H^{2k} on \mathbb{S}^1 is defined as

$$\|f\|_{H^{2k}} = \sqrt{\int_{\mathbb{S}^1} ((1 - \Delta)^k f(x))^2 dx} = 2\pi \left(\sum_{n \in \mathbb{Z}} (1 + n^2)^{2k} |\hat{f}(n)|^2 \right)^{\frac{1}{2}}, \quad (17)$$

where the last representation uses the Fourier series $f(x) = \sum_n \hat{f}(n) e^{inx}$. By the Sobolev injections, $H^2(\mathbb{S}^1)$ injects continuously in $C^0(\mathbb{S}^1)$. In other words, there is a constant $C > 0$ such that $\|f\|_{C^0} \leq C \|f\|_{H^2}$ for every $f \in C^\infty(\mathbb{S}^1)$.

Now we take a function $g \in C^\infty(\mathbb{R})$ such that $g(t) = 1$ when $t \leq 1/3C^2$ and $g(t) = 0$ when $t \geq 1/2C^2$ and we define $\chi : C^\infty(\mathbb{S}^1) \rightarrow \mathbb{R}$ by composing g with the square of the Sobolev norm.

$$\chi(f) = g(\|1 - f\|_{H^2}^2).$$

If $\|1 - f\|_{C^0} \geq 1$ (in particular, if $f \in \mathcal{I}$ by Lemma 5.1) the Sobolev injection leads to:

$$1 \leq \|1 - f\|_{C^0} \leq C \|1 - f\|_{H^2} \implies \|1 - f\|_{H^2}^2 \geq \frac{1}{C^2}$$

hence $g(\|1 - f\|_{H^2}^2) = 0$ by definition of g .

On the other hand $\|1 - f\|_{H^2} \leq \frac{1}{\sqrt{3}C}$ means that f belongs to the neighborhood of the constant function $f = 1$ defined by $V = \{f; \|1 - f\|_{H^2} \leq 1/\sqrt{3}C\}$. On this neighborhood, $g(\|1 - f\|_{H^2}^2) = 1$. The smoothness of χ is an immediate consequence of the chain rule, the smoothness of g and of the squared Sobolev norm $\|\cdot\|_{H^2(\mathbb{S}^1)}^2$. \square

We are now ready to define our counterexample:

Theorem 5.3. *The functional F_{nl} on $C^\infty(\mathbb{S}^1)$ defined for any integer $N > 1$ by*

$$F_{\text{nl}}(f) = (1 - \chi(f)) \int_{\mathbb{S}^1} f(x) dx + \chi(f) \left(\int_{\mathbb{S}^1} f(x) dx \right)^N, \quad (18)$$

is partially additive but not local.

Proof. For every $(\varphi_1, \varphi_2) \in C^\infty(\mathbb{S}^1)^2$ whose supports are disjoint, $f = \varphi_1 + \varphi_2 \in \mathcal{I}$ hence $\chi(f) = 0$ by Lemma 5.2. Moreover, we saw that, if $\text{supp } \varphi_1 \cap \text{supp } \varphi_2 = \emptyset$, then there is a point $x \in \mathbb{S}^1$ such that $\varphi_1(x) = \varphi_2(x) = 0$. Thus, $\|1 - \varphi_1\|_{C^0} \geq 1$ and $\|1 - \varphi_2\|_{C^0} \geq 1$. As a consequence, $\chi(\varphi_1) = \chi(\varphi_2) = 0$ by Lemma 5.2 and $F_{\text{nl}}(\varphi_1 + \varphi_2) = \int_{\mathbb{S}^1} (\varphi_1(x) + \varphi_2(x)) dx = F_{\text{nl}}(\varphi_1) + F_{\text{nl}}(\varphi_2)$.

On the other hand, in the neighborhood V of $f = 1$ given by Lemma 5.2, $\chi(f) = 1$ hence $F_{\text{nl}}(f) = \left(\int_{\mathbb{S}^1} f(x) dx \right)^N$ which is not local. It is even a typical example of a multilocal functional [14]. \square

Since partial additivity is equivalent to locality for polynomial functions, the non-locality of F_{nl} can be considered to be non perturbative. Moreover, the fact that the derivatives $D^n F_{\text{nl}}$ calculated at $f = 0$ are supported in the thin diagonal of $(\mathbb{S}^1)^n$, although F_{nl} is not local, means that locality cannot be controlled by the support of differentials taken at a single function f . We come now to the property that is relevant for quantum field theory.

5.3 Additive functionals

In 1965, Chacon and Friedman [84] introduced a more stringent concept of additivity which meets our needs:

Definition 5.4. *We say that a Bastiani smooth map $F : C^\infty(M) \rightarrow \mathbb{K}$ is additive if, for every triple $(\varphi_1, \varphi_2, \varphi_3)$ of smooth functions on M , the property $\text{supp } \varphi_1 \cap \text{supp } \varphi_3 = \emptyset$ implies the property*

$$F(\varphi_1 + \varphi_2 + \varphi_3) = F(\varphi_1 + \varphi_2) + F(\varphi_2 + \varphi_3) - F(\varphi_2). \quad (19)$$

In the literature, the additivity equation (19) is also called the *Hammerstein property* [85, 86, 87, 88, 89, 90]. The additivity property is equivalent to the fact that the functional derivatives are supported on the thin diagonal $D_n = \{(x_1, \dots, x_n) \in M^n; x_1 = \dots = x_n\}$ [15, 83].

Proposition 5.5. *A smooth functional F on $C^\infty(M)$ is additive iff $\text{supp } F_\varphi^{(2)} \subset D_2$ for every $\varphi \in C^\infty(M)$, where $D_2 = \{(x, y) \in M^2; x = y\}$. Moreover, if F is an additive functional, then $\text{supp } F_\varphi^{(n)} \subset D_n$ for every $\varphi \in C^\infty(M)$, where $D_n = \{(x_1, \dots, x_n) \in M^n; x_1 = \dots = x_n\}$.*

Proof. We first prove that the second derivative of an additive functional is localized on the diagonal [15]. If we use the additivity property with $\varphi_1 = \lambda\psi$, $\varphi_3 = \mu\chi$ and $\text{supp } \psi \cap \text{supp } \chi = \emptyset$, then

$$F(\lambda\psi + \varphi_2 + \mu\chi) = F(\lambda\psi + \varphi_2) + F(\varphi_2 + \mu\chi) - F(\varphi_2).$$

Since no term on the right hand side of this equation depends on both λ and μ , we have

$$\frac{\partial^2 F(\lambda\psi + \varphi_2 + \mu\chi)}{\partial \lambda \partial \mu} = D^2 F_{\lambda\psi + \varphi_2 + \mu\chi}(\psi, \chi) = F_{\lambda\psi + \varphi_2 + \mu\chi}^{(2)}(\psi \otimes \chi) = 0.$$

This equation, being true for every φ_2 , can be written $F_\varphi^{(2)}(\psi \otimes \chi) = 0$ for every φ and every pair (ψ, χ) with disjoint supports. Now for every point $(x, y) \in M^2$ such that $x \neq y$, there are two open sets U_x containing x and U_y containing y such that $U_x \cap U_y = \emptyset$. Then, any pair of functions ψ and χ supported in U_x and U_y satisfies $F_\varphi^{(2)}(\psi \otimes \chi) = 0$. Since the functions $\psi \otimes \chi$ are dense in $\mathcal{D}(M^2)$, this implies that every test functions $f \in \mathcal{D}(M^2)$ supported in $U_x \times U_y$ satisfies $F_\varphi^{(2)}(f) = 0$. Thus $(x, y) \notin \text{supp } F_\varphi^{(2)}$ and $\text{supp } F_\varphi^{(2)} \subset D_2$. To determine the support of $F_\varphi^{(n)}$, consider a point (x_1, \dots, x_n) which is not in D_n . Then, there are two indices i and j such that $x_i \neq x_j$. Denote by U_x an open neighborhood of x_i and by U_y an open neighborhood of x_j and repeat the previous proof to obtain $F_\varphi^{(2)}(\psi \otimes \chi) = 0$ for every φ and every pair (ψ, χ) with supports in U_x and U_y . Now, rewrite $\varphi = \varphi_0 + \sum \lambda_k \psi_k$, where $\psi_k(x_k) \neq 0$ and the sum is over all integers from 1 to n except i and j . Then, the derivatives with respect to λ_k are all zero and we find again with the same argument that (x_1, \dots, x_n) is not in the support of $F_\varphi^{(n)}$ for every φ .

Conversely [83, 17], assume that $\text{supp } F_\varphi^{(2)} \subset D_2$ for every φ . As we have seen in the first part of the proof, this means that, if ψ and χ have disjoint support, then $D^2 F_\varphi(\psi, \chi) = F_\varphi^{(2)}(\psi \otimes \chi) = 0$. By the fundamental theorem of calculus,

$$\begin{aligned} F(\varphi + \psi + \chi) &= F(\varphi + \psi) + \int_0^1 d\mu \frac{d}{d\mu} F(\varphi + \psi + \mu\chi), \\ F(\varphi + \psi + \mu\chi) &= F(\varphi + \mu\chi) + \int_0^1 d\lambda \frac{d}{d\lambda} F(\varphi + \lambda\psi + \mu\chi). \end{aligned}$$

Thus,

$$F(\varphi + \psi + \chi) = F(\varphi + \psi) + \int_0^1 d\mu \frac{d}{d\mu} F(\varphi + \mu\chi) + \int_0^1 d\lambda \int_0^1 d\mu \frac{\partial^2}{\partial \lambda \partial \mu} F(\varphi + \lambda\psi + \mu\chi).$$

The last term is zero because $D^2F_\varphi(\psi, \chi) = 0$ and the second term is $F(\varphi + \chi) - F(\varphi)$. We recover the additivity condition. \square

Finally, additivity is stronger than partial additivity because the latter corresponds to the case $\varphi_2 = 0$ and $F(0) = 0$. It is strictly stronger because F_{nl} is not additive.

6 Characterization of smooth local functionals

In this section, we give a characterization of local functionals inspired by the topology described in section 4.5. In the sequel, we shall deal with compactly supported distributions u with empty wavefront sets. We repeat the definition of local functionals in terms of jets:

Definition 6.1. *Let U be an open subset of $C^\infty(M)$. A Bastiani smooth functional $F : U \rightarrow \mathbb{K}$ is said to be local if, for every $\varphi \in U$, there is a neighborhood V of φ , an integer k , an open subset $\mathcal{V} \subset J^k M$ and a smooth function $f \in C^\infty(\mathcal{V})$ such that $x \in M \mapsto f(j_x^k \psi)$ is supported in a compact subset $K \subset M$ and*

$$F(\varphi + \psi) = F(\varphi) + \int_M f(j_x^k \psi) dx,$$

whenever $\varphi + \psi \in V$ and where $j_x^k \psi$ denotes the k -jet of ψ at x .

We invite the reader not familiarized with jet bundles to have a look at section 6.1, where these objects are carefully defined. Note that the representation of F by f is not unique: adding the total derivative of a function does not change the result. We shall see that f belongs to a unique cohomology class for some specific cohomology theory on the space of local functionals.

Before we state the main Theorem of this section, let us start by some useful definition-lemma :

Lemma 6.2. *Let U be an open subset of $C^\infty(M)$ and $F : U \rightarrow \mathbb{K}$ be Bastiani smooth. For every φ such that the distribution $DF_\varphi \in \mathcal{E}'(M)$ has empty wave front set, there exists a **unique function** $\nabla F_\varphi \in \mathcal{D}(M)$ such that*

$$DF_\varphi[h] = \int_M \nabla F_\varphi(x) h(x) dx. \quad (20)$$

Proof. Once a density dx is fixed on M , functions in $L^1_{loc}(M)$ (in particular in $C^\infty(M)$) can be identified with distributions by the map :

$$f \in L^1_{loc}(M) \mapsto \left(\phi \mapsto \int_M f \phi dx \right)$$

and [29, Theorem 1.2.4] shows that the distribution is uniquely defined when f is continuous hence when f is smooth.

Since $WF(DF_\varphi) = \emptyset$, there exists a unique C^∞ function ∇F_φ which represents the distribution $DF_\varphi \in \mathcal{E}'(M)$ by integration on M against dx . \square

The main theorem of this section is

Theorem 6.3. *Let U be an open subset of $C^\infty(M)$ and $F : U \rightarrow \mathbb{K}$ be Bastiani smooth. Then, F is local if and only if the following two conditions are satisfied:*

1. F is additive,
2. for every $\varphi \in U$, the differential $DF_\varphi = F_\varphi^{(1)}$ of F at φ has an empty wave front set and the map $\varphi \mapsto \nabla F_\varphi$ is Bastiani smooth from U to $\mathcal{D}(M)$.

Note that our definition of locality is strictly more general than the usual one because the counterexample described in Lemma 3.12 is local in our sense but not in the sense of Eq. (1) since its order is infinite.

The proof is delayed to Section 6.3. Since this theorem deals with jets, we start with a short presentation of the jet bundle. Our point of view on jets is based on the concept of infinitesimal neighborhoods due to Grothendieck and is closely related to several expositions in the literature [91, 92, 93].

6.1 The manifold of jets of functions on a manifold

Let M be a manifold. For every smooth real-valued function φ on M , we call k -jet of φ at a point $x \in M$ the class $j_x^k(\varphi)$ of φ in the quotient $C^\infty(M)/I_x^{k+1}$, with the understanding that I_x^{k+1} stands for the $(k+1)$ -th power of the ideal I_x of smooth functions on M vanishing at $x \in M$. Recall that I_x^{k+1} coincides with the ideal of smooth functions on M whose $k+1$ first derivatives vanish at the point x .

For all $x \in M$, the space $J_x^k(M)$ of all k -jets of functions on M at x coincides with $C^\infty(M)/I_x^{k+1}$ and is called the *space of k -jets at x* . It is clearly a vector space. The disjoint union $J^k(M) := \coprod_{x \in M} J_x^k(M)$ is a smooth vector bundle over M called the bundle of k -th jets. Consider the map:

$$\begin{aligned} \mathfrak{J}_\Delta : C^\infty(M \times M) &\rightarrow \Gamma(J^k(M)) \\ \psi &\mapsto x \mapsto j_x^k(\mathfrak{i}_x^* \psi), \end{aligned}$$

where $\mathfrak{i}_x : M \rightarrow M \times M$ is the map $y \mapsto (x, y)$. It is known that \mathfrak{J}_Δ is surjective onto the space of smooth sections of $J^k(M)$ and its kernel is the $(k+1)$ -th power of the ideal \mathcal{I}_Δ of functions on $M \times M$ vanishing on the diagonal.

Last, the projection $p_1 : M \times M \rightarrow M$ onto the first component dualizes in an algebra morphism $\varphi \mapsto p_1^* \varphi$ from $C^\infty(M)$ to $C^\infty(M \times M)$ which endows $C^\infty(M \times M)$ with a $C^\infty(M)$ -module structure. The space of sections of $J^k(M)$ is also a $C^\infty(M)$ -module, and it is routine to check that \mathfrak{J}_Δ is a morphism of $C^\infty(M)$ -modules. Therefore, the space of sections of $J^k(M)$ is, as a $C^\infty(M)$ -module, isomorphic to the quotient $C^\infty(M \times M)/\mathcal{I}_\Delta^{k+1}$ and $\Gamma(J^k(M))$ fits into the following exact sequence of $C^\infty(M)$ -modules:

$$0 \rightarrow \mathcal{I}_\Delta^{k+1} \rightarrow C^\infty(M \times M) \rightarrow \Gamma(J^k(M)) \simeq C^\infty(M \times M)/\mathcal{I}_\Delta^{k+1} \rightarrow 0.$$

And the map that to f associates its k -jet reads:

$$f \in C^\infty(M) \mapsto [(p_1^* f)] \in C^\infty(M \times M)/\mathcal{I}_\Delta^{k+1}.$$

The purpose of the rest of this section is to prove the technical Proposition 6.4, the statement of which we now explain. For all integer $k \in \mathbb{N}$, there is a natural vector bundle morphism \mathfrak{j}^k from the trivial bundle over M with typical fiber $E = C^\infty(M)$ to the bundle $J^k(M) \rightarrow M$ of k -jets. This morphism simply consists in assigning to a pair (f, x) in $E \times M \rightarrow M$ the k -jet of f at x . In equation:

$$\begin{aligned} \mathfrak{j}^k : E \times M &\mapsto j^k(M) \\ (f, x) &\rightarrow j_x^k(f). \end{aligned} \tag{21}$$

The result goes as follows.

Proposition 6.4. *Let $E = C^\infty(M)$ and $V \subset E$ be an open subset and $k \in \mathbb{N}$ an integer.*

1. *The subset $\mathfrak{j}^k(V \times M)$ is an open subset of $j^k(M)$.*
2. *Let c be a smooth \mathbb{K} -valued function on $V \times M$, with $V \subset E$ an open subset. Assume that $c(f, x)$ depends only on the k -jet of the function f at the point x . Then there exists a unique smooth \mathbb{K} -valued function \tilde{c} on the open subset $\mathfrak{j}^k(V \times M) \subset j^k(M)$ that makes the following diagram commutative:*

$$\begin{array}{ccc} V \times M & \xrightarrow{c} & \mathbb{K}. \\ \downarrow \mathfrak{j}^k & \nearrow \tilde{c} & \\ \mathfrak{j}^k(V \times M) & & \end{array} \tag{22}$$

i.e., such that the relation $c(f, x) = \tilde{c}(j_x^k(f))$ holds for all $f \in V$ and $x \in M$.

When $V = E$, Proposition 6.4 specializes to the following easier statement:

Corollary 6.5. *Let c be a smooth function from $E \times M$ to \mathbb{K} . Assume that there exists an integer k such that $c(f, x)$ depends only on the k -jet of the function f at the point x . Then there exists a unique smooth \mathbb{K} -valued function \tilde{c} on $j^k(M)$ such that the following diagram commutes:*

$$\begin{array}{ccc} E \times M & \xrightarrow{c} & \mathbb{K}. \\ \downarrow j^k & \nearrow \tilde{c} & \\ j^k(M) & & \end{array} \quad (23)$$

i.e., such that the relation $c(f, x) = \tilde{c}(j_x^k(f))$ holds for all $f \in V$ and $x \in M$.

Before establishing these results, we shall need several lemmas.

Lemma 6.6. *The vector bundle morphism j^k described in (21) is surjective and admits a smooth section \mathfrak{s}^k .*

Proof. The section \mathfrak{s}^k , when it exists, being by construction a right inverse of j^k , the latter is surjective. It suffices therefore to prove the existence of \mathfrak{s}^k .

We first prove that the lemma holds true for M an open subset V of \mathbb{R}^n . In that case, the bundle of k -jets $j_x^k(V)$ is isomorphic to the trivial bundle over V with typical fiber the space of polynomials of degree less or equal to k . There is an obvious candidate for the section of j^k : it consists in mapping $\alpha_x \in j_x^k(V)$ to the unique polynomial of degree k whose k -jet at $x \in V$ is α . The henceforth obtained assignment, that we denote by \mathfrak{s}_V^k , is a smooth vector bundle morphism from $j^k(V) \rightarrow V$ to the trivial bundle $C^\infty(V) \times V \rightarrow V$. It is by construction a section of j^k .

We now go back to the general case of an arbitrary manifold M . For every point $x \in M$, choose V_x a coordinate neighbourhood and let χ be a smooth function with compact support on V_x which is identically equal to 1 in a neighbourhood $V'_x \subset V_x$ of x . Since V_x is a coordinate neighbourhood, it can be identified with an open subset of \mathbb{R}^n , which allows to consider

$$\mathfrak{s}_{V_x}^k : j^k(V_x) \mapsto C^\infty(V_x) \times V_x$$

as in the previous paragraph. We can then consider the composition of vector bundle morphisms over V_x :

$$j^k(M)|_{V_x} \simeq j^k(V_x) \xrightarrow{\mathfrak{s}_{V_x}^k} C^\infty(V_x) \times V_x \xrightarrow{m_\chi \times id} E \times V_x$$

where $j^k(M)|_{V_x} \simeq j^k(V_x)$ is the obvious identification of the k -jet bundle of V_x to the restriction to V_x of the k -jet bundle on M , and where m_χ is the smooth linear map from $C^\infty(V_x)$ to $E = C^\infty(M)$ defined by $m_\chi(f) = f\chi$. Since $\chi = 1$ identically equal to 1 on V'_x , the restriction to V'_x of this vector bundle morphism is by construction a section of the restriction of j^k to V'_x .

Since the manifold M is paracompact³, the latter point implies that the manifold M can be covered by open subsets $(U_i)_{i \in I}$ such that the restriction of j^k to U_i admits a section \mathfrak{s}_i^k . Without any loss of generality, we can assume the existence of a smooth partition of unity $(\chi_i)_{i \in I}$ relative to this open cover. A global smooth section of j^k is then given by the explicit formula $\mathfrak{s}^k = \sum_{i \in I} \chi_i \mathfrak{s}_i^k$, as follows from the obvious computation:

$$j^k \circ \mathfrak{s}^k = \sum_{i \in I} \chi_i j^k \circ \mathfrak{s}_i^k = \sum_{i \in I} \chi_i \text{id}_{j^k(M)} = \text{id}_{j^k(M)}$$

where we used the fact that j^k commutes with multiplications by χ_i since $j^k : E \times M \mapsto J^k M$ is a vector bundle morphism. This completes the proof. \square

³ Since all manifolds in this article are assumed to be paracompact, for every property T on the set of all open subsets on M , provided that (i) open subsets of subsets satisfying T satisfy T , and (ii) every point of M admits a neighbourhood that satisfies property T , then there exists an open cover $(U_i)_{i \in I}$ made of subsets satisfying the property T that admits a partition of unity $(\chi_i)_{i \in I}$ relative to it.

Since $c : V \times M \rightarrow \mathbb{K}$ is only defined on the open subset $V \times M$ of $E \times M$, we need the following refinement of Lemma 6.6 where the local sections \mathfrak{t}_x^k of j^k are valued in $V \times M$:

Lemma 6.7. *For every $(f, x) \in E \times M$, the vector bundle morphism j^k described in (21) admits a smooth section \mathfrak{t}^k through⁴ (f, x) .*

Proof. Notice that Lemma 6.7 can be derived from Lemma 6.6 for any vector bundle morphism over the identity of M . A careful check shows that the arguments below are absolutely general and indeed show that for any two vector bundle E_1, E_2 over M , any vector bundle morphism $E_1 \mapsto E_2$ over M that admits a section is a submersion, and admits a section through every point of E_1 .

We prefer to do it, however, in our particular setting – since one of the bundles is infinite dimensional and requires careful attention.

Let \mathfrak{s}^k be a section of j^k as in Lemma 6.6. Consider the smooth map defined at all point $y \in M$ by

$$\begin{aligned} \mathfrak{t}_y^k : j_y^k(M) &\mapsto (E \times M)_y \simeq E \\ \beta &\rightarrow \mathfrak{s}_y^k(\beta) + (f - \mathfrak{s}_y^k \circ j_y^k(f)) \end{aligned}$$

This map is smooth by construction. It is again a section j^k , as follows from the following computation, valid for all $y \in M, \beta \in j_y^k(M)$:

$$\begin{aligned} j_y^k \circ \mathfrak{t}_y^k(\beta) &= j_y^k(\mathfrak{s}_y^k(\beta) + (f - \mathfrak{s}_y^k \circ j_y^k(f))) \\ &= j_y^k \circ \mathfrak{s}_y^k(\beta) + j_y^k(f) - j_y^k \circ \mathfrak{s}_y^k \circ j_y^k(f) \\ &= \beta + j_y^k(f) - j_y^k(f) = \beta. \end{aligned}$$

Then the section \mathfrak{t}^k above satisfies by construction:

$$\mathfrak{t}_x^k \circ j_x^k(f) = \mathfrak{s}_x^k \circ j_x^k(f) + f - \mathfrak{s}_x^k \circ j_x^k(f) = f.$$

This completes the proof. \square

Lemma 6.7 has the following immediate consequence.

Lemma 6.8. *The vector bundle morphism j^k described in (21) is a submersion.*

Proof. For every (f, x) in $E \times M$, let us choose \mathfrak{t}^k to be a section through (f, x) as in Lemma 6.7. By construction, the differential of j^k at (f, x) admits the differential of \mathfrak{t}^k at $j^k(f, x)$ as right inverse, so it is surjective. \square

We can now prove Proposition 6.4.

Proof. Since the vector bundle morphism j^k described in (21) is a submersion by Lemma 6.8 and since $V \times M$ is open in $E \times M$, the subset $j^k(V \times M)$ is an open subset of $j^k(M)$. This proves the first item in Proposition 6.4.

Let us now prove the second item. Assume that we are given a function $c : V \times M \mapsto \mathbb{K}$ such that the value $c(f, x)$ at an arbitrary $f \in E$ and $x \in M$ depends only the k -jet of f at x . The existence of an unique function \tilde{c} from $j^k(M)$ to \mathbb{K} making the diagram (22) commute is simply a set-theoretic property: the difficulty is to show that this function \tilde{c} is smooth.

When $V = E$ (i.e. under the assumptions of Corollary 6.5), the smoothness of \tilde{c} follows directly from Lemma 6.6, which implies that the commutative diagram (23) can be completed to

$$\begin{array}{ccc} E \times M & \xrightarrow{c} & \mathbb{K}. \\ \uparrow \mathfrak{s}^k & \nearrow j^k \tilde{c} & \\ j^k(M) & & \end{array}$$

⁴i.e. such that $\mathfrak{t}_x^k \circ j_x^k(f) = f$.

which amounts to say that the following relation holds

$$\tilde{c} = c \circ \mathfrak{s}^k. \quad (24)$$

The latter formula and the smoothness of \mathfrak{s}^k implies that, when c is assumed to be a smooth function, so is the function \tilde{c} by composition. This proves Corollary 6.5.

For the general case, we have to choose, for all $\alpha \in \mathfrak{j}^k(V \times M)$ a section \mathfrak{t}^k of \mathfrak{j}^k such that $\mathfrak{t}^k(\alpha) \in V \times M$. Such a section \mathfrak{t}^k always exists by Lemma 6.7. Since \mathfrak{t}^k is smooth, there exists a neighbourhood W_α of α in $\mathfrak{j}^k(V \times M)$ on which \mathfrak{t}^k takes values in the domain of definition $V \times M$ of c , which implies that the commutative diagram (22) can be completed to

$$\begin{array}{ccc} (V \times M) \cap (\mathfrak{j}^k)^{-1}(W_\alpha^c) & \longrightarrow & \mathbb{K} \\ \mathfrak{t}^k \left(\begin{array}{c} \uparrow \\ \downarrow \\ \mathfrak{j}^k \end{array} \right) & \nearrow \tilde{c} & \\ W_\alpha & & \end{array}$$

In turn, the commutativity of this diagram gives the explicit description of \tilde{c} through the following formula, valid on W_α :

$$\tilde{c} = c \circ \mathfrak{t}^k. \quad (25)$$

Formula (25) and the smoothness of \mathfrak{t}^k imply that, when c is assumed to be a smooth function, so is, by composition, the restriction to W_α of the function \tilde{c} . Since every $\alpha \in \mathfrak{j}^k(V \times M)$ admits a neighbourhood on which the restriction of \tilde{c} is smooth, the function \tilde{c} is a smooth function. This completes the proof. \square

6.2 Properties of $F_\varphi^{(2)}$

We first show that the two assumptions of our theorem are equivalent to some strong assumptions on the second derivative of F :

Lemma 6.9. *Let U be an open subset of $C^\infty(M)$ and $F : U \rightarrow \mathbb{K}$ be Bastiani smooth. Assume that for every $\varphi \in U$, the differential $DF_\varphi = F_\varphi^{(1)}$ of F at φ has no wave front set, i.e. $\text{WF}(F_\varphi^{(1)}) = \emptyset$. Then the two following properties are equivalent:*

1. F is additive and the map $\varphi \mapsto \nabla F_\varphi$ is Bastiani smooth from U to $\mathcal{D}(M)$.
2. For every $\varphi_0 \in U$, there is a neighborhood V of φ_0 , a compact $K \subset M$ and a finite family of Bastiani smooth maps $f_\alpha : V \rightarrow \mathcal{D}(K)$ with $|\alpha| \leq k$, such that in any system of local coordinates (x, y) on M^2 :

$$F_\varphi^{(2)}(x, y) = \sum_{|\alpha| \leq k} f_\alpha(\varphi)(x) \partial_y^\alpha \delta(x - y), \quad (26)$$

for every $\varphi \in V$.

In particular, both conditions imply that $D^2 F_\varphi$ is represented by a distribution $F_\varphi^{(2)}$ whose wave front set is the conormal bundle of the diagonal in M^2 [94, p. 32].

In the sequel, we shall often use the following simple lemma :

Lemma 6.10. *Let E, F and G be locally convex spaces. If $f : E \rightarrow F$ is Bastiani smooth and $\ell : F \rightarrow G$ is linear and continuous, then $\ell \circ f : E \rightarrow G$ is Bastiani smooth and $D^k(\ell \circ f) = \ell \circ D^k f$.*

Proof. This is a consequence of three facts: the map ℓ is Bastiani smooth because it is linear and continuous, $\ell \circ f$ is Bastiani smooth because it is the composition of two Bastiani smooth maps and the chain rule. \square

We also need the following Lemma in the proof of Lemma 6.9:

Lemma 6.11. *Let U be a convex open subset of $E = C^\infty(M)$ containing the origin and $F : U \rightarrow E$ a Bastiani smooth map. Then, $G : U \rightarrow E$ defined by $G(\varphi) = \int_0^1 F(s\varphi)ds$ is Bastiani smooth.*

Proof. The first step is to define a candidate for the Bastiani differential $D^k G$ by determining $D^k G(x)$ pointwise in $x \in M$. For every $(t_1, \dots, t_k, x) \in [0, 1]^k \times M$ and $(\varphi, \psi_1, \dots, \psi_k) \in U \times E^k$, the function $(t_1, \dots, t_k, x) \mapsto \int_0^1 ds F(s(\varphi + t_1\psi_1 + \dots + t_k\psi_k))(x)$ is smooth in (t_1, \dots, t_k, x) by dominated convergence theorem since x can always be restricted to some compact subset $K \subset M$ to obtain uniform bounds. We can differentiate in (t_1, \dots, t_k) outside and inside the integral and both differentials coincide. Therefore, for every $x \in M$, the Bastiani k th-differential $D^k G(x)$ of $G(x) = \int_0^1 ds F(s\varphi(x))$ exists and satisfies the relation $D^k G_\varphi(\psi_1, \dots, \psi_k)(x) = \int_0^1 ds s^k D^k F_{s\varphi}(\psi_1, \dots, \psi_k)(x)$. Let us show that $D^k G : U \times E^k \mapsto E$ is jointly continuous in $(\varphi, \psi_1, \dots, \psi_k)$.

We know that the map $\chi : (s, \varphi, \psi_1, \dots, \psi_k) \in [0, 1] \times U \times E^k \mapsto s^k D^k F_{s\varphi}(\psi_1, \dots, \psi_k) \in E$ is continuous by joint continuity of $D^k F : U \times E^k \mapsto E$ and composition of the continuous maps

$$(s, \varphi, \psi_1, \dots, \psi_k) \mapsto (s\varphi, \psi_1, \dots, \psi_k) \mapsto s^k D^k F_{s\varphi}(\psi_1, \dots, \psi_k).$$

Then by [36, Thm 2.1.5 p. 72] applied to the function χ , the integrated map $(\varphi, \psi_1, \dots, \psi_k) \mapsto \int_0^1 ds s^k D^k F_{s\varphi}(\psi_1, \dots, \psi_k)$ is continuous and the proof is complete because continuity holds true for every k . \square

Let us now prove Lemma 6.9.

Proof. First of all, by proposition 5.5, F is additive iff its second derivative is represented by a distribution supported in the diagonal. We start by proving the direct sense assuming that $\varphi \in U \mapsto \nabla F_\varphi \in C^\infty(M)$ is Bastiani smooth.

We first show that item 1 implies item 2 in Lemma 6.9. Since F is Bastiani smooth that for any $\varphi_0 \in U$, we already know by Proposition 3.4 that there is some neighborhood V of φ_0 on which $F|_V$ has fixed compact support that we denote by K . Therefore, ∇F_φ belongs to $\mathcal{D}(K)$ for every $\varphi \in V$ and $F_\varphi^{(2)}$ is supported in $K \times K$. Since $F_\varphi^{(2)}$ is also supported in the diagonal of M^2 by proposition 5.5, the support of $F_\varphi^{(2)}$ is contained in the diagonal of $K \times K$ which can be identified with K itself.

Since DF_φ has an empty wavefront set by assumption, its singular support is empty and it can be represented by a *unique* smooth compactly supported function ∇F_φ [29, p. 37] such that

$$\frac{d}{dt} F(\varphi + th)|_{t=0} = DF_\varphi(h) = \int_M \nabla F_\varphi(x) h(x) dx. \quad (27)$$

The main step is to represent $F_\varphi^{(2)}$ as the Bastiani differential of ∇F_φ by calculating the second derivatives in two different ways. The Bastiani smoothness of F yields:

$$\begin{aligned} D^2 F_\varphi(g, h) &= \frac{d^2}{dt_1 dt_2} F(\varphi + t_1 h + t_2 g)|_{t_1=t_2=0} \\ &= \frac{d}{dt_2} \left(\frac{d}{dt_1} F(\varphi + t_1 h + t_2 g)|_{t_1=0} \right) |_{t_2=0} \\ &= \frac{d}{dt_2} \left(\int_M \nabla F_{\varphi+t_2 g}(x) h(x) dx \right) |_{t_2=0}, \end{aligned}$$

where we used the Schwarz lemma and Equation (27). To justify switching $\frac{d}{dt_2}$ and integration over M , observe that the map $\varphi \in U \mapsto \nabla F_\varphi \in \mathcal{D}(M)$ is Bastiani smooth hence C^1 . It follows by the chain rule that $t \mapsto \frac{d}{dt} \nabla F_{\varphi+tg}$ is a C^0 map valued in $\mathcal{D}(M)$. Since ∇F_φ is actually in $\mathcal{D}(K)$ for every $\varphi \in V$ and the topology induced by $\mathcal{D}(M)$ on $\mathcal{D}(K)$ is the usual Fréchet topology of $\mathcal{D}(K)$, the map ∇F is smooth from V to the Fréchet space $\mathcal{D}(K)$.

Since $\mathcal{D}(K)$ injects continuously in $(C^0(K), \pi_{0,K})$, this implies that $(t, x) \in [-1, 1] \times K \mapsto \frac{d}{dt} \nabla F_{\varphi+tg}(x) \in C^0([-1, 1] \times K)$. Hence the integrand $\frac{d}{dt_2} \nabla F_{\varphi+t_2g}(x)h(x)$ is in $C^0([-1, 1] \times K)$ and is bounded on the integration domain⁵.

By the dominated convergence theorem, we can differentiate under the integral sign:

$$\begin{aligned} D^2 F_{\varphi}(g, h) &= \frac{d}{dt_2} \left(\int_M \nabla F_{\varphi+t_2g}(x)h(x)dx \right) \Big|_{t_2=0} \\ &= \int_M \left(\frac{d}{dt_2} \nabla F_{\varphi+t_2g}(x) \Big|_{t_2=0} \right) h(x)dx. \end{aligned}$$

By definition $\frac{d}{dt_2} \nabla F_{\varphi+t_2g} \Big|_{t_2=0}$ is only the Bastiani derivative

$$D\nabla F : (\varphi, g) \in V \times C^\infty(M) \mapsto D\nabla F_{\varphi}[g] \in \mathcal{D}(K), \quad (28)$$

where $D\nabla F$ is a Bastiani smooth map since ∇F is Bastiani smooth.

Note also that by theorem 3.9, the second derivative $D^2 F_{\varphi}(g, h)$ can be represented by a map $\varphi \in U \mapsto F_{\varphi}^{(2)} \in \mathcal{E}'(M^2)$ such that $\forall (\varphi, g, h) \in U \times C^\infty(M)^2$, $D^2 F_{\varphi}(g, h) = \langle F_{\varphi}^{(2)}, g \otimes h \rangle$ which means that $F_{\varphi}^{(2)}$ is the distributional kernel of the second derivative $D^2 F_{\varphi}$. We now arrive at the following equality which identifies two different representations of the second derivative

$$\forall (\varphi, g, h) \in U \times C^\infty(M)^2, \quad \langle F_{\varphi}^{(2)}, g \otimes h \rangle = \int_M D\nabla F_{\varphi}[g](y)h(y)dx. \quad (29)$$

By the same theorem 3.9 and the chain rule, $D\nabla F_{\varphi}[g](y) = ev_y D\nabla F_{\varphi}[g]$ is linear continuous in $g \in C^\infty(M)$, hence there is a distribution, denoted by $D\nabla F_{\varphi}(x, y)$, such that $\int_M D\nabla F_{\varphi}(x, y)g(x)dx = D\nabla F_{\varphi}[g](y)$ and $\int_M D\nabla F_{\varphi}(x, y)g(x)dx$ is in $\mathcal{D}(K)$ by Eq. (28). Since the above identity holds for all $(g, h) \in C^\infty(M)^2$, we have in the sense of distributions that $F_{\varphi}^{(2)}(x, y) = D\nabla F_{\varphi}(x, y)$ where the map

$$(\varphi, g) \in V \times C^\infty(M) \mapsto \int_M F_{\varphi}^{(2)}(x, \cdot)g(x)dx \in \mathcal{D}(K), \quad (30)$$

is Bastiani smooth.

It suffices to do the last part of the proof, which is local in nature, on $M = \mathbb{R}^d$. We now represent $F_{\varphi}^{(2)}(x, y)$ as a $C^\infty(M)$ -linear combination of derivatives of Dirac distributions concentrated on the diagonal. By Proposition 5.5, the additive property satisfied by F implies that the distribution $F^{(2)}$ associated to the second derivative $D^2 F$ is supported in the diagonal $D_2 \subset M \times M$. By Proposition 3.11, the kernel $F_{\varphi}^{(2)}(x, y) \in \mathcal{E}'(M \times M)$ has bounded distributional order uniformly in $\varphi \in V$. It follows by a theorem of L. Schwartz [24, p. 101] that in local coordinates, there exists a finite sequence of distributions $(\varphi \in V \mapsto f_{\alpha}(\varphi, \cdot) \in \mathcal{D}(K))_{|\alpha| \leq k}$ such that $F_{\varphi}^{(2)} = \sum_{|\alpha| \leq k} f_{\alpha}(\varphi, x) \partial_y^{\alpha} \delta(x - y)$. We denote the distributions $f_{\alpha}(\varphi)$ by $f_{\alpha}(\varphi, x)$ because we shall show that $\varphi \mapsto f_{\alpha}(\varphi)$ is Bastiani smooth from V to $\mathcal{D}(K)$.

By Equation (30), we know that

$$(\varphi, g) \in V \times C^\infty(M) \mapsto \int_M F_{\varphi}^{(2)}(x, \cdot)g(x)dx = \sum_{|\alpha| \leq k} (-1)^{|\alpha|} f_{\alpha}(\varphi, \cdot) \partial^{\alpha} g(\cdot) \in \mathcal{D}(K)$$

is smooth. Choosing g to be equal to the Fourier oscillatory function $e^{-i\langle \xi, x \rangle}$, we obtain by the

⁵A continuous map $u : t \in [-1, 1] \mapsto u(t, \cdot) \in (C^0(K), \pi_{0,K})$ corresponds to a map also denoted by $u \in C^0([-1, 1] \times K)$. Indeed for every convergent sequence $(t_n, x_n) \xrightarrow{n \rightarrow \infty} (t, x)$ in $[-1, 1] \times K$, the simple estimate $|u(t, x) - u(t_n, x_n)| \leq |u(t, x) - u(t, x_n)| + |u(t, x_n) - u(t_n, x_n)| \leq |u(t, x) - u(t, x_n)| + \pi_{0,K}(u(t_n, \cdot) - u(t, \cdot))$ shows that $u(t_n, x_n) \xrightarrow{n \rightarrow \infty} u(t, x)$.

chain rule the following map:

$$\begin{aligned} (\varphi, \xi) \mapsto \int_M F_\varphi^{(2)}(x, y) e^{-i\langle \xi, x \rangle} dy &= \int_M \sum_{|\alpha| \leq k} (-1)^{|\alpha|} f_\alpha(\varphi, y) \delta(x - y) \partial_y^\alpha e^{-i\langle \xi, x \rangle} dy \\ &= \sum_{|\alpha| \leq k} (-1)^{|\alpha|} f_\alpha(\varphi, x) (-i\xi)^\alpha, \end{aligned}$$

is Bastiani smooth. Moreover, since the image of the map in Eq. (30) is in $\mathcal{D}(K)$ for every smooth g , we obtain that $\sum_{|\alpha| \leq k} (-1)^{|\alpha|} f_\alpha(\varphi, \cdot) (-i\xi)^\alpha$ is in $\mathcal{D}(K)$ for every ξ . This is only possible if $f_\alpha(\varphi) \in \mathcal{D}(K)$ for every $|\alpha| \leq k$. Therefore $\varphi \mapsto f_\alpha(\varphi) = (i \frac{d}{d\xi})^\alpha \int_M F_\varphi^{(2)}(\cdot, y) e^{-i\langle \xi, y \rangle} dy|_{\xi=0}$ is Bastiani smooth from V to $\mathcal{D}(K)$ and the proof of the direct sense is complete.

Conversely, we want to prove that if there is a neighborhood V of φ_0 , a compact $K \subset M$ and a finite family of smooth maps $\varphi \mapsto f_\alpha(\varphi) \in \mathcal{D}(K)$, $|\alpha| \leq k$ such that in any system of local coordinates (x, y) on M^2 :

$$F_\varphi^{(2)}(x, y) = \sum_{|\alpha| \leq k} f_\alpha(\varphi)(x) \partial_y^\alpha \delta(x - y),$$

then $\varphi \mapsto \nabla F_\varphi \in \mathcal{D}(M)$ is Bastiani smooth. Without loss of generality, we assume that V is convex. By the Taylor formula with remainder for Bastiani smooth functions, for every $(\varphi, \psi_1, \psi_2) \in V \times C^\infty(M)^2$:

$$D^2 F_{\varphi+s_1\psi_1+s_2\psi_2}(\psi_1, \psi_2) = \partial_{s_1} \partial_{s_2} F(\varphi + s_1\psi_1 + s_2\psi_2) = \partial_{s_2} D F_{\varphi+s_1\psi_1+s_2\psi_2}(\psi_1),$$

for s_1 and s_2 small enough. It follows by the fundamental theorem of calculus and by evaluating at $s_1 = 0$ the previous relation that

$$D F_{\varphi+t\psi_2}(\psi_1) = D F_\varphi(\psi_1) + \int_0^t \partial_{s_2} D F_{\varphi+s_2\psi_2}(\psi_1) ds_2 = D F_\varphi(\psi_1) + \int_0^t D^2 F_{\varphi+s\psi_2}(\psi_1, \psi_2) ds,$$

where by assumption $D F_\varphi(\psi_1)$ is represented by integration against a smooth function

$$\begin{aligned} D F_\varphi(\psi_1) &= \int_M \nabla F_\varphi(x) \psi_1(x) dx, \\ D^2 F_{\varphi+s\psi_2}(\psi_1, \psi_2) &= \int_{M \times M} F_{\varphi+s\psi_2}^{(2)}(x, y) \psi_1(x) \psi_2(y) dx dy, \end{aligned}$$

and $F_{\varphi+s\psi}^{(2)}$ is supported on a subset of the diagonal $D_2 \subset M \times M$ that can be identified with K . Hence, for $\psi \in C^\infty(M)$ such that $\varphi + \psi \in V$:

$$\begin{aligned} \nabla F_{\varphi+\psi}(x) &= \nabla F_\varphi(x) + \int_0^1 \left(\int_M F_{\varphi+s\psi}^{(2)}(x, y) \psi(y) dy \right) ds \\ &= \nabla F_\varphi(x) + \sum_{|\alpha| \leq k} (-1)^{|\alpha|} \partial^\alpha \psi(x) \int_0^1 f_\alpha(\varphi + s\psi)(x) ds. \end{aligned}$$

To show that the map $\chi : V \rightarrow \mathcal{D}(K)$ defined by $\chi(\psi) = \nabla F_{\varphi+\psi}$ is smooth, we notice that, according to the last equation, $\nabla F_{\varphi+\psi}$ is the sum of the constant ∇F_φ and a finite linear combination of products of $\psi \mapsto \partial^\alpha \psi$ by an integral over s . The integrand $f_\alpha(\varphi + s\psi)$ is smooth by assumption. Therefore, the map $\psi \in (V - \varphi) \mapsto \int_0^1 f_\alpha(\varphi + s\psi)(x) ds \in \mathcal{D}(K)$ is smooth by Lemma 6.11 and the fact that the topology induced on $\mathcal{D}(K)$ by the topology of $C^\infty(M)$ is the standard topology of $\mathcal{D}(K)$. The map $\psi \mapsto \partial^\alpha \psi$ is smooth because it is linear and continuous. Finally, the product of the integral by $\partial^\alpha \psi$ is smooth by a trivial extension of Lemma 6.15. This completes the proof of Lemma 6.9. \square

We are now ready to prove Theorem 6.3 characterizing local functionals.

6.3 Proof of theorem 6.3

Let us start by proving the converse sense where we assume that F is the integral of some local function on jet space. Let $\varphi \in U$ and V some neighborhood of φ such that $F(\varphi + \psi) = \int_M f(x, j_x^k \psi) dx$ for every $\psi \in V$ where $j_x^k \psi$ is the k -jet of ψ at x and where f is smooth and compactly supported in the variable x in some fixed compact $K \subset M$. Without loss of generality, we can restrict the support K of f by a smooth partition of unity and assuming that K is contained in some open chart of M , we may reduce to the same problem for $f \in C^\infty(\Omega)$ where Ω is some open set in \mathbb{R}^d and $K \subset \Omega$.

We choose a smooth compactly supported function $\chi \in \mathcal{D}(\Omega)$ such that $\chi = 1$ on a compact neighborhood of K with $\text{supp } \chi \subset \Omega$ and we observe that

$$\Psi : \psi \in C^\infty(\Omega) \mapsto (\partial^\alpha \psi)_{|\alpha| \leq k} \chi \in \mathcal{D}(\text{supp}(\chi))^{\frac{(d+k)!}{d!}},$$

is linear continuous hence Bastiani smooth. We need a simple

Lemma 6.12. *Let Ω be an open set in \mathbb{R}^d then the map*

$$\Phi : \varphi \in C^\infty(\Omega, \mathbb{R}^r) \mapsto \{x \mapsto (x, \varphi(x))\} \in C^\infty(\Omega, \mathbb{R}^d \times \mathbb{R}^r),$$

is Bastiani smooth.

Proof. The first Bastiani differential $D\Phi_\varphi(h)$ can be identified with the smooth function $x \mapsto (0, h(x))$, which is linear continuous in h and does not depend on φ . It is thus smooth and so is Φ . \square

Therefore, the composition $\Phi \circ \Psi : C^\infty(\Omega) \rightarrow C^\infty(\Omega, \mathbb{R}^d \times \mathbb{R}^{\frac{(d+k)!}{d!}})$ defined by $\Phi \circ \Psi(\psi) : x \mapsto (x, \chi \partial^\alpha \psi(x)_{|\alpha| \leq k})$, is Bastiani smooth and finally

$$\psi \in C^\infty(M) \mapsto f(\cdot, j_x^k \psi(\cdot)) \in \mathcal{D}(K) \mapsto \int_\Omega f(x, j_x^k \psi) dx,$$

is Bastiani smooth by the chain rule and since the last integration map is linear continuous thus Bastiani smooth.

Now let us prove the direct sense, where we start from a functional characterization of F and end up with a representation as a function $F(\varphi + \psi) = \int_M f(x, j_x^k \psi) dx$ on jet space, for $\varphi + \psi$ in a neighborhood V of φ , that we assume convex. We start by deriving a candidate for the function f . According to the fundamental theorem of calculus,

$$F(\varphi + \psi) = F(\varphi) + \int_0^1 dt DF_{\varphi+t\psi}(\psi) dx. \quad (31)$$

As discussed at the beginning of this section, since we assume that $\text{WF}(F_\varphi^{(1)}) = \emptyset$ for every $\varphi \in U$, there exists a unique smooth compactly supported function $x \mapsto \nabla F_\varphi(x)$ such that:

$$F_\varphi^{(1)}(\psi) = \int_M dx \nabla F_\varphi(x) \psi(x). \quad (32)$$

Therefore equation (31) reads:

$$F(\varphi + \psi) = F(\varphi) + \int_0^1 dt \int_M \nabla F_{\varphi+t\psi}(x) \psi(x) dx. \quad (33)$$

We show that Fubini's theorem can be applied to the function $\chi : (x, t) \mapsto \nabla_{\varphi+t\psi} F(x) \psi(x)$. By Prop. 3.4, $F^{(1)}$ is locally compactly supported, so that there is a convex neighborhood V of φ and a compact subset K of Ω such that $F_{\varphi+\psi}^{(1)}$ is supported in K for every $\varphi + \psi \in V$. The function χ is defined on $[0, 1] \times K$ and supported on K for fixed $t \in [0, 1]$. Moreover, by imposing the additional assumption carried by item 2 in theorem 6.3, namely that $\varphi \mapsto \nabla F_\varphi$ be Bastiani

smooth from U to $\mathcal{D}(M)$, the support property of F implies that the image of $\nabla F_{\varphi+t\psi}$ is actually in $\mathcal{D}(K)$ and ∇F is smooth from V to $\mathcal{D}(K)$ because the topology induced on $\mathcal{D}(K)$ by $\mathcal{D}(M)$ is the Fréchet topology of $\mathcal{D}(K)$ determined by the seminorms $\pi_{m,K}$ [27, p. 172]. Since $\mathcal{D}(K)$ injects continuously in $(C^0(K), \pi_{0,K})$, $\varphi \mapsto D_\varphi F$ is a continuous $(C^0(K), \pi_{0,K})$ -valued map. This implies that $(t, x) \mapsto \nabla F_{\varphi+t\psi}(x)$ is continuous as a \mathbb{K} -valued function on $[0, 1] \times K$. Hence so is the integrand of (33), Fubini theorem holds and we obtain:

Lemma 6.13. *Let U be an open subset of $E = C^\infty(M)$ and $F : E \rightarrow \mathbb{K}$ be Bastiani smooth. Assume that for every $\varphi \in U$, $\text{WF}(F_\varphi^{(1)}) = \emptyset$ and $F^{(1)} : U \rightarrow \mathcal{D}(M)$ is Bastiani smooth, then, for every $\varphi \in U$, there is a convex neighborhood V of φ such that, if $\varphi + \psi \in V$, then*

$$F(\varphi + \psi) = F(\varphi) + \int_M dx \int_0^1 \nabla F_{\varphi+t\psi}(x) \psi(x) dt. \quad (34)$$

From now on, we consider $\varphi \in U$ to be fixed. Our candidate for $f(j_x^k \psi)$ is

$$c_\psi(x) = \int_0^1 \nabla F_{\varphi+t\psi}(x) dt \psi(x). \quad (35)$$

By definition and lemma 6.13, for all ψ such that $\varphi + \psi \in V$,

$$F(\varphi + \psi) = F(\varphi) + \int_M c_\psi(x) dx.$$

To show that $c_\psi(x)$ is the right candidate we first need

Proposition 6.14. *The function c_ψ depends only on a finite jet of ψ . More precisely, for every $\varphi \in U$, there is a convex neighborhood V of φ and an integer $k \geq 0$ such that, for all $x \in M$, for every ψ_1 and ψ_2 such that $\varphi + \psi_1$ and $\varphi + \psi_2$ are in V and $j_x^k \psi_1 = j_x^k \psi_2$, then $c_{\psi_1}(x) = c_{\psi_2}(x)$.*

The beginning of the proof is inspired by Ref. [17]. For fixed $\varphi \in U$, by Proposition 3.11, there exists an integer k , a compact K and a convex neighborhood V of φ such that the order of $F_{\varphi+\psi}^{(2)}$ is smaller than k and the support of $DF_{\varphi+\psi}$ is in K if $\varphi + \psi \in V$.

Let us choose some point $x_0 \in M$. Consider a pair ψ_1, ψ_2 of smooth functions such that $\psi_1(x_0) = \psi_2(x_0)$. Then,

$$\begin{aligned} c_{\psi_1}(x_0) - c_{\psi_2}(x_0) &= \int_0^1 dt (\nabla F_{\varphi+t\psi_2}(x_0) \psi_2(x_0) - \nabla F_{\varphi+t\psi_1}(x_0) \psi_1(x_0)) \\ &= \psi_1(x_0) \int_0^1 dt (\nabla F_{\varphi+t\psi_2}(x_0) - \nabla F_{\varphi+t\psi_1}(x_0)). \end{aligned}$$

We use the fundamental theorem of analysis again for $DF_\varphi(h) = \int_M dx \nabla F_\varphi(x) h(x)$ for an arbitrary $h \in C^\infty(M)$ to get

$$DF_{\varphi+t\psi_2}(h) - DF_{\varphi+t\psi_1}(h) = t \int_0^1 ds \langle F_{\varphi+t\psi_1+st(\psi_2-\psi_1)}^{(2)}, (\psi_2 - \psi_1) \otimes h \rangle.$$

Now we take a sequence of smooth functions $(h_n)_{n \in \mathbb{N}}$ which converges to δ_{x_0} in $\mathcal{D}'(M)$ when n goes to infinity and show that both the left and right hand side have limits. For the left hand side, the distribution $DF_{\varphi+t\psi_i}$ being smooth, it defines the continuous form $u \mapsto DF_{\varphi+t\psi_i}(u)$ on $\mathcal{D}'(M)$ by duality pairing. By continuity, $DF_{\varphi+t\psi_i}(h_n) \rightarrow DF_{\varphi+t\psi_i}(\delta_{x_0})$ and Eq. (32) yields

$$\begin{aligned} DF_{\varphi+t\psi_2}(\delta_{x_0}) - DF_{\varphi+t\psi_1}(\delta_{x_0}) &= \int_M dx (\nabla F_{\varphi+t\psi_2}(x) - \nabla F_{\varphi+t\psi_1}(x)) \delta(x - x_0) \\ &= \nabla F_{\varphi+t\psi_2}(x_0) - \nabla F_{\varphi+t\psi_1}(x_0). \end{aligned}$$

For the right hand side, we know by Lemma 6.9 that for every $s \in [0, 1]$, the wave front set of the distribution $F_{\varphi+t\psi_1+st(\psi_2-\psi_1)}^{(2)}$ is in the conormal C_2 , while the sequence $(\psi_2 - \psi_1) \otimes h_n$ converges

to $(\psi_2 - \psi_1) \otimes \delta_{x_0}$ in $\mathcal{D}'_{N^*(M \times \{x_0\})}$, where $N^*(M \times \{x_0\})$ is the conormal of the submanifold $M \times \{x_0\} \subset M \times M$ in $T^*(M \times M)$. Therefore, by transversality of the wave front sets and hypocontinuity of the duality pairings [95], $\lim_n \langle F_{\varphi+t\psi_1+st(\psi_2-\psi_1)}^{(2)}, (\psi_2 - \psi_1) \otimes h_n \rangle$ exists.

Moreover, still by Lemma 6.9, we have for $\varphi + \psi \in V$:

$$\langle F_{\varphi+\psi}^{(2)}, g \otimes h \rangle = \sum_{|\alpha| \leq k} (-1)^{|\alpha|} \int_M dx \theta_\psi^\alpha(x) g(x) \partial^\alpha h(x),$$

for every $(g, h) \in C^\infty(M)^2$ and all θ_ψ^α belong to $\mathcal{D}(K)$. An integration by parts yields

$$\langle F_{\varphi+\psi}^{(2)}, g \otimes h \rangle = \sum_{|\alpha| \leq k} \int_M dx f_\psi^\alpha(x) h(x) \partial^\alpha g(x),$$

where $f_\psi^\alpha = \sum_\beta \binom{\beta}{\alpha} \partial^{\beta-\alpha} \theta_\psi^\beta$ where the sum is over the multi-indices such that $\beta \geq \alpha$ and $|\beta| \leq k$.

As a consequence, for $\varphi + \psi_1 + \psi_2$ in the convex neighborhood V :

$$\langle F_{\varphi+t\psi_1+st(\psi_2-\psi_1)}^{(2)}, (\psi_2 - \psi_1) \otimes \delta_{x_0} \rangle = \sum_{|\alpha| \leq k} f_{t\psi_1+st(\psi_2-\psi_1)}^\alpha(x_0) \partial^\alpha (\psi_2 - \psi_1)(x_0).$$

If, at the point x_0 , $j_{x_0}^k \psi_1 = j_{x_0}^k \psi_2$, then

$$c_{\psi_1}(x_0) - c_{\psi_2}(x_0) = \psi_1(x_0) \sum_{|\alpha| \leq k} \int_0^1 t dt \int_0^1 ds f_{t\psi_1+st(\psi_2-\psi_1)}^\alpha(x_0) (\partial^\alpha \psi_2(x_0) - \partial^\alpha \psi_1(x_0)) = 0.$$

We showed that c_ψ depends only on the k -jet of ψ at x_0 . Moreover, the number k depends only on V and not on x_0 , so that c_ψ depends on the k -jet for every $x \in M$. In other words there is an integer k and a function f such that $c_\psi(x) = f(x, \psi(x), \dots, \partial^\alpha \psi(x))$ for every $x \in M$, where $1 \leq |\alpha| \leq k$.

We want to show that f is smooth in its arguments hence we now investigate in which manner c_ψ depends on ψ . This suggests to study the regularity of the $\mathcal{D}(K)$ -valued function $\psi \mapsto \nabla F_{\varphi+t\psi} \psi$. More precisely, we need to show that the map $\psi \mapsto \nabla F_{\varphi+t\psi} \psi$ is Bastiani smooth from U to $\mathcal{D}(K)$. This is not completely trivial because the map $x \mapsto \nabla F_{\varphi+t\psi}(x)$ is in $\mathcal{D}(K)$ and ψ in $C^\infty(M)$ and we must check that the product of a function in $\mathcal{D}(K)$ by a function in $C^\infty(M)$ is continuous [24, p. 119].

Lemma 6.15. *If U is an open set in $C^\infty(M)$ and $F : U \rightarrow \mathcal{D}(K)$ is a compactly supported Bastiani-smooth map, then the function $G : U \rightarrow \mathcal{D}(M)$ defined by $G(\varphi) = F(\varphi)\varphi$ is compactly supported Bastiani-smooth with the same support as F .*

Proof. Yoann Dabrowski pointed out to us the following fact. For any compact subset K of Ω , both $\mathcal{D}(\Omega)$ and $C^\infty(\Omega)$ induce on $\mathcal{D}(K)$ the usual Fréchet topology of $\mathcal{D}(K)$ [27, p. 172]. Thus to establish the smoothness of G , it suffices to show that the multiplication $(u, v) \in \mathcal{D}(K) \times C^\infty(M) \mapsto \mathcal{D}(K)$ is continuous then it would be Bastiani smooth and by the chain rule it follows that $\varphi \mapsto (F(\varphi), \varphi) \mapsto F(\varphi)\varphi$ is smooth. Since both $\mathcal{D}(K)$ and $C^\infty(M)$ are Fréchet, the product $\mathcal{D}(K) \times C^\infty(M)$ endowed with the product topology is metrizable and it is enough to prove that the product is sequentially continuous. Indeed, let $(u_n, v_n) \rightarrow (u, v)$ in $\mathcal{D}(K) \times C^\infty(M)$, we can find some cut-off function $\chi \in \mathcal{D}(M)$ such that $\chi = 1$ on the support of all u_n , and for all m , by [76, p. 1351]

$$\begin{aligned} \pi_{m,K}(uv - u_n v_n) &\leq \pi_{m,K}((u - u_n)v\chi) + \pi_{m,K}(u_n \chi(v - v_n)) \\ &\leq 2^m (\pi_{m,K}(u - u_n)\pi_{m,K}(\chi v) + \pi_{m,K}(u_n)\pi_{m,K}((v - v_n)\chi)) \rightarrow 0. \end{aligned}$$

Hence G is smooth. \square

This implies that $\psi \mapsto c_\psi = \int_0^1 \nabla F_{\varphi+t\psi} \psi dt$ is smooth since the above Lemma shows the smoothness of $a(t, \psi) \mapsto \nabla F_{\varphi+t\psi} \psi$ and integration over t conserves smoothness by Lemma 6.11. At this point, Theorem 6.3 follows directly from Proposition 6.4.

6.4 Representation theory of local functionals.

In this section, we discuss the issue of representation of our local functionals and the relations between the functionals $(c_\psi, \nabla F, f(j_x^k \psi))$ which are defined or constructed in the course of our proof of Theorem 6.3. In the sequel, we assume that our manifold M is connected, oriented without boundary, hence we can fix a density dx on M which is also a differential form on M of top degree.

In the sequel, we shall work out all explicit formulas in local charts which means without loss of generality that we work on \mathbb{R}^d and the reference density dx is chosen to be the standard Lebesgue measure. We will denote by $(x, u, u^\alpha)_{|\alpha| \leq k}$ where α are multi-indices, some local coordinates on the jet bundle $J^k(\mathbb{R}^d)$. Introduce the vertical Euler vector field $\rho = \sum u^{(\alpha)} \frac{\partial}{\partial u^{(\alpha)}}$ on the bundle $J^k(\mathbb{R}^d)$. In the manifold case if we work on $J^k(M)$, this vector field is intrinsic since it generates scaling in the fibers of $J^k(M)$. For all multiindex $(\alpha) = (\alpha_1 \dots \alpha_p)$, $\alpha_i \in \{1, \dots, d\}$, introduce the operators $\partial^{(\alpha)} = \partial^{\alpha_1} \dots \partial^{\alpha_p}$ where $\partial^i = \frac{\partial}{\partial x^i} + \sum_\alpha u^{(\alpha i)} \frac{\partial}{\partial u^{(\alpha)}}$ and the Euler–Lagrange operator $EL = u \frac{\partial}{\partial u} + \sum_\alpha (-1)^{|\alpha|} \partial^{(\alpha)} u^{(\alpha)} \frac{\partial}{\partial u^{(\alpha)}}$. Let us discuss the nature of the objects involved, ρ is a vertical vector field and acts on $C^\infty(J^k M)$ as a $C^\infty(M)$ linear map, for $\chi \in C^\infty(\mathbb{R}^d)$, $f \in C^\infty(J^k \mathbb{R}^d)$, $\rho(\chi f) = \chi(\rho f)$. For every $i \in \{1, \dots, d\}$, ∂^i is a vector field on $J^k \mathbb{R}^d$ but it has a horizontal component, therefore it is not $C^\infty(\mathbb{R}^d)$ linear and the Euler–Lagrange operator is not $C^\infty(\mathbb{R}^d)$ –linear either.

What follows is a definition–proposition where we give an intrinsic and global definition of the Euler–Lagrange operator in terms of the operator ∇F associated to a functional.

Proposition 6.16 (Euler-Lagrange operator is intrinsic). *Let U be an open subset of $C^\infty(M)$ and $F : U \rightarrow \mathbb{K}$ a Bastiani smooth local functional. For $\varphi \in U$, if there is an integer k , a neighborhood V of φ , an open subset \mathcal{V} of $J^k M$ and $f \in C^\infty(\mathcal{V})$ such that $x \mapsto f(j_x^k \psi_x)$ is compactly supported and*

$$F(\varphi + \psi) = F(\varphi) + \int_M f(j_x^k \psi) dx$$

whenever $\varphi + \psi \in V$ then in every local chart

$$\nabla F_\varphi = EL(f)(j^k \psi) \tag{36}$$

where $EL(f)(\psi) = \sum_{|\alpha| \leq k} (-1)^{|\alpha|} \left(\partial^{(\alpha)} \left(\frac{\partial f}{\partial u^{(\alpha)}} \right) \right) (j_x^k \psi)$ is the Euler–Lagrange operator and $EL(f)(\psi)$ is uniquely determined by F .

The above proposition means that $EL(f)$ does not depend on the choice of representative f and is intrinsic (i.e. it does not depend on the choice of a local chart).

Proof. Indeed, assume that we make a small perturbation $\varphi + \psi$ of the background field φ by ψ which is compactly supported in some open chart U of M . Then a local calculation yields

$$\begin{aligned} DF_\varphi(\psi) &= \sum_\alpha \int_M \frac{\partial f}{\partial u^{(\alpha)}(x)} \psi^{(\alpha)}(x) dx \\ &= \sum_\alpha (-1)^{|\alpha|} \int_M \psi(x) \sum_{|\alpha| \leq k} (-1)^{|\alpha|} \left(\partial^{(\alpha)} \left(\frac{\partial f}{\partial u^{(\alpha)}} \right) \right) (j_x^k \psi) dx, \end{aligned}$$

where we used an integration by parts to recover the Euler-Lagrange operator and all boundary terms vanish since f is compactly supported in x and $\psi \in \mathcal{D}(U)$. We have just proved that for all open chart $U \subset M$, $\nabla F_\varphi|_U = EL(f)|_U$. But ∇F_φ is intrinsically defined on M therefore so is $EL(f)$ and we have the equality $\nabla F = EL(f)$. The unique determination of ∇F_φ follows from Lemma 6.2. \square

Theorem 6.17. [Global Poincaré] *Assume that M is a smooth, connected, oriented manifold without boundary. Let U be an open subset of $C^\infty(M)$ and $F : U \rightarrow \mathbb{K}$ a Bastiani smooth local functional. Then the following statements are equivalent:*

- two functions $(f_1, f_2) \in C^\infty(\mathcal{V})$ for \mathcal{V} an open subset of the jet space $J^k M$, are two representations of F in a neighborhood V of $\varphi \in U$:

$$F(\varphi + \psi) = F(\varphi) + \int_M f_1(j_x^k \psi) dx = F(\varphi) + \int_M f_2(j_x^k \psi) dx$$

whenever $\overline{\varphi + \psi} \in V$

- for all $\psi \in V - \varphi$:

$$f_1(j_x^k \psi) dx - f_2(j_x^k \psi) dx = d\beta(j_x^{2k} \psi), \quad (37)$$

where $\beta(j_x^{2k} \psi) \in \Omega_c^{d-1}(M)$ is a differential form of degree $d - 1$ whose value at a point x depends only on the $2k$ -jet of ψ at x .

Let us stress that we do not need to constraint the topology of M in the above Theorem only the compactness of the support of $f_i(j_x^k \psi) dx, i \in \{1, 2\}$ really matters.

Proof. One sense of the equivalence is trivial since the integral of a compactly supported exact form on M always vanishes. By Proposition 6.4, we know that the map $\psi \in V - \varphi \mapsto j^k \psi$ has an open image in $J^k(M)$ denoted \mathcal{V} and we only need (f_1, f_2) to be defined on \mathcal{V} . We denote by $(x, u, u^\alpha)_{|\alpha| \leq k}$ where α are multi-indices, some local coordinates on the jet bundle $J^k(\mathbb{R}^d)$. We use the vertical Euler vector field $\rho = \sum u^{(\alpha)} \frac{\partial}{\partial u^{(\alpha)}}$ on the bundle $J^k(\mathbb{R}^d)$. For every multiindex $(\alpha) = (\alpha_1 \dots \alpha_p), \alpha_i \in \{1, \dots, d\}$, introduce the operators $\partial^{(\alpha)} = \partial^{\alpha_1} \dots \partial^{\alpha_p}$ where $\partial^i = \frac{\partial}{\partial x^i} + \sum_\alpha u^{(\alpha i)} \frac{\partial}{\partial u^{(\alpha)}}$ and the Euler-Lagrange operator reads $EL = u \frac{\partial}{\partial u} + \sum_\alpha (-1)^{|\alpha|} \partial^{(\alpha)} u^{(\alpha)} \frac{\partial}{\partial u^{(\alpha)}}$.

We shall prove two related identities, in local chart

$$(\rho f)(j^k \psi) dx = (uEL(f))(j^k \psi) dx + d \left(\sum_{\mu=1}^d j_\mu(j^{2k} \psi) \frac{\partial}{\partial x^\mu} \lrcorner dx \right) \quad (38)$$

$$\begin{aligned} f(j^k(\psi_1 + \psi_2)) dx &= f(j^k \psi_1) dx + \int_0^1 dt \psi_2 EL(f)(j^k(\psi_1 + t\psi_2)) dx \\ &+ d \left(\int_0^1 \frac{dt}{t} j_\mu(j^{2k}(\psi_1 + t\psi_2)) \partial_{x^\mu} \lrcorner dx \right). \end{aligned} \quad (39)$$

For all $(f, g) \in C^\infty(J^k \mathbb{R}^d)^2$ and all multiindices α , the generalized Leibniz-like identity holds true:

$$(\partial^{\alpha_1} \dots \partial^{\alpha_p} f)g = \underbrace{\sum_{i=1}^p (-1)^{i+1} \partial^{\alpha_i} ((\partial^{\alpha_{i+1}} \dots \partial^{\alpha_p} f) \partial^{\alpha_{i-1}} \dots \partial^{\alpha_1} g)} + (-1)^p f(\partial^{\alpha_p} \dots \partial^{\alpha_1} g)$$

where the underbraced terms are all total derivatives. Using this we derive the following key identity which is valid on jet spaces. For all $f dx \in C^\infty(J^k \mathbb{R}^d) \otimes \Omega^d(\mathbb{R}^d)$:

$$\begin{aligned} (\rho f) dx &= \sum u^{(\alpha)} \frac{\partial f}{\partial u^{(\alpha)}} dx = u \frac{\partial f}{\partial u} + \sum_{|\alpha| \geq 1} u^{(\alpha)} \frac{\partial f}{\partial u^{(\alpha)}} dx \\ &= uEL(f) dx + \sum_{\mu=1}^d \partial^\mu j_\mu(j^{2k} \psi) dx = uEL(f) dx + d \left(\sum_{\mu=1}^d j_\mu(j^{2k} \psi) \frac{\partial}{\partial x^\mu} \lrcorner dx \right) \end{aligned}$$

where $j_\mu \in C^\infty(J^{2k} \mathbb{R}^d)$ is a local functional.

To prove the second identity, we shall use the fundamental Theorem of calculus and the first identity :

$$\begin{aligned} f(j^k(\psi_1 + \psi_2)) dx &= f(j^k \psi_1) dx + \int_0^1 \frac{dt}{t} (\rho f)(j^k(\psi_1 + t\psi_2)) dx \\ &= f(j^k \psi_1) dx + \int_0^1 dt \psi_2 EL(f)(j^k(\psi_1 + t\psi_2)) dx \\ &+ d \left(\int_0^1 \frac{dt}{t} j_\mu(j^{2k}(\psi_1 + t\psi_2)) \partial_{x^\mu} \lrcorner dx \right) \end{aligned}$$

To prove the claim of the Lemma is equivalent to show the following statement: if a local functional F is locally constant i.e. $F(\varphi + \psi) = F(\varphi)$ whenever $\varphi + \psi \in V$, then $F(\varphi + \psi) = F(\varphi) + \int_M d\beta(j_x^{2k}\psi)$ and $\beta(j_x^{2k}\psi) \in \Omega_c^{n-1}(M)$ is a compactly supported $n - 1$ form.

$$\begin{aligned} \forall \psi \in V - \varphi, F(\varphi + t\psi) = F(\varphi) &\implies \int_{\mathbb{R}^d} \int_0^1 \frac{dt}{t} (\rho f)(j_x^k(t\psi)) dx = 0, \forall \psi \in V - \varphi \\ &\implies \int_{\mathbb{R}^d} \int_0^1 dt (\psi EL(f)(j_x^k(t\psi))) dx = 0, \forall \psi \in V - \varphi. \end{aligned}$$

This means that $EL(f) = 0$ therefore on any **open chart** U (U is contractible), Eq. (39) yields

$$f(j_x^p(\psi)) = f(0) + d \int_0^1 dt \left(\sum_{\mu} j_{\mu}(j_x^{2p}(t\psi)) \partial_{x^{\mu}} \right) dx.$$

We want to prove that $EL(f) = 0 \implies f(j_x^k(\psi))dx - f(0)dx|_{M_{p+1}} = d\beta(j^{2k}\psi)$ where $\beta \in C^{\infty}(J^{2p}(M))$ knowing that this holds true on any local chart, and that $EL(f) = 0$ is equivalent to assuming that $F(\varphi + \psi) := \int_M f(j_x^k\psi)dx$ is locally constant. We cover M by some countable union $\cup_{i \in \mathbb{N}} U_i$ of contractible open charts such that every element $x \in M$ belongs to a **finite number** of charts U_i , set $M_p = (U_1 \cup \dots \cup U_p)$ and we arrange the cover in such a way that $M_p \cap U_{p+1} \neq \emptyset$ for all p which is always possible. Assume by induction on p that

$$EL(f) = 0, \text{supp}(f) \subset M_p \implies f(j_x^k(\psi))dx - f(0)dx|_{M_p} = d\beta(j^{2k}\psi)$$

where $\beta \in C^{\infty}(J^{2p}(M)) \otimes \Omega_c^{d-1}(M_p)$.

We want to prove that $EL(f) = 0, \text{supp}(f) \subset M_{p+1} \implies f(j_x^k(\psi))dx - f(0)dx|_{M_{p+1}} = d\beta(j^{2k}\psi)$ where $\beta \in C^{\infty}(J^{2p}(M)) \otimes \Omega_c^{d-1}(M_{p+1})$. Choose a partition of unity $(\chi, 1 - \chi)$ subordinated to $M_p \cup U_{p+1}$, the key idea is to decompose the variation ψ of the background field φ as the sum of two components $\chi\psi + (1 - \chi)\psi$ where $\chi\psi$ (resp $(1 - \chi)\psi$) vanishes outside U_p (resp U_{p+1}) which yields :

$$f(j^k\psi) = f(j^k(\chi\psi + (1 - \chi)\psi)) - f(j^k((1 - \chi)\psi)) + f(j^k((1 - \chi)\psi)) - f(0) + f(0).$$

The second idea is to note that for every fixed ψ , the new functional

$$\phi \mapsto \tilde{f}(j^k\phi) = f(j^k(\chi\phi + (1 - \chi)\psi)) - f(j^k((1 - \chi)\psi))$$

has **trivial** Euler–Lagrange equation $EL(\tilde{f})(j^k\phi) = EL(f)(j^k(\chi\phi + (1 - \chi)\psi)) = 0$ since $EL(f) = 0$ and its support is contained in M_p . Therefore :

$$f(j^k\psi) = \tilde{f}(j^k\psi) + f(j^k((1 - \chi)\psi)) - f(0) + f(0) = d\tilde{\beta}(j^{2k}\psi) + \underbrace{f(j^k((1 - \chi)\psi)) - f(0)}_{\text{bracket}} + f(0)$$

by the inductive assumption. To treat the term under brace, define a new functional $\psi \mapsto g(j^k\psi) = f(j^k((1 - \chi)\psi)) - f(0)$ whose support is contained in U_{p+1} and whose Euler-Lagrange equation vanishes, $EL(g) = 0$ again by the fact that $EL(f) = 0$. Since U_{p+1} is contractible we know that $f(j^k((1 - \chi)\psi)) - f(0) = d\alpha(j^{2k}\psi)$ where $\alpha \in C^{\infty}(J^{2k}M) \otimes \Omega_c^{d-1}(U_{p+1})$ and therefore we found that

$$f(j^k\psi)dx = d\beta(j^{2k}\psi) + f(0)dx$$

$\beta \in C^{\infty}(J^{2k}M) \otimes \Omega_c^{d-1}(M_{p+1})$. Therefore for all $\psi \in V - \varphi$, $f(j_x^k(\psi))dx = d\beta(j^{2k}\psi) + f(0)dx$. Now we conclude by using the fact that F is a constant functional thus $0 = F(\varphi + \psi) - F(\varphi) = \int_M (f(0)dx + d\beta(j^{2k}\psi)) = \int_M f(0)dx$. But $f(0)dx$ is a top form in $\Omega_c^d(M)$ which does not depend on ψ and whose integral over M vanishes hence $f(0)dx = dk$ for some $k \in \Omega_c^{d-1}(M)$ since $H_c^d(M, \mathbb{R}) \simeq \mathbb{R}$ for the top de Rham cohomology with compact support when M is **connected** [96, Theorem 17.30 p. 454]. \square

The next Theorem summarizes the above results :

Theorem 6.18. *Let U be an open subset of $C^\infty(M)$ and $F : U \rightarrow \mathbb{K}$ a Bastiani smooth local functional. For $\varphi \in U$, if there is an integer k , a neighborhood V of φ , an open subset \mathcal{V} of $J^k M$ and $f \in C^\infty(\mathcal{V})$ such that $x \mapsto f(j^k \psi_x)$ compactly supported and*

$$F(\varphi + \psi) = F(\varphi) + \int_M f(j_x^k \psi) dx$$

whenever $\varphi + \psi \in V$ then in every local chart

$$\nabla F_\varphi = EL(f)(j^k \psi) \quad (40)$$

where $EL(f)(\psi) = \sum_{|\alpha| \leq k} (-1)^{|\alpha|} \left(\partial^{(\alpha)} \left(\frac{\partial f}{\partial u^{(\alpha)}} \right) \right) (j_x^k \psi)$ is the Euler–Lagrange operator and $EL(f)(\psi)$ is uniquely determined by F .

Furthermore, we find that :

$$F(\varphi + \psi) = F(\varphi) + \int_M f(j_x^k \psi) dx = F(\varphi) + \int_M \left(\int_0^1 dt \psi EL(f)(t\psi) \psi \right) dx \quad (41)$$

where $f(j_x^k \psi) - \left(\int_0^1 dt \psi EL(f)(t\psi) \psi \right) dx = d\beta(j_x^{2k} \psi)$ and $\beta(j_x^{2k} \psi) \in \Omega_c^{d-1}(M)$ is a compactly supported $d-1$ form.

6.4.1 Explicit forms

In this section we derive the explicit expression of ∇F_φ and $F^{(\alpha)}(\varphi)$ in terms of f when $M = \mathbb{R}^d$. Since the general expression is not very illuminating, let us start with the following simple example:

$$F(\varphi) = \int_M h(x) \varphi^4(x) + g^{\mu\nu}(x) \partial_\mu \varphi(x) \partial_\nu \varphi(x) dx,$$

where h and $g^{\mu\nu}$ are smooth and compactly supported and $g^{\mu\nu}$ is symmetric. We compute

$$\begin{aligned} DF_\varphi(u) &= 2 \int_M dx 2h(x) \varphi^3(x) u(x) + g^{\mu\nu}(x) \partial_\mu \varphi(x) \partial_\nu u(x) \\ &= 2 \int_M dx \left(2h(x) \varphi^3(x) - \partial_\nu (g^{\mu\nu}(x) \partial_\mu \varphi(x)) \right) u(x), \end{aligned}$$

where we used integration by parts. Thus,

$$\nabla F_\varphi(x) = 4h(x) \varphi^3(x) - 2\partial_\nu (g^{\mu\nu}(x) \partial_\mu \varphi(x)).$$

Moreover,

$$D^2 F_\varphi(u, v) = 2 \int_M dx u(x) \left(6h(x) \varphi^2(x) v(x) - \partial_\nu (g^{\mu\nu}(x) \partial_\mu v(x)) \right).$$

To write this as a distribution, we need to integrate over two variables:

$$D^2 F_\varphi(u, v) = 2 \int_{M^2} dx dy u(x) \delta(x-y) \left(6h(y) \varphi^2(y) v(y) - \partial_\nu (g^{\mu\nu}(y) \partial_\mu v(y)) \right).$$

Now we can use integration by parts over y to recover $v(y)$:

$$D^2 F_\varphi(u, v) = \sum_\alpha \int_{M^2} dx dy u(x) v(y) f^\alpha(\varphi)(y) \partial_y^\alpha \delta(x-y),$$

where the non-zero $f^\alpha(\varphi)$ are

$$\begin{aligned} f^0(\varphi)(y) &= 12\varphi^2(y), \\ f^\mu(\varphi)(y) &= -\partial_\nu g^{\mu\nu}(y), \\ f^{\mu\nu}(\varphi)(y) &= -g^{\mu\nu}(y). \end{aligned}$$

More generally

Proposition 6.19. *If*

$$F(\varphi) = \int_M f(\varphi^{(\alpha)}(x))dx,$$

then

$$F^\alpha(\varphi)(x) = \sum_{\beta \leq \gamma} (-1)^{|\beta|} \binom{\beta}{\gamma} \partial_{y^{\beta-\gamma}} \frac{\partial^2 f}{\partial \varphi^{(\alpha-\gamma)}(x) \partial \varphi^{(\beta)}(x)}.$$

Proof. The proof is a straightforward generalization of the example. Indeed,

$$DF_\varphi(u) = \sum_\alpha \int_M \frac{\partial f}{\partial \varphi^{(\alpha)}(x)} u^{(\alpha)}(x) dx = \sum_\alpha (-1)^{|\alpha|} \int_M u(x) \frac{\partial^{|\alpha|}}{\partial x^\alpha} \frac{\partial f}{\partial \varphi^{(\alpha)}(x)} dx,$$

where we used an integration by parts to recover the Euler-Lagrange operator. The second derivative is

$$D^2 F_\varphi(u, v) = \sum_{\alpha\beta} (-1)^{|\alpha|} \int_M u(x) \frac{\partial^{|\alpha|}}{\partial x^\alpha} \left(\frac{\partial f^2}{\partial \varphi^{(\alpha)}(x) \partial \varphi^{(\beta)}(x)} v^{(\beta)}(x) \right) dx.$$

We write this as a double integral

$$D^2 F_\varphi(u, v) = \sum_{\alpha\beta} (-1)^{|\alpha|} \int_{M^2} u(x) \delta(x-y) \partial_{y^\alpha} \left(\frac{\partial f^2}{\partial \varphi^{(\alpha)}(y) \partial \varphi^{(\beta)}(y)} v^{(\beta)}(y) \right) dx dy.$$

A first integration by parts gives us

$$D^2 F_\varphi(u, v) = \sum_{\alpha\beta} \int_{M^2} u(x) \left(\frac{\partial f^2}{\partial \varphi^{(\alpha)}(y) \partial \varphi^{(\beta)}(y)} v^{(\beta)}(y) \right) \partial_{y^\alpha} \delta(x-y) dx dy.$$

A second integration by parts isolates $v(y)$:

$$D^2 F_\varphi(u, v) = \sum_{\alpha\beta} (-1)^{|\beta|} \sum_{\gamma \leq \beta} \binom{\beta}{\gamma} \int_{M^2} u(x) v(y) \left(\partial_{y^{\beta-\gamma}} \frac{\partial f^2}{\partial \varphi^{(\alpha)}(y) \partial \varphi^{(\beta)}(y)} \right) \partial_{y^{\alpha+\gamma}} \delta(x-y) dx dy.$$

□

If we calculate higher differentials $D^k F_\varphi(u_1, \dots, u_k)$ we see that we always obtain products of smooth functions by derivatives of products of delta functions. This shows that the wavefront set of $F_\varphi^{(k)}$ is in the conormal C_k .

7 Peetre theorem for local and multilocal functionals

In this section, we propose an alternative characterization of local functionals in terms of a non-linear Peetre theorem. We do not characterize the locality of the action F but the locality of the Lagrangian density, that we denoted ∇F in the previous section. We first state our theorems for local functionals, and then we prove them for the case of multilocal functionals, which are a natural generalization of local functionals in quantum field theory. Our proof is inspired by recent works on the Peetre theorem [97, 98], however it is formulated in the language of Bastiani smoothness and uses simpler assumptions than Slovák's paper [99].

7.1 Peetre theorem for local functionals

Let Ω be some open set in a manifold M . We first begin with an alternative definition of a local map from $C^\infty(\Omega)$ to itself, that we call *Peetre local*.

Definition 7.1. *A map $F : C^\infty(\Omega) \mapsto C^\infty(\Omega)$ is Peetre local if for every $x \in \Omega$, if $\varphi_1 = \varphi_2$ on some neighborhood of x then $F(\varphi_1)(x) = F(\varphi_2)(x)$.*

The relation with the additivity condition is given by

Proposition 7.2. *Let $F : C^\infty(\Omega) \mapsto C^\infty(\Omega)$ be Peetre local. For every $(\varphi_1, \varphi_2) \in C^\infty(\Omega)^2$ if $\text{supp } \varphi_1$ and $\text{supp } \varphi_2$ do not meet then*

$$\forall x \in \Omega, \forall \varphi, F(\varphi_1 + \varphi_2 + \varphi)(x) = F(\varphi_1 + \varphi)(x) + F(\varphi_2 + \varphi)(x) - F(\varphi)(x). \quad (42)$$

Proof. If $x \notin (\text{supp } \varphi_1 \cup \text{supp } \varphi_2)$ then $\varphi_1 = \varphi_2 = 0$ in some neighborhood of x , it follows that $F(\varphi_1 + \varphi_2 + \varphi)(x) = F(0 + 0 + \varphi)(x) = F(\varphi)(x)$ and $F(\varphi_1 + \varphi)(x) + F(\varphi_2 + \varphi)(x) - F(\varphi)(x) = 2F(\varphi)(x) - F(\varphi)(x) = F(\varphi)(x)$ hence Eq. (42) holds true.

If $x \in \text{supp } \varphi_1$ then necessarily there is some neighborhood U of x on which $\varphi_2|_U = 0$ hence $\varphi_1 + \varphi_2 + \varphi|_U = \varphi_1 + \varphi|_U$ and $F(\varphi_1 + \varphi_2 + \varphi)(x) = F(\varphi_1 + \varphi)(x)$. Also $F(\varphi_1 + \varphi)(x) + F(\varphi_2 + \varphi)(x) - F(\varphi)(x) = F(\varphi_1 + \varphi)(x) + F(\varphi)(x) - F(\varphi)(x) = F(\varphi_1 + \varphi)(x)$ hence again Eq. (42) holds true. The case where $x \in \text{supp } \varphi_2$ can be treated by similar methods which yields the final result. \square

The Peetre theorem for local functionals is

Theorem 7.3. *Let $F : C^\infty(\Omega) \rightarrow C^\infty(\Omega)$ be a Bastiani smooth Peetre local map. Then, for every $\varphi \in C^\infty(\Omega)$ there is a neighborhood V of φ in $C^\infty(\Omega)$ and an integer k such that for all ψ such that $\varphi + \psi \in V$, $F(\varphi + \psi)(x) = c(j^k \psi_x)$ for some smooth function c on $J^k \Omega$.*

In other words, if F is a Bastiani smooth Peetre local map, then for every $g \in \mathcal{D}(M)$, $\int_M F(\varphi)g$ is a Bastiani smooth local map in the sense of the rest of the paper. This relation between a priori different concepts of locality strongly supports the idea that our definition is a natural one.

If F is only assumed to be a continuous local map, then a similar theorem exists for which the function c is not necessarily smooth. These theorems are proved in the next section for the more general case of multilocal functionals.

7.2 Multilocal functionals and first Peetre theorem

By generalizing Definition 7.1 of local maps, we can define multilocal maps. These maps appear naturally in quantum field theory as the product of several Lagrangian densities $\mathcal{L}(x_1) \dots \mathcal{L}(x_k)$.

Definition 7.4. *Let k be an integer. A map $F : C^\infty(\Omega) \mapsto C^\infty(\Omega^k)$ is k -local if for every $(x_1, \dots, x_k) \in \Omega^k$, if $\varphi_1 = \varphi_2$ on some neighborhood of $\{x_1, \dots, x_k\} \subset \Omega$ then $F(\varphi_1)(x_1, \dots, x_k) = F(\varphi_2)(x_1, \dots, x_k)$.*

The *multilocal maps* are the maps that are k -local for some k . We emphasize that **Peetre local** maps in the sense of definition 7.1 correspond with **1-local maps** in the above sense. In the sequel, for M a smooth manifold, we denote by $J^p M^{\boxtimes k}$ the bundle over M^k whose fiber over a k -uple of points $(x_1, \dots, x_k) \in M^k$ is $J^p M_{x_1} \times \dots \times J^p M_{x_k}$.

Theorem 7.5. *Let $F : C^\infty(\Omega) \mapsto C^\infty(\Omega^k)$ be a continuous k -local map. Then, for every $\varphi \in C^\infty(\Omega)$ there is a neighborhood V of φ in $C^\infty(\Omega)$, $p \in \mathbb{N}$ such that for all ψ such that $\varphi + \psi \in V$, $F(\varphi + \psi)(x_1, \dots, x_k) = c(j^p \psi_{x_1}, \dots, j^p \psi_{x_k})$ for some function $c : J^p M^{\boxtimes k}|_{(M^k \setminus D_k)} \rightarrow M^k$, where $M^k \setminus D_k$ denotes the configuration space M^k minus all diagonals.*

Proof. Fix a k -tuple of points $(x_1, \dots, x_k) \in \Omega^k$ and some compact neighborhood K of (x_1, \dots, x_k) in Ω^k . Continuity of F implies that for all $\varepsilon > 0$, there exists $\eta > 0$ and a seminorm $\pi_{m,K'}$ of $C^\infty(\Omega)$ such that

$$\pi_{m,K'}(\varphi_1 - \varphi_2) \leq \eta \implies \sup_{(y_1, \dots, y_k) \in K} |F(\varphi_1)(y_1, \dots, y_k) - F(\varphi_2)(y_1, \dots, y_k)| \leq \varepsilon.$$

Assume that (φ_1, φ_2) have same $(m+1)$ -jets at $\{x_1, \dots, x_k\}$. Let $(\chi_\lambda)_\lambda$ be the family of compactly supported cut-off functions equal to 1 in some neighborhood of $X = \{x_1, \dots, x_k\}$ defined in lemma 7.6. It follows that $\varphi_{1,\lambda} = \varphi_1 \chi_\lambda$ (resp. $\varphi_{2,\lambda} = \varphi_2 \chi_\lambda$) coincides with φ_1 (resp. φ_2) near $\{x_1, \dots, x_k\}$. Hence, for all $\lambda > 0$, $F(\varphi_{1,\lambda})(x_1, \dots, x_k) = F(\varphi_1)(x_1, \dots, x_k)$ and $F(\varphi_{2,\lambda})(x_1, \dots, x_k) = F(\varphi_2)(x_1, \dots, x_k)$. Set $\varepsilon_n = \frac{1}{2^n}$ then there exists η_n such that

$$\pi_{m,K'}(\psi_1 - \psi_2) \leq \eta_n \implies \sup_{(y_1, \dots, y_k) \in K} |F(\psi_1)(y_1, \dots, y_k) - F(\psi_2)(y_1, \dots, y_k)| \leq \frac{1}{2^n}.$$

Therefore it suffices to find some sequence $\lambda_n \rightarrow 0$ such that $\pi_{m,K'}(\varphi_{1,\lambda_n} - \varphi_{2,\lambda_n}) \leq \eta_n$. Since $\varphi_1 - \varphi_2$ vanishes at order $m+1$ on the set $X = \{x_1, \dots, x_k\}$, Lemma 7.6 yields the estimate

$$|\pi_{m,K'}(\varphi_{1,\lambda} - \varphi_{2,\lambda})| \leq \tilde{C} \lambda \pi_{m+1,K}(\varphi_1 - \varphi_2),$$

which implies that $\lim_{\lambda \rightarrow 0} \pi_{m,K'}(\varphi_{1,\lambda} - \varphi_{2,\lambda}) = \lim_{\lambda \rightarrow 0} \pi_{m,K'}((\varphi_1 - \varphi_2)\chi_\lambda) = 0$.

Finally, we obtain that if φ_1, φ_2 have same $(m+1)$ -jet at $X = \{x_1, \dots, x_k\}$ then for all $n > 0$:

$$|F(\varphi_1)(x_1, \dots, x_k) - F(\varphi_2)(x_1, \dots, x_k)| = |F(\varphi_{1,\lambda_n})(x_1, \dots, x_k) - F(\varphi_{2,\lambda_n})(x_1, \dots, x_k)| \leq \frac{1}{2^n} \quad (43)$$

which implies that $F(\varphi_1)(x_1, \dots, x_k) = F(\varphi_2)(x_1, \dots, x_k)$. \square

Lemma 7.6. *Let X be any closed subset of \mathbb{R}^d . Let $\mathcal{I}^{m+1}(X, \mathbb{R}^d)$ denote the closed ideal of functions of regularity C^{m+1} which vanish at order $m+1$ on X . Then there is a function $\chi_\lambda \in C^\infty(\mathbb{R}^d)$ parametrized by $\lambda \in (0, 1]$ s.t. $\chi_\lambda = 1$ (resp $\chi_\lambda = 0$) when $d(x, X) \leq \frac{\lambda}{8}$ (resp $d(x, X) \geq \lambda$) such that for all compact subset $K \subset \mathbb{R}^d$,*

$$\exists \tilde{C}, \forall \lambda \in (0, 1], \forall \varphi \in \mathcal{I}^{m+1}(X, \mathbb{R}^d), \pi_{m,K}(\chi_\lambda \varphi) \leq \tilde{C} \lambda \pi_{m+1, K \cap \{d(x, X) \leq \lambda\}}(\varphi). \quad (44)$$

Proof. Choose $\phi \geq 0$ s.t. $\int_{\mathbb{R}^d} \phi(x) d^d x = 1$ and $\phi = 0$ if $|x| \geq \frac{3}{8}$. Then set $\phi_\lambda = \lambda^{-d} \phi(\lambda^{-1} \cdot)$ and set α_λ to be the characteristic function of the set $\{x \text{ s.t. } d(x, X) \leq \frac{\lambda}{2}\}$ then the convolution product $\chi_\lambda = \phi_\lambda * \alpha_\lambda$ satisfies $\chi_\lambda(x) = 1$ if $d(x, X) \leq \frac{\lambda}{8}$ and $\chi_\lambda(x) = 0$ if $d(x, X) \geq \lambda$. Since by

Leibniz rule one has $\partial^\alpha(\chi_\lambda \varphi)(x) = \sum_{|k| \leq |\alpha|} \binom{\alpha}{k} \partial^k \chi_\lambda \partial^{\alpha-k} \varphi(x)$, it suffices to estimate each term

$\partial^k \chi_\lambda \partial^{\alpha-k} \varphi(x)$ of the above sum. For every multi-index k , there is some constant C_k such that $\forall x \in \mathbb{R}^d \setminus X, |\partial_x^k \chi_\lambda| \leq \frac{C_k}{\lambda^{|k|}}$ and $\text{supp } \partial_x^k \chi_\lambda \subset \{d(x, X) \leq \lambda\}$. Therefore for all $\varphi \in \mathcal{I}^{m+1}(X, \mathbb{R}^d)$, for all $x \in \text{supp } \partial_x^k \chi_\lambda \partial^{\alpha-k} \varphi$, for $y \in X$ such that $d(x, X) = |x-y|$, we find that $\partial^{\alpha-k} \varphi$ vanishes at y at order $|k|+1$. Indeed φ vanishes at order $m+1$ hence $\partial^{\alpha-k} \varphi$ vanishes at order $m+1-|\alpha|+k \geq k+1$ since $|\alpha| \leq m$. Therefore:

$$\partial_x^{\alpha-k} \varphi(x) = \sum_{|\beta|=|k|+1} (x-y)^\beta R_\beta(x),$$

where the right hand side is just the integral remainder in Taylor's expansion of $\partial^{\alpha-k} \varphi$ around y . Hence:

$$|\partial^k \chi_\lambda \partial^{\alpha-k} \varphi(x)| \leq \frac{C_k}{\lambda^{|k|}} \sum_{|\beta|=|k|+1} |(x-y)^\beta R_\beta(x)|.$$

It is easy to see that R_β only depends on the jets of φ of order $\leq m+1$. Hence

$$|\partial^k \chi_\lambda \partial^{\alpha-k} \varphi(x)| \leq C_k \lambda \sup_{x \in K, d(x, X) \leq \lambda} \sum_{|\beta|=|k|+1} |R_\beta(x)|$$

and the conclusion follows easily. \square

7.3 The second Peetre Theorem.

Theorem 7.7. *Let $F : C^\infty(\Omega) \mapsto C^\infty(\Omega^k)$ be a Bastiani smooth k -local map. Then, for every $\varphi \in C^\infty(\Omega)$ there is a neighborhood V of φ in $C^\infty(\Omega)$, $p \in \mathbb{N}$ such that for all ψ such that $\varphi + \psi \in V$, $F(\varphi + \psi)(x_1, \dots, x_k) = c(j^p \psi_{x_1}, \dots, j^p \psi_{x_k})$ for some smooth function c on $J^p M^{\boxtimes k}|_{(M^k \setminus D_k)}$ where $M^k \setminus D_k$ denotes the configuration space M^k minus all diagonals.*

Proof. Without loss of generality, we may assume that $M = \mathbb{R}^d$ and to go back to arbitrary manifolds, we use partitions of unity as in the proof of Lemma 6.6. The coordinates on the jet space $J^p(\mathbb{R}^d)$ are denoted by $(x, p^\alpha)_{|\alpha| \leq p}$. Let (U_1, \dots, U_k) be two by two disjoint open subsets of \mathbb{R}^d , then $U_1 \times \dots \times U_k$ is an open subset of $(\mathbb{R}^d)^k \setminus D_k$. We define the smooth map: $\Phi : (x_1, \dots, x_k; p_1, \dots, p_k) \in J^p(\mathbb{R}^d)^{\boxtimes k}|_{U_1 \times \dots \times U_k} \mapsto (\sum_{1 \leq i \leq k} \frac{p_i \alpha}{\alpha!} (\cdot - x_i)^\alpha \chi_i(\cdot - x_i)) \in C^\infty(\mathbb{R}^d)$ where the functions $\chi_i \in C_c^\infty(\mathbb{R}^d)$ are cut-off functions equal to 1 near 0 and such that for all $(x_1, \dots, x_k) \in U_1 \times \dots \times U_k$, the support of the functions $\chi_i(\cdot - x_i)$ are disjoint on \mathbb{R}^d . Then,

$$(x_1, \dots, x_k; p_1, \dots, p_k), (y_1, \dots, y_k) \mapsto F(\varphi + \Phi(x_1, \dots, x_k; p_1, \dots, p_k))(y_1, \dots, y_k)$$

is smooth by smoothness of F and Φ . Hence, its pull-back on the diagonal $x_1 = y_1, \dots, x_k = y_k$ is also smooth and reads

$$F(\varphi + \Phi(x_1, \dots, x_k; p_1, \dots, p_k))(x_1, \dots, x_k) = c(x_1, \dots, x_k; p_1, \dots, p_k)$$

as the composition of smooth functions and it follows that c is smooth on $J^p M^{\boxtimes k}|_{U_1 \times \dots \times U_k}$. \square

8 Multi-vector fields and graded functionals

In the quantum theory of gauge fields, especially in the Batalin-Vilkovisky approach, it is necessary to deal, not only with functionals as discussed above, but also with multi-vector fields on the configuration space E (assumed to be the space of sections of some vector bundle B) [14]. Such multi-vector fields can be seen as functionals on the graded space $T^*[1]E \doteq E \oplus E^*[1]$, where $E^* \doteq \Gamma(M, B^*)$ is the space of smooth sections. To make this notion precise, we use the ideas presented in [100] and characterize the ‘‘odd’’ space $E^*[1]$ through the space of functions on it, understood as multilinear smooth, totally antisymmetric, functionals. Then we shall make a conjectural claim on the meaning of locality in that context.

8.0.1 Locality of functionals on graded space.

We consider a graded space $E_0 \oplus E_1[1]$, where $E_0 = \Gamma(M, B_0)$ and $E_1 = \Gamma(M, B_1)$ are spaces of smooth sections of finite rank vector bundles B_0 and B_1 over M respectively. Before giving formal definitions, let us explain the idea of our construction. We will first define the space $\mathcal{O}(E_0 \oplus E_1[1])$ to be space of maps from E_0 to \mathcal{A} , where

$$\mathcal{A} \doteq \prod_{k=0}^{\infty} \mathcal{A}^k \doteq \prod_{k=0}^{\infty} \Gamma'_a(M^k, B_1^{\boxtimes k}),$$

satisfying an appropriate smoothness condition. Let us clarify the notation Γ'_a . We first define the iterated wedge product of k elements u_1, \dots, u_k of the space of distributional sections $\Gamma'(M, B_1)$ by

$$\langle u_1 \wedge \dots \wedge u_k, h_1 \otimes \dots \otimes h_k \rangle = \sum_{\sigma} (-1)^\sigma \langle u_1, h_{\sigma(1)} \rangle \dots \langle u_k, h_{\sigma(k)} \rangle,$$

where h_1, \dots, h_k are sections in $\Gamma(M, B_1)$ and σ runs over the permutations of $\{1, \dots, k\}$. Then, the k -th exterior power $\Lambda^k \Gamma'(M, B_1)$ is the vector space of finite sums of such iterated wedge products and $\Gamma'_a(M^k, B_1^{\boxtimes k})$ is the completion of $\Lambda^k \Gamma'(M, B_1)$ with respect to the topology of $\Gamma'(M, B_1)^{\hat{\otimes} \pi^k} \cong \Gamma'(M^k, B_1^{\boxtimes k})$ where all the duals are strong. The subscript ‘‘a’’ stands for antisymmetry.

In the case of multilinear symmetric functions, we can identify a k -linear map $f(h_1, \dots, h_k)$ of k variables with a polynomial map of one variable $f(h, \dots, h)$ by using the polarization identity. There is no polarization identity in the antisymmetric case and we must consider a function $F : E_0 \rightarrow \mathcal{A}^k$ as a function of one variable φ_0 in E_0 and k variables (h_1, \dots, h_k) in E_1 (or a variable in $H \in E_1^{\hat{\otimes} \pi^k}$). Then, we can identify a function $F : E_0 \rightarrow \mathcal{A}^k$ and the function $\tilde{F} : E_0 \times E_1^{\hat{\otimes} \pi^k} \rightarrow \mathbb{K}$ defined by

$$\tilde{F}(\varphi_0; h_1 \otimes \cdots \otimes h_k) = F(\varphi_0)(h_1 \otimes \cdots \otimes h_k).$$

This motivates the following

Definition 8.1. Let M be a smooth manifold, (B_0, B_1) are smooth vector bundles on M and $E_0 = \Gamma(M, B_0)$, $E_1 = \Gamma(M, B_1)$ are spaces of smooth sections of the respective bundles. We say that a function F from E_0 to \mathcal{A}^k is an element of $\mathcal{O}^k(E_0 \oplus E_1[1])$ if there exists a Bastiani smooth map $\tilde{F} : E_0 \times E_1^{\hat{\otimes} \pi^k} \rightarrow \mathbb{K}$ which is linear in $E_1^{\hat{\otimes} \pi^k}$ and **antisymmetric** w.r.t. the natural action of permutations on $E_1^{\hat{\otimes} \pi^k}$ such that :

$$\tilde{F}(\varphi_0; h_1 \otimes \cdots \otimes h_k) = F(\varphi_0)(h_1 \otimes \cdots \otimes h_k). \quad (45)$$

We denote by $\mathcal{O}(E_0 \oplus E_1[1])$ the direct product of all $\mathcal{O}^k(E_0 \oplus E_1[1])$, over $k \in \mathbb{N}_0$ and set $\mathcal{O}^0(E_0 \oplus E_1[1]) \equiv \mathbb{K}$.

Let us now discuss the notion of derivative for the type of functionals introduced above. Clearly, if F belongs to $\mathcal{O}(E_0 \oplus E_1[1])$, there are two natural ways to differentiate it. In the first instance we can differentiate \tilde{F} in the sense of Bastiani in the first variable ($\varphi \in E_0$) and we denote this derivative as

$$D_0 F_{(\varphi; u)}(g) \doteq D\tilde{F}_{(\varphi, u)}(g, 0),$$

where $u \in E_1^{\hat{\otimes} \pi^k} \rightarrow \mathbb{K}$, $g \in E_0$ or $\frac{\delta}{\delta \varphi_0} F$.

8.0.2 The contraction operation.

Let us now consider contraction of the graded part with some $h \in E_1$, sometimes referred to as derivations with respect to odd variables. This concept is needed in order to define the Koszul complex and the Chevalley-Eilenberg complex in the Batalin–Vilkovisky formalism in infinite dimension. The definition is spelled out below.

Definition 8.2. Let $F \in \mathcal{O}^k(E_0 \oplus E_1[1])$, $h \in E_1$. The contraction of F by h is defined, for every integer $k > 0$ and $u \in E_1^{\hat{\otimes} k-1}$, by

$$\begin{aligned} \langle \iota_h F, u \rangle &= \tilde{F}(h \otimes u), \\ \text{and } \iota_h F &= 0 \quad \text{if } F \in \mathcal{A}^0. \end{aligned}$$

In particular, $\iota_h F = \langle \tilde{F}, h \rangle$ if $F \in \mathcal{A}^1$. We extend this definition to \mathcal{A} by linearity.

In view of (45) and the definition of $\mathcal{O}^k(E_0 \oplus E_1[1])$, it is clear that $\iota_h F \in \mathcal{O}^{k-1}(E_0 \oplus E_1[1])$ for all $F \in \mathcal{O}^k(E_0 \oplus E_1[1])$. Equation (45) allows also to make sense of a second important operation on $\mathcal{O}(E_0 \oplus E_1[1])$:

Definition 8.3. The wedge product $\wedge : \mathcal{O}^k(E_0 \oplus E_1[1]) \times \mathcal{O}^{k'}(E_0 \oplus E_1[1]) \rightarrow \mathcal{O}^{k+k'}(E_0 \oplus E_1[1])$ is defined by

$$\left(\widetilde{F \wedge G} \right) (u_1, \dots, u_{k+k'}) = \sum_{\sigma} \text{sgn}(\sigma) \tilde{F}(u_{\sigma(1)}, \dots, u_{\sigma(k)}) \tilde{G}(u_{\sigma(k+1)}, \dots, u_{\sigma(k+k')})$$

(where the sum runs over $k-k'$ shuffles) and extended by linearity on $\mathcal{O}(E_0 \oplus E_1[1]) \times \mathcal{O}(E_0 \oplus E_1[1])$.

Again, in view of (45) and the definition of $\mathcal{O}(E_0 \oplus E_1[1])$, it is clear that the wedge product of an element in $\mathcal{O}^k(E_0 \oplus E_1[1])$ with an element in $\mathcal{O}^{k'}(E_0 \oplus E_1[1])$ is an element in $\mathcal{O}^{k+k'}(E_0 \oplus E_1[1])$. The contraction and wedge product satisfy the following relation on $\mathcal{O}(E_0 \oplus E_1[1])$:

Lemma 8.4. *The contraction satisfies the graded Leibniz rule: if $F \in \mathcal{O}^k(E_0 \oplus E_1[1])$, $G \in \mathcal{O}(E_0 \oplus E_1[1])$ and $h \in E_1$, then*

$$\iota_h(F \wedge G) = (\iota_h F) \wedge G + (-1)^k F \wedge \iota_h G.$$

Let us now discuss the notion of support which is the appropriate generalization of the notion of support for graded functionals, generalizing the definitions in section 3.1.

Definition 8.5. *Let $F \in \mathcal{O}^k(U \oplus E_1[1])$ be a graded functional, with U an open subset of E_0 . The support of F is defined by*

$$\bigcup_{(h_1, \dots, h_k) \in E_1^k} \text{supp}(\varphi \mapsto (\iota_{h_1} \dots \iota_{h_k} F)(\varphi)) \cup \bigcup_{\varphi \in U, (h_1, \dots, h_{k-1}) \in E_1^{k-1}} \text{supp}(h \mapsto (\iota_{h_1} \dots \iota_{h_{k-1}} F(\varphi, h))).$$

8.0.3 Some conjectures on local graded functionals.

Let $F \in \mathcal{O}^k(E_0 \oplus E_1[1])$ be such that the WF set of both $(\iota_{h_1} \dots \iota_{h_k} F)_\varphi^{(1)}$ and $\iota_{h_1} \dots \iota_{h_{k-1}} F(\varphi, \cdot)$ is empty for all $\varphi \in U$ and $(h_1, \dots, h_k) \in E_1^k$. We conjecture that some version of lemma 6.2 and 6.9 should hold in the graded case. The ‘‘standard’’ characterization of locality for a functional $F \in \mathcal{O}^k(E_0 \oplus E_1[1])$ is the requirement that F is compactly supported and for each $(\varphi; u_1, \dots, u_k) \in E_0 \times E_1^k$ there exists $i_0, \dots, i_k \in \mathbb{N}$ such that

$$F(\varphi; u_1, \dots, u_k) = \int_M \alpha(j_x^{i_0}(\varphi), j_x^{i_1}(u_1), \dots, j_x^{i_k}(u_k)), \quad (46)$$

where α is a density-valued function on the jet bundle. To conclude, we **conjecture** some graded analogue of Theorem 6.3 whose formulation would be as follows :

Let U be an open subset of E_0 and $F \in \mathcal{O}^k(U \oplus E_1[1])$ be a graded functional. Assume that

1. F is additive in some suitable sense, still to be written with care (conceivably this would be additivity of \tilde{F} as a function of several variables).
2. $(\iota_{h_1} \dots \iota_{h_k} F)_\varphi^{(1)}$ and $\iota_{h_1} \dots \iota_{h_{k-1}} F(\varphi, \cdot)$ have empty wave front set for all $\varphi \in U$ and $(h_1, \dots, h_k) \in E_1^k$ and the maps $(\varphi, u) \mapsto (\iota_{h_1} \dots \iota_{h_k} F)_\varphi^{(1)}, \iota_{h_1} \dots \iota_{h_{k-1}} F(\varphi, \cdot)$ are Bastiani smooth from $U \times \bigoplus_{k \in \mathbb{N}} E_1^{\hat{\otimes} \pi^k}$ to $\Gamma_c(M, B_0^*)$ and $\Gamma_c(M, B_1^*)$, respectively. Here B_0^* and B_1^* denote dual bundles.

Then, for every $\varphi \in U$, $u \in \bigoplus_{k \in \mathbb{N}} E_1^{\hat{\otimes} \pi^k}$, there is a neighborhood V of the origin in E_0 , an integer N and a smooth \mathbb{K} -valued function f on the N -jet bundle such that

$$F(\varphi + \psi; v_1 \otimes \dots \otimes v_k) = \int_M \alpha(j_x^{i_0}(\psi), j_x^{i_1}(v_1), \dots, j_x^{i_k}(v_k)), \quad (47)$$

for every $\psi \in V$ and some $i_0, \dots, i_k < N$.

9 Acknowledgements

We are very grateful to the Institut Poincaré for making this work possible through the grant and facilities it offered us within the framework of its ‘‘Research in Paris’’ program. We thank the CNRS Groupement de Recherche Renormalisation for its financial support. We are extremely grateful to Yoann Dabrowski for his very generous help during all the phases of this work. We thank Romeo Brunetti and Pedro Ribeiro for very useful discussions.

References

- [1] J. Schwinger. On the Green's functions of quantized fields. I. *Proc. Nat. Acad. Sci.*, 37:452–5, 1951.
- [2] J. Schwinger. On the Green's functions of quantized fields. II. *Proc. Nat. Acad. Sci.*, 37:455–9, 1951.
- [3] W. Kohn and L. J. Sham. Self-consistent equations including exchange and correlation effects. *Phys. Rev.*, 140:A1133–8, 1965.
- [4] E. Engel and R. M. Dreizler. *Density Functional Theory. An Advanced Course*. Springer, Heidelberg, 2011.
- [5] K. Rejzner. *Perturbative Algebraic Quantum Field Theory. An Introduction for Mathematicians*. Springer, Cham, 2016.
- [6] L. Lempert and N. Zhang. Dolbeault cohomology of a loop space. *Acta. Math.*, 193:241–268, 2004.
- [7] L Dickey. *Soliton equations and Hamiltonian systems*, volume 26. World Scientific, 2003.
- [8] J. Lee and R. M. Wald. Local symmetries and constraints. *J. Math. Phys.*, 31:725–43, 1990.
- [9] R. M. Wald. *General Relativity*. The University of Chicago Press, Chicago, 1984.
- [10] J. D. Stasheff. The (secret?) homological algebra of the Batalin-Vilkovisky approach. *Contemp. Math.*, 219:195–210, 1998.
- [11] G. Barnich, F. Brandt, and M. Henneaux. Local BRST cohomology in gauge theories. *Phys. Repts*, 338:439–569, 2000.
- [12] R. Brunetti, K. Fredenhagen, and K. Rejzner. Quantum gravity from the point of view of locally covariant quantum field theory. *Commun. Math. Phys.*, 345:741–79, 2016.
- [13] I. Khavkine. Local and gauge invariant observables in gravity. *Class. Quant. Grav.*, 32:185019, 2015.
- [14] K. Fredenhagen and K. Rejzner. Batalin-Vilkovisky formalism in the functional approach to classical field theory. *Commun. Math. Phys.*, 314:93–127, 2012.
- [15] R. Brunetti, M. Dütsch, and K. Fredenhagen. Perturbative algebraic quantum field theory and the renormalization groups. *Adv. Theor. Math. Phys.*, 13:1541–99, 2009.
- [16] R. Brunetti and K. Fredenhagen. Quantum field theory on curved backgrounds. In C. Bär and K. Fredenhagen, editors, *Quantum Field Theory on Curved Spacetimes: Concepts and Mathematical Foundations*, volume 786 of *Lecture Notes in Physics*, pages 129–155, Berlin, 2009. Springer.
- [17] R. Brunetti, K. Fredenhagen, and P. L. Ribeiro. Algebraic structure of classical field theory. I. Kinematics and linearized dynamics for real scalar fields. 2012. arXiv:1209.2148.
- [18] Y. Dabrowski. Functional properties of generalized Hörmander spaces of distributions I: Duality theory, completions and bornologifications. 2014. arXiv:1411.3012.
- [19] Y. Dabrowski. Functional properties of generalized Hörmander spaces of distributions II: Multilinear maps and applications to spaces of functionals with wave front set conditions. 2014. arXiv:1412.1749.
- [20] H. Epstein and V. Glaser. The role of locality in perturbation theory. *Ann. Inst. Henri Poincaré*, 19:211–95, 1973.

- [21] A. S. Wightman. How it was learned that quantized fields are operator-valued distributions. *Fortschr. Phys.*, 44:143–78, 1996.
- [22] M. Martellini. Quantum field theory on curved space-time: An axiomatic approach. *Nuovo Cimento A*, 67:305–55, 1982.
- [23] M. Dütsch and K. Fredenhagen. Algebraic quantum field theory, perturbation theory and the loop expansion. *Commun. Math. Phys.*, 219:5–30, 2001.
- [24] L. Schwartz. *Théorie des distributions*. Hermann, Paris, second edition, 1966.
- [25] A. Kriegl and P. W. Michor. *The Convenient Setting of Global Analysis*. American Mathematical Society, Providence, 1997.
- [26] E. C. G. Stueckelberg. Relativistic quantum theory for finite time intervals. *Phys. Rev.*, 81:130–3, 1951.
- [27] J. Horváth. *Topological Vector Spaces and Distributions*. Addison-Wesley, Reading, 1966.
- [28] F. Trèves. *Topological Vector Spaces, Distributions and Kernels*. Dover, New York, 2007.
- [29] L. Hörmander. *The Analysis of Linear Partial Differential Operators I. Distribution Theory and Fourier Analysis*. Springer Verlag, Berlin, second edition, 1990.
- [30] A. Shapiro. On concepts of directional differentiability. *J. Optim. Theo. Appl.*, 66:477–87, 1990.
- [31] G. A. Ladas and V. Laskhmiathanam. *Differential Equations in Abstract Spaces*. Academic Press, New York, 1972.
- [32] A. J. Kurdila and M. Zabaranin. *Convex Functional Analysis*. Birkhäuser, Basel, 2005.
- [33] H. Sagan. *Introduction to the Calculus of Variations*. Dover, New York, 1992.
- [34] S. Yamamuro. *Differential Calculus in Topological Linear Spaces*, volume 374 of *Lecture Notes in Mathematics*. Springer, Berlin, 1974.
- [35] A. Bastiani. Applications différentiables et variétés différentiables de dimension infinie. *J. Anal. Math.*, 13:1–114, 1964.
- [36] R. S. Hamilton. The inverse function theorem of Nash and Moser. *Bull. Am. Math. Soc.*, 7:65–222, 1982.
- [37] A. Bastiani. *Différentiabilité dans les espaces localement convexes. Distructures*. Université de Paris, 1962. Ph.D. thesis.
- [38] K.-H. Neeb. Infinite-dimensional groups and their representations. In A. Huckleberry and T. Wurzbacher, editors, *Infinite Dimensional Kähler Manifolds*, pages 131–78, Basel, 2001. Birkhäuser.
- [39] J.-P. Penot. Calcul différentiel sur les espaces vectoriels topologiques. *Studia Math.*, 47(1):1–23, 1973.
- [40] H. H. Keller. *Differential Calculus in Locally Convex Spaces*, volume 417 of *Lecture Notes in Mathematics*. Springer, Berlin, 1974.
- [41] V. I. Averbukh and O. G. Smolyanov. The various definitions of the derivative in linear topological spaces. *Russ. Math. Surv.*, 23(4):67–113, 1968.
- [42] W. Gähler. *Grundstrukturen der Analysis*, volume 2. Birkhäuser, Basel, 1978.

- [43] P. ver Eecke. *Fondements du Calcul Différentiel*. Presses Universitaires de France, Paris, 1983.
- [44] M. Z. Nashed. Differentiability and related properties of nonlinear operators: some aspects of the role of differentials in nonlinear functional analysis. In L. B. Rall, editor, *Nonlinear Functional Analysis and Applications*, pages 103–359, New York, 1971. Academic Press.
- [45] M. C. Abbati and A. Manià. On differential structure for projective limits of manifolds. *J. Geom. Phys.*, 29:35–63, 1999.
- [46] P. W. Michor. *Manifolds of Differentiable Mappings*. Shiva Publishing Ltd, Orpington, 1980.
- [47] J. Milnor. Remarks on infinite-dimensional Lie groups. In B. S. DeWitt and R. Stora, editors, *Relativity, Groups, and Topology II*, volume 40 of *Les Houches*, pages 1007–57, Amsterdam, 1983. North Holland.
- [48] B. Khesin and R. Wendt. *The Geometry of Infinite-Dimensional Groups*. Springer, Berlin, 2009.
- [49] W. Bertram. *Differential Geometry, Lie Groups and Symmetric Spaces over General Base Fields and Rings*, volume 192 of *Mem. Amer. Math. Soc.* American Mathematical Society, Providence, 2008.
- [50] J. Szilasi and R. L. Lovas. Some aspects of differential theories. In D. Krupka and D. Saunders, editors, *Handbook of Global Analysis*, pages 1069–114, Oxford, 2008. Oxford University Press.
- [51] R. Dahmen. Peetre’s theorem in the locally convex setting. *Forum Math.*, 27:2957–79, 2015.
- [52] A. D. Michal. Differential calculus in linear topological spaces. *Proc. Nat. Acad. Sci.*, 24:340–2, 1938.
- [53] H. Massam and S. Zlobec. Various definitions of the derivative in mathematical programming. *Math. Program.*, 7:144–61, 1974.
- [54] A. Ehresmann. Des espèces de structures locales aux distructures et systèmes guidables. In *Category Theory Yesterday, Today (and Tomorrow?): A colloquium in Honour of Jean Benabou*, ENS, Paris, 2011.
- [55] J. L. Kelley. *General Topology*, volume 27 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, revised third edition, 1955.
- [56] N. Bourbaki. *Elements of Mathematics. General Topology. Part 2*. Addison-Wesley, Reading, 1966.
- [57] N. Bourbaki. *Elements of Mathematics. Topological Vector Spaces*. Springer, Berlin, 2003.
- [58] M. Reed and B. Simon. *Methods of Modern Mathematical Physics. I Functional Analysis*. Academic Press, New York, second edition, 1980.
- [59] K.-H. Neeb. On analytic vectors for unitary representations of infinite dimensional Lie groups. *Ann. Inst. Fourier*, 61:1839–74, 2011.
- [60] N. Bourbaki. *Éléments de Mathématique. Intégration. Chapitres 1 à 4*. Hermann, Paris, second edition, 1965.
- [61] N. Bourbaki. *Éléments de Mathématique. Intégration. Chapitre 6. Intégration vectorielle*. Hermann, Paris, 1959.
- [62] H. Glöckner. Infinite-dimensional Lie groups without completeness restrictions. In *Orlicz Centenary Volume*, volume 55 of *Banach Center Publ.*, pages 43–59, Warsaw, 2002. Polish Acad. Sci.

- [63] K.-H. Neeb. *Monastir Summer School: Infinite-dimensional Lie groups*. TU Darmstadt Preprint 2433, Darmstadt, 2006.
- [64] H. Glöckner. *Infinite-dimensional Lie Groups*. Lecture Notes. TU Darmstadt, Darmstadt, 2005.
- [65] N. N. Tarkhanov. *The Analysis of Solutions of Elliptic Equations*. Kluwer Academic Publishers, Dordrecht, 1997.
- [66] M. Grosser. A note on distribution spaces on manifolds. *Novi Sad J. Math*, 38:121–8, 2008.
- [67] H. Glöckner. Lie group structures on quotient groups and universal complexifications for infinite-dimensional Lie groups. *J. Funct. Anal.*, 194:347–409, 2002.
- [68] C. Wockel. *Central extensions of gauge groups*. Diplomarbeit, Technische Universität Darmstadt, 2003.
- [69] C. Wockel. *Infinite-dimensional Lie theory for gauge groups*. PhD Thesis, Technische Universität Darmstadt, 2006.
- [70] Ch. Brouder, N. V. Dang, and F. Hélein. A smooth introduction to the wavefront set. *J. Phys. A: Math. Theor.*, 47:443001, 2014.
- [71] J.-F. Colombeau and R. Meise. C^∞ -functions on locally convex and on bornological vector spaces. In *Functional Analysis, Holomorphy, and Approximation Theory*, volume 843 of *Lecture Notes in Mathematics*, pages 195–216. Springer, Berlin, 1981.
- [72] K. Fredenhagen and K. Rejzner. Batalin-Vilkovisky formalism in perturbative algebraic quantum field theory. *Commun. Math. Phys.*, 317:697–725, 2013.
- [73] K. Fredenhagen and K. Rejzner. Perturbative algebraic quantum field theory. *Lecture Notes*, 2012. arXiv:1208.1428.
- [74] C. Bär and R. Tagne Wafo. Initial value problems for wave equations on manifolds. *Math. Phys. Anal. Geom.*, 18:7, 2015.
- [75] T. Hirai, H. Shimomura, N. Tatsuuma, and E. Hirai. Inductive limit of topologies, their direct products, and problems related to algebraic structures. *J. Math. Kyoto Univ.*, 41:475–505, 2001.
- [76] Y. Dabrowski and Ch. Brouder. Functional properties of Hörmander’s space of distributions having a specified wavefront set. *Commun. Math. Phys.*, 332:1345–80, 2014.
- [77] S. Hollands and R. M. Wald. Local Wick polynomials and time ordered products of quantum fields in curved spacetime. *Commun. Math. Phys.*, 223:289–326, 2001.
- [78] K. Rejzner. *Batalin-Vilkovisky formalism in locally covariant field theory*. Ph.D. thesis, University of Hamburg, 2011.
- [79] M. M. Rao. Local functionals. In *Measure Theory Oberwolfach 1979*, volume 794 of *Lecture Notes in Mathematics*, pages 484–96. Springer, Berlin, 1980.
- [80] A. G. Pinsker. La représentation analytique de quelques fonctionnelles partiellement additives. *Comptes Rendus (Doklady) Acad. Sci. URSS*, 18:399–403, 1938.
- [81] I. M. Gel’fand and N. Ya. Vilenkin. *Generalized Functions: Volume IV - Applications of Harmonic Analysis*. Academic Press, New York, 1964.
- [82] M. Dütsch and K. Fredenhagen. Causal perturbation theory in terms of retarded products, and a proof of the Action Ward Identity. *Rev. Math. Phys.*, 16:1291–348, 2004.

- [83] K. J. Keller. Euclidean Epstein-Glaser renormalization. *J. Math. Phys.*, 50:103503, 2009.
- [84] R. V. Chacon and N. Friedman. Additive functionals. *Arch. Rational Mech. Anal.*, 18:230–40, 1965.
- [85] J. Batt. Nonlinear integral operators on $C(S, E)$. *Studia Math.*, 48:146–77, 1973.
- [86] S. H. Fesmiré. Representation of nonlinear operators with the Hammerstein property. In *Functional Analysis and its Applications*, volume 399 of *Lecture Notes in Mathematics*, pages 129–48. Springer, Berlin, 1974.
- [87] A. de Korvin and C. E. Roberts Jr. Convergence theorems for some scalar valued integrals when the measure is Nemytskii. *Pacific J. Math.*, 92:329–43, 1981.
- [88] K. Lermer. The Grothendieck-Pietsch domination principle for nonlinear summing integral operators. *Studia Math.*, 129:97–112, 1998.
- [89] Z. Ercan. A generalisation of a theorem of Koldunov with an elementary proof. *Czech. Math. J.*, 49:187–90, 1999.
- [90] H. G. R. Millington. Riesz integral representation theory. 2007. arXiv:0706.4281.
- [91] J. A. Navarro. Differential spaces of finite type. *Math. Proc. Cambridge Phil. Soc.*, 117:371–384, 1995.
- [92] R. Moosa. Jet spaces in complex analytic geometry: An exposition. *arXiv preprint math/0405563*, 2004.
- [93] J.-M. Kantor. Formes et opérateurs différentiels sur les espaces analytiques complexes. *Mém. Soc. Math. France*, 53:5–80, 1977.
- [94] N. V. Dang. The extension of distributions on manifolds, a microlocal approach. *Ann. Henri Poincaré*, 17:819–59, 2016.
- [95] Ch. Brouder, N. V. Dang, and F. Hélein. Continuity of the fundamental operations on distributions having a specified wavefront set (with a counterexample by Semyon Alesker). *Studia Math.*, 232:201–26, 2016. arXiv:1409.7662.
- [96] J. M. Lee. *Introduction to Smooth Manifolds*. Springer, New York, 2003.
- [97] J. Navarro and J. B. Sancho. Peetre-Slovák’s theorem revisited. 2014. arXiv:1411.7499.
- [98] J.-P. Brasselet and M. J. Pflaum. On the homology of algebras of Whitney functions over subanalytic sets. *Annals of mathematics*, pages 1–52, 2008.
- [99] J. Slovák. Peetre theorem for nonlinear operators. *Ann. Glob. Anal. Geom.*, 6:273–283, 1988.
- [100] K. Rejzner. Fermionic fields in the functional approach to classical field theory. *Rev. Math. Phys.*, 23:1009–33, 2011.