

Ramsey properties and extending partial automorphisms for classes of finite structures

David M. Evans*, Jan Hubička†, Jaroslav Nešetřil‡

Abstract

We show that every free amalgamation class of finite structures with relations and (set-valued) functions is a Ramsey class when enriched by a free linear ordering of vertices. This is a common strengthening of the Nešetřil-Rödl Theorem and the second and third authors' Ramsey theorem for finite models (that is, structures with both relations and functions). We also find subclasses with the ordering property. For languages with relational symbols and unary functions we also show the extension property for partial automorphisms (EPPA) of free amalgamation classes. These general results solve several conjectures and provide an easy Ramsey-ness test for many classes of structures.

1 Introduction

In this paper we discuss three related concepts — Ramsey classes, the ordering (or lift or expansion) property and the extension property for partial automorphisms (EPPA). The main novelty of our results is that they hold for free amalgamation classes of finite structures with both relations and *set-valued functions*. This provides a useful tool for proving these types of results for some classes of structures which naturally carry a closure operation. We will explain below what we mean by this; examples and applications are given at the end of the paper.

As is well known, all three of these concepts about classes of finite structures are related to issues in topological dynamics and this relationship provides much of the motivation for what we do. For example, by [22] the automorphism group of the Fraïssé limit of a Ramsey class \mathcal{R} is extremely amenable. Moreover, if the Ramsey class \mathcal{R} has the ordering property with respect to some amalgamation class \mathcal{K} , then it determines the universal minimal flow of the automorphism group of the Fraïssé limit of \mathcal{K} . Thus, our results Theorem 1.3 and Theorem 1.4

*Department of Mathematics, Imperial College London, London SW7 2AZ, UK, david.evans@imperial.ac.uk

†Supported by project 18-13685Y of the Czech Science Foundation (GAČR), Center for Foundations of Modern Computer Science (Charles University project UNCE/SCI/004), by the PRIMUS/17/SCI/3 project of Charles University and by ERC Synergy grant DYNASNET. Department of Applied Mathematics (KAM), Charles University, Malostranské nám. 25, 11800 Praha, Czech Republic, 118 00 Praha 1, hubicka@kam.mff.cuni.cz

‡Supported ERC Synergy grant DYNASNET and by European Associated Laboratory (LEA STRUCO). Computer Science Institute of Charles University (IUUK), Charles University, Malostranské nám. 25, 11800 Praha, Czech Republic, 118 00 Praha 1, nesetřil@iuuk.mff.cuni.cz

about the Ramsey and ordering properties give new examples of this correspondence. By [23] and our Theorem 1.7 about EPPA, the automorphism group of the Fraïssé limit of every free amalgamation class \mathcal{K} in a language where all functions are unary is amenable. However we note that the same conclusion, without the restriction that the (set-valued) functions are unary, also follows from our Ramsey Theorem 1.3 and Proposition 9.3 of [2] (note that the assumption of the ordering property in the statement of Proposition 9.3 of [2] is not necessary, as was observed by Pawliuk and Sokić in [33]).

To generalise naturally the known results about relational structures, we need to define carefully what we mean by a structure and substructure, because none of these results holds in the context of free amalgamation classes with strong embeddings (as discussed in [11], see also the remarks preceding Theorem 1.3). In the Ramsey theory setting it is common to work with ‘incomplete’ structures. Thus we have to modify the standard model-theoretic notion of structures (see e.g. [15]), where functions are required to be total and thus complete in some sense. Before stating the main results, we give the basic (model-theoretic) setting of this paper. We find it useful to introduce a variant of the usual model-theoretic structures, allowing set-valued functions, which is well tailored both to the Ramsey and EPPA setting.

Let $L = L_{\mathcal{R}} \cup L_{\mathcal{F}}$ be a language with relational symbols $R \in L_{\mathcal{R}}$ and function symbols $F \in L_{\mathcal{F}}$ each having its arity denoted by $\text{arity}(R)$ for relations and $\text{arity}(F)$ for functions. Denote by $\mathcal{P}(A)$ the set of all subsets of A . An L -structure \mathbf{A} is a structure with vertex set A , functions $F_{\mathbf{A}}: A^{\text{arity}(F)} \rightarrow \mathcal{P}(A)$, $F \in L_{\mathcal{F}}$ and relations $R_{\mathbf{A}} \subseteq A^{\text{arity}(R)}$, $R \in L_{\mathcal{R}}$. Set-valued functions permits an explicit description of algebraic closures in the Fraïssé limits without changing the automorphism group which is necessary for some applications discussed in Section 5. Because the image of a tuple can be empty set it also gives a natural meaning to the notion of free amalgamation for structures in languages containing functions of arity greater or equal to 2 and it simplifies some of the notation below.

The language is usually fixed and understood from the context (and it is in most cases denoted by L). If the set A is finite we call \mathbf{A} a *finite structure*. We consider only structures with finitely or countably infinitely many vertices. If the language L contains no function symbols, we call L a *relational language* and say that an L -structure is a *relational L -structure*. A function symbol F such that $\text{arity}(F) = 1$ is a *unary function*.

The notions of embedding, isomorphism, homomorphisms and free amalgamation classes are natural generalisations of the corresponding notions on relational structures and are formally introduced in Section 1.4. Considering function symbols has important consequences for what we consider a substructure. An L -structure \mathbf{A} is a *substructure* of \mathbf{B} if $A \subseteq B$ and all relations and functions of \mathbf{B} restricted to A are precisely relations and functions of \mathbf{A} . In particular for every $F \in L$ and every tuple $(v_1, v_2, \dots, v_{\text{arity}(F)})$ of vertices of A it also holds that $F_{\mathbf{B}}(v_1, v_2, \dots, v_{\text{arity}(F)}) \subseteq A$. This implies that \mathbf{B} does not induce a substructure on every subset of \mathbf{B} (but only on ‘closed’ sets, to be defined later).

Building on these model theoretic notions we now outline the contents of this paper. We proceed to the three main directions—Ramsey theory, the ordering property and the extension property for partial automorphisms (EPPA). In each of these directions we now state the main result (stated below as Theorems 1.3,

1.4 and 1.7). This can be summarised by saying that for free amalgamation classes we have strong positive theorems in each of these areas. There is more to it than just meets the eye: for the first time we demonstrate affinity of all these directions (for the ordering property and Ramsey this is known, but for EPPA much less so).

This has a number of applications to special classes of structures. In Section 5 we give several examples which have received recent attention: k -orientations, bowtie-free graphs and Steiner systems. These all are easy consequences of our main result. Finally, let us remark that our results further narrow the gap for the project of characterisation of Ramsey classes [25].

1.1 Ramsey classes

For structures \mathbf{A}, \mathbf{B} denote by $\binom{\mathbf{B}}{\mathbf{A}}$ the set of all sub-structures of \mathbf{B} , which are isomorphic to \mathbf{A} . Using this notation the definition of a Ramsey class has the following form. A class \mathcal{C} is a *Ramsey class* if for every two objects \mathbf{A} and \mathbf{B} in \mathcal{C} and for every positive integer k there exists a structure \mathbf{C} in \mathcal{C} such that the following holds: For every partition $\binom{\mathbf{C}}{\mathbf{A}}$ into k classes there exists a $\tilde{\mathbf{B}} \in \binom{\mathbf{C}}{\mathbf{B}}$ such that $\binom{\tilde{\mathbf{B}}}{\mathbf{A}}$ belongs to one class of the partition. It is usual to shorten the last part of the definition to $\mathbf{C} \rightarrow (\mathbf{B})_k^{\mathbf{A}}$.

We are motivated by the following, now classical, result.

Theorem 1.1 (Nešetřil-Rödl Theorem [27, 31]). *Let L be a relational language, (\mathbf{A}, \leq) and (\mathbf{B}, \leq) be ordered L -structures. Then there exists an ordered L -structure (\mathbf{C}, \leq) such that $(\mathbf{C}, \leq) \rightarrow (\mathbf{B}, \leq)_2^{(\mathbf{A}, \leq)}$.*

Moreover, if \mathbf{F} is an irreducible L -structure (see Section 2 for the definition of irreducibility) and \mathbf{A} and \mathbf{B} do not contain \mathbf{F} as a substructure, then \mathbf{C} may be chosen with the same property.

In our setting this result may be reformulated as follows: Given a language L , denote by \vec{L} the language L extended by one binary relation \leq . Given an L -structure \mathbf{A} , an *ordering of \mathbf{A}* is an \vec{L} -structure extending \mathbf{A} by an arbitrary linear ordering $\leq_{\mathbf{A}}$ of the vertices. For brevity we denote such ordered \mathbf{A} as $\vec{\mathbf{A}}$. Given a class \mathcal{K} of L -structures, denote by $\vec{\mathcal{K}}$ the class of all orderings of structures in \mathcal{K} . We sometimes say that $\vec{\mathcal{K}}$ arises by taking *free orderings* of structures in \mathcal{K} . Note that minimal relational structures which do not belong to a free amalgamation class are all irreducible. Thus Theorem 1.1 can now be re-formulated using basic notions of Fraïssé theory (which will be introduced in Section 1.4) as follows:

Theorem 1.2 (Nešetřil-Rödl Theorem for free amalgamation classes). *Let L be a relational language and \mathcal{K} be a free amalgamation class of relational L -structures. Then $\vec{\mathcal{K}}$ is a Ramsey class.*

The more recent connection between Ramsey classes and extremely amenable groups [22] has motivated a systematic search for new examples of Ramsey classes. It became apparent that it is important to consider structures with both relations and functions or, equivalently, classes of structures with “strong embeddings”. This led to [21] which provides a sufficient structural condition for a subclass of a Ramsey class of structures to be Ramsey and also generalises this approach to classes of structures (containing relations and functions) with

formally-described closures. It is however clear that considering classes with closures (especially non-unary closures) leads to many technical difficulties. In fact, a recent example given by first author based on Hrushovski's predimension construction [11] not only answers one of the main questions in the area (about the existence of precompact Ramsey expansions), but also shows that there is no direct generalisation of the Nešetřil-Rödl Theorem to a free amalgamation classes with strong embeddings. However, perhaps surprisingly, we show that if closures can be explicitly represented by means of set-valued functions such that strong embeddings are just embeddings, such a statement is true. We prove:

Theorem 1.3. *Let L be a language (involving relational symbols and set-valued functions) and let \mathcal{K} be a free amalgamation class of L -structures. Then $\vec{\mathcal{K}}$ is a Ramsey class.*

This yields an alternative proof of the Ramsey property for some recently-discovered Ramsey classes (such as ordered partial Steiner systems [4], bowtie-free graphs [19], bouquet-free graphs [8]) and also for new classes: most importantly a Ramsey expansion of the class of 2-orientations of a Hrushovski predimension construction which is elaborated in [11] and which was one of the main motivations for this paper.

1.2 Ordering property

A class $\mathcal{O} \subseteq \vec{\mathcal{K}}$ has the *ordering property (with respect to \mathcal{K})* if for every $\mathbf{A} \in \mathcal{K}$ there exists $\mathbf{B} \in \mathcal{K}$ such that every ordering $\vec{B} \in \mathcal{O}$ of \mathbf{B} contains a copy of every ordering $\vec{A} \in \mathcal{O}$ of \mathbf{A} . It is well known that for every free amalgamation class \mathcal{K} of relational structures the class $\vec{\mathcal{K}}$ has ordering property. This fact follows by an application of Theorem 1.2 but can also be shown by more general methods based on hypergraphs of large girth [30, 28]. This shows that there are many classes \mathcal{K} of relational structures for which $\vec{\mathcal{K}}$ has the ordering property (with respect to \mathcal{K}) but \mathcal{K} itself is not a Ramsey class.

For languages containing function symbols, the situation is more complicated. To see that some extra restriction on our class \mathcal{O} is required, we note the following example. Denote by \mathcal{T} the class of all finite forests represented by a single unary function F connecting a vertex to its father. Let \mathbf{A} be a structure containing two vertices a, b and $F_{\mathbf{A}}(a) = \{b\}$. A vertex c is a *root* if $F_{\mathbf{A}}(c) = \emptyset$. Any structure \mathbf{B} can be ordered in increasing order according to the distance from a root vertex. It follows that such an ordering never contains the ordering of \mathbf{A} given by $a \leq_{\mathbf{A}} b$ and consequently $\vec{\mathcal{T}}$ does not have the ordering property.

Nevertheless, we show the following:

Theorem 1.4. *Let L be a language (involving relational symbols and set-valued functions) and let \mathcal{K} be a free amalgamation class of L -structures. Then there exists an explicitly-defined amalgamation class $\mathcal{O} \subseteq \vec{\mathcal{K}}$ of admissible orderings such that:*

1. every $\mathbf{A} \in \mathcal{K}$ has an ordering in \mathcal{O} ;
2. \mathcal{O} is a Ramsey class; and,
3. \mathcal{O} has the ordering property (with respect to \mathcal{K}).

The details of the admissible orderings are technical and are described in full in Definition 3.2. The existence of a subclass $\mathcal{O} \subseteq \vec{\mathcal{K}}$ with the above three properties follows directly from Theorem 10.7 of [22] and Theorem 1.3. The novelty in Theorem 1.4 is the explicit description of a class \mathcal{O} . The proof of Theorem 1.4 is a combination of the Ramsey methods used to show the ordering property of classes of relational structures and the methods used to show the ordering property of classes with unary functions.

1.3 Extension property for partial automorphisms – EPPA

A *partial automorphism* of an L -structure \mathbf{A} is an isomorphism $f : \mathbf{D} \rightarrow \mathbf{E}$ of substructures \mathbf{D}, \mathbf{E} of \mathbf{A} . We say that a class of finite L -structures \mathcal{K} has the *extension property for partial automorphisms* (*EPPA*, sometimes called the *Hrushovski extension property*) if whenever $\mathbf{A} \in \mathcal{K}$ there is $\mathbf{B} \in \mathcal{K}$ such that \mathbf{A} is substructure of \mathbf{B} and every partial automorphism of \mathbf{A} extends to an automorphism of \mathbf{B} , see [18, 13, 14, 17, 36, 38]. In the following we will simply call \mathbf{B} with the property above an *EPPA-witness* of \mathbf{A} .

For relational languages, the extension property for partial automorphisms of free amalgamation classes can be derived from the following strengthening of the extension property for partial automorphisms:

Theorem 1.5 (Hodkinson-Otto [17]). *Let L be a relational language, then for every finite L -structure \mathbf{A} there exists a finite and clique faithful EPPA-witness \mathbf{B} .*

A *clique faithful EPPA-witness* \mathbf{B} is an EPPA-witness of \mathbf{A} with the additional property that for every clique \mathbf{C} in the Gaifman graph of \mathbf{B} there exists an automorphism g of \mathbf{B} such that $g(C) \subseteq A$. (For a relational L -structure \mathbf{A} the *Gaifman graph* is the graph $\mathbf{G}_{\mathbf{A}}$ with vertices A and edges those pairs of vertices contained in a tuple of a relation of \mathbf{A} : $\mathbf{G}_{\mathbf{A}} = (V, E)$ where $\{x, y\} \in E$ if and only if $x \neq y$ and there exists tuple $\vec{t} \in R_{\mathbf{A}}, R \in L$ such that $x, y \in \vec{t}$.)

It is a well known fact that free amalgamation classes can be equivalently described by forbidden embeddings from a family of structures whose Gaifman graph is a clique and consequently Theorem 1.5 implies that every free amalgamation class of relational structures has EPPA (see, for example, [35]).

The notion of *irreducibility* of a structure (given in Definition 2.1) is a natural generalisation to the context of functional languages of the above notion of a clique in a graph. We say that an EPPA-witness \mathbf{B} of \mathbf{A} is *irreducible substructure faithful* if for every irreducible substructure \mathbf{C} of \mathbf{B} there exists an automorphism g of \mathbf{B} such that $g(C) \subseteq A$.

Theorem 1.5 was further strengthened by Siniora and Solecki in the following form.

Theorem 1.6 (Siniora-Solecki [35]). *Let L be relational language. Then for every finite relational L -structure \mathbf{A} there exists a finite clique faithful and coherent EPPA-witness \mathbf{B} .*

Let X be a set and \mathcal{P} be a family of partial bijections between subsets of X . A triple (f, g, h) from \mathcal{P} is called a *coherent triple* if $\text{Dom}(f) = \text{Dom}(h)$, $\text{Range}(f) = \text{Dom}(g)$, $\text{Range}(g) = \text{Range}(h)$ and $h = g \circ f$.

Let X and Y be sets, and \mathcal{P} and \mathcal{Q} be families of partial bijections between subsets of X and between subsets of Y , respectively. A function $\varphi : \mathcal{P} \rightarrow \mathcal{Q}$ is

said to be a *coherent map* if for each coherent triple (f, g, h) from \mathcal{P} , its image $\varphi(f), \varphi(g), \varphi(h)$ in \mathcal{Q} is coherent.

An EPPA-witness \mathbf{B} of \mathbf{A} is *coherent* if every partial automorphism f extends to some $\hat{f} \in \text{Aut}(\mathbf{B})$ with the property that the map φ from partial automorphisms of \mathbf{A} to automorphisms of \mathbf{B} given by $\varphi(f) = \hat{f}$ is coherent.

Our third main result is a strengthening of all of the above results to classes of structures with unary functions. We do not know how to strengthen this theorem for higher arities.

Theorem 1.7. *Let L be a language such that every function symbol $F \in L$ is unary. Then for every finite L -structure \mathbf{A} there exists a finite, irreducible substructure faithful, coherent EPPA-witness \mathbf{B} . Consequently every free amalgamation class \mathcal{K} of finite L -structures has the coherent extension property for partial automorphisms.*

1.4 Further background and notation

We now review some standard graph-theoretic and model-theoretic notions (see e.g. [15]).

A *homomorphism* $f : \mathbf{A} \rightarrow \mathbf{B}$ is a mapping $f : A \rightarrow B$ such that:

- (a) for every relation symbol $R \in L_{\mathcal{R}}$ of arity a it holds:

$$(x_1, x_2, \dots, x_a) \in R_{\mathbf{A}} \implies (f(x_1), f(x_2), \dots, f(x_a)) \in R_{\mathbf{B}},$$

- (b) for every function symbol $F \in L_{\mathcal{F}}$ of arity a it holds:

$$f(F_{\mathbf{A}}(x_1, x_2, \dots, x_a)) \subseteq F_{\mathbf{B}}(f(x_1), f(x_2), \dots, f(x_a)).$$

Here, for a subset $A' \subseteq A$ we denote by $f(A')$ the set $\{f(x) : x \in A'\}$ and by $f(\mathbf{A})$ the homomorphic image of a structure \mathbf{A} .

If f is injective, then f is called a *monomorphism*. A monomorphism f is an *embedding* if it holds:

- (a) for every relation symbol $R \in L_{\mathcal{R}}$ of arity a it holds:

$$(x_1, x_2, \dots, x_a) \in R_{\mathbf{A}} \iff (f(x_1), f(x_2), \dots, f(x_a)) \in R_{\mathbf{B}},$$

- (b) for every function symbol $F \in L_{\mathcal{F}}$ of arity a it holds:

$$f(F_{\mathbf{A}}(x_1, x_2, \dots, x_a)) = F_{\mathbf{B}}(f(x_1), f(x_2), \dots, f(x_a)).$$

If f is an embedding which is an inclusion then \mathbf{A} is a *substructure* (or *subobject*) of \mathbf{B} . For an embedding $f : \mathbf{A} \rightarrow \mathbf{B}$ we say that \mathbf{A} is *isomorphic* to $f(\mathbf{A})$ and $f(\mathbf{A})$ is also called a *copy* of \mathbf{A} in \mathbf{B} . Thus $\binom{\mathbf{B}}{\mathbf{A}}$ is defined as the set of all copies of \mathbf{A} in \mathbf{B} .

Given $\mathbf{A} \in \mathcal{K}$ and $B \subset A$, the *closure of B in \mathbf{A}* , denoted by $\text{Cl}_{\mathbf{A}}(B)$, is the smallest substructure of \mathbf{A} containing B . Closure in \mathbf{A} is *unary* if $\text{Cl}_{\mathbf{A}}(B) = \bigcup_{v \in B} \text{Cl}_{\mathbf{A}}(v)$ for all $B \subset A$.

Let \mathbf{A} , \mathbf{B}_1 and \mathbf{B}_2 be structures, α_1 an embedding of \mathbf{A} into \mathbf{B}_1 and α_2 an embedding of \mathbf{A} into \mathbf{B}_2 . Then every structure \mathbf{C} with embeddings $\beta_1 : \mathbf{B}_1 \rightarrow \mathbf{C}$




Figure 1: An amalgamation of \mathbf{B}_1 and \mathbf{B}_2 over \mathbf{A} .

and $\beta_2 : \mathbf{B}_2 \rightarrow \mathbf{C}$ such that $\beta_1 \circ \alpha_1 = \beta_2 \circ \alpha_2$ is called an *amalgamation* of \mathbf{B}_1 and \mathbf{B}_2 over \mathbf{A} with respect to α_1 and α_2 . See Figure 1. We will call \mathbf{C} simply an *amalgamation* of \mathbf{B}_1 and \mathbf{B}_2 over \mathbf{A} (as in most cases α_1 and α_2 can be chosen to be inclusion embeddings). We say that such an amalgamation is *free* if $\beta_1(x_1) = \beta_2(x_2)$ if and only if $x_1 \in \alpha_1(A)$ and $x_2 \in \alpha_2(A)$, $C = \beta_1(B_1) \cup \beta_2(B_2)$ and whenever a tuple \vec{t} of vertices of \mathbf{C} contains vertices of both $\beta_1(B_1 \setminus \alpha_1(A))$ and $\beta_2(B_2 \setminus \alpha_2(A))$, then \vec{t} is in no relation of \mathbf{C} , and for every function $F \in L$ with $\text{arity}(F) = |\vec{t}|$ it holds that $F_{\mathbf{C}}(\vec{t}) = \emptyset$.

Definition 1.1. An *amalgamation class* is a class \mathcal{K} of finite structures satisfying the following three conditions:

1. *Hereditary property:* For every $\mathbf{A} \in \mathcal{K}$ and a substructure \mathbf{B} of \mathbf{A} we have $\mathbf{B} \in \mathcal{K}$;
2. *Joint embedding property:* For every $\mathbf{A}, \mathbf{B} \in \mathcal{K}$ there exists $\mathbf{C} \in \mathcal{K}$ such that \mathbf{C} contains both \mathbf{A} and \mathbf{B} as substructures;
3. *Amalgamation property:* For $\mathbf{A}, \mathbf{B}_1, \mathbf{B}_2 \in \mathcal{K}$ and α_1 embedding of \mathbf{A} into \mathbf{B}_1 , α_2 embedding of \mathbf{A} into \mathbf{B}_2 , there is $\mathbf{C} \in \mathcal{K}$ which is an amalgamation of \mathbf{B}_1 and \mathbf{B}_2 over \mathbf{A} with respect to α_1 and α_2 .

If the \mathbf{C} in the amalgamation property can always be chosen as the free amalgamation, then \mathcal{K} is a *free amalgamation class*.

We will give examples of free amalgamation classes (in the case where the language L is not relational) in Section 5.

2 Free amalgamation classes are Ramsey

The proof of Theorem 1.3 is a variation of the *Partite Construction* introduced by Nešetřil and Rödl for classes of hypergraphs and relational structures (see [29]) which was recently extended to classes with unary [19] and later general closures (or functions) [21]. The Partite Construction is a machinery which allows one to transform one Ramsey class into another, more special, Ramsey class. What follows is a Partite Construction proof (as used previously, for example, in [26]) done in the context of structures involving set-valued functions.

The basic Ramsey class in our construction will be provided by the following result about the Ramsey property of ordered structures with relations and (total) functions.

Theorem 2.1 (Ramsey theorem for finite models, Theorem 2.19 of [21]). *Suppose L is a language containing a relational symbol \leq and let $\overrightarrow{\text{Str}}(L)$ be the class of all finite L -structures where \leq is a linear ordering of the vertices. Then $\overrightarrow{\text{Str}}(L)$ is a Ramsey class.*

This is a strengthening of the theorem giving the Ramsey property of ordered relational structures proved independently by Nešetřil-Rödl [31] and Abramson-Harrington [1] in 1970s. Considering functions is a rather difficult task and the proof of Theorem 2.1 involves a recursive nesting of the Partite Constructions to establish valid non-unary closures. Building on Theorem 2.1 our task is significantly easier and we only concentrate on further refining the Ramsey structure given by Theorem 2.1 into one belonging to a given free amalgamation class. This is done by tracking all irreducible substructures of the object constructed.

We shall remark that recent paper of second two authors [21] defines structures with partial function to singletons only. In the Ramsey context this is however equivalent because the structures are ordered. Thus it is possible to replace a function F by symbols F^1, F^2, F^3, \dots and put $F^i(\vec{t})$ to be the i -th element of $F(\vec{t})$ whenever it is defined. With this it may be seen then that Theorem 2.11 of [21] implies Theorem 1.3. However the proof presented here is cleaner and much simpler.

As mentioned before, a relational structure \mathbf{A} is *irreducible* if every pair of vertices belongs to some tuple in a relation of \mathbf{A} . It is well known that every free amalgamation class \mathcal{K} of relational structures can be equivalently described as a class of finite structures that contains no copies of structures from a fixed family \mathcal{F} of irreducible relational structures (see, for example, [21]). In fact the family \mathcal{F} consists of all minimal structures not belonging to class \mathcal{K} . This easy observation explains the correspondence of Theorem 1.1 and Theorem 1.2.

Our construction is based on the following refinement of the notion of irreducible structure in the context of structures with functions (which allows us to strengthen the construction in [21]):

Definition 2.1. An L -structure \mathbf{A} is *irreducible* if it cannot be created as a free amalgamation of any two of its proper substructures.

Example. Consider the language L consisting of one binary relation R and one unary function F . An example of an irreducible structure is a structure \mathbf{A} (depicted in Figure 2) on vertices $A = \{a, b, c, d\}$ where $(a, b) \in R_{\mathbf{A}}$, $F_{\mathbf{A}}(a) = \{c\}$, $F_{\mathbf{A}}(b) = \{d\}$ and $F_{\mathbf{A}}(a) = F_{\mathbf{A}}(b) = \emptyset$. This structure is reducible if F is seen as a relation rather than a function.

The basic part of our construction of Ramsey objects is a variant of the Partite Lemma (introduced in [29] and refined for closures in [21]) which deals with the following objects.

Definition 2.2 (**A-partite system**). Let L be a language and \mathbf{A} an L -structure. Assume $A = \{1, 2, \dots, a\}$. An **A-partite L -system** is a tuple $(\mathbf{A}, \mathcal{X}_{\mathbf{B}}, \mathbf{B})$ where \mathbf{B} is an L -structure and $\mathcal{X}_{\mathbf{B}} = \{X_{\mathbf{B}}^1, X_{\mathbf{B}}^2, \dots, X_{\mathbf{B}}^a\}$ is a partition of the vertex set of \mathbf{B} into a many classes $X_{\mathbf{B}}^i$, called *parts* of \mathbf{B} , such that




Figure 2: An example of an irreducible structure with a binary relation R and a unary function F .

1. the mapping π which maps every $x \in X_{\mathbf{B}}^i$ to i , $i = 1, 2, \dots, a$, is a homomorphism $\mathbf{B} \rightarrow \mathbf{A}$ (π is called the *projection*);
2. every tuple in every relation of \mathbf{B} meets every class $X_{\mathbf{B}}^i$ in at most one element (i.e. these tuples are called *transversal* with respect to the partition).
3. for every function $F \in L_{\mathcal{F}}$ it holds that for every $\vec{t} \in A^{\text{arity}(F)}$ such that $F_{\mathbf{A}}(\vec{t}) \neq \emptyset$ the tuple created by concatenation of \vec{t} and $F_{\mathbf{B}}(\vec{t})$ (in any order) is transversal.

The *isomorphisms* and *embeddings* of \mathbf{A} -partite systems, say of \mathbf{B}_1 into \mathbf{B}_2 , are defined as the isomorphisms and embeddings of structures together with the condition that all parts are being preserved (the part $X_{\mathbf{B}_1}^i$ is mapped to $X_{\mathbf{B}_2}^i$ for every $i = 1, 2, \dots, a$). Of course, \mathbf{A} itself can be considered as an \mathbf{A} -partite system.

We say that an \mathbf{A} -partite L -system is *transversal* if all of its parts consist of at most one vertex. Thus \mathbf{A} -partite L -systems are naturally ordered by ordering of its parts.

Lemma 2.2 (Partite Lemma with relations and functions). *Let L be a language, \mathbf{A} be a finite L -structure, and \mathbf{B} be a finite \mathbf{A} -partite L -system. Then there exists a finite \mathbf{A} -partite L -system \mathbf{C} such that*

$$\mathbf{C} \longrightarrow (\mathbf{B})_2^{\mathbf{A}}.$$

Moreover if every irreducible subsystem of \mathbf{B} is transversal, then we can also ensure that every irreducible subsystem of \mathbf{C} is transversal.

(Compare the Partite Lemma in [21]. Here we introduce the statement about transversality of irreducible substructures. Note that the embeddings considered in the Ramsey statement are all as \mathbf{A} -partite systems.)

Advancing the proof of Lemma 2.2, for completeness, we briefly recall the Hales-Jewett Theorem [12]. Consider the family of functions $f : \{1, 2, \dots, N\} \rightarrow \Sigma$ for some finite alphabet Σ . A *combinatorial line* \mathcal{L} is a pair (ω, h) where $\emptyset \neq \omega \subseteq \{1, 2, \dots, N\}$ and h is a function from $\{1, 2, \dots, N\} \setminus \omega$ to Σ . The combinatorial line \mathcal{L} describes the family of all those functions $f : \{1, 2, \dots, N\} \rightarrow \Sigma$ that are constant on ω and $f(i) = h(i)$ otherwise. The Hales-Jewett Theorem guarantees, for sufficiently large N , that for every 2-colouring of the functions $f : \{1, 2, \dots, N\} \rightarrow \Sigma$ there exists a monochromatic combinatorial line.

Proof of Lemma 2.2. Assume without loss of generality $A = \{1, 2, \dots, a\}$ and denote by $\mathcal{X}_{\mathbf{B}} = \{X_{\mathbf{B}}^1, X_{\mathbf{B}}^2, \dots, X_{\mathbf{B}}^a\}$ the parts of \mathbf{B} . We take N sufficiently large (that will be specified later) and construct an \mathbf{A} -partite L -system \mathbf{C} with parts $\mathcal{X}_{\mathbf{C}} = \{X_{\mathbf{C}}^1, X_{\mathbf{C}}^2, \dots, X_{\mathbf{C}}^a\}$ as follows:

1. For every $1 \leq i \leq a$ let $X_{\mathbf{C}}^i$ be the set of all functions

$$f : \{1, 2, \dots, N\} \rightarrow X_{\mathbf{B}}^i.$$

2. For every relational symbol $R \in L_{\mathcal{R}}$, put

$$(f_1, f_2, \dots, f_{\text{arity}(R)}) \in R_{\mathbf{C}}$$

if and only if for every $1 \leq j \leq N$ it holds that

$$(f_1(j), f_2(j), \dots, f_{\text{arity}(R)}(j)) \in R_{\mathbf{B}}.$$

3. For every function symbol $F \in L_{\mathcal{F}}$ we put

$$F_{\mathbf{C}}(f_1, f_2, \dots, f_{\text{arity}(F)}) \neq \emptyset$$

if and only if there exists $r \geq 1$ and a set $\{i_1, i_2, \dots, i_r\} \subseteq A$ such that for every $1 \leq j \leq N$ it holds that $F_{\mathbf{B}}(f_1(j), f_2(j), \dots, f_{\text{arity}(F)}(j))$ is defined and its value contains vertices precisely in parts i_1, i_2, \dots, i_r . In this case the value of $F_{\mathbf{C}}(f_1, f_2, \dots, f_{\text{arity}(F)})$ consists of one vertex in each of parts i_1, i_2, \dots, i_r . The vertex in part i_k is the function $f_k : \{1, 2, \dots, N\} \rightarrow X_{\mathbf{B}}^{i_k}$ defined by putting $f_k(j')$ to be the (unique) vertex in $F_{\mathbf{B}}(f_1(j'), f_2(j'), \dots, f_{\text{arity}(F)}(j')) \cup X_{\mathbf{B}}^{i_k}$ (for each $1 \leq j' \leq N$).

This completes the construction of \mathbf{C} . It is easy to check that \mathbf{C} is indeed an \mathbf{A} -partite L -system with parts $\mathcal{X}_{\mathbf{C}} = \{X_{\mathbf{C}}^1, X_{\mathbf{C}}^2, \dots, X_{\mathbf{C}}^a\}$.

We verify that, if N is large enough, $\mathbf{C} \rightarrow (\mathbf{B})_2^{\mathbf{A}}$. Let $\tilde{\mathbf{A}}_1, \tilde{\mathbf{A}}_2, \dots, \tilde{\mathbf{A}}_t$ be an enumeration of all subsystems of \mathbf{B} which are isomorphic to \mathbf{A} . Put $\Sigma = \{1, 2, \dots, t\}$ which we consider as an alphabet. Each combinatorial line $\mathcal{L} = (\omega, h)$ in Σ^N corresponds to an embedding $e_{\mathcal{L}} : \mathbf{B} \rightarrow \mathbf{C}$ which assigns to every vertex $v \in X_{\mathbf{B}}^p$ a function $e_{\mathcal{L}}(v) : \{1, 2, \dots, N\} \rightarrow X_{\mathbf{B}}^p$ (i.e. a vertex of $X_{\mathbf{C}}^p$) such that:

$$e_{\mathcal{L}}(v)(j) = \begin{cases} v & \text{for } j \in \omega, \text{ and,} \\ \text{the unique vertex in } \tilde{\mathbf{A}}_{h(j)} \cap X_{\mathbf{B}}^p & \text{otherwise.} \end{cases}$$

It follows from the construction of \mathbf{C} and from the fact that \mathbf{B} has a projection to \mathbf{A} that $e_{\mathcal{L}}$ is an embedding.

Let N be the Hales-Jewett number guaranteeing a monochromatic line in any 2-colouring of the N -dimensional cube over an alphabet Σ . Now assume that $\mathcal{A}_1, \mathcal{A}_2$ is a 2-colouring of all copies of \mathbf{A} in \mathbf{C} . Using the construction of \mathbf{C} we see that among copies of \mathbf{A} are copies induced by an N -tuple $(\tilde{\mathbf{A}}_{u(1)}, \tilde{\mathbf{A}}_{u(2)}, \dots, \tilde{\mathbf{A}}_{u(N)})$ of copies of \mathbf{A} in \mathbf{B} for every function $u : \{1, 2, \dots, N\} \rightarrow \{1, 2, \dots, t\}$. However such copies are coded by the elements of the cube $\{1, 2, \dots, t\}^N$ and thus there is a monochromatic combinatorial line \mathcal{L} . The monochromatic copy of \mathbf{B} is then $e_{\mathcal{L}}(\mathbf{B})$.

Finally we verify that if every irreducible subsystem of \mathbf{B} is transversal then also every irreducible subsystem of \mathbf{C} is transversal. Assume the contrary and denote by \mathbf{D} a non-transversal irreducible subsystem of \mathbf{C} . Denote by f_1, f_2, \dots, f_n an enumeration of all distinct vertices of \mathbf{D} . By non-transversality




Figure 3: The construction of \mathbf{P}_0 .

assume that f_1 and f_2 are in the same part. For every $j \in \{1, 2, \dots, N\}$ denote by \mathbf{D}_j the substructure of \mathbf{B} on vertices $f_1(j), f_2(j), \dots, f_n(j)$. Because \mathbf{D}_j is a homomorphic image of \mathbf{D} it is irreducible and thus it follows that \mathbf{D}_j is transversal. Consequently $f_1(j) = f_2(j)$. Because this holds for every choice of j , we have $f_1 = f_2$. A contradiction. \square

Proof of Theorem 1.3. Given $\mathbf{A}, \mathbf{B} \in \vec{\mathcal{K}} \subseteq \vec{\text{Str}}(\vec{L})$ use Theorem 2.1 to obtain

$$\mathbf{C}_0 \longrightarrow (\mathbf{B})_2^{\mathbf{A}}.$$

Enumerate all copies of \mathbf{A} in \mathbf{C}_0 as $\{\tilde{\mathbf{A}}_1, \tilde{\mathbf{A}}_2, \dots, \tilde{\mathbf{A}}_b\}$. We will define \mathbf{C}_0 -partite systems (‘pictures’) $\mathbf{P}_0, \mathbf{P}_1, \dots, \mathbf{P}_b$ such that

- (i) every irreducible subsystem \mathbf{E} of \mathbf{P}_k is transversal and (if seen as a structure) is isomorphic to some substructure of \mathbf{B} , and,
- (ii) in any 2-colouring of $(\mathbf{P}_k)_{\tilde{\mathbf{A}}}$, $1 \leq k \leq b$, there exists a copy $\tilde{\mathbf{P}}_{k-1}$ of \mathbf{P}_{k-1} such that all copies of \mathbf{A} with a projection to $\tilde{\mathbf{A}}_k$ are monochromatic.

We then show that putting \mathbf{C} to be \mathbf{P}_b (seen as a structure) with the linear order completed arbitrarily (extending the order of the parts), we have the desired Ramsey property $\mathbf{C} \longrightarrow (\mathbf{B})_2^{\mathbf{A}}$.

We first verify that if (i) holds, then $\mathbf{C} \in \vec{\mathcal{K}}$. Assume the contrary. Denote by \mathbf{F} the minimal substructure of \mathbf{C} such that $\mathbf{F} \notin \vec{\mathcal{K}}$. Because \mathcal{K} is a free amalgamation class, we know that the unordered reduct of \mathbf{F} is irreducible. By (i) however \mathbf{F} is a substructure of $\mathbf{B} \in \mathcal{K}$. A contradiction to \mathcal{K} being hereditary.

It remains to prove (i) and (ii). Put $C_0 = \{1, \dots, c\}$ and $\mathcal{X}_{\mathbf{P}_k} = \{X_k^1, X_k^2, \dots, X_k^c\}$. We proceed by induction on k .

1. The picture \mathbf{P}_0 is constructed as a disjoint union of copies of \mathbf{B} . For every copy $\tilde{\mathbf{B}}$ of \mathbf{B} in \mathbf{C}_0 we include a copy $\tilde{\mathbf{B}}'$ of $\tilde{\mathbf{B}}$ in \mathbf{P}_0 which projects onto $\tilde{\mathbf{B}}$. The copies corresponding to different $\tilde{\mathbf{B}}$ are disjoint (see Figure 3). This indeed satisfies (i).
2. Suppose the picture \mathbf{P}_k is already constructed. Let \mathbf{B}_k be the substructure of \mathbf{P}_k induced by \mathbf{P}_k on vertices which project to $\tilde{\mathbf{A}}_{k+1}$. We use the Partite Lemma 2.2 to obtain an $\tilde{\mathbf{A}}_{k+1}$ -partite system \mathbf{D}_{k+1} with $\mathbf{D}_{k+1} \longrightarrow (\mathbf{B}_k)_2^{\tilde{\mathbf{A}}_{k+1}}$. Now consider all copies of \mathbf{B}_k in \mathbf{D}_{k+1} and extend each of these structures to a copy of \mathbf{P}_k (using free amalgamation of

\mathbf{C}_0 -partite systems). These copies are disjoint outside \mathbf{D}_{k+1} . In this extension we preserve the parts of all the copies. The result of this multiple amalgamation is \mathbf{P}_{k+1} . Because $\mathbf{D}_{k+1} \rightarrow (\mathbf{B}_k)_2^{\mathbf{A}^{k+1}}$ we know that \mathbf{P}_{k+1} satisfies (ii).

From the ‘Moreover’ part of Lemma 2.2, we may assume that \mathbf{D}_{k+1} satisfies (i). Because \mathbf{P}_{k+1} is created by a series of free amalgamations, it follows that \mathbf{P}_{k+1} also satisfies (i).

Put $\mathbf{C} = \mathbf{P}_b$. It follows easily that $\mathbf{C} \rightarrow (\mathbf{B})_2^{\mathbf{A}}$. Indeed, by a backward induction on k one proves that in any 2-colouring of $\binom{\mathbf{C}}{\mathbf{A}}$ there exists a copy $\tilde{\mathbf{P}}_0$ of \mathbf{P}_0 such that the colour of a copy of \mathbf{A} in $\tilde{\mathbf{P}}_0$ depends only on its projection to \mathbf{C}_0 . As this in turn induces a colouring of the copies of \mathbf{A} in \mathbf{C}_0 , we obtain a monochromatic copy of \mathbf{B} in $\tilde{\mathbf{P}}_0$. \square

3 Ordering property

For many Ramsey classes \mathcal{K} , the class $\vec{\mathcal{K}}$ of free orderings of structures in \mathcal{K} has the ordering property. In fact, for free amalgamation classes we can give a full characterisation by means of the following easy (and folklore) proposition.

Proposition 3.1. *Let \mathcal{K} be a free amalgamation class of L -structures. Then $\vec{\mathcal{K}}$ has the ordering property if and only if all closures $\text{Cl}_{\mathbf{A}}(u)$ of vertices are mutually isomorphic single element structures.*

Remark. It does not follow from Proposition 3.1 that the class \mathcal{K} must have only trivial closures in order for $\vec{\mathcal{K}}$ to have the ordering property. Consider, for example, a class of structures with one binary function F such that the images of tuples with duplicated vertices are empty. Here all vertices are closed, but pairs of vertices have non-trivial closures. A related example is discussed in Section 5.2.

Proof. Assume that there is $\vec{\mathbf{A}} \in \mathcal{K}$ and $u, v \in \vec{\mathbf{A}}$ such that $\text{Cl}_{\mathbf{A}}(u)$ is not isomorphic to $\text{Cl}_{\mathbf{A}}(v)$. Then one can choose an ordering $\vec{\mathbf{A}}$ of \mathbf{A} so that the set of all vertices $v' \in A$ such that $\text{Cl}_{\mathbf{A}}(v')$ is isomorphic to $\text{Cl}_{\mathbf{A}}(v)$ forms an initial segment. Now assume, to the contrary, that there exists $\mathbf{B} \in \mathcal{K}$ such that every ordering of \mathbf{B} contains a copy of $\vec{\mathbf{A}}$. This is clearly not possible because one can choose an ordering of \mathbf{B} such that all vertices $u' \in B$ such that $\text{Cl}_{\mathbf{A}}(u')$ is isomorphic to $\text{Cl}_{\mathbf{A}}(u)$ forms an initial segment. This is a contradiction to $\vec{\mathcal{K}}$ having the ordering property.

Now consider the case that all closures of vertices are isomorphic, but not trivial. Because the intersection of two closures is also closed, it follows that closures of vertices are disjoint and thus it is always possible to order structures in a way that all vertex closures form intervals. It follows that there is no \mathbf{B} witnessing the ordering property for any $\vec{\mathbf{A}} \in \mathcal{K}$ which is ordered so that some vertex closure is not an interval.

Finally assume that \mathcal{K} is a free amalgamation class where all closures of vertices are trivial and mutually isomorphic. We may assume that there are no unary relations. Given $\vec{\mathbf{A}} \in \vec{\mathcal{K}}$ we construct $\vec{\mathbf{B}}_0$ from a disjoint copy of $\vec{\mathbf{A}}$ and $\vec{\mathbf{A}}^{\text{op}}$ (by this we mean a structure created from $\vec{\mathbf{A}}$ by reversing the

linear ordering of vertices). Now extend $\vec{\mathbf{B}}_0$ to $\vec{\mathbf{B}}_1$ by adding, between every neighbouring pair of vertices $u \leq_{\vec{\mathbf{B}}_0} v \in B_0$ a new vertex $n_{u,v}$. Extend the order so $u \leq_{\vec{\mathbf{B}}_1} n_{u,v} \leq_{\vec{\mathbf{B}}_1} v$. By free amalgamation, $\vec{\mathbf{B}}_1 \in \vec{\mathcal{K}}$.

Denote by $\vec{\mathbf{I}} \in \vec{\mathcal{K}}$ a structure consisting of two vertices $u \leq_{\vec{\mathbf{I}}} v$ and no relations or functions containing both of them besides $\leq_{\vec{\mathbf{I}}}$ (such a structure can be created by means of the free amalgamation of two copies of the unique single vertex structure in \mathbf{K} over an empty set with order completed arbitrarily and thus clearly $\vec{\mathbf{I}} \in \vec{\mathcal{K}}$). Find $\vec{\mathbf{B}} \in \vec{\mathcal{K}}$ with $\vec{\mathbf{B}} \rightarrow (\vec{\mathbf{B}}_1)_{\vec{\mathbf{I}}}$ by application of Theorem 1.3.

We verify that $\vec{\mathbf{B}}$ has the desired property. Let $\vec{\mathbf{C}}$ be any re-ordering of $\vec{\mathbf{B}}$. This re-ordering induces a coloring of $(\vec{\mathbf{B}})_{\vec{\mathbf{I}}}$: if the order of the points in some $\vec{\mathbf{I}}' \in (\vec{\mathbf{B}})_{\vec{\mathbf{I}}}$ agrees with the order in $\vec{\mathbf{C}}$, then color $\vec{\mathbf{I}}'$ red, and blue otherwise. The monochromatic copy of $\vec{\mathbf{B}}_1$ will have the property that it is either ordered in the same way as $\vec{\mathbf{B}}_1$ or the order is reversed. By construction of $\vec{\mathbf{B}}_0$ it follows that in both alternatives there is a copy of $\vec{\mathbf{A}}$. \square

In the following we generalize the main idea of this proof (the idea of which goes back to [30]) to classes with non-trivial closures of vertices. Free orderings do not suffice anymore and we have to define carefully the admissible orderings.

3.1 Admissible orderings

Definition 3.1. Let \mathbf{A} be an L -structure. If $a, b \in A$ we write $a \sim_A b$ if $\text{Cl}_{\mathbf{A}}(a) = \text{Cl}_{\mathbf{A}}(b)$. This is an equivalence relation on A and we refer to the classes as the *closure-components* of \mathbf{A} . The class containing a will be denoted by $\text{Cc}_{\mathbf{A}}(a)$.

If $a \in A$ we define the *level* $l_{\mathbf{A}}(a)$ of a in \mathbf{A} inductively. We say that $l_{\mathbf{A}}(a) = 0$ in the case where $\text{Cc}_{\mathbf{A}}(a) = \text{Cl}_{\mathbf{A}}(a)$; otherwise $l_{\mathbf{A}}(a) = l_{\mathbf{A}}(b) + 1$ where b is a vertex of the maximal level in \mathbf{A} amongst vertices in $\text{Cl}_{\mathbf{A}}(a) \setminus \text{Cc}_{\mathbf{A}}(a)$.

We say that $\mathbf{A} \in \mathcal{K}$ is a *closure-extension* at level k if there is a unique closure-component C of vertices of level k in \mathbf{A} and $\text{Cl}_{\mathbf{A}}(a) = \mathbf{A}$ for every $a \in C$. In this case, we write $A^\circ = A \setminus C$. Every closure of a vertex is a closure-extension.

We say that two closure-components C and C' of $\vec{\mathbf{A}}$ (or their closures) are *homologous* if $\text{Cl}_{\mathbf{A}}(C)$ and $\text{Cl}_{\mathbf{A}}(C')$ are isomorphic and this isomorphism is an identity on $\text{Cl}_{\mathbf{A}}(C) \setminus C = \text{Cl}_{\mathbf{A}}(C') \setminus C'$. Note that if $C \neq C'$, then $\text{Cl}_{\mathbf{A}}(C) \setminus C = \text{Cl}_{\mathbf{A}}(C) \cap \text{Cl}_{\mathbf{A}}(C')$ is closed in \mathbf{A} .

Example. Consider the class \mathcal{T} of forests represented by a single unary function (the predecessor relation) described in Section 1.2. In this class the closure of a vertex is the path to a root vertex. Every vertex thus forms a trivial closure component and its level is determined by the distance to the root vertex. In this particular case a structure is a closure-extension if and only if \mathbf{A} is an oriented path (that is structure on vertex set $\mathbf{A} = \{v_1, v_2, \dots, v_n\}$ and $F_{\mathbf{A}}(v_i) = v_{i+1}$ for every $1 \leq i < n$).

Observe also that for a closure-extension \mathbf{A} (depicted in Figure 4), the set of vertices A° is not necessarily closed. Consider a language with a function F^1 arity 1 a function F^2 of arity 2. The structure \mathbf{A} with $A = \{a, b, c\}$, $F_{\mathbf{A}}^1(c) =$




Figure 4: Example of a closure-extension \mathbf{A} where A° is not a substructure.

$\{a, b\}$, $F_{\mathbf{A}}^2(a, b) = \{c\}$ is a closure-extension of level 1: $l(a) = l(b) = 0$ and $l(c) = 1$, the set A° is $\{a, b\}$, however $\text{Cl}_{\mathbf{A}}\{a, b\} = \{a, b, c\}$.

Suppose $\mathbf{A}, \mathbf{B} \in \mathcal{K}$ are closure-extensions and $\vec{\mathbf{A}}, \vec{\mathbf{B}} \in \vec{\mathcal{K}}$ are orderings. We say that these are *similar* if there is an isomorphism $\alpha : \mathbf{A} \rightarrow \mathbf{B}$ which is also order-preserving when seen as a mapping $\alpha : A^\circ \rightarrow B^\circ$. This is an equivalence relation and in our admissible orderings, we choose a fixed representative from each similarity-type of ordered closure-extension.

In what follows we shall assume that we have fixed some total ordering $\vec{\mathbf{A}} \trianglelefteq \vec{\mathbf{B}}$ between isomorphism types of orderings of closure-extensions such that

$$\text{S1 } |A| < |B| \text{ implies } \vec{\mathbf{A}} \trianglelefteq \vec{\mathbf{B}}.$$

(In particular, \trianglelefteq is a well ordering.)

First we define a preorder of vertices which we will later refine to a linear order. Given two vertices $u \neq v \in \vec{\mathbf{A}}$ we write $u \preceq_{\vec{\mathbf{A}}} v$ if one of the following holds:

- P1 $\text{Cl}_{\vec{\mathbf{A}}}(u) \trianglelefteq \text{Cl}_{\vec{\mathbf{A}}}(v)$ and they are not isomorphic;
- P2 $\text{Cl}_{\vec{\mathbf{A}}}(u)$ is isomorphic to $\text{Cl}_{\vec{\mathbf{A}}}(v)$ but $\text{Cl}_{\vec{\mathbf{A}}}(u) \setminus \text{Cc}_{\vec{\mathbf{A}}}(v)$ is lexicographically before $\text{Cl}_{\vec{\mathbf{A}}}(v) \setminus \text{Cc}_{\vec{\mathbf{A}}}(v)$ considering the order $\leq_{\vec{\mathbf{A}}}$;
- P3 $\text{Cl}_{\vec{\mathbf{A}}}(u)$ and $\text{Cl}_{\vec{\mathbf{A}}}(v)$ are homologous closure-extensions.

Note that this is indeed a preorder on the vertices of \mathbf{A} . We can now describe our class of admissible orderings.

Definition 3.2. Suppose \mathcal{K} is a free amalgamation class. We say that $\mathcal{O} \subseteq \vec{\mathcal{K}}$ is a *class of admissible orderings* of structures in \mathcal{K} if the following conditions hold.

- A1 If $\mathbf{A} \in \mathcal{K}$, then there is some ordering $\leq_{\vec{\mathbf{A}}}$ of \mathbf{A} in \mathcal{O} .
- A2 \mathcal{O} is closed for substructures.
- A3 For every $\vec{\mathbf{A}} \in \mathcal{O}$, the ordering $\leq_{\vec{\mathbf{A}}}$ refines $\preceq_{\vec{\mathbf{A}}}$.
- A4 For every $\vec{\mathbf{A}} \in \mathcal{O}$, the closure-components form linear intervals in $\leq_{\vec{\mathbf{A}}}$.
- A5 For every $\mathbf{B} \in \mathcal{K}$, if $\mathbf{A}_1, \mathbf{A}_2, \dots, \mathbf{A}_n$ is a family of substructures and \leq is a linear order of $A = \cup_{1 \leq i \leq n} A_i$ such that

- (a) \leq satisfies the conclusions of A3 and A4; and
- (b) each substructure of \mathbf{B} contained in A is admissibly ordered by \leq ;

then there exists $\vec{\mathbf{B}} \in \mathcal{O}$ such that $\leq_{\vec{\mathbf{B}}}$ restricted to A is \leq .

A6 Suppose that $\vec{\mathbf{A}}, \vec{\mathbf{B}} \in \mathcal{O}$ are similar ordered closure-extensions. Then $\vec{\mathbf{A}}$ is isomorphic to $\vec{\mathbf{B}}$.

Example. Consider $\mathbf{A} \in \mathcal{T}$ where \mathcal{T} is the class of forests discussed in Section 1.2. Because the closure-extensions are all formed by oriented paths, the order \leq requires vertices to be ordered according to their levels (in particular all root vertices come first and can be ordered arbitrarily). The sons of a vertex $v \in \mathbf{A}$ are trivial homologous components and thus they are required to always form an interval and the order amongst these intervals is given by order of their fathers.

There is some flexibility in the definition of admissibility. For example it is possible to order forests in a way that every tree (and recursively every subtree) forms an interval. The particular choice is however not very important as it can be shown that they are all equivalent up to bi-definability (this follows as \mathcal{O} is a Ramsey class, see [22]).

Proposition 3.2. *Suppose \mathcal{K} is a free amalgamation class. Then there is a class \mathcal{O} of admissible orderings of \mathcal{K} .*

Proof. We proceed by induction on $\vec{\mathcal{K}}$ ordered arbitrarily in order of increasing number of vertices. In this order, for every $\vec{\mathbf{C}} \in \vec{\mathcal{K}}$ we decide if $\vec{\mathbf{C}} \in \mathcal{O}$ or not by a variant of a greedy algorithm. In the induction step assume that we already decided the presence in \mathcal{O} for every proper substructure of $\vec{\mathbf{C}}$.

We put $\vec{\mathbf{C}} \in \mathcal{O}$ if and only if:

- O1 $\vec{\mathbf{C}}$ satisfies A3 and A4,
- O2 every proper substructure of $\vec{\mathbf{C}}$ is in \mathcal{O} , and,
- O3 if $\vec{\mathbf{C}}$ is an ordered closure-extension, then there is no similar but non-isomorphic $\vec{\mathbf{D}} \in \mathcal{O}$.

This finishes the description of \mathcal{O} . We verify that the conditions of Definition 3.2 are satisfied.

First we check A5. Assume, to the contrary, that there are $\mathbf{B} \in \mathcal{K}$, substructures $\mathbf{A}_1, \mathbf{A}_2, \dots, \mathbf{A}_n$ and a linear order \leq on $A = \cup_{1 \leq i \leq n} A_i$ without such an extension. From all counter-examples choose one minimizing $|B|$ and among those, minimize $|B| - |A|$.

We consider three cases:

1. Suppose \mathbf{B} is not a closure-extension and $B = A$. It follows that the order \leq satisfies both O1 and O2 and thus we can put $\leq_{\vec{\mathbf{B}}} = \leq$ and obtain $\vec{\mathbf{B}} \in \mathcal{O}$, a contradiction.
2. Suppose \mathbf{B} is a closure-extension and $B^\circ = A$. Extend \leq to an order of $\vec{\mathbf{B}}$ in a way it satisfies O1. In this case $\vec{\mathbf{B}}$ satisfies O2 because no proper substructure contains $B \setminus B^\circ$. Furthermore, we may assume that O3 holds. This is a contradiction to $\vec{\mathbf{B}} \notin \mathcal{O}$.

3. Suppose neither of the previous cases apply. Then there is a proper closure-extension $\mathbf{C} \subseteq \mathbf{B}$ such that $C \not\subseteq A$. Denote by \leq' the order \leq restricted to $C \cap A$ and consider the structures $\mathbf{C}_i = \mathbf{A}_i \cap \mathbf{C}$, $1 \leq i \leq n$. From the minimality of the counter-example and because $|C| < |B|$ we know that order of \leq' can be extended to an admissible order $\vec{\mathbf{C}}$ of \mathbf{C} . Now extend to $C \cup A$ the orders \leq and $\leq_{\vec{\mathbf{C}}}$ in such a way that A3 and A4 are satisfied. This combined order along with the family of substructures $\mathbf{C}, \mathbf{A}_1, \mathbf{A}_2, \dots, \mathbf{A}_n$ contradicts the second assumption about the minimality of the counter-example.

This finishes proof of A5.

Condition A1 is implied by A5. Conditions A2, A3, A4 and A6 follows directly from the construction of \mathcal{O} . \square

3.2 Proof of Theorem 1.4

Advancing the proof of Theorem 1.4 we show two lemmas which generalize the main ideas of proof of Proposition 3.1.

Lemma 3.3. *Let \mathcal{R} be a Ramsey class of ordered structures. Then for every $\vec{\mathbf{A}} \in \mathcal{R}$ there exists $\vec{\mathbf{B}} \in \mathcal{R}$ such that every re-ordering $\vec{\mathbf{C}} \in \mathcal{R}$ of $\vec{\mathbf{B}}$ contains a re-ordering $\vec{\mathbf{A}}$ of $\vec{\mathbf{A}}$ such that every two isomorphic substructures of $\vec{\mathbf{A}}$ are also isomorphic in the order of $\vec{\mathbf{A}}$.*

Proof. Let $\vec{\mathbf{A}}$ be a structure and $\vec{\mathbf{A}}_0$ a substructure of $\vec{\mathbf{A}}$. Denote by n the number of possible re-orderings of $\vec{\mathbf{A}}_0$ (i.e. $n = |A_0|!$). By the Ramsey property construct $\vec{\mathbf{B}}_0$ such that $\vec{\mathbf{B}}_0 \rightarrow (\vec{\mathbf{A}})_{\vec{\mathbf{A}}_0}^n$. Every re-ordering of $\vec{\mathbf{B}}_0$ induces an n -coloring of copies of $\vec{\mathbf{A}}_0$ in $\vec{\mathbf{B}}_0$ and so by the Ramsey property there exists a copy $\vec{\mathbf{A}}$ of re-ordered $\vec{\mathbf{A}}$ in $\vec{\mathbf{B}}_0$ having the property that all copies of $\vec{\mathbf{A}}_0$ in $\vec{\mathbf{A}}$ are ordered the same way. The statement follows by iterating this argument for every substructure of $\vec{\mathbf{A}}$. \square

Lemma 3.4. *Let \mathcal{K} by a free amalgamation class. Then for every $\vec{\mathbf{A}} \in \mathcal{K}$ in which every closure-component forms an interval (in the order of $\vec{\mathbf{A}}$) there exists $\vec{\mathbf{B}}$ such that every re-ordering $\vec{\mathbf{C}}$ of $\vec{\mathbf{B}}$ where every closure-component forms an interval (in the order of $\vec{\mathbf{C}}$) contains a copy of a re-ordering of $\vec{\mathbf{A}}$ where the order between vertices in distinct homologous closure-components is preserved.*

Proof. For simplicity we show the construction for a given $\vec{\mathbf{A}}$ and two distinct homologous closure-components $C_1 \leq_{\vec{\mathbf{A}}} C_2$. The full statement can be shown by iterating the argument for every such pair. The construction is schematically depicted in Figure 5.

First observe that C_1 and C_2 are not in the closure of $\text{Cl}_{\mathbf{A}}(C_1) \setminus C_1 = \text{Cl}_{\mathbf{A}}(C_2) \setminus C_2$ because this would imply $C_1 = C_2$.

Now, by a free amalgamation construct $\vec{\mathbf{A}}_1$ extending $\vec{\mathbf{A}}$ with a new closure-component C homologous to both C_1 and C_2 where the linear order is extended so that $C_1 \leq_{\vec{\mathbf{A}}} C \leq_{\vec{\mathbf{A}}} C_2$. Denote by $\vec{\mathbf{A}}_1^{\text{op}}$ the structure created from $\vec{\mathbf{A}}_1$ by re-ordering closure-components C_1, C and C_2 so that $C_2 \leq_{\vec{\mathbf{A}}_1^{\text{op}}} C \leq_{\vec{\mathbf{A}}_1^{\text{op}}} C_1$. Denote by $\vec{\mathbf{A}}_2$ the free amalgamation of $\vec{\mathbf{A}}_1$ and $\vec{\mathbf{A}}_1^{\text{op}}$ over $\text{Cl}_{\mathbf{A}}(C_1) \setminus C_1$. Because \mathcal{K} is a free amalgamation class we have $\vec{\mathbf{A}}_1, \vec{\mathbf{A}}_1^{\text{op}}, \vec{\mathbf{A}}_2 \in \mathcal{K}$.




Figure 5: Construction used in the proof of Lemma 3.4. Pairs of closure components which are not necessarily free are connected by dotted lines.

Note that the addition of C into \mathbf{A}_1 is necessary only when $\text{Cl}(\vec{\mathbf{A}}(C_1 \cup C_2))$ is not a result of free amalgamation of two copies of $\text{Cl}_{\vec{\mathbf{A}}}(C_1)$. In general there may be relations spanning C_1 and C_2 which would make use of the Ramsey argument below impossible.

Put $\vec{\mathbf{I}} = \text{Cl}_{\vec{\mathbf{A}}_1}(C_1 \cup C)$ and use Theorem 1.3 to find $\vec{\mathbf{B}} \in \vec{\mathcal{K}}$ containing $\vec{\mathbf{A}}_2$ such that $\vec{\mathbf{B}} \rightarrow (\vec{\mathbf{A}}_2)_{\vec{\mathbf{I}}}$. Now every re-ordering $\vec{\mathbf{C}}$ of $\vec{\mathbf{B}}$ induces a 2-coloring of the copies of $\vec{\mathbf{I}}$ in $\vec{\mathbf{B}}$ which in turn leads to the existence of a copy of a re-ordering of $\vec{\mathbf{A}}$ where the order of C_1 and C_2 is preserved. \square

Now we are finally ready to prove the main result of this section.

Proof of Theorem 1.4. Given $\vec{\mathbf{A}} \in \mathcal{O}$ we construct $\vec{\mathbf{B}} \in \vec{\mathcal{K}}$ with $\vec{\mathbf{A}}$ as a substructure such that every admissible re-ordering $\vec{\mathbf{C}} \in \mathcal{O}$ of $\vec{\mathbf{B}}$ contains a copy of $\vec{\mathbf{A}}$.

By application of Lemmas 3.4 and 3.3 it is enough to construct an admissibly ordered $\vec{\mathbf{B}}_0$ such that that every re-ordering $\vec{\mathbf{C}}_0 \in \mathcal{O}$ of $\vec{\mathbf{B}}_0$ with the following two properties contains a copy of $\vec{\mathbf{A}}$:

- (a) the order of vertices in distinct homologous components is preserved;
- (b) the order in $\vec{\mathbf{C}}_0$ on substructures which are closures of vertices depends only on their isomorphism type in $\vec{\mathbf{B}}_0$.

Given $\vec{\mathbf{A}}$ denote by $\vec{\mathbf{A}}_1, \vec{\mathbf{A}}_2, \dots, \vec{\mathbf{A}}_n$ all admissible re-orderings of $\vec{\mathbf{A}}$ having properties (a) and (b). So these structures all have the same domain, and only differ in their orderings. Put $\vec{\mathbf{B}}_0$ to be the disjoint union of $\vec{\mathbf{A}}_1, \vec{\mathbf{A}}_2, \dots, \vec{\mathbf{A}}_n$ with the order completed arbitrarily. Let $\vec{\mathbf{C}}_0$ be an admissible reordering of $\vec{\mathbf{B}}_0$ satisfying (a) and (b). Denote by $\alpha_i : \vec{\mathbf{A}} \rightarrow \vec{\mathbf{C}}_0$ the map which sends $a \in A$ to $a \in A_i$, for $i \leq n$. It is enough to prove the following.

Claim: For all $k \geq 0$, there is some $i \leq n$ such that the map α_i preserves the ordering on vertices in $\vec{\mathbf{A}}$ of level at most k .

We prove this by induction on k . Denote by $A|_k$ the set of vertices in A at level $\leq k$ (this is not necessarily a substructure or even a structure in \mathcal{K} , but it still makes sense to discuss admissible orderings of it as it contains the closures

of all of its vertices, and the criteria for being an admissible ordering depend only on these). By setting $A|_{-1} = \emptyset$, we can incorporate the proof for the base case $k = 0$ into the general argument.

Step 1: Every admissible ordering of vertices in $A|_{k-1}$ satisfying (a) and (b) extends to one of $A|_k$. If \mathbf{X} is the closure of a vertex of level k in \mathbf{A} , then any two such closure-extensions differ on \mathbf{X} by a permutation in $\text{Aut}(\mathbf{X}/X^\circ)$ (that is, automorphisms of \mathbf{X} fixing all vertices of level less than k).

Indeed, we need only to say how to define the ordering on \mathbf{X} . But, given the ordering on X° , this is determined by condition A6 in Definition 3.2, up to the action of $\text{Aut}(\mathbf{X}/X^\circ)$. It is easy to see that the resulting ordering on $A|_k$ is admissible.

Step 2: Suppose the claim holds up to level $k - 1$. Let I denote the set of $i \leq n$ for which α_i restricted to $A|_{k-1}$ is order-preserving. So this is non-empty. Let $i \in I$ and let \mathbf{X} be as in Step 1.

There is $\beta_i \in \text{Aut}(\mathbf{X}/X^\circ)$ such that $\alpha_i \circ \beta_i$ preserves the ordering on \mathbf{X} and so is an isomorphism between $\vec{\mathbf{X}}$, the structure on \mathbf{X} in $\vec{\mathbf{A}}$ and the substructure $\alpha_i(\mathbf{X})$ in $\vec{\mathbf{C}}_0$. By Step 1, there is some $j \leq n$ such that the map β_i^{-1} , regarded as a map from $\vec{\mathbf{A}}_i$ to $\vec{\mathbf{A}}_i$ (in $\vec{\mathbf{B}}_0$) is order-preserving and all vertices in $A|_{k-1}$ have the same ordering in $\vec{\mathbf{A}}_i$ and $\vec{\mathbf{A}}_j$. By condition (b), it follows that this map between the corresponding subsets of $\vec{\mathbf{C}}_0$ is order-preserving (as all orderings are determined by what happens in closures of vertices). Thus $j \in I$ and α_j is order-preserving on X . Repeating this argument for other vertices at level k , we complete the inductive step.

We have verified that the class of admissible orderings \mathcal{O} has the ordering property. The Ramsey property follows from Theorem 1.3 and A5: Given $\vec{\mathbf{A}}, \vec{\mathbf{B}} \in \mathcal{O}$ and $\vec{\mathbf{C}} \in \vec{\mathcal{K}}$ such that $\vec{\mathbf{C}} \rightarrow (\vec{\mathbf{B}})_{\vec{\mathbf{A}}}$. Construct a new order \leq of $\vec{\mathbf{C}}$ in a way that \leq agrees with $\leq_{\vec{\mathbf{C}}}$ on every copy of $\vec{\mathbf{B}}$ in $\vec{\mathbf{C}}$. Complete \leq so it satisfies A3 and A4. By A5 it follows that \mathbf{C} ordered by \leq is in \mathcal{O} . \square

4 Irreducible substructure faithful EPPA

The proof of Theorem 1.7 is a variant of the proof of clique-faithful EPPA by Hodkinson and Otto [17] strengthened for languages with unary functions. As in [17] our starting point is the following construction giving EPPA for relational structures and we verify coherency as in [34, 35].

Theorem 4.1 (Herwig [13], coherency verified by Solecki [37]). *For any relational language L , the class of all finite L -structures has coherent EPPA.*

To apply this construction to structures with functions we will temporarily interpret functions as relational symbols.

Definition 4.1. Suppose L is a language where all function symbols are unary. Given an L -structure \mathbf{A} we denote by \mathbf{A}^- its *relational reduct* constructed as follows. The language L^- of \mathbf{A}^- is a relational language containing all relational symbols of \mathbf{A} and additionally containing for every function symbol $F \in L_{\mathcal{F}}$ a relation symbol $R^F \in L_{\mathcal{R}}^-$ of arity 2. The vertex set of \mathbf{A} is the same as the vertex set of \mathbf{A}^- . For every $R \in L_{\mathcal{R}}$ we have $R_{\mathbf{A}} = R_{\mathbf{A}^-}$ and for every $F \in L_{\mathcal{F}}$ it holds that $(t_1, t_2) \in R_{\mathbf{A}^-}^F$ if and only if $t_2 \in F_{\mathbf{A}}(t_1)$.

Note that in the above, the structures \mathbf{A} and \mathbf{A}^- have the same automorphisms and any partial automorphism of \mathbf{A} is a partial automorphism of \mathbf{A}^- (but of course not conversely).

The proof of Theorem 1.7 will occupy the rest of this section.

Proof of Theorem 1.7. Given \mathbf{A} we invoke Theorem 4.1 to obtain a coherent EPPA-witness \mathbf{B}^- of \mathbf{A}^- (with respect to all partial isomorphisms of \mathbf{A}^-). We use the following terminology, following an exposition by Hodkinson in [16, 17]. The difference between our proof and that in [17] and [35] (which give clique faithful EPPA for structures in relational languages) is that whereas in these papers, the vertices in the EPPA-witness are of the form (v, χ_v) (as in the notation below), our vertices will actually be sets of such vertices (carrying an L -structure).

A set $S \subseteq B^-$ is called *small* if there is some $g \in \text{Aut}(\mathbf{B}^-)$ such that $g(S) \subseteq A$. Otherwise S is called *big*. Denote by \mathcal{U} the set of all big subsets of B^- and note that this is preserved by $\text{Aut}(\mathbf{B}^-)$. Given $b \in B^-$ a map $\chi : \mathcal{U} \rightarrow \mathbb{N}$ is a *b-valuation function* if $\chi(S) = 0$ for all $b \notin S \in \mathcal{U}$ and $1 \leq \chi(S) < |S|$ otherwise.

Given vertices $a, b \in B^-$ and their valuation functions, χ_a and χ_b we say that pairs (a, χ_a) and (b, χ_b) are *generic* if either $(a, \chi_a) = (b, \chi_b)$, or $a \neq b$ and for every $S \in \mathcal{U}$ such that $a, b \in S$ it holds that $\chi_a(S) \neq \chi_b(S)$.

The key construction in the proof is the following “local covering” construction of a *b-valuation L-structure* \mathbf{V}_b . Fix $b \in B^-$ and an automorphism $\alpha : B^- \rightarrow B^-$ such that $\alpha(b) \in \mathbf{A}^-$ (we can assume such an automorphism always exists — all other vertices can be removed from \mathbf{B}^-). Now consider the L -substructure $\mathbf{V}_b^\alpha = \text{Cl}_{\mathbf{A}}(\alpha(b))$ of \mathbf{A} . Suppose that for every $v \in \alpha^{-1}(V_b^\alpha)$ we have a v -valuation function χ_v such that the assigned valuation functions are generic for every pair of vertices in $\alpha^{-1}(V_b^\alpha)$. (Such a choice of valuation functions always exists and we will show how to obtain it later when we define an embedding ϕ of \mathbf{A} .) Denote by V_b the set of all such pairs (v, χ_v) , $v \in \alpha^{-1}(V_b^\alpha)$. On the set V_b we consider the L -structure \mathbf{V}_b , called a *b-valuation*, which is defined in such a way that the composition of mappings α and $\pi(v, \chi_v) = v$ forms an embedding $\alpha \circ \pi : \mathbf{V}_b \rightarrow \mathbf{V}_b^\alpha$. (This is a standard construction, we use the 1-1 mapping $\alpha \circ \pi$ to pull back the structure \mathbf{V}_b^α to V_b ; note that the isomorphism type of structure here does not depend on the choice of α .) Observe that then $\text{Cl}_{\mathbf{V}_b}((b, \chi_b)) = \mathbf{V}_b$.

Notice that for every b there are multiple choices of b -valuations \mathbf{V}_b (depending on particular choice of valuation functions assigned to vertices, but not depending on the choice of α). The sets V_b and structures \mathbf{V}_b will form a “cover of \mathbf{B}^- ” and we find it convenient to make the following definitions.

Definition 4.2. Recalling that all functions of L are unary, we say that a pair of valuations \mathbf{V}_a and \mathbf{V}_b is *generic* if

- (i) every pair of vertices $(u, \chi_u) \in V_a$ and $(v, \chi_v) \in V_b$ is generic;
- (ii) for every $(u, \chi_u) \in V_a$ and $(v, \chi_v) \in V_b$ and $F \in L_{\mathcal{F}}$ it holds that
 - (a) if $(u, v) \in R_{\mathbf{B}^-}^F$, then $(v, \chi_v) \in V_a$ and
 - (b) if $(v, u) \in R_{\mathbf{B}^-}^F$, then $(u, \chi_u) \in V_b$;
- (iii) if $(u, \chi_u) \in V_a \cap V_b$, then $\text{Cl}_{\mathbf{V}_a}((u, \chi_u)) = \text{Cl}_{\mathbf{V}_b}((u, \chi_u))$.

We also say that a set S of valuations is *generic* if every pair of valuations in S is generic.

Now we construct an L -structure \mathbf{C} :

1. The vertices of \mathbf{C} are all b -valuation L -structures \mathbf{V}_b , for $b \in B^-$.
2. For every relation $R \in L_{\mathcal{R}}$ put $(\mathbf{V}_{v_1}, \mathbf{V}_{v_2}, \dots, \mathbf{V}_{v_{\text{arity}(R)}}) \in R_{\mathbf{C}}$ if and only if $(v_1, v_2, \dots, v_{\text{arity}(R)}) \in R_{\mathbf{B}^-}$ and the set $\{\mathbf{V}_{v_i} : 1 \leq i \leq \text{arity}(R)\}$ is generic.
3. For every function $F \in L_{\mathcal{F}}$ put $F_{\mathbf{C}}(\mathbf{V}_{v_1}) = \{\mathbf{V}_{v_2}, \mathbf{V}_{v_3}, \dots, \mathbf{V}_{v_s}\}$ for some $s > 1$ if and only if
 - (a) $\mathbf{V}_2, \dots, \mathbf{V}_{s+1}$ are substructures of \mathbf{V}_1 ,
 - (b) $F_{\mathbf{V}_{v_1}}((v_1, \chi_1)) = \{(v_2, \chi_2), (v_3, \chi_3), \dots, (v_s, \chi_s)\}$ where for every $1 \leq i \leq s+1$ we denote by χ_i is the unique v_i -valuation such that $(v_i, \chi_i) \in V_{v_i}$,
 - (c) $(v_1, v_l) \in R_{\mathbf{B}^-}^F$ for every $2 \leq l \leq s+1$.
 - (d) the set $\{\mathbf{V}_{v_i} : 1 \leq i \leq s+1\}$ is generic.

Put $F_{\mathbf{C}}(\mathbf{V}_{v_1}) = \emptyset$ if the conditions above are not satisfied.

We give an embedding $\phi : \mathbf{A} \rightarrow \mathbf{C}$ with generic image. For every big set $S \in \mathcal{U}$ choose $f_S : S \rightarrow \{0, 1, 2, \dots, |S| - 1\}$ to be a function such that $f_S(v) > 0$ if and only if $v \in A \cap S$ and for every pair of vertices $u, v \in A \cap S$ it holds that $f_S(u) \neq f_S(v)$. Such a function exists because $A \cap S$ is always a proper subset of S . Given a vertex $a \in A$ we put $\phi(a)$ to be an a -valuation constructed from $\text{Cl}_{\mathbf{A}}(a)$ by mapping every vertex $v \in \text{Cl}_{\mathbf{A}}(a)$ to (v, χ_v) where $\chi_v(S) = f_S(v)$. It is easy to verify that this is indeed an embedding from \mathbf{A} to \mathbf{C} and $\phi(\mathbf{A})$ is generic.

We aim to show that \mathbf{C} is an EPPA-witness of $\phi(\mathbf{A})$. We first take time to introduce a terminology and prove a lemma which will allow us to use the fact that \mathbf{B}^- is an EPPA-extension of \mathbf{A}^- .

Denote by \mathcal{V} the union of all vertex sets of $\mathbf{V}_v \in \mathbf{C}$. If $g \in \text{Aut}(\mathbf{B}^-)$, we say that the partial map $q : \mathcal{V} \rightarrow \mathcal{V}$ is *g -compatible* if for all $(a, \chi) \in \text{Dom}(q)$ there exists a $g(a)$ -valuation function χ' such that $q((a, \chi)) = (g(a), \chi')$.

Similarly, let $g \in \text{Aut}(\mathbf{B}^-)$ and $p : \mathbf{C} \rightarrow \mathbf{C}$ be a partial automorphism. We say that p is *g -compatible* if there exists a g -compatible map $q : \mathcal{V} \rightarrow \mathcal{V}$ such that for all $\mathbf{V}_v \in \text{Dom}(p)$ q restricted to V_v is an isomorphism of \mathbf{V}_v and $p(\mathbf{V}_v)$.

Denote by π the homomorphism (called a *projection*) $\mathbf{C}^- \rightarrow \mathbf{B}^-$ defined by $\pi(\mathbf{V}_v) = v$.

Lemma 4.2. *Let $p : \mathbf{C} \rightarrow \mathbf{C}$ be a partial automorphism with generic domain and range, $g \in \text{Aut}(\mathbf{B}^-)$, and suppose that p is g -compatible. Then p extends to some g -compatible $\hat{p} \in \text{Aut}(\mathbf{C})$.*

Proof. As $\text{Dom}(p)$ is generic, for every $v \in \pi(\text{Dom}(p))$ there is precisely one v -valuation function χ_v such that the pair (v, χ_v) is a vertex of some valuation $V_b \in \text{Dom}(p)$. Denote by D the set of all such pairs (so $D = \bigcup \text{Dom}(p)$). The same is true for the range and denote by R all pairs appearing as vertices in

valuations in $p(\text{Dom}(p))$. It follows that p uniquely defines a g -compatible map $q : D \rightarrow R$. Fix a big set $S \in \mathcal{U}$. Then the set of pairs

$$\left\{ \left(\chi_b(S), \chi'_{g(b)}(g(S)) \right) : (b, \chi_b) \in D, q(b, \chi_b) = (g(b), \chi'_{g(b)}) \right\}$$

is the graph of a partial permutation of $\{0, 1, 2, 3, \dots, |S| - 1\}$ fixing 0 if defined on it. Extend it to a permutation θ_S^p of $\{0, 1, 2, 3, \dots, |S| - 1\}$ fixing 0.

Now we define $\hat{q} : \mathcal{V} \rightarrow \mathcal{V}$ by mapping $(b, \chi) \in \mathcal{V}$ to $(g(b), \chi')$ such that $\chi'(g(S)) = \theta_S^p(\chi(S))$ and $\hat{p} : \mathbf{V}_v \rightarrow \mathbf{V}_{g(v)}$ where $\mathbf{V}_{g(v)}$ is created from \mathbf{V}_v by mapping every vertex $(b, \chi) \in V_v$ to $\hat{q}(b, \chi)$.

It is easy to verify that \hat{q} is a well defined permutation of \mathcal{V} which extends q and is g -compatible. Moreover it preserves the relation of genericity between elements of \mathcal{V} . Therefore also \hat{p} is a well-defined permutation of C which preserves generic sets, extends p and is g -compatible. Consequently \hat{p} is an automorphism of \mathbf{C} . \square

By Lemma 4.2 the extension property of \mathbf{C} for partial isomorphisms of $\phi(\mathbf{A})$ follows easily. Let p be a partial isomorphism of $\phi(\mathbf{A})$. We extend it to $\hat{p} \in \text{Aut}(\mathbf{C})$. First extend $\phi^{-1} \circ p$ (which is an partial automorphism of \mathbf{A}) to automorphism $g \in \text{Aut}(\mathbf{B}^-)$. Clearly p is g -compatible and because domain and range are generic. By Lemma 4.2 p extends to g -compatible $\hat{p} \in \text{Aut}(\mathbf{C})$. This shows that \mathbf{C} is indeed an extension of $\phi(\mathbf{A})$.

For coherence, we use a similar argument to that in [34, 35]. Given a coherent triple (f_0, g_0, h_0) of partial automorphisms of \mathbf{A} we first extend this to a coherent triple (f, g, h) of automorphisms of \mathbf{B}^- . We let (f_1, g_1, h_1) be the coherent triple of partial automorphisms of $\phi(\mathbf{A})$ induced by (f_0, g_0, h_0) . Using Lemma 4.2 we extend f_1 to an f -compatible $\hat{f} \in \text{Aut}(\mathbf{C})$. Similarly we obtain extensions \hat{g}, \hat{h} of g, h . In order to ensure that the triple $(\hat{f}, \hat{g}, \hat{h})$ is coherent, we only need to ensure that, in the proof of Lemma 4.2, the permutations θ_S^p can be chosen coherently. More precisely, we want to ensure that $\theta_S^g \theta_S^f = \theta_S^h$. As in [35], if we extend any partial permutation α on $\{1, \dots, s\}$ to a permutation by mapping $\{1, \dots, s\} \setminus \text{Dom}(\alpha)$ to $\{1, \dots, s\} \setminus \alpha(\text{Dom}(\alpha))$ in an order-preserving way, then we obtain the required coherence.

Finally we verify that \mathbf{C} is faithful for irreducible substructures. Let \mathbf{D} be an irreducible substructure of \mathbf{C} . We first show that \mathbf{D} is generic. Suppose not and that $\mathbf{V}_a, \mathbf{V}_b \in D$ form a non-generic pair of vertices. Let $\mathbf{E}_a = \{\mathbf{V}_v \in D : \mathbf{V}_a \notin \text{Cl}_{\mathbf{D}}(\mathbf{V}_v)\}$. As closures are unary, this is a (proper) substructure of \mathbf{D} . Similarly define \mathbf{E}_b . Note that $\mathbf{E}_a \cup \mathbf{E}_b = \mathbf{D}$: otherwise, there is $\mathbf{V}_v \in \mathbf{D}$ with $\mathbf{V}_a, \mathbf{V}_b \in \text{Cl}_{\mathbf{D}}(\mathbf{V}_v)$ and then $\mathbf{V}_a, \mathbf{V}_b \subseteq \mathbf{V}_v$, so form a generic pair. Moreover, no relation of \mathbf{C} can involve a vertex $\mathbf{V}_u \in \mathbf{E}_a \setminus \mathbf{E}_b$ and a vertex $\mathbf{V}_v \in \mathbf{E}_b \setminus \mathbf{E}_a$ as $\mathbf{V}_b \subseteq \mathbf{V}_u$ and $\mathbf{V}_a \subseteq \mathbf{V}_v$, which implies that $\mathbf{V}_u, \mathbf{V}_v$ is not a generic pair. Thus \mathbf{D} is a free amalgam of the substructures \mathbf{E}_a and \mathbf{E}_b , which is a contradiction to its irreducibility. So \mathbf{D} is generic.

Because \mathbf{D} is generic it follows that $S = \pi(D)$ is small. Indeed, for each $u \in S$, there is a u -valuation χ_u such that the set of pairs $\{(u, \chi_u) : u \in S\}$ is generic. If S were big, this would imply that $\{\chi_u(S) : u \in S\}$ has size $|S|$, which is impossible (its elements j satisfy $1 \leq j < |S|$).

It follows that there is $g \in \text{Aut}(\mathbf{B}^-)$ such that $g(\pi(\mathbf{D})) \subseteq A$. The map $p : \mathbf{D} \rightarrow \phi(\mathbf{A})$ given by $p(\mathbf{V}_u) = \phi(g(u))$ is a g -compatible partial automorphism

of \mathbf{C} with generic domain and range. By Lemma 4.2, p extends to $\hat{p} \in \text{Aut}(\mathbf{C})$ and $\hat{p}(\mathbf{D}) \subseteq \phi(\mathbf{A})$. This completes the proof that \mathbf{C} is faithful for irreducible substructures. \square

Remark. The construction above adds an extra tool to the existing constructions of EPPA-witness and can be thus used as an additional layer in the construction of EPPA-witness for non-free amalgamation classes based on application of Herwig-Lascar theorem [14, 37, 32]. An example of such application is given in [3] giving EPPA for some classes of antipodal metric spaces.

5 Applications

In this section we discuss how some previously-studied classes of structures can be viewed naturally as free amalgamation classes of structures with set-valued functions.

Before doing this, we mention an alternative viewpoint for classes of structures with closures which we used in [11]. In the following examples we will show how this is related to our definitions.

Consider a class \mathcal{K} of finite L -structures, closed under isomorphisms, and a distinguished class \sqsubseteq of embeddings between elements of \mathcal{K} , called *strong embeddings*. We shall assume \sqsubseteq is closed under composition and contains all isomorphisms. In this case, we refer to $(\mathcal{K}; \sqsubseteq)$ as a *strong class*. If \mathbf{A} is a substructure of $\mathbf{B} \in \mathcal{K}$ and the inclusion map $\mathbf{A} \rightarrow \mathbf{B}$ is in \sqsubseteq , then we say that \mathbf{A} is a *strong substructure* of \mathbf{B} and write $\mathbf{A} \sqsubseteq \mathbf{B}$. In the other words, a strong class is a subcategory of \mathcal{K} with the strong embeddings.

The Ramsey property and amalgamation property can be defined analogously to the Ramsey property and amalgamation property of classes of L -structures, but considering only strong substructures and strong embedding. Most of the Fraïssé theory remains valid in this setting (see [11] for more details).

5.1 k -orientations

For a fixed natural number k , a *k -orientation* is an oriented (that is, directed) graph such that the out-degree of every vertex is at most k . We say that a substructure $G_1 = (V_1, E_1)$ of a k -orientation $G_2 = (V_2, E_2)$ is *successor closed* if there is no edge from V_1 to $V_2 \setminus V_1$ in G_2 .

Denote by \mathcal{D}_k the class of all finite k -orientations. This is a hereditary class closed for free amalgamation over successor-closed subgraphs and thus the successor-closedness plays the rôle of strong substructure, so \mathcal{D}_k can be considered as a class with corresponding strong embeddings. We show how to turn \mathcal{D}_k into a free amalgamation class in the sense of Definition 1.1.

Given an oriented graph $G = (V, E) \in \mathcal{D}_k$ denote by \mathbf{G}^+ the structure with vertex set V and (partial) unary function F . The function F maps every vertex to its out-neighborhood. Denote by \mathcal{D}_k^+ the class of all structures \mathbf{G}^+ for $G \in \mathcal{D}_k$. Because \mathbf{G}_1^+ is a substructure of \mathbf{G}_2^+ if and only if G_1 is successor closed in G_2 it follows that \mathcal{D}_k^+ is a free amalgamation class. We immediately obtain:




Figure 6: An amalgamation of $B_1, B_2 \in \mathcal{C}_2$ over A .

Theorem 5.1. *The class $\vec{\mathcal{D}}_k^+$ is Ramsey and there exists a class $\mathcal{O}_k \subseteq \vec{\mathcal{D}}_k^+$ with the ordering property (with respect to \mathcal{D}_k^+). The class \mathcal{D}_k^+ has the extension property for partial automorphisms.*

Let us briefly discuss what is the structure of \mathcal{O}_k . Given $\vec{\mathbf{A}} \in \mathcal{O}_k$, the closure-components (recall Definition 3.1) of $\vec{\mathbf{A}}$ corresponds to strongly connected components in the underlying oriented graph and $\vec{\mathbf{A}}$ is an ordered closure-extension of level 0 if and only if the underlying graph is strongly connected. More generally $\vec{\mathbf{A}}$ is an ordered closure-extension of level k if it contains a single strongly connected component C of level k and all other vertices of $\vec{\mathbf{A}}$ are reachable from C via an oriented path. Condition A6 of Definition 3.2 thus requires that the ordering of C is determined by the isomorphism type of \mathbf{A} (the underlying oriented graph) and the ordering of $A \setminus C$. Thus in \mathcal{O}_k , vertices are ordered primarily by the number of vertices in their closure. Every closure-component forms an interval where the order within this interval is fixed by the similarity type of corresponding closure-extension. The relative order of closure-components is given by their isomorphism type and the ordering of closure-components reachable from them. This can be seen as a generalization of the order of oriented forests described in Section 3.

Theorem 5.1 can be seen as the most elementary use of Theorems 1.3, 1.4, and 1.7, but it has important consequences. Denote by \mathcal{C}_k the undirected reducts of oriented graphs in \mathcal{D}_k , that is, the class of all unoriented graphs which can be oriented to an k -orientation. Given a graph $G = (V, E)$, its *predimension* is $\delta(G) = k|V| - |E|$. It is the heart of Hrushovski predimension construction that the class \mathcal{C}_k forms a free amalgamation class for the following notion of strong subgraph. Given a graph $G \in \mathcal{C}_k$ its subgraph H is a *self sufficient* or *strong subgraph* if for every subgraph H' of G containing H it holds that $\delta(H) \leq \delta(H')$.

The connection between the Hrushovski predimension construction and orientability follows by the Marriage Theorem and was first introduced in [9, 10]. Its consequences in Ramsey theory are the main topic of [11] and they are out of scope of this paper. We however point out why this free amalgamation class over strong subgraphs does not translate to a free amalgamation class when enriched by set-valued functions representing the smallest self-sufficient subgraph of a given set. Consider a graph in \mathcal{C}_2 created as amalgamation depicted in Figure 6. While in both B_1 and B_2 the vertices denoted by circles forms a self-sufficient substructures, it is not the case in the free amalgamation. The predimension of the 4 independent vertices is 8, while the predimension of the whole amalgam is 6. It follows that in order to represent self-sufficient substructures by means of set-valued functions, a new function from the vertices denoted by circles would need to be added. This makes the amalgamation non-




Figure 7: Fano plane (Steiner (2,3)-system).

free in our representation and this is the reason why additional information about orientation of the edges is needed.

5.2 Steiner systems

It was established in [4, 21, 20] that the class of finite partial Steiner systems is Ramsey with respect to strong subsystems. Moreover, the ordering property follows from techniques of [30]. We derive both results by a re-interpretation of partial Steiner systems as a free amalgamation class in a functional language.

This is an example where non-unary functions are necessary.

For fixed integers $r \geq t \geq 2$, by a *partial Steiner (r, t) -system* we mean an r -uniform hypergraph $G = (V, E)$ with the property that every t -element subset of V is contained in at most one edge of G . (If there is always exactly one such edge, we have a Steiner system.) Abusing terminology somewhat, we shall refer to this simply as a Steiner (r, t) -system. Given two Steiner (r, t) -systems G and H , we say that G is a *strongly induced subsystem* of H if

1. G is an induced subhypergraph of H ; and,
2. every hyperedge of H which is not a hyperedge of G intersects G in at most $t - 1$ vertices.

In [4] the Ramsey property was formulated with respect to strongly induced subsystems. We, equivalently, use set-valued functions to represent this.

Definition 5.1. Denote by $\mathcal{S}_{r,t}$ the class of all finite structures \mathbf{A} with one function F from t -tuples with the following properties:

1. If $F_{\mathbf{A}}(\vec{x}) \neq \emptyset$ then \vec{x} has no repeated vertices, $|F_{\mathbf{A}}(\vec{x})| = r$ and every vertex of \vec{x} is in $F_{\mathbf{A}}(\vec{x})$.
2. Whenever \vec{x} is an t -tuple of vertices of \mathbf{A} , \vec{x}_2 is an t -tuple consisting of distinct vertices in $F_{\mathbf{A}}(\vec{x})$ it holds that $F_{\mathbf{A}}(\vec{x}) = F_{\mathbf{A}}(\vec{x}_2)$.

It is easy to see that $\mathcal{S}_{r,t}$ is a free amalgamation class.

Given a Steiner (r, t) -system $G = (V, E)$ we can interpret it as a structure $\mathbf{S}_G \in \mathcal{S}_{r,t}$ with vertex set V and function $F(\vec{x})$ defined for every t -tuple \vec{x} of distinct vertices such that there is hyperedge $A \in E$ containing all vertices of \vec{x} . In this case we put $F(\vec{x}) = A$.

Observe that if G is a strong subsystem of H if and only if \mathbf{S}_G is a substructure of \mathbf{S}_H . It follows that Steiner (r, t) -systems are in 1-1 correspondence to




Figure 8: The bowtie graph.

structures \mathbf{S}_G and moreover this correspondence maps subsystems to substructures.

We obtain an alternative proof of the following main result of [4]:

Theorem 5.2. *The class $\vec{\mathcal{S}}_{r,t}$ is a Ramsey class with the ordering property.*

Proof. The Ramsey property follows directly from Theorem 1.3. For the ordering property, note that single vertices are closed in structures in $\mathcal{S}_{r,t}$, so all orderings are admissible, in the sense of Theorem 1.4. \square

In [20] we obtained further corollaries to this approach. We remark that EPPA is presently open for the class of partial Steiner systems. Of course, our results in Section 4 does not apply in this case, as the function introduced is not unary.

5.3 Bowtie-free graphs

A *bowtie* B is a graph consisting of two triangles with one vertex identified (see Figure 8). A graph G is *bowtie-free* if there is no monomorphism from B to G . The existence of a universal graph in the class of all countable bowtie-free graphs was shown in [24]. The paper [7] gave a far reaching generalization by giving a condition for the existence of ω -categorical universal graphs for classes defined by forbidden monomorphisms (which we refer to as to as *Cherlin-Shelah-Shi classes*). This led to several new classes being identified [8, 6, 5]. Bowtie-free graphs represent a key example of a class that is not a free amalgamation class by itself, but can be turned into one by means of unary functions. This analysis was carried in [19] where we gave an explicit characterisation of the ultrahomogeneous lift and the Ramseyness of this lift. The presentation can be greatly simplified by considering structures with set-valued functions and moreover we show the extension property for partial automorphisms (See also [34] for related results on ample generics).

While not all Cherlin-Shelah-Shi classes give rise to free amalgamation classes (see the more detailed analysis in [21]), what follows can be generalized to many of the other block-path examples given by [6] and [5].

We review the main observations about the structure of bowtie-free graphs from [19]. For completeness we include the (easy) proofs.

Definition 5.2 (Chimneys). For $n \geq 2$, an *n-chimney graph*, Ch_n , is a free amalgamation of n triangles over one common edge. A *chimney graph* is any graph Ch_n for some $n \geq 2$.

Chimneys together with K_4 (a clique on 4 vertices) will form the only components of bowtie-free graphs formed by triangles. The assumption $n \geq 2$ for chimney is a technical assumption to avoid isolated triangles. Note also that Ch_2 is not an induced subgraph of K_4 .

Definition 5.3 (Good bowtie-free graphs). A bowtie-free graph $G = (V, E)$ is *good* if every vertex is contained either in a copy of chimney or a copy of the complete graph K_4 .

The structure of bowtie-free graphs is captured by means of the following three lemmas:

Lemma 5.3 ([19]). *Every bowtie-free graph G is an induced subgraph of some good bowtie-free graph G' .*

Proof. The graph G can be extended in the following way:

1. For every vertex v not contained in a triangle add a new copy of Ch_2 and identify the vertex v with one of the vertices of Ch_2 .
2. For every triangle v_1, v_2, v_3 that is not part of a 2-chimney nor a K_4 , add a new vertex v_4 and the triangle v_1, v_2, v_4 turning the original triangle into Ch_2 .

It is easy to see that step 1 cannot introduce a new bowtie.

Assume, to the contrary, that step 2 introduced a new bowtie. Further assume that v_1 is the unique vertex of degree 4 of this new bowtie and consequently there is another triangle on vertex v_1 in G . Because G is bowtie-free, this triangle must share a common edge with the triangle v_1, v_2, v_3 and therefore the triangle v_1, v_2, v_3 is already part of a K_4 or a 2-chimney in the original graph G . A contradiction. \square

Now we are ready to describe how to turn the class of bowtie-free graphs into a free amalgamation class. Our language L will consist of one binary relation R and unary function F .

Definition 5.4. For every good bowtie-free graph $G = (V, E)$ denote by \mathbf{G}^+ the L -structure with vertex set V and relations and functions defined as follows:

1. $(u, v) \in R_{\mathbf{G}}$ if and only if $\{u, v\}$ is an edge of G .
2. $F_{\mathbf{G}}(v) = u$ if and only if $\{u, v\}$ is contained in at least two triangles of a chimney.
3. $F_{\mathbf{G}}(v) = \{u_1, u_2\}$ if and only if $\{v, u_1, u_2\}$ is a triangle of a chimney and v is not contained in multiple triangles.
4. $F_{\mathbf{G}}(v) = \{u_1, u_2, u_3\}$ if and only if $\{v, u_1, u_2, u_3\}$ forms a 4-clique in G .

Denote by \mathcal{B} the class of all \mathbf{A} -partite substructures of structures \mathbf{G}^+ where G is a good bowtie-free graph.

Theorem 5.4. *\mathcal{B} is a free amalgamation class.*

Proof. Let $\mathbf{A}, \mathbf{B}, \mathbf{B}' \in \mathcal{B}$. Assume that \mathbf{A} is a substructure of both \mathbf{B} and \mathbf{B}' . We show that the free amalgamation \mathbf{C} of \mathbf{B}_1 and \mathbf{B}_2 over \mathbf{A} is in \mathcal{B} .

There are good bowtie free graphs G_1 and G_2 such that $\mathbf{B} \subseteq \mathbf{G}_1^+$ and $\mathbf{B}' \subseteq \mathbf{G}_2^+$. We claim that that the free amalgamation H of G_1 and G_2 over A is a good bowtie-free graph. Consequently \mathbf{H}^+ (given by Definition 5.4) is the free amalgam of \mathbf{G}_1^+ and \mathbf{G}_2^+ over \mathbf{A} . As \mathbf{C} is a substructure of \mathbf{H}^+ , it then follows that $\mathbf{C} \in \mathcal{B}$.




Figure 9: All isomorphism types of closure-extensions in \mathcal{B} .

Because \mathbf{A} is a substructure of both \mathbf{G}_1^+ and \mathbf{G}_2^+ , the function F ensure that the free amalgamation preserves the structure of chimneys: if a vertex of a chimney in G_1 is identified with a vertex of a chimney in G_2 (because it is in A) then also the bases (i.e. the edges in multiple triangles) of these chimneys are contained in A , so are identified in H and the result is again a chimney. Similarly F makes sure that a 4-clique containing a vertex of A is in A . Finally free amalgamation cannot introduce any new triangles and thus the free amalgamation is a good bowtie-free graph H . Consequently \mathbf{H}^+ is the free amalgam of \mathbf{G}_1^+ and \mathbf{G}_2^+ over \mathbf{A} . \square

Corollary 5.5. *The class \mathcal{B} has irreducible-structure faithful EPPA; $\vec{\mathcal{B}}$ is a Ramsey class and there is $\mathcal{B}' \subseteq \vec{\mathcal{B}}$ with the ordering property (with respect to \mathcal{B}).*

The class \mathcal{B}' can be easily derived from the Definition 3.2. There are only three types of closure-extensions in \mathcal{B} depicted in Figure 9 (with arrows representing function F and circles denoting the vertices of maximal level). It follows that vertices are ordered by size of their closures (here we make use of the definition of \preceq refining the order given by number of vertices). That is vertices in bases of chimney are first, vertices in the top of chimneys next and vertices in 4-cliques last. Every vertex-closure forms an interval. Pair of vertices of level 1 (in the top of chimneys) form homologous extensions if and only if they belong to the same chimney. It follows that for every chimney the set of its top vertices forms an interval and the relative order of these intervals corresponds to the relative order of corresponding bases.

Remark. The Ramsey property and an explicit description of the admissible ordering was given in [19]. The relational language used there is however more complicated and does not preserve all automorphisms of the Fraïssé limit of \mathcal{B} . This makes it unsuitable for the extension property for partial automorphisms. The formulation here is a more optimized version.

The EPPA for bowtie-free graphs was states as a problem in Siniora's thesis [34] and is attributed to Macpherson.

The argument above together with the observation that in the Fraïssé limit \mathbf{B} of \mathcal{B} we have that for every finite $S \subseteq B$, $|\text{Cl}_{\mathbf{B}}(S)| \leq 3|S|$, also gives a compact proof for the existence of an ω -categorical countable universal bowtie-free graph. This bound follows from the fact that function F cannot cascade. The ω -categoricity follows from the fact that the orbit of S in $\text{Aut}(\mathbf{B})$ is fully determined by the isomorphism type of $\text{Cl}_{\mathbf{B}}(S)$ and there are only finitely many closures for every finite S . This, of course, is just a re-formulation of the argument in [7].

6 Concluding remarks

1. It would be interesting to extend Theorem 1.7 to a class of structures which include non-unary functions. Perhaps this is too much to ask as EPPA is presently open even in the case of Steiner triple systems (as we remark in Section 5.2). However note that our structures involve set-valued functions and thus the EPPA may be easier to prove. However even for partial triple systems the EPPA seems to be presently open.

2. On the structural Ramsey theory side open problems include Ramsey properties of finite lattices and other algebraic structures where the axioms (such as associativity) are difficult to control in an amalgamation procedure. See [21, 3] for results on Ramsey classes.

Acknowledgements: We would like to thank to Daoud Siniora for several useful discussions concerning clique faithful EPPA and the notion of coherency. We are also grateful to the anonymous referee or detailed report which improved presentation of this paper.

References

- [1] Fred G. Abramson and Leo A. Harrington. Models without indiscernibles. *Journal of Symbolic Logic*, 43:572–600, 1978.
- [2] Omer Angel, Alexander S Kechris, and Russell Lyons. Random orderings and unique ergodicity of automorphism groups. *Journal of the European Math. Society*, 16:2059–2095, 2014.
- [3] Andres Aranda, David Bradley-Williams, Jan Hubička, Miltiadis Karamanlis, Michael Kompatscher, Matěj Konečný, and Micheal Pawliuk. Ramsey expansions of metrically homogeneous graphs. Submitted, arXiv:1707.02612, 2017.
- [4] Vindya Bhat, Jaroslav Nešetřil, Christian Reiher, and Vojtěch Rödl. A Ramsey class for Steiner systems. *Journal of Combinatorial Theory, Series A*, 154:323 – 349, 2018.
- [5] Gregory Cherlin and Saharon Shelah. Universal graphs with one forbidden subgraph: the generic case. In preparation.
- [6] Gregory Cherlin and Saharon Shelah. Universal graphs with a forbidden subgraph: block path solidity. *Combinatorica*, pages 1–16, 2013.
- [7] Gregory Cherlin, Saharon Shelah, and Niandong Shi. Universal graphs with forbidden subgraphs and algebraic closure. *Advances in Applied Mathematics*, 22(4):454–491, 1999.
- [8] Gregory Cherlin and Lasse Tallgren. Universal graphs with a forbidden near-path or 2-bouquet. *Journal of Graph Theory*, 56(1):41–63, 2007.
- [9] David M. Evans. Ample dividing. *The Journal of Symbolic Logic*, 68(04):1385–1402, 2003.

- [10] David M. Evans. Trivial stable structures with non-trivial reducts. *Journal of the London Mathematical Society (2)*, 72(2):351–363, 2005.
- [11] David M. Evans, Jan Hubička, and Jaroslav Nešetřil. Automorphism groups and Ramsey properties of sparse graphs. *Proceedings of the London Mathematical Society*, 119(2):515–546, 2019.
- [12] Alfred W. Hales and Robert I. Jewett. Regularity and positional games. *Transactions of the American Mathematical Society*, 106:222–229, 1963.
- [13] Bernhard Herwig. Extending partial isomorphisms on finite structures. *Combinatorica*, 15(3):365–371, 1995.
- [14] Bernhard Herwig and Daniel Lascar. Extending partial automorphisms and the profinite topology on free groups. *Transactions of the American Mathematical Society*, 352(5):1985–2021, 2000.
- [15] Wilfrid Hodges. *Model theory*, volume 42. Cambridge University Press, 1993.
- [16] Ian Hodkinson. Finite model property for guarded fragments. slides available at <http://www.cllc.vuw.ac.nz/LandCtalks/imhslides.pdf>, 2002.
- [17] Ian Hodkinson and Martin Otto. Finite conformal hypergraph covers and Gaifman cliques in finite structures. *Bulletin of Symbolic Logic*, 9(03):387–405, 2003.
- [18] Ehud Hrushovski. Extending partial isomorphisms of graphs. *Combinatorica*, 12(4):411–416, 1992.
- [19] Jan Hubička and Jaroslav Nešetřil. Bowtie-free graphs have a Ramsey lift. *Advances in Applied Mathematics*, 96:286 – 311, 2018.
- [20] Jan Hubička and Jaroslav Nešetřil. Ramsey classes with closure operations (selected combinatorial applications). In Steve Butler, Joshua Cooper, and Glenn Hurlbert, editors, *Connections in Discrete Mathematics: A Celebration of the Work of Ron Graham*, page 240–258. Cambridge University Press, 2018.
- [21] Jan Hubička and Jaroslav Nešetřil. All those Ramsey classes (Ramsey classes with closures and forbidden homomorphisms). *Advances in Mathematics*, 356C:106791, 2019.
- [22] Alexander S. Kechris, Vladimir G. Pestov, and Stevo Todorčević. Fraïssé limits, Ramsey theory, and topological dynamics of automorphism groups. *Geometric and Functional Analysis*, 15(1):106–189, 2005.
- [23] Alexander S. Kechris and Christian Rosendal. Turbulence, amalgamation, and generic automorphisms of homogeneous structures. *Proceedings of the London Mathematical Society*, 94(2):302–350, 2007.
- [24] Péter Komjáth. Some remarks on universal graphs. *Discrete Mathematics*, 199(1):259–265, 1999.

- [25] Jaroslav Nešetřil. Ramsey classes and homogeneous structures. *Combinatorics, probability and computing*, 14(1-2):171–189, 2005.
- [26] Jaroslav Nešetřil. Metric spaces are Ramsey. *European Journal of Combinatorics*, 28(1):457–468, 2007.
- [27] Jaroslav Nešetřil and Vojtěch Rödl. A structural generalization of the Ramsey theorem. *Bulletin of the American Mathematical Society*, 83(1):127–128, 1977.
- [28] Jaroslav Nešetřil and Vojtěch Rödl. On a probabilistic graph-theoretical method. *Proceedings of the American Mathematical Society*, 72(2):417–421, 1978.
- [29] Jaroslav Nešetřil and Vojtěch Rödl. The partite construction and Ramsey set systems. *Discrete Mathematics*, 75(1):327–334, 1989.
- [30] Jaroslav Nešetřil and Vojtěch Rödl. Partitions of subgraphs. In Miroslav Fiedler, editor, *Recent advances in graph theory*, pages 413–423. Prague, 1975.
- [31] Jaroslav Nešetřil and Vojtěch Rödl. Partitions of finite relational and set systems. *Journal Combinatorial Theory, Series A*, 22(3):289–312, 1977.
- [32] Martin Otto. Finite groupoids, finite coverings and symmetries in finite structures. arXiv:1404.4599, 2014.
- [33] Micheal Pawliuk and Miodrag Sokić. Amenability and unique ergodicity of automorphism groups of countable homogeneous directed graphs. *Ergodic Theory and Dynamical Systems*, page 1–51, 2018.
- [34] Daoud Siniora. *Automorphism Groups of Homogeneous Structures*. PhD thesis, University of Leeds, March 2017.
- [35] Daoud Siniora and Sławomir Solecki. Coherent extension of partial automorphisms, free amalgamation, and automorphism groups. *The Journal of Symbolic Logic*, 2019.
- [36] Sławomir Solecki. Extending partial isometries. *Israel Journal of Mathematics*, 150(1):315–331, 2005.
- [37] Sławomir Solecki. Notes on a strengthening of the Herwig–Lascar extension theorem. Unpublished note, 2009.
- [38] Anatoly M. Vershik. Globalization of the partial isometries of metric spaces and local approximation of the group of isometries of Urysohn space. *Topology and its Applications*, 155(14):1618–1626, 2008.