

ESTIMATES OF PRODUCTS OF NORMS FOR FLOWS AWAY FROM HOMOCLINIC TANGENCY

QIANYING XIAO

ABSTRACT. We give a detailed proof for the estimates of products of norms for flows away from homoclinic tangency. The estimates we can prove is weaker than the expectation of experts.

1. **Introduction.** Assume $X \in \mathcal{X}^1(M)$ is away from homoclinic tangencies. ϕ_t is the flow generated by X , ψ_t is the linear Poincaré flow. There exists a neighborhood \mathcal{U} of X such that any $Y \in \mathcal{U}$ has no homoclinic tangencies.

Wen [2] shows that there exist $\mathcal{U}_2 \subset \mathcal{U}$, $\gamma > 0$, $N > 0$, $\lambda > 0$, $\delta > 0$, and $T > 0$, any $Y \in \mathcal{U}$, any periodic orbits $orb(x)$ of Y with period $\tau > T$,

1. $\psi_\tau(x)$ has at most one eigenvalue with modulo in $[(1+\delta)^{-\tau}, (1+\delta)^\tau]$, i.e. there exists eigenspace splitting $\mathcal{N}_x = V^s(x) + V^c(x) + V^u(x)$ such that $V^c(x)$ is the eigenspace corresponding to the eigenvalue with modulo in $[(1+\delta)^{-\tau}, (1+\delta)^\tau]$, and $\dim V^c(x) \leq 1$.
2. $\angle(V^s(x), V^c(x) + V^u(x)) > \gamma$, $\angle(V^s(x) + V^c(x), V^u(x)) > \gamma$.
3. The splitting is dominated:

$$\|\psi_t|V^s(x)\| \|\psi_{-t}|V^c(\phi_t(x)) + V^u(\phi_t(x))\| \leq Ne^{-\lambda t},$$

$$\|\psi_t|V^s(x) + V^c(x)\| \|\psi_{-t}|V^u(\phi_t(x))\| \leq Ne^{-\lambda t}.$$

4. Estimation of norms of products:

$$\|\psi_\tau|V^s(x)\| \leq N(1+\delta)^{-\tau},$$

$$\|\psi_{-\tau}|V^u(x)\| \leq N(1+\delta)^{-\tau}.$$

The above estimates are improved by Wen [3]. In [3, Lemma 3.4], Wen gives the estimates of products of norms, which are stronger than the previous result. Wen regards the proofs are standard in Liao's and Mane's work(for instance see [1, Page 528]), and he doesn't give details of the proofs there. We try to prove it as an exercise, and it turns out we are not able to prove the Item 2 of [3, Lemma 3.4]. Instead we prove a weaker version, and it works for our purpose to prove Theorem 2.1 of our work [4]. The following ideas of Liao: minimal nonhyperbolic set, (local) star flows and the Selecting lemma, which are stressed by Wen extensively, help us to avoid a direct use of the non-proved estimates.

Proposition 1. *There exists neighborhood $\mathcal{U}_3 \subset \mathcal{U}_2$ of X , $0 < \lambda_1 < 1$, $T' > 0$, any $Y \in \mathcal{U}_3$, any periodic point x of Y with period $\tau > \max\{T, T'\}$, any partition of $[0, \tau]$:*

$$0 = t_0 < t_1 < \cdots < t_\ell = \tau$$

such that $t_{i+1} - t_i \geq T'$ for $0 \leq i \leq \ell - 1$, the following are satisfied: either

$$\prod_{i=0}^{\ell-1} \|\psi_{t_{i+1}-t_i} |V^s(\phi_{t_i})\| \leq N\lambda_1^\tau,$$

or

$$\prod_{i=0}^{\ell-1} \|\psi_{t_i-t_{i+1}} |V^s(\phi_{t_{i+1}})\| \leq N\lambda_1^\tau,$$

2. The proofs. Let us give the details of estimating the products of norms. For the sake of completeness, let us insert a well-known fact in linear algebra. Let E^n denotes the n -dimensional Euclidean space. $A : E^n \rightarrow E^n$ is a linear map.

Lemma 2.1. [1, Page 528] *Any $\epsilon > 0$, $v \in E^n$ with $\|v\| = 1$, there exist $Q : E^n \rightarrow E^n$ such that $\|Q - \text{id}\| \leq \epsilon$, and that $\|A \circ Q(v)\| \geq \epsilon n^{-\frac{3}{2}} \|A\|$.*

Proof. Let $\{e_1, \dots, e_n\}$ be an orthonormal basis, $v = (v_1, \dots, v_n)$. We may as well assume $|v_1| \geq n^{-\frac{1}{2}} \|v\|$.

$a_{ji} = (Ae_i)_j$ such that

$$\|A\| \leq n^{\frac{1}{2}} \|Ae_i\| \leq n|a_{ji}|$$

$Q : E^n \rightarrow E^n$ such that $Q(e_1) = e_1 + \epsilon t e_i$, $t = 1$ or -1 , and $Q(e_k) = e_k$ for $k > 1$. Obviously $\|Q - \text{id}\| \leq \epsilon$.

$$\|A \circ Q(v)\| \geq |(A \circ Q(v))_j| = |(A(v))_j + \epsilon t a_{ji} v_1|.$$

Choose t such that $(A(v))_j + \epsilon t a_{ji} v_1 \geq 0$, then

$$\|A \circ Q(v)\| \geq \epsilon |a_{ji} v_1| \geq \epsilon n^{-\frac{3}{2}} \|A\|.$$

□

Proof of the Proposition. Let $\epsilon > 0$, $\mathcal{U}_3 \subset \mathcal{U}_2$ be determined by Frank's lemma. K is an upperbound of $\|\psi_t^Y\|$ for $Y \in \mathcal{U}_3$, $1 \leq t \leq 2$.

$\epsilon' = \epsilon K^{-1}$, $\epsilon_1 > 0$ such that

$$\frac{1+\gamma}{\gamma} 2\epsilon_1 < \epsilon'.$$

$$T' > \frac{\frac{3}{2} \ln n - \ln \epsilon_1}{\ln(1 + \frac{1}{2}\delta)}.$$

Given any $Y \in \mathcal{U}_3$, any periodic point x of Y with periodic $\tau \geq \max\{T', T\}$, any partition of $[0, \tau]$:

$$0 = t_0 < t_1 < \dots < t_\ell = \tau,$$

such that $t_{i+1} - t_i \geq T'$ for $0 \leq i \leq \ell - 1$.

Case 1 Assume $\dim V^c(x) = 1$. Denote $\mathcal{N}_{\phi_{t_i}(x)}$ as N_i , $V_{\phi_{t_i}(x)}^\sigma$ as V_i^σ for $\sigma = s, c, u$, $i = 0, \dots, \ell - 1$.

Define $Q_1 : N_1 \rightarrow N_1$ such that $Q_1|V_1^c + V_1^u = \text{id}$, $Q_1(V_1^s) = V_1^s$, $\|Q_1|V_1^s - \text{id}\| < \epsilon_1$, $v_1 \in V_1^s$ such that $\|v_1\| = \|\psi_{t_1}|V^s(x)\|$, and

$$\|\psi_{t_2-t_1} \circ Q_1(v_1)\| \geq \epsilon_1 n^{-\frac{3}{2}} \|\psi_{t_2-t_1}|V_1^s\| \|\psi_{t_1}|V^s(x)\|.$$

Moreover, $\|Q_1 - \text{id}\| < \frac{1+\gamma}{\gamma} \epsilon_1 < \epsilon'$.

Similarly, we can define $Q_2 : N_2 \rightarrow N_2$ such that $\|Q_2 - \text{id}\| < \epsilon'$, $Q_2|V_2^c + V_2^u = \text{id}$, $Q_2(V_2^s) = V_2^s$, and that

$$\begin{aligned} & \|\psi_{t_3-t_2} \circ Q_2 \circ \psi_{t_2-t_1} \circ Q_1 \circ \psi_{t_1}|V^s(x)\| \\ & \geq (\epsilon_1 n^{-\frac{3}{2}})^2 \|\psi_{t_3-t_2}|V_1^s\| \|\psi_{t_2-t_1}|V_1^s\| \|\psi_{t_1}|V^s(x)\|. \end{aligned}$$

.....

$Q_{\ell-1} : N_{\ell-1} \rightarrow N_{\ell-1}$, such that $\|Q_{\ell-1} - \text{id}\| < \epsilon'$, $Q_{\ell-1}|V_{\ell-1}^c + V_{\ell-1}^u = \text{id}$, $Q_{\ell-1}(V_{\ell-1}^s) = V_{\ell-1}^s$.

$$\|(\prod_{i=1}^{\ell-1} \psi_{t_{i+1}-t_i}|V_i^s \circ Q_i) \circ \psi_{t_1}|V^s(x)\| \quad (1)$$

$$\geq (\epsilon_1 n^{-\frac{3}{2}})^{\ell-1} \prod_{i=0}^{\ell-1} \|\psi_{t_{i+1}-t_i}|V_i^s\| \quad (2)$$

$$\geq (\epsilon_1 n^{-\frac{3}{2}})^{\frac{\tau}{T'}} \prod_{i=0}^{\ell-1} \|\psi_{t_{i+1}-t_i}|V_i^s\| \quad (3)$$

Denote $L_{i,r} = \psi_{t_{i+1}-t_i} \circ (\text{id} + r(Q_i - \text{id}))$ for $i = 1, \dots, \ell-1$, $L_r = L_{\ell-1,r} \circ \dots \circ L_{1,r} \circ L_{0,r}$ ($0 \leq r \leq 1$).

Claim. *The spectral radius of $L_1|V^s(x)$ is less than $(1+\delta)^{-\tau}$. Otherwise, for some r , L_r admits at least two eigenvalues with modulo in $[(1+\delta)^{-\tau}, (1+\delta)^\tau]$. By Frank's lemma, there exists $Z \in \mathcal{U}_2$ such that $\psi_\tau^Z(x) = L_r$, a contradiction.*

Apply Frank's lemma, there exist $Z' \in \mathcal{U}_2$ such that $Z' = Y$ along $\text{orb}(x)$, $\psi_\tau^{Z'}(p) = L_1$, $V_{Z'}^\sigma(x) = V^\sigma(x)$, $\sigma = s, c, u$. Hence

$$\|L_1|V^s(x)\| \leq N(1+\delta)^{-\tau}$$

According to inequality 1, and by $T' > \frac{\frac{3}{2} \ln n - \ln \epsilon_1}{\ln(1+\frac{1}{2}\delta)}$,

$$\prod_{i=0}^{\ell-1} \|\psi_{t_{i+1}-t_i}|V_i^s\| \leq N[(1+\delta)^{-1}(\epsilon_1 n^{-\frac{3}{2}})]^{-\frac{1}{T'}} < N\left(\frac{1+\frac{1}{2}\delta}{1+\delta}\right)^\tau$$

Let $\lambda_1 = \frac{1+\frac{1}{2}\delta}{1+\delta}$, and we are done.

Case 2 Assume $\dim V^c(x) = 0$. As in case 1, we have Q_i for $i = 1, \dots, \ell-1$, such that $Q_i|V_i^u = \text{id}$, $Q_i(V_i^s) = V_i^s$, and $\|Q_i|V_i^s - \text{id}\| < \epsilon_1$. Moreover,

$$\begin{aligned} & \|\prod_{i=1}^{\ell-1} \psi_{t_{i+1}-t_i}|V_i^s \circ Q_i) \circ \psi_{t_1}|V^s(x)\| \\ & \geq (\epsilon_1 n^{-\frac{3}{2}})^{\frac{\tau}{T'}} \prod_{i=0}^{\ell-1} \|\psi_{t_{i+1}-t_i}|V_i^s\| \end{aligned}$$

Similarly, we have P_i for $i = 1, \dots, \ell-1$, such that $P_i|V_i^s = \text{id}$, $P_i(V_i^u) = V_i^u$, and $\|P_i|V_i^u - \text{id}\| < \epsilon_1$, and

$$\|\psi_{-t_1} \circ P_1 \circ \dots \circ \psi_{t_{\ell-2}-t_{\ell-1}} \circ P_{\ell-1} \circ \psi_{t_{\ell-1}-t_\ell}|V^u(x)\|$$

$$\geq (\epsilon_1 n^{-\frac{3}{2}})^{\frac{\tau}{T'}} \prod_{i=0}^{\ell-1} \|\psi_{t_i-t_{i+1}}|V_{i+1}^u\|,$$

$$\|P_1^{-1} \circ Q_i - \text{id}\| \leq \frac{1+\gamma}{\gamma} 2\epsilon_1 < \epsilon'.$$

For $0 \leq \alpha \leq 1$, $0 \leq \beta \leq 1$, define $L_{i,\alpha,\beta} = \psi_{t_{i+1}-t_i} \circ (\text{id} + \alpha(P_i^{-1} - \text{id})) \circ (\text{id} + \beta(Q_i - \text{id}))$.

Denote $L_{\alpha,\beta} = L_{\ell-1,\alpha,\beta} \circ \cdots \circ L_{0,\alpha,\beta}$.

Claim. *Either the spectral radius of $L_{1,1}|V^s(x)$ is less than $(1+\delta)^{-\tau}$, or the spectral radius of $L_{1,1}|V^u(x)$ is greater than $(1+\delta)^\tau$. Otherwise some $L_{\alpha,\beta}$ admits two eigenvalues (counting multiplicity) with modulo in $[(1+\delta)^{-\tau}, (1+\delta)^\tau]$. Apply Frank's lemma, there exists $Z \in \mathcal{U}_2$ such that $\psi_\tau^Z(x)$ has two eigenvalues with modulo in $[(1+\delta)^{-\tau}, (1+\delta)^\tau]$, a contradiction.*

By Frank's lemma, there exists $Z' \in \mathcal{U}_2$, such that $\psi_\tau^{Z'}(x) = L_{1,1}$, therefore $\|L_{1,1}|V^s(x)\| \leq N(1+\delta)^{-\tau}$ or $\|L_{1,1}^{-1}|V^u(x)\| \leq N(1+\delta)^{-\tau}$.

Note that $L_{1,1}|V^s(x) = (\prod_{i=0}^{\ell-1} \psi_{t_{i+1}-t_i} \circ Q_i)|V^s(x)$, $L_{1,1}^{-1}|V^u(x) = \psi_{-t_1} \circ \prod_{i=\ell-1}^1 P_i \circ \psi_{t_i-t_{i+1}}|V^u(x)$.

Hence

$$(\epsilon_1 n^{-\frac{3}{2}})^{\frac{\tau}{T'}} \prod_{i=0}^{\ell-1} \|\psi_{t_{i+1}-t_i}|V_i^s\| \leq N(1+\delta)^{-\tau},$$

or

$$(\epsilon_1 n^{-\frac{3}{2}})^{\frac{\tau}{T'}} \prod_{i=0}^{\ell-1} \|\psi_{t_i-t_{i+1}}|V_{i+1}^u\| \leq N(1+\delta)^{-\tau}.$$

Let $\lambda_1 = \frac{1+\frac{1}{2}\delta}{1+\delta}$, then

$$\prod_{i=0}^{\ell-1} \|\psi_{t_{i+1}-t_i}|V_i^s\| \leq N\lambda_1^\tau,$$

or

$$\prod_{i=0}^{\ell-1} \|\psi_{t_i-t_{i+1}}|V_{i+1}^u\| \leq N\lambda_1^{-\tau}.$$

□

REFERENCES

- [1] R. Mane, An ergodic closing lemma, *Ann. Math.*, **116** (1982), 503–540.
- [2] L. Wen, Homoclinic tangencies and dominated splittings, *Nonlinearty*, **15** (2002), 1445–1469.
- [3] L. Wen, Generic diffeomorphisms away from homoclinic tangencies and heterodimensional cycles, *Bull. Braz. Math. Soc.(N.S.)*, **35** (2004), 419–452.
- [4] Q. Xiao and Z. Zheng, C^1 weak Palis conjecture for nonsingular flows, [arXiv:1507.07781](https://arxiv.org/abs/1507.07781)