

# NONCOMMUTATIVE MAXIMAL ERGODIC INEQUALITIES ASSOCIATED WITH DOUBLING CONDITIONS

GUIXIANG HONG, BEN LIAO AND SIMENG WANG

ABSTRACT. This paper is devoted to the study of noncommutative maximal inequalities and ergodic theorems for group actions on von Neumann algebras. Consider a locally compact group  $G$  of polynomial growth with a symmetric compact subset  $V$ . Let  $\alpha$  be a continuous action of  $G$  on a von Neumann algebra  $\mathcal{M}$  by trace-preserving automorphisms. We then show that the operators defined by

$$A_n x = \frac{1}{m(V^n)} \int_{V^n} \alpha_g x \, dm(g), \quad x \in L_p(\mathcal{M}), n \in \mathbb{N}, 1 \leq p \leq \infty,$$

are of weak type  $(1, 1)$  and of strong type  $(p, p)$  for  $1 < p < \infty$ . Consequently, the sequence  $(A_n x)_{n \geq 1}$  converges almost uniformly for  $x \in L_p(\mathcal{M})$  for  $1 \leq p < \infty$ . Also, we establish the noncommutative maximal and individual ergodic theorems associated with more general doubling conditions, and we prove the corresponding results for general actions on one fixed noncommutative  $L_p$ -space which are beyond the class of Dunford–Schwartz operators considered previously by Junge and Xu. As key ingredients, we also obtain the Hardy–Littlewood maximal inequality on metric spaces with doubling measures in the operator-valued setting. After the groundbreaking work of Junge and Xu on the noncommutative Dunford–Schwartz maximal ergodic inequalities, this is the first time that more general maximal inequalities are proved beyond Junge and Xu’s setting. Our approach is based on quantum probabilistic methods as well as random walk theory.

## 1. INTRODUCTION

This article studies maximal inequalities and ergodic theorems for group actions on noncommutative  $L_p$ -spaces. The connection between ergodic theory and von Neumann algebras goes back to the very beginning of the theory of operator algebras. However, the study of individual ergodic theorems in the noncommutative setting only took off with Lance’s pioneering work [Lan76] in 1976. The topic was then extensively investigated in a series of works of Conze, Dang-Ngoc, Kümmerer, Yeadon, and others (see [CDN78, Küm78, Yea77, Jaj85] and references therein). Among them, Yeadon [Yea77] obtained a maximal ergodic theorem in the preduals of semifinite von Neumann algebras. But the corresponding maximal inequalities in  $L_p$ -spaces remained open until the celebrated work of Junge and Xu [JX07], which established the noncommutative analogue of the Dunford–Schwartz maximal ergodic theorem. This breakthrough has motivated further research on noncommutative ergodic theorems, such as [AD06, Bek08, HS16, Hu08], and [Lit14]. Note that all these works essentially remain in the class of Dunford–Schwartz operators, that is, do not go beyond Junge and Xu’s setting.

On the other hand, in classical ergodic theory, a number of significant developments related to individual ergodic theorems for group actions have been established in recent years. In particular, Breuillard [Bre14] and Tessera [Tes07] studied the balls in groups of polynomial growth; they proved that for any invariant metric quasi-isometric to a word metric (such as invariant Riemannian metrics on connected nilpotent Lie groups), the balls are asymptotically invariant and satisfy the doubling condition, and hence satisfy the individual ergodic theorem. This settled a long-standing problem in ergodic theory since Calderón’s classical paper [Cal53] in 1953. Also, Lindenstrauss [Lin01] proved the individual ergodic theorem for tempered Følner sequences, which resolves the

---

*Date:* November 3, 2020.

*Key words and phrases.* Maximal ergodic theorems, individual ergodic theorems, noncommutative  $L_p$ -spaces, Hardy–Littlewood maximal inequalities, transference principles.

problem of constructing pointwise ergodic sequences on an arbitrary amenable group. (We refer to [Nev06] for more details.)

Thus, it is natural to extend Junge and Xu's work to actions of more general amenable groups rather than the integer group. As in the classical case, the first natural step would be to establish the maximal ergodic theorems for doubling conditions. However, since we do not have an appropriate analogue of covering lemmas in the noncommutative setting, no significant progress has been made in this direction. In this paper we provide a new approach to this problem. It is based on both classical and quantum probabilistic methods, and allows us to go beyond the class of Dunford–Schwartz operators considered by Junge and Xu.

Our main results establish the noncommutative maximal and individual ergodic theorems for ball averages under the doubling condition. Let  $G$  be a locally compact group equipped with a right Haar measure  $m$ . Recall that for an invariant metric  $d$  on  $G$ ,<sup>1</sup> we say that  $(G, d)$  satisfies the *doubling condition* if the balls  $B_r := \{s \in G : d(s, e) \leq r\}$  satisfy

$$(1.1) \quad m(B_{2r}) \leq Cm(B_r), \quad r > 0,$$

where  $C$  is a constant independent of  $r$ . We say that the balls are *asymptotically invariant* under right translation if for every  $s \in G$ ,

$$(1.2) \quad \lim_{r \rightarrow \infty} \frac{m((B_r s) \triangle B_r)}{m(B_r)} = 0,$$

where  $\triangle$  denotes the usual symmetric difference of subsets. To state the noncommutative ergodic theorems, we consider a von Neumann algebra  $\mathcal{M}$  equipped with a normal semifinite trace  $\tau$ . We also consider an action  $\alpha$  of  $G$  on the associated noncommutative  $L_p$ -spaces  $L_p(\mathcal{M})$ , under some mild assumptions clarified in later sections. In particular, if  $\alpha$  is a continuous action of  $G$  on  $\mathcal{M}$  by  $\tau$ -preserving automorphisms of  $\mathcal{M}$ , then  $\alpha$  extends to isometric actions on the spaces  $L_p(\mathcal{M})$ . The following is one of our main results.

**Theorem 1.1.** *Assume that  $(G, d)$  satisfies (1.1) and (1.2). Let  $\alpha$  be a continuous action of  $G$  on  $\mathcal{M}$  by  $\tau$ -preserving automorphisms. Let  $A_r$  be the averaging operators*

$$A_r x = \frac{1}{m(B_r)} \int_{B_r} \alpha_s x \, dm(s), \quad x \in \mathcal{M}, r > 0.$$

*Then  $(A_r)_{r>0}$  is of weak type (1,1) and of strong type  $(p,p)$  for  $1 < p < \infty$ . Moreover, for all  $1 \leq p < \infty$ , the sequence  $(A_r x)_{r>0}$  converges almost uniformly for  $x \in L_p(\mathcal{M})$ .*

Here we refer to Section 2.1 for the notion of weak and strong type  $(p,p)$  inequalities in the noncommutative setting. Also, the notion of almost uniform convergence is a noncommutative analogue of the notion of almost everywhere convergence. We refer to Definition 6.1 for the relevant definitions.

There exist a number of examples satisfying assumptions (1.1) and (1.2) of the above theorem, for which we refer to the previously mentioned works [Bre14, Nev06], and [Tes07]. In particular, if we take  $G$  to be the integer group  $\mathbb{Z}$  and  $d$  to be the usual word metric, then we recover the usual ergodic average  $A_n = \frac{1}{2n+1} \sum_{k=-n}^n T^k$  for an invertible operator  $T$ , as is treated in [JX07]. More generally, we may consider groups of polynomial growth.

**Theorem 1.2.** *Assume that  $G$  is generated by a symmetric compact subset  $V$  and is of polynomial growth.*

- (1) *Fix  $1 < p < \infty$ . Let  $\alpha$  be a strongly continuous and uniformly bounded action of  $G$  on  $L_p(\mathcal{M})$  such that  $\alpha_s$  is a positive map for each  $s \in G$ . Then the operators defined by*

$$A_n x = \frac{1}{m(V^n)} \int_{V^n} \alpha_s x \, dm(s), \quad x \in L_p(\mathcal{M}), n \in \mathbb{N}$$

*are of strong type  $(p,p)$ . The sequence  $(A_n x)_{n \geq 1}$  converges bilaterally almost uniformly for  $x \in L_p(\mathcal{M})$ .*

---

<sup>1</sup>In this paper, we always assume that  $d$  is a measurable function on  $G \times G$  and  $m$  is a Radon Borel measure with respect to  $(G, d)$ .

- (2) Let  $\alpha$  be a strongly continuous action of  $G$  on  $\mathcal{M}$  by  $\tau$ -preserving automorphisms. Then the operators defined by

$$A_n x = \frac{1}{m(V^n)} \int_{V^n} \alpha_s x \, dm(s), \quad x \in \mathcal{M}, n \in \mathbb{N}$$

are of weak type  $(1, 1)$  and of strong type  $(p, p)$  for all  $1 < p < \infty$ . The sequence  $(A_n x)_{n \geq 1}$  converges almost uniformly for  $x \in L_p(\mathcal{M})$  for all  $1 \leq p < \infty$ .

The theorems rely on several key results obtained in this work. We address the following subjects.

(i) *Noncommutative transference principles.* Our first key ingredient is a noncommutative variant of Calderón's transference principle in [Cal68] (see also [CW76, Fen98]), given in Theorems 3.1 and 3.3. More precisely, we prove that for actions by an amenable group, in order to establish the noncommutative maximal ergodic inequalities, it suffices to show the inequalities for translation actions on operator-valued functions. We remark that the particular case of certain actions by  $\mathbb{R}$  is also discussed in [Hon17] by the first author.

(ii) *Noncommutative Hardy–Littlewood maximal inequalities on metric measure spaces.* As for the second key ingredient, we prove in Theorem 4.1 a noncommutative extension of Hardy–Littlewood maximal inequalities on metric measure spaces. For a doubling metric measure space  $(X, d, \mu)$ , denote by  $B(x, r)$  the ball with center  $x$  and radius  $r$  with respect to the metric  $d$ . Our result asserts that the Hardy–Littlewood averaging operators on the  $L_p(\mathcal{M})$ -valued functions

$$A_r f(x) = \frac{1}{\mu(B(x, r))} \int_{B(x, r)} f \, d\mu, \quad f \in L_p(X; L_p(\mathcal{M})), x \in X, r > 0$$

satisfy the weak type  $(1, 1)$  and strong type  $(p, p)$  inequalities. We remark that the classical argument via covering lemmas does not seem to fit into this operator-valued setting. Instead, our approach is based on the study of random dyadic systems by Naor and Tao in [NT10] and Hytönen and Kairema in [HK12]. The key idea is to relate the desired inequality to noncommutative martingales, and to use the available results in quantum probability developed in [Cuc71] and [Jun02]. The approach is inspired by Mei's famous work in [Mei03] and [Mei07], which asserts that the usual continuous BMO space is the intersection of several dyadic BMO spaces.

(iii) *Domination by Markov operators.* In the study of ergodic theorems for actions by free groups or free abelian groups, it is a key fact that the associated ergodic averages can be dominated by the standard averaging operators of the form  $\frac{1}{n} \sum_{k=1}^n T^k$  for some map  $T$  (see [Bru73, NS94]). Also, in [SS83], Stein and Strömberg apply the Markov semigroup with heat kernels to estimate the maximal inequalities on Euclidean spaces with large dimensions. In this paper, we build a similar result for groups of polynomial growth. Our approach is new and the construction follows easily from some typical Markov chains on these groups. More precisely, we show in Proposition 4.8 that for a group  $G$  of polynomial growth with a symmetric compact generating subset  $V \subset G$ , and for an action  $\alpha$  of  $G$ , there exists a constant  $c$  such that

$$\frac{1}{m(V^n)} \int_{V^n} \alpha_s x \, dm(s) \leq \frac{c}{n^2} \sum_{k=1}^{2n^2} T^k x, \quad x \geq 0,$$

where  $T = \frac{1}{m(V)} \int_V \alpha_s \, dm(s)$ . The result will help us to improve the weak type inequalities in Theorem 1.2.

(iv) *Individual ergodic theorems for  $L_p$  representations.* In the classical setting, the individual ergodic theorem holds for positive contractions on  $L_p$ -spaces with one fixed  $p \in (1, \infty)$  (see [IT64, Akc75]). The results can be also generalized for positive power-bounded operators and more general Lamperti operators (see, e.g., [Kan78, MRDIT88, Tem15]). However, in the noncommutative setting, the individual ergodic theorems on  $L_p$ -spaces were only known for operators which can be extended to  $L_1 + L_\infty$ . In Section 6 we will develop some new methods to prove the individual ergodic theorems for operators on one fixed  $L_p$ -space.

Apart from the above approach, we also provide in Section 5 an alternative proof of Theorem 1.1 for discrete groups of polynomial growth. Compared to the previous approach, this proof is much

more group-theoretical and has its own interests. It relies essentially on the concrete structure of groups of polynomial growth discovered by Bass, Gromov, and Wolf.

We remark that although our results are stated in the setting of tracial  $L_p$ -spaces, a large number of the results can be extended to the general nontracial case without difficulty. Since the standard methods for these generalizations are already well developed in [HJX10] and [JX07], we leave the details to the reader and restrict our attention to the semifinite case for simplicity of exposition.

We end this Introduction with a brief description of the organization of the paper. In the next section, we recall some basics on noncommutative maximal operators as well as actions by amenable groups. Section 3 is devoted to the proof of the noncommutative variant of Calderón's transference principle. In Section 4, we prove the Hardy–Littlewood maximal inequalities mentioned above and deduce the maximal inequalities in Theorem 1.1. We also use similar ideas to establish the ergodic theorems for increasing sequences of compact subgroups (Theorem 4.7). In the last part of the section, we will provide an approach based on the random walk theory, which relates the ball averages to the classical ergodic averages of Markov operators. In Section 5, we provide an alternative group-theoretical approach to Theorem 1.2. In Section 6, we discuss the individual ergodic theorems, which prove the bilateral almost uniform convergences in Theorem 1.1. Also, we give new results on almost uniform convergences associated with actions on one fixed  $L_p$ -space.

## 2. PRELIMINARIES

**2.1. Noncommutative  $L_p$ -spaces and noncommutative maximal norms.** Throughout the paper, unless explicitly stated otherwise,  $\mathcal{M}$  will always denote a semifinite von Neumann algebra equipped with a normal semifinite trace  $\tau$ . Let  $\mathcal{S}_+$  denote the set of all  $x \in \mathcal{M}_+$  such that  $\tau(\text{supp } x) < \infty$ , where  $\text{supp } x$  denotes the support of  $x$ . Let  $\mathcal{S}$  be the linear span of  $\mathcal{S}_+$ . Given  $1 \leq p < \infty$ , we define

$$\|x\|_p = [\tau(|x|^p)]^{1/p}, \quad x \in \mathcal{S},$$

where  $|x| = (x^*x)^{1/2}$  is the modulus of  $x$ . Then  $(\mathcal{S}, \|\cdot\|_p)$  is a normed space, whose completion is the noncommutative  $L_p$ -space associated with  $(\mathcal{M}, \tau)$ , denoted by  $L_p(\mathcal{M})$ . As usual, we set  $L_\infty(\mathcal{M}) = \mathcal{M}$  equipped with the operator norm. Let  $L_0(\mathcal{M})$  denote the space of all closed densely defined operators on  $H$  measurable with respect to  $(\mathcal{M}, \tau)$  ( $H$  being the Hilbert space on which  $\mathcal{M}$  acts). Then  $L_p(\mathcal{M})$  can be viewed as closed densely defined operators on  $H$ . We denote by  $L_0^+(\mathcal{M})$  the positive part of  $L_0(\mathcal{M})$ , and set  $L_p^+(\mathcal{M}) = L_0^+(\mathcal{M}) \cap L_p(\mathcal{M})$ . We refer to [PX03] for more information on noncommutative  $L_p$ -spaces.

For a  $\sigma$ -finite measure space  $(X, \Sigma, \mu)$ , we consider the von Neumann algebraic tensor product  $L_\infty(X) \bar{\otimes} \mathcal{M}$  equipped with the trace  $\int \otimes \tau$ , where  $\int$  denotes the integral against  $\mu$ . For  $1 \leq p < \infty$ , the space  $L_p(L_\infty(X) \bar{\otimes} \mathcal{M})$  isometrically coincides with  $L_p(X; L_p(\mathcal{M}))$ , the usual  $L_p$ -space of  $p$ -integrable functions from  $X$  to  $L_p(\mathcal{M})$ . In this paper we will not distinguish these two notions unless specified otherwise.

Maximal norms in the noncommutative setting require a specific definition. The subtlety is that  $\sup_n |x_n|$  does not make any sense for a sequence  $(x_n)_n$  of arbitrary operators. This difficulty is overcome by considering the spaces  $L_p(\mathcal{M}; \ell_\infty)$ , which are the noncommutative analogues of the usual Bochner spaces  $L_p(X; \ell_\infty)$ . These vector-valued  $L_p$ -spaces were first introduced by Pisier [Pis98] for injective von Neumann algebras and then extended to general von Neumann algebras by Junge [Jun02]. The descriptions and properties below can be found in [JX07, Section 2]. Given  $1 \leq p \leq \infty$ ,  $L_p(\mathcal{M}; \ell_\infty)$  is defined as the space of all sequences  $x = (x_n)_{n \geq 0}$  in  $L_p(\mathcal{M})$  which admit a factorization of the following form: there are  $a, b \in L_{2p}(\mathcal{M})$  and a bounded sequence  $y = (y_n) \subset L_\infty(\mathcal{M})$  such that

$$x_n = ay_n b, \quad \forall n \geq 0.$$

We then define

$$\|x\|_{L_p(\mathcal{M}; \ell_\infty)} = \inf \left\{ \|a\|_{2p} \sup_{n \geq 0} \|y_n\|_\infty \|b\|_{2p} \right\},$$

where the infimum runs over all factorizations as above. We will adopt the convention that the norm  $\|x\|_{L_p(\mathcal{M}; \ell_\infty)}$  is denoted by  $\|\sup_n^+ x_n\|_p$ . As an intuitive description, we remark that a positive sequence  $(x_n)_{n \geq 0}$  of  $L_p(\mathcal{M})$  belongs to  $L_p(\mathcal{M}; \ell_\infty)$  if and only if there exists a positive  $a \in L_p(\mathcal{M})$  such that  $x_n \leq a$  for any  $n \geq 0$  and in this case,

$$\|\sup_n^+ x_n\|_p = \inf\{\|a\|_p : a \in L_p(\mathcal{M}), a \geq 0 \text{ and } x_n \leq a \text{ for any } n \geq 0\}.$$

Also, we denote by  $L_p(\mathcal{M}; c_0)$  the closure of finite sequences in  $L_p(\mathcal{M}; \ell_\infty)$  for  $1 \leq p < \infty$ . On the other hand, we may also define the space  $L_p(\mathcal{M}; \ell_\infty^c)$ , which is the space of all sequences  $x = (x_n)_{n \geq 0}$  in  $L_p(\mathcal{M})$  which admit a factorization of the following form: there are  $a \in L_p(\mathcal{M})$  and  $y = (y_n) \subset L_\infty(\mathcal{M})$  such that

$$x_n = y_n a, \quad \forall n \geq 0.$$

And we define

$$\|x\|_{L_p(\mathcal{M}; \ell_\infty^c)} = \inf\{\sup_{n \geq 0} \|y_n\|_\infty \|a\|_p\},$$

where the infimum runs over all factorizations as above. Similarly, we denote by  $L_p(\mathcal{M}; c_0^c)$  the closure of finite sequences in  $L_p(\mathcal{M}; \ell_\infty^c)$ . (We refer to [DJ04] and [Mus03] for more information.)

Indeed, for any index  $I$ , we can define the spaces  $L_p(\mathcal{M}; \ell_\infty(I))$  of families  $(x_i)_{i \in I}$  in  $L_p(\mathcal{M})$  with similar factorizations as above. We omit the details and we will simply denote the spaces by the same notation  $L_p(\mathcal{M}; \ell_\infty)$  and  $L_p(\mathcal{M}; c_0)$  if no confusion can occur.

The following properties will be useful here.

**Proposition 2.1.**

- (1) A family  $(x_i)_{i \in I} \subset L_p(\mathcal{M})$  belongs to  $L_p(\mathcal{M}; \ell_\infty)$  if and only if

$$\sup_{J \text{ finite}} \|\sup_{i \in J}^+ x_i\|_p < \infty,$$

and in this case

$$\|\sup_{i \in I}^+ x_i\|_p = \sup_{J \text{ finite}} \|\sup_{i \in J}^+ x_i\|_p.$$

- (2) Let  $1 \leq p_0 < p_1 \leq \infty$ , and let  $0 < \theta < 1$ . Then we have isometrically

$$(2.1) \quad L_p(\mathcal{M}; \ell_\infty) = (L_{p_0}(\mathcal{M}; \ell_\infty), L_{p_1}(\mathcal{M}; \ell_\infty))_\theta,$$

where  $\frac{1}{p} = \frac{1-\theta}{p_0} + \frac{\theta}{p_1}$ . If additionally  $p_0 \geq 2$ , then we have isometrically

$$(2.2) \quad L_p(\mathcal{M}; \ell_\infty^c) = (L_{p_0}(\mathcal{M}; \ell_\infty^c), L_{p_1}(\mathcal{M}; \ell_\infty^c))_\theta,$$

where  $\frac{1}{p} = \frac{1-\theta}{p_0} + \frac{\theta}{p_1}$ .

Based on these notions we can discuss the noncommutative maximal inequalities.

**Definition 2.2.** Let  $1 \leq p \leq \infty$ , and let  $S = (S_i)_{i \in I}$  be a family of maps from  $L_p^+(\mathcal{M})$  to  $L_0^+(\mathcal{M})$ .

- (1) For  $p < \infty$ , we say that  $S$  is of *weak type*  $(p, p)$  with constant  $C$  if there exists a constant  $C > 0$  such that, for all  $x \in L_p^+(\mathcal{M})$  and  $\lambda > 0$ , there is a projection  $e \in \mathcal{M}$  satisfying

$$\tau(1 - e) \leq \frac{C^p}{\lambda^p} \|x\|_p^p, \quad e S_i(x) e \leq \lambda e, \quad i \in I.$$

- (2) For  $1 \leq p \leq \infty$ , we say that  $S$  is of *strong type*  $(p, p)$  with constant  $C$  if there exists a constant  $C > 0$  such that

$$\|\sup_{i \in I}^+ S_i x\|_p \leq C \|x\|_p, \quad x \in L_p(\mathcal{M}).$$

We will also need a reduction below for weak type inequalities.

**Lemma 2.3.** *If for all finite subsets  $J \subset I$ ,  $(S_i)_{i \in J}$  is of weak type  $(p, p)$  with constant  $C$ , then  $(S_i)_{i \in I}$  is of weak type  $(p, p)$  with constant  $2^{1/p} C$ .*

*Proof.* Indeed, the proof of [Hon17, Lemma 3.2] shows the following general property: if a family  $(x_i)_{i \in I} \subset L_0^+(\mathcal{M})$  and a constant  $c > 0$  satisfy that for all  $\lambda > 0$  and all finite subsets  $J \subset I$ , there exists a projection  $e_J \in \mathcal{M}$  with

$$\tau(1 - e_J) \leq c^p \lambda^{-p}, \quad \|e_J x_i e_J\|_\infty \leq \lambda, \quad i \in J,$$

then there exists a projection  $e \in \mathcal{M}$  with

$$\tau(1 - e) \leq 2c^p \lambda^{-p}, \quad \|e x_i e\|_\infty \leq \lambda, \quad i \in I.$$

To see this, we note that  $\mathcal{S}_+$  is dense in  $L_0^+(\mathcal{M})$  with respect to the topology of convergence in measure. So we may assume without loss of generality that  $(x_i)_{i \in I} \subset \mathcal{S}_+$ . Then it suffices to take a  $w^*$ -accumulation point  $u$  of  $(e_J)_J$  and set the spectral projection  $e = \chi_{[\frac{1}{2}, 1]}(u)$ . In this case we have

$$\begin{aligned} \|e x_i e\|_\infty &= \|e u x_i u e\|_\infty \\ &\leq \|u x_i u\|_\infty = \|x_i^{1/2} u\|_\infty^2 = \sup_{y \in L_1(\mathcal{M}) \cap \mathcal{M}} \frac{|\tau(x_i^{1/2} u y)|^2}{\|y\|_1^2} \\ &= \sup_{y \in L_1(\mathcal{M})} \sup_J \frac{|\tau(x_i^{1/2} e_J y)|^2}{\|y\|_1^2} \\ &\leq \sup_J \|x_i^{1/2} e_J\|_\infty^2 \leq \lambda \end{aligned}$$

and

$$e = \chi_{[\frac{1}{2}, 1]}(u) \leq 2u, \quad 1 - e \leq 2(1 - u).$$

Then we obtain the claimed property. The lemma follows by taking  $x_i = S_i(x)$  and  $c = C\|x\|_p$ .  $\square$

The following noncommutative Doob inequalities will play a crucial role in our proof.

**Lemma 2.4** ([Cuc71, Proposition 6], [Jun02, Theorem 0.2]). *Let  $(\mathcal{M}_n)_{n \in \mathbb{Z}}$  be an increasing sequence of von Neumann subalgebras of  $\mathcal{M}$  such that  $\bigcup_{n \in \mathbb{Z}} \mathcal{M}_n$  is  $w^*$ -dense in  $\mathcal{M}$ . Denote by  $\mathbb{E}_n$  the  $\tau$ -preserving conditional expectation from  $L_p(\mathcal{M})$  onto  $L_p(\mathcal{M}_n)$ . Then  $(\mathbb{E}_n)_{n \in \mathbb{Z}}$  is of weak type  $(1, 1)$  with a universal constant and is of strong type  $(p, p)$  with constants depending only on  $p$  for all  $1 < p < \infty$ .*

Note that only monotone sequences of the form  $(\mathbb{E}_n)_{n \geq 0}$  are concerned in the statement of [Cuc71] and [Jun02]. However, we can easily deduce the general result in Lemma 2.4 from the previous case. Indeed, for a sequence of the form  $(\mathbb{E}_n)_{n \in \mathbb{Z}}$  in the above lemma, the statement in [Cuc71] and [Jun02] yields that for all  $k \in \mathbb{Z}$ , the sequence  $(\mathbb{E}_n)_{n \geq k}$  is of weak type  $(1, 1)$  and strong type  $(p, p)$  with constants independent of  $k$  by reindexing the sequence. In particular,  $(\mathbb{E}_n)_{-k \leq n \leq k}$  satisfies the same maximal inequalities. Then by Proposition 2.1(1) and Lemma 2.3, we see that the same property holds for the sequence  $(\mathbb{E}_n)_{n \in \mathbb{Z}}$  as well.

**2.2. Actions by amenable groups.** Unless explicitly stated otherwise, throughout  $G$  will denote a locally compact group with neutral element  $e$ , equipped with a fixed right invariant Haar measure  $m$ . For a Banach space  $E$ , we say that

$$\alpha : G \rightarrow B(E), \quad s \mapsto \alpha_s$$

is an action if  $\alpha_s \circ \alpha_h = \alpha_{st}$  for all  $s, t \in G$ . Let  $(\mathcal{M}, \tau)$  be as before. For a fixed  $1 \leq p \leq \infty$ , we will be interested in actions  $\alpha = (\alpha_s)_{s \in G}$  on  $L_p(\mathcal{M})$  with the following conditions:

- (A<sub>1</sub><sup>p</sup>) Continuity: for all  $x \in L_p(\mathcal{M})$ , the map  $s \mapsto \alpha_s x$  from  $G$  to  $L_p(\mathcal{M})$  is continuous. Here we take the norm topology on  $L_p(\mathcal{M})$  if  $1 \leq p < \infty$  and the  $w^*$ -topology if  $p = \infty$ .
- (A<sub>2</sub><sup>p</sup>) Uniform boundedness:  $\sup_{s \in G} \|\alpha_s : L_p(\mathcal{M}) \rightarrow L_p(\mathcal{M})\| < \infty$ .
- (A<sub>3</sub><sup>p</sup>) Positivity: for all  $s \in G$ ,  $\alpha_s x \geq 0$  if  $x \geq 0$  in  $L_p(\mathcal{M})$ .

As a natural example, if  $\alpha$  is an action on  $\mathcal{M}$  satisfying the condition:

- (A') for all  $x \in \mathcal{M}$ , the map  $s \mapsto \alpha_s x$  from  $G$  to  $\mathcal{M}$  is continuous with respect to the  $w^*$ -topology on  $\mathcal{M}$ ; and for all  $s \in G$ ,  $\alpha_s$  is an automorphism of  $\mathcal{M}$  (in the sense of  $*$ -algebraic structures) such that  $\tau = \tau \circ \alpha_s$ ,

then  $\alpha$  extends naturally to actions on  $L_p(\mathcal{M})$  with conditions  $(\mathbf{A}_1^p)$ – $(\mathbf{A}_3^p)$  for all  $1 \leq p \leq \infty$ , still denoted by  $\alpha$  (see, e.g., [JX07, Lemma 1.1]). In this case for each  $s \in G$ ,  $\alpha_s$  is an isometry on  $L_p(\mathcal{M})$ . We refer to [Bek15, Oli12], and [Oli13] for other natural examples of group actions on noncommutative  $L_p$ -spaces.

Recall that  $G$  is said to be *amenable* if  $G$  admits a Følner net, that is, a net  $(F_i)_{i \in I}$  of measurable subsets of  $G$  with  $m(F_i) < \infty$  such that for all  $s \in G$ ,

$$\lim_i \frac{m((F_i s) \triangle F_i)}{m(F_i)} = 0.$$

Note that the above condition is a reformulation of the asymptotic invariance (1.2) for the general setting. It is known that  $(F_i)_{i \in I}$  is a Følner net if for all compact measurable subsets  $K \subset G$ ,

$$(2.3) \quad \lim_i \frac{m(F_i K)}{m(F_i)} = 1.$$

Recall that  $G$  is a compactly generated group of *polynomial growth* if the compact generating subset  $V \subset G$  satisfies

$$m(V^n) \leq kn^r, \quad n \geq 1,$$

where  $k > 0$  and  $r \in \mathbb{N}$  are constants independent of  $n$ . It is well known that any group of polynomial growth is amenable and the sequence  $(V^n)_{n \geq 1}$  satisfies the above Følner condition (see, e.g., [Bre14, Tes07]). We refer to [Pat88] for more information on amenable groups.

Now let  $G$  be amenable, and let  $(F_i)_{i \in I}$  be a Følner net in  $G$ . Let  $1 < p < \infty$ . Let  $\alpha = (\alpha_s)_{s \in G}$  be an action of  $G$  on  $L_p(\mathcal{M})$  satisfying  $(\mathbf{A}_1^p)$ – $(\mathbf{A}_3^p)$ . Denote by  $A_i$  the corresponding averaging operators

$$A_i x = \frac{1}{m(F_i)} \int_{F_i} \alpha_s x \, dm(s), \quad x \in L_p(\mathcal{M}).$$

According to the mean ergodic theorem for amenable groups (see, e.g., [ADAB<sup>+</sup>10, Théorème 2.2.7]), we have a canonical splitting on  $L_p(\mathcal{M})$ :

$$L_p(\mathcal{M}) = \mathcal{F}_p \oplus \mathcal{F}_p^\perp$$

with

$$(2.4) \quad \begin{aligned} \mathcal{F}_p &= \{x \in L_p(\mathcal{M}) : \alpha_s x = x, s \in G\}, \\ \mathcal{F}_p^\perp &= \overline{\text{span}}\{x - \alpha_s x : s \in G, x \in L_p(\mathcal{M})\}. \end{aligned}$$

Let  $P$  be the bounded positive projection from  $L_p(\mathcal{M})$  onto  $\mathcal{F}_p$ . Then  $(A_i x)_i$  converges to  $Px$  in  $L_p(\mathcal{M})$  for all  $x \in L_p(\mathcal{M})$ .

Assume additionally that  $\alpha$  extends to an action on  $\bigcup_{1 \leq p \leq \infty} L_p(\mathcal{M})$  satisfying  $(\mathbf{A}_1^p)$ – $(\mathbf{A}_3^p)$  for every  $1 \leq p \leq \infty$ . Note that the convergence in  $L_p(\mathcal{M})$  yields the convergence in measure in  $L_0(\mathcal{M})$ , and in particular for  $x \in L_1^+(\mathcal{M}) \cap \mathcal{M}_+$  and for  $p_0 = 1$  or  $p_0 = \infty$ ,

$$\|Px\|_{p_0} \leq \liminf_{n \rightarrow \infty} \|A_n x\|_{p_0} \leq \sup_{s \in G} \|\alpha_s\|_{B(L_{p_0}(\mathcal{M}))} \|x\|_{p_0},$$

so by [JX07, Lemma 1.1] and [Yea77, Proposition 1],  $P$  admits a continuous extension on  $L_1(\mathcal{M})$  and  $\mathcal{M}$ , still denoted by  $P$ . The splitting (2.4) is also true in this case. Note then, however, that  $\mathcal{F}_\infty^\perp$  is the  $w^*$ -closure of the space spanned by  $\{x - \alpha_s x : s \in G, x \in \mathcal{M}\}$ .

### 3. NONCOMMUTATIVE CALDERÓN'S TRANSFERENCE PRINCIPLE

In this section, we discuss a noncommutative variant of Calderón's transference principle. Fix  $1 \leq p < \infty$ . Let  $G$  be a locally compact group, and let  $\alpha$  be an action satisfying  $(\mathbf{A}_1^p)$ – $(\mathbf{A}_3^p)$  in the

previous section. Let  $(\mu_n)_{n \geq 1}$  be a sequence of Radon probability measures on  $G$ . We consider the following averages:

$$(3.1) \quad A_n x = \int_G \alpha_s x d\mu_n(s), \quad x \in L_p(\mathcal{M}), n \geq 1.$$

Let us also consider the natural translation action of  $G$  on itself. We are interested in the following averages: for all  $f \in L_p(G; L_p(\mathcal{M}))$ ,

$$(3.2) \quad A'_n f(s) = \int_G f(st) d\mu_n(t), \quad s \in G, n \geq 1,$$

where the integration denotes the usual integration of Banach space-valued functions.

**3.1. Strong type inequalities.** We begin with the transference principle for strong type  $(p, p)$  inequalities.

**Theorem 3.1.** *Assume that  $G$  is amenable. Fix  $1 \leq p < \infty$ . If there exists a constant  $C > 0$  such that*

$$\left\| \sup_n^+ A'_n f \right\|_p \leq C \|f\|_p, \quad f \in L_p(G; L_p(\mathcal{M})),$$

*then there exists a constant  $C' > 0$  depending on  $\alpha$  such that*

$$\left\| \sup_n^+ A_n x \right\|_p \leq CC' \|x\|_p, \quad x \in L_p(\mathcal{M}).$$

*Proof.* Note that we may take an increasing net of compact subsets  $K_i \subset G$  such that  $\lim_i \mu_n(K_i) = \mu_n(G)$ . Then for

$$A_{n,i} x = \int_G \alpha_s x \chi_{K_i}(s) d\mu_n(s), \quad x \in L_p(\mathcal{M}), n \geq 1,$$

we have

$$A_{n,i} x \leq A_n x, \quad \lim_{i \rightarrow \infty} \|A_n x - A_{n,i} x\|_p = 0, \quad x \in L_p^+(\mathcal{M}).$$

So for  $x, y \in L_p^+(\mathcal{M})$ ,

$$A_n x \leq y \quad \text{iff} \quad \forall i, \quad A_{n,i} x \leq y.$$

Hence  $\| \sup_{n,i}^+ A_{n,i} x_{n,i} \|_p = \| \sup_n^+ A_n x_n \|_p$ . So without loss of generality we may assume that the  $\mu_n$ 's are of compact support.

We fix  $x \in L_p(\mathcal{M})$  and  $N \geq 1$ . Choose a compact subset  $K \subset G$  such that  $\mu_n$  is supported in  $K$  for all  $1 \leq n \leq N$ . Since  $\alpha_s : L_p(\mathcal{M}) \rightarrow L_p(\mathcal{M})$  is positive for all  $s \in G$ , we see that  $(\alpha_s \otimes \text{Id})_{s \in G}$  extends to a uniformly bounded family of maps on  $L_p(\mathcal{M}; \ell_\infty)$  (see, e.g., [HJX10, Proposition 7.3]). So we may choose a constant  $C' > 0$  such that

$$\left\| \sup_{1 \leq n \leq N}^+ A_n x \right\|_p = \left\| \sup_{1 \leq n \leq N}^+ \alpha_{s^{-1}} \alpha_s A_n x \right\|_p \leq C' \left\| \sup_{1 \leq n \leq N}^+ \alpha_s A_n x \right\|_p, \quad s \in G.$$

Let  $F$  be a compact subset. Then we have

$$(3.3) \quad \left\| \sup_{1 \leq n \leq N}^+ A_n x \right\|_p^p \leq C'^p \frac{1}{m(F)} \int_F \left\| \sup_{1 \leq n \leq N}^+ \alpha_s A_n x \right\|_p^p dm(s).$$

We define a function  $f \in L_p(G; L_p(\mathcal{M}))$  as

$$f(h) = \chi_{FK}(h) \alpha_h x, \quad h \in G.$$

Then for all  $s \in F$ ,

$$(3.4) \quad \alpha_s A_n x = \int_K \alpha_{st} x d\mu_n(t) = \int_K f(st) d\mu_n(t) = A'_n f(s).$$

We consider  $(A'_n f)_{1 \leq n \leq N} \in L_p(L_\infty(G) \bar{\otimes} \mathcal{M}; \ell_\infty)$ , and for any  $\varepsilon > 0$  we take a factorization  $A'_n f = a F_n b$  such that  $a, b \in L_{2p}(L_\infty(G) \bar{\otimes} \mathcal{M})$ ,  $F_n \in L_\infty(G) \bar{\otimes} \mathcal{M}$ , and

$$\|a\|_{2p} \sup_{1 \leq n \leq N} \|F_n\|_\infty \|b\|_{2p} \leq \left\| (A'_n f)_{1 \leq n \leq N} \right\|_{L_p(L_\infty(G) \bar{\otimes} \mathcal{M}; \ell_\infty)} + \varepsilon.$$

Then we have

$$\begin{aligned} \int_G \left\| \sup_{1 \leq n \leq N}^+ A'_n f(s) \right\|_p^p dm(s) &\leq \int_G \|a(s)\|_{2p}^p \sup_{1 \leq n \leq N} \|F_n(s)\|_\infty^p \|b(s)\|_{2p}^p dm(s) \\ &\leq \|a\|_{2p}^p \sup_{1 \leq n \leq N} \|F_n\|_\infty^p \|b\|_{2p}^p \\ &\leq (\|(A'_n f)_{1 \leq n \leq N}\|_{L_p(L_\infty(G) \otimes \mathcal{M}; \ell_\infty)} + \varepsilon)^p. \end{aligned}$$

Since  $\varepsilon$  is arbitrarily chosen, we obtain

$$\int_G \left\| \sup_{1 \leq n \leq N}^+ A'_n f(s) \right\|_p^p dm(s) \leq \left\| \sup_{1 \leq n \leq N}^+ A'_n f \right\|_p^p.$$

Thus, together with (3.3), (3.4), and the assumption, we see that

$$\begin{aligned} \left\| \sup_{1 \leq n \leq N}^+ A_n x \right\|_p^p &\leq \frac{C'^p}{m(F)} \int_F \left\| \sup_{1 \leq n \leq N}^+ A'_n f(s) \right\|_p^p dm(s) \leq \frac{C'^p}{m(F)} \left\| \sup_{1 \leq n \leq N}^+ A'_n f \right\|_p^p \\ &\leq \frac{C^p C'^p}{m(F)} \|f\|_p^p = \frac{C^p C'^p}{m(F)} \int_{FK} \|\alpha_h x\|_p^p dm(h) \\ &\leq \frac{C^p C'^p m(FK)}{m(F)} \|x\|_p^p. \end{aligned}$$

Since  $G$  is amenable, for any  $\varepsilon > 0$  we may choose the above subset  $F$  such that  $m(FK)/m(F) \leq 1 + \varepsilon$ . Therefore, we obtain

$$\left\| \sup_{1 \leq n \leq N}^+ A_n x \right\|_p \leq CC'(1 + \varepsilon) \|x\|_p.$$

Note that  $N, \varepsilon, x$  are all arbitrarily chosen, so we establish the theorem.  $\square$

*Remark 3.2.* Applying the same argument, we may obtain several variants of the above theorem.

- (1) The sequence of measures  $(\mu_n)_{n \geq 1}$  can be replaced by any family  $(\mu_i)_{i \in I}$  of Radon probability measures for an arbitrary index set  $I$ .
- (2) The positivity of the action  $\alpha$  can be replaced by more general assumptions. It suffices to assume that

$$\sup_{s \in G} \|\alpha_s \otimes \text{Id}\|_{B(L_p(\mathcal{M}; \ell_\infty))} < \infty.$$

If  $\mathcal{M}$  is commutative, then this is equivalent to saying that the operators  $(\alpha_s)_{s \in G}$  are regular with uniformly bounded regular norm (see [MN91]). In the noncommutative setting, one may assume that  $(\alpha_s)_{s \in G}$  are uniformly bounded decomposable maps, and we refer to [JR04] and [Pis95] for more details.

- (3) One may also state similar properties for transference of linear operators; in this case the assumption on positivity of  $\alpha$  can be ignored, and the semigroup actions can be included. We have the following noncommutative analogue of the transference result in [CW76, Theorem 2.4]. Assume that  $G$  and  $\alpha$  satisfy one of the following conditions:
  - (a)  $G$  is an amenable locally compact group, and  $\alpha$  satisfies  $(\mathbf{A}_1^p)$  and  $(\mathbf{A}_2^p)$ ;
  - (b)  $G$  is a discrete amenable semigroup or  $G = \mathbb{R}_+$ ,  $\alpha$  satisfies  $(\mathbf{A}_1^p)$ , and each  $\alpha_s$  is an isometry on  $L_p(\mathcal{M})$  (or more generally, there exist  $K_1, K_2 > 0$  such that for all  $s \in G$ , we have  $K_1 \|x\|_p \leq \|\alpha_s x\|_p \leq K_2 \|x\|_p$ ).

Let  $\mu$  be a bounded Radon measure on  $G$ . Define

$$T_\mu(f)(s) = \int_G f(st) d\mu(t), \quad f \in L_p(G), s \in G,$$

and

$$\tilde{T}_\mu(x) = \int_G \alpha_s x d\mu(s), \quad x \in L_p(\mathcal{M}).$$

Then we have

$$\|\tilde{T}_\mu\|_{B(L_p(\mathcal{M}))} \leq \sup_{s \in G} \|\alpha_s\|_{B(L_p(\mathcal{M}))} \|T_\mu \otimes \text{Id}\|_{B(L_p(L_\infty(G) \otimes \mathcal{M}))}.$$

**3.2. Weak type inequalities.** Now we discuss the transference principle for weak type  $(p, p)$  inequalities. In this case we will only consider the special case of group actions on von Neumann algebras. We assume that  $\alpha$  is given by an action on  $\mathcal{M}$  satisfying the condition **(A')** in Section 2.2.

**Theorem 3.3.** *Assume that  $G$  is amenable. Let  $(A_n)_{n \geq 1}$  and  $(A'_n)_{n \geq 1}$  be the associated sequences of maps given in (3.1) and (3.2). Fix  $1 \leq p < \infty$ . If the sequence  $(A'_n)_{n \geq 1}$  is of weak type  $(p, p)$ , then  $(A_n)_{n \geq 1}$  is of weak type  $(p, p)$  too.*

*Proof.* As in the last subsection, we may assume without loss of generality that  $\mu_n$  is of compact support. Assume that the sequence  $(A'_n)_{n \geq 1}$  is of weak type  $(p, p)$  with constant  $C$ . By Lemma 2.3, it suffices to show that there exists a constant  $C' > 0$  such that for all  $\lambda > 0$ ,  $x \in L_p^+(\mathcal{M})$ ,  $N \geq 1$ , there exists a projection  $e \in \mathcal{M}$  such that

$$\tau(e^\perp) \leq \frac{C'^p}{\lambda^p} \|x\|_p^p, \quad e(A_n x)e \leq \lambda, \quad 1 \leq n \leq N.$$

We fix  $\lambda > 0$ ,  $x \in L_p(\mathcal{M})$ , and  $N \geq 1$ . Choose a compact subset  $K \subset G$  such that  $\mu_n$  is supported in  $K$  for all  $1 \leq n \leq N$ . Let  $F$  be a compact subset. We define a function  $f \in L_p(G; L_p(\mathcal{M}))$  as

$$f(t) = \chi_{FK}(t)\alpha_t x, \quad t \in G.$$

Then for all  $s \in F$ ,

$$(3.5) \quad \alpha_s A_n x = \int_K \alpha_{st} x d\mu_n(t) = \int_K f(st) d\mu_n(t) = A'_n f(s).$$

Since the sequence  $(A'_n)_{n \geq 1}$  is of weak type  $(p, p)$  with constant  $C$ , we may choose a projection  $e \in L_\infty(G) \bar{\otimes} \mathcal{M}$  such that

$$\int_G \tau(e(s)^\perp) ds \leq \left( C \frac{\|f\|_{L_p(G; L_p(\mathcal{M}))}}{\lambda} \right)^p \quad \text{and} \quad e(A'_n f)e \leq \lambda, \quad 1 \leq n \leq N,$$

where we regard  $e$  as a measurable operator-valued function on  $G$ . Therefore, by (3.5) for  $s \in G$ , we have

$$(3.6) \quad (\alpha_{s^{-1}} e(s))(A_n x)(\alpha_{s^{-1}} e(s)) \leq \lambda, \quad n \geq 1.$$

Recall that each  $\alpha_{s^{-1}}$  is a unital  $\tau$ -preserving automorphism of  $\mathcal{M}$ . In particular, for an arbitrary  $\varepsilon > 0$ , we may choose  $s_0 \in G$  and a projection  $\tilde{e} := \alpha_{s_0^{-1}} e(s_0) \in \mathcal{M}$  such that

$$\tau(\tilde{e}^\perp) \leq \inf_{s \in G} \tau(e(s)^\perp) + \varepsilon \quad \text{and} \quad \tilde{e}(A_n x)\tilde{e} \leq \lambda, \quad n \geq 1.$$

Then we have

$$\begin{aligned} \tau(\tilde{e}^\perp) &\leq \frac{1}{m(F)} \int_F \tau(e(s)^\perp) ds + \varepsilon \leq \frac{C^p}{\lambda^p m(F)} \|f\|_{L_p(G; L_p(\mathcal{M}))}^p + \varepsilon \\ &= \frac{C^p}{\lambda^p m(F)} \int_{FK} \|\alpha_h x\|_p^p dm(h) + \varepsilon \\ &\leq \frac{C^p m(FK)}{\lambda^p m(F)} \|x\|_p^p + \varepsilon. \end{aligned}$$

Since  $G$  is amenable, for any  $\varepsilon > 0$  we may choose the above subset  $F$  such that  $m(FK)/m(F) \leq 1 + \varepsilon$ . Therefore, we obtain

$$\tau(\tilde{e}^\perp) \leq \frac{C^p(1 + \varepsilon)}{\lambda^p} \|x\|_p^p + \varepsilon, \quad \tilde{e}(A_n x)\tilde{e} \leq \lambda, \quad 1 \leq n \leq N.$$

Note that  $N, \varepsilon, x$  are all arbitrarily chosen, so we establish the theorem.  $\square$

*Remark 3.4.* We remark that the above result also holds for other index sets. For example,  $(\mu_n)_{n \geq 1}$  can be replaced by a one-parameter family  $(\mu_r)_{r \in \mathbb{R}_+}$  such that  $r \mapsto A_r x$  is continuous for all  $x \in L_p(\mathcal{M})$ . The only ingredient needed in the proof is that the condition (3.6) hold true almost everywhere on  $G$ .

## 4. MAXIMAL INEQUALITIES: PROBABILISTIC APPROACH

This section is devoted to the proof of the maximal inequalities in Theorem 1.1. To this end, we will first establish noncommutative Hardy–Littlewood maximal inequalities on doubling metric spaces.

**4.1. Hardy–Littlewood maximal inequalities on metric measure spaces.** Throughout, a metric measure space  $(X, d, \mu)$  refers to a metric space  $(X, d)$  equipped with a Radon measure  $\mu$ . We denote  $B(x, r) = \{y \in X : d(y, x) \leq r\}$ , and we say that  $\mu$  satisfies the *doubling condition* if there exists a constant  $K > 0$  such that

$$(4.1) \quad \forall r > 0, x \in X, \quad \mu(B(x, 2r)) \leq K\mu(B(x, r)).$$

In the sequel, we always assume the nondegeneracy property  $0 < \mu(B(x, r)) < \infty$  for all  $r > 0$ . The following theorem can be regarded as an operator-valued analogue of the Hardy–Littlewood maximal inequalities.

**Theorem 4.1.** *Let  $(X, d, \mu)$  be a metric measure space. Suppose that  $\mu$  satisfies the doubling condition. Let  $1 \leq p < \infty$ , and let  $A_r$  be the averaging operators*

$$A_r f(x) = \frac{1}{\mu(B(x, r))} \int_{B(x, r)} f d\mu, \quad f \in L_p(X; L_p(\mathcal{M})), x \in X, r > 0.$$

Then  $(A_r)_{r \in \mathbb{R}_+}$  is of weak type  $(1, 1)$  and of strong type  $(p, p)$  for  $1 < p < \infty$ .

The key ingredient of the proof is the following construction of dyadic systems of metric measure spaces, which is established in [HK12, Corollary 7.4].

**Lemma 4.2.** *Let  $(X, d)$  be a metric space, and let  $\mu$  be a Radon measure on  $X$  satisfying the doubling condition. Then there exists a finite collection of families  $\mathcal{P}^1, \mathcal{P}^2, \dots, \mathcal{P}^N$ , where each  $\mathcal{P}^i := (\mathcal{P}_k^i)_{k \in \mathbb{Z}}$  is a sequence of partitions of  $X$ , such that the following conditions hold true:*

- (1) *for each  $1 \leq i \leq N$  and for each  $k \in \mathbb{Z}$ , the partition  $\mathcal{P}_{k+1}^i$  is a refinement of the partition  $\mathcal{P}_k^i$ ;*
- (2) *there exists a constant  $C > 0$  such that for all  $x \in X$  and  $r > 0$ , there exist  $1 \leq i \leq N$ ,  $k \in \mathbb{Z}$  and an element  $Q \in \mathcal{P}_k^i$  such that*

$$B(x, r) \subset Q, \quad \mu(Q) \leq C\mu(B(x, r)).$$

*Remark 4.3.* The lemma dates back to the construction of dyadic systems in the case of  $X = \mathbb{R}^d$ , which is due to Mei [Mei03, Mei07] (see also [Pis16]). We remark that Mei’s construction also works for the discrete space  $\mathbb{Z}^d$  as follows. For  $0 \leq i \leq d$  and  $k \geq 0$ , we set  $\tilde{\mathcal{P}}_{-k}^i$  to be the following family of intervals in  $\mathbb{Z}$ :

$$\tilde{\mathcal{P}}_{-k}^i = \{[\alpha_k^{(i)} + m2^k, \alpha_k^{(i)} + (m+1)2^k) \cap \mathbb{Z} : m \geq 1\},$$

where  $\alpha_k^{(i)} = \sum_{j=0}^{k-1} 2^j \xi_j^{(i)}$  modulo  $2^k$ , with

$$\xi_{(d+1)n+l}^{(i)} = \delta_{i,l}, \quad n \geq 0, 0 \leq l \leq d.$$

And we set  $\tilde{\mathcal{P}}_k^i = \tilde{\mathcal{P}}_0^i$  for all  $k \geq 0$ . Consider the usual word metric  $d$  and the counting measure  $\mu$  on  $X = \mathbb{Z}^d$ . Then the partitions

$$\mathcal{P}_k^i := (\tilde{\mathcal{P}}_k^i)^d, \quad k \in \mathbb{Z}, 0 \leq i \leq d$$

satisfy the conditions in Lemma 4.2, with constant  $C \leq 2^{3d(d+2)}$ .

*Proof of Theorem 4.1.* Let  $\Sigma$  be the  $\sigma$ -algebra of Borel sets on  $X$ . For  $1 \leq i \leq N$  and  $k \in \mathbb{Z}$ , we define  $\Sigma_k^i \subset \Sigma$  to be the  $\sigma$ -subalgebra generated by the elements of  $\mathcal{P}_k^i$ . Denote by  $\mathbb{E}_k^i$  the conditional expectation from  $L_\infty(X, \Sigma, \mu) \bar{\otimes} \mathcal{M}$  to  $L_\infty(X, \Sigma_k^i, \mu|_{\Sigma_k^i}) \bar{\otimes} \mathcal{M}$ . For each  $x \in X$ , let  $\mathcal{P}_k^i(x)$  be the unique element of  $\mathcal{P}_k^i$  which contains  $x$ . Then we have

$$\mathbb{E}_k^i g(x) = \frac{1}{\mu(\mathcal{P}_k^i(x))} \int_{\mathcal{P}_k^i(x)} g d\mu, \quad x \in X, g \in L_\infty(X, \Sigma, \mu) \bar{\otimes} \mathcal{M}.$$

By Lemma 2.4, there exists a constant  $C > 0$  such that for  $\lambda > 0$  and  $f \in L_1^+(X; L_1(\mathcal{M}))$ , there exists a projection  $e_i \in L_\infty(X, \Sigma, \mu) \otimes \mathcal{M}$  satisfying

$$\tau(e_i^\perp) \leq \frac{C}{\lambda} \|f\|_{L_1(X; L_1(\mathcal{M}))}, \quad e_i(\mathbb{E}_k^i f) e_i \leq \lambda, \quad k \in \mathbb{Z}.$$

Take  $e = \bigwedge_i e_i$  to be the infimum of  $(e_i)_{i=1}^N$ , that is, the projection onto  $\bigcap_i e_i H$  ( $H$  being the Hilbert space on which  $\mathcal{M}$  acts). Note that  $(\bigwedge_i e_i)^\perp \leq \sum_i e_i^\perp$ . Then we have

$$(4.2) \quad \tau(e^\perp) \leq \frac{CN}{\lambda} \|f\|_{L_1(X; L_1(\mathcal{M}))}, \quad e(\mathbb{E}_k^i f) e \leq \lambda, \quad 1 \leq i \leq N, k \in \mathbb{Z}.$$

By Lemma 4.2, there exists a constant  $C' > 0$  such that for each  $x \in X$  and  $r > 0$ , there exist  $1 \leq i \leq N$  and  $k \in \mathbb{Z}$  such that  $B(x, r) \subset \mathcal{P}_k^i(x)$ ,  $\mu(\mathcal{P}_k^i(x)) \leq C' \mu(B(x, r))$ ; in particular,

$$(4.3) \quad \begin{aligned} A_r f(x) &= \frac{1}{\mu(B(x, r))} \int_{B(x, r)} f d\mu \leq C' \frac{1}{\mu(\mathcal{P}_k^i(x))} \int_{\mathcal{P}_k^i(x)} f d\mu \\ &= C' \mathbb{E}_k^i f(x). \end{aligned}$$

Then together with (4.2) we see that

$$e(A_r f) e \leq C' \lambda, \quad r > 0.$$

Therefore,  $(A_r)_{r>0}$  is of weak type  $(1, 1)$ .

On the other hand, for  $1 < p < \infty$ , according to the proof of (4.3), we have for  $f \in L_p^+(X; L_p(\mathcal{M}))$ ,

$$\begin{aligned} \|(A_r f)_{r>0}\|_{L_p(\mathcal{M} \otimes L_\infty(X); \ell_\infty)} &\leq C' \left\| \left( \sum_{i=1}^N \mathbb{E}_k^i f \right)_{k \in \mathbb{Z}} \right\|_{L_p(\mathcal{M} \otimes L_\infty(X); \ell_\infty)} \\ &\leq C' \sum_{i=1}^N \left\| (\mathbb{E}_k^i f)_{k \in \mathbb{Z}} \right\|_{L_p(\mathcal{M} \otimes L_\infty(X); \ell_\infty)}. \end{aligned}$$

Since each  $(\mathbb{E}_k^i)_{k \in \mathbb{Z}}$  on the right-hand side is of strong type  $(p, p)$  by Lemma 2.4, we see that  $(A_r)_{r \in \mathbb{R}_+}$  is of strong type  $(p, p)$ , as desired.  $\square$

*Remark 4.4.* There is another approach to random dyadic systems of metric measure spaces, which is proved by Naor and Tao [NT10, Lemma 3.1]. The construction is motivated by the study of Hardy–Littlewood maximal inequalities on large-dimensional doubling spaces. Their result replaces the families  $\mathcal{P}^1, \mathcal{P}^2, \dots, \mathcal{P}^N$  in Lemma 4.2 by an infinite random collection  $(\mathcal{P}^\omega)_{\omega \in \Omega}$  for a probability space  $(\Omega, P)$ , and assumes a positive probability for the coverings of balls. In this case we may find a random family of martingales  $\{(\mathbb{E}_k^\omega)_{k \in \mathbb{Z}}, \omega \in \Omega\}$  such that for some  $k : \mathbb{R}_+ \rightarrow \mathbb{Z}$  and for some fixed constant  $c$ , we have

$$A_r f \leq c \int_{\Omega} \mathbb{E}_{k(r)}^\omega f(x) dP(\omega), \quad f \in L_p^+(X; L_p(\mathcal{M})).$$

This yields as well the strong type  $(p, p)$  inequalities of  $(A_r)_{r \in \mathbb{R}_+}$  for  $1 < p < \infty$ . We omit the details.

**4.2. Maximal ergodic inequalities.** Based on the previous result, we are now ready to deduce the following maximal ergodic theorems. We say that a metric  $d$  on  $G$  is *invariant* if  $d(e, g) = d(h, hg)$  for all  $g, h \in G$ . We denote  $B_r = B(e, r)$  for  $r > 0$ . As before we consider an action  $\alpha$  on  $L_p(\mathcal{M})$  satisfying the conditions  $(\mathbf{A}_1^p)$ – $(\mathbf{A}_3^p)$  for a fixed  $p$  in Section 2.2. The following result establishes the maximal inequalities in Theorem 1.1.

**Theorem 4.5.** *Let  $G$  be an amenable locally compact group, and let  $d$  be an invariant metric on  $G$ . Assume that  $(G, d)$  satisfies the doubling condition (1.1). Fix  $1 \leq p < \infty$ . Let  $A_r$  be the averaging operators*

$$A_r x = \frac{1}{m(B_r)} \int_{B_r} \alpha_s x dm(s), \quad x \in L_p(\mathcal{M}), r > 0.$$

Then  $(A_r)_{r>0}$  is of strong type  $(p, p)$  if  $1 < p < \infty$ . If, in addition,  $\alpha$  satisfies the condition  $(A')$ , then  $(A_r)_{r>0}$  is of weak type  $(1, 1)$ .

*Proof.* By Theorems 3.1 and 3.3 and the remarks following them, it suffices to prove the maximal inequalities for the averaging operators

$$A'_r f(s) = \frac{1}{m(B_r)} \int_{B_r} f(st) dm(t), \quad f \in L_p(G; L_p(\mathcal{M})), s \in G, r > 0.$$

Since the condition (1.1) holds,  $G$  must be unimodular (see, e.g., [Cal53]). In other words, the measure  $m$  is also invariant under left translation. Note that  $sB_r = B(s, r)$  by the invariance of  $d$ . Thus  $m(B_r) = m(sB_r) = m(B(s, r))$  and

$$A'_r f(s) = \frac{1}{m(B(s, r))} \int_{B(s, r)} f(t) dm(t).$$

And by Theorem 4.1, the right-hand side is of weak type  $(1, 1)$  and of strong type  $(p, p)$  for  $1 < p < \infty$ . Thus  $(A'_r)_{r>0}$  is of weak type  $(1, 1)$  and of strong type  $(p, p)$  for  $1 < p < \infty$  as well. The theorem is proved.  $\square$

**Example 4.6.** The theorem comprises noncommutative variants of classical results due to Wiener [Wie39], Calderón [Cal53], and Nevo [Nev06]. If  $G$  is a compactly generated group of polynomial growth, then the theorem applies to a large class of invariant metrics on  $G$ , such as distance functions derived from an invariant Riemann metric or word metrics. We refer to [Nev06, Sections 4 and 5] for more examples. Here, we list several typical examples satisfying the doubling condition.

- (1) Let  $G$  be a compactly generated group of polynomial growth, and let  $V$  be a symmetric compact generating subset. The *word metric* defined by

$$\forall s, t \in G, \quad d(s, t) = \inf\{n \in \mathbb{N}, s^{-1}h \in V^n\}$$

satisfies (1.1) and (1.2). Note that the integer groups and finitely generated nilpotent groups are of polynomial growth.

- (i) The averaging operators

$$A_n x = \frac{1}{m(V^n)} \int_{V^n} \alpha_s x dm(s), \quad x \in L_p(\mathcal{M}), n \in \mathbb{N}$$

are of strong type  $(p, p)$  if  $1 < p < \infty$ . If, in addition,  $\alpha$  satisfies the condition  $(A')$ , then  $(A_n)_{n \geq 1}$  is of weak type  $(1, 1)$ . This in particular establishes the maximal inequalities in Theorem 1.2.

- (ii) Let  $T : L_p(\mathcal{M}) \rightarrow L_p(\mathcal{M})$  be a positive invertible operator with positive inverse such that  $\sup_{n \in \mathbb{Z}} \|T^n\| < \infty$ . Then

$$A_n = \frac{1}{2n+1} \sum_{k=-n}^n T^k, \quad n \in \mathbb{N}$$

is of strong type  $(p, p)$  if  $1 < p < \infty$ . If  $T$  is an automorphism of  $\mathcal{M}$  which leaves  $\tau$  invariant, then  $(A_n)_{n \geq 1}$  is of weak type  $(1, 1)$ .

- (2) Let  $G$  be a compactly generated group of polynomial growth, and let  $d$  be a metric on  $G$ . If  $d$  is invariant under a cocompact subgroup of  $G$  and if  $d$  satisfies a weak kind of the existence of geodesics axiom (see [Bre14, Definition 4.1]), then  $(G, d)$  satisfies (1.1) and (1.2).

We remark that a natural generalization of the doubling condition (1.1) is given by Tempelman [Tem67] as follows. A sequence  $(F_k)_{k \geq 1}$  of sets of finite measure in  $G$  satisfies *Tempelman's regular condition* if

$$m(F_k^{-1} F_k) < C m(F_k)$$

for some  $C > 0$  independent of  $k$ . We refer to [Tem92, Chapter 5] for more details. It is unclear for us how to establish the noncommutative maximal inequalities in this setting. In the following, we provide a typical example for which the inequalities hold true.

**Theorem 4.7.** *Let  $G$  be an increasing union of compact subgroups  $(G_n)_{n \geq 1}$ . Fix  $1 \leq p < \infty$ . Let  $(A_n)_{n \geq 1}$  be the averaging operators*

$$A_n x = \frac{1}{m(G_n)} \int_{G_n} \alpha_s x \, dm(s), \quad x \in L_p(\mathcal{M}).$$

*Then  $(A_n)_{n \geq 1}$  is of strong type  $(p, p)$  if  $1 < p < \infty$ . If, in addition,  $\alpha$  satisfies the condition  $(\mathbf{A}')$ , then  $(A_n)_{n \geq 1}$  is of weak type  $(1, 1)$ .*

*Proof.* By Theorems 3.1 and 3.3, it suffices to prove the maximal inequalities for the averaging operators

$$A'_n f(s) = \frac{1}{m(G_n)} \int_{G_n} f(st) \, dm(t), \quad f \in L_p(G; L_p(\mathcal{M})), s \in G, n \geq 1.$$

Set  $\Sigma$  to be the  $\sigma$ -algebra of Borel sets on  $G$ . For each  $n \geq 1$ , we define  $\Sigma_n \subset \Sigma$  to be the  $\sigma$ -subalgebra generated by the cosets of  $G_n$

$$\{sG_n : s \in G\}.$$

We see that  $\Sigma_{n+1} \subset \Sigma_n$  for all  $n \geq 1$ . Let  $\mathbb{E}_n$  be the conditional expectation from  $L_\infty(G, \Sigma, m) \bar{\otimes} \mathcal{M}$  to  $L_\infty(G, \Sigma_n, m|_{\Sigma_n}) \bar{\otimes} \mathcal{M}$ . Then it is easy to see that

$$\mathbb{E}_n = A'_n, \quad n \geq 1.$$

According to Lemma 2.4, we see that  $(\mathbb{E}_n)_{n \geq 1}$  is of weak type  $(1, 1)$  and of strong type  $(p, p)$  for  $1 < p < \infty$ . This yields the desired inequalities.  $\square$

**4.3. A random walk approach.** In this subsection, we provide an alternative approach to maximal inequalities for groups of polynomial growth. This approach is based on a Gaussian lower bound of random walks on groups (see [HSC93]). Independent of the previous approaches, in this method we do not need the results on dyadic decompositions of the group, nor do we use the transference principle. The key observation is that we may relate the ball averages on groups with the ergodic averages of a Markov operator.

**Proposition 4.8.** *Let  $G$  be a locally compact group of polynomial growth, and let  $V$  be a compact generating set. Let  $\alpha$  be a strongly continuous action of  $G$  on an ordered Banach space  $E$  such that  $\alpha_s x \geq 0$  for all  $s \in G$  and  $x \in E_+$ . Define an operator  $T$  on  $E$  by*

$$Tx = \frac{1}{m(V)} \int_V \alpha_s x \, dm(s), \quad x \in E.$$

*Then there exists a constant  $c$  depending only on  $G$  such that*

$$\frac{1}{m(V^n)} \int_{V^n} \alpha_s x \, dm(s) \leq \frac{c}{n^2} \sum_{k=1}^{2n^2} T^k x, \quad x \in E_+.$$

We remark that for actions by abelian semigroups, it is known by Brunel [Bru73] that the ergodic averages of multioperators can be related to averages of some Markov operators; Nevo and Stein [NS94] showed that similar observations hold for spherical averages of free group actions. These results play an essential role in the proof of ergodic theorems therein. In the case of groups of polynomial growth, our construction of Markov operators is different from theirs, which is inspired by [SS83]. The argument is relatively easy and based on the Markov chains on groups of polynomial growth.

To prove the proposition, we consider a locally compact group  $G$  and a measure  $\mu$  on  $G$ . For an integer  $k$ , we denote by  $\mu^{*k}$  the  $k$ th convolution of  $\mu$ , that is, the unique measure  $\nu$  on  $G$  satisfying

$$\int_G f \, d\nu = \int_{\prod_{i=1}^k G} f(s_1 \cdots s_k) \, d\mu(s_1) \cdots d\mu(s_k), \quad f \in C_0(G).$$

If  $f$  is the density function of  $\mu$ , then we still denote by  $f^{*k}$  the density function of  $\mu^{*k}$ .

In the following  $d$  will denote the word metric with respect to  $V$  introduced in Example 4.6(1), and  $B_r = \{x \in G : d(e, x) \leq r\}$  for  $r > 0$ .

**Lemma 4.9** ([HSC93, Theorem 5.1]). *Let  $G$  be a locally compact group of polynomial growth, and let  $V$  be a compact generating set. Let  $f$  be the density function of a symmetric continuous probability measure on  $G$  such that  $\text{supp}(f)$  is bounded and  $V \subset \text{supp}(f)$ . Then there exists a constant  $c > 0$  such that for any integer  $k$ ,*

$$f^{*k}(s) \geq \frac{ce^{-d(e,s)^2/k}}{m(B_{\sqrt{k}})}, \quad s \in B_k.$$

**Lemma 4.10.** *Let  $G$ ,  $V$ , and  $f$  be as in the previous lemma. Then there exists a constant  $c > 0$  such that for any integer  $n$ ,*

$$\frac{\chi_{B_n}}{m(B_n)} \leq \frac{c}{2n^2} \sum_{k=1}^{2n^2} f^{*k}.$$

*Proof.* It suffices to prove the inequality in the lemma for sufficiently large  $n$ . By the previous lemma, there exists  $c > 0$  such that

$$f^{*k}(s) \geq \frac{c}{m(B_{\sqrt{k}})}, \quad s \in B_{\sqrt{k}}(e).$$

Therefore, for  $s \in B_n(e)$ ,

$$\begin{aligned} \frac{1}{2n^2} \sum_{k=1}^{2n^2} f^{*k}(s) &\geq \frac{1}{2n^2} \sum_{k=(n+1)^2}^{2n^2} f^{*k}(s) \geq \frac{1}{2n^2} \sum_{k=(n+1)^2}^{2n^2} \frac{c}{m(B_{\sqrt{k}})} \\ &\geq \frac{c(n^2 - 2n - 1)}{2n^2 m(B_{\sqrt{2n}})} \geq \frac{c'}{m(B_n)}, \end{aligned}$$

where  $c' > 0$  is a constant depending only on the doubling condition of  $G$ .  $\square$

*Proof of Proposition 4.8.* We apply Lemma 4.10 with  $f = \chi_V/m(V)$ . Then we obtain for  $x \in E_+$ ,

$$\frac{1}{m(V^n)} \int_{V^n} \alpha_s x \, dm(s) \leq \frac{c}{2n^2} \sum_{k=1}^{2n^2} \int_G \alpha_s x f^{*k}(s) \, dm(s).$$

By the definition of  $f^{*k}$ , we have

$$\int_G \alpha_s x f^{*k}(s) \, dm(s) = \int_{\prod_{i=1}^k G} \alpha_{s_1 \dots s_k} x f(s_1) \cdots f(s_k) \, dm(s_1) \cdots dm(s_k).$$

Recall that  $\alpha$  is a group action and  $f = \chi_V/m(V)$ , so we obtain

$$\int_G \alpha_s x f^{*k}(s) \, dm(s) = \frac{1}{m(V)^k} \int_{\prod_{i=1}^k V} \alpha_{s_1} \cdots \alpha_{s_k} x \, dm(s_1) \cdots dm(s_k) = T^k x.$$

We have therefore established the desired inequality.  $\square$

Proposition 4.8 allows us to deduce maximal ergodic theorems for group actions from well-known ergodic theorems for Markov operators. As a corollary, we may obtain the maximal inequalities in Example 4.6(1). Moreover, for maximal ergodic inequalities on  $L_1$ , it is not necessary to assume that  $\alpha$  is an action by automorphisms as in  $(\mathbf{A}')$ .

**Corollary 4.11.** *Let  $G$  and  $V$  be as above. Let  $\alpha$  be a continuous  $\tau$ -preserving action of  $G$  on  $\mathcal{M}$  such that  $\alpha_s$  is a positive isometry on  $\mathcal{M}$  for each  $s \in G$ . Then  $\alpha$  extends to an action on  $L_1(\mathcal{M})$ . The operators defined by*

$$A_n x = \frac{1}{m(V^n)} \int_{V^n} \alpha_s x \, dm(s), \quad x \in L_1(\mathcal{M}), n \in \mathbb{N}$$

*are of weak type  $(1, 1)$ .*

*Proof.* Note that the operator

$$Tx = \frac{1}{m(V)} \int_V \alpha_s x \, dm(s), \quad x \in \mathcal{M}$$

is a positive contraction on  $\mathcal{M}$ , which preserves  $\tau$ . Then it is well known that the averages  $\frac{1}{n} \sum_{k=1}^n T^k$  are of weak type (1,1) (see [Yea77]). Thus, by Proposition 4.8,  $(A_n)_{n \geq 0}$  is of weak type (1,1) as well.  $\square$

## 5. MAXIMAL INEQUALITIES: GROUP-THEORETIC APPROACH

In this section we provide an alternative approach to Theorem 4.5 in the case where  $G$  is a finitely generated discrete group of polynomial growth,  $d$  is the word metric, and  $p \neq 1$ . The argument follows from a structural study of nilpotent groups.

We first recall some well-known facts on the structure of nilpotent groups. Let  $G$  be a discrete finitely generated nilpotent group with lower central series

$$G = G_1 \supset G_2 \supset \cdots \supset G_K \supset G_{K+1} = \{e\}.$$

Each quotient group  $G_i/G_{i+1}$  is an abelian group of rank  $r_i$ , that is, there is a group isomorphism

$$\pi_i : G_i/G_{i+1} \rightarrow F_i \times \mathbb{Z}^{r_i}$$

with a finite abelian group  $F_i$ . It was shown in [Bas72] that  $G$  is of polynomial growth. We summarize below some facts in the argument of [Bas72]. We may choose a finite generating set  $T$  of  $G$  such that

$$[T, T] = \{s^{-1}t^{-1}st : s, t \in T\} \subset T$$

and take

$$T_i = G_i \cap T, \quad 1 \leq i \leq K+1.$$

Then

$$G_i = \langle T_i \rangle, \quad 1 \leq i \leq K+1.$$

For each  $1 \leq j \leq K$ , we order the elements in  $T_j \setminus T_{j+1}$  as

$$(5.1) \quad T_j \setminus T_{j+1} = \{t_1^{(j)}, t_2^{(j)}, \dots, t_{r_j}^{(j)}, t_{r_j+1}^{(j)}, \dots, t_{l_j}^{(j)}\}$$

so that  $\pi_j([t_1^{(j)}]), \dots, \pi_j([t_{r_j}^{(j)}])$  are the generators of  $\mathbb{Z}^{r_j}$ . Let  $N_j$  be the index of the subgroup  $\langle t_1^{(j)}, t_2^{(j)}, \dots, t_{r_j}^{(j)} \rangle G_{j+1}$  in  $G_j$ .

By a *word* in a subset  $T' \subset T$  we mean a sequence of elements  $w = (s_1, s_2, \dots, s_n)$  with  $s_1, s_2, \dots, s_n \in T'$ , and we denote by  $|w| = s_1 s_2 \cdots s_n \in G$  the resulting group element in  $G$ . If  $w = (s_1, s_2, \dots, s_n)$  is a word in  $T'$  and if  $T'' \subset T'$ , then we let  $\deg_{T''}(w)$  be the cardinality of  $\{k : 1 \leq k \leq n, s_k \in T''\}$ . We say that

$$\deg^{(j)}(w) \leq (d_j, d_{j+1}, \dots, d_K) := d$$

if we have  $\deg_{T_j \setminus T_{i+1}}(w) \leq d_i$  for all  $j \leq i \leq K$ . Denote by  $G_j(d)$  the set of all words  $w$  in  $T_j$  such that  $\deg^{(j)}(w) \leq d$ , and denote by  $G'_j(d)$  the subset of the words  $w \in G_j(d)$  of the form

$$w = (t_1^{(j)}, \dots, t_1^{(j)}, \dots, t_{l_j}^{(j)}, \dots, t_{l_j}^{(j)}, v),$$

where  $v$  is a word in  $T_{j+1}$  and the element  $t_k^{(j)}$  does not appear more than  $N_j$  times in  $w$  for  $r_j < k \leq l_j$ .

The key observation in [Bas72] for proving the polynomial growth of  $G$  is as follows (see the assertions (6) and (7) in [Bas72, p. 613]).

**Lemma 5.1.** *Let  $c > 0$  be a constant, and let  $m \geq 1$ . For each  $1 \leq j \leq K$ , we have*

$$\begin{aligned} & \{g \in G : g = |w|, w \in G_j(cm^j, \dots, cm^K)\} \\ & \subset \{g \in G : g = |w|, w \in G'_j(c'm^j, \dots, c'm^K)\}, \end{aligned}$$

where  $c' > 0$  is a constant depending only on  $c$  and  $G_j$ .

In particular, we take  $g \in T^n$ . Hence  $g$  corresponds to a word  $w$  in  $T = T_1$  such that

$$g = |w|, \quad w \in G_1(m_1, \dots, m_K), m := \max\{m_1, \dots, m_K\} \leq n.$$

Using the lemma inductively, we may find another word  $w' = (w_1, \dots, w_K)$  in  $T$  (where each  $w_j$  in the bracket stands for a subword in  $T_j \setminus T_{j+1}$ ) and a constant  $c > 0$  such that

$$g = |w'|, \quad (w_j, \dots, w_K) \in G'_j(cn, \dots, cn^K), 1 \leq j \leq K.$$

In other words, we have the following observation.

**Lemma 5.2.** *Let  $n \in \mathbb{N}$ . Each element  $g \in T^n$  can be written in the form*

$$g = (t_1^{(1)})^{n_{11}} \dots (t_{l_1}^{(1)})^{n_{1l_1}} \dots (t_1^{(K)})^{n_{K1}} \dots (t_{l_K}^{(K)})^{n_{Kl_K}},$$

where there exists a constant  $c > 0$  such that for  $1 \leq j \leq K$ ,

$$n_{jk} \leq cn^j, \quad \text{if } 1 \leq k \leq r_j,$$

and

$$n_{jk} \leq N_j, \quad \text{if } r_j < k \leq l_j.$$

In [Bas72] and [Wol68] it is proved that  $G$  satisfies the following strict polynomial growth condition.

**Lemma 5.3.** *We have two constants  $c_1, c_2 > 0$  such that*

$$c_1 n^{d(G)} \leq |T^n| \leq c_2 n^{d(G)},$$

where  $||$  denotes the cardinality of a subset and

$$d(G) = \sum_{j=1}^K jr_j.$$

Note that the upper bound in the above lemma follows directly from Lemma 5.2.

Now we will prove the following maximal inequalities, which are particular cases studied in Theorem 4.5 and Example 4.6(1).

**Proposition 5.4.** *Let  $G$  be a finitely generated discrete group of polynomial growth, and let  $S \subset G$  be a finite generating set. Fix  $1 < p < \infty$ , and let  $\alpha$  be an action  $\alpha = (\alpha_s)_{s \in G}$  of  $G$  on  $L_p(\mathcal{M})$  which satisfies  $(\mathbf{A}_1^p)$ – $(\mathbf{A}_3^p)$ . We consider the averaging operators*

$$A_n = \frac{1}{|S^n|} \sum_{s \in S^n} \alpha_s, \quad n \geq 1,$$

where  $||$  denotes the cardinality of a subset. Then  $(A_n)_{n \geq 1}$  is of strong type  $(p, p)$ .

The proposition relies on the following characterization of groups of polynomial growth by Gromov [Gro81].

**Lemma 5.5.** *Any finitely generated discrete group of polynomial growth contains a finitely generated nilpotent subgroup of finite index.*

We also need the following fact.

**Lemma 5.6.** *Let  $G$  be a finitely generated group of polynomial growth. Let  $H$  be a normal subgroup of  $G$  of finite index. Then  $H$  is finitely generated. Let  $U \subset G$  be a finite system of representatives of the cosets  $G/H$  with  $e \in U$ . Let  $T \subset H$  be a finite generating set of  $H$ . Write  $V = U \cup T$ . Then there exists an integer  $N$  such that*

$$\forall m \in \mathbb{N}, \quad V^m \subset UT^{(3N+1)m}.$$

*Proof.* This is given in the proof of [Wol68, Theorem 3.11]. Let  $N$  be an integer large enough such that for all  $u_1^\epsilon, u_2^\eta \in U$  with  $\epsilon, \eta \in \{\pm 1\}$ , there exist  $u \in U$  and  $t \in T^N$  satisfying  $u_1^\epsilon u_2^\eta = ut$ . Then  $N$  satisfies the desired condition.  $\square$

Now we deduce the desired result.

*Proof of Proposition 5.4.* By Theorem 3.1, it suffices to consider the case where  $\alpha$  is an action on  $L_p(G;$

$L_p(\mathcal{M}))$  by translation. By Lemma 5.5, we may find a nilpotent subgroup  $H \subset G$  of finite index. As is explained in [Wol68, Theorem 3.11],  $H$  can be taken normal by replacing  $H$  with  $\bigcap_{s \in G} sHs^{-1}$ . Now let  $T = \{t_k^{(j)} : 1 \leq k \leq l_j, 1 \leq j \leq K\}$  be a finite generating set of the nilpotent group  $H$ , where  $T$  and the indices  $k, j$  are chosen in the same manner as in (5.1). Also, let  $U$  and  $V$  be given as in the previous lemma. Consider  $x \in L_p^+(G; L_p(\mathcal{M}))$ , and write

$$\tilde{A}_n x = \frac{1}{|V^n|} \sum_{s \in V^n} \alpha_s x, \quad n \in \mathbb{N}.$$

Since the operators  $\alpha_s$  extend to positive operators on  $L_p(G; L_p(\mathcal{M}))$ , by Lemma 5.2 and Lemma 5.6, there exists a constant  $c > 0$  for all  $n \in \mathbb{N}$  such that

$$\begin{aligned} \sum_{s \in V^n} \alpha_s x &\leq \sum_{h \in U} \alpha_h \sum_{t \in T^{(3N+1)n}} \alpha_t x \\ &\leq \sum_{h \in U} \alpha_h \sum_{1 \leq j \leq K} \sum_{\substack{1 \leq n_{jk} \leq cn^j \\ 1 \leq k \leq r_j}} \sum_{\substack{1 \leq n_{jk} \leq N_j \\ r_j < k \leq l_j}} \alpha_{t_1^{(1)}}^{n_{11}} \cdots \alpha_{t_1^{(1)}}^{n_{1l_1}} \cdots \alpha_{t_1^{(K)}}^{n_{K1}} \cdots \alpha_{t_1^{(K)}}^{n_{Kl_K}} x. \end{aligned}$$

Recall that by Lemma 5.3 we may find a constant  $c' > 0$  such that

$$|V^n| \geq c' n^{\sum_{j=1}^K j r_j}.$$

So we may find a constant  $c'' > 0$  satisfying

$$\begin{aligned} \tilde{A}_n x &\leq c'' \sum_{h \in U} \alpha_h \frac{1}{n^{\sum_{j=1}^K j r_j}} \\ &\quad \times \sum_{1 \leq j \leq K} \sum_{\substack{1 \leq n_{jk} \leq cn^j \\ 1 \leq k \leq r_j}} \sum_{\substack{1 \leq n_{jk} \leq N_j \\ r_j < k \leq l_j}} \alpha_{t_1^{(1)}}^{n_{11}} \cdots \alpha_{t_1^{(1)}}^{n_{1l_1}} \cdots \alpha_{t_1^{(K)}}^{n_{K1}} \cdots \alpha_{t_1^{(K)}}^{n_{Kl_K}} x. \end{aligned}$$

Note that by [JX07, Theorem 4.1], for each  $1 \leq j \leq K$  and  $1 \leq k \leq r_j$  there exists a constant  $C_p$  depending only on  $p$  such that

$$\left\| \sup_n^+ \frac{1}{cn^j} \sum_{l=1}^{cn^j} \alpha_{t_k^{(j)}}^l x \right\|_p \leq C_p \|x\|_p, \quad x \in L_p(G; L_p(\mathcal{M})).$$

Applying the inequality iteratively, we obtain a constant  $C'_p > 0$  such that

$$\left\| \sup_n^+ \tilde{A}_n x \right\|_p \leq C'_p \|x\|_p, \quad x \in L_p(G; L_p(\mathcal{M})).$$

Since  $S$  and  $V$  are both finite, we may find two integers  $k$  and  $k'$  with

$$S \subset V^k, \quad V \subset S^{k'}.$$

So the strong type  $(p, p)$  inequality for  $A_n$  follows as well.  $\square$

## 6. INDIVIDUAL ERGODIC THEOREMS

In this section, we apply the maximal inequalities to study the pointwise ergodic convergence in Theorems 1.1 and 1.2.

We will use the following analogue for the noncommutative setting of the usual almost everywhere convergence. The definition is introduced by Lance [Lan76] (see also [Jaj85]).

**Definition 6.1.** Let  $\mathcal{M}$  be a von Neumann algebra equipped with a normal semifinite faithful trace  $\tau$ . Let  $x_n, x \in L_0(\mathcal{M})$ . We say that  $(x_n)_{n \geq 1}$  converges *bilaterally almost uniformly* (b.a.u. for short) to  $x$  if for every  $\varepsilon > 0$  there is a projection  $e \in \mathcal{M}$  such that

$$\tau(e^\perp) < \varepsilon \quad \text{and} \quad \lim_{n \rightarrow \infty} \|e(x_n - x)e\|_\infty = 0,$$

and that it converges *almost uniformly* (a.u. for short) to  $x$  if for every  $\varepsilon > 0$  there is a projection  $e \in \mathcal{M}$  such that

$$\tau(e^\perp) < \varepsilon \quad \text{and} \quad \lim_{n \rightarrow \infty} \|(x_n - x)e\|_\infty = 0.$$

In the case of classical probability spaces, the definition above is equivalent to the usual almost everywhere convergence in terms of Egorov's theorem.

Now let  $G$  be an amenable locally compact group, and let  $(F_n)_{n \geq 1}$  be a Følner sequence in  $G$ . Let  $1 \leq p \leq \infty$ . Assume that  $\alpha = (\alpha_s)_{s \in G}$  is an action on  $L_p(\mathcal{M})$  which satisfies  $(\mathbf{A}_1^p)$ – $(\mathbf{A}_3^p)$ . Denote by  $A_n$  the corresponding averaging operators

$$A_n x = \frac{1}{m(F_n)} \int_{F_n} \alpha_s x \, dm(s), \quad x \in L_p(\mathcal{M}).$$

We keep the notation  $\mathcal{F}_p \subset L_p(\mathcal{M})$  and  $P$  introduced in Section 2.2.

We first consider the case where  $\alpha$  extends to an action on  $L_1(\mathcal{M}) + \mathcal{M}$ . In this case we use a standard argument for b.a.u. convergences adapted from [Hon17, JX07], and [Yea77]. The following lemma from [DJ04] will be useful.

**Lemma 6.2.** *Let  $1 \leq p < \infty$ . If  $(x_n) \in L_p(\mathcal{M}; c_0)$ , then  $x_n$  converges b.a.u. to 0. If  $(x_n) \in L_p(\mathcal{M}; c_0^c)$  with  $2 \leq p < \infty$ , then  $x_n$  converges a.u. to 0.*

We will also use the following noncommutative analogue of the Banach principle given by [Lit17] and [CL16, Theorem 3.1].

**Lemma 6.3.** *Let  $1 \leq p < \infty$ , and let  $S = (S_n)_{n \geq 1}$  be a sequence of additive maps from  $L_p^+(\mathcal{M})$  to  $L_0^+(\mathcal{M})$ . Assume that  $S$  is of weak type  $(p, p)$ . Then the set*

$$\mathcal{C} = \{x \in L_p(\mathcal{M}) : (S_n x)_{n \geq 1} \text{ converges a.u.}\}$$

*is closed in  $L_p(\mathcal{M})$ .*

**Proposition 6.4.** *Assume that  $\alpha = (\alpha_s)_{s \in G}$  is an action well defined on  $\bigcup_{1 \leq p \leq \infty} L_p(\mathcal{M})$  which satisfies  $(\mathbf{A}_1^p)$ – $(\mathbf{A}_3^p)$  for every  $1 \leq p \leq \infty$ . Let  $(A_n)_{n \geq 1}$  be as above, and let  $1 \leq p_0 < p_1 \leq \infty$ . Assume that  $(A_n)_{n \geq 1}$  is of strong type  $(p, p)$  for all  $p_0 < p < p_1$ .*

- (1) *For all  $x \in L_p(\mathcal{M})$  with  $p_0 < p \leq 2p_0$ ,  $(A_n x - Px)_{n \geq 1} \in L_p(\mathcal{M}; c_0)$ , and hence  $(A_n x)_{n \geq 1}$  converges b.a.u. to  $Px$ .*
- (2) *For all  $x \in L_p(\mathcal{M})$  with  $2p_0 < p < p_1$ ,  $(A_n x - Px)_{n \geq 1} \in L_p(\mathcal{M}; c_0^c)$ , and hence  $(A_n x)_{n \geq 1}$  converges a.u. to  $Px$ .*

*Proof.* According to the splitting (2.4) and the discussion after it, we know that

$$S = \text{span}\{x - \alpha_s x : s \in G, x \in L_1(\mathcal{M}) \cap \mathcal{M}\}$$

is dense in  $\overline{(\text{Id} - P)(L_p(\mathcal{M}))}$  for all  $1 \leq p < \infty$ . Also, observe that for all  $x \in S$ ,

$$(6.1) \quad \lim_{n \rightarrow \infty} A_n x = 0 \text{ a.u., } \quad x \in S.$$

To see this, take an arbitrary  $x \in S$  of the form  $x = y - \alpha_{s_0} y$  for some  $s_0 \in G$  and  $y \in L_1(\mathcal{M}) \cap \mathcal{M}$ . Then

$$\begin{aligned} A_n x &= \frac{1}{m(F_n)} \int_{F_n} (\alpha_s y - \alpha_{s s_0} y) \, dm(s) \\ &= \frac{1}{m(F_n)} \int_{F_n \setminus (F_n s_0 \cap F_n)} \alpha_s y \, dm(s) - \frac{1}{m(F_n)} \int_{F_n s_0 \setminus (F_n s_0 \cap F_n)} \alpha_s y \, dm(s). \end{aligned}$$

Therefore, according to  $(\mathbf{A}_2^\infty)$ ,

$$(6.2) \quad \|A_n x\|_\infty \leq \frac{m(F_n \triangle F_n s_0)}{m(F_n)} \|y\|_\infty \sup_{s \in G} \|\alpha_s\|_{B(\mathcal{M})},$$

which converges to 0 as  $n \rightarrow \infty$  according to the Følner condition. This therefore yields the a.u. convergence of  $(A_n x)_n$  in (6.1), as desired.

Now we prove assertion (1). Take  $x \in L_p(\mathcal{M})$ . Since  $S$  is dense in  $(\text{Id} - P)(L_p(\mathcal{M}))$ , there are  $x_k \in S$  such that

$$\lim_{k \rightarrow \infty} \|x - Px - x_k\|_p = 0.$$

Since  $(A_n)_{n \geq 1}$  is of strong type  $(p, p)$ , there exists a constant  $C > 0$  independent of  $x$  such that

$$\|(A_n x - Px - A_n x_k)_n\|_{L_p(\mathcal{M}; \ell_\infty)} \leq C \|x - Px - x_k\|_p.$$

Thus,

$$\lim_{k \rightarrow \infty} (A_n x_k)_n = (A_n x - Px)_n \quad \text{in } L_p(\mathcal{M}; \ell_\infty).$$

Since  $L_p(\mathcal{M}; c_0)$  is closed in  $L_p(\mathcal{M}; \ell_\infty)$ , it suffices to show that  $(A_n x_k)_n \in L_p(\mathcal{M}; c_0)$  for all  $k$ . To this end, we take an arbitrary  $z \in S$  of the form  $z = y - \alpha_{s_0} y$  for some  $s_0 \in G$  and  $y \in L_1(\mathcal{M}) \cap \mathcal{M}$ . Take some  $p_0 < q < p$ . Note that  $z \in L_q(\mathcal{M})$  and that  $(A_n)_n$  is of strong type  $(q, q)$  by assumption, so  $(A_n(z))_n$  belongs to  $L_q(\mathcal{M}; \ell_\infty)$ . Then by (2.1) and (6.2), for any  $m < n$ ,

$$\begin{aligned} \|\sup_{m \leq j \leq n}^+ A_j z\|_p &\leq \sup_{m \leq j \leq n} \|A_j z\|_\infty^{1-\frac{q}{p}} \|\sup_{m \leq j \leq n}^+ A_j z\|_q^{\frac{q}{p}} \\ &\leq \sup_{m \leq j \leq n} \left( \frac{m(F_j \Delta F_j s_0)}{m(F_j)} \|y\|_\infty \right)^{1-\frac{q}{p}} \|\sup_{m \leq j \leq n}^+ A_j z\|_q^{\frac{q}{p}}. \end{aligned}$$

Thus,  $\|\sup_{j \geq m}^+ A_j z\|_p$  tends to 0 as  $m \rightarrow \infty$ . Therefore, the finite sequence  $(A_1 z, \dots, A_m z, 0, \dots)$  converges to  $(A_n z)_n$  in  $L_p(\mathcal{M}; \ell_\infty)$  as  $m \rightarrow \infty$ . As a result  $(A_n(z))_n \in L_p(\mathcal{M}; c_0)$ , as desired.

Assertion (2) is similar. It suffices to note that by the classical Kadison inequality in [Kad52],

$$(A_n x)^2 \leq A_n(x^2) \sup_{s \in G} \|\alpha_s\|_{B(\mathcal{M})}, \quad x \in L_p(\mathcal{M}) \cap \mathcal{M}, x \geq 0,$$

and hence by the strong type  $(p, p)$  inequality and the definition of  $L_p(\mathcal{M}; \ell_\infty^c)$ , there exists a constant  $C$  such that

$$\|(A_n x)_{n \geq 1}\|_{L_p(\mathcal{M}; \ell_\infty^c)} \leq \|((A_n x^2)_{n \geq 1})\|_{L_{p/2}(\mathcal{M}; \ell_\infty)}^{1/2} \leq C \|x\|_p, \quad x \in L_p^+(\mathcal{M}).$$

Then a similar argument yields that  $(A_n x - Px)_{n \geq 1} \in L_p(\mathcal{M}; c_0^c)$ .  $\square$

*Remark 6.5.* The above argument certainly works as well for a Følner sequence  $(F_r)_{r > 0}$  indexed by  $r \in \mathbb{R}_+$ , provided that  $r \mapsto A_r x$  is continuous for  $x \in L_p(\mathcal{M})$ .

As a corollary, we obtain the individual ergodic theorems for actions on  $L_1(\mathcal{M}) + \mathcal{M}$ . We complete the proof of Theorem 1.1.

**Theorem 6.6.** *Let  $d$  be an invariant metric on  $G$ . Assume that  $(G, d)$  satisfies (1.1) and (1.2). Let  $\alpha$  be an action of  $G$  well defined on  $\bigcup_{1 \leq p \leq \infty} L_p(\mathcal{M})$  which satisfies  $(\mathbf{A}_1^p)$ – $(\mathbf{A}_3^p)$  for every  $1 \leq p \leq \infty$ . Denote*

$$A_r x = \frac{1}{m(B_r)} \int_{B_r} \alpha_s x \, dm(s), \quad x \in L_p(\mathcal{M}), r > 0.$$

*Then  $(A_r x)_{r > 0}$  converges a.u. to  $Px$  as  $r \rightarrow \infty$  for all  $1 < p < \infty$ .*

*Moreover, if  $\alpha$  is a continuous action of  $G$  on  $\mathcal{M}$  by  $\tau$ -preserving automorphisms (i.e.,  $\alpha$  satisfies  $(\mathbf{A}')$ ), then  $(A_r x)_{r > 0}$  converges a.u. to  $Px$  as  $r \rightarrow \infty$  for all  $x \in L_1(\mathcal{M})$ .*

*Proof.* Note that for  $1 \leq p \leq 2$  and  $2 < p' < \infty$ ,  $L_p(\mathcal{M}) \cap L_{p'}(\mathcal{M})$  is dense in  $L_p(\mathcal{M})$ , and  $(A_r x)_{r > 0}$  converges a.u. to  $Px$  for all  $x \in L_{p'}(\mathcal{M})$  according to Proposition 6.4. Then the theorem is an immediate consequence of Lemma 6.3 and Theorem 4.5.  $\square$

For word metrics on groups of polynomial growth, it is well known that the associated balls satisfy the Følner condition (see [Bre14, Tes07]). Together with Corollary 4.11 we obtain the following result. This also proves the a.u. convergence on  $L_1(\mathcal{M})$  stated in Theorem 1.2.

**Theorem 6.7.** *Assume that  $G$  is of polynomial growth, and is generated by a symmetric compact subset  $V$ . Let  $\alpha$  be an action of  $G$  well defined on  $\bigcup_{1 \leq p \leq \infty} L_p(\mathcal{M})$  which satisfies  $(\mathbf{A}_1^p)$ – $(\mathbf{A}_3^p)$  for every  $1 \leq p \leq \infty$ . Denote*

$$A_n x = \frac{1}{m(V^n)} \int_{V^n} \alpha_s x \, dm(s), \quad x \in L_1(\mathcal{M}), n \in \mathbb{N}.$$

*Then  $(A_n x)$  converges a.u. to  $Px$  for all  $1 < p < \infty$ .*

*Moreover, if  $\alpha$  is a continuous  $\tau$ -preserving action of  $G$  on  $\mathcal{M}$  such that  $\alpha_s$  is a positive isometry on  $\mathcal{M}$  for each  $s \in G$ , then  $(A_n x)$  converges a.u. to  $Px$  for all  $x \in L_1(\mathcal{M})$ .*

Also, it is obvious that an increasing sequence of compact subgroups always satisfies the Følner condition. Together with Theorem 4.7 we obtain the following.

**Theorem 6.8.** *Let  $G$  be an increasing union of compact subgroups  $(G_n)_{n \geq 1}$ . Let  $\alpha$  be an action of  $G$  well defined on  $\bigcup_{1 \leq p \leq \infty} L_p(\mathcal{M})$  which satisfies  $(\mathbf{A}_1^p)$ – $(\mathbf{A}_3^p)$  for every  $1 \leq p \leq \infty$ . Denote*

$$A_n x = \frac{1}{m(G_n)} \int_{G_n} \alpha_s x \, dm(s), \quad x \in L_p(\mathcal{M}), n \in \mathbb{N}.$$

*Then  $(A_n x)$  converges a.u. to  $Px$  for all  $1 < p < \infty$ .*

*Moreover, if  $\alpha$  is a continuous action of  $G$  on  $\mathcal{M}$  by  $\tau$ -preserving automorphisms (i.e.,  $\alpha$  satisfies  $(\mathbf{A}')$ ), then  $(A_n x)$  converges a.u. to  $Px$  for all  $x \in L_1(\mathcal{M})$ .*

Note that all the above arguments rely on the assumption that the action  $\alpha$  extends to a uniformly bounded action on  $L_\infty(\mathcal{M})$  with conditions  $(\mathbf{A}_1^\infty)$ – $(\mathbf{A}_3^\infty)$ , though our strong type  $(p, p)$  inequalities in previous sections do not require this assumption. Also, in general this assumption does not hold for bounded representations on one fixed  $L_p$ -space. In Theorem 6.11 we will give a stronger result for Følner sequences associated with doubling conditions. This also completes the proof of Theorem 1.2.

**Lemma 6.9.** *Let  $(X, d, \mu)$  be a metric measure space satisfying the doubling condition (4.1). Take  $i \in \mathbb{N}$  and  $k \leq 2^i$ . Then there exists  $2^i \leq r_i < 2^{i+1}$  such that*

$$\mu(B(x, r_i + k) \setminus B(x, r_i)) \leq Ck\mu(B(x, r_i))/r_i,$$

*where  $C$  depends only on the doubling constant.*

*Proof.* The result and the argument are adapted from [Tes07, Proposition 17]. For each  $r \in \mathbb{N}$ , we denote

$$S(x, r) = B(x, r + k) \setminus B(x, r).$$

Then

$$\bigcup_{n=0}^{\lfloor \frac{2^i}{k} \rfloor} S(x, 2^i + nk) \subset B(x, 2^{i+1}).$$

Therefore,

$$\frac{2^i}{k} \inf_{0 \leq n \leq \lfloor \frac{2^i}{k} \rfloor} \mu(S(x, 2^i + nk)) \leq \mu(B(x, 2^{i+1})).$$

Thus, the lemma follows thanks to the doubling condition (4.1).  $\square$

**Lemma 6.10** ([Bre14, Theorem 1.1], [Tes07, Corollary 10]). *Let  $G$  be a locally compact group of polynomial growth, generated by a symmetric compact subset  $V$ . Then*

$$\lim_{n \rightarrow \infty} \frac{m(V^n)}{n^{d(G)}} = c,$$

*where  $d(G)$  is the rank of  $G$  and  $c$  is a constant depending on  $V$ . And there exist  $\delta > 0$  and a constant  $C$  such that*

$$\frac{m(V^{n+1} \setminus V^n)}{m(V^n)} \leq Cn^{-\delta}, \quad n \geq 1.$$

**Theorem 6.11.** *Fix  $1 < p < \infty$ . Let  $\alpha = (\alpha_s)_{s \in G}$  be an action on  $L_p(\mathcal{M})$  which satisfies  $(\mathbf{A}_1^p)$ – $(\mathbf{A}_3^p)$ .*

- (1) Assume that there exists an invariant metric  $d$  on  $G$  and that  $(G, d)$  satisfies (1.1) and (1.2). Denote

$$A_r x = \frac{1}{m(B_r)} \int_{B_r} \alpha_s x \, dm(s), \quad x \in L_p(\mathcal{M}), r > 0.$$

Then there exists a lacunary sequence  $(r_k)_{k \geq 1}$  with  $2^k \leq r_k < 2^{k+1}$  such that  $(A_{r_k} x)_{k \geq 1}$  converges b.a.u. to  $Px$  for all  $x \in L_p(\mathcal{M})$ . If additionally  $p \geq 2$ , then  $(A_{r_k} x)_{k \geq 1}$  converges a.u. to  $Px$  for all  $x \in L_p(\mathcal{M})$ .

- (2) Assume that  $G$  is a locally compact group of polynomial growth, generated by a symmetric compact subset  $V$ . Then the sequence

$$A_n x = \frac{1}{m(V^n)} \int_{V^n} \alpha_s x \, dm(s), \quad n \in \mathbb{N},$$

converges b.a.u. to  $Px$  for all  $x \in L_p(\mathcal{M})$ .

- (3) Assume that  $G$  is an increasing union of compact subgroups  $(G_n)_{n \geq 1}$ . Then the sequence

$$A_n x = \frac{1}{m(G_n)} \int_{G_n} \alpha_s x \, dm(s), \quad n \in \mathbb{N},$$

converges a.u. to  $Px$  for all  $x \in L_p(\mathcal{M})$ .

*Proof.* (1) By Lemma 6.9, there exists  $(r_i)_{i \geq 1}$  such that  $2^i \leq r_i \leq 2^{i+1}$  and such that

$$m(B_{r_i} \setminus B_{r_i - (3/2)^i}) \leq C(3/2)^i m(B_{r_i}) / r_i.$$

That is to say,

$$(6.3) \quad \frac{m(B_{r_i} \setminus B_{r_i - (3/2)^i})}{m(B_{r_i})} \leq C(3/4)^i.$$

We show that  $(A_{r_i} x)_{i \geq 1}$  converges b.a.u. to  $Px$ . By (2.4) and Lemma 6.2, it suffices to show that  $(A_{r_i} x)_{i \geq 1} \in L_p(\mathcal{M}; c_0)$  for  $x \in \mathcal{F}_p^\perp$ . By Theorem 4.5, it is enough to consider the case where  $x = y - \alpha_{s_0} y$  with  $y \in L_p^+(\mathcal{M})$  and  $s_0 \in G$ . Indeed, if  $(A_{r_i} x)_{i \geq 1} \in L_p(\mathcal{M}; c_0)$  for  $x$ 's of the aforementioned form  $y - \alpha_{s_0} y$ , then the same holds for all  $x \in \text{span}\{x - \alpha_s x : s \in G, x \in L_p(\mathcal{M})\}$ . Now by the definition of  $\mathcal{F}_p^\perp$ , for any  $\varepsilon > 0$  and  $x \in \mathcal{F}_p^\perp$ , we may find an element  $y \in \text{span}\{x - \alpha_s x : s \in G, x \in L_p(\mathcal{M})\}$  with  $\|x - y\|_p < \varepsilon$ . So, by Theorem 4.5, we see that

$$\|(A_{r_i} x - A_{r_i} y)\|_{L_p(\mathcal{M}; \ell_\infty)} \leq C\|x - y\|_p < C\varepsilon$$

with a constant  $C$  independent of  $x$  and  $y$ . Thus,  $(A_{r_i} x)$  belongs to  $L_p(\mathcal{M}; c_0)$  since  $L_p(\mathcal{M}; c_0)$  is a Banach space. In the following we prove the claim  $(A_{r_i} x)_{i \geq 1} \in L_p(\mathcal{M}; c_0)$  with  $x = y - \alpha_{s_0} y$ . Denote  $|s_0| = d(e, s_0)$ . Note that

$$A_r x = A_r^1 y - A_r^2 y,$$

where

$$A_r^1 y = \frac{1}{m(B_r)} \int_{B_r \setminus (B_r \cap B_{r, s_0})} \alpha_s y \, dm(s),$$

$$A_r^2 y = \frac{1}{m(B_r)} \int_{(B_r, s_0) \setminus (B_r \cap B_{r, s_0})} \alpha_s y \, dm(s).$$

By (6.3) we have for  $i$  so that  $(3/2)^i \geq |s_0|$ ,

$$\|A_{r_i}^1 y\|_p \leq \frac{m(B_{r_i} \setminus B_{r_i - |s_0|})}{m(B_{r_i})} \|y\|_p \leq C(3/4)^i \|y\|_p.$$

On the other hand, for any  $m \leq j \leq n$ ,

$$A_{r_j}^1 y = [(A_{r_j}^1 y)^p]^{1/p} \leq \left[ \sum_{m \leq i \leq n} (A_{r_i}^1 y)^p \right]^{1/p},$$

and by the previous argument

$$\left\| \left[ \sum_{m \leq i \leq n} A_{r_i}^1 y^p \right]^{1/p} \right\|_p \leq \left[ \sum_{m \leq i \leq n} \|A_{r_i}^1 y\|_p^p \right]^{1/p} \leq C(3/4)^m \|y\|_p.$$

Hence,  $\|(A_{r_j}^1 y)_{m \leq j \leq n}\|_{L_p(\mathcal{M}; c_0)}$  tends to 0 as  $m, n \rightarrow \infty$ . Similarly,  $(A_{r_i}^2 y)_{i \geq 1}$  converges in the same manner. Therefore,  $(A_{r_i} x)_{i \geq 1} \in L_p(\mathcal{M}; c_0)$ , as desired.

Moreover, if  $p \geq 2$ , then

$$(A_{r_j}^1 y)^2 = [(A_{r_j}^1 y)^p]^{2/p} \leq \left[ \sum_{m \leq i \leq n} (A_{r_i}^1 y)^p \right]^{2/p},$$

and hence we can find contractions  $u_j \in L_\infty(\mathcal{M})$  such that for  $m$  large enough,

$$A_{r_j}^1 y = u_j \left[ \sum_{m \leq i \leq n} (A_{r_i}^1 y)^p \right]^{1/p} \quad \text{with} \quad \left\| \left[ \sum_{m \leq i \leq n} (A_{r_i}^1 y)^p \right]^{1/p} \right\|_p \leq C(3/4)^m \|y\|_p.$$

Therefore,  $\|(A_{r_j}^1 y)_{m \leq j \leq n}\|_{L_p(\mathcal{M}; c_0^c)}$  tends to 0 as  $m, n \rightarrow \infty$ , and  $(A_{r_i}^1 y)_{i \geq 1}$  converges a.u. to 0 according to Lemma 6.2. Similarly,  $(A_{r_i}^2 y)_{i \geq 1}$  converges in the same manner. Thus, we obtain that  $(A_{r_i} x)_{i \geq 1}$  converges a.u. to 0. Then by (2.4) and Lemma 6.3,  $(A_{r_i} x)_{i \geq 1}$  converges a.u. for all  $x \in L_p(\mathcal{M})$ .

(2) We keep the notation  $x, y, s_0$  in (1), and denote as before

$$A_k^1 y = \frac{1}{m(V^k)} \int_{(V^k) \setminus (V^k \cap V^{k s_0})} \alpha_s y \, dm(s), \quad y \in L_p(\mathcal{M}), k \geq 1.$$

Write  $\delta' = [\delta^{-1}] + 1$ . Note that by Lemma 6.10, there exists a constant  $C > 0$  such that for  $y \in L_p(\mathcal{M})$  and  $k \geq 1$ ,

$$(6.4) \quad \|A_{k^{\delta'}}^1 y\|_p \leq \frac{m(V^{k^{\delta'}} \setminus V^{k^{\delta'} - |s_0|})}{m(V^{k^{\delta'}})} \|y\|_p \leq \frac{C|s_0| \|y\|_p}{k},$$

where  $|s_0| = d(e, s_0)$  and  $d$  refers to the word metric defined in Example 4.6(1). Hence,

$$\left\| \left[ \sum_{m \leq k \leq n} A_{k^{\delta'}}^1 y^p \right]^{1/p} \right\|_p \leq \left[ \sum_{m \leq k \leq n} \|A_{k^{\delta'}}^1 y\|_p^p \right]^{1/p} \leq C|s_0| \left( \sum_{m \leq k \leq n} \frac{1}{k^p} \right)^{1/p} \|y\|_p.$$

Then, by an argument similar to that in (1), we see that  $(A_{n^{\delta'}} x)_{n \geq 1}$  converges b.a.u. to  $Px$  for all  $x \in L_p(\mathcal{M})$ , and if  $p \geq 2$ , then  $(A_{n^{\delta'}} x)_{n \geq 1}$  converges a.u. to  $Px$ .

For the general case, we consider  $x \in L_p^+(\mathcal{M})$ . For each  $k$ , let  $n(k)$  be the number such that  $n(k)^{\delta'} \leq k < (n(k) + 1)^{\delta'}$ . Then

$$\frac{m(V^{n(k)^{\delta'}})}{m(V^k)} A_{n(k)^{\delta'}} x \leq A_k x \leq \frac{m(V^{(n(k)+1)^{\delta'}})}{m(V^k)} A_{(n(k)+1)^{\delta'}} x.$$

Also note that according to Lemma 6.10,  $m(V^{n(k)^{\delta'}})/m(V^k)$  tends to 1. Therefore, it is easy to see from the definition of b.a.u. convergence that  $A_k x$  converges b.a.u. to  $Px$ .

(3) Note that for  $x = y - \alpha_{s_0} y$  with  $y \in L_p(\mathcal{M})$  and  $s_0 \in G$ , we have for  $n$  large enough so that  $G_n \ni s_0$ ,

$$A_n x = \frac{1}{m(G_n)} \int_{G_n} \alpha_s y \, dm(s) - \int_{G_n} \alpha_{s s_0} y \, dm(s) = 0.$$

That is to say,  $A_n x$  converges a.u. to 0 as  $n \rightarrow \infty$ . Then by (2.4) and Lemma 6.3, we see that  $A_n x$  converges a.u. for all  $x \in L_p(\mathcal{M})$ .  $\square$

In particular, the above arguments give the individual ergodic theorem for positive invertible operators on  $L_p$ -spaces.

**Corollary 6.12.** *Let  $1 < p < \infty$ . Let  $T : L_p(\mathcal{M}) \rightarrow L_p(\mathcal{M})$  be a positive invertible operator with positive inverse such that  $\sup_{n \in \mathbb{Z}} \|T^n\| < \infty$ . Denote*

$$A_n = \frac{1}{2n+1} \sum_{k=-n}^n T^k, \quad n \in \mathbb{N}.$$

*Then  $(A_n x)_{n \geq 1}$  converges b.a.u. to  $Px$  for all  $x \in L_p(\mathcal{M})$ . If additionally  $p \geq 2$ , then  $(A_n x)_{n \geq 1}$  converges a.u. to  $Px$ .*

*Proof.* The assertion follows from the proof of Theorem 6.11(2) for the case where  $G$  equals the integer group  $\mathbb{Z}$ . It suffices to notice that in this case we may choose  $\delta = 1$  in Lemma 6.10 and take  $\delta' = 1$  in (6.4).  $\square$

*Remark 6.13.* Note that the above result is not true for  $p = 1$ , even for positive invertible isometries on classical  $L_1$ -spaces (see, e.g., [IT64]). So it is natural to assume that  $p \neq 1$  in the above discussions.

The following conjecture for mean bounded maps is still open. The result for classical  $L_p$ -spaces is given by [MRDIT88].

**Conjecture 6.14.** *Let  $1 < p < \infty$ . Let  $T : L_p(\mathcal{M}) \rightarrow L_p(\mathcal{M})$  be a positive invertible operator with positive inverse such that  $\sup_{n \in \mathbb{Z}} \|\frac{1}{2n+1} \sum_{k=-n}^n T^k\| < \infty$ . Denote*

$$A_n = \frac{1}{2n+1} \sum_{k=-n}^n T^k, \quad n \in \mathbb{N}.$$

*Then  $(A_n x)_{n \geq 1}$  converges b.a.u. to  $Px$  for all  $x \in L_p(\mathcal{M})$ . If additionally  $p \geq 2$ , then  $(A_n x)_{n \geq 1}$  converges a.u. to  $Px$ .*

**Acknowledgment.** The subject of this paper came from a suggestion made several years ago to the authors by Professor Quanhua Xu, whom we thank for many fruitful discussions. We would also like to thank the anonymous referees for their helpful comments and suggestions. Part of the work was done during the Summer Working Seminar on Noncommutative Analysis at Wuhan University (2015) and the Institute for Advanced Study in Mathematics at Harbin Institute of Technology (2016 and 2017).

This work was supported by National Science Foundation (NSF) of China grant 11431011. Hong's work was partially supported by NSF of China grants 11601396 and 12071355. Liao's work was partially supported by the National Science Foundation and by Coalition for National Science Funding grant 11420101001. Wang's work was partially supported by the European Research Council Advanced Grant "Non-Commutative Distributions in Free Probability" No. 339760, Agence Nationale de la Recherche grant ANR-19-CE40-0002, and a public grant as part of the Fondation Mathématique Jacques Hadamard.

#### REFERENCES

- [AD06] C. Anantharaman-Delaroche. On ergodic theorems for free group actions on noncommutative spaces. *Probab. Theory Related Fields*, 135(4):520–546, 2006.
- [ADAB<sup>+</sup>10] C. Anantharaman-Delaroche, J.-P. Anker, M. Babilot, A. Bonami, B. Demange, S. Grellier, F. Havard, P. Jaming, E. Lesigne, P. Maheux, J.-P. Otal, B. Schapira, and J.-P. Schreiber. *Théorèmes ergodiques pour les actions de groupes*, volume 41 of *Monographies de L'Enseignement Mathématique [Monographs of L'Enseignement Mathématique]*. L'Enseignement Mathématique, Geneva, 2010. With a foreword in English by Amos Nevo.
- [Akc75] M. A. Akcoglu. A pointwise ergodic theorem in  $L_p$ -spaces. *Canad. J. Math.*, 27(5):1075–1082, 1975.
- [Bas72] H. Bass. The degree of polynomial growth of finitely generated nilpotent groups. *Proc. London Math. Soc. (3)*, 25:603–614, 1972.
- [Bek08] T. N. Bekjan. Noncommutative maximal ergodic theorems for positive contractions. *J. Funct. Anal.*, 254(9):2401–2418, 2008.
- [Bek15] B. Bekka. Local rigidity for actions of Kazhdan groups on noncommutative  $L_p$ -spaces. *Internat. J. Math.*, 26(8):1550064, 2015.
- [Bre14] E. Breuillard. Geometry of locally compact groups of polynomial growth and shape of large balls. *Groups Geom. Dyn.*, 8(3):669–732, 2014.

- [Bru73] A. Brunel. Théorème ergodique ponctuel pour un semi-groupe commutatif finiment engendré de contractions de  $L^1$ . *Ann. Inst. H. Poincaré Sect. B (N.S.)*, 9:327–343, 1973.
- [Cal53] A. P. Calderón. A general ergodic theorem. *Ann. of Math. (2)*, 58:182–191, 1953.
- [Cal68] A. P. Calderón. Ergodic theory and translation-invariant operators. *Proc. Nat. Acad. Sci. U.S.A.*, 59:349–353, 1968.
- [CDN78] J.-P. Conze and N. Dang-Ngoc. Ergodic theorems for noncommutative dynamical systems. *Invent. Math.*, 46(1):1–15, 1978.
- [CL16] V. Chilin and S. Litvinov. Individual ergodic theorems for semifinite von Neumann algebras (preprint). Available at arXiv:1607.03452, 2016.
- [Cuc71] I. Cuculescu. Martingales on von Neumann algebras. *J. Multivariate Anal.*, 1(1):17–27, 1971.
- [CW76] R. R. Coifman and G. Weiss. *Transference methods in analysis*. American Mathematical Society, Providence, R.I., 1976. Conference Board of the Mathematical Sciences Regional Conference Series in Mathematics, No. 31.
- [DJ04] A. Defant and M. Junge. Maximal theorems of Menchoff-Rademacher type in non-commutative  $L_q$ -spaces. *J. Funct. Anal.*, 206(2):322–355, 2004.
- [Fen98] G. Fendler. On dilations and transference for continuous one-parameter semigroups of positive contractions on  $L^p$ -spaces. *Ann. Univ. Sarav. Ser. Math.*, 9(1):iv+97, 1998.
- [Gro81] M. Gromov. Groups of polynomial growth and expanding maps. *Inst. Hautes Études Sci. Publ. Math.*, (53):53–73, 1981.
- [HJX10] U. Haagerup, M. Junge, and Q. Xu. A reduction method for noncommutative  $L_p$ -spaces and applications. *Trans. Amer. Math. Soc.*, 362(4):2125–2165, 2010.
- [HK12] T. Hytönen and A. Kairema. Systems of dyadic cubes in a doubling metric space. *Colloq. Math.*, 126(1):1–33, 2012.
- [Hon17] G. Hong. Noncommutative ergodic averages of balls and spheres over Euclidean spaces. *Ergod. Th. Dynam. Sys.*, doi:10.1017/etds.2018.40.
- [HS16] G. Hong and M. Sun. Noncommutative multi-parameter Wiener-Wintner type ergodic theorem. *J. Funct. Anal.*, 275:1100–1137, 2018.
- [HSC93] W. Hebisch and L. Saloff-Coste. Gaussian estimates for Markov chains and random walks on groups. *Ann. Probab.*, 21(2):673–709, 1993.
- [Hu08] Y. Hu. Maximal ergodic theorems for some group actions. *J. Funct. Anal.*, 254(5):1282–1306, 2008.
- [IT64] A. Ionescu Tulcea. Ergodic properties of isometries in  $L^p$  spaces,  $1 < p < \infty$ . *Bull. Amer. Math. Soc.*, 70:366–371, 1964.
- [Jaj85] R. Jajte. *Strong limit theorems in noncommutative probability*, volume 1110 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 1985.
- [JR04] M. Junge and Z.-J. Ruan. Decomposable maps on non-commutative  $L_p$ -spaces. In *Operator algebras, quantization, and noncommutative geometry*, volume 365 of *Contemp. Math.*, pages 355–381. Amer. Math. Soc., Providence, RI, 2004.
- [Jun02] M. Junge. Doob’s inequality for non-commutative martingales. *J. Reine Angew. Math.*, 549:149–190, 2002.
- [JX07] M. Junge and Q. Xu. Noncommutative maximal ergodic theorems. *J. Amer. Math. Soc.*, 20(2):385–439, 2007.
- [Kad52] R. V. Kadison. A generalized Schwarz inequality and algebraic invariants for operator algebras. *Ann. of Math. (2)*, 56:494–503, 1952.
- [Kan78] C. H. Kan. Ergodic properties of Lamperti operators. *Canad. J. Math.*, 30(6):1206–1214, 1978.
- [Küm78] B. Kümmerer. A non-commutative individual ergodic theorem. *Invent. Math.*, 46(2):139–145, 1978.
- [Lan76] E. C. Lance. Ergodic theorems for convex sets and operator algebras. *Invent. Math.*, 37(3):201–214, 1976.
- [Lin01] E. Lindenstrauss. Pointwise theorems for amenable groups. *Invent. Math.*, 146(2):259–295, 2001.
- [Lit14] S. Litvinov. A non-commutative Wiener-Wintner theorem. *Illinois J. Math.*, 58(3):697–708, 2014.
- [Lit17] S. Litvinov. Almost uniform convergence in the noncommutative Dunford–Schwartz ergodic theorem. *C. R. Math. Acad. Sci. Paris*, 355(9):977–980, 2017.
- [Mei03] T. Mei. BMO is the intersection of two translates of dyadic BMO. *C. R. Math. Acad. Sci. Paris*, 336(12):1003–1006, 2003.
- [Mei07] T. Mei. Operator valued Hardy spaces. *Mem. Amer. Math. Soc.*, 188(881):vi+64, 2007.
- [MN91] P. Meyer-Nieberg. *Banach lattices*. Universitext. Springer-Verlag, Berlin, 1991.
- [MRDIT88] F. J. Martin-Reyes and A. De la Torre. The dominated ergodic estimate for mean bounded, invertible, positive operators. *Proc. Amer. Math. Soc.*, 104(1):69–75, 1988.
- [Mus03] M. Musat. Interpolation between non-commutative BMO and non-commutative  $L_p$ -spaces. *J. Funct. Anal.*, 202(1):195–225, 2003.
- [Nev06] A. Nevo. Pointwise ergodic theorems for actions of groups. In *Handbook of dynamical systems. Vol. 1B*, pages 871–982. Elsevier B. V., Amsterdam, 2006.
- [NS94] A. Nevo and E. M. Stein. A generalization of Birkhoff’s pointwise ergodic theorem. *Acta Math.*, 173(1):135–154, 1994.

- [NT10] A. Naor and T. Tao. Random martingales and localization of maximal inequalities. *J. Funct. Anal.*, 259(3):731–779, 2010.
- [Oli12] B. Olivier. Kazhdan’s property  $(T)$  with respect to non-commutative  $L_p$ -spaces. *Proc. Amer. Math. Soc.*, 140(12):4259–4269, 2012.
- [Oli13] B. Olivier. *Rigidity and non-rigidity of group actions on non-commutative  $L_p$  spaces*. PhD thesis, 2013. Thèse de doctorat, Université de Rennes 1, Université européenne de Bretagne.
- [Pat88] A. L. T. Paterson. *Amenability*, volume 29 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 1988.
- [Pis95] G. Pisier. Regular operators between non-commutative  $L_p$ -spaces. *Bull. Sci. Math.*, 119(2):95–118, 1995.
- [Pis98] G. Pisier. Non-commutative vector valued  $L_p$ -spaces and completely  $p$ -summing maps. *Astérisque*, (247):vi+131, 1998.
- [Pis16] G. Pisier. Banach space valued  $H^1$  and BMO. In *Martingales in Banach Spaces*, Cambridge Studies in Advanced Mathematics, pages 263–298. Cambridge University Press, 2016.
- [PX03] G. Pisier and Q. Xu. Non-commutative  $L^p$ -spaces. In *Handbook of the geometry of Banach spaces, Vol. 2*, pages 1459–1517. North-Holland, Amsterdam, 2003.
- [SS83] E. M. Stein and J.-O. Strömberg. Behavior of maximal functions in  $\mathbf{R}^n$  for large  $n$ . *Ark. Mat.*, 21(2):259–269, 1983.
- [Tem67] A. Tempelman. Ergodic theorems for general dynamical systems. *Dokl. Akad. Nauk SSSR*, 176:790–793, 1967.
- [Tem92] A. Tempelman. *Ergodic theorems for group actions*, volume 78 of *Mathematics and its Applications*. Kluwer Academic Publishers Group, Dordrecht, 1992. Informational and thermodynamical aspects, Translated and revised from the 1986 Russian original.
- [Tem15] A. Tempelman. Pointwise ergodic theorems for bounded Lamperti representations of amenable groups. *Proc. Amer. Math. Soc.*, 143(11):4989–5004, 2015.
- [Tes07] R. Tessera. Volume of spheres in doubling metric measured spaces and in groups of polynomial growth. *Bull. Soc. Math. France*, 135(1):47–64, 2007.
- [Wie39] N. Wiener. The ergodic theorem. *Duke Math. J.*, 5(1):1–18, 1939.
- [Wol68] J. A. Wolf. Growth of finitely generated solvable groups and curvature of Riemannian manifolds. *J. Differential Geometry*, 2:421–446, 1968.
- [Yea77] F. J. Yeadon. Ergodic theorems for semifinite von Neumann algebras. I. *J. London Math. Soc. (2)*, 16(2):326–332, 1977.

SCHOOL OF MATHEMATICS AND STATISTICS, WUHAN UNIVERSITY, 430072 WUHAN, CHINA AND HUBEI KEY LABORATORY OF COMPUTATIONAL SCIENCE, WUHAN UNIVERSITY, 430072 WUHAN, CHINA.

*Email address:* guixiang.hong@whu.edu.cn

DEPARTMENT OF MATHEMATICS, TEXAS A&M UNIVERSITY, COLLEGE STATION, TEXAS 77843-3368, USA

*Current address:* Tencent Quantum Laboratory, Shenzhen, China.

*Email address:* liao@hotmail.co.uk

UNIVERSITÄT DES SAARLANDES, FACHRICHTUNG MATHEMATIK, POSTFACH 151150, 66041 SAARBRÜCKEN, GERMANY

*Current address:* Université Paris-Saclay, CNRS, Laboratoire de mathématiques d’Orsay, 91405, Orsay, France.

*Email address:* simeng.wang@universite-paris-saclay.fr