

On the computation of the n th power of a matrix

Nikolaos Halidias
Department of Mathematics
University of the Aegean
Karlovassi 83200 Samos, Greece
email: nikoshalidias@hotmail.com

March 22, 2018

Abstract

In this note we discuss the problem of finding the n th power of a matrix which is strongly connected to the study of Markov chains and others mathematical topics. We observe the known fact (but maybe not well known) that the Cayley-Hamilton theorem is of key importance to this goal. We also demonstrate the classical Gauss elimination technique as a tool to compute the minimum polynomial of a matrix without necessarily know the characteristic polynomial.

Keywords matrix, n th power, Cayley-Hamilton theorem, minimum polynomial

2010 Mathematics Subject Classification

1 Using Cayley-Hamilton theorem to find the n th power of a matrix

Let $A_{m \times m}$ be a real matrix and suppose that we want to compute the n th power. Let us denote by $\delta(k)$ the characteristic polynomial of A . Then it holds that

$$k^n = \delta_A(k)\pi(k) + v(k) \quad (1)$$

where $v(k)$ is a polynomial of degree less or equal to $m - 1$. Using the Cayley-Hamilton theorem we see that $A^n = v(A)$ since $\delta(A) = 0$. Therefore, in order to find the form of A^n we have to find the coefficients of $v(k)$ which are at most m . In this direction we will use the eigenvalues k_1, \dots, k_n (not necessarily distinct) of the matrix A since $\delta(k_i) = 0$. Setting $k = k_i$ we produce some equations involving the unknown coefficients of v . Of course if some of the eigenvalues are of multiplicity two or more then we have to produce some more equations. This can be done by differentiating equation 1 and setting $k = k_i$. We differentiate this equation $m_i - 1$ times (where m_i is the multiplicity of k_i eigenvalue) and each time we set $k = k_i$ in order to produce one more equation. We do this for any eigenvalue k_j with multiplicity m_j . In this fashion we produce m different equations in order to determine the m unknown coefficients of v . One can easily observe that the computation of the coefficients of $v(k)$ is in fact an interpolation problem which has a unique solution (see [2], Thm. 2.2.2). Therefore, we can always find the n th power of matrix by using the Cayley-Hamilton theorem even if the matrix is not diagonalizable. Below, we give some examples.

Example 1 We will compute the n th power of the matrix

$$P = \begin{pmatrix} -3 & 6 & 0 \\ 2 & 1 & 0 \\ 0 & 0 & 3 \end{pmatrix}$$

The eigenvalues are $k_1 = 3$ with multiplicity two and $k_2 = -5$ that is the characteristic polynomial is $\delta_P(k) = (k - 3)^2(k + 5)$.

Therefore, it holds that $k^n = t(k)\delta_P(k) + v(k)$ where $v(k) = ak^2 + bk + c$. We have to produce three equations in order to determine the unknown coefficients. Setting $k_1 = 3$ we obtain the first equation $3^n = 9a + 3b + c$ and setting $k = -5$ we produce another one which is $(-1)^n 5 = 25a - 5b + c$. In order to have one more equation we differentiate the equality $k^n = t(k)\delta_P(k) + v(k)$ and then we set again $k = 3$ because this eigenvalue has multiplicity two. Therefore, we obtain the third equation which is $3^{n-1}n = 6a + b$. Next we solve for a, b, c and thus

$$P^n = aP^2 + bP + c = \begin{pmatrix} 21a - 3a + c & -12a + 6b & 0 \\ -4a + 2b & 13a + b + c & 0 \\ 0 & 0 & 9a + 3b + c \end{pmatrix}$$

Finally, one can verify the result by induction.

Example 2 We will compute the n th power of the following matrix

$$P = \begin{pmatrix} 0 & 1 & 0 \\ 0 & \frac{2}{3} & \frac{1}{3} \\ \frac{1}{3} & 0 & \frac{2}{3} \end{pmatrix}$$

This matrix has the following eigenvalues, $k_1 = 1$, $k_2 = \frac{1+i\sqrt{3}}{6}$ and $k_3 = \frac{1-i\sqrt{3}}{6}$. We have the following equation $k^n = \delta_P(k)t(k) + v(k)$ where $v(k) = ak^2 + bk + c$. Next, we will produce three equations in order to evaluate the unknown coefficients. We transform every complex number in the polar form, that is $x + iy = r(\cos \phi + i \sin \phi)$ where $r = \sqrt{x^2 + y^2}$ and $\phi = \arctan \frac{y}{x}$. Therefore we have that

$$\begin{aligned} k_2 &= \frac{1}{3} \left(\cos \frac{\pi}{3} + i \sin \frac{\pi}{3} \right) \\ k_3 &= \frac{1}{3} \left(\cos \frac{\pi}{3} - i \sin \frac{\pi}{3} \right) \end{aligned}$$

We obtain the following equations by setting $k = k_i$ in the equation $k^n = \delta_P(k)t(k) + v(k)$

$$\begin{aligned} 1^n &= a + b + c \\ \frac{1}{3^n} \left(\cos \frac{n\pi}{3} + i \sin \frac{n\pi}{3} \right) &= a \frac{1}{3^{2n}} \left(\cos \frac{2\pi}{3} + i \sin \frac{2\pi}{3} \right) + b \frac{1}{3} \left(\cos \frac{\pi}{3} + i \sin \frac{\pi}{3} \right) + c \\ \frac{1}{3^n} \left(\cos \frac{n\pi}{3} - i \sin \frac{n\pi}{3} \right) &= a \frac{1}{3^{2n}} \left(\cos \frac{2\pi}{3} - i \sin \frac{2\pi}{3} \right) + b \frac{1}{3} \left(\cos \frac{\pi}{3} - i \sin \frac{\pi}{3} \right) + c \end{aligned}$$

Adding the second equation to the third we produce the equality

$$\frac{1}{3^n} \cos \frac{n\pi}{3} = a \frac{1}{3^{2n}} \cos \frac{2\pi}{3} + b \frac{1}{3} \cos \frac{\pi}{3} + c$$

while subtracting the third equation from the second we produce the following

$$\frac{1}{3^n} \sin \frac{n\pi}{3} = a \frac{1}{3^2} \sin \frac{2\pi}{3} + b \frac{1}{3} \sin \frac{\pi}{3}$$

Finally, we have the following system of equations

$$\begin{aligned} 1^n &= a + b + c \\ \frac{1}{3^n} \cos \frac{n\pi}{3} &= a \frac{1}{3^2} \cos \frac{2\pi}{3} + b \frac{1}{3} \cos \frac{\pi}{3} + c \\ \frac{1}{3^n} \sin \frac{n\pi}{3} &= a \frac{1}{3^2} \sin \frac{2\pi}{3} + b \frac{1}{3} \sin \frac{\pi}{3} \end{aligned}$$

Solving for a, b, c we obtain

$$\begin{aligned} a &= -\frac{3\sqrt{3}}{7} \left(\sqrt{3} \frac{1}{3^n} \cos \frac{n\pi}{3} - \sqrt{3} + 5 \frac{1}{3^n} \sin \frac{n\pi}{3} \right) \\ b &= \frac{\sqrt{3}}{7} \left(\sqrt{3} \frac{1}{3^n} \cos \frac{n\pi}{3} - \sqrt{3} + 19 \frac{1}{3^n} \sin \frac{n\pi}{3} \right) \\ c &= \frac{\sqrt{3}}{21} \left(6\sqrt{3} \frac{1}{3^n} \cos \frac{n\pi}{3} + \sqrt{3} - 12 \frac{1}{3^n} \sin \frac{n\pi}{3} \right) \end{aligned}$$

and therefore $P^n = aP^2 + bP + cI_{2 \times 2}$. One can verify the result by induction.

2 Minimum polynomial

Recall that the minimum polynomial of A is the polynomial $q(k)$ of less degree such that $q(A) = 0$. Therefore one can use the minimum polynomial rather than the characteristic one in order to compute the n th power of the matrix A .

We will demonstrate the classical Gauss elimination procedure in order to find the minimum polynomial.

Let the minimum polynomial has the form

$$q(k) = k^r + a_{r-1}k^{r-1} + \dots + a_0$$

$r \leq m$.

Obviously the relation

$$b_{r-1}A^{r-1} + \dots + b_0I_{m \times m} = 0_{m \times m} \quad (2)$$

drive us to the conclusion that $b_{r-1} = b_{r-2} = \dots = b_0 = 0$ otherwise there will be a polynomial \hat{q} such that $\hat{q}(A) = 0$ with less degree than the minimum polynomial which is a contradiction. That means that the matrices $I_{m \times m}, A, \dots, A^{r-1}$ are linearly independent while the matrices $I_{n \times n}, A, \dots, A^{r-1}, A^r$ are linearly dependent. Obviously the matrices

$$I_{m \times m}, A, \dots, A^{r-1}, A^r, A^{r+1}, \dots, A^m$$

are also linearly dependent.

Let $C_{m^2 \times r}$ the matrix that has in its first column the identity matrix I , that is at the first m places of the first column of C we put the first column of I , next at the next m places of the first column of C we put the second column of I and so on. We do the same for the matrices A, A^2, \dots, A^{r-1} . This matrix is the matrix of the system 2. Since the matrices $I, A, A^2, \dots, A^{r-1}$ are linearly dependent then the reduced row echelon form of C will have the following form

$$\hat{C} = \begin{pmatrix} 1 & 0 & 0 & \cdots \\ 0 & 1 & 0 & \cdots \\ 0 & 0 & 1 & \cdots \\ \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

where the number of leading 1 equals to r .

Let now $D_{m^2 \times (m+1-r)}$ the matrix that has in its columns the matrices A^r, \dots, A^m with the same fashion as with C . We construct next the matrix $B = (C|D)$ (which is in fact the matrix of system 3 below) and we compute the reduced row echelon form. Obviously, the number of the leading 1 equals r again and these are located at the r first columns of B .

That means that the system

$$a_n A^n + a_{n-1} A^{n-1} + \cdots + a_0 I_{n \times n} = 0_{n \times n} \quad (3)$$

has infinite many solutions with $n + 1 - r$ free parameters. Setting $a_r = 1, a_{r+1} = a_{r+2} = \cdots = a_n = 0$ and solving for the others we obtain the minimum polynomial

$$q(k) = a_0 + a_1 k + \cdots + a_{r-1} k^{r-1} + k^r$$

One can easily verify that the minimum polynomial is as follows

$$q(k) = k^r - \hat{B}_{r,r+1} k^{r-1} - \hat{B}_{r-1,r+1} k^{r-2} - \cdots - \hat{B}_{1,r+1}$$

Example 3 *Let the matrix*

$$A = \begin{pmatrix} -4 & 2 & 0 \\ -2 & -1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

We will compute the minimum polynomial.

We construct the matrix B by using the matrices I, A, A^2, A^3 . Then

$$B = \begin{pmatrix} 1 & -4 & 12 & -28 \\ 0 & -2 & 10 & -34 \\ 0 & 0 & 0 & 0 \\ 0 & 2 & -10 & 34 \\ 1 & -1 & -3 & 23 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 1 & 1 & 1 & 1 \end{pmatrix}$$

end the reduced row echelon matrix is the following

$$\hat{B} = \begin{pmatrix} 1 & 0 & 0 & 8 \\ 0 & 1 & 0 & -3 \\ 0 & 0 & 1 & -4 \\ 0 & 0 & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots \end{pmatrix}$$

The number of the leading 1 is 3 and that means that the minimum polynomial is of third degree. Thus,

$$q(k) = \delta(k) = k^3 + 4k^2 + 3k - 8$$

and in this case coincides with the characteristic polynomial.

Example 4 Let the matrix

$$A = \begin{pmatrix} -3 & 6 & 0 \\ 2 & 1 & 0 \\ 0 & 0 & 3 \end{pmatrix}$$

We construct the matrix B as before and therefore we have

$$B = \begin{pmatrix} 1 & -3 & 21 & -87 \\ 0 & 2 & -4 & 38 \\ 0 & 0 & 0 & 0 \\ 0 & 6 & -12 & 114 \\ 1 & 1 & 13 & -11 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 1 & 3 & 9 & 27 \end{pmatrix}$$

The reduced row echelon matrix of B is

$$\hat{B} = \begin{pmatrix} 1 & 0 & 15 & -30 \\ 0 & 1 & -2 & 19 \\ 0 & 0 & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots \end{pmatrix}$$

The number of the leading 1 is 2 therefore the degree of the minimum polynomial is 2. Setting $a_2 = 1$, $a_3 = 0$ and solving for the other we get

$$q(k) = k^2 + 2k - 15 = (k + 5)(k - 3)$$

That means that the matrix A has the $k_1 = -5$ and $k_2 = 3$ as eigenvalues.

References

- [1] S. Bialas - M. Bialas, *An algorithm for the calculation of the minimal polynomial*, Bulletin Polish Academy Sciences, vol. 56, 2008.
- [2] P. Davis, *Interpolation and Approximation*, Dover, 1975.
- [3] S. Lang, *Introduction to Linear Algebra*, Springer, 1986.