

Algebraic New Foundations

Paul Gorbow

University of Gothenburg
 Department of Philosophy, Linguistics, and Theory of Science
 Box 200, 405 30 GÖTEBORG, Sweden
 paul.gorbow@gu.se

November 16, 2018

1 Introduction

New Foundations (NF) is a set theory obtained from naive set theory by putting a stratification constraint on the comprehension schema; for example, it proves that there is a universal set V , and the natural numbers are implemented in the Fregean way (i.e. n is implemented as the set of all sets n many elements). NFU (NF with atoms) is known to be consistent through its remarkable connection with models of conventional set theory that admit automorphisms. This connection was discovered by Jensen, who established the equiconsistency of NFU with a weak fragment of ZF, and its consistency with the axiom of choice [Jensen, 1969].

This paper aims to lay the ground for an algebraic approach to the study of NF. A first-order theory, ML_{CAT} , in the language of categories is introduced and proved to be equiconsistent to NF (analogous results are obtained for intuitionistic and classical NF with and without atoms). ML_{CAT} is intended to capture the categorical content ML, the predicative class theory of NF. NF is interpreted in ML_{CAT} through the categorical semantics. Thus, the result enables application of category theoretic techniques to meta-mathematical problems about NF-style set theory. For example, an immediate corollary is that NF is equiconsistent to $\text{NFU} + |V| = |\mathcal{P}(V)|$. This has already been proved in [Crabbé, 2000], but becomes intuitively obvious in the categorical setting of the present paper.

Just like a category of classes has a distinguished subcategory of small morphisms (cf. [ABSS, 2014]), a category modelling ML_{CAT} has a distinguished subcategory of type-level morphisms. This corresponds to the distinction between sets and proper classes in NF. With this in place, the axiom of power objects familiar from topos theory can be appropriately formulated for NF. It turns out that the subcategory of type-level morphisms contains a topos as a natural subcategory.

This paper is organized as follows. Section 2 explains the motivation behind the paper and in section 3 some further background is given. Section 4 introduces the set theories $(\text{I})\text{NF}(\text{U})_{\text{SET}}$ and the class theories $(\text{I})\text{ML}(\text{U})_{\text{CLASS}}$ (the I stands for the intuitionistic version and the U stands for the version with atoms, so four versions of the theories are considered in parallel), and includes a proof of their respective equiconsistency. In section 5, Heyting categories are introduced and the axioms of the novel categorical theory of this paper, $(\text{I})\text{ML}(\text{U})_{\text{CAT}}$, are given. Section 6 consists of an interpretation of $(\text{I})\text{ML}(\text{U})_{\text{CAT}}$ in $(\text{I})\text{ML}(\text{U})_{\text{CLASS}}$. It is only after this that the main original results of the paper are proved. Most importantly, in section 7 category

theoretic reasoning is used to validate the axioms of $(\text{I})\text{NF}(\text{U})_{\text{SET}}$ in the internal language of $(\text{I})\text{ML}(\text{U})_{\text{CAT}}$ through the categorical semantics. This means that $(\text{I})\text{NF}(\text{U})_{\text{SET}}$ is interpretable in $(\text{I})\text{ML}(\text{U})_{\text{CAT}}$. The equiconsistency of NF and $\text{NFU} + |V| = |\mathcal{P}(V)|$ is obtained as a corollary. In section 8, it is shown that every $(\text{I})\text{ML}(\text{U})_{\text{CAT}}$ -category contains a topos as a subcategory. Section 9 poses some questions and suggests further developments.

2 Motivation

NF corresponds closely with the simple theory of types, TST, an extensional version of higher order logic which Chwistek and Ramsey independently formulated as a simplification of Russel and Whitehead’s system in Principia Mathematica. It was from contemplation of TST that Quine introduced NF [Quine, 1937]. Essentially, NF is obtained from TST by forgetting the typing of the relations while retaining the restriction on comprehension induced by the typing (thus avoiding Russel’s paradox). This results in the notion of stratification, see Definition 1 below. Thus NF and NFU resolve an aspect of type theory which may be considered philosophically dissatisfying: Ontologically, it is quite reasonable to suppose that there are relations which can take both individuals and relations as relata. The simplest example is probably the relation of identity. But in type theory, it is not possible to relate entities of different types. We cannot even say that they are unequal. Since the universe of NF or NFU is untyped, such issues disappear. It is therefore not surprising that stratified set theory has attracted attention from philosophers. For example, Cocchiarella applied these ideas to repair Frege’s system [Cocchiarella, 1985] (a similar result is obtained in [Holmes, 2015]), and Cantini applied them to obtain an interesting type-free theory of truth [Cantini, 2015]. Along a similar line of thought, the categorical version of NF and NFU brought forth in this paper may well be helpful for transferring the ideas of stratified set theory to research in formal ontology. In a formal ontology, one may account for what individuals, properties, relations and tropes exist, where properties and relations are considered in an intensional rather than an extensional sense. Roughly, in category theory the objects are non-extensional, but the morphisms are extensional, and this is arguably fitting to the needs of formal ontology.

NFU is also intimately connected with the highly active research on nonstandard models of arithmetic and set theory, as is further explained in the next section. Out of this connection, Feferman proposed a version of NFU as a foundation for category theory, allowing for such unlimited categories as the category of all sets, the category of all groups, the category of all topological spaces, the category of all categories, etc [Feferman, 2006]. This line of research was further pursued by Enayat, McKenzie and the author in [EGM, 2017]. In short, conventional category theory works perfectly fine in a subdomain of the NFU-universe, but the unlimited categories live outside of this subdomain, and their category theoretic properties are unconventional. Even though they are unconventional (usually failing to be cartesian closed), one might argue that nothing is lost by including them in our mathematical universe. These categories remain to be systematically studied.

The need for a categorical understanding of stratified set theory is especially pressing since very little work has been done in this direction. McLarty showed in [McLarty, 1992] that the “category of sets and functions” in NF is not cartesian closed. However, several positive results concerning this category were proved in an unpublished paper by Forster, Lewicki and Vidrine [FLV, 2014]: In particular, they showed that it has a property they call “pseudo-Cartesian closedness”. Similarly, Thomas showed in [Thomas, 2017] that it has a property he calls “stratified Cartesian closedness”. The moral is that it is straight forward to show in INFU, that if A and B are sets, which are respectively isomorphic to sets of singletons A' and B' , then the set of

functions from $\bigcup A'$ to $\bigcup B'$ is an exponential object of A and B . (V is not isomorphic to any set of singletons.) In [FLV, 2014] a generalization of the notion of topos was proposed, with “the category of sets and functions” of NF as an instance. It has however not been proved that the appropriate extension T of this theory (which NF interprets) satisfies $\text{Con}(T) \Rightarrow \text{Con}(\text{NF})$. Using the results of Section 7 of this paper, it seems within reach to obtain that result by canonically extending a model of T to a model of the categorical theory ML_{CAT} introduced here. That line of research would also help carve out exactly what axioms of T are necessary for that result. Moreover, in [FLV, 2014] it was conjectured that any model of T has a subcategory which is a topos. In Section 8 of this paper, it is proved that every model of $(\text{I})\text{ML}(\text{U})_{\text{CAT}}$ has a subcategory which is a topos.

A related direction of research opened up by the present paper is to generalize the techniques of automorphisms of models of conventional set theory, in order to study automorphisms of topoi. The author expects that a rich landscape of models of $(\text{I})\text{ML}(\text{U})_{\text{CAT}}$ would be uncovered from such an enterprise. For example, just like there is a topos in which every function on the reals is continuous, a similar result may be obtainable for IMLU_{CAT} by finding such a topos with an appropriate automorphism. Given the intriguing prospects for founding category theory in stratified set theory, this would open up interesting possibilities for stratified category theoretic foundation of mathematics.

The categorical approach to NF is also promising for helping the metamathematical study of NF. As stated in the introduction, the main result of this paper has the immediate corollary that NF is equiconsistent to $\text{NFU} + |V| = |\mathcal{P}(V)|$. A major open question in the metamathematics of NF is whether NF (or even its intuitionistic counterpart, which has not been shown to be equiconsistent to NF) is consistent relative to a system of conventional set theory (independent proof attempts by Gabbay and Holmes have recently been put forth). So yet a motivation for introducing ML_{CAT} is simply that the flexibility of category theory may make it easier to construct models of ML_{CAT} , than to construct models of NF, thus aiding efforts to prove and/or simplify proofs of $\text{Con}(\text{NF})$.

Since categorical model theory tends to be richer in the intuitionistic setting, an intriguing line of research is to investigate the possibilities for stratified dependent type theory. Dependent type theory is commonly formulated with a hierarchy of universes. In a sense, this hierarchy is inelegant and seemingly redundant, since any proof on one level of the hierarchy can be shifted to a proof on other levels of the hierarchy. Model-theoretically, this can be captured in a model with an automorphism. Since the semantics of type theory tends to be naturally cast in category theory, the understanding arising from the present paper would be helpful in such an effort.

In conclusion, “categorification” tends to open up new possibilities, as forcefully shown by the fruitfulness of topos theory as a generalization of set theory. In the present paper it has already resulted in a simple intuitive proof of the old result of Crabbé stated above. So given the relevance of NF and NFU to type theory, philosophy, nonstandard models of conventional set theory and the foundations of category theory, it is important to investigate how NF and NFU can be expressed as theories in the language of category theory.

3 Background

Algebraic set theory, categorical semantics and categorical logic more generally, have been developed by a large number of researchers. An early pioneering paper of categorical logic is [Lawvere, 1963]; Joyal and Moerdijk started out algebraic set theory with [JM, 1991] and wrote a short book on the subject [JM, 1995]. The present paper is largely influenced by the comprehensive work of Awodey, Butz, Simpson and Streicher embodied in [ABSS, 2014]. It parallels its

approach to an algebraic set theory of categories of classes, but for the NF context. A category of classes \mathbf{C} is a (Heyting) Boolean category with a subcategory \mathbf{S} , satisfying various axioms capturing the notion of ‘smallness’, and with a universal object U , such that every object is a subobject of U . While the axiomatization of categories of classes naturally focuses on the notion of smallness, the axiomatization in this paper focuses on the notion of type level stratification. Like in [ABSS, 2014], a restricted notion of powerobject is obtained, which facilitates interpretation of set theory in the categorical semantics.

NF may be axiomatized as Extensionality + Stratified Comprehension (Ext + SC); it avoids the Russell paradox of naive set theory, by restricting comprehension to stratified formulas, a notion to be defined below. An introduction to NF is given in [Forster, 1995]. For any basic claims about NF in this paper, we implicitly refer to that monograph.

Though the problem of proving the consistency of NF in terms of a traditional ZF-style set theory turned out to be difficult, Jensen proved the consistency of the subsystem NFU, where Ext is weakened to Ext’ (Ext’ is extensionality for non-empty sets, thus inviting atoms), in [Jensen, 1969]. Jensen used Ramsey’s theorem to obtain a particular model of Mac Lane set theory with an automorphism, and it is relatively straight forward to obtain a model of NFU + Infinity + Choice from that model. There are various interesting axioms that can be added to NFU to increase its consistency strength. As the understanding of automorphisms of non-standard models of ZF-style set theories has increased, several results on the consistency strength of such extensions of NFU have been obtained in the work of Solovay [Solovay, 1997], Enayat [Enayat, 2004] and McKenzie [McKenzie, 2015]. In this paper we define NFU as a theory in the language $\{\in, S, \langle -, - \rangle\}$, where S is a predicate distinguishing sets from atoms and $\langle -, - \rangle$ is a primitive pairing function, axiomatized as $\text{Ext}_S + \text{SC}_S + \text{P} + \text{Sethood}$, where Ext_S and SC_S are extensionality and stratified comprehension for sets, P regulates $\langle -, - \rangle$ and Sethood regulates S , to be specified below. $\text{NFU} = \text{Ext}_S + \text{SC}_S + \text{P} + \text{Sethood}$ proves Infinity and is equiconsistent with Mac Lane set theory; $\text{Ext}_S + \text{SC}_S$ is weaker and does not prove Infinity. In this paper, NF is defined as $\text{NFU} + \text{“everything is a set”}$, which (in classical logic) is equivalent to $\text{Ext} + \text{SC}$. An introduction to NFU and extended systems is given in [Holmes, 1998]. For any basic claims about NFU in this paper, we implicitly refer to that monograph.

The theories NF and NFU in intuitionistic logic will be referred to as INF and INFU, respectively. Note that the way NFU and NF are axiomatized in this paper, the intuitionistic versions INFU and INF also satisfy e.g. the axiom of ordered pair. But if INF were axiomatized as $\text{Ext} + \text{SC}$ with intuitionistic logic, as done e.g. in [Dzierzowski, 1995], it is not clear that the resulting intuitionistic theory would be as strong.

NF and NFU also have finite axiomatizations, as shown in [Hailperin, 1944], which clarify that their “categories of sets and functions” are Boolean categories. In this paper certain extensions of the theories of (Heyting) Boolean categories (in the language of category theory) are proved equiconsistent to (I)NF(U), respectively.

4 Stratified set theory and class theory

Let $\mathcal{L}_{\text{SET}} = \{\in, S, \langle -, - \rangle\}$ be the language of set theory augmented with a unary predicate symbol S of “sethood” and a binary function symbol $\langle -, - \rangle$ of “ordered pair”. We introduce notation for the “set-many quantifier”:

$$\mathcal{E}z.\phi \text{ abbreviates } \exists x.(S(x) \wedge \forall z.(z \in x \leftrightarrow \phi(z))),$$

where x is chosen fresh, i.e. not free in ϕ .

Definition 1. Let ϕ be an \mathcal{L}_{SET} -formula. ϕ is *stratified* iff there is a function $s : \text{term}(\phi) \rightarrow \mathbb{N}$, where $\text{term}(\phi)$ is the set of terms occurring in ϕ , such that for any $u, v, w \in \text{term}(\phi)$ and any atomic subformula θ of ϕ ,

- (i) if $u \equiv \langle v, w \rangle$, then $s(u) = s(v) = s(w)$,
- (ii) if $\theta \equiv (u = v)$, then $s(u) = s(v)$,
- (iii) if $\theta \equiv (u \in v)$, then $s(u) + 1 = s(v)$,

where \equiv denotes literal equality (of terms or formulas). Such an s is called a *stratification* of ϕ . $s(u)$ is called the *type* of u . Clearly, if ϕ is stratified, then there is a *minimal* stratification in the sense that $s(v) = 0$ for some variable v occurring in ϕ . Also note that the formula $\langle v, w \rangle = \{\{v\}, \{v, w\}\}$, stipulating that the ordered pair is the Kuratowski ordered pair, is not stratified. Therefore, it is condition (i), read as “type-level ordered pair”, that gives power to axiom P below.

Notation 2. In the axiomatizations below, NFU is the theory thus axiomatized in classical logic, while INFU is the theory thus axiomatized in intuitionistic logic. For brevity I simply write (I)NFU, and similarly for (I)NF, to talk about the intuitionistic and classical theories in parallel. More generally, when I write that (I)XX(U)_K relates to (I)YY(U)_L in some way, I mean that each of the four theories IXXU_K, XXU_K, IXX_K, XX_K relates in that way to IXXU_L, XXU_L, IXX_L, XX_L, respectively. Since we will be proving equiconsistency results between theories in different languages, the language will sometimes be emphasized as a subscript to the name of the theory, thus I write (I)NF(U)_{SET} for the set theoretic theory (I)NF(U).

Axioms 3 ((I)NFU_{SET}).

Extensionality for Sets (Ext _S)	$(S(x) \wedge S(y) \wedge \forall z. z \in x \leftrightarrow z \in y) \rightarrow x = y$
Stratified Comprehension Schema (SC _S)	For all stratified ϕ : $\exists z. \phi(z, y)$
Ordered Pair (P)	$\langle x, y \rangle = \langle x', y' \rangle \leftrightarrow (x = x' \wedge y = y')$
Sethood	$z \in x \rightarrow S(x)$

In order to keep the treatment uniform, I axiomatize (I)NF_{SET} as (I)NFU_{SET} + $\forall x. S(x)$. Obviously, (I)NF can be axiomatized in the language without the predicate S , simply as Ext + SC + P (where Ext and SC are as Ext_S and SC_S, respectively, but without the S -conjuncts). Less obviously, NF proves the negation of Choice [Specker, 1953], which entails the axiom of Infinity, which in turn enables implementation of type-level ordered pairs. So NF can be axiomatized as Ext + SC in the plain language $\{\in\}$ of set theory.

Note that SC_S implies the existence of a universal set, denoted V . In the context of the sethood predicate, it is natural to restrict the definition of subset to sets. So define

$$x \subseteq y \Leftrightarrow_{\text{df}} S(x) \wedge S(y) \wedge \forall z. (z \in x \rightarrow z \in y).$$

The power set, $\mathcal{P}y$, of y is defined as $\{z \mid z \subseteq y\}$, and exists by SC_S. Therefore, only sets are elements of power sets. An important special case of this is that $S(x) \leftrightarrow x \in \mathcal{P}V$. So the axiom $\forall x. S(x)$, yielding (I)NF, may alternatively be written $V = \mathcal{P}V$. In the meta-theory \subseteq and \mathcal{P} are defined in the standard way. When proving the existence of functions (coded as sets of ordered pairs) in (I)NF(U), the type-level requirement of ordered pairs means that the defining formula (in addition to being stratified) needs to have the argument- and value-variable at the same type.

(I)ML(U)_{CLASS} is the impredicative class theory corresponding to (I)NF(U)_{SET}. ML was introduced by Quine in his book [Quine, 1940]. Apparently ML stands for “Mathematical Logic”

(the title of that book). There is both a predicative and an impredicative version of ML, and both are equiconsistent with NF_{SET} , as proved in [Wang, 1950]. One obtains a model of ML simply by taking the power set of a model of NF, along with a natural interpretation that suggests itself, so the proof requires enough strength in the meta-theory to handle sets of the size of the continuum. (The equiconsistency between predicative ML and NF can be proved in a weaker meta-theory that is only strong enough to handle countable sets.) Without difficulty, the proof extends to equiconsistency between each of the theories $(\text{I})\text{ML}(\text{U})_{\text{CLASS}}$ and $(\text{I})\text{NF}(\text{U})_{\text{SET}}$, respectively. For the purpose of completeness, a proof of $\text{Con}((\text{I})\text{NF}(\text{U})_{\text{SET}}) \Rightarrow \text{Con}((\text{I})\text{ML}(\text{U})_{\text{CLASS}})$ is provided below.

The theory $(\text{I})\text{ML}(\text{U})_{\text{CAT}}$, which I introduce in this paper as an algebraic set theory of NF, probably corresponds better to predicative ML. The difficult and interesting direction of my proof of equiconsistency between $(\text{I})\text{ML}(\text{U})_{\text{CAT}}$ and $(\text{I})\text{NF}(\text{U})_{\text{SET}}$ is my interpretation of $(\text{I})\text{NF}(\text{U})_{\text{SET}}$ in $(\text{I})\text{ML}(\text{U})_{\text{CAT}}$; the easy but tedious part is to interpret $(\text{I})\text{ML}(\text{U})_{\text{CAT}}$ in $(\text{I})\text{ML}(\text{U})_{\text{CLASS}}$. The latter could have been done using the predicative class theory instead, but the impredicative version requires slightly less attention to details which are not related to the original results of this paper.

In order to axiomatize $(\text{I})\text{ML}(\text{U})_{\text{CLASS}}$, let us use a two-sorted language $\mathcal{L}_{\text{CLASS}}$ specified as follows. Upper-case variables range over the sort Clatom, and lower-case variables range over the sort Setom (these words are portmanteaus for class-or-atom and set-or-atom, respectively). Setom is a sub-sort of Clatom, i.e. the domain of Clatom includes the domain of Setom. The symbols are as in \mathcal{L}_{SET} plus a new predicate C of classhood. But we need sort-specifications for all symbols: \in is a binary relation symbol on the sort $\text{Setom} \times \text{Clatom}$; $\langle -, - \rangle$ is a function symbol of sort $\text{Setom} \times \text{Setom} \rightarrow \text{Setom}$; S is a predicate symbol on the sort Setom; and C is a predicate symbol on the sort Clatom. Since Setom is a sub-sort of Clatom, the lower-case variables can be used in any place where the upper-case variables are used. We say that an $\mathcal{L}_{\text{CLASS}}$ -formula ϕ has *small variables* iff every variable is lower-case.

Axioms 4 $((\text{I})\text{MLU}_{\text{CLASS}})$.

Extensionality for Classes (Ext_C)	$(C(X) \wedge C(Y) \wedge \forall z.z \in X \leftrightarrow z \in Y) \rightarrow X = Y$
Class Comprehension Schema (CC_C)	For all ϕ : $\exists X.(C(X) \wedge \forall z.(z \in X \leftrightarrow \phi(z)))$
Stratified Comprehension Schema (SC_S)	For all stratified ϕ with small variables: $\exists z.\phi(z)$
Ordered Pair (P)	$\langle x, y \rangle = \langle x', y' \rangle \leftrightarrow (x = x' \wedge y = y')$
Classhood	$z \in X \rightarrow C(X)$
Sethood	$z \in x \rightarrow S(x)$
Small Classhood	$S(x) \leftrightarrow C(x)$

In CC_C , we assume that X is fresh, i.e. not free in ϕ . If $C(x)$, then we say that x is a *class*. If $S(x)$, then we say that x is a *set*. As before, we obtain $(\text{I})\text{ML}_{\text{CLASS}}$ by adding the axiom $\forall x.C(x)$. There is an equivalent axiomatization without the predicate S and the axioms of Sethood and Small Classhood. I have chosen this more detailed presentation because it has the property: If $\mathcal{M} \models (\text{I})\text{ML}(\text{U})_{\text{CLASS}}$, then by first restricting \mathcal{M} to the sort Setom, and second taking the reduct to the language \mathcal{L}_{SET} , we obtain a model of $(\text{I})\text{NF}(\text{U})_{\text{SET}}$. This is easily seen by comparing Axioms 3 to Axioms 4. Thus, it is obvious that $(\text{I})\text{ML}(\text{U})$ interprets $(\text{I})\text{NF}(\text{U})$. We now proceed to establish that the consistency of $(\text{I})\text{NF}(\text{U})$ implies the consistency of $(\text{I})\text{ML}(\text{U})$. The proof is actually trivial, but involves some detail of presentation.

Theorem 5. $\text{Con}((\text{I})\text{NF}(\text{U})_{\text{SET}}) \Leftrightarrow \text{Con}((\text{I})\text{ML}(\text{U})_{\text{CLASS}})$

Proof. The \Leftarrow direction was established above, so we show that $\text{Con}((\text{I})\text{NF}(\text{U})) \Rightarrow \text{Con}((\text{I})\text{ML}(\text{U}))$. We concentrate on the case $\text{Con}(\text{INFU}) \Rightarrow \text{Con}(\text{IMLU})$. Afterwards it will be easy to see the modifications required for the other three cases.

Let \mathbb{P} be a poset (i.e. an intuitionistic Kripke frame), partially ordered by \leq , and let $\mathcal{N} =_{\text{df}} \langle \mathcal{N}_p \mid p \in \mathbb{P} \rangle$ be a Kripke model of INFU, where each \mathcal{N}_p is a classical \mathcal{L}_{SET} -structure $\langle N_p, \in_p^{\mathcal{N}}, S_p^{\mathcal{N}}, P_p^{\mathcal{N}} \rangle$. Define a Kripke model $\mathcal{M} =_{\text{df}} \langle \mathcal{M}_p \mid p \in \mathbb{P} \rangle$, where each \mathcal{M}_p is a classical $\mathcal{L}_{\text{CLASS}}$ -structure $\langle \text{Clatom}_p^{\mathcal{M}}, \text{Setom}_p^{\mathcal{M}}, \in_p^{\mathcal{M}}, S_p^{\mathcal{N}}, C_p^{\mathcal{N}}, P_p^{\mathcal{N}} \rangle$, naturally defined as follows. Let $p \in \mathbb{P}$ be arbitrary. For each $x \in N_p$, let $t_p(x) =_{\text{df}} \{z \in N_p \mid z \in_p^{\mathcal{N}} x\}$.

$$\begin{aligned} \text{Clatom}_p^{\mathcal{M}} &=_{\text{df}} \mathcal{P}(N_p) \\ \text{Setom}_p^{\mathcal{M}} &=_{\text{df}} \{t_p(x) \mid x \in N_p\} \\ X \in_p^{\mathcal{M}} Y &\Leftrightarrow_{\text{df}} \exists x \in Y. t_p(x) = X \\ S_p^{\mathcal{M}} &=_{\text{df}} \{t_p(x) \mid x \in S_p^{\mathcal{N}}\} \\ C_p^{\mathcal{M}} &=_{\text{df}} S_p^{\mathcal{M}} \cup (\text{Clatom}_p^{\mathcal{M}} - \text{Setom}_p^{\mathcal{M}}) \\ P_p^{\mathcal{M}} &=_{\text{df}} \{\langle t_p(x), t_p(y) \rangle, t_p(z) \mid P_p^{\mathcal{N}}(x, y) = z\} \end{aligned}$$

Since \mathcal{N} is a Kripke model, it is easily seen from the construction that \mathcal{M} is a Kripke structure in $\mathcal{L}_{\text{CLASS}}$; more precisely, $\text{Setom}_p^{\mathcal{M}} \subseteq \text{Clatom}_p^{\mathcal{M}}$ for each $p \in \mathbb{P}$, the interpretations of the symbols match their sort-specifications, and if $p \leq q \in \mathbb{P}$, then $\text{Symbol}_p^{\mathcal{M}} \subseteq \text{Symbol}_q^{\mathcal{M}}$ for each $\text{Symbol} \in \{\text{Clatom}, \text{Setom}, \in, S, C, P\}$.

Note that t_p witnesses $\langle N_p, \in_p^{\mathcal{N}}, S_p^{\mathcal{N}}, P_p^{\mathcal{N}} \rangle \cong \langle \text{Setom}_p^{\mathcal{M}}, \in_p^{\mathcal{M}} \upharpoonright_{\text{Setom}_p^{\mathcal{M}}}, S_p^{\mathcal{M}}, P_p^{\mathcal{M}} \rangle$. Thus, \mathcal{M} satisfies SC_S , P and Sethood. Moreover, it is immediate from the construction that \mathcal{M} satisfies Ext_C and Setom Classhood. Hence, it only remains to verify that \mathcal{M} satisfies CC. We show even that each \mathcal{M}_p satisfies CC. Let $\phi(x)$ be an $\mathcal{L}_{\text{CLASS}}$ -formula with free small variable x . Let

$$A =_{\text{df}} \{u \in N_p \mid \mathcal{M}_p \models \phi(t_p(u))\}.$$

The following sequence of implications completes the proof of $\text{Con}(\text{INFU}) \Rightarrow \text{Con}(\text{IMLU})$.

$$\begin{aligned} &A = \{u \in N_p \mid \mathcal{M}_p \models \phi(t_p(u))\} \\ \Rightarrow &\forall u \in N_p. (u \in A \leftrightarrow \mathcal{M}_p \models \phi(t_p(u))) \\ \Rightarrow &\forall z \in \text{Setom}_p^{\mathcal{M}}. (z \in_p^{\mathcal{M}} A \leftrightarrow \mathcal{M}_p \models \phi(z)) \\ \Rightarrow &\forall z \in \text{Setom}_p^{\mathcal{M}}. (\mathcal{M}_p \models (z \in A \leftrightarrow \phi(z))) \\ \Rightarrow &\mathcal{M}_p \models \exists X. \forall z. (z \in X \leftrightarrow (\phi(z))) \end{aligned}$$

$\text{Con}(\text{INF}) \Rightarrow \text{Con}(\text{IML})$ follows from $\langle N_p, \in_p^{\mathcal{N}}, S_p^{\mathcal{N}}, P_p^{\mathcal{N}} \rangle \cong \langle \text{Setom}_p^{\mathcal{M}}, \in_p^{\mathcal{M}} \upharpoonright_{\text{Setom}_p^{\mathcal{M}}}, S_p^{\mathcal{M}}, P_p^{\mathcal{M}} \rangle$. The classical cases $\text{Con}(\text{NF}(\text{U})) \Rightarrow \text{Con}(\text{ML}(\text{U}))$ follow by letting \mathbb{P} be a singleton. \square

In the predicative version of (I)ML(U), the axiom CC_C is modified so that no quantified Clatom-variables are allowed in ϕ . To obtain a model of this theory, it suffices to consider the set of definable subsets of a model of (I)NF(U). Thus, a slightly modified version of the above proof can be carried out for the predicative case in the usual set theory of countable sets.

5 Category theoretic axiomatization

Categories may be viewed as structures in the basic language of category theory. Traditionally, a theory in the first order language of category theory (or an expansion of that language) is formulated as a definition of a class of models. Such definitions, that can be turned into first order axiomatizations, are called elementary. The definitions of classes of categories made in this section are all easily seen to be elementary.

Now follows a presentation of the categorical semantics of first order logic in Heyting (intuitionistic logic) and Boolean (classical logic) categories. A full account can be found e.g. in [Johnstone, 2002, pp. 807-859].

A morphism f is a *cover* iff whenever $f = m \circ g$ for a mono m , then m is an isomorphism. A morphism f has an *image* iff it factors as $f = m \circ e$, where m is a mono with the universal property that if $f = m' \circ e'$ is some factorization with m' mono, then there is a unique k such that $m = m' \circ k$.

Definition 6. A category is a *Heyting category* iff it satisfies the following axioms (HC).

- (F1) It has finite limits.
- (F2) It has images.
- (F3) Covers are stable under pullback.
- (F4) Each Sub_X is a sup-semilattice.
- (F5) For each $f : X \rightarrow Y$, the inverse image functor $f^* : \text{Sub}_Y \rightarrow \text{Sub}_X$ preserves finite suprema and has left and right adjoints: $\exists_f \dashv f^* \dashv \forall_f$.

We call this theory HC. Sub_X and f^* are explained below. One can prove from these axioms that that each Sub_X is a Heyting algebra. A *Boolean category* is a Heyting category such that each Sub_X is a Boolean algebra. We call that theory BC.

A *Heyting (Boolean) functor*, is a functor between Heyting (Boolean) categories that preserves the structure above. \mathbf{C} is a *Heyting (Boolean) subcategory* of \mathbf{D} iff it is a subcategory and the inclusion functor is Heyting (Boolean).

A Heyting category \mathbf{C} has a terminal object 1 and an initial object 0 , as well as a diagonal mono $\Delta_X : X \rightarrow X \times X$, for each X . A *subobject* of an object X is an isomorphism class of monos $m : Y \rightarrow X$ in the slice category \mathbf{C}/X . Sometimes such a subobject is simply denoted Y , although it is an abuse of notation. The axioms (F1)–(F5) ensure that for any object X , the partial order of subobjects of X , denoted $\text{Sub}(X)$, is a Heyting algebra, with constants \perp_X , \top_X and operations \wedge_X , \vee_X , \rightarrow_X (we suppress the subscript when it is clear from the context). Given a morphism $f : X \rightarrow Y$ in \mathbf{C} , the functor $f^* : \text{Sub}(Y) \rightarrow \text{Sub}(X)$ is defined by sending any subobject of Y , represented by $m_B : B \rightarrow Y$, say, to the subobject of X represented by the pullback of m_B along f .

A *structure* (or *model*) \mathcal{M} , in the categorical semantics of \mathbf{C} , in a sorted signature \mathcal{S} , is an assignment of sorts, relation symbols and function symbols of \mathcal{S} to objects, subobjects and morphisms of \mathbf{C} , respectively: Thus, a relation symbol R on the sort $S_1 \times \dots \times S_n$ is assigned to a subobject $\mathcal{M}R \leq \mathcal{M}S_1 \times \dots \times \mathcal{M}S_n$; in particular, the equality symbol $=_S$ on the sort $S \times S$ is assigned to the subobject of $\mathcal{M}S \times \mathcal{M}S$ determined by $\Delta_{\mathcal{M}S} : \mathcal{M}S \rightarrow \mathcal{M}S \times \mathcal{M}S$; and a function symbol $f : S_1 \times \dots \times S_n \rightarrow T$ is assigned to a morphism $\mathcal{M}f : \mathcal{M}S_1 \times \dots \times \mathcal{M}S_n \rightarrow \mathcal{M}T$.

The \mathcal{M} -*interpretation* $\llbracket \vec{x} \mid t \rrbracket$ of a term t of sort T in context \vec{x} of sort $S_1 \times \dots \times S_n$ is a morphism $S_1 \times \dots \times S_n \rightarrow \mathcal{M}T$ defined recursively:

$$\begin{aligned} \llbracket \vec{x} \mid x_k \rrbracket &=_{\text{df}} \pi_k : \mathcal{M}S_1 \times \dots \times \mathcal{M}S_n \rightarrow \mathcal{M}S_k \\ \llbracket \vec{x} \mid f(t_1, \dots, t_m) \rrbracket &=_{\text{df}} \mathcal{M}f \circ \langle \llbracket \vec{x} \mid t_1 \rrbracket, \dots, \llbracket \vec{x} \mid t_m \rrbracket \rangle : \mathcal{M}S_1 \times \dots \times \mathcal{M}S_n \rightarrow \mathcal{M}W, \end{aligned}$$

where t_1, \dots, t_m are terms of sorts T_1, \dots, T_m , respectively, and f is a function symbol of sort $T_1 \times \dots \times T_m \rightarrow W$. The \mathcal{M} -*interpretation* $\llbracket \vec{x} \mid \phi \rrbracket$ of a formula ϕ in context \vec{x} of sort $S_1 \times \dots \times S_n$ is defined recursively:

$$\begin{aligned} \llbracket \vec{x} \mid \perp \rrbracket &=_{\text{df}} [\perp \rightarrow \mathcal{M}S_1 \times \dots \times \mathcal{M}S_n] \\ \llbracket \vec{x} \mid R(\vec{t}) \rrbracket &=_{\text{df}} \llbracket \vec{x} \mid \vec{t} \rrbracket^*(\mathcal{M}R), \text{ where } R \text{ is a relation symbol and } \vec{t} \text{ are terms.} \\ \llbracket \vec{x} \mid \chi \odot \psi \rrbracket &=_{\text{df}} \llbracket \vec{x} \mid \chi \rrbracket \odot \llbracket \vec{x} \mid \psi \rrbracket, \text{ where } \odot \in \{\wedge, \vee, \rightarrow\}. \\ \llbracket \vec{x} \setminus \{x_k\} \mid \nabla x_k. \psi \rrbracket &=_{\text{df}} \langle \pi_1, \dots, \pi_{k-1}, \pi_{k+1}, \dots, \pi_n \rangle_{\nabla} (\llbracket \vec{x} \mid \psi \rrbracket), \text{ where } \nabla \in \{\forall, \exists\}. \end{aligned}$$

We say that $\phi(\vec{x})$ is *valid* in \mathcal{M} , and write $\mathcal{M} \models \phi$, whenever $\llbracket \vec{x} \mid \phi(\vec{x}) \rrbracket$ equals the maximal element in $\text{Sub}(\mathcal{M}S_1 \times \dots \times \mathcal{M}S_n)$, where $S_1 \times \dots \times S_n$ are the sorts of x_1, \dots, x_n , respectively.

When an interpretation \mathcal{M} of \mathcal{S} in a Heyting category \mathbf{C} is given, we will often simply write “ $\mathbf{C} \models \phi$ ”. Sometimes it is convenient to extend \mathcal{S} with some objects, morphisms and subobjects of \mathbf{C} as new sorts, function symbols and relation symbols, respectively.

Definition 7. Let \mathbf{C} be a Heyting category and let \mathbf{D} be a subcategory of \mathbf{C} with finite products. We define the *\mathbf{D} -signature with respect to \mathbf{C}* , denoted $\mathcal{S}_{\mathbf{D}}^{\mathbf{C}}$, as the following signature.

- **Sorts:** For each object A of \mathbf{D} , A is a sort in $\mathcal{S}_{\mathbf{D}}^{\mathbf{C}}$.
- **Function symbols:** For each morphism $f : A \rightarrow B$ of \mathbf{D} , $f : A \rightarrow B$ is a function symbol in $\mathcal{S}_{\mathbf{D}}^{\mathbf{C}}$ from the sort A to the sort B .
- **Relation symbols:** For each $n \in \mathbb{N}$, and for each morphism $m : A \rightarrow B_1 \times \dots \times B_n$ of \mathbf{D} , such that m is monic in \mathbf{C} , m is an n -ary relation symbol in $\mathcal{S}_{\mathbf{D}}^{\mathbf{C}}$ on the sort $B_1 \times \dots \times B_n$.

Given $\mathcal{S}_{\mathbf{D}}^{\mathbf{C}}$, the *natural $\mathcal{S}_{\mathbf{D}}^{\mathbf{C}}$ -structure* is defined by assigning each sort A to the object A , assigning each function symbol f to the morphism f , and assigning each relation symbol m on the sort $B_1 \times \dots \times B_n$ to the subobject of $B_1 \times \dots \times B_n$ in \mathbf{C} determined by m . Let ϕ be a $\mathcal{S}_{\mathbf{D}}^{\mathbf{C}}$ -formula. We write $\mathbf{C} \models \phi$ for the statement that ϕ is satisfied in the natural $\mathcal{S}_{\mathbf{D}}^{\mathbf{C}}$ -structure.

The importance of Heyting categories lies in this result:

Theorem 8 (Completeness for categorical semantics). *Intuitionistic and classical first order logic are sound and complete for the categorical semantics of Heyting and Boolean categories, respectively.*

We now proceed to introduce a new categorical theory, intended to characterize the categorical content of the predicative class theory ML. For comparison, let us first recall the definition of topos.

Definition 9. A *topos* is a category with finite limits and power objects. A *power object* of an object A , is an object $\mathbf{P}A$ along with a mono $m : \in \rightarrow A \times \mathbf{P}A$ such that for any mono $r : R \rightarrow A \times \mathbf{P}A$, there is a unique morphism $\chi : B \rightarrow \mathbf{P}A$ making this a pullback square:

$$\begin{array}{ccc} R & \longrightarrow & \in \\ r \downarrow & & \downarrow m \\ A \times B & \xrightarrow{\text{id} \times \chi} & A \times \mathbf{P}A \end{array}$$

It is a well-known fact that every topos is a Heyting category, see e.g. [Johnstone, 2002].

We need a relativized notion of power object, for the axiomatization to be presented below:

Definition 10. Let \mathbf{C} be a category, and let \mathbf{D} be a subcategory of \mathbf{C} . A *power object in \mathbf{C} with respect to \mathbf{D}* , of an object A in \mathbf{D} , is defined as in Definition 9, except that r is assumed to be in \mathbf{D} and m, χ are required to be in \mathbf{D} . More precisely, it is an object $\mathbf{P}A$ along with a morphism $m : \in \rightarrow A \times \mathbf{P}A$ in \mathbf{D} which is monic in \mathbf{C} , such that for any $r : R \rightarrow A \times \mathbf{P}A$ in \mathbf{D} which is monic in \mathbf{C} , there is a morphism $\chi : B \rightarrow \mathbf{P}A$ in \mathbf{D} , which is the unique morphism in \mathbf{C} making this a pullback square in \mathbf{C} :

$$\begin{array}{ccc} R & \longrightarrow & \in \\ r \downarrow & & \downarrow m \\ A \times B & \xrightarrow{\text{id} \times \chi} & A \times \mathbf{P}A \end{array}$$

We need a couple of more definitions: A functor $\mathbf{F} : \mathbf{C} \rightarrow \mathbf{D}$ is *conservative* iff for any morphism f in \mathbf{C} , if $\mathbf{F}(f)$ is an isomorphism then f is an isomorphism. A subcategory is *conservative* iff its inclusion functor is conservative. A *universal object* in a category \mathbf{C} is an object X , such that for every object Y there is a mono $f : Y \rightarrow X$. The theory IMLU_{CAT} is axiomatized as follows.

Definition 11 (IMLU_{CAT}). An IMLU-category is a pair of Heyting categories $\langle \mathbf{M}, \mathbf{N} \rangle$, such that

- \mathbf{N} is a conservative Heyting subcategory of \mathbf{M} ,
- there is an object U in \mathbf{N} which is universal in \mathbf{N} ,
- there is an endofunctor \mathbf{T} on \mathbf{M} , restricting to an endofunctor of \mathbf{N} (also denoted \mathbf{T}), along with a natural isomorphism $\iota : \text{id}_{\mathbf{M}} \xrightarrow{\sim} \mathbf{T}$ on \mathbf{M} ,
- there is an endofunctor \mathbf{P} on \mathbf{N} , such that for each object A in \mathbf{N} , $\mathbf{T}A$ has a power object $\mathbf{P}A$, $m_{\subseteq_A^1} : \subseteq_A^1 \rightarrow \mathbf{T}A \times \mathbf{P}A$ in \mathbf{M} with respect to \mathbf{N} ; spelling this out:
 - $m_{\subseteq_A^1}$ is a morphism in \mathbf{N} which is monic in \mathbf{M} , such that
 - for any $r : R \rightarrow \mathbf{T}A \times B$ in \mathbf{N} which is monic in \mathbf{M} , there is $\chi : B \rightarrow \mathbf{P}A$ in \mathbf{N} , which is the unique morphism in \mathbf{M} making this a pullback square in \mathbf{M} :

$$\begin{array}{ccc}
R & \longrightarrow & \subseteq_A^1 \\
r \downarrow & & \downarrow m_{\subseteq_A^1} \\
\mathbf{T}A \times B & \xrightarrow{\text{id} \times \chi} & \mathbf{T}A \times \mathbf{P}A
\end{array} \tag{PT}$$

- there is a natural isomorphism $\mu : \mathbf{P} \circ \mathbf{T} \xrightarrow{\sim} \mathbf{T} \circ \mathbf{P}$ on \mathbf{N} .

In order to carry over some intuitions from set theory, $\mathbf{T}A$ may be thought of as $\{\{x\} \mid x \in A\}$ and $\mathbf{P}A$ may be thought of as $\{X \mid X \subseteq A\}$. Now \subseteq_A^1 corresponds to the subset relation on $\mathbf{T}A \times \mathbf{P}A$; its superscript 1 is intended to convey that its left relata are “like singletons”. Note that on this picture, \subseteq^1 is very similar to the \in -relation. Thus (PT) is intended to be the appropriate variant for NF of the power object axiom of topos theory. These intuitions are made precise in the next section, where we interpret $(\text{I})\text{ML}(\text{U})_{\text{CAT}}$ in $(\text{I})\text{ML}(\text{U})_{\text{CLASS}}$.

It is easily seen that this axiomatization is elementary, i.e. it corresponds to a theory in a first order language \mathcal{L}_{CAT} . Its precise specification involves quite some detail. Suffice to say that the language of category theory is augmented with relation symbols \mathbf{M}_{Ob} , \mathbf{M}_{Mor} , \mathbf{N}_{Ob} , and \mathbf{N}_{Mor} ; a constant symbol U ; and function symbols \mathbf{T}_{Ob} , \mathbf{T}_{Mor} , ι , μ , \mathbf{P}_{Ob} and \mathbf{P}_{Mor} (using the same names for the symbols and their interpretations).

If “Heyting” is replaced with “Boolean” we obtain the theory MLU_{CAT} . If the axiom $U \cong \mathbf{P}U$ is added to $(\text{I})\text{MLU}_{\text{CAT}}$, we obtain the theories $(\text{I})\text{ML}_{\text{CAT}}$, respectively.

Note that the definition can easily be generalized, so that we merely require that \mathbf{N} is a Heyting category that is mapped into \mathbf{M} by a faithful conservative Heyting functor $\mathbf{F} : \mathbf{N} \rightarrow \mathbf{M}$. This would not hinder any of the results below. We choose the more specific definition in terms of a subcategory because it simplifies the statements of the results.

We shall now collect a few useful properties of $(\text{I})\text{ML}(\text{U})$ -categories. First a definition: A functor $\mathbf{F} : \mathbf{B} \rightarrow \mathbf{C}$ *reflects finite limits* iff for any finite diagram $\mathbf{D} : \mathbf{I} \rightarrow \mathbf{B}$ and for any cone Λ of \mathbf{D} in \mathbf{B} , if $\mathbf{F}\Lambda$ is a limit in \mathbf{C} of $\mathbf{F} \circ \mathbf{D} : \mathbf{I} \rightarrow \mathbf{C}$, then Λ is a limit of $\mathbf{D} : \mathbf{I} \rightarrow \mathbf{B}$ in \mathbf{B} .

Proposition 12. *Let $\langle \mathbf{M}, \mathbf{N} \rangle$ along with U , \mathbf{T} , ι and \mathbf{P} be an IMLU-category.*

- (a) For any morphism $f : A \rightarrow B$ in \mathbf{N} , f is monic in \mathbf{N} iff f is monic in \mathbf{M} .
- (b) The inclusion functor of \mathbf{N} (as a subcategory) into \mathbf{M} reflects finite limits.
- (c) $\mathbf{P}A$ along with $m_{\subseteq_A^1}$, as in (PT) above, is a power object of $\mathbf{T}A$ in \mathbf{N} , for any A in \mathbf{N} .
- (d) $\mathbf{T} : \mathbf{M} \rightarrow \mathbf{M}$ is a Heyting endofunctor. If $\langle \mathbf{M}, \mathbf{N} \rangle$ is an MLU-category, then $\mathbf{T} : \mathbf{M} \rightarrow \mathbf{M}$ is a Boolean endofunctor.
- (e) $\mathbf{T} : \mathbf{N} \rightarrow \mathbf{N}$ preserves finite limits.

Proof. (a) (\Leftarrow) follows immediately from that \mathbf{N} is a subcategory of \mathbf{M} . (\Rightarrow) follows from that \mathbf{N} is a Heyting subcategory of \mathbf{M} , and that Heyting functors preserve images.

(b) Let L , along with some morphisms in \mathbf{N} , be a cone in \mathbf{N} of a finite diagram $\mathbf{D} : \mathbf{I} \rightarrow \mathbf{N}$, such that this cone is a limit of $\mathbf{D} : \mathbf{I} \rightarrow \mathbf{N}$ in \mathbf{M} . Let K be a limit in \mathbf{N} of $\mathbf{D} : \mathbf{I} \rightarrow \mathbf{N}$. Since \mathbf{N} is a Heyting subcategory of \mathbf{M} , K is also such a limit in \mathbf{M} . Let $f : L \rightarrow K$ be the universal morphism in \mathbf{N} obtained from the limit property of K in \mathbf{N} . By the limit properties of K and L in \mathbf{M} , f is an isomorphism in \mathbf{M} . Since \mathbf{N} is a conservative subcategory of \mathbf{M} , f is also an isomorphism in \mathbf{N} , whence L is a limit in \mathbf{N} of $\mathbf{D} : \mathbf{I} \rightarrow \mathbf{N}$, as desired.

(c) By (a), we do not need to worry about whether the morphisms involved are monic. Let A be an object of \mathbf{N} . By (b), (PT) is a pullback square in \mathbf{N} . Suppose that χ' in \mathbf{N} makes (PT) a pullback in \mathbf{N} (in place of χ). Since \mathbf{N} is a Heyting subcategory of \mathbf{M} , χ' also makes (PT) a pullback square in \mathbf{M} . So by the uniqueness property in \mathbf{M} , $\chi' = \chi$.

(d) Since $\mathbf{T} : \mathbf{M} \rightarrow \mathbf{M}$ is naturally isomorphic to the identity functor, it is a Heyting (Boolean) endofunctor of \mathbf{M} .

(e) Let L be a limit in \mathbf{N} of a finite diagram $\mathbf{D} : \mathbf{I} \rightarrow \mathbf{N}$. By (d), $\mathbf{T} : \mathbf{M} \rightarrow \mathbf{M}$ preserves limits, so $\mathbf{T}L$ is a limit in \mathbf{M} of $\mathbf{T} \circ \mathbf{D} : \mathbf{I} \rightarrow \mathbf{N}$. By (b), $\mathbf{T}L$ is also a limit in \mathbf{N} of $\mathbf{T} \circ \mathbf{D} : \mathbf{I} \rightarrow \mathbf{N}$. \square

6 Interpretation of the CAT-theories in the CLASS-theories

This section is devoted to establishing $\text{Con}((\mathbf{I})\text{NF}(\mathbf{U})_{\text{CLASS}}) \Rightarrow \text{Con}((\mathbf{I})\text{ML}(\mathbf{U})_{\text{CAT}})$. This is the easy and less interesting part of the equiconsistency proof, but it has the beneficial spin-off of showing how the axioms of $(\mathbf{I})\text{ML}(\mathbf{U})_{\text{CAT}}$ correspond to set theoretic intuitions. Given Theorem 5, it suffices to find an interpretation of $(\mathbf{I})\text{ML}(\mathbf{U})_{\text{CAT}}$ in $(\mathbf{I})\text{ML}(\mathbf{U})_{\text{CLASS}}$, as is done in the proof below. It is easy to see how this proof can be modified to show that $(\mathbf{I})\text{ML}(\mathbf{U})_{\text{CAT}}$ can be interpreted in the predicative version of ML, but since the impredicative theory is of the same consistency strength, that is not needed for the equiconsistency result.

Theorem 13. $(\mathbf{I})\text{ML}(\mathbf{U})_{\text{CAT}}$ is interpretable in $(\mathbf{I})\text{ML}(\mathbf{U})_{\text{CLASS}}$.

Proof. We go through the case of IMLU in detail, and then explain the modifications required for the other cases. The interpretation proceeds as follows.

1. Interpret $\mathbf{M}_{\text{Ob}}(x)$ as “ x is a class”.
2. Interpret $\mathbf{M}_{\text{Mor}}(m)$ as “ m is a disjoint union of three classes A , B and f , such that f codes a function with domain A and codomain B ”.¹

¹A disjoint union of three classes may be implemented as a class using the formula $\langle i, x \rangle \in m \leftrightarrow ((i = 1 \wedge x \in A) \vee (i = 2 \wedge x \in B) \vee (i = 3 \wedge x \in f))$. In order to be able to interpret the domain and codomain function symbols, we need to include information about the domain class and codomain class in the interpretation of the morphisms. Otherwise, the same functional class will often interpret many morphisms with different co-domains.

3. Interpret the remaining symbols of the language of category theory in the obvious way. The resulting interpretations of the axioms of basic category theory are now easily verified for \mathbf{M} .
4. Interpret $\mathbf{N}_{\text{Ob}}(x)$ as “ x is a set”; and interpret $\mathbf{N}_{\text{Mor}}(m)$ as “[insert the interpretation of $\mathbf{M}_{\text{Mor}}(m)$] and m is a set”. The axioms of basic category theory are now easily verified for \mathbf{N} , by restricting the quantifiers in their interpretation to sets.
5. We need to show that the interpretation of the axioms of Heyting categories hold for \mathbf{M} and \mathbf{N} . Ext_C ensures the uniqueness conditions in the axioms. Existence conditions will be supported by class/set abstracts $\{x \mid \phi(x)\}$. In the case of \mathbf{M} we rely on CC_C , and we need to check that the free variable is small. In the case of \mathbf{N} we rely on SC_S , and we need to check that the formula is stratified. These verifications are simple and left to the reader. It is also easy for the reader to verify, for both \mathbf{M} and \mathbf{N} , that subobjects are represented by subclasses/subsets, and that a morphism m is monic iff injective, and is a cover iff surjective. We now proceed to verify the HC axioms.
 - (F1) Finite limits: $\{\emptyset\}$ is a terminal object. $D =_{\text{df}} \{\langle x, y \rangle \in A \times B \mid m(x) = n(y)\}$, along with projection morphisms $\pi_1 : D \rightarrow A$ and $\pi_2 : D \rightarrow C$, is a pullback of morphisms $m : A \rightarrow B$ and $n : C \rightarrow B$.
 - (F2) Images: $\{m(x) \mid x \in A\}$, along with its inclusion function into B is the image of $m : A \rightarrow B$.
 - (F3) Covers stable under pullback: Suppose m is surjective in (5). Then, for any $c \in C$, there is $a \in A$ such that $m(a) = n(c)$. Hence, the projection $D \rightarrow C$ is surjective.
 - (F4) Each Sub_X is a sup-semilattice: Binary union is given by $\{z \mid z \in A \vee z \in B\}$.
 - (F5) For each $f : X \rightarrow Y$, the functor $f^* : \text{Sub}_Y \rightarrow \text{Sub}_X$ preserves finite suprema and has left and right adjoints, $\exists_f \dashv f^* \dashv \forall_f$: Here f^* is the inverse image functor, which clearly preserves finite unions. \exists_f is simply the image functor and \forall_f is the functor mapping $X' \subseteq X$ to $\{y \in Y \mid \forall x \in X. (f(x) = y \rightarrow x \in X')\} \subseteq Y$. Let $X' \subseteq X$ and $Y' \subseteq Y$. It is easily seen that $\exists_f(X') \subseteq Y' \iff X' \subseteq f^*(Y')$, i.e. $\exists_f \dashv f^*$. It is also easily seen that $f^*(Y') \subseteq X' \iff Y' \subseteq \forall_f(X')$, i.e. $f^* \dashv \forall_f$.
6. In the verification of the HC axioms above, when the objects and morphisms are in \mathbf{N} , the same witnesses are used regardless of whether the HC axioms are verified for \mathbf{M} or \mathbf{N} . It follows that \mathbf{N} is a Heyting subcategory of \mathbf{M} .
7. In both \mathbf{M} and \mathbf{N} , a morphism is an isomorphism iff it is bijective. Hence, \mathbf{N} is a conservative subcategory of \mathbf{M} .
8. Interpret U as V , which is a superset of every set, and hence universal in \mathbf{N} .
9. Interpret $\mathbf{T}_{\text{Ob}}(x)$ as “the class of singletons of members of x ”; and interpret $\mathbf{T}_{\text{Mor}}(m)$ as “the class coding the morphism $(\{x\} \mapsto \{f(x)\}) : \mathbf{T}A \rightarrow \mathbf{T}B$, where m codes $f : A \rightarrow B$ ”. It is easily verified that the interpreted axioms of a functor hold. By the singleton image axiom, \mathbf{T} restricts appropriately to \mathbf{N} .
10. Interpret ι_x as the code of the morphism $(z \mapsto \{z\}) : x \rightarrow \mathbf{T}(x)$, which is a class. Since the inverse of ι_x is similarly interpretable, we obtain that the interpretation of ι_x is an isomorphism in the category theoretic sense. That ι is a natural isomorphism is clear from its definition and the definition of \mathbf{T} .

11. Interpret $\mathbf{P}_{\text{Ob}}(x)$ as $\mathcal{P}x$. Interpret $\mathbf{P}_{\text{Mor}}(m)$ as “the set coding the morphism $(x \mapsto \{f(z) \mid z \in x\}) : \mathcal{P}A \rightarrow \mathcal{P}B$, where m codes $f : A \rightarrow B$ ”. It is easily seen that this makes \mathbf{P} an endofunctor on \mathbf{N} .
12. When interpreting $\mu_x : \mathbf{P}\mathbf{T}x \rightarrow \mathbf{T}\mathbf{P}x$, note that $\mathbf{P}\mathbf{T}x = \mathcal{P}\{\{z\} \mid z \in x\}$. Let μ_x be the set coding the morphism $(u \mapsto \{\cup u\}) : \mathcal{P}\{\{z\} \mid z \in y\} \rightarrow \{\{v\} \mid v \in \mathcal{P}y\}$. Union and singleton are defined by stratified formulas. Because the union operation lowers type by one and the singleton operation raises type by one, argument and value are type-level, so μ_x exists as a set. It is easily seen from the definitions of \mathbf{T} , \mathbf{P} and μ , that μ is a natural isomorphism.
13. Define $x \subseteq^1 y$ set theoretically by $\exists u.(x = \{u\} \wedge u \in y)$. We interpret $\subseteq_A^1 \hookrightarrow \mathbf{T}A \times \mathbf{P}A$ as $\{\langle x, y \rangle \in \mathbf{T}A \times \mathbf{P}A \mid x \subseteq^1 y\} \subseteq \mathbf{T}A \times \mathbf{P}A$ for the diagram (PT).
14. We proceed to verify the property (PT). Suppose that $r : R \rightarrow \mathbf{T}A \times B$ in \mathbf{N} is monic in \mathbf{M} . In both \mathbf{N} and \mathbf{M} , a morphism is monic iff it is injective, so r is monic in \mathbf{N} . Define $\chi : B \rightarrow \mathcal{P}A$ by $y \mapsto \{u \mid \exists c \in R.r(c) = \langle \{u\}, y \rangle\}$. Since this is a stratified definition, where argument and value have equal type, χ is a morphism in \mathbf{N} . The proof that χ is the unique morphism making (PT) a pullback in \mathbf{M} is just like the standard proof in conventional set theory; it proceeds as follows. We may assume that $R \subseteq A \times B$ and r is the inclusion function. Then χ is $(y \mapsto \{u \mid \{u\}Ry\})$. For the top arrow in (PT) we choose $(\text{id} \times \chi) \upharpoonright_R$. Since $\forall \langle \{u\}, y \rangle \in R.\{u\} \subseteq^1 \chi(y)$, (PT) commutes.

For the universal pullback property: Suppose that $\langle f, g \rangle : Q \rightarrow \mathbf{T}A \times B$ and $\langle d, e \rangle : Q \rightarrow \subseteq_A^1$ are morphisms in \mathbf{M} making the diagram commute. Let $q \in Q$ be arbitrary. Then $f(q) = d(q)$ and $\chi(g(q)) = e(q)$, so $f(q) \subseteq^1 \chi(g(q))$ and $f(q)Rg(q)$. Thus, $(q \mapsto \langle f(q), g(q) \rangle)$ defines the unique morphism in \mathbf{M} , witnessing the universal pullback property.

It remains to show that if χ' is a morphism in \mathbf{M} that (in place of χ) makes (PT) a pullback in \mathbf{M} , then $\chi' = \chi$. Let χ' be such that it makes (PT) a pullback, and let $u \in A$ and $y \in B$; we need to show that $\{u\}Ry \iff \{u\} \subseteq^1 \chi'(y)$. By commutativity $\{u\}Ry \Rightarrow \{u\} \subseteq^1 \chi'(y)$. Conversely, applying the universal pullback property to $\{\langle \{u\}, y \rangle \hookrightarrow \mathbf{T}A \times B\}$, we find that $\{u\} \subseteq^1 \chi'(y) \Rightarrow \{u\}Ry$.

This completes the interpretation of IMLU_{CAT} in $\text{IMLU}_{\text{CLASS}}$. For MLU , simply observe that $\text{MLU}_{\text{CLASS}} \vdash \forall X.\forall X' \subseteq X.X' \cup (X - X') = X$, so each Sub_X is Boolean. For (I)ML, the fact that $V = \mathcal{P}V$ ensures that the interpretation of $U \cong \mathbf{P}U$ holds. \square

7 Interpretation of the SET-theories in the CAT-theories

For the rest of the paper, fix an IMLU-category $\langle \mathbf{M}, \mathbf{N} \rangle$ – along with U in \mathbf{N} ; $\mathbf{T} : \mathbf{M} \rightarrow \mathbf{M}$ (restricting to an endofunctor of \mathbf{N}); $\iota : \mathbf{id} \xrightarrow{\sim} \mathbf{T}$ in \mathbf{M} ; $\mathbf{P} : \mathbf{N} \rightarrow \mathbf{N}$; $\mu : \mathbf{P} \circ \mathbf{T} \xrightarrow{\sim} \mathbf{T} \circ \mathbf{P}$; and $\subseteq_X^1 \hookrightarrow \mathbf{T}X \times \mathbf{P}X$ (for each object X) in \mathbf{N} – all satisfying the conditions in Definition 11. Moreover, fix an object 1 which is terminal in both \mathbf{M} and \mathbf{N} and fix a product functor \times on \mathbf{M} which restricts to a product functor on \mathbf{N} . This can be done since \mathbf{N} is a Heyting subcategory of \mathbf{M} . Given an $n \in \mathbb{N}$ and a product P of n objects, the i -th projection morphism, for $i = 1, \dots, n$, is denoted π_P^i .

In this section, we shall establish that $\text{Con}((\text{I})\text{ML}(U)_{\text{CAT}}) \Rightarrow \text{Con}((\text{I})\text{NF}(U)_{\text{SET}})$. We do so by proving that the axioms of $(\text{I})\text{NF}(U)_{\text{SET}}$ can be interpreted in the internal language of $(\text{I})\text{ML}(U)_{\text{CAT}}$. In particular, we construct a structure in the categorical semantics of \mathbf{M} which satisfies the axioms of $(\text{I})\text{NF}(U)_{\text{SET}}$. The variation between the intuitionistic and the classical case is handled by Theorem 8, so we will concentrate on proving $\text{Con}(\text{IMLU}_{\text{CAT}}) \Rightarrow \text{Con}(\text{INFU}_{\text{SET}})$,

and $\text{Con}(\text{MLU}_{\text{CAT}}) \Rightarrow \text{Con}(\text{NFU}_{\text{SET}})$ is thereby obtained as well, simply by assuming that $\langle \mathbf{M}, \mathbf{N} \rangle$ is an MLU-category. Lemma 19 below shows that this suffices for also establishing $\text{Con}((\text{I})\text{ML}_{\text{CAT}}) \Rightarrow \text{Con}((\text{I})\text{NF}_{\text{SET}})$.

Construction 14. For each object A of \mathbf{N} let \in_A , along with m_{\in_A} , be this pullback in \mathbf{M} :

$$\begin{array}{ccc} \in_A & \xrightarrow{\sim} & \subseteq_A^1 \\ \downarrow m_{\in_A} & & \downarrow \\ A \times \mathbf{P}A & \xrightarrow[\iota \times \text{id}]{\sim} & \mathbf{T}A \times \mathbf{P}A \end{array}$$

In order to avoid confusing the \in_A defined above with the membership symbol of \mathcal{L}_{SET} , the latter is replaced by the symbol ε .

Construction 15. This \mathcal{L}_{SET} -structure, in the categorical semantics of \mathbf{M} , is denoted \mathcal{U} :

1. The single sort of \mathcal{L}_{SET} is assigned to the universal object U of \mathbf{N} .
2. Fix a mono $m_S : \mathbf{P}U \rightarrow U$ in \mathbf{N} . The sethood predicate symbol S is identified with the predicate symbol m_S in $\mathcal{S}_{\mathbf{N}}^{\mathbf{M}}$, and is assigned to the subobject of U determined by m_S .
3. Fix the mono $m_\varepsilon =_{\text{df}} (\text{id}_U \times m_S) \circ m_{\in_U} : \in_U \rightarrow U \times \mathbf{P}U \rightarrow U \times U$. The membership symbol ε is identified with the symbol m_ε in $\mathcal{S}_{\mathbf{M}}^{\mathbf{M}}$, and is assigned to the subobject of $U \times U$ determined by m_ε .
4. Fix a mono $m_P : U \times U \rightarrow U$ in \mathbf{N} . The function symbol $\langle -, - \rangle$ is identified with the symbol m_P in $\mathcal{S}_{\mathbf{N}}^{\mathbf{M}}$ and is assigned to the subobject of U determined by m_P .

By the identifications of symbols, the signature of \mathcal{L}_{SET} is a subsignature of $\mathcal{S}_{\mathbf{M}}^{\mathbf{M}}$.

We will usually omit subscripts such as in \in_A and \subseteq_A^1 , as they tend to be obvious. Similarly, sort declarations are sometimes omitted when considering formulas of the internal language. Note that the symbol ε in \mathcal{L}_{SET} is interpreted by the subobject of $U \times U$ determined by m_ε , not by the subobject of $U \times \mathbf{P}U$ determined by m_{\in_U} . In the categorical setting it tends to be more natural to have a membership relation of sort $A \times \mathbf{P}A$ for each object A , while in the set-theoretical setting it tends to be more natural to have just one sort, say Universe, and just one membership relation of sort Universe \times Universe.

To prove $\text{Con}((\text{I})\text{ML}(\text{U})_{\text{CAT}}) \Rightarrow \text{Con}((\text{I})\text{NF}(\text{U})_{\text{SET}})$, we need to establish that \mathcal{U} satisfies Axioms 3. $\mathcal{U} \models \phi$ is the statement that the \mathcal{L}_{SET} -structure \mathcal{U} satisfies $\phi \in \mathcal{L}_{\text{SET}}$, in the categorical semantics of \mathbf{M} . For the major axioms, Extensionality and Stratified Comprehension, we will first prove the more general (and more naturally categorical) statements in terms of the \in_A , and second obtain the required statements about ε as corollaries. The general results will be stated in the form $\mathbf{M} \models \phi$, where ϕ is a formula in the language of $\mathcal{S}_{\mathbf{M}}^{\mathbf{M}}$ or some subsignature of it.

The following proposition is the expression of Construction 14 in the categorical semantics.

Proposition 16. *Let X be an object of \mathbf{N} .*

$$\mathbf{M} \models \forall x : X. \forall y : \mathbf{P}X. (x \in y \leftrightarrow \iota x \subseteq^1 y).$$

Proof. $\llbracket x, y \mid \iota x \subseteq^1 y \rrbracket = (\iota \times \text{id})^* \llbracket u, y \mid u \subseteq^1 y \rrbracket = \llbracket x, y \mid x \in y \rrbracket$. □

Let us start the proof of $\mathcal{U} \models \text{INFU}_{\text{SET}}$ with the easy axioms of Sethood and Ordered Pair.

Proposition 17 (Sethood). $\mathcal{U} \models \forall z. \forall x. (z \varepsilon x \rightarrow S(x))$

Proof. $\llbracket z, x : U \mid z \varepsilon x \rrbracket \leq U \times \mathbf{P}U \cong \llbracket z, x : U \mid S(x) \rrbracket$. \square

Proposition 18 (Ordered Pair). $\mathcal{U} \models \forall x, x', y, y'. (\langle x, y \rangle = \langle x', y' \rangle \leftrightarrow (x = x' \wedge y = y'))$

Proof. The result follows immediately from that $\langle -, - \rangle$ is interpreted by a mono $U \times U \rightarrow U$. \square

The following Lemma shows that we get $\text{Con}((\mathbf{I})\text{ML}_{\text{CAT}}) \Rightarrow \text{Con}((\mathbf{I})\text{NF}_{\text{SET}})$ for free, if we successfully prove that $\mathcal{U} \models (\mathbf{I})\text{NF}_{\text{SET}}$.

Lemma 19 ((I)NF for free). *If $U \cong \mathbf{P}U$, then we can choose $m_S : \mathbf{P}U \rightarrow U$ (i.e. the interpretation of the predicate symbol S) to be an isomorphism. If so, then $\mathcal{U} \models \forall x.S(x)$.*

Proof. Since m_S is an isomorphism, $m_s : \mathbf{P}U \rightarrow U$ and $\text{id} : U \rightarrow U$ represent the same subobject of U . \square

Note that we do not need $U = \mathbf{P}U$ for this result; $U \cong \mathbf{P}U$ suffices. This means that our results will actually give us that $(\mathbf{I})\text{NF}_{\text{SET}}$ is equiconsistent with $(\mathbf{I})\text{NF}_{\text{SET}} + (|V| = |\mathcal{P}(V)|)$, with essentially no extra work. See Corollary 35 below. This result has been proved previously in [Crabbé, 2000] using the conventional set-theoretical semantics, but in the present categorical setting the result is intuitively obvious.

Proposition 20 (Extensionality). *Let Z be an object of \mathbf{N} .*

$$\mathbf{M} \models \forall x : \mathbf{P}Z. \forall y : \mathbf{P}Z. [(\forall z : Z. (z \in x \leftrightarrow z \in y)) \rightarrow x = y].$$

Proof. We use the fact that \mathbf{N} is a Heyting subcategory of \mathbf{M} . By Proposition 16, it suffices to establish that in \mathbf{N} :

$$\llbracket x : \mathbf{P}Z, y : \mathbf{P}Z \mid \forall z : \mathbf{T}Z. (z \subseteq^1 x \leftrightarrow z \subseteq^1 y) \rrbracket \leq \llbracket x, y \mid x = y \rrbracket.$$

Let $\langle a, b \rangle : E \rightarrow \mathbf{P}Z \times \mathbf{P}Z$ represent $\llbracket x, y \mid \forall z. (z \subseteq^1 x \leftrightarrow z \subseteq^1 y) \rrbracket$. We need to show that $a = b$.

Consider $\llbracket w, u, v \mid w \subseteq^1 u \rrbracket$ and $\llbracket w, u, v \mid w \subseteq^1 v \rrbracket$ as subobjects of $\mathbf{T}Z \times \mathbf{P}Z \times \mathbf{P}Z$. We calculate their pullbacks along $\text{id} \times \langle a, b \rangle$ to be equal subobjects of $\llbracket x, y \mid \forall z. (z \subseteq^1 x \leftrightarrow z \subseteq^1 y) \rrbracket$:

$$\begin{aligned} & (\text{id} \times \langle a, b \rangle)^* \llbracket w, u, v \mid w \subseteq^1 u \rrbracket \\ &= \llbracket w, u, v \mid w \subseteq^1 u \wedge \forall z. (z \subseteq^1 u \leftrightarrow z \subseteq^1 v) \rrbracket \\ &= \llbracket w, u, v \mid w \subseteq^1 u \wedge w \subseteq^1 v \wedge \forall z. (z \subseteq^1 u \leftrightarrow z \subseteq^1 v) \rrbracket \\ &= \llbracket w, u, v \mid w \subseteq^1 v \wedge \forall z. (z \subseteq^1 u \leftrightarrow z \subseteq^1 v) \rrbracket \\ &= (\text{id} \times \langle a, b \rangle)^* \llbracket w, u, v \mid w \subseteq^1 v \rrbracket \end{aligned}$$

From inspection of the chain of pullbacks

$$\begin{array}{ccccc} (\text{id} \times \langle a, b \rangle)^* \llbracket w, u, v \mid w \subseteq^1 u \rrbracket & \xrightarrow{\quad} & \llbracket w, u, v \mid w \subseteq^1 u \rrbracket & \longrightarrow & \llbracket w, t \mid w \subseteq^1 t \rrbracket \\ \downarrow f & & \downarrow & & \downarrow \\ \llbracket z, x, y \mid \forall z. (z \subseteq^1 x \leftrightarrow z \subseteq^1 y) \rrbracket & \xrightarrow{\text{id} \times \langle a, b \rangle} & \mathbf{T}Z \times \mathbf{P}Z \times \mathbf{P}Z & \xrightarrow{\langle \pi^1, \pi^2 \rangle} & \mathbf{T}Z \times \mathbf{P}Z, \\ & \searrow & & & \nearrow \\ & & \text{id} \times a & & \end{array}$$

it is evident that

$$(\text{id} \times a)^* (\llbracket w, t \mid w \subseteq^1 t \rrbracket) = (\text{id} \times \langle a, b \rangle)^* \llbracket w, u, v \mid w \subseteq^1 u \rrbracket.$$

Similarly,

$$(\text{id} \times b)^* (\llbracket w, t \mid w \subseteq^1 t \rrbracket) = (\text{id} \times \langle a, b \rangle)^* \llbracket w, u, v \mid w \subseteq^1 v \rrbracket.$$

So

$$(\text{id} \times a)^* (\llbracket w, t \mid w \subseteq^1 t \rrbracket) = (\text{id} \times b)^* (\llbracket w, t \mid w \subseteq^1 t \rrbracket),$$

and f represents them as a subobject of $\llbracket z, x, y \mid \forall z. (z \subseteq^1 x \leftrightarrow z \subseteq^1 y) \rrbracket$.

By Proposition 12(c) and uniqueness of χ in (PT), we conclude that $a = b$. \square

Corollary 21. $\mathcal{U} \models \text{Ext}_S$

Proof. By Lemma 20, $\mathbf{M} \models \forall x : \mathbf{P}\mathcal{U}. \forall y : \mathbf{P}\mathcal{U}. [(\forall z : \mathcal{U}. (z \in x \leftrightarrow z \in y)) \rightarrow x = y]$. Now

$$\begin{aligned} & \mathbf{M} \models \forall x, y : \mathbf{P}\mathcal{U}. [(\forall z : \mathcal{U}. (z \in x \leftrightarrow z \in y)) \rightarrow x = y] \\ \iff & \mathbf{M} \models \forall x, y : \mathbf{P}\mathcal{U}. [(\forall z : \mathcal{U}. (z \varepsilon m_S(x) \leftrightarrow z \varepsilon m_S(y))) \rightarrow x = y] \\ \iff & \mathbf{M} \models \forall x', y' : \mathcal{U}. [(S(x') \wedge S(y')) \rightarrow ((\forall z : \mathcal{U}. (z \varepsilon x' \leftrightarrow z \varepsilon y')) \rightarrow x' = y')]. \end{aligned}$$

So $\mathcal{U} \models \text{Ext}_S$. \square

The only axiom of INFU left to validate is SC_S (i.e. Stratified Comprehension). In order to approach this, we first need to construct some signatures and define stratification for an appropriate internal language:

Definition 22. Let $\mathcal{S}_{N, \in}^M$ be the subsignature of \mathcal{S}_M^M containing \mathcal{S}_N^M and the relation symbol \in_A , which is identified with m_{\in_A} , for each object A in \mathbf{N} .

Stratification in the language of $\mathcal{S}_{N, \in}^M$ is defined analogously as in Definition 1. Thus we require that the stratification function s , of an $\mathcal{S}_{N, \in}^M$ -formula ϕ , satisfies these conditions:

- (i) if $u \equiv v(w_1, \dots, w_n)$, then $s(u) = s(w_1) = \dots = s(w_n)$,
- (ii) if $\theta \equiv R(w_1, \dots, w_n)$, then $s(w_1) = \dots = s(w_n)$,
- (iii) if $\theta \equiv (u \in_A w)$, then $s(u) + 1 = s(w)$,

where \equiv is literal syntactic equality; $n \in \mathbb{N}$; u, v, w, w_1, \dots, w_n are $\mathcal{S}_{N, \in}^M$ -terms in ϕ ; θ is an atomic subformula of ϕ ; R is a relation symbol in \mathcal{S}_N^M ; and A is an object in \mathbf{N} .

Let $\mathcal{S}_{N, \iota}^M$ be the subsignature of \mathcal{S}_M^M containing \mathcal{S}_N^M and the function symbol ι_A for each object A in \mathbf{N} .

If ϕ is a formula in either of these languages, then $\mathbf{M} \models \phi$ is to be understood as satisfaction in the natural \mathcal{S}_M^M -structure.

We start by verifying a form of comprehension for \mathcal{S}_N^M -formulas:

Proposition 23. *If $\phi(w, y)$ is an \mathcal{S}_N^M -formula, with context $w : \mathbf{T}Z, y : Y$ for Z, Y in \mathbf{N} , and in which x is not free, then*

$$\mathbf{M} \models \forall y : Y. \exists x : \mathbf{P}Z. \forall w : \mathbf{T}Z. (w \subseteq^1 x \leftrightarrow \phi(w, y)).$$

Proof. This is a familiar property of power objects. Considering this instance of (PT) in \mathbf{N} :

$$\begin{array}{ccc} \llbracket w : \mathbf{T}Z, y : Y \mid \phi(w, y) \rrbracket & \longrightarrow & \subseteq_Z^1 \\ \downarrow & & \downarrow \\ \mathbf{T}Z \times Y & \xrightarrow{\text{id} \times \chi} & \mathbf{T}Z \times \mathbf{P}Z \end{array} \quad (1)$$

This pullback along $\text{id} \times \chi$ can also be expressed as $\llbracket w : \mathbf{T}Z, y : Y \mid w \subseteq_Z^1 \chi(y) \rrbracket$ in \mathbf{N} . So since \mathbf{N} is a Heyting subcategory of \mathbf{M} ,

$$\mathbf{M} \models \forall y : Y. \forall w : \mathbf{T}Z. (w \subseteq^1 \chi(y) \leftrightarrow \phi(w, y)), \text{ and}$$

$$\mathbf{M} \models \forall y : Y. \exists x : \mathbf{P}Z. \forall w : \mathbf{T}Z. (w \subseteq^1 x \leftrightarrow \phi(w, y)),$$

as desired. \square

To obtain stratified comprehension for $\mathcal{S}_{\mathbf{N}, \epsilon}^{\mathbf{M}}$ -formulas, we need to establish certain coherence conditions. The facts that \mathbf{N} is a Heyting subcategory of \mathbf{M} and that \mathbf{T} preserves limits as an endofunctor of \mathbf{N} , enable us to prove that certain morphisms constructed in \mathbf{M} also exist in \mathbf{N} , as in the proposition and lemmata below. This is useful when applying (PT), since the relation R is required to be in \mathbf{N} (see Definition 11).

Proposition 24. *Let $n \in \mathbb{N}$.*

1. $\iota_1 : 1 \rightarrow \mathbf{T}1$ is an isomorphism in \mathbf{N} .
2. For any objects A, B of \mathbf{N} , $(\iota_A \times \iota_B) \circ \iota_{A \times B}^{-1} : \mathbf{T}(A \times B) \xrightarrow{\sim} A \times B \xrightarrow{\sim} \mathbf{T}A \times \mathbf{T}B$ is an isomorphism in \mathbf{N} .
3. For any objects A_1, \dots, A_n in \mathbf{N} ,

$$(\iota_{A_1} \times \dots \times \iota_{A_n}) \circ \iota_{A_1 \times \dots \times A_n}^{-1} : \mathbf{T}(A_1 \times \dots \times A_n) \xrightarrow{\sim} A_1 \times \dots \times A_n \xrightarrow{\sim} \mathbf{T}A_1 \times \dots \times \mathbf{T}A_n$$

is an isomorphism in \mathbf{N} .

Proof. 1. Since \mathbf{T} preserves limits, $\mathbf{T}1$ is terminal in \mathbf{N} , and since \mathbf{N} is a Heyting subcategory of \mathbf{M} , both are terminal in \mathbf{M} as well. So by the universal property of terminal objects, 1 and $\mathbf{T}1$ are isomorphic in \mathbf{N} , and the isomorphisms must be ι_1 and ι_1^{-1} .

2. Since \mathbf{T} preserves limits, $\mathbf{T}(A \times B)$ is a product of $\mathbf{T}A$ and $\mathbf{T}B$ in \mathbf{N} , and since \mathbf{N} is a Heyting subcategory of \mathbf{M} , it is such a product in \mathbf{M} as well. Now note that

$$\pi_{\mathbf{T}A \times \mathbf{T}B}^1 \circ (\iota_A \times \iota_B) \circ \iota_{A \times B}^{-1} = \iota_A \circ \pi_{A \times B}^1 \circ \iota_{A \times B}^{-1} = \mathbf{T}\pi_{A \times B}^1,$$

and similarly for the second projection. The right equality follows from that ι is a natural isomorphism. This means that

$$(\iota_A \times \iota_B) \circ \iota_{A \times B}^{-1} : \mathbf{T}(A \times B) \xrightarrow{\sim} \mathbf{T}A \times \mathbf{T}B$$

is the unique universal morphism provided by the definition of product. Hence, it is an isomorphism in \mathbf{N} .

3. This follows from the two items above by induction on n . \square

Lemma 25. *Let $n \in \mathbb{N}$. If $u : A_1 \times \dots \times A_n \rightarrow B$ is a morphism in \mathbf{N} , then there is a morphism $v : \mathbf{T}A_1 \times \dots \times \mathbf{T}A_n \rightarrow \mathbf{T}B$ in \mathbf{N} , such that*

$$\iota_B \circ u = v \circ (\iota_{A_1} \times \dots \times \iota_{A_n}).$$

Proof. Since ι is a natural transformation,

$$\iota_B \circ u = (\mathbf{T}u) \circ \iota_{A_1 \times \dots \times A_n}.$$

Since ι is a natural isomorphism,

$$\iota_{A_1 \times \dots \times A_n} = \iota_{A_1 \times \dots \times A_n} \circ (\iota_{A_1} \times \dots \times \iota_{A_n})^{-1} \circ (\iota_{A_1} \times \dots \times \iota_{A_n}).$$

Thus, by letting $v = (\mathbf{T}u) \circ \iota_{A_1 \times \dots \times A_n} \circ (\iota_{A_1} \times \dots \times \iota_{A_n})^{-1}$, the result is obtained from Proposition 24. \square

Construction 26. Let $n \in \mathbb{N}$, and suppose that $m_R : R \rightarrow A_1 \times \dots \times A_n$ is a morphism in \mathbf{N} that is monic (and hence determines a relation) in \mathbf{M} ; i.e. m_R is a relation symbol in $\mathcal{S}_{\mathbf{N}}^{\mathbf{M}}$. Using the isomorphism obtained in Proposition 24, we can construct $\hat{\mathbf{T}}m_R : \hat{\mathbf{T}}R \rightarrow \mathbf{T}A_1 \times \dots \times \mathbf{T}A_n$ in \mathbf{N} as the pullback of $\mathbf{T}m_R$ along that isomorphism:

$$\begin{array}{ccc} \hat{\mathbf{T}}R & \xrightarrow{\sim} & \mathbf{T}R \\ \hat{\mathbf{T}}m_R \downarrow & & \downarrow \mathbf{T}m_R \\ \mathbf{T}A_1 \times \dots \times \mathbf{T}A_n & \xrightarrow{\sim} & \mathbf{T}(A_1 \times \dots \times A_n) \end{array}$$

Lemma 27. Let $m_R : R \rightarrow A_1 \times \dots \times A_n$ be as in Construction 26.

$$\mathbf{M} \models \forall x_1 : A_1 \dots \forall x_n : A_n. ((\mathbf{T}m_R)(\iota_{A_1 \times \dots \times A_n}(x_1, \dots, x_n)) \leftrightarrow (\hat{\mathbf{T}}m_R)(\iota_{A_1}(x_1), \dots, \iota_{A_n}(x_n))).$$

Proof. Everything commutes in this diagram:

$$\begin{array}{ccc} \hat{\mathbf{T}}R & \xrightarrow{\sim} & \mathbf{T}R \\ \hat{\mathbf{T}}m_R \downarrow & & \downarrow \mathbf{T}m_R \\ \mathbf{T}A_1 \times \dots \times \mathbf{T}A_n & \xrightarrow{\sim} & \mathbf{T}(A_1 \times \dots \times A_n) \\ & \searrow \sim & \swarrow \sim \\ & A_1 \times \dots \times A_n & \end{array}$$

Therefore, $(\mathbf{T}m_R) \circ \iota_{A_1 \times \dots \times A_n}$ and $(\hat{\mathbf{T}}m_R) \circ (\iota_{A_1} \times \dots \times \iota_{A_n})$ represent the same subobject of $A_1 \times \dots \times A_n$. \square

Let $n \in \mathbb{N}$. We define iterated application of \mathbf{P} , \mathbf{T} and $\hat{\mathbf{T}}$ in the usual way, as \mathbf{P}^n , etc. The iterated application of ι requires a special definition. We define $\iota_A^n : A \xrightarrow{\sim} \mathbf{T}^n A$ recursively by $\iota_A^0 = \text{id}_A$ and $\iota_A^{k+1} = \iota_{\mathbf{T}^k A} \circ \iota_A^k : A \xrightarrow{\sim} \mathbf{T}^k A \xrightarrow{\sim} \mathbf{T}^{k+1} A$, where $k \in \mathbb{N}$.

Lemma 28. Let $n, k \in \mathbb{N}$. Let $m_R : R \rightarrow A_1 \times \dots \times A_n$ be as in Construction 26.

$$\mathbf{M} \models \forall x_1 : A_1 \dots \forall x_n : A_n. (m_R(x_1, \dots, x_n) \leftrightarrow (\hat{\mathbf{T}}^k m_R)(\iota^k x_1, \dots, \iota^k x_n)).$$

Proof. Since ι is a natural isomorphism, we have by induction that $\iota^n : \text{id} \xrightarrow{\sim} \mathbf{T}^n$ is a natural isomorphism, so that

$$\mathbf{M} \models \forall x_1 : A_1 \dots \forall x_n : A_n. (m_R(x_1, \dots, x_n) \leftrightarrow (\mathbf{T}^k m_R)(\iota_{A_1 \times \dots \times A_n}^k(x_1, \dots, x_n))).$$

By iterating Lemma 27, we obtain by induction that

$$\mathbf{M} \models \forall x_1 : A_1 \dots \forall x_n : A_n. ((\mathbf{T}^k m_R)(\iota_{A_1 \times \dots \times A_n}^k(x_1, \dots, x_n)) \leftrightarrow (\hat{\mathbf{T}}^k m_R)(\iota_{A_1}^k x_1, \dots, \iota_{A_n}^k x_n)).$$

The result now follows by combining the two. \square

Construction 29. For any stratified ϕ in the language of $\mathcal{S}_{\mathbf{N},\in}^M$:

- Let $s_\phi : \{x_0, x_1, \dots, x_n\} \rightarrow \mathbb{N}$ be the minimal stratification of ϕ , and let \max_ϕ be the maximum value attained by s_ϕ .
- Let ϕ^t be the $\mathcal{S}_{\mathbf{N},\iota}^M$ -formula obtained from ϕ by the construction below. Each atomic subformula θ of ϕ is replaced as specified in the following two mutually exclusive and jointly exhaustive cases.
 1. Suppose that $\theta \equiv m_R(x_1, \dots, x_n)$, where x_1, \dots, x_n are variables and $m_R : R \rightarrow A_1 \times \dots \times A_n$ is in $\mathcal{S}_{\mathbf{N}}^M$. Note that by stratification, $s_\phi(x_1) = \dots = s_\phi(x_n)$. Let $k = \max_\phi - s_\phi(x_1)$. In ϕ , replace θ by

$$(\hat{\mathbf{T}}^k m_R)(\iota^k x_1, \dots, \iota^k x_n).$$

2. Suppose that $\theta \equiv u(x_1, \dots, x_n) \in_A v(y_1, \dots, y_m)$, where A is an object in \mathbf{N} . Note that by stratification, $s_\phi(x_i) + 1 = s_\phi(y_j)$, for each $1 \leq i \leq n$ and $1 \leq j \leq m$. Let $k_x = \max_\phi - s_\phi(x_1)$ and $k_y = \max_\phi - s_\phi(y_1)$. Replace θ by

$$u'(\iota^{k_x} x_1, \dots, \iota^{k_x} x_n)(\hat{\mathbf{T}}^{k_y} \subseteq^1)v'(\iota^{k_y} y_1, \dots, \iota^{k_y} y_m),$$

where u' and v' are the morphisms in \mathbf{N} obtained from u and v , respectively, by applying Lemma 25.

- Let ϕ^{\subseteq^1} be the formula in the language of $\mathcal{S}_{\mathbf{N}}^M$ obtained from ϕ^t as follows.
 - Replace each term of the form $\iota^{\max_\phi - s_\phi(x)}(x)$ (where x is a variable of sort A) by a fresh variable x' (of sort $\mathbf{T}^{\max_\phi - s_\phi(x)}A$).
 - Replace each quantifier scope declaration $x : A$ by $x' : \mathbf{T}^{\max_\phi - s_\phi(x)}A$.

It is easily seen that ϕ^t is an $\mathcal{S}_{\mathbf{N},\iota}^M$ -formula. Moreover, note that every variable x in ϕ^t occurs in a term $\iota^{\max_\phi - s_\phi(x)}x$, and conversely, every occurrence of ι in ϕ^t is in such a term $\iota^{\max_\phi - s_\phi(x)}x$. Therefore, ϕ^{\subseteq^1} is an $\mathcal{S}_{\mathbf{N}}^M$ -formula.

Proposition 30. *If ϕ is a stratified $\mathcal{S}_{\mathbf{N},\in}^M$ -sentence, then $\mathbf{M} \models \phi \Leftrightarrow \mathbf{M} \models \phi^t \Leftrightarrow \mathbf{M} \models \phi^{\subseteq^1}$.*

Proof. We start by showing that $\mathbf{M} \models \phi \Leftrightarrow \phi^t$. By induction, we only need to show that every atomic subformula of ϕ is replaced by an equivalent subformula.

Case 1: Suppose that $\theta \equiv m_R(x_1, \dots, x_n)$, where $m_R : R \rightarrow A_1 \times \dots \times A_n$ is in $\mathcal{S}_{\mathbf{N}}^M$. By Lemma 28,

$$\mathbf{M} \models m_R(x_1, \dots, x_n) \Leftrightarrow (\hat{\mathbf{T}}^k m_R)(\iota^k x_1, \dots, \iota^k x_n).$$

Case 2: Suppose that $\theta \equiv u(x_1, \dots, x_n) \in_A v(y_1, \dots, y_m)$, where A is an object in \mathbf{N} . Let k_x, k_y, u', v' be as in Construction 29. By Proposition 16,

$$\mathbf{M} \models u(x_1, \dots, x_n) \in_A v(y_1, \dots, y_m) \Leftrightarrow \iota(u(x_1, \dots, x_n)) \subseteq_A^1 v(y_1, \dots, y_m).$$

Since $k_x = k_y + 1$, we have by Lemma 28 that

$$\iota(u(x_1, \dots, x_n)) \subseteq_A^1 v(y_1, \dots, y_m) \Leftrightarrow \iota^{k_x}(u(x_1, \dots, x_n))(\hat{\mathbf{T}}^{k_y} \subseteq_A^1) \iota^{k_y} v(y_1, \dots, y_m).$$

By Lemma 25,

$$\mathbf{M} \models \iota^{k_x}(u(x_1, \dots, x_n))(\hat{\mathbf{T}}^{k_y} \subseteq_A^1) \iota^{k_y} v(y_1, \dots, y_m) \Leftrightarrow u'(\iota^{k_x} x_1, \dots, \iota^{k_x} x_n)(\hat{\mathbf{T}}^{k_y} \subseteq_A^1) v'(\iota^{k_y} y_1, \dots, \iota^{k_y} y_m).$$

Putting the three together, we obtain

$$\mathbf{M} \models u(x_1, \dots, x_n) \in_A v(y_1, \dots, y_m) \leftrightarrow u'(\iota^{k_x}(x_1), \dots, \iota^{k_x}(x_n))(\hat{\mathbf{T}}^{k_y} \subseteq_A^1) v'(\iota^{k_y}(y_1), \dots, \iota^{k_y}(y_m)).$$

Thus, we have established $\mathbf{M} \models \phi \Leftrightarrow \mathbf{M} \models \phi^\iota$.

To see that $\mathbf{M} \models \phi^\iota \Leftrightarrow \mathbf{M} \models \phi^{\subseteq^1}$, note that for each variable x (say of sort A) in ϕ^ι , $\iota^{\max_\phi - s_\phi(x)}$ is an isomorphism from A to $\mathbf{T}^{\max_\phi - s_\phi(x)} A$. So the equivalence follows by Construction 29 from the categorical semantics. \square

Proposition 31 (Stratified Comprehension). *For every stratified $\mathcal{S}_{\mathbf{N}, \in}^{\mathbf{M}}$ -formula $\phi(z, y)$ with only z and y free, in context $z : Z, y : Y$ for Z, Y in \mathbf{M} ,*

$$\mathbf{M} \models \forall y : Y. \exists x : \mathbf{P}Z. \forall z : Z. (z \in x \leftrightarrow \phi(z, y)).$$

Proof. By Proposition 30, we have

$$\begin{aligned} & \mathbf{M} \models \forall y : Y. \exists x : \mathbf{P}Z. \forall z : Z. (z \in_Z x \leftrightarrow \phi(z, y)) \\ \Leftrightarrow & \mathbf{M} \models \forall y' : \mathbf{T}^m Y. \exists x' : \mathbf{T}^k \mathbf{P}Z. \forall z' : \mathbf{T}^{k+1} Z. (z' (\hat{\mathbf{T}}^k \subseteq_Z^1) x' \leftrightarrow \phi^{\subseteq^1}(z', y')), \end{aligned} \quad (\dagger)$$

for some $k, m \in \mathbb{N}$, where x', y', z' are fresh variables.

In order to apply Proposition 23, we need to move the \mathbf{T} :s through the \mathbf{P} and transform the $\hat{\mathbf{T}}^k \subseteq_Z^1$ into a $\subseteq_{\mathbf{T}^k Z}^1$. Since $\mu : \mathbf{P}\mathbf{T} \rightarrow \mathbf{T}\mathbf{P}$ is a natural isomorphism,

$$\nu =_{\text{df}} \mathbf{T}^{k-1}(\mu_Z) \circ \mathbf{T}^{k-2}(\mu_{\mathbf{T}Z}) \circ \dots \circ \mathbf{T}(\mu_{\mathbf{T}^{k-2}Z}) \circ \mu_{\mathbf{T}^{k-1}Z} : \mathbf{P}\mathbf{T}^k Z \xrightarrow{\sim} \mathbf{T}^k \mathbf{P}Z,$$

is an isomorphism making this a pullback square:

$$\begin{array}{ccc} \subseteq_{\mathbf{T}^k Z}^1 & \xrightarrow{\sim} & \hat{\mathbf{T}}^k \subseteq_Z^1 \\ \downarrow m_{\subseteq_{\mathbf{T}^k Z}^1} & & \downarrow \hat{\mathbf{T}}^k m_{\subseteq_Z^1} \\ \mathbf{T}^{k+1} Z \times \mathbf{P}\mathbf{T}^k Z & \xrightarrow[\text{id} \times \nu]{\sim} & \mathbf{T}^{k+1} Z \times \mathbf{T}^k \mathbf{P}Z \end{array}$$

So introducing a fresh variable $x'' : \mathbf{P}\mathbf{T}^k Z$, (\dagger) is equivalent to

$$\mathbf{M} \models \forall y' : \mathbf{T}^m Y. \exists x'' : \mathbf{P}\mathbf{T}^k Z. \forall z' : \mathbf{T}^{k+1} Z. (z' \subseteq_{\mathbf{T}^k Z}^1 x'' \leftrightarrow \phi^{\subseteq^1}(z', y')).$$

By Proposition 23 we are done. \square

Corollary 32. $\mathcal{U} \models \text{SC}_S$

Proof. Let $\phi(z, y)$ be a stratified formula in \mathcal{L}_{SET} , with only z, y free. By Proposition 31, $\mathbf{M} \models \forall y : U. \exists x : \mathbf{P}U. \forall z : U. (z \in x \leftrightarrow \phi(z, y))$. Now,

$$\begin{aligned} & \llbracket y : U \mid \exists x : \mathbf{P}U. \forall z : U. (z \in x \leftrightarrow \phi(z, y)) \rrbracket \\ = & \llbracket y : U \mid \exists x : \mathbf{P}U. \forall z : U. (z \varepsilon m_S(x) \leftrightarrow \phi(z, y)) \rrbracket \\ = & \llbracket y : U \mid \exists x' : U. (S(x') \wedge \forall z : U. (z \varepsilon x' \leftrightarrow \phi(z, y))) \rrbracket. \end{aligned}$$

So $\mathcal{U} \models \text{SC}_S$. \square

Theorem 33. $\mathcal{U} \models (\text{I})\text{NFU}$, and if $U \cong \mathbf{P}U$, then $\mathcal{U} \models (\text{I})\text{NF}$. Thus, each of $(\text{I})\text{NF}(U)_{\text{SET}}$ is interpretable in $(\text{I})\text{ML}(U)_{\text{CAT}}$, respectively.

Proof. The cases of (I)NFU are settled by the results above on Sethood, Ordered Pair, Extensionality and Stratified Comprehension. The cases of (I)NF now follow from Lemma 19. \square

Theorem 34. *The following theories are, respectively, equiconsistent.*

- (I)NF(U)_{SET}
- (I)ML(U)_{CLASS}
- (I)ML(U)_{CAT}

Moreover, (I)ML(U)_{CLASS} interprets (I)ML(U)_{CAT}, which in turn interprets (I)NF(U)_{SET}.

Proof. Combine Theorems 5, 13 and 33. \square

Corollary 35. (I)NF_{SET} is equiconsistent with (I)NFU_{SET} + ($|V| = |\mathcal{P}(V)|$).

Proof. Only \Leftarrow is non-trivial. By the proofs above, $\text{Con}((\text{I)NFU}_{\text{SET}} + (|V| = |\mathcal{P}(V)|)) \Rightarrow \text{Con}((\text{I)MLU}_{\text{CLASS}} + (|V| = |\mathcal{P}(V)|)) \Rightarrow \text{Con}((\text{I)ML}_{\text{CAT}}) \Rightarrow \text{Con}((\text{I)NF}_{\text{SET}})$. \square

8 The subtopos of strongly Cantorian objects

Definition 36. An object X in \mathbf{N} is *Cantorian* iff $X \cong \mathbf{T}X$ in \mathbf{N} , and is *strongly Cantorian* iff $\iota_X : X \xrightarrow{\sim} \mathbf{T}X$ is an isomorphism in \mathbf{N} . Define $\mathbf{SCan}_{\langle \mathbf{M}, \mathbf{N} \rangle}$ as the full subcategory of \mathbf{N} on the set of strongly Cantorian objects. When the subscript $\langle \mathbf{M}, \mathbf{N} \rangle$ is clear from the context, we simply write \mathbf{SCan} .

Proposition 37. $\mathbf{SCan}_{\langle \mathbf{M}, \mathbf{N} \rangle}$ has finite limits.

Proof. Let L be a limit in \mathbf{N} of a finite diagram $\mathbf{D} : \mathbf{I} \rightarrow \mathbf{SCan}$. Since \mathbf{N} is a Heyting subcategory of \mathbf{M} , L is a limit of \mathbf{D} in \mathbf{M} ; and since \mathbf{T} preserves limits, $\mathbf{T}L$ is a limit of $\mathbf{T} \circ \mathbf{D}$ in \mathbf{M} and in \mathbf{N} . But also, since \mathbf{D} is a diagram in \mathbf{SCan} , $\mathbf{T}L$ is a limit of \mathbf{D} , and L is a limit of $\mathbf{T} \circ \mathbf{D}$, in \mathbf{M} and in \mathbf{N} . So there are unique morphisms in \mathbf{N} back and forth between L and $\mathbf{T}L$ witnessing the universal property of limits. Considering these as morphisms in \mathbf{M} we see that they must be ι_L and ι_L^{-1} . Now L is a limit in \mathbf{SCan} by fullness. \square

Before we can show that $\mathbf{SCan}_{\langle \mathbf{M}, \mathbf{N} \rangle}$ has power objects, we need to establish results showing that $\mathbf{SCan}_{\langle \mathbf{M}, \mathbf{N} \rangle}$ is a “nice” subcategory of \mathbf{N} .

Corollary 38. *The inclusion functor of $\mathbf{SCan}_{\langle \mathbf{M}, \mathbf{N} \rangle}$ into \mathbf{N} preserves and reflects limits.*

Proof. To see that it reflects limits, simply repeat the proof of Proposition 37. We proceed to show that it preserves limits.

Let L be a limit in \mathbf{SCan} of a finite diagram $\mathbf{D} : \mathbf{I} \rightarrow \mathbf{SCan}$. Let L' be a limit of this diagram in \mathbf{N} . By the proof of Proposition 37, L' is also such a limit in \mathbf{SCan} , whence L is isomorphic to L' in \mathbf{SCan} , and in \mathbf{N} . So L is a limit of \mathbf{D} in \mathbf{N} as well. \square

Corollary 39. $m : A \rightarrow B$ is monic in $\mathbf{SCan}_{\langle \mathbf{M}, \mathbf{N} \rangle}$ iff it is monic in \mathbf{N} .

Proof. (\Leftarrow) follows from that \mathbf{SCan} is a subcategory of \mathbf{N} .

(\Rightarrow) Assume that $m : A \rightarrow B$ is monic in \mathbf{SCan} . Then A along with $\text{id}_A : A \rightarrow A$ and $\text{id}_A : A \rightarrow A$ is a pullback of m and m in \mathbf{SCan} . By Corollary 38, this is also a pullback in \mathbf{N} , from which it follows that m is monic in \mathbf{N} . \square

Proposition 40. *If $m : A \rightarrow B$ is monic in \mathbf{N} and B is in $\mathbf{SCan}_{\langle \mathbf{M}, \mathbf{N} \rangle}$, then A is in $\mathbf{SCan}_{\langle \mathbf{M}, \mathbf{N} \rangle}$.*

Proof. Let $m : A \rightarrow B$ be a mono in \mathbf{N} and assume that ι_B is in \mathbf{N} . Let P be this pullback in \mathbf{N} , which is also a pullback in \mathbf{M} since \mathbf{N} is a Heyting subcategory:

$$\begin{array}{ccc} P & \xrightarrow{n} & B \\ \langle p, q \rangle \downarrow & & \downarrow \langle \text{id}_B, \iota_B \rangle \\ A \times \mathbf{T}A & \xrightarrow{m \times \mathbf{T}m} & B \times \mathbf{T}B \end{array}$$

We shall now establish that the following square is also a pullback in \mathbf{M} :

$$\begin{array}{ccc} A & \xrightarrow{m} & B \\ \langle \text{id}_A, \iota_A \rangle \downarrow & & \downarrow \langle \text{id}_B, \iota_B \rangle \\ A \times \mathbf{T}A & \xrightarrow{m \times \mathbf{T}m} & B \times \mathbf{T}B \end{array}$$

The square commutes since ι is a natural isomorphism. So since P is a pullback, it suffices to find $f : P \rightarrow A$ such that $\langle p, q \rangle = \langle \text{id}_A, \iota_A \rangle \circ f$ and $n = m \circ f$. Let $f' = p$ and let $f'' = \iota_A^{-1} \circ q$. We shall show that $f' = f''$ and that this is the desired f . By commutativity of the former square, $n = m \circ f'$ and $\iota_B \circ n = \mathbf{T}m \circ \iota_A \circ f''$, whence

$$\iota_B \circ m \circ f' = \mathbf{T}m \circ \iota_A \circ f''.$$

Note that $\iota_B \circ m$ is monic, and since ι is a natural transformation it is equal to $\mathbf{T}m \circ \iota_A$. Hence, $f' = f''$. Let $f = f' = f''$. We have already seen that $n = m \circ f$. That $\langle p, q \rangle = \langle \text{id}_A, \iota_A \rangle \circ f$ is immediately seen by plugging the definitions of f' and f'' in place of f . Since P is a pullback, it follows that f is an isomorphism in \mathbf{M} and that the latter square is a pullback.

We can now establish that $f : P \rightarrow A$ and $\iota_A \circ f : P \rightarrow \mathbf{T}A$ are in \mathbf{N} , by noting that

$$\begin{aligned} \pi_{A \times \mathbf{T}A}^1 \circ \langle p, q \rangle &= \pi_{A \times \mathbf{T}A}^1 \circ \langle \text{id}_A, \iota_A \rangle \circ f = f \\ \pi_{A \times \mathbf{T}A}^2 \circ \langle p, q \rangle &= \pi_{A \times \mathbf{T}A}^2 \circ \langle \text{id}_A, \iota_A \rangle \circ f = \iota_A \circ f \end{aligned}$$

Since f is an isomorphism in \mathbf{M} , and \mathbf{N} is a conservative subcategory, f is an isomorphism in \mathbf{N} , whence $\iota_A = \iota_A \circ f \circ f^{-1}$ is in \mathbf{N} , as desired. \square

Proposition 41. *$\mathbf{SCan}_{\langle \mathbf{M}, \mathbf{N} \rangle}$ has power objects.*

Proof. Let A be in \mathbf{SCan} . We shall show that $\mathbf{P}A$ along with $(\iota_A^{-1} \times \text{id}_{\mathbf{P}A}) \circ m_{\subseteq_A^1} : \subseteq_A^1 \rightarrow A \times \mathbf{P}A$ is a power object of A in \mathbf{SCan} . In Step 1 we show that $(\iota_A^{-1} \times \text{id}_{\mathbf{P}A}) \circ m_{\subseteq_A^1} : \subseteq_A^1 \rightarrow A \times \mathbf{P}A$ is in \mathbf{SCan} , and in Step 2 we show that it satisfies the power object property.

Step 1: It actually suffices to show that $\mathbf{P}A$ is in \mathbf{SCan} . Because then, by Proposition 37 and Corollary 38, $A \times \mathbf{P}A$ is in \mathbf{SCan} (and is such a product in both \mathbf{N} and \mathbf{SCan}), so that by Proposition 40 and fullness of \mathbf{SCan} , $(\iota_A^{-1} \times \text{id}_{\mathbf{P}A}) \circ m_{\subseteq_A^1} : \subseteq_A^1 \rightarrow A \times \mathbf{P}A$ is in \mathbf{SCan} .

Since $\mathbf{P}A$ along with $m_{\subseteq_{\mathbf{T}A}^1} : \subseteq_{\mathbf{T}A}^1 \rightarrow \mathbf{T}A \times \mathbf{P}A$ is a power object of $\mathbf{T}A$, we have that $\mathbf{P}A$ along with $(\iota_{\mathbf{T}A}^{-1} \times \text{id}_{\mathbf{P}A}) \circ m_{\subseteq_{\mathbf{T}A}^1} : \subseteq_{\mathbf{T}A}^1 \rightarrow \mathbf{T}A \times \mathbf{P}A$ is a power object of $\mathbf{T}A$ in \mathbf{N} . Moreover, $\mathbf{P}A$ is a power object of $\mathbf{T}A$ in \mathbf{N} . Therefore, using the natural isomorphism $\mu : \mathbf{P}\mathbf{T} \rightarrow \mathbf{T}\mathbf{P}$, we obtain an isomorphism $\alpha : \mathbf{P}A \xrightarrow{\sim} \mathbf{P}A \xrightarrow{\sim} \mathbf{T}\mathbf{P}A$ in \mathbf{N} . This results in the following two-way pullback in both \mathbf{M} and \mathbf{N} :

$$\begin{array}{ccc} \subseteq^1 & \xleftarrow{\sim} & \subseteq^1 \\ \downarrow & & \downarrow \\ \mathbf{T}A \times \mathbf{P}A & \xrightarrow{\text{id} \times \alpha} & \mathbf{T}A \times \mathbf{T}\mathbf{P}A \\ & \xleftarrow{\sim} & \\ & \text{id} \times \alpha^{-1} & \end{array}$$

Since this can be filled with $\iota_{\mathbf{P}A}$ in place of α , the uniqueness property in \mathbf{M} implies that $\iota_{\mathbf{P}A} = \alpha$ is in \mathbf{N} , which means that $\mathbf{P}A$ is in \mathbf{SCan} .

Step 2: Let $r : R \rightarrow A \times B$ be a mono in \mathbf{SCan} . By Corollary 39, r is also monic in \mathbf{N} . So since ι_A is an isomorphism in \mathbf{N} , there is a unique χ in \mathbf{N} such that this is a pullback in \mathbf{N} :

$$\begin{array}{ccc} R & \longrightarrow & \subseteq_A^1 \\ \downarrow r & & \downarrow (\iota_A^{-1} \times \text{id}_{\mathbf{P}A}) \circ m_{\subseteq_A^1} \\ A \times B & \xrightarrow{\text{id}_A \times \chi} & A \times \mathbf{P}A \end{array}$$

By Step 1, by fullness and by Corollary 38, it is also a pullback in \mathbf{SCan} . To see uniqueness of χ in \mathbf{SCan} , suppose that χ' were some morphism in \mathbf{SCan} making this a pullback in \mathbf{SCan} (in place of χ). Then by Corollary 38, it would also make it a pullback in \mathbf{N} , whence $\chi = \chi'$. \square

Theorem 42. $\mathbf{SCan}_{(\mathbf{M}, \mathbf{N})}$ is a topos.

Proof. A category with finite limits and power objects is a topos. \square

9 Where to go from here?

$(\mathbf{I}ML(\mathbf{U}))_{\text{CAT}}$ has been shown, respectively, to interpret $(\mathbf{I}NF(\mathbf{U}))_{\text{SET}}$, and has conversely been shown to be interpretable in $(\mathbf{I}ML(\mathbf{U}))_{\text{CLASS}}$, thus yielding equiconsistency results. Since the axioms of a Heyting category can be obtained from the axioms of topos theory, it is natural to ask:

Question 43. Can the axioms of $(\mathbf{I}ML(\mathbf{U}))_{\text{CAT}}$ be simplified? In particular, is it necessary to include the axioms of Heyting categories or do these follow from the other axioms?

To be able to interpret the set theory in the categorical semantics, this paper introduces the axiomatization $(\mathbf{I}ML(\mathbf{U}))_{\text{CAT}}$ roughly corresponding the predicative class theory of NF called ML. This is analogous to the categories of classes for conventional set theory studied e.g. in [ABSS, 2014]. But it remains to answer:

Question 44. How should the speculative theory $(\mathbf{I}NF(\mathbf{U}))_{\text{CAT}}$ naturally be axiomatized? I.e. what is the natural generalization of $(\mathbf{I}NF(\mathbf{U}))_{\text{SET}}$ to category theory, analogous to topos theory as the natural generalization of conventional set theory? Moreover, can any category modelling this theory be canonically extended to a model of $(\mathbf{I}ML(\mathbf{U}))_{\text{CAT}}$, or under what conditions are such extensions possible?

Closely intertwined with this question, is the potential project of generalizing to topos theory the techniques of automorphisms and self-embeddings of nonstandard models of set theory. In particular, the endofunctor \mathbf{T} considered in this paper should arise from an automorphism or self-embedding of a topos. This would be a natural approach to constructing a rich variety of categories modelling the speculative theory $(\mathbf{I}NF(\mathbf{U}))_{\text{CAT}}$, many of which would presumably be extendible to models of $(\mathbf{I}ML(\mathbf{U}))_{\text{CAT}}$.

References

[ABSS, 2014] Steve Awodey, Carsten Butz, Alex Simpson and Thomas Streicher. *Relating first-order set theories, toposes and categories of classes*. Annals of Pure and Applied Logic, Vol. 165 (2014), pp. 428-502.

- [Cantini, 2015] Andrea Cantini. *On Stratified Truth*. In *Unifying the Philosophy of Truth*, Springer (2015), pp. 369-389.
- [Cocchiarella, 1985] Nino Cocchiarella. *Frege's Double-Correlation Thesis and Quine's Set Theories NF and ML*. *Journal of Philosophical Logic*, Vol. 14, No. 1 (1985), pp. 1-39.
- [Crabbé, 2000] Marcel Crabbé. *On the Set of Atoms*. *Logic Journal of the IGPL*, Vol. 8, No. 6 (2000), pp. 751-759.
- [Dzierzowski, 1995] Daniel Dzierzowski. *Models of Intuitionistic TT and NF*. *Journal of Symbolic Logic*, Vol. 60 (1995), pp. 640-653.
- [Enayat, 2004] Ali Enayat. *Automorphisms, Mahlo Cardinals, and NFU*. In *Nonstandard Models of Arithmetic and Set Theory, Contemporary Mathematics*, vol. 361, American Mathematical Society (2004), pp. 37-59.
- [EGM, 2017] Ali Enayat, Paul Gorbow, and Zachiri McKenzie. *Feferman's forays into the foundations of category theory*. In *Feferman on Foundations: Logic, Mathematics, Philosophy, Outstanding Contributions to Logic Series*, Springer Verlag, in press.
- [Feferman, 2006] Solomon Feferman. *Enriched Stratified systems for the Foundations of Category Theory*. In *What is category theory?*, *Polimetrika* (2006). Preprint on-line: <http://math.stanford.edu/~feferman/papers/ess.pdf>
- [Forster, 1995] Thomas Forster. *Set Theory with a Universal Set*. Oxford University Press, 2:nd edition (1995).
- [FLV, 2014] Thomas Forster, Adam Lewicki and Alice Vidrine. *The category of sets in stratifiable set theories*. Unpublished (2014).
- [Hailperin, 1944] Theodore Hailperin. *A set of axioms for logic*. *Journal of Symbolic Logic*, Vol. 9 (1944), pp. 1-19.
- [Holmes, 1998] M. Randall Holmes. *Elementary Set Theory with a Universal Set*. *Cahiers du Centre de logique*, Vol. 10 (1998). Preprint of revised and corrected version on-line: <http://math.boisestate.edu/~holmes/holmes/head.pdf>
- [Holmes, 2015] M. Randall Holmes. *Repairing Frege's Logic*. Online: <https://math.boisestate.edu/~holmes/holmes/frege-note.pdf> (2015)
- [Jensen, 1969] Ronald B. Jensen. *On the consistency of a slight (?) modification of Quine's NF*. *Synthese*, Vol. 19 (1969), pp. 250-263.
- [Johnstone, 2002] Peter T. Johnstone. *Sketches of an Elephant*. Vol. 2. Oxford University Press (2002).
- [JM, 1991] André Joyal and Ieke Moerdijk. *A categorical theory of cumulative hierarchies of sets*. *Comptes Rendus Mathématiques de l'Académie des Science*, Vol. 13 (1991), pp. 55-58.
- [JM, 1995] André Joyal and Ieke Moerdijk. *Algebraic Set Theory*. Cambridge University Press (1995).
- [Lawvere, 1963] F. William Lawvere. *Functorial Semantics of Algebraic Theories*. In *Proceedings of the National Academy of Sciences*, Vol. 50, No. 5 (1963), pp. 869-872.

- [McKenzie, 2015] Zachiri McKenzie. *Automorphisms of models of set theory and extensions of NFU*. *Annals of Pure and Applied Logic*, Vol. 166 (2015), pp. 601-638.
- [McLarty, 1992] Colin McLarty. *Failure of Cartesian Closedness in NF*. *The Journal of Symbolic Logic*, Vol. 57, No. 2 (1992), pp. 555-556.
- [MM, 1992] Saunders Mac Lane and Ieke Moerdijk. *Sheaves in Geometry and Logic*. Springer-Verlag (1992).
- [Quine, 1937] Willard V. Quine. *New foundations for mathematical logic*. *American Mathematical Monthly*, Vol. 44 (1937), pp. 111-115.
- [Quine, 1940] Willard V. Quine. *Mathematical logic*. New York, 1940.
- [Specker, 1953] Ernst P. Specker. *The axiom of choice in Quine's 'New Foundations for Mathematical Logic'*". *Proceedings of the National Academy of Sciences of the U. S. A.*, Vol. 39 (1953), pp. 972-975.
- [Solovay, 1997] Robert Solovay. *The consistency strength of NFUB*. arXiv:math/9707207 [math.LO] (1997).
- [Thomas, 2017] Morgan Thomas. *Approximating Cartesian Closed Categories in NF-Style Set Theories*. *Journal of Philosophical Logic* (2017). doi:10.1007/s10992-017-9425-2
- [Wang, 1950] Hao Wang. *A Formal System of Logic*. *The Journal of Symbolic Logic*, Vol. 15, No. 1 (1950), pp. 25-32.