PIERRE-PAUL ROMAGNOLI

ABSTRACT. In this paper, we introduce two measure theoretical notions of conditional entropy for finite measurable covers conditioned to a finite measurable partition and prove that they are equal. Using this we state a local variational principle with respect to the notion of conditional entropy defined by Misiurewicz in [M] for the case of open covers. This in particular extends the work done in [R] and [HYZ].

1. INTRODUCTION

Topological dynamics and Ergodic Theory exhibit a remarkable parallelism. It is usual to find counterparts in both theories as is the case of transitivity, weak mixing and strong mixing and in particular with the notions that we study in this paper, this is topological entropy and measure theoretical entropy. Although, even if the notions and results are similar the methods to prove them are quite different. In many cases the connection between these notions is through a variational principle by taking suprema over all invariant measures to obtain the topological notion. For the case of \mathbb{Z} actions Measure-theoretical Entropy for an invariant measure was introduced in 1958 in [K] and Topological Entropy in 1965 in [AKM]. In 1969 and 1970 Goodman [G] and Goodwyn [Gw] in two separate papers proved the first variational principle. M. Misiurewicz introduced the notion of topological conditional entropy in [M] and the variational principle was established by F. Ledrappier in [L].

This paper is inserted in the so called local theory of entropy for Topological Dynamical Systems that started in the early 90's with the work of François Blanchard (see [B] and [BL]) that is interesting by itself but has also proven to be fundamental to many other related areas. For instance the existence of topological Pinsker factors, zero entropy factors, disjointness theorems, characterizations of positive entropy, entropy pairs and tuples and so on.

In [BGH] while extending the notion of topological entropy pairs to a measure theoretical setting the authors proved a *variational principle for open covers* for a topological dynamical system (X, T). More precisely, for every open cover there exists a *T*-invariant measure μ such that the topological entropy of the cover is bounded above by the μ -entropy of every partition finer than the cover.

In [R] the author gave a new approach to the variational principle given in [BGH] extending the measure theoretical notions from partitions to covers and thus proposing two measure theoretical notions of entropy for open covers denoted h^+_{μ} and h^-_{μ} . In doing so he was then able to state and prove a local variational principle for h^-_{μ} . Namely, for every open cover there exists a *T*-invariant measure

PIERRE-PAUL ROMAGNOLI

 μ such that the topological entropy is equal to the h_{μ}^{-} -entropy of the cover. The notion h_{μ}^{-} verifies most of the relevant properties of the usual notion of entropy for measurable partitions. It is also proven that the infimum that appears in the definition of h_{μ}^{-} is attained. This fact implies some good properties, for instance it implies that h_{μ}^{-} is preserved by measure-theoretic extension and as a function of μ , h_{μ}^{-} is upper semicontinuos (see [HY]). The main open question left in [R] was to know if the two notions were equal.

In [HMRY] the authors proved that these notions were positive simultaneously and, using the Jewett-Krieger Theorem they proved that the equality between h^+_{μ} and h^-_{μ} is equivalent to the existence of a local variational principle for h^+_{μ} (similar to the one proven for h^-_{μ}). In [GW] such a variational principle was established for h^+_{μ} and thus the equality holds. An alternative definition for measure entropy for covers was given in [S] giving a new proof of the variational principle and at the same time proving the equality of h^+_{μ} and h^-_{μ} once again.

In [HY] the authors applied the same ideas to extend the notion of topological pressure to a measure theoretical setting proving again a local variational principle of the same topological pressure and in [HYZ] a similar one in a conditional version with respect to a factor.

For the existence of a variational principle the requirement of the cover to be open is unavoidable. However the equality between h^+_{μ} and h^-_{μ} can be stated in general for measurable covers extending the topological notion to this kind of covers since it is turns out to be mainly combinatorial. This was well known even in the first known proofs of the global variational principle. The global bound of the variational principle fails even for nice non open covers as is the case of closed covers where for example the existence of one non recurrent point implies that the suprema of the entropy over all closed covers is infinite. This was reviewed and explained by Goodwyn in [Gw2] (see example 3) although it was already mentioned by him in [Gw].

Since then, the research has mainly focused in extending the notions of entropy and pressure to more general actions. In [HYZ2] it was done for the case countable discrete amenable groups by using the tool of Følner sequences making the techniques and the proofs very similar. The case of continuous bundle random dynamical systems of an infinite countable discrete amenable group action has been proven in [DZ] and for sofic group actions in [Z].

However the main idea that gives the natural bridge between the measurable and the topological notions in every one of the versions of variational principles discussed so far is the one given in [R]. This is, extended the measurable formulas that apply to partitions to covers by considering the infima over all partitions finer than the cover. Then for measurable covers the natural "+" and "-" definitions prove that they are equal for all measurable covers and then use this to prove a variational principle for open covers for the suitable topological equivalent.

Unfortunately the real local variational problem is still open, this is, with respect to a fixed open cover conditioned to another fixed open cover. The missing ingredient is a clear way to extend the notion with respect to the conditioning partition. Extending the definition with respect to the conditioning variable in the same way by taking infima is not the way to go for a local definition and even for a fixed partition using the conditional measurable definition of partitions is not enough.

Considering the alternative definition of conditional entropy as an average of the entropy of the first partition with respect to the induced measures over the atoms of the conditioning partition and using infima in a clever way solves the issue giving once again two different definitions that turn out to be the same. In this paper we address this for the first time proving the local conditional version with respect to a finite measurable partition.

More precisely, in this paper we propose two notions of conditional measure theoretical entropy for measurable finite covers conditioned to a fixed measurable partition that extends the notions h^-_{μ} and h^+_{μ} given in [R] and [HYZ] and proving in general that they coincide for every finite measurable cover. In the case of finite open covers we prove that they satisfy a local variational principle with the notion of conditional entropy defined by Misiurewicz in [M] once again extending the results in [R] and [HYZ].

2. Basic definitions and results

We will define a series of notions as limits that are also infima using the classic subadditive lemma that we state without proof:

Lemma 1. For any subadditive sequence $\{a_N\}_{N \in \mathbb{N}} \subseteq \mathbb{R}^+$, this is $a_{N+M} \leq a_N + a_M$ for any $N, M \in \mathbb{N}$ we have:

(1)
$$\lim_{N \to \infty} \frac{1}{N} a_N = \inf_{N \in \mathbb{N}} \frac{1}{N} a_N.$$

Proof: See [D].

Let us introduce the basic notation and definitions used in this article. For more details on measure-theoretical and topological entropy including the most recent results of the beginning of the year 2000 we refer to the deeply insightful and inspirational textbook by Tomasz Downarowicz [D].

 \square

Let (X,T) be a topological dynamical system (TDS). This is, X a compact metric space and $T: X \to X$ a homeomorphism. A TDS (X,T) is 0-dimensional when the space X has a countable open-closed (clopen) topological basis. The set $\mathcal{M}_T(X)$ of T-invariant Borel probability measures is a convex, compact (in the weak topology) and nonempty set. We denote $\mathcal{M}_T^e(X)$ the set of ergodic measures. A measure theoretic dynamical system (MDS), (X, \mathcal{B}, μ, T) , is a probability space (X, \mathcal{B}, μ) and a bi-measurable bijection $T: X \to X$ that preserves the measure μ . So a TDS (X, T) gives a family of MDS indexed by the set $\mathcal{M}_T(X)$ where \mathcal{B} is the σ -algebra of the borel sets of X.

In this article a cover of X is a finite cover of Borel subsets of X. A cover is said to be an *open cover* if it consists only of open sets. A *partition of* X is a cover by pairwise disjoint sets. Let \mathcal{P}_X denote the set of partitions of X, \mathcal{C}_X the set of covers of X and \mathcal{C}_X^o the set of open covers of X.

Given two covers $\mathcal{U}, \mathcal{V} \in \mathcal{C}_X, \mathcal{U}$ is said to be finer than \mathcal{V} $(\mathcal{U} \succeq \mathcal{V})$ if for every $U \in \mathcal{U}$, there is $V \in \mathcal{V}$ such that $U \subseteq V$. Let $\mathcal{U} \lor \mathcal{V} = \{U \cap V / U \in \mathcal{U}, V \in \mathcal{V}\}$.

It is clear that $\mathcal{U} \vee \mathcal{V} \succeq \mathcal{U}$ and $\mathcal{U} \vee \mathcal{V} \succeq \mathcal{V}$. However, $\mathcal{V} \succeq \mathcal{U}$ does not imply that $\mathcal{U} \vee \mathcal{V} = \mathcal{V}$. Given integers $M \leq N$ and $\mathcal{U} \in \mathcal{C}_X$, one sets, $\mathcal{U}_M^N = \bigvee_{n=M}^N T^{-n}\mathcal{U}$. For $\mathcal{U} = \{U_1, \ldots, U_M\}$ define $\mathcal{U}^* = \{\alpha = \{A_1, \ldots, A_M\} \in \mathcal{P}_X / \forall m \in [M], A_m \subseteq U_m\}.$

The set \mathcal{U}^* is useful to simplify several proofs by using lemma 2 in [HMRY] that we state without proof.

Lemma 2. For any measure space (X, \mathcal{B}, μ) and $G: \mathcal{P}_X \to \mathbb{R}$ such that $\forall \alpha, \beta \in \mathcal{P}_X$, $\alpha \succeq \beta$ implies that $G(\alpha) \ge G(\beta)$ one has that for any $\mathcal{U} \in \mathcal{C}_X$:

(2)
$$\inf_{\substack{\alpha \succeq \mathcal{U} \\ \alpha \in \mathcal{P}_X}} G(\alpha) = \inf_{\alpha \in \mathcal{U}^*} G(\alpha)$$

In general one can define the μ -conditional entropy of $\alpha \in \mathcal{P}_X$ conditioned by a sub σ -algebra $\mathcal{A} \subseteq \mathcal{B}$ as:

(3)
$$H_{\mu}(\alpha|\mathcal{A}) := \sum_{A \in \alpha} \int_{X} \phi[\mathbb{E}_{\mu}(\mathbb{1}_{A}|\mathcal{A})(x)] d\mu(x).$$

Where $\phi:[0,1] \to \mathbb{R}^+$ is defined as $\phi(x) := x \log x$ for x > 0 and $\phi(0) = 0$.

Given $\alpha, \beta \in \mathcal{P}_X$ the entropy of α conditioned by β is given by $H_{\mu}(\alpha|\beta) = H_{\mu}(\alpha|\overline{\beta})$ where $\overline{\beta}$ is the σ -algebra generated by β . In this case there are simpler formulas to compute it, for instance when $\beta = \{X\}$ this gives the entropy of α and $H_{\mu}(\alpha) := -\sum_{A \in \alpha} \phi(\mu(A))$. In general for any $\beta \in \mathcal{P}_X$:

(4)
$$H_{\mu}(\alpha|\beta) := H_{\mu}(\alpha \vee \beta) - H_{\mu}(\beta) = \sum_{B \in \beta} \mu(B) H_{\mu_B}(\alpha)$$

where μ_B denotes the conditional measure induced by μ on B (zero if $\mu(B) = 0$). The function $H_{\cdot}(\alpha|\mathcal{A})$ is concave on $\mathcal{M}_T(X)$ (since $-x \log x$ is concave).

REMARK: The formula $H_{\mu}(\alpha)$ can be applied to any disjoint family of measurable sets even if it does not cover X.

Given $\alpha, \beta \in \mathcal{P}_X$ or any sub- σ -algebra $A \subseteq \mathcal{B}$ by lemma 1 the μ -entropy of α conditioned to the partition $\beta \in \mathcal{P}_X$ (or some *T*-invariant σ -algebra \mathcal{A}) with respect to *T* is well defined as:

(5)
$$h_{\mu}(\alpha|\beta,T) := \lim_{N \to \infty} \frac{1}{N} H_{\mu}(\alpha_{0}^{N-1}|\beta_{0}^{N-1}) = \inf_{N \in \mathbb{N}} \frac{1}{N} H_{\mu}(\alpha_{0}^{N-1}|\beta_{0}^{N-1}),$$
$$h_{\mu}(\alpha|\mathcal{A},T) := \lim_{N \to \infty} \frac{1}{N} H_{\mu}(\alpha_{0}^{N-1}|\mathcal{A}) = \inf_{N \in \mathbb{N}} \frac{1}{N} H_{\mu}(\alpha_{0}^{N-1}|\mathcal{A}).$$

When $\beta = \{X\}$ this yields the standard μ -entropy of α with respect to T that we denote as $h_{\mu}(\alpha, T) := h_{\mu}(\alpha | \{X\}, T)$.

For any
$$\mathcal{U} = \{U_1, \dots, U_M\}, \mathcal{V} = \{V_1, \dots, V_M\} \in \mathcal{C}_X$$
 define $\mu(\mathcal{U}\Delta\mathcal{V}) = \sum_{m=1}^M \mu(U_m\Delta V_m).$

This definition gives a notion of distance between covers and partitions that is compatible with the conditional measure entropy as the following lemma whose proof is taken from Peter Walter's classic textbook [W].

Lemma 3. Fix $M \in \mathbb{N}$ and $\epsilon > 0$ then there exists $\delta > 0$ such that $\forall \alpha, \alpha', \beta \in \mathcal{P}_X$ with $|\alpha| = |\alpha'| = M$ if $\mu(\alpha \Delta \alpha') < \delta$ one has that $|h_{\mu}(\alpha | \beta, T) - h_{\mu}(\alpha' | \beta, T)| \leq \epsilon$.

We use the same definition of chapter 6.3 in [D] but for C_X and $\beta \in \mathcal{P}_X$:

(6)
$$\mathcal{N}(\mathcal{U}|\beta) := \max_{B \in \beta} \mathcal{N}(\mathcal{U} \cap B), \ \mathcal{N}(\mathcal{U} \cap B) := \min\{|\mathcal{V}|/\mathcal{V} \subseteq \mathcal{U}, B \subseteq \bigcup_{V \in \mathcal{V}} V\}.$$

From the same ideas used in Fact 6.3.2 in [D] and other simple calculations:

Proposition 4. Let (X,T) be a TDS and $\mathcal{U}, \mathcal{V} \in \mathcal{C}_X$ and $\alpha, \beta, \gamma \in \mathcal{P}_X$ then

(1) $\mathcal{N}(\mathcal{U}|\beta) = 0$ iff $\beta \succeq \mathcal{U}$. (2) $\mathcal{N}(T^{-1}\mathcal{U}|T^{-1}\beta) = \mathcal{N}(\mathcal{U}|\beta)$. (3) $\mathcal{N}(\mathcal{U}|\beta) \ge \mathcal{N}(\mathcal{V}|\beta)$ if $\mathcal{U} \succeq \mathcal{V}$. (4) $\mathcal{N}(\mathcal{U}|\beta) \le \mathcal{N}(\mathcal{U}|\gamma)$ for $\beta \succeq \gamma$. (5) $\mathcal{N}(\mathcal{U} \lor \mathcal{V}|\beta) \le \mathcal{N}(\mathcal{U}|\beta) \cdot \mathcal{N}(\mathcal{V}|\beta)$.

Proof: Part (2) is trivial and for all the others use Fact 6.3.2. with F = X and consider \mathcal{V} and \mathcal{W} as partitions when required.

Proposition 4 parts (2),(4) and (5) imply that the sequence $\{\log \mathcal{N}(\mathcal{U}_0^{N-1}|\beta_0^{N-1})\}_{N\in\mathbb{N}}$ is subadditive, and using lemma 1 as in Definition 6.3.14 in [D] we obtain the combinatorial (topological if \mathcal{U} is open) entropy of the cover $\mathcal{U} \in \mathcal{C}_X$ conditioned by partition $\beta \in \mathcal{P}_X$ with respect to T as:

(7)
$$h(\mathcal{U}|\beta,T) := \lim_{N \to \infty} \frac{1}{N} \log \mathcal{N}(\mathcal{U}_0^{N-1}|\beta_0^{N-1}) = \inf_{N \in \mathbb{N}} \frac{1}{N} \log \mathcal{N}(\mathcal{U}_0^{N-1}|\beta_0^{N-1}).$$

When $\beta = \{X\}$ we recover the classical topological entropy and denote $\mathcal{N}(\mathcal{U}|\{X\}) = \mathcal{N}(\mathcal{U})$ and $h(\mathcal{U}|\{X\}, T) = h(\mathcal{U}, T)$.

In [R] we defined for $\mathcal{U} \in \mathcal{C}_X^o$ and $\mu \in \mathcal{M}_T(X)$, $H_\mu(\mathcal{U}) := \inf_{\substack{\alpha \succeq \mathcal{U} \\ \alpha \in \mathcal{P}_X}} H_\mu(\alpha)$. To extend this idea we define for any $\mathcal{U} \in \mathcal{C}_X$ and $\beta \in \mathcal{P}_X$ given $B \in \beta$, $H_{\mu_B}(\mathcal{U})$ as for $H_\mu(\mathcal{U})$ (even if μ_B is not *T*-invariant) and using the decomposition shown in (4):

(8)
$$H_{\mu}(\mathcal{U}|\beta) := \sum_{B \in \beta} \mu(B) H_{\mu_B}(\mathcal{U}), \ H^*_{\mu}(\mathcal{U}|\beta) := \inf_{\alpha \succeq \mathcal{U}} H_{\mu}(\alpha|\beta).$$

It is straightforward that $H_{\mu}(\mathcal{U}|\beta) \leq H_{\mu}^*(\mathcal{U}|\beta)$ and when $\beta = \{X\}$ and $\mathcal{U} \in \mathcal{C}_X^\circ$ we recover in both cases the definition given in [R] since $H_{\mu}^*(\mathcal{U}|\{X\}) = H_{\mu}(\mathcal{U}|\{X\}) =$ $H_{\mu}(\mathcal{U})$. Nevertheless we will prove that they are actually always equal. This result and some basic properties for the conditional entropy will extend to $H_{\mu}(\mathcal{U}|\beta)$ with respect to any fixed $\beta \in C_X$ with the use of two key facts. From [R] as mentioned in Fact 8.3.5 in [D],

(9)
$$H_{\mu}(\mathcal{U}) = \min_{\alpha \in Ext(\mathcal{U})} H_{\mu_B}(\alpha).$$

With $Ext(\mathcal{U})$ the set of partitions of the form $\{U_1, U_2 \setminus U_1, \ldots, U_d \setminus (U_1 \cup \cdots \cup U_{d-1})\}$ where $\{U_1, \ldots, U_d\}$ is an ordering of \mathcal{U} . This is also true for any $B \in \beta$ for $H_{\mu_B}(\mathcal{U})$ since the proof does not need μ to be *T*-invariant.

When A and B are disjoint then:

(10)
$$\mu(A \cup B)H_{\mu_{A \cup B}}(\alpha) \ge \mu(A)H_{\mu_{A}}(\alpha) + \mu(B)H_{\mu_{B}}(\alpha) \ge \mu(A)H_{\mu_{A}}(\alpha).$$

Thus if $A \subseteq B$ then $\mu(A)H_{\mu_{A}}(\mathcal{U}) \le \mu(B)H_{\mu_{B}}(\mathcal{U}).$

Lemma 5. For any $\mathcal{U} \in \mathcal{C}_X$, $\beta \in \mathcal{P}_X$, $H_{\mu}(\mathcal{U}|\beta) = H^*_{\mu}(\mathcal{U}|\beta)$.

Proof: Fix $\mathcal{U} \in \mathcal{C}_X$ and $\beta \in \mathcal{P}_X$, we only need to prove that $H^*_{\mu}(\mathcal{U}|\beta) \leq H_{\mu}(\mathcal{U}|\beta)$. From fact 9 for any $B \in \beta$ there exists $\alpha_B \in \mathcal{P}_X$ such that $H_{\mu_B}(\mathcal{U}) = H_{\mu_B}(\alpha_B)$ so $H^*_{\mu}(\mathcal{U}|\beta) = \sum_{B \in \beta} \mu(B) H_{\mu_B}(\alpha_B)$.

Define $\alpha = \bigvee_{B \in \beta} [\alpha_B \cap B] \cup \{B^c\}$. By definition for any any $\tilde{\alpha} \succeq \mathcal{U}$ and $B \in \beta$, $H_{\mu_B}(\tilde{\alpha}) \geq H_{\mu_B}(\alpha_B) = H_{\mu_B}([\alpha_B \cap B] \cup \{B^c\}) = H_{\mu_B}(\alpha)$ so multiplying by $\mu(B)$ and adding up over $B \in \beta$ we conclude that $H_{\mu}(\mathcal{U}|\beta) = H_{\mu}(\alpha|\beta) = H_{\mu}^*(\mathcal{U}|\beta)$.

REMARK: From now we will use as definition of $H_{\mu}(\mathcal{U}|\beta)$ the most suitable of these two formulas as needed in the proofs.

Now we prove that this notion has the standard properties of a static entropy.

Proposition 6. Let (X, \mathcal{B}, μ, T) be a MDS and $\mathcal{U}, \mathcal{V}, \mathcal{W} \in \mathcal{C}_X$ and $\alpha, \beta, \gamma \in \mathcal{P}_X$ then

(1) $0 \leq H_{\mu}(\mathcal{U}|\beta) \leq \log \mathcal{N}(\mathcal{U}|\beta) \text{ and } H_{\mu}(\mathcal{U}|\beta) = 0 \text{ if } \beta \succeq \mathcal{U}.$ (2) $H_{\mu}(T^{-1}\mathcal{U}|T^{-1}\beta) = H_{\mu}(\mathcal{U}|\beta).$ (3) $H_{\mu}(\mathcal{U}|\beta) \leq H_{\mu}(\mathcal{U}|\gamma) \text{ for } \beta \succeq \gamma.$ (4) $H_{\mu}(\mathcal{U} \lor \mathcal{V}|\beta) \leq H_{\mu}(\mathcal{U}|\beta) + H_{\mu}(\mathcal{V}|\beta).$

Proof: Fix $\mathcal{U}, \mathcal{V}, \mathcal{W} \in \mathcal{C}_X$ and $\alpha, \beta, \gamma \in \mathcal{P}_X$.

(1) For any $B \in \beta$, $H_{\mu_B}(\alpha) \leq \log |\alpha \cap B|$ and $\inf_{\alpha \succeq \mathcal{U}} |\alpha \cap B| = \mathcal{N}(\mathcal{U} \cap B)$. So:

(11)
$$H_{\mu}(\mathcal{U}|\beta) = \sum_{B \in \beta} \mu(B) H_{\mu_B}(\mathcal{U}) \le \sum_{B \in \beta} \mu(B) \log \mathcal{N}(\mathcal{U} \cap B) \le \log \mathcal{N}(\mathcal{U}|\beta).$$

If $\beta \succeq \mathcal{U}, H_{\mu_B}(\mathcal{U}) \le H_{\mu_B}(\beta) = 0$ for any $B \in \beta$.

(2) For any $B \in \beta$ from equation (9):

$$H_{\mu_{T^{-1}B}}(T^{-1}\mathcal{U}) = \min_{\alpha \in Ext(T^{-1}\mathcal{U})} H_{\mu_{T^{-1}B}}(\alpha) = \min_{\alpha \in T^{-1}Ext(\mathcal{U})} H_{\mu_{T^{-1}B}}(\alpha),$$

= $\min_{\alpha \in Ext(\mathcal{U})} H_{\mu_{T^{-1}B}}(T^{-1}\alpha) = \min_{\alpha \in Ext(\mathcal{U})} H_{\mu_B}(\alpha) = H_{\mu_B}(\mathcal{U}).$

Adding up over all $B \in \beta$ and since $\mu(T^{-1}B) = \mu(B)$ we conclude the result.

(3) Let $\beta = \{B_1, \ldots, B_N\}$ and $\gamma = \{C_1, \ldots, C_M\}$ if $\beta \succeq \gamma$ then there exists a partition $\{K_n\}_{n=1}^N$ of $\{1, \ldots, N\}$ such that $B_n = \bigcup_{k \in K_n} C_k$ for any $1 \le n \le N$.

From equation (10) for any $1 \le n \le N$:

(12)
$$\mu(B_n)H_{\mu_{B_n}}(\mathcal{U}) \ge \sum_{k \in K_n} \mu(C_n)H_{\mu_{C_n}}(\mathcal{U}).$$

Adding up over n concludes the result.

(4) For any $\alpha \succeq \mathcal{U}$ and $\gamma \succeq \mathcal{V}$ from basic properties of H_{μ} for partitions:

(13)
$$H_{\mu}(\mathcal{U} \vee \mathcal{V}|\beta) \le H_{\mu}(\alpha \vee \gamma|\beta) \le H_{\mu}(\alpha|\beta) + H_{\mu}(\gamma|\beta).$$

Taking infima over all $\alpha \succeq \mathcal{U}$ and $\gamma \succeq \mathcal{V}$ concludes the result.

This allows us to define two measure theoretical notions of dynamical entropy for a measurable cover conditioned to a fixed partition.

Proposition 6 parts (2),(3) and (4) imply that the sequence $\{H_{\mu}(\mathcal{U}_{0}^{N-1}|\beta_{0}^{N-1})\}_{N\in\mathbb{N}}$ is subadditive, and so by lemma 1 we define two notions of μ -entropy of the cover $\mathcal{U} \in \mathcal{C}_{X}^{\circ}$ conditioned by the partition $\beta \in \mathcal{P}_{X}$ with respect to T as:

(14)
$$h_{\mu}^{-}(\mathcal{U}|\beta,T) := \lim_{N \to \infty} \frac{1}{N} H_{\mu}(\mathcal{U}_{0}^{N-1}|\beta_{0}^{N-1}) = \inf_{N \in \mathbb{N}} \frac{1}{N} H_{\mu}(\mathcal{U}_{0}^{N-1}|\beta_{0}^{N-1}),$$
$$h_{\mu}^{+}(\mathcal{U}|\beta,T) := \inf_{\substack{\alpha \geq \mathcal{U} \\ \alpha \in \mathcal{P}_{X}}} h_{\mu}(\alpha|\beta,T).$$

Notice that when $\beta = \{X\}$ we recover the notions in [R].

The following lemma is extremely useful to extend properties between h^+_{μ} and h^-_{μ} and finally prove they coincide and it will be used many times in the sequel.

Lemma 7. Let (X, \mathcal{B}, μ, T) a MDS and $\mathcal{U} \in \mathcal{C}_X$ and $\beta \in \mathcal{P}_X$. Then:

 $\begin{array}{ll} (1) & h_{\mu}^{-}(\mathcal{U}|\beta,T) \leq h_{\mu}^{+}(\mathcal{U}|\beta,T) \ and \ h_{\mu}^{-}(\mathcal{U}|\beta,T) \leq h(\mathcal{U}|\beta,T). \\ (2) & \forall M \in \mathbb{N}, \ h_{\mu}^{-}(\mathcal{U}|\beta,T) = \frac{1}{M}h_{\mu}^{-}(\mathcal{U}_{0}^{M-1}|\beta_{0}^{M-1},T^{M}). \\ (3) & h_{\mu}^{-}(\mathcal{U}|\beta,T) = \lim_{M \to \infty} \frac{1}{M}h_{\mu}^{+}(\mathcal{U}_{0}^{M-1}|\beta_{0}^{M-1},T^{M}). \end{array}$

Proof: Fix $\mathcal{U} \in \mathcal{C}_X$ and $\beta \in \mathcal{P}_X$.

(1) By Proposition 6 part (1), $H_{\mu}(\mathcal{U}|\beta) \leq \log \mathcal{N}(\mathcal{U}|\beta)$ so:

$$h_{\mu}^{-}(\mathcal{U}|\beta,T) = \lim_{N \to \infty} \frac{1}{N} H_{\mu}(\mathcal{U}_{0}^{N-1}|\beta_{0}^{N-1}),$$

$$\leq \lim_{N \to \infty} \frac{1}{N} \log \mathcal{N}(\mathcal{U}_{0}^{N-1}|\beta_{0}^{N-1}) = h(\mathcal{U}|\beta,T).$$

For any $N \in \mathbb{N}$, $H_{\mu}(\mathcal{U}_0^{N-1}|\beta_0^{N-1}) \leq H_{\mu}^*(\mathcal{U}_0^{N-1}|\beta_0^{N-1}) \leq \inf_{\alpha \succeq \mathcal{U}} H_{\mu}(\alpha_0^{N-1}|\beta_0^{N-1}).$ Dividing by N and taking limit proves that $h_{\mu}^-(\mathcal{U}|\beta, T) \leq h_{\mu}^+(\mathcal{U}|\beta, T).$

(2) For every $M \in \mathbb{N}$ just by definition:

$$h_{\mu}^{-}(\mathcal{U}|\beta, T) = \lim_{N \to \infty} \frac{1}{NM} H_{\mu}(\mathcal{U}_{0}^{NM-1}|\beta_{0}^{NM-1}),$$

$$= \frac{1}{M} \lim_{N \to \infty} \frac{1}{N} H_{\mu} \left(\bigvee_{n=0}^{N-1} T^{-nM} \mathcal{U}_{0}^{M-1} | \bigvee_{n=0}^{N-1} T^{-nM} \beta_{0}^{M-1} \right),$$

$$= \frac{1}{M} h_{\mu}^{-}(\mathcal{U}_{0}^{M-1}|\beta_{0}^{M-1}, T^{M}).$$

(3) From parts (1) and (2) for every $M \in \mathbb{N}$:

(15)
$$h_{\mu}^{-}(\mathcal{U}|\beta,T) = \frac{1}{M}h_{\mu}^{-}(\mathcal{U}_{0}^{M-1}|\beta_{0}^{M-1},T^{M}) \leq \frac{1}{M}h_{\mu}^{+}(\mathcal{U}_{0}^{M-1}|\beta_{0}^{M-1},T^{M}).$$

Also for every $N \in \mathbb{N}$ using Proposition 6 parts (3) and (4):

$$\frac{1}{M}h_{\mu}^{+}(\mathcal{U}_{0}^{M-1}|\beta_{0}^{M-1},T^{M}) = \lim_{N \to \infty} \frac{1}{NM} \inf_{\alpha \succeq \mathcal{U}_{0}^{M-1}} H_{\mu}\left(\bigvee_{n=0}^{N-1} T^{-nM}\alpha|\beta_{0}^{NM-1}\right), \\
\leq \lim_{N \to \infty} \frac{1}{NM} \inf_{\alpha \succeq \mathcal{U}_{0}^{M-1}} \sum_{n=0}^{N-1} H_{\mu}\left(T^{-nM}\alpha|T^{-nM}\beta_{0}^{M-1}\right), \\
\leq \frac{1}{M} \inf_{\alpha \succeq \mathcal{U}_{0}^{M-1}} H_{\mu}\left(\alpha|\beta_{0}^{M-1}\right) = \frac{1}{M}H_{\mu}\left(\mathcal{U}_{0}^{M-1}|\beta_{0}^{M-1}\right).$$

Taking limit as M goes to infinity concludes the proof.

Proposition 8. Let (X,T) and (Y,S) be TDS, $\mu \in \mathcal{M}_T(X)$ and $\nu \in \mathcal{M}_S(Y)$.Let $\varphi: (X,T,\mu) \to (Y,S,\nu)$ be a measure-theoretical factor map, $\mathcal{U} \in \mathcal{C}_Y(S)$ and $\beta \in \mathcal{P}_Y$. Then $h_{\nu}^{-}(\mathcal{U}|\beta,S) = h_{\mu}^{-}(\varphi^{-1}\mathcal{U}|\varphi^{-1}\beta,T)$ and $h(\mathcal{U}|\beta,S) = h(\varphi^{-1}\mathcal{U}|\varphi^{-1}\beta,T)$.

Proof: Fix $\mathcal{U} \in \mathcal{C}_Y(S)$ and $\beta \in \mathcal{P}_Y$. Clearly for any $N \in \mathbb{N}$, $\mathcal{N}(\varphi^{-1}\mathcal{U}_0^{N-1}|\varphi^{-1}\beta_0^{N-1}) = \mathcal{N}(\mathcal{U}_0^{N-1}|\beta_0^{N-1})$ so taking log diving by N and taking limit as N tends to infinity proves that $h(\mathcal{U}|\beta, S) = h(\varphi^{-1}\mathcal{U}|\varphi^{-1}\beta, T)$.

In Proposition 6 in [R] it is proven that $H_{\mu}(\varphi^{-1}\mathcal{U}) = H_{\nu}(\mathcal{U})$ and that part of the proof does not require for μ to be *T*-invariant only that $\mu \circ \varphi^{-1} = \nu$. Since for any measurable set *B* it is straightforward to see that $\mu_{\varphi^{-1}B} \circ \varphi^{-1} = \nu_B$ thus the same proof applies to prove that $H_{\mu_{\varphi^{-1}B}}(\varphi^{-1}\mathcal{U}) = H_{\nu_B}(\mathcal{U})$. Taking the sum over all $B \in \beta$ we conclude that:

(16)
$$H_{\mu}(\varphi^{-1}\mathcal{U}|\varphi^{-1}\beta) = H_{\nu}(\mathcal{U}|\beta).$$

Using equation (16) since $S \circ \varphi = \varphi \circ T$ for any $N \in \mathbb{N}$ we have that:

(17)
$$H_{\mu}\left(\bigvee_{n=0}^{N-1}T^{-n}(\varphi^{-1}\mathcal{U})|\bigvee_{n=0}^{N-1}T^{-n}\varphi^{-1}\beta\right) = H_{\nu}\left(\bigvee_{n=0}^{N-1}S^{-n}\mathcal{U}|\bigvee_{n=0}^{N-1}S^{-n}\beta\right).$$

Dividing and taking the limit when N goes to infinity concludes the proof.

3. Local Variational Principles

In this section we prove the local variational principle for both notions. The techniques and proofs are a mix of the work done in [R], [GW], [HMRY] and [HYZ].

Lemma 9. Let (X,T) a TDS, $\mathcal{U} \in \mathcal{C}_X$ and $\beta \in \mathcal{P}_X$. For every family $\{\alpha_l\}_{l=1}^K$ of finite partitions finer than \mathcal{U} , for every $N \in \mathbb{N}$ choose $B \in \beta_0^{N-1}$ such that the maximum of $\mathcal{N}(\mathcal{U}_0^{N-1}|\beta_0^{N-1})$ is attained.

There exists a finite subset $P \subseteq B$ such that every element of $(\alpha_l)_0^{N-1}$ contains at most one point of P for every $1 \le l \le K$ and $|P| \ge \frac{\mathcal{N}(\mathcal{U}_0^{N-1}|\beta_0^{N-1})}{K}$.

Proof: Choose $\mathcal{U} \in \mathcal{C}_X$, $\beta \in \mathcal{P}_X$, $N \in \mathbb{N}$ and $B \in \beta_0^{N-1}$ as stated.

For $x \in X$ and $1 \leq l \leq K$, let $A_l(x)$ be the element in $(\alpha_l)_0^{N-1}$ that contains x. Take $x_1 \in B$. If $B \subseteq \bigcup_{l=1}^K A_l(x_1)$ then let $P = \{x_1\}$. Every element in $(\alpha_l)_0^{N-1}$

contains at most one point of P for every $1 \le l \le K$ and $|P|K \ge \mathcal{N}(\mathcal{U}_0^{N-1}|\beta_0^{N-1})$. If $B \nsubseteq \bigcup_{l=1}^K A_l(x_1)$ let $B_1 = B \setminus \bigcup_{l=1}^K A_l(x_1)$ and take $x_2 \in B_1$. If $B_1 \subseteq \bigcup_{l=1}^K A_l(x_2)$ then $P = \{x_1, x_2\}$. Every element in $(\alpha_l)_0^{N-1}$ contains at most one point of P for every $1 \le l \le K$ and $|P_N|K \ge \mathcal{N}(\mathcal{U}_0^{N-1}|\beta_0^{N-1})$.

Otherwise, $B_2 = B_1 \setminus \bigcup_{l=1}^K A_l(x_2)$. Since this is a finite procedure we obtain a

set $\{x_1, \ldots, x_m\}$ such that $B \subseteq \bigcup_{j=1}^m \bigcup_{l=1}^K A_l(x_j)$. Let $P = \{x_1, \ldots, x_m\}$. By construction $(\alpha_l)_0^{N-1}$ contains at most one point of P for every $1 \leq l \leq K$ and $|P|K = |\bigcup_{j=1}^m \bigcup_{l=1}^K A_l(x_j)| \geq \mathcal{N}(\mathcal{U}_0^{N-1}|\beta_0^{N-1})$.

Theorem 10. For every TDS(X,T), $\mathcal{U} \in \mathcal{C}_X^o$ and $\beta \in \mathcal{P}_X$ there exists $\mu \in \mathcal{M}_T(X)$ such that $h^+_{\mu}(\mathcal{U}|\beta,T) \ge h(\mathcal{U}|\beta,T)$ and $h^-_{\mu}(\mathcal{U}|\beta,T) = h(\mathcal{U}|\beta,T)$.

Proof: First we prove in the 0-dimensional case that for every $\mathcal{U} = \{U_1, \ldots, U_d\} \in$ \mathcal{C}_X and $\beta \in \mathcal{P}_\mu$ there exists $\mu \in \mathcal{M}_T(X)$ such that $h^+_\mu(\mathcal{U}|\beta, T) \ge h(\mathcal{U}|\beta, T)$.

We will then use the now classic technique used in [BGH], that works only for $h_{\mu}^{-}(\mathcal{U}|\beta,T)$ in our case. This is we prove the result first in the zero-dimensional case and then extend it to the general case.

The set of clopen sets of a zero dimensional set X is a countable set, thus the family of partitions in \mathcal{U}^* consisting of clopen sets is countable and lets enumerate it as $\{\alpha_l / l \in \mathbb{N}\}$.

Now fix $n \in \mathbb{N}$ and use lemma (9) with $N = n^2$, K = n and the family $\{T^{-n}\alpha_l\}_{l=1}^n$ applied to the cover $T^{-n}\mathcal{U}$ and the partition $T^{-n}\beta$. We obtain $C_n \in \beta_n^{n^2+n-1}$ and $P_n \subseteq C_n$ such that $|P_n| \geq \frac{\mathcal{N}(\mathcal{U}_n^{n^2+n-1}|\beta_n^{n^2+n-1})}{n}$ and every element of $(\alpha_l)_n^{n^2+n-1}$ contains at most one point of P_n for every $1 \leq l \leq n$. To simplify the size of the formulas denote $N = n^2 + n$.

Denote as ν_n the counting measure over P_n and choose $1 \le i, l \le n$. By definition and basic properties:

(18)

$$H_{T^{i}\nu_{n}}((\alpha_{l})_{0}^{N-1}|\beta_{0}^{N-1}) \geq H_{T^{i}\nu_{n}}((\alpha_{l})_{0}^{N-1}|\beta_{n}^{N-1}) - H_{T^{i}\nu_{n}}(\beta_{0}^{n-1}),$$

$$= H_{T^{i}\nu_{n}}((\alpha_{l})_{0}^{N-1}) - H_{T^{i}\nu_{n}}(\beta_{0}^{n-1}),$$

$$\geq H_{T^{i}\nu_{n}}((\alpha_{l})_{0}^{N-1}) - \log |\beta_{0}^{n-1}|.$$

From proposition 4 parts (2), (4) and (5):

(19)
$$\mathcal{N}(\mathcal{U}_{0}^{N-1}|\beta_{0}^{N-1}) \leq \mathcal{N}(\mathcal{U}_{0}^{n-1}|\beta_{0}^{N-1}) \cdot \mathcal{N}(\mathcal{U}_{n}^{N-1}|\beta_{0}^{N-1}),$$
$$\leq \mathcal{N}(\mathcal{U}_{0}^{n-1}|\beta_{0}^{n-1}) \cdot \mathcal{N}(\mathcal{U}_{n}^{N-1}|\beta_{n}^{N-1}),$$
$$\leq d^{n} \cdot \mathcal{N}(\mathcal{U}_{n}^{N-1}|\beta_{n}^{N-1}).$$

Since $T^{-i}(\alpha_l)_0^{N-1} \succeq (\alpha_l)_n^{N-1}$ then every element contains at most one atom of the discrete measure ν_n and so using equations (18) and (19):

(20)
$$H_{T^{i}\nu_{n}}((\alpha_{l})_{0}^{N-1}|\beta_{0}^{N-1}) \geq H_{\nu_{n}}(T^{-i}(\alpha_{l})_{0}^{N-1}) - \log|\beta_{0}^{n-1}|,$$
$$\geq \log\left[\frac{\mathcal{N}(\mathcal{U}_{n}^{N-1}|\beta_{n}^{N-1})}{n}\right] - \log|\beta_{0}^{n-1}|,$$
$$\geq \log\left[\frac{\mathcal{N}(\mathcal{U}_{0}^{N-1}|\beta_{0}^{N-1})}{nd^{n}}\right] - \log|\beta_{0}^{n-1}|,$$

Fix $m \in \mathbb{N}$ with $m \leq n$ and decompose N = km + b with $0 \leq b \leq m - 1$. Then:

$$\begin{aligned} H_{T^{i}\nu_{n}}((\alpha_{l})_{0}^{N-1}|\beta_{0}^{N-1}) &= H_{T^{i}\nu_{n}}\left(\bigvee_{j=0}^{k-1}T^{-mj}(\alpha_{l})_{0}^{m-1}\vee(\alpha_{l})_{km}^{N-1}|\beta_{0}^{N-1}\right),\\ &\leq \sum_{j=0}^{k-1}H_{T^{i}\nu_{n}}(T^{-mj}(\alpha_{l})_{0}^{m-1}|\beta_{0}^{N-1}) + H_{T^{i}\nu_{n}}((\alpha_{l})_{km}^{N-1}),\\ &\leq \sum_{j=0}^{k-1}H_{T^{i}\nu_{n}}(T^{-mj}(\alpha_{l})_{0}^{m-1}|T^{-mj}\beta_{0}^{m-1}) + \log|(\alpha_{l})_{km}^{N-1}|,\\ &\leq \sum_{j=0}^{k-1}H_{T^{i+mj}\nu_{n}}((\alpha_{l})_{0}^{m-1}|\beta_{0}^{m-1}) + m\log d.\end{aligned}$$

Adding up for every $0 \le i \le m - 1$ from equation (20) we obtain:

(21)

$$m \log\left[\frac{\mathcal{N}(\mathcal{U}_{0}^{N-1}|\beta_{0}^{N-1})}{nd^{n}}\right] - m \log|\beta_{0}^{n-1}| \leq \sum_{i=0}^{m-1} H_{T^{i}\nu_{n}}((\alpha_{l})_{0}^{N-1}|\beta_{0}^{N-1}),$$

$$\leq \sum_{i=0}^{m-1} \sum_{j=0}^{k-1} H_{T^{i}\nu_{n}}((\alpha_{l})_{0}^{m-1}|\beta_{0}^{m-1}) + m^{2}\log d.$$

$$\leq \sum_{i=0}^{N-1} H_{T^{i}\nu_{n}}((\alpha_{l})_{0}^{m-1}|\beta_{0}^{m-1}) + m^{2}\log d.$$

Denote as μ_n the Cesaro Mean Measure of $\{T^i\nu_n\}_{i=0}^{N-1}$. By concavity and using equation (21) we get for any $0 \le l \le n$:

(22)
$$H_{\mu_n}((\alpha_l)_0^{m-1}|\beta_0^{m-1}) \ge \frac{1}{N} \sum_{i=0}^{N-1} H_{T^i\nu_n}((\alpha_l)_0^{m-1}|\beta_0^{m-1}),$$
$$\ge \frac{m}{N} \left(\log \left[\frac{\mathcal{N}(\mathcal{U}_0^{N-1}|\beta_0^{N-1})}{nd^n} \right] - m \log d - \log |\beta_0^{n-1}| \right).$$

We can assume by taking a subsequence that there exists $\mu \in \mathcal{M}(X)$ such that in the weak-* topology $\lim_{n \to \infty} \mu_n = \mu$. Also by construction $\mu \in \mathcal{M}_T(X)$. Clearly:

(23)
$$h(\mathcal{U}|\beta, T) = \lim_{n \to \infty} \frac{m}{N} \left(\log \left[\frac{\mathcal{N}(\mathcal{U}_0^{N-1}|\beta_0^{N-1})}{nd^n} \right] - m \log d - \log |\beta_0^{n-1}| \right).$$

Since all elements of $(\alpha_l)_0^{m-1}$ are clopen by upper semicontinuity and equation (22) divided by m we obtain for any $l, m \in \mathbb{N}$:

(24)
$$\frac{1}{m}H_{\mu}((\alpha_{l})_{0}^{m-1}|\beta_{0}^{m-1}) \geq \limsup_{n \to \infty} \frac{1}{m}H_{\mu_{n}}((\alpha_{l})_{0}^{m-1}|\beta_{0}^{m-1}) = h(\mathcal{U}|\beta, T).$$

Taking limit as m tends to infinity proves that $h_{\mu}(\alpha_{l}|\beta, T) \geq h(\mathcal{U}|\beta, T)$. Since $\{\alpha_{l}\}_{l\in\mathbb{N}}$ is dense in L^{1}_{μ} in the set of borel partitions finer than \mathcal{U} this proves that $h^{+}_{\mu}(\mathcal{U}|\beta, T) \geq h(\mathcal{U}|\beta, T)$. By lemma 7 part (3) this proves that $h^{-}_{\mu}(\mathcal{U}|\beta, T) \geq h(\mathcal{U}|\beta, T)$ but since for any $\mu \in \mathcal{M}_{T}(X), h^{-}_{\mu}(\mathcal{U}|\beta, T) \leq h(\mathcal{U}|\beta, T)$ we conclude that $h^{-}_{\mu}(\mathcal{U}|\beta, T) = h(\mathcal{U}|\beta, T)$.

As in [BGH] we use the existence of a zero dimensional extension (Y, S) of (X, T), this is a continuous surjective map $\varphi: Y \to X$ such that $\varphi \circ S = T \circ \varphi$ where (Y, S)is a zero-dimensional TDS.

Now choose $\mathcal{U} \in \mathcal{C}_X$, $\beta \in \mathcal{P}_X$ and apply the previous case for (Y, S), the cover $\varphi^{-1}\mathcal{U}$ and the partition $\varphi^{-1}\beta$ to obtain $\nu \in \mathcal{M}_S(Y)$ such that $h_{\nu}^{-}(\varphi^{-1}\mathcal{U}|\varphi^{-1}\beta, S) = h(\varphi^{-1}\mathcal{U}|\varphi^{-1}\beta, S)$.

Define $\mu = \nu \circ \varphi$ from proposition 8 we conclude that:

PIERRE-PAUL ROMAGNOLI

(25)
$$h_{\mu}^{-}(\mathcal{U}|\beta, S) = h_{\nu}^{-}(\varphi^{-1}\mathcal{U}|\varphi^{-1}\beta, S) = h(\varphi^{-1}\mathcal{U}|\varphi^{-1}\beta, S) = h(\mathcal{U}|\beta, T)$$

In order to prove the inequality for h^+_{μ} just notice that it is always true that $h^+_{\mu}(\mathcal{U}|\beta, T) \ge h^-_{\mu}(\mathcal{U}|\beta, T)$.

4. Ergodic Decomposition and the equality of + and -

There are several approaches to define the ergodic decomposition of the measure theoretical entropy and we choose the simple and clear approach taken in [D]. The idea is to use the notion of disintegration of the measure μ with respect to a borel sub sigma-algebra $\mathcal{D} \subseteq \mathcal{B}_X$. There is a one to one correspondence between a sub sigma-algebra \mathcal{D} and a measure theoretical φ -factor (Y, \mathcal{D}', ν) where the elements $y \in Y$ are the atoms of \mathcal{D} and the map $\varphi : X \to Y$ is defined by inclusion with $\varphi^{-1}\mathcal{D}' = \mathcal{D}$ and $\nu = \mu \circ \varphi^{-1}$.

For any $B \in \mathcal{B}$, the function $\mathbb{E}_{\mu}(\mathbb{1}_{B}|\mathcal{D}): X \to \mathbb{R}$ for ν -almost every $y \in Y$, is μ -almost surely constant in $\varphi^{-1}y \in Y$ so we can consider that $\mathbb{E}_{\mu}(\mathbb{1}_{B}|\mathcal{D})$ is defined over Y. Using this we define the disintegration of μ as the ν -almost everywhere defined assignment $y \to \mu_{y}(\cdot) := \mathbb{E}_{\mu}(\mathbb{1}_{(\cdot)}|\mathcal{D})(y) \in \mathcal{M}(X)$ where μ_{y} is a probability measure on \mathcal{B} supported by the atom y. By simple properties of the conditional expectation:

(26)
$$\forall B \in \mathcal{B}, \mu(B) = \mathbb{E}_{\nu}(\mathbb{E}_{\mu}(\mathbb{1}_{B}|\mathcal{D})) = \int_{Y} \mu_{y}(B) \, d\nu(y) = \mathbb{E}_{\nu}(\mathbb{E}_{\mu}(\mathbb{1}_{B}|\mathcal{D})).$$

In our case we consider $\mathcal{D} = \{B \in \mathcal{B}/T^{-1}B = B\}$. For the case of any *T*-invariant σ -algebra we can also define a φ -factor (Y, \mathcal{D}, ν, S) such that for any $y \in Y$, $\mu_{Sy} = \mu_y \circ T^{-1}$. In our choice of \mathcal{D} we also have that for any $y \in Y$, $\mu_y \in \mathcal{M}_T^e(X)$.

From Theorem 2.6.4 in [D] for any $\alpha \in \mathcal{P}_X$:

(27)
$$h_{\mu}(\alpha,T) = \int_{Y} h_{\mu_{y}}(\alpha,T) \, d\nu(y).$$

For any $\beta \in \mathcal{P}_X$ since $h_{\mu}(\alpha|\beta,T) = h_{\mu}(\alpha \vee \beta,T) - h_{\mu}(\beta,T)$ we also have that:

(28)
$$h_{\mu}(\alpha|\beta,T) = \int_{Y} h_{\mu_{y}}(\alpha|\beta,T) \, d\nu(y).$$

Using the same ideas given in [HMRY] we prove that:

Lemma 11. For any $\mathcal{U} \in \mathcal{C}_X$, $\beta \in \mathcal{P}_X$

(29)
$$h^+_{\mu}(\mathcal{U}|\beta, T) = \int_{Y} h^+_{\mu_y}(\mathcal{U}|\beta, T) \, d\nu(y), \ h^-_{\mu}(\mathcal{U}|\beta, T) = \int_{Y} h^-_{\mu_y}(\mathcal{U}|\beta, T) \, d\nu(y).$$

Proof: Let $\mathcal{U} = \{U_1, \ldots, U_M\} \in \mathcal{C}_X$ and $\beta \in \mathcal{P}_X$. Since X is a compact metric space there exists a sequence $\{\alpha_k\}_{k\in\mathbb{N}}$ in \mathcal{U}^* that is L^1_{ν} dense in \mathcal{U}^* for every $\eta \in \mathcal{M}(X)$.

So in particular we have that for any $\eta \in \mathcal{M}_T(X)$:

(30)
$$h_{\eta}^{+}(\mathcal{U}|\beta,T) = \inf_{k \in \mathbb{N}} h_{\eta}(\alpha_{k}|\beta,T).$$

Denote for any $k \in \mathbb{N}$, $\alpha_k = \{A_1^k, \ldots, A_M^k\}$ where $A_m^k \subseteq U_m$ for any $1 \le m \le M$. By equations (28) and (30) and Fatou's Lemma we have that:

(31)
$$h_{\mu}^{+}(\mathcal{U}|\beta,T) = \inf_{k \in \mathbb{N}} h_{\mu}(\alpha_{k}|\beta,T) = \inf_{k \in \mathbb{N}} \int_{Y} h_{\mu_{y}}(\alpha|\beta,T) \, d\nu(y),$$
$$\geq \int_{Y} \inf_{k \in \mathbb{N}} h_{\mu_{y}}(\alpha|\beta,T) \, d\nu(y) = \int_{Y} h_{\mu_{y}}^{+}(\mathcal{U}|\beta,T) \, d\nu(y).$$

For every $\epsilon > 0$ and $n \in \mathbb{N}$ define $B_n^{\epsilon} = \{y \in Y/h_{\mu_y}(\alpha|\beta, T) < h_{\mu_y}^+(\mathcal{U}|\beta, T) + \epsilon\}$. By equation (30) we know that $\mu(Y\Delta \bigcup_{n\in\mathbb{N}} B_n^{\epsilon}) = 0$ and so there exists $\{Y_n\}_{n\in\mathbb{N}}$ a ν -partition of X, $\mu(Y_n) > 0$ and a subsequence of partitions $\{\alpha_{k_n}\}_{k\in\mathbb{N}}$ such that for any $n \in \mathbb{N}$ and ν -almost every $y \in Y_n$, $h_{\mu_y}(\alpha_{k_n}|\beta, T) < h_{\mu_y}^+(\mathcal{U}|\beta, T) + \epsilon$.

Now define for any $n \in \mathbb{N}$ a measure $\mu_n \in \mathcal{M}_T(X)$ as:

(32)
$$\forall B \in \mathcal{B}, \mu_n(B) = \frac{1}{\mu(Y_n)} \int\limits_Y \mu_y(B \cap Y_n) \, d\nu(y).$$

By definition:

(33)
$$h_{\mu_n}(\alpha_{k_n}|\beta, T) = \frac{1}{\mu(Y_n)} \int_Y h_{\mu_y}(\alpha_{k_n}|\beta, T) \, d\nu(y),$$
$$\leq \frac{1}{\mu(Y_n)} \int_Y h_{\mu_y}^+(\mathcal{U}|\beta, T) \, d\nu(y) + \epsilon$$

Notice that for any $n, m \in \mathbb{N}$, $\mu_n(Y_m) = \mathbb{1}_{\{n=m\}}$. For any $1 \leq m \leq M$ define $A_m = \bigcup_{n \in \mathbb{N}} (Y_n \cap A_m^{k_n})$ and $\alpha = \{A_1, \ldots, A_M\} \in \mathcal{U}^*$. By construction:

(34)
$$h_{\mu}^{+}(\mathcal{U}|\beta,T) \leq h_{\mu}(\alpha|\beta,T) = \sum_{n \in \mathbb{N}} \mu(Y_{n})h_{\mu_{n}}(\alpha,T),$$
$$= \sum_{n \in \mathbb{N}} \mu(Y_{n})h_{\mu_{n}}(\alpha_{k_{n}},T) \leq \int_{Y} h_{\mu_{y}}^{+}(\mathcal{U}|\beta,T) d\nu(y) + \epsilon.$$

Since this is true for any $\epsilon > 0$ this proves the reverse inequality for $h^+_{\mu}(\mathcal{U}|\beta, T)$. Finally using lemma 7 part (3) proves the result for $h^-_{\mu}(\mathcal{U}|\beta, T)$.

The final step in this procedure is to prove equality between h^+_{μ} and h^-_{μ} we state a universal version of Rohlin's Lemma following the ideas in [GW].

Lemma 12. Let (X,T) an invertible TDS. Then $\forall N \in \mathbb{N}$ and $\epsilon > 0$, there exists $D \in \mathcal{B}(X)$ such that $D, TD, \ldots, T^{N-1}D$ are pairwise disjoint and $\mu(\bigcup_{n=0}^{N-1} T^n D) > 1 - \epsilon$ for every non atomic measure $\mu \in \mathcal{M}_T^e(X)$.

Theorem 13. For every TDS(X,T), $\mathcal{U} \in \mathcal{C}_X$, $\beta \in \mathcal{P}_X$ we have that $\inf_{\alpha \succeq \mathcal{U}} \sup_{\mu \in \mathcal{M}_T(X)} h_{\mu}(\alpha|\beta,T) \leq 1$

 $h(\mathcal{U}|\beta, T).$

Proof:

By Lemma (11) we just need to prove it for ergodic measures moreover just for non atomic ergodic measures since atomic ergodic measures have zero measure theoretical entropy.

Let $\mathcal{U} = \{U_1, \ldots, U_k\}$ and for any $N \in \mathbb{N}$ let $\alpha' \succeq \mathcal{U}_0^{N-1}$ such that $\mathcal{N}(\alpha'|\beta_0^{N-1}) = \mathcal{N}(\mathcal{U}_0^{N-1}|\beta_0^{N-1})$. Every element of α' is of the form

$$P = P_{i_0 i_1 \dots i_{N-1}} \subseteq \bigcap_{j=0}^{N-1} T^{-j} U_{i_j}.$$

Define the partition of D, $\alpha'_D = \{A' \cap D : A' \in \alpha'\}$. For any $\epsilon > 0$ there exists $N \in \mathbb{N}$ big enough such that:

(35)
$$\mathcal{N}(\mathcal{U}_0^{N-1}|\beta_0^{N-1}) \le e^{N(h(\mathcal{U}|\beta)+\epsilon)}, \\ -\frac{1}{N}\log\frac{1}{N} - \left(1 - \frac{1}{N}\right)\log\left(1 - \frac{1}{N}\right) < \epsilon.$$

Let $0 < \delta < 1$ small enough so that $2\sqrt{\delta} \log 2k < \epsilon$.

By Lemma 12 $\exists D \subseteq X$ with $D, TD, \ldots, T^{N-1}D$ pairwise disjoint such that $\mu(\bigcup_{i=0}^{N-1} T^i D) > 1 - \delta$ for every $\mu \in \mathcal{M}_T^e(X)$ non atomic.

We use the partition α'_D to define a partition of cardinality at most 2k where we define k atoms $\{A_1, \ldots, A_k\}$ over the set $\bigcup_{i=0}^{N-1} T^i D$ assigning to the set A_i all sets

$$T^{j}P$$
 con $P = P_{i_{0}i_{1}...i_{j}...i_{N-1}}$ where $i_{j} = i$ and $j = 0, ..., N-1$.

On the rest of the space α we use any partition that refines \mathcal{U} that can be done with at most k atoms. By construction, it is not difficult to see that $\alpha_0^{N-1} \cap D \preceq \alpha'_D$. So, for any $B \in \beta_0^{N-1}$:

(36)
$$\begin{aligned} |\alpha_0^{N-1} \cap D \cap B| &\leq |\alpha_D' \cap B| \leq |\alpha' \cap B|, \\ &\leq \mathcal{N}(\alpha'|\beta_0^{N-1}) = \mathcal{N}(\mathcal{U}_0^{N-1}|\beta_0^{N-1}). \end{aligned}$$

Let us fix $\mu \in \mathcal{M}^e_T(X)$. We will show that

(37)
$$h_{\mu}(\alpha|\beta, T) \le h(\mathcal{U}|\beta, T) + 3\epsilon.$$

Let
$$E = \bigcup_{i=0}^{N-1} T^i D$$
 such that $\mu(E) > 1 - \delta$. For $n > N$, define:

$$G_n = \{ x \in X / \frac{1}{n} \sum_{i=0}^{n-1} \mathbb{1}_E(T^i x) > 1 - \sqrt{\delta} \}.$$

By definition $\int_X \frac{1}{n} \sum_{i=0}^{n-1} \mathbb{1}_E(T^i x) d\mu(x) > 1 - \delta$ and since $\frac{1}{n} \sum_{i=0}^{n-1} \mathbb{1}_E(T^i x) \leq 1$ then:

011C11.

$$\mu(G_n) + (1 - \sqrt{\delta})(1 - \mu(G_n)) \ge \int_X \frac{1}{n} \sum_{i=0}^{n-1} \mathbb{1}_E(T^i x) d\mu(x) > 1 - \delta \Rightarrow \mu(G_n) > 1 - \sqrt{\delta}.$$

For $x \in G_n$, define $S_n(x) = \{i \in \{0, 1, \dots, n-1\}/T^i x \in D\}$. By definition for any $i = 0, 1, \dots, n-1, T^i x \notin E$ if and only if $\forall j \in \{0, 1, \dots, N-1\}, i \notin S_n(x) + j$ thus:

(38)
$$\left| \{0, 1, \dots, n-1\} \setminus \bigcup_{j=0}^{N-1} (S_n(x)+j) \right| \le n\sqrt{\delta}.$$

Let $\mathcal{F}_n = \{S_n(x)/x \in G_n\}$. By disjointness of the *T*-iterations of *D* for any $F = \{s_1, \ldots, s_l\} \in \mathcal{F}_n, F \cap (F+i) = \emptyset, i = 1, \ldots, N-1$, so $|F| \leq \frac{n}{N} + 1$. Define $a_n = \lfloor \frac{n}{N} \rfloor + 1$ then:

$$|\mathcal{F}_n| \le \sum_{j=1}^{a_n} \binom{n}{j} \le a_n \binom{n}{a_n} \le n \binom{n}{a_n}.$$

By Stirling's formula and equation(35):

$$\lim_{n \to \infty} \frac{1}{n} \log\left(n \binom{n}{a_n}\right) = -\frac{1}{N} \log \frac{1}{N} - \left(1 - \frac{1}{N}\right) \log\left(1 - \frac{1}{N}\right) < \epsilon.$$

This implies that:

(39)
$$\limsup_{n \to \infty} \frac{1}{n} \log |\mathcal{F}_n| \le \lim_{n \to \infty} \frac{1}{n} \log \left(n \binom{n}{a_n} \right) \le \epsilon.$$

For $F \in \mathcal{F}_n$ define $B_F = \{x \in G_n / S_n(x) = F\}$ and $\gamma = \{B_F / F \in \mathcal{F}_n\}$ a measurable partition of G_n .

Let
$$H_F = \{0, 1, \dots, n-1\} \setminus \bigcup_{j=0}^{N-1} (F+j)$$
 by (38), $|H_F| \le n\sqrt{\delta}$.
For any $B = \bigcap_{i=0}^{n-1} T^{-i} B_i \in \beta_0^{n-1}$ define $\forall j = 1, \dots, l-1$:
 $B^{s_j} = \bigcap_{t=0}^{N-1} T^{-t} B_{s_j+t}$ and $B^{s_l} = \bigcap_{t=0}^{n-1-s_l} T^{-t} B_{s_l+t}$
Then $B = \bigcap_{j=1}^{l-1} T^{-s_j} B(s_j) \cap T^{-s_l} B(s_l) \cap \bigcap_{r \in H_F} T^{-r} B_r$.

Since for all j = 1, ..., l - 1, $B(s_j) \in \beta_j \subseteq \beta_0^{n-1}$. Since $B_F \subseteq \bigcap_{j=1}^l T^{-s_j} D$ and equation (36):

$$\begin{aligned} \left| \alpha_{0}^{n-1} \cap B_{F} \cap B \right| &\leq |\bigvee_{j=1}^{l} T^{-s_{j}} (\alpha_{0}^{N-1} \cap D \cap B^{s_{j}}) \vee \bigvee_{r \in H_{F}} T^{-r} (\alpha \cap B_{r})|, \\ &\leq \prod_{j=1}^{l} |\alpha_{0}^{N-1} \cap D \cap B^{s_{j}}| \cdot \prod_{r \in H_{F}} |\alpha \cap B_{r}|, \\ &\leq \mathcal{N} (\mathcal{U}_{0}^{N-1} |\beta_{0}^{N-1})^{l-1} |\alpha|^{|H_{F}|+N} \leq \mathcal{N} (\mathcal{U}_{0}^{N-1} |\beta_{0}^{N-1})^{\frac{n}{N}} (2k)^{n\sqrt{\delta}+N}. \end{aligned}$$

Then, adding over $F \in \mathcal{F}_n$, we have for every $B \in \beta_0^{n-1}$

(40)
$$\sum_{F \in \mathcal{F}_n} |\alpha_0^{n-1} \cap B_F \cap B| \le |\mathcal{F}_n| \mathcal{N}(\mathcal{U}_0^{N-1}|\beta_0^{N-1})^{\frac{n}{N}} (2k)^{n\sqrt{\delta}+N}.$$

Then by equation (40):

(41)

$$H_{\mu_B}(\alpha_0^{n-1} \vee \gamma) \leq \sum_{F \in \mathcal{F}_n} \sum_{C \in \alpha_0^{n-1} \cap B_F} \phi(\mu_B(C)) + \phi(\mu_B(X \setminus G_n)),$$

$$\leq \log(\sum_{F \in \mathcal{F}_n} |\alpha_0^{n-1} \cap B_F \cap B| + 1),$$

$$\leq \log(|\mathcal{F}_n| \mathcal{N}(\mathcal{U}_0^{N-1} |\beta_0^{N-1})^{\frac{n}{N}} (2k)^{n\sqrt{\delta}+N} + 1).$$

A simple calculation shows that:

(42)

$$\mu(B)H_{\mu_B}(\alpha_0^{n-1} \cap \{X \setminus G_n\}) = \mu((X \setminus G_n) \cap B)H_{\mu_{((X \setminus G_n) \cap B)}}(\alpha_0^{n-1}) + \mu(B)H_{\mu_B}(\{X \setminus G_n\}),$$

$$\leq \mu((X \setminus G_n) \cap B)\log(2k)^n + \mu(B)\log 2.$$

By equations (41) and (42):

(43)
$$\begin{aligned} H_{\mu}(\alpha_{0}^{n-1}|\beta_{0}^{n-1}) &\leq H_{\mu}(\alpha_{0}^{n-1} \vee (\gamma \cup \{X \setminus G_{n}\})|\beta_{0}^{n-1}), \\ &= \sum_{B \in \beta_{0}^{n-1}} \mu(B) \left[H_{\mu_{B}}(\alpha_{0}^{n-1} \vee \gamma) + H_{\mu_{B}}(\alpha_{0}^{n-1} \cap \{X \setminus G_{n}\}) \right], \\ &\leq \log(|\mathcal{F}_{n}|\mathcal{N}(\mathcal{U}_{0}^{N-1}|\beta_{0}^{N-1})^{\frac{n}{N}}(2k)^{n\sqrt{\delta}+N} + 1) + \sqrt{\delta}\log(2k)^{n} + \log 2. \end{aligned}$$

By the form in which we choose δ and equations (39) and (43):

$$h_{\mu}(\alpha|\beta, T) = \lim_{n \to \infty} \frac{1}{n} H_{\mu}(\alpha_0^{n-1}|\beta_0^{n-1}),$$

$$\leq \limsup_{n \to \infty} \frac{1}{n} \left(\log |\mathcal{F}_n| + \frac{n}{N} \log \mathcal{N}(\mathcal{U}_0^{N-1}|\beta_0^{N-1}) + (2n\sqrt{\delta} + N) \log 2k + \log 2 \right),$$

$$\leq \frac{1}{N} \log \mathcal{N}(\mathcal{U}_0^{N-1}|\beta_0^{N-1}) + 2\epsilon.$$

Finally by equation (25), $h_{\lambda}(\alpha|\beta, T) \leq h(\mathcal{U}|\beta, T) + 2\epsilon.$

Finally by equation (35), $h_{\mu}(\alpha|\beta, T) \leq h(\mathcal{U}|\beta, T) + 3\epsilon$.

Since this is true $\forall \mu \in \mathcal{M}_T^e(X)$ and ϵ is arbitrary we conclude that:

$$\inf_{\alpha \succeq \mathcal{U}} \sup_{\mu \in \mathcal{M}_T^e(X)} h_{\mu}(\alpha | \beta, T) \le h(\mathcal{U} | \beta, T).$$

Fix $\beta \in \mathcal{P}_X$ and $\mathcal{U} \in \mathcal{C}_X$ as in [GW] define:

(44)
$$\check{h}(\mathcal{U}|\beta,T) = \sup_{\mu \in \mathcal{M}_T(X)} \inf_{\alpha \succeq \mathcal{U}} h_{\mu}(\alpha|\beta,T) = \sup_{\mu \in \mathcal{M}_T(X)} h_{\mu}^+(\mathcal{U}|\beta,T),$$

(45)
$$\hat{h}(\mathcal{U}|\beta,T) = \inf_{\alpha \succeq \mathcal{U}} \sup_{\mu \in \mathcal{M}_T(X)} h_{\mu}(\alpha|\beta,T).$$

Clearly $\check{h}(\mathcal{U}|\beta, T) \leq \hat{h}(\mathcal{U}|\beta, T)$. By Theorem 10 $h(\mathcal{U}|\beta, T) \leq \check{h}(\mathcal{U}|\beta, T)$ and by Theorem 13 $\hat{h}(\mathcal{U}|\beta, T) \leq h(\mathcal{U}|\beta, T)$. So $h(\mathcal{U}|\beta, T) = \check{h}(\mathcal{U}|\beta, T) = \hat{h}(\mathcal{U}|\beta, T)$.

We can state an analogue of lemma 9 in [HMRY]:

Lemma 14. For any $\beta \in \mathcal{P}_X$, for every $M \in \mathbb{N}$ and $\epsilon > 0$ there exists $\delta > 0$ such that for every pair of measurable covers $\mathcal{U} = \{U_1, \ldots, U_M\}$ and $\mathcal{V} = \{V_1, \ldots, V_M\}$ such that $\mu(\mathcal{U}\Delta\mathcal{V}) < \delta$ one has that $|h^+_{\mu}(\mathcal{U}|\beta, T) - h^+_{\mu}(\mathcal{V}|\beta, T)| \leq \epsilon$.

Proof: Fix $M \in \mathbb{N}$ and $\epsilon > 0$ and choose $\delta > 0$ given by lemma 3. Now fix two measurable covers $\mathcal{U} = \{U_1, \ldots, U_M\}$ and $\mathcal{V} = \{V_1, \ldots, V_M\}$ such that $\mu(\mathcal{U}\Delta\mathcal{V}) < \delta$.

First we prove that $\alpha = \{A_1, \ldots, A_M\} \in \mathcal{U}^*$ there exists $\alpha' \in \mathcal{V}^*$ such that:

(46)
$$h_{\mu}(\alpha|\beta,T) \ge h_{\mu}(\alpha'|\beta,T) - \epsilon.$$

Define $\alpha' = \{A'_1, \dots, A'_M\}$ as:

$$A'_{1} = V_{1} \setminus \left(\bigcup_{m > 1} A_{m} \cap V_{m} \right),$$

$$A'_{m} = V_{m} \setminus \left(\bigcup_{k > m} A_{k} \cap V_{k} \cup \bigcup_{l < m} A'_{l} \right) \text{ for } 1 < m \le M.$$

For $1 \leq m \leq M$ by construction $A_m \cap V_m \subseteq A'_m$, $A_m \setminus A'_m \subseteq U_m \setminus V_m$ and $A'_m \setminus A_m \subseteq \bigcup_{k \neq m} U_k \setminus V_k$. This implies that $A_m \Delta A'_m \subseteq \bigcup_{m=1}^M U_m \Delta V_m$ and so $\mu(\alpha \Delta \beta) < M \cdot \mu(\mathcal{U}\Delta \mathcal{V}) < \delta$ that proves $h_\mu(\alpha|\beta, T) \geq h_\mu(\alpha'|\beta, T) - \epsilon$.

Since for every $\alpha' \succeq \mathcal{V}$, $h_{\mu}(\alpha'|\beta, T) \ge h_{\mu}^+(\mathcal{V}|\beta, T)$ from equation (46) we conclude that $h_{\mu}(\alpha|\beta, T) \ge h_{\mu}^+(\mathcal{V}|\beta, T) - \epsilon$. Taking infima over all $\alpha \in \mathcal{U}^*$ from we conclude that $h_{\mu}^+(\mathcal{U}|\beta, T) \ge h_{\mu}^+(\mathcal{V}|\beta, T) - \epsilon$.

Exchanging the roles of \mathcal{U} and \mathcal{V} shows that $h^+_{\mu}(\mathcal{V}|\beta, T) \ge h^+_{\mu}(\mathcal{U}|\beta, T) - \epsilon$ and so $|h^+_{\mu}(\mathcal{U}|\beta, T) - h^+_{\mu}(\mathcal{V}|\beta, T)| \le \epsilon$.

Theorem 15. For every TDS (X,T), $\mathcal{U} \in \mathcal{C}_X$, $\beta \in \mathcal{P}_X$ and $\forall \mu \in \mathcal{M}_T(X)$, $h^+_{\mu}(\mathcal{U}|\beta,T) = h^-_{\mu}(\mathcal{U}|\beta,T)$.

Proof:

First consider the case when (X, T) is uniquely ergodic. Fix $\mathcal{V} \in \mathcal{C}_X$ and define $R = |\mathcal{V}|$ by lemma 14 there exists $\delta_1 > 0$ such that for every $\mathcal{U}_1 \in \mathcal{C}_X$, $|\mathcal{U}_1| = R$ and $\mu(\mathcal{U}_1 \Delta \mathcal{V}) < \delta_1$ one has:

(47)
$$h^+_{\mu}(\mathcal{V}|\beta, T) \le h^+_{\mu}(\mathcal{U}_1|\beta, T) + \epsilon.$$

By lemma 7 part (3) we can choose $M \in \mathbb{N}$ such that:

(48)
$$\frac{1}{M}h^+_{\mu}(\mathcal{V}^{M-1}_0|\beta, T^M) \le h^-_{\mu}(\mathcal{U}|\beta, T) + \frac{\epsilon}{2}$$

Once again by lemma 14 there exists $\delta_2 > 0$ such that for every $\mathcal{U}_2 \in \mathcal{C}_X$, $|\mathcal{U}_2| = R^M$ and $\mu(\mathcal{U}_2 \Delta \mathcal{V}_0^{M-1}) < \delta_1$ one has:

(49)
$$\frac{1}{M}h_{\mu}^{+}(\mathcal{V}_{0}^{M-1}|\beta,T^{M}) \leq \frac{1}{M}h_{\mu}^{+}(\mathcal{U}_{2}|\beta,T^{M}) + \frac{\epsilon}{2}.$$

Now choose any $\mathcal{U} \in \mathcal{C}_X^o$ with $|\mathcal{U}| = R$ by unique ergodicity and Theorems 10 and 13 we have that:

(50)
$$h^+_{\mu}(\mathcal{U}|\beta,T) = h(\mathcal{U}|\beta,T) = h^-_{\mu}(\mathcal{U}|\beta,T).$$

Choose $\delta < \min\{\delta_1, \frac{\delta_2}{R^M}\}$. By simple calculations:

(51)
$$\mu(\mathcal{U}_0^{M-1}\Delta\mathcal{V}_0^{M-1}) \le R^M \mu(\mathcal{U}\Delta\mathcal{V}) < \delta_2$$

By equations (48) and (49):

(52)
$$\frac{1}{M}h_{\mu}^{+}(\mathcal{U}_{0}^{M-1}|\beta, T^{M}) \leq \frac{1}{M}h_{\mu}^{+}(\mathcal{V}_{0}^{M-1}|\beta, T^{M}) + \frac{\epsilon}{2} \leq h_{\mu}^{-}(\mathcal{V}|\beta, T) + \epsilon.$$

By equations (48), (49), (50) and lemma lemma 7 part (2):

(53)
$$h^{+}_{\mu}(\mathcal{V}|\beta, T) \leq h^{+}_{\mu}(\mathcal{U}|\beta, T) + \epsilon = h^{-}_{\mu}(\mathcal{U}|\beta, T) + \epsilon = \frac{1}{M}h^{-}_{\mu}(\mathcal{U}^{M-1}_{0}|\beta, T^{M}) + \epsilon,$$
$$\leq \frac{1}{M}h^{+}_{\mu}(\mathcal{U}^{M-1}_{0}|\beta, T^{M}) + \epsilon \leq h^{-}_{\mu}(\mathcal{V}|\beta, T) + 2\epsilon.$$

Since this is for any $\epsilon > 0$ this proves that $h^+_{\mu}(\mathcal{V}|\beta, T) = h^-_{\mu}(\mathcal{V}|\beta, T)$.

Now by lemma 11 we just need to prove the equality for $\mu \in \mathcal{M}^e_T(X)$.

By the Jewett-Krieger theorem (see [W]) there exists a φ -measure theoretical isomorphism with a uniquely ergodic TDS (Y, S). In general $h^+_{\nu}(\varphi^{-1}\mathcal{U}|\varphi^{-1}\beta, S) \leq h^+_{\mu}(\mathcal{U}|\beta, T)$ but since φ is an isomorphism it is also true that $h^+_{\nu}(\varphi^{-1}\mathcal{U}|\varphi^{-1}\beta, S) \geq h^+_{\mu}(\mathcal{U}|\beta, T)$. By lemma 8 $h^-_{\mu}(\mathcal{U}|\beta, T) = h^-_{\nu}(\varphi^{-1}\mathcal{U}|\varphi^{-1}\beta, S) = h^+_{\nu}(\varphi^{-1}\mathcal{U}|\varphi^{-1}\beta, S) = h^+_{\mu}(\mathcal{U}|\beta, T)$.

5. Relation with other Variational Principles

This obviously extendes the work done in [R] and [GW] just by considering the case $\beta = \{X\}$. Another direct application is the work done in [HYZ] where they consider a fixed φ -factor (Y, \mathcal{D}, S) and denote $\mathcal{B}_{\varphi} = \varphi^{-1}\mathcal{B}_X$ to define for any $\mathcal{U} \in \mathcal{C}_X$:

$$H_{\mu}(\mathcal{U}|Y) := H_{\mu}(\mathcal{U}|\mathcal{B}_{\varphi}), \ \mathcal{N}(\mathcal{U}|Y) := \sup_{y \in Y} \mathcal{N}(\mathcal{U} \cap \varphi^{-1}\{y\}),$$
$$h_{\mu}^{-}(\mathcal{U}|Y,T) := \lim_{N \to \infty} \frac{1}{N} H_{\mu}(\mathcal{U}_{0}^{N-1}|\mathcal{B}_{\varphi}), \\ h_{\mu}^{+}(\mathcal{U}|Y,T) := \inf_{\alpha \succeq \mathcal{U}} h_{\mu}(\alpha|\mathcal{B}_{\varphi},T),$$
$$h(\mathcal{U}|Y,T) := \lim_{N \to \infty} \frac{1}{N} \log \mathcal{N}(\mathcal{U}_{0}^{N-1}|\mathcal{B}_{\varphi}).$$

By some classic and simple calculations (see REMARK 4 in [DS] for instance) $H_{\mu}(\mathcal{U}|\mathcal{B}_{\varphi}) = \inf_{\beta \in \mathcal{P}_X} H_{\mu}(\mathcal{U}|\varphi^{-1}\beta) \text{ and } \mathcal{N}(\mathcal{U}|Y) = \inf_{\beta \in \mathcal{P}_X} \mathcal{N}(\mathcal{U}|\varphi^{-1}\beta).$

So Theorems 10, 13 and 15 extend to this notion as proven in [HYZ].

6. CONCLUSION AND FINAL REMARKS

The results of this paper show that the variational principle once again is more local than was previously stated, now with respect to the conditioning variable. This takes us one step closer to the ultimate local variational principle, this is, for an open cover conditioned with respect to a fixed open cover. Once again, the technique developed in [R] to define a "+" and "-" definitions of measure theoretical entropy and proving their equality remains crucial and unavoidable and works for all measurable covers.

The requirement of the cover to be open is once more proven to be necessary for the existence of a local variational principle. This fails even for closed covers as the pioneers of this ideas knew from the beginning for the more global principles. This shows that this is not a merely combinatorial result and the topological structure is crucial to link measure and topology. In the local case, an equality is attained for a specific invariant measure and not just an equality for the suprema over all invariant measures as for the global principles but little is known about that measure. However in many cases for the global variational principles much is known about the measure that attains the equality when it exists.

The extension of this result for more general actions requires extra work but at least for the case of countable discrete amenable groups by using the tool of Følner sequences it seems clear how to do it.

6.1. Acknowledgment. The author would like to thank Daniel Coronel for a lot of work and some of the main ideas in this work and François Blanchard, Tomasz Downarovicz, Alejandro Maass and Karl Petersen for valuable remarks and discussions. The author acknowledges the support of Programa Basal PFB 03, CMM, Universidad de Chile.

PIERRE-PAUL ROMAGNOLI

References

- [AKM] R. L. Adler, A. G. Konheim and M. H. McAndrew. Topological entropy. Trans. Amer. Math. Soc. 114 (1965), 309-319.
- [B] F. Blanchard . Fully positive topological entropy and topological mixing. Symbolic Dynamics and its Applications, 135 (Contemporary Mathematics). American Mathematical Society, Providence, RI, 1992, pp. 95-105.
- [BL] F. Blanchard and Y. Lacroix . Zero-entropy factors of topological flows. Proc. Amer. Math. Soc. 119 (1993), 985-992.
- [BGH] F. Blanchard, E. Glasner, B. Host, A variation on the variational principle and applications to entropy pairs, Ergodic Th. Dynam. Sys. 17 (1997) 29-43.
- [D] T. Downarovicz, Entropy in Dynamical Systems, New Mathematical Monographs, Cambridge University, Vol 18.
- [DS] T. Downarowicz, J. Serafin, Fiber entropy and conditional variational principles in compact non-metrizable spaces, Fund. Math., 172 (3) (2002), pp. 217-247
- [DZ] Dooley, A., Zhang, G. (2015). Local entropy theory of a random dynamical system (Vol. 233, No. 1099). American Mathematical Society.
- [G] T. N. T. Goodman. Relating topological entropy and measure entropy. Bull. London Math. Soc. 3 (1971), 176-180.
- [Gw] L. W. Goodwyn. Topological entropy bounds measure-theoretic entropy. Proc. Amer. Math. Soc. 23 (1969), 679-688.
- [Gw2] L. W. Goodwyn, Some counter-examples in topological entropy. Topology 11 (1972), 377-385.
- [GW] E. Glasner and B. Weiss, On the interplay between measurable and topological dynamics, in Handbook of Dynamical Systems, Vol. 1 B(eds. B. Hasselblatt and A. Katok), Elsevier B. V., Amsterdam, 2006, 597-648.
- [HMRY] W. Huang, A. Maass, P. P. Romagnoli, and X. Ye, Entropy pairs and a local Abramov formula for a measure theoretical entropy of open covers. Ergodic Th. Dynam. Sys. 24 (2004) 1127-1153.
- [HY] W. Huang and Y. F. Yi, A local variational principle for pressure and its applications to equilibrium states, Israel Journal of Mathematics, 161 (2007), 29-74.
- [HYZ] W. Huang, X. Ye, and G. Zhang, A local variational principle for conditional entropy.Ergodic Th. Dynam. Sys. Volume 26(2006), 219-245.
- [HYZ2] W. Huang, X. Ye, G.H. Zhang, Local entropy theory for a countable discrete amenable group action, J. Funct Anal. 261 (4) (2011) 1028-1082.
- [K] A. N. Kolmogorov. A new metric invariant of transient dynamical systems and automorphisms of Lebesgue spaces. Dokl. Akad. Sci. SSSR 119 (1958), 861-864 (Russian).
- [L] F. Ledrappier, A variational principle for the topological conditional entropy, Ergodic theory (Proc. Conf., Math. Forschungsinst., Oberwolfach, 1978), Lecture Notes in Math., vol. 729, Springer, Berlin, 1979, pp. 78-88. MR550412 (80j:54011).
- [M] M. Misiurewicz, Topological conditional entropy, Studia Math. 55 (1976) 176-200.
- [R] P.P Romagnoli, A local variational principle for the topological entropy, Ergodic Th. and Dynam. Sys. 23 (2003) 1601-1610.
- [S] U. Shapira, Isr. J. Math. (2007) 158: 225. doi:10.1007/s11856-007-0012-z
- [W] P. Walters, 1982. An Introduction to Ergodic Theory. Graduate Texts in Mathematics, vol. 79. New York: Springer-Verlag.
- [Z] Zhang, Guohua., Local variational principle concerning entropy of a sofic group action, Journal of Functional Analysis 262.4 (2012): 1954-1985.

DEPARTAMENTO DE MATEMÁTICAS, UNIVERSIDAD ANDRES BELLO, REPÚBLICA 252, SANTIAGO, CHILE

E-mail address: promagnoli@unab.cl