

The large- N limit for two-dimensional Yang–Mills theory

Brian C. Hall*

University of Notre Dame
Department of Mathematics
Notre Dame, IN 46556 USA
bhall@nd.edu

Abstract

The analysis of the large- N limit of $U(N)$ Yang–Mills theory on a surface proceeds in two stages: the analysis of the Wilson loop functional for a simple closed curve and the reduction of more general loops to a simple closed curve. In the case of the 2-sphere, the first stage has been treated rigorously in recent work of Dahlqvist and Norris, which shows that the large- N limit of the Wilson loop functional for a simple closed curve in S^2 exists and that the associated variance goes to zero.

We give a rigorous treatment of the second stage of analysis in the case of the 2-sphere. Dahlqvist and Norris independently performed such an analysis, using a similar but not identical method. Specifically, we establish the existence of the limit and the vanishing of the variance for arbitrary loops with (a finite number of) simple crossings. The proof is based on the Makeenko–Migdal equation for the Yang–Mills measure on surfaces, as established rigorously by Driver, Gabriel, Hall, and Kemp, together with an explicit procedure for reducing a general loop in S^2 to a simple closed curve. The methods used here also give a new proof of these results in the plane case, as a variant of the methods used by Lévy.

We also consider loops on an arbitrary surface Σ . We put forth two natural conjectures about the behavior of Wilson loop functionals for topologically trivial simple closed curves in Σ . Under the weaker of the conjectures, we establish the existence of the limit and the vanishing of the variance for topologically trivial loops with simple crossings that satisfy a “smallness” assumption. Under the stronger of the conjectures, we establish the same result without the smallness assumption.

Contents

1 Introduction and main results

2

*Supported in part by NSF Grant DMS-1301534.

1.1	The Makeenko–Migdal equation in two dimensions	2
1.2	The master field in two dimensions	4
1.3	The master field on the sphere	5
1.4	The reduction procedure	6
1.5	The Wilson loop for simple closed curve in S^2	9
2	Tools for the proof	11
2.1	Variation of the Wilson loop and of the variance	11
2.2	Variance estimates	14
3	Examples	15
3.1	The figure eight	15
3.2	The trefoil	19
4	Analysis of a general loop	20
4.1	The strategy	20
4.2	Winding numbers	21
4.3	The span of the Makeenko–Migdal vectors	23
4.4	Shrinking all but two of the faces	23
4.5	Analyzing the limiting case	24
4.6	The induction	28
4.7	The n -fold circle	30
5	The plane case, revisited	34
5.1	The general result	34
5.2	Example computations in the plane case	36
6	General compact surfaces	37
6.1	Introduction	37
6.2	Statements	39
6.3	Proofs	40
7	Acknowledgments	44

1 Introduction and main results

1.1 The Makeenko–Migdal equation in two dimensions

Let us fix a connected compact Lie group K together with an Ad-invariant inner product on its Lie algebra, \mathfrak{k} . The path integral for Euclidean Yang–Mills theory over a manifold M is supposed to describe a probability measure on the space of connections for a principal K -bundle over M . One of the main objects of study in such a theory is the *Wilson loop functional*, namely the expectation value of the trace (in some fixed representation of K) of the holonomy of the connection around a loop. The Makeenko–Migdal equation is an identity for the variation of Wilson loop functionals with respect to a variation in the loop. The




Figure 1: The labeling of the faces surrounding v

original version of this equation, in any number of dimensions, was proposed by Makeenko and Migdal in [MM]. A version specific to the two-dimensional case was then developed by Kazakov and Kostov in [KK, Eq. (24)]. (See also [K, Eq. (9)] and [GG, Eq. (6.4)].)

A special feature of the two-dimensional Yang–Mills measure is its invariance under area-preserving diffeomorphisms. Suppose we fix the topological type of a loop L in a surface Σ and consider the faces of L , that is, the connected components of the complement of L in Σ . Then the Wilson loop functional depends only on the areas of the faces of L . Let us now take $K = U(N)$ with the inner product on the Lie algebra $u(N)$ given by the scaled Hilbert–Schmidt inner product,

$$\langle X, Y \rangle := N \text{Trace}(X^*Y). \quad (1)$$

It is then convenient to express the Wilson loop functionals in terms of the normalized trace,

$$\text{tr}(X) := \frac{1}{N} \text{Trace}(X). \quad (2)$$

We now consider a loop L with (a finite number of) simple crossings, and we let v be one such crossing. We label the four faces of L adjacent to the crossing in cyclic order as F_1, \dots, F_4 , with F_1 denoting the face whose boundary contains the two outgoing edges of L . We then let t_1, \dots, t_4 denote the areas of these faces. (See Figure 1.) We also let L_1 denote the loop from the beginning to the first return to v and let L_2 denote the loop from the first return to the end. (See Figure 2.)

The two-dimensional version of the Makeenko–Migdal equation, in the $U(N)$ case, is then as follows:

$$\left(\frac{\partial}{\partial t_1} - \frac{\partial}{\partial t_2} + \frac{\partial}{\partial t_3} - \frac{\partial}{\partial t_4} \right) \mathbb{E}\{\text{tr}(\text{hol}(L))\} = \mathbb{E}\{\text{tr}(\text{hol}(L_1))\text{tr}(\text{hol}(L_2))\}, \quad (3)$$

where $\text{hol}(\cdot)$ denotes the holonomy. Although the curves L_1 and L_2 occurring on the right-hand side of (3) are simpler than the loop L , the right-hand side




Figure 2: The loops L_1 (black) and L_2 (dashed) for the loop in Figure 1

of (3) involves the *expectation of the product* of the traces, rather than the product of the expectations. Thus, even if one has already computed the Wilson loop functionals $\mathbb{E}\{\text{tr}(\text{hol}(L_1))\}$ and $\mathbb{E}\{\text{tr}(\text{hol}(L_2))\}$, the right-hand side of (3) cannot be regarded as a known quantity. In the large- N limit, however, we will see that the Makeenko–Migdal equation becomes an effective tool for inductive computation of Wilson loop functionals.

The original argument of Makeenko and Migdal for the equation that bears their names was based on heuristic manipulations of the path integral. In the plane case, Lévy then gave a rigorous proof of the Makeenko–Migdal equation in [Lév2]. (See Eq. (159) in Proposition 9.2.2 of [Lév2].) Subsequent proofs of the planar Makeenko–Migdal equation were then provided by Dahlqvist [Dahl] and Driver–Hall–Kemp [DHK2].

Meanwhile, in [DGHK], Driver, Gabriel, Hall, and Kemp gave a rigorous derivation of the Makeenko–Migdal equation for $U(N)$ Yang–Mills theory over an arbitrary surface. Actually, the proof given in [DHK2] in the plane case extends with minor modifications to the case of a general surface.

1.2 The master field in two dimensions

In the paper [’t H], ’t Hooft proposed that Yang–Mills theory for $U(N)$ in any dimension should simplify in the limit as $N \rightarrow \infty$. In particular, it is expected that in this limit, the path integral should concentrate onto a *single* connection (modulo gauge transformations), known as the master field. The concentration phenomenon for the Yang–Mills measure has an important implication for the form of the two-dimensional Makeenko–Migdal equation. Specifically, in the limit, there should be no difference between the expectation of a product of traces and the product of the associated expectations: both $\mathbb{E}\{fg\}$ and $\mathbb{E}\{f\}\mathbb{E}\{g\}$ should become $f(M_0)g(M_0)$, where M_0 is the master field.

If, therefore, the large- N limit of $U(N)$ Yang–Mills theory exists on a surface

Σ , we expect it to satisfy a Makeenko–Migdal equation of the form

$$\left(\frac{\partial}{\partial t_1} - \frac{\partial}{\partial t_2} + \frac{\partial}{\partial t_3} - \frac{\partial}{\partial t_4} \right) W(L) = W(L_1)W(L_2), \quad (4)$$

where $W(L)$ is the limiting value of $\mathbb{E}\{\text{tr}(\text{hol}(L))\}$. Note that the loops L_1 and L_2 on the right-hand side of (4) have fewer crossings than L , since neither L_1 nor L_2 has a crossing at v . Thus, one may hope that the large- N Makeenko–Migdal equation may allow one to reduce computations of Wilson loop functionals for general curves to simpler ones, until one eventually reaches a simple closed curve. Of course, since a simple closed curve has no crossings, the Makeenko–Migdal equation gives no information about the Wilson loop for such a curve.

In the plane case, the structure of the master field was worked out by Singer [Si], Gopakumar and Gross [GG, Gop], Xu [Xu], Sengupta [Sen4], Anshelevich and Sengupta [AS], and then in greater detail by Lévy [Lév2]. In particular, the expected concentration phenomenon was verified in detail in the plane case in [Lév2]. (See the explicit variance estimate in Theorem 6.3.1 of [Lév2].) A generalization of the master field on the plane was then constructed by Cébron, Dahlqvist, and Gabriel in [CDG].

In [Lév2], Lévy shows that the large- N limit of the Wilson loop functional for a loop in the plane with simple crossings is completely determined by (4), together with another, simpler condition. This simpler condition—given as Axiom Φ_4 on p. 11 of [Lév2] and called the “unbounded face condition” in [DHK2, Theorem 2.3]—gives a simple formula for the derivative of the Wilson loop functional with respect to the area of any face of L that adjoins the unbounded face.

1.3 The master field on the sphere

The existence of a large- N limit of Yang–Mills theory on a general surface Σ is currently unknown. There has, however, been much interest in the problem because of connections with string theory, as developed by Gross and Taylor [Gr, GT1, GT2].

The S^2 case, meanwhile, has been extensively studied at varying levels of rigor. The analysis proceeds in two stages. First, one studies the large- N limit of the Wilson loop functional for a simple closed curve. Second, one attempts to use the large- N Makeenko–Migdal equation to reduce Wilson loop functionals for all other loops with simple crossings to the simple closed curve.

In the first stage of analysis, a formula was proposed in the physics literature for the Wilson loop functional for a simple closed curve. (See Section 1.5 for more information.) A notable feature of this formula is the presence of a phase transition. If the total area of the sphere is less than π^2 , the Wilson loop for a simple closed curve is expressible in terms of the semicircular distribution from random matrix theory. If, however, the total area is greater than π^2 , the Wilson loop is much more complicated. In addition to the proposed formula for the limiting Wilson loop functional, it is expected that the limit should be

deterministic, in keeping with the idea of the master field. This brings us to the following recent rigorous result of Dahlqvist and Norris [DN].

Theorem 1 (Dahlqvist–Norris) *If C is a simple closed curve on S^2 then the limit*

$$\lim_{N \rightarrow \infty} \mathbb{E} \{ \text{tr}(\text{hol}(C)) \} \tag{5}$$

exists and depends continuously on the areas of the two faces of C . Furthermore, the associated variance tends to zero:

$$\lim_{N \rightarrow \infty} \text{Var} \{ \text{tr}(\text{hol}(C)) \} = 0. \tag{6}$$

The method of proof used in [DN] is discussed briefly in Section 1.5.

Notation 2 *We denote the large- N limit of the Wilson loop functional for a simple closed curve by W_1 :*

$$W_1(a, b) = \lim_{N \rightarrow \infty} \mathbb{E} \{ \text{tr}(\text{hol}(C)) \},$$

where C is a simple closed curve and where a and b are the areas of the faces of C .

In the second stage of analysis, it has been claimed by Daul and Kazakov that, “All averages for self-intersecting loops can be reproduced from the average for a simple (non-self-intersecting) loop by means of loop equations.” (See the abstract of [DaK]. The loop equations referred to are the large- N Makeenko–Migdal equation (4).) It should be noted, however, that Daul and Kazakov analyze only two examples, and it is not obvious how to extend their analysis to general loops; see Section 3. Furthermore, they assume that the large- N limit exists and satisfies the large- N Makeenko–Migdal equation.

1.4 The reduction procedure

In this paper, we give a rigorous treatment of the second stage of the analysis of the large- N limit for Yang–Mills theory on S^2 , as well as results for the plane and general surfaces. (Dahlqvist and Norris also give treat the S^2 case by a similar but not identical method, as discussed further in Section 1.4.1.)

1.4.1 On the sphere

Specifically, we establish the following results in the sphere case: (1) the existence of the large- N limit of Wilson loop functionals for arbitrary loops with simple crossings; (2) the vanishing of the associated variance; and (3) the large- N Makeenko–Migdal equation for the limiting theory. In particular, we give a concrete procedure for reducing the Wilson loop functional for general loops in S^2 to the Wilson loop functional for a simple closed curve.

Here are some notable features of our approach.




Figure 3: A checkerboard variation of the areas

- We do not assume the existence of the large- N limit ahead of time, except for a simple closed curve (Theorem 1).
- We do not assume ahead of time that the limiting theory satisfies the large- N Makeenko–Migdal equation. Rather, we assume only the finite- N Makeenko–Migdal equation in (3), as established rigorously in [DGHK]. We then prove that the limiting theory satisfies a large- N version of the equation.
- We give a constructive procedure for reducing the Wilson loop functional an arbitrary loop in S^2 with simple crossings inductively to that for a simple closed curve. Specifically, we show that any loop can first be reduced to one that winds n times around a simple closed curve, which can then be reduced to a simple closed curve.

The just-referred-to procedure relies on a result (Proposition 10) that says that it is possible to perform a combination of Makeenko–Migdal variations at all of the vertices, with the effect that the areas of *all but two* of the faces shrink to zero, with the areas of the remaining two faces remaining non-negative. Furthermore, it is possible to choose one of the “unshrunk” faces arbitrarily.

After the first version of this paper was posted to the arXiv, I became aware of a preprint of Dahlqvist and Norris [DN], which had been posted approximately two months earlier. The paper of Dahlqvist and Norris proves Theorem 1, which I stated as a conjecture in the first version of this paper. In addition, [DN] gives a reduction procedure that is similar to, but not identical to, the one I use here. The main difference between the two approaches is the just-mentioned freedom in my approach to arbitrarily choose one of the faces whose area does not shrink to zero. This flexibility is exploited crucially to give results on arbitrary surfaces, as discussed in Sections 1.4.2 and 6.

Our main result on the sphere may be stated as follows.

Theorem 3 *If L is a closed curve traced out on a graph in S^2 and having only simple crossings, the following results hold. First, the limit*

$$W(L) := \lim_{N \rightarrow \infty} \mathbb{E} \{ \text{tr}(\text{hol}(L)) \} \quad (7)$$

exists and depends continuously on the areas of the faces of L . Second, the associated variance goes to zero:

$$\lim_{N \rightarrow \infty} \text{Var} \{ \text{tr}(\text{hol}(L)) \} = 0. \quad (8)$$

Third, the limiting expectation values satisfy the following large- N Makeenko–Migdal equation. Let us vary the areas of the faces surrounding a crossing v in a checkerboard pattern as in Figure 3, resulting in a family of curves $L(t)$. Then

$$\frac{d}{dt} W(L(t)) = W(L_1(t))W(L_2(t)), \quad (9)$$

where $L_1(t)$ and $L_2(t)$ are derived from $L(t)$ in the usual way.

In Figure 3, we do not assume the four faces are distinct. If, say, the two faces labeled as $+t$ are the same, we are then increasing the area of that face by $2t$.

The reason for stating the Makeenko–Migdal equation in the form in (9) is that we have not established the differentiability of the large- N Wilson loop functional $W(L)$ with respect to the area of an individual face. If this differentiability property turns out to hold, we can then apply the chain rule to express the derivative on the left-hand side of (9) in the usual form as an alternating sum of such derivatives. This issue is of little consequence, since the result in (9) is the way one applies the Makeenko–Migdal equation in all applications.

1.4.2 On the plane and on arbitrary surfaces

We also provide a new proof of Theorem 3 in the plane case, as a variant of the methods used by Lévy in [Lév2]. In the plane case, the result is not dependent on results of [DN], since the analog of Theorem 1 for \mathbb{R}^2 is a simple computation; see Section 5.1. In Lévy’s analysis in [Lév2], the structure of the master field on \mathbb{R}^2 is based on two main axioms, the large- N version of the Makeenko–Migdal equation and a second condition, labeled as Axiom Φ_4 in [Lév2, Section 0] and called the “unbounded face condition” in [DHK2, Theorem 2.3], which gives a formula for the derivative of a Wilson loop functional with respect to the area of any face that adjoins the unbounded face. (There are also some continuity and invariance properties.) We show that the master field on \mathbb{R}^2 can alternatively be characterized by the large- N Makeenko–Migdal equation together with the (simple) formula for the Wilson loop for a simple closed curve. See Section 5.

Finally, we consider Yang–Mills theory on an arbitrary compact surface Σ . Let us call a loop in Σ “topologically trivial” if it is contained in a topological disk $U \subset \Sigma$. We put forth two natural conjectures regarding topologically trivial

simple closed curves in Σ . The first, Conjecture 16, is simply the obvious analog of Theorem 1 for topologically trivial simple closed curves. (No such result is known for surfaces other than the plane and the sphere.) The second, Conjecture 19, asserts also similar results for a loop that winds n times around a simple closed curve, $n \in \mathbb{Z}$. Assuming the first conjecture, we establish the analog of Theorem 3 for topologically trivial loops with simple crossings that satisfy a “smallness” assumption. Assuming the second conjecture, we establish the analog of Theorem 3 for *all* topologically trivial loops with simple crossings. See Section 6.

Our results for the plane and for arbitrary surfaces depend crucially on an extra level of flexibility in the reduction process that is not present in [DN]. This paper and [DN] both use an approach in which the areas of *all but two* of the faces of the curve are shrunk to zero. In the procedure in [DN, Section 4.5], one generically has no choice regarding which two faces remain unshrunk; they are specified by conditions on the winding numbers. In our approach, by contrast, *one* of the unshrunk faces can be chosen arbitrarily. In the plane case, we choose one of the unshrunk faces to be the unbounded face, while in the case of an arbitrary surface, we choose one of the unshrunk faces to be the one containing the complement of the topological disk U .

1.5 The Wilson loop for simple closed curve in S^2

In this section, we describe three approaches (at varying levels of rigor) to analyzing the Wilson loop functional for a simple closed curve in the sphere. If C is a simple closed curve on S^2 and the areas of the two faces of C are a and b , Sengupta’s formula [Sen1] reads

$$\mathbb{E} \{ \text{tr}(\text{hol}(C)) \} = \frac{1}{Z} \int_{U(N)} \text{tr}(U) \rho_a(U) \rho_b(U) dU, \quad (10)$$

where $Z = \rho_{a+b}(\text{id})$ is a normalization factor. Here ρ_a is the heat kernel on $U(N)$, based at the identity and evaluated at “time” a . The probability measure

$$\frac{1}{Z} \rho_a(U) \rho_b(U) dU \quad (11)$$

is precisely the distribution at time a of a Brownian bridge on $U(N)$, starting at the origin and returning to the origin at time $a + b$.

In the first approach, one writes the heat kernels in (10) as sums over the characters of the irreducible representations of $U(N)$. In the large- N limit, one attempts to find the “most probable representation,” that is, the one whose character contributes the most to the sum. The representations, meanwhile, are labeled by certain diagrams; the objective is then to determine the limiting shape of the diagram for the most probable representation. Using this method, physicists have found different shapes in the small-area phase (namely $a + b < \pi^2$) and the large-area phase (namely $a + b > \pi^2$). (See works by Douglas and Kazakov [DoK] and Boulatov [Bou].)

At a rigorous level, Boutet de Monvel and Shcherbina [BS] and Lévy and Maïda [LM] have analyzed the partition function (i.e., the normalization factor $Z = \rho_{a+b}(\text{id})$) by this method and confirmed the existence of a phase transition at $a + b = \pi^2$. Then, recently, Dahlqvist and Norris [DN, Section 3] have given a rigorous analysis of the Wilson loop functional using a rigorous version of the arguments in [DoK] and [Bou], leading to Theorem 1.

In the second approach, one writes the heat kernels in (10) as a sum over all geodesics connecting the identity to U , using a formula developed by Ęskin [Ęs] and rediscovered by Urakawa [Ur]. (This formula is a Poisson-summed version of the formula as a sum of characters.) When the quantities a and b in (10) are small, the contribution of the shortest geodesic dominates. Recall that we are using the scaled Hilbert–Schmidt inner product (1) on the Lie algebra $u(N)$. Since the Laplacian scales oppositely to the inner product, the Laplacian on $U(N)$ is scaled by a factor of $1/N$ compared to the Laplacian for the unscaled Hilbert–Schmidt inner product. Thus, at a heuristic level, the large- N limit ought to be pushing us toward the small-time regime for the heat kernels ρ_a and ρ_b . It is therefore possible that in the large- N limit, one can simply “neglect the winding terms,” that is, include only the contribution from the shortest geodesic.

The contribution of the shortest geodesic, meanwhile, is a Gaussian integral of the sort that arises in the Gaussian unitary ensemble (GUE) in random matrix theory. Thus, *if* it is valid to keep only the contribution from the shortest geodesic, the Wilson loop functional may be computed using results from GUE theory. (See the work of Daul and Kazakov in [DaK].) On the other hand, a consistency argument indicates that neglecting the winding terms can only be valid in the small area phase. Little work has been done, however, in estimating the size of the winding terms.

In the third approach, one may, as we have noted, recognize the probability measure in (11) as the distribution of a Brownian bridge on $U(N)$. Forrester, Majumdar, and Schehr have then developed a method [FMS] to represent the partition function for Yang–Mills theory in terms of a collection of N nonintersecting Brownian bridges on the unit circle. (That is, we consider N Brownian motions in the unit circle, starting at 1. We then constrain them to return to 1 at time $T = a + b$ and to be nonintersecting for all times $0 < t < T$.) In fact, the distribution of the eigenvalues of the Brownian bridge in $U(N)$ is precisely the distribution of these nonintersecting Brownian bridges. This claim is presumably well known to experts—I learned it from Thierry Lévy—and is explained in the notes [Ha2]. (The claim is analogous to the well-known result that the eigenvalues of a Brownian motion in the space of $N \times N$ Hermitian matrices are described by the “Dyson Brownian motion” [Dys] in \mathbb{R}^N . See Section 3.1 of [Tao].) Thus, not just the partition function, but also the Wilson loop functional for a simple closed curve can be expressed in terms of nonintersecting Brownian bridges.

Meanwhile, Liechty and Wang [LW1, LW2] have obtained various rigorous results about the large- N behavior of the nonintersecting Brownian bridges in S^1 . In particular, they confirm the existence of a phase transition: When the

lifetime $a+b$ of the bridge is less than π^2 , the nonintersecting Brownian motions do not wind around the circle, whereas for lifetime greater than π^2 they do. It is possible that one could establish Theorem 1 rigorously in the small-area phase using results from [LW1]. (Theorem 1.2 of [LW1] would be relevant.) In the large-area phase, however, [LW1] does not provide information about the distribution of eigenvalues when t is close to half the lifetime of the bridge. (See the restrictions on θ in Theorem 1.5(a) of [LW1].)

2 Tools for the proof

In this section, we review some prior results that will allow us to prove our main theorem. Our main tool, besides the crucial result of Dahlqvist and Norris in Theorem 1, is the Makeenko–Migdal equation for $U(N)$ Yang–Mills theory on compact surfaces, which was established at a rigorous level in [DGHK]. More precisely, we require not only the standard Makeenko–Migdal equation in (3), but also an “abstract” Makeenko–Migdal equation, which allows us to compute the alternating sum of derivatives of expectation values of more general functions. We also require an estimate on the variance of the product of two bounded random variables, as described in Section 2.2.

2.1 Variation of the Wilson loop and of the variance

Rigorous constructions of the two-dimensional Yang–Mills measure with structure group K from a continuum perspective were given in the plane case by Gross, King, and Sengupta [GKS] and by Driver [Dr], and in the case of a compact surface, possibly with boundary, by Sengupta [Sen1, Sen2, Sen3]. (See also [Lév1], which, among other things, extends the analysis to rectifiable loops with infinitely many self-intersections.) In particular, suppose \mathbb{G} is an “admissible” graph in a surface Σ , meaning that \mathbb{G} contains the boundary of Σ and that each face of \mathbb{G} is a topological disk. Let e denote the number of unoriented edges of \mathbb{G} and let g be a gauge-invariant function of the connection that depends only on the parallel transports x_1, \dots, x_e along the edges of \mathbb{G} . Then Driver (in the plane case) and Sengupta (in the general case) give a formula for the expectation value of g with respect to the Yang–Mills measure. The formulas of Driver and Sengupta correspond to what is known as the *heat kernel action* in the physics literature, as developed by Menotti and Onofri [MO] and others.

Let $\rho_t : K \rightarrow \mathbb{R}$ denote the heat kernel on K , based at the identity. Then we have, explicitly,

$$\mathbb{E}\{g\} = \frac{1}{Z} \int_{K^e} g(x_1, \dots, x_e) \prod_i \rho_{|F_i|}(\text{hol}(F_i)) dx_1 \cdots dx_e, \quad (12)$$

where dx_i denotes the normalized Haar measure on K , $|F_i|$ is the area of the i th face, and $\text{hol}(F_i)$ is the product of edge variables going around the boundary of F_i . Here Z is a normalization constant. Since ρ is invariant under conjugation

and inversion, the formula does not depend on the starting point or orientation of the boundary of F_i . If the boundary of Σ is nonempty, it is possible to incorporate into (12) constraints on the holonomies around the boundary components; the proof of the Makeenko–Migdal equation in [DGHK] holds in this more general context.

Remark 4 *In the rest of the paper, when we speak about “varying the areas” of the faces of graph, we mean more precisely that we replace the numbers $\{|F_i|\}$ by some other collection of positive real numbers $\{t_i\}$ in Sengupta’s formula (12). If the sum of the t_i ’s equals the sum of the $|F_i|$ ’s, it may be possible to implement this variation “geometrically,” by continuously deforming the graph, but this is not necessary. In particular, the Makeenko–Migdal equation (3) was proved under such an “analytic” (i.e., not necessarily geometric) variation of the area.*

If we have a fixed loop L and we let the areas of the faces of L depend on a parameter t , we will (in a small abuse of notation) denote that pair consisting of L and the collection of areas by $L(t)$.

Suppose now that L is a loop that can be traced out on an oriented graph in Σ and let \mathbb{G} be a minimal graph on which L can be traced. We now explain what it means for L to have a simple crossing at a vertex v . First, we assume that \mathbb{G} has exactly four edges incident to v , where we count an edge e twice if both the initial and final vertices of e are equal to v . Second, we assume that L , when viewed as a map of the circle into the plane, passes through v exactly twice. Third, we assume that each time L passes through v , it comes in along one edge and passes “straight across” to the cyclically opposite edge. Last, we assume that L traverses two of the edges on one pass through v and the remaining two edges on the other pass through v .

Under these assumptions, Theorem 1 of [DGHK] gives a rigorous derivation of the Makeenko–Migdal equation in (3). We now restate the Makeenko–Migdal equation for $U(N)$, in the S^2 case, in a way that facilitates the large- N limit. In addition, we derive a similar result for the variation of the variance of $\text{tr}(\text{hol}(L))$, where for a complex-valued random variable X , we define

$$\text{Var}(X) = \mathbb{E}\{|X - \mathbb{E}\{X\}|^2\} = \mathbb{E}\{|X|^2\} - |\mathbb{E}\{X\}|^2.$$

Proposition 5 *Let L be a loop traced out on a graph in S^2 and having only simple crossings. Let v be one such crossing and let L_1 and L_2 be obtained from L as usual in the Makeenko–Migdal equation. Then we have*

$$\begin{aligned} & \left(\frac{\partial}{\partial t_1} - \frac{\partial}{\partial t_2} + \frac{\partial}{\partial t_3} - \frac{\partial}{\partial t_4} \right) \mathbb{E}\{\text{tr}(\text{hol}(L))\} \\ &= \mathbb{E}\{\text{tr}(\text{hol}(L_1))\} \mathbb{E}\{\text{tr}(\text{hol}(L_2))\} \\ &+ \text{Cov}\{\text{tr}(\text{hol}(L_1)), \text{tr}(\text{hol}(L_2))\} \end{aligned} \tag{13}$$

and

$$\begin{aligned}
& \left(\frac{\partial}{\partial t_1} - \frac{\partial}{\partial t_2} + \frac{\partial}{\partial t_3} - \frac{\partial}{\partial t_4} \right) \text{Var} \{ \text{tr}(\text{hol}(L)) \} \\
&= 2 \text{Re} \left[\text{Cov} \left\{ \text{tr}(\text{hol}(L_1)) \text{tr}(\text{hol}(L_2)), \overline{\text{tr}(\text{hol}(L))} \right\} \right] \\
& - \frac{1}{N^2} \mathbb{E} \{ \text{tr}(\text{hol}(L_3)) \} - \frac{1}{N^2} \mathbb{E} \{ \text{tr}(\text{hol}(L_4)) \}, \tag{14}
\end{aligned}$$

where L_3 and L_4 are the composite curves $L_3 = L_1 L_2 L_1^{-1} L_2^{-1}$ and $L_4 = L_2 L_1 L_2^{-1} L_1^{-1}$. Here Cov denotes the covariance, defined as $\text{Cov}\{f, g\} = \mathbb{E}\{fg\} - \mathbb{E}\{f\}\mathbb{E}\{g\}$.

Proof. Equation (13) is simply the Makeenko–Migdal equation (3) rewritten using the definition of the covariance. To establish (14) we need to use the abstract Makeenko–Migdal equation established in Theorem 2 of [DGHK]. (This result generalizes the abstract Makeenko–Migdal equation formulated and proved for the plane case by Lévy in [Lév2, Proposition 9.1.3].) Following the argument in Section 2.3 of [DHK2], we express the loop L as

$$L = e_1 A e_4^{-1} e_2 B e_3^{-1},$$

where A and B are words in edges other than e_1, \dots, e_4 . Then $L_1 = e_1 A e_4^{-1}$ and $L_2 = e_2 B e_3^{-1}$. If a_1, \dots, a_4 are the edge variables corresponding to e_1, \dots, e_4 , we then have (following the convention that parallel transport is order reversing)

$$\text{hol}(L) = a_3^{-1} \beta a_2 a_4^{-1} \alpha a_1,$$

where α and β are words in the edge variables other than a_1, \dots, a_4 .

Now, the abstract Makeenko–Migdal equation reads

$$\left(\frac{\partial}{\partial t_1} - \frac{\partial}{\partial t_2} + \frac{\partial}{\partial t_3} - \frac{\partial}{\partial t_4} \right) \mathbb{E}\{f\} = -\mathbb{E}\{\nabla^{a_1} \cdot \nabla^{a_2} f\}, \tag{15}$$

whenever f has “extended gauge invariance” at the vertex v . When the edges e_1, \dots, e_4 are distinct, extended gauge invariance at v means that

$$f(a_1 x, a_2, a_3 x, a_4, \mathbf{b}) = f(a_1, a_2 x, a_3, a_4 x, \mathbf{b}) = f(a_1, a_2, a_3, a_4, \mathbf{b})$$

for all $x \in K$, where \mathbf{b} is the collection of edge variables other than a_1, \dots, a_4 . (See Section 4 of [DHK2] for a discussion of extended gauge invariance when the edges are not distinct.)

Let us apply (15) to the function $f = |g|^2$, where

$$g(a_1, a_2, a_3, a_4, \mathbf{b}) = \text{tr}(\text{hol}(L)) = \text{tr}(a_3^{-1} \beta a_2 a_4^{-1} \alpha a_1).$$

We also use the notation

$$\begin{aligned}
g_1(a_1, a_2, a_3, a_4, \mathbf{b}) &= \text{tr}(\text{hol}(L_1)) = \text{tr}(a_4^{-1} \alpha a_1) \\
g_2(a_1, a_2, a_3, a_4, \mathbf{b}) &= \text{tr}(\text{hol}(L_2)) = \text{tr}(a_3^{-1} \beta a_2).
\end{aligned}$$

We then note that

$$\begin{aligned}\nabla^{a_1} \cdot \nabla^{a_2} f &= (\nabla^{a_1} \cdot \nabla^{a_2} g) \bar{g} + g(\nabla^{a_1} \cdot \nabla^{a_2} \bar{g}) \\ &\quad + (\nabla^{a_1} g) \cdot (\nabla^{a_2} \bar{g}) + (\nabla^{a_2} g) \cdot (\nabla^{a_1} \bar{g}).\end{aligned}$$

Now, as verified in [DHK2, Eqs. (2.13)–(2.15)], we have $\nabla^{a_1} \cdot \nabla^{a_2} g = -g_1 g_2$. Meanwhile, using $\overline{\operatorname{tr}(U)} = \operatorname{tr}(U^{-1})$ for $U \in U(N)$ and computing as in the second example in [DHK2, Section 2.5], we have that

$$\begin{aligned}(\nabla^{a_1} g) \cdot (\nabla^{a_2} \bar{g}) &= \frac{1}{N^2} \operatorname{tr}(\operatorname{hol}(L_1 L_2 L_1^{-1} L_2^{-1})) \\ (\nabla^{a_2} g) \cdot (\nabla^{a_1} \bar{g}) &= \frac{1}{N^2} \operatorname{tr}(\operatorname{hol}(L_2 L_1 L_2^{-1} L_1^{-1})).\end{aligned}$$

Thus,

$$\begin{aligned}&\left(\frac{\partial}{\partial t_1} - \frac{\partial}{\partial t_2} + \frac{\partial}{\partial t_3} - \frac{\partial}{\partial t_4} \right) \mathbb{E}\{|g|^2\} \\ &= \mathbb{E}\{g_1 g_2 \bar{g}\} + \mathbb{E}\{g \overline{g_1 g_2}\} - \mathbb{E}\{\operatorname{tr}(\operatorname{hol}(L_3))\} - \mathbb{E}\{\operatorname{tr}(\operatorname{hol}(L_4))\}.\end{aligned}\quad (16)$$

Meanwhile, by the ordinary Makeenko–Migdal equation (3), we have

$$\begin{aligned}&\left(\frac{\partial}{\partial t_1} - \frac{\partial}{\partial t_2} + \frac{\partial}{\partial t_3} - \frac{\partial}{\partial t_4} \right) |\mathbb{E}\{g\}|^2 \\ &= \mathbb{E}\{g_1 g_2\} \mathbb{E}\{\bar{g}\} + \mathbb{E}\{g\} \mathbb{E}\{\overline{g_1 g_2}\}.\end{aligned}\quad (17)$$

Subtracting (17) from (16) gives the claimed result. ■

2.2 Variance estimates

For a complex-valued random variable X , we define the **variance** of X by

$$\operatorname{Var}(X) := \mathbb{E}\{|X - \mathbb{E}\{X\}|^2\} = \mathbb{E}\{|X|^2\} - |\mathbb{E}\{X\}|^2.$$

In particular,

$$\operatorname{Var}(X) \leq \mathbb{E}\{|X|^2\}.\quad (18)$$

We then define the **standard deviation** of X , denoted σ_X , by $\sigma_X = \sqrt{\operatorname{Var}(X)}$.

We observe that for any two random variables X and Y , we have

$$\sigma_{X+Y} \leq \sigma_X + \sigma_Y,\quad (19)$$

and similarly for any number of random variables. (It is harmless to assume that the expectation values of X and Y —and therefore $X + Y$ —are zero, in which case (19) is the triangle inequality for the L^2 norm.) We also consider the covariance of two random variables, defined as

$$\operatorname{Cov}\{X, Y\} = \mathbb{E}\{(X - \mathbb{E}\{X\})(Y - \mathbb{E}\{Y\})\}\quad (20)$$

$$= \mathbb{E}\{XY\} - \mathbb{E}\{X\} \mathbb{E}\{Y\}.\quad (21)$$

(For our purposes, it is convenient not to put a complex conjugate into this definition; the reader should note that $\text{Cov}\{X, X\} \neq \text{Var}(X)$.) We record the elementary inequality

$$|\text{Cov}\{X, Y\}| \leq \sigma_X \sigma_Y, \quad (22)$$

which follows from (20) and the Cauchy–Schwarz inequality.

We now establish a simple estimate on the standard deviation of the product of two bounded random variables.

Proposition 6 *Suppose X and Y are two complex-valued random variables satisfying $|X| \leq 1$ and $|Y| \leq 1$. Then*

$$\sqrt{\sigma_{XY}} \leq \sqrt{\sigma_X} + \sqrt{\sigma_Y}.$$

Proof. For simplicity of notation, let $\bar{X} = \mathbb{E}\{X\}$ and let $\dot{X} = X - \bar{X}$. Thus, $\text{Var}(X) = \mathbb{E}\{|\dot{X}|^2\}$. Then since $|X| \leq 1$, we have $|\bar{X}| \leq 1$ and $|\dot{X}| \leq 2$. Then by (18) and the Cauchy–Schwarz inequality, we have

$$\begin{aligned} \text{Var}(\dot{X}\dot{Y}) &\leq \mathbb{E}\{|\dot{X}|^2|\dot{Y}|^2\} \\ &\leq \sqrt{\mathbb{E}\{|\dot{X}|^4\}\mathbb{E}\{|\dot{Y}|^4\}} \\ &\leq 4\sigma_X\sigma_Y, \end{aligned}$$

since $|\dot{X}|^4 = |\dot{X}|^2|\dot{X}|^2 \leq 4|\dot{X}|^2$ and similarly for $|\dot{Y}|^4$. Now,

$$\begin{aligned} \text{Var}(XY) &= \text{Var}(\dot{X}\dot{Y} + \bar{X}\dot{Y} + \dot{X}\bar{Y} + \bar{X}\bar{Y}) \\ &= \text{Var}(\dot{X}\dot{Y} + \bar{X}\dot{Y} + \dot{X}\bar{Y}) \end{aligned}$$

because adding a constant does not change the variance. Thus, by (19),

$$\begin{aligned} \sigma_{XY} &\leq \sigma_{\dot{X}\dot{Y}} + |\bar{X}|\sigma_Y + |\bar{Y}|\sigma_X \\ &\leq 2\sqrt{\sigma_X\sigma_Y} + \sigma_Y + \sigma_X \\ &= (\sqrt{\sigma_X} + \sqrt{\sigma_Y})^2, \end{aligned}$$

which reduces to the claimed formula. ■

3 Examples

Before developing a general method for analyzing a general loop in S^2 with simple crossings, we consider two illustrative examples, the same two that are considered in [DaK].

3.1 The figure eight

Although we consider loops on S^2 , we can draw them as a loops on the plane by picking a face and placing a puncture in that face, so that what is left of S^2 is




Figure 4: Two views of a loop on S^2

identifiable with \mathbb{R}^2 . We need to keep in mind, however, that the “unbounded” face in such a drawing is actually a bounded face (with finite area) on S^2 . Furthermore, by placing the puncture in different faces, the same loop on S^2 can give inequivalent loops on \mathbb{R}^2 . As a simple example, consider the loop in Figure 4, which we refer to as the figure eight. Figure 4 gives two different views of this loop coming from puncturing two different faces.

We write the holonomy around the figure eight as

$$\text{hol}_L(a, b, c)$$

to indicate the dependence of the Wilson loop functional on the areas. In this case, the loops L_1 and L_2 occurring on the right-hand side of the Makeenko–Migdal equation are both simple closed curves. The loop L_1 (the outer loop on the left-hand side of Figure 4, which is the lower loop on the right-hand side of the figure) encloses areas $a + b$ and c . The loop L_2 (the inner loop on the left-hand side of the figure, which is the upper loop on the right-hand side of the figure) encloses areas $b + c$ and a .

Theorem 7 *The limit*

$$W_L(a, b, c) := \lim_{N \rightarrow \infty} \mathbb{E} \{ \text{tr}(\text{hol}_L(a, b, c)) \}$$

exists and satisfies the large- N Makeenko–Migdal equation in the form

$$\left. \frac{d}{dt} W_L(a - t, b + 2t, c - t) \right|_{t=0} = W_1(a + b, c) W_1(a, b + c),$$

where W_1 is as in Notation 2. Furthermore, we have

$$\lim_{N \rightarrow \infty} \text{Var} \{ \text{tr}(\text{hol}_L(a, b, c)) \} = 0.$$

See (26) for a formula for $W_L(a, b, c)$ in terms of the function $W_1(\cdot, \cdot)$. If the partial derivatives of $W_L(a, b, c)$ with respect a , b , and c exist and are continuous, it follows from the chain rule that

$$\left. \frac{d}{dt} W_L(a - t, b + 2t, c - t) \right|_{t=0} = \left(2 \frac{\partial}{\partial b} - \frac{\partial}{\partial a} - \frac{\partial}{\partial c} \right) W_L(a, b, c).$$

The following proof, however, does not establish the existence or continuity of the partial derivatives of $W_L(a, b, c)$.

Proof. We denote the holonomies for the two loops L_1 and L_2 as $\text{hol}_{L_1}(a+b, c)$ and $\text{hol}_{L_2}(a, b+c)$. The face labeled as F_1 should be the one bounded by the two outgoing edges of L at v , which is the face with area b . Then F_3 coincides with F_1 , while F_2 and F_4 are the faces with areas a and c (in either order). We assume $c \geq a$, with the case $c < a$ being entirely similar.

Proposition 5 takes the form

$$\begin{aligned} & \frac{d}{dt} \mathbb{E} \{ \text{tr} (\text{hol}_L(a-t, b+2t, c-t)) \} \\ &= \left(2 \frac{\partial}{\partial b} - \frac{\partial}{\partial a} - \frac{\partial}{\partial c} \right) \mathbb{E} \{ \text{tr} (\text{hol}_L(a-t, b+2t, c-t)) \} \\ &= \mathbb{E} \{ \text{tr} (\text{hol}_{L_1}(a+b+t, c-t)) \} \mathbb{E} \{ \text{tr} (\text{hol}_{L_2}(a-t, b+c+t)) \} \\ &+ \text{Cov} \{ \text{tr} (\text{hol}_{L_1}(a+b+t, c-t)), \text{tr} (\text{hol}_{L_2}(a-t, b+c+t)) \}. \quad (23) \end{aligned}$$

Let us denote $\mathbb{E} \{ \text{tr} (\text{hol}_L(a-t, b+2t, c-t)) \}$ by $F(t)$ (with a, b , and c fixed). We then write $F(0) = F(a-\varepsilon) - \int_0^{a-\varepsilon} F'(t) dt$; that is,

$$\begin{aligned} & \mathbb{E} \{ \text{tr} (\text{hol}_L(a, b, c)) \} \\ &= \mathbb{E} \{ \text{tr} (\text{hol}_L(\varepsilon, 2a+b-2\varepsilon, c-a+\varepsilon)) \} \\ &- \int_0^{a-\varepsilon} \mathbb{E} \{ \text{tr} (\text{hol}_{L_1}(a+b+t, c-t)) \} \mathbb{E} \{ \text{tr} (\text{hol}_{L_2}(a-t, b+c+t)) \} dt \\ &- \int_0^{a-\varepsilon} \text{Cov} \{ \text{tr} (\text{hol}_{L_1}(a+b+t, c-t)), \text{tr} (\text{hol}_{L_2}(a-t, b+c+t)) \} dt. \quad (24) \end{aligned}$$

Now, it should be clear geometrically that if we let the area a in the figure eight tend to zero, the result is a simple closed curve. That is to say, we expect that

$$\begin{aligned} & \lim_{\varepsilon \rightarrow 0} \mathbb{E} \{ \text{tr} (\text{hol}_L(\varepsilon, 2a+b-2\varepsilon, c-a+\varepsilon)) \} \\ &= \mathbb{E} \{ \text{tr} (\text{hol}_{L_0}(2a+b, c-a)) \}, \quad (25) \end{aligned}$$

where $L_0(\alpha, \beta)$ is a simple closed curve on S^2 enclosing areas α and β . Analytically, (25) follows easily from Sengupta's formula, using that $\rho_a(\cdot)$ converges to a δ -measure on K as a tends to zero. Thus, letting ε tend to zero, we obtain

$$\begin{aligned} & \mathbb{E} \{ \text{tr} (\text{hol}_L(a, b, c)) \} \\ &= \mathbb{E} \{ \text{tr} (\text{hol}_{L_0}(2a+b, c-a)) \} \\ &- \int_0^a \mathbb{E} \{ \text{tr} (\text{hol}_{L_1}(a+b+t, c-t)) \} \mathbb{E} \{ \text{tr} (\text{hol}_{L_2}(a-t, b+c+t)) \} dt \\ &- \int_0^a \text{Cov} \{ \text{tr} (\text{hol}_{L_1}(a+b+t, c-t)), \text{tr} (\text{hol}_{L_2}(a-t, b+c+t)) \} dt. \quad (26) \end{aligned}$$




Figure 5: Trefoil with one lobe shrunk to area zero

Note that all holonomies on the right-hand side of (26) are of simple closed curves. If we use (6) in Theorem 1 together with the inequality (22) and dominated convergence, we find that the last term in (26) tends to zero as N tends to infinity. Then using (5) in Theorem 1 along with dominated convergence, we may let $N \rightarrow \infty$ to obtain

$$\begin{aligned} \lim_{N \rightarrow \infty} \mathbb{E} \{ \text{tr}(\text{hol}_L(a, b, c)) \} &= W_1(2a + b, c - a) \\ &\quad - \int_0^a W_1(a + b + t, c - t) W_1(a - t, b + c + t) dt, \end{aligned} \tag{27}$$

for $c \geq a$, where W_1 is as in Notation 2. (Note that the normalized trace defined in (2) satisfies $|\text{tr}(U)| \leq 1$ for all $U \in U(N)$, so that dominated convergence applies in both integrals in (26).) This result establishes the first claim in the theorem.

If we now subtract the value of (27) at $(a - s, b + 2s, c - s)$ and the value at (a, b, c) , we obtain

$$\begin{aligned} &W_L(a - s, b + 2s, c - s) - W_L(a, b, c) \\ &= \int_0^s W_1(a + b + t, c - t) W_1(a - t, b + c + t) dt. \end{aligned}$$

Dividing this relation by s and letting s tend to zero gives

$$\left. \frac{\partial}{\partial s} W_L(a - s, b + 2s, c - s) \right|_{s=0} = W_1(a + b + t, c - t) W_1(a - t, b + c + t)$$

by the continuity of W_1 . This relation is the desired large- N Makeenko–Migdal equation.




Figure 6: This loop is a perturbation of the loop on the right-hand side of Figure 5

Meanwhile, by the second relation in Proposition 5, we have

$$\begin{aligned}
\text{Var} \{ \text{tr}(\text{hol}_L(a, b, c)) \} &= \text{Var} \{ \text{tr}(\text{hol}_{L_0}(b + 2a, c - a)) \} \\
&- 2 \int_0^a \text{Cov} \{ \text{tr}(\text{hol}(L_1)) \text{tr}(\text{hol}(L_2)), \text{tr}(\text{hol}(L)) \} dt \\
&- \frac{2}{N^2} \int_0^a \mathbb{E} \{ \text{tr}(\text{hol}(L_3)) \} dt.
\end{aligned} \tag{28}$$

The first term on the right-hand side tends to zero as N tends to infinity, by Theorem 1. In the second term on the right-hand side, L_1 and L_2 are simple closed curves. Furthermore, the normalized trace satisfies $|\text{tr}(U)| \leq 1$ for all $U \in U(N)$. Thus, using (22) and Proposition 6 together with Theorem 1, we see that the second term on the right-hand side of (28) tends to zero. Finally, since $|\text{tr}(U)| \leq 1$, the last term on the right-hand side manifestly goes to zero. ■

3.2 The trefoil

In this section, we briefly outline an analysis of the trefoil loop by a method similar to the one in the previous subsection. Later we will develop a systematic method for analyzing any loop; this will provide an alternative analysis of the trefoil. We label the areas of the faces as in Figure 5, where we may assume without loss of generality that $c \leq b$. We now perform a Makeenko–Migdal variation at the vertex in the top middle of the figure. In this case, the loops L_1 and L_2 turn out to be simple closed curves.

Let us denote by L' the loop on the right-hand side of Figure 5. Then L' is the limit as ε tends to zero of the loops L''_ε in Figure 6, where ε denotes the distance between the two nearby arcs of the loop. Specifically, in Figure 6, we let ε tend to zero, while keeping all of the areas of the faces fixed to the values indicated, using the continuity properties of the Wilson loop functional developed in [Lév1, Theorem 2.58]. But, $\mathbb{E} \{ \text{tr}(\text{hol}(L''_\varepsilon)) \}$ is independent of ε and we conclude that $\mathbb{E} \{ \text{tr}(\text{hol}(L')) \} = \mathbb{E} \{ \text{tr}(\text{hol}(L''_\varepsilon)) \}$. But since the loop L''_ε




Figure 7: A more complicated loop

is of the type analyzed in Section 3, we may already know the large- N behavior of $\mathbb{E} \{\text{tr}(\text{hol}(L'))\}$. The argument then proceeds much as in Section 3.1; since we will develop later a systematic method for analyzing arbitrary loops, we omit the details of this analysis.

Other examples are not quite so easy to simplify by using the Makeenko–Migdal equation at a single vertex. In the loop in Figure 7, for example, it is not evident how shrinking any one of the faces to zero simplifies the problem.

4 Analysis of a general loop

4.1 The strategy

Given an arbitrary loop L traced out on a graph in S^2 with simple crossings, we will consider a linear combination of Makeenko–Migdal variations of the areas over all the vertices of L . We will show that it is possible to make such a variation, depending on a parameter t , in such a way that as t tends to 1, *all but two* of the areas of the faces tend to zero. Thus, in the $t \rightarrow 1$ limit, we effectively have a loop with only two faces. Indeed, we will show that the $t \rightarrow 1$ limit of the Wilson loop functional is the Wilson loop functional for a loop L^n that winds n times around a simple closed curve. Here n is an integer determined by the topology of the original loop L . (A similar but not identical procedure is used in [DN]. See Section 4.4 for a comparison.)

Consider, for example, the trefoil loop of Section 3.2. If we vary the areas by the amounts shown in Figure 8, the net effect on the areas is:

$$\begin{aligned} b &\mapsto b - tb & a &\mapsto a + t(b + c + d)/2 \\ c &\mapsto c - tc & e &\mapsto e + t(b + c + d)/2 . \\ d &\mapsto d - td \end{aligned}$$




Figure 8: We can shrink the areas of the three lobes of the trefoil to zero, while increasing the areas of the other two faces

Thus, as t varies from 0 to 1, the areas b , c , and d shrink simultaneously to zero, while the areas of the two remaining faces increase. The limiting curve is shown in Figure 9. If the areas of the three lobes are zero, the curve becomes a circle traversed twice.

Let $L(t)$ denote the trefoil with areas varying as above. If we differentiate $\mathbb{E}\{\text{tr}(\text{hol}(L(t)))\}$ with respect to t , then by the chain rule, we will get a linear combination of terms of the form

$$\mathbb{E}\{\text{tr}(\text{hol}(L_{1,j}(t)))\} \mathbb{E}\{\text{tr}(\text{hol}(L_{2,j}(t)))\},$$

where $L_{1,j}(t)$ and $L_{2,j}(t)$ represent the loop $L(t)$ cut at the j th vertex of the trefoil ($j = 1, 2, 3$), along with some covariance terms. Each $L_{1,j}(t)$ and $L_{2,j}(t)$ is actually a simple closed curve, but in any case, these curves have fewer crossings than the original trefoil. It is then a straightforward matter to let N tend to infinity to get the limiting Wilson loop functional, and similarly for the variance.

For an arbitrary loop L with k crossings, we will show that we can deform L into a loop L_n that winds n times around a simple closed curve. The variation of the Wilson loop functional along this path will be a linear combination of products of Wilson loop functionals for curves with at most $k - 1$ crossings. In an inductive argument then, it remains only to analyze L_n , which we do by another induction, this time reducing L_n to L_{n-1} , and so on.

4.2 Winding numbers

We consider S^2 with a fixed orientation. We then consider a loop L traced out on a graph in S^2 and having only simple crossings. We consider the faces of L , that is, the connected components of the complement of L in S^2 . If we




Figure 9: The trefoil with the lobes shrunk to area zero

pick a face F_0 , we can puncture F_0 , thus turning S^2 topologically into \mathbb{R}^2 . The orientation on S^2 gives an orientation on \mathbb{R}^2 . Thus, for each face F , we may speak about the winding number of L around F . Since this winding number depends on which face F_0 we puncture, we denote it thus:

$$w_{F_0}(F),$$

so that $w_{F_0}(F_0) = 0$. It is important to understand how the winding number changes if the location of the puncture changes.

Proposition 8 *If F_0 , F'_0 , and F are faces, then*

$$w_{F_0}(F) - w_{F'_0}(F) = w_{F_0}(F'_0).$$

In particular, the difference between $w_{F_0}(F)$ and $w_{F'_0}(F)$ does not depend on F .

In particular, if we change the location of the puncture, all the winding numbers change *by the same amount*.

Proof. Let us fix points x , y , and z in F_0 , F'_0 , and F , respectively. Let us put our puncture initially in x , regarding $S^2 \setminus \{x\}$ as the plane. Let us then assume that y is at the origin and z is at the point $(2,0)$. We may then regard L as an element of $\pi_1(\mathbb{R}^2 \setminus \{y, z\})$ with base point at $(1,0)$, which is a free group on two generators e_1 and e_2 . These generators may be identified with circles of radius one centered at y and z , respectively, traversed in the counter-clockwise direction. Then $w_{F_0}(F)$ is the number of times L winds around z , which is the number of occurrences of the generator e_2 in the representation of L as a word in e_1 and e_2 .

Suppose we now shift our puncture from x to y . This shift amounts to composing L with the inversion map in the complex plane, $\mathbb{C} = \mathbb{R}^2$, that is, the map $z \mapsto 1/z$. After this process, the generator e_1 traverses the unit circle in the opposite direction, while the generator e_2 is now inside the unit circle (Figure 10). Thus, $w_{F'_0}(F)$ is the number of occurrences of e_2 minus the number of occurrences of e_1 , giving the claimed result. ■




Figure 10: Transformation of the winding numbers under a change of puncture

4.3 The span of the Makeenko–Migdal vectors

Let L be a loop traced out on a graph in S^2 and having only simple crossings. We consider vectors assigning real numbers to the faces of L . For each vertex v of L , we define the **Makeenko–Migdal vector** associated to v , denoted MM_v , as

$$\text{MM}_v = \sum_{i=1}^4 (-1)^{i+1} \delta_{F_i},$$

where F_1, \dots, F_4 are the four faces surrounding v . If, for example, F_1, \dots, F_4 are distinct, then MM_v is the vector that assigns the numbers $1, -1, 1, -1$ to F_1, \dots, F_4 , respectively, and zero to all other faces.

Theorem 9 (T. Lévy) *Fix a face F_0 of L and let u be a vector assigning a real number to each face of L . Then u belongs to the span of the Makeenko–Migdal vectors if and only if (1) u is orthogonal to the constant vector $\mathbf{1} := (1, 1, \dots, 1)$, and (2) u is orthogonal to the winding-number vector $w_{F_0}(\cdot)$.*

This result is the $r = 1$ case of Lemma 9.4.3 in [Lév2]. Note that by Proposition 8, the winding number vector $w_{F'_0}$ associated to some other face F'_0 differs from w_{F_0} by a multiple of the constant vector $\mathbf{1}$. Thus, if u is orthogonal to $\mathbf{1}$, then u is orthogonal to $w_{F_0}(\cdot)$ if and only if u is orthogonal to $w_{F'_0}(\cdot)$. Thus, the condition in the theorem is independent of the choice of F_0 .

4.4 Shrinking all but two of the faces

Let f denote the number of faces of L . We now choose one face arbitrarily and denote it by F_0 . We will show that there is another face F_1 such that we can perform a Makeenko–Migdal variation of the areas in which the areas of the faces F_2, \dots, F_{f-1} shrink simultaneously to zero, while the area of F_0 increases or remains the same. In the generic case, the area of F_1 also remains positive, while in a certain borderline case, the area of F_1 tends to zero as well.

The freedom we have to choose one of the “unshrunk” faces arbitrarily will be essential in applications to the plane and to arbitrary surfaces. This freedom is a difference between our approach and the one used in [DN], where the two unshrunk faces in Section 4.5 are chosen to have maximal and minimal winding numbers.

Proposition 10 *Assume L has at least three faces. Choose one face F_0 arbitrarily and let all winding numbers be computed relative to a puncture in F_0 , so that $w(F_0) = 0$. Let $\mathbf{a} = (a_0, a_1, \dots, a_{f-1})$ denote the vector of areas of the faces and let $\mathbf{w} = (0, w_1, \dots, w_{f-1})$ be the vector of winding numbers. Suppose $\mathbf{a} \cdot \mathbf{w} \neq 0$ and adjust the labeling of F_1, \dots, F_{f-1} so that w_1 is maximal among the winding numbers if $\mathbf{a} \cdot \mathbf{w} > 0$ and w_1 is minimal if $\mathbf{a} \cdot \mathbf{w} < 0$. Then there exists \mathbf{b} in the span of the Makeenko–Migdal vectors such that (1) all the entries of $\mathbf{a} + t\mathbf{b}$ are non-negative for $0 \leq t \leq 1$ and (2) $\mathbf{a}' := \mathbf{a} + \mathbf{b}$ has the form $\mathbf{a}' = (a'_0, a'_1, 0, \dots, 0)$ with $a'_0 \geq a_0$ and $a'_1 > 0$.*

Meanwhile, if $\mathbf{a} \cdot \mathbf{w} = 0$, there exists \mathbf{b} in the span of the Makeenko–Migdal vectors such that (1) all the entries of $\mathbf{a} + t\mathbf{b}$ are non-negative for $0 \leq t \leq 1$ and (2) $\mathbf{a}' := \mathbf{a} + \mathbf{b}$ has the form $\mathbf{a}' = (a'_0, 0, \dots, 0)$ with $a'_0 > 0$.

The proposition says, briefly, that we can shrink a_2, \dots, a_{f-1} to zero without decreasing a_0 and while keeping a_1 non-negative.

In the case of the trefoil loop, for example, suppose we take F_0 to be the “unbounded” face in Figure 8 and we orient the loop in the counter-clockwise direction. Then the winding numbers are 1 for the three lobes of the trefoil and 2 for the central region. Since all winding numbers are positive in this case, we always have $\mathbf{a} \cdot \mathbf{w} > 0$, in which case we take F_1 to be the central region. Figures 8 and 9 then illustrate Proposition 10.

Proof. Assume first that $\mathbf{a} \cdot \mathbf{w} > 0$, in which case, the maximal winding number w_1 must be positive. In light of Theorem 9, two vectors \mathbf{a} and \mathbf{a}' differ by a vector in the span of the Makeenko–Migdal vectors if \mathbf{a} and \mathbf{a}' have the same inner products with the constant vector $\mathbf{1}$ and the winding number vector \mathbf{w} . To achieve these conditions, we first set a'_1 equal to $\mathbf{a} \cdot \mathbf{w} / w_1$ and a'_2, \dots, a'_{f-1} equal to 0. Since $w_0 = 0$, we then have $\mathbf{a}' \cdot \mathbf{w} = \mathbf{a} \cdot \mathbf{w}$, regardless of the value of a'_0 . Next, we choose $a'_0 = -a'_1 + a_0 + a_1 + \dots + a_{f-1}$ to achieve the condition $\mathbf{a}' \cdot \mathbf{1} = \mathbf{a} \cdot \mathbf{1}$. Now, $\mathbf{a}' \cdot \mathbf{w} = a_1 w_1 = \mathbf{a} \cdot \mathbf{w}$. Then since w_1 is maximal, we have

$$\begin{aligned} a'_1 w_1 &= a_1 w_1 + \dots + a_{f-1} w_{f-1} \\ &\leq w_1 (a_1 + \dots + a_{f-1}) \end{aligned}$$

Thus, $a'_1 \leq a_1 + \dots + a_{f-1}$, which shows that $a'_0 \geq a_0$. (In particular, $a'_0 > 0$.) Since both \mathbf{a} and \mathbf{a}' have non-negative entries, so does every point along the line segment joining them. The analysis of the case $\mathbf{a} \cdot \mathbf{w} < 0$ is similar.

Finally, if $\mathbf{a} \cdot \mathbf{w} = 0$, we may set $a'_0 = a_0 + a_1 + \dots + a_{f-1}$ and all other entries of \mathbf{a}' equal to zero. In that case, $\mathbf{a}' \cdot \mathbf{w} = \mathbf{a} \cdot \mathbf{w} = 0$ and $\mathbf{a}' \cdot \mathbf{1} = a'_0 = \mathbf{a} \cdot \mathbf{1}$, showing that \mathbf{a}' differs from \mathbf{a} by a vector in the span of the Makeenko–Migdal vectors. ■

4.5 Analyzing the limiting case

As a consequence of Proposition 10, we may start with an arbitrary loop L and perform a linear combination of Makeenko–Migdal variations at each vertex, obtaining a family $L(t)$, $0 \leq t < 1$, of loops with the same topological type with

all but two of the areas shrinking to zero as $t \rightarrow 1$. We now analyze the behavior of the Wilson loop functional in the limit $t \rightarrow 1$.

We will analyze the limit “analytically,” using Sengupta’s formula for the finite- N case. (Recall Remark 4.) For this result, the structure group can be an arbitrary connected compact Lie group K .

Theorem 11 *Let L be a loop traced out on a graph in S^2 and having only simple crossings. Denote the number of faces of L by f and label the faces as $F_0, F_1, F_2, \dots, F_{f-1}$. Suppose we vary the areas of the faces as a function of a parameter $t \in [0, 1)$ in such a way that as $t \rightarrow 1$, the areas of F_2, \dots, F_{f-1} tend to zero, while the areas of F_0 and F_1 approach positive real numbers a and c , respectively. Then*

$$\lim_{t \rightarrow 1} \mathbb{E}\{\text{tr}(\text{hol}(L))\} = \frac{1}{Z} \int_K \text{tr}(h^n) \rho_a(h) \rho_c(h) dh, \quad (29)$$

where $n = w_{F_0}(F_1)$ is the winding number of L around F_1 , relative to a puncture in F_0 , and Z is a normalization constant. Meanwhile, if the area of F_1 also tends to zero in the $t \rightarrow 1$ limit, then

$$\lim_{t \rightarrow 1} \mathbb{E}\{\text{tr}(\text{hol}(L))\} = 1.$$

The integral on the right-hand side of (29) is just the Wilson loop functional for a loop L_n that winds n times around a simple closed curve enclosing areas a and c . Note that the winding number of L_n around F_0 is the same as the winding number of the original loop L around F_0 . Note also that by Theorem 9, the value of $\mathbf{a} \cdot \mathbf{w}$ for the loop L_t is independent of t for $t \in [0, 1)$. Since $w_0 = 0$ and a_2 through a_{f-1} are tending to zero, this means that a_1 must be tending to the value $\mathbf{a} \cdot \mathbf{w}/n$, where $n = w_1$ is the winding number of L around F_1 . It then follows that the limiting loop L_n has the same value of $\mathbf{a} \cdot \mathbf{w}$, even though L_n has a different topological type from L .

Proof. Let \mathbb{G} be a minimal oriented graph on which L can be traced. We think of \mathbb{G} as a graph in the plane, by placing a puncture into F_0 . If e denotes the number of edges of \mathbb{G} , then we may consider two different measures on K^e : the Yang–Mills measure $\mu_{\text{plane}}^{\mathbb{G}}$ for \mathbb{G} viewed as a graph in the plane and the Yang–Mills measure $\mu_{\text{sphere}}^{\mathbb{G}}$ for \mathbb{G} viewed as a graph in the sphere. By comparing Sengupta’s formula [Sen1, Sect. 5] in the sphere case to Driver’s formula [Dr, Theorem 6.4] in the plane case, we see that

$$d\mu_{\text{sphere}}^{\mathbb{G}}(\mathbf{x}) = \frac{1}{Z} \rho_{|F_0|}(\text{hol}_{F_0}(\mathbf{x})) d\mu_{\text{plane}}^{\mathbb{G}}(\mathbf{x}),$$

where $\mathbf{x} \in K^e$ is the collection of edge variables, $|F_0|$ is the area of F_0 as a face in S^2 , and hol_{F_0} is the product of edge variables around the boundary of F_0 . Here Z is a normalization constant.

We may rewrite both of the Yang–Mills measures using “loop variables” as follows. Let us fix a vertex v and a spanning tree T for \mathbb{G} . Then in Section 5.3




Figure 11: The example loop L (left) decomposes as $L_1L_2L_3L_1^{-1}L_4^{-1}L_3^{-1}$

of [Lév2], Lévy gives a procedure associating to the faces F_1, \dots, F_{f-1} certain loops L_1, \dots, L_{f-1} in \mathbb{G} that constitute free generators for $\pi_1(\mathbb{G})$, based at v . (Note that there is no generator associated to the face F_0 .) Each L_i is a word in the edges of \mathbb{G} . We may then associate to each of the faces F_i , $i = 1, \dots, f-1$, of \mathbb{G} a loop *variable*, which is the product of edge variables in L_i (in the reverse order, since parallel transport reverses order). The map sending the collection of edge variables to the collection of loop variables defines a map $\Gamma : K^e \rightarrow K^{f-1}$, where e is the number of edges of \mathbb{G} .

According to Proposition 5.3.3 of [Lév2], the loop variables are independent heat-kernel distributed random variables with respect to $\mu_{\text{plane}}^{\mathbb{G}}$. (See also Proposition 12.7 of [Ga].) That is to say, the push-forward of $\mu_{\text{plane}}^{\mathbb{G}}$ under Γ is just the product of heat kernels, at times equal to the areas of the faces. Although there is no generator associated to the face F_0 , the L_1, \dots, L_{f-1} generate $\pi_1(\mathbb{G})$. Suppose, therefore, that L_0 is a loop that starts at v , travels along a path P to a vertex in ∂F_0 , then around the boundary of F_0 , and then back to v along P^{-1} . Then L_0 is expressible as a word in L_1, \dots, L_{f-1} and therefore $\text{hol}_{F_0}(\mathbf{x})$ is expressible as a word in the loop variables: $\text{hol}_{F_0}(\mathbf{x}) = w_0(\Gamma(\mathbf{x}))$ for some function w_0 on K^{f-1} . (Although $\text{hol}_{F_0}(\mathbf{x})$ depends on the choice of v and P , the invariance of the heat kernel under conjugation guarantees that $\rho_{|F_0|}(\text{hol}_{F_0}(\mathbf{x}))$ is well defined.) It then follows from the measure-theoretic change of variables theorem that the push-forward of $\mu_{\text{sphere}}^{\mathbb{G}}$ is given by

$$\begin{aligned} & d\Gamma_*(\mu_{\text{sphere}}^{\mathbb{G}})(h_1, \dots, h_{f-1}) \\ &= \frac{1}{Z} \rho_{|F_0|}(w_0(h_1, \dots, h_{f-1})) \left(\prod_{i=1}^{f-1} \rho_{|F_i|}(h_i) \right) dh_1 \dots dh_{f-1}. \end{aligned}$$

Meanwhile, the loop L (whose Wilson loop functional we are considering) is also expressible as a word w_1 in the generators L_1, \dots, L_{f-1} . (See Figure 11.)

Thus,

$$\begin{aligned} \int_{K^e} \text{tr}(\text{hol}(L)) d\mu_{\text{sphere}}^{\mathbb{G}} &= \frac{1}{Z} \int_{K^{f-1}} \text{tr}(w_1(h_1, \dots, h_{f-1})) \\ &\times \rho_{|F_0|}(w_0(h_1, \dots, h_{f-1})) \left(\prod_{i=1}^{f-1} \rho_{|F_i|}(h_i) \right) dh_1 \dots dh_{f-1}. \end{aligned} \quad (30)$$

It is now straightforward to take a limit in (30) as $t \rightarrow 1$, that is, as all areas other than $|F_0|$ and $|F_1|$ tend to zero. In this limit, each heat kernel associated to h_i , $i \geq 2$, becomes a δ -measure, so we simply evaluate each such h_i to the identity element of K , giving

$$\begin{aligned} \lim_{t \rightarrow 1} \int_{K^e} \text{tr}(\text{hol}(L)) d\mu_{\text{sphere}}^{\mathbb{G}} \\ = \frac{1}{Z} \int_K \text{tr}(w_1(h_1, \text{id}, \dots, \text{id})) \rho_a(w_0(h_1, \text{id}, \dots, \text{id})) \rho_c(h_1) dh_1. \end{aligned} \quad (31)$$

In the sphere case, the normalization factor is given by $Z = \rho_A(\text{id})$, where A is the total area of the sphere. Although Z may depend on t (since we do not currently assume that the area of the sphere is fixed), it has a limit as $t \rightarrow 1$.

If the limiting value a of $|F_1|$ is also zero, then $\rho_{|F_1|}$ becomes also becomes a δ -function. Since the normalized trace of the identity matrix equals 1, the right-hand side of (31) becomes $\rho_c(\text{id})/Z$. But the total area of the sphere in this limit is just c , so $Z = \rho_c(\text{id})$ and $\rho_c(\text{id})/Z = 1$.

Finally, if the limiting value a of $|F_1|$ is nonzero, we must understand the effect on w_0 and w_1 of evaluating h_i , $i \geq 2$, to the identity. Recall that the words w_0 and w_1 arise from representing the boundary of F_0 and the loop L as words in L_1, \dots, L_{f-1} , which are free generators for $\pi_1(\mathbb{G})$. Now, $\pi_1(\mathbb{G})$ is naturally isomorphic to $\pi_1(\mathbb{R}^2 \setminus \{x_1, \dots, x_{f-1}\})$, where x_i is an arbitrarily chosen element of F_i . There is then a homomorphism from $\pi_1(\mathbb{G})$ to $\pi_1(\mathbb{R}^2 \setminus \{x_1\})$ induced by the inclusion of $\mathbb{R}^2 \setminus \{x_1, \dots, x_{f-1}\}$ into $\mathbb{R}^2 \setminus \{x_1\}$. Since $\pi_1(\mathbb{R}^2 \setminus \{x_1\})$ is just a free group on the single generator L_1 , this homomorphism is computed by mapping each of the generators L_2, \dots, L_{f-1} to the identity, leaving only powers of L_1 . Thus, if we write, say, ∂F_0 as a word in the generators L_1, \dots, L_{f-1} and apply the just-mentioned homomorphism, the result will be $L_1^{n_0}$, where n_0 is the winding number of ∂F_0 around x_1 (i.e., around F_1). By the Jordan curve theorem, this winding number is 1, assuming that we traverse ∂F_0 in the counter-clockwise direction. (In Figure 12, for example, the outer boundary of the loop in Figure 11 decomposes as $L_2 L_3 L_4 L_1$; if we set any three of the four generators to the identity, the remaining generator will occur to the power 1.) Expressing this result in terms of the loop *variables*, rather than the loops themselves, we conclude that $w_0(h_1, \text{id}, \dots, \text{id})$ is h_1 . Similarly, $w_1(h_1, \text{id}, \dots, \text{id}) = h_1^n$, where n is the winding number of L around F_1 . ■




Figure 12: The outer boundary of the loop in Figure 11 is computed as $L_2L_3L_4L_1$

4.6 The induction

In this section, we prove Theorem 3 by induction on the number of crossings. Our strategy is to deform an arbitrary loop L with k crossings into a loop L_n that winds n times around a simple closed curve. The variation of the Wilson loop functional along this deformation will be a linear combination of products of Wilson loop functionals for curves with fewer crossings, plus a covariance term. We begin by recording a result for a loop that winds n times around a simple closed curve.

Theorem 12 *Let $L_n(a, c)$ denote a loop that winds n times around a simple closed curve enclosing areas a and c . Then for all a and c , the limit*

$$W_n(a, c) := \mathbb{E} \{ \text{tr}(\text{hol}(L_n(a, c))) \}$$

exists and depends continuously on a and c . Furthermore, the associated variance tends to zero:

$$\lim_{N \rightarrow \infty} \text{Var} \{ \text{tr}(\text{hol}(L_n(a, c))) \} = 0.$$

The proof of this result is given in Section 4.7. Assuming Theorem 12 for the moment, we are now ready for the proof of our main result.

Proof of Theorem 3. Let k denote the number of crossings of L . We will proceed by induction on k . When $k = 0$, the result is precisely Theorem 1. Assume, then, that (7) and (8) and the continuity condition hold for all loops with $l < k$ crossings and consider a loop L with k crossings. Using Proposition 10, we may make a combination of Makeenko–Migdal variations at the vertices of L , giving loops $L(t)$ in which $L(0) = L$ and so that all but two of the faces shrink to zero as $t \rightarrow 1$. By Theorem 11, the limit as $t \rightarrow 1$ of the Wilson loop functional of $L(t)$ is the Wilson loop functional of a loop $L^n(a, c)$ that winds n times around a simple closed curve enclosing areas a and c , where a and

c are the areas of the two remaining faces. We may differentiate the Wilson loop functional of $L(t)$ using the chain rule and the first part of Proposition 5. Integrating the derivative then gives

$$\begin{aligned} \mathbb{E}\{\mathrm{tr}(\mathrm{hol}(L))\} &= \mathbb{E}\{\mathrm{tr}(\mathrm{hol}(L_n(a, c)))\} \\ &\quad - \sum_j c_j \int_0^1 \mathbb{E}\{\mathrm{tr}(\mathrm{hol}(L_{1,j}(t)))\} \mathbb{E}\{\mathrm{tr}(\mathrm{hol}(L_{2,j}(t)))\} dt \\ &\quad - \sum_j c_j \mathrm{Cov}\{\mathrm{tr}(\mathrm{hol}(L_{1,j}(t))), \mathrm{tr}(\mathrm{hol}(L_{2,j}(t)))\} dt, \end{aligned} \quad (32)$$

where the constants c_j are the ones expressing the vector \mathbf{b} in Proposition 10 as a linear combination of the Makeenko–Migdal vectors MM_{v_j} . Here $L_{1,j}(t)$ and $L_{2,j}(t)$ are the loops obtained by applying the Makeenko–Migdal equation at the vertex v_j to the loop $L(t)$. In particular, since neither of these loops has a crossing at v_j , we see that $L_{1,j}(t)$ and $L_{2,j}(t)$ have at most $k - 1$ crossings.

By our induction hypothesis, the limit

$$W(L_{i,j}(t)) := \lim_{N \rightarrow \infty} \mathbb{E}\{\mathrm{tr}(\mathrm{hol}(L_{i,j}(t)))\}$$

exists for each $t \in [0, 1)$, each $i \in \{1, 2\}$, and each j . Our induction hypothesis also tells us that the variance of $\mathrm{tr}(\mathrm{hol}(L_{i,j}(t)))$ goes to zero as $N \rightarrow \infty$; the inequality (22) then tells us that the covariances on the last line of (32) also tend to zero. Now, $|\mathrm{tr}(\mathrm{hol}(U))| \leq 1$ for all $U \in U(N)$, from which it follows that $|\mathrm{Var}(\mathrm{tr}(\mathrm{hol}(U)))| \leq 1$. Thus, we may apply dominated convergence to all integrals in (32), along with Theorem 12, to obtain

$$\begin{aligned} \lim_{N \rightarrow \infty} \mathbb{E}\{\mathrm{tr}(\mathrm{hol}(L))\} &= W(L_n(a, c)) \\ &\quad - \sum_j c_j \int_0^1 W(L_{1,j}(t))W(L_{2,j}(t)) dt, \end{aligned} \quad (33)$$

establishing the existence of the limit in (7) of Theorem 3.

We now establish the claimed continuity of $W(L)$ with respect to the areas of the faces. For each topological type of loop L , we fix an arbitrary face to play the role of F_0 in the proof of Proposition 10. We then fix two other faces with maximal and minimal winding numbers, respectively, relative to a puncture in F_0 . The first key observation is that with these choices made, the vector \mathbf{a}' in the proof of the proposition can be chosen to depend continuously on the areas of the faces. If we compute \mathbf{a}' by the formula in the proof, the only case in which continuity is not obvious is in the case in which (in the notation of the proof) $\mathbf{a} \cdot \mathbf{w} = 0$. But a moment's thought will confirm that as $\mathbf{a} \cdot \mathbf{w}$ tends to zero, \mathbf{a}' tends to the vector $(a_0, 0, \dots, 0)$, where a_0 is the area of S^2 , which is just the value of \mathbf{a}' when $\mathbf{a} \cdot \mathbf{w} = 0$.

Now, when we deform our original loop L into a loop of the form $L_n(a, c)$, the values of a and c depend continuously on the areas of the faces of L ; indeed, c and a are the first two entries of the vector \mathbf{a}' . The continuity of \mathbf{a}' also gives

the continuity of $\mathbf{b} = \mathbf{a}' - \mathbf{a}$ and thus of the coefficients c_j in (33), which are the coefficients of the expansion of \mathbf{b} in terms of the Makeenko–Migdal vectors. Thus, the continuity of the second term on the right-hand side of (33) follows from our induction hypothesis and dominated convergence.

To establish the second claim (8) of Theorem 3, we use the second part of Proposition 5. We then need to bound the covariance term appearing in (14). We first use (22) and then use Proposition 6 to bound the variance of the product of $\text{tr}(\text{hol}(L_1))$ and $\text{tr}(\text{hol}(L_2))$. The argument is then similar to the proof of (7).

Finally, we establish the large- N Makeenko–Migdal equation (9) for the limiting Wilson loop functionals. If we vary the areas in a checkerboard pattern along a path $L(t)$ as in Figure 3, we have

$$\begin{aligned} \mathbb{E} \{ \text{tr}(\text{hol}(L(t))) \} &= \mathbb{E} \{ \text{tr}(\text{hol}(L(t_0))) \} \\ &+ \int_{t_0}^t \mathbb{E} \{ \text{tr}(\text{hol}(L_1(s))) \} \mathbb{E} \{ \text{tr}(\text{hol}(L_2(s))) \} ds \\ &+ \int_{t_0}^t \text{Cov} \{ \text{tr}(\text{hol}(L_1(s))), \text{tr}(\text{hol}(L_2(s))) \} ds. \end{aligned}$$

Using the first two points (7) and (8) in Theorem 3, we can let N tend to infinity to obtain

$$W(L(t)) = W(L(t_0)) + \int_{t_0}^t W(L_1(s))W(L_2(s)) ds. \quad (34)$$

Since we have shown that $W(L)$ depends continuously on the areas of L , we see that $W(L_1(s))$ and $W(L_2(s))$ depend continuously on s . Thus, we can apply the fundamental theorem of calculus to differentiate (34) with respect to t to obtain the large- N Makeenko–Migdal equation (9). ■

4.7 The n -fold circle

In this section, we analyze the Wilson loop functional for a loop that winds n times around a simple closed curve, establishing Theorem 12. Similar results were obtained by Dahlgvist and Norris in [DN, Section 4.5].

Recall from (11) that the distribution of the parallel transport around a simple closed curve is the distribution of a Brownian bridge in $U(N)$. Assuming that the eigenvalue distribution of a Brownian bridge in $U(N)$ has a deterministic large- N limit, the calculations here provide a recursive method for computing the higher moments of the limiting distribution, in terms of the first moment. (But the formula for the first moment is complicated, especially in the large-lifetime phase, $a + b > \pi^2$.) A similar, but not identical, recursion for the moments of the free unitary Brownian motion—i.e., the large- N limit of a Brownian *motion* in $U(N)$, rather than a Brownian bridge—was established by Biane. See the bottom of p. 16 of [Bi1] and p. 266 of [Bi2].

We approximate a loop that winds n times around a simple closed curve by a loop with simple crossings, specifically, by a curve of the “loop within a loop




Figure 13: The loop $L_5(a, b, c)$

within a loop” form, as in Figure 13. If there are a total of n loops, then there are $n - 1$ crossings and a total of $n + 1$ faces. The faces divide into the innermost circle, the outer region, and $n - 1$ annular regions. Shifting the puncture from the innermost to outermost regions gives another curve of the same sort, with the order of the annular regions reversed.

Now, nothing prevents us from letting the areas of one or more of the annular regions equal zero. Specifically, we may understand the limit as some of the areas tend to zero by the same method as in the proof of Theorem 11. The only difference is that in the expression (30), we set an arbitrary subset of the loop variables equal to the identity. We may similarly make a Makeenko–Migdal variation at one or more of the vertices and then take a limit on both sides of the resulting equation as some of the areas tend to zero.

If we set the areas of the $n - 1$ annular regions to zero, the curve will simply wind n times around the outermost circle. More generally, we will show that it is possible to shrink the central region and the outer region, while increasing the innermost annular region and keeping the areas of the outermost $n - 2$ annular regions equal to zero, as in Figure 13. This observation motivates consideration of the following class of loops.

Definition 13 Let $L_n(a, c)$ denote the loop that winds n times around a simple closed curve enclosing areas a and c . Now let $L_1(a+b, c)$ be a simple closed curve having two faces F_1 and F_2 with areas $a + b$ and c , respectively. For $n \geq 2$, let $L_n(a, b, c)$ denote the loop that winds $n - 1$ times around $L_1(a + b, c)$ and then winds once around a region of area a inside F_1 .

When $n = 1$, we interpret $L_1(a, b, c)$ as being $L_1(a, b + c)$ (the loop that winds zero times around around $L_1(a + b, c)$ and then once around a region of area a).

We may consider two different limiting cases of $L_n(a, b, c)$, the case where a tends to zero and the case where c tends to zero. If a tends to zero, we are left with $L_{n-1}(b, c)$, while if c tends to zero, we are left with $L_1(a, b)$. (To see this, it may be useful to shift the puncture from the c face to the b face, as on the right-hand side of Figure 4.) We now give an inductive procedure for computing the limiting Wilson loop functionals for loops of the form $L_n(a, c)$ and $L_n(a, b, c)$.

Theorem 14 *For all $n \geq 1$ and all non-negative real numbers a, b , and c , the limits*

$$W_n(a, c) := \lim_{N \rightarrow \infty} \mathbb{E} \{ \text{tr}(\text{hol}(L_n(a, c))) \} \quad (35)$$

and

$$W_n(a, b, c) := \lim_{N \rightarrow \infty} \mathbb{E} \{ \text{tr}(\text{hol}(L_n(a, b, c))) \} \quad (36)$$

exist, and the associated variances tend to zero. Furthermore, for all $n \geq 1$, we have the following inductive formulas. If $c \geq a/n$, we have

$$\begin{aligned} W_{n+1}(a, b, c) &= W_n \left(\frac{n+1}{n}a + b, c - \frac{a}{n} \right) \\ &- \sum_{k=1}^n k \int_0^{a/n} W_k(a+b+t, c-t) W_{n+1-k}(a-nt, b+(n+1)t, c-t) dt. \end{aligned} \quad (37)$$

If, on the other hand, $c < a/n$, we have

$$\begin{aligned} W_{n+1}(a, b, c) &= W_1(a - nc, b + (n+1)c) \\ &- \sum_{k=1}^n k \int_0^c W_k(a+b+t, c-t) W_{n+1-k}(a-nt, b+(n+1)t, c-t) dt. \end{aligned} \quad (38)$$

Our main interest is in the quantity $W_n(a, c)$, which is the same as $W_n(a, 0, c)$. Note, however, that even if we put $b = 0$ on the left-hand side of (37) or (38), the right-hand side of the equation will still involve $W_{n+1-k}(a', b', c')$ with nonzero values of b' . On the other hand, the inductive procedure ultimately expresses $W_n(a, b, c)$ —and thus, in particular, $W_n(a, c) = W_n(a, 0, c)$ —in terms of $W_1(\cdot, \cdot)$.

Suppose, for example, that we wish to compute $W_3(a, c) = W_3(a, 0, c)$ in terms of $W_1(\cdot, \cdot)$. Since $W_3(a, c)$ is symmetric in a and c , it is harmless to assume that $c \geq a$, so that (37) applies. We first compute $W_2(a, b, c)$ in terms of $W_1(\cdot, \cdot)$, by applying (37) with $n = 1$ (since $c \geq a$) giving

$$\begin{aligned} W_2(a, b, c) &= W_1(2a + b, c - a) \\ &- \int_0^a W_1(a + b + t, c - t) W_1(a - t, b + c + t) dt, \quad c \geq a. \end{aligned} \quad (39)$$

(This formula is just what we obtained in (27) in Section 3.1.) We then apply

(37) with $n = 2$ and $b = 0$, giving

$$\begin{aligned}
W_3(a, c) &= W_2(3a/2, c - a/2) \\
&\quad - \int_0^{a/2} W_1(a + t, c - t)W_2(a - 2t, 3t, c - t) dt \\
&\quad - 2 \int_0^{a/2} W_2(a + t, c - t)W_1(a - 2t, c + 2t) dt. \tag{40}
\end{aligned}$$

Last, we plug into (40) the values of $W_2(a - 2t, 3t, c - t)$ and $W_2(a + t, c - t) = W_2(a + t, 0, c - t)$ computed in (39).

In the case of $W_2(a - 2t, 3t, c - t)$, the assumption that $c \geq a$ guarantees that $c - t \geq a - 2t$, so that we may directly apply (39). In the case of $W_2(a + t, c - t)$, there may be some values of t for which we need to use the symmetry of $W_2(a, c)$ with respect to a and c before applying (39). If, however, $c \geq 2a$, we may directly apply (39) to the computation of $W_2(a + t, c - t)$ for all $0 \leq t \leq a/2$.

Some explicit computations of these expressions in the plane case (i.e., the $c = \infty$ case) are given in Section 5.2.

Proof of Theorem 14. The argument is similar to the proof of Theorem 3, except with a different variation of the areas. We assume, inductively, that the limits in (35) and (36) exist for all $k \leq n$. We also assume that the corresponding variances go to zero. (When $n = 1$, this claim is just Theorem 1.) We then prove these results for $n + 1$, while also proving the inductive formulas (37) and (38).

We consider the loop $L_n(a, b, c)$. We realize this loop by starting with an n -fold “loop within a loop within a loop,” with the areas of the $n - 2$ outermost annular regions set equal to ε , and then letting ε tend to zero. We then order the $n - 1$ crossings from outermost to innermost. We then make a Makeenko–Migdal variation at the k th crossing, scaled by a factor of k . (See Figure 14.) The resulting change in the areas is

$$\begin{aligned}
a &\mapsto a - (n - 1)t \\
b &\mapsto b + nt \\
c &\mapsto c - t,
\end{aligned}$$

while the areas of the outer $n - 2$ annular regions remain unchanged.

If $c \geq a/n$, the area of the outermost region will remain non-negative as the area of the innermost region shrinks to zero. Thus, we may apply the Makeenko–Migdal equation (13) with the $n - 2$ annular areas equal to ε and then let ε tend




Figure 14: The variation in the areas of $L_5(a, b, c)$

to zero, giving a finite- N version of (37):

$$\begin{aligned}
& W_{n+1}^N(a, b, c) \\
&= W_n^N \left(\frac{n+1}{n}a + b, c - \frac{a}{n} \right) \\
&- \sum_{k=1}^n k \int_0^{a/n} W_k^N(a+b+t, c-t) W_{n+1-k}^N(a-nt, b+(n+1)t, c-t) dt \\
&+ \text{Cov.} \tag{41}
\end{aligned}$$

Here $W_n^N(a, b, c) = \mathbb{E} \{ \text{tr}(\text{hol}(L_n(a, b, c))) \}$ and $W_n^N(a, b) = \mathbb{E} \{ \text{tr}(\text{hol}(L_n(a, b))) \}$ and ‘‘Cov’’ is a covariance term.

Then, as in the proof of the first two points of Theorem 3, our induction hypothesis, together with (22) and Proposition 6, allows us to take the $N \rightarrow \infty$ limit with the covariance term going to zero. Thus, the large- N limit of $W_{n+1}^N(a, b, c)$ exists for $c \geq a/n$ and satisfies (37). The vanishing of the variance then follows by the same argument as in the proof of Theorem 3.

An entirely similar argument applies when $c < a/n$, using a finite- N version of (38), establishing existence of the limit and vanishing of the variance for $W_{n+1}(a, b, c)$ for all non-negative values of a , b , and c . The limit in (35) and vanishing of the variance then follow by setting $b = 0$. ■

5 The plane case, revisited

5.1 The general result

In [Lév2], Thierry Lévy establishes the large- N limit of Wilson loop functionals for Yang–Mills theory in the plane, together with the vanishing of the associ-

ated variances. (See Theorem 9.3.1 in [Lév2].) The limiting expectation values for loops with simple crossings are then characterized in [Lév2] by the large- N Makeenko–Migdal equation and another condition, labeled as Axiom Φ_4 in Section 0 of [Lév2] and termed the *unbounded face condition* in Theorem 2.3 of [DHK2].

The methods of the present paper give a new proof of some of these results, making it possible to treat the cases of the plane and the sphere in a uniform way. Specifically, the proof of Theorem 3 applies with minor changes in the plane case. Furthermore, the analog of Theorem 1 is easily established by direct computation in the plane case, as we will demonstrate shortly. We thus obtain a characterization of the master field on the plane slightly different from the one in [Lév2]; both characterizations use the large- N Makeenko–Migdal equation, but ours replaces the unbounded face condition with the (simple!) formula for the Wilson loop functional of a simple closed curve in the plane.

Theorem 15 *Theorem 3 holds also in the plane case.*

Proof. We follow the same argument as in the case of the sphere, with a few minor modifications. First, in Section 4.4, we should choose F_0 to be the unbounded face, so that the areas a_1, \dots, a_{f-1} are finite numbers. Theorem 11 still holds, with the same proof, in the plane case, provided we interpret $\rho_{|F_0|}$ as being the constant function $\mathbf{1}$. (The normalization factor Z is then also equal to 1.) All other proofs go through without change—except that in Section 4.7, the condition $c \geq a/n$ always holds because $c = \infty$. ■

We now supply the proof of the plane case of Theorem 1. Of course, the claim follows from results in [Lév2] (see Theorem 6.3.1), but we can give a elementary proof as follows.

Proof of Theorem 1 for \mathbb{R}^2 . The function $\text{tr}(U)$ is an eigenfunction for the Laplacian on $U(N)$ —with respect to the scaled Hilbert–Schmidt metric in (1)—with eigenvalue -1 . (See, e.g., Remark 3.4 in [DHK1].) It follows that for a simple closed curve C enclosing area a , we have

$$\mathbb{E} \{ \text{tr}(\text{hol}(C)) \} = e^{a\Delta/2}(\text{tr}(U)) \Big|_{U=I} = e^{-a/2}.$$

It is also possible to compute the (complex) variance of $\text{tr}(\text{hol}(C))$ explicitly. Using results from Section 3 of [DHK1], we may easily compute that

$$\Delta[\text{tr}(U)\overline{\text{tr}(U)}] = -2\text{tr}(U)\overline{\text{tr}(U)} - \frac{2}{N^2}.$$

(We have used that $\overline{\text{tr}(U)} = \text{tr}(U^*) = \text{tr}(U^{-1})$.) Thus, the space of functions spanned by $\text{tr}(U)\overline{\text{tr}(U)}$ and 1 is invariant under Δ , and the action of Δ on this space is represented by the matrix

$$\begin{pmatrix} -2 & 0 \\ -2/N^2 & 0 \end{pmatrix}.$$

We may exponentiate $a/2$ times this matrix using elementary formulas for the exponential of 2×2 matrices (e.g., Exercise 5 or Exercises 6 and 7 in [Ha1, Chapter 2]), with the result that

$$e^{a\Delta/2} \left[\text{tr}(U) \overline{\text{tr}(U)} \right] = e^{-a} \text{tr}(U) \overline{\text{tr}(U)} + \frac{e^{-a} - 1}{N^2},$$

so that

$$\begin{aligned} \mathbb{E} \left\{ |\text{tr}(U)|^2 \right\} &= e^{a\Delta/2} \left[\text{tr}(U) \overline{\text{tr}(U)} \right] \Big|_{U=I} \\ &= e^{-a} + \frac{e^{-a} - 1}{N^2}. \end{aligned}$$

Thus, the complex variance of $\text{tr}(\text{hol}(C))$ is

$$\mathbb{E} \left\{ |\text{tr}(\text{hol}(C))|^2 \right\} - |\mathbb{E} \{ \text{tr}(\text{hol}(C)) \}|^2 = e^{-a} + \frac{e^{-a} - 1}{N^2} - e^{-a},$$

which goes to zero as N tends to infinity.

More generally, Theorem 1.20 in [DHK1] identifies the leading-order term in the large- N asymptotics of the $U(N)$ Laplacian, acting on “trace polynomials” (that is, sums of products of traces of powers of U). This leading term satisfies a *first-order* product rule. Thus, the limiting heat operator is multiplicative, from which it follows that the variance of any trace polynomial vanishes as $N \rightarrow \infty$. (See the last displayed formula on p. 2620 of [DHK1].) ■

5.2 Example computations in the plane case

We now illustrate the computation of large- N Wilson loop functionals in the plane case. We first compute the large- N Wilson loop functionals $W_n(a, b, c)$ from Section 4.7 in the plane case, which corresponds to $c = \infty$, for $n \leq 3$. The condition $c \geq a/n$ is always satisfied in that case, and, as noted in the previous subsection, the Wilson loop functional for a simple closed curve is $W_1(a, \infty) = e^{-a/2}$.

The two-fold circle. When $c = \infty$, (37) takes the following form:

$$\begin{aligned} W_2(a, b, \infty) &= W_1(2a + b, \infty) - \int_0^a W_1(a + b + t, \infty) W_1(a - t, \infty) dt \\ &= e^{-a-b/2} - \int_0^a e^{-(a+b+t)/2} e^{-(a-t)/2} dt \\ &= e^{-a-b/2} - e^{-a-b/2} \int_0^a dt \\ &= e^{-a-b/2} (1 - a), \end{aligned}$$

which agrees with Lévy’s formula for the “loop within a loop” in the appendix of [Lév2]. When $b = 0$, we get $W_2(a, \infty) = e^{-a}(1 - a)$, which is the second

moment of Biane's distribution for the free unitary Brownian motion. (The moments may be found in Lemma 3 of [Bi1] or the Remark on p. 267 of [Bi2].)

The three-fold circle. When $n = 3$ and $c = \infty$, (40) becomes

$$\begin{aligned} W_3(a, b, \infty) &= e^{-(b+3a/2)}(1 - b - 3a/2) \\ &\quad - \int_0^{a/2} e^{-(a+b+t)/2} e^{-(a-2t)-(b+3t)/2} (1 - a + 2t) dt \\ &\quad - 2 \int_0^a e^{-(a+b+t)} (1 - a - b - t) e^{-(a-2t)/2} dt. \end{aligned}$$

All the t -terms in the exponents cancel, leaving us with $e^{-(b+3a/2)}$ times polynomials. Direct computation of the integral then gives

$$e^{-(b+3a/2)} \left(1 - 3a + \frac{3}{2}a^2 - b(1 - a) \right),$$

which agrees with the $s = 0$ case of the last of Lévy's examples of loops with two crossings in the appendix of [Lév2]. When $b = 0$, we get $e^{-3a/2}(1 - 3a + 3a^2/2)$, which is the third moment of Biane's distribution.

The trefoil. We now analyze the large- N Wilson loop functional for the trefoil loop in the plane case, using the strategy outlined in Section 4.1. In this example, the loops $L_{1,j}$ and $L_{2,j}$ occurring on the right-hand side of the Makeenko–Migdal equation are simple closed curves for all $j = 1, 2, 3$, and the product of the Wilson loop functionals simplifies as

$$W(L_{1,j}(t))W(L_{2,j}(t)) = e^{-a-(b+c+d)/2},$$

where the dependence on t and j drops out. Thus, when we integrate from $t = 0$ to $t = 1$, we obtain simply the constant value of the integrand. Now, the signs in the Makeenko–Migdal variations in Figure 8 are reversed from usual labeling. After adjusting for this and noting that the coefficients of the variations in Figure 8 add to $(b + c + d)/2$, we obtain

$$W(L) = W_2(a + (b + c + d)/2, \infty) + \left(\frac{b + c + d}{2} \right) e^{-a-(b+c+d)/2}.$$

Using the value for $W_2(a + (b + c + d)/2, \infty)$ computed above and simplifying gives

$$W(L) = e^{-a-(b+c+d)/2}(1 - a),$$

which agrees with the value of the master field for the trefoil in the appendix of [Lév2].

6 General compact surfaces

6.1 Introduction

We now consider the Yang–Mills measure on a compact surface Σ , possibly with boundary. (See [Sen3] and also [Lév1].) If the boundary of Σ is nonempty,

we may optionally impose constraints on the holonomy around the boundary components. The Makeenko–Migdal equation in this setting was established rigorously in [DGHK].

Let us say that a loop in Σ is **topologically trivial** if it is contained in an (open) topological disk $U \subset \Sigma$. All results in this section pertain only to topologically trivial loops. For a topologically trivial simple closed curve in Σ , one may reasonably expect that the analog of Theorem 1 will hold, but the author is not aware of any results in this direction.

Suppose, however, that we simply assume that the analog of Theorem 1 holds for topologically trivial simple closed curves in Σ . We will then establish the analog of Theorem 3—not for all topologically trivial loops, but only for those satisfying a “smallness” condition. The reason for the smallness assumption is as follows. Suppose L is a loop in a topological disk U and let F_0 denote the face of L that contains the complement of U in Σ . Then in the first stage of deformation, following the procedure in Section 4.4, we perform a Makeenko–Migdal variation of L to a loop that winds n times around a simple closed curve. This first variation can be done in such a way that *the area of F_0 does not decrease*, so that the deformed loop can be chosen to remain in U . In the second stage of deformation, we try to reduce—as in Section 4.7—from a curve that winds n times around a simple closed curve to a curve that winds only once around a simple closed curve. In this second variation, the area of F_0 will unavoidably decrease and may become zero, unless the original loop is “small.” If the area of F_0 becomes zero, the deformation process becomes undefined, because the limit of the Wilson loop as $|F_0|$ tends to zero is not easily evaluated (except when F_0 is a topological disk, i.e., when Σ is a sphere).

On the other hand, suppose we assume that a natural strengthening of Theorem 1 holds for topologically trivial loops in Σ . Specifically, suppose we simply *assume* that Theorem 1 holds not only for topologically trivial simple closed curves in Σ , but also for curves that wind n times around a topologically trivial simple closed curve. Then the second stage of deformation in the previous paragraph is not needed. Under this assumption, therefore, we will be able to prove Theorem 3 for *all* topologically trivial loops in Σ .

Let us consider what the previous discussion means for the study of the master field on a general compact surface Σ . The analysis will have two stages, as in the S^2 case: A direct calculation for the case of simple closed curves, and a reduction of general curves to the simple closed case using the Makeenko–Migdal equation. If we hope to use the just-discussed results for general surface, *more must be put into the direct calculation stage*.

For a simple closed curve C in Σ , Sengupta’s formula gives a probability measure μ_C on $U(N)$, which describes the distribution of the holonomy of a random connection around C . If we ultimately wish to analyze *all* topologically trivial loops with simple crossing, then in the direct calculation stage of analysis, we will have to look not just at the first moment of μ_C ,

$$\int_{U(N)} \text{tr}(U) d\mu_C(U), \quad (42)$$

but also at the higher moments,

$$\int_{U(N)} \text{tr}(U^n) d\mu_C(U), \quad (43)$$

which are just the Wilson loop functionals for curves that wind n times around C . If one could establish directly (i.e., without using the Makeenko–Migdal equation) the existence of the large- N limit and the vanishing of the variance for the quantities in (42) and (43), then the results of this section would apply to give the existence of the limit and vanishing of the variance for all topologically trivial loops with simple crossings.

6.2 Statements

We begin by stating the analog of Theorem 1 for topologically trivial loops in Σ as a conjecture.

Conjecture 16 *Consider a surface Σ of area $\text{area}(\Sigma)$. Let $U \subset \Sigma$ be a topological disk and consider a simple closed curve C in U . In Sengupta’s formula for the Yang–Mills measure on Σ , let us assign area a to the interior of C and area*

$$c := \text{area}(\Sigma) - a$$

to the exterior of C , for any $a < \text{area}(\Sigma)$. Then the limit

$$\lim_{N \rightarrow \infty} \mathbb{E} \{ \text{tr}(\text{hol}(C)) \}$$

exists and depends continuously on a and c , and

$$\lim_{N \rightarrow \infty} \text{Var} \{ \text{tr}(\text{hol}(C)) \} = 0.$$

We now introduce a notion of “smallness” for a general loop L in U with simple crossings. We may cut a loop L in U at a crossing, obtaining two loops L_1 and L_2 . We may then cut either L_1 or L_2 at one of its crossings, and so on. We refer to any loop that can be obtained by a finite sequence of such cuts as a **subloop** of L . In particular, L is a subloop of itself corresponding to making zero cuts. We label the faces of L as F_0, \dots, F_{f-1} where F_0 is the face containing the complement U^c of U .

For a loop L in U and a point x that is in U but not in L , we define the winding number of L around x by considering the homotopy class of L in $U \setminus \{x\}$, with respect to a fixed orientation on U . Next, we define, for each face F_j of L ,

$$|w|_{\max}(F_j) = \max_{L', j} |w(L', F_j)|, \quad (44)$$

where the maximum ranges over all subloops L' of L and all $j = 1, \dots, f-1$, and where $w(L', F)$ is the winding number of L' around F . Finally, if a_j is the area of F_j , we define

$$A = a_1 + a_2 + \dots + a_{f-1} = \text{area}(\Sigma) - \text{area}(F_0). \quad (45)$$

Remark 17 *One may bound $|w|_{\max}$ as follows. Suppose for some k , one can travel from any face of L to F_0 while crossing L at most k times. Then the same is true of any subloop L' of L . Since the winding number of F_0 is zero and $w(L', F)$ changes by one each time we cross L' , we conclude that $|w|_{\max}$ is at most k .*

Theorem 18 *Assume Conjecture 16. Let L be a loop traced out on a graph in a topological disk $U \subset \Sigma$ with simple crossings. Then Theorem 3 holds for L , provided L satisfies the “smallness” assumption*

$$A |w|_{\max} < \text{area}(\Sigma), \tag{46}$$

where $|w|_{\max}$ and A are defined in (44) and (45), respectively.

Note that since L is contained in a disk, L is certainly homotopically trivial in Σ . Thus, Theorem 18 does not tell us anything about homotopically nontrivial loops. Since U is a topological disk, Theorem 9 applies in this context, provided we compute the winding numbers of the faces of L by regarding L as a loop in U .

We next state a natural extension of Conjecture 16.

Conjecture 19 *Continuing with the notation of Conjecture 16, let $C^{(n)}$ denote the loop obtained by traveling n times around C . Then for all $n \in \mathbb{Z}$, the limit*

$$\lim_{N \rightarrow \infty} \mathbb{E} \left\{ \text{tr}(\text{hol}(C^{(n)})) \right\}$$

exists and depends continuously on a and c , and

$$\lim_{N \rightarrow \infty} \text{Var} \left\{ \text{tr}(\text{hol}(C^{(n)})) \right\} = 0.$$

Assuming Conjecture 19 holds, we can prove Theorem 3 for topologically trivial loops in Σ , *without* imposing a smallness assumption.

Theorem 20 *Assume Conjecture 19. Let L be a loop with only simple crossing, traced out on a graph in a topological disk U in Σ . Then Theorem 3 holds for L , without any smallness assumption on L .*

6.3 Proofs

Let U be a topological disk in Σ and let L be a loop traced out on a graph in U and having only simple crossings. We follow the deformation process in Section 4.4, with the face F_0 whose area does not decrease taken to be the face of L containing $\Sigma \setminus U$. The obvious analog of Theorem 11 continues to hold (see Lemma 22 below), so that at the end of the deformation process, we obtain a loop that winds n times around a simple closed curve in U . We may apply the same deformation process to all the subloops generated by the Makeenko–Migdal equation, and again the area of F_0 will not decrease. Thus, if we assume

the stronger assertion in Conjecture 19, we may prove Theorem 20 precisely as in Section 4.6, but using Conjecture 19 in place of Theorem 12.

If, on the other hand, we wish to assume only the weaker assertion in Conjecture 16, we must proceed to the second stage of analysis, deforming the loop that winds n times around a simple closed curve to one that winds once around a simple closed curve, as in Section 4.7. In this stage of analysis, the area of F_0 will unavoidably decrease, unless $|n| = 1$. If the area of F_0 becomes zero during the deformation, the whole process fails, because there is no easy way to compute the limit of a Wilson loop functional as the area of F_0 goes to zero. (See (50), in which the limit as $|F_0|$ goes to zero is not easily computed. The only case where this limit is easy to evaluate is when F_0 is a topological disk, that is, when Σ is a sphere.)

In the case of a curve winding n times around a simple closed curve, there is a simple condition (Theorem 21) on the n and the enclosed area guaranteeing that the area of F_0 will remain positive. We must then show that the smallness condition in Theorem 18 guarantees that each “ n -fold circle” generated by L will satisfy the hypotheses of Theorem 21.

Theorem 21 *Let the notation be as in Conjecture 16. For each nonzero integer n , let $L_n(a, c)$ denote the loop that winds n times around C . Assuming Conjecture 16, Theorem 3 holds for $L_n(a, c)$, provided that*

$$|n|a < \text{area}(\Sigma). \quad (47)$$

Proof. It is harmless to assume $n > 0$. Following the proof of Theorem 12 in the sphere case, we deform $L_n(a, c)$ into a loop of the form $L_n(a, b, c)$. Let $\mathbf{a} = (a, b, c)$ denote the vector of areas of the faces of $L_n(a, b, c)$ and let $\mathbf{w} = (w_1, w_2, w_3)$ be the associated vector of winding numbers, viewing $L_n(a, b, c)$ as a loop in the disk U , with $w_3 = 0$. We will actually prove Theorem 21 for the loops $L_n(a, b, c)$ by induction on $n > 0$, under the assumption that

$$\mathbf{a} \cdot \mathbf{w} < \text{area}(\Sigma), \quad (48)$$

which reduces to (47) when $n > 0$ and $b = 0$. When $n = 1$, the loop $L_1(a, b, c)$ is (by definition) the same as $L_1(a, b + c)$, so that the desired result is just Conjecture 16. For $n \geq 2$, we may follow the same sort of inductive argument as in the sphere case, provided that we never shrink the area of the “ c ” region to zero.

Take $n \geq 2$ and assume that Theorem 21 holds for loops of the form $L_k(a, b, c)$ satisfying (48), with $k < n$. Consider a loop $L_n(a, b, c)$ satisfying (48) and deform it into the loop $L_n(a(t), b(t), c(t))$ with $0 \leq t \leq a/(n-1)$, as in Section 4.7. As we vary the values of a , b , and c , the values of $\text{area}(\Sigma)$ and $\mathbf{a} \cdot \mathbf{w}$ remains constant—as can be seen explicitly or as a consequence of Theorem 9. Thus, by (48), we have

$$\mathbf{a} \cdot \mathbf{w} = na(t) + (n-1)b(t) < a(t) + b(t) + c(t). \quad (49)$$

Since $n \geq 2$, (49) tells us that $c(t) > (n-1)a(t) + (n-2)b(t) > 0$.

Thus, $c(t)$ remains positive as t approaches $a/(n-1)$, when we obtain a loop of the form $L_{n-1}(a', c')$, with

$$a' = b(t)|_{t=a/(n-1)} = b + \frac{n}{n-1}a.$$

We can then see explicitly that the value of $\mathbf{a} \cdot \mathbf{w}$ for $L_{n-1}(a', c')$ is the same as for $L_n(a, b, c)$, namely $na + (n-1)b$. Thus, by induction, Theorem 18 holds for $L_{n-1}(a', c')$. Furthermore, the loops $L_{1,j}(t)$ and $L_{2,j}(t)$ obtained from the Makeenko–Migdal equation will be $L_k(a(t), b(t), c(t))$ or $L_{n-k}(a(t) + b(t), c(t))$, with $1 \leq k < n$. It is then easy to see that the value of $\mathbf{a} \cdot \mathbf{w}$ for these loops is no bigger than for $L_n(a(t), b(t), c(t))$, which is the same as for $L_n(a, b, c)$. Thus, $L_{1,j}(t)$ and $L_{2,j}(t)$ satisfy (48) and by induction, Theorem 18 holds for these loops as well.

From this point, the argument is the same as in the sphere case. In particular, since we have ensured that $c(t)$ remains positive as we deform the areas of $L_n(a, b, c)$, we may apply (41) to compute the Wilson loop functional for $L_n(a, b, c)$. ■

We now prove Theorem 18. Following the logic in Section 4, we first deform L into a loop of the form $L_n(a, c)$. Since the area of F_0 increases during this process, there is no obstruction to carrying out this first step in the analysis of L . Nevertheless, we require a smallness assumption on L that will ensure that the limiting loop $L_n(a, c)$ will satisfy the hypothesis of Theorem 21. This smallness assumption must also be inherited by the loops $L_{1,j}(t)$ and $L_{2,j}(t)$ occurring on the right-hand side of the Makeenko–Migdal equation, so that these loops can be analyzed by induction on the number of crossings.

Proof of Theorem 18. We proceed by induction on the number k of crossings. If $k = 0$, the result is Conjecture 16. Assume, then, that the result holds for loops with fewer than k crossings and consider a loop L with k crossings. As in Lemma 22, we deform L into a loop $L_n(a, c)$. By the Makeenko–Migdal equation, the variation of the Wilson loop functional will involve loops of the form $L_{1,j}(t)$ and $L_{2,j}(t)$, all of which have fewer than k crossings. We need to verify (1) that $L_n(a, c)$ satisfies $|n|a < \text{area}(\Sigma)$ and (2) that $L_{1,j}(t)$ and $L_{2,j}(t)$ both satisfy the smallness assumption (46). If so, we may apply Theorem 21 to the loops $L_n(a, c)$ and our induction hypothesis to $L_{1,j}(t)$ and $L_{2,j}(t)$ and the argument is then the same as in the sphere case.

For Point (1), we note that the value of n is (Lemma 22) the winding number of L around F_1 , while the value of a is the limiting value of $a_1(t)$ (the area of F_1) as t approaches 1. Now, on the one hand, the value of $\mathbf{a} \cdot \mathbf{w}$ for $L(t)$ is independent of t , by Theorem 9. On the other hand, $\mathbf{a} \cdot \mathbf{w}$ approaches the value na as t approaches 1, since $a_1(t) \rightarrow a$ and $a_j(t) \rightarrow 0$ for $j \geq 2$. Thus, $na = \mathbf{a} \cdot \mathbf{w}$. But from the definitions (45) and (44) of A and $|w|_{\max}$, we have $|\mathbf{a} \cdot \mathbf{w}| \leq A|w|_{\max}$ and thus

$$|n|a = |\mathbf{a} \cdot \mathbf{w}| \leq A|w|_{\max} < \text{area}(\Sigma).$$

For Point (2), we note that for the loops $L(t)$, the area of F_0 is always increasing (Proposition 10), meaning that $A(t)$ is decreasing. Thus, for all i and

j , we have

$$A_{L_{i,j}(t)} \leq A_{L(t)} \leq A.$$

Furthermore, since every subloop of $L_{i,j}$ is also a subloop of L , we see that $|w_{L_{i,j}}|_{\max} \leq |w_L|_{\max}$. Thus, we have

$$A_{L_{i,j}(t)} |w_{L_{i,j}}|_{\max} \leq A |w_L|_{\max} < \text{area}(\Sigma).$$

Having verified these two points, the proof now proceeds as in the sphere case.

■

It remains only to verify that the analog of Theorem 11 holds for loops in $U \subset \Sigma$.

Lemma 22 *Consider the Yang–Mills measure on Σ for an arbitrary connected compact Lie group K . Let L be a loop traced out on a graph in $U \subset \Sigma$ and having only simple crossings. Denote the number of faces of L by f and label the faces as $F_0, F_1, F_2, \dots, F_{f-1}$, where F_0 is the face containing U^c . Suppose we vary the areas of the faces as a function of a parameter $t \in [0, 1)$ in such a way that as $t \rightarrow 1$, the areas of F_2, \dots, F_{f-1} tend to zero, while the areas of F_0 and F_1 approach non-negative real numbers c and a , respectively, with $a > 0$. Then*

$$\lim_{t \rightarrow 1} \mathbb{E}\{\text{tr}(\text{hol}(L))\} = \mathbb{E}\{\text{tr}(\text{hol}(L_n(a, c)))\},$$

where n is the winding number of L around F_1 .

Proof. Let \mathbb{G} be a minimal graph (necessarily connected) in which L can be traced. Before we can apply Sengupta’s formula, we must embed \mathbb{G} into an admissible graph \mathbb{G}' , that is, one that contains the boundary of Σ and each of whose faces is a topological disk. Actually, by the Jordan curve theorem, all the faces of \mathbb{G} other than F_0 will automatically be disks. It is then possible to construct \mathbb{G}' by adding new edges entirely in the closure of F_0 . (See Section 1.2 of [Lév1].) Thus, the faces of \mathbb{G}' may be chosen to be of the form $F'_0, F_1, \dots, F_{f-1}$, where F'_0 is a subset of F_0 having the same area as F_0 . Let us divide the edge variables for \mathbb{G}' into the edge variables \mathbf{x} corresponding to the original graph \mathbb{G} and the remaining edge variables \mathbf{y} . Then integration with respect to the Yang–Mills measure for the graph \mathbb{G}' in Σ may be written as

$$\int_{K^{e'}} f(\mathbf{x}, \mathbf{y}) d\mu_{\Sigma}^{\mathbb{G}'} = \frac{1}{Z} \int_{K^{e'-e}} \rho_{|F_0|}(\text{hol}_{F'_0}(\mathbf{x}, \mathbf{y})) \int_{K^e} f(\mathbf{x}, \mathbf{y}) d\mu_{\text{plane}}^{\mathbb{G}}(\mathbf{x}) d\mathbf{y}, \quad (50)$$

where e and e' are the number of edges of \mathbb{G} and \mathbb{G}' , respectively. Furthermore, we may write

$$\text{hol}_{F'_0}(\mathbf{x}, \mathbf{y}) = \text{hol}_{F_0}(\mathbf{x})g(\mathbf{y})$$

for some word $g(\mathbf{y})$ in the \mathbf{y} variables. Here, for notational simplicity, we consider the unconditional Yang–Mills measure, but a similar argument applies if there are constraints on the holonomies around the boundary components of Σ .

In the case that f is the trace of the holonomy of L , we may imitate the proof of Theorem 11 to obtain

$$\begin{aligned} \mathbb{E} \{ \text{tr}(\text{hol}(L)) \} &= \frac{1}{Z} \int_{K^{e'-e}} \int_{K^{f-1}} \text{tr}(w_1(h_1, \dots, h_{f-1})) \\ &\times \rho_{|F_0|}(w_0(h_1, \dots, h_{f-1})g(\mathbf{y})) \left(\prod_{i=1}^{f-1} \rho_{|F_i|}(h_i) \right) dh_1 \dots dh_{f-1} d\mathbf{y}. \end{aligned}$$

As in that proof, if we let $t \rightarrow 1$, we obtain

$$\lim_{t \rightarrow 1} \mathbb{E} \{ \text{tr}(\text{hol}(L)) \} = \frac{1}{Z} \int_{K^{e'-e}} \int_{K^{f-1}} \text{tr}(h_1^n) \rho_c(h_1 g(\mathbf{y})) \rho_a(h_1) dh_1 d\mathbf{y}, \quad (51)$$

where n is the winding number of L around F_1 . But the right-hand side of (51) is just Sengupta's formula for the Wilson loop functional for the loop that winds n times around the boundary of F_0 (i.e., the outer boundary of \mathbb{G}), enclosing areas a and c . ■

7 Acknowledgments

The author thanks Bruce Driver, David Galvin, Todd Kemp, Thierry Lévy, Karl Liechty, Andy Putman, Ambar Sengupta, and Steve Zelditch for valuable discussions. The author is also grateful to Franck Gabriel for a detailed reading of the entire manuscript, along with many corrections and helpful suggestions. Finally, the author thanks the two referees, whose careful reading and thoughtful comments have improved the manuscript immensely.

References

- [AS] M. Anshelevich and A. N. Sengupta, Quantum free Yang-Mills on the plane. *J. Geom. Phys.* **62** (2012), 330–343.
- [Bi1] P. Biane, Free Brownian motion, free stochastic calculus and random matrices. *In: Free probability theory* (Waterloo, ON, 1995), 1–19, Fields Inst. Commun. **12**, Amer. Math. Soc., Providence, RI, 1997.
- [Bi2] P. Biane, Segal-Bargmann transform, functional calculus on matrix spaces and the theory of semi-circular and circular systems, *J. Funct. Anal.* **144** (1997), 232–286.
- [Bou] D. V. Boulatov, Wilson loop on a sphere, *Mod. Phys. Lett. A* **9** (1994), 365–374.
- [BS] A. Boutet de Monvel and M. Shcherbina, On free energy in two-dimensional $U(N)$ -gauge field theory on the sphere. (Russian) *Theoret. Mat. Fiz.* **115** (1998), 389–401; translation in *Theoret. and Math. Phys.* **115** (1998), 670–679.

- [CDG] G. Cébron, A. Dahlqvist, and F. Gabriel, The generalized master fields, *J. Geom. Phys.* **119** (2017), 34–53.
- [Dahl] A. Dahlqvist, Free energies and fluctuations for the unitary Brownian motion, *Comm. Math. Phys.* **348** (2016), 395–444.
- [DN] A. Dahlqvist and J. Norris, Yang-Mills measure and the master field on the sphere, preprint arXiv:1703.10578 [math.PR].
- [DaK] J.-M. Daul and V. A. Kazakov, Wilson loop for large N Yang-Mills theory on a two-dimensional sphere, *Physics Letters B* **335** (1994), 371–376.
- [DoK] M. R. Douglas and V. A. Kazakov, Large N phase transition in continuum QCD₂, *Phys. Let. B* **319** (1993), 219–230.
- [Dr] B. K. Driver, YM₂: continuum expectations, lattice convergence, and lassos. *Comm. Math. Phys.* **123** (1989), 575–616.
- [DHK1] B. K. Driver, B. C. Hall, and T. Kemp, The large- N limit of the Segal–Bargmann transform on $U(N)$, *J. Funct. Anal.* **265** (2013), 2585–2644.
- [DHK2] B. K. Driver, B. C. Hall, and T. Kemp, Three proofs of the Makeenko–Migdal equation for Yang–Mills theory on the plane, *Comm. Math. Phys.* **351** (2017), 741–774.
- [DGHK] B. Driver, F. Gabriel, B. C. Hall, and T. Kemp, The Makeenko–Migdal equation for Yang–Mills theory on compact surfaces, *Comm. Math. Phys.* **352** (2017), 967–978.
- [Dys] F. J. Dyson, A Brownian-motion model for the eigenvalues of a random matrix, *J. of Math. Phys.* **3** (1962), 1191–1198.
- [Ès] L. D. Èskin, Heat equation on Lie groups (Russian). In 1964 In Memoriam: N. G. Čebotarev (Russian) pp. 113–132, *Izdat. Kazan. Univ.*, Kazan.
- [FMS] P. J. Forrester, S. N. Majumdar, G. Schehr, Non-intersecting Brownian walkers and Yang–Mills theory on the sphere, *Nucl. Phys. B* **844** (2011), 500–526.
- [Ga] F. Gabriel, Planar Markovian holonomy fields, preprint arXiv:1501.05077 [math-ph].
- [Gop] R. Gopakumar, The master field in generalised QCD₂, *Nuclear Phys. B* **471** (1996), 246–260.
- [GG] R. Gopakumar and D. Gross, Mastering the master field, *Nuclear Phys. B* **451** (1995), 379–415.

- [Gr] D. Gross, Two Dimensional QCD as a String Theory, *Nucl. Phys. B* **400** (1993), 181-208.
- [GT1] D. Gross and W. Taylor, Two-dimensional QCD is a String Theory, *Nucl. Phys. B* **400** (1993) 161-180.
- [GT2] D. Gross and W. Taylor, Twists and loops in the string theory of two dimensional QCD, *Nucl. Phys. B* 403 (1993), 395- 452.
- [GM] D. J. Gross and A. Matytsin, Some properties of large- N two-dimensional Yang-Mills theory, *Nuclear Phys. B* **437** (1995), 541–584.
- [GKS] L. Gross, C. King, and A. N. Sengupta, Two-dimensional Yang-Mills theory via stochastic differential equations, *Ann. Physics* **194** (1989), 65–112.
- [Hal1] B. C. Hall, Lie Groups, Lie Algebras, and Representations. An Elementary Introduction. Graduate Texts in Mathematics, #222, Second Edition, Springer, 2015.
- [Ha2] B. C. Hall, The eigenvalue process for Brownian motion in $U(N)$, currently unpublished note, available on the author’s web page www.nd.edu/~bhall/.
- [K] A. Kazakov, Wilson loop average for an arbitrary contour in two-dimensional $U(N)$ gauge theory, *Nuclear Physics* **B179** (1981) 283-292.
- [Lév1] T. Lévy, Yang-Mills measure on compact surfaces, *Mem. Amer. Math. Soc.* **166** (2003), no. 790, xiv+122 pp.
- [Lév2] T. Lévy, The master field on the plane, *Astérisque* No. 388 (2017), ix+201 pp.
- [LM] T. Lévy and M. Maïda, On the Douglas-Kazakov phase transition. Weighted potential theory under constraint for probabilists. *Modélisation Aléatoire et Statistique—Journées MAS 2014*, 89–121, ESAIM Proc. Surveys, 51, EDP Sci., Les Ulis, 2015.
- [LW1] K. Liechty and D. Wang, Nonintersecting Brownian motions on the unit circle: noncritical cases, arXiv:1312.7390v3 [math.PR]. (This is a preprint version of [LW2] that contains material not in the final version.)
- [LW2] K. Liechty and D. Wang, Nonintersecting Brownian motions on the unit circle, *Ann. Probab.* **44** (2016), 1134–1211.
- [KK] V. A. Kazakov and I. K. Kostov, Non-linear strings in two-dimensional $U(\infty)$ gauge theory, *Nucl. Phys. B* **176** (1980), 199-215.

- [MM] Y. M. Makeenko and A. A. Migdal, Exact equation for the loop average in multicolor QCD, *Physics Letters* **88B** (1979), 135-137.
- [MO] P. Menotti and E. Onofri, The action of $SU(N)$ lattice gauge theory in terms of the heat kernel on the group manifold, *Nuclear Phys. B* **190**, 288–300 (1981).
- [Mig] A. A. Migdal, Recursion equations in gauge field theories, *Sov. Phys. JETP* **42**, 413–418 (1975).
- [Sen1] A. N. Sengupta, Quantum gauge theory on compact surfaces, *Ann. Physics* **221** (1993), 17–52.
- [Sen2] A. N. Sengupta, Gauge theory on compact surfaces, *Mem. Amer. Math. Soc.* **126** (1997), no. 600, viii+85 pp.
- [Sen3] A. N. Sengupta, Yang-Mills on surfaces with boundary: quantum theory and symplectic limit, *Comm. Math. Phys.* **183** (1997), 661–705.
- [Sen4] A. N. Sengupta, Traces in two-dimensional QCD: the large- N limit. Traces in number theory, geometry and quantum fields, 193–212, *Aspects Math.*, E38, Friedr. Vieweg, Wiesbaden, 2008.
- [Si] I. M. Singer, On the master field in two dimensions, *In: Functional analysis on the eve of the 21st century*, Vol. 1 (New Brunswick, NJ, 1993), 263–281, *Progr. Math.*, 131, Birkhäuser Boston, Boston, MA, 1995.
- [’t H] G. ’t Hooft, A planar diagram theory for strong interactions, *Nuclear Physics B* **72** (1974), 461-473.
- [Tao] T. Tao, Topics in random matrix theory. Graduate Studies in Mathematics, 132. American Mathematical Society, Providence, RI, 2012.
- [Ur] H. Urakawa, The heat equation on compact Lie group, *Osaka J. Math.* **12** (1975), 285–297.
- [Xu] F. Xu, A random matrix model from two-dimensional Yang-Mills theory, *Comm. Math. Phys.* **190** (1997), 287 – 307.