

# A Global div-curl-Lemma for Mixed Boundary Conditions in Weak Lipschitz Domains and a Corresponding Generalized $A_0^*$ - $A_1$ -Lemma in Hilbert Spaces

DIRK PAULY

ABSTRACT. We prove global and local versions of the so-called div-curl-lemma, a crucial result in the homogenization theory of partial differential equations, for mixed boundary conditions on bounded weak Lipschitz domains in 3D with weak Lipschitz interfaces. We will generalize our results using an abstract Hilbert space setting, which shows corresponding results to hold in arbitrary dimensions as well as for various differential operators. The crucial tools and the core of our arguments are Hilbert complexes and related compact embeddings.

## CONTENTS

1. Introduction	1
2. Definitions and Preliminaries	4
3. The div-rot-Lemma	6
4. Generalizations	7
4.1. Functional Analysis Toolbox	7
4.2. The $A_0^*$ - $A_1$ -Lemma	9
4.3. Generalizations of the $A_0^*$ - $A_1$ -Lemma	10
5. Applications	16
5.1. The div-rot-Lemma Revisited	16
5.2. Generalized Electro-Magnetics	20
5.3. Biharmonic Equation, General Relativity, and Gravitational Waves	22
5.4. Linear Elasticity	24
References	26

## 1. INTRODUCTION

The classical div-curl-lemma by Murat [17] and Tartar [32], a famous and crucial result in the homogenization theory of partial differential equations and often used for so-called compensated compactness, reads as follows:

**Theorem I** (classical div-curl-lemma). *Let  $\Omega \subset \mathbb{R}^3$  be an open set and let  $(E_n), (H_n) \subset L^2(\Omega)$  be two sequences bounded in  $L^2(\Omega)$  such that both  $(\widetilde{\text{curl}} E_n)$  and  $(\widetilde{\text{div}} H_n)$  are relatively compact in  $H^{-1}(\Omega)$ . Then there exist  $E, H \in L^2(\Omega)$  as well as subsequences, again denoted by  $(E_n)$  and  $(H_n)$ , such that the sequence of scalar products  $(E_n \cdot H_n)$  converges in the sense of distributions, i.e.,*

$$\forall \varphi \in \dot{C}^\infty(\Omega) \quad \int_{\Omega} \varphi (E_n \cdot H_n) \rightarrow \int_{\Omega} \varphi (E \cdot H).$$

Here,  $H^{-1}(\Omega)$  denotes the dual space of  $\dot{H}^1(\Omega)$  and the distributional extensions

$$\widetilde{\text{curl}} : L^2(\Omega) \rightarrow H^{-1}(\Omega), \quad \widetilde{\text{div}} : L^2(\Omega) \rightarrow H^{-1}(\Omega)$$

of curl and div, respectively, are defined for  $E \in L^2(\Omega)$  by

$$\widetilde{\text{curl}} E(\Phi) := \langle \text{curl } \Phi, E \rangle_{L^2(\Omega)}, \quad \Phi \in \dot{H}^1(\Omega),$$

---

*Date:* December 12, 2018.

*1991 Mathematics Subject Classification.* 35B27, 35Q61, 47B07, 46B50.

*Key words and phrases.* div-curl-lemma, compensated compactness, mixed boundary conditions, weak Lipschitz domains, Maxwell's equations.

$$\widetilde{\operatorname{div}} E(\varphi) := -\langle \nabla \varphi, E \rangle_{L^2(\Omega)}, \quad \varphi \in \dot{H}^1(\Omega).$$

We will prove a global version of the div-curl-lemma stating that under certain (mixed tangential and normal) boundary conditions and (very weak) regularity assumptions on a domain  $\Omega \subset \mathbb{R}^3$ , see Section 2, the following holds:

**Theorem II** (global div-curl-lemma). *Let  $\Omega \subset \mathbb{R}^3$  be a bounded weak Lipschitz domain with boundary  $\Gamma$  and weak Lipschitz boundary parts  $\Gamma_t$  and  $\Gamma_n$ . Let  $(E_n)$  and  $(H_n)$  be two sequences bounded in  $L^2(\Omega)$ , such that  $(\operatorname{curl} E_n)$  and  $(\operatorname{div} H_n)$  are also bounded in  $L^2(\Omega)$  and  $\nu \times E_n = 0$  on  $\Gamma_t$  and  $\nu \cdot H_n = 0$  on  $\Gamma_n$ . Then there exist subsequences, again denoted by  $(E_n)$  and  $(H_n)$ , such that  $(E_n)$ ,  $(\operatorname{curl} E_n)$  and  $(H_n)$ ,  $(\operatorname{div} H_n)$  converge weakly to  $E$ ,  $\operatorname{curl} E$  and  $H$ ,  $\operatorname{div} H$  in  $L^2(\Omega)$ , respectively, and the inner products converge as well, i.e.,*

$$\int_{\Omega} E_n \cdot H_n \rightarrow \int_{\Omega} E \cdot H.$$

A local version similar to the classical div-curl-lemma from Theorem I (distributional like convergence for arbitrary domains and no boundary conditions needed) is then immediately implied.

**Corollary III** (local div-curl-lemma). *Let  $\Omega \subset \mathbb{R}^3$  be an open set. Let  $(E_n)$  and  $(H_n)$  be two sequences bounded in  $L^2(\Omega)$ , such that  $(\operatorname{curl} E_n)$  and  $(\operatorname{div} H_n)$  are also bounded in  $L^2(\Omega)$ . Then there exist subsequences, again denoted by  $(E_n)$  and  $(H_n)$ , such that  $(E_n)$ ,  $(\operatorname{curl} E_n)$  and  $(H_n)$ ,  $(\operatorname{div} H_n)$  converge weakly to  $E$ ,  $\operatorname{curl} E$  and  $H$ ,  $\operatorname{div} H$  in  $L^2(\Omega)$ , respectively, and the inner products converge in the distributional sense as well, i.e., for all  $\varphi \in \dot{C}^\infty(\Omega)$  it holds*

$$\int_{\Omega} \varphi (E_n \cdot H_n) \rightarrow \int_{\Omega} \varphi (E \cdot H).$$

For details see Theorem 3.1, Corollary 3.2, Theorem 5.2, and Theorem 5.6.

We will also generalize these results to a natural Hilbert complex setting. For this, let

$$A_0 : D(A_0) \subset H_0 \rightarrow H_1, \quad A_1 : D(A_1) \subset H_1 \rightarrow H_2$$

be two (possibly unbounded) densely defined and closed linear operators on three Hilbert spaces  $H_0$ ,  $H_1$ ,  $H_2$  with Hilbert space adjoints

$$A_0^* : D(A_0^*) \subset H_1 \rightarrow H_0, \quad A_1^* : D(A_1^*) \subset H_2 \rightarrow H_1.$$

Moreover, let the complex property  $A_1 A_0 = 0$  be satisfied, i.e.,

$$R(A_0) \subset N(A_1).$$

In Theorem 4.7 we present our central result of this contribution which reads as follows:

**Theorem IV** (generalized div-curl-lemma:  $A_0^*$ - $A_1$ -lemma). *Let  $D(A_1) \cap D(A_0^*) \hookrightarrow H_1$  be compact. If  $(x_n) \subset D(A_1)$  and  $(y_n) \subset D(A_0^*)$  are two  $D(A_1)$ -bounded resp.  $D(A_0^*)$ -bounded sequences, then there exist  $x \in D(A_1)$  and  $y \in D(A_0^*)$  as well as subsequences, again denoted by  $(x_n)$  and  $(y_n)$ , such that  $(x_n)$  and  $(y_n)$  converge weakly in  $D(A_1)$  and  $D(A_0^*)$  to  $x$  and  $y$ , respectively, together with the convergence of the inner products*

$$\langle x_n, y_n \rangle_{H_1} \rightarrow \langle x, y \rangle_{H_1}.$$

**Remark V.** *The compact embedding  $D(A_1) \cap D(A_0^*) \hookrightarrow H_1$  reads in Theorem II as*

$$\{E \in L^2(\Omega) : \operatorname{curl} E \in L^2(\Omega), \operatorname{div} E \in L^2(\Omega), \nu \times E|_{\Gamma_t} = 0, \nu \cdot E|_{\Gamma_n} = 0\} \hookrightarrow L^2(\Omega),$$

which is known as Weck's selection theorem, see Lemma 2.1.

In Theorem 4.14 the latter theorem is even generalized to a distributional version as follows:

**Theorem VI** (generalized div-curl-lemma: generalized  $A_0^*$ - $A_1$ -lemma). *Let the ranges  $R(A_0)$  and  $R(A_1)$  be closed and let  $N(A_1) \cap N(A_0^*)$  be finite-dimensional. Moreover, let  $(x_n), (y_n) \subset H_1$  be two bounded sequences such that  $(\widetilde{A_1} x_n)$  and  $(\widetilde{A_0^*} y_n)$  are relatively compact in  $D(A_1^*)'$  and  $D(A_0)'$ , respectively. Then there exist  $x, y \in H_1$  as well as subsequences, again denoted by  $(x_n)$  and  $(y_n)$ , such that  $(x_n)$  and  $(y_n)$  converge weakly in  $H_1$  to  $x$  and  $y$ , respectively, together with the convergence of the inner products*

$$\langle x_n, y_n \rangle_{H_1} \rightarrow \langle x, y \rangle_{H_1}.$$

Here, the distributional extensions

$$\widetilde{A_1} : H_1 \rightarrow D(A_1^*)', \quad \widetilde{A_0^*} : H_1 \rightarrow D(A_0)'$$

of  $A_1$  and  $A_0^*$ , respectively, are defined for  $x \in H_1$  by

$$\begin{aligned} \widetilde{A_1} x(\phi) &:= \langle A_1^* \phi, x \rangle_{H_1}, \quad \phi \in D(A_1^*), \\ \widetilde{A_0^*} x(\varphi) &:= \langle A_0 \varphi, x \rangle_{H_1}, \quad \varphi \in D(A_0). \end{aligned}$$

In Section 5 we apply these results to various differential operators in 3D and ND, appearing, e.g., in classical and generalized electro-magnetics, for the biharmonic equation, in general relativity, for gravitational waves, and in the theory of linear elasticity and plasticity. We obtain also an interesting additional version of the global div-curl-lemma, compare to Theorem 5.9.

**Theorem VII** (alternative global div-curl-lemma). *Let  $\Omega \subset \mathbb{R}^3$  be a bounded strong Lipschitz domain with trivial topology. Moreover, let  $(E_n), (H_n) \subset L^2(\Omega)$  be two bounded sequences such that either  $(\widehat{\text{curl}} E_n)$  and  $(\widehat{\text{div}} H_n)$  are relatively compact in  $\dot{H}^{-1}(\Omega)$  and  $H^{-1}(\Omega)$ , respectively, or  $(\widehat{\text{curl}} E_n)$  and  $(\widehat{\text{div}} H_n)$  are relatively compact in  $H^{-1}(\Omega)$  and  $\dot{H}^{-1}(\Omega)$ , respectively. Then there exist  $E, H \in L^2(\Omega)$  as well as subsequences, again denoted by  $(E_n)$  and  $(H_n)$ , such that  $E_n$  and  $H_n$  converge weakly in  $L^2(\Omega)$ , respectively, together with the convergence of the inner products*

$$\langle E_n, H_n \rangle_{L^2(\Omega)} \rightarrow \langle E, H \rangle_{L^2(\Omega)}.$$

Here,  $\dot{H}^{-1}(\Omega) := H^1(\Omega)'$  and the distributional extensions

$$\widehat{\text{curl}} : L^2(\Omega) \rightarrow \dot{H}^{-1}(\Omega), \quad \widehat{\text{div}} : L^2(\Omega) \rightarrow \dot{H}^{-1}(\Omega)$$

of curl and div, respectively, are defined for  $E \in L^2(\Omega)$  by

$$\begin{aligned} \widehat{\text{curl}} E(\Phi) &:= \langle \text{curl } \Phi, E \rangle_{L^2(\Omega)}, \quad \Phi \in H^1(\Omega), \\ \widehat{\text{div}} E(\varphi) &:= -\langle \nabla \varphi, E \rangle_{L^2(\Omega)}, \quad \varphi \in H^1(\Omega). \end{aligned}$$

The div-curl-lemma, which serves as a central result in the theory of compensated compactness, see the original papers by Murat [17] and Tartar [32] with crucial applications in [8] or [10, 31], and its variants and extensions have plenty of important applications. For an extensive discussion and a historical overview of the div-curl-lemma see [33]. More recent discussions can be found, e.g., in [6, 34] as well as in [7] and in the nice paper [35] of Marcus Waurick. The latter two contributions utilize a Hilbert/Banach space setting as well, but from different perspectives. In [35] Waurick achieved closely related results using different methods and proofs, see Section 4.3. Interesting applications to homogenization of partial differential equations have recently been given in [36]. From our personal<sup>i</sup> point of view, although the results of [7, 35] are slightly more general, our methods and proofs are easier and more canonical and hence give deeper insight into the underlying structure and the core of the main result and thus of all div-curl-type lemmas.

<sup>i</sup>The idea of this paper came up a few years ago in 2012, when Sören Bartels asked the author about the div-curl-lemma and for a simpler proof. Moreover, in 2016, the div-curl-lemma in a form similar to the one in this article was subject of lots of discussions with Marcus Waurick, when he as well as the author were lecturing Special Semester Courses on Maxwell's equations and related topics invited by Ulrich Langer at the Johann Radon Institute for Computational and Applied Mathematics (RICAM) in Linz.

The div-curl-lemma is widely used in the theory of homogenization of (nonlinear) partial differential equations, see, e.g., [31]. Compensated compactness has many important applications in nonlinear partial differential equations and calculus of variations, e.g., in the partial regularity theory of stationary harmonic maps, see, e.g., [12, 11, 28]. Numerical applications can be found, e.g., in [2]. It is further a crucial tool in the homogenization of stochastic partial differential equations, especially with certain random coefficients, see, e.g., the survey [1] and the literature cited therein, e.g., [13].

Let us also mention that the div-curl-lemma is particularly useful to treat homogenization of problems arising in plasticity, see, e.g., a recent contribution on this topic [29], for which [30] provides the important key div-curl-lemma. As in [30, 29]  $H^1(\Omega)$ -potentials are used, these contributions are restricted to smooth, e.g.,  $C^2$  or convex, domains and to full boundary conditions. This clearly shows that the more general and stronger div-curl-lemma results presented in the contribution at hand are of great importance and so far unknown to the community. The same  $H^1(\Omega)$ -detour as in [30, 29] is used in the recent contribution [15] where div-curl-type lemmas are presented which also allow for inhomogeneous boundary conditions. This unnecessarily high regularity assumption of  $H^1(\Omega)$ -fields excludes results like [15, 30, 29] to be applied to important applications which are stated, e.g., in Lipschitz domains.

Generally, for problems related to Maxwell's equations the detour over  $H^1(\Omega)$  and using Rellich's selection theorem instead of using Weck's selection theorem, see Lemma 2.1, seems to be the wrong way to deal with such equations. Most of the arguments simply fail, and if not, the results are usually limited to smooth domains and trivial topologies. Mixed boundary conditions cannot be treated properly. Since the early 1970's, see the original paper by Weck [38] for Weck's selection theorem, it is well-known, that the  $H^1(\Omega)$ -detour is often not helpful and does not lead to satisfying results. Surprisingly, this fact appears to be unknown to a wider community.

## 2. DEFINITIONS AND PRELIMINARIES

Let  $\Omega \subset \mathbb{R}^3$  be a bounded weak Lipschitz domain, see [3, Definition 2.3] for details, with boundary  $\Gamma := \partial\Omega$ , which is divided into two relatively open weak Lipschitz subsets  $\Gamma_t$  and  $\Gamma_n := \Gamma \setminus \overline{\Gamma_t}$  (its complement), see [3, Definition 2.5] for details. Note that strong Lipschitz (graph of Lipschitz functions) implies weak Lipschitz (Lipschitz manifolds) for the boundary as well as the interface. Throughout this section we shall assume the latter regularity on  $\Omega$  and  $\Gamma_t$ .

Recently, in [3], Weck's selection theorem, also known as the Maxwell compactness property, has been shown to hold for such bounded weak Lipschitz domains and mixed boundary conditions. More precisely, the following holds:

**Lemma 2.1** (Weck's selection theorem). *The embedding  $\mathring{R}_{\Gamma_t}(\Omega) \cap \mathring{D}_{\Gamma_n}(\Omega) \hookrightarrow L^2(\Omega)$  is compact.*

For a proof see [3, Theorem 4.7]. A short historical overview of Weck's selection theorem is given in the introduction of [3], see also the original paper [38] and [27, 37, 9, 39, 14, 16] for simpler proofs and generalizations.

Here the usual Lebesgue and Sobolev spaces are denoted by  $L^2(\Omega)$  and  $H^1(\Omega)$  as well as

$$R(\Omega) := \{E \in L^2(\Omega) : \operatorname{rot} E \in L^2(\Omega)\}, \quad D(\Omega) := \{E \in L^2(\Omega) : \operatorname{div} E \in L^2(\Omega)\},$$

where we prefer to write  $\operatorname{rot}$  instead of  $\operatorname{curl}$ .  $R(\Omega)$  and  $D(\Omega)$  are also written as  $H(\operatorname{rot}, \Omega)$ ,  $H(\operatorname{curl}, \Omega)$  and  $H(\operatorname{div}, \Omega)$  in the literature. With the help of test functions and test vector fields

$$\mathring{C}_{\Gamma_t}^\infty(\Omega) := \{\varphi|_\Omega : \varphi \in \mathring{C}^\infty(\mathbb{R}^3), \operatorname{dist}(\operatorname{supp} \varphi, \Gamma_t) > 0\}$$

we define the closed subspaces

$$(2.1) \quad \mathring{H}_{\Gamma_t}^1(\Omega) := \overline{\mathring{C}_{\Gamma_t}^\infty(\Omega)}^{H^1(\Omega)}, \quad \mathring{R}_{\Gamma_t}(\Omega) := \overline{\mathring{C}_{\Gamma_t}^\infty(\Omega)}^{R(\Omega)}, \quad \mathring{D}_{\Gamma_n}(\Omega) := \overline{\mathring{C}_{\Gamma_n}^\infty(\Omega)}^{D(\Omega)}$$

as closures of test functions and vector fields, respectively. If  $\Gamma_t = \Gamma$  we skip the index  $\Gamma$  and write

$$\mathring{C}^\infty(\Omega) = \mathring{C}_\Gamma^\infty(\Omega), \quad \mathring{H}^1(\Omega) = \mathring{H}_\Gamma^1(\Omega), \quad \mathring{R}(\Omega) = \mathring{R}_\Gamma(\Omega), \quad \mathring{D}(\Omega) = \mathring{D}_\Gamma(\Omega).$$

In (2.1) homogeneous scalar, tangential and normal traces on  $\Gamma_t$  and  $\Gamma_n$  are generalized. For the pathological case  $\Gamma_t = \emptyset$ , we put

$$\mathring{H}_\emptyset^1(\Omega) := H^1(\Omega) \cap \mathbb{R}^{\perp L^2(\Omega)} = \left\{ u \in H^1(\Omega) : \int_\Omega u = 0 \right\}$$

in order to still have a Poincaré estimate for  $u \in \mathring{H}_\emptyset^1(\Omega)$ . Let us emphasize that our assumptions also allow for Rellich's selection theorem, i.e., the embedding

$$(2.2) \quad \mathring{H}_{\Gamma_t}^1(\Omega) \hookrightarrow L^2(\Omega)$$

is compact, see, e.g., [3, Theorem 4.8]. By density we have the two rules of integration by parts

$$(2.3) \quad \forall u \in \mathring{H}_{\Gamma_t}^1(\Omega) \quad \forall H \in \mathring{D}_{\Gamma_n}(\Omega) \quad \langle \nabla u, H \rangle_{L^2(\Omega)} = -\langle u, \operatorname{div} H \rangle_{L^2(\Omega)},$$

$$(2.4) \quad \forall E \in \mathring{R}_{\Gamma_t}(\Omega) \quad \forall H \in \mathring{R}_{\Gamma_n}(\Omega) \quad \langle \operatorname{rot} E, H \rangle_{L^2(\Omega)} = \langle E, \operatorname{rot} H \rangle_{L^2(\Omega)}.$$

We emphasize that, besides Weck's selection theorem, the resulting Maxwell estimates (Friedrichs/Poincaré type estimates), Helmholtz decompositions, closed ranges, continuous and compact inverse operators, and an appropriate electro-magneto static solution theory for bounded weak Lipschitz domains and mixed boundary conditions, another important result has been shown in [3]. It holds

$$(2.5) \quad \begin{aligned} \mathring{H}_{\Gamma_t}^1(\Omega) &= \{ u \in H^1(\Omega) : \langle \nabla u, \Phi \rangle_{L^2(\Omega)} = -\langle u, \operatorname{div} \Phi \rangle_{L^2(\Omega)} \text{ for all } \Phi \in \mathring{C}_{\Gamma_n}^\infty(\Omega) \}, \\ \mathring{R}_{\Gamma_t}(\Omega) &= \{ E \in R(\Omega) : \langle \operatorname{rot} E, \Phi \rangle_{L^2(\Omega)} = \langle E, \operatorname{rot} \Phi \rangle_{L^2(\Omega)} \text{ for all } \Phi \in \mathring{C}_{\Gamma_n}^\infty(\Omega) \}, \\ \mathring{D}_{\Gamma_n}(\Omega) &= \{ H \in D(\Omega) : \langle \operatorname{div} H, \varphi \rangle_{L^2(\Omega)} = -\langle H, \nabla \varphi \rangle_{L^2(\Omega)} \text{ for all } \varphi \in \mathring{C}_{\Gamma_t}^\infty(\Omega) \}, \end{aligned}$$

i.e., strong and weak definitions of boundary conditions coincide, see [3, Theorem 4.5]. Furthermore, we define the closed subspaces of irrotational and solenoidal vector fields

$$R_0(\Omega) := \{ E \in R(\Omega) : \operatorname{rot} E = 0 \}, \quad D_0(\Omega) := \{ E \in D(\Omega) : \operatorname{div} E = 0 \},$$

respectively, as well as

$$\mathring{R}_{\Gamma_t,0}(\Omega) := \mathring{R}_{\Gamma_t}(\Omega) \cap R_0(\Omega), \quad \mathring{D}_{\Gamma_n,0}(\Omega) := \mathring{D}_{\Gamma_n}(\Omega) \cap D_0(\Omega).$$

A direct consequence of Lemma 2.1 is the compactness of the unit ball in

$$\mathcal{H}(\Omega) := \mathring{R}_{\Gamma_t,0}(\Omega) \cap \mathring{D}_{\Gamma_n,0}(\Omega),$$

the space of so-called Dirichlet-Neumann fields. Hence  $\mathcal{H}(\Omega)$  is finite-dimensional. Another immediate consequence of Weck's selection theorem, Lemma 2.1, using a standard indirect argument, is the so-called Maxwell estimate, i.e., there exists  $c_m > 0$  such that

$$(2.6) \quad \forall E \in \mathring{R}_{\Gamma_t}(\Omega) \cap \mathring{D}_{\Gamma_n}(\Omega) \cap \mathcal{H}(\Omega)^{\perp L^2(\Omega)} \quad |E|_{L^2(\Omega)} \leq c_m (|\operatorname{rot} E|_{L^2(\Omega)} + |\operatorname{div} E|_{L^2(\Omega)})$$

or, equivalently,

$$(2.7) \quad \forall E \in \mathring{R}_{\Gamma_t}(\Omega) \cap \mathring{D}_{\Gamma_n}(\Omega) \quad |E - \pi E|_{L^2(\Omega)} \leq c_m (|\operatorname{rot} E|_{L^2(\Omega)} + |\operatorname{div} E|_{L^2(\Omega)}),$$

see [3, Theorem 5.1], where  $\pi : L^2(\Omega) \rightarrow \mathcal{H}(\Omega)$  denotes the  $L^2(\Omega)$ -orthonormal projector onto the Dirichlet-Neumann fields. Recent estimates for the Maxwell constant  $c_m$  can be found in [18, 19, 20]. Analogously, Rellich's selection theorem (2.2) shows the Friedrichs/Poincaré estimate

$$(2.8) \quad \exists c_{f,p} > 0 \quad \forall u \in \mathring{H}_{\Gamma_t}^1(\Omega) \quad |u|_{L^2(\Omega)} \leq c_{f,p} |\nabla u|_{L^2(\Omega)},$$

see [3, Theorem 4.8]. By the projection theorem, applied to the densely defined and closed (unbounded) linear operator

$$\nabla : \mathring{H}_{\Gamma_t}^1(\Omega) \subset L^2(\Omega) \longrightarrow L^2(\Omega)$$

with (Hilbert space) adjoint

$$\nabla^* = -\operatorname{div} : \mathring{D}_{\Gamma_n}(\Omega) \subset L^2(\Omega) \longrightarrow L^2(\Omega),$$

where we have used (2.5), we get the simple Helmholtz decomposition

$$(2.9) \quad \mathbf{L}^2(\Omega) = \nabla \mathring{H}_{\Gamma_t}^1(\Omega) \oplus_{\mathbf{L}^2(\Omega)} \mathring{D}_{\Gamma_n,0}(\Omega),$$

see [3, Theorem 5.3 or (13)], which immediately implies

$$(2.10) \quad \mathring{R}_{\Gamma_t}(\Omega) = \nabla \mathring{H}_{\Gamma_t}^1(\Omega) \oplus_{\mathbf{L}^2(\Omega)} (\mathring{R}_{\Gamma_t}(\Omega) \cap \mathring{D}_{\Gamma_n,0}(\Omega))$$

as  $\nabla \mathring{H}_{\Gamma_t}^1(\Omega) \subset \mathring{R}_{\Gamma_t,0}(\Omega)$ . Here  $\oplus_{\mathbf{L}^2(\Omega)}$  in the decompositions (2.9) and (2.10) denotes the orthogonal sum in the Hilbert space  $\mathbf{L}^2(\Omega)$ . By (2.8), the range  $\nabla \mathring{H}_{\Gamma_t}^1(\Omega)$  is closed in  $\mathbf{L}^2(\Omega)$ , see also [3, Lemma 5.2]. Note that we call (2.9) a simple Helmholtz decomposition, since the refined Helmholtz decomposition

$$\mathbf{L}^2(\Omega) = \nabla \mathring{H}_{\Gamma_t}^1(\Omega) \oplus_{\mathbf{L}^2(\Omega)} \mathcal{H}(\Omega) \oplus_{\mathbf{L}^2(\Omega)} \text{rot } \mathring{R}_{\Gamma_n}(\Omega)$$

holds as well, see [3, Theorem 5.3], where also  $\text{rot } \mathring{R}_{\Gamma_n}(\Omega)$  is closed in  $\mathbf{L}^2(\Omega)$  as a consequence of (2.6), see [3, Lemma 5.2].

### 3. THE DIV-ROT-LEMMA

From now on we use synonymously the notion div-curl-lemma and div-rot-lemma. Let  $\Omega \subset \mathbb{R}^3$  be a bounded weak Lipschitz domain with weak Lipschitz interfaces as introduced in Section 2.

**Theorem 3.1** (global div-rot-lemma). *Let  $(E_n) \subset \mathring{R}_{\Gamma_t}(\Omega)$  and  $(H_n) \subset \mathring{D}_{\Gamma_n}(\Omega)$  be two sequences bounded in  $\mathbf{R}(\Omega)$  and  $\mathbf{D}(\Omega)$ , respectively. Then there exist  $E \in \mathring{R}_{\Gamma_t}(\Omega)$  and  $H \in \mathring{D}_{\Gamma_n}(\Omega)$  as well as subsequences, again denoted by  $(E_n)$  and  $(H_n)$ , such that*

- $E_n \rightharpoonup E$  in  $\mathring{R}_{\Gamma_t}(\Omega)$ ,
- $H_n \rightharpoonup H$  in  $\mathring{D}_{\Gamma_n}(\Omega)$ ,
- $\langle E_n, H_n \rangle_{\mathbf{L}^2(\Omega)} \rightarrow \langle E, H \rangle_{\mathbf{L}^2(\Omega)}$ .

*Proof.* We pick subsequences, again denoted by  $(E_n)$  and  $(H_n)$ , such that  $(E_n)$  and  $(H_n)$  converge weakly in  $\mathring{R}_{\Gamma_t}(\Omega)$  and  $\mathring{D}_{\Gamma_n}(\Omega)$  to  $E \in \mathring{R}_{\Gamma_t}(\Omega)$  and  $H \in \mathring{D}_{\Gamma_n}(\Omega)$ , respectively. By the simple Helmholtz decomposition (2.10), we have the orthogonal decomposition  $\mathring{R}_{\Gamma_t}(\Omega) \ni E_n = \nabla u_n + \tilde{E}_n$  with some  $u_n \in \mathring{H}_{\Gamma_t}^1(\Omega)$  and  $\tilde{E}_n \in \mathring{R}_{\Gamma_t}(\Omega) \cap \mathring{D}_{\Gamma_n,0}(\Omega)$ . Then  $(u_n)$  is bounded in  $\mathbf{H}^1(\Omega)$  by orthogonality and the Friedrichs/Poincaré estimate (2.8).  $(\tilde{E}_n)$  is bounded in  $\mathbf{R}(\Omega) \cap \mathbf{D}(\Omega)$  by orthogonality and  $\text{rot } \tilde{E}_n = \text{rot } E_n$ ,  $\text{div } \tilde{E}_n = 0$ . Hence, using Rellich's and Weck's selection theorems, i.e., (2.2) and Lemma 2.1, there exist  $u \in \mathring{H}_{\Gamma_t}^1(\Omega)$  and  $\tilde{E} \in \mathring{R}_{\Gamma_t}(\Omega) \cap \mathring{D}_{\Gamma_n,0}(\Omega)$  and we can extract two subsequences, again denoted by  $(u_n)$  and  $(\tilde{E}_n)$ , such that  $u_n \rightarrow u$  in  $\mathring{H}_{\Gamma_t}^1(\Omega)$  and  $u_n \rightarrow u$  in  $\mathbf{L}^2(\Omega)$  as well as  $\tilde{E}_n \rightharpoonup \tilde{E}$  in  $\mathring{R}_{\Gamma_t}(\Omega) \cap \mathring{D}_{\Gamma_n,0}(\Omega)$  and  $\tilde{E}_n \rightarrow \tilde{E}$  in  $\mathbf{L}^2(\Omega)$ . We have  $E = \nabla u + \tilde{E}$ , giving the simple Helmholtz decomposition for  $E$ , as, e.g., for all  $\varphi \in \mathring{C}^\infty(\Omega)$

$$\langle E, \varphi \rangle_{\mathbf{L}^2(\Omega)} \leftarrow \langle E_n, \varphi \rangle_{\mathbf{L}^2(\Omega)} = \langle \nabla u_n, \varphi \rangle_{\mathbf{L}^2(\Omega)} + \langle \tilde{E}_n, \varphi \rangle_{\mathbf{L}^2(\Omega)} \rightarrow \langle \nabla u, \varphi \rangle_{\mathbf{L}^2(\Omega)} + \langle \tilde{E}, \varphi \rangle_{\mathbf{L}^2(\Omega)}.$$

Then by (2.3)

$$\begin{aligned} \langle E_n, H_n \rangle_{\mathbf{L}^2(\Omega)} &= \langle \nabla u_n, H_n \rangle_{\mathbf{L}^2(\Omega)} + \langle \tilde{E}_n, H_n \rangle_{\mathbf{L}^2(\Omega)} = -\langle u_n, \text{div } H_n \rangle_{\mathbf{L}^2(\Omega)} + \langle \tilde{E}_n, H_n \rangle_{\mathbf{L}^2(\Omega)} \\ &\rightarrow -\langle u, \text{div } H \rangle_{\mathbf{L}^2(\Omega)} + \langle \tilde{E}, H \rangle_{\mathbf{L}^2(\Omega)} = \langle \nabla u, H \rangle_{\mathbf{L}^2(\Omega)} + \langle \tilde{E}, H \rangle_{\mathbf{L}^2(\Omega)} = \langle E, H \rangle_{\mathbf{L}^2(\Omega)}, \end{aligned}$$

completing the proof.  $\square$

**Corollary 3.2** (local div-rot-lemma). *Let  $(E_n) \subset \mathbf{R}(\Omega)$  and  $(H_n) \subset \mathbf{D}(\Omega)$  be two sequences bounded in  $\mathbf{R}(\Omega)$  and  $\mathbf{D}(\Omega)$ , respectively. Then there exist  $E \in \mathbf{R}(\Omega)$  and  $H \in \mathbf{D}(\Omega)$  as well as subsequences, again denoted by  $(E_n)$  and  $(H_n)$ , such that  $E_n \rightharpoonup E$  in  $\mathbf{R}(\Omega)$  and  $H_n \rightharpoonup H$  in  $\mathbf{D}(\Omega)$  together with the distributional convergence*

$$\forall \varphi \in \mathring{C}^\infty(\Omega) \quad \langle \varphi E_n, H_n \rangle_{\mathbf{L}^2(\Omega)} \rightarrow \langle \varphi E, H \rangle_{\mathbf{L}^2(\Omega)}.$$

*Proof.* Let  $\Gamma_t := \Gamma$  and hence  $\Gamma_n = \emptyset$ .  $(\varphi E_n)$  is bounded in  $\mathring{R}_\Gamma(\Omega)$  and  $(H_n)$  is bounded in  $D(\Omega)$ . Theorem 3.1 shows the assertion.  $\square$

**Remark 3.3.** We note that the boundedness of  $(E_n)$  and  $(H_n)$  in local spaces is sufficient for Corollary 3.2 to hold. Hence, no regularity or boundedness assumptions on  $\Omega$  are needed, i.e., Corollary 3.2 holds for an arbitrary open set  $\Omega \subset \mathbb{R}^3$ . Moreover,  $\varphi \in \mathring{C}^\infty(\Omega)$  may be replaced by  $\varphi \in \mathring{C}^1(\Omega)$  or even  $\varphi \in \mathring{C}^{0,1}(\Omega)$ , the space of Lipschitz continuous functions vanishing in a neighbourhood of  $\Gamma$ .

#### 4. GENERALIZATIONS

The idea of the proof of Theorem 3.1 can be generalized.

**4.1. Functional Analysis Toolbox.** Let  $A : D(A) \subset H_1 \rightarrow H_2$  be a (possibly unbounded) closed and densely defined linear operator on two Hilbert spaces  $H_1$  and  $H_2$  with adjoint  $A^* : D(A^*) \subset H_2 \rightarrow H_1$ . Note  $(A^*)^* = \overline{A} = A$ , i.e.,  $(A, A^*)$  is a dual pair. By the projection theorem the Helmholtz type decompositions

$$(4.1) \quad H_1 = N(A) \oplus_{H_1} \overline{R(A^*)}, \quad H_2 = N(A^*) \oplus_{H_2} \overline{R(A)}$$

hold, where we introduce the notation  $N$  for the kernel (or null space) and  $R$  for the range of a linear operator. We can define the reduced operators

$$\begin{aligned} \mathcal{A} &:= A|_{\overline{R(A^*)}} : D(\mathcal{A}) \subset \overline{R(A^*)} \rightarrow \overline{R(A)}, & D(\mathcal{A}) &:= D(A) \cap N(A)^{\perp_{H_1}} = D(A) \cap \overline{R(A^*)}, \\ \mathcal{A}^* &:= A^*|_{\overline{R(A)}} : D(\mathcal{A}^*) \subset \overline{R(A)} \rightarrow \overline{R(A^*)}, & D(\mathcal{A}^*) &:= D(A^*) \cap N(A^*)^{\perp_{H_2}} = D(A^*) \cap \overline{R(A)}, \end{aligned}$$

which are also closed and densely defined linear operators. We note that  $\mathcal{A}$  and  $\mathcal{A}^*$  are indeed adjoint to each other, i.e.,  $(\mathcal{A}, \mathcal{A}^*)$  is a dual pair as well. Now the inverse operators

$$\mathcal{A}^{-1} : R(\mathcal{A}) \rightarrow D(\mathcal{A}), \quad (\mathcal{A}^*)^{-1} : R(\mathcal{A}^*) \rightarrow D(\mathcal{A}^*)$$

exist and are bijective, since  $\mathcal{A}$  and  $\mathcal{A}^*$  are injective by definition. Furthermore, by (4.1) we have the refined Helmholtz type decompositions

$$(4.2) \quad D(A) = N(A) \oplus_{H_1} D(\mathcal{A}), \quad D(A^*) = N(A^*) \oplus_{H_2} D(\mathcal{A}^*)$$

and thus we obtain for the ranges

$$(4.3) \quad R(A) = R(\mathcal{A}), \quad R(A^*) = R(\mathcal{A}^*).$$

By the closed range theorem and the closed graph theorem we get immediately the following.

**Lemma 4.1.** *The following assertions are equivalent:*

- (i)  $\exists c_A \in (0, \infty) \quad \forall x \in D(\mathcal{A}) \quad |x|_{H_1} \leq c_A |Ax|_{H_2}$
- (i\*)  $\exists c_{A^*} \in (0, \infty) \quad \forall y \in D(\mathcal{A}^*) \quad |y|_{H_2} \leq c_{A^*} |A^*y|_{H_1}$
- (ii)  $R(A) = R(\mathcal{A})$  is closed in  $H_2$ .
- (ii\*)  $R(A^*) = R(\mathcal{A}^*)$  is closed in  $H_1$ .
- (iii)  $\mathcal{A}^{-1} : R(\mathcal{A}) \rightarrow D(\mathcal{A})$  is continuous and bijective.
- (iii\*)  $(\mathcal{A}^*)^{-1} : R(\mathcal{A}^*) \rightarrow D(\mathcal{A}^*)$  is continuous and bijective.

In case that one of the latter assertions is true, e.g., (ii),  $R(A)$  is closed, we have

$$\begin{aligned} H_1 &= N(A) \oplus_{H_1} R(A^*), & H_2 &= N(A^*) \oplus_{H_2} R(A), \\ D(A) &= N(A) \oplus_{H_1} D(\mathcal{A}), & D(A^*) &= N(A^*) \oplus_{H_2} D(\mathcal{A}^*), \\ D(\mathcal{A}) &= D(A) \cap R(A^*), & D(\mathcal{A}^*) &= D(A^*) \cap R(A), \end{aligned}$$

and

$$\mathcal{A} : D(\mathcal{A}) \subset R(A^*) \rightarrow R(A), \quad \mathcal{A}^* : D(\mathcal{A}^*) \subset R(A) \rightarrow R(A^*).$$

**Remark 4.2.** For the “best” constants  $c_A, c_{A^*}$  the following holds: The Rayleigh quotients

$$\frac{1}{c_A} := \inf_{0 \neq x \in D(\mathcal{A})} \frac{|Ax|_{\mathbf{H}_2}}{|x|_{\mathbf{H}_1}}, \quad \frac{1}{c_{A^*}} := \inf_{0 \neq y \in D(\mathcal{A}^*)} \frac{|A^*y|_{\mathbf{H}_1}}{|y|_{\mathbf{H}_2}}$$

coincide, i.e.,  $c_A = c_{A^*} \in (0, \infty]$ .

**Lemma 4.3.** The following assertions are equivalent:

- (i)  $D(\mathcal{A}) \Leftrightarrow \mathbf{H}_1$  is compact.
- (i\*)  $D(\mathcal{A}^*) \Leftrightarrow \mathbf{H}_2$  is compact.
- (ii)  $\mathcal{A}^{-1} : R(\mathcal{A}) \rightarrow R(\mathcal{A}^*)$  is compact.
- (ii\*)  $(\mathcal{A}^*)^{-1} : R(\mathcal{A}^*) \rightarrow R(\mathcal{A})$  is compact.

If one of these assertions holds true, e.g., (i),  $D(\mathcal{A}) \Leftrightarrow \mathbf{H}_1$  is compact, then the assertions of Lemma 4.1 and Remark 4.2 hold with  $c_A = c_{A^*} \in (0, \infty)$ . Especially, the Friedrichs/Poincaré type estimates hold, all ranges are closed and the inverse operators

$$\mathcal{A}^{-1} : R(\mathcal{A}) \rightarrow R(\mathcal{A}^*), \quad (\mathcal{A}^*)^{-1} : R(\mathcal{A}^*) \rightarrow R(\mathcal{A})$$

are compact with norms  $|\mathcal{A}^{-1}|_{R(\mathcal{A}), R(\mathcal{A}^*)} = |(\mathcal{A}^*)^{-1}|_{R(\mathcal{A}^*), R(\mathcal{A})} = c_A$ .

*Proof.* As the other assertions are easily proved or immediately clear by symmetry, we just show that (i), i.e., the compactness of

$$D(\mathcal{A}) = D(\mathcal{A}) \cap \overline{R(\mathcal{A}^*)} \Leftrightarrow \mathbf{H}_1,$$

implies (i\*) as well as Lemma 4.1 (i).

(i) $\Rightarrow$ Lemma 4.1 (i): For this we use a standard indirect argument. If Lemma 4.1 (i) were wrong, there would exist a sequence  $(x_n) \subset D(\mathcal{A})$  with  $|x_n|_{\mathbf{H}_1} = 1$  and  $Ax_n \rightarrow 0$ . As  $(x_n)$  is bounded in  $D(\mathcal{A})$  we can extract a subsequence, again denoted by  $(x_n)$ , with  $x_n \rightarrow x \in \mathbf{H}_1$  in  $\mathbf{H}_1$ . Since  $\mathcal{A}$  is closed, we have  $x \in D(\mathcal{A})$  and  $Ax = 0$ , hence  $x \in N(\mathcal{A}) = \{0\}$ , in contradiction to  $1 = |x_n|_{\mathbf{H}_1} \rightarrow |x|_{\mathbf{H}_1} = 0$ .

(i) $\Rightarrow$ (i\*): Let  $(y_n) \subset D(\mathcal{A}^*)$  be a bounded sequence. Utilizing Lemma 4.1 (i) and (ii) we obtain  $D(\mathcal{A}^*) = D(\mathcal{A}^*) \cap R(\mathcal{A})$  and thus  $y_n = Ax_n$  with  $(x_n) \subset D(\mathcal{A})$ , which is bounded in  $D(\mathcal{A})$  by Lemma 4.1 (i). Hence we may extract a subsequence, again denoted by  $(x_n)$ , converging in  $\mathbf{H}_1$ . Therefore with  $x_{n,m} := x_n - x_m$  and  $y_{n,m} := y_n - y_m$  we see

$$|y_{n,m}|_{\mathbf{H}_2}^2 = \langle y_{n,m}, Ax_{n,m} \rangle_{\mathbf{H}_2} = \langle A^*(y_{n,m}), x_{n,m} \rangle_{\mathbf{H}_1} \leq c|x_{n,m}|_{\mathbf{H}_1},$$

and hence  $(y_n)$  is a Cauchy sequence in  $\mathbf{H}_2$ .  $\square$

Now, let  $A_0 : D(A_0) \subset \mathbf{H}_0 \rightarrow \mathbf{H}_1$  and  $A_1 : D(A_1) \subset \mathbf{H}_1 \rightarrow \mathbf{H}_2$  be (possibly unbounded) closed and densely defined linear operators on three Hilbert spaces  $\mathbf{H}_0, \mathbf{H}_1$ , and  $\mathbf{H}_2$  with adjoints  $A_0^* : D(A_0^*) \subset \mathbf{H}_1 \rightarrow \mathbf{H}_0$  and  $A_1^* : D(A_1^*) \subset \mathbf{H}_2 \rightarrow \mathbf{H}_1$  as well as reduced operators  $\mathcal{A}_0, \mathcal{A}_0^*$ , and  $\mathcal{A}_1, \mathcal{A}_1^*$ . Furthermore, we assume the sequence or complex property of  $A_0$  and  $A_1$ , that is,  $A_1 A_0 = 0$ , i.e.,

$$(4.4) \quad R(A_0) \subset N(A_1).$$

Then also  $A_0^* A_1^* = 0$ , i.e.,  $R(A_1^*) \subset N(A_0^*)$ . From the Helmholtz type decompositions (4.1) for  $A = A_0$  and  $A = A_1$  we get in particular

$$(4.5) \quad \mathbf{H}_1 = \overline{R(A_0)} \oplus_{\mathbf{H}_1} N(A_0^*), \quad \mathbf{H}_1 = \overline{R(A_1^*)} \oplus_{\mathbf{H}_1} N(A_1),$$

and the following result for Helmholtz type decompositions:

**Lemma 4.4.** Let  $N_{0,1} := N(A_1) \cap N(A_0^*)$ . The refined Helmholtz type decompositions

$$(4.6) \quad N(A_1) = \overline{R(A_0)} \oplus_{\mathbf{H}_1} N_{0,1}, \quad D(A_1) = \overline{R(A_0)} \oplus_{\mathbf{H}_1} (D(A_1) \cap N(A_0^*)), \quad R(A_0) = R(\mathcal{A}_0),$$

$$(4.7) \quad N(A_0^*) = \overline{R(A_1^*)} \oplus_{\mathbf{H}_1} N_{0,1}, \quad D(A_0^*) = \overline{R(A_1^*)} \oplus_{\mathbf{H}_1} (D(A_0^*) \cap N(A_1)), \quad R(A_1^*) = R(\mathcal{A}_1^*),$$

and

$$(4.8) \quad \mathbf{H}_1 = \overline{R(A_0)} \oplus_{\mathbf{H}_1} N_{0,1} \oplus_{\mathbf{H}_1} \overline{R(A_1^*)}$$

hold, which can be further refined and specialized, e.g., to

$$(4.9) \quad \begin{aligned} D(\mathcal{A}_1) &= \overline{R(\mathcal{A}_0)} \oplus_{\mathbf{H}_1} N_{0,1} \oplus_{\mathbf{H}_1} D(\mathcal{A}_1), \\ D(\mathcal{A}_0^*) &= D(\mathcal{A}_0^*) \oplus_{\mathbf{H}_1} N_{0,1} \oplus_{\mathbf{H}_1} \overline{R(\mathcal{A}_1^*)}, \\ D(\mathcal{A}_1) \cap D(\mathcal{A}_0^*) &= D(\mathcal{A}_0^*) \oplus_{\mathbf{H}_1} N_{0,1} \oplus_{\mathbf{H}_1} D(\mathcal{A}_1). \end{aligned}$$

*Proof.* By (4.5) and the complex properties we see (4.6) and (4.7), yielding directly (4.8) and (4.9).  $\square$

We observe

$$\begin{aligned} D(\mathcal{A}_1) &= D(\mathcal{A}_1) \cap \overline{R(\mathcal{A}_1^*)} \subset D(\mathcal{A}_1) \cap N(\mathcal{A}_0^*) \subset D(\mathcal{A}_1) \cap D(\mathcal{A}_0^*), \\ D(\mathcal{A}_0^*) &= D(\mathcal{A}_0^*) \cap \overline{R(\mathcal{A}_0)} \subset D(\mathcal{A}_0^*) \cap N(\mathcal{A}_1) \subset D(\mathcal{A}_0^*) \cap D(\mathcal{A}_1), \end{aligned}$$

and using the refined Helmholtz type decompositions of Lemma 4.4 as well as the results of Lemma 4.1, Lemma 4.3, and Lemma 4.5, we immediately see:

**Lemma 4.5.** *The following assertions are equivalent:*

- (i)  $D(\mathcal{A}_0) \overset{\Leftrightarrow}{\mathbf{H}_0}$ ,  $D(\mathcal{A}_1) \overset{\Leftrightarrow}{\mathbf{H}_1}$ , and  $N_{0,1} \overset{\Leftrightarrow}{\mathbf{H}_1}$  are compact.
- (ii)  $D(\mathcal{A}_1) \cap D(\mathcal{A}_0^*) \overset{\Leftrightarrow}{\mathbf{H}_1}$  is compact.

In this case, the cohomology group  $N_{0,1}$  has finite dimension.

We summarize:

**Theorem 4.6.** *Let  $D(\mathcal{A}_1) \cap D(\mathcal{A}_0^*) \overset{\Leftrightarrow}{\mathbf{H}_1}$  be compact. Then  $D(\mathcal{A}_0) \overset{\Leftrightarrow}{\mathbf{H}_0}$ ,  $D(\mathcal{A}_1) \overset{\Leftrightarrow}{\mathbf{H}_1}$ , as well as  $D(\mathcal{A}_0^*) \overset{\Leftrightarrow}{\mathbf{H}_1}$ ,  $D(\mathcal{A}_1^*) \overset{\Leftrightarrow}{\mathbf{H}_2}$  are compact,  $\dim N_{0,1} < \infty$ , all ranges  $R(\mathcal{A}_0)$ ,  $R(\mathcal{A}_0^*)$ , and  $R(\mathcal{A}_1)$ ,  $R(\mathcal{A}_1^*)$  are closed, and the corresponding Friedrichs/Poincaré type estimates hold, i.e. there exists positive constants  $c_{\mathcal{A}_0}, c_{\mathcal{A}_1}$  such that*

$$(4.10) \quad \begin{aligned} \forall z \in D(\mathcal{A}_0) & \quad |z|_{\mathbf{H}_0} \leq c_{\mathcal{A}_0} |A_0 z|_{\mathbf{H}_1}, \\ \forall x \in D(\mathcal{A}_0^*) & \quad |x|_{\mathbf{H}_1} \leq c_{\mathcal{A}_0} |A_0^* x|_{\mathbf{H}_0}, \\ \forall x \in D(\mathcal{A}_1) & \quad |x|_{\mathbf{H}_1} \leq c_{\mathcal{A}_1} |A_1 x|_{\mathbf{H}_2}, \\ \forall y \in D(\mathcal{A}_1^*) & \quad |y|_{\mathbf{H}_2} \leq c_{\mathcal{A}_1} |A_1^* y|_{\mathbf{H}_1}. \end{aligned}$$

Moreover, all refined Helmholtz type decompositions of Lemma 4.4 hold with closed ranges, especially

$$(4.11) \quad D(\mathcal{A}_1) = R(\mathcal{A}_0) \oplus_{\mathbf{H}_1} (D(\mathcal{A}_1) \cap N(\mathcal{A}_0^*)).$$

*Proof.* Apply the latter lemmas and remarks to  $\mathbf{A} = \mathbf{A}_0$  and  $\mathbf{A} = \mathbf{A}_1$ .  $\square$

**4.2. The  $\mathbf{A}_0^*$ - $\mathbf{A}_1$ -Lemma.** Let  $\mathbf{A}_0$  and  $\mathbf{A}_1$  be as introduced before satisfying the complex property (4.4), i.e.,  $\mathbf{A}_1 \mathbf{A}_0 = 0$  or  $R(\mathbf{A}_0) \subset N(\mathbf{A}_1)$ . In other words, the primal and dual sequences

$$(4.12) \quad \begin{aligned} D(\mathbf{A}_0) \subset \mathbf{H}_0 & \xrightarrow{\mathbf{A}_0} D(\mathbf{A}_1) \subset \mathbf{H}_1 \xrightarrow{\mathbf{A}_1} \mathbf{H}_2, \\ \mathbf{H}_0 & \xleftarrow{\mathbf{A}_0^*} D(\mathbf{A}_0^*) \subset \mathbf{H}_1 \xleftarrow{\mathbf{A}_1^*} D(\mathbf{A}_1^*) \subset \mathbf{H}_2 \end{aligned}$$

are Hilbert complexes of closed and densely defined linear operators. The additional assumption that the ranges  $R(\mathbf{A}_0)$  and  $R(\mathbf{A}_1)$  are closed (and then also the ranges  $R(\mathbf{A}_0^*)$  and  $R(\mathbf{A}_1^*)$ ) is equivalent to the closedness of the Hilbert complexes. Moreover, the complexes are exact if and only if  $N_{0,1} = \{0\}$ .

As our main result, the following generalized global div-curl-lemma holds.

**Theorem 4.7** ( $\mathbf{A}_0^*$ - $\mathbf{A}_1$ -lemma). *Let  $D(\mathbf{A}_1) \cap D(\mathbf{A}_0^*) \overset{\Leftrightarrow}{\mathbf{H}_1}$  be compact. Moreover, let  $(x_n) \subset D(\mathbf{A}_1)$  and  $(y_n) \subset D(\mathbf{A}_0^*)$  be two sequences bounded in  $D(\mathbf{A}_1)$  and  $D(\mathbf{A}_0^*)$ , respectively. Then there exist  $x \in D(\mathbf{A}_1)$  and  $y \in D(\mathbf{A}_0^*)$  as well as subsequences, again denoted by  $(x_n)$  and  $(y_n)$ , such that*

- $x_n \rightharpoonup x$  in  $D(\mathbf{A}_1)$ ,
- $y_n \rightharpoonup y$  in  $D(\mathbf{A}_0^*)$ ,
- $\langle x_n, y_n \rangle_{\mathbf{H}_1} \rightarrow \langle x, y \rangle_{\mathbf{H}_1}$ .

*Proof.* Note that Theorem 4.6 can be applied. We pick subsequences, again denoted by  $(x_n)$  and  $(y_n)$ , such that  $(x_n)$  and  $(y_n)$  converge weakly in  $D(A_1)$  and  $D(A_0^*)$  to  $x \in D(A_1)$  and  $y \in D(A_0^*)$ , respectively. By (4.11) we get the orthogonal decomposition

$$D(A_1) \ni x_n = A_0 z_n + \tilde{x}_n, \quad z_n \in D(\mathcal{A}_0), \quad \tilde{x}_n \in D(A_1) \cap N(A_0^*).$$

$(z_n)$  is bounded in  $D(\mathcal{A}_0)$  by orthogonality and the Friedrichs/Poincaré type estimate (4.10).  $(\tilde{x}_n)$  is bounded in  $D(A_1) \cap D(A_0^*)$  by orthogonality and  $A_1 \tilde{x}_n = A_1 x_n$ ,  $A_0^* \tilde{x}_n = 0$ . Using the compact embeddings  $D(\mathcal{A}_0) \hookrightarrow H_0$  and  $D(A_1) \cap D(A_0^*) \hookrightarrow H_1$ , there exist  $z \in D(\mathcal{A}_0)$  and  $\tilde{x} \in D(A_1) \cap N(A_0^*)$  and we can extract two subsequences, again denoted by  $(z_n)$  and  $(\tilde{x}_n)$ , such that  $z_n \rightharpoonup z$  in  $D(\mathcal{A}_0)$  and  $z_n \rightarrow z$  in  $H_0$  as well as  $\tilde{x}_n \rightharpoonup \tilde{x}$  in  $D(A_1) \cap D(A_0^*)$  and  $\tilde{x}_n \rightarrow \tilde{x}$  in  $H_1$ . We have  $x = A_0 z + \tilde{x}$ , giving the Helmholtz type decomposition for  $x$ , as, e.g., for all  $\varphi \in H_1$

$$\langle x, \varphi \rangle_{H_1} \leftarrow \langle x_n, \varphi \rangle_{H_1} = \langle A_0 z_n, \varphi \rangle_{H_1} + \langle \tilde{x}_n, \varphi \rangle_{H_1} \rightarrow \langle A_0 z, \varphi \rangle_{H_1} + \langle \tilde{x}, \varphi \rangle_{H_1}.$$

Finally, we see

$$\begin{aligned} \langle x_n, y_n \rangle_{H_1} &= \langle A_0 z_n, y_n \rangle_{H_1} + \langle \tilde{x}_n, y_n \rangle_{H_1} = \langle z_n, A_0^* y_n \rangle_{H_0} + \langle \tilde{x}_n, y_n \rangle_{H_1} \\ &\rightarrow \langle z, A_0^* y \rangle_{H_0} + \langle \tilde{x}, y \rangle_{H_1} = \langle A_0 z, y \rangle_{H_1} + \langle \tilde{x}, y \rangle_{H_1} = \langle x, y \rangle_{H_1}, \end{aligned}$$

completing the proof.  $\square$

**4.3. Generalizations of the  $A_0^*$ - $A_1$ -Lemma.** In this section we present and discuss some variants of Theorem 4.7 using weaker assumptions, which are taken from the nice paper [35] of Marcus Waurick. We start with the following remarks.

**Remark 4.8.** *By Lemma 4.5 the crucial assumption, i.e.,  $D(A_1) \cap D(A_0^*) \hookrightarrow H_1$  is compact, holds, if and only if  $D(\mathcal{A}_0) \hookrightarrow H_0$ ,  $D(A_1) \hookrightarrow H_1$  are compact and  $N_{0,1}$  is finite-dimensional. Moreover, as Banach space adjoints we have*

$$H'_0 \hookrightarrow D(\mathcal{A}_0)' \quad \Leftrightarrow \quad D(\mathcal{A}_0) \hookrightarrow H_0 \quad \Leftrightarrow \quad D(\mathcal{A}_0^*) \hookrightarrow H_1 \quad \Leftrightarrow \quad H'_1 \hookrightarrow D(\mathcal{A}_0^*)',$$

and

$$H'_1 \hookrightarrow D(A_1)' \quad \Leftrightarrow \quad D(A_1) \hookrightarrow H_1 \quad \Leftrightarrow \quad D(A_1^*) \hookrightarrow H_2 \quad \Leftrightarrow \quad H'_2 \hookrightarrow D(A_1^*)'.$$

*In particular, the assumption on the compactness of  $D(A_1) \cap D(A_0^*) \hookrightarrow H_1$  is equivalent to the assumptions that  $\dim N_{0,1} < \infty$  and  $H_0 \cong H'_0 \hookrightarrow D(\mathcal{A}_0)'$ ,  $H_2 \cong H'_2 \hookrightarrow D(A_1^*)'$  are compact. Thus we observe that the assumptions of Theorem 4.7 are stronger but closely related to those of [35, Theorem 2.4]. Recall that by Theorem 4.6 both ranges  $R(A_0)$  and  $R(A_1)$  are closed and that  $\dim N_{0,1} < \infty$  if  $D(A_1) \cap D(A_0^*) \hookrightarrow H_1$  is compact. We emphasize that we have provided a different proof under stronger assumptions, which is from our personal point of view and taste easier and more canonical.*

Let us discuss the relations to [35], in particular to [35, Theorem 2.4], in more detail. First we note that Theorem 4.7 is equivalent to [35, Theorem 2.5] and that the assumptions of Theorem 4.7 are stronger but closely related to those of [35, Theorem 2.4].

A closer inspection of the proof of Theorem 4.7 shows that we can deal with slightly weaker assumptions. For this, let  $R(A_0)$  and  $R(A_1)$  be closed (which automatically would be implied by the compact embedding  $D(A_1) \cap D(A_0^*) \hookrightarrow H_1$ , see Theorem 4.6), and let  $(x_n) \subset D(A_1)$  and  $(y_n) \subset D(A_0^*)$  be two sequences bounded in  $H_1$ . By (4.9) we have

$$(4.13) \quad \begin{aligned} D(A_1) \ni x_n &= A_0 z_n + \hat{x}_n + A_1^* w_n \in R(\mathcal{A}_0) \oplus_{H_1} N_{0,1} \oplus_{H_1} D(A_1), \\ D(A_0^*) \ni y_n &= A_0 u_n + \hat{y}_n + A_1^* v_n \in D(\mathcal{A}_0^*) \oplus_{H_1} N_{0,1} \oplus_{H_1} R(A_1^*), \end{aligned}$$

with  $(z_n)$  and  $(v_n)$  bounded in  $D(\mathcal{A}_0)$  and  $D(A_1^*)$  by Lemma 4.1, respectively. W.l.o.g. we can assume that  $(z_n)$  and  $(v_n)$  already converge weakly in  $D(\mathcal{A}_0)$  and  $D(A_1^*)$ , respectively. Orthogonality shows

$$(4.14) \quad \begin{aligned} \langle x_n, y_n \rangle_{H_1} &= \langle A_0 z_n, y_n \rangle_{H_1} + \langle \hat{x}_n, \hat{y}_n \rangle_{H_1} + \langle x_n, A_1^* v_n \rangle_{H_1} \\ &= \langle z_n, A_0^* y_n \rangle_{H_0} + \langle \hat{x}_n, \hat{y}_n \rangle_{H_1} + \langle A_1 x_n, v_n \rangle_{H_2}. \end{aligned}$$

Hence, we observe that after extracting subsequences,  $(\langle x_n, y_n \rangle_{H_1})$  converges, provided that  $N_{0,1}$  is finite-dimensional and  $(A_0^* y_n)$  and  $(A_1 x_n)$  are relatively compact in  $D(\mathcal{A}_0)'$  and  $D(\mathcal{A}_1)'$ , respectively. This is almost the statement of [35, Theorem 2.4], still with stronger assumptions.

4.3.1. *More Generalizations.* The latter idea can be generalized and, indeed, in [35, Theorem 2.4] a more general situation is considered as  $(x_n) \subset D(A_1)$  and  $(y_n) \subset D(A_0^*)$  are not assumed to hold. In fact, these conditions are replaced by corresponding canonical distributional versions making the respective operators continuous on certain natural dual spaces. For this we need some preliminaries and new notations.

Dual pairs  $(A, A^*)$ ,  $(\mathcal{A}, \mathcal{A}^*)$  of densely defined and closed (unbounded) linear operators (as discussed in the latter sections) with domains of definitions  $D(A)$ ,  $D(\mathcal{A})$  and  $D(A^*)$ ,  $D(\mathcal{A}^*)$ , which are Hilbert spaces equipped with the respective graph norms, and closed ranges  $R(A) = R(\mathcal{A})$  and  $R(A^*) = R(\mathcal{A}^*)$  can also be considered as bounded linear operators. More precisely,

$$\begin{aligned} A : D(A) &\rightarrow H_2, & A^* : D(A^*) &\rightarrow H_1, \\ \mathcal{A} : D(\mathcal{A}) &\rightarrow R(\mathcal{A}) = R(A), & \mathcal{A}^* : D(\mathcal{A}^*) &\rightarrow R(\mathcal{A}^*) = R(A^*) \end{aligned}$$

are bounded with bounded Banach space adjoints

$$\begin{aligned} A' : H_2' &\rightarrow D(A)', & (A^*)' : H_1' &\rightarrow D(A^*)', \\ \mathcal{A}' : R(A)' &\rightarrow D(\mathcal{A})', & (\mathcal{A}^*)' : R(A^*)' &\rightarrow D(\mathcal{A}^*)', \end{aligned}$$

defined as usual by

$$\begin{aligned} A' y'(\varphi) &:= y'(A \varphi), & y' \in H_2', & \varphi \in D(A), \\ (A^*)' x'(\phi) &:= x'(A^* \phi), & x' \in H_1', & \phi \in D(A^*), \\ \mathcal{A}' y'(\varphi) &:= y'(\mathcal{A} \varphi), & y' \in R(A)', & \varphi \in D(\mathcal{A}), \\ (\mathcal{A}^*)' x'(\phi) &:= x'(\mathcal{A}^* \phi), & x' \in R(A^*)', & \phi \in D(\mathcal{A}^*). \end{aligned}$$

Moreover, we introduce the standard Riesz isomorphisms

$$\mathcal{R}_{H_n} : H_n \rightarrow H_n', \quad \mathcal{R}_{R(A)} : R(A) \rightarrow R(A)', \quad \mathcal{R}_{R(A^*)} : R(A^*) \rightarrow R(A^*)'$$

by  $x \mapsto \langle \cdot, x \rangle_{H_n}$ . Note that the closed ranges are itself Hilbert spaces with the inner products of  $H_n$ . Using the latter operators we define linear extensions of  $A$ ,  $\mathcal{A}$  and  $A^*$ ,  $\mathcal{A}^*$  by

$$\begin{aligned} \widetilde{A} &:= (A^*)' \mathcal{R}_{H_1} : H_1 \rightarrow D(A^*)', & \widetilde{A}^* &:= A' \mathcal{R}_{H_2} : H_2 \rightarrow D(A)', \\ \widetilde{\mathcal{A}} &:= (\mathcal{A}^*)' \mathcal{R}_{R(A^*)} : R(A^*) \rightarrow D(\mathcal{A}^*)', & \widetilde{\mathcal{A}}^* &:= \mathcal{A}' \mathcal{R}_{R(A)} : R(A) \rightarrow D(\mathcal{A})', \end{aligned}$$

with actions given by

$$\begin{aligned} \widetilde{A} x(\phi) &= (A^*)' \mathcal{R}_{H_1} x(\phi) = \mathcal{R}_{H_1} x(A^* \phi) = \langle A^* \phi, x \rangle_{H_1}, & x \in H_1, & \phi \in D(A^*), \\ \widetilde{\mathcal{A}} x(\phi) &= (\mathcal{A}^*)' \mathcal{R}_{R(A^*)} x(\phi) = \mathcal{R}_{R(A^*)} x(\mathcal{A}^* \phi) = \langle \mathcal{A}^* \phi, x \rangle_{H_1}, & x \in R(A^*), & \phi \in D(\mathcal{A}^*), \\ \widetilde{A}^* y(\varphi) &= A' \mathcal{R}_{H_2} y(\varphi) = \mathcal{R}_{H_2} y(A \varphi) = \langle A \varphi, y \rangle_{H_2}, & y \in H_2, & \varphi \in D(A), \\ \widetilde{\mathcal{A}}^* y(\varphi) &= \mathcal{A}' \mathcal{R}_{R(A)} y(\varphi) = \mathcal{R}_{R(A)} y(\mathcal{A} \varphi) = \langle \mathcal{A} \varphi, y \rangle_{H_2}, & y \in R(A), & \varphi \in D(\mathcal{A}). \end{aligned}$$

Introducing the canonical embeddings and their adjoints

$$\begin{aligned} \iota_{D(A)} : D(A) &\hookrightarrow H_1, & \iota'_{D(A)} : H_1' &\hookrightarrow D(A)', \\ \iota_{D(\mathcal{A})} : D(\mathcal{A}) &\hookrightarrow R(A^*), & \iota'_{D(\mathcal{A})} : R(A^*)' &\hookrightarrow D(\mathcal{A})', \\ \iota_{D(A^*)} : D(A^*) &\hookrightarrow H_2, & \iota'_{D(A^*)} : H_2' &\hookrightarrow D(A^*)', \\ \iota_{D(\mathcal{A}^*)} : D(\mathcal{A}^*) &\hookrightarrow R(A), & \iota'_{D(\mathcal{A}^*)} : R(A)' &\hookrightarrow D(\mathcal{A}^*)' \end{aligned}$$

we emphasize that for all  $x \in D(A)$  and for all  $\phi \in D(A^*)$

$$\widetilde{A}x(\phi) = \langle A^* \phi, x \rangle_{H_1} = \langle \phi, Ax \rangle_{H_2} = \langle \iota_{D(A^*)} \phi, Ax \rangle_{H_2} = \mathcal{R}_{H_2} Ax(\iota_{D(A^*)} \phi) = \iota'_{D(A^*)} \mathcal{R}_{H_2} Ax(\phi)$$

holds and therefore

$$\widetilde{A}|_{D(A)} := \widetilde{A} \iota_{D(A)} = \iota'_{D(A^*)} \mathcal{R}_{H_2} A : D(A) \rightarrow D(A^*)'.$$

Thus, in this sense,  $\widetilde{A}$  is indeed an extension of  $A$ . In the same way we see that

$$\widetilde{A}|_{D(A)} = \iota'_{D(A^*)} \mathcal{R}_{R(A)} A, \quad \widetilde{A}^*|_{D(A^*)} = \iota'_{D(A)} \mathcal{R}_{H_1} A^*, \quad \widetilde{A}^*|_{D(A^*)} = \iota'_{D(A)} \mathcal{R}_{R(A^*)} A^*$$

are extensions as well.

**Lemma 4.9** ([35, Theorem 2.2]). *Let  $R(A)$  be closed. Then*

- (i)  $\mathcal{A}, (\mathcal{A}^*)', \widetilde{\mathcal{A}}$  are topological isomorphisms,
- (i\*)  $\mathcal{A}^*, \mathcal{A}', \widetilde{\mathcal{A}}^*$  are topological isomorphisms,
- (ii)  $N(\widetilde{\mathcal{A}}) = N(A)$ ,
- (ii\*)  $N(\widetilde{\mathcal{A}}^*) = N(A^*)$ ,
- (iii)  $\mathcal{A}', \mathcal{A}^*$  are surjective if and only if  $N(A) = 0$ ,
- (iii\*)  $(\mathcal{A}^*)', \widetilde{\mathcal{A}}$  are surjective if and only if  $N(A^*) = 0$ .

*Proof.*  $\mathcal{A}$  and  $\mathcal{A}^*$  are a topological isomorphisms by the bounded inverse theorem or the considerations from the previous sections. If  $\mathcal{A}'y' = 0$  for  $y' \in R(A)'$ , then  $\mathcal{A}'y'(z) = \widetilde{y}'(\mathcal{A}z) = 0$  for all  $z \in D(\mathcal{A})$ . Hence  $y' = 0$  on  $R(\mathcal{A}) = R(A)$ , i.e.,  $y' = 0$ . Thus  $\mathcal{A}'$  is injective and so is  $\widetilde{\mathcal{A}}^* = \mathcal{A}' \mathcal{R}_{R(A)}$  as  $\mathcal{R}_{R(A)}$  is an isomorphism. For  $f \in D(\mathcal{A})'$  we obtain by Riesz' representation theorem a unique  $z \in D(\mathcal{A})$  such that

$$\forall \varphi \in D(\mathcal{A}) \quad \langle A\varphi, Az \rangle_{H_2} = f(\varphi).$$

Note that  $\langle A \cdot, A \cdot \rangle_{H_2}$  is an inner product for  $D(\mathcal{A})$  by Lemma 4.1. Thus with  $y := Az \in R(A)$  we see

$$\forall \varphi \in D(\mathcal{A}) \quad f(\varphi) = \langle A\varphi, y \rangle_{H_2} = \widetilde{\mathcal{A}}^* y(\varphi),$$

i.e.,  $f = \widetilde{\mathcal{A}}^* y$ . Hence  $\widetilde{\mathcal{A}}^*$  is surjective and so is  $\mathcal{A}' = \widetilde{\mathcal{A}}^* \mathcal{R}_{R(A)}^{-1}$  as  $\mathcal{R}_{R(A)}$  is an isomorphism. By the bounded inverse theorem both  $\mathcal{A}'$  and  $\widetilde{\mathcal{A}}^*$  are topological isomorphisms. Analogously we show the assertions for  $(\mathcal{A}^*)'$  and  $\widetilde{\mathcal{A}}$ , which shows (i) and (i\*). For (ii) we observe  $x \in N(\widetilde{\mathcal{A}})$  if and only if

$$\forall \phi \in D(A^*) \quad 0 = \widetilde{A}x(\phi) = \langle A^* \phi, x \rangle_{H_1},$$

if and only if  $x \in N(A)$ . Similarly we see  $N(\widetilde{\mathcal{A}}^*) = N(A^*)$ , proving (ii\*). Let  $N(A) = \{0\}$  and  $f \in D(\mathcal{A})'$ . Then  $D(\mathcal{A}) = D(A)$  and following the argument for  $\mathcal{A}'$  from above we obtain  $y \in R(A) \subset H_2$  with  $f = \widetilde{\mathcal{A}}^* y$ . Hence  $\widetilde{\mathcal{A}}^*$  is surjective and so is  $\mathcal{A}' = \widetilde{\mathcal{A}}^* \mathcal{R}_{H_2}^{-1}$  as  $\mathcal{R}_{H_2}$  is an isomorphism. On the other hand,  $\widetilde{\mathcal{A}}^*$  is surjective if and only if  $\mathcal{A}'$  is surjective, and in this case for any  $\varphi \in N(A)$  we can represent  $f := \iota'_{D(A)} \mathcal{R}_{H_1} \iota_{N(A)} \varphi \in D(\mathcal{A})'$  by  $\widetilde{\mathcal{A}}^* y = f$  with some  $y \in H_2$ . Hence

$$0 = \langle A\varphi, y \rangle_{H_2} = \widetilde{\mathcal{A}}^* y(\varphi) = f(\varphi) = \mathcal{R}_{H_1} \iota_{N(A)} \varphi(\iota_{D(A)} \varphi) = \langle \iota_{D(A)} \varphi, \iota_{N(A)} \varphi \rangle_{H_1} = \langle \varphi, \varphi \rangle_{H_1},$$

showing  $N(A) = \{0\}$ , i.e., (iii). Analogously, we show (iii\*) for  $(\mathcal{A}^*)'$  and  $\widetilde{\mathcal{A}}$ , completing the proof.  $\square$

**Remark 4.10.** *Another, even shorter proof using annihilators is possible. It holds*

$$N(\mathcal{A}') = R(\mathcal{A})^\circ = \{0\}, \quad R(\mathcal{A}') = N(\mathcal{A})^\circ = \{0\}^\circ = D(\mathcal{A})',$$

the latter by the closed range theorem. Hence  $\mathcal{A}'$  is a topological isomorphism by the bounded inverse theorem. The same applies to  $(\mathcal{A}^*)'$ . The Riesz mappings are topological isomorphisms, so are  $\tilde{\mathcal{A}}$ ,  $\tilde{\mathcal{A}}^*$ . Moreover,

$$R(\tilde{\mathcal{A}}^*) = R(\mathcal{A}') = N(\mathcal{A})^\circ, \quad R(\tilde{\mathcal{A}}) = R((\mathcal{A}^*)') = N(\mathcal{A}^*)^\circ.$$

Note that also  $N(\mathcal{A}') = R(\mathcal{A})^\circ$  and  $N((\mathcal{A}^*)') = R(\mathcal{A}^*)^\circ$  hold.

Using Hilbert space adjoints we introduce the canonical embeddings and projections

$$\begin{aligned} \iota_{R(\mathcal{A})} : R(\mathcal{A}) &\rightarrow \mathbf{H}_2, & \iota_{R(\mathcal{A})}^* : \mathbf{H}_2 &\rightarrow R(\mathcal{A}), & \pi_{R(\mathcal{A})} &:= \iota_{R(\mathcal{A})} \iota_{R(\mathcal{A})}^* : \mathbf{H}_2 \rightarrow \mathbf{H}_2, \\ \iota_{R(\mathcal{A}^*)} : R(\mathcal{A}^*) &\rightarrow \mathbf{H}_1, & \iota_{R(\mathcal{A}^*)}^* : \mathbf{H}_1 &\rightarrow R(\mathcal{A}^*), & \pi_{R(\mathcal{A}^*)} &:= \iota_{R(\mathcal{A}^*)} \iota_{R(\mathcal{A}^*)}^* : \mathbf{H}_1 \rightarrow \mathbf{H}_1. \end{aligned}$$

**Remark 4.11.** Indeed,  $\pi_{R(\mathcal{A})}$  and  $\pi_{R(\mathcal{A}^*)}$  are the corresponding projections. To see this, let us consider, e.g.,  $\pi_{R(\mathcal{A})}$ . For  $x \in D(\iota_{R(\mathcal{A})}^*) = \mathbf{H}_2$  with  $\iota_{R(\mathcal{A})}^* x \in R(\mathcal{A})$  and all  $\phi \in D(\iota_{R(\mathcal{A})}) = R(\mathcal{A})$  it holds

$$\langle \phi, x \rangle_{\mathbf{H}_2} = \langle \iota_{R(\mathcal{A})} \phi, x \rangle_{\mathbf{H}_2} = \langle \phi, \iota_{R(\mathcal{A})}^* x \rangle_{R(\mathcal{A})} = \langle \phi, \pi_{R(\mathcal{A})} x \rangle_{\mathbf{H}_2}.$$

Hence  $\pi_{R(\mathcal{A})} x \in R(\mathcal{A})$  and  $(1 - \pi_{R(\mathcal{A})})x \in R(\mathcal{A})^\perp_{\mathbf{H}_2}$ . Moreover, since  $\pi_{R(\mathcal{A})} x \in D(\iota_{R(\mathcal{A})}^*) = \mathbf{H}_2$  the latter computation shows for all  $\phi \in R(\mathcal{A})$

$$\langle \phi, x \rangle_{\mathbf{H}_2} = \langle \phi, \pi_{R(\mathcal{A})} x \rangle_{\mathbf{H}_2} = \langle \phi, \pi_{R(\mathcal{A})} \pi_{R(\mathcal{A})} x \rangle_{\mathbf{H}_2},$$

i.e.,  $\pi_{R(\mathcal{A})} \pi_{R(\mathcal{A})} x = \pi_{R(\mathcal{A})} x$  on  $R(\mathcal{A})$ . Finally,  $\pi_{R(\mathcal{A})}$  is self-adjoint.

Furthermore, we need

$$\begin{aligned} \iota_{R(\mathcal{A})}^* \iota_{D(\mathcal{A}^*)} : D(\mathcal{A}^*) &\rightarrow D(\mathcal{A}^*), & (\iota_{R(\mathcal{A})}^* \iota_{D(\mathcal{A}^*)})' : D(\mathcal{A}^*)' &\rightarrow D(\mathcal{A}^*)', \\ \iota_{R(\mathcal{A}^*)}^* \iota_{D(\mathcal{A})} : D(\mathcal{A}) &\rightarrow D(\mathcal{A}), & (\iota_{R(\mathcal{A}^*)}^* \iota_{D(\mathcal{A})})' : D(\mathcal{A})' &\rightarrow D(\mathcal{A})'. \end{aligned}$$

We also emphasize that for  $x \in \mathbf{H}_1$  it holds  $(1 - \pi_{R(\mathcal{A}^*)})x \in R(\mathcal{A}^*)^\perp_{\mathbf{H}_1} = N(\mathcal{A})$  and thus

$$x = \pi_{R(\mathcal{A}^*)} x + (1 - \pi_{R(\mathcal{A}^*)})x \in R(\mathcal{A}^*) \oplus_{\mathbf{H}_1} N(\mathcal{A})$$

is the Helmholtz decomposition for  $x$ . Analogously for  $y \in \mathbf{H}_2$  the Helmholtz decomposition is given by

$$y = \pi_{R(\mathcal{A})} y + (1 - \pi_{R(\mathcal{A})})y \in R(\mathcal{A}) \oplus_{\mathbf{H}_2} N(\mathcal{A}^*).$$

Hence for  $x \in D(\mathcal{A})$  and  $y \in D(\mathcal{A}^*)$  we identify

$$(4.15) \quad \pi_{R(\mathcal{A}^*)} x = \iota_{R(\mathcal{A}^*)}^* \iota_{D(\mathcal{A})} x \in D(\mathcal{A}), \quad \pi_{R(\mathcal{A})} y = \iota_{R(\mathcal{A})}^* \iota_{D(\mathcal{A}^*)} y \in D(\mathcal{A}^*).$$

**Lemma 4.12.** Let  $R(\mathcal{A})$  be closed. Then

- (i)  $\tilde{\mathcal{A}} = (\iota_{R(\mathcal{A})}^* \iota_{D(\mathcal{A}^*)})' \tilde{\mathcal{A}} \iota_{R(\mathcal{A}^*)}^*$  and
 
$$|\tilde{\mathcal{A}} x|_{D(\mathcal{A}^*)'} = |\tilde{\mathcal{A}} \iota_{R(\mathcal{A}^*)}^* x|_{D(\mathcal{A}^*)'}, \quad |\tilde{\mathcal{A}}|_{\mathbf{H}_1 \rightarrow D(\mathcal{A}^*)'} = |\tilde{\mathcal{A}}|_{R(\mathcal{A}^*) \rightarrow D(\mathcal{A}^*)'},$$
- (ii)  $\tilde{\mathcal{A}}^* = (\iota_{R(\mathcal{A}^*)}^* \iota_{D(\mathcal{A})})' \tilde{\mathcal{A}}^* \iota_{R(\mathcal{A})}^*$  and
 
$$|\tilde{\mathcal{A}}^* x|_{D(\mathcal{A})'} = |\tilde{\mathcal{A}}^* \iota_{R(\mathcal{A})}^* x|_{D(\mathcal{A})'}, \quad |\tilde{\mathcal{A}}^*|_{\mathbf{H}_2 \rightarrow D(\mathcal{A})'} = |\tilde{\mathcal{A}}^*|_{R(\mathcal{A}) \rightarrow D(\mathcal{A})'}.$$

*Proof.* For  $x \in \mathbf{H}_1$  and  $\phi \in D(\mathcal{A}^*)$  we have  $\pi_{R(\mathcal{A})} \phi = \iota_{R(\mathcal{A})}^* \iota_{D(\mathcal{A}^*)} \phi \in D(\mathcal{A}^*)$  and

$$\begin{aligned} \tilde{\mathcal{A}} x(\phi) &= \langle \mathcal{A}^* \phi, x \rangle_{\mathbf{H}_1} = \langle \pi_{R(\mathcal{A}^*)} \mathcal{A}^* \pi_{R(\mathcal{A})} \phi, x \rangle_{\mathbf{H}_1} = \langle \mathcal{A}^* \pi_{R(\mathcal{A})} \phi, \pi_{R(\mathcal{A}^*)} x \rangle_{\mathbf{H}_1} \\ &= \langle \mathcal{A}^* \iota_{R(\mathcal{A})}^* \iota_{D(\mathcal{A}^*)} \phi, \iota_{R(\mathcal{A}^*)}^* x \rangle_{R(\mathcal{A}^*)} = \tilde{\mathcal{A}} \iota_{R(\mathcal{A}^*)}^* x (\iota_{R(\mathcal{A})}^* \iota_{D(\mathcal{A}^*)} \phi) \\ &= (\iota_{R(\mathcal{A})}^* \iota_{D(\mathcal{A}^*)})' \tilde{\mathcal{A}} \iota_{R(\mathcal{A}^*)}^* x(\phi). \end{aligned}$$

Moreover, by the latter computations for  $x \in \mathbf{H}_1$

$$\begin{aligned} |\tilde{\mathcal{A}}x|_{D(A^*)'} &= \sup_{\substack{\phi \in D(A^*) \\ |\phi|_{D(A^*)} \leq 1}} \langle \tilde{\mathcal{A}}^* \phi, x \rangle_{\mathbf{H}_1} = \sup_{\substack{\phi \in D(A^*) \\ |\phi|_{D(A^*)} \leq 1}} \langle \tilde{\mathcal{A}}^* \pi_{R(A)} \phi, \pi_{R(A^*)} x \rangle_{\mathbf{H}_1} \\ &= \sup_{\substack{\psi \in D(A^*) \\ |\psi|_{D(A^*)} \leq 1}} \langle \tilde{\mathcal{A}}^* \psi, \iota_{R(A^*)}^* x \rangle_{\mathbf{H}_1} = |\tilde{\mathcal{A}} \iota_{R(A^*)}^* x|_{D(A^*)'}, \end{aligned}$$

and thus

$$|\tilde{\mathcal{A}}|_{\mathbf{H}_1 \rightarrow D(A^*)'} = \sup_{\substack{x \in \mathbf{H}_1 \\ |x|_{\mathbf{H}_1} \leq 1}} |\tilde{\mathcal{A}}x|_{D(A^*)'} = \sup_{\substack{x \in \mathbf{H}_1 \\ |x|_{\mathbf{H}_1} \leq 1}} |\tilde{\mathcal{A}} \iota_{R(A^*)}^* x|_{D(A^*)'} = \sup_{\substack{z \in R(A^*) \\ |z|_{\mathbf{H}_1} \leq 1}} |\tilde{\mathcal{A}}z|_{D(A^*)'} = |\tilde{\mathcal{A}}|_{R(A^*) \rightarrow D(A^*)'}.$$

The assertions in (ii) follow analogously.  $\square$

The next result from [35] is crucial for the further considerations. We give a slightly modified version.

**Lemma 4.13** ([35, Corollary 2.6]). *Let  $R(A)$  be closed.*

(i) *For  $(x_n) \subset \mathbf{H}_1$  the following statements are equivalent:*

- (i<sub>1</sub>)  $(\tilde{\mathcal{A}}x_n)$  *is relatively compact in  $D(A^*)'$ .*
- (i<sub>2</sub>)  $(\tilde{\mathcal{A}} \iota_{R(A^*)}^* x_n)$  *is relatively compact in  $D(A^*)'$ .*
- (i<sub>3</sub>)  $(\iota_{R(A^*)}^* x_n)$  *is relatively compact in  $R(A^*)$ .*
- (i<sub>4</sub>)  $(\pi_{R(A^*)} x_n)$  *is relatively compact in  $\mathbf{H}_1$ .*
- (i<sub>5</sub>)  $(\mathcal{R}_{R(A^*)} \iota_{R(A^*)}^* x_n)$  *is relatively compact in  $R(A^*)'$ .*

*If  $x_n \rightharpoonup x \in \mathbf{H}_1$  in  $\mathbf{H}_1$ , then either of the latter conditions (i<sub>1</sub>)-(i<sub>5</sub>) implies  $\iota_{R(A^*)}^* x_n \rightarrow \iota_{R(A^*)}^* x$  in  $R(A^*)$  and  $\pi_{R(A^*)} x_n \rightarrow \pi_{R(A^*)} x$  in  $\mathbf{H}_1$ .*

(ii) *For  $(y_n) \subset \mathbf{H}_2$  the following statements are equivalent:*

- (ii<sub>1</sub>)  $(\tilde{\mathcal{A}}^* y_n)$  *is relatively compact in  $D(A)'$ .*
- (ii<sub>2</sub>)  $(\tilde{\mathcal{A}}^* \iota_{R(A)}^* y_n)$  *is relatively compact in  $D(A)'$ .*
- (ii<sub>3</sub>)  $(\iota_{R(A)}^* y_n)$  *is relatively compact in  $R(A)$ .*
- (ii<sub>4</sub>)  $(\pi_{R(A)} y_n)$  *is relatively compact in  $\mathbf{H}_2$ .*
- (ii<sub>5</sub>)  $(\mathcal{R}_{R(A)} \iota_{R(A)}^* y_n)$  *is relatively compact in  $R(A)'$ .*

*If  $y_n \rightharpoonup x \in \mathbf{H}_2$  in  $\mathbf{H}_2$ , then either of the latter conditions (ii<sub>1</sub>)-(ii<sub>5</sub>) implies  $\iota_{R(A)}^* y_n \rightarrow \iota_{R(A)}^* y$  in  $R(A)$  and  $\pi_{R(A)} y_n \rightarrow \pi_{R(A)} y$  in  $\mathbf{H}_2$ .*

*Proof.* By Lemma 4.9 (i)  $\tilde{\mathcal{A}} = (\mathcal{A}^*)' \mathcal{R}_{R(A^*)} : R(A^*) \rightarrow D(\mathcal{A}^*)'$  is a topological isomorphism. Hence (i<sub>2</sub>)-(i<sub>5</sub>) are equivalent. The equivalence of (i<sub>1</sub>) and (i<sub>2</sub>) follows by Lemma 4.12 (i). If  $x_n \rightharpoonup x$  in  $\mathbf{H}_1$ , then  $\iota_{R(A^*)}^* x_n \rightharpoonup \iota_{R(A^*)}^* x$  in  $R(A^*)$  and  $\pi_{R(A^*)} x_n \rightharpoonup \pi_{R(A^*)} x$  in  $\mathbf{H}_1$ . By a subsequence argument we see that, e.g., (i<sub>3</sub>) implies  $\iota_{R(A^*)}^* x_n \rightarrow \iota_{R(A^*)}^* x$  in  $R(A^*)$  and hence  $\pi_{R(A^*)} x_n \rightarrow \pi_{R(A^*)} x$  in  $\mathbf{H}_1$ . Analogously we show (ii).  $\square$

With this latter key observation we can prove a general (distributional)  $A_0^*A_1$ -lemma. For this, we introducing two bounded linear operators  $A_0 : D(A_0) \rightarrow \mathbf{H}_1$ ,  $A_1 : D(A_1) \rightarrow \mathbf{H}_2$  satisfying the complex property  $A_1 A_0 = 0$  and recall the linear extensions of  $A_1$ ,  $\mathcal{A}_1$  and  $A_0^*$ ,  $\mathcal{A}_0^*$

$$\begin{aligned} \tilde{A}_1 &:= (A_1^*)' \mathcal{R}_{\mathbf{H}_1} : \mathbf{H}_1 \rightarrow D(A_1^*)', & \tilde{A}_0^* &:= A_0' \mathcal{R}_{\mathbf{H}_1} : \mathbf{H}_1 \rightarrow D(A_0)', \\ \tilde{\mathcal{A}}_1 &:= (\mathcal{A}_1^*)' \mathcal{R}_{R(A_1^*)} : R(A_1^*) \rightarrow D(\mathcal{A}_1^*)', & \tilde{\mathcal{A}}_0^* &:= \mathcal{A}_0' \mathcal{R}_{R(A_0)} : R(A_0) \rightarrow D(\mathcal{A}_0)'. \end{aligned}$$

**Theorem 4.14** (generalized  $A_0^*A_1$ -lemma, [35, Theorem 2.4]). *Let the ranges  $R(A_0)$  and  $R(A_1)$  be closed and let  $N_{0,1}$  be finite-dimensional. Moreover, let  $(x_n), (y_n) \subset \mathbf{H}_1$  be two bounded sequences such that*

- $(\widetilde{A}_1 x_n)$  is relatively compact in  $D(A_1^*)'$ ,
- $(\widetilde{A}_0^* y_n)$  is relatively compact in  $D(A_0)'$ .

Then there exist  $x, y \in H_1$  as well as subsequences, again denoted by  $(x_n)$  and  $(y_n)$ , such that

- $x_n \rightharpoonup x$  in  $H_1$ ,
- $y_n \rightharpoonup y$  in  $H_1$ ,
- $\langle x_n, y_n \rangle_{H_1} \rightarrow \langle x, y \rangle_{H_1}$ .

**Remark 4.15.** By Lemma 4.13 the assumptions on the relative compactness can be replaced equivalently by the assumptions that  $(\widetilde{A}_1 \iota_{R(A_1^*)}^* x_n)$  is relatively compact in  $D(A_1^*)'$  and that  $(\widetilde{A}_0^* \iota_{R(A_0)}^* y_n)$  is relatively compact in  $D(A_0)'$ .

*Proof of Theorem 4.14.* Let  $(x_n), (y_n) \subset H_1$  be two bounded sequences. W.l.o.g. let  $x_n \rightharpoonup x$  and  $y_n \rightharpoonup y$  in  $H_1$ . By Lemma 4.13  $\pi_{R(A_1^*)} x_n \rightarrow \pi_{R(A_1^*)} x$  and  $\pi_{R(A_0)} y_n \rightarrow \pi_{R(A_0)} y$  in  $H_1$ . By Lemma 4.4, in particular (4.8) (compare to (4.13)), we have the Helmholtz decompositions

$$(4.16) \quad \begin{aligned} x_n &= \pi_{R(A_0)} x_n + \pi_{N_{0,1}} x_n + \pi_{R(A_1^*)} x_n \in R(A_0) \oplus_{H_1} N_{0,1} \oplus_{H_1} R(A_1^*), \\ y_n &= \pi_{R(A_0)} y_n + \pi_{N_{0,1}} y_n + \pi_{R(A_1^*)} y_n \in R(A_0) \oplus_{H_1} N_{0,1} \oplus_{H_1} R(A_1^*), \end{aligned}$$

yielding (compare to (4.14))

$$(4.17) \quad \langle x_n, y_n \rangle_{H_1} = \langle \pi_{R(A_1^*)} x_n, y_n \rangle_{H_1} + \langle \pi_{N_{0,1}} x_n, y_n \rangle_{H_1} + \langle x_n, \pi_{R(A_0)} y_n \rangle_{H_1}.$$

Similar to (4.16) we can decompose  $x$  and  $y$  and w.l.o.g. we can assume that  $\pi_{N_{0,1}} x_n \rightarrow \pi_{N_{0,1}} x$  as  $N_{0,1}$  has finite dimension. Finally it follows

$$\langle x_n, y_n \rangle_{H_1} \rightarrow \langle \pi_{R(A_1^*)} x, y \rangle_{H_1} + \langle \pi_{N_{0,1}} x, y \rangle_{H_1} + \langle x, \pi_{R(A_0)} y \rangle_{H_1} = \langle x, y \rangle_{H_1},$$

completing the proof.  $\square$

Now, we make the connection to Theorem 4.7 and show that the assumptions in Theorem 4.7 imply those of Theorem 4.14.

**Lemma 4.16** ([35, Corollary 2.7]). *Let either  $A: D(A) \subset H_1 \rightarrow H_2$  be a densely defined and closed linear operator or  $A: D(A) \rightarrow H_2$  be a continuous linear operator. Moreover, let  $D(A) \Leftrightarrow H_1$  be compact.*

- (i) *Let  $(x_n) \subset D(A)$  be bounded in  $D(A)$ . Then  $(\pi_{R(A^*)} x_n)$  is relatively compact in  $H_1$ . Equivalently,  $(\widetilde{A} x_n)$  is relatively compact in  $D(A^*)'$ .*
- (ii) *Let  $(y_n) \subset D(A^*)$  be bounded in  $D(A^*)$ . Then  $(\pi_{R(A)} y_n)$  is relatively compact in  $H_2$ . Equivalently,  $(\widetilde{A}^* y_n)$  is relatively compact in  $D(A)'$ .*

*Proof.* By Lemma 4.3 the compactness of  $D(A) \Leftrightarrow H_1$  yields the closedness of  $R(A)$ . Hence Lemma 4.13 is applicable. Let  $(x_n) \subset D(A)$  be bounded in  $D(A)$ . Then by (4.15), see also (4.2),  $(\pi_{R(A^*)} x_n) \subset D(A)$  is bounded in  $D(A)$ . Hence it contains a subsequence converging in  $H_1$ . Lemma 4.13 shows the equivalence to the second relative compactness. Analogously we prove the assertions in (ii).  $\square$

For two linear operators  $A_0$  and  $A_1$  as in Lemma 4.16, i.e., bounded or unbounded, densely defined and closed, satisfying the complex property  $A_1 A_0 = 0$  we obtain the following results.

**Lemma 4.17.** *Let  $D(A_1) \cap D(A_0^*) \Leftrightarrow H_1$  be compact. Moreover, let  $(x_n) \subset D(A_1)$  and  $(y_n) \subset D(A_0^*)$  be two sequences bounded in  $D(A_1)$  and  $D(A_0^*)$ , respectively. Then:*

- (i)  *$(\pi_{R(A_1^*)} x_n)$  is relatively compact in  $H_1$ . Equivalently,  $(\widetilde{A}_1 x_n)$  is relatively compact in  $D(A_1^*)'$ .*
- (ii)  *$(\pi_{R(A_0)} y_n)$  is relatively compact in  $H_1$ . Equivalently,  $(\widetilde{A}_0^* y_n)$  is relatively compact in  $D(A_0)'$ .*
- (iii)  *$(\pi_{N_{0,1}} x_n)$  and  $(\pi_{N_{0,1}} y_n)$  are relatively compact in  $H_1$ .*

*Proof.* By Lemma 4.5  $D(A_0) \Leftrightarrow H_0$ ,  $D(A_1) \Leftrightarrow H_1$ ,  $N_{0,1} \Leftrightarrow H_1$  are compact, in particular,  $N_{0,1}$  is finite dimensional, showing (iii). Lemma 4.16 yields (i) and (ii).  $\square$

**Remark 4.18.** By Lemma 4.3 and Lemma 4.5 the compactness of  $D(A_1) \cap D(A_0^*) \Leftrightarrow H_1$  implies the closedness of the ranges  $R(A_0)$  and  $R(A_1)$  and the finite dimensionality of  $N_{0,1}$ . Thus Lemma 4.17 shows that the proof of Theorem 4.14 provides another and different proof for Theorem 4.7.

The above considerations lead to the following insight, which is interesting on its own right.

**Lemma 4.19.** Let  $R(A_0)$  and  $R(A_1)$  be closed. For a sequence  $(x_n) \subset H_1$  the following assertions are equivalent:

(i)  $(x_n)$  is relatively compact in  $H_1$ .

(ii)  $(\pi_{R(A_1^*)}x_n)$ ,  $(\pi_{R(A_0)}x_n)$ , and  $(\pi_{N_{0,1}}x_n)$  are relatively compact in  $H_1$ .

(iii)  $(\widetilde{A_0^*}x_n)$ ,  $(\widetilde{A_1}x_n)$ , and  $(\pi_{N_{0,1}}x_n)$  are relatively compact in  $D(A_0)'$ ,  $D(A_1^*)'$ , and  $H_1$ , respectively.

Moreover, if  $(x_n) \subset D(A_1) \cap D(A_0^*)$  is bounded in  $D(A_1) \cap D(A_0^*)$  and  $D(A_1) \cap D(A_0^*) \Leftrightarrow H_1$  is compact, then (i), (ii), and (iii) hold.

*Proof.* By the continuity of the projections and the Helmholtz decompositions (4.16), i.e.,

$$x_n = \pi_{R(A_0)}x_n + \pi_{N_{0,1}}x_n + \pi_{R(A_1^*)}x_n \in R(A_0) \oplus_{H_1} N_{0,1} \oplus_{H_1} R(A_1^*),$$

the relative compactness of  $(x_n)$  in  $H_1$  is equivalent to (ii), which is equivalent to (iii) by Lemma 4.13. The last assertion follows by definition.  $\square$

## 5. APPLICATIONS

Whenever closed Hilbert complexes like (4.12) together with the corresponding compact embedding  $D(A_1) \cap D(A_0^*) \Leftrightarrow H_1$  occur, we can apply the general  $A_0^*$ - $A_1$ -lemma, i.e., Theorem 4.7. In three dimensions we typically have three closed and densely defined linear operators  $A_0$ ,  $A_1$ , and  $A_2$ , satisfying the complex properties  $R(A_0) \subset N(A_1)$  and  $R(A_1) \subset N(A_2)$ , i.e.,

$$(5.1) \quad \begin{aligned} D(A_0) \subset H_0 &\xrightarrow{A_0} D(A_1) \subset H_1 \xrightarrow{A_1} D(A_2) \subset H_2 \xrightarrow{A_2} H_3, \\ H_0 &\xleftarrow{A_0^*} D(A_0^*) \subset H_1 \xleftarrow{A_1^*} D(A_1^*) \subset H_2 \xleftarrow{A_2^*} D(A_2^*) \subset H_3, \end{aligned}$$

together with the crucial compact embeddings

$$(5.2) \quad D(A_1) \cap D(A_0^*) \Leftrightarrow H_1, \quad D(A_2) \cap D(A_1^*) \Leftrightarrow H_2.$$

With slightly weaker assumptions we can apply Theorem 4.14.

Recalling our general assumptions on the underlying domain from Section 2, throughout this application section  $\Omega$  can be a

- weak Lipschitz domain with boundary  $\Gamma$ ,
- weak Lipschitz domain with boundary  $\Gamma$  and weak Lipschitz interfaces  $\Gamma_t$  and  $\Gamma_n$ ,
- strong Lipschitz domain with boundary  $\Gamma$ ,
- strong Lipschitz domain with boundary  $\Gamma$  and strong Lipschitz interfaces  $\Gamma_t$  and  $\Gamma_n$ .

We extend this definition to  $\Omega \subset \mathbb{R}^N$  or Riemannian manifolds  $\Omega$ .

**5.1. The div-rot-Lemma Revisited.** Let  $\Omega \subset \mathbb{R}^3$ . The first example is given by the classical operators from vector analysis

$$\begin{aligned} A_0 &:= \mathring{\nabla}_{\Gamma_t} : \mathring{H}_{\Gamma_t}^1(\Omega) \subset L^2(\Omega) \longrightarrow L_\epsilon^2(\Omega); & u &\mapsto \nabla u, \\ A_1 &:= \mu^{-1} \mathring{\text{rot}}_{\Gamma_t} : \mathring{R}_{\Gamma_t}(\Omega) \subset L_\epsilon^2(\Omega) \longrightarrow L_\mu^2(\Omega); & E &\mapsto \mu^{-1} \text{rot } E, \\ A_2 &:= \mathring{\text{div}}_{\Gamma_t} \mu : \mu^{-1} \mathring{D}_{\Gamma_t}(\Omega) \subset L_\mu^2(\Omega) \longrightarrow L^2(\Omega); & H &\mapsto \text{div } \mu H. \end{aligned}$$

$A_0$ ,  $A_1$ , and  $A_2$  are unbounded, densely defined, and closed linear operators with adjoints

$$\begin{aligned} A_0^* &= \mathring{\nabla}_{\Gamma_t}^* = -\mathring{\text{div}}_{\Gamma_n} \epsilon : \epsilon^{-1} \mathring{D}_{\Gamma_n}(\Omega) \subset L_\epsilon^2(\Omega) \longrightarrow L^2(\Omega); & H &\mapsto -\text{div } \epsilon H, \\ A_1^* &= (\mu^{-1} \mathring{\text{rot}}_{\Gamma_t})^* = \epsilon^{-1} \mathring{\text{rot}}_{\Gamma_n} : \mathring{R}_{\Gamma_n}(\Omega) \subset L_\mu^2(\Omega) \longrightarrow L_\epsilon^2(\Omega); & E &\mapsto \epsilon^{-1} \text{rot } E, \\ A_2^* &= (\mathring{\text{div}}_{\Gamma_t} \mu)^* = -\mathring{\nabla}_{\Gamma_n} : \mathring{H}_{\Gamma_n}^1(\Omega) \subset L^2(\Omega) \longrightarrow L_\mu^2(\Omega); & u &\mapsto -\nabla u. \end{aligned}$$

Here,  $\epsilon, \mu : \Omega \rightarrow \mathbb{R}^{3 \times 3}$  are symmetric and uniformly positive definite  $L^\infty(\Omega)$ -tensor fields. Moreover, the Hilbert-Lebesgue space  $L_\epsilon^2(\Omega)$  is defined as the standard Lebesgue space  $L^2(\Omega)$  but with an equivalent inner product  $\langle \cdot, \cdot \rangle_{L_\epsilon^2(\Omega)} := \langle \epsilon \cdot, \cdot \rangle_{L^2(\Omega)}$ . Analogously we define  $L_\mu^2(\Omega)$ . The complex properties hold as

$$\begin{aligned} R(A_0) &= \mathring{\nabla}_{\Gamma_t} \mathring{H}_{\Gamma_t}^1(\Omega) \subset \mathring{R}_{\Gamma_t,0}(\Omega) = N(A_1), & R(A_1^*) &= \epsilon^{-1} \mathring{\text{rot}}_{\Gamma_n} \mathring{R}_{\Gamma_n}(\Omega) \subset \epsilon^{-1} \mathring{D}_{\Gamma_n,0}(\Omega) = N(A_0^*), \\ R(A_1) &= \mu^{-1} \mathring{\text{rot}}_{\Gamma_t} \mathring{R}_{\Gamma_t}(\Omega) \subset \mu^{-1} \mathring{D}_{\Gamma_t,0}(\Omega) = N(A_2), & R(A_2^*) &= \mathring{\nabla}_{\Gamma_n} \mathring{H}_{\Gamma_n}^1(\Omega) \subset \mathring{R}_{\Gamma_n,0}(\Omega) = N(A_1^*). \end{aligned}$$

Hence, the sequences (5.1) read

$$\begin{aligned} \mathring{H}_{\Gamma_t}^1(\Omega) \subset L^2(\Omega) &\xrightarrow{A_0 = \mathring{\nabla}_{\Gamma_t}} \mathring{R}_{\Gamma_t}(\Omega) \subset L_\epsilon^2(\Omega) \xrightarrow{A_1 = \mu^{-1} \mathring{\text{rot}}_{\Gamma_t}} \mu^{-1} \mathring{D}_{\Gamma_t}(\Omega) \subset L_\mu^2(\Omega) \xrightarrow{A_2 = \mathring{\text{div}}_{\Gamma_t} \mu} L^2(\Omega), \\ L^2(\Omega) &\xleftarrow{A_0^* = -\mathring{\text{div}}_{\Gamma_n} \epsilon} \epsilon^{-1} \mathring{D}_{\Gamma_n}(\Omega) \subset L_\epsilon^2(\Omega) \xleftarrow{A_1^* = \epsilon^{-1} \mathring{\text{rot}}_{\Gamma_n}} \mathring{R}_{\Gamma_n}(\Omega) \subset L_\mu^2(\Omega) \xleftarrow{A_2^* = -\mathring{\nabla}_{\Gamma_n}} \mathring{H}_{\Gamma_n}^1(\Omega) \subset L^2(\Omega). \end{aligned}$$

These are the well-known Hilbert complexes for electro-magnetics, which are also known as de Rham complexes. Typical equations arising from the de Rham complex are systems of electro-magneto statics, e.g.,

$$\begin{aligned} A_1 E &= \mu^{-1} \mathring{\text{rot}}_{\Gamma_t} E = F, \\ A_0^* E &= -\mathring{\text{div}}_{\Gamma_n} \epsilon E = f, \end{aligned}$$

or simply the Dirichlet-Neumann Laplacians and rot rot systems, e.g.,

$$\begin{aligned} A_0^* A_0 u &= -\mathring{\text{div}}_{\Gamma_n} \epsilon \mathring{\nabla}_{\Gamma_t} u = f, & A_1^* A_1 E &= \epsilon^{-1} \mathring{\text{rot}}_{\Gamma_n} \mu^{-1} \mathring{\text{rot}}_{\Gamma_t} E = F, \\ & & A_0^* E &= -\mathring{\text{div}}_{\Gamma_n} \epsilon E = f. \end{aligned}$$

The crucial embeddings (5.2) are compact by Weck's selection theorem, compare to Lemma 2.1.

**Lemma 5.1** (Weck's selection theorem). *Let  $\Omega \subset \mathbb{R}^3$  be a weak Lipschitz domain with weak Lipschitz interfaces. Then the embeddings*

$$\begin{aligned} D(A_1) \cap D(A_0^*) &= \mathring{R}_{\Gamma_t}(\Omega) \cap \epsilon^{-1} \mathring{D}_{\Gamma_n}(\Omega) \hookrightarrow L^2(\Omega), \\ D(A_2) \cap D(A_1^*) &= \mu^{-1} \mathring{D}_{\Gamma_t}(\Omega) \cap \mathring{R}_{\Gamma_n}(\Omega) \hookrightarrow L^2(\Omega) \end{aligned}$$

are compact.

Note that by interchanging the boundary conditions and  $\epsilon, \mu$  the latter two compact embeddings are equal. A proof can be found in [3, Theorem 4.7]. Indeed, Weck's selection theorems are independent of the material law tensors  $\epsilon$  or  $\mu$ . Choosing the pair  $(A_0, A_1)$  we get by Theorem 4.7 the following:

**Theorem 5.2** (global div  $\epsilon\mu^{-1}$  rot-lemma). *Let  $\mathring{R}_{\Gamma_t}(\Omega) \cap \epsilon^{-1} \mathring{D}_{\Gamma_n}(\Omega) \hookrightarrow L^2(\Omega)$  be compact. Moreover, let  $(E_n) \subset \mathring{R}_{\Gamma_t}(\Omega)$  and  $(H_n) \subset \epsilon^{-1} \mathring{D}_{\Gamma_n}(\Omega)$  be two sequences bounded in  $R(\Omega)$  and  $\epsilon^{-1} D(\Omega)$ , respectively. Then there exist  $E \in \mathring{R}_{\Gamma_t}(\Omega)$  and  $H \in \epsilon^{-1} \mathring{D}_{\Gamma_n}(\Omega)$  as well as subsequences, again denoted by  $(E_n)$  and  $(H_n)$ , such that*

- $E_n \rightharpoonup E$  in  $\mathring{R}_{\Gamma_t}(\Omega)$ ,
- $H_n \rightharpoonup H$  in  $\epsilon^{-1} \mathring{D}_{\Gamma_n}(\Omega)$ ,
- $\langle E_n, H_n \rangle_{L_\epsilon^2(\Omega)} \rightarrow \langle E, H \rangle_{L_\epsilon^2(\Omega)}$ .

**Remark 5.3.** *We note:*

- (i) *Considering  $(E_n)$  and  $(\epsilon H_n)$  shows that Theorem 5.2 is equivalent to the global div-rot-lemma Theorem 3.1.*
- (ii) *Theorem 5.2 has a corresponding local version similar to the local div-rot-lemma Corollary 3.2 and Remark 3.3, which holds with no regularity or boundedness assumptions on  $\Omega$ .*

The generalization given in Theorem 4.14 reads as follows.

**Theorem 5.4** (generalized/distributional global div  $\epsilon\mu^{-1}$  rot-lemma). *Let  $\mathring{\nabla} \mathring{H}_{\Gamma_t}^1(\Omega)$  and  $\mathring{\text{rot}} \mathring{R}_{\Gamma_t}(\Omega)$  be closed and let the Dirichlet-Neumann fields  $\mathring{R}_{\Gamma_t,0}(\Omega) \cap \epsilon^{-1} \mathring{D}_{\Gamma_n,0}(\Omega)$  be finite-dimensional. Moreover, let  $(E_n), (H_n) \subset L_\epsilon^2(\Omega)$  be two bounded sequences such that*

- $(\widetilde{\mu^{-1}\mathring{\text{rot}}_{\Gamma_t} E_n})$  is relatively compact in  $\mathring{\mathbb{R}}_{\Gamma_n}(\Omega)'$ ,
- $(\mathring{\text{div}}_{\Gamma_n} \epsilon H_n)$  is relatively compact in  $\mathring{\mathbb{H}}_{\Gamma_t}^1(\Omega)'$ .

Then there exist  $E, H \in \mathbb{L}_\epsilon^2(\Omega)$  as well as subsequences, again denoted by  $(E_n)$  and  $(H_n)$ , such that

- $E_n \rightharpoonup E$  in  $\mathbb{L}_\epsilon^2(\Omega)$ ,
- $H_n \rightharpoonup H$  in  $\mathbb{L}_\epsilon^2(\Omega)$ ,
- $\langle E_n, H_n \rangle_{\mathbb{L}_\epsilon^2(\Omega)} \rightarrow \langle E, H \rangle_{\mathbb{L}_\epsilon^2(\Omega)}$ .

**Remark 5.5.** We emphasize:

- By Lemma 5.1 and Lemma 4.18, Theorem 5.2 and Theorem 5.4 hold for weak Lipschitz domains  $\Omega \subset \mathbb{R}^3$  with weak Lipschitz interfaces.
- Choosing the pair  $(A_1, A_2)$  we get by Theorem 4.7 a variant of Theorem 5.2, shortly stating, that for bounded sequences  $(E_n) \subset \mu^{-1}\mathring{\mathbb{D}}_{\Gamma_t}(\Omega)$  and  $(H_n) \subset \mathring{\mathbb{R}}_{\Gamma_n}(\Omega)$  it holds (after picking subsequences)  $\langle E_n, H_n \rangle_{\mathbb{L}_\mu^2(\Omega)} \rightarrow \langle E, H \rangle_{\mathbb{L}_\mu^2(\Omega)}$ . Similarly, we get a variant of Theorem 5.4.

5.1.1. *The Classical div-rot-Lemma.* The classical div-rot-lemma (or div-curl-lemma) by Murat [17] and Tartar [32] reads as a slightly weaker version of Corollary III (local div-curl-lemma) from the introduction and uses only the standard dual space

$$\mathbb{H}^{-1}(\Omega) := \mathring{\mathbb{H}}^1(\Omega)'.$$

**Theorem 5.6** (classical div-rot-lemma). *Let  $\Omega \subset \mathbb{R}^3$  be an open set and let  $(E_n), (H_n) \subset \mathbb{L}^2(\Omega)$  be two sequences bounded in  $\mathbb{L}^2(\Omega)$  such that both  $(\widetilde{\text{rot}} E_n)$  and  $(\widetilde{\text{div}} H_n)$  are relatively compact in  $\mathbb{H}^{-1}(\Omega)$ . Then there exist  $E, H \in \mathbb{L}^2(\Omega)$  as well as subsequences, again denoted by  $(E_n)$  and  $(H_n)$ , such that the sequence of scalar products  $(E_n \cdot H_n)$  converges in the sense of distributions, i.e.,*

$$\forall \varphi \in \mathring{C}^\infty(\Omega) \quad \int_{\Omega} \varphi (E_n \cdot H_n) \rightarrow \int_{\Omega} \varphi (E \cdot H).$$

Here, we recall the linear extensions of  $A$  and  $A^*$  (tilde-operators) from Section 4.3.1

$$\widetilde{A} = (A^*)' \mathcal{R}_{H_1} : H_1 \rightarrow D(A^*)', \quad \widetilde{A}^* = A' \mathcal{R}_{H_2} : H_2 \rightarrow D(A)'$$

and consider the bounded linear operators and their adjoints

$$\begin{aligned} \mathring{\nabla} : \mathring{\mathbb{H}}^1(\Omega) &\longrightarrow \mathbb{L}^2(\Omega), & -\widetilde{\text{div}} = \mathring{\nabla}' \mathcal{R} : \mathbb{L}^2(\Omega) &\longrightarrow \mathring{\mathbb{H}}^1(\Omega)' = \mathbb{H}^{-1}(\Omega), \\ \mathring{\text{rot}} : \mathring{\mathbb{R}}(\Omega) &\longrightarrow \mathbb{L}^2(\Omega), & \widetilde{\text{rot}} = \mathring{\text{rot}}' \mathcal{R} : \mathbb{L}^2(\Omega) &\longrightarrow \mathring{\mathbb{R}}(\Omega)', \\ \mathring{\text{div}} : \mathring{\mathbb{D}}(\Omega) &\longrightarrow \mathbb{L}^2(\Omega), & -\widetilde{\nabla} = \mathring{\text{div}}' \mathcal{R} : \mathbb{L}^2(\Omega) &\longrightarrow \mathring{\mathbb{D}}(\Omega)', \\ \nabla : \mathbb{H}^1(\Omega) &\longrightarrow \mathbb{L}^2(\Omega), & -\widetilde{\text{div}} = \nabla' \mathcal{R} : \mathbb{L}^2(\Omega) &\longrightarrow \mathbb{H}^1(\Omega)' =: \mathring{\mathbb{H}}^{-1}(\Omega), \\ \text{rot} : \mathbb{R}(\Omega) &\longrightarrow \mathbb{L}^2(\Omega), & \widetilde{\text{rot}} = \text{rot}' \mathcal{R} : \mathbb{L}^2(\Omega) &\longrightarrow \mathbb{R}(\Omega)', \\ \text{div} : \mathbb{D}(\Omega) &\longrightarrow \mathbb{L}^2(\Omega), & -\widetilde{\nabla} = \text{div}' \mathcal{R} : \mathbb{L}^2(\Omega) &\longrightarrow \mathbb{D}(\Omega)', \end{aligned}$$

where  $\mathcal{R} := \mathcal{R}_{\mathbb{L}^2(\Omega)} : \mathbb{L}^2(\Omega) \rightarrow \mathbb{L}^2(\Omega)'$  denotes the (scalar or vector valued) Riesz isomorphism of  $\mathbb{L}^2(\Omega)$ . Note that the embeddings

$$\begin{aligned} \mathring{\mathbb{R}}(\Omega)', \mathring{\mathbb{D}}(\Omega)' &\subset \mathring{\mathbb{H}}^1(\Omega)' = \mathbb{H}^{-1}(\Omega), \\ \mathbb{R}(\Omega)', \mathbb{D}(\Omega)' &\subset \mathbb{H}^1(\Omega)' = \mathring{\mathbb{H}}^{-1}(\Omega), & \mathring{\mathbb{H}}^{-1}(\Omega) &= \mathbb{H}^1(\Omega)' \subset \mathring{\mathbb{H}}^1(\Omega)' = \mathbb{H}^{-1}(\Omega) \end{aligned}$$

justify the formulations in Theorem 5.6.

A typical application of Theorem 5.6 in homogenization of partial differential equations is given by the following problem: Let  $(u_n) \subset \mathring{\mathbb{H}}^1(\Omega)$  be the sequence of unique solutions of the Dirichlet-Laplace problems

$$-\widetilde{\text{div}} \Theta_n \mathring{\nabla} u_n = f \in \mathbb{H}^{-1}(\Omega),$$

with some tensor (matrix) fields  $\Theta_n$  having appropriate properties. Note that for all  $\varphi \in \mathring{H}^1(\Omega)$  we have the variational formulation

$$f(\varphi) = \mathring{\nabla}' \mathcal{R} \Theta_n \mathring{\nabla} u_n(\varphi) = \mathcal{R} \Theta_n \mathring{\nabla} u_n(\mathring{\nabla} \varphi) = \langle \mathring{\nabla} \varphi, \Theta_n \mathring{\nabla} u_n \rangle_{L^2(\Omega)}.$$

Setting

$$E_n := \mathring{\nabla} u_n \in \mathring{R}_0(\Omega) = N(A_1) \subset L^2(\Omega), \quad H_n := \Theta_n E_n \in L^2(\Omega)$$

we see

$$\widetilde{\text{rot}} E_n = \text{rot} E_n = 0 \in H^{-1}(\Omega), \quad \widetilde{\text{div}} H_n = -f \in H^{-1}(\Omega)$$

and thus both  $(\widetilde{\text{rot}} E_n)$  and  $(\widetilde{\text{div}} H_n)$  are trivially relatively compact in  $H^{-1}(\Omega)$  as they are even constant. Hence Theorem 5.6 yields for all  $\varphi \in \mathring{C}^\infty(\Omega)$  the convergence of

$$\int_{\Omega} \varphi (E_n \cdot H_n) = \int_{\Omega} \varphi (\mathring{\nabla} u_n \cdot \Theta_n \mathring{\nabla} u_n).$$

Let us conclude that in view of Theorem 5.4 ( $\epsilon = \mu = \text{id}$ ) the proper assumptions for  $(E_n), (H_n) \subset L^2(\Omega)$  in Theorem 5.6 are given either by (Dirichlet-Laplace)

- $(\widetilde{\text{rot}} E_n)$  is relatively compact in  $R(\Omega)'$ ,
- $(\widetilde{\text{div}} H_n)$  is relatively compact in  $\mathring{H}^1(\Omega)' = H^{-1}(\Omega)$ ,

or (Neumann-Laplace)

- $(\widetilde{\text{rot}} E_n)$  is relatively compact in  $\mathring{R}(\Omega)'$ ,
- $(\widetilde{\text{div}} H_n)$  is relatively compact in  $H^1(\Omega)' = \mathring{H}^{-1}(\Omega)$ ,

additionally to the closedness of the ranges  $\mathring{\nabla} \mathring{H}^1(\Omega)$ ,  $\nabla H^1(\Omega)$  and  $\mathring{\text{rot}} \mathring{R}(\Omega)$ ,  $\text{rot} R(\Omega)$  as well as the finite dimension of the Dirichlet fields  $\mathring{R}_0(\Omega) \cap \mathring{D}_0(\Omega)$  and the Neumann fields  $R_0(\Omega) \cap \mathring{D}_0(\Omega)$ , which is a topological property of the underlying domain  $\Omega$ , see [24, 25, 26]. Note that Theorem 5.4 implies the stronger convergence

$$\int_{\Omega} E_n \cdot H_n = \langle E_n, H_n \rangle_{L^2(\Omega)} \rightarrow \langle E, H \rangle_{L^2(\Omega)}.$$

**Remark 5.7.** Let  $\Omega \subset \mathbb{R}^3$  be a bounded strong Lipschitz domain with trivial topology. Then

$$\begin{aligned} \mathring{R}(\Omega)' &= \mathring{D}^{-1}(\Omega) := \{F \in H^{-1}(\Omega) : \widehat{\text{div}} F \in H^{-1}(\Omega)\}, \\ \mathring{D}(\Omega)' &= \mathring{R}^{-1}(\Omega) := \{F \in H^{-1}(\Omega) : \widehat{\text{rot}} F \in H^{-1}(\Omega)\} \end{aligned}$$

hold with equivalent norms, see [23] or for the two-dimensional analog [5]. We conjecture that the duals of  $R(\Omega)$  and  $D(\Omega)$  are given by

$$\begin{aligned} R(\Omega)' &= \mathring{D}^{-1}(\Omega) := \{F \in \mathring{H}^{-1}(\Omega) : \widehat{\text{div}} F \in \mathring{H}^{-1}(\Omega)\}, \\ D(\Omega)' &= \mathring{R}^{-1}(\Omega) := \{F \in \mathring{H}^{-1}(\Omega) : \widehat{\text{rot}} F \in \mathring{H}^{-1}(\Omega)\} \end{aligned}$$

with equivalent norms. Here,  $\widehat{\text{div}}$  and  $\widehat{\text{rot}}$  act as operators from  $H^{-1}(\Omega)$  to  $H^{-2}(\Omega)$  and  $\mathring{\widehat{\text{div}}}$  and  $\mathring{\widehat{\text{rot}}}$  act as operators from  $\mathring{H}^{-1}(\Omega)$  to  $\mathring{H}^{-2}(\Omega)$ .

We observe the following.

**Lemma 5.8.** Let the assertions in Remark 5.7 hold. Then for  $E \in L^2(\Omega)$  and  $(E_n) \subset L^2(\Omega)$  it holds:

- (i)  $\mathring{\widehat{\text{div}}} \widetilde{\text{rot}} E = 0$
- (i')  $\widetilde{\text{rot}} E \in \mathring{R}(\Omega)' \Leftrightarrow \widetilde{\text{rot}} E \in H^{-1}(\Omega)$
- (i'')  $(\widetilde{\text{rot}} E_n)$  relatively compact in  $\mathring{R}(\Omega)'$   $\Leftrightarrow$   $(\widetilde{\text{rot}} E_n)$  relatively compact in  $H^{-1}(\Omega)$
- (ii)  $\mathring{\widehat{\text{div}}} \mathring{\widetilde{\text{rot}}} E = 0$
- (ii')  $\mathring{\widetilde{\text{rot}}} E \in R(\Omega)' \Leftrightarrow \mathring{\widetilde{\text{rot}}} E \in \mathring{H}^{-1}(\Omega)$
- (ii'')  $(\mathring{\widetilde{\text{rot}}} E_n)$  relatively compact in  $R(\Omega)'$   $\Leftrightarrow$   $(\mathring{\widetilde{\text{rot}}} E_n)$  relatively compact in  $\mathring{H}^{-1}(\Omega)$

*Proof.* For  $F := \widetilde{\text{rot}} E \in \mathring{R}(\Omega)' \subset \mathbf{H}^{-1}(\Omega)$  we have  $\widehat{\text{div}} F = 0 \in \mathbf{H}^{-1}(\Omega)$  as for all  $\varphi \in \mathring{H}^2(\Omega)$

$$-\widehat{\text{div}} \widetilde{\text{rot}} E(\varphi) = \mathring{\text{rot}}' \mathcal{R} E(\nabla \varphi) = \mathcal{R} E(\text{rot } \nabla \varphi) = 0,$$

which shows (i), (i'), (i'') by Remark 5.7. Analogously we see (ii), (ii'), (ii'').  $\square$

Finally, we obtain a refined version of Theorem 5.4 in the case of full boundary conditions, compare to Theorem 5.6.

**Theorem 5.9** (improved classical div-rot-lemma). *Let  $\Omega \subset \mathbb{R}^3$  be a bounded strong Lipschitz domain with trivial topology. Moreover, let  $(E_n), (H_n) \subset \mathbf{L}^2(\Omega)$  be two bounded sequences such that either*

- $(\widetilde{\text{rot}} E_n)$  is relatively compact in  $\mathring{H}^{-1}(\Omega)$ ,
- $(\widetilde{\text{div}} H_n)$  is relatively compact in  $\mathbf{H}^{-1}(\Omega)$

or

- $(\widetilde{\text{rot}} E_n)$  is relatively compact in  $\mathbf{H}^{-1}(\Omega)$ ,
- $(\mathring{\text{div}} H_n)$  is relatively compact in  $\mathring{H}^{-1}(\Omega)$ .

Then there exist  $E, H \in \mathbf{L}^2(\Omega)$  as well as subsequences, again denoted by  $(E_n)$  and  $(H_n)$ , such that

- $E_n \rightharpoonup E$  in  $\mathbf{L}^2(\Omega)$ ,
- $H_n \rightharpoonup H$  in  $\mathbf{L}^2(\Omega)$ ,
- $\langle E_n, H_n \rangle_{\mathbf{L}^2(\Omega)} \rightarrow \langle E, H \rangle_{\mathbf{L}^2(\Omega)}$ .

We emphasize that the assumptions on  $\Omega$  in the latter theorem imply that  $\mathring{\nabla} \mathring{H}^1(\Omega)$ ,  $\nabla \mathbf{H}^1(\Omega)$ ,  $\mathring{\text{rot}} \mathring{R}(\Omega)$ ,  $\mathring{\text{rot}} \mathbf{R}(\Omega)$  are closed and that the Dirichlet fields  $\mathring{R}_0(\Omega) \cap \mathring{D}_0(\Omega)$  and the Neumann fields  $\mathring{R}_0(\Omega) \cap \mathring{D}_0^*(\Omega)$  are finite-dimensional, even trivial.

A more detailed discussion with nice results on the connections to the classical div-rot-lemma can be found in [35].

**5.2. Generalized Electro-Magnetics.** Let  $\Omega \subset \mathbb{R}^N$  or let  $\Omega$  even be a smooth Riemannian manifold with Lipschitz boundary  $\Gamma$  (Lipschitz submanifold) having (interface) Lipschitz submanifolds  $\Gamma_t, \Gamma_n$ . Using the calculus of alternating differential  $q$ -forms,  $q = 0, \dots, N$ , we define the exterior derivative  $d$  and co-derivative  $\delta = \pm * d *$  in the weak sense by

$$\mathbf{D}^q(\Omega) := \{E \in \mathbf{L}^{2,q}(\Omega) : dE \in \mathbf{L}^{2,q+1}(\Omega)\}, \quad \Delta^{q+1}(\Omega) := \{H \in \mathbf{L}^{2,q+1}(\Omega) : \delta H \in \mathbf{L}^{2,q}(\Omega)\},$$

where  $\mathbf{L}^{2,q}(\Omega)$  denotes the standard Lebesgue space of square integrable  $q$ -forms. To introduce boundary conditions we define

$$\mathring{d}_{\Gamma_t}^q : \mathring{D}_{\Gamma_t}^q(\Omega) := \overline{\mathring{C}_{\Gamma_t}^{\infty,q}(\Omega)}^{\mathbf{D}^q(\Omega)} \subset \mathbf{L}^{2,q}(\Omega) \longrightarrow \mathbf{L}^{2,q+1}(\Omega); \quad E \mapsto dE$$

as closure of the classical exterior derivative  $d$  acting on test  $q$ -forms.  $\mathring{d}_{\Gamma_t}^q$  is an unbounded, densely defined, and closed linear operator with adjoint

$$(\mathring{d}_{\Gamma_t}^q)^* = -\mathring{\delta}_{\Gamma_n}^{q+1} : \mathring{\Delta}_{\Gamma_n}^{q+1}(\Omega) := \overline{\mathring{C}_{\Gamma_n}^{\infty,q+1}(\Omega)}^{\Delta^{q+1}(\Omega)} \subset \mathbf{L}^{2,q+1}(\Omega) \longrightarrow \mathbf{L}^{2,q}(\Omega); \quad H \mapsto -\delta H.$$

Let us introduce

$$A_0 := \mathring{d}_{\Gamma_t}^{q-1}, \quad A_1 := \mathring{d}_{\Gamma_t}^q, \quad A_0^* = -\mathring{\delta}_{\Gamma_n}^q, \quad A_1^* = -\mathring{\delta}_{\Gamma_n}^{q+1}.$$

The complex properties hold as, e.g.,

$$R(A_0) = \mathring{d}_{\Gamma_t}^{q-1} \mathring{D}_{\Gamma_t}^{q-1}(\Omega) \subset \mathring{D}_{\Gamma_t,0}^q(\Omega) = N(A_1), \quad R(A_1^*) = \mathring{\delta}_{\Gamma_n}^{q+1} \mathring{\Delta}_{\Gamma_n}^{q+1}(\Omega) \subset \mathring{\Delta}_{\Gamma_n,0}^q(\Omega) = N(A_0^*)$$

by the classical properties  $\delta \delta = \pm * d d * = 0$ . Hence, the sequences (5.1) read

$$\begin{aligned} \mathring{D}_{\Gamma_t}^{q-1}(\Omega) \subset \mathbf{L}^{2,q-1}(\Omega) &\xrightarrow{A_0 = \mathring{d}_{\Gamma_t}^{q-1}} \mathring{D}_{\Gamma_t}^q(\Omega) \subset \mathbf{L}^{2,q}(\Omega) \xrightarrow{A_1 = \mathring{d}_{\Gamma_t}^q} \mathbf{L}^{2,q+1}(\Omega), \\ \mathbf{L}^{2,q-1}(\Omega) &\xleftarrow{A_0^* = -\mathring{\delta}_{\Gamma_n}^q} \mathring{\Delta}_{\Gamma_n}^q(\Omega) \subset \mathbf{L}^{2,q}(\Omega) \xleftarrow{A_1^* = -\mathring{\delta}_{\Gamma_n}^{q+1}} \mathring{\Delta}_{\Gamma_n}^{q+1}(\Omega) \subset \mathbf{L}^{2,q+1}(\Omega), \end{aligned}$$

which are the well-known Hilbert complexes for generalized electro-magnetics, i.e., the de Rham complexes. Typical equations arising from the de Rham complex are systems of generalised electro-magneto statics, e.g.,

$$\begin{aligned} A_1 E &= \mathring{d}_{\Gamma_t}^q E = F, \\ A_0^* E &= -\mathring{\delta}_{\Gamma_n}^q E = G, \end{aligned}$$

or systems of generalized Dirichlet-Neumann Laplacians, e.g.,

$$\begin{aligned} A_1^* A_1 E &= -\mathring{\delta}_{\Gamma_n}^{q+1} \mathring{d}_{\Gamma_t}^q E = F, & (A_1^* A_1 + A_0 A_0^*) E &= -(\mathring{\delta}_{\Gamma_n}^{q+1} \mathring{d}_{\Gamma_t}^q + \mathring{d}_{\Gamma_n}^{q-1} \mathring{\delta}_{\Gamma_t}^q) E = F, \\ A_0^* E &= -\mathring{\delta}_{\Gamma_n}^q E = G. \end{aligned}$$

The crucial embeddings (5.2) are compact by (a generalization) Weck's selection theorem, compare to Lemma 2.1.

**Lemma 5.10** (Weck's selection theorem). *Let  $\Omega \subset \mathbb{R}^N$  be a weak Lipschitz domain with weak Lipschitz interfaces or even a Riemannian manifold with Lipschitz boundary and Lipschitz interfaces. Then for all  $q$  the embeddings*

$$D(A_1) \cap D(A_0^*) = \mathring{D}_{\Gamma_t}^q(\Omega) \cap \mathring{\Delta}_{\Gamma_n}^q(\Omega) \hookrightarrow \mathbb{L}^2(\Omega)$$

are compact.

A proof can be found in [4, Theorem 4.9], see also the fundamental papers of Weck [38] (strong Lipschitz) and Picard [27] (weak Lipschitz) for full boundary conditions. Again, Weck's selection theorems are independent of possible material law tensors  $\epsilon$  or  $\mu$ . Theorem 4.7 shows the following result:

**Theorem 5.11** (global  $\delta$ -d-lemma). *Let the embedding  $\mathring{D}_{\Gamma_t}^q(\Omega) \cap \mathring{\Delta}_{\Gamma_n}^q(\Omega) \hookrightarrow \mathbb{L}^2(\Omega)$  be compact. Moreover, let  $(E_n) \subset \mathring{D}_{\Gamma_t}^q(\Omega)$  and  $(H_n) \subset \mathring{\Delta}_{\Gamma_n}^q(\Omega)$  be two sequences bounded in  $\mathbb{D}^q(\Omega)$  and  $\Delta^q(\Omega)$ , respectively. Then there exist  $E \in \mathring{D}_{\Gamma_t}^q(\Omega)$  and  $H \in \mathring{\Delta}_{\Gamma_n}^q(\Omega)$  as well as subsequences, again denoted by  $(E_n)$  and  $(H_n)$ , such that*

- $E_n \rightharpoonup E$  in  $\mathring{D}_{\Gamma_t}^q(\Omega)$ ,
- $H_n \rightharpoonup H$  in  $\mathring{\Delta}_{\Gamma_n}^q(\Omega)$ ,
- $\langle E_n, H_n \rangle_{\mathbb{L}^{2,q}(\Omega)} \rightarrow \langle E, H \rangle_{\mathbb{L}^{2,q}(\Omega)}$ .

**Remark 5.12.** *We note:*

- (i) For  $N = 3$  and  $q = 1$  (or  $q = 2$ ) we obtain by Theorem 5.11 again the global div-rot-lemma Theorem 3.1.
- (ii) For  $q = 0$  (or  $q = N$ ) as well as identifying  $\mathring{d}_{\Gamma_t}^0 = \mathring{\nabla}_{\Gamma_t}$  and  $\mathring{\Delta}_{\Gamma_n}^0(\Omega) = 0$  (or  $\mathring{d}_{\Gamma_t}^N = 0$  and  $\mathring{\Delta}_{\Gamma_n}^N(\Omega) = \mathring{\nabla}_{\Gamma_n}$ ) we get by Theorem 5.11 the following trivial (by Rellich's selection theorem) result: For all bounded sequences  $(u_n) \subset \mathring{H}_{\Gamma_t}^1(\Omega)$  and  $(v_n) \subset \mathbb{L}^2(\Omega)$  there exist  $u \in \mathring{H}_{\Gamma_t}^1(\Omega)$  and  $v \in \mathbb{L}^2(\Omega)$  as well as subsequences, again denoted by  $(u_n)$  and  $(v_n)$ , such that  $(u_n)$  and  $(v_n)$  converge weakly in  $\mathring{H}_{\Gamma_t}^1(\Omega)$  (or  $\mathbb{L}^2(\Omega)$ ) to  $u$  and  $v$ , respectively, together with the convergence of the inner products  $\langle u_n, v_n \rangle_{\mathbb{L}^2(\Omega)} \rightarrow \langle u, v \rangle_{\mathbb{L}^2(\Omega)}$ .
- (iii) Theorem 5.11 has a corresponding local version similar to the local div-rot-lemma Corollary 3.2 and Remark 3.3, which holds with no regularity or boundedness assumptions on  $\Omega$ .
- (iv) Material law tensors  $\epsilon$  and  $\mu$  different from the identities can be handled as well.

The generalization given in Theorem 4.14 reads as follows.

**Theorem 5.13** (generalized/distributional global  $\delta$ -d-lemma). *Let  $d\mathring{D}_{\Gamma_t}^{q-1}(\Omega)$  and  $d\mathring{D}_{\Gamma_t}^q(\Omega)$  be closed and let the generalized Dirichlet-Neumann fields  $\mathring{D}_{\Gamma_t,0}^q(\Omega) \cap \mathring{\Delta}_{\Gamma_n,0}^q(\Omega)$  be finite-dimensional. Moreover, let  $(E_n), (H_n) \subset \mathbb{L}^{2,q}(\Omega)$  be two bounded sequences such that*

- $(\mathring{d}_{\Gamma_t}^q E_n)$  is relatively compact in  $\mathring{\Delta}_{\Gamma_n}^{q+1}(\Omega)'$ ,

- $(\widetilde{\delta_{\Gamma_n}^q} H_n)$  is relatively compact in  $\mathring{D}_{\Gamma_n}^{q-1}(\Omega)'$ .

Then there exist  $E, H \in \mathbb{L}^{2,q}(\Omega)$  as well as subsequences, again denoted by  $(E_n)$  and  $(H_n)$ , such that

- $E_n \rightharpoonup E$  in  $\mathbb{L}^{2,q}(\Omega)$ ,
- $H_n \rightharpoonup H$  in  $\mathbb{L}^{2,q}(\Omega)$ ,
- $\langle E_n, H_n \rangle_{\mathbb{L}^{2,q}(\Omega)} \rightarrow \langle E, H \rangle_{\mathbb{L}^{2,q}(\Omega)}$ .

**Remark 5.14.** By Lemma 5.10 and Lemma 4.18, Theorem 5.11 and Theorem 5.13 hold for weak Lipschitz domains  $\Omega \subset \mathbb{R}^N$  with weak Lipschitz interfaces or even for Riemannian manifolds  $\Omega$ .

**5.3. Biharmonic Equation, General Relativity, and Gravitational Waves.** Let  $\Omega \subset \mathbb{R}^3$ . We introduce symmetric and deviatoric (trace-free) square integrable tensor fields in  $\mathbb{L}^2(\Omega; \mathbb{S})$  and  $\mathbb{L}^2(\Omega; \mathbb{T})$  and as closures of the Hessian  $\nabla \nabla$ , and Rot, Div (row-wise rot, div), applied to test functions or test tensor fields, the linear operators

$$\begin{aligned} A_0 &:= \nabla \nabla : \mathring{H}^2(\Omega) := \overline{\mathring{C}^\infty(\Omega)}^{\mathring{H}^2(\Omega)} \subset \mathbb{L}^2(\Omega) \longrightarrow \mathbb{L}^2(\Omega; \mathbb{S}); & u &\mapsto \nabla \nabla u, \\ A_1 &:= \mathring{\text{Rot}}_{\mathbb{S}} : \mathring{R}(\Omega; \mathbb{S}) := \overline{\mathring{C}^\infty(\Omega; \mathbb{S})}^{\mathring{R}(\Omega)} \subset \mathbb{L}^2(\Omega; \mathbb{S}) \longrightarrow \mathbb{L}^2(\Omega; \mathbb{T}); & S &\mapsto \text{Rot } S, \\ A_2 &:= \mathring{\text{Div}}_{\mathbb{T}} : \mathring{D}(\Omega; \mathbb{T}) := \overline{\mathring{C}^\infty(\Omega; \mathbb{T})}^{\mathring{D}(\Omega)} \subset \mathbb{L}^2(\Omega; \mathbb{T}) \longrightarrow \mathbb{L}^2(\Omega); & T &\mapsto \text{Div } T. \end{aligned}$$

$A_0, A_1$ , and  $A_2$  are unbounded, densely defined, and closed linear operators with adjoints

$$\begin{aligned} A_0^* &= (\nabla \nabla)^* = \text{div Div}_{\mathbb{S}} : \mathring{D}(\Omega; \mathbb{S}) \subset \mathbb{L}^2(\Omega; \mathbb{S}) \longrightarrow \mathbb{L}^2(\Omega); & S &\mapsto \text{div Div } S, \\ A_1^* &= \mathring{\text{Rot}}_{\mathbb{S}}^* = \text{sym Rot}_{\mathbb{T}} : \mathring{R}_{\text{sym}}(\Omega; \mathbb{T}) \subset \mathbb{L}^2(\Omega; \mathbb{T}) \longrightarrow \mathbb{L}^2(\Omega; \mathbb{S}); & T &\mapsto \text{sym Rot } T, \\ A_2^* &= \mathring{\text{Div}}_{\mathbb{T}}^* = -\text{dev } \nabla : \mathring{H}^1(\Omega) \subset \mathbb{L}^2(\Omega) \longrightarrow \mathbb{L}^2(\Omega; \mathbb{T}); & v &\mapsto -\text{dev } \nabla v, \end{aligned}$$

where  $\mathring{H}^1(\Omega), \mathring{H}^2(\Omega)$  denote the usual Sobolev spaces and

$$\begin{aligned} \mathring{R}(\Omega) &:= \{S \in \mathbb{L}^2(\Omega) : \text{Rot } S \in \mathbb{L}^2(\Omega)\}, & \mathring{R}(\Omega; \mathbb{S}) &:= \mathring{R}(\Omega) \cap \mathbb{L}^2(\Omega; \mathbb{S}), \\ \mathring{D}(\Omega) &:= \{T \in \mathbb{L}^2(\Omega) : \text{Div } T \in \mathbb{L}^2(\Omega)\}, & \mathring{D}(\Omega; \mathbb{T}) &:= \mathring{D}(\Omega) \cap \mathbb{L}^2(\Omega; \mathbb{T}), \\ \mathring{D}(\Omega) &:= \{S \in \mathbb{L}^2(\Omega) : \text{div Div } S \in \mathbb{L}^2(\Omega)\}, & \mathring{D}(\Omega; \mathbb{S}) &:= \mathring{D}(\Omega) \cap \mathbb{L}^2(\Omega; \mathbb{S}), \\ \mathring{R}_{\text{sym}}(\Omega) &:= \{T \in \mathbb{L}^2(\Omega) : \text{sym Rot } T \in \mathbb{L}^2(\Omega)\}, & \mathring{R}_{\text{sym}}(\Omega; \mathbb{T}) &:= \mathring{R}_{\text{sym}}(\Omega) \cap \mathbb{L}^2(\Omega; \mathbb{T}), \end{aligned}$$

see [22] for details. Note that  $u, v$ , and  $S, T$  are scalar, vector, and tensor (matrix) fields, respectively. Moreover, for  $S \in \mathring{R}(\Omega; \mathbb{S})$  it holds  $\text{Rot } S \in \mathbb{L}^2(\Omega; \mathbb{T})$ . The complex properties hold as

$$\begin{aligned} R(A_0) &= \nabla \nabla \mathring{H}^2(\Omega) \subset \mathring{R}_0(\Omega; \mathbb{S}) = N(A_1), \\ R(A_1^*) &= \text{sym Rot}_{\mathbb{T}} \mathring{R}_{\text{sym}}(\Omega; \mathbb{T}) \subset \mathring{D}(\Omega; \mathbb{S}) = N(A_0^*), \\ R(A_1) &= \mathring{\text{Rot}}_{\mathbb{S}} \mathring{R}(\Omega; \mathbb{S}) \subset \mathring{D}_0(\Omega; \mathbb{T}) = N(A_2), \\ R(A_2^*) &= \text{dev } \nabla \mathring{H}^1(\Omega) \subset \mathring{R}_{\text{sym},0}(\Omega; \mathbb{T}) = N(A_1^*), \end{aligned}$$

see again [22]. The sequences (5.1) read

$$\begin{aligned} \mathring{H}^2(\Omega) \subset \mathbb{L}^2(\Omega) &\xrightarrow{A_0 = \nabla \nabla} \mathring{R}(\Omega; \mathbb{S}) \subset \mathbb{L}^2(\Omega; \mathbb{S}) \xrightarrow{A_1 = \mathring{\text{Rot}}_{\mathbb{S}}} \mathring{D}(\Omega; \mathbb{T}) \subset \mathbb{L}^2(\Omega; \mathbb{T}) \xrightarrow{A_2 = \mathring{\text{Div}}_{\mathbb{T}}} \mathbb{L}^2(\Omega), \\ \mathbb{L}^2(\Omega) &\xleftarrow{A_0^* = \text{div Div}_{\mathbb{S}}} \mathring{D}(\Omega; \mathbb{S}) \subset \mathbb{L}^2(\Omega; \mathbb{S}) \xleftarrow{A_1^* = \text{sym Rot}_{\mathbb{T}}} \mathring{R}_{\text{sym}}(\Omega; \mathbb{T}) \subset \mathbb{L}^2(\Omega; \mathbb{T}) \xleftarrow{A_2^* = -\text{dev } \nabla} \mathring{H}^1(\Omega) \subset \mathbb{L}^2(\Omega). \end{aligned}$$

These are the so-called Gradgrad and divDiv complexes, appearing, e.g., in biharmonic problems or general relativity, see [22] for details. Typical equations arising from the Gradgrad complex are systems of general relativity, e.g.,

$$\begin{aligned} A_1 S &= \mathring{\text{Rot}}_{\mathbb{S}} S = F, & A_2 T &= \mathring{\text{Div}}_{\mathbb{T}} T = g, \\ A_0^* S &= \text{div Div}_{\mathbb{S}} S = f, & A_1^* T &= \text{sym Rot}_{\mathbb{T}} T = G, \end{aligned}$$

or simply biharmonic equations and related second order systems, e.g.,

$$\begin{aligned} A_0^* A_0 u &= \operatorname{div} \operatorname{Div}_{\mathbb{S}} \nabla \nabla u = f, & A_1^* A_1 S &= \operatorname{sym} \operatorname{Rot}_{\mathbb{T}} \mathring{\operatorname{Rot}}_{\mathbb{S}} S = G, \\ & & A_0^* S &= \operatorname{div} \operatorname{Div}_{\mathbb{S}} S = f. \end{aligned}$$

The crucial embeddings (5.2) are compact, compare to Lemma 2.1.

**Lemma 5.15** (biharmonic selection theorems). *Let  $\Omega \subset \mathbb{R}^3$  be a strong Lipschitz domain. Then the embeddings*

$$\begin{aligned} D(A_1) \cap D(A_0^*) &= \mathring{R}(\Omega; \mathbb{S}) \cap \operatorname{DD}(\Omega; \mathbb{S}) \hookrightarrow L^2(\Omega; \mathbb{S}), \\ D(A_2) \cap D(A_1^*) &= \mathring{D}(\Omega; \mathbb{T}) \cap \operatorname{R}_{\operatorname{sym}}(\Omega; \mathbb{T}) \hookrightarrow L^2(\Omega; \mathbb{T}), \end{aligned}$$

are compact.

A proof can be found in [22, Lemma 3.22]. Again, the biharmonic selection theorems are independent of possible material law tensors  $\epsilon$  or  $\mu$ . Choosing the pair  $(A_0, A_1)$  we get by Theorem 4.7 the following:

**Theorem 5.16** (global div Div-Rot- $\mathbb{S}$ -lemma). *Let  $\mathring{R}(\Omega; \mathbb{S}) \cap \operatorname{DD}(\Omega; \mathbb{S}) \hookrightarrow L^2(\Omega; \mathbb{S})$  be compact. Moreover, let  $(S_n) \subset \mathring{R}(\Omega; \mathbb{S})$  and  $(T_n) \subset \operatorname{DD}(\Omega; \mathbb{S})$  be two sequences bounded in  $R(\Omega)$  and  $\operatorname{DD}(\Omega)$ , respectively. Then there exist  $S \in \mathring{R}(\Omega; \mathbb{S})$  and  $T \in \operatorname{DD}(\Omega; \mathbb{S})$  as well as subsequences, again denoted by  $(S_n)$  and  $(T_n)$ , such that*

- $S_n \rightharpoonup S$  in  $\mathring{R}(\Omega; \mathbb{S})$ ,
- $T_n \rightharpoonup T$  in  $\operatorname{DD}(\Omega; \mathbb{S})$ ,
- $\langle S_n, T_n \rangle_{L^2(\Omega; \mathbb{S})} \rightarrow \langle S, T \rangle_{L^2(\Omega; \mathbb{S})}$ .

For the pair  $(A_1, A_2)$  Theorem 4.7 implies:

**Theorem 5.17** (global sym Rot-Div- $\mathbb{T}$ -lemma). *Let  $\mathring{D}(\Omega; \mathbb{T}) \cap \operatorname{R}_{\operatorname{sym}}(\Omega; \mathbb{T}) \hookrightarrow L^2(\Omega; \mathbb{T})$  be compact. Moreover, let  $(S_n) \subset \mathring{D}(\Omega; \mathbb{T})$  and  $(T_n) \subset \operatorname{R}_{\operatorname{sym}}(\Omega; \mathbb{T})$  be two sequences bounded in  $D(\Omega)$  and  $\operatorname{R}_{\operatorname{sym}}(\Omega)$ , respectively. Then there exist  $S \in \mathring{D}(\Omega; \mathbb{T})$  and  $T \in \operatorname{R}_{\operatorname{sym}}(\Omega; \mathbb{T})$  as well as subsequences, again denoted by  $(S_n)$  and  $(T_n)$ , such that*

- $S_n \rightharpoonup S$  in  $\mathring{D}(\Omega; \mathbb{T})$ ,
- $T_n \rightharpoonup T$  in  $\operatorname{R}_{\operatorname{sym}}(\Omega; \mathbb{T})$ ,
- $\langle S_n, T_n \rangle_{L^2(\Omega; \mathbb{T})} \rightarrow \langle S, T \rangle_{L^2(\Omega; \mathbb{T})}$ .

**Remark 5.18.** *Material law tensors  $\epsilon$  and  $\mu$  different from the identities can be handled as well. Theorem 5.16 and Theorem 5.17 have corresponding local versions similar to the local div-rot-lemma Corollary 3.2 and Remark 3.3, which hold with no regularity or boundedness assumptions on  $\Omega$ . We note that the local version of Theorem 5.16 is a bit more involved as standard localization techniques (multiplication by test functions) fail due to the second order nature of the Sobolev space  $\operatorname{DD}(\Omega; \mathbb{S})$ . This additional difficulty can be overcome with the help of a non-standard Helmholtz type decomposition, see [22, Lemma 3.21] and the proof of [22, Lemma 3.22].*

The generalizations from Theorem 4.14 read as follows.

**Theorem 5.19** (generalized/distributional global div Div-Rot- $\mathbb{S}$ -lemma). *Let  $\nabla \nabla \mathring{H}^2(\Omega)$  and  $\operatorname{Rot} \mathring{R}(\Omega; \mathbb{S})$  be closed and let the generalized Dirichlet-Neumann fields  $\mathring{R}_0(\Omega; \mathbb{S}) \cap \operatorname{DD}_0(\Omega; \mathbb{S})$  be finite-dimensional. Moreover, let  $(S_n), (T_n) \subset L^2(\Omega; \mathbb{S})$  be two bounded sequences such that*

- $(\widetilde{\operatorname{Rot}}_{\mathbb{S}} S_n)$  is relatively compact in  $\operatorname{R}_{\operatorname{sym}}(\Omega; \mathbb{T})'$ ,
- $(\operatorname{div} \operatorname{Div}_{\mathbb{S}} T_n)$  is relatively compact in  $\mathring{H}^2(\Omega)' = H^{-2}(\Omega)$ .

Then there exist  $S, T \in L^2(\Omega; \mathbb{S})$  as well as subsequences, again denoted by  $(S_n)$  and  $(T_n)$ , such that

- $S_n \rightharpoonup S$  in  $L^2(\Omega; \mathbb{S})$ ,
- $T_n \rightharpoonup T$  in  $L^2(\Omega; \mathbb{S})$ ,

- $\langle S_n, T_n \rangle_{\mathbf{L}^2(\Omega, \mathbb{S})} \rightarrow \langle S, T \rangle_{\mathbf{L}^2(\Omega, \mathbb{S})}$ .

**Theorem 5.20** (generalized/distributional global sym Rot-Div-T-lemma). *Let the ranges  $\text{Rot } \mathring{R}(\Omega; \mathbb{S})$  and  $\text{Div } \mathring{D}(\Omega; \mathbb{T})$  be closed and let the generalized Dirichlet-Neumann fields  $\mathring{D}_0(\Omega; \mathbb{T}) \cap \mathbf{R}_{\text{sym}, 0}(\Omega; \mathbb{T})$  be finite-dimensional. Moreover, let  $(S_n), (T_n) \subset \mathbf{L}^2(\Omega, \mathbb{T})$  be two bounded sequences such that*

- $(\widetilde{\text{Div}}_{\mathbb{T}} S_n)$  is relatively compact in  $\mathbf{H}^1(\Omega)' = \mathring{H}^{-1}(\Omega)$ ,
- $(\text{sym Rot}_{\mathbb{T}} T_n)$  is relatively compact in  $\mathring{R}(\Omega; \mathbb{S})'$ .

Then there exist  $S, T \in \mathbf{L}^2(\Omega, \mathbb{T})$  as well as subsequences, again denoted by  $(S_n)$  and  $(T_n)$ , such that

- $S_n \rightharpoonup S$  in  $\mathbf{L}^2(\Omega, \mathbb{T})$ ,
- $T_n \rightharpoonup T$  in  $\mathbf{L}^2(\Omega, \mathbb{T})$ ,
- $\langle S_n, T_n \rangle_{\mathbf{L}^2(\Omega, \mathbb{T})} \rightarrow \langle S, T \rangle_{\mathbf{L}^2(\Omega, \mathbb{T})}$ .

**Remark 5.21.** By Lemma 5.15 and Lemma 4.18, Theorem 5.16, Theorem 5.17, and Theorem 5.19, Theorem 5.20 hold for strong Lipschitz domains  $\Omega \subset \mathbb{R}^3$ .

**5.4. Linear Elasticity.** Let  $\Omega \subset \mathbb{R}^3$  and let

$$\begin{aligned} A_0 &:= \text{sym } \mathring{\nabla} : \mathring{H}^1(\Omega) \subset \mathbf{L}^2(\Omega) \longrightarrow \mathbf{L}^2(\Omega; \mathbb{S}); & v &\mapsto \text{sym } \nabla v, \\ A_1 &:= \text{Rot } \mathring{\text{Rot}}_{\mathbb{S}}^{\top} : \mathring{\text{RR}}^{\top}(\Omega; \mathbb{S}) := \overline{\mathring{C}^{\infty}(\Omega; \mathbb{S})}^{\text{RR}^{\top}(\Omega)} \subset \mathbf{L}^2(\Omega; \mathbb{S}) \longrightarrow \mathbf{L}^2(\Omega; \mathbb{S}); & S &\mapsto \text{Rot } \text{Rot}^{\top} S, \\ A_2 &:= \mathring{\text{Div}}_{\mathbb{S}} : \mathring{D}(\Omega; \mathbb{S}) := \overline{\mathring{C}^{\infty}(\Omega; \mathbb{S})}^{\mathring{D}(\Omega)} \subset \mathbf{L}^2(\Omega; \mathbb{S}) \longrightarrow \mathbf{L}^2(\Omega); & T &\mapsto \text{Div } T. \end{aligned}$$

$A_0, A_1,$  and  $A_2$  are unbounded, densely defined, and closed linear operators with adjoints

$$\begin{aligned} A_0^* &= (\text{sym } \mathring{\nabla})^* = -\text{Div}_{\mathbb{S}} : \mathring{D}(\Omega; \mathbb{S}) \subset \mathbf{L}^2(\Omega; \mathbb{S}) \longrightarrow \mathbf{L}^2(\Omega); & S &\mapsto -\text{Div } S, \\ A_1^* &= (\text{Rot } \mathring{\text{Rot}}_{\mathbb{S}}^{\top})^* = \text{Rot } \text{Rot}_{\mathbb{S}}^{\top} : \mathring{\text{RR}}^{\top}(\Omega; \mathbb{S}) \subset \mathbf{L}^2(\Omega; \mathbb{S}) \longrightarrow \mathbf{L}^2(\Omega; \mathbb{S}); & T &\mapsto \text{Rot } \text{Rot}^{\top} T, \\ A_2^* &= \mathring{\text{Div}}_{\mathbb{S}}^* = -\text{sym } \mathring{\nabla} : \mathring{H}^1(\Omega) \subset \mathbf{L}^2(\Omega) \longrightarrow \mathbf{L}^2(\Omega; \mathbb{S}); & v &\mapsto -\text{sym } \nabla v, \end{aligned}$$

where  $\mathring{D}(\Omega; \mathbb{S}) := \mathring{D}(\Omega) \cap \mathbf{L}^2(\Omega; \mathbb{S})$  and

$$\mathring{\text{RR}}^{\top}(\Omega) := \{S \in \mathbf{L}^2(\Omega) : \text{Rot } \text{Rot}^{\top} S \in \mathbf{L}^2(\Omega)\}, \quad \mathring{\text{RR}}^{\top}(\Omega; \mathbb{S}) := \mathring{\text{RR}}^{\top}(\Omega) \cap \mathbf{L}^2(\Omega; \mathbb{S}).$$

Moreover, for  $S \in \mathring{\text{RR}}^{\top}(\Omega; \mathbb{S})$  it holds  $\text{Rot } \text{Rot}^{\top} S \in \mathbf{L}^2(\Omega; \mathbb{S})$ . Note that  $v$  and  $S, T$  are vector and tensor (matrix) fields, respectively. The complex properties hold as

$$\begin{aligned} R(A_0) &= \text{sym } \mathring{\nabla} \mathring{H}^1(\Omega) \subset \mathring{\text{RR}}_0^{\top}(\Omega; \mathbb{S}) = N(A_1), \\ R(A_1^*) &= \text{Rot } \text{Rot}_{\mathbb{S}}^{\top} \mathring{\text{RR}}^{\top}(\Omega; \mathbb{S}) \subset \mathring{D}_0(\Omega; \mathbb{S}) = N(A_0^*), \\ R(A_1) &= \text{Rot } \mathring{\text{Rot}}_{\mathbb{S}}^{\top} \mathring{\text{RR}}^{\top}(\Omega; \mathbb{S}) \subset \mathring{D}_0(\Omega; \mathbb{S}) = N(A_2), \\ R(A_2^*) &= \text{sym } \mathring{\nabla} \mathring{H}^1(\Omega) \subset \mathring{\text{RR}}_0^{\top}(\Omega; \mathbb{S}) = N(A_1^*). \end{aligned}$$

The sequences (5.1) read

$$\begin{aligned} \mathring{H}^1(\Omega) \subset \mathbf{L}^2(\Omega) &\xrightarrow{A_0 = \text{sym } \mathring{\nabla}} \mathring{\text{RR}}^{\top}(\Omega; \mathbb{S}) \subset \mathbf{L}^2(\Omega; \mathbb{S}) \xrightarrow{A_1 = \text{Rot } \mathring{\text{Rot}}_{\mathbb{S}}^{\top}} \mathring{D}(\Omega; \mathbb{S}) \subset \mathbf{L}^2(\Omega; \mathbb{S}) \xrightarrow{A_2 = \mathring{\text{Div}}_{\mathbb{S}}} \mathbf{L}^2(\Omega), \\ \mathbf{L}^2(\Omega) &\xleftarrow{A_0^* = -\text{Div}_{\mathbb{S}}} \mathring{D}(\Omega; \mathbb{S}) \subset \mathbf{L}^2(\Omega; \mathbb{S}) \xleftarrow{A_1^* = \text{Rot } \text{Rot}_{\mathbb{S}}^{\top}} \mathring{\text{RR}}^{\top}(\Omega; \mathbb{S}) \subset \mathbf{L}^2(\Omega; \mathbb{S}) \xleftarrow{A_2^* = -\text{sym } \mathring{\nabla}} \mathring{H}^1(\Omega) \subset \mathbf{L}^2(\Omega). \end{aligned}$$

These are the so-called Rot Rot complexes, appearing, e.g., in linear elasticity, see [22]. Typical equations arising from the Rot Rot complex are systems of generalized linear elasticity, e.g.,

$$\begin{aligned} A_1 S &= \text{Rot } \mathring{\text{Rot}}_{\mathbb{S}}^{\top} S = F, \\ A_0^* S &= -\text{Div}_{\mathbb{S}} S = f, \end{aligned}$$

or simply linear elasticity and related fourth order Rot Rot Rot Rot systems, e.g.,

$$\begin{aligned} A_0^* A_0 v &= -\operatorname{Div}_{\mathbb{S}} \operatorname{sym} \overset{\circ}{\nabla} v = f, & A_1^* A_1 S &= \operatorname{Rot} \operatorname{Rot}_{\mathbb{S}}^{\top} \operatorname{Rot} \overset{\circ}{\operatorname{Rot}}_{\mathbb{S}}^{\top} S = G, \\ & & A_0^* S &= -\operatorname{Div}_{\mathbb{S}} S = f. \end{aligned}$$

The crucial embeddings (5.2) are compact, compare to Lemma 2.1.

**Lemma 5.22** (elasticity selection theorems). *Let  $\Omega \subset \mathbb{R}^3$  be a strong Lipschitz domain. Then the embeddings*

$$\begin{aligned} D(A_1) \cap D(A_0^*) &= \overset{\circ}{\operatorname{RR}}^{\top}(\Omega; \mathbb{S}) \cap D(\Omega; \mathbb{S}) \hookrightarrow L^2(\Omega; \mathbb{S}), \\ D(A_2) \cap D(A_1^*) &= \overset{\circ}{D}(\Omega; \mathbb{S}) \cap \overset{\circ}{\operatorname{RR}}^{\top}(\Omega; \mathbb{S}) \hookrightarrow L^2(\Omega; \mathbb{S}), \end{aligned}$$

are compact.

A proof can be done by the same techniques showing [22, Lemma 3.22], see [21]. Again, the elasticity selection theorems are independent of possible material law tensors  $\epsilon$  or  $\mu$ . Choosing the pair  $(A_0, A_1)$  we get by Theorem 4.7 the following:

**Theorem 5.23** (global Div-Rot Rot<sup>⊤</sup>- $\mathbb{S}$ -lemma). *Let  $\overset{\circ}{\operatorname{RR}}^{\top}(\Omega; \mathbb{S}) \cap D(\Omega; \mathbb{S}) \hookrightarrow L^2(\Omega; \mathbb{S})$  be compact. Moreover, let  $(S_n) \subset \overset{\circ}{\operatorname{RR}}^{\top}(\Omega; \mathbb{S})$  and  $(T_n) \subset D(\Omega; \mathbb{S})$  be two sequences bounded in  $\overset{\circ}{\operatorname{RR}}^{\top}(\Omega)$  and  $D(\Omega)$ , respectively. Then there exist  $S \in \overset{\circ}{\operatorname{RR}}^{\top}(\Omega; \mathbb{S})$  and  $T \in D(\Omega; \mathbb{S})$  as well as subsequences, again denoted by  $(S_n)$  and  $(T_n)$ , such that*

- $S_n \rightharpoonup S$  in  $\overset{\circ}{\operatorname{RR}}^{\top}(\Omega; \mathbb{S})$ ,
- $T_n \rightharpoonup T$  in  $D(\Omega; \mathbb{S})$ ,
- $\langle S_n, T_n \rangle_{L^2(\Omega; \mathbb{S})} \rightarrow \langle S, T \rangle_{L^2(\Omega; \mathbb{S})}$ .

For the pair  $(A_1, A_2)$  we obtain:

**Theorem 5.24** (global Rot Rot<sup>⊤</sup>-Div- $\mathbb{S}$ -lemma). *Let  $\overset{\circ}{D}(\Omega; \mathbb{S}) \cap \overset{\circ}{\operatorname{RR}}^{\top}(\Omega; \mathbb{S}) \hookrightarrow L^2(\Omega; \mathbb{S})$  be compact. Moreover, let  $(S_n) \subset \overset{\circ}{D}(\Omega; \mathbb{S})$  and  $(T_n) \subset \overset{\circ}{\operatorname{RR}}^{\top}(\Omega; \mathbb{S})$  be two sequences bounded in  $D(\Omega)$  and  $\overset{\circ}{\operatorname{RR}}^{\top}(\Omega)$ , respectively. Then there exist  $S \in \overset{\circ}{D}(\Omega; \mathbb{S})$  and  $T \in \overset{\circ}{\operatorname{RR}}^{\top}(\Omega; \mathbb{S})$  as well as subsequences, again denoted by  $(S_n)$  and  $(T_n)$ , such that*

- $S_n \rightharpoonup S$  in  $\overset{\circ}{D}(\Omega; \mathbb{S})$ ,
- $T_n \rightharpoonup T$  in  $\overset{\circ}{\operatorname{RR}}^{\top}(\Omega; \mathbb{S})$ ,
- $\langle S_n, T_n \rangle_{L^2(\Omega; \mathbb{S})} \rightarrow \langle S, T \rangle_{L^2(\Omega; \mathbb{S})}$ .

**Remark 5.25.** *Let us note:*

- (i) *The Rot Rot complexes of linear elasticity have a strong symmetry.*
- (ii) *Theorem 5.23 and Theorem 5.24 are the same results just with interchanged boundary conditions.*
- (iii) *Theorem 5.23 and Theorem 5.24 have corresponding local versions similar to the local div-rot-lemma Corollary 3.2 and Remark 3.3, which hold with no regularity or boundedness assumptions on  $\Omega$ . As in Remark 5.18 we note that the local versions of Theorem 5.23 and Theorem 5.24 are more involved as well, here due to the second order nature of the Sobolev spaces  $\overset{\circ}{\operatorname{RR}}^{\top}(\Omega; \mathbb{S})$  and  $\overset{\circ}{\operatorname{RR}}^{\top}(\Omega; \mathbb{S})$ . A corresponding non-standard Helmholtz type decomposition similar to [22, Lemma 3.21] is needed to overcome these difficulties.*
- (iv) *Material law tensors  $\epsilon$  and  $\mu$  different from the identities can be handled as well.*

The generalizations in Theorem 4.14 read as follows.

**Theorem 5.26** (generalized/distributional global Div-Rot Rot<sup>⊤</sup>- $\mathbb{S}$ -lemma). *Let the ranges  $\operatorname{sym} \overset{\circ}{\nabla} \overset{\circ}{H}^1(\Omega)$  and  $\operatorname{Rot} \operatorname{Rot}_{\mathbb{S}}^{\top} \overset{\circ}{\operatorname{RR}}^{\top}(\Omega; \mathbb{S})$  be closed and let the generalized Dirichlet-Neumann fields  $\overset{\circ}{\operatorname{RR}}_0^{\top}(\Omega; \mathbb{S}) \cap \overset{\circ}{D}_0(\Omega; \mathbb{S})$  be finite-dimensional. Moreover, let  $(S_n), (T_n) \subset L^2(\Omega; \mathbb{S})$  be two bounded sequences such that*

- $(\widetilde{\text{Rot Rot}}_{\mathbb{S}}^{\top} S_n)$  is relatively compact in  $\text{RR}^{\top}(\Omega; \mathbb{S})'$ ,
- $(\widetilde{\text{Div}}_{\mathbb{S}} T_n)$  is relatively compact in  $\mathring{\text{H}}^1(\Omega)' = \text{H}^{-1}(\Omega)$ .

Then there exist  $S, T \in \text{L}^2(\Omega, \mathbb{S})$  as well as subsequences, again denoted by  $(S_n)$  and  $(T_n)$ , such that

- $S_n \rightharpoonup S$  in  $\text{L}^2(\Omega, \mathbb{S})$ ,
- $T_n \rightharpoonup T$  in  $\text{L}^2(\Omega, \mathbb{S})$ ,
- $\langle S_n, T_n \rangle_{\text{L}^2(\Omega, \mathbb{S})} \rightarrow \langle S, T \rangle_{\text{L}^2(\Omega, \mathbb{S})}$ .

**Theorem 5.27** (generalized/distributional global  $\text{Rot Rot}^{\top}$ -Div- $\mathbb{S}$ -lemma). *Let  $\text{Rot Rot}^{\top} \text{RR}^{\top}(\Omega; \mathbb{S})$  and  $\text{Div } \mathring{\text{D}}(\Omega; \mathbb{S})$  be closed and let the generalized Dirichlet-Neumann fields  $\mathring{\text{D}}_0(\Omega; \mathbb{S}) \cap \text{RR}_0^{\top}(\Omega; \mathbb{S})$  be finite-dimensional. Moreover, let  $(S_n), (T_n) \subset \text{L}^2(\Omega, \mathbb{S})$  be two bounded sequences such that*

- $(\widetilde{\text{Div}}_{\mathbb{S}} S_n)$  is relatively compact in  $\text{H}^1(\Omega)' = \mathring{\text{H}}^{-1}(\Omega)$ ,
- $(\text{Rot Rot}_{\mathbb{S}}^{\top} T_n)$  is relatively compact in  $\text{RR}^{\top}(\Omega; \mathbb{S})'$ .

Then there exist  $S, T \in \text{L}^2(\Omega, \mathbb{S})$  as well as subsequences, again denoted by  $(S_n)$  and  $(T_n)$ , such that

- $S_n \rightharpoonup S$  in  $\text{L}^2(\Omega, \mathbb{S})$ ,
- $T_n \rightharpoonup T$  in  $\text{L}^2(\Omega, \mathbb{S})$ ,
- $\langle S_n, T_n \rangle_{\text{L}^2(\Omega, \mathbb{S})} \rightarrow \langle S, T \rangle_{\text{L}^2(\Omega, \mathbb{S})}$ .

**Remark 5.28.** *By Lemma 5.22 and Lemma 4.18, Theorem 5.23, Theorem 5.24, and Theorem 5.26, Theorem 5.27 hold for strong Lipschitz domains  $\Omega \subset \mathbb{R}^3$ .*

**ACKNOWLEDGEMENTS** The author is grateful to Sören Bartels for bringing up the topic of the div-curl-lemma, and especially to Marcus Waurick for lots of inspiring discussions on the div-curl-lemma and for his substantial contributions to the Special Semester at RICAM in Linz late 2016.

## REFERENCES

- [1] A. Alexanderian. Expository paper: a primer on homogenization of elliptic pdes with stationary and ergodic random coefficient functions. *Rocky Mountain J. Math.*, 45(3):703–735, 2015.
- [2] S. Bartels. Numerical analysis of a finite element scheme for the approximation of harmonic maps into surfaces. *Math. Comp.*, 79(271):1263–1301, 2010.
- [3] S. Bauer, D. Pauly, and M. Schomburg. The Maxwell compactness property in bounded weak Lipschitz domains with mixed boundary conditions. *SIAM J. Math. Anal.*, 48(4):2912–2943, 2016.
- [4] S. Bauer, D. Pauly, and M. Schomburg. The Maxwell compactness property in bounded weak Lipschitz domains with mixed boundary conditions in ND. *Maxwell's Equations: Analysis and Numerics (Radon Series on Computational and Applied Mathematics)*, De Gruyter, 2019.
- [5] D. Braess. *Finite elements: Theory, fast solvers, and applications in elasticity theory*. Cambridge University Press, Cambridge, third edition, 2007.
- [6] M. Briane, J. Casado-Dáz, and F. Murat. The div-curl lemma “trente ans après”: an extension and an application to the g-convergence of unbounded monotone operators. *J. Math. Pures Appl.*, (9) 91(5), 2009.
- [7] G.-Q.G. Chen and S. Li. Global weak rigidity of the Gauss-Codazzi-Ricci equations and isometric immersions of Riemannian manifolds with lower regularity. *J. Geom. Anal.*, 28(3):1957–2007, 2018.
- [8] R. Coifman, P.-L. Lions, Y. Meyer, and S. Semmes. Compensated compactness and Hardy spaces. *J. Math. Pures Appl.*, 72(9)(3):247–286, 1993.
- [9] M. Costabel. A remark on the regularity of solutions of Maxwell's equations on Lipschitz domains. *Math. Methods Appl. Sci.*, 12(4):365–368, 1990.
- [10] L.C. Evans. *Weak convergence methods for nonlinear partial differential equations*. American Mathematical Society, Providence, 1990.
- [11] L.C. Evans. Partial regularity for stationary harmonic maps into spheres. *Arch. Rational Mech. Anal.*, 116(2):101–113, 1991.
- [12] A. Freire, S. Müller, and M. Struwe. Weak compactness of wave maps and harmonic maps. *Ann. Inst. H. Poincaré Anal., Non Linéaire* 15(6):725–754, 1998.
- [13] A. Gloria, S. Neukamm, and F. Otto. Quantification of ergodicity in stochastic homogenization: optimal bounds via spectral gap on glauher dynamics. *Invent. Math.*, 199(2):455–515, 2015.

- [14] F. Jochmann. A compactness result for vector fields with divergence and curl in  $L^q(\Omega)$  involving mixed boundary conditions. *Appl. Anal.*, 66:189–203, 1997.
- [15] H. Kozono and T. Yanagisawa. Global compensated compactness theorem for general differential operators of first order. *Arch. Ration. Mech. Anal.*, 207(3):879–905, 2013.
- [16] R. Leis. *Initial Boundary Value Problems in Mathematical Physics*. Teubner, Stuttgart, 1986.
- [17] F. Murat. Compacité par compensation. *Annali della Scuola Normale Superiore di Pisa-Classe di Scienze*, 5(3):489–507, 1978.
- [18] D. Pauly. On constants in Maxwell inequalities for bounded and convex domains. *Zapiski POMI*, 435:46-54, 2014, *È J. Math. Sci. (N.Y.)*, 210(6):787-792, 2015.
- [19] D. Pauly. On Maxwell’s and Poincaré’s constants. *Discrete Contin. Dyn. Syst. Ser. S*, 8(3):607–618, 2015.
- [20] D. Pauly. On the Maxwell constants in 3D. *Math. Methods Appl. Sci.*, 40(2):435–447, 2017.
- [21] D. Pauly, M. Schomburg, and W. Zulehner. The RotRot-complex and applications to elasticity. *submitted*, 2019.
- [22] D. Pauly and W. Zulehner. The divDiv-complex and applications to biharmonic equations. *Appl. Anal.*, 2019.
- [23] D. Pauly and W. Zulehner. On the duals of Sobolev spaces for vector fields. *submitted*, 2019.
- [24] R. Picard. Zur Theorie der harmonischen Differentialformen. *Manuscripta Math.*, 1:31–45, 1979.
- [25] R. Picard. Randwertaufgaben der verallgemeinerten Potentialtheorie. *Math. Methods Appl. Sci.*, 3:218–228, 1981.
- [26] R. Picard. On the boundary value problems of electro- and magnetostatics. *Proc. Roy. Soc. Edinburgh Sect. A*, 92:165–174, 1982.
- [27] R. Picard. An elementary proof for a compact imbedding result in generalized electromagnetic theory. *Math. Z.*, 187:151–164, 1984.
- [28] T. Rivière. Conservation laws for conformally invariant variational problems. *Invent. Math.*, 168(1):1–22, 2007.
- [29] M. Röger and B. Schweizer. Strain gradient visco-plasticity with dislocation densities contributing to the energy. *Math. Models Methods Appl. Sci.*, 27(4):2595–2629, 2017.
- [30] B. Schweizer. On Friedrichs inequality, Helmholtz decomposition, vector potentials, and the div-curl lemma. *Trends in applications of mathematics to mechanics, Springer INdAM Ser., Cham*, 27:65–79, 2018.
- [31] M. Struwe. *Variational methods. Applications to nonlinear partial differential equations and Hamiltonian systems*. Springer, Berlin, 2008.
- [32] L. Tartar. Compensated compactness and applications to partial differential equations. *Nonlinear analysis and mechanics, Heriot-Watt symposium*, 4:136–211, 1979.
- [33] L. Tartar. *The general theory of homogenization. A personalized introduction*. Springer, Berlin, 2009.
- [34] L. Tartar. Compensated compactness with more geometry. *Springer Proc. Math. Stat.*, 137:74–101, 2015.
- [35] M. Waurick. A functional analytic perspective to the div-curl lemma. *J. Operator Theory*, 80(1):95–111, 2018.
- [36] M. Waurick. Nonlocal H-convergence. *Calc. Var. Partial Differential Equations*, 57(6):Art. 159, 46 pp., 2018.
- [37] C. Weber. A local compactness theorem for Maxwell’s equations. *Math. Methods Appl. Sci.*, 2:12–25, 1980.
- [38] N. Weck. Maxwell’s boundary value problems on Riemannian manifolds with nonsmooth boundaries. *J. Math. Anal. Appl.*, 46:410–437, 1974.
- [39] K.-J. Witsch. A remark on a compactness result in electromagnetic theory. *Math. Methods Appl. Sci.*, 16:123–129, 1993.