

**OPTIMAL QUADRATURE FORMULAS FOR THE
CAUCHY TYPE SINGULAR INTEGRAL IN THE
SOBOLEV SPACE $L_2^{(2)}(0,1)$**

D.M. AKHMEDOV

ABSTRACT. In the present paper in $L_2^{(2)}(0,1)$ S.L.Sobolev space the optimal quadrature formula is constructed for approximate calculation of Cauchy type singular integral.

MSC: 65D32.

Keywords: optimal quadrature formulas; the error functional; the extremal function; Hilbert space; optimal coefficients.

1. Introduction

Many problems of sciences and technics is naturally reduced to singular integral equations. Moreover (see. [1]) plane problems are reduced to one dimensional singular equations. The theory of one-dimensional singular integral equations is given in [2, 3]. It is known that the solutions of such integral equations are expressed by singular integrals. Therefore approximate calculation of the singular integrals with high exactness is actual problem of numerical analysis.

For the singular integral of Cauchy type $\int_0^1 \frac{\varphi(x)}{x-t} dx$ we consider the following quadrature formula

$$\int_0^1 \frac{\varphi(x)}{x-t} dx \cong \sum_{\beta=0}^N C_{\beta} \varphi(x_{\beta}), \quad (1.1)$$

where $0 < t < 1$, C_{β} are the coefficients, x_{β} are the nodes, $N = 2, 3, 4, \dots$, φ is a function of the space $L_2^{(m)}(0,1)$. Here $L_2^{(m)}(0,1)$ is the Sobolev space of functions with a square integrable m th generalized derivative.

To the problem of approximate integration of the Cauchy type singular integrals are devoted many works (see, for instance, [1, 4, 5, 6, 7] and references therein).

In the work [8] in the space for the coefficients of the weight optimal quadrature formulas of the form

$$\int_0^1 p(x) \varphi(x) \cong \sum_{\beta=0}^N C[\beta] \varphi(h\beta) \quad (1.2)$$

the following system of linear equations is obtained

$$\sum_{\gamma=0}^N C[\gamma] \cdot \frac{|h\beta - h\gamma|^{2m-1}}{2(2m-1)!} + P_{m-1}[\beta] = f[\beta], \quad [\beta] \in [0, 1], \quad (1.3)$$

$$\sum_{\beta=0}^N C[\beta] \cdot [\beta]^\alpha = \int_0^1 p(x)x^\alpha dx, \quad \alpha = 0, 1, \dots, m-1, \quad (1.4)$$

where $f[\beta] = \int_0^1 p(x) \frac{|x-h\beta|^{2m-1}}{2(2m-1)!} dx$, $p(x)$ is a weight function, $[\beta] = h\beta$, $h = \frac{1}{N}$,

$C[\beta]$ are unknown coefficients, $P_{m-1}[\beta]$ is a polynomial of degree $m-1$.

In particular, from (1.2) for $p(x) = \frac{1}{x-t}$, $0 < t < 1$ we get quadrature formula (1.1) when $x_\beta = h\beta = [\beta]$ and correspondingly from system (1.3)-(1.4) when $m = 2$, $p(x) = \frac{1}{x-t}$, $0 < t < 1$ we have the following system of linear equations for the coefficients of optimal quadrature formula in the sense of Sard in the form (1.1) when $x_\beta = h\beta$ in the space $L_2^{(2)}(0, 1)$

$$\sum_{\gamma=0}^N C([\gamma], t) \cdot \frac{|h\beta - h\gamma|^3}{12} + p_1 \cdot [\beta] + p_0 = f([\beta], t), \quad \beta = 0, 1, 2, \dots, N, \quad (1.5)$$

$$\sum_{\gamma=0}^N C([\gamma], t) = g_0, \quad (1.6)$$

$$\sum_{\gamma=0}^N C([\gamma], t) \cdot [\gamma] = g_1, \quad (1.7)$$

where

$$f(h\beta, t) = \frac{1}{12} \int_0^1 \frac{|x-h\beta|^3}{x-t} dx = \frac{1}{12} \left(-\frac{11}{3}(h\beta)^3 + (5t+3)(h\beta)^2 - (2t^2+3t+1,5)(h\beta) + (t^2 + \frac{t}{2} + \frac{1}{3}) + (t-h\beta)^3(-2 \ln|h\beta-t| + \ln(t-t^2)) \right), \quad (1.8)$$

$$g_0 = \int_0^1 \frac{dx}{x-t} = \ln \frac{1-t}{t}, \quad (1.9)$$

$$g_1 = \int_0^1 \frac{x}{x-t} dx = 1 + t \ln \frac{1-t}{t}. \quad (1.10)$$

$C([\gamma], t)$, $\gamma = \overline{0, N}$ and p_1, p_0 are unknowns.

The rest of the paper is organized as follows. In Section 2 we give some auxiliary results and definitions. Section 3 is devoted to calculation of the optimal coefficients, i.e. we solve system (1.5)-(1.7).

2. Auxiliary results

In the process of solution of system (1.5)-(1.7) the Sobolev method of construction of optimal quadrature formulas (see [9]) is used. In addition, here we use the concept of discrete argument functions and operations on them. The theory of discrete argument functions is given in [9]. For completeness we give some definitions.

Definition 2.1. The function $\varphi[\beta] = \varphi(h\beta)$ is a *function of discrete argument* if it is given on some set of integer values of β .

Definition 2.2. The *inner product* of two discrete functions $\varphi[\beta]$ and $\psi[\beta]$ is given by

$$[\varphi, \psi] = \sum_{\beta \in B} \varphi[\beta] \cdot \psi[\beta], \quad (2.1)$$

if the series on the right hand side of the last equality converges absolutely.

Definition 2.3. The convolution of two functions $\varphi[\beta]$ and $\psi[\beta]$ is the inner product

$$\chi[\beta] = \varphi[\beta] * \psi[\beta] = [\varphi[\gamma], \psi[\beta - \gamma]] = \sum_{\gamma=-\infty}^{\infty} \varphi[\gamma] \psi[\beta - \gamma]. \quad (2.2)$$

Moreover, we need the discrete analogue $D_2(h\beta)$ of the differential operator d^4/dx^4 , which is defined by the following formula (see [10])

$$D_2(h\beta) = \frac{3!}{h^4} \begin{cases} 3!\sqrt{3}q^{|\beta|}, & |\beta| \geq 2, \\ 19 - 12\sqrt{3}, & |\beta| = 1, \\ 6\sqrt{3} - 8, & \beta = 0, \end{cases} \quad (2.3)$$

where $q = \sqrt{3} - 2$.

We give some properties of the discrete function $D_2(h\beta)$ from [9]:

$$D_2(h\beta) * (h\beta)^\alpha = 0, \quad 0 \leq \alpha < 4, \quad (2.4)$$

$$hD_2(h\beta) * \frac{|h\beta|^3}{12} = \delta(h\beta), \quad (2.5)$$

where $\delta(h\beta)$ is the discrete delta function and $\delta(h\beta) = \begin{cases} 0, & \beta \neq 0, \\ 1, & \beta = 0. \end{cases}$

3. Main result

In the present section we solve system (1.5)-(1.7).

Suppose $C([\beta], t) = 0$ when $\beta < 0$ and $\beta > N$. Then, using Definition 2.3, we rewrite system(1.5)-(1.7) in the following convolution form

$$C([\beta], t) * \frac{|h\beta|^3}{12} + p_1 \cdot (h\beta) + p_0 = f(h\beta, t), \quad \beta = \overline{0, N}, \quad (3.1)$$

$$\sum_{\beta=0}^N C([\beta], t) = g_0, \quad (3.2)$$

$$\sum_{\beta=0}^N C([\beta], t) \cdot (h\beta) = g_1, \quad (3.3)$$

where $f(h\beta)$, g_0 and g_1 are defined by (1.8), (1.9) and (1.10), respectively.

We denote

$$v(h\beta) = C([\beta], t) * \frac{|h\beta|^3}{12}, \quad (3.4)$$

$$u(h\beta) = v(h\beta) + p_1 \cdot (h\beta) + p_0. \quad (3.5)$$

Using the properties (2.4), (2.5) of the operator $D_2(h\beta)$ from (2.3) and (3.5) we have

$$C([\beta], t) = hD_2(h\beta) * u(h\beta). \quad (3.6)$$

But for calculation of the convolution (3.6) we need to define the function $u(h\beta)$ in all integer values of β . When $\beta = 0, 1, 2, \dots, N$ from (3.1) we have $u(h\beta) = f(h\beta, t)$. Therefore it is sufficient to determine the function $u(h\beta)$ when $\beta < 0$ and $\beta > N$.

Further, we define the form of the function $u(h\beta)$ for $\beta \leq 0$ and $\beta \geq N$. From (3.4) taking into account (3.2), (3.3) we have

$$v(h\beta) = \begin{cases} -\frac{1}{12}(h\beta)^3 g_0 + \frac{1}{4}(h\beta)^2 g_1 - (h\beta) \cdot p_1^{(0)} - p_0^{(0)}, & \beta \leq 0, \\ \frac{1}{12}(h\beta)^3 g_0 - \frac{1}{4}(h\beta)^2 g_1 + (h\beta) \cdot p_1^{(0)} + p_0^{(0)}, & \beta \geq N. \end{cases}$$

into account the last equality, from (3.5) for $u(h\beta)$ we obtain

$$u(h\beta) = \begin{cases} -\frac{1}{12}(h\beta)^3 g_0 + \frac{1}{4}(h\beta)^2 g_1 + a_1^-(h\beta) + a_0^-, & \beta \leq 0, \\ f(h\beta, t), & 0 \leq \beta \leq N, \\ \frac{1}{12}(h\beta)^3 g_0 - \frac{1}{4}(h\beta)^2 g_1 + a_1^+(h\beta) + a_0^+, & \beta \geq N, \end{cases} \quad (3.7)$$

where $a_1^-, a_0^-, a_1^+, a_0^+$ are unknowns and

$$\begin{aligned} a_1^- &= p_1 - p_1^{(0)}, & a_0^- &= p_0 - p_0^{(0)}, \\ a_1^+ &= p_1 + p_1^{(0)}, & a_0^+ &= p_0 + p_0^{(0)}. \end{aligned} \quad (3.8)$$

Thus, keeping in mind (2.3), (3.6) and (3.7) we get the following problem

Problem. Find the solution of the equation

$$D_2(h\beta) * u(h\beta) = 0 \text{ where } \beta < 0, \beta > N \quad (3.9)$$

in the form (3.7).

If we find unknowns $a_1^-, a_0^-, a_1^+, a_0^+$, then from (3.8) we have

$$\begin{aligned} p_1 &= \frac{1}{2}(a_1^- + a_1^+), & p_0 &= \frac{1}{2}(a_0^- + a_0^+), \\ p_1^{(0)} &= \frac{1}{2}(a_1^+ - a_1^-), & p_0^{(0)} &= \frac{1}{2}(a_0^+ - a_0^-). \end{aligned} \quad (3.10)$$

Unknowns $a_1^-, a_0^-, a_1^+, a_0^+$ will be found from (3.9), using (2.3) and (3.7). Then we get the explicit form of the function $u(h\beta)$ and from (3.6) we find optimal coefficients $C([\beta], t)$. Moreover, from (3.10) can be found p_1, p_0 . Thus, the stated problem will be solved completely.

The following holds

Theorem 1. *Assume $t \neq h\beta$, $\beta = \overline{0, N}$. Then the coefficients of the optimal quadrature formula (1.1) in the space $L_2^{(2)}(0, 1)$ have the form*

$$C([0], t) = \frac{6}{h^3} \left[\frac{g_0}{12} (h^3 - 3q^N (h^2 + h(q+2))) + \frac{g_1 q^N}{4} (h^2 + 2h(q+2)) + a_1^- h(q+1) + f(0, t)(3q+2) - f(h, t)(12q+5) - q^N (3f(1, t) \times (q+1) + a_1^+ h(q+2)) + 6(q+2) \sum_{\gamma=2}^N q^\gamma f(h\gamma, t) \right], \quad (3.11)$$

$$C([\beta], t) = \frac{6}{h^3} \left[6(q+2) \sum_{\gamma=0}^{\beta-2} q^{\beta-\gamma} f(h\gamma, t) - (12q+5) \left(f(h(\beta-1), t) + f(h(\beta+1), t) \right) + (6q+4)f(h\beta, t) + 6(q+2) \sum_{\gamma=\beta+2}^N q^{\gamma-\beta} f(h\gamma, t) + \frac{g_1}{4} \left(q^{N-\beta} (2h(q+2) + h^2) - q^\beta h^2 \right) + q^\beta \left(a_1^- h(q+2) - 3f(0, t) \times (q+1) \right) - q^{N-\beta} (3f(1, t)(q+1) + \frac{g_0}{4} (h(q+2) + h^2) + a_1^+ h(q+2)) \right], \quad \beta = \overline{1, N-1}, \quad (3.12)$$

$$C([N], t) = \frac{6}{h^3} \left[-\frac{g_0}{12} (3h(q+1) - h^3) + \frac{g_1}{4} (2h(q+1) - q^N h^2) + q^N \left(a_1^- h(q+2) - 3f(0, t)(q+1) \right) - a_1^+ h(q+1) + f(1, t)(3q+2) - f(1-h, t) \times (12q+5) + 6(q+2) \sum_{\gamma=0}^{N-2} q^{N-\gamma} f(h\gamma, t) \right]. \quad (3.13)$$

where

$$a_1^- = \frac{\Delta_1}{\Delta}, \quad (3.14)$$

$$a_1^+ = \frac{\Delta_2}{\Delta}, \quad (3.15)$$

$$a_0^- = f(0, t), \quad (3.16)$$

$$a_0^+ = f(1, t) - \frac{1}{12} g_0 + \frac{1}{4} g_1 - \frac{\Delta_2}{\Delta}, \quad (3.17)$$

$$\begin{aligned}
\Delta &= B_2^2 - A_2^2, \\
\Delta_1 &= A_2 \left[-F_1 - \frac{1}{12}g_0(B_1 + 3B_2 + 3B_3) + \frac{1}{4}g_1(B_1 + 2B_2 + B_3 - A_3) - \right. \\
&\quad \left. -A_1f(0, t) - B_1 \left(f(1, t) - \frac{1}{12}g_0 + \frac{1}{4}g_1 \right) \right] - \\
&\quad -B_2 \left[-F_2 - \frac{1}{12}g_0(A_1 + 3A_2 + 3A_3) + \frac{1}{4}g_1(A_1 + 2A_2 + A_3 - B_3) - \right. \\
&\quad \left. -B_1f(0, t) - A_1 \left(f(1, t) - \frac{1}{12}g_0 + \frac{1}{4}g_1 \right) \right], \\
\Delta_2 &= -A_2 \left[-F_2 - \frac{1}{12}g_0(A_1 + 3A_2 + 3A_3) + \frac{1}{4}g_1(A_1 + 2A_2 + A_3 - B_3) - \right. \\
&\quad \left. -B_1f(0, t) - A_1 \left(f(1, t) - \frac{1}{12}g_0 + \frac{1}{4}g_1 \right) \right] + \\
&\quad +B_2 \left[-F_1 - \frac{1}{12}g_0(B_1 + 3B_2 + 3B_3) + \frac{1}{4}g_1(B_1 + 2B_2 + B_3 - A_3) - \right. \\
&\quad \left. -A_1f(0, t) - B_1 \left(f(1, t) - \frac{1}{12}g_0 + \frac{1}{4}g_1 \right) \right], \\
F_1 &= 6(q+2) \sum_{\gamma=1}^N q^{\gamma+1} f(h\gamma, t) - (12q+5)f(0, t), \\
F_2 &= 6(q+2) \sum_{\gamma=0}^{N-1} q^{N+1-\gamma} f(h\gamma, t) - (12q+5)f(1, t),
\end{aligned}$$

$$\begin{aligned}
A_1 &= 3q+2, & B_1 &= 3q^N(3q+1) \\
A_2 &= h(2q+1), & B_2 &= hq^N(2q+1) \\
A_3 &= h^2q, & B_3 &= -h^2q^{N+1},
\end{aligned}$$

$$q = \sqrt{3} - 2.$$

Proof. From (3.7) for $\beta = 0$ and $\beta = N$ we get (3.16) and (3.17), i.e.

$$a_0^- = f(0, t), \quad (3.18)$$

$$a_0^+ = f(1, t) - \frac{1}{12}g_0 + \frac{1}{4}g_1 - a_1^+. \quad (3.19)$$

From (3.9), using (2.3) and (3.7) for $\beta = -1$ and $\beta = N+1$ we have

$$\begin{aligned}
&-a_1^- \sum_{\gamma=1}^{\infty} D(h\gamma - h)(h\gamma) + a_1^+ \sum_{\gamma=1}^{\infty} D(h(N + \gamma) + h)(h\gamma) = \\
&= - \sum_{\gamma=0}^N D(h\gamma + h)f(h\gamma, t) - \frac{1}{12}g_0 \sum_{\gamma=1}^{\infty} D(h\gamma - h)(h\gamma)^3 - \\
&- \frac{1}{4}g_1 \sum_{\gamma=1}^{\infty} D(h\gamma - h)(h\gamma)^2 - \frac{1}{12}g_0 \sum_{\gamma=1}^{\infty} D(h(N + \gamma) + h)(1 + h\gamma)^3 + \quad (3.20) \\
&+ \frac{1}{4}g_1 \sum_{\gamma=1}^{\infty} D(h(N + \gamma) + h)(1 + h\gamma)^2 - \left(f(1, t) - \frac{1}{12}g_0 + \frac{1}{4}g_1 \right) \times \\
&\times \sum_{\gamma=1}^{\infty} D(h(N + \gamma) + h) - f(0, t) \sum_{\gamma=1}^{\infty} D(h\gamma - h),
\end{aligned}$$

$$\begin{aligned}
& -a_1^- \sum_{\gamma=1}^{\infty} D(h(N+\gamma)+h)(h\gamma) + a_1^+ \sum_{\gamma=1}^{\infty} D(h\gamma-h)(h\gamma) = \\
& = - \sum_{\gamma=0}^N D(h(N+1)-h\gamma)f(h\gamma, t) - \frac{1}{12}g_0 \sum_{\gamma=1}^{\infty} D(h(N+\gamma)+h)(h\gamma)^3 - \\
& - \frac{1}{4}g_1 \sum_{\gamma=1}^{\infty} D(h(N+\gamma)+h)(h\gamma)^2 - \frac{1}{12}g_0 \sum_{\gamma=1}^{\infty} D(h\gamma-h)(1+h\gamma)^3 + \\
& + \frac{1}{4}g_1 \sum_{\gamma=1}^{\infty} D(h\gamma-h)(1+h\gamma)^2 - \left(f(1, t) - \frac{1}{12}g_0 + \frac{1}{4}g_1 \right) \times \\
& \times \sum_{\gamma=1}^{\infty} D(h\gamma-h) - f(0, t) \sum_{\gamma=1}^{\infty} D(h(N+\gamma)+h).
\end{aligned} \tag{3.21}$$

Thus, for unknowns a_1^- , a_1^+ , a_0^- , a_0^+ we obtained the system of linear equations (3.18)-(3.21). Solving this system we obtain (3.14)-(3.17). This means we obtained the explicit form of the function $u(h\beta)$.

Next, using (2.3) and (3.7), from (3.6) calculating the convolution $C([\beta], t) = hD_2(h\beta) * u(h\beta)$ for $\beta = \overline{0, N}$ we respectively get (3.11)-(3.13). Theorem is proved. \square

REFERENCES

- [1] Lifanov I.K. The method of singular equations and numerical experiments. Moscow, TOO "Yanus", 1995. -520 p. (in Russian)
- [2] Gakhov F.D. Boundary problems. -Moscow, Nauka, 1977. -640 p. (in Russian)
- [3] Muskhelishvili N.I. Singular Integral Equations. -M.: Nauka, 1968. -512 p. (in Russian)
- [4] Belotserkovskiy S.M., Lifanov I.K. Numerical methods i singular integral equations. -Moscow, Nauka, 1985. -256 p.
- [5] Gabdulkaev B.G. Cubature formulas for multidimensional singular integrals I // Izvestiya Vuzov. Mathematics. 1975. 4. 3-13. (in Russian)
- [6] Israilov M.I., Shadimetov Kh.M. Weight optimal quadrature formulas for singular integrals of the Cauchy type. Doklady AN RUz -1991, 8, 10-11.
- [7] Shadimetov Kh.M. Optimal quadrature formulas for singular integrals of the Cauchy type. Doklady AN RUz -1987, 6, -Pp.9-11. (in Russian)
- [8] Shadimetov Kh.M. Coefficients of the weight optimal quadrature formulas in the space// Proceedings of the V-the International Conference "Cubature Formulas and Their Applications", -Krasnoyarsk, 1999. -Pp.207-214. (in Russian)
- [9] Sobolev S.L. Introduction to the Theory of Cubature Formulas, Moscow, Nauka, 1974, 808 p. (in Russian).
- [10] Shadimetov Kh.M. The discrete analogue of the operator d^{2m}/dx^{2m} and its properties // Voprosy vichislitelnoy i prikladnoy matematiki. Tashkent, 1985. -Pp.22-35. www.Arxiv.org 2010, January(math.NA).

D.M. AKHMEDOV, INSTITUTE OF MATHEMATICS AND INFORMATION TECHNOLOGIES, TASHKENT, UZBEKISTAN.

E-mail address: hayotov@mail.ru, axmedovdilshod@mail.ru