## A free boundary problem in biological selection models

Jimyeong Lee\*

Gran Sasso Science Institute, Via. F. Crispi 7, 67100 L'Aquila, Italy

July 27, 2021

#### Abstract

We prove local existence for classical solutions of a free boundary problem which arises in one of the biological selection models proposed by Brunet and Derrida, [3]. The problem we consider describes the limit evolution of branching brownian particles on the line with death of the leftmost particle at each creation time as studied in [7]. We use extensively results in [1] and [2].

## 1 Introduction

Brunet and Derrida [3], have proposed a class of models for biological selection processes including the one we consider here. This is a system of N brownian particles on the line which branch independently at rate 1 creating a new brownian particle on the same position of the father; simultaneously the leftmost particle (which is the less fit) disappears. Thus the total number of particles does not change.

In [7] it is proved that the particles density has a limit as N diverges (under suitable assumptions on the initial datum). It is also proved that if the following FBP has a "classical solution" then this is the same as the limit density of the branching brownians, similar results have been obtained for other FBP, see [6].

Let  $b \in \mathbb{R}$ , say b > 0, and  $\rho_0 \in C_c^2([b,\infty))$  such that  $\rho_0(b) = 0$ ,  $\frac{d}{dx}\rho_0(b) = 2$ ,  $\int_b^\infty \rho_0(x)dx = 1$ . The free boundary problem FBP (\*) that we consider is: fixed b and  $\rho_0$  as above find a continuous curve  $L_t$ ,  $t \ge 0$ , starting from  $L_0 = b$ , and  $\rho(x,t)$ ,  $x \ge L_t$ ;  $t \ge 0$ , so that

$$(*) \begin{cases} \rho_t(x,t) = \frac{1}{2}\rho_{xx}(x,t) + \rho(x,t), & \text{if } L_t < x, \ t > 0, \\ \rho(L_t,t) = 0, & \text{if } t \ge 0, \\ \rho(x,0) = \rho_0(x), & \text{if } b \le x, \\ \int_{L_t}^{\infty} \rho(x,t) dx = \int_b^{\infty} \rho_0(x) dx = 1, & \text{if } t > 0. \end{cases}$$

Classical solutions of the FBP (\*) are defined in the next section. In this paper we prove the local in time existence of a classical solution of the FBP (\*), observing that uniqueness follows from the results in [7]. Traveling wave solutions and the fine asymptotics of  $L_t$  are studied in [8] for a large class of FBP including our FBP (\*).

<sup>\*</sup>E-mail: ljm9667@gmail.com

## 2 Main results

By classical solutions we mean the following:

**Definition.** The pair  $(L, \rho)$  is a classical solution of the FBP (\*) in the time interval [0, T], T > 0, if:

- $L \in C^1([0,T]), L_0 = b$
- $\rho \in C(\overline{D_{L,T}}) \cap C^{2,1}(D_{L,T})$ , where  $D_{L,T} = \{(x,t) : L_t < x, \ 0 < t < T\}$
- $(L, \rho)$  satisfies the equations in (\*).

(

**Theorem 1** There is T > 0 so that the FBP (\*) has a classical solution in the time interval [0,T] (denoted by  $(L,\rho)$ ).

We will prove Theorem 1 in the next sections. We will also prove that the derivative  $\rho_x(x,t)$  has a limit when  $x \to L_t$  denoted by  $\rho_x(L_t,t)$  and that it vanishes as  $x \to \infty$ . This together with the conservation law implies  $\rho_x(L_t,t) = 2$  for all  $t \in [0,T]$ . Furthermore if also  $\rho_{xx}(L_t,t)$  exists, by differentiating  $\rho(L_t,t) = 0$  we get  $\dot{L}_t = -\frac{1}{4}\rho_{xx}(L_t,t)$ .

Given  $(L, \rho)$  as above we define  $v(x, t) := e^{-t}\rho_x(x, t)$  and observe that (L, v) satisfies the following free boundary problem FBP (\*\*):

$$\begin{aligned}
(**) \begin{cases}
v_t(x,t) &= \frac{1}{2}v_{xx}(x,t), & \text{if } L_t < x, \quad t > 0, \\
v(L_t,t) &= 2e^{-t}, & \text{if } t > 0, \\
v(x,0) &= h(x), & h \in C_c^1([b,\infty)) \\
\dot{L}_t &= -\frac{1}{4}e^t v_x(L_t,t), & \text{if } t > 0,
\end{aligned}$$

To prove Theorem 1 we will first prove the existence of a classical solution of the FBP (\*\*) in [0,T]:

**Theorem 2** There is T > 0 and a pair (L, v) which satisfies the FBP (\*\*) in [0, T] with:  $L \in C^{1}([0, T]), L_{0} = b$ , and  $v \in C^{2,1}(D_{L,T})$ , where  $D_{L,T} = \{(x, t) : L_{t} < x, 0 < t \leq T\}$ .

We did not find a proof of Theorem 2 in the existing literature, see for instance [4], [5] and references therein. Our proof exploits the one dimensionality of the problem and uses extensively the Cannon estimates, [1], following the strategy proposed by Fasano in [2]. In the next sections we will prove Theorem 2 and then, taking h equal to the space derivative of  $\rho_0$ , we will prove Theorem 1 as a corollary of Theorem 2.

## 3 Strategy of the proof of Theorem 2

The idea is to reduce the analysis of the FBP (\*\*) to a fixed point problem:

- Take a curve  $L_t, t \in [0, T]$  and find v which solves from the first equation to the third one of (\*\*) in the domain  $[L_t, \infty), t \in [0, T]$
- Construct a new curve  $K[L](t) = b \frac{1}{4} \int_0^t e^{\tau} v_x(L_{\tau}, \tau) d\tau$  for  $0 \le t \le T$
- Find L so that K[L] = L and prove that the corresponding pair (L, v) solves the FBP (\*\*)

The first task is to prove existence and smoothness of v, that we do in this section using the lemmas below, see [1]. Let A > 0 and

$$\Sigma(A,T) := \left\{ L \in C[0,T] : L_0 = b, \left| \frac{L_{t_2} - L_{t_1}}{t_2 - t_1} \right| \le A \quad \text{for } 0 \le t_1 < t_2 \le T \right\}.$$

We define  $C_{(1)}^0((0,T])$  as the subspace of C((0,T]) that consists of those functions  $\varphi$  such that

$$\|\varphi\|_{T}^{(1)} = \sup_{0 < t \le T} |\varphi(t)| < \infty.$$
(3.1)

Thus  $C_{(1)}^0((0,T])$  is a Banach space under the norm  $\|\cdot\|_T^{(1)}$ . Let us denote by  $G(x,t;\xi,\tau)$  the Gaussian kernel  $\frac{1}{\sqrt{2\pi(t-\tau)}} \exp\left\{-\frac{|x-\xi|^2}{2(t-\tau)}\right\}$  which satisfies  $G_t = \frac{1}{2}G_{xx}$  and  $G_\tau = -\frac{1}{2}G_{\xi\xi}$ .

**Lemma 1 (jump relation)** Let  $L \in \Sigma(A,T)$  and  $\varphi \in C^0_{(1)}((0,T])$ . Define

$$w_{\varphi}(x,t;L) := \int_0^t G(x,t;L_{\tau},\tau)\varphi(\tau)d\tau.$$

Then

$$\lim_{x \to L_t^+} \frac{\partial w_{\varphi}}{\partial x}(x,t;L) = -\varphi(t) + \int_0^t G_x(L_t,t;L_\tau,\tau)\varphi(\tau)d\tau.$$
(3.2)

**Proof** See Lemma 14.2.5. of [1].

**Lemma 2** Let  $L \in \Sigma(A,T)$ , then there exists a unique  $\varphi \in C^0_{(1)}((0,T])$  such that  $\int_b^\infty h(\xi)G(L_t,t;\xi,0)d\xi - \varphi(t) + \int_0^t G_x(L_t,t;L_\tau,\tau)\varphi(\tau)d\tau = 2e^{-t} \text{ for all } t \in (0,T].$  **Proof** Since  $|G_x(L_t, t; L_\tau, \tau)| \leq \frac{C}{\sqrt{t-\tau}}$  and  $\int_b^\infty h(\xi)G(L_t, t; \xi, 0)d\xi \in C^0_{(1)}((0,T])$ , we have a Volterra integral equation:

$$\varphi(t) = \psi(t) + \int_0^t G_x(L_t, t; L_\tau, \tau)\varphi(\tau)d\tau$$
(3.3)

where  $\psi(t) = -2e^{-t} + \int_b^\infty h(\xi)G(L_t,t;\xi,0)d\xi \in C^0_{(1)}((0,T])$ . As in [1] p.247 the right hand side of 3.3 defines a contraction map from  $C^0_{(1)}((0,T])$  into itself for T small. Thus there is a unique  $\varphi \in C^0_{(1)}((0,T])$  which satisfies (3.3).

**Proposition 1** For any  $L \in \Sigma(A,T)$  let  $\varphi$  as in Lemma 2. Define

$$v(x,t) := \int_b^\infty h(\xi) G(x,t;\xi,0) d\xi + \int_0^t G_x(x,t;L_\tau,\tau)\varphi(\tau) d\tau$$
(3.4)

Then  $v \in C^{2,1}(D_{L,T})$  satisfies from the first equation to the third one of (\*\*) and  $\lim_{x \to \infty} \sup_{0 \le t \le T} |v(x,t)| = 0.$  In addition, v has the right derivative at the boundary  $v_x(L_t,t) \in C^0_{(1)}((0,T]).$ 

### Proof

By (3.4) v satisfies the heat equation with initial datum h and boundary conditions  $v(L_t,t) = 2e^{-t}$ ,  $\lim_{x\to\infty} \sup_{0\le t\le T} |v(x,t)| = 0$ . Let us choose  $R > \sup_{t\in[0,T]} L_t$ . Then we have  $v(R,t) \in C^0_{(1)}((0,T])$  such that by Lemma 14.4.4. and Theorem 14.4.1. of [1], we also obtain  $v_x(L_t,t) \in C^0_{(1)}((0,T])$ .

# 4 Proof of Theorem 2

Theorem 2 is proved at the end of the section.

**Lemma 3** Let  $L \in \Sigma(A,T)$  and v as in (3.4), then there are positive constants  $C_1, C_2, C_3$  such that

$$|v_x(L_t, t)| \le C_1 \int_0^t \frac{|v_x(L_\tau, \tau)|}{\sqrt{t - \tau}} d\tau + (C_2 \sqrt{t} + C_3).$$
(4.5)

•

Moreover, we have  $v_x(L_t, t) \in C([0, T])$ .

### Proof

Let us fix  $(x,t) \in D_{L,T}$  and let us define  $D_{\epsilon,R}^{(t)} := \{(\xi,\tau) : L_{\tau} + \epsilon < \xi < R, \ \epsilon < \tau < t - \epsilon\}$  for each  $\epsilon > 0$  and  $R \in \mathbb{R}$ . By the Green's identity, we have

$$\frac{1}{2}(v_{\xi}G - vG_{\xi})_{\xi} - (vG)_{\tau} = 0 \implies \oint_{\partial D_{\epsilon,R}^{(t)}} \frac{1}{2}(v_{\xi}G - vG_{\xi})d\tau + (vG)d\xi = 0.$$
(4.6)

Hence we obtain another representation of v by letting  $\epsilon \to 0, R \to \infty$ ,

$$v(x,t) = \int_{b}^{\infty} h(\xi)G(x,t;\xi,0)d\xi - \frac{1}{2}\int_{0}^{t} G(x,t;L_{\tau},\tau)v_{\xi}(L_{\tau},\tau)d\tau + \int_{0}^{t} e^{-\tau}G_{\xi}(x,t;L_{\tau},\tau)d\tau.$$
(4.7)

Differentiating both sides of (4.7) with respect to x and integrating by parts, we get

$$v_x(x,t) = \int_b^\infty h_{\xi}(\xi) G(x,t;\xi,0) d\xi - \frac{1}{2} \int_0^t G_x(x,t;L_{\tau},\tau) v_x(L_{\tau},\tau) d\tau + 2 \int_0^t e^{-\tau} G(x,t;L_{\tau},\tau) d\tau,$$
(4.8)

$$\frac{1}{2}v_x(L_t,t) = \int_b^\infty h_\xi(\xi)G(L_t,t;\xi,0)d\xi - \frac{1}{2}\int_0^t G_x(L_t,t;L_\tau,\tau)v_x(L_\tau,\tau)d\tau + 2\int_0^t e^{-\tau}G(L_t,t;L_\tau,\tau)d\tau.$$
(4.9)

Since  $|G_x(L_t, t; L_\tau, \tau)| \leq \frac{C}{\sqrt{t-\tau}}$  and there are constants  $C_2$  and  $C_3$  such that

$$\left| 2 \int_0^t e^{-\tau} G(x,t;L_{\tau},\tau) d\tau \right| \le C_2 \sqrt{t}, \quad \left| \int_b^\infty h_{\xi}(\xi) G(x,t;\xi,0) d\xi \right| \le \|h_{\xi}\|_{\infty} \le C_3, \quad (4.10)$$

then

$$|v_x(L_t, t)| \le (C_2\sqrt{t} + C_3) + C_1 \int_0^t \frac{|v_x(L_\tau, \tau)|}{\sqrt{t - \tau}} d\tau.$$
(4.11)

Using (4.9), we have

$$v_x(L_t, t) \in C([0, T]).$$
 (4.12)

•

**Lemma 4** Let  $A > \frac{C_3}{4}$ ,  $C_3$  as in Lemma 3. Then for all sufficiently small T,  $K[L](t) := b - \frac{1}{4} \int_0^t e^{\tau} v_x(L_{\tau}, \tau) d\tau$ ,  $0 \le t \le T$ , maps  $K : \Sigma(A, T) \longrightarrow \Sigma(A, T)$ .

**Proof** By Lemma 17.7.1. in [1] applied to (4.11), we obtain

$$|v_x(L_t, t)| \le [1 + 2C_1\sqrt{T}] \exp\{\pi C_1^2 T\} (C_2\sqrt{t} + C_3).$$
(4.13)

Then  $\left|\frac{d}{dt}K[L](t)\right| = \frac{1}{4}\left|e^t v_x(L_t,t)\right| \le \frac{1}{4}e^T[1+2C_1\sqrt{T}]\exp\{\pi C_1^2 T\}(C_2\sqrt{T}+C_3) \le A$  for all sufficiently small T so that the map K is well defined.

Hereafter  $A > \frac{C_3}{4}$  is fixed. We will show that K is continuous, then, since  $\Sigma(A, T)$  is convex and compact we can apply the Schauder fixed point theorem to conclude that K has a fixed point.

**Lemma 5** For any sufficiently small T > 0 the map K defined in Lemma 4 is continuous on  $\Sigma(A, T)$  with sup norm.

### Proof

Let T > 0 be such that 0 < b - AT and so small that K is well defined. For  $L \in \Sigma(A, T)$ , we have  $0 < b - AT \leq \inf L \leq \sup L \leq b + AT$ . Let v be the function determined by L via Proposition 1. Then v satisfies Green's identity as follows; for  $D_{\epsilon,R}^{(t)}$  defined in the proof of Lemma 3,

$$\oint_{\partial D_{\epsilon,R}^{(t)}} [\xi v d\xi + \frac{1}{2} (\xi v_{\xi} - v) d\tau] = 0.$$
(4.14)

Let  $v^{(1)}, v^{(2)}$  be the functions which correspond to  $L^{(1)}, L^{(2)} \in \Sigma(A, T)$ . We denote  $K[L^{(i)}] = \sigma^{(i)}, i = 1, 2$ , we apply (4.14) and let  $\epsilon \to 0, R \to \infty$ . We get

$$\begin{split} \int_0^t 2e^{-\tau} L_{\tau}^{(1)} \left[ \frac{d}{d\tau} (\sigma_{\tau}^{(1)} - \sigma_{\tau}^{(2)}) \right] d\tau + \int_0^t 2e^{-\tau} \left[ \frac{d}{d\tau} \sigma_{\tau}^{(2)} \right] (L_{\tau}^{(1)} - L_{\tau}^{(2)}) d\tau \\ &= \int_{L_t^{(1)}}^\infty \xi v^{(1)}(\xi, t) d\xi - \int_{L_t^{(2)}}^\infty \xi v^{(2)}(\xi, t) d\xi. \end{split}$$

By integration by parts we have

$$2e^{-t}L_{t}^{(1)}[\sigma_{t}^{(1)} - \sigma_{t}^{(2)}] = \int_{0}^{t} (-2e^{-\tau}L_{\tau}^{(1)} + 2e^{-\tau}\frac{d}{d\tau}L_{\tau}^{(1)})(\sigma_{\tau}^{(1)} - \sigma_{\tau}^{(2)})d\tau$$
$$-\int_{0}^{t} 2e^{-\tau} \left[\frac{d}{d\tau}\sigma_{\tau}^{(2)}\right] (L_{\tau}^{(1)} - L_{\tau}^{(2)})d\tau + \left[\int_{L_{t}^{(1)}}^{\infty} \xi v^{(1)}(\xi, t)d\xi - \int_{L_{t}^{(2)}}^{\infty} \xi v^{(2)}(\xi, t)d\xi\right]. \quad (4.15)$$

To control  $\int_{L_t^{(1)}}^{\infty} \xi v^{(1)}(\xi, t) d\xi - \int_{L_t^{(2)}}^{\infty} \xi v^{(2)}(\xi, t) d\xi$ , using (4.7) and Fubini's theorem, we obtain for i = 1, 2:

Taking the difference for the first term of  $\left(4.16\right)$  , we have

$$\begin{split} I_1 &:= \int_b^\infty h(\xi) t[G(L_t^{(1)}, t; \xi, 0) - G(L_t^{(2)}, t; \xi, 0)] d\xi + \int_{L_t^{(1)}}^{L_t^{(2)}} \int_b^\infty \xi h(\xi) G(x, t; \xi, 0) d\xi dx. \\ \text{so that } |I_1| &\leq C \sup_{0 \leq \tau \leq T} \left| L_\tau^{(1)} - L_\tau^{(2)} \right|. \end{split}$$

Since

$$-\frac{1}{2}\int_{0}^{t} v_{\xi}^{(i)}(L_{\tau}^{(i)},\tau) \int_{L_{t}^{(i)}}^{\infty} xG(x,t;L_{\tau}^{(i)},\tau) dx d\tau = 2\int_{0}^{t} e^{-\tau} \left[\frac{d}{d\tau}\sigma_{\tau}^{(i)}\right] \int_{L_{t}^{(i)}}^{\infty} xG(x,t;L_{\tau}^{(i)},\tau) dx d\tau,$$

we also have by taking the difference for the second term of (4.16),

$$I_{2} := 2 \int_{0}^{t} e^{-\tau} \frac{d}{d\tau} (\sigma_{\tau}^{(1)} - \sigma_{\tau}^{(2)}) \int_{L_{t}^{(1)}}^{\infty} x G(x, t; L_{\tau}^{(1)}, \tau) dx d\tau$$
$$+ 2 \int_{0}^{t} \left[ \int_{L_{t}^{(1)}}^{\infty} x G(x, t; L_{\tau}^{(1)}, \tau) dx - \int_{L_{t}^{(2)}}^{\infty} x G(x, t; L_{\tau}^{(2)}, \tau) dx \right] e^{-\tau} \frac{d}{d\tau} \sigma_{\tau}^{(2)} d\tau.$$
(4.17)

By integration by parts, we get for the first term on right hand side of (4.17)

$$2\int_0^t e^{-\tau} \frac{d}{d\tau} (\sigma_\tau^{(1)} - \sigma_\tau^{(2)}) \int_{L_t^{(1)}}^\infty x G(x, t; L_\tau^{(1)}, \tau) dx d\tau$$
  
=  $e^{-t} [\sigma_t^{(1)} - \sigma_t^{(2)}] L_t^{(1)} - 2\int_0^t [\sigma_\tau^{(1)} - \sigma_\tau^{(2)}] \frac{\partial}{\partial \tau} \left( e^{-\tau} \int_{L_t^{(1)}}^\infty x G(x, t; L_\tau^{(1)}, \tau) dx \right) d\tau.$ 

so that

$$I_{2} = e^{-t} [\sigma_{t}^{(1)} - \sigma_{t}^{(2)}] L_{t}^{(1)} - 2 \int_{0}^{t} [\sigma_{\tau}^{(1)} - \sigma_{\tau}^{(2)}] \frac{\partial}{\partial \tau} \left( e^{-\tau} \int_{L_{t}^{(1)}}^{\infty} x G(x, t; L_{\tau}^{(1)}, \tau) dx \right) d\tau$$
$$+ 2 \int_{0}^{t} \left[ \int_{L_{t}^{(1)}}^{\infty} x G(x, t; L_{\tau}^{(1)}, \tau) dx - \int_{L_{t}^{(2)}}^{\infty} x G(x, t; L_{\tau}^{(2)}, \tau) dx \right] e^{-\tau} \frac{d}{d\tau} \sigma_{\tau}^{(2)} d\tau.$$
(4.18)

To estimate the last term of (4.18), we use the following identity for i=1,2;

$$\int_{L_t^{(i)}}^{\infty} x G(x,t;L_{\tau}^{(i)},\tau) dx = -(t-\tau) \int_{L_t^{(i)}}^{\infty} G_x(x,t;L_{\tau}^{(i)},\tau) dx + L_{\tau}^{(i)} \int_{L_t^{(i)}}^{\infty} G(x,t;L_{\tau}^{(i)},\tau) dx$$

$$= (t-\tau)G(L_t^{(i)}, t; L_{\tau}^{(i)}, \tau) + L_{\tau}^{(i)} \int_{L_t^{(i)}}^{\infty} G(x, t; L_{\tau}^{(i)}, \tau) dx = (t-\tau)G(L_t^{(i)}, t; L_{\tau}^{(i)}, \tau) + L_{\tau}^{(i)}\Psi\left(\frac{L_t^{(i)} - L_{\tau}^{(i)}}{\sqrt{t-\tau}}\right),$$
  
where  $\Psi(x) = \int_x^{\infty} \frac{1}{\sqrt{2\pi}} \exp\left\{-\frac{|z|^2}{2}\right\} dz.$ 

By applying the mean value theorem on  $\Psi$ , we have

$$\left|\Psi\left(\frac{L_t^{(1)} - L_\tau^{(1)}}{\sqrt{t - \tau}}\right) - \Psi\left(\frac{L_t^{(2)} - L_\tau^{(2)}}{\sqrt{t - \tau}}\right)\right| \le C_1 \left|\frac{L_t^{(1)} - L_\tau^{(1)} - L_t^{(2)} + L_\tau^{(2)}}{\sqrt{t - \tau}}\right| \le \frac{C_1 \sup_{0 \le \eta \le T} \left|L_\eta^{(1)} - L_\eta^{(2)}\right|}{\sqrt{t - \tau}}\right|$$

By applying the mean value theorem on  $\exp\left\{-\frac{|\cdot|^2}{2}\right\}$ , for some  $0 < \theta < 1$ ,

$$\begin{split} |G(L_t^{(1)}, t; L_{\tau}^{(1)}, \tau) - G(L_t^{(2)}, t; L_{\tau}^{(2)}, \tau)| \\ &\leq \frac{1}{\sqrt{2\pi(t-\tau)}} \left| \theta \frac{L_t^{(1)} - L_{\tau}^{(1)}}{\sqrt{t-\tau}} + (1-\theta) \frac{L_t^{(2)} - L_{\tau}^{(2)}}{\sqrt{t-\tau}} \right| \left| \frac{L_t^{(1)} - L_{\tau}^{(1)} - L_t^{(2)} + L_{\tau}^{(2)}}{\sqrt{t-\tau}} \right| \\ &\leq \frac{C_2 \sup_{0 \leq \eta \leq T} \left| L_{\eta}^{(1)} - L_{\eta}^{(2)} \right|}{\sqrt{t-\tau}} \end{split}$$

Thus we obtain

$$\left| \int_{L_t^{(1)}}^{\infty} x G(x,t;L_{\tau}^{(1)},\tau) dx - \int_{L_t^{(2)}}^{\infty} x G(x,t;L_{\tau}^{(2)},\tau) dx \right| \le C_3 \sup_{0 \le \eta \le T} \left| L_{\eta}^{(1)} - L_{\eta}^{(2)} \right| + C_4 \frac{\sup_{0 \le \eta \le T} \left| L_{\eta}^{(1)} - L_{\eta}^{(2)} \right|}{\sqrt{t-\tau}} + C_4 \frac{\sup_{0 \le \eta \le T} \left| L_{\eta}^{(1)} - L_{\eta}^{(2)} \right|}{\sqrt{t-\tau}} + C_4 \frac{\sup_{0 \le \eta \le T} \left| L_{\eta}^{(1)} - L_{\eta}^{(2)} \right|}{\sqrt{t-\tau}} + C_4 \frac{\sup_{0 \le \eta \le T} \left| L_{\eta}^{(1)} - L_{\eta}^{(2)} \right|}{\sqrt{t-\tau}} + C_4 \frac{\sup_{0 \le \eta \le T} \left| L_{\eta}^{(1)} - L_{\eta}^{(2)} \right|}{\sqrt{t-\tau}} + C_4 \frac{\sup_{0 \le \eta \le T} \left| L_{\eta}^{(1)} - L_{\eta}^{(2)} \right|}{\sqrt{t-\tau}} + C_4 \frac{\sup_{0 \le \eta \le T} \left| L_{\eta}^{(1)} - L_{\eta}^{(2)} \right|}{\sqrt{t-\tau}} + C_4 \frac{\sup_{0 \le \eta \le T} \left| L_{\eta}^{(1)} - L_{\eta}^{(2)} \right|}{\sqrt{t-\tau}} + C_4 \frac{\sup_{0 \le \eta \le T} \left| L_{\eta}^{(1)} - L_{\eta}^{(2)} \right|}{\sqrt{t-\tau}} + C_4 \frac{\sup_{0 \le \eta \le T} \left| L_{\eta}^{(1)} - L_{\eta}^{(2)} \right|}{\sqrt{t-\tau}} + C_4 \frac{\sup_{0 \le \eta \le T} \left| L_{\eta}^{(1)} - L_{\eta}^{(2)} \right|}{\sqrt{t-\tau}} + C_4 \frac{\sup_{0 \le \eta \le T} \left| L_{\eta}^{(1)} - L_{\eta}^{(2)} \right|}{\sqrt{t-\tau}} + C_4 \frac{\sup_{0 \le \eta \le T} \left| L_{\eta}^{(1)} - L_{\eta}^{(2)} \right|}{\sqrt{t-\tau}} + C_4 \frac{\sup_{0 \le \eta \le T} \left| L_{\eta}^{(1)} - L_{\eta}^{(2)} \right|}{\sqrt{t-\tau}} + C_4 \frac{\sup_{0 \le \eta \le T} \left| L_{\eta}^{(1)} - L_{\eta}^{(2)} \right|}{\sqrt{t-\tau}} + C_4 \frac{\sup_{0 \le \eta \le T} \left| L_{\eta}^{(1)} - L_{\eta}^{(2)} \right|}{\sqrt{t-\tau}} + C_4 \frac{\sup_{0 \le \eta \le T} \left| L_{\eta}^{(1)} - L_{\eta}^{(2)} \right|}{\sqrt{t-\tau}} + C_4 \frac{\sup_{0 \le T} \left| L_{\eta}^{(1)} - L_{\eta}^{(2)} \right|}{\sqrt{t-\tau}} + C_4 \frac{\sup_{0 \le T} \left| L_{\eta}^{(1)} - L_{\eta}^{(2)} \right|}{\sqrt{t-\tau}} + C_4 \frac{\sup_{0 \le T} \left| L_{\eta}^{(1)} - L_{\eta}^{(2)} \right|}{\sqrt{t-\tau}} + C_4 \frac{\sup_{0 \le T} \left| L_{\eta}^{(1)} - L_{\eta}^{(2)} \right|}{\sqrt{t-\tau}} + C_4 \frac{\sup_{0 \le T} \left| L_{\eta}^{(1)} - L_{\eta}^{(2)} \right|}{\sqrt{t-\tau}} + C_4 \frac{\sup_{0 \le T} \left| L_{\eta}^{(1)} - L_{\eta}^{(2)} \right|}{\sqrt{t-\tau}} + C_4 \frac{\sup_{0 \le T} \left| L_{\eta}^{(1)} - L_{\eta}^{(2)} \right|}{\sqrt{t-\tau}} + C_4 \frac{\sup_{0 \le T} \left| L_{\eta}^{(2)} - L_{\eta}^{(2)} \right|}{\sqrt{t-\tau}} + C_4 \frac{\sup_{0 \le T} \left| L_{\eta}^{(2)} - L_{\eta}^{(2)} \right|}{\sqrt{t-\tau}} + C_4 \frac{\sup_{0 \le T} \left| L_{\eta}^{(2)} - L_{\eta}^{(2)} \right|}{\sqrt{t-\tau}} + C_4 \frac{\sup_{0 \le T} \left| L_{\eta}^{(2)} - L_{\eta}^{(2)} \right|}{\sqrt{t-\tau}} + C_4 \frac{\sup_{0 \le T} \left| L_{\eta}^{(2)} - L_{\eta}^{(2)} \right|}{\sqrt{t-\tau}} + C_4 \frac{\sup_{0 \le T} \left| L_{\eta}^{(2)} - L_{\eta}^{(2)} \right|}{\sqrt{t-\tau}} + C_4 \frac{\sup_{0 \le T} \left| L_{\eta}^{(2)} - L_{\eta}^{(2)} \right|}{\sqrt{t-\tau}}$$

Also we observe

$$\left| \frac{\partial}{\partial \tau} \left( e^{-\tau} \int_{L_t^{(1)}}^{\infty} x G(x,t;L_\tau^{(1)},\tau) dx \right) d\tau \right|$$
$$= \left| \frac{\partial}{\partial \tau} \left[ e^{-\tau} \left\{ (t-\tau) G(L_t^{(1)},t;L_\tau^{(1)},\tau) + L_\tau^{(1)} \Psi\left(\frac{L_t^{(1)} - L_\tau^{(1)}}{\sqrt{t-\tau}}\right) \right\} \right] \right| \le C_5 + \frac{C_6}{\sqrt{t-\tau}}.$$

For the third term of (4.16) we have

$$\int_0^t e^{-\tau} \int_{L_t^{(i)}}^\infty x G_{\xi}(x,t;L_{\tau}^{(i)},\tau) dx d\tau = \int_0^t e^{-\tau} \left[ L_t^{(i)} G(L_t^{(i)},t;L_{\tau}^{(i)},\tau) + \int_{L_t^{(i)}}^\infty G(x,t;L_{\tau}^{(i)},\tau) \right] d\tau.$$

Then we have

$$I_3 := \int_0^t e^{-\tau} \left[ L_t^{(1)} G(L_t^{(1)}, t; L_\tau^{(1)}, \tau) + \int_{L_t^{(1)}}^\infty G(x, t; L_\tau^{(1)}, \tau) - L_t^{(2)} G(L_t^{(2)}, t; L_\tau^{(2)}, \tau) - \int_{L_t^{(2)}}^\infty G(x, t; L_\tau^{(2)}, \tau) \right] d\tau$$

so that

$$|I_3| \le C_7 \sup_{0 \le \eta \le T} \left| L_{\eta}^{(1)} - L_{\eta}^{(2)} \right|.$$

Combining all the results and (4.15) finally we obtain for all sufficiently small T > 0,

$$\begin{aligned} \left| \sigma_t^{(1)} - \sigma_t^{(2)} \right| &\leq C_8 \int_0^t \left| \sigma_\tau^{(1)} - \sigma_\tau^{(2)} \right| d\tau + C_9 \int_0^t \frac{\left| \sigma_\tau^{(1)} - \sigma_\tau^{(2)} \right|}{\sqrt{t - \tau}} d\tau + C_{10} \sup_{0 \leq \tau \leq T} \left| L_\tau^{(1)} - L_\tau^{(2)} \right| \\ &\leq (C_8 + C_9) \int_0^t \frac{\left| \sigma_\tau^{(1)} - \sigma_\tau^{(2)} \right|}{\sqrt{t - \tau}} d\tau + C_{10} \sup_{0 \leq \tau \leq T} \left| L_\tau^{(1)} - L_\tau^{(2)} \right|. \end{aligned}$$

By Lemma 17.7.1 in [1], we get  $\sup_{0 \le \tau \le T} \left| \sigma_t^{(1)} - \sigma_t^{(2)} \right| \le C_{11} \sup_{0 \le \tau \le T} \left| L_{\tau}^{(1)} - L_{\tau}^{(2)} \right|$  so that K is a continuous map.

### Proof of Theorem 2

Let (L, v) as in Lemma 5 then by (4.12) K[L] = L is in  $C^1([0, T])$ . This completes the proof.

## 5 Proof of Theorem 1

Suppose that  $h(x) = \frac{d\rho_0(x)}{dx}$  and let (L, v) be as in Theorem 2 with such h as initial condition. Let  $\rho(x,t) := \int_{L_t}^x e^t v(y,t) dy$ . It can be readily shown that  $\rho$  solves from the first to the third equations in (\*), see Section 1. To prove the last equation in (\*) we first remark that by (3.4)  $v \in L^1([L_t,\infty))$ . We then differentiate  $\int_{L_t}^\infty v(y,t) dy$  with respect to t:

$$\frac{d}{dt} \left( \int_{L_t}^{\infty} v(x,t) dx \right) = -\dot{L}_t v(L_t,t) + \int_{L_t}^{\infty} v_t(x,t) dx$$
$$= -2\dot{L}_t e^{-t} + \int_{L_t}^{\infty} \frac{1}{2} v_{xx}(x,t) = -2\dot{L}_t e^{-t} - \frac{1}{2} v_x(L_t,t) = 0.$$

Since  $\int_{b}^{\infty} h(x)dx = \int_{b}^{\infty} \frac{d\rho_{0}(x)}{dx}dx = 0$ , we obtain that  $\int_{L_{t}}^{\infty} v(x,t)dx = 0$ . By using this, (3.4), and Fubini's theorem, we also have

$$\rho(x,t) = e^t \left[ \int_0^t G(x,t;L_\tau,\tau)\varphi(\tau)d\tau - \int_b^\infty h(\xi) \int_x^\infty G(y,t;\xi,0)dyd\xi \right]$$
(5.19)

so that  $\int_{L_t}^{\infty} \rho(x,t)dx < \infty$ . Similarly, if we differentiate  $\int_{L_t}^{\infty} \rho(x,t)dx$  with respect to t and get  $\frac{d}{dt} \left( \int_{L_t}^{\infty} \rho(x,t)dx \right) = \int_{L_t}^{\infty} \rho_t(x,t)dx = \int_{L_t}^{\infty} \left( \frac{1}{2} \rho_{xx}(x,t) + \rho(x,t) \right) dx =$  $-\frac{1}{2} \rho_x(L_t,t) + \int_{L_t}^{\infty} \rho(x,t)dx = -1 + \int_{L_t}^{\infty} \rho(x,t)dx$ . Since  $\int_b^{\infty} \rho_0(x)dx = 1$ , we obtain that  $\int_{L_t}^{\infty} \rho(x,t)dx = 1$ . The derivative  $\rho_x(x,t)$  has a limit when  $x \to L_t$  and  $\rho_x(L_t,t) = 2$ and vanishes as  $x \to \infty$  by Proposition 1. This completes the proof of Theorem 1; the statements below Theorem 1 also follow from what proved for v(x,t).

Acknowledgments. I thank A. De Masi, P. Ferrari and E. Presutti for useful discussions.

## References

- J.R. Cannon: The One-Dimensional Heat Equation, Addison-Wesley Publishing Company (1984).
- [2] A. Fasano, Mathematical models of some diffusive processes with free boundaries, SIMAI e-Lecture Notes (2008).
- [3] E.Brunet, B.Derrida, Shift in the velocity of a front due to a cutoff, Phys. Rev. E (3) 56 2597-2604. MR1473413 (1997).
- [4] C.M.Brauner, J.Hulshof, A General Approach to Stability in Free Boundary Problems, Journal of Differential Equations 164, 1648 (2000).
- [5] L.A. Caffarelli, J.L.Vazquez, A free-boundary problem for the heat equation arising in flame propagation, Transactions of the American Mathematical Society, Volume 347, Number 2, February (1995).
- [6] G.Carinci, A. De Masi, C.Giardinà, E.Presutti, Free Boundary Problems in PDEs and Particle Systems, Springer (2016).
- [7] A. De Masi, P.Ferrari, E.Presutti, N.Soprano-Loto, Hydrodynamics of the NBBM process, Preprint, https://arxiv.org/abs/1707.00799 (2017).
- [8] J.Berestycki, E.Brunet, B.Derrida, Exact solution and precise asymptotics of a Fisher-KPP type front, Preprint, https://arxiv.org/abs/1705.08416 (2017).