## Linear systems of wave equations on cosmological backgrounds with convergent asymptotics

Hans Ringström KTH Royal Institute of Technology and Institut Mittag-Leffler

January 14, 2021

# Contents

Ι	In	troductory material, results	1
1	Eqι	nations and questions	3
	1.1	Introduction	3
	1.2	Equations	4
	1.3	Metrics with convergent asymptotics	8
	1.4	Questions	14
	1.5	Outline of the introductory part	15
	1.6	Acknowledgements	18
<b>2</b>	On	the notion of balance	19
	2.1	Introduction	19
	2.2	Super exponential growth, first examples	20
	2.3	Gevrey classes	22
	2.4	Unbalanced equations, generic super exponential growth	24
	2.5	A rough idea of the argument in a model case	25
3	A g	eometric perspective	27
	3.1	Introduction	27
	3.2	Conditions on the metric	27
	3.3	The lower order terms	29
	3.4	A rough notion of balance	29
	3.5	Geometric formulation, unbalanced setting	30
4	Sile	nt equations	33
	4.1	Introduction	33
	4.2	Results	35
	4.3	Examples	37
	4.4	Strengths/weaknesses; argument	39
5	Tra	nsparent equations	43
	51		43
	0.1	Introduction	-10

	5.3	Future asymptotics	49
	5.4	Specifying the asymptotics	51
	5.5	An example: the flat Kasner solution	52
	5.6	Outline of the argument	53
6	Equ	ations with a dominant noisy spatial direction	57
	6.1	Introduction	57
	6.2	Results	62
	6.3	Outline of the proof	71
7	The	asymptotically diagonal setting	73
	7.1	Introduction	73
	7.2	Asymptotically diagonal equations	77
	7.3	The constituent growth/decay rates	78
	7.4	Energy estimates	81
	7.5	Outline of the argument, outlook	83
8	Imp	roved asymptotic estimates in the silent setting	87
	8.1	Estimating the asymptotic data in terms of the initial data	87
	8.2	Estimating the initial data in terms of the asymptotic data $\ldots \ldots \ldots \ldots \ldots$	89
	8.3	Outline of the argument, outlook	90
тт	R	ough analysis in the silent and in the transparent settings	93
11	10	ough analysis in the sheat and in the transparent settings	50
9	OD		
	UD.	E's with exponentially decaying error terms	95
	9.1	E's with exponentially decaying error terms Equations and algebraic decompositions	<b>95</b> 96
	9.1 9.2	E's with exponentially decaying error terms Equations and algebraic decompositions	<b>95</b> 96 99
	9.1 9.2 9.3	E's with exponentially decaying error terms Equations and algebraic decompositions	<b>95</b> 96 99 101
	9.1 9.2 9.3 9.4	E's with exponentially decaying error terms Equations and algebraic decompositions	<b>95</b> 96 99 101 104
10	9.1 9.2 9.3 9.4 <b>Asy</b>	E's with exponentially decaying error terms Equations and algebraic decompositions	<ul> <li>95</li> <li>96</li> <li>99</li> <li>101</li> <li>104</li> <li>107</li> </ul>
10	9.1 9.2 9.3 9.4 9 Asy 10.1	E's with exponentially decaying error terms Equations and algebraic decompositions	<ul> <li>95</li> <li>96</li> <li>99</li> <li>101</li> <li>104</li> <li>107</li> <li>107</li> </ul>
10	9.1 9.2 9.3 9.4 • Asy 10.1 10.2	E's with exponentially decaying error terms Equations and algebraic decompositions	<ul> <li>95</li> <li>96</li> <li>99</li> <li>101</li> <li>104</li> <li>107</li> <li>108</li> </ul>
10	9.1 9.2 9.3 9.4 0 Asy 10.1 10.2 10.3	E's with exponentially decaying error terms Equations and algebraic decompositions	<ul> <li>95</li> <li>96</li> <li>99</li> <li>101</li> <li>104</li> <li>107</li> <li>107</li> <li>108</li> <li>109</li> </ul>
10	9.1 9.2 9.3 9.4 9.4 0 Asy 10.1 10.2 10.3 10.4	E's with exponentially decaying error terms Equations and algebraic decompositions	<ul> <li>95</li> <li>96</li> <li>99</li> <li>101</li> <li>104</li> <li>107</li> <li>108</li> <li>109</li> <li>111</li> </ul>
10	<ul> <li>9.1</li> <li>9.2</li> <li>9.3</li> <li>9.4</li> <li>9.4</li> <li>9.4</li> <li>9.1</li> <li>10.1</li> <li>10.2</li> <li>10.3</li> <li>10.4</li> <li>10.5</li> </ul>	E's with exponentially decaying error terms Equations and algebraic decompositions	<ul> <li>95</li> <li>96</li> <li>99</li> <li>101</li> <li>104</li> <li>107</li> <li>108</li> <li>109</li> <li>111</li> <li>114</li> </ul>
10	<ul> <li>9.1</li> <li>9.2</li> <li>9.3</li> <li>9.4</li> <li>9.4</li> <li>9.5</li> <li>10.1</li> <li>10.2</li> <li>10.3</li> <li>10.4</li> <li>10.5</li> <li>10.6</li> </ul>	E's with exponentially decaying error terms         Equations and algebraic decompositions         Rough estimate of solutions         Detailed asymptotics         Specifying data at infinity         Specifying data at infinity         mptotics, the silent setting         Introduction         Weak silence, balance and convergence         The oscillatory regime         Future asymptotics         Specifying the asymptotics	<ul> <li>95</li> <li>96</li> <li>99</li> <li>101</li> <li>104</li> <li>107</li> <li>108</li> <li>109</li> <li>111</li> <li>114</li> <li>118</li> </ul>
10	<ul> <li>9.1</li> <li>9.2</li> <li>9.3</li> <li>9.4</li> <li>9.4</li> <li>9.5</li> <li>10.1</li> <li>10.2</li> <li>10.3</li> <li>10.4</li> <li>10.5</li> <li>10.6</li> <li>0D2</li> </ul>	E's with exponentially decaying error terms         Equations and algebraic decompositions         Rough estimate of solutions         Detailed asymptotics         Detailed asymptotics         Specifying data at infinity         mptotics, the silent setting         Introduction         Weak silence, balance and convergence         The oscillatory regime         Future asymptotics         Specifying the asymptotics         Specifying the asymptotics	<ul> <li>95</li> <li>96</li> <li>99</li> <li>101</li> <li>104</li> <li>107</li> <li>108</li> <li>109</li> <li>111</li> <li>114</li> <li>118</li> <li>121</li> </ul>
10	<ul> <li>9.1</li> <li>9.2</li> <li>9.3</li> <li>9.4</li> <li>9.4</li> <li>9.4</li> <li>9.5</li> <li>10.4</li> <li>10.5</li> <li>10.6</li> <li>0D2</li> <li>11.1</li> </ul>	E's with exponentially decaying error terms         Equations and algebraic decompositions         Rough estimate of solutions         Detailed asymptotics         Detailed asymptotics         Specifying data at infinity         mptotics, the silent setting         Introduction         Weak silence, balance and convergence         The oscillatory regime         Future asymptotics         Specifying the asymptotics         Specifying the asymptotics         Introduction	<ul> <li>95</li> <li>96</li> <li>99</li> <li>101</li> <li>104</li> <li>107</li> <li>108</li> <li>109</li> <li>111</li> <li>114</li> <li>118</li> <li>121</li> <li>121</li> </ul>

1	1.3 Computing projections onto generalised eigenspaces	125
1	1.4 Mode by mode analysis	135
1	1.5 Specifying data at infinity 1	142
1	1.6 Algebraic properties of $A(\iota)$	l45
12 A	Asymptotics, transparent case 1	.49
1	2.1 Introduction $\ldots$	149
1	2.2 Weak transparency, balance and convergence	151
1	2.3 Deriving asymptotics	156
1	2.4 Specifying the asymptotics	169
III	Averaging over oscillations 1	75
13 I	ntroduction and technical preliminaries 1	77
1	3.1 Introduction $\ldots$ $\ldots$ $\ldots$ 1	177
1	3.2 On a class of matrix exponentials	179
1	3.3 Approximating a fundamental solution by a matrix exponential	184
1	3.4 Inhomogeneous equations	186
14 C	Dne period; terminology and preliminaries	.87
1	4.1 Assumptions $\ldots \ldots 1$	187
1	4.2 Change of variables	188
1	4.3 Choosing appropriate time intervals 1	190
1	4.4 Variation of the coefficients during one period	192
1	4.5 Computing the approximate fundamental solution	195
15 C	Dne period; detailed estimates 1	.97
1	5.1 Rough estimate of the evolution during one period	197
1	5.2 First detailed estimate of the evolution over one period $\ldots \ldots \ldots$	200
1	5.3 Iteration $\ldots$ $\ldots$ $\ldots$ $\ldots$ $\ldots$ $\ldots$ $2$	203
IV	Unbalanced equations 2	09
16 U	Inbalanced equations 2	11
1	6.1 Introduction $\ldots$	211
1	6.2 Unbalanced equations, the ODE setting	214
1	6.3 Unbalanced equations, the PDE setting	217
1	6.4 Subdividing the time interval $\ldots \ldots \ldots$	218
1	6.5 Basic properties of the iteration $\ldots$	219
1	6.6 The main estimate	223

iii

	16.8 Super exponential growth of solutions	231	
	16.9 Standard energy estimates		
	16.11Mode analysis	242	
v	Equations with a dominant noisy spatial direction	245	
17	Terminology and basic estimates	<b>247</b>	
	17.1 Introduction	247	
	17.2 Terminology and basic assumptions	248	
	17.3 Oscillation adaptedness and definition of the late time regime	250	
	17.4 Detailed asymptotics of the matrices	251	
	17.5 Reformulating the iteration	254	
	17.6 A rough mode estimate	255	
	17.7 A rough Sobolev estimate	258	
18	Asymptotics for one mode and late times	<b>261</b>	
	18.1 Asymptotics along a time sequence	261	
	18.2 Specifying data at infinity	267	
	18.3 Controlling the solution beyond the time sequence	270	
19	Asymptotics for all future times	277	
	19.1 Introduction	277	
	19.2 The case of scalar homogeneous wave equations	278	
	19.3 A uniqueness result	283	
	19.4 Detailed asymptotics in the noisy setting	285	
	19.5 Specifying data at infinity	295	

## VI Non-degenerate, diagonally dominated, convergent and balanced equations 301

20 Convergent and balanced equations			
	20.1 Introduction	303	
	20.2 Diagonally dominated, convergent and balanced equations	305	
	20.3 Simplifying the matrix coefficients	308	
	20.4 Time intervals, frequency eras	310	
	20.5 Estimating sums in a frequency era	313	
<b>21</b>	21 Non-degenerate equations 3		
	21.1 Introduction	317	
	21.2 Mode analysis in one frequency era	318	

	21.3	Refined estimates in transparent eras	324
	21.4	Analysis for one mode	327
	21.5	The case of unbounded frequency eras $\hdots \hdots \hdot$	335
	21.6	Optimality	337
22	$\mathbf{Esti}$	mating the Sobolev norms of solutions	343
	22.1	Introduction	343
	22.2	Mode estimates for the entire future	343
	22.3	Energy estimates	350
	22.4	Optimality	351
23	The	silent setting	355
	23.1	Introduction	355
	23.2	Future asymptotics, silent setting	356
	23.3	Specifying the asymptotics	363
<b>T</b> 77		A	
V		Appendices	367
<b>24</b>	Spe	ctral aspects of differential operators and matrices	369
	24.1	Dividing the solution into modes $\ldots \ldots \ldots$	369
	24.2	The Jordan normal form	372
25	Geo	metric preliminaries	375
	25.1	Metrics of interest, lapse and shift decomposition $\hdots \hdots \hddots \hdots \hdots\hdots\hdots \hdots \hdots \hdots \hdots \hdots$	375
	25.2	The components of the inverse $\hfill \ldots \ldots$	376
	25.3	The second fundamental form	378
	25.4	Geometric convergence	379
	25.5	Norms	380
	25.6	Quantitative bounds on the shift vector field $\hfill \ldots \ldots \ldots \ldots \ldots \ldots \ldots$	382
	25.7	From geometric to analytical conditions, part I $\ \ldots \ $	382
	25.8	Implications concerning the causal structure	388
26	Rela	ating the analytic and geometric conditions	391
	26.1	Bounds on the coefficients in the silent setting	391
	26.2	Conditions ensuring weak transparency $\hfill \ldots \hfill \ldots$	391
	26.3	A dominant noisy spatial direction $\hdots \ldots \ldots$	394
	26.4	The asymptotically diagonal setting	398

 $\mathbf{V}$ 

# Part I

# Introductory material, results

## Chapter 1

# Equations and questions

## **1.1** Introduction

The subject of these notes is the asymptotic behaviour of solutions to systems of linear wave equations of the following form:

$$\Box_g u + Xu + \zeta u = f. \tag{1.1}$$

Here g is a given Lorentz metric on a manifold M;  $\Box_q$  is the wave operator associated with the Lorentz metric g, defined by  $\Box_q u := \operatorname{div}(\operatorname{grad} u)$ ; X is a smooth  $m \times m$  matrix of vector fields on M whose coefficients with respect to a frame are allowed to be complex (and  $1 \le m \in \mathbb{Z}$ );  $\zeta$  is a smooth  $\mathbf{M}_m(\mathbb{C})$ -valued function on M (where  $\mathbf{M}_m(K)$  denotes the set of  $m \times m$  matrices with entries in the field K); and f is a smooth  $\mathbb{C}^m$ -valued function on M. We are mainly interested in the real (as opposed to the complex) setting, but for technical reasons, it turns out to be convenient to derive results under the above assumptions. The problem is to analyse the asymptotic behaviour of solutions  $u: M \to \mathbb{C}^m$  to (1.1). In particular, we are interested in the problem of deriving optimal estimates for energies naturally associated with solutions to (1.1), and in the problem of calculating the leading order asymptotics. We are only interested in globally hyperbolic Lorentz manifolds (M, q), and in that setting, there is a unique smooth solution to (1.1), given smooth initial data on a Cauchy hypersurface; a justification for this statement is provided, e.g., by [34, Theorem 12.19, p. 144] (though we do not need to appeal to this result in these notes). In fact, we here think of solutions to (1.1) as arising in this way. Our main motivation for studying (1.1)is that we are interested in the linearised Einstein equations. Note, however, that (1.1) is not the most general class of systems relevant in that context. For example, in the study of the Einstein-Euler equations, it is of interest to consider systems for which the symbol is different for the different components of u, a situation not covered by (1.1). Nevertheless, (1.1) represents quite a general class of equations, and, in order to obtain conclusions, we gradually need to impose more and more restrictions on the Lorentz manifolds of interest etc. One assumption we consistently make is that the Cauchy hypersurfaces of (M, g) are closed manifolds. In fact, M is here always of the form  $M := M \times I$ , where M is a closed manifold and  $I = (t_{-}, t_{+})$  is an open interval. In the case of Einstein's equations, this situation is of interest in cosmology (i.e., in the study of the universe as a whole), and the asymptotic regimes associated with  $t \to t_{\pm}$  should be thought of as representing a cosmological singularity (a big bang or a big crunch) or the expanding direction. The cosmological setting should be contrasted with the asymptotically Euclidean or asymptotically hyperbolic situation, which is of interest in the study of isolated systems (isolated galaxies, stars, black holes etc.). In the isolated systems setting, static and stationary solutions (such as the Minkowski and Kerr spacetimes) are the ones of greatest interest. In the cosmological setting, it is more natural to consider solutions with contracting directions (toward a big bang/big crunch) and/or expanding directions. In the following section, we take one more step towards restricting the class of equations defined by (1.1) to the one of interest here.

### 1.2 Equations

In these notes, we consider equations of the form

$$-g^{00}(t)u_{tt} - \sum_{j,l=1}^{d} g^{jl}(t)\partial_{j}\partial_{l}u - 2\sum_{l=1}^{d} g^{0l}(t)\partial_{l}\partial_{t}u - \sum_{r=1}^{R} a_{r}^{-2}(t)\Delta_{g_{r}}u + \alpha(t)u_{t} + \sum_{j=1}^{d} X^{j}(t)\partial_{j}u + \zeta(t)u = f.$$
(1.2)

Here  $1 \leq m \in \mathbb{Z}$ ,  $0 \leq d, R \in \mathbb{Z}$  and  $g^{00}, g^{jl}, g^{0l}, a_r \in C^{\infty}(I, \mathbb{R})$ , where I is an open interval,  $j, l = 1, \ldots, d$  and  $r = 1, \ldots, R$ . In case d = 0, the sum from 1 to d should be thought of as empty, and similarly for the sum from 1 to R. Moreover,  $-g^{00}$  and  $a_r, r = 1, \ldots, R$ , take their values in  $(0, \infty)$  and for each  $t \in I$ ,  $g^{jl}(t), j, l = 1, \ldots, d$ , are the components of a positive definite matrix (in (1.2) we also abuse notation in that we write  $a_r^{-2}(t)$  when we, strictly speaking, mean  $[a_r(t)]^{-2}$ ). In addition,  $(M_r, g_r), r = 1, \ldots, R$ , are closed Riemannian manifolds and  $\Delta_{g_r}$  is the Laplace-Beltrami operator on  $(M_r, g_r)$ . The functions u and f should be smooth functions from  $M := \overline{M} \times I$  to  $\mathbb{C}^m$ , where

$$\bar{M} := \mathbb{T}^d \times M_1 \times \dots \times M_R; \tag{1.3}$$

the differential operators  $\partial_j$  are the standard vector fields on  $\mathbb{T}^d$ ;  $\partial_t$  represents standard differentiation with respect to t (for functions on M); and  $\alpha, X^j, \zeta \in C^{\infty}[I, \mathbf{M}_m(\mathbb{C})]$ . Note that (1.2) is a special case of (1.1) when the Lorentz metric g on M is of the form

$$g = g_{00}dt \otimes dt + g_{0i}dt \otimes dx^i + g_{i0}dx^i \otimes dt + g_{ij}dx^i \otimes dx^j + \sum_{r=1}^R a_r^2 g_r,$$
(1.4)

where we use Einstein's summation convention (to sum over repeated upstairs and downstairs indices) and the coefficients only depend on t. The  $a_r$  appearing in this metric can be read off directly from (1.2) and the  $g_{\gamma\beta}(t), \gamma, \beta \in \{0, \ldots, d\}$ , are the components of the inverse of the matrix with components  $g^{\gamma\beta}(t), \gamma, \beta \in \{0, \ldots, d\}$ . That this inverse exists, and that (1.4) is a Lorentz metric on M, is a consequence of the assumptions; cf. Chapter 25, in particular the beginning of Section 25.1, for a detailed justification. Moreover,  $g_{00}(t) < 0$  and  $g_{ij}(t), i, j \in \{1, \ldots, d\}$ , are the components of a positive definite matrix for all  $t \in I$ . In Chapter 25, we also demonstrate that (M, g) is globally hyperbolic and that each hypersurface

$$\bar{M}_t := \bar{M} \times \{t\},\tag{1.5}$$

 $t \in I$ , is a Cauchy hypersurface; cf. Lemma 25.1.

**Remark 1.1.** There are two important operations that leave the class of equations of the form (1.2) invariant. First, multiplying an equation of the form (1.2) by a strictly positive function of t only yields an equation of the same type. Considering only the leading order derivatives in (1.2), this corresponds to a conformal rescaling of the metric (1.4) (by a conformal factor depending only on t). Secondly, we have the freedom of changing the time coordinate. In what follows, we make use of both of these operations in order to reduce the underlying geometry to a preferred form.

Clearly, the step from (1.1) to (1.2) involves significant restrictions. In particular, the fact that the coefficients of (1.2) only depend on t ensures that (1.2) is separable, an important simplification. Note also that the separability makes it very easy to study solutions under much more general regularity assumptions than smoothness (though we focus on smooth solutions in these notes). On the other hand, our main interest is in cosmological solutions, and in that setting, the standard starting point is the assumption of spatial homogeneity. From this point of view, assuming the coefficients to only depend on time is thus natural.

#### **1.2.1** Separable cosmological model manifolds

Due to our interest in equations of the form (1.2), we restrict our attention to the following class of Lorentz manifolds in what follows.

**Definition 1.2.** A separable cosmological model manifold is a Lorentz manifold (M, g) such that  $M = \overline{M} \times I$ , where  $I = (t_{-}, t_{+})$  is an open interval,  $0 \le d, R \in \mathbb{Z}, \overline{M}$  is given by (1.3) and

$$g = g_{00}dt \otimes dt + g_{0i}dt \otimes dx^i + g_{i0}dx^i \otimes dt + g_{ij}dx^i \otimes dx^j + \sum_{r=1}^R a_r^2 g_r,$$
(1.6)

where the  $g_{\gamma\beta}$  and the  $a_r$  only depend on t; the  $(M_r, g_r)$  are closed Riemannian manifolds;  $g_{\gamma\beta}(t) = g_{\beta\gamma}(t)$  for all t;  $g_{00}(t) < 0$  for all t; for all t, the  $g_{ij}(t)$  are the components of a positive definite matrix; and  $a_r(t) > 0$  for all t.

**Remark 1.3.** In the definition, it is always understood that  $\gamma, \beta \in \{0, \ldots, d\}$ ;  $i, j \in \{1, \ldots, d\}$ ; and  $r \in \{1, \ldots, R\}$ .

**Remark 1.4.** The reason for using the word separable is that the wave equation on (M, g) is separable. The reason for using the word cosmological is that  $\overline{M}$  is closed.

**Remark 1.5.** Sometimes it is convenient to write g as

$$g = -N^2 dt \otimes dt + g_{ij}(\chi^i dt + dx^i) \otimes (\chi^j dt + dx^j) + \sum_{r=1}^R a_r^2 g_r;$$
(1.7)

cf. Subsection 25.1.1 below for an explanation of how to calculate N and  $\chi^i$  in terms of the metric components. When using this notation, N > 0 is called the *lapse function* and  $\chi := \chi^i \partial_i$  the *shift vector field*. Note also that N and  $\chi^i$  only depend on t.

In the case that (M, g) is a separable cosmological model manifold, the induced metrics, second fundamental forms and volumes on the hypersurfaces  $\overline{M}_t$  (cf. (1.5)) are of importance. It is therefore convenient to introduce the following terminology.

**Definition 1.6.** Let (M, g) be a separable cosmological model manifold. Then the metric and second fundamental form of  $\overline{M}_t$  are denoted by  $\overline{g}_t = \overline{g}(t)$  and  $\overline{k}_t = \overline{k}(t)$  respectively (and they are interpreted as symmetric covariant 2-tensor fields on  $\overline{M}$ ). The trace  $\operatorname{tr}_{\overline{g}}\overline{k}$  is referred to as the mean curvature of  $\overline{M}_t$ . Moreover,  $V(t) := \operatorname{vol}_{\overline{g}(t)}(\overline{M})$  denotes the volume of  $\overline{M}$  with respect to  $\overline{g}(t)$ . Finally, U denotes the future directed unit timelike vector field which is normal to all the hypersurfaces  $\overline{M}_t$ .

**Remark 1.7.** The vector field U can be written

$$U := N^{-1} (\partial_t - \chi^i \partial_i). \tag{1.8}$$

Moreover,  $U(\ln V) = \text{tr}_{\bar{q}}\bar{k}$ ; cf. (25.17)–(25.19) below.

In the context of general relativity, the metric and second fundamental form induced on  $\overline{M}_t$  by g are of particular importance, since they constitute the geometric part of the initial data for Einstein's equations.

#### 1.2.2 Examples of separable cosmological model manifolds

In order to illustrate that there are separable cosmological model manifolds of interest, let us give a few examples.

**Minkowski space.** In the study of hyperbolic PDE's on  $\mathbb{R}^{d+1}$ , it is natural to begin with equations whose symbols equal that of the standard wave operator on d + 1-dimensional Minkowski space. One reason for this is that physical theories consistent with special relativity give rise to Lorentz invariant PDE's. From the point of view of general relativity, Minkowski space also plays a central role. This is partly due to the fact that it is a vacuum solution to Einstein's equations describing the geometry in the absence of gravitational fields (and constitutes the weak field limit of general relativity), but also because it is a *stable* solution to Einstein's equations, as has been demonstrated in, e.g., [9, 23, 24, 6]. The Minkowski metric on  $\mathbb{R}^{d+1}$  can, of course, also be considered to be a vacuum solution to Einstein's equations on  $\mathbb{T}^d \times \mathbb{R}$ . However, this solution is less natural. The main reason for this is that it is unstable, as we demonstrate below; cf. also the comments made at the beginning of Section 1.3.

**Bianchi type I.** Consider initial data to Einstein's equations on  $\mathbb{T}^d$ . Assume that they are invariant under the natural action of  $\mathbb{T}^d$  on itself. Here we refer to the corresponding maximal Cauchy developments as *Bianchi type I spacetimes*. In this situation, the spacetime metric can sometimes be written in the form

$$g = -dt \otimes dt + \sum_{i=1}^{d} a_i^2 dx^i \otimes dx^i$$
(1.9)

on  $M = \mathbb{T}^d \times I$ , where I is an open interval; the  $a_i$  are smooth, strictly positive functions on I; t is the coordinate on I; and  $dx^i$  are the standard one-forms on  $\mathbb{T}^d$  (considered as forms on M). We are here mainly interested in the situation that the mean curvature is non-zero (cf. the discussion at the beginning of Section 1.3). In the case of the vacuum Einstein equations, the  $a_i$  then take the form  $\alpha_i t^{p_i}$ , where  $0 < \alpha_i \in \mathbb{R}$  and  $p_i \in \mathbb{R}$  for  $i = 1, \ldots, d$ . Moreover, the sum of the  $p_i$  and the sum of the  $p_i^2$  both equal 1 (these conditions are referred to as the Kasner relations). The corresponding solutions are referred to as the Kasner solutions. Note that when  $a_i(t) = (t/t_0)^{p_i}$ , the initial data induced by (1.9) on the  $t = t_0$  hypersurface take the form

$$\bar{g}_{t_0} := \sum_{j=1}^d d\bar{x}^j \otimes d\bar{x}^j, \quad \bar{k}_{t_0} := \sum_{j=1}^d \frac{p_j}{t_0} d\bar{x}^j \otimes d\bar{x}^j,$$

where  $d\bar{x}^i$  are the standard one-forms on  $\mathbb{T}^d$ . Letting  $t_0 \to \infty$ , it is clear that these initial data converge to those of the Minkowski metric on  $\mathbb{T}^d \times \mathbb{R}$ . Moreover, this convergence is in any  $C^k$ topology. On the other hand, if none of the  $p_j$ 's equal 1, then (1.9) with  $a_i(t) = (t/t_0)^{p_i}$  is an inextendible solution to Einstein's vacuum equations which is past causally geodesically incomplete and exhibits curvature blow up as  $t \to 0+$ . In particular, it is thus clear that d + 1-dimensional Minkowski space, considered as a solution on  $\mathbb{T}^d \times \mathbb{R}$ , is unstable.

Concerning the Kasner solutions, let us remark that even though they can be expected to be unstable (both in the expanding and in the contracting direction), they have traditionally played an important role in general relativity. The reason for this is that they are the basic building blocks in the so-called BKL conjecture (due to Belinskiĭ, Khalatnikov and Lifschitz) concerning spacelike cosmological (big bang or big crunch) singularities. We shall here not discuss this conjecture further, but refer the reader interested in heuristic and numerical perspectives to [11, 13, 18] and [50] respectively, as well as references cited therein. Nevertheless, due to the central role this conjecture plays in the physics literature, it is of interest to understand the behaviour of solutions to the linearised Einstein equations around Kasner solutions.

Bianchi type I, accelerated expansion. Turning to cosmological solutions that are future stable, it is of interest to consider ones with accelerated expansion. The reason for this is partly due to the fact that the currently preferred models of the universe exhibit such expansion. However, it is also related to the cosmic no-hair conjecture, roughly stating that generic cosmological solutions to Einstein's equations with a positive cosmological constant generally asymptote to de Sitter space; cf., e.g., [4] for a more detailed discussion of this topic and a more precise formulation of the cosmic no-hair conjecture. Turning to the results, de Sitter space is stable; cf., e.g., [15]. Here de Sitter space with cosmological constant  $\Lambda > 0$  is the metric

$$-dt \otimes dt + \cosh^2(Ht)g_{\mathbb{S}^3}$$

on  $\mathbb{R} \times \mathbb{S}^3$ , where  $H = (\Lambda/3)^{1/2}$  and  $g_{\mathbb{S}^3_H}$  is the standard metric on the 3-sphere with radius such that  $\operatorname{Ric}[g_{\mathbb{S}^3_H}] = 2H^2 g_{\mathbb{S}^3_H}$ . Similarly,

$$g_{\rm dS} := -dt \otimes dt + \sum_{j=1}^d e^{2Ht} dx^j \otimes dx^j, \tag{1.10}$$

where  $0 < H \in \mathbb{R}$  and  $3 \le d \in \mathbb{Z}$ , is a future stable solution to Einstein's vacuum equations with a positive cosmological constant  $\Lambda = d(d-1)H^2/2$ . Note that (1.10) is a solution of Bianchi type I.

#### 1.2. EQUATIONS

Moreover, the universal covering spacetime of  $(\mathbb{T}^d \times \mathbb{R}, g_{dS})$  is a subset of de Sitter space which, to the future, exhibits the same asymptotics as de Sitter space itself. Moreover, future stability holds in the presence of a large class of different matter models; cf., e.g., [32, 47, 38, 45, 16, 36, 4, 26]. In accordance with the cosmic no-hair conjecture, spatially homogeneous but anisotropic solutions typically asymptote to de Sitter space; cf., e.g., [49, 21] and [36, Part VII]. In fact, there are even classes of solutions that are both highly anisotropic and highly spatially inhomogeneous that asymptote to the geometry described by (1.10); cf., e.g., [48, 4] for two results in the Einstein– Vlasov setting. Due to these results, it is clear that the metric (1.10) is of importance. It also describes the asymptotics of the currently preferred models of the universe.

The Einstein-non-linear scalar field equations with an exponential potential are considered in, e.g., [35, 17, 25]. The motivation for studying this setting is that it can be used to model dark energy, the purpose being to explain the accelerated expansion of the universe. In this setting, the model metrics again take the form (1.9), but with  $a_i(t) = \alpha_i t^p$ , where  $0 \neq \alpha_i \in \mathbb{R}$  and 1 . That these metrics (together with a scalar field corresponding to an appropriate exponential potential) are future stable solutions to the Einstein-non-linear scalar field equations is demonstrated in [35].

Bianchi type I, quiescent singularities. Concerning the spacetimes mentioned above, we focused on the issue of future stability. However, it is also of interest to consider big bang/big crunch type singularities. Due to the BKL conjecture, big bang singularities are typically thought to be 'oscillatory' (in a sense we do not here try to specify). However, there are some special matter models that are expected to neutralise the oscillations. Two examples of such matter models are stiff fluids and scalar fields. In this setting there is support for the BKL picture. The support comes in three main forms. First, there are results in the presence of symmetries (but in the absence of small data conditions); cf., e.g., [28, Section 7]. Second, there are constructions of large classes of solutions with the expected behaviour; cf., e.g., [5, 12]. Third, there are stability results around symmetric solutions; cf., e.g., [39, 40]. The relevant model metrics in this case are, again, of the form (1.9), where  $a_i(t) = \alpha_i t^{p_i}$ ,  $0 < \alpha_i \in \mathbb{R}$  and  $0 < p_i \in \mathbb{R}$ . Moreover, the sum of the  $p_i$  equal 1, but the sum of the  $p_i^2$  is strictly less than 1.

Higher dimensional analogues of de Sitter space. Let  $0 < H \in \mathbb{R}$  and  $(\Sigma, g_{\Sigma})$  be a closed Riemannian manifold of dimension  $n \geq 3$  such that  $\operatorname{Ric}[g_{\Sigma}] = (n-1)H^2g_{\Sigma}$ . Then the Lorentz metric

$$g_{\rm gdS} := -dt \otimes dt + \cosh^2(Ht)g_{\Sigma} \tag{1.11}$$

on  $M_{\text{gdS}} := \Sigma \times \mathbb{R}$  is a solution to Einstein's vacuum equations with a positive cosmological constant  $\Lambda = n(n-1)H^2/2$ . In the special case that n = 3 and  $\Sigma = \mathbb{S}^3$ , the Lorentz manifold  $(M_{\text{gdS}}, g_{\text{gdS}})$  is called *de Sitter space*.

The Nariai spacetimes. The Nariai spacetimes are interesting counterexamples to what is expected to be the generic behaviour of solutions to Einstein's equations with a positive cosmological constant (according to the cosmic no-hair conjecture). For  $0 < H \in \mathbb{R}$ , the Nariai spacetime solving Einstein's vacuum equations with a positive cosmological constant  $\Lambda = H^2$  is given by  $(M_N, g_N)$ , where  $M_N := \mathbb{S}^1 \times \mathbb{S}^2 \times \mathbb{R}$ ,

$$g_{\rm N} := -dt \otimes dt + \cosh^2(Ht) dx^2 + H^{-2} g_{\mathbb{S}^2}, \tag{1.12}$$

and  $g_{\mathbb{S}^2}$  is the standard metric on the unit 2-sphere.

The Milne model. In [3], Andersson and Moncrief demonstrate the future stability of the *Milne* model in the class of vacuum solutions to Einstein's equations. The Milne model is the spacetime  $(\Sigma \times (0, \infty), g_{\rm M})$ , where

$$g_{\rm M} := -dt \otimes dt + t^2 g_{\Sigma}$$

and  $(\Sigma, g_{\Sigma})$  is an appropriate closed hyperbolic manifold. This model plays a central role in a general conjecture concerning the future asymptotics of cosmological solutions to Einstein's vacuum equations due to Arthur Fischer and Vincent Moncrief on the one hand, and Michael Anderson on the other; cf., e.g., [1, 14]. Cosmological solutions with U(1)-symmetry. In [7, 8], Choquet-Bruhat and Moncrief study the future stability of

$$g_{\rm U} := -dt \otimes dt + d\theta \otimes d\theta + t^2 g_{\Sigma}$$

on  $\mathbb{S}^1 \times \Sigma \times (0, \infty)$ , where  $d\theta$  is the standard one-form field on  $\mathbb{S}^1$  and  $(\Sigma, g_{\Sigma})$  is an appropriate closed hyperbolic 2-manifold. In fact, they demonstrated future stability of this solution under U(1)-symmetric perturbations.

**Gowdy and**  $\mathbb{T}^2$ -symmetry. When taking the step from spatially homogeneous solutions to Einstein's equations to spatially inhomogeneous solutions, the setting which is the easiest to consider is that of polarised  $\mathbb{T}^3$ -Gowdy symmetric solutions to Einstein's vacuum equations. In that case, the essential part of the equations is given by one linear, scalar wave equation:

$$P_{tt} - e^{2t} P_{\theta\theta} = 0 \tag{1.13}$$

on  $\mathbb{S}^1 \times \mathbb{R}$ . Here  $t \to \infty$  corresponds to the expanding direction and  $t \to -\infty$  corresponds to the big bang singularity. Again, this is an equation of the form (1.2). Even though (1.13) is a simple equation, deriving all the asymptotic information needed in order to prove strong cosmic censorship in this setting does require an effort; cf. [10, 19]. Relaxing the symmetry assumptions by considering  $\mathbb{T}^3$ -Gowdy symmetric solutions, the equation (1.13) is replaced by a system of non-linear wave equations, but with the same symbol. In this case, analysing the asymptotics is more difficult; cf., e.g., [29, 31, 33]. Moreover, in the expanding direction, the behaviour of solutions can be radically different from that of spatially homogeneous solutions; in fact, spatially homogeneous solutions are unstable. This conclusion is strengthened by the analyses carried out in [22] (in the polarised  $\mathbb{T}^2$ -symmetric setting) and in [37] (in the  $\mathbb{T}^2$ -symmetric setting). Even though we are here interested in the linear setting, a better understanding of linear systems of equations with symbols similar to that of (1.13) can be expected to be useful in interpreting the results of [29, 22, 37], and possibly in extending them to other situations.

**Summary.** As is clear from the above, there is a large number of spacetimes such that studying linear systems of equations of the form (1.2) on these spacetimes is of interest. In several of the examples we have given, stability has already been demonstrated in the non-linear setting. It is therefore reasonable to ask if there is any use in considering linear equations on the corresponding backgrounds. The reason we do so here is that in several of the cases mentioned, the results are limited in that they do not currently apply to the full range of situations one would expect. In fact, in many of the situations discussed above where there are results, it would be desirable to relax the symmetry requirements and/or to extend the results to other matter models. When extending the results, a more detailed understanding of the linear setting can be expected to be valuable. This is particularly true in the case of Einstein's equations, since, even for a fixed solution to Einstein's equations coupled to a specific matter model, there are many different ways of formulating the equations and many different ways of breaking the gauge invariance. In order to be able to make an informed choice on how to break the gauge invariance, it is very valuable to have tools allowing a quick determination of how solutions to the corresponding linearised system behave.

### **1.3** Metrics with convergent asymptotics

In the context of cosmology, the two types of asymptotic regimes that are of greatest interest are expanding directions and big bang/big crunch type singularities. One indicator of expansion is that the volume V tends to infinity, and one indicator of a big bang/big crunch is that the volume tends to zero. It is also natural to assume that, at least asymptotically, the derivative of the volume in the direction normal to the hypersurfaces  $\overline{M}_t$  is non-zero; this would ensure a strict monotonicity of the volume. Due to Remark 1.7 this requirement is equivalent to the assumption that the mean curvature is either strictly positive or strictly negative asymptotically. In what follows, we impose these conditions. However, before stating them, let us make the following comments concerning static metrics. **Remark 1.8** (Static solutions). Due to the importance of static solutions in the asymptotically flat setting, it is perhaps worth commenting on the properties of such solutions in the cosmological setting. We already discussed Minkowski space with  $\mathbb{T}^d$ -spatial topology in Subsection 1.2.2. More generally, we would expect a static cosmological solution to have the following three properties: it should be a causally geodesically complete solution to Einstein's equations; it should be globally hyperbolic with closed spatial topology; and with respect to a suitable foliation by spacelike Cauchy hypersurfaces, say  $\Sigma_t$ , the volume of the leaves  $\Sigma_t$  should be constant. That the volume of the leaves is constant implies that the mean curvature of the leaves vanishes (assuming constant mean curvature). Perturbing the initial data induced on  $\Sigma_t$  should thus yield a sequence of initial data for Einstein's equations with strictly positive mean curvature and converging to those induced on  $\Sigma_t$ . Appealing to Hawking's singularity theorem (assuming the relevant matter fields appearing satisfy the energy conditions needed in order to appeal to Hawking's theorem) yields the conclusion that the maximal Cauchy developments of the members of this sequence are all causally geodesically incomplete; cf., e.g., [27, Theorem 55B, p. 432]. This means that the solution we started with is unstable. For this reason, static solutions are not natural in the context of cosmology. In fact, the asymptotics are typically characterised by expansion and/or contraction in the cosmological setting.

Due to the above observations, we here focus on separable cosmological model manifolds with one of the following properties.

**Definition 1.9.** Let (M, g) be a separable cosmological model manifold.

- If  $V(t) \to \infty$  as  $t \to t_+-$  and there is a  $t_0 \in I$  such that U(V) > 0 for  $t \ge t_0$ , then (M, g) is said to be *future expanding*.
- If  $V(t) \to \infty$  as  $t \to t_- +$  and there is a  $t_0 \in I$  such that U(V) < 0 for  $t \le t_0$ , then (M, g) is said to be *past expanding*.
- If  $V(t) \to 0$  as  $t \to t_+-$  and there is a  $t_0 \in I$  such that U(V) < 0 for  $t \ge t_0$ , then (M, g) is said to have big crunch asymptotics.
- If  $V(t) \to 0$  as  $t \to t_{-}+$  and there is a  $t_0 \in I$  such that U(V) > 0 for  $t \leq t_0$ , then (M, g) is said to have big bang asymptotics.

**Remark 1.10.** Let (M, g) be a separable cosmological model manifold. If it is future expanding, then reversing the time coordinate yields a separable cosmological model manifold which is past expanding (and vice versa). If (M, g) has big crunch asymptotics, then reversing the time coordinate yields a separable cosmological model manifold with big bang asymptotics (and vice versa). For that reason, we here restrict our attention to (M, g)'s which are future expanding or have big crunch asymptotics.

**Remark 1.11.** Due to Remark 1.7,  $\pm U(V) > 0$  is equivalent to  $\pm \text{tr}_{\bar{g}}\bar{k} > 0$ . The conditions involving U(V) could thus be reformulated in terms of  $\text{tr}_{\bar{q}}\bar{k}$ .

The assumptions introduced in Definition 1.9 are not very restrictive. In order to obtain a class of geometries such that we can draw conclusions concerning solutions to (1.2), we need to impose additional conditions. One object on which it is natural to impose conditions is the second fundamental form. In the formulation of the conditions, it is convenient to consider  $\bar{k}$  to be a map from  $T\bar{M}$  to itself. This can be achieved by raising one of the indices of  $\bar{k}$  by  $\bar{g}$ , and we denote the corresponding object by  $\bar{K}$ . Note that  $\bar{K}_i{}^j = \bar{k}_i{}^j$  with respect to local coordinates. In particular,  $\mathrm{tr}\bar{K} = \mathrm{tr}_{\bar{g}}\bar{k}$  (note that  $\mathrm{tr}\bar{K}$  does not depend on a choice of metric, since for each  $p \in \bar{M}$ ,  $(\bar{K})_p \in \mathrm{End}(T_p\bar{M})$ ). In many of the situations of interest,  $\mathrm{tr}\bar{K}$  either tends to zero or  $\pm\infty$ asymptotically. It is therefore natural to normalise  $\bar{K}$ . One way of doing so is to focus on  $\bar{K}/\mathrm{tr}\bar{K}$ . In what follows, we refer to this object as the *expansion normalised second fundamental form*. If the assumptions of Definition 1.9 are satisfied, the expansion normalised second fundamental form is at least asymptotically well defined. In these notes, we are mainly interested in the case that  $\bar{K}/\mathrm{tr}\bar{K}$  converges, a situation we refer to as the convergent setting. However, it also turns out to be important to impose conditions on the normal derivative of the mean curvature. Imposing such restrictions may seem to be less natural. However, in the context of Einstein's vacuum equations, the relevant conditions often follow from a combination of the assumption of convergence and the so-called Raychaudhuri equation. In the presence of matter, additional conditions may need to be imposed. However, the required conditions sometimes follow by combining the assumption of convergence, the Hamiltonian constraint, the Raychaudhuri equation and, possibly, energy conditions. Considering, e.g., solutions to Einstein's vacuum equations corresponding to left invariant initial data on 3-dimensional unimodular Lie groups yields solutions such that the following version of the Raychaudhuri equation holds:

$$U(\mathrm{tr}_{\bar{g}}\bar{k}) = -\bar{k}^{ij}\bar{k}_{ij};$$

cf., e.g., [34, (20.17), p. 218]. This means that if  $\bar{K}/\text{tr}\bar{K}$  is convergent (or bounded), then  $U[(\text{tr}_{\bar{g}}\bar{k})^{-1}]$  is convergent (or bounded). Nevertheless, in these notes, we do not assume (M,g) to be a solution to Einstein's equations. For this reason, we here impose conditions directly on  $U[(\text{tr}_{\bar{g}}\bar{k})^{-1}]$ .

Even though we are mainly interested in the convergent setting, it is sometimes sufficient to assume boundedness. For this reason, we introduce the following terminology.

**Definition 1.12.** Let (M,g) be a separable cosmological model manifold and let  $\varrho$  be a fixed Riemannian metric on  $\overline{M}$ . Then (M,g) is said to have *future bounded geometry* if there is a constant  $0 < C \in \mathbb{R}$  and a  $t_0 \in I$  such that  $|\mathrm{tr}_{\bar{g}}\bar{k}| > 0$  and

$$|\bar{k}|_{\bar{q}}/|\mathrm{tr}_{\bar{q}}\bar{k}| + |U[(\mathrm{tr}_{\bar{q}}\bar{k})^{-1}]| \le C$$

for all  $t \ge t_0$ . If, in addition, there is a 2-tensor field A on  $\overline{M}$  of mixed type, a constant  $a \in \mathbb{R}$  and constants  $0 < C, \eta \in \mathbb{R}$  such that

$$|\bar{K}/\mathrm{tr}\bar{K} - A|_{\varrho} + |U[(\mathrm{tr}_{\bar{g}}\bar{k})^{-1}] - a| \le C \exp[-\eta |\ln V(t)|]$$
(1.14)

for all  $t \ge t_0$ , then (M, g) is said to be *future convergent*.

**Remark 1.13.** It is of course possible to define the notion of future convergence without a rate. However, in these notes we are only interested in convergence in the case that there are quantitative bounds of the form (1.14).

At this stage, it is of interest to make use of the invariance described in Remark 1.1. By conformally rescaling the metric and then changing the time coordinate, we are in a position to reduce it to what we refer to as "canonical form".

**Lemma 1.14.** Let (M,g) be a separable cosmological model manifold, cf. Definition 1.2, and let  $\varrho$  be a fixed Riemannian metric on  $\overline{M}$ . Assume that (M,g) is future expanding and has future bounded geometry; cf. Definitions 1.9 and 1.12. Introduce the metric  $\hat{g} := (\operatorname{tr}_{\bar{g}}\bar{k})^2 g$  and the time coordinate

$$\tau(t) := \ln \frac{V(t)}{V(t_0)}.$$

Then the interval  $[t_0, t_+)$  in t-time corresponds to  $[0, \infty)$  in  $\tau$ -time. Moreover,  $\hat{g}$  is well defined on  $[0, \infty)$  in  $\tau$ -time and can be written

$$\hat{g} = -d\tau \otimes d\tau + \hat{g}_{ij}(\hat{\chi}^{i}d\tau + dx^{i}) \otimes (\hat{\chi}^{j}d\tau + dx^{j}) + \sum_{r=1}^{R} \hat{a}_{r}^{2}g_{r}.$$
(1.15)

If  $\check{g}$  and  $\hat{k}$  are the metric and second fundamental form induced on constant  $\tau$ -hypersurfaces by  $\hat{g}$ , then there is a constant  $0 < C \in \mathbb{R}$  such that  $|\hat{k}|_{\check{g}} \leq C$  for all  $\tau \geq 0$ .

Assuming, in addition to the above, that (M,g) is future convergent, there is a 2-tensor field  $\hat{A}$  on  $\overline{M}$  of mixed type such that

$$\hat{K} - \hat{A}|_{\varrho} \le C e^{-\eta\tau} \tag{1.16}$$

for all  $\tau \geq 0$ , where  $0 < C, \eta \in \mathbb{R}$  and  $\hat{K}$  is obtained from  $\hat{k}$  by raising one of the indices using  $\check{g}$ .

**Remark 1.15.** The proof of this statement is to be found in Section 25.4. Moreover, there is a similar result in the case of big crunch asymptotics.

**Remark 1.16.** The idea to conformally rescale the metric by multiplying it by the square of the mean curvature is not new; cf., e.g., [18] in which this idea is applied in a much more general (cosmological) setting.

**Remark 1.17.** The functions  $\hat{g}_{ij}$ ,  $\hat{a}_r$  and  $\hat{\chi}^i$  only depend on  $\tau$ . Moreover,  $\hat{g}(\partial_{\tau}, \partial_{\tau}) < 0$  by construction. This means that  $\hat{g}_{ij}\hat{\chi}^i\hat{\chi}^j < 1$ .

#### 1.3.1 Canonical separable cosmological model manifolds

With the above observations in mind, it is natural to introduce the following definition.

**Definition 1.18.** A canonical separable cosmological model manifold is a separable cosmological model manifold such that the interval  $I = (t_-, t_+)$  contains  $[0, \infty)$  and such that N appearing in (1.7) equals 1.

**Remark 1.19.** Due to the fact that N = 1, it can be concluded that  $g^{00} = -1$ ; cf. Chapter 25, in particular Lemma 25.2.

**Remark 1.20.** In order to obtain a canonical separable cosmological model manifold, it is sufficient to assume that (M, g) is future expanding; to conformally rescale g according to  $\hat{g} := (\text{tr}_{\bar{g}}\bar{k})^2 g$ ; and to change the time coordinate as in Lemma 1.14. There is a similar statement in the case of big crunch asymptotics.

**Remark 1.21.** We are mainly interested in the case that  $\bar{K}$  converges to a non-trivial mixed 2-tensor field on  $\bar{M}$ ; cf. Lemma 1.14. However, we prefer to impose specific conditions on  $\bar{K}$  in the statements of the results rather than including it in the definition.

It is useful to have some specific examples in mind before proceeding.

Solutions with exponential expansion. Consider the metrics  $g_{dS}$ ,  $g_{gdS}$  and  $g_N$ , introduced in (1.10), (1.11) and (1.12) respectively and defined on  $\mathbb{T}^d \times \mathbb{R}$ ,  $\Sigma \times \mathbb{R}$  and  $\mathbb{S}^1 \times \mathbb{S}^2 \times \mathbb{R}$  respectively. They are all canonical separable cosmological model manifolds. Moreover, in the case of  $(\mathbb{T}^d \times \mathbb{R}, g_{dS})$ ,  $\overline{K} = H \operatorname{Id}_{T\mathbb{T}^d}$ , where we think of  $\overline{K}$  as an element of  $\operatorname{End}(T\mathbb{T}^d)$ . In the case of  $(\Sigma \times \mathbb{R}, g_{dS})$ ,

$$\bar{K} = H \frac{\sinh(Ht)}{\cosh(Ht)} \mathrm{Id}_{T\Sigma} \to \pm H \mathrm{Id}_{T\Sigma}$$

as  $t \to \pm \infty$ . Finally, in the case of  $(\mathbb{S}^1 \times \mathbb{S}^2 \times \mathbb{R}, g_N)$ ,

$$\bar{K} = H \frac{\sinh(Ht)}{\cosh(Ht)} \Pi_{TS^1} \to \pm H \Pi_{TS^1}$$

as  $t \to \pm \infty$ , where  $\Pi_{T\mathbb{S}^1} \in \operatorname{End}[T(\mathbb{S}^1 \times \mathbb{S}^2)]$  is the projection onto  $T\mathbb{S}^1$ .

**Examples, Bianchi type I.** Let us consider a metric of the form (1.9), where  $a_i(t) = \alpha_i t^{p_i}$ ,  $0 < \alpha_i \in \mathbb{R}$  and  $p_i \in \mathbb{R}$ . Metrics of this form arise, e.g., in the vacuum setting, in the study of the Einstein-scalar field equations and in the study of the Einstein-non-linear scalar field equations with an appropriate exponential potential; cf. Subsection 1.2.2 above. In all of these examples, the sum of the  $p_i$ 's, say p, is strictly positive.

The expanding direction. Say now that we want to study the expanding direction of these solutions. According to Lemma 1.14, it is then natural to introduce a time coordinate  $\tau(t) = \ln(t/t_0)^p$ . Choosing  $t_0 = 1$ , this yields  $\tau = p \ln t$ . On the other hand,

$$\bar{k} = \sum_{i=1}^{d} p_i t^{-1} \alpha_i^2 t^{2p_i} dx^i \otimes dx^i$$

so that  $\operatorname{tr}_{\bar{q}}\bar{k} = pt^{-1}$ . Multiplying g (given by (1.9) with  $a_i$  as above) with  $(\operatorname{tr}_{\bar{q}}\bar{k})^2$  thus yields

$$(\operatorname{tr}_{\bar{g}}\bar{k})^{2}g = -p^{2}t^{-2}dt \otimes dt + \sum_{i=1}^{d}p^{2}\alpha_{i}^{2}t^{2(p_{i}-1)}dx^{i} \otimes dx^{i}$$
$$= -d\tau \otimes d\tau + \sum_{i=1}^{d}\beta_{i}^{2}e^{2\gamma_{i}\tau}dx^{i} \otimes dx^{i},$$
(1.17)

where  $\beta_i = p\alpha_i$  and  $\gamma_i = (p_i - 1)/p$ . Note that in the case of the Kasner solutions, p = 1 and the sum of the squares of the  $p_i$  equals one. There are thus two possibilities: either one  $p_i$  equals 1 and all the rest equal zero (the corresponding metrics are referred to as *flat Kasner solutions*, since they have a vanishing Riemann curvature tensor - in fact, they are quotients of a part of d+1-dimensional Minkowski space); or all the  $p_i < 1$ . In the first case, there is one  $\gamma_i = 0$  and all the rest equal -1. In the second case, all the  $\gamma_i < 0$ . In the second case, it is thus clear that there is contraction in all directions, even though the original solution was expanding. Denoting the metric in (1.17) by  $\hat{g}$ , the associated second fundamental form (of the hypersurfaces of constant  $\tau$ ) is given by

$$\hat{k} = \sum_{i=1}^{d} \gamma_i \beta_i^2 e^{2\gamma_i \tau} dx^i \otimes dx^i.$$

In particular, it is clear that  $\hat{K}$  is constant. Moreover, the eigenvalues are given by the  $\gamma_i$ 's.

The contracting direction. Turning to the contracting direction, it is natural to introduce the time coordinate  $\tau = -p \ln t$ . The computation (1.17) is still valid. The only difference in the contracting direction is that  $\gamma_i = (1 - p_i)/p$ . In the case of the Kasner solutions, the typical situation is thus that the metric given by the far right hand side of (1.17) exhibits exponential expansion. In the case of a non-vacuum solution to the Einstein-scalar field equations, the sum of the squares of the  $p_i$ 's is strictly less than 1. For that reason the  $\gamma_i$ 's are always strictly positive, and the metric exhibits exponential expansion. In both cases,  $\hat{K}$  is constant.

**The Milne model.** In this case, the volume is given by  $V(t) = t^d \operatorname{vol}_{g_{\Sigma}}(\Sigma)$ , where d is the dimension of  $\Sigma$ . Fixing  $t_0 = 1$ , it is therefore natural to introduce  $\tau(t) = d \cdot \ln t$ . On the other hand, the second fundamental form is given by  $\bar{k} = tg_{\Sigma}$ , so that  $\operatorname{tr}_{\bar{q}}\bar{k} = d \cdot t^{-1}$ . Thus

$$\hat{g}_{\mathrm{M}} := (\mathrm{tr}_{\bar{g}}\bar{k})^2 g_{\mathrm{M}} = -d^2 t^{-2} dt \otimes dt + d^2 g_{\Sigma} = -d\tau \otimes d\tau + d^2 g_{\Sigma}.$$

In particular, the associated second fundamental form is identically zero, even though the Milne model exhibits expansion.

U(1) symmetric solutions. In this case,  $V(t) = 2\pi t^2 \operatorname{vol}_{g_{\Sigma}}(\Sigma)$ . Letting  $t_0 = 1$  yields  $\tau(t) = 2 \ln t$ . Moreover,  $\operatorname{tr}_{\bar{g}} \bar{k} = 2t^{-1}$ , so that

$$\hat{g}_{\mathrm{U}} := -4t^{-2}dt \otimes dt + 4t^{-2}d\theta \otimes d\theta + 4g_{\Sigma} = -d\tau \otimes d\tau + 4e^{-\tau}d\theta \otimes d\theta + 4g_{\Sigma}.$$

**Model metrics.** Considering the above examples, it is clear that metrics of the following form appear naturally:

$$g = -dt \otimes dt + \sum_{r=1}^{R} \alpha_r^2 e^{2\beta_r t} g_r, \qquad (1.18)$$

where  $0 < \alpha_r \in \mathbb{R}$ ,  $\beta_r \in \mathbb{R}$  and  $(M_r, g_r)$  are closed Riemannian manifolds (here  $(M_r, g_r)$  could equal  $(\mathbb{S}^1, d\theta \otimes d\theta)$ , so that the *d*-torus part appearing in (1.6) should be thought of as being included in (1.18)). In fact, we are mainly interested in metrics that, asymptotically, behave as (1.18). However, in several of the results of these notes we impose much weaker conditions.

#### **1.3.2** Silent, transparent and noisy metrics

The behaviour of solutions to (1.2) is strongly dependent on the asymptotic behaviour of the corresponding metric (after a conformal rescaling and change of time coordinate of the type described in Lemma 1.14). In Chapters 4, 5 and 6 we introduce the notions of silent equations; transparent equations; and equations with a dominant noisy spatial direction. The purpose of this terminology is to allow us to distinguish between asymptotics of fundamentally different type. It would be premature to formulate the corresponding definitions here, but it is useful to define what it means for a model metric (1.18) to be silent, transparent or noisy. **Definition 1.22.** Consider a metric of the form (1.18). If all the  $\beta_r$  are strictly positive, the metric is said to be *silent*; if all the  $\beta_r \ge 0$  but there is one  $\beta_r = 0$ , then the metric is said to be *transparent*; and if there is one  $\beta_r$  which is strictly negative, then the metric is said to be *noisy*.

In order to motivate the terminology, we consider the three cases separately.

Silent metrics. Let g be a silent metric of the form (1.18) and let  $\gamma$  be a causal curve with respect to this metric (so that  $g(\dot{\gamma}, \dot{\gamma}) \leq 0$ ). Note that, in general relativity, light and material objects (or, more generally, information) are assumed to travel along causal curves, so that causal curves are often spoken of as "observers". By a reparametrisation, if necessary, we can assume  $\gamma$ to be of the form  $\gamma(t) = (\bar{\gamma}_1(t), \ldots, \bar{\gamma}_R(t), t)$ , where  $\bar{\gamma}_r$  takes its values in  $M_r$ . The assumption of causality is equivalent to

$$-1 + \sum_{r=1}^{R} \alpha_r^2 e^{2\beta_r t} |\dot{\bar{\gamma}}_r(t)|_{g_r}^2 \le 0.$$

In particular,  $|\dot{\gamma}_r(t)|_{g_r} \leq \alpha_r^{-1} e^{-\beta_r t}$ . This means that  $\bar{\gamma}_r(t)$  converges to a point  $p_r \in M_r$ . If  $\lambda$  is another causal curve, we obtain corresponding points  $q_r \in M_r$ . For two typical curves, there is one  $r \in \{1, \ldots, R\}$  such that  $p_r \neq q_r$ . Given two such curves, there is a  $t_1$  such that for  $t \geq t_1$ , it is not possible to send information from  $\gamma(t)$  to  $\lambda$  (in other words, there is no future directed causal curve  $\xi$ , starting at  $\gamma(t)$  and intersecting  $\lambda$  to the future). Due to this inability of observers to communicate at late times, the metric (1.18) is said to be silent.

**Transparent metrics.** Let g be a transparent metric of the form (1.18) and assume that  $\beta_r = 0$ . Given two distinct points  $p_r, q_r \in M_r$ , let  $\bar{\gamma}_r$  be a smooth simple closed curve in  $M_r$  with constant speed  $\alpha_r^{-1}$  (with respect to  $g_r$ ) such that the image of  $\bar{\gamma}_r$  contains  $p_r$  and  $q_r$ . Assume, moreover, that  $\bar{\gamma}_r(0) = p_r$ . We think of  $\bar{\gamma}_r$  as being defined on  $\mathbb{R}$  and define  $\gamma(t) = (\bar{\gamma}_1(t), \ldots, \bar{\gamma}_R(t), t)$ , where  $\bar{\gamma}_s$  is a constant curve in  $M_s$  for  $s \neq r$ . Then  $\gamma$  is a null curve and the  $M_r$  coordinate of  $\gamma$  equals  $p_r$  and  $q_r$  infinitely many times. In particular, the spatial coordinate of the curve does not converge. Moreover, if  $t_n \ge 0, 1 \le n \in \mathbb{Z}$ , are the non-negative times at which the  $M_r$ coordinate of  $\gamma$  equals  $p_r$ , then  $t_{n+1} - t_n$  is independent of n. We think of  $\gamma$  as representing a signal propagating in M and let  $p \in \overline{M}$  be defined by  $(p, 0) = \gamma(0)$ . Then the signal  $\gamma$  passes through  $(p, t_n)$  for each n. If  $\lambda$  is the observer sitting at p, i.e.  $\lambda(t) = (p, t)$ , then  $\lambda$  receives the signal (propagating along  $\gamma$ ) at times  $t_n$ , and the proper time elapsed between received signals (as measured by  $\lambda$ ) equals  $t_{n+1} - t_n$ . In particular, the proper time elapsed is independent of n. The word transparent is intended to capture the combination of the two properties that information can be sent (along null curves) between points on  $M_r$  and that the time elapsed between received signals remains constant. Note, however, that it is not necessarily possible to send information between an arbitrary pair of points on M. From this point of view, it would perhaps be more natural to say that the metric is partially transparent. However, we here use the terminology introduced above.

**Noisy metrics.** Let g be a noisy metric of the form (1.18) and assume that  $\beta_r < 0$ . Let  $p_r$ ,  $q_r$  and  $\bar{\gamma}_s$ ,  $s \in \{1, \ldots, R\}$ , be as in the transparent setting above, and define

$$\gamma(t) = [\bar{\gamma}_1(-\beta_r^{-1}e^{-\beta_r t}), \dots, \bar{\gamma}_R(-\beta_r^{-1}e^{-\beta_r t}), t].$$

Then  $\gamma$  is a null curve and the  $M_r$  coordinate of  $\gamma$  equals  $p_r$  and  $q_r$  infinitely many times. In particular, the spatial coordinate of the curve does not converge. Letting  $t_n$  be defined as in the transparent setting, there is a constant C such that

$$t_{n+1} - t_n \le C e^{\beta_r t_n}.$$

In particular, the time elapsed between received signals decays exponentially. In short, information can be sent (along null curves) between points on  $M_r$  and the time elapsed between received signals decays exponentially. The word noisy is intended to capture the combination of these two properties. Again, it could be argued that the terminology partially noisy would be more justified.

## 1.4 Questions

**Energy estimates.** Our main goal in these notes is to obtain optimal energy estimates for solutions to subclasses of systems of the form (1.2). In what follows, we are not interested in the most general case, but rather assume the underlying geometry to be a canonical separable cosmological model manifold, so that, in particular,  $g^{00} = -1$ . In other words, we assume that we have already appealed to Lemma 1.14, if necessary, and that the relevant metric and time coordinate are the conformally rescaled metric and logarithmic "volume" time coordinate. In this setting, the main energy of interest is

$$\mathfrak{E}_{\text{bas}}[u](t) = \frac{1}{2} \int_{\bar{M}} \left( |u_t(\cdot, t)|^2 + \sum_{i=1}^m [g^{kl}(t)\partial_k u_i(\cdot, t)\partial_l u_i^*(\cdot, t) + \sum_{r=1}^R a_r^{-2}(t)|\text{grad}_{g_r} u_i(\cdot, t)|_{g_r}^2 \right) + |u(\cdot, t)|^2 \mu_{\text{B}},$$
(1.19)

where  $\overline{M}$  is given by (1.3),

$$\mu_{\rm B} := dx \wedge \mu_{g_1} \wedge \dots \wedge \mu_{g_R}, \tag{1.20}$$

dx is the standard volume form on  $\mathbb{T}^d$  and  $\mu_{g_r}$  and  $\operatorname{grad}_{g_r}$  are the volume form and gradient associated with the Riemannian manifold  $(M_r, g_r)$  respectively; we assume  $M_r$  to be oriented. Moreover, if  $X_j$ , j = 1, 2, are two real vector fields on  $M_r$  and  $X = X_1 + iX_2$  is a complex vector field, then

$$|X|_{g_r}^2 := |X_1|_{g_r}^2 + |X_2|_{g_r}^2.$$

From now on we refer to  $\mathfrak{E}_{\text{bas}}[u]$  as the *basic energy*. There are also associated higher order energies. However, due to the separability of the equation, estimates for the higher order energies are direct consequences of the estimates for the basic energy. Even though  $\mathfrak{E}_{\text{bas}}[u]$  is the energy of greatest interest, it is sometimes convenient to consider

$$\mathfrak{E}_{\text{hom}}[u](t) = \frac{1}{2} \int_{\bar{M}} \left( |u_t(\cdot, t)|^2 + \sum_{i=1}^m [g^{kl}(t)\partial_k u_i(\cdot, t)\partial_l u_i^*(\cdot, t) + \sum_{r=1}^R a_r^{-2}(t)|\text{grad}_{g_r} u_i(\cdot, t)|_{g_r}^2 ] \right) \mu_{\text{B}}.$$
(1.21)

There are several ways of formulating questions related to optimal decay. However, we are here mainly interested in equations such that the basic energy grows or decays exponentially (in some degenerate situations, polynomial factors might also occur, but they represent subdominant behaviour). In such settings it is natural to introduce the following numbers.

**Definition 1.23.** Consider (1.2). Assume that the associated Lorentz manifold (M, g) is a canonical separable cosmological model manifold. Assume, moreover, that f = 0. Let S denote the set of smooth solutions to (1.2) (with f = 0). Assume the set

$$\mathcal{A}_{\rm cr} := \{ \eta \in \mathbb{R} \mid \forall u \in \mathcal{S} \; \exists C \in \mathbb{R} : \mathfrak{E}_{\rm bas}[u](t) \le Ce^{2\eta t} \; \forall t \ge 0 \}$$

to be non-empty and bounded from below. Then  $\eta_{cr}$  is defined to be the infimum of  $\mathcal{A}_{cr}$  and is called the *crude exponential bound* for the growth of the basic energy of solutions to (1.2) with f = 0. Assume the set

$$\mathcal{A}_{\mathrm{nl}} := \{ \eta \in \mathbb{R} \mid \exists C \in \mathbb{R} : \forall u \in \mathcal{S}, \ \mathfrak{E}_{\mathrm{bas}}[u](t) \le C \mathfrak{E}_{\mathrm{bas}}[u](0) e^{2\eta t} \ \forall t \ge 0 \}$$
(1.22)

to be non-empty and bounded from below. Then  $\eta_{nl}$  is defined to be the infimum of  $\mathcal{A}_{nl}$  and is called the *exponential bound* for the growth of the basic energy of solutions to (1.2) with f = 0.

**Remark 1.24.** Note that  $\eta_{cr} \leq \eta_{nl}$ . As we demonstrate below, however, this is the only restriction on the relation between these two numbers.

**Remark 1.25.** Clearly, other definitions are conceivable. For instance, we could replace  $\mathfrak{E}_{bas}[u](0)$  appearing in (1.22) by a higher order energy of u at t = 0, corresponding to, say, the  $H^{s_0+1} \times H^{s_0}$ norm of the initial data. For each  $s_0 \geq 0$ , we then obtain a constant, say  $\alpha_1(s_0)$ , describing the behaviour of the energy. Then  $\eta_{cr} \leq \alpha_1(s_0) \leq \eta_{nl}$ , and  $\alpha_1(s_0)$  is a decreasing function of  $s_0$ .
Nevertheless, we here focus on the two constants  $\eta_{cr}$  and  $\eta_{nl}$ .

Our main goal in these notes is to calculate  $\eta_{nl}$  for large classes of equations. However, we are also interested in obtaining asymptotic information.

**Asymptotics.** In order to give an idea of the type of asymptotic information we would like to derive, let us first give two very simple examples.

The contracting direction of polarised  $\mathbb{T}^3$ -Gowdy. Consider the equation (1.13) with the time reversed; i.e., the equation

$$P_{tt} - e^{-2t} P_{\theta\theta} = 0. (1.23)$$

Then the asymptotics in the direction  $t \to \infty$  correspond to the behaviour close to the big bang singularity of polarised  $\mathbb{T}^3$ -Gowdy symmetric solutions to Einstein's vacuum equations. Given a smooth solution P to (1.23), there are  $v, \psi \in C^{\infty}(\mathbb{S}^1, \mathbb{R})$  such that

$$P(\cdot, t) - vt - \psi, \quad P_t(\cdot, t) - v \tag{1.24}$$

both decay exponentially in any  $C^k$ -norm as  $t \to \infty$ . In fact, one can demonstrate that there is a homeomorphism (in the  $C^{\infty}$ -topology) between initial data to (1.23) and asymptotic data (given by v and  $\psi$  as described above). For a justification of these statements, the reader is referred to Example 4.20 below.

The expanding direction of polarised  $\mathbb{T}^3$ -Gowdy. Returning to (1.13), it is clear that the asymptotics as  $t \to \infty$  are quite different from those in the opposite time direction. However, changing time coordinate to  $\tau = e^t$ , (1.13) becomes

$$P_{\tau\tau} + \frac{1}{\tau} P_{\tau} - P_{\theta\theta} = 0 \tag{1.25}$$

on  $\mathbb{S}^1 \times (0, \infty)$ . Given a solution P to this equation, there are two constants, say a and b, and a solution  $\nu$  to the flat space wave equation with zero mean value over the circle, such that

$$P(\cdot,\tau) - a \ln \tau - b - \tau^{-1/2} \nu(\cdot,\tau), \quad P_{\tau}(\cdot,\tau) - a \tau^{-1} - \tau^{-1/2} \nu_{\tau}(\cdot,\tau)$$

decay to zero as  $\tau^{-3/2}$  with respect to any  $C^k$ -norm. In this case, a, b and  $\nu$  thus constitute the asymptotic data. In fact, in this case there is also a homeomorphism (in the  $C^{\infty}$ -topology) between initial data to (1.25) and the asymptotic data (given by a, b and  $\nu$  as described above, where  $\nu$  is topologised by its initial data). For a justification of these statements, the reader is referred to Example 6.16 below.

The equations (1.13) and (1.23) are two very simple examples of systems of the form (1.2). However, it is of interest to derive analogous asymptotic information, given a solution to (1.2). Moreover, we would like to demonstrate that the map from initial data to asymptotic data is continuous with respect to a suitable topology. Finally, it is of interest to prescribe asymptotic data, and to demonstrate that the map from asymptotic data to initial data is continuous with respect to, say, the  $C^{\infty}$ -topology. These are our goals as far as the asymptotics are concerned.

## **1.5** Outline of the introductory part

The outline of the remainder of Part I is the following.

On the notion of balance. Considering (1.2), it is clear that it is determined by the metric g; the coefficients  $\alpha$ ,  $\zeta$  and  $X^{j}$ ; and the right hand side f. In Section 1.3 above, we introduce conditions

on the metric g. Even though most of the results in these notes require stronger assumptions, these conditions give a good idea of the class of metrics we have in mind. However, at this stage it is unclear what to assume concerning  $\alpha$ ,  $\zeta$  and the  $X^i$ . In Chapter 2, we therefore describe some of the pathologies that can arise if the assumptions concerning these functions are too weak. It is natural to start by focusing on  $\alpha$  and  $\zeta$ . Consider (1.2) with  $g^{00} = -1$  and f = 0; cf. Remark 1.19. The spatially homogeneous solutions to this equation satisfy

$$u_{tt} + \alpha(t)u_t + \zeta(t)u = 0.$$
 (1.26)

If we allow, e.g.,  $\|\alpha\|$  or  $\|\zeta\|$  to grow exponentially, there are corresponding equations whose solutions grow super exponentially (i.e. as  $\exp[a \exp(bt)]$  for some a, b > 0); cf. Section 2.1. There are analogous results if  $\|\alpha\|$  or  $\|\zeta\|$  grow, say, polynomially. Since we here think of the Lorentz manifolds and time coordinates of interest as arising from an application of Lemma 1.14, we think of the volume of the time slices in the original manifold as equalling  $V(t) = V(t_0)e^{t-t_0}$ (assuming the original Lorentz manifold to be future expanding). From this perspective, the natural length scale is proportional to  $e^{t/D}$ , where D is the dimension of  $\overline{M}$ . For this reason, we here consider growth faster than exponential as pathological. As a consequence, we are mainly interested in equations of the form (1.2) with a metric satisfying the assumptions of Definition 1.18 and matrices  $\alpha$  and  $\zeta$  whose norms are bounded to the future. There are similar arguments in the case of Lorentz manifolds with big crunch asymptotics.

Concerning the  $X^i$ , it turns out to be natural to focus on  $Y^i(t) := X^i(t)/[g^{ii}(t)]^{1/2}$  (no summation). If there is a  $\beta > 0$  such that  $e^{-\beta t}Y^i(t)$  converges to a matrix with an eigenvalue with non-zero real part, then the results of Chapter 2 demonstrate that the energies of solutions to (1.2) typically grow super exponentially (additional technical assumptions are needed in order to obtain this conclusion, but we refer the reader to Chapter 2 for the details). In Section 2.2, we state such results for homogeneous equations of the type (1.2) with d = 1 and R = 0. The solutions with super exponential growth that we construct are smooth. However, if  $g^{11}$  decays exponentially, we are not able to prove super exponential growth of solutions corresponding to real analytic initial data. In order to determine where the construction breaks down, it is of interest to consider regularity classes which are intermediate between smoothness and real analyticity. In Section 2.3 we therefore determine to which Gevrey classes the initial data of the constructed solutions belong.

Even though the examples exhibiting super exponential growth are of interest, we would also like to know if they correspond to a generic set of initial data. We turn to this topic in Section 2.4, where we consider equations of the form (1.2) for general d and R, and with an f which is not necessarily zero. We state two genericity results. First, there is a subset of the set of smooth initial data which is generic in the sense of Baire, such that the corresponding solutions exhibit super exponential growth; cf. Proposition 2.18. Second, the set of initial data which do not lead to solutions with super exponential growth has infinite co-dimension; cf. Proposition 2.21. Finally, in Section 2.5 we explain the main ideas behind the proof of the results.

A geometric perspective. On the basis of the observations made in Chapter 2, we are in a position to formulate conditions that exclude pathological behaviour (such as super exponential growth). This is the subject of Chapter 3. It turns out that the conditions quite naturally take a geometric form. To begin with, we assume that the metric satisfies the conditions of Definition 1.18. Turning to the shift vector field, we already know that  $g_{ij}\chi^i\chi^j < 1$ . However, it is convenient to impose conditions of the type  $g_{ij}\chi^i\chi^j \leq a$  for some  $a \in [0, 1)$  and all  $t \geq 0$ ; this ensures that  $-g_{00}$  is uniformly bounded away from zero to the future. In specific results, we impose additional smallness conditions. Concerning the lower order terms, we assume  $\|\alpha\|$  and  $\|\zeta\|$  to be bounded to the future (though there are some exceptions to this; cf. Subsection 6.2.5 below). Turning to the  $X^i$ , it is natural to think of  $\mathcal{X} = X^i \partial_i$  as a matrix of vector fields on M. For a fixed t, it can also be thought of as a matrix of vector fields on  $\mathbb{T}^d$ . It is of interest to consider the norm of  $\mathcal{X}$  with respect to the metric induced on  $\mathbb{T}^d \times \{p\} \times \{t\}$  (considered as a metric on  $\mathbb{T}^d$ ). It turns out that some of the pathologies described in Chapter 2 can be eliminated by assuming the norms of the components of  $\mathcal{X}$  to be bounded to the future with respect to this metric.

In practice, we also need to impose conditions on the derivatives of the second fundamental form,  $\mathcal{X}$  etc. One natural way of doing so is to demand that the Lie derivatives of these objects with respect to U (up to a certain order) are bounded to the future; recall that U is defined by (1.8). We introduce the relevant terminology in Definitions 3.1, 3.4, 3.7 and 3.8. In Section 3.4, we then state conditions ensuring that the energy of solutions to the homogeneous equation cannot grow faster than exponentially; cf. (3.9). Finally, in Section 3.5, we formulate the results of Chapter 2 geometrically.

Silent equations. In Chapter 4, we turn to silent equations. The main condition concerning the metric in this case is that the second fundamental form has a strictly positive lower bound. This means that the metric is expanding exponentially in all directions; cf. Subsection 1.3.2 above. We also assume the shift vector field to be small. Given that these conditions are satisfied and that the norm of  $\mathcal{X}$  is future bounded, the coefficients of the spatial derivatives of u appearing in (1.2) decay exponentially. It is therefore natural to compare solutions to (1.2) with solutions to the equation obtained from (1.2) by dropping all the terms involving spatial derivatives. Additionally assuming  $\alpha$  and  $\zeta$  to converge to constant matrices  $\alpha_{\infty}$  and  $\zeta_{\infty}$  respectively, we obtain a "limit equation" which is a constant coefficient ODE for each spatial point. The first result of Chapter 4 states that the basic energy exhibits the same growth/decay as the basic energy for solutions to the limit equation; cf. (4.6). In particular, the growth/decay is determined by the coefficients of the limit equation (i.e. by  $\alpha_{\infty}$  and  $\zeta_{\infty}$ ) and the right hand side. Next, Proposition 4.11 yields the leading order asymptotics of solutions, as well as the conclusion that these asymptotics coincide with those of solutions to the limit equation. Moreover, according to Proposition 4.16, it is possible to specify the leading order asymptotics. Returning to the questions stated in Section 1.4, we can compute  $\eta_{\rm cr}$ , derive asymptotics and specify asymptotics. However, the results stated in Chapter 4 only yield a lower bound on  $\eta_{nl}$ .

**Transparent equations.** As a next step, it is of interest to consider equations such that the second fundamental form of the associated metric is non-negative, but converges to zero in some direction(s). This is the main assumption concerning the metric in the definition of transparent equations, the subject of Chapter 5. In this situation, the spatial variables are naturally divided into "silent" and "transparent" variables. This division is formalised in Subsection 5.1.2. Additionally, we assume the shift vector field to be small;  $\mathcal{X}$  to be future bounded; the components of  $\mathcal{X}$  corresponding to the transparent variables to converge; and  $\alpha$  and  $\zeta$  to converge. Under these assumptions, it is possible to associate a limit equation with (1.2). The limit equation is a constant coefficient system of linear wave equations in the transparent variables and the time variable for each silent variable. Similarly to the silent setting, the leading order asymptotics coincide with those of the limit equation; cf. Proposition 5.11. We are also in a position to specify the leading order asymptotics; cf. Proposition 5.14.

Equations with a dominant noisy spatial direction. Finally, it is of interest to consider metrics such that the second fundamental form is negative definite in some directions. In the end, we consider, roughly speaking, the following situation: we assume that there is one direction (corresponding to one S<sup>1</sup>-factor in  $\mathbb{T}^d$  or one of the  $M_r$ ) such that  $\bar{k}$  behaves as  $-\beta_n \bar{g}$  in that direction asymptotically, where  $\beta_n > 0$ ; and that  $\bar{k} \ge (-\beta_n + \eta_n)\bar{g}$  in the remaining directions, where  $\eta_n > 0$ . In this sense, there is one "dominant noisy spatial direction". In this setting, the spatial variables can be divided into "noisy" variables and "subdominant" variables. Adding appropriate convergence and boundedness conditions, we again obtain a limit equation. However, as opposed to the previous cases, it is not a constant coefficient equation. In order to obtain conclusions, we need to limit our attention to the modes of the equation that correspond to nontrivial dependence on the noisy variables. It turns out that, for a fixed such mode, the dominant behaviour is oscillatory, and the frequency of the oscillations grows exponentially. However, it turns out to be possible to average over the oscillations, and this yields a well-defined overall growth/decay of solutions. We state the corresponding energy estimate in Proposition 6.8.

In order to describe the asymptotics, we need to separate the oscillatory part from the overall growth/decay. To this end, it is of interest to first focus on solutions to the scalar homogeneous

equation obtained from (1.2) by setting all the lower order terms (i.e., the ones involving  $\alpha$ ,  $\zeta$  and  $X^i$ ) as well as the right hand side f to zero. The resulting equation is given by (6.6) below. For (6.6), we are able not only to derive detailed asymptotics, but in fact to demonstrate that there is a homeomorphism (with respect to the  $C^{\infty}$ -topology) between initial data and asymptotic data. The relevant result is stated as Proposition 6.10. Given these models for the oscillatory behaviour, we derive the leading order asymptotics in Proposition 6.18. The asymptotic form of the solution is given by a sum of terms of the form  $e^{At}u_w$ , where A is a constant matrix and  $u_w$  is a vector valued solution to (6.6). Similarly to silent and transparent equations, it is also possible to specify the leading order asymptotics. The relevant result is stated as Proposition 6.28. Combining the results of Chapter 6, we are in a position to derive and to specify asymptotics. Moreover, restricting to the parts of the solutions corresponding to modes with non-trivial dependence on the noisy variables, we can calculate an analogue of the  $\eta_{cr}$  introduced in Definition 1.23. However, we only obtain a lower bound on  $\eta_{nl}$ .

Energy estimates in the asymptotically diagonal setting. In spite of the detailed asymptotic information obtained in Chapters 4–6, the corresponding results only yield a lower bound on  $\eta_{\rm nl}$ . In order to be able to calculate  $\eta_{\rm nl}$ , we need to make stronger assumptions. In Chapter 7, we assume, roughly speaking, that the metric becomes diagonal asymptotically; i.e., that the S<sup>1</sup>-factors in  $\mathbb{T}^d$  become orthogonal asymptotically (the  $M_r$ -factors are already orthogonal to everything else by construction). Moreover, we assume that the second fundamental form is convergent and that the shift vector field converges to zero. As a consequence, the metric asymptotically takes the form (1.18). Turning to the lower order terms, we assume that  $\alpha$ ,  $\zeta$  and  $X^j_{\mathcal{I}}(g^{jj})^{1/2}$  converge to  $\alpha_{\infty}$ ,  $\zeta_{\infty}$  and  $X^j_{\infty}$  respectively. Finally, we assume the metric to be non-degenerate in the sense that the  $\beta_r$  appearing in (1.18) are distinct. Under these circumstances we are able to calculate  $\eta_{\rm nl}$ ; cf. Theorem 7.21. It is of interest to note that  $\eta_{\rm nl}$  depends on  $\alpha_{\infty}$ ,  $\zeta_{\infty}$ , the  $\beta_r$  and the  $X^j_{\infty}$ .

Improved asymptotic estimates in the silent setting. In Chapter 4, we derive asymptotics, given a solution. Moreover, for given leading order asymptotics, we construct a corresponding solution. Due to the results stated in Chapter 4, we know that the map from initial data to asymptotic data is continuous with respect to the  $C^{\infty}$ -topology. In fact, there is an  $0 \leq s_0 \in \mathbb{R}$  such that the  $H^s$ -norm of the asymptotic data is bounded by the  $H^{s+s_0}$  norm of the initial data. On the other hand, the results stated in Chapter 4 do not allow us to determine  $s_0$  in terms of the coefficients of the equation. Imposing conditions similar to those of Chapter 7, we obtain a specific value for  $s_0$ ; cf. Proposition 8.1. In Proposition 8.9, we state similar results concerning the map from asymptotic data to initial data. It is of interest to note that the map from initial data, where  $s_0 < 0$ .

## **1.6** Acknowledgements

The author would like to acknowledge the support of the Göran Gustafsson Foundation for Research in Natural Sciences and Medicine, the Swedish Research Council (Reference numbers 621-2012-3103 and 2017-03863) and the Erwin Schrödinger Institute. This article was in part written during a stay at the Erwin Schrödinger Institute in Vienna during the ESI Workshop on "Geometric Transport Equations in General Relativity" 2017. Finally, the author would like to thank Ellery Ames for comments on an earlier version of this article. The author is a professor at KTH Royal Institute of Technology and the Deputy Director of the Institut Mittag-Leffler.

## Chapter 2

## On the notion of balance

## 2.1 Introduction

Consider (1.2). The coefficients of this equation can naturally be divided into two groups: those of the highest order derivatives and those of the lower order derivatives. The coefficients of the highest order derivatives correspond to the Lorentz metric (1.6). Even though we impose additional conditions in particular situations, we have already formulated the basic assumptions concerning this metric in Section 1.3. In particular, we, from now on, assume the Lorentz manifold corresponding to (1.2) to satisfy Definition 1.18. Turning to  $\alpha$  and  $\zeta$ , we can already develop a feeling for what types of assumptions are reasonable by considering spatially homogeneous solutions. Let us, to this end, consider (1.26). Assume that there are constants  $C_{cu}, \gamma, \eta_{cu} > 0$ and matrices  $\alpha_{\infty}, \zeta_{\infty} \in \mathbf{M}_m(\mathbb{C})$  such that

$$\|\gamma^{-1}e^{-\gamma t}\alpha(t) - \alpha_{\infty}\| + \|\gamma^{-2}e^{-2\gamma t}\zeta(t) - \zeta_{\infty}\| \le C_{\mathrm{cu}}e^{-\eta_{\mathrm{cu}}t}$$

for all  $t \geq 0$ . Assume, moreover, that one of  $\alpha_{\infty}, \zeta_{\infty}$  is non-zero. Define

$$A_{\infty} := \begin{pmatrix} 0 & \mathrm{Id}_m \\ -\zeta_{\infty} & -\alpha_{\infty} \end{pmatrix}$$
(2.1)

and let  $\kappa_1$  be the largest real part of an eigenvalue of  $A_{\infty}$ . Then solutions to (1.26) behave as  $\exp[\kappa_1 e^{\gamma t}]$  as  $t \to \infty$ ; cf. Lemma 16.4 below for a precise statement. If  $\kappa_1 > 0$ , then solutions grow super exponentially. If  $\kappa_1 < 0$ , then solutions decay super exponentially (meaning that they grow super exponentially to the past). We consider both cases to be pathological. The reason for this is that we think of the Lorentz manifolds of interest as arising from an application of Lemma 1.14. Such Lorentz manifolds have the property that the volume of the hypersurfaces  $\overline{M}_t$  with respect to the original metric is proportional to  $e^{\pm t}$  (in the case of big crunch asymptotics, we obtain an exponentially decreasing volume). As a consequence, the natural length scale is  $e^{\pm t/D}$ , where  $D := \dim M$ . From this point of view, exponential growth/decay is natural, but super exponential growth/decay is not. In case the eigenvalues of  $A_{\infty}$  are purely imaginary, Lemma 16.4 does not yield any conclusions. On the other hand,  $\kappa_1 = 0$  is a sensitive borderline case, which we choose not to consider further (with one exception; cf. Subsection 6.2.5 below). For these reasons, we wish to avoid  $\alpha$  and  $\zeta$  growing exponentially. Another option would be to consider the case that  $\alpha$  and  $\zeta$  grow polynomially. However, this typically yields faster than exponential growth/decay. Due to these observations, we assume  $\|\alpha\|$  and  $\|\zeta\|$  to be future bounded in most of the results of these notes.

Imposing conditions on the matrix  $\mathcal{X}$  of vector fields. Except for the  $X^{j}$ 's, we have developed a rough feeling for what conditions to impose on the coefficients of (1.2). The main purpose of the present chapter is to illustrate some of the pathologies that arise if we do not impose appropriate bounds on  $\mathcal{X}$ . Before turning to a formal statement of the results, let us consider

$$u_{tt} - e^{-4t}u_{\theta\theta} + e^{-t}u_{\theta} + u_t + u = 0$$
(2.2)

on  $\mathbb{S}^1 \times \mathbb{R}$ . One naive way of approaching this equation is to say that the factors in front of the spatial derivatives are exponentially decaying. It should thus be possible to ignore the corresponding terms, and the behaviour should be dominated by the ODE part of the equation. Since the solutions to the corresponding ODE decay exponentially, solutions should in general decay exponentially. Another perspective would be to separate the equation (2.2) and to consider the Fourier coefficients. We discuss Fourier decompositions in Section 24.1 below (and we use the notation introduced in Section 24.1 without further comment in what follows). However, in the particular case of (2.2), we obtain

$$\ddot{z} + \dot{z} + (n^2 e^{-4t} + ine^{-t} + 1)z = 0, \tag{2.3}$$

where z denotes the n'th Fourier coefficient of u. We thus expect z to decay exponentially (just as in the case of the equation obtained by dropping the spatial derivatives). For each fixed Fourier coefficient, this is indeed what happens; Lemma 9.14 below implies that z decays as  $e^{-t/2}$ . On the other hand, the constant C appearing in the estimate  $|z(n,t)| \leq Ce^{-t/2}$  depends on n, indicating that there might be a problem when considering, e.g., smooth solutions to (2.2). Yet another perspective is to study energies. In analogy with the basic energy introduced in (1.19), consider

$$E_{\mathbf{u}}(t) := \frac{1}{2} \int_{\mathbb{S}^1} [|u_t(\theta, t)|^2 + e^{-4t} |u_\theta(\theta, t)|^2 + |u(\theta, t)|^2] d\theta.$$

A crude energy estimate yields the conclusion that  $\dot{E}_{u}(t) \leq e^{t}E_{u}(t)$ , indicating that the growth might be as bad as  $\exp(e^{t})$ ; i.e., super exponential. The different perspectives clearly yield very different conclusions. It is therefore of interest to ask: do solutions decay exponentially or grow super exponentially? Below we demonstrate that for generic smooth solutions to (2.2), the crude energy estimate  $E_{u}(t) \leq C \exp(e^{t})$  is essentially optimal; cf. Example 2.9 and Section 2.4 below. In this particular case, the argument that the spatial derivatives in (2.2) can be dropped since they are multiplied by exponentially decaying factors is not justified. Later on, we introduce the notion of balance, and it will become clear that the super exponential growth of solutions to (2.2) is related to the fact that (2.2) is not balanced.

In the present chapter, the conditions appearing in the results are largely non-geometric in nature. The reason for this is that we, in the next chapter, use the results obtained here to motivate the importance of a geometric perspective. However, in Section 3.5 below, we provide geometric reformulations of the results of the present chapter.

## 2.2 Unbalanced equations, first examples of super exponential growth

To begin with, we state a result concerning equations of the form (1.2) with d = 1, R = 0 and f = 0. In order to be able to state the assumptions in a concise way, it is convenient to introduce the notation

$$\mathfrak{l}(t) := \ln([g^{11}(t)]^{1/2}), \quad Y^1(t) := \frac{X^1(t)}{[g^{11}(t)]^{1/2}}, \quad \varsigma(t) := \frac{g^{01}(t)}{[g^{11}(t)]^{1/2}}.$$
(2.4)

It is possible to take a geometric perspective on these objects (and we encourage the reader to do so). However, in the present chapter, we take an analytic perspective, and only in the next chapter do we interpret the conclusions geometrically. Given the notation (2.4), the main assumptions are the following.

Assumption 2.1. Consider (1.2). Assume the associated metric to be such that (M, g) is a canonical separable cosmological model manifold. Assume, moreover, that d = 1 and that R = 0. Let  $\mathfrak{l}, Y^1$  and  $\varsigma$  be defined by (2.4). Assume that there are real numbers  $\beta_{\mathrm{sh}} < 0$ ;  $\mathfrak{b}, C_{\mathrm{sh}}, C_X, C_{\mathrm{ode}} > 0$ ;  $\beta_X \ge 0$ ; and  $\beta_{\mathrm{ode}}$  such that  $\beta_X > \beta_{\mathrm{ode}}$ ,

$$|\dot{\mathfrak{l}}(t)| + |\ddot{\mathfrak{l}}(t)| \le \mathfrak{b} \tag{2.5}$$

for all  $t \geq 0$ , and

$$|\varsigma(t)| + |\dot{\varsigma}(t)| \le C_{\rm sh} e^{\beta_{\rm sh} t},\tag{2.6}$$

$$||Y^{1}(t)|| + ||\dot{Y}^{1}(t)|| \le C_{X} e^{\beta_{X} t}, \qquad (2.7)$$

$$\|\dot{\alpha}(t)\| + \|\alpha(t)\| + \|\dot{\zeta}(t)\|^{1/2} + \|\zeta(t)\|^{1/2} \le C_{\text{ode}} e^{\beta_{\text{ode}}t}$$
(2.8)

for all  $t \geq 0$ .

**Remark 2.2.** The assumption concerning  $\mathfrak{l}$  guarantees that  $g^{11}$  cannot grow (or decay) faster than exponentially. Combining this observation with (2.6)–(2.8) yields the conclusion that none of the coefficients of the equation grows faster than exponentially. On the other hand, the assumptions still allow both exponential growth and exponential decay of the coefficients.

Assumption 2.1 only involves bounds, and as a consequence there are classes of equations, satisfying the assumptions, whose solutions do not exhibit super exponential growth. In order to proceed, we therefore make the following additional assumption.

Assumption 2.3. Consider (1.2). Assume the associated metric to be such that (M, g) is a canonical separable cosmological model manifold. Assume, moreover, that d = 1 and that R = 0. Let  $Y^1$  be defined by (2.4) and Assumption 2.1 be fulfilled. Assume that there is a  $Y^1_{\infty} \in \mathbf{M}_m(\mathbb{C})$  and constants  $\eta_X > 0$  and  $K_X > 0$  such that

$$\|e^{-\beta_X t} Y^1(t) - Y^1_{\infty}\| \le K_X e^{-\eta_X t}$$
(2.9)

for all  $t \ge 0$ , where  $\beta_X \ge 0$  is the constant appearing in the statement of Assumption 2.1. Finally, assume  $Y_{\infty}^1$  to have an eigenvalue with a non-zero real part.

**Remark 2.4.** If  $\beta_X > 0$ , this assumption ensures that  $Y^1$  grows exponentially. On the other hand, it does not imply that  $X^1$  grows. In fact,  $X^1$  could decay exponentially.

Given that these assumptions hold, we obtain the following result.

**Proposition 2.5.** Consider (1.2). Assume the associated metric to be such that (M,g) is a canonical separable cosmological model manifold. Assume, moreover, that d = 1; that R = 0; and that f = 0. Given that Assumptions 2.1 and 2.3 hold with  $\beta_X > 0$ , let  $\kappa$  be the largest absolute value of a real part of an eigenvalue of  $Y_{\infty}^1/2$  (by assumption,  $\kappa > 0$ ). Let  $\epsilon > 0$ . Then there is a sequence of smooth  $\mathbb{C}^m$ -valued solutions  $v_l$  to (1.2),  $1 \leq l \in \mathbb{Z}$ , and for each  $l \geq 1$ , there is a time sequence  $t_{l,k} \to \infty$  (as  $k \to \infty$ ) such that for each  $s \in \mathbb{R}$ ,

$$\lim_{l \to \infty} \left( \|v_l(\cdot, 0)\|_{(s+1)} + \|\partial_t v_l(\cdot, 0)\|_{(s)} \right) = 0.$$
(2.10)

Moreover,

$$\mathfrak{E}_{\hom}[v_l](t_{l,k}) \ge \exp\left[2\beta_X^{-1}\kappa(1-\epsilon)e^{\beta_X t_{l,k}}\right],\tag{2.11}$$

where  $\mathfrak{E}_{hom}[u]$  is defined by (1.21); i.e., in the present setting,

$$\mathfrak{E}_{\text{hom}}[u](t) := \frac{1}{2} \int_{\mathbb{S}^1} [|u_t(x,t)|^2 + g^{11}(t)|u_x(x,t)|^2] dx.$$
(2.12)

**Remarks 2.6.** Here  $\|\cdot\|_{(s)}$  denotes the norm introduced in (24.10). The combination of the estimates (2.10) and (2.11) demonstrates that the solution u = 0 to (1.2) is future unstable.

Remark 2.7. The statement is a consequence of Lemma 16.27; cf. Remark 16.29.

**Remark 2.8.** It is of interest to compare the lower bound given by (2.11) with the upper bound obtained by standard energy estimates. Due to Lemma 16.42, given a solution u to an equation satisfying the assumptions of the proposition, there is an upper bound on the associated energy of the form  $C_{\epsilon} \exp[\beta_X^{-1} || Y_{\infty}^1 || (1 + \epsilon) e^{\beta_X t}]$ . When  $|| Y_{\infty}^1 || = 2\kappa$  (e.g., when  $Y_{\infty}^1$  is Hermitian), the estimate (2.11) is thus essentially optimal.

Let us apply this result to (2.2).

**Example 2.9.** Consider (2.2). This equation is such that d = 1, R = 0 and f = 0. Moreover,  $g^{11}(t) = e^{-4t}$ ,  $g^{01}(t) = 0$ ,  $\alpha(t) = 1$ ,  $\zeta(t) = 1$  and  $X^1(t) = e^{-t}$ . In particular, the relations  $\mathfrak{l}(t) = -2t$ ,  $Y^1(t) = e^t$  and  $\zeta(t) = 0$  thus hold. As a consequence, Assumption 2.1 holds with  $\beta_{\mathrm{sh}} = -1$ ,  $\mathfrak{b} = 2$ ,  $\eta_{\mathrm{ode}} = 1$ ,  $C_{\mathrm{sh}} = 1$ ,  $C_X = 2$ ,  $C_{\mathrm{ode}} = 2$ ,  $\beta_X = 1$  and  $\beta_{\mathrm{ode}} = 0$ . Moreover, Assumption 2.3 holds with  $Y_1^{\circ} = 1$ ,  $\eta_X = 1$  and  $K_X = 1$ ; clearly  $Y_{\infty}^1/2$  has an eigenvalue with non-zero real part and  $\kappa = 1/2$ . Thus the conclusions of Proposition 2.5 hold, so that, given  $\epsilon > 0$ , there is a sequence of smooth solutions  $v_l$  to (2.2) and time sequences  $\{t_{l,k}\}$  such that  $t_{l,k} \to \infty$  as  $k \to \infty$  and (2.10) and (2.11) hold. In our setting, the estimate (2.11) reads

$$\mathfrak{E}_{\hom}[v_l](t_{l,k}) \ge \exp\left[(1-\epsilon)e^{t_{l,k}}\right].$$

This indicates that the crude energy estimate mentioned in connection with (2.2) is essentially optimal.

The equation in Example 2.9 is rather simple in that the coefficients of (2.2) are either constants or constants times exponential factors. However, similar conclusions hold for, e.g.,

$$u_{tt} - e^{-4t}u_{\theta\theta} + e^{-t}u_{\theta} + \sin(e^{t/2})u_t + e^{t/2}\cos(e^{t/2})u = 0.$$
(2.13)

Here one might naively expect the factors  $\sin(e^{t/2})$  and  $e^{t/2}\cos(e^{t/2})$  to influence the behaviour (due to their growth and the growth of their derivatives). However, as far as the argument is concerned, the corresponding terms in (2.13) are effectively error terms. It is also of interest to consider, e.g.,

$$u_{tt} - e^{-4t}u_{\theta\theta} + e^{-t}u_{\theta} + e^{t/2}u_t + e^t u = 0.$$
(2.14)

In the case of (2.14), initial data with finite frequency content yield solutions with super exponential decay (in fact, such solutions roughly behave as  $\exp(-e^{t/2})$ ; cf. Section 16.2). However, generic smooth solutions exhibit super exponential growth; cf. Section 2.4 below.

Recalling the discussion of the Fourier coefficients of solutions to (2.2), it is clear that solutions to (2.2) with finite frequency content decay exponentially. In particular, there is thus a dense set of initial data such that the corresponding solutions exhibit exponential decay. It is of interest to isolate where the transition between exponential decay and super exponential growth takes place. To this end, it is useful to introduce function spaces interpolating between initial data with finite frequency content and smooth initial data, and to analyse where super exponential growth sets in. In the end, we are not able to demonstrate super exponential growth for solutions with real analytic initial data (at least not for equations such that the coefficients of the spatial derivatives decay exponentially), so that it is sufficient to consider a class of function spaces interpolating between real analyticity and smoothness. This leads us to study Gevrey classes.

## 2.3 Gevrey classes

Let us begin by defining the Gevrey classes in the context of interest here.

**Definition 2.10.** A real valued function f defined on  $\mathbb{T}^d$  is said to be a *Gevrey function of order*  $0 < s \in \mathbb{R}$  if  $f \in C^{\infty}(\mathbb{T}^d, \mathbb{R})$  and there are constants R > 0 and C > 0 such that

$$\sup_{x \in \mathbb{T}^d} |\partial^{\alpha} f(x)| \le RC^{|\alpha|} (\alpha!)^s \tag{2.15}$$

for all *d*-multiindices  $\alpha$ . The class of Gevrey functions of order  $0 < s \in \mathbb{R}$  on  $\mathbb{T}^d$  is denoted  $G^s(\mathbb{T}^d)$ .

**Remarks 2.11.** Here we think of functions defined on  $\mathbb{T}^d$  as being defined on  $\mathbb{R}^d$  and being  $2\pi$ periodic in all of their variables. The definition can be generalised to functions with values in  $\mathbb{R}^m$ or  $\mathbb{C}^m$ , and we use the notation  $G^s(\mathbb{T}^d, \mathbb{R}^m)$  and  $G^s(\mathbb{T}^d, \mathbb{C}^m)$ . Moreover, if  $f \in G^1(\mathbb{T}^d)$ , then f is real analytic. Finally, if  $f \in G^s(\mathbb{T}^d)$ , where 0 < s < 1, then f is entire.

**Remark 2.12.** There is an alternate characterisation of the Gevrey classes; cf. Lemma 16.48 below. In fact, let  $f \in C^{\infty}(\mathbb{T}^d, \mathbb{C})$ ,  $0 < s \in \mathbb{R}$  and  $a_n, n \in \mathbb{Z}^d$ , be the complex Fourier coefficients of f. Then  $f \in G^s(\mathbb{T}^d, \mathbb{C})$  if and only if there are constants  $C_i > 0$ , i = 1, 2, such that

$$|a_n| \le C_1 \exp\left(-C_2 |n|^{1/s}\right)$$
 (2.16)

for all  $n \in \mathbb{Z}^d$ . There is an analogous characterisation in case f takes its values in  $\mathbb{R}^m$  or  $\mathbb{C}^m$ .

When considering solutions to (1.2) with R = 0, it is of interest to verify that Gevrey class regularity is preserved. In that context, the following definition is useful.

**Definition 2.13.** Let  $I \subseteq \mathbb{R}$  be an open interval,  $f \in C^{\infty}(\mathbb{T}^d \times I, \mathbb{C}^m)$  and  $0 < s \in \mathbb{R}$ . Then f is said to belong to  $\mathfrak{G}^s_{\text{loc},u}(I, \mathbb{T}^d, \mathbb{C}^m)$  if, for every compact interval  $J \subset I$ , the following holds: there are constants  $C_1 > 0$  and  $C_2 > 0$  (depending on f and J) such that

$$|\hat{f}(n,t)| \le C_1 \exp\left(-C_2 |n|^{1/s}\right)$$

for all  $n \in \mathbb{Z}^d$  and  $t \in J$ , where  $\hat{f}(n, t)$  is the *n*'th (complex) Fourier coefficient of  $f(\cdot, t)$ .

Consider (1.2). Assume the associated metric to be such that (M, g) is a canonical separable cosmological model manifold. Assume, moreover, that R = 0. If the coefficients are smooth,  $f \in \mathfrak{G}^s_{\mathrm{loc},\mathrm{u}}(\mathbb{R}, \mathbb{T}^d, \mathbb{C}^m)$ , and the initial data at  $t = t_0$  belong to  $G^s(\mathbb{T}^d, \mathbb{C}^m)$ , then the corresponding solution to (1.2) is smooth and such that u and  $u_t$  have the same regularity as f; cf. Lemma 16.51.

Given the above terminology, the statement of Proposition 2.5 can be improved to the following result.

**Proposition 2.14.** Assume that the conditions of Proposition 2.5 are fulfilled. Assume, moreover, that there are constants  $\beta_1 \in \mathbb{R}$  and  $0 < c_{m,1} \in \mathbb{R}$  such that

$$c_{\mathrm{m},1}e^{2\beta_1 t} \le g^{11}(t)$$

for all  $t \ge 0$ . If  $\beta_X - \beta_1 > 0$ , then the sequence of functions  $v_l$  constructed in Proposition 2.5 can be assumed to be such that  $v_l, \partial_t v_l \in \mathfrak{G}^{s_X}_{loc.u}(I, \mathbb{S}^1, \mathbb{C}^m)$  for all l, where

$$s_X := \frac{\beta_X - \beta_1}{\beta_X}.$$
(2.17)

In case  $\beta_X - \beta_1 \leq 0$ , one frequency can be chosen, say  $n_a \in \mathbb{Z}$ , such that the sequence of functions  $v_l$  constructed in Proposition 2.5 has the property that if  $z_l(n,t)$  denotes the n'th Fourier coefficient of  $v_l(\cdot,t)$ , then  $z_l(n,t) = 0$  for all t unless  $n = n_a$ .

**Remark 2.15.** The statement is an immediate consequence of Proposition 16.52.

**Remark 2.16.** Returning to Example 2.9, it is clear that  $\beta_1 = -2$  and  $\beta_X = 1$ . Thus  $s_X = 3$ .

Remark 2.17. Consider

$$u_{tt} - e^{2\beta_1 t} u_{\theta\theta} + e^{(\beta_1 + \beta_X)t} u_{\theta} = 0.$$

If we want the coefficients of the spatial derivatives to decay exponentially, we have to have  $\beta_1 < 0$ and  $\beta_1 + \beta_X < 0$ . If we also want to have super exponential growth, we have to have  $\beta_X > 0$ . This means that  $0 < \beta_X < -\beta_1$ , so that  $s_X > 2$ . In this setting, there is thus a limit on how close we can get to the case of real analytic initial data. If, on the other hand,  $0 < \beta_X \leq \beta_1$ , then there are initial data with finite frequency content such that the corresponding solutions exhibit super exponential growth.

It would be of interest to determine if the value of  $s_X$  appearing in (2.17) has any real significance or is just an articlast of our argument. However, we do not pursue this topic in these notes.

## 2.4 Unbalanced equations, generic super exponential growth

In the previous sections, we demonstrate that there are solutions to equations of the form (1.2) that exhibit super exponential growth. However, it is not clear if these solutions are very special, or if they represent the typical behaviour. When addressing this question, it is of interest to return to general equations of the form (1.2). Let us first note that initial data ensuring super exponential growth are generic in the Baire sense.

**Proposition 2.18.** Consider (1.2). Assume the associated metric to be such that (M, g) is a canonical separable cosmological model manifold. Assume, moreover, that there is a  $j \in \{1, ..., d\}$  such that

$$u_{tt} - g^{jj}(t)\partial_j^2 u - 2g^{0j}(t)\partial_t\partial_j u + \alpha(t)u_t + X^j(t)\partial_j u + \zeta(t)u = 0$$
(2.18)

(no summation on j) satisfies the conditions of Proposition 2.5; note that (2.18) is an equation of the form (1.2) with f = 0, d = 1 and R = 0. Let  $\epsilon > 0$ . Then there is a set of smooth  $\mathbb{C}^m$ -valued initial data to (1.2), say  $\mathcal{A}$ , with the following properties. First,  $\mathcal{A}$  is the intersection of a countable number of open and dense sets (with respect to the  $\mathbb{C}^\infty$  topology). Second, each element of  $\mathcal{A}$  corresponds to a  $\mathbb{C}^m$ -valued solution u to (1.2) such that there is a time sequence  $\{t_k\}, 1 \leq k \in \mathbb{Z}$ , with the properties that  $t_k \to \infty$  and

$$\mathfrak{E}_{\text{hom}}[u](t_k) \ge \exp\left[2\beta_X^{-1}\kappa(1-\epsilon)e^{\beta_X t_k}\right]$$
(2.19)

for all  $1 \leq k \in \mathbb{Z}$ , where  $\mathfrak{E}_{hom}[u](t)$  is defined by (1.21).

**Remark 2.19.** In the present proposition, we do not assume that f = 0, nor do we assume that R = 0 or that d = 1. Moreover, we take it to be understood that the equation (2.18) is obtained from (1.2) by setting f to zero and dropping all terms involving spatial derivatives other than  $\partial_i$ .

Remark 2.20. The result is an immediate consequence of Proposition 16.34.

Even though this result is interesting, we would also like to know that the set of initial data that do not yield solutions with super exponential growth has positive codimension. The next result states that the codimension is in fact infinite.

**Proposition 2.21.** Consider (1.2). Assume the associated metric to be such that (M, g) is a canonical separable cosmological model manifold. Assume, moreover, that there is a  $j \in \{1, ..., d\}$  such that (2.18) (no summation on j) satisfies the conditions of Proposition 2.5. Let  $\epsilon > 0$ . Define  $\mathcal{B}_{\epsilon}$  by the condition that  $\xi \in \mathcal{B}_{\epsilon}$  iff  $\xi$  constitutes smooth  $\mathbb{C}^{m}$ -valued initial data for (1.2) such that if u is the solution corresponding to  $\xi$ , then there is a constant  $C_{\epsilon}$  (depending on the solution) such that

$$\mathfrak{E}_{\text{hom}}[u](t) \le C_{\epsilon} \exp\left[2\beta_X^{-1}\kappa(1-\epsilon)e^{\beta_X t}\right]$$

for all  $t \geq 0$ , where  $\mathfrak{E}_{\text{hom}}$  is defined by (1.21). Define  $\mathcal{A}_{\epsilon}$  by the condition that  $\xi \in \mathcal{A}_{\epsilon}$  iff  $\xi$  constitutes smooth  $\mathbb{C}^m$ -valued initial data for (1.2) such that if u is the solution corresponding to  $\xi$ , then there is a time sequence  $0 \leq t_k \to \infty$  such that

$$\mathfrak{E}_{\hom}[u](t_k) \ge \exp\left[2\beta_X^{-1}\kappa(1-\epsilon/2)e^{\beta_X t_k}\right]$$
(2.20)

for all k. Then  $\mathcal{A}_{\epsilon}$  and  $\mathcal{B}_{\epsilon}$  are disjoint. Moreover, there is an infinite dimensional (linear) subspace  $P_{\epsilon}$  of the vector space of smooth  $\mathbb{C}^m$ -valued initial data for (1.2) such that if  $\xi \in \mathcal{B}_{\epsilon}$  and  $\xi_{\text{per}} \in P_{\epsilon}$ , then  $\xi + \xi_{\text{per}} \in \mathcal{A}_{\epsilon}$  unless  $\xi_{\text{per}} = 0$ .

**Remarks 2.22.** The statement of the proposition can be interpreted as saying that  $\mathcal{B}_{\epsilon}$  has infinite codimension in the set of smooth initial data for (1.2). On the other hand, there are examples of equations such that  $\mathcal{B}_{\epsilon}$ , even though it has infinite codimension, is dense in the set of initial data; this is a consequence of the current proposition and the results of Section 16.11. In fact, there are examples of equations satisfying the conditions of Proposition 2.21 such that all solutions with finite frequency content decay super exponentially. Having one dense set of initial data yielding super exponential growth and one dense set of initial data yielding super exponential growth and one dense set of unter the super exponential growth can be detected by only considering solutions with finite frequency content (even when solutions with finite frequency content exhibit super exponential decay); cf. Subsection 16.8.3.

**Remark 2.23.** By the vector space of smooth  $\mathbb{C}^m$ -valued initial data for (1.2), we here mean the space  $C^{\infty}(\bar{M}, \mathbb{C}^m) \times C^{\infty}(\bar{M}, \mathbb{C}^m)$ , where  $\bar{M}$  is defined by (1.3).

Remark 2.24. The statement is an immediate consequence of Proposition 16.37.

To conclude: solutions to equations satisfying the conditions of Propositions 2.18 and 2.21 typically exhibit super exponential growth. Returning to the conditions, it is thus clear that when  $X^{j}(t)/[g^{jj}(t)]^{1/2}$  exhibits definite exponential growth, the behaviour of solutions to (1.2) might be pathological. The next step of our analysis is to introduce the notion of a balanced equation, the purpose being to exclude this pathological behaviour.

## 2.5 A rough idea of the argument in a model case

In order to obtain a complete proof of the statements of this chapter, it is sufficient to read Parts III and IV of these notes. However, the material in Part III covers much more general situations than are needed to obtain the results of the present chapter. Moreover, the level of abstraction is sufficiently high that extracting the essential idea of the argument from the formal proof is non-trivial. For that reason, it is of interest to develop some intuition by considering a model case. That is the purpose of the present section.

A model case. Let us return to (2.3), the equation for the Fourier coefficients of (2.2). In the coefficients of this equation, there are terms of three different sizes: 1,  $|n|e^{-t}$  and  $n^2e^{-4t}$ . For  $|n| \ge 2$ , the third term dominates initially. Denote the subinterval of  $[0, \infty)$  during which  $n^2e^{-4t}$  is largest by  $I_1$ ; it is of the form  $I := [0, t_{\text{fin}}]$ . Considering the intervals on which the other terms dominate is also of interest, but in order to demonstrate generic super exponential growth, it is sufficient to focus on  $I_1$ . The equation (2.3) can be written  $\dot{w} = Aw$ , where

$$w(n,t) := \begin{pmatrix} \mathfrak{g}(n,t)z(n,t) \\ \dot{z}(n,t) \end{pmatrix}, \quad A(n,t) := \begin{pmatrix} -2 & \mathfrak{g}(n,t) \\ -\mathfrak{g}(n,t) - i\frac{ne^t}{|n|} - \frac{e^{2t}}{|n|} & -1 \end{pmatrix}$$

and  $\mathfrak{g}(n,t) = |n|e^{-2t}$ . The dominant behaviour of solutions in  $I_1$  is oscillatory with a period roughly given by  $T = 2\pi/\mathfrak{g}(n,t)$ .

**Approximating the oscillations.** Since the dominant behaviour is oscillatory, it is natural to divide the analysis of the evolution into two steps. First we consider one period of the oscillations;

the corresponding analysis yields a matrix describing the evolution over the period. Then we draw conclusions concerning the evolution over longer periods of time by multiplying together these matrices. Assuming the oscillations to be sufficiently rapid, the evolution within one period can be approximated by freezing the coefficients of the equation. Freezing the coefficients of A at  $t = t_0$ , the change in the solution from  $t_0$  to  $t_0 + T_0$ , where  $T_0 := 2\pi/\mathfrak{g}(n, t_0)$ , roughly speaking corresponds to multiplication by  $\exp[A(n, t_0)T_0]$ . The matrix  $A(n, t_0)T_0$ , in its turn, is roughly speaking given by  $2\pi B_0$ , where

$$B_0 := \left(\begin{array}{cc} 0 & 1\\ -1 - ir & 0 \end{array}\right)$$

and  $r := n^{-1}e^{3t_0}$ . There are various ways of calculating  $\exp(2\pi B_0)$ . In Section 13.2, we study the matrix exponentials of a general class of matrices that is of interest when approximating oscillations by freezing coefficients. Using these calculations, in particular Lemma 13.3, yields

$$\exp[2\pi B_0] \approx \mathrm{Id}_2 + \pi \left(\begin{array}{cc} 0 & ir \\ -ir & 0 \end{array}\right) \approx \exp\left[\pi \left(\begin{array}{cc} 0 & ir \\ -ir & 0 \end{array}\right)\right].$$

The eigenvalues of the matrix inside the exponential on the right hand side are given by  $\pm \pi r$ . In particular, the largest eigenvalue is given by

$$\pi |r| = \pi |n|^{-1} e^{3t_0} = \frac{1}{2} e^{t_0} T_0.$$

It is thus natural to expect the growth of the solution from  $t_0$  to  $t_0 + T_0$  to be  $\exp(e^{t_0}T_0/2)$ . Here  $T_0$  should be thought of as a small increment in t, so that summing  $e^{t_0}T_0 + e^{t_1}T_1 + \ldots$ (where  $t_1 := t_0 + T_0$ ,  $T_1 := 2\pi/\mathfrak{g}(n, t_1)$  etc.) should roughly speaking yield the integral of  $e^t$ . The change in absolute value of the solution from 0 to t should thus, roughly speaking, correspond to multiplication by  $\exp(e^t/2)$ . This expectation should be compared with the conclusions of Example 2.9.

**Gevrey classes.** The above line of reasoning only concerns one mode. Sooner or later, both  $|n|e^{-t}$  and  $n^2e^{-4t}$  are less than or equal to one. At that point, the ODE behaviour takes over and yields exponential decay of the Fourier coefficient. However, by combining infinitely many modes, we obtain super exponential growth of smooth solutions. The constructed solutions have the property that the frequency of the dominant Fourier coefficient becomes larger as the time increases. On the other hand, in order to obtain super exponential growth, the size of the Fourier coefficients of the initial data must not decay too fast with |n|; otherwise the growth in the intervals of the type  $I_1$  described above is not sufficient to ensure overall growth. This is why we only obtain super exponential growth for the Gevrey indices indicated in Remark 2.16.

Making the argument rigorous. The above discussion provides the intuition behind the proof that there are solutions that grow super exponentially. However, turning the ideas into a rigorous argument requires an effort. The purpose of Part III of these notes is to calculate the evolution over one period to leading order, and to provide an estimate for the error. To this end, we carry out several changes of variables that serve the purpose of isolating the essential behaviour. Moreover, we calculate the leading order terms in the relevant matrix exponentials (and estimate the errors); we estimate the variation of flows associated with the coefficients of (1.2) during one period etc. Once this has been done, we are interested in multiplying together the corresponding matrices in order to derive conclusions concerning the overall behaviour. As |n| grows, so does the number of factors. Proving that we can control the error of the product, even though there is no bound on the number of factors, is non-trivial. Moreover, in the present setting, we are interested in deriving a lower bound on the size of Fourier coefficients corresponding to appropriately chosen initial data. The necessary arguments are presented in Sections 16.3–16.6.

## Chapter 3

## A geometric perspective

## 3.1 Introduction

Before considering the general case, let us return to (2.2), our basic example of an equation with solutions exhibiting super exponential growth. What is the cause of this growth? Considering the assumptions required in order for Proposition 2.5 to apply, in particular (2.9), it is clear that it is related to the fact that  $Y^1$  grows exponentially. In order to obtain a more geometric perspective on the meaning of this object, note that the natural Lorentz metric associated with the equation (2.2) is given by

$$q = -dt \otimes dt + e^{4t} d\theta \otimes d\theta, \tag{3.1}$$

and defined on  $\mathbb{S}^1 \times \mathbb{R}$ . Moreover, for each  $t \in \mathbb{R}$ , g induces a metric on  $\mathbb{S}^1$ :

$$\bar{g} := e^{4t} d\theta \otimes d\theta$$

Turning to the lower order terms,  $e^{-t}u_{\theta}$  can be interpreted as  $\mathcal{X}u$ , where  $\mathcal{X}$  is the vector field  $\mathcal{X} = e^{-t}\partial_{\theta}$ . Thus, even though the coefficient of  $u_{\theta}$  appearing in (2.2) decays exponentially, it is clear that the norm of  $\mathcal{X}$  with respect to  $\bar{g}$  does not. In fact,  $\bar{g}(\mathcal{X}, \mathcal{X}) = e^{2t}$ . When viewed from a more geometric perspective, it is thus clear that  $\mathcal{X}$  is not small. Given the metric (3.1), it is natural to introduce an orthonormal frame  $\{e_0, e_1\}$ , where  $e_0 = \partial_t$  and  $e_1 = e^{-2t}\partial_{\theta}$ . With respect to this frame,  $\mathcal{X} = e^t e_1$ ; i.e.,  $\mathcal{X} = Y^1 e_1$ .

## 3.2 Conditions on the metric

Let us now return to the general case. Consider (1.2). Assume the associated metric to be such that (M,g) is a canonical separable cosmological model manifold (recall that this implies that  $g^{00} = -1$ ; cf. Remark 1.19). Then

$$g = -dt \otimes dt + g_{ij}(\chi^i dt + dx^i) \otimes (\chi^j dt + dx^j) + \sum_{r=1}^R a_r^2 g_r.$$
(3.2)

Note that (3.2) induces a family of metrics on  $\overline{M}$ , given by

$$\bar{g} := g_{ij} dx^i \otimes dx^j + \sum_{r=1}^R a_r^2 g_r,$$

where we normally do not explicitly state the dependence on t. The metric g also induces a family of metrics on  $\mathbb{T}^d$ :

$$\bar{h} := g_{ij} dx^i \otimes dx^j.$$

Again, we normally omit explicit reference to the dependence on t. Next we wish to impose restrictions on the metric (3.2). When studying Einstein's equations, there is a freedom in choosing the gauge. For this reason there is some freedom in imposing conditions on the metric coefficients. In particular, the lapse function and the shift vector field, cf. Remark 1.5, are related to a choice of foliation of the spacetime. In the case of the metric g defined by (3.2), the lapse function equals 1 and the shift vector field is given by  $\chi$ . Even though one typically has a freedom of imposing conditions on the shift vector field, setting it to zero is sometimes problematic. That is the reason why we include it here. On the other hand, it is rarely useful to have a shift vector field with a large norm. Moreover, it is often convenient if the foliation is such that  $\partial_t$  is an approximate normal to the constant t hypersurfaces. Considering (3.2), the future directed unit normal to  $\overline{M}_t$ is

$$U := \partial_t - \chi^i \partial_i. \tag{3.3}$$

To begin with, we therefore impose the following conditions.

**Definition 3.1.** Let (M, g) be a canonical separable cosmological model manifold. If there is an  $0 < \eta_{sh,0} \leq 1$  such that

$$g(\partial_t, \partial_t) \le -\eta_{\mathrm{sh},0}^2 \tag{3.4}$$

for all  $t \ge 0$ , then  $\partial_t$  is said to be *future uniformly timelike*. If, in addition, there is a  $1 \le k \in \mathbb{Z}$ and an  $0 < \eta_{\mathrm{sh},k} \in \mathbb{R}$  such that

$$\sum_{l=1}^{k} |\mathcal{L}_{U}^{l}\chi|_{\bar{g}} \le \eta_{\mathrm{sh},k} \tag{3.5}$$

for all  $t \ge 0$ , then the shift vector field of (3.2) is said to be  $C^k$ -future bounded.

**Remarks 3.2.** The estimate (3.4) is equivalent to  $g_{ij}\chi^i\chi^j \leq 1 - \eta_{\text{sh},0}^2$  (note that if  $g_{ij}\chi^i\chi^j \geq 1$ , then  $\partial_t$  is no longer timelike). Moreover,  $\mathcal{L}_U\chi$  denotes the Lie derivative of  $\chi$  with respect to U.

**Remarks 3.3.** Since  $\mathcal{L}_{\partial_t}\chi = \mathcal{L}_U\chi$ , we could replace  $\mathcal{L}_U\chi$  with  $\mathcal{L}_{\partial_t}\chi$  in (3.5). Note also that since  $\mathcal{L}_U\chi = \dot{\chi}^i\partial_i$ , we sometimes use the notation  $\dot{\chi} := \mathcal{L}_U\chi$ . Finally,

$$|\mathcal{L}_U^l \chi|_{\bar{g}} := \left[ \bar{g}(\mathcal{L}_U^l \chi, \mathcal{L}_U^l \chi) \right]^{1/2},$$

where it is understood that  $\mathcal{L}_{U}^{l}\chi$  and  $\bar{g}$  are both evaluated at the same time t.

Turning to the conditions on  $g_{ij}$  and  $a_r$ , it is convenient to express them in terms of the second fundamental form of the hypersurfaces  $\bar{M}_t$ ; cf. (1.5). The second fundamental form is given by

$$\bar{k}(Z_1, Z_2) := g(\nabla_{Z_1} U, Z_2),$$

where  $\nabla$  is the Levi-Civita connection associated with the Lorentz manifold (M, g), U is given by (3.3) and  $Z_i$ , i = 1, 2, are vector fields tangent to  $\overline{M}_t$ . It can be computed that

$$\bar{k} = \frac{1}{2} \sum_{i,j=1}^{d} \partial_t g_{ij} dx^i \otimes dx^j + \sum_{r=1}^{R} \frac{\dot{a}_r}{a_r} a_r^2 g_r,$$
$$\mathcal{L}_U \bar{k} = \frac{1}{2} \sum_{i,j=1}^{d} \partial_t^2 g_{ij} dx^i \otimes dx^j + \sum_{r=1}^{R} \left[ \frac{\ddot{a}_r}{a_r} + \left( \frac{\dot{a}_r}{a_r} \right)^2 \right] a_r^2 g_r.$$

Again,  $\mathcal{L}_U \bar{k} = \mathcal{L}_{\partial_t} \bar{k}$ , so that in the conditions stated below, we could replace  $\mathcal{L}_U \bar{k}$  with  $\mathcal{L}_{\partial_t} \bar{k}$ . The basic assumptions we make concerning  $\bar{k}$  are the following

**Definition 3.4.** Let (M, g) be a canonical separable cosmological model manifold. Let, moreover,  $0 \le k \in \mathbb{Z}$ . If there is a constant  $0 < C_k \in \mathbb{R}$  such that

$$\sum_{l=0}^{k} |\mathcal{L}_{U}^{l}\bar{k}|_{\bar{g}} \le C_{k}$$

for all  $t \geq 0$ , then the second fundamental form is said to be  $C^k$ -future bounded.
**Remark 3.5.** The notation  $|\bar{k}|_{\bar{g}}$  is defined by

$$|\bar{k}|_{\bar{g}} := \left(\bar{g}^{ab}\bar{g}^{cd}\bar{k}_{ac}\bar{k}_{bd}\right)^{1/2},$$

where the lower case Latin indices from the beginning of the alphabet indicate coordinate components on  $\overline{M}$  and it is understood that both  $\overline{k}$  and  $\overline{g}$  are evaluated at the same time t. In Section 25.5 we develop different perspectives on this norm.

**Remark 3.6.** The motivation for assuming  $\bar{k}$  to be bounded is given in Section 1.3. However, in some of the arguments, it is not sufficient to only assume  $\bar{k}$  to be bounded, we also need to impose bounds on  $\mathcal{L}_U \bar{k}$ . This is the motivation for introducing the terminology of the definition. Note also that the basic examples given in Subsection 1.3.1 are such that the second fundamental form is  $C^k$ -future bounded for any k.

## 3.3 The lower order terms

So far, we introduced bounds on the metric components; cf. Definitions 3.1 and 3.4. However, considering (1.2), we also need to impose restrictions on the  $X^j$ ,  $\alpha$  and  $\zeta$ . Concerning the  $X^j$ 's, it is convenient to combine them and think of the result as a matrix of vector fields on M:

$$\mathcal{X} := X^j(t)\partial_j. \tag{3.6}$$

Due to the results of Chapter 2, it is clear that we need to impose bounds on  $\mathcal{X}$  in order to exclude pathological behaviour of solutions. When stating the relevant bounds, the following notation is convenient:

$$|\mathcal{X}|_{\bar{h}} := \left(\sum_{\varsigma \in \mathfrak{S}} \sum_{i,j=1}^{d} \bar{h}_{ij}\varsigma_i \|X^i\| \cdot \varsigma_j \|X^j\|\right)^{1/2}, \tag{3.7}$$

where it is understood that both  $\mathcal{X}$  and  $\bar{h}$  are evaluated at the same time t and  $\mathfrak{S}$  is the set of elements of  $\mathbb{R}^d$  whose components are plus or minus one; note that  $\mathfrak{S}$  has  $2^d$  elements. We use similar notation for other matrices of vector fields (in the span of the  $\{\partial_j\}$ ). The reader interested in a motivation for the choice (3.7) of norm is referred to the proof of Lemma 25.17, in particular (25.62).

**Definition 3.7.** Consider (1.2). Assume the associated metric to be such that (M, g) is a canonical separable cosmological model manifold. Define  $\mathcal{X}$  by (3.6). If there is a  $0 \le k \in \mathbb{Z}$  and a  $0 < C_k \in \mathbb{R}$  such that

$$\sum_{l=0}^{k} |\mathcal{L}_{U}^{l} \mathcal{X}|_{\bar{h}} \le C_{k} \tag{3.8}$$

for all  $t \geq 0$ , then  $\mathcal{X}$  is said to be  $C^k$ -future bounded.

Finally, let us turn to the matrix valued functions  $\alpha$  and  $\zeta$  appearing in (1.2). Due to the results of Section 16.2, we know that if  $\alpha$  and  $\zeta$  are allowed to grow exponentially, then spatially homogeneous solutions to (1.2) (with f = 0) can exhibit super exponential growth or decay; cf. also the discussion in Section 2.1. We consider both of these types of behaviour to be pathological. For these reasons, we typically impose the conditions that  $\|\alpha(t)\|$ ,  $\|\dot{\alpha}(t)\|$ ,  $\|\zeta(t)\|$  and  $\|\dot{\zeta}(t)\|$  are bounded to the future.

# **3.4** A rough notion of balance

One basic notion of balance, prohibiting the occurrence of super exponential growth, is the following. **Definition 3.8.** Consider (1.2). Assume the associated metric to be such that (M, g) is a canonical separable cosmological model manifold. Let  $0 \le k \in \mathbb{Z}$ . If  $\partial_t$  is future uniformly timelike; there is a constant  $0 < C_k \in \mathbb{R}$  such that

$$\sum_{l=0}^{k} [\|\partial_t^l \alpha(t)\| + \|\partial_t^l \zeta(t)\|] \le C_k$$

for all t > 0 (i.e.,  $\alpha$  and  $\zeta$  are  $C^k$ -future bounded);  $\mathcal{X}$  is  $C^k$ -future bounded; the shift vector field of (3.2) is  $C^{k+1}$ -future bounded; and the second fundamental form is  $C^{k}$ -future bounded, then (1.2) is said to be  $C^{k+1}$ -balanced.

**Remark 3.9.** The bounds on the metric components involve k+1 derivatives, but the conditions on the coefficients of the lower order terms in (1.2) only involve k derivatives.

Given a  $C^1$ -balanced equation, the basic energy cannot grow faster than exponentially (except if the right hand side does). In fact, there is a constant  $0 < \eta_{\text{bal}} \in \mathbb{R}$  (depending only on the constants appearing in the bounds of Definition 3.8) such that

$$\mathfrak{E}_{\rm bas}^{1/2}[u](t_1) \le e^{\eta_{\rm bal}|t_1 - t_0|} \mathfrak{E}_{\rm bas}^{1/2}[u](t_0) + \left| \int_{t_0}^{t_1} e^{\eta_{\rm bal}|t_1 - t|} \|f(\cdot, t)\|_2 dt \right|$$
(3.9)

for all  $0 \le t_0, t_1 \in \mathbb{R}$  and all solutions u to (1.2), where  $||f(\cdot,t)||_2$  denotes the L<sup>2</sup>-norm of  $f(\cdot,t)$ . This statement follows from Lemma 10.15; cf. Remark 10.16. In fact, estimates of the form (3.9) hold under slightly weaker conditions; cf. Remark 10.16. Even though the estimate (3.9) is of interest, it only excludes pathological behaviour. We are also interested in deriving more detailed information concerning the asymptotics of solutions and concerning the behaviour of the basic energy; the constant  $\eta_{\rm bal}$  that results from an application of Lemma 10.15 could very well be quite far from the optimal value. In order to obtain more detailed conclusions, we need to make stronger assumptions.

#### 3.5Geometric formulation of the results in the unbalanced setting

At this stage, it is of interest to return to the unbalanced setting, and to express the conditions geometrically.

**Assumptions 3.10.** Consider (1.2). Assume the associated metric to be such that (M, q) is a canonical separable cosmological model manifold; cf. Definition 1.18. Assume, moreover, that d = 1, that R = 0 and that there is an  $0 < \eta_{sh,0} \le 1$  such that (3.4) holds. Assume that there are constants  $K_{\rm sh}, K_{\bar{k}}, K_{\rm ode}, \bar{K}_X > 0, \beta_{\rm sh} < 0, \beta_{\rm ode}$  and  $\beta_X > 0$  such that  $\beta_X - \beta_{\rm ode} > 0$  and

$$|\chi(t)|_{\bar{q}} + |\dot{\chi}(t)|_{\bar{q}} \le K_{\rm sh} e^{\beta_{\rm sh} t},$$
(3.10)

$$\begin{aligned} |\chi(t)|_{\bar{g}} + |\chi(t)|_{\bar{g}} \leq K_{\rm sh}e^{j\,\sin t}, \\ |\ddot{\chi}(t)|_{\bar{g}} + |\bar{k}(t)|_{\bar{g}} + |\mathcal{L}_U\bar{k}(t)|_{\bar{g}} \leq K_{\bar{k}}, \end{aligned}$$
(3.10)

$$\|\zeta(t)\|^{1/2} + \|\dot{\zeta}(t)\|^{1/2} + \|\alpha(t)\| + \|\dot{\alpha}(t)\| \le K_{\text{ode}}e^{\beta_{\text{ode}}t},$$
(3.12)

$$|\mathcal{X}(t)|_{\bar{h}} + |\dot{\mathcal{X}}(t)|_{\bar{h}} \le \bar{K}_X e^{\beta_X t} \tag{3.13}$$

for all  $t \geq 0$ .

Let  $e_1$  be the unit vector field which is a positive multiple of  $\partial_1$ , define  $\mathcal{Y}^1$  to be the matrix valued function such that  $\mathcal{Y}^1 e_1 = \mathcal{X}$  and assume that there is a matrix  $\mathcal{Y}^1_{\infty} \in \mathbf{M}_m(\mathbb{C})$  and  $\bar{\eta}_X, L_X > 0$ such that

$$\|e^{-\beta_X t} \mathcal{Y}^1(t) - \mathcal{Y}^1_\infty\| \le L_X e^{-\bar{\eta}_X t} \tag{3.14}$$

for all  $t \ge 0$ . Assume, moreover, that  $\mathcal{Y}^1_{\infty}$  has an eigenvalue with a non-zero real part.

Assume, finally, that there are  $\bar{\beta}_j \in \mathbb{R}$ , j = 1, 2, and a continuous function  $0 \leq \mathfrak{e} \in L^1([0, \infty))$  such that

$$(\bar{\beta}_2 - \mathfrak{e})\bar{g} \le \bar{k} \le (\bar{\beta}_1 + \mathfrak{e})\bar{g} \tag{3.15}$$

for all  $t \geq 0$ .

**Remark 3.11.** If Assumptions 3.10 hold, then Assumptions 2.1 and 2.3 hold with  $Y_{\infty}^1 = \mathcal{Y}_{\infty}^1$ ; cf. Lemma 25.19 and Remark 25.20.

**Remark 3.12.** The only quantities we require to converge are the components of  $e^{-\beta_X t} \mathcal{X}$  with respect to the unit vector field  $e_1$ . Moreover, the condition  $\beta_X - \beta_{ode} > 0$  implies that  $\mathcal{Y}^1$  is larger in size than  $\|\alpha(t)\|$  and  $\|\zeta(t)\|^{1/2}$ .

**Remark 3.13.** One consequence of the assumptions is that there is a constant  $c_1 > 0$  such that  $c_1 e^{-2\bar{\beta}_1 t} \leq g^{11}(t)$  for all  $t \geq 0$ ; this follows from Lemma 25.21.

**Proposition 3.14.** Consider (1.2). Assume that d = 1, R = 0 and f = 0. Given that Assumptions 3.10 hold, let  $\kappa > 0$  be the largest absolute value of a real part of an eigenvalue of  $\mathcal{Y}_{\infty}^{1}/2$ . Let  $\epsilon > 0$ . Then there is a sequence of smooth  $\mathbb{C}^{m}$ -valued solutions  $v_{l}$  to (1.2),  $1 \leq l \in \mathbb{Z}$ , and for each  $l \geq 1$ , there is a time sequence  $t_{l,k} \to \infty$  (as  $k \to \infty$ ) such that for each  $s \in \mathbb{R}$ ,

$$\lim_{l \to \infty} \left( \|v_l(\cdot, 0)\|_{(s+1)} + \|\partial_t v_l(\cdot, 0)\|_{(s)} \right) = 0.$$
(3.16)

In addition,

$$\mathfrak{E}_{\hom}[v_l](t_{l,k}) \ge \exp\left[2\beta_X^{-1}\kappa(1-\epsilon)e^{\beta_X t_{l,k}}\right],\tag{3.17}$$

where  $\mathfrak{E}_{\text{hom}}[u]$  is given by (2.12). Moreover, if  $\beta_X + \bar{\beta}_1 > 0$ , then the sequence of functions  $v_l$  can be assumed to be such that  $v_l, \partial_t v_l \in \mathfrak{G}^{s_X}_{\text{loc},u}(I, \mathbb{S}^1, \mathbb{C}^m)$  for all l, where

$$s_X := \frac{\beta_X + \bar{\beta}_1}{\beta_X}.$$
(3.18)

In case  $\beta_X + \overline{\beta}_1 \leq 0$ , one frequency can be chosen, say  $n_a \in \mathbb{Z}$ , such that the sequence of functions  $v_l$  has the property that if  $z_l(n,t)$  denotes the n'th Fourier coefficient of  $v_l(\cdot,t)$ , then  $z_l(n,t) = 0$  for all t unless  $n = n_a$ .

**Remark 3.15.** If  $\mathcal{Y}^1_{\infty}$  is Hermitian, (3.17) is essentially optimal. In fact, in that case there is, given a solution u, a constant  $C_{\epsilon}$  such that  $\mathfrak{E}_{\text{hom}}[u](t) \leq C_{\epsilon} \exp\left[2\beta_X^{-1}\kappa(1+\epsilon)e^{\beta_X t}\right]$  for all  $t \geq 0$ ; cf. Remark 2.8.

**Remark 3.16.** The constituents of the right hand side of (3.18) have a natural interpretation. The constant  $\bar{\beta}_1$  represents the (average) maximal expansion of the spacetime, in the sense that it corresponds to the upper bound in (3.15). Moreover,  $\beta_X$  is the exponential rate of growth of the norm of the vector field  $\mathcal{X}$  (with respect to the induced metric on the constant-*t* hypersurfaces). The statements concerning the regularity can be summarised as follows. If the average maximal expansion is strictly positive (i.e., if  $\bar{\beta}_1 > 0$ ), then  $s_X > 1$ , and the initial data are not analytic. If the average maximal expansion is zero (i.e., if  $\bar{\beta}_1 = 0$ ), then  $s_X = 1$ , and the initial data are analytic. If the average maximal expansion is negative, but strictly greater than minus the growth rate of  $\mathcal{X}$  (i.e., if  $-\beta_X < \bar{\beta}_1 < 0$ ), then  $0 < s_X < 1$ , and the initial data are entire. If the average maximal expansion is less than or equal to minus the growth rate of  $\mathcal{X}$  (i.e., if  $\bar{\beta}_1 \leq -\beta_X$ ), then the initial data have only one non-zero Fourier coefficient.

*Proof.* Due to the assumptions of the proposition, the conditions of Proposition 2.5 are fulfilled. Moreover, due to Remark 3.13, the conditions of Proposition 2.14 hold with  $\beta_1 = -\bar{\beta}_1$ . Combining the corresponding conclusions yields the statement of the proposition.

Next, let us turn to the question of the generic behaviour.

**Proposition 3.17.** Consider (1.2). Assume the associated metric to be such that (M, g) is a canonical separable cosmological model manifold. Assume, moreover, that there is a  $j \in \{1, ..., d\}$  such that

$$u_{tt} - g^{jj}(t)\partial_j^2 u - 2g^{0j}(t)\partial_t\partial_j u + \alpha(t)u_t + X^j(t)\partial_j u + \zeta(t)u = 0$$
(3.19)

(no summation on j) satisfies Assumptions 3.10. Then the conclusions of Propositions 2.18 and 2.21 hold.

**Remark 3.18.** The conclusions of Propositions 2.18 and 2.21 can roughly speaking be summarised as saying that there is a Baire generic set of initial data corresponding to solutions exhibiting super exponential growth; and that the initial data corresponding to solutions that do not exhibit super exponential growth have infinite co-dimension.

*Proof.* The conditions of the proposition ensure that the assumptions required to appeal to Propositions 2.18 and 2.21 are satisfied.  $\Box$ 

It is of interest to contrast this result with conclusions obtained concerning solutions with finite frequency content. In particular, we here consider a situation in which the dominant coefficients in (1.2) are  $\alpha$  and  $\zeta$ . Recalling the notation introduced in Section 24.1, we have the following result.

**Proposition 3.19.** Consider (1.2). Assume the associated metric to be such that (M, g) is a canonical separable cosmological model manifold. Assume, moreover, that f = 0. Assume that there is a  $\gamma > 0$ ;  $\alpha_{\infty}, \zeta_{\infty} \in \mathbf{M}_m(\mathbb{C})$ , one of which is non-zero; an  $\eta_{cu} > 0$ ; and a constant  $C_{cu} > 0$  such that

$$|g^{0l}(t)| + |g^{jl}(t)|^{1/2} + |a_r^{-1}(t)| + ||X^j(t)||^{1/2} \le C_{\rm cu} e^{(\gamma - \eta_{\rm cu})t}, ||e^{-\gamma t}\alpha(t) - \gamma \alpha_{\infty}|| + ||e^{-2\gamma t}\zeta(t) - \gamma^2 \zeta_{\infty}||^{1/2} \le C_{\rm cu} e^{-\eta_{\rm cu}t}$$

for all  $t \ge 0$ , j, l = 1, ..., d, and r = 1, ..., R. Let  $A_{\infty}$  be defined by (2.1) and  $\kappa_1$  be the largest real part of an eigenvalue of  $A_{\infty}$ . Consider a solution u to (1.2) such that if z is defined by (24.14), then  $z(\iota, \cdot) = 0$  for all but a finite number of  $\iota \in \mathcal{I}_{B}$ . Fix  $\epsilon > 0$ . Then there is a constant  $C_{\epsilon} > 0$ such that

$$\mathfrak{E}_{\mathrm{bas}}[u](t) \le C_{\epsilon} \exp[2(\kappa_1 + \epsilon)e^{\gamma t}]\mathfrak{E}_{\mathrm{bas}}[u](0)$$

for all  $t \geq 0$ . Here  $\mathfrak{E}_{\text{bas}}$  is defined by (1.19).

**Remark 3.20.** The constant  $C_{\epsilon}$  only depends on  $C_{cu}$ ; the set of  $\iota \in \mathcal{I}_{B}$  for which  $z(\iota, \cdot)$  does not vanish; the spectrum of the Laplace-Beltrami operator on  $(M_r, g_r)$  for  $r = 1, \ldots, R$ ;  $A_{\infty}$ ;  $\eta_{cu}$ ;  $\gamma$ ; and  $\epsilon$ .

**Remark 3.21.** There are large classes of equations satisfying both the assumptions of Proposition 3.17 and the assumptions of Proposition 3.19 with  $\kappa_1 < 0$ . For the corresponding equations, solutions with finite frequency content decay to zero super exponentially. In other words, there is a dense set of initial data such that the corresponding solutions exhibit super exponential decay. However, generic solutions grow super exponentially.

**Remark 3.22.** Appealing to Lemma 16.59 instead of to Lemma 16.56 yields an analogous result in the case that  $\alpha$  and  $\zeta$  converge exponentially to elements of  $\mathbf{M}_m(\mathbb{C})$ . In that case, assuming the quantity corresponding to  $\kappa_1$  in Proposition 3.19 to be negative, solutions with finite frequency content decay exponentially.

*Proof.* The conclusion is an immediate consequence of Lemma 16.56.

# Chapter 4

# Silent equations

# 4.1 Introduction

Heuristics. Consider (1.23). As already mentioned, the direction  $t \to \infty$  corresponds to the big bang singularity of polarised T<sup>3</sup>-Gowdy symmetric solutions to Einstein's equations. In heuristic arguments concerning cosmological singularities, similar equations appear. In particular, it is often expected, on heuristic grounds, to be possible to approximate the full Einstein equations with equations such that the coefficients multiplying the spatial derivatives of the unknowns decay exponentially. A similar structure appears in the expanding direction when considering solutions to Einstein's equations with a positive cosmological constant. In the physics literature, the exponential decay of the coefficients is often assumed to justify dropping the spatial derivatives. This results in one ODE for each spatial point, a substantially simplified situation. In order to illustrate this line of reasoning, consider (1.23) again. Dropping the term involving the spatial derivatives yields the equation  $P_{tt} = 0$  with solutions  $P(\theta, t) = v(\theta)t + \psi(\theta)$ . This heuristically motivated conclusion should be compared with the statements made in connection with (1.24); see also Example 4.20 below.

Due to the results of Chapter 2, we know that the above line of reasoning does not always work; cf., in particular, Example 2.9. On the other hand, perhaps the situation changes if we assume the equation to be balanced in the sense of Definition 3.8. For this reason, it is of interest to consider balanced systems of equations such that the coefficients of the spatial derivatives decay exponentially. Moreover, the question we wish to ask is: can the above heuristic picture be given a mathematical justification?

**Examples.** Considering the examples of geometries given in Section 1.3, the metrics  $g_{dS}$  and  $g_{gdS}$  yield principal symbols such that the coefficients of the highest order spatial derivatives decay exponentially. Considering the contracting direction of Kasner solutions (with the exception of the flat Kasner solutions), or of non-vacuum Bianchi type I solutions to the Einstein-scalar field equations, yields the same conclusion.

**Geometry.** The above heuristics are formulated in terms of properties of the coefficients of the equations. However, it is desirable to formulate geometric conditions. Exponential decay of the coefficients of the spatial derivatives is related to exponential expansion. Exponential expansion, in its turn, is related to lower bounds on the second fundamental form. In the examples given above, not only do the second fundamental forms have a uniform strictly positive lower bound (for late times), they, in fact, converge. However, in the present chapter, the main assumption is that of a lower bound. It is therefore of interest to introduce the following terminology.

**Definition 4.1.** Consider (1.2). Assume the associated metric to be such that (M, g) is a canonical separable cosmological model manifold. If  $\partial_t$  is future uniformly timelike and there is a  $0 < \mu \in \mathbb{R}$ 

and a continuous non-negative  $\mathfrak{e} \in L^1([0,\infty))$  such that

$$k \ge (\mu - \mathfrak{e})\bar{g}, \quad |\chi|_{\bar{g}} \cdot |\dot{\chi}|_{\bar{g}} \le \mathfrak{e}$$

$$\tag{4.1}$$

for all  $t \ge 0$ ; then (1.2) is said to be  $C^1$ -silent.

**Remark 4.2.** In Subsection 1.3.2, we motivate the terminology "silent metric" in the context of the metrics introduced in (1.18). The motivation in the present setting is similar. In fact, let  $\gamma: J \to M$  be a future directed inextendible causal curve in (M, g), where  $J = (s_-, s_+)$ . Then  $\gamma^0(s) \to \infty$  as  $s \to s_+$ , where  $\gamma^0$  denotes the *t*-coordinate of  $\gamma$ ; this is an immediate consequence of the fact that  $\overline{M}_t$  is a Cauchy hypersurface in (M, g) for each  $t \in \mathbb{R}$  (cf. Lemma 25.1) and the tacit assumption that  $\partial_t$  is future oriented. Moreover, given that the assumptions of the definition are satisfied, the  $\overline{M}$ -coordinate of  $\gamma$ , say  $\overline{\gamma}$ , converges to a point, say  $\overline{p}[\gamma]$  as  $s \to s_+$ . Finally, if two future directed inextendible causal curves, say  $\gamma_i: J_i \to M, i = 1, 2$ , are such that  $\overline{p}[\gamma_1] \neq \overline{p}[\gamma_2]$ , then there are  $s_i \in J_i$  such that

$$J^+[\gamma_1(s_1)] \cap J^+[\gamma_2(s_2)] = \varnothing;$$

the reader interested in an explanation of the terminology  $J^+(p)$  is referred to [27]. Thus, sooner or later, the observers  $\gamma_i$  lose the ability to communicate to the future. In other words, there is silence. For a justification of these statements, the reader is referred to Lemma 25.23 and Remark 25.24.

In the present chapter, we are interested in equations (1.2) that are  $C^1$ -silent in the sense of Definition 4.1; such that  $\mathcal{X}$  is  $C^0$ -future bounded; and such that there are  $\alpha_{\infty}, \zeta_{\infty} \in \mathbf{M}_m(\mathbb{C})$  and  $0 < \eta_{\mathrm{mn}}, C_{\mathrm{mn}} \in \mathbb{R}$  with the property that

$$|\alpha(t) - \alpha_{\infty}|| + ||\zeta(t) - \zeta_{\infty}|| \le C_{\mathrm{mn}} e^{-\eta_{\mathrm{mn}} t}$$

$$\tag{4.2}$$

for all  $t \ge 0$ . Under these assumptions, all the coefficients in (1.2) that multiply spatial derivatives of u decay to zero exponentially; cf. Lemma 26.1. Since (4.2) is satisfied, it is then of interest to compare solutions to (1.2) with solutions to the equation

$$\partial_t \left(\begin{array}{c} v\\ v_t \end{array}\right) = A_{\infty} \left(\begin{array}{c} v\\ v_t \end{array}\right) + \left(\begin{array}{c} 0\\ f \end{array}\right), \tag{4.3}$$

where

$$A_{\infty} := \begin{pmatrix} 0 & \mathrm{Id}_m \\ -\zeta_{\infty} & -\alpha_{\infty} \end{pmatrix}.$$
(4.4)

Note that if one drops the second term on the right hand side of (4.3), then what results is a system of constant coefficient equations in which no spatial derivatives occur. On the other hand, we have not eliminated the dependence of the solution on the spatial variables. Thus (4.3) should be interpreted as a system of ODE's for each point in  $\overline{M}$ . In case f = 0, one would expect solutions to (4.3) to grow as  $\langle t \rangle^{d_1-1} e^{\kappa_1 t}$ , where  $\kappa_1$  is the largest real part of an eigenvalue of  $A_{\infty}$  and  $d_1$  is the largest dimension of a Jordan block corresponding to an eigenvalue of  $A_{\infty}$  with real part  $\kappa_1$ ; cf. Section 24.2 below for a justification. Here we use the notation

$$\langle \xi \rangle := (1 + |\xi|^2)^{1/2} \tag{4.5}$$

for  $\xi \in \mathbb{C}^l$ . Due to the importance of  $\kappa_1$  and  $d_1$  in the description of the asymptotics, it is convenient to introduce the following terminology.

**Definition 4.3.** Given  $A \in \mathbf{M}_k(\mathbb{C})$ , let SpA denote the set of eigenvalues of A. Moreover, let

$$\kappa_{\max}(A) := \sup\{\operatorname{Re}\lambda \mid \lambda \in \operatorname{Sp}A\}, \quad \kappa_{\min}(A) := \inf\{\operatorname{Re}\lambda \mid \lambda \in \operatorname{Sp}A\}.$$

Then RspA, the real eigenvalue spread of A, is defined by RspA :=  $\kappa_{\max}(A) - \kappa_{\min}(A)$ . In addition, if  $\kappa \in \{\operatorname{Re}\lambda \mid \lambda \in \operatorname{Sp}A\}$ , then  $d_{\max}(A, \kappa)$  is defined to be the largest dimension of a Jordan block corresponding to an eigenvalue of A with real part  $\kappa$ . Finally, if  $\kappa \notin \{\operatorname{Re}\lambda \mid \lambda \in \operatorname{Sp}A\}$ , then  $d_{\max}(A, \kappa) := 1$ .

## 4.2 Results

Concerning the energy of solutions to (1.2), we obtain the following conclusion.

**Proposition 4.4.** Consider (1.2). Assume that it is  $C^1$ -silent in the sense of Definition 4.1; that  $\mathcal{X}$  is  $C^0$ -future bounded; and that there are  $\alpha_{\infty}, \zeta_{\infty} \in \mathbf{M}_m(\mathbb{C})$  and  $0 < \eta_{\mathrm{mn}}, C_{\mathrm{mn}} \in \mathbb{R}$  with the property that (4.2) holds for all  $t \geq 0$ . Then there are constants C and  $s_0 \geq 0$ , depending only on the coefficients of the equation (1.2), such that if u is a smooth solution to (1.2), then

$$\mathfrak{E}_{s}^{1/2}[u](t) \leq C\langle t \rangle^{d_{1}-1} e^{\kappa_{1} t} \mathfrak{E}_{s+s_{0}}^{1/2}[u](0) + C \int_{0}^{t} \langle t - \tau \rangle^{d_{1}-1} e^{\kappa_{1}(t-\tau)} \|f(\cdot,\tau)\|_{(s+s_{0})} d\tau$$
(4.6)

for all  $t \ge 0$  and  $s \in \mathbb{R}$ , where  $\kappa_1 := \kappa_{\max}(A_{\infty})$ ,  $d_1 := d_{\max}(A_{\infty}, \kappa_1)$  and  $A_{\infty}$  is given by (4.4); cf. Definition 4.3.

**Remark 4.5.** The proposition is a consequence of Lemma 10.11; cf. Remark 10.14.

**Remarks 4.6.** More detailed information concerning the dependence of the constants is to be found in Remark 10.13. The energy  $\mathfrak{E}_s$  is defined by (24.19). The object  $||f(\cdot,t)||_{(s)}$  is the  $H^s$ -Sobolev norm of the function  $f(\cdot,t)$ ; cf. (24.10).

Next we would like to derive more detailed information concerning the asymptotics. Naively, we would expect a solution to (1.2) to be well approximated by a solution to (4.3). However, there are obstructions. Say, for the sake of argument, that

$$||f||_{A,s} := \int_0^\infty e^{-\kappa_1 \tau} ||f(\cdot, \tau)||_{(s)} d\tau < \infty$$
(4.7)

for  $s \in \mathbb{R}$ , where  $\kappa_1$  is the number defined in the statement of Proposition 4.4. Then the energies associated with solutions u to (1.2) are bounded by  $C\langle t \rangle^{2d_1-2}e^{2\kappa_1 t}$ ; cf. (4.6). Moreover, as is demonstrated below, this estimate is optimal. On the other hand, when taking the step from (1.2) to (4.3), we have ignored terms of the form

$$\chi^{j}\partial_{j}\partial_{t}u, \quad X^{j}\partial_{j}u, \quad (\alpha - \alpha_{\infty})u_{t}, \quad (\zeta - \zeta_{\infty})u,$$
(4.8)

as well as the terms involving second spatial derivatives of u. On the other hand, the first two terms appearing in (4.8) can be expected to be of the order of magnitude  $\langle t \rangle^{d_1-1} e^{(\kappa_1-\mu)t}$ , and the last two terms can be expected to be of the order of magnitude  $\langle t \rangle^{d_1-1} e^{(\kappa_1-\eta_{\rm mn})t}$ ; cf. (4.2) and Lemma 26.1. Moreover, the terms involving second spatial derivatives of u can be expected to be of the order of magnitude  $\langle t \rangle^{d_1-1} e^{(\kappa_1-2\mu)t}$ ; cf. Lemma 26.1. For this reason, it would be optimistic to think that the terms appearing in solutions to (4.3) that are of the order of magnitude  $e^{(\kappa_1-\beta_{\rm rem})t}$ , where

$$\beta_{\rm rem} := \min\{\mu, \eta_{\rm mn}\} \tag{4.9}$$

can be distinguished from the error terms arising when the terms (4.8) are omitted. When considering solutions to the homogeneous version of the equation (4.3), it therefore seems natural to focus on solutions whose norms asymptotically exceed  $e^{(\kappa_1 - \beta_{\text{rem}})t}$ . For this reason, it is natural to introduce the following terminology.

**Definition 4.7.** Let  $1 \leq n \in \mathbb{Z}$ ,  $B \in \mathbf{M}_n(\mathbb{C})$  and  $P_B(X)$  be the characteristic polynomial of B. Then

$$P_B(X) = \prod_{\lambda \in \operatorname{Sp}B} (X - \lambda)^{n_\lambda},$$

where  $1 \leq n_{\lambda} \in \mathbb{Z}$ . Moreover, the generalised eigenspace of B corresponding to  $\lambda$ , denoted  $E_{\lambda}$ , is defined by

$$E_{\lambda} := \ker(B - \lambda \mathrm{Id}_n)^{n_{\lambda}},$$

where  $\operatorname{Id}_n$  denotes the  $n \times n$ -dimensional identity matrix. If  $J \subseteq \mathbb{R}$  is an interval, then the *J*generalised eigenspace of *B*, denoted  $E_{B,J}$ , is the subspace of  $\mathbb{C}^n$  defined to be the direct sum of the generalised eigenspaces of *B* corresponding to eigenvalues with real parts belonging to *J* (in case there are no eigenvalues with real part belonging to *J*, then  $E_{B,J}$  is defined to be  $\{0\}$ ). Finally, given  $0 < \beta \in \mathbb{R}$ , the *first generalised eigenspace in the*  $\beta$ , *B*-decomposition of  $\mathbb{C}^n$ , denoted  $E_{B,\beta}$ , is defined to be  $E_{B,J_\beta}$ , where  $J_\beta := (\kappa - \beta, \kappa]$  and  $\kappa := \kappa_{\max}(B)$ ; cf. Definition 4.3.

**Remark 4.8.** Our definition of a generalised eigenspace coincides with that of [44, p. 51].

**Remark 4.9.** We discuss the significance of the first generalised eigenspace in greater detail in Section 24.2 below. The reason we use the terminology *first* generalised eigenspace is that we, in Section 9.4 below, divide  $\mathbb{C}^n$  into a direct sum of the first *and second* generalised eigenspaces.

**Remark 4.10.** If  $B \in \mathbf{M}_n(\mathbb{R})$ , then  $\lambda \in \operatorname{Sp}B \Rightarrow \lambda^* \in \operatorname{Sp}B$ , where  $\lambda^*$  denotes the conjugate of  $\lambda$ . Moreover,  $n_{\lambda} = n_{\lambda^*}$ , so that  $v \in E_{\lambda}$  implies  $v^* \in E_{\lambda^*}$ . In particular, given an interval  $J \subseteq \mathbb{R}$ , we can thus choose a basis of  $E_{B,J}$  consisting of elements of  $\mathbb{R}^n$ . In short: in the context of real  $n \times n$ -matrices, we can think of  $E_{B,J}$  as a subspace of  $\mathbb{R}^n$ .

Given this terminology, we have the following result concerning the asymptotics.

**Proposition 4.11.** Consider (1.2). Assume that it is  $C^1$ -silent in the sense of Definition 4.1; that  $\mathcal{X}$  is  $C^0$ -future bounded; and that there are  $\alpha_{\infty}, \zeta_{\infty} \in \mathbf{M}_m(\mathbb{C})$  and  $0 < \eta_{\mathrm{mn}}, C_{\mathrm{mn}} \in \mathbb{R}$  with the property that (4.2) holds for all  $t \geq 0$ . Assume, moreover, that f is a smooth function such that for every  $s \in \mathbb{R}$ , (4.7) holds, where  $\kappa_1 := \kappa_{\mathrm{max}}(A_{\infty})$  and  $A_{\infty}$  is given by (4.4). Let  $\beta_{\mathrm{rem}}$ be given by (4.9) and  $E_a$  be the first generalised eigenspace in the  $\beta_{\mathrm{rem}}, A_{\infty}$ -decomposition of  $\mathbb{C}^{2m}$ . Then there are constants C, N and  $s_{\mathrm{hom}}, s_{\mathrm{ih}} \geq 0$ , depending only on the coefficients of the equation (1.2), such that the following holds. Given a smooth solution u to (1.2), there is a unique  $V_{\infty} \in C^{\infty}(\overline{M}, E_a)$  such that

$$\left\| \begin{pmatrix} u(\cdot,t) \\ u_t(\cdot,t) \end{pmatrix} - e^{A_{\infty}t}V_{\infty} - \int_0^t e^{A_{\infty}(t-\tau)} \begin{pmatrix} 0 \\ f(\cdot,\tau) \end{pmatrix} d\tau \right\|_{(s)}$$

$$\leq C \langle t \rangle^N e^{(\kappa_1 - \beta_{\text{rem}})t} \left( \|u_t(\cdot,0)\|_{(s+s_{\text{hom}})} + \|u(\cdot,0)\|_{(s+s_{\text{hom}}+1)} + \|f\|_{A,s+s_{\text{ih}}} \right)$$

$$(4.10)$$

holds for  $t \geq 0$  and all  $s \in \mathbb{R}$ . Moreover,

$$\|V_{\infty}\|_{(s)} \le C\left(\|u_t(\cdot,0)\|_{(s+s_{\text{hom}})} + \|u(\cdot,0)\|_{(s+s_{\text{hom}}+1)} + \|f\|_{A,s+s_{\text{ih}}}\right).$$
(4.11)

Remark 4.12. The proposition is a consequence of Lemma 10.17; cf. Remark 10.18.

**Remark 4.13.** If, in addition to the assumptions of the proposition, there are constants  $K_{der}$ ,  $\beta_{der} > 0$  such that

$$|\chi|_{\bar{g}} + |\mathcal{X}|_{\bar{h}} \le K_{\mathrm{der}} e^{-\beta_{\mathrm{der}} t} \tag{4.12}$$

for all  $t \ge 0$ , then  $\beta_{\text{rem}}$  can be replaced by  $\min\{2\mu, \mu + \beta_{\text{der}}, \eta_{\text{mn}}\}$ . This follows from Remark 10.21.

**Remark 4.14.** If  $\alpha_{\infty}, \zeta_{\infty} \in \mathbf{M}_m(\mathbb{R})$ , then  $E_a$  can be thought of as a subspace of  $\mathbb{R}^{2m}$  due to Remark 4.10

**Remarks 4.15.** Due to the definition of  $E_a$ , the function  $V_{\infty}$  is uniquely determined by the estimate (4.10). Viewing  $V_{\infty}$  as asymptotic data, the estimate (4.11) can be interpreted as saying that the map from initial data to asymptotic data is continuous with respect to the  $C^{\infty}$ -topology. Finally, note that

$$e^{A_{\infty}t}V_{\infty} + \int_0^t e^{A_{\infty}(t-\tau)} \begin{pmatrix} 0\\ f(\cdot,\tau) \end{pmatrix} d\tau$$

is a solution to (4.3).

#### 4.3. EXAMPLES

Proposition 4.11 yields asymptotic data, given a solution. However, it is also of interest to specify the asymptotics. Due to Proposition 4.11, it is sufficient to demonstrate that asymptotic data can be specified in the case of homogeneous equations. For that reason, we focus on the homogeneous setting for the remainder of the section; i.e., we assume f = 0 in (1.2). In order to obtain conclusions, we need to strengthen the assumptions of the previous two propositions. In particular, we need to assume an upper bound on  $\bar{k}$ , analogous to the lower bound of Definition 4.1. The reason for this is that we need to control how the energy evolves when going backwards in time, and an upper bound on  $\bar{k}$  excludes pathologically fast growth.

**Proposition 4.16.** Consider (1.2) with f = 0. Assume that (1.2) is  $C^1$ -silent in the sense of Definition 4.1; that  $\mathcal{X}$  is  $C^0$ -future bounded; and that there are  $\alpha_{\infty}, \zeta_{\infty} \in \mathbf{M}_m(\mathbb{C})$  and  $0 < \eta_{\mathrm{mn}}, C_{\mathrm{mn}} \in \mathbb{R}$  with the property that (4.2) holds for all  $t \ge 0$ . Assume, moreover, that there is a constant  $0 < \mu_+ \in \mathbb{R}$  and a non-negative continuous function  $\mathbf{e}_+ \in L^1([0,\infty))$  such that

$$\bar{k} \le (\mu_+ + \mathfrak{e}_+)\bar{g} \tag{4.13}$$

for all  $t \geq 0$ . Let  $\beta_{\text{rem}}$  be given by (4.9) and  $E_a$  be the first generalised eigenspace in the  $\beta_{\text{rem}}$ ,  $A_{\infty}$ -decomposition of  $\mathbb{C}^{2m}$ . Then there is an injective map

$$\Phi_{\infty}: C^{\infty}(\bar{M}, E_a) \to C^{\infty}(\bar{M}, \mathbb{C}^{2m})$$

such that the following holds. First,

$$\|\Phi_{\infty}(\psi)\|_{(s)} \le C \|\psi\|_{(s+s_{\infty})} \tag{4.14}$$

for all  $s \in \mathbb{R}$  and all  $\psi \in C^{\infty}(\overline{M}, E_a)$ , where the constants C and  $s_{\infty} \geq 0$  only depend on the coefficients of the equation (1.2). Second, if  $\psi \in C^{\infty}(\overline{M}, E_a)$  and u is the solution to (1.2) (with f = 0) such that

$$\begin{pmatrix} u(\cdot,0)\\ u_t(\cdot,0) \end{pmatrix} = \Phi_{\infty}(\psi), \tag{4.15}$$

then

$$\left\| \begin{pmatrix} u(\cdot,t) \\ u_t(\cdot,t) \end{pmatrix} - e^{A_{\infty}t} \psi \right\|_{(s)}$$

$$\leq C \langle t \rangle^N e^{(\kappa_1 - \beta_{\text{rem}})t} \left( \| u_t(\cdot,0) \|_{(s+s_{\text{hom}})} + \| u(\cdot,0) \|_{(s+s_{\text{hom}}+1)} \right)$$

$$(4.16)$$

for all  $t \ge 0$  and  $s \in \mathbb{R}$ , where the constants C, N and  $s_{\text{hom}}$  have the same dependence as in the case of Proposition 4.11,  $\kappa_1 := \kappa_{\max}(A_{\infty})$  and  $A_{\infty}$  is given by (4.4). Finally, if  $E_a = \mathbb{C}^{2m}$ , then  $\Phi_{\infty}$  is surjective.

**Remark 4.17.** The proposition is a consequence of Lemma 10.22; cf. Remark 10.25.

**Remark 4.18.** If, in addition to the assumptions of the proposition, there are constants  $K_{\text{der}}$ ,  $\beta_{\text{der}} > 0$  such that (4.12) holds for all  $t \ge 0$ , then  $\beta_{\text{rem}}$  can be replaced by  $\min\{2\mu, \mu + \beta_{\text{der}}, \eta_{\text{mn}}\}$ . This follows from Remark 10.27.

**Remarks 4.19.** By combining (4.14), (4.15) and (4.16), the Sobolev norms of  $u(\cdot, 0)$  and  $u_t(\cdot, 0)$  appearing on the right hand side of (4.16) can be replaced by a suitable Sobolev norm of  $\psi$ . The estimate (4.14) demonstrates that the map from asymptotic data to initial data is continuous with respect to the  $C^{\infty}$ -topology. Combining Propositions 4.11 and 4.16 yields the conclusion that when  $E_a = \mathbb{C}^{2m}$ , the map from initial data to asymptotic data is a homeomorphism in the  $C^{\infty}$ -topology. Note also that  $\text{Rsp}A_{\infty} < \beta_{\text{rem}}$  implies that  $E_a = \mathbb{C}^{2m}$ .

## 4.3 Examples

In order to illustrate the results, let us consider two simple examples of scalar equations.

**Example 4.20.** Consider (1.23), the relevant equation when studying the singularity in polarised  $\mathbb{T}^3$ -Gowdy. In this case,

$$g = -dt \otimes dt + e^{2t} d\theta \otimes d\theta, \quad \bar{g} = e^{2t} d\theta \otimes d\theta, \quad \bar{k} = \bar{g}, \quad \chi = 0.$$

Moreover,  $\mathcal{X} = 0$ ,  $\alpha = 0$  and  $\zeta = 0$ . It is thus obvious that the equation (1.23) is  $C^1$ -silent with  $\mu = 1$ . In addition, (4.12) holds with  $K_{der} = 1$  and  $\beta_{der} = 1$ . Finally, since  $\zeta = \alpha = 0$ , the matrix  $A_{\infty}$ , introduced in (4.4), is given by

$$A_{\infty} = \left(\begin{array}{cc} 0 & 1\\ 0 & 0 \end{array}\right). \tag{4.17}$$

Thus  $\kappa_1 = 0$ . Note that (4.2) holds with, e.g.,  $\eta_{mn} = 2$  and  $C_{mn} = 1$ . Since  $\beta_{der} = 1$  we can equate  $\beta_{rem}$  with 2; cf. Remarks 4.13 and 4.18. Finally, since  $A_{\infty}$  has only one eigenvalue, the spaces  $E_a$  appearing in the statements of Propositions 4.11 and 4.16 equal  $\mathbb{C}^2$ . To conclude, the assumptions of all the propositions stated in Section 4.2 are fulfilled. Appealing to Propositions 4.11 and 4.16 yields a homeomorphism between initial data to (1.23) and asymptotic data, consisting of  $\psi \in C^{\infty}(\mathbb{S}^1, \mathbb{C}^2)$ . If  $\psi = (\psi_{\infty} \ v_{\infty})^t$ , (4.16) yields

$$\left\| \begin{pmatrix} P(\cdot,t) \\ P_t(\cdot,t) \end{pmatrix} - \begin{pmatrix} v_{\infty}t + \psi_{\infty} \\ v_{\infty} \end{pmatrix} \right\|_{(s)} \le C_s \langle t \rangle^N e^{-2t},$$

where  $C_s$  in this case is allowed to depend on s and the solution. These asymptotics should be compared with (1.24).

**Example 4.21** (The non-flat Kasner solutions). Let us consider the non-flat Kasner solutions in the direction of the singularity. In other words, let us consider a metric g of the form (1.9), where  $a_i(t) = t^{p_i}$  and the  $p_i$  are constants satisfying

$$\sum_{i=1}^{d} p_i = \sum_{i=1}^{d} p_i^2 = 1, \quad p_i < 1.$$

Then

$$\Box_g u = \frac{1}{\sqrt{-\det g}} \partial_\alpha \left( g^{\alpha\beta} \sqrt{-\det g} \ \partial_\beta u \right) = -\frac{1}{t} \partial_t (tu_t) + \sum_{i=1}^d t^{-2p_i} u_{ii}.$$

Changing time coordinate to  $\tau$ , where  $t = e^{-\tau}$ , the Klein-Gordon equation  $\Box_g u - m^2 u = 0$ , where m is a constant, can thus be written

$$-u_{\tau\tau} + \sum_{i=1}^{d} e^{-2\beta_i \tau} u_{ii} - m^2 e^{-2\tau} u = 0, \qquad (4.18)$$

where  $\beta_i = 1 - p_i > 0$ . The metric associated with (4.18) is given by

$$g_{\rm con} = -d\tau \otimes d\tau + \sum_{i=1}^d e^{2\beta_i \tau} dx^i \otimes dx^i$$
(4.19)

on  $\mathbb{T}^d \times \mathbb{R}$ . Let  $\bar{g}$  be the metric and  $\bar{k}$  be the second fundamental form induced on  $\mathbb{T}^d \times \{\tau\}$  by  $g_{\text{con}}$ . Then, beyond the metric (4.19), the functions characterising (4.18) are given by  $\alpha(\tau) = 0$ ,  $\mathcal{X}(\tau) = 0$  and  $\zeta(\tau) = m^2 e^{-2\tau}$ . Moreover,

$$\bar{k} = \sum_{i=1}^{d} \beta_i e^{2\beta_i \tau} dx^i \otimes dx^i;$$

note that  $U = \partial_{\tau}$ . In particular,

$$\bar{k} \ge \min\{\beta_1, \dots, \beta_d\}\bar{g}.$$

Since the shift vector field field  $\chi$  vanishes in the current setting, it is thus clear that (4.18) is  $C^1$ -silent, with  $\mu := \min\{\beta_1, \ldots, \beta_d\} < 1$ ; cf. Definition 4.1. Moreover,  $\mathcal{X}$  is clearly  $C^0$ -future bounded, and (4.2) holds with  $\alpha_{\infty} = 0$ ,  $\zeta_{\infty} = 0$ ,  $C_{\rm mn} = m^2$  and  $\eta_{\rm mn} = 2$ . Finally, (4.12) holds with  $K_{\rm der} = 1$  and  $\beta_{\rm der} = 1$ . Thus Proposition 4.11 applies, and the relevant matrix  $A_{\infty}$  is in the

present setting given by (4.17). Moreover,  $\kappa_1 = 0$ ,  $\beta_{\text{rem}} = 2\mu$  and  $E_a = \mathbb{C}^2$ . Given a solution u to (4.18), appealing to Proposition 4.11 yields  $u_{\infty}, v_{\infty} \in C^{\infty}(\mathbb{T}^d, \mathbb{C})$  such that

$$\left\| \begin{pmatrix} u(\cdot,\tau) \\ u_{\tau}(\cdot,\tau) \end{pmatrix} - \begin{pmatrix} v_{\infty}\tau + u_{\infty} \\ v_{\infty} \end{pmatrix} \right\|_{(s)} \le C_s \langle t \rangle^N e^{-2\mu\tau}, \tag{4.20}$$

where  $C_s$  in this case is allowed to depend on s and the solution. In fact, appealing to Propositions 4.11 and 4.16 yields the conclusion that the map from initial data to asymptotic data is a homeomorphism with respect to the  $C^{\infty}$ -topology.

It is of interest to note that the mass term in the Klein-Gordon equation does not have any influence on the asymptotics.

# 4.4 Strengths and weaknesses of the results; a rough idea of the argument

#### 4.4.1 Strengths and weaknesses

One clear strength of the results contained in Propositions 4.4, 4.11 and 4.16 is that they apply to a general class of equations. Moreover, the consequences are quite far reaching; in favourable situations, combining Propositions 4.11 and 4.16 yields a homeomorphism between initial data and asymptotic data. Turning to the questions concerning the optimal behaviour of the energy, posed in Section 1.4, note that combining Propositions 4.4 and 4.16 yields the conclusion that  $\eta_{\rm cr} = \kappa_1$ , where  $\kappa_1$  is introduced in the statement of Proposition 4.4. In particular, we can calculate  $\eta_{cr}$ . How about  $\eta_{nl}$ ? Considering (4.6), there is clearly one fundamental obstruction to calculating  $\eta_{nl}$ : the fact that  $\mathfrak{E}_{s+s_0}$  appears on the right hand side (where we do not know whether  $s_0 = 0$  or not). In other words, the estimate (4.6) potentially involves a loss of derivatives which makes it impossible to deduce an upper bound on  $\eta_{nl}$ . On the other hand, Proposition 4.16 yields the lower bound  $\eta_{nl} \geq \kappa_1$ . Below we demonstrate that for any  $\eta_1 \geq \eta_2$ , there are equations satisfying the assumptions of Propositions 4.4, 4.11 and 4.16 such that  $\eta_{cr} \leq \eta_2$  and  $\eta_{nl} \geq \eta_1$ ; cf. Example 7.23 below. It is therefore hard to imagine the perspective taken in the present chapter (where we, generally, accept unspecified derivative losses) to be useful in drawing conclusions concerning  $\eta_{nl}$ . On the other hand, when applying the linear theory in the study of non-linear equations,  $\eta_{nl}$  is of much greater interest than  $\eta_{\rm cr}$ .

### 4.4.2 Outline of the argument

The proofs of Propositions 4.4, 4.11 and 4.16 are not very complicated. They are also quite short; the complete arguments are contained in Chapters 9 and 10. In particular, it is of interest to note that an analysis of the type described in Section 2.5, involving a detailed study of the oscillations, is not necessary. In fact, we do not need to appeal to Part III at all (below we describe the contents of Part III in greater detail, but let us here point out that it is the most important part of these notes in that it allows us to study the oscillations in detail and thereby to derive optimal estimates for the energies, our main goal in these notes).

The Fourier coefficients; dividing the time interval. In order to describe the arguments, it is convenient to use the notation introduced in Section 24.1. Consider, in particular, (24.18), the equation for the Fourier coefficients, where  $\sigma$  and X are defined by (24.17). Due to the assumptions on the shift vector field and on  $\mathcal{X}$ , it can be verified that for all  $t \geq 0$ ,  $|\sigma(\iota, t)|$  and  $||X(\iota, t)||$  are bounded by constants which are independent of  $\iota$ ; this is a consequence of Lemma 25.17 below. In this sense, all the dependence on  $\iota$  in (24.18) enters via the function  $\mathfrak{g}(\iota, t)$ , introduced in (24.13). On the other hand, due to the assumptions concerning the second fundamental form,

for all  $t \geq 0$  and  $\iota \in \mathcal{I}_{\mathrm{B}}$ , where  $C \in \mathbb{R}$  is independent of  $\iota$ ,  $\nu(\iota)$  is defined by (24.4) and we use the notation (4.5); this is a consequence of Lemma 25.12 below. Due to (4.21), it is natural to divide  $[0, \infty)$  into two parts:  $[0, T_{\mathrm{ode}}]$  and  $[T_{\mathrm{ode}}, \infty)$ . The division is, roughly speaking, defined by the condition that  $\mathfrak{g}(\iota, t) \leq 1$  for all  $t \geq T_{\mathrm{ode}}$ ; cf. Definition 10.1 below for the details. Given the definition of  $T_{\mathrm{ode}}$ , it is natural to divide the analysis into two parts. First, we analyse the behaviour in  $[T_{\mathrm{ode}}, \infty)$ , which we refer to as the *ODE-regime*. Second, we analyse the behaviour in the interval  $[0, T_{\mathrm{ode}}]$ , which we refer to as the oscillatory regime.

The ODE regime. For  $t \ge T_{\text{ode}}$ ,  $\mathfrak{g}(\iota, \cdot)$  is not only  $\le 1$ , it is also exponentially decaying. In the interval  $[T_{\text{ode}}, \infty)$ , the equation (24.18) can thus be written in the form

$$\dot{v}(\iota, t) = A_{\infty}v(\iota, t) + A_{\text{rem}}(\iota, t)v(\iota, t) + F(\iota, t)$$
(4.22)

for an appropriate choice of v and F, where  $A_{\infty}$  is given by (4.4). Moreover,  $A_{\text{rem}}$  satisfies an estimate of the form  $||A_{\text{rem}}(\iota, t)|| \leq Ce^{-\beta_{\text{rem}}(t-T_{\text{ode}})}$  for all  $t \geq T_{\text{ode}}$ , where  $C \in \mathbb{R}$  is independent of t and  $\iota$ ; and  $\beta_{\text{rem}}$  is given by (4.9). In Chapter 9, we develop the tools necessary for analysing the behaviour of solutions to equations of the form (4.22). The discussion is divided into the following steps.

Algebraic decompositions, energy estimates. The first step of the analysis is to decompose the equation algebraically in order to isolate the leading order behaviour. This is the subject of Section 9.1. The main idea is very simple, namely to, first, change variables in (4.22) so that  $A_{\infty}$  is transformed into its Jordan normal form; second, to change variables by a time dependent unitary matrix in order to eliminate the imaginary parts of the eigenvalues; third, to multiply the variables by a scalar exponential function to normalise the eigenvalues so that the largest one is zero; and, fourth, to multiply the variables by a diagonal matrix so that the Jordan blocks corresponding to a strictly negative eigenvalue become negative definite (after the final change of variables, the Jordan blocks are non-standard in the sense that the non-zero off-diagonal elements all equal a suitably chosen  $0 < \epsilon \in \mathbb{R}$ , small enough). The end result is that (4.22) is transformed into

$$\dot{w}(\iota,t) = Jw(\iota,t) + B_{\rm rem}(\iota,t)w(\iota,t) + G(\iota,t), \tag{4.23}$$

where J is a real matrix, in non-standard Jordan normal form, such that the Jordan blocks corresponding to negative eigenvalues are negative definite;  $B_{\rm rem}$  satisfies the same type of estimate as  $A_{\rm rem}$ ; and  $|G(\iota, t)| \leq Ce^{-\kappa_1 t} |F(\iota, t)|$ , where  $\kappa_1 := \kappa_{\max}(A_{\infty})$  and C is a constant depending only on  $A_{\infty}$ . Moreover, |w| is equivalent to  $e^{-\kappa_1 t} |v|$ , with constants of equivalence depending only on  $A_{\infty}$ . Based on the reformulation (4.23), we derive an energy estimate for  $t \geq T_{\rm ode}$  in Section 9.2.

Deriving asymptotics. Next, we turn to the problem of deriving asymptotics. This is the subject of Section 9.3. Again, the variables w described above play an important role. However, we can, in general, only expect to obtain partial information concerning the asymptotics of w. In Section 4.2 above, we give a heuristic motivation for the analogous statement in the case of  $C^1$ silent equations. We also motivate the necessity of focusing on the spaces  $E_a$  appearing in the statement of Proposition 4.11. This issue already appears in the study of systems of the form (4.23), and makes it necessary to, beyond the algebraic decompositions already discussed, divide the variables w into one group for which we can expect to be able to distinguish the asymptotic behaviour from the error terms arising from  $B_{\rm rem}$ , say  $w_a$ , and one group for which we cannot, say  $w_b$ . Considering the matrix J appearing in (4.23),  $w_a$  corresponds to the Jordan blocks whose eigenvalues are  $> -\beta_{\rm rem}$ , and  $w_b$  corresponds to the remaining Jordan blocks. Concerning  $w_a$ , we need to calculate an asymptotic expression, and to estimate the error. Concerning  $w_b$ , we only need an estimate. Finally, we need to combine the estimates for the two components, and to reinterpret the result in terms of the original variables. Doing so yields Lemma 9.16.

Specifying the asymptotics. Finally, we want to construct solutions with prescribed asymptotics. When doing so, it is sufficient to consider the case that G = 0 in (4.23). Note also that we can only hope to be able to specify the asymptotics for  $w_a$ . The idea of the argument is to fix a sufficiently

#### 4.4. STRENGTHS/WEAKNESSES; ARGUMENT

late time, say  $t_0 \ge T_{\text{ode}}$ ; to specify initial data at  $t_0$  such that  $w_b(t_0) = 0$ ; and to consider the map taking  $w_a(t_0)$  to the asymptotic data for  $w_a$ . For sufficiently late  $t_0$ , it can be demonstrated that this map is injective. Since the relevant vector spaces have the same dimension, this proves that the map is an isomorphism. Inversion yields a map from asymptotic data to initial data. In the end, we also have to estimate the norm of this map, and to translate the estimates into the original variables. The end result is Lemma 9.21.

The oscillatory regime. In Chapter 10, we first consider the interval  $[0, T_{ode}]$  for one individual mode. Note that as  $\langle \nu(\iota) \rangle \to \infty$ ,  $T_{\rm ode} \to \infty$ . Moreover, under the assumptions of the present chapter, we do not have any detailed control of the growth of the energy of the modes in  $[0, T_{ode}]$ . On the other hand, due to the fact that the equation is balanced, we know that the growth is not faster than exponential; cf. Lemma 10.5. Moreover, the constant appearing in the exponential does not depend on  $\iota$ . Finally, due to (4.21), we know that  $T_{ode} \leq C \ln \langle \nu(\iota) \rangle$ . Combining these observations, the energy associated with a mode can, in the interval  $[0, T_{ode}]$ , be estimated by  $\langle \nu(\iota) \rangle^{s_0}$  times the initial energy, for a suitable constant  $s_0$  depending only on the coefficients of the equation. From this point of view, the growth in the interval  $[0, T_{ode}]$  simply corresponds to a loss of derivatives. Combining the rough estimate given by Lemma 10.5 with the estimate (9.12), valid in the ODE-regime, yields Proposition 4.4 after summing over the modes. In order to obtain Proposition 4.11, it is essentially sufficient to combine the estimates in the oscillatory regime with the ones in the ODE-regime. However, writing down the details turns out to be quite technical. Finally, in order to prove Proposition 4.16, we need an additional assumption. The reason is that in order to take the step from initial data at  $t = T_{ode}$  (obtained from asymptotic data by the analysis in the ODE regime) to initial data at t = 0, we need to estimate the energy of a mode at t = 0 in terms of the energy at  $t = T_{ode}$ . In order for the growth (when going backwards in time from  $T_{\text{ode}}$  to 0) not to be worse than exponential (so that it does not correspond to more than a loss of derivatives), it is not sufficient to have a lower bound on k, we also need an upper bound. This is the reason for demanding that (4.13) hold. However, once we have the necessary estimates for the individual modes, it is sufficient to sum them up (even though the details are somewhat technical). This yields Proposition 4.16.

CHAPTER 4. SILENT EQUATIONS

# Chapter 5

# **Transparent** equations

# 5.1 Introduction

In Chapter 4 above, we present results concerning silent equations. Beyond the bounds on  $\mathcal{X}$  and the shift vector field, and the convergence requirements on  $\alpha$  and  $\zeta$  (which we impose in most of these notes), the main assumption in the silent setting is that (4.1) hold. Disregarding the integrable function  $\mathfrak{e}$ , the first inequality appearing in (4.1) implies that  $\overline{k}$  has a strictly positive lower bound. It is of interest to consider the case that  $\overline{k}$  (up to an integrable function) has a nonnegative lower bound, but that there are some directions in which it degenerates asymptotically. This leads to the study of transparent equations. Returning to Subsection 1.3.1, the Nariai spacetimes, the flat Kasner solutions (in the direction of the singularity) and the Milne model (in the expanding direction) are some examples of relevant geometries.

## 5.1.1 Outline

In the transparent setting, it is necessary to divide the spatial variables into "silent" and "transparent" variables; cf. Subsection 5.1.2 below. Given this division, a formal definition of the above requirements can be given; we demand that  $\bar{k}$  converges to zero exponentially in the transparent directions, cf. (5.3) below, and that there is exponential expansion in the silent directions, cf. (5.4)below. The conditions imposed on the shift vector field are described in Definition 5.5. In part, the assumptions are similar to the ones imposed in the silent setting. However, we here also demand that the shift vector field be  $C^1$ -future bounded, and that its transparent components converge to zero exponentially. Due to these assumptions, we can calculate the leading order behaviour of the coefficients of the highest order derivatives in (1.2); cf. (5.7) and (5.8) below. Concerning  $\alpha$  and  $\zeta$ , we assume that they converge exponentially, as in the silent setting. Turning to  $\mathcal{X}$ , the  $C^0$ -future boundedness is sufficient to ensure that the "silent" components of  $\mathcal{X}$  converge to zero exponentially. In order to control the "transparent" components, we explicitly assume that they converge exponentially; cf. (5.9). Adding up all of the above observations we obtain a limit equation, given by (5.10) below. For fixed silent variables, the limit equation can be thought of as a system of linear wave equations in the transparent variables and the time variable. Moreover, this system has constant coefficients.

The Fourier modes. Turning to the Fourier side, cf. Section 24.1, the situation is very similar to the one considered in the previous chapter. However, there is one crucial difference: as opposed to the silent setting, the coefficients of the limit equation here depend on  $\iota$ . This leads to several complications. First, estimating the distance between two solutions by computing the energy of the difference is not so useful, since the energies of different modes typically exhibit different exponential growth/decay. In order to obtain norms such that bounds on these norms have impli-

cations for all the modes, it is necessary to introduce a time and mode dependent normalisation in the definition of the energies/norms. Second, when describing the asymptotics, we need to introduce spaces analogous to the  $E_a$  defined in the statement of Proposition 4.11. However, in the present context, what  $E_a$  is depends on the mode. Third, the proof proceeds similarly to the argument described in Section 4.4. In particular, we wish to make an appropriate algebraic decomposition of the matrices appearing in the equation on the Fourier side. The problem that arises in the transparent setting is that we, in general, need to decompose infinitely many different matrices. This leads to the problem of obtaining uniform control of the norms of the matrices used to transform the equations into the preferred form; we return to this topic in greater detail in Section 5.6 below.

Dividing the equation into silent and transparent parts. Corresponding to the division of the spatial variables into "silent" and "transparent" variables, there is a division of  $\mathcal{I}_{\rm B}$  into  $\mathcal{I}_{\rm B,sil}$  and  $\mathcal{I}_{\rm B,ts}$ ; cf. Definition 5.8 below. We can project (1.2) to the corresponding subspaces of  $L^2(\bar{M}, \mathbb{C}^m)$  (where we tacitly assume the projection to only be in the spatial variables). The projection to the subspace corresponding to  $\mathcal{I}_{\rm B,sil}$ , say  $L^2_{\rm sil}(\bar{M}, \mathbb{C}^m)$ , yields an equation to which the results of the previous chapter apply. We therefore focus on the projection to the subspace corresponding to  $\mathcal{I}_{\rm B,ts}$ , say  $L^2_{\rm ts}(\bar{M}, \mathbb{C}^m)$ , in the present chapter. We refer to the latter equation as the transparent part of (1.2). If the (smooth) initial data for (1.2) are in  $L^2_{\rm ts}(\bar{M}, \mathbb{C}^m) \times L^2_{\rm ts}(\bar{M}, \mathbb{C}^m)$ , then they are said to be transparent. Finally, given a solution to (1.2), say u, we call the projection of u to  $L^2_{\rm ts}(\bar{M}, \mathbb{C}^m)$  the transparent part of the solution u, and we denote it by  $u_{\rm ts}$ .

**Deriving asymptotics.** Before describing the asymptotics, it is of interest to develop a feeling for the solutions to the limit equation. In Subsection 5.2.1, we therefore solve the transparent part of the limit equation, given transparent initial data. Given a solution u to (1.2), our goal is to describe the asymptotics of  $u_{ts}$ . We therefore wish to find transparent initial data such that the corresponding solution to the transparent part of the limit equation, say  $v_{ts}$ , is as "close" to  $u_{ts}$  as possible. The relevant result is formulated as Proposition 5.11 below. Note that it is analogous to Proposition 4.11 above, with the following two main differences. First, the vector spaces to which the asymptotic data belong depend on  $\iota$ , for reasons mentioned above. Second, in order for estimates of the difference between  $u_{ts}$  and  $v_{ts}$  to uniquely determine the asymptotic data, we, in the definition of the norm, need to introduce a time and  $\iota$  dependent normalising factor for each mode. The definition of the norm is given in (5.22) below, and the main estimate of the difference of  $u_{ts}$  and  $v_{ts}$  is given by (5.24) below. As in the silent setting, the map from initial data to asymptotic data is continuous with respect to the  $C^{\infty}$ -topology; cf. (5.25) below.

**Specifying asymptotics, examples and outline.** In Section 5.4 below, we turn to the question of specifying the asymptotics. With the exception of the differences described above, the relevant statement is analogous to Proposition 4.16, the corresponding statement in the silent setting. In particular, we obtain a continuous map from asymptotic data to initial data. In order to illustrate the results, we give a simple example in Section 5.5 below. We end the chapter by giving an outline of the argument in Section 5.6.

#### 5.1.2 Dividing the variables

Before stating the requirements on the second fundamental form, we need to specify in which directions there is expansion and in which directions there is not. This division can be made in the tangent space, or in the cotangent space. The reason for this is that for each hypersurface  $\overline{M}_t$ , the metric  $\overline{g}$  induces an isomorphism between  $T\overline{M}$  and  $T^*\overline{M}$ . We denote this isomorphism by  $\flat$ , and the inverse by  $\sharp$ . In particular,  $(T^*\overline{M})^{\sharp} = T\overline{M}$ . Clearly, the isomorphisms are time dependent, but we omit explicit reference to this dependence in the notation. Given a division into directions, either on the tangent or the cotangent side, the relevant isomorphism, i.e.  $\flat$  or  $\sharp$  respectively, yields a division on the other side. In practice, it turns out to be more natural to make the division on the cotangent side. Later, when we impose requirements on the second fundamental form, it is convenient to state the conditions in terms of the symmetric contravariant

2-tensor fields  $\bar{g}^{\sharp}$  and  $\bar{k}^{\sharp}$ . They are defined by the requirement that their components be given by  $\bar{g}^{ij}$  and  $\bar{k}^{ij}$  respectively. An equivalent, geometric, definition is

$$\bar{g}^{\sharp}(\eta,\xi) := \bar{g}(\eta^{\sharp},\xi^{\sharp}), \quad \bar{k}^{\sharp}(\eta,\xi) := \bar{k}(\eta^{\sharp},\xi^{\sharp})$$

for all  $\eta, \xi \in T^*_{\bar{p}}\bar{M}$  and all  $\bar{p} \in \bar{M}$ .

The following definition yields the relevant division both in the tangent and in the cotangent space.

**Definition 5.1.** Let (M, g) be a canonical separable cosmological model manifold. Assume that a division of the set  $\{1, \ldots, d\}$  into two disjoint sets  $\{j_1, \ldots, j_{d_{ts}}\}$  and  $\{\bar{j}_1, \ldots, \bar{j}_{d_{sil}}\}$  is given (if  $d_{ts} = 0$  or  $d_{sil} = 0$  the corresponding set is empty). These subsets are referred to as the *transparent* and *silent* subsets of  $\{1, \ldots, d\}$  respectively. Assume, moreover, that a division of the set  $\{1, \ldots, R\}$  into the union of two disjoint sets  $\{r_1, \ldots, r_{R_{ts}}\}$  and  $\{\bar{r}_1, \ldots, \bar{r}_{R_{sil}}\}$  is given (if  $R_{ts} = 0$  or  $R_{sil} = 0$  the corresponding set is empty). Again, these subsets are referred to as the *transparent* and *silent* subsets of  $\{1, \ldots, R\}$  respectively. Finally, assume that  $d_{ts} + R_{ts} > 0$ . Given this division, define

$$\bar{M}_{\rm ts} := \mathbb{T}^{d_{\rm ts}} \times M_{r_1} \times \dots \times M_{r_{R_{\rm ts}}}, \quad \bar{M}_{\rm sil} := \mathbb{T}^{d_{\rm sil}} \times M_{\bar{r}_1} \times \dots \times M_{\bar{r}_{R_{\rm sil}}}.$$

These manifolds are referred to as the *transparent* and *silent* manifolds respectively. Given these manifolds, there are associated projections  $\pi_{ts}: \bar{M} \to \bar{M}_{ts}$  and  $\pi_{sil}: \bar{M} \to \bar{M}_{sil}$ , given by

$$\begin{aligned} \pi_{\rm ts}(x_1, \dots, x_d, p_1, \dots, p_R) &:= (x_{j_1}, \dots, x_{j_{d_{\rm ts}}}, p_{r_1}, \dots, p_{r_{R_{\rm ts}}}), \\ \pi_{\rm sil}(x_1, \dots, x_d, p_1, \dots, p_R) &:= (x_{\bar{j}_1}, \dots, x_{\bar{j}_{d_{\rm sil}}}, p_{\bar{r}_1}, \dots, p_{\bar{r}_{R_{\rm sil}}}), \end{aligned}$$

where  $x_j \in \mathbb{S}^1$ ,  $j \in \{1, \ldots, d\}$ , and  $p_r \in M_r$ ,  $r \in \{1, \ldots, R\}$ . These projections induce maps  $\pi_{ts}^* : T^* \overline{M}_{ts} \to T^* \overline{M}$  and  $\pi_{sil}^* : T^* \overline{M}_{sil} \to T^* \overline{M}$ . Let

$$T_{\rm ts}^* \bar{M} := \pi_{\rm ts}^* (T^* \bar{M}_{\rm ts}), \quad T_{\rm sil}^* \bar{M} := \pi_{\rm sil}^* (T^* \bar{M}_{\rm sil}).$$

These sets are referred to as the *transparent* and *silent* subsets of  $T^*\overline{M}$  respectively. Given  $t \ge 0$ , let

$$T_{\mathrm{ts},t}\bar{M} := (T_{\mathrm{ts}}^*\bar{M})^\sharp, \quad T_{\mathrm{sil},t}\bar{M} := (T_{\mathrm{sil}}^*\bar{M})^\sharp,$$

where the operator  $\sharp$  is introduced prior to the statement of the definition. These sets are referred to as the *t*-transparent and *t*-silent subsets of  $T\bar{M}$  respectively.

# 5.1.3 Conditions on the second fundamental form and the shift vector field

Given the above division of the cotangent space into silent and transparent subsets, we are in a position to introduce the relevant conditions on the second fundamental form.

**Definition 5.2.** Let (M, g) be a canonical separable cosmological model manifold. Assume that there is a division of  $\{1, \ldots, d\}$  and  $\{1, \ldots, R\}$  into transparent and silent subsets, as described in Definition 5.1. Assume, moreover, that there are constants  $\eta_{ts} > 0$  and  $C_a > 0$  such that

$$|\bar{k}^{\sharp}(\xi,\xi)| \le C_a e^{-\eta_{\rm ts} t} \bar{g}^{\sharp}(\xi,\xi) \tag{5.1}$$

for all  $\xi \in T^*_{ts}\bar{M}$  and  $t \ge 0$ , where  $T^*_{ts}\bar{M}$  is introduced in Definition 5.1. Then  $\bar{g}^{\sharp}$  is said to be convergent on  $T^*_{ts}\bar{M}$ . Assume that there is a  $\beta_{sil} > 0$  and a continuous non-negative function  $\mathfrak{e}_{sil} \in L^1([0,\infty))$  such that

$$\bar{k}^{\sharp}(\xi,\xi) \ge [\beta_{\rm sil} - \mathfrak{e}_{\rm sil}(t)]\bar{g}^{\sharp}(\xi,\xi) \tag{5.2}$$

for all  $\xi \in T^*_{\text{sil}}\bar{M}$  and  $t \ge 0$ , where  $T^*_{\text{sil}}\bar{M}$  is introduced in Definition 5.1. Then  $\bar{g}^{\sharp}$  is said to be expanding on  $T^*_{\text{sil}}\bar{M}$ .

**Remark 5.3.** The notation  $\bar{g}^{\sharp}$  and  $\bar{k}^{\sharp}$  is introduced at the beginning of Subsection 5.1.2.

**Remark 5.4.** The condition (5.1) is equivalent to the requirement that

$$|\bar{k}(v,v)| \le C_a e^{-\eta_{\rm ts} t} \bar{g}(v,v) \tag{5.3}$$

for all  $v \in T_{ts,t}\overline{M}$  and  $t \ge 0$ , where  $T_{ts,t}\overline{M}$  is introduced in Definition 5.1. Similarly, the condition (5.2) is equivalent to the requirement that

$$\bar{k}(v,v) \ge [\beta_{\rm sil} - \mathfrak{e}_{\rm sil}(t)]\bar{g}(v,v) \tag{5.4}$$

for all  $v \in T_{\text{sil},t}\overline{M}$  and  $t \ge 0$ , where  $T_{\text{sil},t}\overline{M}$  is introduced in Definition 5.1.

In order to be able to draw conclusions, we also need to make assumptions concerning the shift vector field  $\chi$ .

**Definition 5.5.** Let (M, g) be a canonical separable cosmological model manifold. Assume that there is a division of  $\{1, \ldots, d\}$  and  $\{1, \ldots, R\}$  into transparent and silent subsets, as described in Definition 5.1. Assume, in addition, that there is a continuous non-negative function  $\mathfrak{e} \in L^1([0, \infty))$  such that

$$|\chi|_{\bar{g}} \cdot |\dot{\chi}|_{\bar{g}} \le \mathfrak{e} \tag{5.5}$$

holds for all  $t \ge 0$ , and that the shift vector field is  $C^1$ -future bounded; cf. Definition 3.1. Assume, finally, that there are  $C_b > 0$  and  $\eta_{ts} > 0$  such that

$$|\xi(\chi)| \le C_b e^{-\eta_{\rm ts} t} |\xi|_{\bar{q}} \tag{5.6}$$

for all  $\xi \in T_{ts}^* \overline{M}$  and  $t \ge 0$ , where  $T_{ts}^* \overline{M}$  is introduced in Definition 5.1. Then the shift vector field of g is said to be asymptotically negligible.

**Remark 5.6.** The estimate (5.6) corresponds to imposing exponential decay on the shift vector in the transparent directions. This is of course a weaker condition than imposing exponential decay in all directions.

**Remark 5.7.** The constant  $\eta_{ts}$  appears in both (5.1) and (5.6). In situations where we assume both estimates to hold, we tacitly assume the constants  $\eta_{ts}$  appearing in these estimates to coincide.

#### 5.1.4 Consequences for the coefficients of the equations

The highest order coefficients. Consider (1.2). Assume the associated metric to be such that (M, g) is a canonical separable cosmological model manifold. Assume, moreover, that there is a division of  $\{1, \ldots, d\}$  and  $\{1, \ldots, R\}$  into transparent and silent subsets, as described in Definition 5.1. Assume, finally, that Definitions 5.2 and 5.5 are fulfilled. Then the leading order coefficients of (1.2) have the following properties. First, there is a constant C such that

$$|g^{\bar{j}_k\bar{j}_l}(t)| + a_{\bar{r}_j}^{-2}(t) \le Ce^{-2\beta_{\rm sil}t}, \quad |g^{\bar{j}_kj_l}(t)| + |g^{0\bar{j}_k}(t)| \le Ce^{-\beta_{\rm sil}t}$$
(5.7)

for all  $t \ge 0$ , all  $k, l \in \{1, \ldots, d_{\rm sil}\}$ ,  $i \in \{1, \ldots, d_{\rm ts}\}$  and  $j \in \{1, \ldots, R_{\rm sil}\}$ . Moreover, for  $k, l \in \{1, \ldots, d_{\rm ts}\}$  and  $j \in \{1, \ldots, R_{\rm ts}\}$ , there are constants  $g_{\infty, r_j}^{j_k j_l}$  and  $q_{\infty, r_j}$  such that

$$|g_{\infty}^{j_k j_l} - g_{\infty}^{j_k j_l}| + |a_{r_j}^{-2}(t) - q_{\infty, r_j}| + |g^{0j_k}(t)| \le Ce^{-\eta_{ts}t}$$
(5.8)

for all  $t \ge 0$ . Finally,  $q_{\infty,r_j} > 0$  and  $g_{\infty}^{j_k j_l}$  are the components of a positive definite matrix. The justification of the above statements is to be found in Lemma 26.2 below.

The lower order coefficients. In order to obtain control of the coefficients of the lower order derivatives in (1.2), we need to make additional assumptions. To begin with, we assume that  $\mathcal{X}$  is  $C^0$ -future bounded and that there are  $\alpha_{\infty}, \zeta_{\infty} \in \mathbf{M}_m(\mathbb{C})$  and  $0 < \eta_{\mathrm{mn}}, C_{\mathrm{mn}} \in \mathbb{R}$  with the property

47

that (4.2) holds for all  $t \geq 0$ . Finally, assume that there are  $X^{j_k}_{\infty} \in \mathbf{M}_m(\mathbb{C})$  and a constant  $0 < K_{ts} \in \mathbb{R}$  such that

$$\|X^{j_k}(t) - X^{j_k}_{\infty}\| \le K_{\rm ts} e^{-\eta_{\rm ts} t} \tag{5.9}$$

for all  $t \ge 0$  and all  $k \in \{1, \ldots, d_{ts}\}$  (an observation similar to Remark 5.7 is equally relevant in the case of the estimate (5.9)). Under these circumstances, there is a constant C such that

$$\|X^{\overline{j}_k}(t)\| \le Ce^{-\beta_{\rm sil}t}$$

for all  $k \in \{1, \ldots, d_{sil}\}$  and  $t \ge 0$ ; cf. Remark 26.4.

The limit equation. Combining the above assumptions, it is natural to associate the following *limit equation* with (1.2):

$$v_{tt} - \sum_{k,l=1}^{d_{ts}} g_{\infty}^{j_k j_l} \partial_{j_l} v - \sum_{j=1}^{R_{ts}} q_{\infty,r_j} \Delta_{g_{r_j}} v + \alpha_{\infty} v_t + \sum_{k=1}^{d_{ts}} X_{\infty}^{j_k} \partial_{j_k} v + \zeta_{\infty} v = f.$$
(5.10)

Note that this is a constant coefficient equation. Moreover, we can think of it as yielding a system of linear wave equations on  $\overline{M}_{ts} \times I$  for each  $\overline{x} \in \overline{M}_{sil}$ . It is natural to compare the asymptotic behaviour of solutions to (1.2) with that of solutions to (5.10). Before doing so, it is of interest to consider the limit equation in its own right. We turn to this subject next.

# 5.2 The limit equation

In the analysis to follow, it is important to keep track of on which variables the solution depends. In fact, if (1.2) is an equation satisfying the conditions stated in the previous section with f = 0, and if u is a solution arising from initial data depending only on the variables corresponding to  $\overline{M}_{\rm sil}$ , then u is effectively a solution to (1.2) where  $g^{j_k\xi}$ ,  $X^{j_k}$  and  $a_{r_j}^{-2}$  have been set to zero for all  $k \in \{1, \ldots, d_{\rm ts}\}, \xi \in \{0, \ldots, d\}$  and  $j \in \{1, \ldots, R_{\rm ts}\}$ ; we say that the solution only depends on the silent variables and time. Moreover, the conclusions of Propositions 4.4 and 4.11 apply to the corresponding class of solutions; cf. Remark 5.10 below. If, on the other hand, the initial data depend on some of the variables corresponding to  $\overline{M}_{\rm ts}$ , then the asymptotic behaviour can be expected to be different. In order to distinguish the cases that have already been handled from what remains to be done, it is therefore convenient to introduce the following terminology.

**Definition 5.8.** Let (M, g) be a canonical separable cosmological model manifold. Assume that the variables can be divided as described in Definition 5.1. Given  $\iota \in \mathcal{I}_{B}$ , define  $\iota_{ts} \in \mathcal{I}_{B}$  as follows: replace  $n_{\bar{j}_{k}}$  in  $\iota$  by zero for  $k = 1, \ldots, d_{sil}$ ; and replace  $i_{\bar{r}_{k}}$  by 0 for  $k = 1, \ldots, R_{sil}$ . Given  $\iota \in \mathcal{I}_{B}$ , define  $\iota_{sil} \in \mathcal{I}_{B}$  by  $\iota_{sil} := \iota - \iota_{ts}$ . The set of  $\iota \in \mathcal{I}_{B}$  such that  $\iota_{ts} = 0$  ( $\iota_{ts} \neq 0$ ) is denoted by  $\mathcal{I}_{B,sil}$ ( $\mathcal{I}_{B,ts}$ ).

**Remark 5.9.** It is important to keep in mind that the set of  $\iota_{ts}$  for  $\iota \in \mathcal{I}_B$  is in general different from  $\mathcal{I}_{B,ts}$ , but that the set of  $\iota_{sil}$  for  $\iota \in \mathcal{I}_B$  equals  $\mathcal{I}_{B,sil}$ .

**Remark 5.10.** Let (1.2) be an equation satisfying the conditions stated in the previous section with f = 0. Then the conclusions of Propositions 4.4 and 4.11 hold for the class of solutions depending only on the silent variables and time. The reason for this is that, due to Lemma 26.2, in particular (26.4); Lemma 25.17; and (4.2); the conditions of Definition 10.1 are satisfied. In particular, Lemmas 10.11 and 10.17 thus apply. Finally, assuming  $\bar{k}$  to be  $C^0$ -future bounded and appealing to Lemma 25.11, we conclude that Lemma 10.22 applies as well. Thus the conclusions of Proposition 4.16 hold.

#### 5.2.1 Solving the limit equation

Let us consider the limit equation (5.10). Decomposing it into modes as described in Subsection 24.1.3 yields

$$\ddot{z}(\iota, t) + \mathfrak{g}_{\infty}^{2}(\iota)z(\iota, t) + \alpha_{\infty}\dot{z}(\iota, t) + \sum_{l=1}^{d_{\mathrm{ts}}} in_{j_{l}} X_{j_{l}}^{j_{l}} z(\iota, t) + \zeta_{\infty} z(\iota, t) = \hat{f}(\iota, t),$$
(5.11)

where, applying Einstein's summation convention to  $j_k$  and  $j_l$ ,

$$\mathfrak{g}_{\infty}(\iota) = \lim_{t \to \infty} \mathfrak{g}(\iota, t) = \left( g_{\infty}^{j_k j_l} n_{j_k} n_{j_l} + \sum_{j=1}^{R_{\rm ts}} q_{\infty, r_j} \nu_{r_j, i_{r_j}}^2 \right)^{1/2}.$$
 (5.12)

Assuming that  $\iota \in \mathcal{I}_{B,ts}$  (or, equivalently, that  $\mathfrak{g}_{\infty}(\iota) \neq 0$ ), (5.11) can be written

$$\partial_t w(\iota, t) = A(\iota)w(\iota, t) + F(\iota, t), \tag{5.13}$$

where

$$A(\iota) := \begin{pmatrix} 0 & \mathfrak{g}_{\infty}(\iota) \mathrm{Id}_{m} \\ -\mathfrak{g}_{\infty}(\iota) \mathrm{Id}_{m} - i[\mathfrak{g}_{\infty}(\iota)]^{-1} n_{j_{l}} X_{\infty}^{j_{l}} - [\mathfrak{g}_{\infty}(\iota)]^{-1} \zeta_{\infty} & -\alpha_{\infty} \end{pmatrix}.$$
 (5.14)

Moreover,

$$w(\iota,t) := \begin{pmatrix} \mathfrak{g}_{\infty}(\iota)z(\iota,t) \\ \dot{z}(\iota,t) \end{pmatrix}, \quad F(\iota,t) := \begin{pmatrix} 0 \\ \hat{f}(\iota,t) \end{pmatrix}.$$
(5.15)

The solutions to (5.13) can be written

$$w(\iota, t) = e^{A(\iota)t}w(\iota, 0) + \int_0^t e^{A(\iota)(t-\tau)}F(\iota, \tau)d\tau.$$
(5.16)

Returning to (5.10), let  $f_{ts}$  denote the function obtained from f by setting  $\hat{f}(\iota_{sil}, t)$  to zero for all  $t \in I$  and all  $\iota \in \mathcal{I}_B$ . Define  $v_{ts}$  similarly and introduce

$$V_{\rm ts}(\cdot,t) := \begin{pmatrix} (-\Delta_{\infty})^{1/2} v_{\rm ts}(\cdot,t) \\ \partial_t v_{\rm ts}(\cdot,t) \end{pmatrix}.$$
(5.17)

Here  $\Delta_{\infty}$  is the differential operator given by

$$\Delta_{\infty} := g_{\infty}^{j_k j_l} \partial_{j_k} \partial_{j_l} + \sum_{j=1}^{R_{\rm ts}} q_{\infty, r_j} \Delta_{g_{r_j}}.$$
(5.18)

Moreover, if  $\psi \in C^{\infty}(\overline{M}, \mathbb{C}^k)$  is such that  $\hat{\psi}(\iota) = 0$  for all  $\iota \in \mathcal{I}_{B,sil}$  and  $a \in \mathbb{R}$ , then  $(-\Delta_{\infty})^a \psi$  is the function whose  $\iota$ 'th Fourier coefficient is given by 0 in case  $\iota \in \mathcal{I}_{B,sil}$  and by  $[\mathfrak{g}_{\infty}(\iota)]^{2a}\hat{\psi}(\iota)$  otherwise. Then (5.16) can formally be summarised into

$$V_{\rm ts}(\cdot,t) = e^{\mathcal{A}_{\rm ts}t} V_{\rm ts}(\cdot,0) + \int_0^t e^{\mathcal{A}_{\rm ts}(t-\tau)} F_{\rm ts}(\cdot,\tau) d\tau, \qquad (5.19)$$

where

$$F_{\rm ts}(\cdot,t) := \begin{pmatrix} 0 \\ f_{\rm ts}(\cdot,t) \end{pmatrix},$$
$$\mathcal{A}_{\rm ts} := \begin{pmatrix} 0 & \operatorname{Id}_m(-\Delta_{\infty})^{1/2} \\ -\operatorname{Id}_m(-\Delta_{\infty})^{1/2} - (X^{j_l}_{\infty}\partial_{j_l} + \zeta_{\infty})(-\Delta_{\infty})^{-1/2} & -\alpha_{\infty} \end{pmatrix}.$$
(5.20)

#### 5.2.2 Describing the asymptotics

Let (1.2) be an equation satisfying the assumptions of Section 5.1. If u is a corresponding solution, we can divide u into  $u = u_{ts} + u_{sil}$ , where  $u_{ts}$  is defined as above. Assuming f to satisfy the appropriate conditions,  $u_{sil}$  solves an equation such that Propositions 4.4 and 4.11 apply; cf. Remark 5.10. We therefore focus on  $u_{ts}$ . When describing the asymptotics of  $u_{ts}$ , it is convenient to define  $U_{ts}$  to be the right hand side of (5.17) with  $v_{ts}$  replaced by  $u_{ts}$ . Keeping (5.19) in mind, it is then of interest to try to find a  $U_{\infty}$  (whose  $\iota$ 'th Fourier coefficient vanishes for  $\iota \in \mathcal{I}_{B,sil}$ ) such that

$$U_{\rm ts}(\cdot,t) - e^{\mathcal{A}_{\rm ts}t} U_{\infty} - \int_0^t e^{\mathcal{A}_{\rm ts}(t-\tau)} F_{\rm ts}(\cdot,\tau) d\tau$$
(5.21)

is small. In practice, we carry out this comparison on the level of the Fourier coefficients. In other words, we compare solutions to (24.12) with solutions to (5.11). This is very similar to the analysis carried out in the case of silent equations. However, there are two important differences.

- Consider the equation (5.11). For  $\iota \in \mathcal{I}_{B,sil}$ , the coefficients of this equation do not depend on  $\iota$ . However, for  $\iota \in \mathcal{I}_{B,ts}$ , they do. In fact, (5.11) corresponds to infinitely many different equations. In particular, as opposed to the silent setting, the asymptotic behaviour depends on the mode. This yields complications when making the statement that (5.21) is small precise.
- Just as in the case of silent equations, we can only expect to be able to obtain conclusions concerning the leading order behaviour of solutions to (5.11); cf. the discussion adjacent to (4.8). Moreover, in order to isolate the leading order behaviour, we need to transform the matrix  $A(\iota)$  appearing in (5.13) in such a way that the leading order part is separated from the remainder. The problem in the current setting is that there are infinitely many different matrices of the form  $A(\iota)$  (as opposed to a single matrix in the silent setting), but we would like to have uniform control in the estimates (independent of  $\iota$ ).

The first problem can be handled by defining an appropriate norm. To be more precise, assume that the maximal growth of solutions to (5.13) with  $F(\iota, t) = 0$ , up to polynomial factors, is  $e^{\kappa_{\iota} t}$ for some  $\kappa_{\iota} \in \mathbb{R}$ . We then, roughly speaking, define a norm in which the  $\iota$ 'th mode is multiplied with  $e^{-\kappa_{\iota} t}$ . That such a norm decays exponentially then gives relevant information for every mode. The second problem is more difficult to deal with. Here we address it by introducing additional restrictions.

Consider the matrix  $A(\iota)$  introduced in (5.14). If  $X_{\infty}^{j_{\ell}} = 0$  for all  $l \in \{1, \ldots, d_{ts}\}$ , then  $A(\iota)$  is such that as  $\mathfrak{g}_{\infty}(\iota) \to \infty$ , there is an asymptotic expansion of the eigenvalues of  $A(\iota)$ ; cf. Chapter 11, in particular Sections 11.2 and 11.3. In addition, for a fixed  $0 < \mu_0 \in \mathbb{R}$ , the set

$$\{A(\iota)|\iota\in\mathcal{I}_{\mathrm{B,ts}}:\mathfrak{g}_{\infty}(\iota)\leq\mu_0\}$$

is finite. Analysing the equation (5.13) for  $\iota \in \mathcal{I}_{B,ts}$  such that  $\mathfrak{g}_{\infty}(\iota) \leq \mu_0$  is therefore not a problem; it is essentially sufficient to appeal to the results of Chapter 4. Similarly, if we allow  $X_{\infty}^{j_k} \neq 0$  for some  $k \in \{1, \ldots, d_{ts}\}$ , but insist that  $d_{ts} = 1$  and  $R_{ts} = 0$ , then the matrices  $A(\iota)$  have the same properties. For that reason, we, from now on, restrict to these two situations.

## 5.3 Future asymptotics

Before stating a result concerning the future asymptotics, we need to give a formal definition of a norm of the type described in Subsection 5.2.2. Given  $\iota \in \mathcal{I}_{B,ts}$ , let  $\kappa_{\iota} := \kappa_{\max}[A(\iota)]$  and  $\delta_{\iota} := d_{\max}[A(\iota), \kappa_{\iota}]$ ; cf. Definition 4.3. Then we expect generic solutions to the homogeneous version of (5.13) to behave as  $\langle t \rangle^{\delta_{\iota}-1} e^{\kappa_{\iota} t}$ . We therefore introduce the norm

$$\|\psi\|_{t,s} := \left(\sum_{\iota \in \mathcal{I}_{\mathrm{B,ts}}} \langle \nu(\iota) \rangle^{2s} \langle t \rangle^{-2\delta_{\iota}+2} e^{-2\kappa_{\iota}t} |\hat{\psi}(\iota)|^2 \right)^{1/2}.$$
(5.22)

1 10

Clearly,  $\|\cdot\|_{t,s}$  is not a norm on  $C^{\infty}(\overline{M}, \mathbb{C}^k)$ . To justify the terminology, we can consider two functions to be equivalent if their difference, say v, is such that  $\hat{v}(\iota) = 0$  for all  $\iota \in \mathcal{I}_{B,ts}$ . Then  $\|\cdot\|_{t,s}$  is a norm on the corresponding vector space of equivalence classes. In the case of a solution to the homogeneous version of the limit equation (5.10), we expect the norm  $\|\cdot\|_{t,s}$  to be bounded for  $t \geq 0$ , but not better. Next, we are interested in situations where the behaviour of solutions is not dominated by the contribution from the inhomogeneity. In the case of silent equations, this assumption led us to the introduction of the norm  $\|\cdot\|_{A,s}$ ; cf. (4.7). Given  $0 < \beta_{mar} \in \mathbb{R}$ , we introduce an analogous norm in the present setting:

$$\|f\|_{\mathrm{ts},s} := \int_0^\infty \left( \sum_{\iota \in \mathcal{I}_{\mathrm{B},\mathrm{ts}}} \langle \nu(\iota) \rangle^{2s} e^{-2(\kappa_\iota - \beta_{\mathrm{mar}})t} |\hat{f}(\iota, t)|^2 \right)^{1/2} dt;$$
(5.23)

in order for  $\|\cdot\|_{\text{ts},s}$  to be a norm, we need to give it an interpretation similar to the one given to  $\|\cdot\|_{t,s}$  above. When using the notation (5.23), the value of the constant  $\beta_{\text{mar}} > 0$  should be clear from the context. The reason for including  $\beta_{\text{mar}}$  in (5.23) is related to the complications described in Subsection 5.2.2. Due to the fact that there are infinitely many matrices of the form  $A(\iota)$ , infinitely many different matrices are needed in order to transform all the  $A(\iota)$  into their preferred form. On the other hand, we need to have uniform bounds on the norms of the transformation matrices and their inverses. Under certain circumstances, the latter requirement is not compatible with obtaining as detailed an algebraic decomposition of the  $A(\iota)$ 's as the decomposition of the matrix A described in Subsection 4.4.2. As a consequence, we are not always able to, in the algebraic decompositions, resolve the real parts of the eigenvalues exactly. This necessitates the margin represented by  $\beta_{\text{mar}}$ .

**Proposition 5.11.** Consider (1.2). Assume the associated metric to be such that (M, g) is a canonical separable cosmological model manifold. Assume that there is a division of  $\{1, \ldots, d\}$  and  $\{1, \ldots, R\}$  into transparent and silent subsets, as described in Definition 5.1. Assume, moreover, that Definitions 5.2 and 5.5 are fulfilled; that  $\mathcal{X}$  and the second fundamental form are  $C^0$ -future bounded; that there are  $\alpha_{\infty}, \zeta_{\infty} \in \mathbf{M}_m(\mathbb{C})$  and  $0 < \eta_{\mathrm{mn}}, C_{\mathrm{mn}} \in \mathbb{R}$  with the property that (4.2) holds for all  $t \geq 0$ ; that there are  $X^{j_k}_{\infty} \in \mathbf{M}_m(\mathbb{C})$ ,  $k = 1, \ldots, d_{\mathrm{ts}}$ , and a constant  $0 < K_{\mathrm{ts}} \in \mathbb{R}$  with the property that (5.9) holds for all  $t \geq 0$  and all  $k \in \{1, \ldots, d_{\mathrm{ts}}\}$ ; and that if there is an  $X^{j_k}_{\infty} \neq 0$ , then  $d_{\mathrm{ts}} = 1$  and  $R_{\mathrm{ts}} = 0$ . Fix a  $0 < \beta_{\mathrm{mar}} \in \mathbb{R}$  which is small enough, the bound depending only on the coefficients of the operator  $\Delta_{\infty}$  introduced in (5.18);  $X^{j_k}_{\infty}, k = 1, \ldots, d_{\mathrm{ts}}; \alpha_{\infty}, \zeta_{\infty}$ ; and the constant

$$\beta_{\rm ts} := \min\{\eta_{\rm ts}, \beta_{\rm sil}, \eta_{\rm mn}\}.$$

Assume f to be such that  $||f||_{ts,s} < \infty$  for all  $s \in \mathbb{R}$ , where  $||\cdot||_{ts,s}$  is defined in (5.23). Then there are constants  $\sigma_{hom}, \sigma_{ih} \in [0, \infty), 0 \le N \in \mathbb{Z}$  and  $0 < C \in \mathbb{R}$  such that the following holds. Given a solution u to (1.2), there is a unique  $U_{\infty} \in C^{\infty}(\overline{M}, \mathbb{C}^{2m})$  with the following properties

- the  $\iota$ 'th Fourier coefficient of  $U_{\infty}$  vanishes unless  $\iota \in \mathcal{I}_{B,ts}$ ,
- if  $\iota \in \mathcal{I}_{B,ts}$ , then the  $\iota$ 'th Fourier coefficient of  $U_{\infty}$  belongs to  $E_{\iota}$ , where  $E_{\iota}$  is the first generalised eigenspace in the  $(\beta_{ts} \beta_{mar}), A(\iota)$ -decomposition of  $\mathbb{C}^{2m}$ , and  $A(\iota)$  is given by (5.14),
- the estimate

$$\left\| \begin{pmatrix} (-\Delta_{\infty})^{1/2} u(\cdot,t) \\ \partial_t u(\cdot,t) \end{pmatrix} - e^{\mathcal{A}_{ts}t} U_{\infty} - \int_0^t e^{\mathcal{A}_{ts}(t-\tau)} \begin{pmatrix} 0 \\ f(\cdot,\tau) \end{pmatrix} d\tau \right\|_{t,s}$$

$$\leq C \langle t \rangle^N e^{-(\beta_{ts}-\beta_{mar})t} [\|u_t(\cdot,0)\|_{(s+\sigma_{hom})} + \|u(\cdot,0)\|_{(s+\sigma_{hom}+1)} + \|f\|_{ts,s+\sigma_{ih}}]$$

$$(5.24)$$

holds for all  $t \ge 0$  and  $s \in \mathbb{R}$ , where  $\Delta_{\infty}$  is defined in (5.18),  $\mathcal{A}_{ts}$  is defined in (5.20) and  $\|\cdot\|_{t,s}$  is defined in (5.22).

Moreover,

$$\|U_{\infty}\|_{(s)} \le C \|u_t(\cdot, 0)\|_{(s+\sigma_{\text{hom}})} + C \|u(\cdot, 0)\|_{(s+\sigma_{\text{hom}}+1)} + C \|f\|_{\text{ts},s+\sigma_{\text{ih}}}$$
(5.25)

for all  $s \in \mathbb{R}$ .

**Remark 5.12.** The constant *C* appearing in (5.24) and (5.25) only depends on the coefficients of (1.2), the spectrum of the Riemannian manifolds  $(M_{r_j}, g_{r_j})$ ,  $j = 1, \ldots, R_{ts}$ , and  $\beta_{mar}$ . The constants  $\sigma_{hom}$  and  $\sigma_{ih}$  have the same dependence and *N* only depends on *m*.

*Proof.* The proposition is a consequence of Proposition 12.22; cf. Remark 12.24. The statements of Remark 5.12 follow from Remark 12.23.

Due to the proposition, we can think of the transparent part of solutions to (1.2) as being well approximated by the transparent part of solutions to the limit equation (5.10). Just as in the case of silent equations, we obtain the leading order asymptotics. However, there is a limit to how detailed asymptotics we can obtain. Moreover, the limit is set by the size of the discrepancy between the actual equation and the limit equation (quantitatively, the limit is determined by the number  $\beta_{ts}$ ). Finally, (5.25) implies that the map from initial data to asymptotic data is continuous.

The reader interested in an application of this result is referred to Section 5.5 below.

# 5.4 Specifying the asymptotics

In the previous section, we derived asymptotics, given a solution. However, it is also of interest to specify the leading order asymptotics, just as in the case of silent equations. In analogy with Proposition 4.16, it is sufficient to focus on the homogeneous case. In this section, we therefore assume that f = 0 in (1.2). When comparing the present context with that of Proposition 4.16, there is one important difference: we cannot assume the asymptotic data to take its values in one specific subspace of  $\mathbb{C}^{2m}$ ; cf. the statement of Proposition 5.11. For this reason, it is convenient to introduce the following terminology.

**Definition 5.13.** Let (M, g) be a canonical separable cosmological model manifold. Assume that the variables can be divided as described in Definition 5.1. Let  $E_{ts}$  be a function from  $\mathcal{I}_{B,ts}$  to the set of vector subspaces of  $\mathbb{C}^{2m}$ . Then  $\psi \in C^{\infty}(\bar{M}, \mathbb{C}^{2m})$  is said to be  $E_{ts}$ -adapted if the  $\iota$ 'th Fourier coefficient of  $\psi$  vanishes for  $\iota \in \mathcal{I}_{B,sil}$  and belongs to  $E_{ts}(\iota)$  for  $\iota \in \mathcal{I}_{B,ts}$ . The set of  $E_{ts}$ -adapted elements of  $C^{\infty}(\bar{M}, \mathbb{C}^{2m})$  is denoted by  $C^{\infty}(\bar{M}, \mathbb{C}^{2m}; E_{ts})$ . Finally,  $C_{ts}^{\infty}(\bar{M}, \mathbb{C}^{2m})$  denotes the set of elements of  $C^{\infty}(\bar{M}, \mathbb{C}^{2m})$  whose  $\iota$ 'th Fourier coefficients vanish for  $\iota \in \mathcal{I}_{B,sil}$ .

Next, we demonstrate that we can specify the leading order asymptotics.

**Proposition 5.14.** Assume that the conditions of Proposition 5.11 are fulfilled. Let  $\beta_{\text{mar}}$  and  $\beta_{\text{ts}}$  be as in the statement of Proposition 5.11. Assume, finally, that f = 0. Then there is a function  $E_{\text{ts}}$  from  $\mathcal{I}_{\text{B,ts}}$  to the set of subspaces of  $\mathbb{C}^{2m}$  such that if  $E_{\iota} := E_{\text{ts}}(\iota)$ , then the spaces  $E_{\iota}$  have the properties stated in Proposition 5.11. Moreover, there are constants  $C_{\Phi} > 0$  and  $s_{\infty} \geq 0$ , and an injective linear map  $\Phi_{\infty}$  from  $C^{\infty}(\bar{M}, \mathbb{C}^{2m}; E_{\text{ts}})$  to  $C_{\text{ts}}^{\infty}(\bar{M}, \mathbb{C}^{2m})$  with the following properties. First,

$$\|\Phi_{\infty}(\psi)\|_{(s)} \le C_{\Phi} \|\psi\|_{(s+s_{\infty})}$$
(5.26)

for all  $s \in \mathbb{R}$  and  $\psi \in C^{\infty}(\overline{M}, \mathbb{C}^{2m}; E_{ts})$ . Second, if  $\psi \in C^{\infty}(\overline{M}, \mathbb{C}^{2m}; E_{ts})$  and u is the solution to (1.2) (with f = 0) such that

$$\begin{pmatrix} u(\cdot,0)\\ u_t(\cdot,0) \end{pmatrix} = \Phi_{\infty}(\psi), \tag{5.27}$$

then

$$\left\| \begin{pmatrix} (-\Delta_{\infty})^{1/2} u(\cdot,t) \\ u_t(\cdot,t) \end{pmatrix} - e^{\mathcal{A}_{ts}t} \psi \right\|_{t,s}$$

$$\leq C \langle t \rangle^N e^{-(\beta_{ts} - \beta_{mar})t} \left( \| u_t(\cdot,0) \|_{(s+\sigma_{hom})} + \| u(\cdot,0) \|_{(s+\sigma_{hom}+1)} \right),$$
(5.28)

where the constants C, N and  $\sigma_{\text{hom}}$  have the same dependence as in the case of Proposition 5.11; cf. Remark 5.12. Finally, if there is an  $\epsilon > 0$  such that  $\text{Rsp}[A(\iota)] < \beta_{\text{ts}} - \epsilon$  for all  $\iota \in \mathcal{I}_{\text{B,ts}}$ , cf. Definition 4.3, then  $E_{\text{ts}}$  can be chosen to be such that  $E_{\text{ts}}(\iota) = \mathbb{C}^{2m}$  for all  $\iota \in \mathcal{I}_{\text{B,ts}}$ . In that case,  $\Phi_{\infty}$  is surjective.

**Remark 5.15.** The constant  $C_{\Phi}$  only depends on the coefficients of (1.2), the spectrum of the Riemannian manifolds  $(M_{r_j}, g_{r_j})$ ,  $j = 1, \ldots, R_{\text{ts}}$ ,  $\beta_{\text{mar}}$  and the supremum of  $\kappa_{\iota}$  for  $\iota \in \mathcal{I}_{\text{B,ts}}$ . The constant  $s_{\infty} \geq 0$  only depends on the coefficients of (1.2) and the supremum of  $\kappa_{\iota}$  for  $\iota \in \mathcal{I}_{\text{B,ts}}$ .

**Remark 5.16.** By combining (5.26), (5.27) and (5.28), the norms of  $u(\cdot, 0)$  and  $u_t(\cdot, 0)$  appearing on the right hand side of (5.28) can be replaced by a suitable Sobolev norm of  $\psi$ .

**Remark 5.17.** In order to obtain a similar result in the case of inhomogeneous equations, it is sufficient to combine Propositions 5.11 and 5.14.

**Remark 5.18.** Due to the estimate (5.26), the map from asymptotic data to initial data is continuous.

*Proof.* The statement follows from Proposition 12.25 and Remark 12.24. The statements of Remark 5.15 follow from Remark 12.26.  $\hfill \square$ 

Due to Proposition 5.14, we are in a position to calculate  $\eta_{\rm cr}$ . The reason for this is the following. First, we can divide a solution, say u, to the homogeneous version of (1.2) into two parts:  $u_{\rm ts}$  and  $u_{\rm sil}$ . Due to Proposition 4.4, the energy of  $u_{\rm sil}$  does not grow faster than  $\langle t \rangle^{2d_1-2}e^{2\kappa_1 t}$ , where  $\kappa_1$  and  $d_1$  are defined in the statement of Proposition 4.4. Similarly, due to Proposition 5.11, for every  $\epsilon > 0$ , the energy of  $u_{\rm ts}$  does not grow faster than  $e^{2(\kappa+\epsilon)t}$ , where  $\kappa$  is the supremum of the  $\kappa_{\iota}$ 's introduced at the beginning of Section 5.3. Thus  $\eta_{\rm cr} \leq \max\{\kappa_1, \kappa\}$ . Combining this observation with Propositions 4.16 and 5.14 yields the conclusion that  $\eta_{\rm cr} = \max\{\kappa_1, \kappa\}$ ; cf. Remark 5.10 for a justification of the statement that Proposition 4.16 applies. On the other hand, we do not obtain any conclusions concerning  $\eta_{\rm nl}$ .

# 5.5 An example: the flat Kasner solution

In Example 4.21, we consider the Klein-Gordon equation for all the non-flat Kasner solutions. In the case of the flat Kasner solution, the Klein-Gordon equation also takes the form (4.18) (though in what follows, we relabel  $\tau$  to t). However, all the  $\beta_i$  except one equal 1 and the exceptional exponent, say  $\beta_d$ , equals 0. In the case of solutions that only depend on the first d-1 variables, an analysis similar to the one provided in Example 4.21 yields the conclusion that for such solutions, there is a homeomorphism from initial data to asymptotic data. Moreover, an estimate of the form (4.20) holds, where  $\mu = 1$ . Let us therefore focus on the part of the solution that corresponds to modes with  $n_d \neq 0$ ;

$$u_{\rm ts}(x,t) = \sum_{n \in \mathbb{Z}^d, n_d \neq 0} z(n,t) (2\pi)^{-d/2} e^{in \cdot x}.$$

In accordance with Definition 5.1, there is a division of  $\{1, \ldots, d\}$  into a silent subset (in our case given by  $\{1, \ldots, d-1\}$ ) and a transparent subset (in our case given by  $\{d\}$ ). Moreover, Definition 5.2 applies with  $C_a = 1$ ,  $\eta_{ts} = 2$ ,  $\beta_{sil} = 1$  and  $\mathfrak{e}_{sil} = 0$ . Since the shift vector field vanishes in the present setting, it is clear that it is asymptotically negligible in the sense of

Definition 5.5. Since  $\mathcal{X} = 0$  and (4.2) holds, with  $C_{\rm mn} = m^2$ ,  $\eta_{\rm mn} = 2$ ,  $\alpha_{\infty} = 0$  and  $\zeta_{\infty} = 0$ , the limit equation (5.10) takes the form

$$v_{tt} - v_{dd} = 0. (5.29)$$

Next, note that  $\mathcal{I}_{\rm B} = \mathbb{Z}^d$  in the present setting. For this reason, we here use the notation n instead of  $\iota$ . The function  $\mathfrak{g}_{\infty}$ , introduced in (5.12), is here given by  $\mathfrak{g}_{\infty}(n) = |n_d|$ . Turning to  $A(\iota)$ ,  $w(\iota, t)$  and  $F(\iota, t)$ , introduced in (5.14) and (5.15), they are given by

$$A(n) = \begin{pmatrix} 0 & |n_d| \\ -|n_d| & 0 \end{pmatrix}, \quad w(n,t) = \begin{pmatrix} |n_d|z(n,t) \\ \dot{z}(n,t) \end{pmatrix}$$

and F(n,t) = 0. In particular,

$$e^{A(n)t} = \begin{pmatrix} \cos(|n_d|t) & \sin(|n_d|t) \\ -\sin(|n_d|t) & \cos(|n_d|t) \end{pmatrix}.$$

If  $U_{\infty} \in C^{\infty}(\mathbb{T}^d, \mathbb{C}^2)$  is a function with Fourier coefficients  $\hat{U}_{\infty}$  such that  $\hat{U}_{\infty}(n) = 0$  if  $n_d = 0$ , then

$$(e^{\mathcal{A}_{ts}t}U_{\infty})(x,t) = \sum_{n \in \mathbb{Z}^d, n_d \neq 0} \begin{pmatrix} \cos(|n_d|t) & \sin(|n_d|t) \\ -\sin(|n_d|t) & \cos(|n_d|t) \end{pmatrix} \hat{U}_{\infty}(n)(2\pi)^{-d/2}e^{in \cdot x}.$$

In practice, the variables  $x^1, \ldots, x^{d-1}$  are thus frozen, and we solve the linear wave equation (5.29) in the  $tx^d$ -directions.

Let us now return to Proposition 5.11. Note that in the present setting, the assumptions of this proposition are satisfied. In addition,  $\beta_{ts} = 1$ . Fix  $0 < \beta_{mar} \in \mathbb{R}$  satisfying the restrictions of Proposition 5.11. Given a solution u to (4.18), there is a unique  $U_{\infty} \in C^{\infty}(\mathbb{T}^d, \mathbb{C}^2)$  with the property that  $\hat{U}_{\infty}(n) = 0$  if  $n_d = 0$ , and the property that

$$\left\| \begin{pmatrix} |\partial_d| u_{\rm ts}(\cdot, t) \\ \partial_t u_{\rm ts}(\cdot, t) \end{pmatrix} - e^{\mathcal{A}_{\rm ts}t} U_{\infty} \right\|_{(s)} \le C_s \langle t \rangle^N e^{-(1-\beta_{\rm mar})t}$$
(5.30)

for some constant  $0 < C_s \in \mathbb{R}$  and all  $t \ge 0$ . Here the constant  $C_s$  is allowed to depend on s, the solution and the equation, and  $|\partial_d| := (-\partial_d^2)^{1/2}$ . In fact, appealing to Proposition 5.14, the map from initial data to  $U_{\infty}$  is a homeomorphism with respect to the  $C^{\infty}$  topology. In order to obtain this conclusion, we use the fact that the eigenvalues of A(n) are purely imaginary. Turning to the function  $u_{ts}$  itself, note that (5.30) implies that

$$\sum_{n \in \mathbb{Z}^d, n_d \neq 0} \langle n \rangle^{2s} \left| |n_d| z(n,t) - \cos(|n_d|t) \hat{U}_1(n) - \sin(|n_d|t) \hat{U}_2(n) \right|^2 \le C_s \langle t \rangle^{2N} e^{-2(1-\beta_{\max})t}$$

for all  $t \ge 0$ , where  $\hat{U}_i$ , i = 1, 2, are the components of  $\hat{U}_{\infty}$ . Combining this estimate with (5.30) yields the conclusion that there is a function  $U_{\text{ts}} \in C^{\infty}(\mathbb{T}^d \times \mathbb{R}, \mathbb{C})$  which solves (5.29); has the property that  $\hat{U}_{\text{ts}}(n, \cdot) = 0$  for all  $n \in \mathbb{Z}^d$  such that  $n_d = 0$ ; and is such that

$$\|u_{\rm ts}(\cdot,t) - U_{\rm ts}(\cdot,t)\|_{(s)} + \|\partial_t u_{\rm ts}(\cdot,t) - \partial_t U_{\rm ts}(\cdot,t)\|_{(s)} \le C_s \langle t \rangle^N e^{-(1-\beta_{\rm mar})t}$$

for all  $t \geq 0$ .

# 5.6 Outline of the argument

The strengths and weaknesses of the results are quite similar to those in the silent setting; cf. Subsection 4.4.1. Here, we therefore focus on an outline of the proof. The rough structure of the argument in the transparent setting is similar to that of the argument in the silent setting. In particular, Subsection 4.4.2 gives a general idea of how to proceed. The main difference is

due to the fact that the set of  $A(\iota)$  is infinite. This difference gives rise to difficulties that are purely algebraic, but somewhat technical to resolve. In order to describe the complications, let us consider the matrices  $A(\iota)$  in greater detail.

**Model matrices.** Consider the case that  $X_{\infty}^{j_k} \neq 0$  for some  $k \in \{1, \ldots, d_{ts}\}$ . Then, due to the assumptions of Propositions 5.11 and 5.14,  $d_{ts} = 1$  and  $R_{ts} = 0$ . Thus  $\mathfrak{g}_{\infty}(\iota) = (g_{\infty}^{j_1 j_1})^{1/2} |n_{j_1}|$ , so that  $A(\iota)$  equals

$$A_{\mu} := \begin{pmatrix} 0 & \mu \mathrm{Id}_{m} \\ -\mu \mathrm{Id}_{m} + iV + 2\mu^{-1}W & U \end{pmatrix},$$
 (5.31)

where  $\mu := \mathfrak{g}_{\infty}(\iota); U := -\alpha_{\infty}; W := -\zeta_{\infty}/2;$  and

$$V := -\frac{n_{j_1}}{|n_{j_1}|} \frac{X_{\infty}^{j_1}}{(g_{\infty}^{j_1 j_1})^{1/2}}$$

For  $\iota \in \mathcal{I}_{B,ts}$ ,  $n_{j_1} \neq 0$ , so that there are only two possibilities for V. In that sense, it is sufficient to focus on matrices of the form (5.31) for fixed matrices  $U, V, W \in \mathbf{M}_m(\mathbb{C})$ . In case all the  $X^{j_k}_{\infty}$  vanish,  $A(\iota)$  also takes the form (5.31), this time with V = 0.

Asymptotic expansions for the eigenvalues. Let us consider  $A_{\mu}$  in greater detail. Conjugating  $A_{\mu}$  by a matrix independent of  $\mu$  yields a matrix  $N_{\mu}$  of the form

$$N_{\mu} = \begin{pmatrix} -i\mu \mathrm{Id}_{m} + R_{\mu}^{-} & Q_{\mu}^{+} \\ Q_{\mu}^{-} & i\mu \mathrm{Id}_{m} + R_{\mu}^{+} \end{pmatrix},$$
(5.32)

where

$$R^{\pm}_{\mu} = \gamma_{\pm} \mp i \mu^{-1} W_{\pm}, \quad Q^{\pm}_{\mu} = \pm i \delta_{\pm} + \mu^{-1} V_{\pm}.$$

Here  $\gamma_{\pm}$ ,  $\delta_{\pm}$ ,  $V_{\pm}$  and  $W_{\pm}$  are independent of  $\mu$ . Moreover, the  $\gamma_{\pm}$  are Jordan normal forms of the matrices  $(U \pm V)/2$ . Let  $\lambda_{j,\pm}$ ,  $j = 1, \ldots, p_{\pm}$ , be the distinct eigenvalues of  $\gamma_{\pm}$ , and  $m_{j,\pm}$  be the corresponding multiplicities. It can then be demonstrated that there are constants  $0 < c_a, \mu_a \in \mathbb{R}$  such that for  $\mu \ge \mu_a$  and  $j = 1, \ldots, p_{\pm}$ , there are  $m_{j,\pm}$  eigenvalues of  $N_{\mu}$  in a ball of radius  $c_a \mu^{-1/m_{j,\pm}}$  and centre  $\pm i\mu + \lambda_{j,\pm}$ ; cf. Lemma 11.6. In other words, there are asymptotic expansions for the eigenvalues.

Asymptotic partial diagonalisation. Ideally, we would like to diagonalise  $N_{\mu}$ . However, the potential multiplicities of the eigenvalues of  $\gamma_{\pm}$  cause problems. In the end, we therefore only obtain a partial diagonalisation. In fact, for sufficiently large  $\mu$ , there are matrices  $T_{\mu} \in \mathbf{M}_{2m}(\mathbb{C})$  such that  $||T_{\mu}||, ||T_{\mu}^{-1}|| \leq 2$  and such that

$$T_{\mu}^{-1}N_{\mu}T_{\mu} = \operatorname{diag}\{N_{\mu,1}^{-}, \dots, N_{\mu,p_{-}}^{-}, N_{\mu,1}^{+}, \dots, N_{\mu,p_{+}}^{+}\},$$
(5.33)

where

$$||N_{\mu,j}^{\pm} \mp i\mu \mathrm{Id}_{m_{j,\pm}} - \gamma_{\pm,j}|| \le C\mu^{-1}.$$

Here  $\gamma_{\pm,j}$ ,  $j = 1, \ldots, p_{\pm}$ , is the matrix collecting all the Jordan blocks in  $\gamma_{\pm}$  corresponding to the eigenvalue  $\lambda_{j,\pm}$ . The justification for these statements is to be found in Lemma 11.14. The proof is quite long, and requires some background material, which we develop in Section 11.3 below.

Dividing the matrices according to frequency. Fix  $0 \leq \mu_0 \in \mathbb{R}$ . Due to the assumptions, the set of  $\mathfrak{g}_{\infty}(\iota)$  satisfying  $\mathfrak{g}_{\infty}(\iota) \leq \mu_0$  is finite. Considering the family of matrices  $A(\iota)$ , there is thus a natural way to divide it into two subsets: the matrices corresponding to  $\mathfrak{g}_{\infty}(\iota) \leq \mu_0$  and the matrices corresponding to  $\mathfrak{g}_{\infty}(\iota) > \mu_0$ . The use of this division is due to the fact that for large  $\mu_0$ , there is a partial diagonalisation of the form (5.33). Choosing  $\mu_0$  large enough, the behaviour of the modes satisfying  $\mathfrak{g}_{\infty}(\iota) > \mu_0$  can therefore be analysed (up to, possibly, some small error). Since the remaining modes only correspond to a finite number of matrices  $A(\iota)$ , their behaviour can be analysed by appealing to the results of the previous chapter.

Asymptotic analysis for the modes. Let us now focus on the Fourier mode corresponding to a  $\iota \in \mathcal{I}_{B,ts}$ . In analogy with the discussion in Subsection 4.4.2, it is natural to divide  $[0, \infty)$  into two

subsets:  $[0, T_{ts}]$  and  $(T_{ts}, \infty)$ . Here  $T_{ts}$  is defined so that the error terms (i.e., the terms that make up the difference between the actual equation and the limit equation) are bounded by  $Ce^{-\beta_{ts}\bar{t}}$  for  $t \geq T_{ts}$ , where C is a constant independent of  $\iota$  and  $\bar{t} := t - T_{ts}$ . Moreover,  $T_{ts}$  is logarithmic in  $\langle \nu(\iota) \rangle$ . Just as in the silent setting, the analysis in  $[0, T_{ts}]$  consists of a crude energy estimate. Due to the fact that  $T_{ts}$  is logarithmic in  $\langle \nu(\iota) \rangle$ , this is sufficient; our lack of detailed knowledge in  $[0, T_{ts}]$  only corresponds to a finite loss of derivatives. How we analyse the asymptotics in the interval  $(T_{ts}, \infty)$  depends on the division of the modes into low and high frequencies. The parameter defining the division is  $\mu_0$ . A large  $\mu_0$  yields more detailed control of the eigenvalues (and corresponds to a smaller  $\beta_{mar}$ ; cf. Section 5.3). On the other hand, the larger the  $\mu_0$ , the larger the number of cases to which we need to apply the analysis of the previous chapter (and, thereby, the larger the constants appearing in the estimates). Given the partial diagonalisation (5.33), we are in a position to carry out an analysis for high frequencies and  $t \geq T_{ts}$ . This is the subject of Section 11.4. In Section 11.5, we then turn to the problem of specifying the asymptotic data for an individual high frequency mode.

Summing up. Given the analysis described above and the results of the previous chapter, we are in a position to derive the desired conclusions concerning transparent equations. This is the subject of Chapter 12. We start, in Sections 12.1 and 12.2, by describing the equations of interest; estimating the discrepancy between the actual equation and the limit equation for fixed modes; and defining  $T_{\rm ts}$ . The derivation of the asymptotics is divided into two parts. In Lemma 12.18, we derive conclusions for the part of the solution corresponding to all the high frequency modes. Combining this result with the methods developed in the silent setting yields the desired conclusions concerning the asymptotics for the transparent part of the solution; cf. Proposition 12.22. In particular, we obtain a continuous map from initial data to asymptotic data. We end the chapter in Section 12.4 by demonstrating that we can specify the leading order asymptotics; cf. Proposition 12.25. Again, the map from asymptotic data to initial data is continuous.

# Chapter 6

# Equations with a dominant noisy spatial direction

# 6.1 Introduction

In Chapter 4 we consider the case that  $\bar{k}$  has a strictly positive lower bound and in Chapter 5 we consider the case that  $\bar{k}$  asymptotically vanishes in some directions. As a next step, it is natural to allow  $\bar{k}$  to be negative in some directions. Note that this corresponds to contraction. As far as the asymptotic behaviour is concerned, the direction in which the contraction is the strongest is the most important. Moreover, assuming that there is one direction in which the contraction dominates simplifies the analysis, and we restrict our attention to this case. For reasons mentioned in Subsection 1.3.2, we refer to the corresponding geometries (and equations) as noisy. Two examples of noisy geometries are given by the Kasner solutions and the U(1)-symmetric solutions (in the expanding direction); cf. Subsection 1.3.1. In the study of the expanding direction of polarised Gowdy solutions, we also obtain noisy equations.

### 6.1.1 Outline

Division of the variables and basic assumptions. In order to give a formal definition of what it means for an equation to have a dominant noisy spatial direction, we need to divide the cotangent space into directions in which the spacetime contracts maximally and the remaining directions. The formal definition of this division is similar to the definition in the transparent setting, cf. Definition 5.1, and is provided in Subsection 6.1.2. In Subsection 6.1.2, we also describe the requirements on the second fundamental form, the shift vector field, the  $X^{l}$ 's,  $\alpha$  and  $\zeta$ . Concerning the second fundamental form, the main assumption is that there is a  $0 < \beta_n \in \mathbb{R}$  such that  $\bar{k}^{\sharp} + \beta_{n} \bar{g}^{\sharp}$  converges to zero exponentially in the dominant noisy spatial direction. Concerning the remaining directions, we only need to assume that the contraction is slower, with a margin. Turning to the coefficients of the lower order derivatives, we only need to impose convergence conditions on the  $X^{l}$ 's corresponding to the dominant noisy direction and on  $\alpha$ . Beyond these requirements, we demand that (1.2) be  $C^2$ -balanced and that the  $C^1$ -norm of the shift vector field decay exponentially. In other words, the conditions are weaker than the ones imposed in the silent and transparent settings, in the sense that we do not require  $\zeta$  to converge. However, they are stronger in that we require control over a higher number of derivatives. In both cases, the reason for the difference is the fact that the solutions exhibit oscillatory behaviour, with a frequency that grows exponentially. Due to the exponentially growing frequency,  $\zeta$  is effectively an error term, as long as it is future bounded in  $C^1$ . On the other hand, in order to control the evolution in this highly oscillatory setting, we need to approximate the oscillations, just as for the equations considered in Chapter 2; cf. Section 2.5. In order for the relevant methods, developed in Part III, to work, we need to impose bounds on a larger number of derivatives than in the silent and transparent settings.

The limit equation. Just as in the case of silent and transparent equations, there is a limit equation in the noisy setting. On the Fourier side, cf. Section 24.1, the homogeneous version of this equation reads

$$\ddot{z} + \nu_{\mathbf{n}}^2(\iota)e^{2\beta_{\mathbf{n}}t}z + i\nu_{\mathbf{n}}(\iota)e^{\beta_{\mathbf{n}}t}\bar{X}_{\mathbf{n}}(\iota)z + \alpha_{\infty}\dot{z} = 0;$$
(6.1)

cf. (6.16) below. Here  $\nu_n(\iota)$  is the limit of  $e^{-\beta_n t}\mathfrak{g}(\iota, t)$  (which exists due to the assumptions). Moreover,  $\alpha_{\infty}$  is the limit of  $\alpha$ , and  $\bar{X}_n(\iota)$  is either zero or, up to a sign, given by a limit that exists due to the assumptions; cf. (6.13)–(6.15) below.

At this point, let us remark that the assumptions that we make in the present chapter only allow us to draw conclusions for the  $\iota \in \mathcal{I}_B$  such that  $\nu_n(\iota) \neq 0$ . The Fourier modes with  $\nu_n(\iota) = 0$ correspond to solutions to the so-called *subdominant equation*. This equation is obtained by setting all the coefficients in front of the derivatives with respect to the dominant noisy spatial variables to zero. In order to obtain asymptotic information concerning solutions to the subdominant equation, we need to make additional assumptions. For instance, if the subdominant equation is transparent in the sense of the previous chapter, then the results of the previous two chapters apply; if it is silent, the results of Chapter 4 apply; and if it has a dominant noisy spatial direction in the sense of the present chapter, then we can proceed inductively.

Averaging over the oscillations. Consider a solution to (6.1) corresponding to a  $\iota \in \mathcal{I}_B$  such that  $\nu_n(\iota) \neq 0$ . Since  $\nu_n(\iota)e^{\beta_n t} \to \infty$ , it is clear that the dominant behaviour of solutions is oscillatory. However, it is also of interest to understand the overall behaviour over longer periods of time. To this end, it is of interest to consider one period of the oscillations in detail and then to derive conclusions concerning the overall behaviour by combining the conclusions concerning the individual periods; cf. the discussion in Section 2.5. In order to develop a feeling for the evolution over one period, fix a  $t_0 \geq 0$ . Then the corresponding period of the oscillations is roughly speaking  $T := 2\pi/\mathfrak{g}_{\infty}(\iota, t_0)$ , where  $\mathfrak{g}_{\infty}(\iota, t) := \nu_n(\iota)e^{\beta_n t}$ . Moreover, (6.1) can be rewritten as

$$\dot{w}(\iota, t) = A_{\infty}(\iota, t)w(\iota, t), \tag{6.2}$$

where

$$w(\iota, t) := e^{-\beta_{n}(t-t_{0})/2} \begin{pmatrix} e^{\alpha_{\infty}(t-t_{0})/2} \mathfrak{g}_{\infty}(\iota, t) z(\iota, t) \\ e^{\alpha_{\infty}(t-t_{0})/2} \dot{z}(\iota, t) \end{pmatrix},$$
  

$$A_{\infty}(\iota, t) := \begin{pmatrix} (\beta_{n} \mathrm{Id}_{m} + \alpha_{\infty})/2 & \mathfrak{g}_{\infty}(\iota, t) \mathrm{Id}_{m} \\ -\mathfrak{g}_{\infty}(\iota, t) \mathrm{Id}_{m} - i \overline{Y}_{n}(\iota, t) & -(\beta_{n} \mathrm{Id}_{m} + \alpha_{\infty})/2 \end{pmatrix}$$
(6.3)

and

$$\bar{Y}_{n}(\iota, t) := e^{\alpha_{\infty}(t-t_{0})/2} \bar{X}_{n}(\iota) e^{-\alpha_{\infty}(t-t_{0})/2}.$$
(6.4)

Note that the choice of variables ensures that the sum of the matrices on the diagonal of  $A_{\infty}$  equals zero. A naive way of approximating the evolution of w from  $t_0$  to  $t_0 + T$  is to say that it corresponds to multiplication by  $\exp[A_{\infty}(\iota, t_0)T]$ . However, due to the fact that T decays exponentially as  $t_0$  grows, we expect this approximation to gradually become better and better. The matrix  $A_{\infty}(\iota, t_0)T$  has a very special form. First, the dominant part corresponds to the evolution over exactly one full period of the oscillations. Second, the sum of the matrices on the diagonal vanishes (so that, intuitively, the mean contribution from the diagonal matrices over one period should vanish). Third, the only matrix that remains beyond this is

$$-iY_{\rm n}(\iota, t_0)T = -iX_{\rm n}(\iota)T.$$

In Section 13.2, we develop methods for calculating exponentials of matrices of exactly this form. The result is

$$\exp[A_{\infty}(\iota, t_0)T] \approx \operatorname{Id}_{2m} + \frac{1}{2} \begin{pmatrix} 0 & i\bar{X}_{n}(\iota) \\ -i\bar{X}_{n}(\iota) & 0 \end{pmatrix} T.$$

Combining the above definitions and observations yields the conclusion that if  $t_1 := t_0 + T$ , then

$$\begin{pmatrix} \mathfrak{g}_{\infty}(\iota,t_1)z(\iota,t_1) \\ \dot{z}(\iota,t_1) \end{pmatrix} \approx e^{\beta_{n}T/2} \begin{pmatrix} e^{-\alpha_{\infty}T/2} & 0 \\ 0 & e^{-\alpha_{\infty}T/2} \end{pmatrix} \exp[A_{\infty}(\iota,t_0)T]w(\iota,t_0) \\ \approx [\mathrm{Id}_{2m} + B_{\infty}(\iota)T]w(\iota,t_0),$$

where

$$B_{\infty}(\iota) := \frac{1}{2} \begin{pmatrix} \beta_{n} \mathrm{Id}_{m} - \alpha_{\infty} & i\bar{X}_{n}(\iota) \\ -i\bar{X}_{n}(\iota) & \beta_{n} \mathrm{Id}_{m} - \alpha_{\infty} \end{pmatrix}.$$

Thus

$$\left( \begin{array}{c} \mathfrak{g}_{\infty}(\iota,t_1)z(\iota,t_1) \\ \dot{z}(\iota,t_1) \end{array} \right) \approx e^{B_{\infty}(\iota)T} \left( \begin{array}{c} \mathfrak{g}_{\infty}(\iota,t_0)z(\iota,t_0) \\ \dot{z}(\iota,t_0) \end{array} \right)$$

Conjugating  $B_{\infty}(\iota)$  by

$$S := \frac{1}{2} \left( \begin{array}{cc} \mathrm{Id}_m & i \mathrm{Id}_m \\ i \mathrm{Id}_m & \mathrm{Id}_m \end{array} \right)$$

yields

$$SB_{\infty}(\iota)S^{-1} = \frac{1}{2} \begin{pmatrix} \beta_{n} \mathrm{Id}_{m} - \alpha_{\infty} + \bar{X}_{n}(\iota) & 0\\ 0 & \beta_{n} \mathrm{Id}_{m} - \alpha_{\infty} - \bar{X}_{n}(\iota) \end{pmatrix}.$$
 (6.5)

Naively, it is thus reasonable to expect the overall evolution to be determined by the diagonal elements of this matrix. Note that  $\bar{X}_n(\iota)$  can be written as  $\pm V$  for a fixed matrix V (independent of  $\iota$ ). For this reason, the set consisting of the matrices on the diagonal of (6.5) is independent of  $\iota$ . In particular, the growth of generic solutions to  $\dot{v} = B_{\infty}(\iota)v$  is independent of  $\iota$ .

**Outline, results.** After describing the conditions we impose in the present chapter, cf. Subsection 6.1.2; and the limit equation, cf. Subsection 6.1.3; we turn to the results in Section 6.2. First, we derive an energy estimate in Subsection 6.2.1. The growth corresponds exactly to the generic growth of solutions to  $\dot{v} = B_{\infty}(\iota)v$  (which, as we noted above, is independent of  $\iota$ ). However, just as in the silent and transparent settings, the stated energy estimate involves a loss of derivatives; cf. Proposition 6.8.

Understanding the oscillations. Our second goal is to understand the oscillatory character of the solutions. In order to do so, we focus on the scalar equation obtained from (1.2) by dropping the right hand side and only keeping the terms on the left hand side that involve second order derivatives of u, i.e.,

$$u_{tt} - \sum_{j,l=1}^{d} g^{jl}(t) \partial_j \partial_l u - 2 \sum_{l=1}^{d} g^{0l}(t) \partial_l \partial_l u - \sum_{r=1}^{R} a_r^{-2}(t) \Delta_{g_r} u = 0.$$
(6.6)

Considering (6.5), it seems reasonable to expect oscillations with an overall growth of  $\mathfrak{E}[u]$  of the form  $e^{\beta_n t}$ . This is indeed what happens. Moreover, not only are we in a position to derive detailed asymptotics of solutions, we actually obtain a homeomorphism between initial data and asymptotic data. Due to this fact, we can consider solutions to (6.6) as models for the oscillatory behaviour. The result is stated as Proposition 6.10. As an illustration of the result, we consider the future asymptotics of polarised vacuum Gowdy solutions.

Deriving/specifying the asymptotics. Outline of the proof. Let us return to (1.2). In order to describe the future asymptotics of a solution, it turns out to be convenient to compare it with a sum of terms of the form  $e^{Rt}u$ , where R is a constant matrix and u is a vector valued solution to (6.6). However, describing the detailed combination of matrices and solutions to (6.6) needed is somewhat technical, and is left to Subsection 6.2.3, in particular Proposition 6.18. Finally, similarly to the silent and transparent settings, we can specify the leading order asymptotics. Moreover, in favourable circumstances, we obtain a homeomorphism between initial data and asymptotic data. The relevant result is stated as Proposition 6.28. In Section 6.3 we end the chapter by giving an outline of the proofs of the results.

#### 6.1.2 Dividing the cotangent space

Just as in the transparent setting, we here need to divide the cotangent space into the direct sum of two subspaces. We refer to the relevant subspaces as the dominant noisy spatial direction and the subdominant directions. In parallel with the introduction of these subspaces, we state the main convergence requirements. Since the formulation of the conditions depends on whether the dominant noisy spatial direction corresponds to an  $\mathbb{S}^1$ -factor or an  $M_r$ -factor, we state the requirements in two separate definitions. We start by considering the  $\mathbb{S}^1$ -case.

**Definition 6.1.** Let (M, g) be a canonical separable cosmological model manifold. Fix  $j \in \{1, \ldots, d\}$  and let  $\pi_n$  be the map from  $\overline{M}$  to  $\mathbb{S}^1$  corresponding to projection onto the j'th  $\mathbb{S}^1$ -factor in  $\mathbb{T}^d$ . Let  $\pi_{sub}$  be the map from  $\overline{M}$  to

$$\bar{M}_{\mathrm{sub}} := \mathbb{T}^{d-1} \times M_1 \times \dots \times M_R$$

corresponding to the projection onto what remains after removing the j'th  $S^1$ -factor. Define

$$T^*_{\mathrm{n}}\bar{M} := \pi^*_{\mathrm{n}}(T^*\mathbb{S}^1), \quad T^*_{\mathrm{sub}}\bar{M} := \pi^*_{\mathrm{sub}}(T^*\bar{M}_{\mathrm{sub}}).$$

The metric g is said to have a geometric dominant noisy spatial  $\mathbb{S}^1$ -direction corresponding to j if there are  $0 < \beta_n, \eta_n, C_n \in \mathbb{R}$  and a continuous non-negative function  $\mathfrak{e} \in L^1([0,\infty))$  such that the following holds:

$$|\bar{k}^{\sharp}(\xi,\xi) + \beta_{n}\bar{g}^{\sharp}(\xi,\xi)| \le C_{n}e^{-\eta_{n}t}\bar{g}^{\sharp}(\xi,\xi)$$
(6.7)

for all  $\xi \in T_n^* \overline{M}$  and all  $t \ge 0$ , and

$$\bar{k}^{\sharp}(\xi,\xi) \ge [-\beta_{\mathrm{n}} + \eta_{\mathrm{n}} - \mathfrak{e}(t)]\bar{g}^{\sharp}(\xi,\xi)$$
(6.8)

for all  $\xi \in T^*_{\text{sub}}\overline{M}$  and all  $t \ge 0$ . Consider (1.2), where (M, g) is such that the above conditions hold. If there are constants  $\eta_{\text{mn}} > 0$ ,  $C_{\text{mn}} > 0$  and matrices  $X^j_{\infty}, \alpha_{\infty} \in \mathbf{M}_m(\mathbb{C})$  such that

$$\|e^{-\beta_{\rm n}t}X^{j}(t) - X_{\infty}^{j}\| + \|\alpha(t) - \alpha_{\infty}\| \le C_{\rm mn}e^{-\eta_{\rm mn}t}$$
(6.9)

for all  $t \ge 0$ ; then (1.2) is said to be such that the dominant coefficients are convergent.

**Remark 6.2.** The definitions of  $\bar{g}^{\sharp}$  and  $\bar{k}^{\sharp}$  are given at the beginning of Subsection 5.1.2.

**Remark 6.3.** In Remark 5.4, we reformulate the conditions (5.1) and (5.2) on the second fundamental form in the transparent setting to (5.3) and (5.4). Using the notation  $\sharp$  introduced at the beginning of Subsection 5.1.2, and the notation

$$T_{n,t}\bar{M} := [\pi_n^*(T^*\mathbb{S}^1)]^{\sharp}, \quad T_{\mathrm{sub},t}\bar{M} := [\pi_{\mathrm{sub}}^*(T^*\bar{M}_{\mathrm{sub}})]^{\sharp},$$

there is a similar reformulation of the conditions (6.7) and (6.8).

We use similar terminology, and identical notation, when the dominant noisy spatial direction corresponds to one of the  $M_r$ .

**Definition 6.4.** Let (M, g) be a canonical separable cosmological model manifold. Fix  $r \in \{1, \ldots, R\}$  and let  $\pi_n$  be the projection from  $\overline{M}$  to the  $M_r$ -factor in  $\overline{M}$  and let  $\pi_{sub}$  be the map from  $\overline{M}$  to

$$\bar{M}_{sub} := \mathbb{T}^d \times M_1 \times \cdots \times \hat{M}_r \times \cdots \times M_R$$

corresponding to the projection onto what remains after removing  $M_r$ ; here a hat signifies omission. Define

$$T_{n}^{*}\overline{M} := \pi_{n}^{*}(T^{*}M_{r}), \quad T_{sub}^{*}\overline{M} := \pi_{sub}^{*}(T^{*}\overline{M}_{sub}).$$

The metric g is said to have a geometric dominant noisy spatial direction corresponding to  $M_r$ if there are  $0 < \beta_n, \eta_n, C_n \in \mathbb{R}$  and a continuous non-negative function  $\mathfrak{e} \in L^1([0,\infty))$  such that (6.7) holds for all  $\xi \in T_n^* \overline{M}$  and all  $t \ge 0$ , and such that (6.8) holds for all  $\xi \in T_{sub}^* \overline{M}$  and all  $t \ge 0$ . Consider (1.2), where (M, g) is such that the above conditions hold. If there are constants  $\eta_{mn} > 0$ ,  $C_{mn} > 0$  and a matrix  $\alpha_{\infty} \in \mathbf{M}_m(\mathbb{C})$  such that

$$\|\alpha(t) - \alpha_{\infty}\| \le C_{\rm mn} e^{-\eta_{\rm mn}t} \tag{6.10}$$

for all  $t \ge 0$ ; then (1.2) is said to be such that the dominant coefficients are convergent.

In what follows, it is convenient to make additional assumptions. In order for the statements of the results not to be too cumbersome, we therefore introduce the following terminology.

**Definition 6.5.** Consider (1.2). Assume the associated metric to be such that (M, g) is a canonical separable cosmological model manifold. Assume, moreover, the metric g to have a geometric dominant noisy spatial direction and the dominant coefficients of (1.2) to be convergent; i.e., either all the requirements of Definition 6.1 or all the requirements of Definition 6.4 are satisfied. Assume, in addition, that (1.2) is  $C^2$ -balanced and that there are constants  $C_{\rm sh}, \eta_{\rm sh} > 0$  such that

$$|\chi(t)|_{\bar{q}} + |\dot{\chi}(t)|_{\bar{q}} \le C_{\rm sh} e^{-\eta_{\rm sh}t} \tag{6.11}$$

for all  $t \ge 0$ . Then (1.2) is said to be  $C^2$ -balanced with a geometric dominant noisy spatial direction, convergent dominant coefficients and a negligible shift vector field.

Before turning to the results, note that given assumptions of the form stated in Definitions 6.1 and 6.4, we cannot say much about modes such that there is no spatial variation in the dominant noisy spatial direction. For that reason, we, from now on, focus on  $\iota \in \mathcal{I}_{B,n}$ , where  $\mathcal{I}_{B,n}$  is defined as follows.

**Definition 6.6.** Assume that (1.2) is  $C^2$ -balanced with a geometric dominant noisy spatial direction, convergent dominant coefficients and a negligible shift vector field; cf. Definition 6.5. Define

$$\nu_{\mathbf{n}}(\iota) := \lim_{t \to \infty} e^{-\beta_{\mathbf{n}} t} \mathfrak{g}(\iota, t), \tag{6.12}$$

where  $\beta_n$  is given in Definitions 6.1 and 6.4. Then the set  $\mathcal{I}_{B,n}$  is defined to consist of the  $\iota \in \mathcal{I}_B$  such that  $\nu_n(\iota) \neq 0$ .

**Remark 6.7.** The function  $\mathfrak{g}(\iota, t)$  is defined by (24.13). Due to Remarks 17.2 and 17.4, the limit (6.12) exists. Moreover, if g has a geometric dominant noisy spatial S<sup>1</sup>-direction corresponding to j, then  $\iota \in \mathcal{I}_{\mathrm{B}}$  if and only if  $\nu_{\mathrm{T},j}(\iota) \neq 0$ ; cf. (24.5). Similarly, if g has a geometric dominant noisy spatial corresponding to  $M_r$ , then  $\iota \in \mathcal{I}_{\mathrm{B}}$  if and only if  $\nu_{\mathrm{r},i_r}(\iota) \neq 0$ .

#### 6.1.3 The limit equation

Assume that (1.2) is  $C^2$ -balanced with a geometric dominant noisy spatial direction, convergent dominant coefficients and a negligible shift vector field; cf. Definition 6.5. Consider (24.16), the equation for the Fourier coefficients of a solution to (1.2), and assume that  $\iota \in \mathcal{I}_{B,n}$ . In analogy with earlier chapters, it is of interest to isolate a limit equation. Considering (6.12), it seems natural to, as a first approximation, replace  $\mathfrak{g}(\iota, t)$  with  $\nu_n(\iota)e^{\beta_n t}$ . Moreover, due to (6.9) and (6.10), it seems reasonable to replace  $\alpha$  with  $\alpha_{\infty}$ . Turning to the term involving the  $X^l$ , it can be verified that it, to leading order, is given by  $i\bar{X}_n(\iota)\nu_n(\iota)e^{\beta_n t}z$  (cf. Lemma 17.15 for a detailed justification). Here

$$\bar{X}_{n}(\iota) := \operatorname{sgn}_{n}(\iota) \tilde{X}_{n,\infty}.$$
(6.13)

Concerning the definition of the expressions appearing on the right hand side, there are two cases to consider. If (1.2) has a geometric dominant noisy spatial  $S^1$ -direction corresponding to j, and  $\iota \in \mathcal{I}_{B,n}$ , then

$$\tilde{X}_{\infty,n} := (g_{\infty}^{jj})^{-1/2} X_{\infty}^{j}, \quad \text{sgn}_{n}(\iota) := \frac{n_{j}}{|n_{j}|}$$
(6.14)

(no summation on j); here  $g_{\infty}^{jj} := \lim_{t\to\infty} e^{-2\beta_{n}t}g^{jj}(t)$  (that this limit exists and is strictly positive is a consequence of Remark 17.2). If (1.2) has a geometric dominant noisy spatial direction corresponding to  $M_r$  and  $\iota \in \mathcal{I}_{B,n}$ , then

$$X_{\infty,n} := 0, \quad \text{sgn}_n(\iota) := 1.$$
 (6.15)

Leaving the significance of  $\zeta$  aside for a moment, the preliminary, homogeneous version of the limit equation is given by

$$\ddot{z} + \nu_{\mathrm{n}}^2(\iota)e^{2\beta_{\mathrm{n}}t}z + i\nu_{\mathrm{n}}(\iota)e^{\beta_{\mathrm{n}}t}\bar{X}_{\mathrm{n}}(\iota)z + \alpha_{\infty}\dot{z} + \zeta(t)z = 0,$$

where the absence of the term involving the shift vector field is justified by (6.11). This equation can be written as a first order system for a vector valued function whose first m components are given by  $\nu_n(\iota)e^{\beta_n t}z$  and whose last m components are given by  $\dot{z}$ . If one does so, it becomes clear that  $\zeta$  only appears divided by  $\nu_n(\iota)e^{\beta_n t}$ . As a consequence, it seems reasonable to expect  $\zeta$  to have a negligible influence on the leading order asymptotics. In the end, this expectation turns out to be justified. Summing up yields the homogeneous limit equation

$$\ddot{z} + \nu_{\rm n}^2(\iota) e^{2\beta_{\rm n} t} z + i\nu_{\rm n}(\iota) e^{\beta_{\rm n} t} \bar{X}_{\rm n}(\iota) z + \alpha_{\infty} \dot{z} = 0.$$
(6.16)

Since  $\beta_n > 0$ , it is clear that the leading order behaviour is oscillatory, with a frequency that grows exponentially. However, there is also an overall exponential growth/decay. Considering the limit equation, it is natural to expect the growth/decay to be determined by  $\beta_n$ ,  $\tilde{X}_{\infty,n}$  and  $\alpha_{\infty}$ . In fact, we expect the matrices appearing on the diagonal on the right hand side of (6.5) to determine the growth/decay of solutions. It is therefore natural to introduce the notation

$$R_{\mathbf{n},+}^{\pm} := \frac{1}{2} (-\alpha_{\infty} + \beta_{\mathbf{n}} \mathrm{Id}_{m} \pm \tilde{X}_{\infty,\mathbf{n}}).$$

$$(6.17)$$

Let  $R_{n,+} := \text{diag}(R_{n,+}^+, R_{n,+}^-)$ ,  $\kappa_{n,+} := \kappa_{\max}(R_{n,+})$  and  $d_{n,+} := d_{\max}(R_{n,+}, \kappa_{n,+})$ ; cf. Definition 4.3.

# 6.2 Results

Turning to the results, it is natural to begin by stating a rough Sobolev estimate. However, for reasons mentioned above, we can only expect to be able to estimate the projection of solutions to the subspace corresponding to  $\mathcal{I}_{B,n}$ . This is the subject of Subsection 6.2.1. After that, we turn to the problem of deriving asymptotics. As mentioned above, the asymptotic behaviour can roughly speaking be divided into two pieces: an oscillatory part and an overall growth/decay. The oscillatory part can be expressed in terms of solutions to (6.6); i.e., the homogeneous wave equation obtained by dropping the right hand side of (1.2) as well as all the lower order terms on the left hand side. In Subsection 6.2.2, we state the asymptotics for solutions to this equation. Moreover, we note that there is a homeomorphism between initial data and asymptotic data. An important observation that arises in the study of (6.6) is that solutions (with frequency content contained in  $\mathcal{I}_{B,n}$ ) can be divided into two pieces, one of which is said to be positively oriented, and the other of which is said to be negatively oriented.

Turning to the asymptotics of solutions to (1.2), we restrict our attention to the situation that the right hand side exhibits slower growth than solutions to the homogeneous equation. Then the solutions can be approximated by a sum of functions of the form  $e^{At}u_w$ , where A is a constant matrix and  $u_w$  is a solution to (6.6). However, there are two types of orientation that influence what the matrix A is: the orientation of solutions to (6.6) mentioned above; and the sign of  $n_j$  (assuming the metric g associated with (1.2) has a geometric dominant noisy spatial S<sup>1</sup>-direction corresponding to j). We describe the relevant results in Subsection 6.2.3. Finally, in Subsection 6.2.4, we turn to the problem of specifying the asymptotics.

#### 6.2. RESULTS

#### 6.2.1 A rough Sobolev estimate

Next, we formulate a rough estimate of Sobolev type energies. In the statement, we use the following notation. If  $f \in C^{\infty}(M, \mathbb{C}^m)$ ,

$$f_{n}(p,t) := \sum_{\iota \in \mathcal{I}_{B,n}} \hat{f}(\iota,t)\varphi_{\iota}(p), \qquad (6.18)$$

where  $\varphi_{\iota}$  and  $\hat{f}$  are defined by (24.6) and (24.15) respectively. We define  $u_n$  similarly, and  $\mathfrak{E}_s$  by (24.19).

**Proposition 6.8.** Assume that (1.2) is  $C^2$ -balanced with a geometric dominant noisy spatial direction, convergent dominant coefficients and a negligible shift vector field; cf. Definition 6.5. Then there are constants C > 0 and  $s_n \ge 0$  such that if u is a solution to (1.2), then

$$\mathfrak{E}_{s}^{1/2}[u_{n}](t) \leq C\langle t \rangle^{d_{n,+}-1} e^{\kappa_{n,+}t} \mathfrak{E}_{s+s_{n}}^{1/2}[u_{n}](0) + \int_{0}^{t} C\langle t - t' \rangle^{d_{n,+}-1} e^{\kappa_{n,+}(t-t')} \|f_{n}(\cdot,t')\|_{(s+s_{n})} dt'$$
(6.19)

for all  $t \ge 0$  and  $s \in \mathbb{R}$ , where  $f_n$  and  $u_n$  are defined by (6.18) and the adjacent text. Finally,  $\kappa_{n,+}$  and  $d_{n,+}$  are defined in the text adjacent to (6.17).

**Remarks 6.9.** The proposition is a consequence of Remark 17.23. Moreover, C and  $s_n$  only depend on the Riemannian manifolds  $(M_r, g_r)$ ,  $r = 1, \ldots, R$ , and the coefficients of the equation (1.2).

### 6.2.2 A model for the oscillations

As a preparation for the description of the asymptotics of solutions to (6.6), we need to introduce some terminology. Assume, first of all, that (6.6) is  $C^2$ -balanced with a geometric dominant noisy spatial direction, convergent dominant coefficients and a negligible shift vector field; cf. Definition 6.5. We then define  $\eta_A$  by

$$\eta_A := \min\{\eta_n/2, \eta_{\rm sh}, \eta_{\rm mn}, \beta_n\},\tag{6.20}$$

where  $\beta_n$ ,  $\eta_n$  and  $\eta_{mn}$  are the constants appearing in Definitions 6.1 and 6.4; and  $\eta_{sh}$  is the constant appearing in Definition 6.5. Note that, in the present setting,  $\eta_{mn}$  could be removed from the definition of  $\eta_A$ , since  $\alpha = 0$  and  $X^l = 0$  for (6.6). However, we also wish to use the definition (6.20) in the study of solutions to (1.2), and in that setting, we need to include  $\eta_{mn}$ . In order to describe the oscillations of the solutions, the following two functions are of central importance:

$$\omega_{\rm sh}(\iota, t) := \int_0^t \sigma(\iota, t') \mathfrak{g}(\iota, t') dt', \qquad (6.21)$$

$$\varphi_{\text{tot}}(\iota, t) := \int_0^t [1 + \sigma^2(\iota, t')]^{1/2} \mathfrak{g}(\iota, t') dt', \qquad (6.22)$$

where  $\mathfrak{g}$  is defined by (24.13) and  $\sigma$  is defined by (24.17). The main result concerning the asymptotics of solutions to (6.6) is the following.

**Proposition 6.10.** Consider the equation (6.6). Assume that it is  $C^2$ -balanced with a geometric dominant noisy spatial direction, convergent dominant coefficients and a negligible shift vector field; cf. Definition 6.5. Let  $\eta_A$  be defined by (6.20) and  $\chi_{\pm}(\iota) \in \mathbb{C}$ ,  $\iota \in \mathcal{I}_{B,n}$ , be such that for every  $0 \leq k \in \mathbb{Z}$ , there is a constant  $C_k < \infty$  with the property that

$$\sum_{\iota \in \mathcal{I}_{B,n}} \langle \nu(\iota) \rangle^{2k} [|\chi_{+}(\iota)|^{2} + |\chi_{-}(\iota)|^{2}] \le C_{k}.$$
(6.23)

Define  $z_{app}$  and  $u_{app}$  by

$$z_{\rm app}(\iota,t) := \frac{e^{\beta_{\rm n}t/2}}{\mathfrak{g}(\iota,t)} \left( \chi_+(\iota) e^{i[\omega_{\rm sh}(\iota,t) - \varphi_{\rm tot}(\iota,t)]} + \chi_-(\iota) e^{i[\omega_{\rm sh}(\iota,t) + \varphi_{\rm tot}(\iota,t)]} \right)$$
(6.24)

$$u_{\rm app}(p,t) := \sum_{\iota \in \mathcal{I}_{\rm B,n}} z_{\rm app}(\iota,t) \varphi_{\iota}(p), \tag{6.25}$$

where  $\iota \in \mathcal{I}_{B,n}$  in (6.24);  $\omega_{sh}$  is defined by (6.21);  $\varphi_{tot}$  is defined by (6.22); and  $\mathfrak{g}(\iota, t)$  is defined by (24.13). Then there is a unique smooth solution u to (6.6) such that the following holds:  $u = u_n$ , where  $u_n$  is defined in analogy with (6.18); and there are constants  $s \in \mathbb{R}$ ,  $\eta > 0$  and C > 0 such that

$$\mathfrak{E}_s[u - u_{\mathrm{app}}](t) \le C e^{(\beta_n - \eta)t} \tag{6.26}$$

for all  $t \geq 0$ . Moreover, this unique solution has the property that

$$\mathfrak{E}_{s}[u - u_{\mathrm{app}}](t) \le C_{A} e^{(\beta_{\mathrm{n}} - 2\eta_{A})t} \sum_{\iota \in \mathcal{I}_{\mathrm{B},\mathrm{n}}} \langle \nu(\iota) \rangle^{2(s+s_{\mathrm{m}})} [|\chi_{+}(\iota)|^{2} + |\chi_{-}(\iota)|^{2}]$$
(6.27)

for all  $t \ge 0$  and  $s \in \mathbb{R}$ , where  $0 \le s_m \in \mathbb{R}$  and  $0 < C_A \in \mathbb{R}$  only depend on the coefficients of the equation (6.6) and the Riemannian manifolds  $(M_r, g_r), r = 1, \ldots, R$ ; and  $\mathfrak{E}_s$  is defined by (24.19). Finally,

$$\mathfrak{E}_{s}^{1/2}[u](0) \le C_{A} \left( \sum_{\iota \in \mathcal{I}_{\mathrm{B},\mathrm{n}}} \langle \nu(\iota) \rangle^{2(s+s_{\mathrm{m}})} [|\chi_{+}(\iota)|^{2} + |\chi_{-}(\iota)|^{2}] \right)^{1/2}$$
(6.28)

for all  $s \in \mathbb{R}$ , where  $C_A$  and  $s_m$  have the same dependence as in the case of (6.27).

Conversely, assume that u is a smooth solution to (6.6) such that  $u = u_n$ . Then there are uniquely determined  $\chi_{\pm}(\iota)$ ,  $\iota \in \mathcal{I}_{B,n}$ , such that (6.23) holds, and such that if  $u_{app}$  is defined by (6.24) and (6.25), then (6.26) holds. Moreover, this  $u_{app}$  is such that (6.27) holds. Finally,

$$\left(\sum_{\iota \in \mathcal{I}_{\mathrm{B,n}}} \langle \nu(\iota) \rangle^{2s} [|\chi_{+}(\iota)|^{2} + |\chi_{-}(\iota)|^{2}]\right)^{1/2} \le C_{A} \mathfrak{E}_{s+s_{\mathrm{m}}}^{1/2} [u](0)$$
(6.29)

for all  $s \in \mathbb{R}$ , where  $C_A$  and  $s_m$  have the same dependence as in the case of (6.27).

**Remark 6.11.** The proposition is a consequence of Remark 19.2.

**Remark 6.12.** Specifying  $\chi_{\pm}(\iota) \in \mathbb{C}$ ,  $\iota \in \mathcal{I}_{B,n}$ , such that (6.23) holds for all  $k \geq 0$  corresponds to specifying two smooth  $\mathbb{C}$ -valued functions on  $\overline{M}$  with frequency content contained in  $\mathcal{I}_{B,n}$ . These two functions can be thought of as the asymptotic data. Moreover, from this point of view, the estimates (6.28) and (6.29) imply that the map from initial data to asymptotic data is a homeomorphism with respect to the  $C^{\infty}$ -topology.

**Remark 6.13.** Since  $\mathfrak{g}(\iota, t) \approx \nu_n(\iota)e^{\beta_n t}$ , (6.24) implies that  $z_{app}(\iota, t)$  decays as  $e^{-\beta_n t/2}$ . However, the energy of  $u_{app}$  grows as  $e^{\beta_n t}$ . The reason for this is the following. First of all,

$$\dot{z}_{\rm app}(\iota,t) \approx e^{\beta_{\rm n}t/2} \left( -i\chi_{+}(\iota)e^{i[\omega_{\rm sh}(\iota,t)-\varphi_{\rm tot}(\iota,t)]} + i\chi_{-}(\iota)e^{i[\omega_{\rm sh}(\iota,t)+\varphi_{\rm tot}(\iota,t)]} \right)$$
$$\mathfrak{g}(\iota,t)z_{\rm app}(\iota,t) = e^{\beta_{\rm n}t/2} \left( \chi_{+}(\iota)e^{i[\omega_{\rm sh}(\iota,t)-\varphi_{\rm tot}(\iota,t)]} + \chi_{-}(\iota)e^{i[\omega_{\rm sh}(\iota,t)+\varphi_{\rm tot}(\iota,t)]} \right).$$

Thus

$$|\dot{z}_{\rm app}(\iota,t)|^2 + \mathfrak{g}^2(\iota,t)|z_{\rm app}(\iota,t)|^2 \approx 2e^{\beta_{\rm n}t}[|\chi_+(\iota)|^2 + |\chi_-(\iota)|^2].$$

In particular, the contribution of the  $\iota$ 'th mode to  $\mathfrak{E}_s[u_{app}]$  grows as  $e^{\beta_n t}$  unless the  $\chi_{\pm}(\iota)$  vanish.

**Remark 6.14.** The estimate (6.27) immediately yields an estimate for the  $L^2$ -norm of the difference between u and  $u_{app}$ ; cf. (24.19). However, this estimate is not very good. In order to improve it, assume that there is a non-negative continuous function  $\mathfrak{e}_{low} \in L^1([0,\infty))$  such that

$$\bar{k} \ge -[\beta_{\rm n} + \mathfrak{e}_{\rm low}(t)]\bar{g} \tag{6.30}$$
#### 6.2. RESULTS

for all  $t \ge 0$  (in addition to the assumptions of the proposition). Then there is a constant  $0 < C \in \mathbb{R}$ (depending only on the spectrum of the Riemannian manifold corresponding to the dominant noisy spatial direction and the coefficients of the equation (6.6)) such that  $e^{\beta_n t}[\mathfrak{g}(\iota, t)]^{-1} \le C$  for all  $t \ge 0$ and all  $\iota \in \mathcal{I}_{B,n}$ ; cf. Remarks 26.7 and 26.10. As a consequence,

$$e^{2\beta_{n}t}\sum_{\iota\in\mathcal{I}_{B}}\langle\nu(\iota)\rangle^{2s}|z(\iota,t)-z_{app}(\iota,t)|^{2}$$
$$=\sum_{\iota\in\mathcal{I}_{B}}e^{2\beta_{n}t}[\mathfrak{g}(\iota,t)]^{-2}\langle\nu(\iota)\rangle^{2s}\mathfrak{g}^{2}(\iota,t)|z(\iota,t)-z_{app}(\iota,t)|^{2}\leq C\mathfrak{E}_{s}[u-u_{app}].$$

Combining this estimate with (6.27) yields

$$\sum_{\iota \in \mathcal{I}_{B}} \langle \nu(\iota) \rangle^{2s} |z(\iota, t) - z_{app}(\iota, t)|^{2} \\ \leq C_{A} e^{-(\beta_{n} + 2\eta_{A})t} \sum_{\iota \in \mathcal{I}_{B,n}} \langle \nu(\iota) \rangle^{2(s+s_{m})} [|\chi_{+}(\iota)|^{2} + |\chi_{-}(\iota)|^{2}]$$
(6.31)

for all  $t \ge 0$  and  $s \in \mathbb{R}$ , where  $C_A$  and  $s_m$  have the same dependence as in the case of (6.27).

Due to this proposition, we can introduce a notion of orientation of a solution to (6.6).

**Definition 6.15.** Consider the equation (6.6). Assume that it is  $C^2$ -balanced with a geometric dominant noisy spatial direction, convergent dominant coefficients and a negligible shift vector field; cf. Definition 6.5. Assume that u is a smooth solution to (6.6) such that  $u = u_n$ , where  $u_n$  is defined in analogy with (6.18). Let  $u_{app}$  be the uniquely associated function of the form (6.24) and (6.25), whose existence is guaranteed by Proposition 6.10. Consider (6.24) and (6.25). If  $\chi_{-}(\iota) = 0$  for all  $\iota \in \mathcal{I}_{B,n}$ , then u is said to be *positively oriented*. If  $\chi_{+}(\iota) = 0$  for all  $\iota \in \mathcal{I}_{B,n}$ , then u is said to be *negatively oriented*.

In order to illustrate the use of Proposition 6.10, let us consider a special case.

**Example 6.16.** Consider (1.13). In the case of this equation,

$$g = -dt \otimes dt + e^{-2t} d\theta \otimes d\theta, \quad \bar{g} = e^{-2t} d\theta \otimes d\theta, \quad \bar{k} = -\bar{g}, \quad \mathcal{L}_U \bar{k} = 2\bar{g}.$$

Moreover, the shift vector field vanishes (as does  $\alpha$ ,  $\zeta$  and  $\mathcal{X}$ ). Thus (1.13) is  $C^2$ -balanced. Since  $\bar{k} + \bar{g} = 0$ , it is clear that (1.13) has a geometric dominant noisy spatial S<sup>1</sup>-direction corresponding to 1. Moreover,  $\beta_n = 1$  and we can choose  $C_n = 1$  and  $\eta_n = 2$ . That the dominant coefficients are convergent is obvious (since  $\mathcal{X}$  and  $\alpha$  vanish), and we can choose  $\eta_{mn} = 2$ . Finally, (1.13) has a negligible shift vector field, and we can choose  $\eta_{sh} = 2$ . Considering (6.20), it is clear that  $\eta_A = 1$ . Turning to  $\omega_{sh}$  and  $\varphi_{tot}$ , note that  $\omega_{sh} = 0$ ,  $\mathfrak{g}(n,t) = e^t |n|$  and  $\varphi_{tot}(n,t) = (e^t - 1)|n|$ . Thus  $z_{app}$  appearing in (6.24) can be written

$$z_{\rm app}(n,t) = |n|^{-1} e^{-t/2} \left( \chi_+(n) \exp[-i(e^t - 1)|n|] + \chi_-(n) \exp[i(e^t - 1)|n|] \right).$$

This means that  $u_{app}$  can be written in the form

$$u_{\rm app}(\theta, t) = e^{-t/2} u_{\rm w}(\theta, e^t),$$

where  $u_{\rm w}$  is a solution to the flat space wave equation  $u_{\tau\tau} - u_{\theta\theta} = 0$  with mean value zero for all  $\tau$  (in the present setting, the requirement that  $u_{\rm w}$  have mean value zero corresponds to the frequency content being contained in  $\mathcal{I}_{\rm B,n}$ ). Moreover, there is a homeomorphism (in the  $C^{\infty}$ topology) between the initial data for P (assuming that the mean value of P over  $\mathbb{S}^1$  vanishes) and the initial data for  $u_{\rm w}$ . Turning to the estimates of the difference between  $u_{\rm app}$  and P, note that (6.31) implies that every Sobolev norm of  $P(\cdot, t) - u_{\rm app}(\cdot, t)$  decays as  $e^{-3t/2}$ . Consider

$$P_t(\theta, t) - \partial_t u_{\rm app}(\theta, t) = P_t(\theta, t) - e^{t/2} (\partial_\tau u_{\rm w})(\theta, e^t) + \frac{1}{2} e^{-t/2} u_{\rm w}(\theta, e^t).$$
(6.32)

Due to (6.27), the Sobolev norm of this difference decays as  $e^{-t/2}$ . However, the Sobolev norm of the last term on the right hand side of (6.32) also decays as  $e^{-t/2}$ , so that the sum of the first two

terms on the right hand side of (6.32) decays as  $e^{-t/2}$ . Changing time coordinate to  $\tau = e^t$ , we obtain, by abuse of notation,

$$\|P(\cdot,\tau) - \tau^{-1/2} u_{\mathbf{w}}(\cdot,\tau)\|_{C^{k}} + \|P_{\tau}(\cdot,\tau) - \tau^{-1/2} \partial_{\tau} u_{\mathbf{w}}(\cdot,\tau)\|_{C^{k}} \le C_{k} \tau^{-3/2}$$

for all  $0 \le k \in \mathbb{Z}$  and all  $\tau \ge 1$ . Moreover, there is a homeomorphism between the initial data for P and the initial data for  $u_w$ . Adding to these observations the spatially homogeneous solutions to (1.25) yields the conclusions stated in connection with (1.25). Finally, let us note that the fact that there is a bijection follows from [20, 30].

#### 6.2.3 Deriving asymptotics

Next, we consider the asymptotics of solutions to (1.2). As already mentioned, a given solution can be approximated by a sum of terms consisting of an oscillatory part and an overall growth/decay. However, what the overall growth/decay is depends on the orientation of the oscillatory part, as well as the sign of  $n_j$  (assuming (1.2) has a geometric dominant noisy spatial S<sup>1</sup>-direction corresponding to j); here the oscillatory part is modelled by a solution to (6.6) and the notion of an orientation for a solution to (6.6) is given by Definition 6.15. It is therefore convenient to introduce the following notation.

**Definition 6.17.** Assume that (1.2) is  $C^2$ -balanced with a geometric dominant noisy spatial direction, convergent dominant coefficients and a negligible shift vector field; cf. Definition 6.5. In case (1.2) has a geometric dominant noisy spatial  $S^1$ -direction corresponding to j, let

$$\mathcal{I}_{\mathrm{B},\mathrm{n}}^{+} := \{ \iota \in \mathcal{I}_{\mathrm{B},\mathrm{n}} : \nu_{\mathrm{T},j}(\iota) < 0 \}, \quad \mathcal{I}_{\mathrm{B},\mathrm{n}}^{-} := \{ \iota \in \mathcal{I}_{\mathrm{B},\mathrm{n}} : \nu_{\mathrm{T},j}(\iota) > 0 \};$$
(6.33)

cf. (24.5). In case (1.2) has a geometric dominant noisy spatial direction corresponding to, say,  $M_r$ , let

$$\mathcal{I}_{B,n}^{+} := \mathcal{I}_{B,n}, \quad \mathcal{I}_{B,n}^{-} := \varnothing.$$
(6.34)

**Proposition 6.18.** Assume that (1.2) is  $C^2$ -balanced with a geometric dominant noisy spatial direction, convergent dominant coefficients and a negligible shift vector field; cf. Definition 6.5. Let  $\eta_A$  be defined by (6.20) and assume that there is an  $0 < \eta_B \leq \eta_A$  such that

$$\int_{0}^{\infty} e^{-(\kappa_{n,+} - \eta_B)t} \|f_n(\cdot, t)\|_{(s)} dt < \infty$$
(6.35)

for all s, where  $\kappa_{n,+}$  is defined by the text adjacent to (6.17);  $f_n$  is defined by (6.18); and  $\|\cdot\|_{(s)}$  is defined by (24.10). Let

$$A_{\mathbf{n},\pm} := -\frac{1}{2} \left( \alpha_{\infty} \pm \tilde{X}_{\infty,\mathbf{n}} \right), \quad \kappa_{\mathbf{n},\mathbf{m}} := \max\{\kappa_{\max}(A_{\mathbf{n},+}), \kappa_{\max}(A_{\mathbf{n},-})\},$$

where  $\tilde{X}_{\infty,n}$  is defined by (6.14) and (6.15). Finally, let  $I_n := (\kappa_{n,m} - \eta_B, \kappa_{n,m}]$  and  $E_{n,\pm} := E_{A_{n,\pm},I_n}$ ; cf. Definition 4.7.

If the metric g has a geometric dominant noisy spatial direction corresponding to, say,  $M_r$ , cf. Definition 6.4, then  $E_{n,+} = E_{n,-} =: E_n$  and  $A_{n,+} = A_{n,-} =: A_n$ . Moreover, if u is a solution to (1.2), there is a unique solution  $u_{0,w} \in C^{\infty}(M, E_n)$  to (6.6) such that

- if  $z_{0,w}(\iota, t)$  denotes the  $\iota$ 'th Fourier coefficient of  $u_{0,w}(\cdot, t)$ , then  $z_{0,w}(\iota, \cdot) = 0$  if  $\iota \notin \mathcal{I}_{B,n}$ ,
- if  $u_w$  is defined by

$$u_{\rm w}(p,t) = e^{A_{\rm n}t} u_{0,\rm w}(p,t), \tag{6.36}$$

then there are real constants C, N and s such that

$$\mathfrak{E}_s^{1/2}[u_{\mathrm{n}} - u_{\mathrm{w}}](t) \leq C \langle t \rangle^N e^{(\kappa_{\mathrm{n},+} - \eta_B)t}$$
(6.37)

for all  $t \ge 0$ , where  $u_n$  is defined in analogy with (6.18).

#### 6.2. RESULTS

If the metric g has a geometric dominant noisy spatial  $\mathbb{S}^1$ -direction corresponding to j and u is a solution to (1.2), then there are unique solutions  $u_{\pm,w} \in C^{\infty}(M, E_{n,\pm})$  to (6.6) such that the following holds:

•  $u_{+,w} = u_{+,w}^+ + u_{-,w}^-$  and  $u_{-,w} = u_{+,w}^- + u_{-,w}^+$ , where  $u_{\tau,w}^{\upsilon}$ ,  $\tau, \upsilon \in \{+, -\}$ , are smooth solutions to (6.6) whose *i*'th Fourier coefficients are denoted by  $z_{\tau,w}^{\upsilon}(\iota, t)$  and are such that

$$\begin{aligned} - z_{\pm,\mathrm{w}}^+(\iota,\cdot) &= 0 \text{ in case } \iota \notin \mathcal{I}_{\mathrm{B,n}}^+ \text{ and } z_{\pm,\mathrm{w}}^-(\iota,\cdot) = 0 \text{ in case } \iota \notin \mathcal{I}_{\mathrm{B,n}}^-, \\ - u_{+,\mathrm{w}}^{\pm} \text{ are positively oriented and } u_{-,\mathrm{w}}^{\pm} \text{ are negatively oriented solutions to (6.6),} \end{aligned}$$

• if  $u_w$  is defined by

$$u_{\rm w}(p,t) = e^{A_{\rm n,+}t} u_{+,\rm w}(p,t) + e^{A_{\rm n,-}t} u_{-,\rm w}(p,t), \tag{6.38}$$

then there are real constants C, N and s such that

$$\mathfrak{E}_s^{1/2}[u_{\mathbf{n}} - u_{\mathbf{w}}](t) \leq C \langle t \rangle^N e^{(\kappa_{\mathbf{n},+} - \eta_B)t}$$
(6.39)

for all  $t \geq 0$ .

Moreover, in both cases there are constants  $C_B > 0$  and  $s_B \ge 0$  (depending only on  $\eta_B$ , the Riemannian manifolds  $(M_r, g_r)$ , r = 1, ..., R, and the coefficients of the equation (1.2)); and a non-negative integer N (depending only on m) such that the following holds. If u and  $u_w$  are as above, then

$$\mathfrak{E}_{s}^{1/2}[u_{n}-u_{w}](t) \leq C_{B}\langle t\rangle^{N}e^{(\kappa_{n,+}-\eta_{B})t}\left(\mathfrak{E}_{s+s_{B}}^{1/2}[u_{n}](0)+\int_{0}^{\infty}e^{-(\kappa_{n,+}-\eta_{B})t'}\|f_{n}(\cdot,t')\|_{(s+s_{B})}dt'\right)$$
(6.40)

for all  $t \ge 0$  and all  $s \in \mathbb{R}$ . In addition, if the metric g has a geometric dominant noisy spatial direction corresponding to, say,  $M_r$ , then

$$\mathfrak{E}_{s}^{1/2}[u_{0,\mathbf{w}}](0) \le C_{B}\left(\mathfrak{E}_{s+s_{B}}^{1/2}[u_{n}](0) + \int_{0}^{\infty} e^{-(\kappa_{n,+}-\eta_{B})t'} \|f_{n}(\cdot,t')\|_{(s+s_{B})} dt'\right)$$
(6.41)

for all  $s \in \mathbb{R}$ , where  $C_B$  and  $s_B$  have the same dependence as in the case of (6.40). Finally, if the metric g has a geometric dominant noisy spatial  $\mathbb{S}^1$ -direction corresponding to j, then

$$\mathfrak{E}_{s}^{1/2}[u_{\pm,\mathbf{w}}](0) \le C_{B}\left(\mathfrak{E}_{s+s_{B}}^{1/2}[u_{n}](0) + \int_{0}^{\infty} e^{-(\kappa_{n,+}-\eta_{B})t'} \|f_{n}(\cdot,t')\|_{(s+s_{B})} dt'\right)$$
(6.42)

for all  $s \in \mathbb{R}$ , where  $C_B$  and  $s_B$  have the same dependence as in the case of (6.40).

Remark 6.19. The result is a consequence of Lemma 19.11; cf. Remark 19.12.

**Remark 6.20.** The asymptotics can be characterised in terms of functions such as  $u_{app}$  introduced in (6.24) and (6.25) (as opposed to solutions to (6.6)); cf. Remark 19.19.

**Remark 6.21.** Note that  $\kappa_{n,+}$  and  $\kappa_{n,m}$  are related according to  $\kappa_{n,+} = \kappa_{n,m} + \beta_n/2$ .

**Remarks 6.22.** In the statement of the proposition, two equations play an important role: (1.2) and (6.6). Note that it is here taken for granted that (6.6) is the equation obtained from (1.2) by setting  $\alpha$ ,  $\zeta$ ,  $X^j$  and f to zero. In the statement of the present proposition, we speak of  $\eta_A$ , defined by (6.20). Note that this  $\eta_A$  need not necessarily coincide with the  $\eta_A$  associated with the equation (6.6), say  $\eta_{A,\text{hom}}$ , since  $\eta_{\text{mn}}$  can be omitted from the right hand side of (6.20) in the definition of  $\eta_{A,\text{hom}}$ . On the other hand, it is clear that  $\eta_{A,\text{hom}} \geq \eta_A$ , so that when we appeal to Proposition 6.10, the discrepancy does not cause complications. Turning to the constant  $\beta_n$ , note that it is the same for the two equations; cf. Definitions 6.1 and 6.4.

**Remark 6.23.** If (1.2) has a dominant noisy spatial generalised direction, then  $A_n = -\alpha_{\infty}/2$ , so that (6.36) can be written

$$u_{\mathbf{w}}(p,t) = e^{-\alpha_{\infty}t/2}u_{0,\mathbf{w}}(p,t).$$

**Remark 6.24.** The constants C, s and N appearing in (6.37) and (6.39) are allowed to depend on the coefficients of the equation (1.2),  $\eta_B$ , the functions  $u_n$ ,  $u_w$ ,  $f_n$  etc.

**Remark 6.25.** If all the Jordan blocks of the matrices  $A_{n,\pm}$  are trivial, then the N appearing in (6.40) can be replaced by 1. If  $E_{n,+} = E_{n,-} = \mathbb{C}^m$ , then N can be replaced by  $d_{n,+} - 1$ , where  $d_{n,+}$  is defined in connection with (6.17).

#### 6.2.4 Specifying the asymptotics

The result of the previous subsection yields conclusions concerning the asymptotics, given a solution. In fact, (6.41) and (6.42) demonstrate that there is a continuous map from initial data to asymptotic data. It is then of interest to ask if it is possible to construct a solution, given asymptotic data. If there is a corresponding map, it is also natural to ask whether it is continuous. In the preparation for the statement of such a result, it is convenient to introduce the following terminology.

**Definition 6.26.** Consider the equation (6.6). Assume that it is  $C^2$ -balanced with a geometric dominant noisy spatial direction, convergent dominant coefficients and a negligible shift vector field; cf. Definition 6.5. Let  $1 \le m \in \mathbb{Z}$  and V be a vector subspace of  $\mathbb{C}^m$ . Then

$$W_{\rm n}(\bar{M}, V) \subset C^{\infty}(\bar{M}, V) \times C^{\infty}(\bar{M}, V)$$

denotes the set of initial data at t = 0 for solutions u to (6.6) such that  $u \in C^{\infty}(M, V)$  and the  $\iota$ 'th Fourier coefficient of  $u(\cdot, t)$  vanishes for all t if  $\iota \notin \mathcal{I}_{B,n}$ . Moreover,  $W_{n,+}(\bar{M}, V)$  ( $W_{n,-}(\bar{M}, V)$ ) denotes the subset of  $W_n(\bar{M}, V)$  corresponding to positively (negatively) oriented solutions to (6.6). In addition,  $W_{n,\pm}^+(\bar{M}, V)$  ( $W_{n,\pm}^-(\bar{M}, V)$ ) denotes the subset of  $W_{n,\pm}(\bar{M}, V)$  consisting of functions whose  $\iota$ 'th Fourier coefficients vanish if  $\iota \notin \mathcal{I}_{B,n}^+$  ( $\iota \notin \mathcal{I}_{B,n}^-$ ).

**Remark 6.27.** By initial data we here mean  $[u(\cdot, 0), u_t(\cdot, 0)]$ .

**Proposition 6.28.** Assume that (1.2) is  $C^2$ -balanced with a geometric dominant noisy spatial direction, convergent dominant coefficients and a negligible shift vector field; cf. Definition 6.5. Assume, moreover, that f = 0. Let  $\eta_A$  be defined by (6.20) and  $0 < \eta_B \le \eta_A$ . Define  $A_{n,\pm}$  and  $E_{n,\pm}$  as in the statement of Proposition 6.18.

If the metric g has a geometric dominant noisy spatial direction corresponding to, say,  $M_r$ , then  $E_{n,+} = E_{n,-} =: E_n$ , and there is an injective linear map

$$\Phi_{n,g}: W_n(\bar{M}, E_n) \to W_n(\bar{M}, \mathbb{C}^m)$$

such that if  $\psi \in W_n(\bar{M}, E_n)$ ;  $u_{0,w}$  is the solution to (6.6) corresponding to the initial data  $\psi$  at t = 0;  $u_w$  is defined by (6.36); and u is the solution to (1.2) (with f = 0) corresponding to the initial data  $\Phi_{n,g}(\psi)$  at t = 0, then  $u = u_n$  and

$$\mathfrak{E}_s^{1/2}[u-u_{\mathrm{w}}](t) \le C_B \langle t \rangle^N e^{(\kappa_{\mathrm{n},+}-\eta_B)t} \|\psi\|_{(s+s_B)}$$
(6.43)

for all  $t \ge 0$  and  $s \in \mathbb{R}$ , where  $C_B$ ,  $s_B$  and N have the same dependence as in the case of (6.40). If the Jordan blocks of the matrix  $A_n$  are trivial, the N appearing in (6.43) can be replaced by 1. Moreover, if  $E_n = \mathbb{C}^m$ , then N can be replaced by  $d_{n,+} - 1$ , where  $d_{n,+}$  is defined in connection with (6.17). If  $E_n = \mathbb{C}^m$ , then  $\Phi_{n,g}$  is surjective. Finally, there are constants  $C_B > 0$  and  $s_B \ge 0$ such that

$$\|\Phi_{n,g}(\psi)\|_{(s)} \le C_B \|\psi\|_{(s+s_B)} \tag{6.44}$$

for all  $s \in \mathbb{R}$  and all  $\psi \in W_n(\overline{M}, E_n)$ , where  $C_B$  and  $s_B$  have the same dependence as in the case of (6.40).

If the metric g has a geometric dominant noisy spatial  $\mathbb{S}^1$ -direction corresponding to j, let

$$\mathcal{W}_{n}^{+} := W_{n,+}^{+}(\bar{M}, E_{n,+}) \oplus W_{n,-}^{-}(\bar{M}, E_{n,+}),$$
  
$$\mathcal{W}_{n}^{-} := W_{n,+}^{-}(\bar{M}, E_{n,-}) \oplus W_{n,-}^{+}(\bar{M}, E_{n,-}).$$

Then there is an injective linear map

$$\Phi_{\mathrm{n,T}}: \mathcal{W}_{\mathrm{n}}^{+} \times \mathcal{W}_{\mathrm{n}}^{-} \to W_{\mathrm{n}}(\bar{M}, \mathbb{C}^{m})$$

such that if  $\chi_{\pm} \in \mathcal{W}_{n}^{\pm}$ ;  $u_{\pm,w}$  is the solution to (6.6) corresponding to the initial data  $\chi_{\pm}$  at t = 0;  $u_{w}$  is defined by (6.38); and u is the solution to (1.2) (with f = 0) corresponding to the initial data  $\Phi_{n,T}(\chi_{+}, \chi_{-})$  at t = 0, then  $u = u_{n}$  and

$$\mathfrak{E}_{s}^{1/2}[u-u_{\mathbf{w}}](t) \leq C_{B}\langle t \rangle^{N} e^{(\kappa_{\mathbf{n},+}-\eta_{B})t}[\|\chi_{+}\|_{(s+s_{B})} + \|\chi_{-}\|_{(s+s_{B})}]$$
(6.45)

for all  $t \ge 0$  and  $s \in \mathbb{R}$ , where  $C_B$ ,  $s_B$  and N have the same dependence as in the case of (6.40). If the Jordan blocks of the matrices  $A_{n,\pm}$  are trivial, the N appearing in (6.45) can be replaced by 1. Moreover, if  $E_{n,+} = E_{n,-} = \mathbb{C}^m$ , then N can be replaced by  $d_{n,+} - 1$ , where  $d_{n,+}$  is defined in connection with (6.17). If  $E_{n,+} = E_{n,-} = \mathbb{C}^m$ , then  $\Phi_{n,T}$  is surjective. Finally, there are constants  $C_B > 0$  and  $s_B \ge 0$  such that

$$\|\Phi_{\mathbf{n},\mathbf{T}}(\chi_{+},\chi_{-})\|_{(s)} \le C_{B}[\|\chi_{+}\|_{(s+s_{B})} + \|\chi_{-}\|_{(s+s_{B})}]$$
(6.46)

for all  $s \in \mathbb{R}$  and all  $(\chi_+, \chi_-) \in \mathcal{W}_n^+ \times \mathcal{W}_n^-$ , where  $C_B$  and  $s_B$  have the same dependence as in the case of (6.40).

Remark 6.29. The result is a consequence of Lemma 19.22; cf. Remark 19.24.

**Remark 6.30.** The asymptotic data can be specified in terms of functions such as  $u_{app}$  introduced in (6.24) and (6.25) (as opposed to solutions to (6.6)); cf. Remark 19.26.

**Remark 6.31.** Remarks 6.22 are equally relevant in the present setting.

## 6.2.5 Reinterpreting oscillatory ODE behaviour; equations with unbounded $\zeta$

In most of these notes, we assume  $\zeta$  to be future bounded. However, there are some situations in which this requirement can be omitted. To be more specific, assume that there is a real valued function  $\xi_2 \in C^{\infty}(I, \mathbb{R})$  and real numbers  $\xi_{2,\infty}, \eta_{\text{aux}}, C_{\text{aux}} > 0$  such that

$$|\xi_2(t) - \xi_{2,\infty}| + |\partial_t \xi_2(t)| \le C_{\text{aux}} e^{-\eta_{\text{aux}} t}$$
(6.47)

for all  $t \geq 0$ . Assume, moreover,  $\zeta$  to be of the form

$$\zeta(t) = \zeta_1(t) + \zeta_2(t) \operatorname{Id}_m, \quad \zeta_2(t) := \xi_2(t) e^{2\beta_{\operatorname{aux}} t}$$

for all  $t \ge 0$ , where  $\beta_{aux} > 0$  is a constant. Here  $\zeta_1$  should be thought of as being such that  $\|\zeta_1\|$ and  $\|\partial_t\zeta_1\|$  are bounded (however,  $\zeta_1$  need not converge). By redefining  $\zeta_1$  and  $\zeta_2$  (in such a way that they remain unchanged in a neighbourhood of  $t = \infty$ ), it can be ensured that  $\xi_2 \ge \xi_{2,\infty}/2$ for all  $t \ge 0$ . For that reason, we from now on assume this inequality to be satisfied. Returning to (1.2) with  $g^{00} = -1$ , note that it can be written

$$u_{tt} - g^{jl}\partial_j\partial_l u - 2g^{0l}\partial_l\partial_t u - \sum_{r=1}^R a_r^{-2}(t)\Delta_{g_r} u + \alpha u_t + X^j\partial_j u + \zeta_1 u + \zeta_2 u = f.$$
(6.48)

On the other hand, we can view this equation as being the equation for a Fourier coefficient of a solution to

$$U_{tt} - g^{jl} \partial_j \partial_l U - \zeta_2 \partial_{d+1}^2 U - 2g^{0l} \partial_l \partial_t U - \sum_{r=1}^R a_r^{-2}(t) \Delta_{g_r} U + \alpha U_t + X^j \partial_j U + \zeta_1 U = F \quad (6.49)$$

on  $M_{\text{aux}} := \overline{M}_{\text{aux}} \times I$ , where  $\overline{M}_{\text{aux}} := \mathbb{T}^{d+1} \times M_1 \times \cdots \times M_R$ . In fact, given a solution u to (6.48),  $U = ue^{ix^{d+1}}$  satisfies (6.49) with  $F = fe^{ix^{d+1}}$  (where  $x^{d+1}$  denotes the (d+1)'th coordinate on the (d+1)-torus). If we can derive asymptotic information concerning solutions to (6.49), we can thus do the same for solutions to (6.48). In fact, it is sufficient if we can derive asymptotic information concerning the parts of solutions to (6.49) corresponding to  $\iota$ 's such that  $n_{d+1} \neq 0$ . There are many situations in which the results of these notes could be applied to (6.49). However, for the sake of definiteness, let us consider one such situation.

**Dominant ODE oscillations.** Note that (6.49) is an equation of the form (1.2), where the metric is given by

$$g_{\mathrm{aux}} := g + \zeta_2^{-1} dx^{d+1} \otimes dx^{d+1}$$

on  $M_{\text{aux}}$ . Moreover,  $(M_{\text{aux}}, g_{\text{aux}})$  is a canonical separable cosmological model manifold, and if  $\bar{k}_{\text{aux}}$  is the second fundamental form associated with  $g_{\text{aux}}$ , then

$$\bar{k}_{aux} = \bar{k} - \left(\beta_{aux} + \frac{1}{2}\xi_2^{-1}\dot{\xi}_2\right)\zeta_2^{-1}dx^{d+1} \otimes dx^{d+1},$$

where  $\bar{k}$  is the second fundamental form of g. Assume that there is a constant  $\eta_n > 0$  and a continuous  $\mathfrak{e} \in L^1([0,\infty))$  such that

$$\bar{k}(v,v) \ge [-\beta_{\text{aux}} + \eta_{\text{n}} - \mathfrak{e}(t)]\bar{g}(v,v)$$
(6.50)

for all  $v \in T\bar{M}$  and  $t \geq 0$ . Define  $\bar{M}_{sub} := \bar{M}; \pi_n : \bar{M}_{aux} \to \mathbb{S}^1$  to be the projection onto the (d+1)'th  $\mathbb{S}^1$ -factor in  $\bar{M}_{aux}; \pi_{sub}$  to be the map from  $\bar{M}_{aux}$  to  $\bar{M}_{sub}$  corresponding to the projection onto what remains after removing the (d+1)'th  $\mathbb{S}^1$ -factor; and let  $\beta_n := \beta_{aux}$ . Then (6.47) ensures that (6.7) holds. Moreover, (6.50) ensures that (6.8) holds. Thus  $g_{aux}$  has a geometric dominant noisy spatial  $\mathbb{S}^1$ -direction corresponding to d+1. Since  $X^{d+1} = 0$  in (6.49), the requirement that (6.9) hold is equivalent to the requirement that

$$\|\alpha(t) - \alpha_{\infty}\| \le C_{\mathrm{mn}} e^{-\eta_{\mathrm{mn}} t} \tag{6.51}$$

for all  $t \ge 0$ . From now on, we therefore assume (6.51) to hold, so that the dominant coefficients are convergent; cf. Definition 6.1. Turning to the shift vector field, note that it is the same for gand  $g_{\text{aux}}$ , and assume that (6.11) holds for all  $t \ge 0$ . What remains to be verified in order to ensure that the conditions of Definition 6.5 are satisfied is that (6.49) is  $C^2$ -balanced. Assuming the shift vector field of g to be  $C^2$ -future bounded;  $\alpha$  and  $\zeta_1$  to be  $C^1$ -future bounded;  $\mathcal{X}$  to be  $C^1$ -future bounded;  $\bar{k}$  to be  $C^1$ -future bounded; and (in addition to (6.47)),  $\ddot{\xi}_2$  to be future bounded, then (6.49) is  $C^2$ -balanced. Making appropriate assumptions concerning f, the results of the present section thus apply to (6.49).

**Example 6.32.** Consider the Klein-Gordon equation in the expanding direction of Kasner spacetimes. Due to the calculations carried out in Example 4.21, the Klein-Gordon equation can be written

$$t\partial_t(tu_t) - \sum_{i=1}^d t^{2\beta_i} u_{ii} + m^2 t^2 u = 0$$

where  $\beta_i = 1 - p_i$ . Letting  $\tau = \ln t$ , this equation becomes

$$u_{\tau\tau} - \sum_{i=1}^{d} e^{2\beta_i \tau} u_{ii} + m^2 e^{2\tau} u = 0$$

In the discussion above, this equation corresponds to (6.48). The corresponding version of (6.49) is

$$U_{\tau\tau} - \sum_{i=1}^{d} e^{2\beta_i \tau} U_{ii} - m^2 e^{2\tau} U_{DD} = 0, \qquad (6.52)$$

where D := d + 1. If the  $\beta_i$  are all distinct and different from 1, we can think of (6.52) as being defined on  $\mathbb{T}^D \times \mathbb{R}$ . If there are several  $\beta_i$ 's equalling, say,  $\beta_1$ , then we can combine the corresponding circles to a  $\mathbb{T}^{d_1} =: M_1$  etc. Due to these observations, we can apply the methods developed so far in these notes to (6.52). Doing so yields a homeomorphism between initial data and asymptotic data (we leave the details of the verification of this statement to the reader).

#### 6.3 Outline of the proof

Fix a Fourier mode  $\iota \in \mathcal{I}_{B,n}$ . In one respect, the analysis of the asymptotics of a corresponding Fourier coefficient is similar to the analysis in the silent and transparent cases:

- For large t, say  $t \ge T_n$ ,  $\mathfrak{g}(\iota, t)$  is well approximated by  $\nu_n(\iota)e^{\beta_n t}$ , and (24.16) can effectively be replaced by a limit equation.
- For small t, i.e.  $t \in [0, T_n]$ , the solution can be controlled by a crude energy estimate.

The main difference between the noisy setting on the one hand, and the silent and transparent settings on the other, is that the limit equation in the noisy setting is not a constant coefficient equation, so that quite different methods are needed in order to analyse the behaviour of solutions for  $t \ge T_n$ . The fact that a crude energy estimate is sufficient to handle the interval  $[0, T_n]$  is a strength in the sense that the conditions needed to obtain the results are weak. On the other hand, the crude analysis leads to a loss of derivatives, which makes it impossible to calculate  $\eta_{nl}$ .

The noisy regime. Due to the above observations, it is clear that the main difficulty consists in analysing the behaviour of solutions in the regime  $[T_n, \infty)$ . In the introduction to the present chapter, we give a rough idea of how to approximate the corresponding oscillatory behaviour in the case of (6.1). In Part III, we develop general methods for approximating the evolution over one period of the oscillations. However, the methods developed in Part III of these notes apply not only to the equations considered in Chapter 2 and the equations considered in the present chapter. If we make additional assumptions concerning the equations, they also apply in regions of the form  $[0, T_{ode}]$ ,  $[0, T_{ts}]$  and  $[0, T_n]$  in the silent, transparent and noisy settings respectively. Moreover, in Chapter 7 below, we demonstrate that it is possible to calculate  $\eta_{nl}$  on the basis of such an application. To summarise: since the full range of the applications of the results of Part III only becomes clear at the end of the next chapter, we only give an outline of the methods developed in Part III at the end of Chapter 7. However, let us describe the main outcome of the analysis. The goal is to estimate the evolution of solutions in intervals where the behaviour is oscillatory. To this end, we introduce a time sequence  $\{t_k\}$  such that  $[t_k, t_{k+1}]$ , roughly speaking, corresponds to one period of the oscillations. The final result of Part III is (15.63), an equation that yields

$$\psi_{k+1} = A_k^+ \psi_k + A_k^+ \int_{t_k}^{t_{k+1}} \hat{F}_k(t) dt.$$
(6.53)

Here  $\psi_k$  represents the initial data for the Fourier coefficient at  $t_k$  (the exact relation is somewhat technical, and we refer the reader interested in the details to Part III). Moreover,  $A_k^+$  is the matrix taking initial data at  $t_k$  to initial data at  $t_{k+1}$  in the homogeneous setting. Finally,  $\hat{F}_k(t)$  is a reformulated version of the contribution from  $\hat{f}(\iota, t)$ . That a relation of the form (6.53) holds is an immediate consequence of the fact that the underlying equation is a linear system of ODE's. However, in Part III, we provide approximations for  $A_k^+$ , and estimate the error. Moreover, the approximations can be expressed in terms of the coefficients of (1.2).

**Estimates along a time sequence.** A rough Sobolev estimate. The main part of the arguments needed to prove the results of the present chapter is contained in Parts III and V of these notes. Focusing on Part V, we begin, in Sections 17.2 and 17.3, by providing an analytic version of the definition of a noisy equation (the conditions are such that they are implied by the

72

geometric conditions imposed in the current chapter). Moreover, we verify that the conditions needed in order to apply the results of Part III are satisfied, and we define  $T_n$ . In Section 17.4, we adapt the general framework of Part III to the context of interest here. In particular, we estimate  $A_k^+$  appearing in (6.53) in terms of  $\alpha_{\infty}$ ,  $\beta_n$  and  $\bar{X}_n(\iota)$ . Given this approximation, we reformulate the iteration (6.53) in order to isolate the leading order behaviour; this is the subject of Section 17.5. On the basis of this reformulation, we are then in a position to estimate the growth of the individual Fourier coefficients. This is the subject of Section 17.6. Combining this estimate for the Fourier coefficients with a rough energy estimate in the interval  $[0, T_n]$  and summing up over the modes yields the Sobolev estimate (6.19); cf. Section 17.7.

Asymptotics along a time sequence. In Section 18.1, we turn to the problem of deriving the asymptotic behaviour along a time sequence. The argument is quite long and technical, but in some respects, it is a discrete version of the corresponding argument in the silent setting. Note, however, that the analysis is only relevant for one mode and along a time sequence contained in  $[T_n, \infty)$ . In Section 18.2, we turn to the problem of specifying the asymptotic data along the time sequence. Again, the argument is long and technical, but roughly speaking a discrete version of the corresponding argument in the silent setting. Deriving and specifying asymptotics along a time sequence is interesting. However, there are two drawbacks: the time sequence is not canonically defined and depends on  $\iota$ ; and the conclusions do not tell us what happens between the times belonging to the sequence. In Section 18.3 we begin addressing these issues by estimating the evolution for all  $t \geq T_n$ . In order to obtain the desired conclusion, we, again, need to appeal to the results of Part III.

The results. Given the analysis described above, we are in a position to derive the main results of the present chapter; Propositions 6.10, 6.18 and 6.28. In Section 19.2 we demonstrate Proposition 6.10. Even though most of the ingredients are already in place, we need to demonstrate that we can specify the asymptotics for the individual modes (in the particular case of interest); we need to sum up over the modes; and, given a solution, we need to derive asymptotics. As a next step, we demonstrate a uniqueness result; cf. Section 19.3. The purpose of this result is to ensure that the uniqueness statement made in Proposition 6.18 holds. Proving this uniqueness statement is more complicated than proving the corresponding result in the silent setting. The reason for this is that in the silent setting, the decay rate on the right hand side of the estimate is strong enough to ensure uniqueness. However, in the noisy setting, the fact that uniqueness holds is dependent on the orientation of the solutions, and this complicates the proof. In Section 19.4 we prove Proposition 6.18: we verify the uniqueness statement; we derive estimates for the individual modes and late times; we extend the results to early times; we consider the approximate solutions and use the approximate solutions to construct solutions to the model equation; and finally we derive the desired estimates. We end by demonstrating that the asymptotic data can be specified in Section 19.5.

## Chapter 7

# Energy estimates in the asymptotically diagonal setting

#### 7.1 Introduction

Weaknesses of the results. The results of Chapters 4–6 may, superficially, seem quite satisfactory. First of all, they yield optimal energy estimates in the sense that they allow us to calculate  $\eta_{cr}$ . Second, they yield detailed asymptotics and, third, they allow us to specify the leading order asymptotics. In fact, we sometimes obtain a homeomorphism between initial data and asymptotic data. In other words, if one is only interested in the linear systems of equations discussed in these notes, then the results are quite informative. However, our main motivation for studying these equations is that they arise as linearised versions of Einstein's equations. Moreover, in the original non-linear setting, the question of stability is of interest. Assume, therefore, that the homogeneous version of (1.2) is the linearised version of a non-linear equation. We would then like to know that if u is a solution with initially small energy, then the energy of u remains small. In order to evaluate whether the results of, say, Chapter 4 allow us to draw such conclusions, say that the conditions of Proposition 4.4 are satisfied and that f = 0. Assume, moreover, that  $d_1 = 1$ and that  $\kappa_1 = -4711$ . For a smooth solution u, Proposition 4.4 then guarantees that there is a  $0 < C \in \mathbb{R}$  such that

$$\mathfrak{E}^{1/2}[u](t) \le C e^{-4711t} \tag{7.1}$$

for all  $t \ge 0$ ; recall that  $\mathfrak{E} = \mathfrak{E}_0 = \mathfrak{E}_{\text{bas}}$ . In particular, the energy is decaying exponentially, so that the solution will eventually be small. Let  $0 < \epsilon \in \mathbb{R}$  be given, and assume that  $\mathfrak{E}^{1/2}[u](0) \le \epsilon$ . Assuming  $\epsilon$  to be small, we know  $\mathfrak{E}^{1/2}[u]$  to be small initially and to decay exponentially in the sense that there is a constant such that (7.1) holds. Does the combination of these two observations ensure that  $\mathfrak{E}$  is always small? Unfortunately, the answer to this question is no. Given two constants  $0 < \epsilon$ ,  $N \in \mathbb{R}$ , there is an equation satisfying the above restrictions and a corresponding solution u such that  $\mathfrak{E}^{1/2}[u](0) \le \epsilon$ ; such that (7.1) holds for some C and all  $t \ge 0$ ; and such that there is a  $0 < t_1 \in \mathbb{R}$  such that  $\mathfrak{E}^{1/2}[u](t_1) \ge N$ . In this sense, the results of Chapter 4 are not very useful when addressing the issue of stability. On the other hand, assuming  $\mathfrak{E}_{s_0}[u](0) \le C^{-2}\epsilon^2$ , where  $s_0$  and C are the constants appearing in the statement of Proposition 4.4, the estimate  $\mathfrak{E}^{1/2}[u](t) \le \epsilon e^{-4711t}$  holds for all  $t \ge 0$ . However, this result involves a loss of derivatives, and, depending on the context, this may substantially reduce its use. For a justification of the above statements, we refer the reader to Example 7.23 below.

**Calculating**  $\eta_{\rm nl}$ . The deficiencies mentioned above are related to our inability to calculate  $\eta_{\rm nl}$  in the previous chapters. Moreover, this inability is related to the loss of derivatives in, e.g., Propositions 4.4 and 6.8. Finally, this loss of derivatives is related to the rough energy estimates we carry out in time intervals of the form  $[0, T_{\rm ode}]$ ,  $[0, T_{\rm ts}]$  and  $[0, T_{\rm n}]$  for a fixed mode; cf. the

descriptions of the arguments in the previous chapters. In order to calculate  $\eta_{nl}$ , we need to improve the analysis in these intervals. However, in order to be able to do so, we need to make stronger assumptions. In Section 7.2 we introduce the relevant conditions. However, it is useful to consider a few examples before stating the results; this is the subject of Subsection 7.1.1. We also give an outline of the present chapter in Subsection 7.1.2.

#### 7.1.1 Examples

In order to contrast the results of Chapter 4 and the results of the present chapter with standard energy estimates, let us consider the following two examples.

Example 7.1. Consider

$$u_{tt} - e^{-2t}u_{\theta\theta} + 2e^{-t}u_{\theta} + u_t + u = 0.$$
(7.2)

This equation is such that the results of Chapter 4 apply. In fact,  $\bar{k} = \bar{g}$  and  $\chi = 0$ , so that (7.2) is  $C^1$ -future silent in the sense of Definition 4.1, with  $\mu = 1$ . Moreover,  $\mathcal{X} = 2e^{-t}\partial_{\theta}$  is  $C^0$ -future bounded, and since  $\alpha = 1$  and  $\zeta = 1$ , (4.2) holds with  $C_{\rm mn} = \eta_{\rm mn} = 1$ . In particular, the eigenvalues of  $A_{\infty}$  introduced in (4.4) all have real part  $\kappa_1 = -1/2$  and the corresponding Jordan blocks are trivial. Moreover, Proposition 4.4 applies and yields the conclusion that there are constants C and  $s_0$  such that

$$\mathfrak{E}_{s}^{1/2}[u](t) \le Ce^{-t/2}\mathfrak{E}_{s+s_{0}}^{1/2}[u](0) \tag{7.3}$$

for all  $t \ge 0$  and all  $s \in \mathbb{R}$ . Since the energy of spatially homogeneous solutions decays as  $e^{-t}$ , this estimate is optimal as far as the time dependence is concerned, and  $\eta_{cr} = -1/2$ .

On the other hand, considering the problem from the point of view of standard energy estimates, let

$$\mathfrak{E}_{\text{bas}}[u](t) := \frac{1}{2} \int_{\mathbb{S}^1} [|u_t(\theta, t)|^2 + e^{-2t} |u_\theta(\theta, t)|^2 + |u(\theta, t)|^2] d\theta.$$
(7.4)

It can then be calculated that

$$\frac{d\mathfrak{E}_{\text{bas}}[u]}{dt} = -\int_{\mathbb{S}^1} |u_t + e^{-t} u_\theta|^2 d\theta.$$
(7.5)

Clearly,  $\mathfrak{E}_{\text{bas}}$  is a decreasing function. On the other hand, we can chose initial data so that the right hand side of (7.5) vanishes. In general, we therefore cannot obtain a better estimate than  $\partial_t \mathfrak{E}_{\text{bas}} \leq 0$ . From this point of view, it seems optimistic to even expect decay. Clearly, there is a tension between this observation and the estimate (7.3). However, it turns out that  $\eta_{nl} = 0$ ; this is a consequence of Theorem 7.21 below. In other words, if  $a \in \mathbb{R}$  and  $0 < C \in \mathbb{R}$  are such that

$$\mathfrak{E}_{\mathrm{bas}}[u](t) \le C e^{2at} \mathfrak{E}_{\mathrm{bas}}[u](0)$$

for all  $t \ge 0$  and all solutions u to (7.2), then  $a \ge 0$ . To conclude, what the decay rate is depends on how many derivatives one is prepared to lose. The estimate (7.3) is the best as far as decay is concerned, but the estimate  $\mathfrak{E}_{\text{bas}}[u](t) \le \mathfrak{E}_{\text{bas}}[u](0)$  for all  $t \ge 0$  is the best estimate if one is not prepared to lose derivatives.

In the case of (7.2), the equality (7.5) yields a good indication of the value of  $\eta_{nl}$ . In order to illustrate that the situation is sometimes more complicated, we consider the following example.

Example 7.2. Consider the equation

$$u_{tt} - e^{-2t}u_{\theta\theta} + ae^{-t}u_{\theta} + bu_t = 0, (7.6)$$

where  $a, b \in \mathbb{R}$  and b < 0. In what follows, we focus on real valued solutions. Due to the absence of an undifferentiated u, it is convenient to consider the energy

$$\mathfrak{E}_{\hom}[u](t) := \frac{1}{2} \int_{\mathbb{S}^1} [u_t^2(\theta, t) + e^{-2t} u_{\theta}^2(\theta, t)] d\theta.$$

#### 7.1. INTRODUCTION

Due to (7.6),

$$\frac{d\mathfrak{E}_{\text{hom}}[u]}{dt} = \int_{\mathbb{S}^1} (-ae^{-t}u_\theta u_t - bu_t^2 - e^{-2t}u_\theta^2)d\theta.$$

The integrand is a quadratic form in  $u_t$  and  $e^{-t}u_{\theta}$ . The largest eigenvalue of the corresponding matrix is

$$\lambda_{\max} = \frac{1}{2} \left[ -(b+1) + \sqrt{(b-1)^2 + a^2} \right].$$

Note that if b < 0 and  $a \neq 0$ , then  $\lambda_{\max} > -b$ . For a given  $t_0$ , there are non-vanishing initial data at  $t_0$  such that

$$\partial_t \mathfrak{E}_{\text{hom}}(t_0) = 2\lambda_{\max} \mathfrak{E}_{\text{hom}}(t_0).$$

Hoping for a better estimate than  $\mathfrak{E}_{\text{hom}}(t) \leq e^{2\lambda_{\max}t}\mathfrak{E}_{\text{hom}}(0)$  thus seems optimistic. On the other hand, it can be calculated that for the equation (7.6),

$$\eta_{\rm nl} = \frac{1}{2} \max\{-b - 1 + |a|, -2b\},\tag{7.7}$$

assuming b < 0; cf. Example 7.24 below. In fact, we demonstrate that there is a constant C such that for every solution u to (7.6), the energy  $\mathfrak{E}_{\text{bas}}$  introduced in (7.4) satisfies

$$\mathfrak{E}_{\mathrm{bas}}[u](t) \le C e^{2\eta_{\mathrm{nl}}t} \mathfrak{E}_{\mathrm{bas}}[u](0) \tag{7.8}$$

for all  $t \ge 0$ . Note that, as long as b < 0 and  $a \ne 0$ ,  $\eta_{nl} < \lambda_{max}$ . Since  $\mathfrak{E}_{hom} \le \mathfrak{E}_{bas}$ , it follows that  $\mathfrak{E}_{hom}(t) \le Ce^{2\eta_{nl}t}\mathfrak{E}_{bas}(0)$ . This yields a better estimate for the energy  $\mathfrak{E}_{hom}$  than we would naively have expected. Of course, there is a loss involved in that  $\mathfrak{E}_{bas}(0) \ge \mathfrak{E}_{hom}(0)$ . On the other hand,  $\mathfrak{E}_{bas}(0)$  involves the same number of derivatives of the initial data as  $\mathfrak{E}_{hom}(0)$ . In this sense, the loss is not severe. Even though this argument does not prove that energy estimates cannot be used to obtain optimal conclusions, they do indicate that crude energy estimates sometimes do not suffice.

#### 7.1.2 Outline

As indicated by the title of the chapter, we are here interested in metrics that become diagonal asymptotically. A precise explanation of what this means is given in Definition 7.3 below. However, loosely speaking, there are two main conditions. First, in each  $\mathbb{S}^1$  and  $M_r$  direction, the second fundamental form converges; cf. (7.15) and (7.16) below. Second, the different  $\mathbb{S}^1$ -directions are becoming exponentially more orthogonal with time in the sense that (7.17) and (7.18) below hold for some suitable choice of k. In the present chapter, we also assume that the shift vector field becomes negligible asymptotically, in the sense that (7.19) below holds. Adding to these conditions on the metric the assumption that the lower order coefficients are convergent, in the sense that (7.20) below holds, and the assumption that (1.2) is  $C^2$ -balanced, we obtain the conclusion that the equation (1.2) asymptotically takes the form

$$u_{tt} - \sum_{j} g_{\infty}^{jj} e^{2\beta_j t} u_{jj} - \sum_{r} a_{r,\infty}^{-2} e^{2\beta_{\mathrm{Ri},r} t} \Delta_{g_r} u + \sum_{j} e^{\beta_j t} X_{\infty}^j u_j + \alpha_{\infty} u_t + \zeta_{\infty} u = f,$$
(7.9)

where  $u_t = \partial_t u$ ,  $u_j = \partial_j u$  etc.; cf. Section 7.2 for a justification of this statement. Here  $0 < g_{\infty}^{jj}, a_{r,\infty} \in \mathbb{R}, \beta_j, \beta_{\text{Ri},r} \in \mathbb{R}$  and  $X_{\infty}^j, \alpha_{\infty}, \zeta_{\infty} \in \mathbf{M}_m(\mathbb{C})$ . In order to develop some intuition for the results, let us consider the limit equation (7.9).

The Fourier modes. Introducing

$$\mathfrak{g}_{\infty}(\iota, t) := \left(\sum_{j} g_{\infty}^{jj} e^{2\beta_{j}t} n_{j}^{2} + \sum_{r} a_{r,\infty}^{-2} e^{2\beta_{\mathrm{Ri},r}t} \nu_{r,i_{r}}^{2}(\iota)\right)^{1/2},\tag{7.10}$$

the equation for the Fourier coefficients of solutions to the limit equation can be written

$$\ddot{z} + \mathfrak{g}_{\infty}^2 z + \sum_j i n_j e^{\beta_j t} X_{\infty}^j z + \alpha_{\infty} \dot{z} + \zeta_{\infty} z = \hat{f}.$$

$$(7.11)$$

In the present chapter, we focus on the non-degenerate setting, where the set consisting of the union of the  $\beta_j$ 's and the  $\beta_{\text{Ri},r}$ 's contains Q := d + R distinct elements. Under this assumption, the interval  $[0, \infty)$  can be divided into, at most, Q + 1 intervals. Each interval corresponds to one of the functions

1, 
$$(g_{\infty}^{jj})^{1/2} |n_j| e^{\beta_j t}$$
,  $a_{r,\infty}^{-1} \nu_{r,i_r}(\iota) e^{\beta_{\mathrm{Ri},r} t}$ 

being larger than all the others, and we refer to these intervals as *eras*. Let us now assume that  $(g_{\infty}^{jj})^{1/2}|n_j|e^{\beta_j t}$  dominates. Naively, it then seems reasonable to replace (7.11) by

$$\ddot{z} + g_{\infty}^{jj} e^{2\beta_j t} n_j^2 z + i n_j e^{\beta_j t} X_{\infty}^j z + \alpha_{\infty} \dot{z} = \hat{f}$$

$$(7.12)$$

(no summation on j). Here we have discarded the term  $\zeta_{\infty} z$  since  $(g_{\infty}^{jj})^{1/2} |n_j| e^{\beta_j t} \ge 1$  by assumption. When f = 0, the equation (7.12) is identical to (6.1) if we introduce the notation

$$\nu_{\mathbf{n}}(\iota) := (g_{\infty}^{jj})^{1/2} |n_j|, \quad \beta_{\mathbf{n}} := \beta_j, \quad \bar{X}_{\mathbf{n}}(\iota) := \frac{n_j}{|n_j|} (g_{\infty}^{jj})^{-1/2} X_{\infty}^j$$

If  $\beta_j > 0$ , we can thus appeal to the line of reasoning presented in Subsection 6.1.1 in order to conclude that the growth/decay of solutions should be determined by the matrices

$$\frac{1}{2} \left( \beta_j \mathrm{Id}_m - \alpha_\infty \pm (g_\infty^{jj})^{-1/2} X_\infty^j \right); \tag{7.13}$$

cf. (6.5). In fact, the same line of reasoning works in the case that  $\beta_j < 0$ . In other words, as long as  $\beta_j \neq 0$ , the largest real part of an eigenvalue of a matrix of the form (7.13) can naively be expected to determine the overall growth of the Fourier coefficient in the era during which  $(g_{\infty}^{jj})^{1/2}|n_j|e^{\beta_j t}$  dominates. In the case that  $a_{r,\infty}^{-1}\nu_{r,i_r}(\iota)e^{\beta_{\text{Ri},r}t}$  dominates with  $\beta_{\text{Ri},r} \neq 0$ , a similar argument applies. In an interval where 1 dominates, the largest real part of an eigenvalue of the matrix  $A_{\infty}$  introduced in (4.4) determines the overall growth/decay.

The transparent eras. Consider an era corresponding to  $\beta_j \neq 0$ ,  $\beta_{\text{Ri},r} \neq 0$  or 1. Then there is one matrix that determines the overall growth/decay rate. Now consider an era with  $\beta_j = 0$  or  $\beta_{\text{Ri},r} = 0$ ; from now on, we refer to such eras as transparent eras. In the case of a transparent era, the term  $\zeta_{\infty} z$  has to be included in the limit equation. In the interval where, say,  $(g_{\infty}^{jj})^{1/2} |n_j| e^{\beta_j t}$ dominates (and  $\beta_j = 0$ ), the limit equation thus becomes

$$\ddot{z} + g_{\infty}^{jj} n_j^2 z + i n_j X_{\infty}^j z + \alpha_{\infty} \dot{z} + \zeta_{\infty} z = \hat{f}.$$
(7.14)

This is a family of infinitely many different systems of constant coefficient ODE's. For each member of the family, there is an associated generic growth rate for solutions to the corresponding homogeneous equation. On the basis of the results of the previous chapters, we expect the supremum of these growth rates over all  $0 \neq n_i \in \mathbb{Z}$  to be a lower bound for  $\eta_{nl}$ .

**Outline of the chapter.** In Section 7.2 below, we introduce the conditions that characterise the asymptotically diagonal equations. First, we impose conditions on the geometry, cf. Definition 7.3; and then we impose conditions on the lower order coefficients, cf. Definition 7.5. However, in order for the results of the present chapter to apply, we also need to make a non-degeneracy assumption; cf. Definition 7.6. Finally, we need to assume that the equation is balanced. For convenience, we summarise the assumptions in Definition 7.8.

The growth/decay rates corresponding to the different eras. In Section 7.3, we introduce the constituent growth/decay rates that determine  $\eta_{nl}$ . The constituents are of three types. First, there is the ODE part, corresponding to the maximal growth rate of solutions to the homogeneous version of (4.3), where  $A_{\infty}$  is given by (4.4). Second, there are the growth/decay rates given by the greatest real parts of eigenvalues of matrices of the form (7.13) (for *j*'s such that  $\beta_j \neq 0$ ). In addition, there are similar contributions from the corresponding matrices for *r*'s such that  $\beta_{\text{Ri},r} \neq 0$ . Third, there are the growth/decay rates of the solutions to the homogeneous versions of the supremum of the growth/decay rates of the solutions to the homogeneous versions.

of (7.14) (or the relevant analogous equations in case  $\beta_{\text{Ri},r} = 0$ ). Combining these definitions leads to the introduction of  $\kappa_{\text{tot},+}$ , cf. Definition 7.16. In the end, we are able to demonstrate that  $\eta_{\text{nl}} = \kappa_{\text{tot},+}$ ; cf. Theorem 7.21 below. In the energy estimates, we, in general, have to count on a loss of the form  $e^{\epsilon t}$ ; cf. (7.30). However, if (1.2) satisfies some additional non-degeneracy requirements, this loss can be removed; cf. (7.31), (7.32) and (7.33). In Definition 7.16, we introduce not only  $\kappa_{\text{tot},+}$ , but also the relevant non-degeneracy conditions.

Results/outline of the argument. Once we have introduced the necessary terminology, we are in a position to formulate the main results. In Section 7.4 we state the main energy estimates; cf. Theorem 7.19. Moreover, we demonstrate optimality and calculate  $\eta_{nl}$  in Section 7.4.1; cf. Theorem 7.21. We end the chapter, in Section 7.5, by giving an outline of the argument.

#### 7.2 Asymptotically diagonal equations

In addition to the requirements made in previous chapters, the main assumption concerning the metric in the present chapter is that it becomes asymptotically diagonal and is diagonally convergent. These notions are defined as follows.

**Definition 7.3.** Let (M, g) be a canonical separable cosmological model manifold. For each  $l \in \{1, \ldots, d\}$  let  $\pi_{\mathrm{T},l}$  be the map from  $\overline{M}$  to  $\mathbb{S}^1$  corresponding to projection onto the *l*'th  $\mathbb{S}^1$ -factor in  $\mathbb{T}^d$ . For each  $r \in \{1, \ldots, R\}$ , let  $\pi_{\mathrm{Ri},r}$  be the map from  $\overline{M}$  to  $M_r$  corresponding to projection onto  $M_r$ . Then  $\overline{k}$  is said to be *diagonally convergent* if there are real numbers  $\beta_j$  and  $\beta_{\mathrm{Ri},r}$ , where  $j \in \{1, \ldots, R\}$ , and constants  $C_d, \kappa_d > 0$  such that the following holds:

$$|\bar{k}^{\sharp}(v_j, v_j) + \beta_j \bar{g}^{\sharp}(v_j, v_j)| \leq C_{\mathrm{d}} e^{-\kappa_{\mathrm{d}} t} \bar{g}^{\sharp}(v_j, v_j), \tag{7.15}$$

$$\left|\bar{k}^{\sharp}(w_r, w_r) + \beta_{\mathrm{Ri},r}\bar{g}^{\sharp}(w_r, w_r)\right| \leq C_{\mathrm{d}} e^{-\kappa_{\mathrm{d}}t}\bar{g}^{\sharp}(w_r, w_r)$$
(7.16)

for all  $t \ge 0, j \in \{1, \dots, d\}, v_j \in \pi^*_{\mathrm{T}, j}(T^* \mathbb{S}^1), r \in \{1, \dots, R\}$ , and  $w_r \in \pi^*_{\mathrm{Ri}, r}(T^* M_r)$ .

In addition,  $\bar{g}$  is said to be  $C^0$ -asymptotically diagonal if there are constants  $C_{od}$ ,  $\kappa_{od} > 0$  such that

$$|\bar{g}^{\sharp}(v_j, v_l)| \le C_{\text{od}} e^{-\kappa_{\text{od}}t} |v_j|_{\bar{g}} |v_l|_{\bar{g}}$$

$$(7.17)$$

for all  $t \ge 0$ ;  $j, l \in \{1, \ldots, d\}$  such that  $j \ne l$ ;  $v_j \in \pi^*_{\mathrm{T},j}(T^*\mathbb{S}^1)$ ; and  $v_l \in \pi^*_{\mathrm{T},l}(T^*\mathbb{S}^1)$ . Similarly,  $\bar{g}$  is said to be  $C^k$ -asymptotically diagonal,  $1 \le k \in \mathbb{Z}$ , if it is  $C^0$ -asymptotically diagonal and there are constants  $C_{\mathrm{od},k}, \kappa_{\mathrm{od}} > 0$  such that

$$\sum_{i=0}^{k-1} |(\mathcal{L}_{U}^{i}\bar{k})^{\sharp}(v_{j},v_{l})| \leq C_{\mathrm{od},k}e^{-\kappa_{\mathrm{od}}t}|v_{j}|_{\bar{g}}|v_{l}|_{\bar{g}}$$
(7.18)

for all  $t \ge 0$ ;  $j, l \in \{1, ..., d\}$  such that  $j \ne l$ ;  $v_j \in \pi^*_{T,j}(T^*\mathbb{S}^1)$ ; and  $v_l \in \pi^*_{T,j}(T^*\mathbb{S}^1)$ .

Finally, if there are constants  $C_{\rm sh}$ ,  $\kappa_{\rm sh} > 0$  such that

$$|\chi(t)|_{\bar{g}} + |\dot{\chi}(t)|_{\bar{g}} \le C_{\rm sh} e^{-\kappa_{\rm sh}t}$$
(7.19)

for all  $t \ge 0$ , then the shift vector field is said to be *negligible*.

**Remark 7.4.** In (7.15)–(7.18), the symbol  $\sharp$  indicates raising all indices using  $\bar{g}$ ; in particular,  $\bar{g}^{\sharp}$ ,  $\bar{k}^{\sharp}$  and  $(\mathcal{L}_U \bar{k})^{\sharp}$  are the contravariant 2-tensor fields whose components are given by  $\bar{g}^{ij}$ ,  $\bar{k}^{ij}$  and  $(\mathcal{L}_U \bar{k})^{ij}$  respectively. Moreover, if  $\eta \in T^*M$ , then  $|\eta|_{\bar{g}} := (\bar{g}^{ij}\eta_i\eta_j)^{1/2}$ .

Beyond making assumptions concerning the metric appearing in (1.2), we also need to make assumptions concerning the lower order coefficients.

**Definition 7.5.** Consider (1.2). Assume the associated metric to be such that (M, g) is a canonical separable cosmological model manifold. Assume, moreover, the associated second fundamental

form  $\bar{k}$  to be diagonally convergent; cf. Definition 7.3. If there are constants  $0 < C_{\rm mn}, \kappa_{\rm mn} \in \mathbb{R}$ and matrices  $X^j_{\infty}, \alpha_{\infty}, \zeta_{\infty} \in \mathbf{M}_m(\mathbb{C}), j \in \{1, \ldots, d\}$ , such that

$$\|e^{-\beta_j t} X^j(t) - X^j_{\infty}\| + \|\alpha(t) - \alpha_{\infty}\| + \|\zeta(t) - \zeta_{\infty}\| \le C_{\rm mn} e^{-\kappa_{\rm mn} t}$$
(7.20)

for all  $t \ge 0$  and all  $j \in \{1, \ldots, d\}$ , then (1.2) is said to be such that the main coefficients are convergent; here the  $\beta_j$  appearing in (7.20) are the constants whose existence is guaranteed by Definition 7.3.

Consider (1.2). Assume the associated metric to be such that (M, g) is a canonical separable cosmological model manifold. Assume (1.2) to be  $C^2$ -balanced; cf. Definition 3.8. Assume, moreover, the shift vector field to be negligible,  $\bar{k}$  to be diagonally convergent and  $\bar{g}$  to be  $C^2$ -asymptotically diagonal; cf. Definition 7.3. Finally, assume (1.2) to be such that the main coefficients are convergent; cf. Definition 7.5. Then Lemma 26.12 applies. In particular,  $e^{-2\beta_j t}g^{jj}(t)$  converges to  $g_{\infty}^{jj} > 0$ ;  $e^{-2\beta_{\text{Ri},r}t}a_r^{-2}(t)$  converges to  $a_{r,\infty}^{-2} > 0$ ;  $e^{-\beta_j t}X^j(t)$  converges to  $X_{\infty}^j$ ;  $\alpha(t)$  converges to  $\alpha_{\infty}$ ; and  $\zeta(t)$  converges to  $\zeta_{\infty}$ . Moreover, the shift vector field  $g^{0l}$  and the off-diagonal components  $g^{jl}, j \neq l$ , are relatively speaking negligible. This leads to the limit equation (7.9).

Treating the  $\beta_j$  and the  $\beta_{\text{Ri},r}$  separately is inconvenient in the discussions to follow. We therefore introduce the following terminology.

**Definition 7.6.** Consider (1.2). Assume the associated metric to be such that (M, g) is a canonical separable cosmological model manifold. Assume, moreover,  $\bar{k}$  to be diagonally convergent; cf. Definition 7.3. Define  $\bar{\beta}_1 < \bar{\beta}_2 < \cdots < \bar{\beta}_Q$  to be the distinct elements of the union of the sets  $\{\beta_1, \ldots, \beta_d\}$  and  $\{\beta_{\text{Ri},1}, \ldots, \beta_{\text{Ri},R}\}$ , where  $\beta_j$  and  $\beta_{\text{Ri},r}$  are the numbers whose existence is guaranteed by Definition 7.3. If Q = R + d, the equation (1.2) is said to be *asymptotically non-degenerate*.

**Remark 7.7.** The numbers  $\beta_1, \ldots, \beta_d, \beta_{\text{Ri},1}, \ldots, \beta_{\text{Ri},R}$  are distinct if and only if the equality Q = d + R holds.

In what follows, we are mainly interested in asymptotically non-degenerate equations. In order to simplify the statement of the assumptions, we therefore introduce the following terminology.

**Definition 7.8.** The equation (1.2) is said to be geometrically non-degenerate, diagonally dominated, balanced and convergent if the following assumptions hold. First, the associated metric is such that (M, g) is a canonical separable cosmological model manifold. Second, (1.2) is  $C^2$ balanced; cf. Definition 3.8. Third, the shift vector field is negligible,  $\bar{k}$  is diagonally convergent and  $\bar{g}$  is  $C^2$ -asymptotically diagonal; cf. Definition 7.3. Fourth, (1.2) is such that the main coefficients are convergent; cf. Definition 7.5. Fifth, (1.2) is asymptotically non-degenerate; cf. Definition 7.6.

#### 7.3 The constituent growth/decay rates

Consider (1.2). Assume that it is geometrically non-degenerate, diagonally dominated, balanced and convergent; cf. Definition 7.8. It is then convenient to introduce the following terminology: For  $1 \leq q \leq Q$  and  $\iota \in \mathcal{I}_B$ , let

$$\bar{\nu}_{q}(\iota) := \left(\sum_{\{j:\beta_{j}=\bar{\beta}_{q}\}} g_{\infty}^{jj} n_{j}^{2} + \sum_{\{r:\beta_{\mathrm{Ri},r}=\bar{\beta}_{q}\}} a_{r,\infty}^{-2} \nu_{r,i_{r}}^{2}(\iota)\right)^{1/2}.$$
(7.21)

The following objects play a central role in the analysis to follow.

**Definition 7.9.** Assume that (1.2) is geometrically non-degenerate, diagonally dominated, balanced and convergent; cf. Definition 7.8. Given  $q \in \{1, \ldots, Q\}$ , define  $\tilde{X}_{\infty}^{q}$  as follows. If  $\bar{\beta}_{q} = \beta_{j}$  for some  $j \in \{1, \ldots, d\}$ , then j is uniquely determined by q and

$$\tilde{X}^{q}_{\infty} := (g^{jj}_{\infty})^{-1/2} X^{j}_{\infty}.$$
(7.22)

Moreover, if  $\iota \in \mathcal{I}_{\mathrm{B}}$  is such that  $\nu_{\mathrm{T},j}(\iota) = n_j \neq 0$ , then

$$\bar{X}_{\text{diag}}^q(\iota) := \frac{n_j}{|n_j|} \tilde{X}_{\infty}^q.$$

If  $\bar{\beta}_q = \beta_{\mathrm{Ri},r}$  for some  $r \in \{1, \ldots, R\}$ , then r is uniquely determined by q and  $\tilde{X}^q_{\infty} := 0$ . Moreover,  $\bar{X}^q_{\mathrm{diag}}(\iota) := 0$ . For  $q \in \{1, \ldots, Q\}$ , let

$$R_{q,\pm}^{+} := \pm \frac{1}{2} (-\alpha_{\infty} + \bar{\beta}_{q} \mathrm{Id}_{m} + \tilde{X}_{\infty}^{q}), \quad R_{q,\pm}^{-} := \pm \frac{1}{2} (-\alpha_{\infty} + \bar{\beta}_{q} \mathrm{Id}_{m} - \tilde{X}_{\infty}^{q}), \tag{7.23}$$

and let

$$\kappa_{q,\pm} := \max\{\kappa_{\max}(R_{q,\pm}^+), \kappa_{\max}(R_{q,\pm}^-)\},$$
(7.24)

$$d_{q,\pm} := d_{\max}(\operatorname{diag}\{R_{q,\pm}^+, R_{q,\pm}^-\}, \kappa_{q,\pm});$$
(7.25)

cf. Definition 4.3. Finally, let

$$\kappa_{\pm} := \max_{q} \kappa_{q,\pm}. \tag{7.26}$$

**Remark 7.10.** It is of interest to interpret the constituents of  $R_{q,+}^{\pm}$  geometrically. First of all, the  $\bar{\beta}_q$  are the asymptotic eigenvalues of the second fundamental form. Second, note that

$$\mathcal{Y} := \alpha \partial_t + X^j \partial_j$$

is the matrix of vector fields appearing in (1.2). Let  $E_j$  be the unit one-form field which is a positive multiple of  $dx^j$ . Then

$$\lim_{t \to \infty} U^{\flat}(\mathcal{Y}) = -\alpha_{\infty}, \quad \lim_{t \to \infty} E_j(\mathcal{Y}) = (g_{\infty}^{jj})^{-1/2} X_{\infty}^j.$$

Note that  $U^{\flat}$  is metrically equivalent to the future directed unit normal to the hypersurfaces  $\overline{M}_t$ . Moreover,  $\pm E_j$  are the two unit one-form fields generating  $\pi^*_{T,j}(T^*\mathbb{S}^1)$ . Since the sign corresponds to a choice of orientation of the j'th  $\mathbb{S}^1$ -factor, it is not surprising that  $\kappa_{\pm}$  does not depend on it.

**Remark 7.11.** The matrices appearing in (7.23) should be compared with those appearing in (7.13).

**Remark 7.12.** At this stage, it is not so clear why it would be of interest to consider  $R_{q,-}^{\pm}$ . However, the argument demonstrating that we can specify asymptotic data is based on an analysis of the evolution when going backwards in time. In that context, the matrices  $R_{q,-}^{\pm}$  are of importance.

The number  $\kappa_+$  plays an important role in determining the growth rate of the basic energy. However, in the case of equations with a  $q \in \{1, \ldots, Q\}$  such that  $\bar{\beta}_q = 0$ , we need to complement  $\kappa_+$  with additional information; cf. the discussion of transparent eras in Subsection 7.1.2.

**Definition 7.13.** Consider the equation (1.2). Assume that it is geometrically non-degenerate, diagonally dominated, balanced and convergent; cf. Definition 7.8. If there is a  $q \in \{1, \ldots, Q\}$  such that  $\bar{\beta}_q = 0$ , say  $q_{\rm ts}$ , define the sets  $\mathcal{K}_{q_{\rm ts},\pm}$  by

$$\mathcal{K}_{q_{\mathrm{ts}},\pm} := \left\{ \kappa_{q_{\mathrm{ts}},\iota,\pm} \mid \iota \in \mathcal{I}_{\mathrm{B}}, \ \bar{\nu}_{q_{\mathrm{ts}}}(\iota) \neq 0 \right\},\$$

where  $\bar{\nu}_q(\iota)$  is defined by (7.21),  $\kappa_{q_{\rm ts},\iota,\pm}$  is defined by

$$\kappa_{q_{\rm ts},\iota,\pm} := \kappa_{\rm max}[\pm A_{q_{\rm ts},\infty}(\iota)] \tag{7.27}$$

and

$$A_{q_{\rm ts},\infty}(\iota) := \begin{pmatrix} 0 & \bar{\nu}_{q_{\rm ts}}(\iota) \mathrm{Id}_m \\ -\bar{\nu}_{q_{\rm ts}}(\iota) \mathrm{Id}_m - i\bar{X}_{\mathrm{diag}}^{q_{\rm ts}}(\iota) - [\bar{\nu}_{q_{\rm ts}}(\iota)]^{-1}\zeta_\infty & -\alpha_\infty \end{pmatrix}.$$
 (7.28)

Define  $\bar{\kappa}_{q_{\rm ts},\pm}$  to be the limit of  $\kappa_{q_{\rm ts},\iota,\pm}$  as  $\bar{\nu}_{q_{\rm ts}}(\iota) \to \infty$  and define

$$\kappa_{q_{\rm ts},{\rm ts},\pm} := \sup \mathcal{K}_{q_{\rm ts},\pm}.$$

It is also convenient to introduce the notation

$$\kappa_{\mathrm{ts},\pm} := \max\{\kappa_{\pm}, \kappa_{q_{\mathrm{ts}},\mathrm{ts},\pm}\}\tag{7.29}$$

**Remark 7.14.** The matrix  $A_{q_{ts},\infty}(\iota)$  introduced in (7.28) should be compared with  $A(\iota)$  introduced in (5.14); cf. also (7.14).

**Remark 7.15.** That the stated limit,  $\bar{\kappa}_{q_{ts},\pm}$ , exists is a consequence of Remark 21.15. In this remark, we also calculate  $\bar{\kappa}_{q_{ts},\pm}$  in terms of  $\alpha_{\infty}$  and  $\tilde{X}_{\infty}^{q_{ts}}$ .

In Definition 7.9, we introduce the growth/decay rates associated with the eras with  $\bar{\beta}_q \neq 0$ ; and in Definition 7.13 we introduce the growth/decay rate associated with the transparent eras. Combining these definitions with the growth/decay rate corresponding to the spatially homogeneous solutions to (1.2) yields a number  $\kappa_{\text{tot},+}$ ; cf. Definition 7.16 below. Naively, we would hope  $\kappa_{\text{tot},+}$ to equal  $\eta_{\text{nl}}$ . In the end this turns out to be the case; cf. Theorem 7.21 below.

In some situations, it is possible to not only calculate  $\eta_{nl}$ , but also to obtain the correct polynomial rate; cf. (7.31)–(7.33) below. However, to obtain such conclusions, we need to make suitable non-degeneracy assumptions. The relevant conditions are the following.

**Definition 7.16.** Consider the equation (1.2). Assume that it is geometrically non-degenerate, diagonally dominated, balanced and convergent; cf. Definition 7.8. If there is a  $q_{ts} \in \{1, \ldots, Q\}$  such that  $\bar{\beta}_{q_{ts}} = 0$ , let  $\kappa_{tot,+} := \max\{\kappa_{ts,+}, \kappa_{sil,+}\}$ , where  $\kappa_{ts,+}$  is defined in (7.29);  $\kappa_{sil,+} := \kappa_{\max}(A_{\infty})$ ; and  $A_{\infty}$  is defined in (4.4). If there is no such  $q_{ts}$ , let  $\kappa_{tot,+} := \max\{\kappa_{+}, \kappa_{sil,+}\}$ , where  $\kappa_{sil,+}$  is defined as before and  $\kappa_{+}$  is defined by (7.26).

The equation (1.2) is said to exhibit subdominant block non-degeneracy provided that

- if there is a  $q_{ts} \in \{1, \ldots, Q\}$  such that  $\bar{\beta}_{q_{ts}} = 0$ , then  $\bar{\kappa}_{q_{ts},+} < \kappa_{tot,+}$ ,
- for  $q \in \{1, \ldots, Q\}$  such that  $\bar{\beta}_q < \max\{0, \bar{\beta}_Q\}$ , all the Jordan blocks of  $R_{q,+}^+$  and  $R_{q,+}^-$  corresponding to the eigenvalues with real part  $\kappa_{\text{tot},+}$  are trivial, and
- if  $\bar{\beta}_Q > 0$  and there is a  $q_{ts} \in \{1, \ldots, Q\}$  such that  $\bar{\beta}_{q_{ts}} = 0$ , then the Jordan blocks of  $A_{q_{ts},\infty}(\varsigma)$  corresponding to the eigenvalues with real part  $\kappa_{tot,+}$  are trivial for all  $\varsigma \in \mathcal{I}_B$  such that  $\bar{\nu}_{q_{ts}}(\varsigma) \neq 0$ .

**Remark 7.17.** Note that  $\bar{\beta}_Q \geq \bar{\beta}_q$  for all  $q \in \{1, \ldots, Q\}$ .

**Remark 7.18.** In Remark 22.5 below, we give an intuitive motivation for the conditions that characterise subdominant block non-degeneracy.

#### 7.4 Energy estimates

We are now in a position to state the main result of these notes.

**Theorem 7.19.** Consider the equation (1.2). Assume that it is geometrically non-degenerate, diagonally dominated, balanced and convergent; cf. Definition 7.8. Fix  $\epsilon > 0$ . Then there is a constant  $0 < C_{\epsilon} \in \mathbb{R}$ , depending only on  $\epsilon$ , the spectra of the Riemannian manifolds  $(M_r, g_r)$ ,  $r \in \{1, \ldots, R\}$ , and the coefficients of the equation (1.2), such that

$$\mathfrak{E}_{s}^{1/2}[u](t) \le C_{\epsilon} e^{(\kappa_{\text{tot},+}+\epsilon)t} \mathfrak{E}_{s}^{1/2}[u](0) + C_{\epsilon} \int_{0}^{t} e^{(\kappa_{\text{tot},+}+\epsilon)(t-t')} \|f(\cdot,t')\|_{(s)} dt'$$
(7.30)

for all  $t \ge 0$ , all solutions u to (1.2) and all  $s \in \mathbb{R}$ , where  $\kappa_{tot,+}$  is given by Definition 7.16. If, in addition to the above, the equation (1.2) exhibits subdominant block non-degeneracy, there are the following improvements of (7.30):

• If  $\bar{\beta}_Q < 0$ , then the estimate

$$\mathfrak{E}_{s}^{1/2}[u](t) \le C\langle t \rangle^{d_{\mathrm{sil},+}-1} e^{\kappa_{\mathrm{tot},+}t} \mathfrak{E}_{s}^{1/2}[u](0) + C \int_{0}^{t} \langle t - t' \rangle^{d_{\mathrm{sil},+}-1} e^{\kappa_{\mathrm{tot},+}(t-t')} \|f(\cdot,t')\|_{(s)} dt'$$
(7.31)

holds for all  $t \ge 0$ , all solutions u to (1.2) and all  $s \in \mathbb{R}$ . Here  $d_{\operatorname{sil},+} = d_{\max}(A_{\infty}, \kappa_{\operatorname{tot},+})$ ;  $A_{\infty}$  is defined in (4.4); and the constant C only depends on the spectra of the Riemannian manifolds  $(M_r, g_r), r \in \{1, \ldots, R\}$ , and the coefficients of the equation (1.2).

• If  $\bar{\beta}_Q = 0$ , then

$$\mathfrak{E}_{s}^{1/2}[u](t) \le C\langle t \rangle^{d_{\mathrm{sts},+}-1} e^{\kappa_{\mathrm{tot},+}t} \mathfrak{E}_{s}^{1/2}[u](0) + C \int_{0}^{t} \langle t - t' \rangle^{d_{\mathrm{sts},+}-1} e^{\kappa_{\mathrm{tot},+}(t-t')} \|f(\cdot,t')\|_{(s)} dt'$$
(7.32)

holds for all  $t \ge 0$ , all solutions u to (1.2) and all  $s \in \mathbb{R}$ . Here  $d_{\text{sts},+} := \max\{d_{\text{sil},+}, d_{t,+}\};$  $d_{\text{sil},+}$  is defined as in the case of (7.31);  $d_{t,+}$  is the largest dimension of a non-trivial Jordan block of a matrix of the form  $A_{Q,\infty}(\varsigma)$  (where  $\varsigma \in \mathcal{I}_{\text{B}}$  is such that  $\bar{\nu}_Q(\varsigma) \neq 0$ ) corresponding to an eigenvalue with real part  $\kappa_{\text{tot},+}$ ; and the constant C only depends on the spectra of the Riemannian manifolds  $(M_r, g_r), r \in \{1, \ldots, R\}$ , and the coefficients of the equation (1.2).

• If  $\bar{\beta}_Q > 0$ , then

$$\mathfrak{E}_{s}^{1/2}[u](t) \leq C\langle t \rangle^{d_{\text{tot},+}-1} e^{\kappa_{\text{tot},+}t} \mathfrak{E}_{s}^{1/2}[u](0) + C \int_{0}^{t} \langle t - t' \rangle^{d_{\text{tot},+}-1} e^{\kappa_{\text{tot},+}(t-t')} \|f(\cdot,t')\|_{(s)} dt'$$
(7.33)

holds for all  $t \ge 0$ , all solutions u to (1.2) and all  $s \in \mathbb{R}$ . Here  $d_{tot,+} := \max\{d_{sil,+}, d_{n,+}\}; d_{sil,+}$  is defined as in the case of (7.31);

$$d_{n,+} := \max\{d_{\max}(R_{Q,+}^+, \kappa_{\text{tot},+}), d_{\max}(R_{Q,+}^-, \kappa_{\text{tot},+})\};$$

and the constant C only depends on the spectra of the Riemannian manifolds  $(M_r, g_r), r \in \{1, \ldots, R\}$ , and the coefficients of the equation (1.2).

Remark 7.20. The statement follows from Theorem 22.8, cf. Remark 22.10.

#### 7.4.1 Optimality

Even though it is of interest to know that the estimate (7.30) holds, its value is dependent on whether it is optimal or not. For this reason, we now address the issue of optimality.

**Theorem 7.21.** Consider the equation (1.2) with f = 0. Assume that it is geometrically nondegenerate, diagonally dominated, balanced and convergent; cf. Definition 7.8. Then  $\eta_{nl}$ , given by Definition 1.23, satisfies  $\eta_{nl} = \kappa_{tot,+}$ . Assume, in addition to the above, that the equation exhibits subdominant block non-degeneracy. If  $\bar{\beta}_Q < 0$  and  $\kappa_{sil,+} = \kappa_{tot,+}$ , then there is a constant  $0 < C \in \mathbb{R}$  and a solution  $u \neq 0$  to (1.2) with f = 0 such that

$$\mathfrak{E}^{1/2}[u](t) \ge C\langle t \rangle^{d_{\mathrm{sil},+}-1} e^{\kappa_{\mathrm{tot},+}t} \mathfrak{E}^{1/2}[u](0)$$
(7.34)

for all  $t \ge 0$ . In this setting, the estimate (7.31) is thus optimal for f = 0. If  $\bar{\beta}_Q = 0$  and either  $\kappa_{\text{sil},+} = \kappa_{\text{tot},+}$  or  $\kappa_{Q,\text{ts},+} = \kappa_{\text{tot},+}$  holds, then there is a constant  $0 < C \in \mathbb{R}$  and a solution  $u \ne 0$  to (1.2) with f = 0 such that

$$\mathfrak{E}^{1/2}[u](t) \ge C\langle t \rangle^{d_{\mathrm{sts},+}-1} e^{\kappa_{\mathrm{tot},+}t} \mathfrak{E}^{1/2}[u](0)$$
(7.35)

for all  $t \ge 0$ . In this setting, the estimate (7.32) is thus optimal for f = 0. Finally, if  $\beta_Q > 0$  and either  $\kappa_{sil,+} = \kappa_{tot,+}$  or  $\kappa_{Q,+} = \kappa_{tot,+}$  holds, then there is a constant  $0 < C \in \mathbb{R}$  and a solution  $u \ne 0$  to (1.2) with f = 0 such that

$$\mathfrak{E}^{1/2}[u](t) \ge C\langle t \rangle^{d_{\text{tot},+}-1} e^{\kappa_{\text{tot},+}t} \mathfrak{E}^{1/2}[u](0)$$
(7.36)

for all  $t \ge 0$ . In this setting, the estimate (7.33) is thus optimal for f = 0.

Remark 7.22. The statement follows from Theorem 22.13, cf. Remark 22.10.

#### 7.4.2 Examples

In order to illustrate the results, we give two examples.

**Example 7.23.** Consider the equation

$$u_{tt} - e^{-2t}u_{\theta\theta} + e^{-t}Xu_{\theta} + \alpha u_t + \zeta u = 0$$
(7.37)

on  $\mathbb{S}^1 \times \mathbb{R}$ , where  $X, \alpha, \zeta \in \mathbb{R}$ . The relevant metric in this case is

$$g = -dt \otimes dt + e^{2t} d\theta \otimes d\theta.$$

In particular,  $\bar{g} = e^{2t} d\theta \otimes d\theta$  and  $\bar{k} = \bar{g}$ . Moreover,  $\chi = 0$ . In particular, (7.37) is  $C^1$ -silent in the sense of Definition 4.1. Note also that  $\mathcal{X} := e^{-t} X \partial_{\theta}$  is  $C^0$ -future bounded and that (4.2) holds with  $\alpha_{\infty} = \alpha$ ,  $\zeta_{\infty} = \zeta$ ,  $C_{\rm mn} = 1$  and  $\eta_{\rm mn} = 1$ . In particular, Proposition 4.4 applies. In order to justify that the results of the present chapter are applicable, note that  $\bar{k}$  is diagonally convergent (with  $\beta_1 = -1$ ) and that  $\bar{g}$  is  $C^2$ -asymptotically diagonal. Moreover, the shift vector field is negligible, and the main coefficients are convergent;  $X_{\infty}^1 = X$ ,  $\alpha_{\infty} = \alpha$  and  $\zeta_{\infty} = \zeta$ . Finally, (7.37) is  $C^2$ -balanced and asymptotically non-degenerate. To conclude, (7.37) is geometrically non-degenerate, diagonally dominated, balanced and convergent; cf. Definition 7.8. Thus the results of the present chapter apply.

Assume that  $\alpha > 0$ ,  $\zeta = \alpha^2/4 + 1$  and that  $X = 2\alpha + 1$ . Then Proposition 4.4 yields the conclusion that there are constants C and  $s_0$  such that

$$\mathfrak{E}_s[u](t) \le C \mathfrak{E}_{s+s_0}[u](0) e^{-\alpha t}$$

for all  $t \ge 0$ . Choosing  $\alpha := 2 \cdot 4711$ , it is thus clear that (7.1) holds. On the other hand,

$$R_{1,+}^{\pm} = \frac{1}{2}(-\alpha + \beta_1 \pm \tilde{X}_{\infty}^1).$$

Moreover,  $\beta_1 = -1$ ,  $g_{\infty}^{11} = 1$  and  $X_{\infty}^1 = X$ . In particular,  $R_{1,+}^+ = \alpha/2$  and  $\kappa_{\text{tot},+} = \alpha/2$ . Thus, Theorem 7.21 yields the conclusion that  $\eta_{\text{nl}} = \alpha/2$ . Let  $\alpha := 2 \cdot 4711$  and fix  $0 < \epsilon, N \in \mathbb{R}$ . Then

 $\eta_{\rm nl} = 4711$ . Let  $\mathcal{A}_{\rm nl}$  be defined by (1.22). Clearly,  $4710 \notin \mathcal{A}_{\rm nl}$ . In other words, regardless of the choice of C > 0, there is a solution  $0 \neq u \in \mathcal{S}$  and a  $t \geq 0$  such that  $\mathfrak{E}[u](t) \geq C\mathfrak{E}[u](0)e^{9420t}$ . Choosing  $C = \epsilon^{-2}N^2$  and normalising u so that  $\mathfrak{E}[u](0) = \epsilon^2$ , it is clear that

$$\mathfrak{E}^{1/2}[u](0) = \epsilon, \quad \mathfrak{E}^{1/2}[u](t) \ge N.$$

This justifies the statements made in the introduction.

**Example 7.24.** Consider the equation (7.6). It is of the same type as (7.37), so that the results of the present chapter apply. In the case of (7.6),

$$\kappa_{\mathrm{sil},+} = -b, \quad \kappa_{+} = \frac{1}{2}(-b - 1 + |a|), \quad \kappa_{\mathrm{tot},+} = \frac{1}{2}\max\{-b - 1 + |a|, -2b\},$$

where we assume that b < 0. Appealing to Theorem 7.21 yields the conclusion that (7.7) holds. Moreover, since (7.6) exhibits subdominant block non-degeneracy, we know that (7.8) holds; cf. Theorem 7.19.

#### 7.5 Outline of the argument, outlook

The proofs of the results of the present chapter are provided in Part VI, in particular Chapters 20–22. However, they build upon some of the results of the previous chapters, in particular on those of Part III. We therefore begin by describing the contents of Part III.

#### 7.5.1 Averaging over oscillations

Consider a solution u to (1.2). Let  $z(\iota, t)$  be the  $\iota$ 'th Fourier coefficient of  $u(\cdot, t)$ . Then z satisfies (24.18). For at least some part of the interval  $[0, \infty)$ , the behaviour of z can be expected to be oscillatory. The purpose of Part III is to develop methods for estimating how z evolves in such oscillatory regimes. Before turning to the formal details, it is useful to develop some intuition. However, a rough idea of the perspective developed in Part III is given by the argument presented in Subsection 6.1.1. Turning to the formal details, the first step is to calculate the matrix exponential of a special class of matrices; one example of an element of this class is given by  $A_{\infty}(\iota, t_0)T$  appearing in Subsection 6.1.1.

Matrix exponentials and approximations. In order to estimate the overall evolution of solutions during one period of the oscillations, it is useful to reformulate the equation; in the case of the limit equation in the noisy setting, the relevant reformulation is given by (6.2)-(6.4). In the general case, the reformulation is given in Section 14.2 below. Freezing the coefficients in the reformulated equation, one is led to the problem of calculating the matrix exponential  $\exp(2\pi P)$ , where P is a matrix of the form

$$P = \begin{pmatrix} 0 & \mathrm{Id}_m \\ -\mathrm{Id}_m & 0 \end{pmatrix} + \begin{pmatrix} i\xi \mathrm{Id}_m + R_{11} & 0 \\ R_{21} & -i\xi \mathrm{Id}_m - R_{11} \end{pmatrix} + E.$$
(7.38)

Here E should be thought of as an error term and  $\xi$  and  $R_{ij}$  should be thought of as being small; cf. Section 13.2 for a detailed description of the assumptions. Note that P has a very special structure. First of all, the exponential of  $2\pi$  times the first term on the right hand side of (7.38) equals Id<sub>2m</sub>. Second, the second matrix on the right hand side of (7.38) is trace free, block triangular and of lower order. In Section 13.2, we calculate  $\exp(2\pi sP)$  up to an error term for  $s \in [-1, 1]$ . In order to understand the overall evolution over longer periods of time, it is sufficient to focus on the case that s = 1. However, we are sometimes interested in going backwards in time (for instance when specifying asymptotics), and in that case, it is of interest to know what happens for s = -1. Finally, in the noisy setting, we are also interested in knowing what happens during one period of the oscillations, and this is the motivation for taking an interest in all the  $s \in [-1, 1]$ .

*Freezing the coefficients.* To estimate the evolution of solutions to non-constant coefficient equations (over one period), we freeze the coefficients and calculate the solution to the resulting constant coefficient equation. In order to estimate the error involved, we need to compare the matrix exponential (which arises when freezing the coefficients) with the appropriate fundamental solution of the actual system of equations. This is the topic of Section 13.3.

Assumptions and preliminary estimates. Turning to equations of the form (1.2), we state the relevant assumptions in Section 14.1. The assumptions are restrictive enough that we can analyse the behaviour over one period of the oscillations. However, they are also general enough that the results apply in the situations considered in Chapters 2 and 6, as well as in those considered in the present chapter. In Section 14.2 we perform a change of variables that allows us to isolate the leading order behaviour during one period of the oscillations. It also allows us to appeal to the matrix exponential calculations of Section 13.2.

Choosing time intervals. Above, we speak of "one period of the oscillations" etc. In general, this notion is not uniquely defined. However, for practical purposes, we here take one period of the oscillations to mean an interval  $[t_a, t_b]$  such that

$$\int_{t_a}^{t_b} \mathfrak{g}(\iota, t) dt = 2\pi.$$

Variation within a period; a first approximation of the fundamental solution. Before appealing to the results of Sections 13.2 and 13.3, we need to estimate the variation of the coefficients of the equation within one period of the oscillations. This is the purpose of Sections 14.3 and 14.4. Given the relevant estimates, we are then in a position to give a first approximation of the fundamental solution; this is the purpose of Section 14.5.

**Evolution over one period; iteration.** In Chapter 15, we consider the evolution over one period of the oscillations in greater detail. First, we derive a rough estimate in Section 15.1. This estimate is then refined to a detailed estimate in Section 15.2.

Iteration. In order to estimate the evolution of the solution over longer periods of time, it is convenient to introduce a time sequence  $\{t_k\}$  such that  $[t_k, t_{k+1}]$  corresponds to one period of the oscillations. Introducing such a time sequence and changing the variables appropriately yields an iteration relating the solution at  $t_{k+1}$  to the solution at  $t_k$ . Moreover, the matrices involved in the relation can be well approximated by matrices that can be calculated from the coefficients of the equation. The iteration is the basic tool for analysing the behaviour of solutions over longer periods of time. The reader interested in the details is referred to Section 15.3.

#### 7.5.2 The asymptotically diagonal setting

**Preliminaries.** In Part VI we turn to a study of asymptotically diagonal equations. To begin with, we give an analytical definition of what this means in Section 20.2; in Part VII we verify that the requirements of the analytical definition follow from the geometric conditions introduced in Definition 7.8. In Section 20.2, we also work out some of the consequences of the definition. In particular, we verify that the basic assumptions of Part III are satisfied. In Section 20.3, we then turn to the problem of simplifying the matrix coefficients that appear in the iteration mentioned above. Using this simplification, we are in a position to introduce a notion of "frequency eras"; cf. Definition 20.10. In the non-degenerate setting, a frequency era is a time interval in which one of the terms inside the parenthesis on the right hand side of (7.10) dominates. In Section 20.4, we simplify the matrix coefficients further in frequency eras. Finally, in Section 20.5, we estimate sums of the error terms that appear in the iteration.

Analysis for one mode. Given the preliminaries, we are in a position to analyse how the energy associated with one mode evolves over time. This is the subject of Chapter 21, a chapter in

which we restrict our attention to the asymptotically non-degenerate setting; cf. Definition 7.6. To begin with, we devote Section 21.2 to estimating the evolution of the energy of one mode in one frequency era. The basic tool for obtaining the estimates is the iteration. In the end, we therefore need to estimate the norm of a matrix product (where the number of factors depends on the frequency). The argument begins by a derivation of as good estimates of the constituent matrices as possible. As a second step, we simplify the constituent matrices by conjugating them with appropriate matrices. Finally, we estimate the norm of the matrix products; cf. the proof of Lemma 21.7 for the details.

Refined estimates in transparent eras. Lemma 21.7 applies both when going forward and when going backwards in time. In the case that  $\bar{\beta}_q \neq 0$ , the estimate is essentially optimal (with, possibly, the exception of a loss  $e^{\epsilon t}$  in the decay). However, in the case that  $\bar{\beta}_q = 0$ , there is potentially a substantial loss. We therefore devote Section 21.3 to deriving refined estimates in transparent eras. The main idea in this case is that for  $\bar{\nu}_q(\iota)$  large, the results of Lemma 21.7 yield good estimates. On the other hand, fixing a  $0 < K_{\rm co} \in \mathbb{R}$ , there are only finitely many types of frequency eras with  $\bar{\beta}_q = 0$  and  $\bar{\nu}_q(\iota) \leq K_{\rm co}$ . For the latter class of frequency eras, a refined analysis can therefore be carried out, and the loss in the estimates is at worst of the form  $e^{\epsilon t}$ .

*Estimates for one mode.* Given the analysis described above, we are in a position to derive estimates for one full mode. This is the purpose of Section 21.4. The desired conclusion is essentially obtained by combining previous results.

Unbounded frequency eras. As already mentioned, many of the estimates involve a loss of the type  $e^{\epsilon t}$ . However, in some cases, this loss can be avoided. In particular, in the case of a frequency era of the form  $[t_0, \infty)$ . In this case the loss  $e^{\epsilon t}$  can sometimes be replaced by a polynomial loss. Justifying this statement is the subject of Section 21.5; cf., in particular, Lemma 21.29.

*Optimality.* Finally, in Section 21.6, we turn to the question of optimality. In order to obtain the desired conclusions, it is sufficient to consider one frequency era. Moreover, we only need to derive a lower bound on the norm of the matrix products, say  $P_k$  appearing in the iteration. However, deriving such a lower bound is somewhat technical. First, we have to conjugate the constituent matrices in order to isolate the dominant part; then we have to choose the initial data  $\xi_0$  appropriately to ensure maximal growth of  $|P_k\xi_0|$ ; and, finally, we have to verify that the error terms do not cause problems.

**Optimal energy estimates.** In Chapter 22, we are finally in a position to prove the results stated in the present chapter. The argument essentially consists of summarising the estimates derived in previous chapters. In Section 22.2, we begin by dividing  $\mathcal{I}_{\rm B}$  into different subsets. Then, in Proposition 22.6, we derive estimates for individual modes, based on this division of  $\mathcal{I}_{\rm B}$ . Applying Minkowski's inequality to the estimates for the individual modes yields Theorem 7.19; cf. Section 22.3 below. Finally, we prove optimality in Section 22.4.

#### 7.5.3 Outlook

**Degeneracy.** There are several ways in which one could think of improving the results of the present chapter. First of all, it would be desirable to remove the non-degeneracy condition. It is then to be expected that the one matrix  $\tilde{X}_{\infty}^q$  appearing in (7.23) is replaced by a compact set of matrices; cf.  $\bar{X}_{\text{diag}}^q(\iota)$  introduced in (20.37) below. This causes complications, but it should be possible to derive conclusions nonetheless. For a non-degenerate equation with no transparent eras, the growth/decay rate is determined by finitely many matrices. In the degenerate setting, it is to be expected that the growth/decay rate is determined by the supremum of the real parts of the eigenvalues of matrices belonging to a compact set. Moreover, there would be infinitely many different matrices appearing in the estimates for the frequency modes. Nevertheless, it should be possible to carry out a corresponding analysis.

**Detailed estimates for a single mode.** Considering a fixed mode, the time interval  $[0, \infty)$  can be divided into several parts. First, there is the initial interval  $[0, T_{ini}]$ , where  $T_{ini}$  is, roughly

speaking, defined by the condition that (7.11) is a good approximation of the equation for  $t \geq T_{\text{ini}}$ . The interval  $[T_{\text{ini}}, \infty)$  is, in its turn, divided into frequency eras. Due to the methods developed in Part VI of these notes, we have a good understanding of how the solutions behave in a frequency era. However, it would be of interest to give a description of how the Fourier coefficient behaves for all  $t \in [T_{\text{ini}}, \infty)$ . In the present chapter we focus on estimates, but it would be of interest to derive more information. A first step is taken in Subsection 21.4.1, but it should be possible to do better.

## Chapter 8

# Improved asymptotic estimates in the silent setting

Even though Proposition 4.11 gives detailed information concerning the asymptotic behaviour of solutions in the silent setting, there is one fundamental problem: we have essentially no information concerning the values of  $s_{\text{hom}}$  and  $s_{\text{ih}}$ . Proposition 4.16 suffers from a similar deficiency. In the present chapter, we state results analogous to these propositions, but with specific values for the loss of derivatives involved in the estimates.

# 8.1 Estimating the asymptotic data in terms of the initial data

Let us start by giving an estimate of the asymptotic data in terms of the initial data, with a loss of derivatives that can be calculated in terms of the coefficients of the equation.

**Proposition 8.1.** Consider the equation (1.2). Assume that it is geometrically non-degenerate, diagonally dominated, balanced and convergent; cf. Definition 7.8. Assume that  $\bar{\beta}_Q < 0$  and that f is a smooth function such that for every  $s \in \mathbb{R}$ ,

$$\|f\|_{\mathrm{sil},s} := \int_0^\infty e^{-\kappa_{\mathrm{sil},+}t} \|f(\cdot,t)\|_{(s)} dt < \infty,$$
(8.1)

where  $\kappa_{\text{sil},+} := \kappa_{\max}(A_{\infty})$  and  $A_{\infty}$  is defined by (4.4). Let  $\beta_{\text{rem}} := \min\{-\bar{\beta}_Q, \kappa_{\text{mn}}\}$ , where  $\kappa_{\text{mn}}$ is the constant appearing in (7.20). Fix  $0 < \beta \leq \beta_{\text{rem}}$  and  $\epsilon > 0$ . Let  $E_a$  be the first generalised eigenspace in the  $\beta, A_{\infty}$ -decomposition of  $\mathbb{C}^{2m}$ ; cf. Definition 4.7. Then there are constants  $C_{\epsilon,\beta}$ and N, where  $C_{\epsilon,\beta}$  only depends on  $\epsilon, \beta$ , the coefficients of the equation (1.2) and the spectra of the Riemannian manifolds  $(M_r, g_r), r \in \{1, \ldots, R\}$ ; and N only depends on m, such that the following holds. Given a smooth solution u to (1.2), there is a  $V_{\infty} \in C^{\infty}(\overline{M}, E_a)$  such that

$$\left\| \begin{pmatrix} u(\cdot,t) \\ u_t(\cdot,t) \end{pmatrix} - e^{A_{\infty}t} V_{\infty} - \int_0^t e^{A_{\infty}(t-\tau)} \begin{pmatrix} 0 \\ f(\cdot,\tau) \end{pmatrix} d\tau \right\|_{(s)}$$

$$(8.2)$$

 $\leq C_{\epsilon,\beta} \langle t \rangle^N e^{(\kappa_{\mathrm{sil},+}-\beta)t} \left( \|u_t(\cdot,0)\|_{(s+s_{\mathrm{h},\beta,+}+\epsilon)} + \|u(\cdot,0)\|_{(s+s_{\mathrm{h},\beta,+}+\epsilon+1)} + \|f\|_{\mathrm{sil},s+s_{\mathrm{ih},\beta}+\epsilon} \right)$ holds for all  $t \geq 0$  and all  $s \in \mathbb{R}$ . Here

$$s_{\mathrm{h},\beta,+} := \max_{1 \le q \le Q} \max\left\{0, -\frac{\beta + \kappa_{q,+} - \kappa_{\mathrm{sil},+}}{\bar{\beta}_q}\right\},\tag{8.3}$$

$$s_{\mathrm{ih},\beta} := \max\left\{-\frac{\kappa_{\mathrm{ode}}}{\bar{\beta}_Q}, -\frac{\beta}{\bar{\beta}_Q} + \max_{1 \le q \le Q} \max\left\{0, -\frac{\kappa_{q,+} - \kappa_{\mathrm{sil},+}}{\bar{\beta}_q}\right\}\right\},\tag{8.4}$$

where  $\kappa_{ode} := \operatorname{Rsp} A_{\infty}$ , cf. Definition 4.3, and  $\kappa_{q,+}$  is defined by (7.24). Moreover,

$$\|V_{\infty}\|_{(s)} \le C_{\epsilon,\beta} \left( \|u_t(\cdot,0)\|_{(s+s_{h,\beta}+\epsilon)} + \|u(\cdot,0)\|_{(s+s_{h,\beta}+\epsilon+1)} + \|f\|_{A,s+s_{h,\beta}+\epsilon} \right),$$
(8.5)

where

$$s_{\mathrm{h},\beta} := \max_{1 \le q \le Q} \left( -\frac{\beta + \kappa_{q,+} - \kappa_{\mathrm{sil},+}}{\bar{\beta}_q} \right).$$
(8.6)

Remark 8.2. The proposition is a consequence of Proposition 23.1 and Remark 23.2.

**Remark 8.3.** The constants  $\bar{\beta}_q$ ,  $q \in \{1, \ldots, Q\}$ , are introduced in Definition 7.6.

**Remark 8.4.** The function  $V_{\infty}$  is uniquely determined by the fact that it satisfies the estimate (8.2).

**Remark 8.5.** Both  $s_{h,\beta,+}$  and  $s_{ih,\beta}$  are non-negative. However,  $s_{h,\beta}$  could be strictly negative. In this sense, the function  $V_{\infty}$  could be more regular than the initial data.

**Remark 8.6.** As opposed to (4.10),  $\beta$  need not equal  $\beta_{\text{rem}}$  in (8.2). The reason for allowing  $\beta$  to vary in the range  $(0, \beta_{\text{rem}}]$  is that a higher  $\beta$  yields more detailed asymptotics, but a lower  $\beta$  yields a lower loss of derivatives.

**Remark 8.7.** Under some circumstances, the  $\epsilon$  appearing in (8.2) and (8.5) can be removed. See Remarks 23.8 and 23.9 for more details.

In order to illustrate the result, let us consider an example.

**Example 8.8.** Consider the equation (7.37). It satisfies the conditions of Proposition 8.1. Moreover, Q = 1;  $\bar{\beta}_1 = -1$ ; (7.20) holds with  $\kappa_{mn} = 1$ ; and  $\beta_{rem} = 1$ . Let

$$\zeta = \frac{1}{4}\alpha^2 \pm \frac{1}{4}Y^2,$$

where  $Y \in \mathbb{R}$ . There are two cases to consider.

Assuming, for the sake of argument, that  $\zeta \geq \alpha^2/4$ , we know that  $\kappa_{\rm sil,+} = -\alpha/2$  and that  $\kappa_{\rm ode} = 0$ . Moreover,

$$\kappa_{1,+} = \frac{1}{2}(-\alpha + \bar{\beta}_1 + |X|).$$

Thus

$$s_{\mathrm{h},\beta} = \beta - \frac{1}{2}(1 - |X|).$$

In particular, for small  $\beta$  and |X|, we know that  $s_{h,\beta}$  is negative. Note also that, regardless of how small  $\beta$  is,  $E_a = \mathbb{C}^2$ , and the map from initial data to asymptotic data is a homeomorphism. If X = 0, (8.5) thus implies that for every  $\epsilon > 0$ , there is a constant  $C_{\epsilon}$  such that

$$\|V_{\infty}\|_{(s)} \le C_{\epsilon} \left( \|u_t(\cdot, 0)\|_{(s-1/2+\epsilon)} + \|u(\cdot, 0)\|_{(s+1/2+\epsilon)} \right).$$

Moreover, for  $\beta \leq 1/2$ ,  $s_{\mathrm{h},\beta,+} = 0$ , so that (8.2) only involves an  $\epsilon$ -loss in derivatives. On the other hand, the time dependence of the right hand side is then not optimal. In fact, we are allowed to choose  $\beta = 1$ , and then the right hand side of (8.2) has a time dependence of the form  $\langle t \rangle^N e^{(\kappa_{\mathrm{sil},+}-1)t}$  as opposed to  $\langle t \rangle^N e^{(\kappa_{\mathrm{sil},+}-1/2)t}$ . However, the price for the improvement as far as the time dependence is concerned is a loss of derivatives.

Assume now that  $\zeta \leq \alpha^2/4$ . Then  $\kappa_{\rm sil,+} = -\alpha/2 + |Y|/2$ . On the other hand,  $\kappa_{1,+}$  is the same as before so that

$$s_{\mathrm{h},\beta} = \beta - \frac{1}{2}(1 + |Y| - |X|).$$

In this case, it is thus clear that  $s_{h,\beta}$  can be made arbitrarily negative, by making |Y| large enough. On the other hand,  $\operatorname{Rsp} A_{\infty} = \kappa_{\text{ode}} = |Y|$ . Thus, if  $\beta \leq |Y|$ ,  $E_a$  is (complex) one dimensional. If we wish  $E_a$  to be two dimensional, we have to have  $\beta > |Y|$ , in which case

$$s_{\mathrm{h},\beta} > \frac{1}{2}(|Y| + |X| - 1) \ge -\frac{1}{2}.$$

If we want complete asymptotic information, we can thus not hope for  $s_{h,\beta} \leq -1/2$ . Note also that if  $|Y| \geq 1$ , we, in the present context, do not obtain a homeomorphism between initial data and asymptotic data.

# 8.2 Estimating the initial data in terms of the asymptotic data

Next, we improve Proposition 4.16 by expressing the derivative losses in terms of the coefficients of the equation.

**Proposition 8.9.** Consider the equation (1.2) with f = 0. Assume that it is geometrically non-degenerate, diagonally dominated, balanced and convergent; cf. Definition 7.8. Assume that  $\bar{\beta}_Q < 0$  and let  $\beta_{\text{rem}} := \min\{-\bar{\beta}_Q, \kappa_{\text{mn}}\}$ , where  $\kappa_{\text{mn}}$  is the constant appearing in (7.20). Fix  $0 < \beta \leq \beta_{\text{rem}}$  and  $\epsilon > 0$ . Let  $E_a$  be the first generalised eigenspace in the  $\beta, A_{\infty}$ -decomposition of  $\mathbb{C}^{2m}$ ; cf. Definition 4.7. Then there is a linear injective map

$$\Phi_{\beta,\infty}: C^{\infty}(\bar{M}, E_a) \to C^{\infty}(\bar{M}, \mathbb{C}^{2m})$$

such that the following holds. First, if  $\Phi_{\beta,\infty}^j : C^{\infty}(\bar{M}, E_a) \to C^{\infty}(\bar{M}, \mathbb{C}^m)$ , j = 1, 2, are defined by the condition that

$$\Phi_{\beta,\infty}(\psi) = \begin{pmatrix} \Phi^1_{\beta,\infty}(\psi) \\ \Phi^2_{\beta,\infty}(\psi) \end{pmatrix}$$

for all  $\psi \in C^{\infty}(\overline{M}, E_a)$ , then

$$\|\Phi^{1}_{\beta,\infty}(\psi)\|_{(s+1)} + \|\Phi^{2}_{\beta,\infty}(\psi)\|_{(s)} \le C_{\epsilon} \|\psi\|_{(s+s_{\mathrm{h},-}+\epsilon)}$$
(8.7)

for all  $s \in \mathbb{R}$  and all  $\psi \in C^{\infty}(\overline{M}, E_a)$ , where  $C_{\epsilon}$  only depends on  $\epsilon$ , the coefficients of the equation (1.2) and the spectra of  $(M_r, g_r)$ ,  $r \in \{1, \ldots, R\}$ , and

$$s_{\mathrm{h},-} := \max_{1 \le q \le Q} \left( -\frac{\kappa_{q,-} + \kappa_{\mathrm{sil},+}}{\bar{\beta}_q} \right).$$
(8.8)

Second, if  $\psi \in C^{\infty}(\overline{M}, E_a)$  and u is the solution to (1.2) (with f = 0) such that

$$\begin{pmatrix} u(\cdot,0)\\ u_t(\cdot,0) \end{pmatrix} = \Phi_{\beta,\infty}(\psi), \tag{8.9}$$

then

$$\left\| \begin{pmatrix} u(\cdot,t) \\ u_t(\cdot,t) \end{pmatrix} - e^{A_{\infty}t} \psi \right\|_{(s)}$$

$$\leq C_{\epsilon,\beta} \langle t \rangle^N e^{(\kappa_{\rm sil,+}-\beta)t} \left( \| u_t(\cdot,0) \|_{(s+s_{\rm h,\beta,+}+\epsilon)} + \| u(\cdot,0) \|_{(s+s_{\rm h,\beta,+}+\epsilon+1)} \right)$$

$$(8.10)$$

holds for all  $t \ge 0$  and all  $s \in \mathbb{R}$ , where  $C_{\epsilon,\beta}$  only depends on  $\epsilon$ ,  $\beta$ , the coefficients of the equation (1.2) and the spectra of the Riemannian manifolds  $(M_r, g_r)$ ,  $r \in \{1, \ldots, R\}$ , and N only depends on m. Finally, if  $E_a = \mathbb{C}^{2m}$  (i.e., if  $\beta > \operatorname{Rsp} A_{\infty}$ ; cf. Definition 4.3), then  $\Phi_{\beta,\infty}$  is surjective.

**Remark 8.10.** The proposition is a consequence of Proposition 23.10 and Remark 23.11.

**Remark 8.11.** Again, the  $\epsilon$  appearing in (8.7) can be removed under appropriate circumstances; cf. Remark 23.12.

Again, let us illustrate the result with an example.

**Example 8.12.** Consider the equation (7.37). It satisfies the conditions of Proposition 8.9 and in what follows, we use the notation introduced in Example 8.8. As before, there are two cases to consider.

Assume that  $\zeta \geq \alpha^2/4$ . Then  $\kappa_{\rm sil,+} = -\alpha/2$  and

$$\kappa_{1,-} = \frac{1}{2}(\alpha - \bar{\beta}_1 + |X|),$$

so that

$$s_{\mathrm{h},-} = \frac{1}{2}(1+|X|).$$

Note that |X| only leads to a loss. Comparing with Example 8.8, we see that if X = 0, we can, roughly speaking, gain 1/2 derivative when going from initial data to asymptotic data, but we lose roughly 1/2 derivative when going back. Introducing  $X \neq 0$  leads to an additional loss in both directions.

Assume that  $\zeta \leq \alpha^2/4$ . In this case,  $\kappa_{1,-}$  remains the same, but  $\kappa_{\rm sil,+}$  changes. In fact,

$$s_{\mathrm{h},-} = \frac{1}{2}(1 + |X| + |Y|).$$

In other words, X and Y only yield a loss of derivatives. Comparing with Example 8.8, we see that the (potential) gain caused by Y when going from initial data to asymptotic data is lost when going back.

#### 8.3 Outline of the argument, outlook

The proofs of Propositions 8.1 and 8.9 build on Part III and Chapters 20–22. However, the argument is based on a more refined version of the analysis of the Fourier coefficients than the one used to obtain optimal energy estimates. In fact, in order to obtain optimal energy estimates, it is sufficient to isolate the frequency era corresponding to the least amount of decay. However, in order to calculate the loss of derivatives of interest in the present chapter, we need to consider all the frequency eras for a given mode.

**Refined analysis.** Consider a Fourier coefficient z corresponding to a solution to (1.2) and a  $\iota \in \mathcal{I}_{\mathrm{B}}$ . Under the assumptions of Propositions 8.1 and 8.9, there is a  $T_{\mathrm{ode}}$  such that the ODEbehaviour is dominant in  $[T_{\mathrm{ode}}, \infty)$ . The interval  $[0, T_{\mathrm{ode}}]$  can, in its turn, be divided into frequency eras, say  $[0, T_1], \ldots, [T_l, T_{\mathrm{ode}}]$ . In each frequency era, there is a worst case scenario as far as the growth/decay is concerned, say  $\lambda_j$  in  $[T_j, T_{j+1}]$ . This leads, roughly speaking, to a growth of the form

$$\exp[\lambda_0 T_1 + \dots + \lambda_l (T_{\text{ode}} - T_l) + \kappa_{\text{sil},+} (t - T_{\text{ode}})]$$

for  $t \ge T_{\text{ode}}$ . On the other hand,  $T_{l+1} - T_l$  can be estimated in terms of  $\iota$ . Moreover, we, in the end, wish to compare this growth with  $e^{\kappa_{\text{sil},+}t}$ . This leads to the problem of estimating, e.g.,

$$\exp[(\lambda_0 - \kappa_{\rm sil,+})T_1 + \dots + (\lambda_l - \kappa_{\rm sil,+})(T_{\rm ode} - T_l)].$$
(8.11)

In particular, we are interested in the worst case scenario as we vary  $\iota$ . Carrying out the corresponding analysis leads to the conclusion that expressions of the form (8.11) can be estimated by  $C\langle\nu(\iota)\rangle^{s_a}$  for some  $s_a \in \mathbb{R}$  depending on the  $\bar{\beta}_q$ , the  $\kappa_{q,+}$  and  $\kappa_{\mathrm{sil},+}$ . This is, roughly speaking, the argument by which one arrives at the number  $s_{\mathrm{h},\beta}$ . Note that all the  $\lambda_j - \kappa_{\mathrm{sil},+}$  could be negative;

this is the origin of the (potential) gain of derivatives in the estimate for  $V_{\infty}$ . In the case of  $s_{h,\beta,+}$ , the estimate has to be valid for all  $t \ge 0$ , and it is therefore clear that  $s_{h,\beta,+}$  has to be bounded from below by zero.

**Outline.** The main structure of the proofs of Propositions 8.1 and 8.9 is similar to that of the proofs of Propositions 4.11 and 4.16; cf. Subsection 4.4.2 for a rough description of the arguments in the case of the latter propositions. However, in order to obtain the desired estimates, we need to appeal to the refined analysis described above. The details are provided in Sections 23.2 and 23.3.

#### 8.3.1 Outlook

In the present chapter, we only consider the silent setting. It would, however, also be of interest to derive similar conclusions in the transparent and noisy settings. Using the methods developed in these notes, it should be possible to do so. However, the required arguments can be expected to be lengthy.

## Part II

# Rough analysis in the silent and in the transparent settings

## Chapter 9

# ODE's with exponentially decaying error terms

**Introduction.** Consider (24.16), obtained by decomposing (1.2) into Fourier modes. Introducing additional variables, it can be reformulated as a first order system

$$\dot{v}(t) = B(t)v(t) + F(t).$$
 (9.1)

The function F arises from  $\hat{f}$  appearing on the right hand side of (24.16) and depends on the frequency  $\iota \in \mathcal{I}_{\mathrm{B}}$  and t. In general, B also depends on both  $\iota$  and t. However, in the silent setting, the matrix can be written as a sum  $B(t) = A + A_{\mathrm{rem}}(t)$ , where A is a constant matrix and  $A_{\mathrm{rem}}$  decays exponentially. Moreover, A does not depend on  $\iota$ . This is the simplest situation, and the one we consider in the present chapter. The case of transparent equations is similar. However, in that case, A depends on  $\iota$ , a more complicated situation which we discuss in Chapter 11. Even though the division of B into A and  $A_{\mathrm{rem}}$  corresponds to a division into a dominant part A and an error term  $A_{\mathrm{rem}}$ , there is one problem:  $A_{\mathrm{rem}}$  depends on  $\iota$ , and  $||A_{\mathrm{rem}}(0)||$  typically tends on  $\infty$  as  $|\nu(\iota)| \to \infty$ . In that sense,  $A_{\mathrm{rem}}$  cannot be considered to be negligible. On the other hand, the equations of interest are such that for each  $\iota$ , there is a time, say  $T_{\mathrm{ode}}$  (depending on  $\iota$ ), such that

$$||A_{\rm rem}(t)|| \le C_{\rm rem} e^{-\beta_{\rm rem}(t-T_{\rm ode})} \tag{9.2}$$

for  $t \ge T_{\text{ode}}$ . Moreover,  $C_{\text{rem}} > 0$  and  $\beta_{\text{rem}} > 0$  are independent of  $\iota$ . Here we refer to the interval  $[T_{\text{ode}}, \infty)$  as the *non-oscillatory* or *ODE era* of the relevant Fourier mode, and this is the interval we focus on in the present chapter.

Asymptotics. In this chapter we analyse the asymptotics of solutions to

$$\dot{v}(t) = Av(t) + A_{\rm rem}(t)v(t) + F(t), \tag{9.3}$$

where  $A_{\text{rem}}$  satisfies (9.2) and A is constant. To begin with, it can be demonstrated that, roughly speaking, solutions do not grow faster than solutions to  $\dot{v}(t) = Av(t) + F(t)$ ; cf. Lemma 9.14. In order to obtain detailed asymptotics, it is necessary to make more detailed assumptions concerning F. If F grows more quickly (or decays less slowly) than the fastest growing solution to  $\dot{v} = Av$ , then F can be expected to have a dominant influence on the asymptotics. However, there might be cancellations associated with, e.g., oscillatory behaviour in F. In order to obtain detailed asymptotics in such a situation, it can thus be expected to be necessary to make detailed assumptions concerning F. This is not our main interest here, and we therefore restrict our attention to the case that the fastest growing solutions to  $\dot{v} = Av$  grow more quickly than F. To be more precise, we assume that

$$||F||_A := \int_0^\infty e^{-\kappa_1 s} |F(s)| ds < \infty,$$
(9.4)

where  $\kappa_1 := \kappa_{\max}(A)$ ; cf. Definition 4.3. Under these circumstances, Lemma 9.14 ensures that solutions to (9.3) do not grow faster than the fastest growing solutions to  $\dot{v} = Av$ ; i.e.,

$$|v(t)| \le C \langle t - T_{\text{ode}} \rangle^{d_1 - 1} e^{\kappa_1 (t - T_{\text{ode}})}$$

$$\tag{9.5}$$

for  $t \ge T_{\text{ode}}$ , where we have used the notation (4.5) and  $d_1 := d_{\max}(A, \kappa_1)$ ; cf. Definition 4.3. At this stage, it would seem natural to return to (9.3) in order to calculate the leading order terms of the asymptotics. Compute, to this end,

$$\frac{d}{dt}\left(e^{-At}v - \int_{T_{\text{ode}}}^{t} e^{-As}F(s)ds\right) = e^{-At}A_{\text{rem}}(t)v(t).$$
(9.6)

Under ideal circumstances, the right hand side decays exponentially, so that we can integrate this equality. This would lead to the conclusion that the expression inside the parenthesis on the left hand side converges exponentially. Thus v would, to good approximation, behave as a solution to (9.3) with  $A_{\rm rem}$  set to zero. However, even though  $A_{\rm rem}$  and v satisfy (9.2) and (9.5) respectively, there is no reason why the right hand side of (9.6) should decay exponentially. Whether we obtain exponential decay or not is determined by the spread of the real parts of the eigenvalues of A; cf. Definition 4.3. If  $\text{Rsp}A < \beta_{\text{rem}}$ , then the right hand side of (9.6) decays exponentially. However, if  $\text{Rsp}A \geq \beta_{\text{rem}}$  we need to proceed differently. By conjugating A by a suitable  $T \in \mathbf{GL}_k(\mathbb{C})$ , the resulting matrix is in block form;  $T^{-1}AT = \text{diag}\{A_a, A_b\}$ . Moreover,  $\kappa_{\max}(A_a) = \kappa_{\max}(A)$ ,  $\text{Rsp}A_a < \beta_{\text{rem}}$  and  $\kappa_{\max}(A_b) \leq \kappa_{\max}(A) - \beta_{\text{rem}}$ ; cf. Definition 4.3 for an explanation of the notation. This follows from transforming A to Jordan normal form and arranging the Jordan blocks in a suitable way. Letting  $w(t) := T^{-1}v(t)$  and letting  $w_a$  and  $w_b$  denote the components of w corresponding to the blocks  $A_a$  and  $A_b$  respectively, (9.3) can be divided into the two equations

$$\dot{w}_a(t) = A_a w_a(t) + [T^{-1}A_{\text{rem}}(t)v(t)]_a + [T^{-1}F(t)]_a, \qquad (9.7)$$

$$\dot{w}_b(t) = A_b w_b(t) + [T^{-1}A_{\rm rem}(t)v(t)]_b + [T^{-1}F(t)]_b.$$
(9.8)

Here  $\xi_a$  and  $\xi_b$  denote the components of  $\xi \in \mathbb{C}^k$  corresponding to the blocks  $A_a$  and  $A_b$  respectively. In the case of  $w_a$ , we can carry out a computation similar to (9.6) by appealing to (9.7). In this way we obtain the leading order behaviour of  $w_a$ . However, in the case of  $w_b$  we only obtain an estimate. This is not so surprising, since the second term on the right hand side of (9.8) is, on the one hand, an error term (the only information we have concerning  $A_{\text{rem}}$  is an estimate). On the other hand, this term could potentially grow more quickly than the fastest growing solution to  $\dot{w}_b = A_b w_b$ . For this reason, it is not realistic to hope to extract asymptotic information concerning  $w_b$ .

**Outline.** The outline of the present chapter is as follows. In Section 9.1, we state the equations of interest and describe in greater detail the algebraic decomposition leading to (9.7) and (9.8). Then we proceed to derive a rough estimate for solutions to (9.3). This is the subject of Section 9.2. Given this information, it is then possible to derive detailed asymptotics concerning  $w_a$  in the above decomposition; cf. Section 9.3. This gives a map from initial data for (9.3) to asymptotic data for  $w_a$ . However, it is also of interest to go in the other direction; i.e., to start with asymptotic data for  $w_a$  and to construct initial data for (9.3) such that the corresponding solution yields the desired asymptotic data for  $w_a$ . In Section 9.4 we demonstrate that this is possible. In all of the arguments, it is of great importance to keep track of the constants involved, since we, in the end, are interested in an infinite sequence of equations and a corresponding infinite sequence of times  $T_{\text{ode}}$ .

#### 9.1 Equations and algebraic decompositions

In this chapter we are concerned with the non-oscillatory, or ODE, era of a Fourier mode of an asymptotically silent equation. However, it is convenient to formulate general results that do not refer to the PDE origin of the problem. Let us start by describing the class of equations we are interested in.

#### 9.1.1 Class of equations

We are interested in equations of the form (9.3) for a  $\mathbb{C}^k$ -valued function  $v, 1 \leq k \in \mathbb{Z}$ , where F is a given smooth  $\mathbb{C}^k$ -valued function on  $\mathbb{R}$ ;  $A \in \mathbf{M}_k(\mathbb{C})$ ; and  $A_{\text{rem}}$  is a smooth function from  $\mathbb{R}$  to  $\mathbf{M}_k(\mathbb{C})$ . We assume, moreover, that there are constants  $C_{\text{rem}}, T_{\text{ode}} \geq 0$  and  $\beta_{\text{rem}} > 0$  such that

$$\|A_{\rm rem}(t)\| \le C_{\rm rem} e^{-\beta_{\rm rem}\bar{t}} \tag{9.9}$$

for  $t \geq T_{\text{ode}}$ , where  $\bar{t} := t - T_{\text{ode}}$ . We refer to  $\beta_{\text{rem}}$  as the *ODE margin*.

#### 9.1.2 Motivation and goal

The equation (9.3) should be thought of as a model equation for the Fourier modes of an asymptotically silent equation. In that setting, there are infinitely many different modes (corresponding to infinitely many different matrices  $A_{\rm rem}$ ). However, there are universal constants  $\beta_{\rm rem}$  and  $C_{\rm rem}$  such that (9.9) holds for  $T_{\rm ode}$  large enough. Note, however, that  $T_{\rm ode}$  does depend on the mode. Our goal is to estimate the difference between v and an appropriate solution to the equation that results by removing the second term on the right hand side from (9.3). The estimate should be valid for  $t \geq T_{\rm ode}$ . Since A,  $\beta_{\rm rem}$  and  $C_{\rm rem}$  should be thought of as universal (independent of mode), it is not of any greater importance to spell out the dependence of constants (appearing in the estimates) on  $C_{\rm rem}$ ,  $\beta_{\rm rem}$  and A explicitly. However, we need to keep very careful track of the dependence on  $T_{\rm ode}$ .

#### **9.1.3** Decomposition of A

The complex  $k \times k$ -matrix A plays a central role in (9.3). Many of the results below will be phrased in terms of different algebraic decompositions of A. For the sake of clarity, we here collect all the details of these decompositions in one subsection; see also Section 24.2 below for a summary of some of the basic facts concerning the Jordan normal form and the matrix exponential. Our motivation for decomposing A is described in connection with (9.7) and (9.8). In particular, the starting point, motivating the decomposition, is the fact that the ODE margin might be strictly smaller than the real eigenvalue spread of A; i.e., that the inequality  $\beta_{\rm rem} < \text{Rsp}A$  might hold.

**Lemma 9.1.** Given  $1 \leq k \in \mathbb{Z}$ ,  $0 < \beta \in \mathbb{R}$  and  $A \in \mathbf{M}_k(\mathbb{C})$ , there is a  $T \in \mathbf{GL}_k(\mathbb{C})$  such that  $A_J := T^{-1}AT$  has the following properties. To begin with,

$$A_J = T^{-1}AT = \operatorname{diag}\{A_a, A_b\},\$$

where  $A_a \in \mathbf{M}_{k_a}(\mathbb{C})$ ;  $A_b \in \mathbf{M}_{k_b}(\mathbb{C})$ ;  $1 \leq k_a \in \mathbb{Z}$  and  $0 \leq k_b \in \mathbb{Z}$  are such that  $k_a + k_b = k$ ; and  $A_a$  and  $A_b$  consist of Jordan blocks. Moreover,  $\operatorname{Rsp} A_a < \beta$ ;  $\kappa_{\max}(A_a) = \kappa_{\max}(A)$ ; and  $\kappa_{\max}(A_b) \leq \kappa_{\max}(A) - \beta$  (assuming  $k_b \geq 1$ ).

**Remark 9.2.** See Section 24.2 below for the definition of Jordan blocks and a summary of some of the basic facts concerning the Jordan normal form.

**Remarks 9.3.** If  $k_b = 0$ , then diag $\{A_a, A_b\}$  should be interpreted as equalling  $A_a$ . Moreover, the notation  $\kappa_{\max}(A)$  is introduced in Definition 4.3. The matrices T,  $A_a$  etc. are not unique. However, the dimensions  $k_a$  and  $k_b$  are well defined.

*Proof.* The statement follows by transforming A to Jordan normal form and then arranging the Jordan blocks appropriately.

**Definition 9.4.** Given  $1 \leq k \in \mathbb{Z}$ ,  $A \in \mathbf{M}_k(\mathbb{C})$  and  $0 < \beta \in \mathbb{R}$ , let  $k_a$  and  $k_b$  be the integers obtained by appealing to Lemma 9.1. Below,  $k_a$  and  $k_b$  are referred to as the dimensions of the first and second subspaces (respectively) of the  $\beta$ , A-decomposition of  $\mathbb{C}^k$ . If  $B \in \mathbf{M}_k(\mathbb{C})$  can be

written  $B = \text{diag}\{B_a, B_b\}$ , where  $B_a \in \mathbf{M}_{k_a}(\mathbb{C})$  and  $B_b \in \mathbf{M}_{k_b}(\mathbb{C})$ , then  $B_a$   $(B_b)$  is referred to as the first (second) block of the  $\beta$ , A-decomposition of B. If  $\xi \in \mathbb{C}^k$  and  $\xi_a \in \mathbb{C}^{k_a}$  collects the first  $k_a$  components of  $\xi$  and  $\xi_b \in \mathbb{C}^{k_b}$  collects the last  $k_b$  components of  $\xi$ , then  $\xi_a$   $(\xi_b)$  is referred to as the first (second) component in the  $\beta$ , A-decomposition of  $\xi$ . Finally, if T is obtained by appealing to Lemma 9.1,  $E_a := T(\mathbb{C}^{k_a} \times \{0\}^{k_b})$  and  $E_b := T(\{0\}^{k_a} \times \mathbb{C}^{k_b})$ , then  $E_a$   $(E_b)$  is referred to as the first (second) generalised eigenspace in the  $\beta$ , A-decomposition of  $\mathbb{C}^k$ . The first generalised eigenspace in the  $\beta$ , A-decomposition of  $\mathbb{C}^k$  is also denoted by  $E_{A,\beta}$ .

**Remark 9.5.** The vector space  $E_a$  ( $E_b$ ) is the direct sum of the generalised eigenspaces of A corresponding to eigenvalues in Sp $A_a$  (Sp $A_b$ ); here  $A_a$  and  $A_b$  are obtained by appealing to Lemma 9.1. In particular, the spaces  $E_a$  and  $E_b$  are uniquely determined by A and  $\beta > 0$ . Moreover, dim  $E_a = k_a$  and dim  $E_b = k_b$ . Finally, note that the present definition is consistent with Definition 4.7.

In our applications, the following estimate will be of interest.

**Lemma 9.6.** Given  $1 \leq k \in \mathbb{Z}$ ,  $A \in \mathbf{M}_k(\mathbb{C})$  and  $0 < \beta \in \mathbb{R}$ , let  $E_a$  ( $E_b$ ) be the first (second) generalised eigenspace in the  $\beta$ , A-decomposition of  $\mathbb{C}^k$ . Let  $\Pi_a : \mathbb{C}^k \to E_a$  and  $\Pi_b : \mathbb{C}^k \to E_b$  be defined by the condition that  $x = \Pi_a x + \Pi_b x$  for all  $x \in \mathbb{C}^k$ . Let  $\kappa_b := \kappa_{\max}(A_b)$  and  $d_b := d_{\max}(A_b, \kappa_b)$ , where  $A_b$  is the restriction of A to  $E_b$ ; cf. Definition 4.3. Then

$$\|e^{At}\Pi_b\| \le C_A \langle t \rangle^{d_b - 1} e^{\kappa_b t}$$

for all  $t \ge 0$ , where  $C_A$  only depends on A.

*Proof.* Let T,  $k_a$  and  $k_b$  be obtained by appealing to Lemma 9.1 and let  $\pi_b : \mathbb{C}^k \to \{0\}^{k_a} \times \mathbb{C}^{k_b}$  be the projection onto the last  $k_b$  elements of  $\mathbb{C}^k$ . Then  $\Pi_b = T\pi_b T^{-1}$ , so that

$$e^{At}\Pi_b = TT^{-1}e^{At}T\pi_b T^{-1} = T\operatorname{diag}\{0_{k_a}, e^{A_b t}\}T^{-1},$$

where  $A_b$  here denotes the matrix appearing in the statement of Lemma 9.1. In view of the observations concerning the matrix exponential made in Section 24.2 below, the statement of the lemma follows.

Below, the following terminology will be important.

**Definition 9.7.** Given  $1 \leq k \in \mathbb{Z}$ ,  $A \in \mathbf{M}_k(\mathbb{C})$  and  $0 < \beta \in \mathbb{R}$ , let  $A_{J,i} := i \operatorname{Im} \{A_J\}$ , where  $A_J$  is obtained by appealing to Lemma 9.1. Then  $A_{J,i}$  is diagonal and purely imaginary, and the smooth function  $R_A : \mathbb{R} \to \mathbf{M}_k(\mathbb{C})$  defined by  $R_A(t) := \exp(-A_{J,i}t)$  is referred to as a *rotation matrix* associated with A.

**Remark 9.8.** Note that  $||R_A(t)|| = ||[R_A(t)]^{-1}|| = 1$ .

In our setting, we are given a matrix  $A \in \mathbf{M}_k(\mathbb{C})$  and a  $0 < \beta_{\text{rem}} \in \mathbb{R}$ . Appealing to Lemma 9.1 with  $\beta$  replaced by  $\beta_{\text{rem}}$  then yields a matrix  $A_J$ , which we here refer to as  $A_{\text{pre}}$ . Let  $\kappa_1 := \kappa_{\max}(A)$ , cf. Definition 4.3, and consider  $A_{\text{inter}} := \text{Re}\{A_{\text{pre}}\} - \kappa_1 \text{Id}_k$ . This is a real  $k \times k$ -matrix consisting of Jordan blocks, all of whose diagonal elements are non-positive. It would be convenient if the Jordan blocks in  $A_{\text{inter}}$  with strictly negative eigenvalues were negative definite. However, this need not be the case. On the other hand, it can be arranged.

**Lemma 9.9.** Let  $J \in \mathbf{M}_k(\mathbb{R})$  be a Jordan block whose diagonal components equal  $\lambda \in \mathbb{R}$ . For every  $0 < \epsilon \in \mathbb{R}$ , there is then a diagonal matrix  $D \in \mathbf{GL}_k(\mathbb{R})$  such that for every  $x \in \mathbb{R}^k$ ,

$$x^t D^{-1} J D x \le (\lambda + \epsilon) |x|^2$$

Moreover,  $||D|| = \max\{1, \epsilon^{k-1}\}$  and  $||D^{-1}|| = \max\{1, \epsilon^{-(k-1)}\}.$ 

#### 9.2. ROUGH ESTIMATE OF SOLUTIONS

**Remark 9.10.** The difference between J and  $D^{-1}JD$  is that the off-diagonal 1's in J have been replaced by  $\epsilon$ . Below, it will be convenient to refer to a matrix of the form  $D^{-1}JD$  as a generalised Jordan block. We also use this terminology when J is complex.

*Proof.* The matrix D can be chosen according to 
$$D = \text{diag}\{1, \epsilon, \dots, \epsilon^{k-1}\}$$
.

Applying this to the individual Jordan blocks leads to the following definition.

**Definition 9.11.** Given  $1 \leq k \in \mathbb{Z}$ ,  $A \in \mathbf{M}_k(\mathbb{C})$  and  $0 < \beta \in \mathbb{R}$ , let T be a matrix of the form obtained in Lemma 9.1. Let, moreover,  $\kappa_1 := \kappa_{\max}(A)$ . Then there is a diagonal matrix  $D \in \mathbf{GL}_k(\mathbb{R})$  such that if  $T_A := TD$ , then  $T_A^{-1}AT_A$  consists of generalised Jordan blocks. Moreover, all the generalised Jordan blocks of  $J_A := \operatorname{Re}\{T_A^{-1}AT_A\} - \kappa_1\operatorname{Id}_k$  corresponding to negative eigenvalues are negative definite. A matrix  $T_A$  obtained in this way is referred to as a  $\beta$ -normalised A-conjugation matrix. Moreover, the matrix  $J_A$  is referred to as a real  $\beta$ -normalised Jordan decomposition of A.

**Remarks 9.12.** The notion of a generalised Jordan block is introduced in Remark 9.10. Note that the definition of  $T_A$  leads to  $||T_A||$  and  $||T_A^{-1}||$  depending on the distance between the largest and the second largest real part of an eigenvalue of A. On the other hand,  $||T_A||$  and  $||T_A^{-1}||$  depend only on A (partly through this distance).

**Remarks 9.13.** Note that the matrices  $R_A(t)$ ,  $T_A$  and  $J_A$  are dependent on a number of choices. For this reason we speak of a rotation matrix instead of the rotation matrix etc. However, we assume the different choices to be consistent with each other in the sense that they are related to each other as described above. Then  $R_A(t)$  and  $A_{J,i}$  (cf. Definition 9.7) commute with  $J_A$ , since for each generalised Jordan block in  $J_A$ , the corresponding diagonal components in  $A_{J,i}$  are constant.

For future reference, it is of interest to note that  $e^{At}$  can be expressed in terms of the above decomposition of A. In fact, given  $A \in \mathbf{M}_k(\mathbb{C})$  and  $0 < \beta \in \mathbb{R}$ , let T and J be the  $T_A$  and  $J_A$  (respectively) constructed in Definition 9.11. Let, moreover, R be the matrix valued function  $R_A$  constructed in Definition 9.7. Then

$$T^{-1}AT = A_{J,i} + J + \kappa_1 \mathrm{Id}_k, \tag{9.10}$$

so that

$$e^{At} = e^{\kappa_1 t} T[R(t)]^{-1} e^{Jt} T^{-1}; (9.11)$$

note that R(t) commutes with J, cf. Remarks 9.13. Alternately, since  $T^{-1}AT = \text{diag}\{A_a, A_b\}$  (this is true irrespective of whether T is the matrix obtained in Lemma 9.1 or if T is the matrix  $T_A$  introduced in Definition 9.11),

$$e^{At} = T \operatorname{diag}\{e^{A_a t}, e^{A_b t}\}T^{-1}.$$

#### 9.2 Rough estimate of solutions

The first step in the analysis of the asymptotics is to derive an estimate for |v(t)| for  $t \ge T_{\text{ode}}$ , where v is a solution to (9.3). This is the subject of the present section.

**Lemma 9.14.** Consider the equation (9.3), where A,  $A_{\text{rem}}$  and F satisfy the assumptions described in Subsection 9.1.1. There is a constant C, depending only on  $C_{\text{rem}}$ ,  $\beta_{\text{rem}}$  and the matrix A, such that if v is a solution to (9.3), then

$$|v(t)| \le C \langle \bar{t} \rangle^{d_1 - 1} e^{\kappa_1 \bar{t}} |v(T_{\text{ode}})| + C \int_{T_{\text{ode}}}^t \langle t - s \rangle^{d_1 - 1} e^{\kappa_1 (t - s)} |F(s)| ds$$
(9.12)

for  $t \ge T_{\text{ode}}$ , where  $\bar{t} = t - T_{\text{ode}}$ ,  $\kappa_1 := \kappa_{\max}(A)$  and  $d_1 := d_{\max}(A, \kappa_1)$ ; cf. Definition 4.3.

**Remark 9.15.** The constant C may depend on, e.g., the reciprocal of the difference between the largest and the second largest real part of an eigenvalue of A. If one is interested in applying the result for an infinite set of matrices A, this has to be kept in mind.

*Proof.* Denoting the sum of A and  $A_{\text{rem}}(t)$  by B(t), the equation (9.3) takes the same form as (9.1). Let  $\Phi(t; t_0)$  denote the solution to the initial value problem

$$\frac{d\Psi}{dt}(t) = B(t)\Psi(t), \quad \Psi(t_0) = \mathrm{Id}_k.$$

Then the solution to (9.3) with initial data  $v_0$  at  $T_{ode}$  can be written

$$v(t) = \Phi(t; T_{\text{ode}})v_0 + \int_{T_{\text{ode}}}^t \Phi(t; s)F(s)ds.$$
(9.13)

It is thus of interest to estimate  $\|\Phi(t;s)\|$  for  $T_{ode} \leq s \leq t$ . Since  $|\Phi(t;s)\xi| = |v(t)|$ , where v is the solution to the initial value problem

$$\frac{dv}{dt}(t) = B(t)v(t), \quad v(s) = \xi,$$
(9.14)

it is clear that we only need to estimate  $|v(t)|/|\xi|$  for solutions v to (9.14).

Given A and  $\beta_{\text{rem}}$ , let  $R(t) := R_A(t)$ ,  $T := T_A$  and  $J := J_A$  be the function/matrices obtained by appealing to Definitions 9.7 and 9.11. Recall that (9.10) holds and that R(t) and J commute due to Remarks 9.13. Let v be the solution to (9.14) and define w by

$$w(t) := e^{-\kappa_1 t} R(\bar{t}) T^{-1} v(t).$$
(9.15)

Then

$$\dot{w} = Jw + A_{\text{rest}}w,\tag{9.16}$$

where

$$\|A_{\text{rest}}(t)\| \le C e^{-\beta_{\text{rem}}t} \tag{9.17}$$

for  $t \geq T_{\text{ode}}$ , and the constant C only depends on  $C_{\text{rem}}$ ,  $\beta_{\text{rem}}$ , ||T|| and  $||T^{-1}||$ . Let  $J_{\text{tr}}$  be the truncated version of J, obtained from J by setting the Jordan blocks with zero eigenvalue to zero. Let  $J_{\text{diff}} = J - J_{\text{tr}}$  and define u and  $B_{\text{rest}}$  by

$$u = e^{-J_{\text{diff}}(t-s)}w, \quad B_{\text{rest}}(t) = e^{-J_{\text{diff}}(t-s)}A_{\text{rest}}(t)e^{J_{\text{diff}}(t-s)}.$$
 (9.18)

Then

$$\dot{u} = J_{\rm tr} u + B_{\rm rest} u, \tag{9.19}$$

where

$$\|B_{\text{rest}}(t)\| \le C\langle t-s\rangle^{2(d_1-1)} e^{-\beta_{\text{rem}}\overline{t}}$$
(9.20)

for  $t \geq T_{\text{ode}}$ , where the constant C only depends on  $C_{\text{rem}}$ ,  $\beta_{\text{rem}}$  and A, and we have used the notation (4.5). The advantage of the version (9.19) of the equation is that  $J_{\text{tr}}$  is negative semidefinite. Moreover, even in the case that  $B_{\text{rest}}$  in (9.19) equals zero, a solution u to (9.19) is typically bounded, but no more. Estimate

$$\frac{d}{dt}|u|^2 = \langle u, \dot{u} \rangle + \langle \dot{u}, u \rangle \le 2 \|B_{\text{rest}}\| \|u\|^2.$$

Integrating this estimate yields

$$|u(t)| \le C|u(s)|,$$
 (9.21)

for all  $T_{\text{ode}} \leq s \leq t$ , where C only depends on  $C_{\text{rem}}$ ,  $\beta_{\text{rem}}$  and A. Returning to (9.15) and (9.18), the estimate (9.21) yields

$$|v(t)| \le C \langle t-s \rangle^{d_1 - 1} e^{\kappa_1 (t-s)} |v(s)|,$$

where C has the same dependence as in the case of (9.21). In particular,

$$\|\Phi(t;s)\| \le C \langle t-s \rangle^{d_1-1} e^{\kappa_1(t-s)}$$

for  $T_{\text{ode}} \leq s \leq t$ , where C has the same dependence as in the case of (9.21). Combining this estimate with (9.13) yields the conclusion of the lemma.
## 9.3 Detailed asymptotics

Consider (9.12). If F = 0, this estimate is optimal (strictly speaking, we only demonstrate that this is true in Section 9.4 below, but in the case that  $A_{\rm rem} = 0$ , it is obvious). Moreover, the first term on the right hand side of (9.12) exhibits the asymptotic growth associated with solutions to  $\dot{v} = Av$ . On the other hand, if |F| grows very quickly, the second term on the right hand side of (9.12) might be dominant. Before proceeding to a derivation of more detailed information concerning the asymptotics, it is necessary to decide which type of situation is of greatest interest: the case where F dominates the evolution, or the case when the contribution from F is, at worst, comparable with solutions to the homogeneous system associated with (9.3). Here, we are mainly interested in the latter case, and for that reason, we assume that (9.4) holds. Then (9.12) implies that the growth of v is bounded by  $\langle \bar{t} \rangle^{d_1-1} e^{\kappa_1 t}$ . This is a natural starting point for deriving more detailed information concerning the asymptotics, the subject of the present section.

**Lemma 9.16.** Consider the equation (9.3), where A,  $A_{\text{rem}}$  and F satisfy (9.4) and the assumptions described in Subsection 9.1.1. Given notation as in Subsection 9.1.1, there is then a constant C and a non-negative integer N, where C only depends on  $C_{\text{rem}}$ ,  $\beta_{\text{rem}}$  and A, and N only depends on k, such that the following holds. Fix  $0 < \beta \leq \beta_{\text{rem}}$ . For every solution v to (9.3), there is a unique vector  $v_{\infty} \in E_{A,\beta}$  such that

$$\left| v(t) - e^{At} v_{\infty} - \int_{T_{\text{ode}}}^{t} e^{A(t-s)} F(s) ds \right|$$

$$\leq C \langle \bar{t} \rangle^{N} e^{(\kappa_{1}-\beta)\bar{t}} \left[ |v(T_{\text{ode}})| + e^{\kappa_{1} T_{\text{ode}}} \|F\|_{A} \right]$$

$$(9.22)$$

for all  $t \geq T_{\text{ode}}$ , where  $\kappa_1 := \kappa_{\max}(A)$  and the notation  $E_{A,\beta}$  is introduced in Definition 9.4. Moreover,  $v_{\infty}$  can be written

$$v_{\infty} = e^{-AT_{\rm ode}} u_{\infty},\tag{9.23}$$

where  $u_{\infty} \in E_{A,\beta}$ ,

$$|u_{\infty}| \le C \left[ |v(T_{\text{ode}})| + e^{\kappa_1 T_{\text{ode}}} \|F\|_A \right],$$
(9.24)

and C only depends on  $C_{\text{rem}}$ ,  $\beta_{\text{rem}}$  and A.

**Remark 9.17.** In the ODE setting, it may seem unnatural not to assume  $\beta = \beta_{\text{rem}}$ . The reason for this is that, the larger the  $\beta$ , the more detailed the asymptotics. However, we wish to apply this lemma in the PDE setting, in which case we are interested in an infinite number of ODE's (one for each mode). In that setting, a larger  $\beta$  is not only associated with more detailed information concerning the asymptotics, it is also associated with a larger loss of derivatives. In some applications, it might be important to minimise the loss of derivatives (while still retaining a certain amount of asymptotic information). This is the reason for not fixing the value of  $\beta$  to equal  $\beta_{\text{rem}}$ .

**Remark 9.18.** The constants  $C_{\text{rem}}$ ,  $\beta_{\text{rem}}$  and  $T_{\text{ode}}$  are the ones appearing in connection with the assumption (9.9).

**Remark 9.19.** Remark 9.15 is equally relevant here.

**Remark 9.20.** Combining (9.23) and (9.24) yields a crude estimate of  $|v_{\infty}|$ . However, it is of interest to obtain more detailed information. This is the reason for including (9.23), (9.24) and the statement that  $u_{\infty} \in E_{A,\beta}$ .

*Proof.* Uniqueness. Let v be a solution to (9.3). Assume that there are  $v_{\infty,i} \in E_{A,\beta}$ , i = 1, 2, such that

$$\left| v(t) - e^{At} v_{\infty,i} - \int_{T_{\text{ode}}}^{t} e^{A(t-s)} F(s) ds \right| \le C_i \langle \bar{t} \rangle^{N_i} e^{(\kappa_1 - \beta)\bar{t}}$$

for all  $t \ge T_{\text{ode}}$  and some constants  $C_i$  and  $N_i \ge 0$  which are allowed to depend on v, F,  $T_{\text{ode}}$  as well as  $C_{\text{rem}}$ ,  $\beta_{\text{rem}}$  and A. Then

$$|e^{At}(v_{\infty,1} - v_{\infty,2})| \le (C_1 + C_2) \langle \bar{t} \rangle^{N_1 + N_2} e^{(\kappa_1 - \beta)\bar{t}}$$

for all  $t \ge T_{\text{ode}}$ . However, if  $v_{\infty,1} \ne v_{\infty,2}$ , then the left hand side of this inequality grows more quickly than the right hand side, which leads to a contradiction. Therefore  $v_{\infty,1} = v_{\infty,2}$ , and  $v_{\infty}$  is uniquely determined by the estimate (9.22).

**Existence.** To begin with, we introduce notation which is similar to, but not exactly the same as, the one introduced in the proof of Lemma 9.14. Given A and  $\beta$ , let  $R(t) := R_A(t)$ ,  $T := T_A$  and  $J := J_A$  be the function/matrices obtained by appealing to Definitions 9.7 and 9.11. Recall that ||T|| and  $||T^{-1}||$  only depend on A; that (9.10) holds; and that R(t) and J commute due to Remarks 9.13. Note, moreover, that T, J and R can be chosen independently of  $\beta$ . However, what  $A_a$  and  $A_b$  are depends on  $\beta$ . Define w and G by

$$w(t) := e^{-\kappa_1 t} R(\bar{t}) T^{-1} v(t), \quad G(t) := e^{-\kappa_1 t} R(\bar{t}) T^{-1} F(t), \tag{9.25}$$

where  $\bar{t} := t - T_{\text{ode}}$ . Then

$$\dot{w} = Jw + A_{\text{rest}}w + G,\tag{9.26}$$

where

$$\|A_{\text{rest}}\| \le C e^{-\beta_{\text{rem}}t} \tag{9.27}$$

for  $t \ge T_{\text{ode}}$ , and the constant C only depends on  $C_{\text{rem}}$  and A. Let  $J_{\text{tr}}$  be the truncated version of J, obtained from J by setting the Jordan blocks with zero eigenvalue to zero. Let  $J_{\text{diff}} = J - J_{\text{tr}}$  and define u, B and H by

$$u = e^{-J_{\text{diff}}\bar{t}}w, \quad B(t) = e^{-J_{\text{diff}}\bar{t}}A_{\text{rest}}(t)e^{J_{\text{diff}}\bar{t}}, \quad H(t) = e^{-J_{\text{diff}}\bar{t}}G(t).$$
(9.28)

Then

$$\dot{u} = J_{\rm tr}u + Bu + H,\tag{9.29}$$

where

$$|B(t)|| \le C\langle \bar{t} \rangle^{2(d_1-1)} e^{-\beta_{\rm rem}\bar{t}}$$

$$\tag{9.30}$$

for  $t \geq T_{\text{ode}}$ , where the constant C only depends on  $C_{\text{rem}}$  and A. For future reference, it is convenient to keep in mind that (9.12), (9.25) and (9.28) yield

$$|u(t)| \le C\langle \bar{t} \rangle^{2(d_1 - 1)} |w(T_{\text{ode}})| + C\langle \bar{t} \rangle^{2(d_1 - 1)} ||F||_A$$
(9.31)

for  $t \geq T_{\text{ode}}$ , where C only depends on  $C_{\text{rem}}$ ,  $\beta_{\text{rem}}$  and A.

Estimating the first component. Let  $u_a(t)$  denote the first component in the  $\beta$ , A-decomposition of u(t); cf. Definition 9.4 for an explanation of the terminology. We similarly use the notation  $w_a$ ,  $(Bu)_a$ ,  $G_a$ ,  $F_a$  etc. Let  $J_a$  and  $J_{tr,a}$  denote the first blocks in the  $\beta$ , A-decomposition of J and  $J_{tr}$ respectively; cf. Definition 9.4. Integrating the *a*'th component of (9.29) yields

$$e^{-J_a\bar{t}}w_a(t) = w_a(T_{\text{ode}}) + \int_{T_{\text{ode}}}^t e^{-J_{\text{tr},a}\bar{s}}(Bu)_a(s)ds + \int_{T_{\text{ode}}}^t e^{-J_a\bar{s}}G_a(s)ds.$$
(9.32)

Define

$$w_{\infty,a} := w_a(T_{\text{ode}}) + \int_{T_{\text{ode}}}^{\infty} e^{-J_{\text{tr},a}\bar{s}} (Bu)_a(s) ds.$$
(9.33)

Then (9.32) implies

$$w_{a}(t) - e^{J_{a}\bar{t}}w_{\infty,a} - \int_{T_{ode}}^{t} e^{J_{a}(t-s)}G_{a}(s)ds \bigg| \le \bigg| \int_{t}^{\infty} e^{-J_{a}(s-t)}e^{J_{diff,a}\bar{s}}(Bu)_{a}(s)ds \bigg|, \qquad (9.34)$$

#### 9.3. DETAILED ASYMPTOTICS

<

where  $J_{\text{diff},a}$  is defined similarly to  $J_{\text{tr},a}$ . Note that (9.30) and (9.31) yield

$$\left| \int_{T_{\text{ode}}}^{\infty} e^{-J_{\text{tr},a}\bar{s}} (Bu)_a(s) ds \right| \le C[|w(T_{\text{ode}})| + ||F||_A], \tag{9.35}$$

where C only depends on  $C_{\text{rem}}$ ,  $\beta_{\text{rem}}$  and A. Note, however, that the dependence is via, e.g.,  $(\beta_{\text{rem}} + \kappa_{\min})^{-1}$  (where  $\kappa_{\min} := \kappa_{\min}(A_a) - \kappa_1$ ; cf. Definition 4.3), a quantity which tends to infinity as  $|\kappa_{\min}|$  tends to  $\beta_{\text{rem}}$ . Combining this estimate with (9.33) yields

$$|w_{\infty,a}| \le C[|w(T_{\text{ode}})| + ||F||_A], \tag{9.36}$$

where C has the same dependence as in (9.35). Returning to (9.34), we need to estimate the right hand side. Due to (9.30) and (9.31), it can be bounded by

$$C \int_{t}^{\infty} e^{-\kappa_{\min}(s-t)} \langle \bar{s} \rangle^{N} e^{-\beta_{\operatorname{rem}} \bar{s}} [|w(T_{\operatorname{ode}})| + ||F||_{A}] ds$$
$$\leq C \langle \bar{t} \rangle^{N} e^{-\beta_{\operatorname{rem}} \bar{t}} [|w(T_{\operatorname{ode}})| + ||F||_{A}]$$

for  $t \ge T_{\text{ode}}$ , where C only depends on  $C_{\text{rem}}$ ,  $\beta_{\text{rem}}$  and A, and N is a non-negative integer depending only on k. Combining this estimate with (9.34) and (9.25) yields

$$\left| w_{a}(t) - e^{J_{a}\bar{t}} w_{\infty,a} - \int_{T_{\text{ode}}}^{t} e^{J_{a}(t-s)} G_{a}(s) ds \right|$$

$$\leq C \langle \bar{t} \rangle^{N} e^{-\beta_{\text{rem}}\bar{t}} [e^{-\kappa_{1}T_{\text{ode}}} |v(T_{\text{ode}})| + \|F\|_{A}]$$

$$(9.37)$$

for  $t \geq T_{ode}$ , where C and N have the same dependence as before.

Estimating the second component. Let us now turn to  $u_b(t)$ , the second component in the  $\beta$ , A-decomposition of u(t); cf. Definition 9.4 for an explanation of the terminology. Similarly to the above, we use the notation  $w_b$ ,  $(Bu)_b$ ,  $G_b$ ,  $F_b$  etc. In this case, (9.29) yields

$$w_b(t) = e^{J_b \bar{t}} w_b(T_{\text{ode}}) + \int_{T_{\text{ode}}}^t e^{J_b(t-s)} (Bu)_b(s) ds + \int_{T_{\text{ode}}}^t e^{J_b(t-s)} G_b(s) ds;$$
(9.38)

note that  $J_{tr,b} = J_b$ , that  $u_b = w_b$  and that  $H_b = G_b$ . To begin with

$$|e^{J_b\bar{t}}w_b(T_{\text{ode}})| \le C\langle \bar{t} \rangle^N e^{\kappa_b\bar{t}} |w_b(T_{\text{ode}})| \le C\langle \bar{t} \rangle^N e^{-\beta\bar{t}} |w_b(T_{\text{ode}})|,$$
(9.39)

where  $\kappa_b := \kappa_{\max}(A_b) - \kappa_1$ , C only depends on A and N only depends on k; note that, by definition,  $\kappa_b \leq -\beta$ . Next,

$$\left| \int_{T_{\text{ode}}}^{t} e^{J_{b}(t-s)} (Bu)_{b}(s) ds \right| \leq C \langle \bar{t} \rangle^{N} \int_{T_{\text{ode}}}^{t} e^{\kappa_{b}(t-s)} e^{-\beta_{\text{rem}}\bar{s}} ds [|w(T_{\text{ode}})| + ||F||_{A}]$$

$$\leq C \langle \bar{t} \rangle^{N} e^{-\beta \bar{t}} [|w(T_{\text{ode}})| + ||F||_{A}],$$
(9.40)

where C only depends on  $C_{\text{rem}}$ ,  $\beta_{\text{rem}}$  and A; N is a non-negative integer depending only on k; and we have appealed to (9.30) and (9.31). Combining (9.38), (9.39) and (9.40) yields

$$\left| w_b(t) - \int_{T_{\text{ode}}}^t e^{J_b(t-s)} G_b(s) ds \right| \le C \langle \bar{t} \rangle^N e^{-\beta \bar{t}} [|w(T_{\text{ode}})| + ||F||_A]$$
(9.41)

for  $t \geq T_{ode}$ , where C only depends on  $C_{rem}$ ,  $\beta_{rem}$  and A, and N is a non-negative integer depending only on k.

Summing up. Let  $w_{\infty} \in \mathbb{C}^k$  be such that the first component in the  $\beta$ , A-decomposition of  $w_{\infty}$  is given by  $w_{\infty,a}$ , defined in (9.33), and the second component in the  $\beta$ , A-decomposition of  $w_{\infty}$  vanishes. Then (9.37) and (9.41) yield

$$\left| w(t) - e^{J\bar{t}} w_{\infty} - \int_{T_{\text{ode}}}^{t} e^{J(t-s)} G(s) ds \right|$$

$$\leq C \langle \bar{t} \rangle^{N} e^{-\beta \bar{t}} [e^{-\kappa_{1} T_{\text{ode}}} |v(T_{\text{ode}})| + \|F\|_{A}]$$

$$(9.42)$$

for  $t \geq T_{\text{ode}}$ , where C only depends on  $C_{\text{rem}}$ ,  $\beta_{\text{rem}}$  and A; and N is a non-negative integer depending only on k. Appealing to (9.11) and (9.25) yields (9.22), where  $v_{\infty}$  is given by (9.23) and  $u_{\infty} \in E_{A,\beta}$  is defined by

$$u_{\infty} := T e^{\kappa_1 T_{\text{ode}}} w_{\infty}. \tag{9.43}$$

Combining this definition with (9.25) and (9.36) yields (9.24), and the lemma follows.

# 9.4 Specifying data at infinity

Lemma 9.16 contains a statement concerning the leading order behaviour of solutions. However, it is also of interest to prove that, given asymptotic data of the form  $v_{\infty}$  (described in the statement of Lemma 9.16), there is a solution with asymptotics as in (9.22). In order to obtain such a result, it is sufficient to consider the homogeneous version of the equation (9.3).

Lemma 9.21. Consider the equation

$$\dot{v}(t) = Av(t) + A_{\rm rem}(t)v(t),$$
(9.44)

where  $A \in \mathbf{M}_k(\mathbb{C})$  and  $A_{\text{rem}} \in C^{\infty}[\mathbb{R}, \mathbf{M}_k(\mathbb{C})]$  satisfies the estimate (9.9) for some choice of  $T_{\text{ode}} \geq 0$ ,  $C_{\text{rem}}$  and  $\beta_{\text{rem}} > 0$ . Let  $0 < \beta \leq \beta_{\text{rem}}$  and  $E_{A,\beta}$  be the first generalised eigenspace in the  $\beta$ , A-decomposition of  $\mathbb{C}^k$ . Then there is a linear injective map  $\Psi_{\infty} : E_{A,\beta} \to \mathbb{C}^k$  such that if  $v_{\infty} \in E_{A,\beta}$  and v is the solution to (9.44) satisfying  $v(T_{\text{ode}}) = \Psi_{\infty}(v_{\infty})$ , then v satisfies (9.22) with F = 0. Moreover, for  $u_{\infty} \in E_{A,\beta}$ 

$$|\Psi_{\infty}(e^{-AT_{\text{ode}}}u_{\infty})| \le C|u_{\infty}|,\tag{9.45}$$

where C only depends on  $C_{\text{rem}}$ ,  $\beta_{\text{rem}}$  and A.

**Remark 9.22.** In the statement of the lemma, as well as the proof, we use the terminology introduced in Subsection 9.1.3.

Remark 9.23. The result can be used to derive conclusions concerning the inhomogeneous equation (9.3). In order to justify this statement, assume that  $||F||_A < \infty$  and let  $v_{\text{part}}$  be the solution to (9.3) corresponding to the initial data  $v_{\text{part}}(T_{\text{ode}}) = 0$ . Let  $v_{\text{part},\infty} \in E_{A,\beta}$  be such that (9.22) holds with v and  $v_{\infty}$  replaced by  $v_{\text{part}}$  and  $v_{\text{part},\infty}$  respectively. Given  $v_{\infty} \in E_{A,\beta}$ , let  $v_{\text{hom}}$  be the solution to (9.44) corresponding to initial data  $v_{\text{hom}}(T_{\text{ode}}) = \Psi_{\infty}(v_{\infty} - v_{\text{part},\infty})$ . Then  $v := v_{\text{part}} + v_{\text{hom}}$  solves (9.3) and satisfies (9.22). In other words, we are also allowed to specify the asymptotic data  $v_{\infty} \in E_{A,\beta}$  in the case of the inhomogeneous equation (9.3), assuming  $||F||_A < \infty$ . In order to obtain estimates, let  $u_{\text{part},\infty} \in E_{A,\beta}$  be such that (9.23) and (9.24) hold with  $v, v_{\infty}$  and  $u_{\infty}$  replaced by  $v_{\text{part}}, v_{\text{part},\infty}$  and  $u_{\text{part},\infty}$  respectively. Due to (9.24),  $|u_{\text{part},\infty}| \leq Ce^{\kappa_1 T_{\text{ode}}} ||F||_A$ , where C only depends on  $C_{\text{rem}}, \beta_{\text{rem}}$  and A. Due to (9.23),  $v_{\text{part},\infty} = e^{-AT_{\text{ode}}} u_{\text{part},\infty}$ . Representing  $v_{\infty}$  according to  $v_{\infty} = e^{-AT_{\text{ode}}} u_{\infty}$ , it follows that

$$|v(T_{\text{ode}})| \leq |v_{\text{part}}(T_{\text{ode}})| + |v_{\text{hom}}(T_{\text{ode}})| = |\Psi_{\infty}(v_{\infty} - v_{\text{part},\infty})|$$
  
$$= |\Psi[e^{-AT_{\text{ode}}}(u_{\infty} - u_{\text{part},\infty})]|$$
  
$$\leq C|u_{\infty} - u_{\text{part},\infty}| \leq C(|u_{\infty}| + e^{\kappa_{1}T_{\text{ode}}}||F||_{A}),$$
  
(9.46)

where C is a constant depending only on  $C_{\text{rem}}$ ,  $\beta_{\text{rem}}$  and A.

**Remark 9.24.** The equality  $E_{A,\beta} = \mathbb{C}^k$  holds if  $\beta > \operatorname{Rsp}A$ ; cf. Definition 4.3. Moreover, if  $E_{A,\beta} = \mathbb{C}^k$ , then the map  $\Psi_{\infty}$  is an isomorphism. This means that the solution is uniquely specified in terms of the "data at infinity",  $v_{\infty}$ . Due to Remark 9.23, the same holds for inhomogeneous equations such that  $||F||_A < \infty$ . Finally,  $|v(T_{\text{ode}})|$  can be estimated in terms of the asymptotic data and F; cf. (9.46).

#### 9.4. SPECIFYING DATA AT INFINITY

*Proof.* Let T, J and R be defined as in the proof of Lemma 9.16. Define w as in (9.25) and let  $J_a$  be the first block in the  $\beta$ , A-decomposition of J; cf. Definition 9.4. Let

$$J_{a,0} = \text{diag}\{J_a, 0\}, \quad \bar{B}(t) = e^{-J_{a,0}\bar{t}}A_{\text{rest}}(t)e^{J_{a,0}\bar{t}},$$
$$\bar{J} = J - J_{a,0}, \quad \bar{u}(t) = e^{-J_{a,0}\bar{t}}w(t),$$

where  $A_{\text{rest}}$  is the matrix appearing in (9.26). Then (9.44) can be written

$$\dot{\bar{u}} = \bar{J}\bar{u} + \bar{B}\bar{u},\tag{9.47}$$

where we have appealed to (9.26) with G = 0. Due to (9.27),

$$\|\bar{B}(t)\| \le C \langle \bar{t} \rangle^N e^{-\beta_{\min}\bar{t}} \tag{9.48}$$

for  $t \geq T_{\text{ode}}$ , where  $\beta_{\min} = \beta_{\text{rem}} + \kappa_{\min}$ ;  $\kappa_{\min} := \kappa_{\min}(A_a) - \kappa_1$ , where  $A_a$  is the first block in the  $\beta$ , A-decomposition of  $T^{-1}AT$ ; C is a constant depending only on  $C_{\text{rem}}$ ,  $\beta_{\text{rem}}$  and A; and N is a non-negative integer depending only on k. Note that  $\beta_{\min} > 0$  and that  $\overline{J}$  is negative semi-definite. Let  $t_0 \geq T_{\text{ode}}$ . Due to (9.47) and the fact that  $\overline{J}$  is negative semi-definite,

$$\frac{d}{dt}|\bar{u}|^2 \le 2\|\bar{B}\| \cdot |\bar{u}|^2$$

so that

$$|\bar{u}(t)| \le \exp\left(\int_{t_0}^t \|\bar{B}(s)\| ds\right) |\bar{u}(t_0)|$$
(9.49)

for  $t \ge t_0$ . Note, in particular, that by choosing  $t_0$  to be large enough, the first factor on the right hand side of (9.49) can be assumed to be as close to 1 as we wish. Let  $\bar{u}_a(t)$  and  $\bar{u}_b(t)$  denote the first and second components in the  $\beta$ , A-decomposition of  $\bar{u}(t)$  respectively (we use similar notation for the same decomposition of other vectors and vector valued functions). Then

$$\bar{u}_a(t) = \bar{u}_a(t_0) + \int_{t_0}^t (\bar{B}\bar{u})_a(s)ds, \qquad (9.50)$$

so that

$$|\bar{u}_a(t) - \bar{u}_a(t_0)| \le C \int_{t_0}^t \langle \bar{s} \rangle^N e^{-\beta_{\min}\bar{s}} ds \exp\left(\int_{t_0}^\infty \|\bar{B}(s)\| ds\right) |\bar{u}(t_0)|$$

for  $t \ge t_0$ , where we have appealed to (9.48) and (9.49); C only depends on  $C_{\text{rem}}$ ,  $\beta_{\text{rem}}$  and A; and N only depends on k. For any  $\epsilon > 0$ , it is thus possible to choose  $t_0$  so that

$$|\bar{u}_a(t) - \bar{u}_a(t_0)| \le \epsilon |\bar{u}(t_0)| \tag{9.51}$$

for  $t \ge t_0$  (where  $t_0$  only depends on  $C_{\text{rem}}$ ,  $\beta_{\text{rem}}$ , A and  $\epsilon$ ). Given a  $t_0$  such that (9.51) holds with  $\epsilon$  replaced by 1/2, define the map  $L_a : \mathbb{C}^{k_a} \to \mathbb{C}^{k_a}$  as follows (here  $k_a$  is the dimension of the first subspace of the  $\beta$ , A-decomposition of  $\mathbb{C}^k$ ; cf. Definition 9.4). Given  $\eta \in \mathbb{C}^{k_a}$ , let  $\xi \in \mathbb{C}^k$  be such that  $\xi_a = \eta$  and  $\xi_b = 0$ . Let  $\bar{u}$  be the solution to (9.47) with  $\bar{u}(t_0) = \xi$  and define  $L_a \eta$  by

$$L_a \eta := \lim_{t \to \infty} \bar{u}_a(t). \tag{9.52}$$

Due to (9.48), (9.49) and (9.50), it is clear that the limit on the right hand side of (9.52) exists. Thus  $L_a\eta$  is well defined. Moreover, it is clear that  $L_a$  is linear. Due to the fact that (9.51) holds with  $\epsilon$  replaced by 1/2 and the fact that  $\bar{u}_b(t_0) = 0$ ,

$$|L_a\eta - \bar{u}_a(t_0)| \le \frac{1}{2}|\bar{u}_a(t_0)|.$$

In particular,  $|\eta|/2 \leq |L_a\eta|$ , so that  $L_a$  is injective. Since  $L_a$  is a linear map between spaces of the same finite dimension, it follows that  $L_a$  is an isomorphism. Moreover,

$$|L_a^{-1}\eta| \le 2|\eta|. \tag{9.53}$$

Note that  $L_a^{-1}$  (essentially) maps data at infinity to data at  $t_0$ . In the end, we wish to translate the data at  $t_0$  to data at  $T_{ode}$ . To this end, we wish to solve (9.47) backwards. Since  $t_0 - T_{ode}$  can be bounded by a constant depending only on  $C_{rem}$ ,  $\beta_{rem}$  and A, there is a constant C (with the same dependence) such that for all solutions  $\bar{u}$  to (9.47),

$$|\bar{u}(T_{\text{ode}})| \le C |\bar{u}(t_0)|.$$
 (9.54)

Before defining the map  $\Psi_{\infty}$  appearing in the statement of the lemma, we need to clarify how the limit of  $\bar{u}_a(t)$  is related to the  $v_{\infty} \in E_{A,\beta}$  appearing in (9.22). However, returning to (9.32) and (9.33) (and recalling that G = 0 in the present setting), it is clear that

$$\lim_{t \to \infty} \bar{u}_a(t) = \lim_{t \to \infty} e^{-J_a \bar{t}} w_a(t) = w_{\infty,a}.$$
(9.55)

On the other hand, (9.23) and (9.43) imply that

$$v_{\infty} = e^{-AT_{\text{ode}}} u_{\infty} = e^{-AT_{\text{ode}}} T e^{\kappa_1 T_{\text{ode}}} \iota_a(w_{\infty,a}),$$

where  $\iota_a : \mathbb{C}^{k_a} \to \mathbb{C}^k$  is defined as follows: if  $\eta \in \mathbb{C}^{k_a}$ , then  $\iota_a(\eta)$  is the vector  $\xi \in \mathbb{C}^k$  such that  $\xi_a = \eta$  and  $\xi_b = 0$ . Note also that for each  $v_{\infty} \in E_{A,\beta}$ , there is a unique  $\eta \in \mathbb{C}^{k_a}$  such that

$$v_{\infty} = e^{-AT_{\text{ode}}} T e^{\kappa_1 T_{\text{ode}}} \iota_a(\eta) \tag{9.56}$$

holds. Define  $\Psi_{\infty}(v_{\infty})$  to be the composition of the following maps: first the map which takes  $v_{\infty} \in E_{A,\beta}$  to the  $\eta \in \mathbb{C}^{k_a}$  such that (9.56) holds; second, the map which takes  $\eta$  to  $\xi := \iota_a \circ L_a^{-1}(\eta)$ ; and, third, the map which takes  $\xi \in \mathbb{C}^k$  to

$$v_0 := e^{\kappa_1 T_{\text{ode}}} T \bar{u}(T_{\text{ode}}), \qquad (9.57)$$

where  $\bar{u}$  is the solution to (9.47) with initial data  $\bar{u}(t_0) = \xi$ . Note that  $\Psi_{\infty}$  is linear and injective, and that if (9.23) holds, then

$$\begin{aligned} |\Psi_{\infty}(e^{-AT_{\text{ode}}}u_{\infty})| &\leq Ce^{\kappa_{1}T_{\text{ode}}}|\bar{u}(T_{\text{ode}})| \leq Ce^{\kappa_{1}T_{\text{ode}}}|\bar{u}(t_{0})| \leq Ce^{\kappa_{1}T_{\text{ode}}}|\eta| \\ &\leq C|u_{\infty}|, \end{aligned}$$

where C only depends on  $C_{\text{rem}}$ ,  $\beta_{\text{rem}}$  and A; and we have appealed to (9.53), (9.54) and (9.57). Thus (9.45) holds. Assume now that v is the solution to (9.44) corresponding to  $v(T_{\text{ode}}) = \Psi_{\infty}(v_{\infty})$ . Then v satisfies (9.22) with F = 0 and  $v_{\infty}$  replaced by some  $\bar{v}_{\infty} \in E_{A,\beta}$ . Moreover, Lemma 9.16, in particular (9.23) and (9.43), yields  $\bar{v}_{\infty} = e^{-AT_{\text{ode}}}Te^{\kappa_1 T_{\text{ode}}}\iota_a(\bar{w}_{\infty,a})$  for some  $\bar{w}_{\infty,a} \in \mathbb{C}^{k_a}$ . In addition,

$$\lim_{t \to \infty} \bar{u}_a(t) = \bar{w}_{\infty,a},$$

where  $\bar{u}$  is defined in terms of v in the same way as in the beginning of the present proof; cf. (9.55). This means that

$$\iota_a \circ L_a^{-1}(\bar{w}_{\infty,a}) = \bar{u}(t_0),$$

due to the definition of  $L_a$ . Going through the steps defining  $\Psi_{\infty}$ , it is thus clear that  $\Psi_{\infty}(\bar{v}_{\infty}) = v(T_{\text{ode}}) = \Psi_{\infty}(v_{\infty})$ . Due to the injectivity of  $\Psi_{\infty}$ , it follows that  $v_{\infty} = \bar{v}_{\infty}$ . The lemma follows.  $\Box$ 

# Chapter 10

# Asymptotics for weakly silent, balanced and convergent equations

# 10.1 Introduction

In the present chapter, we focus on the simplest setting of interest in these notes, namely that of weakly silent, balanced and convergent equations. The goal is to derive asymptotics, as well as to obtain a (partial) characterisation of the solutions in terms of their asymptotics. The formal definition of weak silence, balance and convergence is provided in Definition 10.1 below. Let us briefly justify the terminology

Weak silence. Recall that we justified the use of the word silent in, e.g., Chapter 4. The conditions imposed in the present chapter are implied by the requirements appearing in, e.g., Proposition 4.4. In that sense, the conditions appearing here are weaker than the conditions imposed in Chapter 4. We therefore speak of weak silence.

Weak balance and weak convergence. The notion of weak balance involves bounds on  $\chi$ ,  $\mathcal{X}$ ,  $\alpha$  and  $\zeta$ . The bounds on the shift vector field and  $\mathcal{X}$  are formulated in terms of  $\sigma$  and X introduced in (24.17). Moreover, the bounds on  $\sigma$  and X follow from the requirements that  $\chi$  be future uniformly timelike and that  $\mathcal{X}$  be  $C^0$ -future bounded; cf. Lemma 25.17 and Definition 10.1 below. We also demand that the norms of  $\alpha$  and  $\zeta$  be future bounded. This is to exclude super exponential growth; cf. Section 2.1. Finally, we are mainly interested in weakly convergent equations. This means that  $\alpha$  and  $\zeta$  converge to limits. The reason we speak of *weak* balance and *weak* convergence is that we later impose similar conditions which also involve bounds on the first derivatives of the relevant objects.

Sketch of the analysis. Consider (24.16). Say that  $\mathfrak{g}$ ,  $g^{0l}$  and  $X^l$  decay to zero exponentially, and that  $\alpha$  and  $\zeta$  converge exponentially to some limits. Say, moreover, that the decay rate of  $\mathfrak{g}$  is independent of the frequency. For each frequency  $\nu(\iota)$ , there is then a  $T_{\text{ode}} \geq 0$  such that (24.16) can be reformulated as an equation of the form (9.3), where  $A_{\text{rem}}$  satisfies an estimate of the form (9.2);  $C_{\text{rem}}$  and  $\beta_{\text{rem}}$  are independent of the frequency; and  $A \in \mathbf{M}_{2m}(\mathbb{C})$ . In the case of weakly silent, balanced and convergent equations, the details of this reformulation is provided in (10.15)–(10.17) below. Due to the results of Chapter 9, we can analyse the asymptotics to the future of  $T_{\text{ode}}$ . However, in order for this analysis to be meaningful, we need to control the behaviour in the interval  $[0, T_{\text{ode}}]$ . How long is this interval? How much can the solution grow in it? Can the growth be estimated in a way which does not destroy the asymptotic analysis to the future of  $T_{\text{ode}}$ ? The assumption of weak silence implies that  $\dot{\mathfrak{g}}/\mathfrak{g} \leq -\mathfrak{b}_{\rm s} + \mathfrak{e}$ , where  $\mathfrak{b}_{\rm s} > 0$  and  $0 \leq \mathfrak{e} \in L^1([0,\infty))$  are independent of the frequency. Thus  $\mathfrak{g}(t) \leq C \exp[-\mathfrak{b}_{\rm s}(t - T_{1,\text{ode}})]$  for  $t \geq T_{1,\text{ode}}$ , where C is a constant independent of the frequency and  $T_{1,\text{ode}} = \mathfrak{b}_{\rm s}^{-1} \ln \mathfrak{g}(0)$ . Similar conclusions can be derived

for  $n_l g^{0l}$  and  $n_l X^l$ , given that  $g^{0l}$  and  $X^l$  decay exponentially. This leads to times  $T_{i,\text{ode}}$ , i = 2, 3. In comparison with the above, we can then choose  $T_{\text{ode}}$  to be the largest of the  $T_{i,\text{ode}}$ , i = 1, 2, 3. The next problem is then to estimate how the norm of the solution evolves on the interval  $[0, T_{\text{ode}}]$ . Say, for the sake of argument, that  $T_{\text{ode}}$  can be chosen to equal  $T_{1,\text{ode}}$ , and that the norm of the solution cannot grow faster than exponentially on  $[0, T_{\text{ode}}]$ , where the rate is independent of the frequency. Then the norm cannot grow more than by a factor of  $e^{CT_{\text{ode}}}$ , where the constant Cis independent of the frequency. However, since  $T_{\text{ode}}$  is a constant multiple of  $\ln \mathfrak{g}(0)$ , this means that  $e^{CT_{\text{ode}}}$  is simply a power of  $\mathfrak{g}(0)$  (moreover, the power is independent of the frequency). In other words, we can interpret the growth on the interval  $[0, T_{\text{ode}}]$  as a loss of regularity. In terms of Sobolev spaces, this loss corresponds to having to estimate the initial data in  $H^{s+s_0}$  (for some  $s_0 \geq 0$ ) in order to obtain an estimate concerning the asymptotics in  $H^s$ . In short, we can appeal to Chapter 9 in order to analyse the behaviour to the future of  $T_{\text{ode}}$ , and we can interpret the possible growth occurring in the interval  $[0, T_{\text{ode}}]$  as a loss of derivatives.

Purpose of the conditions. The question remains: how do we ensure that  $T_{ode} = T_{1,ode}$ , and that the exponential growth rate in the interval  $[0, T_{ode}]$  is independent of the frequency? Consider (24.18), a reformulation of (24.16). It is clear that if  $|\sigma|$  and ||X|| appearing in (24.18) are bounded by constants independent of  $\iota$ , then the essential dependence on  $\iota$  is contained in  $\mathfrak{g}$ . Moreover, we can choose  $T_{i,ode}$  to equal  $T_{1,ode}$  for i = 2, 3. This means that  $T_{ode} = T_{1,ode}$ . If we, in addition to the assumption of weak silence, assume  $|\sigma|$ , ||X||,  $||\alpha||$  and  $||\zeta||$  to be bounded by a constant independent of  $\iota$  (these requirements correspond to demanding that (1.2) be weakly balanced), then the exponential growth rate of the norm is bounded in  $[0, T_{ode}]$  by a constant independent of  $\iota$ . Thus the above line of reasoning can be carried out.

Failure of the conditions. If we would not impose the condition of weak balance, then the norm of solutions to (24.16) could grow super exponentially on the interval  $[0, T_{ode}]$ , even in situations when  $\mathfrak{g}$ ,  $g^{0l}$  and  $X^l$  all decay to zero exponentially; cf. Chapter 2. Again, we could interpret this growth as a loss of regularity, but the loss would be of the form  $\exp[C\mathfrak{g}(0)^{\eta}]$  for some  $C, \eta > 0$ . Clearly, this corresponds to more than a finite loss of regularity. In particular, note that simply appealing to Cauchy stability in order to estimate the evolution in the interval  $[0, T_{ode}]$  is not sufficient: Cauchy stability does not exclude the possibility of super exponential growth.

**Outline.** In Section 10.2 we give precise definitions of the notions of weak silence, balance and convergence. As indicated above, for a given frequency this leads to a division of  $[0, \infty)$  into the intervals  $[0, T_{ode}]$  and  $[T_{ode}, \infty)$ . In Chapter 9 we already considered the interval  $[T_{ode}, \infty)$ . The subject of Section 10.3 is to analyse the evolution in the interval  $[0, T_{ode}]$  for one fixed frequency. Combining this analysis with the results of Chapter 9 yields a basic energy estimate for how the  $H^s$ -norm of the solution evolves. This is the subject of Section 10.4; cf., in particular, Lemma 10.11. Combining the estimates on the interval  $[0, T_{ode}]$  with the analysis presented in Section 9.3 leads to detailed asymptotics in the future direction. This topic is treated in Section 10.5. Finally, appealing to the results of Section 9.4 leads to a partial characterisation of the solution in terms of the asymptotic data (in fact, in some situations there is a homeomorphism between initial data and the asymptotic data). This is the subject of Section 10.6.

# 10.2 Weak silence, balance and convergence

In order to give a precise meaning to the notion of a weakly silent, balanced and convergent equation, we need to introduce some terminology. Consider the equation (1.2). Assume the associated metric to be such that (M,g) is a canonical separable cosmological model manifold. Given a  $0 \neq \iota \in \mathcal{I}_{\mathrm{B}}$ , where  $\mathcal{I}_{\mathrm{B}}$  is defined in connection with (24.3), define  $\mathfrak{g}(\iota, t)$  by (24.13) and  $\ell$ ,  $\sigma$  and X by

$$\ell(\iota, t) := \ln[\mathfrak{g}(\iota, t)], \quad \sigma(\iota, t) := \frac{n_l g^{0l}(t)}{\mathfrak{g}(\iota, t)}, \quad X(\iota, t) := \frac{n_l X^l(t)}{\mathfrak{g}(\iota, t)}, \tag{10.1}$$

where  $n_j = \nu_{T,j}(\iota)$ ; cf. (24.5). Most of the time, it is convenient to omit the argument  $\iota$ . Sometimes we also omit reference to the argument t.

**Definition 10.1.** Consider (1.2). Assume the associated metric to be such that (M, g) is a canonical separable cosmological model manifold. For  $0 \neq \iota \in \mathcal{I}_{\mathrm{B}}$ , define  $\mathfrak{g}(\iota, t)$  by (24.13) and  $\ell$ ,  $\sigma$  and X by (10.1). If there is a constant  $C_{\mathrm{coeff}}$  such that

$$|\sigma(\iota, t)| + ||X(\iota, t)|| + ||\alpha(t)|| + ||\zeta(t)|| \le C_{\text{coeff}}$$
(10.2)

for all  $0 \neq \iota \in \mathcal{I}_{\mathrm{B}}$  and all  $t \geq 0$ , the equation (1.2) is said to be *weakly balanced*. If there is a constant  $\mathfrak{b}_{\mathrm{s}} > 0$  and a continuous non-negative function  $\mathfrak{e} \in L^{1}([0,\infty))$  such that

$$\dot{\ell}(\iota, t) \le -\mathfrak{b}_{\mathrm{s}} + \mathfrak{e}(t) \tag{10.3}$$

for all  $0 \neq \iota \in \mathcal{I}_{\mathrm{B}}$  and all  $t \geq 0$ , the equation (1.2) is said to be *weakly silent*. In case (1.2) is weakly silent,  $T_{\mathrm{ode}}$  is defined as follows. If  $c_{\mathfrak{e}} := \|\mathfrak{e}\|_1$  and  $\mathfrak{g}(\iota, 0) \leq e^{-c_{\mathfrak{e}}}$ , then  $T_{\mathrm{ode}} := 0$ . If  $\mathfrak{g}(\iota, 0) > e^{-c_{\mathfrak{e}}}$ ,  $T_{\mathrm{ode}}$  is defined as the first  $t \geq 0$  such that  $\mathfrak{g}(\iota, t) = e^{-c_{\mathfrak{e}}}$ . Finally, if there are  $\alpha_{\infty}, \zeta_{\infty} \in \mathbf{M}_m(\mathbb{C})$ and constants  $\eta_{\mathrm{mn}} > 0$  and  $C_{\mathrm{mn}}$  such that

$$\|\alpha(t) - \alpha_{\infty}\| + \|\zeta(t) - \zeta_{\infty}\| \le C_{\mathrm{mn}} e^{-\eta_{\mathrm{mn}}t}$$

$$(10.4)$$

for all  $t \ge 0$ , then (1.2) is said to be *weakly convergent*.

**Remark 10.2.** It is to be expected that the requirement that  $\mathfrak{e}$  be continuous can be avoided at the expense of additional technicalities in the proofs below.

**Remark 10.3.** If the conditions of one of Propositions 4.4, 4.11 and 4.16 are satisfied, then (1.2) is weakly balanced, weakly silent and weakly convergent. The reasons for this are the following. First of all, the norms of  $\alpha$  and  $\zeta$  are bounded due to the fact that (4.2) holds, an estimate which also yields (10.4). That (1.2) is weakly silent is a consequence of Lemma 25.12 and the assumptions of any one of the propositions mentioned at the beginning of the remark. Finally, the desired bounds on  $|\sigma(\iota, t)|$  and  $||X(\iota, t)||$  follow from Lemma 25.17.

**Remark 10.4.** Some of the estimates derived below can be improved if we, in addition to the requirements of Definition 10.1, demand that there be constants  $C_{der}$ ,  $\beta_{der} > 0$  such that

$$|\sigma(\iota, t)| + ||X(\iota, t)|| \le C_{\operatorname{der}} e^{-\beta_{\operatorname{der}} t}$$
(10.5)

for all  $0 \neq \iota \in \mathcal{I}_{\mathrm{B}}$  and all  $t \geq 0$ .

# 10.3 The oscillatory regime

For a weakly silent and balanced equation, it is, given a solution to (24.16), natural to divide  $[0, \infty)$  into the intervals  $[0, T_{ode}]$  and  $[T_{ode}, \infty)$ , where  $T_{ode}$  is given by Definition 10.1. The methods developed in Chapter 9 can be used to analyse the behaviour in the interval  $[T_{ode}, \infty)$ . In the present section, we estimate how the norm of the solution evolves in the interval  $[0, T_{ode}]$ .

**Lemma 10.5.** Assume that (1.2) is weakly balanced and weakly silent. Let  $\iota \in \mathcal{I}_B$ ,  $\mathfrak{g}$  be given by (24.13) and  $T_{ode}$  by Definition 10.1. Let z be a solution to (24.16) and define E by

$$E(t) := \frac{1}{2} [|\dot{z}(t)|^2 + \mathfrak{g}^2(t)|z(t)|^2].$$
(10.6)

Then

$$E^{1/2}(t) \le e^{c_{\mathfrak{e}} + \eta_{\text{ode}} t} E^{1/2}(0) + e^{c_{\mathfrak{e}}} \int_0^t e^{\eta_{\text{ode}}(t-s)} |\hat{f}(s)| ds$$
(10.7)

for  $t \in [0, T_{ode}]$ , where  $c_{\mathfrak{e}} := \|\mathfrak{e}\|_1$ ;  $\mathfrak{e}$  is the continuous non-negative  $L^1$ -function appearing in the definition of weak silence; and  $\eta_{ode} > 0$  is a constant depending only on  $c_{\mathfrak{e}}$  and the constant  $C_{coeff}$  appearing in the definition of weak balance.

**Remark 10.6.** The estimate is independent of the size of  $|\sigma|$ . In this respect, the assumptions can be weakened. However, in later estimates, we need control of  $\sigma$ . Moreover, the estimate (10.2) turns out to be convenient.

Proof. Estimate

$$\frac{dE}{dt} = \frac{1}{2}(\dot{z} \cdot \ddot{z}^* + \ddot{z} \cdot \dot{z}^*) + \dot{\ell}\mathfrak{g}^2|z|^2 + \frac{1}{2}\mathfrak{g}^2(z \cdot \dot{z}^* + \dot{z} \cdot z^*) \\
\leq \frac{1}{2}(\ddot{z}^* + \mathfrak{g}^2 z^*) \cdot \dot{z} + \frac{1}{2}(\ddot{z} + \mathfrak{g}^2 z) \cdot \dot{z}^* + \mathfrak{e}\mathfrak{g}^2|z|^2 \\
= \frac{1}{2}(\ddot{z}^* + 2i\sigma\mathfrak{g}\dot{z}^* + \mathfrak{g}^2 z^*) \cdot \dot{z} + \frac{1}{2}(\ddot{z} - 2i\sigma\mathfrak{g}\dot{z} + \mathfrak{g}^2 z) \cdot \dot{z}^* + \mathfrak{e}\mathfrak{g}^2|z|^2.$$
(10.8)

Combining this estimate with (24.18) yields

$$\frac{dE}{dt} \le \|\alpha\| |\dot{z}|^2 + \|X\| E + \frac{1}{2} \|\zeta\| (|\dot{z}|^2 + |z|^2) + |\dot{z}| |\hat{f}| + \mathfrak{eg}^2 |z|^2.$$
(10.9)

On the other hand,  $\mathfrak{g}(t) \geq e^{-c_{\mathfrak{e}}}$  on  $[0, T_{ode}]$ , where  $c_{\mathfrak{e}} := \|\mathfrak{e}\|_1$ . Thus (10.9) yields

$$\frac{dE}{dt} \le (C+2\mathfrak{e})E + \sqrt{2}E^{1/2}|\hat{f}|$$

for  $t \in [0, T_{ode}]$ , where the constant C only depends on  $C_{coeff}$  and  $c_{\mathfrak{e}}$ . As a consequence, it can be deduced that (10.7) holds.

For future reference, it is of interest to note that the following related estimate holds.

**Lemma 10.7.** Assume that (1.2) is weakly balanced. Assume, moreover, that there is a constant  $C_{\ell} > 0$  and a continuous non-negative function  $\mathfrak{e} \in L^1([0,\infty))$  such that

$$|\dot{\ell}(\iota,t)| \le C_{\ell} + \mathfrak{e}(t) \tag{10.10}$$

for all  $0 \neq \iota \in \mathcal{I}_B$  and all  $t \geq 0$ . Let  $\iota \in \mathcal{I}_B$  and  $\mathfrak{g}$  be given by (24.13). Let z be a solution to (24.16) and define  $\mathcal{E}$  by

$$\mathcal{E}(\iota, t) := \frac{1}{2} [|\dot{z}(\iota, t)|^2 + \mathfrak{g}^2(\iota, t)|z(\iota, t)|^2 + |z(\iota, t)|^2].$$
(10.11)

Then

$$\mathcal{E}^{1/2}(\iota, t_1) \le e^{c_{\mathfrak{e}} + \eta_{\text{bal}}|t_1 - t_0|} \mathcal{E}^{1/2}(\iota, t_0) + e^{c_{\mathfrak{e}}} \left| \int_{t_0}^{t_1} e^{\eta_{\text{bal}}|t_1 - t|} |\hat{f}(\iota, t)| dt \right|$$
(10.12)

for  $t_0, t_1 \ge 0$ , where  $c_{\mathfrak{e}} := \|\mathfrak{e}\|_1$ ; and  $\eta_{\text{bal}} > 0$  is a constant depending only on  $C_{\ell}$  and the constant  $C_{\text{coeff}}$  appearing in the definition of weak balance.

**Remark 10.8.** The reason for including the absolute value sign around the integral on the far right hand side of (10.12) is that  $t_1$  might be smaller than  $t_0$ .

*Proof.* By arguments similar to (10.8) and (10.9),

$$\left|\frac{d\mathcal{E}}{dt}\right| \leq (C+2\mathfrak{e})\mathcal{E} + \sqrt{2}|\hat{f}|\mathcal{E}^{1/2}$$

where C only depends on  $C_{\text{coeff}}$  and  $C_{\ell}$ . This estimate can be used to deduce the conclusion of the lemma.

Consider (24.18). Assume that there is a  $\mu \in \mathbb{R}$  such that for each  $\iota \in \mathcal{I}_{\mathrm{B}}$ , there is a  $T \geq 0$  with the property that solutions behave as  $e^{\mu t}$  for  $t \geq T$ . We would then like to derive an estimate corresponding to this behaviour which is valid for all  $t \geq 0$ . This can be done with the help of (10.12), but at the price of introducing a T-dependent constant.

**Corollary 10.9.** Given assumptions and notation as in Lemma 10.7, let  $\mu_l \in \mathbb{R}$ , l = 1, 2, and  $0 \leq T \in \mathbb{R}$ . Then

$$\mathcal{E}^{1/2}(\iota,t) \le e^{c_{\mathfrak{e}} + \eta_1 T} e^{\mu_1 t} \mathcal{E}^{1/2}(\iota,0) + e^{c_{\mathfrak{e}} + \eta_2 T} e^{\mu_1 t} \int_0^t e^{\mu_2(t-t')} |\hat{f}(\iota,t')| dt'$$
(10.13)

for  $t \in [0,T]$ , where  $c_{\mathfrak{e}}$  and  $\eta_{\mathrm{bal}} > 0$  have the same dependence as in the case of (10.12); and

$$\eta_1 := \max\{0, \eta_{\text{bal}} - \mu_1\}, \quad \eta_2 := \max\{0, -\mu_1\} + \max\{0, \eta_{\text{bal}} - \mu_2\}.$$

Similarly,

$$\mathcal{E}^{1/2}(\iota, T) \le e^{c_{\mathfrak{e}} + \eta_a T} e^{\mu_1 T} \mathcal{E}^{1/2}(\iota, 0) + e^{c_{\mathfrak{e}} + \eta_b T} e^{\mu_1 T} \int_0^T e^{\mu_2 (T-t)} |\hat{f}(\iota, t)| dt,$$
(10.14)

where  $c_{\mathfrak{e}}$  and  $\eta_{\text{bal}} > 0$  have the same dependence as in the case of (10.12); and

 $\eta_a := \eta_{\text{bal}} - \mu_1, \quad \eta_b := -\mu_1 + \max\{0, \eta_{\text{bal}} - \mu_2\}.$ 

**Remark 10.10.** Note that the constants  $\mu_l \in \mathbb{R}$  are arbitrary. Moreover, the estimates (10.13) and (10.14) differ in the sense that the first estimate applies in the interval [0, T], but the second estimate only applies at T. On the other hand, the second estimate is better than the estimate obtained by inserting t = T in the first estimate. The T-dependent constant should be thought of as a loss of derivatives.

*Proof.* Note that (10.12) implies that

$$\mathcal{E}^{1/2}(\iota,t) \leq e^{c_{\mathfrak{e}}+\eta_{\mathrm{bal}}t} \mathcal{E}^{1/2}(\iota,0) + e^{c_{\mathfrak{e}}} \int_0^t e^{\eta_{\mathrm{bal}}(t-t')} |\hat{f}(\iota,t')| dt'.$$

At this stage, we can estimate  $\eta_{\text{bal}}t = (\eta_{\text{bal}} - \mu_1)t + \mu_1t \leq \eta_1T + \mu_1t$  and

$$\eta_{\text{bal}}(t-t') = -\mu_1 t + \mu_1 t + (\eta_{\text{bal}} - \mu_2)(t-t') + \mu_2(t-t') \le \eta_2 T + \mu_1 t + \mu_2(t-t')$$

for  $0 \le t' \le t \le T$ . This yields (10.13) and the proof of (10.14) is similar.

### 10.4 The basic energy estimate

Assume that (1.2) is weakly silent, balanced and convergent. In the present section, we then derive energy estimates for solutions to (1.2) for  $t \ge 0$ . We do so by carrying out a mode by mode analysis, and then summing up the resulting estimates. Let, therefore, z be a solution to (24.16) corresponding to  $\iota \in \mathcal{I}_{\rm B}$ , and let  $T_{\rm ode}$  be given by Definition 10.1. In order to analyse the behaviour for  $t \ge T_{\rm ode}$ , we wish to appeal to the results derived in Section 9.2. Before doing so, note that (10.4) holds. Moreover,  $\mathfrak{g}(t) \le e^{-\mathfrak{b}_s \overline{t}}$  for  $t \ge T_{\rm ode}$ , where

$$\bar{t} := t - T_{\text{ode}},$$

so that (10.2) yields

$$n_l g^{0l}(t) | + ||n_l X^l(t)|| \le C_{\text{coeff}} e^{-\mathfrak{b}_s t}$$

for  $t \ge T_{\text{ode}}$ . When (1.2) is weakly silent, balanced and convergent, the equation (24.16) can thus be reformulated

$$\dot{v} = A_{\infty}v + A_{\rm rem}v + F,\tag{10.15}$$

where

$$v = \begin{pmatrix} z \\ \dot{z} \end{pmatrix}, \quad A_{\infty} = \begin{pmatrix} 0 & \mathrm{Id}_m \\ -\zeta_{\infty} & -\alpha_{\infty} \end{pmatrix}, \quad F = \begin{pmatrix} 0 \\ \hat{f} \end{pmatrix}.$$
 (10.16)

Moreover,  $A_{\text{rem}}$  is a smooth function taking its values in  $\mathbf{M}_{2m}(\mathbb{C})$  and satisfying

$$\|A_{\rm rem}(t)\| \le C_{\rm rem} e^{-\beta_{\rm rem}\bar{t}} \tag{10.17}$$

for  $t \geq T_{\text{ode}}$ , where  $\beta_{\text{rem}} = \min\{\mathfrak{b}_{\text{s}}, \eta_{\text{mn}}\}$  and the constant  $C_{\text{rem}}$  only depends on  $C_{\text{coeff}}$  and  $C_{\text{mn}}$ . The equation of interest here thus falls into the category described in Subsection 9.1.1. If we, in addition to the above, demand that (10.5) hold, then  $\beta_{\text{rem}} = \min\{2\mathfrak{b}_{\text{s}}, \mathfrak{b}_{\text{s}} + \beta_{\text{der}}, \eta_{\text{mn}}\}$  and  $C_{\text{rem}}$  only depends on  $C_{\text{coeff}}, C_{\text{mn}}$  and  $C_{\text{der}}$ .

Before stating the results, let us introduce the following terminology. Let  $\iota \in \mathcal{I}_{B}$ . For a solution z to (24.16) and an  $s \in \mathbb{R}$ , define

$$\mathcal{E}_{s}(\iota,t) := \frac{1}{2} \langle \nu(\iota) \rangle^{2s} [|\dot{z}(\iota,t)|^{2} + \mathfrak{g}^{2}(\iota,t)|z(\iota,t)|^{2} + |z(\iota,t)|^{2}],$$

where  $\nu(\iota)$  is given by (24.4); note that  $\nu(\iota)$  is an element of  $\mathbb{R}^{d+R}$ , and that  $\langle \nu(\iota) \rangle = [1+|\nu(\iota)|^2]^{1/2}$ . Let, also,  $\mathcal{E} := \mathcal{E}_0$ , in accordance with (10.11). If u is a solution to (1.2), z are the corresponding Fourier modes and  $s \in \mathbb{R}$ , consider (24.19):

$$\mathfrak{E}_{s}[u](t) = \frac{1}{2} \sum_{\iota \in \mathcal{I}_{B}} \langle \nu(\iota) \rangle^{2s} \left[ |\dot{z}(\iota, t)|^{2} + \mathfrak{g}^{2}(\iota, t)|z(\iota, t)|^{2} + |z(\iota, t)|^{2} \right].$$
(10.18)

Finally, in analogy with (24.10), define

$$\|f(\cdot,t)\|_{(s)} := \left(\sum_{\iota \in \mathcal{I}_{B}} \langle \nu(\iota) \rangle^{2s} |\hat{f}(\iota,t)|^{2}\right)^{1/2},$$
(10.19)

where  $\hat{f}(\iota, t)$  is given by (24.15). Let us begin by deriving two simple estimates.

**Lemma 10.11.** Assume that (1.2) is weakly silent, weakly balanced and weakly convergent. Then there are constants C and  $s_0 \ge 0$ , depending only on the coefficients of the equation (1.2), such that if z is a solution to (24.16) corresponding to  $\iota \in \mathcal{I}_{\mathrm{B}}$ , then

$$\mathcal{E}_{s}^{1/2}(\iota,t) \leq C\langle t \rangle^{d_{1}-1} e^{\kappa_{1}t} \mathcal{E}_{s+s_{0}}^{1/2}(\iota,0) + C\langle \nu(\iota) \rangle^{s+s_{0}} \int_{0}^{t} \langle t-\tau \rangle^{d_{1}-1} e^{\kappa_{1}(t-\tau)} |\hat{f}(\iota,\tau)| d\tau$$

$$(10.20)$$

for all  $t \ge 0$ , where  $\kappa_1 := \kappa_{\max}(A_{\infty})$ ,  $d_1 := d_{\max}(A_{\infty}, \kappa_1)$ , the matrix  $A_{\infty}$  is defined in (10.16) and  $\hat{f}(\iota, t)$  is given by (24.15). Moreover, if  $\iota$  is a smooth solution to (1.2), then

$$\mathfrak{E}_{s}^{1/2}[u](t) \leq C\langle t \rangle^{d_{1}-1} e^{\kappa_{1}t} \mathfrak{E}_{s+s_{0}}^{1/2}[u](0) + C \int_{0}^{t} \langle t-\tau \rangle^{d_{1}-1} e^{\kappa_{1}(t-\tau)} \|f(\cdot,\tau)\|_{(s+s_{0})} d\tau$$
(10.21)

for all  $t \geq 0$  and  $s \in \mathbb{R}$ .

**Remark 10.12.** The functions  $\kappa_{\text{max}}$  and  $d_{\text{max}}$  are introduced in Definition 4.3.

**Remark 10.13.** The constant C appearing in the statement of the lemma only depends on  $\mathfrak{b}_s$ ,  $\eta_{\rm mn}$ ,  $C_{\rm coeff}$ ,  $C_{\rm mn}$ ,  $c_{\mathfrak{e}}$ ,  $A_{\infty}$ ,  $g^{ij}(0)$  and  $a_r(0)$ , where  $i, j = 1, \ldots, d$  and  $r = 0, \ldots, R$ . Moreover,  $s_0$  only depends on  $c_{\mathfrak{e}}$ ,  $C_{\rm coeff}$ ,  $\kappa_1$  and  $\mathfrak{b}_s$ .

**Remark 10.14.** If the assumptions of Proposition 4.4 are satisfied, then the assumptions of the present lemma are fulfilled; cf. Remark 10.3.

#### 10.4. THE BASIC ENERGY ESTIMATE

Proof. In the proof of (10.20), there are two cases to consider:  $T_{ode} = 0$  and  $T_{ode} > 0$ . In the first case, we need only appeal to Lemma 9.14; since  $T_{ode} = 0$  implies  $\mathfrak{g}(t) \leq 1$  for all  $t \geq 0$ , the expressions |v| and  $\mathcal{E}^{1/2}$  are equivalent in this case. Assume that  $T_{ode} > 0$ . In the interval  $[0, T_{ode}]$ , we can appeal to Lemma 10.5 and for  $t \geq T_{ode}$ , we can appeal to the combination of Lemmas 9.14 and 10.5. When  $T_{ode} > 0$ ,  $\mathcal{E}(T_{ode})$  and  $\mathcal{E}(T_{ode})$  are equivalent (with a constant of equivalence depending only on the constant  $c_{\mathfrak{e}}$  introduced in Lemma 10.5). Combining Lemmas 9.14 and 10.5 thus yields

$$\begin{split} \mathcal{E}^{1/2}(t) \leq & C \langle \bar{t} \rangle^{d_1 - 1} e^{\kappa_1 \bar{t}} E^{1/2}(T_{\text{ode}}) + C \int_{T_{\text{ode}}}^t \langle t - \tau \rangle^{d_1 - 1} e^{\kappa_1 (t - \tau)} |\hat{f}(\tau)| d\tau \\ \leq & C \langle t \rangle^{d_1 - 1} e^{\kappa_1 t} e^{\eta + T_{\text{ode}}} E^{1/2}(0) + C e^{\eta + T_{\text{ode}}} \int_0^{T_{\text{ode}}} \langle t - \tau \rangle^{d_1 - 1} e^{\kappa_1 (t - \tau)} |\hat{f}(\tau)| d\tau \\ & + C \int_{T_{\text{ode}}}^t \langle t - \tau \rangle^{d_1 - 1} e^{\kappa_1 (t - \tau)} |\hat{f}(\tau)| d\tau \end{split}$$

for  $t \geq T_{ode}$ , where  $\eta_{+} = \max\{\eta_{ode} - \kappa_1, 0\}$ . In particular,

$$\mathcal{E}^{1/2}(t) \le C \langle t \rangle^{d_1 - 1} e^{\kappa_1 t} e^{\eta_+ T_{\text{ode}}} \mathcal{E}^{1/2}(0) + C e^{\eta_+ T_{\text{ode}}} \int_0^t \langle t - \tau \rangle^{d_1 - 1} e^{\kappa_1 (t - \tau)} |\hat{f}(\tau)| d\tau$$
(10.22)

for  $t \ge T_{\text{ode}}$ . Due to (10.7), an even better estimate holds for  $t \in [0, T_{\text{ode}}]$ . In particular, (10.22) thus holds for all  $t \ge 0$ . However, in order for this estimate to be of any interest, we need to estimate  $e^{\eta_+ T_{\text{ode}}}$ . Note, to this end, that (10.3) and the definition of  $T_{\text{ode}}$  yield

$$e^{-c_{\mathfrak{e}}} = \mathfrak{g}(T_{\text{ode}}) \le \mathfrak{g}(0)e^{-\mathfrak{b}_{s}T_{\text{ode}}+c_{\mathfrak{e}}}.$$
(10.23)

Thus  $T_{\text{ode}} \leq \mathfrak{b}_{s}^{-1} \ln \mathfrak{g}(0) + 2\mathfrak{b}_{s}^{-1}c_{\mathfrak{e}}$ . In particular,

$$e^{\eta_+ T_{\text{ode}}} \le C \exp\left[\eta_+ \mathfrak{b}_{\text{s}}^{-1} \ln \mathfrak{g}(0)\right] \le C |\nu(\iota)|^{\eta_+/\mathfrak{b}_{\text{s}}},$$

where C only depends on  $g^{ij}(0)$ ,  $a_r(0)$ ,  $\mathfrak{b}_s$ ,  $c_{\mathfrak{c}}$ ,  $C_{\text{coeff}}$  and  $A_{\infty}$ . Thus, letting  $s_0 := \eta_+/\mathfrak{b}_s$ , the fact that (10.22) holds for all  $t \ge 0$  implies that (10.20) holds. Applying Minkowski's inequality to this estimate yields (10.21).

For future reference, it is of interest to note that the following related estimate holds.

**Lemma 10.15.** Assume that (1.2) is weakly balanced. Assume, moreover, that there is a constant  $C_{\ell} > 0$  and a continuous non-negative function  $\mathfrak{e} \in L^1([0,\infty))$  such that (10.10) holds for all  $0 \neq \iota \in \mathcal{I}_B$  and all  $t \geq 0$ . Let u be a solution to (1.2). Then

$$\mathfrak{E}_{s}^{1/2}[u](t_{1}) \leq e^{c_{\mathfrak{e}} + \eta_{\mathrm{bal}}|t_{1} - t_{0}|} \mathfrak{E}_{s}^{1/2}[u](t_{0}) + e^{c_{\mathfrak{e}}} \left| \int_{t_{0}}^{t_{1}} e^{\eta_{\mathrm{bal}}|t_{1} - t|} \|f(\cdot, t)\|_{(s)} dt \right|$$
(10.24)

for  $t_0, t_1 \ge 0$  and  $s \in \mathbb{R}$ , where  $c_{\mathfrak{e}} := \|\mathfrak{e}\|_1$ ; and  $\eta_{\text{bal}} > 0$  is a constant depending only on  $C_{\ell}$  and the constant  $C_{\text{coeff}}$  appearing in the definition of weak balance.

**Remark 10.16.** Note that if (1.2) is  $C^1$ -balanced in the sense of Definition 3.8, then it satisfies the conditions of the lemma with  $\mathfrak{e} = 0$ ; this is a consequence of Lemmas 25.11 and 25.17. In fact, the conditions of the lemma also hold if we relax the assumptions in Definition 3.8 by replacing the conditions that

- the shift vector field be  $C^1$ -future bounded, and
- $|\bar{k}|_{\bar{g}} \leq C_{\bar{k}}$  for  $t \geq 0$

by the requirements that  $\partial_t$  be future uniformly timelike and

$$|k|_{\bar{g}} + |\chi|_{\bar{g}} \cdot |\dot{\chi}|_{\bar{g}} \le C_a + \mathfrak{e}(t)$$

for all  $t \ge 0$ , where  $0 < C_a \in \mathbb{R}$  and  $\mathfrak{e} \in L^1([0,\infty))$  is continuous and non-negative. This is a consequence of Lemmas 25.15 and 25.17.

*Proof.* The conclusion follows from (10.12) and Minkowski's inequality.

# 10.5 Future asymptotics

As a next step, we wish to derive asymptotics by appealing to Lemma 9.16; note that the equation (10.15) is of a form such that this lemma is applicable, assuming f to satisfy a suitable integrability condition.

**Lemma 10.17.** Assume that (1.2) is weakly silent, weakly balanced and weakly convergent. Assume, moreover, that f is a smooth function such that for every  $s \in \mathbb{R}$ ,

$$||f||_{A,s} := \int_0^\infty e^{-\kappa_1 \tau} ||f(\cdot, \tau)||_{(s)} d\tau < \infty$$
(10.25)

holds, where  $\kappa_1 := \kappa_{\max}(A_{\infty})$  and  $A_{\infty}$  is defined in (10.16); cf. Definition 4.3. Let  $\beta_{\text{rem}} := \min\{\mathfrak{b}_s, \eta_{\text{mn}}\}$ , where  $\mathfrak{b}_s$  and  $\eta_{\text{mn}}$  are the constants appearing in the definition of weak silence and weak convergence; cf. Definition 10.1. Let, moreover,  $E_a$  be the first generalised eigenspace in the  $\beta_{\text{rem}}, A_{\infty}$ -decomposition of  $\mathbb{C}^{2m}$ ; cf. Definition 9.4. Then there are constants C, N and  $s_{\text{hom}}, s_{\text{ih}} \geq 0$ , depending only on the coefficients of the equation (1.2), such that the following holds. Given a smooth solution u to (1.2), there is a unique  $V_{\infty} \in C^{\infty}(\overline{M}, E_a)$  such that

$$\left\| \begin{pmatrix} u(\cdot,t) \\ u_t(\cdot,t) \end{pmatrix} - e^{A_{\infty}t} V_{\infty} - \int_0^t e^{A_{\infty}(t-\tau)} \begin{pmatrix} 0 \\ f(\cdot,\tau) \end{pmatrix} d\tau \right\|_{(s)}$$

$$\leq C \langle t \rangle^N e^{(\kappa_1 - \beta_{\text{rem}})t} \left( \| u_t(\cdot,0) \|_{(s+s_{\text{hom}})} + \| u(\cdot,0) \|_{(s+s_{\text{hom}}+1)} + \| f \|_{A,s+s_{\text{ih}}} \right)$$

$$(10.26)$$

holds for  $t \geq 0$  and all  $s \in \mathbb{R}$ ; recall that  $\overline{M}$  is given by (1.3). Moreover,

$$\|V_{\infty}\|_{(s)} \le C\left(\|u_t(\cdot,0)\|_{(s+s_{\text{hom}})} + \|u(\cdot,0)\|_{(s+s_{\text{hom}}+1)} + \|f\|_{A,s+s_{\text{ih}}}\right).$$
(10.27)

**Remark 10.18.** If the assumptions of Proposition 4.11 are satisfied, then the assumptions of the present lemma are fulfilled; cf. Remark 10.3. Note also that the fact that the  $\beta_{\text{rem}}$  appearing in the statement of the present lemma coincides with the  $\beta_{\text{rem}}$  appearing in the statement of Proposition 4.11 is a consequence of the fact that, in the context of Proposition 4.11,  $\mathfrak{b}_{s} = \mu$ ; cf. (25.52).

**Remark 10.19.** Recall that the notation  $C_{\text{coeff}}$ ,  $C_{\text{mn}}$ ,  $\mathfrak{b}_{s}$ ,  $\eta_{\text{mn}}$  and  $\mathfrak{e}$  is introduced in Definition 10.1. Let  $c_{\mathfrak{e}} := \|\mathfrak{e}\|_{1}$ . Then the constant C only depends on  $c_{\mathfrak{e}}$ ,  $C_{\text{coeff}}$ ,  $C_{\text{mn}}$ ,  $\mathfrak{b}_{s}$ ,  $\eta_{\text{mn}}$ ,  $\eta_{\text{ode}}$ ,  $A_{\infty}$ ,  $g^{ij}(0)$  and  $a_{r}(0)$ , where  $\eta_{\text{ode}}$  is the constant appearing in Lemma 10.5. The constant N only depends on m. Finally,  $s_{\text{hom}} = (\beta_{\text{rem}} + \eta_{+})/\mathfrak{b}_{s}$  where  $\eta_{+} := \max\{\eta_{\text{ode}} - \kappa_{1}, 0\}$ ; and  $s_{\text{ih}} := \max\{\beta_{\text{rem}} + \eta_{+}, \kappa + 1\}/\mathfrak{b}_{s}$ , where  $\kappa := \text{Rsp}A_{\infty}$  and the notation  $\text{Rsp}A_{\infty}$  is introduced in Definition 4.3.

**Remark 10.20.** If, in addition to the assumptions of the lemma, the estimate (10.5) holds, then  $\beta_{\text{rem}}$  can be replaced by  $\min\{2\mathfrak{b}_s, \mathfrak{b}_s + \beta_{\text{der}}, \eta_{\text{mn}}\}$ ; cf. the comments made in connection with (10.17). The dependence of the constants then changes in that they also depend on  $C_{\text{der}}$  and  $\beta_{\text{der}}$ .

**Remark 10.21.** If the assumptions of Proposition 4.11 are satisfied and there are constants  $K_{\text{der}}$  and  $\beta_{\text{der}}$  such that (4.12) holds for all  $t \ge 0$ , then the conclusions of Lemma 10.17 hold with  $\beta_{\text{rem}}$  replaced by  $\min\{2\mu, \mu + \beta_{\text{der}}, \eta_{\text{mn}}\}$ . The reason for this is that (4.12) and Lemma 25.17 imply that (10.5) holds so that Remark 10.20 applies.

*Proof.* Uniqueness. The proof of uniqueness is essentially identical to the argument presented at the beginning of the proof of Lemma 9.16. We therefore omit the details here.

**Strategy, existence.** In order to prove the statement, we derive estimates for each mode and then sum up the result. For a given  $\iota \in \mathcal{I}_{\mathrm{B}}$ , it is natural to divide the interval  $[0, \infty)$  into two parts:  $[0, T_{\mathrm{ode}}]$  and  $[T_{\mathrm{ode}}, \infty)$ . On the latter interval, we can appeal to Lemma 9.16. However, we have no similar estimate on the interval  $[0, T_{\mathrm{ode}}]$ . In fact, there is no reason to expect an expression such as the left hand side of (9.22) to be small on  $[0, T_{\mathrm{ode}}]$ . The way to overcome this problem is to

- observe that the solution cannot grow more than exponentially on  $[0, T_{ode}]$ ; cf. Lemma 10.5;
- note that this exponential growth cannot exceed a constant multiple of  $\langle \nu(\iota) \rangle^{\kappa}$  for some  $\kappa$ , so that the growth can be reinterpreted in terms of a loss of regularity.

**Preliminary observations; the objects that need to be estimated.** Consider (9.22). We wish to reformulate this estimate to one in which  $T_{ode}$  does not appear, and we wish to prove that it is valid for all  $t \ge 0$  (note that  $v, v_{\infty}$  and F appearing in this estimate are all dependent on  $\iota$  even though this is not explicitly stated). Introducing

$$v_{\infty,\text{mod}} := v_{\infty} - \int_0^{T_{\text{ode}}} e^{-A_{\infty}\tau} F(\tau) d\tau, \qquad (10.28)$$

(9.22) can be written

$$\left| v(t) - e^{A_{\infty}t} v_{\infty, \text{mod}} - \int_{0}^{t} e^{A_{\infty}(t-\tau)} F(\tau) d\tau \right|$$

$$\leq C \langle t \rangle^{N} e^{(\kappa_{1}-\beta_{\text{rem}})t} e^{\beta_{\text{rem}}T_{\text{ode}}} \left[ e^{-\kappa_{1}T_{\text{ode}}} |v(T_{\text{ode}})| + \|F\|_{A} \right];$$

$$(10.29)$$

recall that this estimate holds on  $[T_{ode}, \infty)$  and that the notation  $||F||_A$  is introduced in (9.4). Note also that the constant C only depends on  $C_{rem}$ ,  $\beta_{rem}$  and  $A_{\infty}$ ; due to the comments made in connection with (10.17), this means that the constant only depends on  $C_{coeff}$ ,  $C_{mn}$ ,  $\mathfrak{b}_s$ ,  $\eta_{mn}$  and  $A_{\infty}$ . In accordance with the above strategy, we wish to estimate each term appearing on the left hand side separately on the interval  $[0, T_{ode}]$ . Moreover, we wish to estimate the right hand side in terms of initial data and  $\iota$ , but we do not wish to have any dependence on  $|v(T_{ode})|$  or  $T_{ode}$ . In practice, it turns out to be convenient to estimate

$$e^{(\beta_{\rm rem}-\kappa_1)t}|v(t)|, \quad e^{(\beta_{\rm rem}-\kappa_1)t}|e^{A_{\infty}t}v_{\infty}|, \quad \left|e^{(\beta_{\rm rem}-\kappa_1)t}\int_t^{T_{\rm ode}}e^{A_{\infty}(t-\tau)}F(\tau)d\tau\right| \tag{10.30}$$

for  $t \in [0, T_{ode}]$ ; this yields the desired estimate on the interval  $[0, T_{ode}]$ . In order to obtain a good estimate for  $t \ge T_{ode}$ , we need to estimate

$$e^{(\beta_{\rm rem} - \kappa_1)T_{\rm ode}} |v(T_{\rm ode})|, \quad e^{\beta_{\rm rem}T_{\rm ode}} ||F||_A.$$
 (10.31)

Finally, we wish to estimate  $v_{\infty,\text{mod}}$  introduced in (10.28). Note that if we have estimated the first expression appearing in (10.30) on  $[0, T_{\text{ode}}]$ , then we have also estimated the first expression appearing in (10.31).

Estimating the size of the solution. Assume that  $T_{ode} > 0$ . Then we can appeal to Lemma 10.5 in order to obtain

$$e^{(\beta_{\rm rem}-\kappa_1)t}|v(t)| \le \sqrt{2}e^{2c_{\mathfrak{e}}}e^{\eta_{\rm hom,1}t}E^{1/2}(0) + \sqrt{2}e^{2c_{\mathfrak{e}}}e^{\eta_{\rm ih,1}t}\int_0^t e^{-\kappa_1\tau}|\hat{f}(\tau)|d\tau|^{1/2}$$

for  $t \in [0, T_{\text{ode}}]$ , where  $\eta_{\text{hom},1} := \max\{\beta_{\text{rem}} + \eta_{\text{ode}} - \kappa_1, 0\}$ ,  $\eta_{\text{ih},1} := \beta_{\text{rem}} + \max\{\eta_{\text{ode}} - \kappa_1, 0\}$ ,  $\mathfrak{e}$  is the function appearing in the definition of weak silence and  $c_{\mathfrak{e}} := \|\mathfrak{e}\|_1$  (for future reference, it is

convenient to keep in mind that the same estimate holds with  $\beta_{\text{rem}}$  replaced by 0). Due to (10.23), it then follows that

$$\langle \nu(\iota) \rangle^{s} \sup_{t \in [0, T_{\text{ode}}]} e^{(\beta_{\text{rem}} - \kappa_{1})t} |\nu(t)|$$

$$\leq C \mathcal{E}_{s+s_{\text{hom},1}}^{1/2}(0) + C \langle \nu(\iota) \rangle^{s+s_{\text{ih},1}} \int_{0}^{T_{\text{ode}}} e^{-\kappa_{1}\tau} |\hat{f}(\tau)| d\tau,$$

$$(10.32)$$

where  $s_{\text{hom},1} := \eta_{\text{hom},1}/\mathfrak{b}_s$ ,  $s_{\text{ih},1} := \eta_{\text{ih},1}/\mathfrak{b}_s$  and the constant C only depends on  $c_{\mathfrak{e}}$ ,  $\mathfrak{b}_s$ ,  $\eta_{\text{mn}}$ ,  $\eta_{\text{ode}}$ ,  $\kappa_1$ ,  $g^{ij}(0)$  and  $a_r(0)$  (and, again, the analogous estimate with  $\beta_{\text{rem}}$  replaced by 0 holds). One particular consequence of this estimate is

$$\langle \nu(\iota) \rangle^{s} e^{(\beta_{\rm rem} - \kappa_{1})T_{\rm ode}} |\nu(T_{\rm ode})| + \langle \nu(\iota) \rangle^{s} e^{\beta_{\rm rem}T_{\rm ode}} \|F\|_{A}$$

$$\leq C \mathcal{E}_{s+s_{\rm hom,1}}^{1/2}(0) + C \langle \nu(\iota) \rangle^{s+s_{\rm ih,1}} \|\hat{f}\|_{A},$$

$$(10.33)$$

where the constant C has the same dependence as in the case of (10.32) and we have used the notation (9.4). Moreover, the analogous estimate with  $\beta_{\text{rem}}$  replaced by 0 holds. Clearly, there are various ways in which (10.33) could be improved in terms of the power of  $\langle \nu(\iota) \rangle$ . However, we do not aim for an optimal estimate in that regard here.

Estimating the limit. Let us consider the vector  $v_{\infty}$  appearing in (9.22) in greater detail. To begin with, it is given by (9.23), where  $u_{\infty}$  satisfies the estimate (9.24), in which the constant C only depends on  $C_{\text{coeff}}$ ,  $C_{\text{mn}}$ ,  $\mathfrak{b}_{s}$ ,  $\eta_{\text{mn}}$  and  $A_{\infty}$ . Note that due to (9.24) and estimates of the above type (in fact, the estimate (10.33) where  $\beta_{\text{rem}}$  has been set to zero),

$$\langle \nu(\iota) \rangle^s |u_{\infty}| \le C e^{\kappa_1 T_{\text{ode}}} \mathcal{E}_{s+s_0}^{1/2}(0) + C e^{\kappa_1 T_{\text{ode}}} \langle \nu(\iota) \rangle^{s+s_0} \|\hat{f}\|_A,$$
(10.34)

where  $s_0 := \eta_+/\mathfrak{b}_s$ ,  $\eta_+ := \max\{\eta_{ode} - \kappa_1, 0\}$  and C only depends on  $c_{\mathfrak{e}}$ ,  $C_{coeff}$ ,  $C_{mn}$ ,  $\mathfrak{b}_s$ ,  $\eta_{mn}$ ,  $\eta_{ode}$ ,  $A_{\infty}$ ,  $g^{ij}(0)$  and  $a_r(0)$  (since we appeal to (9.24), the constant depends on  $C_{rem}$ ). Since  $v_{\infty}$  is given by (9.23) and since  $u_{\infty} \in E_{A_{\infty},\beta_{rem}}$ , it follows that

$$\langle \nu(\iota) \rangle^s |v_{\infty}| \le C \mathcal{E}_{s+s_{\text{hom},2}}^{1/2}(0) + C \langle \nu(\iota) \rangle^{s+s_{\text{ih},2}} \|\hat{f}\|_A,$$
(10.35)

where  $s_{\text{hom},2} := s_{\text{ih},1}$ ,  $s_{\text{ih},2} := s_{\text{ih},1}$  and the constant C has the same dependence as in the case of (10.34). In order to estimate the second expression in (10.30), note that (9.23) yields

$$e^{(\beta_{\rm rem}-\kappa_1)t}e^{A_{\infty}t}v_{\infty} = e^{(\beta_{\rm rem}-\kappa_1)t}e^{A_{\infty}\bar{t}}u_{\infty}.$$
(10.36)

This equation, the fact that  $u_{\infty} \in E_{A_{\infty},\beta_{\text{rem}}}$  and the fact that  $t \leq T_{\text{ode}}$  lead to the conclusion that

$$|e^{(\beta_{\rm rem}-\kappa_1)t}e^{A_{\infty}t}v_{\infty}| \le Ce^{(\beta_{\rm rem}-\kappa_1)T_{\rm ode}}|u_{\infty}|$$

for  $t \in [0, T_{ode}]$ , where C only depends on  $\beta_{rem}$  and  $A_{\infty}$ . Combining this estimate with (10.34) yields

$$\langle \nu(\iota) \rangle^{s} \sup_{t \in [0, T_{\text{ode}}]} |e^{(\beta_{\text{rem}} - \kappa_{1})t} e^{A_{\infty}t} v_{\infty}| \le C \mathcal{E}_{s+s_{\text{hom},2}}^{1/2}(0) + C \langle \nu(\iota) \rangle^{s+s_{\text{ih},2}} \|\hat{f}\|_{A},$$
(10.37)

where the constant C has the same dependence as in the case of (10.34).

Estimating the contribution of the right hand side of (1.2). Let us turn to

$$e^{(\beta_{\rm rem}-\kappa_1)t} \int_t^{T_{\rm ode}} e^{A_{\infty}(t-\tau)} F(\tau) d\tau$$

$$= \int_t^{T_{\rm ode}} e^{\beta_{\rm rem}\tau} e^{(A_{\infty}-\kappa_1 {\rm Id}_{2m}+\beta_{\rm rem} {\rm Id}_{2m})(t-\tau)} e^{-\kappa_1 \tau} F(\tau) d\tau.$$
(10.38)

The situation is similar to (10.36). However, there is a fundamental difference in that we do not know that F takes its values in  $E_{A_{\infty},\beta_{\text{rem}}}$ . In order to estimate the right hand side of (10.38), it is convenient to consider the cases  $\beta_{\text{rem}} - \kappa > 0$  and  $\beta_{\text{rem}} - \kappa \le 0$  separately, where  $\kappa := \text{Rsp}A_{\infty}$ and the notation  $\text{Rsp}A_{\infty}$  is introduced in Definition 4.3. In the second case, there is potentially a polynomial loss, which we estimate crudely by an exponentially growing factor. This yields

$$\langle \nu(\iota) \rangle^s \sup_{t \in [0, T_{\text{ode}}]} \left| e^{(\beta_{\text{rem}} - \kappa_1)t} \int_t^{T_{\text{ode}}} e^{A_\infty(t-\tau)} F(\tau) d\tau \right| \le C \langle \nu(\iota) \rangle^{s+s_{\text{ih},3}} \|\hat{f}\|_A,$$
(10.39)

where  $s_{ih,3} = \max\{\beta_{rem}, \kappa+1\}/\mathfrak{b}_s$  and C only depends on  $\mathfrak{b}_s, \eta_{mn}, c_{\mathfrak{e}}, A_{\infty}, g^{ij}(0)$  and  $a_r(0)$ .

**Combining the estimates.** Since the second term on the right hand side of (10.28) (multiplied by  $\langle \nu(\iota) \rangle^s$ ) can be estimated by the left hand side of (10.39), we have all the estimates we need. To begin with, (10.28), (10.35) and (10.39) yield

$$\langle \nu(\iota) \rangle^s |v_{\infty,\text{mod}}| \le C \mathcal{E}_{s+s_{\infty,\text{hom}}}^{1/2}(0) + C \langle \nu(\iota) \rangle^{s+s_{\infty,\text{ih}}} \|\hat{f}\|_A,$$
(10.40)

where  $s_{\infty,\text{hom}} := (\beta_{\text{rem}} + \eta_+)/\mathfrak{b}_s$ ,  $s_{\infty,\text{ih}} := \max\{\beta_{\text{rem}} + \eta_+, \kappa + 1\}/\mathfrak{b}_s$  and C has the same dependence as in the case of (10.34). Moreover, combining (10.32), (10.37) and (10.39) yields

$$\langle \nu(\iota) \rangle^{s} \left| v(t) - e^{A_{\infty}t} v_{\infty, \text{mod}} - \int_{0}^{t} e^{A_{\infty}(t-\tau)} F(\tau) d\tau \right|$$

$$\leq C e^{(\kappa_{1} - \beta_{\text{rem}})t} \left( \mathcal{E}_{s+s_{\text{hom},4}}^{1/2}(0) + \langle \nu(\iota) \rangle^{s+s_{\text{ih},4}} \| \hat{f} \|_{A} \right)$$

$$(10.41)$$

for  $0 \le t \le T_{\text{ode}}$ , where  $s_{\text{hom},4} := (\beta_{\text{rem}} + \eta_+)/\mathfrak{b}_s$ ,  $s_{\text{ih},4} := \max\{\beta_{\text{rem}} + \eta_+, \kappa + 1\}/\mathfrak{b}_s$  and C has the same dependence as in the case of (10.34). Combining (10.29) and (10.33) yields an estimate similar to (10.41) which is valid for  $t \ge T_{\text{ode}}$ . Combining the two yields

$$\langle \nu(\iota) \rangle^{s} \left| v(t) - e^{A_{\infty}t} v_{\infty, \text{mod}} - \int_{0}^{t} e^{A_{\infty}(t-\tau)} F(\tau) d\tau \right|$$

$$\leq C \langle t \rangle^{N} e^{(\kappa_{1} - \beta_{\text{rem}})t} \left( \mathcal{E}_{s+s_{\text{hom}}}^{1/2}(0) + \langle \nu(\iota) \rangle^{s+s_{\text{ih}}} \| \hat{f} \|_{A} \right)$$

$$(10.42)$$

for  $t \ge 0$ , where  $s_{\text{hom}} := (\beta_{\text{rem}} + \eta_+)/\mathfrak{b}_s$ ;  $s_{\text{ih}} := \max\{\beta_{\text{rem}} + \eta_+, \kappa + 1\}/\mathfrak{b}_s$ ; *C* has the same dependence as in the case of (10.34); and *N* is a non-negative integer depending only on *m*. Moreover, in case  $T_{\text{ode}} = 0$ , (10.42) is an immediate consequence of (9.22).

**Projecting the data at infinity.** At this stage, we would like to sum up the estimate (10.42) over the different modes in order to obtain (10.26). Comparing (10.42) and (10.26), it seems reasonable to choose  $V_{\infty}$  so that its *i*'th mode is given by  $v_{\infty, \text{mod}}(\iota)$ . However, there is one problem with doing so: we do not know that  $v_{\infty, \text{mod}}(\iota)$  belongs to  $E_a$ , since it is not clear that the second term on the right hand side of (10.28) belongs to this space. In fact, we first need to project  $v_{\infty, \text{mod}}(\iota)$ to  $E_a$ . This leaves an error term we need to estimate. It is given by

$$e^{A_{\infty}t}\Pi_b\left(\int_0^{T_{\text{ode}}} e^{-A_{\infty}\tau}F(\tau)d\tau\right),\tag{10.43}$$

where  $\Pi_b$  denotes the projection to the second generalised eigenspace in the  $\beta_{\text{rem}}$ ,  $A_{\infty}$ -decomposition of  $\mathbb{C}^{2m}$ . Appealing to Lemma 9.6 and (10.39) yields

$$\left| e^{A_{\infty}t} \Pi_{b} \left( \int_{0}^{T_{\text{ode}}} e^{-A_{\infty}\tau} F(\tau) d\tau \right) \right| \leq C_{A} \langle t \rangle^{d_{b}-1} e^{\kappa_{b}t} \left| \int_{0}^{T_{\text{ode}}} e^{-A_{\infty}\tau} F(\tau) d\tau \right|$$
$$\leq C \langle t \rangle^{d_{b}-1} e^{\kappa_{b}t} \langle \nu(\iota) \rangle^{s_{\text{in},3}} \|\hat{f}\|_{A},$$

where  $d_b$  and  $\kappa_b$  are given in the statement of Lemma 9.6 and C has the same dependence as in the case of (10.39). Since  $\kappa_b \leq \kappa_1 - \beta_{\text{rem}}$ , this estimate yields

$$\langle \nu(\iota) \rangle^s | e^{A_\infty t} \Pi_b v_{\infty, \text{mod}} | \le C \langle t \rangle^N e^{(\kappa_1 - \beta_{\text{rem}})t} \langle \nu(\iota) \rangle^{s+s_{\text{ih}}} \| \hat{f} \|_A,$$

where C and N have the same dependence as in the case of (10.42). Thus (10.42) holds with  $v_{\infty,\text{mod}}$  replaced by  $\prod_a v_{\infty,\text{mod}}$ .

Summing up. Given  $p = (x, p_1, \ldots, p_R) \in \overline{M}$ , where  $\overline{M}$  is given by (1.3), define  $V_{\infty}(p)$  by

$$V_{\infty}(p) := \sum_{\iota \in \mathcal{I}_{\mathrm{B}}} \prod_{a} [v_{\infty, \mathrm{mod}}(\iota)] \varphi_{\iota}(p),$$

where  $\varphi_{\iota}$  is given by (24.6). In order to justify that the sum on the right hand side makes sense and yields a smooth function, it is sufficient to appeal to the fact that u is smooth; the assumption that (10.25) holds for all s; (10.40); and the Minkowski inequality. Then (10.42) (with  $v_{\infty,\text{mod}}$ replaced by  $\Pi_a v_{\infty,\text{mod}}$ ) implies that

$$\left\| \begin{pmatrix} u(\cdot,t) \\ u_t(\cdot,t) \end{pmatrix} - e^{A_{\infty}t}V_{\infty} - \int_0^t e^{A_{\infty}(t-\tau)} \begin{pmatrix} 0 \\ f(\cdot,\tau) \end{pmatrix} d\tau \right\|_{(s)}$$
  
 
$$\leq C \langle t \rangle^N e^{(\kappa_1 - \beta_{\text{rem}})t} \left( \mathfrak{E}_{s+s_{\text{hom}}}^{1/2} [u](0) + \|f\|_{A,s+s_{\text{ih}}} \right)$$

for  $t \ge 0$ , where we have appealed to the Minkowski inequality and C has the same dependence as in the case of (10.34). Thus (10.26) holds. Note also that, due to (10.40),

$$\|V_{\infty}\|_{(s)} \le C \left(\mathfrak{E}_{s+s_{\text{hom}}}^{1/2}[u](0) + \|f\|_{A,s+s_{\text{ih}}}\right)$$

where C has the same dependence as in the case of (10.34). Thus (10.27) holds.

# **10.6** Specifying the asymptotics

As a next step, it is of interest to go in the direction opposite to that of Lemma 10.17; i.e., to specify  $V_{\infty}$  and to find a solution such that (10.26) holds. Since it is sufficient to consider the case of homogeneous equations, let us assume that f = 0.

**Lemma 10.22.** Assume that (1.2) is weakly balanced, weakly silent and weakly convergent. Assume, moreover, that f = 0 and that there is a constant  $\mathfrak{b}_{low} > 0$  and a non-negative continuous function  $\mathfrak{e}_{low} \in L^1([0,\infty))$  such that

$$\ell(\iota, t) \ge -\mathfrak{b}_{\text{low}} - \mathfrak{e}_{\text{low}}(t) \tag{10.44}$$

for all  $t \ge 0$  and all  $0 \ne \iota \in \mathcal{I}_{\mathrm{B}}$ ; recall that  $\ell$  is defined in (10.1). Let  $A_{\infty}$  be defined by (10.16) and let  $\beta_{\mathrm{rem}} = \min\{\eta_{\mathrm{mn}}, \mathfrak{b}_{\mathrm{s}}\}$ , where  $\mathfrak{b}_{\mathrm{s}}$  and  $\eta_{\mathrm{mn}}$  are the constants appearing in (10.3) and (10.4) respectively. Finally, let  $E_a$  be the first generalised eigenspace in the  $\beta_{\mathrm{rem}}, A_{\infty}$ -decomposition of  $\mathbb{C}^{2m}$ ; cf. Definition 9.4. Then there is an injective map

$$\Phi_{\infty}: C^{\infty}(\bar{M}, E_a) \to C^{\infty}(\bar{M}, \mathbb{C}^{2m})$$

such that the following holds. First,

$$\|\Phi_{\infty}(\chi)\|_{(s)} \le C \|\chi\|_{(s+s_{\infty})} \tag{10.45}$$

for all  $s \in \mathbb{R}$  and all  $\chi \in C^{\infty}(\overline{M}, E_a)$ , where the constants C and  $s_{\infty} \geq 0$  only depend on  $C_{\text{coeff}}$ ,  $C_{\text{mn}}$ ,  $\mathfrak{b}_s$ ,  $\mathfrak{b}_{\text{low}}$ ,  $\eta_{\text{mn}}$ ,  $A_{\infty}$ ,  $\|\mathfrak{e}\|_1$ ,  $\|\mathfrak{e}_{\text{low}}\|_1$ ,  $g^{ij}(0)$ ,  $i, j = 1, \ldots, d$ , and  $a_r(0)$ ,  $r = 1, \ldots, R$ . Secondly, if  $\chi \in C^{\infty}(\overline{M}, E_a)$  and u is the solution to (1.2) (with f = 0) such that

$$\begin{pmatrix} u(\cdot,0)\\ u_t(\cdot,0) \end{pmatrix} = \Phi_{\infty}(\chi), \tag{10.46}$$

then

$$\left\| \begin{pmatrix} u(\cdot,t) \\ u_t(\cdot,t) \end{pmatrix} - e^{A_{\infty}t} \chi \right\|_{(s)}$$

$$\leq C \langle t \rangle^N e^{(\kappa_1 - \beta_{\text{rem}})t} \left( \| u_t(\cdot,0) \|_{(s+s_{\text{hom}})} + \| u(\cdot,0) \|_{(s+s_{\text{hom}}+1)} \right)$$

$$(10.47)$$

for all  $t \ge 0$  and all  $s \in \mathbb{R}$ , where the constants C, N and shom have the same dependence as in the case of Lemma 10.17; cf. Remark 10.19. Finally, if  $E_a = \mathbb{C}^{2m}$  (i.e., if  $\beta_{\text{rem}} > \text{Rsp}A_{\infty}$ ; cf. Definition 4.3), then  $\Phi_{\infty}$  is surjective.

**Remark 10.23.** By combining (10.45), (10.46) and (10.47), the norms of  $u(\cdot, 0)$  and  $u_t(\cdot, 0)$  appearing on the right hand side of (10.47) can be replaced by a suitable Sobolev norm of  $\chi$ .

**Remark 10.24.** In order to obtain a similar result in the case of inhomogeneous equations, it is sufficient to combine Lemmas 10.17 and 10.22; cf. Remark 9.23.

**Remark 10.25.** If the assumptions of Proposition 4.16 hold, then the conditions of the lemma are fulfilled. The reasons for this are the following. First of all, (1.2) is weakly balanced, weakly silent and weakly convergent due to Remark 10.3. The lower bound on  $\dot{\ell}$  is a consequence of the assumptions of Proposition 4.16 and Remark 25.14. The fact that the  $\beta_{\text{rem}}$  appearing in the statement of the present lemma coincides with the  $\beta_{\text{rem}}$  appearing in the statement of Proposition 4.16 is a consequence of the fact that, in the context of Proposition 4.16,  $\mathfrak{b}_{s} = \mu$ ; cf. (25.52).

**Remark 10.26.** If, in addition to the assumptions of the lemma, the estimate (10.5) holds, then  $\beta_{\text{rem}}$  can be replaced by  $\min\{2\mathfrak{b}_s, \mathfrak{b}_s + \beta_{\text{der}}, \eta_{\text{mn}}\}$ ; cf. the comments made in connection with (10.17). The dependence of the constants then changes in that they also depend on  $C_{\text{der}}$  and  $\beta_{\text{der}}$ .

**Remark 10.27.** If the assumptions of Proposition 4.16 are satisfied and there are constants  $K_{\text{der}}$  and  $\beta_{\text{der}}$  such that (4.12) holds for all  $t \ge 0$ , then the conclusions of Lemma 10.22 hold with  $\beta_{\text{rem}}$  replaced by  $\min\{2\mu, \mu + \beta_{\text{der}}, \eta_{\text{mn}}\}$ . The reason for this is that (4.12) and Lemma 25.17 imply that (10.5) holds so that Remark 10.26 applies.

*Proof.* Let  $\chi \in C^{\infty}(\overline{M}, E_a)$ ,  $\iota \in \mathcal{I}_B$  and  $\hat{\chi}(\iota) := \langle \chi, \varphi_\iota \rangle_B \in E_a$ ; cf. the notation introduced in Subsection 24.1.2. Let u be a solution to (1.2) and let  $z(\iota, t)$  be given by (24.14). Note that the  $\iota$ 'th mode of the expression inside the norm on the left hand side of (10.47) can be written

$$v - e^{A_{\infty}t}\hat{\chi},\tag{10.48}$$

where we use the notation (10.16). On the other hand, v satisfies (10.15) with F = 0, and  $A_{\text{rem}}$  satisfies (10.17) for  $t \geq T_{\text{ode}}$ . Thus the conditions needed to apply Lemma 9.21 are satisfied. In fact, specifying v by imposing the initial condition  $v(T_{\text{ode}}) := \Psi_{\infty}[\hat{\chi}(\iota)]$  yields the conclusion that (10.48) is small asymptotically; here  $\Psi_{\infty}$  is the map constructed in Lemma 9.21. Note that  $\Psi_{\infty}$  depends on  $\iota$ . However, the estimates involving  $\Psi_{\infty}$  that we need do not. In order to estimate  $|v(T_{\text{ode}})|$  in terms of  $|\hat{\chi}(\iota)|$ , define  $u_{\infty} \in E_a$  by

$$e^{-A_{\infty}T_{\text{ode}}}u_{\infty} = \hat{\chi}(\iota).$$

Then  $|u_{\infty}| = |e^{A_{\infty}T_{\text{ode}}}\hat{\chi}(\iota)|$ , so that (9.45) yields the estimate

$$|v(T_{\text{ode}})| = |\Psi_{\infty}(e^{-A_{\infty}T_{\text{ode}}}u_{\infty})| \le C|u_{\infty}| = C|e^{A_{\infty}T_{\text{ode}}}\hat{\chi}(\iota)|$$

$$< Ce^{(\kappa_1+1)T_{\text{ode}}}|\hat{\chi}(\iota)|$$
(10.49)

where C only depends on  $C_{\text{rem}}$ ,  $\beta_{\text{rem}}$  and  $A_{\infty}$ . In order to estimate  $\mathcal{E}_s(0)$ , it is natural to divide the analysis into two cases. To begin with, assume that  $T_{\text{ode}} = 0$ . Then

$$\mathcal{E}_s^{1/2}(0) \le \langle \nu(\iota) \rangle^s |v(0)| = \langle \nu(\iota) \rangle^s |v(T_{\text{ode}})| \le C \langle \nu(\iota) \rangle^s |\hat{\chi}(\iota)|$$
(10.50)

for all s, where C only depends on  $C_{\text{rem}}$ ,  $\beta_{\text{rem}}$  and  $A_{\infty}$  and we have appealed to (10.49) and the fact that  $T_{\text{ode}} = 0$ . In case  $T_{\text{ode}} > 0$ , it is natural to proceed as in the case of (10.8). Since we wish to estimate dE/dt from below, we need a lower bound on  $\dot{\ell}$ , which is provided by (10.44). An estimate similar to (10.8) yields the conclusion that  $dE/dt \ge -2(\eta_{\text{low}} + \mathfrak{e}_{\text{low}})E$  on  $[0, T_{\text{ode}}]$ , where the constant  $\eta_{\text{low}}$  only depends on  $C_{\text{coeff}}$ ,  $c_{\mathfrak{e}}$  and  $\mathfrak{b}_{\text{low}}$ , where  $c_{\mathfrak{e}} := \|\mathfrak{e}\|_1$ ,  $\mathfrak{e}$  is the function appearing in (10.3) and we have used the fact that  $\mathfrak{g}(t) \geq e^{-c_{\mathfrak{e}}}$  on  $[0, T_{\text{ode}}]$ . Using the notation  $c_{\text{low}} := \|\mathfrak{e}_{\text{low}}\|_1$  and appealing to (10.49) yields

$$E^{1/2}(0) \leq e^{\eta_{\text{low}}T_{\text{ode}}+c_{\text{low}}}E^{1/2}(T_{\text{ode}}) = 2^{-1/2}e^{\eta_{\text{low}}T_{\text{ode}}+c_{\text{low}}}|v(T_{\text{ode}})|$$

$$\leq Ce^{(\eta_{\text{low}}+\kappa_{1}+1)T_{\text{ode}}}|\hat{\chi}(\iota)|,$$
(10.51)

where C only depends on  $C_{\text{rem}}$ ,  $\beta_{\text{rem}}$ ,  $c_{\text{low}}$  and  $A_{\infty}$ . Thus

$$\mathcal{E}_s^{1/2}(0) \le C \langle \nu(\iota) \rangle^{s+s_\infty} |\hat{\chi}(\iota)| \tag{10.52}$$

for all  $s \in \mathbb{R}$ , where C only depends on  $C_{\text{coeff}}$ ,  $C_{\text{mn}}$ ,  $\mathfrak{b}_{\text{low}}$ ,  $\mathfrak{b}_{\text{s}}$ ,  $\eta_{\text{mn}}$ ,  $c_{\text{low}}$ ,  $c_{\mathfrak{e}}$ ,  $A_{\infty}$ ,  $g^{ij}(0)$   $(i, j = 1, \ldots, d)$  and  $a_r(0)$   $(r = 0, \ldots, R)$ ;  $s_{\infty} \ge 0$  only depends on  $C_{\text{coeff}}$ ,  $\mathfrak{b}_{\text{low}}$ ,  $\mathfrak{b}_{\text{s}}$ ,  $c_{\mathfrak{e}}$  and  $A_{\infty}$  (it can be chosen to equal max{ $\eta_{\text{low}} + \kappa_1 + 1, 0$ }/ $\mathfrak{b}_{\text{s}}$ ); and we have appealed to (10.23). Combining (10.50) and (10.52) yields the conclusion that (10.52) holds regardless of whether  $T_{\text{ode}} = 0$  or not.

Let  $\Phi_{\infty}$  be the map taking  $\chi \in C^{\infty}(\bar{M}, E_a)$  to the element of  $C^{\infty}(\bar{M}, \mathbb{C}^{2m})$  whose  $\iota$ 'th mode is given by v(0), where v is constructed as above. Then (10.52) implies that (10.45) holds; note that this implies that  $\Phi_{\infty}$  maps  $\chi \in C^{\infty}(\bar{M}, E_a)$  to an element of  $C^{\infty}(\bar{M}, \mathbb{C}^{2m})$ . Since  $\Phi_{\infty}$  is injective on the level of Fourier coefficients (this is a consequence of Lemma 9.21), it follows that  $\Phi_{\infty}$  is injective. Next, let  $\chi \in C^{\infty}(\bar{M}, E_a)$  and let u be the solution to (1.2) (with f = 0) satisfying (10.46). Due to Lemma 10.17, we know that (10.47) holds with  $\chi$  replaced by some  $V_{\infty} \in C^{\infty}(\bar{M}, E_a)$ . Moreover, the constants C, N and  $s_{\text{hom}}$  have the dependence stated in Lemma 10.17. In order to determine  $V_{\infty}$  in terms of  $\chi$ , note that (10.47) (with  $\chi$  replaced by  $V_{\infty}$ ) implies that

$$\left|v(t) - e^{A_{\infty}t}v_{\infty}\right| \le C\langle t \rangle^{N} e^{(\kappa_{1} - \beta_{\text{rem}})t}$$
(10.53)

where C depends on the initial data and  $\iota$ ; v is defined as in (10.16) where z is the  $\iota$ 'th mode of u; and  $v_{\infty}$  is the  $\iota$ 'th mode of  $V_{\infty}$ . Due to the construction of  $\Phi_{\infty}$ , we know that the estimate (10.53) holds with  $v_{\infty}$  replaced by  $\hat{\chi}(\iota)$ . Thus

$$\left|e^{A_{\infty}t}[\hat{\chi}(\iota)-v_{\infty}]\right| \leq C\langle t \rangle^{N} e^{(\kappa_{1}-\beta_{\mathrm{rem}})t}$$

for  $t \ge 0$ . Since  $\hat{\chi}(\iota) - v_{\infty} \in E_a$ , this estimate implies that  $\hat{\chi}(\iota) - v_{\infty} = 0$ . Thus  $V_{\infty} = \chi$ , and the lemma follows, except for the statement concerning surjectivity.

Assume that  $E_a = \mathbb{C}^{2m}$  and let  $\psi \in C^{\infty}(\overline{M}, \mathbb{C}^{2m})$ . We wish to demonstrate that  $\psi$  is in the image of  $\Phi_{\infty}$ . Let u be the solution to (1.2) with initial data given by (10.46), where the right hand side has been replaced by  $\psi$ . Appealing to Lemma 10.17 yields a  $V_{\infty} \in C^{\infty}(\overline{M}, \mathbb{C}^{2m})$  such that (10.26) holds (with f = 0). Let  $\overline{u}$  be the solution to (1.2) with initial data given by (10.46), where the right hand side has been replaced by  $\Phi_{\infty}(V_{\infty})$ . Then, by the above arguments, (10.26) holds (with f = 0) and u replaced by  $\overline{u}$ . In particular, there is thus a constant C, depending on  $u, \overline{u}$ etc., such that

$$\left\| \begin{pmatrix} u(\cdot,t) \\ u_t(\cdot,t) \end{pmatrix} - \begin{pmatrix} \bar{u}(\cdot,t) \\ \bar{u}_t(\cdot,t) \end{pmatrix} \right\|_{(s)} \le C \langle t \rangle^N e^{(\kappa_1 - \beta_{\rm rem})t}$$
(10.54)

for all  $t \ge 0$  and all  $s \in \mathbb{R}$ . Let z and  $\bar{z}$  be the Fourier coefficients of u and  $\bar{u}$  respectively, let v and  $\bar{v}$  be defined in analogy with (10.16), starting with z and  $\bar{z}$  respectively. Finally, let  $V = v - \bar{v}$ . If  $V(\iota, 0) = 0$  for all  $\iota \in \mathcal{I}_{\mathrm{B}}$ , then  $u = \bar{u}$  (since  $V(\iota, \cdot)$  solves a homogeneous equation,  $V(\iota, 0) = 0$  for all  $\iota \in \mathcal{I}_{\mathrm{B}}$  implies that  $V(\iota, t) = 0$  for all  $\iota \in \mathcal{I}_{\mathrm{B}}$  and all  $t \in I$ ), so that  $\Phi_{\infty}(V_{\infty}) = \psi$ . Thus  $\psi$  is in the image of  $\Phi_{\infty}$ . Assume now that there is a  $\iota \in \mathcal{I}_{\mathrm{B}}$  such that  $V(\iota, 0) \neq 0$ . Note that  $V(\iota, \cdot)$  is a solution to an equation to which Lemma 9.21 applies. Moreover, applying Lemma 9.21 yields a  $\Psi_{\infty}$  which is linear and bijective. Since  $V(\iota, 0) \neq 0$ , there is a  $0 \neq \chi \in \mathbb{C}^{2m}$  such that  $\Psi_{\infty}(\chi) = V(\iota, T_{\mathrm{ode}})$ , where  $T_{\mathrm{ode}}$  is given by Definition 10.1 (for the relevant  $\iota$  under consideration). This means that

$$|V(\iota, t) - e^{A_{\infty}t}\chi| \le C\langle t \rangle^N e^{(\kappa_1 - \beta_{\rm rem})t}$$

for all  $t \ge T_{\text{ode}}$  and a constant C depending on V,  $\iota$  etc. On the other hand, (10.54) implies that the same estimate holds with  $\chi$  set to zero. Due to the fact that  $\text{Rsp}A_{\infty} < \beta_{\text{rem}}$ , these estimates are contradictory. To conclude, every element of  $C^{\infty}(\bar{M}, \mathbb{C}^{2m})$  is in the image of  $\Phi_{\infty}$ . The lemma follows.

# Chapter 11

# ODE analysis in the transparent setting

# 11.1 Introduction

Consider (1.2). In the previous chapter we consider the silent setting. Then  $g^{ij}$ ,  $X^i$ ,  $g^{0i}$ ,  $i, j = 1, \ldots, d$ , and  $a_r^{-2}$ ,  $r = 1, \ldots, R$ , converge to zero exponentially. Next, we wish to consider the transparent setting. In that case, these functions are all bounded. Moreover, some of them converge to zero exponentially. However, some converge to non-zero functions. Let

$$g_{\infty}^{ij} := \lim_{t \to \infty} g^{ij}(t), \quad g_{\infty}^{0i} := \lim_{t \to \infty} g^{0i}(t), \quad q_{\infty,r} := \lim_{t \to \infty} a_r^{-2}(t), \quad X_{\infty}^i := \lim_{t \to \infty} X^i(t).$$
(11.1)

We here assume that these limits exist. We also assume the convergence to be exponential. Note that  $q_{r,\infty} \ge 0$ , and that  $g_{\infty}^{ij}$  are the components of a positive semi-definite matrix. Given  $\iota \in \mathcal{I}_{\mathrm{B}}$ , let

$$\mathfrak{g}_{\infty}(\iota) := \lim_{t \to \infty} \mathfrak{g}(\iota, t) = \left( \sum_{j,l=1}^{d} g_{\infty}^{jl} n_{j} n_{l} + \sum_{r=1}^{R} q_{\infty,r} \nu_{r,i_{r}}^{2}(\iota) \right)^{1/2}.$$
 (11.2)

Assume that  $\alpha$  and  $\zeta$  converge, exponentially, to  $\alpha_{\infty}$  and  $\zeta_{\infty}$  respectively and that  $\mathfrak{g}_{\infty}(\iota) \neq 0$ . Then (24.16) can be written

$$\dot{v} = A(\iota)v + A_{\iota,\text{rem}}v + F_{\iota}, \qquad (11.3)$$

where

$$A(\iota) := \begin{pmatrix} 0 & \mathfrak{g}_{\infty} \mathrm{Id}_m \\ -\mathfrak{g}_{\infty} \mathrm{Id}_m - i \mathfrak{g}_{\infty}^{-1} n_l X_{\infty}^l - \mathfrak{g}_{\infty}^{-1} \zeta_{\infty} & 2i n_l g_{\infty}^{0l} \mathrm{Id}_m - \alpha_{\infty} \end{pmatrix}.$$
 (11.4)

Moreover,

$$v(\iota,t) := \begin{pmatrix} \mathfrak{g}_{\infty}(\iota)z(\iota,t) \\ \dot{z}(\iota,t) \end{pmatrix}, \quad F_{\iota}(t) := \begin{pmatrix} 0 \\ \hat{f}(\iota,t) \end{pmatrix}.$$
(11.5)

Finally,

$$||A_{\iota,\text{rem}}(t)|| \le C_{\text{ts}} e^{-\beta_{\text{ts}}\bar{t}}$$
(11.6)

for  $t \geq T_{\rm ts}$ . Here  $0 < C_{\rm ts}, \beta_{\rm ts} \in \mathbb{R}$  are independent of  $\iota$ , but  $T_{\rm ts}$  does depend on  $\iota$ . Moreover,  $\overline{t} := t - T_{\rm ts}$ . Formally, the equation (11.3) is identical to (9.3). However, there is a fundamental difference. In (9.3), the matrix A is fixed once and for all, but in (11.3) it depends on  $\iota$ . In the present chapter, we would, essentially, like to repeat the analysis presented in Chapter 9. However, this involves dividing  $\mathbb{C}^{2m}$  into the first and second generalised eigenspace in the  $\beta_{\rm ts}, A(\iota)$ decomposition of  $\mathbb{C}^{2m}$ . As opposed to the case studied in Chapter 9, there are potentially infinitely many different such generalised eigenspaces in the present setting. Similarly, there are potentially infinitely many different matrices  $T_{A(\iota)}$  with the properties stated in Definition 9.11; one for each  $\iota$ . This is a problem, since if we choose the matrices  $T_{A(\iota)}$  so that their norms are uniformly bounded, there is no reason why  $||T_{A(\iota)}^{-1}||$  should be uniformly bounded for  $\iota \in \mathcal{I}_{B}$ . Since  $||T_{A(\iota)}^{-1}||$  appears in the estimates, this could potentially cause a problem. In some of the estimates, an even more subtle dependence on  $A(\iota)$  appears. As a consequence, it is clear that we, in the present setting, need to keep very detailed track of how the algebraic constructions carried out in Section 9.1 depend on  $A(\iota)$ .

Consider  $A(\iota)$  defined by (11.4). The bottom left component of this matrix contains the expression  $-i\mathfrak{g}_{\infty}^{-1}n_lX_{\infty}^l$ . By varying  $\iota$ , this expression gives rise to a large number of linear combinations of the matrices  $X_{\infty}^l$ . In fact, one would typically expect the closure of the set of  $-i\mathfrak{g}_{\infty}^{-1}n_lX_{\infty}^l$ ,  $\iota \in \mathcal{I}_B$ , to be *d*-dimensional. Dealing with such a situation can be expected to be quite difficult, and we do not attempt to do so here. In fact, we consider, exclusively, the following two restricted situations.

**Definition 11.1.** Consider (1.2). Assume the associated metric to be such that (M, g) is a canonical separable cosmological model manifold. Then (1.2) is said to be an *X*-non-trivial weakly transparent equation if the following holds. The limits (11.1) exist and there is a  $j \in \{1, \ldots, d\}$  such that  $X_{\infty}^{j} \neq 0$ . Moreover,  $X_{\infty}^{l} = 0$  for  $l \neq j$ ;  $q_{\infty,r} = 0$  for  $r = 1, \ldots, R$ ;  $g_{\infty}^{jj} > 0$ ;  $g_{\infty}^{kl} = 0$  unless k = l = j; and  $g_{\infty}^{0l} = 0$  for  $l = 1, \ldots, d$ . Given an X-non-trivial weakly transparent equation (1.2), let  $d_{\text{ts}} := 1$  and  $j_1 := j$ .

**Remarks 11.2.** If (1.2) is an X-non-trivial weakly transparent equation, then  $\mathfrak{g}_{\infty} = (g_{\infty}^{jj})^{1/2} |n_j|$ . Thus, assuming  $n_j \neq 0$ ,

$$-i\mathfrak{g}_{\infty}^{-1}n_l X_{\infty}^l = -i(g_{\infty}^{jj})^{-1/2}\operatorname{sgn}(n_j)X_{\infty}^j.$$

In other words, the left hand side can only be one of two possible matrices. Note also that if we would allow  $q_{\infty,r} > 0$ , then the set of  $-i\mathfrak{g}_{\infty}^{-1}n_l X_{\infty}^l$  could potentially have every point on the line between  $-i(g_{\infty}^{jj})^{-1/2}X_{\infty}^j$  and  $i(g_{\infty}^{jj})^{-1/2}X_{\infty}^j$  as an accumulation point. This leads to complications we wish to avoid.

**Definition 11.3.** Consider (1.2). Assume the associated metric to be such that (M, g) is a canonical separable cosmological model manifold. Then (1.2) is said to be an *X*-degenerate weakly transparent equation if the following holds. The limits (11.1) exist,  $X_{\infty}^{j} = 0$  and  $g_{\infty}^{0j} = 0$  for  $j = 1, \ldots, d$ . Moreover,  $\{1, \ldots, d\}$  can be written as the union of two disjoint sets  $\{j_{1}, \ldots, j_{d_{\text{ts}}}\}$  and  $\{\bar{j}_{1}, \ldots, \bar{j}_{d_{\text{sil}}}\}$  (if  $d_{\text{ts}} = 0$  or  $d_{\text{sil}} = 0$  the corresponding set is empty) such that the spatial limit metric is positive definite on  $P_{\infty} := \text{span}\{e_{j_{1}}, \ldots, e_{j_{d_{\text{ts}}}}\}$  and vanishes on  $N_{\infty} := \text{span}\{e_{\bar{j}_{1}}, \ldots, e_{\bar{j}_{d_{\text{sil}}}}\}$ . Finally, if  $d_{\text{ts}} = 0$ , there is an  $r \in \{1, \ldots, R\}$  such that  $q_{\infty, r} > 0$ .

**Remarks 11.4.** In the definition, the spatial limit metric corresponds to the matrix with components  $g_{\infty}^{jk}$ . Moreover, the statement that the spatial limit metric vanishes on  $N_{\infty}$  means that if  $v \in \mathbb{R}^d$  and  $w \in N_{\infty}$ , then  $g_{\infty}^{jk}v_jw_k = 0$ .

**Remarks 11.5.** One purpose of the restriction on the spatial limit metric is to ensure that the only accumulation point of  $\{\mathfrak{g}_{\infty}(\iota) : \iota \in \mathcal{I}_{\mathrm{B}}\}$  is  $\infty$ . To see that this is a consequence of the restriction, fix R > 0 and assume that  $\mathfrak{g}_{\infty}(\iota) \leq R$ . If  $q_{\infty,r} > 0$ , this means that the  $\nu_{r,i}$  component of  $\nu(\iota)$  is bounded; cf. (24.4) for an explanation of the terminology. Thus  $\nu_{r,i}$  takes one of a finite number of values. Similarly,  $n_{j_l}$ ,  $l = 1, \ldots, d_{\mathrm{ts}}$ , can only take one of a finite number of values due to the positive definiteness of the limit metric on  $P_{\infty}$ . In other words, the only elements of  $\nu(\iota)$  that contribute to  $\mathfrak{g}_{\infty}(\iota)$  are restricted to belong to a finite set if  $\mathfrak{g}_{\infty}(\iota) \leq R$ .

Returning to (11.4), it is clear that if one of the Definitions 11.1 and 11.3 is satisfied, then the matrices  $A(\iota)$  are of the form

$$A_{\mu} := \begin{pmatrix} 0 & \mu \mathrm{Id}_{m} \\ -\mu \mathrm{Id}_{m} + iV + 2\mu^{-1}W & U \end{pmatrix},$$
(11.7)

where  $U, V, W \in \mathbf{M}_m(\mathbb{C})$ . Moreover,  $\mu \in \mathcal{M}$ , where  $\mathcal{M}$  is a countable subset of  $(0, \infty)$  with  $\infty$  as its only accumulation point. The factor 2 is included for future convenience.

# 11.2 Asymptotic eigenvalue calculation

The first step of our analysis is to calculate the asymptotic values of the eigenvalues of the matrix  $A_{\mu}$  introduced in (11.7). Let, to this end,

$$U_{\pm} := \frac{1}{2} (U \pm V), \quad S_1 := \begin{pmatrix} i \operatorname{Id}_m & \operatorname{Id}_m \\ \operatorname{Id}_m & i \operatorname{Id}_m \end{pmatrix}.$$
(11.8)

Then

$$S_1^{-1}A_{\mu}S_1 = \begin{pmatrix} -i\mu \mathrm{Id}_m + U_- + i\mu^{-1}W & iU_+ + \mu^{-1}W \\ -iU_- + \mu^{-1}W & i\mu \mathrm{Id}_m + U_+ - i\mu^{-1}W \end{pmatrix}$$

Let  $S_{\pm} \in \mathbf{GL}_m(\mathbb{C})$  be such that

$$\gamma_{\pm} := (S_{\pm})^{-1} U_{\pm} S_{\pm} \tag{11.9}$$

is of Jordan normal form. Letting  $S_2 := \text{diag}\{S_-, S_+\}$  and  $S_{12} := S_1S_2$  then yields

$$N_{\mu} := S_{12}^{-1} A_{\mu} S_{12} = \begin{pmatrix} -i\mu \mathrm{Id}_{m} + \gamma_{-} + i\mu^{-1}W_{-} & i\delta_{+} + \mu^{-1}W_{-+} \\ -i\delta_{-} + \mu^{-1}W_{+-} & i\mu \mathrm{Id}_{m} + \gamma_{+} - i\mu^{-1}W_{+} \end{pmatrix},$$
(11.10)

where

$$W_{\pm} := (S_{\pm})^{-1}WS_{\pm}, \quad W_{+-} := (S_{+})^{-1}WS_{-}, \quad W_{-+} := (S_{-})^{-1}WS_{+}.$$

Moreover,  $\delta_{\pm}$  is defined by

$$\delta_+ := (S_-)^{-1} U_+ S_+, \quad \delta_- := (S_+)^{-1} U_- S_-$$

and the  $\gamma_{\pm}$  are defined in (11.9). For the sake of brevity, it is convenient to introduce

$$R_{\mu}^{-} := \gamma_{-} + i\mu^{-1}W_{-}, \quad R_{\mu}^{+} := \gamma_{+} - i\mu^{-1}W_{+}, \tag{11.11}$$

$$Q_{\mu}^{+} := i\delta_{+} + \mu^{-1}W_{-+}, \quad Q_{\mu}^{-} := -i\delta_{-} + \mu^{-1}W_{+-}.$$
(11.12)

Then

$$N_{\mu} = \begin{pmatrix} -i\mu \mathrm{Id}_{m} + R_{\mu}^{-} & Q_{\mu}^{+} \\ Q_{\mu}^{-} & i\mu \mathrm{Id}_{m} + R_{\mu}^{+} \end{pmatrix}.$$
 (11.13)

**Lemma 11.6.** Consider the matrix  $N_{\mu}$  given by (11.13). Let  $\lambda_{j,\pm}$ ,  $j = 1, \ldots, p_{\pm}$ , be the distinct eigenvalues of  $\gamma_{\pm}$ , and  $m_{j,\pm}$  be the corresponding algebraic multiplicities, where the matrices  $\gamma_{\pm}$  are introduced in (11.9). There are then constants  $0 < c_a, \mu_a \in \mathbb{R}$  such that for  $\mu \geq \mu_a$  and  $j = 1, \ldots, p_{\pm}$ , there are  $m_{j,\pm}$  eigenvalues of  $N_{\mu}$  in a ball of radius  $c_a \mu^{-1/m_{j,\pm}}$  and centre  $\pm i\mu + \lambda_{j,\pm}$ .

**Remark 11.7.** Since the sum of the  $m_{j,+}$  equal m, and similarly for the  $m_{j,-}$ , each eigenvalue of  $N_{\mu}$  has to belong to a ball of radius  $c_a \mu^{-1/m_{j,\pm}}$  and centre  $\pm i\mu + \lambda_{j,\pm}$  for some choice of j and  $\pm$ .

**Remark 11.8.** The constants  $c_a$  and  $\mu_a$  only depend on U, V and W.

*Proof.* In order to obtain estimates for the eigenvalues, we proceed step by step. Moreover, it is convenient to consider

$$N_{\mu,r} = \begin{pmatrix} -i\mu \mathrm{Id}_m + R^-_{\mu,r} & Q^+_{\mu,r} \\ Q^-_{\mu,r} & i\mu \mathrm{Id}_m + R^+_{\mu,r} \end{pmatrix}$$
(11.14)

for  $r \in [0, 1]$ . Here

$$R_{\mu,r}^{-} := \gamma_{-} + ir\mu^{-1}W_{-}, \quad R_{\mu,r}^{+} := \gamma_{+} - ir\mu^{-1}W_{+}, \quad Q_{\mu,r}^{+} := rQ_{\mu}^{+}, \quad Q_{\mu,r}^{-} := rQ_{\mu}^{-}.$$

Note that  $N_{\mu,1} = N_{\mu}$ . To begin with, we appeal to Gershgorin's Theorem; cf., e.g., [44, Proposition 5.12, p. 102]. For  $j = 1, \ldots, 2m$ , let

$$\rho_j := \sum_{l \neq j} |N_{\mu,r,lj}| \le \varrho_0 + \mu^{-1} \varrho_1,$$

where  $0 \leq \varrho_l \in \mathbb{R}$ , l = 0, 1, are independent of  $\mu$  and  $r \in [0, 1]$ , and  $N_{\mu,r,lj}$  is the lj'th component of  $N_{\mu,r}$ . Then the eigenvalues of  $N_{\mu,r}$  are contained in the discs  $B_{\rho_j}(N_{\mu,r,jj})$  (no summation on j). In particular, there are thus constants  $0 < \mu_0, R \in \mathbb{R}$  such that for  $\mu \geq \mu_0$  and  $r \in [0, 1]$ , the two balls  $B_R(\pm i\mu)$  are disjoint and contain the eigenvalues of  $N_{\mu,r}$ . Assume that  $\mu \geq \mu_0$ , that  $r \in [0, 1]$  and that  $\lambda$  is an eigenvalue belonging to  $B_R(-i\mu)$ . Consider

$$N_{\mu,r} - \lambda \mathrm{Id}_{2m} = \begin{pmatrix} N_{\mu,r,\lambda}^- & Q_{\mu,r}^+ \\ Q_{\mu,r}^- & N_{\mu,r,\lambda}^+ \end{pmatrix}, \qquad (11.15)$$

where

$$N_{\mu,r,\lambda}^{\pm} := -(\lambda \mp i\mu) \mathrm{Id}_m + R_{\mu,r}^{\pm}.$$

Due to the assumptions, the diagonal elements of  $N^+_{\mu,r,\lambda}$  are contained in a ball centred at  $2i\mu$  and with radius bounded by a constant independent of  $\mu$  and  $r \in [0, 1]$  (for  $\mu$  large enough). There is thus a  $\mu_1 \geq \mu_0$  such that for  $\mu \geq \mu_1$  and  $r \in [0, 1]$ ,  $N^+_{\mu,r,\lambda}$  is invertible and

$$\|(N_{\mu,r,\lambda}^+)^{-1}\| \le c_0 \mu^{-1} \tag{11.16}$$

for some constant  $c_0$  independent of  $\mu$  and  $r \in [0, 1]$ . In particular,

$$0 = \det(N_{\mu,r} - \lambda \mathrm{Id}_{2m}) = \det \left( \begin{array}{cc} N_{\mu,r,\lambda}^{-} - Q_{\mu,r}^{+} (N_{\mu,r,\lambda}^{+})^{-1} Q_{\mu,r}^{-} & Q_{\mu,r}^{+} \\ 0 & N_{\mu,r,\lambda}^{+} \end{array} \right).$$

Since  $N^+_{\mu,r,\lambda}$  is invertible, this implies

$$\det \left[ N^{-}_{\mu,r,\lambda} - Q^{+}_{\mu,r} (N^{+}_{\mu,r,\lambda})^{-1} Q^{-}_{\mu,r} \right] = 0.$$

On the other hand,

$$N_{\mu,r,\lambda}^{-} - Q_{\mu,r}^{+} (N_{\mu,r,\lambda}^{+})^{-1} Q_{\mu,r}^{-} = -(\lambda + i\mu) \mathrm{Id}_{m} + \gamma_{-} + \mathrm{P}_{\mu,r}, \qquad (11.17)$$

where  $P_{\mu,r}$  is a matrix satisfying

$$\|\mathbf{P}_{\mu,r}\| \le c_0 \mu^{-1} \tag{11.18}$$

for  $\mu \ge \mu_1$  and  $r \in [0, 1]$ , where  $c_0$ , again, is independent of  $\mu$  and  $r \in [0, 1]$ . Note that the norms of the first two terms on the right hand side of (11.17) are (individually) bounded by a constant independent of  $\mu$  and r. Moreover, the third term on the right hand side of (11.17) satisfies the bound (11.18). Note that we can consider the determinant of the right hand side of (11.17) as a multilinear function, say Det, of the corresponding columns:

$$det[-(\lambda + i\mu)Id_m + \gamma_- + P_{\mu,r}] = Det(K_{\mu,1} + P_{\mu,r,1}, \dots, K_{\mu,m} + P_{\mu,r,m})$$
  
= Det(K<sub>\mu,1</sub>, \dots, K<sub>\mu,m</sub>) + Det(K<sub>\mu,1</sub>, \dots, K<sub>\mu,m-1</sub>, P<sub>\mu,r,m</sub>)  
+ Det(K<sub>\mu,1</sub>, \dots, P<sub>\mu,r,m-1</sub>, K<sub>\mu,m</sub> + P<sub>\mu,r,m</sub>) + \dots + Det(P<sub>\mu,r,1</sub>, \dots, K<sub>\mu,m</sub> + P<sub>\mu,r,m</sub>), (11.19)

where  $K_{\mu,l}$  denotes the *l*'th column of the sum of the first two matrices on the right hand side of (11.17) and  $P_{\mu,r,l}$  denotes the *l*'th column of  $P_{\mu,r}$ . We know that the left hand side of (11.19) is zero. Moreover, due to (11.18) and the above observations, all the terms on the far right hand side of (11.19), except for the first one, can be bounded by a  $c_0\mu^{-1}$  for some constant  $c_0$  independent of  $\mu$  and  $r \in [0, 1]$ . Thus

$$|\det[-(\lambda + i\mu)\mathrm{Id}_m + \gamma_-]| \le c_0\mu^{-1}$$

for  $\mu \ge \mu_1$  and some constant  $c_0$  independent of  $\mu$  and  $r \in [0, 1]$ . With notation as in the statement of the lemma, it follows that

$$|(\lambda - \lambda_{1,-} + i\mu)^{m_{1,-}} \cdots (\lambda - \lambda_{p_{-,-}} + i\mu)^{m_{p_{-,-}}}| \le c_0\mu^{-1}$$
(11.20)

for  $\mu \ge \mu_1$ ; recall that  $\gamma_-$  is of Jordan normal form. Due to this estimate, it is clear that the smallest of the  $|\lambda - \lambda_{s,-} + i\mu|$  has to be bounded by  $c_0\mu^{-1/m}$ . In particular, there are

constants  $\mu_2 \ge \mu_1$  and  $0 < c_0 \in \mathbb{R}$  such that for  $\mu \ge \mu_2$  and  $r \in [0, 1]$ , there is one and only one  $j \in \{1, \ldots, p_-\}$  such that  $|\lambda - \lambda_{j,-} + i\mu| \le c_0 \mu^{-1/m}$ . In order to justify the uniqueness of j, we use the fact that if  $|\lambda - \lambda_{j,-} + i\mu| \le c_0 \mu^{-1/m}$ , then, for  $l \ne j$ ,

$$|\lambda - \lambda_{l,-} + i\mu| \ge |\lambda_{l,-} - \lambda_{j,-}| - |\lambda - \lambda_{j,-} + i\mu| \ge |\lambda_{l,-} - \lambda_{j,-}| - c_0 \mu^{-1/m}.$$

Since the first term on the far right hand side has a strictly positive lower bound for all  $l \neq j$ (independent of  $\mu$  and r), and since the second term converges to zero as  $\mu \to \infty$ , it is clear that the desired statement holds for  $\mu$  large enough. We also obtain the conclusion that for  $\mu \geq \mu_2$ , there is a  $c_1 > 0$ , independent of  $\mu$  and  $r \in [0, 1]$ , such that for  $l \in \{1, \ldots, p_-\}$  with  $l \neq j$ ,  $|\lambda - \lambda_{l,-} + i\mu| \geq c_1$ . Combining this observation with (11.20) yields

$$|\lambda - \lambda_{j,-} + i\mu| \le c_2 \mu^{-1/m_{j,-}} \tag{11.21}$$

for  $\mu \ge \mu_2$  and some constant  $c_2$  independent of  $\mu$  and  $r \in [0, 1]$ .

In order to prove that there are  $m_{j,-}$  eigenvalues of  $N_{\mu}$  in a ball of radius  $c_2 \mu^{-1/m_{j,-}}$  and centre  $-i\mu + \lambda_{j,-}$ , we proceed as in the proof of [44, Theorem 5.7, p. 102]. Let

$$\mathcal{P}_{\mu,r}(z) = \det(N_{\mu,r} - z\mathrm{Id}_{2m}).$$

By the above, we, first of all, know that the zeros of this polynomial are contained in the disjoint balls  $B_R(\pm i\mu)$ . Focusing on the zeros contained in  $B_R(-i\mu)$ , they are contained in the balls of radius  $c_2\mu^{-1/m_{j,-}}$  and centre  $-i\mu + \lambda_{j,-}$  (and for  $\mu$  large enough, we can assume the balls with the same centre and twice the radius to be disjoint). Let  $\Gamma$  be a curve parametrising the boundary circle of the ball of radius  $c_2\mu^{-1/m_{j,-}}$  and centre  $-i\mu + \lambda_{j,-}$ . Define

$$Z_{\mu}(r) := \frac{1}{2\pi i} \int_{\Gamma} \frac{\mathcal{P}'_{\mu,r}(z)}{\mathcal{P}_{\mu,r}(z)} dz.$$

Note that  $\mathcal{P}_{\mu,r}(z)$  is continuous (in r and z) and non-vanishing on  $\Gamma$  for all  $r \in [0, 1]$ , assuming  $\mu$  to be large enough. Moreover,  $Z_{\mu}(r)$  is a continuous function of r. Finally,  $Z_{\mu}(r)$  counts the number of zeros of the polynomial  $\mathcal{P}_{\mu,r}(z)$  inside  $\Gamma$ . Since  $Z_{\mu}(r)$  is integer valued and continuous,  $Z_{\mu}(1) = Z_{\mu}(0)$ . On the other hand, by the definition of  $N_{\mu,r}$ , it is clear that  $Z_{\mu}(0) = m_{j,-}$ . To conclude, there are  $m_{j,-}$  eigenvalues of  $N_{\mu}$  in a ball of radius  $c_a \mu^{-1/m_{j,-}}$  and centre  $-i\mu + \lambda_{j,-}$ . By a similar argument, there are  $m_{j,+}$  eigenvalues of  $N_{\mu}$  in a ball of radius  $c_a \mu^{-1/m_{j,+}}$  and centre  $i\mu + \lambda_{j,+}$ . The lemma follows.

# 11.3 Computing projections onto generalised eigenspaces

In the previous section, we obtained asymptotic estimates for the eigenvalues of the matrix  $A_{\mu}$  introduced in (11.7). The purpose of the present section is to take the first step towards obtaining asymptotic expressions for the generalised eigenspaces of interest when carrying out an analysis similar to that described in Chapter 9. To begin with, we need to recall how to evaluate holomorphic functions on complex square matrices.

#### 11.3.1 Evaluating holomorphic functions on complex square matrices

Let  $A \in \mathbf{M}_m(\mathbb{C})$  and f be a holomorphic function on an open set  $U \subseteq \mathbb{C}$  (which need not be connected). Assume that  $\operatorname{Sp} A \subset U$  and let P be a polynomial such that  $P^{(r)}(\lambda) = f^{(r)}(\lambda)$  for every  $\lambda \in \operatorname{Sp} A$  and every  $0 \leq r \leq m-1$ ; that such a polynomial exists follows, e.g., from [46, Theorem 1, p. 42]. Given this polynomial, f(A) is defined by f(A) := P(A); that f(A) does not depend on the particular polynomial chosen is a consequence of the Cayley-Hamilton theorem. The above discussion is merely a summary of the beginning of [44, Section 5.5], and we refer the reader interested in more details to [44]. Due to [44, Proposition 5.7, pp. 94–95], this definition leads, e.g., to the following natural properties: if M is non-singular, then  $f(M^{-1}AM) = M^{-1}f(A)M$ ; (fg)(A) = f(A)g(A);  $\operatorname{Sp} f(A) = f(\operatorname{Sp} A)$ ;  $(f \circ g)(A) = f(g(A))$  etc. In all of these equalities, we assume f and g to be holomorphic functions defined on open sets  $U \subseteq \mathbb{C}$  and  $V \subseteq \mathbb{C}$  respectively. Moreover, we assume U and V to be such that the expressions formulated above make sense.

Next we want to describe a different way of defining f(A). The corresponding perspective is based on results from complex analysis, and we therefore briefly recall the relevant results and notation.

**Theorem 11.9** (Cauchy's Theorem). Let  $U \subset \mathbb{C}$  be an open set and  $f : U \to \mathbb{C}$  be a holomorphic function. If  $0 \le m \in \mathbb{Z}$  and  $\Gamma$  is a cycle in U such that  $\operatorname{Ind}_{\Gamma}(z) = 0$  for all  $z \notin U$ , then

$$\frac{1}{m!} f^{(m)}(z) \cdot \text{Ind}_{\Gamma}(z) = \frac{1}{2\pi i} \int_{\Gamma} \frac{f(w)}{(w-z)^{m+1}} dw$$

for all  $z \in U - \Gamma^*$ .

**Remarks 11.10.** Here a cycle  $\Gamma$  is a collection of closed paths, say  $\gamma_j$  (j = 1, ..., l), where a path is a piecewise continuously differentiable curve defined on a compact interval (such that the left- and right-hand derivatives always exist, and the number of derivative discontinuities is finite). Moreover,  $\Gamma^*$  is the union of the ranges of the  $\gamma_j$ 's and if h is a continuous function on  $\Gamma^*$ , then

$$\int_{\Gamma} h(z)dz := \sum_{j=1}^{l} \int_{\gamma_j} h(z)dz.$$

Finally, if  $z \notin \Gamma^*$ ,

$$\operatorname{Ind}_{\Gamma}(z) := \frac{1}{2\pi i} \int_{\Gamma} \frac{d\zeta}{\zeta - z}.$$

The theorem is essentially a consequence of [43, Theorem 10.35, pp. 218–219]; cf. also [43, Exercise 7, p. 228].

Combining Cauchy's Theorem and the above definition of f(A) yields the following result (which is merely a slight reformulation of [44, Proposition 5.8, p. 95]). The reader interested in a proof is referred to [44, p. 96].

**Theorem 11.11.** Let  $U \subset \mathbb{C}$  be an open set and  $f : U \to \mathbb{C}$  be a holomorphic function. Assume that  $A \in \mathbf{M}_m(\mathbb{C})$  is such that  $\operatorname{Sp} A \subset U$ . If  $\Gamma$  is a cycle in U such that  $\lambda \notin \Gamma^*$  for all  $\lambda \in \operatorname{Sp} A$ ;  $\operatorname{Ind}_{\Gamma}(\lambda) = 1$  for all  $\lambda \in \operatorname{Sp} A$ ; and  $\operatorname{Ind}_{\Gamma}(z) = 0$  for all  $z \notin U$ , then

$$f(A) = \frac{1}{2\pi i} \int_{\Gamma} f(z) (z \operatorname{Id}_m - A)^{-1} dz.$$

Next we use these observations to derive formulae for the projections onto the generalised eigenspaces of  $A_{\mu}$  that we are interested in.

#### 11.3.2 Formulae for projections onto generalised eigenspaces

Due to Lemma 11.6, we know that the eigenvalues of  $N_{\mu}$  are contained in balls centred at

$$\mu_{j,+} := i\mu + \lambda_{j,+}, \quad \mu_{l,-} := -i\mu + \lambda_{l,-},$$

where  $j = 1, ..., p_+$  and  $l = 1, ..., p_-$ . Let r > 0 be small enough that the balls  $B_{2r}(\lambda_{j,+})$  are disjoint for  $j = 1, ..., p_+$ , and the balls  $B_{2r}(\lambda_{l,-})$  are disjoint for  $l = 1, ..., p_-$ . Then, there is a  $\mu_0 > 0$  such that for  $\mu \ge \mu_0$ , the balls  $B_{2r}(\mu_{j,+})$  and  $B_{2r}(\mu_{l,-})$ ,  $j = 1, ..., p_+$  and  $l = 1, ..., p_-$ , are all disjoint. Moreover, for  $\mu \ge \mu_0$ , all the eigenvalues of  $N_{\mu}$  belong to  $U_{\mu,0}$ , defined to be the union of the  $B_r(\mu_{j,+})$  and  $B_r(\mu_{l,-})$  for  $j = 1, \ldots, p_+$  and  $l = 1, \ldots, p_-$ ; this is due to Lemma 11.6. A final application of Lemma 11.6 allows us to assume that there are  $m_{j,+}$  and  $m_{l,-}$  eigenvalues in  $B_r(\mu_{j,+})$  and  $B_r(\mu_{l,-})$  respectively. Define

$$\gamma_{\mu,j,+}(t) := \mu_{j,+} + re^{2\pi i t}, \quad \gamma_{\mu,l,-}(t) := \mu_{l,-} + re^{2\pi i t}$$

for  $j = 1, \ldots, p_+$ ,  $l = 1, \ldots, p_-$ . Let  $\Gamma_{\mu}$  be the cycle consisting of all the  $\gamma_{\mu,j,+}$  and  $\gamma_{\mu,l,-}$ . Finally, let  $U_{\mu}$  be the union of all the balls  $B_{3r/2}(\mu_{j,+})$ ,  $j = 1, \ldots, p_+$ , and  $B_{3r/2}(\mu_{l,-})$ ,  $l = 1, \ldots, p_-$ ;  $f_{\mu,j,+}: U_{\mu} \to \mathbb{C}$  be defined by  $f_{\mu,j,+}(z) = 1$  for  $z \in B_{3r/2}(\mu_{j,+})$  and  $f_{\mu,j,+}(z) = 0$  for  $z \notin B_{3r/2}(\mu_{j,+})$ ; and let  $f_{\mu,l,-}$  be defined analogously. Let  $\Omega_{\mu,j,\pm}$  be the complement of  $\gamma_{\mu,j,\pm}^*$ , where  $\gamma_{\mu,j,\pm}^*$  denotes the range of  $\gamma_{\mu,j,\pm}$ . Then  $\operatorname{Ind}_{\gamma_{\mu,j,\pm}}$  is an integer valued function on  $\Omega_{\mu,j,\pm}$ which is constant on each component of  $\Omega_{\mu,j,\pm}$  and vanishes on the unbounded component of  $\Omega_{\mu,j,\pm}$ ; cf., e.g., [43, Theorem 10.10, p. 203]. Moreover, it is obvious that  $\operatorname{Ind}_{\gamma_{\mu,j,\pm}}(\mu_{j,\pm}) = 1$ . Thus it is clear that  $\operatorname{Ind}_{\gamma_{\mu,j,\pm}}(z) = 0$  if  $z \notin \overline{B}_r(\mu_{j,\pm})$  and  $\operatorname{Ind}_{\gamma_{\mu,j,\pm}}(z) = 1$  if  $z \in B_r(\mu_{j,\pm})$ . In particular,  $\operatorname{Ind}_{\Gamma_{\mu}}(z) = 1$  if  $z \in U_{\mu,0}$  and  $\operatorname{Ind}_{\Gamma_{\mu}}(z) = 0$  if  $z \notin U_{\mu,0}$ . Thus  $U_{\mu}$  and  $\Gamma_{\mu}$  fulfil the conditions of Theorem 11.11. Moreover, the  $f_{\mu,j,\pm}$  are holomorphic on  $U_{\mu}$ ;  $f_{\mu,j,+}f_{\mu,l,-} = 0$  for all  $j = 1, \ldots, p_+$  and  $l = 1, \ldots, p_-$ ;  $f_{\mu,j,+}f_{\mu,l,+} = 0$  for all  $j, l = 1, \ldots, p_+$  such that  $j \neq l$ ;  $f_{\mu,j,-}f_{\mu,l,-} = 0$  for all  $j, l = 1, \ldots, p_-$  such that  $j \neq l$ ; and  $f_{\mu,j,\pm}^2 = f_{\mu,j,\pm}$  for all  $j = 1, \ldots, p_{\pm}$ . Define  $\pi_{\mu,j,\pm} := f_{\mu,j,\pm}(N_{\mu})$ . Then the  $\pi_{\mu,j,\pm}$ 's have properties similar to the  $f_{\mu,j,\pm}$ . Moreover,

$$\sum_{j=1}^{p_+} \pi_{\mu,j,+} + \sum_{j=1}^{p_-} \pi_{\mu,j,-} = \sum_{j=1}^{p_+} f_{\mu,j,+}(N_\mu) + \sum_{j=1}^{p_-} f_{\mu,j,-}(N_\mu) = \mathrm{Id}_{2m},$$

where the last equality is a consequence of the fact that the sum of the  $f_{\mu,j,\pm}$  equals 1 on Sp $N_{\mu}$ . Letting  $E_{\mu,j,\pm} := \pi_{\mu,j,\pm}(\mathbb{C}^{2m})$ , it is thus clear that  $\mathbb{C}^{2m}$  is the direct sum of the  $E_{\mu,j,\pm}$ . Moreover, since  $\pi_{\mu,j,\pm}$  and  $N_{\mu}$  commute, it is clear that  $N_{\mu}(E_{\mu,j,\pm}) \subseteq E_{\mu,j,\pm}$ . Finally, it is of interest to relate the  $E_{\mu,j,\pm}$ 's to the generalised eigenspaces associated with the eigenvalues of  $N_{\mu}$ . To that end, we appeal to the following lemma.

**Lemma 11.12.** Let each of the sets  $U \subset \mathbb{C}$  and  $V \subset \mathbb{C}$  be a union of a finite number of disjoint open balls with finite radius. Assume, moreover, that for each component of V, its closure is contained in a component of U, and that each component of U contains a unique component of V. Assume that  $A \in \mathbf{M}_k(\mathbb{C})$  is such that  $\operatorname{Sp} A \subset V$ . Let  $V_a \subset U_a$ , be components of Vand U respectively. Let  $f : U \to \mathbb{C}$  be defined by the requirement that f(z) = 1 for  $z \in U_a$  and f(z) = 0 for  $z \notin U_a$ . Then f(A) is the projection onto the direct sum of the generalised eigenspaces corresponding to the eigenvalues in  $\operatorname{Sp} A \cap V_a$ .

**Remark 11.13.** In particular, given the terminology introduced prior to the statement of the lemma,  $\pi_{\mu,j,\pm}$  is the projection onto the direct sum of the generalised eigenspaces of  $N_{\mu}$  corresponding to the eigenvalues in  $B_r(\mu_{j,\pm})$ .

*Proof.* Let  $\Gamma$  consist of the boundary circles of the components of V. Then U, f, A and  $\Gamma$  are such that the conditions of Theorem 11.11 are satisfied. Let  $\gamma_a$  be a curve parametrising the boundary circle  $\partial V_a$ . Then Theorem 11.11 yields

$$\pi := f(A) = \frac{1}{2\pi i} \int_{\gamma_a} (z \operatorname{Id}_k - A)^{-1} dz.$$

Let M be a non-singular matrix such that

$$M^{-1}AM = \operatorname{diag}\{\lambda_1 \operatorname{Id}_{l_1} + N_1, \dots, \lambda_r \operatorname{Id}_{l_r} + N_r\},\tag{11.22}$$

where  $\lambda_1, \ldots, \lambda_r$  are the (distinct) eigenvalues of A,  $l_1, \ldots, l_r$  are the corresponding algebraic multiplicities, and  $N_l$ ,  $l = 1, \ldots, r$ , are upper triangular nilpotent matrices. That such a matrix M exists follows from the Jordan normal decomposition of A. Fixing  $j \in \{1, \ldots, r\}$ , it is clear that

$$M^{-1}(A - \lambda_j \operatorname{Id}_k)^{l_j} M = \operatorname{diag}\{[(\lambda_1 - \lambda_j)\operatorname{Id}_{l_1} + N_1]^{l_j}, \dots, [(\lambda_r - \lambda_j)\operatorname{Id}_{l_r} + N_r]^{l_j}\}.$$

Note that the j'th block vanishes and that all the other blocks are invertible. Thus

$$\ker(A - \lambda_j \mathrm{Id}_k)^{l_j} = M(\{0\}^{l_1 + \dots + l_{j-1}} \times \mathbb{C}^{l_j} \times \{0\}^{l_{j+1} + \dots + l_r}).$$
(11.23)

Thus the generalised eigenspace associated with  $\lambda_j$  is given by the right hand side of (11.23). Let us relate this to  $\pi$ . Due to the fact that  $M^{-1}f(A)M = f(M^{-1}AM)$  for invertible matrices M, it is sufficient to consider the diagonal blocks on the right hand side of (11.22) separately. In fact,

$$M^{-1}\pi M = f(M^{-1}AM)$$
  
=  $\frac{1}{2\pi i} \int_{\gamma_a} \operatorname{diag}\{[(z - \lambda_1)\operatorname{Id}_{l_1} - N_1]^{-1}, \dots, [(z - \lambda_r)\operatorname{Id}_{l_r} - N_r]^{-1}\}dz.$ 

On the other hand,

$$[(z - \lambda_j) \mathrm{Id}_{l_j} - N_j]^{-1} = (z - \lambda_j)^{-1} \mathrm{Id}_{l_j} + (z - \lambda_j)^{-2} N_j + \dots + (z - \lambda_j)^{-l_j} N_j^{l_j - 1}.$$

Combining these observations with Theorem 11.9 yields

$$M^{-1}\pi M = \operatorname{diag}\{\operatorname{Ind}_{\gamma_a}(\lambda_1)\operatorname{Id}_{l_1},\ldots,\operatorname{Ind}_{\gamma_a}(\lambda_r)\operatorname{Id}_{l_r}\}.$$
(11.24)

Assume now that  $\lambda_j \in V_a$  and let  $x \in \ker(A - \lambda_j \operatorname{Id}_k)^{l_j}$ . Then  $\operatorname{Ind}_{\gamma_a}(\lambda_j) = 1$ . Moreover (11.23) implies that x = My, where  $y \in \{0\}^{l_1 + \dots + l_{j-1}} \times \mathbb{C}^{l_j} \times \{0\}^{l_{j+1} + \dots + l_r}$ . Combining these observations with (11.24) yields

$$\pi(x) = \pi(My) = My = x.$$

Similarly, if  $\lambda_j \notin V_a$  and  $x \in \ker(A - \lambda_j \operatorname{Id}_k)^{l_j}$ , then  $\pi(x) = 0$ . Thus  $\pi$  is the identity on the generalised eigenspaces corresponding to eigenvalues belonging to  $V_a$ . Moreover, it annihilates the generalised eigenspaces corresponding to eigenvalues not belonging to  $V_a$ . The lemma follows.  $\Box$ 

#### 11.3.3 Asymptotic partial diagonalisation

Given the above background, we are in a position to prove that there is, asymptotically, a partial diagonalisation of matrices of the form (11.7).

**Lemma 11.14.** Let  $1 \leq m \in \mathbb{Z}$ ,  $U, V, W \in \mathbf{M}_m(\mathbb{C})$  and define  $A_{\mu}$  by (11.7), where  $0 < \mu \in \mathbb{R}$ . Define  $\gamma_{\pm}$  by (11.9) and  $N_{\mu}$  by (11.10). Let  $\lambda_{j,\pm}$ ,  $j = 1, \ldots, p_{\pm}$ , denote the distinct eigenvalues of  $\gamma_{\pm}$ , and let  $m_{j,\pm}$  denote their respective algebraic multiplicities. Finally, let  $\gamma_{\pm,j}$ ,  $j = 1, \ldots, p_{\pm}$ , be the matrix collecting all the Jordan blocks in  $\gamma_{\pm}$  corresponding to the eigenvalue  $\lambda_{j,\pm}$ . Then there is a  $0 < \mu_0 \in \mathbb{R}$  and, for each  $\mu \geq \mu_0$ , a  $T_{\mu} \in \mathbf{GL}_{2m}(\mathbb{C})$  such that  $||T_{\mu}||, ||T_{\mu}^{-1}|| \leq 2$  and such that

$$T_{\mu}^{-1}N_{\mu}T_{\mu} = \operatorname{diag}\{N_{\mu,1}^{-}, \dots, N_{\mu,p-}^{-}, N_{\mu,1}^{+}, \dots, N_{\mu,p+}^{+}\}.$$
(11.25)

Moreover,

 $\|N_{\mu,j}^{\pm} \mp i\mu \operatorname{Id}_{m_{j,\pm}} - \gamma_{\pm,j}\| \le C\mu^{-1}.$ 

Finally, the constants  $\mu_0$  and C only depend on U, V and W.

**Remark 11.15.** Assuming that

 $\gamma_{\pm} = \operatorname{diag}\{\gamma_{\pm,1}, \ldots, \gamma_{\pm,p_{\pm}}\},\$ 

 $T_{\mu}$  can be chosen so that  $T_{\mu} \to \mathrm{Id}_{2m}$  as  $\mu \to \infty$ .

*Proof.* Given the notation introduced in Subsection 11.3.2, we are interested in the projections  $\pi_{\mu,j,\pm}$ , in particular their limits as  $\mu \to \infty$ . The reason is that these projections can be used to calculate the matrix  $T_{\mu}$ , the existence of which is the main conclusion of the lemma. Note that

$$\pi_{\mu,j,\pm} = \frac{1}{2\pi i} \int_{\gamma_{\mu,j,\pm}} (z \operatorname{Id}_{2m} - N_{\mu})^{-1} dz$$
$$= \frac{1}{2\pi i} \int_{\gamma} [(z + \mu_{j,\pm}) \operatorname{Id}_{2m} - N_{\mu}]^{-1} dz.$$

where  $\gamma(t) := re^{2\pi i t}$  and the range of j depends on the sign  $\pm$ . Let us introduce the notation

$$N_{\mu,j,\pm} := N_{\mu} - \mu_{j,\pm} \mathrm{Id}_{2m}$$

Then

$$N_{\mu,j,+} = \begin{pmatrix} -(2i\mu + \lambda_{j,+}) \mathrm{Id}_m + R_{\mu}^- & Q_{\mu}^+ \\ Q_{\mu}^- & -\lambda_{j,+} \mathrm{Id}_m + R_{\mu}^+ \end{pmatrix},$$
(11.26)

$$N_{\mu,l,-} = \begin{pmatrix} -\lambda_{l,-} \mathrm{Id}_m + R_{\mu}^- & Q_{\mu}^+ \\ Q_{\mu}^- & (2i\mu - \lambda_{l,-}) \mathrm{Id}_m + R_{\mu}^+ \end{pmatrix}.$$
 (11.27)

Moreover,

$$\pi_{\mu,j,\pm} = \frac{1}{2\pi i} \int_{\gamma} (z \mathrm{Id}_{2m} - N_{\mu,j,\pm})^{-1} dz, \qquad (11.28)$$

where the range of j is different depending on the sign  $\pm$ . Considering  $N_{\mu,l,-}$ , it is clear that the diagonal components of the bottom right block tend to infinity along the imaginary axis as  $\mu \to \infty$ . Moreover, the remaining components of the bottom right block remain bounded, as well as the remaining blocks. In the case of  $N_{\mu,j,+}$  the situation is similar. In this case it is the diagonal components of the top left block which tend to infinity along the imaginary axis. The same statements are true for  $z \operatorname{Id}_{2m} - N_{\mu,l,-}$  and  $z \operatorname{Id}_{2m} - N_{\mu,j,+}$  respectively. Let us introduce  $A_{\mu,j}^{\pm}(z)$  and  $D_{\mu,j}^{\pm}(z)$  by

$$z \mathrm{Id}_{2m} - N_{\mu,j,\pm} = \begin{pmatrix} A_{\mu,j}^{\pm}(z) & -Q_{\mu}^{+} \\ -Q_{\mu}^{-} & D_{\mu,j}^{\pm}(z) \end{pmatrix}, \qquad (11.29)$$

where the range of j depends on the sign  $\pm$ . Since |z| = r, it is clear that there is a  $0 < \mu_0 \in \mathbb{R}$  such that for  $\mu \ge \mu_0$ ,  $D^-_{\mu,l}(z)$  and  $A^+_{\mu,j}(z)$  are invertible and

$$\|[D_{\mu,l}^{-}(z)]^{-1}\| + \|[A_{\mu,j}^{+}(z)]^{-1}\| \le c_0 \mu^{-1},$$
(11.30)

where  $c_0$  is independent of  $\mu$  but depends on r (though the dependence on r can be eliminated by demanding that  $r \leq 1$ ). In the process of calculating the right hand side of (11.28), it is useful to note that

$$\begin{pmatrix} A_{\mu,l}^{-}(z) & -Q_{\mu}^{+} \\ -Q_{\mu}^{-} & D_{\mu,l}^{-}(z) \end{pmatrix}$$

$$= \begin{pmatrix} B_{\mu,l}^{-}(z) & -Q_{\mu}^{+} \\ 0 & D_{\mu,l}^{-}(z) \end{pmatrix} \begin{pmatrix} \mathrm{Id}_{m} & 0 \\ -[D_{\mu,l}^{-}(z)]^{-1}Q_{\mu}^{-} & \mathrm{Id}_{m} \end{pmatrix},$$

$$(11.31)$$

where

$$B_{\mu,l}^{-}(z) := A_{\mu,l}^{-}(z) - Q_{\mu}^{+} [D_{\mu,l}^{-}(z)]^{-1} Q_{\mu}^{-}.$$
(11.32)

Note that, by construction, the left (and thereby the right) hand side of (11.29) is invertible. Thus the left (and thereby the right) hand side of (11.31) is invertible. In particular, the first factor on the right hand side of (11.31) is invertible. Since this factor is a block diagonal matrix, the individual blocks are invertible. In particular,  $B_{\mu,l}^{-}(z)$  is invertible for the relevant range of z. In the end, we wish to compute the integral (along  $\gamma$ ) of the inverse of the left hand side of (11.31). Moreover, we wish to compute the matrix up to an error of the form  $O(\mu^{-2})$ . Introducing the notation

$$\begin{pmatrix} A_{\mu,l}^{-}(z) & -Q_{\mu}^{+} \\ -Q_{\mu}^{-} & D_{\mu,l}^{-}(z) \end{pmatrix}^{-1} =: \begin{pmatrix} E_{\mu,l}^{-}(z) & F_{\mu,l}^{-}(z) \\ G_{\mu,l}^{-}(z) & H_{\mu,l}^{-}(z) \end{pmatrix},$$
(11.33)

it can be computed that

$$E_{\mu,l}^{-}(z) = [B_{\mu,l}^{-}(z)]^{-1}, \qquad (11.34)$$

$$F_{\mu,l}^{-}(z) = [B_{\mu,l}^{-}(z)]^{-1} Q_{\mu}^{+} [D_{\mu,l}^{-}(z)]^{-1}, \qquad (11.35)$$

$$G_{\mu,l}^{-}(z) = [D_{\mu,l}^{-}(z)]^{-1} Q_{\mu}^{-} [B_{\mu,l}^{-}(z)]^{-1}, \qquad (11.36)$$

$$H_{\mu,l}^{-}(z) = [D_{\mu,l}^{-}(z)]^{-1} Q_{\mu}^{-} [B_{\mu,l}^{-}(z)]^{-1} Q_{\mu}^{+} [D_{\mu,l}^{-}(z)]^{-1} + [D_{\mu,l}^{-}(z)]^{-1}.$$
(11.37)

In what follows, we need a bound on  $||[B^-_{\mu,l}(z)]^{-1}||$ . Note, to this end, that

$$A^{-}_{\mu,l}(z) = (z + \lambda_{l,-}) \mathrm{Id}_m - \gamma_{-} - i\mu^{-1} W_{-}.$$

Combining this observation with (11.30) and (11.32) leads to the conclusion that

$$||B_{\mu,l}^{-}(z) - (z + \lambda_{l,-}) \mathrm{Id}_m + \gamma_{-}|| \le C\mu^{-1},$$

where C is independent of  $\mu$  but depends on r. Since  $(z + \lambda_{l,-}) \operatorname{Id}_m - \gamma_-$  is an upper triangular matrix, we only need to focus on its diagonal elements. They are of the form  $z + \lambda_{l,-} - \lambda_{j,-}$ . By construction, this expression is, in absolute value, bounded from below by a strictly positive constant. Note, however, that the lower bound depends on r. In particular, it depends on a lower bound on r (when taking l = j). To conclude, there is thus a uniform bound on  $\|[B_{\mu,l}^-(z)]^{-1}\|$ which holds for all  $\mu$  large enough.

Next, note that

$$[D_{\mu,l}^{-}(z)]^{-1} = (z - 2i\mu + \lambda_{l,-})^{-1} [\mathrm{Id}_m - (z - 2i\mu + \lambda_{l,-})^{-1} R_{\mu}^+]^{-1},$$
(11.38)

so that

$$\frac{1}{2\pi i} \int_{\gamma} [D^-_{\mu,l}(z)]^{-1} dz = 0$$

for large  $\mu$ . On the other hand, the norm of the first term on the right hand side of (11.37) is  $O(\mu^{-2})$ . Note, however, that the corresponding estimate depends on a lower bound on r (this is due to the fact that the upper bound on  $\|[B^-_{\mu,l}(z)]^{-1}\|$  depends on a lower bound on r). Thus

$$\left\|\frac{1}{2\pi i} \int_{\gamma} H^{-}_{\mu,l}(z) dz\right\| \le c_0 \mu^{-2}.$$
(11.39)

Before turning to (11.35) and (11.36), note that (11.38) yields

$$\left\| [D_{\mu,l}^{-}(z)]^{-1} - \frac{i}{2\mu} \mathrm{Id}_{m} \right\| \le c_{0}\mu^{-2}.$$
(11.40)

Thus

$$\left\|F_{\mu,l}^{-}(z) - \frac{i}{2\mu}[B_{\mu,l}^{-}(z)]^{-1}Q_{\mu}^{+}\right\| \le c_{0}\mu^{-2}.$$
(11.41)

There is a similar estimate concerning  $G^{-}_{\mu,l}(z)$ , so that it is sufficient to focus on the integral of  $[B^{-}_{\mu,l}(z)]^{-1}$  over  $\gamma$ . Due to (11.11), (11.27), (11.29), (11.32) and (11.40)

$$\left\| B_{\mu,l}^{-}(z) - (z + \lambda_{l,-}) \mathrm{Id}_{m} + \gamma_{-} + \mu^{-1} L_{\mu}^{-} \right\| \le c_{0} \mu^{-2},$$

where

$$L_{\mu}^{-} := iW_{-} + \frac{i}{2}Q_{\mu}^{+}Q_{\mu}^{-}.$$

Up to an error of order  $O(\mu^{-2})$ , the inverse of  $B^{-}_{\mu,l}(z)$  is thus given by

$$[\mathrm{Id}_m - [(z + \lambda_{l,-})\mathrm{Id}_m - \gamma_-]^{-1}\mu^{-1}L_{\mu}^{-}]^{-1}[(z + \lambda_{l,-})\mathrm{Id}_m - \gamma_-]^{-1}$$

Again, since we are only interested in computing  $[B^{-}_{\mu,l}(z)]^{-1}$  to order  $O(\mu^{-2})$ , it is sufficient to focus on

$$[(z+\lambda_{l,-})\mathrm{Id}_m-\gamma_{-}]^{-1}+[(z+\lambda_{l,-})\mathrm{Id}_m-\gamma_{-}]^{-1}\mu^{-1}L_{\mu}^{-}[(z+\lambda_{l,-})\mathrm{Id}_m-\gamma_{-}]^{-1}.$$
 (11.42)

Recall that  $\gamma_{-}$  is in Jordan normal form and compute

$$K_{\lambda}(z) := \begin{pmatrix} z - \lambda & -1 & 0 & \cdots & 0 \\ 0 & z - \lambda & -1 & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & z - \lambda \end{pmatrix}^{-1} \\ = \begin{pmatrix} (z - \lambda)^{-1} & (z - \lambda)^{-2} & (z - \lambda)^{-3} & \cdots & (z - \lambda)^{-l} \\ 0 & (z - \lambda)^{-1} & (z - \lambda)^{-2} & \cdots & (z - \lambda)^{-l+1} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & (z - \lambda)^{-1} \end{pmatrix},$$
(11.43)

where l is the dimension of the Jordan block. Assuming  $\lambda \notin \gamma^*$ , there are thus two possibilities: If  $\lambda \in B_r(0)$ ,

$$\frac{1}{2\pi i} \int_{\gamma} K_{\lambda}(z) dz = \mathrm{Id}_l.$$
(11.44)

If  $\lambda \notin \overline{B}_r(0)$ , then the integral vanishes. As a consequence,

$$\frac{1}{2\pi i} \int_{\gamma} [(z+\lambda_{l,-})\mathrm{Id}_m - \gamma_-]^{-1} dz = \Pi_{l,-},$$

where  $\Pi_{l,-}$  is the projection to the subspace of  $\mathbb{C}^m$  corresponding to the diagonal components of  $\gamma_-$  that equal  $\lambda_{l,-}$ . What we need to focus on is thus the second term on the right hand side of (11.42). Before proceeding, it is useful to introduce some notation. Recall that  $\lambda_{l,-}$ ,  $l = 1, \ldots, p_-$ , are the distinct eigenvalues of  $\gamma_-$ , and that  $m_{l,-}$  is the algebraic multiplicity of  $\lambda_{l,-}$ . We may thus assume, without loss of generality, that  $\gamma_- = \text{diag}\{\gamma_{-,11}, \ldots, \gamma_{-,p-p-}\}$ , where  $\gamma_{-,jj} \in \mathbf{M}_{m_{j,-}}(\mathbb{C})$ ,  $j = 1, \ldots, p_-$ . Moreover,  $\gamma_{-,jj}$  consists of all the Jordan blocks corresponding to the eigenvalue  $\lambda_{j,-}$ . If  $M \in \mathbf{M}_m(\mathbb{C})$  and  $r, s \in \{1, \ldots, p_-\}$ , we define  $M_{rs} \in \mathbf{M}_{m_{r,-} \times m_{s,-}}(\mathbb{C})$  by

$$M =: \begin{pmatrix} M_{11} & M_{12} & \cdots & M_{1p_{-}} \\ M_{21} & M_{22} & \cdots & M_{2p_{-}} \\ \vdots & \vdots & \vdots & \vdots \\ M_{p_{-}1} & M_{p_{-}2} & \cdots & M_{p_{-}p_{-}} \end{pmatrix}.$$

This notation is consistent with the notation  $\gamma_{-,jj}$ . Moreover,  $\gamma_{-,rs} = 0$  if  $r \neq s$ . Let  $\mathcal{K}$  be defined by

$$\mathcal{K} := \frac{1}{2\pi i} \int_{\gamma} [(z+\lambda_{l,-}) \mathrm{Id}_m - \gamma_-]^{-1} L^-_{\mu} [(z+\lambda_{l,-}) \mathrm{Id}_m - \gamma_-]^{-1} dz.$$

Then

$$\mathcal{K}_{rs} = \frac{1}{2\pi i} \int_{\gamma} [(z+\lambda_{l,-}) \mathrm{Id}_{m_{r,-}} - \gamma_{-,rr}]^{-1} L^{-}_{\mu,rs} [(z+\lambda_{l,-}) \mathrm{Id}_{m_{s,-}} - \gamma_{-,ss}]^{-1} dz.$$
(11.45)

If  $r \neq l$  and  $s \neq l$ , it is clear that the integrand appearing on the right hand side of (11.45) is analytic in  $B_r(0)$ , so that  $\mathcal{K}_{rs} = 0$ . If r = s = l, it is clear (keeping (11.43) in mind) that the components of the integrand on the right hand side of (11.45) are (finite) power series in  $z^{-1}$ , with the lowest power being 2. Combining this observation with Theorem 11.9 yields  $\mathcal{K}_{ll} = 0$ . The only non-zero components of  $\mathcal{K}$  are thus  $\mathcal{K}_{rl}$  and  $\mathcal{K}_{lr}$  for  $r \neq l$ . Summing up the above observations, it is clear that

$$\left\|\frac{1}{2\pi i} \int_{\gamma} [B_{\mu,l}^{-}(z)]^{-1} dz - \Pi_{l,-} - \mu^{-1} \mathcal{K}\right\| \le c_0 \mu^{-2}.$$
(11.46)

Combining this estimate with (11.41) and the analogous estimate for  $G^{-}_{\mu,l}(z)$  yields

$$\left\|\frac{1}{2\pi i} \int_{\gamma} F_{\mu,l}^{-}(z) dz - \frac{i}{2\mu} \Pi_{l,-} Q_{\mu}^{+}\right\| \le c_{0} \mu^{-2}, \tag{11.47}$$

$$\left\|\frac{1}{2\pi i} \int_{\gamma} G_{\mu,l}^{-}(z) dz - \frac{i}{2\mu} Q_{\mu}^{-} \Pi_{l,-}\right\| \le c_0 \mu^{-2}.$$
(11.48)

Combining (11.28), (11.29), (11.33), (11.34), (11.39), (11.46), (11.47) and (11.48)

$$\left\| \pi_{\mu,l,-} - \left( \begin{array}{cc} \Pi_{l,-} + \mu^{-1} \mathcal{K} & \frac{i}{2\mu} \Pi_{l,-} Q_{\mu}^{+} \\ \frac{i}{2\mu} Q_{\mu}^{-} \Pi_{l,-} & 0 \end{array} \right) \right\| \le c_{0} \mu^{-2}.$$

Let  $e_{l,k}^{-}$ ,  $k = 1, \ldots, m_{l,-}$ , be the standard basis for the image of  $\Pi_{l,-}$ . Then

$$\left| \pi_{\mu,l,-} \begin{pmatrix} e_{l,k}^{-} \\ 0 \end{pmatrix} - \begin{pmatrix} e_{l,k}^{-} + \mu^{-1} \mathcal{K} e_{l,k}^{-} \\ \frac{i}{2\mu} Q_{\mu}^{-} e_{l,k}^{-} \end{pmatrix} \right\| \le c_0 \mu^{-2}.$$

Thus

$$\hat{e}_{l,k}^{-} := \pi_{\mu,l,-} \left( \begin{array}{c} e_{l,k}^{-} \\ 0 \end{array} \right)$$

is a basis for  $E_{\mu,l,-}$  for  $\mu$  large enough. Since we know that  $N_{\mu}$  leaves  $E_{\mu,l,-}$  invariant, there is a matrix  $N_{\mu,l}^{-} \in \mathbf{M}_{m_{l,-}}(\mathbb{C})$  such that

$$N_{\mu}\hat{e}_{l,r}^{-} = \sum_{s=1}^{m_{l,-}} N_{\mu,l,sr}^{-}\hat{e}_{l,s}^{-}.$$
(11.49)

We wish to compute  $N_{\mu,l}^-$ . Consider, to this end,

$$\begin{split} N_{\mu}\hat{e}_{l,k}^{-} &= \begin{pmatrix} -i\mu \mathrm{Id}_{m} + R_{\mu}^{-} & Q_{\mu}^{+} \\ Q_{\mu}^{-} & i\mu \mathrm{Id}_{m} + R_{\mu}^{+} \end{pmatrix} \begin{pmatrix} e_{l,k}^{-} + \mu^{-1}\mathcal{K}e_{l,k}^{-} \\ \frac{i}{2\mu}Q_{\mu}^{-}e_{l,k}^{-} \end{pmatrix} + O(\mu^{-1}) \\ &= \begin{pmatrix} -i\mu e_{l,k}^{-} + \gamma_{-}e_{l,k}^{-} - i\mathcal{K}e_{l,k}^{-} \\ \frac{1}{2}Q_{\mu}^{-}e_{l,k}^{-} \end{pmatrix} + O(\mu^{-1}). \end{split}$$

Let us take the scalar product of this equality with  $\hat{e}_{l,r}^-$  (when we say scalar product, we here mean the ordinary scalar product and not the Hermitian inner product). This yields

$$(N_{\mu}\hat{e}_{l,k}^{-}) \cdot \hat{e}_{l,r}^{-} = (-i\mu e_{l,k}^{-} + \gamma_{-} e_{l,k}^{-} - i\mathcal{K}e_{l,k}^{-}) \cdot (e_{l,r}^{-} + \mu^{-1}\mathcal{K}e_{l,r}^{-}) + O(\mu^{-1})$$

$$= -i\mu\delta_{kr} + (\gamma_{-}e_{l,k}^{-}) \cdot e_{l,r}^{-} + O(\mu^{-1}),$$

$$(11.50)$$

where we have used the fact that  $e_{l,r}^- \cdot (\mathcal{K}e_{l,k}^-) = 0$ . On the other hand,

$$\hat{e}_{l,r}^{-} \cdot \hat{e}_{l,s}^{-} = \delta_{rs} + O(\mu^{-2}), \qquad (11.51)$$

where we have, again, used the fact that  $e_{l,r}^- \cdot (\mathcal{K}e_{l,k}^-) = 0$ . Combining (11.49), (11.50) and (11.51) yields

$$N^{-}_{\mu,l,rs} = -i\mu\delta_{rs} + (\gamma_{-}e^{-}_{l,s}) \cdot e^{-}_{l,r} + O(\mu^{-1}).$$

There is a similar analysis concerning  $E_{\mu,j,+}$  (which we provide below), and as a consequence, there is  $0 < \mu_0 \in \mathbb{R}$ , and, for  $\mu \ge \mu_0$ , a matrix  $T_{\mu}$  (which converges to  $\mathrm{Id}_{2m}$  as  $\mu \to \infty$ ) such that (11.25) holds. The lemma follows. Analysis in the case of  $E_{\mu,j,+}$ . The analysis in the case of  $E_{\mu,j,+}$  is similar, but we include it for the sake of completeness. In the process of calculating the right hand side of (11.28), it is useful to note that

$$\begin{pmatrix} A_{\mu,l}^{+}(z) & -Q_{\mu}^{+} \\ -Q_{\mu}^{-} & D_{\mu,l}^{+}(z) \end{pmatrix} = \begin{pmatrix} A_{\mu,l}^{+}(z) & 0 \\ -Q_{\mu}^{-} & B_{\mu,l}^{+}(z) \end{pmatrix} \begin{pmatrix} \mathrm{Id}_{m} & -[A_{\mu,l}^{+}(z)]^{-1}Q_{\mu}^{+} \\ 0 & \mathrm{Id}_{m} \end{pmatrix},$$
(11.52)

where

$$B_{\mu,l}^+(z) := D_{\mu,l}^+(z) - Q_{\mu}^-[A_{\mu,l}^+(z)]^{-1}Q_{\mu}^+.$$
(11.53)

Introducing the notation

$$\begin{pmatrix} A_{\mu,l}^+(z) & -Q_{\mu}^+ \\ -Q_{\mu}^- & D_{\mu,l}^+(z) \end{pmatrix}^{-1} =: \begin{pmatrix} E_{\mu,l}^+(z) & F_{\mu,l}^+(z) \\ G_{\mu,l}^+(z) & H_{\mu,l}^+(z) \end{pmatrix},$$
(11.54)

it can be computed that

$$E_{\mu,l}^{+}(z) = [A_{\mu,l}^{+}(z)]^{-1}Q_{\mu}^{+}[B_{\mu,l}^{+}(z)]^{-1}Q_{\mu}^{-}[A_{\mu,l}^{+}(z)]^{-1} + [A_{\mu,l}^{+}(z)]^{-1}, \qquad (11.55)$$

$$F_{\mu,l}(z) = [A_{\mu,l}(z)]^{-1} Q_{\mu} [B_{\mu,l}(z)]^{-1}, \qquad (11.56)$$

$$G_{\mu,l}^+(z) = [B_{\mu,l}^+(z)]^{-1} Q_{\mu}^-[A_{\mu,l}^+(z)]^{-1}, \qquad (11.57)$$

$$H_{\mu,l}^+(z) = [B_{\mu,l}^+(z)]^{-1}.$$
(11.58)

Note that

$$[A_{\mu,l}^+(z)]^{-1} = (z + 2i\mu + \lambda_{l,+})^{-1} [\mathrm{Id}_m - (z + 2i\mu + \lambda_{l,+})^{-1} R_{\mu}^-]^{-1},$$
(11.59)

so that

$$\frac{1}{2\pi i} \int_{\gamma} [A^+_{\mu,l}(z)]^{-1} dz = 0$$

for large  $\mu$ . On the other hand, the norm of the first term on the right hand side of (11.55) is  $O(\mu^{-2})$ . Thus

$$\left\|\frac{1}{2\pi i} \int_{\gamma} E_{\mu,l}^{+}(z) dz\right\| \le c_0 \mu^{-2}.$$
(11.60)

Before turning to (11.56) and (11.57), note that (11.59) yields

$$\left\| [A_{\mu,l}^+(z)]^{-1} + \frac{i}{2\mu} \mathrm{Id}_m \right\| \le c_0 \mu^{-2}.$$
(11.61)

Thus

$$\left\|F_{\mu,l}^{+}(z) + \frac{i}{2\mu}Q_{\mu}^{+}[B_{\mu,l}^{+}(z)]^{-1}\right\| \le c_{0}\mu^{-2}.$$
(11.62)

There is a similar estimate concerning  $G^+_{\mu,l}(z)$ , so that it is sufficient to focus on the integral of  $[B^+_{\mu,l}(z)]^{-1}$  over  $\gamma$ . Due to (11.11), (11.26), (11.29), (11.53) and (11.61)

$$\left\| B_{\mu,l}^{+}(z) - (z + \lambda_{l,+}) \mathrm{Id}_{m} + \gamma_{+} - \mu^{-1} L_{\mu}^{+} \right\| \le c_{0} \mu^{-2},$$

where

$$L^+_{\mu} := iW_+ + \frac{i}{2}Q^-_{\mu}Q^+_{\mu}.$$

Up to an error of order  $O(\mu^{-2})$ , the inverse of  $B^+_{\mu,l}(z)$  is thus given by

$$[\mathrm{Id}_m + [(z + \lambda_{l,+})\mathrm{Id}_m - \gamma_+]^{-1} \mu^{-1} L_{\mu}^+]^{-1} [(z + \lambda_{l,+})\mathrm{Id}_m - \gamma_+]^{-1}.$$

Again, since we are only interested in computing  $[B^+_{\mu,l}(z)]^{-1}$  to order  $O(\mu^{-2})$ , it is sufficient to focus on

$$[(z+\lambda_{l,+})\mathrm{Id}_m - \gamma_+]^{-1} - [(z+\lambda_{l,+})\mathrm{Id}_m - \gamma_+]^{-1}\mu^{-1}L^+_{\mu}[(z+\lambda_{l,+})\mathrm{Id}_m - \gamma_+]^{-1}.$$
 (11.63)

Due to (11.43) and (11.44),

$$\frac{1}{2\pi i} \int_{\gamma} [(z+\lambda_{l,+}) \mathrm{Id}_m - \gamma_+]^{-1} dz = \Pi_{l,+},$$

where  $\Pi_{l,+}$  is the projection to the subspace of  $\mathbb{C}^m$  corresponding to the diagonal components of  $\gamma_+$  that equal  $\lambda_{l,+}$ . What we need to focus on is thus the second term appearing in (11.63). Before proceeding, it is useful to introduce some notation. Recall that  $\lambda_{l,+}$ ,  $l = 1, \ldots, p_+$ , are the distinct eigenvalues of  $\gamma_+$ , and that  $m_{l,+}$  is the algebraic multiplicity of  $\lambda_{l,+}$ . We may thus assume, without loss of generality, that  $\gamma_+ = \text{diag}\{\gamma_{+,11}, \ldots, \gamma_{+,p_+p_+}\}$ , where  $\gamma_{+,jj} \in \mathbf{M}_{m_{j,+}}(\mathbb{C})$ ,  $j = 1, \ldots, p_+$ . Moreover,  $\gamma_{+,jj}$  consists of all the Jordan blocks corresponding to the eigenvalue  $\lambda_{j,+}$ . If  $M \in \mathbf{M}_m(\mathbb{C})$  and  $r, s \in \{1, \ldots, p_+\}$ , we define  $M_{rs} \in \mathbf{M}_{m_{r,+} \times m_{s,+}}(\mathbb{C})$  by

$$M =: \begin{pmatrix} M_{11} & M_{12} & \cdots & M_{1p_+} \\ M_{21} & M_{22} & \cdots & M_{2p_+} \\ \vdots & \vdots & \vdots & \vdots \\ M_{p_+1} & M_{p_+2} & \cdots & M_{p_+p_+} \end{pmatrix}$$

This notation is consistent with the notation  $\gamma_{+,jj}$ . Moreover,  $\gamma_{+,rs} = 0$  if  $r \neq s$ . Let  $\mathcal{L}$  be defined by

$$\mathcal{L} := \frac{1}{2\pi i} \int_{\gamma} [(z+\lambda_{l,+}) \mathrm{Id}_m - \gamma_+]^{-1} L^+_{\mu} [(z+\lambda_{l,+}) \mathrm{Id}_m - \gamma_+]^{-1} dz$$

Then

$$\mathcal{L}_{rs} = \frac{1}{2\pi i} \int_{\gamma} [(z + \lambda_{l,+}) \mathrm{Id}_{m_{r,+}} - \gamma_{+,rr}]^{-1} L^{+}_{\mu,rs} [(z + \lambda_{l,+}) \mathrm{Id}_{m_{s,+}} - \gamma_{+,ss}]^{-1} dz.$$
(11.64)

If  $r \neq l$  and  $s \neq l$ , it is clear that the integrand appearing on the right hand side of (11.45) is analytic in  $B_r(0)$ , so that  $\mathcal{L}_{rs} = 0$ . If r = s = l, it is clear (keeping (11.43) in mind) that the components of the integrand on the right hand side of (11.45) are (finite) power series in  $z^{-1}$ , with the lowest power being 2. Combining this observation with Theorem 11.9 yields  $\mathcal{L}_{ll} = 0$ . The only non-zero components of  $\mathcal{L}$  are thus  $\mathcal{L}_{rl}$  and  $\mathcal{L}_{lr}$  for  $r \neq l$ . Summing up the above observations, it is clear that

$$\left\|\frac{1}{2\pi i}\int_{\gamma} [B_{\mu,l}^+(z)]^{-1}dz - \Pi_{l,+} + \mu^{-1}\mathcal{L}\right\| \le c_0\mu^{-2}.$$
(11.65)

Combining this estimate with (11.62) and the analogous estimate for  $G^+_{\mu l}(z)$  yields

$$\left\|\frac{1}{2\pi i}\int_{\gamma}F_{\mu,l}^{+}(z)dz + \frac{i}{2\mu}Q_{\mu}^{+}\Pi_{l,+}\right\| \leq c_{0}\mu^{-2},$$
(11.66)

$$\left\|\frac{1}{2\pi i} \int_{\gamma} G^{+}_{\mu,l}(z) dz + \frac{i}{2\mu} \Pi_{l,+} Q^{-}_{\mu}\right\| \le c_0 \mu^{-2}.$$
(11.67)

Combining (11.28), (11.29), (11.54), (11.58), (11.60), (11.65), (11.66) and (11.67)

$$\left\| \pi_{\mu,l,+} - \left( \begin{array}{cc} 0 & -\frac{i}{2\mu} Q_{\mu}^{+} \Pi_{l,+} \\ -\frac{i}{2\mu} \Pi_{l,+} Q_{\mu}^{-} & \Pi_{l,+} - \mu^{-1} \mathcal{L} \end{array} \right) \right\| \le c_{0} \mu^{-2}.$$

Let  $e_{l,k}^+$ ,  $k = 1, \ldots, m_{l,+}$ , be the standard basis for the image of  $\Pi_{l,+}$ . Then

$$\left\| \pi_{\mu,l,+} \begin{pmatrix} 0 \\ e_{l,k}^+ \end{pmatrix} - \begin{pmatrix} -\frac{i}{2\mu} Q_{\mu}^+ e_{l,k}^+ \\ e_{l,k}^+ - \mu^{-1} \mathcal{L} e_{l,k}^+ \end{pmatrix} \right\| \le c_0 \mu^{-2}.$$

Thus

$$\hat{e}_{l,k}^+ := \pi_{\mu,l,+} \left( \begin{array}{c} e_{l,k}^+ \\ 0 \end{array} \right)$$

is a basis for  $E_{\mu,l,+}$  for  $\mu$  large enough. Since we know that  $N_{\mu}$  leaves  $E_{\mu,l,+}$  invariant, there is a matrix  $N_{\mu,l}^+ \in \mathbf{M}_{m_{l,+}}(\mathbb{C})$  such that

$$N_{\mu}\hat{e}^{+}_{l,r} = \sum_{s=1}^{m_{l,+}} N^{+}_{\mu,l,sr}\hat{e}^{+}_{l,s}.$$
(11.68)

We wish to compute  $N_{\mu,l}^+$ . Consider, to this end,

$$\begin{split} N_{\mu} \hat{e}^{+}_{l,k} &= \begin{pmatrix} -i\mu \mathrm{Id}_{m} + R^{-}_{\mu} & Q^{+}_{\mu} \\ Q^{-}_{\mu} & i\mu \mathrm{Id}_{m} + R^{+}_{\mu} \end{pmatrix} \begin{pmatrix} -\frac{i}{2\mu} Q^{+}_{\mu} e^{+}_{l,k} \\ e^{+}_{l,k} - \mu^{-1} \mathcal{L} e^{+}_{l,k} \end{pmatrix} + O(\mu^{-1}) \\ &= \begin{pmatrix} \frac{1}{2} Q^{+}_{\mu} e^{+}_{l,k} \\ i\mu e^{+}_{l,k} + \gamma_{+} e^{+}_{l,k} - i\mathcal{L} e^{+}_{l,k} \end{pmatrix} + O(\mu^{-1}). \end{split}$$

Let us take the scalar product of this equality with  $\hat{e}_{l,r}^+$ . This yields

$$(N_{\mu}\hat{e}^{+}_{l,k}) \cdot \hat{e}^{+}_{l,r} = (i\mu e^{+}_{l,k} + \gamma_{+} e^{+}_{l,k} - i\mathcal{L}e^{+}_{l,k}) \cdot (e^{+}_{l,r} - \mu^{-1}\mathcal{L}e^{+}_{l,r}) + O(\mu^{-1})$$

$$= i\mu\delta_{kr} + (\gamma_{+}e^{+}_{l,k}) \cdot e^{+}_{l,r} + O(\mu^{-1}),$$

$$(11.69)$$

where we have used the fact that  $e_{l,r}^+ \cdot (\mathcal{L}e_{l,k}^+) = 0$ . On the other hand,

$$\hat{e}_{l,r}^{+} \cdot \hat{e}_{l,s}^{+} = \delta_{rs} + O(\mu^{-2}), \qquad (11.70)$$

where we have, again, used the fact that  $e_{l,r}^+ \cdot (\mathcal{L}e_{l,k}^+) = 0$ . Combining (11.68), (11.69) and (11.70) yields

$$N_{\mu,l,rs}^{+} = i\mu\delta_{rs} + (\gamma_{+}e_{l,s}^{+}) \cdot e_{l,r}^{+} + O(\mu^{-1}).$$

This completes the analysis.

# 11.4 Mode by mode analysis

Consider the index set  $\mathcal{I}_{\mathrm{B}}$  introduced in connection with (24.3) and fix  $1 \leq m \in \mathbb{Z}$ . Assume that for each  $\iota \in \mathcal{I}_{\mathrm{B}}$ , there is a matrix  $A(\iota) \in \mathbf{M}_{2m}(\mathbb{C})$  and functions  $A_{\iota,\mathrm{rem}} \in C^{\infty}(\mathbb{R}, \mathbf{M}_{2m}(\mathbb{C}))$  and  $F_{\iota} \in C^{\infty}(\mathbb{R}, \mathbb{C}^{2m})$ . We are then interested in the equation

$$\dot{v} = A(\iota)v + A_{\iota,\text{rem}}v + F_{\iota}.$$
(11.71)

Here the matrix valued function  $A_{\iota,\text{rem}}$  is assumed to satisfy the estimate

$$\|A_{\iota,\mathrm{rem}}(t)\| \le C_{\mathrm{ts}} e^{-\beta_{\mathrm{ts}}\bar{t}} \tag{11.72}$$

for  $t \geq T_{\rm ts}$ , where  $0 < C_{\rm ts}$ ,  $\beta_{\rm ts} \in \mathbb{R}$  are independent of  $\iota$ ;  $T_{\rm ts}$  depends on  $\iota$ ; and  $\bar{t} := t - T_{\rm ts}$ . In order to restrict the possible matrices  $A(\iota)$ , we think of (11.71) as arising from an equation of the form (1.2) satisfying one of Definitions 11.1 and 11.3. In the present setting, it is convenient to make assumptions that follow from these definitions without referring to them or to (1.2). One way of doing this is by introducing a function  $\mu : \mathcal{I}_{\rm B} \to [0, \infty)$  with the property that if  $\mathcal{M} := \mu(\mathcal{I}_{\rm B})$ , then the only accumulation point of  $\mathcal{M}$  is  $\infty$ . Note that if (1.2) satisfies one of Definitions 11.1 and 11.3, and if  $\mu(\iota) := \mathfrak{g}_{\infty}(\iota)$ , then this requirement is satisfied. Given the function  $\mu$ , we restrict our attention to matrices  $A(\iota)$  of the form

$$A(\iota) := A_{\mu(\iota)} = \begin{pmatrix} 0 & \mu(\iota) \mathrm{Id}_m \\ -\mu(\iota) \mathrm{Id}_m + iV + 2[\mu(\iota)]^{-1}W & U \end{pmatrix},$$
(11.73)

where  $V, W, U \in \mathbf{M}_m(\mathbb{C})$  and we have used the notation (11.7). Below we frequently omit reference to the argument  $\iota$  and simply write  $\mu$  instead of  $\mu(\iota)$ . When we do,  $\mu^{-1}$  should be interpreted as  $[\mu(\iota)]^{-1}$ . For a fixed  $\mu_a \in \mathcal{M}$ , we can analyse the asymptotics of solutions to (11.71) for  $\iota \in \mu^{-1}(\mu_a)$ . In fact, we can appeal to the results obtained in Sections 9.3 and 9.4. However, the constants appearing in the results depend on  $A(\iota) = A_{\mu_a}$ . Appealing to the analysis in these sections is thus acceptable for a finite number of  $\mu$ -values. In the present setting, this is not sufficient, and we need to proceed differently. Considering the analysis carried out in Sections 11.2 and 11.3, it is clear that there is an asymptotic partial diagonalisation of the matrices  $A_{\mu}$  as  $\mu \to \infty$ . Moreover, the real parts of the eigenvalues of  $A_{\mu}$  converge to specific values as  $\mu \to \infty$ . Since the only accumulation point of  $\mathcal{M}$  is  $\infty$ , this leads to the following approach to obtaining estimates for all  $\iota \in \mathcal{I}_{B}$ . First, we appeal to the asymptotic partial diagonalisation in order to obtain estimates for  $\mu$  large enough, say  $\mu \geq \mu_0$ . Second, we note that the remaining  $\mu$ -values are finite in number. Thus we can handle them by appealing to Sections 9.3 and 9.4. By increasing  $\mu_0$ , the asymptotic analysis becomes more and more refined. However, there is a price to pay for this refinement: the supremum of the constants involved in the estimates for the remaining  $\mu$ values can be expected to tend to infinity as  $\mu_0$  tends to infinity. What is preferable might depend on the context. For that reason, it is natural to introduce a parameter  $0 < \beta_{\text{mar}} \in \mathbb{R}$  which quantifies the quality of the asymptotic analysis; a small  $\beta_{mar}$ -value corresponds to good control in the asymptotic region where  $\mu$  is large. However, the constants appearing in the estimates can be expected to tend to infinity as  $\beta_{\text{mar}} \rightarrow 0+$ .

Just as in the case of Lemma 9.16, we need to make assumptions concerning the integrability properties of  $F_{\iota}$  in order to obtain detailed information concerning the asymptotics of solutions to (11.71). In the next lemma, we therefore assume that the following norm of  $F_{\iota}$  is bounded:

$$\|F_{\iota}\|_{\text{lma}} := \int_{0}^{\infty} e^{-(\kappa_{\iota} - \beta_{\text{mar}})s} |F_{\iota}(s)| ds, \qquad (11.74)$$

where  $\kappa_{\iota} := \kappa_{\max}[A(\iota)]$  and the choice of  $\beta_{\max} > 0$  should be clear from the context. Note that we abuse notation slightly here, in that the norm depends on  $\iota$  without this being explicitly stated.

In Lemma 9.16, the first generalised eigenspace in the  $\beta_{\text{rem}}$ , A-decomposition of  $\mathbb{C}^{2m}$  played an important role. Similar algebraic constructions can be expected to be important in the present setting as well. However, since we need to have uniform (i.e., independent of  $\iota$ ) bounds on the norms of the matrices, say  $T_{\iota}$  and  $T_{\iota}^{-1}$ , putting  $A(\iota)$  into Jordan normal form, in order for the  $\beta_{\text{rem}}$ ,  $A(\iota)$ decomposition of  $\mathbb{C}^{2m}$  to be of use, and since such such uniform bounds are not to be expected in the present setting, the algebraic constructions we use here have to be somewhat more rough. On the other hand, we can appeal to the partial diagonalisation derived in Subsection 11.3.3. This leads to the following asymptotic substitute for the notion of a first generalised eigenspace in the  $\beta_{\text{rem}}$ , A-decomposition of  $\mathbb{C}^{2m}$ .

**Definition 11.16.** Let  $1 \leq m \in \mathbb{Z}$ ,  $\mathcal{I}_{\mathrm{B}}$  be the set introduced in connection with (24.3) and  $\mu$  be a function  $\mu : \mathcal{I}_{\mathrm{B}} \to [0, \infty)$ . Let  $U, V, W \in \mathbf{M}_m(\mathbb{C})$  and define  $A(\iota)$  by (11.73). Let  $\lambda_{j,\pm}$ ,  $j = 1, \ldots, p_{\pm}$ , be the eigenvalues of  $U_{\pm} = (U \pm V)/2$ , and let  $\kappa_{\infty}$  be the largest real part of an element of  $\mathrm{Sp}(U_+) \cup \mathrm{Sp}(U_-)$ . Let  $0 < r_a, \mu_a \in \mathbb{R}$  (depending only on U, V and W) be such that if  $\mu \geq \mu_a$ , then the closed balls with radius  $r_a$  and centre  $\pm i\mu + \lambda_{j,\pm}$  are disjoint and a distance at least  $r_a$  apart. Due to Lemma 11.6, there is a  $\mu_0 \geq \mu_a$  such that if  $\mu(\iota) \geq \mu_0$ , then the set  $\mathrm{Sp}[A(\iota)]$  is contained in the disjoint union of the open balls of radius  $r_a$  and centre  $\pm i\mu(\iota) + \lambda_{j,\pm}$ . Given a  $0 < \beta \in \mathbb{R}$  and a  $\iota \in \mathcal{I}_{\mathrm{B}}$  such that  $\mu(\iota) \geq \mu_0$ , the vector space  $E_{\iota,\beta}^{\infty}$  is defined to be the direct sum of the generalised eigenspaces of  $A(\iota)$  corresponding to eigenvalues in the balls of radius  $r_a$  and centre  $\pm i\mu(\iota) + \lambda_{j,\pm}$  for j such that  $\kappa_{\infty} - \mathrm{Re}(\lambda_{j,\pm}) < \beta$ .

**Lemma 11.17.** Let  $1 \leq m \in \mathbb{Z}$ ,  $\mathcal{I}_{\rm B}$  be the set introduced in connection with (24.3) and  $\mu$  be a function  $\mu : \mathcal{I}_{\rm B} \to [0, \infty)$ . Let  $\mathcal{I}_{\rm B,ts} \subseteq \mathcal{I}_{\rm B}$  be the set of  $\iota \in \mathcal{I}_{\rm B}$  such that  $\mu(\iota) > 0$ . Let  $U, V, W \in \mathbf{M}_m(\mathbb{C})$  and define  $A(\iota)$  by (11.73) for  $\iota \in \mathcal{I}_{\rm B,ts}$ . Assume that for each  $\iota \in \mathcal{I}_{\rm B,ts}$ , there are functions  $A_{\iota,\text{rem}} \in C^{\infty}(\mathbb{R}, \mathbf{M}_{2m}(\mathbb{C}))$  and  $F_{\iota} \in C^{\infty}(\mathbb{R}, \mathbb{C}^{2m})$ . The function  $A_{\iota,\text{rem}}$  is assumed to be such that there are constants  $0 < \beta_{\text{ts}}, C_{\text{ts}} \in \mathbb{R}$  (independent of  $\iota$ ) and a constant  $T_{\text{ts}}$  (which
is allowed to depend on  $\iota$ ) such that (11.72) holds for  $t \geq T_{ts}$ . Fix a  $0 < \beta_{mar} \in \mathbb{R}$  which is small enough, the bound depending only on U, V and  $\beta_{ts}$ . Assume the functions  $F_{\iota}$  to be such that  $\|F_{\iota}\|_{lma} < \infty$  for  $\iota \in \mathcal{I}_{B,ts}$ ; cf. (11.74). Then there are constants  $0 < \mu_0 \in \mathbb{R}$  and  $0 < C \in \mathbb{R}$ , where  $\mu_0$  only depends on U, V, W and  $\beta_{mar}$ , and C only depends on U, V,  $\beta_{ts}$ ,  $\beta_{mar}$  and  $C_{ts}$ , such that the following holds. If  $\iota \in \mu^{-1}([\mu_0, \infty))$  and v is a solution to (11.71), then there is a  $v_{\infty} \in E_{\iota,\beta_{ts}}^{\infty}$ such that

$$\left| v(t) - e^{A(\iota)t} v_{\infty} - \int_{T_{\rm ts}}^{t} e^{A(\iota)(t-s)} F_{\iota}(s) ds \right|$$

$$\leq C e^{(\kappa_{\iota} - \beta_{\rm ts} + \beta_{\rm mar})\bar{t}} [|v(T_{\rm ts})| + e^{\kappa_{\iota} T_{\rm ts}} \|F_{\iota}\|_{\rm Ima}]$$

$$(11.75)$$

for  $t \geq T_{ts}$ , where  $E_{\iota,\beta_{ts}}^{\infty}$  is given by Definition 11.16 and  $\kappa_{\iota} := \kappa_{\max}[A(\iota)]$ ; cf. Definition 4.3. Moreover,

$$v_{\infty} := e^{-A(\iota)T_{\rm ts}} u_{\infty} \tag{11.76}$$

for a  $u_{\infty} \in E^{\infty}_{\iota,\beta_{ts}}$  satisfying the estimate

$$|u_{\infty}| \le C[|v(T_{\rm ts})| + e^{\kappa_{\iota} T_{\rm ts}} ||F_{\iota}||_{\rm Ima}].$$
(11.77)

Here C has the same dependence as in the case of (11.75).

**Remark 11.18.** Note that if  $A_{\mu}$  is given by (11.7), then  $\kappa_{\max}(A_{\mu}) \to \kappa_{\infty}$  as  $\mu \to \infty$ , where  $\kappa_{\infty}$  is introduced in Definition 11.16; this is a consequence of Lemma 11.6. In fact, we could replace  $\kappa_{\iota}$  by  $\kappa_{\infty}$  in the statement of the lemma.

*Proof.* It is convenient to divide the proof into several steps. To begin with we derive several reformulations of the equations.

First reformulation of the equations; partial diagonalisation. Due to (11.10) and Lemma 11.14, we know that there is a  $0 < \mu_0 \in \mathbb{R}$  such that if  $\mu(\iota) \ge \mu_0$ , then (11.71) can be reformulated to

$$\dot{v}_{\rm pre} = \bar{A}(\iota)v_{\rm pre} + \bar{A}_{\iota,\rm rem}v_{\rm pre} + \bar{F}_{\iota},$$

where  $T_{\iota} := S_{12}T_{\mu(\iota)}$  and

$$v_{\rm pre} := T_{\iota}^{-1} v, \quad \bar{A}(\iota) := T_{\iota}^{-1} A(\iota) T_{\iota}, \quad \bar{A}_{\iota,\rm rem} := T_{\iota}^{-1} A_{\iota,\rm rem} T_{\iota}, \quad \bar{F}_{\iota} := T_{\iota}^{-1} F_{\iota}.$$
(11.78)

Note that  $||T_{\iota}||$  and  $||T_{\iota}^{-1}||$  are bounded by numerical constants. In addition,  $\bar{A}(\iota)$  is given by the right hand side of (11.25) with  $\mu$  replaced by  $\mu(\iota)$ . Moreover,

$$N_{\mu(\iota),j}^{\pm} = \pm i\mu(\iota) \mathrm{Id}_{m_{j,\pm}} + \gamma_{\pm,j} + \mathcal{R}_{\iota,j}^{\pm}, \qquad (11.79)$$

where

$$\|\mathcal{R}_{\iota,j}^{\pm}\| \le c_0 [\mu(\iota)]^{-1} \tag{11.80}$$

for  $\mu(\iota) \ge \mu_0$ , where  $c_0$  and  $\mu_0$  only depend on U, V and W. Letting  $M_{\iota,j}^{\pm}$  denote the sum of the first two terms on the right hand side of (11.79), let

$$M_{\iota} := \operatorname{diag}\{M_{\iota,1}^{-}, \dots, M_{\iota,p_{-}}^{-}, M_{\iota,1}^{+}, \dots, M_{\iota,p_{+}}^{+}\}.$$

Second reformulation of the equations; eliminating the imaginary parts. Note that the imaginary part of  $M_{\iota,j}^{\pm}$  is a multiple of the identity, so that the imaginary part of  $M_{\iota}$  commutes with  $\bar{A}(\iota)$ . Introducing  $M_{\iota,\text{im}} := \text{Im}(M_{\iota})$  and

$$v_{\rm fin} := \exp\left(-iM_{\iota,\rm im}\bar{t}\right)v_{\rm pre}$$

thus yields

$$\dot{v}_{\rm fin} = \tilde{A}(\iota) v_{\rm fin} + \tilde{A}_{\iota,\rm rem} v_{\rm fin} + \tilde{F}_{\iota},$$

where

$$\bar{A}_{\iota,\mathrm{rem}} := \exp\left(-iM_{\iota,\mathrm{im}}\bar{t}\right)\bar{A}_{\iota,\mathrm{rem}}\exp\left(iM_{\iota,\mathrm{im}}\bar{t}\right), \quad \bar{F}_{\iota} := \exp\left(-iM_{\iota,\mathrm{im}}\bar{t}\right)\bar{F}_{\iota}.$$

Moreover,

$$\tilde{A}(\iota) = \tilde{J} + \tilde{\mathcal{R}}_{\iota}, \quad \tilde{J} := \operatorname{diag}\{\tilde{J}_1, \dots, \tilde{J}_K\}, \quad \tilde{\mathcal{R}}_{\iota} := \operatorname{diag}\{\tilde{\mathcal{R}}_{\iota,1}, \dots, \tilde{\mathcal{R}}_{\iota,K}\},$$

where dim  $\tilde{\mathcal{R}}_{\iota,k}$  = dim  $\tilde{J}_k$  and the  $\tilde{\mathcal{R}}_{\iota,k}$  satisfy estimates similar to (11.80). Finally, the  $\tilde{J}_k$  are the real parts of the  $\gamma_{\pm,j}$ .

Third reformulation of the equations; normalising the real parts. Let  $\kappa_{\infty}$  be the constant introduced in Definition 11.16. Note that this means that  $\kappa_{\infty}$  is the largest real part of an eigenvalue of  $\gamma_{\pm}$ . Define

$$w := e^{-\kappa_{\infty}t} D v_{\text{fin}},$$

where D is a real non-degenerate diagonal matrix that remains to be determined. Then

$$\dot{w} = B(\iota)w + B_{\iota,\text{rem}}w + G_{\iota},\tag{11.81}$$

where

$$B(\iota) := D\tilde{A}(\iota)D^{-1} - \kappa_{\infty} \mathrm{Id}_{2m}, \quad B_{\iota,\mathrm{rem}} := D\tilde{A}_{\iota,\mathrm{rem}}D^{-1}, \quad G_{\iota} := e^{-\kappa_{\infty}t}D\tilde{F}_{\iota}.$$

Moreover,

$$B(\iota) = J + \mathcal{R}_{\iota}, \quad J := \operatorname{diag}\{J_1, \dots, J_K\}, \quad \mathcal{R}_{\iota} := \operatorname{diag}\{\mathcal{R}_{\iota,1}, \dots, \mathcal{R}_{\iota,K}\},\$$

where dim  $\mathcal{R}_{\iota,k}$  = dim  $J_k$  and the  $\mathcal{R}_{\iota,k}$  satisfy estimates similar to (11.80) (however, in this case the estimate depends on D). Here the  $J_k$  consist of generalised Jordan blocks (their non-zero offdiagonal components need not equal 1) and their diagonal components are non-positive. Moreover, we choose D so that if the diagonal components of  $J_k$  are strictly negative, then  $J_k$  is negative definite; cf. Lemma 9.9 and Remark 9.10. The price we need to pay in order to achieve this is that D depends on  $\gamma_{\pm}$ . For the resulting  $J_k$ , we can then ensure that the symmetrisation of  $J_k + \mathcal{R}_{\iota,k}$ is negative definite by demanding that  $\mu(\iota)$  be large enough; the symmetrisation of  $A \in \mathbf{M}_k(\mathbb{C})$  is given by  $(A + A^{\dagger})/2$ , where  $A^{\dagger}$  is the transpose of the complex conjugate of A. The lower bound on  $\mu(\iota)$  required to achieve this, say  $\mu_0$  depends only on U, V and W. Finally, we can ensure that if the diagonal component of  $J_k$  is  $\kappa_k$ , then

$$\|J_k + \mathcal{R}_{\iota,k} - \kappa_k \mathrm{Id}_{d_k}\| + \|J_k - \kappa_k \mathrm{Id}_{d_k}\| \le \beta_{\mathrm{mar}}/4, \tag{11.82}$$

where  $d_k$  is the dimension of  $J_k$ . In order to ensure that this estimate holds, we need to choose a D depending only on  $\gamma_{\pm}$  and  $\beta_{\text{mar}}$ . Moreover, we need to demand that  $\mu(\iota) \geq \mu_0$ , where  $\mu_0$ only depends on U, V, W and  $\beta_{\text{mar}}$ . At this stage, it is of interest to note that there is a  $\mu_0 > 0$ (depending only on U, V and W) such that for  $\mu(\iota) \geq \mu_0$ ,

$$\|B_{\iota,\mathrm{rem}}(t)\| \le C e^{-\beta_{\mathrm{ts}}\bar{t}} \tag{11.83}$$

for  $t \ge T_{\rm ts}$ , where C only depends on U, V,  $\beta_{\rm mar}$  and  $C_{\rm ts}$ ; here we use the fact that  $||T_{\iota}^{-1}||$  and  $||T_{\iota}||$  are bounded by numerical constants for  $\mu(\iota) \ge \mu_0$ . Similarly, for  $\mu(\iota) \ge \mu_0$  and  $t \ge 0$ ,

$$|G_{\iota}| \le C e^{-\kappa_{\infty} t} |F_{\iota}|, \tag{11.84}$$

where C only depends on U, V and  $\beta_{\text{mar}}$ , and  $\mu_0$  only depends on U, V and W.

Fourth reformulation of the equations; obtaining negative semi-definiteness. If the diagonal components of  $J_k$  are zero, let  $\bar{J}_k = \bar{\mathcal{R}}_{\iota,k} = 0$ . If the diagonal components of  $J_k$  are non-zero, let  $\bar{J}_k = J_k$  and  $\bar{\mathcal{R}}_{\iota,k} = \mathcal{R}_{\iota,k}$ . Finally, let

$$\bar{J} := \operatorname{diag}\{\bar{J}_1, \dots, \bar{J}_K\}, \quad \bar{\mathcal{R}}_\iota := \operatorname{diag}\{\bar{\mathcal{R}}_{\iota,1}, \dots, \bar{\mathcal{R}}_{\iota,K}\}, \quad \bar{B}(\iota) = \bar{J} + \bar{\mathcal{R}}_\iota$$

138

and  $B_{\iota,\text{diff}} := B(\iota) - \bar{B}(\iota)$ ; note that  $B_{\iota,\text{diff}}$ ,  $B(\iota)$  and  $\bar{B}(\iota)$  all commute. By construction there is a  $0 < \mu_0 \in \mathbb{R}$  (depending only on U, V, W and  $\beta_{\text{mar}}$ ) such that for  $\mu(\iota) \ge \mu_0$ ,

$$||B_{\iota,\mathrm{diff}}|| \leq \beta_{\mathrm{mar}}/4.$$

Let

$$u(t) := \exp\left(-B_{\iota, \text{diff}}\bar{t}\right) w(t).$$

Then

$$\dot{u} = \bar{B}(\iota)u + \bar{B}_{\iota,\text{rem}}u + H_{\iota}, \qquad (11.85)$$

where

$$\bar{B}_{\iota,\mathrm{rem}}(t) := \exp\left(-B_{\iota,\mathrm{diff}}\bar{t}\right) B_{\iota,\mathrm{rem}}(t) \exp\left(B_{\iota,\mathrm{diff}}\bar{t}\right), \quad H_{\iota}(t) := \exp\left(-B_{\iota,\mathrm{diff}}\bar{t}\right) G_{\iota}(t).$$

Note that for  $\mu(\iota) \geq \mu_0$ ,

$$\|\bar{B}_{\iota,\mathrm{rem}}(t)\| \leq C e^{-(\beta_{\mathrm{ts}}-\beta_{\mathrm{mar}}/2)\bar{t}}$$

for  $t \ge T_{\rm ts}$ , where C has the same dependence as in the case of (11.83) and  $\mu_0$  only depends on U, V, W and  $\beta_{\rm mar}$ . Similarly, if  $\mu(\iota) \ge \mu_0$ ,

$$|H_{\iota}| \le C e^{-\kappa_{\infty} t + \beta_{\max} \overline{t}/4} |F_{\iota}|, \qquad (11.86)$$

where C has the same dependence as in the case of (11.84) and  $\mu_0$  depends only on U, V, W and  $\beta_{\text{mar}}$ .

**Rough estimate of the solutions.** The main advantage of (11.85) is that the symmetrisation of  $\bar{B}(\iota)$  is negative semi-definite. Thus

$$\frac{d}{dt}|u|^2 = \langle \dot{u}, u \rangle + \langle u, \dot{u} \rangle \leq \langle [\bar{B}(\iota) + \bar{B}(\iota)^{\dagger}]u, u \rangle + 2\|\bar{B}_{\iota, \text{rem}}\||u|^2 + 2|H_{\iota}||u|$$
$$\leq 2\|\bar{B}_{\iota, \text{rem}}\||u|^2 + 2|H_{\iota}||u|.$$

Define

$$\sigma_{\iota}(t) := -\int_{T_{\rm ts}}^t \|\bar{B}_{\iota,\rm rem}(s)\| ds.$$

Then

$$\frac{d}{dt}(e^{2\sigma_{\iota}}|u|^2) \le 2e^{2\sigma_{\iota}}|H_{\iota}||u|.$$

This estimate can be used to deduce that

$$e^{\sigma_{\iota}(t)}|u(t)| \leq |u(T_{\rm ts})| + \int_{T_{\rm ts}}^{t} e^{\sigma_{\iota}(s)}|H_{\iota}(s)|ds$$

for  $t \geq T_{\rm ts}$ . Assuming  $\beta_{\rm mar} \leq \beta_{\rm ts}/2$ , there is thus a constant *C*, depending only on *U*, *V*,  $\beta_{\rm mar}$ ,  $\beta_{\rm ts}$  and  $C_{\rm ts}$ , such that

$$|u(t)| \le C|u(T_{\rm ts})| + C \int_{T_{\rm ts}}^t e^{-\kappa_\infty s + \beta_{\rm mar}\bar{s}/4} |F_\iota(s)| ds$$

for  $t \ge T_{ts}$  and  $\iota$  such that  $\mu(\iota) \ge \mu_0$ . Here  $\mu_0$  has the same dependence as in the case of (11.86) and we have appealed to (11.86). Combining the above definitions, this leads to the estimate

$$|v(t)| \le Ce^{(\kappa_{\infty} + \beta_{\max}/4)\overline{t}} |v(T_{\rm ts})| + Ce^{\beta_{\max}\overline{t}/4} \int_{T_{\rm ts}}^{t} e^{\kappa_{\infty}(t-s) + \beta_{\max}\overline{s}/4} |F_{\iota}(s)| ds$$
(11.87)

for  $t \ge T_{ts}$  and  $\mu(\iota) \ge \mu_0$ , where C depends only on U, V,  $\beta_{mar}$ ,  $\beta_{ts}$  and  $C_{ts}$ , and  $\mu_0$  depends only on U, V, W and  $\beta_{mar}$ . It is of interest to note that this estimate implies

$$|v(t)| \le C e^{(\kappa_{\iota} + \beta_{\max}/2)\bar{t}} |v(T_{ts})| + C e^{\beta_{\max}\bar{t}/2} \int_{T_{ts}}^{t} e^{\kappa_{\iota}(t-s) + \beta_{\max}\bar{s}/2} |F_{\iota}(s)| ds$$
(11.88)

for  $t \ge T_{ts}$  and  $\mu(\iota) \ge \mu_0$ , where  $\kappa_{\iota} = \kappa_{max}[A(\iota)]$  and C and  $\mu_0$  have the same dependence as in the case of (11.87). For future reference, it is also of interest to keep in mind that

$$|w(t)| \le C e^{\beta_{\max}\bar{t}/4} \left[ |w(T_{ts})| + \int_{T_{ts}}^{t} e^{-\kappa_{\infty}s + \beta_{\max}\bar{s}/4} |F_{\iota}(s)| ds \right]$$
(11.89)

for  $t \ge T_{\rm ts}$  and  $\mu(\iota) \ge \mu_0$ , where C and  $\mu_0$  have the same dependence as in the case of (11.87).

Splitting the equations. Let  $\mathcal{K}_{\pm}$  denote the set of real parts of elements of  $\operatorname{Sp}(\gamma_{\pm})$ . Then  $\kappa_{\infty}$  is the largest element of  $\mathcal{K} := \mathcal{K}_{+} \cup \mathcal{K}_{-}$ . Given  $\beta_{\mathrm{ts}}$ , there are two possibilities. Either  $\kappa_{\infty} - \beta_{\mathrm{ts}} \in \mathcal{K}$  or  $\kappa_{\infty} - \beta_{\mathrm{ts}} \notin \mathcal{K}$ . In either case, we let  $\kappa_{\min}$  be the smallest element of  $\mathcal{K}$  which is strictly larger than  $\kappa_{\infty} - \beta_{\mathrm{ts}}$ . Note that the distance between  $\kappa_{\min}$  and  $\kappa_{\infty} - \beta_{\mathrm{ts}}$  has a positive lower bound which only depends on U, V and  $\beta_{\mathrm{ts}}$ . Let  $\kappa_{\max} := \kappa_{\infty} - \kappa_{\min}$ . For  $k = 1, \ldots, K$ , consider  $J_k$ . If the diagonal component of  $J_k$  is in the interval  $[-\kappa_{\max}, 0]$ , let  $J_{k,a} := J_k, J_{k,b} := 0, \mathcal{R}_{k,a} := \mathcal{R}_{\iota,k}, \mathcal{R}_{k,b} := 0, \Pi_{k,a} := \operatorname{Id}_{d_k}$  and  $\Pi_{k,b} := 0$ . Similarly, if the diagonal component of  $J_k$  is in the interval  $(-\infty, -\beta_{\mathrm{ts}}]$ , let  $J_{k,a} := 0, J_{k,b} := J_k, \mathcal{R}_{k,a} := 0, \mathcal{R}_{k,b} := \mathcal{R}_{\iota,k}, \Pi_{k,a} := 0$  and  $\Pi_{k,b} := \operatorname{Id}_{d_k}$ . Let, moreover,

$$J_a := \operatorname{diag}\{J_{1,a}, \dots, J_{K,a}\}, \quad J_b := \operatorname{diag}\{J_{1,b}, \dots, J_{K,b}\}.$$
(11.90)

Define  $\mathcal{R}_a$ ,  $\mathcal{R}_b$ ,  $\Pi_a$  and  $\Pi_b$  similarly, and let  $B_{\iota,a} := J_a + \mathcal{R}_a$  and  $B_{\iota,b} := J_b + \mathcal{R}_b$ . For a vector  $\xi$  with values in  $\mathbb{C}^{2m}$  we similarly divide its components into  $\xi_a, \xi_b \in \mathbb{C}^{2m}$  such that  $\xi = \xi_a + \xi_b$ ; in fact,  $\xi_a = \Pi_a(\xi)$  and  $\xi_b = \Pi_b(\xi)$ . For future reference, it is of interest to keep in mind that if  $t \ge 0$ , then

$$\|e^{-B_{\iota,a}t}\| \le e^{(\kappa_{\max} + \beta_{\max}/4)t} \le e^{(\beta_{ts} - \beta_{\max}/2)t},$$
(11.91)

where we have appealed to (11.82), and the last inequality is based on the assumption that  $\beta_{\text{mar}}$  is small enough, the bound depending only on  $\beta_{\text{ts}}$  and  $\gamma_{\pm}$ . Similarly, for  $t \ge 0$ ,

$$\|e^{B_{\iota,b}t}\Pi_b\| \le e^{-(\beta_{\rm ts} - \beta_{\rm mar}/2)t}.$$
(11.92)

Let us return to (11.81). Given the above terminology, it can be divided into two components

$$\dot{w}_a = B_{\iota,a} w_a + (B_{\iota,\text{rem}} w)_a + G_{\iota,a},$$
(11.93)

$$\dot{w}_b = B_{\iota,b} w_b + (B_{\iota,\text{rem}} w)_b + G_{\iota,b};$$
 (11.94)

note that while  $B_{\iota}$  respects the division into components, the matrix  $B_{\iota,\text{rem}}$  cannot be expected to do so.

Analysis of the first equation. Integrating (11.93) yields

$$e^{-B_{\iota,a}\bar{t}}w_a(t) = w_a(T_{\rm ts}) + \int_{T_{\rm ts}}^t e^{-B_{\iota,a}\bar{s}}(B_{\iota,\rm rem}w)_a(s)ds + \int_{T_{\rm ts}}^t e^{-B_{\iota,a}\bar{s}}G_{\iota,a}(s)ds.$$
(11.95)

Define

$$w_{\infty,a} := w_a(T_{\rm ts}) + \int_{T_{\rm ts}}^{\infty} e^{-B_{\iota,a}\bar{s}} (B_{\iota,\rm rem}w)_a(s) ds.$$
(11.96)

Then (11.95) implies

$$\left| w_{a}(t) - e^{B_{\iota,a}\bar{t}} w_{\infty,a} - \int_{T_{ts}}^{t} e^{B_{\iota,a}(t-s)} G_{\iota,a}(s) ds \right| \\
\leq \left| \int_{t}^{\infty} e^{-B_{\iota,a}(s-t)} (B_{\iota,\text{rem}} w)_{a}(s) ds \right|.$$
(11.97)

Appealing to (11.83), (11.89) and (11.91) yields

$$\left| \int_{T_{\rm ts}}^{\infty} e^{-B_{\iota,a}\bar{s}} (B_{\iota,\rm rem}w)_a(s) ds \right| \le C \left[ |w(T_{\rm ts})| + e^{-\beta_{\rm mar}T_{\rm ts}/4} \int_{T_{\rm ts}}^{\infty} e^{-(\kappa_{\infty} - \beta_{\rm mar}/4)s} |F_{\iota}(s)| ds \right].$$
(11.98)

This estimate holds assuming  $\mu(\iota) \geq \mu_0$ , where  $\mu_0$  only depends on U, V, W and  $\beta_{\text{mar}}$ . Moreover, the constant C only depends on  $U, V, \beta_{\text{mar}}, \beta_{\text{ts}}$  and  $C_{\text{ts}}$ . Finally,  $\beta_{\text{mar}}$  has to be small enough, the bound depending only on  $\gamma_{\pm}$  and  $\beta_{\text{ts}}$ . In the end, we wish to replace  $\kappa_{\infty}$  by  $\kappa_{\iota}$  in (11.98). This can be achieved by replacing  $\beta_{\text{mar}}/4$  by  $\beta_{\text{mar}}$  in  $e^{-(\kappa_{\infty}-\beta_{\text{mar}}/4)s}$ . Moreover, it is convenient to introduce the notation (11.74). Combining (11.96) and (11.98) then yields

$$|w_{\infty,a}| \le C[|w(T_{\rm ts})| + e^{-\beta_{\rm mar}T_{\rm ts}/4} ||F_{\iota}||_{\rm lma}]$$
(11.99)

for  $\mu(\iota) \ge \mu_0$ , where C and  $\mu_0$  have the same dependence as in (11.98), and  $\beta_{\text{mar}}$  has to satisfy the same bound as in the case of (11.98). Returning to (11.97), we need to estimate the right hand side. Appealing to (11.83), (11.89) and (11.91) again yields the conclusion that it can be bounded by

$$Ce^{-(\beta_{\rm ts}-\beta_{\rm mar}/4)\overline{t}}[|w(T_{\rm ts})| + e^{-\beta_{\rm mar}T_{\rm ts}/4}||F_{\iota}||_{\rm lma}]$$

for  $t \ge T_{ts}$  and  $\mu(\iota) \ge \mu_0$ , where C and  $\mu_0$  have the same dependence as in (11.98), and  $\beta_{mar}$  has to satisfy the same bound as in the case of (11.98). Combining this estimate with (11.97) yields

$$\left| w_{a}(t) - e^{B_{\iota,a}\bar{t}} w_{\infty,a} - \int_{T_{ts}}^{t} e^{B_{\iota,a}(t-s)} G_{\iota,a}(s) ds \right|$$

$$\leq C e^{-(\beta_{ts} - \beta_{mar}/4)\bar{t}} [e^{-\kappa_{\infty}T_{ts}} |v(T_{ts})| + e^{-\beta_{mar}T_{ts}/4} \|F_{\iota}\|_{lma}]$$

$$(11.100)$$

for  $t \ge T_{ts}$  and  $\mu(\iota) \ge \mu_0$ , where C and  $\mu_0$  have the same dependence as in (11.98), and  $\beta_{mar}$  has to satisfy the same bound as in the case of (11.98).

Analysis of the second equation. Let us now turn to (11.94). Integrating this equation yields

$$w_b(t) = e^{B_{\iota,b}\bar{t}} w_b(T_{\rm ts}) + \int_{T_{\rm ts}}^t e^{B_{\iota,b}(t-s)} (B_{\iota,\rm rem}w)_b(s) ds + \int_{T_{\rm ts}}^t e^{B_{\iota,b}(t-s)} G_{\iota,b}(s) ds.$$
(11.101)

Appealing to (11.92) yields

$$|e^{B_{\iota,b}\bar{t}}w_b(T_{\rm ts})| \le e^{-(\beta_{\rm ts}-\beta_{\rm mar}/2)\bar{t}}|w_b(T_{\rm ts})|.$$
(11.102)

Next, (11.83), (11.89) and (11.92) yield

$$\left| \int_{T_{\rm ts}}^{t} e^{B_{\iota,b}(t-s)} (B_{\iota,\rm rem}w)_b(s) ds \right| \le C e^{-(\beta_{\rm ts}-\beta_{\rm mar}/2)\bar{t}} [|w(T_{\rm ts})| + e^{-\beta_{\rm mar}T_{\rm ts}/4} ||F_{\iota}||_{\rm lma}]$$
(11.103)

for  $t \ge T_{ts}$  and  $\mu(\iota) \ge \mu_0$ , where C and  $\mu_0$  have the same dependence as in (11.98), and  $\beta_{mar}$  has to satisfy the same bound as in the case of (11.98). Combining (11.101), (11.102) and (11.103) yields

$$\left| w_{b}(t) - \int_{T_{\rm ts}}^{t} e^{B_{\iota,b}(t-s)} G_{\iota,b}(s) ds \right| \le C e^{-(\beta_{\rm ts} - \beta_{\rm mar}/2)\bar{t}} [|w(T_{\rm ts})| + e^{-\beta_{\rm mar} T_{\rm ts}/4} ||F_{\iota}||_{\rm lma}]$$
(11.104)

for  $t \ge T_{ts}$  and  $\mu(\iota) \ge \mu_0$ , where C and  $\mu_0$  have the same dependence as in (11.98), and  $\beta_{mar}$  has to satisfy the same bound as in the case of (11.98).

**Combining the estimates.** Let  $w_{\infty} \in \mathbb{C}^{2m}$  be defined by  $w_{\infty} := w_{\infty,a} + w_{\infty,b}$ , where  $w_{\infty,b} = 0$ . Then (11.100) and (11.104) yield

$$\left| w(t) - e^{B(\iota)\bar{t}} w_{\infty} - \int_{T_{\rm ts}}^{t} e^{B(\iota)(t-s)} G_{\iota}(s) ds \right|$$

$$\leq C e^{-(\beta_{\rm ts} - \beta_{\rm mar}/2)\bar{t}} [e^{-\kappa_{\infty} T_{\rm ts}} |v(T_{\rm ts})| + e^{-\beta_{\rm mar} T_{\rm ts}/4} \|F_{\iota}\|_{\rm lma}]$$

$$(11.105)$$

for  $t \ge T_{ts}$  and  $\mu(\iota) \ge \mu_0$ , where C and  $\mu_0$  have the same dependence as in (11.98), and  $\beta_{mar}$  has to satisfy the same bound as in the case of (11.98). This estimate can be reformulated as

$$\left| v(t) - e^{A(\iota)t} v_{\infty} - \int_{T_{\rm ts}}^{t} e^{A(\iota)(t-s)} F_{\iota}(s) ds \right|$$

$$\leq C e^{\kappa_{\infty} t - (\beta_{\rm ts} - \beta_{\rm mar}/2)\bar{t}} [e^{-\kappa_{\infty} T_{\rm ts}} |v(T_{\rm ts})| + e^{-\beta_{\rm mar} T_{\rm ts}/4} \|F_{\iota}\|_{\rm Ima}]$$

$$(11.106)$$

for  $t \ge T_{ts}$  and  $\mu(\iota) \ge \mu_0$ , where C and  $\mu_0$  have the same dependence as in (11.98), and  $\beta_{mar}$  has to satisfy the same bound as in the case of (11.98). Moreover,

$$v_{\infty} := e^{\kappa_{\infty} T_{\rm ts}} e^{-A(\iota)T_{\rm ts}} T_{\iota} D^{-1} w_{\infty}.$$
(11.107)

Sacrificing a part of the margin yields (11.75) for  $t \ge T_{\rm ts}$  and  $\mu(\iota) \ge \mu_0$ , where C and  $\mu_0$  have the same dependence as in (11.98), and  $\beta_{\rm mar}$  has to satisfy the same bound as in the case of (11.98). Finally, let  $u_{\infty} := e^{\kappa_{\infty} T_{\rm ts}} T_{\iota} D^{-1} w_{\infty}$ . Then  $u_{\infty} \in E_{\iota,\beta_{\rm ts}}^{\infty}$  due to the definition of  $w_{\infty}$ , D and  $T_{\iota}$ . Moreover, the estimate (11.99) yields the conclusion that (11.77) holds. The lemma follows.

### 11.5 Specifying data at infinity

The present section is quite analogous to Section 9.4; the goal is to construct a solution to (11.71) corresponding to the asymptotic data  $v_{\infty}$  appearing in (11.75). However, in the present setting the fact that the matrix  $A(\iota)$  appearing in (11.71) depends on the mode number  $\iota$  causes additional complications. Fortunately, it turns out that these complications can be handled using the methods developed in the previous section. Just as in the case of Section 9.4, it is, moreover, sufficient to consider homogeneous equations.

**Lemma 11.19.** Let  $1 \leq m \in \mathbb{Z}$ ,  $\mathcal{I}_{B}$  be the set introduced in connection with (24.3) and  $\mu$  be a function  $\mu : \mathcal{I}_{B} \to [0, \infty)$ . Let  $\mathcal{I}_{B,ts} \subseteq \mathcal{I}_{B}$  be the set of  $\iota \in \mathcal{I}_{B}$  such that  $\mu(\iota) > 0$ . Let  $U, V, W \in \mathbf{M}_{m}(\mathbb{C})$  and define  $A(\iota)$  by (11.73) for  $\iota \in \mathcal{I}_{B,ts}$ . Assume that for each  $\iota \in \mathcal{I}_{B,ts}$ , there is a function  $A_{\iota,rem} \in C^{\infty}(\mathbb{R}, \mathbf{M}_{2m}(\mathbb{C}))$  with the property that there are constants  $0 < \beta_{ts}, C_{ts} \in \mathbb{R}$ (independent of  $\iota$ ) and a constant  $T_{ts}$  (which is allowed to depend on  $\iota$ ) such that (11.72) holds for  $t \geq T_{ts}$ . Fix a  $0 < \beta_{mar} \in \mathbb{R}$  which is small enough, the bound depending only on U, V and  $\beta_{ts}$ . Then there are constants  $0 < \mu_0 \in \mathbb{R}$  and  $0 < C \in \mathbb{R}$ , where  $\mu_0$  only depends on U, V, W and  $\beta_{mar}$ , and C only depends on  $U, V, \beta_{ts}, \beta_{mar}$  and  $C_{ts}$ , such that the following holds. If  $\iota \in \mu^{-1}([\mu_0, \infty))$ , there is a linear injective map  $\Psi_{\iota,\infty} : E_{\iota,\beta_{ts}}^{\infty} \to \mathbb{C}^{2m}$  such that if  $v_{\infty} \in E_{\iota,\beta_{ts}}^{\infty}$  and v is the solution to

$$\dot{v} = A(\iota)v + A_{\iota,\text{rem}}v \tag{11.108}$$

with initial data  $v(T_{ts}) = \Psi_{\iota,\infty}(v_{\infty})$ , then v satisfies (11.75) (with  $F_{\iota} = 0$ ) for  $t \ge T_{ts}$ . Moreover,

$$|\Psi_{\iota,\infty}[e^{-A(\iota)T_{\rm ts}}\eta]| \le C|\eta| \tag{11.109}$$

for  $\eta \in E^{\infty}_{\iota,\beta_{ts}}$ , where C only depends on  $U, V, \beta_{mar}, \beta_{ts}$  and  $C_{ts}$ .

**Remark 11.20.** The result can be used to derive conclusions concerning the inhomogeneous equation (11.71). In order to justify this statement, assume that  $||F_{\iota}||_{\text{Ima}} < \infty$ , where  $\beta_{\text{mar}}$  and  $\iota$  are such that the assumptions of Lemmas 11.17 and 11.19 are satisfied. Let  $v_{\text{part}}$  be the solution to (11.71) corresponding to the initial data  $v_{\text{part}}(T_{\text{ts}}) = 0$ . Let  $v_{\text{part},\infty} \in E_{\iota,\beta_{\text{ts}}}^{\infty}$  be such that (11.75) holds with v and  $v_{\infty}$  replaced by  $v_{\text{part}}$  and  $v_{\text{part},\infty}$  respectively. Given  $v_{\infty} \in E_{\iota,\beta_{\text{ts}}}^{\infty}$ , let  $v_{\text{hom}}$  be the solution to (11.108) corresponding to initial data  $v_{\text{hom}}(T_{\text{ts}}) = \Psi_{\iota,\infty}(v_{\infty} - v_{\text{part},\infty})$ . Then  $v := v_{\text{part}} + v_{\text{hom}}$  solves (11.71) and satisfies (11.75). In other words, we are also allowed to specify the asymptotic data  $v_{\infty} \in E_{\iota,\beta_{\text{ts}}}^{\infty}$  in the case of the inhomogeneous equation (11.71), assuming  $||F_{\iota}||_{\text{Ima}} < \infty$ . In order to obtain estimates, note first of all that (11.76) and (11.77) hold due to Lemma 11.17, where  $u_{\infty} \in E_{\iota,\beta_{\text{ts}}}^{\infty}$  is defined by the first equation. There is a similar

statement with  $v_{\infty}$  and  $u_{\infty}$  replaced by  $v_{\text{part},\infty}$  and  $u_{\text{part},\infty}$ . Keeping in mind that  $v_{\text{part}}(T_{\text{ts}}) = 0$ , this yields

$$|v(T_{\rm ts})| = |v_{\rm hom}(T_{\rm ts})| = |\Psi_{\iota,\infty}(v_{\infty} - v_{\rm part,\infty})| = |\Psi_{\iota,\infty}[e^{-A(\iota)T_{\rm ts}}(u_{\infty} - u_{\rm part,\infty})]|$$
  

$$\leq C|u_{\infty} - u_{\rm part,\infty}| \leq C[|u_{\infty}| + e^{\kappa_{\iota}T_{\rm ts}}||F_{\iota}||_{\rm Ima}],$$
(11.110)

where we have appealed to (11.109) and the constant C only depends on  $U, V, \beta_{\text{mar}}, \beta_{\text{ts}}$  and  $C_{\text{ts}}$ .

**Remark 11.21.** The equality  $E_{\iota,\beta_{ts}}^{\infty} = \mathbb{C}^{2m}$  holds if  $\beta_{ts}$  is strictly larger than the difference between the largest and the smallest real parts of elements of  $\operatorname{Sp}(U_+) \cup \operatorname{Sp}(U_-)$ ; cf. Definition 11.16. According to the statement, the map  $\Psi_{\iota,\infty}$  is then an isomorphism. This means that the solution is uniquely specified in terms of the "data at infinity",  $v_{\infty}$ . Due to Remark 11.20, the same holds for inhomogeneous equations such that  $||F_{\iota}||_{\operatorname{Ima}} < \infty$ . Finally,  $|v(T_{ts})|$  can be estimated in terms of the asymptotic data and  $F_{\iota}$ ; cf. (11.110).

*Proof.* The argument is based on the proof of Lemma 11.17. We therefore use the notation introduced in that proof without further comment. A natural starting point is the third reformulation of the equations, as given in the proof of Lemma 11.17. Let

$$\tilde{v}(t) := e^{-B_{\iota,a}\bar{t}}w(t).$$

Then (11.81) with  $G_{\iota} = 0$  implies

$$\tilde{\tilde{w}} = B_{\iota,b}\tilde{w} + \tilde{B}_{\iota,\text{rem}}\tilde{w},\tag{11.111}$$

where

$$\tilde{B}_{\iota,\mathrm{rem}}(t) := e^{-B_{\iota,a}\bar{t}} B_{\iota,\mathrm{rem}}(t) e^{B_{\iota,a}\bar{t}}$$

Due to (11.82), (11.83) and the definition of  $B_{\iota,a}$ ,

$$\|\tilde{B}_{\iota,\mathrm{rem}}(t)\| \le C e^{-(\beta_{\mathrm{ts}}-\kappa_{\mathrm{mar}}-\beta_{\mathrm{mar}}/2)\bar{t}},$$

where C only depends on  $U, V, \beta_{\text{mar}}$  and  $C_{\text{ts}}$ . However, to obtain this estimate, we need to demand that  $\mu(\iota) \geq \mu_0$ , where  $\mu_0$  only depends on U, V, W and  $\beta_{\text{mar}}$ . Note that  $\beta_{\text{ts}} - \kappa_{\text{mar}} > 0$  and that  $\kappa_{\text{mar}}$  is determined by the eigenvalues of  $U_{\pm}$  (or, equivalently,  $\gamma_{\pm}$ ) and  $\beta_{\text{ts}}$ . Assuming  $\beta_{\text{mar}}$  to be small enough, the bound depending only on U, V and  $\beta_{\text{ts}}$ , we thus have

$$\|\tilde{B}_{\iota,\mathrm{rem}}(t)\| \le C e^{-(\beta_{\mathrm{ts}} - \kappa_{\mathrm{mar}})\bar{t}/2},\tag{11.112}$$

where C only depends on  $U, V, \beta_{\text{mar}}$  and  $C_{\text{ts}}$ . Note also that the symmetrisation of  $B_{\iota,b}$  is negative semi-definite. Let  $t_0 \geq T_{\text{ts}}$ . By an argument similar to the derivation of (9.49),

$$|\tilde{w}(t)| \le \exp\left(\int_{t_0}^t \|\tilde{B}_{\iota,\text{rem}}(s)\|ds\right) |\tilde{w}(t_0)|$$
(11.113)

for  $t \ge t_0$ . Note, in particular, that by choosing  $t_0 - T_{\rm ts}$  to be large enough (the bound depending only on  $U, V, \beta_{\rm mar}, \beta_{\rm ts}$  and  $C_{\rm ts}$ ), the first factor on the right hand side of (11.113) can be assumed to be as close to 1 as we wish. Let  $w_a$  and  $w_b$  be defined as in the proof of Lemma 11.17 and define

$$\tilde{w}_a(t) := e^{-B_{\iota,a}\bar{t}} w_a(t), \quad \tilde{w}_b(t) := e^{-B_{\iota,a}\bar{t}} w_b(t).$$
 (11.114)

Note that  $\tilde{w}_b = w_b$ . Then appealing to (11.93) with  $G_{\iota,a} = 0$  yields

$$\dot{\tilde{w}}_a = e^{-B_{\iota,a}\tilde{t}} (B_{\iota,\text{rem}} w)_a = (e^{-B_{\iota,a}\tilde{t}} B_{\iota,\text{rem}} w)_a = (\tilde{B}_{\iota,\text{rem}} \tilde{w})_a.$$

Thus

$$\tilde{w}_a(t) = \tilde{w}_a(t_0) + \int_{t_0}^t (\tilde{B}_{\iota,\text{rem}}\tilde{w})_a(s)ds,$$
(11.115)

so that

$$|\tilde{w}_a(t) - \tilde{w}_a(t_0)| \le C \int_{t_0}^t e^{-(\beta_{\rm ts} - \kappa_{\rm mar})\bar{s}/2} ds \exp\left(\int_{t_0}^\infty \|\tilde{B}_{\iota,\rm rem}(s)\| ds\right) |\tilde{w}(t_0)|$$

for  $t \ge t_0$ , where we have appealed to (11.112) and (11.113) and C only depends on  $U, V, \beta_{\text{mar}}$  and  $C_{\text{ts}}$ . For any  $\epsilon > 0$ , it is thus possible to choose  $t_0$  so that

$$|\tilde{w}_{a}(t) - \tilde{w}_{a}(t_{0})| \le \epsilon |\tilde{w}(t_{0})| \tag{11.116}$$

for  $t \geq t_0$  (where  $\bar{t}_0 := t_0 - T_{\rm ts}$  only depends on  $U, V, \beta_{\rm mar}, \beta_{\rm ts}, C_{\rm ts}$  and  $\epsilon$ ). Before proceeding, let  $V_a := \Pi_a(\mathbb{C}^{2m})$  and  $V_b := \Pi_b(\mathbb{C}^{2m})$ , where  $\Pi_a$  and  $\Pi_b$  are introduced in connection with (11.90). Given a  $t_0$  such that (11.116) holds with  $\epsilon$  replaced by 1/2, define the map  $L_a : V_a \to V_a$  as follows. Given  $\eta \in V_a$ , let  $\tilde{w}$  be the solution to (11.111) with  $\tilde{w}(t_0) = \eta$  and define  $L_a \eta$  by

$$L_a \eta := \lim_{t \to \infty} \tilde{w}_a(t).$$

Due to (11.112), (11.113) and (11.115), it is clear that the limit on the right hand side exists. Thus  $L_a\eta$  is well defined. Moreover, it is clear that  $L_a$  is linear. Due to the fact that (11.116) holds with  $\epsilon$  replaced by 1/2 and the fact that  $\tilde{w}_b(t_0) = 0$ ,

$$|L_a\eta - \tilde{w}_a(t_0)| \le \frac{1}{2} |\tilde{w}_a(t_0)|.$$

In particular,  $|\eta|/2 \leq |L_a\eta|$ , so that  $L_a$  is injective. Since  $L_a$  is a linear map between spaces of the same finite dimension, it follows that  $L_a$  is an isomorphism. Moreover,

$$|L_a^{-1}\eta| \le 2|\eta|. \tag{11.117}$$

Note that  $L_a^{-1}$  (essentially) maps data at infinity to data at  $t_0$ . In the end, we wish to translate the data at  $t_0$  to data at  $T_{ts}$ . To this end, we wish to solve (11.111) backwards. Since  $\bar{t}_0$  can be bounded by a constant depending only on  $U, V, \beta_{mar}, \beta_{ts}$  and  $C_{ts}$ , there is a constant C (with the same dependence) such that for all solutions  $\tilde{w}$  to (11.111)

$$|\tilde{w}(T_{\rm ts})| \le C |\tilde{w}(t_0)|.$$
 (11.118)

Before defining the map  $\Psi_{\iota,\infty}$  appearing in the statement of the lemma, we need to clarify how the limit of  $\tilde{w}_a(t)$  is related to the  $v_{\infty} \in E^{\infty}_{\iota,\beta_{ts}}$  appearing in (11.75). However, returning to (11.95) and (11.96) (and recalling that  $G_{\iota,a} = 0$  in the present setting), it is clear that

$$\lim_{t \to \infty} \tilde{w}_a(t) = \lim_{t \to \infty} e^{-B_{\iota,a}\bar{t}} w_a(t) = w_{\infty,a}.$$
(11.119)

On the other hand, (11.107) and the fact that  $w_{\infty} = w_{\infty,a}$  imply that

$$v_{\infty} = e^{\kappa_{\infty} T_{\rm ts}} e^{-A(\iota)T_{\rm ts}} T_{\iota} D^{-1} w_{\infty,a}.$$

Note also that for each  $v_{\infty} \in E^{\infty}_{\iota,\beta_{ts}}$ , there is a unique  $\eta \in V_a$  such that

$$v_{\infty} = e^{\kappa_{\infty} T_{\rm ts}} e^{-A(\iota)T_{\rm ts}} T_{\iota} D^{-1} \eta \tag{11.120}$$

holds. Define  $\Psi_{\iota,\infty}(v_{\infty})$  to be the composition of the following maps: first the map which takes  $v_{\infty} \in E_{\iota,\beta_{\text{ts}}}^{\infty}$  to the  $\eta \in V_a$  such that (11.120) holds; second, the map which takes  $\eta$  to  $\xi := L_a^{-1}(\eta)$ ; and, third, the map which takes  $\xi \in V_a$  to

$$v_0 := e^{\kappa_\infty T_{\rm ts}} T_\iota D^{-1} \tilde{w}(T_{\rm ts}), \tag{11.121}$$

where  $\tilde{w}$  is the solution to (11.111) with initial data  $\tilde{w}(t_0) = \xi$ . Note that  $\Psi_{\iota,\infty}$  is linear and injective, and that for  $\eta \in E^{\infty}_{\iota,\beta_{ts}}$ 

$$|\Psi_{\iota,\infty}[e^{-A(\iota)T_{\rm ts}}\eta]| \le C|\eta|,$$

144

where C only depends on  $U, V, \beta_{\text{mar}}, \beta_{\text{ts}}$  and  $C_{\text{ts}}$ , and we have appealed to (11.117), (11.118), (11.120) and (11.121). Thus (11.109) holds.

Assume now that v is the solution to (11.108) corresponding to  $v(T_{ts}) = \Psi_{\iota,\infty}(v_{\infty})$  for some  $v_{\infty} \in E^{\infty}_{\iota,\beta_{ts}}$ . Then v satisfies (11.75) with  $F_{\iota} = 0$  and  $v_{\infty}$  replaced by some  $\bar{v}_{\infty} \in E^{\infty}_{\iota,\beta_{ts}}$ . Moreover, the proof of Lemma 11.17, in particular (11.107), yields

$$\bar{v}_{\infty} := e^{\kappa_{\infty} T_{\rm ts}} e^{-A(\iota)T_{\rm ts}} T_{\iota} D^{-1} \bar{w}_{\infty}$$

where  $\bar{w}_{\infty} \in V_a$ . In addition,

$$\lim_{t \to \infty} \tilde{w}_a(t) = \bar{w}_\infty,$$

where  $\tilde{w}_a$  is defined by (11.114). This means that

$$L_a^{-1}(\bar{w}_\infty) = \tilde{w}(t_0),$$

due to the definition of  $L_a$  and the fact that  $\tilde{w}(t_0) = \tilde{w}_a(t_0)$ . Going through the steps defining  $\Psi_{\iota,\infty}$ , it is thus clear that  $\Psi_{\iota,\infty}(\bar{v}_{\infty}) = v(T_{\rm ts}) = \Psi_{\iota,\infty}(v_{\infty})$ . Due to the injectivity of  $\Psi_{\iota,\infty}$ , it follows that  $v_{\infty} = \bar{v}_{\infty}$ . The lemma follows.

#### **11.6** Algebraic properties of $A(\iota)$

As a final section of the present chapter, it is convenient to collect some of the algebraic properties of  $A(\iota)$  that we need in what follows.

**Lemma 11.22.** Let  $1 \leq m \in \mathbb{Z}$ ,  $\mathcal{I}_{B}$  be the set introduced in connection with (24.3) and  $\mu$  be a function  $\mu : \mathcal{I}_{B} \to [0, \infty)$ . Let  $\mathcal{I}_{B,ts} \subseteq \mathcal{I}_{B}$  be the set of  $\iota \in \mathcal{I}_{B}$  such that  $\mu(\iota) > 0$ . Let  $U, V, W \in \mathbf{M}_{m}(\mathbb{C})$  and define  $A(\iota)$  by (11.73) for  $\iota \in \mathcal{I}_{B,ts}$ . Let  $\beta > 0$ . Then there are constants C and  $\mu_{0} > 0$  (depending only on U, V, W and  $\beta$ ) such that

$$|e^{-A(\iota)t}\eta| \le Ce^{-(\kappa_{\iota}-\beta)t}|\eta|$$

for all  $\iota$  such that  $\mu(\iota) \geq \mu_0$ ; all  $\eta \in E_{\iota,\beta}^{\infty}$ ; and all  $t \geq 0$ . Using the notation introduced in Definition 11.16, let  $e_{\iota,\beta}^{\infty}$  be the direct sum of the generalised eigenspaces of  $A(\iota)$  corresponding to eigenvalues in the balls of radius  $r_a$  and centre  $\pm i\mu(\iota) + \lambda_{j,\pm}$  for j such that  $\kappa_{\infty} - \operatorname{Re}(\lambda_{j,\pm}) \geq \beta$ . Given  $\epsilon > 0$ , there are constants C and  $\mu_0 > 0$  (depending only on U, V, W and  $\epsilon$ ) such that

$$|e^{A(\iota)t}\eta| \le C e^{(\kappa_b + \epsilon)t} |\eta|$$

for all  $\iota$  such that  $\mu(\iota) \ge \mu_0$ ; all  $\eta \in e_{\iota,\beta}^{\infty}$ ; and all  $t \ge 0$ , where  $\kappa_b$  is the largest element of  $\{\operatorname{Re}(\lambda_{j,\pm})\}\$  which is  $\le \kappa_{\infty} - \beta$ .

*Proof.* Let  $\mu_0 > 0$  be larger than the  $\mu_0$  appearing in Definition 11.16 and let  $\iota$  be such that  $\mu(\iota) \ge \mu_0, \eta \in E^{\infty}_{\iota,\beta}$  and  $t \ge 0$ . Keeping the change of variables introduced at the beginning of the proof of Lemma 11.17 in mind,

$$e^{-A(\iota)t}\eta = T_{\iota}T_{\iota}^{-1}e^{-A(\iota)t}T_{\iota}T_{\iota}^{-1}\eta = T_{\iota}e^{-\bar{A}(\iota)t}T_{\iota}^{-1}\eta, \qquad (11.122)$$

where

$$\bar{A}(\iota) = \operatorname{diag}\{N_{\iota,1}^{-}, \dots, N_{\iota,p_{-}}^{-}, N_{\iota,1}^{+}, \dots, N_{\iota,p_{+}}^{+}\};$$
(11.123)

cf. (11.25). The blocks appearing on the right hand side of (11.123) are given by

$$N_{\iota,j}^{\pm} = \pm i\mu(\iota) \mathrm{Id}_{m_{j,\pm}} + \gamma_{\pm,j} + \mathcal{R}_{\iota,j}^{\pm}, \qquad (11.124)$$

where

$$\|\mathcal{R}_{\iota,i}^{\pm}\| \le c_0[\mu(\iota)]^{-1} \tag{11.125}$$

for  $\mu(\iota) \geq \mu_0$ , where  $c_0$  and  $\mu_0$  only depend on U, V and W. Recall that the notation  $\gamma_{\pm,j}$  is introduced in Lemma 11.14. The matrices  $\gamma_{\pm,j}$  are defined in terms of  $\gamma_{\pm}$ , which are defined in terms of U and V. Recall, finally, that  $\gamma_{\pm,j}$  consists of the Jordan blocks with diagonal components  $\lambda_{j,\pm}$ . The blocks appearing on the right hand side of (11.123) correspond to a division of the generalised eigenspaces of  $A(\iota)$ ; cf. Definition 11.16. The blocks of greatest interest are the ones corresponding to eigenvalues that belong to the sets described in Definition 11.16, since they correspond to the only non-zero components of  $T_{\iota}^{-1}\eta$ . Considering (11.122), it is thus clear that we only need to estimate

$$||T_{\iota}|| \cdot ||T_{\iota}^{-1}|| \cdot ||\exp(-N_{\iota,j}^{\pm}t)|| \cdot |\eta|$$

in the case that the real parts of the diagonal components of  $\gamma_{\pm,j}$  are strictly larger than  $\kappa_{\infty} - \beta$ , where the notation is taken from Definition 11.16. Note that

$$\exp(-N_{\iota,j}^{\pm}t) = \exp[\mp i\mu(\iota)t - i\operatorname{Im}(\lambda_{j,\pm})t] \exp[-\operatorname{Re}(\lambda_{j,\pm})t] \cdot \exp[-(\gamma_{\pm,j} - \lambda_{j,\pm}\operatorname{Id}_{m_{j,\pm}} + \mathcal{R}_{\iota,j}^{\pm})t].$$
(11.126)

Note that the first factor on the right hand side has absolute value one, and that

$$\exp[-\operatorname{Re}(\lambda_{j,\pm})t] \le e^{-(\kappa_{\infty}-\beta)t}.$$
(11.127)

In fact, the latter estimate can be improved; since there are only finitely many eigenvalues  $\lambda_{j,\pm}$  and  $\operatorname{Re}(\lambda_{j,\pm}) > \kappa_{\infty} - \beta$  for all the eigenvalues of relevance here, there is a gap between the smallest  $\operatorname{Re}(\lambda_{j,\pm})$  satisfying  $\operatorname{Re}(\lambda_{j,\pm}) > \kappa_{\infty} - \beta$ , and  $\kappa_{\infty} - \beta$  (and the gap only depends on  $\beta$ , U and V). What remains to estimate is the norm of the last factor on the right hand side of (11.126). In order to simplify the notation, note that this factor can be written  $\exp[-(N+R)t]$ , where N consists of Jordan blocks corresponding to the eigenvalue zero (in other words, the only potentially non-zero elements in N are ones immediately above the diagonal) and ||R|| is bounded by the right hand side of (11.125). Given  $\eta_{\max} > 0$  it is possible to conjugate N by a diagonal matrix, say  $S_{\max} \in \operatorname{\mathbf{GL}}_{m_{j,\pm}}(\mathbb{R})$ , such that  $||S_{\max}NS_{\max}^{-1}|| \leq \eta_{\max}/2$ ; cf. Lemma 9.9 and its proof. The maximal element of  $\{||S_{\max}||, ||S_{\max}^{-1}||\}$  then depends only on  $\eta_{\max}$ . Given such an  $\eta_{\max}$ , there is then a  $\mu_0 > 0$ , depending only on  $\eta_{\max}$ , U, V and W, such that if  $\mu(\iota) \geq \mu_0$ , then  $||S_{\max}RS_{\max}^{-1}|| \leq \eta_{\max}/2$ . Thus

$$\|\exp[-(N+R)t]\| \le \|S_{\max}^{-1}\exp[-S_{\max}(N+R)S_{\max}^{-1}t]S_{\max}\| \le Ce^{\eta_{\max}t}$$

where the constant only depends on  $\eta_{\text{mar}}$ . Combining this estimate with (11.126), (11.127) and the comments made in connection with (11.127), it is clear that by first fixing  $\eta_{\text{mar}} > 0$  small enough (in fact, we specify  $\eta_{\text{mar}}$  in terms of U, V and  $\beta$ ) and then requiring  $\mu_0$  to be large enough, the bound depending only on U, V, W and  $\beta$ , we can ensure that

$$\|\exp(-N_{\iota,i}^{\pm}t)\| \le Ce^{-(\kappa_{\iota}-\beta)t},$$

where C only depends on U, V, W and  $\beta$ . In order to obtain this conclusion, we have also appealed to the fact that  $|\kappa_{\infty} - \kappa_{\iota}| \leq c_0 [\mu(\iota)]^{1/m}$ , where  $c_0$  only depends on U, V and W; cf. Lemma 11.6. Combining this estimate with previous observations yields the first statement of the lemma. The proof of the second statement is similar.

**Lemma 11.23.** Let  $1 \leq m \in \mathbb{Z}$ ,  $\mathcal{I}_{B}$  be the set introduced in connection with (24.3) and  $\mu$  be a function  $\mu : \mathcal{I}_{B} \to [0, \infty)$ . Let  $\mathcal{I}_{B,ts} \subseteq \mathcal{I}_{B}$  be the set of  $\iota \in \mathcal{I}_{B}$  such that  $\mu(\iota) > 0$ . Let  $U, V, W \in \mathbf{M}_{m}(\mathbb{C})$  and define  $A(\iota)$  by (11.73) for  $\iota \in \mathcal{I}_{B,ts}$ . Let  $\epsilon > 0$ . Then there are constants C and  $\mu_{0} > 0$  (depending only on U, V, W and  $\epsilon$ ) such that

$$\|e^{-A(\iota)t}\| < Ce^{-(\kappa_{\min,\iota}-\epsilon)t},\tag{11.128}$$

$$\|e^{A(\iota)t}\| < Ce^{(\kappa_\iota + \epsilon)t} \tag{11.129}$$

for all  $\iota$  such that  $\mu(\iota) \ge \mu_0$ ; all  $\eta \in \mathbb{C}^{2m}$ ; and all  $t \ge 0$ . Here  $\kappa_{\iota} := \kappa_{\max}[A(\iota)]$  and  $\kappa_{\min,\iota} := \kappa_{\min}[A(\iota)]$ .

**Remark 11.24.** In (11.128) and (11.129),  $\kappa_{\min,\iota}$  and  $\kappa_{\iota}$  can be replaced by  $\kappa_b$  and  $\kappa_a$  respectively, where  $\kappa_a$  ( $\kappa_b$ ) is the largest (smallest) real part of an element of  $\operatorname{Sp}(U_+) \cup \operatorname{Sp}(U_-)$ ; cf. the notation introduced in Definition 11.16.

*Proof.* The proof is based on arguments similar to those of the proof of Lemma 11.22. We therefore leave the details to the reader.  $\Box$ 

# Chapter 12

# Asymptotics for weakly transparent, balanced and convergent equations

## 12.1 Introduction

In the present chapter, we consider the asymptotics of solutions to so-called weakly transparent, balanced and convergent equations; cf. Definition 12.8 below for a clarification of the terminology. The arguments and results are quite analogous to those of Chapter 10; in particular, we derive asymptotics and provide a partial characterisation of solutions in terms of asymptotic data. The intuitive reason why we refer to the equations studied here as transparent is that the spatial coordinate of a causal curve no-longer needs to converge; cf. the discussion in Subsection 1.3.2. The relevance of the notions of balance and convergence is the same in the present setting as in the case of Chapter 10. The main steps of the analysis are the following.

Silent and transparent variables. One important difference when comparing the present setting with silent equations is that for transparent equations, it is natural to divide the spatial variables into silent and transparent variables; cf. Definitions 12.1 and 12.3 below. Roughly speaking, the silent variables correspond to the subspace (of the cotangent space of  $\overline{M}$ ) on which the spatial part of the inverse metric degenerates as  $t \to \infty$ , and the transparent variables correspond to the subspace on which the spatial part of the inverse metric converges to a positive definite bilinear form as  $t \to \infty$ . Projecting a solution onto the modes that correspond to dependence only on the silent variables leads to a situation analogous to that considered in Chapter 10. For that reason we here focus on the situation that results when projecting onto the modes that correspond to dependence on at least one of the transparent variables.

**Defining the notion of weak transparency.** Before proceeding to the analysis of the asymptotics, we have to give a precise meaning to the notion of weak transparency, and this is done in Definition 12.8 below. The first criterion is that one of Definitions 11.1 and 11.3 hold. The reason for making this assumption is discussed in the introduction of Chapter 11. The function  $g^2(\iota, t)$  can then be divided into three parts; one corresponding to the silent variables, one corresponding to the transparent variables and one corresponding to a mixture. The one corresponding to the silent variables should satisfy a condition similar to (10.3). In fact, we demand that (12.3) hold. Concerning the part of  $g^2(\iota, t)$  corresponding to the transparent variables we demand that it converge exponentially; this is one consequence of (12.4). However, the estimate (12.4) also implies that  $g^{0j}$  converges to zero exponentially if j corresponds to a transparent variable. Concerning the mixture, it turns out that it can be considered to be an error term. Finally, we require the  $X^j$ 

to converge exponentially when j corresponds to a transparent variable; cf. (12.5). To summarise, it is clear that due to the fact that there is a division of the variables and that the asymptotic behaviour of the inverse metric is qualitatively different on the corresponding subspaces of the cotangent space, the definition of weak transparency is somewhat more involved than the definition of weak silence. It also requires more of an effort to arrive at (11.3)–(11.6), in particular to calculate the exact dependence of the constants  $C_{\rm ts}$  and  $\beta_{\rm ts}$  on the constants and functions appearing in the definition of weak transparency, balance and convergence. We derive the estimates we need in Section 12.2.

Analysis prior to the transparent era. In Chapter 11 we obtain conclusions starting from the assumption that (11.3)-(11.6) hold. However, this analysis is only valid at late times; i.e., for  $t \ge T_{ts}$ , where  $T_{ts}$  depends on the mode under consideration. Here we need estimates for all  $t \ge 0$ . We therefore have to take the step from t = 0 to  $t = T_{ts}$ . This is achieved by appealing to Lemma 10.7. The proof of this lemma is very simple. On the other hand, the corresponding part of the argument is the source of the greatest weakness of the results. The reason for this is that we only obtain a very rough estimate of the number of derivatives lost in the estimates of the asymptotics.

**Deriving and specifying asymptotics.** Finally, we both derive and specify asymptotics; cf. Sections 12.3 and 12.4. Partly, the presentation parallels that of Sections 10.5 and 10.6. However, there are two fundamental differences. One is that in the case of silent equations, all the modes have the same asymptotic behaviour. This is not true for transparent equations. In fact, if v is a solution to (11.3) and  $F_{\iota}$  satisfies appropriate bounds, we expect v to behave asymptotically as  $\langle t \rangle^{\delta_{\iota}-1} e^{\kappa_{\iota} t}$ , where  $\kappa_{\iota} := \kappa_{\max}[A(\iota)]$  and  $\delta_{\iota} := d_{\max}[A(\iota), \kappa_{\iota}]$ ; cf. Definition 4.3. This leads to a problem when estimating the difference between a solution to (1.2) and a solution to the limit equation (obtained by replacing the coefficients (but not the right hand side) of (1.2) by their limits as  $t \to \infty$ ). In fact, to say that the difference satisfies a given bound says more for some modes than for others. It is therefore natural to apply an operator which multiplies the Fourier coefficient corresponding to a mode  $\iota$  with  $\langle t \rangle^{-\delta_{\iota}+1} e^{-\kappa_{\iota}t}$  before estimating the difference with respect to a Sobolev norm. A related issue is that of the optimal size of the difference. In Section 10.5 we obtain (10.26). Normalising this estimate as above corresponds to multiplying both sides with  $\langle t \rangle^{-d_1+1} e^{-\kappa_1 t}$ , using the terminology introduced in Lemma 10.17. The time dependent part of the right hand side then takes the form of a polynomial times  $e^{-\beta_{\text{rem}}t}$ . It would be optimistic to hope for more than this (though it is of course possible to discuss the optimal degree of the polynomial). In the transparent setting, it is even optimistic to hope for a time dependence on the right hand side of the form  $\langle t \rangle^N e^{-\beta_{\rm ts} t}$ , where  $\beta_{\rm ts}$  is the constant appearing in (11.6); for weakly transparent, balanced and convergent equations,  $\beta_{ts}$  is defined by (12.23) below. The reason for this is that we have to handle an infinite number of matrices  $A(\iota)$ . This leads to complications that make it natural to replace the polynomial by a factor of the form  $e^{\beta_{\text{mar}}t}$ . On the other hand,  $\beta_{\rm mar} > 0$  can be chosen freely. However, the constants appearing in the estimates depend on  $\beta_{\rm mar}$ , and they can be expected to blow up as  $\beta_{mar} \rightarrow 0$ . A second difference between the silent and transparent settings is that we here have to divide the analysis into two parts. First we need to consider the modes for which  $\mathfrak{g}_{\infty}(\iota)$ , introduced in (11.2), is large. The analysis in that case is based on Sections 11.4 and 11.5. How large  $\mathfrak{g}_{\infty}(\iota)$  has to be depends, among other things, on  $\beta_{\text{mar}}$ . Once the lower bound, say  $\mu_0$ , on  $\mathfrak{g}_{\infty}(\iota)$  has been fixed, it remains to consider the modes satisfying  $0 < \mathfrak{g}_{\infty}(\iota) < \mu_0$  (the modes satisfying  $\mathfrak{g}_{\infty}(\iota) = 0$  correspond to the silent part of the equation and can be analysed as in Chapter 10). However, for these modes, we can appeal to the analysis in Chapter 9. The reason this is possible is that our definition of weak transparency is such that there are only a finite number of matrices  $A(\iota)$  for  $\iota$  satisfying  $0 < \mathfrak{g}_{\infty}(\iota) < \mu_0$ .

**Geometric perspective.** In the present chapter, we derive conclusions based on assumptions that are purely analytic in nature. However, the relevant assumptions follow from quite natural geometric conditions. In Chapter 5, we describe the corresponding geometric perspective, and the reader is encouraged to read that chapter in parallel with the present one.

#### 12.2 Weak transparency, balance and convergence

As indicated in the previous section, several of the arguments and definitions of the present chapter are based on a division of the spatial variables into two groups. We therefore begin by defining this division. In the present chapter we only do so in case (1.2) is an X-non-trivial weakly transparent equation or an X-degenerate weakly transparent equation; cf. Definitions 11.1 and 11.3.

**Definition 12.1.** Assume that (1.2) is an X-non-trivial weakly transparent equation. Assume, moreover, that the integer  $j \in \{1, \ldots, d\}$  has the properties stated in Definition 11.1 and that  $\overline{M}$  is given by (1.3). Then the *silent variables* are all the variables in  $\overline{M}$  except the one corresponding to the j'th S<sup>1</sup>-factor in  $\mathbb{T}^d$ . The *transparent variables* are the variable corresponding to the j'th S<sup>1</sup>-factor in  $\mathbb{T}^d$ . Finally, define  $d_{sil} := d - 1$ ;  $d_{ts} := 1$ ;  $R_{sil} := R$ ;  $R_{ts} := 0$ ;  $j_1 := j$ ;  $\overline{j}_i := i$  for i < j;  $\overline{j}_i = i + 1$  for  $i \ge j$ ; and  $\overline{r}_i = j$  for  $j = 1, \ldots, R$ .

**Remark 12.2.** The reasons for introducing the notation  $d_{\rm sil}$ ,  $d_{\rm ts}$  etc. becomes clear in Definition 12.3 below.

**Definition 12.3.** Assume that (1.2) is an X-degenerate weakly transparent equation. Assume, moreover, that the integers  $j_i$ ,  $i = 1, \ldots, d_{\rm ts}$ , and  $\overline{j}_i$ ,  $i = 1, \ldots, d_{\rm sil}$ , have the properties stated in Definition 11.3, and that  $\overline{M}$  is given by (1.3). Then the *silent variables* are all the variables in  $\overline{M}$  corresponding to the  $\overline{j}_i$ 'th S<sup>1</sup>-factors in  $\mathbb{T}^d$ ,  $i = 1, \ldots, d_{\rm sil}$ , together with all the variables corresponding to  $M_r$  for r's such that  $q_{\infty,r} = 0$ ; the corresponding r's are denoted  $\overline{r}_j$  for  $j = 1, \ldots, R_{\rm sil}$ . Moreover, the *transparent variables* are all the variables in  $\overline{M}$  corresponding to the  $j_i$ 'th S<sup>1</sup>-factors in  $\mathbb{T}^d$ ,  $i = 1, \ldots, d_{\rm ts}$ , together with all the variables or responding to  $M_r$  for r's such that  $q_{\infty,r} > 0$ ; the corresponding to  $M_r$  for r's such that  $q_{\infty,r} > 0$ ; the corresponding to  $M_r$  for r's such that  $q_{\infty,r} > 0$ ; the corresponding to  $M_r$  for r's such that  $q_{\infty,r} > 0$ ; the corresponding to  $M_r$  for r's such that  $q_{\infty,r} > 0$ ; the corresponding r's are denoted  $r_j$  for  $j = 1, \ldots, R_{\rm ts}$ .

Given that (1.2) is an X-non-trivial or an X-degenerate weakly transparent equation, we can introduce the following terminology:

$$\mathfrak{g}_{\rm ts}(\iota,t) := \left(\sum_{k,l=1}^{d_{\rm ts}} g^{j_k j_l}(t) n_{j_k} n_{j_l} + \sum_{j=1}^{R_{\rm ts}} a_{r_j}^{-2}(t) \nu_{r_j,i_{r_j}}^2(\iota)\right)^{1/2},\tag{12.1}$$

$$\mathfrak{g}_{\rm sil}(\iota,t) := \left(\sum_{k,l=1}^{d_{\rm sil}} g^{\bar{j}_k \bar{j}_l}(t) n_{\bar{j}_k} n_{\bar{j}_l} + \sum_{j=1}^{R_{\rm sil}} a_{\bar{r}_j}^{-2}(t) \nu_{\bar{r}_j, i_{\bar{r}_j}}^2(\iota) \right)^{1/2}.$$
(12.2)

In case the upper limit of a sum is zero, it should be understood as being empty. Note that, by assumption,

$$\lim_{t \to \infty} \mathfrak{g}_{\mathrm{ts}}(\iota, t) = \left( \sum_{k,l=1}^{d_{\mathrm{ts}}} g_{\infty}^{j_k j_l} n_{j_k} n_{j_l} + \sum_{j=1}^{R_{\mathrm{ts}}} q_{\infty, r_j} \nu_{r_j, i_{r_j}}^2(\iota) \right)^{1/2} = \mathfrak{g}_{\infty}(\iota),$$

where  $\mathfrak{g}_{\infty}(\iota)$  is given by (11.2). Moreover,  $g_{\infty}^{j_k j_l}$ ,  $k, l = 1, \ldots, d_{ts}$ , are the components of a positive definite matrix, and  $q_{\infty,r_j} > 0$  for  $j = 1, \ldots, R_{ts}$ . It is also of interest to note that  $\mathfrak{g}_{sil}(\iota, t) \to 0$  as  $t \to \infty$ . It would be very convenient if the quantity

$$\mathfrak{g}(\iota,t) - [\mathfrak{g}_{ts}^2(\iota,t) + \mathfrak{g}_{sil}^2(\iota,t)]^{1/2}$$

were zero. However, this is unfortunately not the case, due to the occurrence of cross terms of the form  $g^{j_i \bar{j}_i}(t) n_{j_i} n_{\bar{j}_i}$  in  $\mathfrak{g}^2(\iota, t)$ . Such terms have to be dealt with separately.

It is convenient to divide  $\iota$  into a transparent and a silent part.

**Definition 12.4.** Assume that (1.2) is an X-non-trivial or an X-degenerate weakly transparent equation, and divide the variables of  $\overline{M}$  as described in Definitions 12.1 and 12.3 respectively. Given  $\iota \in \mathcal{I}_{\mathrm{B}}$ , define  $\iota_{\mathrm{ts}} \in \mathcal{I}_{\mathrm{B}}$  as follows: replace  $n_{\overline{j}_k}$  in  $\iota$  by zero for  $k = 1, \ldots, d_{\mathrm{sil}}$ ; and replace  $i_{\overline{r}_k}$  by 0 for  $k = 1, \ldots, R_{\mathrm{sil}}$ . Given  $\iota \in \mathcal{I}_{\mathrm{B}}$ , define  $\iota_{\mathrm{sil}} \in \mathcal{I}_{\mathrm{B}}$  by  $\iota_{\mathrm{sil}} := \iota - \iota_{\mathrm{ts}}$ . The set of  $\iota \in \mathcal{I}_{\mathrm{B}}$  such that  $\iota_{\mathrm{ts}} = 0$  ( $\iota_{\mathrm{ts}} \neq 0$ ) is denoted by  $\mathcal{I}_{\mathrm{B,sil}}$  ( $\mathcal{I}_{\mathrm{B,ts}}$ ).

**Remark 12.5.** It is important to keep in mind that the set of  $\iota_{ts}$  for  $\iota \in \mathcal{I}_B$  is in general *different* from  $\mathcal{I}_{B,ts}$ , but that the set of  $\iota_{sil}$  for  $\iota \in \mathcal{I}_B$  equals  $\mathcal{I}_{B,sil}$ .

With this notation,

$$\mathfrak{g}_{\mathrm{ts}}(\iota, t) = \mathfrak{g}(\iota_{\mathrm{ts}}, t), \quad \mathfrak{g}_{\mathrm{sil}}(\iota, t) = \mathfrak{g}(\iota_{\mathrm{sil}}, t).$$

It is convenient to introduce projections associated with this division of the variables.

**Definition 12.6.** Assume that (1.2) is an X-non-trivial or an X-degenerate weakly transparent equation and define the sets  $\mathcal{I}_{B,sil}$  and  $\mathcal{I}_{B,ts}$  as in Definition 12.4. Let  $s \in \mathbb{R}$ . Then  $\Pi_{ts}, \Pi_{sil} : H^s(\bar{M}) \to H^s(\bar{M})$  are defined as follows. Given  $u \in H^s(\bar{M})$  with Fourier coefficients  $\hat{u}(\iota)$  determined by (24.8),  $\Pi_{ts}u$  is the function whose  $\iota$ 'th Fourier coefficient equals zero if  $\iota \in \mathcal{I}_{B,sil}$  and equals  $\hat{u}(\iota)$  if  $\iota \in \mathcal{I}_{B,ts}$ . Similarly,  $\Pi_{sil}u$  is the function whose  $\iota$ 'th Fourier coefficient equals zero if  $\hat{u}(\iota)$  if  $\iota \in \mathcal{I}_{B,sil}$  and equals zero if  $\iota \in \mathcal{I}_{B,ts}$ .

**Remarks 12.7.** Clearly, the norms of  $\Pi_{ts}$  and  $\Pi_{sil}$  are bounded by 1. Moreover, these operators map smooth functions to smooth functions and we can apply them to functions with values in  $\mathbb{C}^k$ . If  $u \in C^{\infty}(\bar{M} \times I, \mathbb{C}^k)$ , then  $u_{ts}, u_{sil}$  are defined by  $u_{ts}(\cdot, t) = \Pi_{ts}[u(\cdot, t)]$  and  $u_{sil}(\cdot, t) = \Pi_{sil}[u(\cdot, t)]$ . It can then be verified that  $u_{ts}, u_{sil} \in C^{\infty}(\bar{M} \times I, \mathbb{C}^k)$ .

Finally, we are in a position to define what is meant by a weakly transparent, balanced and convergent equation.

**Definition 12.8.** Consider (1.2). Assume the associated metric g to be such that (M, g) is a canonical separable cosmological model manifold. Then (1.2) is said to be *weakly transparent* if the following conditions hold. First, (1.2) is either an X-non-trivial or an X-degenerate weakly transparent equation; cf. Definitions 11.1 and 11.3. Second, there is a constant  $\beta_{sil} > 0$  and a continuous non-negative function  $\mathfrak{e}_{sil} \in L^1([0,\infty))$  such that

$$\dot{\ell}(\iota_{\rm sil}, t) \le -\beta_{\rm sil} + \mathfrak{e}_{\rm sil}(t) \tag{12.3}$$

for all  $\iota \in \mathcal{I}_{\mathrm{B}}$  such that  $\iota_{\mathrm{sil}} \neq 0$  and all  $t \geq 0$ ; here the notation  $\ell$  and  $\iota_{\mathrm{sil}}$  is introduced in (10.1) and Definition 12.4 respectively. Third, there are constants  $K_{\mathrm{ts}}, \eta_{\mathrm{ts}} > 0$  such that

$$|\dot{\ell}(\iota_{\rm ts},t)| + |\sigma(\iota_{\rm ts},t)| \le K_{\rm ts}e^{-\eta_{\rm ts}t} \tag{12.4}$$

for all  $\iota \in \mathcal{I}_{\mathrm{B}}$  such that  $\iota_{\mathrm{ts}} \neq 0$  and all  $t \geq 0$ . Fourth,

$$\|X^{j_k}(t) - X^{j_k}_{\infty}\| \le K_{\rm ts} e^{-\eta_{\rm ts} t}$$
(12.5)

for all  $t \ge 0$  and all  $k = 1, \ldots, d_{ts}$ , where the notation  $j_k$  and  $d_{ts}$  is introduced in Definitions 12.1 12.3.

**Remark 12.9.** As in the case of Definition 10.1, it is to be expected that the requirement that  $\mathfrak{e}_{\rm sil}$  be continuous can be avoided at the expense of additional technicalities in the proofs below. For future reference, we use the notation  $c_{\rm sil} := \|\mathfrak{e}_{\rm sil}\|_1$ .

**Remark 12.10.** The equation (1.2) is said to be weakly transparent, balanced and convergent, if it is weakly transparent in the sense of Definition 12.8, weakly balanced in the sense of Definition 10.1 and weakly convergent in the sense of Definition 10.1.

It is of interest to note that if (1.2) is weakly transparent, balanced and convergent, then restricting (1.2) to solutions which only depend on the silent variables (cf. Definitions 12.1 and 12.3) yields an equation which is weakly silent, balanced and convergent. In other words, the results of Chapter 10 apply. What we need to do in the present chapter is thus to analyse the modes such that  $\iota_{ts} \neq 0$ . Comparing Definition 10.1 with Definition 12.8, it is clear that we still need to define the time corresponding to  $T_{ode}$  appearing in Definition 10.1. Before doing so, let us introduce some terminology.

**Definition 12.11.** Assume that (1.2) is weakly transparent in the sense of Definition 12.8. Let  $\mathcal{I}_{\rm B}$  be the set introduced in connection with (24.3). Define  $C_{\rm ini} \geq 1$  to be the smallest constant such that

$$C_{\text{ini}}^{-1}|\nu(\iota)| \le \mathfrak{g}(\iota,0) \le C_{\text{ini}}|\nu(\iota)|$$
(12.6)

for all  $\iota \in \mathcal{I}_{\mathrm{B}}$ ; here  $\nu(\iota)$  is defined by (24.4). Define  $S_{\mathrm{ts}}$  by

$$S_{ts} := \left[\inf\{\mathfrak{g}(\iota, 0) : \iota \in \mathcal{I}_{B, ts}\}\right]^{-1}, \qquad (12.7)$$

where the notation  $\mathcal{I}_{B,ts}$  is introduced in Definition 12.4.

**Remark 12.12.** The constant  $C_{\text{ini}}$  only depends on  $g^{ij}(0)$  and  $a_r(0)$ . However, the constant  $S_{\text{ts}}$  depends not only on these quantities, but also on the spectrum of the Laplace-Beltrami operator on  $(M_{r_k}, g_{r_k})$  for  $k = 1, \ldots, R_{\text{ts}}$ .

Before defining the constant  $T_{\rm ts}$ , which here plays a role analogous to that of  $T_{\rm ode}$  in Chapter 10, it is convenient to rewrite (24.16) in the form (11.3) for suitable choices of  $A(\iota)$ ,  $A_{\iota,\rm rem}$  and  $F_{\iota}$ , and to derive a general estimate for  $||A_{\iota,\rm rem}(t)||$ .

**Lemma 12.13.** Assume that (1.2) is weakly transparent in the sense of Definition 12.8, weakly balanced in the sense of Definition 10.1 and weakly convergent in the sense of Definition 10.1. Fix  $a \ \iota \in \mathcal{I}_{B,ts}$ . Then (24.16) can be formulated as (11.3), where  $A(\iota)$  is given by

$$A(\iota) := \begin{pmatrix} 0 & \mathfrak{g}_{\infty} \mathrm{Id}_{m} \\ -\mathfrak{g}_{\infty} \mathrm{Id}_{m} - i \mathfrak{g}_{\infty}^{-1} n_{l} X_{\infty}^{l} - \mathfrak{g}_{\infty}^{-1} \zeta_{\infty} & -\alpha_{\infty} \end{pmatrix};$$
(12.8)

v and  $F_{\iota}$  are given by (11.5);  $\mathfrak{g}_{\infty}$  is given by (11.2);  $X_{\infty}^{l}$  is defined by (11.1); and  $\alpha_{\infty}$  and  $\zeta_{\infty}$  are defined by (10.4). Moreover,  $A_{\iota,\text{rem}}$  satisfies the estimate

$$||A_{\iota,\text{rem}}(t)|| \leq Ce^{-\min\{\eta_{\text{ts}},\eta_{\text{ms}}\}t} + Ce^{-\min\{\beta_{\text{sil}},\eta_{\text{ts}}\}t}\mathfrak{g}(\iota_{\text{ts}},0) + Ce^{-\beta_{\text{sil}}t}\mathfrak{g}(\iota_{\text{sil}},0) + Ce$$

for  $t \geq 0$ , where the constant C only depends on  $K_{\rm ts}, C_{\rm mn}, C_{\rm coeff}, C_{\rm ini}, c_{\rm sil}, S_{\rm ts}$  and  $\eta_{\rm ts}$ .

**Remark 12.14.** Here  $K_{ts}$  and  $\eta_{ts}$  are the constants appearing in (12.4) and (12.5);  $C_{mn}$  and  $\eta_{mn}$  are the constants appearing in (10.4);  $C_{coeff}$  is the constant appearing in (10.2); the notation  $C_{ini}$  and  $S_{ts}$  is introduced in Definition 12.11;  $\beta_{sil}$  is the constant appearing in (12.3); and  $c_{sil} := \|\mathbf{e}_{sil}\|_1$ , where  $\mathbf{e}_{sil}$  is the function appearing in (12.3).

**Remarks 12.15.** Given that the assumptions of Lemma 12.13 are satisfied, the only possibility for  $X_{\infty}^{l}$  to be non-zero is that (1.2) is an X-non-trivial weakly transparent equation. Then there is one (and only one)  $j \in \{1, \ldots, d\}$  such that  $X_{\infty}^{j} \neq 0$ , and  $-i\mathfrak{g}_{\infty}^{-1}n_{l}X_{\infty}^{l}$  is one of two possible matrices, irrespective of the value of  $\iota \in \mathcal{I}_{B,ts}$ ; cf. Remarks 11.2. Thus, if the assumptions of Lemma 12.13 are satisfied,  $A(\iota)$  is of the form (11.7), where  $\mu = \mathfrak{g}_{\infty}(\iota)$ ; V is one of the three matrices 0,  $(g_{\infty}^{j_{1}j_{1}})^{-1/2}X_{j_{1}}^{j_{1}}$  or  $-(g_{\infty}^{j_{1}j_{1}})^{-1/2}X_{j_{1}}^{j_{1}}$  (the first possibility only occurs if (1.2) is an Xdegenerate weakly transparent equation, and the latter two possibilities only occur when (1.2) is an X-non-trivial weakly transparent equation, in which case  $j_{1}$  is the number appearing in Definition 12.1);  $W := -\zeta_{\infty}/2$ ; and  $U := -\alpha_{\infty}$ . In addition, due to the requirements appearing in Definitions 11.1 and 11.3, the set  $\mathcal{M} := \{\mathfrak{g}_{\infty}(\iota) : \iota \in \mathcal{I}_{B}\}$  has  $\infty$  as its only accumulation point. Finally, the real parts of the eigenvalues of  $A(\iota)$  converge as  $\mathfrak{g}_{\infty}(\iota) \to \infty$ ; cf. Lemma 11.6.

*Proof.* Due to (12.4),

$$C^{-1}\mathfrak{g}(\iota_{\rm ts},0) \le \mathfrak{g}(\iota_{\rm ts},t) \le C\mathfrak{g}(\iota_{\rm ts},0) \tag{12.10}$$

for  $t \ge 0$ , where  $C \ge 1$  only depends on  $\eta_{ts}$  and  $K_{ts}$ . Taking the limit as  $t \to \infty$  and keeping in mind that  $\mathfrak{g}(\iota_{ts}, t)$  converges to  $\mathfrak{g}_{\infty}(\iota_{ts})$  yields

$$C^{-1}\mathfrak{g}(\iota_{\rm ts},0) \le \mathfrak{g}_{\infty}(\iota_{\rm ts}) \le C\mathfrak{g}(\iota_{\rm ts},0), \tag{12.11}$$

where  $C \ge 1$  only depends on  $\eta_{ts}$  and  $K_{ts}$ . Appealing to (12.3) yields

$$\mathfrak{g}(\iota_{\rm sil}, t) \le e^{-\beta_{\rm sil}t + c_{\rm sil}} \mathfrak{g}(\iota_{\rm sil}, 0) \tag{12.12}$$

for  $t \ge 0$ , where  $c_{\rm sil} := \|\mathbf{e}_{\rm sil}\|_1$ . Before proceeding, note that, using the notation introduced in (24.5) and Definition 12.4,

$$n = \nu_{\rm T}(\iota) = \nu_{\rm T}(\iota_{\rm ts}) + \nu_{\rm T}(\iota_{\rm sil}), \quad n_l g^{0l}(t) = [\nu_{\rm T}(\iota_{\rm ts})]_l g^{0l}(t) + [\nu_{\rm T}(\iota_{\rm sil})] g^{0l}(t).$$

On the other hand (24.17) yields

$$\sigma(\iota_{\rm ts}, t) = \frac{[\nu_{\rm T}(\iota_{\rm ts})]_l g^{0l}(t)}{\mathfrak{g}(\iota_{\rm ts}, t)}, \quad \sigma(\iota_{\rm sil}, t) = \frac{[\nu_{\rm T}(\iota_{\rm sil})]_l g^{0l}(t)}{\mathfrak{g}(\iota_{\rm sil}, t)}$$

Combining the last four equalities yields

$$n_l g^{0l}(t) = \sigma(\iota_{\rm ts}, t) \mathfrak{g}(\iota_{\rm ts}, t) + \sigma(\iota_{\rm sil}, t) \mathfrak{g}(\iota_{\rm sil}, t).$$

Due to this observation, (12.4), (12.10), (12.12) and the fact that the equation is weakly balanced, it follows that

$$|n_l g^{0l}(t)| \leq |\sigma(\iota_{\rm ts}, t)| \cdot \mathfrak{g}(\iota_{\rm ts}, t) + |\sigma(\iota_{\rm sil}, t)| \cdot \mathfrak{g}(\iota_{\rm sil}, t)$$
  
$$\leq C e^{-\eta_{\rm ts} t} \mathfrak{g}(\iota_{\rm ts}, 0) + C e^{-\beta_{\rm sil} t} \mathfrak{g}(\iota_{\rm sil}, 0)$$
(12.13)

for  $t \ge 0$ , where C depends only on  $K_{\rm ts}$ ,  $C_{\rm coeff}$ ,  $c_{\rm sil}$  and  $\eta_{\rm ts}$ , and  $C_{\rm coeff}$  is the constant appearing in (10.2); note that the fact that the equation is weakly balanced implies that  $|\sigma(\iota_{\rm sil}, t)| \le C_{\rm coeff}$ . In case  $g^{0j_k} \equiv 0$  for  $k = 1, \ldots, d_{\rm ts}$ , the first term on the far right of (12.13) can be removed. Similarly, we can estimate

$$\begin{aligned} \|n_{l}X^{l}(t) - n_{l}X_{\infty}^{l}\| &\leq \sum_{k=1}^{d_{\rm ts}} \|n_{j_{k}}X^{j_{k}}(t) - n_{j_{k}}X_{\infty}^{j_{k}}\| + \|X(\iota_{\rm sil},t)\| \cdot \mathfrak{g}(\iota_{\rm sil},t) \\ &\leq K_{\rm ts}e^{-\eta_{\rm ts}t}\sum_{k=1}^{d_{\rm ts}} |n_{j_{k}}| + Ce^{-\beta_{\rm sil}t}\mathfrak{g}(\iota_{\rm sil},0) \end{aligned}$$
(12.14)

where C only depends on  $C_{\text{coeff}}$  and  $c_{\text{sil}}$ ; we have appealed to (12.5) and the fact that  $||X(\iota_{\text{sil}}, t)|| \leq C_{\text{coeff}}$ , an estimate which holds due to the fact that the equation is weakly balanced. In the final estimate, we need to divide both sides of (12.14) by  $\mathfrak{g}_{\infty}(\iota_{\text{ts}})$ . Note, to this end, that

$$[\mathfrak{g}_{\infty}(\iota_{\rm ts})]^{-1} \sum_{k=1}^{d_{\rm ts}} |n_{j_k}| \le C[\mathfrak{g}(\iota_{\rm ts}, 0)]^{-1} \sum_{k=1}^{d_{\rm ts}} |n_{j_k}| \le C,$$

where we have appealed to (12.6) and (12.11) and the constant C only depends on  $K_{\rm ts}$ ,  $C_{\rm ini}$  and  $\eta_{\rm ts}$ . When dividing the second term on the far right hand side of (12.14) by  $\mathfrak{g}_{\infty}(\iota_{\rm ts})$ , we can similarly appeal to (12.7) and (12.11). Summing up yields

$$[\mathfrak{g}_{\infty}(\iota_{\rm ts})]^{-1} \| n_l X^l(t) - n_l X^l_{\infty} \| \le C[e^{-\eta_{\rm ts}t} + e^{-\beta_{\rm sil}t}\mathfrak{g}(\iota_{\rm sil}, 0)]$$
(12.15)

for  $t \ge 0$ , where C only depends on  $K_{\rm ts}, C_{\rm ini}, S_{\rm ts}, c_{\rm sil}, C_{\rm coeff}$  and  $\eta_{\rm ts}$ . In this estimate we tacitly assume  $\iota$  to be such that  $\iota_{\rm ts} \ne 0$ , since it does not make sense otherwise. An alternate version of (12.15) is obtained by replacing  $\mathfrak{g}(\iota_{\rm sil}, 0)$  appearing on the right hand side by  $[\mathfrak{g}(\iota_{\rm ts}, 0)]^{-1}\mathfrak{g}(\iota_{\rm sil}, 0)$ . In that case the constant C appearing in (12.15) does not depend on  $S_{\rm ts}$ . Finally, let us consider

$$\mathfrak{g}^{2}(\iota,t) - \mathfrak{g}^{2}_{\infty}(\iota) = \mathfrak{g}^{2}(\iota_{\rm sil},t) + \mathfrak{g}^{2}(\iota_{\rm ts},t) + \mathfrak{G}_{\rm rem}(\iota,t) - \mathfrak{g}^{2}_{\infty}(\iota), \qquad (12.16)$$

where  $\mathfrak{g}_{\infty}(\iota)$  is defined in (11.2) and

$$\mathfrak{G}_{\rm rem}(\iota,t) := 2 \sum_{k=1}^{d_{\rm ts}} \sum_{l=1}^{d_{\rm sil}} g^{j_k \overline{j}_l} n_{j_k} n_{\overline{j}_l}.$$

The first term on the right hand side of (12.16) can be estimated using (12.12). In order to estimate  $\mathfrak{G}_{rem}$ , note that the Cauchy-Schwarz inequality yields

$$|\mathfrak{G}_{\rm rem}(\iota, t)| \le 2\mathfrak{g}(\iota_{\rm ts}, t)\mathfrak{g}(\iota_{\rm sil}, t) \tag{12.17}$$

154

for all  $t \ge 0$ . In particular

$$|\mathfrak{G}_{\rm rem}(\iota,t)| \le e^{\beta_{\rm sil}t}\mathfrak{g}^2(\iota_{\rm sil},t) + e^{-\beta_{\rm sil}t}\mathfrak{g}^2(\iota_{\rm ts},t)$$
(12.18)

for all  $t \ge 0$ . Combining this observation with (12.16) yields

$$|\mathfrak{g}^{2}(\iota,t) - \mathfrak{g}^{2}_{\infty}(\iota)| \leq |\mathfrak{g}^{2}(\iota_{\mathrm{ts}},t) - \mathfrak{g}^{2}_{\infty}(\iota)| + Ce^{-\beta_{\mathrm{sil}}t}\mathfrak{g}^{2}(\iota_{\mathrm{sil}},0) + Ce^{-\beta_{\mathrm{sil}}t}\mathfrak{g}^{2}(\iota_{\mathrm{ts}},0)$$
(12.19)

for  $t \ge 0$ , where C only depends on  $\eta_{ts}$ ,  $K_{ts}$  and  $c_{sil}$  and we have appealed to (12.10) and (12.12). The first term on the right of (12.19) can be estimated using (12.4). In fact, if  $\iota_{ts} \in \mathcal{I}_B$  is such that  $\iota_{ts} \ne 0$ , then (12.4) can be used to deduce that

$$\left|\frac{\mathfrak{g}_{\infty}(\iota_{\mathrm{ts}})}{\mathfrak{g}(\iota_{\mathrm{ts}},t)}-1\right| \leq Ce^{-\eta_{\mathrm{ts}}t},$$

where C only depends on  $K_{\rm ts}$  and  $\eta_{\rm ts}$ . Combining this estimate with (12.11) yields

$$|\mathfrak{g}^{2}(\iota_{\mathrm{ts}},t) - \mathfrak{g}_{\infty}^{2}(\iota_{\mathrm{ts}})| \leq C e^{-\eta_{\mathrm{ts}}t} \mathfrak{g}^{2}(\iota_{\mathrm{ts}},0)$$
(12.20)

for  $t \ge 0$ , where C only depends on  $K_{\rm ts}$  and  $\eta_{\rm ts}$ . Clearly, this estimate holds even if  $\iota_{\rm ts} = 0$ . To conclude,

$$\mathfrak{g}^{2}(\iota,t) - \mathfrak{g}_{\infty}^{2}(\iota)| \leq Ce^{-\beta_{\mathrm{sil}}t}\mathfrak{g}^{2}(\iota_{\mathrm{sil}},0) + Ce^{-\min\{\beta_{\mathrm{sil}},\eta_{\mathrm{ts}}\}t}\mathfrak{g}^{2}(\iota_{\mathrm{ts}},0)$$

for  $t \ge 0$ , where we have appealed to (12.19) and (12.20); the constant only depends on  $K_{\rm ts}, c_{\rm sil}$ and  $\eta_{\rm ts}$ ; and we have used the fact that  $\mathfrak{g}_{\infty}(\iota) = \mathfrak{g}_{\infty}(\iota_{\rm ts})$ . Dividing this estimate by  $\mathfrak{g}_{\infty}(\iota_{\rm ts})$  and appealing to (12.7) and (12.11) yields

$$[\mathfrak{g}_{\infty}(\iota_{\rm ts})]^{-1}|\mathfrak{g}^{2}(\iota,t) - \mathfrak{g}_{\infty}^{2}(\iota)| \le Ce^{-\beta_{\rm sil}t}\mathfrak{g}^{2}(\iota_{\rm sil},0) + Ce^{-\min\{\beta_{\rm sil},\eta_{\rm ts}\}t}\mathfrak{g}(\iota_{\rm ts},0)$$
(12.21)

for  $t \geq 0$ , where the constant *C* only depends on  $K_{ts}$ ,  $c_{sil}$ ,  $S_{ts}$  and  $\eta_{ts}$  and we, again, assume  $\iota_{ts} \neq 0$ . An alternate version of (12.21) is obtained by replacing  $\mathfrak{g}^2(\iota_{sil}, 0)$  appearing on the right hand side by  $[\mathfrak{g}(\iota_{ts}, 0)]^{-1}\mathfrak{g}^2(\iota_{sil}, 0)$ . In that case the constant *C* appearing in (12.21) does not depend on  $S_{ts}$ . Finally, note that

$$\left[\mathfrak{g}_{\infty}(\iota_{\mathrm{ts}})\right]^{-1} \left\|\zeta(t) - \zeta_{\infty}\right\| \le C e^{-\eta_{\mathrm{mn}}t} \tag{12.22}$$

for  $t \ge 0$ , where C only depends on  $K_{\rm ts}$ ,  $\eta_{\rm ts}$ ,  $C_{\rm mn}$  and  $S_{\rm ts}$ .

Next, the equation (24.16) can be formulated as (11.3), where A is given by (12.8) and v and  $F_{\iota}$  are given by (11.5). Due to (10.4), (12.13), (12.15), (12.21) and (12.22),  $A_{\iota,\text{rem}}$  satisfies the estimate (12.9). The lemma follows.

Finally, we are in a position to define constants  $C_{ts} > 0$  and  $\beta_{ts} > 0$  such that an estimate of the form (11.6) holds.

**Definition 12.16.** Assume that (1.2) is weakly transparent in the sense of Definition 12.8, weakly balanced in the sense of Definition 10.1 and weakly convergent in the sense of Definition 10.1. Fix a  $\iota \in \mathcal{I}_{B,ts}$ . Then  $\beta_{ts}$  and  $T_{ts}$  are defined by

$$\beta_{\rm ts} := \min\{\eta_{\rm ts}, \beta_{\rm sil}, \eta_{\rm mn}\},\tag{12.23}$$

$$T_{\rm ts} := \frac{1}{\min\{\beta_{\rm sil}, \eta_{\rm ts}\}} \ln[1 + |\nu(\iota_{\rm ts})|] + \frac{2}{\beta_{\rm sil}} \ln[1 + |\nu(\iota_{\rm sil})|].$$
(12.24)

Remark 12.17. Combining Lemma 12.13 and Definition 12.16, it is clear that

$$\|A_{\iota,\mathrm{rem}}(t)\| \le C_{\mathrm{ts}} e^{-\beta_{\mathrm{ts}}t} \tag{12.25}$$

for all  $t \ge T_{\rm ts}$ , where  $\bar{t} := t - T_{\rm ts}$  and  $C_{\rm ts}$  only depends on  $K_{\rm ts}, C_{\rm mn}, C_{\rm coeff}, C_{\rm ini}, c_{\rm sil}, S_{\rm ts}$  and  $\eta_{\rm ts}$ .

155

Given a solution u to (1.2), we can divide it into two parts,  $u = u_{\rm sil} + u_{\rm ts}$ ; cf. Remarks 12.7. The results of Chapter 10 apply to  $u_{\rm sil}$ . We can therefore focus on  $u_{\rm ts}$ . Let  $\iota \in \mathcal{I}_{\rm B,ts}$ . Then, due to Lemma 12.13 and Definition 12.16, we can rewrite (24.16) as (11.3), where  $A(\iota)$  is given by (12.8), v and  $F_{\iota}$  are given by (11.5), and  $A_{\iota,\rm rem}$  satisfies the estimate (12.25). It is thus natural to divide the interval  $[0, \infty)$  into  $[0, T_{\rm ts}]$  and  $[T_{\rm ts}, \infty)$ . On the interval  $[T_{\rm ts}, \infty)$ , we can appeal to the results of Chapter 11. What remains to do is thus, first, to analyse how the solution to (24.16) develops in the time period  $[0, T_{\rm ts}]$ ; and, second, to sum up the estimates over all the  $\iota \in \mathcal{I}_{\rm B,ts}$ . On the other hand, in order to be able to estimate the solution in the interval  $[0, T_{\rm ts}]$ , we need to make stronger assumptions than those contained in Definition 12.8. The reason for this is that the conditions (12.3) and (12.4) do not impose any restrictions on, e.g.,  $\dot{g}^{j_k \bar{j}_l}$ . On the other hand, the assumptions made in the statement of Lemma 10.7 are sufficient, since they allow us to apply this lemma.

#### **12.3** Deriving asymptotics

As a next step, we wish to derive asymptotics for solutions to (1.2) for all  $t \ge 0$ . Doing so is more complicated in the present setting than in Section 10.5. The reason for this is that in Section 10.5, the equations of interest are such that all modes have the same asymptotic behaviour. For weakly transparent equations, the asymptotic behaviour is mode dependent, and how to formulate estimates is therefore less obvious; cf. the discussion in the introduction to the present chapter. In order to develop some intuition, it is useful to consider the limit equation defined by (12.26) below. Moreover, it is useful to divide solutions u to (1.2) according to  $u = u_{\rm sil} + u_{\rm ts}$ , where  $u_{\rm sil}$  and  $u_{\rm ts}$  are defined as in Remarks 12.7.

The limit equation. If u is a solution to (1.2), then it is of interest to compare the asymptotics of  $u_{ts}$  with the asymptotics of an appropriately chosen solution to the limit equation. By the *limit equation*, we here mean the equation obtained by letting the coefficients appearing in (1.2) tend to infinity. It can be written

$$u_{tt} - \sum_{j,l=1}^{d} g_{\infty}^{jl} \partial_j \partial_l u - \sum_{r=1}^{R} q_{\infty,r} \Delta_{g_r} u + \alpha_{\infty} u_t + \sum_{j=1}^{d} X_{\infty}^j \partial_j u + \zeta_{\infty} u = f.$$
(12.26)

In order to arrive at this equation, we assume the metric associated with (1.2) to be such that (M,g) is a canonical separable cosmological model manifold; the limits (11.1) to exist; (10.4) to hold; and  $g_{\infty}^{0j}$  to equal zero for  $j = 1, \ldots, d$ . Note that  $g_{\infty}^{0j} = 0, j = 1, \ldots, d$ , holds if one of Definitions 11.1 and 11.3 is satisfied; i.e., if, e.g., (1.2) is a weakly transparent equation. In this context, it is also convenient to introduce a *limiting Laplace-Beltrami operator*:

$$\Delta_{\infty} := \sum_{j,l=1}^{d} g_{\infty}^{jl} \partial_{j} \partial_{l} + \sum_{r=1}^{R} q_{\infty,r} \Delta_{g_{r}}$$
(12.27)

on  $\overline{M}$ . Needless to say, this operator is typically not elliptic on  $\overline{M}$ , since it has degeneracies associated with, e.g., coefficients of the form  $q_{\infty,r}$  that vanish. When considering a solution uto (12.26), it is convenient to divide it into  $u = u_{\rm sil} + u_{\rm ts}$  (as described in Remark 12.7), just as in the case of solutions to (1.2). On the Fourier side,  $\Delta_{\infty}$  corresponds to multiplication with  $-\mathfrak{g}_{\infty}^2(\iota)$ . For  $\iota \in \mathcal{I}_{\rm B,ts}$ , this expression is strictly negative. For that reason, it makes sense to apply  $(-\Delta_{\infty})^{-1/2}$  to  $u_{\rm ts}$  (on the Fourier side, this corresponds to multiplication by  $[\mathfrak{g}_{\infty}(\iota)]^{-1}$ ).

Solving the limit equation. Assume the conditions of Lemma 12.13 to be fulfilled and let (12.26) be the associated limit equation. Let u be a solution to (12.26). Then

$$U_{\rm sil} = A_{\infty} U_{\rm sil} + F_{\rm sil},\tag{12.28}$$

where  $A_{\infty}$  is the matrix given by (10.16),  $U_{\rm sil}$  is the vector valued function with components  $u_{\rm sil}$ and  $\partial_t u_{\rm sil}$ , and  $F_{\rm sil}$  is the vector valued function with components 0 and  $f_{\rm sil}$ , where  $f_{\rm sil}$  is obtained from f as described in Remark 12.7. Solutions to (12.28) can be written

$$U_{\rm sil}(p,t) = e^{A_{\infty}t}U_{\rm sil}(p,0) + \int_0^t e^{A_{\infty}(t-\tau)}F_{\rm sil}(p,\tau)d\tau,$$

#### 12.3. DERIVING ASYMPTOTICS

where  $p \in \overline{M}$ . In particular, the behaviour of  $U_{\rm sil}$  is independent of the Fourier mode. In the case of  $u_{\rm ts}$ , the situation is more complicated. Given  $\iota \in \mathcal{I}_{\rm B,ts}$ , let

$$v_{\rm ts}(\iota,t) := \begin{pmatrix} \mathfrak{g}_{\infty}(\iota)\hat{u}_{\rm ts}(\iota,t) \\ \partial_t\hat{u}_{\rm ts}(\iota,t) \end{pmatrix}, \quad F_{\rm ts,\iota}(t) := \begin{pmatrix} 0 \\ \hat{f}_{\rm ts}(\iota,t) \end{pmatrix}$$

and let  $A(\iota)$  be given by (12.8) (keeping Remarks 12.15 in mind). Here  $f_{ts}$  is obtained from f as described in Remark 12.7 and we use the notation introduced in (24.8). Then

$$v_{\rm ts}(\iota, t) = e^{A(\iota)t} v_{\rm ts}(\iota, 0) + \int_0^t e^{A(\iota)(t-\tau)} F_{\rm ts,\iota}(\tau) d\tau.$$

A formal way of writing this equality so that it is valid for all  $\iota \in \mathcal{I}_{B,ts}$  at once is

$$U_{\rm ts}(\cdot,t) = e^{\mathcal{A}_{\rm ts}t} U_{\rm ts}(\cdot,0) + \int_0^t e^{\mathcal{A}_{\rm ts}(t-\tau)} F_{\rm ts}(\cdot,\tau) d\tau,$$

where

$$F_{\rm ts}(\cdot,t) := \begin{pmatrix} 0 \\ f_{\rm ts}(\cdot,t) \end{pmatrix},$$

$$U_{\rm ts}(\cdot,t) := \begin{pmatrix} (-\Delta_{\infty})^{1/2} u_{\rm ts}(\cdot,t) \\ \partial_t u_{\rm ts}(\cdot,t) \end{pmatrix},$$

$$\mathcal{A}_{\rm ts} := \begin{pmatrix} 0 & \operatorname{Id}_m(-\Delta_{\infty})^{1/2} \\ -\operatorname{Id}_m(-\Delta_{\infty})^{1/2} - (X^l_{\infty}\partial_l + \zeta_{\infty})(-\Delta_{\infty})^{-1/2} & -\alpha_{\infty} \end{pmatrix}.$$
(12.29)

**Norms.** When measuring the norm of, e.g.,  $U_{ts}$ , it is convenient to rescale the individual Fourier coefficients before applying Sobolev norms. The purpose of this rescaling is to normalise the size of the coefficients so that they are asymptotically comparable. Since the analysis of the asymptotics is divided into two parts according to the size of  $\mathfrak{g}_{\infty}(\iota)$ , cf. the introduction, it is also convenient to restrict attention to certain ranges of the frequencies. Given  $1 \leq k \in \mathbb{Z}$ ,  $\psi \in C^{\infty}(\overline{M}, \mathbb{C}^k)$ ,  $s \in \mathbb{R}$ ,  $0 \leq t \in \mathbb{R}$  and  $0 < \mu_0 \in \mathbb{R}$ , we therefore introduce the semi-norm

$$\|\psi\|_{t,s,\mu_0,\pm} := \left(\sum_{\iota \in \mathcal{I}_{\mathsf{B},\mu_0}^{\pm}} \langle \nu(\iota) \rangle^{2s} \langle t \rangle^{-2\delta_{\iota}+2} e^{-2\kappa_{\iota}t} |\hat{\psi}(\iota)|^2 \right)^{1/2}, \tag{12.30}$$

where  $\hat{\psi}$  is defined in analogy with (24.8);  $\kappa_{\iota} := \kappa_{\max}[A(\iota)]$  and  $\delta_{\iota} := d_{\max}[A(\iota), \kappa_{\iota}]$ , cf. Definition 4.3;  $\mathcal{I}^{+}_{\mathrm{B},\mu_{0}}$  is the set of  $\iota \in \mathcal{I}_{\mathrm{B}}$  such that  $\mathfrak{g}_{\infty}(\iota) \geq \mu_{0}$ ; and  $\mathcal{I}^{-}_{\mathrm{B},\mu_{0}}$  is the set of  $\iota \in \mathcal{I}_{\mathrm{B},\mathrm{ts}}$  such that  $\mathfrak{g}_{\infty}(\iota) \geq \mu_{0}$ . Similarly,

$$\|\psi\|_{t,s} := \left(\sum_{\iota \in \mathcal{I}_{\mathrm{B},\mathrm{ts}}} \langle \nu(\iota) \rangle^{2s} \langle t \rangle^{-2\delta_{\iota}+2} e^{-2\kappa_{\iota}t} |\hat{\psi}(\iota)|^2 \right)^{1/2}.$$
(12.31)

In the case of the right hand side of the equation, we also use the semi-norm

$$\|f\|_{\text{ts},s} := \int_0^\infty \left( \sum_{\iota \in \mathcal{I}_{\text{B,ts}}} \langle \nu(\iota) \rangle^{2s} e^{-2(\kappa_\iota - \beta_{\text{mar}})t} |\hat{f}(\iota, t)|^2 \right)^{1/2} dt,$$
(12.32)

where the value of the constant  $\beta_{mar} > 0$  should be clear from the context.

When estimating  $u_{ts}$ , it is convenient to divide the elements of  $\mathcal{I}_{B,ts}$  into the two sets  $\mathcal{I}_{B,\mu_0}^{\pm}$ . For appropriately chosen  $\mu_0$  we can appeal to Lemma 11.17 for  $\mathfrak{g}_{\infty}(\iota) \geq \mu_0$ . For the remaining  $\iota$  there are only a finite number of matrices  $A(\iota)$ . To derive corresponding estimates, it is thus sufficient to appeal to the results of Chapter 10 a finite number of times. We start by combining Lemmas 11.17 and 10.7.

**Lemma 12.18.** Assume that (1.2) is weakly transparent in the sense of Definition 12.8, weakly balanced in the sense of Definition 10.1 and weakly convergent in the sense of Definition 10.1. Assume, finally, that there is a constant  $0 < C_{\ell} \in \mathbb{R}$  and a continuous non-negative function  $\mathfrak{e} \in L^1([0,\infty))$  such that (10.10) holds for all  $0 \neq \iota \in \mathcal{I}_B$  and all  $t \geq 0$ . Fix a  $0 < \beta_{mar} \in \mathbb{R}$  which is small enough, the bound depending only on  $\alpha_{\infty}$ ,  $\zeta_{\infty}$ , the limits appearing in (11.1) and the constant  $\beta_{ts}$  introduced in Definition 12.16. Assume f to be such that  $||f||_{ts,s} < \infty$  for all  $s \in \mathbb{R}$ , where  $|| \cdot ||_{ts,s}$  is defined in (12.32). Then there are constants  $s_{\infty,hom}, s_{\infty,ih} \in [0,\infty), 0 < \mu_0 \in \mathbb{R}$ and  $0 < C \in \mathbb{R}$  such that the following holds. Given a solution u to (1.2), there is a unique  $U_{\infty} \in C^{\infty}(\overline{M}, \mathbb{C}^{2m})$  with the following properties

- the *i*'th Fourier coefficient of  $U_{\infty}$  vanishes unless  $\iota \in \mathcal{I}_{B,\mu_0}^+$ , where the notation  $\mathcal{I}_{B,\mu_0}^+$  is introduced in connection with (12.30),
- if  $\iota \in \mathcal{I}^+_{\mathrm{B},\mu_0}$ , then the  $\iota$ 'th Fourier coefficient of  $U_\infty$  belongs to  $E_\iota$ , where  $E_\iota$  is the first generalised eigenspace in the  $(\beta_{\mathrm{ts}} \beta_{\mathrm{mar}}), A(\iota)$ -decomposition of  $\mathbb{C}^{2m}$ ,
- the estimate

$$\left\| \begin{pmatrix} (-\Delta_{\infty})^{1/2} u(\cdot,t) \\ \partial_t u(\cdot,t) \end{pmatrix} - e^{\mathcal{A}_{ts}t} U_{\infty} - \int_0^t e^{\mathcal{A}_{ts}(t-\tau)} \begin{pmatrix} 0 \\ f(\cdot,\tau) \end{pmatrix} d\tau \right\|_{t,s,\mu_0,+}$$

$$\leq C e^{-(\beta_{ts}-\beta_{mar})t} [\|u_t(\cdot,0)\|_{(s+s_{\infty,hom})} + \|u(\cdot,0)\|_{(s+s_{\infty,hom}+1)} + \|f\|_{ts,s+s_{\infty,ih}}]$$

$$(12.33)$$

holds for  $t \ge 0$  and  $s \in \mathbb{R}$ , where  $\Delta_{\infty}$  is defined in (12.27),  $\mathcal{A}_{ts}$  is defined in (12.29) and  $\|\cdot\|_{t,s,\mu_0,+}$  is defined in (12.30).

Moreover,

$$\|U_{\infty}\|_{(s)} \le C \|u_t(\cdot, 0)\|_{(s+s_{\infty, \text{hom}})} + C \|u(\cdot, 0)\|_{(s+s_{\infty, \text{hom}}+1)} + C \|f\|_{\text{ts}, s+s_{\infty, \text{ih}}}$$
(12.34)

for all  $s \in \mathbb{R}$ .

**Remark 12.19.** The constant *C* only depends on  $C_{\text{ini}}$ ,  $C_{\text{mn}}$ ,  $C_{\text{coeff}}$ ,  $S_{\text{ts}}$ ,  $K_{\text{ts}}$ ,  $\eta_{\text{ts}}$ ,  $c_{\mathfrak{e}}$ ,  $\mathcal{C}_{\ell}$ ,  $\beta_{\text{sil}}$ , the limits appearing in (11.1),  $\alpha_{\infty}$ ,  $\zeta_{\infty}$ ,  $\beta_{\text{ts}}$  and  $\beta_{\text{mar}}$ ;  $\mu_0$  only depends on  $\beta_{\text{ts}}$ ,  $\beta_{\text{mar}}$ ,  $\alpha_{\infty}$ ,  $\zeta_{\infty}$  and the limits appearing in (11.1); and  $s_{\infty,\text{hom}}$  and  $s_{\infty,\text{ih}}$  only depend on  $C_{\text{coeff}}$ ,  $C_{\ell}$ ,  $\eta_{\text{ts}}$ ,  $\eta_{\text{mn}}$ ,  $\beta_{\text{sil}}$ ,  $\alpha_{\infty}$ ,  $\zeta_{\infty}$  and the limits appearing in (11.1). Here  $c_{\mathfrak{e}} := \|\mathfrak{e}\|_1$  and  $c_{\text{sil}} := \|\mathfrak{e}_{\text{sil}}\|_1$ .

Remark 12.20. The terms

$$e^{\mathcal{A}_{\rm ts}t}U_{\infty} + \int_0^t e^{\mathcal{A}_{\rm ts}(t-\tau)} \begin{pmatrix} 0\\ f(\cdot,\tau) \end{pmatrix} d\tau$$

appearing on the left hand side of (12.33) correspond to a solution to the limit equation (12.26).

**Remark 12.21.** Due to the statement of the lemma,  $U_{\infty}$  is uniquely determined by the solution. Moreover, the map from the initial data of u to  $U_{\infty}$  is continuous in the  $C^{\infty}$ -topology.

Proof. The proof is based on a combination of Lemmas 10.7 and 11.17. However, the justification that we are allowed to appeal to Lemma 11.17 depends on which of Definitions 11.1 and 11.3 is satisfied. If (1.2) is an X-non-trivial weakly transparent equation, then there are only two possibilities for  $-\mathfrak{g}_{\infty}^{-1}n_l X_{\infty}^l$  appearing in the bottom left component of the right hand side of (12.8), namely  $\pm (g_{\infty}^{j_1j_1})^{-1/2} X_{\infty}^{j_1}$ ; cf. Remark 12.15. When appealing to Lemma 11.17, the matrices U, V, W are thus given by  $U := -\alpha_{\infty}, V := \mp (g_{\infty}^{j_1j_1})^{-1/2} X_{\infty}^{j_1}$  and  $W := -\zeta_{\infty}/2$ . Since there are two possibilities for V, we, strictly speaking, need to apply Lemma 11.17 twice. If (1.2) is an X-degenerate weakly transparent equation, then  $U := -\alpha_{\infty}, V := 0$  and  $W := -\zeta_{\infty}/2$ , and we only need to appeal to Lemma 11.17 once. However, it is of interest to note that, irrespective of whether (1.2) is an X-non-trivial weakly transparent equation or an X-degenerate weakly transparent equation, the matrices U, V and W only depend on  $\alpha_{\infty}$ ,  $\zeta_{\infty}$  and the limits appearing in (11.1).

When appealing to Lemma 11.17, we let  $\mu(\iota) := \mathfrak{g}_{\infty}(\iota)$  for  $\iota \in \mathcal{I}_{\mathrm{B}}$ . Note that then  $\mu(\iota)$  is strictly positive if and only if  $\iota \in \mathcal{I}_{\mathrm{B,ts}}$ . Thus the notation  $\mathcal{I}_{\mathrm{B,ts}}$  used in the present chapter coincides with the notation  $\mathcal{I}_{\mathrm{B,ts}}$  used in the statement of Lemma 11.17. We have already clarified what the matrices U, V and W are, and due to Remark 12.15, it is clear that  $A(\iota)$  is of the form (11.73) for  $\iota \in \mathcal{I}_{\mathrm{B,ts}}$ . Turning to the  $A_{\iota,\mathrm{rem}}$  and  $F_{\iota}$  appearing in Lemma 11.17, we have already defined them in the case of  $\iota \in \mathcal{I}_{\mathrm{B,ts}}$ ; cf. Lemma 12.13. The constants  $\beta_{\mathrm{ts}}$ ,  $C_{\mathrm{ts}}$  and  $T_{\mathrm{ts}}$  are all introduced in Definition 12.16 and Remark 12.17. The upper bound on the constant  $\beta_{\mathrm{mar}}$ appearing in Lemma 11.17 only depends on U, V and  $\beta_{\mathrm{ts}}$ . In the present setting, that means that it only depends on  $\alpha_{\infty}, \zeta_{\infty}$ , the limits appearing in (11.1) and  $\beta_{\mathrm{ts}}$ , as stated in the present lemma. Finally, the condition  $||F_{\iota}||_{\mathrm{Ima}} < \infty$  clearly holds for all  $\iota \in \mathcal{I}_{\mathrm{B,ts}}$ . Thus Lemma 11.17 applies and we can appeal to the conclusions of that lemma. If (1.2) is an X-non-trivial weakly transparent equation, we apply Lemma 11.17 twice, and there are two corresponding  $\mu_0$  values, and two corresponding constants C. In that case, the constant  $\mu_0$  appearing in the statement of the present lemma is the largest of these two  $\mu_0$ -values, and similarly for C.

**Rough estimate.** Let us start by obtaining a rough estimate. Due to (11.88),

$$|v(\iota,t)| \le C e^{(\kappa_{\iota} + \beta_{\max}/2)\bar{t}} |v(\iota,T_{\rm ts})| + C e^{\beta_{\max}\bar{t}/2} \int_{T_{\rm ts}}^{t} e^{\kappa_{\iota}(t-s) + \beta_{\max}\bar{s}/2} |F_{\iota}(s)| ds$$
(12.35)

for  $t \geq T_{\rm ts}$  and  $\mathfrak{g}_{\infty}(\iota) \geq \mu_0$ , where *C* and  $\mu_0$  have the same dependence as in Remark 12.19. Note that it is not immediately obvious that |v| and  $\mathcal{E}^{1/2}$  are equivalent on  $[T_{\rm ts}, \infty)$ , where the notation  $\mathcal{E}$  is introduced in (10.11). In order to see that this is the case, note first that (12.10) and (12.11) hold, where the constant *C* only depends on  $K_{\rm ts}$  and  $\eta_{\rm ts}$ . Thus there is a constant  $C \geq 1$ , depending only on  $S_{\rm ts}$ ,  $K_{\rm ts}$  and  $\eta_{\rm ts}$ , such that

$$C^{-1}[|\dot{z}|^2 + \mathfrak{g}_{\rm ts}^2|z|^2 + |z|^2] \le |v|^2 \le C[|\dot{z}|^2 + \mathfrak{g}_{\rm ts}^2|z|^2 + |z|^2]$$
(12.36)

on  $[T_{ts}, \infty)$ . Moreover, for  $t \ge T_{ts}$ , (12.6), (12.7), (12.10), (12.12) and (12.24) yield

$$\mathfrak{g}(\iota_{\mathrm{sil}},t) \leq e^{-\beta_{\mathrm{sil}}t+c_{\mathrm{sil}}}\mathfrak{g}(\iota_{\mathrm{sil}},0) \leq e^{-\beta_{\mathrm{sil}}t+c_{\mathrm{sil}}}C_{\mathrm{ini}}|\nu(\iota_{\mathrm{sil}})| \leq C_{\mathrm{ini}}e^{c_{\mathrm{sil}}} \leq C_{\mathrm{ini}}e^{c_{\mathrm{sil}}}S_{\mathrm{ts}}\mathfrak{g}(\iota_{\mathrm{ts}},0) \leq C\mathfrak{g}(\iota_{\mathrm{ts}},t),$$
(12.37)

where C only depends on  $C_{\rm ini}, c_{\rm sil}, K_{\rm ts}, S_{\rm ts}$  and  $\eta_{\rm ts}$ . Appealing to (12.17) and (12.37) yields

$$\begin{aligned} \mathfrak{g}^{2}(\iota,t) &= \mathfrak{g}^{2}(\iota_{\mathrm{ts}},t) + \mathfrak{g}^{2}(\iota_{\mathrm{sil}},t) + \mathfrak{G}_{\mathrm{rem}}(\iota,t) \\ &\geq \frac{1}{2}\mathfrak{g}^{2}(\iota_{\mathrm{ts}},t) - \mathfrak{g}^{2}(\iota_{\mathrm{sil}},t) \geq \frac{1}{2}\mathfrak{g}^{2}(\iota_{\mathrm{ts}},t) - C \end{aligned}$$

for all  $t \geq T_{ts}$ , where C only depends on  $C_{ini}$  and  $c_{sil}$ . In particular,

$$\mathfrak{g}^2(\iota_{\mathrm{ts}},t) + 1 \le C[\mathfrak{g}^2(\iota,t) + 1]$$

for all  $t \ge T_{ts}$ , where C only depends on  $C_{ini}$  and  $c_{sil}$ . Combining this estimate with (12.36) and (12.37) yields a constant  $C \ge 1$  such that

$$C^{-2}\mathcal{E}(t) \le |v(t)|^2 \le C^2 \mathcal{E}(t)$$
 (12.38)

for  $t \ge T_{\rm ts}$ , where C has the same dependence as in the case of (12.37). Combining this estimate with (12.35) yields

$$\mathcal{E}^{1/2}(t) \le C e^{(\kappa_{\iota} + \beta_{\max}/2)\overline{t}} \mathcal{E}^{1/2}(T_{\rm ts}) + C e^{\beta_{\max}\overline{t}/2} \int_{T_{\rm ts}}^{t} e^{\kappa_{\iota}(t-s) + \beta_{\max}\overline{s}/2} |F_{\iota}(s)| ds \tag{12.39}$$

for  $t \geq T_{ts}$  and  $\mathfrak{g}_{\infty}(\iota) \geq \mu_0$ , where C and  $\mu_0$  have the same dependence as in Remark 12.19. Appealing to (10.12) yields

$$Ce^{(\kappa_{\iota}+\beta_{\max}/2)\overline{t}}\mathcal{E}^{1/2}(T_{ts}) \leq Ce^{c_{\mathfrak{e}}}e^{(\kappa_{\iota}+\beta_{\max}/2)t}e^{\delta_{ts}T_{ts}}\mathcal{E}^{1/2}(0) + Ce^{c_{\mathfrak{e}}}e^{\delta_{ts}T_{ts}}\int_{0}^{T_{ts}}e^{(\kappa_{\iota}+\beta_{\max}/2)(t-s)}|F_{\iota}(s)|ds,$$

$$(12.40)$$

where  $c_{\mathfrak{e}} := \|\mathfrak{e}\|_1$ , C has the same dependence as in the case of (12.39), and

$$\delta_{\iota,\mathrm{ts}} := \max\{\eta_{\mathrm{bal}} - \kappa_{\iota}, 0\}, \quad \delta_{\mathrm{ts}} := \sup_{\iota \in \mathcal{I}_{\mathrm{B},\mathrm{ts}}} \delta_{\iota,\mathrm{ts}}.$$

Note also that  $\eta_{\text{bal}}$  depends only on  $C_{\ell}$  and  $C_{\text{coeff}}$ . In order to estimate  $e^{\delta_{\text{ts}}T_{\text{ts}}}$  from above, note that (12.24) implies that

$$e^{\delta_{\rm ts}T_{\rm ts}} = [1 + |\nu(\iota_{\rm ts})|]^{\delta_{\rm ts}/\min\{\eta_{\rm ts},\beta_{\rm sil}\}} [1 + |\nu(\iota_{\rm sil})|]^{2\delta_{\rm ts}/\beta_{\rm sil}} \le C \langle \nu(\iota) \rangle^{s_0}, \tag{12.41}$$

where C only depends on  $C_{\ell}$ ,  $C_{\text{coeff}}$ ,  $\alpha_{\infty}$ ,  $\zeta_{\infty}$ , the limits appearing in (11.1),  $\beta_{\text{sil}}$  and  $\eta_{\text{ts}}$ ; and

$$s_0 := [(\min\{\eta_{\rm ts}, \beta_{\rm sil}\})^{-1} + 2\beta_{\rm sil}^{-1}]\delta_{\rm ts}.$$

In particular,  $s_0$  only depends on  $\eta_{\text{ts}}$ ,  $\beta_{\text{sil}}$ ,  $C_{\ell}$ ,  $C_{\text{coeff}}$ , the limits appearing in (11.1),  $\alpha_{\infty}$  and  $\zeta_{\infty}$ . Combining (12.39), (12.40) and (12.41) yields

$$\mathcal{E}^{1/2}(t) \leq C e^{(\kappa_{\iota} + \beta_{\max}/2)t} \langle \nu(\iota) \rangle^{s_0} \mathcal{E}^{1/2}(0) + C e^{(\kappa_{\iota} + \beta_{\max}/2)t} \langle \nu(\iota) \rangle^{s_0} \int_0^t e^{-(\kappa_{\iota} - \beta_{\max}/2)s} |F_{\iota}(s)| ds$$
(12.42)

for  $t \ge T_{\rm ts}$  and  $\mathfrak{g}_{\infty}(\iota) \ge \mu_0$ , where C and  $\mu_0$  have the same dependence as in Remark 12.19. In order to obtain an estimate for  $0 \le t \le T_{\rm ts}$ , we appeal to (10.12) again. This yields

$$\begin{aligned} \mathcal{E}^{1/2}(t) &\leq e^{c_{\mathfrak{e}} + \eta_{\text{bal}} t} \mathcal{E}^{1/2}(0) + e^{c_{\mathfrak{e}}} \int_{0}^{t} e^{\eta_{\text{bal}}(t-u)} |F_{\iota}(u)| du \\ &\leq e^{c_{\mathfrak{e}}} e^{\delta_{\text{ts}} T_{\text{ts}}} e^{(\kappa_{\iota} + \beta_{\text{mar}}/2)t} \mathcal{E}^{1/2}(0) + e^{c_{\mathfrak{e}}} e^{\delta_{\text{ts}} T_{\text{ts}}} \int_{0}^{t} e^{(\kappa_{\iota} + \beta_{\text{mar}}/2)(t-u)} |F_{\iota}(u)| du \end{aligned}$$

for  $0 \le t \le T_{\text{ts}}$ . Keeping (12.41) in mind, this estimate yields the conclusion that (12.42) holds for  $0 \le t \le T_{\text{ts}}$ . Thus (12.42) holds for all  $t \ge 0$ .

Asymptotics, introduction. In order to obtain asymptotics, we wish to appeal to Lemma 11.17. However, before we do so, we need to impose restrictions on  $\beta_{\text{mar}}$ . Let, to this end,  $\lambda_{j,\pm}$  be given by Definition 11.16, where U and V are constructed from  $\alpha_{\infty}$ ,  $\zeta_{\infty}$  and the limits appearing in (11.1) as described in the beginning of the proof. Let  $\kappa_{j,\pm}$  denote the real parts of  $\lambda_{j,\pm}$  and let  $\kappa_{\infty}$ denote the largest of the  $\kappa_{j,\pm}$ . Given  $\beta_{\text{ts}} > 0$ , there are two possibilities. Either there is a  $\kappa_{j,\pm}$ , say  $\kappa_c$ , such that  $\kappa_c = \kappa_{\infty} - \beta_{\text{ts}}$ , or there is no such  $\kappa_c$ . Regardless, there is a  $\beta_{\text{spe}} > 0$  (depending only on  $\alpha_{\infty}$ ,  $\zeta_{\infty}$ , the limits appearing in (11.1) and  $\beta_{\text{ts}}$ ) such that there is no  $\kappa_{j,\pm}$  in the interval  $(\kappa_{\infty} - \beta_{\text{ts}}, \kappa_{\infty} - \beta_{\text{ts}} + 2\beta_{\text{spe}})$ . Fix  $0 < \beta_{\text{mar}} < \beta_{\text{spe}}$  (we here also assume  $\beta_{\text{mar}}$  to be small enough that Lemma 11.17 can be applied). Then, for  $\mu_0$  large enough (the bound depending only on  $\alpha_{\infty}$ ,  $\zeta_{\infty}$ , the limits appearing in (11.1),  $\beta_{\text{ts}}$  and  $\beta_{\text{mar}}$ ) there is a constant C (with the same dependence) such that if  $\mathfrak{g}_{\infty}(\iota) \geq \mu_0$ , then

- the generalised eigenspaces of  $A(\iota)$  of which  $E_{\iota,\beta_{ts}}^{\infty}$  consists are all the generalised eigenspaces such that the corresponding eigenvalues have real parts  $> \kappa_{\iota} \beta_{ts} + \beta_{mar}$ ,
- if  $e_{\iota,\beta_{ts}}^{\infty}$  is the space introduced in the statement of Lemma 11.22, then

$$|e^{A(\iota)t}\eta| \le Ce^{(\kappa_\iota - \beta_{\rm ts} + \beta_{\rm mar}/2)t}|\eta| \tag{12.43}$$

for all  $t \geq 0$  and all  $\eta \in e_{\iota,\beta_{ts}}^{\infty}$ .

160

Appealing to Lemma 11.17 yields a  $v_{\infty} \in E_{\iota,\beta_{\text{ts}}}^{\infty}$  such that (11.75) holds for  $t \geq T_{\text{ts}}$ , where  $E_{\iota,\beta_{\text{ts}}}^{\infty}$  is given by Definition 11.16 (even though  $v_{\infty}$  depends on  $\iota$ , we omit reference to this dependence for the sake of brevity). Introducing

$$v_{\infty,\text{mod}} := v_{\infty} - \int_{0}^{T_{\text{ts}}} e^{-A(\iota)\tau} F_{\iota}(\tau) d\tau,$$
 (12.44)

the estimate (11.75) can be reformulated as

$$\left| v(t) - e^{A(\iota)t} v_{\infty, \text{mod}} - \int_0^t e^{A(\iota)(t-s)} F_\iota(s) ds \right|$$
  
$$\leq C e^{\kappa_\iota t - (\beta_{\text{ts}} - \beta_{\text{mar}})\bar{t}} [e^{-\kappa_\iota T_{\text{ts}}} |v(T_{\text{ts}})| + \|F_\iota\|_{\text{lma}}]$$
(12.45)

for  $t \geq T_{ts}$  and  $\mathfrak{g}_{\infty}(\iota) \geq \mu_0$ , where  $\mu_0$  and C have the same dependence as in Remark 12.19. As in the proof of Lemma 10.17, we need to project  $v_{\infty,mod}$  onto the correct subspace. However, we only write down the corresponding estimates at the end of the argument.

The objects to be estimated. In order to obtain the desired conclusion, we proceed as in the proof of Lemma 10.17. As in that proof, we thus need to estimate

$$e^{(\beta_{\rm ts}-\kappa_{\iota})t}|v(t)|, \quad e^{(\beta_{\rm ts}-\kappa_{\iota})t}|e^{A(\iota)t}v_{\infty}|, \quad \left|e^{(\beta_{\rm ts}-\kappa_{\iota})t}\int_{t}^{T_{\rm ts}}e^{A(\iota)(t-\tau)}F_{\iota}(\tau)d\tau\right|$$
(12.46)

for  $t \in [0, T_{ts}]$ ; this yields the desired estimate on the interval  $[0, T_{ts}]$ . In order to obtain a good estimate for  $t \ge T_{ts}$ , we need to estimate

$$e^{(\beta_{\rm ts}-\kappa_\iota)T_{\rm ts}}|v(T_{\rm ts})|, \quad e^{\beta_{\rm ts}T_{\rm ts}}\|F_\iota\|_{\rm lma}.$$
(12.47)

Finally, we wish to estimate  $v_{\infty,\text{mod}}$  introduced in (12.44). Note that if we have estimated the first expression appearing in (12.46) on  $[0, T_{\text{ts}}]$ , then we have also estimated the first expression appearing in (12.47).

Estimating the solution. Note, to begin with, that

$$\mathfrak{g}_{\infty}^{2}(\iota) \leq C\mathfrak{g}^{2}(\iota_{\mathrm{ts}}, t) \leq C[\mathfrak{g}^{2}(\iota, t) + \mathfrak{g}^{2}(\iota_{\mathrm{sil}}, t)]$$
(12.48)

for  $t \ge 0$ , where C only depends on  $K_{\rm ts}$  and  $\eta_{\rm ts}$  and we have appealed to (12.10), (12.11) and the fact that  $\mathfrak{g}_{\infty}(\iota) = \mathfrak{g}_{\infty}(\iota_{\rm ts})$ . We have also used the fact that (12.17) yields

$$|\mathfrak{G}_{\rm rem}(\iota,t)| \le \frac{1}{2}\mathfrak{g}^2(\iota_{\rm ts},t) + 2\mathfrak{g}^2(\iota_{\rm sil},t)$$
(12.49)

for  $t \ge 0$ ; note that (12.49) implies that

$$\mathfrak{g}^{2}(\iota,t) \geq \frac{1}{2}\mathfrak{g}^{2}(\iota_{\mathrm{ts}},t) - \mathfrak{g}^{2}(\iota_{\mathrm{sil}},t)$$

for  $t \ge 0$ . The estimate (12.48) yields

$$|v(\iota,t)| \le C[1 + \mathfrak{g}^2(\iota_{\rm sil},t)]^{1/2} \mathcal{E}^{1/2}(\iota,t) \le C\langle \nu(\iota) \rangle \mathcal{E}^{1/2}(\iota,t)$$

for  $t \ge 0$ , where C only depends on  $K_{\rm ts}$ ,  $C_{\rm ini}$ ,  $c_{\rm sil}$  and  $\eta_{\rm ts}$ . Combining this estimate with Lemma 10.7 yields

$$e^{(\beta_{\rm ts}-\kappa_{\iota})t}|v(\iota,t)| \leq Ce^{c_{\mathfrak{c}}}\langle\nu(\iota)\rangle[e^{\gamma_{\rm hom,1}(\iota)t}\mathcal{E}^{1/2}(\iota,0) + e^{\gamma_{\rm ih,1}(\iota)t}\int_{0}^{t}e^{-\kappa_{\iota}\tau}|\hat{f}(\iota,\tau)|d\tau]$$
(12.50)

for  $t \in [0, T_{ts}]$ , where  $\gamma_{\text{hom},1}(\iota) := \max\{\beta_{ts} + \eta_{\text{bal}} - \kappa_{\iota}, 0\}$ ;  $\gamma_{\text{ih},1}(\iota) := \beta_{ts} + \max\{\eta_{\text{bal}} - \kappa_{\iota}, 0\}$ ; and the constant C only depends on  $K_{ts}$ ,  $\eta_{ts}$ ,  $C_{\text{ini}}$  and  $c_{\text{sil}}$ . For future reference, it is useful to keep in

mind that the same estimate holds with  $\beta_{ts}$  replaced by 0. Due to estimates of the form (12.41), it then follows that

$$\langle \nu(\iota) \rangle^{s} \sup_{t \in [0, T_{\rm ts}]} e^{(\beta_{\rm ts} - \kappa_{\iota})t} |\nu(\iota, t)|$$

$$\leq C \langle \nu(\iota) \rangle^{s + s_{\rm hom, 1}} \mathcal{E}^{1/2}(\iota, 0) + C \langle \nu(\iota) \rangle^{s + s_{\rm ih, 1}} \int_{0}^{T_{\rm ts}} e^{-\kappa_{\iota}\tau} |\hat{f}(\iota, \tau)| d\tau$$

$$(12.51)$$

for  $\iota \in \mathcal{I}^+_{\mathrm{B},\mu_0}$ , where

$$s_{\text{hom},1} := 1 + [(\min\{\eta_{\text{ts}}, \beta_{\text{sil}}\})^{-1} + 2\beta_{\text{sil}}^{-1}] \sup_{\iota \in \mathcal{I}_{B,\mu_0}^+} \gamma_{\text{hom},1}(\iota),$$
  
$$s_{\text{ih},1} := 1 + [(\min\{\eta_{\text{ts}}, \beta_{\text{sil}}\})^{-1} + 2\beta_{\text{sil}}^{-1}] \sup_{\iota \in \mathcal{I}_{B,\mu_0}^+} \gamma_{\text{ih},1}(\iota)$$

and the constant C has the same dependence as in Remark 12.19. Again, this estimate also holds when  $\beta_{ts}$  is replaced by zero. One particular consequence of this estimate is

$$\langle \nu(\iota) \rangle^{s} e^{(\beta_{\rm ts} - \kappa_{\iota})T_{\rm ts}} | v(\iota, T_{\rm ts}) | + \langle \nu(\iota) \rangle^{s} e^{\beta_{\rm ts}T_{\rm ts}} \| F_{\iota} \|_{\rm lma}$$

$$\leq C \langle \nu(\iota) \rangle^{s+s_{\rm hom,1}} \mathcal{E}^{1/2}(\iota, 0) + C \langle \nu(\iota) \rangle^{s+s_{\rm ih,1}} \| \widehat{f} \|_{\rm lma}$$

$$(12.52)$$

for  $\iota \in \mathcal{I}^+_{\mathrm{B},\mu_0}$ , where the constant *C* has the same dependence as in Remark 12.19 and we have used the notation (11.74). Moreover, the analogous estimate with  $\beta_{\mathrm{ts}}$  replaced by 0 holds.

Estimating the limiting value. Next, let us turn to the problem of estimating the second expression appearing in (12.46). To begin with,  $v_{\infty}$  is such that (11.76) and (11.77) are satisfied, where  $u_{\infty} \in E_{\iota,\beta_{\text{ts}}}^{\infty}$ . Appealing to Lemma 11.22, it is clear that for  $\mu_0$  large enough, the bound depending only on the limits appearing in (11.1),  $\alpha_{\infty}, \zeta_{\infty}$  and  $\beta_{\text{ts}}$ ,

$$|v_{\infty}| \le C e^{-(\kappa_{\iota} - \beta_{\rm ts})T_{\rm ts}} |u_{\infty}| \tag{12.53}$$

for  $\iota \in \mathcal{I}_{\mathrm{B},\mu_0}^+$ , where *C* only depends on the limits appearing in (11.1),  $\alpha_{\infty}$ ,  $\zeta_{\infty}$  and  $\beta_{\mathrm{ts}}$ . On the other hand,  $u_{\infty} \in E_{\iota,\beta_{\mathrm{ts}}}^{\infty}$  satisfies the estimate

$$|u_{\infty}| \le C[|v(T_{\rm ts})| + e^{\kappa_{\iota} T_{\rm ts}} ||F_{\iota}||_{\rm lma}], \qquad (12.54)$$

where C has the same dependence as in Remark 12.19. Combining (12.52), (12.53) and (12.54) yields

$$\langle \nu(\iota) \rangle^s |v_{\infty}| \le C \langle \nu(\iota) \rangle^{s+s_{\text{hom},1}} \mathcal{E}^{1/2}(0) + C \langle \nu(\iota) \rangle^{s+s_{\text{ih},1}} \|\hat{f}\|_{\text{lma}}$$
(12.55)

for  $\iota \in \mathcal{I}^+_{\mathrm{B},\mu_0}$ , where C has the same dependence as in Remark 12.19. Moreover, the constants  $s_{\mathrm{hom},1}$  and  $s_{\mathrm{ih},1}$  are defined in connection with (12.51).

In order to estimate the second expression in (12.46), it is of interest to note that

$$e^{A(\iota)t}v_{\infty} = e^{-A(\iota)(T_{\rm ts}-t)}u_{\infty}.$$
(12.56)

Here we are only interested in the case that  $t \in [0, T_{ts}]$ . The argument which led to the estimate (12.53) thus yields

$$e^{(\beta_{\rm ts}-\kappa_\iota)t}|e^{A(\iota)t}v_{\infty}| \le e^{(\beta_{\rm ts}-\kappa_\iota)t}Ce^{-(\kappa_\iota-\beta_{\rm ts})(T_{\rm ts}-t)}|u_{\infty}|$$

for  $\iota \in \mathcal{I}_{\mathrm{B},\mu_0}^+$  and  $t \in [0, T_{\mathrm{ts}}]$ , where C and the lower bound on  $\mu_0$  only depend on the limits appearing in (11.1),  $\alpha_{\infty}$ ,  $\zeta_{\infty}$  and  $\beta_{\mathrm{ts}}$ . Combining this estimate with (12.52) and (12.54) yields

$$\langle \nu(\iota) \rangle^s \sup_{t \in [0, T_{\rm ts}]} |e^{(\beta_{\rm ts} - \kappa_\iota)t} e^{A(\iota)t} v_\infty| \le C \langle \nu(\iota) \rangle^{s + s_{\rm hom,1}} \mathcal{E}^{1/2}(0) + C \langle \nu(\iota) \rangle^{s + s_{\rm ih,1}} \|\hat{f}\|_{\rm lma}$$
(12.57)

for  $\iota \in \mathcal{I}^+_{\mathrm{B},\mu_0}$ , where C and  $\mu_0$  have the same dependence as in Remark 12.19.

Estimating the right hand side. Let us turn to

$$\int_{t}^{T_{\rm ts}} e^{A(\iota)(t-\tau)} F_{\iota}(\tau) d\tau$$

Due to Lemma 11.23 and Remark 11.24, we can, for a fixed  $\eta_{mar} > 0$ , estimate

$$\|e^{A(\iota)(t-\tau)}\| < Ce^{(\kappa_b - \eta_{\max})(t-\tau)}$$

for  $t \leq \tau$  and  $\mathfrak{g}_{\infty}(\iota) \geq \mu_0$ , where  $\kappa_b$  ( $\kappa_a$ ) is the smallest (largest) real part of an element in the set of eigenvalues  $\lambda_{j,\pm}$ , and C and  $\mu_0$  only depend on  $\eta_{\text{mar}}$ , the limits appearing in (11.1),  $\alpha_{\infty}$  and  $\zeta_{\infty}$ .

When estimating the last expression appearing in (12.46), it is convenient to divide the situation into two cases;  $\beta_{ts} > \kappa_a - \kappa_b$  and  $\beta_{ts} \le \kappa_a - \kappa_b$ . In the first case, we choose  $\eta_{mar}$  small enough, the bound depending only on  $\beta_{ts}$ ,  $\alpha_{\infty}$ ,  $\zeta_{\infty}$  and the limits appearing in (11.1), and then a  $\mu_0$  large enough, with the same dependence, such that  $\kappa_b - \kappa_\iota + \beta_{ts} - \eta_{mar} > 0$  for  $\iota \in \mathcal{I}^+_{B,\mu_0}$ . In the second case, we choose  $\eta_{mar} = \beta_{mar}/2$  and then  $\mu_0$  large enough, the bound depending only on  $\beta_{mar}$ ,  $\alpha_{\infty}$ ,  $\zeta_{\infty}$  and the limits appearing in (11.1). Summing up yields

$$\langle \nu(\iota) \rangle^{s} \sup_{t \in [0, T_{\rm ts}]} \left| e^{(\beta_{\rm ts} - \kappa_{\iota})t} \int_{t}^{T_{\rm ts}} e^{A(\iota)(t-\tau)} F_{\iota}(\tau) d\tau \right| \le C \langle \nu(\iota) \rangle^{s+s_{\rm ih,2}} \|\hat{f}\|_{\rm lma}$$
(12.58)

for  $\mathfrak{g}_{\infty}(\iota) \geq \mu_0$ , where

$$s_{\text{ih},2} = \max\{\beta_{\text{ts}}, \kappa_a - \kappa_b\}[(\min\{\eta_{\text{ts}}, \beta_{\text{sil}}\})^{-1} + 2\beta_{\text{sil}}^{-1}]$$

and C and  $\mu_0$  have the same dependence as in Remark 12.19.

**Combining the estimates.** Since the the absolute value of the second term on the right hand side of (12.44) (multiplied by  $\langle \nu(\iota) \rangle^s$ ) can be estimated by the left hand side of (12.58), we have all the estimates we need. To begin with, (12.44), (12.55) and (12.58) yield

$$\langle \nu(\iota) \rangle^{s} |v_{\infty,\text{mod}}| \le C \langle \nu(\iota) \rangle^{s+s_{\infty,\text{hom}}} \mathcal{E}^{1/2}(0) + C \langle \nu(\iota) \rangle^{s+s_{\infty,\text{ih}}} \|\hat{f}\|_{\text{lma}}$$
(12.59)

for  $\mathfrak{g}_{\infty}(\iota) \geq \mu_0$ , where  $s_{\infty,\text{hom}} := s_{\text{hom},1}$ ;  $s_{\infty,\text{ih}} := \max\{s_{\text{ih},1}, s_{\text{ih},2}\}$ ; and C and  $\mu_0$  have the same dependence as in Remark 12.19. Moreover, combining (12.51), (12.57) and (12.58) yields

$$\langle \nu(\iota) \rangle^{s} \left| v(t) - e^{A(\iota)t} v_{\infty, \text{mod}} - \int_{0}^{t} e^{A(\iota)(t-\tau)} F_{\iota}(\tau) d\tau \right|$$

$$\leq C e^{(\kappa_{\iota} - \beta_{\text{ts}})t} \left( \langle \nu(\iota) \rangle^{s+s_{\infty, \text{hom}}} \mathcal{E}^{1/2}(0) + \langle \nu(\iota) \rangle^{s+s_{\infty, \text{ih}}} \|\hat{f}\|_{\text{lma}} \right)$$

$$(12.60)$$

for  $0 \le t \le T_{ts}$  and  $\mathfrak{g}_{\infty}(\iota) \ge \mu_0$ , where C and  $\mu_0$  have the same dependence as in Remark 12.19. Combining (12.45) and (12.52) yields

$$\langle \nu(\iota) \rangle^{s} \left| v(t) - e^{A(\iota)t} v_{\infty, \text{mod}} - \int_{0}^{t} e^{A(\iota)(t-\tau)} F_{\iota}(\tau) d\tau \right|$$

$$\leq C e^{(\kappa_{\iota} - \beta_{\text{ts}} + \beta_{\text{mar}})t} \left( \langle \nu(\iota) \rangle^{s+s_{\infty, \text{hom}}} \mathcal{E}^{1/2}(0) + \langle \nu(\iota) \rangle^{s+s_{\infty, \text{ih}}} \|\hat{f}\|_{\text{lma}} \right)$$

$$(12.61)$$

for  $\mathfrak{g}_{\infty}(\iota) \geq \mu_0$  and  $t \geq T_{\text{ts}}$ , where C and  $\mu_0$  have the same dependence as in Remark 12.19. Combining (12.60) and (12.61), it is clear that the latter estimate holds for all  $t \geq 0$ .

**Projection onto the right subspace.** Next we would like to define a function by the condition that its Fourier coefficients are given by  $v_{\infty,\text{mod}}$  and then sum up the estimates (12.61) in order to obtain (12.33). However, we do not yet know that  $v_{\infty,\text{mod}}$  belongs to the correct space. We

therefore have to project it to  $E_{\iota,\beta_{ts}}^{\infty}$ . Let  $\Pi_{a,\iota}$  denote projection onto  $E_{\iota,\beta_{ts}}^{\infty}$  and  $\Pi_{b,\iota}$  denote projection onto  $e_{\iota,\beta_{ts}}^{\infty}$ . Then we need to estimate

$$\left| e^{A(\iota)t} \Pi_{b,\iota} \left( \int_0^{T_{\rm ts}} e^{-A(\iota)\tau} F_\iota(\tau) d\tau \right) \right|$$
  
$$\leq C e^{(\kappa_\iota - \beta_{\rm ts} + \beta_{\rm mar}/2)t} \left| \int_0^{T_{\rm ts}} e^{-A(\iota)\tau} F_\iota(\tau) d\tau \right|$$

where we have appealed to (12.43) and the constant only depends on  $\alpha_{\infty}$ ,  $\zeta_{\infty}$ , the limits appearing in (11.1) and  $\beta_{\text{mar}}$ ; note that the norm of  $\Pi_{b,\iota}$  is bounded by a constant depending only on  $\alpha_{\infty}$ ,  $\zeta_{\infty}$  and the limits appearing in (11.1). Combining this estimate with (12.58) yields

$$\langle \nu(\iota) \rangle^s \left| e^{A(\iota)t} \Pi_{b,\iota} \left( \int_0^{T_{\rm ts}} e^{-A(\iota)\tau} F_\iota(\tau) d\tau \right) \right| \le C \langle \nu(\iota) \rangle^{s+s_{\rm ih},2} e^{(\kappa_\iota - \beta_{\rm ts} + \beta_{\rm mar}/2)t} \|\hat{f}\|_{\rm lma}$$

for all  $t \ge 0$  and all s, where C has the same dependence as in Remark 12.19. Combining this estimate with (12.61), it is clear that we can replace  $v_{\infty,\text{mod}}$  with  $\prod_{a,\iota} [v_{\infty,\text{mod}}]$  on the left hand side of (12.61).

Summing up. As indicated previously, the derivation of the asymptotics proceeds in several steps. In the present proof, we derive asymptotics for the part of the solution that corresponds to  $\iota \in \mathcal{I}_{\rm B}$  such that  $\mathfrak{g}_{\infty}(\iota)$  is large. Due to the above arguments, there are  $\mu_0$  and C, with dependence as in Remark 12.19, such that if u is a solution to (1.2), then the modes of u corresponding to  $\iota \in \mathcal{I}_{{\rm B},\mu_0}^+$  satisfy (12.61) for  $t \geq 0$ , where  $v_{\infty,{\rm mod}}(\iota)$  can be replaced by  $\prod_{a,\iota} [v_{\infty,{\rm mod}}(\iota)]$ . In particular, for each such mode, we have constructed an amplitude  $\prod_{a,\iota} [v_{\infty,{\rm mod}}(\iota)]$ . Combining these amplitudes then yields a function, defined as follows. Given  $p \in \overline{M}$ , define  $U_{\infty}(p)$  by

$$U_{\infty}(p) := \sum_{\iota \in \mathcal{I}_{\mathrm{B},\mu_0}^+} \Pi_{a,\iota}[v_{\infty,\mathrm{mod}}(\iota)]\varphi_{\iota}(p),$$

where  $\varphi_{\iota}$  is given by (24.6). In order to justify that the sum on the right hand side makes sense and yields a smooth function, it is sufficient to appeal to the fact that u is smooth; the assumption that  $||f||_{ts,s} < \infty$  for all  $s \in \mathbb{R}$ ; (12.59); and the Minkowski inequality. In fact, (12.34) holds, where C and  $\mu_0$  have the same dependence as in Remark 12.19. Similarly, (12.61) implies that (12.33) holds, where C and  $\mu_0$  have the same dependence as in Remark 12.19. That the  $\iota$ 'th Fourier coefficient belongs to a space of the stated type follows from the observations made above concerning  $E_{\iota,\beta_{ts}}^{\infty}$ .

**Uniqueness.** In order to justify the uniqueness, let us assume that there are two functions  $U_{\infty,i}$ , i = 1, 2, satisfying the assumptions of the lemma. Assume, moreover, that there is a  $\iota \in \mathcal{I}_{B,\mu_0}^+$  such that the corresponding Fourier coefficients, say  $v_{\infty,i}(\iota)$ , i = 1, 2, differ. Then (12.33) implies that

$$|e^{A(\iota)t}[v_{\infty,1}(\iota) - v_{\infty,2}(\iota)]| \le C\langle t \rangle^{\delta_{\iota} - 1} e^{(\kappa_{\iota} - \beta_{\rm ts} + \beta_{\rm mar})t}$$
(12.62)

for  $t \ge 0$ , where the constant C depends on the solution and f. On the other hand, the left hand side grows faster than the right hand side due to the assumption that  $v_{\infty,i}(\iota) \in E_{\iota}$ , where  $E_{\iota}$  has properties stated in the lemma. This yields the desired contradiction and the lemma follows.  $\Box$ 

Finally, we are in a position to consider all the  $\iota \in \mathcal{I}_{B,ts}$  simultaneously. We do so by combining the results of Chapter 10 with Lemma 12.18.

**Proposition 12.22.** Assume that (1.2) is weakly transparent in the sense of Definition 12.8, weakly balanced in the sense of Definition 10.1 and weakly convergent in the sense of Definition 10.1. Assume, finally, that there is a constant  $0 < C_{\ell} \in \mathbb{R}$  and a continuous non-negative function  $\mathfrak{e} \in L^1([0,\infty))$  such that (10.10) holds for all  $0 \neq \iota \in \mathcal{I}_B$  and all  $t \geq 0$ . Fix a  $0 < \beta_{mar} \in \mathbb{R}$ which is small enough, the bound depending only on  $\alpha_{\infty}$ ,  $\zeta_{\infty}$ , the limits appearing in (11.1) and the constant  $\beta_{\text{ts}}$  introduced in Definition 12.16. Assume f to be such that  $||f||_{\text{ts},s} < \infty$  for all  $s \in \mathbb{R}$ , where  $|| \cdot ||_{\text{ts},s}$  is defined in (12.32). Then there are constants  $\sigma_{\text{hom}}, \sigma_{\text{ih}} \in [0, \infty), 0 \leq N \in \mathbb{Z}$ and  $0 < C \in \mathbb{R}$  such that the following holds. Given a solution u to (1.2), there is a unique  $U_{\infty} \in C^{\infty}(\bar{M}, \mathbb{C}^{2m})$  with the following properties

- the  $\iota$ 'th Fourier coefficient of  $U_{\infty}$  vanishes unless  $\iota \in \mathcal{I}_{B,ts}$ ,
- if  $\iota \in \mathcal{I}_{B,ts}$ , then the  $\iota$ 'th Fourier coefficient of  $U_{\infty}$  belongs to  $E_{\iota}$ , where  $E_{\iota}$  is the first generalised eigenspace in the  $(\beta_{ts} \beta_{mar}), A(\iota)$ -decomposition of  $\mathbb{C}^{2m}$ , and  $A(\iota)$  is given by (12.8),
- the estimate

$$\left\| \begin{pmatrix} (-\Delta_{\infty})^{1/2} u(\cdot,t) \\ \partial_t u(\cdot,t) \end{pmatrix} - e^{\mathcal{A}_{\rm ts}t} U_{\infty} - \int_0^t e^{\mathcal{A}_{\rm ts}(t-\tau)} \begin{pmatrix} 0 \\ f(\cdot,\tau) \end{pmatrix} d\tau \right\|_{t,s}$$

$$\leq C \langle t \rangle^N e^{-(\beta_{\rm ts} - \beta_{\rm mar})t} [\|u_t(\cdot,0)\|_{(s+\sigma_{\rm hom})} + \|u(\cdot,0)\|_{(s+\sigma_{\rm hom}+1)} + \|f\|_{\rm ts,s+\sigma_{\rm ih}}]$$

$$(12.63)$$

holds for  $t \ge 0$  and  $s \in \mathbb{R}$ , where  $\Delta_{\infty}$  is defined in (12.27),  $\mathcal{A}_{ts}$  is defined in (12.29) and  $\|\cdot\|_{t,s}$  is defined in (12.31).

Moreover,

$$\|U_{\infty}\|_{(s)} \le C \|u_t(\cdot, 0)\|_{(s+\sigma_{\text{hom}})} + C \|u(\cdot, 0)\|_{(s+\sigma_{\text{hom}}+1)} + C \|f\|_{\text{ts}, s+\sigma_{\text{ih}}}$$
(12.64)

for all  $s \in \mathbb{R}$ .

**Remark 12.23.** The constants C, N,  $\sigma_{\text{hom}}$  and  $\sigma_{\text{ih}}$  have the following dependence: C only depends on  $C_{\text{ini}}$ ,  $g^{ij}(0)$ ,  $a_r(0)$ ,  $\alpha_{\infty}$ ,  $\zeta_{\infty}$ , the limits appearing in (11.1),  $S_{\text{ts}}$ ,  $\beta_{\text{sil}}$ ,  $\beta_{\text{mar}}$ ,  $\eta_{\text{mn}}$ ,  $\eta_{\text{ts}}$ ,  $c_{\text{sil}}$ ,  $K_{\text{ts}}$ ,  $C_{\ell}$ ,  $c_{\epsilon}$ ,  $C_{\text{coeff}}$ ,  $C_{\text{mn}}$  and the spectra of the Laplace-Beltrami operators on the Riemannian manifolds  $(M_{r_j}, g_{r_j})$ ,  $j = 1, \ldots, R_{\text{ts}}$ ; N only depends on m; and  $\sigma_{\text{hom}}$  and  $\sigma_{\text{ih}}$  only depend on  $\eta_{\text{mn}}$ ,  $\eta_{\text{ts}}$ ,  $\beta_{\text{sil}}$ ,  $\beta_{\text{mar}}$ ,  $c_{\text{sil}}$ ,  $C_{\text{coeff}}$ ,  $C_{\ell}$ ,  $K_{\text{ts}}$ ,  $\alpha_{\infty}$ ,  $\zeta_{\infty}$ , the limits appearing in (11.1) and the spectra of the Laplace-Beltrami operators on the Riemannian manifolds  $(M_{r_i}, g_{r_i})$ ,  $j = 1, \ldots, R_{\text{ts}}$ .

Remark 12.24. If the conditions of Proposition 5.11 are satisfied, then the assumptions of the present proposition are satisfied. In order to justify this statement, note first that since Definitions 5.2 and 5.5 are fulfilled, the conclusions of Lemma 26.2 hold. Thus the first three limits appearing in (11.1) exist. Turning to the fourth limit, it also exists due to (5.9), Remark 26.4 and the fact that  $\mathcal{X}$  is  $C^0$ -future bounded. Moreover, it is clear that if there is an  $X^{j_k}_{\infty} \neq 0$ , then Definition 11.1 is satisfied. If, on the other hand,  $X_{\infty}^{j_k} = 0$  for all the  $j_k$ , then Definition 11.3 is satisfied; cf. the conclusions of Lemma 26.2. Thus the first condition of Definition 12.8 is satisfied. That the second and third conditions are satisfied follows from Lemma 26.2. The fourth condition follows from (5.9). Thus (1.2) is weakly transparent in the sense of Definition 12.8. Combining the fact that (4.2) holds; the fact that  $\mathcal{X}$  is  $C^0$ -future bounded; the fact that Definition 5.5 is satisfied; and Lemma 25.17, yields the conclusion that (1.2) is weakly balanced and weakly convergent; cf. Definition 10.1. Finally, there is a constant  $0 < C_{\ell} \in \mathbb{R}$  such that (10.10) (with  $\mathfrak{e} = 0$ ) holds for all  $0 \neq \iota \in \mathcal{I}_{B}$  and all  $t \geq 0$ ; this is a consequence of the fact that the second fundamental form is  $C^0$ -future bounded and Definition 5.5; cf. Lemma 25.11. In addition to the above, it is of interest to note that the constants  $\beta_{sil}$ ,  $\eta_{ts}$  and  $\eta_{mn}$  appearing in Proposition 5.11 are the same as the  $\beta_{sil}$ ,  $\eta_{\rm ts}$  and  $\eta_{\rm mn}$  appearing in the statement of Proposition 12.22.

*Proof.* Consider a solution u to (1.2). In Lemma 12.18, we derive asymptotics for the Fourier modes such that  $\mathfrak{g}_{\infty}(\iota) \geq \mu_0$  for a suitably chosen  $0 < \mu_0 \in \mathbb{R}$ . What remains is to analyse the asymptotics for Fourier modes such that  $\mathfrak{g}_{\infty}(\iota) \leq \mu_0$ . To this end, it is convenient to partially Fourier transform (1.2). In fact, in analogy with the terminology introduced in (24.6)–(24.7), we here introduce the following notation. First,

$$\bar{M}_{\rm ts} := \mathbb{T}^{d_{\rm ts}} \times \prod_{k=1}^{R_{\rm ts}} M_{r_k},\tag{12.65}$$

where  $d_{\rm ts}$ ,  $R_{\rm ts}$  are introduced in Definitions 12.1 and 12.3 (below we use additional notation introduced in these definitions without further comment). Note that the manifold  $\bar{M}_{\rm ts}$  corresponds to the transparent variables. Analogously, one can define  $\bar{M}_{\rm sil}$  corresponding to the silent variables. In analogy with (24.6), we introduce

$$\varphi_{\iota, \text{ts}}(p) := (2\pi)^{-d_{\text{ts}}/2} \prod_{k=1}^{d_{\text{ts}}} e^{in_{j_k} \cdot x^{j_k}} \cdot \prod_{k=1}^{R_{\text{ts}}} \varphi_{r_k, i_{r_k}}(p_{r_k})$$
(12.66)

for  $p \in M$  and  $\iota \in \mathcal{I}_{\mathrm{B}}$  (note, however, that the right hand side only depends on  $\iota_{\mathrm{ts}}$ ). Note that we can consider  $\varphi_{\iota,\mathrm{ts}}$  to be a function on  $\bar{M}_{\mathrm{ts}}$ . Introduce the  $(d_{\mathrm{sil}} + R_{\mathrm{sil}})$ -form  $\mu_{\mathrm{B,sil}}$  on  $\bar{M}$  by

$$\mu_{\rm B,sil} := \bigwedge_{k=1}^{d_{\rm sil}} dx^{\bar{j}_k} \wedge \bigwedge_{k=1}^{R_{\rm sil}} \mu_{g_{\bar{r}_k}}$$
(12.67)

and the  $(d_{ts} + R_{ts})$ -form  $\mu_{B,ts}$  on  $\overline{M}$  by

$$\mu_{\rm B,ts} := \pm \bigwedge_{k=1}^{d_{\rm ts}} dx^{j_k} \wedge \bigwedge_{k=1}^{R_{\rm ts}} \mu_{g_{r_k}}, \qquad (12.68)$$

where the sign is chosen so that

$$\mu_{\rm B,ts} \wedge \mu_{\rm B,sil} = \mu_{\rm B}.$$

It is convenient to introduce

$$\langle u, v \rangle_{\mathrm{B,ts}} := \int_{\bar{M}_{\mathrm{ts}}} u v^* \mu_{\mathrm{B,ts}} \tag{12.69}$$

for  $u \in C^{\infty}(\overline{M}, \mathbb{C}^k)$  and  $v \in C^{\infty}(\overline{M}, \mathbb{C})$ , where the star denotes complex conjugation. Note that  $\langle u, v \rangle_{B,ts}$  is a function on  $\overline{M}_{sil}$ . In particular, given a solution u to (1.2), we define (with slight abuse of notation)

$$\tilde{u}(\iota_{\rm ts}, p_{\rm sil}, t) := \langle u(\cdot, t), \varphi_{\iota_{\rm ts}, \rm ts} \rangle_{\rm B, ts}$$

This function satisfies the equation

$$\tilde{u}_{tt} - \sum_{k,l=1}^{d_{\rm sil}} g^{\bar{j}_k \bar{j}_l}(t) \partial_{\bar{j}_k} \partial_{\bar{j}_l} \tilde{u} - 2 \sum_{l=1}^{d_{\rm sil}} g^{0\bar{j}_l}(t) \partial_{\bar{j}_l} \partial_t \tilde{u} - \sum_{k=1}^{R_{\rm sil}} a_{\bar{r}_k}^{-2}(t) \Delta_{g_{\bar{r}_k}} \tilde{u} + \alpha_{\rm mod}(t) \tilde{u}_t + \sum_{k=1}^{d_{\rm sil}} X_{\rm mod}^{\bar{j}_k}(t) \partial_{\bar{j}_k} \tilde{u} + \zeta_{\rm mod}(t) \tilde{u} = \tilde{f},$$
(12.70)

where  $\tilde{f}$  is defined in the same way as  $\tilde{u}$  and

$$\begin{aligned} \alpha_{\rm mod}(t) &:= \alpha(t) - 2 \sum_{k=1}^{d_{\rm ts}} i n_{j_k} g^{0j_k}(t) \mathrm{Id}_m, \\ \zeta_{\rm mod}(t) &:= \zeta(t) + \mathfrak{g}^2(\iota_{\rm ts}, t) \mathrm{Id}_m + i \sum_{k=1}^{d_{\rm ts}} n_{j_k} X^{j_k}(t), \\ X_{\rm mod}^{\bar{j}_k}(t) &:= X^{\bar{j}_k}(t) - 2 \sum_{l=1}^{d_{\rm ts}} i n_{j_l} g^{j_l \bar{j}_k}(t) \mathrm{Id}_m. \end{aligned}$$

Next, we wish to verify that (12.70) is an equation such that the results of Chapter 10 apply. We do so in several steps.

**Properties of the metric.** As a first step, we verify that the metric on  $M_{\rm sil} := M_{\rm sil} \times I$ , say  $g_{\rm sil}$ , associated with the equation (12.70) is such that  $(M_{\rm sil}, g_{\rm sil})$  is a canonical separable cosmological model manifold. Note that  $g_{\rm sil}^{00}(t) = -1$  and that  $g_{\rm sil}^{kl}(t) = g^{j_k \bar{j}_l}(t)$  are the components of a positive definite matrix for all  $t \in I$ . Finally,  $g_{\rm sil}^{0l}(t) = g^{0\bar{j}_l}(t)$ . As a consequence, there are smooth functions  $g_{\rm sil,\alpha\beta}$ ,  $\alpha, \beta \in \{0, \ldots, d_{\rm sil}\}$ , from I to  $\mathbb{R}$  such that  $g_{\rm sil,\alpha\beta}(t)$  are the components of the inverse of the matrix with components  $g_{\rm sil}^{\alpha\beta}(t)$ . Moreover,  $g_{\rm sil,00}(t) < 0$  and  $g_{\rm sil,ij}(t)$  are the components of a positive definite matrix for all t. These statements are a consequence of [34, Lemma 8.5, p. 72]. To conclude,  $g_{\rm sil}$  can be written

$$g_{\rm sil} = g_{\rm sil,00} dt \otimes dt + g_{\rm sil,0i} dt \otimes dx^i + g_{\rm sil,i0} dx^i \otimes dt + g_{\rm sil,ij} dx^i \otimes dx^j + \sum_{i=1}^{R_{\rm sil}} a_{\bar{r}_i}^2 g_{\bar{r}_i}$$

Moreover, it is clear that  $(M_{\rm sil}, g_{\rm sil})$  is a separable cosmological model manifold; that  $g_{\rm sil}$  is a Lorentz metric follows from [34, Lemma 8.5, p. 72]. Due to Remark 25.3, it is clear that the lapse function equals one. Finally, I contains  $[0, \infty)$ , so that  $(M_{\rm sil}, g_{\rm sil})$  is a canonical separable cosmological model manifold.

Estimating the coefficients of the equation. Let us start by considering  $\alpha_{\text{mod}}$  and  $\zeta_{\text{mod}}$ . To begin with,

$$\|\alpha_{\rm mod}(t)\| \le \|\alpha(t)\| + 2|\sigma(\iota_{\rm ts}, t)| \cdot \mathfrak{g}(\iota_{\rm ts}, t) \le C(1+\mu_0)$$
(12.71)

for all  $t \ge 0$  and all  $\iota \in \mathcal{I}_{\mathrm{B},\mu_0}^-$ , where the notation  $\mathcal{I}_{\mathrm{B},\mu_0}^-$  is introduced in connection with (12.30). Here *C* only depends on  $C_{\mathrm{coeff}}$  appearing in (10.2) and  $K_{\mathrm{ts}}$  and  $\eta_{\mathrm{ts}}$  appearing in (12.4). Moreover, we have appealed to the estimates (12.10), (12.11) and the bound  $\mathfrak{g}_{\infty}(\iota) \le \mu_0$ . Similarly,

$$\|\zeta_{\text{mod}}(t)\| \le \|\zeta(t)\| + \mathfrak{g}^2(\iota_{\text{ts}}, t) + \|X(\iota_{\text{ts}}, t)\| \cdot \mathfrak{g}(\iota_{\text{ts}}, t) \le C(1 + \mu_0^2)$$
(12.72)

for all  $t \ge 0$  and all  $\iota \in \mathcal{I}_{B,\mu_0}^-$ , where *C* only depends on  $C_{\text{coeff}}$ ,  $K_{\text{ts}}$  and  $\eta_{\text{ts}}$ , and we have appealed to the estimates (12.10), (12.11) and the bound  $\mathfrak{g}_{\infty}(\iota) \le \mu_0$ . Finally, for  $\iota \in \mathcal{I}_{B,\mu_0}^-$  such that  $\iota_{\text{sil}} \ne 0$ ,

$$\|X_{\text{mod}}(\iota_{\text{sil}}, t)\| = \frac{\|n_{\bar{j}_k} X_{\text{mod}}^{j_k}(t)\|}{\mathfrak{g}(\iota_{\text{sil}}, t)} \le \frac{\|n_{\bar{j}_k} X^{\bar{j}_k}(t)\|}{\mathfrak{g}(\iota_{\text{sil}}, t)} + 2\frac{|n_{\bar{j}_k} n_{j_l} g^{j_l \bar{j}_k}(t)|}{\mathfrak{g}(\iota_{\text{sil}}, t)}$$

The first term on the far right hand side is bounded by  $C_{\text{coeff}}$  for  $t \ge 0$ . Due to (12.17), the second term on the far right hand side is bounded by  $2\mathfrak{g}(\iota_{\text{ts}}, t)$ . Thus

$$\|X_{\text{mod}}(\iota_{\text{sil}}, t)\| \le C_{\text{coeff}} + 2\mathfrak{g}(\iota_{\text{ts}}, t) \le C(1+\mu_0)$$
(12.73)

for all  $t \geq 0$  and all  $\iota \in \mathcal{I}_{B,\mu_0}^-$  such that  $\iota_{sil} \neq 0$ . Here C only depends on  $C_{coeff}$ ,  $K_{ts}$  and  $\eta_{ts}$ . Introducing

$$\begin{aligned} \alpha_{\mathrm{mod},\infty} &:= \alpha_{\infty}, \\ \zeta_{\mathrm{mod},\infty} &:= \zeta_{\infty} + \mathfrak{g}_{\infty}^{2}(\iota_{\mathrm{ts}}) \mathrm{Id}_{m} + i \sum_{k=1}^{d_{\mathrm{ts}}} n_{j_{k}} X_{\infty}^{j_{k}}, \end{aligned}$$

it can similarly be verified that

$$\|\alpha_{\rm mod}(t) - \alpha_{\rm mod,\infty}\| \le C(1+\mu_0)e^{-\eta_{\rm mod,mn}t},$$
(12.74)

$$\|\zeta_{\text{mod}}(t) - \zeta_{\text{mod},\infty}\| \le C(1 + \mu_0^2) e^{-\eta_{\text{mod},\text{mn}}t}$$
(12.75)

for all  $t \ge 0$  and all  $\iota \in \mathcal{I}_{B,\mu_0}^-$ , where *C* only depends on  $C_{mn}$ ,  $C_{ini}$ ,  $K_{ts}$  and  $\eta_{ts}$ ;  $\eta_{mod,mn} := \min\{\eta_{mn}, \eta_{ts}\}$ ; and we have appealed to (10.4), (12.4), (12.5), (12.6), (12.10), (12.11) and (12.20). It is thus clear that (12.70) is weakly convergent in the sense of Definition 10.1; in (10.4) we simply have to replace  $\eta_{mn}$  by  $\eta_{mod,mn}$  and  $C_{mn}$  by  $C_{mod,mn} := C(2 + \mu_0 + \mu_0^2)$ , where *C* is the constant appearing in (12.74) and (12.75).

Weak balance; weak silence; and generalised eigenspaces. Before turning to the topic of weak balance, it is important to note that there is one  $\mathfrak{g}, \sigma$  and X associated with the equation (1.2) and one  $\mathfrak{g}$ ,  $\sigma$  and X associated with the equation (12.70) (say  $\mathfrak{g}_{mod}$ ,  $\sigma_{mod}$  and  $X_{mod}$ ), and it is important not to confuse them with one another. Moreover, the set  $\mathcal{I}_{\mathrm{B}}$  is different for the two equations. However, we can think of  $\mathcal{I}_{B,sil}$  as the set  $\mathcal{I}_B$  corresponding to the equation (12.70). Moreover, in practice,  $\mathfrak{g}_{mod} = \mathfrak{g}$  and  $\sigma_{mod} = \sigma$ , assuming that we only evaluate  $\mathfrak{g}$  and  $\sigma$  at  $\iota \in \mathcal{I}_{B,sil}$ . Finally, we should consider  $\iota_{ts} \neq 0$  to be fixed once and for all. Combining (10.2) (which holds for (1.2) by assumption) with (12.71), (12.72) and (12.73), it follows that (12.70)is weakly balanced (and the constant  $C_{\text{coeff}}$  is replaced by a constant depending only on  $C_{\text{coeff}}$ ,  $K_{\rm ts}$ ,  $\eta_{\rm ts}$  and  $\mu_0$ ). Finally, that (12.70) is weakly silent is an immediate consequence of (12.3). When appealing to Lemma 10.17, it is important to keep track of what  $\kappa_1$ ,  $\beta_{\rm rem}$  and  $E_a$  are. In the present context,  $\kappa_1 = \kappa_i$ , where the notation  $\kappa_i$  is introduced in connection with (12.30); this is justified by (12.76)–(12.78) below. Moreover,  $\beta_{\text{rem}} = \min\{\eta_{\text{mod},\text{mn}}, \beta_{\text{sil}}\} = \min\{\eta_{\text{mn}}, \eta_{\text{ts}}, \beta_{\text{sil}}\},$ so that  $\beta_{\rm rem} = \beta_{\rm ts}$ ; cf. (12.23). Comparing the statement of the present proposition with the statement of Lemma 10.17, it is clear that we would like to appeal to Lemma 10.17 with  $\beta_{\rm rem}$ replaced by  $\beta_{\rm ts} - \beta_{\rm mar}$ . Formally, this can be achieved by, in the application of Lemma 10.17, replacing  $\eta_{\text{mod},\text{mn}}$  by  $\beta_{\text{ts}} - \beta_{\text{mar}}$ . Appealing to Lemma 10.17 with this modification,  $E_a$  is the

direct sum of the generalised eigenspaces of  $A_{\text{mod}}$  (defined below) corresponding to eigenvalues with real part >  $\kappa_{\iota} - (\beta_{\text{ts}} - \beta_{\text{mar}})$ .

Applying the results of Chapter 10. When comparing with the results of Chapter 10, the matrix  $A_{\infty}$  is replaced by

$$A_{\text{mod}} := \begin{pmatrix} 0 & \text{Id}_m \\ -\zeta_\infty - \mathfrak{g}_\infty^2 \text{Id}_m - i \sum_{k=1}^{d_{\text{ts}}} n_{j_k} X_\infty^{j_k} & -\alpha_\infty \end{pmatrix}.$$
 (12.76)

Introducing the matrix

$$D_{\infty} := \operatorname{diag}(\mathfrak{g}_{\infty}^{-1} \operatorname{Id}_{m}, \operatorname{Id}_{m}), \qquad (12.77)$$

it can be verified that

$$D_{\infty}^{-1}A_{\mathrm{mod}}D_{\infty} = A(\iota), \qquad (12.78)$$

where  $A(\iota)$  is given by (12.8). Due to this observation, it is clear that the requirement (10.25) can be reformulated to

$$\int_0^\infty e^{-\kappa_\iota \tau} \|\tilde{f}(\iota_{\rm ts},\cdot,\tau)\|_{(s)} d\tau < \infty.$$
(12.79)

Moreover, due to the requirement  $||f||_{ts,s} < \infty$  for all  $s \in \mathbb{R}$  and the definition (12.32), it is clear that (12.79) holds. Let  $\beta_{\text{rem}} = \beta_{\text{ts}} - \beta_{\text{mar}}$  be defined as above and  $E_a$  be the first generalised eigenspace in the  $\beta_{\text{rem}}, A_{\text{mod}}$ -decomposition of  $\mathbb{C}^{2m}$  (note that  $D_{\infty}^{-1}E_a$  is the first generalised eigenspace in the  $(\beta_{ts} - \beta_{mar}), A(\iota)$ -decomposition of  $\mathbb{C}^{2m}$ ). Then Lemma 10.17 is applicable, and there are constants C, N,  $s_{\text{hom}}$  and  $s_{\text{ih}}$  such that, given a solution  $\tilde{u}$  to (12.70), there is a  $V_{\infty} \in C^{\infty}(\bar{M}_{\rm sil}, E_a)$  such that (10.26) and (10.27) hold with u, f and  $A_{\infty}$  replaced by  $\tilde{u}, \tilde{f}$  and  $A_{\rm mod}$  respectively. Moreover, the constants have the following dependence: the constant C only depends on  $c_{\rm sil}$ ,  $C_{\rm coeff}$ ,  $K_{\rm ts}$ ,  $\eta_{\rm ts}$ ,  $\mu_0$ ,  $\beta_{\rm mar}$ ,  $C_{\rm mn}$ ,  $C_{\rm ini}$ ,  $\beta_{\rm sil}$ ,  $\eta_{\rm mn}$ ,  $A_{\rm mod}$ ,  $g^{ij}(0)$  and  $a_r(0)$ . Since there are only finitely many  $\iota_{\rm ts}$  satisfying  $\mathfrak{g}_{\infty}(\iota_{\rm ts}) \leq \mu_0$ , we only need to consider a finite number of matrices of the form (12.76). The dependence on  $A_{\rm mod}$  can thus be replaced by a dependence on  $\alpha_{\infty}, \zeta_{\infty}$ , the limits appearing in (11.1),  $\mu_0$  and the spectra of the Laplace-Beltrami operators on the Riemannian manifolds  $(M_{r_j}, g_{r_j}), j = 1, \ldots, R_{ts}$ . This yields a constant for which Lemma 10.17 is applicable irrespective of  $\iota \in \mathcal{I}_{\mathrm{B},\mu_0}^-$ . The constant N only depends on m. Finally, by a similar argument,  $s_{\text{hom}} \geq 0$  and  $s_{\text{ih}} \geq 0$  can be chosen to depend only  $\eta_{\text{mn}}$ ,  $\eta_{\text{ts}}$ ,  $\beta_{\text{sil}}$ ,  $\beta_{\text{mar}}$ ,  $C_{\text{coeff}}$ ,  $K_{\text{ts}}$ ,  $c_{\rm sil}, \alpha_{\infty}, \zeta_{\infty}$ , the limits appearing in (11.1), the spectra of the Laplace-Beltrami operators on the Riemannian manifolds  $(M_{r_i}, g_{r_i}), j = 1, \ldots, R_{ts}$ , and  $\mu_0$  (in particular, they are thus independent of  $\iota \in \mathcal{I}_{B,\mu_0}^-$ ). Thus, given a solution  $\tilde{u}$  to (12.70), there is a  $V_{\infty}(\iota_{ts}, \cdot) \in C^{\infty}(\bar{M}_{sil}, E_a)$  such that

$$\left\| \begin{pmatrix} \tilde{u}(\iota_{\rm ts},\cdot,t) \\ \tilde{u}_{t}(\iota_{\rm ts},\cdot,t) \end{pmatrix} - e^{A_{\rm mod}t}V_{\infty}(\iota_{\rm ts},\cdot) - \int_{0}^{t} e^{A_{\rm mod}(t-\tau)} \begin{pmatrix} 0 \\ \tilde{f}(\iota_{\rm ts},\cdot,\tau) \end{pmatrix} d\tau \right\|_{(s)} \\
\leq C\langle t \rangle^{N} e^{(\kappa_{\iota}-\beta_{\rm ts}+\beta_{\rm mar})t} \left( \|\tilde{u}_{t}(\iota_{\rm ts},\cdot,0)\|_{(s+s_{\rm hom})} + \|\tilde{u}(\iota_{\rm ts},\cdot,0)\|_{(s+s_{\rm hom}+1)} \\
+ \|\tilde{f}(\iota_{\rm ts},\cdot)\|_{A,s+s_{\rm ih}} \right)$$
(12.80)

holds for  $t \ge 0$ ; recall that  $\overline{M}_{sil}$  is defined in connection with (12.65). Moreover,

$$V_{\infty}(\iota_{\rm ts}, \cdot)\|_{(s)} \leq C \left( \|\tilde{u}_t(\iota_{\rm ts}, \cdot, 0)\|_{(s+s_{\rm hom})} + \|\tilde{u}(\iota_{\rm ts}, \cdot, 0)\|_{(s+s_{\rm hom}+1)} + \|\tilde{f}(\iota_{\rm ts}, \cdot)\|_{A, s+s_{\rm ih}} \right).$$
(12.81)

Appealing to (12.77) and (12.78), the estimate (12.80) can be reformulated to

$$\left\| \begin{pmatrix} \mathfrak{g}_{\infty}(\iota)\tilde{u}(\iota_{\mathrm{ts}},\cdot,t) \\ \tilde{u}_{t}(\iota_{\mathrm{ts}},\cdot,t) \end{pmatrix} - e^{A(\iota)t}W_{\infty}(\iota_{\mathrm{ts}},\cdot) - \int_{0}^{t} e^{A(\iota)(t-\tau)} \begin{pmatrix} 0 \\ \tilde{f}(\iota_{\mathrm{ts}},\cdot,\tau) \end{pmatrix} d\tau \right\|_{(s)} \\ \leq C\langle t \rangle^{N} e^{(\kappa_{\iota}-\beta_{\mathrm{ts}}+\beta_{\mathrm{mar}})t} \left( \|\tilde{u}_{t}(\iota_{\mathrm{ts}},\cdot,0)\|_{(s+s_{\mathrm{hom}})} + \|\tilde{u}(\iota_{\mathrm{ts}},\cdot,0)\|_{(s+s_{\mathrm{hom}}+1)} \\ + \|\tilde{f}(\iota_{\mathrm{ts}},\cdot)\|_{A,s+s_{\mathrm{ih}}} \right),$$
(12.82)

168

#### 12.4. SPECIFYING THE ASYMPTOTICS

where C, N,  $s_{\text{hom}}$  and  $s_{\text{ih}}$  have the same dependence as in the case of (12.80). Moreover,  $W_{\infty}$ satisfies an estimate of the form (12.81) and takes its values in the first generalised eigenspace of the  $(\beta_{\text{ts}} - \beta_{\text{mar}}), A(\iota)$ -decomposition of  $\mathbb{C}^{2m}$ , a space having the properties required of  $E_{\iota}$  in the statement of the proposition. Note that (12.81) and (12.82) hold for the individual modes. One way to see this is to replace f on the right hand side of (1.2) by a function, say  $f_1$ , whose  $\iota$ 'th mode is that of f and all of whose remaining modes vanish. Replace u by the solution, say  $u_1$ , to (1.2) with f replaced by  $f_1$  and initial data given by the functions whose  $\iota$ 'th modes are those of  $u(\cdot, 0)$  and  $u_t(\cdot, 0)$  respectively (and all of whose remaining modes vanish). Then it is clear that (12.81) and (12.82) hold for the  $\iota$ 'th mode. In particular, (12.82) implies that

$$\begin{vmatrix} \left( \mathfrak{g}_{\infty}(\iota)\hat{u}(\iota,t) \\ \hat{u}_{t}(\iota,t) \right) - e^{A(\iota)t}\hat{U}_{\infty}(\iota) - \int_{0}^{t} e^{A(\iota)(t-\tau)} \left( \begin{array}{c} 0 \\ \hat{f}(\iota,\tau) \end{array} \right) d\tau \\ \leq C\langle t \rangle^{N} e^{(\kappa_{\iota} - \beta_{\mathrm{ts}} + \beta_{\mathrm{mar}})t} \left( \langle \nu(\iota_{\mathrm{sil}}) \rangle^{s_{\mathrm{hom}}} |\hat{u}_{t}(\iota,0)| + \langle \nu(\iota_{\mathrm{sil}}) \rangle^{s_{\mathrm{hom}}+1} |\hat{u}(\iota,0)| \\ + \langle \nu(\iota_{\mathrm{sil}}) \rangle^{s_{\mathrm{ih}}} \| \hat{f}(\iota,\cdot) \|_{A} \end{matrix} \right) \tag{12.83}$$

for  $\iota \in \mathcal{I}_{\mathrm{B}, \iota_0}^-$ , where  $\hat{U}_{\infty}(\iota)$  is the  $\iota_{\mathrm{sil}}$ 'th mode of  $W_{\infty}(\iota_{\mathrm{ts}}, \cdot)$ . Thus

$$\left\| \left( \begin{array}{c} (-\Delta_{\infty})^{1/2} u(\cdot,t) \\ \partial_{t} u(\cdot,t) \end{array} \right) - e^{\mathcal{A}_{ts}t} U_{\infty} - \int_{0}^{t} e^{\mathcal{A}_{ts}(t-\tau)} \left( \begin{array}{c} 0 \\ f(\cdot,\tau) \end{array} \right) d\tau \right\|_{t,s,\mu_{0},-}$$

$$\leq C \langle t \rangle^{N} e^{-(\beta_{ts}-\beta_{mar})t} [\|u_{t}(\cdot,0)\|_{(s+s_{hom},\mu_{0},-)} + \|u(\cdot,0)\|_{(s+s_{hom}+1,\mu_{0},-)} + \|f\|_{ts,s+s_{h},\mu_{0},-}],$$

$$(12.84)$$

where C, N,  $s_{\text{hom}}$  and  $s_{\text{ih}}$  have the same dependence as in the case of (12.80). Moreover,

$$\|U_{\infty}\|_{(s,\mu_{0},-)} \leq C \left( \|u_{t}(\cdot,0)\|_{(s+s_{\text{hom}},\mu_{0},-)} + \|u(\cdot,0)\|_{(s+s_{\text{hom}}+1,\mu_{0},-)} + \|f\|_{\text{ts},s+s_{\text{ih}},\mu_{0},-} \right),$$
(12.85)

and

$$\begin{split} \|\psi\|_{(s,\mu_{0},-)} &:= \left(\sum_{\iota \in \mathcal{I}_{\mathrm{B},\mu_{0}}^{-}} \langle \nu(\iota) \rangle^{2s} |\hat{\psi}(\iota)|^{2}\right)^{1/2}, \\ \|f\|_{\mathrm{ts},s,\mu_{0},-} &:= \int_{0}^{\infty} \left(\sum_{\iota \in \mathcal{I}_{\mathrm{B},\mu_{0}}^{-}} \langle \nu(\iota) \rangle^{2s} \langle t \rangle^{2\delta_{\iota}-2} e^{-2\kappa_{\iota} t} |\hat{f}(\iota,t)|^{2}\right)^{1/2} dt. \end{split}$$

Summarising. To summarise, the conclusions of the proposition are a consequence of the following argument. First, we fix a  $\beta_{\text{mar}}$  as in the statement of the proposition. In particular,  $\beta_{\text{mar}} > 0$ should be small enough that Lemma 12.18 is applicable. Second, we choose a  $\mu_0$  as in the statement of Lemma 12.18. Assume, moreover,  $\mu_0$  to have the dependence stated in Remark 12.19. Third, for the given  $\beta_{\text{mar}}$  and  $\mu_0$ , we appeal to Lemma 12.18 and the above arguments of the present proof. Fourth, we define  $U_{\infty}$  by combining the function constructed in Lemma 12.18 with the function constructed above. As a consequence of the construction, the  $\iota$ 'th Fourier coefficient of  $U_{\infty}$  vanishes if  $\iota \notin \mathcal{I}_{\text{B,ts}}$  and takes its values in  $E_{\iota}$  for  $\iota \in \mathcal{I}_{\text{B,ts}}$ . Moreover,  $E_{\iota}$  is the first generalised eigenspace in the ( $\beta_{\text{ts}} - \beta_{\text{mar}}$ ),  $A(\iota)$ -decomposition of  $\mathbb{C}^{2m}$ . Combining (12.33) and (12.84) yields the conclusion that (12.63) holds. Combining (12.34) and (12.85) yields the conclusion that (12.64) holds. Moreover, the constants have the dependence stated in Remark 12.23. Finally, the proof of uniqueness is in practice identical to the proof of uniqueness in Lemma 12.18. The proposition follows.

### **12.4** Specifying the asymptotics

In the present section, we derive results similar to those of Section 10.6, but for weakly transparent equations. Considering the statement of Proposition 12.22, it is clear that there are some differences between the present setting and that considered in Section 10.6. In particular, we cannot

expect the limiting function  $U_{\infty}$  appearing in (12.63) to take its values in one specific subspace of  $\mathbb{C}^{2m}$ . In fact, each mode of  $U_{\infty}$  can be expected to belong to a different subspace. For that reason, it is convenient to introduce the following terminology. Let  $E_{\rm ts}$  be a map from  $\mathcal{I}_{\rm B,ts}$  to the set of vector subspaces of  $\mathbb{C}^{2m}$ . Then we say that  $\chi \in C^{\infty}(\bar{M}, \mathbb{C}^{2m})$  is  $E_{\rm ts}$ -adapted if the *i*'th Fourier coefficient of  $\chi$  vanishes for  $\iota \in \mathcal{I}_{\rm B,sil}$  and belongs to  $E_{\rm ts}(\iota)$  for  $\iota \in \mathcal{I}_{\rm B,ts}$ . We denote the set of  $E_{\rm ts}$ -adapted elements of  $C^{\infty}(\bar{M}, \mathbb{C}^{2m})$  by  $C^{\infty}(\bar{M}, \mathbb{C}^{2m}; E_{\rm ts})$ . It is also convenient to introduce the notation  $C_{\rm ts}^{\infty}(\bar{M}, \mathbb{C}^{2m})$  to denote the set of elements of  $C^{\infty}(\bar{M}, \mathbb{C}^{2m})$  whose *i*'th Fourier coefficients vanish for  $\iota \in \mathcal{I}_{\rm B,sil}$ .

**Proposition 12.25.** Assume that (1.2) is weakly transparent in the sense of Definition 12.8, weakly balanced in the sense of Definition 10.1 and weakly convergent in the sense of Definition 10.1. Assume, finally, that there is a constant  $0 < C_{\ell} \in \mathbb{R}$  and a continuous non-negative function  $\mathfrak{e} \in L^1([0,\infty))$  such that (10.10) holds for all  $0 \neq \iota \in \mathcal{I}_B$  and all  $t \geq 0$ . Fix a  $0 < \beta_{\max} \in \mathbb{R}$ which is small enough, the bound depending only on  $\alpha_{\infty}, \zeta_{\infty}$ , the limits appearing in (11.1) and the constant  $\beta_{ts}$  introduced in Definition 12.16. Assume, moreover, that f = 0. Then there is a function  $E_{ts}$  from  $\mathcal{I}_{B,ts}$  to the set of subspaces of  $\mathbb{C}^{2m}$  such that if  $E_{\iota} := E_{ts}(\iota)$ , then the spaces  $E_{\iota}$  have the properties stated in Proposition 12.22. Moreover, there are constants  $C_{\Phi} > 0$  and  $s_{\infty} \geq 0$ , and an injective linear map  $\Phi_{\infty}$  from  $C^{\infty}(\bar{M}, \mathbb{C}^{2m}; E_{ts})$  to  $C_{ts}^{\infty}(\bar{M}, \mathbb{C}^{2m})$  with the following properties. First,

$$\|\Phi_{\infty}(\chi)\|_{(s)} \le C_{\Phi} \|\chi\|_{(s+s_{\infty})} \tag{12.86}$$

for all  $s \in \mathbb{R}$  and all  $\chi \in C^{\infty}(\overline{M}, \mathbb{C}^{2m}; E_{ts})$ . Second, if  $\chi \in C^{\infty}(\overline{M}, \mathbb{C}^{2m}; E_{ts})$  and u is the solution to (1.2) (with f = 0) such that

$$\begin{pmatrix} u(\cdot,0)\\ u_t(\cdot,0) \end{pmatrix} = \Phi_{\infty}(\chi), \qquad (12.87)$$

then

$$\left\| \begin{pmatrix} (-\Delta_{\infty})^{1/2} u(\cdot,t) \\ u_t(\cdot,t) \end{pmatrix} - e^{\mathcal{A}_{ts}t} \chi \right\|_{t,s}$$

$$\leq C \langle t \rangle^N e^{-(\beta_{ts} - \beta_{mar})t} \left( \| u_t(\cdot,0) \|_{(s+\sigma_{hom})} + \| u(\cdot,0) \|_{(s+\sigma_{hom}+1)} \right)$$

$$(12.88)$$

for all  $t \geq 0$  and all  $s \in \mathbb{R}$ , where the constants C, N and  $\sigma_{\text{hom}}$  have the same dependence as in the case of Proposition 12.22; cf. Remark 12.23. Finally, if there is an  $\epsilon > 0$  such that  $\text{Rsp}[A(\iota)] < \beta_{\text{ts}} - \epsilon$  for all  $\iota \in \mathcal{I}_{\text{B,ts}}$ , then  $E_{\text{ts}}$  can be chosen to be such that  $E_{\text{ts}}(\iota) = \mathbb{C}^{2m}$  for all  $\iota \in \mathcal{I}_{\text{B,ts}}$ . Then  $\Phi_{\infty}$  is surjective.

**Remark 12.26.** The constant  $C_{\Phi}$  only depends on  $\beta_{\text{mar}}$ ,  $\beta_{\text{sil}}$ ,  $\alpha_{\infty}$ ,  $\zeta_{\infty}$ , the limits appearing in (11.1),  $C_{\text{ini}}$ ,  $C_{\ell}$ ,  $C_{\text{mn}}$ ,  $C_{\text{coeff}}$ ,  $c_{\text{sil}} := \|\mathbf{e}_{\text{sil}}\|_1$ ,  $c_{\mathbf{e}} := \|\mathbf{e}\|_1$ ,  $S_{\text{ts}}$ ,  $\eta_{\text{ts}}$ ,  $\eta_{\text{mn}}$ ,  $K_{\text{ts}}$ , the supremum of  $\kappa_{\iota}$  for  $\iota \in \mathcal{I}_{\text{B,ts}}$  and the spectra of the Laplace-Beltrami operators on the Riemannian manifolds  $(M_{r_j}, g_{r_j}), j = 1, \ldots, R_{\text{ts}}$ ; and  $s_{\infty} \geq 0$  only depends on  $C_{\text{coeff}}$ ,  $C_{\ell}$ ,  $\beta_{\text{sil}}$ ,  $\eta_{\text{ts}}$  and the supremum of  $\kappa_{\iota}$  for  $\iota \in \mathcal{I}_{\text{B,ts}}$ .

**Remark 12.27.** By combining (12.86), (12.87) and (12.88), the norms of  $u(\cdot, 0)$  and  $u_t(\cdot, 0)$  appearing on the right hand side of (12.88) can be replaced by a suitable Sobolev norm of  $\chi$ .

**Remark 12.28.** In order to obtain a similar result in the case of inhomogeneous equations, it is sufficient to combine Propositions 12.22 and 12.25; cf. Remark 9.23.

Proof. Let  $\chi \in C^{\infty}(\overline{M}, \mathbb{C}^{2m})$ ,  $\iota \in \mathcal{I}_{B,ts}$  and  $\hat{\chi}(\iota) := \langle \chi, \varphi_{\iota} \rangle_{B}$ ; cf. the notation introduced in Subsection 24.1.2. Let u be a solution to (1.2) (with f = 0) and let  $z(\iota, t)$  be given by (24.14). Note that the  $\iota$ 'th Fourier coefficient of the expression inside the norm on the left hand side of (12.88) can be written

$$v(\iota, t) - e^{A(\iota)t}\hat{\chi}(\iota),$$
 (12.89)

where we use the notation (11.5) and (12.8). Clearly, we want to construct a v such that this object, after multiplication with  $e^{-\kappa_{\iota}t}$ , decays exponentially. On the other hand, v satisfies (11.3)

with  $F_{\iota} = 0$ , where  $A_{\iota,\text{rem}}$  satisfies (12.25) for  $t \ge T_{\text{ts}}$ ; cf. Definition 12.16 for an explanation of the terminology used. In order to construct v, it thus seems natural to appeal to Lemmas 9.21 and 11.19.

Due to the above observations, the strategy of the proof is the following. In order to construct the initial data corresponding to  $\chi$ , we consider the different Fourier modes, as above, and appeal to Lemmas 9.21 and 11.19. There are three things to keep in mind when appealing to these lemmas. First, Lemma 11.19 only applies for  $\mathfrak{g}_{\infty}(\iota)$  sufficiently large; second, when appealing to Lemma 9.21, the constants appearing depend on  $A(\iota)$ ; and, third, in order to be allowed to appeal to Lemmas 9.21 and 11.19,  $\hat{\chi}(\iota)$  has to belong to an appropriate vector subspace of  $\mathbb{C}^{2m}$ , say  $E_{ts}(\iota)$ (what space this is depends on the lemma to which we appeal). It is therefore natural to begin by appealing to Lemma 11.19, and, in the process, to fix a choice of  $\mu_0 > 0$  and to make a choice of  $E_{\rm ts}(\iota)$  for  $\mathfrak{g}_{\infty}(\iota) \geq \mu_0$ . For the  $\iota \in \mathcal{I}_{\rm B,ts}$  to which Lemma 11.19 does not apply, we then appeal to Lemma 9.21 and make a corresponding choice of  $E_{ts}(\iota)$ . Since the number of matrices  $A(\iota)$  for  $0 < \mathfrak{g}_{\infty}(\iota) < \mu_0$  is finite, the dependence of the constants on  $A(\iota)$  is manageable. The outcome of the analysis is a construction of a map  $E_{\rm ts}$  from  $\mathcal{I}_{\rm B,ts}$  to the set of subspaces of  $\mathbb{C}^{2m}$ ; given  $\iota \in \mathcal{I}_{\rm B,ts}$ and  $\hat{\chi}(\iota) \in E_{\rm ts}(\iota)$ , a construction of  $v(\iota, T_{\rm ts})$ ; estimates of  $|v(\iota, T_{\rm ts})|$  in terms of the asymptotic data  $\hat{\chi}(\iota)$ ; and an asymptotic estimate of (12.89). Once  $v(\iota, T_{\rm ts})$  has been constructed, the next step is to estimate  $v(\iota, 0)$  in terms of  $\hat{\chi}(\iota)$ . We do so by appealing to Lemma 10.7. Finally, some additional arguments are required to verify that all the conclusions of the proposition hold.

From asymptotic data to  $T_{\rm ts}$ . Note, to begin with, that the terminology of the present setting is related to that of Lemma 11.19 as described at the beginning of the proof of Lemma 12.18. Fixing  $\beta_{\rm mar} > 0$  small enough, the bound depending only on  $\beta_{\rm ts}$ ,  $\alpha_{\infty}$ ,  $\zeta_{\infty}$  and the limits appearing in (11.1), there are then constants  $\mu_0 > 0$  and C > 0 such that the conclusions of Lemma 11.19 hold. Moreover,  $\mu_0$  only depends on  $\beta_{\rm mar}$ ,  $\alpha_{\infty}$ ,  $\zeta_{\infty}$  and the limits appearing in (11.1), and Conly depends on  $C_{\rm ts}$ ,  $\beta_{\rm ts}$ ,  $\beta_{\rm mar}$ ,  $\alpha_{\infty}$ ,  $\zeta_{\infty}$  and the limits appearing in (11.1). Let  $\Psi_{\iota,\infty}$  be the map constructed in Lemma 11.19. Note that this map depends on  $\iota$ , but that the constants appearing in the estimates involving  $\Psi_{\iota,\infty}$  do not. Note, moreover, that  $\Psi_{\iota,\infty}$  maps  $E_{\iota,\beta_{\rm ts}}^{\infty}$  to  $\mathbb{C}^{2m}$ . It is thus clear that we have to choose  $E_{\rm ts}(\iota) = E_{\iota,\beta_{\rm ts}}^{\infty}$  for  $\mathfrak{g}_{\infty}(\iota) \geq \mu_0$ . Assuming  $\beta_{\rm mar}$  to be small enough, the bound depending only on  $\beta_{\rm ts}$ ,  $\alpha_{\infty}$ ,  $\zeta_{\infty}$  and the limits appearing in (11.1), and then  $\mu_0$  large enough, the bound depending only on  $\beta_{\rm mar}$ ,  $\beta_{\rm ts}$ ,  $\alpha_{\infty}$ ,  $\zeta_{\infty}$  and the limits appearing in (11.1), it can be ensured that the spaces  $E_{\iota} := E_{\rm ts}(\iota)$  have the properties stated in Proposition 12.22 for  $\mathfrak{g}_{\infty}(\iota) \geq \mu_0$ . This is justified by the argument presented in the proof of Lemma 12.18. Given  $\iota \in \mathcal{I}^+_{\mathrm{B},\mu_0}$ , let us specify v by imposing the initial condition  $v(\iota, T_{\rm ts}) := \Psi_{\iota,\infty}[\hat{\chi}(\iota)]$ . Then (12.89) is small asymptotically. In order to estimate  $|v(\iota, T_{\rm ts})|$  in terms of  $|\hat{\chi}(\iota)|$ , define  $u_{\infty}(\iota) \in E_{\iota,\beta_{\rm ts}}^{\infty}$  by

$$e^{-A(\iota)T_{\rm ts}}u_{\infty}(\iota) = \hat{\chi}(\iota). \tag{12.90}$$

Then  $|u_{\infty}| = |e^{A(\iota)T_{\text{ts}}}\hat{\chi}(\iota)|$ , so that (11.109) yields the estimate

$$|v(\iota, T_{\rm ts})| = |\Psi_{\iota,\infty}(e^{-A(\iota)T_{\rm ts}}u_{\infty})| \le C|u_{\infty}(\iota)| = C|e^{A(\iota)T_{\rm ts}}\hat{\chi}(\iota)|$$
(12.91)

where C only depends on  $\beta_{\text{mar}}$ ,  $\beta_{\text{ts}}$ ,  $C_{\text{ts}}$ ,  $\alpha_{\infty}$ ,  $\zeta_{\infty}$  and the limits appearing in (11.1). In order to estimate the far right hand side, note that Lemma 11.23 implies that there are  $\mu_0$  and C, depending only on  $\alpha_{\infty}$ ,  $\zeta_{\infty}$  and the limits appearing in (11.1), such that

$$\|\exp[A(\iota)T_{\rm ts}]\| \le C \exp[(\kappa_{\iota} + 2)T_{\rm ts}]$$

for all  $\iota \in \mathcal{I}^+_{B,\mu_0}$ . Combining this estimate with (12.91) yields

$$|v(\iota, T_{\rm ts})| \le C \exp[(\kappa_{\iota} + 2)T_{\rm ts}]|\hat{\chi}(\iota)|, \qquad (12.92)$$

where C only depends on  $\beta_{\text{mar}}$ ,  $\beta_{\text{ts}}$ ,  $C_{\text{ts}}$ ,  $\alpha_{\infty}$ ,  $\zeta_{\infty}$  and the limits appearing in (11.1).

For  $\iota \in \mathcal{I}_{B,ts}$  such that  $\mathfrak{g}_{\infty}(\iota) < \mu_0$ , we appeal to Lemma 9.21. When doing so, we replace A,  $A_{rem}$  and  $E_{A,\beta}$  by  $A(\iota)$ ,  $A_{\iota,rem}$  and  $E_{A(\iota),\beta_{ts}-\beta_{max}}$  respectively. Moreover, we replace k,  $T_{ode}$ ,  $\beta_{rem}$  and

 $C_{\text{rem}}$  by 2m,  $T_{\text{ts}}$ ,  $\beta_{\text{ts}}$  and  $C_{\text{ts}}$  respectively. Finally, the map  $\Psi_{\infty}$  appearing in Lemma 9.21 we here refer to as  $\Psi_{\iota,\infty}: E_{A(\iota),\beta_{\text{ts}}-\beta_{\text{mar}}} \to \mathbb{C}^{2m}$ . In particular, it is thus clear that for  $\iota \in \mathcal{I}_{\text{B,ts}}$  such that  $\mathfrak{g}_{\infty}(\iota) < \mu_0$ , we let  $E_{\text{ts}}(\iota) := E_{A(\iota),\beta_{\text{ts}}-\beta_{\text{mar}}}$ ; note that this space has the desired properties. For  $\iota \in \mathcal{I}_{\text{B,ts}}$  such that  $\mathfrak{g}_{\infty}(\iota) < \mu_0$ , let  $v(\iota, T_{\text{ts}}) := \Psi_{\iota,\infty}[\hat{\chi}(\iota)]$  as before and define  $u_{\infty}(\iota)$  by (12.90). Then (9.45) yields

$$|v(\iota, T_{\rm ts})| \le C \exp[(\kappa_{\iota} + 2)T_{\rm ts}]|\hat{\chi}(\iota)|, \qquad (12.93)$$

where C only depends on  $C_{\text{ts}}$ ,  $\beta_{\text{ts}}$  and  $A(\iota)$ . However, since there are only finitely many matrices  $A(\iota)$  for  $\mathfrak{g}_{\infty}(\iota) < \mu_0$ , the dependence on  $A(\iota)$  can be replaced by a dependence on  $\mu_0$ ,  $\alpha_{\infty}$ ,  $\zeta_{\infty}$ , the limits appearing in (11.1) and the spectra of the Laplace-Beltrami operators on the Riemannian manifolds  $(M_{r_j}, g_{r_j}), j = 1, \ldots, R_{\text{ts}}$ . Defining  $\kappa$  to be the supremum of the set of  $\kappa_{\iota}$  for  $\iota \in \mathcal{I}_{\text{B,ts}}$  (note that this supremum is finite), (12.92) and (12.93) yield

$$|v(\iota, T_{\rm ts})| \le C \exp[(\kappa + 2)T_{\rm ts}]|\hat{\chi}(\iota)| \tag{12.94}$$

for all  $\iota \in \mathcal{I}_{B,ts}$ , where C only depends on  $\beta_{mar}$ ,  $\beta_{ts}$ ,  $C_{ts}$ ,  $\mu_0$ ,  $\alpha_{\infty}$ ,  $\zeta_{\infty}$ , the limits appearing in (11.1) and the spectra of the Laplace-Beltrami operators on the Riemannian manifolds  $(M_{r_j}, g_{r_j})$ ,  $j = 1, \ldots, R_{ts}$ . In practice we fix  $\mu_0$  to a value depending only on  $\beta_{mar}$ ,  $\beta_{ts}$ ,  $\alpha_{\infty}$ ,  $\zeta_{\infty}$  and the limits appearing in (11.1). Thus the dependence of C on  $\mu_0$  can be eliminated. To conclude, we have defined  $E_{ts}(\iota)$  for  $\iota \in \mathcal{I}_{B,ts}$  and this set has the desired properties; we have defined  $v(\iota, T_{ts})$  in terms of  $\hat{\chi}(\iota)$ ; we know that (12.94) holds; and, finally, we know that if  $v(\iota, T_{ts}) := \Psi_{\iota,\infty}[\hat{\chi}(\iota)]$ , then

$$|v(\iota, t) - e^{A(\iota)t}\hat{\chi}(\iota)| \le C\langle t \rangle^N e^{(\kappa_\iota - \beta_{\rm ts} + \beta_{\rm mar})\overline{t}} |v(\iota, T_{\rm ts})|$$
(12.95)

for  $t \geq T_{\rm ts}$ . If  $\mathfrak{g}_{\infty}(\iota) \geq \mu_0$ , then (12.95) is a consequence of (11.75) with  $F_{\iota} = 0$ ; recall that (11.75) holds as a consequence of Lemma 11.19. If  $\mathfrak{g}_{\infty}(\iota) < \mu_0$ , then (12.95) is a consequence of (9.22) with F = 0; recall that (9.22) holds as a consequence of Lemma 9.21. Finally, the constant C appearing in (12.95) only depends on  $C_{\rm ts}$ ,  $\beta_{\rm ts}$ ,  $\beta_{\rm mar}$ ,  $\mu_0$ ,  $\alpha_{\infty}$ ,  $\zeta_{\infty}$ , the limits appearing in (11.1) and the spectra of the Laplace-Beltrami operators on the Riemannian manifolds  $(M_{r_j}, g_{r_j})$ ,  $j = 1, \ldots, R_{\rm ts}$ . In practice we fix  $\mu_0$  to a value depending only on  $\beta_{\rm mar}$ ,  $\beta_{\rm ts}$ ,  $\alpha_{\infty}$ ,  $\zeta_{\infty}$  and the limits appearing in (11.1). Thus the dependence of C on  $\mu_0$  can be eliminated. From now on, we assume  $\mu_0$  to have been fixed in this way, so that the constants appearing in (12.94) and (12.95) are independent of  $\mu_0$ .

From  $T_{\rm ts}$  to t = 0. Next we wish to estimate  $|v(\iota, 0)|$ . Due to the definition of  $T_{\rm ts}$ , we know that  $T_{\rm ts} > 0$ ; cf. (12.24). Note also that (12.38) holds, where C only depends on  $C_{\rm ini}, c_{\rm sil}, K_{\rm ts}, S_{\rm ts}, \eta_{\rm ts}$ , and  $c_{\rm sil} := \|\mathbf{e}_{\rm sil}\|_1$ . Thus, appealing to (10.12) and (12.38),

$$\mathcal{E}^{1/2}(\iota,0) \le e^{c_{\mathfrak{e}} + \eta_{\mathrm{bal}} T_{\mathrm{ts}}} \mathcal{E}^{1/2}(\iota,T_{\mathrm{ts}}) \le C e^{\eta_{\mathrm{bal}} T_{\mathrm{ts}}} |v(\iota,T_{\mathrm{ts}})|,$$

where C only depends on  $C_{\text{ini}}, c_{\text{sil}}, K_{\text{ts}}, S_{\text{ts}}, \eta_{\text{ts}}$  and  $c_{\mathfrak{e}} := \|\mathfrak{e}\|_1$ ; and  $\eta_{\text{bal}}$  only depends on  $C_{\text{coeff}}$ and  $C_{\ell}$ . Combining this estimate with (12.94) yields

$$\mathcal{E}^{1/2}(\iota, 0) \le C \exp[(\kappa + \eta_{\text{bal}} + 2)T_{\text{ts}}]|\hat{\chi}(\iota)|,$$
(12.96)

where C only depends on  $C_{\text{ini}}, c_{\text{sil}}, K_{\text{ts}}, S_{\text{ts}}, \eta_{\text{ts}}, c_{\mathfrak{e}}, \beta_{\text{mar}}, \beta_{\text{ts}}, C_{\text{ts}}, \mu_0, \alpha_{\infty}, \zeta_{\infty}$ , the limits appearing in (11.1) and the spectra of the Laplace-Beltrami operators on the Riemannian manifolds  $(M_{r_i}, g_{r_i}), j = 1, \ldots, R_{\text{ts}}$ . Thus

$$\langle \nu(\iota) \rangle^{s} (|z(\iota,0)|^{2} + |\dot{z}(\iota,0)|^{2})^{1/2} \le C \langle \nu(\iota) \rangle^{s+s_{\infty}} |\hat{\chi}(\iota)|$$
 (12.97)

for  $\iota \in \mathcal{I}_{B,ts}$ , where  $s_{\infty} := (\kappa + 2 + \eta_{bal})[2\beta_{sil}^{-1} + (\min\{\beta_{sil}, \eta_{ts}\})^{-1}]$  and C only depends on  $C_{ini}, c_{sil}, K_{ts}, S_{ts}, \eta_{ts}, c_{\epsilon}, \beta_{mar}, \beta_{ts}, \beta_{sil}, \kappa, C_{\ell}, C_{coeff}, C_{ts}, \mu_0, \alpha_{\infty}, \zeta_{\infty}$ , the limits appearing in (11.1) and the spectra of the Laplace-Beltrami operators on the Riemannian manifolds  $(M_{r_j}, g_{r_j}), j = 1, \ldots, R_{ts}$ . Let  $\Phi_{\infty}$  be the map taking  $\chi \in C^{\infty}(\bar{M}, \mathbb{C}^{2m}; E_{ts})$  to the distribution whose  $\iota$ 'th mode is given by  $v_0(\iota, 0)$ , where

$$v_0(\iota,t) := \left(\begin{array}{c} z(\iota,t) \\ \dot{z}(\iota,t) \end{array}\right)$$
and z is constructed as above. Then (12.97) implies that (12.86) holds; note that this implies that  $\Phi_{\infty}$  maps  $\chi \in C^{\infty}(\bar{M}, \mathbb{C}^{2m}; E_{ts})$  to an element of  $C_{ts}^{\infty}(\bar{M}, \mathbb{C}^{2m})$ . Since  $\Phi_{\infty}$  is injective on the level of Fourier coefficients (this is a consequence of Lemmas 9.21 and 11.19), it follows that  $\Phi_{\infty}$ is injective.

Verification of the properties of  $\Phi_{\infty}$ . Fix  $\beta_{\text{mar}} > 0$  such that Proposition 12.22 and the above arguments apply. Let, moreover  $\chi \in C^{\infty}(\bar{M}, \mathbb{C}^{2m}; E_{\text{ts}})$  and let u be the solution to (1.2) (with f = 0) satisfying (12.87). Due to Proposition 12.22, we know that (12.88) holds with  $\chi$  replaced by some  $U_{\infty} \in C^{\infty}(\bar{M}, \mathbb{C}^{2m}; E_{\text{ts}})$ . Moreover, the constants C, N and  $\sigma_{\text{hom}}$  have the dependence stated in Proposition 12.22. In order to determine  $U_{\infty}$  in terms of  $\chi$ , note that (12.88) (with  $\chi$ replaced by  $U_{\infty}$ ) implies that

$$\left| v(\iota, t) - e^{A(\iota)t} u_{\infty}(\iota) \right| \le C \langle t \rangle^{N} e^{(\kappa_{\iota} - \beta_{\rm ts} + \beta_{\rm mar})t}$$
(12.98)

where C depends on the initial data and  $\iota$ ; v is defined as in (11.5) where  $z(\iota, \cdot)$  is the  $\iota$ 'th mode of u; and  $u_{\infty}(\iota)$  is the  $\iota$ 'th mode of  $U_{\infty}$ . Due to the construction of  $\Phi_{\infty}$ , we know that the estimate (12.95) holds. Thus

$$\left| e^{A(\iota)t} [\hat{\chi}(\iota) - u_{\infty}(\iota)] \right| \le C \langle t \rangle^{N} e^{(\kappa_{\iota} - \beta_{\rm ts} + \beta_{\rm mar})t}$$

for  $t \geq T_{\text{ts}}$ . Since  $\hat{\chi}(\iota) - u_{\infty}(\iota) \in E_{\text{ts}}(\iota) = E_{\iota}$ , and since  $E_{\iota}$  has the properties stated in Proposition 12.22, this estimate implies that  $\hat{\chi}(\iota) - u_{\infty}(\iota) = 0$ . Thus  $U_{\infty} = \chi$ , and the lemma follows, except for the statement concerning surjectivity.

If there is an  $\epsilon > 0$  such that  $\operatorname{Rsp}[A(\iota)] < \beta_{\operatorname{ts}} - \epsilon$  for all  $\iota \in \mathcal{I}_{\operatorname{B,ts}}$ , then, choosing  $\beta_{\operatorname{mar}}$  to be small enough,  $E_{\operatorname{ts}}(\iota) = \mathbb{C}^{2m}$  for all  $\iota \in \mathcal{I}_{\operatorname{B,ts}}$ . Thus  $\Phi_{\infty}$  is a map from  $C_{\operatorname{ts}}^{\infty}(\bar{M}, \mathbb{C}^{2m})$  to itself. Let  $\psi \in C_{\operatorname{ts}}^{\infty}(\bar{M}, \mathbb{C}^{2m})$ . We wish to demonstrate that  $\psi$  is in the image of  $\Phi_{\infty}$ . Let u be the solution to (1.2) (with f = 0) with initial data given by (12.87), where the right hand side has been replaced by  $\psi$ . Appealing to Proposition 12.22 yields a  $U_{\infty} \in C_{\operatorname{ts}}^{\infty}(\bar{M}, \mathbb{C}^{2m})$  such that (12.63) holds (with f = 0). Let  $\bar{u}$  be the solution to (1.2) with initial data given by (12.87), where the right hand side has been replaced by  $\Phi_{\infty}(U_{\infty})$ . Then, by the above arguments, (12.63) holds (with f = 0) and ureplaced by  $\bar{u}$ . In particular, there is thus a constant C, depending on u,  $\bar{u}$  etc., such that

$$\left\| \begin{pmatrix} (-\Delta_{\infty})^{1/2} u(\cdot,t) \\ u_t(\cdot,t) \end{pmatrix} - \begin{pmatrix} (-\Delta_{\infty})^{1/2} \bar{u}(\cdot,t) \\ \bar{u}_t(\cdot,t) \end{pmatrix} \right\|_{t,s} \le C \langle t \rangle^N e^{-(\beta_{\rm ts} - \beta_{\rm mar})t}$$
(12.99)

for all  $t \ge 0$  and all  $s \in \mathbb{R}$ . Let z and  $\bar{z}$  be the Fourier coefficients of u and  $\bar{u}$  respectively, let v and  $\bar{v}$  be defined in analogy with (11.5), starting with z and  $\bar{z}$  respectively. Finally, let  $V = v - \bar{v}$ . If  $V(\iota, 0) = 0$  for all  $\iota \in \mathcal{I}_{B,ts}$ , then  $u = \bar{u}$  (since  $V(\iota, \cdot)$  solves a homogeneous equation,  $V(\iota, 0) = 0$  for all  $\iota \in \mathcal{I}_{B,ts}$  implies that  $V(\iota, t) = 0$  for all  $\iota \in \mathcal{I}_{B,ts}$  and all  $t \in I$ ), so that  $\Phi_{\infty}(U_{\infty}) = \psi$ . Thus  $\psi$  is in the image of  $\Phi_{\infty}$ . Assume now that there is a  $\iota \in \mathcal{I}_{B,ts}$  such that  $V(\iota, 0) \neq 0$ . Note that  $V(\iota, \cdot)$  is a solution to an equation to which either Lemma 9.21 or Lemma 11.19 applies; cf. the above. In fact, due to the above arguments, for each  $\iota \in \mathcal{I}_{B,ts}$ , there is a map  $\Psi_{\iota,\infty}$  which is linear and bijective. Since  $V(\iota, 0) \neq 0$ , there is a  $0 \neq \chi \in \mathbb{C}^{2m}$  such that  $\Psi_{\iota,\infty}(\chi) = V(\iota, T_{ts})$ , where  $T_{ts}$  is given by (12.24) (for the relevant  $\iota$  under consideration). Due to the above arguments,

$$|V(\iota, t) - e^{A(\iota)t}\chi| \le C \langle t \rangle^N e^{[\kappa_\iota - (\beta_{\rm ts} - \beta_{\rm mar})]t}$$

for all  $t \ge T_{\rm ts}$  and a constant C depending on V,  $\iota$  etc. Here  $\kappa_{\iota} := \kappa_{\rm max}[A(\iota)]$ . On the other hand, (12.99) implies that the same estimate holds with  $\chi$  set to zero. Due to the fact that  $\operatorname{Rsp}A(\iota) < \beta_{\rm ts} - \beta_{\rm mar}$ , these estimates are contradictory. To conclude, every element of  $C^{\infty}_{\rm ts}(\bar{M}, \mathbb{C}^{2m})$  is in the image of  $\Phi_{\infty}$ . The lemma follows.

### Part III

# Averaging over oscillations

### Chapter 13

# Introduction and technical preliminaries

#### 13.1 Introduction

When (1.2) is weakly silent or weakly transparent, the limit equation (obtained by replacing the coefficients in (1.2) by their limits as  $t \to \infty$ ) is either a system of second order ODE's or a system of linear PDE's with constant coefficients. The solutions to the limit equation can then be calculated by solving linear algebra problems. However, it is also of interest to consider situations in which some of the coefficients of the equation exhibit exponential growth asymptotically. In that setting, there is no constant coefficient limit equation with which to compare, and it is less obvious how to proceed. Moreover, if some of the  $q^{ij}$  asymptotically grow exponentially, then the Fourier coefficients of a solution can typically be expected to oscillate with a frequency that grows exponentially. This may seem to be an unpleasant situation to handle. However, when considering such equations in greater detail, it becomes clear that an exponentially growing frequency in some respects simplifies the analysis. The reason for this is the following. Consider a Fourier coefficient of a solution to the homogeneous version of (1.2). This coefficient solves a (typically non-constant coefficient) system of homogeneous ODE's. In the case that the frequency of the oscillations grows exponentially, it is natural to try to approximate the evolution by considering one period of the oscillations at a time. If  $[t_a, t_b]$  is one period of the oscillations, one would like to calculate the matrix, say  $\Phi(t_b; t_a)$ , that maps initial data at  $t_a$  to the corresponding solution at  $t_b$ . Here we are not able to do so. On the other hand, we can approximate  $\Phi(t_b; t_a)$  by calculating the corresponding matrix obtained by freezing the coefficients at  $t_a$  in the original equation. In order for this calculation to be of interest, we need to estimate the error. However, due to the fact that the frequency of the oscillations increases exponentially, the size of the interval  $[t_a, t_b]$  decreases exponentially. Under rather mild conditions on the coefficients of (1.2), the approximation of  $\Phi(t_b;t_a)$  therefore becomes better and better. In the end, we are interested in the asymptotic behaviour as  $t \to \infty$ . We then need to calculate the matrix product of a sequence of matrices of the form  $\Phi(t_b; t_a)$  (and the number of factors in the product tends to infinity asymptotically). In practice we only calculate  $\Phi(t_b; t_a)$  up to some error term, and it is thus necessary to verify that this error term is manageable even when the number of factors in the matrix product tends to infinity. Again, due to the exponential decay of the length of the periods of the oscillations, this can be done.

In the present part of these notes, we develop methods that are suitable for dealing with situations of the above type. However, as we argue in Subsection 13.1.1 below, the arguments are also useful in other contexts. The analysis is quite technical, but we give an overview of the arguments involved in Subsection 13.1.2 below.

#### 13.1.1 Reformulating the equation

The above description gives a rough idea of the argument in the case of an exponentially growing frequency. However, the methods described above are also useful in the case of, e.g., weakly silent equations. In particular, under suitable assumptions they can be used to obtain better (in fact, optimal) estimates for how the energy E of a Fourier coefficient develops over a period of the form  $[0, T_{ode}]$ ; cf. Section 10.3. In order to understand why this is the case, as well as why the methodology described above is useful even more generally, let us reformulate (24.16). Introducing the notation X and  $\sigma$  as in (10.1), the equation (24.16) can be written

$$\ddot{z} + \mathfrak{g}^2 z - 2i\sigma \mathfrak{g} \dot{z} + \alpha \dot{z} + iX\mathfrak{g} z + \zeta z = \hat{f}.$$
(13.1)

Assuming  $\hat{f} = 0$ , (13.1) can be written

$$\dot{v} = Av, \tag{13.2}$$

where

$$v := \begin{pmatrix} \mathfrak{g}z\\ \dot{z} \end{pmatrix}, \quad A := \begin{pmatrix} \dot{\ell} \mathrm{Id}_m & \mathfrak{g} \mathrm{Id}_m\\ -\mathfrak{g} \mathrm{Id}_m - iX - \zeta/\mathfrak{g} & -\alpha + 2i\sigma \mathfrak{g} \mathrm{Id}_m \end{pmatrix}$$
(13.3)

and  $\ell := \ln \mathfrak{g}$ . Say now that the equation is weakly balanced in the sense that (10.2) holds for  $t \geq 0$ , where  $C_{\text{coeff}}$  is independent of  $\iota$ . Assume, moreover, that  $|\sigma|$  decays exponentially and that  $\dot{\ell}$  is bounded. Then it is clear that when  $\mathfrak{g}$  is large, the dominant feature of the evolution associated with (13.2) is oscillations with a period of (roughly)  $2\pi/\mathfrak{g}$ . In fact, it is not even necessary to assume that  $\|\alpha\|$ ,  $\|\zeta\|$  and  $\|X\|$  are bounded; it is sufficient that they are small in comparison with  $\mathfrak{g}$ . Thus considering one period at a time can be expected to be of interest for unbalanced equations. Moreover, even in the case of a weakly silent equation (for which  $\mathfrak{g}$  converges to zero exponentially), there are time intervals during which  $\mathfrak{g}$  is large, and the behaviour is oscillatory. In such situations it therefore also makes sense to consider one period of the oscillations at a time.

#### 13.1.2 Main steps of the analysis

Determining a setting in which it is meaningful to average over one period of the oscillations. Due to the discussion of the previous subsection, it is clear that in regions where  $\mathfrak{g}$  is dominant, the evolution associated with (13.2) is essentially oscillatory. It is therefore of interest to consider the evolution over one period. Moreover, this perspective is of interest even for equations that violate (10.2); i.e., for equations that are not weakly balanced. This heuristic insight is the basis for the present part of these notes. Turning to the actual analysis, it is natural to start by stating the assumptions needed. We do so at the beginning of Chapter 14 below. It is of interest to note that the assumptions are stronger than those made in, e.g., Chapter 10 in some respects, but weaker in other respects. In particular, in Chapter 14 we make assumptions concerning the time derivatives of  $\ell$ , X,  $\alpha$  and  $\zeta$ , even though there are no corresponding assumptions in Definition 10.1. On the other hand, the assumptions stated in Chapter 14 do not exclude the possibility that X,  $\alpha$  and  $\zeta$  might grow exponentially. The reason for making assumptions concerning the time derivatives of, e.g., X,  $\alpha$  and  $\zeta$  is that, in the present setting, we do not want these quantities to vary too much during one period of the oscillations. In Chapter 10, on the other hand, we do not consider the oscillations in detail, and therefore we do not need to control the variation of X,  $\alpha$ and  $\zeta$  during a period of the oscillations.

**Reformulating the equations and deriving rough estimates for one period of the oscillations.** Once suitable assumptions have been stated, it is convenient to reformulate the equations. The purpose of the reformulation is twofold. First, it is convenient to extract the overall growth (or decay) during one period. This is achieved by reformulating the equation to, say,  $\dot{w} = Bw + F$ , where the real part of the trace of B is (essentially) zero. Second, it is convenient to change the time coordinate so that the dominant coefficients of the equation become time independent. Deriving such a reformulation is the subject of Section 14.2 below. Once the reformulation is derived, we define what is meant by one period of the oscillations. In fact, we define one period to be an interval  $[t_a, t_b]$  such that

$$\int_{t_a}^{t_b} \mathfrak{g}(\iota, t) dt = 2\pi$$

In preparation for the analysis of the evolution over one period, we need to estimate how the coefficients of the equation vary in an interval of the form  $[t_a, t_b]$ ; this is the subject of Sections 14.3 and 14.4 below. Once we have gotten this far, it is natural to proceed as described at the beginning of the present introduction. In other words, we, on the one hand, freeze the coefficients and calculate the evolution over one period. Moreover, we, on the other hand, estimate the error associated with this procedure. In the situation of interest in Chapter 14, the formulae for the coefficients of the equation are quite involved; cf. (14.15)-(14.18) below. Carrying out the calculations using these specific coefficients is not very illuminating; it is not so clear what the relevant structure is. Moreover, the calculations involved are quite cumbersome. In order to estimate the evolution associated with freezing the coefficients in the equation, we therefore present an abstract argument in Section 13.2 below. The disadvantage of this procedure is of course that the motivation for the assumptions made in Section 13.2 only becomes clear in Chapter 14. A second important technical step is to compare the evolution associated with freezing the coefficients with the actual evolution. Again, doing this in the particular case of interest makes the analysis obscure and cumbersome. We therefore present an abstract argument in Section 13.3 below. Combining the above ingredients yields rough estimates of the evolution over a period. We write down the result in Section 14.5.

Variables adapted to the evolution over long periods of time. Setting up an iteration. The above analysis is adapted to one particular period of the oscillations. In fact, the variables we use depend on the starting point, say  $t_a$ , of the period  $[t_a, t_b]$  of interest. However, our main interest is in the long term behaviour of solutions. In that setting, it is important to derive estimates for variables which do not depend on the particular period. In Section 15.1 we introduce suitable variables and derive a rough estimate for how they evolve during one period. In fact, the variables are

$$w(\iota,t) := \exp\left(-i\int_0^t \sigma(\iota,s)\mathfrak{g}(\iota,s)ds\right) \left(\begin{array}{c}\mathfrak{g}(\iota,t)z(\iota,t)\\\dot{z}(\iota,t)\end{array}\right).$$

In particular,  $|w(\iota, t)|^2/2$  equals the energy E introduced in (10.6). To first approximation, the evolution over one period is simply a rotation; cf. Lemma 15.1. Even though this observation is of interest, we need more detailed information. We therefore proceed, in Section 15.2, by deriving a more detailed estimate. Once this main estimate has been derived, it is natural to define an iteration. The iteration corresponds to a time sequence  $\{t_k\}$  such that  $[t_k, t_{k+1}]$  is the interval of one period of the oscillations. The purpose of the iteration is to enable us to estimate the evolution over long periods of time. In connection with the introduction of the iteration, we also change the variables slightly, the goal being to make the change in norm over one period of the oscillations as clear as possible. The resulting variables are equivalent to w at the times  $t_k$ . The relevant statements are given in Lemmas 15.7 and 15.11. The final result of the arguments is an iteration on which a more detailed analysis can be based. However, in order to be able to draw more detailed conclusions, we need to estimate matrix products in situations where the number of factors tends to infinity. In order to be able to estimate such products, we need to make additional assumptions. Since there are different types of assumptions that lead to interesting conclusions, the presentation, after Section 15.3, branches off into Parts IV, V and VI.

#### 13.2 On a class of matrix exponentials

As mentioned in the introduction, we are here interested in solutions to ODE's that exhibit oscillatory behaviour. Moreover, we wish to estimate the evolution over one period of the oscillations. The analysis is divided into two steps. First, we estimate the evolution associated with freezing the coefficients of the equations. Second, we estimate the error associated with freezing the coefficients. In the present section, we consider the case of frozen coefficients. However, the statements of the relevant results are abstract in nature; there is no reference to the equation in which we are interested in in the end. The reader interested in a motivation for the particular assumptions we make here is therefore referred to Chapter 14.

We begin the present section with an elementary lemma. The result is not very deep and could be derived under much more general circumstances. However, the precise statement is adapted to the applications of it in the proof of the main result of this section, Lemma 13.3. After the statement and proof of the preliminary Lemma 13.1, we introduce the terminology specifying the matrices we are interested in here; cf. (13.4)-(13.8) below. Given this terminology, we are in a position to state and prove the main result, Lemma 13.3.

**Lemma 13.1.** Let  $a_k \in \mathbb{C}$ ,  $0 \leq k \in \mathbb{Z}$  be a sequence of complex numbers with the following property: there is a constant C > 0 and an integer  $0 \leq l \in \mathbb{Z}$  such that  $|a_k| \leq C \langle k \rangle^l$  for all  $k \geq 0$ . Given  $1 \leq p \in \mathbb{Z}$ ,  $A \in \mathbf{M}_p(\mathbb{C})$  and a real number a, define f(A) and  $f_{abs}(a)$  by

$$f(A) := \sum_{k=0}^{\infty} \frac{a_k}{k!} A^k, \quad f_{abs}(a) := \sum_{k=0}^{\infty} \frac{|a_{k+1}|}{k!} a^k.$$

Let  $A, R \in \mathbf{M}_p(\mathbb{C})$ . Then

$$||f(A+R) - f(A)|| \le f_{abs}(||A|| + ||R||) \cdot ||R||.$$

**Remark 13.2.** Note that if  $a_k = 1$ , then  $f = \exp$  and  $f_{abs} = \exp$ , so that

$$|\exp(A+R) - \exp(A)|| \le \exp(||A|| + ||R||) \cdot ||R||.$$

*Proof.* Note that

$$f(A+R) - f(A) = \sum_{k=1}^{\infty} \frac{a_k}{k!} \int_0^1 \frac{d}{ds} (A+sR)^k ds.$$

Since

$$\left\|\frac{d}{ds}(A+sR)^{k}\right\| = \|R(A+sR)^{k-1} + \dots + (A+sR)^{k-1}R\| \le k\|R\|(\|A\|+\|R\|)^{k-1},$$

we conclude that

$$\begin{split} \|f(A+R) - f(A)\| &\leq \sum_{k=1}^{\infty} \frac{|a_k|}{k!} k \|R\| (\|A\| + \|R\|)^{k-1} \\ &= \|R\| \sum_{k=1}^{\infty} \frac{|a_k|}{(k-1)!} (\|A\| + \|R\|)^{k-1} = \|R\| f_{\text{abs}}(\|A\| + \|R\|). \end{split}$$

The lemma follows.

Next, we wish to calculate the matrix exponential of a complex  $2m \times 2m$ -matrix P. However, we are interested in matrices of a special form. To begin with

$$P := \begin{pmatrix} P_{11} & P_{12} \\ P_{21} & P_{22} \end{pmatrix}, \tag{13.4}$$

where

$$P_{11} := i\xi \mathrm{Id}_m + R_{11} + E_{11}, \quad P_{12} := \mathrm{Id}_m + E_{12}, \tag{13.5}$$

$$P_{21} := -\operatorname{Id}_m + R_{21} + E_{21}, \quad P_{22} := -i\xi \operatorname{Id}_m - R_{11} + E_{22}. \tag{13.6}$$

In these expressions,  $\xi \in \mathbb{R}$  and  $R_{ij}, E_{ij} \in \mathbf{M}_m(\mathbb{C})$ ,  $i, j \in \{1, 2\}$ . Intuitively,  $R_{ij}$  and  $\xi$  should be thought of as "first order terms" that we need to keep track of. However, the  $E_{ij}$  should be thought of as "second order terms" that we don't need to keep track of. As a first, basic, assumption, we require a universal bound on the first and second order terms:

$$|\xi| + ||R_{11}|| + ||R_{21}|| + \sum_{i,j=1}^{2} ||E_{ij}|| \le C_1$$
(13.7)

for some constant  $C_1$ . We here consider the  $E_{ij}$  to be small; they should be thought of as error terms. Similarly,  $||R_{ij}||^2$  should be thought of as small. Finally,  $|\xi|(||R_{11}|| + ||R_{21}||)$  should be thought of as small. Define  $\mathcal{E}$  by

$$\mathcal{E} := \|R_{11}\|^2 + \|R_{21}\|^2 + |\xi|(\|R_{11}\| + \|R_{21}\|) + \sum_{i,j=1}^2 \|E_{ij}\|.$$
(13.8)

Any term which can be bounded by a constant times  $\mathcal{E}$  should be thought of as an error term. Given the above terminology and assumptions, the following lemma holds.

**Lemma 13.3.** Let P,  $P_{ij}$ ,  $\xi$ ,  $R_{ij}$ ,  $E_{ij}$  and  $\mathcal{E}$  be defined as above. Assume, moreover, that (13.7) is satisfied for some  $0 < C_1 \in \mathbb{R}$ . Then

$$\left\| \exp(2\pi sP) - \cos(s\omega)\operatorname{Id}_{2m} - \nu^{-1}\sin(s\omega) \begin{pmatrix} i\xi\operatorname{Id}_m & \operatorname{Id}_m \\ -\operatorname{Id}_m & -i\xi\operatorname{Id}_m \end{pmatrix} - \pi s\sin(s\omega) \begin{pmatrix} R_{21} & 0 \\ 0 & R_{21} \end{pmatrix} - \frac{1}{2}\sin(s\omega) \begin{pmatrix} 2R_{11} & R_{21} \\ R_{21} & -2R_{11} \end{pmatrix} + \pi s\cos(s\omega) \begin{pmatrix} 0 & R_{21} \\ -R_{21} & 0 \end{pmatrix} \right\| \le C\mathcal{E}$$

$$(13.9)$$

for all  $s \in [-1,1]$ , where  $\nu := (1+\xi^2)^{1/2}$ ,  $\omega := 2\pi\nu$  and the constant C only depends on  $C_1$ . In particular,

$$\left\| \exp(\pm 2\pi P) - \cos(\omega) \operatorname{Id}_{2m} \mp \nu^{-1} \sin(\omega) \begin{pmatrix} i\xi \operatorname{Id}_m & \operatorname{Id}_m \\ -\operatorname{Id}_m & -i\xi \operatorname{Id}_m \end{pmatrix} \right\| \\ \pm \pi \begin{pmatrix} 0 & R_{21} \\ -R_{21} & 0 \end{pmatrix} \right\| \leq C\mathcal{E},$$

$$(13.10)$$

where the constant C only depends on  $C_1$ .

**Remark 13.4.** In case  $\xi = 0$ , (13.10) can be simplified to

$$\left\| \exp(\pm 2\pi P) - \operatorname{Id}_{2m} \pm \pi \begin{pmatrix} 0 & R_{21} \\ -R_{21} & 0 \end{pmatrix} \right\| \le C\mathcal{E},$$

where the constant C only depends on  $C_1$ .

**Remark 13.5.** In the course of the proof below, it becomes clear that the structure of P is advantageous when it comes to computing matrix exponentials. In fact, ignoring the error terms  $E_{ij}$  in P yields a matrix Q such that  $Q^2$  is block triangular; cf. (13.12) below. Moreover, ignoring error terms in  $Q^2$ , it is block diagonal. This fact significantly simplifies the calculation of  $Q^{2k}$ , and thereby  $Q^{2k+1}$ . On the other hand, these are the matrices we need to calculate in order to compute the matrix exponential. The main reason why  $Q^2$  has this nice structure is that the sum of the matrices on the diagonal of Q equal zero, the matrix in the top right corner of Q (i.e.,  $Q_{12}$ , using notation analogous to the above) is  $\mathrm{Id}_m$ , and the matrix on the bottom left is  $-\mathrm{Id}_m$  plus lower order terms. It is also of interest to note that  $R_{11}$  does not contribute to the relevant terms in  $Q^2$ .

*Proof.* Note that P = Q + E, where E is defined in terms of the  $E_{ij}$  in the same way as P is defined in terms of the  $P_{ij}$ ; cf. (13.4). Due to Lemma 13.1,

$$\|\exp(2\pi sP) - \exp(2\pi sQ)\| \le 2\pi \exp(2\pi \|Q\| + 2\pi \|E\|) \|E\| \le C\mathcal{E}$$
(13.11)

for  $s \in [-1, 1]$ , where the constant C only depends on  $C_1$ . Here we are only interested in calculating  $\exp(2\pi sP)$  up to an error term. For that reason, it is sufficient to calculate  $\exp(2\pi sQ)$ . Letting

$$Q_{11} = i\xi \mathrm{Id}_m + R_{11}, \quad Q_{12} = \mathrm{Id}_m, \quad Q_{21} = -\mathrm{Id}_m + R_{21}, \quad Q_{22} = -Q_{11},$$

Q is defined in terms of the  $Q_{ij}$  in the same way as P is defined in terms of the  $P_{ij}$ . Note also that

$$Q^{2} = \begin{pmatrix} Q_{11}^{2} + Q_{21} & 0\\ [Q_{21}, Q_{11}] & Q_{11}^{2} + Q_{21} \end{pmatrix}.$$
 (13.12)

Let us calculate

$$Q_{11}^2 + Q_{21} = -(1+\xi^2) \mathrm{Id}_m + R_{21} + 2i\xi R_{11} + R_{11}^2.$$

The last two terms on the right hand side should be thought of as error terms. Moreover,  $[Q_{21}, Q_{11}] = [R_{21}, R_{11}]$ , which is an error term. Thus

$$Q^2 = \left(\begin{array}{cc} L & 0\\ 0 & L \end{array}\right) + F,$$

where

$$L := -(1+\xi^2) \mathrm{Id}_m + R_{21}, \quad F := \begin{pmatrix} 2i\xi R_{11} + R_{11}^2 & 0\\ [R_{21}, R_{11}] & 2i\xi R_{11} + R_{11}^2 \end{pmatrix}$$

By an argument similar to the proof of Lemma 13.1,

$$\sum_{k=0}^{\infty} \frac{1}{(2k)!} (2\pi s Q)^{2k} = \sum_{k=0}^{\infty} \frac{(2\pi s)^{2k}}{(2k)!} \begin{pmatrix} L^k & 0\\ 0 & L^k \end{pmatrix} + O(\mathcal{E})$$
(13.13)

for  $s \in [-1, 1]$ , where the constant implicit in the ordo expression only depends on  $C_1$ . Similarly,

$$\sum_{k=0}^{\infty} \frac{1}{(2k+1)!} (2\pi s Q)^{2k+1} = \sum_{k=0}^{\infty} \frac{(2\pi s)^{2k+1}}{(2k+1)!} \begin{pmatrix} L^k & 0\\ 0 & L^k \end{pmatrix} Q + O(\mathcal{E})$$
(13.14)

for  $s \in [-1, 1]$ , where the constant implicit in the ordo expression only depends on  $C_1$ . Before calculating  $L^k$ , let us note that, using the notation  $\nu$  introduced in the statement of the lemma,  $L = -\nu^2 \mathrm{Id}_m + R_{21}$ . Moreover,

$$L^{k} = (-1)^{k} \nu^{2k} \mathrm{Id}_{m} + k(-1)^{k-1} \nu^{2(k-1)} R_{21} + \sum_{l=2}^{k} \binom{k}{l} (-1)^{k-l} \nu^{2(k-l)} R_{21}^{l}.$$

Let us denote the terms on the right hand side by  $L_{i,k}$ , i = 0, 1, 2, where  $L_{1,0} = L_{2,0} = L_{2,1} = 0$ . For  $k \ge 2$ ,

$$\begin{aligned} \|L_{2,k}\| &\leq \sum_{l=2}^{k} \binom{k}{l} \nu^{2(k-l)} \|R_{21}\|^{l} \leq \|R_{21}\|^{2} \sum_{l=2}^{k} \binom{k}{l} \nu^{2(k-l)} (\max\{1, \|R_{21}\|\})^{l} \\ &\leq \|R_{21}\|^{2} (\nu^{2} + \max\{1, \|R_{21}\|\})^{k}. \end{aligned}$$

Combining this estimate with (13.13) and (13.14) yields

$$\sum_{k=0}^{\infty} \frac{1}{(2k)!} (2\pi sQ)^{2k} = \sum_{k=0}^{\infty} \frac{(2\pi s)^{2k}}{(2k)!} \begin{pmatrix} L_{0,k} + L_{1,k} & 0\\ 0 & L_{0,k} + L_{1,k} \end{pmatrix} + O(\mathcal{E}),$$
$$\sum_{k=0}^{\infty} \frac{1}{(2k+1)!} (2\pi sQ)^{2k+1} = \sum_{k=0}^{\infty} \frac{(2\pi s)^{2k+1}}{(2k+1)!} \begin{pmatrix} L_{0,k} + L_{1,k} & 0\\ 0 & L_{0,k} + L_{1,k} \end{pmatrix} Q + O(\mathcal{E}).$$

Let us calculate

$$\sum_{k=0}^{\infty} \frac{(2\pi s)^{2k}}{(2k)!} L_{0,k} = \sum_{k=0}^{\infty} \frac{(-1)^k}{(2k)!} (s\omega)^{2k} \mathrm{Id}_m = \cos(s\omega) \mathrm{Id}_m,$$
$$\sum_{k=0}^{\infty} \frac{(2\pi s)^{2k+1}}{(2k+1)!} L_{0,k} = \frac{1}{\nu} \sum_{k=0}^{\infty} \frac{(-1)^k}{(2k+1)!} (s\omega)^{2k+1} \mathrm{Id}_m = \nu^{-1} \sin(s\omega) \mathrm{Id}_m$$

Next, let us calculate

$$\sum_{k=0}^{\infty} \frac{(2\pi s)^{2k}}{(2k)!} L_{1,k} = \sum_{k=1}^{\infty} \frac{(2\pi s)^{2k}}{(2k)!} k(-1)^{k-1} \nu^{2(k-1)} R_{21}$$
$$= \frac{\pi s}{\nu} \sum_{k=1}^{\infty} \frac{(-1)^{k-1}}{(2k-1)!} (s\omega)^{2k-1} R_{21} = \frac{\pi s}{\nu} \sin(s\omega) R_{21}.$$

Moreover,

$$\sum_{k=0}^{\infty} \frac{(2\pi s)^{2k+1}}{(2k+1)!} L_{1,k} = \sum_{k=0}^{\infty} \frac{(2\pi s)^{2k+1}}{(2k+1)!} k(-1)^{k-1} \nu^{2(k-1)} R_{21}$$
$$= \frac{1}{2} \sum_{k=0}^{\infty} \frac{(2\pi s)^{2k+1}}{(2k+1)!} (2k+1)(-1)^{k-1} \nu^{2(k-1)} R_{21}$$
$$- \frac{1}{2} \sum_{k=0}^{\infty} \frac{(2\pi s)^{2k+1}}{(2k+1)!} (-1)^{k-1} \nu^{2(k-1)} R_{21}.$$

Similarly to earlier computations, this can be simplified to

$$\sum_{k=0}^{\infty} \frac{(2\pi s)^{2k+1}}{(2k+1)!} L_{1,k} = -\frac{\pi s}{\nu^2} \cos(s\omega) R_{21} + \frac{1}{2\nu^3} \sin(s\omega) R_{21}.$$

Adding up the above computations yields

$$\exp(2\pi sQ) = \cos(s\omega)\operatorname{Id}_{2m} + \frac{\pi s}{\nu}\sin(s\omega)\begin{pmatrix} R_{21} & 0\\ 0 & R_{21} \end{pmatrix} + \nu^{-1}\sin(s\omega)Q + \left[-\frac{\pi s}{\nu^2}\cos(s\omega) + \frac{1}{2\nu^3}\sin(s\omega)\right]\begin{pmatrix} R_{21} & 0\\ 0 & R_{21} \end{pmatrix}Q + O(\mathcal{E}).$$
(13.15)

Note that

$$\frac{1}{\nu^l} = 1 + O(\xi^2)$$

for l = 1, 2, 3 and the constant implicit in the ordo-expression is numerical. Note also that

$$\begin{pmatrix} R_{21} & 0\\ 0 & R_{21} \end{pmatrix} Q = \begin{pmatrix} 0 & R_{21}\\ -R_{21} & 0 \end{pmatrix} + O(\mathcal{E}).$$

Inserting this information into (13.15) yields

$$\exp(2\pi sQ) = \cos(s\omega)\operatorname{Id}_{2m} + \nu^{-1}\sin(s\omega) \begin{pmatrix} i\xi \operatorname{Id}_m & \operatorname{Id}_m \\ -\operatorname{Id}_m & -i\xi \operatorname{Id}_m \end{pmatrix} \\ + \pi s\sin(s\omega) \begin{pmatrix} R_{21} & 0 \\ 0 & R_{21} \end{pmatrix} + \frac{1}{2}\sin(s\omega) \begin{pmatrix} 2R_{11} & R_{21} \\ R_{21} & -2R_{11} \end{pmatrix} \\ - \pi s\cos(s\omega) \begin{pmatrix} 0 & R_{21} \\ -R_{21} & 0 \end{pmatrix} + O(\mathcal{E}).$$

Combining this calculation with (13.11) yields (13.9). In order to derive (13.10), note that

$$\sin(\pm\omega) = \pm \sin[2\pi(\nu-1)] = O(\xi^2), \quad \cos(\pm\omega) = \cos[2\pi(\nu-1)] = 1 + O(\xi^4).$$
(13.16)

In case  $s = \pm 1$ , several of the terms in (13.9) can thus be dropped (or simplified). This yields the estimate (13.10).

# 13.3 Approximating a fundamental solution by a matrix exponential

As described in Subsection 13.1.2, the strategy for estimating the evolution over one period is to first estimate the evolution for the equation with frozen coefficients; this we did in the previous section. The second step is then to estimate the error associated with freezing the coefficients. This is the subject of the present section.

Consider the initial value problem

$$\frac{d\Phi}{ds}(s) = M(s)\Phi(s), \tag{13.17}$$

$$\Phi(s_a) = \mathrm{Id}_k,\tag{13.18}$$

where  $M \in C^{\infty}[I, \mathbf{M}_k(\mathbb{C})]$ ,  $1 \leq k \in \mathbb{Z}$  and I is an open interval containing  $s_a$ . We denote the solution (when evaluated at s) by  $\Phi(s; s_a)$ . It is also of interest to consider

$$\frac{d\Psi}{ds}(s) = \Psi(s)M(s), \tag{13.19}$$

$$\Psi(s_a) = \mathrm{Id}_k,\tag{13.20}$$

with M and  $s_a$  as above. In this case, we denote the solution (when evaluated at s) by  $\Psi(s; s_a)$ . Fix a compact interval J such that  $s_a \in J \subset I$  (the length of J should be thought of as being small; in practice it will correspond to one period of the oscillations). It is then of interest to estimate the differences

$$\|\exp[M(s_a)(s-s_a)] - \Phi(s;s_a)\|, \quad \|\exp[M(s_a)(s-s_a)] - \Psi(s;s_a)\|$$

for  $s \in J$ , and to obtain similar estimates for the inverses.

**Lemma 13.6.** Let  $s_a \in \mathbb{R}$ ,  $1 \leq k \in \mathbb{Z}$ , I be an open interval containing  $s_a$  and  $M \in C^{\infty}[I, \mathbf{M}_k(\mathbb{C})]$ . Let  $\Phi(s; s_a)$  denote the solution to (13.17) and (13.18), evaluated at s, and let  $\Psi(s; s_a)$  denote the solution to (13.19) and (13.20), evaluated at s. Let  $J \subset I$  be a compact interval such that  $s_a \in J$  and assume that there is a constant  $0 < C_1 \in \mathbb{R}$  such that

$$\|M\|_{C(J)} \cdot |J| \le C_1, \tag{13.21}$$

where |J| denotes the length of the interval J. Then

$$\|\exp[M(s_a)(\cdot - s_a)] - \Phi(\cdot; s_a)\|_{C(J)} \le C \|M - M(s_a)\|_{C(J)} \cdot |J|,$$
(13.22)

$$\|\exp[M(s_a)(\cdot - s_a)] - \Psi(\cdot; s_a)\|_{C(J)} \le C\|M - M(s_a)\|_{C(J)} \cdot |J|,$$
(13.23)

where the constant C only depends on an upper bound on  $C_1$ . Moreover,

$$\|\exp[-M(s_a)(\cdot - s_a)] - [\Phi(\cdot; s_a)]^{-1}\|_{C(J)} \le C\|M - M(s_a)\|_{C(J)} \cdot |J|,$$
(13.24)

$$\|\exp[-M(s_a)(\cdot - s_a)] - [\Psi(\cdot; s_a)]^{-1}\|_{C(J)} \le C\|M - M(s_a)\|_{C(J)} \cdot |J|,$$
(13.25)

where the constant C only depends on an upper bound on  $C_1$ .

**Remark 13.7.** In this lemma, we use the notation

$$||M||_{C(J)} = \sup_{s \in J} ||M(s)||$$

*Proof.* Note that if we define  $\Phi$  by

$$\tilde{\Phi}(s;s_a) := \exp[-M(s_a)(s-s_a)]\Phi(s;s_a),$$

184

then

$$\Phi(s; s_a) = \exp[M(s_a)(s - s_a)] + \exp[M(s_a)(s - s_a)][\tilde{\Phi}(s; s_a) - \mathrm{Id}_k].$$
(13.26)

It is therefore of interest to estimate  $\|\tilde{\Phi}(s;s_a) - \mathrm{Id}_k\|$ . Note, to this end, that

$$\frac{d\bar{\Phi}}{ds}(s;s_a) = \exp[-M(s_a)(s-s_a)][M(s) - M(s_a)] \exp[M(s_a)(s-s_a)]\tilde{\Phi}(s;s_a)$$

Due to (13.21),

$$\left\|\tilde{\Phi}(s;s_{a}) - \mathrm{Id}_{k}\right\| \leq \left|\int_{s_{a}}^{s} \left\|\frac{d\tilde{\Phi}}{dt}(t;s_{a})\right\| dt\right| \leq C \left|\int_{s_{a}}^{s} \|M(t) - M(s_{a})\| \cdot \|\tilde{\Phi}(t;s_{a})\| dt\right|$$
(13.27)

for  $s \in J$ , where the constant C only depends on an upper bound on  $C_1$ ; note that

$$\|\exp[M(s_a)(s-s_a)]\| \le \exp[\|M(s_a)\| \cdot |J|] \le e^{C_1}$$
(13.28)

for  $s \in J$  etc. due to (13.21), and that s could be strictly less than  $s_a$  (this is the reason we include the absolute value signs outside the integrals even though the integrands are non-negative). The estimate (13.27) implies

$$\|\tilde{\Phi}(s;s_a)\| \le 1 + C \left| \int_{s_a}^s \|M(t) - M(s_a)\| \cdot \|\tilde{\Phi}(t;s_a)\| dt \right|.$$

Appealing to Grönwall's lemma then yields

$$\|\tilde{\Phi}(s;s_a)\| \le \exp\left(C\left|\int_{s_a}^s \|M(t) - M(s_a)\|dt\right|\right) \le \exp(2CC_1)$$

for  $s \in J$ , where C is a constant which only depends on an upper bound on  $C_1$ . Returning to (13.27), we conclude that

$$\sup_{s\in J} \|\tilde{\Phi}(s;s_a) - \mathrm{Id}_k\| \le C \sup_{s\in J} \|M(s) - M(s_a)\| \cdot |J|,$$

where the constant C only depends on an upper bound on  $C_1$ . Combining this estimate with (13.26) and (13.28) yields (13.22). Turning to (13.23), note that if we use the symbol  $\dagger$  to indicate the conjugate transpose of a square complex matrix, then  $\Psi^{\dagger}$  satisfies (13.17) and (13.18) with M replaced by  $M^{\dagger}$ . Moreover, since  $\|L^{\dagger}\| = \|L\|$  for any complex square matrix L, it is clear that (13.21) is satisfied with M replaced by  $M^{\dagger}$ . We are thus allowed to appeal to (13.22) in order to obtain

$$\sup_{s \in J} \|\exp[M^{\dagger}(s_a)(s-s_a)] - \Psi^{\dagger}(s;s_a)\| \le C \sup_{s \in J} \|M^{\dagger}(s) - M^{\dagger}(s_a)\| \cdot |J|.$$

Since

$$\exp[M^{\dagger}(s_{a})(s-s_{a})] - \Psi^{\dagger}(s;s_{a}) = \left\{\exp[M(s_{a})(s-s_{a})] - \Psi(s;s_{a})\right\}^{\dagger}$$

and since the conjugate transpose is an isometry, we conclude that (13.23) holds. In order to prove (13.24) and (13.25), note that

$$\frac{d\Phi^{-1}}{ds} = -\Phi^{-1}\frac{d\Phi}{ds}\Phi^{-1} = -\Phi^{-1}M, \quad \frac{d\Psi^{-1}}{ds} = -\Psi^{-1}\frac{d\Psi}{ds}\Psi^{-1} = -M\Psi^{-1}.$$

Appealing to (13.22) and (13.23) with M replaced by -M yields (13.24) and (13.25).

#### 13.4 Inhomogeneous equations

In the following chapters, we are interested in estimating the evolution of solutions to inhomogeneous equations. In the present section, we indicate how the above analysis can be used to that end.

Consider an equation of the form

$$\frac{dv}{ds} = M(s)v + F(s). \tag{13.29}$$

Here v is  $\mathbb{C}^{2m}$  valued, and v and M can be written

$$v = \begin{pmatrix} x \\ y \end{pmatrix}, \quad M = \begin{pmatrix} M_{11} & M_{12} \\ M_{21} & M_{22} \end{pmatrix}$$

where x and y each take their values in  $\mathbb{C}^m$  and the  $M_{ij}$  take their values in  $\mathbf{M}_m(\mathbb{C})$ . Fix  $s_a$  in the domain of definition, say I, of M and let  $J \subset I$  be a compact interval. The dominant term in  $M(s_a)$  should be thought of as being

$$\left(\begin{array}{cc} 0 & \nu_0 \mathrm{Id}_m \\ -\nu_0 \mathrm{Id}_m & 0 \end{array}\right).$$

where  $\nu_0$  is a large positive number, and the interval J should be thought of as roughly corresponding to one period of the associated oscillation (with  $s_a$  being either the left or the right endpoint of J). Let  $\Phi(s; s_a)$  denote the solution to (13.17) and (13.18), evaluated at s. Then the solution to (13.29) can be written

$$v(s) = \Phi(s; s_a)v(s_a) + \Phi(s; s_a) \int_{s_a}^{s} [\Phi(u; s_a)]^{-1} F(u) du.$$

We are mainly interested in the value of v at  $s_b$ , where  $s_b$  is the endpoint of J which is different from  $s_a$ . We are therefore particularly interested in estimating  $\Phi(s_b; s_a)$  and  $[\Phi(u; s_a)]^{-1}$  for  $u \in J$ . However, we can appeal to Lemma 13.6 in order to approximate these matrices using matrix exponentials. Moreover, the matrix exponentials of interest are of the form considered in Lemma 13.3.

### Chapter 14

## One period of the oscillations; terminology and preliminary estimates

In the present part of these notes, we are interested in solutions to (24.16) in situations where the behaviour is oscillatory. Moreover, due to the applications we have in mind, the arguments should apply even when estimates of the form (10.2) are violated. It is therefore necessary to formulate assumptions that are different from the ones we have made previously. We do so in Section 14.1 below. Once the appropriate assumptions have been made, we need to reformulate (24.16) to a form suitable for appealing to the results of Section 13.2. We take a first step in this direction in Section 14.2. Most of the remaining part of the chapter is devoted to verifying that not only can the equation be reformulated to an appropriate form, its coefficients also satisfy bounds such that the results of Section 13.3 apply.

The assumptions stated in Section 14.1 below involve bounds on, e.g.,  $\dot{\ell}$  and  $\ddot{\ell}$ , where  $\ell$  is introduced in (10.1). Such bounds are of central importance, since we are considering situations in which the leading order behaviour is oscillatory, and the frequency of the oscillations is roughly speaking given by  $\mathfrak{g}$ . As an initial step in the analysis, we therefore first define what we mean by one period of the oscillations, and then analyse the variation of expressions involving  $\mathfrak{g}$  over one such period. This is the subject of Section 14.3. Given this information, we are then in a position to analyse the variation of the remaining coefficients of the equation; cf. Section 14.4. To end the chapter, we, in Section 14.5, obtain a first rough estimate of the evolution over one period.

#### 14.1 Assumptions

Definitions 10.1 and 12.8 are two examples of assumptions that we have made concerning the coefficients of (1.2) so far. However, they are not appropriate in the present context. The reason for this is that, on the one hand, we are here interested in unbalanced equations, which are excluded by Definition 10.1; in that sense the assumptions in Definition 10.1 are too strong. On the other hand, we also need to control the variation of the coefficients over one period of the oscillations. We therefore need to bound, e.g.,  $\|\dot{\alpha}\|$ . In that sense, Definition 10.1 is too weak. Even though we here admit unbalanced equations, we do not want, e.g.,  $\|\alpha\|$  to grow too quickly. For that reason, we impose restrictions that imply that  $\|\alpha\|$  etc. cannot grow faster than exponentially.

**Definition 14.1.** Consider (1.2). Assume the associated metric to be such that (M, g) is a canonical separable cosmological model manifold. Define  $\mathfrak{g}$  by (24.13) and  $\ell$ ,  $\sigma$  and X by (10.1). Then (1.2) is said to be *oscillation adapted* if there are functions  $\mathfrak{f}_{\mathrm{sh}}, \mathfrak{f}_X, \mathfrak{f}_{\mathrm{ode}} \in C^{\infty}(\mathbb{R}, \mathbb{R}_+)$ , where

 $\mathbb{R}_+ := (0, \infty)$ , and a constant  $0 < \mathfrak{b} \in \mathbb{R}$  such that the following holds. First,

$$|\dot{\ell}(\iota,t)| + |\ddot{\ell}(\iota,t)| \le \mathfrak{b} \tag{14.1}$$

for all  $t \geq 0$  and all  $0 \neq \iota \in \mathcal{I}_{B}$ . Second,

$$|\sigma(\iota, t)| + |\dot{\sigma}(\iota, t) \leq \mathfrak{f}_{\rm sh}(t), \tag{14.2}$$

$$\|X(\iota, t)\| + \|\dot{X}(\iota, t)\| \le \mathfrak{f}_X(t), \tag{14.3}$$

$$\|\dot{\alpha}(t)\| + \|\alpha(t)\| \le \mathfrak{f}_{\text{ode}}(t),\tag{14.4}$$

$$\|\zeta(t)\| + \|\zeta(t)\| \le \mathfrak{f}_{ode}^2(t) \tag{14.5}$$

for all  $t \geq 0$  and all  $0 \neq \iota \in \mathcal{I}_{B}$ . Third,

$$\frac{|\dot{\mathfrak{f}}_{\mathrm{sh}}(t)|}{\mathfrak{f}_{\mathrm{sh}}(t)} + \frac{|\dot{\mathfrak{f}}_X(t)|}{\mathfrak{f}_X(t)} + \frac{|\dot{\mathfrak{f}}_{\mathrm{ode}}(t)|}{\mathfrak{f}_{\mathrm{ode}}(t)} \le \mathfrak{b}$$
(14.6)

for all  $t \geq 0$ .

**Remark 14.2.** The constant  $\mathfrak{b}$  and the functions  $\mathfrak{f}_{sh}, \mathfrak{f}_X, \mathfrak{f}_{ode}$  are all independent of  $\iota \in \mathcal{I}_B$ .

**Remarks 14.3.** In order to motivate the requirements, note that we do not want the coefficients  $g^{jl}$  and  $a_r^{-2}$  to vary too quickly over time. This is the reason for the bound on  $\dot{\ell}$ ; it ensures that  $\mathfrak{g}$  does not grow or decay faster than exponentially. The bounds concerning  $\sigma$ , X,  $\alpha$  and  $\zeta$  are still quite general. However, it is clear from (14.6) that the right hand sides of (14.2)–(14.5) are not allowed to grow faster than exponentially. Moreover, the bound (14.6), in the end, ensures that the variation of the functions  $\mathfrak{f}_{sh}, \mathfrak{f}_X, \mathfrak{f}_{ode}$  over one period of the oscillations is negligible. Nevertheless, we impose more detailed restrictions on these functions below.

#### 14.2 Change of variables

In order to analyse how the solution evolves over one period of the oscillations, it is convenient to change variables (so that we can apply, e.g., Lemma 13.3). It would be preferable if it were possible to change the variables once and for all. However, that turns out to lead to technical difficulties. For that reason we here fix one time, say  $0 \le t_a \in \mathbb{R}$ , and then adapt the variables so that we are in a position to analyse the variation of solutions in a small neighbourhood of  $t_a$ .

**Change of variables.** Fix  $t_a \ge 0$ , and let  $U(t; t_a)$  denote the matrix valued solution (evaluated at t) to the initial value problem

$$U_t(t;t_a) = \frac{1}{2}\alpha(t)U(t;t_a), \quad U(t_a;t_a) = \mathrm{Id}_m.$$

Let

$$\begin{pmatrix} x_1\\ y_1\\ F_1 \end{pmatrix} := \exp\left(-i\int_0^t \sigma(s)\mathfrak{g}(s)ds\right) \begin{pmatrix} \mathfrak{g}z\\ \dot{z}\\ \hat{f} \end{pmatrix}, \qquad (14.7)$$

where z is a solution to (24.16) and we suppress all the arguments but the variable of integration. Define x, y and  $F_2$  by

$$\begin{pmatrix} x(\iota,t) \\ y(\iota,t) \\ F_2(\iota,t) \end{pmatrix} := \exp\left[-\frac{1}{2}\int_{t_a}^t \dot{\ell}(\iota,s)ds\right] \begin{pmatrix} U(t;t_a)x_1(\iota,t) \\ U(t;t_a)y_1(\iota,t) \\ U(t;t_a)F_1(\iota,t) \end{pmatrix}.$$
(14.8)

Given x and y defined by (14.8), let v be defined by

$$v := \begin{pmatrix} x \\ y \end{pmatrix}. \tag{14.9}$$

#### 14.2. CHANGE OF VARIABLES

Appealing to (24.16), it can be computed that

$$\dot{x} = \frac{1}{2} (\alpha + \dot{\ell} \mathrm{Id}_m - 2i\sigma \mathfrak{g} \mathrm{Id}_m) x + \mathfrak{g} y, \qquad (14.10)$$

$$\dot{y} = -\mathfrak{g}x - \frac{1}{2}(\alpha + \dot{\ell}\mathrm{Id}_m - 2i\sigma\mathfrak{g}\mathrm{Id}_m)y + [\alpha, U(\cdot; t_a)][U(\cdot; t_a)]^{-1}y$$
(14.11)

$$-U(\cdot;t_a)(iX+Z)[U(\cdot;t_a)]^{-1}x+F_2,$$

where  $Z(\iota, t) := \zeta(t)/\mathfrak{g}(\iota, t)$ . At this stage, it is natural to pause to consider what has been gained by this reformulation. Freezing the coefficients of x and y in (14.10) and (14.11) at  $t = t_a$ , interpreting the frozen coefficients as the components of a matrix in  $\mathbf{M}_{2m}(\mathbb{C})$ , and dividing the matrix by  $\mathfrak{g}(t_a)$  yields

$$P_a := \begin{pmatrix} R_{11} + i\xi \mathrm{Id}_m & \mathrm{Id}_m \\ -\mathrm{Id}_m + R_{21} + E_{21} & -R_{11} - i\xi \mathrm{Id}_m \end{pmatrix},$$
(14.12)

where

$$R_{11} := \frac{1}{2\mathfrak{g}(\iota, t_a)} [\alpha(t_a) + \dot{\ell}(\iota, t_a) \mathrm{Id}_m], \quad R_{21} := -i\frac{X(\iota, t_a)}{\mathfrak{g}(\iota, t_a)}, \quad E_{21} := -\frac{Z(\iota, t_a)}{\mathfrak{g}(\iota, t_a)}$$

and  $\xi := -\sigma(\iota, t_a)$ . Thus  $P_a$  has the structure considered in Section 13.2. The advantage of this structure is discussed in Remark 13.5; cf. also the statement of Lemma 13.3.

**Change of time coordinate.** The above discussion demonstrates that freezing the coefficients of (14.10) and (14.11) yields a matrix of a form appropriate for appealing to the results of Section 13.2. However, when estimating the error associated with freezing the coefficients, it is convenient to change the time coordinate so that the largest coefficients appearing in the equations for x and y are constant. To that end, we introduce a new time coordinate

$$\tau(\iota, t) := \int_{t_0}^t \frac{\mathfrak{g}(\iota, s)}{\mathfrak{g}(\iota, t_0)} ds.$$
(14.13)

Here  $0 \leq t_0 \in \mathbb{R}$  is a parameter which is typically different from  $t_a$ . The purpose of introducing such a parameter is to indicate a starting point at which  $\mathfrak{g}$  is sufficiently large for the behaviour to be oscillatory. On the other hand, we wish to use the same coordinate  $\tau$  for many periods of the oscillations close to  $t_0$ . That is the reason we do not necessarily require  $t_0$  to equal  $t_a$ . Roughly speaking,  $t_0$  fixes an oscillatory "era", whereas  $t_a$  fixes the starting point of one period of the oscillations. In what follows, it is convenient to suppress the dependence of  $\tau$  on  $\iota$ , and we often do so in what follows. Moreover, we abuse notation in the way it is normally done when changing coordinates; we think of  $\tau$  as being a function and a real number interchangeably; and we use the notation  $\alpha$ , e.g., for both the function mapping t to  $\alpha(t)$  and the function mapping  $\tau$  to  $\alpha[t(\tau)]$ etc. Changing time coordinate according to (14.13) yields

$$\begin{split} \frac{dx}{d\tau} &= \frac{\mathfrak{g}_0}{2\mathfrak{g}} (\alpha + \dot{\ell} \mathrm{Id}_m - 2i\sigma \mathfrak{g} \mathrm{Id}_m) x + \mathfrak{g}_0 y, \\ \frac{dy}{d\tau} &= -\mathfrak{g}_0 x - \frac{\mathfrak{g}_0}{2\mathfrak{g}} (\alpha + \dot{\ell} \mathrm{Id}_m - 2i\sigma \mathfrak{g} \mathrm{Id}_m) y + \frac{\mathfrak{g}_0}{\mathfrak{g}} \left[ \alpha, U \right] U^{-1} y \\ &- U \frac{\mathfrak{g}_0}{\mathfrak{g}} (iX + Z) U^{-1} x + F_3, \end{split}$$

where  $\mathfrak{g}_0(\iota) := \mathfrak{g}(\iota, t_0),$ 

$$F_3 := \frac{\mathfrak{g}_0}{\mathfrak{g}} F_2 \tag{14.14}$$

and we, in this case, have omitted all the arguments. Moreover, we think of  $\alpha$  as the function of  $\tau$  mapping  $\tau$  to  $\alpha[t(\tau)]$  etc. Finally, the dots appearing still refer to the time coordinate t.

Matrix formulation of the equations. Define

$$A_{11} := \frac{1}{2} \frac{\alpha \mathfrak{g}_0}{\mathfrak{g}} + \frac{1}{2} \frac{\dot{\ell} \mathfrak{g}_0}{\mathfrak{g}} \mathrm{Id}_m - i\sigma \mathfrak{g}_0 \mathrm{Id}_m, \qquad (14.15)$$

$$A_{12} := \mathfrak{g}_0 \mathrm{Id}_m, \tag{14.16}$$

$$A_{21} := -\mathfrak{g}_0 \mathrm{Id}_m - U \frac{\mathfrak{g}_0}{\mathfrak{g}} (iX + Z) U^{-1}, \qquad (14.17)$$

$$A_{22} := -A_{11} + \frac{\mathfrak{g}_0}{\mathfrak{g}} [\alpha, U] U^{-1}.$$
(14.18)

Let A and F be defined by

$$A := \begin{pmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{pmatrix}, \quad F := \begin{pmatrix} 0 \\ F_3 \end{pmatrix}$$
(14.19)

With this notation, (24.16) yields

$$\frac{dv}{d\tau} = Av + F. \tag{14.20}$$

#### 14.3 Choosing appropriate time intervals

Our long term goal is to estimate how the energy of solutions to (14.20) evolve over time. To this end, it is natural to divide the interval  $[0, \infty)$  into different regions. Consider the matrix A defined by (14.19) and (14.15)–(14.18). In intervals where the expression  $\mathfrak{g}_0 \mathrm{Id}_m$ , appearing in  $A_{12}$  and  $A_{21}$ , dominates, the behaviour can be expected to be oscillatory; below, we loosely speak of such intervals as "oscillatory regions" or "oscillatory eras" (though we do not provide a formal definition of the meaning of this terminology). Considering the homogeneous equation associated with (14.20), we then, roughly speaking, expect oscillations with a period of  $2\pi/\mathfrak{g}_0$ . It is therefore of interest to analyse how solutions to (14.20) evolve over one such period. Keeping (14.13) in mind, a period of  $2\pi/\mathfrak{g}_0$  in  $\tau$ -time corresponds to an interval  $[t_a, t_b]$  (or  $[t_b, t_a]$ ) in t-time such that  $t_a, t_b \geq 0$  and such that

$$\left| \int_{t_a}^{t_b} \mathfrak{g}(\iota, t) dt \right| = 2\pi, \tag{14.21}$$

where we include the absolute value sign in order to allow for the possibility that  $t_b$  might be strictly smaller than  $t_a$ . To begin with, it is thus natural to verify that there are time intervals with this property and to analyse the variation of quantities such as  $\mathfrak{g}$  in such intervals.

**Lemma 14.4.** Assume that (1.2) is oscillation adapted, let  $0 \neq \iota \in \mathcal{I}_B$  and define  $\mathfrak{g}$  by (24.13). Assume that there is a  $t_a \geq 0$  such that

$$\mathfrak{g}(\iota, t_a) \ge \max\{4\pi\mathfrak{b}, 2\},\tag{14.22}$$

where  $\mathfrak{b}$  is the constant appearing in (14.1). Then there is a  $t_b \geq t_a$  such that (14.21) holds. If, in addition to (14.22),  $t_a \geq 2\pi$ , then there is a  $0 \leq t_b < t_a$  such that (14.21) holds. Fix a  $t_b \geq 0$ such that (14.21) holds and let  $I_a$  be the interval with endpoints  $t_a$  and  $t_b$ . Then  $\mathfrak{g}(\iota, t) \geq 1$  and

$$\frac{1}{2} \le \frac{\mathfrak{g}(\iota, t)}{\mathfrak{g}(\iota, t_a)} \le 2, \tag{14.23}$$

$$\frac{\pi}{\mathfrak{g}(\iota, t_a)} \le |I_a| \le \frac{4\pi}{\mathfrak{g}(\iota, t_a)} \tag{14.24}$$

for all  $t \in I_a$ , where  $|I_a|$  denotes the length of  $I_a$ . Finally,

$$\left|\frac{1}{\mathfrak{g}(\iota,t)} - \frac{1}{\mathfrak{g}(\iota,t_a)}\right| \le \frac{C}{\mathfrak{g}^2(\iota,t_a)},\tag{14.25}$$

$$\left|\frac{\dot{\mathfrak{g}}(\iota,t)}{\mathfrak{g}^{2}(\iota,t)} - \frac{\dot{\mathfrak{g}}(\iota,t_{a})}{\mathfrak{g}^{2}(\iota,t_{a})}\right| \leq \frac{C}{\mathfrak{g}^{2}(\iota,t_{a})},\tag{14.26}$$

$$\left| |I_a| - \frac{2\pi}{\mathfrak{g}(\iota, t_a)} \right| \le \frac{C}{\mathfrak{g}^2(\iota, t_a)}, \tag{14.27}$$

$$\left| \int_{I_a} \frac{\dot{\mathfrak{g}}(\iota, s)}{\mathfrak{g}(\iota, s)} ds - \frac{2\pi \dot{\mathfrak{g}}(\iota, t_a)}{\mathfrak{g}^2(\iota, t_a)} \right| \le \frac{C}{\mathfrak{g}^2(\iota, t_a)}$$
(14.28)

for all  $t \in I_a$ , where the constant C only depends on the parameter  $\mathfrak{b}$  appearing in the estimate (14.1).

Remark 14.5. In the statement of the lemma, we use the convention that

$$\int_{t_a}^{t_b} f(s)ds = \operatorname{sgn}(t_b - t_a) \int_{I_a} f(s)ds,$$

where sgn(t) denotes the sign of t.

**Remark 14.6.** In the proof, we omit reference to the argument  $\iota$  for the sake of brevity.

*Proof.* To begin with, (14.1) implies that  $\mathfrak{g}(t) \ge \mathfrak{g}(t_a) \exp[-\mathfrak{b}|t - t_a|]$ . Thus

$$\left|\int_{t_a}^t \mathfrak{g}(s)ds\right| \ge \left|\int_{t_a}^t \mathfrak{g}(t_a) \exp[-\mathfrak{b}|s - t_a|]ds\right| = \mathfrak{b}^{-1}\mathfrak{g}(t_a)[1 - e^{-\mathfrak{b}|t - t_a|}]$$
(14.29)

for  $t \ge 0$ . Assuming  $\mathfrak{g}(t_a) \ge 4\pi\mathfrak{b}$  and  $t \ge 0$  to be such that  $\mathfrak{b}|t - t_a| \ge \ln 2$ , it is clear that the right hand side is bounded from below by  $2\pi$ . In particular, there is thus a  $t_b \ge t_a$  such that (14.21) holds. A more crude way to estimate the left hand side of (14.29) is the following:

$$\left|\int_{t_a}^{t} \mathfrak{g}(s)ds\right| \ge |t - t_a|\mathfrak{g}(t_a)e^{-\mathfrak{b}|t - t_a|}.$$
(14.30)

Combining (14.29) and (14.30) yields the following conclusion. Assume that (14.22) holds and that  $t_a \ge 2\pi$ . Then there are two possibilities. Either  $\mathfrak{b}t_a \ge \ln 2$ , in which case (14.29) implies that there is a  $0 \le t_b < t_a$  such that (14.21) holds, or  $\mathfrak{b}t_a \le \ln 2$ , in which case (14.30) yields the same conclusion. The statements concerning the existence of  $t_b$  follow.

Fix a  $t_b \ge 0$  such that (14.21) holds and define  $I_a$  as in the statement of the lemma. Then (14.29) with  $t = t_b$  yields

$$2\pi = \left| \int_{t_a}^{t_b} \mathfrak{g}(t) dt \right| \ge \mathfrak{b}^{-1} \mathfrak{g}(t_a) [1 - e^{-\mathfrak{b} l_a}] \ge 4\pi [1 - e^{-\mathfrak{b} l_a}],$$

where  $l_a := |I_a|$ . Thus  $e^{-\mathfrak{b}l_a} \ge 1/2$ . Combining this estimate with (14.1) yields

$$\frac{1}{2} \le e^{-\mathfrak{b}l_a} \le \frac{\mathfrak{g}(t)}{\mathfrak{g}(t_a)} \le e^{\mathfrak{b}l_a} \le 2 \tag{14.31}$$

for  $t \in I_a$ . This estimate implies (14.23). Moreover, combining (14.22) with (14.23) yields the conclusion that  $\mathfrak{g}(t) \geq 1$  for all  $t \in I_a$ . The estimate (14.31) also yields

$$\frac{1}{2}\mathfrak{g}(t_a)l_a \le \left|\int_{t_a}^{t_b}\mathfrak{g}(t)dt\right| \le 2\mathfrak{g}(t_a)l_a$$

Combining this estimate with (14.21) yields (14.24).

We are now in a position to turn to the variation of  $\mathfrak g$  and  $\dot{\mathfrak g}$  etc. during one period. To begin with, note that

$$\begin{aligned} \left| \frac{1}{\mathfrak{g}(t)} - \frac{1}{\mathfrak{g}(t_a)} \right| &= \frac{1}{\mathfrak{g}(t_a)} \left| \frac{\mathfrak{g}(t_a)}{\mathfrak{g}(t)} - 1 \right| \le \frac{1}{\mathfrak{g}(t_a)} \left( e^{\mathfrak{b}|t - t_a|} - 1 \right) \le \frac{1}{\mathfrak{g}(t_a)} \left( e^{\mathfrak{b}l_a} - 1 \right) \\ &\le \frac{1}{\mathfrak{g}(t_a)} \left( \exp\left[\frac{4\pi\mathfrak{b}}{\mathfrak{g}(t_a)}\right] - 1 \right) \le \frac{C}{\mathfrak{g}^2(t_a)} \end{aligned}$$

for  $t \in I_a$ , where the constant C only depends on the parameter  $\mathfrak{b}$  appearing in the assumption (14.1). In particular, (14.25) holds. Let us turn to the variation of  $\dot{\mathfrak{g}}/\mathfrak{g}^2$ . Note, to begin with, that

$$\left|\frac{\dot{\mathfrak{g}}(t)}{\mathfrak{g}(t)} - \frac{\dot{\mathfrak{g}}(t_a)}{\mathfrak{g}(t_a)}\right| \le \left|\int_{t_a}^t \ddot{\ell}(s)ds\right| \le \mathfrak{b}l_a \le \frac{C}{\mathfrak{g}(t_a)} \tag{14.32}$$

for  $t \in I_a$ , where the constant C only depends on the parameter  $\mathfrak{b}$  appearing in the assumption (14.1). Due to this estimate,

$$\left|\frac{\dot{\mathfrak{g}}(t)}{\mathfrak{g}^{2}(t)} - \frac{\dot{\mathfrak{g}}(t_{a})}{\mathfrak{g}^{2}(t_{a})}\right| \leq \left|\frac{1}{\mathfrak{g}(t)} \left(\frac{\dot{\mathfrak{g}}(t)}{\mathfrak{g}(t)} - \frac{\dot{\mathfrak{g}}(t_{a})}{\mathfrak{g}(t_{a})}\right)\right| + \left|\left(\frac{1}{\mathfrak{g}(t)} - \frac{1}{\mathfrak{g}(t_{a})}\right) \frac{\dot{\mathfrak{g}}(t_{a})}{\mathfrak{g}(t_{a})}\right| \leq \frac{C}{\mathfrak{g}^{2}(t_{a})}$$

for  $t \in I_a$ , where the constant C only depends on the parameter  $\mathfrak{b}$  appearing in the assumption (14.1), and we have used (14.25) and (14.23). In particular, (14.26) holds. Next, let us derive a better estimate for  $l_a$ . To begin with,

$$2\pi = \int_{I_a} \mathfrak{g}(t)dt = \int_{I_a} \mathfrak{g}(t)\mathfrak{g}(t_a) \left[\frac{1}{\mathfrak{g}(t_a)} - \frac{1}{\mathfrak{g}(t)}\right]dt + l_a\mathfrak{g}(t_a).$$

Thus

$$\left|l_a - \frac{2\pi}{\mathfrak{g}(t_a)}\right| \leq \int_{I_a} \mathfrak{g}(t) \left|\frac{1}{\mathfrak{g}(t_a)} - \frac{1}{\mathfrak{g}(t)}\right| dt \leq \frac{C}{\mathfrak{g}^2(t_a)},$$

where the constant C only depends on the parameter  $\mathfrak{b}$  appearing in the assumption (14.1) and we have appealed to (14.23)–(14.25). Thus (14.27) holds. Next, note that

$$\left|\int_{I_a} \frac{\dot{\mathfrak{g}}(s)}{\mathfrak{g}(s)} ds - \frac{\dot{\mathfrak{g}}(t_a)}{\mathfrak{g}(t_a)} l_a\right| \leq \int_{I_a} \left| \frac{\dot{\mathfrak{g}}(s)}{\mathfrak{g}(s)} - \frac{\dot{\mathfrak{g}}(t_a)}{\mathfrak{g}(t_a)} \right| ds \leq \frac{C}{\mathfrak{g}^2(t_a)},$$

where the constant C only depends on the parameter  $\mathfrak{b}$  appearing in the assumption (14.1) and we have appealed to (14.24) and (14.32). Combining this estimate with (14.1) and (14.27) yields (14.28). The lemma follows.

#### 14.4 Variation of the coefficients during one period

In order to estimate the behaviour of solutions to (14.20), it is convenient to appeal to Lemma 13.6. In order to be allowed to do so, we need to verify two things. First, we need an estimate of the form (13.21) and second, we need an estimate of the variation of A in the interval of interest. To obtain such estimates is the purpose of the next lemma.

**Lemma 14.7.** Assume that (1.2) is oscillation adapted and let  $0 \neq \iota \in \mathcal{I}_{B}$ . Given that, in addition, the assumptions of Lemma 14.4 are satisfied, let  $t_{a}$ ,  $t_{b}$  and  $I_{a}$  be defined as in the statement of Lemma 14.4. Finally, let  $\tau_{a} := \tau(t_{a})$ ,  $\tau_{b} := \tau(t_{b})$  and

$$\varepsilon_a := \frac{\mathfrak{f}_{\mathrm{sh}}(t_a)}{\mathfrak{g}(t_a)} [1 + \mathfrak{f}_{\mathrm{ode}}(t_a) + \mathfrak{f}_X(t_a)] + \frac{1 + \mathfrak{f}_X^2(t_a) + \mathfrak{f}_{\mathrm{ode}}^2(t_a)}{\mathfrak{g}^2(t_a)}.$$
 (14.33)

Then if A is defined as in Section 14.2,

$$\|A\|_{C(J_a)} \cdot |J_a| \le C_{a,\mathfrak{b}} \varepsilon_a^{1/2} + C_{\mathfrak{b}} \mathfrak{f}_{\mathrm{sh}}(t_a) + C, \qquad (14.34)$$

where  $J_a$  is the interval in  $\tau$ -time corresponding to  $I_a$ ; C is a numerical constant;  $C_{\mathfrak{b}}$  only depends on an upper bound on  $\mathfrak{b}$ ; and  $C_{a,\mathfrak{b}}$  only depends on an upper bound on  $\varepsilon_a$  and  $\mathfrak{b}$ . Moreover,

$$\|A - A(\tau_a)\|_{C(J_a)} \cdot |J_a| \le C_{a,b}\varepsilon_a, \tag{14.35}$$

where  $C_{a,\mathfrak{b}}$  only depends on an upper bound on  $\varepsilon_a$  and the parameter  $\mathfrak{b}$  appearing in (14.1).

*Proof.* Let  $I_a$  and  $J_a$  be as in the statement of the lemma. We consider the different variables to be functions of t or functions of  $\tau$  as convenient. In particular, instead of computing the supremum of ||A|| over  $J_a$ , we can consider ||A|| to be a function of t and compute the supremum over  $I_a$ . In what follows, we wish to express all the estimates in terms of  $\mathfrak{g}(t_a)$ ,  $\mathfrak{f}_{sh}(t_a)$ ,  $\mathfrak{f}_X(t_a)$  and  $\mathfrak{f}_{ode}(t_a)$ . In order to be allowed to do so, we need to estimate the variation of  $\mathfrak{f}_{sh}$ ,  $\mathfrak{f}_X$  and  $\mathfrak{f}_{ode}$  in  $I_a$ . Due to (14.6), (14.22) and (14.24), it is clear that

$$\mathfrak{f}_{\mathrm{sh}}(t) \le e^{2\pi\mathfrak{b}}\mathfrak{f}_{\mathrm{sh}}(t_a), \quad \mathfrak{f}_X(t) \le e^{2\pi\mathfrak{b}}\mathfrak{f}_X(t_a), \quad \mathfrak{f}_{\mathrm{ode}}(t) \le e^{2\pi\mathfrak{b}}\mathfrak{f}_{\mathrm{ode}}(t_a)$$
(14.36)

for all  $t \in I_a$ .

Estimating the norm of A. Most of the functions of which A is constructed can immediately be estimated using (14.1)–(14.6). However, this is not true of ||U|| and  $||U^{-1}||$ . In order to estimate these objects, we appeal to Lemma 13.6 with M,  $s_a$  and J replaced by  $\alpha/2$ ,  $t_a$  and  $I_a$  respectively. Then

$$\frac{1}{2} \|\alpha\|_{C(I_a)} |I_a| \le 2\pi e^{2\pi \mathfrak{b}} \frac{\mathfrak{f}_{\text{ode}}(t_a)}{\mathfrak{g}(t_a)} \le 2\pi e^{2\pi \mathfrak{b}} \varepsilon_a^{1/2}, \tag{14.37}$$

where we have used (14.4), (14.24) and (14.36). Thus Lemma 13.6 applies with  $C_1$  replaced by the right hand side of (14.37), so that

$$\begin{aligned} \|\exp[\alpha(t_a)(\cdot - t_a)/2] - U(\cdot; t_a)\|_{C(I_a)} &\leq C \|\alpha - \alpha(t_a)\|_{C(I_a)} |I_a| \\ &\leq Ce^{2\pi\mathfrak{b}}\mathfrak{f}_{\text{ode}}(t_a)|I_a|^2 \leq C_{a,\mathfrak{b}}\frac{\mathfrak{f}_{\text{ode}}(t_a)}{\mathfrak{g}^2(t_a)} \end{aligned}$$
(14.38)

where the constant C only depends on an upper bound on  $C_1$  (and thus only on an upper bound on  $\mathfrak{b}$  and  $\varepsilon_a$ );  $C_{a,\mathfrak{b}}$  only depends on an upper bound on  $\mathfrak{b}$  and  $\varepsilon_a$ ; and we have used (14.4), (14.24) and (14.36). The estimate for  $U^{-1}$  is analogous. Note, in particular, that

$$\|U(\cdot;t_a) - \mathrm{Id}_m\|_{C(I_a)} + \|[U(\cdot;t_a)]^{-1} - \mathrm{Id}_m\|_{C(I_a)} \le C_{a,\mathfrak{b}} \frac{\mathfrak{f}_{\mathrm{ode}}(t_a)}{\mathfrak{g}(t_a)},$$
(14.39)

where the constant  $C_{a,b}$  only depends on an upper bound on  $\mathfrak{b}$  and  $\varepsilon_a$ .

Let us now estimate the norm of the components of A (after multiplication by  $|J_a|$ ). Note, to begin with, that  $|J_a| = 2\pi/\mathfrak{g}_0$ ; cf. (14.13). Thus

$$\|A_{11}(\tau)\| \cdot |J_a| \le \pi \left[\frac{\mathfrak{f}_{\text{ode}}(t)}{\mathfrak{g}(t)} + \frac{\mathfrak{b}}{\mathfrak{g}(t)}\right] + 2\pi \mathfrak{f}_{\text{sh}}(t) \le C_{\mathfrak{b}}[\varepsilon_a^{1/2} + \mathfrak{f}_{\text{sh}}(t_a)]$$
(14.40)

for  $\tau \in J_a$ , where  $C_{\mathfrak{b}}$  only depends on an upper bound on  $\mathfrak{b}$  and we have used (14.1), (14.2), (14.4), (14.23) and (14.36). Next,

$$\|A_{12}(\tau)\| \cdot |J_a| = 2\pi \tag{14.41}$$

for all  $\tau$ . Turning to  $A_{21}$ ,

$$\|A_{21}(\tau)\| \cdot |J_a| \le 2\pi \left[1 + C_{a,\mathfrak{b}}\left(\frac{\mathfrak{f}_X(t)}{\mathfrak{g}(t)} + \frac{\mathfrak{f}_{ode}^2(t)}{\mathfrak{g}^2(t)}\right)\right] \le 2\pi + C_{a,\mathfrak{b}}\varepsilon_a^{1/2} \tag{14.42}$$

for  $\tau \in J_a$ , where  $C_{a,b}$  only depends on an upper bound on  $\varepsilon_a$  and  $\mathfrak{b}$ , and we have used (14.3), (14.5), (14.23), (14.36) and (14.39). Before turning to  $A_{22}$ , we need to estimate the last term on the right hand side of (14.18). Estimate

$$\begin{split} & \left\| \frac{\mathfrak{g}_{0}}{\mathfrak{g}(t)} \left[ \alpha(t), U(t; t_{a}) \right] \left[ U(t; t_{a}) \right]^{-1} \right\| \\ \leq & \left\| \frac{\mathfrak{g}_{0}}{\mathfrak{g}(t)} \left[ \alpha(t), U(t; t_{a}) - \exp(\alpha(t_{a})(t - t_{a})/2) \right] \left[ U(t; t_{a}) \right]^{-1} \right\| \\ & + \left\| \frac{\mathfrak{g}_{0}}{\mathfrak{g}(t)} \left[ \alpha(t) - \alpha(t_{a}), \exp(\alpha(t_{a})(t - t_{a})/2) \right] \left[ U(t; t_{a}) \right]^{-1} \right\| . \end{split}$$

Appealing to (14.4), (14.23), (14.24), (14.36), (14.38) and (14.39) yields

.....

.

$$\left\|\frac{\mathfrak{g}_{0}}{\mathfrak{g}(t)}\left[\alpha(t), U(t; t_{a})\right]\left[U(t; t_{a})\right]^{-1}\right\| \leq C_{a, \mathfrak{b}}\mathfrak{g}_{0}\frac{\mathfrak{f}_{\mathrm{ode}}(t_{a})}{\mathfrak{g}^{2}(t_{a})} \leq C_{a, \mathfrak{b}}\mathfrak{g}_{0}\varepsilon_{a}$$
(14.43)

for all  $t \in I_a$ , where the constant  $C_{a,b}$  only depends on an upper bound on  $\mathfrak{b}$  and  $\varepsilon_a$ . Combining this estimate with (14.40) yields

$$\|A_{22}(\tau)\| \cdot |J_a| \le C_{a,\mathfrak{b}} \varepsilon_a^{1/2} + C_{\mathfrak{b}}\mathfrak{f}_{\mathrm{sh}}(t_a) \tag{14.44}$$

for  $\tau \in J_a$ , where  $C_{a,b}$  only depends on an upper bound on  $\mathfrak{b}$  and  $\varepsilon_a$ , and  $C_{\mathfrak{b}}$  only depends on an upper bound on  $\mathfrak{b}$ . Combining (14.40), (14.41), (14.42) and (14.44) yields (14.34).

The variation of A. Let us now turn to the variation of the  $A_{ij}$ . Since the  $A_{ij}$  are given by (14.15)-(14.18), we need to estimate the variation of

$$\frac{\alpha \mathfrak{g}_0}{\mathfrak{g}}, \quad \frac{\ell \mathfrak{g}_0}{\mathfrak{g}}, \quad \sigma \mathfrak{g}_0, \quad \frac{\mathfrak{g}_0}{\mathfrak{g}} U X U^{-1}, \quad \frac{\mathfrak{g}_0}{\mathfrak{g}} U Z U^{-1}, \quad \frac{\mathfrak{g}_0}{\mathfrak{g}} \left[ \alpha, U \right] U^{-1}$$

in the interval  $I_a$ . Estimate

$$\left\|\frac{\alpha(t)\mathfrak{g}_0}{\mathfrak{g}(t)} - \frac{\alpha(t_a)\mathfrak{g}_0}{\mathfrak{g}(t_a)}\right\| \le C_{\mathfrak{b}}\frac{\mathfrak{f}_{\text{ode}}(t_a)\mathfrak{g}_0}{\mathfrak{g}^2(t_a)} \le C_{\mathfrak{b}}\mathfrak{g}_0\varepsilon_a \tag{14.45}$$

for all  $t \in I_a$ , where we have used (14.4), (14.23), (14.24), (14.25) and (14.36) and the constant  $C_{\mathfrak{b}}$  only depends on an upper bound on the parameter  $\mathfrak{b}$  appearing in (14.1). Moreover,

...

$$\left\|\frac{\ell(t)\mathfrak{g}_0}{\mathfrak{g}(t)} - \frac{\ell(t_a)\mathfrak{g}_0}{\mathfrak{g}(t_a)}\right\| \le C_{\mathfrak{b}}\frac{\mathfrak{g}_0}{\mathfrak{g}^2(t_a)} \le C_{\mathfrak{b}}\mathfrak{g}_0\varepsilon_a \tag{14.46}$$

for all  $t \in I_a$ , where the constant  $C_{\mathfrak{b}}$  only depends on an upper bound on  $\mathfrak{b}$ , and we have used the estimate (14.26). Appealing to (14.2), (14.24) and (14.36),

$$\|\sigma(t)\mathfrak{g}_0 - \sigma(t_a)\mathfrak{g}_0\| \le C_{\mathfrak{b}}\mathfrak{g}_0\frac{\mathfrak{f}_{\mathrm{sh}}(t_a)}{\mathfrak{g}(t_a)} \le C_{\mathfrak{b}}\mathfrak{g}_0\varepsilon_a \tag{14.47}$$

for all  $t \in I_a$ , where the constant  $C_{\mathfrak{b}}$  only depends on an upper bound on  $\mathfrak{b}$ . Next, let us estimate

$$\left\|\frac{\mathfrak{g}_{0}}{\mathfrak{g}(t)}U(t;t_{a})X(t)[U(t;t_{a})]^{-1} - \frac{\mathfrak{g}_{0}}{\mathfrak{g}(t_{a})}X(t_{a})\right\|$$

$$\leq C_{a,\mathfrak{b}}\mathfrak{g}_{0}\frac{1+\mathfrak{f}_{ode}^{2}(t_{a}) + \mathfrak{f}_{X}^{2}(t_{a})}{\mathfrak{g}^{2}(t_{a})} \leq C_{a,\mathfrak{b}}\mathfrak{g}_{0}\varepsilon_{a}$$
(14.48)

for all  $t \in I_a$ , where the constant only depends on an upper bound on  $\mathfrak{b}$  and  $\varepsilon_a$  and we have used (14.3), (14.23), (14.24), (14.25), (14.36) and (14.39). Similarly,

$$\left\|\frac{\mathfrak{g}_0}{\mathfrak{g}(t)}U(t;t_a)Z(t)[U(t;t_a)]^{-1} - \frac{\mathfrak{g}_0}{\mathfrak{g}(t_a)}Z(t_a)\right\| \le C_{a,\mathfrak{b}}\mathfrak{g}_0\frac{\mathfrak{f}_{\text{ode}}^2(t_a)}{\mathfrak{g}^2(t_a)} \le C_{a,\mathfrak{b}}\mathfrak{g}_0\varepsilon_a \tag{14.49}$$

for all  $t \in I_a$ , where the constant only depends on an upper bound on  $\mathfrak{b}$  and  $\varepsilon_a$  and we have used (14.5), (14.23), (14.24), (14.25), (14.36) and (14.39). Combining (14.45), (14.46), (14.47), (14.48), (14.49) with (14.43) yields (14.35). The lemma follows.

#### 14.5 Computing the approximate fundamental solution

Due to Lemma 14.7, we have the estimates needed for appealing to Lemma 13.6. In particular,

$$\|\exp[A(\tau_a)(\cdot - \tau_a)] - \Phi(\cdot; \tau_a)\|_{C(J_a)} \le C_{a,\mathfrak{b},\mathrm{sh}}\varepsilon_a, \tag{14.50}$$

$$\|\exp[-A(\tau_a)(\cdot - \tau_a)] - [\Phi(\cdot; \tau_a)]^{-1}\|_{C(J_a)} \leq C_{a,\mathfrak{b},\mathrm{sh}}\varepsilon_a,$$
(14.51)

where  $J_a$ ,  $\varepsilon_a$  and  $\tau_a$  are defined as in the statement of Lemma 14.7; and the constant  $C_{a,\mathfrak{b},\mathrm{sh}}$  only depends on an upper bound on  $\varepsilon_a$ , the parameter  $\mathfrak{b}$  appearing in (14.1) and  $\mathfrak{f}_{\mathrm{sh}}(t_a)$ . Moreover,  $\Phi(\cdot;\tau_a)$  is the solution to the initial value problem

$$\frac{d\Phi}{d\tau}(\tau) = A(\tau)\Phi(\tau), \quad \Phi(\tau_a) = \mathrm{Id}_{2m}, \tag{14.52}$$

where A is the matrix defined in Section 14.2. Since the evolution of a solution to (14.20) over a period can be written

$$v(\tau_b) = \Phi(\tau_b; \tau_a) v(\tau_a) + \Phi(\tau_b; \tau_a) \int_{\tau_a}^{\tau_b} [\Phi(\tau; \tau_a)]^{-1} F(\tau) d\tau, \qquad (14.53)$$

where  $\tau_b$  is defined as in the statement of Lemma 14.7, we are mainly interested in approximating  $\Phi(\tau_b; \tau_a)$  and  $[\Phi(\tau'; \tau_a)]^{-1}$  for  $\tau' \in J_a$ . This is the purpose of the next lemma.

**Lemma 14.8.** Assume that (1.2) is oscillation adapted and let  $0 \neq \iota \in \mathcal{I}_{B}$ . Given that, in addition, the assumptions of Lemma 14.4 are satisfied, let  $t_{a}$ ,  $t_{b}$ ,  $\tau_{a}$ ,  $\tau_{b}$ ,  $I_{a}$  and  $J_{a}$  be defined as in the statement of Lemma 14.7. Let A and  $\tau$  be defined as in Section 14.2 and let  $\Phi(\tau; \tau_{a})$  be the solution to (14.52). Then

$$\|\Phi_{\mathrm{app}}(\tau_b;\tau_a) - \Phi(\tau_b;\tau_a)\| + \|\Phi_{\mathrm{app}}^{\mathrm{I}}(\cdot;\tau_a) - [\Phi(\cdot;\tau_a)]^{-1}\|_{C(J_a)} \le C_{a,\mathfrak{b},\mathrm{sh}}\varepsilon_a,$$
(14.54)

where the constant  $C_{a,\mathfrak{b},\mathrm{sh}}$  only depends on an upper bound on  $\varepsilon_a$  defined in (14.33); the parameter  $\mathfrak{b}$  appearing in (14.1); and  $\mathfrak{f}_{\mathrm{sh}}(t_a)$ . In this estimate,

$$\Phi_{\rm app}(\tau_b;\tau_a) := \cos(\omega) \operatorname{Id}_{2m} + \operatorname{sgn}(t_b - t_a) \cdot \nu^{-1} \sin(\omega) \begin{pmatrix} i\xi \operatorname{Id}_m & \operatorname{Id}_m \\ -\operatorname{Id}_m & -i\xi \operatorname{Id}_m \end{pmatrix} + \operatorname{sgn}(t_b - t_a) \cdot \frac{i\pi}{\mathfrak{g}(t_a)} \begin{pmatrix} 0 & X(t_a) \\ -X(t_a) & 0 \end{pmatrix}$$
(14.55)

where  $\xi := -\sigma(t_a)$ ;  $\nu := (1 + \xi^2)^{1/2}$ ; and  $\omega := 2\pi\nu$ . Moreover,

$$\Phi_{\rm app}^{\rm I}(\tau;\tau_a) := \cos(s\omega) \operatorname{Id}_{2m} + \nu^{-1} \sin(s\omega) \begin{pmatrix} i\xi \operatorname{Id}_m & \operatorname{Id}_m \\ -\operatorname{Id}_m & -i\xi \operatorname{Id}_m \end{pmatrix} + \pi s \sin(s\omega) \begin{pmatrix} R_{21} & 0 \\ 0 & R_{21} \end{pmatrix} + \frac{1}{2} \sin(s\omega) \begin{pmatrix} 2R_{11} & R_{21} \\ R_{21} & -2R_{11} \end{pmatrix}$$
(14.56)  
$$- \pi s \cos(s\omega) \begin{pmatrix} 0 & R_{21} \\ -R_{21} & 0 \end{pmatrix},$$

where  $\xi$ ,  $\nu$  and  $\omega$  are defined as above,  $R_{11}$  and  $R_{21}$  are defined in (14.58) below, and

$$s := -\frac{\mathfrak{g}_0}{2\pi}(\tau - \tau_a), \tag{14.57}$$

where  $\mathfrak{g}_0 := \mathfrak{g}(t_0)$  and  $t_0$  is introduced in connection with the definition of  $\tau$ ; cf. (14.13).

**Remark 14.9.** When convenient, we write

$$\Phi_{\rm app} = \Phi^0_{\rm app} + \Phi^1_{\rm app}, \quad \Phi^{\rm I}_{\rm app} = \Phi^{\rm I,0}_{\rm app} + \Phi^{\rm I,1}_{\rm app}$$

where  $\Phi_{app}^{0}$  and  $\Phi_{app}^{I,0}$  are the sum of the first two terms on the right hand sides of (14.55) and (14.56) respectively.

**Remark 14.10.** It is important to keep in mind that, regardless of the choice of  $s, \xi \in \mathbb{R}$ , the matrix  $\Phi_{\text{app}}^{\text{I},0}$  is an isometry of  $\mathbb{C}^{2m}$  (with respect to the standard inner product); i.e.,  $\Phi_{\text{app}}^{\text{I},0}$  is unitary.

*Proof.* Due to (14.50) and (14.51), it is sufficient to calculate  $\exp[A(\tau_a)T_0s]$ , for  $s \in [-1, 1]$ , where  $T_0 := 2\pi/\mathfrak{g}_0$ . Returning to Section 13.2, in particular Lemma 13.3, we are thus interested in calculating  $\exp(2\pi sP)$ , where  $P = A(\tau_a)/\mathfrak{g}_0$ . Note that the P defined in this way is also given by (13.4), (13.5) and (13.6), where  $E_{11}$ ,  $E_{12}$  and  $E_{22}$  are defined to be zero, and

$$\xi := -\sigma(t_a), \quad R_{11} := \frac{1}{2} \frac{\alpha(t_a)}{\mathfrak{g}(t_a)} + \frac{1}{2} \frac{\ell(t_a)}{\mathfrak{g}(t_a)} \mathrm{Id}_m, \quad R_{21} := -i \frac{X(t_a)}{\mathfrak{g}(t_a)}, \quad E_{21} := -\frac{Z(t_a)}{\mathfrak{g}(t_a)}.$$
(14.58)

The next step is to apply Lemma 13.3. To be allowed to do so, we need to verify that the bound (13.7) holds. Estimate

$$|\xi| + ||R_{11}|| + ||R_{21}|| + \sum_{i,j} ||E_{ij}|| \le \mathfrak{f}_{\mathrm{sh}}(t_a) + C_{a,\mathfrak{b}}\varepsilon_a^{1/2}, \qquad (14.59)$$

where the constant  $C_{a,b}$  only depends on an upper bound on  $\mathfrak{b}$  and  $\varepsilon_a$ . Thus Lemma 13.3 is applicable with  $C_1$  replaced by the right hand side of (14.59). It is of interest to express  $\mathcal{E}$ , defined in (13.8), in terms of  $\mathfrak{g}(t_a)$  etc. Note, to this end, that in our situation

$$\|R_{11}\|^{2} \leq \frac{1}{2} \frac{\mathfrak{f}_{\text{ode}}^{2}(t_{a}) + \mathfrak{b}^{2}}{\mathfrak{g}^{2}(t_{a})}, \quad \|R_{21}\|^{2} \leq \frac{\mathfrak{f}_{X}^{2}(t_{a})}{\mathfrak{g}^{2}(t_{a})}, \quad \|E_{21}\| \leq \frac{\mathfrak{f}_{\text{ode}}^{2}(t_{a})}{\mathfrak{g}^{2}(t_{a})}.$$

Moreover,

$$|\xi|(||R_{11}|| + ||R_{21}||) \le \frac{\mathfrak{f}_{\mathrm{sh}}(t_a)}{\mathfrak{g}(t_a)}[\mathfrak{b} + \mathfrak{f}_{\mathrm{ode}}(t_a) + \mathfrak{f}_X(t_a)].$$

Thus  $\mathcal{E} \leq C_{\mathfrak{b}}\varepsilon_a$  for some constant  $C_{\mathfrak{b}}$  only depending on an upper bound on  $\mathfrak{b}$ . Summing up the above observations, we conclude that (13.9) and (13.10) are applicable, and that the constant C appearing in these estimates only depends on an upper bound on  $\varepsilon_a$ , the parameter  $\mathfrak{b}$  appearing in (14.1) and  $\mathfrak{f}_{\mathrm{sh}}(t_a)$ . Moreover,  $\mathcal{E}$  can be replaced by  $\varepsilon_a$ . Thus (14.54) holds, where  $\Phi_{\mathrm{app}}$  and  $\Phi_{\mathrm{app}}^{\mathrm{I}}$  are given by (14.55) and (14.56) respectively. Moreover, the constant  $C_{a,\mathfrak{b},\mathrm{sh}}$  has the properties stated in the lemma.

### Chapter 15

## One period of the oscillations; reformulation and detailed estimates

The variables introduced in Section 14.2 are adapted to one period of the oscillations. However, we are interested in drawing conclusions concerning the evolution over longer periods of time. This requires, first of all, a change of variables, and it turns out to be convenient to work with w introduced in (15.1) below. Using (14.53), we can derive an expression for how w evolves over one period of the oscillations; cf. (15.3) below. The main point of the initial part of the present chapter is to estimate the matrices appearing in this expression; in Section 15.1 we derive a rough, and in Section 15.2 a detailed, estimate. With these estimates as a starting point, we are in a position to set up an iteration; cf. Section 15.3. The purpose of the iteration is to make it possible to estimate how w develops over longer periods of time. As a starting point, it is natural to consider how  $w(t_k)$  develops, where  $\{t_k\}$  is the time sequence associated with the iteration. However, in order to simplify the formulae as much as possible, it is convenient to modify  $w(t_k)$  somewhat. To this end, we introduce  $w_{\text{pre},k}$  and  $w_{\text{fin},k}$  in (15.36) and (15.53) respectively. The norms of these variables are equivalent to that of  $w(t_k)$ . In Lemmas 15.7 and 15.11, we derive formulae for how  $w_{\text{pre},k}$  and  $w_{\text{fin},k}$  evolve. In the end, we obtain (15.63) and (15.64), which are the basis for the analysis to follow.

#### 15.1 Rough estimate of the evolution during one period

The motivation for considering the function v introduced in (14.9) is the desire to simplify the analysis of how the solution evolves over one period. By a period, we here mean an interval  $I_a$  of the form described in Lemma 14.4. In order to consider the evolution of the solution over longer time intervals, we, however, need to consider other functions. In what follows, we focus our attention on w, defined by

$$w(t) := \begin{pmatrix} x_1(t) \\ y_1(t) \end{pmatrix} = \exp\left[\frac{1}{2} \int_{t_a}^t \dot{\ell}(t') dt'\right] \begin{pmatrix} [U(t;t_a)]^{-1} & 0 \\ 0 & [U(t;t_a)]^{-1} \end{pmatrix} v(t).$$
(15.1)

Here the first equality is a definition and the second is a consequence of (14.8). Due to (15.1) it is perhaps not surprising that the following matrix will play an important role in the analysis:

$$\Xi(\tau;\tau_a) = \exp\left[\frac{1}{2}\int_{t_a}^t \dot{\ell}(t')dt'\right] \begin{pmatrix} [U(t;t_a)]^{-1} & 0\\ 0 & [U(t;t_a)]^{-1} \end{pmatrix} \Phi(\tau;\tau_a),$$
(15.2)

where  $\Phi(\tau; \tau_a)$  is the solution to (14.52); in this formula, we take it for granted that t and  $\tau$  are related according to  $\tau(t) = \tau$ . Since (14.53) holds with  $\tau_b$  replaced by  $\tau$ ,

$$w(t) = \Xi(\tau; \tau_a) w(t_a) + \Xi(\tau; \tau_a) \int_{\tau_a}^{\tau} [\Phi(\tau'; \tau_a)]^{-1} F(\tau') d\tau',$$
(15.3)

where we have used the fact that  $w(t_a) = v(\tau_a)$ . Here, and in the formulae below,  $\tau_a$ ,  $\tau_b$  and  $J_a$  are defined as in the statement of Lemma 14.7. We are mainly interested in calculating the right hand side of (15.3) when  $\tau = \tau_b$ . However, in some situations, it is of interest to obtain a rough approximation of the right hand side when  $\tau \in J_a$ . For that reason, we here derive a rough estimate for  $\Xi(\tau; \tau_a)$ .

**Lemma 15.1.** Assume that (1.2) is oscillation adapted and let  $0 \neq \iota \in \mathcal{I}_{B}$ . Given that, in addition, the assumptions of Lemma 14.4 are satisfied, let  $t_{a}$ ,  $t_{b}$ ,  $\tau_{a}$ ,  $\tau_{b}$ ,  $I_{a}$  and  $J_{a}$  be defined as in the statement of Lemma 14.7. Let A and  $\tau$  be defined as in Section 14.2 and let  $\Phi(\tau; \tau_{a})$  be the solution to (14.52). Finally, let  $\Xi(\tau; \tau_{a})$  be defined by (15.2). Then, for  $\tau \in J_{a}$ ,

$$\begin{aligned} &\|\Xi(\tau;\tau_a) - \mathfrak{D}_a(t)\| + \left\| [\Xi(\tau;\tau_a)]^{-1} - [\mathfrak{D}_a(t)]^T \right\| \\ \leq C_{a,\mathfrak{b},\mathrm{sh}} \frac{1 + \mathfrak{f}_{\mathrm{ode}}(t_a) + \mathfrak{f}_X(t_a)}{\mathfrak{g}(t_a)} + C_{a,\mathfrak{b},\mathrm{sh}}\varepsilon_a + C_{a,\mathfrak{b},\mathrm{sh}}\mathfrak{f}_{\mathrm{sh}}(t_a), \end{aligned}$$
(15.4)

where  $C_{a,\mathfrak{b},\mathrm{sh}}$  only depends on an upper bound on  $\varepsilon_a$  defined in (14.33); the parameter  $\mathfrak{b}$  appearing in (14.1); and  $\mathfrak{f}_{\mathrm{sh}}(t_a)$ . Moreover,  $\mathfrak{D}_a(t)$  is defined by

$$\varphi_a(t) := \int_{t_a}^t \mathfrak{g}(t')dt', \quad \mathfrak{D}_a(t) := \begin{pmatrix} \cos\varphi_a(t)\mathrm{Id}_m & \sin\varphi_a(t)\mathrm{Id}_m \\ -\sin\varphi_a(t)\mathrm{Id}_m & \cos\varphi_a(t)\mathrm{Id}_m \end{pmatrix}$$
(15.5)

and  $[\mathfrak{D}_a(t)]^T$  denotes the transpose of the matrix  $\mathfrak{D}_a(t)$ .

**Remark 15.2.** In (15.4) we take it for granted that t and  $\tau$  are related according to  $\tau(t) = \tau$ , where  $\tau(t)$  is defined in (14.13).

**Remark 15.3.** Due to (15.5), it is clear that the leading order behaviour of  $\Xi(\tau; \tau_a)$  over one period of the oscillations is just a rotation.

*Proof.* Consider  $\Xi(\tau; \tau_a)$ , defined by (15.2). The first factor on the right hand side of (15.2) is given by

$$\exp\left[\frac{1}{2}\int_{t_a}^t \dot{\ell}(t')dt'\right] = \frac{\mathfrak{g}^{1/2}(t)}{\mathfrak{g}^{1/2}(t_a)}$$

Appealing to (14.23) and (14.25) thus yields

$$\left| \exp\left[\frac{1}{2} \int_{t_a}^t \dot{\ell}(t') dt'\right] - 1 \right| \le \frac{C_{\mathfrak{b}}}{\mathfrak{g}(t_a)}$$
(15.6)

for  $t \in I_a$ , where  $C_{\mathfrak{b}}$  only depends on the parameter  $\mathfrak{b}$  appearing in (14.1). Turning to  $[U(t;t_a)]^{-1}$ , note that (14.39) holds. Combining this estimate with (15.6) yields

$$\left\| \exp\left[\frac{1}{2} \int_{t_a}^t \dot{\ell}(t') dt'\right] \begin{pmatrix} [U(t;t_a)]^{-1} & 0\\ 0 & [U(t;t_a)]^{-1} \end{pmatrix} - \mathrm{Id}_{2m} \right\| \le \frac{C_{\mathfrak{b}}}{\mathfrak{g}(t_a)} + C_{a,\mathfrak{b}} \frac{\mathfrak{f}_{\mathrm{ode}}(t_a)}{\mathfrak{g}(t_a)}$$
(15.7)

for  $t \in I_a$ , where the constant  $C_{a,b}$  only depends on an upper bound on  $\mathfrak{b}$  and  $\varepsilon_a$ ; here  $\varepsilon_a$  is given by (14.33). Similarly,

$$\left\| \exp\left[ -\frac{1}{2} \int_{t_a}^t \dot{\ell}(t') dt' \right] \begin{pmatrix} U(t; t_a) & 0\\ 0 & U(t; t_a) \end{pmatrix} - \mathrm{Id}_{2m} \right\| \le \frac{C_{\mathfrak{b}}}{\mathfrak{g}(t_a)} + C_{a,\mathfrak{b}} \frac{\mathfrak{f}_{\mathrm{ode}}(t_a)}{\mathfrak{g}(t_a)}.$$
(15.8)

In the present setting, we consider the right hand side of (15.7) to be an error term. As a consequence, the first two factors appearing in the definition of  $\Xi(\tau;\tau_a)$  can be replaced by  $\mathrm{Id}_{2m}$ . Next, let us approximate  $\Phi(\tau;\tau_a)$ . In order to do so, it is convenient to return to Section 14.5, in particular (14.50) and the proof of Lemma 14.8. Due to the proof of Lemma 14.8, we know that (13.9) holds with  $P = A(\tau_a)/\mathfrak{g}_0$ . Moreover, the constant C appearing in this estimate only depends on an upper bound on  $\varepsilon_a$ , the parameter  $\mathfrak{b}$  appearing in (14.1) and  $\mathfrak{f}_{\mathrm{sh}}(t_a)$ . In addition,  $\mathcal{E}$  can be replaced by  $\varepsilon_a$ . Finally, since  $R_{11}$  and  $R_{21}$  are given by (14.58), we know that

$$||R_{11}|| + ||R_{21}|| \le \frac{\mathfrak{b} + \mathfrak{f}_{\text{ode}}(t_a) + \mathfrak{f}_X(t_a)}{\mathfrak{g}(t_a)}$$

Appealing to (13.9) with P replaced by  $A(\tau_a)/\mathfrak{g}_0$  yields

$$\left\| \exp(2\pi sP) - \cos(s\omega_a) \operatorname{Id}_{2m} - \nu_a^{-1} \sin(s\omega_a) \begin{pmatrix} i\xi_a \operatorname{Id}_m & \operatorname{Id}_m \\ -\operatorname{Id}_m & -i\xi_a \operatorname{Id}_m \end{pmatrix} \right\|$$

$$\leq C_{\mathfrak{b}} \frac{1 + \mathfrak{f}_{\operatorname{ode}}(t_a) + \mathfrak{f}_X(t_a)}{\mathfrak{g}(t_a)} + C_{a,\mathfrak{b},\operatorname{sh}}\varepsilon_a,$$

$$(15.9)$$

where  $\xi_a = -\sigma(t_a)$ ;  $\nu_a = (1 + \xi_a^2)^{1/2}$ ;  $\omega_a = 2\pi\nu_a$ ;  $C_{\mathfrak{b}}$  only depends on  $\mathfrak{b}$ ; and  $C_{a,\mathfrak{b},\mathrm{sh}}$  only depends on an upper bound on  $\varepsilon_a$ , the parameter  $\mathfrak{b}$  appearing in (14.1), and  $\mathfrak{f}_{\mathrm{sh}}(t_a)$ . In (15.9), s takes its values in [-1, 1], so that

$$|\cos(s\omega_a) - \cos(2\pi s)| \le |s| \cdot |\omega_a - 2\pi| \le 2\pi |\nu_a - 1| \le \pi \mathfrak{f}_{\mathrm{sh}}^2(t_a).$$

Similarly,

$$|\sin(s\omega_a) - \sin(2\pi s)| + |\nu_a^{-1} - 1| \le (\pi + 1)\mathfrak{f}_{\mathrm{sh}}^2(t_a).$$

Finally,  $|\xi_a| \leq \mathfrak{f}_{\rm sh}(t_a)$ , so that (15.9) yields

$$\left\| \exp(2\pi sP) - \cos(2\pi s) \operatorname{Id}_{2m} - \sin(2\pi s) \begin{pmatrix} 0 & \operatorname{Id}_m \\ -\operatorname{Id}_m & 0 \end{pmatrix} \right\|$$
  
$$\leq C_{\mathfrak{b}} \frac{1 + \mathfrak{f}_{\text{ode}}(t_a) + \mathfrak{f}_X(t_a)}{\mathfrak{g}(t_a)} + C_{a,\mathfrak{b},\text{sh}}\varepsilon_a + C_{\text{sh}}\mathfrak{f}_{\text{sh}}(t_a),$$
(15.10)

where  $C_{\mathfrak{b}}$  and  $C_{a,\mathfrak{b},\mathrm{sh}}$  have the same dependence as before and  $C_{\mathrm{sh}}$  only depends on an upper bound on  $\mathfrak{f}_{\mathrm{sh}}(t_a)$ . In this estimate,  $s = s_{\pm}$ , where

$$2\pi s_{\pm} = \pm 2\pi \frac{\tau - \tau_a}{T_0} = \pm \mathfrak{g}_0 \int_{t_a}^t \frac{\mathfrak{g}(t')}{\mathfrak{g}_0} dt' = \pm \int_{t_a}^t \mathfrak{g}(t') dt', \qquad (15.11)$$

and  $T_0 = 2\pi/\mathfrak{g}_0$ . Here the plus sign is the relevant sign when estimating  $\Xi(\tau; \tau_a)$  and the minus sign is the relevant sign when estimating  $[\Xi(\tau; \tau_a)]^{-1}$ . Define  $\varphi_a$  and  $\mathfrak{D}_a$  by (15.5). Then (15.10) yields

$$\|\exp(2\pi s_+P) - \mathfrak{D}_a(t)\| \le C_{\mathfrak{b}} \frac{1 + \mathfrak{f}_{\text{ode}}(t_a) + \mathfrak{f}_X(t_a)}{\mathfrak{g}(t_a)} + C_{a,\mathfrak{b},\text{sh}}\varepsilon_a + C_{\text{sh}}\mathfrak{f}_{\text{sh}}(t_a)$$
(15.12)

for  $t \in I_a$ . An analogous estimate holds for  $s_-$ . Combining these estimates with (14.50) and (14.51) yields

$$\begin{split} & \left\| \Phi(\tau;\tau_a) - \mathfrak{D}_a(t) \right\| + \left\| [\Phi(\tau;\tau_a)]^{-1} - [\mathfrak{D}_a(t)]^T \right\| \\ \leq C_{\mathfrak{b}} \frac{1 + \mathfrak{f}_{\text{ode}}(t_a) + \mathfrak{f}_X(t_a)}{\mathfrak{g}(t_a)} + C_{a,\mathfrak{b},\text{sh}}\varepsilon_a + C_{\text{sh}}\mathfrak{f}_{\text{sh}}(t_a) \end{split}$$
(15.13)

for  $t \in I_a$ , where  $\tau$  and t are related according to (14.13); the constants have the same dependence as before; and  $[\mathfrak{D}_a(t)]^T$  denotes the transpose of the matrix  $\mathfrak{D}_a(t)$ . Finally, combining (15.7), (15.8) and (15.13) yields (15.4). The lemma follows.

#### 15.1.1 Estimating solutions to inhomogeneous equations

For future reference, it is of interest to make the following observation concerning the problem of estimating solutions to inhomogeneous equations. To begin with, (24.16), (14.7) and (15.1) yield

$$\dot{w}(\iota, t) = \mathcal{A}(\iota, t)w(\iota, t) + \mathcal{F}(\iota, t), \qquad (15.14)$$

where

$$\begin{aligned} \mathcal{A}(\iota,t) &:= \begin{pmatrix} \dot{\ell}(\iota,t) \mathrm{Id}_m - i\sigma(\iota,t) \mathfrak{g}(\iota,t) \mathrm{Id}_m & \mathfrak{g}(\iota,t) \mathrm{Id}_m \\ -\mathfrak{g}(\iota,t) \mathrm{Id}_m - iX(\iota,t) - [\mathfrak{g}(\iota,t)]^{-1} \zeta(t) & i\sigma(\iota,t) \mathfrak{g}(\iota,t) \mathrm{Id}_m - \alpha(t) \end{pmatrix}, \\ \mathcal{F}(\iota,t) &:= \exp\left(-i \int_0^t \sigma(\iota,t') \mathfrak{g}(\iota,t') dt'\right) \begin{pmatrix} 0 \\ \hat{f}(\iota,t) \end{pmatrix}. \end{aligned}$$

Given  $t', t \in I$ , let  $\Phi_{\iota}(t; t')$  be the solution to the initial value problem

$$\frac{d\Phi}{dt} = \mathcal{A}\Phi, \quad \Phi(t') = \mathrm{Id}_{2m},$$

evaluated at t. Then solutions to (15.14) satisfy

$$w(\iota, t) = \Phi_{\iota}(t; t_a)w(\iota, t_a) + \int_{t_a}^{t} \Phi_{\iota}(t; t')\mathcal{F}(\iota, t')dt'.$$
(15.15)

In order to estimate  $|w(\iota, t)|$  on, say,  $[t_a, t_b]$  in terms of  $|w(\iota, t_a)|$  and an integral involving  $|\hat{f}|$ , it is thus sufficient to estimate  $||\Phi_{\iota}(t; t')||$  for  $t_a \leq t' \leq t \leq t_b$ . There is a similar statement if one wants the estimate to be in terms of  $|w(\iota, t_b)|$  and an integral involving  $|\hat{f}|$ .

#### 15.2 First detailed estimate of the evolution over one period

In the previous section, we derived a rough estimate for  $\Xi(\tau; \tau_a)$ . However, in order to obtain information concerning the long time behaviour of solutions, we need to derive detailed estimates. The purpose of the present section is to take a first step in that direction.

**Lemma 15.4.** Assume that (1.2) is oscillation adapted and let  $0 \neq \iota \in \mathcal{I}_{B}$ . Given that, in addition, the assumptions of Lemma 14.4 are satisfied, let  $t_{a}$ ,  $t_{b}$ ,  $\tau_{a}$ ,  $\tau_{b}$ ,  $I_{a}$  and  $J_{a}$  be defined as in the statement of Lemma 14.7. Let A and  $\tau$  be defined as in Section 14.2 and let  $\Xi(\tau;\tau_{a})$  be defined by (15.2). Then

$$w(t_b) = \Xi(\tau_b; \tau_a) w(t_a) + \Xi(\tau_b; \tau_a) \int_{t_a}^{t_b} [\Xi(\tau(t); \tau_a)]^{-1} \exp\left(-i \int_0^t \sigma(t') \mathfrak{g}(t') dt'\right) \begin{pmatrix} 0\\ \hat{f}(t) \end{pmatrix} dt,$$
(15.16)

where w is defined in (15.1) and z is a solution to (24.16). Moreover,

$$\|\Xi(\tau_b;\tau_a) - \Xi_{\mathrm{app}}(\tau_b;\tau_a)\| + \|[\Xi(\cdot;\tau_a)]^{-1} - \Xi_{\mathrm{app}}^{\mathrm{I}}(\cdot;\tau_a)\|_{C(J_a)} \le C_{a,\mathfrak{b},\mathrm{sh}}\varepsilon_a,$$
(15.17)

where the constant  $C_{a,\mathfrak{b},\mathrm{sh}}$  only depends on an upper bound on  $\varepsilon_a$ ; the parameter  $\mathfrak{b}$  appearing in (14.1); and  $\mathfrak{f}_{\mathrm{sh}}(t_a)$ . In (15.17),  $\Xi_{\mathrm{app}}$  is given by

$$\Xi_{\rm app}(\tau_b;\tau_a) = \cos(\omega) \operatorname{Id}_{2m} + \operatorname{sgn}(t_b - t_a) \cdot \nu^{-1} \sin(\omega) \begin{pmatrix} i\xi \operatorname{Id}_m & \operatorname{Id}_m \\ -\operatorname{Id}_m & -i\xi \operatorname{Id}_m \end{pmatrix} + \operatorname{sgn}(t_b - t_a) \cdot \frac{\pi}{\mathfrak{g}(t_a)} \begin{pmatrix} \dot{\ell}(t_a) \operatorname{Id}_m - \alpha(t_a) & iX(t_a) \\ -iX(t_a) & \dot{\ell}(t_a) \operatorname{Id}_m - \alpha(t_a) \end{pmatrix},$$
(15.18)

where  $\xi = -\sigma(t_a)$ ,  $\nu = (1 + \xi^2)^{1/2}$  and  $\omega = 2\pi\nu$ . Finally, the  $\Xi^{I}_{app}(\tau; \tau_a)$  appearing in (15.17) is given by

$$\Xi_{\mathrm{app}}^{\mathrm{I}}(\tau;\tau_{a}) = \cos(s\omega)\mathrm{Id}_{2m} + \nu^{-1}\sin(s\omega) \begin{pmatrix} i\xi\mathrm{Id}_{m} & \mathrm{Id}_{m} \\ -\mathrm{Id}_{m} & -i\xi\mathrm{Id}_{m} \end{pmatrix} + \pi s\sin(s\omega) \begin{pmatrix} R_{21} & R_{\mathrm{aux}} \\ -R_{\mathrm{aux}} & R_{21} \end{pmatrix} + \frac{1}{2}\sin(s\omega) \begin{pmatrix} 2R_{11} & R_{21} \\ R_{21} & -2R_{11} \end{pmatrix} (15.19) + \pi s\cos(s\omega) \begin{pmatrix} R_{\mathrm{aux}} & -R_{21} \\ R_{21} & R_{\mathrm{aux}} \end{pmatrix},$$

where  $R_{11}$  and  $R_{21}$  are defined in (14.58);  $R_{aux}$  is defined in (15.31) below; and s is defined by (14.57).

*Proof.* Let us start by deriving (15.16). Note that (15.3) yields

$$w(t_b) = \Xi(\tau_b; \tau_a) w(t_a) + \Xi(\tau_b; \tau_a) \int_{\tau_a}^{\tau_b} [\Phi(\tau; \tau_a)]^{-1} F(\tau) d\tau.$$
(15.20)

Let us consider the second factor in the last term on the right hand side. It can be written

$$\int_{\tau_{a}}^{\tau_{b}} [\Phi(\tau;\tau_{a})]^{-1} F(\tau) d\tau = \int_{t_{a}}^{t_{b}} [\Phi(\tau(t);\tau_{a})]^{-1} F[\tau(t)] \frac{\mathfrak{g}(t)}{\mathfrak{g}_{0}} dt \\
= \int_{t_{a}}^{t_{b}} [\Phi(\tau(t);\tau_{a})]^{-1} \begin{pmatrix} 0 \\ F_{2}(t) \end{pmatrix} dt \\
= \int_{t_{a}}^{t_{b}} [\Xi(\tau(t);\tau_{a})]^{-1} \exp\left(-i \int_{0}^{t} \sigma(t') \mathfrak{g}(t') dt'\right) \begin{pmatrix} 0 \\ \hat{f}(t) \end{pmatrix} dt.$$
(15.21)

Combining (15.20) and (15.21) yields (15.16).

**Calculating**  $\Xi_{app}(\tau_b; \tau_a)$ . The next step is to calculate  $\Xi_{app}(\tau_b; \tau_a)$ . Considering (15.2), it is clear that we need to estimate  $[U(t_b; t_a)]^{-1}$ . By an estimate similar to (14.38),

$$\|\exp[-\alpha(t_a)(t_b - t_a)/2] - [U(t_b; t_a)]^{-1}\| \le C_{a, \mathfrak{b}} \frac{\mathfrak{f}_{ode}(t_a)}{\mathfrak{g}^2(t_a)},$$
(15.22)

where  $C_{a,b}$  only depends on an upper bound on  $\mathfrak{b}$  and  $\varepsilon_a$ . In particular,

$$\left\| [U(t_b; t_a)]^{-1} - \mathrm{Id}_m + \mathrm{sgn}(t_b - t_a) \cdot \frac{\pi \alpha(t_a)}{\mathfrak{g}(t_a)} \right\| \le C_{a, \mathfrak{b}} \frac{\mathfrak{f}_{\mathrm{ode}}^2(t_a) + 1}{\mathfrak{g}^2(t_a)} \le C_{a, \mathfrak{b}} \varepsilon_a, \tag{15.23}$$

where  $C_{a,b}$  only depends on an upper bound on  $\mathfrak{b}$  and  $\varepsilon_a$ , and we have used (14.27) and (14.37). Appealing to (14.28),

$$\left| \exp\left[\frac{1}{2} \int_{t_a}^{t_b} \dot{\ell}(t') dt'\right] - 1 - \operatorname{sgn}(t_b - t_a) \cdot \frac{\pi \dot{\ell}(t_a)}{\mathfrak{g}(t_a)} \right| \le \frac{C_{\mathfrak{b}}}{\mathfrak{g}^2(t_a)} \le C_{\mathfrak{b}}\varepsilon_a, \tag{15.24}$$

where the constant only depends on an upper bound on  $\mathfrak{b}$ . Combining (15.23) and (15.24) with (14.54), (14.55) and (15.2) yields

$$\|\Xi(\tau_b;\tau_a) - \Xi_{\rm app}(\tau_b;\tau_a)\| \le C_{a,\mathfrak{b},{\rm sh}}\varepsilon_a,$$

where  $\Xi_{app}(\tau_b; \tau_a)$  is given by (15.18) and and the constant  $C_{a,\mathfrak{b},sh}$  only depends on an upper bound on  $\varepsilon_a$ ; the parameter  $\mathfrak{b}$  appearing in (14.1); and  $\mathfrak{f}_{sh}(t_a)$ . Note, however, that in order to obtain this estimate, we have used the fact that

$$|\sin(2\pi\nu)| = |\sin[2\pi(\nu-1)]| \le C\mathfrak{f}_{\rm sh}^2(t_a), \quad |\cos(2\pi\nu) - 1| \le C\mathfrak{f}_{\rm sh}^4(t_a),$$

where C is a numerical constant; cf. (13.16).

**Calculating**  $\Xi_{app}^{I}(\tau;\tau_a)$ . Next we need to calculate  $\Xi_{app}^{I}(\tau;\tau_a)$ . Note to this end that (14.38) yields

$$\|\exp[\alpha(t_a)(t-t_a)/2] - U(t;t_a)\| \le C_{a,\mathfrak{b}} \frac{\mathfrak{f}_{ode}(t_a)}{\mathfrak{g}^2(t_a)}$$
(15.25)

for  $t \in I_a$ , where  $C_{a,\mathfrak{b}}$  only depends on an upper bound on  $\mathfrak{b}$  and  $\varepsilon_a$ . Thus

$$\left\| U(t;t_a) - \mathrm{Id}_m - \frac{1}{2}\alpha(t_a)(t-t_a) \right\| \le C_{a,\mathfrak{b}} \frac{\mathfrak{f}_{\mathrm{ode}}^2(t_a) + 1}{\mathfrak{g}^2(t_a)} \le C_{a,\mathfrak{b}}\varepsilon_a$$
(15.26)

for  $t \in I_a$ , where  $C_{a,b}$  only depends on an upper bound on  $\mathfrak{b}$  and  $\varepsilon_a$ . By an argument which is essentially identical to the first step of the proof of (14.28),

$$\left|\int_{t_a}^t \dot{\ell}(t')dt' - \dot{\ell}(t_a)(t - t_a)\right| \le \frac{C_{\mathfrak{b}}}{\mathfrak{g}^2(t_a)} \tag{15.27}$$

for  $t \in I_a$ , where  $C_{\mathfrak{b}}$  only depends on an upper bound on  $\mathfrak{b}$ . It is of interest to express the  $t - t_a$  appearing in this formula in terms of the *s* defined by (14.57). Note, to this end, that (14.13) and (14.57) yield

$$s = -\frac{1}{2\pi} \int_{t_a}^t \mathfrak{g}(t') dt' = -\frac{1}{2\pi} \int_{t_a}^t \mathfrak{g}(t') \mathfrak{g}(t_a) \left(\frac{1}{\mathfrak{g}(t_a)} - \frac{1}{\mathfrak{g}(t')}\right) dt' - \frac{1}{2\pi} \mathfrak{g}(t_a)(t - t_a).$$

Using the results of Lemma 14.4 to estimate the first term on the right hand side, it can then be verified that

$$\left| t - t_a + \frac{2\pi s}{\mathfrak{g}(t_a)} \right| \le \frac{C_{\mathfrak{b}}}{\mathfrak{g}^2(t_a)}$$

for  $t \in I_a$ , where  $C_{\mathfrak{b}}$  only depends on an upper bound on  $\mathfrak{b}$ . Combining this estimate with (15.26) and (15.27) yields

$$\left\| \exp\left[ -\frac{1}{2} \int_{t_a}^t \dot{\ell}(t') dt' \right] U(t; t_a) - \mathrm{Id}_m - \frac{\pi s \dot{\ell}(t_a)}{\mathfrak{g}(t_a)} \mathrm{Id}_m + \frac{\pi s}{\mathfrak{g}(t_a)} \alpha(t_a) \right\| \le C_{a, \mathfrak{b}} \varepsilon_a \tag{15.28}$$

for  $t \in I_a$ , where  $C_{a,\mathfrak{b}}$  only depends on an upper bound on  $\mathfrak{b}$  and  $\varepsilon_a$ . On the other hand, (15.2) implies that

$$[\Xi(\tau;\tau_a)]^{-1} = [\Phi(\tau;\tau_a)]^{-1} \exp\left[-\frac{1}{2}\int_{t_a}^t \dot{\ell}(t')dt'\right] \begin{pmatrix} U(t;t_a) \\ 0 & U(t;t_a) \end{pmatrix}$$

Combining this formula with (14.54) and (15.28) yields

$$\|\Xi_{\text{inter}}^{\text{I}}(\tau;\tau_a) - [\Xi(\tau;\tau_a)]^{-1}\| \le C_{a,\mathfrak{b},\text{sh}}\varepsilon_a$$
(15.29)

for  $\tau \in J_a$ , where the constant  $C_{a,\mathfrak{b},\mathrm{sh}}$  only depends on an upper bound on  $\mathfrak{b}$ ,  $\varepsilon_a$  and  $\mathfrak{f}_{\mathrm{sh}}(t_a)$ . Moreover,

$$\Xi_{\text{inter}}^{\text{I}}(\tau;\tau_{a}) := \Phi_{\text{app}}^{\text{I}}(\tau;\tau_{a}) \left[ \text{Id}_{2m} + \pi s \begin{pmatrix} R_{\text{aux}} & 0\\ 0 & R_{\text{aux}} \end{pmatrix} \right]$$

$$= \Phi_{\text{app}}^{\text{I}}(\tau;\tau_{a}) + \pi s \begin{pmatrix} \cos(s\omega)R_{\text{aux}} & \sin(s\omega)R_{\text{aux}}\\ -\sin(s\omega)R_{\text{aux}} & \cos(s\omega)R_{\text{aux}} \end{pmatrix} + \Xi_{\text{rem}}^{\text{I}}(\tau;\tau_{a})$$
(15.30)

for  $\tau \in J_a$ , where

$$R_{\text{aux}} := \frac{1}{\mathfrak{g}(t_a)} \left[ \dot{\ell}(t_a) \mathrm{Id}_m - \alpha(t_a) \right]$$
(15.31)

and

$$\|\Xi_{\rm rem}^{\rm I}(\tau;\tau_a)\| \le C_{a,\mathfrak{b},\rm sh}\varepsilon_a \tag{15.32}$$

for  $\tau \in J_a$ , where the constant  $C_{a,\mathfrak{b},\mathrm{sh}}$  only depends on an upper bound on  $\mathfrak{b}$ ,  $\varepsilon_a$  and  $\mathfrak{f}_{\mathrm{sh}}(t_a)$ . In order to justify the second equality in (15.30), note that

$$\left\| \Phi_{\mathrm{app}}^{\mathrm{I},1}(\tau;\tau_a) \begin{pmatrix} R_{\mathrm{aux}} & 0\\ 0 & R_{\mathrm{aux}} \end{pmatrix} \right\| \le C \|R_{\mathrm{aux}}\|(\|R_{11}\| + \|R_{21}\|) \le C_{\mathfrak{b}}\varepsilon_a$$

for  $\tau \in J_a$ , where the constant C is numerical and the constant  $C_{\mathfrak{b}}$  only depends on an upper bound on  $\mathfrak{b}$ . Moreover,

$$|\nu^{-1} - 1| \|R_{\mathrm{aux}}\| + |\xi| \|R_{\mathrm{aux}}\| \le C_{\mathfrak{b}} \varepsilon_{\mathfrak{a}}$$

for  $\tau \in J_a$ , where the constant  $C_{\mathfrak{b}}$  only depends on an upper bound on  $\mathfrak{b}$ .

Combining (15.29), (15.30) and (15.32) with (14.56) yields

$$\|[\Xi(\cdot;\tau_a)]^{-1} - \Xi^{\mathrm{I}}_{\mathrm{app}}(\cdot;\tau_a)\|_{C(J_a)} \le C_{a,\mathfrak{b},\mathrm{sh}}\varepsilon_a,$$

where  $\Xi_{app}^{1}(\tau;\tau_{a})$  is given by (15.19) and the constant  $C_{a,\mathfrak{b},\mathrm{sh}}$  only depends on an upper bound on  $\varepsilon_{a}$ ; the parameter  $\mathfrak{b}$  appearing in (14.1); and  $\mathfrak{f}_{\mathrm{sh}}(t_{a})$ .

#### 15.3 Iteration

Previous sections contain a description of how solutions evolve over one period of the oscillations; cf., in particular, Lemma 15.4. In what follows, we wish to iterate the corresponding estimate. To begin with, it is therefore natural to define a time sequence. Since we are interested in going both forward and backward in time, we in fact define two sequences.

**Definition 15.5.** Assume that (1.2) is oscillation adapted and let  $0 \neq \iota \in \mathcal{I}_B$ . Assume, moreover,  $t_0 \in [0, \infty)$  to be such that (14.22) holds with  $t_a$  replaced by  $t_0$ . Assume that  $t_k \geq t_0$  has been defined for some  $0 \leq k \in \mathbb{Z}$  and that (14.22) holds with  $t_a$  replaced by  $t_k$ . Then  $t_{k+1}$  is defined by the condition that

$$\int_{t_k}^{t_{k+1}} \mathfrak{g}(\iota, t) dt = 2\pi.$$
(15.33)

If there is a final k such that (14.22) holds with  $t_a$  replaced by  $t_k$ , then this k is denoted by  $k_{\text{fin}}$ . Moreover,  $t_{\text{fin}} := t_{k_{\text{fin}}}$ . If there is no such final k, then  $k_{\text{fin}} := \infty$  and  $t_{\text{fin}} := \infty$ . If, in addition to the above,  $t_0 \ge 2\pi$ , then  $t_{-1}$  is defined by the condition that (15.33) holds with k = -1. Assume  $t_k \le t_0$  has been defined for some  $0 \ge k \in \mathbb{Z}$ ; that (14.22) holds with  $t_a$  replaced by  $t_k$ ; and that  $t_k \ge 2\pi$ . Then  $t_{k-1}$  is defined by the condition that (15.33) holds with k replaced by k-1. Let  $k_{\text{ini}}$ be the first  $0 \ge k \in \mathbb{Z}$  such that either  $0 \le t_k < 2\pi$  or  $\mathfrak{g}(t_k) < \max\{4\pi\mathfrak{b}, 2\}$ . Finally,  $t_{\text{ini}} := t_{k_{\text{ini}}}$ .

**Remark 15.6.** That the definition of  $t_k$  for  $k_{ini} \le k \le k_{fin}$  makes sense is ensured by Lemma 14.4.

In order to obtain a clearer picture of the development of the solution, it is convenient to introduce the transformation

$$T_{\text{pre},k} := \frac{1}{2} \begin{pmatrix} (\nu_k - \xi_k) \text{Id}_m & i \text{Id}_m \\ i(\nu_k + \xi_k) \text{Id}_m & \text{Id}_m \end{pmatrix},$$
(15.34)

$$T_{\text{pre},k}^{-1} = \frac{1}{\nu_k} \begin{pmatrix} \text{Id}_m & -i\text{Id}_m \\ -i(\xi_k + \nu_k)\text{Id}_m & (\nu_k - \xi_k)\text{Id}_m \end{pmatrix}.$$
 (15.35)

Here and below we use the notation  $\xi_k := -\sigma(t_k)$ ,  $\nu_k := (1 + \xi_k^2)^{1/2}$  and  $\omega_k := 2\pi\nu_k$ . Note that if  $\mathfrak{f}_{sh}(t)$  is bounded by a constant independent of  $0 \le t \in \mathbb{R}$ , then the norms of both of the matrices

appearing in (15.34) and (15.35) are bounded by constants independent of k and  $0 \neq \iota \in \mathcal{I}_B$ . It turns out that the analysis is simplified by considering the variable

$$w_{\mathrm{pre},k} := T_{\mathrm{pre},k} w(t_k) \tag{15.36}$$

instead of  $w(t_k)$ .

**Lemma 15.7.** Assume that (1.2) is oscillation adapted and let  $0 \neq \iota \in \mathcal{I}_{B}$ . Assume  $t_{0} \in [0, \infty)$  to be such that (14.22) holds with  $t_{a}$  replaced by  $t_{0}$ , and define  $t_{k}$  and  $w_{\text{pre},k}$  as in Definition 15.5 and (15.36) respectively. Then there is a matrix  $\Xi_{\text{pre}}^{k,+}$  for  $0 \leq k < k_{\text{fin}}$  and, in case  $t_{0} \geq 2\pi$ , a matrix  $\Xi_{\text{pre}}^{k,-}$  for  $k_{\text{ini}} < k \leq 0$ . Moreover, there is a matrix valued function  $\Xi_{\text{pre}}(\cdot;\tau_{k})$  such that

$$w_{\text{pre},k\pm 1} = \Xi_{\text{pre}}^{k,\pm} w_{\text{pre},k} + \Xi_{\text{pre}}^{k,\pm} \int_{t_k}^{t_{k\pm 1}} [\Xi_{\text{pre}}(\tau(t);\tau_k)]^{-1} \exp\left(-i\int_0^t \sigma(t')\mathfrak{g}(t')dt'\right) T_{\text{pre},k} \begin{pmatrix} 0\\ \hat{f}(t) \end{pmatrix} dt$$
(15.37)

where  $0 \le k < k_{\text{fin}}$  for the equality with a plus sign and  $k_{\text{ini}} < k \le 0$  for the equality with a minus sign (in the case of a minus sign, it is also assumed that  $t_0 \ge 2\pi$ ). Moreover, there is a matrix  $\Xi_{\text{pre,app}}^{k,\pm}$  and a matrix valued function  $\Xi_{\text{pre,app}}^{\text{I}}(\cdot;\tau_k)$  such that

$$\|\Xi_{\text{pre}}^{k,\pm} - \Xi_{\text{pre,app}}^{k,\pm}\| + \|[\Xi_{\text{pre}}(\cdot;\tau_k)]^{-1} - \Xi_{\text{pre,app}}^{\text{I}}(\cdot;\tau_k)\|_{C(J_{k,\pm})} \le C_{\varepsilon,\mathfrak{b},\mathfrak{sh}}\varepsilon_k,$$
(15.38)

where  $J_{k,\pm}$  is the interval in  $\tau$ -time corresponding to the interval with endpoints  $t_k$  and  $t_{k\pm 1}$ ; and the constant  $C_{\varepsilon,\mathfrak{b},\mathrm{sh}}$  only depends on an upper bound on  $\mathfrak{b}$ ,  $\varepsilon_k$  and  $\mathfrak{f}_{\mathrm{sh}}(t_k)$  (here  $\varepsilon_k$  is defined to be the right hand side of (14.33) with  $t_a$  replaced by  $t_k$ ). Finally, in (15.38),  $\Xi_{\mathrm{pre,app}}^{k,\pm}$  is defined by

$$\Xi_{\text{pre,app}}^{k,\pm} := \begin{pmatrix} e^{\mp i\omega_k} \text{Id}_m & 0\\ 0 & e^{\pm i\omega_k} \text{Id}_m \end{pmatrix} \pm \pi \begin{pmatrix} R_{\text{pre},k}^{10} & 0\\ 0 & R_{\text{pre},k}^{11} \end{pmatrix}$$
(15.39)

and  $\Xi^{I}_{pre,app}$  is defined by

$$\Xi_{\text{pre,app}}^{\text{I}}(\tau;\tau_k) := \begin{pmatrix} e^{-is\omega_k} \text{Id}_m & 0\\ 0 & e^{is\omega_k} \text{Id}_m \end{pmatrix} + \pi s \begin{pmatrix} e^{-is\omega_k} R_{\text{pre},k}^{10} & 0\\ 0 & e^{is\omega_k} R_{\text{pre},k}^{11} \end{pmatrix} - \frac{i}{2} \sin(s\omega_k) \begin{pmatrix} 0 & R_{\text{pre},k}^{00}\\ -R_{\text{pre},k}^{01} & 0 \end{pmatrix},$$
(15.40)

where s is defined in (14.57) (with  $\tau_a$  replaced by  $\tau_k$ ) and

$$R_{\text{pre},k}^{ab} := \frac{1}{\mathfrak{g}(t_k)} [(-1)^a \alpha(t_k) + \dot{\ell}(t_k) \text{Id}_m + (-1)^b X(t_k)]$$
(15.41)

for a, b = 0, 1.

**Remark 15.8.** All the equalities and the estimates in the lemma which refer to the intervals  $[t_{k-1}, t_k], k_{\text{ini}} < k \leq 0$ , are based on the assumption that  $t_0 \geq 2\pi$ .

**Remark 15.9.** The  $t_0$  appearing in (14.13) can be assumed to equal the  $t_0$  appearing in the assumptions of the lemma.

**Remark 15.10.** In case  $k_{\text{fin}} \ge 1$ ,  $\Xi_{\text{pre}}^{k,-}$  can be defined for finite k such that  $1 \le k \le k_{\text{fin}}$ . Similarly, in case  $k_{\text{ini}} \le -1$ ,  $\Xi_{\text{pre}}^{k,+}$  can be defined for  $k_{\text{ini}} + 1 \le k \le 0$ . The remaining statements of the lemma are then valid for the corresponding k and  $\mp$ .

*Proof.* The main purpose of introducing  $T_{\text{pre},k}$  is that it has the property that

$$T_{\text{pre},k} \begin{bmatrix} \cos(s\omega_k) \text{Id}_{2m} + \nu_k^{-1} \sin(s\omega_k) \begin{pmatrix} i\xi_k \text{Id}_m & \text{Id}_m \\ -\text{Id}_m & -i\xi_k \text{Id}_m \end{pmatrix} \end{bmatrix} T_{\text{pre},k}^{-1}$$

$$= \begin{pmatrix} e^{-is\omega_k} \text{Id}_m & 0 \\ 0 & e^{is\omega_k} \text{Id}_m \end{pmatrix}.$$
(15.42)

#### 15.3. ITERATION

Note also that

$$\left\| T_{\operatorname{pre},k} - \frac{1}{2} \begin{pmatrix} \operatorname{Id}_m & i \operatorname{Id}_m \\ i \operatorname{Id}_m & \operatorname{Id}_m \end{pmatrix} \right\| + \left\| T_{\operatorname{pre},k}^{-1} - \begin{pmatrix} \operatorname{Id}_m & -i \operatorname{Id}_m \\ -i \operatorname{Id}_m & \operatorname{Id}_m \end{pmatrix} \right\|$$

$$\leq C(|\nu_k - 1| + |\xi_k|) \leq C \mathfrak{f}_{\operatorname{sh}}(t_k),$$

$$(15.43)$$

where the constant C is numerical. Thus

$$\left\| \frac{\pi}{\mathfrak{g}(t_k)} T_{\operatorname{pre},k} \begin{pmatrix} \dot{\ell}(t_k) \operatorname{Id}_m - \alpha(t_k) & iX(t_k) \\ -iX(t_k) & \dot{\ell}(t_k) \operatorname{Id}_m - \alpha(t_k) \end{pmatrix} T_{\operatorname{pre},k}^{-1} \\ - \frac{\pi}{\mathfrak{g}(t_k)} \begin{pmatrix} \dot{\ell}(t_k) \operatorname{Id}_m - \alpha(t_k) + X(t_k) & 0 \\ 0 & \dot{\ell}(t_k) \operatorname{Id}_m - \alpha(t_k) - X(t_k) \end{pmatrix} \right\| \leq C_{\mathfrak{b}, \mathrm{sh}} \varepsilon_k,$$

$$(15.44)$$

where  $\varepsilon_k$  is defined by (14.33) with  $t_a$  replaced by  $t_k$  and  $C_{b,sh}$  only depends on an upper bound on  $\mathfrak{b}$  and  $\mathfrak{f}_{sh}(t_k)$ . The relations (15.42) and (15.44) demonstrate that  $T_{\text{pre},k}$  diagonalises the matrices of interest. However, there is one problem. In practice, the first factor on the left hand side of (15.42) should be  $T_{\text{pre},k\pm 1}$ , not  $T_{\text{pre},k}$ . For that reason, it is of interest to estimate  $T_{\text{pre},k\pm 1}T_{\text{pre},k}^{-1} - \text{Id}_{2m}$ . Note that

$$T_{\mathrm{pre},k\pm 1}T_{\mathrm{pre},k}^{-1} - \mathrm{Id}_{2m} = \frac{1}{2\nu_k} \begin{pmatrix} \mathfrak{q}_{k,\pm}\mathrm{Id}_m & -i\mathfrak{q}_{k,\pm}\mathrm{Id}_m \\ i\mathfrak{r}_{k,\pm}\mathrm{Id}_m & \mathfrak{r}_{k,\pm}\mathrm{Id}_m \end{pmatrix},$$
(15.45)

where

$$\mathfrak{q}_{k,\pm} = \nu_{k\pm 1} - \nu_k + \xi_k - \xi_{k\pm 1}, \quad \mathfrak{r}_{k,\pm} = \nu_{k\pm 1} - \nu_k + \xi_{k\pm 1} - \xi_k$$

It is thus sufficient to estimate  $|\nu_{k\pm 1} - \nu_k|$  and  $|\xi_{k\pm 1} - \xi_k|$ . In fact, since

$$|\nu_{k\pm 1} - \nu_k| = \frac{|\nu_{k\pm 1}^2 - \nu_k^2|}{\nu_{k\pm 1} + \nu_k} = \frac{|\xi_{k\pm 1} - \xi_k| |\xi_{k\pm 1} + \xi_k|}{\nu_{k\pm 1} + \nu_k} \le |\xi_{k\pm 1} - \xi_k|,$$
(15.46)

it is sufficient to estimate  $|\xi_{k\pm 1} - \xi_k|$ . On the other hand

$$|\xi_{k\pm 1} - \xi_k| = \left| \int_{t_k}^{t_{k\pm 1}} \dot{\sigma}(t') dt' \right| \le e^{2\pi\mathfrak{b}} \mathfrak{f}_{\mathrm{sh}}(t_k) |t_{k\pm 1} - t_k| \le 4\pi e^{2\pi\mathfrak{b}} \frac{\mathfrak{f}_{\mathrm{sh}}(t_k)}{\mathfrak{g}(t_k)},\tag{15.47}$$

where we have used (14.2), (14.24) and (14.36); note that the last two estimates are applicable in the present context with  $t_a$  replaced by  $t_k$ . Thus

$$\|T_{\operatorname{pre},k\pm 1}T_{\operatorname{pre},k}^{-1} - \operatorname{Id}_{2m}\| \le C_{\mathfrak{b}}\frac{\mathfrak{f}_{\operatorname{sh}}(t_k)}{\mathfrak{g}(t_k)},\tag{15.48}$$

where  $C_{\mathfrak{b}}$  is a constant only depending on an upper bound on  $\mathfrak{b}$ .

Next we wish to compute the relation between  $w_{\text{pre},k}$  and  $w_{\text{pre},k\pm 1}$ . Note, to this end, that

$$w_{\mathrm{pre},k\pm 1} = \Xi_{\mathrm{pre}}^{k,\pm} w_{\mathrm{pre},k} + \Xi_{\mathrm{pre}}^{k,\pm} \int_{t_k}^{t_{k\pm 1}} [\Xi_{\mathrm{pre}}(\tau(t);\tau_k)]^{-1} \exp\left(-i\int_0^t \sigma(t')\mathfrak{g}(t')dt'\right) T_{\mathrm{pre},k} \begin{pmatrix} 0\\\hat{f}(t) \end{pmatrix} dt,$$

where we have appealed to (15.16) and

$$\Xi_{\mathrm{pre}}^{k,\pm} := T_{\mathrm{pre},k\pm 1} \Xi(\tau_{k\pm 1};\tau_k) T_{\mathrm{pre},k}^{-1}, \quad \Xi_{\mathrm{pre}}(\tau;\tau_k) := T_{\mathrm{pre},k} \Xi(\tau;\tau_k) T_{\mathrm{pre},k}^{-1}$$

To begin with, we wish to compute an approximation for the matrix  $\Xi_{\text{pre}}^{k,\pm}$ . Due to (15.17), (15.18), (15.43) and (15.48),

$$\|\Xi_{\mathrm{pre}}^{k,\pm} - T_{\mathrm{pre},k}\Xi_{\mathrm{app}}(\tau_{k\pm 1};\tau_k)T_{\mathrm{pre},k}^{-1}\| \le C_{\varepsilon,\mathfrak{b},\mathrm{sh}}\varepsilon_k,$$

where the constant  $C_{\varepsilon,\mathfrak{b},\mathrm{sh}}$  only depends on an upper bound on  $\mathfrak{b}$ ,  $\varepsilon_k$  and  $\mathfrak{f}_{\mathrm{sh}}(t_k)$ . Combining this estimate with (15.42) and (15.44) yields

$$\|\Xi_{\rm pre}^{k,\pm} - \Xi_{\rm pre,app}^{k,\pm}\| \le C_{\varepsilon,\mathfrak{b},\rm sh}\varepsilon_k,\tag{15.49}$$

where the constant  $C_{\varepsilon,\mathfrak{b},\mathrm{sh}}$  only depends on an upper bound on  $\mathfrak{b}$ ,  $\varepsilon_k$  and  $\mathfrak{f}_{\mathrm{sh}}(t_k)$ . In (15.49),  $\Xi_{\mathrm{pre,app}}^{k,\pm}$  is defined by (15.39). Similarly,

$$\|[\Xi_{\rm pre}(\cdot;\tau_k)]^{-1} - \Xi^{\rm I}_{\rm pre,app}(\cdot;\tau_k)\| \le C_{\varepsilon,\mathfrak{b},\rm sh}\varepsilon_k,\tag{15.50}$$

where the constant  $C_{\varepsilon,\mathfrak{b},\mathrm{sh}}$  only depends on an upper bound on  $\mathfrak{b}$ ,  $\varepsilon_k$  and  $\mathfrak{f}_{\mathrm{sh}}(t_k)$ . In (15.50),  $\Xi^{\mathrm{I}}_{\mathrm{pre,app}}$  is defined by (15.40). The lemma follows.

Let

$$\mathcal{D}_k(\iota) := \exp\left[i\int_0^{t_k} [1+\sigma^2(\iota,t)]^{1/2}\mathfrak{g}(\iota,t)dt\right],\tag{15.51}$$

$$D_k(\iota) := \begin{pmatrix} \mathcal{D}_k(\iota) \mathrm{Id}_m & 0\\ 0 & [\mathcal{D}_k(\iota)]^{-1} \mathrm{Id}_m \end{pmatrix},$$
(15.52)

$$w_{\mathrm{fin},k} := D_k w_{\mathrm{pre},k}.\tag{15.53}$$

With these definitions, we obtain the following simplification.

**Lemma 15.11.** Assume that (1.2) is oscillation adapted and let  $0 \neq \iota \in \mathcal{I}_{\mathrm{B}}$ . Assume  $t_0 \in [0, \infty)$  to be such that (14.22) holds with  $t_a$  replaced by  $t_0$ , and define  $t_k$  and  $w_{\mathrm{fin},k}$  as in Definition 15.5 and (15.53) respectively. Then there is a matrix  $\Xi_{\mathrm{fin}}^{k,+}$  for  $0 \leq k < k_{\mathrm{fin}}$  and, in case  $t_0 \geq 2\pi$ , a matrix  $\Xi_{\mathrm{fin}}^{k,-}$  for  $k_{\mathrm{ini}} < k \leq 0$ . Moreover, there is a matrix valued function  $\Xi_{\mathrm{fin}}(\cdot;\tau_k)$  such that

$$w_{\text{fin},k\pm 1} = \Xi_{\text{fin}}^{k,\pm} w_{\text{fin},k} + \Xi_{\text{fin}}^{k,\pm} \int_{t_k}^{t_{k\pm 1}} \hat{F}_k(t) dt, \qquad (15.54)$$

where  $0 \le k < k_{\text{fin}}$  for the equality with a plus sign;  $k_{\text{ini}} < k \le 0$  for the equality with a minus sign (in the case of a minus sign, it is also assumed that  $t_0 \ge 2\pi$ );

$$\hat{F}_k(t) := [\Xi_{\text{fin}}(\tau(t);\tau_k)]^{-1} \exp\left(-i \int_0^t \sigma(t')\mathfrak{g}(t')dt'\right) T_{\text{fin},k} \begin{pmatrix} 0\\ \hat{f}(t) \end{pmatrix};$$
(15.55)

and  $T_{\text{fin},k} := D_k T_{\text{pre},k}$ . Moreover, there is a matrix  $\Xi_{\text{fin},\text{app}}^{k,\pm}$  and a matrix valued function  $\Xi_{\text{fin},\text{app}}^{\text{I}}(\cdot;\tau_k)$  such that

$$\|\Xi_{\text{fin}}^{k,\pm} - \Xi_{\text{fin},\text{app}}^{k,\pm}\| + \|[\Xi_{\text{fin}}(\cdot;\tau_k)]^{-1} - \Xi_{\text{fin},\text{app}}^{\text{I}}(\cdot;\tau_k)\|_{C(J_{k,\pm})} \le C_{\varepsilon,\mathfrak{b},\text{sh}}\varepsilon_k,$$
(15.56)

where  $J_{k,\pm}$  is the interval in  $\tau$ -time corresponding to the interval with endpoints  $t_k$  and  $t_{k\pm 1}$ ; and the constant  $C_{\varepsilon,\mathfrak{b},\mathrm{sh}}$  only depends on an upper bound on  $\mathfrak{b}$ ,  $\varepsilon_k$  and  $\mathfrak{f}_{\mathrm{sh}}(t_k)$  (here  $\varepsilon_k$  is defined to be the right hand side of (14.33) with  $t_a$  replaced by  $t_k$ ). Finally, in (15.56),  $\Xi_{\mathrm{fin,app}}^{k,\pm}$  is defined by

$$\Xi_{\rm fin,app}^{k,\pm} := \mathrm{Id}_{2m} \pm \pi \left( \begin{array}{cc} R_{\rm pre,k}^{10} & 0\\ 0 & R_{\rm pre,k}^{11} \end{array} \right), \tag{15.57}$$

where  $R^{ab}_{\text{pre},k}$  is defined in (15.41). Moreover,  $\Xi^{\text{I}}_{\text{fin,app}}$  is defined by

$$\Xi_{\mathrm{fin,app}}^{\mathrm{I}}(\tau;\tau_k) := \begin{pmatrix} e^{-is\omega_k} \mathrm{Id}_m & 0\\ 0 & e^{is\omega_k} \mathrm{Id}_m \end{pmatrix} + \pi s \begin{pmatrix} e^{-is\omega_k} R_{\mathrm{pre},k}^{10} & 0\\ 0 & e^{is\omega_k} R_{\mathrm{pre},k}^{11} \end{pmatrix} - \frac{i}{2} \sin(s\omega_k) \begin{pmatrix} 0 & \mathcal{D}_k^2 R_{\mathrm{pre},k}^{00}\\ -\mathcal{D}_k^{-2} R_{\mathrm{pre},k}^{01} & 0 \end{pmatrix},$$
(15.58)

where s is defined in (14.57) (with  $\tau_a$  replaced by  $\tau_k$ ),  $\mathcal{D}_k$  is defined in (15.51) and  $R_{\text{pre},k}^{ab}$  is defined in (15.41).

**Remark 15.12.** It is of interest to note that  $\Xi_{\text{fin,app}}^{k,\pm}$  is independent of the functions  $g^{0l}$  and  $\zeta$  appearing in the equation (1.2).

206

#### 15.3. ITERATION

Remark 15.13. Remark 15.8 is equally relevant in the present setting.

**Remark 15.14.** In case  $k_{\text{fin}} \ge 1$ ,  $\Xi_{\text{fin}}^{k,-}$  can be defined for finite k such that  $1 \le k \le k_{\text{fin}}$ . Similarly, in case  $k_{\text{ini}} \le -1$ ,  $\Xi_{\text{fin}}^{k,+}$  can be defined for  $k_{\text{ini}} + 1 \le k \le 0$ . The remaining statements of the lemma are then valid for the corresponding k and  $\mp$ .

*Proof.* Note that  $D_k$ , defined in (15.52), and its inverse both have norm 1. Note also that

$$\pm \omega_k = \pm 2\pi [1 + \sigma^2(t_k)]^{1/2} = \int_{t_k}^{t_{k\pm 1}} [1 + \sigma^2(t_k)]^{1/2} \mathfrak{g}(t) dt;$$

recall that (14.21) holds with  $t_a$  and  $t_b$  replaced by  $t_k$  and  $t_{k\pm 1}$  respectively. Thus

$$\mathcal{D}_{k\pm 1}\mathcal{D}_{k}^{-1} = \exp\left[i\int_{t_{k}}^{t_{k\pm 1}} [1+\sigma^{2}(t)]^{1/2}\mathfrak{g}(t)dt\right]$$

$$=e^{\pm i\omega_{k}}\exp\left[i\int_{t_{k}}^{t_{k\pm 1}} \left([1+\sigma^{2}(t)]^{1/2} - [1+\sigma^{2}(t_{k})]^{1/2}\right)\mathfrak{g}(t)dt\right].$$
(15.59)

However, by estimates similar to (15.46) and (15.47),

$$\left| [1 + \sigma^2(t)]^{1/2} - [1 + \sigma^2(t_k)]^{1/2} \right| \le C_{\mathfrak{b}} \frac{\mathfrak{f}_{\mathrm{sh}}(t_k)}{\mathfrak{g}(t_k)}$$

for  $t \in I_{k,\pm}$ , where  $I_{k,\pm}$  is the interval with endpoints  $t_k$  and  $t_{k\pm 1}$ , and the constant  $C_{\mathfrak{b}}$  only depends on an upper bound on  $\mathfrak{b}$ . Combining this estimate with (15.59) yields

$$\left|\mathcal{D}_{k\pm 1}\mathcal{D}_{k}^{-1} - e^{\pm i\omega_{k}}\right| + \left|\mathcal{D}_{k\pm 1}^{-1}\mathcal{D}_{k} - e^{\mp i\omega_{k}}\right| \le C_{\mathfrak{b}}\frac{\mathfrak{f}_{\mathrm{sh}}(t_{k})}{\mathfrak{g}(t_{k})},\tag{15.60}$$

where the constant  $C_{\mathfrak{b}}$  only depends on an upper bound on  $\mathfrak{b}$ . Similarly,

$$|\exp(\pm i\omega_k) - 1| = |\exp(\pm 2\pi i(\nu_k - 1)) - 1| \le 2\pi |\nu_k - 1| \le 2\pi \mathfrak{f}_{\rm sh}(t_k).$$
(15.61)

Defining

$$\Xi_{\text{fin}}^{k,\pm} := D_{k\pm 1} \Xi_{\text{pre}}^{k,\pm} D_k^{-1}, \quad \Xi_{\text{fin}}(\tau;\tau_k) := D_k \Xi_{\text{pre}}(\tau;\tau_k) D_k^{-1}, \quad T_{\text{fin},k} := D_k T_{\text{pre},k}$$
(15.62)

yields (15.54). Combining these definitions with Lemma 15.7, (15.60) and (15.61) yields the conclusions of the lemma.  $\hfill \Box$ 

Since we appeal to (15.54) repeatedly in applications, it is convenient to simplify the notation. We therefore write (15.54) as

$$\psi_{k\pm 1} = A_k^{\pm} \psi_k + A_k^{\pm} \int_{t_k}^{t_{k\pm 1}} \hat{F}_k(t) dt, \qquad (15.63)$$

where  $\psi_k := w_{\text{fin},k}$  and  $A_k^{\pm} := \Xi_{\text{fin}}^{k,\pm}$ . Applying this equality iteratively yields, e.g.,

$$\psi_{k+1} = A_k^+ \cdots A_0^+ \psi_0 + A_k^+ \int_{t_k}^{t_{k+1}} \hat{F}_k(t) dt + \cdots + A_k^+ \cdots A_0^+ \int_{t_0}^{t_1} \hat{F}_0(t) dt.$$
(15.64)

Due to equalities of this type, it is clearly of interest to calculate matrix products such as  $A_k^+ \cdots A_{k-l}^+$ . This is an important part of the analysis in the next two parts of these notes.
# Part IV

# Unbalanced equations

# Chapter 16

# Unbalanced equations

# 16.1 Introduction

Consider the requirement (10.2) for an equation to be weakly balanced. There are various ways of violating this condition. However, we here focus on two situations. First, there is the possibility that the norms of  $\alpha(t)$  and  $\zeta(t)$  are unbounded. Then pathological behaviour might arise even on the level of spatially homogeneous solutions. Another option is that the norm of  $X(\iota, t)$  is unbounded. In that case, we clearly need to consider spatially inhomogeneous solutions in order to see something interesting. Finally, there is the possibility that  $\sigma(\iota, t)$  is not bounded. This is not a situation we consider in greater detail here. One reason for this is that the main motivation for studying (1.2) is that it arises when linearising Einstein's equations around certain cosmological solutions. In that context, the size of  $\sigma(\iota, t)$  can be influenced by the choice of gauge. In fact, we expect it to be possible to ensure that  $\sigma(\iota, t)$  is small by an appropriate choice of gauge. For this reason, we do not consider the case that  $\sigma(\iota, t)$  is large here.

In order to be able to obtain results, we need to assume more than that, say, the norms of  $\alpha(t)$  and  $\zeta(t)$  are unbounded. In fact, if these quantities are unbounded and strongly oscillatory, it can be expected to be hard to draw conclusions. In the present chapter we therefore consider situations in which at least one of the following two assumptions holds:

- $e^{-\beta_{\alpha}t}\alpha(t)$  and  $e^{-2\beta_{\zeta}t}\zeta(t)$  converge exponentially for some real numbers  $\beta_{\alpha}$  and  $\beta_{\zeta}$ , where at least one of  $\beta_{\alpha}$  and  $\beta_{\zeta}$  is strictly positive and the corresponding limit is non-zero,
- $e^{-\beta_X t} X^j(t) / [g^{jj}(t)]^{1/2}$  converges to a matrix  $Y_{\infty}^j$  for some  $j \in \{1, \ldots, d\}$  and some  $\beta_X > 0$ , where  $Y_{\infty}^j$  has an eigenvalue with non-zero real part.

In practice, we need to make additional assumptions, but these requirements are the essential ones. Below, we informally refer to the first situation as the ODE-unbalanced setting, and the second situation as the PDE-unbalanced setting.

The ODE-unbalanced setting. Dealing with the ODE-unbalanced setting is fairly straightforward, and we record the corresponding results in Section 16.2. In practice, we consider equations of the form

$$\ddot{z} + \alpha(t)\dot{z} + \zeta(t)z = 0, \tag{16.1}$$

where  $\alpha, \zeta \in C^{\infty}[\mathbb{R}, \mathbf{M}_m(\mathbb{C})]$ , and  $\alpha$  and  $\zeta$  converge after rescaling by an appropriate exponential factor. The conclusion we obtain in the ODE-unbalanced setting is that solutions typically either grow or decay super exponentially; cf. Lemma 16.4. This is one reason for considering such equations to be pathological. At the end of the chapter, in Section 16.11, we consider the behaviour of the individual modes, and provide conditions ensuring super exponential decay; cf. Lemma 16.56. This result is based on the analysis in Section 16.2.

The PDE-unbalanced setting. The PDE-unbalanced setting is substantially more complicated than the ODE-unbalanced setting. The reason for this is that in order to be able to derive conclusions in the former case, we need to appeal to the results of Part III of these notes. On the other hand, one simplifying aspect is the fact that in order to demonstrate super exponential growth of the energy, it is sufficient to consider (1.2) in the case that d = 1, R = 0 and f = 0. The reason for this is that if we consider the natural energy associated with (1.2) for general dand R, then this energy is bounded from below by the energy of the part of the solution which only depends on t and, say,  $x^j$ , the j'th "coordinate" on the d-torus. If we can demonstrate super exponential growth of such solutions, we are thus done. The steps of the analysis can then roughly be divided as follows.

Specifying the class of equations, preliminary steps. In Section 16.3 we introduce conditions such that the results of Part III apply, and such that we, in the end, are able to obtain super exponential growth. Under these assumptions, we then consider the modes of the solutions. Since the solutions of interest are defined on  $I \times \mathbb{S}^1$ , we only need to keep track of one frequency, say  $n_1$ . For a given equation and a given  $n_1$ , we gradually focus our attention on a more and more restricted time interval. The reason for doing so is that we want to construct a time sequence of the form introduced in Definition 15.5 and to appeal to Lemma 15.11. In fact, by appealing to Lemma 15.11, we can estimate how the solution evolves along the time sequence. For this to be useful we, however, need to restrict the time interval so that the time sequence is well defined and so that the the error terms arising in Lemma 15.11 are small. Moreover, even though Lemma 15.11 yields interesting information concerning the evolution of the solution, we, e.g., need to know that the constituents of the matrix  $R^{ab}_{\text{pre},k}$  appearing on the right hand side of (15.57) have some definite behaviour. Due to the form of our assumptions, we can only expect this to be the case in an asymptotic regime, where t is large. Summing up, it is thus necessary to restrict the value of  $n_1$  in such a way that there is a time interval in which the behaviour is oscillatory and so that the definition of the time sequence makes sense. Second, it is necessary to restrict the time sequence from above so that the error terms appearing in Lemma 15.11 are small. Third, t has to be large enough that we are in the asymptotic regime. Since there are bounds on t both from above and below, there is a risk that the different conditions contradict each other. However, this can be avoided by simply increasing the frequency  $n_1$ . In fact, it turns out that the upper bounds (when they exist) increase at least logarithmically with  $n_1$ . On the other hand, the lower bounds are independent of  $n_1$ ; they only depend on the coefficients of (1.2). In Section 16.4 we derive upper bounds on t ensuring that the error terms appearing in Lemma 15.11 are small and that the sequence introduced in Definition 15.5 is well defined. In Section 16.5, we then introduce the condition that the quotient  $e^{-\beta_X t} X^j(t) / [g^{jj}(t)]^{1/2}$  should converge to a matrix  $Y^j_\infty$  for some  $j \in \{1, \ldots, d\}$  and some  $\beta_X > 0$ , where  $Y_{\infty}^j$  has an eigenvalue with non-zero real part. Moreover, we reformulate the iteration derived in Lemma 15.11 so that the source of the super exponential growth becomes more apparent. Finally, we derive an estimate for the matrix appearing in the new iteration; cf. Lemma 16.17. This result is the basis for the proof of super exponential growth.

The main estimate. Once we have taken the preliminary steps, we are in a position to prove super exponential growth of the individual modes. This is the subject of Section 16.6. The main tool is the iteration derived previously. In practice we would thus like to estimate a matrix product. On the other hand, we are interested in situations where the number of factors tends to infinity. Moreover, we do not know the individual factors exactly; we only have an estimate. Finally, the product can be expected to contract some directions and expand others. On the other hand, we only need to find some appropriate initial datum for the iteration that leads to super exponential growth.

The first step of the analysis is to demonstrate that the sums of the error terms (that appear when estimating how solutions to the iteration grow) can be approximated by integrals which, in their turn, can be computed. The relevant statement is given in Lemma 16.22. In Lemma 16.23 we then state the main estimate. The proof of this lemma is based, first, on a restriction of the time interval as described above. Second, given an  $\epsilon > 0$ , we reformulate the iteration so as to ensure

that the influence of the off-diagonal parts of the Jordan blocks on the growth of the solution is bounded by an error term whose size is quantified by  $\epsilon$ . Once these steps have been taken, we divide the components, say  $\check{\rho}_k$ , of the iteration, into two parts, say  $\check{\rho}_{k,a}$  and  $\check{\rho}_{k,b}$ . The idea is then that  $\check{\rho}_{k,a}$  corresponds to the components of  $\check{\rho}_k$  with maximal growth. However, we modify the iteration so that the maximal growth corresponds to no growth at all. In other words, we wish to demonstrate that  $\check{\rho}_{k,a}$  does not decay too much, whereas  $\check{\rho}_{k,b}$  is much smaller in size than  $|\check{\rho}_{k,a}|$ . In the end, it turns out to be possible to prove that if  $|\check{\rho}_{k,b}|$  is much smaller than  $|\check{\rho}_{k,a}|$  initially (where initially here means late enough that we are in the asymptotic regime), then it remains much smaller for a long period of time. Moreover, during this period,  $\check{\rho}_{k,a}$  does not decay too much. Returning to the original variables, this then yields super exponential growth.

Super exponential instability. Given the main estimate, we are in a position to prove a super exponential instability result. In fact, given assumptions of the type described above, the solution u = 0 to (1.2) (where d = 1, R = 0 and f = 0) is unstable in the following sense. Fix  $\epsilon > 0$ . Then there is a sequence  $v_l$ ,  $1 \leq l \in \mathbb{Z}$ , of solutions to (1.2) such that the corresponding initial data at t = 0 converge to zero with respect to every Sobolev norm. Moreover, for each  $1 \leq l \in \mathbb{Z}$ , there is a sequence  $t_{l,k} \to \infty$  (as  $k \to \infty$ ) with the property that

$$\mathfrak{E}_{\hom}[v_l](t_{l,k}) \ge \exp\left[2\beta_X^{-1}\kappa(1-\epsilon)e^{\beta_X t_{l,k}}\right],$$

where  $\beta_X > 0$  is the constant appearing above;  $\kappa$  is the largest absolute value of a real part of an eigenvalue of the matrix  $Y_{\infty}^1/2$  (cf. the above); and

$$\mathfrak{E}_{\text{hom}}[v](t) := \frac{1}{2} \int_{\mathbb{S}^1} [|v_t(x,t)|^2 + g^{11}(t)|v_x(x,t)|^2] dx.$$

The relevant result is stated in Lemma 16.27.

Generic super exponential growth, general equations. As a next step, it is natural to consider general equations of the form (1.2). The only assumption we need to make is that there is one  $j \in \{1, \ldots, d\}$  such that

$$u_{tt} - g^{jj}(t)\partial_j^2 u - 2g^{0j}(t)\partial_t\partial_j u + \alpha(t)u_t + X^j(t)\partial_j u + \zeta(t)u = 0$$
(16.2)

satisfies the assumptions guaranteeing super exponential instability; cf. the above. Proving density of initial data yielding super exponential growth is then fairly straightforward. However, as we demonstrate in Section 16.11, the set of initial data yielding super exponential decay is sometimes also dense. It is therefore of interest to prove that super exponential growth represents the generic behaviour. In the end we prove two corresponding results; Propositions 16.34 and 16.37. Proposition 16.34 states that there is a set of initial data, say  $\mathcal{A}$ , which is a countable intersection of open and dense sets (with respect to the  $C^{\infty}$ -topology) and such that solutions corresponding to initial data in  $\mathcal{A}$  exhibit super exponential growth. Proposition 16.37 states, roughly speaking, that the set of initial data whose corresponding solutions do not exhibit super exponential growth has infinite codimension. Finally, in Subsection 16.8.3 we also demonstrate that the super exponential growth of solutions can be detected by solutions with finite frequency content; cf. Proposition 16.40. This is of interest due to the fact that solutions with finite frequency content sometime exhibit super exponential decay.

Energy estimates. In Section 16.9 we derive upper bounds on the growth of the energy under the same assumptions that we make when proving the super exponential instability. In particular, we consider (16.2) with j = 1. In the end we prove that if  $Y_{\infty}^1$  is Hermitian, then the energy of solutions does not grow faster than  $\exp\left[2\beta_X^{-1}\kappa(1+\epsilon)e^{\beta_X t}\right]$ ; cf. Lemma 16.42 and Remark 16.43. In that setting, the previously derived lower bounds on the growth of the energy are thus essentially optimal.

Gevrey classes. In Section 16.10, we consider solutions corresponding to initial data in Gevrey classes. The reason for taking an interest in this topic is the following. There are equations of the form (16.2) (with j = 1) such that there are smooth solutions exhibiting super exponential

growth and such that solutions with finite frequency content exhibit super exponential decay. It is therefore of interest to consider function classes interpolating between smooth and real analytic solutions and to maximise the degree of regularity of the solutions exhibiting super exponential growth. This is the topic of Section 16.10. To begin with, we provide two different characterisations of Gevrey classes; a standard characterisation and one more adapted to the problem we are interested in here. We also demonstrate that Gevrey type regularity is preserved under the evolution, given appropriate assumptions. Finally, we then calculate a Gevrey index characterising the best regularity (we can demonstrate) of the solutions exhibiting super exponential growth; cf. Proposition 16.52.

*Mode analysis.* Finally, in Section 16.11, we analyse how the modes of the solution develop over time.

# 16.2 Unbalanced equations, the ODE setting

Consider the homogeneous version of (1.2) when  $g^{00} = -1$ . The simplest special case of this equation is obtained by considering spatially homogeneous solutions. The relevant equation is then given by

$$\ddot{x} + \alpha(t)\dot{x} + \zeta(t)x = 0, \tag{16.3}$$

where  $\alpha, \zeta \in C^{\infty}[I, \mathbf{M}_m(\mathbb{C})].$ 

**Definition 16.1.** Let  $\alpha, \zeta \in C^{\infty}[I, \mathbf{M}_m(\mathbb{C})]$ , where *I* is an open interval and  $[0, \infty) \subset I$ . Then (16.3) is said to be *convergent and unbalanced* if the following holds:

• there are constants  $C_{cu}, \gamma, \eta_{cu} > 0$  and matrices  $\alpha_{\infty}, \zeta_{\infty} \in \mathbf{M}_m(\mathbb{C})$  such that

$$\|\gamma^{-1}e^{-\gamma t}\alpha(t) - \alpha_{\infty}\| + \|\gamma^{-2}e^{-2\gamma t}\zeta(t) - \zeta_{\infty}\| \le C_{\mathrm{cu}}e^{-\eta_{\mathrm{cu}}t}$$

for  $t \geq 0$ ,

• one of  $\alpha_{\infty}, \zeta_{\infty}$  is non-zero.

**Remark 16.2.** Due to the requirements, the norm of at least one of the matrices  $\alpha(t)$  and  $\zeta(t)$  is unbounded as  $t \to \infty$ .

**Remark 16.3.** The case that  $e^{-\gamma t}\alpha(t)$  and  $e^{-2\gamma t}\zeta(t)$  converge exponentially for some real number  $\gamma \leq 0$  can be dealt with as in Chapter 9. We therefore do not consider this case here.

Given that (16.3) is convergent and unbalanced, the matrix

$$A_{\infty} := \begin{pmatrix} 0 & \mathrm{Id}_m \\ -\zeta_{\infty} & -\alpha_{\infty} \end{pmatrix}$$
(16.4)

determines the dominant part of the asymptotic behaviour.

**Lemma 16.4.** Consider the equation (16.3). Assume that it is convergent and unbalanced, cf. Definition 16.1, and define  $A_{\infty}$  by (16.4). Let  $\kappa_1 := \kappa_{\max}(A_{\infty})$ ; cf. Definition 4.3. For every  $\epsilon > 0$ , there is a constant  $C_{+,\epsilon} > 0$  such that for every solution x to (16.3)

$$|x(t)| + |\dot{x}(t)| \le C_{+,\epsilon} \exp[(\kappa_1 + \epsilon)e^{\gamma t}][|x(0)| + |\dot{x}(0)|]$$
(16.5)

for all  $t \ge 0$ . Moreover, for every  $\epsilon > 0$ , there is a solution x to (16.3) and a constant  $C_{-,\epsilon} > 0$  such that

$$C_{-,\epsilon} \exp[(\kappa_1 - \epsilon)e^{\gamma t}] \le |x(t)| + |\dot{x}(t)| \tag{16.6}$$

for all  $t \geq 0$ .

**Remark 16.5.** The lemma illustrates that convergent and unbalanced equations typically have solutions with pathological properties. In fact, if  $\kappa_1 > 0$ , there are solutions that grow super exponentially. If  $\kappa_1 < 0$ , there are solutions that decay super exponentially (meaning that they grow super exponentially towards the past). Both of these situations we consider to be pathological.

**Remark 16.6.** The constants  $C_{+,\epsilon}$  and  $C_{-,\epsilon}$  only depend on  $C_{cu}$ ,  $A_{\infty}$ ,  $\eta_{cu}$ ,  $\gamma$  and  $\epsilon$ 

**Remarks 16.7.** If  $\kappa_1$  is non-zero, (16.5) and (16.6) yield a rough idea of the behaviour of solutions. However, this is not true if  $\kappa_1 = 0$ . In that case, a further analysis is necessary. The solutions with a lower bound of the form (16.6) (that we construct in the proof) are such that they are stable under perturbations in the sense that a small perturbation of the corresponding initial data yields solutions satisfying a bound of the form (16.6). Moreover, given a solution to (16.3) satisfying

$$|x(t)| + |\dot{x}(t)| \le C \exp[(\kappa_1 - \eta)e^{\gamma t}]$$

for some constants  $C, \eta > 0$  and all  $t \ge 0$ , it can be perturbed to a solution satisfying (16.6); this is an immediate consequence of the existence of solutions satisfying (16.6).

*Proof.* Letting  $x(t) = y(e^{\gamma t})$ , (16.3) can be written

$$y''(e^{\gamma t}) + e^{-\gamma t}y'(e^{\gamma t}) + \gamma^{-1}e^{-\gamma t}\alpha(t)y'(e^{\gamma t}) + \gamma^{-2}e^{-2\gamma t}\zeta(t)y(e^{\gamma t}) = 0.$$
 (16.7)

Introducing the time coordinate  $\tau = e^{\gamma t}$ , (16.7) can be written

$$y'' + \tau^{-1}y' + \alpha_{\rm eff}(\tau)y' + \zeta_{\rm eff}(\tau)y = 0,$$

where

$$\alpha_{\text{eff}}(\tau) := \gamma^{-1} \tau^{-1} \alpha(\gamma^{-1} \ln \tau), \quad \zeta_{\text{eff}}(\tau) := \gamma^{-2} \tau^{-2} \zeta(\gamma^{-1} \ln \tau).$$

Moreover,

$$\|\alpha_{\rm eff}(\tau) - \alpha_{\infty}\| + \|\zeta_{\rm eff}(\tau) - \zeta_{\infty}\| \le C_{\rm cu}\tau^{-\eta_{\rm cu}/\gamma}$$

for  $\tau \geq 1$ , where  $\alpha_{\infty}$  and  $\zeta_{\infty}$  are the matrices given in Definition 16.1. Introduce

$$v := \begin{pmatrix} y \\ y' \end{pmatrix}, \quad A_{\text{rem}}(\tau) := \begin{pmatrix} 0 & 0 \\ -\zeta_{\text{eff}}(\tau) + \zeta_{\infty} & -\tau^{-1} \text{Id}_m - \alpha_{\text{eff}}(\tau) + \alpha_{\infty} \end{pmatrix}$$

and let  $A_{\infty}$  be as in the statement of the lemma. Then

$$v' = A_{\infty}v + A_{\rm rem}(\tau)v. \tag{16.8}$$

Upper bound on the solution. Let T be a complex matrix such that  $B_{\infty} := TA_{\infty}T^{-1}$  consists of generalised Jordan blocks, where the blocks are ordered in such a way that the ones corresponding to eigenvalues with maximal real part are on the top left; say that the first  $m_{\rm s}$  elements on the diagonal of  $B_{\infty}$  have maximal real part and define  $m_{\rm u}$  so that  $m_{\rm s} + m_{\rm u} = 2m$ . Note that, given  $\epsilon > 0$ , T can be chosen so that the generalised Jordan blocks are such that the non-zero off-diagonal terms equal  $\epsilon/2$ ; cf. Lemma 9.9 and its proof. Note, moreover, that T only depends on  $A_{\infty}$  and  $\epsilon$ . The equation (16.8) can then be rewritten

$$w' = B_{\infty}w + B_{\rm rem}(\tau)w,\tag{16.9}$$

where  $B_{\rm rem}(\tau) := TA_{\rm rem}(\tau)T^{-1}$  and w := Tv. Note that the norm of  $B_{\rm rem}(\tau)$  is bounded by  $C_{\rm cu,\epsilon}\langle \tau \rangle^{-\eta_{\rm min}}$  for all  $\tau \geq 1$ , where  $\eta_{\rm min} = \min\{1, \eta_{\rm cu}/\gamma\}$  and the constant  $C_{\rm cu,\epsilon}$  only depends on  $C_{\rm cu}$ ,  $A_{\infty}$  and  $\epsilon$ . Due to (16.9),

$$\frac{d}{d\tau}|w|^2 \le 2\left(\kappa_1 + \epsilon/2 + C_{\mathrm{cu},\epsilon}\langle \tau \rangle^{-\eta_{\min}}\right)|w|^2.$$

Given  $\epsilon > 0$  and a solution w, there is thus a constant  $C_{up,\epsilon}$  such that

$$|w(\tau)| \le C_{\mathrm{up},\epsilon} \exp[(\kappa_1 + 3\epsilon/4)\tau]|w(1)|$$

for  $\tau \geq 1$ , where  $C_{up,\epsilon}$  only depends on  $C_{cu}$ ,  $A_{\infty}$ ,  $\eta_{cu}$ ,  $\gamma$  and  $\epsilon$ . Since

$$|x(t)| + |\dot{x}(t)| = |y(e^{\gamma t})| + \gamma e^{\gamma t} |y'(e^{\gamma t})| \le C_{\text{fin},\epsilon} e^{\gamma t} |w(e^{\gamma t})|,$$
(16.10)

(16.5) holds, where  $C_{\text{fin},\epsilon}$  and  $C_{+,\epsilon}$  only depend on  $C_{\text{cu}}$ ,  $A_{\infty}$ ,  $\eta_{\text{cu}}$ ,  $\gamma$  and  $\epsilon$ .

**Lower bound on solutions.** In order to obtain a lower bound, it is convenient to divide the components of w into two parts, say  $w_s$  and  $w_u$ , where  $w_s$  and  $w_u$  take their values in  $\mathbb{C}^{m_s}$  and  $\mathbb{C}^{m_u}$  respectively (in case  $m_u = 0$ , it is trivial to prove the stated lower bound, so we here assume that  $m_u > 0$ ). It is convenient to rewrite (16.9) as

$$w'_{\rm s} = B_{\infty,11} w_{\rm s} + B_{\rm rem,11}(\tau) w_{\rm s} + B_{\rm rem,12}(\tau) w_{\rm u}, \qquad (16.11)$$

$$w'_{\rm u} = B_{\infty,22} w_{\rm u} + B_{\rm rem,21}(\tau) w_{\rm s} + B_{\rm rem,22}(\tau) w_{\rm u}.$$
(16.12)

Let  $\kappa_2$  be the second largest real part of an eigenvalue of  $A_{\infty}$ . Then  $\kappa_2 < \kappa_1$ . Note that  $\|B_{\text{rem},ij}(\tau)\|$  satisfies the same bound as  $\|B_{\text{rem}}(\tau)\|$ . Thus

$$\frac{d}{d\tau}|w_{\rm s}|^2 \ge 2\kappa_1|w_{\rm s}|^2 - \epsilon|w_{\rm s}|^2 - C_{\rm cu,\epsilon}\langle\tau\rangle^{-\eta_{\rm min}}(3|w_{\rm s}|^2 + |w_{\rm u}|^2),\tag{16.13}$$

$$\frac{d}{d\tau}|w_{\rm u}|^2 \le 2\kappa_2|w_{\rm u}|^2 + \epsilon|w_{\rm u}|^2 + C_{\rm cu,\epsilon}\langle\tau\rangle^{-\eta_{\rm min}}(|w_{\rm s}|^2 + 3|w_{\rm u}|^2).$$
(16.14)

Assume that  $|w_s(\tau)| \neq 0$  and compute

$$\frac{d}{d\tau} \frac{|w_{\rm u}|^2}{|w_{\rm s}|^2} \le 2(\kappa_2 - \kappa_1 + \epsilon) \frac{|w_{\rm u}|^2}{|w_{\rm s}|^2} + K_{\rm cu,\epsilon} \langle \tau \rangle^{-\eta_{\rm min}} \left(1 + \frac{|w_{\rm u}|^2}{|w_{\rm s}|^2} + \frac{|w_{\rm u}|^4}{|w_{\rm s}|^4}\right),\tag{16.15}$$

where  $K_{cu,\epsilon}$  has the same dependence as  $C_{cu,\epsilon}$ . Without loss of generality, we may assume  $\epsilon$  to be such that  $\epsilon \leq (\kappa_1 - \kappa_2)/2$ . Given that such an  $\epsilon > 0$  has been fixed, fix  $\tau_0$  so that if  $K_{cu,\epsilon}$  is the constant appearing on the right hand side of (16.15), then  $K_{cu,\epsilon}\langle \tau \rangle^{-\eta_{\min}} \leq (\kappa_1 - \kappa_2)/8$  for  $\tau \geq \tau_0$ . Fix initial data for  $w_s$  and  $w_u$  at  $\tau_0$  in such a way that  $|w_u(\tau_0)|^2/|w_s(\tau_0)|^2 \leq 1/4$ . Assume, in order to obtain a contradiction, that there is a  $\tau_1 \geq \tau_0$  such that  $|w_u(\tau_1)|^2/|w_s(\tau_1)|^2 = 1/2$ . Then, at  $\tau = \tau_1$ ,

$$\frac{d}{d\tau}\frac{|w_{\mathbf{u}}|^2}{|w_{\mathbf{s}}|^2} \le \frac{1}{4}(\kappa_2 - \kappa_1) < 0.$$

Thus  $|w_{\rm u}|^2/|w_{\rm s}|^2$  is strictly decreasing at  $\tau = \tau_1$ , which means that  $|w_{\rm u}(\tau)|^2/|w_{\rm s}(\tau)|^2 \leq 1/2$  for all  $\tau \geq \tau_0$ . Choose initial data at  $\tau_0$  so that  $w_{\rm u}(\tau_0) = 0$  and  $|w_{\rm s}(\tau_0)| = 1$ . Returning to (16.13), it is then clear that for any  $\epsilon > 0$ , there is a constant  $C_{\rm low,\epsilon} > 0$  such that

$$C_{\log,\epsilon} e^{(\kappa_1 - \epsilon)\tau} \le |w(\tau)|$$

for all  $\tau \geq \tau_0$ , where  $C_{\text{low},\epsilon}$  only depends on  $C_{\text{cu}}$ ,  $A_{\infty}$ ,  $\eta_{\text{cu}}$ ,  $\gamma$  and  $\epsilon$ . Since we can change the upper bound in (16.10) to a lower bound by simply removing the factor  $e^{\gamma t}$  on the far right hand side, and changing the value of  $C_{\text{fin},\epsilon}$ , it is clear that the constructed solution satisfies (16.6) for  $t \geq t_0 := \gamma^{-1} \ln \tau_0$ . Note also that the solution is stable under perturbations of the initial data. In order to ensure that the estimate holds in  $[0, t_0]$ , note that  $t_0$  only depends on  $C_{\text{cu}}$ ,  $A_{\infty}$ ,  $\epsilon$ ,  $\eta_{\text{cu}}$  and  $\gamma$ . Combining this observation with Definition 16.1 yields the conclusion that

$$\|\alpha(t)\| + \|\zeta(t)\| \le C_{0,\epsilon}$$

for  $0 \le t \le t_0$ , where  $C_{0,\epsilon}$  only depends on  $C_{cu}$ ,  $A_{\infty}$ ,  $\epsilon$ ,  $\eta_{cu}$  and  $\gamma$ . Using this estimate, it can be verified that (16.6) is valid for all  $t \ge 0$ .

#### 16.3Unbalanced equations, the PDE setting

Let us turn to the case that the equation (1.2) is not balanced due to a lack of boundedness of the norm of  $X(\iota, t)$ . Since we are only interested in producing counterexamples, it turns out to be sufficient to consider solutions that depend on only one of the spatial coordinates. In fact, we here assume that d = 1 and that R = 0. Then (1.2) is an equation on  $\mathbb{S}^1 \times I$ . When considering the corresponding equation for the Fourier coefficients, (24.16), it turns out to be sufficient to assume that  $n_1 > 0$  (and we therefore typically make this assumption from now on). In this context, the definitions (24.13) and (10.1) become

$$\mathfrak{g}(n_1,t) = [g^{11}(t)]^{1/2} |n_1|, \quad \ell(n_1,t) = \ln \mathfrak{g}(n_1,t)$$

and

$$X(n_1,t) = \frac{n_1 X^1(t)}{[g^{11}(t)]^{1/2} |n_1|} = \frac{X^1(t)}{[g^{11}(t)]^{1/2}}, \quad \sigma(n_1,t) = \frac{g^{01}(t)}{[g^{11}(t)]^{1/2}}.$$
 (16.16)

In what follows, it will also be convenient to introduce the notation

$$\mathfrak{l}(t) := \ln[g^{11}(t)]^{1/2}, \quad Y^1(t) := \frac{X^1(t)}{[g^{11}(t)]^{1/2}}, \quad \varsigma(t) := \frac{g^{01}(t)}{[g^{11}(t)]^{1/2}}.$$
(16.17)

The reason we introduce this notation is that the relations (16.16) do not hold in general; they are based on the assumption that  $n_1 > 0$ . Note also that  $Y^1$ ,  $\varsigma$  and  $\mathfrak{l}$  are independent of  $n_1$  and that  $\mathfrak{l} = \ell$ . Next, we want to state conditions that ensure that (1.2) is oscillation adapted; cf. Definition 14.1.

**Lemma 16.8.** Consider (1.2). Assume the associated metric to be such that (M,g) is a canonical separable cosmological model manifold. Assume, moreover, that d = 1 and that R = 0. Let  $\mathfrak{l}, Y^1$ and  $\varsigma$  be defined by (16.17). Assume that there are real numbers  $\beta_{\rm sh} < 0$ ,  $\mathfrak{b} > 0$ ,  $\eta_{\rm ode} > 0$ ,  $C_{\rm sh} > 0$ ,  $C_X > 0, C_{ode} > 0, \beta_X \ge 0$  and  $\beta_{ode}$  such that

$$\beta_X - \beta_{\text{ode}} = \eta_{\text{ode}},\tag{16.18}$$

$$|\beta_{\rm sh}| + |\beta_X| + |\beta_{\rm ode}| \le \mathfrak{b}, \tag{16.19}$$

$$|\mathfrak{l}(t)| + |\mathfrak{l}(t)| \le \mathfrak{b} \tag{16.20}$$

for all  $t \geq 0$ , and

$$\begin{aligned} |\varsigma(t)| + |\dot{\varsigma}(t)| &\leq C_{\rm sh} e^{\beta_{\rm sh} t}, \\ |t|| + ||\dot{Y}^1(t)|| &\leq C_X e^{\beta_X t}. \end{aligned}$$
(16.21)

$$||Y^{1}(t)|| + ||Y^{1}(t)|| \le C_{X} e^{\beta_{X} t}, \qquad (16.22)$$

$$\|\dot{\alpha}(t)\| + \|\alpha(t)\| \le C_{\text{ode}} e^{\beta_{\text{ode}} t},\tag{16.23}$$

$$\|\dot{\zeta}(t)\| + \|\zeta(t)\| \le C_{\text{ode}}^2 e^{2\beta_{\text{ode}}t} \tag{16.24}$$

for all  $t \ge 0$ . Then (1.2) is oscillation adapted, with  $f_{\rm sh}$ ,  $f_X$  and  $f_{\rm ode}$  replaced by  $C_{\rm sh}e^{\beta_{\rm sh}t}$ ,  $C_Xe^{\beta_X t}$ and  $C_{\text{ode}}e^{\beta_{\text{ode}}t}$  respectively.

**Remark 16.9.** The statement is of course true regardless of the sign of  $\beta_{sh}$  and  $\beta_X$ , and regardless of whether the left hand side of (16.18) is positive or not. However, we are here only interested in the case  $\beta_{\rm sh} < 0$ ,  $\beta_X \ge 0$  and  $\eta_{\rm ode} > 0$ , which is why we immediately restrict to that case.

*Proof.* The statement follows from the facts that  $\dot{\mathfrak{l}} = \dot{\ell}$ ,  $\varsigma = \pm \sigma$  and  $Y^1 = \pm X$  (depending on whether  $n_1$  is positive or negative). 

### 16.4 Subdividing the time interval

Before turning to the actual estimates, it is natural to restrict the time interval so that the results of Chapter 15 can be applied. This is the purpose of the present section. In what follows, we are only interested in the case that  $n_1$  is very large. In that context, there is a time interval  $[0, t_{\text{max}}]$  such that the conditions of Lemma 14.4 are satisfied. In order to obtain a more quantitative control over the size of this interval, it is convenient to make the following assumption.

Assumption 16.10. Consider (1.2). Assume the associated metric to be such that (M, g) is a canonical separable cosmological model manifold. Assume, moreover, that d = 1 and that R = 0. Assume, finally, that there are  $\beta_1, \beta_2 \in \mathbb{R}$  and constants  $c_{m,l} > 0, l = 1, 2$ , such that

$$c_{m,1}e^{2\beta_1 t} \le g^{11}(t) \le c_{m,2}e^{2\beta_2 t} \tag{16.25}$$

for all  $t \geq 0$ .

**Remark 16.11.** The estimate (16.25) is an immediate consequence of (16.20). Nevertheless, we state it explicitly here in order to introduce the terminology  $\beta_l$  and  $c_{m,l}$ .

When considering an interval  $I_a$  of the form constructed in Lemma 14.4, it is of interest to bound the quantity  $\varepsilon_a$  appearing in (14.33):

$$\varepsilon_a := \frac{\mathfrak{f}_{\mathrm{sh}}(t_a)}{\mathfrak{g}(t_a)} [1 + \mathfrak{f}_{\mathrm{ode}}(t_a) + \mathfrak{f}_X(t_a)] + \frac{1 + \mathfrak{f}_X^2(t_a) + \mathfrak{f}_{\mathrm{ode}}^2(t_a)}{\mathfrak{g}^2(t_a)}.$$
 (16.26)

Given that the assumptions of Lemma 16.8 hold and that Assumption 16.10 holds,

$$\varepsilon_a \le C_{\text{par}} \frac{e^{(\beta_X + \beta_{\text{sh}})t_a}}{|n_1|e^{\beta_1 t_a}} + C_{\text{par}} \frac{e^{2\beta_X t_a}}{n_1^2 e^{2\beta_1 t_a}},\tag{16.27}$$

where  $C_{\text{par}}$  is a constant depending only on  $\mathbf{c}_{\text{par}}$  defined by

$$\mathbf{c}_{\text{par}} := (\beta_{\text{sh}}, \mathfrak{b}, \eta_{\text{ode}}, C_{\text{sh}}, C_X, C_{\text{ode}}, \beta_X, \beta_{\text{ode}}, \beta_1, \beta_2, c_{\text{m},1}, c_{\text{m},2}).$$
(16.28)

In the analysis to follow, there are two things we want to be fulfilled. First of all, we need to know that the conditions of Lemma 14.4 are fulfilled. Second, we wish to ensure that the right hand side of (16.27) is small. The following lemma provides assumptions that ensure that these conditions hold.

**Lemma 16.12.** Consider (1.2). Assume the associated metric to be such that (M, g) is a canonical separable cosmological model manifold. Assume, moreover, that d = 1 and that R = 0. Assume, finally, that the conditions of Lemma 16.8 are fulfilled and that Assumption 16.10 holds. Let  $0 < \epsilon \leq 1$ . If  $\beta_X - \beta_1 > 0$ , there are constants  $\nu_{\text{par},\epsilon}$  and  $C_{\text{par},\epsilon}$ , depending only on  $\mathbf{c}_{\text{par}}$ , defined in (16.28), and  $\epsilon$ , such that if  $|n_1| \geq \nu_{\text{par},\epsilon}$ , then

$$T_{\text{par},\epsilon} := \frac{1}{\beta_X - \beta_1} \ln |n_1| + C_{\text{par},\epsilon} > 0$$
(16.29)

and

$$\varepsilon_a \le \epsilon, \quad \mathfrak{g}(n_1, t) \ge \max\{4\pi\mathfrak{b}, 2\}, \quad \frac{e^{\beta_X t}}{\mathfrak{g}(n_1, t)} \le \epsilon$$
(16.30)

for  $0 \leq t, t_a \leq T_{\text{par},\epsilon}$ , where  $\varepsilon_a$  is defined in (16.26). If  $\beta_X - \beta_1 \leq 0$ , there is a constant  $\nu_{\text{par},\epsilon}$ , depending only on  $\mathbf{c}_{\text{par}}$  and  $\epsilon$ , such that if  $|n_1| \geq \nu_{\text{par},\epsilon}$ , then (16.30) holds for all  $t, t_a \geq 0$ .

**Remark 16.13.** In what follows, we assume  $n_1$  to be such that (16.30) holds with  $\epsilon = 1$  for  $t \in [0, t_{\max})$ , where  $t_{\max} = \infty$  in case  $\beta_X - \beta_1 \leq 0$  and  $t_{\max} = T_{\max,1}$  in case  $\beta_X - \beta_1 > 0$ ; here  $T_{\max,\epsilon}$  is defined in (16.29). Note that it is sufficient to assume that  $|n_1| \geq \nu_{\max}$  (where  $\nu_{\max}$  only depends on  $\mathbf{c}_{\max}$ ) in order to obtain this conclusion.

**Remark 16.14.** Given that the assumptions of Remark 16.13 are satisfied, it is clear that we can construct a sequence  $\{t_k\}$  as in Definition 15.5 with  $0 \le t_0 < t_{\text{max}}$ . Then  $t_{\text{fin}} \ge t_{\text{max}} - 2\pi$ , where  $t_{\text{max}}$  is introduced in Remark 16.13.

*Proof.* The three conditions we want to hold are (16.30). Let us, to begin with, focus on the second condition. Due to Assumption 16.10, it is ensured by requiring

$$c_{\mathrm{m},1}e^{2\beta_1 t}|n_1|^2 \ge (\max\{4\pi\mathfrak{b},2\})^2.$$
 (16.31)

As a consequence, there is a  $0 < \nu_{\text{par}} \in \mathbb{R}$ , only depending on  $\mathbf{c}_{\text{par}}$ , such that the following holds. If  $\beta_1 \geq 0$  and  $n_1$  satisfies  $|n_1| \geq \nu_{\text{par}}$ , then (16.31) holds for all  $t \geq 0$ . If  $\beta_1 < 0$  and  $n_1$  satisfies  $|n_1| \geq \nu_{\text{par}}$ , then (16.31) holds for all  $t \geq 0$  such that

$$t \le -\frac{1}{\beta_1} \ln |n_1| + C_{\text{par}},\tag{16.32}$$

where  $C_{\text{par}}$  only depends on  $\mathbf{c}_{\text{par}}$ , and the right hand side is strictly positive if  $|n_1| \ge \nu_{\text{par}}$ .

As a second step, we wish to ensure that the first and third estimates in (16.30) hold. To prove that the first estimate holds, it is, due to (16.27), sufficient to assume that right hand side of (16.27) is bounded by  $\epsilon$ . Given  $0 < \epsilon \leq 1$ , there is a  $C_{\text{par}}$  (depending only on  $\mathbf{c}_{\text{par}}$ ) such that if

$$\frac{e^{\beta_X t_a}}{|n_1|e^{\beta_1 t_a}} \le C_{\text{par}}\epsilon,\tag{16.33}$$

then the right hand side of (16.27) is bounded from above by  $\epsilon$  (note that a condition of this type is also sufficient in order to ensure that the third estimate in (16.30) holds). In this case, given  $0 < \epsilon \leq 1$ , there is a  $\nu_{\text{par},\epsilon} > 0$ , depending only on  $\mathbf{c}_{\text{par}}$  and  $\epsilon$ , such that the following holds. If  $\beta_X - \beta_1 \leq 0$  and  $|n_1| \geq \nu_{\text{par},\epsilon}$ , then (16.33) holds for all  $t_a \geq 0$ . If  $\beta_X - \beta_1 > 0$  and  $|n_1| \geq \nu_{\text{par},\epsilon}$ , then (16.33) holds for all  $t_a \geq 0$  such that

$$t_a \le \frac{1}{\beta_X - \beta_1} \ln |n_1| + C_{\operatorname{par},\epsilon},\tag{16.34}$$

where  $C_{\text{par},\epsilon}$  only depends on  $\mathbf{c}_{\text{par}}$  and  $\epsilon$ , and the right hand side is strictly positive if  $|n_1| \ge \nu_{\text{par},\epsilon}$ .

In order to prove the first statement of the lemma, assume that  $\beta_X - \beta_1 > 0$ . Then there are two possibilities: either  $\beta_1 < 0$  or  $\beta_1 \ge 0$ . If  $\beta_1 < 0$  and  $|n_1| \ge \max\{\nu_{\text{par}}, \nu_{\text{par},\epsilon}\}$ , then it is sufficient to assume that t and  $t_a$  satisfy (16.32) and (16.34) in order for (16.30) to be satisfied. However, these restrictions on t and  $t_a$  can be translated into  $0 \le t, t_a \le T_{\text{par},\epsilon}$ . In case  $\beta_1 \ge 0$  and  $|n_1| \ge \nu_{\text{par}}$ , the second estimate in (16.30) holds for all  $t \ge 0$ . Regardless of whether  $\beta_1 < 0$  or  $\beta_1 \ge 0$ , the first statement of the lemma thus holds.

In order to prove the second statement, assume that  $\beta_X - \beta_1 \leq 0$ . Since  $\beta_X \geq 0$ , this implies that  $\beta_1 \geq 0$ . Due to the above arguments, it is thus sufficient to assume that  $|n_1| \geq \max\{\nu_{\text{par}}, \nu_{\text{par},\epsilon}\}$  in order to ensure that (16.30) holds for all  $t \geq 0$ .

# 16.5 Basic properties of the iteration

As a next step, we wish to apply Lemma 15.11, in particular (15.54); note that if the assumptions of Lemma 16.12 and Remark 16.13 are satisfied, there is a non-empty time sequence of the form given in Definition 15.5, starting at  $0 \le t_0 < t_{\max}$ . We are here mainly interested in homogeneous equations, and in that case, (15.54) implies that  $w_{\text{fin},k+1} = \Xi_{\text{fin}}^{k,+} w_{\text{fin},k}$ . However, before proceeding, it turns out to be convenient to transform  $w_{\text{fin},k}$  slightly. Moreover, since the transformation depends on more detailed knowledge concerning the coefficients of (1.2) than we have thus far assumed, we need to start by making an additional assumption. Assumption 16.15. Consider (1.2). Assume the associated metric to be such that (M, g) is a canonical separable cosmological model manifold. Assume, moreover, that d = 1 and that R = 0. Let  $Y^1$  be defined by (16.17) and the assumptions of Lemma 16.8 be fulfilled. Assume that there is a  $Y^1_{\infty} \in \mathbf{M}_m(\mathbb{C})$  and constants  $\eta_X > 0$  and  $K_X > 0$  such that

$$\|e^{-\beta_X t} Y^1(t) - Y^1_{\infty}\| \le K_X e^{-\eta_X t}$$
(16.35)

for all  $t \ge 0$ , where  $\beta_X \ge 0$  is the constant appearing in the statement of Lemma 16.8.

**Remark 16.16.** In what follows, we are mainly interested in the case that  $Y_{\infty}^1$  has an eigenvalue with a non-zero real part. However, we state this assumption separately in the lemmas to follow.

Let  $R_{\infty}$  be defined by

$$R_{\infty} := \frac{1}{2} \begin{pmatrix} Y_{\infty}^{1} & 0\\ 0 & -Y_{\infty}^{1} \end{pmatrix}.$$
 (16.36)

Let T be a complex  $2m \times 2m$ -matrix such that

$$J := TR_{\infty}T^{-1} \tag{16.37}$$

consists of Jordan blocks  $J_1, \ldots, J_r$ ;  $J = \text{diag}\{J_1, \ldots, J_r\}$ . Denote the eigenvalue corresponding to  $J_s$  by  $\lambda_s = \kappa_s + i\zeta_s$ , where  $\kappa_s$  and  $\zeta_s$  are real. Let  $l_s$  be the dimension of the Jordan block  $J_s$  and define

$$S_k := \operatorname{diag}\{\exp(-i\zeta_1\beta_X^{-1}e^{\beta_X t_k})\operatorname{Id}_{l_1}, \dots, \exp(-i\zeta_r\beta_X^{-1}e^{\beta_X t_k})\operatorname{Id}_{l_r}\}.$$
(16.38)

Here  $t_k$  is the time sequence given by Definition 15.5, starting at  $0 \le t_0 < t_{\text{max}}$ ; note that the sequence is non-empty if the assumptions of Lemma 16.12 and Remark 16.13 are satisfied (cf. Remark 16.14). Moreover, we here assume that  $\beta_X > 0$ , where  $\beta_X$  is the constant introduced in Lemma 16.8 and Assumption 16.15. Defining  $\rho_k$  and  $B_k$  according to

$$\rho_k := S_k T w_{\text{fin},k}, \quad B_k := S_{k+1} T \Xi_{\text{fin}}^{k,+} T^{-1} S_k^{-1}, \tag{16.39}$$

the equation (15.54) becomes  $\rho_{k+1} = B_k \rho_k$  (assuming f = 0). Recall here that  $w_{\text{fin},k}$  is defined in (15.53) and that  $\Xi_{\text{fin}}^{k,+}$  is introduced in the statement of Lemma 15.11; cf. also (15.62). Let  $\kappa := \kappa_{\max}(R_{\infty})$ ; cf. Definition 4.3. Introducing

$$\hat{\rho}_k := \exp\left(-\beta_X^{-1} e^{\beta_X t_k} \kappa\right) \rho_k, \quad \hat{B}_k := \exp\left(-\beta_X^{-1} e^{\beta_X t_{k+1}} \kappa + \beta_X^{-1} e^{\beta_X t_k} \kappa\right) B_k, \tag{16.40}$$

(15.54) can then be written

$$\hat{\rho}_{k+1} = \hat{B}_k \hat{\rho}_k. \tag{16.41}$$

The natural next step is to approximate  $\hat{B}_k$  in order to estimate the product of the matrices.

**Lemma 16.17.** Consider (1.2). Assume the associated metric to be such that (M, g) is a canonical separable cosmological model manifold. Assume, moreover, that d = 1 and that R = 0. Assume that the conditions of Lemma 16.8 are fulfilled with  $\beta_X > 0$  and that Assumptions 16.10 and 16.15 hold. If  $\beta_X - \beta_1 > 0$ , there are constants  $C_{\text{par}}$  and  $\nu_{\text{par}} > 0$  (depending only on  $\mathbf{c}_{\text{par}}$  introduced in (16.28)), and if  $\beta_X - \beta_1 \leq 0$ , there is a constant  $\nu_{\text{par}}$  (depending only on  $\mathbf{c}_{\text{par}}$ ) such that the following holds. If  $|n_1| \geq \nu_{\text{par}}$ , then (16.30), with  $\epsilon$  replaced by 1, holds for  $0 \leq t, t_a < t_{\text{max}}$ , where

$$t_{\max} := \begin{cases} T_{\max,1}, & \beta_X - \beta_1 > 0\\ & \infty, & \beta_X - \beta_1 \le 0 \end{cases}$$
(16.42)

and

$$T_{\text{par},1} := \frac{1}{\beta_X - \beta_1} \ln |n_1| + C_{\text{par},1} > 0;$$

#### 16.5. BASIC PROPERTIES OF THE ITERATION

cf. (16.29). Let  $\{t_k\}$  be a time sequence of the form constructed in Definition 15.5, where  $0 \leq t_0 < t_{\text{max}}$ . Then  $t_{\text{fin}}$ , given in Definition 15.5, satisfies  $t_{\text{fin}} \geq \max\{t_0, t_{\text{max}} - 2\pi\}$ . Assume that  $n_1 \geq \nu_{\text{par}}$ . Then, if  $k \geq 0$  and  $t_k < t_{\text{max}}$ ,

$$\left\|\hat{B}_{k} - \hat{B}_{k,\mathrm{app}}\right\| \leq C_{\mathrm{eep}}\left(\frac{e^{(\beta_{X} - \eta_{\mathrm{mn}})t_{k}}}{\mathfrak{g}(t_{k})} + \frac{e^{2\beta_{X}t_{k}}}{\mathfrak{g}^{2}(t_{k})}\right),\tag{16.43}$$

where  $\eta_{\rm mn} > 0$  is defined by

$$\eta_{\rm mn} := \min\{\eta_X, \eta_{\rm ode}, \beta_X, -\beta_{\rm sh}\}; \tag{16.44}$$

 $\hat{B}_{k,app}$  is given by

$$\hat{B}_{k,\mathrm{app}} := \mathrm{Id}_{2m} + \frac{2\pi e^{\beta_X t_k}}{\mathfrak{g}(t_k)} J_\mathrm{n}, \quad J_\mathrm{n} := \mathrm{Re}\{J\} - \kappa \mathrm{Id}_{2m};$$
(16.45)

J is the matrix introduced in (16.37);  $\kappa := \kappa_{\max}(R_{\infty})$ ; and  $C_{eep}$  only depends on  $\mathbf{c}_{eep}$  defined by

$$\mathbf{c}_{\text{eep}} := (\mathbf{c}_{\text{par}}, \eta_X, K_X, J, T), \tag{16.46}$$

where the J is defined by (16.37) and T is the matrix defined in connection with (16.37) (in (16.46), J and T should be thought of as vectors whose components are the entries of the matrices J and T respectively).

*Proof.* Since the conditions of Lemma 16.8 are fulfilled, we know that (1.2) is oscillation adapted, with  $\mathfrak{f}_{\mathrm{sh}}$  etc. as in the statement of Lemma 16.8. Moreover,  $\iota = n_1 \neq 0$ . Finally,  $t_0$  is such that (14.22) holds with  $t_a$  replaced by  $t_0$ . Thus the conditions of Lemma 15.11 are satisfied.

Let us begin by considering  $\Xi_{\text{fin}}^{k,+}$ . Due to Lemma 15.11, we know that the estimate (15.56) holds. Consider the right hand side. Note that  $\varepsilon_k$  is given by (16.26) with  $t_a$  replaced by  $t_k$ , and that  $\varepsilon_k \leq 1$  due to the definition of  $t_{\text{max}}$  and the fact that  $t_k < t_{\text{max}}$ . Keeping the assumptions of Lemma 16.8 in mind, (15.56) yields

$$\|\Xi_{\text{fin}}^{k,+} - \Xi_{\text{fin,app}}^{k,+}\| \le C_{\text{par}} \left(\frac{e^{(\beta_X + \beta_{\text{sh}})t_k}}{\mathfrak{g}(t_k)} + \frac{e^{2\beta_X t_k}}{\mathfrak{g}^2(t_k)}\right),\tag{16.47}$$

where  $C_{\text{par}}$  only depends on  $\mathbf{c}_{\text{par}}$ ; recall that  $\mathfrak{f}_{\text{sh}}(t) \leq C_{\text{sh}}$  for all  $t \geq 0$ . Next, let us turn to  $\Xi_{\text{fin,app}}^{k,+}$ . Due to (15.57), it is of interest to approximate  $R_{\text{pre},k}^{ab}$ , defined in (15.41). Due to (16.18), (16.20), (16.23) and (16.35),

$$\left\| R_{\text{pre},k}^{ab} - (-1)^b \frac{e^{\beta_X t_k}}{\mathfrak{g}(t_k)} Y_\infty^1 \right\| \le C_{\text{ep}} \frac{e^{(\beta_X - \eta_{\text{mn}})t_k}}{\mathfrak{g}(t_k)}$$
(16.48)

where we have used the fact that  $X = Y^1$  when  $n_1 > 0$ ,  $\eta_{mn}$  is defined by (16.44) and  $C_{ep}$  only depends on

$$\mathbf{c}_{\rm ep} := (\mathbf{c}_{\rm par}, \eta_X, K_X\}. \tag{16.49}$$

In the definition (16.44) of  $\eta_{\rm mn}$ , we could have excluded  $-\beta_{\rm sh}$ ; the estimate (16.48) would still hold. However, in order not to have to introduce additional notation later, we have included  $-\beta_{\rm sh}$ already here. Combining (16.47) and (16.48) with (15.57) and (16.36) yields

$$\|\Xi_{\text{fin}}^{k,+} - \Xi_{\text{app}}^{k,+}\| \le C_{\text{ep}}\left(\frac{e^{(\beta_X - \eta_{\text{mn}})t_k}}{\mathfrak{g}(t_k)} + \frac{e^{2\beta_X t_k}}{\mathfrak{g}^2(t_k)}\right),\tag{16.50}$$

where the constant  $C_{ep}$  only depends on  $\mathbf{c}_{ep}$  and

$$\Xi_{\mathrm{app}}^{k,+} := \mathrm{Id}_{2m} + \frac{2\pi e^{\beta_X t_k}}{\mathfrak{g}(t_k)} R_{\infty}.$$

Due to the definition of T and J, cf. (16.37), it is clear that

$$T\Xi_{\rm app}^{k,+}T^{-1} = \mathrm{Id}_{2m} + \frac{2\pi e^{\beta_X t_k}}{\mathfrak{g}(t_k)}J.$$
 (16.51)

Just as in the second reformulation of the equations in the proof of Lemma 11.17, it is of interest to eliminate the imaginary parts; this is the reason for introducing  $\rho_k$  defined in (16.39). It is also of interest to subtract the largest real part of the eigenvalues of J. This is the reason for introducing  $\hat{\rho}_k$  defined in (16.40). In order to estimate the effect of the transformation involved in defining, e.g.,  $\rho_k$ , let  $\lambda \in \mathbb{C}$  and consider

$$\exp\left(\beta_X^{-1}e^{\beta_X t_{k+1}}\lambda - \beta_X^{-1}e^{\beta_X t_k}\lambda\right) = \exp\left[\beta_X^{-1}e^{\beta_X t_k}\lambda(e^{\beta_X(t_{k+1}-t_k)}-1)\right].$$

Note that

$$\left|e^{\beta_X(t_{k+1}-t_k)} - 1 - \frac{2\pi\beta_X}{\mathfrak{g}(t_k)}\right| \le \frac{C}{\mathfrak{g}^2(t_k)},$$

where the constant C only depends on  $\beta_X$  and the constant  $\mathfrak{b}$  appearing in the estimate (14.1); cf. Lemma 14.4, in particular (14.24) and (14.27). Thus

$$\left|\beta_X^{-1}e^{\beta_X t_{k+1}}\lambda - \beta_X^{-1}e^{\beta_X t_k}\lambda - \frac{2\pi\lambda}{\mathfrak{g}(t_k)}e^{\beta_X t_k}\right| \le C\frac{e^{\beta_X t_k}}{\mathfrak{g}^2(t_k)}$$

where C only depends on  $\lambda$ ,  $\beta_X$  and the constant  $\mathfrak{b}$  appearing in the estimate (14.1). To conclude,

$$\left|\exp\left(\beta_X^{-1}e^{\beta_X t_{k+1}}\lambda - \beta_X^{-1}e^{\beta_X t_k}\lambda\right) - 1 - \frac{2\pi\lambda}{\mathfrak{g}(t_k)}e^{\beta_X t_k}\right| \le C\frac{e^{2\beta_X t_k}}{\mathfrak{g}^2(t_k)},\tag{16.52}$$

where C only depends on  $\lambda$ ,  $\beta_X$ , the constant  $\mathfrak{b}$  appearing in the estimate (14.1), and an upper bound on  $e^{\beta_X t_k}/\mathfrak{g}(t_k)$ . On the other hand, due to (16.30),  $e^{\beta_X t_k}/\mathfrak{g}(t_k) \leq 1$ . Thus C only depends on  $\lambda$ ,  $\beta_X$  and  $\mathfrak{b}$ . Defining  $S_k$  as in (16.38), the estimate (16.52) thus yields

$$\left\|S_{k+1}S_k^{-1} - \mathrm{Id}_{2m} + \frac{2\pi}{\mathfrak{g}(t_k)}e^{\beta_X t_k}\mathrm{diag}\{i\zeta_1\mathrm{Id}_{l_1}, \dots, i\zeta_r\mathrm{Id}_{l_r}\}\right\| \le C_{\mathrm{eep}}\frac{e^{2\beta_X t_k}}{\mathfrak{g}^2(t_k)},\tag{16.53}$$

where  $C_{eep}$  only depends on  $\mathbf{c}_{eep}$  introduced in (16.46). Let

$$B_{k,\text{app}} := S_{k+1}T\Xi_{\text{app}}^{k,+}T^{-1}S_k^{-1} = S_{k+1}\left(\text{Id}_{2m} + \frac{2\pi e^{\beta_X t_k}}{\mathfrak{g}(t_k)}J\right)S_k^{-1},$$

where the last equality is due to (16.51). Due to (16.53) and the fact that  $S_k$  and J commute,

$$\left\| B_{k,\mathrm{app}} - \mathrm{Id}_{2m} - \frac{2\pi e^{\beta_X t_k}}{\mathfrak{g}(t_k)} \mathrm{Re}\{J\} \right\| \le C_{\mathrm{eep}} \frac{e^{2\beta_X t_k}}{\mathfrak{g}^2(t_k)},$$

where  $C_{eep}$  only depends on  $\mathbf{c}_{eep}$ . Combining this estimate with (16.50) yields

$$\left\| B_k - \mathrm{Id}_{2m} - \frac{2\pi e^{\beta_X t_k}}{\mathfrak{g}(t_k)} \mathrm{Re}\{J\} \right\| \le C_{\mathrm{eep}} \left( \frac{e^{(\beta_X - \eta_{\mathrm{mn}})t_k}}{\mathfrak{g}(t_k)} + \frac{e^{2\beta_X t_k}}{\mathfrak{g}^2(t_k)} \right),$$

where  $C_{eep}$  only depends on  $\mathbf{c}_{eep}$ . Applying a similar argument a second time yields the conclusion of the lemma.

### 16.6 The main estimate

On a technical level, the main result of this chapter is Lemma 16.23; the proof of the existence of solutions exhibiting super exponential growth is based on it. The purpose of the present section is to prove this lemma. The main tool we use to obtain the desired result is the iteration (16.41) and the estimate (16.43). As a consequence, it is not surprising that the matrix  $J_n$ , introduced in (16.45), plays an important role. In the analysis to follow, it turns out to be convenient to divide  $J_n$  into a part consisting of the Jordan blocks of  $J_n$  with zero eigenvalue, say  $J_a$ , and a part consisting of the Jordan blocks of  $J_n$  with zero eigenvalue, say  $J_b$ . We can then, without loss of generality, assume that  $J_n = \text{diag}\{J_a, J_b\}$ . Let  $l_a$  and  $l_b$  be the dimensions of  $J_a$  and  $J_b$  respectively (note that  $l_b$  could equal zero, in which case  $J_n = J_a$ ). Before proceeding, it is convenient to introduce the following terminology.

**Definition 16.18.** If  $1 \leq l \in \mathbb{Z}$  and  $M \in \mathbf{M}_l(\mathbb{C})$ , then the matrices  $M_d, M_{od} \in \mathbf{M}_l(\mathbb{C})$  are defined by the following conditions:  $M_d$  is a diagonal matrix; the diagonal components of  $M_{od}$  are zero; and  $M = M_d + M_{od}$ .

**Remark 16.19.** If the matrix M of interest already has one index, say that it is given by  $M = J_n$ , then we write  $J_{n,d} := M_d$  and  $J_{n,od} := M_{od}$  e.g.

It is also useful to divide matrices into blocks according to the following definition.

**Definition 16.20.** Let  $0 \leq j_a, j_b \in \mathbb{Z}$  and  $1 \leq j \in \mathbb{Z}$  be such that  $j = j_a + j_b$ . If  $M \in \mathbf{M}_j(\mathbb{C})$ , it is divided into blocks  $M_{aa}, M_{ab}, M_{ba}$  and  $M_{bb}$  according to the following conditions:  $M_{aa} \in \mathbf{M}_{j_a}(\mathbb{C})$ ;  $M_{ab}$  is a complex  $j_a \times j_b$  matrix;  $M_{ba}$  is a complex  $j_b \times j_a$  matrix;  $M_{bb} \in \mathbf{M}_{j_b}(\mathbb{C})$ ; and

$$M = \left(\begin{array}{cc} M_{aa} & M_{ab} \\ M_{ba} & M_{bb} \end{array}\right)$$

Moreover, if  $\xi \in \mathbb{C}^{j}$ , then we denote the vector consisting of the first  $j_{a}$  and the vector consisting of the last  $j_{b}$  components of  $\xi$  by  $\xi_{a} \in \mathbb{C}^{j_{a}}$  and  $\xi_{b} \in \mathbb{C}^{j_{b}}$  respectively.

**Remarks 16.21.** When writing  $M_{aa}$  etc., the integers j,  $j_a$ ,  $j_b$  and the matrix M should be clear from the context. In practice, we are, in the present section, interested in the case that j = 2m,  $j_a = l_a$  and  $j_b = l_b$ , where  $l_a$  and  $l_b$  are defined immediately prior to Definition 16.18. Concerning vectors and matrices that already have an index, we use a convention similar to that introduced in Remark 16.19.

In the analysis below, we need to estimate the following type of sums.

**Lemma 16.22.** Assume that (1.2) is oscillation adapted, let  $0 \neq \iota \in \mathcal{I}_{B}$  and define  $\mathfrak{g}$  by (24.13). Assume that there is a  $t_0 \geq 0$  such that (14.22) holds with  $t_a$  replaced by  $t_0$ . Let the sequence  $\{t_k\}$  and  $k_{fin}$  be given by Definition 15.5. If  $0 \leq k_0 \leq k_1 \leq k_{fin}$  and  $\lambda \in \mathbb{R}$ , it then follows that

$$\sum_{k=k_0}^{k_1} \frac{e^{\lambda t_k}}{\mathfrak{g}(t_k)} \le C \int_{t_{k_0}}^{t_{k_1+1}} e^{\lambda t} dt$$

where C only depends on  $\lambda$ .

*Proof.* Due to (14.24),

$$\sum_{k=k_0}^{k_1} \frac{e^{\lambda t_k}}{\mathfrak{g}(t_k)} \le \frac{1}{\pi} \sum_{k=k_0}^{k_1} \int_{t_k}^{t_{k+1}} e^{\lambda t_k} dt.$$

On the other hand, for  $t \in [t_k, t_{k+1}]$ ,

$$e^{\lambda t_k} \le e^{\lambda t} e^{|\lambda(t_k - t)|} \le \exp\left(\frac{4\pi|\lambda|}{\max\{2, 4\pi\mathfrak{b}\}}\right) e^{\lambda t} \le e^{2\pi|\lambda|} e^{\lambda t},$$

where we have used (14.24) and the fact that  $\mathfrak{g}(t_k) \geq \max\{2, 4\pi\mathfrak{b}\}$ . The lemma follows.

We are now in a position to prove the main estimate of this section.

**Lemma 16.23.** Consider (1.2). Assume the associated metric to be such that (M, g) is a canonical separable cosmological model manifold. Assume, moreover, that f = 0, d = 1 and R = 0. Assume that the conditions of Lemma 16.8 are fulfilled with  $\beta_X > 0$  and that Assumptions 16.10 and 16.15 hold. Finally, assume that  $\kappa > 0$ , where  $\kappa$  is the largest real part of an eigenvalue of the matrix  $R_{\infty}$  defined by (16.36). Let  $\epsilon > 0$ . If  $\beta_X - \beta_1 > 0$ , there are constants  $K_{\text{eep},\epsilon}$ ,  $N_{\text{eep},\epsilon}$ ,  $T_{0,\epsilon} \geq 0$  and  $T_{\text{med},\epsilon} \geq T_{0,\epsilon}$ , depending only on  $\epsilon$  and  $\mathbf{c}_{\text{eep}}$ , where  $\mathbf{c}_{\text{eep}}$  is introduced in (16.46), and if  $\beta_X - \beta_1 \leq 0$ , there are constants  $N_{\text{eep},\epsilon}$ ,  $T_{0,\epsilon} \geq 0$  and  $T_{\text{med},\epsilon} \geq T_{0,\epsilon}$ , depending only on  $\epsilon$  and  $\mathbf{c}_{\text{eep}}$ , such that the following holds. Let  $n_1 \geq N_{\text{eep},\epsilon}$  and define  $T_{\max,\epsilon}$  by

$$T_{\max,\epsilon} := \begin{cases} T_{\exp,\epsilon}, & \beta_X - \beta_1 > 0\\ \infty, & \beta_X - \beta_1 \le 0 \end{cases},$$
(16.54)

where

$$T_{\text{eep},\epsilon} := \frac{1}{\beta_X - \beta_1} \ln |n_1| + K_{\text{eep},\epsilon}.$$
(16.55)

Then  $T_{\max,\epsilon} > T_{\text{med},\epsilon}$ ; (14.22) holds for  $0 \le t_a < T_{\max,\epsilon}$ ; and if  $\{t_k\}$  is the time sequence given by Definition 15.5 with  $t_0$  replaced by  $T_{0,\epsilon}$ , then there are solutions to (24.16) (corresponding to f = 0 and frequency  $n_1$ ) such that  $|w(T_{0,\epsilon})| = 1$  and such that

$$|w(t_{k+1})| \ge \exp\left[\beta_X^{-1}\kappa(1-\epsilon)e^{\beta_X t_{k+1}}\right]$$
(16.56)

for all k such that  $T_{\text{med},\epsilon} \leq t_k < T_{\max,\epsilon}$ ; recall that if z is a solution to (24.16), then w is defined by (14.7) and (15.1).

**Remark 16.24.** The constant  $K_{\text{eep},\epsilon}$  can be chosen in such a way that  $t_{\text{fin}}$ , given in Definition 15.5, satisfies  $t_{\text{fin}} \geq T_{\max,\epsilon}$ ; cf. Remark 16.14. Moreover, since  $T_{\text{med},\epsilon}$  is fixed (depending only on  $\epsilon$  and  $\mathbf{c}_{\text{eep}}$ ), it is clear that by choosing  $n_1$  to be large enough, the interval  $[T_{\text{med},\epsilon}, T_{\max,\epsilon}]$  can be assumed to contain a large number of elements of the sequence  $\{t_k\}$ .

**Remark 16.25.** The condition  $t \ge T_{0,\epsilon}$  ensures that we are in the asymptotic regime. The time  $T_{\text{med},\epsilon}$  is chosen so that significant growth can take place in the interval  $[T_{0,\epsilon}, T_{\text{med},\epsilon}]$ .

**Remark 16.26.** In terms of the solution z to (24.16), the estimate (16.56) can be written

$$[|\dot{z}(t_{k+1})|^2 + \mathfrak{g}^2(t_{k+1})|z(t_{k+1})|^2]^{1/2} \\ \geq \exp\left[\beta_X^{-1}\kappa(1-\epsilon)e^{\beta_X t_{k+1}}\right] [|\dot{z}(T_{0,\epsilon})|^2 + \mathfrak{g}^2(T_{0,\epsilon})|z(T_{0,\epsilon})|^2]^{1/2},$$

where we have used the fact that  $|w(T_{0,\epsilon})| = 1$  as well as (14.7) and (15.1).

*Proof.* Before turning to the actual proof, it is useful to make a few remarks concerning the constants and parameters appearing in it. In the course of the proof, we bound  $K_{\text{eep},\epsilon}$  from above by conditions that are gradually more restrictive. Similarly, we bound  $N_{\text{eep},\epsilon}$ ,  $T_{0,\epsilon} \ge 0$  and  $T_{\text{med},\epsilon} \ge T_{0,\epsilon}$  from below by conditions that are gradually more restrictive. On the other hand, all of these constants depend only on  $\epsilon$  and  $\mathbf{c}_{\text{eep}}$ . By demanding that  $n_1$  be large enough, it can thus be ensured that  $T_{\max,\epsilon}$  is as much larger than  $T_{0,\epsilon}$  and  $T_{\text{med},\epsilon}$  as we wish.

Turning to the parameter  $\epsilon$ , it is convenient to introduce it only at the end of the proof. To begin with, we instead introduce two parameters  $0 < \epsilon_j \leq 1, j = 1, 2$ . In the end, we describe how the  $\epsilon_j$  are determined by  $\epsilon$  and  $\mathbf{c}_{eep}$ .

In order to obtain a clearer picture of the restrictions on the constants  $K_{\text{eep},\epsilon}$  etc., we proceed step by step, and conclude each step with a description of the restrictions that need to be imposed.

Step 1, dealing with the presence of non-trivial Jordan blocks. If one of the Jordan blocks in  $R_{\infty}$  corresponding to an eigenvalue with real part  $\kappa$  is non-trivial, it is convenient to reformulate the relation (16.41). The reason for this is that we wish to control the influence of

the non-trivial nature of such Jordan blocks on the evolution. In order to quantify this control, we use the parameter  $\epsilon_1$ .

Let  $0 < \epsilon_1 \leq 1$  and recall that  $J_n$  is defined in (16.45). By conjugating  $J_n$  by a suitable diagonal matrix, say  $T_1$ , it can be ensured that

$$\check{J}_{n} := T_{1} J_{n} T_{1}^{-1}$$

$$\|\check{J}_{n,od}\| \le \epsilon_{1},$$
(16.5)

is such that

$$|\check{J}_{n,od}\| \le \epsilon_1, \tag{16.57}$$

where the terminology  $J_{n,od}$  is introduced in Definition 16.18; cf. Lemma 9.9 and its proof. Note, however, that the norms of  $T_1$  and  $T_1^{-1}$  only depend on  $\epsilon_1$ . Define the matrices  $\check{J}_a$  and  $\check{J}_b$  by  $\check{J}_a := \check{J}_{n,aa}$  and  $\check{J}_b := \check{J}_{n,bb}$ , where we use the notation introduced in Definition 16.20 (with  $j_a = l_a, j_b = l_b$  and j = 2m). Then (16.57) yields

$$\|\check{J}_{a,\text{od}}\| \le \|\check{J}_{n,\text{od}}\| \le \epsilon_1, \quad \|\check{J}_{b,\text{od}}\| \le \|\check{J}_{n,\text{od}}\| \le \epsilon_1.$$
 (16.58)

Let  $\nu_{\text{par}}$  and  $C_{\text{par}}$  be defined as in Lemma 16.17. For  $|n_1| \ge \nu_{\text{par}}$ , let  $t_{\text{max}}$  be defined by (16.42). Assume that  $0 \le t_0 < t_{\text{max}}$  has been specified (in the present context,  $t_0$  remains to be determined) and let the sequence  $\{t_k\}$  be given as in Definition 15.5. Then, for  $k \ge 0$  and  $t_k < t_{\max}$  we can appeal to (16.43). Moreover, introducing

$$\check{\rho}_k := T_1 \hat{\rho}_k, \quad \check{B}_k := T_1 \hat{B}_k T_1^{-1}, \quad \check{B}_{k,\text{app}} := T_1 \hat{B}_{k,\text{app}} T_1^{-1},$$
(16.59)

(16.43) yields

$$\left\|\check{B}_{k}-\check{B}_{k,\mathrm{app}}\right\| \leq C_{\mathrm{eep},\epsilon_{1}}\left(\frac{e^{(\beta_{X}-\eta_{\mathrm{mn}})t_{k}}}{\mathfrak{g}(t_{k})}+\frac{e^{2\beta_{X}t_{k}}}{\mathfrak{g}^{2}(t_{k})}\right),\tag{16.60}$$

where  $C_{eep,\epsilon_1}$  only depends on  $\mathbf{c}_{eep}$  defined in (16.46) and  $\epsilon_1$ . In addition,

$$\check{B}_{k,\mathrm{app}} = \mathrm{Id}_{2m} + \frac{2\pi e^{\beta_X t_k}}{\mathfrak{g}(t_k)}\check{J}_{\mathrm{n}}.$$
(16.61)

Finally, (16.41) yields

$$\check{\rho}_{k+1} = B_k \check{\rho}_k. \tag{16.62}$$

Define  $\dot{R}_k$  by

$$\check{B}_k = \check{B}_{k,\mathrm{app}} + \check{R}_k.$$

Then  $||R_k||$  is bounded by the right hand side of (16.60). Using the notation introduced in Definition 16.20 (with  $j_a = l_a$ ,  $j_b = l_b$  and j = 2m), (16.62) can then be written

$$\check{\rho}_{k+1,a} = \check{B}_{k,\text{app},aa}\check{\rho}_{k,a} + (\dot{R}_k\check{\rho}_k)_a, \tag{16.63}$$

$$\check{\rho}_{k+1,b} = \dot{B}_{k,\text{app},bb}\check{\rho}_{k,b} + (\dot{R}_k\check{\rho}_k)_b.$$
(16.64)

**Restrictions, step 1.** Assume that  $T_{0,\epsilon} \geq 0$ ,  $K_{eep,\epsilon} \leq C_{par}$  (in case  $\beta_X - \beta_1 > 0$ ) and  $N_{eep,\epsilon} \geq 0$  $\nu_{\text{par}}$ . Fix  $n_1$  satisfying  $|n_1| \ge N_{\text{eep},\epsilon}$  and define  $T_{\max,\epsilon}$  by (16.54), where  $T_{\text{eep},\epsilon}$  is given by (16.55) in case  $\beta_X - \beta_1 > 0$ . Construct the sequence  $\{t_k\}$  as in Definition 15.5, where the  $t_0$  satisfying  $0 \le t_0 < T_{\max,\epsilon}$  is arbitrary. Then the conclusions of step 1 hold for  $|n_1| \ge N_{\text{eep},\epsilon}$  and  $t_0 \le t_k < \infty$  $T_{\max,\epsilon}$ . Finally, let us remark that if  $K_{eep,\epsilon}$  and  $N_{eep,\epsilon}$  are chosen inappropriately, then  $T_{\max,\epsilon}$ might be negative. However, for a given choice of  $K_{eep,\epsilon}$  (depending only on the  $\epsilon_j$ 's,  $\epsilon$  and  $\mathbf{c}_{eep}$ ), there is a choice of  $N_{\text{eep},\epsilon} \geq \nu_{\text{par}}$  with the same dependence, such that  $T_{\max,\epsilon} > 0$  if  $|n_1| \geq N_{\text{eep},\epsilon}$ .

Step 2, limiting the time sequence. In order to be able to proceed, we need to obtain better control of the right hand side of (16.60). There are two aspects to this. First, we want  $e^{\beta_X t_k}/\mathfrak{g}(t_k)$ to be small. This can be ensured by appealing to Lemma 16.12. Second, we want  $e^{-\eta_{\rm mn}t_k}$  to be small. This can be ensured by assuming that  $t_k$  is sufficiently large. In order to quantify the control, we use the parameter  $\epsilon_2$ .

Let  $0 < \epsilon_2 \leq 1$ . Then Lemma 16.12 ensures the following. If  $\beta_X - \beta_1 > 0$ , then there are constants  $\nu_{\text{par},\epsilon_2}$  and  $C_{\text{par},\epsilon_2}$ , depending only on  $\mathbf{c}_{\text{par}}$  and  $\epsilon_2$ , such that if  $|n_1| \geq \nu_{\text{par},\epsilon_2}$ , then (16.29) holds (with  $\epsilon$  replaced by  $\epsilon_2$ ), and (16.30) holds for  $0 \leq t, t_a \leq T_{\text{par},\epsilon_2}$  (with  $\epsilon$  replaced by  $\epsilon_2$ ). Similarly, if  $\beta_X - \beta_1 \leq 0$ , there is a constant  $\nu_{\text{par},\epsilon_2}$ , depending only on  $\mathbf{c}_{\text{par}}$  and  $\epsilon_2$ , such that if  $|n_1| \geq \nu_{\text{par},\epsilon_2}$ , then (16.30) holds for  $0 \leq t, t_a \leq T_{\text{par},\epsilon_2}$  (with  $\epsilon$  replaced by  $\epsilon_2$ ). Similarly, if  $\beta_X - \beta_1 \leq 0$ , there is a constant  $\nu_{\text{par},\epsilon_2}$ , depending only on  $\mathbf{c}_{\text{par}}$  and  $\epsilon_2$ , such that if  $|n_1| \geq \nu_{\text{par},\epsilon_2}$ , then (16.30) holds for  $0 \leq t, t_a < \infty$  (with  $\epsilon$  replaced by  $\epsilon_2$ ). Defining

$$t_{\max,\epsilon_2} := \begin{cases} T_{\max,\epsilon_2}, & \beta_X - \beta_1 > 0\\ \infty, & \beta_X - \beta_1 \le 0 \end{cases}$$
(16.65)

yields the conclusion that (16.30) (with  $\epsilon$  replaced by  $\epsilon_2$ ) holds for  $0 \leq t, t_a < t_{\max,\epsilon_2}$ . Here we take for granted that  $\nu_{\text{par},\epsilon_2} \geq \nu_{\text{par}}$  and  $C_{\text{par},\epsilon_2} \leq C_{\text{par}}$ , where  $\nu_{\text{par}}$  and  $C_{\text{par}}$  are the constants introduced in Lemma 16.17. As a consequence,  $t_{\max,\epsilon_2} \leq t_{\max}$ . Before turning to specific bounds we wish to impose on  $\epsilon_2$ , let us introduce the notation  $\kappa_{n,\max}$  for the largest eigenvalue of  $-J_n$ . Let, moreover,  $\kappa_n$  be the smallest strictly positive eigenvalue of  $-J_n$ . It is of course possible that there is no strictly positive eigenvalue of  $-J_n$ . However, this case corresponds to  $l_b = 0$ , and only simplifies the analysis; we leave it to the reader to modify the analysis presented below in order to deal with the case  $l_b = 0$ . In what follows, we always assume that

$$\epsilon_2, \epsilon_1 \le \frac{1}{2\pi \max\{\kappa_{n,\max}, 1\}}.$$
(16.66)

One advantage of making this assumption is that, if it holds, then

$$\left\| \operatorname{Id}_{l_b} + \frac{2\pi e^{\beta_X t_k}}{\mathfrak{g}(t_k)} \check{J}_b \right\| \le 1 - \frac{2\pi e^{\beta_X t_k}}{\mathfrak{g}(t_k)} (\kappa_n - \epsilon_1)$$
(16.67)

for  $t_0 \leq t_k < t_{\max,\epsilon_2}$ , where we have appealed to (16.30) (with  $\epsilon$  replaced by  $\epsilon_2$ ) and (16.58). For future reference, it is also of interest to note that

$$1 - \frac{2\pi e^{\beta_X t_k}}{\mathfrak{g}(t_k)} \epsilon_1 \le \left| \left( \mathrm{Id}_{l_a} + \frac{2\pi e^{\beta_X t_k}}{\mathfrak{g}(t_k)} \check{J}_a \right) \xi \right| \le 1 + \frac{2\pi e^{\beta_X t_k}}{\mathfrak{g}(t_k)} \epsilon_1$$
(16.68)

for  $\xi \in \mathbb{C}^{l_a}$  such that  $|\xi| = 1$ .

Fix  $0 < \epsilon_1 \le 1$  satisfying (16.66), and define  $T_1$ ,  $\check{\rho}_k$  etc. as above. Assume that  $0 < \epsilon_2 \le 1$  satisfies (16.66) and

$$C_{\text{eep},\epsilon_1}\epsilon_2 \le \pi\epsilon_1,\tag{16.69}$$

where  $C_{eep,\epsilon_1}$  is the constant appearing on the right hand side of (16.60). Then

$$\|\check{R}_k\| \le C_{\text{eep},\epsilon_1} e^{-\eta_{\text{mn}}t_k} \frac{e^{\beta_X t_k}}{\mathfrak{g}(t_k)} + \pi \epsilon_1 \frac{e^{\beta_X t_k}}{\mathfrak{g}(t_k)}$$
(16.70)

for all k such that  $t_0 \leq t_k < t_{\max,\epsilon_2}$ , where we have appealed to (16.30) (with  $\epsilon$  replaced by  $\epsilon_2$ ) and the fact that  $||\check{R}_k||$  is bounded by the right hand side of (16.60). In (16.70), the factor  $\pi$  in the second term on the right hand side has been inserted for convenience. In order to improve the first term on the right of (16.70), it is necessary to impose a lower bound on  $t_k$ . To this end, let  $t_{\min,\epsilon_1} \geq 0$ , depending only on  $\mathbf{c}_{eep}$  and  $\epsilon_1$ , be such that

$$C_{\text{eep},\epsilon_1} e^{-\eta_{\text{mn}}t} \le \pi \epsilon_1 \tag{16.71}$$

for  $t \ge t_{\min,\epsilon_1}$ , where  $C_{\text{eep},\epsilon_1}$  is the specific constant appearing on the right hand side of (16.70). Construct the sequence  $\{t_k\}$  as in Definition 15.5 but with  $t_{\min,\epsilon_1} \le t_0 < t_{\max,\epsilon_2}$ . Then, for  $t_0 \le t_k < t_{\max,\epsilon_2}$ ,

$$\|\check{R}_k\| \le 2\pi\epsilon_1 \frac{e^{\beta_X t_k}}{\mathfrak{g}(t_k)}.$$
(16.72)

**Restrictions, step 2.** Fix  $0 < \epsilon_1 \le 1$  satisfying (16.66), and fix  $0 < \epsilon_2 \le 1$  satisfying (16.66) and (16.69). Let  $\nu_{\text{par},\epsilon_2} \ge \nu_{\text{par}}$  and  $C_{\text{par},\epsilon_2} \le C_{\text{par}}$  be defined as above; cf. the text adjacent to (16.65).

Finally, let  $t_{\min,\epsilon_1} \geq 0$  be such that (16.71) holds for  $t \geq t_{\min,\epsilon_1}$ . Assume that  $T_{0,\epsilon} \geq t_{\min,\epsilon_1}$ ,  $K_{\text{eep},\epsilon} \leq C_{\text{par},\epsilon_2}$  (in case  $\beta_X - \beta_1 > 0$ ) and  $N_{\text{eep},\epsilon} \geq \nu_{\text{par},\epsilon_2}$ . Fix  $n_1$  satisfying  $|n_1| \geq N_{\text{eep},\epsilon}$  and define  $T_{\max,\epsilon}$  by (16.54), where  $T_{\text{eep},\epsilon}$  is given by (16.55) in case  $\beta_X - \beta_1 > 0$ . Construct the sequence  $\{t_k\}$  as in Definition 15.5, where  $T_{0,\epsilon} \leq t_0 < T_{\max,\epsilon}$  is arbitrary. Then the conclusions of steps 1 and 2 hold for  $t_0 \leq t_k < T_{\max,\epsilon}$ . Finally, note that if  $K_{\text{eep},\epsilon}$  and  $N_{\text{eep},\epsilon}$  are chosen inappropriately, then  $T_{\max,\epsilon}$  might be less than or equal to  $T_{0,\epsilon}$ . However, for a given choice of  $K_{\text{eep},\epsilon}$  and  $T_{0,\epsilon}$  (depending only on the  $\epsilon_j$ 's,  $\epsilon$  and  $\mathbf{c}_{\text{eep}}$ ), there is a choice of  $N_{\text{eep},\epsilon} \geq \nu_{\text{par}}$  with the same dependence, such that  $T_{\max,\epsilon} > T_{0,\epsilon}$  if  $|n_1| \geq N_{\text{eep},\epsilon}$ .

Step 3, controlling the components of the solutions to the difference equation. Assume that the restrictions arising in step 2 are satisfied and let  $t_0$  etc. satisfy the corresponding conditions. Let us return to (16.63) and (16.64). Due to (16.63),

$$\begin{split} |\check{\rho}_{k+1,a}| \geq |\check{B}_{k,\mathrm{app},aa}\check{\rho}_{k,a}| - |(\check{R}_{k}\check{\rho}_{k})_{a}| \\ \geq |\check{B}_{k,\mathrm{app},aa}\check{\rho}_{k,a}| - \epsilon_{1}\frac{2\pi e^{\beta_{X}t_{k}}}{\mathfrak{g}(t_{k})}|\check{\rho}_{k}| \\ \geq |\check{\rho}_{k,a}| - \epsilon_{1}\frac{4\pi e^{\beta_{X}t_{k}}}{\mathfrak{g}(t_{k})}(|\check{\rho}_{k,a}| + |\check{\rho}_{k,b}|), \end{split}$$
(16.73)

where we have appealed to (16.68) and (16.72); from now on we always tacitly assume that k is such that  $T_{0,\epsilon} \leq t_0 \leq t_k < T_{\max,\epsilon}$ . Similarly, (16.64) yields

$$\begin{split} |\check{\rho}_{k+1,b}| \leq &|\check{\rho}_{k,b}| - \frac{2\pi e^{\beta_X t_k}}{\mathfrak{g}(t_k)} (\kappa_n - \epsilon_1) |\check{\rho}_{k,b}| + \epsilon_1 \frac{2\pi e^{\beta_X t_k}}{\mathfrak{g}(t_k)} |\check{\rho}_k| \\ \leq & \left( 1 - \kappa_n \frac{2\pi e^{\beta_X t_k}}{\mathfrak{g}(t_k)} \right) |\check{\rho}_{k,b}| + \epsilon_1 \frac{4\pi e^{\beta_X t_k}}{\mathfrak{g}(t_k)} (|\check{\rho}_{k,b}| + |\check{\rho}_{k,a}|), \end{split}$$
(16.74)

where we have appealed to (16.67) and (16.72). In particular, (16.73) and (16.74) yield

$$\frac{|\check{\rho}_{k+1,a}|}{|\check{\rho}_{k,a}|} \ge 1 - \epsilon_1 \frac{4\pi e^{\beta_X t_k}}{\mathfrak{g}(t_k)} \left(1 + \frac{|\check{\rho}_{k,b}|}{|\check{\rho}_{k,a}|}\right),\tag{16.75}$$

$$\frac{|\check{\rho}_{k+1,b}|}{|\check{\rho}_{k,a}|} \le \left(1 - \kappa_{n} \frac{2\pi e^{\beta_{X} t_{k}}}{\mathfrak{g}(t_{k})}\right) \frac{|\check{\rho}_{k,b}|}{|\check{\rho}_{k,a}|} + \epsilon_{1} \frac{4\pi e^{\beta_{X} t_{k}}}{\mathfrak{g}(t_{k})} \left(1 + \frac{|\check{\rho}_{k,b}|}{|\check{\rho}_{k,a}|}\right).$$
(16.76)

Assuming  $\epsilon_2$  to be small enough ( $\epsilon_2 \leq c_{0,1}$ , where  $c_{0,1} > 0$  is a numerical constant) and the inequality

$$\frac{|\check{\rho}_{k,b}|}{|\check{\rho}_{k,a}|} \le 1 \tag{16.77}$$

to hold, the absolute value of the second term on the right hand side of (16.75) is bounded from above by 1/4. Under these assumptions, (16.75) yields

$$\frac{|\check{\rho}_{k,a}|}{|\check{\rho}_{k+1,a}|} \le 1 + \epsilon_1 \frac{8\pi e^{\beta_X t_k}}{\mathfrak{g}(t_k)} \left(1 + \frac{|\check{\rho}_{k,b}|}{|\check{\rho}_{k,a}|}\right).$$
(16.78)

Thus, if (16.77) holds and  $\epsilon_2 \leq c_{0,1}$ , then (16.76) and (16.78) imply that

$$\frac{|\check{\rho}_{k+1,b}|}{|\check{\rho}_{k+1,a}|} \le \left(1 - \kappa_n \frac{2\pi e^{\beta_X t_k}}{\mathfrak{g}(t_k)}\right) \frac{|\check{\rho}_{k,b}|}{|\check{\rho}_{k,a}|} + c_{0,2}\epsilon_1 \frac{2\pi e^{\beta_X t_k}}{\mathfrak{g}(t_k)},\tag{16.79}$$

where  $c_{0,2} \ge 1$  is a numerical constant. Let us now assume that  $\epsilon_1$  has been chosen so that

$$\epsilon_1 \le \frac{1}{c_{0,2}} \min\left\{\frac{\kappa_n}{4}, \frac{1}{2}\right\}.$$
(16.80)

It is of interest to draw the following two conclusions from (16.79), given that (16.80) is fulfilled. To begin with,

$$\frac{|\check{\rho}_{k+1,b}|}{|\check{\rho}_{k+1,a}|} \le \frac{|\check{\rho}_{k,b}|}{|\check{\rho}_{k,a}|} + \frac{1}{2},$$

where we have used the fact that the first factor in the first term on the right hand side of (16.79) is  $\leq 1$ ; the fact that (16.66) holds; and the fact that (16.30) (with  $\epsilon$  replaced by  $\epsilon_2$ ) holds for  $t = t_k$ . In particular, we obtain the following two conclusions.

Small quotient initially. If  $\epsilon_2 \leq c_{0,1}$ ,  $|\check{\rho}_{k,b}|/|\check{\rho}_{k,a}| \leq 1/2$  and (16.80) holds, then  $|\check{\rho}_{k+1,b}|/|\check{\rho}_{k+1,a}| \leq 1$ . In other words,  $|\check{\rho}_{k,b}|/|\check{\rho}_{k,a}|$  cannot jump from being less than or equal to 1/2 to being strictly larger than 1.

Intermediate quotient initially. Assume now that  $\epsilon_2 \leq c_{0,1}$ ,  $1/2 \leq |\check{\rho}_{k,b}|/|\check{\rho}_{k,a}| \leq 1$  and (16.80) hold. Then

$$\begin{aligned} \frac{|\check{\rho}_{k+1,b}|}{|\check{\rho}_{k+1,a}|} &\leq \left(1 - \kappa_{n} \frac{2\pi e^{\beta_{X}t_{k}}}{\mathfrak{g}(t_{k})}\right) \frac{|\check{\rho}_{k,b}|}{|\check{\rho}_{k,a}|} + \frac{\kappa_{n}}{4} \frac{2\pi e^{\beta_{X}t_{k}}}{\mathfrak{g}(t_{k})} \\ &\leq \left(1 - \kappa_{n} \frac{2\pi e^{\beta_{X}t_{k}}}{\mathfrak{g}(t_{k})}\right) \frac{|\check{\rho}_{k,b}|}{|\check{\rho}_{k,a}|} + \frac{\kappa_{n}}{2} \frac{2\pi e^{\beta_{X}t_{k}}}{\mathfrak{g}(t_{k})} \frac{|\check{\rho}_{k,b}|}{|\check{\rho}_{k,a}|} < \frac{|\check{\rho}_{k,b}|}{|\check{\rho}_{k,a}|}, \end{aligned}$$

where we used (16.79) and (16.80) in the first step and  $|\check{\rho}_{k,b}|/|\check{\rho}_{k,a}| \ge 1/2$  in the second step. In other words, if  $1/2 \le |\check{\rho}_{k,b}|/|\check{\rho}_{k,a}| \le 1$ , then  $|\check{\rho}_{k+1,b}|/|\check{\rho}_{k+1,a}| < |\check{\rho}_{k,b}|/|\check{\rho}_{k,a}|$ .

Combining these two observations, we obtain the following conclusion. If  $\epsilon_2 \leq c_{0,1}$ , (16.80) and  $|\check{\rho}_{0,b}|/|\check{\rho}_{0,a}| \leq 1/2$  hold, then  $|\check{\rho}_{k,b}|/|\check{\rho}_{k,a}| \leq 1$  for all  $k \geq 0$  such that  $t_k < T_{\max,\epsilon}$ .

**Restrictions, step 3.** Fix  $0 < \epsilon_1 \leq 1$  satisfying (16.66) and (16.80), and fix  $0 < \epsilon_2 \leq 1$  satisfying  $\epsilon_2 \leq c_{0,1}$ , (16.66) and (16.69). Given these parameters, impose restrictions as in step 2. Then the conclusions of steps 1, 2 and 3 hold. Moreover, if  $|\check{\rho}_{0,b}|/|\check{\rho}_{0,a}| \leq 1/2$  holds, then  $|\check{\rho}_{k,b}|/|\check{\rho}_{k,a}| \leq 1$  for all  $k \geq 0$  such that  $t_k < T_{\max,\epsilon}$ .

Step 4, growth estimate. Assume that the restrictions arising in step 3 are satisfied and let  $t_0$  etc. satisfy the corresponding conditions. Let us return to (16.73). Given that  $|\check{\rho}_{k,b}|/|\check{\rho}_{k,a}| \leq 1$ , this estimate yields

$$|\check{\rho}_{k+1,a}| \ge \left(1 - \epsilon_1 \frac{8\pi e^{\beta_X t_k}}{\mathfrak{g}(t_k)}\right) |\check{\rho}_{k,a}| \tag{16.81}$$

for  $k \ge 0$  such that  $t_k < T_{\max,\epsilon}$ . Note that when  $l_b = 0$ , the estimate (16.81) is an immediate consequence of (16.73); the issue of the relative sizes of  $|\check{\rho}_{k,b}|$  and  $|\check{\rho}_{k,a}|$  does not arise in that case (needless to say).

Before proceeding, note that  $\ln(1-x) + 2x \ge 0$  for  $0 \le x \le 1/2$ . Assuming  $\epsilon_1$  to be such that  $\epsilon_1 \le 1/8$ , (16.66) yields

$$\epsilon_1 \frac{8\pi e^{\beta_X t_k}}{\mathfrak{g}(t_k)} \le \frac{1}{2}.$$
(16.82)

Combining these observations yields

$$\ln\left[\left(1-\epsilon_1\frac{8\pi e^{\beta_X t_k}}{\mathfrak{g}(t_k)}\right)\cdots\left(1-\epsilon_1\frac{8\pi e^{\beta_X t_0}}{\mathfrak{g}(t_0)}\right)\right]$$
$$\geq -\epsilon_1\sum_{l=0}^k\frac{16\pi e^{\beta_X t_l}}{\mathfrak{g}(t_l)}\geq -C_X\epsilon_1e^{\beta_X t_{k+1}},$$

where we have appealed to Lemma 16.22, and the constant  $C_X$  only depends on  $\beta_X$ . Combining this estimate with (16.81) yields

$$|\check{\rho}_{k+1,a}| \ge \exp\left[-C_X \epsilon_1 e^{\beta_X t_{k+1}}\right] |\check{\rho}_{0,a}|,$$

228

assuming that  $|\check{\rho}_{l,b}|/|\check{\rho}_{l,a}| \leq 1$  for l = 0, ..., k. Choosing  $\check{\rho}_{0,a}$  and  $\check{\rho}_{0,b}$  so that  $|\check{\rho}_{0,a}| > 0$  and  $\check{\rho}_{0,b} = 0$ , it is then clear that  $|\check{\rho}_{k,b}|/|\check{\rho}_{k,a}| \leq 1$  for all  $k \geq 0$  such that  $t_k < T_{\max,\epsilon}$ . Then

$$|\check{\rho}_{k+1}| \ge \exp\left[-C_X \epsilon_1 e^{\beta_X t_{k+1}}\right] |\check{\rho}_0|$$

for all  $k \ge 0$  such that  $t_k < T_{\max,\epsilon}$ . Returning to the definition of  $\check{\rho}_k$  in terms of w (cf., in particular, (15.36), (15.53), (16.39), (16.40) and (16.59)), this estimate yields

$$|w(t_{k+1})| \ge C_{\text{eep},\epsilon_1} \exp\left[\beta_X^{-1} \kappa(e^{\beta_X t_{k+1}} - e^{\beta_X t_0}) - C_X \epsilon_1 e^{\beta_X t_{k+1}}\right] |w(t_0)|$$
(16.83)

for all  $k \ge 0$  such that  $t_k < T_{\max,\epsilon}$ , where  $C_{eep,\epsilon_1} > 0$  only depends on  $\epsilon_1$  and  $\mathbf{c}_{eep}$ . Note, moreover, that

$$|w(t_k)| = [|\dot{z}(t_k)|^2 + \mathfrak{g}^2(t_k)|z(t_k)|^2]^{1/2}.$$

**Restrictions, step 4.** Fix  $0 < \epsilon_1 \leq 1/8$  satisfying (16.66) and (16.80), and fix  $0 < \epsilon_2 \leq 1$  satisfying  $\epsilon_2 \leq c_{0,1}$ , (16.66) and (16.69). Given these parameters, impose restrictions as in step 2. Assume, moreover, that  $|\check{\rho}_{0,b}| = 0$  and  $|\check{\rho}_{0,a}| > 0$ . Then the conclusions of steps 1-4 hold.

Step 5, summing up. Let  $0 < \epsilon \leq 1$ . Let, moreover,  $C_X$  be the constant appearing in (16.83) (depending only on  $\beta_X$ ). Let  $\epsilon_1 > 0$  be the largest constant satisfying  $\epsilon_1 \leq 1/8$ , (16.66), (16.80) and

$$C_X \epsilon_1 \le \frac{\epsilon}{2} \beta_X^{-1} \kappa. \tag{16.84}$$

Recall that  $c_{0,2}$  appearing in (16.80) is a numerical constant and that  $\epsilon_1 \leq 1/8$  ensures that (16.82) holds. In particular, it is thus clear that  $\epsilon_1 > 0$  only depends on  $\epsilon$  and  $\mathbf{c}_{eep}$ . Next, fix  $0 < \epsilon_2 \leq 1$  to be the largest constant such that (16.66) and (16.69) hold, and such that  $\epsilon \leq c_{0,1}$ , where  $c_{0,1}$  is the numerical constant introduced in connection with (16.77). Again,  $\epsilon_2 > 0$  only depends on  $\epsilon$  and  $\mathbf{c}_{eep}$ . Let  $\nu_{par,\epsilon_2} \geq \nu_{par}$  and  $C_{par,\epsilon_2} \leq C_{par}$  be defined as in the text adjacent to (16.65), and define  $t_{\min,\epsilon_1}$  to be the smallest non-negative real number such that (16.71) holds for  $t \geq t_{\min,\epsilon_1}$ . Then  $\nu_{par,\epsilon_2}$ ,  $C_{par,\epsilon_2}$  and  $t_{\min,\epsilon_1}$  only depend on  $\epsilon$  and  $\mathbf{c}_{eep}$ . Let  $K_{eep,\epsilon} := C_{par,\epsilon_2}$  (in case  $\beta_X - \beta_1 > 0$ ) and  $T_{0,\epsilon} := t_{\min,\epsilon_1}$ . Choose, moreover,  $T_{med,\epsilon}$  to be the smallest number such that  $T_{med,\epsilon} \geq T_{0,\epsilon}$ and such that

$$-\ln C_{\operatorname{eep},\epsilon_1} + \beta_X^{-1} \kappa e^{\beta_X T_{0,\epsilon}} \le \frac{\epsilon}{2} \beta_X^{-1} \kappa e^{\beta_X t}$$
(16.85)

for  $t \geq T_{\text{med},\epsilon}$ , where  $C_{\text{eep},\epsilon_1}$  is the constant appearing on the right hand side of (16.83). Note that  $T_{\text{med},\epsilon}$  only depends on  $\epsilon$  and  $\mathbf{c}_{\text{eep}}$ . Finally, let  $N_{\text{eep},\epsilon}$  be the smallest number which is larger than  $\nu_{\text{par},\epsilon_2}$  and such that if  $|n_1| \geq N_{\text{eep},\epsilon}$ , then  $T_{\max,\epsilon}$ , defined by (16.54) (where  $T_{\text{eep},\epsilon}$  is given by (16.55) in case  $\beta_X - \beta_1 > 0$ ), is such that  $T_{\text{med},\epsilon} + 2\pi \leq T_{\max,\epsilon}$ . Then  $N_{\text{eep},\epsilon}$  only depends on  $\epsilon$  and  $\mathbf{c}_{\text{eep}}$ . Finally, let  $t_0 := T_{0,\epsilon}$  and construct the sequence  $\{t_k\}$  as in Definition 15.5.

Consider a solution z to (24.16) such that  $|\check{\rho}_{0,b}| = 0$  and  $|\check{\rho}_{0,a}| > 0$ . Then, since  $\epsilon_1$  and  $\epsilon_2$  satisfy the restrictions imposed in step 4 and  $T_{0,\epsilon}$ ,  $K_{eep}$  and  $N_{eep,\epsilon}$  satisfy the restrictions imposed in step 2, the conclusions of steps 1-4 hold. If  $T_{\text{med},\epsilon} \leq t_k < T_{\max,\epsilon}$ , then (16.83), (16.84) and (16.85) yield

$$|w(t_{k+1})| \ge \exp\left[\beta_X^{-1}\kappa(1-\epsilon)e^{\beta_X t_{k+1}}\right]|w(t_0)|.$$

The lemma follows.

#### 16.7 Lower bounds on the growth of the energy

In the present section, we construct solutions to (1.2) exhibiting super exponential growth, given that the conditions stated in Lemma 16.8 and Assumptions 16.10 and 16.15 are satisfied.

**Lemma 16.27.** Consider (1.2). Assume the associated metric to be such that (M, g) is a canonical separable cosmological model manifold. Assume, moreover, that f = 0, d = 1 and R = 0. Assume that the conditions of Lemma 16.8 are fulfilled with  $\beta_X > 0$  and that Assumptions 16.10 and 16.15

hold. Finally, assume that  $\kappa > 0$ , where  $\kappa$  is the largest real part of an eigenvalue of the matrix  $R_{\infty}$  defined by (16.36). Let  $\epsilon > 0$ . Then there is a sequence of smooth  $\mathbb{C}^m$ -valued solutions  $v_l$  to (1.2) (with d = 1, R = 0 and f = 0),  $1 \leq l \in \mathbb{Z}$ , and for each  $l \geq 1$ , there is a time sequence  $t_{l,k} \to \infty$  (as  $k \to \infty$ ) such that for each  $s \in \mathbb{R}$ ,

$$\lim_{t \to \infty} \left( \|v_l(\cdot, 0)\|_{(s+1)} + \|\partial_t v_l(\cdot, 0)\|_{(s)} \right) = 0.$$
(16.86)

Moreover,

$$\mathfrak{E}_{\hom}[v_l](t_{l,k}) \ge \exp\left[2\beta_X^{-1}\kappa(1-\epsilon)e^{\beta_X t_{l,k}}\right],\tag{16.87}$$

where

$$\mathfrak{E}_{\text{hom}}[u](t) := \frac{1}{2} \int_{\mathbb{S}^1} [|u_t(x,t)|^2 + g^{11}(t)|u_x(x,t)|^2] dx.$$
(16.88)

**Remark 16.28.** In (16.88), we denote the coordinate on  $\mathbb{S}^1$  by x instead of by  $x^1$  for the sake of brevity.

**Remark 16.29.** If the assumptions of Proposition 2.5 are satisfied, then the conditions of Lemma 16.8 are fulfilled with  $\beta_X > 0$  (even though the requirement (16.19) does not appear explicitly in Assumption 2.1, it follows from the remaining assumptions (by increasing  $\mathfrak{b}$ , if necessary)). That Assumption 16.10 holds is an immediate consequence of this; cf. Remark 16.11. That Assumption 16.15 holds is an immediate consequence of Assumption 2.3. Finally, considering the definition of  $R_{\infty}$  (given in (16.36)), it is clear that the  $\kappa$  introduced in the statement of Lemma 16.27 coincides with the  $\kappa$  introduced in the statement of Proposition 2.5. Moreover,  $\kappa > 0$  due to Assumption 2.3. To conclude, if the assumptions of Proposition 2.5 are satisfied, then the assumptions of Lemma 16.27 are satisfied, so that the conclusions of Lemma 16.27 hold.

*Proof.* Considering Lemma 16.23, it is clear that there are two cases to consider:  $\beta_X - \beta_1 \leq 0$  and  $\beta_X - \beta_1 > 0$ .

The case  $\beta_X - \beta_1 \leq 0$ . Fix  $\epsilon > 0$  and apply Lemma 16.23 with  $\epsilon$  replaced by  $\epsilon/2$ . Then there are corresponding constants  $T_{0,\epsilon}$ ,  $T_{\text{med},\epsilon}$  and  $N_{\text{eep},\epsilon}$  (depending only on  $\mathbf{c}_{\text{eep}}$  and  $\epsilon$ ) such that the following holds. For  $n_1 \geq N_{\text{eep},\epsilon}$ , there is a solution z to (24.16) (corresponding to f = 0 and frequency  $n_1$ ) such that

$$|w(t_{k+1})| \ge \exp\left[\beta_X^{-1}\kappa(1-\epsilon/2)e^{\beta_X t_{k+1}}\right]$$
 (16.89)

for all  $k \ge 0$  such that  $t_k \ge T_{\text{med},\epsilon}$ , where  $\{t_k\}$  is the sequence constructed in Definition 15.5 and starting at  $t_0 := T_{0,\epsilon}$ . Moreover,  $|w(t_0)| = 1$ . Consider the solution u to (1.2) (where d = 1, R = 0and f = 0) defined by

$$u(x,t) := (2\pi)^{-1/2} z(n_1,t) e^{in_1 x}; (16.90)$$

for the sake of brevity, we here use x instead of  $x^1$  to denote the variable on  $\mathbb{S}^1$ . The reason for including the factor  $(2\pi)^{-1/2}$  is that we wish the relation (24.14) to hold. Compute

$$\mathfrak{E}_{\text{hom}}[u](t_{k+1}) = \frac{1}{2} |w(t_{k+1})|^2 \ge \frac{1}{2} \exp\left[2\beta_X^{-1}\kappa(1-\epsilon/2)e^{\beta_X t_{k+1}}\right]$$
(16.91)

for  $k \ge 0$  such that  $t_k \ge T_{\text{med},\epsilon}$ , where  $\mathfrak{E}_{\text{hom}}[u]$  is defined in the statement of the lemma. Letting  $s_k = t_{k+1}$  and  $v_l$  be defined by  $v_l(x,t) := u(x,t)/l$  yields

$$\mathfrak{E}_{\text{hom}}[v_l](s_k) \ge \frac{1}{2l^2} \exp\left[\beta_X^{-1} \kappa \epsilon e^{\beta_X s_k}\right] \exp\left[2\beta_X^{-1} \kappa (1-\epsilon) e^{\beta_X s_k}\right].$$

For  $s_k$  large enough (the bound depending only on  $\epsilon$ , l,  $\beta_X$  and  $\kappa$ ), it is thus clear that

$$\mathfrak{E}_{\mathrm{hom}}[v_l](s_k) \ge \exp\left[2\beta_X^{-1}\kappa(1-\epsilon)e^{\beta_X s_k}\right]$$

On the other hand, it is clear that  $\partial_t v_l(\cdot, t_0)$  and  $v_l(\cdot, t_0)$  converge to zero in any  $H^s$ -norm as  $l \to \infty$ . By Cauchy stability, the same is true of  $\partial_t v_l(\cdot, 0)$  and  $v_l(\cdot, 0)$ . The lemma follows in the case that  $\beta_X - \beta_1 \leq 0$ .

The case  $\beta_X - \beta_1 > 0$ . Fix  $\epsilon > 0$  and apply Lemma 16.23 with  $\epsilon$  replaced by  $\epsilon/2$ . Then there are corresponding constants  $T_{0,\epsilon}$ ,  $T_{\text{med},\epsilon}$ ,  $K_{\text{eep},\epsilon}$  and  $N_{\text{eep},\epsilon}$  (depending only on  $\mathbf{c}_{\text{eep}}$  and  $\epsilon$ ) such that the following holds. For  $n_1 \geq N_{\text{eep},\epsilon}$ , there is a solution z to (24.16) (corresponding to f = 0 and frequency  $n_1$ ) such that

$$|w(t_{k+1})| \ge \exp\left[\beta_X^{-1}\kappa(1-\epsilon/2)e^{\beta_X t_{k+1}}\right]$$
(16.92)

for all  $k \ge 0$  such that  $T_{\text{med},\epsilon} \le t_k < T_{\max,\epsilon}$ , where  $\{t_k\}$  is the sequence constructed in Definition 15.5 and starting at  $t_0 := T_{0,\epsilon}$ . Moreover,  $|w(t_0)| = 1$ . Note that

$$T_{\max,\epsilon} = \frac{1}{\beta_X - \beta_1} \ln n_1 + K_{\text{eep},\epsilon}.$$

Note that if  $t_{k_{\max}}$  is the largest admissible  $t_k$ , then, by definition,  $t_{k_{\max}+1} \ge T_{\max,\epsilon}$ . Thus (16.92) yields

$$|w(t_{k_{\max}+1})| \ge \exp\left[\mu_{\operatorname{eep},\epsilon} n_1^{\frac{\beta_X}{\beta_X-\beta_1}}\right] \exp\left[\beta_X^{-1} \kappa (1-\epsilon) e^{\beta_X t_{k_{\max}+1}}\right],$$

where

$$\mu_{\text{eep},\epsilon} := \frac{\epsilon \kappa}{2\beta_X} e^{\beta_X K_{\text{eep},\epsilon}} > 0.$$
(16.93)

Let  $u_{n_1}$  be the  $\mathbb{C}^m$ -valued solution to (1.2) defined by (16.90) and note that

$$\mathfrak{E}_{\mathrm{hom}}[u_{n_1}](t_{k_{\mathrm{max}}+1}) \ge \frac{1}{2} \exp\left[2\mu_{\mathrm{eep},\epsilon} n_1^{\frac{\beta_X}{\beta_X-\beta_1}}\right] \exp\left[2\beta_X^{-1} \kappa (1-\epsilon) e^{\beta_X t_{k_{\mathrm{max}}+1}}\right]$$
(16.94)

and that  $\mathfrak{E}_{\text{hom}}[u_{n_1}](t_0) = 1/2$ . Note that  $k_{\text{max}}$  depends on  $n_1$ . For the sake of clarity, we therefore write  $k_{\text{max}}(n_1)$  in what follows. Define v by

$$v(x,t) := \sum_{n_1 \ge N_{\text{eep},\epsilon}} \sqrt{2} \exp\left[-\mu_{\text{eep},\epsilon} n_1^{\frac{\beta_X}{\beta_X - \beta_1}}\right] u_{n_1}(x,t).$$
(16.95)

Then v is a smooth  $\mathbb{C}^m$ -valued solution to (1.2). In fact, it is more than smooth, as we demonstrate below. Due to (16.94) it is clear that for each  $n_1 \geq N_{\text{eep},\epsilon}$ ,

$$\mathfrak{E}_{\text{hom}}[v](t_{k_{\max}(n_1)+1}) \ge \exp\left[2\beta_X^{-1}\kappa(1-\epsilon)e^{\beta_X t_{k_{\max}(n_1)+1}}\right].$$
(16.96)

Since  $t_{k_{\max}(n_1)+1} \to \infty$  as  $n_1 \to \infty$ , it is clear that there is a sequence  $s_k \to \infty$  such that

$$\mathfrak{E}_{\mathrm{hom}}[v](s_k) \ge \exp\left[2\beta_X^{-1}\kappa(1-\epsilon)e^{\beta_X s_k}\right].$$

Let

$$v_l(x,t) := \sum_{n_1 \ge l, N_{\text{eep},\epsilon}} \sqrt{2} \exp\left[-\mu_{\text{eep},\epsilon} n_1^{\frac{\beta_X}{\beta_X - \beta_1}}\right] u_{n_1}(x,t).$$
(16.97)

Then  $\partial_t v_l(\cdot, t_0)$  and  $v_l(\cdot, t_0)$  converge to zero in any  $H^s$ -norm as  $l \to \infty$ . By Cauchy stability, the same is true of  $\partial_t v_l(\cdot, 0)$  and  $v_l(\cdot, 0)$ . Finally, there is a time sequence  $t_{l,k}$  with the desired properties. The lemma follows.

### 16.8 Super exponential growth of solutions

In this chapter we have, up till now, focused on the equation (1.2) in the case that d = 1, R = 0and f = 0. However, the results we have obtained yield conclusions for more general equations. We turn to this topic next.

#### 16.8.1 Density of solutions yielding super exponential growth

Let us start by proving that the initial data that yield solutions with super exponential growth are dense.

**Proposition 16.30.** Consider (1.2). Assume the associated metric to be such that (M,g) is a canonical separable cosmological model manifold. Assume, moreover, that there is a  $j \in \{1, ..., d\}$  such that

$$u_{tt} - g^{jj}(t)\partial_j^2 u - 2g^{0j}(t)\partial_t\partial_j u + \alpha(t)u_t + X^j(t)\partial_j u + \zeta(t)u = 0$$
(16.98)

(no summation on j) satisfies the conditions of Lemma 16.27. Let  $\epsilon > 0$ . For every  $\mathbb{C}^m$ -valued solution u to (1.2), there is then a sequence of  $\mathbb{C}^m$ -valued solutions  $u_l$ ,  $1 \leq l \in \mathbb{Z}$ , such that the initial data of  $u_l - u$  at t = 0 converge to zero with respect to every Sobolev norm and such that for every  $1 \leq l \in \mathbb{Z}$  there is a sequence  $t_{l,k} \to \infty$  (as  $k \to \infty$ ) such that

$$\mathfrak{E}_{\hom}[u_l](t_{l,k}) \ge \exp\left[2\beta_X^{-1}\kappa(1-\epsilon)e^{\beta_X t_{l,k}}\right],\tag{16.99}$$

where  $\mathfrak{E}_{\text{hom}}$  is given by (1.21).

**Remark 16.31.** Since the initial data of  $u_l - u$  at t = 0 converge to zero with respect to every Sobolev norm, it is clear that they also converge to zero in the  $C^{\infty}$  topology.

**Remark 16.32.** Due to the fact that (M, g) is a canonical separable cosmological model manifold, the metric arising from (16.98) also corresponds to a canonical separable cosmological model manifold. The reason for this is the following. Let  $h^{00}(t) = -1$ ,  $h^{01}(t) = g^{0j}(t)$  and  $h^{11}(t) = g^{jj}(t)$ (no summation on j). Then  $h^{11}$  is strictly positive by assumption, so that the matrix with components  $h^{\alpha\beta}(t)$  has an inverse with components  $h_{\alpha\beta}(t)$ . Moreover,  $h_{00} < 0$ ,  $h_{11} > 0$  and  $h_{\alpha\beta}$ defines a Lorentz metric h on  $M_j := \mathbb{S}^1 \times I$ . The justification of these statements is given at the beginning of Section 25.1. Finally, that  $(M_j, h)$  is a canonical separable cosmological model manifold follows from the fact that I contains  $[0, \infty)$  and the fact that N = 1; the latter statement is a consequence of Lemma 25.2, Remark 25.3 and the fact that  $h^{00} = -1$ .

**Remark 16.33.** Using (24.11), it can be verified that

$$\mathfrak{E}_{\text{hom}}[u](t) = \frac{1}{2} \sum_{\iota \in \mathcal{I}_{\text{B}}} \left( |\dot{z}(\iota, t)|^2 + \mathfrak{g}^2(\iota, t) |z(\iota, t)|^2 \right).$$
(16.100)

*Proof.* Let u be a  $\mathbb{C}^m$ -valued solution to (1.2). There are two possibilities. Either there is a time sequence  $t_k \to \infty$  such that (16.99) holds with  $u_l$  replaced by u and  $t_{l,k}$  replaced by  $t_k$ . In that case we can choose  $u_l = u$  for all  $1 \leq l \in \mathbb{Z}$ . The second possibility is that there is a T > 0 such that

$$\mathfrak{E}_{\text{hom}}[u](t) \le \exp\left[2\beta_X^{-1}\kappa(1-\epsilon)e^{\beta_X t}\right]$$
(16.101)

for all  $t \ge T$ . In this case we apply Lemma 16.27, with  $\epsilon$  replaced by  $\epsilon/2$ , to the equation (16.98). This leads to a sequence of solutions  $v_l$  to (16.98) such that the initial data of  $v_l$  at t = 0 converge to zero with respect to any Sobolev norm and such that (16.87) holds with  $\epsilon$  replaced by  $\epsilon/2$ . We can also, without loss of generality, assume that the sequence  $t_{l,k}$  to be such that  $t_{l,k} \ge T$  for all  $1 \le l, k \in \mathbb{Z}$ . Then

$$\begin{split} \mathfrak{E}_{\mathrm{hom}}^{1/2}[u+v_l](t_{l,k}) \geq \mathfrak{E}_{\mathrm{hom}}^{1/2}[v_l](t_{l,k}) - \mathfrak{E}_{\mathrm{hom}}^{1/2}[u](t_{l,k}) \\ \geq \exp\left[\beta_X^{-1}\kappa(1-\epsilon/2)e^{\beta_X t_{l,k}}\right] - \exp\left[\beta_X^{-1}\kappa(1-\epsilon)e^{\beta_X t_{l,k}}\right]. \end{split}$$

Thus  $u_l := u + v_l$  has the desired properties, and the proposition follows.

#### 16.8.2 Genericity of solutions yielding super exponential growth

Consider an equation satisfying the conditions of Proposition 16.30. Then the set of initial data leading to solutions that exhibit super exponential growth is dense. However, there are also classes of equations satisfying the conditions of Proposition 16.30 such that the set of initial data leading to solutions that exhibit super exponential *decay* is dense; cf. Section 16.11 below. It is therefore of interest to ask what the generic behaviour of solutions is. To address this question, we first have to decide on a notion of genericity. One way is to give a measure theoretic definition. However, that is not so natural in the present situation, since the space of initial data is infinite dimensional. Another option would be to define a set to be generic if it is open and dense. However, in the present setting we are interested in distinguishing between two sets which are both (potentially) dense. For these reasons, we here use the following definition of genericity: a set of initial data is *generic* if it is a countable intersection of open and dense sets with respect to the  $C^{\infty}$  topology. Clearly, one could also consider other function spaces, for instance Sobolev spaces. However, we here only state the results in the smooth setting.

**Proposition 16.34.** Consider (1.2). Assume the associated metric to be such that (M, g) is a canonical separable cosmological model manifold. Assume, moreover, that there is a  $j \in \{1, \ldots, d\}$  such that (16.98) (no summation on j) satisfies the conditions of Lemma 16.27. Let  $\epsilon > 0$ . Then there is a set of smooth  $\mathbb{C}^m$ -valued initial data to (1.2), say  $\mathcal{A}$ , with the following properties. First,  $\mathcal{A}$  is the intersection of a countable number of open and dense sets (with respect to the  $\mathbb{C}^{\infty}$  topology). Second, each element of  $\mathcal{A}$  corresponds to a  $\mathbb{C}^m$ -valued solution u to (1.2) such that there is a time sequence  $\{t_k\}, 1 \leq k \in \mathbb{Z}$ , with the properties that  $t_k \to \infty$  and

$$\mathfrak{E}_{\text{hom}}[u](t_k) \ge \exp\left[2\beta_X^{-1}\kappa(1-\epsilon)e^{\beta_X t_k}\right]$$
(16.102)

for all  $1 \leq k \in \mathbb{Z}$ , where  $\mathfrak{E}_{hom}[u](t)$  is defined by (1.21).

**Remark 16.35.** Due to this statement, it is clear that the generic behaviour of solutions is to exhibit super exponential growth.

**Remark 16.36.** Due to the proof, the sets  $\mathcal{A}_N$  whose intersection yield  $\mathcal{A}$  are open with respect to any Sobolev space topology on the set of initial data. Moreover, the density is demonstrated by perturbing the initial data by a sequence of functions converging to zero in the  $C^{\infty}$ -topology. It should thus not be a problem to prove genericity in Sobolev-type regularity classes of initial data.

Proof. The goal of the proof is to construct the set  $\mathcal{A}$ . We do so by defining sets  $\mathcal{A}_N, N_0 \leq N \in \mathbb{Z}$ , with the following properties. First,  $\mathcal{A}$  is the intersection of the  $\mathcal{A}_N$ . Second, the  $\mathcal{A}_N$  are open and dense (with respect to the  $C^{\infty}$  topology). Third, to each solution u corresponding to initial data in  $\mathcal{A}$  there is a corresponding time sequence  $t_k \to \infty$  such that (16.102) holds. The definition of  $\mathcal{A}_N$  depends on two parameters  $\epsilon$  and  $N_0$  and a sequence  $\{T_k\}$ . The parameter  $\epsilon$  is fixed in the statement of the proposition. However, the definitions of  $N_0$  and the sequence  $T_k$  depend on whether  $\beta_X - \beta_1 \leq 0$  or  $\beta_X - \beta_1 > 0$ . On the other hand, it is convenient to first give a general definition of  $\mathcal{A}_N$  and only later define  $N_0$  and the sequence  $T_k$ .

**Definition of**  $\mathcal{A}_N$ . Fix  $0 < \epsilon \leq 1$  and  $1 \leq N_0 \in \mathbb{Z}$ . Let  $0 < T_k \in \mathbb{R}$ ,  $N_0 \leq k \in \mathbb{Z}$ , be a sequence tending to  $\infty$  as  $k \to \infty$ . For  $N_0 \leq N \in \mathbb{Z}$ , then  $\xi \in \mathcal{A}_N$  if and only if the following holds. First  $\xi$  are smooth  $\mathbb{C}^m$ -valued initial data to (1.2). Second, if u is the solution corresponding to  $\xi$  and  $z(\iota, t)$  is defined by (24.14), then there is a  $k \geq N$  such that

$$|\dot{z}(\iota_k, T_k)|^2 + \mathfrak{g}^2(\iota_k, T_k)|z(\iota_k, T_k)|^2 > 2\exp\left[2\beta_X^{-1}\kappa(1-\epsilon)e^{\beta_X T_k}\right].$$
(16.103)

Here  $\iota_k$  is defined by the conditions that the j'th component of  $\iota_k$  equals k and all the remaining components equal zero.

Let us first prove that the  $\mathcal{A}_N$  are open sets. Let u be a solution corresponding to initial data, say  $\xi$ , in  $\mathcal{A}_N$ . Then there is a  $N \leq k \in \mathbb{Z}$  as above. In particular, (16.103) holds. Since the condition

(16.103) is open, there is an open neighbourhood, say U, of  $\xi$  such that solutions corresponding to initial data in U also have the property that (16.103) holds. Note that U is open with respect to any choice of Sobolev regularity on the initial data, as well as with respect to the  $C^{\infty}$ -topology. Assume now that  $\xi$  is in the intersection of all the  $\mathcal{A}_N$  for  $N_0 \leq N \in \mathbb{Z}$ . For each such N, let k = k(N) be as above and let  $t_N := T_{k(N)}$ . Since  $k(N) \geq N$ , it is clear that  $t_N \to \infty$ . Moreover, due to (16.103) and the fact that (16.100) holds, (16.102) is satisfied for all k. What remains to be done is to prove that the  $\mathcal{A}_N$  are dense for a suitable choice of  $N_0$  and  $\{T_k\}$ . However, to achieve this goal, we need to divide the analysis into the two cases  $\beta_X - \beta_1 > 0$  and  $\beta_X - \beta_1 \leq 0$ , just as in the proof of Lemma 16.27.

The case  $\beta_X - \beta_1 \leq 0$ . Fix  $\epsilon > 0$  and apply Lemma 16.23 with  $\epsilon$  replaced by  $\epsilon/2$  to the equation (16.98). Then there are corresponding constants  $T_{0,\epsilon}$ ,  $T_{\text{med},\epsilon}$  and  $N_{\text{eep},\epsilon}$  (depending only on  $\mathbf{c}_{\text{eep}}$  and  $\epsilon$ ) such that the conclusions stated in the proof of Lemma 16.27 (concerning the case  $\beta_X - \beta_1 \leq 0$ ) hold. Let  $N_0$  be the smallest strictly positive integer which is larger than  $N_{\text{eep},\epsilon}$ . Define  $T_l$ ,  $N_0 \leq l \in \mathbb{Z}$ , such that  $T_l \geq \ln l + 1$  and that  $T_l$  equals  $t_{k+1}$  for some k with the property that  $t_k \geq T_{\text{med},\epsilon}$  (where  $t_k$  is defined as in the proof of Lemma 16.27 in the case that  $n_1 = l$ ). Then  $T_l \to \infty$  as  $l \to \infty$ . Defining  $\mathcal{A}_N$  as above, with this choice of  $\{T_l\}$ , we need to prove that  $\mathcal{A}_N$  is dense. Define  $u_{n_1}$  to be the right hand side of (16.90). Then

$$\mathfrak{E}_{\text{hom}}[u_l](T_l) \ge \frac{1}{2} \exp\left[2\beta_X^{-1}\kappa(1-\epsilon/2)e^{\beta_X T_l}\right]$$

and  $\mathfrak{E}_{\text{hom}}[u_l](t_0) = 1/2$ . Define  $v_l$  by

$$v_l := \sqrt{2} \exp\left[-\beta_X^{-1} \kappa \epsilon e^{\beta_X T_l} / 4\right] u_l$$

Then

$$\mathfrak{E}_{\mathrm{hom}}[v_l](t_0) \leq \exp\left[-\beta_X^{-1}\kappa\epsilon e^{\beta_X T_l}/2\right].$$

Since  $T_l \ge \ln l$ , this implies that every Sobolev norm of the initial data of  $v_l$  induced at  $t_0$  converges to zero as  $l \to \infty$  (and, due to Cauchy stability, the same is true with  $t_0$  replaced by 0). Fix  $N \ge N_0$ and let u be a solution to (1.2). Let z be defined by (24.14). We need to prove that the initial data of u at t = 0, say  $\xi$ , are in the closure of  $\mathcal{A}_N$ . If  $\xi \in \mathcal{A}_N$ , there is nothing to prove, so let us assume that  $\xi \notin \mathcal{A}_N$ . Then for all  $k \ge N$ ,

$$|\dot{z}(\iota_k, T_k)|^2 + \mathfrak{g}^2(\iota_k, T_k)|z(\iota_k, T_k)|^2 \le 2\exp\left[2\beta_X^{-1}\kappa(1-\epsilon)e^{\beta_X T_k}\right].$$

If  $z_k(\iota, t)$  is defined by (24.14) with u replaced by  $u + v_k$ , we thus have

$$[|\dot{z}_{k}(\iota_{k},T_{k})|^{2} + \mathfrak{g}^{2}(\iota_{k},T_{k})|z_{k}(\iota_{k},T_{k})|^{2}]^{1/2}$$

$$\geq \sqrt{2}\mathfrak{E}_{hom}^{1/2}[v_{k}](T_{k}) - [|\dot{z}(\iota_{k},T_{k})|^{2} + \mathfrak{g}^{2}(\iota_{k},T_{k})|z(\iota_{k},T_{k})|^{2}]^{1/2}$$

$$\geq \sqrt{2}\exp\left[\beta_{X}^{-1}\kappa(1-3\epsilon/4)e^{\beta_{X}T_{k}}\right] - \sqrt{2}\exp\left[\beta_{X}^{-1}\kappa(1-\epsilon)e^{\beta_{X}T_{k}}\right].$$
(16.104)

Then (16.104) implies that the inequality (16.103) (with z replaced by  $z_k$ ) is satisfied for  $T_k$  large enough. Thus the initial data of  $u + v_k$  at t = 0, say  $\xi_k$ , belong to  $\mathcal{A}_N$  for k large enough. On the other hand,  $\xi_k$  converges to  $\xi$  with respect to any Sobolev space topology. This proves that  $\mathcal{A}_N$  is not only open, but also dense. The proposition follows in case  $\beta_X - \beta_1 \leq 0$ .

The case  $\beta_X - \beta_1 > 0$ . Fix  $\epsilon > 0$  and apply Lemma 16.23 with  $\epsilon$  replaced by  $\epsilon/2$ . Then there are corresponding constants  $T_{0,\epsilon}$ ,  $T_{\text{med},\epsilon}$ ,  $K_{\text{eep},\epsilon}$  and  $N_{\text{eep},\epsilon}$  (depending only on  $\mathbf{c}_{\text{eep}}$  and  $\epsilon$ ) such that the conclusions stated in the proof of Lemma 16.27 (concerning the case  $\beta_X - \beta_1 > 0$ ) hold. Let  $N_0$  be the smallest strictly positive integer which is larger than  $N_{\text{eep},\epsilon}$ . For  $N_0 \leq l \in \mathbb{Z}$ , define  $T_l := t_{k_{\text{max}}(l)+1}$ , where the sequence  $t_k$  as well as  $k_{\text{max}}(l)$  are defined as in the proof of Lemma 16.27 (in the case that  $n_1 = l$ ). Finally, define  $\mathcal{A}_N$  as above and let  $u_{n_1}$  denote the right hand side of (16.90). Then the function

$$v_l := \sqrt{2} \exp(-\mu_{\operatorname{eep},\epsilon} l^{\beta_X/(\beta_X - \beta_1)}/2) u_l$$

where  $\mu_{\text{eep},\epsilon}$  is given by (16.93), is such that we can finish the proof of density as in the case of  $\beta_X - \beta_1 \leq 0$ . The proposition follows.

Even though Proposition 16.34 does demonstrate the genericity of initial data yielding solutions with super exponential growth, the notion of genericity can be questioned; in the finite dimensional setting, a dense  $G_{\delta}$  set need not have full measure. We therefore here complement Proposition 16.34 with an additional result.

**Proposition 16.37.** Consider (1.2). Assume the associated metric to be such that (M, g) is a canonical separable cosmological model manifold. Assume, moreover, that there is a  $j \in \{1, \ldots, d\}$  such that (16.98) (no summation on j) satisfies the conditions of Lemma 16.27. Let  $\epsilon > 0$ . Define  $\mathcal{B}_{\epsilon}$  by the condition that  $\xi \in \mathcal{B}_{\epsilon}$  iff  $\xi$  constitutes smooth  $\mathbb{C}^{m}$ -valued initial data for (1.2) such that if u is the solution corresponding to  $\xi$ , then there is a constant  $C_{\epsilon}$  (depending on the solution) such that

$$\mathfrak{E}_{\mathrm{hom}}[u](t) \le C_{\epsilon} \exp\left[2\beta_X^{-1}\kappa(1-\epsilon)e^{\beta_X t}\right]$$

for all  $t \ge 0$ . Define  $\mathcal{A}_{\epsilon}$  by the condition that  $\xi \in \mathcal{A}_{\epsilon}$  iff  $\xi$  constitutes smooth  $\mathbb{C}^m$ -valued initial data for (1.2) such that if u is the solution corresponding to  $\xi$ , then there is a time sequence  $0 \le t_k \to \infty$  such that

$$\mathfrak{E}_{\text{hom}}[u](t_k) \ge \exp\left[2\beta_X^{-1}\kappa(1-\epsilon/2)e^{\beta_X t_k}\right]$$
(16.105)

for all k. Then  $\mathcal{A}_{\epsilon}$  and  $\mathcal{B}_{\epsilon}$  are disjoint. Moreover, there is an infinite dimensional (linear) subspace  $P_{\epsilon}$  of the vector space of smooth  $\mathbb{C}^m$ -valued initial data for (1.2) such that if  $\xi \in \mathcal{B}_{\epsilon}$  and  $\xi_{\text{per}} \in P_{\epsilon}$ , then  $\xi + \xi_{\text{per}} \in \mathcal{A}_{\epsilon}$  unless  $\xi_{\text{per}} = 0$ .

**Remark 16.38.** The statement of the proposition can be interpreted as saying that  $\mathcal{B}_{\epsilon}$  has infinite codimension in the set of smooth initial data for (1.2).

**Remark 16.39.** By the vector space of smooth  $\mathbb{C}^m$ -valued initial data for (1.2), we here mean the space  $C^{\infty}(\bar{M}, \mathbb{C}^m) \times C^{\infty}(\bar{M}, \mathbb{C}^m)$ , where  $\bar{M}$  is defined by (1.3).

*Proof.* That  $\mathcal{A}_{\epsilon}$  and  $\mathcal{B}_{\epsilon}$  are disjoint is an immediate consequence of the definitions. In order to construct the subspace  $P_{\epsilon}$ , let us return to the proof of Lemma 16.27. As in the case of that proof, we need to divide the analysis into the two cases  $\beta_X - \beta_1 \leq 0$  and  $\beta_X - \beta_1 > 0$ .

The case  $\beta_X - \beta_1 \leq 0$ . When  $\beta_X - \beta_1 \leq 0$ , we proceed as in the proof of Lemma 16.27, but with  $\epsilon$  replaced by  $\epsilon/2$  (in other words, we, to start with, apply Lemma 16.23 with  $\epsilon$  replaced by  $\epsilon/4$ ). This yields, in particular, numbers  $T_{0,\epsilon}$ ,  $T_{\text{med},\epsilon}$ ,  $N_{\text{eep},\epsilon}$  etc., depending only on  $\mathbf{c}_{\text{eep}}$  and  $\epsilon$ . For  $n_1 \geq N_{\text{eep},\epsilon}$ , we can then define  $u_{n_1}$  by the right hand side of (16.90) (where x should be interpreted as  $x^j$  in the present setting). Moreover, (16.91) with  $\epsilon$  replaced by  $\epsilon/2$  holds and yields

$$\mathfrak{E}_{\text{hom}}[u_{n_1}](t_{k+1}) \ge \frac{1}{2} \exp\left[2\beta_X^{-1}\kappa(1-\epsilon/4)e^{\beta_X t_{k+1}}\right]$$
(16.106)

for all  $k \ge 0$  such that  $t_k \ge T_{\text{med},\epsilon}$  (where the sequence  $t_k$  is constructed as in the proof of Lemma 16.27). Note, however, that the sequence  $t_k$  depends on  $n_1$ . The initial data corresponding to  $u_{n_1}$  for different  $n_1$ 's are linearly independent. Moreover, if

$$u = \sum_{n_1 \ge N_{\text{eep},\epsilon}} a_{n_1} u_{n_1} \tag{16.107}$$

for some complex numbers  $a_{n_1}$  (which we assume to be such that the sum converges to a smooth function), then

$$\mathfrak{E}_{\hom}[u] = \sum_{n_1 \ge N_{\operatorname{eep},\epsilon}} |a_{n_1}|^2 \mathfrak{E}_{\hom}[u_{n_1}].$$
(16.108)

Let  $P_{\epsilon}$  be the closure, in the  $C^{\infty}$ -topology, of the space of smooth  $\mathbb{C}^m$ -valued initial data spanned by the initial data of the  $u_{n_1}$ 's for  $n_1 \geq N_{\text{eep},\epsilon}$ . For future reference, it is of interest to note that elements of  $P_{\epsilon}$  can be considered to be initial data to (16.98). Note also that  $P_{\epsilon}$  is infinite dimensional. Let now  $\xi \in \mathcal{B}_{\epsilon}$ ,  $\xi_{\text{per}} \in P_{\epsilon}$ , v be the solution to (1.2) corresponding to  $\xi$ , and let ube the solution to (16.98) corresponding to  $\xi_{\text{per}}$ . Then v + u is the solution to (1.2) corresponding to  $\xi + \xi_{\text{per}}$ . Moreover,

$$\mathfrak{E}_{\text{hom}}^{1/2}[v+u] \ge \mathfrak{E}_{\text{hom}}^{1/2}[u] - \mathfrak{E}_{\text{hom}}^{1/2}[v] \ge \left(\sum_{n_1 \ge N_{\text{eep},\epsilon}} |a_{n_1}|^2 \mathfrak{E}_{\text{hom}}[u_{n_1}]\right)^{1/2} - \mathfrak{E}_{\text{hom}}^{1/2}[v], \quad (16.109)$$

where we have expressed u as in (16.107). There are two possibilities. Either all the  $a_{n_1}$ 's vanish, in which case there is nothing to prove, or there is one  $n_1 \ge N_{\text{eep},\epsilon}$  such that  $a_{n_1} \ne 0$ . Assume that the latter occurs, and let  $t_k$  be the time sequence associated with  $n_1$  as in the proof of Lemma 16.27. Then (16.106) and (16.109) imply that if  $k \ge 0$  and  $t_k \ge T_{\text{med},\epsilon}$ ,

$$\begin{aligned} \mathfrak{E}_{\text{hom}}^{1/2}[v+u](t_{k+1}) \geq &|a_{n_1}|\mathfrak{E}_{\text{hom}}^{1/2}[u_{n_1}](t_{k+1}) - \mathfrak{E}_{\text{hom}}^{1/2}[v](t_{k+1}) \\ \geq &2^{-1/2}|a_{n_1}|\exp\left[\beta_X^{-1}\kappa(1-\epsilon/4)e^{\beta_X t_{k+1}}\right] \\ &- C_{\epsilon}^{1/2}\exp\left[\beta_X^{-1}\kappa(1-\epsilon)e^{\beta_X t_{k+1}}\right]. \end{aligned}$$

For k large enough, the estimate (16.105) thus holds with u replaced by u + v. Thus  $\xi + \xi_{per} \in \mathcal{A}_{\epsilon}$ . This finishes the proof in case  $\beta_X - \beta_1 \leq 0$ .

The case  $\beta_X - \beta_1 > 0$ . When  $\beta_X - \beta_1 > 0$ , we again proceed as in the proof of Lemma 16.27, but with  $\epsilon$  replaced by  $\epsilon/4$  (in other words, we, to start with, apply Lemma 16.23 with  $\epsilon$  replaced by  $\epsilon/8$ ). This yields, in particular, numbers  $N_{\text{eep},\epsilon}$ ,  $K_{\text{eep},\epsilon}$  etc., depending only on  $\mathbf{c}_{\text{eep}}$  and  $\epsilon$ . Define, moreover,  $\mu_{\text{eep},\epsilon}$  by (16.93) (with  $\epsilon$  replaced by  $\epsilon/4$ ). Let p > 1 be a prime number, and define  $k_{p,\text{eep},\epsilon}$  by

$$k_{p,\operatorname{eep},\epsilon} := \frac{\ln N_{\operatorname{eep},\epsilon}}{\ln p}.$$

We then define  $v_p$  by

$$v_p := \sum_{k \ge k_{p, eep, \epsilon}} \sqrt{2} \exp\left[-\mu_{eep, \epsilon} p^{k \frac{\beta_X}{\beta_X - \beta_1}}\right] u_{p^k},$$

where  $u_{n_1}$  is defined by the right hand side of (16.90); cf. (16.95). Due to an estimate similar to (16.96), there is, for each p, a sequence  $t_{p,l} \to \infty$  as  $k \to \infty$  such that

$$\mathfrak{E}_{\hom}[v_p](t_{p,l}) \ge \exp\left[2\beta_X^{-1}\kappa(1-\epsilon/4)e^{\beta_X t_{p,l}}\right].$$

Moreover, all the  $v_p$  are linearly independent. In fact, formulae analogous to (16.107) and (16.108) hold. We can now finish the proof as in the case  $\beta_X - \beta_1 \leq 0$ . The proposition follows.

#### 16.8.3 Universal bounds on the growth of the energy

Consider the equation (1.2) in case f = 0. Let S be the set of smooth solutions to (1.2), and let  $S_{\text{fin}}$  be the set of smooth solutions to (1.2) with finite frequency content. In other words,  $u \in S_{\text{fin}}$  iff  $z(\iota, \cdot)$  defined by (24.14) is only non-zero for a finite number of  $\iota \in \mathcal{I}_{\text{B}}$ . Define  $\mathfrak{G}, \mathfrak{G}_{\text{fin}} : I \to (0, \infty)$  by

$$\mathfrak{G}(t) := \sup_{0 \neq u \in \mathcal{S}} \frac{\mathfrak{E}[u](t)}{\mathfrak{E}[u](0)}, \quad \mathfrak{G}_{\text{fin}}(t) := \sup_{0 \neq u \in \mathcal{S}_{\text{fin}}} \frac{\mathfrak{E}[u](t)}{\mathfrak{E}[u](0)}, \tag{16.110}$$

where

$$\mathfrak{E}[u](t) := \mathfrak{E}_{\text{hom}}[u](t) + \frac{1}{2} \int_{\bar{M}} |u(\cdot, t)|^2 \mu_{\text{B}}$$

Clearly, an argument is required to justify that  $\mathfrak{G}$  and  $\mathfrak{G}_{\text{fin}}$  take their values in  $(0, \infty)$ . However, since it is sufficient to appeal to standard energy estimates, we omit the justification here. The main point of introducing, e.g.,  $\mathfrak{G}$  is of course that

$$\mathfrak{E}[u](t) \le \mathfrak{G}(t)\mathfrak{E}[u](0)$$

for all  $t \in I$  and all smooth solutions u.

**Proposition 16.40.** Consider (1.2). Assume the associated metric to be such that (M,g) is a canonical separable cosmological model manifold. Assume, moreover, that f = 0 and that there is a  $j \in \{1, \ldots, d\}$  such that (16.98) (no summation on j) satisfies the conditions of Lemma 16.27. Define  $\mathfrak{G}$  and  $\mathfrak{G}_{\text{fin}}$  by (16.110). Then  $\mathfrak{G} = \mathfrak{G}_{\text{fin}}$  and for every  $\epsilon > 0$ , there is a time sequence  $t_k \to \infty$  such that  $\mathfrak{G}(t_k) \ge \exp\left[2\beta_X^{-1}\kappa(1-\epsilon)e^{\beta_X t_k}\right]$ .

**Remark 16.41.** As we remarked at the beginning of Subsection 16.8.2, there are equations of the form considered in the proposition such that there is a dense subset of initial data yielding super exponential growth, and a dense subset of initial data yielding super exponential decay. Moreover, the latter set contains the initial data with finite frequency content. Due to the statement of the proposition, it is, however, clear that the super exponential growth can be detected by only considering solutions that decay super exponentially.

*Proof.* That  $\mathfrak{G} = \mathfrak{G}_{\text{fin}}$  is an immediate consequence of Cauchy stability combined with the fact that the set of initial data with finite frequency content is dense in the set of all smooth initial data. Fix  $\epsilon > 0$ . Let  $v_l$  be the sequence of functions obtained by appealing to Lemma 16.27; by assumption, (16.98) satisfies the required conditions. Then we can consider the  $v_l$  to be solutions to (1.2) with f = 0. Fix an l large enough that  $\mathfrak{E}[v_l](0) \leq 1$ ; that this inequality holds for l large enough is a consequence of (16.86). For this fixed l, (16.87) then implies that

$$\mathfrak{B}(t_{l,k}) \ge \mathfrak{E}_{\mathrm{hom}}[v_l](t_{l,k}) \ge \exp\left[2\beta_X^{-1}\kappa(1-\epsilon)e^{\beta_X t_{l,k}}\right]$$

The proposition follows.

# 16.9 Standard energy estimates

Consider the equation (1.2) in case  $g^{00} = -1$ , d = 1, R = 0 and f = 0; i.e.,

$$u_{tt} - g^{11}(t)\partial_1^2 u - 2g^{01}(t)\partial_1\partial_t u + X^1(t)\partial_1 u + \alpha(t)u_t + \zeta(t)u = 0.$$
(16.111)

The purpose of the present section is to derive energy estimates for solutions to this equation, given that the conditions stated in Lemma 16.8 are fulfilled with  $\beta_X > 0$  and that Assumptions 16.10 and 16.15 hold. Consider the energy

$$E_{\text{bas}}(t) := \frac{1}{2} \int_{\mathbb{S}^1} [|u_t(x,t)|^2 + g^{11}(t)|\partial_1 u(x,t)|^2 + e^{2\beta_{\text{ode}}t}|u(x,t)|^2] dx.$$
(16.112)

**Lemma 16.42.** Consider (16.111). Assume the associated metric to be such that (M, g) is a canonical separable cosmological model manifold. Assume that the conditions stated in Lemma 16.8 are fulfilled with  $\beta_X > 0$  and that Assumptions 16.10 and 16.15 hold. Then there is a constant  $C_{ep}$ , depending only on  $\mathbf{c}_{ep}$  introduced in (16.49), such that the following holds. If u is a  $\mathbb{C}^m$ -valued  $C^2$ -solution to (16.111) and  $E_{bas}$  is defined by (16.112), then

$$E_{\text{bas}}(t) \le \exp\left[\beta_X^{-1} \|Y_\infty^1\| e^{\beta_X t} + C_{\text{ep}}\left(\langle t \rangle + e^{\beta_{\text{ode}} t} + e^{(\beta_X - \eta_X)t}\right)\right] E_{\text{bas}}(0)$$

for all  $t \geq 0$ .

**Remark 16.43.** If  $Y_{\infty}^1$  is Hermitian, then  $||Y_{\infty}^1||/2$  equals  $\kappa$ , defined to be the largest absolute value of an eigenvalue of  $Y_{\infty}^1/2$ . In that case it thus follows that, given  $\epsilon > 0$ , there is a  $T_{\text{eep},\epsilon}$ , depending only  $\epsilon$  and  $\mathbf{c}_{\text{eep}}$  such that

$$E_{\text{bas}}(t) \le \exp\left[2\beta_X^{-1}\kappa(1+\epsilon)e^{\beta_X t}\right]E_{\text{bas}}(0)$$

for all  $t \ge T_{\text{eep},\epsilon}$ . This estimate should be compared with (16.87); note that the  $\kappa$  appearing in (16.87) equals the  $\kappa$  defined in the present remark.

*Proof.* Time differentiation of  $E_{\text{bas}}$  yields

$$\begin{split} \partial_t E_{\mathrm{bas}} =& \frac{1}{2} \int_{\mathbb{S}^1} [\langle u_t, u_{tt} \rangle + \langle u_{tt}, u_t \rangle + (\partial_t g^{11}) |\partial_1 u|^2 + g^{11} \langle \partial_1 u, \partial_1 \partial_t u \rangle \\ &+ g^{11} \langle \partial_1 \partial_t u, \partial_1 u \rangle + 2\beta_{\mathrm{ode}} e^{2\beta_{\mathrm{ode}} t} |u|^2 + e^{2\beta_{\mathrm{ode}} t} \langle u, u_t \rangle + e^{2\beta_{\mathrm{ode}} t} \langle u_t, u \rangle] dx \\ \leq & \int_{\mathbb{S}^1} [\mathfrak{b} g^{11} |\partial_1 u|^2 + \beta_{\mathrm{ode}} e^{2\beta_{\mathrm{ode}} t} |u|^2] dx + \frac{1}{2} e^{\beta_{\mathrm{ode}} t} \int_{\mathbb{S}^1} [|u_t|^2 + e^{2\beta_{\mathrm{ode}} t} |u|^2] dx \\ &+ \frac{1}{2} \int_{\mathbb{S}^1} [\langle u_{tt} - g^{11} \partial_1^2 u, u_t \rangle + \langle u_t, u_{tt} - g^{11} \partial_1^2 u \rangle] dx, \end{split}$$

where we have used (16.17) and (16.20). The first two terms on the far right hand side of this inequality can easily be estimated in terms of the basic energy  $E_{\text{bas}}$ . Let us therefore focus on the last term. It can be written

$$\begin{split} &\frac{1}{2} \int_{\mathbb{S}^1} [\langle u_{tt} - g^{11} \partial_1^2 u, u_t \rangle + \langle u_t, u_{tt} - g^{11} \partial_1^2 u \rangle] dx \\ &= \frac{1}{2} \int_{\mathbb{S}^1} [\langle -X^1 \partial_1 u - \alpha u_t - \zeta u, u_t \rangle + \langle u_t, -X^1 \partial_1 u - \alpha u_t - \zeta u \rangle] dx \\ &+ \int_{\mathbb{S}^1} [\langle g^{01} \partial_1 \partial_t u, u_t \rangle + \langle u_t, g^{01} \partial_1 \partial_t u \rangle] dx \end{split}$$

Note that the last term on the right hand side can be written

$$\int_{\mathbb{S}^1} \partial_1 [g^{01} | u_t |^2] dx = 0,$$

since  $g^{01}$  is independent of the spatial variable. Moreover,

$$-\frac{1}{2}\int_{\mathbb{S}^1} [\langle u_t, \alpha u_t \rangle + \langle \alpha u_t, u_t \rangle] dx \le \|\alpha\| \int_{\mathbb{S}^1} |u_t|^2 dx \le C_{\text{ode}} e^{\beta_{\text{ode}} t} \int_{\mathbb{S}^1} |u_t|^2 dx$$

for  $t \ge 0$ , where we have appealed to (16.23). Similarly, (16.24) yields

$$-\frac{1}{2}\int_{\mathbb{S}^1} [\langle u_t, \zeta u \rangle + \langle \zeta u, u_t \rangle] dx \le \frac{1}{2}C_{\text{ode}}^2 e^{\beta_{\text{ode}}t} \int_{\mathbb{S}^1} [|u_t|^2 + e^{2\beta_{\text{ode}}t}|u|^2] dx$$

for  $t \ge 0$ . Let us focus on

$$-\frac{1}{2} \int_{\mathbb{S}^{1}} [\langle X^{1} \partial_{1} u, u_{t} \rangle + \langle u_{t}, X^{1} \partial_{1} u \rangle] dx$$
  
$$= -\frac{1}{2} \int_{\mathbb{S}^{1}} [\langle (Y^{1} - e^{\beta_{X} t} Y^{1}_{\infty}) e_{1} u, u_{t} \rangle + \langle u_{t}, (Y^{1} - e^{\beta_{X} t} Y^{1}_{\infty}) e_{1} u \rangle] dx \qquad (16.113)$$
  
$$-\frac{1}{2} \int_{\mathbb{S}^{1}} [\langle e^{\beta_{X} t} Y^{1}_{\infty} e_{1} u, u_{t} \rangle + \langle u_{t}, e^{\beta_{X} t} Y^{1}_{\infty} e_{1} u \rangle] dx,$$

where we have used the notation (16.17) and  $e_1 := (g^{11})^{1/2} \partial_1$ . The first term on the right hand side of (16.113) can be estimated by

$$\frac{1}{2}K_X e^{(\beta_X - \eta_X)t} \int_{\mathbb{S}^1} (|u_t|^2 + g^{11} |\partial_1 u|^2) dx$$

for  $t \ge 0$ , where we have appealed to (16.35). Let us focus on the second term on the right hand side of (16.113). It can be estimated by

$$\frac{1}{2} \|Y_{\infty}^{1}\| e^{\beta_{X}t} \int_{\mathbb{S}^{1}} [|u_{t}|^{2} + g^{11}|\partial_{1}u|^{2}] dx.$$

Adding up all the above observations yields

$$\partial_t E_{\text{bas}} \le \|Y_{\infty}^1\| e^{\beta_X t} E_{\text{bas}} + C_{\text{ep}} (1 + e^{\beta_{\text{ode}} t} + e^{(\beta_X - \eta_X)t}) E_{\text{bas}},$$

where the constant  $C_{\rm ep}$  only depends on  $\mathbf{c}_{\rm ep}$  introduced in (16.49). The lemma follows.

#### 

### 16.10 Gevrey classes

Consider the proof of Lemma 16.27, in particular the definition of the sequence  $v_l$ ; cf. (16.97). The fall off of the Fourier coefficients of  $v_l$  is faster than what is required in order for the solutions  $v_l$  to be smooth. It is of interest to introduce terminology quantifying this added fall off. This naturally leads us to the Gevrey classes. If  $\Omega \subseteq \mathbb{R}^d$  is an open set and  $s \geq 1$  is a real number, one way to define the class of Gevrey functions of order s on  $\Omega$  is the following; cf. [41, Definition 1.4.1, p. 19].

**Definition 16.44.** A real valued function f defined on  $\Omega$  is said to be a *Gevrey function of order*  $1 \leq s \in \mathbb{R}$  on  $\Omega$  if  $f \in C^{\infty}(\Omega, \mathbb{R})$  and for each compact set  $K \subset \Omega$ , there are constants R > 0 and C > 0 such that

$$\sup_{x \in K} |\partial^{\alpha} f(x)| \le RC^{|\alpha|} (\alpha!)^s$$

for all d-multiindices  $\alpha$ . The class of Gevrey functions of order  $1 \leq s \in \mathbb{R}$  on  $\Omega$  is denoted  $G^{s}(\Omega)$ .

**Remarks 16.45.** If  $f \in G^1(\Omega)$ , then f is real analytic. Needless to say, the definition can be generalised to functions with values in  $\mathbb{R}^m$  or  $\mathbb{C}^m$ . In this case, we use the notation  $G^s(\Omega, \mathbb{R}^m)$  and  $G^s(\Omega, \mathbb{C}^m)$  respectively.

Here we are interested in functions on  $\mathbb{T}^d$ . We can think of such functions as being defined on  $\mathbb{R}^d$ and being  $2\pi$ -periodic in all of their variables. In that setting, referring to a compact set K, as in Definition 16.44, becomes superfluous. Moreover, in what follows, it turns out to be of interest to generalise the notion of a Gevrey function to the case of 0 < s < 1. This leads us to the following definition.

**Definition 16.46.** A real valued function f defined on  $\mathbb{T}^d$  is said to be a *Gevrey function of order*  $0 < s \in \mathbb{R}$  on  $\mathbb{T}^d$  if  $f \in C^{\infty}(\mathbb{T}^d, \mathbb{R})$  and there are constants R > 0 and C > 0 such that

$$\sup_{x \in \mathbb{T}^d} |\partial^{\alpha} f(x)| \le RC^{|\alpha|} (\alpha!)^s \tag{16.114}$$

for all *d*-multiindices  $\alpha$ . The class of Gevrey functions of order  $0 < s \in \mathbb{R}$  on  $\mathbb{T}^d$  is denoted  $G^s(\mathbb{T}^d)$ .

**Remarks 16.47.** Here we think of functions defined on  $\mathbb{T}^d$  as being defined on  $\mathbb{R}^d$  and being  $2\pi$ -periodic in all of their variables. As before, the definition can be generalised to functions with values in  $\mathbb{R}^m$  or  $\mathbb{C}^m$ , and we use the notation  $G^s(\mathbb{T}^d, \mathbb{R}^m)$  and  $G^s(\mathbb{T}^d, \mathbb{C}^m)$ . Moreover, if  $f \in G^1(\mathbb{T}^d)$ , then f is real analytic. Finally, if  $f \in G^s(\mathbb{T}^d)$ , where 0 < s < 1, then f is entire.

It is sometimes of interest to reformulate the condition (16.114). Clearly, if there are constants R > 0 and C > 0 such that (16.114) holds for all *d*-multiindices  $\alpha$ , then

$$\sup_{x \in \mathbb{T}^d} |\partial^{\alpha} f(x)| \le RC^{|\alpha|} |\alpha|^{s|\alpha|}$$
(16.115)

for all *d*-multiindices  $\alpha$ . However, this implication can also be turned around. More precisely, if there are constants R > 0 and C > 0 such that (16.115) holds for all *d*-multiindices  $\alpha$ , then there are constants  $R_1 > 0$  and  $C_1 > 0$  such that (16.114), with R and C replaced by  $R_1$  and  $C_1$  respectively, holds for all *d*-multiindices  $\alpha$ . A proof of this statement is to be found, e.g., in the proof of [41, Proposition 1.4.2, pp. 19–20]. For completeness, let us, however, note that it follows from Stirling's formula (cf., e.g., [42, (103), p. 194]) and the estimate  $|\alpha|! \leq d^{|\alpha|} \alpha!$ . Note, moreover, that the latter estimate is a consequence of

$$d^{N} = (1 + \dots + 1)^{N} = \sum_{|\alpha|=N} \frac{N!}{\alpha!}.$$

Even though the characterisations (16.114) and (16.115) are of interest, we are here mainly interested in characterisations expressed in terms of the Fourier coefficients of the function f.

**Lemma 16.48.** Let  $f \in C^{\infty}(\mathbb{T}^d, \mathbb{C})$ ,  $0 < s \in \mathbb{R}$  and  $a_n$ ,  $n \in \mathbb{Z}^d$ , be the complex Fourier coefficients of f. Then  $f \in G^s(\mathbb{T}^d, \mathbb{C})$  if and only if there are constants  $C_i > 0$ , i = 1, 2, such that

$$|a_n| \le C_1 \exp\left(-C_2 |n|^{1/s}\right) \tag{16.116}$$

for all  $n \in \mathbb{Z}^d$ .

*Proof.* To begin with, assume that  $f \in G^s(\mathbb{T}^d, \mathbb{C})$  for some s > 0. If  $0 \le k \in \mathbb{Z}$  and  $0 \ne n \in \mathbb{Z}^d$ , then

$$\begin{split} n|^{k}|a_{n}| &\leq d^{k/2}(|n_{1}|^{k} + \dots + |n_{d}|^{k})\frac{1}{(2\pi)^{d}} \left| \int_{\mathbb{T}^{d}} f(x)e^{-in \cdot x} dx \right| \\ &= \frac{d^{k/2}}{(2\pi)^{d}} \sum_{j=1}^{d} \left| \int_{\mathbb{T}^{d}} (-in_{j})^{k} f(x)e^{-in \cdot x} dx \right| \\ &\leq d^{k/2} \sum_{j=1}^{d} \sup_{x \in \mathbb{T}^{d}} |\partial_{j}^{k} f(x)| \leq SB^{k}k^{sk}, \end{split}$$

where we have appealed to (16.115),  $B = \sqrt{dC}$  and  $S = d \cdot R$ ; here C and R are the constants appearing in (16.115). For a given n, it is of interest to choose k so that

$$h(k) := B^k k^{sk} |n|^{-k} = \exp\left(s \cdot (B^{-1}|n|)^{1/s} \frac{k}{(B^{-1}|n|)^{1/s}} \ln\left[\frac{k}{(B^{-1}|n|)^{1/s}}\right]\right)$$

is minimal. Letting  $\xi = k(B^{-1}|n|)^{-1/s}$ , the relevant question is that of minimising  $r(\xi) := \xi \ln \xi$ . Thus the minimum is obtained when  $\xi = \xi_0 := e^{-1}$ , i.e., when  $k = k_0 := (B^{-1}|n|)^{1/s}e^{-1}$ . Of course,  $k_0$  need not be a non-negative integer. Let  $k_1$  be the smallest integer satisfying  $k_1 \ge k_0$  and let  $\xi_1 = k_1(B^{-1}|n|)^{-1/s}$ . Then  $0 \le \xi_1 - \xi_0 \le (B^{-1}|n|)^{-1/s}$ , so that

$$|r(\xi_1) - r(\xi_0)| \le \frac{e}{2} (B^{-1}|n|)^{-2/s}$$

for  $|n| \ge 1$ , where we have used the fact that  $r'(\xi_0) = 0$  and the fact that  $|r''(\xi)| \le e$  for  $\xi \ge \xi_0$ . Thus

$$h(k_1) = \exp\left[s \cdot (B^{-1}|n|)^{1/s} r(\xi_1)\right] = h(k_0) \exp\left(\frac{es}{2} (B^{-1}|n|)^{-1/s}\right) \le Kh(k_0),$$

where K is a constant which is independent of  $n \neq 0$ . Thus

$$|a_n| \le SKh(k_0) = SK \exp\left(-\frac{s}{eB^{1/s}}|n|^{1/s}\right).$$

Assume now that there are constants  $C_i > 0$ , i = 1, 2, such that (16.116) holds for all  $n \in \mathbb{Z}^d$ . Let  $\alpha \neq 0$  be a *d*-multiindex. Then

$$\begin{aligned} |\partial^{\alpha} f(x)| &= \left| \sum_{n \in \mathbb{Z}^{d}} (in)^{\alpha} a_{n} e^{in \cdot x} \right| \\ &\leq \sum_{n \in \mathbb{Z}^{d}} C_{1} \exp\left(-C_{2} |n|^{1/s} / 2\right) |n|^{|\alpha|} \exp\left(-C_{2} |n|^{1/s} / 2\right). \end{aligned}$$
(16.117)

At this stage, it is natural to maximise  $\psi(\xi) = -c_0\xi^{1/s} + k\ln\xi$  for  $\xi > 0$ , where  $c_0 = C_2/2$  and  $k = |\alpha|$ . The maximum is obtained when  $\xi^{1/s} = sk/c_0$ . Thus

$$|n|^k \exp\left(-C_2|n|^{1/s}/2\right) = \exp[\psi(|n|)] \le \left(\frac{s}{ec_0}\right)^{sk} k^{sk}.$$

Combining this estimate with (16.117) yields

$$|\partial^{\alpha} f(x)| \leq \sum_{n \in \mathbb{Z}^d} C_1 \exp\left(-C_2 |n|^{1/s}/2\right) \left(\frac{s}{ec_0}\right)^{sk} k^{sk} \leq RC^{|\alpha|} |\alpha|^{s|\alpha|},$$

for all  $x \in \mathbb{T}^d$ ,  $0 \neq \alpha \in \mathbb{Z}^d$  and a suitable choice of constants R and C.

#### 16.10.1 Preservation of Gevrey class regularity

Next we want to verify that it is meaningful to speak of solutions to (1.2) with Gevrey class regularity (in the case that R = 0). In order for this to be possible, it is, however, necessary to impose conditions on f. This leads us to the following definition.

**Definition 16.49.** Let  $I \subseteq \mathbb{R}$  be an open interval,  $f \in C^{\infty}(\mathbb{T}^d \times I, \mathbb{C}^m)$  and  $0 < s \in \mathbb{R}$ . Then f is said to belong to  $\mathfrak{G}^s_{\text{loc},u}(I, \mathbb{T}^d, \mathbb{C}^m)$  if, for every compact interval  $J \subset I$ , the following holds: there are constants  $C_1 > 0$  and  $C_2 > 0$  (depending on f and J) such that

$$|\hat{f}(n,t)| \le C_1 \exp\left(-C_2 |n|^{1/s}\right)$$

for all  $n \in \mathbb{Z}^d$  and  $t \in J$ , where  $\hat{f}(n,t)$  is the *n*'th (complex) Fourier coefficient of  $f(\cdot,t)$ .

**Remark 16.50.** Note that if  $f \in \mathfrak{G}^s_{\text{loc},u}(I, \mathbb{T}^d, \mathbb{C}^m)$ , then  $f(\cdot, t) \in G^s(\mathbb{T}^d, \mathbb{C}^m)$  for every  $t \in I$ .

**Lemma 16.51.** Let  $1 \leq d, m \in \mathbb{Z}$ , R = 0,  $0 < s \in \mathbb{R}$  and  $I \subseteq \mathbb{R}$  be an open interval. Assume that  $g^{00}, g^{0l}, g^{lj} \in C^{\infty}(I, \mathbb{R})$  are such that for every  $t \in I$ ,  $g^{lj}(t)$  are the components of a positive definite matrix and  $g^{00}(t) = -1$ . Assume, moreover, that  $X^j$ ,  $\alpha, \zeta \in C^{\infty}[I, \mathbf{M}_m(\mathbb{C})]$ . Finally, assume that  $f \in \mathfrak{G}^s_{\mathrm{loc},u}(I, \mathbb{T}^d, \mathbb{C}^m)$ . Let  $t_0 \in I$  and u be the solution to (1.2) (with R = 0) on  $\mathbb{T}^d \times I$  corresponding to initial data  $u(\cdot, t_0), u_t(\cdot, t_0) \in G^s(\mathbb{T}^d, \mathbb{C}^m)$ . Then  $u, u_t \in \mathfrak{G}^s_{\mathrm{loc},u}(I, \mathbb{T}^d, \mathbb{C}^m)$ .

*Proof.* Since the initial data are in  $G^s(\mathbb{T}^d, \mathbb{C}^m)$ , it is clear that the corresponding solution to (1.2) is smooth. Let z(n, t) denote the n'th Fourier coefficient of  $u(\cdot, t)$  and consider (24.16); note that  $\iota = n$  in the present context. Define

$$\begin{split} \mathfrak{H}_n(t) &:= |\dot{z}(n,t)|^2 + [\mathfrak{g}^2(n,t)+1] |z(n,t)|^2, \\ H_n(t) &:= |\dot{z}(n,t)|^2 + \langle n \rangle^2 |z(n,t)|^2, \end{split}$$

where  $\mathfrak{g}(n,t)$  is given by (24.13). In order to prove the statement of the lemma, it is sufficient to estimate  $H_n(t)$  for t in a compact interval J contained in I. In practice we can, without loss of generality, assume that  $J = [t_0, t_1]$  for some  $t_1 > t_0$  (the proof in the case  $J = [t_1, t_0]$  is similar and in the case of a general  $J, J \subseteq [t_a, t_0] \cup [t_0, t_b]$  for suitably chosen  $t_a$  and  $t_b$ ). It is of interest to note that there are constants  $K_j > 0, j = 1, 2$ , depending only on  $g^{ij}(t)$  for  $t \in [t_0, t_1]$ , such that

$$K_1 H_n(t) \le \mathfrak{H}_n(t) \le K_2 H_n(t)$$

for  $t \in [t_0, t_1]$ . It can then be estimated that

$$\dot{\mathfrak{H}}_{n}(t) \leq 2\|\alpha(t)\||\dot{z}(n,t)|^{2} + 2\sum_{j}|n_{j}|\|X^{j}(t)\||z(n,t)||\dot{z}(n,t)| + 2(\|\zeta(t)\|+1)|z(n,t)||\dot{z}(n,t)| + 2|\dot{z}(n,t)||\hat{f}(n,t)| + \dot{g}^{lj}(t)n_{j}n_{l}|z(n,t)|^{2} \leq C\mathfrak{H}_{n}(t) + |\hat{f}(n,t)|^{2}$$
(16.118)

for  $t \in [t_0, t_1]$ , where the constant C > 0 only depends on the coefficients of the equation (1.2) (and not on f). In order to obtain this estimate, we have appealed to the fact that the  $g^{jl}$  are continuously differentiable on I, and that  $\alpha$ , X and  $\zeta$  are continuous. Due to (16.118),

$$\mathfrak{H}_n(t) \le e^{C(t_1 - t_0)} \mathfrak{H}_n(t_0) + \int_{t_0}^{t_1} e^{C(t_1 - \tau)} |\hat{f}(n, \tau)|^2 d\tau$$

for  $t \in [t_0, t_1]$ . Combining this estimate with the assumptions yields the desired conclusion.

#### 16.10.2 Solutions with super exponential growth and Gevrey regularity

The purpose of the present subsection is to make more precise statements concerning the regularity of the solutions constructed in Lemma 16.27.

**Proposition 16.52.** Assume that the conditions of Lemma 16.27 are fulfilled. Let  $\epsilon > 0$ . If  $\beta_X - \beta_1 > 0$ , then the sequence of functions  $v_l$  constructed in Lemma 16.27 can be assumed to such that  $v_l, \partial_t v_l \in \mathfrak{G}^{s_X}_{loc.u}(I, \mathbb{S}^1, \mathbb{C}^m)$  for all l, where

$$s_X := \frac{\beta_X - \beta_1}{\beta_X}.\tag{16.119}$$

In case  $\beta_X - \beta_1 \leq 0$ , one frequency can be chosen, say  $n_a \in \mathbb{Z}$ , such that the sequence of functions  $v_l$  constructed in Lemma 16.27 has the property that if  $z_l(n,t)$  denotes the n'th Fourier coefficient of  $v_l(\cdot,t)$ , then  $z_l(n,t) = 0$  for all t unless  $n = n_a$ .

**Remarks 16.53.** If  $\beta_1 = 0$ , the functions  $v_l$  and  $\partial_t v_l$  are real analytic on constant t hypersurfaces. If  $\beta_1 > 0$ , they are entire on constant t hypersurfaces. In general, the larger the  $\beta_1$ , the better the regularity of the functions  $v_l$ . Considering (16.25), it is clear that improving the lower bound in (16.25) corresponds to improving the regularity of the  $v_l$ . It is of interest to ask if regularity of the form  $\mathfrak{G}^{s_X}_{\text{loc},u}(I,\mathbb{S}^1,\mathbb{C}^m)$ , where  $s_X$  is given by (16.119), is optimal if the parameter  $\beta_1$  appearing in the lower bound in (16.25) is optimal. However, we do not address this question here.

*Proof.* The statement is a consequence of Lemma 16.51 and the proof of Lemma 16.27.  $\Box$ 

# 16.11 Mode analysis

It is of interest to note that in some cases, a mode by mode analysis gives a picture of the future asymptotics which is radically different from that of Lemma 16.27 and Propositions 16.30, 16.34 and 16.37. In the present section, we illustrate this by a sequence of lemmas.

**Lemma 16.54.** Consider (1.2). Assume the associated metric to be such that (M, g) is a canonical separable cosmological model manifold. Assume, moreover, that f = 0. Assume that there is a  $\gamma \geq 0$  and a constant C > 0 such that

$$|g^{0l}(t)| + |g^{jl}(t)|^{1/2} + |a_r^{-1}(t)| + ||X^j(t)||^{1/2} + ||\alpha(t)|| + ||\zeta(t)||^{1/2} \le Ce^{\gamma}$$

for all  $t \ge 0$ , j, l = 1, ..., d, and r = 1, ..., R. Fix  $\iota \in \mathcal{I}_B$ , let  $z(\iota, t)$  be defined by (24.14), where u is a solution to (1.2) with f = 0, and define

$$E_{\iota}(t) := |\dot{z}(\iota, t)|^2 + e^{2\gamma t} |z(\iota, t)|^2.$$

Then there is a constant  $C_{\iota}$ , depending on  $\iota$ , such that

$$E_{\iota}(t) \le \exp\left[C_{\iota}(e^{\gamma t}+t)\right]E_{\iota}(0)$$

for  $t \geq 0$ .

**Remark 16.55.** There is a large class of equations that satisfy both the conditions of Proposition 16.34 and the present lemma with  $\gamma = \beta_{ode}$ . This illustrates that the individual modes can behave quite differently from generic smooth solutions.

*Proof.* Under the assumptions of the lemma, (24.16) with  $\hat{f} = 0$  can be written

$$\ddot{z} + \alpha_{\text{eff}}(t)\dot{z} + \zeta_{\text{eff}}(t)z = 0, \qquad (16.120)$$

where

$$\|\alpha_{\text{eff}}(t)\| + \|\zeta_{\text{eff}}(t)\|^{1/2} \le C_{\iota} e^{\gamma t}$$

for all  $t \ge 0$ . Note, however, that  $C_{\iota}$  depends on the frequency  $\iota$ . It can then be estimated that

$$\partial_t E_{\iota}(t) \leq C_{\iota} e^{\gamma t} E_{\iota}(t)$$

for all  $t \ge 0$ . The lemma follows.

In order to obtain more detailed conclusions, we need to make more detailed assumptions.

**Lemma 16.56.** Consider (1.2). Assume the associated metric to be such that (M, g) is a canonical separable cosmological model manifold. Assume, moreover, that f = 0. Assume that there is a  $\gamma > 0$ ;  $\alpha_{\infty}, \zeta_{\infty} \in \mathbf{M}_m(\mathbb{C})$ , one of which is non-zero; an  $\eta_{cu} > 0$ ; and a constant  $C_{cu} > 0$  such that

$$|g^{0l}(t)| + |g^{jl}(t)|^{1/2} + |a_r^{-1}(t)| + ||X^j(t)||^{1/2} \le C_{\rm cu} e^{(\gamma - \eta_{\rm cu})t} ||e^{-\gamma t} \alpha(t) - \gamma \alpha_{\infty}|| + ||e^{-2\gamma t} \zeta(t) - \gamma^2 \zeta_{\infty}||^{1/2} \le C_{\rm cu} e^{-\eta_{\rm cu}t}$$

for all  $t \ge 0$ , j, l = 1, ..., d, and r = 1, ..., R. Let  $A_{\infty}$  be defined by (16.4) and  $\kappa_1 := \kappa_{\max}(A_{\infty})$ ; cf. Definition 4.3. Fix  $\iota \in \mathcal{I}_B$  and  $\epsilon > 0$ . Then there is a constant  $C_{+,\epsilon} > 0$  such that the following holds. If  $z(\iota, t)$  is a solution to (24.16) with  $\hat{f} = 0$ , then

$$|\dot{z}(\iota,t)| + |z(\iota,t)| \le C_{+,\epsilon} \exp[(\kappa_1 + \epsilon)e^{\gamma t}][|\dot{z}(\iota,0)| + |z(\iota,0)|]$$

for all  $t \geq 0$ .

**Remark 16.57.** The constant  $C_{+,\epsilon}$  only depends on  $C_{cu}$ ,  $\iota$ , the spectrum of the Laplace-Beltrami operator on  $(M_r, g_r)$  for  $r = 1, \ldots, R, A_{\infty}, \eta_{cu}, \gamma$  and  $\epsilon$ .

**Remark 16.58.** There is a large class of equations satisfying the conditions of Propositions 16.34, 16.37 and the present lemma with  $\gamma = \beta_{ode}$  and  $\kappa_1 < 0$ . For such equations, the energy of generic smooth solutions grows super exponentially. However, the energies of the individual modes decay to zero super exponentially. In fact, there is a set of initial data, say  $\mathcal{B}$ , such that the corresponding solutions decay super exponentially and  $\mathcal{B}$  is dense in the set of initial data. In fact, we can take  $\mathcal{B}$  to be the set of initial data with finite frequency content. In that setting, it is thus clear that we cannot improve the conclusions of Proposition 16.34 to say that  $\mathcal{A}$  is open and dense.

*Proof.* As in the proof of Lemma 16.54, the equation (24.16) with  $\hat{f} = 0$  is equivalent to (16.120). However, under the assumptions of the present lemma,

$$\|e^{-\gamma t}\alpha_{\text{eff}}(t) - \gamma \alpha_{\infty}\| + \|e^{-2\gamma t}\zeta_{\text{eff}}(t) - \gamma^{2}\zeta_{\infty}\|^{1/2} \le C_{\iota,\text{cu,spec}}e^{-\eta_{\text{cu}}t}$$

for all  $t \ge 0$ , where  $C_{\iota, \text{cu,spec}}$  only depends on  $\iota$ ,  $C_{\text{cu}}$  and the spectrum of the Laplace-Beltrami operator on the Riemannian manifolds  $(M_r, g_r)$ ,  $r = 1, \ldots, R$ . Given this information, the statement follows from Lemma 16.4.

The case where  $\gamma = 0$  is also of interest, but it requires a slightly different formulation.

**Lemma 16.59.** Consider (1.2). Assume the associated metric to be such that (M, g) is a canonical separable cosmological model manifold. Assume, moreover, that f = 0. Assume that there are  $\alpha_{\infty}, \zeta_{\infty} \in \mathbf{M}_m(\mathbb{C})$ ; an  $\eta_{cu} > 0$ ; and a constant  $C_{cu} > 0$  such that

$$|g^{0l}(t)| + |g^{jl}(t)| + |a_r^{-2}(t)| + ||X^j(t)|| \le C_{\rm cu}e^{-\eta_{\rm cu}t},$$
  
$$||\alpha(t) - \alpha_{\infty}|| + ||\zeta(t) - \zeta_{\infty}|| \le C_{\rm cu}e^{-\eta_{\rm cu}t}$$

for all  $t \ge 0$ , j, l = 1, ..., d, and r = 1, ..., R. Let  $A_{\infty}$  be defined by (16.4),  $\kappa_1 := \kappa_{\max}(A_{\infty})$  and  $d_1 := d_{\max}(A_{\infty}, \kappa_1)$ ; cf. Definition 4.3. Fix  $\iota \in \mathcal{I}_B$ . Then there is a constant  $C_+ > 0$  such that the following holds. If  $z(\iota, t)$  is a solution to (24.16) with  $\hat{f} = 0$ , then

$$|\dot{z}(\iota,t)| + |z(\iota,t)| \le C_+ \langle t \rangle^{d_1 - 1} e^{\kappa_1 t} (|\dot{z}(\iota,0)| + |z(\iota,0)|)$$

for all  $t \geq 0$ .

**Remark 16.60.** The constant  $C_+$  only depends on  $C_{cu}$ ,  $\iota$ , the spectrum of the Laplace-Beltrami operator on  $(M_r, g_r)$  for  $r = 1, \ldots, R$ ,  $A_{\infty}$  and  $\eta_{cu}$ .

*Proof.* As before, (24.16) with  $\hat{f} = 0$  is equivalent to (16.120). However, here

 $\|\alpha_{\text{eff}}(t) - \alpha_{\infty}\| + \|\zeta_{\text{eff}}(t) - \zeta_{\infty}\| \le C_{\text{cu},\iota,\text{spec}} e^{-\eta_{\text{cu}} t}$ 

for all  $t \ge 0$ , where  $C_{\iota,cu,spec}$  only depends on  $\iota$ ,  $C_{cu}$  and the spectrum of the Laplace-Beltrami operator on the Riemannian manifolds  $(M_r, g_r)$ ,  $r = 1, \ldots, R$ . Thus (24.16) can be reformulated to

$$\dot{v}(t) = A_{\infty}v(t) + A_{\text{rem}}(t)v(t),$$
(16.121)

where

$$v(t) := \begin{pmatrix} z(\iota, t) \\ \dot{z}(\iota, t) \end{pmatrix}, \quad \|A_{\operatorname{rem}}(t)\| \le C_{\operatorname{cu},\iota,\operatorname{spec}} e^{-\eta_{\operatorname{cu}} t}$$

for all  $t \ge 0$ . Thus (16.121) is an equation satisfying the conditions described in Subsection 9.1.1 with  $T_{\text{ode}} = 0$ . The desired conclusion is a consequence of Lemma 9.14.
## $\mathbf{Part}~\mathbf{V}$

# Equations with a dominant noisy spatial direction

## Chapter 17

## Terminology and basic estimates

## 17.1 Introduction

The notion of a dominant noisy spatial direction. In Chapters 10 and 12, we consider silent and transparent equations. In the present part of these notes, we turn to so-called noisy equations. We give a formal definition of what we mean by noisy equations in Definition 17.1 below. However, considering the model equation

$$u_{tt} - \sum_{j=1}^{d} g_{\infty}^{jj} e^{2\beta_j t} \partial_j^2 u - \sum_{r=1}^{R} a_{r,\infty}^{-2} e^{2\beta_{\mathrm{Ri},r} t} \Delta_{g_r} u + \alpha_{\infty} u_t + \sum_{j=1}^{d} X_{\infty}^j e^{\beta_j t} \partial_j u + \zeta_{\infty} u = f$$

(where  $g_{\infty}^{jj} > 0$  and  $a_{r,\infty} > 0$ ), the conditions in Definition 17.1 amount to requiring one of the exponents, i.e. one of the elements of

$$\{\beta_1,\ldots,\beta_d,\beta_{\mathrm{Ri},1},\ldots,\beta_{\mathrm{Ri},R}\},\$$

to be strictly positive and strictly larger than all the other elements. In practice, it turns out that when there is one spatial direction which is noisy (meaning that the corresponding exponent is strictly positive) and dominant (meaning that the corresponding exponent is strictly larger than all the other exponents), then the detailed behaviour of the remaining coefficients of the highest order spatial derivatives is not so important; cf. Definition 17.1 below. Turning to the coefficients of the lower order derivatives, the coefficient in front of u does not have to converge, as long as it and its time derivative are bounded. The coefficient in front of  $u_t$  does, however, need to converge exponentially. In case  $g^{jj}$  is the dominant coefficient in front of the second order spatial derivatives, and  $e^{-2\beta_j t}g^{jj}(t)$  converges to a strictly positive number, then we also require  $e^{-\beta_j t}X^j(t)$  to converge. Again, we refer to Definition 17.1 below for the precise list of requirements.

Averaging over oscillations. The main tools we use in order to analyse the asymptotic behaviour of solutions to noisy equations are the ones developed in Part III of these notes. However, in order to be allowed to appeal to the relevant results, we need to make additional assumptions. In particular, we require the equation to be strongly balanced and to have a negligible shift vector field; cf. Definition 17.5 below. Given that the requirements of both Definition 17.1 and Definition 17.5 below are satisfied, it can be verified that the equation is oscillation adapted; cf. Lemma 17.9. Thus the results of Part III apply.

**Defining the asymptotic regime.** In order to be able to draw conclusions, we need to focus on Fourier modes such that the coefficients corresponding to the dominant noisy spatial direction appear in (24.16). This corresponds to focusing on a subset, say  $\mathcal{I}_{B,n}$ , of  $\mathcal{I}_{B}$ ; cf. Definition 17.3 below for a formal definition. However, even for  $\iota \in \mathcal{I}_{B,n}$ , the behaviour of solutions to (24.16) need not initially be determined by the coefficients corresponding to the dominant noisy spatial direction. For that reason, it is necessary to, given  $\iota \in \mathcal{I}_{B,n}$ , determine a late time regime in which these coefficients have a dominant influence. After some preliminary observations concerning the asymptotic behaviour of  $\mathfrak{g}(\iota, t)$ , we, given  $\iota \in \mathcal{I}_{B,n}$ , define a number  $T_n \geq 0$ , cf. Definition 17.12 below, characterising the late time regime.

Behaviour along a time sequence, rough estimates. Once the notion of a late time regime has been defined, we introduce a time sequence  $\{t_k\}$ , given by Definition 15.5 and the requirement that  $t_0 = T_n$ . In order to analyse the behaviour along this time sequence, we appeal to the results of Part III, in particular Lemma 15.11. Considering Lemma 15.11, it is clear that it is of interest to derive detailed estimates for the right hand side of (15.57). This is the purpose of Section 17.4, and leads to an asymptotic expansion for the matrix  $A_k^{\pm}$  appearing in (15.63); cf. Lemma 17.16. Given this asymptotic expansion, we are in a position to analyse the asymptotics of solutions. However, the analysis proceeds in several steps. First, it is convenient to reformulate the equations. This is the subject of Section 17.5, and the goal is to isolate the leading order behaviour. Given this reformulation, we are in a position to derive an estimate for how the energy associated with a mode develops in the late time regime; cf. Section 17.6. Combining this estimate with Lemma 10.7 then yields an estimate for the Sobolev norm of the part of the solution associated with modes  $\iota \in \mathcal{I}_{B,n}$ ; cf. Section 17.7.

### 17.2 Terminology and basic assumptions

We begin by defining the notion of a dominant noisy spatial direction.

**Definition 17.1.** Consider (1.2). Assume the associated metric to be such that (M, g) is a canonical separable cosmological model manifold. Assume, moreover, that there are constants  $\beta_n > 0$ ,  $\eta_{mar} > 0$  and  $C_n > 0$  such that one of the following conditions holds:

• there is one  $j \in \{1, \ldots, d\}$  and a  $g_{\infty}^{jj} \in (0, \infty)$  such that

$$\begin{aligned} |e^{-2\beta_{n}t}g^{jj}(t) - g_{\infty}^{jj}| + |e^{-2\beta_{n}t}\partial_{t}g^{jj}(t) - 2\beta_{n}g_{\infty}^{jj}| \\ + e^{-2\beta_{n}t}\left(\sum_{k,l\neq j}|g^{kl}(t)| + \sum_{l\neq j}|g^{jl}(t)| + \sum_{r=1}^{R}a_{r}^{-2}(t)\right) \\ + e^{-2\beta_{n}t}\left(\sum_{k,l\neq j}|\partial_{t}g^{kl}(t)| + \sum_{l\neq j}|\partial_{t}g^{jl}(t)| + \sum_{r=1}^{R}|\partial_{t}(a_{r}^{-2})(t)|\right) \leq C_{n}e^{-\eta_{mar}t} \end{aligned}$$

for all  $t \ge 0$ , or

• there is one  $r_n \in \{1, \ldots, R\}$  and a  $q_{\infty,n} \in (0, \infty)$  such that

$$\begin{aligned} |e^{-2\beta_{n}t}a_{r_{n}}^{-2}(t) - q_{\infty,n}| + |e^{-2\beta_{n}t}\partial_{t}(a_{r_{n}}^{-2})(t) - 2\beta_{n}q_{\infty,n}| \\ + e^{-2\beta_{n}t}\left(\sum_{k,l=1}^{d} |g^{kl}(t)| + \sum_{r \neq r_{n}} a_{r}^{-2}(t)\right) \\ + e^{-2\beta_{n}t}\left(\sum_{k,l=1}^{d} |\partial_{t}g^{kl}(t)| + \sum_{r \neq r_{n}} |\partial_{t}(a_{r}^{-2})(t)|\right) \leq C_{n}e^{-\eta_{mar}t} \end{aligned}$$

for all  $t \ge 0$ .

Then (1.2) is said to have a dominant noisy spatial direction. In the first case, (1.2) is said to have a dominant noisy spatial torus direction and j is said to correspond to the torus direction. Moreover, the spectrum of the Laplace-Beltrami operator of the flat metric on  $\mathbb{S}^1$  is referred to as the spectrum of the Riemannian manifold corresponding to the dominant noisy spatial direction. In the second case, (1.2) is said to have a dominant noisy spatial generalised direction and  $r_n$  is said to correspond to the generalised direction. Moreover, the spectrum of the Laplace-Beltrami operator of  $(M_{r_n}, g_{r_n})$  is referred to as the spectrum of the Riemannian manifold corresponding to the dominant noisy spatial direction. If (1.2) has a dominant noisy spatial torus direction; *j* corresponds to the torus direction; and there are constants  $\eta_{mn} > 0$ ,  $C_{mn} > 0$  and matrices  $X^j_{\infty}, \alpha_{\infty} \in \mathbf{M}_m(\mathbb{C})$  such that

$$\|e^{-\beta_{\rm n}t}X^{j}(t) - X^{j}_{\infty}\| + \|\alpha(t) - \alpha_{\infty}\| \le C_{\rm mn}e^{-\eta_{\rm mn}t}$$
(17.1)

for all  $t \ge 0$ ; then (1.2) is said to be such that the dominant coefficients are convergent. If (1.2) has a dominant noisy spatial generalised direction; and there are constants  $\eta_{mn} > 0$ ,  $C_{mn} > 0$  and a matrix  $\alpha_{\infty} \in \mathbf{M}_m(\mathbb{C})$  such that

$$\|\alpha(t) - \alpha_{\infty}\| \le C_{\rm mn} e^{-\eta_{\rm mn}t} \tag{17.2}$$

for all  $t \ge 0$ ; then (1.2) is said to be such that the dominant coefficients are convergent.

**Remark 17.2.** Assume that (1.2) is  $C^2$ -balanced with a geometric dominant noisy spatial direction, convergent dominant coefficients and a negligible shift vector field; cf. Definition 6.5. Then the conditions of Definition 17.1 are satisfied. This is a consequence of Lemmas 26.5 and 26.8.

In order not to have to discuss the two cases described in Definition 17.1 separately, it is convenient to introduce the following terminology.

**Definition 17.3.** Assume that (1.2) has a dominant noisy spatial direction. Define

$$\nu_{\mathbf{n}}(\iota) := \lim_{t \to \infty} e^{-\beta_{\mathbf{n}} t} \mathfrak{g}(\iota, t), \tag{17.3}$$

where  $\beta_n$  is given by Definition 17.1. Then the set  $\mathcal{I}_{B,n}$  is defined to consist of the  $\iota \in \mathcal{I}_B$  such that  $\nu_n(\iota) \neq 0$ .

**Remark 17.4.** Due to Definition 17.1, the limit on the right hand side of (17.3) exists.

In what follows, we restrict our attention to balanced equations such that the shift components are negligible.

**Definition 17.5.** Consider (1.2). Assume the associated metric to be such that (M, g) is a canonical separable cosmological model manifold. Then (1.2) is said to be *strongly balanced* if there is a constant C such that

$$\begin{aligned} |\ell(\iota, t)| + |\ell(\iota, t)| + |\sigma(\iota, t)| + |\dot{\sigma}(\iota, t)| + \|X(\iota, t)\| \\ + \|\dot{X}(\iota, t)\| + \|\dot{\alpha}(t)\| + \|\alpha(t)\| + \|\dot{\zeta}(t)\| + \|\zeta(t)\| \le C \end{aligned}$$

for all  $t \ge 0$  and all  $0 \ne \iota \in \mathcal{I}_{B}$ , where  $\ell$ ,  $\sigma$  and X are defined by (10.1). If there are constants C and  $\eta_{sh} > 0$  such that

$$|\sigma(\iota, t)| + |\dot{\sigma}(\iota, t)| \le C e^{-\eta_{\rm sh} t} \tag{17.4}$$

for all  $t \ge 0$  and all  $0 \ne \iota \in \mathcal{I}_{B}$ , then (1.2) is said to have a negligible shift vector field.

**Remark 17.6.** Consider (1.2). Assume the associated metric to be such that (M, g) is a canonical separable cosmological model manifold. Assume, moreover, that (1.2) is  $C^2$ -balanced and that there are constants  $C_{\rm sh}$ ,  $\eta_{\rm sh} > 0$  such that

$$|\chi(t)|_{\bar{q}} + |\dot{\chi}(t)|_{\bar{q}} \le C_{\rm sh}e^{-\eta_{\rm sh}t}$$

for all  $t \ge 0$ . Then the conditions of Definition 17.5 are satisfied. This is a consequence of Lemmas 25.11 and 25.17.

The following objects will play a central role in the analysis

..

**Definition 17.7.** Assume that (1.2) has a dominant noisy spatial direction and is such that the dominant coefficients are convergent. If (1.2) has a dominant noisy spatial torus direction, say j, and  $\iota \in \mathcal{I}_{B,n}$ , define

$$\tilde{X}_{\infty,\mathbf{n}} := (g_\infty^{jj})^{-1/2} X_\infty^j, \quad \mathrm{sgn}_\mathbf{n}(\iota) := \frac{n_j}{|n_j|}.$$

If (1.2) has a dominant noisy spatial generalised direction and  $\iota \in \mathcal{I}_{B,n}$ , define

$$\ddot{X}_{\infty,n} := 0, \quad \operatorname{sgn}_n(\iota) := 1.$$

Define  $\bar{X}_{n}(\iota) := \operatorname{sgn}_{n}(\iota)\tilde{X}_{\infty,n}$ . Moreover,

$$R_{n,\pm}^{+} := \pm \frac{1}{2} (-\alpha_{\infty} + \beta_{n} \mathrm{Id}_{m} + \tilde{X}_{\infty,n}), \quad R_{n,\pm}^{-} := \pm \frac{1}{2} (-\alpha_{\infty} + \beta_{n} \mathrm{Id}_{m} - \tilde{X}_{\infty,n}).$$
(17.5)

Finally,

$$\kappa_{n,\pm} := \max\{\kappa_{\max}(R_{n,\pm}^+), \kappa_{\max}(R_{n,\pm}^-)\},\tag{17.6}$$

$$\kappa_{\mathbf{n}} := \kappa_{\mathbf{n},+} + \kappa_{\mathbf{n},-}. \tag{17.7}$$

**Remark 17.8.** The quantities  $R_{n,\pm}^+$ ,  $R_{n,\pm}^-$  and  $\kappa_{n,\pm}$  are all independent of  $\iota$ .

# 17.3 Oscillation adaptedness and definition of the late time regime

If the conditions of Definition 17.5 are satisfied, then (1.2) is oscillation adapted; cf. Definition 14.1. In fact, the following holds.

**Lemma 17.9.** Assume that (1.2) is strongly balanced and has a negligible shift vector field. Then (1.2) is oscillation adapted; cf. Definition 14.1. Moreover,

$$\mathfrak{f}_X(t) := C, \quad \mathfrak{f}_{ode}(t) := C, \quad \mathfrak{f}_{sh}(t) := C e^{-\eta_{sh} t}, \tag{17.8}$$

where C > 0 only depends on the coefficients of the equation (1.2).

**Remark 17.10.** Dependence on the coefficients includes dependence on the Riemannian manifolds  $(M_r, g_r), r = 1, \ldots, R$ . For future reference, it is of interest to keep in mind that one particular consequence of this observation is that dependence on the minimum of  $\nu_n(\iota)$  for  $\iota \in \mathcal{I}_{B,n}$  is included in dependence on the coefficients of the equation (1.2).

The following observation is also of interest in what follows.

**Lemma 17.11.** Assume that (1.2) has a dominant noisy spatial direction. Then, if  $\iota \in \mathcal{I}_{B,n}$ ,

$$\left|\frac{\mathfrak{g}(\iota,t)}{\nu_{\mathrm{n}}(\iota)e^{\beta_{\mathrm{n}}t}} - 1\right| \le K_{\mathrm{n}}|\nu(\iota)|^{2}e^{-\eta_{\mathrm{mar}}t}$$
(17.9)

for  $t \ge 0$ , where  $K_n \ge 1$  only depends on the coefficients of the equation (1.2).

*Proof.* Since (1.2) has a dominant noisy spatial direction,

$$|\mathfrak{g}^{2}(\iota,t)-\nu_{\mathrm{n}}^{2}(\iota)e^{2\beta_{\mathrm{n}}t}| \leq C|\nu(\iota)|^{2}e^{(2\beta_{\mathrm{n}}-\eta_{\mathrm{mar}})t},$$

where C only depends on the coefficients of the equation (1.2). For  $\iota \in \mathcal{I}_{B,n}$ , this estimate implies that (17.9) holds.

In order to be able to draw conclusions, given assumptions of the type introduced in Definition 17.1, we need to restrict our attention to modes such that  $\nu_n(\iota) \neq 0$ . Moreover, for a given  $\iota \in \mathcal{I}_{B,n}$ , we need to focus on a time interval, say  $[T_n, \infty)$  in which  $\nu_n^2(\iota)e^{2\beta_n t}$  is the dominant part of  $\mathfrak{g}^2(\iota, t)$ . Finally, we wish  $T_n$  to be such that a time sequence of the type introduced in Definition 15.5 can be defined on  $[T_n, \infty)$ ; the reason for this is that we wish to use the results of Chapter 15 in order to analyse the asymptotics. Next we demonstrate that such a  $T_n$  can be defined.

**Definition 17.12.** Assume that (1.2) is strongly balanced, has a negligible shift vector field, a dominant noisy spatial direction and is such that the dominant coefficients are convergent. Let  $K_n$  be the constant appearing in the estimate (17.9). Given  $\iota \in \mathcal{I}_{B,n}$ , define  $T_{n,1}$  by

$$K_{\rm n}|\nu(\iota)|^2 e^{-\eta_{\rm mar}T_{\rm n,1}} = \frac{1}{2}.$$
(17.10)

Let  $\mathfrak{b}$  be the constant appearing in (14.1) and (14.6) (that these estimates hold is a consequence of Lemma 17.9). Given  $\iota \in \mathcal{I}_{B,n}$ , define  $T_{n,2}$  by

$$\nu_{\rm n}(\iota)e^{\beta_{\rm n}T_{{\rm n},2}} = 2\max\{4\pi\mathfrak{b}, 2, \pi(\kappa_{\rm n}+1)\}.$$
(17.11)

Define

$$T_{n} := \max\{T_{n,1}, T_{n,2}, 0\}.$$
(17.12)

Given this terminology, the following statements hold.

**Lemma 17.13.** Assume that (1.2) is strongly balanced, has a negligible shift vector field, a dominant noisy spatial direction and is such that the dominant coefficients are convergent. Given  $\iota \in \mathcal{I}_{B,n}$ , define  $T_n$  by (17.12). Then

$$\mathfrak{g}(\iota, t) \ge \max\{4\pi\mathfrak{b}, 2\} \tag{17.13}$$

for all  $t \geq T_n$ . Moreover,

$$\left|\frac{\mathfrak{g}(\iota,t)}{\nu_{n}(\iota)e^{\beta_{n}t}}-1\right| \leq \frac{1}{2}e^{-\eta_{\max}\bar{t}}, \quad \left|\frac{\nu_{n}(\iota)e^{\beta_{n}t}}{\mathfrak{g}(\iota,t)}-1\right| \leq e^{-\eta_{\max}\bar{t}}$$
(17.14)

for all  $t \geq T_n$ , where  $\bar{t} := t - T_n$ .

**Remark 17.14.** Due to (17.13), it is clear that we can start a sequence  $\{t_k\}$  of the type introduced in Definition 15.5 at  $t_0 := T_n$ . Moreover,  $t_{\text{fin}} = \infty$ , where  $t_{\text{fin}}$  is given by Definition 15.5.

*Proof.* Due to (17.9), (17.10) and (17.12),

$$\frac{1}{2} \le \frac{\mathfrak{g}(\iota, t)}{\nu_{\mathrm{n}}(\iota)e^{\beta_{\mathrm{n}}t}} \le 2 \tag{17.15}$$

for all  $t \ge T_n$ . Combining this estimate with (17.11) and (17.12) yields the conclusion that (17.13) holds for all  $t \ge T_n$ . Combining (17.9), (17.10), (17.12) and (17.15) yields (17.14).

# 17.4 Detailed asymptotics of the matrices appearing in the iteration

Next, let us estimate the constituents of  $R^{ab}_{\text{pre},k}$ ; cf. (15.41) and (15.57).

**Lemma 17.15.** Assume that (1.2) is strongly balanced, has a negligible shift vector field, a dominant noisy spatial direction, and is such that the dominant coefficients are convergent. Let  $\iota \in \mathcal{I}_{B,n}$ and define  $T_n$  by (17.12). Then

$$\left\|\frac{\alpha(t)}{\mathfrak{g}(\iota,t)} - \frac{\alpha_{\infty}}{\nu_{n}(\iota)e^{\beta_{n}t}}\right\| \leq \frac{Ce^{-\eta_{1}\overline{\iota}}}{\nu_{n}(\iota)e^{\beta_{n}t}}$$
(17.16)

for all  $t \ge T_n$ , where C only depends on the coefficients of the equation (1.2) and  $\eta_1 := \min\{\eta_{\max}, \eta_{\max}\}$ . Moreover,

$$\left|\frac{\dot{\ell}(\iota,t)}{\mathfrak{g}(\iota,t)} - \frac{\beta_{\mathrm{n}}}{\nu_{\mathrm{n}}(\iota)e^{\beta_{\mathrm{n}}t}}\right| \le \frac{Ce^{-\eta_{\mathrm{mar}}\bar{t}}}{\nu_{\mathrm{n}}(\iota)e^{\beta_{\mathrm{n}}t}}$$
(17.17)

for all  $t \ge T_n$ , where C only depends on the coefficients of the equation (1.2). Finally

$$\left\|\frac{X(\iota,t)}{\mathfrak{g}(\iota,t)} - \frac{\bar{X}_{\mathrm{n}}(\iota)}{\nu_{\mathrm{n}}(\iota)e^{\beta_{\mathrm{n}}t}}\right\| \le C \frac{e^{-\eta_{2}t}}{\nu_{\mathrm{n}}(\iota)e^{\beta_{\mathrm{n}}t}}$$
(17.18)

for all  $t \geq T_n$ , where C only depends on the coefficients of the equation (1.2) and

$$\eta_2 := \min\{\eta_{\rm mn}, \eta_{\rm mar}/2\}. \tag{17.19}$$

*Proof.* That (17.16) holds is a consequence of (17.1), (17.2), (17.14) and the fact that the equation is strongly balanced.

Estimating  $\dot{\ell}/\mathfrak{g}$ . Note that

$$\begin{aligned} \left| \partial_t \mathfrak{g}^2 - 2\beta_n \nu_n^2 e^{2\beta_n t} \right| &= \left| \sum_{j,l=1}^d \partial_t g^{jl}(t) n_j n_l + \sum_{r=1}^R \partial_t (a_r^{-2})(t) \nu_{r,i_r}^2 - 2\beta_n \nu_n^2 e^{2\beta_n t} \right| \\ &\leq C |\nu|^2 e^{(2\beta_n - \eta_{\max})t} \end{aligned}$$

for all  $t \ge 0$ , where C only depends on the coefficients of the equation (1.2) and we have appealed to the assumptions of Definition 17.1. Thus,

$$\left|\frac{\partial_t \mathfrak{g}^2}{\nu_{\mathbf{n}}^3 e^{3\beta_{\mathbf{n}}t}} - \frac{2\beta_{\mathbf{n}}}{\nu_{\mathbf{n}} e^{\beta_{\mathbf{n}}t}}\right| \le \frac{C}{\nu_{\mathbf{n}} e^{\beta_{\mathbf{n}}t}} |\nu|^2 e^{-\eta_{\mathrm{mar}}t}$$

for all  $t \ge 0$ , where C only depends on the coefficients of the equation (1.2); recall Remark 17.10. Keeping (17.10) and (17.12) in mind, this estimate yields

$$\left|\frac{\partial_t \mathfrak{g}^2}{\nu_n^3 e^{3\beta_n t}} - \frac{2\beta_n}{\nu_n e^{\beta_n t}}\right| \le \frac{C e^{-\eta_{\max} \bar{t}}}{\nu_n e^{\beta_n t}},\tag{17.20}$$

for all  $t \ge T_n$ , where C only depends on the coefficients of the equation (1.2). Estimate

$$\begin{aligned} \left| \frac{\dot{\ell}}{\mathfrak{g}} - \frac{\beta_{\mathrm{n}}}{\nu_{\mathrm{n}} e^{\beta_{\mathrm{n}} t}} \right| &= \left| \frac{1}{2\mathfrak{g}^{3}} \partial_{t} \mathfrak{g}^{2} - \frac{\beta_{\mathrm{n}}}{\nu_{\mathrm{n}} e^{\beta_{\mathrm{n}} t}} \right| \\ &\leq \left| \frac{\nu_{\mathrm{n}}^{3} e^{3\beta_{\mathrm{n}} t}}{2\mathfrak{g}^{3}} \left( \frac{\partial_{t} \mathfrak{g}^{2}}{\nu_{\mathrm{n}}^{3} e^{3\beta_{\mathrm{n}} t}} - \frac{2\beta_{\mathrm{n}}}{\nu_{\mathrm{n}} e^{\beta_{\mathrm{n}} t}} \right) \right| + \left| \frac{\nu_{\mathrm{n}}^{3} e^{3\beta_{\mathrm{n}} t}}{2\mathfrak{g}^{3}} \frac{2\beta_{\mathrm{n}}}{\nu_{\mathrm{n}} e^{\beta_{\mathrm{n}} t}} - \frac{\beta_{\mathrm{n}}}{\nu_{\mathrm{n}} e^{\beta_{\mathrm{n}} t}} \right|. \end{aligned}$$

Due to (17.15) and (17.20), we can estimate the first term on the right hand side. Due to (17.14) and (17.15) we can estimate the second term on the right hand side. Summing up yields (17.17).

Estimating  $X/\mathfrak{g}$ . Finally, let us turn to the expression  $X/\mathfrak{g}$  appearing in (15.41). Assume first that the equation (1.2) has a dominant noisy spatial generalised direction. Fix a  $k \in \{1, \ldots, d\}$ , assume that  $n_k \neq 0$  and estimate

$$\left\| n_k X^k(t) \right\| \le \left\| [g^{kk}(t)n_k^2]^{-1/2} n_k X^k(t) \right\| [g^{kk}(t)n_k^2]^{1/2} \le C[g^{kk}(t)n_k^2]^{1/2}$$

(no summation on k) for all  $t \ge 0$ , where we have appealed to the fact that equation is strongly balanced. Here C only depends on the coefficients of the equation (1.2). This estimate yields

$$\left\|\frac{X(t)}{\mathfrak{g}(t)}\right\| \leq C \sum_{k=1}^{d} \frac{[g^{kk}(t)n_k^2]^{1/2}}{\nu_{\mathrm{n}}e^{\beta_{\mathrm{n}}t}} \frac{1}{\nu_{\mathrm{n}}e^{\beta_{\mathrm{n}}t}} \leq C e^{-\eta_{\mathrm{mar}}\bar{t}/2} \leq C e^{-\eta_{\mathrm{mar}}\bar{t}/2} \qquad (17.21)$$

#### 17.4. DETAILED ASYMPTOTICS OF THE MATRICES

for all  $t \ge T_n$ , where we have appealed to (17.10), (17.12), (17.15) and the assumptions of Definition 17.1. Moreover, C only depends on the coefficients of the equation (1.2). Thus (17.18) holds in this setting.

Assume now that (1.2) has a dominant noisy spatial torus direction, say j. Then

$$\tilde{X}_{\infty,n} = (g_{\infty}^{jj})^{-1/2} X_{\infty}^{j}, \quad \text{sgn}_{n}(\iota) = \frac{n_{j}}{|n_{j}|}.$$

Estimate

$$\frac{n_k X^k(t)}{\mathfrak{g}^2(t)} - \frac{\bar{X}_{\mathbf{n}}}{\nu_{\mathbf{n}} e^{\beta_{\mathbf{n}} t}} \Bigg\| \leq \sum_{k \neq j} \frac{\|n_k X^k(t)\|}{\mathfrak{g}^2(t)} + \Bigg\| \frac{n_j X^j(t)}{\mathfrak{g}^2(t)} - \frac{n_j X^j_\infty e^{\beta_{\mathbf{n}} t}}{\nu_{\mathbf{n}}^2 e^{2\beta_{\mathbf{n}} t}} \Bigg\|$$

(no summation on j). The first term on the right hand side we can estimate by an argument similar to the one leading to (17.21). The second term we can estimate by appealing to (17.1) and (17.14). Summing up yields

$$\left\|\frac{n_k X^k(t)}{\mathfrak{g}^2(t)} - \frac{\bar{X}_n}{\nu_n e^{\beta_n t}}\right\| \le C \frac{e^{-\eta_2 \bar{t}}}{\nu_n e^{\beta_n t}} \tag{17.22}$$

for all  $t \ge T_n$ , where C only depends on the coefficients of the equation (1.2) and  $\eta_2$  is given by (17.19). Thus (17.18) holds in this setting, and the lemma follows.

Next we wish to approximate the matrix  $\Xi_{\text{fin}}^{k,\pm}$  appearing in the statement of Lemma 15.11. Before doing so, recall (17.5) and define, for  $\iota \in \mathcal{I}_{\text{B,n}}$ ,

$$R_{\rm n}(\iota) := \begin{pmatrix} R_{\rm n,+}^{\upsilon(\iota)} & 0\\ 0 & R_{\rm n,+}^{-\upsilon(\iota)} \end{pmatrix},$$
(17.23)

where  $v(\iota) = +$  if  $\operatorname{sgn}_n(\iota) = 1$  and  $v(\iota) = -$  if  $\operatorname{sgn}_n(\iota) = -1$ . Similarly,  $-v(\iota) = -$  if  $\operatorname{sgn}_n(\iota) = 1$ and  $-v(\iota) = +$  if  $\operatorname{sgn}_n(\iota) = -1$ . Note that the only effect of varying  $\iota$  in (17.23) is that the blocks appearing on the right hand side might change place. For future reference, it is also of interest to note that

$$R_{n}(\iota) = \frac{1}{2} \begin{pmatrix} -\alpha_{\infty} + \beta_{n} \mathrm{Id}_{m} + X_{n}(\iota) & 0\\ 0 & -\alpha_{\infty} + \beta_{n} \mathrm{Id}_{m} - \bar{X}_{n}(\iota) \end{pmatrix}.$$

**Lemma 17.16.** Assume that (1.2) is strongly balanced, has a negligible shift vector field, a dominant noisy spatial direction, and is such that the dominant coefficients are convergent. Let  $\iota \in \mathcal{I}_{B,n}$ and define  $T_n$  by (17.12). Let  $\{t_k\}$  be the time sequence given by Definition 15.5 and the condition that  $t_0 := T_n$ . Then

$$\left\| A_k^{\pm} - \operatorname{Id}_{2m} \mp \frac{2\pi}{\nu_{\mathrm{n}}(\iota)e^{\beta_{\mathrm{n}}t_k}} R_{\mathrm{n}}(\iota) \right\| \le \frac{Ce^{-\eta_A \bar{t}_k}}{\nu_{\mathrm{n}}(\iota)e^{\beta_{\mathrm{n}}t_k}}$$
(17.24)

for  $k \ge 0$  in the case of a plus sign and  $k \ge 1$  in the case of a minus sign, where  $A_k^{\pm} := \Xi_{\text{fin}}^{k,\pm}$ ;  $\Xi_{\text{fin}}^{k,\pm}$  is introduced in the statement of Lemma 15.11;  $\bar{t}_k := t_k - T_n$ ;

$$\eta_A := \min\{\eta_{\mathrm{mar}}/2, \eta_{\mathrm{sh}}, \eta_{\mathrm{mn}}, \beta_{\mathrm{n}}\}; \qquad (17.25)$$

and C only depends on the coefficients of the equation (1.2); cf. Remark 17.10.

**Remark 17.17.** In order to obtain the conclusions for  $A_k^-$  for  $k \ge 1$ , we have appealed to Remark 15.14.

Proof. Appealing to Lemma 17.15,

$$\left\| R^{ab}_{\text{pre},k} - \frac{1}{\nu_{n}(\iota)e^{\beta_{n}t_{k}}} \left[ (-1)^{a}\alpha_{\infty} + \beta_{n}\text{Id}_{m} + (-1)^{b}\text{sgn}_{n}(\iota)\tilde{X}_{\infty,n} \right] \right\| \leq \frac{Ce^{-\eta_{2}t_{k}}}{\nu_{n}(\iota)e^{\beta_{n}t_{k}}}$$

for  $t_k \ge T_n$ , where C only depends on the coefficients of the equation (1.2) and  $\eta_2$  is given by (17.19). Combining this observation with (15.57) and (17.23) yields

$$\left\|\Xi_{\text{fin,app}}^{k,\pm} - \text{Id}_{2m} \mp \frac{2\pi}{\nu_{n}(\iota)e^{\beta_{n}t_{k}}} R_{n}(\iota)\right\| \le \frac{Ce^{-\eta_{2}t_{k}}}{\nu_{n}(\iota)e^{\beta_{n}t_{k}}}$$
(17.26)

for  $t_k \ge T_n$ , where C only depends on the coefficients of the equation (1.2). As a final step, we need to appeal to (15.56). We therefore need to estimate the right hand side of this estimate. Note, to begin with, that

$$\varepsilon_k \le C \frac{e^{-\eta_{\rm sh} t_k}}{\mathfrak{g}(t_k)} + \frac{C}{\mathfrak{g}^2(t_k)},\tag{17.27}$$

where C only depends on the coefficients of the equation (1.2). This is due to the fact that  $\varepsilon_k$  is given by the right hand side of (14.33), with  $t_a$  replaced by  $t_k$ , and Lemma 17.9. In particular, it is thus clear that  $\mathfrak{b}$ ,  $\varepsilon_k$  and  $\mathfrak{f}_{\mathrm{sh}}(t_k)$  are bounded by constants depending only on the coefficients of the equation (1.2). Combining this observation with (15.56), (17.26) and (17.27) yields the conclusion of the lemma.

## 17.5 Reformulating the iteration

The purpose of the present section is to reformulate the iteration given by (15.63) in such a way that it is easier to derive estimates.

**Transforming to generalised Jordan normal form.** Fix  $\epsilon > 0$  and let  $T_{n,\epsilon}$  be such that

$$J_{\mathbf{n},\epsilon} := T_{\mathbf{n},\epsilon}^{-1} R_{\mathbf{n}}(\iota) T_{\mathbf{n},\epsilon}$$

is in generalised Jordan normal form with the non-zero off-diagonal components equal to  $\epsilon$ ; cf. Remark 9.10. Due to the remarks made in connection with (17.23), it is clear that  $T_{n,\epsilon}$  can be chosen so that  $J_{n,\epsilon}$  is independent of  $\iota$ . Moreover, even though  $T_{n,\epsilon}$  depends on  $\iota$ , the difference between different  $T_{n,\epsilon}$ 's corresponds to a permutation of the blocks of which  $R_n(\iota)$  consists. Therefore the norms of  $T_{n,\epsilon}$  and  $T_{n,\epsilon}^{-1}$  are independent of  $\iota$ . Introducing

$$A_{k,n,\epsilon}^{\pm} := T_{n,\epsilon}^{-1} A_k^{\pm} T_{n,\epsilon}, \qquad (17.28)$$

the following relations hold:

$$T_{n,\epsilon}^{-1}A_{k_1}^+ \cdots A_{k_0}^+ T_{n,\epsilon} = A_{k_1,n,\epsilon}^+ \cdots A_{k_0,n,\epsilon}^+,$$
(17.29)

$$T_{n,\epsilon}^{-1} A_{k_0}^- \cdots A_{k_1}^- T_{n,\epsilon} = A_{k_0,n,\epsilon}^- \cdots A_{k_1,n,\epsilon}^-.$$
(17.30)

Moreover, (17.24) yields

$$\|A_{k,\mathbf{n},\epsilon}^{\pm} - A_{k,\mathbf{n},\epsilon}^{\mathrm{app},\pm}\| \le \frac{C_{\epsilon}e^{-\eta_{A}t_{k}}}{\nu_{\mathbf{n}}(\iota)e^{\beta_{\mathbf{n}}t_{k}}}$$

where  $C_{\epsilon}$  only depends on  $\epsilon$  and the coefficients of (1.2). Finally,

$$A_{k,\mathbf{n},\epsilon}^{\mathrm{app},\pm} := \mathrm{Id}_{2m} \pm \frac{2\pi}{\nu_{\mathbf{n}}(\iota)e^{\beta_{\mathbf{n}}t_{k}}} J_{\mathbf{n},\epsilon}$$

#### Adapting to the leading order exponential behaviour. Note that

$$e^{-\lambda(t_{1,+}-t_{0,+})}A^+_{k_1,n,\epsilon}\cdots A^+_{k_0,n,\epsilon} = \hat{A}^+_{k_1,n,\epsilon}\cdots \hat{A}^+_{k_0,n,\epsilon},$$
(17.31)

$$e^{-\lambda(t_{0,-}-t_{1,-})}A^{-}_{k_{0},n,\epsilon}\cdots A^{-}_{k_{1},n,\epsilon} = \hat{A}^{-}_{k_{0},n,\epsilon}\cdots \hat{A}^{-}_{k_{1},n,\epsilon},$$
(17.32)

where

$$t_{1,+} := t_{k_1+1}, \quad t_{0,+} := t_{k_0}, \quad t_{1,-} := t_{k_1}, \quad t_{0,-} := t_{k_0-1},$$

 $\lambda$  is a real number and

$$\hat{A}_{k,\mathbf{n},\epsilon}^{\pm} := e^{-\lambda(t_{k\pm1}-t_k)} A_{k,\mathbf{n},\epsilon}^{\pm}.$$
(17.33)

Moreover,

$$\left|e^{-\lambda(t_{k\pm 1}-t_k)}-1+\lambda(t_{k\pm 1}-t_k)\right| \le \frac{C_\lambda}{\mathfrak{g}^2(\iota,t_k)}$$

where the constant  $C_{\lambda}$  only depends on  $\lambda$  and we have appealed to (14.24); for future reference, it is of interest to keep in mind that a similar estimate holds when  $\lambda$  is complex. Combining this estimate with (14.27) and (17.14) yields

$$\left| e^{-\lambda(t_{k\pm 1} - t_k)} - 1 \pm \frac{2\pi\lambda}{\nu_n(\iota)e^{\beta_n t_k}} \right| \le \frac{C_\lambda e^{-\eta_{\max} t_k}}{\nu_n(\iota)e^{\beta_n t_k}} + \frac{C_\lambda}{\mathfrak{g}^2(\iota, t_k)},\tag{17.34}$$

where  $C_{\lambda}$  only depends on  $\lambda$  and the coefficients of (1.2); again, a similar estimate holds when  $\lambda$  is complex. Thus

$$\|\hat{A}_{k,\mathbf{n},\epsilon}^{\pm} - \hat{A}_{k,\mathbf{n},\epsilon}^{\mathrm{app},\pm}\| \le \frac{C_{\epsilon,\lambda}e^{-\eta_A \bar{t}_k}}{\nu_{\mathbf{n}}(\iota)e^{\beta_{\mathbf{n}}t_k}},$$

where

$$\hat{A}_{k,\mathbf{n},\epsilon}^{\mathrm{app},\pm} := \mathrm{Id}_{2m} \pm \frac{2\pi}{\nu_{\mathbf{n}}(\iota)e^{\beta_{\mathbf{n}}t_{k}}}[J_{\mathbf{n},\epsilon} - \lambda \mathrm{Id}_{2m}]$$

Moreover,  $C_{\epsilon,\lambda}$  only depends on the coefficients of (1.2),  $\epsilon$  and  $\lambda$ .

Eliminating the imaginary parts. Before proceeding, it is of interest to eliminate the imaginary parts of the eigenvalues of  $J_{n,\epsilon}$ . Note that  $J_{n,\epsilon} - \lambda Id_{2m}$  consists of generalised Jordan blocks. Assume that the blocks have dimension  $l_1, \ldots, l_D$ . Let  $\zeta_r$  be the imaginary part of the eigenvalue corresponding to the *r*'th generalised Jordan block. Define  $S_k$  and  $\check{A}_{k,n,\epsilon}^{\pm}$  via

$$S_k := \operatorname{diag}\{e^{-i\zeta_1 t_k} \operatorname{Id}_{l_1}, \dots, e^{-i\zeta_D t_k} \operatorname{Id}_{l_D}\}, \quad \check{A}_{k,n,\epsilon}^{\pm} := S_{k\pm 1} \hat{A}_{k,n,\epsilon}^{\pm} S_k^{-1}.$$
(17.35)

Note that  $||S_k|| = ||S_k^{-1}|| = 1$  and that

$$S_{k_1+1}\hat{A}^+_{k_1,n,\epsilon}\cdots\hat{A}^+_{k_0,n,\epsilon}S^{-1}_{k_0}=\check{A}^+_{k_1,n,\epsilon}\cdots\check{A}^+_{k_0,n,\epsilon},$$
(17.36)

$$S_{k_0-1}\hat{A}^{-}_{k_0,n,\epsilon}\cdots\hat{A}^{-}_{k_1,n,\epsilon}S^{-1}_{k_1}=\check{A}^{-}_{k_0,n,\epsilon}\cdots\check{A}^{-}_{k_1,n,\epsilon}.$$
(17.37)

Moreover,

$$\|\check{A}_{k,n,\epsilon}^{\pm} - \check{A}_{k,n,\epsilon}^{\operatorname{app},\pm}\| \le \frac{C_{\epsilon,\lambda}e^{-\eta_A t_k}}{\nu_n(\iota)e^{\beta_n t_k}}$$
(17.38)

for all  $k \ge 0$ , where

$$\check{\mathbf{A}}_{k,\mathbf{n},\epsilon}^{\mathrm{app},\pm} := \mathrm{Id}_{2m} \pm \frac{2\pi}{\nu_{\mathbf{n}}(\iota)e^{\beta_{\mathbf{n}}t_{k}}} [\mathrm{Re}\{J_{\mathbf{n},\epsilon}\} - \lambda \mathrm{Id}_{2m}];$$
(17.39)

this is a consequence of an estimate similar to (17.34). Finally, the constant  $C_{\epsilon,\lambda}$  appearing in (17.38) only depends on  $\epsilon$ ,  $\lambda$  and the coefficients of the equation (1.2).

## 17.6 A rough mode estimate

As a first application of the framework developed above, we derive an estimate for the growth of the energy of solutions in the late time regime.

**Lemma 17.18.** Assume that (1.2) is strongly balanced, has a negligible shift vector field, a dominant noisy spatial direction, and is such that the dominant coefficients are convergent. Then there is a constant C, depending only on the coefficients of the equation (1.2), such that the following holds. Let  $\iota \in \mathcal{I}_{B,n}$  and define  $T_n$  by (17.12). If z is a solution to (24.16) and w is defined by (15.1) and (14.7), then

$$|w(\iota, t)| \leq C \langle t - t_a \rangle^{d_{n,+} - 1} e^{\kappa_{n,+}(t - t_a)} |w(\iota, t_a)| + \int_{t_a}^t C \langle t - t' \rangle^{d_{n,+} - 1} e^{\kappa_{n,+}(t - t')} |\hat{f}(\iota, t')| dt'$$
(17.40)

for all  $T_n \leq t_a \leq t$ . Here  $\kappa_{n,+}$  is defined by (17.6) and  $d_{n,+} := d_{\max}(R_n(\iota), \kappa_{n,+})$ , where  $R_n(\iota)$  is defined by (17.23); cf. Definition 4.3.

**Remark 17.19.** The constant  $d_{n,+}$  is independent of  $\iota \in \mathcal{I}_{B,n}$ ; cf. the comments made in connection with (17.23).

**Remark 17.20.** Dependence on the coefficients includes dependence on the Riemannian manifolds  $(M_r, g_r), r = 1, \ldots, R$ ; cf. Remark 17.10.

*Proof.* Reducing the problem. Recall the observations made in Subsection 15.1.1. Due to these observations, (17.40) is a consequence of the estimate

$$\|\Phi_{\iota}(t;t')\| \le C\langle t-t'\rangle^{d_{n,+}-1} e^{\kappa_{n,+}(t-t')}$$
(17.41)

for all  $T_n \leq t' \leq t$ , where C only depends on the coefficients of the equation (1.2). The remainder of the proof is devoted to demonstrating (17.41). In order to prove this estimate, it is sufficient to prove that

$$|w(\iota, t)| \le C \langle t - t' \rangle^{d_{n, +} - 1} e^{\kappa_{n, +} (t - t')} |w(\iota, t')|$$
(17.42)

for every solution to (24.16) with  $\hat{f} = 0$  and all  $T_n \leq t' \leq t$ , where C has the same dependence as in the case of (17.41).

**Notation.** In the remainder of the proof, we use the notation introduced in Section 17.5. Recall (17.6) and note that  $\kappa_{n,+} = \kappa_{\max}[R_n(\iota)]$ , where  $\kappa_{n,+}$  is independent of  $\iota$ ; cf. Remark 17.8. Let, moreover,

$$J_{R,n,\epsilon} := \operatorname{Re}\{J_{n,\epsilon}\} - \kappa_{n,+} \operatorname{Id}_{2m}, \qquad (17.43)$$

 $J_{\rm tr}$  be the matrix obtained from  $J_{R,n,\epsilon}$  by setting the generalised Jordan blocks with zero eigenvalue to zero, and  $J_{\rm diff} := J_{R,n,\epsilon} - J_{\rm tr}$ . Note that the diagonal components of  $J_{R,n,\epsilon}$  are elements of  $[-\kappa_n, 0]$ , where  $\kappa_n$  is introduced in (17.7). Finally, comparing with Section 17.5, we take  $\lambda$  to equal  $\kappa_{n,+}$  from now on.

Estimates for times belonging to the sequence, preliminaries. Fix an integer  $k_0 \ge 0$ . To begin with, we are interested in proving that (17.42) holds for  $t' = t_{k_0}$  and  $t = t_{k+1}$ , where  $0 \le k_0 \le k$ . Consider (15.63), where  $\psi_k = w_{\text{fin},k}$ ,  $A_k^{\pm} = \Xi_{\text{fin}}^{k,\pm}$  and  $\hat{F}_k$  is replaced by 0. This equality implies that

$$\rho_{k+1} = \bar{A}^+_{k,n,\epsilon} \rho_k, \tag{17.44}$$

where

$$\rho_k := e^{-\kappa_{n,+}t_k} e^{-J_{\text{diff}}\tilde{t}_k} S_k T_{n,\epsilon}^{-1} \psi_k, \qquad (17.45)$$
$$\bar{A}_{k,n,\epsilon}^+ := e^{-J_{\text{diff}}\tilde{t}_{k+1}} \check{A}_{k,n,\epsilon}^+ e^{J_{\text{diff}}\tilde{t}_k},$$

 $\tilde{t}_k := t_k - t_{k_0}, \lambda = \kappa_{n,+}$  and  $S_k$  and  $T_{n,\epsilon}$  are defined as above. Thus, if  $k \ge k_0$ ,

$$\rho_{k+1} = \bar{A}^+_{k,n,\epsilon} \cdots \bar{A}^+_{k_0,n,\epsilon} \rho_{k_0}. \tag{17.46}$$

Compute

$$\bar{A}_{k,\mathbf{n},\epsilon}^{+} = e^{-J_{\text{diff}}\tilde{t}_{k+1}} (\check{A}_{k,\mathbf{n},\epsilon}^{+} - \check{A}_{k,\mathbf{n},\epsilon}^{\text{app},+}) e^{J_{\text{diff}}\tilde{t}_{k}} + e^{-J_{\text{diff}}(t_{k+1}-t_{k})} \check{A}_{k,\mathbf{n},\epsilon}^{\text{app},+};$$
(17.47)

#### 17.6. A ROUGH MODE ESTIMATE

note that  $J_{\rm diff}$  and  $\check{A}_{k,n,\epsilon}^{\rm app,+}$  commute. On the other hand,

$$e^{-J_{\text{diff}}(t_{k+1}-t_k)}\check{A}_{k,n,\epsilon}^{\text{app},+} = \check{A}_{k,n,\epsilon}^{\text{app},+} - J_{\text{diff}}(t_{k+1}-t_k) + \frac{2\pi}{\nu_n(\iota)e^{\beta_n t_k}}J_{R,n,\epsilon}[e^{-J_{\text{diff}}(t_{k+1}-t_k)} - \text{Id}_{2m}] + [e^{-J_{\text{diff}}(t_{k+1}-t_k)} - \text{Id}_{2m} + J_{\text{diff}}(t_{k+1}-t_k)].$$

Introducing

$$\bar{A}_{k,\mathbf{n},\epsilon}^{\mathrm{app},+} := \mathrm{Id}_{2m} + \frac{2\pi}{\nu_{\mathbf{n}}(\iota)e^{\beta_{\mathbf{n}}t_{k}}} J_{\mathrm{tr}}$$
(17.48)

and appealing to (14.27) and (17.14) yields

$$\|e^{-J_{\text{diff}}(t_{k+1}-t_k)}\check{A}_{k,\mathbf{n},\epsilon}^{\text{app},+} - \bar{A}_{k,\mathbf{n},\epsilon}^{\text{app},+}\| \le \frac{C_{\epsilon}e^{-\eta_A t_k}}{\nu_{\mathbf{n}}(\iota)e^{\beta_{\mathbf{n}}t_k}}$$

for all  $k \ge k_0$ , where  $C_{\epsilon}$  only depends on  $\epsilon$  and the coefficients of the equation (1.2). Combining this estimate with (17.38) and (17.47) yields

$$\|\bar{A}_{k,n,\epsilon}^{+} - \bar{A}_{k,n,\epsilon}^{\text{app},+}\| \le \frac{C_{\epsilon}e^{-\eta_{A}t_{k}}}{\nu_{n}(\iota)e^{\beta_{n}t_{k}}} \langle \tilde{t}_{k} \rangle^{N}$$
(17.49)

for all  $k \ge k_0$ , where  $C_{\epsilon}$  only depends on  $\epsilon$  and the coefficients of the equation (1.2), and N is a non-negative integer depending only on m.

Estimating the matrix products. Consider (17.48). By construction, all the diagonal elements of the second term on the right hand side are non-positive. Moreover, they are strictly smaller than 1 in absolute value due to (17.11). Finally, all non-trivial generalised Jordan blocks in  $J_{\rm tr}$  have the property that the corresponding diagonal components are strictly negative. By choosing  $\epsilon$  sufficiently small, the bound depending only on the coefficients of the equation, it can be ensured that

$$\|\bar{A}_{k,n,\epsilon}^{\operatorname{app},+}\| \le 1 \tag{17.50}$$

for all  $k \ge k_0$ . Fixing such an  $\epsilon$ , the estimates below are independent of  $\epsilon$ . One particular consequence of (17.50) is that for  $0 \le k_0 \le k_1$ ,

$$\|\bar{A}_{k_{1},n,\epsilon}^{+}\cdots\bar{A}_{k_{0},n,\epsilon}^{+}\| \leq (1+\|\bar{A}_{k_{1},n,\epsilon}^{+}-\bar{A}_{k_{1},n,\epsilon}^{app,+}\|)\cdots(1+\|\bar{A}_{k_{0},n,\epsilon}^{+}-\bar{A}_{k_{0},n,\epsilon}^{app,+}\|).$$

Since  $\ln(1+x) \le x$  for  $x \ge 0$ , the logarithm of the right hand side is bounded by

$$\sum_{k=k_0}^{k_1} \|\bar{A}_{k,n,\epsilon}^+ - \bar{A}_{k,n,\epsilon}^{\text{app},+}\| \le \sum_{k=k_0}^{k_1} \frac{Ce^{-\eta_A \bar{t}_k}}{\nu_n(\iota)e^{\beta_n t_k}} \langle \tilde{t}_k \rangle^N \le \int_{t_{k_0}}^{t_{k_1+1}} C \langle t - t_{k_0} \rangle^N e^{-\eta_A \bar{t}} dt \le Ce^{-\eta_A(t_{k_0} - T_n)},$$

where C only depends on the coefficients of the equation (1.2). In order to obtain this estimate, we appealed to Lemma 14.4, (17.15) and (17.49). In particular, for  $0 \le k_0 \le k$ ,

$$\|\bar{A}_{k,n,\epsilon}^+\cdots\bar{A}_{k_0,n,\epsilon}^+\| \le C,\tag{17.51}$$

where C only depends on the coefficients of the equation (1.2). Combining this estimate with (17.46) yields

$$|\rho_{k+1}| \le C|\rho_{k_0}| \tag{17.52}$$

for all  $k \ge k_0$ , where the constant C has the same dependence as in the case of (17.51). Returning to the definition of  $\rho_k$  in terms of  $\psi_k$ , cf. (17.45), and the definition of  $\psi_k = w_{\text{fin},k}$ , cf. (15.53), this estimate yields

$$|w(\iota, t_{k+1})| \le C e^{\kappa_{n,+} t_{k+1}} \langle \tilde{t}_{k+1} \rangle^{d_{n,+}-1} |w(\iota, t_{k_0})|, \qquad (17.53)$$

where the constant C has the same dependence as in the case of (17.51). Here  $d_{n,+}$  is the largest dimension of a Jordan block of  $R_n(\iota)$  corresponding to an eigenvalue with real part  $\kappa_{n,+}$ . Thus (17.42) holds if  $t' = t_{k_0}$  and  $t = t_{k+1}$ .

**General** t' and t. In general,  $t' \in [t_{k_0}, t_{k_0+1}]$  and  $t \in [t_k, t_{k+1}]$  for some  $0 \le k_0 \le k$ . Appealing to Lemma 10.7 in order to estimate the evolution between t' and  $t_{k_0}$  and between  $t_k$  and t yields the conclusion that (17.42) holds for all  $t_0 \le t' \le t$ . Moreover, C has the same dependence as in the case of (17.51).

## 17.7 A rough Sobolev estimate

Assume that (1.2) is strongly balanced, has a negligible shift vector field, a dominant noisy spatial direction, and is such that the dominant coefficients are convergent. Due to, e.g., Lemma 17.18 above, it is then clear that we can draw conclusions concerning the behaviour for  $\iota \in \mathcal{I}_{B,n}$ . What happens for  $\iota \notin \mathcal{I}_{B,n}$  is, however, less clear. For this reason, it is natural to divide (1.2) into two parts, one corresponding to  $\mathcal{I}_{B,n}$ , and one corresponding to the complement of  $\mathcal{I}_{B,n}$ . If  $f \in C^{\infty}(M, \mathbb{C}^m)$ , it is therefore convenient to introduce the notation

$$f_{\mathbf{n}}(p,t) := \sum_{\iota \in \mathcal{I}_{\mathbf{B}_{\mathbf{n}}}} \hat{f}(\iota,t)\varphi_{\iota}(p), \qquad (17.54)$$

where  $\varphi_{\iota}$  and  $\hat{f}$  are defined by (24.6) and (24.15) respectively. Noting that  $\mathfrak{E}_s$  is defined by (24.19), we have the following result.

**Lemma 17.21.** Assume that (1.2) is strongly balanced, has a negligible shift vector field, a dominant noisy spatial direction, and is such that the dominant coefficients are convergent. Then there are constants C and  $s_n$ , depending only on the coefficients of the equation (1.2), such that if u is a solution to (1.2), then

$$\mathfrak{E}_{s}^{1/2}[u_{n}](t) \leq C\langle t \rangle^{d_{n,+}-1} e^{\kappa_{n,+}t} \mathfrak{E}_{s+s_{n}}^{1/2}[u_{n}](0) + \int_{0}^{t} C\langle t-t' \rangle^{d_{n,+}-1} e^{\kappa_{n,+}(t-t')} \|f_{n}(\cdot,t')\|_{(s+s_{n})} dt'$$
(17.55)

for all  $t \ge 0$  and  $s \in \mathbb{R}$ , where  $f_n$  and  $u_n$  are defined by (17.54) and the adjacent text. Finally,  $\kappa_{n,+}$  and  $d_{n,+}$  are defined as in the statement of Lemma 17.18.

**Remark 17.22.** Dependence on the coefficients includes dependence on the Riemannian manifolds  $(M_r, g_r), r = 1, \ldots, R$ ; cf. Remark 17.10.

**Remark 17.23.** Assume that (1.2) is  $C^2$ -balanced with a geometric dominant noisy spatial direction, convergent dominant coefficients and a negligible shift vector field; cf. Definition 6.5. Then the assumptions of the lemma are satisfied; cf. Remarks 17.2 and 17.6.

*Proof.* Due to Lemma 17.18, we know that

$$|w(\iota,t)| \le C\langle \bar{t} \rangle^{d_{n,+}-1} e^{\kappa_{n,+}\bar{t}} |w(\iota,T_n)| + \int_{T_n}^t C\langle t-t' \rangle^{d_{n,+}-1} e^{\kappa_{n,+}(t-t')} |\hat{f}(\iota,t')| dt'$$
(17.56)

for all  $\iota \in \mathcal{I}_{B,n}$  and  $t \ge T_n$ , where  $\overline{t} := t - T_n$  and C only depends on the coefficients of the equation (1.2). Since  $\mathfrak{g}(\iota, t) \ge 2$  for  $t \ge T_n$ , cf. (17.13), it is clear that  $\mathcal{E}^{1/2}(\iota, t)$  and  $|w(\iota, t)|$  are equivalent for  $t \ge T_n$ , the constants of equivalence being numerical. One particular consequence of (17.56) is thus that

$$\mathcal{E}^{1/2}(\iota,t) \le C\langle \bar{t} \rangle^{d_{n,+}-1} e^{\kappa_{n,+}\bar{t}} \mathcal{E}^{1/2}(\iota,T_n) + \int_{T_n}^t C\langle t-t' \rangle^{d_{n,+}-1} e^{\kappa_{n,+}(t-t')} |\hat{f}(\iota,t')| dt'$$
(17.57)

for all  $\iota \in \mathcal{I}_{B,n}$  and  $t \ge T_n$ , where C only depends on the coefficients of the equation (1.2). On the other hand, Lemma 10.7 yields

$$\mathcal{E}^{1/2}(\iota, t) \le e^{\eta_{\text{bal}} t} \mathcal{E}^{1/2}(\iota, 0) + \int_0^t e^{\eta_{\text{bal}}(t - t')} |\hat{f}(\iota, t')| dt'$$
(17.58)

for all  $t \ge 0$ , where  $\eta_{\text{bal}} \ge 0$  only depends on the coefficients of the equation (1.2). Letting

$$\eta_{\mathrm{mod}} := \max\{\eta_{\mathrm{bal}} - \kappa_{\mathrm{n},+}, 0\},\$$

it is of interest to estimate

$$\eta_{\text{mod}} T_{\text{n}} \le \frac{2\eta_{\text{mod}}}{\eta_{\text{mar}}} \ln |\nu(\iota)| + C,$$

where C only depends on the coefficients of the equation (1.2); cf. Definition 17.12. Combining this estimate with (17.58) yields

$$\mathcal{E}^{1/2}(\iota,t) \le C e^{\kappa_{n,+}t} \langle \nu(\iota) \rangle^{s_n} \mathcal{E}^{1/2}(\iota,0) + C \langle \nu(\iota) \rangle^{s_n} \int_0^t e^{\kappa_{n,+}(t-t')} |\hat{f}(\iota,t')| dt'$$
(17.59)

for all  $0 \le t \le T_n$ , where  $s_n := 2\eta_{mod}/\eta_{mar}$  and C only depends on the coefficients of the equation (1.2). Combining (17.57) and (17.59) yields

$$\mathcal{E}_{s}^{1/2}(\iota,t) \leq C\langle t \rangle^{d_{n,+}-1} e^{\kappa_{n,+}t} \mathcal{E}_{s+s_{n}}^{1/2}(\iota,0) + C\langle \nu(\iota) \rangle^{s+s_{n}} \int_{0}^{t} \langle t-t' \rangle^{d_{n,+}-1} e^{\kappa_{n,+}(t-t')} |\hat{f}(\iota,t')| dt'$$

for all  $t \ge 0$ ,  $s \in \mathbb{R}$  and  $\iota \in \mathcal{I}_{B,n}$ , where C only depends on the coefficients of the equation (1.2). Appealing to Minkowski's inequality yields the conclusion of the lemma.

## Chapter 18

## Asymptotics for one mode and late times

Given the framework developed in the previous chapter, we are in a position to derive information concerning the asymptotics. In the present chapter, we focus on one single mode  $\iota \in \mathcal{I}_{B,n}$ . Moreover, we only derive conclusions in the late time regime  $[T_n, \infty)$ , where  $T_n$  is introduced in Definition 17.12. To begin with, we derive asymptotic information by analysing how the solution behaves along the time sequence  $\{t_k\}$ ; this is the subject of Section 18.1. In Section 18.2 we then turn to the problem of specifying the asymptotic data. However, the considerations are, again, restricted to a time sequence in the interval  $[T_n, \infty)$ . Finally, in Section 18.3, we extend the estimates derived along the time sequence to asymptotic estimates valid for all  $t \geq T_n$ .

### 18.1 Asymptotics along a time sequence

Lemma 17.18 yields an estimate for the growth of the energy of one mode. In what follows, we want to derive more detailed information. On the interval  $[T_n, \infty)$ , it is natural to carry out the analysis in two steps. First, we consider how the solution evolves along the sequence  $\{t_k\}$ . Second, we analyse the development during one period. In this section begin by analysing the asymptotics along the time sequence.

**Lemma 18.1.** Assume that (1.2) is strongly balanced, has a negligible shift vector field, a dominant noisy spatial direction, and is such that the dominant coefficients are convergent. Let  $\iota \in \mathcal{I}_{B,n}$  and define  $T_n$  by (17.12). Let  $\{t_k\}$  be the time sequence given by Definition 15.5 and the condition that  $t_0 := T_n$ . Let  $\eta_A$  be defined by (17.25) and assume that there is an  $0 < \eta_B \leq \eta_A$  such that

$$\int_0^\infty e^{-(\kappa_{\mathrm{n},+}-\eta_B)t} |\hat{f}(\iota,t)| dt < \infty,$$

where  $\kappa_{n,+}$  is defined by (17.6). Then there is a constant  $C_B$ , depending only on  $\eta_B$  and the coefficients of the equation (1.2); and a non-negative integer N, depending only on m, such that the following holds. If z is a solution to (24.16), then there is a  $\psi_{\infty}$  belonging to the first generalised eigenspace of the  $\eta_B$ ,  $R_n(\iota)$ -decomposition of  $\mathbb{C}^{2m}$  (cf. Definition 9.4) and satisfying

$$|e^{R_{n}(\iota)T_{n}}\psi_{\infty}| \leq C_{B}\left(|w(\iota,T_{n})| + e^{\kappa_{n,+}T_{n}}\int_{T_{n}}^{\infty} e^{-(\kappa_{n,+}-\eta_{B})t}|\hat{f}(\iota,t)|dt\right),$$
(18.1)

such that

$$\left| \begin{pmatrix} \mathfrak{g}(\iota, t_k) z(\iota, t_k) \\ \dot{z}(\iota, t_k) \end{pmatrix} - \exp\left(i \int_0^{t_k} \sigma(\iota, t') \mathfrak{g}(\iota, t') dt'\right) T_{\mathrm{pre}, k}^{-1} [D_k(\iota)]^{-1} e^{R_n(\iota) t_k} \psi_{\infty} \right| \\
\leq C_B \langle \bar{t}_k \rangle^N e^{(\kappa_{n, +} - \eta_B) \bar{t}_k} \left( |w(\iota, T_n)| + e^{\kappa_{n, +} T_n} \int_0^\infty e^{-(\kappa_{n, +} - \eta_B) t} |\hat{f}(\iota, t)| dt \right) \tag{18.2}$$

holds for all  $k \ge 0$ , where  $\bar{t}_k := t_k - t_0$ . In these estimates, w is given by (15.1) and (14.7);  $\mathfrak{g}$  is given by (24.13);  $\sigma$  is given by (10.1);  $T_{\text{pre},k}$  is given by (15.34);  $D_k(\iota)$  is given by (15.52); and  $R_n(\iota)$  is given by (17.23).

Remark 18.2. The norms of the matrix

$$\exp\left(i\int_0^{t_k}\sigma(t')\mathfrak{g}(t')dt'\right)T_{\mathrm{pre},k}^{-1}D_k^{-1}$$

and its inverse are bounded by constants depending only on the coefficients of the equation (1.2). In particular,  $\psi_{\infty}$  is thus uniquely determined by the estimate (18.2) and the condition that it belong to the first generalised eigenspace of the  $\eta_B$ ,  $R_n(\iota)$ -decomposition of  $\mathbb{C}^{2m}$ .

**Remark 18.3.** In case all the Jordan blocks of  $R_n(\iota)$  are trivial, the *N* appearing in (18.2) can be replaced by 1. Moreover, if the first generalised eigenspace of the  $\eta_B$ ,  $R_n(\iota)$ -decomposition of  $\mathbb{C}^{2m}$  equals  $\mathbb{C}^{2m}$ , then the *N* appearing in (18.2) can be replaced by  $d_{n,+} - 1$ , where  $d_{n,+} := d_{\max}(R_n(\iota), \kappa_{n,+})$ .

**Remark 18.4.** Dependence on the coefficients includes dependence on the Riemannian manifolds  $(M_r, g_r), r = 1, \ldots, R$ ; cf. Remark 17.10.

*Proof.* The proof is based on the reformulation of the iteration introduced in Section 17.5. Moreover, we here let  $\lambda = \kappa_{n,+}$ . Correspondingly, we introduce

$$\zeta_k := e^{-\kappa_{\mathbf{n},+}t_k} S_k T_{\mathbf{n},\epsilon}^{-1} \psi_k, \tag{18.3}$$

$$\check{F}_k(t) := e^{-\kappa_{\mathbf{n},+}t_k} S_k T_{\mathbf{n},\epsilon}^{-1} \hat{F}_k(t), \qquad (18.4)$$

where  $\psi_k = w_{\text{fin},k}$  and  $\hat{F}_k$  is given by (15.55). As opposed to the proof of Lemma 17.18, the value of  $\epsilon > 0$  is not important in the present proof. We therefore set it to 1 here. Nevertheless, we keep the  $\epsilon$ 's that appear in the notation of Section 17.5 in order not to have to introduce additional symbols. Given the definitions (18.3) and (18.4), equation (15.63) yields

$$\zeta_{k+1} = \check{A}_{k,n,\epsilon}^{+} \zeta_{k} + \check{A}_{k,n,\epsilon}^{+} \int_{t_{k}}^{t_{k+1}} \check{F}_{k}(t) dt.$$
(18.5)

Due to (17.38) and (17.39), it is clear that

$$\|\check{A}_{k,\mathbf{n},\epsilon}^{+} - \check{A}_{k,\mathbf{n},\epsilon}^{\mathrm{app},+}\| \le \frac{Ce^{-\eta_{A}t_{k}}}{\nu_{\mathbf{n}}(\iota)e^{\beta_{\mathbf{n}}t_{k}}},\tag{18.6}$$

where C only depends on the coefficients of the equation (1.2);  $\eta_A$  is given by (17.25);  $\bar{t}_k := t_k - T_n = t_k - t_0$ ; and

$$\check{A}_{k,\mathbf{n},\epsilon}^{\mathrm{app},+} := \mathrm{Id}_{2m} + \frac{2\pi}{\nu_{\mathbf{n}}(\iota)e^{\beta_{\mathbf{n}}t_k}} J_{R,\mathbf{n},\epsilon},$$

where  $J_{R,n,\epsilon}$  is given by (17.43). Moreover, the estimate (17.40) implies

$$|\zeta_k| \le C \langle \bar{t}_k \rangle^{d_{n,+}-1} |\zeta_0| + \int_{t_0}^{t_k} C \langle t_k - t' \rangle^{d_{n,+}-1} e^{-\kappa_{n,+}t'} |\hat{f}(\iota, t')| dt'$$
(18.7)

#### 18.1. ASYMPTOTICS ALONG A TIME SEQUENCE

for all  $k \ge 0$ , where *C* only depends on the coefficients of the equation (1.2). Let  $\eta_B$  be as in the statement of the lemma. In the analysis of the future asymptotics of  $\zeta_k$ , it is convenient to divide the components of  $\zeta_k$  into  $\zeta_{k,a}$  and  $\zeta_{k,b}$ . Here  $\zeta_{k,a}$  and  $\zeta_{k,b}$  denote the first and second component in the  $\eta_B$ ,  $J_{R,n,\epsilon}$ -decomposition of  $\zeta_k$  respectively; cf. Definition 9.4. Similarly, we denote the first and second blocks of the  $\eta_B$ ,  $J_{R,n,\epsilon}$ -decomposition of  $J_{R,n,\epsilon}$  by  $J_{R,a}$  and  $J_{R,b}$  respectively. Let us start by considering  $\zeta_{k,a}$ .

Estimating the first component. With notation as described above, (18.5) yields

$$e^{-J_{R,a}\bar{t}_{k+1}}\zeta_{k+1,a} = e^{-J_{R,a}\bar{t}_{k+1}}[\check{A}^+_{k,n,\epsilon}\zeta_k]_a + e^{-J_{R,a}\bar{t}_{k+1}}\int_{t_k}^{t_{k+1}}[\check{A}^+_{k,n,\epsilon}\check{F}_k(t)]_a dt,$$
(18.8)

where  $\chi_a$  is the first component in the  $\eta_B$ ,  $J_{R,n,\epsilon}$ -decomposition of the vector  $\chi \in \mathbb{C}^{2m}$ . Let us rewrite the first term on the right hand side according to

$$e^{-J_{R,a}\bar{t}_{k+1}}[\check{A}^{+}_{k,n,\epsilon}\zeta_{k}]_{a} = e^{-J_{R,a}\bar{t}_{k+1}}[\check{A}^{\text{app},+}_{k,n,\epsilon}]_{a}\zeta_{k,a} + e^{-J_{R,a}\bar{t}_{k+1}}[(\check{A}^{+}_{k,n,\epsilon} - \check{A}^{\text{app},+}_{k,n,\epsilon})\zeta_{k}]_{a}$$
(18.9)
$$= e^{-J_{R,a}\bar{t}_{k}}\zeta_{k,a} + e^{-J_{R,a}\bar{t}_{k}}[B_{k}\zeta_{k}]_{a},$$

where  $B_k$  is a matrix valued function such that

$$\|B_k\| \le \frac{Ce^{-\eta_A \bar{t}_k}}{\nu_n(\iota)e^{\beta_n t_k}},\tag{18.10}$$

where C only depends on the coefficients of the equation (1.2). Letting

$$\chi_k := e^{-J_{R,a}\bar{t}_k} \zeta_{k,a},\tag{18.11}$$

the equalities (18.8) and (18.9) yield

$$\chi_{k+1} = \chi_k + e^{-J_{R,a}\bar{t}_k} [B_k \zeta_k]_a + \int_{t_k}^{t_{k+1}} e^{-J_{R,a}\bar{t}_{k+1}} [\check{A}^+_{k,n,\epsilon}\check{F}_k(t)]_a dt$$

$$= \chi_0 + \sum_{l=0}^k e^{-J_{R,a}\bar{t}_l} [B_l \zeta_l]_a + \sum_{l=0}^k \int_{t_l}^{t_{l+1}} e^{-J_{R,a}\bar{t}_{l+1}} [\check{A}^+_{l,n,\epsilon}\check{F}_l(t)]_a dt.$$
(18.12)

At this stage it is of interest to divide the right hand side into a limiting value and an error term. Let us begin by defining the limiting value:

$$\zeta_{\infty,a} := \zeta_{0,a} + \sum_{l=0}^{\infty} e^{-J_{R,a}\bar{t}_l} [B_l\zeta_l]_a + \sum_{l=0}^{\infty} \int_{t_l}^{t_{l+1}} e^{-J_{R,a}\bar{t}_{l+1}} [\check{A}_{l,n,\epsilon}^+\check{F}_l(t)]_a dt;$$
(18.13)

note that the sums appearing on the right hand side are convergent (we justify this statement in (18.16) and (18.17) below). Given the definition (18.13), the equality (18.12) can be written

$$\zeta_{k+1,a} = e^{J_{R,a}\bar{t}_{k+1}} \zeta_{\infty,a} - \sum_{l=k+1}^{\infty} e^{-J_{R,a}(t_l - t_{k+1})} [B_l \zeta_l]_a - \sum_{l=k+1}^{\infty} \int_{t_l}^{t_{l+1}} e^{-J_{R,a}(t_{l+1} - t_{k+1})} [\check{A}^+_{l,n,\epsilon}\check{F}_l(t)]_a dt.$$
(18.14)

The remainder of the analysis concerning the first component is naturally divided into two parts. First, we need to estimate the limiting value  $\zeta_{\infty,a}$ . Second, we need to estimate the error terms appearing in (18.14); i.e., the last two terms on the right hand side of (18.14). Estimating the limiting value. Let us estimate the terms appearing on the right hand side of (18.13) one by one. To begin with,

$$|\zeta_{0,a}| \le |\zeta_0| \le Ce^{-\kappa_{n,+}t_0} |\psi_0| \le Ce^{-\kappa_{n,+}t_0} |w(\iota, t_0)|;$$
(18.15)

cf. (18.3), (15.36), (15.53) and recall that  $\psi_k = w_{\text{fin},k}$ . Here, C only depends on the coefficients of the equation (1.2). Next,

$$\left|\sum_{l=0}^{\infty} e^{-J_{R,a}\bar{t}_{l}} [B_{l}\zeta_{l}]_{a}\right| \leq C \sum_{l=0}^{\infty} \langle \bar{t}_{l} \rangle^{N} e^{-\eta_{\min,a}\bar{t}_{l}} \frac{e^{-\eta_{A}\bar{t}_{l}}}{\nu_{n}(\iota)e^{\beta_{n}t_{l}}} \\ \cdot \left[ \langle \bar{t}_{l} \rangle^{d_{n,+}-1} |\zeta_{0}| + \int_{t_{0}}^{t_{l}} \langle t_{l} - t' \rangle^{d_{n,+}-1} e^{-\kappa_{n,+}t'} |\hat{f}(\iota,t')| dt' \right] \qquad (18.16)$$
$$\leq C_{B} \left( |\zeta_{0}| + \int_{t_{0}}^{\infty} e^{-\kappa_{n,+}t} |\hat{f}(\iota,t)| dt \right),$$

where we have appealed to (18.7) and (18.10);  $C_B$  only depends on  $\eta_B$  and the coefficients of the equation (1.2); and N is a non-negative integer depending only on m. Moreover,  $\eta_{\min,a}$  is the smallest eigenvalue of  $J_{R,a}$  and we have used the fact that  $-\eta_{\min,a} < \eta_B \leq \eta_A$ . Finally,

$$\left|\sum_{l=0}^{\infty} \int_{t_l}^{t_{l+1}} e^{-J_{R,a}\bar{t}_{l+1}} [\check{A}^+_{l,n,\epsilon}\check{F}_l(t)]_a dt\right| \leq C \int_{t_0}^{\infty} \langle \bar{t} \rangle^N e^{-\eta_{\min,a}\bar{t}} e^{-\kappa_{n,+}t} |\hat{f}(\iota,t)| dt$$

$$\leq C_B \int_{t_0}^{\infty} e^{-(\kappa_{n,+}-\eta_B)t} |\hat{f}(\iota,t)| dt,$$
(18.17)

where  $C_B$  only depends on  $\eta_B$  and the coefficients of the equation (1.2). Combining (18.15)–(18.17) yields the conclusion

$$|\zeta_{\infty,a}| \le C_B \left( e^{-\kappa_{n,+}t_0} |w(\iota,t_0)| + \int_{t_0}^{\infty} e^{-(\kappa_{n,+}-\eta_B)t} |\hat{f}(\iota,t)| dt \right),$$
(18.18)

where  $C_B$  has the same dependence as in the case of (18.17).

Estimating the error terms. Let us estimate the last two terms on the right hand side of (18.14). The second term on the right hand side of (18.14) can be estimated as follows:

$$\left| \sum_{l=k+1}^{\infty} e^{-J_{R,a}(t_{l}-t_{k+1})} [B_{l}\zeta_{l}]_{a} \right| \\
\leq \sum_{l=k+1}^{\infty} C \langle t_{l} - t_{k+1} \rangle^{N} e^{-\eta_{\min,a}(t_{l}-t_{k+1})} \frac{e^{-\eta_{A}\bar{t}_{l}}}{\nu_{n}(\iota)e^{\beta_{n}t_{l}}} \\
\cdot \left( \langle \bar{t}_{l} \rangle^{d_{n,+}-1} |\zeta_{0}| + \int_{t_{0}}^{t_{l}} \langle t_{l} - t' \rangle^{d_{n,+}-1} e^{-\kappa_{n,+}t'} |\hat{f}(\iota,t')| dt' \right) \\
\leq \int_{t_{k+1}}^{\infty} C \langle t - t_{k+1} \rangle^{N} e^{-\eta_{\min,a}(t-t_{k+1})} e^{-\eta_{A}\bar{t}} \langle \bar{t} \rangle^{d_{n,+}-1} dt \\
\cdot \left( |\zeta_{0}| + \int_{t_{0}}^{\infty} e^{-\kappa_{n,+}t'} |\hat{f}(\iota,t')| dt' \right) \\
\leq C_{B} \langle \bar{t}_{k+1} \rangle^{d_{n,+}-1} e^{-\eta_{A}\bar{t}_{k+1}} \left( |\zeta_{0}| + \int_{t_{0}}^{\infty} e^{-\kappa_{n,+}t'} |\hat{f}(\iota,t')| dt' \right),$$
(18.19)

where  $\eta_{\min,a}$  is the smallest eigenvalue of  $J_{R,a}$  and we have appealed to (18.7) and (18.10); note that  $-\eta_{\min,a} < \eta_A$ . Moreover,  $C_B$  only depends on  $\eta_B$  and the coefficients of the equation (1.2).

#### 18.1. ASYMPTOTICS ALONG A TIME SEQUENCE

Let us estimate the third term on the right hand side of (18.14):

$$\left| \sum_{l=k+1}^{\infty} \int_{t_{l}}^{t_{l+1}} e^{-J_{R,a}(t_{l+1}-t_{k+1})} [\check{A}_{l,n,\epsilon}^{+}\check{F}_{l}(t)]_{a} dt \right|$$
  

$$\leq C \int_{t_{k+1}}^{\infty} \langle t - t_{k+1} \rangle^{N} e^{-\eta_{\min,a}(t-t_{k+1})} e^{-\kappa_{n,+}t} |\hat{f}(\iota,t)| dt \qquad (18.20)$$
  

$$\leq C_{B} e^{-\eta_{B} t_{k+1}} \int_{0}^{\infty} e^{-(\kappa_{n,+}-\eta_{B})t} |\hat{f}(\iota,t)| dt,$$

where  $C_B$  only depends on  $\eta_B$  and the coefficients of the equation (1.2). Combining (18.14), (18.19) and (18.20) yields

$$\begin{aligned} |\zeta_{k+1,a} - e^{J_{R,a}\bar{t}_{k+1}}\zeta_{\infty,a}| &\leq C_B \langle \bar{t}_{k+1} \rangle^{d_{n,+}-1} e^{-\eta_B \bar{t}_{k+1}} \\ &\cdot \left( |\zeta_0| + \int_0^\infty e^{-(\kappa_{n,+}-\eta_B)t} |\hat{f}(\iota,t)| dt \right), \end{aligned}$$
(18.21)

where  $C_B$  has the same dependence as in (18.20). Moreover, in case  $\eta_A > \eta_B$ , the polynomial factor can be removed. Before reformulating this estimate, note that if

$$S(t) := \operatorname{diag}\{e^{-i\zeta_1 t} \operatorname{Id}_{l_1}, \dots, e^{-i\zeta_D t} \operatorname{Id}_{l_D}\},\$$

cf. (17.35), then

$$e^{J_{R,n,\epsilon}t} = e^{-\kappa_{n,+}t}S(t)T_{n,\epsilon}^{-1}\exp[R_n(\iota)t]T_{n,\epsilon}$$

where  $R_{\rm n}(\iota)$  is defined in (17.23). Thus

$$\exp[R_{n}(\iota)t] = e^{\kappa_{n,+}t} T_{n,\epsilon}[S(t)]^{-1} e^{J_{R,n,\epsilon}t} T_{n,\epsilon}^{-1}.$$
(18.22)

With these observations in mind, it can be verified that the estimate (18.21) implies that

$$|[T_{\mathbf{n},\epsilon}^{-1}\psi_{k+1}]_{a} - [T_{\mathbf{n},\epsilon}^{-1}\exp[R_{\mathbf{n}}(\iota)\bar{t}_{k+1}]T_{\mathbf{n},\epsilon}]_{a}\chi_{\infty,a}|$$
  

$$\leq C_{B}\langle \bar{t}_{k+1}\rangle^{d_{\mathbf{n},+}-1}e^{\kappa_{\mathbf{n},+}t_{k+1}}e^{-\eta_{B}\bar{t}_{k+1}}\left(|\zeta_{0}| + \int_{0}^{\infty}e^{-(\kappa_{\mathbf{n},+}-\eta_{B})t}|\hat{f}(\iota,t)|dt\right),$$
(18.23)

where  $C_B$  has the same dependence as in (18.20),

$$\chi_{\infty,a} := e^{\kappa_{n,+}t_0} S_{0,a}^{-1} \zeta_{\infty,a} \tag{18.24}$$

and  $S_{k,a}$  denotes the first block in the  $\eta_B, J_{R,n,\epsilon}$ -decomposition of  $S_k$ . Finally, if  $\eta_A > \eta_B$ , then the polynomial factor can be removed. Note that, due to (18.18),

$$|\chi_{\infty,a}| \le C_B \left( |w(\iota, t_0)| + e^{\kappa_{n,+} t_0} \int_{t_0}^{\infty} e^{-(\kappa_{n,+} - \eta_B)t} |\hat{f}(\iota, t)| dt \right),$$
(18.25)

where  $C_B$  has the same dependence as in (18.20).

Estimating the second component. Let us turn to the second component in the  $\eta_B, J_{R,n,\epsilon}$ -decomposition of  $\zeta_k$ . To begin with, (18.5) yields

$$e^{-J_{R,b}\bar{t}_{k+1}}\zeta_{k+1,b} = e^{-J_{R,b}\bar{t}_{k+1}}[\check{A}^+_{k,n,\epsilon}\zeta_k]_b + e^{-J_{R,b}\bar{t}_{k+1}}\int_{t_k}^{t_{k+1}}[\check{A}^+_{k,n,\epsilon}\check{F}_k(t)]_b dt,$$
(18.26)

where  $\zeta_{k,b}$  denotes the second component in the  $\eta_B$ ,  $J_{R,n,\epsilon}$ -decomposition of  $\zeta_k$ . Let us rewrite the first term on the right hand side according to

$$e^{-J_{R,b}\bar{t}_{k+1}}[\check{A}^{+}_{k,n,\epsilon}\zeta_{k}]_{b} = e^{-J_{R,b}\bar{t}_{k+1}}[\check{A}^{\text{app},+}_{k,n,\epsilon}]_{b}\zeta_{k,b} + e^{-J_{R,b}\bar{t}_{k+1}}[(\check{A}^{+}_{k,n,\epsilon} - \check{A}^{\text{app},+}_{k,n,\epsilon})\zeta_{k}]_{b}$$
(18.27)
$$= e^{-J_{R,b}\bar{t}_{k}}\zeta_{k,b} + e^{-J_{R,b}\bar{t}_{k}}[B_{k}\zeta_{k}]_{b},$$

where  $B_k$  is a matrix valued function satisfying (18.10). Letting  $\phi_k := e^{-J_{R,b}\bar{t}_k}\zeta_{k,b}$ , the equalities (18.26) and (18.27) yield

$$\phi_{k+1} = \phi_k + e^{-J_{R,b}\bar{t}_k} [B_k\zeta_k]_b + \int_{t_k}^{t_{k+1}} e^{-J_{R,b}\bar{t}_{k+1}} [\check{A}^+_{k,n,\epsilon}\check{F}_k(t)]_b dt$$

$$= \phi_0 + \sum_{l=0}^k e^{-J_{R,b}\bar{t}_l} [B_l\zeta_l]_b + \sum_{l=0}^k \int_{t_l}^{t_{l+1}} e^{-J_{R,b}\bar{t}_{l+1}} [\check{A}^+_{l,n,\epsilon}\check{F}_l(t)]_b dt.$$
(18.28)

Thus

$$\zeta_{k+1,b} = e^{J_{R,b}\bar{t}_{k+1}}\zeta_{0,b} + \sum_{l=0}^{k} e^{J_{R,b}(t_{k+1}-t_l)} [B_l\zeta_l]_b + \sum_{l=0}^{k} \int_{t_l}^{t_{l+1}} e^{J_{R,b}(t_{k+1}-t_{l+1})} [\check{A}^+_{l,n,\epsilon}\check{F}_l(t)]_b dt.$$
(18.29)

Let us estimate the terms on the right hand side one by one. To begin with,

$$|e^{J_{R,b}\bar{t}_{k+1}}\zeta_{0,b}| \le C\langle \bar{t}_{k+1}\rangle^N e^{\eta_{\max,b}\bar{t}_{k+1}}|\zeta_{0,b}| \le C\langle \bar{t}_{k+1}\rangle^N e^{-\eta_B\bar{t}_{k+1}}|\zeta_0|,$$
(18.30)

where  $\eta_{\max,b}$  is the largest eigenvalue of  $J_{R,b}$  (note that  $\eta_{\max,b} \leq -\eta_B$  due to the definition of  $J_{R,b}$ ), C only depends on the coefficients of the equation (1.2), and N is a non-negative integer depending only on m. Note that if all the Jordan blocks of  $R_n(\iota)$  are trivial, then N = 0. Turning to the second term on the right of (18.29),

$$\left| \sum_{l=0}^{k} e^{J_{R,b}(t_{k+1}-t_{l})} [B_{l}\zeta_{l}]_{b} \right| \leq C \int_{t_{0}}^{t_{k+1}} \langle t_{k+1}-t \rangle^{N} e^{\eta_{\max,b}(t_{k+1}-t)} e^{-\eta_{A}\bar{t}} \langle \bar{t} \rangle^{d_{n,+}-1} dt \cdot \left( |\zeta_{0}| + \int_{t_{0}}^{t_{k+1}} e^{-\kappa_{n,+}t} |\hat{f}(\iota,t)| dt \right)$$

$$\leq C \langle \bar{t}_{k+1} \rangle^{N} e^{-\eta_{B}\bar{t}_{k+1}} \left( |\zeta_{0}| + \int_{t_{0}}^{t_{k+1}} e^{-\kappa_{n,+}t} |\hat{f}(\iota,t)| dt \right),$$
(18.31)

where we have appealed to (18.7) and (18.10). Moreover, C only depends on the coefficients of the equation (1.2), and N is a non-negative integer depending only on m. In addition, N = 1 if all the Jordan blocks of  $R_n(\iota)$  are trivial. Finally, let us estimate the third term on the right hand side of (18.29):

$$\left| \sum_{l=0}^{k} \int_{t_{l}}^{t_{l+1}} e^{J_{R,b}(t_{k+1}-t_{l+1})} [\check{A}^{+}_{l,n,\epsilon} \check{F}_{l}(t)]_{b} dt \right| \\
\leq C \int_{t_{0}}^{t_{k+1}} \langle t_{k+1} - t \rangle^{N} e^{\eta_{\max,b}(t_{k+1}-t)} e^{-\kappa_{n,+}t} |\hat{f}(\iota,t)| dt \qquad (18.32) \\
\leq C \langle \bar{t}_{k+1} \rangle^{N} e^{-\eta_{B} t_{k+1}} \int_{t_{0}}^{t_{k+1}} e^{-(\kappa_{n,+}-\eta_{B})t} |\hat{f}(\iota,t)| dt,$$

where C and N have the same dependence as in the case of (18.31). Moreover, N = 0 if all the Jordan blocks of  $R_n(\iota)$  are trivial. Combining (18.29), (18.30), (18.31) and (18.32) yields

$$|[T_{\mathbf{n},\epsilon}^{-1}\psi_{k+1}]_b| \le C\langle \bar{t}_{k+1}\rangle^N e^{\kappa_{\mathbf{n},+}t_{k+1}} e^{-\eta_B \bar{t}_{k+1}} \left( |\zeta_0| + \int_0^\infty e^{-(\kappa_{\mathbf{n},+}-\eta_B)t} |\hat{f}(\iota,t)| dt \right),$$
(18.33)

where C and N have the same dependence as in the case of (18.31). Moreover, N = 1 if all the Jordan blocks of  $R_n(\iota)$  are trivial.

**Combining the estimates.** Let  $\chi_{\infty}$  be the vector such that the first component in the  $\eta_B$ ,  $J_{R,n,\epsilon}$ -decomposition of  $\chi_{\infty}$  is given by  $\chi_{\infty,a}$ , cf. (18.24), and the second component is zero. Define

$$\bar{\psi}_{\infty} := T_{\mathbf{n},\epsilon} \chi_{\infty}, \quad \psi_{\infty} := e^{-R_{\mathbf{n}}(\iota)T_{\mathbf{n}}} \bar{\psi}_{\infty}.$$
(18.34)

Then (18.25) yields (18.1), and it is clear that  $\bar{\psi}_{\infty}$  and  $\psi_{\infty}$  belong to the first generalised eigenspace of the  $\eta_B$ ,  $R_n(\iota)$ -decomposition of  $\mathbb{C}^{2m}$ .

In order to prove (18.2), note that (18.23) and (18.33) can be combined to yield

$$\begin{aligned} |\psi_{k+1} - e^{R_{n}(\iota)\bar{t}_{k+1}}\bar{\psi}_{\infty}| &= |\psi_{k+1} - e^{R_{n}(\iota)t_{k+1}}\psi_{\infty}| \\ \leq C_{B}\langle \bar{t}_{k+1}\rangle^{N}e^{(\kappa_{n,+}-\eta_{B})\bar{t}_{k+1}} \left( |w(\iota,t_{0})| + e^{\kappa_{n,+}t_{0}}\int_{0}^{\infty} e^{-(\kappa_{n,+}-\eta_{B})t} |\hat{f}(\iota,t)| dt \right) \end{aligned}$$
(18.35)

for  $k \geq 0$ , where  $C_B$  has the same dependence as in the case of (18.17) and N is a non-negative integer depending only on m. Moreover, N = 1 if all the Jordan blocks of  $R_n(\iota)$  are trivial. In addition, if the first generalised eigenspace of the  $\eta_B$ ,  $R_n(\iota)$ -decomposition of  $\mathbb{C}^{2m}$  equals  $\mathbb{C}^{2m}$ , then the N appearing in (18.35) can be replaced by  $d_{n,+} - 1$ , where  $d_{n,+} := d_{\max}(R_n(\iota), \kappa_{n,+})$ . It can also be verified that (18.35) holds for k = -1. Retracing the definitions,

$$\psi_k = D_k(\iota) T_{\text{pre},k} \exp\left(-i \int_0^{t_k} \sigma(\iota, t') \mathfrak{g}(\iota, t') dt'\right) \begin{pmatrix} \mathfrak{g}(\iota, t_k) z(\iota, t_k) \\ \dot{z}(\iota, t_k) \end{pmatrix}.$$
 (18.36)

Thus (18.35) yields (18.2). The lemma follows.

## 18.2 Specifying data at infinity

As a next step, we wish to address the problem of specifying data at infinity, proceeding along an appropriate time sequence. We here restrict our attention to the case of homogeneous equations.

**Lemma 18.5.** Assume that (1.2) is strongly balanced, has a negligible shift vector field, a dominant noisy spatial direction, and is such that the dominant coefficients are convergent. Assume, moreover, that f = 0. Let  $\iota \in \mathcal{I}_{B,n}$  and define  $T_n$  by (17.12). Let  $\{t_k\}$  be the time sequence given by Definition 15.5 and the condition that  $t_0 := T_n$ . Let  $\eta_A$  be defined by (17.25) and fix an  $0 < \eta_B \leq \eta_A$ . Let  $E_a$  denote the first generalised eigenspace of the  $\eta_B$ ,  $R_n(\iota)$ -decomposition of  $\mathbb{C}^{2m}$ , where  $R_n(\iota)$  is given by (17.23); cf. Definition 9.4. Then there is an injective linear map  $\Psi_\infty : E_a \to \mathbb{C}^{2m}$  such that if  $\chi \in E_a$  and z is the solution to the homogeneous version of (24.16) with initial data at  $T_n$  determined by

$$\Psi_{\infty}(\chi) = \begin{pmatrix} \mathfrak{g}(\iota, T_{n}) z(\iota, T_{n}) \\ \dot{z}(\iota, T_{n}) \end{pmatrix}, \qquad (18.37)$$

then (18.2) holds with  $\psi_{\infty}$  replaced by  $\chi$  and  $\hat{f}(\iota, \cdot)$  replaced by 0. Moreover,

$$|\Psi_{\infty}(e^{-R_{n}(\iota)T_{n}}\chi)| \le C|\chi| \tag{18.38}$$

for all  $\chi \in E_a$ , where C only depends on the coefficients of the equation (1.2).

**Remark 18.6.** In analogy with Remark 9.23, it is possible to combine Lemmas 18.1 and 18.5 in order to obtain analogous conclusions in the inhomogeneous setting.

**Remark 18.7.** Dependence on the coefficients includes dependence on the Riemannian manifolds  $(M_r, g_r), r = 1, \ldots, R$ ; cf. Remark 17.10.

*Proof.* Notation and preliminary estimates. In what follows, we use the notation introduced in Section 17.5 without further comment. Moreover, we let  $\lambda = \kappa_{n,+}$  and define  $J_{R,n,\epsilon}$  by (17.43).

Denote the first and second blocks of the  $\eta_B$ ,  $J_{R,n,\epsilon}$ -decomposition of  $J_{R,n,\epsilon}$  by  $J_{R,a}$  and  $J_{R,b}$  respectively, and let

$$J_a := \text{diag}\{J_{R,a}, 0\}, \quad J_b := \text{diag}\{0, J_{R,b}\}, \quad \tilde{A}^+_{k,n,\epsilon} := e^{-J_a \bar{t}_{k+1}} \check{A}^+_{k,n,\epsilon} e^{J_a \bar{t}_k}$$

be  $2m \times 2m$ -matrices, where  $\bar{t}_k := t_k - t_0$ . Compute

$$\tilde{A}_{k,n,\epsilon}^{+} = e^{-J_a \bar{t}_{k+1}} [\check{A}_{k,n,\epsilon}^{+} - \check{A}_{k,n,\epsilon}^{\text{app},+}] e^{J_a \bar{t}_k} + e^{-J_a (t_{k+1} - t_k)} \check{A}_{k,n,\epsilon}^{\text{app},+};$$
(18.39)

note that  $J_a$  and  $\check{A}^{\mathrm{app},+}_{k,\mathrm{n},\epsilon}$  commute. On the other hand,

$$e^{-J_{a}(t_{k+1}-t_{k})}\check{A}_{k,n,\epsilon}^{\mathrm{app},+} = \check{A}_{k,n,\epsilon}^{\mathrm{app},+} - J_{a}(t_{k+1}-t_{k}) + \frac{2\pi}{\nu_{n}(\iota)e^{\beta_{n}t_{k}}}J_{R,n,\epsilon}[e^{-J_{a}(t_{k+1}-t_{k})} - \mathrm{Id}_{2m}] + [e^{-J_{a}(t_{k+1}-t_{k})} - \mathrm{Id}_{2m} + J_{a}(t_{k+1}-t_{k})].$$

Introducing

$$\tilde{A}_{k,\mathbf{n},\epsilon}^{\mathrm{app},+} := \mathrm{Id}_{2m} + \frac{2\pi}{\nu_{\mathbf{n}}(\iota)e^{\beta_{\mathbf{n}}t_k}}J_b \tag{18.40}$$

and appealing to (14.27) and (17.14) yields

$$\|e^{-J_a(t_{k+1}-t_k)}\check{A}_{k,\mathbf{n},\epsilon}^{\mathrm{app},+} - \check{A}_{k,\mathbf{n},\epsilon}^{\mathrm{app},+}\| \le \frac{C_{\epsilon}e^{-\eta_A t_k}}{\nu_{\mathbf{n}}(\iota)e^{\beta_{\mathbf{n}}t_k}}$$

for all  $k \ge 0$ , where  $C_{\epsilon}$  only depends on  $\epsilon$  and the coefficients of the equation (1.2). Combining this estimate with (17.38) and (18.39) yields

$$\|\tilde{A}_{k,n,\epsilon}^{+} - \tilde{A}_{k,n,\epsilon}^{\mathrm{app},+}\| \le \frac{C_{\epsilon}}{\nu_{n}(\iota)e^{\beta_{n}t_{k}}} \langle \bar{t}_{k} \rangle^{N} e^{-\eta_{A,a}\bar{t}_{k}}$$
(18.41)

for all  $k \geq 0$ , where  $C_{\epsilon}$  only depends on  $\epsilon$  and the coefficients of the equation (1.2); N is a non-negative integer depending only on m; and  $\eta_{A,a}$  is the difference between  $\eta_A$  and the largest eigenvalue of the matrix  $-J_a$  (note that this difference is strictly positive due to the definition of  $J_a$ ). Note also that, by the definition of  $J_b$ ,  $\|\tilde{A}_{k,n,\epsilon}^{app,+}\| \leq 1$  (in order to obtain this estimate, we, however, need to assume  $\epsilon$  to be small enough, the bound mentioned in connection with (17.50) being sufficient; from now on we fix such an  $\epsilon$ , depending only on the coefficients of the equation (1.2), so that dependence on  $\epsilon$  from now on can be replaced by dependence on the coefficients of the equation (1.2)). In particular, if  $0 \leq k_a \leq k_b$  are integers, then

$$\|\tilde{A}_{k_b,\mathbf{n},\epsilon}^+\cdots\tilde{A}_{k_a,\mathbf{n},\epsilon}^+\| \le C \tag{18.42}$$

for a constant C depending only on the coefficients of the equation (1.2); the proof of this statement is similar to the proof of (17.51).

Asymptotics of the leading order component. Let us return to (15.63) in case  $\hat{f} = 0$ . Then  $\psi_{k+1} = A_k^+ \psi_k$ . Introducing

$$\tilde{\rho}_k := e^{-\kappa_{\mathbf{n},+}t_k} e^{-J_a \bar{t}_k} S_k T_{\mathbf{n},\epsilon}^{-1} \psi_k, \qquad (18.43)$$

we thus have

$$\tilde{\rho}_{k+1} = A_{k,n,\epsilon}^+ \tilde{\rho}_k. \tag{18.44}$$

Due to (18.42),

$$|\tilde{\rho}_k| \le C |\tilde{\rho}_{k_0}| \tag{18.45}$$

for  $0 \le k_0 \le k$ , where C has the same dependence as in the case of (18.42). Letting  $\tilde{\rho}_{k,a}$  denote the first component in the  $\eta_B, J_{R,n,\epsilon}$ -decomposition of  $\tilde{\rho}_k$ , (18.44) implies

$$\tilde{\rho}_{k+1,a} = \tilde{\rho}_{k,a} + [(\tilde{A}^+_{k,n,\epsilon} - \tilde{A}^{\mathrm{app},+}_{k,n,\epsilon})\tilde{\rho}_k]_a = \tilde{\rho}_{k_0,a} + \sum_{l=k_0}^k [(\tilde{A}^+_{l,n,\epsilon} - \tilde{A}^{\mathrm{app},+}_{l,n,\epsilon})\tilde{\rho}_l]_a$$

#### 18.2. SPECIFYING DATA AT INFINITY

for every  $0 \le k_0 \le k$ . Combining this equality with (18.41) and (18.45) yields

$$\begin{aligned} |\tilde{\rho}_{k+1,a} - \tilde{\rho}_{k_0,a}| &\leq \sum_{l=k_0}^{k} \frac{C}{\nu_{n}(\iota)e^{\beta_{n}t_{l}}} \langle \bar{t}_{l} \rangle^{N} e^{-\eta_{A,a}\bar{t}_{l}} |\tilde{\rho}_{k_0}| \\ &\leq C \int_{t_{k_0}}^{t_{k+1}} \langle \bar{t} \rangle^{N} e^{-\eta_{A,a}\bar{t}} dt |\tilde{\rho}_{k_0}| \leq C \langle \bar{t}_{k_0} \rangle^{N} e^{-\eta_{A,a}\bar{t}_{k_0}} |\tilde{\rho}_{k_0}| \end{aligned}$$
(18.46)

for all  $0 \le k_0 \le k$ , where C has the same dependence as in the case of (18.42). In particular, given an  $\varepsilon > 0$ , we can choose a  $\bar{t}_{k_0}$  (depending only on  $\varepsilon$  and the coefficients of the equation (1.2)) such that

$$\left|\tilde{\rho}_{k+1,a} - \tilde{\rho}_{k_0,a}\right| \le \varepsilon |\tilde{\rho}_{k_0}| \tag{18.47}$$

for all  $0 \le k_0 \le k$ .

From asymptotic data to a finite time hypersurface. From now on, the structure of the argument is quite similar to the end of the proof of Lemma 9.21, though some of the details are different. Let  $k_0$  be such that (18.47) holds with  $\varepsilon$  replaced by 1/2 and let  $l_a$  denote the dimension of the first subspace of the  $\eta_B, J_{R,n,\epsilon}$ -decomposition of  $\mathbb{C}^{2m}$ . Define the map  $L_a : \mathbb{C}^{l_a} \to \mathbb{C}^{l_a}$  as follows. Given  $\eta \in \mathbb{C}^{l_a}$ , let  $\xi \in \mathbb{C}^{2m}$  be such that  $\xi_a = \eta$  and  $\xi_b = 0$  (where we divide  $\xi$  into components  $\xi_a$  and  $\xi_b$  in the same way as  $\tilde{\rho}_k$ ). Let  $\tilde{\rho}_k$  be the solution to (18.44) such that  $\tilde{\rho}_{k_0} = \xi$  and define  $L_a \eta$  by

$$L_a \eta := \lim_{k \to \infty} \tilde{\rho}_{k,a}. \tag{18.48}$$

Due to estimates such as (18.46), it is clear that  $\tilde{\rho}_{k,a}$  is a Cauchy sequence, so that the limit on the right hand side of (18.48) exists. Thus  $L_a \eta$  is well defined. Moreover, it is clear that  $L_a$  is linear. Due to the fact that (18.47) holds with  $\varepsilon$  replaced by 1/2 and the fact that  $\tilde{\rho}_{k_0,b} = 0$ ,

$$|L_a\eta - \tilde{\rho}_{k_0,a}| \le \frac{1}{2} |\tilde{\rho}_{k_0,a}|.$$

In particular,  $|\eta|/2 \leq |L_a\eta|$ , so that  $L_a$  is injective. Since  $L_a$  is a linear map between spaces of the same finite dimension, it follows that  $L_a$  is an isomorphism. Moreover,

$$|L_a^{-1}\eta| \le 2|\eta|. \tag{18.49}$$

Note that  $L_a^{-1}$  (essentially) maps data at infinity to data at  $t_{k_0}$ . In the end, we wish to translate the data at  $t_{k_0}$  to data at  $t_0$ . To this end, we wish to solve (18.44) backwards.

From asymptotic data to  $t = t_0$ . In order to solve (18.44) backwards, we can proceed similarly to the above; i.e., study the backwards iteration. However, this involves unnecessary complications. In fact, since  $\bar{t}_{k_0}$  is bounded by a constant depending only on the coefficients of the equation (1.2), it is sufficient to appeal to Lemma 10.7. In order to justify this statement, note, to begin with, that (18.43) yields

$$|\tilde{\rho}_0| \le C e^{-\kappa_{n,+} t_0} |w(\iota, t_0)|, \tag{18.50}$$

where C only depends on the coefficients of the equation (1.2). On the other hand,  $\mathcal{E}^{1/2}(\iota, t)$  appearing in (10.12) is equivalent to  $|w(\iota, t)|$  for  $t \geq T_n$ , the constant of equivalence being numerical. Combining (10.12) with (18.50) yields

$$|\tilde{\rho}_0| \le C e^{-\kappa_{\mathrm{n},+}t_0} e^{\eta_{\mathrm{bal}}t_{k_0}} |w(\iota, t_{k_0})|$$

where C only depends on the coefficients of the equation (1.2). On the other hand, combining (18.43) with the fact that  $\bar{t}_{k_0}$  is bounded by a constant depending only on the coefficients of the equation (1.2),

$$|w(\iota, t_{k_0})| \le C e^{\kappa_{\mathbf{n}, +} t_{k_0}} |\tilde{\rho}_{k_0}|,$$

where C only depends on the coefficients of the equation (1.2). Combining the last two estimates yields

$$|\tilde{\rho}_0| \le C e^{\kappa_{\mathrm{n},+} \bar{t}_{k_0}} e^{\eta_{\mathrm{bal}} \bar{t}_{k_0}} |\tilde{\rho}_{k_0}|,$$

where C only depends on the coefficients of the equation (1.2). To conclude,

$$|\tilde{\rho}_0| \le C |\tilde{\rho}_{k_0}|,\tag{18.51}$$

where C only depends on the coefficients of the equation (1.2).

**Defining the map**  $\Psi_{\infty}$ . Given the above information, we are in a position to define the map taking data "at infinity" to data at the finite time  $t_0 = T_n$ . Let  $\Psi_{\infty}$  be the composition of the following maps: first the map which takes  $\chi \in E_a$  to

$$\varsigma_a := L_a^{-1} \left( e^{-\kappa_{n,+} T_n} S_{0,a} [T_{n,\epsilon}^{-1} e^{R_n(\iota) T_n} \chi]_a \right);$$
(18.52)

second, the map which with  $\zeta_a$  associates the vector  $\xi \in \mathbb{C}^{2m}$  such that  $\xi_a = \zeta_a$  and  $\xi_b = 0$ ; third, the map which with  $\xi$  associates  $\tilde{\rho}_0$ , where  $\tilde{\rho}_k$  is the solution to (18.44) such that  $\tilde{\rho}_{k_0} = \xi$ ; and fourth, the map which with  $\tilde{\rho}_0$  associates the element

$$\exp\left(i\int_{0}^{t_{0}}\sigma(\iota,t')\mathfrak{g}(\iota,t')dt'\right)T_{\text{pre},0}^{-1}D_{0}^{-1}e^{\kappa_{n,+}t_{0}}T_{n,\epsilon}S_{0}^{-1}\tilde{\rho}_{0}.$$
(18.53)

Returning to the definitions (14.7), (15.1), (15.36), (15.53), (18.43) and recalling that  $\psi_k = w_{\text{fin},k}$ , it is clear that if, given  $\chi \in E_a$ , we let z be the solution to (24.16) such that

$$\Psi_{\infty}(\chi) = \begin{pmatrix} \mathfrak{g}(\iota, t_0) z(\iota, t_0) \\ \dot{z}(\iota, t_0) \end{pmatrix}$$
(18.54)

holds, then the  $\tilde{\rho}_k$  associated with the solution according to (18.43) satisfies  $\tilde{\rho}_{k_0} = \xi$ , where  $\xi_a = \varsigma_a$ ,  $\xi_b = 0$  and  $\varsigma_a$  is given by (18.52). Note that  $\Psi_{\infty}$  is linear and injective, and that

$$|\Psi_{\infty}(e^{-R_{n}(\iota)T_{n}}\bar{\chi})| \leq C|\bar{\chi}|$$

for all  $\bar{\chi} \in E_a$ , where C only depends on the coefficients of the equation (1.2). In order to obtain this estimate we appeal to (18.51) and the definition of  $\Psi_{\infty}$ .

Let now  $\chi \in E_a$  and z be the solution to (24.16) such that (18.54) holds. Then, by construction,

$$\lim_{k \to \infty} \tilde{\rho}_{k,a} = L_a(\varsigma_a) = e^{-\kappa_{n,+} T_n} S_{0,a} [T_{n,\epsilon}^{-1} e^{R_n(\iota) T_n} \chi]_a.$$
(18.55)

On the other hand, keeping (18.3), (18.11), (18.12), (18.13), (18.24) and (18.43) in mind (in the case that  $\hat{f}(\iota, \cdot) = 0$ ) yields

$$\lim_{k \to \infty} \tilde{\rho}_{k,a} = e^{-\kappa_{n,+}T_n} S_{0,a} \chi_{\infty,a}.$$
(18.56)

Letting  $\chi_{\infty}$  be such that the first component in the  $\eta_B$ ,  $J_{R,n,\epsilon}$ -decomposition of  $\chi_{\infty}$  is given by  $\chi_{\infty,a}$  and the second component vanishes, the equalities (18.55) and (18.56) yield

$$\chi_{\infty} = T_{\mathrm{n},\epsilon}^{-1} e^{R_{\mathrm{n}}(\iota)T_{\mathrm{n}}} \chi_{\mathrm{s}}$$

where we have used the fact that  $\chi \in E_a$ . Recalling (18.34), we conclude that the  $\psi_{\infty}$  appearing in the statement of Lemma 18.1 equals  $\chi$ . The lemma follows.

## 18.3 Controlling the solution beyond the time sequence

The next step is to analyse how the solution behaves in between the times belonging to the time sequence  $\{t_k\}$ . Before doing so, it is convenient to introduce the following terminology:

$$S_{\mathbf{n}}(\iota) := \frac{1}{2} \begin{pmatrix} -\alpha_{\infty} + \beta_{\mathbf{n}} \mathrm{Id}_{m} & iX_{\mathbf{n}}(\iota) \\ -i\bar{X}_{\mathbf{n}}(\iota) & -\alpha_{\infty} + \beta_{\mathbf{n}} \mathrm{Id}_{m} \end{pmatrix},$$
(18.57)

cf. the notation introduced in Definition 17.7, and

$$\omega_{\rm sh}(\iota, t) := \int_0^t \sigma(\iota, t') \mathfrak{g}(\iota, t') dt'.$$
(18.58)

Moreover,

$$\mathfrak{D}_{\rm tot}(\iota,t) := \begin{pmatrix} \cos\varphi_{\rm tot}(\iota,t)\mathrm{Id}_m & \sin\varphi_{\rm tot}(\iota,t)\mathrm{Id}_m \\ -\sin\varphi_{\rm tot}(\iota,t)\mathrm{Id}_m & \cos\varphi_{\rm tot}(\iota,t)\mathrm{Id}_m \end{pmatrix},\tag{18.59}$$

where

$$\varphi_{\text{tot}}(\iota, t) := \int_0^t [1 + \sigma^2(\iota, t')]^{1/2} \mathfrak{g}(\iota, t') dt'.$$
(18.60)

**Lemma 18.8.** Assume that (1.2) is strongly balanced, has a negligible shift vector field, a dominant noisy spatial direction, and is such that the dominant coefficients are convergent. Let  $\iota \in \mathcal{I}_{B,n}$  and define  $T_n$  by (17.12). Let  $\eta_A$  be defined by (17.25) and assume that there is an  $0 < \eta_B \leq \eta_A$  such that

$$\int_0^\infty e^{-(\kappa_{\mathbf{n},+}-\eta_B)t} |\hat{f}(\iota,t)| dt < \infty,$$

where  $\kappa_{n,+}$  is defined by (17.6). Then there is a constant  $C_B$ , depending only on  $\eta_B$  and the coefficients of the equation (1.2); and a non-negative integer N, depending only on m, such that the following holds. If z is a solution to (24.16), then there is a  $z_{\infty}$  belonging to the first generalised eigenspace of the  $\eta_B$ ,  $S_n(\iota)$ -decomposition of  $\mathbb{C}^{2m}$  (cf. Definition 9.4) and satisfying

$$|e^{S_{n}(\iota)T_{n}}z_{\infty}| \leq C_{B}\left(|w(\iota,T_{n})| + e^{\kappa_{n,+}T_{n}}\int_{T_{n}}^{\infty} e^{-(\kappa_{n,+}-\eta_{B})t}|\hat{f}(\iota,t)|dt\right),$$
(18.61)

such that

$$\left| \begin{pmatrix} \mathfrak{g}(\iota,t)z(\iota,t) \\ \dot{z}(\iota,t) \end{pmatrix} - e^{i\omega_{\rm sh}(\iota,t)}\mathfrak{D}_{\rm tot}(\iota,t)e^{S_{\rm n}(\iota)t}z_{\infty} \right| \\ \leq C_B \langle \overline{t} \rangle^N e^{(\kappa_{\rm n,+}-\eta_B)\overline{t}} \left( |w(\iota,T_{\rm n})| + e^{\kappa_{\rm n,+}T_{\rm n}} \int_0^\infty e^{-(\kappa_{\rm n,+}-\eta_B)t} |\hat{f}(\iota,t)| dt \right)$$
(18.62)

holds for all  $t \geq T_n$ , where  $\bar{t} := t - T_n$ .

Remark 18.9. The estimate (18.62) can also be formulated as (18.74).

**Remark 18.10.** In case all the Jordan blocks of  $S_n(\iota)$  are trivial, the *N* appearing in (18.62) can be replaced by 1. Moreover, if the first generalised eigenspace of the  $\eta_B$ ,  $S_n(\iota)$ -decomposition of  $\mathbb{C}^{2m}$  equals  $\mathbb{C}^{2m}$ , then the *N* appearing in (18.62) can be replaced by  $d_{n,+} - 1$ , where  $d_{n,+} := d_{\max}(S_n(\iota), \kappa_{n,+})$ .

**Remark 18.11.** Dependence on the coefficients includes dependence on the Riemannian manifolds  $(M_r, g_r), r = 1, \ldots, R$ ; cf. Remark 17.10.

Proof. Consider (18.2). The first step of the analysis is to simplify the left hand side.

Simplifying (18.2). To begin with, we replace  $T_{\text{pre},k}^{-1}$  with  $T_{\text{pre}}^{-1}$ , where

$$T_{\rm pre} := \frac{1}{2} \begin{pmatrix} \mathrm{Id}_m & i \mathrm{Id}_m \\ i \mathrm{Id}_m & \mathrm{Id}_m \end{pmatrix}$$
(18.63)

In order to justify that this is allowed, note that

$$T_{\text{pre},k}^{-1} - T_{\text{pre}}^{-1} = \frac{1}{\nu_k} \begin{pmatrix} \text{Id}_m & -i\text{Id}_m \\ -i(\xi_k + \nu_k)\text{Id}_m & (\nu_k - \xi_k)\text{Id}_m \end{pmatrix} - \frac{1}{\nu_k} \begin{pmatrix} \text{Id}_m & -i\text{Id}_m \\ -i\text{Id}_m & \text{Id}_m \end{pmatrix} + \left(\frac{1}{\nu_k} - 1\right) \begin{pmatrix} \text{Id}_m & -i\text{Id}_m \\ -i\text{Id}_m & \text{Id}_m \end{pmatrix};$$

cf. (15.34) and (15.35) and the notation introduced in the adjacent text. Thus

$$\|T_{\text{pre},k}^{-1} - T_{\text{pre}}^{-1}\| \le C e^{-\eta_{\text{sh}} t_k} \tag{18.64}$$

for all  $k \ge 0$ , where C only depends on the coefficients of the equation (1.2). As a consequence of this estimate, (18.1) and the fact that the largest real part of an eigenvalue of  $R_n(\iota)$  is  $\kappa_{n,+}$ ,

$$\left| \exp\left( i \int_{0}^{t_{k}} \sigma(\iota, t') \mathfrak{g}(\iota, t') dt' \right) (T_{\text{pre},k}^{-1} - T_{\text{pre}}^{-1}) [D_{k}(\iota)]^{-1} e^{R_{n}(\iota)t_{k}} \psi_{\infty} \right|$$

$$\leq C_{B} e^{-\eta_{\text{sh}}t_{k}} \langle \bar{t}_{k} \rangle^{d_{n,+}-1} e^{\kappa_{n,+}\bar{t}_{k}} \left( |w(\iota, T_{n})| + e^{\kappa_{n,+}T_{n}} \int_{0}^{\infty} e^{-(\kappa_{n,+}-\eta_{B})t} |\hat{f}(\iota, t)| dt \right),$$
(18.65)

where  $C_B$  depends only on  $\eta_B$  and the coefficients of the equation (1.2). Combining this estimate with (18.2) yields

$$\left| \begin{pmatrix} \mathfrak{g}(\iota, t_k) z(\iota, t_k) \\ \dot{z}(\iota, t_k) \end{pmatrix} - \exp\left(i \int_0^{t_k} \sigma(\iota, t') \mathfrak{g}(\iota, t') dt'\right) T_{\text{pre}}^{-1} [D_k(\iota)]^{-1} e^{R_n(\iota) t_k} \psi_{\infty} \right| \\
\leq C_B \langle \bar{t}_k \rangle^N e^{(\kappa_{n,+} - \eta_B) \bar{t}_k} \left( |w(\iota, T_n)| + e^{\kappa_{n,+} T_n} \int_0^{\infty} e^{-(\kappa_{n,+} - \eta_B) t} |\hat{f}(\iota, t)| dt \right),$$
(18.66)

where  $C_B$  has the same dependence as in the case of (18.65) and N only depends on m. Moreover, in case all the Jordan blocks of  $R_n(\iota)$  are trivial, the N appearing in (18.66) can be replaced by 1. In addition, if the first generalised eigenspace of the  $\eta_B$ ,  $R_n(\iota)$ -decomposition of  $\mathbb{C}^{2m}$  equals  $\mathbb{C}^{2m}$ , then the N appearing in (18.66) can be replaced by  $d_{n,+} - 1$ . Let

$$\varphi_{\rm sh}(\iota,t) := \int_0^t \left( [1 + \sigma^2(\iota,t')]^{1/2} - 1 \right) \mathfrak{g}(\iota,t') dt' + \int_0^{t_0} \mathfrak{g}(\iota,t') dt'.$$
(18.67)

Then

$$\varphi_{\rm sh}(\iota, t_k) = \int_0^{t_k} [1 + \sigma^2(\iota, t')]^{1/2} \mathfrak{g}(\iota, t') dt' - 2\pi k;$$

note that by the definition of the time sequence  $\{t_k\}$ ,

$$\int_{t_0}^{t_k} \mathfrak{g}(\iota, t') dt' = 2\pi k.$$

Define

$$\mathfrak{D}_{\rm sh}(\iota,t) := \left(\begin{array}{c} \cos\varphi_{\rm sh}(\iota,t) \mathrm{Id}_m & \sin\varphi_{\rm sh}(\iota,t) \mathrm{Id}_m \\ -\sin\varphi_{\rm sh}(\iota,t) \mathrm{Id}_m & \cos\varphi_{\rm sh}(\iota,t) \mathrm{Id}_m \end{array}\right)$$

Then the equality

$$T_{\rm pre}^{-1}[D_k(\iota)]^{-1}T_{\rm pre} = \mathfrak{D}_{\rm sh}(\iota, t_k)$$

holds, and (18.66) can be rewritten

$$\left| \begin{pmatrix} \mathfrak{g}(\iota, t_k) z(\iota, t_k) \\ \dot{z}(\iota, t_k) \end{pmatrix} - e^{i\omega_{\rm sh}(\iota, t_k)} \mathfrak{D}_{\rm sh}(\iota, t_k) e^{S_{\rm n}(\iota)t_k} z_{\infty} \right| \\ \leq C_B \langle \bar{t}_k \rangle^N e^{(\kappa_{\rm n,+} - \eta_B)\bar{t}_k} \left( |w(\iota, T_{\rm n})| + e^{\kappa_{\rm n,+}T_{\rm n}} \int_0^\infty e^{-(\kappa_{\rm n,+} - \eta_B)t} |\hat{f}(\iota, t)| dt \right)$$
(18.68)

for all  $k \geq 0$ , where  $z_{\infty} := T_{\text{pre}}^{-1}\psi_{\infty}$ ;  $S_n(\iota)$  is defined by (18.57); and  $\omega_{\text{sh}}$  is defined by (18.58). Moreover,  $C_B$  and N have the same dependence as in the case of (18.66). In addition, in case all the Jordan blocks of  $S_n(\iota)$  are trivial, the N appearing in (18.68) can be replaced by 1. Finally, if the first generalised eigenspace of the  $\eta_B$ ,  $S_n(\iota)$ -decomposition of  $\mathbb{C}^{2m}$  equals  $\mathbb{C}^{2m}$ , then the N appearing in (18.68) can be replaced by  $d_{n,+} - 1$ . Note that  $d_{n,+} = d_{\max}(S_n(\iota), \kappa_{n,+})$  and that  $z_{\infty}$  belongs to the first generalised eigenspace of the  $\eta_B$ ,  $S_n(\iota)$ -decomposition of  $\mathbb{C}^{2m}$ . Moreover, given the definition of  $z_{\infty}$ , (18.61) is an immediate consequence of (18.1). Keeping (14.7), (15.1) and (18.58) in mind, the estimate (18.68) can be written

$$\left| w(\iota, t_k) - \mathfrak{D}_{\mathrm{sh}}(\iota, t_k) e^{S_{\mathrm{n}}(\iota)t_k} z_{\infty} \right| \\
\leq C_B \langle \bar{t}_k \rangle^N e^{(\kappa_{\mathrm{n},+} - \eta_B)\bar{t}_k} \left( |w(\iota, T_{\mathrm{n}})| + e^{\kappa_{\mathrm{n},+}T_{\mathrm{n}}} \int_0^\infty e^{-(\kappa_{\mathrm{n},+} - \eta_B)t} |\hat{f}(\iota, t)| dt \right)$$
(18.69)

for all  $k \ge 0$ , where  $C_B$  and N have the same dependence as in the case of (18.66). Next we consider the behaviour of the solution between the  $t_k$ 's.

Estimating the solution for all t's. In order to estimate the solution in the intervals between the  $t_k$ 's, let us recall that for  $t \in [t_k, t_{k+1}]$ 

$$w(\iota, t) = \Xi(\tau; \tau_k) w(\iota, t_k) + \Xi(\tau; \tau_k) \int_{\tau_k}^{\tau} [\Phi(\tau'; \tau_k)]^{-1} F(\iota, \tau') d\tau', \qquad (18.70)$$

where we have appealed to (15.3) and  $\tau$  and  $\tau_k$  are assumed to correspond to t and  $t_k$  according to (14.13). At this stage we would like to appeal to Lemma 15.1. Due to Lemma 17.9 and the definition of the time sequence  $\{t_k\}$ , it is clear that this is allowed, with  $t_a$  replaced by  $t_k$ ,  $t_b$ replaced by  $t_{k+1}$ , where  $k \ge 0$ , etc. Moreover, since  $\varepsilon_a$  is given by (14.33) and Lemmas 17.9 and 17.13 apply, we can replace  $\varepsilon_a$  in the estimates with

$$\varepsilon_k \le \frac{C e^{-\eta_{\mathrm{sh}} t_k}}{\nu_{\mathrm{n}}(\iota) e^{\beta_{\mathrm{n}} t_k}} + \frac{C}{\nu_{\mathrm{n}}^2(\iota) e^{2\beta_{\mathrm{n}} t_k}},$$

where C only depends on the coefficients of the equation (1.2). Keeping in mind that  $\mathfrak{f}_{ode}$ ,  $\mathfrak{f}_X$  and  $\mathfrak{f}_{sh}$  are given by (17.8), (15.4) yields

$$\|\Xi(\tau;\tau_k) - \mathfrak{D}_{\mathrm{m}}(\iota,t)\| \le \frac{C}{\nu_{\mathrm{n}}(\iota)e^{\beta_{\mathrm{n}}t_k}} + Ce^{-\eta_{\mathrm{sh}}t_k}$$
(18.71)

for all  $t \in [t_k, t_{k+1}]$ , where  $\mathfrak{D}_m$  equals  $\mathfrak{D}_a$ , defined in (15.5), with a replaced by k and C only depends on the coefficients of the equation (1.2). Define

$$\varphi_{\mathrm{m}}(\iota,t) := \int_0^t \mathfrak{g}(\iota,t')dt' + \varphi_{\mathrm{m},0}(\iota), \quad \varphi_{\mathrm{m},0}(\iota) := -\int_0^{t_0} \mathfrak{g}(\iota,t')dt'.$$

Then

$$\int_{t_k}^t \mathfrak{g}(\iota, t') dt' = \varphi_{\mathrm{m}}(\iota, t) - 2\pi k,$$

so that

$$\mathfrak{D}_{\mathrm{m}}(\iota,t) = \begin{pmatrix} \cos\varphi_{\mathrm{m}}(\iota,t)\mathrm{Id}_{m} & \sin\varphi_{\mathrm{m}}(\iota,t)\mathrm{Id}_{m} \\ -\sin\varphi_{\mathrm{m}}(\iota,t)\mathrm{Id}_{m} & \cos\varphi_{\mathrm{m}}(\iota,t)\mathrm{Id}_{m} \end{pmatrix}$$

Combining this observation with (18.61), (18.69) and (18.71) and the fact that the largest real part of an eigenvalue of  $S_n(\iota)$  is  $\kappa_{n,+}$ ,

$$\left| \Xi(\tau;\tau_k)w(\iota,t_k) - \mathfrak{D}_{\mathrm{m}}(\iota,t)\mathfrak{D}_{\mathrm{sh}}(\iota,t_k)e^{S_{\mathrm{n}}(\iota)t_k}z_{\infty} \right| \\ \leq C_B \langle \bar{t}_k \rangle^N e^{(\kappa_{\mathrm{n},+}-\eta_B)\bar{t}_k} \left( |w(\iota,T_{\mathrm{n}})| + e^{\kappa_{\mathrm{n},+}T_{\mathrm{n}}} \int_0^\infty e^{-(\kappa_{\mathrm{n},+}-\eta_B)t} |\hat{f}(\iota,t)| dt \right)$$
(18.72)

for all  $k \geq 0$ , where  $C_B$  only depends on  $\eta_B$  and the coefficients of the equation (1.2) and Nonly depends on m. Moreover, in case all the Jordan blocks of  $S_n(\iota)$  are trivial, the N appearing in (18.72) can be replaced by 1. In addition, if the first generalised eigenspace of the  $\eta_B$ ,  $S_n(\iota)$ decomposition of  $\mathbb{C}^{2m}$  equals  $\mathbb{C}^{2m}$ , then the N appearing in (18.72) can be replaced by  $d_{n,+} - 1$ , where  $d_{n,+} = d_{\max}(S_n(\iota), \kappa_{n,+})$ . Next we wish to estimate

$$\Xi(\tau;\tau_k)\int_{\tau_k}^{\tau} [\Phi(\tau';\tau_k)]^{-1}F(\iota,\tau')d\tau'.$$

Note, to this end, that  $\|\Xi(\tau; \tau_k)\|$  is bounded by a constant on the interval of interest, where the constant depends only on the coefficients of the equation (1.2); this is a consequence of (18.71). That  $\|[\Phi(\tau; \tau_k)]^{-1}\|$  can be bounded in the same way on the interval of interest is a consequence of (14.34) and (14.51). Moreover, (14.7), (14.8), (14.13), (14.14) and (14.19) yield the conclusion that

$$\int_{\tau_k}^{\tau} [\Phi(\tau';\tau_k)]^{-1} F(\iota,\tau') d\tau' = \int_{t_k}^{t} [\Phi(\tau(t');\tau_k)]^{-1} \exp\left[-\frac{1}{2} \int_{t_k}^{t'} \dot{\ell}(\iota,t'') dt''\right] \\ \cdot \exp\left(-i \int_0^{t'} \sigma(\iota,t'') \mathfrak{g}(\iota,t'') dt''\right) \begin{pmatrix} 0\\ U(t';t_k) \hat{f}(\iota,t') \end{pmatrix} dt'.$$

That the integral involving  $\dot{\ell}$  is bounded on the interval of interest is a consequence of (14.23) and that  $||U(t';t_k)||$  is bounded on the interval of interest by a constant depending only on the coefficients of the equation (1.2) is a consequence of (14.39). Combining these observations yields the conclusion that

$$\left| \Xi(\tau;\tau_k) \int_{\tau_k}^{\tau} [\Phi(\tau';\tau_k)]^{-1} F(\iota,\tau') d\tau' \right| \leq C \int_{t_k}^{t} |\hat{f}(\iota,t')| dt'$$
$$\leq C e^{(\kappa_{n,+}-\eta_B)\bar{t}_k} e^{\kappa_{n,+}t_0} \int_{t_k}^{t} e^{-(\kappa_{n,+}-\eta_B)t'} |\hat{f}(\iota,t')| dt',$$

where C is a constant depending only on the coefficients of the equation (1.2). Combining this estimate with (18.70) and (18.72) yields

$$\left| w(\iota,t) - \mathfrak{D}_{\mathrm{m}}(\iota,t)\mathfrak{D}_{\mathrm{sh}}(\iota,t_{k})e^{S_{\mathrm{n}}(\iota)t_{k}}z_{\infty} \right| \leq C_{B}\langle \bar{t}_{k}\rangle^{N}e^{(\kappa_{\mathrm{n},+}-\eta_{B})\bar{t}_{k}} \left( |w(\iota,T_{\mathrm{n}})| + e^{\kappa_{\mathrm{n},+}T_{\mathrm{n}}}\int_{0}^{\infty}e^{-(\kappa_{\mathrm{n},+}-\eta_{B})t}|\hat{f}(\iota,t)|dt \right)$$
(18.73)

for all  $k \ge 0$  and  $t \in [t_k, t_{k+1}]$ , where  $C_B$  only depends on  $\eta_B$  and the coefficients of the equation (1.2) and N only depends on m. Moreover, in case all the Jordan blocks of  $S_n(\iota)$  are trivial, the N appearing in (18.73) can be replaced by 1. In addition, if the first generalised eigenspace of the  $\eta_B$ ,  $S_n(\iota)$ -decomposition of  $\mathbb{C}^{2m}$  equals  $\mathbb{C}^{2m}$ , then the N appearing in (18.73) can be replaced by  $d_{n,+} - 1$ , where  $d_{n,+} = d_{\max}(S_n(\iota), \kappa_{n,+})$ .

We would like to change the remaining  $t_k$ 's appearing in (18.73) to t's. Changing  $\bar{t}_k$  to  $\bar{t}$  on the right hand side is clearly allowed. Changing  $t_k$  appearing in  $e^{S_n(\iota)t_k}$  to t is also allowed due to (14.24), (18.61) and the fact that the largest real part of an eigenvalue of  $S_n(\iota)$  is  $\kappa_{n,+}$ . In order to justify that we are allowed to change the argument in  $\mathfrak{D}_{sh}$ , note that, due to (18.67),

$$|\varphi_{\rm sh}(\iota,t) - \varphi_{\rm sh}(\iota,t_k)| \le \frac{1}{2} \int_{t_k}^t \sigma^2(\iota,t') \mathfrak{g}(\iota,t') dt' \le C e^{-2\eta_{\rm sh}t}$$

for all  $t \in [t_k, t_{k+1}]$ , where C only depends on the coefficients of the equation (1.2). Thus (18.73) can be rewritten

$$\left| w(\iota,t) - \mathfrak{D}_{\text{tot}}(\iota,t)e^{S_{n}(\iota)t}z_{\infty} \right| \leq C_{B}\langle \bar{t} \rangle^{N}e^{(\kappa_{n,+}-\eta_{B})\bar{t}} \left( |w(\iota,T_{n})| + e^{\kappa_{n,+}T_{n}} \int_{0}^{\infty} e^{-(\kappa_{n,+}-\eta_{B})t} |\hat{f}(\iota,t)| dt \right)$$
(18.74)

for all  $k \ge 0$  and  $t \in [t_k, t_{k+1}]$ , where  $C_B$  only depends on  $\eta_B$  and the coefficients of the equation (1.2) and N only depends on m. Moreover, in case all the Jordan blocks of  $S_n(\iota)$  are trivial, the

274

N appearing in (18.74) can be replaced by 1. In addition, if the first generalised eigenspace of the  $\eta_B$ ,  $S_n(\iota)$ -decomposition of  $\mathbb{C}^{2m}$  equals  $\mathbb{C}^{2m}$ , then the N appearing in (18.74) can be replaced by  $d_{n,+} - 1$ , where  $d_{n,+} = d_{\max}(S_n(\iota), \kappa_{n,+})$ . Finally,  $\mathfrak{D}_{\text{tot}}$  is given by (18.59), where  $\varphi_{\text{tot}}$  is defined by (18.60). The lemma follows.

## Chapter 19

## Asymptotics for all future times

## **19.1** Introduction

Asymptotics and asymptotic characterisation of solutions for a class of scalar equations. In the previous chapter we fixed one mode  $\iota \in \mathcal{I}_{B,n}$  and analysed the asymptotic behaviour of solutions to (24.16) in the corresponding late time regime  $[T_n, \infty)$ . In the present chapter, we derive estimates valid for all  $t \geq 0$  and all  $\iota \in \mathcal{I}_{B,n}$ . Similarly to the case of silent and transparent equations, our goal is to derive asymptotics, given a solution, and to construct solutions with given asymptotics. However, in the present setting, it is natural to view the asymptotics as composed of two parts. On the one hand, there is the oscillatory behaviour which is already apparent when considering scalar equations such as (19.1) below. Second, there is the overall growth and decay of solutions to systems such as (1.2). In order to analyse the oscillatory part of the behaviour, we consider the equation (19.1) separately in Section 19.2. In particular, we prove that solutions can, in this case, be characterised by asymptotic data, so that there is a bijection between asymptotic data and initial data. In fact, the bijection is even a homeomorphism with respect to the  $C^{\infty}$ topology. In the course of the analysis, it turns out to be natural to define two subclasses of solutions to (19.1). The subclasses appear naturally when considering the asymptotics, and lead to the definitions of positively and negatively oriented solutions in Definition 19.7 below.

Uniqueness of asymptotic representations. Before turning to the asymptotics of solutions to general equations, it is useful to prove a uniqueness result concerning the type of functions we use to represent the asymptotics later on. This is the subject of Section 19.3. It is of interest to note that the notion of an orientation, introduced in Definition 19.7, plays an important role in the formulation and proof of uniqueness. Moreover, the uniqueness result demonstrates that the function representing the asymptotic behaviour of a solution u to (1.2), say  $u_w$ , is uniquely determined by the estimate we derive for the difference between u and  $u_w$ .

Asymptotics for solutions to general equations. Given the above preliminaries, we describe the asymptotics of solutions to general equations in Section 19.4. Interestingly, it turns out that the functions representing the asymptotics can be written  $e^{A_{n,+}t}u_{+,w} + e^{A_{n,-}t}u_{-,w}$ , where the  $A_{n,\pm}$  are constant matrices determined by the coefficients of (1.2), and the  $u_{\pm,w}$  are vector valued solutions to (19.1). In fact, the  $u_{\pm,w}$  take their values in appropriate generalised eigenspaces of  $A_{n,\pm}$ , say  $E_{n,\pm}$ , respectively. In this representation of the asymptotics, the functions  $u_{\pm,w}$  should be thought of as constituting the asymptotic data. However, since there is a bijection between solutions to (19.1) and asymptotic data for solutions to (19.1), we could represent the asymptotic data of solutions to (1.2) in terms of the asymptotic data for the  $u_{\pm,w}$ .

Specifying the asymptotics of solutions to general equations. Finally, in Section 19.5 we demonstrate that, given solutions  $u_{\pm,w}$  to (19.1) taking their values in  $E_{n,\pm}$  respectively, there is a solution u to (1.2) (with f = 0) with the corresponding asymptotic behaviour. Moreover, the

corresponding map from asymptotic data (represented, e.g., by the initial data for the functions  $u_{\pm,w}$ ) to the initial data for u is continuous in the  $C^{\infty}$ -topology. Finally, under favourable circumstances, this map is surjective, giving a homeomorphism from asymptotic data for solutions to (1.2) (with f = 0) to the corresponding initial data.

## 19.2 The case of scalar homogeneous wave equations

Next we turn to the problem of specifying the asymptotic behaviour of solutions to the scalar homogeneous equation

$$u_{tt} - \sum_{j,l=1}^{d} g^{jl}(t) \partial_j \partial_l u - 2 \sum_{l=1}^{d} g^{0l}(t) \partial_l \partial_t u - \sum_{r=1}^{R} a_r^{-2}(t) \Delta_{g_r} u = 0.$$
(19.1)

As in Section 17.7, we focus on  $u_n$ , defined in analogy with (17.54).

**Lemma 19.1.** Consider the equation (19.1). Assume that it is strongly balanced, has a negligible shift vector field, a dominant noisy spatial direction, and is such that the dominant coefficients are convergent. Let  $\eta_A$  be defined by (17.25) and let  $0 < \eta_B \leq \eta_A$ . Let  $\chi_{\pm}(\iota) \in \mathbb{C}$ ,  $\iota \in \mathcal{I}_{B,n}$ , be such that for every  $0 \leq k \in \mathbb{Z}$ , there is a constant  $C_k < \infty$  with the property that

$$\sum_{\iota \in \mathcal{I}_{\mathrm{B,n}}} \langle \nu(\iota) \rangle^{2k} [|\chi_+(\iota)|^2 + |\chi_-(\iota)|^2] \le C_k.$$
(19.2)

Define  $z_{app}$  and  $u_{app}$  by

$$z_{\rm app}(\iota,t) := \frac{e^{\beta_{\rm n}t/2}}{\mathfrak{g}(\iota,t)} \left( \chi_+(\iota) e^{i[\omega_{\rm sh}(\iota,t) - \varphi_{\rm tot}(\iota,t)]} + \chi_-(\iota) e^{i[\omega_{\rm sh}(\iota,t) + \varphi_{\rm tot}(\iota,t)]} \right)$$
(19.3)

$$u_{\rm app}(p,t) := \sum_{\iota \in \mathcal{I}_{{\rm B},{\rm n}}} z_{\rm app}(\iota,t)\varphi_{\iota}(p), \tag{19.4}$$

where  $\iota \in \mathcal{I}_{B,n}$  in (19.3);  $\omega_{sh}$  is defined by (18.58);  $\varphi_{tot}$  is defined by (18.60); and  $\mathfrak{g}(\iota, t)$  is defined by (24.13). Then there is a unique smooth solution u to (19.1) such that the following holds:  $u = u_n$ , where  $u_n$  is defined in analogy with (17.54); and there are constants  $s \in \mathbb{R}$ ,  $\eta > 0$  and C > 0 such that

$$\mathfrak{E}_s[u - u_{\mathrm{app}}](t) \le C e^{(\beta_n - \eta)t}$$
(19.5)

for all  $t \geq 0$ . Moreover, this unique solution has the property that

$$\mathfrak{E}_{s}[u-u_{\mathrm{app}}](t) \leq C_{B} e^{(\beta_{\mathrm{n}}-2\eta_{B})t} \sum_{\iota \in \mathcal{I}_{\mathrm{B},\mathrm{n}}} \langle \nu(\iota) \rangle^{2(s+s_{\mathrm{m}})} [r_{+}^{2}(\iota) + r_{-}^{2}(\iota)]$$
(19.6)

for all  $t \ge 0$  and  $s \in \mathbb{R}$ , where  $C_B$  and  $s_m$  only depend on  $\eta_B$  and the coefficients of the equation (19.1);  $\mathfrak{E}_s$  is defined by (24.19); and  $r_{\pm}(\iota) = |\chi_{\pm}(\iota)|$ . Finally,

$$\mathfrak{E}_{s}^{1/2}[u](0) \le C_{B} \left( \sum_{\iota \in \mathcal{I}_{B,n}} \langle \nu(\iota) \rangle^{2(s+s_{m})} [r_{+}^{2}(\iota) + r_{-}^{2}(\iota)] \right)^{1/2}$$
(19.7)

for all  $s \in \mathbb{R}$ , where  $C_B$  and  $s_m$  have the same dependence as in the case of (19.6).

Conversely, assume that u is a smooth solution to (19.1) such that  $u = u_n$ . Then there are uniquely determined  $\chi_{\pm}(\iota)$ ,  $\iota \in \mathcal{I}_{B,n}$ , such that (19.2) holds, and such that if  $u_{app}$  is defined by (19.3) and (19.4), then (19.5) holds. Moreover, this  $u_{app}$  is such that (19.6) holds. Finally,

$$\left(\sum_{\iota \in \mathcal{I}_{B,n}} \langle \nu(\iota) \rangle^{2s} [|\chi_{+}(\iota)|^{2} + |\chi_{-}(\iota)|^{2}]\right)^{1/2} \le C_{B} \mathfrak{E}_{s+s_{m}}^{1/2} [u](0)$$
(19.8)

for all  $s \in \mathbb{R}$ , where  $C_B$  and  $s_m$  have the same dependence as in the case of (19.6).

**Remark 19.2.** Assume that (19.1) is  $C^2$ -balanced with a geometric dominant noisy spatial direction, convergent dominant coefficients and a negligible shift vector field; cf. Definition 6.5. Then the assumptions of the lemma are satisfied; cf. Remarks 17.2 and 17.6. Moreover, if  $\eta_{\rm sh}$ ,  $\eta_{\rm n}$ ,  $\beta_{\rm n}$  and  $\eta_{\rm mn}$  are the constants arising in Definition 6.5, then  $\eta_{\rm sh}$ ,  $\beta_{\rm n}$  and  $\eta_{\rm mn}$  are the same as in the present context. Moreover,  $\eta_{\rm mar} = \min\{2\eta_{\rm sh}, \eta_{\rm n}\}$  in the case of a geometric dominant noisy spatial direction  $\mathbb{S}^1$ -direction. In addition,  $\eta_{\rm mar} = \eta_{\rm n}$  in the case of a geometric dominant noisy spatial direction corresponding to, say,  $M_r$ . In both cases,  $\eta_A = \min\{\eta_{\rm n}/2, \eta_{\rm sh}, \eta_{\rm mn}, \beta_{\rm n}\}$ .

**Remark 19.3.** The function  $u_{app}$  is smooth and complex valued; the smoothness is an immediate consequence of the definition and (19.2).

**Remark 19.4.** The way the lemma is formulated, there is no apparent gain in letting  $\eta_B$  be strictly smaller than  $\eta_A$ . However, the value of  $s_m$  depends on  $\eta_B$ ; the smaller the  $\eta_B$ , the smaller the  $s_m$ .

**Remark 19.5.** Dependence on the coefficients includes dependence on the Riemannian manifolds  $(M_r, g_r), r = 1, \ldots, R$ ; cf. Remark 17.10.

**Remark 19.6.** The constants C, s and  $\eta$  appearing in (19.5) are allowed to depend on u,  $u_{app}$  etc.

*Proof.* The proof is naturally divided into four parts. First, we demonstrate uniqueness. Second, we consider one mode, constructing a solution corresponding to given asymptotic data. Third, we sum up the estimates for the different modes. Finally, we arrive at asymptotics starting with a solution to (19.1) such that  $u = u_n$ .

**Uniqueness.** Assume that there are two solutions, say  $u_l$ , l = 1, 2, to (19.1) such that  $u_l = (u_l)_n$  and such that (19.5) holds with u replaced by  $u_l$ . Then the difference, say  $u := u_1 - u_2$ , satisfies

$$\mathfrak{E}_{\mathbf{s}}[u](t) < Ce^{(\beta_{n}-\eta)t}$$

for all  $t \ge 0$ . Defining z by (24.14) and letting  $\iota \in \mathcal{I}_{B,n}$ , this implies that

$$|\dot{z}(\iota,t)| + \mathfrak{g}(\iota,t)|z(\iota,t)| < Ce^{(\beta_{n}-\eta)t/2}$$

for all  $t \ge 0$ , where *C* is allowed to depend on  $u_l$ , l = 1, 2,  $\iota$ , etc. We wish to compare this estimate with the conclusions of Lemma 18.1. Note, to this end, that  $R_n(\iota) = \beta_n Id_2/2$  in the present context. Moreover,  $\kappa_{n,+} = \beta_n/2$ . Combining these observations, it is clear that the only way for (18.2) to be satisfied is that  $\psi_{\infty} = 0$ . In fact,  $\psi_{\infty}(\iota) = 0$  for all  $\iota \in \mathcal{I}_{B,n}$ .

Let us now assume that  $u \neq 0$ . Then there is a  $\iota \in \mathcal{I}_{B,n}$  such that  $z(\iota, \cdot) \neq 0$ . In particular, the initial data for  $z(\iota, \cdot)$  at  $T_n$  are non-zero. In order to obtain uniqueness, we demonstrate that this statement contradicts the conclusions of Lemma 18.5. Using the notation of Lemma 18.5, note that m = 1 and that  $E_a = \mathbb{C}^2$  in the present setting. Since  $\Psi_{\infty}$  is injective, we conclude that it is surjective. There is thus a  $\chi \in \mathbb{C}^2$  such that (18.37) holds. Since the right hand side of this equality is non-zero, it is clear that  $\chi$  is non-zero. Moreover, Lemma 18.5 implies that (18.2) holds with  $\psi_{\infty}$  replaced by  $\chi$ . By the above argument, we conclude that  $\chi = 0$ , a contradiction. Thus u = 0, and uniqueness follows.

Uniqueness of the approximation. Let  $u = u_n$  be a solution to (19.1). Assume that there are  $\chi_{\pm,l}(\iota), \iota \in \mathcal{I}_{B,n}, l = 1, 2$ , such that (19.2) holds with  $\chi_{\pm}(\iota)$  replaced by  $\chi_{\pm,l}(\iota)$ , and such that if  $u_{app,l}$  is defined by (19.3) and (19.4) with  $\chi_{\pm}(\iota)$  replaced by  $\chi_{\pm,l}(\iota)$ , then (19.5) holds with  $u_{app}$  replaced by  $u_{app,l}, l = 1, 2$ . Then

$$\mathfrak{E}_s[u_{\mathrm{app}}](t) \le C e^{(\beta_{\mathrm{n}} - \eta)t} \tag{19.9}$$

for all  $t \ge 0$ , where  $u_{\text{app}} := u_{\text{app},1} - u_{\text{app},2}$ . Note that  $u_{\text{app}}$  is defined by (19.3) and (19.4), where  $\chi_{\pm} = \chi_{\pm,1} - \chi_{\pm,2}$ . Fix a  $\iota \in \mathcal{I}_{\text{B,n}}$  and consider  $z_{\text{app}}(\iota, \cdot)$  given by (19.3). Due to (19.9), we know that

$$\lim_{t \to \infty} e^{-\beta_{\mathrm{n}} t/2} \mathfrak{g}(\iota, t) z_{\mathrm{app}}(\iota, t) = 0, \quad \lim_{t \to \infty} e^{-\beta_{\mathrm{n}} t/2} \dot{z}_{\mathrm{app}}(\iota, t) = 0.$$
(19.10)

The first equality implies that

$$\lim_{t \to \infty} \left( \chi_+(\iota) e^{i[\omega_{\rm sh}(\iota,t) - \varphi_{\rm tot}(\iota,t)]} + \chi_-(\iota) e^{i[\omega_{\rm sh}(\iota,t) + \varphi_{\rm tot}(\iota,t)]} \right) = 0.$$
(19.11)

Appealing to the definitions of  $\omega_{\rm sh}$  and  $\varphi_{\rm tot}$ ; the assumed bounds on  $\sigma$  and  $\dot{\ell}$ ; and the fact that  $e^{-\beta_{\rm n}t}\mathfrak{g}(\iota,t)$  converges to a non-zero number (cf. Lemma 17.13), it can be verified that the second equality in (19.10) implies that

$$\lim_{t \to \infty} \left( -\chi_+(\iota) e^{i[\omega_{\rm sh}(\iota,t) - \varphi_{\rm tot}(\iota,t)]} + \chi_-(\iota) e^{i[\omega_{\rm sh}(\iota,t) + \varphi_{\rm tot}(\iota,t)]} \right) = 0.$$
(19.12)

Combining (19.11) and (19.12) yields the conclusion that  $\chi_+(\iota) = \chi_-(\iota) = 0$  for all  $\iota \in \mathcal{I}_{B,n}$ . Thus  $\chi_{\pm,1}(\iota) = \chi_{\pm,2}(\iota)$  for all  $\iota \in \mathcal{I}_{B,n}$ , and the desired uniqueness follows.

**One mode.** We wish to appeal to Lemmas 18.5 and 18.8. Let, to this end, z denote the  $\iota$ 'th Fourier coefficient of a solution u to (19.1). Then z satisfies

$$\ddot{z}(\iota,t) - 2i\sigma(\iota,t)\mathfrak{g}(\iota,t)\dot{z}(\iota,t) + \mathfrak{g}^2(\iota,t)z(\iota,t) = 0.$$
(19.13)

As is clear from the statement of the lemma, we are here only interested in  $\iota \in \mathcal{I}_{B,n}$ . We therefore assume  $\iota \in \mathcal{I}_{B,n}$  from now on. When comparing the present setting with Lemma 18.8, it is clear that  $\alpha = 0$  and X = 0, so that  $S_n(\iota) = \beta_n Id_2/2$ . Thus (18.62) with  $0 < \eta_B \le \eta_A$  yields

$$\begin{vmatrix} \begin{pmatrix} \mathfrak{g}(\iota,t)z(\iota,t) \\ \dot{z}(\iota,t) \end{pmatrix} - e^{i\omega_{\rm sh}(\iota,t)}e^{\beta_{\rm n}t/2}\mathfrak{D}_{\rm tot}(\iota,t)z_{\infty} \\ \leq C_B e^{(\beta_{\rm n}/2 - \eta_B)\bar{t}} \left(|\dot{z}(\iota,T_{\rm n})| + \mathfrak{g}(\iota,T_{\rm n})|z(\iota,T_{\rm n})|\right) \end{aligned}$$
(19.14)

for all  $t \ge T_n$ , where  $\bar{t} := t - T_n$ ;  $z_\infty \in \mathbb{C}^2$ ;  $T_n$  is defined by (17.12); and  $C_B$  only depends on  $\eta_B$ and the coefficients of the equation (19.1). In order to justify the absence of polynomial powers in the estimate (19.14), recall Remark 18.10. Note, in particular, that since the first generalised eigenspace of the  $\eta_B$ ,  $S_n(\iota)$ -decomposition of  $\mathbb{C}^2$  equals  $\mathbb{C}^2$ , N appearing in (18.62) can be replaced by  $d_{n,+} - 1$ . Moreover, since  $d_{n,+} = 1$  in the present setting, N can be replaced by 0. In what follows, it is useful to keep in mind that, due to Lemma 18.5,  $z_\infty$  is freely specifiable. The reason for this is that  $E_a$  appearing in the statement of Lemma 18.5 equals  $\mathbb{C}^2$ . Moreover, the reason why  $E_a = \mathbb{C}^2$  is that  $\operatorname{Rsp}[R_n(\iota)] = 0$ ; cf. Definition 4.3. Recall, finally, that the  $\psi_\infty$  appearing in (18.2) is related to the  $z_\infty$  appearing in (18.62) according to  $z_\infty = T_{\mathrm{pre}}^{-1}\psi_\infty$ . Let now z be the solution to (19.13) corresponding to asymptotic data  $z_\infty$ . Then Lemma 18.5 yields

$$|w(\iota, T_{n})| \le |\dot{z}(\iota, T_{n})| + \mathfrak{g}(\iota, T_{n})|z(\iota, T_{n})| \le Ce^{\beta_{n}T_{n}/2}|z_{\infty}|,$$
(19.15)

where the constant C only depends on the coefficients of the equation (19.1). Denote the components of  $z_{\infty}$  by  $z_1$  and  $z_2$ . Then (19.14) can be written

$$\left| \begin{pmatrix} \mathfrak{g}(\iota,t)z(\iota,t) \\ \dot{z}(\iota,t) \end{pmatrix} - e^{i\omega_{\rm sh}(\iota,t)}e^{\beta_{\rm n}t/2} \begin{pmatrix} z_1\cos\varphi_{\rm tot}(\iota,t) + z_2\sin\varphi_{\rm tot}(\iota,t) \\ -z_1\sin\varphi_{\rm tot}(\iota,t) + z_2\cos\varphi_{\rm tot}(\iota,t) \end{pmatrix} \right|$$

$$\leq C_B e^{(\beta_{\rm n}/2 - \eta_B)\bar{t}}e^{\beta_{\rm n}T_{\rm n}/2}|z_{\infty}|$$

$$(19.16)$$

for all  $t \ge T_n$ , where the constant  $C_B$  has the same dependence as in the case of (19.14). Let

$$z_{\rm app}(\iota,t) := \frac{1}{\mathfrak{g}(\iota,t)} e^{i\omega_{\rm sh}(\iota,t)} e^{\beta_{\rm h}t/2} [z_1 \cos\varphi_{\rm tot}(\iota,t) + z_2 \sin\varphi_{\rm tot}(\iota,t)].$$
(19.17)

Then

$$\dot{z}_{app}(\iota,t) = \left[-\dot{\ell}(\iota,t) + \frac{\beta_{n}}{2} + i\dot{\omega}_{sh}(\iota,t)\right] z_{app}(\iota,t) + \frac{1}{\mathfrak{g}(\iota,t)}\dot{\varphi}_{tot}(\iota,t)e^{i\omega_{sh}(\iota,t)}e^{\beta_{n}t/2}[-z_{1}\sin\varphi_{tot}(\iota,t) + z_{2}\cos\varphi_{tot}(\iota,t)].$$
(19.18)
Due to (18.58) and (18.60),

$$\dot{\varphi}_{\text{tot}}(\iota, t) = [1 + \sigma^2(\iota, t)]^{1/2} \mathfrak{g}(\iota, t), \quad \dot{\omega}_{\text{sh}}(\iota, t) = \sigma(\iota, t) \mathfrak{g}(\iota, t).$$

For this reason, the absolute value of the first term on the right hand side of (19.18) can be bounded by the right hand side of (19.16). Moreover, in the second term on the right hand side of (19.18), we can replace  $\dot{\varphi}_{tot}(\iota, t)$  by  $\mathfrak{g}(\iota, t)$  at the price of an error term whose absolute value can be estimated by the right hand side of (19.16). To conclude, (19.16) can be rewritten

$$\left| \begin{pmatrix} \mathfrak{g}(\iota,t)z(\iota,t) \\ \dot{z}(\iota,t) \end{pmatrix} - \begin{pmatrix} \mathfrak{g}(\iota,t)z_{\mathrm{app}}(\iota,t) \\ \dot{z}_{\mathrm{app}}(\iota,t) \end{pmatrix} \right| \le C_B e^{(\beta_{\mathrm{n}}/2 - \eta_B)\bar{t}} e^{\beta_{\mathrm{n}}T_{\mathrm{n}}/2} |z_{\infty}|$$
(19.19)

for all  $t \ge T_n$ , where the constant  $C_B$  has the same dependence as in the case of (19.14). It is convenient to rewrite (19.17) by introducing  $\chi_{\pm}$  according to

$$\chi_{\pm} = \frac{1}{2}(z_1 \pm i z_2). \tag{19.20}$$

Then (19.17) becomes

$$z_{\rm app}(\iota,t) = \frac{e^{\beta_{\rm n}t/2}}{\mathfrak{g}(\iota,t)} \left( \chi_+ e^{i[\omega_{\rm sh}(\iota,t) - \varphi_{\rm tot}(\iota,t)]} + \chi_- e^{i[\omega_{\rm sh}(\iota,t) + \varphi_{\rm tot}(\iota,t)]} \right)$$

In order for (19.19) to be of use when summing up the modes, we need to know that it holds for all  $t \geq 0$ . We therefore need to consider the interval  $[0, T_n]$  in greater detail. The fact that the equation is strongly balanced implies that Lemma 10.7 is applicable (with  $\mathfrak{e} = 0$ ). Thus, we are allowed to combine (10.12) and (19.15) in order to conclude

$$\mathcal{E}^{1/2}(\iota, t) \leq e^{\eta_{\text{bal}}|t-T_{\text{n}}|} \mathcal{E}^{1/2}(\iota, T_{\text{n}}) \leq C e^{\eta_{\text{bal}}T_{\text{n}}} e^{\beta_{\text{n}}T_{\text{n}}/2} |z_{\infty}| \\
\leq C e^{(\beta_{\text{n}}/2 - \eta_{B})t} e^{\eta_{\text{m}}T_{\text{n}}} |z_{\infty}|$$
(19.21)

for all  $t \in [0, T_n]$ , where  $\eta_{\text{bal}}$  and C only depend on the coefficients of the equation (19.1). Moreover,

$$\eta_{\mathrm{m}} := \max\{\eta_B - \beta_{\mathrm{n}}/2, 0\} + \eta_{\mathrm{bal}} + \beta_{\mathrm{n}}/2.$$

On the other hand, the definition of  $z_{app}$  implies that

$$\mathfrak{g}(\iota,t)|z_{\mathrm{app}}(\iota,t)| + |\dot{z}_{\mathrm{app}}(\iota,t)| \le C e^{(\beta_{\mathrm{n}}/2 - \eta_{B})t} e^{\eta_{B}T_{\mathrm{n}}}|z_{\infty}|$$

for all  $t \in [0, T_n]$ , where C only depends on the coefficients of the equation (19.1). Combining this estimate with (19.19) and (19.21) yields

$$\left| \begin{pmatrix} \mathfrak{g}(\iota,t)z(\iota,t) \\ \dot{z}(\iota,t) \end{pmatrix} - \begin{pmatrix} \mathfrak{g}(\iota,t)z_{\mathrm{app}}(\iota,t) \\ \dot{z}_{\mathrm{app}}(\iota,t) \end{pmatrix} \right| \le C_B e^{(\beta_{\mathrm{n}}/2 - \eta_B)t} e^{\eta_{\mathrm{m}}T_{\mathrm{n}}} |z_{\infty}|$$
(19.22)

for all  $t \ge 0$ , where the constant  $C_B$  has the same dependence as in the case of (19.14).

Summing up the estimates for the individual modes. Let  $\chi_{\pm}(\iota)$  be as in the statement of the lemma, and define  $z_{app}$  and  $u_{app}$  according to (19.3) and (19.4) respectively. In order to construct the solution u with the desired properties, let  $z(\iota, t)$  be the solution to (19.13) corresponding to the asymptotic data specified by  $\chi_{\pm}(\iota)$ . In order to translate (19.22) into an estimate for u, we need to estimate  $e^{\eta_m T_n}$ . Note, to this end, that

$$T_{\rm n} \le \frac{2}{\eta_{\rm mar}} \ln \langle \nu(\iota) \rangle + C,$$
 (19.23)

where C only depends on the coefficients of the equation (19.1); cf. Definition 17.12. Thus

$$e^{\eta_{\mathrm{m}}T_{\mathrm{n}}} \leq C_B \langle \nu(\iota) \rangle^{s_{\mathrm{m},1}},$$

where  $s_{m,1} = 2\eta_m/\eta_{mar}$  and the constant  $C_B$  only depends on  $\eta_B$  and the coefficients of the equation (19.1). Thus (19.22) implies

$$\left| \begin{pmatrix} \mathfrak{g}(\iota,t)z(\iota,t) \\ \dot{z}(\iota,t) \end{pmatrix} - \begin{pmatrix} \mathfrak{g}(\iota,t)z_{\mathrm{app}}(\iota,t) \\ \dot{z}_{\mathrm{app}}(\iota,t) \end{pmatrix} \right| \leq C_B e^{(\beta_n/2 - \eta_B)t} \langle \nu(\iota) \rangle^{s_{\mathrm{m},1}} [r_+(\iota) + r_-(\iota)]$$
(19.24)

for all  $t \ge 0$ , where the constant  $C_B$  has the same dependence as in the case of (19.14) and  $r_{\pm}(\iota) := |\chi_{\pm}(\iota)|$ . Similarly, (19.21) implies that

$$\mathcal{E}^{1/2}(\iota,0) \le C_B \langle \nu(\iota) \rangle^{s_{m,1}} [r_+(\iota) + r_-(\iota)], \qquad (19.25)$$

where the constant  $C_B$  has the same dependence as in the case of (19.14).

In order to prove (19.6), we need to estimate the energy  $\mathcal{E}_s$  of  $z - z_{app}$ . Such an estimate almost follows from (19.24). However, there is one piece missing: we need to estimate  $\mathfrak{g}(\iota, t)$  from below. The estimate  $|\dot{\ell}(\iota, t)| \leq C_{\ell}$  for all  $t \geq 0$  and  $0 \neq \iota \in \mathcal{I}_{B}$ , where  $C_{\ell}$  only depends on the coefficients of the equation, implies that

$$\left|\ln\frac{\mathfrak{g}(\iota,T_{n})}{\mathfrak{g}(\iota,t)}\right| \leq C_{\ell}T_{n}$$

for all  $t \in [0, T_n]$ . In particular,

$$e^{\beta_{\mathbf{n}}t}[\mathfrak{g}(\iota,t)]^{-1} \le e^{(C_{\ell}+\beta_{\mathbf{n}})T_{\mathbf{n}}} \le C\langle\nu(\iota)\rangle^{s_E}$$
(19.26)

for all  $t \in [0, T_n]$ , where C and  $s_E$  only depend on the coefficients of the equation (19.1) and we appealed to (17.13) and (19.23). On the other hand, (17.14) yields the conclusion that  $2\mathfrak{g}(\iota, t) \geq \nu_n(\iota)e^{\beta_n t}$  for all  $t \geq T_n$ . Combining this estimate with (19.26) yields the conclusion that

$$[\mathfrak{g}(\iota,t)]^{-1} \le C \langle \nu(\iota) \rangle^{s_E} e^{-\beta_n t}$$
(19.27)

for all  $t \ge 0$ , where C and  $s_E$  only depend on the coefficients of the equation (19.1). Combining this estimate with (19.24) yields

$$\mathcal{E}_{s}^{1/2}[z - z_{\rm app}](\iota, t) \le C_{B} e^{(\beta_{\rm n}/2 - \eta_{B})t} \langle \nu(\iota) \rangle^{s + s_{\rm m}} [r_{+}(\iota) + r_{-}(\iota)]$$
(19.28)

for all  $t \ge 0$ , where  $C_B$  only depends on  $\eta_B$  and the coefficients of the equation (19.1) and  $s_m = s_{m,1} + s_E$ . Define u by

$$u(p,t) := \sum_{\iota \in \mathcal{I}_{\mathrm{B,n}}} z(\iota,t) \varphi_{\iota}(p).$$

Due to the estimate (19.28), the definition of  $z_{app}$ , (19.2) and the fact that  $z(\iota, t)$  solves (19.13), it is clear that u is a smooth solution to (19.1). Due to (19.28) it is clear that (19.6) holds. Similarly, (19.25) yields (19.7).

**Deriving asymptotics for a given solution.** Finally, let us turn to the converse statement. Let u be a smooth solution to (19.1) such that  $u = u_n$ . Then we can appeal to Lemma 18.8. In particular, if z is the  $\iota$ 'th mode of u, where  $\iota \in \mathcal{I}_{B,n}$ , the estimate (18.62) yields

$$\left| \begin{pmatrix} \mathfrak{g}(\iota,t)z(\iota,t) \\ \dot{z}(\iota,t) \end{pmatrix} - e^{i\omega_{\rm sh}(\iota,t)}\mathfrak{D}_{\rm tot}(\iota,t)e^{\beta_{\rm n}t/2}z_{\infty} \right| \\
\leq C_B e^{(\beta_{\rm n}/2 - \eta_B)\bar{t}} \left( |\dot{z}(\iota,T_{\rm n})| + \mathfrak{g}(\iota,T_{\rm n})|z(\iota,T_{\rm n})| \right),$$
(19.29)

where  $C_B$  only depends on  $\eta_B$  and the coefficients of the equation (1.2). Moreover, combining Lemma 10.7 (with  $\mathfrak{e} = 0$ ) and (18.61) yields

$$|z_{\infty}| \leq C_B e^{-\beta_n T_n/2} \left( |\dot{z}(\iota, T_n)| + \mathfrak{g}(\iota, T_n)| z(\iota, T_n) | \right)$$
  
$$\leq C_B e^{\eta_{\text{bal}} T_n - \beta_n T_n/2} \mathcal{E}^{1/2}(\iota, 0), \qquad (19.30)$$

where  $C_B$  has the same dependence as in the case of (19.29). Define  $\chi_{\pm}(\iota)$  according to (19.20), where  $z_1$ ,  $z_2$  are the components of  $z_{\infty}$ . Then (19.23) and (19.30) yield (19.8). In particular, the  $\chi_{\pm}(\iota)$  satisfy (19.2). Thus we can associate a smooth solution, say  $\bar{u}$ , with the asymptotic data  $\chi_{\pm}(\iota)$  as in the first part of the statement of the lemma. However, since the Fourier coefficients of u satisfy (19.29) and  $\bar{u}$  satisfies (19.6) (with u replaced by  $\bar{u}$ ), the Fourier coefficients of u and  $\bar{u}$  have to coincide, so that  $u = \bar{u}$ ; the reason the Fourier coefficients have to coincide is that they have the same asymptotic data, and that, due to Lemma 18.5, the asymptotic data uniquely determine the solution in the present setting. The lemma follows.

Due to Lemma 19.1, we can assign a meaning to the notion of orientation of a complex valued solution to (19.1).

**Definition 19.7.** Consider the equation (19.1). Assume that it is strongly balanced, has a negligible shift vector field, a dominant noisy spatial direction, and is such that the dominant coefficients are convergent. Assume that u is a smooth solution to (19.1) such that  $u = u_n$ , where  $u_n$  is defined in analogy with (17.54). Let  $u_{app}$  be the uniquely associated function of the form (19.3) and (19.4), whose existence is guaranteed by Lemma 19.1. Consider (19.3) and (19.4). If  $\chi_{-}(\iota) = 0$  for all  $\iota \in \mathcal{I}_{B,n}$ , then u is said to be *positively oriented*. If  $\chi_{+}(\iota) = 0$  for all  $\iota \in \mathcal{I}_{B,n}$ , then u is said to be *negatively oriented*.

#### **19.3** A uniqueness result

Next, we demonstrate a uniqueness result for vector valued solutions to (19.1).

**Lemma 19.8.** Consider the equation (19.1). Assume that it is strongly balanced, has a negligible shift vector field, a dominant noisy spatial direction, and is such that the dominant coefficients are convergent. Let  $1 \le k \in \mathbb{Z}$ ;  $A_{\pm} \in \mathbf{M}_k(\mathbb{C})$ ; and  $0 < \beta \in \mathbb{R}$ . Using the notation of Definitions 4.3 and 4.7, let

$$\kappa_{\rm m} = \max\{\kappa_{\rm max}(A_+), \kappa_{\rm max}(A_-)\};$$

 $J := (\kappa_{\rm m} - \beta, \kappa_{\rm m}]; and E_{\pm} := E_{A_{\pm},J}.$  Let  $u_{\pm} \in C^{\infty}(M, E_{\pm})$  be two solutions to (19.1) and denote the *i*'th Fourier coefficient of  $u_{\pm}(\cdot, t)$  by  $z_{\pm}(\iota, t)$  for  $\iota \in \mathcal{I}_{\rm B}$ . Assume that the following holds:

- $z_{\pm}(\iota, t) = 0$  for all  $\iota \notin \mathcal{I}_{B,n}$  and all  $t \in \mathbb{R}$ ,
- the components of  $u_+$  are positively oriented and the components of  $u_-$  are negatively oriented.

Define u by

$$u(p,t) := e^{A_+ t} u_+(p,t) + e^{A_- t} u_-(p,t).$$
(19.31)

Finally, assume that there is an  $s \in \mathbb{R}$  and constants C and N, which are allowed to depend on the solutions  $u_+$ , such that

$$\mathfrak{E}_{s}^{1/2}[u](t) \le C\langle t \rangle^{N} e^{(\kappa_{\mathrm{m}} + \beta_{\mathrm{n}}/2 - \beta)t}$$
(19.32)

for all  $t \ge 0$ . Then  $u_+ = u_- = 0$ .

**Remarks 19.9.** The notion of positive and negative orientation is introduced in Definition 19.7. The notation  $\mathfrak{E}_s[u](t)$  is introduced in (24.19).

*Proof.* Note, to begin with, that if  $z(\iota, t)$  is the  $\iota$ 'th Fourier coefficient of the function  $u(\cdot, t)$ , defined by (19.31), then

$$z(\iota, t) = e^{A_+ t} z_+(\iota, t) + e^{A_- t} z_-(\iota, t),$$

where the  $z_{\pm}$  are defined in the statement of the lemma. Note also that, due to the assumptions of the lemma, there is no restriction in assuming  $\iota \in \mathcal{I}_{B,n}$ . We therefore do so here and in what follows. Due to (19.32),

$$|\dot{z}(\iota,t)| + \mathfrak{g}(\iota,t)|z(\iota,t)| \le C\langle t \rangle^N e^{(\kappa_{\mathrm{m}} + \beta_{\mathrm{n}}/2 - \beta)t}$$
(19.33)

for all  $t \ge 0$ , where C depends on  $\iota$ , s and the solutions  $u_{\pm}$ . Note that  $e^{A_{\pm}t} z_{\pm}(\iota, t)$  can be written

$$e^{A_{\pm}t} z_{\pm}(\iota, t) = \sum_{r=1}^{k_{\pm}} \phi_{\pm, r}(t) z_{\pm, r}(\iota, t), \qquad (19.34)$$

where the  $\phi_{\pm,r}$  are linearly independent  $E_{\pm}$ -valued solutions to the equation  $\dot{x} = A_{\pm}x$  and  $z_{\pm,r}$ are  $\mathbb{C}$ -valued solutions to (19.13). Moreover, the  $\phi_{\pm,r}$  are of the form

$$\phi_{\pm,r}(t) = e^{\lambda_{\pm,r}t} \psi_{\pm,r}(t), \tag{19.35}$$

where  $\lambda_{\pm,r}$  is an eigenvalue of  $A_{\pm}$  and  $\psi_{\pm,r}$  is an  $E_{\pm}$ -valued function, each of whose components are polynomials (in fact,  $\psi_{\pm,r}$  (and thereby  $\phi_{\pm,r}$ ) takes its values in the generalised eigenspace corresponding to  $\lambda_{\pm,r}$ ). In what follows, we denote the real part of  $\lambda_{\pm,r}$  by  $\kappa_{\pm,r}$ . The ordering of the  $\phi_{\pm,r}$  is such that if  $r_1 < r_2$ , then  $\kappa_{\pm,r_1} \ge \kappa_{\pm,r_2}$ . In addition, the  $z_{+,r}$ 's correspond to positively oriented solutions to (19.1) and the  $z_{-,r}$ 's correspond to negatively oriented solutions to (19.1). Finally, note that we, without loss of generality, can assume that  $\kappa_{\pm,r} > \kappa_{\rm m} - \beta$  for all r. Turning to the  $z_{\pm,r}$ 's, note that, due to Lemma 19.1, they can be approximated by

$$z_{\pm,r,\mathrm{app}}(\iota,t) := \frac{e^{\beta_{\mathrm{n}}t/2}}{\mathfrak{g}(\iota,t)} \chi_{\pm,r}(\iota) e^{i[\omega_{\mathrm{sh}}(\iota,t)\mp\varphi_{\mathrm{tot}}(\iota,t)]}.$$
(19.36)

In fact, there is a constant  $C_{\pm,r}$ , depending on  $z_{\pm,r}$  and  $\iota$ , such that

$$|\dot{z}_{\pm,r}(\iota,t) - \dot{z}_{\pm,r,\mathrm{app}}(\iota,t)| + \mathfrak{g}(\iota,t)|z_{\pm,r}(\iota,t) - z_{\pm,r,\mathrm{app}}(\iota,t)| \le C_{\pm,r}e^{(\beta_n/2 - \eta_B)t}$$
(19.37)

for all  $t \ge 0$ ; cf. Lemma 19.1, in particular (19.6). Thus

$$|\dot{z}_{\pm,r}(\iota,t)| + \mathfrak{g}(\iota,t)|z_{\pm,r}(\iota,t)| \le C_{\pm,r}e^{\beta_{n}t/2}$$
(19.38)

for all  $t \ge 0$ . The argument proceeds by a consideration of several different cases.

**Different**  $\kappa_{\pm,1}$ 's. Say now, for the sake of argument, that  $\kappa_{+,1} > \kappa_{-,1}$ . Assume, moreover, that  $\kappa_{+,r} = \kappa_{+,1}$  for  $r = 1, \ldots, r_a$ , and that  $\kappa_{+,r} < \kappa_{+,1}$  for  $r > r_a$ . Then

$$\lim_{t \to \infty} e^{-(\kappa_{+,1} + \beta_n/2)t} \mathfrak{g}(\iota, t) e^{A_- t} z_-(\iota, t) = 0,$$
(19.39)

where we have appealed to (19.34), (19.35), (19.38) and the fact that  $\kappa_{-,1} < \kappa_{+,1}$ . Combining (19.39) with a similar argument concerning the sum on the right hand side of (19.34) in the case of the plus sign and for  $r > r_a$  yields

$$\lim_{t \to \infty} e^{-(\kappa_{+,1} + \beta_n/2)t} \sum_{r=1}^{r_a} \mathfrak{g}(\iota, t) e^{\lambda_{+,r} t} \psi_{+,r}(t) z_{+,r}(\iota, t) = 0,$$

where we have also appealed to (19.33), (19.35) and the fact that  $\kappa_{\pm,r} > \kappa_{\rm m} - \beta$ . Combining this limit with (19.36) and (19.37) yields

$$\lim_{t \to \infty} \sum_{r=1}^{r_a} e^{(\lambda_{+,r} - \kappa_{+,1})t} \psi_{+,r}(t) \chi_{+,r}(\iota) = 0.$$

Since  $\psi_{+,r}$  takes its values in the generalised eigenspace corresponding to  $\lambda_{+,r}$ , we may focus on each distinct  $\lambda_{+,r}$  separately. In other words, we may, in practice, assume that  $\lambda_{+,r}$  is independent of r. Thus

$$\lim_{t \to \infty} \sum_{r=1}^{r_a} \psi_{+,r}(t) \chi_{+,r}(\iota) = 0.$$
(19.40)

Let the non-negative integers  $m_1 > m_2 > \cdots > m_p$  be such that all the polynomial powers occurring in the functions  $\psi_{+,r}$  belong to  $\{m_1, \ldots, m_p\}$ . Then  $\psi_{+,r}(t)$  can be written

$$\psi_{+,r}(t) = \sum_{j=1}^{p} t^{m_j} v_{r,j},$$

where  $v_{r,j} \in \mathbb{C}^k$ . Due to (19.40) this observation yields

$$0 = \lim_{t \to \infty} t^{-m_1} \sum_{r=1}^{r_a} \psi_{+,r}(t) \chi_{+,r}(\iota) = \sum_{r=1}^{r_a} v_{r,1} \chi_{+,r}(\iota).$$

Proceeding power by power yields

$$\sum_{r=1}^{r_a} v_{r,j} \chi_{+,r}(\iota) = 0.$$

However, this implies that

$$\sum_{r=1}^{r_a} \phi_{+,r}(t) \chi_{+,r}(\iota) = 0$$

for all t. Since the functions  $\phi_{+,r}$  are linearly independent, it follows that  $\chi_{+,r}(\iota) = 0$  for  $1 \le r \le r_a$ . Thus  $z_{+,r,app}(\iota,t) = 0$ , so that  $z_{+,r}(\iota,t) = 0$  for  $1 \le r \le r_a$ .

In case  $\kappa_{+,1} < \kappa_{-,1}$ , the argument is similar. In short, if the  $\kappa_{\pm,1}$ 's are different, we can eliminate the corresponding components of  $z_+$  or  $z_-$ . Proceeding step by step, we are then either finished or left with the situation that the real part of the eigenvalues equal. Without loss of generality, we can assume that the  $\kappa_{\pm,1}$ 's equal.

Equal maximal real parts of the eigenvalues. Assume that the  $\kappa_{\pm,1}$ 's equal. In fact, let us assume that  $\kappa_{\pm,1} = \cdots = \kappa_{\pm,r_{\pm}}$ , but that  $\kappa_{\pm,r_{\pm}+1} < \kappa_{\pm,1}$  (in case  $r_{\pm} < k_{\pm}$ ). Let us introduce the notation

$$\xi_{\pm}(\iota,t) := \sum_{r=1}^{r_{\pm}} \phi_{\pm,r}(t) z_{\pm,r,\mathrm{app}}(\iota,t),$$

where the  $z_{\pm,r,\text{app}}$  are defined by (19.36). Then, due to (19.33), (19.38) and an argument similar to the derivation of (19.39),

$$\lim_{t \to \infty} e^{-(\kappa_{+,1} + \beta_n/2)t} \left( |\dot{\xi}_{-}(\iota, t) + \dot{\xi}_{+}(\iota, t)| + \mathfrak{g}(\iota, t)|\xi_{-}(\iota, t) + \xi_{+}(\iota, t)| \right) = 0.$$
(19.41)

On the other hand,

$$\lim_{t \to \infty} e^{-(\kappa_{+,1} + \beta_n/2)t} \left( |\dot{\xi}_+(\iota, t) + i\mathfrak{g}(\iota, t)\xi_+(\iota, t)| + |\dot{\xi}_-(\iota, t) - i\mathfrak{g}(\iota, t)\xi_-(\iota, t)| \right) = 0.$$
(19.42)

Combining (19.41) and (19.42) yields

$$\lim_{t \to \infty} e^{-(\kappa_{+,1} + \beta_n/2)t} \mathfrak{g}(\iota, t) \left( |\xi_-(\iota, t) + \xi_+(\iota, t)| + |\xi_-(\iota, t) - \xi_+(\iota, t)| \right) = 0.$$

As a consequence, the arguments described to deal with the case of different  $\kappa_{\pm,1}$ 's apply. Thus  $z_{\pm,r}(\iota,t) = 0$  for all  $t \in I$  and  $r = 1, \ldots, r_{\pm}$ . Proceeding step by step, it follows that  $z_{\pm,r}(\iota,t) = 0$  for all  $t \in I$  and  $r = 1, \ldots, k_{\pm}$ . Applying the argument for all  $\iota \in \mathcal{I}_{B,n}$  yields  $u_{+} = u_{-} = 0$ , and the lemma follows.

#### **19.4** Detailed asymptotics in the noisy setting

Next, we turn to the problem of expressing the asymptotics of solutions to (1.2) in terms of solutions to (19.1). Let us begin by introducing the following terminology.

**Definition 19.10.** Assume that (1.2) is strongly balanced, has a negligible shift vector field, a dominant noisy spatial direction, and is such that the dominant coefficients are convergent. In case (1.2) has a dominant noisy spatial torus direction, say j, let

$$\mathcal{I}_{B,n}^{+} := \{ \iota \in \mathcal{I}_{B,n} : \nu_{T,j}(\iota) < 0 \}, \quad \mathcal{I}_{B,n}^{-} := \{ \iota \in \mathcal{I}_{B,n} : \nu_{T,j}(\iota) > 0 \};$$
(19.43)

cf. (24.5). In case (1.2) has a dominant noisy spatial generalised direction, let

$$\mathcal{I}_{B,n}^+ := \mathcal{I}_{B,n}, \quad \mathcal{I}_{B,n}^- := \emptyset.$$
(19.44)

**Lemma 19.11.** Assume that (1.2) is strongly balanced, has a negligible shift vector field, a dominant noisy spatial direction, and is such that the dominant coefficients are convergent. Let  $\eta_A$  be defined by (17.25) and assume that there is an  $0 < \eta_B \leq \eta_A$  such that

$$\int_{0}^{\infty} e^{-(\kappa_{n,+} - \eta_B)t} \|f_n(\cdot, t)\|_{(s)} dt < \infty$$
(19.45)

for all  $s \in \mathbb{R}$ , where  $\kappa_{n,+}$  is defined by (17.6);  $f_n$  is defined by (17.54); and  $\|\cdot\|_{(s)}$  is defined by (24.10). Let

$$A_{n,\pm} := -\frac{1}{2} \left( \alpha_{\infty} \pm \tilde{X}_{\infty,n} \right), \quad \kappa_{n,m} := \max\{\kappa_{\max}(A_{n,+}), \kappa_{\max}(A_{n,-})\},$$

where  $X_{\infty,n}$  is introduced in Definition 17.7. Finally, let  $J := (\kappa_{n,m} - \eta_B, \kappa_{n,m}]$  and  $E_{n,\pm} := E_{A_{n,\pm},J}$ ; cf. Definition 4.7.

If (1.2) has a dominant noisy spatial generalised direction, cf. Definition 17.1, then  $E_{n,+} = E_{n,-} =: E_n$  and  $A_{n,+} = A_{n,-} =: A_n$ . Moreover, if u is a solution to (1.2), there is a unique solution  $u_{0,w} \in C^{\infty}(M, E_n)$  to (19.1) such that

- if  $z_{0,w}(\iota,t)$  denotes the  $\iota$ 'th Fourier coefficient of  $u_{0,w}(\cdot,t)$ , then  $z_{0,w}(\iota,t) = 0$  for all  $t \in I$ and  $\iota \notin \mathcal{I}_{B,n}$ ,
- if  $u_w$  is defined by

$$u_{\rm w}(p,t) = e^{A_{\rm n}t} u_{0,\rm w}(p,t), \tag{19.46}$$

then there are real constants C, N and s such that

$$\mathfrak{E}_s^{1/2}[u_{\mathbf{n}} - u_{\mathbf{w}}](t) \le C\langle t \rangle^N e^{(\kappa_{\mathbf{n},+} - \eta_B)t}$$
(19.47)

for all  $t \geq 0$ , where  $u_n$  is defined in analogy with (17.54).

If (1.2) has a dominant noisy spatial torus direction, say j, and u is a solution to (1.2), then there are unique solutions  $u_{\pm,w} \in C^{\infty}(M, E_{n,\pm})$  to (19.1) such that the following holds:

- $u_{+,w} = u_{+,w}^+ + u_{-,w}^-$  and  $u_{-,w} = u_{+,w}^- + u_{-,w}^+$ , where  $u_{\tau,w}^v$ ,  $\tau, v \in \{+, -\}$ , are smooth solutions to (19.1) whose *i*'th Fourier coefficients are denoted by  $z_{\tau,w}^v(\iota, t)$  and are such that
  - $z_{\pm,w}^+(\iota, \cdot) = 0$  in case  $\iota \notin \mathcal{I}_{B,n}^+$  and  $z_{\pm,w}^-(\iota, \cdot) = 0$  in case  $\iota \notin \mathcal{I}_{B,n}^-$ ,
  - $-u_{+,w}^{\pm}$  are positively oriented and  $u_{-,w}^{\pm}$  are negatively oriented solutions to (19.1),
- if  $u_w$  is defined by

$$u_{\mathbf{w}}(p,t) = e^{A_{\mathbf{n},+}t}u_{+,\mathbf{w}}(p,t) + e^{A_{\mathbf{n},-}t}u_{-,\mathbf{w}}(p,t),$$
(19.48)

then there are real constants C, N and s such that

$$\mathfrak{E}_{s}^{1/2}[u_{\mathrm{n}}-u_{\mathrm{w}}](t) \leq C\langle t \rangle^{N} e^{(\kappa_{\mathrm{n},+}-\eta_{B})t}$$
(19.49)

for all  $t \geq 0$ .

Moreover, in both cases there are constants  $C_B > 0$  and  $s_B \ge 0$  (depending only on  $\eta_B$  and the coefficients of the equation (1.2)); and a non-negative integer N (depending only on m) such that the following holds. If u and  $u_w$  are as above, then

$$\mathfrak{E}_{s}^{1/2}[u_{n} - u_{w}](t) \leq C_{B} \langle t \rangle^{N} e^{(\kappa_{n,+} - \eta_{B})t} \left( \mathfrak{E}_{s+s_{B}}^{1/2}[u_{n}](0) + \int_{0}^{\infty} e^{-(\kappa_{n,+} - \eta_{B})t'} \|f_{n}(\cdot,t')\|_{(s+s_{B})} dt' \right)$$
(19.50)

for all  $t \ge 0$  and all  $s \in \mathbb{R}$ . In addition, if (1.2) has a dominant noisy spatial generalised direction, then

$$\mathfrak{E}_{s}^{1/2}[u_{0,\mathbf{w}}](0) \le C_{B}\left(\mathfrak{E}_{s+s_{B}}^{1/2}[u_{n}](0) + \int_{0}^{\infty} e^{-(\kappa_{n,+}-\eta_{B})t'} \|f_{n}(\cdot,t')\|_{(s+s_{B})} dt'\right)$$
(19.51)

for all  $s \in \mathbb{R}$ , where  $C_B$  and  $s_B$  have the same dependence as in the case of (19.50). Finally, if (1.2) has a dominant noisy spatial torus direction, then

$$\mathfrak{E}_{s}^{1/2}[u_{\pm,\mathbf{w}}](0) \le C_{B}\left(\mathfrak{E}_{s+s_{B}}^{1/2}[u_{\mathbf{n}}](0) + \int_{0}^{\infty} e^{-(\kappa_{\mathbf{n},+}-\eta_{B})t'} \|f_{\mathbf{n}}(\cdot,t')\|_{(s+s_{B})} dt'\right)$$
(19.52)

for all  $s \in \mathbb{R}$ , where  $C_B$  and  $s_B$  have the same dependence as in the case of (19.50).

**Remark 19.12.** Assume that (1.2) is  $C^2$ -balanced with a geometric dominant noisy spatial direction, convergent dominant coefficients and a negligible shift vector field; cf. Definition 6.5. Assume, moreover, that (19.45) holds for all  $s \in \mathbb{R}$ . Then the assumptions of the lemma are satisfied; cf. Remarks 17.2 and 17.6. How to calculate  $\eta_A$  etc. is clarified in Remark 19.2.

**Remark 19.13.** Note that  $\kappa_{n,+}$  and  $\kappa_{n,m}$  are related according to  $\kappa_{n,+} = \kappa_{n,m} + \beta_n/2$ .

**Remarks 19.14.** In the statement of the lemma, two equations play an important role: (1.2) and (19.1). Note that it is here taken for granted that (19.1) is the equation obtained from (1.2) by setting  $\alpha$ ,  $\zeta$ ,  $X^j$  and f to zero. In the statement of the present lemma, we speak of  $\eta_A$ , defined by (17.25). Note that this  $\eta_A$  need not necessarily coincide with the  $\eta_A$  associated with the equation (19.1), say  $\eta_{A,\text{hom}}$ , since  $\eta_{\text{mn}}$  can be omitted from the right hand side of (17.25) in the definition of  $\eta_{A,\text{hom}}$ . On the other hand, it is clear that  $\eta_{A,\text{hom}} \geq \eta_A$ , so that when we appeal to Lemma 19.1, the discrepancy does not cause complications. Turning to the constant  $\beta_n$ , note that it is the same for the two equations; cf. Definition 17.1. Finally, given  $\iota \in \mathcal{I}_{B,n}$ , the  $T_n$  given by Definition 17.12 need not be the same for the two equations (1.2) and (19.1); we obtain, say,  $T_n$  and  $T_{n,\text{hom}}$  respectively. However,  $\eta_{\text{mar}}$  appearing in Definition 17.1 and  $K_n$  appearing in the statement of Lemma 17.11 can be assumed to be the same for the two equations. Moreover, by deteriorating the value of the constant  $\mathfrak{b}$  appearing in (14.1) and (14.6) for one of the equations, it can be assumed that  $T_n = T_{n,\text{hom}}$ .

**Remark 19.15.** If (1.2) has a dominant noisy spatial generalised direction, then  $A_n = -\alpha_{\infty}/2$ , so that (19.46) can be written

$$u_{\rm w}(p,t) = e^{-\alpha_{\infty}t/2} u_{0,{\rm w}}(p,t).$$

**Remark 19.16.** The constants C, s and N appearing in (19.47) and (19.49) are allowed to depend on the coefficients of the equation (1.2),  $\eta_B$ , the functions  $u_n$ ,  $u_w$ ,  $f_n$  etc.

**Remark 19.17.** Dependence on the coefficients includes dependence on the Riemannian manifolds  $(M_r, g_r), r = 1, \ldots, R$ ; cf. Remark 17.10.

**Remark 19.18.** If all the Jordan blocks of the matrices  $A_{n,\pm}$  are trivial, then the N appearing in (19.50) can be replaced by 1. If  $E_{n,+} = E_{n,-} = \mathbb{C}^m$ , then N can be replaced by  $d_{n,+} - 1$ , where  $d_{n,+}$  is defined as in the statement of Lemma 17.18.

**Remark 19.19.** Due to the proof, the solutions  $u_{\pm,w}$  and  $u_{0,w}$  to (19.1) can be replaced by functions of the form described in the statement of Lemma 19.1. To be more precise, let us first consider the case that (1.2) has a dominant noisy spatial generalised direction. If u is a solution to (1.2), there are then uniquely determined  $\chi_{\pm}(\iota) \in E_n$ ,  $\iota \in \mathcal{I}_{B,n}$ , with the following properties: for every  $0 \leq k \in \mathbb{Z}$ , there is a constant  $C_k$  such that (19.2) holds; and if  $u_{0,wa}$  is given by the right hand side of (19.4), where  $z_{app}$  is given by (19.3), then  $u_{wa} = e^{A_n t} u_{0,wa}$  is such that there are constants C, N and s with the property that

$$\mathfrak{E}_s^{1/2}[u_{\mathrm{n}} - u_{\mathrm{wa}}](t) \le C \langle t \rangle^N e^{(\kappa_{\mathrm{n},+} - \eta_B)t}$$
(19.53)

for all  $t \geq 0$ .

Next, assume that (1.2) has a dominant noisy spatial torus direction. If u is a solution to (1.2), there are then uniquely determined  $\zeta_{\pm}^+(\iota)$  and  $\zeta_{\pm}^-(\iota)$ ,  $\iota \in \mathcal{I}_{\mathrm{B}}$ , with the following properties:  $\zeta_{\pm}^-(\iota) = 0$  for all  $\iota \notin \mathcal{I}_{\mathrm{B,n}}^-$  and  $\zeta_{\pm}^+(\iota) = 0$  for all  $\iota \notin \mathcal{I}_{\mathrm{B,n}}^+$ ;  $\zeta_{\pm}^+(\iota), \zeta_{-}^-(\iota) \in E_{\mathrm{n,+}}$  and  $\zeta_{\pm}^-(\iota), \zeta_{-}^+(\iota) \in E_{\mathrm{n,-}}$  for all  $\iota \in \mathcal{I}_{\mathrm{B}}$ ; for every  $0 \leq k \in \mathbb{Z}$ , there is a constant  $C_k$  such that

$$\sum_{\iota \in \mathcal{I}_{\mathrm{B,n}}} \sum_{\pm} \langle \nu(\iota) \rangle^{2k} \left[ |\zeta_{\pm}^{\pm}(\iota)|^2 + |\zeta_{-}^{\pm}(\iota)|^2 \right] \le C_k;$$
(19.54)

and if  $u^{\pm}_{+,\mathrm{wa}}$  and  $u^{\pm}_{-,\mathrm{wa}}$  are the functions whose  $\iota$ 'th Fourier coefficients are given by

$$z_{+,\mathrm{wa}}^{\pm}(\iota,t) = \frac{e^{\beta_{\mathrm{n}}t/2}}{\mathfrak{g}(\iota,t)} e^{i[\omega_{\mathrm{sh}}(\iota,t) - \varphi_{\mathrm{tot}}(\iota,t)]} \zeta_{+}^{\pm}(\iota), \qquad (19.55)$$

$$z_{-,\mathrm{wa}}^{\pm}(\iota,t) = \frac{e^{\beta_{\mathrm{n}}t/2}}{\mathfrak{g}(\iota,t)} e^{i[\omega_{\mathrm{sh}}(\iota,t) + \varphi_{\mathrm{tot}}(\iota,t)]} \zeta_{-}^{\pm}(\iota)$$
(19.56)

for  $\iota \in \mathcal{I}_{B}$  (when  $\iota \notin \mathcal{I}_{B,n}$ , the right hand sides of (19.55) and (19.56) should be interpreted as equalling zero), if  $u_{\pm,wa}$  are given by

$$u_{+,\text{wa}} = u_{+,\text{wa}}^+ + u_{-,\text{wa}}^-, \quad u_{-,\text{wa}} = u_{+,\text{wa}}^- + u_{-,\text{wa}}^+,$$
 (19.57)

and if  $u_{\text{wa}} = e^{A_{n,+}t}u_{+,\text{wa}} + e^{A_{n,-}t}u_{-,\text{wa}}$ , then there are constants C, N and s with the property that (19.53) holds for all  $t \ge 0$ . Note that, by definition,  $u_{\pm,\text{wa}}$  takes its values in  $E_{n,\pm}$ .

In both cases there are constants  $C_B > 0$  and  $s_B \ge 0$  (depending only on  $\eta_B$  and the coefficients of the equation (1.2)) and a non-negative integer N (depending only on m) such that if u and  $u_{wa}$ are as above, then

$$\mathfrak{E}_{s}^{1/2}[u_{n} - u_{wa}](t) \leq C_{B} \langle t \rangle^{N} e^{(\kappa_{n,+} - \eta_{B})t} \left( \mathfrak{E}_{s+s_{B}}^{1/2}[u_{n}](0) + \int_{0}^{\infty} e^{-(\kappa_{n,+} - \eta_{B})t'} \|f_{n}(\cdot,t')\|_{(s+s_{B})} dt' \right)$$
(19.58)

for all  $t \ge 0$  and all  $s \in \mathbb{R}$ . In addition, if all the Jordan blocks of the matrices  $A_{n,\pm}$  are trivial, then the N appearing in (19.58) can be replaced by 1, and if  $E_{n,+} = E_{n,-} = \mathbb{C}^m$ , then N can be replaced by  $d_{n,+} - 1$ . Turning to the  $\zeta_{\pm}^+$  and  $\zeta_{\pm}^-$ , there are constants  $C_B > 0$ ,  $s_a$  and  $s_b$  (depending only on  $\eta_B$  and the coefficients of the equation (1.2)), such that

$$\left(\sum_{\iota \in \mathcal{I}_{B,n}} \sum_{\pm} \langle \nu(\iota) \rangle^{2s} \left[ |\zeta_{\pm}^{\pm}(\iota)|^2 + |\zeta_{-}^{\pm}(\iota)|^2 \right] \right)^{1/2} \\ \leq C_B \left( \mathfrak{E}_{s+s_a}^{1/2} [u_n](0) + \int_0^\infty e^{-(\kappa_{n,+} - \eta_B)t} \|f_n(\cdot,t)\|_{(s+s_b)} dt \right)$$
(19.59)

for all  $s \in \mathbb{R}$ . In the case of a dominant noisy spatial generalised direction, the same estimate holds, assuming we introduce the notation  $\zeta_{\pm}^{-}(\iota) = 0$ ,  $\zeta_{+}^{+}(\iota) = \chi_{+}(\iota)$  and  $\zeta_{-}^{+}(\iota) = \chi_{-}(\iota)$  for all  $\iota \in \mathcal{I}_{\mathrm{B}}$ ; note that with this notation,  $u_{0,\mathrm{wa}} := u_{+,\mathrm{wa}} + u_{-,\mathrm{wa}}$ .

Finally, the functions  $u_{0,w}$  and  $u_{0,wa}$  (appearing in the statement of the lemma and the present remark, respectively) are uniquely associated as described in the statement of Lemma 19.1. Similarly, the functions  $u_{+,w}^{\pm}$  and  $u_{+,wa}^{\pm}$  (as well as the functions  $u_{-,w}^{\pm}$  and  $u_{-,wa}^{\pm}$ ) are uniquely associated as described in the statement of Lemma 19.1.

*Proof.* In order to prove uniqueness, it is convenient to divide the analysis into two different cases.

Uniqueness, dominant noisy spatial generalised direction. Assume that there are two functions, say  $v_l$ , l = 1, 2, with the same properties as the function  $u_{0,w}$  appearing in the statement of the lemma. Letting  $v := v_1 - v_2$  and  $u_{\text{diff}} := e^{A_n t} v$ , we know that

$$\mathfrak{E}_{s}^{1/2}[u_{\text{diff}}](t) \le C\langle t \rangle^{N} e^{(\kappa_{\text{n,m}} + \beta_{\text{n}}/2 - \eta_{B})t}$$
(19.60)

for all  $t \ge 0$ . Since  $u_{\text{diff}}$  can be written

$$u_{\rm diff} = e^{A_{\rm n,+}t} v_+ + e^{A_{\rm n,-}t} v_-,$$

where the components of  $v_{\pm}$  are positively oriented and the components of  $v_{\pm}$  are negatively oriented; the *i*'th Fourier coefficient of  $v_{\pm}(\cdot, t)$  vanishes if  $\iota \notin \mathcal{I}_{B,n}$ ;  $v_{\pm}$  takes its values in  $E_{n,\pm}$ ; and since (19.60) holds, it is clear that Lemma 19.8 is applicable and yields the conclusion that v = 0. The desired uniqueness follows.

Uniqueness, dominant noisy spatial torus direction. Assume now that there are two pairs of functions with the same properties as  $u_{\pm,w}$ , say  $v_{\pm,l}$ , l = 1, 2. Define  $v_{\pm} := v_{\pm,1} - v_{\pm,2}$  and

$$u_{\text{diff}} := e^{A_{\text{n},+}t}v_{+} + e^{A_{\text{n},-}t}v_{-}$$

Define  $u_{\text{diff},\pm}$  to be the projection of  $u_{\text{diff}}$  defined by setting the *i*'th Fourier coefficient of  $u_{\text{diff}}$  to zero for every  $\iota \notin \mathcal{I}_{B,n}^{\pm}$ . We define  $v_{+,\pm}$  and  $v_{-,\pm}$  similarly. Then (19.49) implies that

$$\mathfrak{E}_{s}^{1/2}[u_{\mathrm{diff},\pm}](t) \leq C \langle t \rangle^{N} e^{(\kappa_{\mathrm{n,m}}+\beta_{\mathrm{n}}/2-\eta_{B})t}$$

for all  $t \ge 0$ . Moreover,  $v_{+,+}$  and  $v_{-,-}$  are positively oriented and  $v_{+,-}$  and  $v_{-,+}$  are negatively oriented. Combining these observations with the fact that  $v_{+,\pm}$  takes its values in  $E_{n,+}$ , the fact that  $v_{-,\pm}$  takes its values in  $E_{n,-}$  and Lemma 19.8 yields the conclusion that  $v_{\pm} = 0$ . The desired uniqueness statement thus holds.

**Existence: preliminary estimates for late times.** In order to demonstrate existence, let us return to Lemma 18.8, in particular (18.62). Note that

$$T_{\rm pre}\mathfrak{D}_{\rm tot}(\iota,t)T_{\rm pre}^{-1} = \begin{pmatrix} e^{-i\varphi_{\rm tot}(\iota,t)}\mathrm{Id}_m & 0\\ 0 & e^{i\varphi_{\rm tot}(\iota,t)}\mathrm{Id}_m \end{pmatrix},\tag{19.61}$$

$$T_{\rm pre} e^{S_{\rm n}(\iota)t} T_{\rm pre}^{-1} = e^{\beta_{\rm n}t/2} \begin{pmatrix} e^{-[\alpha_{\infty} - \bar{X}_{\rm n}(\iota)]t/2} & 0\\ 0 & e^{-[\alpha_{\infty} + \bar{X}_{\rm n}(\iota)]t/2} \end{pmatrix},$$
(19.62)

where  $\bar{X}_n(\iota)$  is introduced in Definition 17.7 and  $T_{\text{pre}}$  is defined in (18.63). Introduce  $\psi_{\pm} \in \mathbb{C}^m$  according to

$$\psi_{\infty} = \begin{pmatrix} \psi_+ \\ \psi_- \end{pmatrix} := T_{\text{pre}} z_{\infty}.$$
(19.63)

Note that  $\psi_{\infty}$  is the vector whose existence is guaranteed in the statement of Lemma 18.1; in particular,  $\psi_{\infty}$  belongs to the first generalised eigenspace of the  $\eta_B$ ,  $R_n(\iota)$ -decomposition of  $\mathbb{C}^{2m}$ , where  $R_n(\iota)$  is defined by (17.23) and satisfies  $R_n(\iota) = T_{\text{pre}}S_n(\iota)T_{\text{pre}}^{-1}$ . In other words, if  $J_R :=$  $(\kappa_{n,+} - \eta_B, \kappa_{n,+}]$ , then  $\psi_{\infty}$  takes its values in the  $J_R$ -generalised eigenspace of  $R_n(\iota)$ . However, due to the form of the matrix  $R_n(\iota)$ , cf. (17.23), this is equivalent to the statement that  $\psi_+$ takes its values in the J-generalised eigenspace of  $[-\alpha_{\infty} + \bar{X}_n(\iota)]/2$  and  $\psi_-$  takes its values in the J-generalised eigenspace of  $[-\alpha_{\infty} - \bar{X}_n(\iota)]/2$ . Combining (19.61), (19.62) and (19.63) yields

$$\mathfrak{D}_{\rm tot}(\iota,t)e^{S_{\rm n}(\iota)t}z_{\infty} = e^{\beta_{\rm n}t/2}T_{\rm pre}^{-1} \left(\begin{array}{c} e^{-[\alpha_{\infty}-\bar{X}_{\rm n}(\iota)]t/2 - i\varphi_{\rm tot}(\iota,t)}\psi_{+} \\ e^{-[\alpha_{\infty}+\bar{X}_{\rm n}(\iota)]t/2 + i\varphi_{\rm tot}(\iota,t)}\psi_{-} \end{array}\right).$$

Introduce

$$\begin{aligned} z_{\mathrm{app}}(\iota,t) &:= \frac{e^{\beta_{\mathrm{n}}t/2}}{\mathfrak{g}(\iota,t)} e^{i\omega_{\mathrm{sh}}(\iota,t)} \left( e^{-[\alpha_{\infty} - \bar{X}_{\mathrm{n}}(\iota)]t/2 - i\varphi_{\mathrm{tot}}(\iota,t)} \psi_{+} \right. \\ &\left. -ie^{-[\alpha_{\infty} + \bar{X}_{\mathrm{n}}(\iota)]t/2 + i\varphi_{\mathrm{tot}}(\iota,t)} \psi_{-} \right), \\ p_{\mathrm{app}}(\iota,t) &:= e^{\beta_{\mathrm{n}}t/2 + i\omega_{\mathrm{sh}}(\iota,t)} \left( -ie^{-[\alpha_{\infty} - \bar{X}_{\mathrm{n}}(\iota)]t/2 - i\varphi_{\mathrm{tot}}(\iota,t)} \psi_{+} \right. \\ &\left. + e^{-[\alpha_{\infty} + \bar{X}_{\mathrm{n}}(\iota)]t/2 + i\varphi_{\mathrm{tot}}(\iota,t)} \psi_{-} \right). \end{aligned}$$

Then (18.62) implies that

$$\begin{aligned} |\dot{z}(\iota,t) - p_{\mathrm{app}}(\iota,t)| + \mathfrak{g}(\iota,t)|z(\iota,t) - z_{\mathrm{app}}(\iota,t)| \\ \leq C_B \langle \bar{t} \rangle^N e^{(\kappa_{\mathrm{n},+}-\eta_B)\bar{t}} \left( |w(\iota,T_{\mathrm{n}})| + e^{\kappa_{\mathrm{n},+}T_{\mathrm{n}}} \int_0^\infty e^{-(\kappa_{\mathrm{n},+}-\eta_B)t'} |\hat{f}(\iota,t')| dt' \right) \end{aligned}$$
(19.64)

holds for all  $t \ge T_n$ , where  $\bar{t} := t - T_n$ ;  $C_B$  only depends on  $\eta_B$  and the coefficients of the equation (1.2); and N is a non-negative integer depending only on m. Moreover, N = 1 in case the Jordan blocks of the matrices  $A_{n,\pm}$  are trivial. In addition, if  $E_{n,+} = E_{n,-} = \mathbb{C}^m$ , then N can be replaced by  $d_{n,+} - 1$ , where  $d_{n,+}$  is defined as in the statement of Lemma 17.18. The justification for the latter two statements is to be found in Remark 18.10; recall that  $S_n(\iota)$  and  $R_n(\iota)$  are related by a similarity transformation.

**Reformulating the preliminary estimate.** It would be convenient to replace  $p_{app}$  with  $\dot{z}_{app}$  in (19.64). In order to justify that this can be done, let

$$Z(t) := \frac{1}{\mathfrak{g}(t)} e^{i\omega_{\rm sh}(t)} \mathfrak{D}_{\rm tot}(t) e^{S_{\rm n}t} z_{\infty} = \frac{1}{\mathfrak{g}(t)} \begin{pmatrix} \mathfrak{g}(t) z_{\rm app}(t) \\ p_{\rm app}(t) \end{pmatrix},$$
(19.65)

where we have omitted the argument  $\iota$  for the sake of brevity. Due to (18.61),

$$|e^{S_{n}t}z_{\infty}| \le C_{B}\langle \bar{t} \rangle^{d_{n,+}-1} e^{\kappa_{n,+}\bar{t}} \left( |w(T_{n})| + e^{\kappa_{n,+}T_{n}} \int_{T_{n}}^{\infty} e^{-(\kappa_{n,+}-\eta_{B})t} |\hat{f}(t)| dt \right)$$
(19.66)

for all  $t \ge T_n$ , where  $C_B$  has the same dependence as in the case of (19.64). In particular,  $\mathfrak{g}|Z|$  is thus bounded by the right hand side of (19.66). Compute

$$\dot{Z} = -\dot{\ell}Z + i\sigma\mathfrak{g}Z + \frac{1}{\mathfrak{g}}e^{i\omega_{\rm sh}}\dot{\mathfrak{D}}_{\rm tot}e^{S_{\rm n}t}z_{\infty} + \frac{1}{\mathfrak{g}}e^{i\omega_{\rm sh}}\mathfrak{D}_{\rm tot}S_{\rm n}e^{S_{\rm n}t}z_{\infty}.$$
(19.67)

Combining (19.66); (19.67); (14.1); the fact that  $\mathfrak{g}(t) \geq c_0 e^{\beta_n t}$  for some  $c_0 > 0$  depending only on the coefficients of the equation (1.2) and all  $t \geq T_n$ ; and the fact that  $|\sigma(t)| \leq C e^{-\eta_{\text{sh}}t}$  for some constant C > 0 depending only on the coefficients of the equation (1.2), it follows that

$$\begin{aligned} & \left| \dot{Z}(t) - \dot{\mathfrak{D}}_{\text{tot}}(t) [\mathfrak{D}_{\text{tot}}(t)]^{-1} Z(t) \right| \\ \leq & C_B \langle \bar{t} \rangle^{d_{\text{n},+}-1} e^{(\kappa_{\text{n},+}-\eta_B)\bar{t}} \left( |w(T_{\text{n}})| + e^{\kappa_{\text{n},+}T_{\text{n}}} \int_{T_{\text{n}}}^{\infty} e^{-(\kappa_{\text{n},+}-\eta_B)t} |\hat{f}(t)| dt \right) \end{aligned}$$

for all  $t \ge T_n$ , where  $C_B$  has the same dependence as in the case of (19.64). Since  $\dot{\varphi}_{tot}(t) = [1 + \sigma^2(t)]^{1/2}\mathfrak{g}(t)$  and  $|\sigma(t)| \le Ce^{-\eta_{sh}t}$ , this estimate implies that

$$\begin{aligned} & \left| \dot{Z}(t) - \begin{pmatrix} 0 & \mathrm{Id}_m \\ -\mathrm{Id}_m & 0 \end{pmatrix} \mathfrak{g}(t) Z(t) \right| \\ \leq & C_B \langle \bar{t} \rangle^{d_{\mathrm{n},+}-1} e^{(\kappa_{\mathrm{n},+}-\eta_B)\bar{t}} \left( |w(T_{\mathrm{n}})| + e^{\kappa_{\mathrm{n},+}T_{\mathrm{n}}} \int_{T_{\mathrm{n}}}^{\infty} e^{-(\kappa_{\mathrm{n},+}-\eta_B)t} |\hat{f}(t)| dt \right) \end{aligned}$$

for all  $t \ge T_n$ , where  $C_B$  has the same dependence as in the case of (19.64). In particular, this estimate implies that (19.64) with  $p_{app}$  replaced by  $\dot{z}_{app}$  holds.

**Eliminating**  $|w(T_n)|$ . In order to proceed, it is of interest to estimate the right hand side of (19.64). Appealing to Lemma 10.7,

$$\mathcal{E}^{1/2}(t) \le e^{\eta_{\text{bal}}t} \mathcal{E}^{1/2}(0) + \int_0^t e^{\eta_{\text{bal}}(t-t')} |\hat{f}(t')| dt'$$
(19.68)

for all  $t \ge 0$ . Note that  $\mathcal{E}^{1/2}$  and |w| are equivalent for all  $t \ge T_n$ , the constant of equivalence being numerical; this is due to (17.13). Combining (19.68) (with  $t = T_n$ ) with (19.64) (with  $p_{app}$  replaced by  $\dot{z}_{app}$ ) yields

$$\begin{aligned} &|\dot{z}(t) - \dot{z}_{app}(t)| + \mathfrak{g}(t)|z(t) - z_{app}(t)| \\ \leq & C_B \langle t \rangle^N e^{(\kappa_{n,+} - \eta_B)t} \left( e^{\eta_{1,+}T_n} \mathcal{E}^{1/2}(0) + e^{\eta_{2,+}T_n} \int_0^\infty e^{-(\kappa_{n,+} - \eta_B)t'} |\hat{f}(t')| dt' \right) \end{aligned}$$
(19.69)

for all  $t \ge T_n$ , where  $C_B$  and N have the same dependence as in the case of (19.64), and

$$\eta_{1,+} := \max\{\eta_{\text{bal}} - \kappa_{n,+} + \eta_B, 0\}, \quad \eta_{2,+} := \max\{\eta_{\text{bal}} - \kappa_{n,+} + \eta_B, \eta_B\}.$$

If the Jordan blocks of the matrices  $A_{n,\pm}$  are trivial, the N appearing in (19.69) can be replaced by 1. Moreover, if  $E_{n,+} = E_{n,-} = \mathbb{C}^m$ , then N can be replaced by  $d_{n,+} - 1$ .

**Deriving an estimate for early times.** Next we need to verify that an estimate such as (19.69) holds on  $[0, T_n]$ . To this end, we need to estimate z and  $z_{app}$  separately. Due to (19.68),

$$\mathcal{E}^{1/2}(t) \le C e^{(\kappa_{n,+} - \eta_B)t} \left( e^{\eta_{1,+} T_n} \mathcal{E}^{1/2}(0) + e^{\eta_{1,+} T_n} \int_0^t e^{-(\kappa_{n,+} - \eta_B)t'} |\hat{f}(t')| dt' \right)$$
(19.70)

for all  $t \in [0, T_n]$ , where C is numerical. In order to estimate  $\dot{z}_{app}$  and  $\mathfrak{g}_{zapp}$  for  $t \in [0, T_n]$ , note that

$$\begin{aligned} \mathfrak{g}(t)|Z(t)| &\leq \|e^{S_{n}(t-T_{n})}\| \cdot |e^{S_{n}T_{n}}z_{\infty}| \\ &\leq C_{B}e^{\eta_{s}T_{n}} \left(|w(T_{n})| + e^{\kappa_{n,+}T_{n}} \int_{T_{n}}^{\infty} e^{-(\kappa_{n,+}-\eta_{B})t'} |\hat{f}(t')| dt'\right) \\ &\leq C_{B}e^{(\kappa_{n,+}-\eta_{B})t} \left(e^{\eta_{3,+}T_{n}}\mathcal{E}^{1/2}(0) + e^{\eta_{4,+}T_{n}} \int_{0}^{\infty} e^{-(\kappa_{n,+}-\eta_{B})t'} |\hat{f}(t')| dt'\right) \end{aligned}$$
(19.71)

for all  $t \in [0, T_n]$ , where  $C_B$  has the same dependence as in the case of (19.64);  $\eta_s \ge 0$  only depends on the coefficients of the equation (1.2);

$$\begin{split} \eta_{d,\pm} &:= \max\{0, \pm (\kappa_{n,+} - \eta_B)\}, \\ \eta_{3,+} &:= \eta_s + \eta_{bal} + \eta_{d,-}, \\ \eta_{4,+} &:= \max\{\max\{\eta_s + \kappa_{n,+}, 0\} + \eta_{d,-}, \eta_s + \eta_{1,+} + \eta_{d,+}\}; \end{split}$$

and we appealed to (18.61), (19.65) and (19.68). In order to estimate  $\dot{Z}$ , let us return to (19.67). This equality can be rewritten

$$\dot{Z} = -\dot{\ell}Z + i\sigma \mathfrak{g}Z + \dot{\mathfrak{D}}_{tot}\mathfrak{D}_{tot}^{-1}Z + \mathfrak{D}_{tot}S_{n}\mathfrak{D}_{tot}^{-1}Z.$$

Since  $|\dot{\ell}|$ ,  $|\sigma|$  and  $\mathfrak{g}^{-1} \| \dot{\mathfrak{D}}_{tot} \|$  are bounded for all  $t \geq 0$  by constants depending only on the coefficients of the equation (1.2), this equality yields

$$|\dot{Z}| \le C(1 + \mathfrak{g}^{-1})\mathfrak{g}|Z| \tag{19.72}$$

for all  $t \ge 0$ , where *C* only depends on the coefficients of the equation (1.2) and we used the fact that  $\mathfrak{D}_{tot}$  is unitary. We would like to combine this estimate with (19.71) in order to estimate  $|\dot{Z}|$ . To be allowed to do so, we need to estimate  $\mathfrak{g}$  from below. We already derived such an estimate in the proof of Lemma 19.1; cf. (19.27). The derivation is equally valid in the present setting, and we conclude that (19.27) holds for all  $t \ge 0$ , where *C* and  $s_E$  only depend on the coefficients of the equation (1.2). Combining (19.27), (19.71) and (19.72) yields

$$\begin{aligned} &|\dot{Z}(\iota,t)| + \mathfrak{g}(\iota,t)|Z(\iota,t)| + |Z(\iota,t)| \\ \leq &C_B e^{(\kappa_{n,+}-\eta_B)t} \langle \nu(\iota) \rangle^{s_E} \left( e^{\eta_{3,+}T_n} \mathcal{E}^{1/2}(\iota,0) + e^{\eta_{4,+}T_n} \int_0^\infty e^{-(\kappa_{n,+}-\eta_B)t'} |\hat{f}(\iota,t')| dt' \right) \end{aligned}$$

for all  $t \in [0, T_n]$ , where  $C_B$  has the same dependence as in the case of (19.64). Combining this estimate with (19.69) and (19.70) yields (recall that  $\mathfrak{g}(t) \geq 2$  for  $t \geq T_n$ )

$$\begin{aligned} &|\dot{z}(t) - \dot{z}_{app}(t)| + \mathfrak{g}(t)|z(t) - z_{app}(t)| + |z(t) - z_{app}(t)| \\ &\leq C_B \langle t \rangle^N e^{(\kappa_{n,+} - \eta_B)t} \langle \nu \rangle^{s_E} \left( e^{\eta_h T_n} \mathcal{E}^{1/2}(0) + e^{\eta_{ih} T_n} \int_0^\infty e^{-(\kappa_{n,+} - \eta_B)t'} |\hat{f}(t')| dt' \right) \\ &\leq C_B \langle t \rangle^N e^{(\kappa_{n,+} - \eta_B)t} \left( \langle \nu \rangle^{s_h} \mathcal{E}^{1/2}(0) + \langle \nu \rangle^{s_{ih}} \int_0^\infty e^{-(\kappa_{n,+} - \eta_B)t'} |\hat{f}(t')| dt' \right) \end{aligned}$$
(19.73)

for all  $t \ge 0$ , where  $C_B$  and N have the same dependence as in the case of (19.64) and we appealed to (19.23). If the Jordan blocks of the matrices  $A_{n,\pm}$  are trivial, the N appearing in (19.73) can be replaced by 1. Moreover, if  $E_{n,+} = E_{n,-} = \mathbb{C}^m$ , then N can be replaced by  $d_{n,+} - 1$ . Finally,

$$\begin{split} \eta_{\rm h} &:= \max\{\eta_{1,+}, \eta_{3,+}\}, \quad \eta_{\rm ih} := \max\{\eta_{2,+}, \eta_{4,+}\}, \\ s_{\rm h} &:= \frac{2\eta_{\rm h}}{\eta_{\rm mar}} + s_E, \quad s_{\rm ih} := \frac{2\eta_{\rm ih}}{\eta_{\rm mar}} + s_E. \end{split}$$

Summing up; the approximate solution. Let

$$u_{\mathbf{n}}(p,t) := \sum_{\iota \in \mathcal{I}_{\mathbf{B},\mathbf{n}}} z(\iota,t)\varphi_{\iota}(p), \quad u_{\mathbf{app}}(p,t) := \sum_{\iota \in \mathcal{I}_{\mathbf{B},\mathbf{n}}} z_{\mathbf{app}}(\iota,t)\varphi_{\iota}(p).$$
(19.74)

Then (19.73) yields

$$\mathfrak{E}_{s}^{1/2}[u_{n} - u_{app}](t) \\
\leq C_{B} \langle t \rangle^{N} e^{(\kappa_{n,+} - \eta_{B})t} \left( \mathfrak{E}_{s+s_{h}}^{1/2}[u_{n}](0) + \int_{0}^{\infty} e^{-(\kappa_{n,+} - \eta_{B})t'} \|f_{n}(\cdot,t')\|_{(s+s_{ih})} dt' \right)$$
(19.75)

for all  $s \in \mathbb{R}$  and all  $t \ge 0$ , where  $C_B$  and N have the same dependence as in the case of (19.64). Note also that  $s_h$  and  $s_{ih}$  only depend on  $\eta_B$  and the coefficients of the equation (1.2). Moreover, if the Jordan blocks of the matrices  $A_{n,\pm}$  are trivial, the N appearing in (19.75) can be replaced by 1. Finally, if  $E_{n,+} = E_{n,-} = \mathbb{C}^m$ , then N can be replaced by  $d_{n,+} - 1$ .

Reformulating and estimating the approximate solution. Next, we wish to rewrite  $u_{app}$ . Introduce, to this end,

$$\begin{aligned} z_{+,\mathrm{wa}}(\iota,t) &:= \frac{e^{\beta_{\mathrm{n}}t/2}}{\mathfrak{g}(\iota,t)} e^{i[\omega_{\mathrm{sh}}(\iota,t) - \varphi_{\mathrm{tot}}(\iota,t)]} \psi_{+}(\iota), \\ z_{-,\mathrm{wa}}(\iota,t) &:= -\frac{e^{\beta_{\mathrm{n}}t/2}}{\mathfrak{g}(\iota,t)} e^{i[\omega_{\mathrm{sh}}(\iota,t) + \varphi_{\mathrm{tot}}(\iota,t)]} i\psi_{-}(\iota) \end{aligned}$$

Then  $z_{app}(\iota, t)$  can be written

$$z_{\rm app}(\iota, t) = e^{-[\alpha_{\infty} - \bar{X}_{\rm n}(\iota)]t/2} z_{+,\rm wa}(\iota, t) + e^{-[\alpha_{\infty} + \bar{X}_{\rm n}(\iota)]t/2} z_{-,\rm wa}(\iota, t).$$
(19.76)

Let us estimate the  $\psi_{\pm}$ . Keeping (19.63) in mind, it is sufficient to estimate  $z_{\infty}$ . Moreover,  $z_{\infty}$  can be estimated by appealing to (18.61):

$$\begin{aligned} |\psi_{+}| + |\psi_{-}| &\leq C|z_{\infty}| \leq C \|e^{-S_{n}T_{n}}\| \cdot |e^{S_{n}T_{n}}z_{\infty}| \\ &\leq C_{B} \|e^{-S_{n}(\iota)T_{n}}\| \left( |w(T_{n})| + e^{\kappa_{n,+}T_{n}} \int_{T_{n}}^{\infty} e^{-(\kappa_{n,+}-\eta_{B})t'} |\hat{f}(t')| dt' \right) \\ &\leq C_{B} e^{\eta_{s}T_{n}} \left( e^{\eta_{\text{bal}}T_{n}} \mathcal{E}^{1/2}(0) + e^{\eta_{a}T_{n}} \int_{0}^{T_{n}} e^{-(\kappa_{n,+}-\eta_{B})t'} |\hat{f}(t')| dt' + e^{\kappa_{n,+}T_{n}} \int_{T_{n}}^{\infty} e^{-(\kappa_{n,+}-\eta_{B})t'} |\hat{f}(t')| dt' \right), \end{aligned}$$

where we appealed to (19.68);  $C_B$  has the same dependence as in the case of (19.64);  $\eta_s \ge 0$  only depends on the coefficients of the equation (1.2); and

$$\eta_a := \eta_{\text{bal}} + \max\{0, \kappa_{n,+} - \eta_{\text{bal}} - \eta_B\}.$$

Due to (19.23), this estimate yields

$$|\psi_{+}(\iota)| + |\psi_{-}(\iota)| \le C_B \left( \mathcal{E}_{s_a}^{1/2}(\iota, 0) + \langle \nu(\iota) \rangle^{s_b} \int_0^\infty e^{-(\kappa_{n,+} - \eta_B)t'} |\hat{f}(\iota, t')| dt' \right),$$
(19.77)

where  $C_B$  has the same dependence as in the case of (19.64) and

$$s_a := \frac{2}{\eta_{\text{mar}}} (\eta_{\text{s}} + \eta_{\text{bal}}), \quad s_b := \frac{2}{\eta_{\text{mar}}} (\eta_{\text{s}} + \max\{\eta_a, \kappa_{n,+}\});$$

note that  $s_a$  and  $s_b$  only depend on  $\eta_B$  and the coefficients of the equation (1.2). Thus

$$\left(\sum_{\iota \in \mathcal{I}_{B,n}} \langle \nu(\iota) \rangle^{2s} \left[ |\psi_{+}(\iota)|^{2} + |\psi_{-}(\iota)|^{2} \right] \right)^{1/2} \\ \leq C_{B} \left( \mathfrak{E}_{s+s_{a}}^{1/2} [u_{n}](0) + \int_{0}^{\infty} e^{-(\kappa_{n,+} - \eta_{B})t'} \|f_{n}(\cdot,t')\|_{(s+s_{b})} dt' \right)$$
(19.78)

for all  $s \in \mathbb{R}$ , where  $C_B$  has the same dependence as in the case of (19.64).

Due to the assumptions of the lemma, the right hand side of (19.78) is bounded for all s. In particular, the components of  $\psi_{\pm}(\iota)$  satisfy estimates such that Lemma 19.1 can be applied with  $z_{\rm app}$  replaced with the components of  $z_{\pm,\rm wa}$ . However, there is one problem associated with doing so: in (19.76), there are  $\iota$ -dependent factors multiplying  $z_{\pm,\rm wa}$ . On the other hand, this problem only occurs if the equation has a dominant noisy spatial torus direction and is related to the sign of  $\nu_{\rm T,j}(\iota)$ . It is therefore convenient to use the notation introduced in Definition 19.10; recall also that  $\bar{X}_{\rm n}(\iota) = 0$  for all  $\iota \in \mathcal{I}_{\rm B,n}$  in case the equation has a dominant noisy spatial generalised direction. Define

$$u_{\pm,\text{wa}}^{+}(p,t) := \sum_{\iota \in \mathcal{I}_{B,n}^{+}} z_{\pm,\text{wa}}(\iota,t)\varphi_{\iota}(p), \quad u_{\pm,\text{wa}}^{-}(p,t) := \sum_{\iota \in \mathcal{I}_{B,n}^{-}} z_{\pm,\text{wa}}(\iota,t)\varphi_{\iota}(p).$$
(19.79)

Due to the definition of  $z_{\pm,\text{wa}}(\iota, t)$  and the properties of the  $\psi_{\pm}$ 's, it is clear that  $u^+_{+,\text{wa}}$  and  $u^-_{-,\text{wa}}$  take their values in  $E_{n,+}$  and that  $u^-_{+,\text{wa}}$  and  $u^+_{-,\text{wa}}$  take their values in  $E_{n,-}$ , where the  $E_{n,\pm}$  are defined in the statement of the lemma.

Comparing with Remark 19.19. It is of interest to compare the above conclusions and notation with Remark 19.19. Let us first consider the case of a dominant noisy spatial torus direction. Define  $\zeta_{+}^{\pm}(\iota) = \psi_{+}(\iota)$  for all  $\iota \in \mathcal{I}_{B,n}^{\pm}$  and  $\zeta_{+}^{\pm}(\iota) = 0$  for all  $\iota \notin \mathcal{I}_{B,n}^{\pm}$ . Define, similarly,  $\zeta_{-}^{\pm}(\iota) = -i\psi_{-}(\iota)$  for all  $\iota \in \mathcal{I}_{B,n}^{\pm}$  and  $\zeta_{-}^{\pm}(\iota) = 0$  for all  $\iota \notin \mathcal{I}_{B,n}^{\pm}$ . Define, similarly,  $\zeta_{-}^{\pm}(\iota) = -i\psi_{-}(\iota)$  for all  $\iota \in \mathcal{I}_{B,n}^{\pm}$  and  $\zeta_{-}^{\pm}(\iota) = 0$  for all  $\iota \notin \mathcal{I}_{B,n}^{\pm}$ . Then  $\zeta_{+}^{+}(\iota), \zeta_{-}^{-}(\iota) \in E_{n,+}$  and  $\zeta_{+}^{+}(\iota), \zeta_{-}^{+}(\iota) \in E_{n,-}$  for all  $\iota \in \mathcal{I}_{B}$ . Note also that the estimates (19.54) and (19.59) follow from (19.78). Defining  $z_{+,wa}^{\pm}$  and  $z_{-,wa}^{\pm}$  by (19.55) and (19.56) respectively, it is clear that  $z_{+,wa}^{\pm}$  and  $z_{-,wa}^{\pm}$  are the Fourier coefficients of  $u_{+,wa}^{\pm}$  and  $u_{-,wa}^{\pm}$ , introduced in (19.79), respectively. In short, the  $u_{+,wa}^{\pm}$  and  $u_{-,wa}^{\pm}$  introduced in (19.79) coincide with the  $u_{+,wa}^{\pm}$  and  $u_{-,wa}^{\pm}$  introduced in Remark 19.19. Defining  $u_{\pm,wa}$  by (19.57), the function  $u_{app}$ , introduced in (19.74), can be written

$$u_{\rm app}(p,t) = e^{A_{\rm n,+}t} u_{+,\rm wa}(p,t) + e^{A_{\rm n,-}t} u_{-,\rm wa}(p,t), \qquad (19.80)$$

where  $A_{n,\pm}$  is introduced in the statement of the lemma. In particular,  $u_{app}$  equals the function  $u_{wa}$  introduced in Remark 19.19. Due to (19.75), the estimates (19.53) and (19.58) thus hold. To conclude,  $\zeta^{\pm}_{+}$  and  $\zeta^{\pm}_{+}$  have the properties stated in Remark 19.19, except, possibly, for the uniqueness (which remains to be verified).

Next, let us turn to the case of a dominant noisy spatial generalised direction. Let  $\chi_{+}(\iota) = \psi_{+}(\iota)$ and  $\chi_{-}(\iota) = -i\psi_{-}(\iota)$  for all  $\iota \in \mathcal{I}_{B,n}$ , and let  $\chi_{\pm}(\iota) = 0$  for all  $\iota \notin \mathcal{I}_{B,n}$ . Then  $\chi_{\pm}(\iota) \in E_{n}$  for all  $\iota \in \mathcal{I}_{\rm B}$ . Moreover, (19.2) and (19.59) follow from (19.78) (assuming (19.59) is interpreted as stated immediately below this estimate). If  $u_{0,\rm wa}$  is given by the right hand side of (19.4), where  $z_{\rm app}$  is given by (19.3), then the function  $u_{\rm app}$ , introduced in (19.74), can be written  $u_{\rm app} = e^{A_{\rm n}t}u_{0,\rm wa}$ . In other words, the function  $u_{\rm wa}$ , introduced in Remark 19.19, equals  $u_{\rm app}$ . Due to (19.75), the estimates (19.53) and (19.58) thus hold. To conclude,  $\chi_{\pm}$  have the properties stated in Remark 19.19, except, possibly, for the uniqueness (which remains to be verified).

**Constructing solutions to (19.1).** At this stage, we can appeal to Lemma 19.1. This yields four solutions  $u_{\pm,w}^+$  and  $u_{\pm,w}^-$  to (19.1) in the case of a dominant noisy spatial torus direction. In what follows, we also use the notation

$$u_{+,\mathbf{w}} = u_{+,\mathbf{w}}^+ + u_{-,\mathbf{w}}^-, \quad u_{-,\mathbf{w}} = u_{+,\mathbf{w}}^- + u_{-,\mathbf{w}}^+.$$

In the case of a dominant noisy spatial generalised direction, the  $\chi_{\pm}$  give rise to functions  $u_{\pm,w}$ . In order to be able to discuss the torus setting and the generalised setting at the same time, it is convenient to introduce the notation  $u_{\pm,w}^- := 0$ ,  $u_{\pm,w}^+ := u_{\pm,w}$  and  $u_{0,w} := u_{+,w} + u_{-,w}$  in the generalised setting. Before proceeding, note that the following holds:

- if  $z_{0,w}(\iota, t)$  denotes the  $\iota$ 'th Fourier coefficient of  $u_{0,w}(\cdot, t)$ , then  $z_{0,w}(\iota, \cdot) = 0$  if  $\iota \notin \mathcal{I}_{B,n}$ ,
- $z_{0,w}$ , and thereby  $u_{0,w}$ , takes its values in  $E_n$ ,
- if the *i*'th Fourier coefficients of  $u_{\tau w}^{\upsilon}$ ,  $\tau, \upsilon \in \{+, -\}$ , are denoted by  $z_{\tau w}^{\upsilon}(\iota, \cdot)$ , then
  - $z_{\pm,w}^+(\iota, \cdot) = 0 \text{ in case } \iota \notin \mathcal{I}_{B,n}^+ \text{ and } z_{\pm,w}^-(\iota, \cdot) = 0 \text{ in case } \iota \notin \mathcal{I}_{B,n}^-,$
  - $u^{\pm}_{+,w}$  are positively oriented and  $u^{\pm}_{-,w}$  are negatively oriented solutions to (19.1),
- $u_{+,w}^+$  and  $u_{-,w}^-$  take their values in  $E_{n,+}$  and  $u_{-,w}^+$  and  $u_{-,w}^+$  take their values in  $E_{n,-}$ , so that  $u_{\pm,w}$  takes its values in  $E_{n,\pm}$ .

With the above notation, the conclusions of Lemma 19.1 yield

$$\mathfrak{E}_{s}[u_{\pm,\mathrm{w}} - u_{\pm,\mathrm{wa}}](t) \le C_{B} e^{(\beta_{\mathrm{n}} - 2\eta_{B})t} \sum_{\iota \in \mathcal{I}_{\mathrm{B},\mathrm{n}}} \langle \nu(\iota) \rangle^{2(s+s_{\mathrm{m}})} [r_{\pm}^{2}(\iota) + r_{-}^{2}(\iota)]$$
(19.81)

for all  $t \ge 0$ , where the constants have the dependence stated in Lemma 19.1 and  $r_{\pm}(\iota) = |\psi_{\pm}(\iota)|$ . Combining (19.78) with (19.81) yields

$$\mathfrak{E}_{s}^{1/2}[u_{\pm,\mathrm{w}} - u_{\pm,\mathrm{wa}}](t) \leq C_{B}e^{(\beta_{\mathrm{n}}/2 - \eta_{B})t} \left(\mathfrak{E}_{s+s_{1}}^{1/2}[u_{\mathrm{n}}](0) + \int_{0}^{\infty} e^{-(\kappa_{\mathrm{n},+} - \eta_{B})t'} \|f_{\mathrm{n}}(\cdot,t')\|_{(s+s_{2})} dt'\right)$$
(19.82)

for all  $s \in \mathbb{R}$  and all  $t \ge 0$ , where  $s_1 := s_a + s_m$ ,  $s_2 := s_b + s_m$ , and  $C_B$  has the same dependence as in the case of (19.64). In addition combining (19.7) and (19.78) yields (19.51) and (19.52).

The final estimate. Define  $u_w$  by (19.48) in the case of a dominant noisy spatial torus direction. Define  $u_w$  by (19.46) in the case of a dominant noisy spatial generalised direction. Combining these definitions with (19.80) and (19.82) yields the conclusion

$$\mathfrak{E}_{s}^{1/2}[u_{w} - u_{app}](t) \\
\leq C_{B}\langle t \rangle^{d_{n,+}-1} e^{(\kappa_{n,+}-\eta_{B})t} \left( \mathfrak{E}_{s+s_{1}}^{1/2}[u_{n}](0) + \int_{0}^{\infty} e^{-(\kappa_{n,+}-\eta_{B})t'} \|f_{n}(\cdot,t')\|_{(s+s_{2})} dt' \right)$$
(19.83)

for all  $s \in \mathbb{R}$  and all  $t \ge 0$ , where  $C_B$  has the same dependence as in the case of (19.64). Finally, combining this estimate with (19.75) yields (19.50). Comparing the statements of the lemma with the above constructions, it is clear that the lemma holds.

Uniqueness of the approximation. Finally, let us turn to the uniqueness statements contained in Remark 19.19. Starting with the case of a dominant noisy spatial generalised direction, assume that there are  $\chi_{\pm,l}$ , l = 1, 2, with the properties stated in the remark. This gives rise to two functions  $u_{0,\text{wa},l}$  with values in  $E_n$ . Appealing to Lemma 19.1, we also obtain functions  $u_{0,\text{w},l}$ , l = 1, 2, with values in  $E_n$ . Defining  $u_{\text{wa},l} := e^{A_n t} u_{0,\text{wa},l}$  and  $u_{\text{w},l} := e^{A_n t} u_{0,\text{w},l}$ , (19.53) (with  $u_{\text{wa}}$ replaced by  $u_{\text{wa},l}$ ) and Lemma 19.1 imply

$$\mathfrak{E}_s^{1/2}[u_{\mathbf{n}} - u_{\mathbf{w},l}](t) \le C \langle t \rangle^N e^{(\kappa_{\mathbf{n},+} - \eta_B)t}$$

for all  $t \ge 0$ . Due to the uniqueness part of the lemma, this estimate implies that  $u_{0,w,1} = u_{0,w,2}$ . By the uniqueness part of Lemma 19.1, we conclude that  $\chi_{\pm,1} = \chi_{\pm,2}$ . The proof of uniqueness in the torus setting is similar. The same is true of the final statement in Remark 19.19.

#### **19.5** Specifying data at infinity

If the assumptions of Lemma 19.11 are satisfied, we know that, given a smooth solution u to (1.2), there are uniquely associated solutions  $u_{\pm,w}$  (or  $u_{0,w}$ ) to (19.1) such that  $u_n$  is well approximated by  $u_w$  defined by (19.48) (or (19.46)); cf. (19.50). Even though this result provides some information concerning the asymptotics, it does leave several questions open. The first question is: given that  $u_{\pm,w}$  (or  $u_{0,w}$ ) have the properties stated in Lemma 19.11, except for the estimates concerning the difference between  $u_n$  and  $u_w$ , is there a solution u to (1.2) such that (19.50) holds? The estimates (19.51), (19.52) and (19.59) demonstrate that the map from solutions to (1.2) to the asymptotic data is continuous. Once we have constructed a map from asymptotic data to initial data, we here also wish to demonstrate that it is continuous. As in the case of Lemma 10.22 and Proposition 12.25, we focus our attention on the case of homogeneous equations. Before stating the lemma, it is convenient to introduce the following terminology.

**Definition 19.20.** Let  $1 \leq m \in \mathbb{Z}$  and V be a vector subspace of  $\mathbb{C}^m$ . Then

$$W_{\rm n}(\bar{M}, V) \subset C^{\infty}(\bar{M}, V) \times C^{\infty}(\bar{M}, V)$$

denotes the set of initial data at t = 0 for solutions u to (19.1) such that  $u \in C^{\infty}(M, V)$  and the  $\iota$ 'th Fourier coefficient of  $u(\cdot, t)$  vanishes for all t if  $\iota \notin \mathcal{I}_{B,n}$ . Moreover,  $W_{n,+}(\bar{M}, V)$  ( $W_{n,-}(\bar{M}, V)$ ) denotes the subset of  $W_n(\bar{M}, V)$  corresponding to positively (negatively) oriented solutions to (19.1). In addition,  $W_{n,\pm}^+(\bar{M}, V)$  ( $W_{n,\pm}^-(\bar{M}, V)$ ) denotes the subset of  $W_{n,\pm}(\bar{M}, V)$  consisting of functions whose  $\iota$ 'th Fourier coefficients vanish if  $\iota \notin \mathcal{I}_{B,n}^+$  ( $\iota \notin \mathcal{I}_{B,n}^-$ ).

**Remark 19.21.** By initial data we here mean  $[u(\cdot, 0), u_t(\cdot, 0)]$ .

**Lemma 19.22.** Assume that (1.2) is strongly balanced, has a negligible shift vector field, a dominant noisy spatial direction, and is such that the dominant coefficients are convergent. Assume, moreover, that f = 0. Let  $\eta_A$  be defined by (17.25) and  $0 < \eta_B \leq \eta_A$ . Define  $A_{n,\pm}$  and  $E_{n,\pm}$  as in the statement of Lemma 19.11.

If (1.2) has a dominant noisy spatial generalised direction, then  $E_{n,+} = E_{n,-} =: E_n$  and  $A_{n,+} = A_{n,-} =: A_n$ . Moreover, there is an injective linear map

$$\Phi_{n,g}: W_n(\bar{M}, E_n) \to W_n(\bar{M}, \mathbb{C}^m)$$

such that if  $\chi \in W_n(\overline{M}, E_n)$ ;  $u_{0,w}$  is the solution to (19.1) corresponding to the initial data  $\chi$  at t = 0;  $u_w$  is defined by (19.46); and u is the solution to (1.2) (with f = 0) corresponding to the initial data  $\Phi_{n,g}(\chi)$  at t = 0, then  $u = u_n$  and

$$\mathfrak{C}_{s}^{1/2}[u-u_{w}](t) \le C_{B}\langle t \rangle^{N} e^{(\kappa_{n,+}-\eta_{B})t} \|\chi\|_{(s+s_{B})}$$
(19.84)

for all  $t \ge 0$  and  $s \in \mathbb{R}$ , where  $C_B$ ,  $s_B$  and N have the same dependence as in the case of (19.50). If the Jordan blocks of the matrix  $A_n$  are trivial, the N appearing in (19.84) can be replaced by 1. Moreover, if  $E_n = \mathbb{C}^m$ , then N can be replaced by  $d_{n,+} - 1$ , where  $d_{n,+}$  is defined as in the statement of Lemma 17.18. If  $E_n = \mathbb{C}^m$ , then  $\Phi_{n,g}$  is surjective. Finally, there are constants  $C_B > 0$  and  $s_B \ge 0$  (depending only on  $\eta_B$  and the coefficients of the equation (1.2)) such that

$$\|\Phi_{\mathbf{n},\mathbf{g}}(\chi)\|_{(s)} \le C_B \|\chi\|_{(s+s_B)} \tag{19.85}$$

for all  $s \in \mathbb{R}$  and all  $\chi \in W_n(\overline{M}, E_n)$ .

If (1.2) has a dominant noisy spatial torus direction, let

$$\mathcal{W}_{n}^{+} := W_{n,+}^{+}(\bar{M}, E_{n,+}) \oplus W_{n,-}^{-}(\bar{M}, E_{n,+}),$$
  
$$\mathcal{W}_{n}^{-} := W_{n,+}^{-}(\bar{M}, E_{n,-}) \oplus W_{n,-}^{+}(\bar{M}, E_{n,-}).$$

Then there is an injective linear map

$$\Phi_{n,T}: \mathcal{W}_n^+ \times \mathcal{W}_n^- \to W_n(\bar{M}, \mathbb{C}^m)$$

such that if  $\chi_{\pm} \in \mathcal{W}_{n}^{\pm}$ ;  $u_{\pm,w}$  is the solution to (19.1) corresponding to the initial data  $\chi_{\pm}$  at t = 0;  $u_{w}$  is defined by (19.48); and u is the solution to (1.2) (with f = 0) corresponding to the initial data  $\Phi_{n,T}(\chi_{+}, \chi_{-})$  at t = 0, then  $u = u_{n}$  and

$$\mathfrak{E}_{s}^{1/2}[u-u_{w}](t) \leq C_{B}\langle t \rangle^{N} e^{(\kappa_{n,+}-\eta_{B})t}[\|\chi_{+}\|_{(s+s_{B})} + \|\chi_{-}\|_{(s+s_{B})}]$$
(19.86)

for all  $t \ge 0$  and  $s \in \mathbb{R}$ , where  $C_B$ ,  $s_B$  and N have the same dependence as in the case of (19.50). Moreover, N = 1 in case the Jordan blocks of the matrices  $A_{n,\pm}$  are trivial. In addition, if  $E_{n,+} = E_{n,-} = \mathbb{C}^m$ , then N can be replaced by  $d_{n,+} - 1$ , where  $d_{n,+}$  is defined as in the statement of Lemma 17.18. If  $E_{n,+} = E_{n,-} = \mathbb{C}^m$ , then  $\Phi_{n,T}$  is surjective. Finally, there are constants  $C_B > 0$  and  $s_B \ge 0$  (depending only on  $\eta_B$  and the coefficients of the equation (1.2)) such that

$$\|\Phi_{\mathbf{n},\mathbf{T}}(\chi_{+},\chi_{-})\|_{(s)} \le C_{B}[\|\chi_{+}\|_{(s+s_{B})} + \|\chi_{-}\|_{(s+s_{B})}]$$
(19.87)

for all  $s \in \mathbb{R}$  and all  $(\chi_+, \chi_-) \in \mathcal{W}_n^+ \times \mathcal{W}_n^-$ .

**Remark 19.23.** Dependence on the coefficients includes dependence on the Riemannian manifolds  $(M_r, g_r), r = 1, \ldots, R$ ; cf. Remark 17.10.

**Remark 19.24.** Assume that (1.2) is  $C^2$ -balanced with a geometric dominant noisy spatial direction, convergent dominant coefficients and a negligible shift vector field; cf. Definition 6.5. Assume, moreover, that f = 0. Then the assumptions of the lemma are satisfied; cf. Remarks 17.2 and 17.6. How to calculate  $\eta_A$  etc. is clarified in Remark 19.2.

Remark 19.25. Remarks 19.14 are equally relevant in the present setting.

**Remark 19.26.** It is also possible to specify functions  $u_{\pm,\text{wa}}$  and  $u_{0,\text{wa}}$  as in Remark 19.19, and then to construct a solution u to (1.2) (with f = 0) such that  $u = u_n$  and such that

$$\mathfrak{E}_{s}^{1/2}[u-u_{\mathrm{wa}}](t) \leq C_{B}\langle t \rangle^{N} e^{(\kappa_{\mathrm{n},+}-\eta_{B})t} \mathfrak{E}_{s+s_{B}}^{1/2}[u](0)$$
(19.88)

for all  $t \ge 0$  and all  $s \in \mathbb{R}$ , where the constants  $C_B$ ,  $s_B$  and N have the same dependence as in the case of (19.50);

$$u_{\mathrm{wa}}(p,t) := e^{A_{\mathrm{n}}t} u_{0,\mathrm{wa}}(p,t)$$

in case (1.2) has a dominant noisy spatial generalised direction; and

$$u_{\rm wa}(p,t) := e^{A_{\rm n,+}t} u_{+,\rm wa}(p,t) + e^{A_{\rm n,-}t} u_{-,\rm wa}(p,t)$$
(19.89)

in case (1.2) has a dominant noisy spatial torus direction. Moreover, if the  $u_{\pm,\text{wa}}$  and  $u_{0,\text{wa}}$  are defined in terms of the  $\zeta_{\pm}^+$  and  $\zeta_{\pm}^-$  as described in Remark 19.19, then

$$\mathfrak{E}_{s}^{1/2}[u](0) \le C_{B} \left( \sum_{\iota \in \mathcal{I}_{\mathrm{B,n}}} \sum_{\pm} \langle \nu(\iota) \rangle^{2(s+s_{B})} [|\zeta_{\pm}^{+}(\iota)|^{2} + |\zeta_{\pm}^{-}(\iota)|^{2}] \right)^{1/2}$$
(19.90)

for all  $s \in \mathbb{R}$ , where  $C_B$  and  $s_B$  have the same dependence as in the case of (19.50). Finally, note that by combining (19.88) and (19.90), the last factor on the right hand side of (19.88) can be replaced by the last factor on the right hand side of (19.90).

*Proof.* Note that when (1.2) has a dominant noisy spatial generalised direction, then

$$\mathcal{W}_{n}^{+} = W_{n,+}^{+}(\bar{M}, E_{n,+}), \quad \mathcal{W}_{n}^{-} = W_{n,-}^{+}(\bar{M}, E_{n,-}),$$

so that  $\mathcal{W}_n^+ \times \mathcal{W}_n^- \cong \mathcal{W}_n(\overline{M}, E_n)$  in that case. For that reason, we below assume that (1.2) has a dominant noisy spatial torus direction and leave it to the reader to verify (in the course of the argument), that the case of a dominant noisy spatial generalised direction can be considered to be a special case of this.

**Constructing**  $u_{\pm,w}$  from  $\chi_{\pm}$ . Let  $\chi_{\pm} \in \mathcal{W}_n^{\pm}$  be given. Then, by definition,  $\chi_+$  corresponds to two solutions  $u_{+,w}^+$  and  $u_{-,w}^-$  to (19.1); and  $\chi_-$  corresponds to two solutions  $u_{+,w}^-$  and  $u_{-,w}^+$  to (19.1). Moreover,  $u_{\pm,w}^+$  and  $u_{\pm,w}^-$  have the properties stated in Lemma 19.11. Appealing to Lemma 19.1 yields functions  $u_{\pm,wa}^+$  and  $u_{\pm,wa}^-$  characterised by  $\zeta_{\pm}^+$  and  $\zeta_{\pm}^-$  as described in Remark 19.19. Moreover,

$$\left(\sum_{\iota \in \mathcal{I}_{B,n}} \sum_{\pm} \langle \nu(\iota) \rangle^{2s} [|\zeta_{\pm}^{+}(\iota)|^{2} + |\zeta_{\pm}^{-}(\iota)|^{2}]\right)^{1/2} \le C_{B} [\|\chi_{+}\|_{(s+s_{B})} + \|\chi_{-}\|_{(s+s_{B})}],$$
(19.91)

where  $C_B$  and  $s_B$  have the same dependence as in the case of (19.50); cf. (19.8). In addition,

$$\mathfrak{E}_{s}^{1/2}[u_{\pm,\mathrm{w}} - u_{\pm,\mathrm{wa}}](t) \le C_{B}e^{(\beta_{\mathrm{n}}/2 - \eta_{B})t}[\|\chi_{+}\|_{(s+s_{B})} + \|\chi_{-}\|_{(s+s_{B})}]$$
(19.92)

for all  $t \ge 0$  and all  $s \in \mathbb{R}$ , where  $C_B$  and  $s_B$  have the same dependence as in the case of (19.50), the  $u_{\pm,\text{wa}}$  are defined by (19.57) and the  $u_{\pm,\text{w}}$  are defined analogously; cf. the statement of Lemma 19.11. In order to obtain (19.92), we appealed to (19.6) and (19.91). Defining  $u_{\text{w}}$  by (19.48) and  $u_{\text{wa}}$  by (19.89), the estimate (19.92) yields

$$\mathfrak{E}_{s}^{1/2}[u_{\rm w} - u_{\rm wa}](t) \le C_B \langle t \rangle^{d_{\rm n,+} - 1} e^{(\kappa_{\rm n,+} - \eta_B)t} [\|\chi_+\|_{(s+s_B)} + \|\chi_-\|_{(s+s_B)}]$$
(19.93)

for all  $t \ge 0$  and all  $s \in \mathbb{R}$ , where  $C_B$  and  $s_B$  have the same dependence as in the case of (19.50). Comparing this estimate with the statement of the lemma, it is of interest to construct a solution u to (1.2) (with f = 0) and to estimate the difference between u and  $u_{wa}$  by the right hand side of (19.86), as well as to estimate the initial data for u in terms of  $\chi_{\pm}$ ; cf. also Remark 19.26.

Asymptotics along a time sequence. In order to specify the initial data for u, we wish to appeal to Lemma 18.5. Note that the  $\chi$  appearing in the statement of Lemma 18.5 corresponds to the  $\psi_{\infty}$  appearing in (18.2). However, since  $\psi_{\infty}$  depends on  $\iota$ , we here sometimes write  $\psi_{\infty}(\iota)$ . Due to (18.2), it is of interest to compare

$$\exp\left(i\int_0^{t_k}\sigma(\iota,t')\mathfrak{g}(\iota,t')dt'\right)T_{\mathrm{pre},k}^{-1}[D_k(\iota)]^{-1}e^{R_n(\iota)t_k}\psi_{\infty}(\iota)$$
(19.94)

with the  $\iota$ 'th Fourier coefficient of  $u_{\text{wa}}$ , say  $z_{\text{wa}}(\iota, \cdot)$ . Note that the first factor of the expression (19.94) can be written  $e^{i\omega_{\text{sh}}(\iota,t_k)}$ , and that, due to (18.64),

$$\left| e^{i\omega_{\rm sh}(t_k)} (T_{\rm pre,k}^{-1} - T_{\rm pre}^{-1}) D_k^{-1} e^{R_{\rm n} t_k} \psi_{\infty} \right| \le C \langle t_k \rangle^{d_{\rm n,+} - 1} e^{(\kappa_{\rm n,+} - \eta_B) t_k} |\psi_{\infty}| \tag{19.95}$$

for all  $k \ge 0$ , where C only depends on the coefficients of the equation (1.2). On the other hand, if

$$\psi_{\infty}(\iota) = \begin{pmatrix} \psi_{+}(\iota) \\ \psi_{-}(\iota) \end{pmatrix} = \begin{pmatrix} \psi_{+}^{+}(\iota) + \psi_{-}^{-}(\iota) \\ \psi_{-}^{+}(\iota) + \psi_{-}^{-}(\iota) \end{pmatrix},$$
(19.96)

where  $\psi_{\pm}^{+}(\iota) = 0$  for  $\iota \notin \mathcal{I}_{B,n}^{+}$  and  $\psi_{\pm}^{-}(\iota) = 0$  for  $\iota \notin \mathcal{I}_{B,n}^{-}$ , then

$$\begin{split} e^{i\omega_{\rm sh}(\iota,t_k)} [D_k(\iota)]^{-1} e^{R_{\rm n}(\iota)t_k} \psi_{\infty}(\iota) &= \left(\begin{array}{c} e^{i(\omega_{\rm sh} - \varphi_{\rm tot})(\iota,t_k)} e^{[-\alpha_{\infty} + \beta_{\rm n} {\rm Id}_m + \bar{X}_{\rm n}(\iota)]t_k/2} \psi_+(\iota) \\ e^{i(\omega_{\rm sh} + \varphi_{\rm tot})(\iota,t_k)} e^{[-\alpha_{\infty} + \beta_{\rm n} {\rm Id}_m - \bar{X}_{\rm n}(\iota)]t_k/2} \psi_-(\iota) \end{array}\right) \\ &= \left(\begin{array}{c} e^{i(\omega_{\rm sh} - \varphi_{\rm tot})(\iota,t_k)} [e^{R_{\rm n,+}^- t_k} \psi_+^+(\iota) + e^{R_{\rm n,+}^+ t_k} \psi_-^+(\iota)] \\ e^{i(\omega_{\rm sh} + \varphi_{\rm tot})(\iota,t_k)} [e^{R_{\rm n,+}^+ t_k} \psi_-^+(\iota) + e^{R_{\rm n,+}^- t_k} \psi_-^-(\iota)] \end{array}\right). \end{split}$$

Thus

$$e^{i\omega_{\rm sh}(\iota,t_k)}T_{\rm pre}^{-1}[D_k(\iota)]^{-1}e^{R_{\rm n}(\iota)t_k}\psi_{\infty}(\iota)$$

$$=e^{i(\omega_{\rm sh}-\varphi_{\rm tot})(\iota,t_k)}\begin{pmatrix} e^{R_{\rm n,+}^-t_k}\psi_+^+(\iota) + e^{R_{\rm n,+}^+t_k}\psi_-^-(\iota) \\ -ie^{R_{\rm n,+}^-t_k}\psi_+^+(\iota) - ie^{R_{\rm n,+}^+t_k}\psi_-^-(\iota) \end{pmatrix}$$

$$+e^{i(\omega_{\rm sh}+\varphi_{\rm tot})(\iota,t_k)}\begin{pmatrix} -ie^{R_{\rm n,+}^+t_k}\psi_-^+(\iota) - ie^{R_{\rm n,+}^-t_k}\psi_-^-(\iota) \\ e^{R_{\rm n,+}^+t_k}\psi_-^+(\iota) + e^{R_{\rm n,+}^-t_k}\psi_-^-(\iota) \end{pmatrix}.$$
(19.97)

**Expanding**  $z_{wa}$ . Note that

$$\begin{aligned} z_{\rm wa}(\iota,t) &= e^{A_{\rm n,+}t} z_{+,\rm wa}(\iota,t) + e^{A_{\rm n,-}t} z_{-,\rm wa}(\iota,t) \\ &= e^{A_{\rm n,+}t} [z^+_{+,\rm wa}(\iota,t) + z^-_{-,\rm wa}(\iota,t)] + e^{A_{\rm n,-}t} [z^-_{+,\rm wa}(\iota,t) + z^+_{-,\rm wa}(\iota,t)], \end{aligned}$$

where the  $z_{\pm,\text{wa}}$ ,  $z_{\pm,\text{wa}}^+$  and  $z_{\pm,\text{wa}}^-$  have the properties stated in Remark 19.19. Let us estimate

$$\left| \begin{pmatrix} \mathfrak{g}z_{\mathrm{wa}} \\ \dot{z}_{\mathrm{wa}} \end{pmatrix} - \begin{pmatrix} e^{A_{\mathrm{n},+}t}\mathfrak{g}z_{+,\mathrm{wa}}^+ \\ e^{A_{\mathrm{n},+}t}\dot{z}_{+,\mathrm{wa}}^+ \end{pmatrix} - \begin{pmatrix} e^{A_{\mathrm{n},+}t}\mathfrak{g}z_{-,\mathrm{wa}}^- \\ e^{A_{\mathrm{n},+}t}\dot{z}_{-,\mathrm{wa}}^- \end{pmatrix} - \begin{pmatrix} e^{A_{\mathrm{n},-}t}\mathfrak{g}z_{-,\mathrm{wa}}^- \\ e^{A_{\mathrm{n},-}t}\dot{z}_{+,\mathrm{wa}}^+ \end{pmatrix} - \begin{pmatrix} e^{A_{\mathrm{n},-}t}\mathfrak{g}z_{-,\mathrm{wa}}^- \\ e^{A_{\mathrm{n},-}t}\dot{z}_{+,\mathrm{wa}}^- \end{pmatrix} \right| \le C_B \langle t \rangle^{d_{\mathrm{n},+}-1} e^{(\kappa_{\mathrm{n},+}-\eta_B)t} \langle \nu \rangle^{s_E} \sum_{\pm} (|\zeta_{\pm}^+|+|\zeta_{\pm}^-|)$$

$$(19.98)$$

for all  $t \ge 0$ , where  $C_B$  has the same dependence as in the case of (19.50) and we omitted the arguments  $\iota$  and t for the sake of brevity. Moreover, we appealed to (19.27), which holds in the present setting. By arguments similar to those presented in connection with (19.67), we can replace  $\dot{z}^{\pm}_{+,\text{wa}}$  and  $\dot{z}^{\pm}_{-,\text{wa}}$  appearing in (19.98) by  $-i\mathfrak{g}z^{\pm}_{+,\text{wa}}$  and  $i\mathfrak{g}z^{\pm}_{-,\text{wa}}$  respectively; cf. (19.55) and (19.56). One particular expression it is of interest to consider is thus

$$\begin{pmatrix} e^{A_{\mathrm{n},+}t}\mathfrak{g}z^+_{+,\mathrm{wa}}\\ -ie^{A_{\mathrm{n},+}t}\mathfrak{g}z^+_{+,\mathrm{wa}} \end{pmatrix} = e^{i(\omega_{\mathrm{sh}}-\varphi_{\mathrm{tot}})} \begin{pmatrix} e^{R_{\mathrm{n},+}^-t}\zeta^+_+\\ -ie^{R_{\mathrm{n},+}^-t}\zeta^+_+ \end{pmatrix}.$$

Combining this observation with similar reformulations of the expressions appearing on the left hand side of (19.98) yields

$$\begin{vmatrix} \left( \begin{array}{c} \mathfrak{g}_{z_{\text{wa}}} \\ \dot{z}_{\text{wa}} \end{array} \right) - e^{i(\omega_{\text{sh}} - \varphi_{\text{tot}})} \left( \begin{array}{c} e^{R_{n,+}^{-}t}\zeta_{+}^{+} + e^{R_{n,+}^{+}t}\zeta_{+}^{-} \\ -ie^{R_{n,+}^{-}t}\zeta_{+}^{+} - ie^{R_{n,+}^{+}t}\zeta_{+}^{-} \end{array} \right) \\ - e^{i(\omega_{\text{sh}} + \varphi_{\text{tot}})} \left( \begin{array}{c} e^{R_{n,+}^{-}t}\zeta_{-}^{-} + e^{R_{n,+}^{+}t}\zeta_{-}^{+} \\ ie^{R_{n,+}^{-}t}\zeta_{-}^{-} + ie^{R_{n,+}^{+}t}\zeta_{-}^{+} \end{array} \right) \\ \leq C_{B}\langle t \rangle^{d_{n,+} - 1} e^{(\kappa_{n,+} - \eta_{B})t} \langle \nu \rangle^{s_{E}} \sum_{\pm} (|\zeta_{\pm}^{+}| + |\zeta_{\pm}^{-}|), \qquad (19.99)$$

for all  $t \ge 0$ , where  $C_B$  has the same dependence as in the case of (19.50).

**Comparing the different perspectives.** Comparing (19.97) and (19.99), it is clear that we want the following relations to hold:

$$\psi_{+}^{\pm}(\iota) = \zeta_{+}^{\pm}(\iota), \quad \psi_{-}^{\pm}(\iota) = i\zeta_{-}^{\pm}(\iota).$$
 (19.100)

For a given  $\iota \in \mathcal{I}_{B,n}$ , it is thus clear how to specify  $\psi_{\infty}$  in terms of the  $\zeta_{+}^{\pm}$  and  $\zeta_{-}^{\pm}$ . It is also important to note that, due to the properties of the right hand sides of (19.100),  $\psi_{\infty}(\iota)$  is an element of the first generalised eigenspace of the  $\eta_B$ ,  $R_n(\iota)$ -decomposition of  $\mathbb{C}^{2m}$ . Next, we can then appeal to Lemma 18.5 with  $\chi$  replaced by  $\psi_{\infty}$ . This yields initial data for (24.16) (with vanishing right hand side) at  $T_n$ . In fact, (18.37) and (18.38) hold. Thus

$$|\dot{z}(\iota, T_{\mathrm{n}})| + \mathfrak{g}(\iota, T_{\mathrm{n}})|z(\iota, T_{\mathrm{n}})| \le C|e^{R_{\mathrm{n}}(\iota)T_{\mathrm{n}}}\psi_{\infty}(\iota)| \le C\langle T_{\mathrm{n}}\rangle^{d_{\mathrm{n}, +} - 1}e^{\kappa_{\mathrm{n}, +}T_{\mathrm{n}}}|\psi_{\infty}(\iota)|,$$
(19.101)

where C only depends on the coefficients of the equation (1.2). Next, we need to derive an estimate for the energy of the mode at t = 0. To this end, we combine (19.101) and Lemma 10.7 in order to obtain

$$\mathcal{E}^{1/2}(\iota,0) \le C e^{(\kappa_{\mathrm{n},+}+1+\eta_{\mathrm{bal}})T_{\mathrm{n}}} |\psi_{\infty}(\iota)|,$$

where C only depends on the coefficients of the equation (1.2). Combining this estimate with an estimate for  $T_n$  of the form (19.23) yields

$$\mathcal{E}^{1/2}(\iota,0) \le C\langle\nu(\iota)\rangle^{s_a} |\psi_{\infty}(\iota)|, \tag{19.102}$$

where C and  $s_a$  only depend on the coefficients of the equation (1.2).

**Defining**  $\Phi_{n,T}$  and  $\Phi_{n,g}$ . For reasons already mentioned, we can focus on the case of a dominant noisy spatial torus direction here. Given  $\chi_{\pm} \in \mathcal{W}_n^{\pm}$ , we calculate  $\zeta_{\pm}^+$  and  $\zeta_{\pm}^-$  as described above. Note also that (19.91) holds. Next,  $\psi_{\infty}$  is defined by (19.96) and (19.100). For a given  $\iota \in \mathcal{I}_{B,n}$ , this yields  $z(\iota, T_n)$  and  $\dot{z}(\iota, T_n)$ . Solving (24.16) backwards, this yields initial data for the  $\iota$ 'th mode at t = 0. Combining these maps for the individual modes yields the maps  $\Phi_{n,T}$  and  $\Phi_{n,g}$ . Moreover, combining (19.91) and (19.102) yields (19.85) and (19.87). The injectivity of the maps follows from the injectivity of the maps of which they are composed. Similarly, if  $E_{n,+} = E_{n,-} = \mathbb{C}^m$ , then the surjectivity statement of Lemma 18.5 yields the desired surjectivity for fixed  $\iota \in \mathcal{I}_{B,n}$ .

Asymptotics. What remains to be verified is that (19.84) and (19.86) hold, and that surjectivity holds if  $E_{n,+} = E_{n,-} = \mathbb{C}^m$ . Let  $\chi_{\pm} \in \mathcal{W}_n^{\pm}$  be given. Then we obtain initial data for (1.2) (with f = 0). Solving (1.2) yields a solution u such that  $u = u_n$  (by construction). Moreover, if z denotes the Fourier coefficient of u, then (18.2) holds, where  $\psi_{\infty}(\iota)$  is defined by (19.96) and (19.100). Keeping (19.95) in mind, the estimate

$$\left| \begin{pmatrix} \mathfrak{g}(\iota, t_k) z(\iota, t_k) \\ \dot{z}(\iota, t_k) \end{pmatrix} - e^{i\omega_{\rm sh}(\iota, t_k)} T_{\rm pre}^{-1} [D_k(\iota)]^{-1} e^{R_{\rm n}(\iota)t_k} \psi_{\infty}(\iota) \right| \le C \langle t_k \rangle^N e^{(\kappa_{\rm n,+} - \eta_B)t_k}$$
(19.103)

holds for all  $k \geq 0$ , where *C* is allowed to depend on the solution,  $\iota$  etc. In what follows, it is important to keep in mind that  $\psi_{\infty}(\iota)$  belongs to the first generalised eigenspace of the  $\eta_B$ ,  $R_n(\iota)$ -decomposition of  $\mathbb{C}^{2m}$ .

Next, appealing to Remark 19.19, there are  $\bar{\zeta}^+_{\pm}(\iota)$  and  $\bar{\zeta}^-_{\pm}(\iota)$ ,  $\iota \in \mathcal{I}_{\mathrm{B}}$ , which are uniquely associated with u and satisfy the conditions stated in Remark 19.19. Moreover, we obtain a function  $\bar{u}_{\mathrm{wa}}$ with the properties stated in Remark 19.19. In particular, (19.58) yields

$$\mathfrak{E}_s^{1/2}[u-\bar{u}_{\mathrm{wa}}](t) \le C_B \langle t \rangle^N e^{(\kappa_{\mathrm{n},+}-\eta_B)t} \mathfrak{E}_{s+s_B}^{1/2}[u](0)$$

for all  $t \geq 0$  and all  $s \in \mathbb{R}$ , so that, letting  $\bar{z}_{wa}$  denote the Fourier coefficients of  $\bar{u}_{wa}$ ,

$$\left| \begin{pmatrix} \mathfrak{g}(\iota,t)z(\iota,t) \\ \dot{z}(\iota,t) \end{pmatrix} - \begin{pmatrix} \mathfrak{g}(\iota,t)\bar{z}_{\mathrm{wa}}(\iota,t) \\ \dot{\bar{z}}_{\mathrm{wa}}(\iota,t) \end{pmatrix} \right| \leq C \langle t \rangle^{N} e^{(\kappa_{\mathrm{n},+}-\eta_{B})t}$$

for all  $t \ge 0$ , where *C* is allowed to depend on the solution,  $\iota$  etc. Next we can repeat, verbatim, the argument leading up to (19.99). Thus (19.99) holds with  $z_{wa}$  replaced by  $\bar{z}_{wa}$  and  $\zeta_{\pm}^+$  and  $\zeta_{\pm}^-$  replaced by the analogous barred quantities. Introducing  $\bar{\psi}_{\infty}(\iota)$  in analogy with (19.100) and (19.96) (starting with  $\bar{\zeta}_{\pm}^+(\iota)$  and  $\bar{\zeta}_{\pm}^-(\iota)$ ), it follows that

$$\left| \begin{pmatrix} \mathfrak{g}(\iota, t_k) z(\iota, t_k) \\ \dot{z}(\iota, t_k) \end{pmatrix} - e^{i\omega_{\rm sh}(\iota, t_k)} T_{\rm pre}^{-1} [D_k(\iota)]^{-1} e^{R_{\rm n}(\iota)t_k} \bar{\psi}_{\infty}(\iota) \right| \le C \langle t_k \rangle^N e^{(\kappa_{\rm n,+} - \eta_B)t_k}$$
(19.104)

for all  $k \geq 0$ , where *C* is allowed to depend on the solution,  $\iota$  etc., and we have appealed to the barred analogues of (19.97) and (19.99). Moreover, due to the properties of the  $\bar{\zeta}^+_{\pm}(\iota)$  and  $\bar{\zeta}^-_{\pm}(\iota)$ ,  $\bar{\psi}_{\infty}(\iota)$  belongs to the first generalised eigenspace of the  $\eta_B$ ,  $R_n(\iota)$ -decomposition of  $\mathbb{C}^{2m}$ . Combining (19.103) and (19.104) with the fact that  $\bar{\psi}_{\infty}(\iota)$  and  $\psi_{\infty}(\iota)$  belong to the first generalised eigenspace of the  $\eta_B$ ,  $R_n(\iota)$ -decomposition of  $\mathbb{C}^{2m}$  yields the conclusion that  $\bar{\psi}_{\infty}(\iota) = \psi_{\infty}(\iota)$ . In particular,

$$\bar{\zeta}_{\pm}^{+}(\iota) = \zeta_{\pm}^{+}(\iota), \quad \bar{\zeta}_{\pm}^{-}(\iota) = \zeta_{\pm}^{-}(\iota)$$
(19.105)

for all  $\iota \in \mathcal{I}_{B}$ . Next, we wish to use this information to relate  $u_{w}$ , appearing in the statement of the lemma, with  $\bar{u}_{wa}$ . First of all,  $u_{w}$  is determined by  $u_{\pm,w}^{+}$  and  $u_{\pm,w}^{-}$  (the solutions to (19.1) which are uniquely determined by  $\chi_{\pm}$ ). On the other hand,  $u_{\pm,w}^{+}$  and  $u_{\pm,w}^{-}$  correspond to functions  $u_{\pm,wa}^{+}$  and  $u_{\pm,wa}^{-}$  respectively, as described in Lemma 19.1. Moreover, the latter functions correspond to the  $\zeta_{\pm}^{+}(\iota)$  and  $\zeta_{\pm}^{-}(\iota)$ . Finally, each of these correspondences arise from a bijection. Turning to  $\bar{u}_{wa}$ , it can be written

$$\bar{u}_{wa} = e^{A_{n,+}t}\bar{u}_{+,wa} + e^{A_{n,-}t}\bar{u}_{-,wa}$$

where

$$\bar{u}_{+,\mathrm{wa}} := \bar{u}_{+,\mathrm{wa}}^+ + \bar{u}_{-,\mathrm{wa}}^-, \quad \bar{u}_{-,\mathrm{wa}} := \bar{u}_{+,\mathrm{wa}}^- + \bar{u}_{-,\mathrm{wa}}^+$$

and  $\bar{u}_{\tau,\text{wa}}^{\upsilon}$ ,  $\upsilon, \tau \in \{+, -\}$ , are the functions uniquely determined by  $\bar{\zeta}_{\pm}^{+}(\iota)$  and  $\bar{\zeta}_{\pm}^{-}(\iota)$ . Due to (19.105) it follows that

$$u_{+,\mathrm{wa}} := u_{+,\mathrm{wa}}^+ + u_{-,\mathrm{wa}}^- = \bar{u}_{+,\mathrm{wa}}, \quad u_{-,\mathrm{wa}} := u_{-,\mathrm{wa}}^+ + u_{+,\mathrm{wa}}^- = \bar{u}_{-,\mathrm{wa}}.$$

In particular,

$$u_{\text{wa}} := e^{A_{\text{n},+}t} u_{+,\text{wa}} + e^{A_{\text{n},-}t} u_{-,\text{wa}} = \bar{u}_{\text{wa}}$$

Combining (19.93), (19.58) (which we know to hold with  $u_{wa}$  replaced by  $\bar{u}_{wa}$  and f replaced by 0) and the fact that  $u_{wa} = \bar{u}_{wa}$  yields

$$\begin{aligned} \mathfrak{E}_{s}^{1/2}[u-u_{w}](t) &\leq \mathfrak{E}_{s}^{1/2}[u-u_{wa}](t) + \mathfrak{E}_{s}^{1/2}[u_{wa}-u_{w}](t) \\ &\leq C_{B}\langle t\rangle^{d_{n,+}-1}e^{(\kappa_{n,+}-\eta_{B})t}[\|\chi_{+}\|_{(s+s_{B})} + \|\chi_{-}\|_{(s+s_{B})}] \\ &+ C_{B}\langle t\rangle^{N}e^{(\kappa_{n,+}-\eta_{B})t}\mathfrak{E}_{s+s_{B}}^{1/2}[u](0) \end{aligned}$$

for all  $t \ge 0$  and all  $s \in \mathbb{R}$ . Here  $C_B > 0$  and  $s_B \ge 0$  only depend on  $\eta_B$  and the coefficients of the equation (1.2), and N only depends on m. Moreover, N = 1 in case the Jordan blocks of the matrices  $A_{n,\pm}$  are trivial. In addition, if  $E_{n,+} = E_{n,-} = \mathbb{C}^m$ , then N can be replaced by  $d_{n,+} - 1$ . This is not exactly the estimate we want. However, combining it with (19.87) (which we already demonstrated to hold), it yields (19.84) and (19.86).

**Surjectivity.** In order to demonstrate surjectivity in case  $E_{n,+} = E_{n,-} = \mathbb{C}^m$ , let u be a smooth solution to (1.2) such that  $u = u_n$ . Appealing to Lemma 19.11 yields solutions  $u_{\tau,w}^v$ ,  $v, \tau \in \{+, -\}$ , and  $u_{\pm,w}$  to (19.1) such that the conclusions of Lemma 19.11 hold. Moreover, if  $u_w$  is defined by (19.48), then

$$\mathfrak{E}_s^{1/2}[u-u_{\mathbf{w}}](t) \le C\langle t \rangle^N e^{(\kappa_{\mathbf{n},+}-\eta_B)t}$$
(19.106)

for all  $t \ge 0$  and  $s \in \mathbb{R}$ , where *C* is allowed to depend on the solution, *s*, etc. The initial data for  $u_{\tau,w}^v$  correspond to functions  $\chi_{\pm}$  with properties stated in the present lemma. Letting  $\bar{u}$  be the solution corresponding to initial data  $\Phi_{n,T}(\chi_+, \chi_-)$  at t = 0, the earlier parts of the present proof guarantee that (19.106) holds if we replace u by  $\bar{u}$ . Letting  $v = u - \bar{u}$  and z denote the Fourier coefficients of v, we conclude that for each  $\iota \in \mathcal{I}_{B,n}$ ,

$$\left| \begin{pmatrix} \mathfrak{g}(\iota, t_k) z(\iota, t_k) \\ \dot{z}(\iota, t_k) \end{pmatrix} \right| \le C \langle t_k \rangle^N e^{(\kappa_{n,+} - \eta_B) t_k}$$
(19.107)

for all  $k \geq 0$ . Assume now that the initial data for  $z(\iota, \cdot)$  at  $T_n$  are not zero. Then, due to the surjectivity of the map  $\Psi_{\infty}$  introduced in Lemma 18.5, there is a  $0 \neq \chi \in \mathbb{C}^{2m}$  such that (18.37) holds. Moreover, Lemma 18.5 implies that (18.2) holds with  $\psi_{\infty}$  replaced by  $\chi$ . However, (18.2) (with  $\psi_{\infty} \neq 0$ ) is irreconcilable with (19.107). Thus v = 0 and  $u = \bar{u}$ . Thus the initial data of u are in the image of  $\Phi_{n,T}$ . Since u is an arbitrary smooth solution, the surjectivity of  $\Phi_{n,T}$  follows.

### Part VI

## Non-degenerate, diagonally dominated, convergent and balanced equations

### Chapter 20

## Equations with diagonally dominated, convergent and balanced coefficients

#### 20.1 Introduction

In Part III, we consider Fourier coefficients of solutions to the equation (1.2) under circumstances in which the behaviour is oscillatory. In particular, we derive formulae for how the solution evolves over one period of the oscillations; cf., in particular, (15.54)-(15.58). This analysis is the basis for a further study of the evolution of solutions. However, in order to make progress, we need to make additional assumptions. To be more specific, consider the matrices  $R^{ab}_{\text{pre},k}$  appearing on the right hand side of (15.57). These matrices are of central importance, since they determine the leading order behaviour of the evolution. Returning to (15.41), it is clear that we need to study  $\alpha/\mathfrak{g}$ ,  $\dot{\ell}/\mathfrak{g}$ and  $X/\mathfrak{g}$  in greater detail. Naively, we would like to think of the matrix  $\Xi^{k,\pm}_{\text{fn,app}}$  as follows:

$$\begin{aligned} \Xi_{\text{fin,app}}^{k,\pm} = & \text{Id}_{2m} \pm \frac{1}{2} \frac{2\pi}{\mathfrak{g}(t_k)} \begin{pmatrix} -\alpha(t_k) + \dot{\ell}(t_k) \text{Id}_m + X(t_k) & 0 \\ 0 & -\alpha(t_k) + \dot{\ell}(t_k) \text{Id}_m - X(t_k) \end{pmatrix} \\ \approx & \text{Id}_{2m} \pm \frac{1}{2} \int_{t_k}^{t_{k+1}} \begin{pmatrix} -\alpha(t) + \dot{\ell}(t) \text{Id}_m + X(t) & 0 \\ 0 & -\alpha(t) + \dot{\ell}(t) \text{Id}_m - X(t) \end{pmatrix} dt \\ \approx & \exp\left[ \pm \frac{1}{2} \int_{t_k}^{t_{k+1}} \begin{pmatrix} -\alpha(t) + \dot{\ell}(t) \text{Id}_m + X(t) & 0 \\ 0 & -\alpha(t) + \dot{\ell}(t) \text{Id}_m - X(t) \end{pmatrix} dt \right]. \end{aligned}$$
(20.1)

The justification of the first approximate equality is based on (14.27) and the assumption that  $\alpha$ ,  $\ell$ and X do not vary too much over one period of the oscillations. The second approximate equality is based on the assumption that the integral appearing in the final exponential expression is small. The advantage of the expression on the far right hand side of (20.1) is that it behaves well under multiplication (at least in intervals in which  $\alpha$  and X are essentially constant). It is thus tempting to hope that the product of the  $\Xi_{\text{fin,app}}^{k,\pm}$  can be approximated by the far right hand side of (20.1), where the time interval  $[t_k, t_{k+1}]$  is replaced by the union of the corresponding intervals. However, it is very optimistic to hope for this in general. The reason for this is that the resulting matrix can be expected to contract some directions more than others. Taking a vector corresponding to a direction which is maximally contracted, the small errors associated with the approximations in (20.1) can be expected to generate a small component in the direction which is maximally expanded. The contribution of this small error may then (over long periods of time) yield a bigger contribution to the end result than the original vector. This situation is reminiscent of the one considered already in Chapter 9. In fact, consider (9.3) with F = 0, where A is constant and  $A_{\text{rem}}$  satisfies (9.2). Naively, one might then expect  $\exp(At)$  to be a good approximation of the evolution. However, this expectation is not reasonable if the real eigenvalue spread of A, RspA, exceeds the parameter  $\beta_{\text{rem}} > 0$  appearing in (9.2). To summarise the discussion, (20.1) may give a rough idea of how solutions evolve, even when replacing the interval  $[t_k, t_{k+1}]$  by the union of several periods. However, as a guide to the asymptotic behaviour, it should be treated with care.

Diagonally dominated, non-degenerate, convergent and balanced equations. Due to the above discussion, it is clear that we wish to eliminate the risk of a substantial error arising due to a vector corresponding to a contracting direction being rotated (by a small error) in such a way that it picks up a component corresponding to an expanding direction. In order to avoid problems of this type, it is convenient to keep track of which directions correspond to expansion and which directions correspond to contraction. However, we would preferably like the matrices giving rise to these directions to (in the limit) be finite in number; otherwise error terms arising due to an unbounded number of algebraic manipulations could be significant. One way to fulfil these conditions is to consider situations in which the diagonal components of the metric are dominant and such that they converge after suitable rescaling by an exponential function. We also need to make corresponding assumptions concerning  $X^l$ ,  $\alpha$  and  $\zeta$ . In order to develop some intuition, consider the model equation

$$u_{tt} - \sum_{j=1}^{d} g_{\infty}^{jj} e^{2\beta_j t} \partial_j^2 u + \sum_{j=1}^{d} X_{\infty}^j e^{\beta_j t} \partial_j u + \alpha_{\infty} u_t + \zeta_{\infty} u = 0,$$
(20.2)

where  $0 < g_{\infty}^{jj} \in \mathbb{R}$ ,  $X_{\infty}^{j}, \alpha_{\infty}, \zeta_{\infty} \in \mathbf{M}_{m}(\mathbb{C})$ , j = 1, ..., d. Finally, we assume that  $\beta_{1} < \beta_{2} < \cdots < \beta_{d}$ ; this corresponds to the condition of non-degeneracy. For an equation of the form (20.2),  $\mathfrak{g}$  satisfies the relation

$$\mathfrak{g}^2(\iota,t) = \sum_{j=1}^d g_\infty^{jj} e^{2\beta_j t} n_j^2.$$

For a given  $\iota = n \in \mathbb{Z}^d$ , it is then clear that the time interval  $[0, \infty)$  can be divided into subintervals, which we refer to as *eras*, in which one of the functions  $g_{\infty}^{jj}e^{2\beta_j t}n_j^2$ ,  $j = 1, \ldots, d$ , and 1 dominates (if all the  $\beta_j$  are strictly negative, the function 1 will eventually dominate; we refer to the corresponding interval as the *ODE era*). Let  $I_j$  be an interval in which  $g_{\infty}^{jj}e^{2\beta_j t}n_j^2$  dominates. In  $I_j$ ,  $\mathfrak{g}$  can then effectively be replaced by  $(g_{\infty}^{jj})^{1/2}e^{\beta_j t}|n_j|$  (the exact meaning of this statement, as well as a detailed justification of it, is provided in Lemmas 20.5 and 20.8 below). Moreover, Xcan be replaced by

$$\frac{in_j X_{\infty}^j e^{\beta_j t}}{(g_{\infty}^{jj})^{1/2} |n_j| e^{\beta_j t}} = \pm i (g_{\infty}^{jj})^{-1/2} X_{\infty}^j,$$

where the sign equals the sign of  $n_j$ . Finally,  $\ell$  is effectively  $\beta_j$ . Thus, for  $t_k \in I_j$ ,

$$R_{\text{pre},k}^{ab} \approx \frac{1}{(g_{\infty}^{jj})^{1/2} |n_j| e^{\beta_j t_k}} [(-1)^a \alpha_{\infty} + \beta_j \text{Id}_m \pm i (-1)^b (g_{\infty}^{jj})^{-1/2} X_{\infty}^j],$$
(20.3)

where the sign equals the sign of  $n_j$ . Note that the first factor roughly speaking corresponds to integration over the period of the oscillations under consideration. The essential structure appearing on the right hand side of (20.3) is thus captured by one element of a finite set of matrices. Keeping track of expanding and contracting directions in  $I_j$  is therefore straightforward. To summarise: for a fixed n, there are finitely many eras (never more than d + 1); for each era (except for the ODE era, if it exists), the dominant part of the evolution over one period of the oscillations is determined by one matrix (chosen from a finite set of matrices); and the evolution in the ODE era (if it exists) can be analysed as described in Chapter 9.

Specifying the class of equations of interest. With the above intuition in mind, we introduce the class of equations of interest in the present part of these notes. We do not demand that the equations be of the form (20.2). However, when R = 0, we do demand that this be the dominant part asymptotically. On the other hand, we do not, at this stage, insist on non-degeneracy. For this reason, the material of the present chapter also serves as a basis for the study of degenerate equations (a topic we do not address in these notes). Moreover, as opposed to (20.2), we wish to allow R > 0; cf. (1.2). Giving a precise definition of the corresponding class of equations is the main purpose of Section 20.2. However, we also verify that this class is such that the results of Part III apply; cf. Lemma 20.3.

Simplifying  $R_{\text{pre},k}^{ab}$ . Our next goal is to simplify the expressions  $\alpha/\mathfrak{g}$ ,  $\dot{\ell}/\mathfrak{g}$  and  $X/\mathfrak{g}$  appearing on the right hand side of (15.41). We do this in two steps. In Section 20.3, in particular in Lemma 20.5, we estimate the error associated with ignoring the off-diagonal components of the matrix  $g^{jl}$ . In order to progress further, we need to divide the time interval  $[0,\infty)$  into eras in which particular frequencies and exponential behaviour dominate. Defining such a division and deriving the associated simplified expressions for  $\alpha/\mathfrak{g}$ ,  $\dot{\ell}/\mathfrak{g}$  and  $X/\mathfrak{g}$  is the subject of Section 20.4, in particular Lemma 20.8.

Estimating sums. The final observation in the present chapter concerns the estimation of sums that appear when calculating matrix products of the type appearing in (15.64). In practice, our ambition is not to calculate the product, but rather the leading order behaviour. In order to be able to do so, we first have to divide the interval  $[0, \infty)$  into frequency eras, as described above. In a given frequency era, we can then approximate  $R^{ab}_{\text{pre},k}$  in such a way that the dominant behaviour is determined by one fixed matrix. Transforming the iteration so that this matrix is in Jordan normal form, the problem of estimating the evolution of solutions can then be reduced to estimating certain sums. In Section 20.5 we demonstrate how to estimate the types of sums that appear; cf. Lemma 20.12.

# 20.2 Equations with diagonally dominant, convergent and balanced coefficients

Let us return to the equation (1.2) and impose the conditions of interest in the present part of these notes. The motivation for introducing these conditions is described in the previous section.

**Definition 20.1.** Consider (1.2). Assume the associated metric to be such that (M, g) is a canonical separable cosmological model manifold. The coefficients  $\alpha$ ,  $X^j$  and  $\zeta$ ,  $j = 1, \ldots, d$ , are referred to as the main coefficients of the equation (1.2). The coefficients  $g^{0l}$ ,  $g^{jj}$  and  $g^{jl}$  ( $j \neq l$ ),  $j, l = 1, \ldots, d$ , are referred to as the shift coefficients, the diagonal coefficients and the off-diagonal coefficients of the equation (1.2) respectively. The functions  $a_r^{-2}$ ,  $r = 1, \ldots, R$ , are referred to as the auxiliary diagonal coefficients. The diagonal coefficients are said to be  $C_{\exp}^0$ -convergent and  $C_{\exp}^2$ -bounded if there are real constants  $\beta_j$ , strictly positive real numbers  $g_{\infty}^{ij}$ ,  $j = 1, \ldots, d$ , and constants  $\eta_d, C_d > 0$  such that

$$e^{\eta_{\rm d}t}|e^{-2\beta_j t}g^{jj}(t) - g^{jj}_{\infty}| + |e^{-2\beta_j t}\dot{g}^{jj}(t)| + |e^{-2\beta_j t}\ddot{g}^{jj}(t)| \le C_{\rm d}$$
(20.4)

for all  $t \ge 0$  and  $j = 1, \ldots, d$ . If, in addition,

$$|e^{-2\beta_j t} \dot{g}^{jj}(t) - 2\beta_j g^{jj}_{\infty}| \le C_{\rm d} e^{-\eta_{\rm d} t}$$
(20.5)

for all  $t \ge 0$  and  $j = 1, \ldots, d$ , the diagonal coefficients are said to be  $C_{\exp}^1$ -convergent and  $C_{\exp}^2$ bounded. Similarly, the auxiliary diagonal coefficients are said to be  $C_{\exp}^0$ -convergent and  $C_{\exp}^2$ bounded if there are real constants  $\beta_{\text{Ri},r}$ , strictly positive real numbers  $a_{r,\infty}$ ,  $r = 1, \ldots, R$ , and constants  $\eta_{\text{ad}}, C_{\text{ad}} > 0$  such that

$$e^{\eta_{\rm ad}t}|e^{\beta_{{\rm Ri},r}t}a_r(t) - a_{r,\infty}| + |e^{\beta_{{\rm Ri},r}t}\dot{a}_r(t)| + |e^{\beta_{{\rm Ri},r}t}\ddot{a}_r(t)| \le C_{\rm ad}$$
(20.6)

for all  $t \ge 0$  and  $r = 1, \ldots, R$ . If, in addition,

$$|e^{\beta_{\mathrm{Ri},r}t}\dot{a}_r(t) + \beta_{\mathrm{Ri},r}a_{r,\infty}| \le C_{\mathrm{ad}}e^{-\eta_{\mathrm{ad}}t}$$
(20.7)

for all  $t \ge 0$  and  $j = 1, \ldots, d$ , the auxiliary diagonal coefficients are said to be  $C_{exp}^1$ -convergent and  $C_{exp}^2$ -bounded. Assuming the diagonal coefficients are  $C_{exp}^0$ -convergent and  $C_{exp}^2$ -bounded, the main coefficients are said to be  $C_{exp}^1$ -balanced and bounded if there is a real constant  $C_{mn} > 0$  such that

$$e^{-\beta_j t} \left( \|X^j(t)\| + \|\dot{X}^j(t)\| \right) + \|\alpha(t)\| + \|\dot{\alpha}(t)\| + \|\dot{\zeta}(t)\| + \|\dot{\zeta}(t)\| \le C_{\rm mn}$$
(20.8)

for all  $t \ge 0$  and j = 1, ..., d. If, in addition, there are  $\alpha_{\infty}, X^j_{\infty}, \zeta_{\infty} \in \mathbf{M}_m(\mathbb{C})$  as well as a real constant  $\eta_{mn} > 0$  such that

$$\|e^{-\beta_j t} X^j(t) - X^j_{\infty}\| + \|\alpha(t) - \alpha_{\infty}\| + \|\zeta(t) - \zeta_{\infty}\| \le C_{\rm mn} e^{-\eta_{\rm mn} t}$$
(20.9)

for all  $t \ge 0$  and j = 1, ..., d, the main coefficients are said to be  $C_{\exp}^0$ -balanced and convergent and  $C_{\exp}^1$ -balanced and bounded. The shift coefficients are said to be negligible if there are real constants  $C_{sh}, \eta_{sh} > 0$  such that

$$\sup_{\xi \neq 0} \frac{|g^{0l}(t)\xi_l|}{[g^{jl}(t)\xi_j\xi_l]^{1/2}} + \sup_{\xi \neq 0} \frac{|\dot{g}^{0l}(t)\xi_l|}{[g^{jl}(t)\xi_j\xi_l]^{1/2}} \le C_{\rm sh}e^{-\eta_{\rm sh}t}$$
(20.10)

for all  $t \ge 0$ , where Einstein's summation convention is enforced. The off-diagonal coefficients are said to be *negligible* if there are real constants  $C_{od}$ ,  $\eta_{od} > 0$  such that

$$\sum_{j \neq l} [|g^{jl}(t)\xi_j\xi_l| + |\dot{g}^{jl}(t)\xi_j\xi_l| + |\ddot{g}^{jl}(t)\xi_j\xi_l|] \le C_{\rm od}e^{-\eta_{\rm od}t}\sum_j g^{jj}(t)\xi_j\xi_j$$
(20.11)

for all  $t \ge 0$  and  $\xi \in \mathbb{R}^d$ . If all the above conditions are satisfied, (1.2) is said to be *diagonally* dominated, balanced and convergent.

**Remark 20.2.** In (20.11), the sum is over all distinct j and l, since neither j nor l is fixed from the context. Note, however, that if l were fixed, then the sum appearing in (20.11) should have been interpreted as being over all j different from l.

To begin with, it is of interest to verify that (1.2) is oscillation adapted, cf. Definition 14.1, when the requirements of the previous definition are fulfilled.

**Lemma 20.3.** Consider (1.2). Assume that the diagonal and auxiliary diagonal coefficients are  $C_{\exp}^0$ -convergent and  $C_{\exp}^2$ -bounded; that the main coefficients are  $C_{\exp}^1$ -balanced and bounded; and that the shift coefficients and the off diagonal coefficients are negligible. Then there is a constant C > 1 such that

$$C^{-1}g^{jl}(t)\xi_{j}\xi_{l} \leq \sum_{j}g^{jj}(t)\xi_{j}\xi_{j} \leq Cg^{jl}(t)\xi_{j}\xi_{l}, \qquad (20.12)$$

$$C^{-1}g^{jl}(t)\xi_{j}\xi_{l} \leq \sum_{j} e^{2\beta_{j}t}\xi_{j}\xi_{j} \leq Cg^{jl}(t)\xi_{j}\xi_{l}$$
(20.13)

for all  $t \geq 0$  and  $\xi \in \mathbb{R}^d$ . Moreover,

$$C^{-1} \sum_{r=1}^{R} a_r^{-2}(t) \eta_r^2 \le \sum_{r=1}^{R} e^{2\beta_{\text{Ri},r}t} \eta_r^2 \le C \sum_{r=1}^{R} a_r^{-2}(t) \eta_r^2$$
(20.14)

for all  $\eta \in \mathbb{R}^R$ . Finally, choosing **b** and C large enough and defining

 $\mathfrak{f}_{\mathrm{sh}}(t) := C e^{-\eta_{\mathrm{sh}}t}, \quad \mathfrak{f}_X(t) := C, \quad \mathfrak{f}_{\mathrm{ode}}(t) := C, \tag{20.15}$ 

#### (1.2) is oscillation adapted.

**Remark 20.4.** The constants C and  $\mathfrak{b}$  appearing in the statement of the lemma only depend on the coefficients of the equation (1.2).

*Proof.* Let us begin by demonstrating the norm equivalences (20.12)-(20.14).

Norm equivalences. In order to prove that (20.12) holds, estimate

$$g^{jl}(t)\xi_{j}\xi_{l} = \sum_{j \neq l} g^{jl}(t)\xi_{j}\xi_{l} + \sum_{j} g^{jj}(t)\xi_{j}\xi_{j} \le (1 + C_{\rm od}e^{-\eta_{\rm od}t})\sum_{j} g^{jj}(t)\xi_{j}\xi_{j},$$

where we used (20.11) in the last step. Thus there is a constant C > 1 such that the first inequality in (20.12) holds. Let  $0 \le t_{\rm od} < \infty$  be such that  $C_{\rm od}e^{-\eta_{\rm od}t_{\rm od}} \le 1/2$ . On  $[0, t_{\rm od}]$ , it is clear that there is a constant C > 1 such that the right inequality in (20.12) holds; this is an immediate consequence of the fact that the  $g^{jl}$  are smooth functions and the fact that  $g^{jl}(t)$  are the components of a positive definite matrix for every t. For  $t \ge t_{\rm od}$ ,

$$g^{jl}(t)\xi_{j}\xi_{l} = \sum_{j \neq l} g^{jl}(t)\xi_{j}\xi_{l} + \sum_{j} g^{jj}(t)\xi_{j}\xi_{j} \ge \sum_{j} g^{jj}(t)\xi_{j}\xi_{j}/2,$$

where we used (20.11) and the fact that  $C_{\text{od}}e^{-\eta_{\text{od}}t} \leq 1/2$  for  $t \geq t_{\text{od}}$  in the last step. Thus (20.12) holds. Using (20.4), a similar argument yields the conclusion that there is a constant C > 1 such that

$$C^{-1} \le e^{-2\beta_j t} g^{jj}(t) \le C$$

for all  $t \ge 0$  and  $\xi \in \mathbb{R}^d$ . Combining this estimate with (20.12) yields (20.13). Since  $e^{\beta_{\text{Ri},r}t}a_r(t)$  converges to a strictly positive number, there is similarly a constant C > 1 (depending only on the coefficients of the equation), such that (20.14) holds.

Bounding the derivatives of  $\ell$ . Next, let us estimate

$$\begin{aligned} |\dot{g}^{jl}\xi_{j}\xi_{l}| &\leq \sum_{j\neq l} |\dot{g}^{jl}\xi_{j}\xi_{l}| + \sum_{j} |\dot{g}^{jj}\xi_{j}\xi_{j}| \\ &\leq Ce^{-\eta_{\text{od}}t} \sum_{j} g^{jj}(t)\xi_{j}\xi_{j} + C\sum_{j} e^{2\beta_{j}t}\xi_{j}\xi_{j} \leq Cg^{jl}\xi_{j}\xi_{l} \end{aligned}$$
(20.16)

for all  $t \ge 0$  and  $\xi \in \mathbb{R}^d$ , where we used (20.11) and the fact that  $e^{-2\beta_j t} \dot{g}^{jj}(t)$  is bounded in the second step, and (20.12) and (20.13) in the third step. A similar argument concerning  $\ddot{g}^{jl}$  yields the conclusion that there is a constant C such that

$$|\ddot{g}^{jl}\xi_j\xi_l| \le Cg^{jl}\xi_j\xi_l \tag{20.17}$$

for all  $t \ge 0$  and  $\xi \in \mathbb{R}^d$ . Appealing to (20.6) and (20.14), it can similarly be verified that

$$\left|\sum_{r=1}^{R} \partial_t (a_r^{-2}) \eta_r^2\right| + \left|\sum_{r=1}^{R} \partial_t^2 (a_r^{-2}) \eta_r^2\right| \le C \sum_{r=1}^{R} a_r^{-2} \eta_r^2 \tag{20.18}$$

for all  $\eta \in \mathbb{R}^R$ . Moreover, the constants *C* appearing in (20.16)–(20.18) only depend on the coefficients of the equation (1.2).

Let  $0 \neq \iota \in \mathcal{I}_{B}$ , define  $\mathfrak{g}(\iota, t)$  by (24.13) and let  $\ell(\iota, t) := \ln \mathfrak{g}(\iota, t)$ . Compute

$$\dot{\ell}(\iota,t) = \frac{1}{2\mathfrak{g}^2(\iota,t)} \left( \dot{g}^{jl}(t) n_j n_l + \sum_{r=1}^R \partial_t (a_r^{-2})(t) \nu_{r,i_r}^2(\iota) \right)$$
(20.19)

Due to (20.16) and (20.18), we conclude that  $|\dot{\ell}(\iota, t)|$  is bounded for  $t \ge 0$ . Since

$$\ddot{\ell}(\iota,t) = \frac{1}{2\mathfrak{g}^2(\iota,t)} \left( \ddot{g}^{jl}(t) n_j n_l + \sum_{r=1}^R \partial_t^2(a_r^{-2})(t) \nu_{r,i_r}^2(\iota) \right) - 2\dot{\ell}^2(\iota,t),$$

(20.17) and (20.18) can similarly be used to yield the conclusion that  $|\dot{\ell}(\iota, t)|$  is bounded for  $t \ge 0$ . Thus (14.1) holds for some constant  $\mathfrak{b}$  depending only on the coefficients of the equation (1.2).

Bounding the remaining coefficients. Next, let us consider (14.2). Let  $0 \neq \iota \in \mathcal{I}_{\mathrm{B}}$  and let  $\sigma$  be defined as in (10.1). Due to (20.10) and the fact that  $|\dot{\ell}|$  is bounded to the future, it follows that (14.2) holds with  $\mathfrak{f}_{\mathrm{sh}}(t) = Ce^{-\eta_{\mathrm{sh}}t}$ , for some suitably big constant C, depending only on the coefficients of the equation (1.2).

Turning to (14.3), note that if X is defined as in (10.1) and  $n \neq 0$ , then

$$\|X\| \le \frac{1}{\mathfrak{g}} \sum_{j} |n_{j}| \|X^{j}\| \le C \left( \sum_{j} e^{2\beta_{j} t} n_{j}^{2} \right)^{-1/2} \sum_{j} |n_{j}| e^{\beta_{j} t} \le C$$

for all  $t \ge 0$ , where we have used the fact that (20.8) implies that  $e^{-\beta_j t} ||X^j||$  is bounded, as well as (20.13). Since

$$\dot{X} = \frac{n_l X^l}{\mathfrak{g}} - X\dot{\ell},$$

a similar argument (using (20.8) together with the bounds for ||X|| and  $|\dot{\ell}|$  we have already derived) yields  $||\dot{X}|| \leq C$  for all  $t \geq 0$  and all  $0 \neq n \in \mathbb{Z}^d$ . Thus (14.3) holds with  $\mathfrak{f}_X = C$ , where C is a constant depending only on the coefficients of the equation (1.2).

Finally, (20.8) implies that (14.4) and (14.5) hold for all  $t \ge 0$ , with  $\mathfrak{f}_{ode}(t) = C$ , where C has the same dependence as before. Since  $\mathfrak{f}_{sh}$ ,  $\mathfrak{f}_X$  and  $\mathfrak{f}_{ode}$  are all constants times exponential functions, it is clear that (14.6) holds for a suitably chosen  $\mathfrak{b}$ .

#### 20.3 Simplifying the matrix coefficients

Consider the equation (1.2). Assume that it is diagonally dominated, balanced and convergent; cf. Definition 20.1. Let  $\bar{\beta}_1 < \bar{\beta}_2 < \cdots < \bar{\beta}_Q$  be the distinct elements of the union of the sets  $\{\beta_1, \ldots, \beta_d\}$  and  $\{\beta_{\text{Ri},1}, \ldots, \beta_{\text{Ri},R}\}$ . For  $1 \leq q \leq Q$  and  $\iota \in \mathcal{I}_B$ , let

$$\bar{\nu}_{q}(\iota) := \left( \sum_{\{j:\beta_{j}=\bar{\beta}_{q}\}} g_{\infty}^{jj} n_{j}^{2} + \sum_{\{r:\beta_{\mathrm{Ri},r}=\bar{\beta}_{q}\}} a_{r,\infty}^{-2} \nu_{r,i_{r}}^{2}(\iota) \right)^{1/2}.$$
(20.20)

In the analysis we are about to carry out, it would be convenient to replace  $\mathfrak{g}(t)$  with

$$\mathfrak{g}_{\text{diag}}(\iota, t) := \left(\sum_{j=1}^{d} g_{\infty}^{jj} n_{j}^{2} e^{2\beta_{j}t} + \sum_{r=1}^{R} a_{r,\infty}^{-2} \nu_{r,i_{r}}^{2}(\iota) e^{2\beta_{\text{Ri},r}t}\right)^{1/2} \\ = \left(\sum_{q=1}^{Q} \bar{\nu}_{q}^{2}(\iota) e^{2\bar{\beta}_{q}t}\right)^{1/2}.$$
(20.21)

It is therefore of interest to compare  $\mathfrak{g}(t)$  with  $\mathfrak{g}_{\text{diag}}(t)$ . We are also interested in approximating  $X/\mathfrak{g}$  and  $\alpha/\mathfrak{g}$  in terms of similar asymptotic data; cf. the introduction to the present chapter.

**Lemma 20.5.** Consider the equation (1.2). Assume that it is diagonally dominated, balanced and convergent; cf. Definition 20.1. Then there is a constant C > 1 such that

$$C^{-1}\mathfrak{g}(\iota,t) \le \mathfrak{g}_{\text{diag}}(\iota,t) \le C\mathfrak{g}(\iota,t)$$
(20.22)

for all  $t \geq 0$  and  $\iota \in \mathcal{I}_{\mathrm{B}}$ , where  $\mathfrak{g}_{\mathrm{diag}}$  is defined in (20.21). Moreover,

$$|\mathfrak{g}(\iota, t) - \mathfrak{g}_{\operatorname{diag}}(\iota, t)| \le C e^{-\eta_{\operatorname{diag}} t} \mathfrak{g}(\iota, t)$$
(20.23)

for all  $t \geq 0$  and  $\iota \in \mathcal{I}_{\mathrm{B}}$ , where  $\eta_{\mathrm{diag}} := \min\{\eta_{\mathrm{od}}, \eta_{\mathrm{d}}, \eta_{\mathrm{ad}}\}$ . Letting  $\ell_{\mathrm{diag}} := \ln \mathfrak{g}_{\mathrm{diag}}$ ,

$$\left|\frac{\dot{\ell}(\iota,t)}{\mathfrak{g}(\iota,t)} - \frac{\dot{\ell}_{\mathrm{diag}}(\iota,t)}{\mathfrak{g}_{\mathrm{diag}}(\iota,t)}\right| \le C \frac{e^{-\eta_{\mathrm{diag}}t}}{\mathfrak{g}(\iota,t)}$$
(20.24)

for all  $t \geq 0$  and  $0 \neq \iota \in \mathcal{I}_{B}$ . Letting

$$X_{\text{diag}}(\iota, t) := \frac{1}{\mathfrak{g}_{\text{diag}}(\iota, t)} \sum_{l} n_l e^{\beta_l t} X_{\infty}^l, \qquad (20.25)$$

308

the estimate

$$\left\|\frac{X_{\text{diag}}(\iota,t)}{\mathfrak{g}_{\text{diag}}(\iota,t)} - \frac{X(\iota,t)}{\mathfrak{g}(\iota,t)}\right\| \le C\frac{e^{-\eta_X t}}{\mathfrak{g}(\iota,t)}$$
(20.26)

holds for all  $t \ge 0$  and  $0 \ne \iota \in \mathcal{I}_{\mathrm{B}}$ , where  $\eta_X := \min\{\eta_{\mathrm{diag}}, \eta_{\mathrm{mn}}\}$ . Moreover,

$$\left\|\frac{\alpha_{\infty}}{\mathfrak{g}_{\operatorname{diag}}(\iota,t)} - \frac{\alpha(t)}{\mathfrak{g}(\iota,t)}\right\| \le C \frac{e^{-\eta_{\alpha}t}}{\mathfrak{g}(\iota,t)}$$
(20.27)

for all  $t \ge 0$  and  $0 \ne \iota \in \mathcal{I}_{\mathrm{B}}$ , where  $\eta_{\alpha} := \min\{\eta_{\mathrm{diag}}, \eta_{\mathrm{mn}}\}$ .

**Remark 20.6.** Many of the conclusions of the lemma can be obtained using weaker assumptions.

**Remark 20.7.** The constants C appearing in the statement of the lemma only depend on the coefficients of the equation (1.2).

*Proof.* That there is a constant C > 1, depending only on the coefficients of the equation (1.2), such that (20.22) holds for all  $t \ge 0$  and  $\iota \in \mathcal{I}_{B}$  is a consequence of (20.13) and (20.14). Let us estimate

$$\begin{aligned} &|\mathfrak{g}^{2}-\mathfrak{g}_{\mathrm{diag}}^{2}|\\ \leq &\sum_{j\neq l}\left|g^{jl}n_{j}n_{l}\right|+\sum_{j}e^{2\beta_{j}t}|e^{-2\beta_{j}t}g^{jj}-g^{jj}_{\infty}|n_{j}^{2}\\ &+\sum_{r=1}^{R}|e^{\beta_{\mathrm{Ri},r}t}a_{r}-a_{r,\infty}|(e^{\beta_{\mathrm{Ri},r}t}a_{r}+a_{r,\infty})[e^{\beta_{\mathrm{Ri},r}t}a_{r}]^{-2}a_{r,\infty}^{-2}\nu_{r,i_{r}}^{2}(\iota)e^{2\beta_{\mathrm{Ri},r}t}\\ \leq &Ce^{-\eta_{\mathrm{od}}t}\mathfrak{g}^{2}+Ce^{-\eta_{\mathrm{d}}t}\mathfrak{g}^{2}+Ce^{-\eta_{\mathrm{ad}}t}\mathfrak{g}^{2}\leq Ce^{-\eta_{\mathrm{diag}}t}\mathfrak{g}^{2}\end{aligned}$$

$$(20.28)$$

for all  $t \ge 0$  and  $\iota \in \mathcal{I}_{\mathrm{B}}$ , where C only depends on the coefficients of the equation (1.2) and we have appealed to (20.4), (20.6), (20.11), (20.12), (20.13) and (20.14). Assuming  $\iota \neq 0$  and dividing (20.28) with  $\mathfrak{g} + \mathfrak{g}_{\text{diag}}$  yields (20.23) (if  $\iota = 0$ , this inequality holds trivially).

Let us turn to  $\dot{\ell}/\mathfrak{g}$ . Recalling that  $\dot{\ell}$  is given by (20.19),

.

$$\left| \frac{\dot{\ell}}{\mathfrak{g}} - \frac{\dot{\ell}_{\text{diag}}}{\mathfrak{g}_{\text{diag}}} \right| \leq \frac{1}{2} \left| \frac{\dot{g}^{jl} n_j n_l}{\mathfrak{g}^3} - \frac{1}{\mathfrak{g}^3_{\text{diag}}} \sum_{j=1}^d 2\beta_j g^{jj}_{\infty} n_j^2 e^{2\beta_j t} \right|$$

$$+ \frac{1}{2} \left| \sum_{r=1}^R \left( \frac{1}{\mathfrak{g}^3} \partial_t (a_r^{-2}) - \frac{1}{\mathfrak{g}^3_{\text{diag}}} 2\beta_{\text{Ri},r} a_{r,\infty}^{-2} e^{2\beta_{\text{Ri},r} t} \right) \nu_{r,i_r}^2(t) \right|$$

$$(20.29)$$

The first term on the right hand side of (20.29) can be estimated by

$$\frac{1}{2} \frac{1}{\mathfrak{g}^{3}} \sum_{j \neq l} |\dot{g}^{jl} n_{j} n_{l}| + \frac{1}{2} \frac{1}{\mathfrak{g}^{3}} \sum_{j} e^{2\beta_{j} t} |e^{-2\beta_{j} t} \dot{g}^{jj} - 2\beta_{j} g_{\infty}^{jj} |n_{j}^{2} \\
+ \frac{1}{2} \left| \frac{1}{\mathfrak{g}^{3}} - \frac{1}{\mathfrak{g}_{\text{diag}}^{3}} \right| \sum_{j=1}^{d} g_{\infty}^{jj} n_{j}^{2} 2 |\beta_{j}| e^{2\beta_{j} t} \leq C \frac{e^{-\eta_{\text{diag}} t}}{\mathfrak{g}}$$
(20.30)

for all  $t \ge 0$ , where we appealed to (20.5), (20.11), (20.12), (20.13), (20.22) and (20.23). Before turning to the second term on the right hand side of (20.29), note that

$$\partial_t a_r^{-2} - 2\beta_{\mathrm{Ri},r} a_{r,\infty}^{-2} e^{2\beta_{\mathrm{Ri},r}t} = -2(a_r e^{\beta_{\mathrm{Ri},r}t})^{-3}(e^{\beta_{\mathrm{Ri},r}t}\dot{a}_r + \beta_{\mathrm{Ri},r}a_{r,\infty})e^{2\beta_{\mathrm{Ri},r}t} + 2\beta_{\mathrm{Ri},r}e^{2\beta_{\mathrm{Ri},r}t}a_{r,\infty}[(a_r e^{\beta_{\mathrm{Ri},r}t})^{-3} - a_{r,\infty}^{-3}].$$

Combining this observation with (20.6) and (20.7) yields

$$|\partial_t a_r^{-2} - 2\beta_{\mathrm{Ri},r} a_{r,\infty}^{-2} e^{2\beta_{\mathrm{Ri},r}t}| \le C e^{-\eta_{\mathrm{ad}}t} e^{2\beta_{\mathrm{Ri},r}t}$$

309

for all  $t \ge 0$ , where C only depends on the coefficients of the equation (1.2). Due to this estimate, the second term on the right hand side of (20.29) can be estimated by

$$\begin{split} &\frac{1}{2}\sum_{r=1}^{R}\frac{1}{\mathfrak{g}^{3}}|\partial_{t}a_{r}^{-2}-2\beta_{\mathrm{Ri},r}a_{r,\infty}^{-2}e^{2\beta_{\mathrm{Ri},r}t}|\nu_{r,i_{r}}^{2}(\iota)\\ &+\frac{1}{2}\sum_{r=1}^{R}\left|\frac{1}{\mathfrak{g}^{3}}-\frac{1}{\mathfrak{g}_{\mathrm{diag}}^{3}}\right|2|\beta_{\mathrm{Ri},r}|a_{r,\infty}^{-2}e^{2\beta_{\mathrm{Ri},r}t}\nu_{r,i_{r}}^{2}(\iota)\leq C\frac{e^{-\eta_{\mathrm{diag}}t}}{\mathfrak{g}} \end{split}$$

for all  $t \ge 0$ , where we appealed to (20.14), (20.22) and (20.23), and C only depends on the coefficients of the equation (1.2). Combining this estimate with (20.29) and (20.30) yields (20.24). Next, let us turn to  $X/\mathfrak{g}$  and  $\alpha/\mathfrak{g}$ . To begin with

$$\begin{aligned} \|X_{\text{diag}} - X\| &\leq \frac{1}{\mathfrak{g}_{\text{diag}}} \sum_{l} |n_l| e^{\beta_l t} \|e^{-\beta_l t} X^l - X_{\infty}^l\| + \left|\frac{1}{\mathfrak{g}_{\text{diag}}} - \frac{1}{\mathfrak{g}}\right| \|n_l X^l\| \\ &\leq C e^{-\eta_{\text{mn}} t} + C e^{-\eta_{\text{diag}} t} \|X\| \leq C e^{-\eta_X t} \end{aligned}$$

for all  $t \ge 0$  and  $\iota \in \mathcal{I}_{\mathrm{B}}$ , where we have appealed to (20.8), (20.9), (20.13), (20.22) and (20.23). Combining this estimate with (20.22) and (20.23) yields (20.26). Moreover, C only depends on the coefficients of the equation (1.2). Estimate

$$\left\|\frac{\alpha_{\infty}}{\mathfrak{g}_{\mathrm{diag}}} - \frac{\alpha}{\mathfrak{g}}\right\| \leq \frac{1}{\mathfrak{g}_{\mathrm{diag}}} \|\alpha - \alpha_{\infty}\| + \left|\frac{1}{\mathfrak{g}_{\mathrm{diag}}} - \frac{1}{\mathfrak{g}}\right| \|\alpha\|.$$

Appealing to (20.8), (20.9), (20.22) and (20.23) yields (20.27).

### 20.4 Choosing appropriate time intervals; defining the frequency eras

In order to be able to estimate the matrix products of interest, it is convenient to divide  $[0, \infty)$  into intervals during which the coefficients of the iteration can be simplified. Specifying such intervals and deriving preliminary information concerning the matrix coefficients in them is the main purpose of the next lemma.

**Lemma 20.8.** Consider the equation (1.2). Assume that it is diagonally dominated, balanced and convergent; cf. Definition 20.1. Fix  $0 \neq \iota \in \mathcal{I}_{B}$ . Using the notation introduced in connection with (20.20), let  $I_{q}$  denote the subset of  $[0, \infty)$  such that

$$\bar{\nu}_q(\iota)e^{\beta_q t} \ge \bar{\nu}_p(\iota)e^{\beta_p t} \tag{20.31}$$

for all p = 1, ..., Q. Then  $I_q$  is either empty or a closed interval. Moreover, if  $q, p \in \{1, ..., Q\}$  are such that  $q \neq p$ , then  $I_q \cap I_p$  is either empty or consists of one point. Assuming  $I_q$  to be non-empty, define

$$\mathfrak{H}_{q}^{(s)}(\iota,t) := \sum_{p \neq q} \left( \frac{\bar{\nu}_{p}(\iota) e^{\bar{\beta}_{p}t}}{\bar{\nu}_{q}(\iota) e^{\bar{\beta}_{q}t}} \right)^{s}, \quad \mathfrak{H}_{q}(\iota,t) := \mathfrak{H}_{q}^{(2)}(\iota,t), \tag{20.32}$$

where  $0 < s \in \mathbb{R}$ . Then there is a constant C > 1 such that

$$C^{-1}\mathfrak{g}(\iota,t) \le \bar{\nu}_q(\iota)e^{\beta_q t} \le C\mathfrak{g}(\iota,t) \tag{20.33}$$

for all  $t \in I_q$ . Moreover,

$$\left|\frac{1}{\bar{\nu}_q(\iota)e^{\bar{\beta}_q t}} - \frac{1}{\mathfrak{g}_{\text{diag}}(\iota, t)}\right| \leq \frac{C}{\mathfrak{g}(\iota, t)}\mathfrak{H}_q(\iota, t),$$
(20.34)

$$\left|\frac{1}{\bar{\nu}_q^2(\iota)e^{2\bar{\beta}_q t}} - \frac{1}{\mathfrak{g}_{\text{diag}}^2(\iota, t)}\right| \le \frac{C}{\mathfrak{g}^2(\iota, t)}\mathfrak{H}_q(\iota, t),$$
(20.35)

$$\left|\frac{\dot{\ell}_{\text{diag}}(\iota,t)}{\mathfrak{g}_{\text{diag}}(\iota,t)} - \frac{\bar{\beta}_{q}}{\bar{\nu}_{q}(\iota)e^{\bar{\beta}_{q}t}}\right| \leq \frac{C}{\mathfrak{g}(\iota,t)}\mathfrak{H}_{q}(\iota,t)$$
(20.36)

for all  $t \in I_q$ . Let

$$\bar{X}_{\text{diag}}^q(\iota) := \frac{1}{\bar{\nu}_q(\iota)} \sum_{\{j:\beta_j = \bar{\beta}_q\}} n_j X_{\infty}^j; \qquad (20.37)$$

recall the notation introduced at the beginning of Section 20.3. Then

$$\left\|\frac{\bar{X}_{\text{diag}}^{q}(\iota)}{\bar{\nu}_{q}(\iota)e^{\bar{\beta}_{q}t}} - \frac{X_{\text{diag}}(\iota,t)}{\mathfrak{g}_{\text{diag}}(\iota,t)}\right\| \leq \frac{C}{\mathfrak{g}(\iota,t)}\mathfrak{H}_{q}(\iota,t) + \frac{C}{\mathfrak{g}^{2}(\iota,t)}\sum_{\{j:\beta_{j}\neq\bar{\beta}_{q}\}}|n_{j}|e^{\beta_{j}t}$$
(20.38)

for  $t \in I_q$ , where  $X_{\text{diag}}$  is defined in (20.25). Moreover,

$$\left\|\frac{\alpha_{\infty}}{\bar{\nu}_{q}(\iota)e^{\bar{\beta}_{q}t}} - \frac{\alpha_{\infty}}{\mathfrak{g}_{\text{diag}}(\iota, t)}\right\| \leq \frac{C}{\mathfrak{g}(\iota, t)}\mathfrak{H}_{q}(\iota, t)$$
(20.39)

for  $t \in I_q$ .

**Remark 20.9.** The constants C appearing in the statement of the lemma only depend on the coefficients of the equation (1.2).

*Proof.* Fix  $q \in \{1, \ldots, Q\}$  and assume  $I_q$  to be non-empty. Let  $t_0, t_1 \in I_q$  with  $t_0 \leq t_1$ . Then  $\bar{\nu}_q(\iota)e^{\bar{\beta}_q t}$  is greater than  $\bar{\nu}_p(\iota)e^{\bar{\beta}_p t}$  for all p < q and  $t \geq t_0$ . Moreover,  $\bar{\nu}_q(\iota)e^{\bar{\beta}_q t}$  is greater than  $\bar{\nu}_p(\iota)e^{\bar{\beta}_p t}$  for all p > q and  $t \leq t_1$ . Thus the line connecting  $t_0$  and  $t_1$  is contained in  $I_q$ , so that  $I_q$  is connected. Since  $I_q$  is closed by definition, we conclude that  $I_q$  is a closed interval. By these arguments, the statement concerning the intersection of the different  $I_q$ 's also follows.

In  $I_q$ , it is of interest to write  $\mathfrak{g}_{\text{diag}}$  as

$$\mathfrak{g}_{\text{diag}}(\iota,t) = \bar{\nu}_q(\iota) e^{\bar{\beta}_q t} \mathfrak{p}_q(\iota,t); \quad \mathfrak{p}_q(\iota,t) := [1 + \mathfrak{H}_q(\iota,t)]^{1/2};$$

cf. (20.32). Note that

$$\bar{\nu}_q(\iota)e^{\bar{\beta}_q t} \leq \mathfrak{g}_{\text{diag}}(\iota, t) \leq Q^{1/2}\bar{\nu}_q(\iota)e^{\bar{\beta}_q t}$$

in  $I_q$ . Combining this estimate with (20.22) it is clear that  $\bar{\nu}_q(\iota)e^{\bar{\beta}_q t}$  and  $\mathfrak{g}(t)$  are equivalent in  $I_q$ . Thus (20.33) holds, and the constant C only depends on the coefficients of the equation (1.2). Let us compute

$$\frac{1}{\bar{\nu}_q(\iota)e^{\bar{\beta}_q t}} - \frac{1}{\mathfrak{g}_{\text{diag}}} = \frac{1}{\bar{\nu}_q(\iota)e^{\bar{\beta}_q t}} \frac{\mathfrak{p}_q - 1}{\mathfrak{p}_q} = \frac{1}{\bar{\nu}_q(\iota)e^{\bar{\beta}_q t}} \frac{\mathfrak{p}_q^2 - 1}{\mathfrak{p}_q(\mathfrak{p}_q + 1)}.$$

In particular,

$$\left|\frac{1}{\bar{\nu}_q(\iota)e^{\bar{\beta}_q t}} - \frac{1}{\mathfrak{g}_{\text{diag}}(\iota, t)}\right| \le \frac{1}{2} \frac{1}{\bar{\nu}_q(\iota)e^{\bar{\beta}_q t}} \mathfrak{H}_q(\iota, t)$$
(20.40)

for  $t \in I_q$ , so that (20.34) holds (where C has the same dependence as before). As a consequence,

$$\left|\frac{1}{\bar{\nu}_q^2(\iota)e^{2\bar{\beta}_q t}} - \frac{1}{\mathfrak{g}_{\text{diag}}^2(\iota, t)}\right| \le \frac{1}{\bar{\nu}_q^2(\iota)e^{2\bar{\beta}_q t}}\mathfrak{H}_q(\iota, t)$$
(20.41)

for  $t \in I_q$ , so that (20.35) holds. Again, the constant C only depends on the coefficients of the equation (1.2). Next, let us turn to

$$\dot{\ell}_{\text{diag}} = \frac{1}{\mathfrak{g}_{\text{diag}}^2} \sum_{p=1}^Q \bar{\beta}_p \bar{\nu}_p^2(\iota) e^{2\bar{\beta}_p t} = \bar{\beta}_q + \frac{1}{\mathfrak{p}_q^2} \sum_{p \neq q} (\bar{\beta}_p - \bar{\beta}_q) \left( \frac{\bar{\nu}_p(\iota) e^{\bar{\beta}_p t}}{\bar{\nu}_q(\iota) e^{\bar{\beta}_q t}} \right)^2;$$

recall (20.21). Thus

$$\left|\dot{\ell}_{\text{diag}}(\iota,t) - \bar{\beta}_q\right| \le \max_p |\bar{\beta}_p - \bar{\beta}_q| \cdot \mathfrak{H}_q(\iota,t).$$

Thus

$$\frac{\dot{\ell}_{\mathrm{diag}}(\iota,t)}{\mathfrak{g}_{\mathrm{diag}}(\iota,t)} - \frac{\bar{\beta}_q}{\mathfrak{g}_{\mathrm{diag}}(\iota,t)} \right| \leq \frac{\max_p |\bar{\beta}_p - \bar{\beta}_q|}{\bar{\nu}_q(\iota)e^{\bar{\beta}_q t}} \mathfrak{H}_q(\iota,t).$$

Combining this estimate with (20.40) yields

$$\left|\frac{\dot{\ell}_{\text{diag}}(\iota,t)}{\mathfrak{g}_{\text{diag}}(\iota,t)} - \frac{\bar{\beta}_q}{\bar{\nu}_q(\iota)e^{\bar{\beta}_q t}}\right| \le \frac{1}{\bar{\nu}_q(\iota)e^{\bar{\beta}_q t}} \left(\frac{|\bar{\beta}_q|}{2} + \max_p |\bar{\beta}_p - \bar{\beta}_q|\right) \mathfrak{H}_q(\iota,t).$$
(20.42)

Thus (20.36) holds, where the constant C only depends on the coefficients of the equation (1.2). Next, let us turn to  $X/\mathfrak{a}$ . Note that the norm of  $X_{Y}$  is bounded by a constant depending only

Next, let us turn to  $X/\mathfrak{g}$ . Note that the norm of  $X_{\text{diag}}$  is bounded by a constant depending only on the coefficients of the equation (1.2). Moreover,

$$\begin{split} X_{\text{diag}}(\iota,t) - \bar{X}_{\text{diag}}^{q}(\iota) &= \frac{1}{\mathfrak{g}_{\text{diag}}(\iota,t)} \sum_{l} n_{l} e^{\beta_{l} t} X_{\infty}^{l} - \frac{1}{\bar{\nu}_{q}(\iota)} \sum_{\{j:\beta_{j} = \bar{\beta}_{q}\}} n_{j} X_{\infty}^{j} \\ &= \left(\frac{1}{\mathfrak{g}_{\text{diag}}(\iota,t)} - \frac{1}{\bar{\nu}_{q}(\iota) e^{\bar{\beta}_{q} t}}\right) \sum_{l} n_{l} e^{\beta_{l} t} X_{\infty}^{l} \\ &+ \frac{1}{\bar{\nu}_{q}(\iota) e^{\bar{\beta}_{q} t}} \sum_{\{j:\beta_{j} \neq \bar{\beta}_{q}\}} n_{j} e^{\beta_{j} t} X_{\infty}^{j}. \end{split}$$

Combining this equality with (20.22), (20.33) and (20.34) yields

$$\left\| X_{\text{diag}}(\iota,t) - \bar{X}_{\text{diag}}^q(\iota) \right\| \leq C \mathfrak{H}_q(\iota,t) + \frac{C}{\mathfrak{g}(\iota,t)} \sum_{\{j:\beta_j \neq \bar{\beta}_q\}} |n_j| e^{\beta_j t}$$

for  $t \in I_q$ . Combining this estimate in its turn with (20.33), (20.34) and the previously mentioned bound on the norm of  $X_{\text{diag}}$  yields (20.38). Moreover, the constant C only depends on the coefficients of the equation (1.2). Finally, (20.39) is an immediate consequence of (20.34). The lemma follows.

The intervals  $I_q$  defined in the previous lemma are important in the analysis to follow. It is therefore convenient to introduce the following terminology.

**Definition 20.10.** Given assumptions and terminology as in the statement of Lemma 20.8, let  $I_q$ ,  $q = 1, \ldots, Q$ , be defined as in the statement of that lemma. If  $I_q$  is non-empty, then  $I_q$  is referred to as a *frequency era* of the equation (1.2) corresponding to  $0 \neq \iota \in \mathcal{I}_B$ . If  $\bar{\beta}_q < 0$ , then  $I_q$  is called a *silent era*; if  $\bar{\beta}_q = 0$ , then  $I_q$  is called a *transparent era*; and if  $\bar{\beta}_q > 0$ , then  $I_q$  is called a *noisy era*.

**Remark 20.11.** If the equation and  $\iota$  are understood from the context, we simply speak of a frequency era.

#### 20.5 Estimating sums in a frequency era

When calculating the matrix products appearing in (15.64), it is convenient to restrict one's attention to a frequency era; cf. Definition 20.10. In that context, the following lemma is of interest.

**Lemma 20.12.** Consider the equation (1.2). Assume that it is diagonally dominated, balanced and convergent; cf. Definition 20.1. Fix  $0 \neq \iota \in \mathcal{I}_B$  and let  $I_q$  be a corresponding frequency era; cf. Definition 20.10. Consider a sequence  $\{t_k\}$  defined as in Definition 15.5, assume that there are elements of the sequence belonging to  $I_q$ , let  $k_{q,1}$  be the smallest and  $k_{q,2}$  the largest k such that  $t_k \in I_q$ , and assume that  $k_{q,1} < k_{q,2}$  (if there is no upper bound on the k's for which  $t_k \in I_q$ ,  $k_{q,2}$  is defined to equal  $\infty$ ). Let  $k_0$  and  $k_1$  be such that  $k_{q,1} \leq k_0 \leq k_1 < k_{q,2}$  and define  $\bar{t}_0$  and  $\bar{t}_1$ by  $\bar{t}_0 = t_{k_0}$  and  $\bar{t}_1 = t_{k_1+1}$ . Then

$$\left| \sum_{k=k_{0}}^{k_{1}} \frac{2\pi e^{\lambda t_{k}}}{(\bar{\nu}_{q}(\iota)e^{\bar{\beta}_{q}t_{k}})^{a}} - \int_{\bar{t}_{0}}^{\bar{t}_{1}} \frac{e^{\lambda t}}{(\bar{\nu}_{q}(\iota)e^{\bar{\beta}_{q}t})^{a-1}} dt \right|$$

$$\leq \int_{\bar{t}_{0}}^{\bar{t}_{1}} \left( \frac{Ce^{\lambda t}}{(\bar{\nu}_{q}(\iota)e^{\bar{\beta}_{q}t})^{a-1}} \left[ e^{-\eta_{\text{diag}}t} + \mathfrak{H}_{q}(\iota,t) \right] + \frac{Ce^{\lambda t}}{(\bar{\nu}_{q}(\iota)e^{\bar{\beta}_{q}t})^{a}} \right) dt,$$
(20.43)

where a and  $\lambda$  are real numbers and the constant C only depends on the coefficients of the equation (1.2) and the numbers  $\lambda$  and a. Moreover,

$$\sum_{k=k_0}^{k_1} \frac{2\pi e^{\lambda t_k}}{(\bar{\nu}_q(\iota)e^{\bar{\beta}_q t_k})^a} \le C \int_{\bar{t}_0}^{\bar{t}_1} \frac{e^{\lambda t}}{(\bar{\nu}_q(\iota)e^{\bar{\beta}_q t})^{a-1}} dt,$$
(20.44)

$$\sum_{k=k_0}^{k_1} \frac{2\pi e^{\lambda t_k}}{(\bar{\nu}_q(\iota)e^{\bar{\beta}_q t_k})^a} \mathfrak{H}_q^{(r)}(\iota, t_k) \le C \int_{\bar{t}_0}^{\bar{t}_1} \frac{e^{\lambda t}}{(\bar{\nu}_q(\iota)e^{\bar{\beta}_q t})^{a-1}} \mathfrak{H}_q^{(r)}(\iota, t) dt,$$
(20.45)

where a,  $\lambda$  and r > 0 are real numbers and the constants C only depend on the coefficients of the equation (1.2) and the numbers  $\lambda$ , a and r.

*Proof.* Assume that  $[t_k, t_{k+1}] \subseteq I_q$ . Let  $\lambda \in \mathbb{R}$  and estimate

$$|e^{\lambda t} - e^{\lambda s}| \le e^{\lambda u} |e^{\lambda(t-u)} - e^{\lambda(s-u)}| \le C \frac{e^{\lambda u}}{\bar{\nu}_q(\iota) e^{\bar{\beta}_q u}}$$
(20.46)

for  $s, t, u \in [t_k, t_{k+1}]$ , where we appealed to (14.23), (14.24), (20.33) and the fact that  $\mathfrak{g}(\iota, t_k) \geq 1$ . Note also that C only depends on  $\lambda$  and the coefficients of the equation (1.2). By a similar argument, there is a constant  $C_{\lambda} > 1$ , depending only on  $\lambda$ , such that

$$C_{\lambda}^{-1}e^{\lambda s} \le e^{\lambda t} \le C_{\lambda}e^{\lambda s} \tag{20.47}$$

for  $s, t \in [t_k, t_{k+1}]$ . In addition, (20.46) yields

$$\left|\bar{\nu}_q(\iota)e^{\bar{\beta}_q t} - \bar{\nu}_q(\iota)e^{\bar{\beta}_q t_k}\right| \le C$$

for  $t \in [t_k, t_{k+1}]$ , where C only depends on the coefficients of the equation (1.2). By a similar argument,

$$|\mathfrak{H}_q(\iota,t) - \mathfrak{H}_q(\iota,s)| \le \frac{C}{\bar{\nu}_q(\iota)e^{\bar{\beta}_q u}}\mathfrak{H}_q(\iota,u)$$
(20.48)

for all  $s, t, u \in [t_k, t_{k+1}]$ , where C only depends on the coefficients of the equation (1.2). Note, in particular, that  $\mathfrak{H}_q(\iota, t)$  and  $\mathfrak{H}_q(\iota, s)$  are equivalent for  $s, t \in [t_k, t_{k+1}]$ ; recall that  $\bar{\nu}_q(\iota)e^{\bar{\beta}_q t}$  and  $\mathfrak{g}(\iota, t)$  are equivalent due to (20.33) and that  $\mathfrak{g}(\iota, t) \geq 1$  in  $[t_k, t_{k+1}]$  due to Lemma 14.4. Due to (14.23), (14.25), (20.23), (20.34) and the fact that  $\mathfrak{g}(\iota, t)$ ,  $\mathfrak{g}_{\text{diag}}(\iota, t)$  and  $\bar{\nu}_q(\iota)e^{\beta_q t}$  are equivalent in  $I_q$ 

$$\left|\mathfrak{g}(\iota,t) - \bar{\nu}_q(\iota)e^{\bar{\beta}_q t_k}\right| \le C\mathfrak{g}(\iota,t_k)[e^{-\eta_{\mathrm{diag}}t_k} + \mathfrak{H}_q(\iota,t_k)] + C$$
(20.49)

for all  $t \in [t_k, t_{k+1}]$ , where C only depends on the coefficients of the equation (1.2). For future reference, it is of interest to keep in mind that we could replace the  $t_k$  appearing on the right hand side of (20.49) with any  $s \in [t_k, t_{k+1}]$ ; this is a consequence of (14.23), (20.47) and (20.48). On the other hand,

$$2\pi = \int_{t_k}^{t_{k+1}} \mathfrak{g}(\iota, t) dt = \int_{t_k}^{t_{k+1}} [\mathfrak{g}(\iota, t) - \bar{\nu}_q(\iota) e^{\bar{\beta}_q t_k}] dt + (t_{k+1} - t_k) \bar{\nu}_q(\iota) e^{\bar{\beta}_q t_k}.$$
 (20.50)

Combining (14.24), (20.33), (20.49) and (20.50) yields

$$\left| t_{k+1} - t_k - \frac{2\pi}{\bar{\nu}_q(\iota)e^{\bar{\beta}_q t_k}} \right| \le \frac{C}{\mathfrak{g}(\iota, t_k)} \left[ e^{-\eta_{\text{diag}} t_k} + \mathfrak{H}_q(\iota, t_k) \right] + \frac{C}{\mathfrak{g}^2(\iota, t_k)}, \tag{20.51}$$

where C has the same dependence as before. For future reference, it is of interest to note that  $t_k$  can be replaced by  $t_{k+1}$  on the right hand side and in the third term inside the absolute value sign on the left hand side. One particular consequence of (20.51) is that

$$\begin{aligned} & \left| \frac{2\pi e^{\lambda t_k}}{(\bar{\nu}_q(\iota)e^{\bar{\beta}_q t_k})^a} - \frac{e^{\lambda t_k}}{(\bar{\nu}_q(\iota)e^{\bar{\beta}_q t_k})^{a-1}}(t_{k+1} - t_k) \right| \\ \leq & \frac{Ce^{\lambda t_k}}{(\bar{\nu}_q(\iota)e^{\bar{\beta}_q t_k})^a} \left[ e^{-\eta_{\text{diag}} t_k} + \mathfrak{H}_q(\iota, t_k) \right] + \frac{Ce^{\lambda t_k}}{(\bar{\nu}_q(\iota)e^{\bar{\beta}_q t_k})^{a+1}} \end{aligned}$$

where we appealed to (20.33) and C only depends on the coefficients of the equation (1.2). On the other hand,

$$\left| \int_{t_k}^{t_{k+1}} \frac{e^{\lambda t}}{(\bar{\nu}_q(\iota)e^{\bar{\beta}_q t})^{a-1}} dt - \frac{e^{\lambda t_k}}{(\bar{\nu}_q(\iota)e^{\bar{\beta}_q t_k})^{a-1}} (t_{k+1} - t_k) \right| \le \frac{Ce^{\lambda t_k}}{(\bar{\nu}_q(\iota)e^{\bar{\beta}_q t_k})^{a+1}}$$

where we appealed to (14.24), (20.33) and (20.46), and C only depends on a,  $\lambda$  and the coefficients of the equation (1.2). Combining these two estimates with (14.24), (20.33) and the previously mentioned equivalences yields

$$\begin{split} & \left| \frac{2\pi e^{\lambda t_k}}{(\bar{\nu}_q(\iota)e^{\bar{\beta}_q t_k})^a} - \int_{t_k}^{t_{k+1}} \frac{e^{\lambda t}}{(\bar{\nu}_q(\iota)e^{\bar{\beta}_q t})^{a-1}} dt \right| \\ \leq & \int_{t_k}^{t_{k+1}} \left( \frac{Ce^{\lambda t}}{(\bar{\nu}_q(\iota)e^{\bar{\beta}_q t})^{a-1}} \left[ e^{-\eta_{\text{diag}} t} + \mathfrak{H}_q(\iota, t) \right] + \frac{Ce^{\lambda t}}{(\bar{\nu}_q(\iota)e^{\bar{\beta}_q t})^a} \right) dt, \end{split}$$

an estimate which implies (20.43). Note also that C only depends on a,  $\lambda$  and the coefficients of the equation (1.2). The estimates (20.44) and (20.45) follow by similar, but simpler, arguments.

Next, it is of interest to compute the integrals appearing in (20.43). Before doing so, let us introduce the following terminology.

**Definition 20.13.** Consider the equation (1.2). Assume that it is diagonally dominated, balanced and convergent; cf. Definition 20.1. Fix  $0 \neq \iota \in \mathcal{I}_{B}$  and let  $I_q$  be a corresponding frequency era; cf. Definition 20.10. Let  $\bar{\beta}_q$  and  $\bar{\nu}_q(\iota)$  be defined as at the beginning of Section 20.3. Given  $\lambda, a \in \mathbb{R}$ and  $t \in \mathbb{R}$ , define  $\lambda_{a,q}$  and  $I_{a,q}(\lambda, t)$  by

$$\lambda_{a,q} := \lambda - (a-1)\bar{\beta}_q, \quad I_{a,q}(\lambda,t) := \frac{1}{\bar{\nu}_q^{a-1}(\iota)} \begin{cases} t & \lambda_{a,q} = 0, \\ \lambda_{a,q}^{-1} e^{\lambda_{a,q}t} & \lambda_{a,q} \neq 0. \end{cases}$$

#### 20.5. ESTIMATING SUMS IN A FREQUENCY ERA

Computing the integrals. With the notation of Definition 20.13,

$$\int_{\bar{t}_0}^{\bar{t}_1} \frac{e^{\lambda t}}{(\bar{\nu}_q(\iota)e^{\bar{\beta}_q t})^{a-1}} dt = I_{a,q}(\lambda, \bar{t}_1) - I_{a,q}(\lambda, \bar{t}_0).$$

Most of the integrals appearing in Lemma 20.12 can be computed directly using this formula (with various values of  $\lambda$  and a). However, it is of interest to consider a generalised version of the integral involving  $\mathfrak{H}_q$ . Let  $0 < r \in \mathbb{R}$  and recall the notation introduced in (20.32). Then

$$\int_{\bar{t}_{0}}^{\bar{t}_{1}} \frac{e^{\lambda t}}{(\bar{\nu}_{q}(\iota)e^{\bar{\beta}_{q}t})^{a-1}} \mathfrak{H}_{q}^{(r)}(t) dt 
= \sum_{p \neq q} \bar{\nu}_{p}^{r}(\iota) \left[ I_{a+r,q}(\lambda + r\bar{\beta}_{p}, \bar{t}_{1}) - I_{a+r,q}(\lambda + r\bar{\beta}_{p}, \bar{t}_{0}) \right].$$
(20.52)

Assume first that  $(\lambda + r\bar{\beta}_p)_{a+r,q} \neq 0$  for all  $p \neq q$ . Then

$$\sum_{p \neq q} \bar{\nu}_{p}^{r}(\iota) I_{a+r,q}(\lambda + r\bar{\beta}_{p}, t) = \sum_{p \neq q} \frac{1}{(\lambda + r\bar{\beta}_{p})_{a+r,q}} \frac{\bar{\nu}_{p}^{r}(\iota)}{\bar{\nu}_{q}^{r}(\iota)} \frac{1}{\bar{\nu}_{q}^{a-1}(\iota)} e^{(\lambda + r\bar{\beta}_{p})_{a+r,q}t}$$

$$= \sum_{p \neq q} \frac{1}{(\lambda + r\bar{\beta}_{p})_{a+r,q}} \frac{\bar{\nu}_{p}^{r}(\iota) e^{r\bar{\beta}_{p}t}}{\bar{\nu}_{q}^{r}(\iota) e^{r\bar{\beta}_{q}t}} \frac{1}{(\bar{\nu}_{q}(\iota) e^{\bar{\beta}_{q}t})^{a-1}} e^{\lambda t}.$$
(20.53)

Second, assume that there is one p, say s, such that  $(\lambda + r\bar{\beta}_p)_{a+r,q} = 0$  (note that there cannot be more than one such p, since  $(\lambda + r\bar{\beta}_p)_{a+r,q}$  is strictly increasing with p). Then

$$\sum_{p \neq q} \bar{\nu}_{p}^{r}(\iota) I_{a+r,q}(\lambda + r\bar{\beta}_{p}, t) = \sum_{p \neq q,s} \frac{1}{(\lambda + r\bar{\beta}_{p})_{a+r,q}} \frac{\bar{\nu}_{p}^{r}(\iota) e^{r\beta_{p}t}}{\bar{\nu}_{q}^{r}(\iota) e^{r\bar{\beta}_{q}t}} \frac{1}{(\bar{\nu}_{q}(\iota) e^{\bar{\beta}_{q}t})^{a-1}} e^{\lambda t} + \frac{\bar{\nu}_{s}^{r}(\iota) e^{r\bar{\beta}_{s}t}}{\bar{\nu}_{q}^{r}(\iota) e^{r\bar{\beta}_{q}t}} \frac{1}{(\bar{\nu}_{q}(\iota) e^{\bar{\beta}_{q}t})^{a-1}} t e^{\lambda t}.$$
(20.54)
## Chapter 21

# Non-degenerate, diagonally dominated, convergent and balanced equations

#### 21.1 Introduction

In the present chapter, we consider non-degenerate, diagonally dominated, convergent and balanced equations. The notion of a diagonally dominated, balanced and convergent equation is introduced in Definition 20.1. The requirement of non-degeneracy amounts to demanding that the set consisting of the  $\beta_j$ , j = 1, ..., d, and the  $\beta_{\text{Ri},r}$ , r = 1, ..., R, contain d + R distinct elements; cf. Definition 21.1 below. For solutions to such equations, we are here interested in estimating how the energies of the associated modes evolve over time; in later chapters, we estimate the energy of the entire solution. However, even when considering a single mode, there are normally several frequency eras to take into consideration; cf. Definition 20.10. Moreover, how the energy evolves typically depends on the frequency era. This leads us to divide the analysis into the following parts.

Estimating the energy of a mode during one frequency era. The basic tool for estimating the energy associated with a mode is the equality (15.63). Iterating this equality leads to matrix products of the type appearing in (15.64). Our first goal is to estimate such matrix products. Since the result depends on the frequency era to which the corresponding elements of the time sequence belong, we start, in Section 21.2, to fix one frequency era, say  $I_q$ , and to estimate the norm of matrix products such that each factor corresponds to an element in the time sequence belonging to  $I_q$ . The relevant result is Lemma 21.7. In the end, it turns out that, assuming  $\bar{\beta}_q \neq 0$ , the estimates we obtain are essentially optimal; we demonstrate this in Lemma 21.31. However, when  $\bar{\beta}_q = 0$ , there is room for improvement. We therefore treat this case separately in Section 21.3. When  $\bar{\beta}_q = 0$ , it is natural to divide the analysis into two cases. First, there is the case that  $\bar{\nu}_q(\iota)$  is large. Then the conclusions of Lemma 21.7 are satisfactory. When  $\bar{\nu}_q(\iota)$  is small, basing an analysis on equalities such as (15.63) is, however, not a good idea. In fact, it is better to consider the relevant system of differential equations directly. We derive the corresponding estimates in Lemma 21.11 below.

**Estimating the energy of a mode during several frequency eras.** Given the results of Lemma 21.7 and the refined analysis in the case of transparent eras, it is possible to draw conclusions concerning how the energy evolves over several frequency eras. This is the subject of Section 21.4. In fact, the main result of the present chapter is Corollary 21.17 of Section 21.4. Even though the corresponding estimates are applicable to time intervals containing several frequency

eras, they are, however, still restricted to intervals corresponding to a time sequence of the type introduced in Definition 15.5. In other words, the estimates only apply in situations where the behaviour is oscillatory. On the other hand, the non-oscillatory setting can be dealt with using estimates of the type derived in Chapters 9 and 11. In later chapters, we combine the results relevant in the oscillatory setting with the ones relevant in the non-oscillatory setting in order to obtain a complete picture for one fixed mode  $\iota$  and all future times. Combining the corresponding estimates then yields a complete picture for solutions to (1.2).

Improved estimates for unbounded frequency eras. For  $\bar{\beta}_q \neq 0$ , the estimates derived in Lemma 21.7 are essentially optimal. In fact, the only problem that might occur is that there might be non-trivial Jordan blocks of certain matrices; this leads to an  $\epsilon$ -loss in the estimates. However, for frequency eras of the form  $I_q = [t_0, \infty)$ , this loss can be removed even in the presence of non-trivial Jordan blocks. This is the subject of Section 21.5.

**Optimality.** Finally, in Section 21.6, we turn to the question of optimality. In case  $\beta_q = 0$ , this issue is settled in Chapters 11 and 12. In case  $\bar{\beta}_q \neq 0$ , we demonstrate that the estimates derived in Lemma 21.7 concerning the matrix products are optimal.

#### 21.2 Mode analysis in one frequency era

In the present chapter, we focus on non-degenerate equations, a notion we now define.

**Definition 21.1.** Consider the equation (1.2). Assume that it is diagonally dominated, balanced and convergent; cf. Definition 20.1. If the set consisting of the  $\beta_j$ ,  $j = 1, \ldots, d$ , and the  $\beta_{\text{Ri},r}$ ,  $r = 1, \ldots, R$ , contains d + R distinct elements, then (1.2) is said to be *non-degenerate*, *diagonally* dominated, balanced and convergent.

The following objects will play a central role in the analysis to follow.

**Definition 21.2.** If (1.2) is non-degenerate, diagonally dominated, balanced and convergent, then Q = d + R, using the notation introduced at the beginning of Section 20.3. Given  $q \in \{1, \ldots, Q\}$ , define  $\tilde{X}^q_{\infty}$  as follows. If  $\bar{\beta}_q = \beta_j$  for some  $j \in \{1, \ldots, d\}$ , then j is uniquely determined by q and

$$\tilde{X}^{q}_{\infty} := (g^{jj}_{\infty})^{-1/2} X^{j}_{\infty}.$$
(21.1)

If  $\bar{\beta}_q = \beta_{\mathrm{Ri},r}$  for some  $r \in \{1, \ldots, R\}$ , then r is uniquely determined by q and  $\tilde{X}^q_{\infty} := 0$ . For  $q \in \{1, \ldots, Q\}$ , let

$$R_{q,\pm}^{+} := \pm \frac{1}{2} (-\alpha_{\infty} + \bar{\beta}_{q} \mathrm{Id}_{m} + \tilde{X}_{\infty}^{q}), \quad R_{q,\pm}^{-} := \pm \frac{1}{2} (-\alpha_{\infty} + \bar{\beta}_{q} \mathrm{Id}_{m} - \tilde{X}_{\infty}^{q}), \tag{21.2}$$

and let

$$\kappa_q := \operatorname{Rsp}\left(\operatorname{diag}\{R_{q,+}^+, R_{q,+}^-\}\right), \quad \kappa_{q,\pm} := \max\{\kappa_{\max}(R_{q,\pm}^+), \kappa_{\max}(R_{q,\pm}^-)\}, \quad (21.3)$$

$$d_{q,\pm} := d_{\max}(\operatorname{diag}\{R_{q,\pm}^+, R_{q,\pm}^-\}, \kappa_{q,\pm});$$
(21.4)

cf. Definition 4.3. Finally, let

$$\kappa_{\rm rsp} := \max_{q} \kappa_q, \quad \kappa_{\pm} := \max_{q} \kappa_{q,\pm}, \tag{21.5}$$

**Remark 21.3.** It is of interest to keep in mind that  $\kappa_q = \kappa_{q,+} + \kappa_{q,-}$ .

Time sequences of the type introduced in Definition 15.5 will play a central role in the analysis to follow. However, for technical reasons we need to restrict them slightly.

**Definition 21.4.** Consider the equation (1.2). Assume that it is non-degenerate, diagonally dominated, balanced and convergent; cf. Definition 21.1. Let  $0 \neq \iota \in \mathcal{I}_B$  and assume  $t_0 \in [0, \infty)$  to be such that (14.22) holds with  $t_a$  replaced by  $t_0$ . Assume, moreover, that

$$\max_{\alpha} \bar{\nu}_q(\iota) e^{\beta_q t_0} \ge 2\pi (\kappa_{\rm rsp} + 1), \tag{21.6}$$

where  $\kappa_{rsp}$  is defined by (21.5). Assume that  $t_k \geq t_0$  has been defined for some  $0 \leq k \in \mathbb{Z}$  and that (14.22) and (21.6) hold with  $t_a$  and  $t_0$  replaced by  $t_k$ . Then  $t_{k+1}$  is defined by the condition that (15.33) holds. If there is a final k such that (14.22) and (21.6) hold with  $t_a$  and  $t_0$  replaced by  $t_k$ , then this k is denoted by  $k_{\text{fin,r}}$ . Moreover,  $t_{\text{fin,r}} := t_{k_{\text{fin,r}}}$ . If there is no such final k, then  $k_{\text{fin,r}} := \infty$  and  $t_{\text{fin,r}} := \infty$ . If, in addition to the above,  $t_0 \geq 2\pi$ , then  $t_{-1}$  is defined by the condition that (15.33) holds with k = -1. Assume  $t_k \leq t_0$  has been defined for some  $0 \geq k \in \mathbb{Z}$ ; that (14.22) and (21.6) hold with  $t_0$  and  $t_a$  replaced by  $t_k$ ; and that  $t_k \geq 2\pi$ . Then  $t_{k-1}$  is defined by the condition that (15.33) holds with k replaced by k - 1. Let  $k_{\text{ini,r}}$  be the first  $0 \geq k \in \mathbb{Z}$  such that either  $0 \leq t_k < 2\pi$  or one of the inequalities (14.22) and (21.6) (with  $t_0$  and  $t_a$  replaced by  $t_k$ ) is violated. Finally,  $t_{\text{ini,r}} := t_{k_{\text{ini,r}}}$ .

**Remark 21.5.** Note that since the equation is diagonally dominated, balanced and convergent, it is also oscillation adapted; cf. Lemma 20.3. Given that  $t_0$  satisfies the conditions of the definition, the sequence  $t_k$  is thus simply a restriction of the sequence defined in Definition 15.5. The conclusions derived concerning time sequences of the type introduced in Definition 15.5 thus apply to the time sequences defined here.

**Remark 21.6.** Due to (20.21), it is clear that

$$\max_{q} \bar{\nu}_{q}(\iota) e^{\bar{\beta}_{q}t} \leq \mathfrak{g}_{\text{diag}}(\iota, t) \leq \sqrt{Q} \max_{q} \bar{\nu}_{q}(\iota) e^{\bar{\beta}_{q}t}.$$

Combining this estimate with (20.22), it is clear that the estimate (21.6) is equivalent to a lower bound on  $\mathfrak{g}$  which only depends on the coefficients of the equation (1.2).

Next we turn to the question of how solutions develop in one frequency era. Here we focus on obtaining an upper bound on the norms of matrix products, but we return to the issue of optimality later.

**Lemma 21.7.** Consider the equation (1.2). Assume that it is non-degenerate, diagonally dominated, balanced and convergent; cf. Definition 21.1. Fix  $0 \neq \iota \in \mathcal{I}_{B}$  and let  $I_{q}$  be a corresponding frequency era; cf. Definition 20.10. Let  $\{t_{k}\}$  be a sequence defined as in Definition 21.4 and let  $R_{a,\pm}^{+}$ ,  $R_{a,\pm}^{-}$  and  $\kappa_{q,\pm}$  be given by Definition 21.2. Then the following holds.

• Assume that the Jordan blocks of  $R_{q,+}^+$  and  $R_{q,+}^-$  corresponding to the eigenvalues with real part  $\kappa_{q,+}$  are trivial. Then there are constants  $C_{q,+}$  and  $C_{\deg,q,+}$ , where  $C_{\deg,q,+} = 0$  if  $\bar{\beta}_q \neq 0$ , depending only on the coefficients of the equation (1.2), such that the following holds. If  $t_k \in I_q$  for  $k_0 - 1 \leq k \leq k_1 + 1$ , where  $k_0 \leq k_1$ , then

$$\|A_{k_1}^+ \cdots A_{k_0}^+\| \le C_{q,+} \exp\left[\kappa_{q,+}(\bar{t}_{1,+} - \bar{t}_{0,+}) + \frac{C_{\deg,q,+}}{\bar{\nu}_q(\iota)}(\bar{t}_{1,+} - \bar{t}_{0,+})\right],\tag{21.7}$$

where  $\bar{t}_{0,+} = t_{k_0}$ ,  $\bar{t}_{1,+} = t_{k_1+1}$  and  $A_k^{\pm} := \Xi_{\text{fin}}^{k,\pm}$ ; cf. Lemma 15.11.

• Assume that the Jordan blocks of  $R_{q,-}^+$  and  $R_{q,-}^-$  corresponding to the eigenvalues with real part  $\kappa_{q,-}$  are trivial. Then there are constants  $C_{q,-}$  and  $C_{\deg,q,-}$ , where  $C_{\deg,q,-} = 0$  if  $\bar{\beta}_q \neq 0$ , depending only on the coefficients of the equation (1.2), such that the following holds. If  $t_k \in I_q$  for  $k_0 - 1 \leq k \leq k_1 + 1$ , where  $k_0 \leq k_1$ , then

$$\|A_{k_0}^- \cdots A_{k_1}^-\| \le C_{q,-} \exp\left[\kappa_{q,-}(\bar{t}_{1,-} - \bar{t}_{0,-}) + \frac{C_{\deg,q,-}}{\bar{\nu}_q(\iota)}(\bar{t}_{1,-} - \bar{t}_{0,-})\right],\tag{21.8}$$

where  $\bar{t}_{0,-} := t_{k_0-1}$ ,  $\bar{t}_{1,-} := t_{k_1}$  and  $A_k^{\pm}$  is defined as above.

• Assume that there is a non-trivial Jordan block of  $R_{q,+}^+$  and  $R_{q,+}^-$  corresponding to an eigenvalue with real part  $\kappa_{q,+}$ . For every  $\epsilon > 0$ , there are then constants  $C_{q,\epsilon,+}$  and  $C_{\deg,q,+}$ , where  $C_{\deg,q,+} = 0$  if  $\beta_q \neq 0$ , depending only on the coefficients of the equation (1.2) and  $\epsilon$ , such that the following holds. If  $t_k \in I_q$  for  $k_0 - 1 \leq k \leq k_1 + 1$ , where  $k_0 \leq k_1$ , then

$$\|A_{k_1}^+ \cdots A_{k_0}^+\| \le C_{q,\epsilon,+} \exp\left[(\kappa_{q,+} + \epsilon)(\bar{t}_{1,+} - \bar{t}_{0,+}) + \frac{C_{\deg,q,+}}{\bar{\nu}_q(\iota)}(\bar{t}_{1,+} - \bar{t}_{0,+})\right], \quad (21.9)$$

where  $\bar{t}_{0,+}$ ,  $\bar{t}_{1,+}$  and  $A_k^+$  are defined as above.

• Assume that there is a non-trivial Jordan block of  $R_{q,-}^+$  and  $R_{q,-}^-$  corresponding to an eigenvalue with real part  $\kappa_{q,-}$ . For every  $\epsilon > 0$ , there are then constants  $C_{q,\epsilon,-}$  and  $C_{\deg,q,-}$ , where  $C_{\deg,q,-} = 0$  if  $\beta_q \neq 0$ , depending only on the coefficients of the equation (1.2) and  $\epsilon$ , such that the following holds. If  $t_k \in I_q$  for  $k_0 - 1 \leq k \leq k_1 + 1$ , where  $k_0 \leq k_1$ , then

$$\|A_{k_0}^- \cdots A_{k_1}^-\| \le C_{q,\epsilon,-} \exp\left[(\kappa_{q,-} + \epsilon)(\bar{t}_{1,-} - \bar{t}_{0,-}) + \frac{C_{\deg,q,-}}{\bar{\nu}_q(\iota)}(\bar{t}_{1,-} - \bar{t}_{0,-})\right], \quad (21.10)$$

where  $\bar{t}_{0,-}$ ,  $\bar{t}_{1,-}$  and  $A_k^-$  are defined as above.

**Remark 21.8.** For a given  $0 \neq \iota \in \mathcal{I}_{B}$ , it could be that there is no sequence defined as in Definition 21.4. In that case, the lemma does not yield any conclusions.

**Remark 21.9.** It is important to keep in mind that the orders of the matrices appearing on the left hand sides of (21.7) and (21.8) are different; cf. also (21.9) and (21.10). This is due to the fact that in the case of the plus sign, we are starting at  $t_{k_0}$  and evolving forwards to  $t_{k_1}$ . In the case of the minus sign we are starting at  $t_{k_1}$  and evolving backwards to  $t_{k_0}$ .

**Remark 21.10.** If  $k_0 > k_{\text{ini},r}$  and  $k_1 < k_{\text{fin},r}$ , it is sufficient to demand that  $t_k \in I_q$  for  $k_0 \le k \le k_1$ .

*Proof.* We begin by estimating  $A_k^{\pm}$  for  $t_k \in I_q$ .

**Estimating**  $A_k^{\pm}$ . Since  $A_k^{\pm} = \Xi_{\text{fin}}^{k,\pm}$  and (15.56) holds, it is of interest to simplify the matrix defined in (15.57). We therefore begin by simplifying  $R_{\text{pre},k}^{ab}$ , given by (15.41). Using the notation introduced in (20.37),

$$\bar{X}^q_{\text{diag}}(\iota) = \pm \bar{X}^q_{\infty},\tag{21.11}$$

where  $X_{\infty}^q$  is introduced in Definition 21.2. In order to determine the sign, note that the right hand side of (21.11) is only non-zero when q corresponds to a  $j \in \{1, \ldots, d\}$ . Let us therefore assume this to be the case. In order for  $I_q$  to be non-empty, it is then necessary for  $n_j$  to be non-zero, and

$$\bar{X}_{\text{diag}}^q(\iota) = \frac{n_j}{|n_j|} \tilde{X}_{\infty}^q;$$

cf. (20.37). Assume that  $t_k \in I_q$ . Then (20.27), (20.33) and (20.39) yield

$$\left\|\frac{\alpha(t_k)}{\mathfrak{g}(\iota, t_k)} - \frac{\alpha_{\infty}}{\bar{\nu}_q(\iota)e^{\bar{\beta}_q t_k}}\right\| \le \frac{C}{\bar{\nu}_q(\iota)e^{\bar{\beta}_q t_k}} \left[e^{-\eta_{\alpha} t_k} + \mathfrak{H}_q(\iota, t_k)\right],\tag{21.12}$$

where C only depends on the coefficients of (1.2). Similarly, (20.26), (20.33) and (20.38) yield

$$\left\|\frac{X(\iota, t_k)}{\mathfrak{g}(\iota, t_k)} - \frac{\bar{X}_{\text{diag}}^q(\iota)}{\bar{\nu}_q(\iota)e^{\bar{\beta}_q t_k}}\right\| \le \frac{C}{\bar{\nu}_q(\iota)e^{\bar{\beta}_q t_k}} \left[e^{-\eta_X t_k} + \sum_{s=1}^2 \mathfrak{H}_q^{(s)}(\iota, t_k)\right],\tag{21.13}$$

where C only depends on the coefficients of (1.2). Finally, (20.24), (20.33) and (20.36) yield

$$\frac{\dot{\ell}(\iota, t_k)}{\mathfrak{g}(\iota, t_k)} - \frac{\bar{\beta}_q}{\bar{\nu}_q(\iota)e^{\bar{\beta}_q t_k}} \le \frac{C}{\bar{\nu}_q(\iota)e^{\bar{\beta}_q t_k}} \left[ e^{-\eta_{\text{diag}} t_k} + \mathfrak{H}_q(\iota, t_k) \right],$$
(21.14)

#### 21.2. MODE ANALYSIS IN ONE FREQUENCY ERA

where C only depends on the coefficients of (1.2). Combining (21.12), (21.13) and (21.14) yields

$$\left\| R_{\text{pre},k}^{ab} - \frac{1}{\bar{\nu}_q(\iota)e^{\bar{\beta}_q t_k}} R_{\infty,q}^{ab}(\iota) \right\| \le \frac{C}{\bar{\nu}_q(\iota)e^{\bar{\beta}_q t_k}} \left[ e^{-\eta_R t_k} + \sum_{s=1}^2 \mathfrak{H}_q^{(s)}(\iota, t_k) \right],$$
(21.15)

where C only depends on the coefficients of (1.2),  $R^{ab}_{\text{pre},k}$  is defined in (15.41),

$$R^{ab}_{\infty,q}(\iota) := (-1)^a \alpha_\infty + \bar{\beta}_q \mathrm{Id}_m + (-1)^b \bar{X}^q_{\mathrm{diag}}(\iota)$$

and  $\eta_R := \eta_X = \eta_\alpha \leq \eta_{\text{diag}}$ . Defining  $A_{k,q,\pm}^{\text{app}}$  by

$$A_{k,q,\pm}^{\text{app}} := \text{Id}_{2m} \pm \frac{2\pi}{\bar{\nu}_q(\iota)e^{\bar{\beta}_q t_k}} R_{\infty,q}(\iota), \quad R_{\infty,q}(\iota) := \frac{1}{2} \begin{pmatrix} R_{\infty,q}^{10}(\iota) & 0\\ 0 & R_{\infty,q}^{11}(\iota) \end{pmatrix},$$
(21.16)

Lemma 15.11 and (21.15) imply that

$$\|A_{k}^{\pm} - A_{k,q,\pm}^{\mathrm{app}}\| \le \frac{C}{\bar{\nu}_{q}(\iota)e^{\bar{\beta}_{q}t_{k}}} \left[ e^{-\eta_{A}t_{k}} + \sum_{s=1}^{2} \mathfrak{H}_{q}^{(s)}(\iota, t_{k}) + \frac{1}{\bar{\nu}_{q}(\iota)e^{\bar{\beta}_{q}t_{k}}} \right].$$

where  $A_k^{\pm} = \Xi_{\text{fin}}^{k,\pm}$ ,  $\eta_A := \min\{\eta_R, \eta_{\text{sh}}\}$  and *C* only depends on the coefficients of (1.2). Note, however, that this estimate is only valid for  $t_k \in I_q$ . It is also of interest to note that the set  $\{R_{q,+}^+, R_{q,+}^-\}$  coincides with the set  $\{R_{\infty,q}^{10}(\iota)/2, R_{\infty,q}^{11}(\iota)/2\}$ . In this sense, the sign issue arising in (21.11) does not affect the end result. For future reference, it is of interest to keep in mind that

$$\kappa_{q,+} = \kappa_{\max}[R_{\infty,q}(\iota)], \quad \kappa_{q,-} = -\kappa_{\min}[R_{\infty,q}(\iota)]; \tag{21.17}$$

cf. Definition 4.3.

**Reformulating the matrix products.** As a next step, it is convenient to reformulate the matrix products by conjugating the factors with appropriate matrices. To be more specific, fix  $\epsilon > 0$  and let  $T_{q,\epsilon}$  be such that

$$J_{q,\epsilon} := T_{q,\epsilon} R_{\infty,q}(\iota) T_{q,\epsilon}^{-1}$$

is in generalised Jordan normal form with the non-zero off-diagonal components equal to  $\epsilon$ ; cf. Remark 9.10. For the sake of brevity, we suppress the dependence of  $J_{q,\epsilon}$  and  $T_{q,\epsilon}$  on  $\iota$ ; note, however, that the image of  $R_{\infty,q}(\iota)$ , considered as a function of  $\iota$ , consists of at most two matrices, so that the constants appearing in the estimates below can be chosen to be independent of  $\iota$ . Note also that

$$T_{q,\epsilon}A_{k_1}^+ \cdots A_{k_0}^+ T_{q,\epsilon}^{-1} = A_{k_1,q,\epsilon}^+ \cdots A_{k_0,q,\epsilon}^+, \tag{21.18}$$

$$T_{q,\epsilon}A^{-}_{k_{0}}\cdots A^{-}_{k_{1}}T^{-1}_{q,\epsilon} = A^{-}_{k_{0},q,\epsilon}\cdots A^{-}_{k_{1},q,\epsilon},$$
(21.19)

where

$$A_{k,q,\epsilon}^{\pm} := T_{q,\epsilon} A_k^{\pm} T_{q,\epsilon}^{-1}.$$
(21.20)

Moreover,

$$\|A_{k,q,\epsilon}^{\pm} - A_{k,q,\epsilon}^{\operatorname{app},\pm}\| \le \frac{C_{\epsilon}}{\bar{\nu}_q(\iota)e^{\bar{\beta}_q t_k}} \left[ e^{-\eta_A t_k} + \sum_{s=1}^2 \mathfrak{H}_q^{(s)}(\iota, t_k) + \frac{1}{\bar{\nu}_q(\iota)e^{\bar{\beta}_q t_k}} \right],$$

where  $C_{\epsilon}$  only depends on the coefficients of (1.2) and  $\epsilon$ , and

$$A_{k,q,\epsilon}^{\mathrm{app},\pm} := \mathrm{Id}_{2m} \pm \frac{2\pi}{\bar{\nu}_q(\iota)e^{\bar{\beta}_q t_k}} J_{q,\epsilon}.$$

In order to obtain an upper bound on the norms of the matrix products on the right hand sides of (21.18) and (21.19), it is convenient to consider

$$e^{-\lambda(\bar{t}_{1,+}-\bar{t}_{0,+})}A^+_{k_1,q,\epsilon}\cdots A^+_{k_0,q,\epsilon} = \hat{A}^+_{k_1,q,\epsilon}\cdots \hat{A}^+_{k_0,q,\epsilon},$$
(21.21)

$$e^{-\lambda(\bar{t}_{0,-}-\bar{t}_{1,-})}A^{-}_{k_{0},q,\epsilon}\cdots A^{-}_{k_{1},q,\epsilon} = \hat{A}^{-}_{k_{0},q,\epsilon}\cdots \hat{A}^{-}_{k_{1},q,\epsilon},$$
(21.22)

where  $\lambda$  is a real number and

$$\hat{A}_{k,q,\epsilon}^{\pm} := e^{-\lambda(t_{k\pm 1} - t_k)} A_{k,q,\epsilon}^{\pm}.$$
(21.23)

Note that

$$\left|e^{-\lambda(t_{k\pm 1}-t_k)} - 1 + \lambda(t_{k\pm 1}-t_k)\right| \le \frac{C_\lambda}{\mathfrak{g}^2(\iota,t_k)}$$

where the constant  $C_{\lambda}$  only depends on  $\lambda$  and we appealed to (14.24); for future reference, it is of interest to keep in mind that a similar estimate holds when  $\lambda$  is complex. Combining this estimate with (20.51) (and the comment made in connection with it) yields

$$\left| e^{-\lambda(t_{k\pm 1}-t_k)} - 1 \pm \frac{2\pi\lambda}{\bar{\nu}_q(\iota)e^{\bar{\beta}_q t_k}} \right| \le \frac{C_\lambda}{\mathfrak{g}(\iota, t_k)} \left[ e^{-\eta_{\text{diag}} t_k} + \mathfrak{H}_q(\iota, t_k) \right] + \frac{C_\lambda}{\mathfrak{g}^2(\iota, t_k)}, \tag{21.24}$$

where  $C_{\lambda}$  only depends on the coefficients of (1.2) and  $\lambda$ ; again, a similar estimate holds when  $\lambda$  is complex. Thus

$$|\hat{A}_{k,q,\epsilon}^{\pm} - \hat{A}_{k,q,\epsilon}^{\mathrm{app},\pm}|| \le \frac{C_{\epsilon,\lambda}}{\bar{\nu}_q(\iota)e^{\bar{\beta}_q t_k}} \left[ e^{-\eta_A t_k} + \sum_{s=1}^2 \mathfrak{H}_q^{(s)}(\iota, t_k) + \frac{1}{\bar{\nu}_q(\iota)e^{\bar{\beta}_q t_k}} \right],$$

where

$$\hat{A}_{k,q,\epsilon}^{\mathrm{app},\pm} := \mathrm{Id}_{2m} \pm \frac{2\pi}{\bar{\nu}_q(\iota)e^{\bar{\beta}_q t_k}} [J_{q,\epsilon} - \lambda \mathrm{Id}_{2m}].$$

Moreover,  $C_{\epsilon,\lambda}$  only depends on the coefficients of (1.2),  $\epsilon$  and  $\lambda$ .

Before proceeding, it is of interest to eliminate the imaginary parts of the eigenvalues of  $J_{q,\epsilon}$ . Note that  $J_{q,\epsilon} - \lambda \mathrm{Id}_{2m}$  consists of generalised Jordan blocks. Assume that the blocks have dimension  $l_1, \ldots, l_D$ . Let  $\zeta_r$  be the imaginary part of the eigenvalue corresponding to the r'th generalised Jordan block. Define  $S_k$  and  $\check{A}_{k,q,\epsilon}^{\pm}$  via

$$S_k := \operatorname{diag}\{e^{-i\zeta_1 t_k} \operatorname{Id}_{l_1}, \dots, e^{-i\zeta_D t_k} \operatorname{Id}_{l_D}\}, \quad \check{A}^{\pm}_{k,q,\epsilon} := S_{k\pm 1} \hat{A}^{\pm}_{k,q,\epsilon} S_k^{-1}.$$
(21.25)

Note that  $||S_k|| = ||S_k^{-1}|| = 1$  and that

$$S_{k_1+1}\hat{A}^+_{k_1,q,\epsilon}\cdots\hat{A}^+_{k_0,q,\epsilon}S^{-1}_{k_0} = \check{A}^+_{k_1,q,\epsilon}\cdots\check{A}^+_{k_0,q,\epsilon},$$
(21.26)

$$S_{k_0-1}\hat{A}^{-}_{k_0,q,\epsilon}\cdots\hat{A}^{-}_{k_1,q,\epsilon}S^{-1}_{k_1}=\check{A}^{-}_{k_0,q,\epsilon}\cdots\check{A}^{-}_{k_1,q,\epsilon}.$$
(21.27)

Moreover,

$$\|\check{A}_{k,q,\epsilon}^{\pm} - \check{A}_{k,q,\epsilon}^{\mathrm{app},\pm}\| \le \frac{C_{\epsilon,\lambda}}{\bar{\nu}_q(\iota)e^{\bar{\beta}_q t_k}} \left[ e^{-\eta_A t_k} + \sum_{s=1}^2 \mathfrak{H}_q^{(s)}(\iota, t_k) + \frac{1}{\bar{\nu}_q(\iota)e^{\bar{\beta}_q t_k}} \right],\tag{21.28}$$

where

$$\check{A}_{k,q,\epsilon}^{\mathrm{app},\pm} := \mathrm{Id}_{2m} \pm \frac{2\pi}{\bar{\nu}_q(\iota)e^{\bar{\beta}_q t_k}} [\mathrm{Re}\{J_{q,\epsilon}\} - \lambda \mathrm{Id}_{2m}];$$
(21.29)

this is a consequence of an estimate similar to (21.24). Finally, the constant  $C_{\epsilon,\lambda}$  appearing in (21.28) only depends on  $\epsilon$ ,  $\lambda$  and the coefficients of the equation (1.2).

Estimating the matrix products. Let  $\kappa_{q,+}$  be defined by (21.3) and fix a real number  $\lambda$  such that  $\kappa_{q,+} < \lambda \leq \kappa_{q,+} + 1/2$ . By choosing  $\epsilon > 0$  small enough (depending only on  $\lambda$  and the

coefficients of the equation), it can then be ensured that  $\|\check{A}_{k,q,\epsilon}^{\text{app},+}\| \leq 1$ . The reason for this is the following. Since  $t_k$  is an element of the sequence introduced in Definition 21.4 and since  $t_k \in I_q$ , we know that

$$\bar{\nu}_q(\iota)e^{\bar{\beta}_q t_k} = \max_{o} \bar{\nu}_{\rho}(\iota)e^{\bar{\beta}_{\rho} t_k} \ge 2\pi(\kappa+1).$$

Thus the diagonal components of the second term on the right hand side of (21.29) are bounded in absolute value by  $(\kappa + 1/2)/(\kappa + 1)$ ; recall that (21.17) holds. On the other hand, the diagonal components of  $\operatorname{Re}\{J_{q,\epsilon}\} - \lambda \operatorname{Id}_{2m}$  are bounded from above by  $\kappa_{q,+} - \lambda < 0$ . Choosing  $\epsilon > 0$ small enough (the bound depending only on the coefficients of the equation and  $\lambda$ ), it can then be ensured that in the case of the +-sign, the second term on the right hand side of (21.29) is negative definite and has a norm bounded from above by 1. Moreover,  $\|\check{A}_{k,q,\epsilon}^{\operatorname{app},+}\| \leq 1$ . In case all the generalised Jordan blocks corresponding to eigenvalues with real part  $\kappa_{q,+}$  are trivial,  $\lambda$  can be chosen to equal  $\kappa_{q,+}$  and  $\epsilon > 0$  can be chosen to be a fixed number depending only on the difference between  $\kappa_{q,+}$  and the second largest real part of an eigenvalue of  $J_{q,\epsilon}$  (in case there is no second largest real part, there are no non-trivial generalised Jordan blocks, and then there is no need for  $\epsilon$ ).

By a similar argument, if we fix  $\lambda$  so that  $-\kappa_{q,-} - 1/2 \leq \lambda < -\kappa_{q,-}$ , then there is an  $\epsilon > 0$ (depending only on  $\lambda$  and the coefficients of the equation (1.2)) so that  $\|\check{A}_{k,q,\epsilon}^{\mathrm{app},-}\| \leq 1$ . Again, in case all the generalised Jordan blocks corresponding to eigenvalues with real part  $\kappa_{q,-}$  are trivial,  $\lambda$  can be chosen to equal  $-\kappa_{q,-}$  and  $\epsilon > 0$  can be chosen to be a fixed number depending only on the difference between  $-\kappa_{q,-}$  and the second smallest real part of an eigenvalue of  $J_{q,\epsilon}$ .

With  $\lambda$  and  $\epsilon$  chosen as above,

$$\begin{split} \|\check{A}_{k_{1},q,\epsilon}^{+}\cdots\check{A}_{k_{0},q,\epsilon}^{+}\| &\leq \left(1+\|\check{A}_{k_{1},q,\epsilon}^{+}-\check{A}_{k_{1},q,\epsilon}^{\mathrm{app},+}\|\right)\cdots\left(1+\|\check{A}_{k_{0},q,\epsilon}^{+}-\check{A}_{k_{0},q,\epsilon}^{\mathrm{app},+}\|\right),\\ \|\check{A}_{k_{0},q,\epsilon}^{-}\cdots\check{A}_{k_{1},q,\epsilon}^{-}\| &\leq \left(1+\|\check{A}_{k_{1},q,\epsilon}^{-}-\check{A}_{k_{1},q,\epsilon}^{\mathrm{app},-}\|\right)\cdots\left(1+\|\check{A}_{k_{0},q,\epsilon}^{-}-\check{A}_{k_{0},q,\epsilon}^{\mathrm{app},-}\|\right). \end{split}$$

Since  $\ln(1+x) \le x$  for  $x \ge 0$ , we conclude that the logarithms of the right hand sides are bounded from above by

$$\sum_{k=k_0}^{k_1} \|\check{A}_{k,q,\epsilon}^{\pm} - \check{A}_{k,q,\epsilon}^{\text{app},\pm}\|.$$
(21.30)

Due to (21.28), it is thus of interest to estimate the sum of the right hand side of (21.28). Let us begin with

$$\sum_{k=k_0}^{k_1} \frac{1}{\bar{\nu}_q(\iota)e^{\bar{\beta}_q t_k}} e^{-\eta_A t_k} \le C \int_{\bar{t}_{0,+}}^{\bar{t}_{1,+}} e^{-\eta_A t} dt \le C,$$

where we appealed to (20.44) and the constant C only depends on the coefficients of the equation (1.2). Moreover, (20.44) implies that

$$\sum_{k=k_0}^{k_1} \frac{1}{\bar{\nu}_q^2(\iota) e^{2\bar{\beta}_q t_k}} \le C \int_{\bar{t}_{0,+}}^{\bar{t}_{1,+}} \frac{1}{\bar{\nu}_q(\iota) e^{\bar{\beta}_q t}} dt,$$

where C has the same dependence as before. If  $\bar{\beta}_q \neq 0$ , the right hand side of this inequality is bounded. However, if  $\bar{\beta}_q = 0$ , we only have the bound

$$\sum_{k=k_0}^{k_1} \frac{1}{\bar{\nu}_q^2(\iota) e^{2\bar{\beta}_q t_k}} \le \frac{C}{\bar{\nu}_q(\iota)} (\bar{t}_{1,+} - \bar{t}_{0,+}).$$
(21.31)

On the other hand, the constant C only depends on the coefficients of the equation (1.2). Finally

$$\sum_{k=k_0}^{k_1} \frac{1}{\bar{\nu}_q(\iota)e^{\bar{\beta}_q t_k}} \mathfrak{H}_q^{(s)}(\iota, t_k) \le C \int_{\bar{t}_{0,+}}^{\bar{t}_{1,+}} \mathfrak{H}_q^{(s)}(\iota, t) dt$$

for s = 1, 2, where we appealed to (20.45) and C has the same dependence as before. Note that the right hand side can be computed using (20.52). Since  $\lambda = 0$  and a = 1, we, moreover, have  $(\lambda + s\bar{\beta}_p)_{a+s,q} = s(\bar{\beta}_p - \bar{\beta}_q) \neq 0$  (since  $\bar{\beta}_p \neq \bar{\beta}_q$  for  $p \neq q$  and s > 0). We can therefore combine (20.52) and (20.53) in order to conclude that

$$\int_{\bar{t}_{0,+}}^{\bar{t}_{1,+}} \mathfrak{H}_q^{(s)}(\iota,t) dt = \sum_{p \neq q} \frac{1}{s(\bar{\beta}_p - \bar{\beta}_q)} \left( \frac{\bar{\nu}_p^s(\iota) e^{s\bar{\beta}_p \bar{t}_{1,+}}}{\bar{\nu}_q^s(\iota) e^{s\bar{\beta}_q \bar{t}_{1,+}}} - \frac{\bar{\nu}_p^s(\iota) e^{s\bar{\beta}_p \bar{t}_{0,+}}}{\bar{\nu}_q^s(\iota) e^{s\bar{\beta}_q \bar{t}_{0,+}}} \right) \leq C$$

for s = 1, 2, where *C* has the same dependence as before. To conclude, the sum appearing in (21.30) is bounded if  $\bar{\beta}_q \neq 0$  and is bounded by a constant plus a constant times the right hand side of (21.31) in case  $\bar{\beta}_q = 0$ . In both cases, the constant only depends on  $\epsilon$ ,  $\lambda$  and the coefficients of the equation. Thus  $\|\check{A}_{k_1,q,\epsilon}^+ \cdots \check{A}_{k_0,q,\epsilon}^+\|$  and  $\|\check{A}_{k_0,q,\epsilon}^- \cdots \check{A}_{k_1,q,\epsilon}^-\|$  are bounded if  $\bar{\beta}_q \neq 0$ . In case  $\bar{\beta}_q = 0$ , it is necessary to include a factor of the form

$$\exp\left[\frac{C_{\deg,q,\pm}}{\bar{\nu}_q(\iota)}(\bar{t}_{1,+}-\bar{t}_{0,+})\right].$$

Combining these observations with (21.18), (21.21) and (21.26) yields

$$\begin{aligned} |A_{k_{1}}^{+} \cdots A_{k_{0}}^{+}\| &\leq C_{\epsilon} \|A_{k_{1},q,\epsilon}^{+} \cdots A_{k_{0},q,\epsilon}^{+}\| \leq C_{\epsilon} e^{\lambda(\bar{t}_{1,+} - \bar{t}_{0,+})} \|\hat{A}_{k_{1},q,\epsilon}^{+} \cdots \hat{A}_{k_{0},q,\epsilon}^{+}\| \\ &\leq C_{\epsilon} e^{\lambda(\bar{t}_{1,+} - \bar{t}_{0,+})} \|\check{A}_{k_{1},q,\epsilon} \cdots \check{A}_{k_{0},q,\epsilon}\| \\ &\leq C_{\epsilon,\lambda} \exp\left[\lambda(\bar{t}_{1,+} - \bar{t}_{0,+}) + \frac{C_{\deg,q,+}}{\bar{\nu}_{q}(\iota)}(\bar{t}_{1,+} - \bar{t}_{0,+})\right]. \end{aligned}$$

Here  $C_{\epsilon,\lambda}$  only depends on  $\epsilon$ ,  $\lambda$  and the coefficients of the equation (1.2). Moreover,  $C_{\deg,q,+}$  only depends on  $\epsilon$ ,  $\lambda$  and the coefficients of the equation (1.2), and it vanishes if  $\bar{\beta}_q \neq 0$ . A similar argument yields

$$\|A_{k_0}^- \cdots A_{k_1}^-\| \le C_{\epsilon,\lambda} \exp\left[\lambda(\bar{t}_{0,-} - \bar{t}_{1,-}) + \frac{C_{\deg,q,-}}{\bar{\nu}_q(\iota)}(\bar{t}_{1,-} - \bar{t}_{0,-})\right],$$

where the constants have the same dependence and properties as in the previous estimate. The appearance of  $\bar{t}_{1,-} - \bar{t}_{0,-}$  in the last term inside the exponential requires some justification. However,  $|\bar{t}_{1,+} - \bar{t}_{1,-}|$  and  $|\bar{t}_{0,+} - \bar{t}_{0,-}|$  are bounded by numerical constants. Changing  $\bar{t}_{1,+}$  to  $\bar{t}_{1,-}$  and  $\bar{t}_{0,+}$  to  $\bar{t}_{0,-}$  thus only amounts to a change of the constant  $C_{\epsilon,\lambda}$ . In the case of trivial generalised Jordan blocks we can choose  $\lambda = \kappa_{q,+}$  and  $\lambda = -\kappa_{q,-}$  for the respective cases. Moreover, in those cases,  $\epsilon$  can be chosen to depend only on the coefficients of the equation, as described above. Finally, if the relevant generalised Jordan blocks are non-trivial, we need to include a margin in the form of the  $\epsilon > 0$  appearing in the statement of the lemma. The lemma follows.

In order to justify Remark 21.10, note that  $||A_k^{\pm}||$  is bounded by a constant depending only on the coefficients of the equation.

#### **21.3** Refined estimates in transparent eras

When  $\bar{\beta}_q \neq 0$ , the estimates derived in Lemma 21.7 are satisfactory. When  $\bar{\beta}_q = 0$  and  $\bar{\nu}_q(\iota)$  is large, they are also satisfactory. However, when  $\bar{\beta}_q = 0$  and  $\bar{\nu}_q(\iota)$  is not large, there is room for improvement. We therefore treat this situation separately. In order to develop a feeling concerning how to approximate the solution in this setting, let

$$\begin{pmatrix} \tilde{x}_q(\iota,t) \\ \tilde{y}_q(\iota,t) \end{pmatrix} := \begin{pmatrix} \bar{\nu}_q(\iota)z(\iota,t) \\ \dot{z}(\iota,t) \end{pmatrix}, \quad \varpi_q := \begin{pmatrix} \tilde{x}_q \\ \tilde{y}_q \end{pmatrix}.$$
(21.32)

It can then be calculated that

$$\dot{\varpi}_q(\iota, t) = A_{q,\infty}(\iota) \varpi_q(\iota, t) + A_{q,\text{rem}}(\iota, t) \varpi_q(\iota, t) + F(\iota, t), \qquad (21.33)$$

where

$$A_{q,\infty}(\iota) := \begin{pmatrix} 0 & \bar{\nu}_q(\iota) \mathrm{Id}_m \\ -\bar{\nu}_q(\iota) \mathrm{Id}_m - i \bar{X}_{\mathrm{diag}}^q(\iota) - [\bar{\nu}_q(\iota)]^{-1} \zeta_\infty & -\alpha_\infty \end{pmatrix},$$
(21.34)

$$F(\iota, t) := \begin{pmatrix} 0 \\ \hat{f}(\iota, t) \end{pmatrix}.$$
(21.35)

In order for this observation to be of interest, it is, however, necessary to estimate the norm of the matrix  $A_{q,\text{rem}}(\iota,t)$  appearing in (21.33). We do so in the proof of Lemma 21.11 below. Since  $A_{q,\infty}(\iota)$  is independent of t, approximating the behaviour of solutions should be straightforward. Moreover, assuming  $A_{q,\text{rem}}(\iota,t)$  to be small, the quantities

$$\kappa_{q,\iota,\pm} := \kappa_{\max}[\pm A_{q,\infty}(\iota)] \tag{21.36}$$

can be expected to correspond to the dominant behaviour solutions;  $\kappa_{q,\iota,+}$  for increasing t and  $\kappa_{q,\iota,-}$  for decreasing t.

**Lemma 21.11.** Consider the equation (1.2). Assume that it is non-degenerate, diagonally dominated, balanced and convergent; cf. Definition 21.1. Consider  $0 \neq \iota \in \mathcal{I}_{B}$  such that there is a frequency era  $I_{q}$  with  $\bar{\beta}_{q} = 0$ ; cf. Definition 20.10. Fix  $\epsilon > 0$  and  $K_{co} > 0$ . Then there is a constant  $C_{csp,\epsilon}$  depending only on an upper bound on  $K_{co}$ , the spectrum of the Riemannian manifold corresponding to q (if q corresponds to a  $j \in \{1, \ldots, d\}$ , then the relevant Riemannian manifold is  $\mathbb{S}^{1}$  with the standard Euclidean metric),  $\epsilon$  and the coefficients of the equation (1.2) such that

$$|w(\iota,t)| \le C_{\text{csp},\epsilon} e^{(\kappa_{q,\iota,+}+\epsilon)(t-t_{-})} |w(\iota,t_{-})| + C_{\text{csp},\epsilon} \int_{t_{-}}^{t} e^{(\kappa_{q,\iota,+}+\epsilon)(t-t')} |\hat{f}(\iota,t')| dt'$$
(21.37)

for  $\iota \in \mathcal{I}_{\mathrm{B}}$  such that  $I_q \neq \emptyset$  and  $\overline{\nu}_q(\iota) \leq K_{\mathrm{co}}$ ;  $t_- \in I_q$ ;  $t_- \leq t \in I_q$ ; and z a solution to (24.16). Here w is defined in terms of z as described in (14.7) and (15.1). Moreover,  $\kappa_{q,\iota,+}$  is defined in (21.36) and the  $\epsilon$  can be removed in case the Jordan blocks of  $A_{q,\infty}(\iota)$  corresponding to eigenvalues with real parts equal to  $\kappa_{q,\iota,+}$  are trivial; cf. (21.34) for a definition of  $A_{q,\infty}(\iota)$ . Moreover,

$$|w(\iota, t)| \le C_{\text{csp}, \epsilon} e^{-(\kappa_{q, \iota, -} + \epsilon)(t - t_{+})} |w(\iota, t_{+})| + C_{\text{csp}, \epsilon} \int_{t}^{t_{+}} e^{-(\kappa_{q, \iota, -} + \epsilon)(t - t')} |\hat{f}(\iota, t')| dt'$$

for  $\iota \in \mathcal{I}_{\mathrm{B}}$  such that  $I_q \neq \emptyset$  and  $\bar{\nu}_q(\iota) \leq K_{\mathrm{co}}$ ;  $t_+ \in I_q$ ; and  $t_+ \geq t \in I_q$ . Here  $\kappa_{q,\iota,-}$  is defined in (21.36) and the  $\epsilon$  can be removed in case the Jordan blocks of  $-A_{q,\infty}(\iota)$  corresponding to eigenvalues with real parts equal to  $\kappa_{q,\iota,-}$  are trivial.

**Remark 21.12.** When applying this lemma, it is of interest to keep the following in mind. First, for a given  $0 < K_{co} \in \mathbb{R}$ , there are only finitely many matrices of the form  $A_{q,\infty}(\iota)$  for  $\iota \in \mathcal{I}_{B}$  such that  $\bar{\nu}_{q}(\iota) \neq 0$  and  $\bar{\nu}_{q}(\iota) \leq K_{co}$ ; this is a consequence of the fact that the eigenvalues of the relevant Riemannian manifolds converge to infinity. Second, the matrices  $A_{q,\infty}(\iota)$  are of the form (11.7). Due to Lemma 11.6, the set of real parts of the eigenvalues of  $A_{q,\infty}(\iota)$  for  $\iota \in \mathcal{I}_{B}$  such that  $\bar{\nu}_{q}(\iota) \neq 0$  is thus bounded.

**Remark 21.13.** Note that the set of matrices  $A_{q,\infty}(\iota)$  for  $\iota \in \mathcal{I}_{\mathrm{B}}$  such that  $\bar{\nu}_q(\iota) \neq 0$  and  $\bar{\nu}_q(\iota) \leq K_{\mathrm{co}}$  is determined by three things:  $K_{\mathrm{co}}$ ; the spectrum of the Riemannian manifold corresponding to q; and the coefficients of the equation (1.2). That is why the constants appearing in the estimates have the stated dependence.

*Proof.* As described in the statement of the lemma, we are here interested in situations such that  $\bar{\nu}_q(\iota) \leq K_{co}$  for some constant  $K_{co} > 0$ . Moreover, we below allow the constants appearing in the estimates to depend on an upper bound on  $K_{co}$ .

Estimating the norm of  $A_{q,\text{rem}}(\iota, t)$ . The proof of the lemma is based on an analysis of solutions to (21.33), where the constituents, except for  $A_{q,\text{rem}}(\iota, t)$ , are defined in (21.32), (21.34) and (21.35). As a first step, we need to estimate the norm of  $A_{q,\text{rem}}(\iota, t)$ . Note, to this end, that

$$\mathfrak{g}^2(\iota, t) - \bar{\nu}_q^2(\iota)| \le C_{\mathrm{co}}\bar{\nu}_q(\iota)[\mathfrak{H}_q(\iota, t) + e^{-\eta_{\mathrm{diag}}t}]$$
(21.38)

for all  $t \in I_q$ , where  $C_{co}$  depends only on an upper bound on  $K_{co}$  and the coefficients of the equation (1.2). In order to obtain this estimate, we have appealed to (20.22), (20.23), (20.33) and (20.34). Note also that (20.33) and the present assumptions imply that  $\mathfrak{g}(\iota, t) \leq C_{co}$  for  $t \in I_q$ , where  $C_{co}$  depends only on an upper bound on  $K_{co}$  and the coefficients of the equation (1.2). Similarly,

$$|2i\sigma(\iota, t)\mathfrak{g}(\iota, t)| \le C_{\rm co}e^{-\eta_{\rm sh}t} \tag{21.39}$$

for all  $t \in I_q$ , where  $C_{\rm co}$  has the same dependence as before; recall that the equation is oscillation adapted with  $\mathfrak{f}_{\rm sh}(t) = Ce^{-\eta_{\rm sh}t}$  due to Lemma 20.3. Finally, let us estimate

$$\|n_{l}X^{l}(t) - \bar{\nu}_{q}(\iota)\bar{X}^{q}_{\text{diag}}(\iota)\| = \|n_{l}X^{l}(t) - \sum_{\{j:\beta_{j}=\bar{\beta}_{q}\}}n_{j}X^{j}_{\infty}\|$$

$$\leq C\mathfrak{H}^{(1)}_{q}(\iota,t)\bar{\nu}_{q}(\iota) + Ce^{-\eta_{\text{mn}}t}\bar{\nu}_{q}(\iota)$$
(21.40)

for all  $t \in I_q$ , where we appealed to (20.8), (20.9) and (20.37), and the constants C only depend on the coefficients of the equation (1.2). Combining (20.9) with (21.38), (21.39) and (21.40) yields the conclusion that (21.33) holds, where  $A_{q,\infty}(\iota)$  and  $F(\iota, t)$  are defined by (21.34) and (21.35) respectively. Moreover,

$$||A_{q,\text{rem}}(\iota,t)|| \le C_{\text{co}}\left(\sum_{s=1}^{2} \mathfrak{H}_{q}^{(s)}(\iota,t) + e^{-\eta_{\text{B}}t} + [\bar{\nu}_{q}(\iota)]^{-1}e^{-\eta_{\text{mn}}t}\right)$$

for all  $t \in I_q$ , where  $\eta_{\rm B} := \min\{\eta_{\rm sh}, \eta_{\rm mn}, \eta_{\rm diag}\}$  and  $C_{\rm co}$  depends only on an upper bound on  $K_{\rm co}$  and the coefficients of the equation (1.2).

**Reformulating the equations.** Fix  $\epsilon > 0$  and let  $T_{q,\iota,\epsilon}$  be such that

$$J_{q,\iota,\epsilon} := T_{q,\iota,\epsilon}^{-1} A_{q,\infty}(\iota) T_{q,\iota,\epsilon}$$

consists of generalised Jordan blocks with the non-zero off-diagonal terms equal to  $\epsilon/2$ ; cf. Remark 9.10. Let

$$R_{q,\iota}(t) := \exp[-i\operatorname{Im}\{J_{q,\iota,\epsilon}\}t];$$

note that the imaginary part of  $J_{q,\iota,\epsilon}$  is independent of  $\epsilon$ . Finally, let  $\kappa_{q,\iota,\pm}$  be defined by (21.36) and

$$\xi_{q,\iota}^{\pm}(t) := \exp[\mp(\kappa_{q,\iota,\pm} + \eta)\bar{t}_{\pm}]R_{q,\iota}(\bar{t}_{\pm})T_{q,\iota,\epsilon}^{-1}\varpi_q(\iota,t),$$
(21.41)

where  $\bar{t}_{\pm} := t - t_{\mp}; t_{-}, t_{+} \in I_{q};$  and  $\eta \ge 0$  remains to be determined. Then

$$\dot{\xi}_{q,\iota}^{\pm}(t) = [\operatorname{Re}\{J_{q,\iota,\epsilon}\} \mp (\kappa_{q,\iota,\pm} + \eta)\operatorname{Id}_{2m}]\xi_{q,\iota}^{\pm}(t) + A_{q,\operatorname{rem},\epsilon}^{\pm}(\iota,t)\xi_{q,\iota}^{\pm}(t) + G_{q,\epsilon}^{\pm}(\iota,t), \qquad (21.42)$$

where

$$\begin{aligned} A_{q,\mathrm{rem},\epsilon}^{\pm}(\iota,t) &:= R_{q,\iota}(\bar{t}_{\pm}) T_{q,\iota,\epsilon}^{-1} A_{q,\mathrm{rem}}(\iota,t) T_{q,\iota,\epsilon}[R_{q,\iota}(\bar{t}_{\pm})]^{-1}, \\ G_{q,\epsilon}^{\pm}(\iota,t) &:= \exp[\mp(\kappa_{q,\iota,\pm}+\eta)\bar{t}_{\pm}] R_{q,\iota}(\bar{t}_{\pm}) T_{q,\iota,\epsilon}^{-1} F(\iota,t). \end{aligned}$$

For future reference, it is of interest to note that

$$\|A_{q,\operatorname{rem},\epsilon}^{\pm}(\iota,t)\| \le C_{\operatorname{csp},\epsilon}\left(\sum_{s=1}^{2}\mathfrak{H}_{q}^{(s)}(\iota,t) + e^{-\eta_{\mathrm{B}}t}\right)$$
(21.43)

for all  $t \in I_q$ , where  $C_{csp,\epsilon}$  only depends on an upper bound on  $K_{co}$ , the spectrum of the Riemannian manifold corresponding to q (if q corresponds to a  $j \in \{1, \ldots, d\}$ , then the relevant Riemannian manifold is  $\mathbb{S}^1$  with the standard Euclidean metric),  $\epsilon$  and the coefficients of the equation (1.2). In what follows, we wish to choose  $\epsilon$  and  $\eta$  in such a way that

$$\operatorname{Re}\{J_{q,\iota,\epsilon}\} - (\kappa_{q,\iota,+} + \eta)\operatorname{Id}_{2m}, \quad \operatorname{Re}\{J_{q,\iota,\epsilon}\} + (\kappa_{q,\iota,-} + \eta)\operatorname{Id}_{2m}$$
(21.44)

are negative semi-definite and positive semi-definite respectively; here we say that  $A \in \mathbf{M}_k(\mathbb{C})$ is negative semi-definite if  $\operatorname{Re}\{\langle w, Aw \rangle\} \leq 0$  for all  $w \in \mathbb{C}^k$  etc. In case the Jordan blocks of

#### 21.4. ANALYSIS FOR ONE MODE

 $A_{q,\infty}(\iota)$  corresponding to eigenvalues with real parts equal to  $\kappa_{q,\iota,+}$  are all trivial, we choose  $\eta = 0$ and  $\epsilon$  small enough (the bound depending only on an upper bound on  $K_{co}$ , the spectrum of the Riemannian manifold corresponding to q and the coefficients of the equation (1.2)). Under these conditions, the first matrix appearing in (21.44) is negative semi-definite. In case there is a nontrivial Jordan block of  $A_{q,\infty}(\iota)$  corresponding to an eigenvalue with real part equal to  $\kappa_{q,\iota,+}$ , we choose  $\eta = \epsilon$ . Then the first matrix appearing in (21.44) is negative definite. The choices in the case of the second matrix appearing in (21.44) are similar.

Estimating the energy of the solution. Given the above choices of  $\epsilon$  and  $\eta$ ,

$$\frac{d}{dt}|\xi_{q,\iota}^+|^2 \le 2||A_{q,\mathrm{rem},\epsilon}^+(\iota,\cdot)|||\xi_{q,\iota}^+|^2 + 2|G_{q,\epsilon}^+(\iota,\cdot)||\xi_{q,\iota}^+|$$

for all  $t \in I_q$ , where we appealed to (21.42). Let

$$L_{q,\epsilon}^{+}(\iota,t) := -\int_{t_{-}}^{t} \|A_{q,\text{rem},\epsilon}^{+}(\iota,t')\|dt'.$$
(21.45)

Then

$$\frac{d}{dt} \left( \exp\left[2L_{q,\epsilon}^+(\iota,\cdot)\right] |\xi_{q,\iota}^+|^2 \right) \le 2 \exp\left[2L_{q,\epsilon}^+(\iota,\cdot)\right] |G_{q,\epsilon}^+(\iota,\cdot)| |\xi_{q,\iota}^+|$$

This estimate can be used to deduce that for  $t_{-} \leq t \in I_q$ ,

$$\exp\left[L_{q,\epsilon}^{+}(\iota,t)\right]|\xi_{q,\iota}^{+}(t)| \leq |\xi_{q,\iota}^{+}(t_{-})| + \int_{t_{-}}^{t} \exp\left[L_{q,\epsilon}^{+}(\iota,t')\right]|G_{q,\epsilon}^{+}(\iota,t')|dt'.$$
(21.46)

In order to estimate the right hand side of (21.45), note that (21.43) holds. Combining this estimate with (20.52) and (20.53) yields

$$|L_{q,\epsilon}^+(\iota,t)| \le C_{\operatorname{csp},\epsilon}$$

for all  $t \in I_q$ , where  $C_{csp,\epsilon}$  has the same dependence as in the case of (21.43). Combining this estimate with (21.46) and recalling the definitions of the constituents of (21.46) yields

$$|\varpi_q(\iota,t)| \le C_{\operatorname{csp},\epsilon} e^{(\kappa_{q,\iota,+}+\eta)\bar{t}_+} |\varpi_q(\iota,t_-)| + C_{\operatorname{csp},\epsilon} \int_{t_-}^t e^{(\kappa_{q,\iota,+}+\eta)(t-t')} |F(\iota,t')| dt$$

for  $t_{-} \leq t \in I_q$ , where  $C_{csp,\epsilon}$  has the same dependence as in the case of (21.43). A similar argument yields

$$|\varpi_q(\iota,t)| \le C_{\operatorname{csp},\epsilon} e^{-(\kappa_{q,\iota,-}+\eta)\bar{t}_-} |\varpi_q(\iota,t_+)| + C_{\operatorname{csp},\epsilon} \int_t^{t_+} e^{-(\kappa_{q,\iota,-}+\eta)(t-t')} |F(\iota,t')| dt'$$

for  $t_+ \ge t \in I_q$ , where  $C_{csp,\epsilon}$  has the same dependence as in the case of (21.43). Since  $|\varpi_q(\iota, t)|$  and  $|w(\iota, t)|$  are equivalent for  $t \in I_q$ , where the constant of equivalence only depends on the coefficients of (1.2), the lemma follows.

#### 21.4 Analysis for one mode

In Sections 21.2 and 21.3 we analyse how the energy of one mode evolves over one frequency era. The purpose of the present section is to combine the corresponding results in order to obtain an estimate over intervals that include several frequency eras. Before we state the corresponding result, we, however, need to introduce additional terminology.

**Definition 21.14.** Consider the equation (1.2). Assume that it is non-degenerate, diagonally dominated, balanced and convergent; cf. Definition 21.1. If there is a  $q \in \{1, \ldots, Q\}$  such that  $\bar{\beta}_q = 0$ , say  $q_{\rm ts}$ , define the sets  $\mathcal{K}_{q_{\rm ts},\pm}$  by

$$\mathcal{K}_{q_{\rm ts},\pm} := \{ \kappa_{q_{\rm ts},\iota,\pm} \mid \iota \in \mathcal{I}_{\rm B}, \ \bar{\nu}_{q_{\rm ts}}(\iota) \neq 0 \}$$

where  $\kappa_{q_{ts},\iota,\pm}$  is defined in (21.36). Define  $\bar{\kappa}_{q_{ts},\pm}$  to be the limit of  $\kappa_{q_{ts},\iota,\pm}$  as  $\bar{\nu}_{q_{ts}}(\iota) \to \infty$  and define

$$\kappa_{q_{\rm ts},{\rm ts},\pm} := \sup \mathcal{K}_{q_{\rm ts},\pm}.$$

**Remark 21.15.** That the limits of  $\kappa_{q_{ts},\iota,\pm}$  as  $\bar{\nu}_{q_{ts}}(\iota) \to \infty$  exist is a consequence of the fact that the matrices  $A_{q_{ts},\infty}(\iota)$  are of the form (11.7); Lemma 11.6; the fact that (21.11) holds; and the fact that replacing V by -V in (11.7) does not affect the asymptotic values of the real parts of the eigenvalues of  $A_{\mu}$  (cf. Lemma 11.6). In fact, Lemma 11.6 implies that

$$\bar{\kappa}_{q_{\rm ts},+} = \frac{1}{2} \max\left\{ \kappa_{\rm max} \left( -\alpha_{\infty} + \tilde{X}_{\infty}^{q_{\rm ts}} \right), \kappa_{\rm max} \left( -\alpha_{\infty} - \tilde{X}_{\infty}^{q_{\rm ts}} \right) \right\},\\ \bar{\kappa}_{q_{\rm ts},-} = -\frac{1}{2} \min\left\{ \kappa_{\rm min} \left( -\alpha_{\infty} + \tilde{X}_{\infty}^{q_{\rm ts}} \right), \kappa_{\rm min} \left( -\alpha_{\infty} - \tilde{X}_{\infty}^{q_{\rm ts}} \right) \right\},$$

where we appealed to (21.11) and used the terminology introduced in Definition 4.3. In particular, we thus know that  $\bar{\kappa}_{q_{ts},\pm} = \kappa_{q_{ts},\pm}$ ; cf. Definition 21.2.

**Remark 21.16.** Due to the fact that the limits  $\bar{\kappa}_{q_{ts},\pm}$  exist and the fact that, given  $0 < K \in \mathbb{R}$ , there are only finitely many matrices  $A_{q_{ts},\infty}(\iota)$  for  $0 < \bar{\nu}_{q_{ts}}(\iota) \leq K$ , the sets  $\mathcal{K}_{q_{ts},\pm}$  are bounded. Thus  $\kappa_{q_{ts},ts,\pm}$  are finite numbers.

With the notation introduced in Definition 21.14, Lemmas 21.7 and 21.11 yield the following corollary.

**Corollary 21.17.** Consider the equation (1.2). Assume that it is non-degenerate, diagonally dominated, balanced and convergent; cf. Definition 21.1. Given a  $0 \neq \iota \in \mathcal{I}_{B}$  such that there is a sequence  $\{t_k\}$  of the type described in Definition 21.4, let  $I_r := [t_{\text{ini},r}, t_{\text{fin},r}]$  (or  $I_r := [t_{\text{ini},r}, \infty)$  in case  $t_{\text{fin},r} = \infty$ ), where  $t_{\text{ini},r}$  and  $t_{\text{fin},r}$  are introduced in Definition 21.4. Define  $R_{q,\pm}^+$  and  $R_{q,\pm}^-$  by (21.2) for  $q \in \{1, \ldots, Q\}$ , and define  $\kappa_{\pm}$  by (21.5). Then the following holds.

1. Assume that  $\bar{\beta}_q \neq 0$  for all  $q \in \{1, \ldots, Q\}$  and let  $\epsilon > 0$ . Then there is a constant  $C_{+,\epsilon}$  depending only on  $\epsilon$  and the coefficients of the equation (1.2), such that

$$|w(\iota, t_b)| \le C_{+,\epsilon} e^{(\kappa_+ + \epsilon)(t_b - t_a)} |w(\iota, t_a)| + C_{+,\epsilon} \int_{t_a}^{t_b} e^{(\kappa_+ + \epsilon)(t_b - t')} |\hat{f}(\iota, t')| dt'$$
(21.47)

for  $I_a := [t_a, t_b] \subseteq I_r$  and every solution z to (24.16) corresponding to  $\iota$ . Here w is defined in terms of z as described in (14.7) and (15.1). Moreover, the following improvements hold:

If q<sub>i</sub> ∈ {1,...,Q}, i = 1,...,l, are the only elements q ∈ {1,...,Q} such that I<sub>q</sub> ∩ I<sub>a</sub> ≠ Ø, then κ<sub>+</sub> appearing in (21.47) can be replaced by κ<sub>+,I<sub>a</sub></sub>, defined by

$$\kappa_{\pm,I_a} := \max_{1 \le i \le l} \kappa_{q_i,\pm}, \tag{21.48}$$

where the dependence of  $\kappa_{\pm,I_a}$  on  $\iota$  has been suppressed for the sake of brevity.

• If, in addition, the Jordan blocks of  $R_{q_i,+}^+$  and  $R_{q_i,+}^-$  corresponding to the eigenvalues with real part  $\kappa_{+,I_a}$  are trivial for all i = 1, ..., l, then  $\kappa_+$  appearing in (21.47) can be replaced by  $\kappa_{+,I_a}$ ; the parameter  $\epsilon$  can be replaced by zero; and  $C_{+,\epsilon}$  can be replaced by a constant depending only on the coefficients of the equation (1.2).

#### 21.4. ANALYSIS FOR ONE MODE

2. Assume that  $\bar{\beta}_q \neq 0$  for all  $q \in \{1, \ldots, Q\}$  and let  $\epsilon > 0$ . Then there is a constant  $C_{-,\epsilon}$  depending only on  $\epsilon$  and the coefficients of the equation (1.2), such that

$$|w(\iota, t_a)| \le C_{-,\epsilon} e^{-(\kappa_- + \epsilon)(t_a - t_b)} |w(\iota, t_b)| + C_{-,\epsilon} \int_{t_a}^{t_b} e^{-(\kappa_- + \epsilon)(t_a - t')} |\hat{f}(\iota, t')| dt'$$
(21.49)

for  $I_a := [t_a, t_b] \subseteq I_r$  and every solution z to (24.16) corresponding to  $\iota$ . Moreover, the following improvements hold:

- If  $q_i \in \{1, \ldots, Q\}$ ,  $i = 1, \ldots, l$ , are the only elements  $q \in \{1, \ldots, Q\}$  such that  $I_q \cap I_a \neq \emptyset$ , then  $\kappa_-$  appearing in (21.49) can be replaced by  $\kappa_{-,I_a}$  introduced in (21.48).
- If, in addition, the Jordan blocks of  $R_{q_i,-}^+$  and  $R_{q_i,-}^-$  corresponding to the eigenvalues with real part  $\kappa_{-,I_a}$  are trivial for all  $i = 1, \ldots, l$ , then  $\kappa_-$  appearing in (21.47) can be replaced by  $\kappa_{-,I_a}$ ; the parameter  $\epsilon$  can be replaced by zero; and  $C_{-,\epsilon}$  can be replaced by a constant depending only on the coefficients of the equation (1.2).
- 3. Assume that there is one  $q \in \{1, ..., Q\}$  such that  $\overline{\beta}_q = 0$ , say  $q_{ts}$ , and let  $\epsilon > 0$ . Then there is a constant  $C_{+,\epsilon}$  depending only on  $\epsilon$ , the spectrum of the Riemannian manifold corresponding to  $q_{ts}$  and the coefficients of the equation (1.2), such that

$$|w(\iota, t_b)| \le C_{+,\epsilon} e^{(\kappa_{\text{ts},+} + \epsilon)(t_b - t_a)} |w(\iota, t_a)| + C_{+,\epsilon} \int_{t_a}^{t_b} e^{(\kappa_{\text{ts},+} + \epsilon)(t_b - t')} |\hat{f}(\iota, t')| dt'$$
(21.50)

for  $I_a := [t_a, t_b] \subseteq I_r$  and every solution z to (24.16) corresponding to  $\iota$ , where

$$\kappa_{\mathrm{ts},\pm} := \max\{\kappa_{\pm}, \kappa_{q_{\mathrm{ts}},\mathrm{ts},\pm}\}\tag{21.51}$$

and the  $\kappa_{q,ts,\pm}$  are introduced in Definition 21.14. If  $I_{q_{ts}} \cap I_a = \emptyset$ , then the conclusions stated in case 1 above hold (including the improvements, and with the same dependence of the constants as in case 1). If  $I_{q_{ts}} \cap I_a \neq \emptyset$  and  $\kappa_{q_{ts},+} < \kappa_{ts,+}$ , then the  $\epsilon$  appearing in (21.50) can be removed if

- the Jordan blocks of  $R_{q,+}^+$  and  $R_{q,+}^-$  corresponding to the eigenvalues with real part  $\kappa_{ts,+}$  are trivial for all  $q \in \{1, \ldots, Q\}$ , and
- the Jordan blocks of  $A_{q_{ts},\infty}(\varsigma)$  corresponding to the eigenvalues with real part  $\kappa_{ts,+}$  are trivial for all  $\varsigma \in \mathcal{I}_B$  such that  $\bar{\nu}_{q_{ts}}(\varsigma) \neq 0$ .

When  $\epsilon$  can be removed,  $C_{+,\epsilon}$  can be replaced by a constant depending only on the spectrum of the Riemannian manifold corresponding to  $q_{ts}$  and the coefficients of the equation (1.2).

4. Assume that there is one  $q \in \{1, ..., Q\}$  such that  $\bar{\beta}_q = 0$ , say  $q_{ts}$ , and let  $\epsilon > 0$ . Then there is a constant  $C_{-,\epsilon}$  depending only on  $\epsilon$ , the spectrum of the Riemannian manifold corresponding to  $q_{ts}$  and the coefficients of the equation (1.2) such that

$$|w(\iota, t_a)| \le C_{-,\epsilon} e^{-(\kappa_{\text{ts}, -} + \epsilon)(t_a - t_b)} |w(\iota, t_b)| + C_{-,\epsilon} \int_{t_a}^{t_b} e^{-(\kappa_{\text{ts}, -} + \epsilon)(t_a - t')} |\hat{f}(\iota, t')| dt' \quad (21.52)$$

for  $I_a := [t_a, t_b] \subseteq I_r$  and every solution z to (24.16) corresponding to  $\iota$ , where  $\kappa_{ts,-}$  is defined by (21.51). If  $I_{q_{ts}} \cap I_a = \emptyset$ , then the conclusions stated in case 2 above hold (including the improvements, and with the same dependence of the constants as in case 2). If  $I_{q_{ts}} \cap I_a \neq \emptyset$ and  $\kappa_{q_{ts},-} < \kappa_{ts,-}$ , then the  $\epsilon$  appearing in (21.52) can be removed if

- the Jordan blocks of  $R_{q,-}^+$  and  $R_{q,-}^-$  corresponding to the eigenvalues with real part  $\kappa_{ts,-}$  are trivial for all  $q \in \{1, \ldots, Q\}$ , and
- the Jordan blocks of  $-A_{q_{ts},\infty}(\varsigma)$  corresponding to the eigenvalues with real part  $\kappa_{ts,-}$ are trivial for all  $\varsigma \in \mathcal{I}_B$  such that  $\bar{\nu}_{q_{ts}}(\varsigma) \neq 0$ .

When  $\epsilon$  can be removed,  $C_{-,\epsilon}$  can be replaced by a constant depending only on the spectrum of the Riemannian manifold corresponding to  $q_{ts}$  and the coefficients of the equation (1.2).

**Remark 21.18.** In cases 3 and 4, the  $\kappa_{\pm}$  which enters the definition of  $\kappa_{\text{ts},\pm}$ , cf. (21.51), can be replaced by  $\kappa_{\pm,I_a}$ ; cf. the improvements stated in cases 1 and 2. See also Corollary 21.25 for further improvements.

**Remark 21.19.** It is important to note that the constants appearing in the corollary depend neither on  $\iota$  nor on the solution z.

**Remark 21.20.** Recall that  $|w(\iota, t)|^2$  is given by

$$|w(\iota,t)|^{2} = |x_{1}(\iota,t)|^{2} + |y_{1}(\iota,t)|^{2} = |\dot{z}(\iota,t)|^{2} + \mathfrak{g}^{2}(\iota,t)|z(\iota,t)|^{2}, \qquad (21.53)$$

where z is a solution to (24.16). In this sense,  $|w(\iota, t)|^2$  represents the energy content of the  $\iota$ 'th mode at time t.

*Proof.* We begin by proving (21.47) in the case that both  $t_a$  and  $t_b$  are elements of the sequence  $\{t_k\}$ . Without loss of generality, we can then assume that  $t_a = t_0$  and that  $t_b = t_{l_b+1}$  for some  $0 \le l_b \in \mathbb{Z}$  (this can be achieved by simply relabelling the sequence). Let us begin by estimating

$$|w(t_k)| \le ||T_{\text{pre},k}^{-1}|| \cdot |w_{\text{pre},k}| \le C|w_{\text{fin},k}| = C|\psi_k|, \qquad (21.54)$$

$$|\psi_k| = |w_{\text{pre},k}| \le ||T_{\text{pre},k}|| \cdot |w(t_k)| \le C|w(t_k)|, \tag{21.55}$$

where C only depends on the coefficients of the equation (1.2). In order to arrive at this conclusion, we used the fact that  $\psi_k = w_{\text{fin},k}$ , (15.36), (15.53), the fact that  $D_k$  is an isometry and the fact that the norms of  $T_{\text{pre},k}$  and its inverse are bounded by a constant depending only on the coefficients of the equation (1.2); cf. the comments made in connection with (15.34) and (15.35). On the other hand, recalling (15.64),

$$|\psi_{k+1}| \leq ||A_k^+ \cdots A_0^+|| \cdot |\psi_0| + ||A_k^+|| \int_{t_k}^{t_{k+1}} |\hat{F}_k(t)| dt + \cdots + ||A_k^+ \cdots A_0^+|| \int_{t_0}^{t_1} |\hat{F}_0(t)| dt.$$
(21.56)

Estimating the norm of the matrix products in the non-transparent setting. In order to proceed, we need to estimate  $||A_{k_1}^+ \cdots A_{k_0}^+||$  in cases where  $t_k$  does not necessarily belong to the same frequency era for all  $k_0 \leq k \leq k_1$ . For simplicity, we start by considering the case that there are no transparent eras. Given  $k_0, k_1$  such that  $0 \leq k_0 \leq k_1 \leq l_b < k_{\text{fin},r}$ , there are, say, lsubintervals, denoted  $[k_{0,j}, k_{1,j}], j = 1, \ldots, l$ , such that  $t_k \in I_{q_j}$  for  $k \in [k_{0,j}, k_{1,j}]$  and some choice of  $q_j$ . Moreover,  $k_{0,1} = k_0, k_{1,l} = k_1$  and  $k_{1,j} + 1 = k_{0,j+1}$ . We also assume that none of the intervals  $I_{q_j}$  is a transparent era. Fix  $\epsilon > 0$ . Since  $l \leq Q$ , the following estimate holds:

$$\|A_{k_1}^+ \cdots A_{k_0}^+\| \le \|A_{k_{1,l}}^+ \cdots A_{k_{0,l}}^+\| \cdots \|A_{k_{1,1}}^+ \cdots A_{k_{0,1}}^+\| \le C_{+,\epsilon} e^{(\kappa_+ + \epsilon)(\bar{t}_1 - \bar{t}_0)},\tag{21.57}$$

where  $\bar{t}_1 = t_{k_1+1}$ ,  $\bar{t}_0 = t_{k_0}$  and the constant  $C_{+,\epsilon}$  depends only on  $\epsilon$  and the coefficients of the equation (1.2). In order to obtain the last inequality in (21.57), we appealed to (21.9)  $l \leq Q$  times, as well as Remark 21.10; note that  $C_{\deg,q_{j},+} = 0$  for all  $j = 1, \ldots, l$ , using the notation of (21.9). Assume now that there are no non-trivial Jordan blocks of  $R_{q,+}^+$  and  $R_{q,+}^-$ ,  $q = q_j$ ,  $j = 1, \ldots, l$ , corresponding to eigenvalues with real part  $\kappa_+$ . For a given j, this situation can arise for two reasons. The first possibility is that  $\kappa_+ = \kappa_{q_j,+}$ , but the Jordan blocks of  $R_{q,+}^+$  and  $R_{q_j,+}^-$  and  $R_{q_j,+}^-$  corresponding to eigenvalues with real part  $\kappa_{q_j,+}$  are trivial. For such j, we can appeal to (21.7) with  $C_{\deg,q_j,+} = 0$ . The second possibility is that  $\kappa_+ > \kappa_{q_j,+}$ . In that case, we can appeal to (21.9) with  $\epsilon = \kappa_+ - \kappa_{q_j,+}$  and  $R_{q,+}^-$ ,  $q = q_j$ ,  $j = 1, \ldots, l$ , corresponding to eigenvalues of  $R_{q,+}^+$  and  $r_{q,+}$  are trivial. For such j, we can appeal to (21.9) with  $\epsilon = \kappa_+ - \kappa_{q_j,+}$  and  $R_{q,+}^ q = q_j$ ,  $j = 1, \ldots, l$ , corresponding to eigenvalues of  $R_{q,+}^+$  and  $r_{q,+}^ q = q_j$ ,  $j = 1, \ldots, l$ , corresponding to eigenvalues of  $r_{q,+}^+$  and  $r_{q,+}^ r_{q,+}^-$  and  $r_{q,+}^ r_{q,+}^-$  and  $r_{q,+}^ r_{q,+}^ r_{q,+$ 

#### 21.4. ANALYSIS FOR ONE MODE

with real part  $\kappa_+$ , then the  $\epsilon$  appearing in (21.57) can be removed and  $C_{+,\epsilon}$  can be replaced by a constant  $C_+$  depending only on the coefficients of the equation (1.2). From the above arguments, it is also clear that  $\kappa_+$  can be replaced by  $\kappa_{+,I_{k_0,k_1}}$ , where  $I_{k_0,k_1} := [\bar{t}_0, \bar{t}_1]$ .

**Deriving energy estimates over intervals defined by the sequence.** Returning to Lemma 15.11, in particular (15.55), it is clear that

$$|F_l(t)| \le C|f(t)|$$

in the interval  $[t_l, t_{l+1}]$ , where C only depends on the coefficients of the equation (1.2). Combining this estimate with (21.54), (21.56) and (21.57) yields

$$|w(t_{k+1})| \leq C_{+,\epsilon} e^{(\kappa_{+}+\epsilon)(t_{k+1}-t_{0})} |\psi_{0}| + C_{+,\epsilon} e^{(\kappa_{+}+\epsilon)(t_{k+1}-t_{k})} \int_{t_{k}}^{t_{k+1}} |\hat{f}(t)| dt + \cdots + C_{+,\epsilon} e^{(\kappa_{+}+\epsilon)(t_{k+1}-t_{0})} \int_{t_{0}}^{t_{1}} |\hat{f}(t)| dt$$

$$\leq C_{+,\epsilon} e^{(\kappa_{+}+\epsilon)(t_{k+1}-t_{0})} |\psi_{0}| + C_{+,\epsilon} \int_{t_{0}}^{t_{k+1}} e^{(\kappa_{+}+\epsilon)(t_{k+1}-t)} |\hat{f}(t)| dt,$$
(21.58)

where  $C_{+,\epsilon}$  depends only on  $\epsilon$  and the coefficients of the equation (1.2). In order to obtain this estimate, we used the fact that  $e^{(\kappa_++\epsilon)(t-t_k)} \leq C_{\epsilon}$  for  $t \in [t_k, t_{k+1}]$ , where  $C_{\epsilon}$  depends only on  $\epsilon$  and the coefficients of the equation (1.2). Combining (21.55) with (21.58) yields the conclusion that (21.47) holds for  $t_a = t_{k_0}$  and  $t_b = t_{k_1+1}$ , where  $k_{\text{ini},\text{r}} < k_0 \leq k_1 < k_{\text{fn},\text{r}}$ . Moreover, if there are no non-trivial Jordan blocks of  $R_{q,+}^+$  and  $R_{q,+}^-$ ,  $q = 1, \ldots, Q$ , corresponding to eigenvalues with real part  $\kappa_+$ , then the  $\epsilon$  appearing in (21.47) can be removed and  $C_{+,\epsilon}$  can be replaced by a constant  $C_+$  depending only on the coefficients of the equation (1.2). Finally, the stated improvements also hold.

**Extending the energy estimate beyond the time sequence.** In general, we can assume that  $t_{k_0-1} < t_a \leq t_{k_0}$  and that  $t_{k_1+1} \leq t_b < t_{k_1+2}$ , where  $k_1 + 2 \leq k_{\text{fin,r}}$ . We then need to estimate the change of the solution in the interval  $[t_a, t_{k_0}]$  and in the interval  $[t_{k_1+1}, t_b]$ . In order to obtain such estimates, it is sufficient to appeal to Lemma 10.7. In fact, due to Lemma 20.3, we know that (1.2) is oscillation adapted with  $f_{\text{sh}}$ ,  $f_X$  and  $f_{\text{ode}}$  given by (20.15). Moreover, in (20.15), C only depends on the coefficients of the equation (1.2). Thus Lemma 10.7 applies (with  $\mathfrak{e} = 0$ ). Appealing to (10.12) with  $t_0$  replaced by  $t_{k_1+1}$  and  $t_1$  replaced by  $t_b$  yields

$$|w(t_b)| \le C|w(t_{k_1+1})| + C \int_{t_{k_1+1}}^{t_b} |\hat{f}(t')| dt', \qquad (21.59)$$

where C only depends on the coefficients of the equation (1.2). In order to obtain this conclusion, we used the fact that |w| and  $\mathcal{E}^{1/2}$  are equivalent in the interval of interest, the constant of equivalence being numerical; cf. (10.11), (21.53) and Definition 21.4. A similar estimate holds in the interval  $[t_a, t_{k_0}]$ . Combining these estimates with (21.58) and the fact that  $|\psi_{k_0}|$  can be bounded by a constant times  $|w(t_{k_0})|$  yields (21.47). Moreover, the improvements stated in case 1 also hold.

Energy estimates in the presence of transparent eras. Assume that (1.2) is such that there is a  $q \in \{1, \ldots, Q\}$  with the property that  $\bar{\beta}_q = 0$ , say  $q_{ts}$ . Let  $I_a$  be as in the statement of the corollary. If  $I_{q_{ts}} \cap I_a = \emptyset$ , then the previous analysis applies, so that the conclusions of case 1 hold. In particular, (21.50) holds. When  $I_{q_{ts}} \cap I_a \neq \emptyset$ , either (21.7) or (21.9) holds in intervals of the form  $[k_{0,j}, k_{1,j}]$ ; cf. the division made in connection with (21.57). In the case of (21.9), we are free to replace  $\epsilon$  by  $\epsilon/2$ , and we do so in what follows. In both (21.7) and (21.9), the constant  $C_{\deg,q_{ts},+}$  appears (which only depends on the coefficients of (1.2) and  $\epsilon$ ). This constant gives rise to a "cut off" in  $\bar{\nu}_{q_{ts}}(\iota)$ . In fact, define

If  $\bar{\nu}_{q_{ts}}(\iota) \geq K_{co,\epsilon,+}$ , then (21.7) and (21.9) yield estimates which are good enough that the previous analysis can be applied. Summing up, there are, given  $\iota$  and  $\{t_k\}$  as above, three cases to consider. Either there is no transparent era corresponding to  $\iota$  (or  $I_{q_{ts}} \cap I_a = \emptyset$ ). In that case, the analysis in the absence of transparent eras applies. The second possibility is that  $I_{q_{ts}} \cap I_a \neq \emptyset$ , but that  $\bar{\nu}_{q_{ts}}(\iota) \geq K_{co,\epsilon,+}$ . In that case, (21.7) and (21.9) yield estimates which are good enough that the analysis in the absence of transparent eras applies. The third possibility is that  $I_{q_{ts}} \cap I_a \neq \emptyset$  and  $\bar{\nu}_{q_{ts}}(\iota) \leq K_{co,\epsilon,+}$ . This situation is one we need to consider in greater detail.

Analysis for low frequency transparent eras. Assume that  $I_{q_{ts}} \cap I_a \neq \emptyset$  and  $\bar{\nu}_{q_{ts}}(\iota) \leq K_{co,\epsilon,+}$ . Divide  $I_a$  into three parts:  $I_{a,1} := [t_a, \bar{t}_0], I_{a,2} := [\bar{t}_0, \bar{t}_1]$  and  $I_{a,3} := [\bar{t}_1, t_b]$ , where  $\bar{t}_0$  is the infimum of  $I_{q_{ts}} \cap I_a$  and  $\bar{t}_1$  is the supremum of  $I_{q_{ts}} \cap I_a$ . It could be that one or two of the intervals  $I_{a,j}$  consists of one point. In that case, the corresponding interval(s) can be ignored in the analysis. Here we assume that all the intervals  $I_{a,j}$  have non-empty interior, and leave the remaining cases to the reader. In the interval  $I_{a,1}$ , the analysis in the absence of transparent eras applies. Thus (21.47) (including the improvements) holds when  $t_b$  is replaced by  $\bar{t}_0$ ; i.e.

$$|w(\iota, \bar{t}_0)| \le C_{+,\epsilon} e^{(\kappa_+ + \epsilon)(\bar{t}_0 - t_a)} |w(\iota, t_a)| + C_{+,\epsilon} \int_{t_a}^{\bar{t}_0} e^{(\kappa_+ + \epsilon)(\bar{t}_0 - t')} |\hat{f}(\iota, t')| dt',$$
(21.61)

where  $C_{+,\epsilon}$  only depends on  $\epsilon$  and the coefficients of the equation (1.2). In order to estimate  $|w(\iota, t)|$  for  $t \in I_{a,2}$ , we appeal to Lemma 21.11 with  $K_{\rm co}$  replaced by  $K_{{\rm co},\epsilon,+}$  defined in (21.60). Then the constant  $C_{{\rm csp},\epsilon}$  appearing in (21.37) only depends on  $\epsilon$ , the spectrum of the Riemannian manifold corresponding to  $q_{\rm ts}$  and the coefficients of the equation (1.2). Combining (21.61) and (21.37) (in the latter estimate, we replace  $t_-$  with  $\bar{t}_0$ ) yields

$$\begin{aligned} |w(\iota,t)| &\leq C_{\mathrm{csp},\epsilon} e^{(\kappa_{q_{\mathrm{ts},\iota,+}}+\epsilon)(t-\bar{t}_0)} |w(\iota,\bar{t}_0)| + C_{\mathrm{csp},\epsilon} \int_{\bar{t}_0}^t e^{(\kappa_{q_{\mathrm{ts},\iota,+}}+\epsilon)(t-t')} |\hat{f}(\iota,t')| dt' \\ &\leq C_{+,\epsilon} e^{(\kappa_{\mathrm{ts},+}+\epsilon)(t-t_a)} |w(\iota,t_a)| + C_{+,\epsilon} \int_{t_a}^t e^{(\kappa_{\mathrm{ts},+}+\epsilon)(t-t')} |\hat{f}(\iota,t')| dt' \end{aligned}$$

for  $t \in I_{a,2}$ , where  $C_{+,\epsilon}$  only depends on  $\epsilon$ , the spectrum of the Riemannian manifold corresponding to q and the coefficients of the equation (1.2). Thus (21.50) holds for  $t \in I_{a,1} \cup I_{a,2}$ . In order to extend this estimate to all of  $I_a$ , it is sufficient to appeal to (21.47) in  $I_{a,3}$ . Thus (21.50) holds.

**Removing the**  $\epsilon$ . What remains to be verified is that the  $\epsilon$  appearing in (21.50) can be removed under the circumstances mentioned in the statement of the corollary. Due to the assumption that the Jordan blocks of  $R_{q,+}^+$  and  $R_{q,+}^-$  corresponding to the eigenvalues with real part  $\kappa_{ts,+}$  are trivial for all  $q \in \{1, \ldots, Q\}$ , it is clear that the above analysis in the absence of transparent eras can be modified in that  $\kappa_+$  can be replaced by  $\kappa_{ts,+}$  and the  $\epsilon$  can be removed (the constants are in that case also independent of  $\epsilon$ ). In the presence of transparent eras, we may need to appeal to (21.7) or (21.9). However, we only appeal to these estimates when  $\bar{\nu}_{q_{ts}}(\iota)$  is large. Moreover, how to make a choice of what "large" means is clear in the present setting. In fact, since  $\kappa_{q_{ts},+} < \kappa_{ts,+}$ by assumption, we first fix an  $0 < \epsilon < (\kappa_{ts,+} - \kappa_{q_{ts},+})/2$  (when necessary) and then choose  $K_{co,+}$ according to

$$K_{\rm co,+} := \frac{2C_{\deg,q_{\rm ts},+}}{\kappa_{\rm ts,+} - \kappa_{q_{\rm ts},+}}.$$
(21.62)

Then, for  $\bar{\nu}_{q_{ts}}(\iota) \geq K_{co,+}$ , the exponential factors appearing on the right hand sides of (21.7) and (21.9) can be replaced by  $\exp[\kappa_{ts,+}(\bar{t}_{1,+}-\bar{t}_{0,+})]$ . What remains to be taken into consideration is the finite number of matrices of the form (21.34) that occur for  $\bar{\nu}_{q_{ts}}(\iota) \leq K_{co,+}$ . For those with the property that  $\kappa_{q_{ts},\iota,+} < \kappa_{ts,+}$ , it is sufficient to choose  $\epsilon = \kappa_{ts,+} - \kappa_{q_{ts},\iota,+}$  in (21.37). Then the expression  $\kappa_{q_{ts},\iota,+} + \epsilon$  appearing (twice) on the right hand side of (21.37) can be replaced by  $\kappa_{ts,+}$ . For those matrices  $A_{q_{ts},\infty}(\varsigma)$  for which  $\kappa_{q_{ts},\iota,+} = \kappa_{ts,+}$ , the Jordan blocks of  $A_{q_{ts},\infty}(\varsigma)$ corresponding to the eigenvalues with real part  $\kappa_{ts,+}$  are trivial. In such cases, the  $\epsilon$  appearing on the right hand side of (21.37) can thus be removed. To summarise, in all the estimates needed to obtain the desired conclusion, the  $\epsilon$  can be removed. All the statements of the corollary concerning cases 1 and 3 follow. Analysis in the past direction. The analysis in the past direction is analogous to the above. We therefore leave the details of the proof to the reader.  $\Box$ 

#### 21.4.1 Refinements

Even though Corollary 21.17 is sufficient for many applications, more refined estimates are sometimes needed. In preparation for the statement of such results, it is convenient to introduce the following notation.

**Definition 21.21.** Consider the equation (1.2). Assume that it is non-degenerate, diagonally dominated, balanced and convergent; cf. Definition 21.1. Fix  $\epsilon > 0$  and K > 0. Given  $0 \neq \iota \in \mathcal{I}_{B}$ , consider an interval J = [s, t], where  $0 \leq s \leq t$ . If J is a subset of one frequency era, say  $I_{q_1}$ , let k := 0 and  $I_1 := J$ . Otherwise, let  $s_1$  be the supremum of the set of  $t' \in [s, t]$  such that s and t' belong to the same frequency era, say  $I_{q_1}$ . If  $[s_1, t]$  is contained in one frequency era, say  $I_{q_2}$ , let k := 1,  $I_1 := [s, s_1]$  and  $I_2 := [s_1, t]$ . If  $[s_1, t]$  is not contained in one frequency era, let  $k \geq 2$  and  $s_j$ ,  $j = 2, \ldots, k$ , be defined by the conditions that  $s \leq s_1 \leq \cdots \leq s_k \leq t$ ; that for each  $j = 1, \ldots, k - 1$ ,  $I_{j+1} := [s_j, s_{j+1}]$  equals a frequency era  $I_{q_{j+1}}$  for some  $q_{j+1} \in \{1, \ldots, Q\}$ ; that  $I_1 := [s, s_1]$  is a subset of a frequency era  $I_{q_1}$ ; and that  $I_{k+1} := [s_k, t]$  is a subset of a frequency era  $I_{q_{k+1}}$ . Define

$$\varrho_{\epsilon,K,+}(t,s) := \sum_{j=1}^{k+1} \mu_{j,+} |I_j|,$$

where  $\mu_{j,\pm} := \kappa_{q_j,\pm}$  in case  $q_j \neq 0$  and the Jordan blocks of  $R_{q_j,\pm}^+$  and  $R_{q_j,\pm}^-$  corresponding to eigenvalues with real part  $\kappa_{q_j,\pm}$  are trivial;  $\mu_{j,\pm} := \kappa_{q_j,\pm} + \epsilon$  in case  $q_j \neq 0$  and there is a non-trivial Jordan block of either  $R_{q_j,\pm}^+$  or  $R_{q_j,\pm}^-$  corresponding to an eigenvalue with real part  $\kappa_{q_j,\pm}$ ;  $\mu_{j,\pm} := \kappa_{q_j,\iota,\pm}$  if  $q_j = 0$ ,  $\bar{\nu}_{q_j}(\iota) \leq K$  and the Jordan blocks of  $\pm A_{q_j,\infty}(\iota)$  corresponding to eigenvalues with real part  $\kappa_{q_j,\iota,\pm}$  are trivial; and  $\mu_{j,\pm} := \kappa_{q_j,\iota,\pm} + \epsilon$  otherwise. Similarly,

$$\varrho_{\epsilon,K,-}(s,t) := \sum_{j=1}^{k+1} \mu_{j,-} |I_j|.$$

**Remark 21.22.** If s = t and t belongs to two frequency eras, some parts of the definition are ambiguous. However, this does not affect the end result.

**Remark 21.23.** When there are no transparent eras, the constant K is irrelevant, and the definition can be simplified. In this case, we also use the notation  $\rho_{\epsilon,\pm}$ .

**Remark 21.24.** The functions  $\rho_{\epsilon,K,\pm}$  depend on  $\iota \neq 0$ . However, for the sake of brevity, we omit reference to this dependence. Note also that we only define  $\rho_{\epsilon,K,+}(t,s)$  for  $0 \leq s \leq t$  and  $\rho_{\epsilon,K,-}(t,s)$  for  $0 \leq t \leq s$ .

With this notation, Corollary 21.17 can be improved as follows.

**Corollary 21.25.** Consider the equation (1.2). Assume that it is non-degenerate, diagonally dominated, balanced and convergent; cf. Definition 21.1. Given  $\epsilon > 0$  and K > 0, there is then a constant  $C_{\epsilon,K}$ , depending only on  $\epsilon$ , K, the spectrum of the Riemannian manifold corresponding to the  $q_{ts} \in \{1, \ldots, Q\}$  such that  $\bar{\beta}_{q_{ts}} = 0$  (in case there is such a  $q_{ts}$ ) and the coefficients of the equation, such that the following holds. Given a  $0 \neq \iota \in \mathcal{I}_B$  such that there is a sequence  $\{t_k\}$  of the type described in Definition 21.4, let  $I_r := [t_{ini,r}, t_{fin,r}]$  (or  $I_r := [t_{ini,r}, \infty)$  in case  $t_{fin,r} = \infty$ ), where  $t_{ini,r}$  and  $t_{fin,r}$  are introduced in Definition 21.4. Let  $0 \leq t_a \leq t_b$  be such that  $I_a := [t_a, t_b] \subseteq I_r$ . Then

$$w(\iota, t_b)| \le C_{\epsilon, K} e^{\varrho_{\epsilon, K, +}(t_b, t_a)} |w(\iota, t_a)| + C_{\epsilon, K} \int_{t_a}^{t_b} e^{\varrho_{\epsilon, K, +}(t_b, t')} |\hat{f}(\iota, t')| dt',$$
(21.63)

$$|w(\iota, t_a)| \le C_{\epsilon, K} e^{\varrho_{\epsilon, K, -}(t_a, t_b)} |w(\iota, t_b)| + C_{\epsilon, K} \int_{t_a}^{t_b} e^{\varrho_{\epsilon, K, -}(t_a, t')} |\hat{f}(\iota, t')| dt'$$
(21.64)

for all solutions z to (24.16).

**Remark 21.26.** The constant K should be thought of as a frequency cut off. Considering matrices of the form (21.34), we can only carry out a detailed analysis (transforming the matrix to Jordan normal form etc.) for a finite number of matrices. Here this finite number is determined by the bound  $\bar{\nu}_{q_{ts}}(\iota) \leq K$  (where  $q_{ts}$  is defined by the condition that  $\bar{\beta}_{q_{ts}} = 0$ ); cf. Definition 21.21.

**Remark 21.27.** In case there is no transparent era with non-empty intersection with  $I_a$ , then  $C_{\epsilon,K}$  can be replaced by a constant depending only on  $\epsilon$  and the coefficients of the equation (1.2). In that setting, we also use the notation  $\varrho_{\epsilon,\pm}$  instead of  $\varrho_{\epsilon,K,\pm}$ .

**Remark 21.28.** When the Jordan blocks of  $R_{q,+}^+$  and  $R_{q,+}^-$  corresponding to eigenvalues with real part  $\kappa_{q,+}$  are trivial for all  $q \in \{1, \ldots, Q\}$  such that  $I_q \cap I_a \neq \emptyset$ , and there is no transparent era with non-empty intersection with  $I_a$ , then the  $C_{\epsilon,K}$  appearing in (21.63) can be replaced by a constant depending only on the coefficients of the equation (1.2). A similar statement concerning (21.64) holds.

*Proof.* Let us begin by proving (21.63) in the case that there is no transparent era with non-empty intersection with  $I_a$ . Given  $t_a$  and  $t_b$  as in the statement of the lemma, define the intervals  $I_j$  as in Definition 21.21, where s and t have been replaced by  $t_a$  and  $t_b$  respectively. Let, moreover,  $s_j$ ,  $j = 1, \ldots, k$ , be defined as in Definition 21.21. If k = 0, i.e.,  $I_1 = I_a$ , then Corollary 21.17 yields

$$|w(t_b)| \le C_{\epsilon} e^{(\kappa_{q_1,+}+\epsilon)(t_b-t_a)} |w(t_a)| + C_{\epsilon} \int_{t_a}^{t_b} e^{(\kappa_{q_1,+}+\epsilon)(t_b-t')} |\hat{f}(t')| dt',$$
(21.65)

where  $C_{\epsilon}$  only depends on  $\epsilon$  and the coefficients of the equation (1.2). Moreover, in case the Jordan blocks of  $R_{q_1,+}^+$  and  $R_{q_1,+}^-$  corresponding to the eigenvalues with real part  $\kappa_{q_1,+}$  are trivial, the parameter  $\epsilon$  can be replaced by zero, and  $C_{\epsilon}$  can be replaced by a constant independent of  $\epsilon$ . Thus (21.63) holds if k = 0, i.e., if  $I_1 = I_a$ . Let us now assume, inductively, that (21.63) holds if the number of intervals  $I_j$  into which  $I_a$  is divided (when appealing to Definition 21.21) is bounded from above by k. Assume now that  $I_a$  is divided into k + 1 intervals. Then

$$\begin{split} |w(t_{b})| &\leq C_{\epsilon} e^{(\kappa_{q_{k+1},+}+\epsilon)(t_{b}-s_{k})} |w(s_{k})| + C_{\epsilon} \int_{s_{k}}^{t_{b}} e^{(\kappa_{q_{k+1},+}+\epsilon)(t_{b}-t')} |\hat{f}(t')| dt' \\ &\leq C_{\epsilon} \exp\left[(\kappa_{q_{k+1},+}+\epsilon)(t_{b}-s_{k}) + \varrho_{\epsilon,+}(s_{k},t_{a})\right] |w(t_{a})| \\ &+ C_{\epsilon} \int_{t_{a}}^{s_{k}} \exp\left[(\kappa_{q_{k+1},+}+\epsilon)(t_{b}-s_{k}) + \varrho_{\epsilon,+}(s_{k},t')\right] |\hat{f}(t')| dt' \\ &+ C_{\epsilon} \int_{s_{k}}^{t_{b}} e^{(\kappa_{q_{k+1},+}+\epsilon)(t_{b}-t')} |\hat{f}(t')| dt', \end{split}$$

where we appealed to Corollary 21.17 in the first step and the inductive hypothesis in the second step. Moreover, the  $\epsilon$  can be removed under circumstances analogous to those described in connection with (21.65). By induction, we conclude that (21.63) holds in the absence of transparent eras. Moreover, the  $\epsilon$  and K can both be removed under the circumstances described in Remark 21.28. In each step of the induction process, the constant appearing in the estimate grows. However, this does not lead to problems, since the number of steps is bounded by Q.

**Presence of a transparent era.** In case there is a transparent era, we first assume K to be larger than the  $K_{co,\epsilon,+}$  appearing in (21.60). Moreover, we assume that

$$|\kappa_{q_{\rm ts},\iota,+} - \kappa_{q_{\rm ts},+}| \le \epsilon/4$$

for  $\iota$  such that  $\bar{\nu}_{q_{ts}}(\iota) \geq K$ . Note that we can make these assumptions without loss of generality. Next we wish to use an argument similar to the one presented in connection with (21.60). In particular, we wish to appeal to (21.7) or (21.9). When doing so, we here assume the  $\epsilon$  appearing in (21.9) to have been replaced by  $\epsilon/4$ . Given these assumptions, an argument similar to the one presented in connection with (21.60) yields an estimate for transparent eras such that  $\bar{\nu}_{q_{ts}}(\iota) \geq K$  which is good enough that we can argue as in the absence of transparent eras in order to obtain the desired conclusion. What remains is to consider transparent eras such that  $\bar{\nu}_{q_{ts}}(\iota) \leq K$ . However, the relevant estimate is in that case supplied by Lemma 21.11. This finishes the proof of (21.63), including the improvements described in Remarks 21.27 and 21.28.

The proof of (21.64) is similar, and we leave the details to the reader.

#### 21.5 The case of unbounded frequency eras

Even though Lemma 21.7 yields an essentially optimal estimate (see Subsection 21.6 below), there is room for improvement in the case that there is an unbounded non-transparent era  $I_q$ . In fact, we have the following result.

**Lemma 21.29.** Consider the equation (1.2). Assume that it is non-degenerate, diagonally dominated, balanced and convergent; cf. Definition 21.1. Fix  $0 \neq \iota \in \mathcal{I}_B$  and assume that  $I_q$  is an unbounded and non-transparent era containing  $[t_0, \infty)$  for some  $q \in \{1, \ldots, Q\}$  and  $t_0 \geq 0$ . Assume, moreover, that (14.22) holds with  $t_a$  replaced by  $t_0$  and that (21.6) holds. Assume, finally, that if  $\{t_k\}$  is the sequence given by Definition 21.4 and starting at  $t_0$ , then  $t_{\text{fin},r} = \infty$ . Define  $R_{q,\pm}^+$  and  $R_{q,\pm}^-$  by (21.2); define  $\kappa_{q,\pm}$  by (21.3); and define  $d_{q,\pm}$  by (21.4). Then there is a constant C depending only on the spectrum of the Riemannian manifold corresponding to q and the coefficients of the equation (1.2), such that

$$|w(\iota,t)| \le C e^{\kappa_{q,+}\bar{t}} \langle \bar{t} \rangle^{d_{q,+}-1} |w(\iota,t_a)| + C \int_{t_a}^t \langle t-t' \rangle^{d_{q,+}-1} e^{\kappa_{q,+}(t-t')} |\hat{f}(\iota,t')| dt'$$
(21.66)

for all  $t_0 \leq t_a \leq t$  and all solutions z to (24.16), where  $\bar{t} = t - t_a$ .

**Remark 21.30.** If  $\bar{\beta}_q > 0$ , it should be possible to remove the following assumptions: that (14.22) holds with  $t_a$  replaced by  $t_0$ ; that (21.6) holds; and that  $t_{\text{fin,r}} = \infty$ . The reason for this is that  $\bar{\beta}_q > 0$  implies that  $\mathfrak{g}(\iota, t) \to \infty$  as  $t \to \infty$ . Replacing  $t_0$  by a later time, say  $t_b \ge t_0$ , all of the mentioned assumptions are therefore satisfied. On the other hand, the length of the interval  $[t_0, t_b]$  is bounded by a constant depending only on the spectrum of the Riemannian manifold corresponding to q and the coefficients of the equation (1.2). Deriving a bound for the interval  $[t_0, t_b]$  should therefore not be a problem; cf. Lemma 10.7.

*Proof.* Keeping the observations made in Subsection 15.1.1, in particular (15.15), in mind, it is clear that in order to obtain (21.66), it is sufficient to prove that

$$\|\Phi_{\iota}(t;t')\| \le C\langle t-t'\rangle^{d_{q,+}-1} e^{\kappa_{q,+}(t-t')}$$
(21.67)

for all  $t_0 \leq t' \leq t$ , where C depends only on the spectrum of the Riemannian manifold corresponding to q and the coefficients of the equation (1.2). On the other hand, in order to prove (21.67), it is sufficient to prove that

$$|w(\iota, t)| \le C\langle t - t'\rangle^{d_{q, +} - 1} e^{\kappa_{q, +}(t - t')} |w(\iota, t')|$$
(21.68)

for all  $t_0 \leq t' \leq t$  and all solutions w to the homogeneous version of (15.14). Moreover, C should have the same dependence as in the case of (21.67). The remainder of the proof is devoted to demonstrating (21.68).

**Preliminaries.** To begin with, note that since the frequency era  $I_q$  contains  $[t_0, \infty)$  and since  $t_{\text{fin},r} = \infty$ , it is clear that  $\bar{\beta}_q > 0$  and that  $\bar{\beta}_q > \bar{\beta}_p$  for all  $p \neq q$  such that  $\bar{\nu}_p(\iota) \neq 0$ . Let  $\{t_k\}$  be a sequence of the type described in Definition 21.4, starting at  $t_0$ .

**Improving the matrix estimates.** Consider the proof of Lemma 21.7. Since the assumptions of the present lemma imply that the assumptions of Lemma 21.7 are satisfied, we know that (21.28)

and (21.29) hold. Consider the right hand side of (21.28). The first and the last term inside the parenthesis decay exponentially (since  $\bar{\beta}_q > 0$ ). Let us consider the middle term. Due to (20.32),

$$\mathfrak{H}_{q}^{(s)}(\iota, t_{k}) = \sum_{p \neq q} \frac{\bar{\nu}_{p}^{s}(\iota) e^{s\beta_{p}t_{k}}}{\bar{\nu}_{q}^{s}(\iota) e^{s\bar{\beta}_{q}t_{k}}} = \sum_{p \neq q} \frac{\bar{\nu}_{p}^{s}(\iota) e^{s\beta_{p}t_{0}}}{\bar{\nu}_{q}^{s}(\iota) e^{s\bar{\beta}_{q}t_{0}}} e^{-s(\bar{\beta}_{q} - \bar{\beta}_{p})(t_{k} - t_{0})} \le (Q - 1)e^{-s\beta_{\mathrm{dm}}\bar{t}_{k}}$$
(21.69)

for  $t_k \geq t_0$ , where s > 0;  $\beta_{dm}$  is the difference between  $\bar{\beta}_q$  and the second largest  $\bar{\beta}_p$  for which  $\bar{\nu}_p(\iota) \neq 0$ ;  $\bar{t}_k = t_k - t_0$ ; and we have used the fact that (by the definition of  $t_0$ )  $\bar{\nu}_p(\iota) e^{\bar{\beta}_p t_0} \leq \bar{\nu}_q(\iota) e^{\bar{\beta}_q t_0}$  for all  $p \in \{1, \ldots, Q\}$ . Thus (21.28) yields

$$\|\check{A}_{k,q,\epsilon}^{\pm} - \check{A}_{k,q,\epsilon}^{\mathrm{app},\pm}\| \le \frac{C_{\epsilon,\lambda}}{\bar{\nu}_q(\iota)e^{\bar{\beta}_q t_k}}e^{-\eta_{\mathrm{dm}}\bar{t}_k}$$
(21.70)

for  $t_k \geq t_0$ , where the constant  $C_{\epsilon,\lambda}$  only depends on  $\epsilon$ ,  $\lambda$ , the spectrum of the Riemannian manifold corresponding to q and the coefficients of the equation (1.2);  $\eta_{\rm dm} := \min\{\eta_A, \beta_{\rm dm}, \bar{\beta}_q\}$ ;  $\check{A}_{k,q,\epsilon}^{\rm app,\pm}$  is given by (21.29); and  $\check{A}_{k,q,\epsilon}^{\pm}$  is defined via (21.20), (21.23) and (21.25), where  $\lambda$  remains to be determined. Considering the proof of Lemma 21.7, it is clear that by choosing  $\lambda = \kappa_{q,+}$  and  $\epsilon$  small enough (the bound depending only on the difference between  $\lambda = \kappa_{q,+}$  and the second largest real part of an eigenvalue of the matrix  $R_{\infty,q}(\iota)$  introduced in (21.16)), we can write (21.29) as

$$\check{A}_{k,q,\epsilon}^{\mathrm{app},+} := \mathrm{Id}_{2m} + \frac{2\pi}{\bar{\nu}_q(\iota)e^{\bar{\beta}_q t_k}} J_R,\tag{21.71}$$

where  $J_R \in \mathbf{M}_{2m}(\mathbb{R})$  consists of generalised Jordan blocks (cf. the terminology introduced in Remark 9.10) such that the blocks corresponding to strictly negative eigenvalues are negative definite. Since this choice corresponds to specific values of  $\lambda$  and  $\epsilon$ , depending only on the coefficients of the equation (1.2), we can omit reference to  $\epsilon$  and  $\lambda$  in what follows. Note also that  $J_R$  can be chosen to be independent of  $\iota$ . The reason for this is that there are only two possibilities of what  $R_{\infty,q}(\iota)$  can be, and the difference corresponds to interchanging the blocks in the matrix appearing on the right hand side of the second equality in (21.16).

Estimating the evolution along the time sequence. Fix an integer  $k_0 \ge 0$ . To begin with, we are interested in deriving estimates for  $k \ge k_0$ . Consider (15.63), where  $\psi_k = w_{\text{fin},k}$ ,  $A_k^{\pm} = \Xi_{\text{fin}}^{k,\pm}$ and  $\hat{F}_k$  is replaced by 0. This equality implies that

$$\rho_{k+1} = \bar{A}^+_{k,q,\epsilon} \rho_k, \qquad (21.72)$$

where

$$\rho_k := e^{-\kappa_{q,+}t_k} e^{-J_{\text{diff}}\tilde{t}_k} S_k T_{q,\epsilon} \psi_k, \qquad (21.73)$$
$$\bar{A}^+_{k,q,\epsilon} := e^{-J_{\text{diff}}\tilde{t}_{k+1}} \check{A}^+_{k,q,\epsilon} e^{J_{\text{diff}}\tilde{t}_k},$$

 $\tilde{t}_k := t_k - t_{k_0}$  and  $S_k$  and  $T_{q,\epsilon}$  are defined in the statement of Lemma 21.7. Moreover,  $J_{tr}$  is defined to be the matrix obtained from  $J_R$  by setting the Jordan blocks with zero eigenvalue to zero, and  $J_{\text{diff}} := J_R - J_{\text{tr}}$ . Thus, if  $k \ge k_0$ ,

$$\rho_{k+1} = \bar{A}^+_{k,q,\epsilon} \cdots \bar{A}^+_{k_0,q,\epsilon} \rho_{k_0}.$$
(21.74)

Compute

$$\bar{A}_{k,q,\epsilon}^{+} = e^{-J_{\text{diff}}\tilde{t}_{k+1}} (\check{A}_{k,q,\epsilon}^{+} - \check{A}_{k,q,\epsilon}^{\text{app},+}) e^{J_{\text{diff}}\tilde{t}_{k}} + e^{-J_{\text{diff}}(t_{k+1}-t_{k})} \check{A}_{k,q,\epsilon}^{\text{app},+};$$
(21.75)

note that  $J_{\text{diff}}$  and  $\check{A}_{k,q,\epsilon}^{\text{app},+}$  commute. On the other hand,

$$e^{-J_{\text{diff}}(t_{k+1}-t_k)}\check{A}_{k,q,\epsilon}^{\text{app},+} = \check{A}_{k,q,\epsilon}^{\text{app},+} - J_{\text{diff}}(t_{k+1}-t_k) + \frac{2\pi}{\bar{\nu}_q(\iota)e^{\bar{\beta}_q t_k}}J_R[e^{-J_{\text{diff}}(t_{k+1}-t_k)} - \text{Id}_{2m}] + [e^{-J_{\text{diff}}(t_{k+1}-t_k)} - \text{Id}_{2m} + J_{\text{diff}}(t_{k+1}-t_k)].$$

#### 21.6. OPTIMALITY

Introducing

$$\bar{A}_{k,q}^{\mathrm{app},+} := \mathrm{Id}_{2m} + \frac{2\pi}{\bar{\nu}_{q}(\iota)e^{\bar{\beta}_{q}t_{k}}}J_{\mathrm{tr}}$$
(21.76)

and appealing to (20.51) yields

$$\|e^{-J_{\text{diff}}(t_{k+1}-t_k)}\check{A}_{k,q,\epsilon}^{\text{app},+} - \bar{A}_{k,q}^{\text{app},+}\| \le \frac{C}{\bar{\nu}_q(\iota)e^{\bar{\beta}_q t_k}}e^{-\eta_{\text{dm}}\bar{t}_k}$$

for all  $k \ge k_0$ , where C only depends on the spectrum of the Riemannian manifold corresponding to q and the coefficients of the equation (1.2); note that  $\eta_A \le \eta_{\text{diag}}$ . Combining this estimate with (21.70) and (21.75) yields

$$\|\bar{A}_{k,q,\epsilon}^{+} - \bar{A}_{k,q}^{\mathrm{app},+}\| \le \frac{C}{\bar{\nu}_q(\iota)e^{\bar{\beta}_q t_k}} \langle \tilde{t}_k \rangle^N e^{-\eta_{\mathrm{dm}}\bar{t}_k}$$

for all  $k \ge k_0$ , where *C* only depends on the spectrum of the Riemannian manifold corresponding to *q* and the coefficients of the equation (1.2), and *N* is a non-negative integer depending only on *m*. Note also that  $\epsilon$  has been defined so that  $\|\bar{A}_{k,q}^{\text{app},+}\| \le 1$ . In particular, for  $k \ge k_0$ ,

$$\|\bar{A}_{k,q,\epsilon}^+ \cdots \bar{A}_{k_0,q,\epsilon}^+\| \le C \tag{21.77}$$

for a constant C depending only on the spectrum of the Riemannian manifold corresponding to qand the coefficients of the equation (1.2); the proof of this statement is similar to the argument presented in the proof of Lemma 21.7. Combining this estimate with (21.74) yields

$$|\rho_{k+1}| \le C|\rho_{k_0}| \tag{21.78}$$

for  $k \ge k_0$ , where the constant C has the same dependence as in the case of (21.77). Returning to the definition of  $\rho_k$  in terms of  $\psi_k$ , cf. (21.73), and the definition of  $\psi_k = w_{\text{fin},k}$ , cf. (15.53), this estimate yields

$$|w(\iota, t_{k+1})| \le C e^{\kappa_{q,+} t_{k+1}} \langle \tilde{t}_{k+1} \rangle^{d_{q,+}-1} |w(\iota, t_{k_0})|, \qquad (21.79)$$

where the constant C has the same dependence as in the case of (21.77). Thus (21.68) holds if  $t' = t_{k_0}$  and  $t = t_k$ . In general,  $s \in [t_{k_0-1}, t_{k_0}]$  and  $t \in [t_k, t_{k+1}]$  for some  $0 < k_0 \le k$ . Appealing to Lemma 10.7 in order to estimate the evolution between t' and  $t_{k_0}$  and between  $t_k$  and t yields the conclusion that (21.68) holds for all  $t_0 \le t' \le t$ . Moreover, C has the dependence stated in connection with (21.67). The lemma follows.

#### 21.6 Optimality

It is of interest to determine to what extent the estimates (21.7)-(21.10) are optimal.

**Lemma 21.31.** Consider the equation (1.2). Assume that it is non-degenerate, diagonally dominated, balanced and convergent; cf. Definition 21.1. Fix  $q \in \{1, \ldots, Q\}$  and assume that  $\overline{\beta}_q \neq 0$ . Let  $R_{q,\pm}^+$ ,  $R_{q,\pm}^-$  and  $\kappa_{q,\pm}$  be given by Definition 21.2. For every  $\epsilon > 0$ , there are then constants  $C_{\epsilon} > 0$ ,  $\mathfrak{g}_{\epsilon,\min} > 0$ ,  $t_{\mathrm{as},\epsilon} \geq 0$  and  $t_{\mathrm{mar},\epsilon} \geq 0$ , depending only on the coefficients of the equation (1.2) and  $\epsilon$ , such that the following holds. Fix  $0 \neq \iota \in \mathcal{I}_{\mathrm{B}}$ ; let  $I_q$  be the frequency era corresponding to  $\iota$  and q (cf. Definition 20.10); and define a sequence  $\{t_k\}$  as in Definition 21.4. Then, if  $t_k \in I_q$ and  $\mathfrak{g}(\iota, t_k) \geq \mathfrak{g}_{\epsilon,\min}$  for  $k_- \leq k \leq k_+$ ;  $t_{k_-} \geq t_{\mathrm{as},\epsilon}$ ; and the distance from  $t_{k_{\pm}}$  to the boundary of  $I_q$  is greater than  $t_{\mathrm{mar},\epsilon}$ ,

$$\|A_{k_{\pm}\mp 1}^{\pm}\cdots A_{k_{\mp}}^{\pm}\| \ge C_{\epsilon} \exp\left[\pm(\kappa_{q,\pm}-\epsilon)(t_{k_{\pm}}-t_{k_{\mp}})\right].$$
(21.80)

In the case of the plus sign, the estimate can be improved under the following circumstances. If the Jordan blocks of  $R_{q,+}^+$  and  $R_{q,+}^-$  corresponding to the eigenvalues with real part  $\kappa_{q,+}$  are trivial,

then the constants  $C_{\epsilon} > 0$ ,  $\mathfrak{g}_{\epsilon,\min} > 0$ ,  $t_{as,\epsilon} \ge 0$  and  $t_{mar,\epsilon} \ge 0$  can be chosen to be independent of  $\epsilon$  and the right hand side of (21.80) can be replaced by  $C \exp \left[\kappa_{q,+}(t_{k_+} - t_{k_-})\right]$  for a constant C > 0 depending only on the coefficients of the equation (1.2). The corresponding statement in the case of the minus sign is the following. If the Jordan blocks of  $R_{q,-}^+$  and  $R_{q,-}^-$  corresponding to the eigenvalues with real part  $\kappa_{q,-}$  are trivial, then the constants  $C_{\epsilon} > 0$ ,  $\mathfrak{g}_{\epsilon,\min} > 0$ ,  $t_{as,\epsilon} \ge 0$ and  $t_{mar,\epsilon} \ge 0$  can be chosen to be independent of  $\epsilon$  and the right hand side of (21.80) can be replaced by  $C \exp \left[-\kappa_{q,-}(t_{k_-} - t_{k_+})\right]$  for a constant C > 0 depending only on the coefficients of the equation (1.2).

**Remark 21.32.** Due to (21.80), the estimates (21.7) and (21.8) (with  $C_{\deg,q,\pm} = 0$ ) are optimal in terms of growth/decay in the case of trivial Jordan blocks. In the presence of non-trivial Jordan blocks, there is an  $\epsilon$ -loss; cf. (21.9) and (21.10) (with  $C_{\deg,q,\pm} = 0$ ).

**Remark 21.33.** In case  $\bar{\beta}_q = 0$ , optimality results can be derived using Proposition 12.25; cf. the proof of Theorem 22.13 below.

**Remark 21.34.** The main requirements needed in order to obtain (21.80) have the following intuitive interpretation:

- The inequality  $t_{k_{-}} \geq t_{\mathrm{as},\epsilon}$  corresponds to the requirement that we are in the asymptotic region. Since we do not know much concerning the coefficients before we are in the asymptotic region, such an assumption seems natural.
- That the distance from  $t_{k_{\pm}}$  to the boundary of  $I_q$  is greater than  $t_{\max,\epsilon}$  means that we are sufficiently far away from the boundary of the frequency era  $I_q$ . Since the behaviour close to another frequency era, say  $I_p$ , might as well be approximated by the relevant matrices in  $I_p$ , this is a natural assumption.
- The inequality  $\mathfrak{g}(\iota, t) \geq \mathfrak{g}_{\epsilon,\min}$  corresponds to the requirement that we are well in the oscillatory regime. Since the estimate (21.80) is based on the assumption that there are oscillations and since the corresponding approximations improve with the frequency of the oscillations, this is also a natural assumption.

**Remark 21.35.** The proof also yields a lower bound on the growth of solutions. In fact, define  $w(\iota, t_k)$  by

$$w(\iota, t_k) = T_{\mathrm{pre},k}^{-1} [D_k(\iota)]^{-1} \zeta_k,$$

where  $\zeta_k$  is defined in (21.93) below and  $\xi(k)$  is defined via the iteration (21.82) and a suitable choice of  $\xi(k_a)$  (in the case of the plus sign,  $k_a$  equals  $k_-$  and in the case of the minus sign,  $k_a$ equals  $k_+$ , and the choice of  $\xi(t_{k_a})$  depends on the sign (±)). Then (21.94) yields

$$\frac{|w(\iota, t_{k_{\pm}})|}{|w(\iota, t_{k_{\mp}})|} \ge C_{\epsilon} \exp[\pm(\kappa_{q,\pm} - \epsilon)(t_{k_{\pm}} - t_{k_{\mp}})], \qquad (21.81)$$

where  $C_{\epsilon} > 0$  depends only on  $\epsilon$  and the coefficients of the equation (1.2). Note that there is a solution to the homogeneous version of the equation (24.16) such that  $w(\iota, t_k)$  corresponds to this solution, evaluated at  $t_k$ . Again, in the case of trivial Jordan blocks, the  $\epsilon$  can be removed, so that the growth/decay estimates derived in Corollary 21.17 are optimal in that case (for homogeneous equations and  $\kappa_{q,+} = \kappa_+$  and  $\kappa_{q,-} = \kappa_-$  respectively).

**Remark 21.36.** The optimality statement of Remark 21.35 is limited in the following sense. For every choice of time interval  $[t_{k_-}, t_{k_+}]$  (satisfying the assumptions of the lemma, where q is such that  $\kappa_{q,+} = \kappa_+$ , or such that  $\kappa_{q,-} = \kappa_-$ ), we know that there is a solution such that a lower bound similar to the upper bound in (21.47), or (21.49), holds on that time interval (in the case of homogeneous equations). However, it is not clear that there is a solution such that the lower bound holds on a semi-infinite interval. Nevertheless, if  $\bar{\beta}_q > 0$  and  $\iota$  is such that the only non-zero  $\bar{\nu}_p(\iota)$  is  $\bar{\nu}_q(\iota)$ , then  $I_q$  is semi-infinite and we obtain optimality in the sense that there is a solution to the homogeneous version of the equation such that a lower bound of the desired form holds for all future times. In case  $\bar{\beta}_q = 0$ , optimality results are obtained by appealing to Proposition 12.25. However, in case  $\bar{\beta}_q < 0$  and the spatially homogeneous growth rate is not the largest growth rate, it is not so clear that optimality holds in the sense that we have the desired lower bound for all future times.

*Proof.* Consider the proof of Lemma 21.7. Since the assumptions of the present lemma imply that the assumptions of Lemma 21.7 are satisfied, we know that (21.28) and (21.29) hold. These estimates are the starting point for the analysis. However, in order to be able to demonstrate the desired lower bounds, we need to isolate the component of the solution that corresponds to maximal growth (minimal decay); we need to distinguish between the maximally growing part and the remainder; we need to obtain a lower bound for the growth of the maximally growing component; and, finally, we need to translate the corresponding lower bound to a lower bound on the norm of the matrix products.

Dividing solution vectors into a maximally growing component and a remainder. Assume the top left  $m_+ \times m_+$ -block of  $\operatorname{Re}\{J_{q,\epsilon}\}$  to correspond to the generalised Jordan blocks with eigenvalue  $\kappa_{q,+}$ ; the bottom right  $m_- \times m_-$ -block of  $\operatorname{Re}\{J_{q,\epsilon}\}$  to correspond to the generalised Jordan blocks with eigenvalue  $-\kappa_{q,-}$ ; and collect the remaining blocks in the middle in an  $m_0 \times m_0$ matrix. Note that even if  $J_{q,\epsilon}$  depends on  $\iota$ , the collection of Jordan blocks of  $J_{q,\epsilon}$  does not depend on  $\iota$ . Note, moreover, that the eigenvalues of  $J_{q,\epsilon}$  depend neither on  $\iota$  nor on  $\epsilon$ . Divide vectors  $\xi \in \mathbb{C}^{2m}$  into components  $\xi_{1,\pm}$  and  $\xi_{2,\pm}$ , where  $\xi_{1,+}$  consists of the first  $m_+$  elements of  $\xi$ ;  $\xi_{1,-}$ consists of the last  $m_-$  elements of  $\xi$ ;  $\xi_{2,+}$  consists of the last  $m_0 + m_-$  elements of  $\xi$ ; and  $\xi_{2,-}$ consists of the first  $m_+ + m_0$  elements of  $\xi$ . Let  $\xi(k_a) \in \mathbb{C}^{2m}$  for some  $k_- \leq k_a \leq k_+$ , and define  $\xi(k \pm 1)$  by

$$\xi(k\pm 1) := \check{A}^{\pm}_{k,q,\epsilon}\xi(k), \qquad (21.82)$$

where  $A_{k,q,\epsilon}^{\pm}$  is defined in (21.25). We write the components of  $\xi(k)$  (according to the above division)  $\xi_{1,\pm}(k)$  and  $\xi_{2,\pm}(k)$ . Then appealing to (21.28) and (21.29) with  $\lambda = 0$  yields

$$\begin{aligned} |\xi_{1,\pm}(k\pm1)| &\geq \left(1 + \frac{2\pi}{\bar{\nu}_q(\iota)e^{\bar{\beta}_q t_k}}(\kappa_{q,\pm}-\epsilon)\right) |\xi_{1,\pm}(k)| \\ &- \frac{C_{\epsilon}}{\bar{\nu}_q(\iota)e^{\bar{\beta}_q t_k}} \left[e^{-\eta_A t_k} + \sum_{s=1}^2 \mathfrak{H}_q^{(s)}(\iota,t_k) + \frac{1}{\bar{\nu}_q(\iota)e^{\bar{\beta}_q t_k}}\right] [|\xi_{1,\pm}(k)| + |\xi_{2,\pm}(k)|], \end{aligned}$$
(21.83)

where  $C_{\epsilon}$  depends only on  $\epsilon$  and the coefficients of the equation (1.2). Let us consider the relative sizes of the terms on the right hand side. To begin with, note that, due to (20.33) and the fact that  $t_k \in I_q$ ,

$$\frac{1}{\bar{\nu}_q(\iota)e^{\bar{\beta}_q t_k}} \le \frac{C}{\mathfrak{g}(\iota, t_k)} \le C\mathfrak{g}_{\epsilon, \min}^{-1}, \tag{21.84}$$

where C depends only on the coefficients of the equation (1.2). By assuming  $\mathfrak{g}_{\epsilon,\min}$  to be sufficiently large (depending only on the coefficients of the equation), it is thus clear that we can make the left hand side of (21.84) as small as we wish; from now on we, for the sake of convenience, always assume it to be less than or equal to 1. Thus we can make the first factor in the first term on the right hand side of (21.83) strictly positive by assuming  $\mathfrak{g}_{\epsilon,\min}$  to be sufficiently large, the lower bound depending only on  $\epsilon$  and the coefficients of the equation (1.2) (assuming  $0 < \epsilon < 1$ , which we do below, the lower bound in fact only depends on the coefficients of the equation (1.2)). Consider the second factor in the second term on the right hand side of (21.83). The third term in this factor can be assumed to be as small as we want by demanding  $\mathfrak{g}_{\epsilon,\min}$  to be sufficiently large. Moreover, by assuming  $t_{as,\epsilon}$  to be large enough, we can assume  $e^{-\eta_A t_k}$  to be as small as we wish. Finally, note that

$$\sum_{s=1}^{2} \mathfrak{H}_{q}^{(s)}(\iota, t_{k}) \leq (Q-1)(e^{-\eta_{q} t_{\max,\epsilon}} + e^{-2\eta_{q} t_{\max,\epsilon}})$$

for  $k_{-} \leq k \leq k_{+}$ , where  $\eta_q = |\bar{\beta}_q - \bar{\beta}_p|$ , where  $\bar{\beta}_p$  is the  $\bar{\beta}_q$  which is closest to  $\bar{\beta}_q$ ; cf. (21.69). Again, by assuming  $t_{\max,\epsilon}$  to be large enough, we can assume this expression to be as small as we wish.

**Preliminary conclusions.** Due to the above observations, it is clear that by choosing  $t_{as,\epsilon}$ ,  $t_{mar,\epsilon}$  and  $\mathfrak{g}_{\epsilon,\min}$  to be large enough (the bound depending only on the coefficients of the equation and  $\epsilon$ ), the following holds. If  $|\xi_{1,\pm}(k)| > 0$  and  $|\xi_{1,\pm}(k)| \ge |\xi_{2,\pm}(k)|$ , then the right and side of (21.83) is strictly positive. Note also that we can remove the  $-\epsilon$  appearing in the first row (and replace  $C_{\epsilon}$  by a constant independent of  $\epsilon$  in the second row) in case the relevant generalised Jordan blocks are trivial.

Controlling the relative sizes of the components. Let  $\varkappa_{q,\pm} < \kappa_{q,\pm}$  be the second largest real part of an eigenvalue of  $\pm J_{q,\epsilon}$  (it could be that there is no such  $\varkappa_{q,\pm}$ , but for that to happen the diagonal components of  $\operatorname{Re}\{J_{q,\epsilon}\}$  all have to agree, a situation which is much easier to deal with; in that case, there is no need to control the relative sizes of the components, since there is only one component). Then, assuming  $\mathfrak{g}_{\epsilon,\min}$  to be large enough (the lower bound depending only on the coefficients of the equation),

$$\begin{aligned} |\xi_{2,\pm}(k\pm1)| &\leq \left(1 + \frac{2\pi}{\bar{\nu}_q(\iota)e^{\bar{\beta}_q t_k}}(\varkappa_{q,\pm}+\epsilon)\right) |\xi_{2,\pm}(k)| \\ &+ \frac{C_{\epsilon}}{\bar{\nu}_q(\iota)e^{\bar{\beta}_q t_k}} \left[e^{-\eta_A t_k} + \sum_{s=1}^2 \mathfrak{H}_q^{(s)}(\iota,t_k) + \frac{1}{\bar{\nu}_q(\iota)e^{\bar{\beta}_q t_k}}\right] [|\xi_{1,\pm}(k)| + |\xi_{2,\pm}(k)|], \end{aligned}$$
(21.85)

where  $C_{\epsilon}$  depends only on  $\epsilon$  and the coefficients of the equation (1.2). Let us now assume that  $\xi(k_a) \neq 0$  and that  $|\xi_{2,\pm}(k_a)|/|\xi_{1,\pm}(k_a)| \leq \delta$  for some  $0 < \delta \leq 1$ ; recall that  $k_a$  is introduced in connection with (21.82). Moreover, make the inductive assumption that  $|\xi_{2,\pm}(k)|/|\xi_{1,\pm}(k)| \leq 2\delta$ . Then (21.83) yields

$$\frac{|\xi_{1,\pm}(k\pm1)|}{|\xi_{1,\pm}(k)|} \ge 1 + \frac{2\pi}{\bar{\nu}_q(\iota)e^{\bar{\beta}_q t_k}} (\kappa_{q,\pm} - \epsilon) \\ - \frac{C_\epsilon}{\bar{\nu}_q(\iota)e^{\bar{\beta}_q t_k}} \left[ e^{-\eta_A t_k} + \sum_{s=1}^2 \mathfrak{H}_q^{(s)}(\iota, t_k) + \frac{1}{\bar{\nu}_q(\iota)e^{\bar{\beta}_q t_k}} \right],$$
(21.86)

where  $C_{\epsilon}$  depends only on  $\epsilon$  and the coefficients of the equation (1.2); note that for general  $\delta$ , the constant would also depend on  $\delta$ , but the restriction that  $\delta \leq 1$  allows us to avoid such a dependence. In what follows, we assume  $t_{\mathrm{as},\epsilon}$ ,  $t_{\mathrm{mar},\epsilon}$  and  $\mathfrak{g}_{\epsilon,\min}$  to be large enough that the second and third terms on the right hand side of (21.86) are  $\ll 1$ . For our purposes, it is in fact sufficient to demand that the sum of the second and third terms are, in absolute value, bounded from above by 1/2. The corresponding bounds on  $t_{\mathrm{as},\epsilon}$ ,  $t_{\mathrm{mar},\epsilon}$  and  $\mathfrak{g}_{\epsilon,\min}$  depend only on  $\epsilon$  and the coefficients of the equation (1.2). Due to (21.85), there is an upper bound on  $|\xi_{2,\pm}(k \pm 1)|/|\xi_{1,\pm}(k)|$  which is similar to (21.86). Combining this estimate with (21.86) yields

$$\frac{|\xi_{2,\pm}(k\pm1)|}{|\xi_{1,\pm}(k\pm1)|} \leq \frac{|\xi_{2,\pm}(k)|}{|\xi_{1,\pm}(k)|} + \frac{2\pi}{\bar{\nu}_q(\iota)e^{\bar{\beta}_q t_k}} (\varkappa_{q,\pm} - \kappa_{q,\pm} + 2\epsilon) \frac{|\xi_{2,\pm}(k)|}{|\xi_{1,\pm}(k)|} \\ + \frac{C\epsilon}{\bar{\nu}_q(\iota)e^{\bar{\beta}_q t_k}} \left[ e^{-\eta_A t_k} + \sum_{s=1}^2 \mathfrak{H}_q^{(s)}(\iota, t_k) + \frac{1}{\bar{\nu}_q(\iota)e^{\bar{\beta}_q t_k}} \right],$$
(21.87)

where  $C_{\epsilon}$  has the same dependence as in the case of (21.86). In the end, we would like to argue that if  $|\xi_{2,\pm}(k)|/|\xi_{1,\pm}(k)|$  starts out bounded by  $\delta$ , then it cannot become larger than  $2\delta$ . In order for us to obtain such a conclusion using (21.87), it is necessary for the last factor in the last term on the right hand side of (21.87) to be small. Moreover,  $\epsilon$  has to be small enough that  $\varkappa_{q,\pm} - \kappa_{q,\pm} + 2\epsilon$  is negative. Thus, let us fix

$$0 < \epsilon \leq \frac{1}{6} \min\{\kappa_{q,+} - \varkappa_{q,+}, \kappa_{q,-} - \varkappa_{q,-}\}$$

and  $0 < \delta \leq 1$ . In order for the sum of the last two terms on the right hand side of (21.87) to be strictly negative when  $|\xi_{2,\pm}(k)|/|\xi_{1,\pm}(k)| \geq \delta$ , it is sufficient to assume  $t_{\mathrm{as},\epsilon}$ ,  $t_{\mathrm{mar},\epsilon}$  and  $\mathfrak{g}_{\epsilon,\mathrm{min}}$ to be large enough, where the lower bounds in this case depend only on the coefficients of the equation (1.2),  $\epsilon$  and  $\delta$ . Moreover, again assuming  $t_{\mathrm{as},\epsilon}$ ,  $t_{\mathrm{mar},\epsilon}$  and  $\mathfrak{g}_{\epsilon,\mathrm{min}}$  to be large enough (the constants involved in the restrictions having the same dependence), if  $k_{-} \leq k \leq k_{+}$  and  $|\xi_{2,\pm}(k)| \leq \delta |\xi_{1,\pm}(k)|$ , then

$$\frac{|\xi_{2,\pm}(k\pm1)|}{|\xi_{1,\pm}(k\pm1)|} \le \delta + \frac{2\pi(\kappa_{q,\pm}-\varkappa_{q,\pm})}{\bar{\nu}_q(\iota)e^{\bar{\beta}_q t_k}}\delta$$

Assuming  $\mathfrak{g}_{\epsilon,\min}$  to be large enough, we can assume the second term on the right hand side to be less than  $\delta$ , so that  $|\xi_{2,\pm}(k\pm 1)|/|\xi_{1,\pm}(k\pm 1)| \leq 2\delta$ . Combining the above arguments, it is clear that if  $|\xi_{2,\pm}(k_a)| \leq \delta |\xi_{1,\pm}(k_a)|$ , then the quotient  $|\xi_{2,\pm}(k)|/|\xi_{1,\pm}(k)|$  might reach a value in the interval  $[\delta, 2\delta]$ , but if it does, it will decrease until it is smaller than  $\delta$  etc. In short, an inductive argument demonstrates that  $|\xi_{2,+}(k)|/|\xi_{1,+}(k)| \leq 2\delta$  for all  $k_a \leq k \leq k_+$  and that  $|\xi_{2,-}(k)|/|\xi_{1,-}(k)| \leq 2\delta$ for all  $k_- \leq k \leq k_a$  (though, needless to say, the initial conditions (for  $k = k_a$ ) yielding the first conclusion typically differ from the initial conditions yielding the second conclusion). In the remaining arguments, the particular value of  $\delta$  is not important. We therefore fix  $\delta := 1/4$  from now on.

**Deriving a lower bound on the size of the dominant component.** Returning to (21.83) with the above information concerning the relative sizes of the components in mind, it is clear that

$$\begin{aligned} |\xi_{1,\pm}(k\pm1)| &\geq \left(1 + \frac{2\pi}{\bar{\nu}_q(\iota)e^{\bar{\beta}_q t_k}}(\kappa_{q,\pm} - \epsilon) \\ &- \frac{C_{\epsilon}}{\bar{\nu}_q(\iota)e^{\bar{\beta}_q t_k}} \left[e^{-\eta_A t_k} + \sum_{s=1}^2 \mathfrak{H}_q^{(s)}(\iota, t_k) + \frac{1}{\bar{\nu}_q(\iota)e^{\bar{\beta}_q t_k}}\right] \right) |\xi_{1,\pm}(k)|. \end{aligned}$$
(21.88)

In the case of trivial generalised Jordan blocks, the  $\epsilon$  appearing in the first row can be removed. Moreover, in order to control the relative sizes of the components, it is sufficient to fix an  $\epsilon$  depending only on the coefficients of the equation. For that reason,  $C_{\epsilon}$  appearing on the right hand side can be replaced by a constant C depending only on the coefficients of the equation. In order to estimate  $|\xi_{1,\pm}(k_{\pm})|$  from below in terms of  $|\xi_{1,\pm}(k_a)|$ , it is of interest to estimate the sum of the logarithms of the first factor on the right hand side (for  $k = k_a, \ldots, k_+ - 1$  and  $k = k_- + 1, \ldots, k_a$  respectively). In the case of the plus sign, it can be bounded from below by

$$\begin{split} &\sum_{k=k_a}^{k_+-1} \left( \frac{2\pi}{\bar{\nu}_q(\iota) e^{\bar{\beta}_q t_k}} (\kappa_{q,+} - \epsilon) \right. \\ &\left. - \frac{C_{\epsilon}}{\bar{\nu}_q(\iota) e^{\bar{\beta}_q t_k}} \left[ e^{-\eta_A t_k} + \sum_{s=1}^2 \mathfrak{H}_q^{(s)}(\iota, t_k) + \frac{1}{\bar{\nu}_q(\iota) e^{\bar{\beta}_q t_k}} \right] \right) \\ &\geq & (\kappa_{q,+} - \epsilon)(t_{k_+} - t_{k_a}) - C_{\epsilon}, \end{split}$$

assuming  $\bar{\beta}_q \neq 0$ , where we have appealed to (20.43), (20.44) and (20.45). Moreover,  $C_{\epsilon}$  only depends on  $\epsilon$  and the coefficients of the equation (1.2). Combining this estimate with (21.88) yields

$$|\xi_{1,+}(k_{+})| \ge C_{\epsilon} \exp[(\kappa_{q,+} - \epsilon)(t_{k_{+}} - t_{k_{a}})]|\xi_{1,+}(k_{a})|, \qquad (21.89)$$

where  $C_{\epsilon} > 0$  only depends on  $\epsilon$  and the coefficients of the equation (1.2). A similar argument in the case of the minus sign yields

$$|\xi_{1,-}(k_{-})| \ge C_{\epsilon} \exp[(-\kappa_{q,-} + \epsilon)(t_{k_{-}} - t_{k_{a}})]|\xi_{1,-}(k_{a})|, \qquad (21.90)$$

where the constant has the same dependence as in the case of (21.89). Note, moreover, that the  $\epsilon$  appearing in (21.89) and (21.90) can be removed in case the relevant Jordan blocks are trivial.

One consequence of (21.89) and (21.90) is that

$$|\xi(k_{\pm})| \ge C_{\epsilon} \exp[(\pm \kappa_{q,\pm} \mp \epsilon)(t_{k_{\pm}} - t_{k_a})]|\xi(k_a)|.$$
(21.91)

Here  $C_{\epsilon} > 0$  only depends on  $\epsilon$  and the coefficients of the equation (1.2), and it is important to keep in mind that what  $\xi(k_a)$  is depends on the sign  $(\pm)$ .

**Deriving a lower bound on the norm of the matrix products.** Let us now translate the estimate (21.91) to a lower bound on the norm of the matrix products. By definition,

$$\begin{aligned} \xi(k_{\pm}) &= \check{A}_{k_{\pm}\mp1,q,\epsilon}^{\pm} \cdots \check{A}_{k_{a},q,\epsilon}^{\pm} \xi(k_{a}) = S_{k_{\pm}} \hat{A}_{k_{\pm}\mp1,q,\epsilon}^{\pm} \cdots \hat{A}_{k_{a},q,\epsilon}^{\pm} S_{k_{a}}^{-1} \xi(k_{a}) \\ &= S_{k_{\pm}} A_{k_{\pm}\mp1,q,\epsilon}^{\pm} \cdots A_{k_{a},q,\epsilon}^{\pm} S_{k_{a}}^{-1} \xi(k_{a}) \\ &= S_{k_{\pm}} T_{q,\epsilon} A_{k_{\pm}\mp1}^{\pm} \cdots A_{k_{a}}^{\pm} T_{q,\epsilon}^{-1} S_{k_{a}}^{-1} \xi(k_{a}) \end{aligned}$$
(21.92)

where we have appealed to (21.20), (21.23) and (21.25); recall also that  $\lambda = 0$  in the present setting. In the arguments below, it is, however, important to keep in mind that  $\xi(k_a)$  depends on the sign (±). Letting

$$\zeta_k := T_{q,\epsilon}^{-1} S_k^{-1} \xi(k), \tag{21.93}$$

(21.92) can be written

$$\zeta_{k_{\pm}} = A_{k_{\pm}\mp 1}^{\pm} \cdots A_{k_a}^{\pm} \zeta_{k_a}.$$

In particular, (21.91) yields

$$\|A_{k_{\pm}\mp1}^{\pm}\cdots A_{k_{a}}^{\pm}\| \ge \frac{|\zeta_{k_{\pm}}|}{|\zeta_{k_{a}}|} \ge C_{\epsilon}\frac{|\xi(k_{\pm})|}{|\xi(k_{a})|} \ge C_{\epsilon}\exp[(\pm\kappa_{q,\pm}\mp\epsilon)(t_{k_{\pm}}-t_{k_{a}})],$$
(21.94)

where the  $\epsilon$  can be removed in case the relevant Jordan blocks are trivial. The estimate for the matrix products stated in (21.80) follows and the constants only depend on  $\epsilon$  and the coefficients of the equation; in the case of the plus sign, we can choose  $k_a = k_-$  and in the case of the minus sign, we can choose  $k_a = k_+$ . The lemma follows.

## Chapter 22

# Estimating the Sobolev norms of solutions

#### 22.1 Introduction

Our next goal is to estimate how the Sobolev norms of solutions to (1.2) evolve with time. To be more specific, consider a solution u to (1.2) and define  $\mathfrak{E}_s[u]$  by (24.19), where  $s \in \mathbb{R}$  and  $z(\iota, t)$ is given by (24.14). Then we want to estimate  $\mathfrak{E}_s[u](t), t \geq 0$ , in terms of  $\mathfrak{E}_s[u](0)$  and integrals of appropriate norms of f. However, in the case of homogeneous equations, we are also interested in estimating  $\mathfrak{E}_{s_1}[u](0)$  in terms of  $\mathfrak{E}_{s_2}[u](T)$  for appropriate choices of  $s_1, s_2$  and T > 0. The reason we take an interest in the latter type of estimates is that they are needed when specifying the asymptotics; cf. the proof of Lemma 10.22. In order to obtain estimates, it is natural to consider each mode separately. However, even when considering only one mode, the time intervals under consideration have to be divided into pieces. Due to Corollary 21.17, we can estimate how the energy of one mode evolves in the interval  $[t_{\text{ini},r}, t_{\text{fin},r}]$  (or  $[t_{\text{ini},r}, t_{\text{fin},r})$  in case  $t_{\text{fin},r} = \infty$ ); here  $\{t_k\}$  is a time sequence and  $t_{\text{ini},r}$  and  $t_{\text{fin},r}$  are the corresponding numbers introduced in Definition 21.4. However, we need to proceed differently on  $[t_{\text{fin},r},\infty)$  in case  $t_{\text{fin},r} < \infty$  and on  $[0, t_{\text{ini},r}]$  in case the behaviour is not initially oscillatory. We start by discussing the latter two situations in Section 22.2. We then return to the problem of estimating the energy of the solution in Section 22.3. Finally, in Section 22.4, we verify the optimality of the results.

#### 22.2 Mode estimates for the entire future

When estimating how the energy of a mode develops over time, our main tools are Lemma 9.14, Lemma 10.7 and Corollary 21.17. The subintervals of  $[0,\infty)$  to which we apply these results depend on the  $\iota$  of the mode under consideration. It is therefore convenient to introduce the following classification of the elements of  $\mathcal{I}_{\rm B}$ .

**Definition 22.1.** Consider the equation (1.2). Assume that it is non-degenerate, diagonally dominated, balanced and convergent; cf. Definition 21.1. Then the elements  $\iota$  of  $\mathcal{I}_{\rm B}$  are classified as follows:

- If
- the inequality  $\bar{\beta}_q < 0$  holds for all  $q \in \{1, \ldots, Q\}$  such that  $\bar{\nu}_q(\iota) \neq 0$ , and
- either the inequality (21.6) with  $t_0$  replaced by 0 is violated, or the inequality (14.22) with  $t_a$  replaced by 0 is violated,

then  $\iota$  is said to belong to the low frequency silent sector.

• If

- the inequality  $\bar{\beta}_q < 0$  holds for all  $q \in \{1, \ldots, Q\}$  such that  $\bar{\nu}_q(\iota) \neq 0$ ,
- the inequality (14.22) with  $t_a$  replaced by 0 is satisfied, and
- the inequality (21.6) with  $t_0$  replaced by 0 is satisfied,

then  $\iota$  is said to belong to the high frequency silent sector.

• If

- the inequality  $\bar{\beta}_q \leq 0$  holds for all  $q \in \{1, \ldots, Q\}$  such that  $\bar{\nu}_q(\iota) \neq 0$ ,
- either the inequality (21.6) with  $t_0$  replaced by 0 is violated, or the inequality (14.22) with  $t_a$  replaced by 0 is violated, and
- there is one  $q \in \{1, \ldots, Q\}$  such that  $\bar{\nu}_q(\iota) \neq 0$  and  $\bar{\beta}_q = 0$ ,

then  $\iota$  is said to belong to the *low frequency transparent sector*.

```
• If
```

- the inequality  $\bar{\beta}_q \leq 0$  holds for all  $q \in \{1, \ldots, Q\}$  such that  $\bar{\nu}_q(\iota) \neq 0$ ,
- the inequality (14.22) holds with  $t_a$  replaced by 0,
- the inequality (21.6) with  $t_0$  replaced by 0 is satisfied, and
- there is one  $q \in \{1, \ldots, Q\}$  such that  $\bar{\nu}_q(\iota) \neq 0$  and  $\bar{\beta}_q = 0$ ,

then  $\iota$  is said to belong to the high frequency transparent sector.

• If there is one  $q \in \{1, \ldots, Q\}$  such that  $\bar{\nu}_q(\iota) \neq 0$  and  $\bar{\beta}_q > 0$ , then  $\iota$  is said to belong to the noisy sector.

**Remark 22.2.** Since the equation is non-degenerate, diagonally dominated, balanced and convergent, Lemma 20.3 implies that it is oscillation adapted. Thus (14.1) and (14.6) hold for all  $0 \neq \iota \in \mathcal{I}_{B}$ , and  $\mathfrak{b}$  is the constant appearing in these estimates.

**Remark 22.3.** The element  $\iota = 0$  of  $\mathcal{I}_{B}$  belongs to the low frequency silent sector.

In preparation for the next proposition, it is also convenient to introduce the following terminology.

**Definition 22.4.** Consider the equation (1.2). Assume that it is non-degenerate, diagonally dominated, balanced and convergent; cf. Definition 21.1. If there is a  $q_{ts} \in \{1, \ldots, Q\}$  such that  $\bar{\beta}_{q_{ts}} = 0$ , let  $\kappa_{tot,+} := \max\{\kappa_{ts,+}, \kappa_{sil,+}\}$ , where  $\kappa_{ts,+}$  is defined in (21.51);  $\kappa_{sil,+} := \kappa_{\max}(A_{\infty})$ ; and  $A_{\infty}$  is defined in (10.16). If there is no such  $q_{ts}$ , let  $\kappa_{tot,+} := \max\{\kappa_{+}, \kappa_{sil,+}\}$ , where  $\kappa_{sil,+}$  is defined as before and  $\kappa_{+}$  is defined by (21.5). In addition, (1.2) is said to exhibit subdominant block non-degeneracy if the following holds:

- if there is a  $q_{ts} \in \{1, \ldots, Q\}$  such that  $\bar{\beta}_{q_{ts}} = 0$ , then  $\bar{\kappa}_{q_{ts},+} < \kappa_{tot,+}$ ,
- for  $q \in \{1, \ldots, Q\}$  such that  $\bar{\beta}_q < \max\{0, \bar{\beta}_Q\}$ , all the Jordan blocks of  $R_{q,+}^+$  and  $R_{q,+}^$ corresponding to the eigenvalues with real part  $\kappa_{\text{tot},+}$  are trivial (recall that  $\bar{\beta}_1 < \cdots < \bar{\beta}_Q$ ), and
- if  $\bar{\beta}_Q > 0$  and there is a  $q_{\text{ts}} \in \{1, \dots, Q\}$  such that  $\bar{\beta}_{q_{\text{ts}}} = 0$ , then the Jordan blocks of  $A_{q_{\text{ts}},\infty}(\varsigma)$  corresponding to the eigenvalues with real part  $\kappa_{\text{tot},+}$  are trivial for all  $\varsigma \in \mathcal{I}_{\text{B}}$  such that  $\bar{\nu}_{q_{\text{ts}}}(\varsigma) \neq 0$ .

**Remark 22.5.** The motivation for the introduction of the terminology "subdominant block nondegeneracy" is that, when deriving estimates not involving the loss of an  $\epsilon > 0$ , we wish to exclude the occurrence of frequency eras

- in which the relevant matrices have an eigenvalue with real part  $\kappa_{tot,+}$  and a corresponding non-trivial Jordan block, and
- which are not of the form  $[t_0, \infty)$ .

The reason we wish to exclude such frequency eras is that the polynomial factors in the estimates cause problems at the transition when a new frequency era begins; cf. the estimates appearing in the proof of Lemma 21.7. We also wish to exclude  $\bar{\kappa}_{q_{\text{ts},+}} = \kappa_{\text{tot},+}$ , since the value  $\bar{\kappa}_{q_{\text{ts},+}}$  arises in the limit as  $\bar{\nu}_{q_{\text{ts}}}(\iota) \to \infty$ ; returning to the analysis in Chapter 11, it is clear that there is then a tension between the following two goals: that of decomposing the matrices into Jordan normal form; and that of controlling the norms of the corresponding transformation matrices.

Given the terminology introduced in Definition 22.4, we are in a position to state the main estimates concerning the individual modes of solutions to (1.2).

**Proposition 22.6.** Consider the equation (1.2). Assume that it is non-degenerate, diagonally dominated, balanced and convergent; cf. Definition 21.1. Fix  $\epsilon > 0$ . Then there is a constant  $0 < C_{\epsilon} \in \mathbb{R}$ , depending only on  $\epsilon$ , the spectra of  $(M_r, g_r)$ ,  $r \in \{1, \ldots, R\}$ , and the coefficients of the equation (1.2), such that

$$\mathcal{E}^{1/2}(\iota,t) \le C_{\epsilon} e^{(\kappa_{\text{tot},+}+\epsilon)t} \mathcal{E}^{1/2}(\iota,0) + C_{\epsilon} \int_{0}^{t} e^{(\kappa_{\text{tot},+}+\epsilon)(t-t')} |\hat{f}(\iota,t')| dt'$$
(22.1)

for all  $t \ge 0$ , all solutions z to (24.16) and all  $\iota \in \mathcal{I}_{B}$ , where  $\kappa_{tot,+}$  is given by Definition 22.4. If, in addition to the above, the equation (1.2) exhibits subdominant block non-degeneracy, there are the following improvements of (22.1):

• If  $\bar{\beta}_Q < 0$ , then

$$\mathcal{E}^{1/2}(\iota,t) \le C \langle t \rangle^{d_{\mathrm{s},+}-1} e^{\kappa_{\mathrm{tot},+}t} \mathcal{E}^{1/2}(\iota,0) + C \int_0^t \langle t-t' \rangle^{d_{\mathrm{s},+}-1} e^{\kappa_{\mathrm{tot},+}(t-t')} |\hat{f}(\iota,t')| dt' \quad (22.2)$$

holds for all  $t \ge 0$ , all solutions z to (24.16) and all  $\iota \in \mathcal{I}_B$ , where  $d_{s,+} := d_{\max}(A_{\infty}, \kappa_{tot,+})$ ;  $A_{\infty}$  is defined in (10.16); and the constant C only depends on the spectra of  $(M_r, g_r)$ ,  $r \in \{1, \ldots, R\}$ , and the coefficients of the equation (1.2).

• If  $\bar{\beta}_Q = 0$ , then

$$\mathcal{E}^{1/2}(\iota,t) \le C\langle t \rangle^{d_{\text{sts},+}-1} e^{\kappa_{\text{tot},+}t} \mathcal{E}^{1/2}(\iota,0) + C \int_0^t \langle t-t' \rangle^{d_{\text{sts},+}-1} e^{\kappa_{\text{tot},+}(t-t')} |\hat{f}(\iota,t')| dt' \quad (22.3)$$

holds for all  $t \ge 0$ , all solutions z to (24.16) and all  $\iota \in \mathcal{I}_{\mathrm{B}}$ , where  $d_{\mathrm{sts},+} := \max\{d_{\mathrm{s},+}, d_{\mathrm{t},+}\};$  $d_{\mathrm{s},+}$  is defined as in the case of (22.2);  $d_{\mathrm{t},+}$  is the largest dimension of a non-trivial Jordan block of a matrix of the form  $A_{Q,\infty}(\varsigma)$  (where  $\varsigma \in \mathcal{I}_{\mathrm{B}}$  is such that  $\bar{\nu}_Q(\varsigma) \ne 0$ ) corresponding to an eigenvalue with real part  $\kappa_{\mathrm{tot},+}$ ; and the constant C only depends on the spectra of  $(M_r, g_r), r \in \{1, \ldots, R\}$ , and the coefficients of the equation (1.2).

• If  $\bar{\beta}_Q > 0$ , then

$$\mathcal{E}^{1/2}(\iota,t) \le C\langle t \rangle^{d_{\text{tot},+}-1} e^{\kappa_{\text{tot},+}t} \mathcal{E}^{1/2}(\iota,0) + C \int_0^t \langle t-t' \rangle^{d_{\text{tot},+}-1} e^{\kappa_{\text{tot},+}(t-t')} |\hat{f}(\iota,t')| dt' \quad (22.4)$$

holds for all  $t \ge 0$ , all solutions z to (24.16) and all  $\iota \in \mathcal{I}_B$ , where  $d_{tot,+} := \max\{d_{s,+}, d_{n,+}\}$ ;  $d_{s,+}$  is defined as in the case of (22.2);

$$d_{n,+} := \max\{d_{\max}(R_{O,+}^+, \kappa_{tot,+}), d_{\max}(R_{O,+}^-, \kappa_{tot,+})\};$$

and the constant C only depends on the spectra of  $(M_r, g_r)$ ,  $r \in \{1, \ldots, R\}$ , and the coefficients of the equation (1.2).

**Remark 22.7.** If there is no element of  $\text{Sp}A_{\infty}$  with real part  $\kappa_{\text{tot},+}$ , then the  $d_{\text{s},+}$  appearing in (22.2) equals 1, so that the polynomial expressions appearing in (22.2) can be omitted; cf. Definition 4.3. Similar remarks can be made concerning (22.3) and (22.4).

*Proof.* We divide the proof according to the different cases identified in Definition 22.1.

The low frequency silent sector. Assume that  $\iota$  belongs to the low frequency silent sector. Then either the inequality (21.6) with  $t_0$  replaced by 0 is violated, or the inequality (14.22) with  $t_a$  replaced by 0 is violated. Regardless of which is the case, there is a constant C, depending only on the coefficients of the equation, such that that  $\bar{\nu}_q(\iota) \leq C$  for all q. Combining this observation with (20.21) and (20.22) yields

$$\mathfrak{g}(\iota,t) < C e^{\beta_{\min,-}t}$$

for all  $t \geq 0$ , where *C* only depends on the coefficients of the equation (1.2) and  $\beta_{\min,-}$  is the largest  $\bar{\beta}_q$  which is strictly smaller than zero. In order to obtain this conclusion, we used the fact that  $\bar{\beta}_q < 0$  holds for all  $q \in \{1, \ldots, Q\}$  such that  $\bar{\nu}_q(\iota) \neq 0$ . In analogy with the arguments presented at the beginning of Section 10.4, it is then clear that (24.16) can be written as (10.15), where  $v, A_{\infty}$  and F are as defined in (10.16). Moreover,  $A_{\rm rem}$  is a smooth function taking its values in  $\mathbf{M}_{2m}(\mathbb{C})$  and satisfying (10.17), where, in the present setting,  $C_{\rm rem}$  only depends on the coefficients of the equation (1.2);  $\beta_{\rm rem} := \min\{-\beta_{\min,-}, \eta_{\rm mn}\}$ ; and  $T_{\rm ode} := 0$ . Appealing to Lemma 9.14 yields

$$\mathcal{E}^{1/2}(t) \le C \langle t \rangle^{d_{\mathrm{sil},+}-1} e^{\kappa_{\mathrm{sil},+}t} \mathcal{E}^{1/2}(0) + C \int_0^t \langle t - t' \rangle^{d_{\mathrm{sil},+}-1} e^{\kappa_{\mathrm{sil},+}(t-t')} |\hat{f}(t')| dt'$$
(22.5)

for all  $t \ge 0$ , where C only depends on the coefficients of the equation (1.2);  $\kappa_{\text{sil},+} := \kappa_{\max}(A_{\infty})$ ; and  $d_{\text{sil},+} := d_{\max}(A_{\infty}, \kappa_{\text{sil},+})$ . In order to obtain this conclusion, we used the fact that |v| and  $\mathcal{E}^{1/2}$  are equivalent for  $t \ge 0$ . In particular, it is clear that (22.1) holds.

The high frequency silent sector. Assume that  $\iota$  belongs to the high frequency silent sector. Then we can define a time sequence  $\{t_k\}$  of the type introduced in Definition 21.4, where  $t_0 = 0$ . Since  $\bar{\beta}_q < 0$  for all  $q \in \{1, \ldots, Q\}$  such that  $\bar{\nu}_q(\iota) \neq 0$ , it is clear that  $t_{\text{fin,r}} < \infty$ . Moreover, Corollary 21.17 applies to the interval  $[0, t_{\text{fin,r}}]$ . This yields, for any  $\epsilon > 0$ ,

$$\mathcal{E}^{1/2}(t) \le C_{+,\epsilon} e^{(\kappa_{+}+\epsilon)t} \mathcal{E}^{1/2}(0) + C_{+,\epsilon} \int_{0}^{t} e^{(\kappa_{+}+\epsilon)(t-t')} |\hat{f}(t')| dt'$$
(22.6)

for all  $t \in [0, t_{\text{fin},r}]$ , where  $C_{+,\epsilon}$  only depends on  $\epsilon$  and the coefficients of the equation (1.2) and  $\kappa_+$  is defined by (21.5) (in fact, it is sufficient to compute the supremum over the q's such that  $\bar{\beta}_q < 0$ ). Moreover, in case the Jordan blocks of  $R_{q,+}^+$  and  $R_{q,+}^-$  corresponding to the eigenvalues with real part  $\kappa_+$  are trivial for all  $q \in \{1, \ldots, Q\}$ , the parameter  $\epsilon$  can be replaced by zero, and  $C_{+,\epsilon}$  can be replaced by a constant  $C_+$  independent of  $\epsilon$ . In order to obtain (22.6), we also used the fact that |w| and  $\mathcal{E}^{1/2}$  are equivalent on  $[0, t_{\text{fin},r}]$ . For  $t \geq t_{\text{fin},r}$ , an argument similar to the proof of (22.5) yields

$$\mathcal{E}^{1/2}(t) \leq C \langle \bar{t} \rangle^{d_{\rm sil,+}-1} e^{\kappa_{\rm sil,+}\bar{t}} \mathcal{E}^{1/2}(t_{\rm fin,r}) + C \int_{t_{\rm fin,r}}^{t} \langle t - t' \rangle^{d_{\rm sil,+}-1} e^{\kappa_{\rm sil,+}(t-t')} |\hat{f}(t')| dt'$$
(22.7)

for all  $t \ge t_{\text{fin,r}}$ , where  $\bar{t} := t - t_{\text{fin,r}}$ ; C has the same dependence as in the case of (22.5); and  $\kappa_{\text{sil},+}$  and  $d_{\text{sil},+}$  are defined as in the case of (22.5). When combining (22.6) and (22.7), there are

four different cases to consider. If  $\kappa_{+} < \kappa_{\rm sil,+}$ , then (22.5) holds for all  $t \geq 0$ . If  $\kappa_{+} = \kappa_{\rm sil,+}$ and the  $\epsilon$  appearing in (22.6) can be removed (due to the relevant Jordan blocks being trivial), then (22.5) holds for all  $t \geq 0$ . If  $\kappa_{+} = \kappa_{\rm sil,+}$  and the  $\epsilon$  appearing in (22.6) cannot be removed, then (22.6) holds for all  $t \geq 0$ . If  $\kappa_{+} > \kappa_{\rm sil,+}$ , then (22.6) holds for all  $t \geq 0$ . Moreover, in case the Jordan blocks of  $R_{q,+}^+$  and  $R_{q,+}^-$  corresponding to the eigenvalues with real part  $\kappa_{+}$  are trivial for all  $q \in \{1, \ldots, Q\}$ , the parameter  $\epsilon$  can be replaced by zero, and  $C_{+,\epsilon}$  can be replaced by a constant  $C_{+}$  independent of  $\epsilon$ . Again, it is clear that (22.1) holds.

At this stage, it is convenient to prove (22.2) in case  $\bar{\beta}_Q < 0$  and the equation exhibits subdominant block non-degeneracy. Returning to Definition 22.4, it is clear that the  $\epsilon$  appearing in (22.6) can be removed, if we replace  $\kappa_+$  by  $\kappa_{\text{tot},+}$ . Moreover, if  $\kappa_{\text{sil},+} < \kappa_{\text{tot},+}$ , then  $d_{\text{sil},+}$  can be set equal to 1 in (22.5) and (22.7), if we replace  $\kappa_{\text{sil},+}$  in these estimates with  $\kappa_{\text{tot},+}$ . Considering the above arguments again, it is thus clear that (22.2) holds.

The transparent sectors. Before treating the high/low frequency transparent sectors separately, it is convenient to make some general remarks which apply to both cases. Assume that there are constants  $0 < K_{\rm co} \in \mathbb{R}$  and  $0 \leq t_1 \in \mathbb{R}$  such that  $\mathfrak{g}(\iota, t) \leq K_{\rm co}$ , for  $t \geq t_1$ , where  $\iota$  is in the high or low frequency transparent sector. In this case, there is one element of  $\{1, \ldots, Q\}$ , say  $q_{\rm ts}$ , such that  $\bar{\beta}_{q_{\rm ts}} = 0$  and  $\bar{\nu}_{q_{\rm ts}}(\iota) \neq 0$ . Moreover, there is a constant  $C_{\rm lb} > 0$ , depending only on the coefficients of the equation (1.2) and the spectrum of the Riemannian manifold corresponding to  $q_{\rm ts}$ , such that  $\mathfrak{g}(\iota, t) \geq C_{\rm lb}^{-1}$  for all  $t \geq t_1$ . Keeping this observation in mind, and appealing to (20.21) and (20.23) yields

$$|\mathfrak{g}(\iota,t) - \bar{\nu}_{q_{\mathrm{ts}}}(\iota)| \le C e^{-\beta_{\mathrm{lts},1}t} \tag{22.8}$$

for all  $t \ge t_1$ , where  $\bar{t} := t - t_1$ ,  $\beta_{\text{lts},1} := \min\{-2\beta_{\min,-}, \eta_{\text{diag}}\}$  and C only depends on  $K_{\text{co}}$ , the coefficients of the equation (1.2) and the spectrum of the Riemannian manifold corresponding to  $q_{\text{ts}}$ . Before reformulating the equations as in the beginning of Section 21.3, note that

$$\begin{aligned} \|n_{l}X^{l}(t) - \bar{\nu}_{q_{\rm ts}}(\iota)\bar{X}_{\rm diag}^{q_{\rm ts}}(\iota)\| &= \left\|n_{l}X^{l}(t) - \sum_{\{j:\beta_{j}=0\}}n_{j}X_{\infty}^{j}\right\| \\ &\leq Ce^{\beta_{\min,-}\bar{t}} + \left\|\sum_{\{j:\beta_{j}=0\}}n_{j}[X^{j}(t) - X_{\infty}^{j}]\right\| \end{aligned}$$

for all  $t \ge t_1$ , where the constant only depends on  $K_{\rm co}$  and the coefficients of the equation (1.2). In order to obtain this conclusion, we appealed to (20.8), (20.21), (20.22), (20.37) and the fact that  $\mathfrak{g}(\iota, t) \le K_{\rm co}$  for  $t \ge t_1$ . In particular,

$$\|n_l X^l(t) - \bar{\nu}_{q_{\rm ts}}(\iota) \bar{X}^{q_{\rm ts}}_{\rm diag}(\iota)\| \le C e^{-\beta_{\rm lts,2}\bar{t}}$$

$$\tag{22.9}$$

for all  $t \ge t_1$ , where  $\beta_{lts,2} := \min\{-\beta_{\min,-}, \eta_{mn}\}$  and *C* only depends on  $K_{co}$  and the coefficients of the equation (1.2). Combining (20.9), (22.8) and (22.9) with the fact that the equation is oscillation adapted with  $f_{sh}(t) = Ce^{-\eta_{sh}t}$  yields the conclusion that (24.16) takes the form (21.33) (with  $q = q_{ts}$ ), where  $\varpi_q$  is defined by (21.32);  $A_{q,\infty}(\iota)$  is defined by (21.34);  $F(\iota, t)$  is defined by (21.35); and  $A_{q,rem}(\iota, t)$  satisfies the estimate

$$||A_{q_{\rm ts},\rm rem}(\iota,t)|| \le Ce^{-\beta_{\rm lts}t}$$

for all  $t \geq t_1$ , where

$$eta_{ ext{lts}} := \min\{-eta_{ ext{min},-},\eta_{ ext{mn}},\eta_{ ext{sh}},\eta_{ ext{od}},\eta_{ ext{d}},\eta_{ ext{ad}}\}$$

(recall that  $\eta_{\text{diag}} := \min\{\eta_{\text{od}}, \eta_{\text{d}}, \eta_{\text{ad}}\}$ ) and C only depends on  $K_{\text{co}}$ , the coefficients of the equation (1.2) and the spectrum of the Riemannian manifold corresponding to  $q_{\text{ts}}$ . Introducing

$$\kappa_{\mathrm{ts},\iota}^+ := \kappa_{\max}[A_{q_{\mathrm{ts}},\infty}(\iota)], \quad d_{\mathrm{ts},\iota}^+ := d_{\max}[A_{q_{\mathrm{ts}},\infty}(\iota),\kappa_{\mathrm{ts},\iota}^+],$$

Lemma 9.14 yields the conclusion that

$$|\varpi_{q_{\rm ts}}(\iota,t)| \le C\langle \bar{t} \rangle^{d^+_{{\rm ts},\iota}-1} e^{\kappa^+_{{\rm ts},\iota}\bar{t}} |\varpi_{q_{\rm ts}}(\iota,t_1)| + C \int_{t_1}^t \langle t-t' \rangle^{d^+_{{\rm ts},\iota}-1} e^{\kappa^+_{{\rm ts},\iota}(t-t')} |\hat{f}(\iota,t')| dt'$$
(22.10)

for all  $t \geq t_1$ . However, the constant C in this case depends on  $A_{q_{ts},\infty}(\iota)$  as well as  $K_{co}$ , the coefficients of the equation (1.2) and the spectrum of the Riemannian manifold corresponding to  $q_{ts}$ . On the other hand, since  $\bar{\nu}_{q_{ts}}(\iota) \leq C$  for some C depending only on  $K_{co}$  and the coefficients of the equation (1.2), we need only take a finite number of matrices of the form  $A_{q_{ts},\infty}(\iota)$  into account. For this reason, the constant C appearing in (22.10) can be assumed to only depend on  $K_{co}$ , the coefficients of the equation (1.2) and the spectrum of the Riemannian manifold corresponding to  $q_{ts}$ . Since  $|\varpi_{q_{ts}}(\iota, t)|$  and  $\mathcal{E}^{1/2}(\iota, t)$  are equivalent for  $t \geq t_1$  in the present setting, (22.10) implies that

$$\mathcal{E}^{1/2}(\iota,t) \le C \langle \bar{t} \rangle^{d^+_{\mathrm{ts},\iota} - 1} e^{\kappa^+_{\mathrm{ts},\iota} \bar{t}} \mathcal{E}^{1/2}(\iota,t_1) + C \int_{t_1}^t \langle t - t' \rangle^{d^+_{\mathrm{ts},\iota} - 1} e^{\kappa^+_{\mathrm{ts},\iota}(t-t')} |\hat{f}(\iota,t')| dt'$$
(22.11)

for all  $t \ge t_1$ , where C only depends on  $K_{co}$ , the coefficients of the equation (1.2) and the spectrum of the Riemannian manifold corresponding to  $q_{ts}$ .

The low frequency transparent sector. Assume that  $\iota$  belongs to the low frequency transparent sector. In this case, there is one element of  $\{1, \ldots, Q\}$ , say  $q_{ts}$ , such that  $\bar{\beta}_{q_{ts}} = 0$  and  $\bar{\nu}_{q_{ts}}(\iota) \neq 0$ . Moreover, since either the inequality (21.6) with  $t_0$  replaced by 0 is violated, or the inequality (14.22) with  $t_a$  replaced by 0 is violated, it is clear that there is a constant C, depending only on the coefficients of the equation (1.2) such that  $\mathfrak{g}(\iota, t) \leq C$  for all  $t \geq 0$ . In order to obtain this conclusion, we also used the fact that  $\bar{\beta}_q \leq 0$  for all  $q \in \{1, \ldots, Q\}$  such that  $\bar{\nu}_q(\iota) \neq 0$ . Thus (22.11) holds with  $t_1 = 0$ , and the constant C only depends on the coefficients of the equation (1.2) and the spectrum of the Riemannian manifold corresponding to  $q_{ts}$ . In particular, it is clear that (22.1) holds.

The high frequency transparent sector. Assume that  $\iota$  belongs to the high frequency transparent sector. Then we can construct a time sequence of the type introduced in Definition 21.4, starting at  $t_0 = 0$ . Moreover, there are two possibilities: either  $t_{\text{fin},r} = \infty$  or  $t_{\text{fin},r} < \infty$ . If  $t_{\text{fin},r} = \infty$ , we can appeal to Corollary 21.17 for all  $t \ge 0$ . This yields the conclusion that for every  $\epsilon > 0$ ,

$$\mathcal{E}^{1/2}(\iota,t) \le C_{\epsilon} e^{(\kappa_{\mathrm{ts},+}+\epsilon)t} \mathcal{E}^{1/2}(\iota,0) + C_{\epsilon} \int_{0}^{t} e^{(\kappa_{\mathrm{ts},+}+\epsilon)(t-t')} |\hat{f}(\iota,t')| dt'$$
(22.12)

for all  $t \ge 0$ , where  $C_{\epsilon}$  only depends on  $\epsilon$ , the coefficients of the equation (1.2) and the spectrum of the Riemannian manifold corresponding to  $q_{ts}$  (where  $q_{ts}$  is defined by the condition that  $\bar{\beta}_{q_{ts}} = 0$ ). Moreover,  $\kappa_{ts,+}$  is defined by (21.51), and we used the fact that  $\mathcal{E}^{1/2}$  and |w| are equivalent for  $t \ge 0$ . Note, finally, that the  $\epsilon$  appearing in (22.12) can be removed under certain circumstances. Since these circumstances are described in the statement of Corollary 21.17, we do not repeat them here.

Assume now that  $t_{\text{fin},r} < \infty$ . Then Corollary 21.17 applies in  $[0, t_{\text{fin},r}]$ , so that (22.12) holds for  $t \in [0, t_{\text{fin},r}]$ . As in the low frequency case, since either the inequality (21.6) with  $t_0$  replaced by  $t_{k_{\text{fin},r}+1}$  is violated, or the inequality (14.22) with  $t_a$  replaced by  $t_{k_{\text{fin},r}+1}$  is violated, it is clear that there is a constant C, depending only on the coefficients of the equation (1.2), such that  $\mathfrak{g}(\iota, t) \leq C$  for all  $t \geq t_{\text{fin},r}$ . Moreover, (22.11) yields the conclusion that

$$\mathcal{E}^{1/2}(\iota,t) \leq C \langle \bar{t} \rangle^{d^+_{\mathrm{ts},\iota}-1} e^{\kappa^+_{\mathrm{ts},\iota}\bar{t}} \mathcal{E}^{1/2}(\iota,t_{\mathrm{fin},\mathrm{r}}) + C \int_{t_{\mathrm{fin},\mathrm{r}}}^t \langle t-t' \rangle^{d^+_{\mathrm{ts},\iota}-1} e^{\kappa^+_{\mathrm{ts},\iota}(t-t')} |\hat{f}(\iota,t')| dt'$$

for all  $t \ge t_{\text{fin,r}}$ , where  $\overline{t} := t - t_{\text{fin,r}}$  and C only depends on the coefficients of the equation (1.2) and the spectrum of the Riemannian manifold corresponding to  $q_{\text{ts}}$ . Since  $\kappa_{\text{ts},\iota}^+ \le \kappa_{\text{ts},+}$ , it is thus clear that (22.12) holds for all  $t \ge 0$ , so that (22.1) holds.

Let us now prove that (22.3) holds in case  $\beta_Q = 0$  and the equation exhibits subdominant block non-degeneracy. Due to the analysis in the two silent sectors, we know that the desired conclusion holds if  $\iota$  belongs to either the low or the high frequency silent sector. Due to the analysis presented above for  $\iota$  belonging to the low frequency transparent sector, it also holds for such  $\iota$ . Let us therefore focus on the high frequency transparent sector. By assumption,  $\kappa_{Q,+} < \kappa_{\text{tot},+}$ . Choose a  $K_{\rm co}$ , depending only on the coefficients of the equation (1.2), such that if  $\bar{\nu}_Q(\iota) \geq K_{\rm co}$ , then (14.22) and (21.6) hold for all  $t_a \ge 0$  and  $t_0 \ge 0$  respectively. Defining a time sequence as in Definition 21.4 with  $t_0 = 0$ , it then follows that  $t_{\text{fin,r}} = \infty$ . We are thus allowed to apply conclusion 3 of Corollary 21.17 with  $t_a = 0$  and  $t_b \ge 0$  arbitrary. If  $\kappa_{ts,+} < \kappa_{tot,+}$ , then (21.50) implies (22.3). If  $\kappa_{ts,+} = \kappa_{tot,+}$ , then  $\kappa_{Q,+} = \bar{\kappa}_{Q,+} < \kappa_{tot,+} = \kappa_{ts,+}$ , where the strict inequality is due to the fact that the equation exhibits subdominant block non-degeneracy. Moreover, the Jordan blocks of  $R_{a,+}^+$  and  $R_{a,+}^-$  corresponding to the eigenvalues with real part  $\kappa_{\rm ts,+}$  are trivial for all  $q \in \{1, \ldots, Q\}$ . Considering conclusion 3 of Corollary 21.17, the only obstruction to removing the  $\epsilon$  appearing in (21.50) is that there might be non-trivial Jordan blocks of  $A_{Q,\infty}(\varsigma)$  corresponding to eigenvalues with real part  $\kappa_{ts,+}$  for some  $\varsigma \in \mathcal{I}_B$  such that  $\bar{\nu}_Q(\varsigma) \neq 0$ . Considering the proof of Corollary 21.17, in particular the text adjacent to (21.62), this obstruction can be eliminated if we demand that  $K_{\rm co}$  be large enough, the bound depending only on the coefficients of the equation (1.2) and the spectrum of the Riemannian manifold corresponding to Q. What remains to be considered is thus the case that  $\bar{\nu}_{q_{\rm ts}}(\iota) \leq K_{\rm co}$ . In case  $\iota$  belongs to the low frequency transparent sector, (22.11) with  $t_1$  replaced by 0 yields the desired conclusion. In case  $\iota$  belongs to the high frequency transparent sector, define a time sequence as in Definition 21.4, starting at  $t_0 = 0$ . If  $t_{\rm fin,r} < \infty$ , then a combination of Corollary 21.17 and (22.11) yields the desired conclusion. If  $t_{\text{fin},r} = \infty$ , then there is a  $0 \le t_1 < \infty$  such that  $\mathfrak{g}(\iota, t) \le CK_{\text{co}}$  for all  $t \ge t_1$ , where the constant C only depends on the coefficients of the equation (1.2). Applying Corollary 21.17 in  $[0, t_1]$  and (22.11) in  $[t_1, \infty)$  yields the desired conclusion. To conclude, (22.3) holds.

The noisy sector. Assume that  $\iota$  belongs to the noisy sector. Then there is a  $q_n \in \{1, \ldots, Q\}$  such that  $\bar{\beta}_n := \bar{\beta}_{q_n} > 0$  and  $\bar{\nu}_{q_n}(\iota) \neq 0$ . In particular, there is thus a constant C > 0, depending only on the coefficients of the equation (1.2), such that  $\mathfrak{g}(\iota, t) \geq C\bar{\nu}_{q_n}(\iota)e^{\bar{\beta}_n t}$  for all  $t \geq 0$ . In particular, there is thus a  $t'_0 \geq 0$  such that (14.22) and (21.6) hold for all  $t_a \geq t'_0$  and  $t_0 \geq t'_0$  respectively. Moreover,  $t'_0$  only depends on the coefficients of the equation (1.2) and the spectrum of the Riemannian manifold corresponding to  $q_n$ . Define a time sequence as in Definition 21.4, starting at  $t'_0$ . Then  $t_{\text{fin},\text{r}} = \infty$ . If  $t'_0 = 0$ , we can appeal to Corollary 21.17 with  $t_a = 0$  and  $t_b \geq 0$  arbitrary. If  $t'_0 > 0$ , appealing to Lemma 10.7 yields

$$\mathcal{E}^{1/2}(\iota, t) \le C \mathcal{E}^{1/2}(\iota, 0) + C \int_0^t |\hat{f}(\iota, t')| dt'$$
(22.13)

for all  $t \in [0, t'_0]$ , where C only depends on the coefficients of the equation (1.2) and the spectrum of the Riemannian manifold corresponding to  $q_n$ . Combining this estimate with Corollary 21.17 yields the conclusion that (22.1) holds.

Finally, we wish to prove that (22.4) holds in case  $\beta_Q > 0$  and the equation exhibits subdominant block non-degeneracy. For  $\iota$  in one of the silent sectors, the stated estimate follows from earlier arguments. Assume now that there is a  $q \in \{1, \ldots, Q\}$  such that  $\bar{\beta}_q = 0$  and  $\bar{\nu}_q(\iota) \neq 0$ , say  $q_{\text{ts}}$ . Assume, moreover, that there is no  $q \in \{1, \ldots, Q\}$  such that  $\bar{\beta}_q > 0$  and  $\bar{\nu}_q(\iota) \neq 0$ . As in the transparent setting, there is a  $K_{\text{co}} > 0$ , depending only on the coefficients of the equation (1.2), such that if  $\bar{\nu}_{q_{\text{ts}}}(\iota) \geq K_{\text{co}}$ , then (14.22) and (21.6) hold for all  $t_a \geq 0$  and  $t_0 \geq 0$  respectively. Defining a time sequence as in Definition 21.4 with  $t_0 = 0$ , it then follows that  $t_{\text{fin},\text{r}} = \infty$ . Moreover, there is a  $t_1 \geq 0$  such that  $I_{q_{\text{ts}}} = [t_1, \infty)$ . For any  $\epsilon > 0$ , we can appeal to conclusion 1 of Corollary 21.17 on  $[0, t_1 - \epsilon]$ . Due to the assumption of subdominant block non-degeneracy, this yields an estimate of the form

$$|w(\iota,t)| \le C e^{\kappa_{\text{tot},+}t} |w(\iota,0)| + C \int_0^t e^{\kappa_{\text{tot},+}(t-t')} |\hat{f}(\iota,t')| dt'$$
(22.14)

for all  $t \in [0, t_1 - \epsilon]$ . Moreover, the constant C only depends on the coefficients of the equation (1.2). In particular, it is independent of  $\epsilon$ , so that (22.14) holds on  $[0, t_1]$ . On the interval  $[t_1, \infty)$ , we need to proceed differently. However, assuming  $K_{co}$  to be large enough, the bound depending only on the coefficients of the equation (1.2) and the spectrum of the Riemannian manifold corresponding to  $q_{ts}$ , an argument similar to the one presented in the text adjacent to (21.62) yields the conclusion that

$$|w(\iota,t)| \le C e^{\kappa_{\text{tot},+}(t-t_1)} |w(\iota,t_1)| + C \int_{t_1}^t e^{\kappa_{\text{tot},+}(t-t')} |\hat{f}(\iota,t')| dt'$$

for all  $t \ge t_1$ , where *C* only depends on the coefficients of the equation (1.2) and the spectrum of the Riemannian manifold corresponding to  $q_{\rm ts}$ . Combining this estimate with (22.14) yields (22.4). Next we need to consider the case that  $0 < \bar{\nu}_{q_{\rm ts}}(\iota) < K_{\rm co}$ . If  $\iota$  is in the low frequency transparent sector, (22.11) holds with  $t_1 = 0$ , and the desired conclusion follows. If  $\iota$  is in the high frequency transparent sector, there is a  $t_1$  such that (22.14) holds on  $[0, t_1]$  and such that (22.11) holds for all  $t \ge t_1$ . Keeping in mind that the assumption of subdominant block non-degeneracy is fulfilled, this yields (22.4).

What remains to be considered is the  $\iota \in \mathcal{I}_{B}$  such that  $\bar{\nu}_{q}(\iota) \neq 0$  for some  $q \in \{1, \ldots, Q\}$  with the property that  $\bar{\beta}_{q} > 0$ . Consider such a  $\iota$  and assume that q < Q and that  $\bar{\nu}_{p}(\iota) = 0$  for p > q. Then there is a  $t_{1} \geq 0$  such that  $I_{q} = [t_{1}, \infty)$ . If there is no transparent era, we can appeal to conclusion 1 of Corollary 21.17 combined, possibly, with an estimate of the form (22.13) in order to obtain (22.4). If there is a transparent era, we have to divide the analysis into the two cases  $0 < \bar{\nu}_{q_{\text{ts}}}(\iota) < K_{\text{co}}$  and  $\bar{\nu}_{q_{\text{ts}}}(\iota) \geq K_{\text{co}}$ . If  $K_{\text{co}}$  is large enough, the bound depending only on the coefficients of the equation (1.2) and the spectrum of the Riemannian manifold corresponding to  $q_{\text{ts}}$ , then the arguments presented in the text adjacent to (21.60) yield the conclusion that (22.14) holds for all  $t \geq 0$ . If there is a transparent era with  $0 < \bar{\nu}_{q_{\text{ts}}}(\iota) < K_{\text{co}}$ , then there is a  $t_{\alpha} > 0$ , depending only on the coefficients of the equation (1.2) and the spectra of  $(M_{r}, g_{r}), r \in \{1, \ldots, R\}$ , such that on  $[t'_{0}, \infty)$ , there is a sequence as in Definition 21.4. Moreover, only noisy eras have a non-empty intersection with  $[t'_{0}, \infty)$ . As a consequence, we can appeal to (22.13) on  $[0, t'_{0}]$  and to conclusion 1 of Corollary 21.17 on  $[t'_{0}, \infty)$ . Assume, finally, that  $\bar{\nu}_{Q}(\iota) \neq 0$ . Then there is a  $t_{1} \geq 0$  such that  $I_{Q} = [t_{1}, \infty)$ . In  $[0, t_{1}]$  we can proceed as above in order to obtain the estimate

$$\mathcal{E}^{1/2}(\iota,t) \le C e^{\kappa_{\text{tot},+}t} \mathcal{E}^{1/2}(\iota,0) + C \int_0^t e^{\kappa_{\text{tot},+}(t-t')} |\hat{f}(\iota,t')| dt'$$

for all  $0 \le t \le t_1$ . Finally, on  $[t_1, \infty)$ , we can appeal to Lemma 21.29. Combining these estimates yields the desired conclusion. The proposition follows.

#### 22.3 Energy estimates

Given the estimates derived in Proposition 22.6, it is straightforward to derive Sobolev estimates.

**Theorem 22.8.** Consider the equation (1.2). Assume that it is non-degenerate, diagonally dominated, balanced and convergent; cf. Definition 21.1. Fix  $\epsilon > 0$ . Then there is a constant  $0 < C_{\epsilon} \in \mathbb{R}$ , depending only on  $\epsilon$ , the spectra of  $(M_r, g_r)$ ,  $r \in \{1, \ldots, R\}$ , and the coefficients of the equation (1.2), such that

$$\mathfrak{E}_{s}^{1/2}[u](t) \leq C_{\epsilon} e^{(\kappa_{\text{tot},+}+\epsilon)t} \mathfrak{E}_{s}^{1/2}[u](0) + C_{\epsilon} \int_{0}^{t} e^{(\kappa_{\text{tot},+}+\epsilon)(t-t')} \|f(\cdot,t')\|_{(s)} dt'$$
(22.15)

for all  $t \ge 0$ , all solutions u to (1.2) and all  $s \in \mathbb{R}$ , where  $\kappa_{tot,+}$  is given by Definition 22.4. If, in addition to the above, the equation (1.2) exhibits subdominant block non-degeneracy, there are the following improvements of (22.15):

• If  $\bar{\beta}_Q < 0$ , then

$$\mathfrak{E}_{s}^{1/2}[u](t) \leq C\langle t \rangle^{d_{\mathrm{s},+}-1} e^{\kappa_{\mathrm{tot},+}t} \mathfrak{E}_{s}^{1/2}[u](0) + C \int_{0}^{t} \langle t - t' \rangle^{d_{\mathrm{s},+}-1} e^{\kappa_{\mathrm{tot},+}(t-t')} \|f(\cdot,t')\|_{(s)} dt'$$
(22.16)

holds for all  $t \ge 0$ , all solutions u to (1.2) and all  $s \in \mathbb{R}$ . Here  $d_{s,+} = d_{\max}(A_{\infty}, \kappa_{tot,+})$ ;  $A_{\infty}$  is defined in (10.16); and the constant C only depends on the spectra of  $(M_r, g_r)$ ,  $r \in \{1, \ldots, R\}$ , and the coefficients of the equation (1.2).

350

#### 22.4. OPTIMALITY

• If  $\bar{\beta}_Q = 0$ , then

$$\mathfrak{E}_{s}^{1/2}[u](t) \leq C\langle t \rangle^{d_{\mathrm{sts},+}-1} e^{\kappa_{\mathrm{tot},+}t} \mathfrak{E}_{s}^{1/2}[u](0) + C \int_{0}^{t} \langle t - t' \rangle^{d_{\mathrm{sts},+}-1} e^{\kappa_{\mathrm{tot},+}(t-t')} \|f(\cdot,t')\|_{(s)} dt'$$
(22.17)

holds for all  $t \ge 0$ , all solutions u to (1.2) and all  $s \in \mathbb{R}$ . Here  $d_{\text{sts},+} := \max\{d_{\text{s},+}, d_{\text{t},+}\};$  $d_{\text{s},+}$  is defined as in the case of (22.16);  $d_{\text{t},+}$  is the largest dimension of a non-trivial Jordan block of a matrix of the form  $A_{Q,\infty}(\varsigma)$  (where  $\varsigma \in \mathcal{I}_{\text{B}}$  is such that  $\bar{\nu}_Q(\varsigma) \neq 0$ ) corresponding to an eigenvalue with real part  $\kappa_{\text{tot},+}$ ; and the constant C only depends on the spectra of  $(M_r, g_r), r \in \{1, \ldots, R\}$ , and the coefficients of the equation (1.2).

• If  $\bar{\beta}_Q > 0$ , then

$$\mathfrak{E}_{s}^{1/2}[u](t) \leq C\langle t \rangle^{d_{\text{tot},+}-1} e^{\kappa_{\text{tot},+}t} \mathfrak{E}_{s}^{1/2}[u](0) + C \int_{0}^{t} \langle t - t' \rangle^{d_{\text{tot},+}-1} e^{\kappa_{\text{tot},+}(t-t')} \|f(\cdot,t')\|_{(s)} dt'$$
(22.18)

holds for all  $t \ge 0$ , all solutions u to (1.2) and all  $s \in \mathbb{R}$ . Here  $d_{tot,+} := \max\{d_{s,+}, d_{n,+}\};$  $d_{s,+}$  is defined as in the case of (22.16);

$$d_{n,+} := \max\{d_{\max}(R_{Q,+}^+, \kappa_{\text{tot},+}), d_{\max}(R_{Q,+}^-, \kappa_{\text{tot},+})\};$$

and the constant C only depends on the spectra of  $(M_r, g_r)$ ,  $r \in \{1, \ldots, R\}$ , and the coefficients of the equation (1.2).

**Remark 22.9.** The objects  $\mathfrak{E}_s[u]$  and  $||f(\cdot,t)||_{(s)}$  are defined in (10.18) and (10.19) respectively.

**Remark 22.10.** If (1.2) is geometrically non-degenerate, diagonally dominated, balanced and convergent, then it is non-degenerate, diagonally dominated, balanced and convergent; cf. Remark 26.13.

*Proof.* The statement is an immediate consequence of Proposition 22.6 and Minkowski's inequality.  $\Box$ 

#### 22.4 Optimality

It is of interest to know if the estimates derived in Theorem 22.8 are optimal in the case of homogeneous equations (i.e., when f = 0). There are various ways to define optimality, but here we are mainly interested in comparing  $\kappa_{\text{tot},+}$ , introduced in Definition 22.4, with  $\kappa_{\text{opt},+}$ , defined as follows.

**Definition 22.11.** Consider the equation (1.2) with f = 0. Assume that it is non-degenerate, diagonally dominated, balanced and convergent; cf. Definition 21.1. Let  $\mathcal{A}$  be the set of  $\kappa \in \mathbb{R}$  such that there is a constant  $0 < C \in \mathbb{R}$  with the property that

$$\mathfrak{E}^{1/2}[u](t) \le Ce^{\kappa t} \mathfrak{E}^{1/2}[u](0)$$

for all  $t \ge 0$  and all solutions u to (1.2) with f = 0; here  $\mathfrak{E} = \mathfrak{E}_0$ . Then  $\kappa_{\text{opt},+}$  is defined to be the infimum of  $\mathcal{A}$ .

**Remark 22.12.** Due to Theorem 22.8, we know that  $\mathcal{A}$  is non-empty. Moreover, Lemma 10.15 implies that  $\mathcal{A}$  has a lower bound, so that  $\kappa_{\text{opt},+} \in \mathbb{R}$ .

The main result concerning optimality is the following.

**Theorem 22.13.** Consider the equation (1.2) with f = 0. Assume that it is non-degenerate, diagonally dominated, balanced and convergent; cf. Definition 21.1. Then  $\kappa_{opt,+} = \kappa_{tot,+}$ . Assume, in addition to the above, that the equation exhibits subdominant block non-degeneracy. If  $\bar{\beta}_Q < 0$  and  $\kappa_{sil,+} = \kappa_{tot,+}$ , there is then a constant  $0 < C \in \mathbb{R}$  and a solution  $u \neq 0$  to (1.2) with f = 0 such that

$$\mathfrak{E}^{1/2}[u](t) \ge C\langle t \rangle^{d_{\mathrm{s},+}-1} e^{\kappa_{\mathrm{tot},+}t} \mathfrak{E}^{1/2}[u](0)$$
(22.19)

for all  $t \ge 0$ . In this setting, the estimate (22.16) is thus optimal for f = 0. If  $\bar{\beta}_Q = 0$  and either  $\kappa_{\text{sil},+} = \kappa_{\text{tot},+}$  or  $\kappa_{Q,\text{ts},+} = \kappa_{\text{tot},+}$  holds, then there is a constant  $0 < C \in \mathbb{R}$  and a solution  $u \ne 0$  to (1.2) with f = 0 such that

$$\mathfrak{E}^{1/2}[u](t) \ge C\langle t \rangle^{d_{\mathrm{sts},+}-1} e^{\kappa_{\mathrm{tot},+}t} \mathfrak{E}^{1/2}[u](0)$$
(22.20)

for all  $t \ge 0$ . In this setting, the estimate (22.17) is thus optimal for f = 0. Finally, if  $\overline{\beta}_Q > 0$  and either  $\kappa_{sil,+} = \kappa_{tot,+}$  or  $\kappa_{Q,+} = \kappa_{tot,+}$  holds, then there is a constant  $0 < C \in \mathbb{R}$  and a solution  $u \ne 0$  to (1.2) with f = 0 such that

$$\mathfrak{E}^{1/2}[u](t) \ge C\langle t \rangle^{d_{\text{tot},+}-1} e^{\kappa_{\text{tot},+}t} \mathfrak{E}^{1/2}[u](0)$$
(22.21)

for all  $t \ge 0$ . In this setting, the estimate (22.18) is thus optimal for f = 0.

**Remark 22.14.** The notation  $\kappa_{\text{sil},+}$  and  $\kappa_{\text{tot},+}$  is introduced in Definition 22.4; the notation  $\kappa_{q,\text{ts},+}$  is introduced in Definition 21.14; and the notation  $\kappa_{q,+}$  is introduced in Definition 21.2. The notation  $d_{\text{s},+}$ ,  $d_{\text{sts},+}$  and  $d_{\text{tot},+}$  is introduced in connection with (22.16), (22.17) and (22.18) respectively.

**Remark 22.15.** Note that in the present setting, the  $\kappa_{\text{opt},+}$  introduced in Definition 22.11 equals the  $\eta_{nl}$  introduced in Definition 1.23. In order to justify this statement, note, first of all, that  $\mathfrak{E} = \mathfrak{E}_{\text{bas}}$ ; cf. the comment made in connection with (24.19). Thus  $\mathcal{A}_{nl}$  introduced in Definition 1.23 equals  $\mathcal{A}$  introduced in Definition 22.11. In particular,  $\mathcal{A}_{nl}$  is non-empty and bounded from below, so that  $\eta_{nl}$  is well defined and equals  $\kappa_{\text{opt},+}$ .

*Proof.* Let us begin by proving that  $\kappa_{\text{opt},+} = \kappa_{\text{tot},+}$ . Due to Theorem 22.8 and Definition 22.11, it is clear that  $\kappa_{\text{opt},+} \leq \kappa_{\text{tot},+}$ . What we need to prove is the opposite inequality.

Due to Definition 22.4 and (21.51), we need to consider the following three cases:  $\kappa_{\text{tot},+} = \kappa_{\text{sil},+}$ ;  $\kappa_{\text{tot},+} = \kappa_{q_{\text{ts}},\text{ts},+}$  (assuming there is a  $q \in \{1, \ldots, Q\}$  such that  $\bar{\beta}_q = 0$ , say  $q_{\text{ts}}$ ); and  $\kappa_{\text{tot},+} = \kappa_{q,+}$ for some  $q \in \{1, \ldots, Q\}$  such that  $\bar{\beta}_q \neq 0$ . In the first case, the inequality  $\kappa_{\text{opt},+} \geq \kappa_{\text{tot},+}$  follows from Lemma 9.21; it is sufficient to consider spatially homogeneous solutions to the equations. In the second case, the desired conclusion follows from Proposition 12.25. What remains to be considered is the case that there is a  $q \in \{1, \ldots, Q\}$  such that  $\bar{\beta}_q \neq 0$  and  $\kappa_{q,+} = \kappa_{\text{tot},+}$ . Fix one such q and denote it by  $q_0$ . In this case, we wish to appeal to Lemma 21.31. From now on, we focus on  $\iota \in \mathcal{I}_{\text{B}}$  such that  $\bar{\nu}_q(\iota) = 0$  for all  $q \neq q_0$ . In order to prove the desired lower bound on  $\kappa_{\text{opt},+}$ , it is convenient to assume that  $\kappa_{\text{opt},+} < \kappa_{\text{tot},+} = \kappa_{q_0,+}$ . Let  $\delta := \kappa_{q_0,+} - \kappa_{\text{opt},+} > 0$ . Then there is a constant  $0 < K_{\delta} \in \mathbb{R}$  such that

$$\mathfrak{E}^{1/2}[u](t) \le K_{\delta} e^{(\kappa_{\text{opt},+} + \delta/2)t} \mathfrak{E}^{1/2}[u](0)$$
(22.22)

for all  $t \ge 0$  and all solutions u to (1.2) with f = 0; this is a direct consequence of the definition of  $\kappa_{\text{opt},+}$ . Choose  $\epsilon := \delta/4$  and T > 0. For  $\iota$  such that  $\bar{\nu}_{q_0}(\iota)$  is large enough (the lower bound depending only on the coefficients of the equation (1.2),  $\epsilon$  and T), we can define a sequence as in Definition 21.4, and the sequence satisfies  $t_0 = 0$  and  $t_{\text{fin},r} \ge T$ . Moreover  $[0,T] \subseteq I_{q_0}$ ;  $\mathfrak{g}(\iota,t) \ge$  $\mathfrak{g}_{\epsilon,\min}$  for  $t \in [0,T]$ , where  $\mathfrak{g}_{\epsilon,\min}$  is the constant appearing in the statement of Lemma 21.31; and  $T \ge t_{\mathrm{as},\epsilon} + 2t_{\mathrm{mar},\epsilon}$ , where  $t_{\mathrm{as},\epsilon}$  and  $t_{\mathrm{mar},\epsilon}$  are the constants appearing in the statement of Lemma 21.31. Given  $\iota$  and  $\{t_k\}$  as above, let  $k_-$  be the smallest k such that  $t_k \ge \max\{t_{\mathrm{as},\epsilon}, t_{\mathrm{mar},\epsilon}\}$ and let  $k_+$  be the largest k such that  $t_k \le T - t_{\mathrm{mar},\epsilon}$ . Then there is a solution z to (24.16) such that
where  $t_{\pm} := t_{k_{\pm}}$  and  $C_{\delta} > 0$  only depends on  $\delta$  and the coefficients of the equation (1.2); this is a consequence of (15.63) and Lemma 21.31. Combining this estimate with (10.12), it is clear that

$$|w(\iota, t_{+})| \ge C_{\delta} \exp[(\kappa_{q_{0}, +} - \delta/4)t_{+}]|w(\iota, 0)|, \qquad (22.23)$$

where  $C_{\delta} > 0$  only depends on  $\delta$  and the coefficients of the equation (1.2); note that  $t_{-}$  is bounded by a number depending only on  $\delta$  and the coefficients of the equation (1.2). For T large enough, it is clear that (22.22) and (22.23) are incompatible.

Turning to the estimates (22.19)–(22.21), the first two follow from Lemma 9.21 and Proposition 12.25. The last one is a consequence of the results of Section 19.5. The theorem follows.  $\Box$ 

## Chapter 23

## The silent setting

#### 23.1 Introduction

In Chapter 10, we consider weakly silent, balanced and convergent equations. In particular, we derive detailed asymptotics in Section 10.5, and we demonstrate that it is possible to specify the asymptotics in Section 10.6. Even though the corresponding results are of interest, they suffer from one deficiency: it is not clear how to compute the particular values of the constants  $s_{\text{hom}}$  and  $s_{\text{ih}}$  appearing in the statement of Lemma 10.17; nor is it clear how to compute the value of  $s_{\infty}$  appearing in Lemma 10.22. Our goal in the present chapter is to use the improved estimates derived in Chapter 21 to arrive at specific values for these constants. In order to do so, we here need to make stronger assumptions than in Chapter 10. In fact, in all the results, we consider equations of the form (1.2) which are non-degenerate, diagonally dominated, balanced and convergent; cf. Definition 21.1. Moreover, to ensure that we are in the silent setting, we assume that  $\bar{\beta}_Q < 0$ .

It would be of interest to demonstrate that the numbers  $s_{h,\beta,+}$ ,  $s_{h,\beta}$ ,  $s_{h,\beta}$  and  $s_{h,-}$  appearing in Propositions 23.1 and 23.10 below are optimal. However, we do not attempt to do so here. Nevertheless, in the case of  $s_{h,\beta}$  and  $s_{h,-}$ , we at least conjecture them to be optimal. We also expect methods similar to those used in the proof of Theorem 22.13 to be useful in arriving at such a conclusion. However, there might be a subtlety involved in the transition between the interval in which time sequences of the type introduced in Definition 21.4 are well defined and the remaining interval, say  $[T_{ode}, \infty)$ , in which the ODE behaviour dominates.

**Rough idea of proofs.** The rough structures of the proofs of Propositions 23.1 and 23.10 below are similar to those of Lemmas 10.17 and 10.22. In particular, we consider the different modes  $\iota \in \mathcal{I}_{\rm B}$  one by one, and for each mode, we divide the time interval  $[0, \infty)$  into the subintervals  $[0, T_{\rm ode}]$ and  $[T_{\rm ode}, \infty)$ . However, as opposed to the division made in the proofs of Lemmas 10.17 and 10.22, the time interval  $[0, T_{\rm ode}]$  here equals  $[0, t_{\rm fin,r}]$ , where  $t_{\rm fin,r}$  is introduced in Definition 21.4. In other words, there is a time sequence of the type introduced in Definition 21.4 stretching from t = 0 to  $t = T_{\rm ode}$ . In the interval  $[T_{\rm ode}, \infty)$ , we appeal to Lemmas 9.16 and 9.21, as before. However, in the interval  $[0, T_{\rm ode}]$  we appeal neither to Lemma 10.5, nor to the analogous estimate in the other time direction (cf. the proof of Lemma 10.22). Instead, we make use of the improved knowledge contained in the statement of Corollary 21.25. Considering Definition 21.21 and Corollary 21.25, it is clear that we need to estimate expressions of the form

$$\sum_{j=1}^{k+1} \rho_j |I_j|, \tag{23.1}$$

where  $\rho_j \in \mathbb{R}$ ,  $j = 1, \ldots, k+1$ , are given numbers and  $I_j$  are the intervals into which  $[t', t] \subseteq [0, T_{ode}]$ is divided when appealing to Definition 21.21. In order to obtain useful conclusions, we need to estimate expressions of this type from above. Using the definition of  $T_{ode}$  and the definition of the frequency eras (which enter into the definition of the  $I_j$ ), an upper bound on (23.1) can be derived. It is of the form  $s_{\text{pro}} \ln \langle \nu \rangle + C$ , where  $\nu = \nu(\iota)$ , C is a constant independent of  $\iota$  and  $s_{\text{pro}}$  is a prototype value for the number of derivatives lost in the particular situation under consideration. Moreover,  $s_{\text{pro}}$  can be computed in terms of the  $\rho_j$ 's and the  $\bar{\beta}_q$ 's. In the situations of interest here,  $0 \leq t' \leq t \leq T_{\text{ode}}$ , and we obtain general upper bounds on (23.1) for t', t in this range. Since t' could potentially equal t, it is clear that  $s_{\text{pro}} \geq 0$ . However, in the case that t' = 0 and  $t = T_{\text{ode}}$ , it is sometimes possible to derive bounds of the form  $s_{\text{pro}} \ln \langle \nu \rangle + C$ , where  $s_{\text{pro}} < 0$ . This can happen, in particular, when estimating the asymptotic functions  $V_{\infty}$  appearing in Lemma 10.17 (as well as Proposition 23.1 below), and signals a *gain* in derivatives.

#### 23.2 Future asymptotics, silent setting

As a next step, we wish to strengthen Lemma 10.17 by improving the conclusions concerning the loss of derivatives. In particular, we wish to replace the constants  $s_{\text{hom}}$  and  $s_{\text{ih}}$  appearing in (10.26) and (10.27) by particular values; cf. (23.4), (23.5) and (23.7) below.

**Proposition 23.1.** Consider the equation (1.2). Assume that it is non-degenerate, diagonally dominated, balanced and convergent; cf. Definition 21.1. Assume that  $\bar{\beta}_Q < 0$  and that f is a smooth function such that for every  $s \in \mathbb{R}$ ,

$$||f||_{\mathrm{sil},s} := \int_0^\infty e^{-\kappa_{\mathrm{sil},+}t} ||f(\cdot,t)||_{(s)} dt < \infty,$$
(23.2)

where  $\kappa_{\operatorname{sil},+} := \kappa_{\max}(A_{\infty})$  and  $A_{\infty}$  is defined by (10.16). Let  $\beta_{\operatorname{rem}} := \min\{-\beta_Q, \eta_{\operatorname{mn}}\}$ , where  $\eta_{\operatorname{mn}}$  is the constant appearing in (20.9). Fix  $0 < \beta \leq \beta_{\operatorname{rem}}$  and  $\epsilon > 0$ . Let  $E_a$  be the first generalised eigenspace in the  $\beta, A_{\infty}$ -decomposition of  $\mathbb{C}^{2m}$ ; cf. Definition 4.7. Then there are constants  $C_{\epsilon,\beta}$  and N, where  $C_{\epsilon,\beta}$  only depends on  $\epsilon$ ,  $\beta$ , the coefficients of the equation (1.2) and the spectra of  $(M_r, g_r), r \in \{1, \ldots, R\}$ ; and N only depends on m, such that the following holds. Given a smooth solution u to (1.2), there is a  $V_{\infty} \in C^{\infty}(\overline{M}, E_a)$  such that

$$\left\| \begin{pmatrix} u(\cdot,t) \\ u_t(\cdot,t) \end{pmatrix} - e^{A_{\infty}t}V_{\infty} - \int_0^t e^{A_{\infty}(t-t')} \begin{pmatrix} 0 \\ f(\cdot,t') \end{pmatrix} dt' \right\|_{(s)}$$

$$\leq C_{\epsilon,\beta} \langle t \rangle^N e^{(\kappa_{\mathrm{sil},+}-\beta)t} \left( \|u_t(\cdot,0)\|_{(s+s_{\mathrm{h},\beta,+}+\epsilon)} + \|u(\cdot,0)\|_{(s+s_{\mathrm{h},\beta,+}+\epsilon+1)} + \|f\|_{\mathrm{sil},s+s_{\mathrm{ih},\beta}+\epsilon} \right)$$

$$(23.3)$$

holds for all  $t \ge 0$  and all  $s \in \mathbb{R}$ ; recall that  $\overline{M}$  is given by (1.3). Here

$$s_{\mathrm{h},\beta,+} := \max_{1 \le q \le Q} \max\left\{0, -\frac{\beta + \kappa_{q,+} - \kappa_{\mathrm{sil},+}}{\bar{\beta}_q}\right\},\tag{23.4}$$

$$s_{\mathrm{ih},\beta} := \max\left\{-\frac{\kappa_{\mathrm{ode}}}{\bar{\beta}_Q}, -\frac{\beta}{\bar{\beta}_Q} + \max_{1 \le q \le Q} \max\left\{0, -\frac{\kappa_{q,+} - \kappa_{\mathrm{sil},+}}{\bar{\beta}_q}\right\}\right\},\tag{23.5}$$

where  $\kappa_{ode} := \operatorname{Rsp} A_{\infty}$ , cf. Definition 4.3, and  $\kappa_{q,+}$  is defined by (21.3). Moreover,

$$\|V_{\infty}\|_{(s)} \le C_{\epsilon,\beta} \left( \|u_t(\cdot,0)\|_{(s+s_{h,\beta}+\epsilon)} + \|u(\cdot,0)\|_{(s+s_{h,\beta}+\epsilon+1)} + \|f\|_{\mathrm{sil},s+s_{\mathrm{ih},\beta}+\epsilon} \right),$$
(23.6)

where

$$s_{\mathrm{h},\beta} := \max_{1 \le q \le Q} \left( -\frac{\beta + \kappa_{q,+} - \kappa_{\mathrm{sil},+}}{\bar{\beta}_q} \right).$$
(23.7)

**Remark 23.2.** If the assumptions of Proposition 8.1 are satisfied, then the assumptions of the proposition are satisfied; cf. Remark 22.10.

**Remark 23.3.** The constants  $\beta_q$ ,  $q \in \{1, \ldots, Q\}$ , are introduced at the beginning of Section 20.3.

**Remark 23.4.** The function  $V_{\infty}$  is uniquely determined by the fact that it satisfies the estimate (23.3).

**Remark 23.5.** Both  $s_{h,\beta,+}$  and  $s_{ih,\beta}$  are non-negative. However,  $s_{h,\beta}$  could be strictly negative. In this sense, the function  $V_{\infty}$  could be more regular than the initial data.

**Remark 23.6.** As opposed to (10.26),  $\beta$  need not equal  $\beta_{\text{rem}}$  in (23.3). The reason for allowing  $\beta$  to vary in the range  $(0, \beta_{\text{rem}}]$  is that a higher  $\beta$  yields more detailed asymptotics, but a lower  $\beta$  yields a lower loss of derivatives.

**Remark 23.7.** For particular equations, the value of  $\beta_{\text{rem}}$  can be improved. In fact, if there are constants  $\gamma_i \geq 0$ , i = 1, 2, such that

$$|\sigma(\iota, t)| \le Ce^{-\gamma_1 t}, \quad ||X(\iota, t)|| \le Ce^{-\gamma_2 t}$$
(23.8)

for all  $t \ge 0$  and all  $0 \ne \iota \in \mathcal{I}_{\mathrm{B}}$ , where C only depends on the coefficients of the equation (1.2) and  $\sigma$  and X are defined by (10.1), then  $\beta_{\mathrm{rem}}$  can be replaced by

$$\beta_{\rm rem} := \min\{-2\bar{\beta}_Q, -\bar{\beta}_Q + \gamma_1, -\bar{\beta}_Q + \gamma_2, \eta_{\rm mn}\}.$$
(23.9)

**Remark 23.8.** In the case of homogeneous equations (i.e., when f = 0), the  $\epsilon$  appearing in (23.3) can be removed under the following circumstances. If, in addition to the assumptions of the proposition, all Jordan blocks of  $R_{q,+}^+$  and  $R_{q,+}^-$  corresponding to eigenvalues such that

- the real part of the eigenvalue equals  $\kappa_{q,+}$ , and
- $\kappa_{q,+}$  satisfies

$$-\frac{\beta + \kappa_{q,+} - \kappa_{\mathrm{sil},+}}{\bar{\beta}_q} = s_{\mathrm{h},\beta,+},\tag{23.10}$$

are trivial for  $q = 1, \ldots, Q$ , then

$$\left\| \begin{pmatrix} u(\cdot,t) \\ u_t(\cdot,t) \end{pmatrix} - e^{A_{\infty}t} V_{\infty} \right\|_{(s)}$$

$$\leq C_{\beta} \langle t \rangle^N e^{(\kappa_{\mathrm{sil},+}-\beta)t} \left( \| u_t(\cdot,0) \|_{(s+s_{\mathrm{h},\beta,+})} + \| u(\cdot,0) \|_{(s+s_{\mathrm{h},\beta,+}+1)} \right)$$

$$(23.11)$$

holds for all  $t \ge 0$  and all  $s \in \mathbb{R}$ , where  $C_{\beta}$  only depends on  $\beta$ , the coefficients of the equation (1.2) and the spectra of  $(M_r, g_r), r \in \{1, \ldots, R\}$ ; and N only depends on m. Note also that if there are no eigenvalues of  $R_{q,+}^+$  or  $R_{q,+}^-$  with real part equalling  $\kappa_{q,+}$  and satisfying (23.10),  $q \in \{1, \ldots, Q\}$ , then the conclusion holds.

**Remark 23.9.** Again, in the case of homogeneous equations (i.e., when f = 0), the  $\epsilon$  appearing in (23.6) can be removed under the following circumstances. If, in addition to the assumptions of the proposition, all Jordan blocks of  $R_{q,+}^+$  and  $R_{q,+}^-$  corresponding to eigenvalues such that

- the real part of the eigenvalue equals  $\kappa_{q,+}$ , and
- $\kappa_{q,+}$  satisfies

$$-\frac{\beta + \kappa_{q,+} - \kappa_{\rm sil,+}}{\bar{\beta}_q} = s_{\rm h,\beta},\tag{23.12}$$

are trivial for  $q = 1, \ldots, Q$ , then

$$\|V_{\infty}\|_{(s)} \le C_{\beta} \left( \|u_t(\cdot, 0)\|_{(s+s_{h,\beta})} + \|u(\cdot, 0)\|_{(s+s_{h,\beta}+1)} \right),$$
(23.13)

holds for all  $s \in \mathbb{R}$ , where  $C_{\beta}$  only depends on  $\beta$ , the coefficients of the equation (1.2) and the spectra of  $(M_r, g_r), r \in \{1, \ldots, R\}$ .

*Proof.* Roughly speaking, the proof is quite similar to that of Lemma 10.17. First, for a given  $\iota \in \mathcal{I}_{\mathrm{B}}$ , we divide the time interval  $[0, \infty)$  into  $[0, T_{\mathrm{ode}}]$  and  $[T_{\mathrm{ode}}, \infty)$ . Moreover, we verify that (24.16) can be written as in (10.15), where the constituents are such that the conditions of Subsection 9.1.1 are satisfied on  $[T_{\mathrm{ode}}, \infty)$ . In the interval  $[T_{\mathrm{ode}}, \infty)$  we can appeal to Lemma 9.16. However, we need to reformulate this estimate so that  $T_{\mathrm{ode}}$  does not appear, and we need to derive estimates in the interval  $[0, T_{\mathrm{ode}}]$ . It is at this point that the proof diverges from that of Lemma 10.17; even though the quantities to be estimated are the same as in the proof of Lemma 10.17, we here derive more detailed estimates by appealing to Corollary 21.25 instead of to Lemma 10.5.

**Terminology and preliminary estimates.** Given  $\iota \in \mathcal{I}_{B}$ , it is convenient to divide  $[0, \infty)$  into two subintervals, say  $[0, T_{ode}]$  and  $[T_{ode}, \infty)$ ; cf. the analysis presented in Section 10.5. If  $\iota \in \mathcal{I}_{B}$ is in the low frequency silent sector, cf. Definition 22.1, we let  $T_{ode} := 0$ . If  $\iota \in \mathcal{I}_{B}$  is in the high frequency silent sector, we let  $T_{ode} := t_{fin,r}$ , where  $t_{fin,r}$  results from appealing to Definition 21.4 with  $t_{0} := 0$ . With these definitions,

$$\mathfrak{g}(\iota,t) \le C e^{\beta_Q \bar{t}} \tag{23.14}$$

for all  $t \ge T_{\text{ode}}$ , where  $\bar{t} := t - T_{\text{ode}}$  and C only depends on the coefficients of the equation (1.2). Combining this estimate with (20.10) yields the conclusion that

$$|g^{0l}(t)n_l| \le C e^{(\bar{\beta}_Q - \eta_{\rm sh})\bar{t}}$$

for all  $t \ge T_{\text{ode}}$ , where C only depends on the coefficients of the equation (1.2). Combining (20.8) and (23.14) yields

$$||n_l X^l(t)|| \le C e^{\bar{\beta}_Q \bar{t}}$$

for all  $t \geq T_{\text{ode}}$ , where *C* only depends on the coefficients of the equation (1.2). Finally, recall that (20.9) holds. This yields estimates for  $\alpha - \alpha_{\infty}$  and  $\zeta - \zeta_{\infty}$ . Combining these observations yields the conclusion that the equation (24.16) can be written as in (10.15), where v,  $A_{\infty}$  and F are defined by (10.16). Moreover,  $A_{\text{rem}}$  satisfies the estimate (10.17) for all  $t \geq T_{\text{ode}}$ , where  $\beta_{\text{rem}} := \min\{-\bar{\beta}_Q, \eta_{\text{mn}}\}$  and the constant  $C_{\text{rem}}$  only depends on the coefficients of the equation (1.2). Note also that if (23.8) is satisfied, then (10.17) holds with  $\beta_{\text{rem}}$  given by (23.9).

The expressions to be estimated. Fix  $0 < \beta \leq \beta_{\text{rem}}$ . For every  $\iota \in \mathcal{I}_{\text{B}}$ , we can appeal to Lemma 9.16 for  $t \geq T_{\text{ode}}$ , with this choice of  $\beta$ ; that we are allowed to do so is due to the above observations and the assumption that (23.2) holds. However, just as in the proof of Lemma 10.17, we need to reformulate the estimate so that  $T_{\text{ode}}$  does not appear and to estimate the error in the interval  $[0, T_{\text{ode}}]$ . Introduce, just as in the proof of Lemma 10.17,

$$v_{\infty,\text{mod}} := v_{\infty} - \int_{0}^{T_{\text{ode}}} e^{-A_{\infty}t'} F(t') dt', \qquad (23.15)$$

where we suppress the dependence of the constituents on  $\iota$ . Then (9.22) can be written

$$\left| v(t) - e^{A_{\infty}t} v_{\infty, \text{mod}} - \int_{0}^{t} e^{A_{\infty}(t-t')} F(t') dt' \right|$$

$$\leq C \langle t \rangle^{N} e^{(\kappa_{\text{sil},+}-\beta)t} e^{\beta T_{\text{ode}}} \left[ e^{-\kappa_{\text{sil},+}T_{\text{ode}}} |v(T_{\text{ode}})| + \|F\|_{\text{sil}} \right];$$

$$(23.16)$$

recall that this estimate holds on  $[T_{ode}, \infty)$ . Here we use the notation

$$\|F\|_{\rm sil} := \int_0^\infty e^{-\kappa_{\rm sil,+}t} |F(t)| dt.$$
(23.17)

Note also that the constant C only depends on  $C_{\text{rem}}$ ,  $\beta_{\text{rem}}$  and  $A_{\infty}$ ; i.e., only on the coefficients of the equation (1.2). Similarly to the proof of Lemma 10.17, it is of interest to estimate

$$e^{(\beta-\kappa_{\rm sil,+})t}|v(t)|, \quad e^{(\beta-\kappa_{\rm sil,+})t}|e^{A_{\infty}t}v_{\infty}|, \quad \left|e^{(\beta-\kappa_{\rm sil,+})t}\int_{t}^{T_{\rm ode}}e^{A_{\infty}(t-t')}F(t')dt'\right|$$
(23.18)

for  $t \in [0, T_{ode}]$ ; this yields the desired estimate on the interval  $[0, T_{ode}]$ . In order to obtain a good estimate for  $t \ge T_{ode}$ , we need to estimate

$$e^{(\beta - \kappa_{\rm sil,+})T_{\rm ode}} |v(T_{\rm ode})|, \quad e^{\beta T_{\rm ode}} ||F||_{\rm sil}. \tag{23.19}$$

Finally, we wish to estimate  $v_{\infty,\text{mod}}$  introduced in (23.15).

**Preliminaries, the high frequency case.** Assume that  $T_{ode} > 0$ . Then we can appeal to Corollary 21.25 in  $[0, T_{ode}]$ . Moreover, since there are no transparent eras, we can ignore the parameter K; cf. Remark 21.27. For a given  $\epsilon > 0$ , there is thus a constant  $C_{\epsilon}$ , depending only on  $\epsilon$  and the coefficients of the equation, such that the following holds. If  $0 \le t_a \le t_b$  are such that  $I_a := [t_a, t_b] \subseteq [0, T_{ode}]$ , then

$$|w(t_b)| \le C_{\epsilon} e^{\varrho_{\epsilon,+}(t_b,t_a)} |w(t_a)| + C_{\epsilon} \int_{t_a}^{t_b} e^{\varrho_{\epsilon,+}(t_b,t')} |\hat{f}(t')| dt'$$

for all solutions z to (24.16), where we have omitted reference to  $\iota$  for the sake of brevity. Here  $\rho_{\epsilon,+}$  is introduced in Definition 21.21 (and the omission of the K in the subscript is due to the absence of transparent eras). In particular, since  $|v(t)| \leq C|w(t)|$  for  $t \in [0, T_{\text{ode}}]$  and a constant C only depending on the coefficients of the equation,

$$e^{(\beta-\kappa_{\mathrm{sil},+})t}|v(t)| \leq C_{\epsilon}e^{(\beta-\kappa_{\mathrm{sil},+})t+\varrho_{\epsilon,+}(t,0)}\mathcal{E}^{1/2}(0) + C_{\epsilon}e^{\beta t}\int_{0}^{t}e^{-\kappa_{\mathrm{sil},+}(t-t')+\varrho_{\epsilon,+}(t,t')}e^{-\kappa_{\mathrm{sil},+}t'}|\hat{f}(t')|dt'$$

$$(23.20)$$

for all  $t \in [0, T_{\text{ode}}]$  and all solutions z to (24.16), where we have used the fact that  $|w(t)| \leq \sqrt{2}\mathcal{E}^{1/2}(t)$  and  $C_{\epsilon}$  depends only on  $\epsilon$  and the coefficients of the equation (1.2).

**Deriving estimates for the exponentials.** Due to (23.20), it is clearly of interest to derive upper bounds on

$$(\beta - \kappa_{\rm sil,+})t + \varrho_{\epsilon,+}(t,0), \quad \beta t - \kappa_{\rm sil,+}(t-t') + \varrho_{\epsilon,+}(t,t'), \tag{23.21}$$

where  $0 \leq t' \leq t \leq T_{\text{ode}}$ . Given  $[t',t] \subseteq [0,T_{\text{ode}}]$ , divide it into intervals  $I_1,\ldots,I_{k+1}$  as in Definition 21.21. Then, e.g.,

$$-\kappa_{\rm sil,+}(t-t') + \varrho_{\epsilon,+}(t,t') = \sum_{j=1}^{k+1} (\mu_{j,+} - \kappa_{\rm sil,+}) |I_j|, \qquad (23.22)$$

where the  $\mu_{j,+}$  are introduced in Definition 21.21. Clearly, it is of interest to estimate the right hand side of (23.22). Other expressions appearing in (23.21) can be reformulated similarly. In general, it is of interest to estimate

$$\varsigma(t, t'; \rho) := \sum_{j=1}^{k+1} \rho_j |I_j|,$$

where  $\rho_j \in \mathbb{R}$ ,  $j = 1, \ldots, k+1$ , and  $\rho := (\rho_1, \ldots, \rho_{k+1})$ . Clearly,

$$\varsigma(t, t'; \rho) \le \rho_{\max}(t - t'),$$

where  $\rho_{\max}$  is the largest of the  $\rho_j$ . However, this estimate is too crude for most applications. In order to refine it, note that if  $t' \leq s_1 \leq \cdots \leq s_k \leq t$  corresponds to the division into the intervals  $I_1, \ldots, I_{k+1}$ , then

$$\frac{\bar{\nu}_{q_1}e^{\bar{\beta}_{q_1}t'}}{\bar{\nu}_{q_{k+1}}e^{\bar{\beta}_{q_{k+1}}t}} = \frac{\bar{\nu}_{q_1}e^{\bar{\beta}_{q_1}t'}}{\bar{\nu}_{q_2}e^{\bar{\beta}_{q_2}s_2}} \frac{\bar{\nu}_{q_2}e^{\bar{\beta}_{q_2}s_1}}{\bar{\nu}_{q_{k+1}}e^{\bar{\beta}_{q_{k+1}}s_k}} = \exp\left[-\sum_{j=1}^{k+1}\bar{\beta}_{q_j}|I_j|\right], \quad (23.23)$$

where we suppressed the dependence on  $\iota$  and the numbers  $q_j$  are introduced in Definition 21.21; this equality is a consequence of the definition of the frequency eras. Moreover,

$$\bar{\nu}_{q_{k+1}} e^{\beta_{q_{k+1}}t} \ge 2\pi, \quad \bar{\nu}_{q_1} e^{\beta_{q_1}t'} \le C\langle\nu\rangle,$$
(23.24)

where C only depends on the coefficients of the equation (1.2). Combining (23.23) and (23.24) yields

$$-\sum_{j=1}^{k+1}\bar{\beta}_{q_j}|I_j| \le \ln\langle\nu\rangle + c_0, \qquad (23.25)$$

where  $c_0$  only depends on the coefficients of the equation (1.2). Thus  $\varsigma(t, t'; \rho)$  can be estimated by

$$\varsigma(t,t';\rho) = \sum_{j=1}^{k+1} -\frac{\rho_j}{\bar{\beta}_{q_j}} (-\bar{\beta}_{q_j}|I_j|) \le \max_{1\le j\le k+1} \max\left\{0, -\frac{\rho_j}{\bar{\beta}_{q_j}}\right\} \ln\langle\nu\rangle + C,$$
(23.26)

where C only depends on  $\rho$  and the coefficients of the equation (1.2). In particular, the following estimates hold:

$$(\beta - \kappa_{\operatorname{sil},+})t + \varrho_{\epsilon,+}(t,0) \le \max_{1 \le j \le k+1} \max\left\{0, -\frac{\beta + \mu_{j,+} - \kappa_{\operatorname{sil},+}}{\bar{\beta}_{q_j}}\right\} \ln\langle\nu\rangle + C,$$
(23.27)

$$-\kappa_{\operatorname{sil},+}(t-t') + \varrho_{\epsilon,+}(t,t') \le \max_{1 \le j \le k+1} \max\left\{0, -\frac{\mu_{j,+} - \kappa_{\operatorname{sil},+}}{\bar{\beta}_{q_j}}\right\} \ln\langle\nu\rangle + C,$$
(23.28)

$$\beta t \le -\frac{\beta}{\bar{\beta}_Q} \ln \langle \nu \rangle + C \tag{23.29}$$

for all  $0 \le t' \le t \le T_{\text{ode}}$ , where *C* only depends on the coefficients of the equation (1.2) (that the constant does not depend on  $\epsilon$  is ensured by assuming that  $\epsilon \le 1$  and that it does not depend on  $\beta$  is due to the fact that  $\beta \le \beta_{\text{rem}}$ ). It is of interest to improve the estimate (23.26) when t' = 0 and  $t = T_{\text{ode}}$ . Note, to this end, that when t' = 0, then

$$\bar{\nu}_{q_1} \geq C \langle \nu \rangle$$

where C only depends on the coefficients of the equation (1.2) and the spectrum of the Riemannian manifold corresponding to  $q_1$ . Moreover, when  $t = T_{ode}$ ,

$$\bar{\nu}_{q_{k+1}} e^{\bar{\beta}_{q_{k+1}}t} \le C$$

where  $C \ge 1$  only depends on the coefficients of the equation (1.2). Combining these two estimates with (23.23) yields

$$-\sum_{j=1}^{k+1} \bar{\beta}_{q_j} |I_j| \ge \ln \langle \nu \rangle - c_1,$$
(23.30)

where  $c_1$  only depends on the coefficients of the equation (1.2) and the spectrum of the Riemannian manifold corresponding to  $q_1$ . For this reason, the estimate (23.26) can be improved to

$$\varsigma(T_{\text{ode}}, 0; \rho) \le \max_{1 \le j \le k+1} \left( -\frac{\rho_j}{\bar{\beta}_{q_j}} \right) \ln \langle \nu \rangle + C$$
(23.31)

when t' = 0 and  $t = T_{ode}$ , where C only depends on the coefficients of the equation (1.2),  $\rho$  and the spectrum of the Riemannian manifold corresponding to  $q_1$ . In particular,

$$(\beta - \kappa_{\rm sil,+})T_{\rm ode} + \varrho_{\epsilon,+}(T_{\rm ode},0) \le \max_{1\le j\le k+1} \left(-\frac{\beta + \mu_{j,+} - \kappa_{\rm sil,+}}{\bar{\beta}_{q_j}}\right)\ln\langle\nu\rangle + C,\tag{23.32}$$

where C only depends on the coefficients of the equation (1.2) and the spectrum of the Riemannian manifold corresponding to  $q_1$ .

Estimating v. Combining (23.20) with (23.27)–(23.29) yields

$$e^{(\beta-\kappa_{\rm sil,+})t}|v(t)| \le C_{\epsilon} \langle \nu \rangle^{s_{\rm h,\beta,+}+\epsilon} \mathcal{E}^{1/2}(0) + C_{\epsilon} \langle \nu \rangle^{s_{\rm ih,\beta,1}+\epsilon} \int_0^t e^{-\kappa_{\rm sil,+}t'} |\hat{f}(t')| dt'$$
(23.33)

for all  $t \in [0, T_{ode}]$ , where  $C_{\epsilon}$  only depends on  $\epsilon$  and the coefficients of the equation (1.2). Here  $s_{h,\beta,+}$  is defined by (23.4) and

$$s_{\mathrm{ih},\beta,1} := -\frac{\beta}{\bar{\beta}_Q} + \max_{1 \le q \le Q} \max\left\{0, -\frac{\kappa_{q,+} - \kappa_{\mathrm{sil},+}}{\bar{\beta}_q}\right\};$$

note that  $s_{ih,\beta,1} \leq s_{ih,\beta}$ , where  $s_{ih,\beta}$  is introduced in (23.5). In some situations, the  $\epsilon$  appearing in (23.33) can be removed. Due to complications that arise later in the proof, we here only consider the case of homogeneous equations. If f = 0 and the criteria written down in Remark 23.8 are satisfied, then the  $\epsilon$  appearing in (23.33) can be removed and the constant  $C_{\epsilon}$  can be replaced by one depending only on  $\beta$  and the coefficients of the equation (1.2).

Due to (23.31), the estimate (23.33) can be improved in case  $t = T_{ode}$ . In fact, (23.20) and (23.32) yield

$$e^{(\beta - \kappa_{\rm sil,+})T_{\rm ode}} |v(T_{\rm ode})| \le C_{\epsilon} \langle \nu \rangle^{s_{\rm h,\beta} + \epsilon} \mathcal{E}^{1/2}(0) + C_{\epsilon} \langle \nu \rangle^{s_{\rm ih,\beta,1} + \epsilon} \int_{0}^{T_{\rm ode}} e^{-\kappa_{\rm sil,+}t'} |\hat{f}(t')| dt', \quad (23.34)$$

where  $s_{h,\beta}$  is given by (23.7) and  $C_{\epsilon}$  only depends on  $\epsilon$ , the coefficients of the equation (1.2) and the spectra of  $(M_r, g_r)$ ,  $r \in \{1, \ldots, R\}$ . Moreover, in the case of homogeneous equations, the criterion for removing the  $\epsilon$  appearing in (23.34) is the same as the one stated in Remark 23.9. In that case,  $C_{\epsilon}$  is replaced by a constant  $C_{\beta}$  depending only on  $\beta$ , the coefficients of the equation (1.2) and the spectra of  $(M_r, g_r)$ ,  $r \in \{1, \ldots, R\}$ .

Estimating  $v_{\infty}$ . Let us consider the vector  $v_{\infty}$  appearing in (9.22) in greater detail. To begin with, it is given by (9.23), where  $u_{\infty}$  satisfies the estimate (9.24). Moreover, the constant Cappearing in (9.24) only depends on the coefficients of the equation (1.2). Due to the fact that  $u_{\infty} \in E_a$ ,

$$e^{(\beta-\kappa_{\mathrm{sil},+})t}|e^{A_{\infty}t}v_{\infty}| = e^{(\beta-\kappa_{\mathrm{sil},+})t}|e^{A_{\infty}(t-T_{\mathrm{ode}})}u_{\infty}| \le C_{\beta}e^{(\beta-\kappa_{\mathrm{sil},+})t}e^{-(\kappa_{\mathrm{sil},+}-\beta)(T_{\mathrm{ode}}-t)}|u_{\infty}|$$
$$\le C_{\beta}e^{-(\kappa_{\mathrm{sil},+}-\beta)T_{\mathrm{ode}}}|u_{\infty}| \le C_{\beta}[e^{(\beta-\kappa_{\mathrm{sil},+})T_{\mathrm{ode}}}|v(T_{\mathrm{ode}})| + e^{\beta T_{\mathrm{ode}}}\|\hat{f}\|_{\mathrm{sil}}]$$

for all  $0 \le t \le T_{\text{ode}}$ , where  $C_{\beta}$  only depends on  $\beta$  and the coefficients of the equation (1.2); and we use the notation introduced in (23.17). Combining this estimate with (23.29) and (23.34) yields

$$e^{(\beta - \kappa_{\rm sil,+})t} |e^{A_{\infty}t} v_{\infty}| \le C_{\epsilon,\beta} \langle \nu \rangle^{s_{\rm h,\beta} + \epsilon} \mathcal{E}^{1/2}(0) + C_{\epsilon,\beta} \langle \nu \rangle^{s_{\rm ih,\beta,1} + \epsilon} \|\hat{f}\|_{\rm sil}$$
(23.35)

for all  $0 \le t \le T_{\text{ode}}$ , where  $C_{\epsilon,\beta}$  only depends on  $\epsilon$ ,  $\beta$ , the coefficients of the equation (1.2) and the spectra of  $(M_r, g_r), r \in \{1, \ldots, R\}$ . Moreover, in the case of homogeneous equations, the criterion for removing the  $\epsilon$  appearing in (23.35) is the same as the one stated in Remark 23.9. Note also that in the process of deriving the estimate (23.35), we derived an estimate for the right hand side of (23.16). Note, finally, that setting t to 0 in (23.35) yields an estimate for  $v_{\infty}$ .

#### Estimating the contribution from the inhomogeneity. Let us turn to

$$e^{(\beta - \kappa_{\rm sil,+})t} \int_{t}^{T_{\rm ode}} e^{A_{\infty}(t-t')} F(t') dt'$$

$$= \int_{t}^{T_{\rm ode}} e^{\beta t'} e^{(A_{\infty} - \kappa_{\rm sil,+} \operatorname{Id}_{2m} + \beta \operatorname{Id}_{2m})(t-t')} e^{-\kappa_{\rm sil,+}t'} F(t') dt'.$$
(23.36)

When estimating the right hand side of (23.36), we consider the cases  $\beta - \kappa_{\text{ode}} > 0$  and  $\beta - \kappa_{\text{ode}} \leq 0$ separately; here  $\kappa_{\text{ode}} := \text{Rsp}A_{\infty}$  and the notation RspA is introduced in Definition 4.3. In the second case, there is potentially a polynomial loss, which we estimate crudely by  $e^{-\epsilon \bar{\beta}_Q T_{\text{ode}}}$ . This yields

$$\sup_{t\in[0,T_{\text{ode}}]} \left| e^{(\beta-\kappa_{\text{sil},+})t} \int_{t}^{T_{\text{ode}}} e^{A_{\infty}(t-t')} F(t') dt' \right| \le C_{\epsilon,\beta} \langle \nu \rangle^{s_{\text{ih},\beta,2}+\epsilon} \|\hat{f}\|_{\text{sil}},$$
(23.37)

where

$$s_{\mathrm{ih},\beta,2} := -\max\{\beta,\kappa_{\mathrm{ode}}\}/\bar{\beta}_Q$$

and  $C_{\epsilon,\beta}$  only depends on  $\epsilon$ ,  $\beta$  and the coefficients of the equation (1.2). In order to obtain this conclusion, we appealed to estimates such as (23.29).

Estimating  $v_{\infty,\text{mod}}$ . Combining (23.15), (23.35) (with t set to 0) and (23.37) yields

$$|v_{\infty,\text{mod}}| \le C_{\epsilon,\beta} \langle \nu \rangle^{s_{\text{h},\beta} + \epsilon} \mathcal{E}^{1/2}(0) + C_{\epsilon,\beta} \langle \nu \rangle^{s_{\text{ih},\beta} + \epsilon} \|\hat{f}\|_{\text{sil}}$$
(23.38)

where  $C_{\epsilon,\beta}$  only depends on  $\epsilon$ ,  $\beta$ , the coefficients of the equation (1.2) and the spectra of  $(M_r, g_r)$ ,  $r \in \{1, \ldots, R\}$ . Moreover, in the case of homogeneous equations, the criterion for removing the  $\epsilon$  appearing in (23.38) is the same as the one stated in Remark 23.9.

Analysis for one mode. Consider (23.16). This estimate is valid on  $[T_{ode}, \infty)$ . In order to derive an estimate which is valid for all  $t \ge 0$ , it is of interest to note that

$$e^{-(\kappa_{\mathrm{sil},+}-\beta)t} \left| v(t) - e^{A_{\infty}t} v_{\infty,\mathrm{mod}} - \int_{0}^{t} e^{A_{\infty}(t-t')} F(t') dt' \right|$$

$$\leq C_{\epsilon,\beta} \langle \nu \rangle^{s_{\mathrm{h},\beta,+}+\epsilon} \mathcal{E}^{1/2}(0) + C_{\epsilon,\beta} \langle \nu \rangle^{s_{\mathrm{ih},\beta}+\epsilon} \|\hat{f}\|_{\mathrm{sil}}$$
(23.39)

for all  $t \in [0, T_{\text{ode}}]$ . In order to obtain this estimate, we appealed to (23.33), (23.35) and (23.37). Moreover, the constant  $C_{\epsilon,\beta}$  only depends on  $\epsilon$ ,  $\beta$ , the coefficients of the equation (1.2) and the spectra of  $(M_r, g_r)$ ,  $r \in \{1, \ldots, R\}$ . In the case of homogeneous equations, the criterion for removing the  $\epsilon$  is the same as the one stated in Remark 23.8. Combining (23.16), (23.29) and (23.33) yields

$$\frac{e^{-(\kappa_{\mathrm{sil},+}-\beta)t}\left|v(t)-e^{A_{\infty}t}v_{\infty,\mathrm{mod}}-\int_{0}^{t}e^{A_{\infty}(t-t')}F(t')dt'\right|}{\leq C_{\epsilon}\langle t\rangle^{N}\langle \nu\rangle^{s_{\mathrm{h},\beta,+}+\epsilon}\mathcal{E}^{1/2}(0)+C_{\epsilon}\langle t\rangle^{N}\langle \nu\rangle^{s_{\mathrm{ih},\beta,1}+\epsilon}\|\hat{f}\|_{\mathrm{sil}}}$$
(23.40)

for all  $t \ge T_{\text{ode}}$ , where  $C_{\epsilon}$  only depends on  $\epsilon$  and the coefficients of the equation (1.2). Moreover, N only depends on m. In the case of homogeneous equations, the criterion for removing the  $\epsilon$  is the same as the one stated in Remark 23.8. However,  $C_{\epsilon}$  is then replaced by a constant depending only on  $\beta$  and the coefficients of the equation (1.2). Combining (23.39) and (23.40) yields

$$e^{-(\kappa_{\mathrm{sil},+}-\beta)t} \left| v(t) - e^{A_{\infty}t} v_{\infty,\mathrm{mod}} - \int_{0}^{t} e^{A_{\infty}(t-t')} F(t') dt' \right|$$

$$\leq C_{\epsilon,\beta} \langle t \rangle^{N} \langle \nu \rangle^{s_{\mathrm{h},\beta,+}+\epsilon} \mathcal{E}^{1/2}(0) + C_{\epsilon,\beta} \langle t \rangle^{N} \langle \nu \rangle^{s_{\mathrm{ih},\beta}+\epsilon} \|\hat{f}\|_{\mathrm{sil}}$$
(23.41)

for all  $t \geq 0$ , where  $C_{\epsilon,\beta}$  only depends on  $\epsilon$ ,  $\beta$ , the coefficients of the equation (1.2) and the spectra of  $(M_r, g_r), r \in \{1, \ldots, R\}$ . Again, in the case of homogeneous equations, the criterion for removing the  $\epsilon$  is the same as the one stated in Remark 23.8. In case  $T_{ode} = 0$ , (23.16) immediately yields the conclusion that (23.41) holds (with the factors involving  $\langle \nu \rangle$  removed).

**Projecting the data at infinity.** At this stage we would like to define  $V_{\infty}$  by the requirement that its  $\iota$ 'th Fourier coefficient be given by  $v_{\infty, \text{mod}}(\iota)$ . However, the resulting function need not take its values in  $E_a$ . Just as in the proof of Lemma 10.17, we therefore first need to project  $v_{\infty, \text{mod}}(\iota)$  to the right subspace and to estimate the corresponding error. In particular, we need to estimate

$$e^{-(\kappa_{\mathrm{sil},+}-\beta)t}\left|e^{A_{\infty}t}\Pi_b\left(\int_0^{T_{\mathrm{ode}}}e^{-A_{\infty}t'}F(t')dt'\right)\right|,$$

where  $\Pi_b$  denotes the projection to the second generalised eigenspace in the  $\beta$ ,  $A_{\infty}$ -decomposition of  $\mathbb{C}^{2m}$ ; cf. Lemma 9.6. However, appealing to (23.37) and Lemma 9.6 yields

$$\begin{aligned} & e^{-(\kappa_{\mathrm{sil},+}-\beta)t} \left| e^{A_{\infty}t} \Pi_b \left( \int_0^{T_{\mathrm{ode}}} e^{-A_{\infty}t'} F(t') dt' \right) \right| \\ & \leq C \langle t \rangle^{d_b-1} e^{\kappa_b t} e^{-(\kappa_{\mathrm{sil},+}-\beta)t} \left| \int_0^{T_{\mathrm{ode}}} e^{-A_{\infty}t'} F(t') dt' \right| \\ & \leq C_{\epsilon,\beta} \langle t \rangle^{d_b-1} \langle \nu \rangle^{s_{\mathrm{ih},\beta,2}+\epsilon} \| \widehat{f} \|_{\mathrm{sil}}, \end{aligned}$$

where  $d_b$  and  $\kappa_b$  are given in the statement of Lemma 9.6 and  $C_{\epsilon,\beta}$  only depends on  $\epsilon$ ,  $\beta$  and the coefficients of the equation (1.2). Combining this estimate with (23.41) yields

$$e^{-(\kappa_{\mathrm{sil},+}-\beta)t} \left| v(t) - e^{A_{\infty}t} \Pi_a v_{\infty,\mathrm{mod}} - \int_0^t e^{A_{\infty}(t-t')} F(t') dt' \right|$$

$$\leq C_{\epsilon,\beta} \langle t \rangle^N \langle \nu \rangle^{s_{\mathrm{h},\beta,+}+\epsilon} \mathcal{E}^{1/2}(0) + C_{\epsilon,\beta} \langle t \rangle^N \langle \nu \rangle^{s_{\mathrm{ih},\beta}+\epsilon} \|\hat{f}\|_{\mathrm{sil}}$$
(23.42)

for all  $t \ge 0$ , where  $C_{\epsilon,\beta}$  only depends on  $\epsilon$ ,  $\beta$ , the coefficients of the equation (1.2) and the spectra of  $(M_r, g_r), r \in \{1, \ldots, R\}$ . Here  $\Pi_a$  denotes the projection to the first generalised eigenspace in the  $\beta$ ,  $A_{\infty}$ -decomposition of  $\mathbb{C}^{2m}$ ; cf. Lemma 9.6. Finally, in the case of homogeneous equations, the criterion for removing the  $\epsilon$  in (23.42) is the same as the one stated in Remark 23.8.

**Summing up.** Given  $p = (x, p_1, \ldots, p_R) \in \overline{M}$ , where  $\overline{M}$  is given by (1.3), define  $V_{\infty}(p)$  by

$$V_{\infty}(p) := \sum_{\iota \in \mathcal{I}_{\mathrm{B}}} \prod_{a} [v_{\infty, \mathrm{mod}}(\iota)] \varphi_{\iota}(p),$$

where  $\varphi_{\iota}$  is given by (24.6). In order to justify that the sum on the right hand side makes sense and yields a smooth function, it is sufficient to appeal to the fact that u is smooth; the assumption that (23.2) holds for all s; (23.38); and the Minkowski inequality. Then (23.42) implies that

$$\left\| \begin{pmatrix} u(\cdot,t) \\ u_t(\cdot,t) \end{pmatrix} - e^{A_{\infty}t} V_{\infty} - \int_0^t e^{A_{\infty}(t-t')} \begin{pmatrix} 0 \\ f(\cdot,t') \end{pmatrix} dt' \right\|_{(s)}$$
  
 
$$\leq C_{\epsilon,\beta} \langle t \rangle^N e^{(\kappa_{\mathrm{sil},+}-\beta)t} \left( \mathfrak{E}_{s+s_{\mathrm{h},\beta,+}+\epsilon}^{1/2} [u](0) + \|f\|_{\mathrm{sil},s+s_{\mathrm{ih},\beta}+\epsilon} \right)$$

for all  $t \ge 0$ , where we appealed to the Minkowski inequality and  $C_{\epsilon,\beta}$  and N have the same dependence as in the case of (23.42). Thus (23.3) holds. Moreover, Remark 23.8 holds. Note also that, due to (23.38),

$$\|V_{\infty}\|_{(s)} \le C_{\epsilon,\beta} \left(\mathfrak{e}_{s+s_{\mathrm{h},\beta}+\epsilon}^{1/2} [u](0) + \|f\|_{\mathrm{sil},s+s_{\mathrm{ih},\beta}+\epsilon}\right),$$
(23.43)

where  $C_{\epsilon,\beta}$  has the same dependence as in the case of (23.41). Thus (23.6) holds. Moreover, Remark 23.9 holds. The proposition follows.

#### 23.3 Specifying the asymptotics

As a next step, it is of interest to improve Lemma 10.22 by demonstrating that the conclusions hold for a specific value of  $s_{\infty}$  that can be calculated in terms of the coefficients of the equation.

**Proposition 23.10.** Consider the equation (1.2) with f = 0. Assume that it is non-degenerate, diagonally dominated, balanced and convergent; cf. Definition 21.1. Assume that  $\bar{\beta}_Q < 0$  and let  $\beta_{\text{rem}} := \min\{-\bar{\beta}_Q, \eta_{\text{mn}}\}$ , where  $\eta_{\text{mn}}$  is the constant appearing in (20.9). Fix  $0 < \beta \leq \beta_{\text{rem}}$  and  $\epsilon > 0$ . Let  $E_a$  be the first generalised eigenspace in the  $\beta, A_{\infty}$ -decomposition of  $\mathbb{C}^{2m}$ , where  $A_{\infty}$  is defined by (10.16); cf. Definition 4.7. Then there is a linear injective map

$$\Phi_{\beta,\infty}: C^{\infty}(\bar{M}, E_a) \to C^{\infty}(\bar{M}, \mathbb{C}^{2m})$$

such that the following holds. First, if  $\Phi_{\beta,\infty}^j : C^{\infty}(\bar{M}, E_a) \to C^{\infty}(\bar{M}, \mathbb{C}^m)$ , j = 1, 2, are defined by the condition that

$$\Phi_{\beta,\infty}(\chi) = \begin{pmatrix} \Phi^1_{\beta,\infty}(\chi) \\ \Phi^2_{\beta,\infty}(\chi) \end{pmatrix}$$

for all  $\chi \in C^{\infty}(\overline{M}, E_a)$ , then

$$\|\Phi_{\beta,\infty}^{1}(\chi)\|_{(s+1)} + \|\Phi_{\beta,\infty}^{2}(\chi)\|_{(s)} \le C_{\epsilon}\|\chi\|_{(s+s_{h,-}+\epsilon)}$$
(23.44)

for all  $s \in \mathbb{R}$  and all  $\chi \in C^{\infty}(\overline{M}, E_a)$ , where  $C_{\epsilon}$  only depends on  $\epsilon$ , the coefficients of the equation (1.2) and the spectra of  $(M_r, g_r)$ ,  $r \in \{1, \ldots, R\}$ , and

$$s_{\mathrm{h},-} := \max_{1 \le q \le Q} \left( -\frac{\kappa_{q,-} + \kappa_{\mathrm{sil},+}}{\bar{\beta}_q} \right). \tag{23.45}$$

Secondly, if  $\chi \in C^{\infty}(\overline{M}, E_a)$  and u is the solution to (1.2) (with f = 0) such that

$$\begin{pmatrix} u(\cdot,0)\\ u_t(\cdot,0) \end{pmatrix} = \Phi_{\beta,\infty}(\chi), \tag{23.46}$$

then

$$\left\| \begin{pmatrix} u(\cdot,t) \\ u_t(\cdot,t) \end{pmatrix} - e^{A_{\infty}t} \chi \right\|_{(s)}$$

$$\leq C_{\epsilon,\beta} \langle t \rangle^N e^{(\kappa_{\mathrm{sil},+}-\beta)t} \left( \| u_t(\cdot,0) \|_{(s+s_{\mathrm{h},\beta,+}+\epsilon)} + \| u(\cdot,0) \|_{(s+s_{\mathrm{h},\beta,+}+\epsilon+1)} \right)$$

$$(23.47)$$

for all  $t \ge 0$  and all  $s \in \mathbb{R}$ , where  $C_{\epsilon,\beta}$  only depends on  $\epsilon$ ,  $\beta$ , the coefficients of the equation (1.2) and the spectra of  $(M_r, g_r)$ ,  $r \in \{1, \ldots, R\}$ , and N only depends on m. Finally, if  $E_a = \mathbb{C}^{2m}$  (i.e., if  $\beta > \operatorname{Rsp} A_{\infty}$ ; cf. Definition 4.3), then  $\Phi_{\beta,\infty}$  is surjective.

**Remark 23.11.** If the assumptions of Proposition 8.9 are satisfied, then the assumptions of the proposition are satisfied; cf. Remark 22.10.

**Remark 23.12.** If the Jordan blocks of  $A_{\infty}$  corresponding to eigenvalues with real part  $\kappa_{\rm sil,+}$  are trivial, and if all Jordan blocks of  $R_{q,-}^+$  and  $R_{q,-}^-$  corresponding to eigenvalues such that

- the real part of the eigenvalue equals  $\kappa_{q,-}$ , and
- $\kappa_{q,-}$  satisfies

$$-\frac{\kappa_{q,-}+\kappa_{\rm sil,+}}{\bar{\beta}_q} = s_{\rm h,-},\qquad(23.48)$$

are trivial for  $q = 1, \ldots, Q$ , then the  $\epsilon$  appearing in (23.44) can be removed, so that

$$\|\Phi_{\beta,\infty}^{1}(\chi)\|_{(s+1)} + \|\Phi_{\beta,\infty}^{2}(\chi)\|_{(s)} \le C \|\chi\|_{(s+s_{h,-})},$$

where C only depends on the coefficients of the equation (1.2) and the spectra of  $(M_r, g_r), r \in \{1, \ldots, R\}$ .

**Remark 23.13.** By combining (23.44), (23.46) and (23.47), the norms of  $u(\cdot, 0)$  and  $u_t(\cdot, 0)$  appearing on the right hand side of (23.47) can be replaced by a suitable Sobolev norm of  $\chi$ .

**Remark 23.14.** In order to obtain a similar result in the case of inhomogeneous equations, it is sufficient to combine Propositions 23.1 and 23.10; cf. Remark 9.23.

**Remark 23.15.** The same criteria and conclusions concerning improvements of  $\beta_{\text{rem}}$  that are stated in Remark 23.7 hold in the present proposition.

*Proof.* Let  $\chi \in C^{\infty}(\overline{M}, E_a)$ ,  $\iota \in \mathcal{I}_{\mathrm{B}}$  and  $\hat{\chi}(\iota) := \langle \chi, \varphi_{\iota} \rangle_{\mathrm{B}} \in E_a$ ; cf. the notation introduced in Subsection 24.1.2. Let u be a solution to (1.2) and let  $z(\iota, t)$  be given by (24.14). Note that the  $\iota$ 'th Fourier coefficient of the expression inside the norm on the left hand side of (23.47) can be written

$$v - e^{A_{\infty}t}\hat{\chi},\tag{23.49}$$

where we use the notation (10.16). On the other hand, v satisfies (10.15) with F = 0, and  $A_{\rm rem}$  satisfies (10.17) for  $t \geq T_{\rm ode}$  (where  $T_{\rm ode}$  is defined as in the proof of Proposition 23.1). Moreover,  $\beta_{\rm rem}$  has the value given in the statement of Proposition 23.1 (or in Remark 23.7, if

the conditions of Remark 23.7 apply) and  $C_{\rm rem}$  only depends on the coefficients of the equation (1.2). In particular, the conditions needed to apply Lemma 9.21 are satisfied. In fact, specifying v by imposing the initial condition  $v(T_{\rm ode}) := \Psi_{\infty}[\hat{\chi}(\iota)]$  yields the conclusion that (23.49) is small asymptotically; here  $\Psi_{\infty}$  is the map constructed in Lemma 9.21. Note that  $\Psi_{\infty}$  depends on  $\iota$ . However, the estimates involving  $\Psi_{\infty}$  that we need do not. In order to estimate  $|v(T_{\rm ode})|$  in terms of  $|\hat{\chi}(\iota)|$ , define  $u_{\infty} \in E_a$  by

$$e^{-A_{\infty}T_{\text{ode}}}u_{\infty} = \hat{\chi}(\iota).$$

Thus  $|u_{\infty}| = |e^{A_{\infty}T_{\text{ode}}}\hat{\chi}(\iota)|$ , so that if  $\epsilon > 0$  is fixed, then (9.45) yields the estimate

$$|v(T_{\text{ode}})| = |\Psi_{\infty}(e^{-A_{\infty}T_{\text{ode}}}u_{\infty})| \le C|u_{\infty}| = C|e^{A_{\infty}T_{\text{ode}}}\hat{\chi}(\iota)|$$
  
$$\le C_{\epsilon}e^{(\kappa_{\text{sil},+}+\epsilon)T_{\text{ode}}}|\hat{\chi}(\iota)|$$
(23.50)

where  $C_{\epsilon}$  only depends on  $\epsilon$  and the coefficients of the equation (1.2). Moreover, the  $\epsilon$  can be removed if the Jordan blocks of  $A_{\infty}$  corresponding to eigenvalues with real part  $\kappa_{\rm sil,+}$  are trivial. In order to estimate  $\mathcal{E}_{\rm s}(0)$ , it is natural to divide the analysis into two cases. To begin with, assume that  $T_{\rm ode} = 0$ . Then

$$\mathcal{E}^{1/2}(0) \le C|v(0)| = C|v(T_{\text{ode}})| \le C|\hat{\chi}(\iota)|, \tag{23.51}$$

where C only depends on the coefficients of the equation (1.2) and we have appealed to (23.50) and the fact that  $T_{ode} = 0$ .

The high frequency case. In case  $T_{ode} > 0$ , we appeal to Corollary 21.25 with f = 0. For  $\epsilon_1 > 0$ , this yields

$$\mathcal{E}^{1/2}(0) \le C|w(0)| \le C_{\epsilon_1} e^{\varrho_{\epsilon_1,-}(0,T_{\text{ode}})} |w(T_{\text{ode}})|, \qquad (23.52)$$

where  $C_{\epsilon_1}$  only depends on  $\epsilon_1$  and the coefficients of the equation (1.2), and we have omitted the subscript K due to the absence of transparent eras. Due to the definition of  $T_{\text{ode}}$ , the expression  $|w(T_{\text{ode}})|$  is bounded by  $C|v(T_{\text{ode}})|$ , where C only depends on the coefficients of the equation (1.2). Combining this observation with (23.50) and (23.52) yields

$$\mathcal{E}^{1/2}(0) \le C_{\epsilon,\epsilon_1} e^{\varrho_{\epsilon_1,-}(0,T_{\text{ode}})} e^{(\kappa_{\text{sil},+}+\epsilon)T_{\text{ode}}} |\hat{\chi}(\iota)|, \qquad (23.53)$$

where  $C_{\epsilon,\epsilon_1}$  only depends on  $\epsilon$ ,  $\epsilon_1$  and the coefficients of the equation (1.2). Moreover, the  $\epsilon$  can be removed in case the Jordan blocks of  $A_{\infty}$  corresponding to eigenvalues with real part  $\kappa_{\text{sil},+}$  are trivial. Keeping Definition 21.21 in mind,

$$\rho_{\epsilon_{1,-}}(0, T_{\text{ode}}) + (\kappa_{\text{sil},+} + \epsilon)T_{\text{ode}} = \sum_{j=1}^{k+1} (\mu_{j,-} + \kappa_{\text{sil},+} + \epsilon)|I_{j}|, \qquad (23.54)$$

where the intervals  $I_j$ ,  $j \in \{1, \ldots, k+1\}$ , are the ones into which  $[0, T_{ode}]$  is divided when appealing to Definition 21.21 and the numbers  $\mu_{j,-}$  are introduced in Definition 21.21. Rewriting the right hand side of (23.54) as in the right hand side of the first line of (23.26) and recalling (23.25) (with t' = 0 and  $t = T_{ode}$ ) and (23.30) yields

$$\varrho_{\epsilon_{1},-}(0,T_{\text{ode}}) + (\kappa_{\text{sil},+} + \epsilon)T_{\text{ode}} \le \max_{1\le j\le k+1} \left(-\frac{\mu_{j,-} + \kappa_{\text{sil},+} + \epsilon}{\bar{\beta}_{q_{j}}}\right)\ln\langle\nu\rangle + C,$$
(23.55)

where C only depends on the coefficients of the equation (1.2) and the spectra of  $(M_r, g_r), r \in \{1, \ldots, R\}$  (in order to obtain this conclusion we assume that  $\epsilon, \epsilon_1 \leq 1$ ). Note that the  $\epsilon$  appearing explicitly on the right hand side of (23.55) can be removed in case the Jordan blocks of  $A_{\infty}$  corresponding to eigenvalues with real part  $\kappa_{\text{sil},+}$  are trivial. Let  $s_{\text{h},-}$  be defined by (23.45). Then (23.55) implies

$$\varrho_{\epsilon_{1},-}(0,T_{\text{ode}}) + (\kappa_{\text{sil},+} + \epsilon)T_{\text{ode}} \le (s_{\text{h},-} + \epsilon_{2})\ln\langle\nu\rangle + C$$

which, combined with (23.53), yields

$$\mathcal{E}_{s}^{1/2}(0) \le C_{\epsilon,\epsilon_{1}} \langle \nu \rangle^{s+s_{\mathrm{h},-}+\epsilon_{2}} |\hat{\chi}(\iota)|.$$
(23.56)

Here  $\epsilon_2$  is a linear combination of  $\epsilon$  and  $\epsilon_1$  (where the coefficients only depend on the coefficients of the equation (1.2)). Moreover,  $C_{\epsilon,\epsilon_1}$  only depends on  $\epsilon$  and  $\epsilon_1$ , the coefficients of the equation (1.2) and the spectra of  $(M_r, g_r)$ ,  $r \in \{1, \ldots, R\}$ . In addition, if the criteria stated in Remark 23.12 are satisfied, then  $\epsilon$ ,  $\epsilon_1$  and  $\epsilon_2$  appearing in (23.56) can be removed, so that

$$\mathcal{E}_s^{1/2}(0) \le C \langle \nu \rangle^{s+s_{\mathrm{h},-}} |\hat{\chi}(\iota)|, \qquad (23.57)$$

where C only depends on the coefficients of the equation (1.2) and the spectra of  $(M_r, g_r)$ ,  $r \in \{1, \ldots, R\}$ . Combining (23.51) and (23.56) yields the conclusion that (23.56) holds regardless of whether  $T_{\text{ode}}$  equals zero or not. Moreover, the improvement (23.57) holds in case the conditions stated in Remark 23.12 are fulfilled.

Let  $\Phi_{\beta,\infty}$  be the map taking  $\chi \in C^{\infty}(\bar{M}, E_a)$  to the element of  $C^{\infty}(\bar{M}, \mathbb{C}^{2m})$  whose *i*'th mode is given by v(0), where v is constructed as above. Then (23.56) implies that (23.44) holds; note that this implies that  $\Phi_{\beta,\infty}$  maps  $\chi \in C^{\infty}(\bar{M}, E_a)$  to an element of  $C^{\infty}(\bar{M}, \mathbb{C}^{2m})$ . Moreover, Remark 23.12 holds. Since  $\Phi_{\beta,\infty}$  is injective on the level of Fourier coefficients (this is a consequence of Lemma 9.21), it follows that  $\Phi_{\beta,\infty}$  is injective. Next, let  $\chi \in C^{\infty}(\bar{M}, E_a)$  and let u be the solution to (1.2) (with f = 0) satisfying (23.46). Due to Proposition 23.1, we know that (23.47) holds with  $\chi$  replaced by some  $V_{\infty} \in C^{\infty}(\bar{M}, E_a)$ . Moreover, the constants  $C_{\epsilon,\beta}$  and N have the dependence stated in Proposition 23.1. In order to determine  $V_{\infty}$  in terms of  $\chi$ , note that (23.47) (with  $\chi$  replaced by  $V_{\infty}$ ) implies that

$$|v(t) - e^{A_{\infty}t}v_{\infty}| \le C\langle t \rangle^N e^{(\kappa_{\rm sil,+} - \beta)t}$$
(23.58)

for all  $t \ge 0$ , where *C* depends on the initial data and  $\iota$ ; *v* is defined as in (10.16) where *z* is the  $\iota$ 'th mode of *u*; and  $v_{\infty}$  is the  $\iota$ 'th mode of  $V_{\infty}$ . Due to the construction of  $\Phi_{\beta,\infty}$ , we know that the estimate (23.58) holds with  $v_{\infty}$  replaced by  $\hat{\chi}(\iota)$ . Thus

$$\left|e^{A_{\infty}t}[\hat{\chi}(\iota) - v_{\infty}]\right| \le C \langle t \rangle^{N} e^{(\kappa_{\mathrm{sil},+} - \beta)t}$$

for all  $t \ge 0$ . Since  $\hat{\chi}(\iota) - v_{\infty} \in E_a$ , this estimate implies that  $\hat{\chi}(\iota) - v_{\infty} = 0$ . Thus  $V_{\infty} = \chi$ , and the lemma follows, except for the statement concerning surjectivity.

Assume that  $E_a = \mathbb{C}^{2m}$  and let  $\psi \in C^{\infty}(\bar{M}, \mathbb{C}^{2m})$ . We wish to demonstrate that  $\psi$  is in the image of  $\Phi_{\beta,\infty}$ . Let u be the solution to (1.2) with initial data given by (23.46), where the right hand side has been replaced by  $\psi$ . Appealing to Proposition 23.1 yields a  $V_{\infty} \in C^{\infty}(\bar{M}, \mathbb{C}^{2m})$  such that (23.3) holds (with f = 0). Let  $\bar{u}$  be the solution to (1.2) with initial data given by (23.46), where the right hand side has been replaced by  $\Phi_{\beta,\infty}(V_{\infty})$ . Then, by the above arguments, (23.3) holds (with f = 0) and u replaced by  $\bar{u}$ . In particular, there is thus a constant, depending on u,  $\bar{u}$  etc., such that

$$\left\| \begin{pmatrix} u(\cdot,t) \\ u_t(\cdot,t) \end{pmatrix} - \begin{pmatrix} \bar{u}(\cdot,t) \\ \bar{u}_t(\cdot,t) \end{pmatrix} \right\|_{(s)} \le C_{\epsilon,\beta} \langle t \rangle^N e^{(\kappa_{\mathrm{sil},+}-\beta)t}$$
(23.59)

for all  $t \ge 0$  and all  $s \in \mathbb{R}$ . Let z and  $\bar{z}$  be the Fourier coefficients of u and  $\bar{u}$  respectively, let v and  $\bar{v}$  be defined in analogy with (10.16), starting with z and  $\bar{z}$  respectively. Finally, let  $V = v - \bar{v}$ . If  $V(\iota, 0) = 0$  for all  $\iota \in \mathcal{I}_{\rm B}$ , then  $u = \bar{u}$  (since  $V(\iota, \cdot)$  solves a homogeneous equation,  $V(\iota, 0) = 0$  for all  $\iota \in \mathcal{I}_{\rm B}$  implies that  $V(\iota, t) = 0$  for all  $\iota \in \mathcal{I}_{\rm B}$  and all  $t \in I$ ), so that  $\Phi_{\beta,\infty}(V_{\infty}) = \psi$ . Thus  $\psi$  is in the image of  $\Phi_{\beta,\infty}$ . Assume now that there is a  $\iota \in \mathcal{I}_{\rm B}$  such that  $V(\iota, 0) \neq 0$ . Note that  $V(\iota, \cdot)$  is a solution to an equation to which Lemma 9.21 applies. Moreover, applying Lemma 9.21 yields a  $\Psi_{\infty}$  which is linear and bijective. Since  $V(\iota, 0) \neq 0$ , there is a  $0 \neq \chi \in \mathbb{C}^{2m}$  such that  $\Psi_{\infty}(\chi) = V(\iota, T_{\rm ode})$ , where  $T_{\rm ode}$  is determined by the  $\iota$  under consideration. This means that

$$|V(\iota, t) - e^{A_{\infty}t}\chi| \le C\langle t \rangle^N e^{(\kappa_{\rm sil, +} - \beta)t}$$

for all  $t \ge T_{\text{ode}}$  and a constant C depending on V,  $\iota$  etc. On the other hand, (23.59) implies that the same estimate holds with  $\chi$  set to zero. Due to the fact that  $\operatorname{Rsp} A_{\infty} < \beta$ , these estimates are contradictory. To conclude, every element of  $C^{\infty}(\overline{M}, \mathbb{C}^{2m})$  is in the image of  $\Phi_{\beta,\infty}$ . The lemma follows.

## Part VII Appendices

### Chapter 24

# Spectral aspects of differential operators and matrices

The main object of study in these notes is the equation (1.2). This equation is separable, and the main purpose of the present chapter is to derive the equations satisfied by the modes. We provide the relevant material in Section 24.1. To begin with, we recall the spectral properties of the Laplace-Beltrami operator on a closed Riemannian manifold in Subsection 24.1.1. We then turn to the product setting in Subsection 24.1.2; this is the setting relevant in the study of (1.2). With this background material available, we are in a position to separate the equation. This is the topic of Subsection 24.1.3. Finally, in Subsection 24.1.4, we express the energy in terms of the modes.

The second purpose of the present chapter is to recall some of the basic properties of the Jordan normal form of a matrix, as well as of matrix exponentials of matrices. This is the subject of Section 24.2.

#### 24.1 Dividing the solution into modes

The equation (1.2) is separable. It is therefore possible to analyse the behaviour of solutions by considering the individual modes. In fact, the arguments presented in these notes are based on this perspective, as are the statements of several of the results. Before proceeding, we therefore here present the background relevant to separating (1.2). As a starting point, let us recall some basic facts concerning the Laplace-Beltrami operator on a closed Riemannian manifold.

#### 24.1.1 Spectral properties of the Laplace-Beltrami operator on a closed Riemannian manifold

To begin with, let us recall the following result.

**Theorem 24.1.** Let (M, g) be a closed Riemannian manifold and  $\Delta_g$  be the associated Laplace-Beltrami operator. Then the eigenvalues of  $\Delta_g$  consist of a sequence  $\lambda_i$ ,  $0 \leq i \in \mathbb{Z}$ , such that  $0 = \lambda_0 > \lambda_1 > \cdots$  and  $\lambda_i \to -\infty$  as  $i \to \infty$ . Moreover, if  $\mathcal{E}_i$  is the eigenspace corresponding to  $\lambda_i$ , then  $\mathcal{E}_i$  is finite dimensional and consists of smooth functions. In particular  $\mathcal{E}_0$  is the set of constant functions. Finally, the set of eigenfunctions of  $\Delta_g$  is a basis for  $L^2(M)$ .

**Remark 24.2.** Let (M, g) be an oriented Riemannian manifold. There is an associated volume form  $\mu_q$ . This volume form defines a linear functional  $\Lambda$  on  $C_0(M, \mathbb{C})$  (the set of complex valued

continuous functions on M with compact support):

$$\Lambda(f) = \int_M f\mu_g.$$

Using this linear functional, we can appeal to the Riesz Representation Theorem; cf., e.g., [43, Theorem 2.14, p. 40]. This yields the existence of a positive measure  $\lambda_g$  and a  $\sigma$ -algebra  $\mathcal{A}$  of subsets of M, such that  $(M, \mathcal{A}, \lambda_g)$  is a complete measure space. Moreover,  $\mathcal{A}$  contains all the Borel sets and

$$\int_M f\lambda_g = \int_M f\mu_g$$

for all  $f \in C_0(M, \mathbb{C})$ . Given the measure  $\lambda_g$ , we can define  $L^p(M)$ . When we do so, the relevant underlying Riemannian metric should be understood from the context, and we take it for granted that the functions in question are complex-valued.

**Remark 24.3.** Since each of the eigenspaces is finite dimensional, there is a sequence  $0 = \overline{\lambda}_0 \geq \overline{\lambda}_1 \geq \cdots$ , for  $0 \leq i \in \mathbb{Z}$ , and corresponding orthonormal eigenfunctions  $\varphi_i \in C^{\infty}(M)$ , such that the sequence  $\varphi_i$  is a basis for  $L^2(M)$ . Moreover, there is a sequence  $0 \leq \nu_0 \leq \nu_1 \leq \cdots$  such that  $\overline{\lambda}_i = -\nu_i^2$ . Note that  $\nu_i \to \infty$  as  $i \to \infty$ . In what follows, we assume all closed manifolds to be connected. In that setting  $\nu_0 = 0$  and  $\nu_1 > 0$ .

Let (M, g) be a closed (connected and oriented) Riemannian manifold and  $\varphi_i$  and  $\nu_i$  be defined as in Remark 24.3. Let  $\mu_g$  be the volume form associated with g and  $\lambda_g$  be the measure defined by Remark 24.2. Define the inner product

$$\langle u, v \rangle_g := \int_M u v^* \lambda_g$$

for  $u, v \in L^2(M)$ , where \* denotes complex conjugation; this turns  $L^2(M)$  into a complex Hilbert space. For  $u \in L^2(M)$  and  $0 \leq i \in \mathbb{Z}$ , we also let  $\hat{u}(i) := \langle u, \varphi_i \rangle_g$ . The statement that  $\varphi_i$  is a basis of  $L^2(M)$ , cf. Theorem 24.1, is equivalent to the statement that

$$\int_{M} |u|^2 \lambda_g = \sum_{i=0}^{\infty} |\hat{u}(i)|^2$$
(24.1)

for all  $u \in L^2(M)$ ; cf. [43, Theorem 4.18, p. 85]. If  $u \in C^{\infty}(M, \mathbb{C})$  and  $s \in \mathbb{R}$ , let

$$||u||_{(s)} := \left(\sum_{i=0}^{\infty} \langle \nu_i \rangle^{2s} |\hat{u}(i)|^2\right)^{1/2}.$$
(24.2)

If s = k, where  $0 \le k \in \mathbb{Z}$ , then this norm is equivalent to the standard  $H^k(M)$  Sobolev norm associated with the metric g. From now on, we therefore loosely refer to the  $\|\cdot\|_{(s)}$  as Sobolev norms. Moreover, in these notes,  $H^s(M)$  is defined to be the set of complex valued distributions such that the right hand side of (24.2) is finite, where the Riemannian metric should be understood from the context. We use notation similar to (24.2) if  $u \in C^{\infty}(M, \mathbb{C}^n)$ ; in that case, the  $\hat{u}(i)$  are computed component-wise.

#### 24.1.2 Spectral properties, the product setting

Consider (1.2), where the coefficients have the properties stated in connection with (1.2). For  $r = 1, \ldots, R$ , we apply Remark 24.3 to the Laplace-Beltrami operator  $\Delta_{g_r}$  on  $(M_r, g_r)$ . This yields orthonormal eigenfunctions  $\varphi_{r,i}$ ,  $0 \leq i \in \mathbb{Z}$ , of  $\Delta_{g_r}$  and corresponding eigenvalues  $-\nu_{r,i}^2$ . Let  $n \in \mathbb{Z}^d$ ,  $0 \leq i_r \in \mathbb{Z}$ ,  $r = 1, \ldots, R$ , and

$$\iota := (n, i_1, \dots, i_R); \tag{24.3}$$

#### 24.1. DIVIDING THE SOLUTION INTO MODES

we denote the set of such  $\iota$  by  $\mathcal{I}_{B}$ . For each  $\iota \in \mathcal{I}_{B}$ , there is a uniquely associated  $\nu$  given by

$$\nu(\iota) := (n, \nu_{1,i_1}, \dots, \nu_{R,i_R}).$$
(24.4)

In case (24.4) holds, we also write, by slight abuse of notation,

$$\nu_{\rm T}(\iota) = n, \quad \nu_{{\rm T},j}(\iota) = n_j, \quad \nu_{r,i_r}(\iota) = \nu_{r,i_r}.$$
 (24.5)

Let  $x \in \mathbb{T}^d$  (where we think of  $\mathbb{T}^d$  as  $[0, 2\pi]^d$  with the ends identified),  $p_r \in M_r$ ,  $r = 1, \ldots, R$ , and  $p = (x, p_1, \ldots, p_R)$ . Given  $\iota \in \mathcal{I}_B$ , define  $\varphi_\iota$  by

$$\varphi_{\iota}(p) := (2\pi)^{-d/2} e^{in \cdot x} \varphi_{1,i_1}(p_1) \cdots \varphi_{R,i_R}(p_R).$$
(24.6)

Note that  $\varphi_{\iota}$  is an eigenfunction of the operator  $\Delta_{\rm T} + \Delta_1 + \cdots + \Delta_R$ , and the eigenvalue corresponding to  $\varphi_{\iota}$  is  $-|\nu(\iota)|^2$ ; here  $\Delta_{\rm T}$  is the standard Laplace operator on  $\mathbb{T}^d$ . Introduce the volume form  $\mu_{\rm B}$  on  $\overline{M}$  by (1.20), where  $\overline{M}$  is given by (1.3). Let  $\lambda_{\rm B}$  be obtained from  $\mu_{\rm B}$  in the same way that  $\lambda_g$  is obtained from  $\mu_g$  in Remark 24.2. Using the measure  $\lambda_{\rm B}$ , we can think of  $L^2(\overline{M})$  as a complex Hilbert space, in analogy with the above. We denote the corresponding inner product by

$$\langle u, v \rangle_{\mathcal{B}} := \int_{\bar{M}} u v^* \lambda_{\mathcal{B}}$$
 (24.7)

for  $u, v \in L^2(\overline{M})$ , where the star denotes complex conjugation. If  $\iota_a, \iota_b \in \mathcal{I}_B$ , it is clear that  $\langle \varphi_{\iota_a}, \varphi_{\iota_b} \rangle_B = 0$  unless  $\iota_a = \iota_b$ . Moreover,  $\langle \varphi_{\iota}, \varphi_{\iota} \rangle_B = 1$  for  $\iota \in \mathcal{I}_B$ . Let  $\iota$  be given by (24.3) and  $u \in L^2(\overline{M})$ . Define  $\hat{u}(\iota)$  by

$$\hat{u}(\iota) := \langle u, \varphi_{\iota} \rangle_{\mathcal{B}}.$$
(24.8)

Appealing successively to the fact that  $e^{in \cdot x}$ ,  $n \in \mathbb{Z}^d$ , is a basis of  $L^2(\mathbb{T}^d)$  and the fact that  $\varphi_{r,i}$  is a basis of  $L^2(M_r)$ ,  $r = 1, \ldots, R$ , it can be verified that if  $u \in C^{\infty}(\overline{M}, \mathbb{C})$ ,

$$\int_{\bar{M}} |u|^2 \mu_{\rm B} = \sum_{\iota \in \mathcal{I}_{\rm B}} |\hat{u}(\iota)|^2.$$
(24.9)

In particular, it is thus clear that  $\varphi_{\iota}, \iota \in \mathcal{I}_{\mathrm{B}}$ , is a basis for  $L^{2}(\bar{M})$ . If  $s \in \mathbb{R}$  and  $u \in C^{\infty}(\bar{M}, \mathbb{C})$ , define

$$||u||_{(s)} := \left(\sum_{\iota \in \mathcal{I}_{\mathrm{B}}} \langle \nu(\iota) \rangle^{2s} |\hat{u}(\iota)|^2\right)^{1/2}.$$
(24.10)

If  $0 \leq k \in \mathbb{Z}$ ,  $\|\cdot\|_{(k)}$  is a norm which is equivalent to the standard Sobolev  $H^k$ -norms, and we therefore refer to the  $\|\cdot\|_{(s)}$  as Sobolev norms or  $H^s$ -norms. We also define  $H^s(\bar{M})$  as at the end of Subsection 24.1.1. We use a notation similar to (24.10) for  $u \in C^{\infty}(\bar{M}, \mathbb{C}^n)$ . It is also worth noting that a combination of (24.9) and polarisation type identities yields

$$\langle u, v \rangle_{\mathrm{B}} = \sum_{\iota \in \mathcal{I}_{\mathrm{B}}} \hat{u}(\iota) \hat{v}(\iota)^*;$$
(24.11)

cf. [43, Theorem 4.18, p. 85].

#### 24.1.3 Fourier decomposition of the main equation

Consider (1.2). Assume the associated Lorentz manifold (M, g) to be a canonical separable cosmological model manifold. Taking the inner product of (1.2) with  $\varphi_{\iota}$  (with respect to  $\langle \cdot, \cdot \rangle_{\rm B}$ introduced in (24.7)) yields

$$\ddot{z}(\iota, t) + \mathfrak{g}^{2}(\iota, t)z(\iota, t) - 2\sum_{l=1}^{d} in_{l}g^{0l}(t)\dot{z}(\iota, t) +\alpha(t)\dot{z}(\iota, t) + \sum_{l=1}^{d} in_{l}X^{l}(t)z(\iota, t) + \zeta(t)z(\iota, t) = \hat{f}(\iota, t),$$
(24.12)

where  $n_j := \nu_{\mathrm{T},j}(\iota)$ ,

$$\mathfrak{g}(\iota,t) := \left(\sum_{j,l=1}^{d} g^{jl}(t) n_j n_l + \sum_{r=1}^{R} a_r^{-2}(t) \nu_{r,i_r}^2(\iota)\right)^{1/2}$$
(24.13)

$$z(\iota, t) := \langle u(\cdot, t), \varphi_\iota \rangle_{\mathcal{B}}, \tag{24.14}$$

$$\hat{f}(\iota, t) := \langle f(\cdot, t), \varphi_{\iota} \rangle_{\mathcal{B}}.$$
(24.15)

For the sake of brevity, it is convenient to omit reference to the arguments  $\iota$  and t in (24.12), and to use the Einstein summation convention for sums from 1 to d. Then (24.12) can be written

$$\ddot{z} + \mathfrak{g}^2 z - 2in_l g^{0l} \dot{z} + \alpha \dot{z} + in_l X^l z + \zeta z = \hat{f}.$$
(24.16)

Most of the arguments in these notes are devoted to a study of this equation. For  $0 \neq \iota \in \mathcal{I}_B$ , it is often convenient to introduce the notation

$$\sigma(\iota, t) := \frac{n_l g^{0l}(t)}{\mathfrak{g}(\iota, t)}, \quad X(\iota, t) := \frac{n_l X^l(t)}{\mathfrak{g}(\iota, t)}.$$
(24.17)

Then (24.16) can be written

$$\ddot{z} + \mathfrak{g}^2 z - 2i\sigma \mathfrak{g}\dot{z} + \alpha \dot{z} + iX\mathfrak{g}z + \zeta z = \hat{f}.$$
(24.18)

#### 24.1.4 Higher order energies

In Section 1.4 we introduce the basic energies of interest in these notes. However, it is also of interest to introduce higher order energies. In particular, we define

$$\mathfrak{E}_{s}[u](t) := \frac{1}{2} \sum_{\iota \in \mathcal{I}_{\mathrm{B}}} \langle \nu(\iota) \rangle^{2s} \left[ |\dot{z}(\iota, t)|^{2} + \mathfrak{g}^{2}(\iota, t)|z(\iota, t)|^{2} + |z(\iota, t)|^{2} \right].$$
(24.19)

Here  $u \in C^{\infty}(M, \mathbb{C}^m)$  and z is defined by (24.14). In what follows, we also use the notation  $\mathfrak{E} := \mathfrak{E}_0$ . Note that  $\mathfrak{E} = \mathfrak{E}_{\text{bas}}$ , where  $\mathfrak{E}_{\text{bas}}$  is defined in (1.19).

#### 24.2 The Jordan normal form

Let  $A \in \mathbf{M}_n(\mathbb{C})$ . Then A can be decomposed into Jordan normal form. This means that there is a matrix  $T \in \mathbf{GL}_n(\mathbb{C})$  such that  $J := T^{-1}AT$  is a block diagonal matrix. Moreover, the matrices on the diagonal are Jordan blocks; i.e., they are square matrices of the form

$$J_{\lambda,d} = \begin{pmatrix} \lambda & 1 & 0 & \cdots & 0 & 0 \\ 0 & \lambda & 1 & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & \lambda & 1 \\ 0 & 0 & 0 & \cdots & 0 & \lambda \end{pmatrix}.$$

Equivalently, all the elements on the diagonal of  $J_{\lambda,d}$  are  $\lambda$ , all the elements immediately above the diagonal are 1, and all the remaining elements are 0. Moreover, d is the dimension of the Jordan block; i.e.,  $J_{\lambda,d} \in \mathbf{M}_d(\mathbb{C})$ . Finally,  $\lambda \in \mathrm{Sp}(A)$ , where we use the notation introduced in Definition 4.3.

**Properties of Jordan blocks.** Before considering the matrix A itself, it is useful to describe the properties of isolated Jordan blocks. Let us begin by computing the generalised eigenspace of

#### 24.2. THE JORDAN NORMAL FORM

 $B := J_{\lambda,d}$ . Using the notation of Definition 4.7, it is clear that  $n_{\lambda} = d$ . Moreover  $(B - \lambda \mathrm{Id}_d)^{n_{\lambda}} = 0$ . This means that the generalised eigenspace of B, i.e.  $E_{\lambda}$ , equals  $\mathbb{C}^d$ . Next,  $e^{Bt}$  is given by

$$\exp(J_{\lambda,d}t) = e^{\lambda t} \exp[J_{0,d}t] = e^{\lambda t} \sum_{k=0}^{\infty} \frac{1}{k!} J_{0,d}^k t^k = e^{\lambda t} \sum_{k=0}^{d-1} \frac{1}{k!} J_{0,d}^k t^k;$$
(24.20)

note that  $J_{0,d} = J_{\lambda,d} - \lambda \mathrm{Id}_d$ ; that  $J_{0,d}^{d-1} \neq 0$ ; that  $J_{0,d}^d = 0$ ; and that  $J_{\lambda,d}$  commutes with  $\lambda \mathrm{Id}_d$ . In fact, the k'th power of the matrix  $J_{0,d}$  has ones k steps above the diagonal, and all the other elements are zero. In particular, for generic  $v \in \mathbb{C}^d$ , the leading order term in  $\exp(J_{\lambda,d}t)v$  (as  $t \to \infty$ ) has time dependence of the form  $t^{d-1} \exp(\mathrm{Re}\{\lambda\}t)$ .

Generalised eigenspaces of A. In general, J is of the form

$$J = \operatorname{diag}\{J_{\lambda_1, d_1}, \dots, J_{\lambda_m, d_m}\}.$$

Note that the  $\lambda_i$  need not be distinct. It is convenient to order the blocks. From now on, we assume that  $\operatorname{Re}\{\lambda_i\} \geq \operatorname{Re}\{\lambda_{i+1}\}$  and that  $d_i \geq d_{i+1}$  in case  $\operatorname{Re}\{\lambda_i\} = \operatorname{Re}\{\lambda_{i+1}\}$ . Assume now that  $\lambda$  is an eigenvalue of A and that  $\lambda_l, \ldots, \lambda_k$  are all the  $\lambda_i$  that equal  $\lambda$ , where we can assume that  $1 \leq l \leq k \leq m$ . In that case, it is clear that  $n_{\lambda} = d_l + \cdots + d_k$ . Moreover, the generalised eigenspace of J corresponding to  $\lambda$  equals  $\{0\}^{m_a} \times \mathbb{C}^{n_\lambda} \times \{0\}^{m_b}$ , where  $m_a = d_1 + \cdots + d_{l-1}$  and  $m_b = d_{k+1} + \cdots + d_m$ . On the other hand,  $v \in \mathbb{C}^n$  belongs to the generalised eigenspace of A corresponding to  $\lambda$ . This allows us to calculate the generalised eigenspace of A corresponding to  $\lambda$ .

Matrix exponentials. In order to calculate  $e^{At}$ , given the above decomposition into Jordan normal form, it is sufficient to note that

$$e^{At} = Te^{Jt}T^{-1} = T\text{diag}\{\exp(J_{\lambda_1,d_1}t),\ldots,\exp(J_{\lambda_m,d_m}t)\}T^{-1}.$$

Here  $\exp(J_{\lambda_i,d_i}t)$  can be calculated as in (24.20). In particular, it is clear that for generic  $v \in \mathbb{C}^d$ , the leading order term in  $e^{At}v$  (as  $t \to \infty$ ) has a time dependence of the form  $t^{d_1-1}\exp(\operatorname{Re}\{\lambda_1\}t)$ . This motivates the introduction of  $\kappa_{\max}$  and  $\lambda_{\max}$  in Definition 4.3. In fact,  $\kappa_{\max}(A) = \operatorname{Re}\{\lambda_1\}$ and  $d_{\max}(A, \kappa_{\max}) = d_1$  by construction.

First generalised eigenspace in the  $\beta$ , A-decomposition of  $\mathbb{C}^n$ . In these notes, we often want to solve a system of ODE's for which there is a leading order behaviour corresponding to a model equation  $\dot{x} = Ax$ . However, there are also error terms. As a consequence, it is natural to divide the solutions to the model equation into a leading order part, which can be distinguished from the error terms, and a remainder, which cannot. This division can be made already on the level of the generalised eigenspaces of A, and this is the purpose of Definition 4.7. We already know that the leading order behaviour of solutions is  $t^{d_1-1} \exp(\operatorname{Re}\{\lambda_1\}t)$ . The cut-off, distinguishing between the leading order part and the remainder, is characterised by a positive real number, say  $\beta > 0$ . Roughly speaking we want to keep the part of the solution corresponding to eigenvalues with real part strictly larger than  $\operatorname{Re}\{\lambda_1\} - \beta$ . The remaining part of the solution we do not want to keep. On the transformed, Jordan block, side, we thus want to focus on vectors  $v \in \mathbb{C}^n$  such that the components of v corresponding to the Jordan blocks  $J_{\lambda_i, d_i}$  vanish if  $\operatorname{Re}\{\lambda_i\} \leq \operatorname{Re}\{\lambda_1\} - \beta$ . On the original, untransformed, side, this corresponds to focusing on v in the first generalised eigenspace in the  $\beta$ , A-decomposition of  $\mathbb{C}^n$ . On the Jordan block side, what remains, after removing the leading order part, is vectors  $v \in \mathbb{C}^n$  such that the components of v corresponding to the Jordan blocks  $J_{\lambda_i, d_i}$  vanish if  $\operatorname{Re}\{\lambda_i\} > \operatorname{Re}\{\lambda_1\} - \beta$ . On the original, untransformed, side, this corresponds to focusing on v in the second generalised eigenspace in the  $\beta$ , A-decomposition of  $\mathbb{C}^n$ ; cf. Definition 9.4.

374CHAPTER 24. SPECTRAL ASPECTS OF DIFFERENTIAL OPERATORS AND MATRICES

## Chapter 25

## Geometric preliminaries

The purpose of the present part of these notes is to derive basic conclusions concerning the geometry that arises in the study of equations of the form (1.2). Moreover, since the results of Parts II–VI are based on analytical conditions, we need to relate these analytical conditions with the geometric conditions of Part I.

#### 25.1 Metrics of interest, lapse and shift decomposition

Consider (1.2). By assumption,  $g^{00}(t) < 0$  and  $g^{ij}(t)$ ,  $i, j \in \{1, \ldots, d\}$ , are the components of a positive definite matrix for all  $t \in I$ . As a consequence, there are smooth functions  $g_{\alpha\beta}$ ,  $\alpha, \beta \in \{0, \ldots, d\}$ , from I to  $\mathbb{R}$  such that  $g_{\alpha\beta}(t)$  are the components of the inverse of the matrix with components  $g^{\alpha\beta}(t)$ . Moreover,  $g_{00}(t) < 0$  and  $g_{ij}(t)$  are the components of a positive definite matrix for all t. These statements are a consequence of [34, Lemma 8.5, p. 72]. In [34, Section 8.1], it is also demonstrated that if  $g_{00}(t) < 0$  and  $g_{ij}(t)$  are the components of a positive definite matrix, then the matrix with components  $g_{\alpha\beta}(t)$  has one strictly negative and d strictly positive eigenvalues. Starting with the equation (1.2), we thus arrive at the Lorentz metric (1.4). Next, let us verify that (M, g) is a globally hyperbolic Lorentz manifold.

**Lemma 25.1.** Let (M, g) be a separable cosmological model manifold in the sense of Definition 1.2. Then (M, g) is a globally hyperbolic Lorentz manifold, and each hypersurface  $\overline{M}_t$ , defined by (1.5) for  $t \in I$ , is a Cauchy hypersurface.

*Proof.* It is sufficient to prove that the  $\overline{M}_t$ ,  $t \in I$ , are Cauchy hypersurfaces. Let  $\gamma$  be an inextendible causal curve and let  $\gamma^0$  denote the time component of  $\gamma$ . Then, letting  $\pi_0 : M \to I$  denote projection onto the last factor,

$$\frac{d\gamma^0}{ds} = \frac{d}{ds}\pi_0 \circ \gamma = \gamma'(\pi_0) = d\pi_0(\gamma') = \langle \operatorname{grad}\pi_0, \gamma' \rangle.$$
(25.1)

However,

$$\langle \operatorname{grad} \pi_0, \operatorname{grad} \pi_0 \rangle = g_{\alpha\beta} g^{0\alpha} g^{0\beta} = g^{00} < 0,$$

so that  $\operatorname{grad} \pi_0$  is timelike. For this reason, the right hand side of (25.1) is either always strictly positive or always strictly negative. In particular,  $\gamma^0$  is strictly monotone, so that  $\gamma$  can intersect a given  $\overline{M}_t$  at most once. Moreover, there are two possibilities: either the range of  $\gamma^0$  equals I, or it does not. In order to demonstrate that the second case leads to a contradiction, assume that the range of  $\gamma^0$  is contained in  $(-\infty, t_0), t_0 \in I$ , and that  $\gamma^0$  is strictly increasing (there are of course other cases to consider, but they can be dealt with in a similar fashion). If the domain of  $\gamma$  is  $J = (s_-, s_+)$ , then  $\gamma^0(s)$  converges to some  $t_1$  as  $s \to s_+$ . Moreover,  $t_1 \in I$ . Let  $\pi_1 : M \to \overline{M}$  denote the projection onto the first factor and  $\bar{\gamma} := \pi_1 \circ \gamma$ . If we could prove that  $\bar{\gamma}(s)$  converges as  $s \to s_+$ , then we would know that  $\gamma$  converges as  $s \to s_+$ . This would imply that  $\gamma$  is extendible and contradict the assumptions (and thereby finish the proof). In order to prove that  $\bar{\gamma}$  converges, note that due to the causality of  $\gamma$ ,

$$g_{00}\dot{\gamma}^{0}\dot{\gamma}^{0} + 2g_{0i}\dot{\gamma}^{0}\dot{\gamma}^{i} + g_{ij}\dot{\gamma}^{i}\dot{\gamma}^{j} + \sum_{r}a_{r}^{2}g_{r}(\dot{\gamma}_{r},\dot{\gamma}_{r}) \leq 0,$$

where  $\gamma_r$  denotes the  $M_r$ -component of  $\gamma$ . Using this estimate; the fact that for  $s_0 \in J$ ,  $\gamma^0(s)$  is contained in a compact subset of I for  $s \in [s_0, s_+)$ ; and the fact that  $\gamma^0$  converges, it can be argued that for any  $s_n \to s_+$ ,  $\bar{\gamma}(s_n)$  is a Cauchy sequence with respect to the metric induced by

$$\sum_{i,j} \delta_{ij} dx^i \otimes dx^j + \sum_r g_r. \tag{25.2}$$

Since (25.2) is a complete metric on  $\overline{M}$ , we conclude that  $\overline{\gamma}$  converges. This yields the desired contradiction. Thus the range of  $\gamma^0$  equals I, and  $\gamma^0$  is strictly increasing. In particular, given an inextendible causal curve  $\gamma$ , each  $\overline{M}_t$  is intersected once and only once by  $\gamma$ , so that each  $\overline{M}_t$  is a Cauchy hypersurface.

#### 25.1.1 Lapse and shift

Let (M, g) be a separable cosmological model manifold in the sense of Definition 1.2. The first four terms on the right hand side of (1.6) can be written

$$h := g_{00}dt \otimes dt + g_{0i}dt \otimes dx^{i} + g_{i0}dx^{i} \otimes dt + g_{ij}dx^{i} \otimes dx^{j}$$
  
$$= -N^{2}dt \otimes dt + g_{ij}(\chi^{i}dt + dx^{i}) \otimes (\chi^{j}dt + dx^{j}), \qquad (25.3)$$

where the  $g_{ij}$  are the same functions as in (1.6) and  $\chi^i$  and N > 0 are determined by the relations

$$g_{ij}\chi^j = g_{0i}, \quad N := (-g_{00} + g_{ij}\chi^i\chi^j)^{1/2}.$$
 (25.4)

Since the  $g_{ij}$  are the components of a positive definite matrix, the first equation appearing in (25.4) can be solved in order to yield smooth functions  $\chi^i$ . Given  $\chi^i$ , the second equality in (25.4) defines the strictly positive function N. Note that the  $\chi^i$  can be thought of as the components of a vector field  $\chi = \chi^i \partial_i$  which we refer to as the *shift vector field*. The function N is referred to as the *lapse function*. On the other hand, if we start with a lapse function, a shift vector field and the  $g_{ij}$ 's (which we assume to be the components of a positive definite matrix for every  $t \in I$ ), then we have to assume that

$$g_{ij}\chi^i\chi^j < N^2 \tag{25.5}$$

in order for  $g_{00}$  (now considered to be defined by the second equality in (25.3)) to satisfy  $g_{00} < 0$ .

#### 25.2 The components of the inverse

Let (M, g) be a separable cosmological model manifold in the sense of Definition 1.2, and define h by (25.3). Given  $t \in I$ , we can consider  $h_{\alpha\beta}(t)$ ,  $\alpha, \beta = 0, \ldots, d$ , to be the components of an element of  $\mathbf{M}_{d+1}(\mathbb{R})$ . It is of interest to calculate the components of the inverse in terms of the the lapse function, the shift vector field and  $h_{ij}$ ; considering (25.3), we can consider these three objects to define h. Defining  $\bar{h}$  by

$$\bar{h} := h_{ij} dx^i \otimes dx^j, \tag{25.6}$$

we denote the components of  $\bar{h}$  by  $\bar{h}_{ij} = h_{ij} = g_{ij}$ ; i, j = 1, ..., d. For  $t \in I$ ,  $\bar{h}_{ij}(t)$ , i, j = 1, ..., d, are the components of a positive definite, symmetric element of  $\mathbf{M}_d(\mathbb{R})$ . We denote the components of the inverse by  $\bar{h}^{ij}(t)$ , i, j = 1, ..., d.

**Lemma 25.2.** Let (M, g) be a separable cosmological model manifold in the sense of Definition 1.2, and define h by (25.3). Fix  $t \in I$  and consider  $h_{\alpha\beta}(t)$ ,  $\alpha, \beta = 0, \ldots, d$ , to be the components of an element of  $\mathbf{M}_{d+1}(\mathbb{R})$ . This element is invertible. Denote the components of the inverse by  $h^{\alpha\beta}(t)$ ,  $\alpha, \beta = 0, \ldots, d$ . Then, if the lapse function N satisfies N = 1,

$$h^{00} = -1, \quad h^{0i} = \chi^i, \quad h^{ij} = \bar{h}^{ij} - \chi^i \chi^j,$$
 (25.7)

where  $\bar{h}^{ij}$  is defined immediately prior to the statement of the lemma. Moreover, for  $t \in I$ ,  $h^{ij}(t)$ , i, j = 1, ..., d, are the components of a positive definite matrix. Finally,

$$\det h = -\det \bar{h}$$

where h and  $\bar{h}$  are considered to be matrix valued functions taking  $t \in I$  to the elements of  $\mathbf{M}_{d+1}(\mathbb{R})$ and  $\mathbf{M}_d(\mathbb{R})$  with components  $h_{\alpha\beta}(t)$ ,  $\alpha, \beta = 0, \ldots, d$ , and  $\bar{h}_{ij}(t)$ ,  $i, j = 1, \ldots, d$ , respectively.

**Remark 25.3.** In case  $N \neq 1$ , we can apply the lemma to the metric  $\hat{h} := N^{-2}h$ . This yields

$$h^{00} = -N^{-2}, \quad h^{0i} = N^{-2}\chi^i, \quad h^{ij} = \bar{h}^{ij} - N^{-2}\chi^i\chi^j, \quad \det h = -N^2 \det \bar{h}.$$
 (25.8)

*Proof.* The invertibility follows from [34, Lemma 8.5, p. 72]. Fix  $t \in I$  and let  $a := h_{00}(t), v \in \mathbb{R}^d$  be the vector with components  $v_i = h_{0i}(t)$  and  $\overline{H} \in \mathbf{M}_d(\mathbb{R})$  be the matrix with components  $h_{ij}(t)$ ,  $i, j = 1, \ldots, d$ . Then the matrix with components  $h_{\alpha\beta}(t), \alpha, \beta = 0, \ldots, d$ , can be written

$$H := \left(\begin{array}{cc} a & v^t \\ v & \bar{H} \end{array}\right).$$

Thus

$$\begin{pmatrix} 1 & -v^t \bar{H}^{-1} \\ 0 & \mathrm{Id}_d \end{pmatrix} H \begin{pmatrix} 1 & 0 \\ -\bar{H}^{-1}v & \mathrm{Id}_d \end{pmatrix} = \begin{pmatrix} a - v^t \bar{H}^{-1}v & 0 \\ 0 & \bar{H} \end{pmatrix}.$$
 (25.9)

Note that the components of  $\overline{H}^{-1}v$  are given by

$$\bar{h}^{ij}(t)h_{0j}(t) = \bar{h}^{ij}(t)g_{jl}(t)\chi^{l}(t) = \chi^{i}(t),$$

and that

$$a - v^{t} \bar{H}^{-1} v = h_{00}(t) - \bar{h}^{ij}(t) h_{0i}(t) h_{0j}(t)$$
  
=  $h_{00}(t) - \bar{h}_{ij}(t) \chi^{i}(t) \chi^{j}(t) = -1;$  (25.10)

in the derivation of these equalities, we appealed to (25.3) and the fact that N = 1. Finally, the components of  $\bar{H}^{-1}vv^t\bar{H}^{-1}$  are given by

$$\bar{h}^{il}(t)v_lv_k\bar{h}^{jk}(t) = \chi^i(t)\chi^j(t).$$

Combining these observations with (25.9) yields (25.7). In order to prove that  $h^{ij}(t)$  are the components of a positive definite matrix, let A be a positive definite symmetric matrix such that  $A^2 = \overline{H}$ , let  $x \in \mathbb{R}^d$  be such that  $x^i = \chi^i(t)$  and let  $w \in \mathbb{R}^d$ . Then

$$|\chi^{i}(t)w_{i}| = |x \cdot w| = |A^{-1}w \cdot Ax| \le |A^{-1}w||Ax| = |x|_{\bar{H}}|w|_{\bar{H}^{-1}},$$
(25.11)

where  $|w|_B = (B_{ij}w^iw^j)^{1/2}$  for  $w \in \mathbb{R}^d$  and  $B \in \mathbf{M}_d(\mathbb{R})$  symmetric and positive semi-definite. Moreover,  $|x|_{\bar{H}} < 1$ , since  $h_{00} < 0$  (by assumption) and N = 1. Thus, if  $w \neq 0$  and  $x \neq 0$ ,

$$\chi^{i}(t)\chi^{j}(t)w_{i}w_{j} \leq |x|_{\bar{H}}^{2}|w|_{\bar{H}^{-1}}^{2} < \bar{h}^{lm}(t)w_{l}w_{m}.$$
(25.12)

Combining this estimate with (25.7), it is clear that  $h^{ij}(t)$  are the components of a positive definite matrix. The last statement of the lemma is an immediate consequence of (25.9) and (25.10).

#### 25.3 The second fundamental form

The main purpose of the present section is to derive some basic properties of the second fundamental form. Before proceeding, let us introduce coordinates adapted to the structure of the metric.

**Definition 25.4.** Let  $(\mathcal{O}_{\mathrm{T}}, \bar{x}_{\mathrm{T}})$  be standard local coordinates on  $\mathbb{T}^d$ ,  $(\mathcal{O}_r, \bar{x}_r)$  be local coordinates on  $M_r$  and  $\mathcal{O} := \mathcal{O}_{\mathrm{T}} \times \mathcal{O}_1 \times \cdots \times \mathcal{O}_R \times I$ . Let x be the coordinates mapping  $p = (\xi_{\mathrm{T}}, \xi_1, \ldots, \xi_R, t) \in \mathcal{O}$  to  $[t, \bar{x}_{\mathrm{T}}(\xi_{\mathrm{T}}), \bar{x}_1(\xi_1), \ldots, \bar{x}_R(\xi_R)]$ . Then  $(\mathcal{O}, x)$  are referred to as *canonical local coordinates* on M.

**Remarks 25.5.** By the statement that  $(\mathcal{O}_{\mathrm{T}}, \bar{x}_{\mathrm{T}})$  are standard local coordinates on  $\mathbb{T}^d$ , we mean that  $\partial/\partial \bar{x}_{\mathrm{T}}^1, \ldots, \partial/\partial \bar{x}_{\mathrm{T}}^d$  are the standard vector fields on  $\mathbb{T}^d$ . The components of x are labelled  $x^{\alpha}$ , where  $\alpha = 0, \ldots, D$  and D is the dimension of the manifold  $\bar{M}$  defined by (1.3). In particular,  $\partial_{x^i}, i = 1, \ldots, d$ , are tangent to the torus.

Turning to the second fundamental form, note that

$$U := -(-g^{00})^{-1/2}g^{0\mu}\partial_{\mu} = \frac{1}{N}(\partial_t - \chi^i \partial_i)$$

is the future directed unit normal to the hypersurfaces  $\overline{M}_t$ . If  $\overline{k}$  denotes the second fundamental form of  $\overline{M}_t$ ,  $(\mathcal{O}, x)$  are canonical local coordinates on M (cf. Definition 25.4) and  $\partial_i, \partial_j$  denote coordinate vector fields corresponding to x which are tangent to  $\overline{M}_t$ , then

$$\bar{k}_{ij} := \bar{k}(\partial_i, \partial_j) = \langle \nabla_{\partial_i} U, \partial_j \rangle = -(-g^{00})^{-1/2} g^{0\mu} \langle \nabla_{\partial_i} \partial_\mu, \partial_j \rangle$$
  
=  $-(-g^{00})^{-1/2} g^{0\mu} \frac{1}{2} (\partial_i g_{\mu j} + \partial_\mu g_{ij} - \partial_j g_{i\mu}) = \frac{1}{2} (-g^{00})^{1/2} \partial_t g_{ij}.$  (25.13)

In the last step, we used the following facts: if  $\alpha = 0, \ldots, d$ , then  $\partial_i g_{\alpha\mu} = 0$  for all  $i = 1, \ldots, D$ and all  $\mu = 0, \ldots, D$ ;  $g^{0i} = g_{0i} = 0$  if  $i = d + 1, \ldots, D$ ; and  $\partial_i g_{\mu\nu} = 0$  for all  $i = 1, \ldots, d$  and all  $\mu, \nu = 0, \ldots, D$ . Combining (25.8) and (25.13) yields

$$N\bar{k}_{ij} = \frac{1}{2}\partial_t g_{ij}.$$
(25.14)

Due to this equality, it follows that

$$N\bar{k} = \frac{1}{2} \sum_{i,j=1}^{d} \partial_t \bar{h}_{ij} dx^i \otimes dx^j + \sum_{r=1}^{R} \frac{\dot{a}_r}{a_r} a_r^2 \pi_r^* g_r, \qquad (25.15)$$

where  $\pi_r: \overline{M}_t \to M_r$  denotes the projection onto  $M_r$ ; note that the metric  $\overline{g}$  induced on  $\overline{M}_t$  by g can be written

$$\bar{g} = \sum_{i,j=1}^{d} \bar{h}_{ij} dx^{i} \otimes dx^{j} + \sum_{r=1}^{R} a_{r}^{2} \pi_{r}^{*} g_{r}.$$
(25.16)

Due to (25.15) and (25.16), it is clear that  $\bar{k}$  has a block structure, and that this block structure corresponds to a block structure in  $\bar{g}$ . It is therefore natural to introduce

$$\bar{k}_{\mathrm{T}} := \frac{1}{2N} \sum_{i,j=1}^{d} \partial_t \bar{h}_{ij} dx^i \otimes dx^j, \quad \bar{k}_r := \frac{1}{N} \frac{\dot{a}_r}{a_r} a_r^2 \pi_r^* g_r, \quad \bar{k}_{\mathrm{Rie}} := \sum_{r=1}^{R} \bar{k}_r.$$

Taking the trace of these tensor fields with respect to  $\bar{g}$  yields

$$\theta_{\mathrm{T}} := \mathrm{tr}_{\bar{g}}\bar{k}_{\mathrm{T}} = \frac{1}{2N}\bar{h}^{ij}\partial_t\bar{h}_{ij}, \quad \theta_r := \mathrm{tr}_{\bar{g}}\bar{k}_r = \frac{D_r}{N}\frac{\dot{a}_r}{a_r}, \tag{25.17}$$

where  $D_r$  is the dimension of  $M_r$ . For future reference, it is convenient to note that

$$\partial_t \ln \det \bar{h} = \bar{h}^{ij} \partial_t \bar{h}_{ij} = 2N \operatorname{tr}_{\bar{q}} \bar{k}_{\mathrm{T}} = 2N \theta_{\mathrm{T}}.$$
(25.18)

#### 25.4. GEOMETRIC CONVERGENCE

Summing up the above observations yields

$$\theta := \operatorname{tr}_{\bar{g}}\bar{k} = \frac{1}{N}\partial_t \ln \sqrt{\det \bar{h}} + \frac{D_1}{N}\frac{\dot{a}_1}{a_1} + \dots + \frac{D_R}{N}\frac{\dot{a}_R}{a_R}.$$
(25.19)

For future reference, it is also of interest to note that

$$|\bar{k}|_{\bar{g}}^{2} = |\bar{k}_{\rm T}|_{\bar{h}}^{2} + |\bar{k}_{\rm Rie}|_{\bar{g}}^{2} = |\bar{k}_{\rm T}|_{\bar{h}}^{2} + \sum_{r=1}^{R} \frac{D_{r}}{N^{2}} \left(\frac{\dot{a}_{r}}{a_{r}}\right)^{2}.$$
 (25.20)

Similarly, if N = 1,

$$|\mathcal{L}_U \bar{k}|_{\bar{g}}^2 = |(\mathcal{L}_U \bar{k})_{\mathrm{T}}|_{\bar{h}}^2 + \sum_{r=1}^R D_r \left[\frac{\ddot{a}_r}{a_r} + \left(\frac{\dot{a}_r}{a_r}\right)^2\right]^2.$$
 (25.21)

#### 25.4 Geometric convergence

Let (M, g) be a separable cosmological model manifold in the sense of Definition 1.2. In what follows, we use the terminology introduced in Definition 1.6. Moreover, we focus on manifolds such that Definitions 1.9 and 1.12 are satisfied. The purpose of the present section is to prove Lemma 1.14. There is a similar result in the case of big crunch asymptotics. However, we leave the statement and proof of the corresponding result to the reader.

Lemma 1.14. To begin with, we rescale g by  $e^{2\lambda}$ , for a function  $\lambda$  (depending only on t) yet to be determined. This yields  $\hat{g} = e^{2\lambda}g$ . Letting  $\hat{U} := e^{-\lambda}U$ , we then have

$$\hat{k}_{ij} = \frac{1}{2}\hat{U}\hat{g}_{ij} = \frac{1}{2}e^{-\lambda}U(e^{2\lambda}g_{ij}) = \hat{U}(\lambda)\hat{g}_{ij} + e^{\lambda}\bar{k}_{ij},$$

where  $\hat{k}$  is the second fundamental form induced on  $\bar{M}_t$  by  $\hat{g}$ . Raising the indices of this equality with  $\check{g}$  (the metric induced on  $\bar{M}_t$  by  $\hat{g}$ ) yields

$$\hat{k}^{i}_{\ j} = \hat{U}(\lambda)\delta^{i}_{\ j} + e^{-\lambda}\bar{k}^{i}_{\ j}.$$
(25.22)

Define  $\lambda$  by  $e^{-\lambda} \operatorname{tr}_{\bar{q}} \bar{k} = 1$ . Then,

$$\hat{U}(\lambda) = e^{-\lambda} U(\lambda) = -U(e^{-\lambda}) = -U[(\operatorname{tr}_{\bar{g}}\bar{k})^{-1}],$$

which is bounded by assumption. Thus  $|\hat{k}|_{\check{g}}$  is future bounded, as stated. Turning to the time coordinate  $\tau$ , it is clear by the assumptions that the interval  $[t_0, t_+)$  corresponds to  $[0, \infty)$  in  $\tau$ -time. Moreover, due to (1.8) and the adjacent comments,

$$e^{\lambda} = \mathrm{tr}_{\bar{q}}\bar{k} = U(\tau) = N^{-1}\partial_t \tau$$

In particular,  $Ne^{\lambda}dt = d\tau$ , and  $\hat{g}$  takes the form (1.15). Combining the assumption of convergence with (25.22) and the fact that  $e^{\lambda} = \text{tr}_{\bar{g}}\bar{k}$  yields the conclusion that there is a 2-tensor field  $\hat{A}$  of mixed type such that

$$|\hat{K} - \hat{A}|_{\varrho} \le C \exp[-\eta |\ln V(t)|]$$

for all  $t \ge t_0$ . Due to the definition of  $\tau$ , this estimate implies (1.16) and yields the conclusion of the lemma.

Due to this result, we, from now on, focus on separable cosmological model manifolds such that the lapse function equals 1 and I contains  $[0, \infty)$ .

#### 25.5 Norms

Before proceeding, it is convenient to make some observations of a purely linear algebra nature.

**Lemma 25.6.** Let V be a finite dimensional real vector space; let  $V^*$  denote the dual; and let g be an inner product on V. For  $v \in V$ , let  $|v|_g := [g(v, v)]^{1/2}$ . Let  $0 \le k, l \in \mathbb{Z}, k+l \ge 1$ , and

$$T: V^k \times (V^*)^l \to \mathbb{R}$$

be a multilinear map. Let  $\{e_i\}$  be an orthonormal basis of V and let  $\{e^i\}$  denote the dual basis. Define  $|T|_q$  by

$$|T|_g := \left(\sum |T(e_{i_1}, \dots, e_{i_k}, e^{j_1}, \dots, e^{j_l})|^2\right)^{1/2}.$$
(25.23)

Then  $|T|_g$  is independent of the choice of orthonormal basis. Moreover, if  $v_i \in V$ , i = 1, ..., k, and  $\eta^i \in V^*$ , i = 1, ..., l, then

$$|T(v_1, \dots, v_k, \eta^1, \dots, \eta^l)| \le |T|_g |v_1|_g \cdots |v_k|_g |\eta^1|_g \cdots |\eta^l|_g.$$
(25.24)

**Remark 25.7.** By an *inner product on* V, we mean a positive definite symmetric bilinear form on V.

**Remark 25.8.** Given  $v \in V$ ,  $v^{\flat} \in V^*$  is defined by  $v^{\flat}(w) = g(v, w)$  for all  $w \in V$ . With this notation, if  $\{e_i\}$  is an orthonormal basis, then  $e_i^{\flat} = e^i$  for all i. Note that  $\flat$  is an isomorphism from V to  $V^*$ . The inverse is denoted by  $\sharp$ ; i.e.,  $(v^{\flat})^{\sharp} = v$ . Note, also, that g induces an inner product on  $V^*$ , denoted  $g^{\sharp}$  and defined by  $g^{\sharp}(\eta, \xi) := g(\eta^{\sharp}, \xi^{\sharp})$  for all  $\eta, \xi \in V^*$ . In (25.24),  $|\eta^i|_g$  should be thought of as being defined by (25.23). Note, however, that if  $\{e_i\}$  is an ON basis,  $\{e^i\}$  is the dual basis and  $\eta_i e^i = \eta \in V^*$ , then  $\eta(e_i) = \eta_i$ ,  $\{e^i\}$  is an ON-basis of  $V^*$  with respect to  $g^{\sharp}$ , and

$$g^{\sharp}(\eta,\eta) = \eta_i \eta_j g^{\sharp}(e^i, e^j) = \sum \eta_i^2 = \sum |\eta(e_i)|^2 = |\eta|_g^2,$$

so that  $|\eta|_g = [g^{\sharp}(\eta, \eta)]^{1/2}$ .

*Proof.* We leave it to the reader to verify that  $|T|_g$  is independent of the choice of ON basis. In order to prove (25.24) in the case that k = 2 and l = 0, let  $v, w \in V$ ;  $\{e_i\}$  be an ON basis; and and define  $v^i, w^i$  by  $v = v^i e_i$  and  $w = w^i e_i$ . Then

$$|v|_g = [g(v,v)]^{1/2} = \left(\sum_i (v^i)^2\right)^{1/2}.$$

Moreover,

$$|T(v,w)| = \left|\sum_{i,j} T(e_i, e_j) v^i w^j\right| \le \left(\sum_j \left[\sum_i T(e_i, e_j) v^i\right]^2\right)^{1/2} |w|_g$$
  
$$\le \left(\sum_{i,j} [T(e_i, e_j)]^2\right)^{1/2} |v|_g |w|_g = |T|_g |v|_g |w|_g.$$
(25.25)

The proof of the general case is similar and left to the reader.

Under the assumptions of Lemma 25.6, let  $\{e_i\}$  be a basis of V which is not necessarily orthonormal. Let  $\{e^i\}$  be the dual basis;  $g_{ij} := g(e_i, e_j)$ ; and  $g^{ij} := g^{\sharp}(e^i, e^j)$ . Let  $G^{ij}$  be the components of the inverse of the metric with components  $g_{ij}$ . Note that there are  $a^{ij} \in \mathbb{R}$  such that  $(e^i)^{\sharp} = a^{ij}e_j$ . On the other hand,

$$g[(e^{i})^{\sharp}, e_{j}] = [(e^{i})^{\sharp}]^{\flat}(e_{j}) = e^{i}(e_{j}) = \delta^{i}_{j},$$
  
$$g[(e^{i})^{\sharp}, e_{j}] = g(a^{ik}e_{k}, e_{j}) = a^{ik}g_{kj}.$$

Thus  $a^{ij} = G^{ij}$ . Moreover,

$$g^{ij} = g^{\sharp}(e^i, e^j) = g[(e^i)^{\sharp}, (e^j)^{\sharp}] = G^{ik}G^{jl}g_{kl} = G^{ij}.$$

#### 25.5. NORMS

Let A and B be the symmetric positive definite matrices defined by the conditions that

$$g_{ij} = \sum_{l} B_{il} B_{lj}, \quad g^{ij} = \sum_{l} A^{il} A^{lj}$$

Then A is the inverse of B and  $f_i := A^{ij}e_j$  defines an orthonormal basis whose dual basis is given by  $f^i = B_{ik}e^k$  (we leave the verification of these statements to the reader). Let now T have the properties stated in Lemma 25.6. Introduce the notation

$$T_{i_1\cdots i_k}^{j_1\cdots j_l} = T(e_{i_1},\ldots,e_{i_k},e^{j_1},\ldots,e^{j_l})$$

Then

$$\begin{split} |T|_{g}^{2} &= \sum |T(f_{i_{1}}, \dots, f_{i_{k}}, f^{j_{1}}, \dots, f^{j_{l}})|^{2} \\ &= \sum A^{i_{1}n_{1}} \cdots A^{i_{k}n_{k}} A^{i_{1}r_{1}} \cdots A^{i_{k}r_{k}} B_{j_{1}m_{1}} \cdots B_{j_{l}m_{l}} B_{j_{1}s_{1}} \cdots B_{j_{l}s_{l}} T^{m_{1}\cdots m_{l}}_{n_{1}\cdots n_{k}} T^{s_{1}\cdots s_{l}}_{r_{1}\cdots r_{k}} \\ &= \sum g^{n_{1}r_{1}} \cdots g^{n_{k}r_{k}} g_{m_{1}s_{1}} \cdots g_{m_{l}s_{l}} T^{m_{1}\cdots m_{l}}_{n_{1}\cdots n_{k}} T^{s_{1}\cdots s_{l}}_{r_{1}\cdots r_{k}}. \end{split}$$

An alternate way to define the norm  $|T|_g$  is thus via

$$|T|_{g} = \left(\sum g^{n_{1}r_{1}} \cdots g^{n_{k}r_{k}} g_{m_{1}s_{1}} \cdots g_{m_{l}s_{l}} T^{m_{1}\cdots m_{l}}_{n_{1}\cdots n_{k}} T^{s_{1}\cdots s_{l}}_{r_{1}\cdots r_{k}}\right)^{1/2}.$$
 (25.26)

#### 25.5.1 Tensor estimates

The above analysis yields estimates that are of importance in what follows. In particular, if  $\lambda^i, \kappa_i \in \mathbb{R}, i = 1, ..., d$ , then

$$|\lambda^i \kappa_i| \le (\bar{h}_{ij} \lambda^i \lambda^j)^{1/2} (\bar{h}^{kl} \kappa_k \kappa_l)^{1/2}.$$
(25.27)

Moreover, if  $\kappa$  is a symmetric covariant 2-tensor field on  $\overline{M}$ , then

$$|\kappa(v,w)| \le |\kappa|_{\bar{g}}|v|_{\bar{g}}|w|_{\bar{g}} \tag{25.28}$$

for all  $v, w \in T\overline{M}$  with the same base point.

On a slightly more technical note, let  $\kappa$  be a symmetric covariant 2-tensor field on  $\mathbb{T}^d$ . Define a 2-tensor field T by

$$T_{ij} := \bar{h}^{kl} \kappa_{ik} \kappa_{jl}.$$

Then

$$|T|_{\bar{h}} \le |\kappa|_{\bar{h}}^2. \tag{25.29}$$

In order to prove this statement, note that since  $\kappa$  is symmetric, we can choose an orthonormal basis, say  $\{e_i\}$ , such that  $\kappa$  is diagonal, with diagonal components  $\kappa(e_i, e_i) = \kappa_i$ . Then

$$T(e_i, e_j) = \delta^{kl} \kappa_{ik} \kappa_{jl} = \sum_l \kappa_{il} \kappa_{jl} = \kappa_i^2 \delta_{ij}$$

(no summation on i). Thus

$$|T|_{\bar{h}}^{2} = \sum_{i,j} |T(e_{i}, e_{j})|^{2} = \sum_{i,j} \kappa_{i}^{4} \delta_{ij} = \sum_{i} \kappa_{i}^{4} \le \left(\sum_{i} \kappa_{i}^{2}\right)^{2}.$$
 (25.30)

On the other hand,

$$|\kappa|_{\bar{h}}^{2} = \sum_{i,j} \kappa_{ij}^{2} = \sum_{i,j} \kappa_{i}^{2} \delta_{ij} = \sum_{i} \kappa_{i}^{2}.$$
(25.31)

Combining (25.30) and (25.31) yields (25.29).

#### 25.6 Quantitative bounds on the shift vector field

Due to (25.5), there is a bound on the shift vector field in terms of the lapse function. Even though this estimate is of interest, stronger assumptions lead to more useful conclusions. Drawing some such conclusions is the subject of the next lemma.

**Lemma 25.9.** Let (M,g) be a canonical separable cosmological model manifold. Define h by (25.3). Assume that there is an  $\eta_{sh,0} \in \mathbb{R}$ , satisfying  $0 < \eta_{sh,0} \leq 1$ , such that

$$h(\partial_t, \partial_t) \le -\eta_{\mathrm{sh},0}^2 \tag{25.32}$$

for all  $t \geq 0$ . Then

$$\eta_{\mathrm{sh},0}^2 \bar{h}^{ij} v_i v_j \le h^{ij} v_i v_j \le \bar{h}^{ij} v_i v_j \tag{25.33}$$

for all  $v \in \mathbb{R}^d$  and all  $t \ge 0$ . Moreover,

$$\sup_{t \ge 0} \sup_{0 \ne v \in \mathbb{R}^d} \frac{|\chi^i v_i|}{(h^{ij} v_i v_j)^{1/2}} \le (\eta_{\mathrm{sh},0}^{-2} - 1)^{1/2}.$$
(25.34)

Similarly, if, in addition,  $\lambda^i \in \mathbb{R}$ ,  $i = 1, \ldots, d$ , then

$$\sup_{0 \neq v \in \mathbb{R}^d} \frac{|\lambda^i v_i|}{(h^{ij} v_i v_j)^{1/2}} \le \eta_{\mathrm{sh},0}^{-1} (\bar{h}_{ij} \lambda^i \lambda^j)^{1/2}$$
(25.35)

for all  $t \geq 0$ . In particular, if there is a constant  $0 < \kappa \in \mathbb{R}$  such that

$$\bar{h}_{ij}\lambda^i\lambda^j \le \kappa^2 \tag{25.36}$$

for all  $t \geq 0$ , then

$$\sup_{t \ge 0} \sup_{0 \neq v \in \mathbb{R}^d} \frac{|\lambda^i v_i|}{(h^{ij} v_i v_j)^{1/2}} \le \frac{\kappa}{\eta_{\mathrm{sh},0}}.$$
(25.37)

**Remark 25.10.** The estimate (25.32) is equivalent to

$$\bar{h}_{ij}\chi^i\chi^j \le 1 - \eta_{\rm sh,0}^2.$$
 (25.38)

Proof. Combining (25.7), (25.11) and (25.38) yields

$$\begin{split} h^{ij}v_iv_j = &\bar{h}^{ij}v_iv_j - \chi^i v_i\chi^j v_j \\ \ge &\bar{h}^{ij}v_iv_j - \bar{h}_{ij}\chi^i\chi^j \bar{h}^{lm}v_lv_m \ge \eta_{\mathrm{sh},0}^2 \bar{h}^{ij}v_iv_j. \end{split}$$

Thus (25.33) holds. One particular consequence of this estimate is that if  $v \neq 0$ , then

$$\frac{|\chi^i v_i|}{(h^{ij} v_i v_j)^{1/2}} \le \frac{(\bar{h}_{ij} \chi^i \chi^j)^{1/2} (\bar{h}^{lm} v_l v_m)^{1/2}}{(\eta_{\mathrm{sh},0}^2 \bar{h}^{ij} v_i v_j)^{1/2}} \le (\eta_{\mathrm{sh},0}^{-2} - 1)^{1/2}.$$

Thus (25.34) holds. The arguments justifying the remaining statements of the lemma are similar.  $\hfill \Box$ 

#### 25.7 From geometric to analytical conditions, part I

The conditions appearing in Parts II–VI are phrased in terms of  $\ell$ ,  $\sigma$ , X etc.; cf. (10.1). However, the conditions appearing in Part I are, largely, formulated in terms of the geometry. The purpose of the present part of these notes is to relate the two perspectives. In this section, we begin

by taking the step from conditions on the second fundamental form and the shift vector field to conditions on  $\ell$ ; recall that  $\mathfrak{g}(\iota, t)$  is defined by (24.13) and that  $\ell$  is defined by

$$\ell(\iota, t) := \ln \mathfrak{g}(\iota, t) \tag{25.39}$$

for  $0 \neq \iota \in \mathcal{I}_B$ . In what follows, it is useful to divide  $\mathfrak{g}^2$  into two parts. We therefore introduce the notation

$$\mathfrak{g}_{\mathrm{T}}(\iota, t) := [g^{ij}(t)n_i n_j]^{1/2} = [h^{ij}(t)n_i n_j]^{1/2}$$
(25.40)

for  $\iota \in \mathcal{I}_{\mathrm{B}}$ , where  $n_j = \nu_{\mathrm{T},j}(\iota)$ , cf. (24.5), and h is defined by (25.3).

**Lemma 25.11.** Let (M,g) be a canonical separable cosmological model manifold. Assume that there is an  $0 < \eta_{\text{sh},0} \leq 1$  such that (3.4) holds for all  $t \geq 0$ ; that there is an  $0 < \eta_{\text{sh},1} \in \mathbb{R}$  such that (3.5) holds with k = 1; and that there is a  $0 < C_0 \in \mathbb{R}$  such that  $|\bar{k}|_{\bar{g}} \leq C_0$ . Define  $\ell$  by (25.39). Then there is a constant  $0 < C \in \mathbb{R}$ , depending only on  $C_0$ , and  $\eta_{\text{sh},k}$ , k = 0, 1, such that

$$|\ell(\iota, t)| \le C \tag{25.41}$$

for all  $t \ge 0$  and all  $0 \ne \iota \in \mathcal{I}_{\mathrm{B}}$ . Assume, in addition to the above, that there is an  $0 < \eta_{\mathrm{sh},2} \in \mathbb{R}$ such that (3.5) holds with k = 2 and that there is a  $0 < C_1 \in \mathbb{R}$  such that  $|\mathcal{L}_U \bar{k}|_{\bar{g}} \le C_1$ . Then there is a constant  $0 < C \in \mathbb{R}$ , depending only on  $C_0$ ,  $C_1$ , and  $\eta_{\mathrm{sh},k}$ , k = 0, 1, 2, such that

$$|\hat{\ell}(\iota, t)| \le C \tag{25.42}$$

for all  $t \geq 0$  and all  $0 \neq \iota \in \mathcal{I}_B$ .

*Proof.* Due to Lemma 25.9, we know that (25.33) holds. Moreover, (25.7) yields  $h^{ij} = \bar{h}^{ij} - \chi^i \chi^j$ . Fixing  $n \in \mathbb{Z}^d$  and defining  $\mathfrak{g}_{\mathrm{T}}$  according to (25.40), it can thus be computed that

$$2\mathfrak{g}_{\mathrm{T}}\dot{\mathfrak{g}}_{\mathrm{T}} = -\bar{h}^{il}\bar{h}^{jm}\partial_t\bar{h}_{lm}n_in_j - 2\chi^i n_i \dot{\chi}^j n_j.$$

$$\tag{25.43}$$

Due to (25.27), it can be estimated that

$$|\chi^{i}n_{i}| \leq (\bar{h}_{ij}\chi^{i}\chi^{j})^{1/2}(\bar{h}^{ij}n_{i}n_{j})^{1/2} = |\chi|_{\bar{g}}(\bar{h}^{ij}n_{i}n_{j})^{1/2}.$$
(25.44)

Combining this estimate with a similar estimate for  $\dot{\chi}^j n_j$  yields

$$|\chi^{i} n_{i} \dot{\chi}^{j} n_{j}| \leq |\chi|_{\bar{g}} |\dot{\chi}|_{\bar{g}} \bar{h}^{ij} n_{i} n_{j} \leq \eta_{\mathrm{sh},0}^{-2} |\chi|_{\bar{g}} |\dot{\chi}|_{\bar{g}} \mathfrak{g}_{\mathrm{T}}^{2}, \qquad (25.45)$$

where we appealed to (25.33). Note that  $2\bar{k}_{T,lm} = \partial_t \bar{h}_{lm}$ , so that the first term on the right hand side of (25.43) can be estimated by

$$2|\bar{h}^{il}\bar{h}^{jm}\bar{k}_{\mathrm{T},lm}n_in_j| \le 2|\bar{k}_{\mathrm{T}}|_{\bar{h}}|v|_{\bar{h}}^2,$$

where  $v^i = \bar{h}^{il} n_l$  and we appealed to Lemma 25.6. Note that

$$|v|_{\bar{h}}^2 = \bar{h}_{ij}\bar{h}^{il}\bar{h}^{jm}n_ln_m = \bar{h}^{ij}n_in_j \le \eta_{\mathrm{sh},0}^{-2}\mathfrak{g}_{\mathrm{T}}^2,$$

where we appealed to (25.33). Summing up yields

$$\bar{h}^{il}\bar{h}^{jm}\partial_t h_{lm}n_i n_j | \le 2\eta_{\rm sh,0}^{-2} |\bar{k}_{\rm T}|_{\bar{h}}\mathfrak{g}_{\rm T}^2.$$
(25.46)

Combining (25.43), (25.45) and (25.46) yields

$$|\dot{\mathfrak{g}}_{\mathrm{T}}| \le \eta_{\mathrm{sh},0}^{-2} (|\bar{k}_{\mathrm{T}}|_{\bar{h}} + |\chi|_{\bar{g}} |\dot{\chi}|_{\bar{g}}) \mathfrak{g}_{\mathrm{T}}.$$
(25.47)

Keeping (25.20) in mind,

$$2\mathfrak{g}\mathfrak{g}| = \left| 2\mathfrak{g}_{\mathrm{T}}\mathfrak{g}_{\mathrm{T}} - 2\sum_{r=1}^{R} \frac{\dot{a}_{r}}{a_{r}} a_{r}^{-2} \nu_{r,i_{r}}^{2} \right| \\ \leq 2[\eta_{\mathrm{sh},0}^{-2}(|\bar{k}_{\mathrm{T}}|_{\bar{h}} + |\chi|_{\bar{g}} \cdot |\dot{\chi}|_{\bar{g}}) + |\bar{k}_{\mathrm{Rie}}|_{\bar{g}}]\mathfrak{g}^{2}.$$
(25.48)

Thus (25.41) holds.

Turning to the second derivative, note that differentiating (25.43) and carrying out estimates similar to the above yields

$$|2\dot{\mathfrak{g}}_{\mathrm{T}}^{2} + 2\mathfrak{g}_{\mathrm{T}}\ddot{\mathfrak{g}}_{\mathrm{T}}| \leq 2\eta_{\mathrm{sh},0}^{-2}(|(\mathcal{L}_{U}\bar{k})_{\mathrm{T}}|_{\bar{h}} + 4|\bar{k}_{\mathrm{T}}|_{\bar{h}}^{2} + |\chi|_{\bar{g}} \cdot |\ddot{\chi}|_{\bar{g}} + |\dot{\chi}|_{\bar{g}}^{2})\mathfrak{g}_{\mathrm{T}}^{2},$$
(25.49)

where we also appealed to (25.29). On the other hand,

$$2\dot{\mathfrak{g}}^{2} + 2\mathfrak{g}\ddot{\mathfrak{g}} = 2\dot{\mathfrak{g}}_{\mathrm{T}}^{2} + 2\mathfrak{g}_{\mathrm{T}}\ddot{\mathfrak{g}}_{\mathrm{T}} - 2\sum_{r=1}^{R} \frac{\ddot{a}_{r}}{a_{r}}a_{r}^{-2}\nu_{r,i_{r}}^{2} + 6\sum_{r=1}^{R} \left(\frac{\dot{a}_{r}}{a_{r}}\right)^{2}a_{r}^{-2}\nu_{r,i_{r}}^{2}.$$

The sum of the first two terms can, in absolute value, be bounded by the right hand side of (25.49). Keeping (25.20) and (25.21) in mind, the sum of the last two terms can be bounded in absolute value by

$$2\left(|(\mathcal{L}_U \bar{k})_{\text{Rie}}|_{\bar{g}} + 4|\bar{k}_{\text{Rie}}|_{\bar{g}}^2\right) \sum_{r=1}^R a_r^{-2} \nu_{r,i_r}^2.$$

Adding up yields

$$\begin{aligned} |\mathfrak{g}\ddot{\mathfrak{g}} + \dot{\mathfrak{g}}^{2}| &\leq \eta_{\mathrm{sh},0}^{-2} \left( |(\mathcal{L}_{U}\bar{k})_{\mathrm{T}}|_{\bar{h}} + 4|\bar{k}_{\mathrm{T}}|_{\bar{h}}^{2} + |\chi|_{\bar{g}} \cdot |\ddot{\chi}|_{\bar{g}} + |\dot{\chi}|_{\bar{g}}^{2} \right) \mathfrak{g}^{2} \\ &+ \left( |(\mathcal{L}_{U}\bar{k})_{\mathrm{Rie}}|_{\bar{g}} + 4|\bar{k}_{\mathrm{Rie}}|_{\bar{g}}^{2} \right) \mathfrak{g}^{2}. \end{aligned}$$

Combining this estimate with (25.48) yields the conclusion of the lemma.

#### 25.7.1 Consequences of bounds on the second fundamental form

In Chapter 4, in particular in Definition 4.1, we define the notion of silence in terms of a lower bound on the second fundamental form. However, in Chapter 10, in particular in Definition 10.1, we define the notion of silence in terms of an upper bound on  $\dot{\ell}$ . In the present subsection, we relate the two perspectives.

**Lemma 25.12.** Let (M, g) be a canonical separable cosmological model manifold. Assume, moreover, that there is an  $0 < \eta_{sh,0} \le 1$  such that (3.4) holds for all  $t \ge 0$  and a  $0 < \mu \in \mathbb{R}$  and a continuous non-negative function  $\mathfrak{e} \in L^1([0,\infty))$  such that

$$\bar{k} \ge (\mu - \mathfrak{e})\bar{g},\tag{25.50}$$

$$|\chi|_{\bar{q}} \cdot |\dot{\chi}|_{\bar{q}} \le \mathfrak{e} \tag{25.51}$$

for all  $t \geq 0$ . Then there is a continuous non-negative function  $\mathfrak{e}_{\mathfrak{g}} \in L^1([0,\infty))$  such that

$$\dot{\ell}(\iota, t) \le -\mu + \mathfrak{e}_{\mathfrak{g}}(t) \tag{25.52}$$

for all  $t \geq 0$  and all  $0 \neq \iota \in \mathcal{I}_B$ . Moreover,  $\mathfrak{e}_{\mathfrak{g}} := 2\eta_{sh.0}^{-2}\mathfrak{e}$ .

**Remark 25.13.** The statement that  $\bar{k} \ge \eta \bar{g}$  for all  $t \ge 0$  is equivalent to the statement that  $\bar{k}(\xi,\xi) \ge \eta \bar{g}(\xi,\xi)$  for all  $\xi \in T\bar{M}$  and all  $t \ge 0$ , where we think of  $\bar{g}$  and  $\bar{k}$  as being 2-tensor fields on  $\bar{M}$ .

**Remark 25.14.** If the estimate (25.50) is replaced with

$$\bar{k} \le (\mu_+ + \mathfrak{e}_+)\bar{g},$$

where  $0 < \mu_+ \in \mathbb{R}$  and  $\mathfrak{e}_+ \in L^1([0,\infty))$  is a continuous non-negative function, then the conclusion (25.52) is replaced by

$$\ell(\iota, t) \ge -\mu_{\text{low}} - \mathfrak{e}_{\text{low}}(t) \tag{25.53}$$

for all  $t \ge 0$  and all  $0 \ne \iota \in \mathcal{I}_{\mathrm{B}}$ , where  $\mathfrak{e}_{\mathrm{low}} \in L^1([0,\infty))$  is a continuous non-negative function (depending only on  $\mathfrak{e}, \mathfrak{e}_+$  and  $\eta_{\mathrm{sh},0}$ ) and  $0 < \mu_{\mathrm{low}} \in \mathbb{R}$  depends only on  $\mu_+$  and  $\eta_{\mathrm{sh},0}$ .

*Proof.* The first steps of the proof of Lemma 25.11 still apply. In particular, the estimates up to, and including, (25.45) still hold. On the other hand, appealing to (25.50) and the fact that (25.15) holds yields

$$\bar{h}^{il}\bar{h}^{jm}\partial_t\bar{h}_{lm}n_in_j \ge 2(\mu - \mathfrak{e})\bar{h}^{ij}n_in_j \ge 2(\mu - \eta_{\mathrm{sh},0}^{-2}\mathfrak{e})\mathfrak{g}_{\mathrm{T}}^2$$
(25.54)

for all  $t \ge 0$ ; in the last step, we also appealed to (25.33). Combining (25.43), (25.45), (25.51) and (25.54) yields

$$\dot{\mathfrak{g}}_{\mathrm{T}} \le -(\mu - 2\eta_{\mathrm{sh},0}^{-2}\mathfrak{e})\mathfrak{g}_{\mathrm{T}}$$

$$\tag{25.55}$$

for all  $t \ge 0$ . On the other hand, combining (25.15), (25.16) and (25.50) yields  $\dot{a}_r/a_r \ge \mu - \mathfrak{e}$  for all  $t \ge 0$ . Thus

$$2\mathfrak{g}\dot{\mathfrak{g}} = 2\mathfrak{g}_{\mathrm{T}}\dot{\mathfrak{g}}_{\mathrm{T}} - 2\sum_{r=1}^{R}\frac{\dot{a}_{r}}{a_{r}}a_{r}^{-2}\nu_{r,i_{r}}^{2} \leq -2(\mu - \mathfrak{e}_{\mathfrak{g}})\mathfrak{g}^{2}$$

for all  $t \ge 0$  and all  $0 \ne \iota \in \mathcal{I}_{\mathrm{B}}$ , where  $\mathfrak{e}_{\mathfrak{g}} := 2\eta_{\mathrm{sh},0}^{-2}\mathfrak{e}$ . The lemma follows.

It is also of interest to record the following criteria, ensuring that (10.10) holds.

**Lemma 25.15.** Let (M, g) be a canonical separable cosmological model manifold. Assume, moreover, that there is an  $0 < \eta_{sh,0} \le 1$  such that (3.4) holds for all  $t \ge 0$  and a constant  $0 < C_a \in \mathbb{R}$ and a continuous non-negative function  $\mathfrak{e} \in L^1([0,\infty))$  such that

$$|\bar{k}|_{\bar{g}} + |\chi|_{\bar{g}} \cdot |\dot{\chi}|_{\bar{g}} \le C_a + \mathfrak{e} \tag{25.56}$$

for all  $t \ge 0$ . Then there is a continuous non-negative function  $\mathfrak{e}_{bal} \in L^1([0,\infty))$  and a  $0 < C_{\ell} \in \mathbb{R}$  such that

$$|\ell(\iota, t)| \le C_{\ell} + \mathfrak{e}_{\text{bal}}(t) \tag{25.57}$$

for all  $t \geq 0$  and all  $0 \neq \iota \in \mathcal{I}_{B}$ .

**Remark 25.16.** In the statement of the lemma,

$$C_{\ell} = (1 + \eta_{\mathrm{sh},0}^{-2})C_a, \quad \mathfrak{e}_{\mathrm{bal}} = (1 + \eta_{\mathrm{sh},0}^{-2})\mathfrak{e}.$$

*Proof.* The statement follows from the assumptions and (25.48).

#### **25.7.2** Bounding $\sigma$ and X

Recall that  $\sigma$  and X are defined by (24.17). It is of interest to bound the norms of these functions (and their time derivatives) in terms of the norms of  $\chi$ ,  $\mathcal{X}$  (recall that  $\mathcal{X}$  is introduced in (3.6)) and their time derivatives.

**Lemma 25.17.** Consider (1.2). Assume the associated metric to be such that (M, g) is a canonical separable cosmological model manifold. Define  $\bar{h}$  by (25.6). Assume, moreover, that there is an  $0 < \eta_{sh,0} \leq 1$  such that (3.4) holds for all  $t \geq 0$ . Then

$$|\sigma(\iota, t)| \le \eta_{\mathrm{sh},0}^{-1} |\chi(t)|_{\bar{g}},\tag{25.58}$$

$$\|X(\iota, t)\| \le \eta_{\mathrm{sh}, 0}^{-1} |\mathcal{X}(t)|_{\bar{h}},\tag{25.59}$$

$$|\dot{\sigma}(\iota,t)| \le \eta_{\mathrm{sh},0}^{-1} |\dot{\chi}(t)|_{\bar{g}} + \eta_{\mathrm{sh},0}^{-1} |\chi(t)|_{\bar{g}} |\dot{\ell}(\iota,t)|, \qquad (25.60)$$

$$\|X(\iota, t)\| \le \eta_{\mathrm{sh},0}^{-1} |\mathcal{X}(t)|_{\bar{h}} + \eta_{\mathrm{sh},0}^{-1} |\mathcal{X}(t)|_{\bar{h}} |\ell(\iota, t)|$$
(25.61)

for all  $t \geq 0$  and all  $0 \neq \iota \in \mathcal{I}_{B}$ .

**Remark 25.18.** Recall that  $|\mathcal{X}(t)|_{\bar{h}}$  is defined by (3.7).

Proof. Estimate

$$|\sigma(\iota,t)| = \frac{|n_l \chi^l(t)|}{\mathfrak{g}(\iota,t)} \le \eta_{\mathrm{sh},0}^{-1} \frac{|\chi(t)|_{\bar{g}} \mathfrak{g}_{\mathrm{T}}(\iota,t)}{\mathfrak{g}(\iota,t)} \le \eta_{\mathrm{sh},0}^{-1} |\chi(t)|_{\bar{g}}$$

for all  $t \ge 0$  and all  $0 \ne \iota \in \mathcal{I}_B$ , where we appealed to (25.27) and (25.33). Thus (25.58) holds. Estimate

$$\begin{aligned} \|n_{l}X^{l}(t)\| \leq \sum_{l} |n_{l}| \cdot \|X^{l}(t)\| &= \sum_{l} n_{l} \left( \frac{|n_{l}|}{n_{l}} \|X^{l}(t)\| \right) \\ \leq (\bar{h}^{ij}(t)n_{i}n_{j})^{1/2} \left( \sum_{l,k} \bar{h}_{lk}(t) \frac{|n_{l}|}{n_{l}} \|X^{l}(t)\| \frac{|n_{k}|}{n_{k}} \|X^{k}(t)\| \right)^{1/2} \\ \leq \eta_{\mathrm{sh},0}^{-1} (h^{ij}(t)n_{i}n_{j})^{1/2} |\mathcal{X}(t)|_{\bar{h}} \leq \eta_{\mathrm{sh},0}^{-1} \mathfrak{g}_{\mathrm{T}}(\iota,t) |\mathcal{X}(t)|_{\bar{h}} \end{aligned}$$
(25.62)

for all  $t \ge 0$ , where  $|n_l|/n_l$  should be thought of as equalling 1 when  $n_l = 0$ . Thus

$$\|X(\iota,t)\| \le \frac{\|n_l X^l(t)\|}{\mathfrak{g}(\iota,t)} \le \eta_{\mathrm{sh},0}^{-1} |\mathcal{X}(t)|_{\bar{h}}$$

and (25.59) follows. The proofs of the remaining estimates are similar.

#### 25.7.3 Geometric criteria in the unbalanced setting

Let us return to the conditions stated in Sections 16.3–16.5, in particular in Lemma 16.8 and Assumptions 16.10 and 16.15.

**Lemma 25.19.** Consider (1.2). Assume the associated metric to be such that (M, g) is a canonical separable cosmological model manifold. Assume, moreover, that d = 1, that R = 0 and that there is an  $0 < \eta_{sh,0} \le 1$  such that (3.4) holds for all  $t \ge 0$ . Assume that there are constants  $K_{sh}, K_{\bar{k}}, K_{ode}, \bar{K}_X > 0; \beta_{sh} < 0; \beta_{ode}; and \beta_X \ge 0$  such that

$$|\chi(t)|_{\bar{q}} + |\dot{\chi}(t)|_{\bar{q}} \le K_{\rm sh} e^{\beta_{\rm sh} t},$$
 (25.63)

$$|\ddot{\chi}(t)|_{\bar{g}} + |k(t)|_{\bar{g}} + |\mathcal{L}_U k(t)|_{\bar{g}} \le K_{\bar{k}},$$
(25.64)

$$\|\zeta(t)\|^{1/2} + \|\dot{\zeta}(t)\|^{1/2} + \|\alpha(t)\| + \|\dot{\alpha}(t)\| \le K_{\text{ode}}e^{\beta_{\text{ode}}t},\tag{25.65}$$

$$|\mathcal{X}(t)|_{\bar{h}} + |\dot{\mathcal{X}}(t)|_{\bar{h}} \le \bar{K}_X e^{\beta_X t} \tag{25.66}$$

for all  $t \ge 0$ . Assume, finally, that  $\beta_X - \beta_{ode} > 0$ . Then the assumptions of Lemma 16.8 are satisfied.

Assume, in addition to the above, that  $\beta_X > 0$ . Let  $e_1$  be the unit vector field which is a positive multiple of  $\partial_1$ , define  $\mathcal{Y}^1$  to be the matrix valued function such that  $\mathcal{Y}^1 e_1 = \mathcal{X}$  and assume that there is a matrix  $\mathcal{Y}^1_{\infty} \in \mathbf{M}_m(\mathbb{C})$  and constants  $\bar{\eta}_X, L_X > 0$  such that

$$\|e^{-\beta_X t} \mathcal{Y}^1(t) - \mathcal{Y}^1_\infty\| \le L_X e^{-\bar{\eta}_X t} \tag{25.67}$$

for all  $t \ge 0$ . Then, defining  $Y^1$  by (2.4), there is a constant  $K_X$  such that

$$\|e^{-\beta_X t} Y^1(t) - \mathcal{Y}^1_{\infty}\| \le K_X e^{-\eta_X t}$$
(25.68)

for all  $t \ge 0$ , where  $K_X$  only depends on  $\eta_{\text{sh},0}$ ,  $K_{\text{sh}}$ ,  $\bar{K}_X$ ,  $K_{\bar{k}}$  and  $L_X$ . Moreover,  $\eta_X := \min\{-2\beta_{\text{sh}}, \bar{\eta}_X\}$ . In particular, Assumptions 16.15 are satisfied with  $Y^1_{\infty} = \mathcal{Y}^1_{\infty}$ .

**Remark 25.20.** If the conditions of the lemma are satisfied and  $\mathcal{Y}^1_{\infty}$  has an eigenvalue with a non-zero real part, then Assumptions 2.1 and 2.3 are satisfied with  $Y^1_{\infty} = \mathcal{Y}^1_{\infty}$ ; this is a consequence of the fact that the assumptions of Lemma 16.8 are satisfied and the fact that Assumptions 16.15 hold.

386

*Proof.* Due to (25.63), (25.64) and the fact that there is an  $0 < \eta_{sh,0} \le 1$  such that (3.4) holds for all  $t \ge 0$ , Lemma 25.11 is applicable and yields the conclusion that there is a constant  $C_{\mathfrak{l}}$  such that

$$|\hat{\mathfrak{l}}(t)| + |\hat{\mathfrak{l}}(t)| \le C_{\mathfrak{l}} \tag{25.69}$$

for all  $t \ge 0$ ; note that  $\mathfrak{l}(t) = \ell(\iota, t)$  in the present context. Moreover,  $C_{\mathfrak{l}}$  only depends on  $K_{\bar{k}}$ ,  $\eta_{\mathrm{sh},0}$ and  $K_{\mathrm{sh}}$ . Combining (25.69) with the assumptions of the lemma as well as with Lemma 25.17 yields the conclusion that there are constants  $C_{\mathrm{sh}}$  and  $C_X$  such that

$$|\varsigma(t)| + |\dot{\varsigma}(t)| \le C_{\rm sh} e^{\beta_{\rm sh} t}, \quad ||Y^1(t)|| + ||\dot{Y}^1(t)|| \le C_X e^{\beta_X t}$$
(25.70)

for all  $t \ge 0$ ; note that  $||X(\iota, t)|| = ||Y^1(t)||$ ,  $|\sigma(\iota, t)| = |\varsigma(t)|$  etc. in the present context. In these estimates,  $C_{\rm sh}$  only depends on  $K_{\rm sh}$ ,  $\eta_{\rm sh,0}$  and  $K_{\bar{k}}$ ; and  $C_X$  only depends on  $\bar{K}_X$ ,  $K_{\rm sh}$ ,  $\eta_{\rm sh,0}$  and  $K_{\bar{k}}$ . Returning to the statement of Proposition 16.8, note that by defining  $\eta_{\rm ode} := \beta_X - \beta_{\rm ode}$ , it is clear that  $\eta_{\rm ode} > 0$ . Thus (16.18) holds. Due to the fact that the left hand sides of (16.19) and (16.20) are bounded, it is clear that we can find a constant  $\mathfrak{b} > 0$  such that (16.19) and (16.20) hold. Due to (25.70), it is clear that (16.21) and (16.22) hold. Finally, (16.23) and (16.24) follow from (25.65). Thus the assumptions of Proposition 16.8 are satisfied.

Next, it is of interest to compare  $\mathcal{Y}^1$  with  $Y^1$  introduced in (2.4). By definition,  $e_1 = (g_{11})^{-1/2} \partial_1$ . For this reason,

$$Y^{1} = \frac{X^{1}}{(g^{11})^{1/2}} = \frac{\mathcal{Y}^{1}}{(g_{11}g^{11})^{1/2}}.$$
(25.71)

Due to (25.7), we know that

$$g_{11}g^{11} = g_{11}(g_{11})^{-1} - g_{11}\chi^1\chi^1 = 1 - g_{11}\chi^1\chi^1.$$

Combining this observation with (3.4) yields  $\eta_{\text{sh},0}^2 \leq g_{11}g^{11} \leq 1$ . In particular,

$$\|\mathcal{Y}^{1}(t)\| \le \|Y^{1}(t)\| \le C_{X} e^{\beta_{X} t} \tag{25.72}$$

for all  $t \ge 0$ , where we appealed to (25.70). Moreover,

$$\left| 1 - \frac{1}{(g_{11}g^{11})^{1/2}} \right| = \frac{|g^{11}g_{11} - 1|}{(g_{11}g^{11})^{1/2}[1 + (g_{11}g^{11})^{1/2}]} \\ \leq \eta_{\mathrm{sh},0}^{-1} K_{\mathrm{sh}}^2 e^{2\beta_{\mathrm{sh}}t}$$

for all  $t \ge 0$ . Combining this estimate with (25.71) and (25.72) yields

$$||Y^{1}(t) - \mathcal{Y}^{1}(t)|| \leq \eta_{\mathrm{sh},0}^{-1} K_{\mathrm{sh}}^{2} C_{X} e^{(\beta_{X} + 2\beta_{\mathrm{sh}})t}$$

for all  $t \ge 0$ . Combining this estimate with (25.67) yields (25.68). The lemma follows.

**Lemma 25.21.** Let (M, g) be a canonical separable cosmological model manifold. Assume, moreover, that d = 1, that R = 0 and that there is an  $0 < \eta_{sh,0} \le 1$  such that (3.4) holds for all  $t \ge 0$ . Assume, finally, that there are  $\bar{\beta}_j \in \mathbb{R}$ , j = 1, 2, and a continuous function  $0 \le \mathfrak{e} \in L^1([0, \infty))$ such that

$$(\bar{\beta}_2 - \mathfrak{e})\bar{g} \le \bar{k} \le (\bar{\beta}_1 + \mathfrak{e})\bar{g} \tag{25.73}$$

for all  $t \ge 0$ . Then there are constants  $C_i > 0$ , i = 1, 2, depending only on  $g_{11}(0)$ ,  $C_{\mathfrak{e}} := \|\mathfrak{e}\|_1$  and  $\eta_{\mathrm{sh},0}$  such that

$$C_1 e^{-2\bar{\beta}_1 t} \le g^{11}(t) \le C_2 e^{-2\bar{\beta}_2 t}$$

for all  $t \geq 0$ .

**Remark 25.22.** If (1.2) is such that the the associated Lorentz manifold satisfies the assumptions of Lemma 25.21, then Assumption 16.10 holds with  $\beta_i = -\bar{\beta}_i$ .

*Proof.* Note that (25.73) is equivalent to

$$\bar{\beta}_2 - \mathfrak{e} \le \partial_t \ln(g_{11})^{1/2} \le \bar{\beta}_1 + \mathfrak{e}.$$

Integrating this estimate yields

$$g_{11}(0)\exp(2\bar{\beta}_2 t - 2C_{\mathfrak{e}}) \le g_{11}(t) \le g_{11}(0)\exp(2\bar{\beta}_1 t + 2C_{\mathfrak{e}})$$
(25.74)

for all  $t \ge 0$ , where  $C_{\mathfrak{e}} := \|\mathfrak{e}\|_1$ . Note that  $\eta^2_{\mathrm{sh},0} \le g_{11}(t)g^{11}(t) \le 1$  for all  $t \ge 0$ . This is a consequence of (25.7) and (3.4); cf. the proof of Lemma 25.19. Combining this estimate with (25.74) yields

$$\frac{\eta_{\mathrm{sh},0}^2}{g_{11}(0)}\exp(-2\bar{\beta}_1t - 2C_{\mathfrak{e}}) \le g^{11}(t) \le \frac{1}{g_{11}(0)}\exp(-2\bar{\beta}_2t + 2C_{\mathfrak{e}})$$

for all  $t \ge 0$ . The lemma follows.

#### 25.8 Implications concerning the causal structure

In Definition 4.1, we introduce the notion of  $C^1$ -silence. The purpose of the present section is to justify the use of the word silent.

**Lemma 25.23.** Let (M,g) be a canonical separable cosmological model manifold. Assume that there is a  $0 < \mu \in \mathbb{R}$  and a continuous non-negative function  $\mathfrak{e} \in L^1([0,\infty))$  such that

$$\bar{k} \ge (\mu - \mathfrak{e})\bar{g} \tag{25.75}$$

for all  $t \ge 0$ . Let  $\bar{g}_{ref}$  be a fixed Riemannian metric on  $\bar{M}$ . Then there is a constant  $0 < C \in \mathbb{R}$ , depending only on  $\|\mathfrak{e}\|_1$ ,  $a_r(0)$  (r = 1, ..., R),  $g_{ij}(0)$  (i, j = 1, ..., d) and the reference metric  $\bar{g}_{ref}$ such that if  $\gamma : [0, \infty) \to M$  is a causal curve in (M, g) with  $\gamma(t) = [\bar{\gamma}(t), t]$ , then

$$|\dot{\bar{\gamma}}(t)|_{\bar{g}_{\text{ref}}} \le C e^{-\mu t} \tag{25.76}$$

for all  $t \geq 0$ .

**Remark 25.24.** Due to (25.76), it is clear that  $\bar{\gamma}(t)$  converges exponentially to a point, say  $\bar{p}[\gamma] \in \bar{M}$ , as  $t \to \infty$ . Moreover, if  $\gamma_i : [0, \infty) \to M$ , i = 1, 2, are two future oriented causal curves such that  $\gamma_i(t) = [\bar{\gamma}_i(t), t]$  and  $\bar{p}[\gamma_1] \neq \bar{p}[\gamma_2]$ , then the observers  $\gamma_1$  and  $\gamma_2$  sooner or later lose the ability to communicate. In fact, there are  $t_i \in [0, \infty)$ , i = 1, 2, such that

$$J^+[\gamma_1(t_1)] \cap J^+[\gamma_2(t_2)] = \emptyset.$$

In this sense, there is silence asymptotically.

*Proof.* Given  $\xi \in \mathbb{R}^d$ , let

$$\mathfrak{N}(t,\xi) := \bar{h}_{ij}(t)\xi^i\xi^j.$$

Then

$$\dot{\mathfrak{N}}(t,\xi) = \partial_t \bar{h}_{ij} \xi^i \xi^j = 2\bar{k}_{\mathrm{T},ij} \xi^i \xi^j \ge 2[\mu - \mathfrak{e}(t)]\mathfrak{N}(t,\xi)$$

for all  $t \ge 0$ , where we appealed to (25.15) and (25.75). Integrating this estimate yields

$$\bar{h}_{ij}(t)\xi^i\xi^j \ge C|\xi|^2 e^{2\mu t}$$
(25.77)

for all  $t \ge 0$  and all  $\xi \in \mathbb{R}^d$ , where C > 0 only depends on  $\|\mathfrak{e}\|_1$  and  $g_{ij}(0)$   $(i, j = 1, \ldots, d)$ . Similarly, (25.15), (25.16) and (25.75) yield  $a_r(t) \ge Ce^{\mu t}$  for all  $t \ge 0$ , where the constant C > 0 only depends on  $\|\mathfrak{e}\|_1$  and  $a_r(0)$ . Let  $\gamma : [0, \infty) \to M$  be a causal curve in (M, g) with

$$\gamma(t) = [\bar{\gamma}(t), t] = [\bar{\gamma}_{\mathrm{T}}(t), \bar{\gamma}_{1}(t), \dots, \bar{\gamma}_{R}(t), t],$$
where  $\bar{\gamma}_{\mathrm{T}}$  takes its values in  $\mathbb{T}^d$  and  $\bar{\gamma}_r$  takes its values in  $M_r$ ,  $r = 1, \ldots, R$ . Then

$$1 + \bar{h}_{ij}\chi^{i}\chi^{j} + 2\bar{h}_{ij}\chi^{i}\dot{\bar{\gamma}}_{T}^{j} + \bar{h}_{ij}\dot{\bar{\gamma}}_{T}^{i}\dot{\bar{\gamma}}_{T}^{j} + \sum_{r=1}^{R} a_{r}^{2}|\dot{\bar{\gamma}}_{r}|_{g_{r}}^{2} \le 0$$

for all  $t \ge 0$ . Thus

\_

$$\begin{split} |\dot{\gamma}_{\mathrm{T}}|_{\bar{h}}^{2} + \sum_{r=1}^{R} a_{r}^{2} |\dot{\gamma}_{r}|_{g_{r}}^{2} \leq 1 + 2|\chi|_{\bar{g}} |\dot{\gamma}_{\mathrm{T}}|_{\bar{h}} - |\chi|_{\bar{g}}^{2} \leq 1 + |\chi|_{\bar{g}}^{2} + \frac{1}{2} |\dot{\gamma}_{\mathrm{T}}|_{\bar{h}}^{2} \\ \leq 2 + \frac{1}{2} |\dot{\gamma}_{\mathrm{T}}|_{\bar{h}}^{2}, \end{split}$$

where we used the fact that  $|\chi|_{\bar{g}} \leq 1$ ; recall that (25.5) holds and that N = 1. In particular,

$$|\dot{\bar{\gamma}}_{\mathrm{T}}|_{\bar{h}}^{2} + \sum_{r=1}^{R} a_{r}^{2} |\dot{\bar{\gamma}}_{r}|_{g_{r}}^{2} \le 4$$

for all  $t \ge 0$ . Due to (25.77) and the analogous estimate for the  $a_r$ , we conclude that there is a  $0 < C \in \mathbb{R}$ , depending only on  $\|\mathfrak{e}\|_1$ ,  $a_r(0)$  (r = 1, ..., R), and  $g_{ij}(0)$  (i, j = 1, ..., d), such that

$$|\dot{\bar{\gamma}}_{\mathrm{T}}(t)|^{2} + \sum_{r=1}^{R} |\dot{\bar{\gamma}}_{r}(t)|^{2}_{g_{r}} \leq Ce^{-2\mu t}$$

for all  $t \ge 0$ . The lemma follows.

#### CHAPTER 25. GEOMETRIC PRELIMINARIES

### Chapter 26

# Relating the analytic and geometric conditions

In Part I, we formulate the conditions geometrically. However, in Parts II–VI, the conditions are formulated analytically. The purpose of the present chapter is to relate the two perspectives.

#### 26.1 Bounds on the coefficients in the silent setting

Let us begin by deriving conclusions concerning the coefficients from the conditions appearing in Definition 4.1.

**Lemma 26.1.** Given that the assumptions of Lemma 25.12 are fulfilled, the norm of the matrix with components  $g^{ij}(t)$ , i, j = 1, ..., d, is bounded by  $Ce^{-2\mu t}$  for all  $t \ge 0$ , where C only depends on  $\eta_{sh,0}$ ,  $\|\mathbf{e}\|_1$  and  $g^{ij}(0)$  (i, j = 1, ..., d). Moreover,  $a_r^{-2}(t) \le Ce^{-2\mu t}$  for all  $t \ge 0$ , where C only depends on  $\|\mathbf{e}\|_1$  and  $a_r(0)$ . If, in addition,  $\mathcal{X}$  is  $C^0$ -future bounded, then  $\|X^j(t)\| \le Ce^{-\mu t}$  for all  $t \ge 0$  and all  $j \in \{1, ..., d\}$ , where C only depends on  $\|\mathbf{e}\|_1$ ,  $g_{ij}(0)$  (i, j = 1, ..., d) and the future bound on  $|\mathcal{X}|_{\bar{h}}$ . Finally,  $|\chi(t)| \le Ce^{-\mu t}$  for all  $t \ge 0$ , where C only depends on  $\|\mathbf{e}\|_1$  and  $g_{ij}(0)$  (i, j = 1, ..., d) (note that  $|\chi(t)|$  denotes the Euclidean norm of the vector with components  $\chi^i(t)$ ).

*Proof.* Consider the estimate (25.55). If we would replace  $n \in \mathbb{Z}^d$  in the definition of  $\mathfrak{g}_T$  by  $\xi \in \mathbb{R}^d$ , the result would be the same. In particular, there is thus a constant  $0 < C \in \mathbb{R}$ , depending only on  $\eta_{\text{sh},0}$  and  $\|\mathfrak{e}\|_1$ , such that

$$g^{ij}(t)\xi_i\xi_j \le Cg^{ij}(0)\xi_i\xi_j e^{-2\mu t}$$
(26.1)

for all  $t \ge 0$  and all  $\xi \in \mathbb{R}^d$ . Thus the first statement of the lemma holds. Note that the conditions of Lemma 25.23 are fulfilled. In what follows, we therefore use the conclusions of this lemma freely. The second statement follows from lower bound on  $a_r(t)$  stated in connection with (25.77). The statement concerning  $\mathcal{X}$  follows from (25.77) and the assumption that  $\mathcal{X}$  is  $C^0$ -future bounded. The justification of the bound on  $|\chi(t)|$  is similar.

#### 26.2 Conditions ensuring weak transparency

Let us now, starting with the conditions formulated in Section 5.1, derive conclusions concerning the coefficients of the equation.

**Lemma 26.2.** Let (M,g) be a canonical separable cosmological model manifold. Assume that there is a division of the cotangent space into transparent and silent subsets, as described in

Definition 5.1. Assume, moreover, that Definitions 5.2 and 5.5 are fulfilled. Given  $\iota \in \mathcal{I}_B$ , let  $\iota_{ts}$  and  $\iota_{sil}$  be given by Definition 5.8. Then there are constants  $C_A$  and  $C_B$  such that

$$|\dot{\ell}(\iota_{\rm ts}, t)| \le C_A e^{-\eta_{\rm ts} t},\tag{26.2}$$

$$|\sigma(\iota_{\rm ts}, t)| \le C_B e^{-\eta_{\rm ts} t} \tag{26.3}$$

for all  $t \geq 0$  and all  $\iota \in \mathcal{I}_{B}$  such that  $\iota_{ts} \neq 0$ . Moreover,

$$\dot{\ell}(\iota_{\rm sil}, t) \le -\beta_{\rm sil} + \eta_{\rm sh,0}^{-2} \mathfrak{e}_{\rm sil}(t) + \eta_{\rm sh,0}^{-2} \mathfrak{e}(t)$$
(26.4)

for all  $t \geq 0$  and all  $\iota \in \mathcal{I}_{B}$  such that  $\iota_{sil} \neq 0$ .

Concerning the metric coefficients, there are constants  $g_{\infty}^{j_k j_l}$ ,  $k, l \in \{1, \ldots, d_{ts}\}$ , and  $q_{\infty, r_j}$ ,  $j = 1, \ldots, R_{ts}$ , such that

$$|g^{j_k j_l}(t) - g^{j_k j_l}_{\infty}| \le C e^{-\eta_{ts} t}, \quad |a_{r_j}^{-2}(t) - q_{\infty, r_j}| \le C e^{-\eta_{ts} t}$$
(26.5)

for all  $t \ge 0$ , where C only depends on  $C_A$ ,  $\eta_{ts}$ ,  $g^{ij}(0)$  and  $a_r(0)$ . Moreover,  $g^{j_k j_l}_{\infty}$  are the components of a positive definite matrix, and  $q_{\infty,r_j} > 0$ . In addition,

$$|g^{\bar{j}_k\bar{j}_l}(t)| \le Ce^{-2\beta_{\rm sil}t}, \quad a_{\bar{r}_j}^{-2}(t) \le Ce^{-2\beta_{\rm sil}t}$$
(26.6)

for all  $t \ge 0$ ,  $k, l \in \{1, \ldots, d_{sil}\}$  and  $j = 1, \ldots, R_{sil}$ , where C only depends on  $g^{ij}(0)$ ,  $a_r(0)$ ,  $\eta_{sh,0}$ ,  $c_{sil} := \|\mathbf{e}_{sil}\|_1$  and  $c_{\mathbf{e}} := \|\mathbf{e}\|$ . Moreover,

$$|g^{j_k\bar{j}_l}(t)| \le Ce^{-\beta_{\rm sil}t} \tag{26.7}$$

for all  $t \ge 0$ , all  $k \in \{1, \ldots, d_{ts}\}$  and all  $l \in \{1, \ldots, d_{sil}\}$ , where C only depends on  $C_A$ ,  $\eta_{ts}$ ,  $g^{ij}(0)$ ,  $a_r(0)$ ,  $\eta_{sh,0}$ ,  $c_{sil}$  and  $c_{\mathfrak{e}}$ . Finally,

$$|g^{0j_k}(t)| \le C_1 e^{-\eta_{\rm ts} t}, \quad |g^{0\bar{j}_l}(t)| \le C_2 e^{-\beta_{\rm sil} t}$$
(26.8)

for all  $t \ge 0$ , all  $k \in \{1, \ldots, d_{ts}\}$  and all  $l \in \{1, \ldots, d_{sil}\}$ , where  $C_1$  only depends on  $C_A$ ,  $C_B$ ,  $\eta_{ts}$  and  $g^{ij}(0)$  and  $C_2$  only depends on  $g^{ij}(0)$ ,  $\eta_{sh,0}$ ,  $c_{sil}$  and  $c_{\mathfrak{e}}$ .

**Remark 26.3.** The constant  $C_A$  appearing in (26.2) only depends on  $C_a$ ,  $C_b$  and  $\eta_{\text{sh},k}$ , k = 0, 1. The constant  $C_B$  appearing in (26.3) only depends on  $C_b$  and  $\eta_{\text{sh},0}$ .

**Remark 26.4.** If, in addition to the assumptions of the lemma,  $\mathcal{X}$  is  $C^0$ -future bounded, then

$$\|X^{j_k}(t)\| \le Ce^{-\beta_{\rm sil}t}$$

for all  $t \ge 0$  and all  $k \in \{1, \ldots, d_{sil}\}$ , where C only depends on  $g^{ij}(0)$ ,  $C_0$ ,  $\eta_{sh,0}$ ,  $c_{sil}$  and  $c_{\mathfrak{e}}$ . The reason for this is that if  $n \in \mathbb{Z}^d$  corresponds to an element  $\iota \in \mathcal{I}_B$  such that  $\iota = \iota_{sil}$ , then

$$\|n_j X^j(t)\| \le \eta_{\mathrm{sh},0}^{-1} |\mathcal{X}(t)|_{\bar{h}} \mathfrak{g}_{\mathrm{T}}(\iota_{\mathrm{sil}},t) \le C e^{-\beta_{\mathrm{sil}} t} \mathfrak{g}_{\mathrm{T}}(\iota_{\mathrm{sil}},0)$$

for all  $t \ge 0$ , where C only depends on  $C_0$ ,  $\eta_{\text{sh},0}$ ,  $c_{\text{sil}}$  and  $c_{\mathfrak{e}}$ . Moreover, we appealed to (25.27) in the first step and to (26.21) below in the second step.

#### Proof. The transparent directions. Compute

$$2\mathfrak{g}(\iota_{\rm ts},t)\dot{\mathfrak{g}}(\iota_{\rm ts},t) = 2\mathfrak{g}_{\rm T}(\iota_{\rm ts},t)\dot{\mathfrak{g}}_{\rm T}(\iota_{\rm ts},t) - 2\sum_{j=1}^{R_{\rm ts}}\frac{\dot{a}_{r_j}(t)}{a_{r_j}(t)}a_{r_j}^{-2}(t)\nu_{r_j,i_{r_j}}^2(\iota_{\rm ts}).$$
(26.9)

Let us begin by considering the first term on the right hand side. It can be computed as in (25.43). Consider the first term on the right hand side of (25.43). Since  $\iota = \iota_{ts}$  in the case of interest here,

 $v := n_j dx^j \in T^*_{ts} \overline{M}$ . Moreover,  $\overline{h}^{il} \overline{h}^{jp} \partial_t \overline{h}_{lp}$  is the *ij*'th component of  $2\overline{k}^{\sharp}$ ,  $i, j \in \{1, \ldots, d\}$ . Combining these observations with (5.1) yields

$$|\bar{h}^{il}\bar{h}^{jp}\partial_t\bar{h}_{lp}n_in_j| = 2|\bar{k}^{\sharp}(v,v)| \le 2C_a e^{-\eta_{\rm ts}t}|\bar{h}^{ij}n_in_j| \le 2\eta_{\rm sh,0}^{-2}C_a e^{-\eta_{\rm ts}t}\mathfrak{g}_{\rm T}^2 \tag{26.10}$$

for all  $t \ge 0$ , where we also appealed to (25.33). Consider the second term on the right hand side of (25.43). Note that

$$|\chi^{i} n_{i}| = |v(\chi)| \le C_{b} e^{-\eta_{ts} t} |v|_{\bar{g}} \le \eta_{sh,0}^{-1} C_{b} e^{-\eta_{ts} t} \mathfrak{g}_{T}$$
(26.11)

for all  $t \ge 0$ , where we appealed to (25.33) and (5.6). Since  $\chi$  is C<sup>1</sup>-bounded, we thus obtain

$$|\chi^{i} n_{i} \dot{\chi}^{j} n_{j}| \leq C_{b} \eta_{\mathrm{sh},0}^{-2} \eta_{\mathrm{sh},1} e^{-\eta_{\mathrm{ts}} t} \mathfrak{g}_{\mathrm{T}}^{2}.$$
(26.12)

Combining (26.10) and (26.12) yields

$$2|\mathfrak{g}_{\rm T}(\iota_{\rm ts},t)\dot{\mathfrak{g}}_{\rm T}(\iota_{\rm ts},t)| \le 2\eta_{\rm sh,0}^{-2}(C_a + \eta_{\rm sh,1}C_b)e^{-\eta_{\rm ts}t}\mathfrak{g}_{\rm T}^2(\iota_{\rm ts},t)$$
(26.13)

for all  $t \ge 0$ . Let us now turn to the second term on the right hand side of (26.9). Let  $j \in \{1, \ldots, R_{ts}\}, t \in [0, \infty)$  and  $\xi$  be a unit cotangent vector (with respect to  $\bar{g}$ ) such that  $\xi^{\sharp}$  is tangent to the  $M_{r_i}$ -factor in  $\bar{M}$ . Then (5.1) yields

$$\left|\frac{\dot{a}_{r_j}(t)}{a_{r_j}(t)}\right| \le C_a e^{-\eta_{\rm ts} t};\tag{26.14}$$

note that this estimate holds for all  $t \ge 0$  and all  $j \in \{1, \ldots, R_{ts}\}$ . Thus the second term on the right hand side of (26.9) can be estimated by

$$2\left|\sum_{j=1}^{R_{\rm ts}} \frac{\dot{a}_{r_j}(t)}{a_{r_j}(t)} a_{r_j}^{-2}(t) \nu_{r_j, i_{r_j}}^2(\iota_{\rm ts})\right| \le 2C_a e^{-\eta_{\rm ts}t} \sum_{j=1}^{R_{\rm ts}} a_{r_j}^{-2}(t) \nu_{r_j, i_{r_j}}^2(\iota_{\rm ts})$$
(26.15)

for all  $t \ge 0$ . Combining (26.9), (26.13) and (26.15) yields (26.2) for all  $t \ge 0$ . Note also that (26.11) implies that (26.3) holds for all  $t \ge 0$ .

The silent directions. Next, let us consider (26.9) in the case that  $\iota_{ts}$  is replaced by  $\iota_{sil}$ . Since  $\iota = \iota_{sil}$  in the case of interest here,  $v := n_i dx^i \in T^*_{sil} \overline{M}$ . The first term on the right hand side of (26.9) can be expressed as in (25.43). Consider the first term on the right hand side of (25.43). Due to (5.2) this term can be estimated by

$$-\bar{h}^{il}\bar{h}^{jp}\partial_t\bar{h}_{lp}n_in_j = -2\bar{k}^{\sharp}(v,v) \leq -2(\beta_{\rm sil} - \mathfrak{e}_{\rm sil})\bar{h}^{ij}n_in_j \\ \leq -2(\beta_{\rm sil} - \eta_{\rm sh,0}^{-2}\mathfrak{e}_{\rm sil})\mathfrak{g}_{\rm T}^2,$$
(26.16)

where we appealed to (25.15) in the first step and (25.33) in the last step. Consider the second term on the right of (25.43). It can be estimated by

$$2\eta_{\mathrm{sh},0}^{-2}|\chi|_{\bar{g}}|\dot{\chi}|_{\bar{g}}\mathfrak{g}_{\mathrm{T}}^{2} \leq 2\eta_{\mathrm{sh},0}^{-2}\mathfrak{e}\mathfrak{g}_{\mathrm{T}}^{2},\tag{26.17}$$

where we appealed to (5.5). Finally, due to (25.15) and (5.2),

$$-2\sum_{j=1}^{R_{\rm sil}} \frac{\dot{a}_{\bar{r}_j}(t)}{a_{\bar{r}_j}(t)} a_{\bar{r}_j}^{-2}(t) \nu_{\bar{r}_j, i_{\bar{r}_j}}^2(\iota_{\rm sil}) \leq -2[\beta_{\rm sil} - \mathfrak{e}_{\rm sil}(t)] \sum_{j=1}^{R_{\rm sil}} a_{\bar{r}_j}^{-2}(t) \nu_{\bar{r}_j, i_{\bar{r}_j}}^2(\iota_{\rm sil}).$$
(26.18)

Combining (26.16), (26.17) and (26.18) yields (26.4) for all  $t \ge 0$ .

The coefficients of the equation. Next, let us turn to the coefficients  $g^{ij}$  and  $a_r$ . Note, first of all, that the estimate (26.2) implies that  $\mathfrak{g}(\iota_{ts}, t)$  converges exponentially to a limit, say  $\mathfrak{g}_{\infty}(\iota_{ts})$ , and that

$$\left|\frac{\mathfrak{g}^{2}(\iota_{\mathrm{ts}},t)}{\mathfrak{g}^{2}_{\infty}(\iota_{\mathrm{ts}})} - 1\right| + \left|\frac{\mathfrak{g}^{2}_{\infty}(\iota_{\mathrm{ts}})}{\mathfrak{g}^{2}(\iota_{\mathrm{ts}},t)} - 1\right| \le Ce^{-\eta_{\mathrm{ts}}t}$$
(26.19)

for all  $t \ge 0$  and all  $\iota \in \mathcal{I}_B$  such that  $\iota_{ts} \ne 0$ , where C only depends on  $C_A$  and  $\eta_{ts}$ . This estimate implies the existence of a constant  $C \ge 1$ , depending only on  $C_A$  and  $\eta_{ts}$ , such that

$$C^{-1}\mathfrak{g}^2(\iota_{\rm ts},0) \le \mathfrak{g}^2_{\infty}(\iota_{\rm ts}) \le C\mathfrak{g}^2(\iota_{\rm ts},0) \tag{26.20}$$

for all  $\iota \in \mathcal{I}_{B}$ . Due to this estimate,  $\mathfrak{g}_{\infty}^{2}(\iota_{ts})$ , considered as a quadratic form, is positive definite. Combining (26.19) and (26.20) yields

$$|\mathfrak{g}^{2}(\iota_{\mathrm{ts}},t) - \mathfrak{g}_{\infty}^{2}(\iota_{\mathrm{ts}})| \leq Ce^{-\eta_{\mathrm{ts}}t}\mathfrak{g}^{2}(\iota_{\mathrm{ts}},0)$$

for all  $t \ge 0$  and all  $\iota \in \mathcal{I}_{\mathrm{B}}$ , where C only depends on  $C_A$  and  $\eta_{\mathrm{ts}}$ . This estimate implies that there are  $g_{\infty}^{j_k j_l}$ ,  $k, l \in \{1, \ldots, d_{\mathrm{ts}}\}$ , and  $q_{\infty, r_j}$ ,  $j = 1, \ldots, R_{\mathrm{ts}}$ , such that (26.5) holds for all  $t \ge 0$ , where C only depends on  $C_A$ ,  $\eta_{\mathrm{ts}}$ ,  $g^{ij}(0)$  and  $a_r(0)$ . Note also that

$$g_{\infty}^{2}(\iota_{\rm ts}) = g_{\infty}^{j_{k}j_{l}} n_{j_{k}} n_{j_{l}} + \sum_{j=1}^{R_{\rm ts}} q_{\infty,r_{j}} \nu_{r_{j},i_{r_{j}}}^{2}(\iota_{\rm ts}).$$

Since the left hand side of this equality is a positive definite quadratic form,  $g_{\infty}^{j_k j_l}$  are the components of a positive definite matrix, and  $q_{\infty,r_j} > 0$ .

Turning to the silent directions, note that (26.4) yields

$$\mathfrak{g}^2(\iota_{\rm sil}, t) \le C e^{-2\beta_{\rm sil}t} \mathfrak{g}^2(\iota_{\rm sil}, 0) \tag{26.21}$$

for all  $t \ge 0$ , where C only depends on  $\eta_{\text{sh},0}$ ,  $c_{\text{sil}} := \|\mathbf{e}_{\text{sil}}\|_1$  and  $c_{\mathbf{e}} := \|\mathbf{e}\|$ . Thus (26.6) holds for all  $t \ge 0, k, l \in \{1, \ldots, d_{\text{sil}}\}$  and  $j = 1, \ldots, R_{\text{sil}}$ , where C only depends on  $g^{ij}(0), a_r(0), \eta_{\text{sh},0}, c_{\text{sil}}$  and  $c_{\mathbf{e}}$ . Next, note that the Cauchy-Schwarz inequality yields

$$|g^{j_k \bar{j}_l}(t)| \le [g^{j_k j_k}(t)]^{1/2} [g^{\bar{j}_l \bar{j}_l}(t)]^{1/2} \le C e^{-\beta_{\rm sil} t}$$
(26.22)

for all  $t \ge 0$ , where C only depends on  $C_A$ ,  $\eta_{ts}$ ,  $g^{ij}(0)$ ,  $a_r(0)$ ,  $\eta_{sh,0}$ ,  $c_{sil}$  and  $c_{\mathfrak{e}}$ . Thus (26.7) holds. Next let us turn to the shift vector field  $\chi^l = g^{0l}$ . Note that

$$\sigma(\iota_{\rm ts}, t) = \frac{n_{j_i} g^{0j_i}(t)}{\mathfrak{g}(\iota_{\rm ts}, t)}.$$

Combining this equality with (26.3), (26.19) and (26.20) yields

$$|n_{j_i}g^{0j_i}(t)| \le Ce^{-\eta_{\rm ts}t}\mathfrak{g}(\iota_{\rm ts},0)$$

for all  $t \ge 0$ , where C only depends on  $C_A$ ,  $C_B$  and  $\eta_{ts}$ . Thus the first estimate appearing in (26.8) holds. Finally, if  $n \in \mathbb{Z}^d$  corresponds to a  $\iota = \iota_{sil}$ , then

$$|\chi^{i}(t)n_{i}| \leq \eta_{\mathrm{sh},0}^{-1}(1-\eta_{\mathrm{sh},0}^{2})^{1/2}\mathfrak{g}_{\mathrm{T}}(\iota_{\mathrm{sil}},t)$$

for all  $t \ge 0$ . Combining this estimate with (26.21) yields the second part of (26.8), where  $C_2$  only depends on  $g^{ij}(0)$ ,  $\eta_{\text{sh},0}$ ,  $c_{\text{sil}}$  and  $c_{\mathfrak{e}}$ . The lemma follows.

# 26.3 Conditions ensuring the existence of a dominant noisy spatial direction

The purpose of the present section is to relate the conditions appearing in Chapter 6 with the conditions appearing in Part V.

**Lemma 26.5.** Let (M, g) be a canonical separable cosmological model manifold. Assume that there is an  $l \in \{1, ..., d\}$  such that g has a geometric dominant noisy spatial  $\mathbb{S}^1$ -direction corresponding

to l; cf. Definition 6.1. Assume, moreover, that there is an  $0 < \eta_{sh,0} \leq 1$  such that (3.4) holds for all  $t \geq 0$  and constants  $C_{sh}, \eta_{sh} > 0$  such that

$$|\chi(t)|_{\bar{g}} + |\dot{\chi}(t)|_{\bar{g}} \le C_{\rm sh} e^{-\eta_{\rm sh}t} \tag{26.23}$$

for all  $t \ge 0$ . Finally, assume that  $\bar{k}$  is  $C^0$ -future bounded; i.e., that there is a constant  $C_0 \ge 0$  such that

$$|k|_{\bar{g}} \le C_0 \tag{26.24}$$

for all  $t \geq 0$ . Then there are constants  $h_{\infty}^{ll}, C > 0$  such that

$$|e^{-2\beta_{n}t}h^{ll}(t) - h^{ll}_{\infty}| + |e^{-2\beta_{n}t}\partial_{t}h^{ll}(t) - 2\beta_{n}h^{ll}_{\infty}| \le Ce^{-\eta_{\max}t}$$
(26.25)

for all  $t \ge 0$ , where  $\eta_{mar} := \min\{2\eta_{sh}, \eta_n\}$ . Moreover,

$$e^{-2\beta_{n}t}\sum_{j,k\neq l}|g^{jk}(t)| \le Ce^{-2\eta_{n}t}, \quad e^{-2\beta_{n}t}\sum_{r=1}^{R}a_{r}^{-2}(t) \le Ce^{-2\eta_{n}t}$$
 (26.26)

for all  $t \ge 0$ . In addition,

$$e^{-2\beta_{n}t}\sum_{j\neq l}|g^{jl}(t)| \le Ce^{-\eta_{n}t}$$
 (26.27)

for all  $t \geq 0$ .

Turning to the time derivatives,

$$e^{-2\beta_{n}t}\sum_{i,j\neq l}|\partial_{t}h^{ij}(t)| \leq Ce^{-2\eta_{n}t}, \quad e^{-2\beta_{n}t}\sum_{j\neq l}|\partial_{t}h^{jl}(t)| \leq Ce^{-\eta_{n}t}$$
(26.28)

for all  $t \geq 0$ . Finally,

$$e^{-2\beta_{n}t}|\partial_{t}a_{r}^{-2}(t)| \le Ce^{-2\eta_{n}t}$$
(26.29)

for all  $t \geq 0$ .

**Remark 26.6.** The constant C only depends on  $g^{ij}(0)$ ,  $\bar{g}^{ij}(0)$ ,  $a_r(0)$ ,  $\|\mathfrak{e}\|_1$ ,  $\eta_{sh,0}$ ,  $\eta_{sh}$ ,  $C_{sh}$ ,  $C_n$ ,  $C_0$ ,  $\eta_n$  and  $\beta_n$ 

**Remark 26.7.** If, in addition to the assumptions of the lemma, there is a non-negative continuous function  $\mathfrak{e}_{\text{low}} \in L^1([0,\infty))$  such that (6.30) holds for all  $t \ge 0$ , then there is a constant C > 0 (depending only on  $\beta_n$ ,  $\|\mathfrak{e}_{\text{low}}\|_1$ ,  $\eta_{\text{sh},0}$ ,  $C_{\text{sh}}$  and  $\eta_{\text{sh}}$ ) such that  $e^{-\beta_n t}\mathfrak{g}(\iota,t) \ge C\nu_n(\iota)$  for all  $t \ge 0$  and all  $\iota \in \mathcal{I}_B$ . Here  $\nu_n(\iota)$  is defined by (6.12). In order to justify that the limit appearing in (6.12) exists, note that

$$\begin{split} e^{-2\beta_{n}t}\mathfrak{g}^{2}(\iota,t) = & \sum_{i,j\neq l} e^{-2\beta_{n}t}g^{ij}(t)n_{i}n_{j} + 2\sum_{j\neq l} e^{-2\beta_{n}t}g^{jl}(t)n_{j}n_{l} \\ & + e^{-2\beta_{n}t}g^{ll}(t)n_{l}^{2} + \sum_{r=1}^{R} e^{-2\beta_{n}t}a_{r}^{-2}(t)\nu_{r,i_{r}}^{2}(\iota) \to h_{\infty}^{ll}n_{l}^{2}, \end{split}$$

due to the conclusions of the lemma. In order to justify the lower bound  $e^{-\beta_n t} \mathfrak{g}(\iota, t) \geq C\nu_n(\iota)$ , note that an argument similar to the beginning of the proof of Lemma 25.12 yields

$$\begin{aligned} & 2\mathfrak{g}_{\mathrm{T}}\dot{\mathfrak{g}}_{\mathrm{T}} \leq & 2(\beta_{\mathrm{n}} + \mathfrak{e}_{\mathrm{low}})\bar{h}^{ij}n_{i}n_{j} - 2\chi^{i}\dot{\chi}^{j}n_{i}n_{j} \\ & \leq & 2\beta_{\mathrm{n}}\mathfrak{g}_{\mathrm{T}}^{2} + 2\beta_{\mathrm{n}}\chi^{i}\chi^{j}n_{i}n_{j} + 2\eta_{\mathrm{sh},0}^{-2}\mathfrak{e}_{\mathrm{low}}\mathfrak{g}_{\mathrm{T}}^{2} + 2\eta_{\mathrm{sh},0}^{-2}|\chi|_{\bar{g}}|\dot{\chi}|_{\bar{g}}\mathfrak{g}_{\mathrm{T}}^{2}. \end{aligned}$$

The remaining part of the time derivative of  $\mathfrak{g}^2$  can be estimated more easily. Appealing to (26.23) yields  $\dot{\ell}(\iota, t) \leq \beta_n + \mathfrak{e}_2(t)$  for all  $t \geq 0$  and  $0 \neq \iota \in \mathcal{I}_B$ . Here  $\mathfrak{e}_2 \in L^1([0, \infty))$  is a non-negative continuous function whose  $L^1$ -norm only depends on  $\beta_n$ ,  $\|\mathfrak{e}_{low}\|_1$ ,  $\eta_{sh,0}$ ,  $C_{sh}$  and  $\eta_{sh}$ . Thus

$$\partial_t \ln[e^{-\beta_{\mathbf{n}}t}\mathfrak{g}(\iota,t)] \leq \mathfrak{e}_2(t)$$

for all  $t \ge 0$  and all  $0 \ne \iota \in \mathcal{I}_{\mathrm{B}}$ . Integrating this estimate from t to  $\infty$  yields  $e^{-\beta_{\mathrm{n}}t}\mathfrak{g}(\iota, t) \ge e^{-\|\mathfrak{e}_2\|_1}\nu_{\mathrm{n}}(\iota)$  for all  $t \ge 0$  and all  $0 \ne \iota \in \mathcal{I}_{\mathrm{B}}$ .

*Proof.* The components of the inverse metric in the dominant noisy direction. Note that  $dx^l \in T^*_n \overline{M}$ . Appealing to (6.7) with  $\xi := dx^l$  thus yields

$$\left|\frac{1}{2}\bar{h}^{il}\bar{h}^{jl}\partial_t\bar{h}_{ij} + \beta_{n}\bar{h}^{ll}\right| \le C_{n}e^{-\eta_{n}t}\bar{h}^{ll}$$
(26.30)

(no summation on l) for all  $t \ge 0$ . This estimate is, in its turn, equivalent to

$$|\partial_t (e^{-2\beta_n t} \bar{h}^{ll})| \le 2C_n e^{-\eta_n t} e^{-2\beta_n t} \bar{h}^{ll}$$
(26.31)

for all  $t \ge 0$ . Thus

$$\exp(-2C_{\rm n}/\eta_{\rm n}) \le \frac{e^{-2\beta_{\rm n}t}\bar{h}^{ll}(t)}{\bar{h}^{ll}(0)} \le \exp(2C_{\rm n}/\eta_{\rm n})$$
(26.32)

for all  $t \ge 0$ . Combining (26.31) and (26.32) yields the conclusion that  $e^{-2\beta_n t} \bar{h}^{ll}(t)$  converges to, say,  $\bar{h}^{ll}_{\infty} > 0$ . Moreover,

$$|e^{-2\beta_{n}t}\bar{h}^{ll}(t) - \bar{h}^{ll}_{\infty}| \le C\bar{h}^{ll}(0)e^{-\eta_{n}t}$$
(26.33)

for all  $t \ge 0$ , where C only depends on  $C_n$  and  $\eta_n$ . Combining this estimate with (26.30) and (26.32) yields

$$|e^{-2\beta_{n}t}\partial_{t}\bar{h}^{ll}(t) - 2\beta_{n}\bar{h}^{ll}_{\infty}| \le C\bar{h}^{ll}(0)e^{-\eta_{n}t}$$
(26.34)

for all  $t \ge 0$ , where C only depends on  $C_n$ ,  $\beta_n$  and  $\eta_n$ . In order to estimate  $h^{ll}(t)$ , we also need to estimate  $\chi^l$  and  $\dot{\chi}^l$ . However, combining (26.23) with estimates of the form (25.44) (with  $n_i = \delta_{il}$ ) yields

$$e^{-\beta_{n}t}|\chi^{l}(t)| + e^{-\beta_{n}t}|\dot{\chi}^{l}(t)| \le C[\bar{h}^{ll}(0)]^{1/2}e^{-\eta_{\text{sh}}t}$$

for all  $t \ge 0$ , where we appealed to (26.32) and C only depends on  $C_{\rm sh}$ ,  $C_{\rm n}$  and  $\eta_{\rm n}$ . Combining this estimate with (25.7), (26.33) and (26.34) yields

$$|e^{-2\beta_{n}t}h^{ll}(t) - \bar{h}^{ll}_{\infty}| \le C\bar{h}^{ll}(0)e^{-\eta_{mar}t},$$
(26.35)

$$|e^{-2\beta_{n}t}\partial_{t}h^{ll}(t) - 2\beta_{n}\bar{h}^{ll}_{\infty}| \leq C\bar{h}^{ll}(0)e^{-\eta_{mar}t}$$

$$(26.36)$$

for all  $t \ge 0$ , where  $\eta_{\text{mar}} := \min\{2\eta_{\text{sh}}, \eta_{\text{n}}\}$  and C only depends on  $C_{\text{sh}}, C_{\text{n}}, \eta_{\text{n}}$  and  $\beta_{\text{n}}$ . Thus (26.25) holds.

The subdominant directions. Let us consider the remaining directions. In order to do so, let  $\iota \in \mathcal{I}_{\rm B}$  be such that  $\nu_{{\rm T},l}(\iota) = 0$ . If  $n = \nu_{\rm T}(\iota)$ , we then have  $v := n_i dx^i \in T^*_{\rm sub} \overline{M}$ . In analogy with the proof of Lemma 26.2, let us consider the time derivative of  $\mathfrak{g}^2(\iota, \cdot)$ . As before, this derivative can be divided into several different parts. Let us start by considering the first term on the right hand side of (25.43). It can be estimated by

$$\partial_t \bar{h}^{jk} n_j n_k = -\bar{h}^{ji} \bar{h}^{kp} \partial_t \bar{h}_{ip} n_j n_k = -2\bar{k}^{\sharp}(v,v)$$

$$\leq 2(\beta_n - \eta_n + \mathfrak{e}) \bar{h}^{jk} n_j n_k$$
(26.37)

for all  $t \ge 0$ , where we appealed to (25.15) and (6.8). On the other hand, combining (25.7), (25.27), (25.33) and (26.23) yields

$$|\bar{h}^{jk}(t)n_jn_k - h^{jk}(t)n_jn_k| \le Ce^{-2\eta_{\rm sh}t}\mathfrak{g}_{\rm T}^2(\iota,t)$$

for all  $t \ge 0$ , where C only depends on  $\eta_{sh,0}$  and  $C_{sh}$ . Combining this estimate with (25.33) and (26.37) yields

$$\partial_t \bar{h}^{jk}(t) n_j n_k \le 2(\beta_n - \eta_n) \mathfrak{g}_{\mathrm{T}}^2(\iota, t) + C(\mathfrak{e} + e^{-2\eta_{\mathrm{sh}}t}) \mathfrak{g}_{\mathrm{T}}^2(\iota, t)$$
(26.38)

for all  $t \ge 0$ , where C only depends on  $\eta_n$ ,  $\beta_n$ ,  $\eta_{sh,0}$  and  $C_{sh}$ . Next, the second term on the right hand side of (25.43) can be estimated using (25.45). In fact, combining (25.45) with (26.23) yields

$$2|\chi^{i}(t)n_{i}\dot{\chi}^{j}(t)n_{j}| \leq Ce^{-2\eta_{\rm sh}t}\mathfrak{g}_{\rm T}^{2}(\iota,t)$$

for all  $t \ge 0$ , where C only depends on  $\eta_{\rm sh,0}$  and  $C_{\rm sh}$ . Combining this estimate with (26.38) yields

$$2\mathfrak{g}_{\mathrm{T}}\dot{\mathfrak{g}}_{\mathrm{T}} \leq 2(\beta_{\mathrm{n}} - \eta_{\mathrm{n}} + \mathfrak{e}_{1})\mathfrak{g}_{\mathrm{T}}^{2}$$

for all  $t \ge 0$ , where

$$\mathbf{e}_1(t) = C[\mathbf{e}(t) + e^{-2\eta_{\rm sh}t}]$$

and  $C \ge 1$  only depends on  $\eta_n$ ,  $\beta_n$ ,  $\eta_{sh,0}$  and  $C_{sh}$ . Finally, a similar estimate holds for the time derivative of the remaining part of  $\mathfrak{g}^2$ ; cf. the arguments presented in connection with (26.18). Summing up,

$$2\mathfrak{g}\mathfrak{g}\mathfrak{g} \leq 2(\beta_{\mathrm{n}} - \eta_{\mathrm{n}} + \mathfrak{e}_{1})\mathfrak{g}^{2}$$

for all  $t \ge 0$ . Introducing  $\mathfrak{h}(\iota, t) := e^{-(\beta_n - \eta_n)t}\mathfrak{g}(\iota, t)$ , this estimate can be written

$$\partial_t \mathfrak{h}^2 \leq 2\mathfrak{e}_1 \mathfrak{h}^2$$

for all  $t \ge 0$ . Thus

$$e^{-2(\beta_{n}-\eta_{n})t}\mathfrak{g}^{2}(\iota,t) \leq e^{2c_{1}}\mathfrak{g}^{2}(\iota,0)$$

for all  $t \ge 0$ , where  $c_1 := \|\mathfrak{e}_1\|_1$  only depends on  $\|\mathfrak{e}\|_1$ ,  $\eta_{sh,0}$ ,  $\eta_n$ ,  $\beta_n$ ,  $\eta_{sh}$  and  $C_{sh}$ . This estimate yields (26.26). Finally, similarly to the proof of the first estimate in (26.22),  $j \ne l$  yields

$$e^{-2\beta_{n}t}|g^{jl}(t)| \leq [e^{-2\beta_{n}t}g^{jj}(t)]^{1/2}[e^{-2\beta_{n}t}g^{ll}(t)]^{1/2} \leq Ce^{-\eta_{n}t}$$

for all  $t \ge 0$ , where we appealed to (25.33), (26.32) and (26.26), and C only depends on the constants stated in Remark 26.6.

The time derivatives in the subdominant directions. Due to (25.28),

$$|\partial_t \bar{h}^{ij}| = |\bar{h}^{ip} \bar{h}^{jq} \partial_t \bar{h}_{pq}| = 2|\bar{k}_{pq} \bar{h}^{ip} \bar{h}^{jq}| \le 2|\bar{k}|_{\bar{g}} (\bar{h}^{ii})^{1/2} (\bar{h}^{jj})^{1/2}$$

(no summation on i or j) for all  $t \ge 0$ . Keeping (25.7) and (25.33) in mind, this estimate implies

$$\begin{aligned} \partial_t h^{ij} &|\leq |\partial_t \bar{h}^{ij}| + |\dot{\chi}^i| |\chi^j| + |\dot{\chi}^j| |\chi^i| \\ &\leq 2\eta_{\mathrm{sh},0}^{-2} [|\bar{k}|_{\bar{g}} + |\chi|_{\bar{g}} \cdot |\dot{\chi}|_{\bar{g}}] (h^{ii})^{1/2} (h^{jj})^{1/2} \end{aligned}$$

for all  $t \ge 0$ , where we appealed to (25.27) in the second step. Keeping (25.33), (26.23), (26.24), (26.26) and (26.32) in mind, this estimate implies that (26.28) holds. Finally, note that

$$e^{-2\beta_{\mathbf{n}}t}|\partial_t a_r^{-2}| = 2e^{-2\beta_{\mathbf{n}}t}a_r^{-2}|a_r^{-1}\partial_t a_r| \le 2C_0e^{-2\beta_{\mathbf{n}}t}a_r^{-2} \le Ce^{-2\eta_{\mathbf{n}}t}$$

for all  $t \ge 0$ , where C only depends on the constants stated in Remark 26.6. Thus (26.29) holds, and the lemma follows.

In the case that g has a geometric dominant noisy spatial direction corresponding to  $M_r$ , there is an analogous result.

**Lemma 26.8.** Let (M, g) be a canonical separable cosmological model manifold. Assume that there is an  $r_n \in \{1, \ldots, R\}$  such that g has a geometric dominant noisy spatial direction corresponding to  $M_{r_n}$ ; cf. Definition 6.4. Assume, moreover, that there are constants  $C_{sh}, C_0, \eta_{sh} > 0$  such that (26.23) and (26.24) hold for all  $t \ge 0$ . Then there are constants  $q_{\infty,n}, C > 0$  such that

$$e^{-2\beta_{n}t}a_{r_{n}}^{-2}(t) - q_{\infty,n}| + |e^{-2\beta_{n}t}\partial_{t}a_{r_{n}}^{-2}(t) - 2\beta_{n}q_{\infty,n}| \le Ce^{-\eta_{n}t},$$
(26.39)

for all  $t \geq 0$ . Moreover,

$$e^{-2\beta_{n}t}\sum_{j,k}|g^{jk}(t)| \le Ce^{-2\eta_{n}t}, \quad e^{-2\beta_{n}t}\sum_{r\ne r_{n}}a_{r}^{-2}(t) \le Ce^{-2\eta_{n}t}$$
 (26.40)

for all  $t \geq 0$ . Finally,

$$e^{-2\beta_{\mathbf{n}}t}\sum_{i,j}|\partial_t h^{ij}(t)| \le Ce^{-2\eta_{\mathbf{n}}t},\tag{26.41}$$

$$e^{-2\beta_{n}t}\sum_{r\neq r_{n}}|\partial_{t}a_{r}^{-2}(t)| \leq Ce^{-2\eta_{n}t}$$
 (26.42)

for all  $t \geq 0$ .

**Remark 26.9.** The constant C appearing in the estimates has the dependence described in Remark 26.6.

**Remark 26.10.** If, in addition to the assumptions of the lemma, there is a non-negative continuous function  $\mathfrak{e}_{\text{low}} \in L^1([0,\infty))$  such that (6.30) holds for all  $t \ge 0$ , then there is a constant C > 0 (depending only on  $\beta_n$ ,  $\|\mathfrak{e}_{\text{low}}\|_1$ ,  $\eta_{\text{sh},0}$ ,  $C_{\text{sh}}$  and  $\eta_{\text{sh}}$ ) such that  $e^{-\beta_n t}\mathfrak{g}(\iota,t) \ge C\nu_n(\iota)$  for all  $t \ge 0$  and all  $\iota \in \mathcal{I}_{\text{B}}$ . The justification of this statement is similar to the one given in Remark 26.7.

*Proof.* The statements concerning the subdominant directions are a consequence of the arguments presented in the proof of Lemma 26.5. In order to prove (26.39), note that since (6.7) holds for all  $v \in T_n^* \overline{M}$ ,

$$|\dot{a}_{r_{\mathrm{n}}}/a_{r_{\mathrm{n}}}+\beta_{\mathrm{n}}| \leq C_{\mathrm{n}}e^{-\eta_{\mathrm{n}}t}$$

for all  $t \ge 0$ . Thus

$$|\partial_t (e^{-2\beta_n t} a_{r_n}^{-2})| = |e^{-2\beta_n t} [-2\dot{a}_{r_n}/a_{r_n} - 2\beta_n] a_{r_n}^{-2}| \le 2C_n e^{-\eta_n t} \cdot e^{-2\beta_n t} a_{r_n}^{-2}$$

for all  $t \ge 0$ . In particular, there is a  $q_{\infty,n} > 0$  such that

$$\ln \frac{q_{\infty,n}}{e^{-2\beta_{n}t}a_{r_{n}}^{-2}(t)} \bigg| \le 2C_{n}\eta_{n}^{-1}e^{-\eta_{n}t}$$

for all  $t \ge 0$ . Using this estimate, it can be verified that (26.39) holds. The lemma follows.

#### 26.4 The asymptotically diagonal setting

In the present section, we relate the conditions stated in Section 7.2 with those appearing in Part VI. We begin by deriving conclusions concerning  $\bar{g}_{ij}$ , given the assumption that  $\bar{g}$  is  $C^0$ -asymptotically diagonal; cf. Definition 7.3.

**Lemma 26.11.** Let (M, g) be a canonical separable cosmological model manifold. Assume that the corresponding metric  $\bar{g}$  is  $C^0$ -asymptotically diagonal; cf. Definition 7.3. Then there is a constant C > 1, depending only on the metric g, such that

$$C^{-1} \le \bar{g}^{ii}(t)\bar{g}_{ii}(t) \le C$$
 (26.43)

for all  $t \ge 0$  (no summation on i). Moreover, there is a constant C > 0, depending only on the metric g, such that for  $i \ne j$ ,  $i, j \in \{1, ..., d\}$ ,

$$|[\bar{g}^{ii}(t)]^{1/2}\bar{g}_{ij}(t)[\bar{g}^{jj}(t)]^{1/2}| \le Ce^{-\kappa_{\rm od}t}$$
(26.44)

for all  $t \ge 0$  (no summation on i or j).

Proof. Let  $A \in \mathbf{M}_d(\mathbb{R})$  be the matrix with components  $\bar{g}^{ij}$ ; let D be the matrix obtained from A by setting the off-diagonal components to zero; let R be the remainder (so that A = D + R); and let  $S := -D^{-1/2}RD^{-1/2}$ . If  $S^{ij}$  denotes the ij'th element of S, note that  $S^{ii} = 0$  for  $i \in \{1, \ldots, d\}$  (no summation on i) and that if  $i \neq j, i, j \in \{1, \ldots, d\}$ , then

$$|S^{ij}| = |(\bar{g}^{ii})^{-1/2} \bar{g}^{ij} (\bar{g}^{jj})^{-1/2}| \le C_{\rm od} e^{-\kappa_{\rm od}}$$

(no summation on *i* or *j*) for all  $t \ge 0$ , where we appealed to (7.17). In particular,  $||S|| \le d \cdot C_{\text{od}} e^{-\kappa_{\text{od}}t}$  for all  $t \ge 0$ . On the other hand,

$$A = D + R = D^{1/2} (\mathrm{Id}_d - S) D^{1/2},$$

so that

$$D^{1/2}A^{-1}D^{1/2} = (\mathrm{Id}_d - S)^{-1} = \sum_{l=0}^{\infty} S^l,$$

#### 26.4. THE ASYMPTOTICALLY DIAGONAL SETTING

assuming t to be large enough that  $||S|| \leq 1/2$ . In particular,

$$|D^{1/2}A^{-1}D^{1/2} - \mathrm{Id}_d|| \le 2d \cdot C_{\mathrm{od}}e^{-\kappa_{\mathrm{od}}t}$$

for all t large enough that  $||S|| \le 1/2$ . Since the left hand side of this inequality is always bounded, there is a constant C > 0 such that

$$\|D^{1/2}A^{-1}D^{1/2} - \mathrm{Id}_d\| \le Ce^{-\kappa_{\mathrm{od}}t}$$
(26.45)

for all  $t \ge 0$ , where C only depends on the metric g. One particular consequence of this estimate is that

$$|\bar{g}^{ii}(t)\bar{g}_{ii}(t) - 1| \le Ce^{-\kappa_{\rm od}t}$$

for all  $t \ge 0$  (no summation on *i*), where *C* only depends on the metric *g*. As a consequence of this estimate, there is a constant C > 1 such that (26.43) holds. Finally, (26.45) implies that for  $i \ne j, i, j \in \{1, \ldots, d\}$ , (26.44) holds.

Next, we turn to the problem of verifying that there are geometric conditions ensuring that the (non-geometric) assumptions of Definition 20.1 are satisfied.

**Lemma 26.12.** Consider (1.2). Assume the associated metric to be such that (M, g) is a canonical separable cosmological model manifold. Assume (1.2) to be  $C^2$ -balanced; cf. Definition 3.8. Assume, moreover, the shift vector field to be negligible,  $\bar{k}$  to be diagonally convergent and  $\bar{g}$  to be  $C^2$ -asymptotically diagonal; cf. Definition 7.3. Finally, assume (1.2) to be such that the main coefficients are convergent; cf. Definition 7.5. Then, for each  $j \in \{1, \ldots, d\}$ , there are constants  $g_{\infty}^{jj}, C > 0$  such that

$$|e^{-2\beta_j t}g^{jj}(t) - g^{jj}_{\infty}| \le Ce^{-\kappa_{\max} t},\tag{26.46}$$

$$|e^{-2\beta_j t}\partial_t g^{jj}(t) - 2\beta_j g^{jj}_{\infty}| \le C e^{-\kappa_{\max} t}$$
(26.47)

for all  $t \ge 0$ , where  $\kappa_{\text{mar}} := \min\{2\kappa_{\text{sh}}, \kappa_{\text{d}}\}\ and \ C \ only \ depends \ on \ C_{\text{sh}}, \ C_{\text{d}}, \ \kappa_{\text{d}}, \ \beta_j, \ \chi^j(0) \ and \ g^{jj}(0).$  Moreover, for each  $r \in \{1, \ldots, R\}$ , there are constants  $a_{r,\infty}, C > 0$  such that

$$|e^{\beta_{\mathrm{Ri},r}t}a_r(t) - a_{r,\infty}| \le Ce^{-\kappa_{\mathrm{d}}t},\tag{26.48}$$

$$|e^{\beta_{\mathrm{Ri},r}t}\dot{a}_r(t) + \beta_{\mathrm{Ri},r}a_{r,\infty}| \le Ce^{-\kappa_{\mathrm{d}}t}$$
(26.49)

for all  $t \ge 0$ , where C only depends on  $C_d$ ,  $\kappa_d$ ,  $\beta_{\mathrm{Ri},r}$  and  $a_r(0)$ . There is also a constant C > 0 such that

$$e^{-2\beta_j t} |\dot{g}^{jj}(t)| + e^{-2\beta_j t} |\ddot{g}^{jj}(t)| \le C$$
(26.50)

for all  $t \ge 0$  and all  $j \in \{1, \ldots, d\}$ , where C only depends on the constants appearing in the definition of  $C^2$ -balance,  $C_{\rm sh}$ ,  $C_{\rm d}$ ,  $\kappa_{\rm d}$ ,  $\beta_j$ ,  $\chi^j(0)$  and  $g^{jj}(0)$ . Similarly, there is a constant C > 0 such that

$$e^{\beta_{\mathrm{Ri},r}t}|\dot{a}_r(t)| + e^{\beta_{\mathrm{Ri},r}t}|\ddot{a}_r(t)| \le C$$
(26.51)

for all  $t \ge 0$  and all  $r \in \{1, ..., R\}$ , where C only depends on the constants appearing in the definition of  $C^2$ -balance,  $C_d$ ,  $\kappa_d$ ,  $\beta_{\text{Bi},r}$  and  $a_r(0)$ . There is also a constant C > 0 such that

$$e^{-\beta_j t} \|X^j(t)\| + e^{-\beta_j t} \|\dot{X}^j(t)\| \le C$$
(26.52)

for all  $t \ge 0$ , where C only depends on the constants appearing in the definition of  $C^2$ -balance,  $C_{\rm sh}$ ,  $C_{\rm d}$ ,  $\kappa_{\rm d}$ ,  $\beta_j$ ,  $\chi^j(0)$  and  $g^{jj}(0)$ . Turning to the shift vector field,

$$\sup_{0 \neq \xi \in \mathbb{R}^d} \frac{|g^{0l}(t)\xi_l|}{[g^{jl}(t)\xi_j\xi_l]^{1/2}} + \sup_{0 \neq \xi \in \mathbb{R}^d} \frac{|\dot{g}^{0l}(t)\xi_l|}{[g^{jl}(t)\xi_j\xi_l]^{1/2}} \le \eta_{\mathrm{sh},0}^{-1}C_{\mathrm{sh}}e^{-\kappa_{\mathrm{sh}}t}$$
(26.53)

for all  $t \ge 0$ . Finally, there are constants  $C, \eta_{od} > 0$  such that

$$\sum_{j \neq l} [|g^{jl}(t)\xi_j\xi_l| + |\dot{g}^{jl}(t)\xi_j\xi_l| + |\ddot{g}^{jl}(t)\xi_j\xi_l|] \le Ce^{-\eta_{\rm od}t}\sum_j g^{jj}(t)\xi_j\xi_j$$
(26.54)

for all  $t \ge 0$  and  $\xi \in \mathbb{R}^d$ , where C only depends on the metric g and  $\eta_{\text{od}} := \min\{\kappa_{\text{sh}}, \kappa_{\text{od}}\}$ .

**Remark 26.13.** If the assumptions of the lemma hold, then (1.2) is diagonally dominated, balanced and convergent; cf. Definition 20.1.

**Remark 26.14.** If we, in addition to the assumptions of the lemma, require  $|\ddot{\chi}|_{\bar{g}}$  to be bounded by the right hand side of (7.19), then  $\eta_{\rm od}$  appearing in (26.54) can be improved to  $\eta_{\rm od} := \min\{2\kappa_{\rm sh}, \kappa_{\rm od}\}$ .

*Proof.* That (26.46) and (26.47) hold follows by an argument similar to the beginning of the proof of Lemma 26.5. That (26.48) and (26.49) hold is a consequence of an argument similar to the beginning of the proof of Lemma 26.8.

The time derivatives of the diagonal components of the metric. In order to estimate the first and second time derivatives of  $g^{jj}$ , note that  $g^{jj}(t) = \mathfrak{g}_{\mathrm{T}}^2(\iota, t)$ , where  $\iota$  has been chosen so that  $\nu_{\mathrm{T},j}(\iota) = 1$  and all the other components of  $\iota$  vanish. On the other hand, (25.47) yields

$$|\mathfrak{g}_{\mathrm{T}}\dot{\mathfrak{g}}_{\mathrm{T}}| \leq \eta_{\mathrm{sh},0}^{-2}(|\bar{k}_{\mathrm{T}}|_{\bar{g}} + |\chi|_{\bar{g}} \cdot |\dot{\chi}|_{\bar{g}})\mathfrak{g}_{\mathrm{T}}^2$$

for all  $t \ge 0$ . Thus

$$e^{-2\beta_j t} |\dot{g}^{jj}| \le C e^{-2\beta_j t} g^{jj} \le C \tag{26.55}$$

for all  $t \ge 0$ , where C > 0 only depends on the constants appearing in the definition of  $C^2$ -balance,  $C_{\rm sh}$ ,  $C_{\rm d}$ ,  $\kappa_{\rm d}$ ,  $\beta_j$ ,  $\chi^j(0)$  and  $g^{jj}(0)$ . Turning to  $\ddot{g}^{jj}$ , note that it can be written

$$\ddot{g}^{\jmath\jmath} = 2\mathfrak{g}_{\mathrm{T}}\ddot{\mathfrak{g}}_{\mathrm{T}} + 2\dot{\mathfrak{g}}_{\mathrm{T}}\dot{\mathfrak{g}}_{\mathrm{T}}.$$

On the other hand, the right hand side can be estimated by (25.49). Thus  $e^{-2\beta_j t} |\ddot{g}^{jj}| \leq C$  for all  $t \geq 0$ , where C > 0 has the same dependence as in the case of (26.55). To conclude, (26.50) holds. Due to (25.20), (25.21), (26.48) and the fact that the equation is  $C^2$ -balanced, it is also clear that (26.51) holds.

The main coefficients. The estimate (26.52) follows from the fact that  $\mathcal{X}$  is  $C^1$ -future bounded; (25.27); (25.33); and (26.46).

The shift vector field. Turning to the shift vector field, note that if  $\xi \in \mathbb{R}^d$ , then (25.27) yields

$$|g^{0l}\xi_l| \le |\chi|_{\bar{g}}(\bar{g}^{ij}\xi_i\xi_j)^{1/2}.$$

Combining this estimate with (25.33) and (7.19) yields

$$\frac{|g^{0l}(t)\xi_l|}{[g^{ij}(t)\xi_i\xi_j]^{1/2}} \le \eta_{\mathrm{sh},0}^{-1}C_{\mathrm{sh}}e^{-\kappa_{\mathrm{sh}}t}$$

for all  $0 \neq \xi \in \mathbb{R}^d$  and all  $t \ge 0$ . A similar estimate holds for  $\dot{\chi}$ . In fact, (26.53) holds.

The off-diagonal terms. Finally, let us consider the off-diagonal terms. Due to (7.17), we know that if  $i \neq j, i, j \in \{1, ..., d\}$ , then

$$|\bar{g}^{ij}(t)| = |\bar{g}^{\sharp}(dx^{i}, dx^{j})| \le C_{\rm od} e^{-\kappa_{\rm od} t} [\bar{g}^{ii}(t)]^{1/2} [\bar{g}^{jj}(t)]^{1/2}$$

for all  $t \ge 0$  (no summation on *i* or *j*). Thus

$$\sum_{i \neq j} |\bar{g}^{ij}(t)\xi_i\xi_j| \leq \sum_{i \neq j} C_{\text{od}} e^{-\kappa_{\text{od}}t} [\bar{g}^{ii}(t)\xi_i^2]^{1/2} [\bar{g}^{jj}(t)\xi_j^2]^{1/2} \\ \leq d \cdot \eta_{\text{sh},0}^{-2} C_{\text{od}} e^{-\kappa_{\text{od}}t} \sum_j g^{jj}(t)\xi_j^2$$
(26.56)

for all  $t \ge 0$ , where we appealed to (25.33). Before deriving a similar estimate with  $\bar{g}^{ij}$  on the left hand side replaced by  $g^{ij}$ , note that

$$\begin{aligned} |\chi^{i}\xi_{i}|^{2} \leq |\chi|_{\bar{g}}^{2}\bar{g}^{ij}\xi_{i}\xi_{j} &= |\chi|_{\bar{g}}^{2}\left(\sum_{j}\bar{g}^{jj}\xi_{j}\xi_{j} + \sum_{i\neq j}\bar{g}^{ij}\xi_{i}\xi_{j}\right) \\ \leq \eta_{\mathrm{sh},0}^{-2}C_{\mathrm{sh}}^{2}e^{-2\kappa_{\mathrm{sh}}t}(1 + d \cdot C_{\mathrm{od}}e^{-\kappa_{\mathrm{od}}t})\sum_{j}g^{jj}\xi_{j}\xi_{j}\end{aligned}$$
(26.57)

for all  $t \ge 0$ , where appeal to (25.27) in the first step; and (25.33), (7.19) and (26.56) in the third step. In short

$$|\chi^i \xi_i| \le C e^{-\kappa_{\rm sh} t} \left(\sum_j g^{jj} \xi_j \xi_j\right)^{1/2} \tag{26.58}$$

for all  $t \ge 0$ , where C only depends on  $C_{\rm sh}$ ,  $C_{\rm od}$  and the constants appearing in the definition of  $C^2$ -balance. Combining this estimate with (25.7) and (26.56) yields

$$\sum_{i \neq j} |g^{ij}(t)\xi_i\xi_j| \le Ce^{-\min\{\kappa_{\rm od}, 2\kappa_{\rm sh}\}t} \sum_j g^{jj}(t)\xi_j\xi_j \tag{26.59}$$

for all  $t \ge 0$ , where C has the same dependence as in the case of (26.58).

The first time derivative of the off-diagonal terms. Turning to the time derivative of  $g^{ij}$ , note that  $\partial_t \bar{g}^{ij} = -2\bar{k}^{ij}$ . For  $i \neq j, i, j \in \{1, \ldots, d\}$ ,

$$|\partial_t \bar{g}^{ij}| = 2|\bar{k}^{ij}| = 2|\bar{k}^{\sharp}(dx^i, dx^j)| \le 2C_{\text{od},1}e^{-\kappa_{\text{od}}t}(\bar{g}^{ii})^{1/2}(\bar{g}^{jj})^{1/2}$$

for all  $t \ge 0$ , where we used the fact that  $\bar{g}$  is  $C^2$ -asymptotically diagonal. Consequently, an argument similar to the proof of (26.59) yields

$$\sum_{i \neq j} |\dot{g}^{ij}(t)\xi_i\xi_j| \le Ce^{-\min\{\kappa_{\rm od}, 2\kappa_{\rm sh}\}t} \sum_j g^{jj}(t)\xi_j\xi_j \tag{26.60}$$

for all  $t \ge 0$ , where C only depends on  $C_{\rm sh}$ ,  $C_{\rm od}$ ,  $C_{\rm od,1}$  and the constants appearing in the definition of  $C^2$ -balance.

The second time derivative of the off-diagonal terms. Turning to the second time derivative of  $\bar{g}^{ij}$ , note that

$$\partial_t^2 \bar{g}^{ij} = 8\bar{k}^{il}\bar{k}_l^{\ j} - 2(\mathcal{L}_U\bar{k})^{ij} = 8\sum_{l\neq i}\bar{k}^{il}\bar{k}_l^{\ j} + 8\bar{k}^{ii}\bar{k}_i^{\ j} - 2(\mathcal{L}_U\bar{k})^{ij}$$
(26.61)

(no summation on i in the second term on the far right hand side). Let us estimate

$$\begin{aligned} |\bar{k}_{l}^{j}| \leq & |\bar{g}_{ll}\bar{k}^{lj}| + \sum_{r \neq l} |\bar{g}_{lr}\bar{k}^{rj}| \leq C |\bar{k}|_{\bar{g}}(\bar{g}^{ll})^{-1/2} (\bar{g}^{jj})^{1/2} \\ &+ C \sum_{r \neq l} e^{-\kappa_{\text{od}}t} (\bar{g}^{ll})^{-1/2} (\bar{g}^{rr})^{-1/2} |\bar{k}|_{\bar{g}} (\bar{g}^{rr})^{1/2} (\bar{g}^{jj})^{1/2} \end{aligned}$$

$$(26.62)$$

for all  $t \ge 0$  (no summation on l), where we appeal to (25.28), (26.43) and (26.44). In particular, there is thus a constant C > 0 such that

$$|\bar{k}_l^{\ j}| \le C(\bar{g}^{ll})^{-1/2} (\bar{g}^{jj})^{1/2} \tag{26.63}$$

for all  $t \ge 0$ , where C only depends on the the metric g. However, in the case that  $l \ne j$ , this estimate can be improved by, in the last step of (26.62), appealing to (7.18). Doing so yields

$$|\bar{k}_l^{\ j}| \le C e^{-\kappa_{\rm od} t} (\bar{g}^{ll})^{-1/2} (\bar{g}^{jj})^{1/2} \tag{26.64}$$

for all  $t \ge 0$  and  $l \ne j$   $(l, j \in \{1, \dots, d\})$ , where C only depends on the metric g. Returning to (26.61), consider the first term on the far right hand side. Appealing to (7.18) and (26.63) yields the existence of a constant C > 0 such that

$$\sum_{l \neq i} |8\bar{k}^{il}\bar{k}_l^{\,j}| \le C e^{-\kappa_{\rm od}t} (\bar{g}^{ii})^{1/2} (\bar{g}^{jj})^{1/2} \tag{26.65}$$

for all  $t \ge 0$ , where C only depends on the metric g. Concerning the second term on the far right hand side of (26.61), appealing to (25.28) and (26.64) yields the existence of a constant C > 0such that

$$|8\bar{k}^{ii}\bar{k}_i^{\ j}| \le Ce^{-\kappa_{\rm od}t}(\bar{g}^{ii})^{1/2}(\bar{g}^{jj})^{1/2} \tag{26.66}$$

for all  $t \ge 0$ , where C only depends on the metric g. Combining (26.61), (26.65) and (26.66) with (7.18) yields a constant C > 0 such that

$$|\partial_t^2 \bar{g}^{ij}| \le C e^{-\kappa_{\rm od} t} (g^{ii})^{1/2} (g^{jj})^{1/2}$$

for all  $t \ge 0$  and  $i \ne j$   $(i, j \in \{1, \ldots, d\})$ , where we also appealed to (25.33) and C only depends on the metric g. In order to estimate the second time derivative of  $g^{ij}$ , we also need to estimate  $\dot{\chi}^i \dot{\chi}^j$  and  $\ddot{\chi}^i \chi^j$ . The first of these expressions can be estimated as before. However, in the case of the second expression, we can estimate  $\chi^i$  as before, but in order to estimate  $\ddot{\chi}^j$ , we need to appeal to the fact that the shift vector field is  $C^2$ -future bounded. This yields a constant C > 0such that

$$|\ddot{\chi}^i \chi^j \xi_i \xi_j| \le C e^{-\kappa_{\rm sh} t} \sum_j g^{jj} \xi_j \xi_j$$

for all  $t \ge 0$ , where C only depends on the metric g. Summing up, (26.54) holds and the lemma follows.

# Bibliography

- Anderson, M. T.: On Long-Time Evolution in General Relativity and Geometrization of 3-Manifolds. Commun. Math. Phys., 222, no. 3, 533–567 (2001)
- [2] Andersson, L.: The global existence problem in general relativity. The Einstein equations and the large scale behavior of gravitational fields, Birkhäuser, Basel, 71–120 (2004)
- [3] Andersson, L., Moncrief, V.: Future complete vacuum spacetimes. The Einstein equations and the large scale behavior of gravitational fields, Birkhäuser, Basel, 299–330 (2004)
- [4] Andréasson, H., Ringström, H.: Proof of the cosmic no-hair conjecture in the T<sup>3</sup>-Gowdy symmetric Einstein-Vlasov setting. J. Eur. Math. Soc. (JEMS) 18, no. 7, 1565–1650 (2016)
- [5] Andersson, L., Rendall, A. D.: Quiescent cosmological singularities. Commun. Math. Phys. 218, 479–511 (2001)
- [6] Bieri, L., Zipser, N.: Extensions of the Stability Theorem of the Minkowski Space in General Relativity. AMS-IP. Studies in Advanced Mathematics. Cambridge. MA. (2009)
- [7] Choquet-Bruhat, Y., Moncrief V.: Future Global in Time Einsteinian Spacetimes with U(1)Isometry Group. Ann. Henri Poincaré **2** 1001–64 (2001)
- [8] Choquet-Bruhat, Y.: Future complete U(1) symmetric Einsteinian spacetimes, the unpolarized case. The Einstein equations and the large scale behavior of gravitational fields, Birkhäuser, Basel, 251–298 (2004)
- [9] Christodoulou, D., Klainerman, S.: The global nonlinear stability of the Minkowski space. Princeton University Press, Princeton (1993)
- [10] Chruściel, P.T., Isenberg, J., Moncrief, V.: Strong Cosmic Censorship in Polarized Gowdy Spacetimes. Class. Quantum Grav. 7, 1671–1680 (1990)
- [11] Damour, T., Henneaux, M., Nicolai, H.: Cosmological billiards. Class. Quantum Grav. 20, R145 (2003)
- [12] Damour, T., Henneaux, M., Rendall, A. D., Weaver, M.: Kasner-like behaviour for subcritical Einstein-matter systems. Ann. H. Poincaré 3, 1049–1111 (2002)
- [13] Damour, T., Nicolai, H.: Higher order M theory corrections and the Kac-Moody algebra E10. Class. Quantum Grav. 22 2849 (2005)
- [14] Fischer, A., Moncrief, V.: The reduced Einstein equations and the conformal volume collapse of 3-manifolds. Class. Quantum Grav. 18, 4493–44515 (2001)
- [15] Friedrich, H.: On the existence of n-geodesically complete or future complete solutions of Einstein's field equations with smooth asymptotic structure. Commun. Math. Phys. 107, 587– 609 (1986)

- [16] Hadzic, M., Speck, J.: The Global Future Stability of the FLRW Solutions to the Dust-Einstein System with a Positive Cosmological Constant. Preprint, arXiv:1309.3502
- [17] Heinzle, J. M., Rendall, A. D.: Power-law Inflation in Spacetimes without Symmetry. Commun. Math.Phys. 269, 1–15 (2007)
- [18] Heinzle, J. M., Uggla, C., Niklas Rohr, N.: The cosmological billiard attractor. Adv. Theor. Math. Phys. 13, 293–407 (2009)
- [19] Isenberg, J., Moncrief, V.: Asymptotic behavior of the gravitational field and the nature of singularities in Gowdy spacetimes. Ann. Phys. (N.Y.) 199, 84–122 (1990)
- [20] Jurke, T.: On the future asymptotics of polarized Gowdy T<sup>3</sup>-models. Class. Quantum Grav. 20, 173–191 (2003)
- [21] Lee, H.: Asymptotic behaviour of the Einstein-Vlasov system with a positive cosmological constant. Math. Proc. Camb. Phil. Soc. 137, 495–509 (2004)
- [22] LeFloch, P. G., Smulevici, J.: Future asymptotics and geodesic completeness of polarized T<sup>2</sup>-symmetric spacetimes. Anal. PDE 9, 363–395 (2016)
- [23] Lindblad, H., Rodnianski, I.: Global existence for the Einstein vacuum equations in wave coordinates. Commun. Math. Phys., 256, no. 1, 43–110 (2005)
- [24] Lindblad, H., Rodnianski, I.: The global stability of Minkowski space-time in harmonic gauge. Annals of Math 171, no. 3, 1401–1477 (2010)
- [25] Luo, X., Isenberg, J.: Power law inflation with electromagnetism. Ann. Physics 334, 420–454 (2013)
- [26] Nungesser, E.: Future global non-linear stability of surface symmetric solutions of the Einstein-Vlasov system with a cosmological constant. Preprint, arXiv:1402.7085
- [27] O'Neill, B.: Semi Riemannian Geometry. Academic Press, Orlando (1983)
- [28] Ringström, H.: The Bianchi IX attractor, Annales Henri Poincaré 2, 405–500 (2001)
- [29] Ringström, H.: On a wave map equation arising in general relativity. Commun. Pure Appl. Math. 57, 657–703 (2004)
- [30] Ringström, H.: Data at the moment of infinite expansion for polarized Gowdy. Class. Quantum Grav. 22, No 9, 1647–1653 (2005)
- [31] Ringström, H.: Existence of an asymptotic velocity and implications for the asymptotic behaviour in the direction of the singularity in T<sup>3</sup>-Gowdy. Commun. Pure Appl. Math. 59, 977–1041 (2006)
- [32] Ringström, H.: Future stability of the Einstein non-linear scalar field system. Invent. math. 173, 123–208 (2008)
- [33] Ringström, H.: Strong cosmic censorship in T<sup>3</sup>-Gowdy spacetimes. Ann. Math. 170, 1181– 1240 (2009)
- [34] Ringström, H.: The Cauchy problem in General Relativity, European Mathematical Society, Zürich (2009)
- [35] Ringström, H.: Power law inflation. Commun. Math. Phys. 290, 155–218 (2009)
- [36] Ringström, H.: On the Topology and Future Stability of the Universe, Oxford University Press, Oxford (2013)

- [37] Ringström, H.: Instability of spatially homogeneous solutions in the class of  $T^2$ -symmetric solutions to Einstein's vacuum equations. Commun. Math. Phys., **334**, 1299–1375 (2015)
- [38] Rodnianski, I., Speck, J.: The nonlinear future stability of the FLRW family of solutions to the irrotational Euler-Einstein system with a positive cosmological constant. J. Eur. Math. Soc. (JEMS) 15, no. 6, 2369–2462 (2013)
- [39] Rodnianski, I., Speck, J.: A Regime of Linear Stability for the Einstein-Scalar Field System with Applications to Nonlinear Big Bang Formation. Preprint, arXiv:1407.6293
- [40] Rodnianski, I., Speck, J.: Stable Big Bang Formation in Near-FLRW Solutions to the Einstein-Scalar Field and Einstein-Stiff Fluid Systems. Preprint, arXiv:1407.6298
- [41] Rodino, L.: Linear Partial Differential Operators in Gevrey Spaces, World Scientific Books, Singapore (1993)
- [42] Rudin, W.: Principles of Mathematical Analysis, Third Edition, McGraw-Hill, Singapore (1976)
- [43] Rudin, W. P.: Real and Complex Analysis, McGraw Hill, Singapore (1986)
- [44] Serre, D.: Matrices, Theory and Applications, Second Edition, New York (2010)
- [45] Speck, J.: The nonlinear future stability of the FLRW family of solutions to the Euler-Einstein system with a positive cosmological constant. Selecta Math. (N.S.) **18**, no. 3, 633–715 (2012)
- [46] Spitzbart, A.: A generalization of Hermite's interpolation formula. Amer. Math. Monthly 67, 42–46 (1960)
- [47] Svedberg, C.: Future Stability of the Einstein-Maxwell-Scalar Field System. Ann. Henri Poincaré 12, No. 5, 849–917 (2011)
- [48] Tchapnda, S. B., Rendall, A. D.: Global existence and asymptotic behaviour in the future for the Einstein–Vlasov system with positive cosmological constant. Class. Quantum Grav. 20, 3037–3049 (2003)
- [49] Wald, R.: Asymptotic behaviour of homogeneous cosmological models in the presence of a positive cosmological constant. Phys. Rev. D 28, 2118–2120 (1983)
- [50] Xue, B., Garfinkle, D., Pretorius, F., Steinhardt, P. J.: Nonperturbative analysis of the evolution of cosmological perturbations through a nonsingular bounce. Phys. Rev. D 88, 083509, (2013)

#### BIBLIOGRAPHY

# Index

Notation Auxiliary functions V, 5 $\langle \cdot \rangle$ , 34  $\mathfrak{H}_q(\iota, t), 310$  $\mathfrak{H}_q^{(s)}(\iota,t), 310$  $\omega_{\rm sh}, 63$  $\varphi_{\rm tot},\,63$ Bilinear forms  $\langle \cdot, \cdot \rangle_{\rm B}, 371$  $\langle \cdot, \cdot \rangle_g, \, 370$ Coefficients, equation  $X^j, 4$  $Y^1, 20$  $\alpha, 4$  $\bar{X}_{n}(\iota), 61$ l, 20  $\operatorname{sgn}_n(\iota), 62$  $X_{\infty,n}, 62$  $\varsigma$ , 20  $\zeta, 4$ Coefficients, Fourier side  $X(\iota, t), 109, 372$  $\bar{X}_{\text{diag}}^q(\iota), 79$  $\ell(\iota, t), 109$  $\mathfrak{g}(\iota,t), 372$  $\mathfrak{g}_{\infty}, 75$  $\mathfrak{g}_{\infty}(\iota), 121$  $\mathfrak{g}_{\rm sil}(\iota,t), 151$  $\mathfrak{g}_{ts}(\iota,t), 151$  $\sigma(\iota, t), 109, 372$  $\tilde{X}^q_{\infty}, 79$ Conditions  $C^{0}$ -asymptotically diagonal metric, 77 $C^1$ -silent equation, 34  $C^k$ -balanced equation, 30  $C^k$ -future bounded  $\mathcal{X}$ , 29  $C^k$ -future bounded second fundamental form, 28  $C^k$ -future bounded shift vector field,

#### 28 $E_{\rm ts}$ -adapted function, 51 X-degenerate weakly transparent equation, 122 X-non-trivial weakly transparent equation, 122 $\bar{g}^{\sharp}$ convergent, 45 $\bar{g}^{\sharp}$ expanding, 45 $\mathcal{X} C^k$ -future bounded, 29 Constants $C_{\rm ini}, 153$ R, 4 $\eta_{\rm cr}, 14$ $\eta_{\rm nl}, \, 14$ $S_{ts}, 153$ $\varepsilon_a, 193$ $c_{\rm sil}, 152, 153$ d, 4m, 4Differential operators $\Delta_{\infty}, 48$ $\Delta_{g_r}, 4$ $\mathcal{A}_{\rm ts},\,48$ Eigenfunctions $\varphi_{\iota}, 371$ Eigenvalues $\nu(\iota), 371$ $\nu_{{\rm T},i}(\iota), 371$ $\nu_{\rm T}(\iota), 371$ $\nu_{\rm n}(\iota), 61$ $\nu_{r,i_r}(\iota), 371$ $\bar{\nu}_q(\iota), 78$ Energies $\mathcal{E}_s(\iota, t), 112$ $\mathfrak{E}_{\text{bas}}, 14$ $\mathfrak{E}_{hom}, 14$ $\mathfrak{E}_{s}, 372$ Exponential rates $\beta_X, 21$ $\bar{\beta}_q, 78$

Fourier coefficients

INDEX

 $\hat{f}(\iota, t), 372$  $\hat{u}(\iota), 371$  $z(\iota, t), 372$ Frequencies  $\iota, 371$ Frequency sets  $I_{\rm B}, 371$  $\mathcal{I}^{-}_{\mathrm{B,n}}, \, 66$  $\mathcal{I}_{B,n}^+, 66$  $\mathcal{I}_{B,sil}, 47, 151$  $\mathcal{I}_{B,ts}, 47, 151$ Function spaces  $G^{s}, 23$  $H^{s}(M), 370$  $W_{n,\pm}(\bar{M},V), \, 68$  $W_{\rm n,\pm}^{\pm}(\bar{M},V),\,68$  $W_{\rm n}(\bar{M}, V), \, 68$  $\mathfrak{G}^{s}_{\mathrm{loc},\mathrm{u}}(I,\mathbb{T}^{d},\mathbb{C}^{m}),\,23$ Index  $\operatorname{Ind}_{\Gamma}(z), 126$ Manifolds M, 4 $M_r, 4$  $\overline{M}, 4$  $\overline{M}_{sil}, 45$  $M_{\rm sub}, 60$  $\overline{M}_{\rm ts}, 45$  $\bar{M}_t, 4$ Matrix notation A-conjugation matrix,  $\beta$ -normalised, 99  $A_{\infty}, 19, 34$  $A(\iota), 48$  $A_{q_{\rm ts},\infty}(\iota), 80$  $\beta$ , A-decomposition of a matrix, 98  $\beta$ , A-decomposition of a matrix; first (second) block of, 98  $\beta$ , A-decomposition of a vector, 98  $\beta$ , A-decomposition of a vector; first (second) component of, 98  $\beta$ -normalised A-conjugation matrix, 99 $\delta_{\pm}, 123$  $d_{\rm max}, 34$  $d_{n,+}, 62$  $d_{q,\pm}, 79$  $\gamma_{\pm}, 123$  $\kappa$ , 318  $\kappa_{\iota}, 136$  $\kappa_{\rm max}, 34$  $\kappa_{\min}, 34$  $\kappa_{n,+}, 62$  $\kappa_{\pm}, 79$ 

 $\kappa_q, 318$  $\kappa_{q,\pm}, 79$  $\kappa_{q_{\rm ts},\iota,\pm}, 80$  $\kappa_{q_{\rm ts},{\rm ts},\pm}, 80$  $\kappa_{\rm ts,\pm}, 80$  $\kappa_{\rm sil,+}, 80$  $\kappa_{\text{tot},+}, 80$  $\mathcal{K}_{q_{\rm ts},\pm}, 80$  $\mathbf{M}_m(K), 3$  $N_{\mu}, 123$  $Q^{\pm}_{\mu}, 123 \\ R^{\pm}_{\mu}, 123$  $R'_{n,+}, 62$  $R_{n,+}^{\pm}, 62$  $R_{q,\pm}^{\pm}, 79$ RspA, 34SpA, 34  $U_{\pm}, 123$  $W_{\pm}, 123$  $W_{\pm,\mp}, 123$ Measures  $\mu_{\rm B}, 14$ Metrics N, Lapse function, 5  $\bar{g}, 27$  $\bar{g}^{\sharp}, \, 45$  $\bar{h}, 27$  $a_r, 4$ g, 4, 5, 27 $g^{00}, 4$  $g^{0l}, 4$  $g^{jl}, 4$  $g_r, 4$ Norms  $\|\cdot\|_{(s)}, 371$  $\|\cdot\|_{A}, 96$  $\|\cdot\|_{\mathrm{ts},s}, 50, 157$  $\|\cdot\|_{t,s}, \, 50, \, 157$  $\|\cdot\|_{t,s,\mu_0,\pm},\,157$  $|\bar{k}|_{\bar{q}}, 29$  $\|\cdot\|_{A,s}, 35$  $|\mathcal{X}|_{\bar{h}}, 29$  $\|\cdot\|_{\text{lma}}, 136$ Projected functions  $F_{\rm ts}, 48$  $V_{\rm ts}, \, 48$  $f_{\rm n}, \, 63$  $f_{\rm ts}, \, 48$ Projections  $\Pi_{\rm sil}, 152$  $\Pi_{\rm ts}, 152$ Second fundamental forms  $\bar{k}, 28$ 

408

 $\bar{k}^{\sharp}, 45$ Tangent and cotangent spaces  $T_{\rm n}^* \bar{M}, \, 60$  $T^*_{\rm sil}\bar{M}, 45$  $T_{\rm sub}^* \bar{M}, 60$  $T_{\rm ts}^* \bar{M}, 45$ Vector fields U, 5 $\chi, 5$  $\dot{\chi}, 28$  $\mathcal{X}, 29$ Vector spaces  $E_{\iota,\beta}^{\infty}, 136$  $E_{B,J}^{''}, 36$  $\beta$ , A-decomposition of  $\mathbb{C}^k$ , 97  $\beta$ , A-decomposition of  $\mathbb{C}^k$ ; dimension of first (second) subspace of, 97  $\beta$ , A-decomposition of  $\mathbb{C}^k$ ; first (second) generalised eigenspace, 98 Asymptotically diagonal equations, 18 Asymptotically negligible shift vector field, 46 Asymptotically non-degenerate equation, 78 Auxiliary diagonal coefficients, 305 Balanced strongly, 249 Basic energy, 14 Bianchi type I spacetimes, 6 Big bang asymptotics, 9 Big crunch asymptotics, 9 BKL conjecture, 6 Canonical local coordinates, 378 Canonical separable cosmological model manifold. 11 Coefficients auxiliary diagonal, 305 diagonal, 305 main. 305 off-diagonal, 305 Convergent dominant coefficients, 60, 61, 249 main coefficients, 78 Convergent and unbalanced ODE. 214 Crude exponential bound, 14 Cycle, 126

de Sitter space, 7 Diagonal coefficients, 305 Diagonally convergent second fundamental form, 77 Diagonally dominated, balanced and convergent equation, 306 Direction generalised, 248 Dominant coefficients convergent, 60, 61, 249 Dominant noisy spatial direction, 248 geometric, 60 Dominant noisy spatial generalised direction, 248 Dominant noisy spatial torus direction, 248 Eigenspaces Laplace-Beltrami operator, 370 Eigenvalues Laplace-Beltrami operator, 370 Equation  $C^1$ -silent, 34  $C^2$ -balanced with a geometric dominant noisy spatial direction, convergent dominant coefficients and a negligible shift vector field, 61  $C^k$ -balanced, 30 asymptotically non-degenerate, 78 diagonally dominated, balanced and convergent, 306 geometrically non-degenerate, diagonally dominated, balanced and convergent, 78 non-degenerate, diagonally dominated, balanced and convergent, 318 oscillation adapted, 187 subdominant part, 58 weakly balanced, 109 weakly convergent, 109 weakly silent, 109 weakly transparent, 152 Era noisy, 312 non-oscillatory, 95 ODE, 95 silent, 312 transparent, 312 Expansion normalised second fundamental form, 9 Exponential bound, 14 crude, 14

First generalised eigenspace, 36 Flat Kasner solution, 12 Frequency era, 312 Function  $E_{\rm ts}$ -adapted, 51 Future bounded geometry of a spacetime, 10 convergent spacetime, 10 expanding separable cosmological model manifold, 9 Future uniformly timelike shift vector field, 28 Generalised direction, 248 Jordan block, 99 Generalised eigenspace, 35 corresponding to an eigenvalue, 35 corresponding to an interval, 36 first, 36 Geometric dominant noisy spatial direction, 60 Gevrey class. 23 function, 23 Growth super exponential, 16 High frequency silent sector, 344 transparent sector, 344 Initial data transparent, 44 Jordan block, 372normal form, 372 Jordan block generalised, 99 Kasner flat solution, 12 relations, 6 solutions, 6 Laplace-Beltrami operator, 369 Eigenspaces, 370 Eigenvalues, 370 Lapse function, 5 Limit equation, 17 asymptotically diagonal setting, 75 noisy setting, 58, 62 transparent setting, 47, 156

Limiting Laplace-Beltrami operator, 156 Local coordinates canonical, 378 Low frequency silent sector, 344 transparent sector, 344 Main coefficients, 305 Main coefficients convergent, 78 Manifold silent, 45 transparent, 45 Margin **ODE**, 97 Mean curvature, 5 Milne model, 7 Minkowski space, 5 Model metric, 12 noisy, 13 silent, 13 transparent, 13 Nariai spacetimes, 7 Negatively oriented solution, 65 Negligible shift vector field, 77, 249 Noisy equations, 17 era, 312 model metric, 13 sector, 344 variables, 17 Noisy setting limit equations, 62 Non-degenerate, diagonally dominated, balanced and convergent equation, 318 Non-oscillatory era, 95 ODE convergent and unbalanced, 214 era, 95 margin, 97 Off-diagonal coefficients, 305 Oscillation adapted equation, 187 Past expanding separable cosmological model manifold, 9

#### 410

#### INDEX

Path, 126 Polarised  $\mathbb{T}^3$ -Gowdy symmetric solutions, 8 Positively oriented solution, 65 Raychaudhuri equation, 10 Real  $\beta$ -normalised Jordan decomposition of a matrix, 99 Real eigenvalue spread, 34 Rotation matrix, 98 associated with a matrix, 98 Second fundamental form  $C^k$ -future bounded, 28 diagonally convergent, 77 expansion normalised, 9 Sector noisy, 344 silent, 344 transparent, 344 Separable cosmological model manifold, 5 future expanding, 9 past expanding, 9 Shift vector field, 5  $C^k$ -future bounded, 28 asymptotically negligible, 46 future uniformly timelike, 28 negligible, 77, 249 Silent equations, 17 era, 312 manifold, 45 model metric, 13 sector, 344 subset of  $\{1, ..., d\}, 45$ subset of  $\{1, ..., R\}, 45$ subset of  $T\overline{M}$ , 45 subset of  $T^*\overline{M}$ , 45 variables, 17, 44, 151 Silent sector high frequency, 344 low frequency, 344 Solution negatively oriented, 65 positively oriented, 65 transparent part, 44 Spectrum of the Riemannian manifold corresponding to the dominant noisy spatial direction, 248 Static solutions, 9

Strongly balanced, 249 Subdominant part of equation, 58 variables, 17 Subdominant block non-degeneracy, 80 Subset silent, 45 transparent, 45 Super exponential growth, 16 Transparent equations, 17 era, 312 initial data, 44 limit equation, 47 manifold, 45 model metric, 13 part of solution, 44 sector, 344 subset of  $\{1, ..., d\}, 45$ subset of  $\{1, ..., R\}, 45$ subset of  $T\overline{M}$ , 45 subset of  $T^*\overline{M}$ , 45 variables, 17, 44, 151 Transparent sector high frequency, 344 low frequency, 344  $\mathbb{T}^3$ -Gowdy symmetric solutions, 8  $\mathbb{T}^2$ -symmetric setting, 8 U(1)-symmetry, 8 Variables silent, 44, 151 transparent, 44, 151 Volume of  $\overline{M}_t$ , 5 Weakly balanced equation, 109 Weakly convergent equation, 109 Weakly silent equation, 109 Weakly transparent equation, 152 Weakly transparent equation X-degenerate, 122 X-non-trivial, 122