SAÏD ASSERDA

ABSTRACT. For a smooth strictly plurisubharmonic function u on a open set $\Omega \subset \mathbb{C}^n$ and $F \neq C^1$ nondecreasing function on \mathbb{R}^*_+ , we investigate the complex partial differential equations

 $\Delta_g \log \det(u_{i\bar{j}}) = F(\det(u_{i\bar{j}})) \|\nabla_g \log \det(u_{i\bar{j}})\|_g^2,$

where Δ_g , $\|.\|_g$ and ∇_g are the Laplacian, tensor norm and the Levi-Civita connexion, respectively, with respect to the Kähler metric $g = \partial \bar{\partial} u$. We show that the above PDE's has a Bernstein property, i.e $\det(u_{i\bar{j}}) = \text{constant on } \Omega$, provided that g is complete, the Ricci curvature of g is bounded below and Fsatisfies $\inf_{t \in \mathbf{R}^+} (2tF'(t) + \frac{F(t)^2}{n}) > \frac{1}{4}$ and $F(\max_{B(R)} \det u_{i\bar{j}}) = o(R)$.

1. INTRODUCTION

In this note, we investigate a class of fourth order complex nonlinear partial differential equations for a strictly plurisubharmonic function u on a open set $\Omega \subset \mathbb{C}^n$:

$$\Delta_g \log \det(u_{i\bar{j}}) = F(\det(u_{i\bar{j}})) \|\nabla_g \log \det(u_{i\bar{j}})\|_g^2 \tag{1.1}$$

where Δ_g , $\|.\|_g$ and ∇_g are the Laplacian, tensor norm and the Levi-Civita connexion, respectively, with respect to the Kähler metric $g = \partial \bar{\partial} u$. For $n \geq 2$, the equation (1.1) is the Euler-Lagrange equation of the variational problem of the functional

$$\mathcal{V}_{\Phi}(u) = \int_{\Omega} \Phi(\det(u_{i\bar{j}})) dV_e \tag{1.2}$$

for some $\Phi \in C^4(\mathbf{R}^*_+)$. If $\Phi(t) = t^{1-\beta}$, $\beta \neq 1$, the Euler-Lagrange equation of \mathcal{V}_{Φ} is

$$\Delta_g \log \det(u_{i\bar{j}}) = \beta \|\nabla_g \log \det(u_{i\bar{j}})\|_g^2 \tag{1.3}$$

In [1], Chen and Li showed the following Bernstein property of the solution of (3)

Theorem 1.1. Let $\Omega \subset \mathbb{C}^n$ and $u \in C^2(\Omega)$ a strictly plurisubharmonic solution of (3). Assume that

(i) (Ω, g) is complete, (ii) $Ric(g) \ge -C_1$ $(K \ge 0)$, (ii) $\det(u_{i\bar{j}}) < C_2$ where $C - C \ge 0$ are real constants. If $\beta \ge \sqrt{9\pi}$

where $C_1, C_2 > 0$ are real constants. If $\beta > \frac{\sqrt{9n^2 + 4n} + 3n}{4}$ then $\det(u_{i\bar{i}}) = const \text{ on } \Omega.$

Date: September 24, 2018.

2010 Mathematics Subject Classification. Primary 58E20, Secondary 53C21. Key words and phrases. Ricci curvature, Bernstein property.

For more details about Bernstein property, one can see the paper [1] and references therein. In this note, we will show a Bernstein property of the solutions of (1.1). Our main result is the following.

Theorem 1.2. Let $\Omega \subset \mathbb{C}^n$ be an open set and $u \in C^2(\Omega)$ a strictly plurisubharmonic solution of (1). Assume that

(i) (Ω, g) is complete,

(ii) $Ric(g) \ge -K \ (K \ge 0),$

(ii) $F \in C^1(]0, +\infty[)$ nondecreasing satisfying

$$\inf_{t>0} \left(2tF'(t) + \frac{F(t)^2}{n} \right) > \frac{1}{4} \quad and \quad F(\max_{B(R)} \det(u_{i\bar{j}})) = o(R) \tag{1.4}$$

where $B(R) = \{z \in \Omega, d_g(z, z_0) < R\}$. Then $det(u_{i\bar{j}})$ is constant on Ω .

If $F(t) = \beta$, the following corollary extends theorem (1.1).

Corollary 1.3. Let $\Omega \subset \mathbb{C}^n$ be an open set and $u \in C^2(\Omega)$ a strictly plurisubharmonic solution of (3). Assume that (Ω, g) is complete and $\operatorname{Ric}(g) \geq -K$ $(K \geq 0)$. If $|\beta| > \frac{\sqrt{n}}{2}$ then $\det(u_{i\overline{i}})$ is constant on Ω .

Example. Example of function F which satisfies conditions of Theorem 1.2 : 1- $F(t) = \log^{2\alpha} t$ where $\alpha \in]\alpha_0(n), \frac{1}{2}], \quad \alpha_0(n) > 0.$ 2- $F(t) = (\log t) \log(\alpha + (\log t)^2)$ where $\alpha > e^{\frac{1}{4}}.$ 3- $F(t) = f(\log t)$ where f is a solution of the Riccati equation $2y' + \frac{y^2}{n} = \alpha, \ \alpha > \frac{1}{4}.$

2. Proof of theorem 1.2

Proof. We denote $\Psi = |\nabla U|^2$ where $U = \log \det(u_{i\bar{j}})$ and $f(t) = F(e^t)$. By Bochner formulae

$$\Delta_g \Psi = 2f'(U)\Psi^2 + f(U)(\langle \nabla_g \Psi, \nabla_g U \rangle + \langle \nabla_g U, \nabla_g \Psi \rangle) + \sum u^{i\bar{j}} u^{k\bar{l}} U_{i\bar{l}} U_{\bar{j}\bar{l}} + \sum u^{i\bar{j}} u^{k\bar{l}} U_{i\bar{l}} U_{\bar{j}k} + \sum u^{i\bar{j}} u^{m\bar{s}} U_{i\bar{l}} U_m U_{\bar{j}} R_{i\bar{s}}$$
(2.1)

where $(u^{i\bar{j}})$ is the inverse matrix of $(u_{i\bar{j}})$ and $R_{i\bar{s}}$ is the Ricci curvature of g

$$R_{i\bar{s}} = -\frac{\partial^2 \log \det(u_{k\bar{l}})}{\partial z_i \partial \bar{z}_s} = \frac{\partial^2 \log U}{\partial z_i \partial \bar{z}_s}$$

Let $z \in \Omega$ fixed, we can choose a normal coordinates (z_1, \dots, z_n) at z such that

$$u_{i\bar{j}} = \delta_{ij}, \quad U_1 = U_{\bar{1}}, \quad U_i = U_{\bar{i}} = 0, \quad \forall i > 1$$

At z the formulae (2.1) become

$$\frac{\Delta_g \Psi}{\Psi} = 2f'(U)\Psi + (f(U) - \frac{1}{2})(\frac{\Psi_1}{\Psi}U_{\bar{1}} + U_1\frac{\Psi_{\bar{1}}}{\Psi}) + \frac{1}{2}(\frac{\Psi_1}{\Psi}U_{\bar{1}} + U_1\frac{\Psi_{\bar{1}}}{\Psi}) \\ + \frac{\sum U_{ik}U_{\bar{j}\bar{k}}}{\Psi} + \frac{\sum U_{i\bar{k}}U_{\bar{i}k}}{\Psi} - \frac{U_{1\bar{1}}U_1U_{\bar{1}}}{\Psi}$$
(2.2)

Since $\sum U_{ik}U_{\bar{i}\bar{k}} \ge U_{11}U_{\bar{1}\bar{1}}$ and

$$\frac{\frac{1}{2}(\frac{\Psi_{1}}{\Psi}U_{\bar{1}} + U_{1}\frac{\Psi_{\bar{1}}}{\Psi}) - \frac{U_{1\bar{1}}U_{1}U_{\bar{1}}}{\Psi} = \frac{1}{2\Psi}2\Re(U_{11}U_{\bar{1}}U_{\bar{1}}) \\
\geq \frac{U_{11}U_{\bar{1}\bar{1}}}{\Psi} - \frac{1}{4}\Psi$$

we obtain

$$\frac{1}{2}\left(\frac{\Psi_{1}}{\Psi}U_{\bar{1}} + U_{1}\frac{\Psi_{\bar{1}}}{\Psi}\right) - \frac{U_{1\bar{1}}U_{1}U_{\bar{1}}}{\Psi} + \frac{\sum U_{ik}U_{\bar{i}\bar{k}}}{\Psi} \ge -\frac{1}{4}\Psi$$
(2.3)

Let G be the function defined on $B(R) = B_g(z_0, R)$ by

$$G(z) = (R^2 - r(z)^2)^2 \Psi$$

where $r(z) = d_g(z, z_0)$. Since G vanishes at the boundary $\partial B(R)$, the maximum of G is attained on $w \in B(R)$. By maximum principle

$$\nabla_g G(w) = 0 \Delta_g G(w) \leq 0$$

A computation at w gives

$$\frac{\Psi_i}{\Psi} = \frac{4rr_i}{R^2 - r^2}$$
(2.4)
$$\Delta_g G = 2\Psi \|\nabla_g (R^2 - r^2)\|^2 + 2(R^2 - r^2)\Psi \Delta_g (R^2 - r^2) + 2(R^2 - r^2)\sum_{i}(R^2 - r^2)_i\Psi_i + (R^2 - r^2)^2\Delta_g\Psi \\
= 2\Psi r^2 \|\nabla_g r\|^2 - 2\Psi (R^2 - r^2)\Delta_g r^2 + (R^2 - r^2)^2\Delta_g\Psi \\
-4r(R^2 - r^2)\sum_i r_i\Psi_i \le 0$$
(2.5)

Since $\|\nabla_g r\|_g = 1$, (2.4) and (2.5) implies

$$\frac{\Delta_g \Psi}{\Psi} \le \frac{2\Delta_g r^2}{R^2 - r^2} + \frac{14r^2}{(R^2 - r^2)^2}$$
(2.6)

Let $\eta > \text{fixed such that } \inf_{t \in \mathbb{R}} \left(2f'(t) + \frac{f(t)^2}{n} \right) > 2\eta + \frac{1}{4}$. By Shwarz inequality

$$(f(U) - \frac{1}{2})(\frac{\Psi_1}{\Psi}U_{\bar{1}} + U_1\frac{\Psi_{\bar{1}}}{\Psi}) = 2\Re(U_1\frac{4rr_1(f(U) - \frac{1}{2})}{R^2 - r^2})$$

$$\geq -\eta\Psi - \frac{16(f(U) - \frac{1}{2})^2r^2}{\eta(^2 - r^2)^2}$$
(2.7)

Since $\Delta U = f(U)$ we have

$$\frac{\sum U_{i\bar{k}}U_{\bar{i}k}}{\Psi} \ge \frac{\sum (U_{i\bar{i}})^2}{\Psi} \ge \frac{(\sum U_{i\bar{i}})^2}{n\Psi} = \frac{(\Delta U)^2}{n\Psi} = \frac{f(U))^2}{n}\Psi$$
(2.8)

The inequalities (2.2), (2.7) and (2.8) give

$$\frac{\Delta_g \Psi}{\Psi} \ge \eta \Psi - \frac{16(f(U) - \frac{1}{2})^2 r^2}{\eta (R^2 - r^2)^2}$$
(2.9)

By Laplacian comparison theorem [3]: $\Delta_g r^2 \leq 2n + 2(n-1)\sqrt{Kr}$. Since $|f(U) - \frac{1}{2}|^2 \leq 2f(U)^2 + \frac{1}{2}$ and f is nondecreasing, the inequalities (2.6) and (2.9) give on B(R)

$$(R^2 - r^2)^2 \Psi \le G(w) \le C_1 (1+R)R^2 + C_2 (f(\max_{B(R)} U))^2 R^2$$

Hence if $z \in B(R/2)$, we have

$$\Psi(z) \le C_1 \frac{1+R}{R^2} + C_2 \left(\frac{f(\max_{B(R)} U)}{R}\right)^2$$

Since (Ω, g) is complete, letting R to infinity, we obtain $\Psi(z) = \|\nabla_g U(z)\|^2 = 0$ i.e det $(u_{i\bar{j}}) = C$ on Ω .

References

- Chen,G, Sheng,L, A Bernstein property of a class of fourth order complex partial differential equations. Results.Math. 58 (2010), 81-92.
- Chen,G, Sheng,L, Erratum to A Bernstein property of a class of fourth order complex partial differential equations. Results.Math. 58 (2010)
- [3] Greene, R.E., Wu, H, Faction theory on manifolds which posses a pole. Lecture Notes In Math. 699 (1979).

Ibn tofail university , faculty of sciences, department of mathematics, PO 242 Kenitra Morroco

E-mail address: asserda-said@univ-ibntofail.ac.ma