GLOBAL SOLVABILITY OF THE NAVIER-STOKES EQUATIONS WITH A FREE SURFACE IN THE MAXIMAL L_p - L_q REGULARITY CLASS

HIROKAZU SAITO

ABSTRACT. We consider the motion of incompressible viscous fluids bounded above by a free surface and below by a solid surface in the N-dimensional Euclidean space for $N \geq 2$. The aim of this paper is to show the global solvability of the Naiver-Stokes equations with a free surface, describing the above-mentioned motion, in the maximal L_p - L_q regularity class. Our approach is based on the maximal L_p - L_q regularity with exponential stability for the linearized equations, and also it is proved that solutions to the original nonlinear problem are exponentially stable.

1. INTRODUCTION

This paper is concerned with the global solvability of the Navier-Stokes equations with a free surface, describing the motion of incompressible viscous fluids bounded above by a free surface and below by a solid surface in the N-dimensional Euclidean space for $N \ge 2$, in the maximal L_p - L_q regularity class (cf. [36] for the class). Such equations were mathematically treated by Beale [6] for the first time. He proved, in an L_2 -in-time and L_2 -in-space setting with the gravity, the local solvability for large initial data in [6], whereas we prove in the maximal L_p - L_q regularity class the global solvability for small initial data in the case where the gravity is not taken into account in the present paper.

The problem is stated as follows: We are given an initial domain $\Omega \subset \mathbf{R}^N$, occupied by an incompressible viscous fluid, such that

$$\Omega = \{\xi = (\xi', \xi_N) \mid \xi' = (\xi_1, \dots, \xi_{N-1}) \in \mathbf{R}^{N-1}, 0 < \xi_N < d\} \quad (d > 0),$$

as well as an initial velocity field $\mathbf{a} = \mathbf{a}(\xi) = (a_1(\xi), \dots, a_N(\xi))^{\mathsf{T}1}$ of the fluid on Ω . The symbols Γ , S denote the boundaries of Ω such that

$$\Gamma = \{\xi = (\xi', \xi_N) \mid \xi' = (\xi_1, \dots, \xi_{N-1}) \in \mathbf{R}^{N-1}, \xi_N = d\},\$$

$$S = \{\xi = (\xi', \xi_N) \mid \xi' = (\xi_1, \dots, \xi_{N-1}) \in \mathbf{R}^{N-1}, \xi_N = 0\}.$$

We wish to find for each $t \in (0, \infty)$ a transformation $\Theta = \Theta(\cdot, t) : \Omega \ni \xi \mapsto x = \Theta(\xi, t) \in \mathbf{R}^N$, a velocity field $\mathbf{v} = \mathbf{v}(x, t) = (v_1(x, t), \dots, v_N(x, t))^{\mathsf{T}}$ of the fluid, and a pressure field $\pi = \pi(x, t)$ of the fluid so that

(1.1) $\partial_t \Theta = \mathbf{v} \circ \Theta, \quad \Theta(\xi, 0) = \xi, \quad \xi \in \Omega,$

(1.2)
$$\Omega(t) = \Theta(\Omega, t), \quad \Gamma(t) = \Theta(\Gamma, t), \quad S = \Theta(S, t),$$

²⁰¹⁰ Mathematics Subject Classification. Primary: 35Q30; Secondary: 76D05.

Key words and phrases. Global solvability, Navier-Stokes equations, Free surfaces, Maximal regularity, L_p - L_q framework, Exponential stability, Infinite layers.

¹⁾ \mathbf{M}^{T} denotes the transposed \mathbf{M} .

HIROKAZU SAITO

- (1.3) $\partial_t \mathbf{v} + (\mathbf{v} \cdot \nabla) \mathbf{v} = \text{Div} \mathbf{T}(\mathbf{v}, \pi), \quad x \in \Omega(t),$
- (1.4) $\operatorname{div} \mathbf{v} = 0, \quad x \in \Omega(t),$
- (1.5) $\mathbf{T}(\mathbf{v},\pi)\mathbf{n} = -\pi_0\mathbf{n}, \quad x \in \Gamma(t),$
- $\mathbf{v} = 0, \quad x \in S,$
- (1.7) $\mathbf{v}|_{t=0} = \mathbf{a}, \quad \xi \in \Omega,$

where $\mathbf{v} \circ \Theta = (\mathbf{v} \circ \Theta)(\xi, t) = \mathbf{v}(\Theta(\xi, t), t).$

Here the density of the fluid have been set to 1; **n** is the unit outward normal to $\Gamma(t)$; the constant π_0 is the atmospheric pressure, and it is assumed in this paper that $\pi_0 = 0$ without loss of generality. The stress tensor $\mathbf{T}(\mathbf{v}, \pi)$ is given by $\mathbf{T}(\mathbf{v}, \pi) = \mu \mathbf{D}(\mathbf{v}) - \pi \mathbf{I}$, where μ is a positive constant and denotes the viscosity coefficient of the fluid; **I** is the $N \times N$ identity matrix; $\mathbf{D}(\mathbf{v}) = \nabla \mathbf{v} + (\nabla \mathbf{v})^{\mathsf{T}}$ is the doubled strain tensor. Here and subsequently, we use the following notation for differentiations: Let f = f(x), $\mathbf{g} = (g_1(x), \dots, g_N(x))^{\mathsf{T}}$, and $\mathbf{M} = (M_{ij}(x))$ be a scalar-, a vector-, and an $N \times N$ matrix-valued function on a domain of \mathbf{R}^N , respectively, and then for $\partial_j = \partial/\partial x_j$

$$\nabla f = (\partial_1 f, \dots, \partial_N f)^{\mathsf{T}}, \quad \Delta f = \sum_{j=1}^N \partial_j^2 f, \quad \Delta \mathbf{g} = (\Delta g_1, \dots, \Delta g_N)^{\mathsf{T}},$$

div $\mathbf{g} = \sum_{j=1}^N \partial_j g_j, \quad \nabla^2 \mathbf{g} = \{\partial_i \partial_j g_k \mid i, j, k = 1, \dots, N\},$
$$\nabla \mathbf{g} = \begin{pmatrix} \partial_1 g_1 & \dots & \partial_N g_1 \\ \vdots & \ddots & \vdots \\ \partial_1 g_N & \dots & \partial_N g_N \end{pmatrix}, \quad (\mathbf{g} \cdot \nabla) \mathbf{g} = \left(\sum_{j=1}^N g_j \partial_j g_1, \dots, \sum_{j=1}^N g_j \partial_j g_N\right)^{\mathsf{T}},$$

Div $\mathbf{M} = \left(\sum_{j=1}^N \partial_j M_{1j}, \dots, \sum_{j=1}^N \partial_j M_{Nj}\right)^{\mathsf{T}}.$

Let $\mathbf{u}(\xi, t) = (\mathbf{v} \circ \Theta)(\xi, t)$, which is the so-called *Lagrangian velocity*, and then the solution Θ to (1.1) is represented as

(1.8)
$$\Theta(\xi,t) = \xi + \int_0^t \mathbf{u}(\xi,s) \, ds \quad (\xi \in \Omega, \, t > 0).$$

We now write the equations (1.3)-(1.7) in the Lagrangian formulation. Thus our unknowns will be the Lagrangian velocity $\mathbf{u}(\xi,t) = \mathbf{v}(\Theta(\xi,t),t)$ and pressure $\mathfrak{p}(\xi,t) = \pi(\Theta(\xi,t),t)$ for $(\xi,t) \in \Omega \times (0,\infty)$. If we substitute the new unknowns $\mathbf{u} = \mathbf{u}(\xi,t)$ and $\mathfrak{p} = \mathfrak{p}(\xi,t)$ in (1.3)-(1.7), then the equations turn into²

- (1.9) $\partial_t \mathbf{u} \operatorname{Div} \mathbf{T}(\mathbf{u}, \mathbf{p}) = \mathbf{F}(\mathbf{u}) \quad \text{in } \Omega, \, t > 0,$
- (1.10) $\operatorname{div} \mathbf{u} = G(\mathbf{u}) = \operatorname{div} \mathbf{G}(\mathbf{u}) \quad \text{in } \Omega, \, t > 0,$
- (1.11) $\mathbf{T}(\mathbf{u}, \mathbf{p})\mathbf{e}_N = \mathbf{H}(\mathbf{u})\mathbf{e}_N \quad \text{on } \Gamma, t > 0,$
- (1.12) $\mathbf{u} = 0$ on S, t > 0,
- (1.13) $\mathbf{u}|_{t=0} = \mathbf{a} \quad \text{in } \Omega,$

²⁾The derivation of (1.9)-(1.11) is discussed in the appendix.

where $\mathbf{e}_N = (0, \dots, 0, 1)^{\mathsf{T}}$. Here $\mathbf{F}(\mathbf{u})$, $G(\mathbf{u})$, $\mathbf{G}(\mathbf{u})$, and $\mathbf{H}(\mathbf{u})$ are nonlinear terms, with respect to \mathbf{u} , of the forms:

$$(1.14) \quad \mathbf{F}(\mathbf{u}) = \mathbf{U}_{1} \left(\int_{0}^{t} \nabla \mathbf{u}(\xi, s) \, ds \right) \partial_{t} \mathbf{u} + \mathbf{V} \left(\int_{0}^{t} \nabla \mathbf{u}(\xi, s) \, ds \right) \nabla^{2} \mathbf{u} \\ + \left[\mathbf{W} \left(\int_{0}^{t} \nabla \mathbf{u}(\xi, s) \, ds \right) \int_{0}^{t} \nabla^{2} \mathbf{u}(\xi, s) \, ds \right] \nabla \mathbf{u}, \\ G(\mathbf{u}) = \mathbf{U}_{2} \left(\int_{0}^{t} \nabla \mathbf{u}(\xi, s) \, ds \right) : \nabla \mathbf{u}, \quad \mathbf{G}(\mathbf{u}) = \mathbf{U}_{3} \left(\int_{0}^{t} \nabla \mathbf{u}(\xi, s) \, ds \right) \mathbf{u} \\ \mathbf{H}(\mathbf{u}) = \mathbf{D}(\mathbf{u}) \mathbf{U}_{4} \left(\int_{0}^{t} \nabla \mathbf{u}(\xi, s) \, ds \right) + (\nabla \mathbf{u})^{\mathsf{T}} \mathbf{U}_{5} \left(\int_{0}^{t} \nabla \mathbf{u}(\xi, s) \, ds \right) \\ + \mathbf{U}_{6} \left(\int_{0}^{t} \nabla \mathbf{u}(\xi, s) \, ds \right) (\nabla \mathbf{u})^{\mathsf{T}} \left(\mathbf{I} + \mathbf{U}_{7} \left(\int_{0}^{t} \nabla \mathbf{u}(\xi, s) \, ds \right) \right),$$
where $\mathbf{U} : \mathbf{R}^{N \times N} \Rightarrow \mathbf{R}^{N \times N} (i = 1, ..., 7)$

where $\mathbf{U}_i: \mathbf{R}^{N \times N} \to \mathbf{R}^{N \times N}$ $(i = 1, \dots, 7),$

$$\mathbf{V}(\cdot)\nabla^{2}\mathbf{u} = \left(\sum_{i,j,k=1}^{N} V_{ijk}^{1}(\cdot)\partial_{i}\partial_{j}u_{k}, \dots, \sum_{i,j,k=1}^{N} V_{ijk}^{N}(\cdot)\partial_{i}\partial_{j}u_{k}\right)^{\mathsf{T}},$$
$$\left[\mathbf{W}(\cdot)\int_{0}^{t}\nabla^{2}\mathbf{u}\,ds\right]\nabla\mathbf{v} = \left(\sum_{i,j,k,l,m=1}^{N} W_{ijklm}^{1}(\cdot)\int_{0}^{t}\partial_{i}\partial_{j}u_{k}\,ds\,\partial_{l}v_{m}, \dots, \sum_{i,j,k,l,m=1}^{N} W_{ijklm}^{N}(\cdot)\int_{0}^{t}\partial_{i}\partial_{j}u_{k}\,ds\,\partial_{l}v_{m}\right)^{\mathsf{T}},$$

for some $V_{ijk}^1, \ldots, V_{ijk}^N, W_{ijklm}^1, \ldots, W_{ijklm}^N : \mathbf{R}^{N \times N} \to \mathbf{R}$ and for any *N*-vectors $\mathbf{u} = (u_1, \ldots, u_N)^{\mathsf{T}}, \mathbf{v} = (v_1, \ldots, v_N)^{\mathsf{T}}$. Note that, for $N \times N$ matrices $\mathbf{A} = (A_{ij}), \mathbf{B} = (B_{ij})$, we have set

$$\mathbf{A}: \mathbf{B} = \sum_{i,j=1}^{N} A_{ij} B_{ij}.$$

One has the following information about \mathbf{U}_i , \mathbf{V} , \mathbf{W} : Let $\mathbf{X} = (X_{mn})$ be $N \times N$ matrices. Then all the components of $\mathbf{U}_i(\mathbf{X})$ (i = 1, ..., 7), $\mathbf{V}(\mathbf{X})$, and $\mathbf{W}(\mathbf{X})$ are polynomials with respect to X_{mn} for m, n = 1, ..., N. Furthermore,

(1.15)
$$\mathbf{U}_i(\mathbf{O}) = \mathbf{O} \quad (i = 1, \dots, 7), \quad V_{ijk}^l(\mathbf{O}) = 0 \quad (i, j, k, l = 1, \dots, N),$$

where ${\bf O}$ denotes the $N\times N$ zero matrix.

Let us introduce historical remarks and key ideas of the present paper at this point. If we consider free boundary problems, then we first usually transform them to nonlinear problems on given domains, independent of time t, by using a suitable transformation. Roughly speaking, such transformations are divided into

- Lagrangian transformation;
- Eulerian approaches (e.g. Beale's transformation in [7], [8]; Hanzawa's transformation in [19]).

Lagrangian transformation denotes the transformation Θ of (1.8). It is quite useful to show the local solvability for a lot of situations. In fact, by using the Lagrangian transformation, Shibata [36] and Enomoto et al. [17] proved, respectively, the local solvability of the Navier-Stokes equations with a free surface for

HIROKAZU SAITO

incompressible viscous fluids and for compressible viscous fluids in the case where the initial domain Ω has uniform $W_q^{2^{-1/q}}$ regularity (cf. their papers for the definition). Here half-spaces, bent half-spaces, layer-like domains, cylinder-like domains, bounded domains, and exterior domains are typical examples of uniform $W_q^{2^{-1/q}}$ domains. As for the local solvability with the Lagrangian transformation, we also refer e.g. to the following papers: Solonnikov [42, 43] and references therein, Mogilevskii and Solonnikov [28], Mucha and Zajączkowski [29], Shibata and Shimizu [38] for incompressible viscous fluids in bounded domains; Beale [6], Allain [4], Tani [50], Abels [1] for incompressible viscous fluids in layer-like domains; Tani [48], Solonnikov and Tani [44, 45], Secchi and Valli [35], Secchi [32, 33, 34], Zajączkowski [55, 56, 57], Zadrzyńska and Zajączkowski [52, 53, 54], Ströhmer and Zajączkowski [46], Denisova and Solonnikov [15] for compressible viscous fluids in layer-like domains; Tani [49], Denisova [9, 10, 11, 12, 13], Denisova and Solonnikov [14] for the motion of two fluids separated by a closed free surface.

The advantage of the Lagrangian transformation for the local solvability is that $\int_0^t \nabla \mathbf{u}(\xi, s) \, ds$ appears in nonlinear terms (cf. (1.14)). By choosing T > 0 small enough, we can see $\int_0^t \nabla \mathbf{u}(\xi, s) \, ds$ $(t \in (0, T))$ as a small coefficient, and thus the nonlinear terms would be small with suitable norms. This enables us to show the local solvability by using the contraction mapping theorem.

In order to prove the global solvability, we need the integrability of $\nabla \mathbf{u}(\xi, t)$ with respect to time $t \in (0, \infty)$ because of $\int_0^t \nabla \mathbf{u}(\xi, s) ds$. If we are in an L_p -intime and L_q -in-space setting, then a key idea to guarantee the time integrability is exponential stability of solutions for the linearized equations as was seen e.g. in Shibata-Shimizu [39], Shibata [36]. These two papers tell us that the exponential stability can be proved for bounded domains by some abstract approach. However, for unbounded domains containing Ω of the present paper, it is not true in general that the exponential stability holds. This is one of main difficulties to prove the global solvability in our situation.

Eulerian approaches are useful to show the large-time behavior of solutions to the Navier-Stokes equations with a free surface, because $\int_0^t \mathbf{u}(\xi, s) \, ds$ as above does not appear. We here introduce e.g. the following papers as references of the Eulerian approaches: Beale and Nishida [8] (cf. also [20] for the detailed proof), Hataya and Kawashima [21] proved polynomial decay of solutions for layer-like domains in an L_2 -in-time and L_2 -in-space setting; Köhne, Prüss, and Wilke [22], Solonnikov [41] proved exponential stability of solutions for bounded domains in an L_p -in-time and L_p -in-time and Shibata [31] showed L_q - L_r estimates of the Stokes semigroup with surface tension and gravity on the half-space.

The key idea of this paper is to prove the maximal L_p - L_q regularity with exponential stability for the linearized equations of (1.9)-(1.13). As mentioned above, we can prove such an exponential stability for bounded domains, but the technique can not be applied to our situation because our domain Ω is unbounded. To overcome this difficulty, we make use of Abels [2] and Saito [30] in this paper. We then apply the maximal L_p - L_q regularity with exponential stability and the contraction mapping theorem to (1.9)-(1.13) in order to prove their global solvability.

This paper is organized as follows: The next section first introduce notation and function spaces that are used throughout this paper. Secondly, we state main results of the global solvability for (1.9)-(1.13) and the maximal L_p - L_q regularity with exponential stability for the linearized equations. Section 3 introduces some results concerning a time-shifted problem, a variational problem, and the Helmholtz decomposition on Ω . Section 4 shows the generation of an analytic C_0 -semigroup associated with the linearized equations. In Section 5, we prove the maximal L_p - L_q regularity with exponential stability for the linearized equations by means of results introduced in Sections 3, 4. Section 6 proves the global solvability of (1.9)-(1.13) based on the contraction mapping theorem together with the maximal L_p - L_q regularity with exponential stability proved in Section 5. In Section 7, we show the global existence and uniqueness of solutions for the original problem (1.1)-(1.7), and also their exponential stability.

2. NOTATION AND MAIN RESULTS

In this section, we first introduce notation and function spaces that are used throughout this paper. Next our main results are stated.

2.1. Notation. The set of all natural numbers, real numbers, and complex numbers are denoted by N, R, and C, respectively, and $N_0 = N \cup \{0\}$.

Let $m, n \in \mathbf{N}$ and G be a domain of \mathbf{R}^n . We set $\mathbf{a} \cdot \mathbf{b} = \sum_{j=1}^m a_j b_j$ for m-vectors $\mathbf{a} = (a_1, \dots, a_m)^\mathsf{T}$ and $\mathbf{b} = (b_1, \dots, b_m)^\mathsf{T}$, while we set $(\mathbf{f}, \mathbf{g})_G = \int_G \mathbf{f}(x) \cdot \mathbf{g}(x) \, dx = \sum_{j=1}^m \int_G f_j(x) g_j(x) \, dx$ for m-vector functions $\mathbf{f} = \mathbf{f}(x) = (f_1(x), \dots, f_m(x))^\mathsf{T}$, $\mathbf{g} = \mathbf{g}(x) = (g_1(x), \dots, g_m(x))^\mathsf{T}$ on G. In addition, $C_0^\infty(G)$ denotes the set of all C^∞ functions on \mathbf{R}^n whose supports are compact and contained in G.

Let X and Y be Banach spaces, and let $1 \leq p \leq \infty$. The Banach space of all bounded linear operators from X to Y is denoted by $\mathcal{L}(X, Y)$, and $\mathcal{L}(X) = \mathcal{L}(X, X)$. $L_p(G, X)$ and $W_p^m(G, X)$ denote, respectively, the standard X-valued Lebesgue spaces on G and the standard X-valued Sobolev spaces on G, and $W_p^0(G, X) =$ $L_p(G, X)$. If $X = \mathbf{R}$ or $X = \mathbf{C}$, then $L_p(G, X)$, $W_p^m(G, X)$, and $W_p^0(G, X)$ are denoted by $L_p(G)$, $W_p^m(G)$, and $W_p^0(G)$, respectively, for short.

The symbol C(G, X) stands for the set of all X-valued continuous function on G, while $C^m(G, X)$ is the set of all *m*-times continuously differentiable functions on G with values in X. Let BUC(G, X) be the Banach space of all X-valued uniformly continuous and bounded functions on G. In addition,

$$BUC^{m}(G,X) = \{ f \in C^{m}(G,X) \mid \partial^{\alpha} f \in BUC(G,X) \text{ for } |\alpha| = 0, 1, \dots, m \},\$$

where we have set $\partial^{\alpha} f = \partial_1^{\alpha_1} \dots \partial_n^{\alpha_n} f$ and $|\alpha| = \alpha_1 + \dots + \alpha_n$ for multi-index $\alpha = (\alpha_1, \dots, \alpha_n) \in \mathbf{N}_0^n$. Here C(G), $C^m(G)$, BUC(G), and $BUC^m(G)$ are defined similarly as above, and also G can be replaced by the closure \overline{G} of G.

Let $\gamma \in \mathbf{R}$. We then define functions spaces with exponential weights as

$$L_{p,\gamma}(\mathbf{R},X) = \{ f \in L_{p,\text{loc}}(\mathbf{R},X) \mid e^{-\gamma t} f(t) \in L_p(\mathbf{R},X) \},$$

$$W_{p,\gamma}^1(\mathbf{R},X) = \{ f \in W_{p,\text{loc}}^1(\mathbf{R},X) \mid e^{-\gamma t} \partial_t^k f(t) \in L_p(\mathbf{R},X) \text{ for } k = 0,1 \}$$

On the other hand, one sets for $\mathbf{R}_{+} = (0, \infty)$

$${}_{0}W_{p}^{1}(\mathbf{R}_{+}, X) = \{ f \in W_{p}^{1}(\mathbf{R}_{+}, X) \mid f|_{t=0} = 0 \}.$$

In order to define Bessel potential spaces of order 1/2, we introduce the Fourier transform and its inverse transform as follows: Let f = f(t) and $g = g(\tau)$ be

functions defined on \mathbf{R} , and then

$$\mathcal{F}[f](\tau) = \int_{\mathbf{R}} e^{-it\tau} f(t) \, dt, \quad \mathcal{F}_{\tau}^{-1}[g](t) = \frac{1}{2\pi} \int_{\mathbf{R}} e^{it\tau} g(\tau) \, d\tau.$$

The Bessel potential spaces are given by

$$H_p^{1/2}(\mathbf{R}, X) = \{ f \in L_p(\mathbf{R}, X) \mid ||f||_{H_p^{1/2}(\mathbf{R}, X)} < \infty \},$$

$$\|f\|_{H_p^{1/2}(\mathbf{R}, X)} = \|\mathcal{F}_{\tau}^{-1}[(1 + \tau^2)^{1/4}\mathcal{F}[f](\tau)]\|_{L_p(\mathbf{R}, X)},$$

$$H_{p, \gamma}^{1/2}(\mathbf{R}, X) = \{ f \in L_{p, \gamma}(\mathbf{R}, X) \mid ||e^{-\gamma t}f||_{H_p^{1/2}(\mathbf{R}, X)} < \infty \},$$

and furthermore,

$$H_p^{1/2}(\mathbf{R}_+, X) = [L_p(\mathbf{R}_+, X), W_p^1(\mathbf{R}_+, X)]_{1/2},$$

$${}_0H_p^{1/2}(\mathbf{R}_+, X) = [L_p(\mathbf{R}_+, X), {}_0W_p^1(\mathbf{R}_+, X)]_{1/2},$$

where $[\cdot, \cdot]_{\theta}$ is the complex interpolation functor with $0 < \theta < 1$. For notational convenience, we set for $1 < p, q < \infty$ and $Z \in \{H, {}_{0}H\}$

$$\begin{split} H^{1,1/2}_{q,p,\gamma}(\Omega \times \mathbf{R}) &= H^{1/2}_{p,\gamma}(\mathbf{R}, L_q(\Omega)) \cap L_{p,\gamma}(\mathbf{R}, W^1_q(\Omega)), \\ H^{1,1/2}_{q,p,\gamma}(\Omega \times \mathbf{R}) &= H^{1/2}_p(\mathbf{R}, L_q(\Omega)) \cap L_p(\mathbf{R}, W^1_q(\Omega)), \\ Z^{1,1/2}_{q,p}(\Omega \times \mathbf{R}_+) &= H^{1/2}_p(\mathbf{R}_+, L_q(\Omega)) \cap L_p(\mathbf{R}_+, W^1_q(\Omega)), \\ W^{2,1}_{q,p,\gamma}(\Omega \times \mathbf{R}) &= W^1_{p,\gamma}(\mathbf{R}, L_q(\Omega)) \cap L_{p,\gamma}(\mathbf{R}, W^2_q(\Omega)), \\ W^{2,1}_{q,p}(\Omega \times \mathbf{R}_+) &= W^1_p(\mathbf{R}_+, L_q(\Omega)) \cap L_p(\mathbf{R}_+, W^2_q(\Omega)). \end{split}$$

Let $1 < q < \infty$ and q' = q/(q-1). A closed subspace $W^1_{q,\Gamma}(\Omega)$ of $W^1_q(\Omega)$ is defined as $W^1_{q,\Gamma}(\Omega) = \{f \in W^1_q(\Omega) \mid f = 0 \text{ on } \Gamma\}$. Then the solenoidal space $J_q(\Omega)$ is given by

$$J_q(\Omega) = \{ \mathbf{f} \in L_q(\Omega)^N \mid (\mathbf{f}, \nabla \varphi)_\Omega = 0 \text{ for all } \varphi \in W^1_{q', \Gamma}(\Omega) \}$$

Here we set $D_{q,p}(\Omega) = (J_q(\Omega), D(A_q))_{1-1/p,p}^{3}$, where $(\cdot, \cdot)_{\theta,p}$ is the real interpolation functor with $0 < \theta < 1$.

Remark 2.1. The interpolation space $D_{q,p}(\Omega)$ is characterized as follows⁴:

$$D_{q,p}(\Omega) = \begin{cases} \{\mathbf{u} \in J_q(\Omega) \cap B_{q,p}^{2-2/p}(\Omega)^N \mid (\mu \mathbf{D}(\mathbf{u})\mathbf{e}_N)_\tau = 0 \text{ on } \Gamma, \, \mathbf{u} = 0 \text{ on } S \} \\ \text{when } 2 - 2/p > 1 + 1/q, \\ \{\mathbf{u} \in J_q(\Omega) \cap B_{q,p}^{2-2/p}(\Omega)^N \mid \mathbf{u} = 0 \text{ on } S \} \\ \text{when } 1/q < 2 - 2/p < 1 + 1/q, \\ J_q(\Omega) \cap B_{q,p}^{2-2/p}(\Omega)^N \text{ when } 2 - 2/p < 1/q, \end{cases}$$

where we have set $B_{q,p}^{2-2/p}(\Omega) = (L_q(\Omega), W_q^2(\Omega))_{1-1/p,p}$ and $\mathbf{v}_{\tau} = \mathbf{v} - \mathbf{e}_N(\mathbf{e}_N \cdot \mathbf{v})$ for any N-vector \mathbf{v} .

Throughout this paper, the letter C denotes generic constants and $C_{a,b,c,\ldots}$ means that the constant depends on the quantities a, b, c, \ldots . The values of constants C and $C_{a,b,c,\ldots}$ may change from line to line.

³⁾ $D(A_q)$ is the domain of the Stokes operator A_q associated with the linearized equations of (1.9)-(1.13). They are discussed in Section 4 below in more detail, especially in (4.5).

 $^{^{(4)}}$ We refer e.g. to [36, page 4133].

2.2. Main results. This subsection introduces our main results of this paper. First the global solvability of (1.9)-(1.13) is stated as follows:

Theorem 2.2. Let p and q be exponents satisfying

(2.1)
$$2$$

Then there exist positive constants γ_0 , δ_0 , and ε_0 such that, for any $\mathbf{a} \in D_{q,p}(\Omega)$ with $\|\mathbf{a}\|_{D_{q,p}(\Omega)} \leq \varepsilon_0$, the equations (1.9)-(1.13) admit a unique solution

$$(\mathbf{u}, \mathbf{p}) \in W^{2,1}_{q,p}(\Omega \times \mathbf{R}_+)^N \times L_p(\mathbf{R}_+, W^1_q(\Omega)),$$

with $\lim_{t\to 0+} \|\mathbf{u}(t) - \mathbf{a}\|_{B^{2-2/p}_{q,p}(\Omega)} = 0$, satisfying the estimate:

(2.2)
$$\|e^{\gamma_0 t}(\partial_t \mathbf{u}, \mathbf{u}, \nabla \mathbf{u}, \nabla^2 \mathbf{u})\|_{L_p(\mathbf{R}_+, L_q(\Omega))} + \|e^{\gamma_0 t}\theta\|_{L_p(\mathbf{R}_+, W_q^1(\Omega))} \le \delta_0.$$

Remark 2.3. We discuss the equations (1.1)-(1.7) in Section 7 below.

Next we introduce the maximal L_p - L_q regularity with exponential stability for the following linearized system associated with (1.9)-(1.13):

(2.3)
$$\begin{cases} \partial_t \mathbf{u} - \operatorname{Div} \mathbf{T}(\mathbf{u}, \mathbf{p}) = \mathbf{f} & \text{in } \Omega, \ t > 0, \\ \operatorname{div} \mathbf{u} = g & \text{in } \Omega, \ t > 0, \\ \mathbf{T}(\mathbf{u}, \mathbf{p}) \mathbf{e}_N = \mathbf{h} & \text{on } \Gamma, \ t > 0, \\ \mathbf{u} = 0 & \text{on } S, \ t > 0, \\ \mathbf{u}|_{t=0} = \mathbf{a} & \text{in } \Omega. \end{cases}$$

To this end, following [25], we introduce some function spaces related to the solvability of the divergence equation div $\mathbf{u} = g$ in Ω with boundary condition $\mathbf{u} \cdot (-\mathbf{e}_N) = 0$ on S. Let $1 < q < \infty$ and q' = q/(q-1). Noting [2, Lemma 2.3], we can regard $W^1_{q',\Gamma}(\Omega)$ as a Banach space with norm $\|\nabla \cdot\|_{L_{q'}(\Omega)}$, which is denoted by $\widehat{W}^1_{q',\Gamma}(\Omega)$. Assume that $g \in L_q(\Omega)$, and then one has

$$|(g,\varphi)_{\Omega}| \le C ||g||_{L_q(\Omega)} ||\nabla\varphi||_{L_{q'}(\Omega)} \quad \text{for any } \varphi \in \widehat{W}^1_{q',\Gamma}(\Omega),$$

with some positive constant C independent of g and φ . This inequality implies that g is an element of $\widehat{W}_{q,\Gamma}^{-1}(\Omega)$, where $\widehat{W}_{q,\Gamma}^{-1}(\Omega)$ is the dual space of $\widehat{W}_{q',\Gamma}^{1}(\Omega)$. Here we see g as a functional on $\{\nabla \varphi \mid \varphi \in \widehat{W}_{q',\Gamma}^{1}(\Omega)\} \subset L_{q'}(\Omega)^N$, which, combined with Hahn-Banach's theorem, furnishes that there is a $\mathbf{G} \in L_q(\Omega)^N$ such that

$$\|g\|_{\widehat{W}_{q,\Gamma}^{-1}(\Omega)} = \|\mathbf{G}\|_{L_q(\Omega)}, \quad (g,\varphi)_{\Omega} = -(\mathbf{G},\nabla\varphi)_{\Omega} \quad \text{for all } \varphi \in \widehat{W}_{q',\Gamma}^1(\Omega).$$

Let $[\mathbf{G}] = {\mathbf{G} + \mathbf{f} \mid \mathbf{f} \in J_q(\Omega)} \in L_q(\Omega)^N / J_q(\Omega)$. Then $L_q(\Omega) \ni g \mapsto [\mathbf{G}] \in L_q(\Omega)^N / J_q(\Omega)$ is well-defined, so that we denote $[\mathbf{G}]$ by $\mathcal{G}(g)$. One especially notes that $\|\mathcal{G}(g)\|_{L_q(\Omega)^N / J_q(\Omega)} = \|g\|_{\widehat{W}_{q,\Gamma}^{-1}(\Omega)}$. Thus, for any $g \in L_q(\Omega)$ and any representative \mathfrak{g} of $\mathcal{G}(g)$ regular enough, we have

$$(\operatorname{div} \mathfrak{g}, \varphi)_{\Omega} - (\mathfrak{g} \cdot (-\mathbf{e}_N), \varphi)_S = (g, \varphi)_{\Omega} \text{ for all } \varphi \in W^1_{q', \Gamma}(\Omega),$$

which implies that $\mathbf{u} = \mathbf{g}$ solves the divergence equation mentioned above.

Now we state the main result for (2.3) as follows:

HIROKAZU SAITO

Theorem 2.4. Let $1 < p, q < \infty$ with $2/p + 1/q \neq 1$ and $2/p + 1/q \neq 2$. Then there exists a positive constant σ_0 such that, for every \mathbf{f} , g, \mathbf{h} , and \mathbf{a} satisfying

$$e^{\sigma_0 t} \mathbf{f} \in L_p(\mathbf{R}_+, L_q(\Omega))^N, \quad e^{\sigma_0 t} g \in {}_0H^{1,1/2}_{q,p}(\mathbf{R}_+ \times \Omega) \cap {}_0W^1_p(\mathbf{R}_+, \widehat{W}^{-1}_{q,\Gamma}(\Omega)),$$
$$e^{\sigma_0 t} \mathbf{h} \in {}_0H^{1,1/2}_{q,p}(\mathbf{R}_+ \times \Omega)^N, \quad \mathbf{a} \in D_{q,p}(\Omega),$$

the system (2.3) admits a unique solution $(\mathbf{u}, \mathbf{p}) \in W^{2,1}_{q,p}(\Omega \times \mathbf{R}_+)^N \times L_p(\mathbf{R}_+, W^1_q(\Omega))$ with $\lim_{t\to 0+} \|\mathbf{u}(t) - \mathbf{a}\|_{B^{2-2/p}_{q,p}(\Omega)} = 0$. In addition, the solution (\mathbf{u}, \mathbf{p}) satisfies

$$\begin{split} \|e^{\sigma_{0}t}(\partial_{t}\mathbf{u},\mathbf{u},\nabla\mathbf{u},\nabla^{2}\mathbf{u})\|_{L_{p}(\mathbf{R}_{+},L_{q}(\Omega))} + \|e^{\sigma_{0}t}\mathfrak{p}\|_{L_{p}(\mathbf{R}_{+},W_{q}^{1}(\Omega))} \\ &\leq c_{0}\Big(\|e^{\sigma_{0}t}\mathbf{f}\|_{L_{p}(\mathbf{R}_{+},L_{q}(\Omega))} + \|e^{\sigma_{0}t}g\|_{W_{p}^{1}(\mathbf{R}_{+},\widehat{W}_{q,\Gamma}^{-1}(\Omega))} \\ &+ \|e^{\sigma_{0}t}(g,\mathbf{h})\|_{0H_{q,p}^{1,1/2}(\Omega\times\mathbf{R}_{+})} + \|\mathbf{a}\|_{D_{q,p}(\Omega)}\Big), \end{split}$$

with some positive constant $c_0 \geq 1$ depending only on N, d, p, q, μ , and σ_0 .

3. Preliminaries

This section introduces some results concerning a time-shifted problem for (2.3), a variational problem, and the Helmholtz decomposition on Ω .

3.1. A time-shifted problem. We consider in this subsection the following time-shifted linear system:

(3.1)
$$\begin{cases} \partial_t \mathbf{u} + 2\delta \mathbf{u} - \operatorname{Div} \mathbf{T}(\mathbf{u}, \mathbf{p}) = \mathbf{f} & \text{in } \Omega, \ t \in \mathbf{R}, \\ & \operatorname{div} \mathbf{u} = g & \text{in } \Omega, \ t \in \mathbf{R}, \\ & \mathbf{T}(\mathbf{u}, \mathbf{p})\mathbf{e}_N = \mathbf{h} & \text{on } \Gamma, \ t \in \mathbf{R}, \\ & \mathbf{u} = 0 & \text{on } S, \ t \in \mathbf{R}, \end{cases}$$

where δ is a positive constant. More precisely, we prove

Proposition 3.1. Let $1 < p, q < \infty$ and $\delta > 0$. Then, for every

$$\begin{split} \mathbf{f} &\in L_{p,-\delta}(\mathbf{R}, L_q(\Omega))^N, \quad g \in H^{1,1/2}_{q,p,-\delta}(\Omega \times \mathbf{R}) \cap W^1_{p,-\delta}(\mathbf{R}, \widehat{W}^{-1}_{q,\Gamma}(\Omega)), \\ \mathbf{h} &\in H^{1,1/2}_{q,p,-\delta}(\Omega \times \mathbf{R})^N, \end{split}$$

the system (3.1) admits a unique solution (\mathbf{u}, \mathbf{p}) with

$$\mathbf{u} \in W^{2,1}_{q,p,-\delta}(\Omega \times \mathbf{R})^N, \quad \mathfrak{p} \in L_{p,-\delta}(\mathbf{R}, W^1_q(\Omega)).$$

 $\label{eq:intro:intro} In \ addition, \ the \ following \ assertions \ hold \ true.$

(1) The solution (\mathbf{u}, \mathbf{p}) satisfies the estimate:

$$\begin{split} \|e^{\delta t}(\partial_{t}\mathbf{u},\mathbf{u},\nabla\mathbf{u},\nabla^{2}\mathbf{u})\|_{L_{p}(\mathbf{R},L_{q}(\Omega))}+\|e^{\delta t}\mathfrak{p}\|_{L_{p}(\mathbf{R},W_{q}^{1}(\Omega))}\\ &\leq C\Big(\|e^{\delta t}\mathbf{f}\|_{L_{p}(\mathbf{R},L_{q}(\Omega))}+\|e^{\delta t}(\partial_{t}g,g)\|_{L_{p}(\mathbf{R},\widehat{W}_{q,\Gamma}^{-1}(\Omega))}+\|e^{\delta t}(g,\mathbf{h})\|_{H^{1,1/2}_{q,p}(\Omega\times\mathbf{R})}\Big), \end{split}$$

with some positive constant $C = C_{N,d,p,q,\delta,\mu}$.

(2) If \mathbf{f} , g, and \mathbf{h} vanish for t < 0, then \mathbf{u} also vanishes for t < 0.

Remark 3.2. Proposition 3.1 (2) implies $\mathbf{u}|_{t=0} = 0$ in $L_q(\Omega)^N$, provided that \mathbf{f} , g, and \mathbf{h} vanish for t < 0.

To prove Proposition 3.1, we start with

(3.2)
$$\begin{cases} \partial_t \mathbf{U} - \operatorname{Div} \mathbf{T}(\mathbf{U}, P) = \mathbf{F} & \text{in } \Omega, t \in \mathbf{R}, \\ \operatorname{div} \mathbf{U} = G & \text{in } \Omega, t \in \mathbf{R}, \\ \mathbf{T}(\mathbf{U}, P) \mathbf{e}_N = \mathbf{H} & \text{on } \Gamma, t \in \mathbf{R}, \\ \mathbf{U} = 0 & \text{on } S, t \in \mathbf{R}. \end{cases}$$

Concerning this system, we have

Lemma 3.3. Let $1 < p, q < \infty$ and $\gamma > 0$. Then, for every

$$\begin{aligned} \mathbf{F} &\in L_{p,\gamma}(\mathbf{R}, L_q(\Omega))^N, \quad G \in H^{1,1/2}_{q,p,\gamma}(\Omega \times \mathbf{R}) \cap W^1_{p,\gamma}(\mathbf{R}, \widehat{W}^{-1}_{q,\Gamma}(\Omega)), \\ \mathbf{H} &\in H^{1,1/2}_{q,p,\gamma}(\Omega \times \mathbf{R})^N, \end{aligned}$$

the system (3.2) admits a unique solution (\mathbf{U}, P) with

$$\mathbf{U} \in W^{2,1}_{q,p,\gamma}(\Omega \times \mathbf{R})^N, \quad P \in L_{p,\gamma}(\mathbf{R}, W^1_q(\Omega)).$$

In addition, the following assertions hold true.

(1) The solution (\mathbf{U}, P) satisfies the estimate:

$$\begin{split} \|e^{-\gamma t}(\partial_{t}\mathbf{U},\mathbf{U},\nabla\mathbf{U},\nabla\mathbf{U}^{2})\|_{L_{p}(\mathbf{R},L_{q}(\Omega))} + \|e^{-\gamma t}P\|_{L_{p}(\mathbf{R},W_{q}^{1}(\Omega))} \\ &\leq C\Big(\|e^{-\gamma t}\mathbf{F}\|_{L_{p}(\mathbf{R},L_{q}(\Omega))} + \|e^{-\gamma t}(\partial_{t}G,G)\|_{L_{p}(\mathbf{R},\widehat{W}_{q,\Gamma}^{-1}(\Omega))} \\ &+ \|e^{-\gamma t}(G,\mathbf{H})\|_{H_{q,p}^{1,1/2}(\Omega\times\mathbf{R})}\Big), \end{split}$$

with some positive constant $C = C_{N,d,p,q,\gamma,\mu}$. (2) If **F**, G, and **H** vanish for t < 0, then **U** also vanishes for t < 0.

Proof. This lemma was proved in [30, Theorem 2.1].

Proof of Proposition 3.1. In order to apply Lemma 3.3 with $\gamma = \delta$ to (3.1), we set $\mathbf{F} = e^{2\delta t} \mathbf{f}$, $G = e^{2\delta t} g$, and $\mathbf{H} = e^{2\delta t} \mathbf{h}$. It then is clear that

$$\begin{aligned} \mathbf{F} &\in L_{p,\delta}(\mathbf{R}, L_q(\Omega))^N, \quad G \in H^{1,1/2}_{q,p,\delta}(\Omega \times \mathbf{R}) \cap W^1_{p,\delta}(\mathbf{R}, \widehat{W}^{-1}_{q,\Gamma}(\Omega)), \\ \mathbf{H} &\in H^{1,1/2}_{q,p,\delta}(\Omega \times \mathbf{R})^N. \end{aligned}$$

Thus, by Lemma 3.3, we have a solution $(\mathbf{U}, P) \in W^{2,1}_{q,p,\delta}(\Omega \times \mathbf{R})^N \times L_{p,\delta}(\mathbf{R}, W^1_q(\Omega))$ to (3.2), which satisfies

$$\begin{aligned} \|e^{-\delta t}(\partial_t \mathbf{U}, \mathbf{U}, \nabla \mathbf{U}, \nabla \mathbf{U}^2)\|_{L_p(\mathbf{R}, L_q(\Omega))} + \|e^{-\delta t}P\|_{L_p(\mathbf{R}, W_q^1(\Omega))} \\ &\leq C\left(\|e^{\delta t}\mathbf{f}\|_{L_p(\mathbf{R}, L_q(\Omega))} + \|e^{\delta t}(\partial_t g, g)\|_{L_p(\mathbf{R}, \widehat{W}_{q, \Gamma}^{-1}(\Omega))} + \|e^{\delta t}(g, \mathbf{h})\|_{H_{q, p}^{1, 1/2}(\Omega \times \mathbf{R})}\right) \end{aligned}$$

for some positive constant $C = C_{N,d,p,q,\delta,\mu}$. Let $(\mathbf{u}, \mathbf{p}) = (e^{-2\delta t}\mathbf{U}, e^{-2\delta t}P)$, and then (\mathbf{u}, \mathbf{p}) solves the system (3.1) and satisfies the required estimate of Proposition 3.1 (1) by the last inequality. The other assertions immediately follow from Lemma 3.3, which completes the proof of Proposition 3.1.

3.2. A variational problem. Let $1 < q < \infty$ and q' = q/(q-1). This subsection is concerned with the following variational problem:

(3.3)
$$(\nabla u, \nabla \varphi)_{\Omega} = (\mathbf{f}, \nabla \varphi)_{\Omega} \text{ for all } \varphi \in W^1_{q', \Gamma}(\Omega),$$

which is the so-called *weak Dirichlet-Neumann problem*. Our aim in this subsection is to prove

Proposition 3.4. Let $1 < q < \infty$ and q' = q/(q-1). Then, for every $\mathbf{f} \in L_q(\Omega)^N$, the variational problem (3.3) admits a unique solution $u \in W^1_{q,\Gamma}(\Omega)$, and also $\|u\|_{W^1_q(\Omega)} \leq C_{N,d,q} \|\mathbf{f}\|_{L_q(\Omega)}$ for a positive constant $C_{N,d,q}$. In this case, we define the solution operator Q_q from $L_q(\Omega)^N$ to $W^1_{q,\Gamma}(\Omega)$ by $Q_q \mathbf{f} = u$.

Proof. Since $C_0^{\infty}(\Omega)$ is dense in $L_q(\Omega)$, it suffices to consider the case where $\mathbf{f} = (f_1, \ldots, f_N)^{\mathsf{T}} \in C_0^{\infty}(\Omega)^N$.

We first consider a strong problem associated with (3.3) as follows:

(3.4)
$$\begin{cases} \Delta u = \operatorname{div} \mathbf{f} & \operatorname{in} \Omega, \\ u = 0 & \operatorname{on} \Gamma, \\ \partial_N u = 0 & \operatorname{on} S. \end{cases}$$

For $g = g(x', x_N)$ defined on \mathbf{R}^N_+ , let g^o and g^e be the odd extension of g and the even extension of g, respectively, i.e.

$$g^{o} = \begin{cases} g(x', x_{N}) & (x_{N} > 0), \\ -g(x', -x_{N}) & (x_{N} < 0) \end{cases}, \quad g^{e} = \begin{cases} g(x', x_{N}) & (x_{N} > 0), \\ g(x', -x_{N}) & (x_{N} < 0). \end{cases}$$

Regarding **f** as an element of $C_0^{\infty}(\mathbf{R}^N_+)^N$, we set

(3.5)
$$\mathbf{F} = (F_1, \dots, F_N)^{\mathsf{T}} = (f_1^e, \dots, f_{N-1}^e, f_N^o)^{\mathsf{T}}.$$

Let $\mathcal{F}[g](\xi)$ be the Fourier transform of g = g(x) and $\mathcal{F}_{\xi}^{-1}[h](x)$ the inverse Fourier transform of $h = h(\xi)$, i.e.

$$\mathcal{F}[g](\xi) = \int_{\mathbf{R}^N} e^{-ix\cdot\xi} g(x) \, dx, \quad \mathcal{F}_{\xi}^{-1}[h](x) = \frac{1}{(2\pi)^N} \int_{\mathbf{R}^N} e^{ix\cdot\xi} h(\xi) \, d\xi.$$

Now we define a function v = v(x) on \mathbf{R}^N by

(3.6)
$$v = -\mathcal{F}_{\xi}^{-1} \left[\frac{1}{|\xi|^2} \mathcal{F}[\operatorname{div} \mathbf{F}](\xi) \right](x) = -\sum_{j=1}^N \mathcal{F}_{\xi}^{-1} \left[\frac{i\xi_j}{|\xi|^2} \mathcal{F}[F_j](\xi) \right](x).$$

It then holds that $\Delta v = \operatorname{div} \mathbf{F}$ in \mathbf{R}^N , which implies that $\Delta v = \operatorname{div} \mathbf{f}$ in Ω . On the other hand, one has for $k, l = 1, \ldots, N$

$$\partial_k v = \sum_{j=1}^N \mathcal{F}_{\xi}^{-1} \left[\frac{\xi_j \xi_k}{|\xi|^2} \mathcal{F}[F_j](\xi) \right] (x), \quad \partial_k \partial_l v = \mathcal{F}_{\xi}^{-1} \left[\frac{\xi_k \xi_l}{|\xi|^2} \mathcal{F}[\operatorname{div} \mathbf{F}](\xi) \right] (x),$$

which, combined with the Fourier multiplier theorem of Mikhlin (cf. [27, Appendix Theorem 2]), furnishes that

(3.7)
$$\|\nabla v\|_{L_q(\mathbf{R}^N)} \le C_{N,q} \|\mathbf{f}\|_{L_q(\Omega)}, \quad \|\nabla^2 v\|_{L_q(\mathbf{R}^N)} \le C_{N,q} \|\nabla \mathbf{f}\|_{L_q(\Omega)},$$

for some positive constant $C_{N,q}$.

Next we estimate $||v||_{L_q(\Omega)}$. Let $\widehat{g}(\xi', x_N)$ be the partial Fourier transform of $g = g(x', x_N)$ with respect to x' and $\mathcal{F}_{\xi'}^{-1}[h(\xi', x_N)](x')$ the inverse partial Fourier transform of $h = h(\xi', x_N)$ with respect to ξ' , i.e.

$$\widehat{g}(\xi', x_N) = \int_{\mathbf{R}^{N-1}} e^{-ix'\cdot\xi'} g(x', x_N) \, dx',$$

$$\mathcal{F}_{\xi'}^{-1}[h(\xi', x_N)](x') = \frac{1}{(2\pi)^{N-1}} \int_{\mathbf{R}^{N-1}} e^{ix'\cdot\xi'} h(\xi', x_N) \, d\xi'.$$

We then observe that, for $j = 1, \ldots, N - 1$,

$$\mathcal{F}[F_j](\xi) = \int_0^d \left(e^{-iy_N \xi_N} + e^{iy_N \xi_N} \right) \widehat{f_j}(\xi', y_N) \, dy_N,$$

and that

$$\mathcal{F}[F_N](\xi) = \int_0^d \left(e^{-iy_N\xi_N} - e^{iy_N\xi_N} \right) \widehat{f}_N(\xi', y_N) \, dy_N.$$

Inserting these formulas into (3.6) yields

$$v = v(x', x_N) = -\sum_{j=1}^{N-1} \int_0^d \mathcal{F}_{\xi'}^{-1} \left[\left(\frac{1}{2\pi} \int_{-\infty}^\infty \frac{e^{i(x_N - y_N)\xi_N} + e^{(x_N + y_N)\xi_N}}{|\xi|^2} \right) d\xi_N i\xi_j \widehat{f}_j(\xi', y_N) \right] (x') dy_N - \int_0^d \mathcal{F}_{\xi'}^{-1} \left[\left(\frac{1}{2\pi} \int_{-\infty}^\infty \frac{i\xi_N (e^{i(x_N - y_N)\xi_N} + e^{(x_N + y_N)\xi_N})}{|\xi|^2} \right) d\xi_N \widehat{f}_N(\xi', y_N) \right] (x') dy_N.$$

On the other hand, it holds by the residue theorem that for $a \in \mathbf{R} \setminus \{0\}$

(3.8)
$$\int_{-\infty}^{\infty} \frac{e^{ia\xi_N}}{|\xi|^2} d\xi_N = \frac{\pi e^{-|a||\xi'|}}{|\xi'|}, \quad \int_{-\infty}^{\infty} \frac{i\xi_N e^{ia\xi_N}}{|\xi|^2} d\xi_N = -\pi e^{-|a||\xi'|} \operatorname{sign}(a),$$

where sign(a) = 1 when a > 0 and sign(a) = -1 when a < 0. We combine (3.8) with the above formula of v in order to obtain

$$v = -\frac{1}{2} \sum_{j=1}^{N-1} \int_0^d \mathcal{F}_{\xi'}^{-1} \left[\frac{i\xi_j}{|\xi'|} \left(e^{-|x_N - y_N||\xi'|} + e^{-(x_N + y_N)|\xi'|} \right) \widehat{f}_j(\xi', y_N) \right] (x') \, dy_N \\ + \frac{1}{2} \int_0^d \mathcal{F}_{\xi'}^{-1} \left[\left(\operatorname{sign}(x_N - y_N) e^{-|x_N - y_N||\xi'|} + e^{-(x_N + y_N)|\xi'|} \right) \widehat{f}_N(\xi', y_N) \right] (x') \, dy_N,$$

which implies that $\|v\|_{L_q(\Omega)} \leq C_{N,d,q} \|\mathbf{f}\|_{L_q(\Omega)}$ in the same manner as in the proof of [30, pages 1897-1898]. Hence, together with (3.7), one has

(3.9)
$$\|v\|_{W_q^1(\Omega)} \le C_{N,d,q} \|\mathbf{f}\|_{L_q(\Omega)}, \quad \|v\|_{W_q^2(\Omega)} \le C_{N,d,q} \|\mathbf{f}\|_{W_q^1(\Omega)}$$

We here prove that $\partial_N v = 0$ on \mathbf{R}_0^N . Noting div $\mathbf{F} = (\operatorname{div} \mathbf{f})^e$ by the definition (3.5) and setting $z = \operatorname{div} \mathbf{f}$, we have

$$\mathcal{F}[\operatorname{div} \mathbf{F}](\xi) = \mathcal{F}[z^e](\xi) = \int_0^d \left(e^{-iy_N \xi_N} + e^{iy_N \xi_N} \right) \widehat{z}(\xi', y_N) \, dy_N.$$

By this formula and (3.6), one obtains

$$\partial_N v = (\partial_N v)(x', x_N) = -\int_0^d \mathcal{F}_{\xi'}^{-1} \left[\left(\frac{1}{2\pi} \int_{-\infty}^\infty \frac{i\xi_N (e^{i(x_N - y_N)\xi_N} + e^{i(x_N + y_N)\xi_N})}{|\xi|^2} d\xi_N \right) \hat{z}(\xi', y_N) \right] (x') \, dy_N,$$

which, combined with (3.8), furnishes $(\partial_N v)(x', 0) = 0$. Let u = v + w in (3.4), and thus (3.4) is reduced to

(3.10)
$$\begin{cases} \Delta w = 0 & \text{in } \Omega, \\ w = -v & \text{on } \Gamma, \\ \partial_N w = 0 & \text{on } S. \end{cases}$$

From now on, we solve the system (3.10). Applying the partial Fourier transform to (3.10) yields

$$\begin{cases} (\partial_N^2 - |\xi'|^2)\widehat{w}(\xi', x_N) = 0 & (0 < x_N < d), \\ \widehat{w}(\xi', d) = -\widehat{v}(\xi', d), \\ \partial_N \widehat{w}(\xi', 0) = 0. \end{cases}$$

One then solves this system as ordinary differential equations with respect to \boldsymbol{x}_N in order to obtain

$$w = w(x', x_N) = -\sum_{k=1}^2 \mathcal{F}_{\xi'}^{-1} \left[\frac{e^{-|\xi'|(d+(-1)^k x_N)}}{1+e^{-2|\xi'|d}} \widehat{v}(\xi', d) \right] (x').$$

Let $\varphi = \varphi(s)$ be a function in $C^{\infty}(\mathbf{R})$ such that $0 \leq \varphi \leq 1$ and

$$\varphi(s) = \begin{cases} 1 & \text{for } s \ge 2d/3, \\ 0 & \text{for } s \le d/3, \end{cases}$$

and note that for functions $f(x_N)$, $g(x_N)$ and for k = 1, 2

$$f(x_N)g(d) = \int_0^d \frac{d}{dy_N} \left\{ \varphi(y_N) f(x_N + (-1)^k (y_N - d)) g(y_N) \right\} \, dy_N.$$

Combining these identities with the above formula of w, we obtain

$$\begin{split} w &= -\sum_{k=1}^{2} \int_{0}^{d} \mathcal{F}_{\xi'}^{-1} \left[\frac{\dot{\varphi}(y_{N})e^{-|\xi'|(y_{N}+(-1)^{k}x_{N})}}{1+e^{-2|\xi'|d}} \widehat{v}(\xi',y_{N}) \right] (x') \, dy_{N} \\ &+ \sum_{k=1}^{2} \int_{0}^{d} \mathcal{F}_{\xi'}^{-1} \left[\frac{\varphi(y_{N})|\xi'|e^{-|\xi'|(y_{N}+(-1)^{k}x_{N})}}{1+e^{-2|\xi'|d}} \widehat{v}(\xi',y_{N}) \right] (x') \, dy_{N} \\ &- \sum_{k=1}^{2} \int_{0}^{d} \mathcal{F}_{\xi'}^{-1} \left[\frac{\varphi(y_{N})e^{-|\xi'|(y_{N}+(-1)^{k}x_{N})}}{1+e^{-2|\xi'|d}} \widehat{\partial_{N}v}(\xi',y_{N}) \right] (x') \, dy_{N} \\ &=: I_{1} + I_{2} + I_{3}, \end{split}$$

where $\dot{\varphi}(y_N) = (d\varphi/dy_N)(y_N)$. Noting $|\xi'| = |\xi'|^2/|\xi'| = -\sum_{j=1}^{N-1} (i\xi_j)^2/|\xi'|$, we can write I_2 as

$$I_{2} = -\sum_{j=1}^{N-1} \sum_{k=1}^{2} \int_{0}^{d} \mathcal{F}_{\xi'}^{-1} \left[\frac{\varphi(y_{N})i\xi_{j}e^{-|\xi'|(y_{N}+(-1)^{k}x_{N})}}{|\xi'|(1+e^{-2|\xi'|d})} \widehat{\partial_{j}v}(\xi',y_{N}) \right] (x') \, dy_{N}.$$

The following lemma was essentially proved in Lemma [30, Lemma 5.5].

Lemma 3.5. Let $1 < q < \infty$. Assume that $m(\xi')$ satisfies

$$|\partial_{\xi'}^{\alpha'} m(\xi')| \le C_{\alpha'} |\xi'|^{-|\alpha'|} \quad (\xi' \in \mathbf{R}^{N-1} \setminus \{0\})$$

for any multi-index $\alpha' \in \mathbf{N}_0^{N-1}$, and set for k = 1, 2

$$[L_k f](x) = \int_0^d \mathcal{F}_{\xi'}^{-1} \left[\dot{\varphi}(y_N) m(\xi') e^{-|\xi'|(y_N + (-1)^k x_N)} \widehat{f}(\xi', y_N) \right](x') \, dy_N,$$

$$[M_k f](x) = \int_0^d \mathcal{F}_{\xi'}^{-1} \left[\varphi(y_N) m(\xi') e^{-|\xi'|(y_N + (-1)^k x_N)} \widehat{f}(\xi', y_N) \right](x') \, dy_N.$$

Then, for any $f \in L_q(\Omega)$, we have

$$||(L_k f, M_k f)||_{W^1_q(\Omega)} \le C_{N,d,q} ||f||_{L_q(\Omega)} \quad (k = 1, 2).$$

It is known by e.g. [40, Section 5] that for any multi-index $\alpha' \in \mathbf{N}_0^{N-1}$

(3.11)
$$\left| \partial_{\xi'}^{\alpha'}(i\xi_j|\xi'|^{-1}) \right| + \left| \partial_{\xi'}^{\alpha'}e^{-a|\xi'|} \right| \le C_{\alpha'}|\xi'|^{-|\alpha'|} \quad (\xi' \in \mathbf{R}^{N-1} \setminus \{0\}),$$

where j = 1, ..., N - 1 and $a \ge 0$, with some positive constant $C_{\alpha'}$ independent of ξ' and a. In order to estimate $(1 + e^{-2|\xi'|d})^{-1}$, we introduce Bell's formula for derivatives of the composite function of f(t) and $t = g(\xi')$ as follows: For any multi-index $\alpha' \in \mathbf{N}_0^{N-1}$,

$$\partial_{\xi'}^{\alpha'} f(g(\xi')) = \sum_{l=1}^{|\alpha'|} f^{(l)}(g(\xi')) \sum_{\substack{\alpha'_1 + \dots + \alpha'_l = \alpha', \\ |\alpha'_m| \ge 1}} \Gamma_{\alpha'_1,\dots,\alpha'_l}^{\alpha'}(\partial_{\xi'}^{\alpha'_1}g(\xi')) \dots (\partial_{\xi'}^{\alpha'_l}g(\xi'))$$

with suitable coefficients $\Gamma_{\alpha'_1,\ldots,\alpha'_l}^{\alpha'}$, where $f^{(l)}(t)$ is the *l*th derivative of f(t). By Bell's formula with $f(t) = t^{-1}$ and $t = g(\xi') = 1 + e^{-2|\xi'|d}$ and by (3.11),

(3.12)
$$\left|\partial_{\xi'}^{\alpha'}(1+e^{-2|\xi'|d})^{-1}\right| \le C_{\alpha'}\sum_{l=1}^{|\alpha'|} \left|1+e^{-2|\xi'|d}\right|^{-(l+1)} |\xi'|^{-|\alpha'|} \le C_{\alpha'} |\xi'|^{-|\alpha'|}$$

for any $\xi' \in \mathbf{R}^{N-1} \setminus \{0\}$ and any multi-index $\alpha' \in \mathbf{N}_0^{N-1}$. One thus obtains $\|(I_1, I_3)\|_{W^1_q(\Omega)} \leq C_{N,d,q} \|v\|_{W^1_q(\Omega)}$ by Lemma 3.5 and (3.12). In addition, it holds by Leibniz's rule, (3.11), and (3.12) that

$$\left|\partial_{\xi'}^{\alpha'}\left(\frac{i\xi_j}{|\xi'|(1+e^{-2|\xi'|d})}\right)\right| \le C_{\alpha'}|\xi'|^{-|\alpha'|} \quad (\xi' \in \mathbf{R}^{N-1} \setminus \{0\}),$$

which, combined with Lemma 3.5, furnishes $||I_2||_{W_q^1(\Omega)} \leq C_{N,d,q}||v||_{W_q^1(\Omega)}$. Analogously, we can prove that $||\nabla(I_1, I_2, I_3)||_{W_q^1(\Omega)} \leq C_{N,d,q}||\nabla v||_{W_q^1(\Omega)}$. Summing up these inequalities for I_1 , I_2 , and I_3 , one has by (3.9)

(3.13)
$$\|w\|_{W_q^1(\Omega)} \le C_{N,d,q} \|\mathbf{f}\|_{L_q(\Omega)}, \quad \|w\|_{W_q^2(\Omega)} \le C_{N,d,q} \|\mathbf{f}\|_{W_q^1(\Omega)}.$$

It now holds that u = v + w solves (3.4) and satisfies $||u||_{W_q^1(\Omega)} \leq C_{N,d,q} ||\mathbf{f}||_{L_q(\Omega)}$ by (3.9) and (3.13). Clearly, such an u is a solution to (3.3).

Finally, we prove the uniqueness of solutions to (3.3). Let $u \in W^1_{q,\Gamma}(\Omega)$ be a solution to (3.3) with $\mathbf{f} = 0$, and let $\Phi \in C_0^{\infty}(\Omega)^N$. Since $\Phi \in L_{q'}(\Omega)^N$, there is a $v \in W^1_{q',\Gamma}(\Omega)$ such that

$$(\nabla v, \nabla \psi)_{\Omega} = (\Phi, \nabla \psi)_{\Omega}$$
 for all $\psi \in W^1_{q,\Gamma}(\Omega)$.

In this equation, we set $\psi = u$ in order to obtain

$$(\Phi, \nabla u)_{\Omega} = (\nabla v, \nabla u)_{\Omega} = 0,$$

which implies that u is a constant. Hence, we have u = 0, because u = 0 on Γ . This completes the proof of the Proposition 3.4.

3.3. Helmholtz decomposition on Ω . We here introduce the Helmholtz decomposition on Ω . Let $G_q(\Omega) = \{\nabla \theta \mid \theta \in W^1_{q,\Gamma}(\Omega)\}$, and then one has

Proposition 3.6. Let $1 < q < \infty$. Then the following assertions hold true. (1) $L_q(\Omega)^N = J_q(\Omega) \oplus G_q(\Omega).$

(2) Let P_q be the projection from $L_q(\Omega)^N$ to $J_q(\Omega)$, and let Q_q be the solution operator of Proposition 3.4 from $L_q(\Omega)^N$ to $W_{q,\Gamma}^1(\Omega)$. Then, for any $\mathbf{f} \in L_q(\Omega)^N$, we have $\mathbf{f} = P_q \mathbf{f} + \nabla Q_q \mathbf{f} \in J_q(\Omega) \oplus G_q(\Omega)$ and

(3.14)
$$\|P_{q}\mathbf{f}\|_{L_{q}(\Omega)} + \|Q_{q}\mathbf{f}\|_{W_{q}^{1}(\Omega)} \le C_{N,d,q}\|\mathbf{f}\|_{L_{q}(\Omega)}$$

for some positive constant $C_{N,d,q}$.

Proof. (1). Let $\mathbf{f} \in L_q(\Omega)^N$. By Proposition 3.4, we have a unique solution u = $Q_q \mathbf{f} \in W^1_{a,\Gamma}(\Omega)$ to the variational problem:

$$(\nabla u, \nabla \varphi)_{\Omega} = (\mathbf{f}, \nabla \varphi)_{\Omega} \text{ for all } \varphi \in W^1_{q', \Gamma}(\Omega),$$

where q' = q/(q-1). Setting $\mathbf{g} = \mathbf{f} - \nabla Q_q \mathbf{f}$, we observe that $\mathbf{g} \in J_q(\Omega)$ and $\mathbf{f} = \mathbf{g} + \nabla Q_q \mathbf{f} \in J_q(\Omega) + G_q(\Omega).$ Next we prove that $L_q(\Omega)^N = J_q(\Omega) \oplus G_q(\Omega)$. To this end, let $\mathbf{f} \in J_q(\Omega) \cap G_q(\Omega)$.

We then have

$$(\nabla Q_q \mathbf{f}, \nabla \varphi)_{\Omega} = (\mathbf{f}, \nabla \varphi)_{\Omega} = 0 \text{ for all } \varphi \in W^1_{q', \Gamma}(\Omega),$$

which, combined with the uniqueness of Proposition 3.4, furnishes $Q_q \mathbf{f} = 0$. On the other hand, since there is a $\theta \in W^1_{q,\Gamma}(\Omega)$ such that $\mathbf{f} = \nabla \theta$, we have

$$(\nabla Q_q \mathbf{f}, \nabla \varphi)_{\Omega} = (\mathbf{f}, \nabla \varphi)_{\Omega} = (\nabla \theta, \nabla \varphi)_{\Omega} \quad \text{for all } \varphi \in W^1_{q', \Gamma}(\Omega).$$

By the uniqueness of Proposition 3.4 again, it holds that $\theta = Q_q \mathbf{f} = 0$. Thus $\mathbf{f} = 0$, which implies $L_q(\Omega)^N = J_q(\Omega) \oplus G_q(\Omega)$.

(2). For any $\mathbf{f} \in L_q(\Omega)^N$, the projection P_q is given by $P_q \mathbf{f} = \mathbf{f} - \nabla Q_q \mathbf{f} \in J_q(\Omega)$ as was seen in the proof of (1). Thus the first assertion clearly holds true, and also the second assertion (3.14) follows from Proposition 3.4 immediately. This completes the proof of Proposition 3.6.

4. Generation of the Stokes semigroup

Our aim in this section is to construct an analytic C_0 -semigroup associated with

(4.1)
$$\begin{cases} \partial_t \mathbf{u} - \operatorname{Div} \mathbf{T}(\mathbf{u}, \mathbf{p}) = 0 & \text{in } \Omega, \ t > 0, \\ \operatorname{div} \mathbf{u} = 0 & \operatorname{in} \Omega, \ t > 0, \\ \mathbf{T}(\mathbf{u}, \mathbf{p}) \mathbf{e}_N = 0 & \operatorname{on} \Gamma, \ t > 0, \\ \mathbf{u} = 0 & \operatorname{on} S, \ t > 0, \\ \mathbf{u}|_{t=0} = \mathbf{a} & \operatorname{in} \Omega. \end{cases}$$

To this end, we start with the following resolvent problem:

(4.2)
$$\begin{cases} \lambda \mathbf{v} - \operatorname{Div} \mathbf{T}(\mathbf{v}, \mathbf{q}) = \mathbf{f} & \text{in } \Omega, \\ \operatorname{div} \mathbf{v} = 0 & \text{in } \Omega, \\ \mathbf{T}(\mathbf{v}, \mathbf{q}) \mathbf{e}_N = 0 & \text{on } \Gamma, \\ \mathbf{v} = 0 & \text{on } S, \end{cases}$$

with the resolvent parameter $\lambda \in \Sigma_{\varepsilon} = \{\omega \in \mathbf{C} \setminus \{0\} \mid |\arg \omega| < \pi - \varepsilon\}$ for $0 < \varepsilon < \pi/2$. The following lemma was proved in [2, Theorem 1.1].

Lemma 4.1. Let $1 < q < \infty$ and $0 < \varepsilon < \pi/2$. Then there exists a positive number ω_1 such that, for every $\lambda \in \Sigma_{\varepsilon} \cup \{\omega \in \mathbf{C} \mid |\omega| < \omega_1\}$ and $\mathbf{f} \in L_q(\Omega)^N$, there is a unique solution $(\mathbf{v}, \mathbf{q}) \in W_q^2(\Omega)^N \times W_q^1(\Omega)$ to (4.2). In addition,

$$|\lambda| \|\mathbf{v}\|_{L_{q}(\Omega)} + |\lambda|^{1/2} \|\nabla \mathbf{v}\|_{L_{q}(\Omega)} + \|\mathbf{v}\|_{W_{q}^{2}(\Omega)} + \|\mathbf{q}\|_{W_{q}^{1}(\Omega)} \le C \|\mathbf{f}\|_{L_{q}(\Omega)}$$

for any $\lambda \in \Sigma_{\varepsilon} \cup \{\omega \in \mathbf{C} \mid |\omega| < \omega_1\}$ with a positive constant $C = C_{N,d,q,\varepsilon,\mu,\omega_1}$.

Let $1 < q < \infty$ and q' = q/(q-1). By Proposition 3.4, we see that, for any $\mathbf{f} \in L_q(\Omega)^N$ and $g \in W_q^{1-1/q}(\Gamma)$, there exists a unique solution $\tilde{\mathbf{q}} \in W_{q,\Gamma}^1(\Omega)$ to the variational problem:

$$(\nabla \widetilde{\mathfrak{q}}, \varphi)_{\Omega} = (\mathbf{f} - \nabla \widetilde{g}, \nabla \varphi)_{\Omega} \text{ for all } \varphi \in W^1_{a', \Gamma}(\Omega),$$

where \tilde{g} is an extension of g satisfying $\|\tilde{g}\|_{W_q^1(\Omega)} \leq C \|g\|_{W_q^{1-1/q}(\Gamma)}$ for some positive constant C independent of g, \tilde{g} . Furthermore, the solution $\tilde{\mathfrak{q}}$ satisfies

$$\begin{aligned} \|\widetilde{\mathfrak{q}}\|_{W_q^1(\Omega)} &\leq C_{N,d,q}(\|\mathbf{f}\|_{L_q(\Omega)} + \|\nabla\widetilde{g}\|_{L_q(\Omega)}) \\ &\leq C_{N,d,q}(\|\mathbf{f}\|_{L_q(\Omega)} + \|g\|_{W_q^{1-1/q}(\Gamma)}). \end{aligned}$$

Setting $\mathfrak{q} = \widetilde{\mathfrak{q}} + \widetilde{g} \in W^1_{q,\Gamma}(\Omega) + W^1_q(\Omega)$ yields that

$$(\nabla \mathfrak{q}, \nabla \varphi)_{\Omega} = (\mathbf{f}, \nabla \varphi)_{\Omega} \quad \text{for all } \varphi \in W^1_{q', \Gamma}(\Omega), \quad \mathfrak{q} = g \quad \text{on } \Gamma,$$

and also $\|\mathbf{q}\|_{W_q^1(\Omega)} \leq C_{N,d,q}(\|\mathbf{f}\|_{L_q(\Omega)} + \|g\|_{W_q^{1-1/q}(\Gamma)})$. From this viewpoint, we define an operator $K: W_q^2(\Omega)^N \ni \mathbf{v} \mapsto K(\mathbf{v}) \in W_{q,\Gamma}^1(\Omega) + W_q^1(\Omega)$ as follows:

$$(\nabla K(\mathbf{v}), \nabla \varphi)_{\Omega} = (\operatorname{Div}(\mu \mathbf{D}(\mathbf{v})) - \nabla \operatorname{div} \mathbf{v}, \nabla \varphi)_{\Omega} \quad \text{for all } \varphi \in W^{1}_{q', \Gamma}(\Omega)$$

$$K(\mathbf{v}) = \mathbf{e}_{N} \cdot (\mu \mathbf{D}(\mathbf{v}) \mathbf{e}_{N}) - \operatorname{div} \mathbf{v} \quad \text{on } \Gamma.$$

Then $K(\mathbf{v})$ satisfies $||K(\mathbf{v})||_{W_q^1(\Omega)} \leq C_{N,d,q,\mu} ||\mathbf{v}||_{W_q^2(\Omega)}$ with some positive constant $C_{N,q,d,\mu}$ independent of \mathbf{v} and φ .

At this point, we introduce a result concerning the weak Dirichlet-Neumann problem with resolvent parameter λ .

Proposition 4.2. Let $1 < q < \infty$ and $0 < \varepsilon < \pi/2$. Then there exists a positive number ω_2 such that, for every $\lambda \in \Sigma_{\varepsilon} \cup \{\omega \in \mathbf{C} \mid |\omega| < \omega_2\}$ and $\mathbf{f} \in L_q(\Omega)^N$, there is a unique solution $u \in W^1_{q,\Gamma}(\Omega)$ to the variational problem:

(4.3)
$$(\lambda u, \varphi)_{\Omega} + (\nabla u, \nabla \varphi)_{\Omega} = (\mathbf{f}, \nabla \varphi)_{\Omega} \quad \text{for all } \varphi \in W^{1}_{q', \Gamma}(\Omega),$$

where q' = q/(q-1).

Proof. The case $\lambda = 0$ was already proved in Proposition 3.4. Then, by a small perturbation method, we can prove that there exists a positive constant ω_2 such that, for every $\lambda \in \{\omega \in \mathbb{C} \mid |\omega| < \omega_2\}$ and $\mathbf{f} \in L_q(\Omega)^N$, (4.3) admits a unique solution $u \in W^1_{q,\Gamma}(\Omega)$. In the case $\lambda \in \Sigma_{\varepsilon}$ with $|\lambda| \geq \omega_2/2$, we consider the strong problem with resolvent parameter λ for (3.4). One can construct solutions to the strong problem in a similar way to Proposition 3.4 (cf. also [30]). The uniqueness follows from the existence of solutions for a dual problem as was seen in the proof of Proposition 3.4. This completes the proof of the proposition.

We now consider the reduced resolvent problem:

(4.4)
$$\begin{cases} \lambda \mathbf{v} - \operatorname{Div} \mathbf{T}(\mathbf{v}, K(\mathbf{v})) = \mathbf{f} & \text{in } \Omega, \\ \mathbf{T}(\mathbf{v}, K(\mathbf{v})) \mathbf{e}_N = 0 & \text{on } \Gamma, \\ \mathbf{v} = 0 & \text{on } S. \end{cases}$$

Then the following proposition holds.

Proposition 4.3. Let $1 < q < \infty$ and $0 < \varepsilon < \pi/2$. Assume that ω_1 and ω_2 are, respectively, the same positive constants as in Lemma 4.1 and Proposition 4.2, and set $\omega_0 = \min(\omega_1, \omega_2)$. Then (4.2) is equivalent to (4.4) for every $\lambda \in \Sigma_{\varepsilon} \cup \{\omega \in \mathbf{C} \mid |\omega| < \omega_0\}$ and $\mathbf{f} \in J_q(\Omega)$, which means that the following assertions hold true: $(\mathbf{v}, \mathbf{q}) = (\mathbf{v}, K(\mathbf{v})) \in W_q^2(\Omega)^N \times W_q^1(\Omega)$ is a unique solution to (4.2) if $\mathbf{v} \in W_q^2(\Omega)^N$ is a solution to (4.4), and conversely, $\mathbf{v} \in W_q^2(\Omega)^N$ is a unique solution to (4.4) if $(\mathbf{v}, \mathbf{q}) \in W_q^2(\Omega)^N \times W_q^1(\Omega)$ is a solution to (4.2).

Proof. Suppose that $\mathbf{v} \in W^2_q(\Omega)^N$ is a solution to (4.4). Let $\varphi \in W^1_{q',\Gamma}(\Omega)$ with 1/q + 1/q' = 1. We then see, by the definition of K, that div $\mathbf{v} = 0$ on Γ and

$$0 = -(\mathbf{f}, \nabla \varphi)_{\Omega} = -(\lambda \mathbf{v} - \operatorname{Div} \mathbf{T}(\mathbf{v}, K(\mathbf{v})), \nabla \varphi)_{\Omega} = -(\lambda \mathbf{v} - \nabla \operatorname{div} \mathbf{v}, \nabla \varphi)_{\Omega}$$
$$= (\lambda \operatorname{div} \mathbf{v}, \varphi)_{\Omega} + (\nabla \operatorname{div} \mathbf{v}, \nabla \varphi)_{\Omega}.$$

Hence, div $\mathbf{v} = 0$ by Proposition 4.2 when $\lambda \in \Sigma_{\varepsilon} \cup \{\omega \in \mathbf{C} \mid |\omega| < \omega_0\}$, and therefore setting $\mathbf{q} = K(\mathbf{v})$ implies that $(\mathbf{v}, \mathbf{q}) \in W_q^2(\Omega)^N \times W_q^1(\Omega)$ solves (4.2). The uniqueness of solutions to (4.2) follows from Lemma 4.1.

Next we show the opposite direction. Suppose that $(\mathbf{v}, \mathbf{q}) \in W_q^2(\Omega)^N \times W_q^1(\Omega)$ is a solution to (4.2). Let $\varphi \in W_{q',\Gamma}^1(\Omega)$, and then we see, by the definition of Kand div $\mathbf{v} = 0$ in Ω , that

$$0 = (\mathbf{f}, \nabla \varphi)_{\Omega} = (\nabla \mathbf{q} - \operatorname{Div}(\mu \mathbf{D}(\mathbf{v})), \nabla \varphi)_{\Omega} = (\nabla (\mathbf{q} - K(\mathbf{v})), \nabla \varphi)_{\Omega},$$

$$0 = \mathbf{q} - K(\mathbf{v}) \quad \text{on } \Gamma,$$

where we have used $(\lambda \mathbf{v}, \nabla \varphi)_{\Omega} = 0$ by div $\mathbf{v} = 0$ in Ω and $\mathbf{v} = 0$ on S. Combining these two equations with the uniqueness of Proposition 3.4 implies $\mathbf{q} = K(\mathbf{v})$. Thus \mathbf{v} is a solution to (4.4). The uniqueness of solutions to (4.4) follows from the first half of this proof, which completes the proof of the proposition.

In view of (4.4), we set the Stokes operator A_q as $A_q \mathbf{v} = \text{Div } \mathbf{T}(\mathbf{v}, K(\mathbf{v}))$ with the domain $D(A_q)$:

(4.5)
$$D(A_q) = W_{q,\mathcal{B}}^2(\Omega) \cap J_q(\Omega),$$
$$W_{q,\mathcal{B}}^2(\Omega) = \{ \mathbf{v} \in W_q^2(\Omega)^N \mid (\mu \mathbf{D}(\mathbf{v})\mathbf{e}_N)_\tau = 0 \text{ on } \Gamma, \quad \mathbf{v} = 0 \text{ on } S \}.$$

The system (4.4) then can be written as $\lambda \mathbf{v} - A_q \mathbf{v} = \mathbf{f}$, and one has

Lemma 4.4. Let $1 < q < \infty$ and $0 < \varepsilon < \pi/2$. Suppose that ω_0 is the same positive constant as in Proposition 4.3. Then there exists a positive constant $C = C_{N,d,q,\varepsilon,\mu,\omega_1,\omega_2}$ such that, for every $\lambda \in \Sigma_{\varepsilon} \cup \{\omega \in \mathbf{C} \mid |\omega| < \omega_0\}$,

$$\|(\lambda - A_q)^{-1}\|_{\mathcal{L}(J_q(\Omega))} \le \frac{C}{1+|\lambda|}.$$

In addition, A_q is a densely defined closed operator on $J_q(\Omega)$.

Proof. The required estimate follows from Lemma 4.1 and Proposition 4.3.

Let $\mathbf{v} \in D(A_q)$. Then $(\operatorname{div} \mathbf{v}, \varphi)_{\Omega} = -(\mathbf{v}, \nabla \varphi)_{\Omega} = 0$ for any $\varphi \in C_0^{\infty}(\Omega)$, because $\mathbf{v} \in J_q(\Omega)$ and $C_0^{\infty}(\Omega) \subset W^1_{q',\Gamma}(\Omega)$ with q' = q/(q-1). This implies $\operatorname{div} \mathbf{v} = 0$ in Ω . It thus holds by the definition of K that

$$(A_q \mathbf{v}, \nabla \varphi)_{\Omega} = (\nabla \operatorname{div} \mathbf{v}, \nabla \varphi)_{\Omega} = 0 \text{ for all } \varphi \in W^1_{q', \Gamma}(\Omega),$$

which furnishes $A_q \mathbf{v} \in J_q(\Omega)$. Then, following the proof of [39, Lemma 3.7], we can show the last assertion of Lemma 4.4. This completes the proof of the lemma. \Box

The following proposition follows from Lemma 4.4 and the standard theory of analytic C_0 -semigroups.

Proposition 4.5. Let $1 < q < \infty$. Then A_q generates an analytic C_0 -semigroup $\{e^{A_q t}\}_{t \ge 0}$ on $J_q(\Omega)$. Furthermore, there exist positive constants σ_0 and $C = C_{N,d,q,\mu,\sigma_0}$ such that for any t > 0

$$\|e^{A_q t} \mathbf{a}\|_{J_q(\Omega)} \le C e^{-2\sigma_0 t} \|\mathbf{a}\|_{J_q(\Omega)} \qquad (\mathbf{a} \in J_q(\Omega)),$$

$$\|\partial_t e^{A_q t} \mathbf{a}\|_{J_q(\Omega)} \le C t^{-1} e^{-2\sigma_0 t} \|\mathbf{a}\|_{J_q(\Omega)} \qquad (\mathbf{a} \in J_q(\Omega)),$$

$$\|\partial_t e^{A_q t} \mathbf{a}\|_{J_q(\Omega)} \le C e^{-2\sigma_0 t} \|\mathbf{a}\|_{D(A_q)} \qquad (\mathbf{a} \in D(A_q)),$$

where $\|\mathbf{a}\|_{D(A_q)} = \|\mathbf{a}\|_{J_q(\Omega)} + \|A_q\mathbf{a}\|_{J_q(\Omega)}.$

Recall $D_{q,p}(\Omega) = (J_q(\Omega), D(A_q))_{1-1/p,p}$ for $1 < p, q < \infty$. We have a corollary of Proposition 4.5 that can be proved in the same manner as in [39, Theorem 3.9].

Corollary 4.6. Let $1 < p, q < \infty$ and σ_0 be the same positive constant as in Proposition 4.5. Then, for every $\mathbf{a} \in D_{q,p}(\Omega)$, $(\mathbf{u}, \mathbf{p}) = (e^{A_q t} \mathbf{a}, K(e^{A_q t} \mathbf{a}))$ is a unique solution to (4.1), and also

$$\|e^{\sigma_0 t}(\partial_t \mathbf{u}, \mathbf{u}, \nabla \mathbf{u}, \nabla^2 \mathbf{u})\|_{L_p(\mathbf{R}_+, L_q(\Omega))} + \|e^{\sigma_0 t} \mathfrak{p}\|_{L_p(\mathbf{R}_+, W^1_q(\Omega))} \le C \|\mathbf{a}\|_{D_{q, p}(\Omega)}$$

for a positive constant $C = C_{N,d,p,q,\mu,\sigma_0}$.

Remark 4.7. By [24, Propositions 2.2.2, 2.2.8], it holds that $\lim_{t\to 0+} ||e^{A_q t} \mathbf{a} - \mathbf{a}||_{D_{q,p}(\Omega)} = 0$ for any $\mathbf{a} \in D_{q,p}(\Omega)$. Then, under the conditions $2/p + 1/q \neq 1$ and $2/p + 1/q \neq 2$, we observe by Remark 2.1 that $\lim_{t\to 0+} ||e^{A_q t} \mathbf{a} - \mathbf{a}||_{B^{2-2/p}_{q,p}(\Omega)} = 0$ for any $\mathbf{a} \in D_{q,p}(\Omega)$.

5. Proof of Theorem 2.4

This section proves Theorem 2.4. Assume that σ_0 is the same positive constant as in Proposition 4.5 in what follows.

Step 1. The aim of this step is to decompose (\mathbf{u}, \mathbf{p}) of (2.3). Let

$$\mathbf{u} = \mathbf{u}_1 + \mathbf{u}_2 + \widetilde{\mathbf{u}}, \quad \mathbf{p} = \mathbf{p}_1 + \mathbf{p}_2 + \mathbf{p}$$

where each term on the right-hand sides satisfies the following systems:

(5.1)
$$\begin{cases} \partial_t \mathbf{u}_1 - \operatorname{Div} \mathbf{T}(\mathbf{u}_1, \mathbf{p}_1) = 0 & \text{in } \Omega, \ t > 0, \\ \operatorname{div} \mathbf{u}_1 = 0 & \operatorname{in} \Omega, \ t > 0, \\ \mathbf{T}(\mathbf{u}_1, \mathbf{p}_1)\mathbf{e}_N = 0 & \text{on } \Gamma, \ t > 0, \\ \mathbf{u}_1 = 0 & \text{on } S, \ t > 0, \\ \mathbf{u}_1|_{t=0} = \mathbf{a} & \operatorname{in} \Omega, \end{cases}$$

HIROKAZU SAITO

(5.2)
$$\begin{cases} \partial_t \mathbf{u}_2 + 2\sigma_0 \mathbf{u}_2 - \operatorname{Div} \mathbf{T}(\mathbf{u}_2, \mathbf{p}_2) = 0 & \text{in } \Omega, \ t > 0, \\ \operatorname{div} \mathbf{u}_2 = g & \text{in } \Omega, \ t > 0, \\ \mathbf{T}(\mathbf{u}_2, \mathbf{p}_2) \mathbf{e}_N = \mathbf{h} & \text{on } \Gamma, \ t > 0, \\ \mathbf{u}_2 = 0 & \text{on } S, \ t > 0, \\ \mathbf{u}_2|_{t=0} = 0 & \text{in } \Omega, \end{cases} \\\begin{cases} \partial_t \widetilde{\mathbf{u}} - \operatorname{Div} \mathbf{T}(\widetilde{\mathbf{u}}, \widetilde{\mathbf{p}}) = \mathbf{f} + 2\sigma_0 \mathbf{u}_2 & \text{in } \Omega, \ t > 0, \\ \operatorname{div} \widetilde{\mathbf{u}} = 0 & \text{in } \Omega, \ t > 0, \\ \operatorname{div} \widetilde{\mathbf{u}} = 0 & \text{on } S, \ t > 0, \\ \widetilde{\mathbf{u}}(\widetilde{\mathbf{u}}, \widetilde{\mathbf{p}}) \mathbf{e}_N = 0 & \text{on } \Gamma, \ t > 0, \\ \widetilde{\mathbf{u}} = 0 & \text{on } S, \ t > 0, \\ \widetilde{\mathbf{u}}|_{t=0} = 0 & \text{in } \Omega. \end{cases}$$

Let P_q and Q_q be the operators studied in Subsection 3.3. We then have $\mathbf{f} + 2\sigma_0 \mathbf{u}_2 = P_q(\mathbf{f} + 2\sigma_0 \mathbf{u}_2) + \nabla Q_q(\mathbf{f} + 2\sigma_0 \mathbf{u}_2),$

$$T + 2\sigma_0 \mathbf{u}_2 = P_q(\mathbf{f} + 2\sigma_0 \mathbf{u}_2) + \nabla Q_q(\mathbf{f} + 2\sigma_0 \mathbf{u}_2)$$

which gives further decompositions of $(\widetilde{\mathbf{u}}, \widetilde{\mathfrak{p}})$ as follows:

$$\widetilde{\mathbf{u}} = \mathbf{u}_3 + \mathbf{u}_4, \quad \widetilde{\mathbf{p}} = \mathbf{p}_3 + \mathbf{p}_4,$$

where

$$(5.3) \begin{cases} \partial_{t} \mathbf{u}_{3} + 2\sigma_{0} \mathbf{u}_{3} - \operatorname{Div} \mathbf{T}(\mathbf{u}_{3}, \mathbf{p}_{3}) = \nabla Q_{q}(\mathbf{f} + 2\sigma_{0} \mathbf{u}_{2}) & \text{in } \Omega, t > 0, \\ \operatorname{div} \mathbf{u}_{3} = 0 & \operatorname{in} \Omega, t > 0, \\ \mathbf{T}(\mathbf{u}_{3}, \mathbf{p}_{3}) \mathbf{e}_{N} = 0 & \operatorname{on} \Gamma, t > 0, \\ \mathbf{u}_{3} = 0 & \operatorname{on} S, t > 0, \\ \mathbf{u}_{3}|_{t=0} = 0 & \operatorname{in} \Omega, \end{cases} \\ (5.4) \begin{cases} \partial_{t} \mathbf{u}_{4} - \operatorname{Div} \mathbf{T}(\mathbf{u}_{4}, \mathbf{p}_{4}) = P_{q}(\mathbf{f} + 2\sigma_{0} \mathbf{u}_{2}) + 2\sigma_{0} \mathbf{u}_{3} & \operatorname{in} \Omega, t > 0, \\ \operatorname{div} \mathbf{u}_{4} = 0 & \operatorname{in} \Omega, t > 0, \\ \operatorname{div} \mathbf{u}_{4} = 0 & \operatorname{on} S, t > 0, \\ \mathbf{u}_{4}|_{t=0} = 0 & \operatorname{on} S, t > 0, \\ \mathbf{u}_{4}|_{t=0} = 0 & \operatorname{in} \Omega. \end{cases} \end{cases}$$

By Corollary 4.6 and Remark 4.7, there exists a solution $(\mathbf{u}_1, \mathbf{p}_1)$ of (5.1) such that $\lim_{t\to 0+} \|\mathbf{u}_1(t) - \mathbf{a}\|_{B^{2-2/p}_{q,p}(\Omega)} = 0$ and

(5.5)
$$\|e^{\sigma_0 t}(\partial_t \mathbf{u}_1, \mathbf{u}_1, \nabla \mathbf{u}_1, \nabla^2 \mathbf{u}_1)\|_{L_p(\mathbf{R}_+, L_q(\Omega))} + \|e^{\sigma_0 t} \mathfrak{p}_1\|_{L_p(\mathbf{R}_+, W^1_q(\Omega))}$$
$$\leq C_{N, d, p, q, \mu, \sigma_0} \|\mathbf{a}\|_{D_{q, p}(\Omega)}.$$

Step 2. We consider (5.2) in this step. To this end, one extends g and \mathbf{h} to functions defined on the whole line with respect to time t. Let E_0 be the zero extension operator and X be a Banach space. Then,

(5.6)
$$E_0 \in \mathcal{L}(L_p(\mathbf{R}_+, X), L_p(\mathbf{R}, X)) \cap \mathcal{L}({}_0W_p^1(\mathbf{R}_+, X), W_p^1(\mathbf{R}, X))$$

Especially, by (5.6) with $X = L_q(\Omega)$ and the complex interpolation method,

(5.7)
$$E_0 \in \mathcal{L}({}_0H_p^{1/2}(\mathbf{R}_+, L_q(\Omega)), H_p^{1/2}(\mathbf{R}, L_q(\Omega))),$$

where we have used the fact that

$$[L_p(\mathbf{R}, L_q(\Omega)), W_p^1(\mathbf{R}, L_q(\Omega))]_{1/2} = H_p^{1/2}(\mathbf{R}, L_q(\Omega))$$

(cf. e.g. [16, Theorem 1.56]). By (5.6) and (5.7), one has

 $E_0(e^{\sigma_0 t}g) \in H^{1,1/2}_{q,p}(\Omega \times \mathbf{R}) \cap W^1_p(\mathbf{R}, \widehat{W}^{-1}_{q,\Gamma}(\Omega)), \quad E_0(e^{\sigma_0 t}\mathbf{h}) \in H^{1,1/2}_{q,p}(\Omega \times \mathbf{R})^N,$

together with the estimates:

$$\begin{aligned} &\|E_0(e^{\sigma_0 t}g)\|_{H^{1,1/2}_{q,p}(\Omega\times\mathbf{R})} \le C_{p,q} \|e^{\sigma_0 t}g\|_{0H^{1,1/2}_{q,p}(\Omega\times\mathbf{R}_+)}, \\ &\|E_0(e^{\sigma_0 t}g)\|_{W^1_p(\mathbf{R},\widehat{W}^{-1}_{q,\Gamma}(\Omega))} \le C_{p,q} \|e^{\sigma_0 t}g\|_{W^1_p(\mathbf{R}_+,\widehat{W}^{-1}_{q,\Gamma}(\Omega))}, \\ &\|E_0(e^{\sigma_0 t}\mathbf{h})\|_{H^{1,1/2}_{q,p}(\Omega\times\mathbf{R})} \le C_{p,q} \|e^{\sigma_0 t}\mathbf{h}\|_{0H^{1,p/2}_{q,\Gamma}(\Omega\times\mathbf{R}_+)}. \end{aligned}$$

Thus, setting $G = e^{-\sigma_0 t} E_0(e^{\sigma_0 t}g)$ and $\mathbf{H} = e^{-\sigma_0 t} E_0(e^{\sigma_0 t}\mathbf{h})$ yields

$$\begin{split} & G \in H^{1,1/2}_{q,p,-\sigma_0}(\Omega \times \mathbf{R}) \cap W^1_{p,-\sigma_0}(\mathbf{R},\widehat{W}^{-1}_{q,\Gamma}(\Omega)), \quad \mathbf{H} \in H^{1,1/2}_{q,p,-\sigma_0}(\Omega \times \mathbf{R})^N, \\ & G = \begin{cases} g \quad (t > 0), \\ 0 \quad (t < 0), \end{cases} \quad \mathbf{H} = \begin{cases} \mathbf{h} \quad (t > 0), \\ 0 \quad (t < 0), \end{cases} \end{split}$$

and also

$$\begin{split} \|e^{\sigma_{0}t}G\|_{H^{1,1/2}_{q,p}(\Omega\times\mathbf{R})} &\leq C_{p,q} \|e^{\sigma_{0}t}g\|_{{}_{0}H^{1,1/2}_{q,p}(\Omega\times\mathbf{R}_{+})},\\ \|e^{\sigma_{0}t}(\partial_{t}G,G)\|_{L_{p}(\mathbf{R},\widehat{W}^{-1}_{q,\Gamma}(\Omega))} &\leq C_{p,q} \|e^{\sigma_{0}t}g\|_{W^{1}_{p}(\mathbf{R}_{+},\widehat{W}^{-1}_{q,\Gamma}(\Omega))},\\ \|e^{\sigma_{0}t}\mathbf{H}\|_{H^{1,1/2}_{q,p}(\Omega\times\mathbf{R})} &\leq C_{p,q} \|e^{\sigma_{0}t}\mathbf{h}\|_{{}_{0}H^{1,1/2}_{q,p}(\Omega\times\mathbf{R}_{+})}. \end{split}$$

Combining these properties with Proposition 3.1 for $\delta = \sigma_0$ furnishes that there is a solution $(\mathbf{U}, P) \in W^{2,1}_{q,p,-\sigma_0}(\Omega \times \mathbf{R})^N \times L_{p,-\sigma_0}(\mathbf{R}, W^1_q(\Omega))$ to

$$\begin{cases} \partial_t \mathbf{U} + 2\sigma_0 \mathbf{U} - \operatorname{Div} \mathbf{T}(\mathbf{U}, P) = 0 & \text{in } \Omega, \ t \in \mathbf{R}, \\ \operatorname{div} \mathbf{U} = G & \operatorname{in} \Omega, \ t \in \mathbf{R}, \\ \mathbf{T}(\mathbf{U}, P) \mathbf{e}_N = \mathbf{H} & \text{on } \Gamma, \ t \in \mathbf{R}, \\ \mathbf{U} = 0 & \text{on } S, \ t \in \mathbf{R}, \end{cases}$$

and furthermore, **U** vanishes for t < 0 and

$$\begin{aligned} &\|e^{\sigma_0 t}(\partial_t \mathbf{U}, \mathbf{U}, \nabla \mathbf{U}, \nabla^2 \mathbf{U})\|_{L_p(\mathbf{R}, L_q(\Omega))} + \|e^{\sigma_0 t} P\|_{L_p(\mathbf{R}, W_q^1(\Omega))} \\ &\leq C_{N, d, p, q, \mu, \sigma_0} \left(\|e^{\sigma_0 t} g\|_{W_p^1(\mathbf{R}_+, \widehat{W}_{q, \Gamma}^{-1}(\Omega))} + \|e^{\sigma_0 t}(g, \mathbf{h})\|_{0H_{q, p}^{1, 1/2}(\Omega \times \mathbf{R}_+)} \right). \end{aligned}$$

Therefore, $(\mathbf{u}_2, \mathbf{p}_2) = (\mathbf{U}, P)$ solves (5.2) and satisfies

(5.8)
$$\|e^{\sigma_0 t}(\partial_t \mathbf{u}_2, \mathbf{u}_2, \nabla \mathbf{u}_2, \nabla^2 \mathbf{u}_2)\|_{L_p(\mathbf{R}_+, L_q(\Omega))} + \|e^{\sigma_0 t} \mathfrak{p}_2\|_{L_p(\mathbf{R}_+, L_q(\Omega))}$$

 $\leq C_{N,d,p,q,\mu,\sigma_0} \left(\|e^{\sigma_0 t}g\|_{W_p^1(\mathbf{R}_+, \widehat{W}_{q,\Gamma}^{-1}(\Omega))} + \|e^{\sigma_0 t}(g, \mathbf{h})\|_{_0H_{q,p}^{1,1/2}(\Omega \times \mathbf{R}_+)} \right).$

Step 3. We consider (5.3) in this step. By Proposition 3.6 and (5.8),

$$\begin{split} \| e^{\sigma_0 t} E_0 \nabla Q_q (\mathbf{f} + 2\sigma_0 \mathbf{u}_2) \|_{L_p(\mathbf{R}, L_q(\Omega))} &\leq C \| e^{\sigma_0 t} (\mathbf{f} + 2\sigma_0 \mathbf{u}_2) \|_{L_p(\mathbf{R}_+, L_q(\Omega))} \\ &\leq C_{N, d, p, q, \mu, \sigma_0} \Big(\| e^{\sigma_0 t} \mathbf{f} \|_{L_p(\mathbf{R}_+, L_q(\Omega))} + \| e^{\sigma_0 t} g \|_{W_p^1(\mathbf{R}_+, \widehat{W}_{q, \Gamma}^{-1}(\Omega))} \\ &+ \| e^{\sigma_0 t} (g, \mathbf{h}) \|_{_0 H_{q, p}^{1, 1/2}(\Omega \times \mathbf{R}_+)} \Big), \end{split}$$

which implies $E_0 \nabla Q_q (\mathbf{f} + 2\sigma_0 \mathbf{u}_2) \in L_{p,-\sigma_0}(\mathbf{R}, L_q(\Omega))^N$. One thus observes by Proposition 3.1 with $\delta = \sigma_0$ that there is a solution $(\mathbf{V}, Q) \in W^{2,1}_{q,p,-\sigma_0}(\Omega \times \mathbf{R})^N \times$

$$\begin{split} L_{p,-\sigma_0}(\mathbf{R},W_q^1(\Omega)) \text{ to} \\ \left\{ \begin{array}{ll} \partial_t \mathbf{V} + 2\sigma_0 \mathbf{V} - \operatorname{Div} \mathbf{T}(\mathbf{V},Q) &= E_0 \nabla Q_q(\mathbf{f} + 2\sigma_0 \mathbf{u}_2) & \text{ in } \Omega, \ t \in \mathbf{R}, \\ & \operatorname{div} \mathbf{V} = 0 & \text{ in } \Omega, \ t \in \mathbf{R}, \\ & \mathbf{T}(\mathbf{V},Q)\mathbf{e}_N = 0 & \text{ on } \Gamma, \ t \in \mathbf{R}, \\ & \mathbf{V} = 0 & \text{ on } S, \ t \in \mathbf{R}, \end{array} \right. \end{split}$$

and furthermore, **V** vanishes for t < 0 and

$$\begin{split} \|e^{\sigma_0 t}(\partial_t \mathbf{V}, \mathbf{V}, \nabla \mathbf{V}, \nabla^2 \mathbf{V})\|_{L_p(\mathbf{R}, L_q(\Omega))} + \|e^{\sigma_0 t}Q\|_{L_p(\mathbf{R}, W_q^1(\Omega))} \\ &\leq C_{N, d, p, q, \mu, \sigma_0} \Big(\|e^{\sigma_0 t} \mathbf{f}\|_{L_p(\mathbf{R}_+, L_q(\Omega))} + \|e^{\sigma_0 t}g\|_{W_p^1(\mathbf{R}_+, \widehat{W}_{q, \Gamma}^{-1}(\Omega))} \\ &+ \|e^{\sigma_0 t}(g, \mathbf{h})\|_{_0 H_{q, p}^{1, 1/2}(\Omega \times \mathbf{R}_+)} \Big). \end{split}$$

It is then clear that $(\mathbf{u}_3, \mathbf{p}_3) = (\mathbf{V}, Q)$ solves (5.3) and satisfies

(5.9)
$$\|e^{\sigma_{0}t}(\partial_{t}\mathbf{u}_{3},\mathbf{u}_{3},\nabla\mathbf{u}_{3},\nabla^{2}\mathbf{u}_{3})\|_{L_{p}(\mathbf{R}_{+},L_{q}(\Omega))} + \|e^{\sigma_{0}t}\mathfrak{p}_{3}\|_{L_{p}(\mathbf{R}_{+},W_{q}^{1}(\Omega))}$$

$$\leq C_{N,d,p,q,\mu,\sigma_{0}} \Big(\|e^{\sigma_{0}t}\mathbf{f}\|_{L_{p}(\mathbf{R}_{+},L_{q}(\Omega))} + \|e^{\sigma_{0}t}g\|_{W_{p}^{1}(\mathbf{R}_{+},\widehat{W}_{q,\Gamma}^{-1}(\Omega))}$$

$$+ \|e^{\sigma_{0}t}(g,\mathbf{h})\|_{_{0}H_{q,p}^{1,1/2}(\Omega\times\mathbf{R}_{+})} \Big).$$

Step 4. We consider (5.4) in this step. Let $\mathbf{F} = P_q(\mathbf{f} + 2\sigma_0\mathbf{u}_2) + 2\sigma_0\mathbf{u}_3$. Then, by Proposition 3.6, (5.8), and (5.9), one has $e^{\sigma_0 t}\mathbf{F} \in L_p(\mathbf{R}_+, J_q(\Omega))$ with

(5.10)
$$\|e^{\sigma_0 t} \mathbf{F}\|_{L_p(\mathbf{R}_+, L_q(\Omega))} \leq C_{N, d, p, q, \mu, \sigma_0} \Big(\|e^{\sigma_0 t} \mathbf{f}\|_{L_p(\mathbf{R}_+, L_q(\Omega))} + \|e^{\sigma_0 t} g\|_{W_p^1(\mathbf{R}_+, \widehat{W}_{q, \Gamma}^{-1}(\Omega))} + \|e^{\sigma_0 t}(g, \mathbf{h})\|_{_0 H_{q, p}^{1, 1/2}(\Omega \times \mathbf{R}_+)} \Big).$$

Such an **F** can be approximated by an element of $C_0^{\infty}(\mathbf{R}_+, J_q(\Omega))$ under the norm $\|e^{\sigma_0 t} \cdot \|_{L_p(\mathbf{R}_+, L_q(\Omega))}$, so that it suffices to consider the case $\mathbf{F} \in C_0^{\infty}(\mathbf{R}_+, J_q(\Omega))$.

As the first step, we estimate the solution \mathbf{u}_4 to (5.4) given by Duhamel's formula:

$$\mathbf{u}_4(t) = \int_0^t e^{A_q(t-s)} \mathbf{F}(s) \, ds \quad (t>0),$$

which is also written as

$$e^{\sigma_0 t} \mathbf{u}_4(t) = \int_0^t e^{\sigma_0(t-s)} e^{A_q(t-s)} (e^{\sigma_0 s} \mathbf{F}(s)) \, ds.$$

Let χ_A be the characteristic function of $A \subset \mathbf{R}$, and then by Proposition 4.5

$$\begin{aligned} \|e^{\sigma_{0}t}\mathbf{u}_{4}(t)\|_{L_{q}(\Omega)} &\leq C_{N,d,q,\mu,\sigma_{0}} \int_{0}^{t} e^{-\sigma_{0}(t-s)} \|e^{\sigma_{0}s}\mathbf{F}(s)\|_{L_{q}(\Omega)} \, ds \\ &= C_{N,d,q,\mu,\sigma_{0}} \int_{-\infty}^{t} e^{-\sigma_{0}(t-s)} \|e^{\sigma_{0}s}\mathbf{F}(s)\|_{L_{q}(\Omega)} \, ds \\ &= C_{N,d,q,\mu,\sigma_{0}} \int_{-\infty}^{\infty} \chi_{(0,\infty)}(t-s) e^{-\sigma_{0}(t-s)} \|e^{\sigma_{0}s}\mathbf{F}(s)\|_{L_{q}(\Omega)} \, ds \\ &= C_{N,d,q,\mu,\sigma_{0}} (\chi_{(0,\infty)}(\cdot)e^{-\sigma_{0}\cdot} * (e^{\sigma_{0}\cdot}\mathbf{F}(\cdot)))(t), \end{aligned}$$

where $(f * g)(t) = \int_{\mathbf{R}} f(t - s)g(s) ds$. Combining this inequality with Young's inequality $||f * g||_{L_p(\mathbf{R})} \le ||f||_{L_1(\mathbf{R})} ||g||_{L_p(\mathbf{R})}$ furnishes that

(5.11)
$$\|e^{\sigma_0 t} \mathbf{u}_4\|_{L_p(\mathbf{R}_+, L_q(\Omega))} \le C_{N, d, q, \mu, \sigma_0} \|\chi_{(0, \infty)}(\cdot) e^{-\sigma_0 \cdot}\|_{L_1(\mathbf{R})} \|e^{\sigma_0 t} \mathbf{F}\|_{L_p(\mathbf{R}, L_q(\Omega))}$$

$$\leq C_{N,d,q,\mu,\sigma_0} \| e^{\sigma_0 t} \mathbf{F} \|_{L_p(\mathbf{R}_+,L_q(\Omega))}.$$

As the second step, we write (5.4) as

$$\begin{aligned} \left(\begin{array}{cc} \partial_t \mathbf{u}_4 + 2\sigma_0 \mathbf{u}_4 - \operatorname{Div} \mathbf{T}(\mathbf{u}_4, \mathbf{p}_4) = \mathbf{F} + 2\sigma_0 \mathbf{u}_4 & \text{in } \Omega, \, t > 0, \\ & \operatorname{div} \mathbf{u}_4 = 0 & \operatorname{in} \Omega, \, t > 0, \\ \mathbf{T}(\mathbf{u}_4, \theta_4) \mathbf{e}_N = 0 & \operatorname{on} \Gamma, \, t > 0, \\ & \mathbf{u}_4 = 0 & \operatorname{on} S, \, t > 0, \\ & \mathbf{u}_4 |_{t=0} = 0 & \operatorname{in} \Omega. \end{aligned} \end{aligned}$$

Then, similarly to Step 3, one observes by (5.11) that

$$\begin{aligned} &\|e^{\sigma_0 t}(\partial_t \mathbf{u}_4, \mathbf{u}_4, \nabla \mathbf{u}_4, \nabla^2 \mathbf{u}_4)\|_{L_p(\mathbf{R}_+, L_q(\Omega))} + \|e^{\sigma_0 t} \mathfrak{p}_4\|_{L_p(\mathbf{R}_+, W_q^1(\Omega))} \\ &\leq C_{N,d,p,q,\mu,\sigma_0} \|e^{\sigma_0 t}(\mathbf{F}, \mathbf{u}_4)\|_{L_p(\mathbf{R}_+, L_q(\Omega))} \\ &\leq C_{N,d,p,q,\mu,\sigma_0} \|e^{\sigma_0 t} \mathbf{F}\|_{L_p(\mathbf{R}_+, L_q(\Omega))}, \end{aligned}$$

which, combined with (5.10), furnishes

(5.12)
$$\|e^{\sigma_0 t}(\partial_t \mathbf{u}_4, \mathbf{u}_4, \nabla \mathbf{u}_4, \nabla^2 \mathbf{u}_4)\|_{L_p(\mathbf{R}_+, L_q(\Omega))} + \|e^{\sigma_0 t} \mathfrak{p}_4\|_{L_p(\mathbf{R}_+, W^1_q(\Omega))}$$

$$\leq C_{N, d, p, q, \mu, \sigma_0} \Big(\|e^{\sigma_0 t} \mathbf{f}\|_{L_p(\mathbf{R}_+, L_q(\Omega))} + \|e^{\sigma_0 t} g\|_{L_p(\mathbf{R}_+, \widehat{W}^{-1}_{q, \Gamma}(\Omega))}$$

$$+ \|e^{\sigma_0 t}(g, \mathbf{h})\|_{_0 H^{1, 1/2}_{q, \Omega}(\Omega \times \mathbf{R}_+)} \Big).$$

Step 5. Let $\mathfrak{p} = \sum_{j=1}^{4} \mathfrak{p}_j$ and $\mathbf{u} = \mathbf{u}_1 + \mathbf{v}$ for $\mathbf{v} = \sum_{j=2}^{4} \mathbf{u}_j$, and then $(\mathbf{u}, \mathfrak{p})$ is a solution to (2.3) and satisfies the required estimate of Theorem 2.4 by (5.5), (5.8), (5.9), and (5.12).

Finally, we consider the initial condition and the uniqueness of solutions. For two Banach spaces X, Y, the symbol $X \hookrightarrow Y$ means that X is continuously injected in Y, i.e. there is a positive constant C such that $||f||_Y \leq C||f||_X$ for any $f \in X$. It is then well-known by e.g. [5, Theorem 4.10.2] that

(5.13)
$$W_p^1(\mathbf{R}_+, X_0) \cap L_p(\mathbf{R}_+, X_1) \hookrightarrow BUC([0, \infty), (X_0, X_1)_{1-1/p, p})$$

for Banach spaces X_0 , X_1 satisfying the properties: $X_1 \hookrightarrow X_0$ and X_1 is dense in X_0 . One makes use of (5.13) with $X_0 = L_q(\Omega)$ and $X_1 = W_q^2(\Omega)$ in order to obtain

(5.14)
$$W^{2,1}_{q,p}(\Omega \times \mathbf{R}_+) \hookrightarrow BUC([0,\infty), B^{2-2/p}_{q,p}(\Omega)),$$

which implies that $\mathbf{v}|_{t=0} = 0$ in $B_{q,p}^{2-2/p}(\Omega)^N$. We thus conclude $\lim_{t\to 0+} \|\mathbf{u}(t) - \mathbf{a}\|_{B_{q,p}^{2-2/p}(\Omega)} = 0$. The uniqueness of solutions to (2.3) follows from the existence of solutions for a dual problem (cf. e.g. [30, Subsection 7.2]), which completes the proof of Theorem 2.4.

6. Proof of Theorem 2.2

This section proves Theorem 2.2. To this end, we set for $1 < q < \infty$

$$B_q(\Omega \times \mathbf{R}_+) = W^1_{\infty}(\mathbf{R}_+, L_{\infty}(\Omega)) \cap L_{\infty}(\mathbf{R}_+, W^1_q(\Omega))$$

and start with

Lemma 6.1. Let $1 < p, q < \infty$. Then the following assertions hold true. (1) For any $f \in W^1_{\infty}(\mathbf{R}_+, L_{\infty}(\Omega))$ and $g \in H^{1/2}_p(\mathbf{R}_+, L_q(\Omega))$,

 $\|fg\|_{H_p^{1/2}(\mathbf{R}_+, L_q(\Omega))} \le \|f\|_{W_{\infty}^1(\mathbf{R}_+, L_{\infty}(\Omega))} \|g\|_{H_p^{1/2}(\mathbf{R}_+, L_q(\Omega))}.$

(2) Let q > N. Then there exists a positive constant $C_{N,q}$ such that for any $f, g \in$ $B_q(\Omega \times \mathbf{R}_+)$ and $h \in H^{1,1/2}_{q,p}(\Omega \times \mathbf{R}_+),$

 $\|fg\|_{B_{q}(\Omega \times \mathbf{R}_{+})} \leq C_{N,q} \|f\|_{B_{q}(\Omega \times \mathbf{R}_{+})} \|g\|_{B_{q}(\Omega \times \mathbf{R}_{+})},$ $h\|_{M,Q} \leq C_{N,q} \|f\|_{B_{q}(\Omega \times \mathbf{R}_{+})} \|h\|_{H^{1,1/2}(\Omega \times \mathbf{R}_{+})}.$ +).

$$\|J^{II}\|_{H^{1,1/2}_{q,p}}(\Omega \times \mathbf{R}_{+}) \geq \mathbb{C}_{N,q}\|J\|_{B_{q}}(\Omega \times \mathbf{R}_{+})\|I^{II}\|_{H^{1,1/2}_{q,p}}(\Omega \times \mathbf{R}_{+})\|I^{I}$$

(3) For any $f \in H_p^{1/2}(\mathbf{R}_+, W_q^1(\Omega)),$

$$\|\nabla f\|_{H_p^{1/2}(\mathbf{R}_+, L_q(\Omega))} \le \|f\|_{H_p^{1/2}(\mathbf{R}_+, W_q^1(\Omega))}$$

Proof. (1). It clearly holds that

$$\|fg\|_{L_p(\mathbf{R}_+,L_q(\Omega))} \le \|f\|_{W^1_{\infty}(\mathbf{R}_+,L_{\infty}(\Omega))} \|g\|_{L_p(\mathbf{R}_+,L_q(\Omega))},$$

$$\|fg\|_{W^1_{n}(\mathbf{R}_+,L_q(\Omega))} \le \|f\|_{W^1_{\infty}(\mathbf{R}_+,L_{\infty}(\Omega))} \|g\|_{W^1_{n}(\mathbf{R}_+,L_q(\Omega))}.$$

Let $T_f g = fg$, and then $T_f \in \mathcal{L}(L_p(\mathbf{R}_+, L_q(\Omega)) \cap \mathcal{L}(W_p^1(\mathbf{R}_+, L_q(\Omega)))$ with

 $||T_f||_{\mathcal{L}(L_p(\mathbf{R}_+, L_q(\Omega)))} \le ||f||_{W^1_{\infty}(\mathbf{R}_+, L_{\infty}(\Omega))},$ $||T_f||_{\mathcal{L}(W^1_n(\mathbf{R}_+, L_q(\Omega)))} \le ||f||_{W^1_\infty(\mathbf{R}_+, L_\infty(\Omega))}.$

Combining these properties with the complex interpolation method furnishes that

$$\begin{aligned} \|T_f g\| [L_p(\mathbf{R}_+, L_q(\Omega), W_p^1(\mathbf{R}_+, L_q(\Omega)))]_{1/2} \\ &\leq \|f\|_{W_\infty^1(\mathbf{R}_+, L_\infty(\Omega))} \|g\| [L_p(\mathbf{R}_+, L_q(\Omega), W_p^1(\mathbf{R}_+, L_q(\Omega)))]_{1/2}. \end{aligned}$$

Noting $[L_p(\mathbf{R}_+, L_q(\Omega), W_p^1(\mathbf{R}_+, L_q(\Omega)))]_{1/2} = H_p^{1/2}(\mathbf{R}_+, L_q(\Omega))$ by the definition, we complete the proof of Lemma 6.1(1).

(2). Since $W_q^1(\Omega)$ is a Banach algebra for q > N, we have

(6.1)
$$\|fg\|_{L_{\infty}(\mathbf{R}_{+},W^{1}_{q}(\Omega))} \leq C_{N,q} \|f\|_{L_{\infty}(\mathbf{R}_{+},W^{1}_{q}(\Omega))} \|g\|_{L_{\infty}(\mathbf{R}_{+},W^{1}_{q}(\Omega))}, \|fh\|_{L_{p}(\mathbf{R}_{+},W^{1}_{q}(\Omega))} \leq C_{N,q} \|f\|_{L_{\infty}(\mathbf{R}_{+},W^{1}_{q}(\Omega))} \|h\|_{L_{p}(\mathbf{R}_{+},W^{1}_{q}(\Omega))}.$$

It is clear that $||fg||_{W^1_{\infty}(\mathbf{R}_+, L_{\infty}(\Omega))} \leq ||f||_{W^1_{\infty}(\mathbf{R}_+, L_{\infty}(\Omega))} ||g||_{W^1_{\infty}(\mathbf{R}_+, L_{\infty}(\Omega))}$, so that one has, together with the first inequality of (6.1),

$$\|fg\|_{B_q(\Omega\times\mathbf{R}_+)} \le C_{N,q} \|f\|_{B_q(\Omega\times\mathbf{R}_+)} \|g\|_{B_q(\Omega\times\mathbf{R}_+)}.$$

The other estimate of Lemma 6.1(2) follows from Lemma 6.1(1) and the second inequality of (6.1).

(3). The required inequality follows from the complex interpolation method immediately, which completes the proof of Lemma 6.1 (3).

At this point, we introduce some embedding properties.

Lemma 6.2. Let p, q satisfy (2.1). Then there exists a positive constant $M_1 \ge 1$ such that the following assertions hold true.

 $(1) \ \|f\|_{BUC(\overline{\Omega})} \le M_1 \|f\|_{W^1_q(\Omega)} \ \text{for any} \ f \in W^1_q(\Omega).$

(2) $\|f\|_{BUC([0,\infty),BUC^1(\overline{\Omega}))} \leq M_1 \|f\|_{W^{2,1}_{q,p}(\Omega \times \mathbf{R}_+)}$ for any $f \in W^{2,1}_{q,p}(\Omega \times \mathbf{R}_+)$.

(3)
$$\|f\|_{H_p^{1/2}(\mathbf{R}_+, W_q^1(\Omega))} \le M_1 \|f\|_{W_{q,p}^{2,1}(\Omega \times \mathbf{R}_+)}$$
 for any $f \in W_{q,p}^{2,1}(\Omega \times \mathbf{R}_+)$.

Proof. (1). See e.g. [3, Theorem 4.12].

(2). It follows from (5.14) and $B_{q,p}^{2-2/p}(\Omega) \hookrightarrow BUC^1(\overline{\Omega})$ under the condition (2.1) (cf. [51, Theorem 4.6.1]).

(3). See e.g. [26, Proposition 3.2, Remark 3.3], [37].

We prove Theorem 2.2 in the remaining part of this section by using Theorem 2.4 and the contraction mapping theorem. Let $\gamma_0 = \sigma_0$, where σ_0 is the same positive constant as in Theorem 2.4. We here define a closed set $X_{q,p}^{\gamma_0}(R)$ by

$$\begin{aligned} X_{q,p}^{\gamma_0}(R) &= \{ (\mathbf{u}, \mathfrak{p}) \in W_{q,p}^{2,1}(\Omega \times \mathbf{R}_+)^N \times L_p(\mathbf{R}_+, W_q^1(\Omega)) \mid \| (\mathbf{u}, \mathfrak{p}) \|_{X_{q,p}^{\gamma_0}} \leq R, \\ \mathbf{u} &= 0 \text{ on } S, \ \lim_{t \to 0+} \| \mathbf{u} - \mathbf{a} \|_{B_{q,p}^{2-2/p}(\Omega)} = 0 \}, \end{aligned}$$

where 0 < R < 1 is a positive constant and $\|(\mathbf{u}, \mathfrak{p})\|_{X^{\gamma_0}_{q,p}} = \|\mathbf{u}\|_{Y^{\gamma_0}_{q,p}} + \|\mathfrak{p}\|_{Z^{\gamma_0}_{q,p}}$ with

$$\|\mathbf{u}\|_{Y^{\gamma_0}_{q,p}} = \|e^{\gamma_0 t}(\partial_t \mathbf{u}, \mathbf{u}, \nabla \mathbf{u}, \nabla^2 \mathbf{u})\|_{L_p(\mathbf{R}_+, L_q(\Omega))}, \quad \|\mathfrak{p}\|_{Z^{\gamma_0}_{q,p}} = \|e^{\gamma_0 t}\mathfrak{p}\|_{L_p(\mathbf{R}_+, W^1_q(\Omega))}.$$

In addition, by e.g. [23], we know the following characterization:

(6.2)
$${}_{0}H^{1,1/2}_{q,p}(\Omega \times \mathbf{R}_{+}) = \{ f \in H^{1,1/2}_{q,p}(\Omega \times \mathbf{R}_{+}) \mid f|_{t=0} = 0 \text{ in } L_{q}(\Omega) \}$$

under the condition (2.1).

Let $(\mathbf{v}, \mathbf{q}) \in X_{q,p}^{\gamma_0}(R)$, and one considers

(6.3)
$$\begin{cases} \partial_t \mathbf{u} - \operatorname{Div} \mathbf{T}(\mathbf{u}, \mathbf{p}) = \mathbf{F}(\mathbf{v}) & \text{in } \Omega, \ t > 0, \\ \operatorname{div} \mathbf{u} = G(\mathbf{u}) = \operatorname{div} \mathbf{G}(\mathbf{v}) & \text{in } \Omega, \ t > 0, \\ \mathbf{T}(\mathbf{u}, \mathbf{p}) \mathbf{e}_N = \mathbf{H}(\mathbf{v}) \mathbf{e}_N & \text{on } \Gamma, \ t > 0, \\ \mathbf{u} = 0 & \text{on } S, \ t > 0, \\ \mathbf{u}|_{t=0} = \mathbf{a} & \text{in } \Omega. \end{cases}$$

Here we prove

Lemma 6.3. Let p, q satisfy (2.1), and let γ_0 be as above. Then there exists positive constants $M_3 \ge M_2 \ge 1$ such that the following assertions hold true. (1) For any \mathbf{v}_r with $\|\mathbf{v}_r\|_{Y^{\gamma_0}_{q,p}} \le 1$ (r = 1, 2), we have

$$\begin{split} \left\| \mathbf{U}_{1} \left(\int_{0}^{t} \nabla \mathbf{v}_{2}(\xi, s) \, ds \right) - \mathbf{U}_{1} \left(\int_{0}^{t} \nabla \mathbf{v}_{1}(\xi, s) \, ds \right) \right\|_{L_{\infty}(\mathbf{R}_{+}, L_{\infty}(\Omega))} \\ &\leq M_{2} \| \mathbf{v}_{2} - \mathbf{v}_{1} \|_{Y_{q,p}^{\gamma_{0}}}, \\ &\sum_{i,j,k,l=1}^{N} \left\| V_{ijk}^{l} \left(\int_{0}^{t} \nabla \mathbf{v}_{2}(\xi, s) \, ds \right) - V_{ijk}^{l} \left(\int_{0}^{t} \nabla \mathbf{v}_{1}(\xi, s) \, ds \right) \right\|_{L_{\infty}(\mathbf{R}_{+}, L_{\infty}(\Omega))} \\ &\leq M_{2} \| \mathbf{v}_{2} - \mathbf{v}_{1} \|_{Y_{q,p}^{\gamma_{0}}}, \\ &\sum_{i,j,k,l,n=1}^{N} \left\| W_{ijklm}^{n} \left(\int_{0}^{t} \nabla \mathbf{v}_{2}(\xi, s) \, ds \right) - W_{ijklm}^{n} \left(\int_{0}^{t} \nabla \mathbf{v}_{1}(\xi, s) \, ds \right) \right\|_{L_{\infty}(\mathbf{R}_{+}, L_{\infty}(\Omega))} \\ &\leq M_{2} \| \mathbf{v}_{2} - \mathbf{v}_{1} \|_{Y_{q,p}^{\gamma_{0}}}, \\ &\left\| \mathbf{U}_{r} \left(\int_{0}^{t} \nabla \mathbf{v}_{2}(\xi, s) \, ds \right) - \mathbf{U}_{r} \left(\int_{0}^{t} \nabla \mathbf{v}_{1}(\xi, s) \, ds \right) \right\|_{B_{q}(\Omega \times \mathbf{R}_{+})} \\ &\leq M_{2} \| \mathbf{v}_{2} - \mathbf{v}_{1} \|_{Y_{q,p}^{\gamma_{0}}} \quad (r = 2, \dots, 7). \end{split}$$

$$(2) For any \mathbf{v} with \| \mathbf{v} \|_{Y_{q,p}^{\gamma_{0}}} \leq 1, we have$$

$$\left\| \mathbf{U}_1\left(\int_0^t \nabla \mathbf{v}(\xi, s) \, ds \right) \right\|_{L_\infty(\mathbf{R}_+, L_\infty(\Omega))} \le M_2 \|\mathbf{v}\|_{Y^{\gamma_0}_{q, p}},$$

HIROKAZU SAITO

$$\sum_{i,j,k,l=1}^{N} \left\| V_{ijk}^{l} \left(\int_{0}^{t} \nabla \mathbf{v}(\xi,s) \, ds \right) \right\|_{L_{\infty}(\mathbf{R}_{+},L_{\infty}(\Omega))} \leq M_{2} \| \mathbf{v} \|_{Y_{q,p}^{\gamma_{0}}}, \\ \left\| \mathbf{U}_{r} \left(\int_{0}^{t} \nabla \mathbf{v}(\xi,s) \, ds \right) \right\|_{B_{q}(\Omega \times \mathbf{R}_{+})} \leq M_{2} \| \mathbf{v} \|_{Y_{q,p}^{\gamma_{0}}} \quad (r = 2, \dots, 7).$$

(3) For any **v** with $\|\mathbf{v}\|_{Y^{\gamma_0}_{q,p}} \leq 1$, we have

$$\sum_{i,j,k,l,m,n=1}^{N} \left\| W_{ijklm}^{n} \left(\int_{0}^{t} \nabla \mathbf{v}(\xi,s) \, ds \right) \right\|_{L_{\infty}(\mathbf{R}_{+},L_{\infty}(\Omega))} \leq M_{3},$$

Proof. For $F(\mathbf{X})$ with $\mathbf{X} = (X_{\alpha\beta}) \in \mathbf{R}^{N \times N}$, we denote the derivative of $F(\mathbf{X})$ with respect to $X_{\alpha\beta}$ by $\partial_{(\alpha,\beta)}F(\mathbf{X})$ for $(\alpha,\beta) \in \mathcal{N} := \{1,\ldots,N\} \times \{1,\ldots,N\}$. In addition, we choose $m_2 \geq 1$ large enough so that for $r_1 = 1,\ldots,7$ and $r_2 = 2,\ldots,7$

$$\max\left\{\sum_{(\alpha,\beta)\in\mathcal{N}} \left|\partial_{(\alpha,\beta)}\mathbf{U}_{r_{1}}\left(\mathbf{X}\right)\right| \mid |\mathbf{X}| \leq 2M_{1}(p'\gamma_{0})^{-1/p'}\right\} \leq m_{2},\\ \max\left\{\sum_{(\alpha',\beta')\in\mathcal{N}}\sum_{(\alpha,\beta)\in\mathcal{N}} \left|\partial_{(\alpha',\beta')}\partial_{(\alpha,\beta)}\mathbf{U}_{r_{2}}\left(\mathbf{X}\right)\right| \mid |\mathbf{X}| \leq 2M_{1}(p'\gamma_{0})^{-1/p'}\right\} \leq m_{2},\\ \max\left\{\sum_{(\alpha,\beta)\in\mathcal{N}}\sum_{i,j,k,l=1}^{N} \left|\partial_{(\alpha,\beta)}V_{ijk}^{l}\left(\mathbf{X}\right)\right| \mid |\mathbf{X}| \leq 2M_{1}(p'\gamma_{0})^{-1/p'}\right\} \leq m_{2},\\ \max\left\{\sum_{(\alpha,\beta)\in\mathcal{N}}\sum_{i,j,k,l,m,n=1}^{N} \left|\partial_{(\alpha,\beta)}W_{ijklm}^{n}\left(\mathbf{X}\right)\right| \mid |\mathbf{X}| \leq 2M_{1}(p'\gamma_{0})^{-1/p'}\right\} \leq m_{2},\right\}$$

where p' = p/(p-1) and M_1 is the same positive constant as in Lemma 6.2. (1). First, we consider \mathbf{U}_1 . Let us write $\mathbf{v}_r = (v_{r1}, \dots, v_{rN})^{\mathsf{T}}$ for r = 1, 2. It then holds that

(6.4)
$$\mathbf{U}_{1}\left(\int_{0}^{t} \nabla \mathbf{v}_{2}(\xi, s) \, ds\right) - \mathbf{U}_{1}\left(\int_{0}^{t} \nabla \mathbf{v}_{1}(\xi, s) \, ds\right)$$
$$= \sum_{(\alpha, \beta) \in \mathcal{N}} \int_{0}^{1} (\partial_{(\alpha, \beta)} \mathbf{U}_{1}) \left(\theta \int_{0}^{t} \nabla \mathbf{v}_{2}(\xi, s) \, ds + (1 - \theta) \int_{0}^{t} \nabla \mathbf{v}_{1}(\xi, s) \, ds\right) \, d\theta$$
$$\cdot \left(\int_{0}^{t} \partial_{\beta} v_{2\alpha}(\xi, s) \, ds - \int_{0}^{t} \partial_{\beta} v_{1\alpha}(\xi, s) \, ds\right)$$

and that by Lemma 6.2(1)

(6.5)
$$\left\| \theta \int_{0}^{t} \nabla \mathbf{v}_{2}(\xi, s) \, ds + (1 - \theta) \int_{0}^{t} \nabla \mathbf{v}_{1}(\xi, s) \, ds \right\|_{L_{\infty}(\mathbf{R}_{+}, L_{\infty}(\Omega))}$$

$$\leq \sum_{r=1}^{2} \int_{0}^{\infty} \| \nabla \mathbf{v}_{r}(\cdot, s) \|_{L_{\infty}(\Omega)} \, ds \leq M_{1} \sum_{r=1}^{2} \int_{0}^{\infty} \| \mathbf{v}_{r}(\cdot, s) \|_{W_{q}^{2}(\Omega)} \, ds$$

$$\leq M_{1} \sum_{r=1}^{2} \left(\int_{0}^{\infty} e^{-p' \gamma_{0} s} \, ds \right)^{1/p'} \| \mathbf{v}_{r} \|_{Y_{q,p}^{\gamma_{0}}} \leq 2M_{1} (p' \gamma_{0})^{-1/p'}.$$

One similarly obtains

(6.6)
$$\left\| \int_{0}^{t} \nabla \mathbf{y}(\xi, s) \, ds \right\|_{L_{\infty}(\mathbf{R}_{+}, L_{\infty}(\Omega))} \leq M_{1}(p'\gamma_{0})^{-1/p'} \|\mathbf{y}\|_{Y^{\gamma_{0}}_{q, p}},$$

(6.7)
$$\left\| \int_0^t \nabla \mathbf{y}(\xi, s) \, ds \right\|_{L_{\infty}(\mathbf{R}_+, W^1_q(\Omega))} \le (p'\gamma_0)^{-1/p'} \|\mathbf{y}\|_{Y^{\gamma_0}_{q, p}}.$$

Combining (6.4) with (6.5) and (6.6) furnishes that

$$\left\| \mathbf{U}_1\left(\int_0^t \nabla \mathbf{v}_2(\xi, s) \, ds\right) - \mathbf{U}_1\left(\int_0^t \nabla \mathbf{v}_1(\xi, s) \, ds\right) \right\|_{L_{\infty}(\mathbf{R}_+, L_{\infty}(\Omega))}$$

$$\leq m_2 M_1(p'\gamma_0)^{-1/p'} \|\mathbf{v}_2 - \mathbf{v}_1\|_{Y^{\gamma_0}_{q, p}}.$$

Thus we choose $M_2 \geq 1$ large enough so that $M_2 \geq m_2 M_1 (p'\gamma_0)^{-1/p'}$ in order to obtain the required estimate for \mathbf{U}_1 . Analogously, one can prove that the required estimates hold true for V_{ijk}^l and W_{ijklm}^n . Next, we consider \mathbf{U}_r for $r = 2, \ldots, 7$. Following (6.4), we observe that for

 $\partial \in \{\partial_t, \partial_1, \ldots, \partial_N\}$

$$\begin{split} &\partial \left\{ \mathbf{U}_r \left(\int_0^t \nabla \mathbf{v}_2(\xi, s) \, ds \right) - \mathbf{U}_r \left(\int_0^t \nabla \mathbf{v}_1(\xi, s) \, ds \right) \right\} \\ &= \sum_{(\alpha', \beta') \in \mathcal{N}} \sum_{(\alpha, \beta) \in \mathcal{N}} \\ &\int_0^1 (\partial_{(\alpha', \beta')} \partial_{(\alpha, \beta)} \mathbf{U}_r) \left(\theta \int_0^t \nabla \mathbf{v}_2(\xi, s) \, ds + (1 - \theta) \int_0^t \nabla \mathbf{v}_1(\xi, s) \, ds \right) \\ &\cdot \partial \left(\theta \int_0^t \partial_{\beta'} v_{2\alpha'}(\xi, s) \, ds + (1 - \theta) \int_0^t \partial_{\beta'} v_{1\alpha'}(\xi, s) \, ds \right) \, d\theta \\ &\cdot \left(\int_0^t \partial_{\beta} v_{2\alpha}(\xi, s) \, ds - \int_0^t \partial_{\beta} v_{1\alpha}(\xi, s) \, ds \right) \\ &+ \sum_{(\alpha, \beta) \in \mathcal{N}} \int_0^1 (\partial_{(\alpha, \beta)} \mathbf{U}_r) \left(\theta \int_0^t \nabla \mathbf{v}_2(\xi, s) \, ds + (1 - \theta) \int_0^t \nabla \mathbf{v}_1(\xi, s) \, ds \right) \, d\theta \\ &\cdot \partial \left(\int_0^t \partial_{\beta} v_{2\alpha}(\xi, s) \, ds - \int_0^t \partial_{\beta} v_{1\alpha}(\xi, s) \, ds \right). \end{split}$$

By (6.5), (6.6), (6.7), and Lemma 6.2 (2), it holds that

$$\begin{aligned} \left\| \partial_t \left\{ \mathbf{U}_r \left(\int_0^t \nabla \mathbf{v}_2(\xi, s) \, ds \right) - \mathbf{U}_r \left(\int_0^t \nabla \mathbf{v}_1(\xi, s) \, ds \right) \right\} \right\|_{L_{\infty}(\mathbf{R}_+, L_{\infty}(\Omega))} \\ &\leq \left(2m_2 M_1^2 (p'\gamma_0)^{-1/p'} + m_2 M_1 \right) \|\mathbf{v}_2 - \mathbf{v}_1\|_{Y_{q,p}^{\gamma_0}}, \\ \left\| \partial_j \left\{ \mathbf{U}_r \left(\int_0^t \nabla \mathbf{v}_2(\xi, s) \, ds \right) - \mathbf{U}_r \left(\int_0^t \nabla \mathbf{v}_1(\xi, s) \, ds \right) \right\} \right\|_{L_{\infty}(\mathbf{R}_+, L_q(\Omega))} \\ &\leq \left(2m_2 M_1 (p'\gamma_0)^{-2/p'} + m_2 (p'\gamma_0)^{-1/p'} \right) \|\mathbf{v}_2 - \mathbf{v}_1\|_{Y_{q,p}^{\gamma_0}}, \end{aligned}$$

for $j = 1, \ldots, N$. Similarly to the case \mathbf{U}_1 , we also have

$$\begin{aligned} \left\| \mathbf{U}_r \left(\int_0^t \nabla \mathbf{v}_2(\xi, s) \, ds \right) - \mathbf{U}_r \left(\int_0^t \nabla \mathbf{v}_1(\xi, s) \, ds \right) \right\|_{L_{\infty}(\mathbf{R}_+, L_{\infty}(\Omega))} \\ &\leq m_2 M_1(p'\gamma_0)^{-1/p'} \| \mathbf{v}_2 - \mathbf{v}_1 \|_{Y^{\gamma_0}_{q,p}}, \\ \left\| \mathbf{U}_r \left(\int_0^t \nabla \mathbf{v}_2(\xi, s) \, ds \right) - \mathbf{U}_r \left(\int_0^t \nabla \mathbf{v}_1(\xi, s) \, ds \right) \right\|_{L_{\infty}(\mathbf{R}_+, L_q(\Omega))} \\ &\leq m_2(p'\gamma_0)^{-1/p'} \| \mathbf{v}_2 - \mathbf{v}_1 \|_{Y^{\gamma_0}_{q,p}}. \end{aligned}$$

One thus obtains the required estimates for \mathbf{U}_r (r = 2, ..., 7) by choosing a larger $M_2 \ge 1$ if necessary. This completes the proof of Lemma 6.3 (1).

(2). The estimates follows from (1.15) and (1) with $(\mathbf{v}_2, \mathbf{v}_1) = (\mathbf{v}, 0)$ immediately. This completes the proof of Lemma 6.3 (2).

(3). One sets $(\mathbf{v}_2, \mathbf{v}_1) = (\mathbf{v}, 0)$ in the inequality proved in (1) in order to obtain

$$\sum_{\substack{i,j,k,l,\\m,n=1}}^{N} \left\| W_{ijklm}^{n} \left(\int_{0}^{t} \nabla \mathbf{v}(\xi,s) \, ds \right) - W_{ijklm}^{n} \left(\mathbf{O} \right) \right\|_{L_{\infty}(\mathbf{R}_{+},L_{\infty}(\Omega))} \le M_{2} \| \mathbf{v} \|_{Y_{q,p}^{\gamma_{0}}} \le M_{2},$$

which implies that

$$\sum_{i,j,k,l,m,n=1}^{N} \left\| W_{ijklm}^{n} \left(\int_{0}^{t} \nabla \mathbf{v}(\xi,s) \, ds \right) \right\|_{L_{\infty}(\mathbf{R}_{+},L_{\infty}(\Omega))}$$

$$\leq M_{2} + \sum_{i,j,k,l,m,n=1}^{N} \left\| W_{ijklm}^{n}(\mathbf{O}) \right\|_{L_{\infty}(\mathbf{R}_{+},L_{\infty}(\Omega))}.$$

Noting $W_{ijklm}^{n}(\mathbf{O}) \in L_{\infty}(\mathbf{R}_{+}, L_{\infty}(\Omega))$, we complete the proof of Lemma 6.3 (3) with $M_{3} = M_{2} + \sum_{i,j,k,l,m,n=1}^{N} \|W_{ijklm}^{n}(\mathbf{O})\|_{L_{\infty}(\mathbf{R}_{+}, L_{\infty}(\Omega))}$.

Let M_1 , M_2 , and M_3 be the same positive constants as in Lemmas 6.2, Lemma 6.3, and also assume that $(\mathbf{v}, \mathbf{q}), (\mathbf{v}_1, \mathbf{q}_1), (\mathbf{v}_2, \mathbf{q}_2) \in X_{q,p}^{\gamma_0}(1)$ in what follows. **Estimates of F(v).** One observes that

$$\begin{split} e^{\gamma_0 t} (\mathbf{F}(\mathbf{v}_2) - \mathbf{F}(\mathbf{v}_1)) \\ &= \left\{ \mathbf{U}_1 \left(\int_0^t \nabla \mathbf{v}_2 \, ds \right) - \mathbf{U}_1 \left(\int_0^t \nabla \mathbf{v}_1 \, ds \right) \right\} e^{\gamma_0 t} \partial_t \mathbf{v}_2 \\ &+ \mathbf{U}_1 \left(\int_0^t \nabla \mathbf{v}_1 \, ds \right) e^{\gamma_0 t} \partial_t (\mathbf{v}_2 - \mathbf{v}_1) \\ &+ \left\{ \mathbf{V} \left(\int_0^t \nabla \mathbf{v}_2 \, ds \right) - \mathbf{V} \left(\int_0^t \nabla \mathbf{v}_1 \, ds \right) \right\} e^{\gamma_0 t} \nabla^2 \mathbf{v}_2 \\ &+ \mathbf{V} \left(\int_0^t \nabla \mathbf{v}_1 \, ds \right) e^{\gamma_0 t} \nabla^2 (\mathbf{v}_2 - \mathbf{v}_1) \\ &+ \left[\left\{ \mathbf{W} \left(\int_0^t \nabla \mathbf{v}_2 \, ds \right) - \mathbf{W} \left(\int_0^t \nabla \mathbf{v}_1 \, ds \right) \right\} \int_0^t \nabla^2 \mathbf{v}_2 \, ds \right] e^{\gamma_0 t} \nabla \mathbf{v}_2 \\ &+ \left[\mathbf{W} \left(\int_0^t \nabla \mathbf{v}_1 \, ds \right) \int_0^t \nabla^2 (\mathbf{v}_2 - \mathbf{v}_1) \, ds \right] e^{\gamma_0 t} \nabla \mathbf{v}_2 \end{split}$$

+
$$\left[\mathbf{W} \left(\int_0^t \nabla \mathbf{v}_1 \, ds \right) \int_0^t \nabla^2 \mathbf{v}_1 \, ds \right] e^{\gamma_0 t} \nabla (\mathbf{v}_2 - \mathbf{v}_1)$$

=: $I_1 + \dots + I_7$.

By Lemma 6.3, we easily have

$$\|I_{j}\|_{L_{p}(\mathbf{R}_{+},L_{q}(\Omega))} \leq M_{2}\left(\|\mathbf{v}_{2}\|_{Y_{q,p}^{\gamma_{0}}} + \|\mathbf{v}_{1}\|_{Y_{q,p}^{\gamma_{0}}}\right)\|\mathbf{v}_{2} - \mathbf{v}_{1}\|_{Y_{q,p}^{\gamma_{0}}} \quad (j = 1, \dots, 4).$$

On the other hand, by Lemma 6.2(1)

$$\left\|e^{\gamma_0 t} \nabla \mathbf{v}_2\right\|_{L_p(\mathbf{R}_+, L_{\infty}(\Omega))} \leq M_1 \|e^{\gamma_0 t} \nabla \mathbf{v}_2\|_{L_p(\mathbf{R}_+, W^1_q(\Omega))} \leq M_1 \|\mathbf{v}_2\|_{Y^{\gamma_0}_{q, p}} \leq M_1,$$
which, combined with Lemma 6.3 and (6.7), furnishes

$$\begin{aligned} \|I_5\|_{L_p(\mathbf{R}_+,L_q(\Omega))} &\leq \left\| \mathbf{W} \left(\int_0^t \nabla \mathbf{v}_2 \, ds \right) - \mathbf{W} \left(\int_0^t \nabla \mathbf{v}_1 \, ds \right) \right\|_{L_\infty(\mathbf{R}_+,L_\infty(\Omega))} \\ &\cdot \left\| \int_0^t \nabla^2 \mathbf{v}_2 \, ds \right\|_{L_\infty(\mathbf{R}_+,L_q(\Omega))} \left\| e^{\gamma_0 t} \nabla \mathbf{v}_2 \right\|_{L_p(\mathbf{R}_+,L_\infty(\Omega))} \\ &\leq (p'\gamma_0)^{-1/p'} M_1 M_2 \left(\|\mathbf{v}_2\|_{Y^{\gamma_0}_{q,p}} + \|\mathbf{v}_1\|_{Y^{\gamma_0}_{q,p}} \right) \|\mathbf{v}_2 - \mathbf{v}_1\|_{Y^{\gamma_0}_{q,p}}. \end{aligned}$$

Analogously, it holds that for j = 6, 7

$$\|I_{j}\|_{L_{p}(\mathbf{R}_{+},L_{q}(\Omega))} \leq (p'\gamma_{0})^{-1/p'} M_{1}M_{3}\left(\|\mathbf{v}_{2}\|_{Y_{q,p}^{\gamma_{0}}} + \|\mathbf{v}_{1}\|_{Y_{q,p}^{\gamma_{0}}}\right)\|\mathbf{v}_{2} - \mathbf{v}_{1}\|_{Y_{q,p}^{\gamma_{0}}}.$$

Summing up the above estimates for I_1, \ldots, I_7 , we have achieved

(6.8)
$$\|e^{\gamma_0 t} (\mathbf{F}(\mathbf{v}_2) - \mathbf{F}(\mathbf{v}_1))\|_{L_p(\mathbf{R}_+, L_q(\Omega))}$$

$$\leq \left(4M_2 + (p'\gamma_0)^{-1/p'} M_1 M_2 + 2(p'\gamma_0)^{-1/p'} M_1 M_3 \right)$$

$$\cdot \left(\|\mathbf{v}_2\|_{Y^{\gamma_0}_{q,p}} + \|\mathbf{v}_1\|_{Y^{\gamma_0}_{q,p}} \right) \|\mathbf{v}_2 - \mathbf{v}_1\|_{Y^{\gamma_0}_{q,p}}.$$

Especially, setting $(\mathbf{v}_2, \mathbf{v}_1) = (\mathbf{v}, 0)$ in (6.8) yields

(6.9)
$$\|e^{\gamma_0 t} \mathbf{F}(\mathbf{v})\|_{L_p(\mathbf{R}_+, L_q(\Omega))} \leq \left(4M_2 + (p'\gamma_0)^{-1/p'} M_1 M_2 + 2(p'\gamma_0)^{-1/p'} M_1 M_3\right) \|\mathbf{v}\|_{Y_{q,p}^{\gamma_0}}^2.$$

Estimates of $G(\mathbf{v})$, $\mathbf{G}(\mathbf{v})$. One observes that

$$e^{\gamma_0 t}(G(\mathbf{v}_2) - G(\mathbf{v}_1)) = \left\{ \mathbf{U}_2\left(\int_0^t \nabla \mathbf{v}_2 \, ds\right) - \mathbf{U}_2\left(\int_0^t \nabla \mathbf{v}_1 \, ds\right) \right\} : e^{\gamma_0 t} \nabla \mathbf{v}_2$$
$$+ \mathbf{U}_2\left(\int_0^t \nabla \mathbf{v}_1 \, ds\right) : e^{\gamma_0 t}(\nabla \mathbf{v}_2 - \nabla \mathbf{v}_1) =: I_8 + I_9.$$

By Lemmas 6.1, 6.3, we have

$$\begin{split} &\|I_8\|_{H^{1,1/2}_{q,p}(\Omega \times \mathbf{R}_+)} \\ &\leq \left\|\mathbf{U}_2\left(\int_0^t \nabla \mathbf{v}_2 \, ds\right) - \mathbf{U}_2\left(\int_0^t \nabla \mathbf{v}_1 \, ds\right)\right\|_{B_q(\Omega \times \mathbf{R}_+)} \|e^{\gamma_0 t} \nabla \mathbf{v}_2\|_{H^{1,1/2}_{q,p}(\Omega \times \mathbf{R}_+)} \\ &\leq M_2 \|\mathbf{v}_2 - \mathbf{v}_1\|_{Y^{\gamma_0}_{q,p}} \left(\|e^{\gamma_0 t} \mathbf{v}_2\|_{H^{1/2}_p(\mathbf{R}_+,W^1_q(\Omega))} + \|e^{\gamma_0 t} \mathbf{v}_2\|_{L_p(\mathbf{R}_+,W^2_q(\Omega))}\right). \end{split}$$

Since it holds by Lemma 6.2 (3) that

$$\|e^{\gamma_0 t} \mathbf{v}_2\|_{H_p^{1/2}(\mathbf{R}_+, W_q^1(\Omega))} \le M_1 \|e^{\gamma_0 t} \mathbf{v}_2\|_{W_{q,p}^{2,1}(\Omega \times \mathbf{R}_+)}$$

$$\leq M_1(\gamma_0+1) \|\mathbf{v}_2\|_{Y^{\gamma_0}_{q,p}},$$

the above inequality for I_8 yields

$$\|I_8\|_{H^{1,1/2}_{q,p}(\Omega \times \mathbf{R}_+)} \le 2M_1 M_2(\gamma_0 + 1) \|\mathbf{v}_2\|_{Y^{\gamma_0}_{q,p}} \|\mathbf{v}_2 - \mathbf{v}_1\|_{Y^{\gamma_0}_{q,p}}.$$

Analogously, we have

$$\|I_9\|_{H^{1,1/2}_{q,p}(\Omega \times \mathbf{R}_+)} \le 2M_1 M_2(\gamma_0 + 1) \|\mathbf{v}_1\|_{Y^{\gamma_0}_{q,p}} \|\mathbf{v}_2 - \mathbf{v}_1\|_{Y^{\gamma_0}_{q,p}},$$

and therefore

(6.10)
$$\|e^{\gamma_0 t} (G(\mathbf{v}_2) - G(\mathbf{v}_1))\|_{H^{1,1/2}_{q,p}(\Omega \times \mathbf{R}_+)} \\ \leq 2M_1 M_2(\gamma_0 + 1) \left(\|\mathbf{v}_2\|_{Y^{\gamma_0}_{q,p}} + \|\mathbf{v}_1\|_{Y^{\gamma_0}_{q,p}} \right) \|\mathbf{v}_2 - \mathbf{v}_1\|_{Y^{\gamma_0}_{q,p}}.$$

In addition, it is clear that $e^{\gamma_0 t}(G(\mathbf{v}_2) - G(\mathbf{v}_1))|_{t=0} = 0$ in $L_q(\Omega)$, which, combined with (6.2) and (6.10), furnishes

(6.11)
$$e^{\gamma_0 t}(G(\mathbf{v}_2) - G(\mathbf{v}_1)) \in {}_0H^{1,1/2}_{q,p}(\Omega \times \mathbf{R}_+).$$

Especially, setting $(\mathbf{v}_2, \mathbf{v}_1) = (\mathbf{v}, 0)$ in (6.11) and (6.10) yields, respectively,

(6.12)
$$e^{\gamma_0 t} G(\mathbf{v}) \in {}_0H^{1,1/2}_{q,p}(\Omega \times \mathbf{R}_+), \\ \|e^{\gamma_0 t} G(\mathbf{v})\|_{H^{1,1/2}_{q,p}(\Omega \times \mathbf{R}_+)} \le 2M_1 M_2(\gamma_0 + 1) \|\mathbf{v}\|^2_{Y^{\gamma_0}_{q,p}}$$

Next we show $e^{\gamma_0 t} G(\mathbf{v}) \in {}_0W_p^1(\mathbf{R}_+, \widehat{W}_{q,\Gamma}^{-1}(\Omega))$ and its estimate. For any $\varphi \in W^1_{q',\Gamma}(\Omega)$ with q' = q/(q-1), we have for k = 0, 1

$$(\partial_t^k (G(\mathbf{v}_2) - G(\mathbf{v}_1)), \varphi)_{\Omega} = (\partial_t^k \operatorname{div}(\mathbf{G}(\mathbf{v}_2) - \mathbf{G}(\mathbf{v}_1)), \varphi)_{\Omega}$$
$$= -(\partial_t^k (\mathbf{G}(\mathbf{v}_2) - \mathbf{G}(\mathbf{v}_1)), \nabla \varphi)_{\Omega},$$

where we have used $\mathbf{G}(\mathbf{v}_l) = 0$ on S for l = 1, 2. Thus $\partial_t^k(\mathbf{G}(\mathbf{v}_2) - \mathbf{G}(\mathbf{v}_1)) \in \mathcal{G}(\partial_t^k(G(\mathbf{v}_2) - G(\mathbf{v}_1)))$ as was discussed in Subsection 2.2, and also

$$\|\partial_t^k(G(\mathbf{v}_2) - G(\mathbf{v}_1))\|_{\widehat{W}_{q,\Gamma}^{-1}(\Omega)} \le \|\partial_t^k(\mathbf{G}(\mathbf{v}_2) - \mathbf{G}(\mathbf{v}_1))\|_{L_q(\Omega)}$$

This inequality, together with $e^{\gamma_0 t}(\mathbf{G}(\mathbf{v}_2) - \mathbf{G}(\mathbf{v}_1))|_{t=0} = 0$ in $L_q(\Omega)^N$, implies that

(6.13)
$$e^{\gamma_0 t} (G(\mathbf{v}_2) - G(\mathbf{v}_1))|_{t=0} = 0 \quad \text{in } \widehat{W}_{q,\Gamma}^{-1}(\Omega), \\ \|e^{\gamma_0 t} \partial_t^k (G(\mathbf{v}_2) - G(\mathbf{v}_1))\|_{L_p(\mathbf{R}_+, \widehat{W}_{q,\Gamma}^{-1}(\Omega))} \\ \leq \|e^{\gamma_0 t} \partial_t^k (\mathbf{G}(\mathbf{v}_2) - \mathbf{G}(\mathbf{v}_1))\|_{L_p(\mathbf{R}_+, L_q(\Omega))} \quad (k=0,1).$$

From now on, we estimate the right-hand side of the inequality in (6.13). To this end, we set

$$e^{\gamma_0 t} (\mathbf{G}(\mathbf{v}_2) - \mathbf{G}(\mathbf{v}_1)) = \left(\mathbf{U}_3 \left(\int_0^t \nabla \mathbf{v}_2 \, ds \right) - \mathbf{U}_3 \left(\int_0^t \nabla \mathbf{v}_1 \, ds \right) \right) e^{\gamma_0 t} \mathbf{v}_2 + \mathbf{U}_3 \left(\int_0^t \nabla \mathbf{v}_1 \, ds \right) e^{\gamma_0 t} (\mathbf{v}_2 - \mathbf{v}_1) =: I_{10} + I_{11}, e^{\gamma_0 t} \partial_t (\mathbf{G}(\mathbf{v}_2) - \mathbf{G}(\mathbf{v}_1)) = \left\{ \partial_t \left(\mathbf{U}_3 \left(\int_0^t \nabla \mathbf{v}_2 \, ds \right) - \mathbf{U}_3 \left(\int_0^t \nabla \mathbf{v}_1 \, ds \right) \right) \right\} e^{\gamma_0 t} \mathbf{v}_2 + \left(\mathbf{U}_3 \left(\int_0^t \nabla \mathbf{v}_2 \, ds \right) - \mathbf{U}_3 \left(\int_0^t \nabla \mathbf{v}_1 \, ds \right) \right) e^{\gamma_0 t} \partial_t \mathbf{v}_2$$

$$+ \left\{ \partial_t \mathbf{U}_3 \left(\int_0^t \nabla \mathbf{v}_1 \, ds \right) \right\} e^{\gamma_0 t} (\mathbf{v}_2 - \mathbf{v}_1) \\ + \mathbf{U}_3 \left(\int_0^t \nabla \mathbf{v}_1 \, ds \right) e^{\gamma_0 t} \partial_t (\mathbf{v}_2 - \mathbf{v}_1) =: I_{12} + I_{13} + I_{14} + I_{15}.$$

By Lemma 6.3, we see that for $j = 10, \ldots, 15$

$$\|I_{j}\|_{L_{p}(\mathbf{R}_{+},L_{q}(\Omega))} \leq M_{2}\left(\|\mathbf{v}_{2}\|_{Y_{q,p}^{\gamma_{0}}} + \|\mathbf{v}_{1}\|_{Y_{q,p}^{\gamma_{0}}}\right)\|\mathbf{v}_{2} - \mathbf{v}_{1}\|_{Y_{q,p}^{\gamma_{0}}}.$$

It thus holds that

$$\begin{split} &\sum_{k=0}^{1} \|e^{\gamma_{0}t}\partial_{t}^{k}(\mathbf{G}(\mathbf{v}_{2})-\mathbf{G}(\mathbf{v}_{1}))\|_{L_{p}(\mathbf{R}_{+},L_{q}(\Omega))} \\ &\leq 6M_{2}\left(\|\mathbf{v}_{2}\|_{Y_{q,p}^{\gamma_{0}}}+\|\mathbf{v}_{1}\|_{Y_{q,p}^{\gamma_{0}}}\right)\|\mathbf{v}_{2}-\mathbf{v}_{1}\|_{Y_{q,p}^{\gamma_{0}}}, \end{split}$$

which, combined with (6.13), furnishes

(6.14)
$$e^{\gamma_{0}t}(G(\mathbf{v}_{2}) - G(\mathbf{v}_{1})) \in {}_{0}W_{p}^{1}(\mathbf{R}_{+}, \widehat{W}_{q,\Gamma}^{-1}(\Omega)),$$
$$\|e^{\gamma_{0}t}(G(\mathbf{v}_{2}) - G(\mathbf{v}_{1}))\|_{W_{p}^{1}(\mathbf{R}_{+}, \widehat{W}_{q,\Gamma}^{-1}(\Omega))}$$
$$\leq 6M_{2}(\gamma_{0} + 1) \left(\|\mathbf{v}_{2}\|_{Y_{q,p}^{\gamma_{0}}} + \|\mathbf{v}_{1}\|_{Y_{q,p}^{\gamma_{0}}}\right) \|\mathbf{v}_{2} - \mathbf{v}_{1}\|_{Y_{q,p}^{\gamma_{0}}}.$$

Especially, setting $(\mathbf{v}_2, \mathbf{v}_1) = (\mathbf{v}, 0)$ in (6.14) yields

(6.15)
$$e^{\gamma_0 t} G(\mathbf{v}) \in {}_0 W_p^1(\mathbf{R}_+, \widehat{W}_{q,\Gamma}^{-1}(\Omega)), \\ \|e^{\gamma_0 t} G(\mathbf{v})\|_{W_p^1(\mathbf{R}_+, \widehat{W}_{q,\Gamma}^{-1}(\Omega))} \le 6M_2(\gamma_0 + 1) \|\mathbf{v}\|_{Y_{q,p}^{\gamma_0}}^2.$$

Etimates of $H(\mathbf{v})$. In the same manner as in the case $G(\mathbf{v})$, we have

(6.16)
$$e^{\gamma_{0}t}(\mathbf{H}(\mathbf{v}_{2})\mathbf{e}_{N} - \mathbf{H}(\mathbf{v}_{1})\mathbf{e}_{N}), e^{\gamma_{0}t}\mathbf{H}(\mathbf{v})\mathbf{e}_{N} \in {}_{0}H_{q,p}^{1,1/2}(\Omega \times \mathbf{R}_{+})^{N}, \\ \left\| e^{\gamma_{0}t}(\mathbf{H}(\mathbf{v}_{2})\mathbf{e}_{N} - \mathbf{H}(\mathbf{v}_{1})\mathbf{e}_{N}) \right\|_{H_{q,p}^{1,1/2}(\Omega \times \mathbf{R}_{+})} \\ \leq \left(8M_{1}M_{2}(\gamma_{0}+1) + 6M_{1}M_{2}^{2}(\gamma_{0}+1) \right) \\ \cdot \left(\left\| \mathbf{v}_{2} \right\|_{Y_{q,p}^{\gamma_{0}}} + \left\| \mathbf{v}_{2} \right\|_{Y_{q,p}^{\gamma_{0}}} \right) \left\| \mathbf{v}_{2} - \mathbf{v}_{1} \right\|_{Y_{q,p}^{\gamma_{0}}}, \\ \left\| e^{\gamma_{0}t}\mathbf{H}(\mathbf{v})\mathbf{e}_{N} \right\|_{H_{q,p}^{1,1/2}(\Omega \times \mathbf{R}_{+})} \\ \leq \left(8M_{1}M_{2}(\gamma_{0}+1) + 6M_{1}M_{2}^{2}(\gamma_{0}+1) \right) \left\| \mathbf{v} \right\|_{Y_{q,p}^{\gamma_{0}}}^{2}.$$

Let $M_4 \ge 1$ be a positive constant defined as

(6.17)
$$M_4 = 4M_2 + (p'\gamma_0)^{-1/p'} M_1 M_2 + 2(p'\gamma_0)^{-1/p'} M_1 M_3 + 2M_1 M_2(\gamma_0 + 1) + 6M_2(\gamma_0 + 1) + 8M_1 M_2(\gamma_0 + 1) + 6M_1 M_2^2(\gamma_0 + 1).$$

Summing up (6.8)-(6.12) and (6.14)-(6.16), we have obtained

(6.18)
$$e^{\gamma_0 t}(\mathbf{F}(\mathbf{v}_2) - \mathbf{F}(\mathbf{v}_1)), e^{\gamma_0 t} \mathbf{F}(\mathbf{v}) \in L_p(\mathbf{R}_+, L_q(\Omega))^N,$$
$$e^{\gamma_0 t}(G(\mathbf{v}_2) - G(\mathbf{v}_1)), e^{\gamma_0 t}G(\mathbf{v}) \in {}_0H^{1,1/2}_{q,p}(\Omega \times \mathbf{R}_+) \cap {}_0W^1_p(\mathbf{R}_+, \widehat{W}^{-1}_{q,\Gamma}(\Omega)),$$
$$e^{\gamma_0 t}(\mathbf{H}(\mathbf{v}_2)\mathbf{e}_N - \mathbf{H}(\mathbf{v}_1)\mathbf{e}_N), e^{\gamma_0 t}\mathbf{H}(\mathbf{v})\mathbf{e}_N \in {}_0H^{1,1/2}_{q,p}(\Omega \times \mathbf{R}_+)^N,$$

together with the estimates:

(6.19) $\left\| e^{\gamma_0 t} (\mathbf{F}(\mathbf{v}_2) - \mathbf{F}(\mathbf{v}_1)) \right\|_{L_p(\mathbf{R}_+, L_q(\Omega))}$

$$+ \left\| e^{\gamma_{0}t} (\mathbf{H}(\mathbf{v}_{2})\mathbf{e}_{N} - \mathbf{H}(\mathbf{v}_{1})\mathbf{e}_{N}) \right\|_{H^{1,1/2}_{q,p}(\Omega \times \mathbf{R}_{+})} + \left\| e^{\gamma_{0}t} (G(\mathbf{v}_{2}) - G(\mathbf{v}_{1})) \right\|_{H^{1,1/2}_{q,p}(\Omega \times \mathbf{R}_{+})} + \left\| e^{\gamma_{0}t} (G(\mathbf{v}_{2}) - G(\mathbf{v}_{1})) \right\|_{W^{1}_{p}(\mathbf{R}_{+},\widehat{W}^{-1}_{q,\Gamma}(\Omega))} \leq 4M_{4} \left(\left\| \mathbf{v}_{2} \right\|_{Y^{\gamma_{0}}_{q,p}} + \left\| \mathbf{v}_{1} \right\|_{Y^{\gamma_{0}}_{q,p}} \right) \left\| \mathbf{v}_{2} - \mathbf{v}_{1} \right\|_{Y^{\gamma_{0}}_{q,p}}.$$

$$(6.20) \quad \left\| e^{\gamma_{0}t} \mathbf{F}(\mathbf{v}) \right\|_{L_{p}(\mathbf{R}_{+},L_{q}(\Omega))} + \left\| e^{\gamma_{0}t} \mathbf{H}(\mathbf{v}) \mathbf{e}_{N} \right\|_{H^{1,1/2}_{q,p}(\Omega \times \mathbf{R}_{+})} + \left\| e^{\gamma_{0}t} G(\mathbf{v}) \right\|_{H^{1,1/2}_{q,p}(\Omega \times \mathbf{R}_{+})} + \left\| e^{\gamma_{0}t} G(\mathbf{v}) \right\|_{W^{1}_{p}(\mathbf{R}_{+},\widehat{W}^{-1}_{q,\Gamma}(\Omega)} \leq 4M_{4} \| \mathbf{v} \|^{2}_{Y^{\gamma_{0}}_{q,p}}.$$

We now construct a solution to (6.3) by using Thorem 2.4. For any $(\mathbf{v}, \mathbf{q}) \in X_{q,p}^{\gamma_0}(\delta_0)$ with $0 < \delta_0 < 1$, we have by (6.18) a unique solution $(\mathbf{u}, \mathbf{p}) \in W_{q,p}^{2,1}(\Omega \times \mathbf{R}_+)^N \times L_p(\mathbf{R}_+, W_q^1(\Omega))$ to (6.3) and have by (6.20)

$$\|(\mathbf{u},\mathfrak{p})\|_{X_{q,p}^{\gamma_0}} \le c_0(\|\mathbf{a}\|_{B_{q,p}^{2-2/p}(\Omega)} + 4M_4 \|\mathbf{v}\|_{Y_{q,p}^{\gamma_0}}^2) \le c_0(\varepsilon_0 + 4M_4\delta_0^2).$$

One here chooses δ_0 , ε_0 as follows:

(6.21)
$$4c_0 M_4 \delta_0^2 \le \frac{\delta_0}{4}, \quad c_0 \varepsilon_0 \le \frac{\delta_0}{2},$$

which enable us to define the operator:

$$\Phi: X_{q,p}^{\gamma_0}(\delta_0) \ni (\mathbf{v}, \mathfrak{q}) \mapsto \Phi(\mathbf{v}, \mathfrak{q}) = (\mathbf{u}, \mathfrak{p}) \in X_{q,p}^{\gamma_0}(\delta_0).$$

Let $(\mathbf{v}_i, \mathfrak{q}_i) \in X_{q,p}^{\gamma_0}(\delta_0)$ and $(\mathbf{u}_i, \mathfrak{p}_i) = \Phi(\mathbf{v}_i, \mathfrak{q}_i)$ for i = 1, 2. Setting $\bar{\mathbf{u}} = \mathbf{u}_2 - \mathbf{u}_1$ and $\bar{\mathfrak{p}} = \mathfrak{p}_2 - \mathfrak{p}_1$, we observe that

$$\begin{cases} \partial_t \bar{\mathbf{u}} - \operatorname{Div} \mathbf{T}(\bar{\mathbf{u}}, \bar{\mathbf{p}}) = \mathbf{F}(\mathbf{v}_2) - \mathbf{F}(\mathbf{v}_1) & \text{in } \Omega, t > 0, \\ \operatorname{div} \bar{\mathbf{u}} = G(\mathbf{v}_2) - G(\mathbf{v}_1) = \operatorname{div}(\mathbf{G}(\mathbf{v}_2) - \mathbf{G}(\mathbf{v}_1)) & \text{in } \Omega, t > 0, \\ \mathbf{T}(\bar{\mathbf{u}}, \bar{\mathbf{p}})\mathbf{e}_N = (\mathbf{H}(\mathbf{v}_2) - \mathbf{H}(\mathbf{v}_1))\mathbf{e}_N & \text{on } \Gamma, t > 0, \\ \bar{\mathbf{u}} = 0 & \text{on } S, t > 0, \\ \bar{\mathbf{u}}|_{t=0} = 0 & \text{in } \Omega. \end{cases}$$

Together with (6.18), (6.19), and (6.21), one has by Thorem 2.4

$$\|(\bar{\mathbf{u}},\bar{\mathfrak{p}})\|_{X^{\gamma_0}_{q,p}} \le 4c_0 M_4(\delta_0 + \delta_0) \|\mathbf{v}_2 - \mathbf{v}_1\|_{Y^{\gamma_0}_{q,p}} \le \frac{1}{2} \|(\mathbf{v}_2,\mathfrak{q}_2) - (\mathbf{v}_1,\mathfrak{q}_1)\|_{X^{\gamma_0}_{q,p}},$$

which implies that Φ is a contraction mapping on $X_{q,p}^{\gamma_0}(\delta_0)$. The contraction mapping theorem thus proves that there exists a unique fixed point $(\mathbf{u}_*, \mathbf{p}_*)$ of Φ in $X_{q,p}^{\gamma_0}$. Such a $(\mathbf{u}_*, \mathbf{p}_*)$ is a unique solution to (1.9)-(1.13) in $X_{q,p}^{\gamma_0}(\delta_0)$. This completes the proof of Theorem 2.2.

7. Original nonlinear problem

This section is concerned with the global solvability of the original nonlinear problem (1.1)-(1.7). We first introduce the definition of L_p - L_q solutions to (1.1)-(1.7). Next we show the global existence and uniqueness of such solutions, and also their exponential stability. Let M_1 , M_2 , M_3 , and M_4 be the same positive constants as in Section 6, and let c_0 be the positive constant given by Theorem 2.4.

7.1. Definition of L_p - L_q solutions. Following [18], we introduce the definition of L_p - L_q solutions for (1.1)-(1.7) in this subsection.

One first recalls the definition of L_p - L_q solutions for the equations (1.9)-(1.13).

Definition 7.1. We call a pair (\mathbf{u}, \mathbf{p}) an L_p - L_q solution global in time to (1.9)-(1.13) if $(\mathbf{u}, \mathbf{p}) \in W^{2,1}_{q,p}(\Omega \times \mathbf{R}_+)^N \times L_p(\mathbf{R}_+ W^1_q(\Omega))$ and if (\mathbf{u}, \mathbf{p}) satisfies (1.9)-(1.13) in the L_p - L_q sense for some $1 < p, q < \infty$ and $\mathbf{a} \in B^{2-2/p}_{q,p}(\Omega)^N$.

Remark 7.2. Due to [36], the maximal L_p - L_q regularity class means the function space of (\mathbf{u}, \mathbf{p}) in Definition 7.1.

Then we can define L_p - L_q solutions to (1.1)-(1.7) as follows:

Definition 7.3. We call a triplet $(\Theta, \mathbf{v}, \pi)$, where for $\Omega(t) = \Theta(\Omega, t)$

$$\mathbf{v}: \bigcup_{t \in (0,\infty)} \left(\Omega(t) \times \{t\} \right) \to \mathbf{R}^N, \quad \pi: \bigcup_{t \in (0,\infty)} \left(\Omega(t) \times \{t\} \right) \to \mathbf{R},$$

an L_p - L_q solution global in time to (1.1)-(1.7) if the following assertions hold true for some $1 < p, q < \infty$ and $\mathbf{a} \in B_{q,p}^{2-2/p}(\Omega)^N$:

- (1) $\Theta = \Theta(\xi, t)$ is a solution to (1.1) in the classical sense.
- (2) $\Theta(\cdot,t)$ is a C^1 -diffeomorphism from Ω onto $\Omega(t)$ for each t > 0.

(3) $(\mathbf{u}, \mathbf{p}) = (\mathbf{v} \circ \Theta, \pi \circ \Theta)$ is an $L_p - L_q$ solution global in time to (1.9)-(1.13).

7.2. Global solvability and exponential stability. We here prove

Theorem 7.4. Let p, q satisfy (2.1). Suppose that ε_0 is the same positive number as in Theorem 2.2 and that $\mathbf{a} \in D_{q,p}(\Omega)$ with $\|\mathbf{a}\|_{D_{q,p}(\Omega)} \leq \varepsilon_0$. Then there exists an L_p - L_q solution (Θ, \mathbf{v}, π) global in time to (1.1)-(1.7), which is unique. In addition, $\|\mathbf{v}(t)\|_{L_q(\Omega(t))} = O(e^{-\gamma_0 t})$ as $t \to \infty$, where γ_0 is the same positive constant as in Theorem 2.2.

Proof. By Theorem 2.2, we have an L_p - L_q solution (\mathbf{u}, \mathbf{p}) to (1.9)-(1.13). Let us define for t > 0

$$\Theta(\xi,t) = \xi + \int_0^t \mathbf{u}(\xi,s) \, ds \quad (\xi \in \Omega), \quad \Omega(t) = \Theta(\Omega,t),$$

where we note that $\mathbf{u} \in BUC([0,\infty), BUC^1(\overline{\Omega}))$ by Lemma 6.2 (2).

Step 1. In this step, we prove the existence of L_p - L_q solutions global in time to the equations (1.1)-(1.7).

Let $x_1, x_2 \in \Omega(t)$ with $x_1 = x_2$ for

$$x_i = \xi_i + \int_0^t \mathbf{u}(\xi_i, s) \, ds \quad (i = 1, 2),$$

where $\xi_1, \xi_2 \in \Omega$. Since it holds that by Lemma 6.2 (1), (6.17), and (6.21)

$$(7.1) \qquad \int_0^t \|\nabla \mathbf{u}(\cdot,s)\|_{BUC(\overline{\Omega})} \, ds \leq \left(\int_0^\infty e^{-p'\gamma_0 s} \, ds\right)^{1/p'} \|e^{\gamma_0 t} \nabla \mathbf{u}\|_{L_p(\mathbf{R}_+, BUC(\overline{\Omega}))}$$
$$\leq (p'\gamma_0)^{-1/p'} M_1 \|e^{\gamma_0 t} \nabla \mathbf{u}\|_{L_p(\mathbf{R}_+, W_q^1(\Omega))}$$
$$\leq (p'\gamma_0)^{-1/p'} M_1 \delta_0 \leq (p'\gamma_0)^{-1/p'} M_1 M_2 \delta_0 \leq 4c_0 M_4 \delta_0 \leq \frac{1}{4}$$

for any t > 0, one observes that

$$0 = |x_1 - x_2| \ge |\xi_1 - \xi_2| - \int_0^t |\mathbf{u}(\xi_1, s) - \mathbf{u}(\xi_2, s)| \, ds$$
$$\ge |\xi_1 - \xi_2| - |\xi_1 - \xi_2| \int_0^t \|\nabla \mathbf{u}(\cdot, s)\|_{BUC(\overline{\Omega})} \, ds$$
$$\ge \frac{3}{4} |\xi_1 - \xi_2|.$$

This inequality implies $\xi_1 = \xi_2$, and thus $\Theta(\cdot, t)$ is bijective from Ω onto $\Omega(t)$ for each t > 0. We denote inverse mapping of $\Theta(\cdot, t)$ by $\Theta^{-1}(\cdot, t)$ in what follows. Here $\Theta(\cdot, t) : \Omega \to \Omega(t)$ is a C^1 function, so that we see by the inverse function theorem that $\Theta^{-1}(\cdot, t) : \Omega(t) \to \Omega$ is also a C^1 function. Hence, $\Theta(\cdot, t)$ satisfies Definition 7.3 (2), while it is clear that

(7.2)
$$\mathbf{v}(x,t) = \mathbf{u}(\Theta^{-1}(x,t),t), \quad \pi(x,t) = \mathfrak{p}(\Theta^{-1}(x,t),t) \quad (x \in \Omega(t), t > 0)$$

satisfies Definition 7.3 (3) and Θ is a solution to (1.1).

Step 2. Let \mathbf{J}_{Θ} be the Jacobian matrix of Θ . Since $\mathbf{J}_{\Theta} = \mathbf{I} + \int_{0}^{t} \nabla \mathbf{u}(\xi, s) ds$, one has $|\mathbf{J}_{\Theta}| \leq c_{1}$ by (7.1) for some positive constant c_{1} . It then holds by (7.2) that

$$\begin{aligned} \|\mathbf{v}(t)\|_{L_q(\Omega(t))} &= \left(\int_{\Omega} |\mathbf{v}(\Theta(\xi, t), t)|^q |\mathbf{J}_{\Theta}| \, d\xi\right)^{1/q} \\ &= \left(\int_{\Omega} |\mathbf{u}(\xi, t)|^q |\mathbf{J}_{\Theta}| \, d\xi\right)^{1/q} \leq (c_1)^{1/q} \|\mathbf{u}(t)\|_{L_q(\Omega)} \\ &\leq (c_1)^{1/q} e^{-\gamma_0 t} \|e^{\gamma_0 t} \mathbf{u}\|_{BUC([0,\infty), L_q(\Omega))} \\ &\leq (c_1)^{1/q} e^{-\gamma_0 t} \|e^{\gamma_0 t} \mathbf{u}\|_{BUC([0,\infty), B^{2-2/p}_{q, 0}(\Omega))}.\end{aligned}$$

This inequality, together with (2.2) and (5.14), furnishes the exponential stability of the solution **v**.

Step 3. We prove the uniqueness of solutions in this step, Let i = 1, 2 and $(\Theta_i, \mathbf{v}_i, \pi_i)$ be solutions to (1.1)-(1.7) with $\mathbf{a} \in D_{q,p}(\Omega)$ satisfying $\|\mathbf{a}\|_{D_{q,p}(\Omega)} \leq \varepsilon_0$. Then $(\mathbf{u}_i, \mathbf{p}_i) = (\mathbf{v}_i \circ \Theta_i, \pi_i \circ \Theta_i)$ are L_p - L_q solutions global in time to (1.9)-(1.13). By Theorem 2.2, we see that $\mathbf{u}_1 = \mathbf{u}_2$ and $\mathbf{p}_1 = \mathbf{p}_2$. One integrates (1.1) with respect to time t in order to obtain

$$\Theta_1(\xi, t) = \xi + \int_0^t \mathbf{v}_1(\Theta_1(\xi, s), s) \, ds = \xi + \int_0^t \mathbf{u}_1(\xi, s) \, ds$$

= $\xi + \int_0^t \mathbf{u}_2(\xi, s) \, ds = \xi + \int_0^t \mathbf{v}_2(\Theta_2(\xi, s), s) \, ds = \Theta_2(\xi, t).$

Furthermore, for t > 0 and $x \in \Theta_1(\Omega, t) = \Theta_2(\Omega, t)$, we observe that

$$\mathbf{v}_1(x,t) = \mathbf{v}_1(\Theta_1(\xi,t),t) = \mathbf{u}_1(\xi,t) = \mathbf{u}_2(\xi,t) = \mathbf{v}_2(\Theta_2(\xi,t),t) = \mathbf{v}_2(x,t)$$

and that $\pi_1(x,t) = \pi_2(x,t)$. This completes the proof of Theorem 7.4.

Remark 7.5. One can prove similarly further properties of the solution $(\Theta, \mathbf{v}, \pi)$ as follows:

(1) Let $\Gamma(t) = \Theta(\Gamma, t)$ for t > 0. Then $\Theta(\cdot, t)$ is a C^1 -diffeomorphism from Γ onto $\Gamma(t)$ for each t > 0. In addition, it holds that $\Theta(\overline{\Omega}, t) = \overline{\Omega(t)}$.

- (2) Let $\widetilde{\Theta}(\xi, t) = (\Theta(\xi, t), t)$ for $(\xi, t) \in \Omega \times \mathbf{R}_+$. Then $\widetilde{\Theta}$ is a C^1 -diffeomorphism from $\Omega \times \mathbf{R}_+$ onto $\bigcup_{t \in (0,\infty)} (\Omega(t) \times \{t\})$.
- (3) $\|\nabla \mathbf{v}(t)\|_{L_q(\Omega(t))} = O(e^{-\gamma_0 t})$ as $t \to \infty$.

Acknowledgments. This research was partly supported by JSPS Grant-in-aid for Young Scientists (B) #17K14224, JSPS Japanese-German Graduate Externship at Waseda University, and Waseda University Grant for Special Research Projects (Project number: 2017K-176).

Α.

Following [38, Appendix], we derive the equations (1.9)-(1.11) in this appendix. To this end, we assume that the equations (1.1)-(1.7) have a sufficiently regular solution $(\Theta, \mathbf{v}, \pi)$ in the following argumentation. Let $x = \Theta(\xi, t)$ with (1.8), and recall that $\mathbf{u}(\xi, t) = \mathbf{v}(\Theta(\xi, t), t) = \mathbf{v}(x, t)$ and $\mathfrak{p}(\xi, t) = \pi(\Theta(\xi, t), t) = \pi(x, t)$. If we write $\mathbf{M} = (M_{ij})$, then \mathbf{M} is an $N \times N$ matrix whose (i, j)-component is M_{ij} . In addition, δ_{ij} denotes Kronecker's delta defined by the formula: $\delta_{ij} = 1$ when i = jand $\delta_{ij} = 0$ when $i \neq j$.

Case (1.9). It is clear that

(A.1)
$$\frac{\partial}{\partial t}\mathbf{u}(\xi,t) = \frac{\partial}{\partial t}\mathbf{v}(x,t) + \sum_{j=1}^{N} \frac{\partial x_j}{\partial t} \frac{\partial}{\partial x_j} \mathbf{v}(x,t) = \frac{\partial}{\partial t}\mathbf{v} + (\mathbf{v}\cdot\nabla)\mathbf{v}.$$

We here set

$$\mathbf{A} = \left(\frac{\partial x_i}{\partial \xi_j}\right) = \mathbf{I} + \mathbf{B}, \quad \mathbf{B} = (B_{ij}), \quad B_{ij} = \int_0^t \frac{\partial u_i}{\partial \xi_j}(\xi, s) \, ds.$$

Let \mathcal{A} be the cofactor matrix of \mathbf{A} , i.e. $\mathbf{A}^{-1} = (\det \mathbf{A})^{-1} \mathcal{A}$. Then $\mathcal{A} = \mathbf{A}^{-1}$ because det $\mathbf{A} = 1$ (cf. e.g. [38, page 271]), and also one observes by $\delta_{ij} = \partial \xi_i / \partial \xi_j = \sum_{k=1}^{N} \frac{\partial \xi_i}{\partial x_k} \frac{\partial x_k}{\partial \xi_j}$ that

$$\mathcal{A} = \mathbf{A}^{-1} = \begin{pmatrix} \frac{\partial \xi_1}{\partial x_1} & \cdots & \frac{\partial \xi_1}{\partial x_N} \\ \vdots & \ddots & \vdots \\ \frac{\partial \xi_N}{\partial x_1} & \cdots & \frac{\partial \xi_N}{\partial x_N} \end{pmatrix}.$$

In addition,

$$\mathcal{A} = \mathbf{I} + \mathcal{B} \quad \text{for some } N \times N \text{ matrix } \mathcal{B} = (\mathcal{B}_{ij}) = \mathcal{B}\left(\int_0^t \nabla \mathbf{u}(\xi, s) \, ds\right),$$

where $\mathcal{B} : \mathbf{R}^{N \times N} \to \mathbf{R}^{N \times N}$ with $\mathcal{B}(\mathbf{O}) = \mathbf{O}$ and the (i, j)-component $\mathcal{B}_{ij}(\mathbf{X})$ of $\mathcal{B}(\mathbf{X}), \mathbf{X} = (X_{kl})$, are polynomials with respect to X_{kl} for $k, l = 1, \ldots, N$.

Now it holds, by direct calculations and by [14, page 771], that

(A.2)
$$\nabla_x = \mathcal{A}^{\mathsf{T}} \nabla_{\xi}, \quad \operatorname{div}_x = (\mathcal{A}^{\mathsf{T}} : \nabla_{\xi} \cdot) = \operatorname{div}_{\xi} (\mathcal{A} \cdot),$$
$$\nabla_x \operatorname{div}_x = \mathcal{A}^{\mathsf{T}} \nabla_{\xi} \operatorname{div}_{\xi} + \mathcal{A}^{\mathsf{T}} \nabla_{\xi} (\mathcal{B}^{\mathsf{T}} : \nabla_{\xi} \cdot),$$

where the subscripts x and ξ denote their variables, and also

(A.3)
$$\Delta_x = \operatorname{div}_x \nabla_x = (\operatorname{div}_{\xi} + \mathcal{B}^{\mathsf{T}} : \nabla_{\xi} \cdot)(\mathbf{I} + \mathcal{B}^{\mathsf{T}})\nabla_{\xi}$$
$$= \Delta_{\xi} + \operatorname{div}_{\xi} \mathcal{B}^{\mathsf{T}} \nabla_{\xi} + \mathcal{B}^{\mathsf{T}} : \nabla_{\xi} \nabla_{\xi} + \mathcal{B}^{\mathsf{T}} : \nabla_{\xi} \mathcal{B}^{\mathsf{T}} \nabla_{\xi}$$

$$= \Delta_{\xi} + \sum_{i,j,k=1}^{N} \mathcal{B}_{ji} \left(2\delta_{ik} + \mathcal{B}_{ki} \right) \frac{\partial^2}{\partial \xi_j \partial \xi_k} + \sum_{i,j,k=1}^{N} \left(\delta_{ij} + \mathcal{B}_{ji} \right) \left(\frac{\partial \mathcal{B}_{ki}}{\partial \xi_j} \right) \frac{\partial}{\partial \xi_k}$$

Then we can write

(A.4)
$$\sum_{i,j,k=1}^{N} \mathcal{B}_{ji} \left(2\delta_{ik} + \mathcal{B}_{ki} \right) \frac{\partial^2}{\partial \xi_j \partial \xi_k} = \sum_{i,j,k=1}^{N} \mathcal{V}_{ijk} \left(\int_0^t \nabla \mathbf{u}(\xi, s) \, ds \right) \frac{\partial^2}{\partial \xi_j \partial \xi_k},$$
$$\sum_{i,j,k=1}^{N} \left(\delta_{ij} + \mathcal{B}_{ji} \right) \left(\frac{\partial \mathcal{B}_{ki}}{\partial \xi_j} \right) \frac{\partial}{\partial \xi_k}$$
$$= \sum_{i,j,k=1}^{N} \left[\mathcal{W}_{ijk} \left(\int_0^t \nabla \mathbf{u}(\xi, s) \, ds \right) \int_0^t \nabla^2 \mathbf{u}(\xi, s) \, ds \right] \frac{\partial}{\partial \xi_k}$$

by $\mathcal{V}_{ijk}(\cdot) : \mathbf{R} \to \mathbf{R}$ with $\mathcal{V}_{ijk}(\mathbf{O}) = 0$ and by $\mathcal{W}_{ijk}(\cdot) : \mathbf{R}^{N^3} \to \mathbf{R}$. Note that both $\mathcal{V}_{ijk}(\mathbf{X})$ and $\mathcal{W}_{ijk}(\mathbf{X})$, $\mathbf{X} = (X_{lm})$, are polynomials with respect to X_{lm} for $l, m = 1, \ldots, N$. Since Div $\mathbf{T}(\mathbf{v}, \pi) = \mu(\Delta \mathbf{v} + \nabla \operatorname{div} \mathbf{v}) - \nabla \pi$, we insert (A.1) and (A.2)-(A.4) into (1.3) in order to obtain

$$\partial_{t}\mathbf{u} - \mu \left(\Delta \mathbf{u} + \mathcal{A}^{\mathsf{T}}\nabla \operatorname{div}\mathbf{u} + \mathcal{A}^{\mathsf{T}}\nabla(\mathcal{B}^{\mathsf{T}}:\nabla\mathbf{u})\right) + \mathcal{A}^{\mathsf{T}}\nabla\mathfrak{p}$$
$$= \mu \left(\sum_{i,j,k=1}^{N} \mathcal{V}_{ijk} \left(\int_{0}^{t} \nabla \mathbf{u}(\xi,s) \, ds\right) \frac{\partial^{2}}{\partial \xi_{j} \partial \xi_{k}} \mathbf{u} + \sum_{i,j,k=1}^{N} \left[\mathcal{W}_{ijk} \left(\int_{0}^{t} \nabla \mathbf{u}(\xi,s) \, ds\right) \int_{0}^{t} \nabla^{2} \mathbf{u}(\xi,s) \, ds\right] \frac{\partial}{\partial \xi_{k}} \mathbf{u}\right)$$

Let $\mathcal{A}^{-\mathsf{T}} = (\mathcal{A}^{\mathsf{T}})^{-1} = (\mathcal{A}^{-1})^{\mathsf{T}}$, and multiply the last equation by $\mathcal{A}^{-\mathsf{T}} = (\mathbf{I} + \mathbf{B})^{\mathsf{T}}$ from the left-hand side. Thus,

.

$$\begin{aligned} \partial_t \mathbf{u} &- \mu (\Delta \mathbf{u} + \nabla \operatorname{div} \mathbf{u}) + \nabla \mathbf{p} = -\mathbf{B}^{\mathsf{T}} \partial_t \mathbf{u} + \mu \mathbf{B}^{\mathsf{T}} \Delta \mathbf{u} + \mu \nabla (\mathcal{B}^{\mathsf{T}} : \nabla \mathbf{u}) \\ &+ \mu \left(\mathbf{I} + \mathbf{B}^{\mathsf{T}} \right) \sum_{i,j,k=1}^N \mathcal{V}_{ijk} \left(\int_0^t \nabla \mathbf{u}(\xi, s) \, ds \right) \frac{\partial^2}{\partial \xi_j \partial \xi_k} \mathbf{u} \\ &+ \mu \left(\mathbf{I} + \mathbf{B}^{\mathsf{T}} \right) \sum_{i,j,k=1}^N \left[\mathcal{W}_{ijk} \left(\int_0^t \nabla \mathbf{u}(\xi, s) \, ds \right) \int_0^t \nabla^2 \mathbf{u}(\xi, s) \, ds \right] \frac{\partial}{\partial \xi_k} \mathbf{u}, \end{aligned}$$

which completes Case (1.9).

Case (1.10). The equation (1.10) follows from (1.4) and the second relation of (A.2) immediately. This completes Case (1.10).

Case (1.11). Let $F(\xi) = \xi_N - d$, and then $\Gamma = \{\xi \in \mathbf{R}^N \mid F(\xi) = 0\}$. We can regard $\xi \in \Gamma$ as $\xi = \xi(x, t) = \Theta^{-1}(x, t)$ for $x \in \Gamma(t)$, where $\Theta^{-1}(\cdot, t)$ is the inverse mapping of $\Theta(\cdot, t)$. Let $G(x, t) = F(\xi(x, t))$. Since $\Gamma(t)$ is defined by G(x, t) = 0, we see that the unit outward normal \mathbf{n} to $\Gamma(t)$ is given by

(A.5)
$$\mathbf{n} = \frac{\nabla G(x,t)}{|\nabla G(x,t)|}.$$

Now it holds that

$$\nabla G(x,t) = \left(\sum_{j=1}^{N} \frac{\partial \xi_j}{\partial x_1} \frac{\partial F}{\partial \xi_j}, \dots, \sum_{j=1}^{N} \frac{\partial \xi_j}{\partial x_N} \frac{\partial F}{\partial \xi_j}\right)^{\mathsf{T}} = \mathcal{A}^{\mathsf{T}} \mathbf{e}_N,$$

and thus we have by (A.5) the relation:

(A.6)
$$\mathbf{n} = \frac{\mathcal{A}^{\mathsf{T}} \mathbf{e}_N}{|\mathcal{A}^{\mathsf{T}} \mathbf{e}_N|}.$$

On the other hand, we see by (A.2) that

$$\mathbf{\Gamma}(\mathbf{v}, \pi) = -\mathbf{\mathfrak{p}}\mathbf{I} + \mu \left(\mathcal{A}^{\mathsf{T}} \nabla \mathbf{u} + (\nabla \mathbf{u})^{\mathsf{T}} \mathcal{A} \right),$$

which, combined with (A.6) and inserted into (1.11), furnishes

$$-\mathfrak{p}\frac{\mathcal{A}^{T}\mathbf{e}_{N}}{|\mathcal{A}^{T}\mathbf{e}_{N}|} + \mu\left(\mathcal{A}^{T}\nabla\mathbf{u} + (\nabla\mathbf{u})^{\mathsf{T}}\mathcal{A}\right)\frac{\mathcal{A}^{\mathsf{T}}\mathbf{e}_{N}}{|\mathcal{A}^{\mathsf{T}}\mathbf{e}_{N}|} = 0.$$

The last equation is multiplied by $|\mathcal{A}^{\mathsf{T}}\mathbf{e}_N|\mathcal{A}^{\mathsf{T}}$ from the left-hand side to give

$$-\mathbf{p}\mathbf{e}_N + \mu \left(\nabla \mathbf{u} + \mathcal{A}^{-\mathsf{T}} (\nabla \mathbf{u})^T \mathcal{A}\right) \mathcal{A}^{\mathsf{T}} \mathbf{e}_N = 0,$$

which, combined with $\mathcal{A}^{-\mathsf{T}} = \mathbf{I} + \mathbf{B}^T$ and $\mathcal{A} = \mathbf{I} + \mathcal{B}$, implies

$$\mathbf{T}(\mathbf{u}, \mathbf{p})\mathbf{e}_N = -\mu \left(\mathbf{D}(\mathbf{u}) \mathbf{\mathcal{B}}^{\mathsf{T}} + \left\{ (\nabla \mathbf{u})^{\mathsf{T}} \mathbf{\mathcal{B}} + \mathbf{B}^{\mathsf{T}} (\nabla \mathbf{u})^{\mathsf{T}} (\mathbf{I} + \mathbf{\mathcal{B}}) \right\} (\mathbf{I} + \mathbf{\mathcal{B}}^{\mathsf{T}}) \right) \mathbf{e}_N.$$

This completes Case (1.11).

References

- [1] H. Abels. The initial-value problem for the Navier-Stokes equations with a free surface in L_q -Sobolev spaces. Adv. Differential Equations, 10(1):45–64, 2005.
- [2] H. Abels. Generalized Stokes resolvent equations in an infinite layer with mixed boundary conditions. Math. Nachr., 279(4):351–367, 2006.
- [3] R.A. Adams and J.J.F. Fournier. Sobolev spaces, volume 140 of Pure and Applied Mathematics. Elsevier/Academic Press, Amsterdam, 2nd edition, 2003.
- [4] G. Allain. Small-time existence for the Navier-Stokes equations with a free surface. Appl. Math. Optim., 16(1):37–50, 1987.
- [5] H. Amann. Linear and quasilinear parabolic problems. Vol. I. Abstract linear theory, volume 89 of Monographs in Mathematics. Birkhäuser Boston, Inc., Boston, MA, 1995.
- [6] J.T. Beale. The initial value problem for the Navier-Stokes equations with a free surface. Comm. Pure Appl. Math., 34(3):359–392, 1981.
- [7] J.T. Beale. Large-time regularity of viscous surface waves. Arch. Rational Mech. Anal., 84(4):307–352, 1983/84.
- [8] J.T. Beale and T. Nishida. Large-time behavior of viscous surface waves. In *Recent Topics in Nonlinear PDE II*, volume 128 of *North-Holland Math. Stud.*, pages 1–14. North-Holland, Amsterdam, 1985.
- [9] I.V. Denisova. Problem of the motion of two viscous incompressible fluids separated by a closed free interface. Acta Appl. Math., 37(1-2):31–40, 1994.
- [10] I.V. Denisova. Problem of the motion of two compressible fluids separated by a closed free surface. Zap. Nauchn. Sem. S.-Peterburg. Otdel. Mat. Inst. Steklov. (POMI), 243:61–86; English transl: J. Math. Sci. (N. Y.) 99 (2000), no. 1, 837–853, 1997.
- [11] I.V. Denisova. Evolution of compressible and incompressible fluids separated by a closed interface. *Interfaces Free Bound.*, 2(3):283–312, 2000.
- [12] I.V. Denisova. Evolution of a closed interface between two liquids of different types. In European Congress of Mathematics, Vol. II (Barcelona, 2000), volume 202 of Progr. Math., pages 263–272. Birkhäuser, Basel, 2001.
- [13] I.V. Denisova. Solvability in weighted Hölder spaces for a problem governing the evolution of two compressible fluids. Zap. Nauchn. Sem. S.-Peterburg. Otdel. Mat. Inst. Steklov. (POMI), 295:57–89; English transl.: J. Math. Sci. (N. Y.) **127** (2005), no. 2, 1849–1868, 2003.

HIROKAZU SAITO

- [14] I.V. Denisova and V.A. Solonnikov. Classical solvability of the problem of the motion of two viscous incompressible fluids. *Algebra i Analiz*, 7(5):101–142; English transl.: St. Petersburg Math. J. 7 (1996), no. 5, 755–786, 1995.
- [15] I.V. Denisova and V.A. Solonnikov. Classical solvability of the problem of the motion of an isolated mass of compressible fluid. *Algebra i Analiz*, 14(1):71–98; English transl.: St. Petersburg Math. J. 14 (2002), no. 1, 53–74, 2002.
- [16] R. Denk and M. Kaip. General Parabolic Mixed Order Systems in L_p and Applications. Oper. Theory Adv. Appl. 239. Birkhäuser/Springer, Cham, 2013.
- [17] Y. Enomoto, L. von Below, and Y. Shibata. On some free boundary problem for a compressible barotropic viscous fluid flow. Ann. Univ. Ferrara Sez. VII Sci. Mat., 60(1):55–89, 2014.
- [18] J. Escher, J. Prüss, and G. Simonett. Analytic solutions for a Stefan problem with Gibbs-Thomson correction. J. Reine Angew. Math., 563:1–52, 2003.
- [19] E. Hanzawa. Classical solutions of the stefan problem. Tôhoku Math. J. (2), 33(3):297–335, 1981.
- [20] Y. Hataya. A remark on Beale-Nishida's paper. Bull. Inst. Math. Acad. Sin. (N. S.), 6(3):293– 303, 2011.
- [21] Y. Hataya and S. Kawashima. Decaying solution of the Navier-Stokes flow of infinite volume without surface tension. *Nonlinear Anal.*, 71(12):e2535–e2539, 2009.
- [22] M. Köhne, J. Prüss, and M. Wilke. Qualitative behaviour of solutions for the two-phase navier-stokes equations with surface tension. *Math. Ann.*, 2(356):737–792, 2013.
- [23] N. Lindemulder, M. Meyries, and M. Veraar. Complex interpolation with Dirichlet boundary conditions on the half line. 2017. preprint (arXiv:1705.11054).
- [24] A. Lunardi. Analytic semigroups and optimal regularity in parabolic problems, volume 16 of Progress in Nonlinear Differential Equations and their Applications. Birkhäuser Verlag, Basel, 1995.
- [25] S. Maryani and H. Saito. On the *R*-boundedness of solution operator families for two-phase Stokes resolvent equations. *Differential Integral Equations*, 30(1-2):1–52, 2017.
- [26] M. Meyries and R. Schnaubelt. Interpolation, embeddings and traces of anisotropic fractional Sobolev spaces with temporal weights. J. Funct. Anal., 262(3):1200–1229, 2012.
- [27] S.G. Mikhlin. Multidimensional Singular Integrals and Integral Equations. Pure and Applied Mathematics Monograph. Pergamon Press, Oxford-New York-Paris, 1965.
- [28] I.S. Mogilevskii and V.A. Solonnikov. On the solvability of an evolution free boundary problem for the Navier-Stokes equations in Hölder spaces of functions. In *Mathematical problems relating to the Navier-Stokes equation*, volume 11 of *Ser. Adv. Math. Appl. Sci.*, pages 105– 181. World Sci. Publ., River Edge, NJ, 1992.
- [29] P.B. Mucha and W.M. Zajączkowski. On local existence of solutions of the free boundary problem for an incompressible viscous self-gravitating fluid motion. *Appl. Math. (Warsaw)*, 27(3):319–333, 2000.
- [30] H. Saito. On the *R*-boundedness of solution operator families of the generalized stokes resolvent problem in an infinite layer. *Math. Methods Appl. Sci.*, 38(9):1888–1925, 2015.
- [31] H. Saito and Y. Shibata. On decay properties of solutions to the Stokes equations with surface tension and gravity in the half space. J. Math. Soc. Japan, 68(4):1559–1614, 2016.
- [32] P. Secchi. On the motion of gaseous stars in the presence of radiation. Comm. Partial Differential Equations, 15(2):185–204, 1990.
- [33] P. Secchi. On the uniqueness of motion of viscous gaseous stars. Math. Methods Appl. Sci., 13(5):391–404, 1990.
- [34] P. Secchi. On the evolution equations of viscous gaseous stars. Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4), 18(2):295–318, 1991.
- [35] P. Secchi and A. Valli. A free boundary problem for compressible viscous fluids. J. Reine Angew. Math., 341:1–31, 1983.
- [36] Y. Shibata. On some free boundary problem of the Navier-Stokes equations in the maximal L_p-L_q regularity class. J. Differential Equations, 258(12):4127–4155, 2015.
- [37] Y. Shibata. On the local wellposendness of free boundary problem for the Navier-Stokes equations in an exterior domain. 2017. preprint.
- [38] Y. Shibata and S. Shimizu. On a free boundary value problem for the Navier-Stokes equations. Differential Integral Equations, 20(3):241–276, 2007.
- [39] Y. Shibata and S. Shimizu. On the L_p-L_q maximal regularity of the Neumann problem for the Stokes equations in a bounded domain. J. Reine Angew. Math., 615:157–209, 2008.

- [40] Y. Shibata and S. Shimizu. On the maximal L_p-L_q regularity of the Stokes problem with first order boundary condition; model problems. J. Math. Soc. Japan, 64(2):561–626, 2012.
- [41] V. A. Solonnikov. l_p-theory of the problem of motion of two incompressible capillary fluids in a container. J. Math. Sci. (N.Y.), 198(6):761–827, 2014.
- [42] V.A. Solonnikov. Lectures on Evolution Free Boundary Problems: Classical Solutions. In Mathematical Aspects of Evolving Interfaces, volume 1812 of Lecture Notes in Math. Springer, Berlin, 2003.
- [43] V.A. Solonnikov. L_q-estimates for a solution to the problem about the evolution of an isolated amount of a fluid. J. Math. Sci. (N. Y.), 117(3):4237–4259, 2003.
- [44] V.A. Solonnikov and A. Tani. A problem with a free boundary for navier-stokes equations for a compressible fluid in the presence of surface tension. Zap. Nauchn. Sem. Leningrad. Otdel. Mat. Inst. Steklov. (LOMI), 182:142–148, 1990. English transl.: J. Soviet Math., 62(3):2814-2818, 1992.
- [45] V.A. Solonnikov and A. Tani. Free boundary problem for a viscous compressible flow with a surface tension. In *Constantin Carathéodory: An international tribute, Vol. II*, pages 1270– 1303. World Sci. Publ., Teaneck, NJ, 1991.
- [46] G. Ströhmer and W.M. Zajączkowski. Local existence of solutions of the free boundary problem for the equations of compressible barotropic viscous self-gravitating fluids. Appl. Math. (Warsaw), 26(1):1–31, 1999.
- [47] N. Tanaka and A. Tani. Surface waves for a compressible viscous fluid. J. Math. Fluid Mech., 5(4):303–363, 2003.
- [48] A. Tani. On the free boundary value problem for compressible viscous fluid motion. J. Math. Kyoto Univ., 21(4):839–859, 1981.
- [49] A. Tani. Two-phase free boundary problem for compressible viscous fluid motion. J. Math. Kyoto Univ., 24(2):243–267, 1984.
- [50] A. Tani. Small-time existence for the three-dimensional Navier-Stokes equations for an incompressible fluid with a free surface. Arch. Rational Mech. Anal., 133(4):299–331, 1996.
- [51] H. Triebel. Interpolation theory, function spaces, differential operators, volume 18 of North-Holland Mathematical Library. North-Holland Publishing Co., Amsterdam-New York, 1978.
- [52] E. Zadrzyńska and W.M. Zajączkowski. On local motion of a general compressible viscous heat conducting fluid bounded by a free surface. Ann. Polon. Math., 59(2):133–170, 1994.
- [53] E. Zadrzyńska and W.M. Zajączkowski. Local existence of solutions of a free boundary problem for equations of compressible viscous heat-conducting fluids. Appl. Math. (Warsaw), 25(2):179–220, 1998.
- [54] E. Zadrzyńska and W.M. Zajączkowski. Local existence of solutions of a free boundary problem for equations of compressible viscous heat-conducting capillary fluids. J. Appl. Anal., 6(2):227–250, 2000.
- [55] W.M. Zajączkowski. On local motion of a compressible barotropic viscous fluid bounded by a free surface. In *Partial differential equations, Part 2 (Warsaw, 1990)*, volume 27, Part 2 of *Banach Center Publ.* Polish Acad. Sci., Warsaw, 1992.
- [56] W.M. Zajączkowski. On nonstationary motion of a compressible barotropic viscous fluid bounded by a free surface. Dissertationes Math. (Rozprawy Mat.), 324:101 pp., 1993.
- [57] W.M. Zajączkowski. Existence of local solutions for free boundary problems for viscous compressible barotropic fluids. Ann. Polon. Math., 60(3):255–287, 1995.

DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE AND ENGINEERING, WASEDA UNIVERSITY, OKUBO 3-4-1, SHINJUKU-KU, TOKYO 169-8555, JAPAN

E-mail address: hsaito@aoni.waseda.jp