On Phragmén-Lindelöf principle for Non-divergence Type Elliptic Equation and Mixed Boundary conditions

Akif Ibraguimov^{*} and Alexander I. Nazarov[†]

Abstract

Paper dedicated to qualitative study of the solution of the Zaremba type problem in Lipschitz domain with respect to the elliptic equation in non-divergent form. Main result is Landis type Growth Lemma in spherical layer for Mixed Boundary Value Problem in the class of "admissible domain". Based on the Growth Lemma Phragmén-Lindelöf theorem is proved at junction point of Dirichlet boundary and boundary over which derivative in non-tangential direction is defined.

1 Introduction

We consider non-divergence elliptic operator

$$\mathcal{L}u := -\sum_{i,j=1}^{n} a_{ij}(x) D_i D_j u \quad \text{in} \quad \Omega.$$
(1.1)

Such operators arise in theory of stochastic processes and various applications.

In (1.1) Ω is a domain in \mathbb{R}^n , $n \geq 3$, and D_i stands for the differentiation with respect to x_i . We suppose that the boundary $\partial\Omega$ is split $\partial\Omega = \Gamma_1 \cup \{\zeta\} \cup \Gamma_2$. Here Γ_1 is support of the Dirichlet condition, and Γ_2 is support of the oblique derivative condition:

$$u(x) = \Phi(x)$$
 on Γ_1 ; $\frac{\partial u}{\partial \ell}(x) := \lim_{\delta \to +0} \frac{u(x) - u(x - \delta \ell)}{\delta} = \Psi(x)$ on Γ_2 ;

where $\ell = \ell(x)$ is a measurable, and uniformly non-tangential outward vector field on Γ_2 . Without loss of generality we can suppose $|\ell| \equiv 1$. We call Γ_1 Dirichlet boundary, and Γ_2 Neumann boundary.

^{*}Email: akif.ibraguimov@ttu.edu. Partially supported by DMS NSF grant 1412796

[†]St.Petersburg Department of Steklov Institute, Fontanka 27, St.Petersburg, 191023, Russia, and St.Petersburg State University, Universitetskii pr. 28, St.Petersburg, 198504, Russia. E-mail: al.il.nazarov@gmail.com. Supported by RFBR grant 15-01-07650.

At point $\zeta \in \overline{\Gamma_1} \cap \overline{\Gamma_2}$ function u is not defined, and we investigate asymptotic properies of the solution at this point.

For divergence type equation in case of Dirichlet Data this type of theorem first was proved in very general case by Mazya in [8]. Criteria for regularity for Zaremba problem first was obtained by Mazya in [7].

Here we consider the case of non-divergence equation in bounded domain Ω where Neumann Γ_2 is Lipschitz in a neighborhood of the point ζ .

In the case $\Gamma_2 = \emptyset$ the similar question was discussed by E.M. Landis (see [1,2]) and sharpened by Yu.A. Alkhutov [6].

We always assume that the matrix of leading coefficients (a_{ij}) is bounded, measurable and symmetric, and satisfies the uniform ellipticity condition:

$$\max_{|\xi|=1} \sup_{x \in \Omega} e(x,\xi) =: e_1 < \infty,$$

where e is the ellipticity function (see [1], [6])

$$e(x,\xi) = \frac{\sum_{i=1}^{n} a_{ii}(x)}{\sum_{i,j=1}^{n} a_{ij}(x)\xi_i\xi_j}.$$

For simplicity we consider the operators without lower-order terms, a more general case can be easily managed.

The paper is organized as follows.

In Sec. 2 we formulate some known results about non-divergence equations: lemma on non-tangential derivatives at point of maximum (minimum) on the boundary in the form of Nadirashvili [9], the Landis Growth Lemma in case $\Gamma_2 = \emptyset$, and Growth Lemma in Krylov's form(see [12]).

The Growth Lemma for elliptic and parabolic equations first was introduced by Landis in [3,4]. Growth Lemma is a fundamental tool to study qualitative properties and regularity of solutions in bounded and unbounded domain. Recent review on Growth Lemma and its applications was published in [11] (see also [13]).

In Sec. 3 we prove strict Growth Lemma near Neumann boundary.

Sec. 4 glues two Growth Lemmas. This result was obtained under some admissibility constraint on the boundary Γ_2 , which is an analog of isoperimetric condition.

In the last Sec. 5, dichotomy theorem is proved for solutions of mixed boundary value problem to non-divergence elliptic equation.

We use the following notation. $x = (x', x_n) = (x_1, \ldots, x_{n-1}, x_n)$ is a point in \mathbb{R}^n . B(x, R) is the ball centered in x with radius R.

2 Preliminary Results

Here we recall some known results and prove auxiliary lemmas for the sub- and supersolution of the equation $\mathcal{L}u = 0$. We call function u sub-elliptic (superelliptic) if $u \in W_n^2(\Omega) \bigcap \mathcal{C}^1(\Omega \cup \Gamma_2)$, and $\mathcal{L}u \leq 0$ (respectively, $\mathcal{L}u \geq 0$).

We say that Γ_2 satisfies inner cone condition (see, e.g., [9]) if there are $0 < \varphi < \pi/2$ and h > 0 such that for any $y \in \Gamma_2$ there exists a right cone $K(y) \subset \Omega$ with the apex at y, apex angle φ and of the height h.



Figure 1: Inner cone condition

In [9] N. Nadirashvili obtained fundamental generalization of Oleinik-Hopf lemma¹, the so-called "lemma on non-tangential derivative":

Lemma 2.1. Let Γ_2 satisfy inner cone condition. Let a non-constant function u be super-elliptic (sub-elliptic) $\mathcal{L}u \geq 0$ ($\mathcal{L}u \leq 0$) in Ω . Suppose that $y \in \Gamma_2$ and $u(y) \leq u(x)$ $(u(y) \geq u(x))$ for all $x \in \Gamma_2$. Then for any neighborhood S of y on Γ_2 and for any $\varepsilon < \varphi$ there exists a point $\widetilde{x} \in S$ s.t.

$$\frac{\partial u}{\partial \ell}(\widetilde{x}) < 0$$
 $\left(\frac{\partial u}{\partial \ell}(\widetilde{x}) > 0\right)$

for any outward direction ℓ s.t. the angle γ between ℓ and the axis of $K(\tilde{x})$ is not greater then $\varphi - \varepsilon$.

From standard maximum principle and Lemma 2.1 follows comparison theorem for mixed boundary value problem.

Lemma 2.2. Let Ω be a bounded domain, $\partial \Omega = \Gamma_1 \cup \Gamma_2$. Let Γ_2 satisfy inner cone condition. Suppose that vector field ℓ satisfies the same condition as in Lemma 2.1. Let functions u and v belong to $W_n^2(\Omega) \bigcap \mathcal{C}^1(\Omega \cup \Gamma_2) \cap \mathcal{C}(\overline{\Omega})$. Then, if $\mathcal{L}u \leq \mathcal{L}v$ in Ω , $u \leq v$ on Γ_1 , and $\frac{\partial u}{\partial \ell} \leq \frac{\partial v}{\partial \ell}$ on Γ_2 then $u \geq v$ in $\overline{\Omega}$.

Definition 2.3. Let Ω be a domain, $\partial \Omega = \Gamma_1 \cup \Gamma_2$. Define "small ball" B(0, R)and "big ball" B(0, aR), a > 1 (see Fig. 2).

We call the function w **barrier** with respect to mixed boundary value problem in these two balls if it posses properties:

> w is sub-elliptic $(\mathcal{L}w \leq 0)$ in the intersection $\Omega \cap B(0, aR)$; (2.1)

¹In [9] classical solutions $u \in \mathcal{C}^2(\Omega) \cap \mathcal{C}^1(\overline{\Omega})$ are used but due to the Aleksandrov-Bakel'man maximum principle it is transferred to $u \in W_n^2(\Omega) \cap \mathcal{C}^1(\Omega \cup \Gamma_2)$.

$$w(x) \le 1 \text{ on } \Gamma_1 \cap B(0, aR); \tag{2.2}$$

$$\frac{\partial w}{\partial \ell} \le 0 \ on \ \Gamma_2 \cap B(0, aR); \tag{2.3}$$

$$w \le 0 \ on \ \overline{\Omega} \cap \partial B(0, aR); \tag{2.4}$$

$$w(x) \ge \eta_0$$
 in the intesection $B(0, R) \cap \Omega$ (2.5)

for some constant η_0 .



Figure 2: Domain G and two balls B(0, R) and B(0, aR), (a > 1)

Now we are in the position to prove the following strict growth property for subsolutions of the mixed boundary value problem.

Lemma 2.4. Let Ω be a domain, $\partial \Omega = \Gamma_1 \cup \Gamma_2$. Suppose that a function u be sub-elliptic in $\Omega \cap B(0, aR)$, u > 0 in Ω , u = 0 on $\Gamma_1 \cap B(0, aR)$ and $\frac{\partial u}{\partial \ell} \leq 0$ on $\Gamma_2 \cap B(0, aR)$. Let Γ_2 satisfy inner cone condition.

Assume that there is a barrier w in balls B(0,R) and B(0,aR).

Then

$$\sup_{\Omega \cap B(0,aR)} u \ge \frac{\sup_{\Omega \cap B(0,R)} u}{1 - \eta_0}.$$
(2.6)

Proof. Let $M = \sup_{\Omega \cap B(0,aR)} u$, and let the barrier w(x) be as in Definition 2.3. Define

$$v(x) = M(1 - w(x)).$$

Obviously $\mathcal{L}v \geq \mathcal{L}u$ in Ω , $v \geq u$ on $\Gamma_1 \cap B(0, aR)$, $\frac{\partial v}{\partial \ell} \geq \frac{\partial u}{\partial \ell}$ on Γ_2 , and $v \geq M \geq u$ on $\partial B(0, aR) \cap \Omega$. Applying comparison Lemma 2.2 to functions v and u in the domain $\Omega \cap B(0, aR)$ we get that $v \geq u$. In the intersection $\Omega \cap B(0, R)$ this gives with regard of (2.5)

$$M(1-\eta_0) \ge M(1-\inf_{\Omega \cap B(0,R)} w) \ge \sup_{\Omega \cap B(0,R)} u.$$

The latter is equivalent to statement in (2.6).

We recall the well-known notion of s-capacity, see, e.g., [1, Sec. I.2].

Definition 2.5. Let H be a Borel set. Let a measure μ be defined on Borel subsets of H. We call μ admissible and write $\mu \in \mathcal{M}(H)$ if

$$\int_{H} \frac{d\mu(y)}{|x-y|^{s}} \le 1, \quad for \quad x \in \mathbb{R}^{n} \setminus H.$$

Then the quantity

$$\mathbf{C}_s(H) = \sup_{\mu \in \mathcal{M}(H)} \mu(H)$$

is called s-capacity of H.

We also recall the following simple statement.

Proposition 2.6. If $s \ge e_1 - 2$ then $L|x|^{-s} \le 0$.

Now we formulate a variant of the Landis Growth Lemma, see [1, Sec. I.4].

Lemma 2.7. Let function u be sub-elliptic in $\Omega \cap B(0, aR)$, u > 0 in Ω , u = 0 on $\Gamma_1 = \partial \Omega \cap B(0, aR)$. Let $s \ge e_1 - 2$. Then there exists $0 < \eta_1 < 1$ depending only on s s.t.

$$\sup_{\Omega \cap B(0,aR)} u \ge \frac{\sup_{\Omega \cap B(0,R)} u}{1 - \eta_1 \mathbf{C}_s(H) R^{-s}}.$$

Here $H = \Gamma_1 \cap B(0, R)$.

Consequently if $B(0,R) \setminus \Omega$ contains a ball with radius δR then

$$\sup_{\Omega\cap B(0,aR)} u \geq \frac{\sup_{\Omega\cap B(0,R)} u}{1-\widetilde{\eta}_1},$$

where the constant $\tilde{\eta}_1$ depend on s and δ .

3 Growth Lemma near Neumann boundary

Here we prove the Growth Lemmas in the domain adjunct to Γ_2 under some assumption on Γ_1 .

We recall that Γ_2 is uniformly Lipschitz in a neighborhood of x^0 . This means that there is $\delta > 0$ s.t. the set $\Gamma_2 \cap B(x^0, \delta)$ is the graph $x_n = f(x')$ in a local Cartesian coordinate system, and the function f is Lipschitz. Moreover, we suppose that its Lipschitz constant does not exceed L. Without loss of generality we assume that $\Omega \cap B(x^0, \delta) \subset \{x_n < f(x')\}$ (see Fig. 3). This implies the inner cone condition if we direct the axis of the cone K along $-x_n$ and set $\varphi = \cot^{-1}(L)$.

Lemma 3.1. Let $\Gamma_2 \cap B(0, R) = \emptyset$, and $x^0 \in \Gamma_2 \cap \partial B(0, R)$, for some $R \leq \frac{\delta}{2}$. Assume that $\Omega \cap B(0, \alpha R) = \emptyset$ for some $0 < \alpha < \frac{1}{2}$ (see Fig. 3).

Suppose that the vector field ℓ satisfies conditions in Lemma 2.1 uniformly on Γ_2 (that is, ε does not depend on $x \in \Gamma_2$).

Let function u be sub-elliptic ($\mathcal{L}u \leq 0$ in Ω), u > 0 in Ω , u = 0 on Γ_1 and $\frac{\partial u}{\partial \ell} \leq 0$ on Γ_2 .



Figure 3: Domain Ω , boundary Γ_2 and balls B(0, R), B(0, aR) and $B(0, \alpha R)$.

Then there exists a > 1 depending on the Lipschitz constant L, ε and ellipticity constant e_1 s.t.

$$\sup_{\Omega \cap B(0,aR)} u \ge \frac{\sup_{\Omega \cap B(0,R)} u}{1 - \eta_2}.$$
(3.1)

Here $\eta_2 \in (0,1)$ is defined by α and a.

Proof. We take $s \ge e_1 - 2$ and set

$$w(x) = \frac{\alpha^s R^s}{|x|^s} - \frac{\alpha^s}{a^s}.$$

We claim that for a sufficiently close to 1 this function satisfies all conditions in Definition 2.3. Indeed:

1. From Proposition 2.6 function w is sub-elliptic, condition (2.1) holds.

2. Evidently w = 0 on $\partial B(0, aR)$, condition (2.4) holds,

3. while $\Omega \cap B(0, \alpha R) = \emptyset$ implies $w \leq 1$ in $\Omega \cap B(0, \alpha R)$ (and therefore on Γ_1) condition (2.2) holds.

Now we check condition (2.3). We introduce the Cartesian coordinate system with axes collinear with those of local coordinate system at x^0 . We observe that the assumption $\Gamma_2 \cap B(0, R) = \emptyset$ and Lipschitz condition imply that for $x \in \Gamma_2 \cap B(0, aR)$

$$|x'| \le \frac{R}{\sqrt{1+L^2}} \left(L + \sqrt{a^2 - 1}\right); \qquad x_n \ge \frac{R}{\sqrt{1+L^2}} \left(1 - L\sqrt{a^2 - 1}\right).$$

Moreover, our assumption on the vector field ℓ means that

$$|\ell'| \le \sin(\cot^{-1}(L) - \varepsilon) \le \frac{1}{\sqrt{1 + L^2}} - \widetilde{\varepsilon};$$
$$\ell_n \ge \cos(\cot^{-1}(L) - \varepsilon) \ge \frac{L}{\sqrt{1 + L^2}} + \widetilde{\varepsilon}$$

where $\tilde{\varepsilon}$ depends only on L and ε .

Therefore, the direct calculation gives

$$\begin{split} \frac{\partial w}{\partial \ell}(x) &= -\frac{s\alpha^s R^s}{|x|^{s+2}} \cdot (x_n \ell_n + \ell' \cdot x') \\ &\leq \frac{s\alpha^s R^s}{|x|^{s+2}} \cdot \frac{R}{\sqrt{1+L^2}} \Big(\sqrt{a^2 - 1} \cdot \left(\sqrt{1+L^2} + \widetilde{\varepsilon}(L-1)\right) - \widetilde{\varepsilon}(L+1)\Big). \end{split}$$

It is easy to see that, given $\tilde{\varepsilon} > 0$, there is a > 1 depending only on $\tilde{\varepsilon}$ and L s.t. $\frac{\partial w}{\partial \ell}(x) \leq 0$, and (2.3) holds.

Finally, for $x \in \Omega \cap B(0, R)$, $w(x) \ge \alpha^s (1 - a^{-s}) =: \eta_2$, and (2.5) holds.

Thus, the claim follows, and w is the barrier in the balls B(0, R), B(0, aR). From Lemma 2.4 we get (3.1).

Growth Lemma in the Spherical Layer 4

In this section we prove Growth Lemma in spherical layer near junction point of interest $\zeta = \overline{\Gamma}_1 \cap \overline{\Gamma}_2$. Without loss of generality we put $\zeta = 0$.

First we will introduce admissible class of domains in the spherical layer.

Definition 4.1. Fix five constants $0 < q_1 < q_2 < q^* < q_3 < q_4$. Define two spherical layers $\hat{U}_R \subset U_R$:

$$U_R = B(0, q_4 R) \setminus B(0, q_1 R);$$
 $\hat{U}_R = B(0, q_3 R) \setminus B(0, q_2 R)$

We call Ω admissible in the layer U_R if for some $\theta > 0$ there is finite set of the balls (see Fig. 4)

$$\mathcal{B} = \{ B^k = B(\xi_k, \theta R) \}_{k=0}^N; \quad B^k \subset \hat{U}_R$$

s.t. the following holds:

1. $\mathbf{C}_{s}(B^{0} \cap \widetilde{\Gamma}_{1}) \geq \varkappa \mathbf{C}_{s}(\Gamma_{1} \cap \widehat{U}_{R}), \text{ for some constant } \varkappa > 0.$ 2. $B^{k} \cap \Gamma_{2} = \emptyset, \ k = 1, ..., N, \ and \ B(\xi_{0}, a\theta R) \cap \Gamma_{2} = \emptyset, \text{ where } a > 1 \text{ is defined}$ in Lemma 3.1. 2

3. There is $\delta \in (0, 1/2)$ s.t. every ball in \mathcal{B} can be connected with B^0 by a subsequence of balls B^j s.t. any intersection $B^j \cap B^{j+1} \cap \Omega$ contains the ball $B(\xi_{j+1}, \delta R).$

4. The set $S_R = \partial B(0, q^*R) \cap \Omega$ is covered by balls in \mathcal{B} .

Fig. 4 schematically illustrate Definition 4.1.

Lemma 4.2. Let function u be sub-elliptic, u > 0 in Ω . Suppose that $u \leq 0$ on Γ_1 and $\frac{\partial u}{\partial \ell} \leq 0$ on Γ_2 . Let domain Ω be admissible in the layer U_R . Then

$$\sup_{\Omega} u \ge \frac{\sup_{S_R} u}{1 - \eta \mathbf{C}_s(H) R^{-s}}$$

²Note that boundaries of some balls B^k may touch Γ_2 .



Figure 4: On the left: domain Ω admissible in Spherical Layer U_R . On the right: domain and layer zoomed near boundary Γ_2 (bold line).

Here $H = \Gamma_1 \cap \hat{U}_R$ while η depends on s, the ellipticity constant e_1 , the Lipschitz constant L, the vector field ℓ , constants θ , \varkappa , δ in Definition 4.1 and the number N of balls in the set \mathcal{B} .

Proof. Without loss of generality we set $\theta = 1$. Let $\sup_{S_R} u =: m = u(y)$, here $y \in \overline{S}_R$. By assumption 4 in Definition 4.1, $y \in \overline{B}^k$ for some k. By assumption 3, we can choose a subsequence B^j connecting B^0 and B^k .

Consider the ball B_0 and the ball $B(\xi_0, aR)$, a > 1, concentric to it. Due to assumptions 1 and 2 in Definition 4.1, we can apply Lemma 2.7 to get:

$$M := \sup_{\Omega} u \ge \sup_{\Omega \cap B(\xi_0, aR)} u \ge \frac{\sup_{B^0 \cap \Omega} u}{1 - \varkappa \eta_1 \mathbf{C}_s(H) R^{-s}}.$$

Suppose that

$$\sup_{B^0 \cap \Omega} u \ge m(1 - \delta_0), \quad \text{where} \quad \delta_0 = \frac{\varkappa \eta_1 \mathbf{C}_s(H) R^{-s}}{2(1 - \varkappa \eta_1 \mathbf{C}_s(H) R^{-s})}.$$
(4.1)

Then after some calculation we get

$$M \ge \frac{m}{1 - \eta_3 \mathbf{C}_s(H) R^{-s}}$$

for some η_3 depending on $\varkappa \eta_1$, and the statement follows.

If (4.1) does not hold, we consider the function

$$u_1(x) = u(x) - m(1 - \delta_0), \qquad (4.2)$$

then $u_1(x) \leq 0$ in $B^0 \cap \Omega$.

By assumption 3, $B^0 \cap B^1 \cap \Omega$ contains a ball of radius δR . Let $\Omega_1 := \{x : u_1(x) > 0\}$. Assume that $B^1 \cap \Omega_1 \neq \emptyset$, otherwise we consider the first ball in the subsequence B^j for which this property holds.

Suppose that

$$\sup_{B^1 \cap \Omega} u_1 \ge m \delta_0 (1 - \tau), \tag{4.3}$$

here the constant τ will be chosen later.

Consider any simply connected component of the domain $B(\xi_1, aR) \cap \Omega_1$ in which the supremum in (4.3) is realised. There are two possibilities:

a) $B(\xi_1, aR) \cap \Gamma_2 = \emptyset;$

b) $B(\xi_1, aR) \cap \Gamma_2 \neq \emptyset$

(recall that $a = a(L, \ell, e_1) > 1$ is defined in Lemma 3.1).

$$\sup_{B(\xi_1, aR) \cap \Omega} u_1 \ge \frac{\sup_{B^1 \cap \Omega} u_1}{1 - \widetilde{\eta}_1} \ge \frac{m\delta_0(1 - \tau)}{1 - \widetilde{\eta}_1}.$$
(4.4)

Using (4.2) and (4.4) we deduce

$$\sup_{B(\xi_1, aR) \cap \Omega} u \ge m \left(1 + \frac{\delta_0(\widetilde{\eta}_1 - \tau)}{1 - \widetilde{\eta}_1} \right).$$

Letting $\tau = \frac{\tilde{\eta}_1}{2}$ we get

$$M \ge \sup_{B(\xi_1, aR) \cap \Omega} u \ge m \left(1 + \frac{\delta_0 \tau}{1 - 2\tau} \right), \tag{4.5}$$

and the statement follows.

In case of (b) we proceed with the same arguments but instead of Lemma 2.7 we apply Lemma 3.1 and put $\tau = \frac{\eta_2}{2}$. Thus, if (4.3) holds with $\tau = \frac{1}{2} \min\{\tilde{\eta}_1, \eta_2\}$ then (4.5) is satisfied in any case, and Lemma is proved.

If (4.3) does not hold then function u satisfies

$$\sup_{B^1 \cap \Omega} u \le m(1 - \delta_0 \tau)$$

As in previous step we consider the function

$$u_2(x) = u(x) - m(1 - \delta_0 \tau),$$

 $u_2(x) \leq 0$ in $B^1 \cap \Omega$.

Repeating previous argument we deduce that if

$$\sup_{B^2 \cap \Omega} u_2 \ge m \delta_0 \tau (1 - \tau) \tag{4.6}$$

then

$$M \ge m \big(1 + \frac{\delta_0 \tau^2}{1 - 2\tau}\big),$$

and Lemma is proved.

If (4.6) does not hold, then

$$\sup_{B^2 \cap \Omega} u \le m(1 - \delta_0 \tau^2).$$

Repeating this process we either prove Lemma or arrive at the inequality

$$\sup_{B^k \cap \Omega} u \le m(1 - \delta_0 \tau^k)$$

that is impossible since $y \in \overline{B}^k$ and u(y) = m.

5 Dichotomy of solutions

In this section we will apply obtained Growth Lemma in spherical layer to prove dichotomy of solutions near point ζ of the junction of Dirichlet and Neumann boundaries. As in previous section we put $\zeta = 0$.

Let $\Omega \subset \{x : x_n < f(x')\}$ and Γ_2 is a graph of the function $x_n = f(x')$, f(0) = 0. Set $R_m = Q^{-m}$ for some Q > 1, $S_m = \partial B(0, q^*R_m)$, and

$$U_m = B(0, q_4 R_m) \setminus B(0, q_1 R_m), \quad U_m = B(0, q_3 R_m) \setminus B(0, q_2 R_m).$$

We fix $N_0 \in \mathbb{N}$ and $q_1 < q_2 < q^* < q_3 < q_4$ s.t. $q^* < q_1 Q$. Suppose that for all $m \ge N_0$ the domain Ω with boundaries Γ_1 and Γ_2 is admissible in the layer U_m in the sense of Definition 4.1 with $R = R_m$, and all constants in Definition 4.1 do not depend on m.

Lemma 5.1. Let function u be sub-elliptic, u > 0 in Ω . Suppose that $u \leq 0$ on $\Gamma_1 \cap B(0, q_4 R_{N_0})$ and $\frac{\partial u}{\partial \ell} \leq 0$ on $\Gamma_2 \cap B(0, q_4 R_{N_0})$. Let domain Ω be admissible in the layers U_m , $m \geq N_0$.

Let $M_m = \sup_{S_m \cap \Omega} u$. Then one of two statements holds:

either $M_{N_1+1} \ge M_{N_1}$ for some N_1 , and for all $m > N_1$

$$M_{m+1} \ge \frac{M_m}{1 - \eta \mathbf{C}_s(H_m)Q^{sm}},\tag{5.1}$$

or for all $m > N_0$

$$M_m \ge \frac{M_{m+1}}{1 - \eta \mathbf{C}_s(H_m)Q^{sm}}.$$
 (5.2)

Here $H_m = \Gamma_1 \cap \hat{U}_m$, and η is the constant from Lemma 4.2.

Proof. Due to Lemma 2.2, there are two possibilities:

(a) if $M_{N_1+1} \ge M_{N_1}$ for some $N_1 > N_0$ then $M(\rho) = \sup_{\partial B(0,\rho) \cap \Omega} u > M_m$, $m > N_1$ for any $\rho < q^* R_m$;

(b) otherwise $M_m > M_{m+1}$ for all $m > N_0$.

Now Lemma 4.2 gives (5.1) in the case (a) and (5.2) in the case (b). \Box

Remark 5.2. Let function u be sub-elliptic, u > 0 in Ω . Suppose that $u \leq 0$ on $\Gamma_1 \cap B(0, \rho_0)$ and $\frac{\partial u}{\partial \ell} \leq 0$ on $\Gamma_2 \cap B(0, \rho_0)$. Then the maximum principle implies the following dichotomy (we recall that $M(\rho) = \sup_{\partial B(0,\rho) \cap \Omega} u$):

either there is $\rho^* \leq \rho_0$ s.t. for $\rho_2 < \rho_1 < \rho^*$ we have $M(\rho_2) > M(\rho_1)$; or $M(\rho_2) < M(\rho_1)$ for all $\rho_2 < \rho_1 < \rho_0$.

Applying recursively alternative in Lemma 5.1 and using Remark 5.2 we get asymptotic dichotomy.

Theorem 5.3. Let the assumptions of Lemma 5.1 be satisfied. Suppose that $\sum_{m=0}^{\infty} \mathbf{C}_s(H_m)Q^{sm} = \infty$, where $H_m = \Gamma_1 \cap \hat{U}_m$. Then one of two statements holds:

either $M(\rho) \to \infty$ as $\rho \to 0$, and

$$\liminf_{\rho \to \infty} M(\rho) \exp \Big(- \widehat{\eta} \sum_{m=0}^{[c \ln \rho]} \mathbf{C}_s(H_m) Q^{sm} \Big) > 0,$$

or $M(\rho) \to 0$ as $\rho \to 0$, and

$$\limsup_{\rho \to \infty} M(\rho) \exp\left(\widehat{\eta} \sum_{m=0}^{[c \ln \rho]} \mathbf{C}_s(H_m) Q^{sm}\right) = 0,$$

Here $\hat{\eta}$ and c depend on the same quantities as η in Lemma 4.2.

References

- Landis, E.M. Second Order Equations of Elliptic and Parabolic Type, Moscow, Nauka, 1971 (Russian). English transl.: Translation of Mathematical Monograph, 171, AMS, Providence, Rhode Island, 1998.
- [2] Landis, E.M. s-capacity and its application to the study of solutions of a second order elliptic equation with discontinuous coefficients, Mat. Sb. (N.S.) 76 (118) (1968), N2, 186–213 (Russian). English transl.: Math. USSR-Sb., 5 (1968), N2, 177–204.
- [3] Landis, E.M. On some properties of the solutions of elliptic equations. Dokl. Akad. Nauk SSSR, 107 (1956), No 4.
- [4] Landis, E.M. Some problems of the qualitative theory of elliptic and parabolic equations, UMN 14 (1959), N1 (85), 21–85 (Russian).
- [5] Landis, E.M. Some problems of the qualitative theory of second order elliptic equations (case of several independent variables), UMN 18 (1963), N1 (109), 3–62 (Russian). English transl.: Russian Math. Surveys, 18 (1963), N1, 1–62.
- [6] Alkhutov, Yu.A. On the regularity of boundary points with respect to the Dirichlet problem for second-order elliptic equations. Mat. Zametki, **30** (1981), N3, 333–342 (Russian). English transl.: Math. Notes, **30** (1981), N3, 655-660.
- [7] Kerimov, T.M., Maz'ya, V.G., Novruzov, A.A., An analogue of the Wiener criterion for the Zaremba problem in a cylindrical domain, Funk. Anal. i Prilozh., 16 (1982), N4, 70–71 (Russian). English transl.: Funct. Analysis and Its Applic., 16 (1982), N4, 301–303.
- [8] Maz'ja, V.G. The behavior near the boundary of the solution of the Dirichlet problem for an elliptic equation of the second order in divergence form. Mat. Zametki 2 (1967), N2, 209–220 (Russian). English transl.: Math. Notes, 2 (1967), N2, 610–617.
- [9] Nadirashvili, N.S. Lemma on the interior derivative and uniqueness of the solution of the second boundary value problem for second-order elliptic equations, Dokl. Akad. Nauk SSSR, 261 (1981), N4, 804–808 (Russian).
- [10] Nadirashvili, N. S. On the question of the uniqueness of the solution of the second boundary value problem for second-order elliptic equations, Mat. Sb. (N.S.) 122 (164) (1983), N3 (11), 341–359 (Russian). English transl.: Math. USSR-Sb. 50 (1985), N2, 325–341.
- [11] Safonov, M.V., Non-divergence Elliptic Equations of Second Order with Unbounded Drift, In: Nonlinear Partial Differential Equations and Related Topics; Dedicated to Nina N. Uraltseva. Advances in Mathematical Sciences-64. AMS Translations, Ser. 2, 229 (2010).
- [12] Krylov, N. V.; Safonov, M. V. A property of the solutions of parabolic equations with measurable coefficients. Izv. Akad. Nauk SSSR Ser. Mat. 44 (1980), no. 1, 161175, 239.

[13] Aimar, H.; Forzani, L.; Toledano, R., Hölder regularity of solutions of PDE's: a geometrical view, Comm. PDE, 26 (2001), N7–8, 1145-1173.