

Characterization of certain sequences of q -polynomials

P. Njionou Sadjang

Abstract. A characterization is given of those sequences of quasi-orthogonal polynomials which form also q -Appell sets.

Mathematics Subject Classification (2010). 33D45; 33C65; 05A40.

Keywords. Quasi-orthogonality, q -Appell sets, Al-Salam Carlitz polynomials.

1. Introduction

A sequence of polynomials $\{Q_n\}$, $n = 0, 1, 2, \dots$, $\deg Q_n(x) = n$ is said to be quasi-orthogonal if there is an interval (a, b) and a non-decreasing function $\alpha(x)$ such that

$$\int_a^b x^m Q_n(x) d\alpha(x) \begin{cases} = 0 & \text{for } 0 \leq m \leq n-2 \\ \neq 0 & \text{for } 0 \leq m = n-1 \\ \neq 0 & \text{for } 0 = m = n. \end{cases}$$

We say that two polynomial sets are related if one set is quasi-orthogonal with respect to the interval and the distribution of the orthogonality of the other set. Riesz [9] and Chihara [3] have shown that a necessary and sufficient condition for the quasi-orthogonality of the $\{Q_n(x)\}$ is that there exist nonzero constants, $\{a_n\}_{n=0}^\infty$ and $\{b_n\}_{n=1}^\infty$, such that

$$\begin{aligned} Q_n(x) &= a_n P_n(x) + b_n P_{n-1}(x), & n \geq 1, \\ Q_0(x) &= a_0 P_0(x) \end{aligned} \quad (1.1)$$

where the $\{P_n(x)\}_{n=0}^\infty$ are the related orthogonal polynomials.

A polynomial set $\{P_n(x)\}$ forms a q -Appell set if the condition [10, 1] given by

$$D_q P_n(x) = [n]_q P_{n-1}(x), \quad n = 0, 1, 2, 3, \dots \quad (1.2)$$

is fulfilled. Here, D_q is the Hahn derivative defined by

$$D_q f(x) = \frac{f(x) - f(qx)}{(1-q)x}, \quad x \neq 0$$

and $[n]_q$ stands for the q -number defined by $[n]_q = \frac{1-q^n}{1-q}$. Such sets were first introduced by Sharma and Chak [10] to generalize the Appell set [2]. Several papers related to q -Appell sets have been recently published, see for instance [5] and the references therein.

In 1967, Al-Salam has given in [1] a characterization of those sequences of orthogonal polynomials $\{P_n(x)\}$ which are also q -Appell sets. More precisely, he gave a characterization of those sequences of orthogonal polynomials for which $D_q P_n(x) = [n]_q P_{n-1}(x)$ for $n = 0, 1, 2, 3, \dots$

The purpose of this paper is to study those classes of polynomial sets $\{P_n(x)\}$ that are at the same time quasi-orthogonal sets and q -Appell sets.

2. Preliminaries results and definitions

Let us introduce the so-called q -Pochhammer symbol

$$(x; q)_n = \begin{cases} (1-x)(1-xq) \cdots (1-xq^{n-1}) & n = 1, 2, \dots \\ 1 & n = 0 \end{cases}$$

and the basic hypergeometric or q -hypergeometric function ${}_r\phi_s$ is defined by the series

$${}_r\phi_s \left(\begin{matrix} a_1, \dots, a_r \\ b_1, \dots, b_s \end{matrix} \middle| q; z \right) := \sum_{k=0}^{\infty} \frac{(a_1, \dots, a_r; q)_k}{(b_1, \dots, b_s; q)_k} \left((-1)^k q^{\binom{k}{2}} \right)^{1+s-r} \frac{z^k}{(q; q)_k},$$

where

$$(a_1, \dots, a_r)_k := (a_1; q)_k \cdots (a_r; q)_k.$$

The Al Salam-Carlitz I polynomials [8, P. 534] have the q -hypergeometric representation

$$U_n^{(a)}(x; q) = (-a)^n q^{\binom{n}{2}} {}_2\phi_1 \left(\begin{matrix} q^{-n}, x^{-1} \\ 0 \end{matrix} \middle| q; \frac{qx}{a} \right)$$

and fulfill the three-term recurrence relation

$$xU_n^{(a)}(x; q) = U_{n+1}^{(a)}(x; q) + (a+1)q^n U_n^{(a)}(x; q) - aq^{n-1}(1-q^n)U_{n-1}^{(a)}(x; q)$$

and the q -derivative rule

$$D_q U_n^{(a)}(x; q) = [n]_q U_{n-1}^{(a)}(x; q).$$

It is therefore clear that the Al-Salam Carlitz I polynomials form a q -Appell set.

Proposition 2.1. (See [4, Theorem 1]) For $\{Q_n(x)\}$ to be a set of polynomials quasi-orthogonal with respect to an interval (a, b) and a distribution $d\alpha(x)$, it is necessary and sufficient that there exist a set of nonzero constants $\{T_k\}_{k=0}^{\infty}$ and a set of polynomials $\{P_n(x)\}$ orthogonal with respect to (a, b) and $d\alpha(x)$ such that

$$P_n(x) = \sum_{k=0}^n T_k Q_k(x), \quad n \geq 0. \quad (2.1)$$

Proposition 2.2. (See [4, Theorem 2]) A necessary and sufficient condition that the set $\{Q_n(x)\}_{n=0}^{\infty}$ where each $Q_n(x)$ is a polynomial of degree precisely n , be quasi-orthogonal is that it satisfies

$$Q_{n+1}(x) = (x + b_n)Q_n(x) - c_n Q_{n-1}(x) + d_n \sum_{k=0}^{n-2} T_k Q_k(x),$$

for all n , with $d_0 = d_1 = 0$.

Proposition 2.3. (See [1, Theorem 4.1]) If $\{Q_n(x)\}_{n=0}^{\infty}$ is a q -Appell set which are also orthogonal, then there exists a non zero constant b such that

$$Q_n(x) = b^n U_n^{(a/b)}\left(\frac{x}{b}; q\right)$$

for all $n \geq 0$.

3. A characterization result

In this section, we characterize quasi-orthogonal polynomial sets that are also q -Appell sets.

Theorem 3.1. If $\{Q_n(x)\}_{n=0}^{\infty}$ is a q -Appell set of quasi-orthogonal polynomials, then there exist three real numbers b, c and λ , such that

$$Q_{n+1}(x) = (x + bq^n)Q_n(x) - cq^{n-1}[n]_q Q_{n-1}(x) + d_n \sum_{k=0}^{n-2} \frac{\lambda^k}{[k]_q!} Q_k(x). \quad (3.1)$$

Proof. Assume that $\{Q_n(x)\}_{n=0}^{\infty}$ is a q -Appell set which is quasi-orthogonal and $\{P_n(x)\}_{n=0}^{\infty}$ the related orthogonal family. From Proposition 2.2, there exist three sequences $\{a_n\}_{n=0}^{\infty}$, $\{b_n\}_{n=0}^{\infty}$ and $\{d_n\}_{n=0}^{\infty}$ with $d_0 = d_1 = 0$ such that

$$Q_{n+1}(x) = (x + b_n)Q_n(x) - c_n Q_{n-1}(x) + d_n \sum_{k=0}^{n-2} T_k Q_k(x). \quad (3.2)$$

If we q -differentiate (3.2) and use the fact that $\{Q_n(x)\}_{n=0}^{\infty}$ is a q -Appell, we get after some simplifications

$$Q_n(x) = \left(x + \frac{b_n}{q}\right) Q_{n-1}(x) - \frac{c_n [n-1]_q}{q [n]_q} Q_{n-2}(x) + \frac{d_n}{q [n]_q} \sum_{k=0}^{n-3} [k+1]_q T_{k+1} Q_k(x). \quad (3.3)$$

Next, if we replace n by $n - 1$ in (3.2), we obtain

$$Q_n(x) = (x + b_{n-1})Q_{n-1}(x) - c_{n-1}Q_{n-2}(x) + d_{n-1} \sum_{k=0}^{n-3} T_k Q_k(x). \quad (3.4)$$

If we compare (3.3) and (3.4), we see that we should have

$$b_n = qb_{n-1}, \quad c_n = q \frac{[n]_q}{[n-1]_q} c_{n-1}, \quad (3.5)$$

and

$$d_n [k+1]_q T_{k+1} = q [n]_q d_{n-1} T_k, \quad k = 0, 1, \dots, n-3. \quad (3.6)$$

Equations (3.5) yield

$$b_n = q^n b_0, \quad \text{and} \quad c_n = q^{n-1} [n]_q c_1.$$

Next, (3.6) gives for $k = 0$ and $k = n - 3$ the relations

$$d_n = \frac{q[n]_q}{T_1} d_{n-1} \quad \text{and} \quad d_n = \frac{q[n]_q T_{n-1}}{[n-2]_q T_{n-2}} d_{n-1}. \quad (3.7)$$

If, for a given $k \geq 2$, $d_k = 0$, it follows from (3.7) that $d_k = 0$ for all k . In this case (3.2) becomes a three-term recurrence relation

$$Q_{n+1}(x) = (x + b_n)Q_n(x) - c_n Q_{n-1}(x). \quad (3.8)$$

In this case, from Proposition 2.3, it is seen that $\{Q_n(x)\}_{n=0}^{\infty}$ is essentially the sequence of Al-Salam Carlitz polynomials. Thus, in this case, $\{Q_n(x)\}_{n=0}^{\infty}$ is not a sequence of quasi-orthogonal polynomials. Thus, we must have $d_k \neq 0$ for $k \geq 2$.

Again, using (3.7), we have for all $n \geq 0$ the identity $\frac{T_{n-1}}{[n]_q T_n} = \frac{1}{T_1}$. This last relation gives $T_n = \frac{T_1^n}{[n]_q!}$. Setting $b_0 = b$, $c_1 = c$ and $T_1 = \lambda$, this ends the proof of the theorem. \square

Theorem 3.2. *Let $\{Q_n(x)\}_{n=0}^{\infty}$ be a monic polynomial set. The following assertions are equivalent:*

1. $\{Q_n(x)\}_{n=0}^{\infty}$ is quasi-orthogonal and is a q -Appell set, $n \geq 1$.
2. There exists three constants α , β and γ ($\beta, \gamma \neq 0$) such that

$$Q_n(x) = \beta^n U_n^{(\alpha/\beta)} \left(\frac{x}{\beta}; q \right) - \frac{\beta^n [n]_q}{\lambda} U_{n-1}^{(\alpha/\beta)} \left(\frac{x}{\beta}; q \right) \quad (n \geq 1),$$

where $U_n^{(a)}(x; q)$ are the Al-Salam Carlitz I polynomials.

Proof. Suppose first that $\{Q_n(x)\}_{n=0}^{\infty}$ is quasi-orthogonal and is a q -Appell set, $n \geq 1$. Then, by Theorem 3.1, the Q_n 's satisfy a recurrence relation of the form (3.1). Let us define the polynomial set $\{P_n(x)\}_{n=0}^{\infty}$ by

$$P_n(x) = \frac{[n]_q!}{\lambda^n} \sum_{k=0}^n \frac{\lambda^k}{[k]_q!} Q_k(x). \quad (3.9)$$

It is not difficult to see that

$$D_q P_n(x) = \frac{[n]_q!}{\lambda^n} \sum_{k=1}^n \frac{\lambda^k}{[k]_q!} [k]_q Q_{k-1}(x) = [n]_q \frac{[n-1]_q!}{\lambda^{n-1}} \sum_{k=0}^{n-1} \frac{\lambda^k}{[k]_q!} Q_k(x) = [n]_q P_{n-1}(x).$$

Hence, $\{P_n(x)\}_{n=0}^\infty$ is a q -Appell set. Moreover, $\{P_n(x)\}_{n=0}^\infty$ is the orthogonal set (see Proposition 2.1) related to $\{Q_n(x)\}_{n=0}^\infty$. By Proposition 2.3, there exist α and β such that

$$P_n(x) = \beta^n U_n^{(\alpha/\beta)} \left(\frac{x}{\beta}; q \right).$$

Next, from (3.9), it follows easily that

$$Q_n(x) = \frac{[n]_q!}{\lambda^n} (P_n(x) - P_{n-1}(x)).$$

The first implication of the theorem follows.

Conversely, assume that there exist three constants α , β and γ ($\beta, \gamma \neq 0$) such that

$$Q_n(x) = \beta^n U_n^{(\alpha/\beta)} \left(\frac{x}{\beta}; q \right) - \frac{\beta^n [n]_q}{\lambda} U_{n-1}^{(\alpha/\beta)} \left(\frac{x}{\beta}; q \right) \quad (n \geq 1).$$

It is easy to see that $\{Q_n(x)\}_{n=0}^\infty$ is a quasi-orthogonal set. It remains to prove that $\{Q_n(x)\}_{n=0}^\infty$ is a q -Appell set. Using the fact that $D_q[f(ax)] = a[D_q f](ax)$, we have

$$D_q U_n^{(\alpha/\beta)} \left(\frac{x}{\beta}; q \right) = \frac{1}{\beta} U_{n-1}^{(\alpha/\beta)} \left(\frac{x}{\beta}; q \right).$$

It follows that $D_q Q_n(x) = [n]_q Q_{n-1}(x)$. This ends the proof of the theorem. \square

References

- [1] W. A. Al-Salam, *q-Appell polynomials*, Ann. Mat. Pura Appl., **4** (1967), pp. 31–45.
- [2] P. Appell, *Une classe de polynomes*, Ann. Sci. Éc. Norm. Supér., **9** (1880), pp. 119–144.
- [3] T. S. Chihara, *On quasi-orthogonal polynomials*, Proc. Amer. Math. Soc., **8** (1957), pp. 765–767.
- [4] D. Dickinson, *On quasi-orthogonal polynomials*, Proc. Amer. Math. Soc., **12** (1961), pp. 185–194.
- [5] T. Ernst, *On certain generalized q-Appell polynomial expansion*, Ann. Univ. Mariae Curie-Skłodowska Sect. A, **68** (2015), pp 27–50.
- [6] V. Kac and P. Cheung, *Quantum Calculus*, Springer, New York, (2001).
- [7] P. J. McCarthy, *Note on quasi-orthogonal polynomials*, Bolletino dell'Unione Matematica Italiana, **16** (1961), pp. 445–448.
- [8] R. Koekoek, P. A. Lesky and R. F. Swarttouw: *Hypergeometric Orthogonal Polynomials and their q-Analogues*, Springer, Berlin, (2010).

- [9] M. Riesz, *Sur le problème des moments. Troisième note*, Arkiv för Matematik Astronomi och Fysik, **17** (1923), pp. 1–52.
- [10] A. Sharma and A. Chak, *The basic analogue of a class of polynomials*, Revisita di Matematica della Università di Parma, **5** (1954), pp. 15–38.

P. Njionou Sadjang
University of Douala,
Faculty of Industrial Engineering
Douala
Cameroon
e-mail: pnjionou@yahoo.fr