

Previous Models of Diffuse Scattering From Protein Crystals Do Not Explain Experimental Observations

Ariana Peck^{1,*}, Frédéric Poitevin^{2,3}, and Thomas J. Lane²

¹*Department of Biochemistry, Stanford University, Stanford, CA, 94305, United States.*

²*SLAC National Accelerator Laboratory, Menlo Park, CA, 94025, United States.*

³*Department of Structural Biology, Stanford University, Stanford, CA, 94305, United States.*

(Dated: May 17, 2022)

X-ray diffuse scattering from macromolecular crystals is frequently assumed to probe biologically important dynamics, but competing models of disorder have been proposed. Here we present a comprehensive comparison of models previously used to interpret diffuse scattering and test their ability to predict the features of three experimental maps. Although simple models of intramolecular, isotropic dynamics consistently outperformed biologically-informed models, none of these models reproduced experimentally observed “speckles” indicative of long-range correlated disorder. Extending a model of liquid-like motions to account for correlations beyond the biological unit, and thus across unit cell boundaries, significantly improved the agreement with experiment but still could not account for all of the observed signal. This analysis highlights the need for new models of disorder that account for correlations across multiple length scales.

I. INTRODUCTION

X-ray diffraction images from macromolecular crystals frequently exhibit a diffuse background between and beneath Bragg peaks [1, 2]. In contrast to the Bragg reflections which arise from coherent diffraction across the crystal, this diffuse signal results from disorder-induced incoherence. Uncorrelated disorder, including from air, solvent, and macromolecular sources, yields a trivial diffuse scattering pattern that is radially symmetric [3]. On the other hand, correlated disorder produces anisotropic diffuse scattering features whose spacing and intensity in reciprocal space are respectively determined by the length scale and amplitudes of the correlated atomic displacements involved [4].

Because *correlated* displacements in macromolecules are likely related to their functional motions, and methods for directly measuring such motions with high spatial resolution are rare, diffuse scattering has routinely been cited as a promising route to understanding biomolecular function [1, 4]. Allostery, signaling, and enzyme catalysis are leading cases where an ability to probe correlated motions in macromolecules would be advantageous for understanding biological function. Indeed, this has motivated the majority of studies of diffuse scattering from macromolecular crystals [5–13], despite the technical challenges of measuring and the computational cost of modeling this signal [1].

A more recently suggested possibility has been the use of diffuse scattering to complement and improve (static) structure inference from crystal diffraction. Chapman and colleagues have suggested that in cases where measurable diffuse scattering extends to higher resolution than the Bragg data, it may be employed for structural inference at that higher resolution [14]. Further, the ability to oversample the diffraction pattern by measuring the continuous diffuse signal raises the possibility of solving the phase problem directly, without resorting to anomalous or isomorphous methods. This approach, how-

ever, requires that diffuse scattering originate from specific types of disorder, such as rigid body motions, which are unlikely to inform on biological function.

Identifying the physical origins of diffuse scattering, and thus its potential for probing biological function or advancing methods, remain challenges for the field. Many types of disorder involve small motions that can be conveniently described by a matrix, whose elements give the covariation between any two atom’s displacements from their mean positions. Such a covariance matrix can directly, but not uniquely, predict diffuse scattering. If the covariance matrix could be inferred directly from experiment, the diffuse signal could be analyzed to determine which regions of the macromolecule move together. There is a fundamental problem with direct inference, however: while the number of observed independent variables is quite large (say, V voxels for a given unit cell volume, assuming the maximum resolution of diffraction is fixed), the number of unknowns is even larger (order V^2 matrix elements, one for each atom pair, assuming the number of atoms scales linearly with the volume). Thus, to infer a covariance matrix one must make simplifying assumptions about the nature of protein motions: a parsimonious model for the protein physics is required.

In this work, we analyze the parsimonious models that have been suggested previously (reviewed in Ref. [15]), and show that they do not satisfactorily explain the observed diffuse scattering – either qualitatively or quantitatively – in a range of systems. The three experimental datasets analyzed here represent both diverse crystalline properties (e.g. cryogenic and room temperature, different space groups, a range of solvent contents) and biological functions: cyclophilin A (CypA), a monomeric proline isomerase; WrpA, a tetrameric flavodoxin-like protein; and a dimeric enzyme, alkaline phosphatase (AP), bound to its transition state analog. Despite this diversity, we consistently found that models of biologically-relevant dynamics were unable to explain the observed diffuse scattering, whereas models of simpler dynamics, includ-

ing rigid body and liquid-like motions, showed modest correlation with the experimental signal. A model allowing liquid-like motions to span neighboring protein units in the crystal significantly improved the agreement both quantitatively and qualitatively, but still did not fully account for the observed signal. Though the variability of macromolecular crystals does not guarantee that these trends are generalizable, this analysis underscores the need for new, likely composite models of disorder to resolve the physical origins of diffuse scattering.

II. METHODS

A. Reconstruction of 3d diffuse scattering maps

For each experimental dataset, Bragg reflections were indexed by XDS [16]. Refined parameters from XDS were then used to index all pixels of each diffraction image. Measured intensities were corrected for polarization of the x-ray beam [17] and the difference in solid angle subtended by pixels at different scattering angles [18]. Per-image scale factors from XDS were applied to correct for differences in overall intensity across the rotation range. For datasets collected on a Pilatus detector, parallax broadening was also accounted for, as implemented in DIALS [19]. No Lorentz correction was applied, as pixel intensities were averaged rather than integrated, and the diffuse features were observed to be roughly constant across the region of measurement (i.e. individual pixels). Under these conditions, the volume of reciprocal space integrated by each pixel needs only to be corrected for the solid angle subtended, and not the arc length the pixel traverses due to rotation of the crystal [20].

After correcting for geometrical distortions, Bragg peaks were removed by implementing the spot prediction algorithm described in Ref. [21]. Pixels predicted to be spanned by a Bragg reflection were masked if their intensity exceeded three standard deviations above the mean intensities of neighboring, unmarked pixels in a 30 x 30 pixel window centered on the reflection. Prior studies have replaced masked pixel intensities by the intensities from adjoining pixels [22] or pre-filtered images [14], but the strategy of masking without replacement was found to maintain an adequate signal to noise ratio for map voxels that coincided with Miller indices (Fig. S1A, cf. solid and dashed lines). To ensure the complete removal of Bragg contaminants, an additional step of masking was performed to eliminate pixels whose intensities exceeded the median radial intensity by more than five times the median absolute deviation of that resolution shell.

The radial intensity profiles of the processed diffraction patterns were then compared for uniformity. In the case of CypA, a radially symmetric peak at $|\mathbf{q}| = 1.3 \text{ \AA}^{-1}$ was variably observed, consistent with scattering from the paratone oil in which the crystal was coated prior to data collection. The contribution of this parasitic scattering was removed by subtracting a scaled paratone

scattering profile from each image. Principal component analysis was performed on the resulting radial intensity profiles, and the first two principal components were used to remove radially asymmetric variance between images. In the case of WrpA, principal component analysis was similarly performed to ensure that radial intensity profiles were consistent across the rotation range. No further corrections were applied to the AP diffraction images.

Diffuse scattering maps were constructed as three-dimensional grids in reciprocal space whose nodes oversample Miller indices by a factor of three along each lattice direction. Corrected intensities were binned into cubic voxels centered on these nodes, and the mean pixel intensity of each voxel was used to estimate the intensity at each node. The signal-to-noise ratio was estimated as the mean divided by the standard deviation of the intensities binned into each voxel and is shown across resolution shells in Fig. S1A. Maps were symmetrized by averaging the intensities of Laue- and Friedel-equivalent voxels. The mean radial intensity was subtracted, followed by addition of a constant value to ensure that all intensities were positive. Code used to generate the maps is available at <https://github.com/apecck12/diffuse>.

B. Bragg data processing

Bragg data were indexed, integrated, and scaled with XDS [16]; statistics are shown in Table 1. For CypA, molecular replacement was performed with Phaser [23] using PDB 2CPL as a search model. This was followed by five macrocycles of refinement in Phenix [24] as previously described [13]. For AP, molecular replacement was performed with Phaser using wild-type AP (PDB 3TG0) stripped of non-protein atoms as the search model. As in Ref. [25], zinc ions at full occupancy and a tungstate ion and water molecules at partial occupancy were manually modeled into the residual electron density in each active site. Automated refinement was performed using CCP4 [26]. For WrpA, molecular replacement was performed with Phaser using PDB 5F51 as a search model [27]. This was followed by alternating rounds of manual refinement in Coot [28] to model a sulfate ion and water molecules and automated refinement in Phenix. The $R_{\text{work}}/R_{\text{free}}$ values of the final refined models were similar to those previously reported [13, 25, 27]. Diffraction images for the CypA, AP, and WrpA datasets are available at the SBGrid Data Bank, with accession numbers 68 [29], 456 [30], and 203 [31], respectively.

C. Diffuse scattering predictions from real-space models of disorder

Experimental diffuse scattering maps were compared to the following set of disorder models:

- Rigid body rotations, in which the atoms in an asymmetric unit rotate as a unit around a ran-

TABLE I. Data collection and map statistics

	CypA	AP	WrpA
<i>Data Collection and Model-free Analysis of Bragg Data</i>			
Space group	P2 ₁ 2 ₁ 2 ₁	P6 ₃ 22	P4 ₂ 22
Unit cell axes			
a, b, c (Å)	42.9 52.4 89.1	161.3, 161.3, 139.4	61.3 61.3 128.7
α, β, γ (°)	90.0 90.0 90.0	90.0, 90.0, 120.0	90.0 90.0 90.0
Wavelength (Å)	0.9795	0.9795	0.9787
Oscillation range (°)	0.5	0.15	1.0
Beam divergence S.D. (°)	0.0417	0.0285	0.0339
Mosaicity (°)	0.106	0.136	0.197
Wilson B	16.1	31.4	64.7
Multiplicity	5.8	6.3	12.7
Completeness (%)	93.6	94.0	99.7
$\langle I/\sigma I \rangle$	11.3	26.2	25.6
R _{merge}	0.113	0.045	0.061
<i>Refined model of Bragg data</i>			
Resolution range (Å)	44.6 - 1.2	50.0 - 2.0	44.4 - 2.5
No. of non-hydrogen atoms	1419	6731	1237
Average B-factor (Å ²)	21.4	46.2	77.0
R _{work} /R _{free}	0.169/0.177	0.192/0.231	0.235/0.277
<i>Diffuse scattering map</i>			
Resolution range (Å)	93.4 - 1.5	75.2 - 2.5	69.3 - 2.1
CC _{Friedel pairs} ^a	0.979 (0.623)	0.954 (0.695)	0.987 (0.683)
CC _{unsym., Friedel-sym.} ^{a,b}	0.996 (0.915)	0.994 (0.957)	0.997 (0.927)
CC _{unsym., Laue-sym.} ^{a,b}	0.993 (0.862)	0.984 (0.887)	0.993 (0.821)

^a For values in parenthesis, the average radial intensity was subtracted prior to applying symmetry operations.

^b Correlation coefficient between the unsymmetrized map and the map after applying the indicated symmetry operations.

dom, isotropically oriented axis, with a normally distributed rotation angle.

- Rigid body translations, in which the atoms in an asymmetric unit translate as a unit. Translations sample an isotropic Gaussian distribution.
- Liquid-like motions, where correlations between atoms decay exponentially as a function of interatomic distance. Two forms of this model were considered: one in which correlations were confined within the boundaries of the asymmetric unit, and a second model “with neighbors” in which correlations could extend between asymmetric units (and therefore cross unit cell boundaries).
- Gaussian elastic network model, a commonly used normal-mode decomposition of the protein motions based on the structure.
- Ensemble models, which model configurational disorder as a discrete set of states. Conformational states were inferred by analyzing the crystal electron density from the Bragg data.

For all models except for the liquid-like motions model with neighbors, correlations were assumed to be confined

within the boundaries of asymmetric units, with no coherence between neighboring protein molecules.

Diffuse scattering maps were simulated using the software package Thor [32]. For consistency with the experimental maps, the average radial intensity was subtracted from the predicted maps. Agreement was assessed by the linear correlation coefficient (CC), with each voxel downweighted by its multiplicity. For visual comparisons, a multiplicative scale factor and constant platform were applied to place the predicted maps on the same intensity scale as the experimental maps unless otherwise noted. Disorder models are described in more mathematical detail in the Supplementary Information.

III. RESULTS

1. The experimental maps exhibit Laue symmetry and significant anisotropic features.

We analyzed three crystallographic datasets collected by the rotation method for which diffuse scattering was visible in the raw diffraction images (Fig. 1, left). The Bragg data were separated and processed by standard protocols, yielding refined structural models similar to




FIG. 1. Reciprocal space maps from experimental diffuse scattering. The diffuse scattering in diffraction images collected for (A) CypA, (B) AP, and (C) WrpA was reconstructed into a reciprocal space map for each system. These maps take the form of three-dimensional grids that are 3x oversampled relative to the Miller indices along each lattice axis. (Left) An example diffraction image from each dataset. (Right) Central slices through reciprocal space are visualized for each unsymmetrized map in the top panels. The lower panels show these slices after symmetrization of Friedel- and Laue-equivalent voxels, followed by subtraction of the average radial intensity profile to highlight anisotropic features. In some cases color maps were thresholded at 0.5-0.7 of the maximum voxel intensity, which saturated a subset of voxels but improved overall contrast.

those previously published [13, 25, 27]. The diffuse scattering was isolated and processed to generate three-dimensional maps in reciprocal space (Fig. 1, upper right panels). The maps oversample the diffuse scattering signal along each lattice direction by a factor of three relative to the Miller indices, which enables these maps to resolve correlations that extend across multiple unit cells.

Overall statistics for the diffuse scattering maps are shown in Table I and by resolution shell in Fig. S1. The intensities of voxels related by Friedel’s law and Laue symmetry showed significant correlation in all cases, supporting symmetrization of the maps by averaging the intensities of these symmetry-equivalent voxels. To remove the intensity contributions from uncorrelated disorder,




FIG. 2. Elastic network models of Gaussian disorder. The highest and lowest magnitude entries in the covariance matrix are overlaid as blue and red cylinders, respectively, on the structures of (A) CypA, (B) AP, and (C) WrpA. The 0kl slices of the predicted diffuse scattering maps are shown on the right, with the overall CC noted in white. The color scales differ from Fig. 1 to enhance visualization of the features.

which includes solvent and air scattering in addition to uncorrelated protein disorder, the average radial intensity was subtracted from each map. The resulting maps were characterized by significant anisotropic features, indicative of correlated disorder (Fig. 1, lower right panels).

2. Models of complex protein dynamics do not correlate with the experimental maps.

Models of disorder that predict idiosyncratic configurational dynamics – the type of motions most likely related to biological function – were assessed for their ability to reproduce the experimental signal. A general class of these models assumes that interatomic displacements are small and sample a Gaussian distribution, and can thus be described by a covariance matrix. Here, covariance matrices were predicted from normal modes analysis of each protein structure in torsion angle space [33]. The predicted distributions of strongly covarying atom pairs were non-uniform and often spatially localized in the protein (Fig. 2, left and Fig. S3), distinguishing these models from those considered in the next section. For all three systems, the predicted diffuse scattering was unable to reproduce the observed signal, apparent both in the low CC and by visual comparison of the predicted and experimental 0kl planes (cf. Figs. 1 and 2).




FIG. 3. Ensemble models inferred from the Bragg data. (A) Multi-conformer modeling of the CypA electron density map predicts a minor population of alternate conformers (purple) that radiate from the active site. (B) CypA datasets collected at 180 K and below show two loop conformations between residues 79 and 83; the visualized loop conformations are those modeled in the 100 K dataset. Above 180 K, the conformation shown in white is not populated. (C) Model of AP assuming that one active site is tungstate-bound (left) while the other is occupied by water molecules that coordinate the active site metal ions (right). The 0kl plane of the predicted map from each ensemble model is shown on the right, with a different color scale from Fig. 1 to enhance visual contrast. The overall CC between the predicted and experimental maps is noted in white.

Non-Gaussian ensemble models inferred from the Bragg data were also evaluated. In the case of CypA, multiconformer modeling of the electron density map revealed a minor population of alternative rotamers for a series of residues that radiate from the active site (Fig. 3A, left) [34]. This observation of a correlated rotameric switch is consistent with prior analysis of CypA crystal structures [34–37]. Another ensemble was generated from the loop conformations populated by residues 79-83 in CypA crystals collected at or below 180 K (Fig. 3B, left) [37]. Although only one of these conformations is populated in the dataset analyzed here (which was collected at 273 K), this model offers a distinct example of a type of configurational disorder that is prevalent in proteins. A third ensemble model was suggested by the occupancy disorder observed in AP, for which the Bragg coordinates were refined with a half-occupied tungstate ion in each active site. However, the Bragg data cannot distinguish between this model of partial occupancy

TABLE II. Refined model parameters

	Model				
	Bragg ^a Wilson B	Rigid body translations	Liquid-like motions ^b		Rigid body rotations
	σ (Å)	σ (Å)	σ (Å)	γ (Å)	σ (°)
CypA	0.45	0.33	0.36 [0.39] ^c	18 [18] ^c	2.9
AP	0.63	0.40	0.40 [0.48]	118 [53]	0.9
WrpA	0.90	0.52	0.54 [0.61]	15 [18]	3.4

^a The Bragg σ was computed from the Wilson B factor.

^b Brackets denote the liquid-like motions with neighbors model.

^c Prior analysis of this dataset found best fit parameters of $\sigma = 0.38$ Å and $\gamma = 7.1$ Å for the model with neighbors [13].

and a model of correlated occupancy in which only one AP monomer can be tungstate-bound at a given time (Fig. 3C, left). Whereas partial occupancy contributes to radially symmetric diffuse scattering, correlated occupancy yields anisotropic features.

Diffuse scattering maps were predicted by Guinier’s equation [38]; the 0kl planes are shown in Fig. 3. The predicted maps from the CypA ensemble models are distinct from one another, but both exhibit features spread over much broader regions of reciprocal space (due to the short length scale of the disorder in real space) than observed experimentally. The map predicted by the correlated occupancy model shows a unique checkered pattern, but these regular features are similarly larger than the features observed in the experimental map for AP. None of these ensemble models show any correlation with the experimental signal (Table III).

3. Simple models of rotational and translational disorder show modest correlation with the experimental maps.

The inability of the elastic network and ensemble models to reproduce the experimental maps prompted us to evaluate other disorder models which predict simpler dynamics of rigid body or liquid-like motions. The models described in this section assume that correlations do not extend beyond the boundaries of individual asymmetric units. Additionally, these models share the symmetrized molecular transform as their basis, which, as discussed below, raises the possibility of using the diffuse scattering signal for static structural inference.

The diffuse scattering produced by rigid body translational disorder is the molecular transform scaled by the Debye-Waller factor [3, 14]. For all three maps, this disorder model showed nontrivial correlation with the experimental maps. Further, the best fit values of the isotropic displacement parameter σ , which reports on the scale of displacement, were within two-fold of the Bragg Wilson B factor (Fig. 4, Table II), suggesting that the diffuse signal could be consistent scattering missing in the Bragg data due to disorder. The fit was modestly improved by imposing exponential decay on interatomic covariances, thereby switching from a rigid body to a liquid-like description of correlated dynamics (Fig. 4). This liquid-like

motions model predicted similar values for the isotropic displacement parameter, but in the case of CypA and WrpA, a correlation length roughly one third to one half the dimensions of the protein molecule (Table II). In the case of AP, the best fit correlation length spanned the longest dimension of the protein, consistent with the lack of improvement in the CC: in the regime of correlation lengths longer than the protein unit, the diffuse scattering predictions of the liquid-like motions and rigid body translational disorder models converge.

A model of rigid body rotational disorder, which blurs features of the molecular transform in concentric shells of \mathbf{q} , also showed modest correlation with the CypA and WrpA maps (Fig. 4). For both maps, the best fit values for the standard deviation of the angle of rotation were on the order of 2-3°, consistent with a blurring effect that spans a few voxels of these reciprocal space maps. The best fit value for the AP map was smaller (0.9°), yielding minimal radial blurring that could be resolved by the coarseness of the map’s voxels, which along with the modest correlation suggested that rotational disorder was inconsistent with the observed signal. Relative to CypA and WrpA, AP has more crystal contacts that may inhibit this type of disorder. It is also possible that the finer slicing during collection of the AP data minimized blurring, but radial blurring due to data collection versus as a result of rotational disorder in the crystal cannot be distinguished by the isotropic model considered here.

4. Speckles indicate long-range correlated disorder that crosses unit cell boundaries.

Of the models considered above, in no case does the correlation coefficient between the predicted and experimental map exceed 0.5 (Table III, Fig. S4). Visual inspection suggests that a feature these models systematically fail to reproduce is the observed “speckles,” periodic spikes in intensity that appear superimposed on diffuse scattering features that span larger volumes in reciprocal space (Fig. 5A-B, insets). Such speckles have previously been noted in studies that analyzed the diffuse scattering signal at fractional Miller indices [22, 39, 40]. Because the length scale of disorder in real space determines the spacing of diffuse features in reciprocal space, the need to oversample the diffuse signal relative to integral Miller indices to observe these speckles indicates that they arise from correlations that extend beyond the boundaries of a single unit cell. The models examined in prior sections assumed correlated disorder confined within asymmetric units, so were unable to generate this type of signal (cf. the size of diffuse features in Figs. 3 and 5, for example).

We therefore considered two models of disorder in which correlations extend across unit cell boundaries. The first of these models is a long-range variation of the liquid-like motions model assessed above, in which the basis is the crystal transform rather than the asymmetric unit-derived molecular transform. For each system, this




FIG. 4. **Comparison of simple isotropic models of disorder.** For the indicated model, parameters that tune the disorder were fit either by a grid scan or least-squares optimization to maximize overall correlation with the experimental map. The predicted 0kl planes for the best fit maps are shown, with the experimental 0kl planes displayed in the left column for comparison. Overall correlation coefficients between the experimental and predicted maps are noted in white.

model of liquid-like motions with neighbors showed considerable agreement with the experimental signal, qualitatively reproducing the speckled features and quantitatively yielding the highest CC of the models considered here (Fig. 4, Table III). The refined model parameters for the two liquid-like motions models were similar in most cases (Table II), indicating that the consistent increase in CC between the short- and long-range models resulted almost exclusively from taking into account correlations across neighboring asymmetric units (and thus unit cell boundaries).

An alternative model proposed to account for speckled features invokes acoustic lattice vibrations produced by phonon-induced inelastic scattering [22, 39, 40]. One prediction of this model is that the diffuse intensity will decrease proportional to the square distance from the center of the Brillouin zone, $\delta\mathbf{q}$. Such a trend is absent from the molecular transform and its derivative models (Fig. 5C), in which disorder is confined within the boundaries of asymmetric units. Though qualitatively the phonon model accounts for the observed halos around Bragg peaks, quantitatively the experimental fall-off in intensity differs from the $1/\delta\mathbf{q}^2$ dependence predicted for single phonon interactions (Fig. 5D-F, dashed red) [39, 40]. More complex phonon models, either from extending the spectrum to include optical modes or accounting multiple-phonon effects, are predicted to cause intensity to vary more slowly with distance from the center of the Brillouin zone [39]. However, there is currently

TABLE III. Correlation coefficients between predicted and experimental maps

	Model				Ensemble
	Rigid body rotations	Rigid body translations	Liquid-like ^a motions	Elastic network	
CypA	0.46	0.41	0.48 [0.71]	0.17	-0.02 ^b , -0.04 ^c
AP	0.26	0.31	0.32 [0.66]	0.09	—
WrpA	0.44	0.36	0.36 [0.67]	0.20	0.00 ^d

^a Brackets denote the liquid-like motions with neighbors model.

^b Rotamer switch model.

^c Disordered loop model.

^d Correlated occupancy model.

no robust method for predicting the diffuse scattering produced by phonons in macromolecular crystals. In the absence of such a method and an established procedure for generating competing acoustic modes, let alone optical modes or multiple phonon interactions, we cannot adequately assess agreement with the phonon model versus other types of long-range disorder.

IV. DISCUSSION

Here we present a unified framework of the principal disorder models that have previously been used to interpret diffuse scattering, and compare their ability to reproduce the signal observed in three experimental datasets.




FIG. 5. **Speckled features of the experimental maps.** Insets of the $0kl$ planes of the (A) CypA and (B) AP maps highlight the characteristic speckles. For (C-F), maps were reconstructed or computed with 5x oversampling along each lattice direction relative to integral Miller indices. Mean voxel intensities were normalized such that the mean value for voxels that coincided with the center of the Brillouin zone was unity, and plotted for twenty resolution shells as a function of distance from the center of the Brillouin zone, δq . The dashed line indicates the best fit of the data points in the lowest resolution shell (where the fall-off in intensity was generally most pronounced) to a curve with $1/\delta q^2$ dependence. For comparison to models whose basis is the molecular transform, analysis for the CypA molecular transform map is shown in (C).

Consistent with prior work, the above analysis finds that rigid body and liquid-like motions models exhibit modest correlation with the experimental maps [5, 13, 14, 41]. Models that predict more complex dynamics were also considered, but showed minimal agreement with experiment (Table III). Experimentally-observed speckles did not fit the profile for phonon-induced lattice dynamics, but could at least in part be reproduced by a model of liquid-like motions in which disorder extended across neighboring asymmetric units and unit cell boundaries. However, none of the models assessed here could fully explain the experimental signal.

Past interest in diffuse scattering has largely stemmed from the premise that these data probe dynamics related to biological function [1, 5–13]. However, the experimental maps showed minimal correlation with the biologically-informed models assessed here, including ensemble models supported by the Bragg electron density. Further, the observation of features indicative of corre-

lations that span neighboring molecules in the crystal cautions against the assumption that the dominant signal originates from the same protein motions that occur under physiological conditions. Models of disorder that account for both intermolecular and intramolecular correlations will thus be needed to resolve the contributions of each to the observed signal, a prerequisite in determining whether diffuse scattering is a useful method for studying dynamics associated with biological function.

On the other hand, the ability of rigid body and liquid-like motions models to reproduce many experimental features suggests that diffuse scattering data could in some cases be useful for resolution extension or phase retrieval. These models share the molecular or crystal transform as their basis, and thus yield a scaled or blurred image of this transform in the diffuse scattering map. The general observation that the diffuse scattering does not *directly* reflect the molecular or crystal transform, but at the very least the convolution of the transform with some blurring

function, must be better understood and taken into account.

Extracting the molecular transform signal will be particularly challenging for maps that exhibit enhanced scattering at the center of the Brillouin zone, as observed here. The model of liquid-like motions with neighbors best accounted for this feature, but in the linear approximation, it is a convolution of the disorder-free diffraction with a kernel that is the Fourier representation of the disorder. Thus, this model reports on the crystal transform, which is non-zero only at integral Miller indices. It does not contain information about the value of the molecular transform at fractional Miller indices. Such information could in principle be employed for resolution extension, but not phase retrieval. Iterative phase retrieval algorithms, such as those recently used by Ayer *et. al.* [14], require non-integral oversampled measurements to uniquely determine unmeasured phases [42]. The liquid-like motions model we present is approximate, and a more rigorous treatment of crystalline disorder may enable measurement of an oversampled molecular transform from the diffuse scattering. Despite this possibility, our results call into question the practice of directly using a diffuse scattering map for either resolution extension or iterative phasing in cases where enhanced scattering at the center of the Brillouin zone is observed – a universal feature in the three model systems we studied. Precisely how to deconvolve useful signals from such maps remains an open area of investigation.

The above analysis highlights a need for new models of diffuse scattering, either to interpret biologically relevant disorder or improve structure solutions. The contribution of different types of disorder to this signal is not trivially additive, so challenges lie ahead both in jointly modeling distinct sources of disorder and in deconvolving weak signals from the dominant features. The search space for such models is intractably large, so the number of acceptable free parameters and constraints will require careful treatment. Ideally, it would be possible to assert a model sophisticated enough to report interesting and idiosyncratic disorder in different systems (such as functional motions), but simple enough (*i.e.* with few independent parameters) to infer directly from the observed data.

An alternative route is detailed forward modeling, such as molecular dynamics, which has previously been used to analyze diffuse scattering [6, 12, 43]. Molecular dynamics concurrently simulates multiple types of disorder, but this method does not lend itself to refining the contri-

butions of different kinds of disorder to fit experimental data. In the common case where such simulations do not satisfactorily reproduce experimental observations, it is challenging to modify them in a principled manner so that they do. Combined with the computational expense of these methods, it seems prudent to seek simple explanations and models for analyzing diffuse scattering before comparing to atomic simulation. The incisive test of any model will come from its predictive power: confirming that a specific physical perturbation of a crystal system results in the predicted change to the diffuse signal.

Here, we sought a phenomenological model of the physics underpinning diffuse scattering observed from protein crystals. Although such models are oversimplifications of what we know to be true of protein motions and anticipate for lattice dynamics, comparison of these models critically addresses the nature and length scale of the disorder involved. Resolving aspects of this correlated disorder by successful modeling is, in turn, necessary for interpreting the diffuse signal either to model functional protein motions or enhance static structure inference. We found that previously considered models did not fully explain experimental observations. One particular model, of liquid-like motions with neighbors, achieved high correlations with the three datasets analyzed (CCs ~ 0.7), but unexplained features in the diffuse signal remain.

ACKNOWLEDGMENTS

We thank Vijay Pande, Soichi Wakatsuki, and Peter Moore for instructive discussions, and Dan Herschlag for the alkaline phosphatase dataset and comments on the manuscript. Michael Levitt is gratefully acknowledged for providing his Normal Mode Analysis code. We also thank the Stanford Research Computing Center for providing the Sherlock cluster and other computational resources that facilitated this work. A.P. was funded by a National Science Foundation Graduate Research Fellowship and supported by Vijay S. Pande. F.P. acknowledges support from the National Institutes of Health (NIH), grant No. 1-R01-GM115749-01 (PI: Michael Levitt). T.L. was supported by the LCLS directorate, SLAC National Accelerator Laboratory, via the U.S. Department of Energy Office of Basic Energy Sciences Contract No. DE-AC02-76SF00515.

-
- [1] M. E. Wall, P. D. Adams, J. S. Fraser, and N. K. Sauter, *Structure* **22**, 182 (2014).
 - [2] T. Welberry and T. Weber, *Crystallography Reviews* **22**, 2 (2016).
 - [3] P. B. Moore, *Structure* **17**, 1307 (2009).
 - [4] J. P. Benoit and J. Doucet, *Q. Rev. Biophys.* **28**, 131 (1995).
 - [5] M. E. Wall, J. B. Clarage, and G. N. Phillips, *Structure* **5**, 1599 (1997).
 - [6] S. Hery, D. Genest, and J. C. Smith, *J. Mol. Biol.* **279**, 303 (1998).
 - [7] A. R. Kolatkar, J. B. Clarage, and G. N. Phillips, *Acta*

- Crystallogr. D Biol. Crystallogr. **50**, 210 (1994).
- [8] L. Meinhold and J. C. Smith, Biophys. J. **88**, 2554 (2005).
- [9] S. Chacko and G. N. Phillips, Biophys. J. **61**, 1256 (1992).
- [10] L. Meinhold and J. C. Smith, Phys. Rev. Lett. **95**, 218103 (2005).
- [11] L. Meinhold and J. C. Smith, Proteins **66**, 941 (2007).
- [12] M. E. Wall, A. H. Van Benschoten, N. K. Sauter, P. D. Adams, J. S. Fraser, and T. C. Terwilliger, Proc. Natl. Acad. Sci. U.S.A. **111**, 17887 (2014).
- [13] A. H. Van Benschoten, L. Liu, A. Gonzalez, A. S. Brewster, N. K. Sauter, J. S. Fraser, and M. E. Wall, Proc. Natl. Acad. Sci. U.S.A. **113**, 4069 (2016).
- [14] K. Ayyer, O. M. Yefanov, D. Oberthur, S. Roy-Chowdhury, L. Galli, V. Mariani, S. Basu, J. Coe, C. E. Conrad, R. Fromme, A. Schaffer, K. Dörner, D. James, C. Kupitz, M. Metz, G. Nelson, P. L. Xavier, K. R. Beylerlein, M. Schmidt, I. Sarrou, J. C. Spence, U. Weierstall, T. A. White, J. H. Yang, Y. Zhao, M. Liang, A. Aquila, M. S. Hunter, J. S. Robinson, J. E. Koglin, S. Boutet, P. Fromme, A. Barty, and H. N. Chapman, Nature **530**, 202 (2016).
- [15] S. P. Meisburger, W. C. Thomas, M. B. Watkins, and N. Ando, Chem. Rev. (2017).
- [16] W. Kabsch, Acta Crystallogr. D Biol. Crystallogr. **66**, 125 (2010).
- [17] G. Hura, J. M. Sorenson, R. M. Glaeser, and T. Head-Gordon, The Journal of Chemical Physics **113**, 9140 (2000).
- [18] M. E. Wall, Methods Mol. Biol. **544**, 269 (2009).
- [19] D. G. Waterman, G. Winter, R. J. Gildea, J. M. Parkhurst, A. S. Brewster, N. K. Sauter, and G. Evans, Acta Crystallogr D Struct Biol **72**, 558 (2016).
- [20] H. Boysen and W. Adlhart, Journal of Applied Crystallography **20**, 200 (1987).
- [21] W. Kabsch, Acta Crystallogr. D Biol. Crystallogr. **66**, 133 (2010).
- [22] Y. S. Polikanov and P. B. Moore, Acta Crystallogr. D Biol. Crystallogr. **71**, 2021 (2015).
- [23] A. J. McCoy, R. W. Grosse-Kunstleve, P. D. Adams, M. D. Winn, L. C. Storoni, and R. J. Read, J Appl Crystallogr **40**, 658 (2007).
- [24] P. D. Adams, P. V. Afonine, G. Bunkoczi, V. B. Chen, I. W. Davis, N. Echols, J. J. Headd, L. W. Hung, G. J. Kapral, R. W. Grosse-Kunstleve, A. J. McCoy, N. W. Moriarty, R. Oeffner, R. J. Read, D. C. Richardson, J. S. Richardson, T. C. Terwilliger, and P. H. Zwart, Acta Crystallogr. D Biol. Crystallogr. **66**, 213 (2010).
- [25] A. Peck, F. Sunden, L. D. Andrews, V. S. Pande, and D. Herschlag, J. Mol. Biol. **428**, 2758 (2016).
- [26] G. N. Murshudov, P. Skubak, A. A. Lebedev, N. S. Pannu, R. A. Steiner, R. A. Nicholls, M. D. Winn, F. Long, and A. A. Vagin, Acta Crystallogr. D Biol. Crystallogr. **67**, 355 (2011).
- [27] J. Herrou, D. M. Czyz, J. W. Willett, H. S. Kim, G. Chhor, G. Babnigg, Y. Kim, and S. Crosson, J. Bacteriol. **198**, 1281 (2016).
- [28] P. Emsley, B. Lohkamp, W. G. Scott, and K. Cowtan, Acta Crystallogr. D Biol. Crystallogr. **66**, 486 (2010).
- [29] J. Fraser, “X-ray diffraction data for: Cyclophilin a. pdb code 4yuo,” (2015).
- [30] A. Peck, S. Ressler, and D. Herschlag, “X-ray diffraction data for: E. coli alkaline phosphatase in complex with tungstate. pdb code 5c66,” (2017).
- [31] J. Herrou and S. Crosson, “X-ray diffraction data for: Structure of b. abortus wrba-related protein a (apowrpa). pdb code 5f51,” (2015).
- [32] T. J. Lane, “Thor,” <https://github.com/tjlane/thor> (2017).
- [33] J. K. Bray, D. R. Weiss, and M. Levitt, Biophys. J. **101**, 2966 (2011).
- [34] H. van den Bedem, A. Dhanik, J. C. Latombe, and A. M. Deacon, Acta Crystallogr. D Biol. Crystallogr. **65**, 1107 (2009).
- [35] J. S. Fraser, M. W. Clarkson, S. C. Degnan, R. Erion, D. Kern, and T. Alber, Nature **462**, 669 (2009).
- [36] J. S. Fraser, H. van den Bedem, A. J. Samelson, P. T. Lang, J. M. Holton, N. Echols, and T. Alber, Proc. Natl. Acad. Sci. U.S.A. **108**, 16247 (2011).
- [37] D. A. Keedy, L. R. Kenner, M. Warkentin, R. A. Woldeyes, J. B. Hopkins, M. C. Thompson, A. S. Brewster, A. H. Van Benschoten, E. L. Baxter, M. Uervirojnangkoorn, S. E. McPhillips, J. Song, R. Alonso-Mori, J. M. Holton, W. I. Weis, A. T. Brunger, S. M. Soltis, H. Lemke, A. Gonzalez, N. K. Sauter, A. E. Cohen, H. van den Bedem, R. E. Thorne, and J. S. Fraser, Elife **4** (2015).
- [38] A. Guinier, *X-ray diffraction in crystals, imperfect crystals, and amorphous bodies* (W. H. Freeman, San Francisco, 1963).
- [39] I. D. Glover, G. W. Harris, J. R. Helliwell, and D. S. Moss, Acta Crystallographica Section B **47**, 960 (1991).
- [40] L. Meinhold, F. Merzel, and J. C. Smith, Phys. Rev. Lett. **99**, 138101 (2007).
- [41] J. Doucet and J. P. Benoit, Nature **325**, 643 (1987).
- [42] D. Sayre, Acta Crystallographica **5**, 843 (1952).
- [43] P. Faure, A. Micu, D. Perahia, J. Doucet, J. C. Smith, and J. P. Benoit, Nat. Struct. Biol. **1**, 124 (1994)
-

Supplemental Materials

I. EXTENDED METHODS

The disorder models assessed in this study are described in more detail below. With the exception of the liquid-like motions with neighbors model, these models assume that correlations are confined within the boundaries of the asymmetric unit. Because of this assumption, the models of rigid body disorder and liquid-like motions without neighbors share the symmetrized molecular transform, I_m , as their basis:

$$I_m = \sum_{asu} |F_{asu}(\mathbf{q})|^2 \quad (\text{S1})$$

where the summation is across all asymmetric units in the unit cell. Thus, this subset of models predicts that the diffuse scattering is a blurred image of the symmetrized molecular transform, with the form of the blurring dependent on the nature of the disorder. This incoherent sum of the asymmetric unit intensities is distinct from the coherent sum of scattered intensities across the crystal or, equivalently, the crystal transform, I_c :

$$I_c = |S(\mathbf{q})|^2 \left| \sum_{asu} F_{asu}(\mathbf{q}) \right|^2 \quad (\text{S2})$$

where the Dirac comb, $|S(\mathbf{q})|^2$, is 1 at integral Miller indices and 0 at all other \mathbf{q} (omitting scaling by the number of unit cells). This is in contrast to the molecular transform, which is characterized by positive, non-zero intensities throughout reciprocal space. This distinction is important for the two variants of liquid-like motions models considered in this work, as noted below.

1. Rigid body rotational disorder

The simplest case of rigid body rotational disorder was evaluated, in which there is no preferred axis of rotation and rotation angles are sampled from a normal distribution. Diffuse scattering maps were predicted from an ensemble of rotated molecules using Guinier's equation [S1]:

$$I_{\text{diffuse}}(\mathbf{q}) \propto \sum_{asu} [\langle |F_{asu,n}(\mathbf{q})|^2 \rangle - \langle |F_{asu,n}(\mathbf{q})| \rangle^2]. \quad (\text{S3})$$

where $F_{asu,n}(\mathbf{q})$ represents the asymmetric unit transform for the n th ensemble member and $\langle \dots \rangle$ indicates the time or ensemble average (which these data cannot distinguish between). In order to focus on the diffuse scattering predicted solely by rotational disorder, the asymmetric unit transforms were not scaled by Debye-Waller factors, which account for translational disorder effects. For each map, the best fit standard deviation of the rotational distribution was determined by scanning over values of this parameter to maximize the correlation coefficient (CC) between the experimental and predicted maps (Fig. S2A).

2. Rigid body translational disorder

The model of rigid body translational disorder assumes that all ordered atoms in the asymmetric unit are displaced as a rigid structural unit by translations that sample an isotropic Gaussian distribution. The displacement covariance between all atom pairs is identical and can be described by a scalar, σ^2 . The expression for the diffuse scattering intensity is:

$$I_{\text{diffuse}}(\mathbf{q}) = \left[1 - e^{-q^2 \sigma^2} \right] I_m \quad (\text{S4})$$

where N is the number of unit cells and I_m is the symmetrized molecular transform (Eq. S1). This expression has previously been derived in Refs. [S2] and [S3]. For each system, the molecular transform was computed from the refined Bragg coordinates, excluding solvent, hydrogen, and crystallographically-unresolved atoms (which were assumed to exhibit uncorrelated disordered behavior and thus contribute to radially symmetric rather than anisotropic diffuse scattering). Best fit values of σ were determined by linear least-squares optimization to minimize the residual between the experimental and predicted maps.

3. Liquid-like motions model

The liquid-like motions model is a special case of Gaussian translational disorder in which correlated motions between atoms decay exponentially with their interatomic distance [S4, S5]. The displacement covariance between atoms is: $V_{ij} = \sigma^2 \Gamma(\mathbf{q}_{ij})$. As with the model of rigid body translational disorder, σ is the global isotropic displacement parameter, while the kernel $\Gamma(\mathbf{q}_{ij})$ is a Fourier representation of interatomic correlations with the following functional form:

$$\Gamma(\mathbf{q}) = \frac{8\pi\gamma^3}{(1 + q^2\sigma^2)^2} \quad (\text{S5})$$

where γ is the correlation length. The diffuse scattering intensity can then be computed as the convolution between the scattering intensity function and the Fourier transform of the correlation kernel:

$$I_{\text{diffuse}}(\mathbf{q}) = q^2 \sigma^2 e^{-q^2 \sigma^2} [I_0 * \Gamma(\mathbf{q})]. \quad (\text{S6})$$

Two variations of the liquid-like motions model were considered:

1. An internally-disordered model, in which the scattering intensity is the symmetrized molecular transform ($I_0 = I_m$). This confines interatomic correlations to be within the boundaries of asymmetric units.
2. A model ‘‘with neighbors,’’ in which the scattering intensity is the crystal transform ($I_0 = I_c$). This allows for correlated disorder to extend between neighboring asymmetric units and across unit cell boundaries.

Prior studies compared the model with neighbors to experimental maps that sampled the observed diffuse scattering intensity at integral Miller indices [S4, S5]. In this study, we compare the predicted signal to experimental maps that sample the diffuse signal at fractional Miller indices, which enables an evaluation of the ability of this model to reproduce observed changes in intensity as a function of distance from the center of the Brillouin zone.

For both liquid-like motions models, best fit values of the isotropic displacement parameter and correlation length were determined by a grid search to maximize the CC between the experimental and predicted maps. As an example, convergence of these parameters for the internally-disordered model is shown in Fig. S2B.

4. Elastic network model

This model assumes that interatomic displacements are small and sample a multivariate normal distribution, with zero mean and covariance $V_{ij} = \langle \delta_i^T \delta_j \rangle$, where δ_i is the instantaneous displacement vector of atom i from its mean position. In the case that correlations do not extend between unit cells, a general form of the predicted diffuse scattering is:

$$I_{\text{diffuse}}(\mathbf{q}) = N \sum_{i,j} f_i f_j e^{-i\mathbf{q} \cdot \mathbf{r}_{ij}} e^{-\frac{1}{2} \mathbf{q}^T V_{ii} \mathbf{q} - \frac{1}{2} \mathbf{q}^T V_{jj} \mathbf{q}} \times \left[e^{\mathbf{q}^T V_{ij} \mathbf{q}} - 1 \right] \quad (\text{S7})$$

where N is the number of unit cells, f_i and V_{ii} are respectively the atomic form factor and anisotropic displacement parameter for atom i , and \mathbf{r}_{ij} is the difference vector between the mean positions of atoms i and j [S2]. When all entries of this covariance matrix are identical, this model predicts rigid body translational disorder, and the expression for diffuse scattering simplifies to Eq. S4 [S2].

Here, we further restricted correlations to be confined within the boundaries of the asymmetric unit, an assumption that renders this model more biologically interpretable. For each system, the covariance matrix was determined from an elastic network model of the ordered atoms in the asymmetric unit. Specifically, the first ten normal modes of each protein system in torsion angle space were summed to generate C_{ij} , the isotropic correlation coefficient between the displacements of asymmetric unit atoms i and j [S6]. Entries in this correlation matrix were converted to covariances using the following formula:

$$V_{ij} = C_{ij} \sqrt{\langle \delta_i^2 \rangle \langle \delta_j^2 \rangle} \quad (\text{S8})$$

where the mean-square atomic displacements, $\langle \delta_i^2 \rangle$, are related to the isotropic B factors by: $B_i = 8\pi^2 \langle \delta_i^2 \rangle$. Thus, the amplitudes of motions described by the covariance matrix are consistent with the refined Bragg models. Diffuse scattering maps were predicted from these covariance matrices using Eq. S7.

5. Ensemble models

Many types of protein disorder involve transitions between discrete states rather than along a continuum of alternate conformations. In real space, this disorder can be modeled as an ensemble of representative “snapshots” of distinct protein configurations. The refined Bragg coordinates of the CypA and AP crystal structures suggested the existence of specific ensembles for these systems, and the diffuse scattering map predicted by each ensemble was computed by Guinier’s equation (Eq. S3) [S1]. The resulting map was scaled by the Debye-Waller factor, $e^{-q^2\sigma^2}$, with the global atomic displacement factor σ computed from the Bragg Wilson B factor. This scaling has the effect of simulating uncorrelated Gaussian translational disorder within each member of the ensemble.

II. EXTENDED FIGURES




FIG. S1. **Diffuse scattering map statistics by resolution shell.** (A) The median voxel intensity, signal to noise ratio, and number of pixels per voxel are shown across resolution bins for the indicated experimental map. The signal to noise ratio for each voxel was estimated as $\langle I \rangle / \sigma(I)$ for the set of pixel intensities binned into each voxel. The solid and dashed lines correspond to the overall values and the values for voxels centered on integral Miller indices, respectively. (B) Correlation coefficients between Friedel pairs (left), the indicated map before and after averaging Friedel pairs (center), and the indicated map before and after averaging Laue-symmetric voxels (right).




FIG. S2. **Convergence of disorder parameters for the rigid body rotational disorder and liquid-like motions models.** (A) Scans over the standard deviation of the rotational distribution, σ , were performed to fit the rigid body rotational disorder model to each experimental map. (B) Grid scans over the global atomic displacement factor, σ , and the correlation length, γ , parameters of the liquid-like motions model. The color indicates the overall correlation coefficient between the experimental and predicted maps.




FIG. S3. **Covariance matrices generated from elastic network models.** The correlation matrix of interatomic Gaussian displacements was predicted for each system by an elastic network model. The covariance matrix was then computed from this correlation matrix and normalized by the refined B factors to ensure consistency with the Bragg data.




FIG. S4. **Correlation coefficients by resolution shell.** The multiplicity-weighted correlation coefficient between the experimental and predicted map for the indicated models is plotted as a function of resolution shell.

-
- [S1] A. Guinier, *X-ray diffraction in crystals, imperfect crystals, and amorphous bodies* (W. H. Freeman, San Francisco, 1963).
[S2] P. B. Moore, *Structure* **17**, 1307 (2009)
[S3] K. Ayyer, O. M. Yefanov, D. Oberthur, S. RoyChowdhury, L. Galli, V. Mariani, S. Basu, J. Coe, C. E. Conrad, R. Fromme, A. Schaffer, K. Dorner, D. James, C. Kupitz, M. Metz, G. Nelson, P. L. Xavier, K. R. Beyerlein, M. Schmidt, I. Sarrou, J. C. Spence, U. Weierstall, T. A. White, J. H. Yang, Y. Zhao, M. Liang, A. Aquila, M. S. Hunter, J. S. Robinson, J. E. Koglin, S. Boutet, P. Fromme, A. Barty, and H. N. Chapman, *Nature* **530**, 202 (2016).
[S4] M. E. Wall, J. B. Clarage, and G. N. Phillips, *Structure* **5**, 1599 (1997).
[S5] J. B. Clarage, M. S. Clarage, W. C. Phillips, R. M. Sweet, and D. L. Caspar, *Proteins* **12**, 145 (1992).
[S6] J. K. Bray, D. R. Weiss, and M. Levitt, *Biophys. J.* **101**, 2966 (2011).