

Regular Potential Games

Brian Swenson^a, Ryan Murray^b, Soumya Kar^c

^a*Department of Electrical Engineering, Princeton University, Princeton, NJ
(bswenson@princeton.edu)*

^b*Department of Mathematics, North Carolina State University, Raleigh, NC
(rwmurray@ncsu.edu)*

^c*Department of Electrical and Computer Engineering, Carnegie Mellon University,
Pittsburgh, PA (soumyyak@andrew.cmu.edu)*

Abstract

A fundamental problem with the Nash equilibrium concept is the existence of certain “structurally deficient” equilibria that (i) lack fundamental robustness properties, and (ii) are difficult to analyze. The notion of a “regular” Nash equilibrium was introduced by Harsanyi. Such equilibria are isolated, highly robust, and relatively simple to analyze. A game is said to be regular if all equilibria in the game are regular. In this paper it is shown that almost all potential games are regular. That is, except for a closed subset with Lebesgue measure zero, all potential games are regular. As an immediate consequence of this, the paper also proves an oddness result for potential games: in almost all potential games, the number of Nash equilibrium strategies is finite and odd. Specialized results are given for weighted potential games, exact potential games, and games with identical payoffs. Applications of the results to game-theoretic learning are discussed.

Keywords: Game theory; Potential games; Generic games; Regular equilibria; Multi-agent systems

1. Introduction

While the notion of Nash equilibrium (NE) is a universally accepted solution concept for games, several shortcomings have been noted over the years. A principal criticism (in addition to non-uniqueness) is that some Nash equilibrium strategies may be undesirable or unreasonable due to a lack of basic robustness properties. As a consequence, many equilibrium refinement con-

cepts have been proposed [1, 2, 3, 4, 5, 6, 7], each attempting to single out subsets of Nash equilibrium strategies that satisfy some desirable criteria.

One of the most stringent refinement concepts, originally proposed by Harsanyi [3], is that a NE strategy be “regular.” In the words of van Damme [4], “regular Nash equilibria possess all the robustness properties that one can reasonably expect equilibria to possess.” Such equilibria are quasi-strict [3, 4], perfect [1], proper [2], strongly stable [6], essential [5], and isolated [4].¹

If all equilibria of a game are regular, then the number of NE strategies in the game has been shown to be finite and, curiously, odd [3, 8]. Regular equilibria have also been studied in the context of games of incomplete information, where, as part of Harsanyi’s celebrated purification theorem [9, 10, 11], they have been shown to be approachable.

A game is said to be regular if all equilibria in the game are regular. Harsanyi [3] showed that almost all² games are regular, and hence, in almost all games, all equilibria possess all the robustness properties we might reasonably hope for.

While this result is a powerful when targeted at general N -player games, there are many important classes of games that have Lebesgue measure zero within the space of all games [12]. Harsanyi’s result tells us nothing about equilibrium properties within such special classes of games. This is the case, for example, in the important class of multi-agent games known as *potential games* [13].

A game is said to be a *potential game* if there exists some underlying function (generally referred to as the *potential function*) that all players implicitly seek to optimize. Potential games have many applications in economics and engineering [14, 13], and are particularly useful in the study of multi-agent systems, e.g., [15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25].

There are several types of potential games—in order of decreasing generality, these include weighted potential games, exact potential games, and games with identical payoffs [13, 26]. Letting WPG, EPG, and GIP denote the set of each of these types of potential games respectively, and letting G

¹See [4] for an in-depth discussion of each of these concepts and their interrelationships.

²Following Harsanyi [3], when we say almost all games satisfy some condition we mean the set of games where the condition fails to hold is a closed set with Lebesgue measure zero. See Section 2.3 for more details.

denote the set of all games, we have the following relationship:³

$$\text{GIP} \subset \text{EPG} \subset \text{WPG} \subset \text{G},$$

where each subset is a low-dimensional (measure-zero) subset within any of its supersets. Harsanyi’s regularity result provides no information on the abundance (or dearth) of regular equilibria within these subclasses of games. Hence, when restricting attention to potential games, as is often done in the study of multi-agent systems, we are deprived of any generic results on the regularity, robustness, or finiteness of the equilibrium set.

We say that a property holds for almost all games in a given class if the subset of games in the class where the property fails to hold is a closed set with Lebesgue measure zero (with the dimension of the Lebesgue measure corresponding to the dimension of the given class of games—see Section 2.3 for more details).

The main result of this paper is the following theorem.

Theorem 1.

- (i) *Almost all weighted potential games are regular.*
- (ii) *Almost all exact potential games are regular.*
- (iii) *Almost all games with identical payoffs are regular.*

We note that this result implies that for almost all games in each of these classes, all equilibria are quasi-strict, perfect, proper, strongly stable, essential, and isolated. Using Harsanyi’s oddness theorem (see [3], Theorem 1), we see that in any regular game, the number of NE strategies is finite and odd. Hence, the following result is an immediate consequence of Theorem 1.

Theorem 2. *In almost all weighted potential games, almost all exact potential games, and almost all games with identical payoffs, the number of NE strategies is finite and odd.*

Regularity may be seen as serving two purposes. First, it ensures that the equilibrium set possesses the desirable structural properties noted above

³More precisely, any finite game of a fixed size (i.e., with a fixed number of players and actions) is uniquely represented as a vector in Euclidean space denoting the payoff received by each player for each pure strategy. The set G of all possible games of a given size is equal to \mathbb{R}^m for some appropriate $m \in \mathbb{N}$ (see, e.g., [28] Section 12.1), and each class of potential games is a lower-dimensional subset of \mathbb{R}^m . See Section 2.3 for more details.

(e.g., equilibria are isolated, robust, and finite in number). Second, it simplifies the analysis of the game near equilibrium points—the important features of players’ utility functions near an equilibrium can be understood by looking only at first- and second-order terms in the associated Taylor series expansion. In this sense, the role of regular equilibria in games is analogous to the role that non-degenerate critical points play in the study of real-valued functions.⁴ This amenable analytic structure can greatly facilitate the study of (for example) game-theoretic learning processes [29, 30] or approachability in games with incomplete information [10].

As an application of these results to learning theory, in the paper [29] we consider the problem of studying continuous best-response dynamics (BR dynamics) [31, 32, 33] in potential games. BR dynamics are fundamental to learning theory—they model various forms of learning in games and underlie many popular game-theoretic learning algorithms including the canonical fictitious play (FP) algorithm [34]. While it is known that BR dynamics converge to the set of NE in potential games, the result is less than satisfactory. BR dynamics can converge to mixed-strategy (saddle-point) Nash equilibria and solutions of BR dynamics may be non-unique. Furthermore, little is understood about transient properties such as the rate of convergence of BR dynamics in potential games. (In fact, due to the non-uniqueness of solutions in potential games, it has been shown that it is impossible to establish convergence rate estimates for BR dynamics that hold at all points [35].)

In [29] we study how regular potential games can be used to address these issues. In particular, it is shown that in any regular potential game, BR dynamics converge generically to pure NE, solutions of BR dynamics are generically unique, and the rate of convergence of BR dynamics is generically exponential. Combined with the results of the present paper, this allows us to show that BR dynamics are “well behaved” in almost all potential games.

Furthermore, in [26] Monderer and Shapley study the convergence of the

⁴A critical point x^* of a function $f : \mathbb{R}^n \rightarrow \mathbb{R}$ is said to be non-degenerate if the Hessian of f at x^* is non-singular. When a critical point is non-degenerate, one can understand the important local properties of f using only the gradient and Hessian of f . If a critical point is *degenerate* then heavy algebraic machinery may be required to understand the local properties of f . With regard to games, if x^* is an interior equilibrium point of a potential game with potential function U , then x^* is regular if and only if x^* is a non-degenerate critical point of f . For non-interior equilibrium points the story is more involved, but the main idea is the same.

closely related FP algorithm in potential games and show convergence to the set of NE. In particular, they show that it is possible for FP to converge to completely mixed NE in potential games, which can be highly problematic for a number of reasons [36, 37, 29]. However, they conjecture that such behavior is exceptional; that is, they conjecture that in generic two-player potential games FP always converges to pure NE (see [26], Section 2).⁵ Regular potential games are well suited to studying this conjecture.⁶ Theorem 1 of the present paper combined with Theorem 1 of [29] shows that for the continuous-time version of FP [35, 39] (which is equivalent to BR dynamics after a time change [35]), this conjecture holds generically for potential games of arbitrary size; that is, in any regular potential game (and hence almost all potential games) continuous-time FP dynamics converge to pure NE from almost all initial conditions.

While regular equilibria have traditionally been studied as an equilibrium selection criterion in games, in potential games, pure strategy NE are naturally selected by virtue of the potential function. We emphasize that our objective in studying regular potential games is not to argue that (possibly mixed) regular equilibria are more natural than pure equilibria in potential games. Rather, we wish to demonstrate that (i) in almost all potential games, all equilibria possess desirable structural properties, (ii) properties that hold for regular potential games are inherently robust to payoff perturbations, and (iii) properties that hold only for irregular potential games are inherently nonrobust to payoff perturbations. We note, however, that our main result does imply that the two considerations (having a pure strategy regular NE) can be aligned in generic (i.e., regular) potential games where a potential maximizer must also be a regular equilibrium. We also remark that, as discussed earlier, regular potential games can be particularly useful in the study of game-theoretic learning processes, and a key aim of the present work is to facilitate the study of such learning processes by establishing the genericity of regular potential games.

The remainder of the paper is organized as follows. In Section 1.1 we outline our strategy for proving Theorem 1. Section 2 sets up notation. Section 3 presents a pair of key non-degeneracy conditions that are equivalent

⁵Since their reasoning relies on the improvement principle [38], which does not hold in games with more than two players, they limit their conjecture to two-player games.

⁶Harsanyi's result (that regular games are generic in the space of all games) does not aid in addressing this conjecture which deals specifically with regular potential games.

to regularity in potential games. Section 4 presents Proposition 13 which states that almost all identical payoff games are second-order non-degenerate (this proposition is the technical core of the paper) and sets up notation for analyzing second-order degeneracy (and regularity). Section 5 takes a brief digression to contrast proof techniques in general games vs potential games. Section 6 proves Proposition 13. Section 7 proves that almost all identical payoff games are first-order non-degenerate. Section 8 proves that almost all exact and weighted potential games are regular. Section 9 concludes the paper.

1.1. Proof Strategy

Our strategy for proving Theorem 1 is as follows. A potential game will be seen to be regular if and only if the corresponding identical payoffs game (with the potential function being the common utility) is regular. Thus, the problem of proving Theorem 1 reduces, by and large, to the following proposition.

Proposition 3. *Almost all games with identical payoffs are regular.*

The bulk of the paper will be devoted to proving this proposition. We will prove it as follows.

An equilibrium in an identical-payoffs (or potential) game can be shown to be regular if and only if the derivatives of the potential function satisfy two simple non-degeneracy conditions which we refer to as first- and second-order non-degeneracy (see Section 3). The first-order condition deals with the gradient of the potential function. (A first-order non-degenerate equilibrium is referred to as a *quasi-strong* equilibrium in [3], and a *quasi-strict* equilibrium in other works [4].)⁷ An equilibrium x^* is second-order non-degenerate if the Hessian of the potential function taken with respect to the support of x^* is invertible.

We say a potential game is first-order (second-order) non-degenerate if all equilibria in the game satisfy the first-order (second-order) non-degeneracy conditions. We will prove Proposition 3 by showing that:

- (i) An equilibrium of an identical payoffs game is regular if and only if it is first- and second order non-degenerate (see Section 3 Lemma 12),

⁷We prefer the term first-order non-degenerate name because it emphasizes the role of the potential function and is consistent with the notion of second-order degeneracy.

- (ii) Almost all games with identical payoffs are second-order non-degenerate (see Section 4.1 Proposition 13),
- (iii) Almost all games with identical payoffs are first-order non-degenerate (see Section 7 Proposition 21).

We remark that in Propositions 13 and 21 we show that the subset of irregular games has (appropriately dimensioned) Lebesgue measure zero. By Remark 5 this is sufficient to establish regularity in almost all identical payoff games.

After proving Proposition 3, we extend the result to exact and weighted potential games. This extension is straightforward and is accomplished in Section 8 (see Proposition 22). Propositions 3 and 22 together prove Theorem 1.

2. Notation

We will outline the notation and terminology used throughout the paper below. A short glossary of additional standard notation used in the paper can be found in Appendix A.

2.1. Normal Form Games

A game in normal form is given by a tuple $\Gamma := (N, (Y_i, u_i)_{i=1, \dots, N})$, where $N \in \{1, 2, \dots\}$ denotes the number of players, $Y_i := \{y_i^1, \dots, y_i^{K_i}\}$ denotes the set of pure strategies (or actions) available to player i , with cardinality $K_i := |Y_i|$, and $u_i : \prod_{j=1}^N Y_j \rightarrow \mathbb{R}$ denotes the utility function of player i .

Given some game Γ , let $Y := \prod_{i=1}^N Y_i$ denote the set of joint pure strategies available to players, and let

$$K := K_1 \times \dots \times K_N \tag{1}$$

denote the number of joint pure strategies. When defining spaces of games (e.g., as in Section 2.3) we will find it convenient to view $u_i = (u_i(y))_{y \in Y}$ as a vector in \mathbb{R}^K ; we will clearly indicate when using this abuse of notation.

The set of mixed strategies of player i is typically defined to be the probability simplex over A_i . It will simplify the presentation to consider an equivalent, but slightly modified, definition of the set of mixed strategies. Let

$$X_i := \{x_i \in \mathbb{R}^{K_i-1} : 0 \leq x_i^k \leq 1 \text{ for } k = 1, \dots, K_i - 1, \text{ and } \sum_{k=1}^{K_i-1} x_i^k \leq 1\}, \tag{2}$$

denote the set of mixed strategies of player i . (This definition will allow us to perform calculations without being directly encumbered by the hyperplane constraint inherent in the probability simplex.) A strategy $x_i \in X_i$ is interpreted as follows: The scalar $1 - \sum_{k=1}^{K_i} x_i^k$ represents the weight placed on the first pure strategy $y_i^1 \in Y_i$, and for $k \in \{2, \dots, K_i - 1\}$, x_i^{k-1} represents the weight placed on the k -th pure strategy y_i^k . In order to later reference this interpretation, the following notation will be useful. Given $x_i \in X_i$, let

$$T_i^1(x_i) := 1 - \sum_{k=1}^{K_i-1} x_i^k \quad \text{and} \quad T_i^k(x_i) := x_i^{k-1}, \quad k \geq 2. \quad (3)$$

Let $X := \prod_{i=1}^N X_i$ denote the set of joint mixed strategies and let $X_{-i} := \prod_{j \neq i} X_j$. When convenient, given a mixed strategy $x = (x_1, \dots, x_N) \in X$, we use the notation x_{-i} to denote the tuple $(x_j)_{j \neq i}$.

Given a mixed strategy $x \in X$, the expected utility of player i is given by

$$U_i(x) = \sum_{k=1}^{K_i-1} x_i^k U_i(y_i^{k+1}, x_{-i}) + \left(1 - \sum_{k=1}^{K_i-1} x_i^k\right) U_i(y_i^1, x_{-i}). \quad (4)$$

A strategy $x \in X$ is said to be a *Nash equilibrium* (or simply an *equilibrium*) if $x_i \in \arg \max_{x'_i \in X_i} U_i(x'_i, x_{-i})$ for all $x'_i \in X_i$, $i = 1, \dots, N$.

We say that a strategy x_i is completely mixed if it lies in the interior of X_i and we say that x_i is a pure strategy if it is a vertex of X_i . Otherwise, we say that x_i is an incompletely mixed strategy.

For $x_i \in X_i$, we define the *carrier set* of x_i to be

$$\text{carr}_i(x_i) := \{y_i^k \in Y_i : T_i^k(x_i) > 0\}, \quad (5)$$

that is, $\text{carr}_i(x_i)$ is the subset of pure strategies in Y_i that receive positive weight under x_i . (This is just a slight modification of the notion of a *support set* to the present context.) For $x = (x_1, \dots, x_N) \in X$ let $\text{carr}(x) := \text{carr}_1(x_1) \cup \dots \cup \text{carr}_N(x_N)$.

Suppose $C = C_1 \cup \dots \cup C_N$, where for each $i = 1, \dots, N$, C_i is a nonempty subset of Y_i . We say that C is the carrier for $x = (x_1, \dots, x_N) \in X$ if $C = \text{carr}(x)$, or equivalently, if $C_i = \text{carr}_i(x_i)$ for $i = 1, \dots, N$.

2.2. Potential Games

Following [13, 26], a game is said to be a *weighted potential game* if there exists a function $u : Y \rightarrow \mathbb{R}$ and a vector of positive weights $(w_i)_{i=1}^N \in \mathbb{R}^N$

such that

$$u_i(y'_i, y_{-i}) - u_i(y''_i, y_{-i}) = w_i(u(y'_i, y_{-i}) - u(y''_i, y_{-i})) \quad (6)$$

A game is said to be an *exact potential game* if (6) holds with $w_i = 1$ for all $i = 1, \dots, N$. A game is said to be an *identical-payoffs game* if there exists a function $u : Y \rightarrow \mathbb{R}$ such that $u_i(y) = u(y)$ for all $y \in Y$, $i = 1, \dots, N$.

When we refer simply to a “potential game” we mean a weighted potential game, which includes the other classes of games as special cases.

Given a potential game, we let $U : X \rightarrow \mathbb{R}$ be the multilinear extension of u defined by

$$U(x) = \sum_{k=1}^{K_i-1} x_i^k U(y_i^{k+1}, x_{-i}) + \left(1 - \sum_{k=1}^{K_i-1} x_i^k\right) U(y_i^1, x_{-i}). \quad (7)$$

We will typically refer to U as the *potential function* and to u as the *pure form of the potential function*.

By way of notation, given a pure strategy $y_i \in Y_i$ and a mixed strategy $x_{-i} \in X_{-i}$, we will write $U(y_i, x_{-i})$ to indicate the value of U when player i uses a mixed strategy placing all weight on the y_i and the remaining players use the strategy $x_{-i} \in X_{-i}$.

2.3. Almost All Games

The following definition specifies the notion of “almost all” to be used in the paper.

Definition 4. *Given some property \mathcal{S} , we say that \mathcal{S} holds for almost all points in some Euclidean space \mathbb{R}^n if the set where \mathcal{S} fails to hold is a closed set with \mathcal{L}^n -measure zero.*

We say that a game Γ has size $(N, (K_i)_{i=1}^N)$ if Γ is a general N -player game and the size of the action space of each player satisfies $|Y_i| = K_i \in \{2, 3, \dots\}$, $i = 1, \dots, N$. Let K be as defined in (1) so that K gives the number of pure strategies in a game of size $(N, (K_i)_{i=1}^N)$.

A game of size $(N, (K_i)_{i=1}^N)$ is uniquely represented by the vector $u := (u_i(y))_{y \in Y, i=1, \dots, N} \in \mathbb{R}^{NK}$ which specifies the utility received by each player for each pure strategy $y \in Y$. We will frequently refer to u as the vector of *utility coefficients*. The set of all games of this size is equivalent to \mathbb{R}^{NK} .

Let $\mathcal{W}, \mathcal{P}, \mathcal{I} \subset \mathbb{R}^{NK}$, denote the subsets of weighted potential games, exact potential games, and identical-payoffs games respectively. (An explicit construction of \mathcal{W} and \mathcal{P} can be found in Section 8.) Let $K_w := \dim \mathcal{W}$ and $K_p := \dim \mathcal{P}$. The sets of weighted and exact potential games are equivalent to the Euclidean spaces \mathbb{R}^{K_w} and \mathbb{R}^{K_p} respectively.

An identical-payoffs game is uniquely represented as a vector $u \in \mathbb{R}^K$ denoting the payoff (identical for all players) received for each action $y \in Y$, and we will represent this set of games as $\mathcal{I} := \mathbb{R}^K$.

For a fixed game size, each class of games discussed above is equivalent to some Euclidean space. We will say that almost all games in a given class are regular if, for any game size, almost all games (per Definition 4) in the class are regular.

Remark 5 (Closedness). *It was shown by Harsanyi [3, 4] that the set of irregular games of (any) size $(N, (K_i)_{i=1}^N)$ is a closed subset in the space of all games of size $(N, (K_i)_{i=1}^N)$. It is straightforward to extend this result to the various subclasses of potential games. In particular, we have:*

- (i) *The set of irregular weighted potential games is closed with respect to \mathcal{W} .*
- (ii) *The set of irregular exact potential games is closed with respect to \mathcal{P} .*
- (iii) *The set of irregular games with identical payoffs is closed with respect to \mathcal{I} .*

For practical purposes this means that in order to verify that almost all potential games of a given class are regular, we need only verify that the subset of irregular games has appropriately dimensioned Lebesgue measure zero.

3. Regular Equilibria in Potential Games

In potential games, regular equilibria have a simple and intuitive meaning.⁸ An equilibrium in a potential game is regular if (i) it is quasi-strict in the sense of [4] (a condition we will refer to as first-order non-degeneracy), and (ii) the Hessian of U (the potential) taken with respect to the support of the equilibrium is invertible (a condition we will refer to as second order non-degeneracy). For equilibria in the interior of the strategy space, this simply

⁸In general games, the definition of a regular equilibrium is somewhat abstruse, relying on the invertibility of the Jacobian of a particular map [4]. See Section 5.1 for more details.

reduces to an equilibrium being regular if and only if it is a non-degenerate critical point of U in the traditional sense.

Since the non-degeneracy conditions given in this section (applicable only within the class of potential games) are simpler to work with than the traditional definition of regularity, we will work with directly with these conditions through the remainder of the paper. However, the traditional definition of regularity can be found in Section 5 (see Definition 14) where we contrast our techniques with those required in general games.

The remainder of the section is organized as follows. In Sections 3.1–3.2 we define the notions of first and second-order degeneracy. In Section 3.3 we show that, in a potential game, these conditions are equivalent to regularity.

3.1. First-Order Degeneracy

Let $C = C_1 \cup \dots \cup C_N$, $C_i \subset Y_i \forall i = 1, \dots, N$ be some carrier set. Let $\gamma_i := |C_i|$ and assume that the strategy set Y_i is reordered so that $\{y_i^1, \dots, y_i^{\gamma_i}\} = C_i$. Under this ordering, the first $\gamma_i - 1$ components of any strategy x_i with $\text{carr}_i(x_i) = C_i$ are free (not constrained to zero by C_i) and the remaining components of x_i are constrained to zero. That is, the subvector $(x_i^k)_{k=1}^{\gamma_i-1}$ is free under C_i and the subvector $(x_i^k)_{k=\gamma_i}^{K_i} = 0$. The set of strategies $\{x \in X : \text{carr}(x) = C\}$ is precisely the interior of the face of X given by

$$X_C := \{x \in X : x_i^k = 0, k = \gamma_i, \dots, K_i - 1, i = 1, \dots, N\}. \quad (8)$$

Definition 6 (First-Order Degenerate Equilibrium). *Suppose Γ is a potential game with potential function U . Suppose $x^* \in X$ is an equilibrium of Γ with carrier C . We say that x^* is a first-order degenerate equilibrium if there exists a pair (i, k) , $i = 1, \dots, N$, $k = \gamma_i, \dots, K_i - 1$ such that*

$$\frac{\partial U(x^*)}{\partial x_i^k} = 0 \quad (9)$$

and we say x^* is first-order non-degenerate otherwise.

We note that the condition (9) in the definition of first-order degeneracy is implicitly dependent on the carrier C (since $\gamma_i = |C_i|$ and we assume that $C_i = \{y_i^1, \dots, y_i^{\gamma_i}\}$).


Definition 7 (First-order Degenerate Game). *We say a game is first-order degenerate if it has an equilibrium that is first-order degenerate, and we say the game is first-order non-degenerate otherwise.*

Example 8. Consider the 3×2 identical-payoffs game with payoff matrices


$$M_1 = \begin{pmatrix} 1 & -4 \\ -1 & 1 \end{pmatrix} \quad M_2 = \begin{pmatrix} -1 & 1 \\ 1 & 2 \end{pmatrix},$$

where player 1 is the “row player”, player 2 is the “column player”, and player 3 is the “matrix player.” Recalling that x_i denotes the probability of player i playing action y_i^2 (since each player has only two strategies, we drop the superscripts on x_i), this game has a first-order degenerate equilibrium at the strategy $x^* = (1/2, 0, 1/2)$.

This may be visualized in terms of the gradient of U . The three (gradient) level surfaces along which $\frac{\partial U(x)}{\partial x_i} = 0$, $i = 1, 2, 3$, are displayed in Figures 1a–1b. The equilibrium x^* has carrier $C = Y_1 \cup \{y_2^1\} \cup Y_3$ and lies on the face of X given by $\{x \in X : x_2 = 0\}$. Yet, x^* also lies at the intersection of all three level surfaces (and thus is a bona fide critical point of U). In particular, x^* lies on the level surface, $\{x : \frac{\partial U(x^*)}{\partial x_2} = 0\}$, making it first-order degenerate.



(a) Level surfaces of $\frac{\partial U(x)}{\partial x_i}$, $i = 1, 2, 3$.



(b) Alternate view of gradient level surfaces.

Remark 9 (Equivalence to Quasi-Strict Equilibria). An equilibrium x^* with carrier C is first order non-degenerate if and only if, for every player i , the set of pure-strategy best responses to x_{-i}^* coincides with C_i . (This is verified using the multi-linearity of U .) We note that, using this later definition, Harsanyi [3] referred to first-order non-degenerate equilibria as quasi-strong equilibria. In other works these have been referred to as quasi-strict equilibria [4]. We prefer to use the term first-order non-degenerate in order to

emphasize that we are concerned with the gradient of the potential function and to keep the nomenclature consistent with the notion of second-order non-degeneracy, introduced next.

3.2. Second-Order Degeneracy

Suppose that Γ is a potential game with potential function U . Let C be some carrier set. Let $\tilde{N} := |\{i = 1, \dots, N : \gamma_i \geq 2\}|$, and assume that the player set is ordered so that $\gamma_i \geq 2$ for $i = 1, \dots, \tilde{N}$. Under this ordering, for strategies with $\text{carr}(x) = C$, the first \tilde{N} players use mixed strategies and the remaining players use pure strategies. Assume that $\tilde{N} \geq 1$ so that any x with carrier C is a genuine mixed (not pure) strategy.

Let the Hessian of U taken with respect to C be given by

$$\tilde{\mathbf{H}}(x) := \left(\frac{\partial^2 U(x)}{\partial x_i^k \partial x_j^\ell} \right)_{\substack{i,j=1,\dots,\tilde{N}, \\ k=1,\dots,\gamma_i-1, \\ \ell=1,\dots,\gamma_j-1}}. \quad (10)$$

Note that this definition of the Hessian restricts attention to the components of x that are free (i.e., unconstrained) under C . That is, $\tilde{\mathbf{H}}(x)$ taken with respect to C is the Hessian of $U|_{X_C}$ at x .

Definition 10 (Second-Order Degenerate Equilibrium). *Let Γ be a potential game with potential function U . We say an equilibrium $x^* \in X$ of Γ is second-order degenerate if the Hessian $\tilde{\mathbf{H}}(x^*)$ taken with respect to $\text{carr}(x^*)$ is singular, and we say x^* is second-order non-degenerate otherwise.*

Definition 11 (Second-Order Degenerate Game). *We say a game is second-order degenerate if it has an equilibrium that is second-order degenerate, and we say the game is second-order non-degenerate otherwise.*

3.3. Regular Equilibria and Non-Degeneracy Conditions

The following lemma shows that within the class of potential games, the first and second-order non-degeneracy conditions defined above are equivalent to regularity.

Lemma 12. *Let Γ be a potential game. Then,*

(i) *If an equilibrium x^* is first-order non-degenerate, then it is second-order non-degenerate if and only if it is regular.*

(ii) *If an equilibrium x^* is regular, then it is first-order non-degenerate.*

In particular, an equilibrium x^ is regular if and only if it is both first and second-order non-degenerate.*

The proof of this lemma follows readily from the definitions of regularity (see Definition 14) and first and second-order degeneracy, and is omitted for brevity.

4. Second-Order Degeneracy in Games with Identical Payoffs

In the previous section we established that a game is regular if and only if it is first- and second-order non-degenerate. In this section we will state our main result regarding second-order non-degeneracy—namely, that the set of identical payoff games that are second-order degenerate has Lebesgue measure zero (see Proposition 13). Later, in Section 7 we will show the analogous result for first-order degenerate games (Proposition 21). By Remark 5, this will be sufficient to prove Proposition 3.

This section is organized as follows. In Section 4.1 we state our main result for second-order degenerate games. In Section 4.2 we outline our strategy for proving Proposition 13. In Section 4.3 we set up notation required for the analysis of second-order non-degeneracy (and the traditional notion of regularity). The proof of Proposition 13 will then be given in Section 6 after a brief interlude to discuss general games.

4.1. Second-Order Degenerate Games

In the following proposition we will assume the number of players N and action-space sizes K_i , $i = 1, \dots, N$ are fixed, and let K be as defined in (1).

Proposition 13. *The set of identical-payoff games that are second-order degenerate has \mathcal{L}^K -measure zero.*

This proposition will be proved in Section 4.3.

4.2. Roadmap of Proof of Proposition 13

Our strategy for proving Proposition 17 will be as follows.

1. Fix a carrier set C with $|C| = \gamma$. Following (8), let

$$X_C^* := \{x : \text{carr}(x) = C\}.$$

We will begin by deducing that for any equilibrium $x^* \in X_C^*$, the key relation (21) holds, where $\mathbf{A}(x)$ is defined in (20).

2. Proposition 17 shows that for every $x \in X_C^*$, the matrix $\mathbf{A}(x)$ has full row rank.

3. Using Proposition 17 we construct a countable cover $(B_\ell)_{\ell \geq 1}$ of X_C^* such that for each ℓ , we may choose γ columns of $\mathbf{A}(x) \in \mathbb{R}^{\gamma \times K}$ so that these same columns are linearly independent for all $x \in B_\ell$. For each ℓ , the associated columns form an invertible square submatrix of $\mathbf{A}(x)$ for all $x \in B_\ell$. Using this invertible submatrix and (21), we construct a mapping $\rho_\ell : B_\ell \times \mathbb{R}^{K-\gamma} \rightarrow \mathbb{R}^K$, which allows us to recover the full vector of potential coefficients $u \in \mathbb{R}^K$ given an equilibrium $x^* \in X_C^*$ and a $(K - \gamma)$ -dimensional subvector of the potential coefficients. (See Lemma 18, (31)–(32), and Corollary 19.)

4. We then demonstrate that if $x^* \in B_\ell$ is a NE of a game with potential-function coefficients $u \in \mathbb{R}^K$, then x^* is second-order degenerate if and only if the Jacobian of ρ_ℓ is non-invertible. (See Lemma 20.) Since the output of ρ_ℓ is the full vector of coefficients u , this implies that u lies in the set of critical values of ρ_ℓ (i.e. the set of outputs of ρ_ℓ at which the Jacobian is not invertible).

5. From here, the proof follows from Sard’s theorem [41]. In particular, applying Sard’s theorem to ρ_ℓ shows that the set of games with second-order degenerate equilibria lying in B_ℓ has measure zero. Repeating this argument over all B_ℓ (of which there is a countable number) and for all carriers C (of which there is a finite number) yields Proposition 13.

We remark that the above proof strategy follows the same general strategy used by Harsanyi in [3] to prove regularity in general games. However, important technical challenges arise in the case of potential games that are not present in general games. This is discussed further in Sections 5.2–5.3.

4.3. Analysis of Second-Order Degeneracy: Setup

Proposition 13 will be proved using Sard’s theorem as outlined above. In this section we introduce some pertinent notation that will allow us to establish (21), which is the key relationship allowing us to analyze second-order degeneracy.⁹

Note that the set of joint pure strategies Y may be expressed as an ordered set $Y = \{y^1, \dots, y^K\}$ where each $y^\tau \in Y$, is an N -tuple of strategies, $\tau \in \{1, \dots, K\}$. We will assume a particular ordering for this set after Proposition 17.

⁹We note that this setup is useful both for analyzing potential games and general games. Indeed, in Section 5 this setup will be reused to establish the key inequality (25), which is the analog of (21) in general games.

In an identical-payoffs game there exists a single function $u : Y \rightarrow \mathbb{R}$ such that $u_i = u$ for all $i = 1, \dots, N$. If we consider the vector of utility coefficients $u = (u(y))_{y \in Y} \in \mathbb{R}^K$ as a variable, then by (7), U is linear in u .¹⁰ At this point we will express U in a more convenient form (see (12)) which more clearly exposes this relationship.

Let $\tau \in \{1, \dots, K\}$, $i \in \{1, \dots, N\}$ and $x_i \in X_i$. We define $q_i^\tau : X_i \rightarrow [0, 1]$ by

$$q_i^\tau(x_i) := T_i^k(x_i) \quad (11)$$

where k corresponds to the action played by player i in the tuple y^τ , i.e., $(y^\tau)_i = y_i^k$, and where $T_i^k(x_i)$ is defined in (3). In words, given some mixed strategy $x_i \in X_i$, $q_i^\tau(x_i)$ gives the weight x_i places on the particular action used by player i in the τ -th action tuple y^τ .

In an abuse of notation, given a pure strategy $y_i^k \in Y_i$, we let $q_i^\tau(y_i^k) = 1$ if $(y^\tau)_i = y_i^k$ and $q_i^\tau(y_i^k) = 0$ otherwise.

Given a fixed vector of utility coefficients $u \in \mathbb{R}^K$, the utility function $U : X \rightarrow \mathbb{R}$ may be expressed as

$$U(x) = \sum_{\tau=1}^K u^\tau \left[\prod_{i=1}^N q_i^\tau(x_i) \right]. \quad (12)$$

Note that this form makes it clear that U is linear in u .

Now, let $C = C_1 \cup \dots \cup C_N$ be some carrier set. The analysis through the remainder of the section will rely on this carrier set being fixed, and many of the subsequent terms are implicitly dependent on the choice of C . In keeping with our prior convention we let $\gamma_i := |C_i|$, and let $\tilde{N} := \{i \in \{1, \dots, N\} : \gamma_i \geq 2\}$, and without loss of generality we assume that the strategies in Y_i are ordered so that $C_i = \{y_i^1, \dots, y_i^{\gamma_i}\}$.

Any x_i with carrier C_i has precisely $\gamma_i - 1$ free components (i.e., not constrained to zero by C_i). The joint strategy $x = (x_1, \dots, x_N)$ is a vector with

$$\gamma := \sum_{i=1}^N (\gamma_i - 1) \quad (13)$$

free components.

¹⁰The potential function U is, of course, a function of both x and u . However, since we will only exploit the dependence on u in this section, we generally stick to the standard game-theoretic convention of writing U as a function of x only [42].

By (7) we have that

$$\frac{\partial U(x)}{\partial x_i^k} = U(y_i^{k+1}, x_{-i}) - U(y_i^1, x_{-i}) =: F_i^k(x, u) \quad (14)$$

for $i = 1, \dots, \tilde{N}$, $k = 1, \dots, \gamma_i - 1$. Let

$$F(x, u) := \left(F_i^k(x, u) \right)_{\substack{i=1, \dots, \tilde{N} \\ k=1, \dots, \gamma_i - 1}}. \quad (15)$$

Given an $x \in X$, it is at times useful to decompose it as $x = (x_p, x_m)$, where $x_m = (x_i^k)_{i=1, \dots, \tilde{N}, k=1, \dots, \gamma_i - 1}$ and x_p contains the remaining components of x . (The subscript of x_m is indicative of “mixed strategy components” of x and x_p indicative of “pure strategy components” of x .) In this decomposition, x_m is a γ -dimensional vector containing the free components of x . Taking the Jacobian of F in terms of the components of x_m we find that

$$D_{x_m} F(x_p, x_m, u) = \tilde{\mathbf{H}}(x), \quad (16)$$

where $D_{x_m} F(x_p, x_m, u) = \left(\frac{\partial F}{\partial x_i^\ell} \right)_{\substack{i=1, \dots, \tilde{N} \\ \ell=1, \dots, \gamma_i - 1}}$.

Let x^* be a mixed equilibrium with carrier C . Differentiating (7) we see that at the equilibrium x^* we have $\frac{\partial U(x^*)}{\partial x_i^k} = 0$, or equivalently,

$$F_i^k(x^*, u) = U(y_i^{k+1}, x_{-i}^*) - U(y_i^1, x_{-i}^*) = 0 \quad (17)$$

for $i = 1, \dots, \tilde{N}$, $k = 1, \dots, \gamma_i - 1$. Using (12) in the above we get

$$F_i^k(x^*, u) = \sum_{\tau=1}^K u^\tau \left[(q_i^\tau(y_i^{k+1}) - q_i^\tau(y_i^1)) \prod_{j \neq i} q_j^\tau(x_j^*) \right] = 0. \quad (18)$$

Note that (18) is a linear in u . We would like to express (18) as a matrix equation (i.e., $\mathbf{A}(x)u = 0$ for some matrix $\mathbf{A}(x)$ dependent only on x). With this in mind, it will be convenient to develop notation relating the ordering of the pure-strategy set Y with the ordering of $(F_i^k)_{i=1, \dots, \tilde{N}, k=1, \dots, \gamma_i - 1}$. Given $i \in \{1, \dots, \tilde{N}\}$, $k \in \{1, \dots, \gamma_i - 1\}$, let

$$s^*(i, k) := \begin{cases} k & \text{for } i = 1, \\ \sum_{j=1}^{i-1} (\gamma_j - 1) + k & \text{for } i \geq 2. \end{cases}$$

Define $i^* : \{1, \dots, \gamma\} \rightarrow \{1, \dots, \tilde{N}\}$ and $k^* : \{1, \dots, \gamma\} \rightarrow \{1, \dots, \max_i \{\gamma_i - 1\}\}$ to be the inverse of s^* ; that is

$$s^*(i^*(s), k^*(s)) = s \quad (19)$$

for all $s = 1, \dots, \gamma$.

The above notation may be understood as follows: Effectively, we have stacked $(F_i^k)_{i=1, \dots, \tilde{N}, k=1, \dots, \gamma_i - 1}$ into a single vector. Given a pair (i, k) , the function $s(i, k) \in \{1, \dots, \gamma\}$ gives the corresponding index in this vector. Conversely, given some $s \in \{1, \dots, \gamma\}$, the functions $i^*(s)$ and $k^*(s)$ simply return the player i or action k corresponding to entry s in the vector.

Given an $x \in X$, let $\mathbf{A}(x) = (a_{s,\tau}(x))_{\substack{s=1, \dots, \gamma_i \\ \tau=1, \dots, K}} \in \mathbb{R}^{\gamma \times K}$ be defined as the matrix with entries

$$a_{s,\tau}(x) := \left(q_{i^*(s)}^\tau(y_i^{k^*(s)+1}) - q_{i^*(s)}^\tau(y_i^1) \right) \prod_{j \neq i^*(s)} q_j^\tau(x_j), \quad (20)$$

Reexpressing (18) using this notation, we see that for any equilibrium x^* with carrier C we have

$$\mathbf{A}(x^*)u = 0. \quad (21)$$

5. General Games: Discussion and Proof Techniques

In this section we take a short digression to discuss the proof that almost all general games are regular, and contrast this with the potential games case. We note that this section may be skipped without loss of continuity.

In Section 5.1 we recall the traditional definition of a regular equilibrium. In Section 5.2 we discuss the strategy used by Harsanyi [3] to prove that almost all general games are regular. In Section 5.3 we discuss the technical challenges arising in the case of potential games and contrast our proof techniques with those used by Harsanyi.

5.1. Regular Equilibria in General Games

We now recall the traditional definition a regular equilibrium as given in [40, 4]. Let the game size $(N, (K_i)_{i=1}^N)$ be fixed.

As discussed in Section 2.3, a game is uniquely defined by a vector $u \in \mathbb{R}^{NK}$ (which we refer to as the utility coefficient vector) specifying the pure-strategy utility received by each player.

Given a strategy $x \in X$ and vector of utility coefficients $u \in \mathbb{R}^{NK}$, let

$$\tilde{F}_i^k(x, u) := T_i^1(x_i)x_i^k[U_i(y_i^{k+1}, x_{-i}) - U_i(y_i^1, x_{-i})] \quad (22)$$

for $i = 1, \dots, N$, $k = 1, \dots, K_i - 1$, and let

$$\tilde{F}(x, u) := \left(\tilde{F}_i^k(x, u) \right)_{\substack{i=1, \dots, N \\ k=1, \dots, K_i-1}}. \quad (23)$$

Definition 14 (Regular Equilibrium). *Let $x^* \in X$ be an equilibrium of a game with utility coefficient vector u . Assume the action set Y_i of each player is reordered so that $y_i^1 \in \text{carr}_i(x_i^*)$. The equilibrium x^* is said to be regular if the Jacobian of $\tilde{F}(x^*, u)$, given by $D_x \tilde{F}(x^*, u)$, is non-singular.*

Remark 15. *We note that if x^* is regular, then the Jacobian of $\tilde{F}(x^*, u)$ can be shown to be nonsingular under any reordering of Y_i in which the reference action satisfies $y_i^1 \in \text{carr}_i(x_i^*)$ for all $i = 1, \dots, N$ (see [40], Theorem 3.8). This justifies the use of an arbitrary reference action $y_i^1 \in \text{carr}_i(x_i^*)$ in the above definition.*

Remark 16. *The notion of a regular equilibrium is traditionally defined by considering mixed strategies directly in the probability simplex rather than X_i [4]. Using the definition of \tilde{F}_i^k and the properties of the determinant of a matrix, it is readily confirmed that the definition of regularity given in Definition 14 coincides with the traditional definition in [4].*

5.2. Almost All General Games are Regular: Proof Strategy

We will now review, at a high level, the classical technique for proving that almost all general games are regular given in [3].

The strategy is as follows.

1. Suppose that a carrier set C is fixed.
2. Let $u \in \mathbb{R}^{NK}$ denote a vector of utility function coefficients, and let u be broken down as $u = (u^*, u^{**})$, where $u^* \in \mathbb{R}^\gamma$, $u^{**} \in \mathbb{R}^{NK-\gamma}$.
3. Construct a C^1 function $\rho^{**} : X_C^* \times \mathbb{R}^{NK-\gamma} \rightarrow \mathbb{R}^\gamma$ such that, given any equilibrium x^* with carrier C and a partial vector of utility coefficients $u^{**} \in \mathbb{R}^{NK-\gamma}$, we have¹¹

$$u^* = \rho^{**}(x^*, u^{**}).$$

¹¹The notation ρ^{**} , ρ^* , u^* and u^{**} is used here to be consistent with the usage in [3].

That is, ρ^{**} allows one to recover u^* given x^* and u^{**} . Given ρ^{**} , it is trivial to construct a C^1 function $\rho^* : X_C^* \times \mathbb{R}^{NK-\gamma} \rightarrow \mathbb{R}^{NK}$ such that

$$u = \rho^*(x^*, u^{**}),$$

so that ρ^* recovers the full vector u given x^* and u^{**} .

4. Show that the set of all irregular games having an equilibrium with carrier C lies in the “critical values” set of ρ^* (i.e, the image of the set of critical points). By Sard’s theorem, this implies that all such games lie in an \mathcal{L}^{NK} -measure zero set.
5. Since the argument holds for arbitrary carrier C and there are a finite number of possible carriers, this completes the proof.

The crucial task in the above proof outline is the construction of the function ρ^{**} which isolates the set of irregular games in a critical values set. For the sake of comparison with the identical payoff games case, we will now review the technique for constructing ρ^{**} in [3] more detail.

Suppose a carrier C is fixed, and suppose x^* is an equilibrium with carrier C . Differentiating (using the same reasoning used to arrive at (17)) we see that at x^* we have

$$U_i(y_i^{k+1}, x_{-i}^*) - U_i(y_i^1, x_{-i}^*) = 0 \tag{24}$$

for $i = 1, \dots, \tilde{N}$, $k = 1, \dots, \gamma_i - 1$. Let $u = (u_1, \dots, u_N) \in \mathbb{R}^{NK}$ represent the vector of all player’s pure-strategy utilities. As with (17), the equality (24) is linear in u and may be expressed as a matrix equation¹²

$$\mathbf{B}(x)u = 0, \tag{25}$$

where $\mathbf{B}(x) \in \mathbb{R}^{\gamma \times NK}$. The matrix $\mathbf{B}(x)$ has particularly convenient structure. To elucidate this structure, first note that, as with (12), the (expected) utility of player i in a general game may be expressed as

$$U_i(x) = \sum_{\tau=1}^K u_i^\tau \left[\prod_{i=1}^N q_i^\tau(x_i) \right].$$

¹²To be consistent with the proof in the potential games case, we have modified the presentation of Harsanyi’s technique to accommodate matrix notation. However, modulo notational differences, the argument discussed here is the same as [3].

Using this form, the equality (24) is expressed as

$$\sum_{\tau=1}^K u_i^\tau \left[(q_i^\tau(y_i^{k+1}) - q_i^\tau(y_i^1)) \prod_{j \neq i} q_j^\tau(x_j^*) \right] = 0, \quad (26)$$

for $i = 1, \dots, \tilde{N}$, $k = 1, \dots, \gamma_i - 1$. For each $i \in \{1, \dots, \tilde{N}\}$, let $\mathbf{B}_i(x) = (b_{i,k,\tau}(x))_{k=1, \dots, \gamma_i-1, \tau=1, \dots, K}$ be the matrix with entries

$$b_{i,k,\tau}(x) := (q_i^\tau(y_i^{k+1}) - q_i^\tau(y_i^1)) \prod_{j \neq i} q_j^\tau(x_j). \quad (27)$$

The equality (26) may now be expressed in the form of (25) with $\mathbf{B}(x)$ given by

$$\mathbf{B}(x) := \begin{pmatrix} \mathbf{B}_1(x) & \mathbf{0} & \cdots & \mathbf{0} \\ \mathbf{0} & \mathbf{B}_2(x) & \cdots & \mathbf{0} \\ \vdots & \ddots & & \vdots \\ \mathbf{0} & \cdots & \mathbf{0} & \mathbf{B}_{\tilde{N}}(x) \end{pmatrix}.$$

The important fact about $\mathbf{B}(x)$ is that there exist γ columns of $\mathbf{B}(x)$ so that these same columns form an invertible diagonal $\gamma \times \gamma$ submatrix of $\mathbf{B}(x)$, for all x with carrier C . (More details can be found in Appendix C.) Thus, without loss of generality, we may reorder the set of pure strategies so that the first γ columns of $\mathbf{B}(x)$ are linearly independent and we may write $\mathbf{B}(x) = [\mathbf{C}(x) \mathbf{D}(x)]$, where $\mathbf{C}(x) \in \mathbb{R}^{\gamma \times \gamma}$ is invertible for any x with carrier C .

Given a vector $u^{**} \in \mathbb{R}^{NK-\gamma}$ and an equilibrium x^* with carrier C , define

$$\rho^{**}(x, u^{**}) := \mathbf{C}(x)^{-1} \mathbf{D}(x) u^{**}. \quad (28)$$

Finally, recalling (25), we observe that ρ^{**} allows one to recover u^* given u^{**} and any equilibrium x^* with support C .

In the following section we will discuss the strategy for constructing an analogous function in the case of identical payoff games, and the challenges that arise in this case.

5.3. Comparison with Proof Technique in Identical-Payoff Games

We will now contrast the proof that almost all general games are regular with the proof that almost all identical payoff games are regular.

The main substantive difference between the proof of Proposition 13 and proof of generic regularity in the case of general games is the construction the functions isolating the subset of irregular games (Step 3 in Section 4.2 and Step 3 in Section 5.2). We will now briefly discuss the manner in which such a function is constructed in the identical-payoffs-game case and contrast this with the approach discussed in Section 5.2.

In the case of an identical-payoff game, at an equilibrium x^* with carrier C we have the key relationship (21) rather than (25). The matrices $\mathbf{A}(x)$ and $\mathbf{B}(x)$ are related via

$$\mathbf{A}(x) = \begin{pmatrix} \mathbf{B}_1(x) \\ \vdots \\ \mathbf{B}_N(x) \end{pmatrix}. \quad (29)$$

Suppose we fix some carrier C , and let γ be as defined in (13). Given a vector of potential function coefficients $u \in \mathbb{R}^K$, decompose it as $u = (u_1, u_2)$, where $u_1 \in \mathbb{R}^{K-\gamma}$, $u_2 \in \mathbb{R}^\gamma$.¹³

Analogous to the general games case, if we can find a function that recovers u_2 given u_1 and an equilibrium x^* with carrier C , then the set of second-order degenerate potential games will lie in the critical values set of this map (see Lemma 20). Thus, the key step in the proof of Proposition 13 is the construction of such a function.

In the case of general N -player games we were conveniently able to choose γ columns of $\mathbf{B}(x)$ that formed a $\gamma \times \gamma$ diagonal submatrix of $\mathbf{B}(x)$ that was invertible for all x with carrier C . Using this diagonal submatrix we were able to construct an appropriate function ρ^{**} in (28).

In the case of identical payoff games, we can (and will) follow a similar approach. If $\mathbf{A}(x)$ can be decomposed as $\mathbf{A}(x) = [\mathbf{E}(x) \mathbf{F}(x)]$, where $\mathbf{E}(x) \in \mathbb{R}^{\gamma \times \gamma}$ is invertible, then (as in (28)) the function $\rho(x, u_1) = \mathbf{E}(x)^{-1} \mathbf{F}(x) u_1$ will recover u_2 . However, in the case of identical payoff games, the problem of finding an invertible submatrix of $\mathbf{A}(x)$ is more challenging than the general games case. Unlike $\mathbf{B}(x)$, there do not exist γ columns of $\mathbf{A}(x)$ that form a invertible diagonal submatrix of $\mathbf{A}(x)$. Moreover, the pure strategies (corresponding to columns of $\mathbf{B}(x)$) used to construct Harsanyi's map ρ^{**} need not correspond to linearly independent columns of $\mathbf{A}(x)$.

¹³In Section 5.2 we used u^* and u^{**} to refer to an analogous breakdown of $u \in \mathbb{R}^{NK}$; this notation was preferred there to avoid confusion with individual player's utility functions.

For example, suppose Γ is a 2×2 game and let C be the carrier containing all pure strategies (so any x with $\text{carr}(x) = C$ is completely mixed). The matrix $\mathbf{A}(x)$ takes the form

$$\mathbf{A}(x) = \begin{pmatrix} -x_2 & -x_2 & (1-x_2) & (1-x_2) \\ -x_1 & (1-x_1) & -x_1 & (1-x_1) \end{pmatrix}. \quad (30)$$

(See Appendix B and, in particular, (B.1)–(B.5) for an in-depth characterization of the structure of $\mathbf{A}(x)$.) In (30) these pure-strategy choices correspond to the middle two columns of $\mathbf{A}(x)$, which are linearly dependent for $x_1 = x_2 = 1/2$. In order to ensure that we may select γ columns of $\mathbf{A}(x)$ that are linearly independent, we will show that for any x with $\text{carr}(x) \subseteq C$, the matrix $\mathbf{A}(x)$ has full rank (see Proposition 17). This will then allow us to construct a set of functions as in Step 3 of Section 4.2 (see also (31)) that recover u_2 given u_1 and x in some subset of the strategy space.

It should be noted that, for the sake of proving Proposition 13, it would be sufficient to prove the forthcoming Proposition 17 under the weaker condition $\text{carr}(x) = C$ (rather than $\text{carr}(x) \subseteq C$). However, we will prove Proposition 17 under the stronger condition that $\text{carr}(x) \subseteq C$ in order to apply the result later in the proof of Proposition 21.

6. Proof of Proposition 13

We will now prove Proposition 13, which states that almost all identical payoff games are second-order non-degenerate.

We begin with the following proposition that establishes the solvability of (21).

Proposition 17. *For any x such that $\text{carr}(x) \subseteq C$, the matrix $\mathbf{A}(x)$ has full row rank.*

Proposition 17 together with (21) will permit us to construct a family of functions $(\rho_\ell)_{\ell \geq 1}$ as outlined in Section 4.2. The complete proof of Proposition 17 can be found in Appendix B.

In broad strokes, the proof of Proposition 17 proceeds as follows: We will consider the “sign pattern” of the matrix $\mathbf{A}(x)$ (i.e., the matrix $\text{sgn}(\mathbf{A}(x))$, with $\text{sgn}(\cdot)$ defined in Appendix A). A matrix is called a *sign-pattern matrix* if its entries take on values from the set $\{-1, 0, 1\}$. The properties of sign-pattern matrices have been well studied in the literature [43, 44]. Of relevance

here, a sign pattern matrix \mathbf{M} is called an L -matrix if every matrix with the same sign pattern as \mathbf{M} has full rank. A relatively simple characterization of L -matrices is given in [43, 44]. Using this characterization, we prove that for any x with $\text{carr}(x) \subseteq C$, $\text{sgn}(\mathbf{A}(x))$ is an L -matrix. This implies the desired result.

Please see Appendix B for complete details of the proof of Proposition 17.

We will now use Proposition 17 to construct a countable cover $(B_\ell)_{\ell \geq 1}$ of the set $\{x : \text{carr}(x) = C\}$ and a family of functions $(\rho_\ell)_{\ell \geq 1}$ as outlined in Section 4.2.

Given the carrier C there are $\binom{K}{\gamma}$ possible combinations (of size γ) of the columns of $\mathbf{A}(x)$. For each $r = 1, \dots, \binom{K}{\gamma}$, let $\mathbf{A}_r(x) \in \mathbb{R}^{\gamma \times \gamma}$ denote a square matrix formed by taking one unique combination of the columns of $\mathbf{A}(x)$. We have the following lemma.

Lemma 18. *There exists a collection of open balls $(B_\ell)_{\ell \geq 1}$, $B_\ell \subset \{x : \text{carr}(x) = C\}$ that satisfy:*

- (i) $\bigcup_{\ell \geq 1} B_\ell = \{x \in X : \text{carr}(x) = C\}$
- (ii) For each $\ell \in \mathbb{N}$ there exists an $r_\ell \in \{1, \dots, \binom{K}{\gamma}\}$ such that $\mathbf{A}_{r_\ell}(x)$ is invertible for all $x \in B_\ell$.

Proof. For $r = 1, \dots, \binom{K}{\gamma}$, let

$$S_r := \{x \in X : \text{carr}(x) = C, \det \mathbf{A}_r(x) \neq 0\}.$$

By Proposition 17, no strategy $x \in X$ with $\text{carr}(x) = C$ may simultaneously be in all S_r^c . Hence $\bigcup_r S_r = \{x : \text{carr}(x) = C\}$. Note also that each S_r is open relative to $\{x \in X : \text{carr}(x) = C\}$. Thus, for each r we may construct a countable cover $(B_{r,\ell})_{\ell \geq 1}$ of S_r , such that $B_{r,\ell} \subset \{x : \text{carr}(x) = C\}$, $\ell \in \mathbb{N}$ and $\bigcup_\ell B_{r,\ell} = S_r$.

By construction we have (i) for any pair (r, ℓ) , the matrix $\mathbf{A}_r(x)$ is invertible for all $x \in B_{r,\ell}$, and (ii) $\bigcup_{r,\ell} B_{r,\ell} = \bigcup_r S_r = \{x : \text{carr}(x) = C\}$. Hence, $(B_{r,\ell})_{r=1, \dots, \binom{K}{\gamma}, \ell \in \mathbb{N}}$ is a countable cover with the desired properties. \square

Fix $\ell \in \{1, 2, \dots\}$. After reordering, $\mathbf{A}(x)$ may be partitioned as $\mathbf{A}(x) = [\tilde{\mathbf{A}}_{r_\ell}(x) \mathbf{A}_{r_\ell}(x)]$, where $\tilde{\mathbf{A}}_{r_\ell}(x)$ is a matrix formed by the columns of $\mathbf{A}(x)$ not used to form $\mathbf{A}_{r_\ell}(x)$. Let the strategy set Y be reordered in the same

way as the columns $\mathbf{A}(x)$.¹⁴ Given a vector of utility coefficients $u \in \mathbb{R}^K$, let it be partitioned as $u = (u_1, u_2)$, where $u_1 = (u^1, \dots, u^{K-\gamma})$ and $u_2 = (u^{K-\gamma+1}, \dots, u^K)$. Define $\tilde{\rho}_\ell : B_\ell \times \mathbb{R}^{K-\gamma} \rightarrow \mathbb{R}^\gamma$ by

$$\tilde{\rho}_\ell(x, u_1) := -\mathbf{A}_{r_\ell}(x)^{-1} \tilde{\mathbf{A}}_{r_\ell}(x) u_1, \quad (31)$$

If $x^* \in B_\ell$ is an equilibrium for some identical-payoffs game with utility coefficient vector u , then by (21) we have $\mathbf{A}(x^*)u = 0$. Since $\mathbf{A}_{r_\ell}(x^*)$ is invertible, this is equivalent to $u_2 = -\mathbf{A}_{r_\ell}(x^*)^{-1} \tilde{\mathbf{A}}_{r_\ell}(x^*) u_1$. Hence, if $x^* \in B_\ell$ is an equilibrium of some identical-payoffs game with utility coefficient vector $u = (u_1, u_2)$, the function $\tilde{\rho}_\ell$ permits us to recover u_2 given u_1 and x^* .

For each $\ell \in \mathbb{N}$, define the function $\rho_\ell : B_\ell \times \mathbb{R}^{K-\gamma} \rightarrow \mathbb{R}^K$ by

$$\rho_\ell(x, u_1) := (u_1, \tilde{\rho}_\ell(x, u_1)). \quad (32)$$

The function ρ_ℓ is a trivial extension of $\tilde{\rho}_\ell$ that recovers the full vector of utility coefficients $u \in \mathbb{R}^K$ given $u_1 \in \mathbb{R}^{K-\gamma}$ and an equilibrium $x^* \in B_\ell$.

We thus get the following corollary to Lemma 18.

Corollary 19. *There exists a countable collection of open balls $(B_\ell)_{\ell \geq 1}$, $B_\ell \subset X_C^*$ such that*

- (i) $\bigcup_{\ell \geq 1} B_\ell = \{x \in X : \text{carr}(x) = C\}$
- (ii) *For each $\ell \in \mathbb{N}$, there exists a differentiable function $\rho_\ell : B_\ell \times \mathbb{R}^{K-\gamma} \rightarrow \mathbb{R}^K$ such that, if $x^* \in B_\ell$ and x^* is an equilibrium of the game with utility coefficients $u = (u_1, u_2)$, $u_1 \in \mathbb{R}^{K-\gamma}$, $u_2 \in \mathbb{R}^\gamma$, then $u = \rho_\ell(x^*, u_1)$.*

We recall that our end goal is to use Sard's theorem to show that the set of second-order degenerate games has \mathcal{L}^K -measure zero. To that end, we will now show that for each $\ell \in \mathbb{N}$, the set of identical payoff games that have a second-order degenerate equilibrium lying in B_ℓ is contained in the critical values set of ρ_ℓ .

Suppose $x \in B_\ell$ and $u_1 \in \mathbb{R}^{K-\gamma}$ are arbitrary. If $u = (u_1, u_2)$ with $u_2 = \tilde{\rho}_\ell(x, u_1)$, then by the definition of $\tilde{\rho}_\ell$ we see that $\mathbf{A}(x)u = 0$. By the definition of $\mathbf{A}(x)$ (see (15)–(20)) this implies

$$F(x, u_1, \tilde{\rho}_\ell(x, u_1)) = 0, \quad \text{for all } x \in B_\ell.$$

¹⁴Note that we previously assumed a specific ordering for Y . However, this was for the purpose of proving Proposition 17, which is unaffected by a reordering of Y at this point.

Thus, taking a partial derivative with respect to x_i^k we get

$$\frac{\partial}{\partial x_i^k} F(x, u_1, \tilde{\rho}_\ell(x, u_1)) = 0, \quad i = 1, \dots, \tilde{N}, \quad k = 1, \dots, \gamma_i - 1. \quad (33)$$

Consider again the decomposition $x = (x_p, x_m)$. Using compact notation, (33) is restated as

$$D_{x_m} F(x_p, x_m, u_1, \tilde{\rho}_\ell(x_p, x_m, u_1)) = 0. \quad (34)$$

Suppose $x^* \in B_\ell$ is an equilibrium of an identical-payoffs game with utility coefficient vector u and $\text{carr}(x^*) = C$. Applying the chain rule in (34), using (16), and using the fact that $F(x, u) = \mathbf{A}(x)u = \tilde{\mathbf{A}}_{r_\ell}(x)u_1 + \mathbf{A}_{r_\ell}(x)u_2$, we find that at x^* there holds¹⁵

$$\tilde{\mathbf{H}}(x^*) = -\mathbf{A}_{r_\ell}(x^*)\tilde{\mathbf{J}}_{\rho_\ell}(x^*, u), \quad (35)$$

where $\tilde{\mathbf{J}}_{\rho_\ell}(x, u) := D_{\tilde{x}_m} \tilde{\rho}_\ell(x_p, \tilde{x}_m, u)|_{\tilde{x}_m=x_m}$ is the Jacobian of $\tilde{\rho}_\ell$ taken with respect to x_m .

Since $\mathbf{A}_{r_\ell}(x)$ is invertible for all $x \in B_\ell$, this means that given any equilibrium $x^* \in B_\ell$, the Hessian $\tilde{\mathbf{H}}(x^*)$ is nonsingular if and only if the Jacobian $\tilde{\mathbf{J}}_{\rho_\ell}(x^*, u)$ is nonsingular.

Note that the Jacobian of ρ_ℓ takes the form

$$\mathbf{J}_{\rho_\ell}(x, u) = \begin{pmatrix} \mathbf{0} & \mathbf{I} \\ \mathbf{J}_{\tilde{\rho}_\ell} & \mathbf{M} \end{pmatrix},$$

for some matrix \mathbf{M} . Clearly, $\det \mathbf{J}_{\rho_\ell} = 0$ if and only if $\det \tilde{\mathbf{J}}_{\rho_\ell} = 0$. Thus, by (35) we get the following result.

Lemma 20. *If $x^* \in B_\ell$ is an equilibrium of an identical-payoffs game with utility coefficient vector u , then*

$$\det \mathbf{J}_{\rho_\ell}(x^*, u) = 0 \quad \iff \quad \det \tilde{\mathbf{H}}(x^*) = 0, \quad (36)$$

where \mathbf{J}_{ρ_ℓ} is the Jacobian of ρ_ℓ .

¹⁵We note that $\tilde{\mathbf{H}}(x^*)$ is also dependent on u . However, we suppress the argument u since it is generally held constant in the context of the Hessian $\tilde{\mathbf{H}}$.

In particular, the above result shows that if u is an identical payoffs game having a second-order degenerate equilibrium $x^* \in B_\ell$, then u is contained in the set of critical values of ρ_ℓ . We now prove Proposition 13.

Proof. Let C be a carrier set. Let $\mathcal{U}(C) \subseteq \mathbb{R}^K$ be the set of identical-payoff games having at least one degenerate equilibrium with carrier set C ; that is,

$$\mathcal{U}(C) := \{u \in \mathbb{R}^K : \exists \text{ degenerate equilibrium } x^* \in \{x \in X : \text{carr}(x) = C\}\}.$$

Let $(B_\ell)_{\ell \in \mathbb{N}}$ and $(\rho_\ell)_{\ell \in \mathbb{N}}$ be as in Corollary 19. We note that B_ℓ and ρ_ℓ , $\ell \in \mathbb{N}$ are implicitly defined with respect to C . For $\ell \in \mathbb{N}$, let $\mathcal{U}(C, \ell) \subseteq \mathbb{R}^K$ be the subset of identical-payoff games having at least one degenerate equilibrium $x^* \in B_\ell$; that is,

$$\mathcal{U}(C, \ell) := \{u \in \mathbb{R}^K : \exists \text{ degenerate equilibrium } x^* \in B_\ell\}.$$

By construction, we have $\bigcup_{\ell \geq 1} B_\ell = \{x \in X : \text{carr}(x) = C\}$, and hence $\mathcal{U}(C) = \bigcup_{\ell \geq 1} \mathcal{U}(C, \ell)$.

By Lemma 20, the set $\mathcal{U}(C, \ell)$ is contained in the set of critical values of ρ_ℓ . By Sard's theorem, we get that $\mathcal{U}(C, \ell)$ is a set with \mathcal{L}^K -measure zero. Since $\mathcal{U}(C)$ is the countable union of sets of \mathcal{L}^K -measure zero, it is itself a set with \mathcal{L}^K -measure zero.

Let $\mathcal{U} \subset \mathbb{R}^K$ denote the subset of identical-payoff games with at least one degenerate equilibrium. The set \mathcal{U} may be expressed as the union $\mathcal{U} = \bigcup_C \mathcal{U}(C)$ taken over all possible carrier sets C . Since there are a finite number of carrier sets C , the set \mathcal{U} has \mathcal{L}^K -measure zero. \square

7. First-Order Degenerate Games

The following proposition shows that, within the set of identical-payoff games, first-order degenerate games form a null set.

Proposition 21. *The set of identical-payoff games which are first-order degenerate has \mathcal{L}^K -measure zero.*

Proof. Fix some set $C = C_1 \cup \dots \cup C_N$ where each C_i is a nonempty subset of Y_i . Let \widehat{C} be any *strict* subset of C . In the context of this proof let $\gamma_i := |C_i|$, let $\gamma := \sum_{i=1}^N (\gamma_i - 1)$, let $\widetilde{N} := |\{i = 1, \dots, N : \gamma_i \geq 2\}|$, and assume Y_i is reordered so that $\{y_i^1, \dots, y_i^{\gamma_i}\} = C_i$. Note that this ordering implies that for any x with $\text{carr}(x) = C$ we have $y_i^1 \in \text{carr}(x)$, $i = 1, \dots, N$.

Given an equilibrium x^* let the *extended carrier* of x^* be defined as

$$\text{ext carr}(x^*) := \text{carr}(x^*) \cup \left(\bigcup_{i=1}^N \{y_i^k \in Y_i : k = 2, \dots, K_i, \frac{\partial U(x^*)}{\partial x_i^{k-1}} = 0\} \right). \quad (37)$$

Suppose that x^* is an equilibrium with *extended carrier* C . By Lemma 30 (see appendix) and the ordering we assumed for Y_i we have

$$F_i^k(x^*, u) = \frac{\partial U(x^*)}{\partial x_i^{k-1}} = 0, \quad i = 1, \dots, \tilde{N}, \quad k = 1, \dots, \gamma_i - 1, \quad (38)$$

where F_i^k is as defined in (14). Thus, if x^* is an equilibrium for some identical-payoffs game with utility coefficient vector $u \in \mathbb{R}^K$, and $\text{ext carr}(x^*) = C$, then by the definition of $\mathbf{A}(x)$ (see (15)–(20)), (38) implies that $\mathbf{A}(x^*)u = 0$, or equivalently,

$$u \in \ker \mathbf{A}(x^*),$$

where the matrix $\mathbf{A}(x) \in \mathbb{R}^{\gamma \times K}$ is defined with respect to C , as in (20).

Let $\mathcal{U}(C, \widehat{C}) \subseteq \mathbb{R}^K$ be the set of identical-payoff games in which there exists an equilibrium x^* with $\text{carr}(x^*) = \widehat{C}$ and $\text{ext carr}(x^*) = C$. Let

$$\widehat{X} := \{x \in X : \text{carr}(x) = \widehat{C}\}.$$

By the above we see that

$$\mathcal{U}(C, \widehat{C}) \subseteq \bigcup_{x \in \widehat{X}} \ker \mathbf{A}(x). \quad (39)$$

For each $x \in \widehat{X}$, let $\text{range } \mathbf{A}(x)^T$ denote the range space of $\mathbf{A}(x)^T$. Each entry of $\mathbf{A}(x)^T$ is a polynomial function in x and hence is Lipschitz continuous over the bounded set \widehat{X} . By Proposition 17 we have $\text{rank } \mathbf{A}(x)^T = \gamma$ for all $x \in \widehat{X}$. Thus, we may choose a set of γ basis vectors $\{\mathbf{b}_1(x), \dots, \mathbf{b}_\gamma(x)\}$ spanning $\text{range } \mathbf{A}(x)^T$ such that each $\mathbf{b}_k(x) \in \mathbb{R}^K$, $k = 1, \dots, \gamma$ is a Lipschitz continuous function in x . Moreover, we may choose a complementary set of $(K - \gamma)$ linearly independent vectors $\{\mathbf{b}_{\gamma+1}(x), \dots, \mathbf{b}_K(x)\}$ forming a basis for the orthogonal complement $(\text{range } \mathbf{A}(x)^T)^\perp$, with each $\mathbf{b}_k(x) \in \mathbb{R}^K$, $k = \gamma + 1, \dots, K$ being a continuous function in x . Let $\mathbf{B}(x) := (\mathbf{b}_{\gamma+1}(x) \cdots \mathbf{b}_K(x)) \in \mathbb{R}^{K \times (K-\gamma)}$. Let $f : \widehat{X} \times \mathbb{R}^{K-\gamma} \rightarrow \mathbb{R}^K$ be given by $f(x, v) := \mathbf{B}(x)v$. Since $\mathbf{B}(x)$ is Lipschitz continuous in x and \widehat{X} is bounded,

f is Lipschitz continuous. By the fundamental theorem of linear algebra, for each $x \in \widehat{X}$, $\ker \mathbf{A}(x) = (\text{range } \mathbf{A}(x)^T)^\perp = \text{range } \mathbf{B}(x)$. Hence,

$$\bigcup_{x \in \widehat{X}} \ker \mathbf{A}(x) = f(\widehat{X} \times \mathbb{R}^{K-\gamma}). \quad (40)$$

Since $\widehat{C} \subsetneq C$, the Hausdorff dimension of \widehat{X} is at most $(\gamma - 1)$ and the Hausdorff dimension of $\widehat{X} \times \mathbb{R}^{K-\gamma}$ is at most $K - 1$. Since f is Lipschitz continuous, this implies (see [45], Section 2.4) that the Hausdorff dimension of $f(\widehat{X} \times \mathbb{R}^{K-\gamma})$ is at most $K - 1$, and in particular, that $f(\widehat{X} \times \mathbb{R}^{K-\gamma})$ has \mathcal{L}^K -measure zero. By (39) and (40), this implies that $\mathcal{U}(C, \widehat{C})$ has \mathcal{L}^K -measure zero.

Let $\mathcal{U} \subseteq \mathbb{R}^K$ denote the set of all identical-payoff games containing a first-order degenerate equilibrium. Since we may represent this set as a finite union of \mathcal{L}^K -measure zero sets,

$$\mathcal{U} = \bigcup_{\substack{\emptyset \neq C_i \subseteq Y_i, \ i=1, \dots, N \\ C = C_1 \cup \dots \cup C_N \\ \widehat{C} \subsetneq C}} \mathcal{U}(C, \widehat{C}),$$

the set \mathcal{U} also has \mathcal{L}^K -measure zero. □

8. Regularity in Exact and Weighted Potential Games

The goal of this section will be to prove the following proposition.

Proposition 22.

- (i) *Almost all weighted potential games are regular.*
- (ii) *Almost all exact potential games are regular.*

Exact and weighted potential games are closely related to games with identical payoffs. By identifying equivalence relationships between identical-payoff games and exact and weighted potential games, this result follows as a relatively simple consequence of Proposition 3.

Let the number of players $N \geq 2$ and action-space size $K_i \geq 2$, $i = 1, \dots, N$ be arbitrary. Define

$$\tilde{\mathcal{I}} := \{u \in \mathcal{I} : \mathbf{1}^T u = 0\},$$

where $\mathcal{I} = \mathbb{R}^K$ is the set of identical-payoff games as given Section 2.3.

The set $\tilde{\mathcal{I}}$ will provide a convenient means of partitioning the sets of exact and weighted potential games.

Note that an identical-payoffs game $u \in \tilde{\mathcal{I}}$ is regular if and only if $\tilde{u} = u + c\mathbf{1}$ is regular for every $c \in \mathbb{R}$. Note also that $\tilde{\mathcal{I}}$ is isomorphic to \mathbb{R}^{K-1} . (Henceforth we will treat $\tilde{\mathcal{I}}$ as $\tilde{\mathcal{I}} = \mathbb{R}^{K-1}$.) This, along with Proposition 3, implies the following lemma.

Lemma 23. *The set of games in $\tilde{\mathcal{I}}$ that are irregular has \mathcal{L}^{K-1} -measure zero.*

The sets of weighted and exact potential games (of size $(N, (K_i)_{i=1}^N)$) are defined explicitly as follows. For convenience, we decompose the vector of utility coefficients $u \in \mathbb{R}^{NK}$ as $u = (u_i)_{i=1}^N$, where $u_i \in \mathbb{R}^K$ gives the pure strategy utility received by player i for each pure strategy $y \in Y$. The set of weighted potential games is given by

$$\mathcal{W} := \left\{ (u_i)_{i=1}^N \in \mathbb{R}^{NK} : \begin{array}{l} (6) \text{ holds for some } u \in \mathbb{R}^K \text{ and some } (w_i)_{i=1}^N, \\ w_i > 0, i = 1, \dots, N \end{array} \right\}.$$

the set of exact potential games is given by

$$\mathcal{P} := \left\{ (u_i)_{i=1}^N \in \mathbb{R}^{NK} : (6) \text{ holds for some } u \in \mathbb{R}^K \text{ with } (w_i)_{i=1}^N = \mathbf{1} \right\},$$

where $\mathbf{1}$ denotes the vector of all ones.

A element $v \in \mathcal{P}$ or $v \in \mathcal{W}$ is a vector in \mathbb{R}^{NK} which we decompose as $v = (v_i)_{i=1}^N$ where $v_i \in \mathbb{R}^K$ represent the pure-strategy utility of player i . For each $u \in \tilde{\mathcal{I}}$, define

$$\mathcal{P}_u := \left\{ v \in \mathbb{R}^{NK} : \begin{array}{l} v_i(y_i, y_{-i}) - v_i(y'_i, y_{-i}) = u(y_i, y_{-i}) - u(y'_i, y_{-i}) \\ \forall i = 1, \dots, N, y_i, y'_i \in Y_i, y_{-i} \in Y_{-i} \end{array} \right\};$$

to be the set of all potential games with potential function $u \in \tilde{\mathcal{I}}$. Using the definition of an exact potential game (see Section 2.2), it is straightforward to verify that $\mathcal{P}_u \cap \mathcal{P}_{\tilde{u}} = \emptyset$ for every $u, \tilde{u} \in \tilde{\mathcal{I}}$, $u \neq \tilde{u}$ and $\bigcup_{u \in \tilde{\mathcal{I}}} \mathcal{P}_u = \mathcal{P}$. Thus $(\mathcal{P}_u)_{u \in \tilde{\mathcal{I}}}$ partitions \mathcal{P} . Likewise, define

$$\mathcal{W}_u := \left\{ v \in \mathbb{R}^{NK} : \begin{array}{l} v_i(y_i, y_{-i}) - v_i(y'_i, y_{-i}) = w_i(u(y_i, y_{-i}) - u(y'_i, y_{-i})) \\ \forall i = 1, \dots, N, y_i, y'_i \in Y_i, y_{-i} \in Y_{-i}, \text{ and for some } w_i > 0 \end{array} \right\}$$

to be set of all weighted potential games with potential function $u \in \tilde{\mathcal{I}}$. As in the case of exact potential games, we see that $(\mathcal{W}_u)_{u \in \tilde{\mathcal{I}}}$ partitions \mathcal{W} .

An equilibrium in a potential game is regular if and only if it is a regular equilibrium in the associated identical payoffs game. Thus, we get the following result.

Lemma 24.

(i) For each $u \in \tilde{\mathcal{I}}$ and each exact potential game $v \in \mathcal{P}_u$, v is regular if and only if u is regular.

(ii) For each $u \in \tilde{\mathcal{I}}$ and each weighted potential game $w \in \mathcal{W}_u$, w is regular if and only if u is regular.

Recall that \mathcal{P} and \mathcal{W} are subspaces of \mathbb{R}^{NK} with dimensions K_p and K_w , respectively (see Section 2.3). Since $(\mathcal{P}_u)_{u \in \tilde{\mathcal{I}}}$ partitions \mathcal{P} , every exact potential game $v \in \mathcal{P}$ may be uniquely represented by a vector (u, z) where $u \in \tilde{\mathcal{I}} = \mathbb{R}^{K-1}$ and $z \in \mathbb{R}^{\tilde{K}_p}$, $\tilde{K}_p := K_p - K + 1$. Likewise, every weighted potential game $v \in \mathcal{W}$ may be uniquely represented by a vector (u, z) where $u \in \tilde{\mathcal{I}} = \mathbb{R}^{K-1}$ and $z \in \mathbb{R}^{\tilde{K}_w}$, $\tilde{K}_w := K_w - K + 1$.

Let $\text{IR}_p \subset \mathcal{P}$ and $\text{IR}_w \subset \mathcal{W}$ denote the subsets of the exact and weighted potential games (respectively) that are irregular. Recall that by Remark 5, IR_p and IR_w are closed and hence measurable. Since $\mathcal{L}^{K-1}(\tilde{\mathcal{I}}) = 0$ we see that $\mathcal{L}^{K_p}(\text{IR}_p) = \mathcal{L}^{K_p}(\tilde{\mathcal{I}} \times \mathbb{R}^{K_p-K+1}) = \mathcal{L}^{K-1}(\tilde{\mathcal{I}}) \mathcal{L}^{K_p-K+1}(\mathbb{R}^{K_p-K+1}) = 0$. By similar reasoning we also see that $\mathcal{L}^{K_w}(\text{IR}_w) = 0$. Since IR_p and IR_w are closed with respect to \mathcal{P} and \mathcal{W} (see Remark 5), this proves Proposition 22.

9. Conclusion

Regular NE are isolated, robust, and have a simple analytic structure. Within the subclass of potential games, regular equilibria have a simple and intuitive meaning in terms of the potential function (see Section 3). The paper showed that almost all weighted potential games, almost all exact potential games, and almost all games with identical payoffs are regular. This result implies that properties holding in regular potential games are inherently robust to payoff perturbations and properties only holding in irregular games are inherently non-robust to payoff perturbations.

Regular potential games can be particularly useful in the study of game-theoretic learning processes. As an example application, in a companion work [29] the authors study BR dynamics in potential games. Regular potential games greatly facilitate the analysis of BR dynamics. Moreover, the results of the present paper allow one to immediately conclude that results obtained

for BR dynamics in [29] hold for almost all potential games, thus addressing open questions regarding the convergence rate and limit set composition of BR dynamics [35, 29, 46, 32]. As another example, [30] studies a form of no-regret learning in potential games. It is shown that if the potential game is regular, then convergence to NE and convergence rate estimates can be established. The present work allows one to immediately conclude that the results of [30] hold for almost all potential games. An important future research direction is to study the application of regular potential games in facilitating the analysis of additional game-theoretic learning processes.

Appendix A.

We use the following standard mathematical notation:

- $\mathbb{N} := \{1, 2, \dots\}$.
- The mapping $\text{sgn} : \mathbb{R}^{n \times m} \rightarrow \mathbb{R}^{n \times m}$ is given by

$$(\text{sgn}(\mathbf{A}))_{i,j} := \begin{cases} 1 & \text{if } a_{i,j} > 0 \\ -1 & \text{if } a_{i,j} < 0 \\ 0 & \text{if } a_{i,j} = 0. \end{cases}$$

- Given two matrices \mathbf{A} and \mathbf{B} of the same dimension, $\mathbf{A} \circ \mathbf{B}$ denotes the Hadamard product (i.e., the entrywise product) of \mathbf{A} and \mathbf{B} .
- Given a set of matrices (or scalars) $\{\mathbf{A}_1, \dots, \mathbf{A}_n\}$, possibly of differing dimensions, the notation $\text{diag}(\mathbf{A}_1, \dots, \mathbf{A}_n)$ denotes the associated block diagonal matrix.
- Suppose $m, n, p \in \mathbb{N}$, $F_i : \mathbb{R}^m \times \mathbb{R}^n \rightarrow \mathbb{R}$, for $i = 1, \dots, p$. Suppose further that $F : \mathbb{R}^m \times \mathbb{R}^n \rightarrow \mathbb{R}^p$ is given by $F(w, z) = (F_i(w, z))_{i=1, \dots, p}$. Then the operator D_w gives the Jacobian of F with respect to the components of $w = (w_k)_{k=1, \dots, m}$; that is

$$D_w F(w, z) = \begin{pmatrix} \frac{\partial F_1(w, z)}{\partial w_1} & \dots & \frac{\partial F_1(w, z)}{\partial w_m} \\ \vdots & \ddots & \vdots \\ \frac{\partial F_p(w, z)}{\partial w_1} & \dots & \frac{\partial F_p(w, z)}{\partial w_m} \end{pmatrix}.$$

- A^c denotes the complement of a set A , and $\overset{\circ}{A}$ denotes the interior of A , and $\text{cl } A$ denotes the closure of A .

- Given a function f , $\mathcal{D}(f)$ refers to the domain of f and $\mathcal{R}(f)$ to the range of f .
- \mathcal{L}^n , $n \in \{1, 2, \dots\}$ refers to the n -dimensional Lebesgue measure.

Appendix B.

This appendix provides the proof of Proposition 17. We will give a short roadmap of the proof before proceeding to the proof itself.

Roadmap of Proof of Proposition 17.

1. We begin by reorganizing the columns of $\mathbf{A}(x)$ and partitioning the matrix as $\mathbf{A}(x) = [\mathbf{A}_1(x) \ \mathbf{A}_2(x)]$, where $\mathbf{A}_1(x)$ has dimensions $\gamma \times \tilde{K}$ with $\tilde{K} \geq \gamma$ (see (B.3)). Since our goal is to prove that $\text{rank}(\mathbf{A}(x)) = \gamma$ it will be sufficient to prove that $\mathbf{A}_1(x)$ has full rank.
2. We will prove Proposition 17 by considering the sign pattern of $\mathbf{A}_1(x)$ (i.e., the matrix $\text{sgn}(\mathbf{A}_1(x))$), where $\text{sgn}(\cdot)$ is defined in Appendix A). To do this, we will leverage known results about sign pattern matrices (i.e., matrices with entries in $\{-1, 0, 1\}$). We will use the following key notion from the theory of sign pattern matrices (see Definition 25): A sign pattern matrix \mathbf{M} is called an L -matrix if every matrix with the same sign pattern as \mathbf{M} has full rank. Using this notion, proving Proposition 17 is equivalent to showing that for any x with $\text{carr}(x) \subseteq C$, $\text{sgn}(\mathbf{A}_1(x))$ is an L -matrix (see Lemma 27). Thus, we will focus our efforts on proving Lemma 27.
3. A convenient characterization of L -matrices known from the literature is given in Lemma 26. We will prove Lemma 27 by directly verifying that it satisfies the L -matrix condition of Lemma 26. This verification process will be simplified by breaking $\mathbf{A}_1(x)$ into component matrices \mathbf{R}_1 and $\mathbf{P}_1(x)$, where \mathbf{R}_1 is a sign pattern matrix independent of x and $\mathbf{P}_1(x)$ is a non-negative matrix depending on x . The decomposition of $\mathbf{A}_1(x)$ into \mathbf{R}_1 and $\mathbf{P}_1(x)$ will be carried out early on in (B.2) before proceeding to the rest of the proof.

We now proceed, following the above roadmap. We begin by assuming a convenient ordering of elements in Y . Let

$$\tilde{K} := \prod_{i=1}^N \gamma_i.$$

For $\tau = 1, \dots, \tilde{K}$, let $\alpha_\tau = (\alpha_\tau^1, \dots, \alpha_\tau^{\tilde{N}})$ be a multi-index associated with the τ -th action tuple in Y , meaning that

$$y^\tau = (y_1^{\alpha_\tau^1}, \dots, y_{\tilde{N}}^{\alpha_\tau^{\tilde{N}}}, y_{\tilde{N}+1}^1, \dots, y_N^1),$$

where y_i^1 , for $i = \tilde{N} + 1, \dots, N$, is, by construction, the pure strategy used by player i at any strategy x with carrier C . Let Y be reordered so that for every $\tau = 1, \dots, \tilde{K}$, $i = 1, \dots, \tilde{N}$ we have $1 \leq \alpha_\tau^i \leq \gamma_i$. This ensures that the first \tilde{K} strategies in Y contain all strategy combinations of the elements of C_1, \dots, C_N , and that for any multi-index $\alpha = (\alpha^1, \dots, \alpha^{\tilde{N}})$, $1 \leq \alpha^i \leq \gamma_i$, there exists a unique $1 \leq \tau \leq \tilde{K}$ such that $y^\tau = (y_1^{\alpha^1}, \dots, y_{\tilde{N}}^{\alpha^{\tilde{N}}}, y_{\tilde{N}+1}^1, \dots, y_N^1)$.

By definition (11) and the ordering we assumed for Y we have that

$$q_i^\tau(y_i^k) = \begin{cases} 1 & \text{if } k = \alpha_\tau^i \\ 0 & \text{otherwise.} \end{cases} \quad (\text{B.1})$$

For $1 \leq s \leq \gamma$ and $1 \leq \tau \leq \tilde{K}$, let

$$r_{s,\tau} := q_i^\tau(y_i^{k+1}) - q_i^\tau(y_i^1) \quad \text{and} \quad p_{s,\tau}(x) := \prod_{j \neq i} q_j^\tau(x_j), \quad (\text{B.2})$$

where $i = i^*(s)$ and $k = k^*(s)$ (see (19)), and note that $a_{s,\tau}(x) = r_{s,\tau} p_{s,\tau}(x)$ (see (20)). We may write $\mathbf{A}(x) = \mathbf{R} \circ \mathbf{P}(x)$, where \circ is the Hadamard product, and \mathbf{R} and $\mathbf{P}(x)$ have entries $r_{s,\tau}$ and $p_{s,\tau}(x)$ respectively.

Partition $\mathbf{A}(x)$, \mathbf{R} and $\mathbf{P}(x)$ as $\mathbf{A}(x) = [\mathbf{A}_1(x) \ \mathbf{A}_2(x)]$, $\mathbf{R} = [\mathbf{R}_1 \ \mathbf{R}_2]$, and $\mathbf{P}(x) = [\mathbf{P}_1(x) \ \mathbf{P}_2(x)]$, where $\mathbf{A}_1(x), \mathbf{R}_1, \mathbf{P}_1(x) \in \mathbb{R}^{\gamma \times \tilde{K}}$ and $\mathbf{A}_2(x), \mathbf{R}_2, \mathbf{P}_2(x) \in \mathbb{R}^{\gamma \times (K - \tilde{K})}$, so we may write

$$\mathbf{A}(x) = [\mathbf{A}_1(x) \ \mathbf{A}_2(x)], \quad \text{with,} \quad \mathbf{A}_1(x) = \mathbf{R}_1 \circ \mathbf{P}_1(x) \quad \text{and} \quad \mathbf{A}_2(x) = \mathbf{R}_2 \circ \mathbf{P}_2(x). \quad (\text{B.3})$$

In order to show that $\mathbf{A}(x)$ has full row rank, it is sufficient to prove that $\mathbf{A}_1(x)$ has full row rank—this is the approach we will take in proving the proposition.

We address this by studying the sign pattern of $\mathbf{A}_1(x)$. Properties of *sign pattern matrices* (i.e., matrices with entries in $\{-1, 0, 1\}$) have been well-studied [43, 44]. We recall the following definition from [43],

Definition 25. A sign pattern matrix $\mathbf{L} \in \mathbb{R}^{m \times n}$ with $n \geq m$ is said to be an *L-matrix* if for every matrix \mathbf{M} with $\text{sgn}(\mathbf{M}) = \text{sgn}(\mathbf{L})$, the matrix \mathbf{M} has full row rank.

The following lemma characterizes *L-matrices* [43, 44].

Lemma 26. *Let $\mathbf{L} \in \mathbb{R}^{m \times n}$ be a sign pattern matrix with $n \geq m$. Then \mathbf{L} is an L -matrix if and only if for every diagonal sign pattern matrix $\mathbf{D} \in \mathbb{R}^{m \times m}$, $\mathbf{D} \neq 0$ there is a nonzero column of \mathbf{DL} in which each nonzero entry has the same sign.*

In light of Definition 25, Proposition 17 is equivalent to the following lemma.

Lemma 27. *For any x such that $\text{carr}(x) \subseteq C$, the matrix $\mathbf{A}_1(x) = (\mathbf{R}_1 \circ \text{sgn}(\mathbf{P}_1(x)))$ is an L -matrix.*

The proof of this lemma relies on showing that $(\mathbf{R}_1 \circ \text{sgn}(\mathbf{P}_1(x)))$ satisfies the L -matrix characterization given in Lemma 26.

Before proving Lemma 27 we introduce some definitions that will be useful in the proof.

Given a diagonal matrix $\mathbf{D} \in \mathbb{R}^{\ell \times \ell}$, $\ell \in \mathbb{N}$, let $\text{diag}(\mathbf{D})$ be the vector in \mathbb{R}^ℓ containing the diagonal elements of \mathbf{D} .

Given a diagonal sign pattern matrix $\mathbf{D} \in \mathbb{R}^{\ell \times \ell}$, $\ell \in \mathbb{N}$, define $\text{idx}(\mathbf{D})$ as follows. If $\text{diag}(\mathbf{D})$ does not contain any ones, then let $\text{idx}(\mathbf{D}) = 1$. Otherwise, let $\text{idx}(\mathbf{D})$ be one more than the first index in $\text{diag}(\mathbf{D})$ containing a 1.¹⁶¹⁷

Given a diagonal matrix $\mathbf{D} \in \mathbb{R}^{\gamma \times \gamma}$, let $\mathbf{D}_i \in \mathbb{R}^{(\gamma_i - 1) \times (\gamma_i - 1)}$, $i = 1, \dots, \tilde{N}$ be the (unique) diagonal matrices satisfying

$$\text{diag}(\mathbf{D}) = (\text{diag}(\mathbf{D}_1), \dots, \text{diag}(\mathbf{D}_N)).$$

We now prove Lemma 27.

Proof. Let x be a strategy satisfying $\text{carr}(x) \subseteq C$. In order to show that $(\mathbf{R}_1 \circ \text{sgn}(\mathbf{P}_1(x)))$ is an L -matrix, it is sufficient (by Lemma 26) to show that and for any diagonal sign pattern matrix $\mathbf{D} \neq 0$, there exists a column of $\mathbf{D}(\mathbf{R}_1 \circ \text{sgn}(\mathbf{P}_1(x)))$ which is nonzero and in which every nonzero entry has the same sign. With this in mind, we begin by giving a characterization of the columns of \mathbf{R}_1 .

Suppose that $i = 1, \dots, \tilde{N}$ and $k = 1, \dots, \gamma_i - 1$ are fixed. Note the following:

- (i) Suppose $\tau \in \{1, \dots, \tilde{K}\}$ is such that $\alpha_\tau^i = k + 1$, where α_τ^i is the i -th index of the multi-index α_τ . Since α_τ is used to define the ordering of actions in Y , we have $q_i^\tau(y_i^{k+1}) = 1$ and $q_i^\tau(y_i^1) = 0$ (see (B.1) and preceding discussion). Hence, $q_i^\tau(y_i^{k+1}) - q_i^\tau(y_i^1) = 1$.

¹⁶Assume indexing starts with one, not zero. For example, if the first time a 1 appears in $\text{diag}(\mathbf{D})$ is at index 2, then $\text{idx}(\mathbf{D}) = 3$.

¹⁷The awkward offset in this definition is needed in order to handle the indexing offset inherent in the definition of X_i , as discussed in Section 2.1.

- (ii) Suppose $\tau \in \{1, \dots, \tilde{K}\}$ is such that $\alpha_\tau^i = 1$. Then $q_i^\tau(y_i^{k+1}) = 0$, and $q_i^\tau(y_i^1) = 1$. Hence, $q_i^\tau(y_i^{k+1}) - q_i^\tau(y_i^1) = -1$.
- (iii) For all other $\tau \in \{1, \dots, \tilde{K}\}$ we have $q_i^\tau(y_i^{k+1}) = 0$, and $q_i^\tau(y_i^1) = 0$, and hence $q_i^\tau(y_i^{k+1}) - q_i^\tau(y_i^1) = 0$.

For $1 \leq \tau \leq \tilde{K}$, let $\mathbf{r}_\tau \in \mathbb{R}^\gamma$ be the τ -th column of \mathbf{R}_1 . Partition this column as

$$\mathbf{r}_\tau = \begin{pmatrix} \mathbf{r}_\tau^1 \\ \vdots \\ \mathbf{r}_\tau^{\tilde{N}} \end{pmatrix}.$$

where $\mathbf{r}_\tau^i \in \mathbb{R}^{\gamma_i-1}$. From (B.2) we see that

$$\mathbf{r}_\tau^i = \begin{pmatrix} q_i^\tau(y_i^2) - q_i^\tau(y_i^1) \\ \vdots \\ q_i^\tau(y_i^{\gamma_i}) - q_i^\tau(y_i^1) \end{pmatrix}.$$

Given the observations (i)–(iii) above we see that for each i we have

$$\mathbf{r}_\tau^i = \begin{cases} -\mathbf{1} & \text{if } \alpha_\tau^i = 1 \\ e_{\alpha_\tau^i-1} & \text{if } 2 \leq \alpha_\tau^i \leq \gamma_i, \end{cases} \quad (\text{B.4})$$

where the symbol $e_{\alpha_\tau^i-1}$ refers to the $(\alpha_\tau^i - 1)$ -th canonical vector in \mathbb{R}^{γ_i-1} and $\mathbf{1} \in \mathbb{R}^{\gamma_i-1}$ is the vector of all ones.

We now characterize the columns of $\text{sgn}(\mathbf{P}_1(x))$. For $i = 1, \dots, \tilde{N}$ we define

$$\text{idx}_i(x) := \{k \in \{1, \dots, \gamma_i\} : T_i^k(x_i) > 0\}.$$

The function $\text{idx}_i(x)$ simply gives the indices of the pure strategies of player i that receive positive weight under x . Since $\text{carr}(x) = C$, the ordering we assumed for Y_i implies that $T_i^k(x) = 0$ for $k \geq \gamma_i + 1$. By the definition of T_i^k , it is not possible to have $T_i^k(x_i) = 0$ for all $k = 1, \dots, \gamma_i$ and hence $\text{idx}_i(x) \neq \emptyset$.

Let \mathbf{p}_τ be the τ -th column of $\mathbf{P}_1(x)$ and let

$$\tilde{\mathbf{p}}_\tau := \text{sgn}(\mathbf{p}_\tau)$$

be the τ -th column of $\text{sgn}(\mathbf{P}_1(x))$.¹⁸ Suppose that $\tau \in \{1, \dots, \tilde{K}\}$ is such that for the multi-index α_τ we have $\alpha_\tau^i \in \text{idx}_i(x)$ for all $i = 1, \dots, \tilde{N}$. Then for each

¹⁸For ease of notation we suppress the argument x when writing the columns of these matrices.

$s = 1, \dots, \gamma$ the (s, τ) -th entry of $\mathbf{P}_1(x)$ is strictly positive (see (11) and (B.2)), and hence \mathbf{p}_τ is positive and $\tilde{\mathbf{p}}_\tau = \mathbf{1}$.

Partition the columns \mathbf{p}_τ and $\tilde{\mathbf{p}}_\tau$ as

$$\mathbf{p}_\tau = \begin{pmatrix} \mathbf{p}_\tau^1 \\ \vdots \\ \mathbf{p}_\tau^{\tilde{N}} \end{pmatrix} \quad \text{and} \quad \tilde{\mathbf{p}}_\tau = \begin{pmatrix} \tilde{\mathbf{p}}_\tau^1 \\ \vdots \\ \tilde{\mathbf{p}}_\tau^{\tilde{N}} \end{pmatrix}, \quad (\text{B.5})$$

where $\mathbf{p}_\tau^i, \tilde{\mathbf{p}}_\tau^i \in \mathbb{R}^{\gamma_i-1}$. Suppose that τ is such that for the multi-index α_τ we have $\alpha_\tau^i \notin \text{idx}_i(x)$ for exactly one subindex $i \in \{1, \dots, \tilde{N}\}$. Then \mathbf{p}_τ^i is positive (see (B.2)) and \mathbf{p}_τ^j is zero for any $j \neq i$. Hence, $\tilde{\mathbf{p}}_\tau^i = \mathbf{1}$ and $\tilde{\mathbf{p}}_\tau^j = 0$ for any $j \neq i$.

Now, let $\mathbf{D} \in \mathbb{R}^{\gamma \times \gamma}$ be a non-zero diagonal sign pattern matrix. We will show that there is a nonzero column of $\mathbf{D}(\mathbf{R}_1 \circ \text{sgn}(\mathbf{P}_1(x)))$ in which each nonzero entry is a 1.

We now consider two possible cases for the structure of \mathbf{D} and show that in each case there is a nonzero column of $\mathbf{D}(\mathbf{R}_1 \circ \text{sgn}(\mathbf{P}_1(x)))$ in which every nonzero entry is 1.

Case 1: Suppose that for all $i \in \{1, \dots, \tilde{N}\}$ such that $\text{idx}(\mathbf{D}_i) \notin \text{idx}_i(x)$ we have $\text{diag}(\mathbf{D}_i) = 0$. Choose τ such that

$$\begin{cases} \alpha_\tau^i = \text{idx}(\mathbf{D}_i) & \text{if } \text{idx}(\mathbf{D}_i) \in \text{idx}_i(x), \\ \alpha_\tau^i \in \text{idx}_i(x) & \text{if } \text{idx}(\mathbf{D}_i) \notin \text{idx}_i(x). \end{cases}$$

Note that $\alpha_\tau^i \in \text{idx}_i(x)$ for all $i = 1, \dots, \tilde{N}$, and hence $\tilde{\mathbf{p}}_\tau = \mathbf{1}$ (see discussion preceding (B.5)). The τ -th column of $\mathbf{D}(\mathbf{R}_1 \circ \text{sgn}(\mathbf{P}_1(x)))$ is given by

$$\mathbf{D}(\mathbf{r}_\tau \circ \tilde{\mathbf{p}}_\tau) = \text{diag}(\mathbf{D}) \circ \mathbf{r}_\tau = (\text{diag}(\mathbf{D}_i) \circ \mathbf{r}_\tau^i)_{i=1}^{\tilde{N}}. \quad (\text{B.6})$$

For any i such that $\text{idx}(\mathbf{D}_i) \notin \text{idx}_i(x)$ we have, by assumption, $\text{diag}(\mathbf{D}_i) = 0$ and hence $\text{diag}(\mathbf{D}_i) \circ \mathbf{r}_\tau^i = 0$. Moreover, note that in this case we have $\text{idx}(\mathbf{D}_i) = 1$ since, by the definition of $\text{idx}(\cdot)$, $\text{diag}(\mathbf{D}_i) = 0$ implies $\text{idx}(\mathbf{D}_i) = 1$.

Suppose now that i is such that $\text{idx}(\mathbf{D}_i) \in \text{idx}_i(x)$. For $i = 1, \dots, \tilde{N}$, if $\alpha_\tau^i = 1$ then $\mathbf{r}_\tau^i = -\mathbf{1}$ (by (B.4)) and $\text{diag}(\mathbf{D}_i)$ contains no ones (this is the definition of $\text{idx}(\mathbf{D}_i) = 1$). In fact, $\text{diag}(\mathbf{D}_i)$ contains only entries with value of 0 or -1 . Hence, $\text{diag}(\mathbf{D}_i) \circ \mathbf{r}_\tau^i = -\text{diag}(\mathbf{D}_i)$, which is a nonnegative vector.

If $2 \leq \alpha_\tau^i \leq \gamma_i$ then $\mathbf{r}_\tau^i = e_{\alpha_\tau^i-1}$ (by (B.4)). Recalling the definition of $\text{idx}(\cdot)$, by our choice of $\alpha_\tau^i = \text{idx}(\mathbf{D}_i)$, the $(\alpha_\tau^i - 1)$ -th entry of $\text{diag}(\mathbf{D}_i)$ is 1. Hence, $\text{diag}(\mathbf{D}_i) \circ \mathbf{r}_\tau^i = e_{\alpha_\tau^i-1}$. In particular, this implies that if $\alpha_\tau^i \neq 1$ then $\text{diag}(\mathbf{D}_i) \circ \mathbf{r}_\tau^i$ is not identically zero and every nonzero entry of $\text{diag}(\mathbf{D}_i) \circ \mathbf{r}_\tau^i$ is 1.

In summary, for $i = 1, \dots, \tilde{N}$, we have $\text{diag}(\mathbf{D}_i) \circ \mathbf{r}_\tau^i \geq 0$, with equality only when $\text{idx}(\mathbf{D}_i) = 1$ and $\mathbf{D}_i = 0$. Hence, by (B.6), the τ -th column of $\mathbf{D}(\mathbf{R}_1 \circ$

$\text{sgn}(\mathbf{P}_1(x))$) satisfies $\mathbf{D}(\mathbf{r}_\tau \circ \tilde{\mathbf{p}}_\tau) \geq 0$, with equality only when $\text{idx}(\mathbf{D}_i) = 1$ and $\mathbf{D}_i = 0$ for all i . But, by assumption $\mathbf{D} \neq 0$, so $\mathbf{D}(\mathbf{r}_\tau \circ \tilde{\mathbf{p}}_\tau) \neq 0$.

Case 2: Suppose that for some $i \in \{1, \dots, \tilde{N}\}$ we have $\text{idx}(\mathbf{D}_i) \notin \text{idx}_i(x)$ and $\text{diag}(\mathbf{D}_i) \neq 0$. Let $\tau \in \{1, \dots, \tilde{K}\}$ be chosen such that $\alpha_\tau^i = \text{idx}(\mathbf{D}_i)$ for exactly one such $i \in \{1, \dots, \tilde{N}\}$ and for all other $j \neq i$ we have $\alpha_\tau^j \in \text{idx}_j$ (this is always possible since $\text{idx}_j \neq \emptyset$). Then we have

$$\tilde{\mathbf{p}}_\tau^i = \mathbf{1}, \quad \text{and} \quad \tilde{\mathbf{p}}_\tau^j = 0, \quad \text{for all } j \neq i$$

(see discussion following (B.5)).

As shown in Case 1, if $\alpha_\tau^i = 1$, then $\mathbf{D}_i \leq 0$ and $r_\tau^i = -1$ which implies that $\text{diag}(\mathbf{D}_i) \circ \mathbf{r}_\tau^i \geq 0$. Moreover, since $\tilde{\mathbf{p}}_\tau^i = \mathbf{1}$ and since, by assumption $\text{diag}(\mathbf{D}_i) \neq 0$ we have $\text{diag}(\mathbf{D}_i) \circ \mathbf{r}_\tau^i \circ \tilde{\mathbf{p}}_\tau^i \neq 0$ and every nonzero entry is 1.

If $2 \leq \alpha_\tau^i \leq \gamma_i$, then, again using the same reasoning as in Case 1, we see that $\text{diag}(\mathbf{D}_i) \circ \mathbf{r}_\tau^i = e_{\alpha_\tau^i - 1}$. Since $\tilde{\mathbf{p}}_\tau^i = \mathbf{1}$ we get that $\text{diag}(\mathbf{D}_i) \circ \mathbf{r}_\tau^i \circ \tilde{\mathbf{p}}_\tau^i = e_{\alpha_\tau^i - 1}$.

For $j \neq i$ we have $\tilde{\mathbf{p}}_\tau^j = 0$, which implies $\text{diag}(\mathbf{D}_j) \circ \mathbf{r}_\tau^j \circ \tilde{\mathbf{p}}_\tau^j = 0$.

All together, this implies that the τ -th column of $\mathbf{D}(\mathbf{R}_1 \circ \text{sgn}(\mathbf{P}_1(x)))$, given by $(\text{diag}(\mathbf{D}_j) \circ \mathbf{r}_\tau^j \circ \tilde{\mathbf{p}}_\tau^j)_{j=1}^{\tilde{N}}$, is nonzero and every nonzero entry is equal to 1.

Since this holds for arbitrary diagonal sign matrix $\mathbf{D} \neq 0$, Lemma 26 implies that $(\mathbf{R}_1 \circ \text{sgn}(\mathbf{P}_1(x)))$ is an L -matrix. Since this holds for any x satisfying $\text{carr}(x) \subseteq C$, we see that the desired result holds. \square

Appendix C.

Lemma 28. *Let $\mathbf{B}(x)$ be as defined in Section 5.2. There are γ columns of $\mathbf{B}(x)$ such that these columns form an invertible diagonal matrix for any x with $\text{carr}(x) = C$.*

Proof. Without loss of generality, suppose $C = C_1 \cup \dots \cup C_N$ and assume that each player's pure strategy set Y_i is ordered so that $y_i^1 \in C_i$, $i = 1, \dots, \tilde{N}$. Now, let $i \in \{1, \dots, \tilde{N}\}$ be arbitrary. Suppose that the set of pure strategies Y is ordered and that the τ 'th pure strategy, $\tau \in \{1, \dots, K\}$, takes the form $(y_1^1, \dots, y_{i-1}^1, y_i^{k+1}, y_{i+1}^1, \dots, y_N^1)$, where $i \in \{1, \dots, \tilde{N}\}$ and $k \in \{1, \dots, \gamma_i - 1\}$. For a given i , there are precisely $\gamma_i - 1$ such strategy profiles. Fixing τ to this value (thus looking at a single column of $\mathbf{B}(x)$) we see from the definition of q_i^τ that

$$q_i^\tau(y_i^\ell) = \begin{cases} 1 & \text{if } \ell = k + 1 \\ 0 & \text{else.} \end{cases}$$

Given our choice of τ (and ordering of Y_i) we also have $\prod_{j \neq i} q_j^\tau(x_j) > 0$ for all x with carrier C . Given the definition of $\mathbf{B}(x)$ in (27), this implies that the τ -th

column of $\mathbf{B}_i(x)$ is composed of zeros except for the k -th entry which takes the positive value $\prod_{j \neq i} q_j^r(x_j) > 0$.

Without loss of generality, we may reorder the set of pure strategies so that the γ combinations of pure strategies taking the aforementioned form constitute the first γ strategies in the set. Under this ordering, we have shown that $\mathbf{B}(x)$ takes the form $\mathbf{B}(x) = [\mathbf{C}(x) \mathbf{D}(x)]$, where $\mathbf{C}(x)$ is a $\gamma \times \gamma$ diagonal matrix (up to a permutation of the columns) with diagonal entries $\prod_{j \neq i} q_j^r(x_j) > 0$. \square

Lemma 29. *Let $x \in X$ and $i = 1, \dots, N$. Assume Y_i is ordered so that $y_i^1 \in BR_i(x_{-i})$. Then: (i) For $k = 1, \dots, K_i - 1$ we have $\frac{\partial U(x)}{\partial x_i^k} \leq 0$, and (ii) For $k = 1, \dots, K_i - 1$, we have $y_i^{k+1} \in BR_i(x_{-i})$ if and only if $\frac{\partial U(x)}{\partial x_i^k} = 0$. In particular, combined with (i) this implies that $y_i^{k+1} \notin BR_i(x_{-i}) \iff \frac{\partial U(x)}{\partial x_i^k} < 0$.*

Proof. (i) Differentiating (7) we find that $\frac{\partial U(x)}{\partial x_i^k} = U(y_i^{k+1}, x_{-i}) - U(y_i^1, x_{-i})$ (i) Since y_i^1 is a best response, we must have $U(y_i^1, x_{-i}) \geq U(y_i^{k+1}, x_{-i})$ for any $k = 1, \dots, K_i - 1$. Hence $\frac{\partial U(x)}{\partial x_i^k} \leq 0$.
(ii) Follows readily from (7). \square

Lemma 30. *Suppose x^* is an equilibrium and $y_i^k \in \text{carr}(x^*)$, $k \geq 2$. Then $\frac{\partial U(x^*)}{\partial x_i^k} = 0$.*

Proof. Since U is multilinear, y_i^k must be a pure-strategy best response to x_{-i}^* . The result then follows from Lemma 29. \square

References

- [1] R. Selten, Reexamination of the perfectness concept for equilibrium points in extensive games, *International Journal of Game Theory* 4 (1) (1975) 25–55.
- [2] R. B. Myerson, Refinements of the Nash equilibrium concept, *International Journal of Game Theory* 7 (2) (1978) 73–80.
- [3] J. C. Harsanyi, Oddness of the number of equilibrium points: A new proof, *International Journal of Game Theory* 2 (1) (1973) 235–250.
- [4] E. Van Damme, *Stability and perfection of Nash equilibria*, Vol. 339, Springer, 1991.

- [5] W. Wen-Tsun, J. Jia-He, Essential equilibrium points of n -person non-cooperative games, *Sci Sinica* 11 (10) (1962) 1307–1322.
- [6] M. Kojima, A. Okada, S. Shindoh, Strongly stable equilibrium points of n -person noncooperative games, *Mathematics of Operations Research* 10 (4) (1985) 650–663.
- [7] A. Okada, On stability of perfect equilibrium points, *International Journal of Game Theory* 10 (2) (1981) 67–73.
- [8] R. Wilson, Computing equilibria of n -person games, *SIAM Journal on Applied Mathematics* 21 (1) (1971) 80–87.
- [9] J. C. Harsanyi, Games with randomly disturbed payoffs: A new rationale for mixed-strategy equilibrium points, *International Journal of Game Theory* 2 (1) (1973) 1–23.
- [10] S. Govindan, P. J. Reny, A. J. Robson, A short proof of harsanyi’s purification theorem, *Games and Economic Behavior* 45 (2) (2003) 369–374.
- [11] S. Morris, “Purification” in *The New Palgrave Dictionary of Economics*, Vol. 6, Palgrave Macmillan Basingstoke, 2008.
- [12] F. Brandt, F. Fischer, On the hardness and existence of quasi-strict equilibria, in: *International Symposium on Algorithmic Game Theory*, Springer, 2008, pp. 291–302.
- [13] D. Monderer, L. Shapley, Potential games, *Games and Economic Behavior* 14 (1) (1996) 124–143.
- [14] J. R. Marden, G. Arslan, J. S. Shamma, Cooperative control and potential games, *IEEE Transactions on Systems, Man, and Cybernetics, Part B (Cybernetics)* 39 (6) (2009) 1393–1407.
- [15] G. Scutari, S. Barbarossa, D. P. Palomar, Potential games: A framework for vector power control problems with coupled constraints, in: *Proceedings of the IEEE International Conference on Acoustics, Speech and Signal Processing*, Vol. 4, IEEE, 2006, pp. 241–244.

- [16] Y. Xu, A. Anpalagan, Q. Wu, L. Shen, Z. Gao, J. Wang, Decision-theoretic distributed channel selection for opportunistic spectrum access: Strategies, challenges and solutions, *IEEE Communications Surveys & Tutorials* 15 (4) (2013) 1689–1713.
- [17] M. Zhu, S. Martínez, Distributed coverage games for energy-aware mobile sensor networks, *SIAM Journal on Control and Optimization* 51 (1) (2013) 1–27.
- [18] C. Ding, B. Song, A. Morye, J. A. Farrell, A. K. Roy-Chowdhury, Collaborative sensing in a distributed PTZ camera network, *IEEE Transactions on Image Processing* 21 (7) (2012) 3282–3295.
- [19] N. Nie, C. Comaniciu, Adaptive channel allocation spectrum etiquette for cognitive radio networks, *Mobile Networks and Applications* 11 (6) (2006) 779–797.
- [20] X. Chu, H. Sethu, Cooperative topology control with adaptation for improved lifetime in wireless ad hoc networks, in: *IEEE International Conference on Computer Communications*, IEEE, 2012, pp. 262–270.
- [21] V. Srivastava, J. O. Neel, A. B. MacKenzie, R. Menon, L. A. DaSilva, J. E. Hicks, J. H. Reed, R. P. Gilles, Using game theory to analyze wireless ad hoc networks, *IEEE Communications Surveys and Tutorials* 7 (4) (2005) 46–56.
- [22] T. J. Lambert, M. A. Epleman, R. L. Smith, A fictitious play approach to large-scale optimization, *Operations Research* 53 (3) (2005) 477–489.
- [23] A. Garcia, D. Reaume, R. L. Smith, Fictitious play for finding system optimal routings in dynamic traffic networks, *Transportation Research Part B: Methodological* 34 (2) (2000) 147–156.
- [24] J. R. Marden, G. Arslan, J. S. Shamma, Connections between cooperative control and potential games, *IEEE Transactions on Systems, Man and Cybernetics. Part B: Cybernetics* 39 (2009) 1393–1407.
- [25] N. Li, J. R. Marden, Designing games for distributed optimization, *IEEE Journal of Selected Topics in Signal Processing* 7 (2) (2013) 230–242.

- [26] D. Monderer, L. S. Shapley, Fictitious play property for games with identical interests, *Journal of Economic Theory* 68 (1) (1996) 258–265.
- [27] M. Voorneveld, Best-response potential games, *Economics Letters* 66 (3) (2000) 289–295.
- [28] D. Fudenberg, D. K. Levine, *The theory of learning in games*, MIT press, 1998.
- [29] B. Swenson, R. Murray, S. Kar, On best-response dynamics in potential games, *SIAM Journal on Control and Optimization* 56 (4) (2018) 2734–2767.
- [30] J. Cohen, A. Héliou, P. Mertikopoulos, Learning with bandit feedback in potential games, in: *Proceedings of the 31th International Conference on Neural Information Processing Systems*, 2017.
- [31] I. Gilboa, A. Matsui, Social stability and equilibrium, *Econometrica: Journal of the Econometric Society* 59 (3) (1991) 859–867.
- [32] J. Hofbauer, Stability for the best response dynamics, Tech. rep., Institut für Mathematik, Universität Wien, Strudlhofgasse 4, A-1090 Vienna, Austria (1995).
- [33] J. Hofbauer, K. Sigmund, Evolutionary game dynamics, *Bulletin of the American Mathematical Society* 40 (4) (2003) 479–519.
- [34] M. Benaïm, J. Hofbauer, S. Sorin, Stochastic approximations and differential inclusions, *SIAM J. Control and Optim.* 44 (1) (2005) 328–348.
- [35] C. Harris, On the rate of convergence of continuous-time fictitious play, *Games and Economic Behavior* 22 (2) (1998) 238–259.
- [36] J. S. Jordan, Three problems in learning mixed-strategy Nash equilibria, *Games and Econ. Behav.* 5 (3) (1993) 368–386.
- [37] D. Fudenberg, Learning mixed equilibria, *Games and Econ. Behav.* 5 (3) (1993) 320–367.
- [38] D. Monderer, A. Sela, Fictitious play and no-cycling conditions.

- [39] J. S. Shamma, G. Arslan, Unified convergence proofs of continuous-time fictitious play, *IEEE Transactions on Automatic Control* 49 (7) (2004) 1137–1141.
- [40] E. van Damme, Regular equilibrium points of n -person games in normal form (1981).
- [41] M. W. Hirsch, *Differential topology*, Springer, 1976.
- [42] D. Fudenberg, J. Tirole, *Game theory*, MIT press, 1991.
- [43] R. A. Brualdi, K. L. Chavey, B. L. Shader, Rectangular L -matrices, *Linear Algebra and its Applications* 196 (1994) 37–61.
- [44] V. Klee, R. Ladner, R. Manber, Signsolvability revisited, *Linear Algebra and its Applications* 59 (1984) 131–157.
- [45] L. C. Evans, R. F. Gariepy, *Measure theory and fine properties of functions*, revised Edition, *Textbooks in Mathematics*, CRC Press, Boca Raton, FL, 2015.
- [46] V. Krishna, T. Sjöström, On the convergence of fictitious play, *Mathematics of Operations Research* 23 (2) (1998) 479–511.