

Formal Embeddings between \mathcal{BSD} -Models

Valentin Burcea

ABSTRACT. It is studied the Classification Problem for Formal (Holomorphic) Embeddings between Shilov Boundaries of Bounded Symmetric Domains of First Type in Complex Spaces of Different Dimensions using techniques of Linear Algebra.

1. Introduction and Main Result

The study of proper holomorphic mappings[25], between unit balls in Complex Spaces, goes back to Webster[26]. For $N > n$, two proper holomorphic mappings $f, g : \mathbb{B}^n \rightarrow \mathbb{B}^N$ are called equivalent if there exist $\sigma \in \text{Aut}(\mathbb{B}^n)$ and $\tau \in \text{Aut}(\mathbb{B}^N)$ such that $g = \tau \circ f \circ \sigma$. Huang[10] proved that any proper holomorphic mapping, between \mathbb{B}^n and \mathbb{B}^N of class \mathcal{C}^2 up to the boundary, is equivalent to

$$(z_1, z_2, \dots, z_n) \longrightarrow (z_1, z_2, \dots, z_n, 0, \dots, 0), \text{ for all } n > 1 \text{ and } N < 2n - 1.$$

Huang-Ji[12] proved that the rational proper holomorphic mappings between \mathbb{B}^n and \mathbb{B}^{2n-1} are equivalent to

$$(z_1, z_2, \dots, z_n) \rightarrow (z_1, z_2, \dots, z_n, 0, \dots, 0), (z_1, z_2, \dots, z_{n-1}, z_n z_1, z_n z_2, \dots, z_n^2), \text{ for all } n \geq 3.$$

The Classification Problem, for proper holomorphic mappings[8],[10],[12] between unit balls in Complex Spaces, is reduced to the study of CR mappings between hyperquadrics (see [1],[2],[19]). More generally, the Classification Problem for C.-R. Embeddings, between Shilov Boundaries of Bounded Symmetric Domains, is also very interesting. Kim-Zaitsev[16] considered this problem using the moving frames method of Cartan for Shilov Boundaries of Bounded Symmetric Domains of First Type with $q < p$, $q' < p'$ such that $p' - q' < 2(p - q)$ and $p - q > 1$. They[16] proved that, up to compositions with suitable automorphisms of Bounded Symmetric Domains of First Type $D_{p,q}$ and $D_{p',q'}$, any Smooth C.-R. Embedding, between their Shilov Boundaries denoted by $S_{p,q}$ and $S_{p',q'}$, is equivalent to

$$(1.1) \quad Z = \begin{pmatrix} z_{11} & z_{12} & \dots & z_{1q} \\ z_{21} & z_{22} & \dots & z_{2q} \\ \vdots & \vdots & \ddots & \vdots \\ z_{p1} & z_{p2} & \dots & z_{pq} \end{pmatrix} \rightarrow \begin{pmatrix} z_{11} & z_{12} & \dots & z_{1q} & 0 & 0 & \dots & 0 \\ z_{21} & z_{22} & \dots & z_{2q} & 0 & 0 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots \\ z_{p1} & z_{p2} & \dots & z_{pq} & 0 & 0 & \dots & 0 \\ 0 & 0 & \dots & 0 & 1 & 0 & \dots & 0 \\ 0 & 0 & \dots & 0 & 0 & 1 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & 0 & 0 & 0 & \dots & 1 \\ 0 & 0 & \dots & 0 & 0 & 0 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & 0 & 0 & 0 & \dots & 0 \end{pmatrix}.$$

Any Bounded Symmetric Domain $D_{p,q}$ of First Type and its Shilov Boundary (see [16],[25]) may be defined by

$$(1.2) \quad D_{p,q} = \left\{ Z \in \mathcal{M}_{p \times q}(\mathbb{C}); \quad I_q - \overline{Z}^t Z > 0 \right\} \text{ and } S_{p,q} = \left\{ Z \in \mathcal{M}_{p \times q}(\mathbb{C}); \quad I_q - \overline{Z}^t Z = 0 \right\}, \text{ for } p > q.$$

It is indicated Chirvasitu[6] for the standard definition of the Shilov Boundary. It generalizes classical models as the hyperquadrics (see [5],[9],[10],[11],[12],[13],[19],[24]). It has a fundamental importance in the study of Holomorphic Isometries in Complex Analysis and Complex Geometry (see [23],[27],[28]) and their properties.

In this, we use the language of matrices in order to establish a normal form type construction (see [2],[3],[29],[30]) for Formal (Holomorphic) Embeddings between Shilov Boundaries of Bounded Symmetric Domains of First Type (see [16],[17],[18],[25]). We make computations using formal power series. In particular, we use suitable linear changes of coordinates in order to implement procedures from Bauendi-Huang[1], Chern-Moser[4], Hamada[9], Huang[10],[11], Huang-Ji[12] and Kim-Zaitsev[16],[17]. It is an alternative to the Method of Cartan applied by Kim-Zaitsev[16],[17]. It detects an analogue of the geometrical rank introduced by Huang[10],[11]. This new invariant is defined by several matrices having identical rank. The main result is the following:

Keywords: Normal Form, Equivalence Problem, Embedding, Bounded Symmetric Domain, Real Submanifold, Shilov Boundary, Power Series.
 This project was supported principally (80%) by CAPES at Federal University of Espirito Santo, being initiated (20%) with Science Foundation Ireland, Grant 10/RFP/MT H2878. Emphasizing that the paper [3] was fully supported by Science Foundation Ireland, Grant 06/RFP/MAT 018.

THEOREM 1.1. *Let $p, p', q, q' \in \mathbb{N}^*$ such that $p' - q' = 2(p - q) > 2$. Up to compositions with suitable automorphisms of Bounded Symmetric Domains $D_{p,q}$ and $D_{p',q'}$, any Formal Embedding, between their Shilov Boundaries, is equivalent to*

$$(1.3) \quad Z = \begin{pmatrix} z_{11} & z_{12} & \dots & z_{1q} \\ z_{21} & z_{22} & \dots & z_{2q} \\ \vdots & \vdots & \ddots & \vdots \\ z_{p-1\ 1} & z_{p-1\ 2} & \dots & z_{p-1\ q} \\ z_{p1}z_{11} & z_{p2}z_{12} & \dots & z_{pq}z_{1q} \\ z_{p1}z_{21} & z_{p2}z_{22} & \dots & z_{pq}z_{2q} \\ \vdots & \vdots & \ddots & \vdots \\ z_{p1}z_{p1} & z_{p2}z_{p2} & \dots & z_{pq}z_{pq} \\ 0 & 0 & \dots & 0 \\ 0 & 0 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & 0 \\ 0 & 0 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & 0 \end{pmatrix} \mapsto \begin{pmatrix} z_{11} & z_{12} & \dots & z_{1q} & 0 & 0 & \dots & 0 \\ z_{21} & z_{22} & \dots & z_{2q} & 0 & 0 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots \\ z_{p-1\ 1} & z_{p-1\ 2} & \dots & z_{p-1\ q} & 0 & 0 & \dots & 0 \\ z_{p1}z_{11} & z_{p2}z_{12} & \dots & z_{pq}z_{1q} & 0 & 0 & \dots & 0 \\ z_{p1}z_{21} & z_{p2}z_{22} & \dots & z_{pq}z_{2q} & 0 & 0 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots \\ z_{p1}z_{p1} & z_{p2}z_{p2} & \dots & z_{pq}z_{pq} & 0 & 0 & \dots & 0 \\ 0 & 0 & \dots & 0 & 1 & 0 & \dots & 0 \\ 0 & 0 & \dots & 0 & 0 & 1 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & 0 & 0 & 0 & \dots & 1 \\ 0 & 0 & \dots & 0 & 0 & 0 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & 0 & 0 & 0 & \dots & 0 \end{pmatrix} \text{ or to (1.1).}$$

The proof of Theorem 1.1 is reduced to the study of Formal Embeddings between \mathcal{BSD} -Models. They are quadratic Models derived from Shilov Boundaries of Bounded Symmetric Domains of First Type using a generalized Cayley Transformation[7]. There are obtained two Classes of Equivalence as it was anticipated by Seo[20],[21]. The first Class of Equivalence is the standard linear Embedding described by Kim-Zaitsev[16]. The second Class of Equivalence is a generalized Whitney type mapping (see [20],[21]). The Classification (1.3) may be seen as an analogue of the Classification Theorem of Huang-Ji[12], especially when $p' = 2p - 1$ and $q' = 2q - 1$.

2. Settings

We work in the coordinates

$$(2.1) \quad (w_{11}, w_{12}, \dots, w_{1q}, \dots, w_{q1}, w_{q2}, \dots, w_{qq}; z_{11}, z_{12}, \dots, z_{1N}, \dots, z_{q1}, z_{q2}, \dots, z_{qN}) \in \mathbb{C}^{qN+q^2}.$$

In particular, we consider matrices using the identifications

$$(2.2) \quad W = \begin{pmatrix} w_{11} & w_{12} & \dots & w_{1q} \\ w_{21} & w_{22} & \dots & w_{2q} \\ \vdots & \vdots & \ddots & \vdots \\ w_{q1} & w_{q2} & \dots & w_{qq} \end{pmatrix} \equiv (w_{11}, w_{12}, \dots, w_{1q}, w_{21}, w_{22}, \dots, w_{2q}, \dots, w_{q1}, w_{q2}, \dots, w_{qq}),$$

$$Z = \begin{pmatrix} z_{11} & z_{12} & \dots & z_{1N} \\ z_{21} & z_{22} & \dots & z_{2N} \\ \vdots & \vdots & \ddots & \vdots \\ z_{q1} & z_{q2} & \dots & z_{qN} \end{pmatrix} \equiv (z_{11}, z_{12}, \dots, z_{1N}, z_{21}, z_{22}, \dots, z_{2N}, \dots, z_{q1}, z_{q2}, \dots, z_{qN}).$$

In particular, we use the identifications

$$(2.3) \quad J := \begin{pmatrix} j_{11} & j_{12} & \dots & j_{1q} \\ j_{21} & j_{22} & \dots & j_{2q} \\ \vdots & \vdots & \ddots & \vdots \\ j_{q1} & j_{q2} & \dots & j_{qq} \end{pmatrix} \equiv (j_{11}, j_{12}, \dots, j_{1q}, j_{21}, j_{22}, \dots, j_{2q}, \dots, j_{q1}, j_{q2}, \dots, j_{qq}) \in \mathbb{N}^{q^2},$$

$$I := \begin{pmatrix} i_{11} & i_{12} & \dots & i_{1N} \\ i_{21} & i_{22} & \dots & i_{2N} \\ \vdots & \vdots & \ddots & \vdots \\ i_{m1} & i_{m2} & \dots & i_{qN} \end{pmatrix} \equiv (i_{11}, i_{12}, \dots, i_{1N}, i_{21}, i_{22}, \dots, i_{2N}, \dots, i_{q1}, i_{q2}, \dots, i_{qN}) \in \mathbb{N}^{qN}.$$

In the light of (2.2) and (2.3), we write

$$(2.4) \quad W^J = w_{11}^{j_{11}} w_{12}^{j_{12}} \dots w_{1q}^{j_{1q}} w_{21}^{j_{21}} w_{22}^{j_{22}} \dots w_{2q}^{j_{2q}} \dots w_{q1}^{j_{q1}} w_{q2}^{j_{q2}} \dots w_{qq}^{j_{qq}},$$

$$Z^I = z_{11}^{i_{11}} z_{12}^{i_{12}} \dots z_{1N}^{i_{1N}} z_{21}^{i_{21}} z_{22}^{i_{22}} \dots z_{2N}^{i_{2N}} \dots z_{q1}^{i_{q1}} z_{q2}^{i_{q2}} \dots z_{qN}^{i_{qN}}.$$

Ignoring the considered numbers, we generalize the standard hermitian inner-product as follows

$$(2.5) \quad \langle L, V \rangle = L \overline{V}^t, \quad \text{provided } L \in \mathcal{M}_{m,n}(\mathbb{C}) \text{ and } V \in \mathcal{M}_{n,p}(\mathbb{C}), \text{ for all } m, n, p \in \mathbb{N}^*.$$

Analogous notations and identifications to (2.2),(2.3) and (2.4) are used in order to work in the coordinates

$$(2.6) \quad (w'_{11}, w'_{12}, \dots, w'_{1q'}, \dots, w'_{q'1}, w'_{q'2}, \dots, w'_{q'q'}; z'_{11}, z'_{12}, \dots, z'_{1N'}, \dots, z'_{q'1}, z'_{q'2}, \dots, z'_{q'N'}) \in \mathbb{C}^{q'N'+q'^2}.$$

In particular, we use matrices denoted by W' and Z' . The rows of the matrices Z and Z' are denoted by

$$(2.7) \quad Z_1, Z_2, \dots, Z_q \text{ and } Z'_1, Z'_2, \dots, Z'_{q'}.$$

3. Formal Embeddings between \mathcal{BSD} -Models

We replace the generalized Cayley transformation (see [7]) in (1.2). It is denoted by

$$(3.1) \quad \mathcal{C}(W, Z) = \left(\frac{[W - \sqrt{-1}I_q, 2Z]}{W + \sqrt{-1}I_q} \right)^t.$$

We obtain the equation of the \mathcal{BSD} -Model

$$(3.2) \quad \text{Im}W := \frac{1}{2\sqrt{-1}} (W - \overline{W}^t) = Z\overline{Z}^t.$$

Any Formal Embedding, denoted by (\tilde{G}, \tilde{F}) , between Shilov Boundaries of Bounded Symmetric Domains of First Type, induces naturally by (3.1) another Formal Embedding, denoted by (G, F) , between the \mathcal{BSD} -Models defined by

$$(3.3) \quad \mathcal{M} : \text{Im}W = Z\overline{Z}^t \subset \mathbb{C}^{qN+q^2} \text{ and } \mathcal{M}' : \text{Im}W' = Z'\overline{Z}'^t \subset \mathbb{C}^{q'N'+q'^2}, \text{ for } N = p - q \text{ and } N' = p' - q'.$$

More precisely, we work with the commutative diagram

$$(3.4) \quad \begin{array}{ccc} \mathcal{M} & \xrightarrow{(G, F)} & \mathcal{M}' \\ \Downarrow & & \Downarrow \\ S_{p, q} & \xrightarrow{(\tilde{G}, \tilde{F})} & S_{p', q'}. \end{array}$$

The above equivalences are defined by transformations as (3.1). The Formal Embedding (G, F) is written as

$$(3.5) \quad G(W, Z) := \begin{pmatrix} G_{11}(W, Z) & G_{12}(W, Z) \\ G_{21}(W, Z) & G_{22}(W, Z) \end{pmatrix} \text{ and } F(W, Z) := \begin{pmatrix} F_1(W, Z) \\ F_2(W, Z) \end{pmatrix},$$

where we have used the following submatrices:

- $G_{11}(W, Z)$ is a $q \times q$ submatrix having formal power series in (W, Z) as entries,
- $G_{21}(W, Z)$ is a $(q' - q) \times q$ submatrix having formal power series in (W, Z) as entries,
- $G_{12}(W, Z)$ is a $q \times (q' - q)$ submatrix having formal power series in (W, Z) as entries,
- $G_{22}(W, Z)$ is a $(q' - q) \times (q' - q)$ submatrix having formal power series in (W, Z) as entries,
- $F_1(W, Z)$ is a $q \times N'$ submatrix having formal power series in (W, Z) as entries,
- $F_2(W, Z)$ is a $(q' - q) \times N'$ submatrix having formal power series in (W, Z) as entries.

Their linear parts in Z are vanishing. We obtain $\text{Im}(G(Z, W)) = F(Z, W) \cdot \overline{F(Z, W)}^t$, or equivalently

$$(3.6) \quad \begin{aligned} G_{11}(W, Z) - \overline{G_{11}(W, Z)}^t &= 2\sqrt{-1} \langle F_1(W, Z), F_1(W, Z) \rangle, \\ G_{22}(W, Z) - \overline{G_{22}(W, Z)}^t &= 2\sqrt{-1} \langle F_2(W, Z), F_2(W, Z) \rangle, \\ G_{12}(W, Z) - \overline{G_{21}(W, Z)}^t &= 2\sqrt{-1} \langle F_1(W, Z), F_2(W, Z) \rangle. \end{aligned}$$

We write using (2.2) the matrices from (3.5), in terms of their entries, as

$$(3.7) \quad G(Z, W) = (g_{kl}(Z, W))_{1 \leq k, l \leq q'} \text{ and } F(Z, W) = (f_{kl}(Z, W))_{\substack{1 \leq l \leq N' \\ 1 \leq k \leq q'}}.$$

The equations (3.6) are used in order to consider transformations preserving \mathcal{BSD} -Models. In particular, we use the product of matrices

$$(3.8) \quad V \otimes Z = \left(\sum_{l=1}^N \sum_{k=1}^q v_{kl}^{ij} z_{kl} \right)_{\substack{1 \leq i \leq q \\ 1 \leq j \leq N}},$$

provided the matrix $V = (v_{\alpha}^{\beta})_{\substack{1 \leq \beta \leq qN \\ 1 \leq \alpha \leq qN}} \in \mathcal{M}_{qN \times qN}(\mathbb{C})$ in the light of the identification

$$V \equiv \begin{pmatrix} \begin{pmatrix} v_{11}^{11} & v_{12}^{11} & \dots & v_{1N}^{11} \\ v_{11}^{12} & v_{12}^{12} & \dots & v_{1N}^{12} \\ \vdots & \vdots & \ddots & \vdots \\ v_{11}^{1N} & v_{12}^{1N} & \dots & v_{1N}^{1N} \end{pmatrix} & \begin{pmatrix} v_{21}^{11} & v_{22}^{11} & \dots & v_{2N}^{11} \\ v_{21}^{12} & v_{22}^{12} & \dots & v_{2N}^{12} \\ \vdots & \vdots & \ddots & \vdots \\ v_{21}^{1N} & v_{22}^{1N} & \dots & v_{2N}^{1N} \end{pmatrix} & \begin{pmatrix} \dots \\ \dots \\ \vdots \\ \dots \end{pmatrix} & \begin{pmatrix} v_{q1}^{11} & v_{q2}^{11} & \dots & v_{qN}^{11} \\ v_{q1}^{12} & v_{q2}^{12} & \dots & v_{qN}^{12} \\ \vdots & \vdots & \ddots & \vdots \\ v_{q1}^{1N} & v_{q2}^{1N} & \dots & v_{qN}^{1N} \end{pmatrix} \\ \begin{pmatrix} v_{11}^{21} & v_{12}^{21} & \dots & v_{1N}^{21} \\ v_{11}^{22} & v_{12}^{22} & \dots & v_{1N}^{22} \\ \vdots & \vdots & \ddots & \vdots \\ v_{11}^{2N} & v_{12}^{2N} & \dots & v_{1N}^{2N} \end{pmatrix} & \begin{pmatrix} v_{21}^{21} & v_{22}^{21} & \dots & v_{2N}^{21} \\ v_{21}^{22} & v_{22}^{22} & \dots & v_{2N}^{22} \\ \vdots & \vdots & \ddots & \vdots \\ v_{21}^{2N} & v_{22}^{2N} & \dots & v_{2N}^{2N} \end{pmatrix} & \begin{pmatrix} \dots \\ \dots \\ \vdots \\ \dots \end{pmatrix} & \begin{pmatrix} v_{q1}^{21} & v_{q2}^{21} & \dots & v_{qN}^{21} \\ v_{q1}^{22} & v_{q2}^{22} & \dots & v_{qN}^{22} \\ \vdots & \vdots & \ddots & \vdots \\ v_{q1}^{2N} & v_{q2}^{2N} & \dots & v_{qN}^{2N} \end{pmatrix} \\ \begin{pmatrix} \vdots & \vdots & \ddots & \vdots \\ \vdots & \vdots & \ddots & \vdots \\ \vdots & \vdots & \ddots & \vdots \end{pmatrix} & \begin{pmatrix} \vdots & \vdots & \ddots & \vdots \\ \vdots & \vdots & \ddots & \vdots \\ \vdots & \vdots & \ddots & \vdots \end{pmatrix} & \begin{pmatrix} \vdots \\ \vdots \\ \vdots \end{pmatrix} & \begin{pmatrix} \vdots & \vdots & \ddots & \vdots \\ \vdots & \vdots & \ddots & \vdots \\ \vdots & \vdots & \ddots & \vdots \end{pmatrix} \\ \begin{pmatrix} v_{11}^{q1} & v_{12}^{q1} & \dots & v_{1N}^{q1} \\ v_{11}^{q2} & v_{12}^{q2} & \dots & v_{1N}^{q2} \\ \vdots & \vdots & \ddots & \vdots \\ v_{11}^{qN} & v_{12}^{qN} & \dots & v_{1N}^{qN} \end{pmatrix} & \begin{pmatrix} v_{21}^{q1} & v_{22}^{q1} & \dots & v_{2N}^{q1} \\ v_{21}^{q2} & v_{22}^{q2} & \dots & v_{2N}^{q2} \\ \vdots & \vdots & \ddots & \vdots \\ v_{21}^{qN} & v_{22}^{qN} & \dots & v_{2N}^{qN} \end{pmatrix} & \begin{pmatrix} \dots \\ \dots \\ \vdots \\ \dots \end{pmatrix} & \begin{pmatrix} v_{q1}^{q1} & v_{q2}^{q1} & \dots & v_{qN}^{q1} \\ v_{q1}^{q2} & v_{q2}^{q2} & \dots & v_{qN}^{q2} \\ \vdots & \vdots & \ddots & \vdots \\ v_{q1}^{qN} & v_{q2}^{qN} & \dots & v_{qN}^{qN} \end{pmatrix} \end{pmatrix}.$$

Motivated by (2.2), (2.3) and (3.8), we use identifications as

$$(3.9) \quad \begin{aligned} \{1, 2, 3, \dots, q^2\} &\equiv \{(1, 1), (1, 2), \dots, (1, q), \\ &\quad (2, 1), (2, 2), \dots, (2, q), \\ &\quad \vdots \quad \vdots \quad \ddots \quad \vdots \\ &\quad (q, 1), (q, 2), \dots, (q, q)\}, \\ \{1, 2, 3, \dots, qN\} &\equiv \{(1, 1), (1, 2), \dots, (1, N), \\ &\quad (2, 1), (2, 2), \dots, (2, N), \\ &\quad \vdots \quad \vdots \quad \ddots \quad \vdots \\ &\quad (q, 1), (q, 2), \dots, (q, N)\}. \end{aligned}$$

In order to organize further computations, we write the matrix V as

$$(3.10) \quad V \equiv \begin{pmatrix} V_{11} & V_{12} & \dots & V_{1q} \\ V_{21} & V_{22} & \dots & V_{2q} \\ \vdots & \vdots & \ddots & \vdots \\ V_{q1} & V_{q2} & \dots & V_{qq} \end{pmatrix}.$$

In particular, we consider using (2.7) the following replacements

$$(3.11) \quad \begin{aligned} w_{11} &= \bar{w}_{11} + 2\sqrt{-1} \langle Z_1, Z_1 \rangle, & w_{12} &= \bar{w}_{21} + 2\sqrt{-1} \langle Z_1, Z_2 \rangle, \dots & w_{1q} &= \bar{w}_{q1} + 2\sqrt{-1} \langle Z_1, Z_q \rangle, \\ w_{21} &= \bar{w}_{12} + 2\sqrt{-1} \langle Z_2, Z_1 \rangle, & w_{22} &= \bar{w}_{22} + 2\sqrt{-1} \langle Z_2, Z_2 \rangle, \dots & w_{2q} &= \bar{w}_{q2} + 2\sqrt{-1} \langle Z_2, Z_q \rangle, \\ &\vdots & & \vdots & & \vdots \\ w_{q1} &= \bar{w}_{1q} + 2\sqrt{-1} \langle Z_q, Z_1 \rangle, & w_{q2} &= \bar{w}_{2q} + 2\sqrt{-1} \langle Z_q, Z_2 \rangle, \dots & w_{qq} &= \bar{w}_{qq} + 2\sqrt{-1} \langle Z_q, Z_q \rangle. \end{aligned}$$

$$(3.12) \quad \begin{aligned} w'_{11} &= \bar{w}'_{11} + 2\sqrt{-1} \langle Z'_1, Z'_1 \rangle, & w'_{12} &= \bar{w}'_{21} + 2\sqrt{-1} \langle Z'_1, Z'_2 \rangle, \dots & w'_{1q'} &= \bar{w}'_{q'1} + 2\sqrt{-1} \langle Z'_1, Z'_{q'} \rangle, \\ w'_{21} &= \bar{w}'_{12} + 2\sqrt{-1} \langle Z'_2, Z'_1 \rangle, & w'_{22} &= \bar{w}'_{22} + 2\sqrt{-1} \langle Z'_2, Z'_2 \rangle, \dots & w'_{2q'} &= \bar{w}'_{q'2} + 2\sqrt{-1} \langle Z'_2, Z'_{q'} \rangle, \\ &\vdots & & \vdots & & \vdots \\ w'_{q'1} &= \bar{w}'_{1q'} + 2\sqrt{-1} \langle Z'_{q'}, Z'_1 \rangle, & w'_{q'2} &= \bar{w}'_{2q'} + 2\sqrt{-1} \langle Z'_{q'}, Z'_2 \rangle, \dots & w'_{q'q'} &= \bar{w}'_{q'q'} + 2\sqrt{-1} \langle Z'_{q'}, Z'_{q'} \rangle. \end{aligned}$$

It is required by (2.1),(2.2),(2.5),(2.6),(2.7),(3.4),(3.5),(3.6),(3.7),(3.8),(3.11) and (3.12) to implement the following:

4. Application of the Normalization Procedure from Bauoendi-Huang[1]

We study the Formal Embedding (3.5) using the commutative diagram

$$(4.1) \quad \begin{array}{ccc} \mathcal{M} & \xrightarrow{(F,G)} & \mathcal{M}' \\ \Downarrow & & \Downarrow \\ \mathcal{M} & \xrightarrow{(F,G)} & \mathcal{M}'. \end{array}$$

Each of the above Equivalences are defined by (3.8) using changes of coordinates preserving BSD -Models. It follows that

PROPOSITION 4.1. *Up to compositions with suitable linear automorphisms of BSD -Models from (3.3), we obtain*

$$(4.2) \quad \begin{pmatrix} G_{11}(Z, W) & G_{12}(Z, W) \\ G_{21}(Z, W) & G_{22}(Z, W) \end{pmatrix} = \begin{pmatrix} W + O(2) & O(2) \\ O(2) & O(2) \end{pmatrix} \text{ and } \begin{pmatrix} F_1(Z, W) \\ F_2(Z, W) \end{pmatrix} = \begin{pmatrix} Z + O(2) & O(2) \\ O(2) & O(2) \end{pmatrix}.$$

PROOF. We write (3.5) as follows

$$(4.3) \quad \begin{aligned} G_{11}(W, Z) &= A \otimes W + O(2), & G_{12}(W, Z) &= B \otimes W + O(2), \\ G_{21}(W, Z) &= C \otimes W + O(2), & G_{22}(W, Z) &= D \otimes W + O(2), \end{aligned}$$

where we have used by (3.8) the following matrices

$$(4.4) \quad \begin{aligned} A &= (A^{ij})_{1 \leq i, j \leq q} = \left(a_{kl}^{ij} \right)_{\substack{1 \leq i, j \leq q \\ 1 \leq k, l \leq q}}, & B &= (B^{ij})_{\substack{1 \leq i \leq q \\ 1 \leq j \leq q' - q}} = \left(b_{kl}^{ij} \right)_{\substack{1 \leq i \leq q' - q \\ 1 \leq k, l \leq q}}, \\ C &= (C^{ij})_{\substack{1 \leq i \leq q \\ 1 \leq j \leq q' - q}} = \left(c_{kl}^{ij} \right)_{\substack{1 \leq i \leq q' - q \\ 1 \leq k, l \leq q}}, & D &= (D^{ij})_{1 \leq i, j \leq q' - q} = \left(d_{kl}^{ij} \right)_{1 \leq k, l \leq q}. \end{aligned}$$

We replace (4.3) in the equations (3.6) in order to identify the terms of bidegree 2 in (Z, \bar{Z}) . We denote by $F_1^{(1)}(Z)$ and $F_2^{(1)}(Z)$ the linear parts in Z of the matrices $F_1(Z, W)$ and $F_2(Z, W)$. We obtain

$$(4.5) \quad \begin{aligned} \frac{A \otimes W - \overline{A \otimes W}^t}{2\sqrt{-1}} &= \langle F_1^{(1)}(Z), F_1^{(1)}(Z) \rangle, & \frac{B \otimes W - \overline{B \otimes W}^t}{2\sqrt{-1}} &= \langle F_1^{(1)}(Z), F_2^{(1)}(Z) \rangle, \\ \frac{C \otimes W - \overline{C \otimes W}^t}{2\sqrt{-1}} &= \langle F_2^{(1)}(Z), F_1^{(1)}(Z) \rangle, & \frac{D \otimes W - \overline{D \otimes W}^t}{2\sqrt{-1}} &= \langle F_2^{(1)}(Z), F_2^{(1)}(Z) \rangle. \end{aligned}$$

We rewrite the diagonal entries separately from the non-diagonal entries in (3.3). We obtain

$$(4.6) \quad \begin{aligned} \frac{w_{kl} - \overline{w_{lk}}}{2\sqrt{-1}} &= \langle Z_k, Z_l \rangle, \quad \text{for all } k \neq l \text{ and } k, l = 1, \dots, q, \\ \text{Im } w_{kk} &= \langle Z_k, Z_k \rangle, \quad \text{for all } k = 1, \dots, q. \end{aligned}$$

The matrix \tilde{V}' must be invertible. Otherwise, it exists a linear non-trivial combination of the rows defining the matrix \tilde{V}' : provided r_1, \dots, r_N the rows of the matrix \tilde{V} , there exist $\beta_{11}, \beta_{12}, \dots, \beta_{1q_1}, \dots, \beta_{q_1 1}, \beta_{q_1 2}, \dots, \beta_{q_1 q_1} \in \mathbb{C}$ not all vanishing such that

$$(4.13) \quad \begin{aligned} & \sum_{i,j=1}^{q_1} \beta_{ij} \cdot (\alpha_{i1} \cdot r_j, \alpha_{i2} \cdot r_j, \dots, \alpha_{iq_1} \cdot r_j) = 0. \text{ In particular, we obtain:} \\ & 0 = \left\langle \sum_{i,j=1}^{q_1} \beta_{ij} \cdot (\alpha_{i1} \cdot r_j, \alpha_{i2} \cdot r_j, \dots, \alpha_{iq_1} \cdot r_j), (\alpha_{11} \cdot r_1, \alpha_{12} \cdot r_1, \dots, \alpha_{1q_1} \cdot r_1) \right\rangle \\ & \quad \left\langle \sum_{i=1}^{q_1} \beta_{i1} \cdot (\alpha_{i1} \cdot r_1, \alpha_{i2} \cdot r_1, \dots, \alpha_{iq_1} \cdot r_1), (\alpha_{11} \cdot r_1, \alpha_{12} \cdot r_1, \dots, \alpha_{1q_1} \cdot r_1) \right\rangle. \end{aligned}$$

Such procedure describes a linear non-trivial combination of columns defining A_1 . Contradiction, because the matrix A_1 is invertible. Therefore the matrix \tilde{V}' must be invertible. They motivate the change of coordinates $(\tilde{W}, \tilde{Z}) = (A_1 \otimes W, \tilde{V}' \otimes Z)$. We obtain

$$(4.14) \quad \begin{aligned} & \frac{\tilde{W} - \overline{\tilde{W}}^t}{2\sqrt{-1}} = \langle F_1^{(1)}(\tilde{V}'^{-1} \otimes \tilde{Z}), F_1^{(1)}(\tilde{V}'^{-1} \otimes \tilde{Z}) \rangle, \quad \frac{\tilde{B} \otimes (\tilde{W} - \overline{\tilde{W}}^t)}{2\sqrt{-1}} = \langle F_1^{(1)}(\tilde{V}'^{-1} \otimes \tilde{Z}), F_2^{(1)}(\tilde{V}'^{-1} \otimes \tilde{Z}) \rangle, \\ & \frac{\tilde{C} \otimes (\tilde{W} - \overline{\tilde{W}}^t)}{2\sqrt{-1}} = \langle F_2^{(1)}(\tilde{V}'^{-1} \otimes \tilde{Z}), F_1^{(1)}(\tilde{V}'^{-1} \otimes \tilde{Z}) \rangle, \quad \frac{\tilde{D} \otimes (\tilde{W} - \overline{\tilde{W}}^t)}{2\sqrt{-1}} = \langle F_2^{(1)}(\tilde{V}'^{-1} \otimes \tilde{Z}), F_2^{(1)}(\tilde{V}'^{-1} \otimes \tilde{Z}) \rangle, \end{aligned}$$

where we have used by (3.10) and (4.4) the following matrices

$$(4.15) \quad \begin{aligned} & \tilde{B} = B \diamond A_1^{-1} \in \mathcal{M}_{q_1(q'-q_1) \times q_1^2}(\mathbb{C}) \text{ of entries } \left\langle \left(\begin{matrix} c_{kl}^{ij} \\ 1 \leq k, l \leq q_1 \\ 1 \leq i, j \leq q' - q_1 \end{matrix} \right), (\overline{A_1^{-1}})^t \right\rangle, \\ & \tilde{C} = C \diamond A_1^{-1} \in \mathcal{M}_{q_1^2 \times q_1(q'-q_1)}(\mathbb{C}) \text{ of entries } \left\langle \left(\begin{matrix} b_{kl}^{ij} \\ 1 \leq k, l \leq q_1 \\ 1 \leq i, j \leq q' - q_1 \end{matrix} \right), (\overline{A_1^{-1}})^t \right\rangle, \\ & \tilde{D} = D \diamond A_1^{-1} \in \mathcal{M}_{q_1(q'-q_1) \times q_1(q'-q_1)}(\mathbb{C}) \text{ of entries } \left\langle \left(\begin{matrix} d_{kl}^{ij} \\ 1 \leq k, l \leq q_1 \\ 1 \leq i, j \leq q' - q_1 \end{matrix} \right), (\overline{A_1^{-1}})^t \right\rangle, \end{aligned}$$

because \diamond defines an obvious rule of multiplication of matrices. They have sense in the light of (2.2), because the matrices $(\overline{A_1^{-1}})^t$, B , C and D may be seen as vectors. Moreover, the matrices (4.15) may be assumed null using a suitable change of coordinates. It suffices to identify terms of similar bi-degrees the second equation in (4.14) in the light of the first equation in (4.14). Provided the rows $\tilde{Z}_1^{(1)}, \dots, \tilde{Z}_1^{(q_1)}$ of the matrix \tilde{Z}_1 , the equations (4.14) transform using the change of coordinates

$$\begin{pmatrix} (\tilde{w}_{ij})_{\substack{1 \leq i \leq q_1 \\ 1 \leq j \leq q_1}} & (\tilde{w}_{ij})_{\substack{1 \leq i \leq q - q_1 \\ 1 \leq j \leq q_1}} \\ (\tilde{w}_{ij})_{\substack{1 \leq i \leq q - q_1 \\ 1 \leq j \leq q_1}} & (\tilde{w}_{ij})_{\substack{1 \leq i \leq q - q_1 \\ 1 \leq j \leq q - q_1}} \end{pmatrix} := \begin{pmatrix} (\tilde{w}_{ij})_{\substack{1 \leq i \leq q_1 \\ 1 \leq j \leq q_1}} & (\tilde{w}_{ij})_{\substack{1 \leq i \leq q - q_1 \\ 1 \leq j \leq q_1}} - \tilde{B} \otimes \tilde{W} \\ (\tilde{w}_{ij})_{\substack{1 \leq i \leq q - q_1 \\ 1 \leq j \leq q_1}} - \tilde{C} \otimes \tilde{W} & (\tilde{w}_{ij})_{\substack{1 \leq i \leq q - q_1 \\ 1 \leq j \leq q - q_1}} - \tilde{D}_1 \otimes \tilde{W} \end{pmatrix}, \quad \begin{pmatrix} \tilde{Z}_1 \\ \tilde{Z}_2 \end{pmatrix} := \begin{pmatrix} \tilde{Z}_1 \\ \tilde{Z}_2 - \mathcal{L}(\tilde{Z}_1) \end{pmatrix},$$

where \tilde{D}_1 is a suitable matrix and $\mathcal{L}(\tilde{Z}_1)$ is the matrix of rows $\mathcal{L}^{(1)}(\tilde{Z}_1), \dots, \mathcal{L}^{(q-q_1)}(\tilde{Z}_1)$ defined by

$$\mathcal{L}^{(k)}(\tilde{Z}_1) = \sum_{l=1}^{q_1} b_{1l}^{lk} \tilde{Z}_1^{(l)}, \quad \text{for all } k = 1, \dots, q - q_1.$$

Let $R_1(\tilde{Z}), \dots, R_{q_1}(\tilde{Z})$ be the rows of the matrix $F_1^{(1)}(\tilde{V}'^{-1} \otimes \tilde{Z})$. We write $R_k(\tilde{Z}) = \sum_{i=1}^{q_1} R_k(\tilde{Z}_i)$, for all $k = 1, \dots, q_1$. We obtain

$$\langle \tilde{Z}_i, \tilde{Z}_j \rangle = \sum_{k,l=1}^{q_1} \langle R_k(\tilde{Z}_j), R_l(\tilde{Z}_j) \rangle, \quad \text{for all } i, j = 1, \dots, q_1.$$

Therefore $R_j(\tilde{Z}) = R_j(\tilde{Z}_j)$ and $\langle R_i(\tilde{Z}_i), R_j(\tilde{Z}_j) \rangle = \delta_i^j \langle \tilde{Z}_i, \tilde{Z}_j \rangle$, for all $i, j = 1, \dots, q_1$. We consider the sets of rows

$$\{\alpha_1(i), \alpha_2(i), \dots, \alpha_N(i)\}_{i=1, \dots, q} \text{ in } \mathbb{C}^N \text{ such that } \langle \alpha_u(i), \alpha_l(j) \rangle = \delta_u^l \cdot \delta_i^j, \quad \text{for all } i, j = 1, \dots, q_1 \text{ and } u, l = 1, \dots, N.$$

The Procedure of Baouendi-Huang[1] provides the sets of rows defined by

$$\left\{ \alpha_1(i), \alpha_2(i), \dots, \alpha_N(i), \alpha_{N+1}^*(i), \dots, \alpha_{2N}^*(i) \right\}_{i=1, \dots, q_1} \text{ defining orthonormal bases in } \mathbb{C}^{2N} \text{ using a set of matrices } \left\{ \tilde{\mathcal{A}}_i \right\}_{i=1, \dots, q_1}.$$

Defining $\tau^*(\tilde{Z}) = \tilde{Z}^*$, we use the composition $F_1 := \tau^* \circ F_1$, provided the matrix \tilde{Z}^* of rows defined by

$$\tilde{Z}_i^* = \tilde{Z}_i \tilde{\mathcal{A}}_i^{-1} \text{ and } \tilde{Z}_{q+j}^* = \tilde{Z}_{q+j}, \quad \text{for all } i = 1, \dots, q_1 \text{ and } j = 1, \dots, q' - q_1.$$

Contradiction, because (F, G) is an Embedding. Therefore $q = q_1$. The proof is finished. \square

5. Analogues of the normalizations (2.5) from Huang[10]

In order to understand the structure of the Formal Embedding (3.5), we write

$$(5.1) \quad \begin{pmatrix} F_1(Z, W) \\ F_2(Z, W) \end{pmatrix} = \begin{pmatrix} Z \\ 0 \end{pmatrix} + A \otimes W + \mathcal{O}(3),$$

with respect to the product of matrices

$$(5.2) \quad A \otimes W = \left(\sum_{k,l=1}^q a_{kl}^{ij} w_{kl} \right)_{\substack{1 \leq i \leq q' \\ 1 \leq j \leq N'}}, \quad \text{provided the matrix } A = \left(a_{\beta}^{\alpha} \right)_{\substack{1 \leq \alpha \leq q'N' \\ 1 \leq \beta \leq qq'}} \in \mathcal{M}_{qq' \times qN'}(\mathbb{C}).$$

Moreover, the Formal Embedding (3.5) depends on the following parameters

$$(5.3) \quad r_{ab}^{ij} = \frac{1}{2} \left(\frac{\partial^2 g_{ji}}{\partial w_{aa} \partial w_{bb}}(0) + \overline{\frac{\partial^2 g_{ij}}{\partial w_{aa} \partial w_{bb}}(0)} \right), \quad \text{for all } a, b, i, j = 1, \dots, q.$$

The matrix $\{r_{ab}^{ij}\}_{\substack{1 \leq i, j \leq q \\ 1 \leq a, b \leq q}}$ induces by (3.8) the matrix

$$(5.4) \quad R(W) = \begin{pmatrix} \begin{pmatrix} r_{11}^{11} \cdot w_{11} & r_{12}^{11} \cdot w_{12} & \dots & r_{1q}^{11} \cdot w_{1q} \\ r_{11}^{12} \cdot w_{11} & r_{12}^{12} \cdot w_{12} & \dots & r_{1q}^{12} \cdot w_{1q} \\ \vdots & \vdots & \ddots & \vdots \\ r_{11}^{1q} \cdot w_{11} & r_{12}^{1q} \cdot w_{12} & \dots & r_{1q}^{1q} \cdot w_{1q} \end{pmatrix} & \begin{pmatrix} \dots \\ \dots \\ \vdots \\ \dots \end{pmatrix} & \begin{pmatrix} r_{q1}^{11} \cdot w_{q1} & r_{q2}^{11} \cdot w_{q2} & \dots & r_{qq}^{11} \cdot w_{qq} \\ r_{q1}^{12} \cdot w_{q1} & r_{q2}^{12} \cdot w_{q2} & \dots & r_{qq}^{12} \cdot w_{qq} \\ \vdots & \vdots & \ddots & \vdots \\ r_{q1}^{1q} \cdot w_{q1} & r_{q2}^{1q} \cdot w_{q2} & \dots & r_{qq}^{1q} \cdot w_{qq} \end{pmatrix} \\ \begin{pmatrix} \vdots & \vdots & \ddots & \vdots \end{pmatrix} & \begin{pmatrix} \vdots \\ \vdots \\ \ddots \\ \vdots \end{pmatrix} & \begin{pmatrix} \vdots & \vdots & \ddots & \vdots \end{pmatrix} \\ \begin{pmatrix} r_{11}^{q1} \cdot w_{11} & r_{12}^{q1} \cdot w_{12} & \dots & r_{1q}^{q1} \cdot w_{1q} \\ r_{11}^{q2} \cdot w_{11} & r_{12}^{q2} \cdot w_{12} & \dots & r_{1q}^{q2} \cdot w_{1q} \\ \vdots & \vdots & \ddots & \vdots \\ r_{11}^{qq} \cdot w_{11} & r_{12}^{qq} \cdot w_{12} & \dots & r_{1q}^{qq} \cdot w_{1q} \end{pmatrix} & \begin{pmatrix} \dots \\ \dots \\ \vdots \\ \dots \end{pmatrix} & \begin{pmatrix} r_{q1}^{q1} \cdot w_{q1} & r_{q2}^{q1} \cdot w_{q2} & \dots & r_{qq}^{q1} \cdot w_{qq} \\ r_{q1}^{q2} \cdot w_{q1} & r_{q2}^{q2} \cdot w_{q2} & \dots & r_{qq}^{q2} \cdot w_{qq} \\ \vdots & \vdots & \ddots & \vdots \\ r_{q1}^{qq} \cdot w_{q1} & r_{q2}^{qq} \cdot w_{q2} & \dots & r_{qq}^{qq} \cdot w_{qq} \end{pmatrix} \end{pmatrix}.$$

Since W may be identified to the first $q^2 \times q^2$ block of the matrix W' in (3.3), we use the identification

$$R(W) \equiv \begin{pmatrix} R(W) & \mathcal{O}_{q^2 \times q^2 - q^2} \\ \mathcal{O}_{q^2 - q^2 \times q^2} & I_{q^2 - q^2} \end{pmatrix}.$$

In order to eliminate the parameters A and $R(W)$ from (3.5), we follow Baouendi-Huang[1] and Chern-Moser[4]:

5.1. Elimination of the matrix R . We search for a transformation as

$$(5.5) \quad T_1(W', Z') = \left(\frac{1}{I_{q^2} + R(W)} \otimes W', V(W', Z') \right), \quad \text{for a fit matrix } V(W', Z').$$

The transformation (5.5) preserves the BSD-Model \mathcal{M}' from (3.3). Provided $\text{Im}W' = Z' \overline{Z'}^t$, we obtain

$$(5.6) \quad \frac{\frac{1}{I_{q^2} + R(W)} \otimes W' - \left(\frac{1}{I_{q^2} + R(W)} \otimes W' \right)^t}{2\sqrt{-1}} = (V(W', Z')) \cdot \overline{(V(W', Z'))}^t.$$

We make fit computations in (5.6). Provided $\text{Im}W' = Z' \overline{Z'}^t$, we obtain

$$\tilde{A} \otimes \frac{\left(\left(\frac{1}{I_{q^2} + R(W)} \right)^t \otimes W' - \left(\frac{1}{I_{q^2} + R(W)} \otimes W' \right)^t \right)}{2\sqrt{-1}} = (V(W', Z')) \cdot \overline{(V(W', Z'))}^t,$$

because the matrix \tilde{A} is hermitian:

$$(5.7) \quad \tilde{A} := \tilde{A}(W', \overline{W}') = \frac{1}{(I_{q^2} + R(W)) \cdot \overline{(I_{q^2} + R(W))}^t}.$$

On the other hand, we have

$$\left(\frac{1}{I_{q^2} + R(W)} \right)^t \otimes W' - \left(\frac{1}{I_{q^2} + R(W)} \otimes W' \right)^t = W' - \overline{W'}^t + \overline{R(W)}^t \otimes W' - \overline{(R(W))^t \otimes W'}^t = W' - \overline{W'}^t,$$

because $\overline{R(W)}^t \otimes W' - \overline{(R(W))^t \otimes W'}^t = 0$ in the light of (5.2) and (5.3). Indeed, we have

$$\overline{(R(W))^t \otimes W'}^t = \left(\sum_{k,l=1}^q \overline{r_{lk}^{ij} w_{lk} w_{kl}} \right)_{1 \leq i, j \leq q^2}^t = \left(\sum_{k,l=1}^q r_{lk}^{ij} w_{lk} \overline{w_{kl}} \right)_{1 \leq i, j \leq q^2} = \overline{R(W)}^t \otimes W'.$$

Provided $\text{Im}W' = Z' \overline{Z'}^t$, the equation (5.6) is equivalent to

$$(5.8) \quad \tilde{A}(W', \overline{W}') \otimes \langle Z', Z' \rangle = (V(W', Z')) \cdot \overline{(V(W', Z'))}^t.$$

In order to find a fit matrix $V(W')$, we write the formal expansion

$$\frac{1}{I_{q/2} + R(W)} = \left(r_{\alpha'}^{\beta'}(W') \right)_{1 \leq \alpha', \beta' \leq q/2}.$$

In the light of (5.8), it suffices to take

$$(5.9) \quad V(W', Z') = \begin{pmatrix} \sum_{k=1}^{q'} r_{k1}^{11}(W') \cdot Z'_k & \sum_{k=1}^{q'} r_{k2}^{11}(W') \cdot Z'_k & \cdots & \sum_{k=1}^{q'} r_{kq'}^{11}(W') \cdot Z'_k \\ \vdots & \vdots & \ddots & \vdots \\ \sum_{k=1}^{q'} r_{k1}^{1q'}(W') \cdot Z'_k & \sum_{k=1}^{q'} r_{k2}^{1q'}(W') \cdot Z'_k & \cdots & \sum_{k=1}^{q'} r_{kq'}^{1q'}(W') \cdot Z'_k \\ \vdots & \vdots & \ddots & \vdots \\ \sum_{k=1}^{q'} r_{k1}^{q'1}(W') \cdot Z'_k & \sum_{k=1}^{q'} r_{k2}^{q'1}(W') \cdot Z'_k & \cdots & \sum_{k=1}^{q'} r_{kq'}^{q'1}(W') \cdot Z'_k \\ \vdots & \vdots & \ddots & \vdots \\ \sum_{k=1}^{q'} r_{k1}^{q'q'}(W') \cdot Z'_k & \sum_{k=1}^{q'} r_{k2}^{q'q'}(W') \cdot Z'_k & \cdots & \sum_{k=1}^{q'} r_{kq'}^{q'q'}(W') \cdot Z'_k \end{pmatrix}.$$

We pursue the following computations

$$(5.10) \quad \begin{aligned} \partial_{w_{aa}} \left(\frac{1}{I_{q/2} + R(W)} \cdot G(W, Z) \right) &= \frac{1}{I_{q/2} + R(W)} \cdot \partial_{w_{aa}}(G(W, Z)) + \partial_{w_{aa}} \left(\frac{1}{I_{q/2} + R(W)} \right) \cdot G(W, Z), \quad \text{for all } a = 1, \dots, q, \\ \partial_{w_{bb}} \left(\frac{1}{I_{q/2} + R(W)} \cdot G(W, Z) \right) &= \frac{1}{I_{q/2} + R(W)} \cdot \partial_{w_{bb}}(G(W, Z)) + \partial_{w_{bb}} \left(\frac{1}{I_{q/2} + R(W)} \right) \cdot G(W, Z), \quad \text{for all } b = 1, \dots, q, \end{aligned}$$

In order to further differentiate the left-hand side in (5.10), we compute

$$(5.11) \quad \begin{aligned} \partial_{w_{bb}} \left(\frac{1}{I_{q/2} + R(W)} \cdot \partial_{w_{aa}}(G(W, Z)) \right) &= \partial_{w_{bb}} \left(\frac{1}{I_{q/2} + R(W)} \right) \cdot \partial_{w_{aa}}(G(W, Z)) + \frac{\partial_{w_{bb}w_{aa}}^2(G(W, Z))}{I_{q/2} + R(W)}, \\ \partial_{w_{aa}} \left(\frac{1}{I_{q/2} + R(W)} \cdot \partial_{w_{bb}}(G(W, Z)) \right) &= \partial_{w_{aa}} \left(\frac{1}{I_{q/2} + R(W)} \right) \cdot \partial_{w_{bb}}(G(W, Z)) + \frac{\partial_{w_{aa}w_{bb}}^2(G(W, Z))}{I_{q/2} + R(W)}, \\ \partial_{w_{bb}} \left(\partial_{w_{aa}} \left(\frac{1}{I_{q/2} + R(W)} \right) \cdot G(W, Z) \right) &= \partial_{w_{bb}w_{aa}}^2 \left(\frac{1}{I_{q/2} + R(W)} \right) \cdot G(W, Z) + \partial_{w_{aa}} \left(\frac{1}{I_{q/2} + R(W)} \right) \cdot \partial_{w_{bb}}(G(W, Z)), \\ \partial_{w_{aa}} \left(\partial_{w_{bb}} \left(\frac{1}{I_{q/2} + R(W)} \right) \cdot G(W, Z) \right) &= \partial_{w_{aa}w_{bb}}^2 \left(\frac{1}{I_{q/2} + R(W)} \right) \cdot G(W, Z) + \partial_{w_{bb}} \left(\frac{1}{I_{q/2} + R(W)} \right) \cdot \partial_{w_{aa}}(G(W, Z)), \end{aligned}$$

according to the differentiation rule of a product of matrices, for all $a, b = 1, \dots, q$. In particular, we obtain

$$(5.12) \quad \partial_{w_{bb}w_{aa}}^2 \left(\frac{1}{I_{q/2} + R(W)} \right) \cdot G(W, Z) \Big|_{\substack{Z=O_{q \times N} \\ W=O_{q \times q}}} = \partial_{w_{aa}w_{bb}}^2 \left(\frac{1}{I_{q/2} + R(W)} \right) \cdot G(W, Z) \Big|_{\substack{Z=O_{q \times N} \\ W=O_{q \times q}}} = O_{q \times q}, \quad \text{for all } a, b = 1, \dots, q.$$

$$(5.13) \quad \partial_{w_{aa}}(G(W, Z)) \Big|_{\substack{Z=O_{q \times N} \\ W=O_{q \times q}}} = \begin{pmatrix} 0 & \cdots & 0 & \cdots & 0 \\ \vdots & \ddots & \vdots & \ddots & \vdots \\ 0 & \cdots & 1 & \cdots & 0 \\ \vdots & \ddots & \vdots & \ddots & \vdots \\ 0 & \cdots & 0 & \cdots & 0 \end{pmatrix} \quad \text{and} \quad \partial_{w_{bb}}(G(W, Z)) \Big|_{\substack{Z=O_{q \times N} \\ W=O_{q \times q}}} = \begin{pmatrix} 0 & \cdots & 0 & \cdots & 0 \\ \vdots & \ddots & \vdots & \ddots & \vdots \\ 0 & \cdots & 1 & \cdots & 0 \\ \vdots & \ddots & \vdots & \ddots & \vdots \\ 0 & \cdots & 0 & \cdots & 0 \end{pmatrix},$$

where the (a, a) -entry and the (b, b) -entry are 1, otherwise the entries are vanishing, for all $a, b = 1, \dots, q$.

We continue using notations as (3.7) in order to define $(G^*, F^*) = T_1 \circ (G, F)$ in the light of (5.5). We obtain

$$(5.14) \quad \left(\frac{\partial^2(g_{ij}^*(Z, W))}{\partial w_{aa} \partial w_{bb}} + \frac{\partial^2(g_{ji}^*(Z, W))}{\partial w_{aa} \partial w_{bb}} \right) \Big|_{\substack{Z=O_{q \times N} \\ W=O_{q \times q}}} = 0, \quad \text{for all } a, b = 1, \dots, q \text{ and } i, j = 1, \dots, q',$$

according to (5.11), (5.12) and (5.13) in the light of the following evaluations

$$\frac{\partial_{w_{bb}w_{aa}}^2(G(W, Z))}{I_{q/2} + R(W)} \Big|_{\substack{Z=O_{q \times N} \\ W=O_{q \times q}}} = \partial_{w_{bb}w_{aa}}^2(G(W, Z)) \Big|_{\substack{Z=O_{q \times N} \\ W=O_{q \times q}}}, \quad \text{for all } a, b = 1, \dots, q,$$

$$\begin{aligned} \overline{\left(\frac{\partial_{w_{aa}w_{bb}}^2 (G(W, Z))}{I_{q'^2} + R(W)} \Big|_{\substack{Z=O_{q \times N} \\ W=O_{q \times q}}} \right)^t} &= \overline{\left(\partial_{w_{aa}w_{bb}}^2 (G(W, Z)) \Big|_{\substack{Z=O_{q \times N} \\ W=O_{q \times q}}} \right)^t}, \quad \text{for all } a, b = 1, \dots, q, \\ \partial_{w_{bb}} \left(\frac{1}{I_{q'^2} + R(W)} \right) \Big|_{W=O_{q \times q}} &= -\partial_{w_{bb}} (R(W)) \Big|_{W=O_{q \times q}}, \quad \text{for all } a, b = 1, \dots, q, \\ \overline{\left(\partial_{w_{aa}} \left(\frac{1}{I_{q'^2} + R(W)} \right) \Big|_{W=O_{q \times q}} \right)^t} &= -\overline{\left(\partial_{w_{aa}} (R(W)) \Big|_{W=O_{q \times q}} \right)^t}, \quad \text{for all } a, b = 1, \dots, q. \end{aligned}$$

5.2. Elimination of the Matrix A: We search for a transformation as

$$(5.15) \quad T_2(W', Z') = \left(\frac{1}{I_{q'^2} + \mathcal{L}(W', Z')} \otimes W', V(W', Z', Z' - A \otimes W') \right), \quad \text{for fit matrices } V(W', Z', Z') \text{ and } \mathcal{L}(W', Z').$$

The matrix $\mathcal{L}(W', Z')$ is chosen such that

$$(5.16) \quad \mathcal{L}(W', Z') = I_{q'} + 2\sqrt{-1}C(Z') - 2\sqrt{-1}B(W'), \quad \text{for fit matrices } C(Z') \text{ and } B(W').$$

The transformation (5.15) preserves the BSD-Model \mathcal{M}' from (3.3). Provided $\text{Im}W' = Z'\overline{Z'}^t$, we obtain

$$(5.17) \quad \frac{\frac{1}{I_{q'^2} + \mathcal{L}(W', Z')} \otimes W' - \overline{\left(\frac{1}{I_{q'^2} + \mathcal{L}(W', Z')} \otimes W' \right)^t}}{2\sqrt{-1}} = (V(W', Z', Z' - A \otimes W')) \cdot \overline{(V(W', Z', Z' - A \otimes W'))^t}.$$

We make fit computations in (5.17). Provided $\text{Im}W' = Z'\overline{Z'}^t$, we obtain

$$(5.18) \quad \tilde{A} \otimes \frac{W' - \overline{W'}^t + \mathcal{L}(W', Z')^t \otimes W' - \mathcal{L}(W', Z') \otimes \overline{W'}^t}{2\sqrt{-1}} = (V(W', Z', Z' - A \otimes W')) \cdot \overline{(V(W', Z', Z' - A \otimes W'))^t},$$

where we have used the hermitian matrix

$$(5.19) \quad \tilde{A} := \tilde{A}(W', Z', \overline{W'}, \overline{Z'}) = \frac{1}{(I_{q'^2} + 2\sqrt{-1}C(Z') - 2\sqrt{-1}B(W')) \cdot \overline{(I_{q'^2} + 2\sqrt{-1}C(Z') - 2\sqrt{-1}B(W'))^t}}.$$

Provided $\text{Im}W' = Z'\overline{Z'}^t$, we search for a matrix $V(W', Z')$ such that

$$(5.20) \quad \tilde{A}(W', Z', \overline{W'}, \overline{Z'}) \otimes \frac{W' - \overline{W'}^t}{2\sqrt{-1}} = (V(W', Z', Z')) \cdot \overline{(V(W', Z', Z'))^t}.$$

In order to find a fit matrix $V(W', Z')$, we write the formal expansion

$$\frac{1}{I_{q'^2} + \mathcal{L}(W', Z')} = \left(l_{\alpha'}^{\beta'}(W', Z') \right)_{\substack{1 \leq \alpha' \leq q'^2 \\ 1 \leq \beta' \leq q'N'}}.$$

In the light of (5.18), it suffices to take

$$(5.21) \quad V(W', Z', Z') = \begin{pmatrix} \sum_{k=1}^{q'} l_{k1}^{11}(W', Z') \cdot Z'_k & \sum_{k=1}^{q'} l_{k2}^{11}(W', Z') \cdot Z'_k & \dots & \sum_{k=1}^{q'} l_{kq'}^{11}(W', Z') \cdot Z'_k \\ \vdots & \vdots & \ddots & \vdots \\ \sum_{k=1}^{q'} l_{k1}^{1q'}(W', Z') \cdot Z'_k & \sum_{k=1}^{q'} l_{k2}^{1q'}(W', Z') \cdot Z'_k & \dots & \sum_{k=1}^{q'} l_{kq'}^{1q'}(W', Z') \cdot Z'_k \\ \vdots & \vdots & \ddots & \vdots \\ \sum_{k=1}^{q'} l_{k1}^{q'1}(W', Z') \cdot Z'_k & \sum_{k=1}^{q'} l_{k2}^{q'1}(W', Z') \cdot Z'_k & \dots & \sum_{k=1}^{q'} l_{kq'}^{q'1}(W', Z') \cdot Z'_k \\ \vdots & \vdots & \ddots & \vdots \\ \sum_{k=1}^{q'} l_{k1}^{N'q'}(W', Z') \cdot Z'_k & \sum_{k=1}^{q'} l_{k2}^{N'q'}(W', Z') \cdot Z'_k & \dots & \sum_{k=1}^{q'} l_{kq'}^{N'q'}(W', Z') \cdot Z'_k \end{pmatrix}.$$

Provided $\text{Im}W' = Z'\overline{Z'}^t$, we search for a matrix $C(Z')$ as

$$(5.22) \quad C(Z') = \left(c_{kl}^{ij}(Z') \right)_{\substack{1 \leq i, j \leq q'^2 \\ 1 \leq k, l \leq q'^2}} \text{ such that } \tilde{A}(W', Z', \overline{W'}, \overline{Z'}) \otimes (C(Z') \otimes \overline{W'}^t) = (V(W', Z', Z')) \cdot \overline{(V(W', Z', Z'))^t}.$$

It suffices to have $C(Z') \otimes \overline{W'}^t = Z' \cdot \overline{(A \otimes W')}^t$ in the light of (5.2) and (5.21). Therefore

$$\sum_{k, l=1}^{q'^2} c_{kl}^{ij}(Z') \overline{w'_{lk}} = \sum_{u=1}^{N'} z'_{iu} \cdot \overline{\left(\sum_{k, l=1}^q a_{kl}^{uj} w'_{kl} \right)} \quad \text{or equivalently } c_{kl}^{ij}(Z') = \sum_{u=1}^{N'} \overline{a_{lk}^{uj}} \cdot z'_{iu}, \quad \text{for all } i, j = 1, \dots, q'.$$

Provided $\text{Im}W' = Z'\overline{Z'}^t$, we search for a matrix $B(W')$ as

$$(5.23) \quad B(W') = \left(\sum_{k',l'=1}^{q'^2} \left(b_{kl}^{ij} \right)_{k'l'} w'_{k'l'} \right)_{1 \leq i,j \leq q'^2}^{1 \leq k,l \leq q'^2} \quad \text{such that } \tilde{A}(W', Z', \overline{W'}, \overline{Z'}) \otimes (B(W') \otimes \overline{W'}^t + \overline{(B(W'))}^t \otimes W'^t) = \\ (V(W', Z', A \otimes W')) \cdot \overline{(V(W', Z', A \otimes W'))}^t.$$

It suffices to have $B(W') \otimes \overline{W'}^t + \overline{(B(W'))}^t \otimes W'^t = (A \otimes W') \cdot \overline{(A \otimes W')}^t$ in the light of (5.2) and (5.21). Therefore

$$(5.24) \quad \left(\sum_{k,l=1}^{q'^2} \sum_{k',l'=1}^{q'^2} \left(\left(b_{kl}^{ij} \right)_{k'l'} w'_{k'l'} \right) \overline{w'_{lk}} + \sum_{k,l=1}^{q'^2} \overline{\left(\sum_{k',l'=1}^{q'^2} \left(b_{lk}^{ji} \right)_{k'l'} w'_{k'l'} \right)} w'_{kl} \right)_{1 \leq i,j \leq q'^2} = \sum_{r=1}^{N'} \left(\sum_{k,l=1}^{q^2} a_{kl}^{ir} w'_{kl} \right) \overline{\left(\sum_{k,l=1}^{q^2} a_{kl}^{rj} w'_{kl} \right)}.$$

We identify similar terms in (5.24). We obtain

$$(5.25) \quad \left(b_{kl}^{ij} \right)_{k'l'} + \overline{\left(b_{lk}^{ji} \right)_{l'k}} = \begin{cases} \sum_{r=1}^{N'} \left(a_{k'l'}^{ir} \cdot \overline{a_{kl}^{rj}} + a_{kl}^{ir} \cdot \overline{a_{k'l'}^{rj}} \right), & \text{for all } i, j, k', l' = 1, \dots, q^2 \text{ and } k, l = 1, \dots, q'^2, \\ 0, & \text{for all } i, j, k, l = 1, \dots, q'^2 \text{ and } k', l' = q^2 + 1, \dots, q'^2. \end{cases}$$

It suffices to summarize (5.20), (5.22) and (5.23) in order to obtain (5.18) using (5.16) in the light of (3.7) and (5.15). In particular, we consider $(G^{**}, F^{**}) = T_2 \circ (G, F)$ in the light of (5.15). We obtain

$$(5.26) \quad \left. \frac{\partial f_{il}^{**}(Z, W)}{\partial w_{ab}} \right|_{\substack{Z=O_{q \times N} \\ W=O_{q \times q}}} = 0, \quad \text{for all } a, b = 1, \dots, q, i = 1, \dots, q' \text{ and } l = 1, \dots, N'.$$

These normalisations (5.14) and (5.26) are crucial in order to study the following:

6. Analogue of the Classical Geometrical Rank

Provided the *BSD*-Models \mathcal{M} and \mathcal{M}' defined by (3.3), we examine using (3.6) the following equations:

$$(6.1) \quad \begin{aligned} \frac{G_{11}^{**}(Z, W) - \overline{G_{11}^{**}(Z, W)}}{2\sqrt{-1}} &= F_1^{**}(Z, W) \overline{F_1^{**}(Z, W)}, & \frac{G_{12}^{**}(Z, W) - \overline{G_{21}^{**}(Z, W)}}{2\sqrt{-1}} &= F_1^{**}(Z, W) \overline{F_2^{**}(Z, W)}, \\ \frac{G_{21}^{**}(Z, W) - \overline{G_{12}^{**}(Z, W)}}{2\sqrt{-1}} &= F_2^{**}(Z, W) \overline{F_1^{**}(Z, W)}, & \frac{G_{22}^{**}(Z, W) - \overline{G_{22}^{**}(Z, W)}}{2\sqrt{-1}} &= F_2^{**}(Z, W) \overline{F_2^{**}(Z, W)}. \end{aligned}$$

The equations (6.1) are studied in order to compute the formal mapping (3.5) from their diagonal entries. In particular, we use standard linearization procedure of constructions of normal forms (see [2],[3],[4],[29],[30]). We extract terms of total degree 4 from (6.1) in respect to suitable linear changes of coordinates. Following Huang-Ji[12], we use the notation

$$(6.2) \quad F_1^{**}(Z, W) = (f^{**}, \varphi^{**})(Z, W).$$

We work with formal power series in (Z, W) defining: the weight of any entry of Z is 1, and the weight to any entry of W is 2. Its weighted degree, denoted by n , is the minimum of the weighted degrees of its homogeneous terms from its formal expansion. We write $H(Z, W) = O(n)$, if $H(Z, W)$ is a formal power series of weighted degree n . More precisely, it follows that

PROPOSITION 6.1. *Up to compositions with suitable linear changes of coordinates preserving \mathcal{M} and \mathcal{M}' , the following holds*

$$(6.3) \quad \left\{ \begin{aligned} f^{**}(Z, W) &= Z + O(3), \\ \varphi^{**}(Z, W) &= O(2), \end{aligned} \right. \quad \text{or the following holds} \quad \left\{ \begin{aligned} f^{**}(Z, W) &= Z + \frac{\sqrt{-1}}{2} \begin{pmatrix} z_{11}w_{11} & 0 & \dots & 0 \\ z_{21}w_{22} & 0 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ z_{q1}w_{qq} & 0 & \dots & 0 \\ 0 & 0 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & 0 \end{pmatrix} + O(4), \\ \varphi^{**}(Z, W) &= \begin{pmatrix} z_{11}z_{11} & z_{11}z_{12} & \dots & z_{11}z_{1N} \\ z_{21}z_{21} & z_{21}z_{22} & \dots & z_{21}z_{2N} \\ \vdots & \vdots & \ddots & \vdots \\ z_{q1}z_{q1} & z_{q1}z_{q2} & \dots & z_{q1}z_{qN} \\ 0 & 0 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & 0 \end{pmatrix} + O(3). \end{aligned} \right.$$

PROOF. We consider using (2.2) homogeneous polynomials of total degree 2 and of degree 1 in Z , denoted by $b_{il}(Z), A_{ij}(Z), (\varphi_{il}^{**}(Z))^{(2)}$ and $a_{ku}^{il}(Z), b_{ku}^{ij}(Z)$, for all $i, j, k, u = 1, \dots, q$ and $l = 1, \dots, N$. We write (6.2) by (2.2), (3.7) and (4.2) as

$$(6.4) \quad \begin{aligned} \varphi_{il}^{**}(Z, W) &= (\varphi_{il}^{**}(Z))^{(2)} + O(3), & \text{for all } i = 1, \dots, q \text{ and } l = 1, \dots, N. \\ f_{il}^{**}(Z, W) &= z_{il} + \sum_{k,u=1}^q a_{ku}^{il}(Z) w_{ku} + b_{il}(Z) + O(4), & \text{for all } i = 1, \dots, q \text{ and } l = 1, \dots, N. \\ g_{ij}^{**}(Z, W) &= w_{ij} + A_{ij}(Z) + \sum_{k,u=1}^q b_{ku}^{ij}(Z) w_{ku} + \sum_{\substack{k,u=1 \\ k',u'=1}}^q D_{kuk'u'}^{ij} w_{ku} w_{k'u'} + O(5), & \text{for all } i, j = 1, \dots, q. \end{aligned}$$

We extract terms up to total degree 4 in (Z, \bar{Z}) from the entries of (6.1) in the light of (6.4). We obtain

$$(6.5) \quad \begin{aligned} \operatorname{Im} \left\{ A_{ij}(Z) + \sum_{k,u=1}^q b_{ku}^{ij}(Z) w_{ku} + \sum_{\substack{k,u=1 \\ k',u'=1}}^q D_{kuk'u'}^{ij} w_{ku} w_{k'u'} - 2\sqrt{-1} \sum_{l=1}^N \bar{z}_{il} \cdot \right. \\ \left. \left(b_{jl}(Z) + \sum_{k,u=1}^q a_{ku}^{jl}(Z) w_{ku} \right) \right\} &= \sum_{l=1}^N (\varphi_{il}^{**}(Z))^{(2)} \overline{(\varphi_{jl}^{**}(Z))^{(2)}}, \quad \text{for all } i, j = 1, \dots, q. \end{aligned}$$

Analysing the homogeneous polynomials of degree 2 in Z from (6.5), we obtain $A_{ij}(Z) = 0$, for all $i, j = 1, \dots, q$. Therefore

$$(6.6) \quad S_1 - S_2 + S_3 + S_4 + S_5 - S_6 - S_7 = \sum_{l=1}^N (\varphi_{il}^{**}(Z))^{(2)} \overline{(\varphi_{jl}^{**}(Z))^{(2)}} + 2\operatorname{Re} \left\{ \sum_{l=1}^N \bar{z}_{il} b_{jl}(Z) \right\}, \quad \text{for all } i, j = 1, \dots, q.$$

We have used the following sums

$$\begin{aligned} S_1 &= \frac{1}{2\sqrt{-1}} \left(\sum_{k,u=1}^q \left(b_{kk}^{ij}(Z) (\operatorname{Re} w_{kk} + \sqrt{-1} \langle Z_k, Z_k \rangle) + D_{kkuu}^{ij} (\operatorname{Re} w_{kk} + \sqrt{-1} \langle Z_k, Z_k \rangle) (\operatorname{Re} w_{uu} + \sqrt{-1} \langle Z_u, Z_u \rangle) \right) \right), \\ S_2 &= \frac{1}{2\sqrt{-1}} \left(\sum_{k,u=1}^q \overline{\left(b_{kk}^{ij}(Z) (\operatorname{Re} w_{kk} + \sqrt{-1} \langle Z_k, Z_k \rangle) + D_{kkuu}^{ij} (\operatorname{Re} w_{kk} + \sqrt{-1} \langle Z_k, Z_k \rangle) (\operatorname{Re} w_{uu} + \sqrt{-1} \langle Z_u, Z_u \rangle) \right)} \right), \\ S_3 &= \frac{1}{2\sqrt{-1}} \left(\sum_{\substack{k,u=1 \\ k \neq u}}^q \left(b_{ku}^{ij}(Z) w_{ku} - \overline{b_{ku}^{ij}(Z)} (w_{uk} - 2\sqrt{-1} \langle Z_u, Z_k \rangle) \right) \right), \\ S_4 &= \frac{1}{2\sqrt{-1}} \left(\sum_{\substack{k,u,k',u'=1 \\ k' \neq u', k \neq u}}^q \left(D_{kuk'u'}^{ij} w_{ku} w_{k'u'} - \overline{D_{kuk'u'}^{ij}} (w_{uk} - 2\sqrt{-1} \langle Z_u, Z_k \rangle) (w_{u'k'} - 2\sqrt{-1} \langle Z_{u'}, Z_{k'} \rangle) \right) \right), \\ S_5 &= \frac{1}{2\sqrt{-1}} \left(\sum_{\substack{k,k',u'=1 \\ k' \neq u'}}^q \left(D_{kkk'u'}^{ij} (\operatorname{Re} w_{kk} + \sqrt{-1} \langle Z_k, Z_k \rangle) w_{k'u'} - \overline{D_{kkk'u'}^{ij}} (\operatorname{Re} w_{kk} + \sqrt{-1} \langle Z_k, Z_k \rangle) (w_{u'k'} - 2\sqrt{-1} \langle Z_{u'}, Z_{k'} \rangle) \right) \right), \\ S_6 &= \sum_{l=1}^N \bar{z}_{il} \left(\sum_{\substack{k,u=1 \\ k \neq u}}^q \left(a_{ku}^{jl}(Z) w_{ku} + a_{kk}^{jl}(Z) (\operatorname{Re} w_{kk} + \sqrt{-1} \langle Z_k, Z_k \rangle) \right) \right), \\ S_7 &= \sum_{l=1}^N z_{il} \left(\sum_{\substack{k,u=1 \\ k \neq u}}^q \left(\overline{a_{ku}^{jl}(Z)} (w_{uk} - 2\sqrt{-1} \langle Z_u, Z_k \rangle) + \overline{a_{kk}^{jl}(Z)} (\operatorname{Re} w_{kk} + \sqrt{-1} \langle Z_k, Z_k \rangle) \right) \right). \end{aligned}$$

Since (5.14) holds and the right-hand side from (6.6) does not depend on products as $(\operatorname{Re} w_{uu}) \cdot (\operatorname{Re} w_{u'u'})$, for all $u, u' = 1, \dots, q$, we focus on the difference $S_1 - S_2$ in the left-hand side from (6.6). We obtain

$$(6.7) \quad D_{uuu'u'}^{ij} = 0, \quad \text{for all } i, j, u, u' = 1, \dots, q,$$

Since the right-hand side in (6.6) does not depend on terms involving Z multiplied by w_{ku} , $(\operatorname{Re} w_{uu})$, for all $k, u = 1, \dots, q$ such that $k \neq u$, the sums S_1, S_2 and S_3 are vanishing in the left-hand side from (6.6). We obtain

$$(6.8) \quad b_{ku}^{ij}(Z) = 0, \quad \text{for all } i, j, k, u = 1, \dots, q.$$

Next, we focus on the second sum from the right-hand side in (6.6). Since the left-hand side from (6.6) does not depend on terms involving terms of bi-degree $(1, 2)$ in (Z, \bar{Z}) , we obtain

$$(6.9) \quad b_{il}(Z) = 0, \quad \text{for all } i = 1, \dots, q \text{ and } l = 1, \dots, N.$$

Since the right-hand side from (6.6) does not depend on products as $w_{u'k'} \cdot \langle Z_u, Z_k \rangle$, for all $k, u, k', u' = 1, \dots, q$ such that $k \neq u, k' \neq u'$, we focus on the fourth sum from left-hand side from (6.6). We obtain

$$(6.10) \quad D_{kk'k'u'}^{ij} = 0, \quad \text{for all } i, j, k, u, k', u' = 1, \dots, q \text{ such that } k \neq u, k' \neq u'.$$

Since the right-hand side from (6.6) does not depend on products as $(\operatorname{Re} w_{kk}) \cdot w_{k'u'}$, for all $k, k', u' = 1, \dots, q$ such that $k' \neq u'$, we obtain

$$(6.11) \quad D_{kkk'u'}^{ij} = \overline{D_{kku'k'}^{ij}}, \quad \text{for all } i, j, k, k', u' = 1, \dots, q \text{ such that } k' \neq u'.$$

Next, we study the coefficient of the product of terms as $\langle Z_k, Z_k \rangle \cdot \langle Z_{u'}, Z_{k'} \rangle$, for all $k, k', u' = 1, \dots, q$ such that $k' \neq u'$, in the both sides from (6.6) in the light of (5.14), (6.10) and (6.11). We obtain

$$(6.12) \quad D_{kkk'u'}^{ij} = 0, \quad \text{for all } i, j, k, k', u' = 1, \dots, q \text{ such that } k' \neq u'.$$

We identify the coefficients of w_{ku} and $\operatorname{Re} w_{kk}$ in S_6 and S_7 from (6.6), for all $k, u = 1, \dots, q$ such that $k \neq u$. We obtain

$$(6.13) \quad \sum_{l=1}^N z_{il} \overline{a_{ku}^{il}(Z)} + \sum_{l=1}^N \overline{z_{jl} a_{uk}^{jl}(Z)} = 0, \quad \text{for all } i, j, k, u = 1, \dots, q.$$

We write $a_{ku}^{il}(Z) = \sum_{i'=1}^q a_{ku}^{il}(Z_{i'})$, for all $i, k, u = 1, \dots, q$ and $l = 1, \dots, N$. The equation (6.13) is equivalent to

$$(6.14) \quad \sum_{l=1}^N \sum_{i'=1}^q z_{il} \overline{a_{ku}^{jl}(Z_{i'})} + \sum_{i'=1}^q \sum_{l=1}^N \overline{z_{jl} a_{uk}^{il}(Z_{i'})} = 0, \quad \text{for all } i, j, k, u = 1, \dots, q.$$

Therefore $a_{ku}^{il}(Z) = a_{ku}^{il}(Z_i)$, for all $i, k, u = 1, \dots, q$ and $l = 1, \dots, N$. The equation (6.6) is equivalent to

$$(6.15) \quad -2\sqrt{-1} \sum_{\substack{k, u=1 \\ k \neq u}}^q \left(\sum_{l=1}^N z_{il} \overline{a_{ku}^{jl}(Z_j)} \right) \langle Z_u, Z_k \rangle + 2\operatorname{Re} \left\{ \sum_{k=1}^q \left(\sum_{l=1}^N z_{il} \sqrt{-1} \cdot \overline{a_{kk}^{jl}(Z_j)} \right) \right\} \cdot \langle Z_k, Z_k \rangle = - \sum_{l=1}^N (\varphi_{il}^{**}(Z))^{(2)} \overline{(\varphi_{jl}^{**}(Z))^{(2)}},$$

for all $i, j = 1, \dots, q$. The first part of (6.3) holds when the right-hand side vanishes in (6.15). Otherwise, we write using (2.7) the expansion

$$(6.16) \quad \varphi_{il}^{**}(Z) = \sum_{i_1, i_2=1}^q \varphi_{il}^{i_1 i_2}(Z_{i_1}, Z_{i_2}), \quad \text{where } \varphi_{il}^{i_1 i_2}(Z_{i_1}, Z_{i_2}) = \sum_{\substack{\alpha, \beta=1 \\ \alpha \leq \beta}}^N b_{\alpha\beta}^{il} z_{i_1\alpha} z_{i_2\beta}, \quad \text{for all } i, i_1, i_2 = 1, \dots, q \text{ and } l = 1, \dots, N.$$

We consider the matrices $\mathcal{B}_{ku}^i = -2\sqrt{-1} (a_{ku}^{i1}, a_{ku}^{i2}, \dots, a_{ku}^{iN})$, for all $k, u, i = 1, \dots, q$. The equation (6.13) is equivalent to

$$(6.17) \quad \langle Z_i, \mathcal{B}_{ku}^j(Z_j) \rangle = \langle \mathcal{B}_{uk}^i(Z_i), Z_j \rangle, \quad \text{for all } k, u, i, j = 1, \dots, q.$$

We collect homogeneous terms from (6.15) using (6.17) and (6.16). We obtain

$$(6.18) \quad \langle Z_i, \mathcal{B}_{ku}^j(Z_j) \rangle \langle Z_k, Z_u \rangle = \sum_{l=1}^N \varphi_{il}^{ik}(Z_i, Z_k) \overline{\varphi_{jl}^{ju}(Z_j, Z_u)}, \quad \text{for all } i, j, k, u = 1, \dots, q.$$

The matrix \mathcal{B}_{kk}^i is hermitian, for all $k, i = 1, \dots, q$. We write the decomposition

$$(6.19) \quad \mathcal{B}_{kk}^i = (U_{kk}^i) \cdot \begin{pmatrix} \alpha_{kk}^{1i} & 0 & \dots & 0 \\ 0 & \alpha_{kk}^{2i} & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & \alpha_{kk}^{Ni} \end{pmatrix} \cdot (U_{kk}^i)^{-1}, \quad \text{provided a suitable unitary matrix } U_{kk}^i, \text{ for all } k, i = 1, \dots, q.$$

In order to implement the strategy of Huang-Ji[12] (see the computations from the pages 226 – 227), we define the vectors

$$B_{\alpha\beta}^k = \begin{pmatrix} b_{\alpha\beta}^{k1} & b_{\alpha\beta}^{k2} & \dots & b_{\alpha\beta}^{kN} \\ ii & ii & & ii \end{pmatrix}, \quad \text{for all } i, k, \alpha, \beta = 1, \dots, q.$$

The set $\{B_{11}^k, B_{12}^k, \dots, B_{1N}^k\}$ is a base in \mathbb{C}^N for some $i \in 1, \dots, q$, for all $k = 1, \dots, q$. It exists a unique $\sigma(i) \in \{1, \dots, q\}$ such that

$$0 \neq \alpha_i := \alpha_{ii}^{1\sigma(i)} = b_{\sigma(i)\sigma(i)}^{i1} \overline{b_{ii}^{i1}} > 0.$$

Otherwise, it would exist two orthogonal bases in \mathbb{C}^N using similar computations in (6.18) like in [12]. In particular, we assume $i = 1$ and $\mathcal{B}_{22}^2 \neq 0$ and $\mathcal{B}_{11}^1 \equiv \dots \mathcal{B}_{qq}^q \equiv 0$. Contradiction, because $\langle Z_2, \mathcal{B}_{22}^2(Z_2) \rangle \cdot \langle Z_2, Z_2 \rangle$ appears only in the right-hand of (6.18). Therefore $\mathcal{B}_{11}^1 \neq 0$. Therefore $\sigma(i) = i$, for all $i = 1, \dots, q$. Moreover, we obtain

$$\begin{aligned} \langle B_{1\beta}^k, B_{1\beta}^{k'} \rangle &> 0, \quad \text{for all } k, k', \beta = 1, \dots, q. \\ \langle B_{1\beta}^k, B_{1\alpha}^{k'} \rangle &= 0, \quad \text{for all } k, k', \alpha, \beta = 1, \dots, q \text{ such that } \alpha \neq \beta. \end{aligned}$$

We return to the equation (6.18). We obtain

$$(6.20) \quad (C_i) \cdot \overline{(C_{i'})^t} = b_{ii}^{i1} \cdot \overline{b_{i'i'}^{i'1}} \cdot I_N, \quad \text{for all } i, i' = 1, \dots, q.$$

We move forward following Huang-Ji[12]. We implement using (6.20) the change of coordinates

$$(W', Z') := \left(W', \begin{pmatrix} (z'_{11}, z'_{12}, \dots, z'_{1N}) \\ (z'_{21}, z'_{22}, \dots, z'_{2N}) \\ \vdots \\ (z'_{q1}, z'_{q2}, \dots, z'_{qN}) \\ (z'_{q+11}, z'_{q+12}, \dots, z'_{q+1N}) \\ \vdots \\ (z'_{q'1}, z'_{q'2}, \dots, z'_{q'N}) \end{pmatrix}, \begin{pmatrix} (z'_{1q+1}, z'_{1q+2}, \dots, z'_{1N}) \cdot \frac{C_1}{\sqrt{\alpha_1}} \\ (z'_{2q+1}, z'_{2q+2}, \dots, z'_{2N}) \cdot \frac{C_2}{\sqrt{\alpha_2}} \\ \vdots \\ (z'_{qq+1}, z'_{qq+2}, \dots, z'_{qN}) \cdot \frac{C_q}{\sqrt{\alpha_q}} \\ (z'_{q+11}, z'_{q+12}, \dots, z'_{q+1N}) \\ \vdots \\ (z'_{q'q+1}, z'_{q'q+2}, \dots, z'_{q'N}) \end{pmatrix} \right).$$

Next, we consider the change of coordinates

$$(W, Z) := \left(\begin{pmatrix} \frac{w_{11}}{\sqrt{\alpha_1} \cdot \sqrt{\alpha_1}} & \frac{w_{12}}{\sqrt{\alpha_1} \cdot \sqrt{\alpha_1}} & \cdots & \frac{w_{1q}}{\sqrt{\alpha_1} \cdot \sqrt{\alpha_q}} \\ \frac{w_{21}}{\sqrt{\alpha_2} \cdot \sqrt{\alpha_1}} & \frac{w_{22}}{\sqrt{\alpha_2} \cdot \sqrt{\alpha_2}} & \cdots & \frac{w_{2q}}{\sqrt{\alpha_2} \cdot \sqrt{\alpha_q}} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{w_{q1}}{\sqrt{\alpha_q} \cdot \sqrt{\alpha_1}} & \frac{w_{q2}}{\sqrt{\alpha_q} \cdot \sqrt{\alpha_2}} & \cdots & \frac{w_{qq}}{\sqrt{\alpha_q} \cdot \sqrt{\alpha_q}} \end{pmatrix}, \begin{pmatrix} \frac{Z_1}{\sqrt{\alpha_1}} \\ \frac{Z_2}{\sqrt{\alpha_2}} \\ \vdots \\ \frac{Z_q}{\sqrt{\alpha_q}} \end{pmatrix} \right).$$

Finally, we consider the change of coordinates

$$(W', Z') = \left(\begin{pmatrix} \sqrt{\alpha_1} \cdot \sqrt{\alpha_1} \cdot w'_{11} & \cdots & \sqrt{\alpha_1} \cdot \sqrt{\alpha_q} \cdot w'_{1q} & \sqrt{\alpha_1} \cdot w'_{1q+1} & \cdots & \sqrt{\alpha_1} \cdot w'_{1q'} \\ \vdots & \ddots & \vdots & \vdots & \ddots & \vdots \\ \sqrt{\alpha_q} \cdot \sqrt{\alpha_1} \cdot w'_{q1} & \cdots & \sqrt{\alpha_q} \cdot \sqrt{\alpha_1} \cdot w'_{qq} & \sqrt{\alpha_q} \cdot w'_{qq+1} & \cdots & \sqrt{\alpha_q} \cdot w'_{qq'} \\ \sqrt{\alpha_1} \cdot w'_{q+11} & \cdots & \sqrt{\alpha_1} \cdot w'_{q+1q} & w'_{q+1q+1} & \cdots & w'_{q+1q'} \\ \vdots & \ddots & \vdots & \vdots & \ddots & \vdots \\ \sqrt{\alpha_1} \cdot w'_{q'1} & \cdots & \sqrt{\alpha_q} \cdot w'_{q'q} & w'_{q'q+1} & \cdots & w'_{q'q'} \end{pmatrix}, \begin{pmatrix} Z'_1 \cdot \sqrt{\alpha_1} \\ \vdots \\ Z'_q \cdot \sqrt{\alpha_q} \\ Z'_{q+1} \\ \vdots \\ Z'_{q'} \end{pmatrix} \right).$$

They preserve the models \mathcal{M} and \mathcal{M}' from (3.3). Their composition provides the second part of (6.3). \square

We are ready to conclude the Classification (1.3) using the following:

7. Application of the Moving Point Trick of Huang[10]

In order to further normalize the embedding (3.5), we recall (2.2),(2.3),(2.4),(3.5) and (3.7). We write the formal expansions

$$(7.1) \quad G(Z, W) = \left(\sum_{J \in \mathbb{N}^{q^2}, I \in \mathbb{N}^{qN}} g_{ij}^{IJ}(Z) W^J \right)_{1 \leq i, j \leq q'} \quad \text{and} \quad F(Z, W) = \left(\sum_{J \in \mathbb{N}^{q^2}, I \in \mathbb{N}^{qN}} f_{kl}^{IJ}(Z) W^J \right)_{\substack{1 \leq l \leq N' \\ 1 \leq k \leq q'}}.$$

Their entries are formal power series, where the coefficients of W are defined by

$$(7.2) \quad g_{ij}^{IJ}(Z) = \sum_{I \in \mathbb{N}^{qN}} c_{ij}^{IJ} Z^I \quad \text{and} \quad f_{kl}^{IJ}(Z) = \sum_{I \in \mathbb{N}^{qN}} d_{kl}^{IJ} Z^I, \quad \text{for all } k, i, j = 1, \dots, q', l = 1, \dots, N' \text{ and } J \in \mathbb{N}^{q^2}.$$

We extract from (3.6) the terms of total degree d in (Z, \bar{Z}) . Provided $\text{Im } W = Z\bar{Z}^t$, we obtain

$$(7.3) \quad \sum_{\substack{J \in \mathbb{N}^{q^2}, I \in \mathbb{N}^{qN} \\ |I|+2|J|=d}} \frac{g_{ij}^{IJ}(Z) W^J - \overline{g_{ji}^{IJ}(Z) W^J}}{2\sqrt{-1}} = \sum_{l=1}^{N'} \sum_{\substack{J_1, J_2 \in \mathbb{N}^{q^2}, I_1, I_2 \in \mathbb{N}^{qN} \\ |I_1|+2|J_1|+|I_2|+2|J_2|=d}} \left(f_{il}^{I_1 J_1}(Z) W^{J_1} \right) \overline{\left(f_{jl}^{I_2 J_2}(Z) W^{J_2} \right)^t}, \quad \text{for all } i, j = 1, \dots, q'.$$

Since the right-hand side from (7.3) is hermitian, we obtain

$$(7.4) \quad g_{ij}^{0J}(Z) = \overline{g_{ji}^{0J}(Z)}, \quad \text{for all } i, j = 1, \dots, q' \text{ and } J \in \mathbb{N}^{q^2}.$$

In particular, (7.4) describes real-valued constants. Following Baouendi-Huang[1] and Huang[10], it follows that

LEMMA 7.1. *Up to compositions with holomorphic automorphisms of \mathcal{M}' , we have*

$$(7.5) \quad G(Z, W) = \begin{pmatrix} W & 0 \\ 0 & 0 \end{pmatrix} \quad \text{and} \quad F(0, W) = 0.$$

PROOF. We define the following transformations

$$\begin{aligned} \sigma_{(Z_0, W_0)}^0(Z, W) &= (Z + Z_0, W + W_0 + 2\sqrt{-1} \langle Z, Z_0 \rangle), \\ \tau_{(Z_0, W_0)}^{(F, G)}(Z^*, W^*) &= \left(Z^* - F(Z_0, W_0), W^* - \overline{G(Z_0, W_0)}^t - 2\sqrt{-1} \langle Z^*, F(Z_0, W_0) \rangle \right). \end{aligned}$$

Provided $P = (Z_0, W_0) \in \mathcal{M}$ close to origin, we define $(F, G)_P = \tau_P^{(F, G)} \circ (F, G) \circ \sigma_P^0 = (F_P, G_P)$. Therefore

$$\sigma_P^0(0) = P, \quad \tau_{(F, G)(P)}^{(F, G)}((F, G)(P)) = 0, \quad \det \left(\frac{\partial G_{11}(W)}{\partial W} \right) (0) \neq 0.$$

We consider using (5.15) the transformation $(\tilde{G}, \tilde{F}) = T_2 \circ (G, F)$, for $T_2 = T_2(P)$. It is defined by subtractions of terms in W from the F -component of the Embedding (3.5). Varying the point $P \in \mathcal{M}$, we obtain

$$(7.6) \quad \tilde{G}(0, W) = \begin{pmatrix} W & 0 \\ 0 & 0 \end{pmatrix} \text{ and } \tilde{F}(0, W) = 0.$$

The decisive argument is motivated by Hamada[9]: we study the coefficients of terms as

$$\begin{aligned} & Z^I (\text{Re } w_{11})^{j_{11}} (w_{12})^{j_{12}} \dots (w_{1q})^{j_{1q}} \cdot \\ & (\bar{w}_{12})^{j_{21}} (\text{Re } w_{22})^{j_{22}} \dots (w_{2q})^{j_{2q}} \cdot \\ & \quad \vdots \quad \quad \quad \ddots \quad \quad \quad \vdots \\ & (\bar{w}_{1q})^{j_{q1}} (\bar{w}_{2q})^{j_{q2}} \dots (\text{Re } w_{qq})^{j_{qq}}, \quad \text{for all } I \in \mathbb{N}^{qN} \text{ and } J \in \mathbb{N}^{q^2}. \end{aligned}$$

We identify coefficients of homogeneous terms in the left-hand side from (7.3) using (7.6). Therefore (7.5) holds. \square

8. Application of the Procedure of Hamada[9]

Before beginning, we write the following weighted formal expansions

- $f_{k1}(Z, W) = z_{k1} + f_{k1}^{(2)}(Z, W) + f_{k1}^{(3)}(Z, W) + \dots + f_{k1}^{(2m)}(Z, W) + f_{k1}^{(2m+1)}(Z, W) + \dots$, for all $k = 1, \dots, q$.
- $f_{kl}(Z, W) = z_{kl} + f_{kl}^{(2)}(Z, W) + f_{kl}^{(3)}(Z, W) + \dots + f_{kl}^{(4m)}(Z, W) + f_{kl}^{(4m+1)}(Z, W) + \dots$, for all $k = 1, \dots, q$ and $l = 2, \dots, N$.
- $\varphi_{kl}(Z, W) = \varphi_{kl}^{(2)}(Z, W) + \varphi_{kl}^{(3)}(Z, W) + \dots + \varphi_{kl}^{(2m)}(Z, W) + \varphi_{kl}^{(2m+1)}(Z, W) + \dots$, for all $k = 1, \dots, q$ and $l = 1, \dots, N$.

In order to compute the F -component from (3.5) using induction, we linearise the diagonal entries in (3.6). Provided $\alpha \in \mathbb{N}^*$, we assume

$$(8.1) \quad \begin{aligned} f_{kl}^{(2)}(Z, W) &= f_{kl}^{(3)}(Z, W) = \dots = f_{kl}^{(\alpha-1)}(Z, W) = 0, \quad \text{for all } k = 1, \dots, q \text{ and } l = 1, \dots, N. \\ \varphi_{kl}^{(2)}(Z, W) &= \varphi_{kl}^{(3)}(Z, W) = \dots = \varphi_{kl}^{(\lfloor \frac{\alpha}{2} \rfloor)}(Z, W) = 0, \quad \text{for all } k = 1, \dots, q \text{ and } l = 1, \dots, N. \end{aligned}$$

We prove that (8.1) holds when α is replaced by $\alpha + 1$. In particular, we write using (2.2) the formal expansions

$$(8.2) \quad f_{kl}^{(\alpha)}(Z, W) = \sum_{\substack{J \in \mathbb{N}^{q^2}, I \in \mathbb{N}^{qN} \\ |I|+2|J|=\alpha}} d_{kl}^{IJ} Z^I W^J \text{ and } \varphi_{kl}^{(\lfloor \frac{\alpha+1}{2} \rfloor)}(Z, W) = \sum_{\substack{J \in \mathbb{N}^{q^2}, I \in \mathbb{N}^{qN} \\ |I|+2|J|=\lfloor \frac{\alpha+1}{2} \rfloor}} \tilde{d}_{kl}^{IJ} Z^I W^J, \quad \text{for all } k = 1, \dots, q \text{ and } l = 1, \dots, N.$$

Since we work with entris of the first BSD -Model from (3.3), we consider the following sums

$$(8.3) \quad \begin{aligned} S_{kl}^\alpha &= \sum_{\substack{J \in \mathbb{N}^{q^2}, I \in \mathbb{N}^{qN} \\ |I|+2|J|=\alpha}} d_{kl}^{IJ} Z^I w_{11}^{j_{11}} w_{12}^{j_{12}} \dots w_{1q}^{j_{1q}} \cdot \\ & w_{21}^{i_{21}} w_{22}^{j_{22}} \dots w_{2q}^{j_{2q}} \cdot \\ & \quad \vdots \quad \quad \quad \ddots \quad \quad \quad \vdots \\ & w_{q1}^{j_{q1}} w_{q2}^{j_{q2}} \dots w_{qq}^{j_{qq}}, \quad \text{for all } k = 1, \dots, q \text{ and } l = 1, \dots, N. \end{aligned}$$

Since we work with substitutions of the first BSD -Model from (3.3), the sums (8.3) reduce to the following sums

$$\begin{aligned} \tilde{S}_{kl}^\alpha &= \sum_{\substack{J \in \mathbb{N}^{q^2}, I \in \mathbb{N}^{qN} \\ |I|+2|J|=\alpha}} d_{kl}^{IJ} Z^I (\text{Re } w_{11})^{j_{11}} (w_{12})^{j_{12}} \dots (w_{1q})^{j_{1q}} \cdot \\ & (\bar{w}_{12})^{j_{21}} (\text{Re } w_{22})^{j_{22}} \dots (w_{2q})^{j_{2q}} \cdot \\ & \quad \vdots \quad \quad \quad \ddots \quad \quad \quad \vdots \\ & (\bar{w}_{1q})^{j_{q1}} (\bar{w}_{2q})^{j_{q2}} \dots (\text{Re } w_{qq})^{j_{qq}}, \quad \text{for all } k = 1, \dots, q \text{ and } l = 1, \dots, N. \end{aligned}$$

Provided $m \in \mathbb{N}^*$, we assume $\alpha \in \{4m, 4m+1, 4m+2, 4m+3\}$ in order to extract weighted terms from (7.3) according to computations similar to (4.2) – (4.9) from Hamada[9] in the light of (3.3). Therefore

$$\begin{aligned} \text{Re} \left\{ \sum_{l=1}^N \bar{z}_{kl} S_{kl}^\alpha \right\} \Big|_{(Z, W) \in \mathcal{M}} &= 0 \text{ and } \text{Re} \left\{ \sum_{l=1}^N \bar{z}_{kl} \cdot \tilde{S}_{kl}^\alpha \Big|_{z_{k1}=0} \right\} = 0, \text{ for all } k = 1, \dots, q. \text{ We obtain} \\ f_{kl}^{(\alpha)}(Z, W) \Big|_{z_{k1}=0} &\equiv \varphi_{kl}^{(\alpha)}(Z, W) \Big|_{z_{k1}=0} \equiv 0, \text{ for all } k = 1, \dots, q \text{ and } l = 1, \dots, N. \end{aligned}$$

In the light of the vanishing of the coefficients of powers of W in (8.2), we obtain (8.1) when (6.3) holds. Therefore $f_{kl}(Z, W) = z_{kl}$ and $\varphi_{kl}(Z, W) = 0$, for all $k = 1, \dots, q$ and $l = 1, \dots, N$. Otherwise, we obtain

- $f_{k1}(Z, W) = z_{k1} \tilde{f}_{k1}(Z, W)$, where $\tilde{f}_{k1}(Z, W)$ are formal mappings, for all $k = 1, \dots, q$.
- $f_{kl}(Z, W) = z_{kl} + z_{k1} \tilde{f}_{kl}(Z, W)$, where $\tilde{f}_{kl}(Z, W)$ are formal mappings, for all $k = 1, \dots, q$ and $l = 2, \dots, N$.
- $\varphi_{kl}(Z, W) = z_{k1} \tilde{\varphi}_{kl}(Z, W)$, where $\tilde{\varphi}_{kl}(Z, W)$ are formal mappings, for all $k = 1, \dots, q$ and $l = 1, \dots, N$.

In articular, we consider the matrices

$$B' = \begin{pmatrix} b_1 & 0 & \dots & 0 \\ 0 & b_2 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & b_{q_1} \end{pmatrix} \in \mathcal{M}_{q_1 \times p}(\mathbb{C}) \text{ and } B'' = \begin{pmatrix} b_1 & 0 & \dots & 0 & 0 & \dots & 0 \\ 0 & b_2 & \dots & 0 & 0 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & b_{q_1} & 0 & \dots & 0 \end{pmatrix} \in \mathcal{M}_{q_1 \times p'}(\mathbb{C}).$$

Following Zhu[31], we introduce the matrices

$$P_{B'}(Z) = \frac{\langle \tilde{Z}, B' \rangle}{B' \overline{B'}^t} B' \text{ and } Q_{B'}(Z) = \tilde{Z} - \frac{\langle \tilde{Z}, B' \rangle}{B' \overline{B'}^t} B',$$

$$P_{B''}(Z') = \frac{\langle \tilde{Z}', B'' \rangle}{B'' \overline{B''}^t} B'' \text{ and } Q_{B''}(Z') = \tilde{Z}' - \frac{\langle \tilde{Z}', B'' \rangle}{B'' \overline{B''}^t} B'',$$

where we have used the matrices

$$\tilde{Z} = \begin{pmatrix} z_{11} & z_{12} & \dots & z_{1N} \\ \vdots & \vdots & \ddots & \vdots \\ z_{q_1 1} & z_{q_1 2} & \dots & z_{q_1 N} \end{pmatrix} \text{ and } \tilde{Z}_0 = \begin{pmatrix} z_{q_1+1 1} & z_{q_1+1 2} & \dots & z_{q_1+1 N} \\ \vdots & \vdots & \ddots & \vdots \\ z_{q 1} & z_{q 2} & \dots & z_{q N} \end{pmatrix},$$

$$\tilde{Z}' = \begin{pmatrix} z'_{11} & z'_{12} & \dots & z'_{1N'} \\ \vdots & \vdots & \ddots & \vdots \\ z'_{q_1 1} & z'_{q_1 2} & \dots & z'_{q_1 N'} \end{pmatrix} \text{ and } \tilde{Z}'_0 = \begin{pmatrix} z'_{q_1+1 1} & z'_{q_1+1 2} & \dots & z'_{q_1+1 N'} \\ \vdots & \vdots & \ddots & \vdots \\ z'_{q' 1} & z'_{q' 2} & \dots & z'_{q' N'} \end{pmatrix}.$$

In particular, we reconsider from [31] the Classes of Involutions

$$\varphi_{B'} : D_{p,q} \rightarrow D_{p,q}, \text{ defined by } (\varphi_{B'}(Z))^t = \left(\frac{B' - P_{B'}(Z) - \sqrt{I_{q_1} - B' \overline{B'}^t} Q_{B'}(Z)}{I_{q_1} - \langle Z, B' \rangle} \right)_{\tilde{Z}_0},$$

$$\varphi_{B''} : D_{p',q'} \rightarrow D_{p',q'}, \text{ defined by } (\varphi_{B''}(Z'))^t = \left(\frac{B'' - P_{B''}(Z') - \sqrt{I_{q_1} - B'' \overline{B''}^t} Q_{B''}(Z')}{I_{q_1} - \langle Z', B'' \rangle} \right)_{\tilde{Z}'_0}.$$

When some numbers are vanishing, the lines from [31] may be implemented in order to work with convenient block of matrices. The proof Lemma 1.1 from [31] may be adapted, because we have considered only diagonal matrices. Let $\tilde{U} \in \mathcal{M}_{pq \times pq}(\mathbb{C})$ preserving $S_{p,q}$ such that $H(Z_1, Z_2) = \tilde{U} \otimes \varphi_B(Z_1, Z_2)$. Therefore

$$(V(Z))^t = \left(\begin{pmatrix} z_{11} & z_{21} & \dots & z_{p-1 1} \\ z_{12} & z_{22} & \dots & z_{p-1 2} \\ \vdots & \vdots & \ddots & \vdots \\ z_{1q} & z_{2q} & \dots & z_{p-1 q} \end{pmatrix} \begin{pmatrix} z_{p1} \\ z_{p2} \\ \vdots \\ z_{pq} \end{pmatrix} \odot (\varphi_B(Z))^t \right).$$

We finally obtain $\varphi'_{B'} \circ V \circ \varphi_B(Z) = \mathcal{W}$. This proof is finished. \square

10. Proof of Theorem 1.1

We obtain the Classes of Equivalence (1.1) and (1.3), because Lemma 9.1 applies in the light of (3.4), since (6.3) holds while the procedure of Hamada[9] was applied in Section 8. The Classes of Equivalence (1.1) and (1.3) are defined using compositions with automorphisms of $S_{p,q}$ and $S_{p',q'}$. They extend to holomorphic automorphisms of $D_{p,q}$ and $D_{p',q'}$ according to Kaup-Zaitsev[14],[15] and Kim-Zaitsev[16],[17].

11. Acknowledgements

Thanks to the Universities Koç, Middle East Technical, Galatasaray and Hacettepe for hospitality prior to my deportation from Istanbul. Special Thanks to my (former) supervisor Prof. Dmitri Zaitsev for important conversations while I was reading [16] (and for his guidance on the main part [3] (supported by Science Foundation Ireland, Grant 06/RFP/MAT 018) of my doctoral thesis).

References

- [1] Baouendi, M.S.; Huang, X. — Super-Rigidity for holomorphic mappings between hyperquadrics with positive signature. *J. Diff. Geom.* **69** (2005), no. 2, 379 – 398.
- [2] Baouendi, M.S.; Ebenfelt, P.; Huang, X. — Holomorphic mappings between hyperquadrics with small signature difference. *American J. Math.* **133** (2011), no. 6, 1633 – 1661.
- [3] Burcea, V. — A normal form for a real 2-codimensional submanifold in \mathbb{C}^{N+1} near a CR singularity. *Adv. in Math.* **243** (2013), 262 – 295.
- [4] Chern, S.S.; Moser, J. — Real hypersurfaces in complex manifolds. *Acta Math.* **133** (1974), 219 – 271.
- [5] Chen, Z.; Ji, S.; Xu, D. — Rational holomorphic maps from \mathbb{B}^2 into \mathbb{B}^N with degree 2. *Science in China: Ser. A Math.* **49** (2006), no. 11, 1504 – 1522.
- [6] Chirvasitu, A. — Shilov boundaries determine irreducible bounded symmetric domains. *Proc. Am. Math. Soc.* **149**, (2021), 3863 – 3872.
- [7] Ezov, V.V.; Schmalz, G. — Automorphisms and holomorphic mappings of standard CR-manifolds and Siegel domains. Complex analysis and representation theory, I. *J. Math. Sci. (New York)* **92** (1998), no. 2, 3712 – 3763.
- [8] Faran, J.J. — Maps from the two ball to the three ball. *Inv. Math.* **68** (1982), no. 3, 441 – 475.
- [9] Hamada, H. — Rational proper holomorphic maps from \mathbb{B}^n into \mathbb{B}^{2n} . *Math. Ann.* **331** (2005), no. 3, 693 – 711.
- [10] Huang, X. — On a linearity problem for proper holomorphic maps between balls in complex spaces of different dimensions. *J. Differential Geom.* **51** (1999), no. 1, 13-33.
- [11] Huang, X. — On a semi-rigidity property for holomorphic maps. *Asian. J. Math.* **7** (2003), no. 4, 463 – 492.
- [12] Huang, X.; Ji, S. — Mapping \mathbb{B}^n into \mathbb{B}^{2n-1} . *Invent. Math.* **145** (2001), no. 2, 219-250.

- [13] **Huang, X.; Ji, S.; Yin, W.**—The third gap for proper holomorphic maps between balls. *Math. Ann.* **358** (2014), no. 1-2, 115 – 142.
- [14] **Kaup, W.; Zaitsev, D.**— On symmetric Cauchy-Riemann manifolds. *Adv. Math.* **149** (2000), no. 2, 145 – 181.
- [15] **Kaup, W.; Zaitsev, D.**— On the CR-structure of compact group orbits associated with bounded symmetric domains. *Invent. Math.* **153** (2003), no. 1, 45 – 104.
- [16] **Kim, S.-Y.; Zaitsev, D.**— Rigidity of CR Maps between Shilov Boundaries of Bounded Symmetric Domains. *Inv. Math.* **193** (2013), no. 2, 409 – 437.
- [17] **Kim, S.-Y.; Zaitsev, D.**— Rigidity of proper holomorphic maps between bounded symmetric domains. *Math. Ann.* **362** (2015), no. 1-2, 639 – 677.
- [18] **Kim, S.-Y.** — Spheres in The Shilov Boundary of Bounded Symmetric Domains *Journal of the Korean Society of Mathematical Education: Series B: Pure and Applied Mathematics* **22** (2015), no. 1, 35 – 56.
- [19] **Lebl, J.; Peters, H.**— Polynomials constant on a hyperplane and CR maps of spheres. *Illinois J. Math.* **56** (2012), no. 1, 155 – 175.
- [20] **Seo, A.** — New examples of proper holomorphic maps among symmetric domains. *Michigan Math. J.* **64** (2015), no. 2, 435 – 448.
- [21] **Seo, A.** — Proper holomorphic polynomial maps between bounded symmetric domains of classical type. *Proc. Amer. Math. Soc.* **144** (2016), no. 2, 739 – 751.
- [22] **Mok, N.** — *Metric rigidity theorems on Hermitian locally symmetric spaces*. Series in Pure Mathematics, **6**, World Scientific Publishing Co., Inc., Teaneck, NJ, 1989. xiv+278
- [23] **Mok, N.** — Extension of germs of holomorphic isometries up to normalizing constants with respect to the Bergman metric *J. Eur. Math.Soc.* **14** (2012), no.5, 1617 – 1656.
- [24] **Ji, S.; Xu, D.**—Maps between \mathbb{B}^n and \mathbb{B}^N with geometrical rank $k_0 \leq n - 2$ and minimum N . *Asian J. Math.* **8** (2004), no. 2, 233 – 258.
- [25] **Tu, Z.**— Rigidity of proper holomorphic mappings between nonequidimensional bounded symmetric domains. *Math. Z.* **240** (2002), no. 1, 13 – 35.
- [26] **Webster, S.M.**— On mapping an n -ball into an $(n + 1)$ -ball in complex spaces. *Pacific J. Math.* **81** (1979), no. 1, 267 – 272.
- [27] **Yuan, Y.** —Local holomorphic isometries, old and new results. *Proceedings of the Seventh International Congress of Chinese Mathematicians*. Vol. II, 409 – 419, Adv. Lect. Math., no. 44, Int. Press, Somerville, MA, (2019).
- [28] **Yuan, Y.; Chan, S.-T.** —Holomorphic Isometries from the Poincare disk into bounded symmetric domains of rank at least two. *Annales de l'Institut Fourier* **69** (2019), no. 5, 2205 – 2240.
- [29] **Zaitsev, D.** — New Normal Forms for Levi-nondegenerate Hypersurfaces. *Several Complex Variables and Connections with PDE Theory and Geometry*. Complex analysis-Trends in Math., Birkhäuser/Springer Basel AG, Basel, (*Special Issue: In the honour of Linda Preiss Rothschild*), pp. 321 – 340, (2010).
- [30] **Zaitsev, D.** — Normal forms of nonintegrable almost CR structures, *Amer. J. Math.* **134** (2012), no.4, 915 – 947.
- [31] **Zhu, K.** — *Spaces of Holomorphic Functions in the Unit Ball* Graduate Texts in Mathematics, **226**, Springer Verlag, New York, USA, 2005.

V. BURCEA: INDEPENDENT
 Email address: vdburcea@gmail.com