Discrete probabilistic and algebraic dynamics: a stochastic commutative Gelfand-Naimark Theorem

Arthur J. Parzygnat

Abstract

We introduce a category of stochastic maps (certain Markov kernels) on compact Hausdorff spaces, construct a stochastic analogue of the Gelfand spectrum functor, and prove a stochastic version of the commutative Gelfand-Naimark Theorem. This relates concepts from algebra and operator theory to concepts from topology and probability theory. For completeness, we review stochastic matrices, their relationship to positive maps on commutative C^* -algebras, and the Gelfand-Naimark Theorem. No knowledge of probability theory nor C^* -algebras is assumed and several examples are drawn from physics.

Contents

1	$\mathbf{A}\mathbf{n}$	algebraic perspective on probability theory	2
	1.1	Brief background and motivation	2
	1.2	Overview of results	3
2	Pos	itive maps and stochastic matrices	5
	2.1	Introduction	5
	2.2	Some categories for finite probability theory	6
	2.3	C^* -algebras and states	11
	2.4	Two types of morphisms of C^* -algebras	14
	2.5	From probability theory to algebra	
	2.6	From algebra to probability theory	
	2.7	Some quantum mechanics	
3	An	equivalence between spaces and algebraic structures	23
	3.1	From spaces to algebras	23
	3.2	Some topological preliminaries	25
	3.3	The spectrum of a commutative C^* -algebra	28
	3.4	The Gelfand transform	31
	3.5	The commutative Gelfand-Naimark equivalence	33

²⁰¹⁰ Mathematics Subject Classification. 60B05 (Primary) 47B65, 18A40 (Secondary).

Key words and phrases. C^* -algebra, positive map, regular measure, Markov kernel, categorical probability, Radon monad, compact Hausdorff, Dynkin π - λ Theorem, states on C^* -algebras, spectrum, Choquet theory.

4	Abs	stract probability theory	35
	4.1	Compact Hausdorff spaces and complex measures	36
	4.2	Probability measures on compact Hausdorff spaces	42
	4.3	Continuous stochastic maps	44
	4.4	From continuous probability theory to algebra	51
	4.5	The stochastic Gelfand spectrum functor	53
	4.6	The stochastic commutative Gelfand-Naimark Theorem	62
5	Clo	sing discussion	64
	5.1	Categories of classical and quantum dynamics	65
	5.2	Categorical probability theory	66
	5.3	The Baire approach	68
In	dev	of notation	70

1 An algebraic perspective on probability theory

That's another thing about categories, not that they give you the proofs, but they tell you what to prove!—Mikhail Gromov¹

1.1 Brief background and motivation

The fact that every compact Hausdorff space can be recovered from its C^* -algebra of complexvalued continuous functions was discovered by Gelfand and Naimark in 1943 (see Lemma 1 in [18]). This involved constructing a space out of a commutative C^* -algebra and a continuous map out of a *-homomorphism of algebras. Extending earlier work of Riesz and Markov (and many others), Kakutani proved that finite regular measures on such spaces correspond to positive linear functionals on the algebras (see Theorem 9 in [23]). One can combine these ideas in the following way. Continuous maps between spaces can be viewed as deterministic assignments in the sense that their values on inputs are specified precisely. Instead, we can imagine a nondeterministic analogue as a "smeared out" function, or more precisely, an assignment sending inputs to probability measures. Such "stochastic maps" correspond to (completely) positive linear maps between the algebras. The latter are well-defined for non-commutative C^* -algebras and describe non-deterministic dynamics in quantum systems [27]. Although this correspondence is well-understood to a large extent [9], [12] some important features are missed without considering the category of such spaces and stochastic maps. For example, how are the compositions of stochastic maps defined, are compositions respected under the correspondence, what is the explicit procedure to go back from a positive map to a stochastic map, and is there a way to formulate all of these dualities in the same context?

Our aim is to introduce these ideas in a suitable language and answer these questions. In the spirit of the Rosetta Stone [4], we will use category theory to do this, though we will only

¹This quote is taken from Mikhail Gromov's Lecture 1 on "Probability, symmetry, linearity" given at the Institut des Hautes Études Scientifiques on October 10, 2014.

use the bare minimum necessary. In particular, we only assume the reader knows what categories, functors, and natural transformations are and how to compose them though the reader may also understand the results without even knowing these definitions verbatim from memory. Using category theory has the immediate benefit of formulating our questions precisely thereby making the goals clear: we must identify the appropriate topological, algebraic, deterministic, and non-deterministic categories, construct mappings between them, and prove the above mentioned equivalences.

This work fits into the general context of categorical probability theory, which was pioneered by Lawvere who first defined a category of stochastic maps in 1962 [26]. Further work in the area of category-theoretic aspects of measure theory was developed by Linton in 1963 [28] and the Polish school in the early 1970's, notably Semadeni [42] and Świrszcz [46]. In the early 1980's, Giry further developed these ideas by describing the monadic structures available in probability theory on a larger class of spaces [19]. More recently, this area was revisited in the area of computer science where functional programming makes use of category-theoretic ideas, particularly monads. The relationship to the category of ordinary relations, used in the theory of computing, was generalized to the stochastic setting by Panangaden in 1998 [31] (an earlier version of this article was titled "Probabilistic Relations") based on earlier work of Kozen in 1981 [24]. A monadic and algebraic viewpoint on the questions raised above were worked out by Furber and Jacobs recently [15]. Our goals are similar, but we approach the problem from an analytic perspective.

1.2 Overview of results

We prove a generalization of the commutative Gelfand-Naimark Theorem (henceforth referred to as just the Gelfand-Naimark Theorem) that is valid for continuous stochastic maps on compact Hausdorff spaces. This is done at three different levels of increasing complication, the first two of which are well known. In Section 2, we review the equivalence between stochastic matrices on finite sets and positive maps on finite-dimensional commutative (unital) C^* -algebras, which are all basically of the form \mathbb{C}^n for some $n \in \mathbb{N}$ with pointwise algebraic structure. Specializing to the case of functions instead of stochastic matrices corresponds to *-homomorphisms on the C^* -algebra side. In Section 3, we review the Gelfand-Naimark Theorem phrased in an appropriate categorical context as an equivalence between the category of compact Hausdorff spaces with continuous maps and the category of commutative unital C^* -algebras and unit-preserving *-homomorphisms. We do not prove the fundamental theorems here since they are well documented in standard references, but we do introduce the necessary concepts needed to understand the statements in what follows.


Finally, in Section 4, we extend the Gelfand-Naimark Theorem to an equivalence between compact Hausdorff spaces with continuous stochastic maps and commutative C^* -algebras with positive unital maps. To even state this theorem, an appropriate category of such spaces and stochastic maps must be defined. Although versions of this category have been defined in the literature (see [19] and [31] for example), the spaces are typically either measurable spaces or Polish spaces. There is an important distinction between these two cases. Briefly, a $\underline{Markov\ kernel}$ from a measurable space (X, \mathcal{M}_X) to another one (Y, \mathcal{M}_Y) is a function $X \times \mathcal{M}_Y \to [0, 1]$ such that fixing the left variable gives a probability measure and fixing the right variable gives a measurable function. A Markov kernel may equivalently be defined as a stochastic map, i.e. a measurable

function $X \to \operatorname{ProbMeas}(Y)$, where $\operatorname{ProbMeas}(Y)$ is the set of all probability measures on Y equipped with the smallest σ -algebra for which the evaluation function ev_E : $\operatorname{ProbMeas}(Y) \to [0,1]$, defined by $\operatorname{ProbMeas}(Y) \ni \nu \mapsto \operatorname{ev}_E(\nu) := \nu(E)$, is measurable for all $E \in \mathcal{M}_Y$. In other words, in the case of measurable spaces, stochastic maps are in one-to-one correspondence with Markov kernels. However, in the case of Polish spaces, many applications demand a more restrictive class of Markov kernels that have additional continuity properties, but demanding that the Markov kernel is continuous is too restrictive in several contexts (see Remarks 4.72 and 4.142). Fortunately, one can use the second perspective and define stochastic maps as continuous functions $X \to \operatorname{ProbMeas}(Y)$, where $\operatorname{ProbMeas}(Y)$ is now equipped with the vague topology. As a result, it is a-priori unclear whether the associated kernel function is Borel measurable after fixing the right variable. Nevertheless, it follows from the fact that the spaces are Polish [19].

If Polish spaces are replaced by compact Hausdorff spaces, one can still define stochastic maps, but the above proof of measurability of the associated kernels fails. To circumvent this difficulty, we restrict our probability measures to be regular (this additional assumption is automatically satisfied for Polish spaces). We first prove that the stochastic maps on such spaces have measurable Markov kernels in Lemmas 4.64 and 4.76. Lemma 4.64 extends results due to Billingsley (see proof of Theorem 2.1 in [5]). This leads to an explicit definition of composition of such stochastic maps in Proposition 4.79 without referring to the Riesz-Markov-Kakutani Representation theorem as is done in [15] and [12]. We then prove that stochastic maps on compact Hausdorff spaces form a category, denoted by **cHausStoch**, in Theorem 4.102.

In addition, we provide an explicit and geometric construction of the Gelfand spectrum functor from commutative C^* -algebras and positive maps to the category of compact Hausdorff spaces and continuous stochastic maps without using the using algebras of functions and without appealing to the Riesz-Markov-Kakutani Representation Theorem. Instead, this "stochastic" Gelfand spectrum functor is constructed using techniques from convex analysis and Choquet theory in Theorems 4.134 and 4.151. Finally, we prove that the stochastic Gelfand spectrum functor exhibits a categorical inverse to the continuous functions functor in Theorem 4.162.

Our results can be succinctly summarized in that we complete the diagrammatic cube (the ^{op} superscript refers to flipping the directionality of all morphisms)



by adding cHausStoch and the functors into and from it. The arrows between the left and right

faces describe the equivalences between topology and analysis on the left and algebra on the right. The arrows from top to bottom describe the generalization of going from deterministic dynamics to stochastic dynamics. The arrows from front to back describe the inclusion of finite systems to possibly infinite systems. The arrows from the left to the right are particularly important in physics for the following reason. Much of classical and quantum dynamics, whether deterministic or non-deterministic, can be formulated in a single category: C^* -algebras and completely positive maps. Although, classical evolution is typically thought of as being quite distinct from quantum evolution, they both take place in the *same* category allowing a precise comparison between dynamics. In particular, it allows mixing classical and quantum systems. For example, measurement of quantum systems by macroscopic (classical) beings can be thought of as a map of C^* -algebras and pulling back states from a non-commutative C^* -algebra to a commutative one (see Example 2.99). The importance of commutative subalgebras in the theory of measurement and their relationship to quantum mechanics has been emphasized and summarized nicely in Heunen's work [21].

In summary, we first analyze the front and right face in Section 2, then the top face in Section 3, and finally the category **cHausStoch** and how it fits into this cube via a stochastic Gelfand-Naimark Theorem in Section 4. Concluding remarks, including a more detailed discussion of the relationship between our work and others, are in Section 5. For the reader who has all the necessary background, our main results are contained in Sections 4.3 through 5.1. The reader may find the Index of notation at the end helpful.

Probability theory evolved before the formulation of measure theory, and, in particular, before the birth of category theory. As such, it has developed its own language and culture, which the author is largely unfamiliar with resulting in a potentially contentious presentation. Caveat lector.

Acknowledgements.

This work is based on a series of lectures given at the Analysis Learning Seminar at the University of Connecticut, Storrs and the Mathematical Physics, Fourier Analysis and Applications Seminar at the CUNY Graduate Center in New York during the Spring 2017 semester. The author is grateful for the invitations by Matthew Badger and Vyron Vellis to the former and by Azita Mayeli and Max Yarmolinsky to the latter. The author has greatly benefited from discussions with Stefan Andronache, Iddo Ben-Ari, Behrang Forghani, Marcelo Nomura, Benjamin Russo, Ambar Sengupta, Scott O. Wilson, and Yun Yang. The author also thanks Markus Haase for several insightful comments and suggestions on an earlier version of this work. Parts of Sections 2 and 3 were worked out when the author was partially supported by the CUNY Graduate Center Capelloni Dissertation Fellowship and NSF grant PHY-1213380.

2 Positive maps and stochastic matrices

2.1 Introduction

We start off with basic probability theory on finite sets and linear algebra. We first learned about this perspective from the work of Baez and Fritz on entropy [2]. We find their notation, particularly distinguishing between deterministic and non-deterministic processes, incredibly lucid. The former are depicted with straight arrows \longrightarrow while the latter with curvy arrows \longrightarrow . We introduce the

concepts of C^* -algebras and states on C^* -algebras. There are two main categories described here both of whose objects consist of C^* -algebras. The difference occurs at the level of morphisms. In one category, the morphisms consist of algebra *-homomorphisms and in the other case they consist of completely positive maps. We will show that there is a correspondence between these categories when restricted to finite-dimensional commutative C^* -algebras and ordinary concepts in probability theory, namely stochastic matrices. Thus, the category of operations in quantum mechanics can be viewed as a non-commutative extension of the category of stochastic (non-deterministic) processes in classical mechanics.

In this section, we introduce

- i. FinSetFun, the category of finite sets and functions (deterministic processes),
- ii. FinSetStoch, the category of finite sets and stochastic matrices (non-deterministic processes),
- iii. $\mathbf{fdcC^*-Alg}$, the category of finite-dimensional commutative \mathbb{C}^* -algebras and *-homomorphisms (the non-commutative analogue of deterministic processes), and
- iv. $\mathbf{fdcC^*}$ -AlgPos, the category of finite-dimensional commutative \mathbb{C}^* -algebras and completely positive maps (the non-commutative analogue of stochastic maps)

and show that there are functors (the op superscript will be explained in Theorem 2.72)

$$\begin{array}{cccc} \mathbf{FinSetFun^{op}} & \longrightarrow \mathbf{fdcC^*-Alg} \\ & & & \downarrow \\ &$$

so that the horizontal functors are fully faithful and so that the diagram commutes.

These concepts will be rigorously defined and sufficient intuition will be provided to illustrate that they are nothing more than simple ideas in probability theory. However, we do assume the reader is (at least vaguely) familiar with the terms category, functor, and natural transformation.

2.2 Some categories for finite probability theory

We begin with probability theory as it might be more intuitive than algebra.

Definition 2.2. A <u>finite probability space</u> is a pair (X, p) consisting of a finite set X and a function $p: X \to \mathbb{R}$ satisfying

$$p(x) \ge 0 \qquad \forall \ x \in X \tag{2.3}$$

and

$$\sum_{x \in X} p(x) = 1. {(2.4)}$$

The elements of X are called <u>events</u> and p is called a <u>probability measure</u> on X. The notation $p_x := p(x)$ will often be used.

Note that p associates a number to any subset $E \subseteq X$ given by

$$p(E) := \sum_{x \in E} p_x \tag{2.5}$$

and is the reason we refer to p as a measure (more on this will be discussed in Section 4). Technically, these subsets are called events.

Definition 2.6. Let (X, p) and (Y, q) be two finite probability spaces. A <u>probability-preserving</u> function from (X, p) to (Y, q) is a function $f: X \to Y$ satisfying

$$q_y = \sum_{x \in f^{-1}(y)} p_x \tag{2.7}$$

for all $y \in Y$.

Composition is given by the usual composition of functions and the composition of two probability-preserving functions is immediately seen to be probability-preserving. The meaning of a probability-preserving function between probability spaces can be seen nicely in the following example.


Example 2.8. Consider the set

$$X := \left\{ \boxed{\bullet}, \boxed{\bullet}, \boxed{\bullet}, \boxed{\bullet}, \boxed{\bullet}, \boxed{\bullet}, \boxed{\bullet} \right\}, \boxed{\bullet} \right\}$$
 (2.9)

with the probability measure given by $\frac{1}{6}$ for each element. Consider the set

$$Y := \{O, E\} \tag{2.10}$$

consisting of just two elements (O stands for "odd" and E stands for "even") with probability measure given by $\frac{1}{2}$ for each element. Also consider the function $f: X \to Y$ defined by



sending \bullet , \bullet , and \bullet to O and sending \bullet , and \bullet to E. Therefore, the probability-preserving function f is associated to the process of rolling a die and considering the likelihood of rolling an odd or an even roll as opposed to rolling a 1, 2, 3, 4, 5, or 6. This describes a deterministic process since we know with certainty that if we see odd, the die must be 1, 3, or 5, and analogously for even.

Notation 2.12. Let **FinProb** be the category whose objects are finite probability spaces (X, p) and whose set of morphisms $\{(X, p) \to (Y, q)\}$ are probability-preserving functions from (X, p) to (Y, q). Similarly, let **FinSetFun** be the category whose objects are finite sets and whose morphisms are functions. Composition in both categories is defined to be the ordinary composition of functions.

The category **FinProb** was recently used by Baez, Fritz, and Leinster to provide a new categorical characterization of entropy in the case of finite sets [3]. Extensions of this result were obtained more recently for relative entropy on finite sets by Baez and Fritz [2] and relative entropy on Polish spaces by Gagné and Panangaden [17]. In all cases, categories of stochastic maps were crucial in these characterizations. While functions take in one input and spit out one output, stochastic maps take in one input and have a "spread" of outputs described by some probability distribution.

Definition 2.13. Let X and Y be two finite sets. Let Pr(Y) denote the set of probability measures on Y. A *stochastic map* from X to Y is a function

$$\begin{array}{c}
X \xrightarrow{f} \Pr(Y) \\
x \mapsto f(x)
\end{array} \tag{2.14}$$

whose evaluation on elements in Y is written as

$$Y \ni y \xrightarrow{f(x)} f_{yx} \in \mathbb{R}_{\geq 0}. \tag{2.15}$$

Note that by definition of f(x) being a probability measure on Y, this means that

$$\sum_{y \in Y} f_{yx} = 1 \tag{2.16}$$

for all $x \in X$. The numbers $\{f_{yx}\}$ labelled by $x \in X$ and $y \in Y$ form what is often called a stochastic matrix. We denote stochastic maps using squiggly arrows $X \leadsto Y$ following the convention of Baez and Fritz [2] to distinguish stochastic maps from ordinary functions. By abuse of notation, we often use the same notation f for both $f: X \to Pr(Y)$ and $f: X \leadsto Y$ and we may often refer to $X \to Pr(Y)$ as the stochastic map (the use of straight and curly arrows avoids potential confusion).

Example 2.17. Let X and Y be finite sets and let $f: X \to Y$ be a function. Associated to f is a canonical stochastic map $f: X \leadsto Y$ defined by

$$X \times Y \ni (x, y) \mapsto f_{yx} := \delta_{yf(x)}. \tag{2.18}$$

Here the function $\delta: Y \times Y \to \mathbb{R}$ is the Kronecker-delta function defined by

$$\delta_{yy'} := \begin{cases} 1 & \text{if } y = y' \\ 0 & \text{otherwise} \end{cases}$$
 (2.19)

In other words, f associates to x the probability measure that is 1 on the element f(x) and 0 elsewhere, i.e. if we have a function, we know with certainty where a point will go.

Example 2.20. Recall Example 2.8 with rolling a die. If instead of interpreting the set Y as the set of odd or even die, suppose that the O and E just stand for boxes labelled by O or E. A probability-preserving stochastic map $X \leadsto Y$ can be interpreted as saying that for any given roll of a die, that particular side has a probability of being placed in box O and a complementary probability of being placed in box E.


Example 2.21. A random walk on a circle is described by the following stochastic map. Let $N \in \mathbb{N}$ be the number of points on a circle, labeled $x_0, x_1, \ldots, x_{N-1}$ with $x_N \equiv x_0$ and so on so that $X := \{x_0, x_1, \ldots, x_{N-1}\}$. The random walk on X is described by the stochastic map $f: X \leadsto X$ defined by

$$f_{ji} := \begin{cases} \frac{1}{2} & \text{if } j \equiv i \pm 1 \mod N \\ 0 & \text{otherwise} \end{cases}$$
 (2.22)

where the shorthand notation f_{ji} is used instead of $f_{x_jx_i}$. The reason this is called the random walk is because starting at any point, you have a $\frac{1}{2}$ probability of moving one step in either direction. A random walk on a line or plane can also be described though the set of points would be infinite, which we are not considering at the moment.

Example 2.23. With such a definition in place, notice that the set of all stochastic maps $\{\bullet\}$ \leadsto X from a single element set $\{\bullet\}$ to a finite set X is isomorphic (bijective) to the set $\Pr(X)$ of all probability measures on X.

Given finite sets X, Y, and Z together with stochastic maps $f: X \leadsto Y$ and $g: Y \leadsto Z$, one would like to iterate these maps. To do so, the probability of $x \in X$ evolving to $z \in Z$ is given by summing over all possible intermediate elements $y \in Y$ weighted by their corresponding probabilities.



This motivates the following definition.

Definition 2.24. Let X, Y, and Z be finite sets. The <u>composition</u> of the stochastic map $f: X \leadsto Y$ followed by the stochastic map $g: Y \leadsto Z$, written as $g \circ f: X \leadsto Z$, is the function $g \circ f: X \to \Pr(Z)$ defined by sending $x \in X$ to the probability measure on Z defined by

$$Z \ni z \xrightarrow{(g \circ f)(x)} (g \circ f)_{zx} := \sum_{y \in Y} g_{zy} f_{yx}. \tag{2.25}$$

Note that $g \circ f$ is indeed a stochastic map because

$$\sum_{z \in Z} (g \circ f)_{zx} \stackrel{\text{def}}{=} \sum_{z \in Z} \sum_{y \in Y} g_{zy} f_{yx} = \sum_{y \in Y} \sum_{z \in Z} g_{zy} f_{yx} = \sum_{y \in Y} f_{yx} = 1.$$
 (2.26)

Notation 2.27. Let FinSetStoch be the category whose objects are finite sets and whose morphisms are stochastic maps. We leave checking the axioms of a category to the reader.

Notice that **FinSetFun** can be viewed as a subcategory of **FinSetStoch** by Example 2.17 and we therefore denote the associated functor by δ : **FinSetFun** \rightarrow **FinSetStoch**.

Remark 2.28. The definition of a stochastic map is closely related to the notion of a stochastic process. A stochastic map $f: X \leadsto X$ from a finite set X to itself can be used to describe the possible evolution of a state. In the special case of a known initial condition, i.e. an element $x_0 \in X$, one can consider the sequence of probability distributions (the first element of which is just the Kronecker delta distribution at x_0) given by $(x_0, f(x_0), f^2(x_0), f^3(x_0), f^4(x_0), \dots)$. Such a sequence is called a Markov chain. If each iteration of f is interpreted as a time-step, then the probability distribution $f^n(x_0)$ is to be interpreted in the following way. The probability of finding the initial condition x_0 to evolve to the position $x_n \in X$ after n time steps is $(f^n)_{x_n x_0}$. More generally, the set of events can change under a given process (as we have described above). An evolution of some initial states can therefore be described by a sequence of stochastic maps

$$X_0 \leadsto X_1 \leadsto X_2 \leadsto X_3 \leadsto \cdots,$$
 (2.29)

where each of the X_i is a set of possible events. A point in X_0 , described by a function $\{\bullet\} \to X_0$, evolves just as a probability measure $\{\bullet\} \leadsto X_0$, namely by precomposing:

$$\{\bullet\} \leadsto X_0 \leadsto X_1 \leadsto X_2 \leadsto X_3 \leadsto \cdots$$
 (2.30)

Example 2.31. Let us model one-dimensional diffusion via a stochastic map on a finite set, thought of as an approximation to diffusion on the interval [-1,1] with periodic boundary conditions. Namely, let X be the finite set given by

$$X := \left\{ \frac{k}{10} : k \in \{-10, -9, \dots, 9\} \right\}, \tag{2.32}$$

where the point 1 is to be interpreted as being identified with -1. According to the heat diffusion equation, after some time step, a particle at a point has the following probability distribution. Approximately, it has a 0.56 likelihood to stay in the same place, 0.21 likelihood to move one unit over in either direction, and a 0.01 likelihood to move two units over in either direction. The stochastic matrix associated to this process is given by

$$\begin{bmatrix} 0.56 & 0.21 & 0.01 & 0 & \cdots & 0.01 & 0.21 \\ 0.21 & 0.56 & 0.21 & 0.01 & \cdots & 0 & 0.01 \\ 0.01 & 0.21 & 0.56 & 0.21 & \cdots & 0 & 0 \\ 0 & 0.01 & 0.21 & 0.56 & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0.01 & 0 & 0 & 0 & \cdots & 0.56 & 0.21 \\ 0.21 & 0.01 & 0 & 0 & \cdots & 0.21 & 0.56 \end{bmatrix}$$

$$(2.33)$$

Under a few iterations of this process (matrix multiplication), the distribution (described by a unit vector) becomes the uniform distribution (as one would expect—systems tend to equilibriate). This is depicted in Figure 1.




Figure 1: A sharp Gaussian representing a definite initial condition at the origin tends towards a uniform distribution after several iterations of the heat flow described by the matrix (2.33). t = n with $n \in \{0\} \cup \mathbb{N}$ stands for the n-th time step.

Note that the formalism described above a-priori only describes discrete stochastic processes. One can adapt these ideas to study continuous processes though we do not discuss this here except for an example from quantum mechanics in Section 2.7. The reader is referred to Giry's work, which is not only a foundational paper on categorical probability theory but also discusses this in some detail [19].

2.3 C^* -algebras and states

There are several definitions that must be provided so that this work can be somewhat self-contained. A more thorough account of many definitions and results can be found in Dixmier's book [11] with additional references provided throughout.

Definition 2.34. An (associative and unital) <u>algebra</u> is a vector space \mathcal{A} (over \mathbb{C}) together with

- (a) a binary multiplication operation $\mathcal{A} \times \mathcal{A} \to \mathcal{A}$,
- (b) an element $1_{\mathcal{A}} \in \mathcal{A} \setminus \{0\}$.

These data must satisfy the following conditions:

- i. (ab)c = a(bc) (associativity of multiplication),
- ii. a(b+c) = ab + ac and (a+b)c = ac + bc (distributivity of multiplication over vector addition),
- iii. $(\lambda a)b = a(\lambda b) = \lambda(ab)$, (distributivity of multiplication over scalar multiplication), and
- iv. $a1_{\mathcal{A}} = 1_{\mathcal{A}}a = a$ ($1_{\mathcal{A}}$ is a unit for the multiplication),

for all $a, b, c \in \mathcal{A}$ and $\lambda \in \mathbb{C}$.

Definition 2.35. An algebra \mathcal{A} together with a norm $\|\cdot\|:\mathcal{A}\to\mathbb{R}_{\geq 0}$ for which \mathcal{A} with this norm is a normed vector space and

$$||ab|| \le ||a|| ||b|| \qquad \forall \ a, b \in \mathcal{A} \tag{2.36}$$

is a <u>normed algebra</u>. A is a <u>Banach algebra</u> iff all Cauchy sequences converge with respect to this norm.

Definition 2.37. An algebra \mathcal{A} together with an anti-automorphism $^*: \mathcal{A} \to \mathcal{A}$, i.e.

$$(\lambda a + b)^* = \lambda^* a^* + b^* \qquad \forall \ \lambda \in \mathbb{C}, \ a, b \in \mathcal{A}$$
 (2.38)

and

$$(ab)^* = b^*a^* \qquad \forall \ a, b \in \mathcal{A}, \tag{2.39}$$

is an <u>involutive algebra</u>. * is known as an <u>involution</u> on A.

Definition 2.40. Let \mathcal{A} be both an involutive algebra and a normed algebra. If, in addition, \mathcal{A} satisfies

$$||a^*|| = ||a|| \qquad \forall \ a \in \mathcal{A} \tag{2.41}$$

then \mathcal{A} is called an <u>normed involutive algebra</u>. If, in addition, all Cauchy sequences in \mathcal{A} converge, then \mathcal{A} is called a <u>involutive Banach algebra</u>.

Definition 2.42. A C^* -algebra is an involutive Banach algebra \mathcal{A} for which

$$||a^*a|| = ||a||^2 \quad \forall \ a \in \mathcal{A}.$$
 (2.43)

A C^* -algebra is <u>commutative</u> iff ab = ba for all $a, b \in \mathcal{A}$.

Remark 2.44. If \mathcal{A} is a normed algebra with an involution for which $||a||^2 \leq ||a^*a||$ for all $a \in \mathcal{A}$, then \mathcal{A} is an involutive algebra. In fact, if \mathcal{A} is a Banach algebra with involution satisfying this condition, then \mathcal{A} is a C^* -algebra [11].

All of the data of these algebraic structures are summarized in the following table (with no reference to their conditions).

data/	vector	algebra	normed	Banach	involutive	normed IA/
structure	space (VS)	argebra	VS	algebra	algebra (IA)	C^* -algebra
0	✓	√	✓	✓	✓	✓
+	✓	√	✓	✓	✓	✓
scalar mult.	✓	✓	✓	✓	✓	✓
×		√		✓	✓	✓
•			✓	✓		✓
*					✓	✓
1		✓		✓	✓	✓

Definition 2.45. Let \mathcal{A} and \mathcal{B} be two C^* -algebras. A *-homorphism of C^* -algebras from \mathcal{B} to \mathcal{A} is a bounded (i.e. continuous) linear map $f: \mathcal{B} \to \mathcal{A}$ such that

i.
$$f(b^*) = f(b)^*$$
,

ii.
$$f(b_1b_2) = f(b_1)f(b_2)$$
, and

iii.
$$f(1_{\mathcal{B}}) = 1_{\mathcal{A}}$$

for all $b, b_1, b_2 \in \mathcal{B}$.

Remark 2.46. The word bounded in the definition of a *-homomorphism of C^* -algebras is redundant. It is a theorem that such a map satisfying conditions i., ii., and iii. is automatically bounded [11]. Thus, the reader may safely ignore that condition.

There are a few basic examples of C^* -algebras that we will focus on in this section.

Example 2.47. \mathbb{C} is a C^* -algebra with its usual structure associated with complex numbers. The involution * is the complex conjugate. This is an example of a commutative C^* -algebra.

This previous example automatically induces a huge class of examples.

Example 2.48. Let X be a finite set and let \mathbb{C}^X denote the set of all functions $X \to \mathbb{C}$. There is a unique algebra structure and involution on \mathbb{C}^X so that the evaluation functions $\operatorname{ev}_x : \mathbb{C}^X \to \mathbb{C}$, defined by sending $\varphi : X \to \mathbb{C}$ to $\varphi(x)$, are *-homomorphisms for all $x \in X$ (these are just the pointwise algebraic structures).² The norm on \mathbb{C}^X is given by

$$\mathbb{C}^X \ni f \mapsto ||f|| := \sup_{x \in X} \{|f(x)|\}. \tag{2.49}$$

The previous examples were commutative C^* -algebras. The following example is the quintessential C^* -algebra of quantum mechanics of a finite system (such as a qubit).

Example 2.50. Fix $n \in \mathbb{N}$. The set $M_n(\mathbb{C})$ of $n \times n$ complex-valued matrices is a C^* -algebra with product given by matrix multiplication and * is the transpose complex conjugate. The norm can be taken to be either the operator norm or the standard Euclidean norm (the topology will be the same so it does not matter).

It is a theorem that every finite-dimensional C^* -algebra is (*-isomorphic to) a direct sum of these (Theorem III.1.1 in [10]).

Example 2.51. More generally, for any Hilbert space \mathcal{H} , the set of bounded operators $\mathcal{B}(\mathcal{H})$ on \mathcal{H} is a C^* -algebra via operator composition, * given by the adjoint, and the norm given by the operator norm.

Notation 2.52. Let C^* -Alg, cC^* -Alg, and $fdcC^*$ -Alg denote the categories whose objects are C^* -algebras, commutative C^* -algebras, and finite-dimensional commutative C^* -algebras, respectively, and whose the morphisms are all *-homomorphisms of C^* -algebras.

²This way of phrasing the natural algebraic structure on \mathbb{C}^X makes precise in what sense it is the "obvious" algebraic structure. Note that X does not have to be finite for this statement. However, one must be careful about the topology when X is infinite—demanding the smallest topology for which ev_x are all continuous would result in the topology of pointwise convergence, which is not the one we will use later.

Definition 2.53. Given a C^* -algebra \mathcal{A} , a <u>state</u> on \mathcal{A} is a bounded (i.e. continuous)³ linear function $\omega : \mathcal{A} \to \mathbb{C}$ such that $\omega(1_{\mathcal{A}}) = 1$ and $\overline{\omega(a^*a)} \ge 0$ for all $a \in \mathcal{A}$.

The following theorem relates states to density matrices and motivates why density matrices are the states of quantum mechanics as opposed to vectors (really, rays) in a Hilbert space [20].

Theorem 2.54. There is a one-to-one correspondence between states on $M_n(\mathbb{C})$ and matrices $\rho \in M_n(\mathbb{C})$ satisfying $\operatorname{tr}(\rho) = 1$ and $\rho^* = \rho$ (such matrices are known as density matrices). More precisely, for any density matrix ρ , the function $\operatorname{tr}(\rho \cdot) : M_n(\mathbb{C}) \to \mathbb{C}$, sending $A \in M_n(\mathbb{C})$ to $\operatorname{tr}(\rho A)$, is a state. Conversely, for any state $\omega : M_n(\mathbb{C}) \to \mathbb{C}$, there exists a unique density matrix ρ such that $\omega = \operatorname{tr}(\rho \cdot)$.

This theorem extends to states on $\mathcal{B}(\mathcal{H})$ with \mathcal{H} a separable Hilbert space and states that satisfy an additional continuity assumption.⁴ In the context of quantum mechanics, $\mathcal{B}(\mathcal{H})$ is the set of physical observables including non self-adjoint operators, such as ladder operators, so that it remains closed under multiplication by complex numbers. A state on $\mathcal{B}(\mathcal{H})$ assigns to every observable a number, interpreted as the expectation value of that observable for that state. States on C^* -algebras and their relationship to the more standard approach of quantum mechanics in terms of Hilbert spaces is described in a categorical framework in [32].

2.4 Two types of morphisms of C^* -algebras

There is another notion of a morphism that is important for C^* -algebras and utilizes less of its structure.

Definition 2.55. Let \mathcal{A} be a C^* -algebra. An element $a \in \mathcal{A}$ is <u>positive</u> iff there exists an element $b \in \mathcal{A}$ such that $a = b^*b$. Positivity of elements induces a partial ordering on \mathcal{A} and the notation $a \geq 0$ will be used to indicate that a is positive ($b \geq a$ iff $b - a \geq 0$).

Definition 2.56. Let \mathcal{A} and \mathcal{B} be two C^* -algebras. A <u>positive (unital) map</u> from \mathcal{B} to \mathcal{A} is a bounded linear function $\varphi : \mathcal{B} \to \mathcal{A}$ such that $\varphi(1_{\mathcal{B}}) = 1_{\mathcal{A}}$ and $\varphi(b^*b) \geq 0$ for all $b \in \mathcal{B}$.

A state is an example of a positive map. Every *-homomorphism of C^* -algebras is a positive map. The converse, however, is far from true. Examples are given in Section 2.7. Non-unital positive maps seem to be the weakest structure needed to make sense of operations and processes on quantum mechanical systems though one may also relax the unit-preserving condition [25].

Notation 2.57. Let \mathcal{A} be an involutive algebra and let $n \in \mathbb{N}$. $M_n(\mathcal{A})$ is the set of of $n \times n$ matrices with coefficients in \mathcal{A} with product given by matrix multiplication and * given by the transpose and applying * in each entry.

 $^{^3}$ The word bounded/continuous here is redundant. It is a theorem that such a map on a C^* -algebra satisfying these conditions is automatically bounded.

⁴Such states go by the name of <u>normal states</u> for those readers who would like to look up more information. However, normal states require a certain notion of continuity not available on arbitrary C^* -algebras, and one requires the notion of a von Neumann algebra, a special type of C^* -algebra that is closed with respect to a certain topology [36]. $\mathcal{B}(\mathcal{H})$ is an example of a von Neumann algebra.

Theorem 2.58. Let \mathcal{A} be a C^* -algebra. Then there exists a unique norm $\|\cdot\|$ on $M_n(\mathcal{A})$ such that

$$||AB|| \le ||A|| ||B|| \quad \forall A, B \in M_n(A)$$
 (2.59)

and

$$||A^*A|| = ||A||^2 \quad \forall A \in M_n(A).$$
 (2.60)

Furthermore, $M_n(A)$ with this structure is a C^* -algebra.

Proof. See Chapter 1 of Paulsen [33] or Chapters 4 and 6 of Warner [49].

Remark 2.61. There are *-isomorphisms

$$M_n(\mathcal{A}) \cong M_n(\mathbb{C}) \otimes \mathcal{A} \cong \mathcal{A} \otimes M_n(\mathbb{C}).$$
 (2.62)

Therefore, all of these involutive algebras have a unique norm giving them the structure of C^* -algebras. The theory of tensor products of C^* -algebras is notoriously complicated. Since we will work with commutative C^* -algebras for the most part, we will avoid such complications.

Definition 2.63. Let \mathcal{A} and \mathcal{B} be two C^* -algebras. A <u>completely positive (unital) map</u> from \mathcal{B} to \mathcal{A} is a function $\varphi: \mathcal{B} \to \mathcal{A}$ such that $\varphi \otimes \mathrm{id}_n : \mathcal{B} \otimes M_n(\mathbb{C}) \to \mathcal{A} \otimes M_n(\mathbb{C})$ is a positive map for all $n \in \mathbb{N}$. The abbreviation "CP map" will often be used for such maps.

Remark 2.64. One motivation for this definition comes from quantum mechanics [27], [25]. The concept of tensoring with the algebra $M_n(\mathbb{C})$ and the positive map with the identity means to adjoin another finite system that does not interact with the original system on \mathcal{B} during the operation. The resulting map $\varphi \otimes \mathrm{id}_n : \mathcal{B} \otimes M_n(\mathbb{C}) \to \mathcal{A} \otimes M_n(\mathbb{C})$ should therefore still be positive. There are examples of positive maps that are not completely positive so requiring this condition to be satisfied is an additional constraint [33], [25]. Unital maps of C^* -algebras correspond to the trace-preserving condition that is often referred to in texts on quantum information [30]. See Example 2.93 for an illustration. All maps in this work are unital so we will not use the more standard notation "CPU" [30].

The composition of CP maps is CP [33].

Theorem 2.65. Let A be a C^* -algebra. Every positive map $A \to \mathbb{C}$ is CP. In particular, a state on A is CP. Furthermore, every positive map between commutative C^* -algebras is CP.

Proof. See Theorems 3.9 and 3.11 in [33].

Notation 2.66. Henceforth, a *-homomorphism from a C^* -algebra \mathcal{B} to another C^* -algebra \mathcal{A} will be denoted by a straight arrow $\mathcal{B} \to \mathcal{A}$. A CP map, on the other hand, will be denoted with a curvy arrow $\mathcal{B} \leadsto \mathcal{A}$.

We now collect all the categories of C^* -algebras that will be needed in the prequel.

Notation 2.67. Let C*-AlgCP, cC*-AlgPos, and fdcC*-AlgPos denote the categories whose objects are C^* -algebras, commutative C^* -algebras, and finite-dimensional commutative C^* -algebras, respectively, and whose the morphisms are all CP maps of C^* -algebras (note that in the latter two cases, CP maps are the same as positive maps).

On a given physical system corresponding to some C^* -algebra, a state provides expectation values for observables. One can define a category of C^* -algebras equipped with a state together with the various flavors of morphisms analogous to the case of finite probability. However, this is tangential to our main goal and will follow as an immediate consequence anyway. Examples are provided in Section 2.7, but first we connect algebra with probability theory.

2.5 From probability theory to algebra

Now, we will construct a functor $C : \mathbf{FinSetStoch} \to \mathbf{fdcC^*-AlgPos}$ and show that it restricts to a well-defined functor $C : \mathbf{FinSetFun} \to \mathbf{fdcC^*-Alg}$. The same notation for these functors will be used as it will be clear from context which one is being referred to. The properties of these functors will be discussed in Section 2.6. Associated to each finite set Y, one has the C^* -algebra \mathbb{C}^Y discussed in Example 2.48. Note that any element $v \in \mathbb{C}^Y$ can be uniquely expressed as

$$v = \sum_{y \in Y} v_y e_y, \tag{2.68}$$

where $e_y: Y \to \mathbb{C}$ is defined by

$$Y \ni y' \mapsto e_y(y') := \delta_{yy'} \tag{2.69}$$

and $v_y \in \mathbb{C}$ is given by $v(y) =: v_y$. Hence, if X is another finite set, any CP map $\varphi : \mathbb{C}^Y \leadsto \mathbb{C}^X$ is determined by its value on the set of elements $\{e_y\}_{y \in Y}$ by linearity. Namely,

$$\varphi(v) = \sum_{y \in Y} v_y \varphi(e_y). \tag{2.70}$$

Hence, to every stochastic map $f: X \leadsto Y$, associate the linear function $C^f: \mathbb{C}^Y \leadsto \mathbb{C}^X$ uniquely determined by

$$C^f(e_y) := \sum_{x \in Y} f_{yx} e_x \tag{2.71}$$

for all $y \in Y$. These assignments define the functor C.

Theorem 2.72. The assignment⁵

FinSetStoch^{op}
$$\xrightarrow{C}$$
 fdcC*-AlgPos
$$X \mapsto \mathbb{C}^{X} \qquad (2.73)$$

$$\left(f: X \leadsto Y\right) \mapsto \left(C^{f}: \mathbb{C}^{Y} \leadsto \mathbb{C}^{X}\right),$$

where C^f is defined in (2.71), is a well-defined functor.

Proof. The proof will be broken into three steps. It will be shown that C^f is CP, the identity stochastic map goes to the identity CP map, and the composition of stochastic maps gets sent to the composition of CP maps.

⁵The op superscript is only meant to keep track that the directionality of the arrows get reversed. This is shown in (2.71), where the domain of C^f is \mathbb{C}^Y instead of \mathbb{C}^X . The directionality change is also shown in (2.77).

i. By Theorem 2.65, it suffices to show that C^f is positive. Every positive element in \mathbb{C}^Y is of the form

$$\sum_{y \in Y} a_y e_y, \quad \text{with } a_y \ge 0 \quad \forall \ y \in Y.$$
 (2.74)

Hence, by linearity, it suffices to show that $C^f(e_y)$, given by (2.71), is positive for every $y \in Y$. By definition of f being a stochastic map, $f_{yx} \ge 0$ for all $y \in Y$ and $x \in X$. Hence, $C^f(e_y)$ is a positive element in \mathbb{C}^X . Furthermore, C^f is unital because

$$C^{f}(1_{Y}) = C^{f}\left(\sum_{y \in Y} e_{y}\right) = \sum_{y \in Y} \left(C^{f}(e_{y})\right) = \sum_{y \in Y} \sum_{x \in X} f_{yx} e_{x} = \sum_{x \in X} \underbrace{\left(\sum_{y \in Y} f_{yx}\right)}_{1} e_{x} = 1_{X}. \quad (2.75)$$

ii. The identity function id: $X \to X$ has the corresponding stochastic map given by the probability measure $\mathrm{id}_{x'x} = \delta_{x'x}$ for all $x, x' \in X$. Therefore,

$$C^{\text{id}}(e_{x'}) = \sum_{x \in X} \text{id}_{x'x} e_{x'} = \sum_{x \in X} \delta_{x'x} e_x = e_{x'}$$
(2.76)

for all $x' \in X$. Thus, C^{id} is the identity CP map.

iii. Now suppose that $f: X \leadsto Y$ and $g: Y \leadsto Z$ are two composable stochastic maps between finite sets. This induces CP maps $C^f: \mathbb{C}^Y \leadsto \mathbb{C}^X$ and $C^g: \mathbb{C}^Z \leadsto \mathbb{C}^Y$. Therefore,

$$C^{f}(C^{g}(e_{z})) = C^{f}\left(\sum_{y \in Y} g_{zy} e_{y}\right)$$

$$= \sum_{y \in Y} g_{zy} C^{f}(e_{y})$$

$$= \sum_{y \in Y} g_{zy} \sum_{x \in X} f_{yx} e_{x}$$

$$= \sum_{x \in X} \left(\sum_{y \in Y} g_{zy} f_{yx}\right) e_{x}$$

$$= C^{g \circ f}(e_{z})$$


$$(2.77)$$

for all $z \in Z$.

Theorem 2.78. The functor $C: \mathbf{FinSetStoch^{op}} \to \mathbf{fdcC}^* - \mathbf{AlgPos}$ from Theorem 2.72 restricts to a well-defined functor $C: \mathbf{FinSetFun^{op}} \to \mathbf{fdcC}^* - \mathbf{Alg}$.

Proof. Using the notation of Theorem 2.72 and its proof, all that is left to show is that C^f : $\mathbb{C}^Y \leadsto \mathbb{C}^X$ is a *-homomorphism when f is a function. By linearity, it suffices to prove that

 $C^f(e_y e_{y'}) = C^f(e_y) C^f(e_{y'})$ for all $y, y' \in Y$. This follows from⁶



2.6 From algebra to probability theory

We will show that the functors of Theorems 2.72 and 2.78 are fully faithful.

Definition 2.80. A functor $F: \mathcal{D} \to \mathcal{E}$ from a category \mathcal{D} to a category \mathcal{E} is <u>fully faithful</u> iff for every pair of objects d and d' in \mathcal{D} , the function $F: \mathcal{D}(d,d') \to \mathcal{E}(F(d),F(d'))$ is a bijection (<u>full</u> means this function is surjective and <u>faithful</u> means it is injective). Here, $\mathcal{D}(d,d')$ denotes the set of morphisms from d to d' in the category \mathcal{D} .

In other words, if there is a fully faithful functor $\mathcal{D} \to \mathcal{E}$ that is also injective on objects, this allows one to study \mathcal{D} in a broader context without altering the relations between its objects. Being faithful means that no information is lost when viewing \mathcal{D} as inside of \mathcal{E} and being full means that no extra information is added to \mathcal{D} when viewing \mathcal{D} inside of \mathcal{E} . In particular, \mathcal{D} when viewed on its own has the same set (if it even is a set) of isomorphism classes of objects as when viewing \mathcal{D} inside of \mathcal{E} .

Theorem 2.81. The functors $C : \mathbf{FinSetStoch^{op}} \to \mathbf{fdcC^*-AlgPos} \ and \ C : \mathbf{FinSetFun^{op}} \to \mathbf{fdcC^*-Alg} \ from \ Theorems 2.72 \ and 2.78 \ are fully faithful and injective on objects.$

Proof. Injectivity on objects is immediate from the definition of C. Fix any two sets X and Y and a CP map $\varphi : \mathbb{C}^Y \leadsto \mathbb{C}^X$. The goal is to show there exists a unique stochastic map $f : X \leadsto Y$

⁶Diagrams like this are to be read from top to bottom going along either direction to recreate the thoughtprocess involved in proving the claim. The reason is because at some point near the bottom, a step which does not seem obvious in one direction is perhaps more intuitive from the other direction (namely, $\delta_{vf(x)}^2 = \delta_{vf(x)}$).

such that $C^f = \varphi$. Furthermore, when φ is a *-homomorphism, then the associated stochastic map f is a function. These two claims are proven in two steps.

i. First note that for any $y \in Y$, $\varphi(e_y)$ is an element of \mathbb{C}^X and, because $\{e_x\}_{x \in X}$ is a basis of \mathbb{C}^X , there exist unique numbers $\{f_{yx} \in \mathbb{C}\}_{x \in X}$ such that

$$\varphi(e_y) = \sum_{x \in X} f_{yx} e_x. \tag{2.82}$$

Positivity of φ demands that

$$f_{yx} \ge 0 \tag{2.83}$$

for all $x \in X$ (and $y \in Y$). Since φ is linear and unital,

$$\sum_{x \in X} e_x = 1_X = \varphi(1_Y) = \varphi\left(\sum_{y \in Y} e_y\right) = \sum_{y \in Y} \varphi(e_y) = \sum_{y \in Y} \sum_{x \in X} f_{yx} e_x = \sum_{x \in X} \left(\sum_{y \in Y} f_{yx}\right) e_x. \tag{2.84}$$

Since $\{e_x\}_{x\in X}$ is a linearly independent set of vectors in \mathbb{C}^X , this implies

$$\sum_{y \in Y} f_{yx} = 1. (2.85)$$

This shows that for every $x \in X$, the assignment

$$Y \ni y \mapsto f_{yx}$$
 (2.86)

is a probability measure on Y. In other words, the numbers $\{f_{yx}\}_{y\in Y,x\in X}$ define a stochastic map $f:X \rightsquigarrow Y$. It is readily checked that $C^f=\varphi$. This proves that C is full.

To see that C is faithful, suppose that $f': X \leadsto Y$ is another stochastic map such that $C^f = C^{f'}$. In particular, this implies that for any $y \in Y$,

$$C^{f}(e_{y})$$

$$C^{f'}(e_{y})$$

$$\sum_{x \in X} f_{yx}e_{x} \qquad \sum_{x \in X} f'_{yx}e_{x}$$

$$(2.87)$$

By linear independence of the set $\{e_x\}_{x\in X}$, this implies

$$f_{yx} = f'_{yx} \qquad \forall \ x \in X. \tag{2.88}$$

Since this calculation was independent of y, this also proves the equality for all $y \in Y$. Hence, f = f' proving that C is faithful. This concludes the proof that C is fully faithful.

ii. Using the same notation as in the previous step, it suffices to show that if $\varphi : \mathbb{C}^Y \to \mathbb{C}^X$ is actually a *-homomorphism, then the corresponding stochastic map $f : X \leadsto Y$ comes from

a function $f: X \to Y$. First notice that for any pair of distinct elements $y, y' \in Y$, because φ is a homomorphism,

$$\varphi(e_{y}e_{y'}) = \varphi(e_{y})\varphi(e_{y'})$$

$$\sum_{x \in X} \sum_{x' \in X} f_{yx}f_{y'x'}e_{x}e_{x'}$$

$$\sum_{x \in X} \sum_{x' \in X} f_{yx}f_{y'x'}\delta_{xx'}e_{x'}$$

$$\sum_{x \in X} f_{yx}f_{y'x}e_{x}$$

$$(2.89)$$

proving that

$$f_{yx}f_{y'x} = 0 \qquad \forall x \in X. \tag{2.90}$$

Putting this result aside for the time being, recall the result (2.85) which, together with the fact that $f_{yx} \ge 0$ for all $y \in Y$ and $x \in X$, implies that for each $x \in X$, there exists a $y \in Y$ such that $f_{yx} > 0$ (for otherwise, the sum over all y would not be 1). For any other $y' \in Y \setminus \{y\}$, the result (2.90) then implies

$$f_{y'x} = 0 \qquad \forall \ y' \in Y \setminus \{y\}. \tag{2.91}$$

Using (2.85) again implies

$$f_{yx} = 1. (2.92)$$

In other words, for every $x \in X$, there exists a unique $y \in Y$ such that $f_{yx} = 1$ and $f_{y'x} = 0$ for all $y' \in Y \setminus \{y\}$. Set f(x) to be this unique element y, i.e. f(x) := y. This assignment then defines a function $f: X \to Y$.

It is a fact that every n-dimensional commutative C^* -algebra is *-isomorphic to \mathbb{C}^X for some set X with cardinality n (this will follow from results in Section 3). This, together with Theorem 2.81, says that **FinSetStoch**^{op} is equivalent to the category **fdcC*-AlgPos** consisting of finite-dimensional commutative C^* -algebras and positive maps. An explicit inverse will be constructed in greater generality in Section 4.5.

2.7 Some quantum mechanics

Completely positive maps represent general processes that are allowed in quantum mechanics. We will provide three different examples illustrating the versatility of the C^* -algebraic approach. The first two examples are actually examples of *-homomorphisms. We thank Stefan Andronache and Marcelo Nomura for discussions leading to these examples.

Example 2.93. Let \mathcal{H} be a finite-dimensional⁷ Hilbert space and H a self-adjoint operator, thought of as a Hamiltonian. For each $t \in \mathbb{R}$, the operator $\exp\left(-\frac{itH}{\hbar}\right)$, where \hbar is the reduced Planck constant, is unitary and describes unitary time evolution in quantum mechanics. Its action on observables is given by the adjoint action (the Heisenberg picture)

$$\mathcal{B}(\mathcal{H}) \to \mathcal{B}(\mathcal{H})$$

$$A \mapsto \exp\left(\frac{itH}{\hbar}\right) A \exp\left(-\frac{itH}{\hbar}\right). \tag{2.94}$$

Infinitesimally, this gives the differential equation that describes the time evolution of an observable

$$\frac{dA}{dt} = \frac{1}{i\hbar}[A, H]. \tag{2.95}$$

Now, given an initial state $\operatorname{tr}(\rho \cdot) : \mathcal{B}(\mathcal{H}) \leadsto \mathbb{C}$ for some density matrix ρ (see Theorem 2.54 for notation) this pulls back to a new state $\operatorname{tr}(\rho \cdot) \circ \operatorname{Ad}_{\exp\left(-\frac{itH}{\hbar}\right)}$ under this evolution

$$\mathcal{B}(\mathcal{H}) \xrightarrow{\operatorname{Ad}_{\exp\left(-\frac{itH}{\hbar}\right)}} \mathcal{B}(\mathcal{H})$$

$$\operatorname{tr}(\rho \cdot) \circ \operatorname{Ad}_{\exp\left(-\frac{itH}{\hbar}\right)} \qquad \qquad (2.96)$$

and is given by the assignment

$$\mathcal{B}(\mathcal{H}) \to \mathbb{C}$$

$$A \mapsto \operatorname{tr}\left(\rho \exp\left(\frac{itH}{\hbar}\right) A \exp\left(-\frac{itH}{\hbar}\right)\right) = \operatorname{tr}\left(\exp\left(-\frac{itH}{\hbar}\right) \rho \exp\left(\frac{itH}{\hbar}\right) A\right) \tag{2.97}$$

by cyclicity of the trace. By the uniqueness of density matrices representing states (Theorem 2.54), this proves that the time evolution of ρ is given by the differential equation

$$\frac{d\rho}{dt} = \frac{1}{i\hbar}[H, \rho]. \tag{2.98}$$

This is the quantum Liouville equation. Notice how it has the opposite sign of the evolution for observables. Also notice that if ρ is a pure state, this reduces to Schrödinger's equation and reproduces the Schrödinger picture. Thus, the C^* -algebraic approach naturally incorporates both perspectives. It is important to notice that the straight horizontal arrow in (2.96) indicates deterministic evolution! The "non-determinacy" of this example only appears in the states and their expectation values.

Example 2.99. Measurement is an example of a CP map, in fact a *-homomorphism. We think of measuring a quantum mechanical system in terms of numbers. For example, in the Stern-Gerlach experiment, we measure a particle (such as a silver atom) to be spin up or spin down depending on its position on a screen after moving through a magnetic field [41]. A state of a quantum-mechanical system is therefore reduced to a probability measure on the set of the eigenvalues of the observable being measured.

⁷We use finite-dimensional Hilbert spaces to avoid technicalities involving domains of operators.

The general (finite-dimensional) situation is as follows.⁸ Let \mathcal{H} be a finite-dimensional Hilbert space, let A be a self-adjoint operator on \mathcal{H} , and let ω be a state on $\mathcal{B}(\mathcal{H})$. Let $\sigma(A)$ denote the spectrum of A (in this case, $\sigma(A)$ is just the set of eigenvalues of A). A is an observable and $\sigma(A)$ are the possible values that an observer will see when trying to measure the observable A. It is a fact of life that an observer can *only* measure elements of $\sigma(A)$ (this is the origin of the "discreteness" of quantum mechanics). In this situation, what is the probability measure that an observer expects to see on $\sigma(A)$? The resulting probability measure is obtained by pulling back ω along the "measurement" map

$$m: \mathbb{C}^{\sigma(A)} \to \mathcal{B}(\mathcal{H})$$

$$e_{\lambda} \mapsto P_{\lambda}, \tag{2.100}$$

where P_{λ} is the projection operator onto the eigenspace associated with the eigenvalue $\lambda \in \sigma(A)$ of A. This produces a state on $\mathbb{C}^{\sigma(A)}$ via pullback

$$\mathbb{C}^{\sigma(A)} \xrightarrow{m} \mathcal{B}(\mathcal{H}) \tag{2.101}$$

By our analysis from Section 2.6, $p := \omega \circ m$ precisely provides us with a probability measure on $\sigma(A)$. This probability measure is interpreted as the probability distribution of measuring the corresponding eigenvalues of A with respect to the state ω . Note that this example does not have an evolution of the form $\mathcal{B}(\mathcal{H}) \to \mathcal{B}(\mathcal{K})$ between observables on Hilbert spaces \mathcal{H} and \mathcal{K} . Instead, it is an "evolution" from the quantum phase space $\mathcal{B}(\mathcal{H})$ to the classical one $\sigma(A)$. Hence, the C^* -algebraic formulation allows one to treat classical and quantum systems in the same category.

In both of the previous situations, the evolution was described by a *-homomorphism. In the following example, we provide completely positive evolution that is not a *-homomorphism. Such evolutions can be used to model noise and measurement among other things [30].

Example 2.102. Let $\{|\uparrow\rangle, |\downarrow\rangle\}$ denote the standard Euclidean basis in \mathbb{C}^2 . Let $|\uparrow\rangle\langle\uparrow|$ denote the projection operator onto span $(|\uparrow\rangle)$ and similarly for $|\downarrow\rangle$. The states on $\mathcal{B}(\mathbb{C}^2)$ given by $\operatorname{tr}(|\uparrow\rangle\langle\uparrow|\cdot)$ and $\operatorname{tr}(|\downarrow\rangle\langle\downarrow|\cdot)$ can be interpreted as a qubit being either in the spin up or spin down state with respect to an appropriately chosen coordinate frame. Consider an operation on the space of all density matrices whose action on these two states is given by

$$|\uparrow\rangle\langle\uparrow|\mapsto p|\uparrow\rangle\langle\uparrow| + (1-p)|\downarrow\rangle\langle\downarrow| |\downarrow\rangle\langle\downarrow|\mapsto (1-p)|\uparrow\rangle\langle\uparrow| + p|\downarrow\rangle\langle\downarrow|$$
 (2.103)

for some fixed $p \in [0,1]$. This is to be interpreted as an operation that has probability 1-p of flipping a spin up state to a spin down state and has probability 1-p of flipping a spin down state to a spin up state. Does there exist a completely positive map $f : \mathcal{B}(\mathbb{C}^2) \leadsto \mathcal{B}(\mathbb{C}^2)$ that

⁸For infinite systems, one must use projection-valued measures [30].

pulls back the pure states above on the left in (2.103) to the states on the right in (2.103)? The answer is yes. Let

$$E := \sqrt{p} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \qquad \& \qquad F := \sqrt{1-p} \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$$
 (2.104)

and define f to be given by

$$\mathcal{B}(\mathbb{C}^2) \ni A \mapsto E^* A E + F^* A F. \tag{2.105}$$

One can check that f is completely positive and is not a *-homomorphism. The induced action on the density matrices can be obtained by looking at the pullback of an initial density matrix ρ

$$\mathcal{B}(\mathbb{C}^2) \ni A \mapsto \operatorname{tr}(\rho f(A)) = \operatorname{tr}(\rho E^* A E + \rho F^* A F) = \operatorname{tr}((E \rho E^* + F \rho F^*) A). \tag{2.106}$$

Hence, the resulting density matrix under such an operation is given by $E\rho E^* + F\rho F^*$. One can check that (2.103) holds under this transformation. So again, the C^* -algebraic framework is able to reproduce familiar operations such as this one, known as the bit flip channel [30], all in the same category of physical processes.

A generalization of Example 2.93 to CP maps gives rise to Lindblad's equation describing non-deterministic dynamics in (open) quantum systems [27].

3 An equivalence between spaces and algebraic structures

On their own, the study of topological spaces and continuous maps is quite different from the study of algebra and homomorphisms. Topological spaces consist of sets equipped with a subset of the power set (known as open sets) and continuous maps are functions whose pre-image function takes open sets to open sets. Algebras are sets equipped with binary operations satisfying certain laws such as associativity and maps between algebras are those that respect this algebraic structure. In the plain context of set theory, it simply does not make sense to compare the two areas of mathematics. However, viewing these two areas as a whole, which is achieved by viewing them as categories, it makes sense to compare them. In fact, under additional suitable restrictions on the respective sides, the two categories are equivalent. There are several other versions of such equivalences between topological categories and algebraic ones [23], [22]. We will discuss one such equivalence, the equivalence between the category of commutative (unital) C^* -algebras and *homomorphisms with the category of compact Hausdorff topological spaces and continuous maps following the expositions in Folland [13], Rudin [39], and Tao's online notes [47]. The references already have well-written proofs of the main results. Our purpose here is to frame these results in the appropriate categorical context and establish certain tools that will be used in Section 4 when we relax the class of morphisms to include non-deterministic processes.

3.1 From spaces to algebras

In Section 2.5, we showed how given a finite set X, the set of functions \mathbb{C}^X can be made into a C^* -algebra in a natural manner. This can also be done if X is replaced with a compact Hausdorff space and the function space is chosen to be the subspace of *continuous* functions from X to \mathbb{C} .

Example 3.1. Let X be a compact Hausdorff space. Then the space of complex-valued continuous functions, denoted by C(X), with the algebraic structure from \mathbb{C}^X and with norm given by the uniform norm, i.e.

$$C(X) \ni \varphi \mapsto \|\varphi\| := \sup_{x \in X} |\varphi(x)|,$$
 (3.2)

is a commutative C^* -algebra. Cauchy completeness follows from the continuous (uniform) limit theorem (see Theorem 7.12 in [37] or Theorem 21.6 in [29]). The constant function 1_X is the unit for C(X). Note that although compactness is needed for C(X) to be a C^* -algebra with this norm, it is not necessary that X be Hausdorff. Proving that C(X) is a C^* -algebra under the only assumption that X is compact is a good exercise. If X is not Hausdorff, there exist points that cannot be separated by continuous complex valued functions so that continuous functions cannot distinguish them (see Remark 3.65 and Theorem 4.102 for further details).

Proposition 3.3. Let X and Y be compact Hausdorff spaces and let $f: X \to Y$ be a continuous function. The function $C^f: C(Y) \to C(X)$ defined by

$$C(Y) \ni \beta \mapsto C^f(\varphi) := \varphi \circ f$$
 (3.4)

 $is \ a *-homomorphism.$

Proof. First note that $C^f(\varphi)$ is continuous since it is the composition of continuous functions. To see that C^f respects the product, let $\psi, \varphi \in C(Y)$ and let $x \in X$. Then

$$(C^{f}(\psi\varphi))(x) = (\psi\varphi)(f(x)) \quad \text{by def'n of } C^{f}$$

$$= (\psi(f(x)))(\varphi(f(x))) \quad \text{by def'n of } \psi\varphi$$

$$= ((\psi\circ f)(x))((\varphi\circ f)(x)) \quad \text{by def'n of } (\phi\circ f)(\varphi\circ f)$$

$$= ((\psi\circ f)(\varphi\circ f))(x) \quad \text{by def'n of } (\psi\circ f)(\varphi\circ f)$$

$$= (C^{f}(\psi)C^{f}(\varphi))(x) \quad \text{by def'n of } C^{f}$$

$$(3.5)$$

proving that $C^f(\psi\varphi) = C^f(\psi)C^f(\varphi)$. Linearity, *-preservation, and preservation of units follow from analogous calculations.

Notation 3.6. Let **cHaus** denote the category whose objects are compact Hausdorff spaces and whose morphisms are continuous maps.

Proposition 3.7. The assignment

$$\mathbf{cHaus}^{\mathrm{op}} \xrightarrow{C} \mathbf{cC} *-\mathbf{Alg}$$

$$X \mapsto C(X)$$

$$\left(X \xrightarrow{f} Y\right) \mapsto \left(C(X) \xleftarrow{C^f} C(Y)\right)$$
(3.8)

from Example 3.1 and Proposition 3.3 defines a functor.

⁹Sketch of proof: start with a Cauchy sequence in C(X), use Cauchy completeness in \mathbb{C} to obtain a candidate function pointwise, show that the resulting function is bounded, then show that the sequence of functions converges to it uniformly, and finally apply the uniform limit theorem to guarantee it is continuous.

Proof. From the definition of C, it is manifest that C is a functor: the identity $\mathrm{id}_X:X\to X$ gets sent to $C^{\mathrm{id}_X}=\mathrm{id}_{C(X)}$ and the composition $X\xrightarrow{f}Y\xrightarrow{g}Z$ gets sent to the function

$$C(Z)\ni\varphi\mapsto C^{g\circ f}(\varphi)=\varphi\circ(g\circ f)=(\varphi\circ g)\circ f=C^f\big(\varphi\circ g\big)=C^f\big(C^g(\varphi)\big)=(C^f\circ C^g)(\varphi)\ \ (3.9)$$

since composition of functions is associative.

3.2 Some topological preliminaries

Given a Banach algebra \mathcal{A} , a certain subspace, denoted by \mathcal{A}^* , of the linear dual space \mathcal{A}^{\vee} has a natural topology (the operator norm topology) coming from the norms on \mathcal{A} and \mathbb{C} . However, this topology has too many open sets for some purposes. Another topology on \mathcal{A}^* that has fewer open sets is the weak* topology. Fewer open sets allows more sets to be compact and more compact sets means that sequences and nets have a better chance of converging (see Tao's notes for an illuminating discussion [47]). There are several ways to describe this topology. The simplest for our purposes is in terms of convergent nets, though it will also be useful to describe it in terms of topological vector spaces whose topology is induced by a family of seminorms. This latter perspective will also become more important in Section 4.

Definition 3.10. A seminorm on a vector space V is a function $\|\cdot\|: V \to \mathbb{R}$ satisfying

- i. $||v|| \ge 0$ for all $v \in V$,
- ii. $\|\lambda v\| = |\lambda| \|v\|$ for all $\lambda \in \mathbb{C}, v \in V$, and
- iii. $||v + w|| \le ||v|| + ||w||$ for all $v, w \in V$.

The only difference between a norm and a seminorm is that a norm satisfies the additional condition that ||v|| = 0 implies v = 0.

Example 3.11. Let V be a vector space and $(W, \| \cdot \|)$ be a normed vector space. For any linear map $\varphi : V \to W$, the function $\| \cdot \|_{\varphi} : V \to \mathbb{R}$ defined by sending $v \in V$ to $\| \varphi(v) \|$ is a seminorm on V. A particular case of interest is when $W := \mathbb{C}$.

A single norm provides a natural topology via open balls of varying radii at different points. While a single seminorm also provides a natural topology in a similar way, it is rarely Hausdorff and does not enjoy many desirable properties (see Example 3.15). However, it does become a topological vector space.

Definition 3.12. A <u>topological vector space</u> consists of a vector space V equipped with a topology in which the addition of vectors $+: V \times V \to V$ and scalar multiplication $\mathbb{C} \times V \to V$ are both continuous.

Definition 3.13. A <u>base</u> on a set X is a collection $\mathfrak{B} \subseteq \mathcal{P}(X)$ of subsets of X satisfying

- i. for each $x \in X$, there exists a $B \in \mathfrak{B}$ such that $x \in B$,
- ii. for every pair $B_1, B_2 \in \mathfrak{B}$ and every $x \in B_1 \cap B_2$, there exists a $B \in \mathfrak{B}$ such that $x \in B \subseteq B_1 \cap B_2$.

A base \mathfrak{B} as above can be used to construct a topology on X by setting $\tau_{\mathfrak{B}}$ to be the smallest topology containing \mathfrak{B} . Explicitly, its elements are given by unions of all possible elements in \mathfrak{B} ,

$$\tau_{\mathfrak{B}} = \left\{ \bigcup_{B \in \mathfrak{C}} B : \mathfrak{C} \subseteq \mathfrak{B} \right\}. \tag{3.14}$$

This topology is called the topology generated by \mathfrak{B} .

Example 3.15. Let $\|\cdot\|$ be a seminorm on a vector space V. Then the set

$${B_r(v) \subseteq V : v \in V \text{ and } r > 0},$$
 (3.16)

where

$$B_r(v) := \{ w \in V : ||v - w|| < r \}, \tag{3.17}$$

forms a base on V. With respect to the topology generated by this base, V is a topological vector space. Furthermore, it is Hausdorff if and only if $\|\cdot\|$ is a norm (see Tao [47]). $B_r(v)$ is called the open ball of radius $r \geq 0$ centered at $v \in V$.

A sufficiently robust *family* of seminorms typically has a more manageable topology that naturally arises frequently enough to merit study.

Notation 3.18. For $n \in \mathbb{N}$, set $\overline{n} := \{1, \dots, n\}$. Let X be a set and $\mathfrak{S} \subseteq \mathcal{P}(X)$ a subset of the power set.¹⁰ The value of a function $S : \overline{n} \to \mathfrak{S}$ at $i \in \overline{n}$ is denoted by S_i instead of S(i).

Definition 3.19. A <u>subbase</u> on a set X is a collection $\mathfrak{S} \subseteq \mathcal{P}(X)$ of subsets of X such that the set

$$\mathfrak{B} := \bigcup_{n \in \{0\} \cup \mathbb{N}} \left\{ \bigcap_{i=1}^{n} S_i : \overline{n} \xrightarrow{S} \mathfrak{S} \right\}$$
 (3.20)

of all finite intersections of elements in \mathfrak{S} is a base for X. Here, $\bigcap_{i=1}^{0} S_i$ is the empty intersection, which is taken to be X itself. The topology generated by \mathfrak{S} is the topology generated by \mathfrak{B} .

It is a fact that any collection of subsets of a set forms a subbase for some topology on that set and that topology is the smallest topology containing those subsets (see Section 5 of [50]).

Proposition 3.21. Let V be a vector space with $\{\|\cdot\|_{\alpha}\}_{\alpha\in A}$ a family of seminorms on V indexed by some set A. Let τ_{α} denote the topology associated to $\|\cdot\|_{\alpha}$. Then $\mathfrak{S}:=\bigcup_{\alpha\in A}\tau_{\alpha}$ is a subbase on V and V with the topology generated by \mathfrak{S} is a topological vector space. Furthermore, V is Hausdorff if and only if for each $v\in V\setminus\{0\}$, there exists an $\alpha\in A$ such that $\|v\|_{\alpha}\neq 0$. Such a family of seminorms is said to separate points of V or is said to be a separating family.

Proof. See Theorem 5.14 and Proposition 5.16 in Folland [14].

 $^{^{10}\}mathfrak{S}$ is a Fraktur "S."

Remark 3.22. The use of seminorms is not necessary in the conclusion that V is a topological vector space. All that is required is that each of the topologies τ_{α} provide V with the structure of a topological vector space [47].

Let V be a vector space and let V^{\vee} denote its linear dual (linear functions from V to \mathbb{C}) In many applications where V has a norm, V^{\vee} is too large to deal with. Instead, one uses the norm to restrict to a certain subspace of linear functionals.

Proposition 3.23. Let V be a normed vector space with norm $\|\cdot\|$. For $\chi \in V^{\vee}$, set (abusing notation slightly)

$$\|\chi\| := \inf \left\{ M \ge 0 : |\chi(v)| \le M \|v\| \quad \forall \ v \in V \right\},$$
 (3.24)

which can be infinite. Set $V^* \subseteq V^{\vee}$ to be the subset given by

$$V^* := \{ \chi \in V^{\vee} : \|\chi\| < \infty \}. \tag{3.25}$$

Then $\|\cdot\|$ is a norm on V^* and V^* equipped with this norm is a Banach space, known as the topological dual space of V. Elements of V^* are called bounded/continuous linear functionals on V.

Proof. This is a general fact about operator norms (see Proposition 5.4 in [14]).

Proposition 3.26. Let V be a vector space. For each $v \in V$, define $\|\cdot\|_v : V^{\vee} \to \mathbb{C}$ by

$$V^{\vee} \ni \chi \mapsto \|\chi\|_v := |\chi(v)|. \tag{3.27}$$

Then $\|\cdot\|_v$ is a seminorm on V^{\vee} . Furthermore, for each $\chi \in V^{\vee} \setminus \{0\}$, there exists a $v \in V$ such that $\chi(v) \neq 0$, i.e. the family of seminorms $\{\|\cdot\|_v\}_{v \in V}$ is separating. The same statement holds for $V^* \subseteq V^{\vee}$ with the same family.

Proof. This immediately follows from the definitions.

Definition 3.28. Let V be a vector space and let $\{\|\cdot\|_v\}_{v\in V}$ be the family of seminorms as in Proposition 3.26. The Hausdorff topology generated by this family as described in Proposition 3.21 is known as the $weak^*$ topology on V^* .

Proposition 3.29. Let V be a normed vector space. For each $\chi \in V^*$, $\epsilon > 0$, $n \in \mathbb{N}$, and $\alpha_n : \overline{n} \to V$, set

$$B_{\epsilon}^{\alpha_n}(\chi) := \bigcap_{j \in \overline{n}} \left\{ \xi \in V^* : \|\chi - \xi\|_{\alpha_n(j)} < \epsilon \right\}. \tag{3.30}$$

Then

$$\mathfrak{B} := \left\{ B_{\epsilon}^{\alpha_n}(\chi) \subseteq V^{\vee} : \epsilon > 0, \ n \in \mathbb{N}, \ \alpha_n : \overline{n} \to V, \ \chi \in V^* \right\}$$
 (3.31)

is a base for the weak* topology on V^* .

Proof. This is a general fact for the topology generated by a separating family of seminorms—see Section 2.4 in [34].

In other words, a base for the weak* topology consists of finite intersections of balls coming from the different seminorms.

Proposition 3.32. Let V be a vector space. A $net^{11} \chi : \Theta \to V^*$ in V^* converges to an element $\lim \chi \in V^*$ in the weak* topology if and only if

$$\lim_{\theta \in \Theta} (\chi_{\theta}(v)) = (\lim \chi)(v) \qquad \forall v \in V.$$
(3.33)

Proof. See Tao [47] or Section 5.4 of Folland [14].

Nets are needed here because not all compact Hausdorff spaces are separable (see Section 4.2).

Proposition 3.34. Let V be a vector space and let V^* denote its topological dual space. The weak* topology on V^* is the weakest topology (meaning that it has the fewest open sets) such that the functions

$$V^* \xrightarrow{\operatorname{ev}_v} \mathbb{C}$$

$$\chi \mapsto \chi(v) \tag{3.35}$$

are continuous for all $v \in V$.

Proof. See Section 3.14 of [39].

Theorem 3.36 (Banach-Alaoglu Theorem). Let V be a Banach space and let V^* denote its topological dual space. The closed unit ball (with respect to the norm on V^*) in V^* is compact with respect to the weak* topology.

This theorem is surprising because the closed (and bounded) unit ball is compact with respect to the norm topology if and only if the normed vector space is finite-dimensional [47].

3.3 The spectrum of a commutative C^* -algebra

It is perhaps not so surprising that one can obtain a commutative C^* -algebra from a space as in Section 3.1. What is more surprising is that there is a way to go back from C^* -algebras to topological spaces. Although C^* -algebras have a norm (and hence a topology), keep in mind that a C^* -algebra is a *linear* space so obtaining a non-trivial topology from such an object will require some work.

Definition 3.37. Let \mathcal{A} be a commutative Banach algebra. A <u>character</u> on \mathcal{A} is a continuous nonzero homomorphism $\chi : \mathcal{A} \to \mathbb{C}$ of Banach algebras, i.e. χ is linear, satisfies $\chi(ab) = \chi(a)\chi(b)$ for all $a, b \in \mathcal{A}$, and satisfies the condition that there exists an $a \in \mathcal{A}$ for which $\chi(a) \neq 0$.

Characters form a subset of \mathcal{A}^* , the topological dual space of \mathcal{A} . This dual space has a natural topology coming from the norms on both \mathcal{A} and \mathbb{C} , but the topology of interest for us is the weak* topology on \mathcal{A}^* .

¹¹A net is a function whose domain is a directed set (see Section 11 of [50]).

Definition 3.38. Let \mathcal{A} be a commutative Banach algebra. The *spectrum of* \mathcal{A} is the set

$$\sigma(\mathcal{A}) := \left\{ \mathcal{A} \xrightarrow{\chi} \mathbb{C} : \chi \text{ is a character} \right\}. \tag{3.39}$$

equipped with the subspace topology coming from \mathcal{A}^* via the weak* topology.

Remark 3.40. The relationship between the spectrum of a commutative C^* -algebra and the spectrum of a certain operator on a Hilbert space is described in Proposition 1.15 in [13]. Briefly, if A is bounded and normal, then the operator-theoretic spectrum of A is canonically homeomorphic with the spectrum of the commutative unital *-subalgebra generated by A. This is the closure of the set of polynomials in A and A^* . By definition of being normal, A and A^* commute so that the order does not matter in which the A's and A^* 's appear in such polynomials guaranteeing that this C^* -algebra is commutative.

Proposition 3.41. Let A be a Banach algebra and let $\chi \in \sigma(A)$ be a character. Then

- i. $\chi(1_A) = 1$,
- ii. $\chi(a) \neq 0$ for all invertible elements $a \in \mathcal{A}$, and
- iii. $|\chi(a)| \leq ||a||$ for all $a \in \mathcal{A}$.

Proof. See Proposition 1.10 in [13].

Proposition 3.42. Let A be a commutative Banach algebra. Then the spectrum $\sigma(A)$ is a nonempty compact Hausdorff space.

Proof. See Theorem 11.9 in [39]. $\sigma(A)$ is nonempty because A is unital.

Proposition 3.43. Let $\mathcal{B} \xrightarrow{f} \mathcal{A}$ be *-homomorphism of commutative Banach algebras. Then the function

$$\sigma(\mathcal{A}) \xrightarrow{\sigma^f} \sigma(\mathcal{B})$$

$$\chi \mapsto \chi \circ f \tag{3.44}$$

is continuous.

Proof. First, note that since

$$(\chi \circ f)(xy) = \chi(f(x)f(y)) = \chi(f(x))\chi(f(y)) = (\chi \circ f)(x)(\chi \circ f)(y)$$
(3.45)

for all $x, y \in \mathcal{B}$, $\chi \circ f$ is indeed an element of $\sigma(\mathcal{B})$. We will give two proofs of continuity, one using open sets and the other using nets.

i. It suffices to show that the inverse image of a base element gets sent to an open set. By Proposition 3.29 and by definition of the subspace topology, a base for the topology on $\sigma(\mathcal{B})$ is given by

$$\sigma(\mathcal{B}) \cap B_{\epsilon}^{\alpha_n}(\xi) = \left\{ \chi \in \sigma(\mathcal{B}) : \|\chi - \xi\|_{\alpha_n(j)} < \epsilon \quad \forall \ j \in \overline{n} \right\}$$
 (3.46)

over all $\xi \in \mathcal{B}^*$, $\alpha_n : \overline{n} \to \mathcal{B}$, $\epsilon > 0$, $n \in \mathbb{N}$. σ^f is continuous if σ^{12} for every $\chi \in \sigma(\mathcal{A})$ and every basic set $\sigma^{13} = \sigma(\mathcal{B}) \cap B_{\epsilon}^{\alpha_n}(\chi \circ f)$ there exists an open set containing χ whose image is contained in this set. In fact, the basic set $\sigma^{f \circ \alpha_n}(\chi)$ accomplishes this goal because

$$\|\chi - \xi\|_{f \circ \alpha_n(j)} = \left| \chi \left(f \left(\alpha_n(j) \right) \right) - \xi \left(f \left(\alpha_n(j) \right) \right) \right| = \|\chi \circ f - \xi \circ f\|_{\alpha_n(j)}, \tag{3.47}$$

which shows that

$$\sigma^{f}\left(\sigma(\mathcal{A}) \cap B_{\epsilon}^{f \circ \alpha_{n}}(\chi)\right) = \left\{\xi \circ f \in \sigma(\mathcal{B}) : \|\chi - \xi\|_{f \circ \alpha_{n}(j)} < \epsilon \quad \forall \ j \in \overline{n}\right\}$$

$$= \left\{\xi \circ f \in \sigma(\mathcal{B}) : \|\chi \circ f - \xi \circ f\|_{\alpha_{n}(j)} < \epsilon \quad \forall \ j \in \overline{n}\right\}$$

$$\subseteq \sigma(\mathcal{B}) \cap B_{\epsilon}^{\alpha_{n}}(\chi \circ f).$$
(3.48)

Thus, σ^f is continuous.

ii. Let $\chi: \Theta \to \sigma(\mathcal{A})$ be a net converging to $\lim \chi$ (in the weak* topology). The goal is to show that the net $\sigma^f \circ \chi: \Theta \to \sigma(\mathcal{B})$ converges to $\sigma^f(\lim \chi) \equiv (\lim \chi) \circ f$. For every $b \in \mathcal{B}$,

$$\lim_{\theta \in \Theta} \left(\sigma^f(\chi_\theta)(b) \right) = \lim_{\theta \in \Theta} \left(\chi_\theta(f(b)) \right) = (\lim \chi) \left(f(b) \right) = \left(\sigma^f(\lim \chi) \right) (b), \tag{3.49}$$

which establishes the required weak* convergence and thus shows that σ^f is continuous.

As the reader may have noticed, the proof was much shorter using nets. The technique of using nets to prove continuity of various functions will be heavily used in Section 4.

Proposition 3.50. The assignment

$$\mathbf{cC}^*\text{-}\mathbf{Alg}^{\mathrm{op}} \xrightarrow{\sigma} \mathbf{cHaus}$$

$$\mathcal{A} \mapsto \sigma(\mathcal{A})$$

$$\left(\mathcal{B} \xrightarrow{f} \mathcal{A}\right) \mapsto \left(\sigma(\mathcal{B}) \xleftarrow{\sigma^f} \sigma(\mathcal{A})\right)$$
(3.51)

from Definition 3.38 and Proposition 3.43 defines a functor.

Proof. The proof is completely analogous to the proof of Proposition 3.7.

¹²Let $f: X \to X'$ be a function and let \mathfrak{B} and \mathfrak{B}' be bases for topologies on X and X', respectively. f is continuous if for every $x \in X$, and for any $B' \in \mathfrak{B}'$ with $f(x) \in B'$, there exists a $B \in \mathfrak{B}$ with $x \in B$ such that $f(B) \subseteq B'$ (this is a simple exercise in the definitions).

¹³Technically, we should have said that for every $\chi \in \sigma(A)$ and every basic set B containing $\chi \circ f$, there exists an open set U containing χ whose image is contained in B. However, due to the seminorms, it suffices to take the basic set B containing $\chi \circ f$ to be centered at $\chi \circ f$ by choosing it to be sufficiently small.

3.4 The Gelfand transform

Up to this point, we have constructed functors

$$\mathbf{cC^*-Alg}^{\mathrm{op}} \xrightarrow{\sigma} \mathbf{cHaus} . \tag{3.52}$$

In general, the diagrams

cHaus
$$\leftarrow$$
 id cHaus cC^* -Alg \leftarrow id cC^* -Alg \leftarrow cC^* -Alg \leftarrow cC^* -Alg \leftarrow cHaus \rightarrow cHa

do not commute so that σ and C are not inverses of each other. However, they are close. In the present section, we will construct natural isomorphisms

indicating a precise sense in which σ and C are close to being inverses of each other. A natural isomorphism between functors is analogous to a homotopy between continuous functions or a unitary intertwiner between representations and what it amounts to more precisely will be described presently. This will show that the categories \mathbf{cC}^* -Alg^{op} and **cHaus** are *equivalent*.

Notation 3.55. Let \mathcal{A} be a commutative Banach algebra and set

$$\mathcal{A} \xrightarrow{\Gamma_{\mathcal{A}}} C(\sigma(\mathcal{A}))$$

$$a \mapsto \left(\chi \mapsto \left(\Gamma_{\mathcal{A}}(a)\right)(\chi) := \chi(a)\right). \tag{3.56}$$

 $\Gamma_{\mathcal{A}}$ is called the <u>Gelfand transform on \mathcal{A} </u>.

Proposition 3.57. The following facts are true regarding the Gelfand transform.

- i. For every commutative Banach algebra \mathcal{A} , the Gelfand transform $\Gamma_{\mathcal{A}}: \mathcal{A} \to C_c(\sigma(\mathcal{A}))$ is a homomorphism of Banach algebras (preserving units).
- ii. If A is a commutative C^* -algebra, then Γ_A is a *-isomorphism (in particular, it is isometric).

Proof. See Theorem 1.13 parts (a) and (d) and Theorem 1.20 in [13] for the first and second claims, respectively.

Remark 3.58. If \mathcal{A} is just an involutive Banach algebra, then $\Gamma_{\mathcal{A}}(a^*)$ is not always equal to $(\Gamma_{\mathcal{A}}(a))^*$ for all $a \in \mathcal{A}$. Hence, $\Gamma_{\mathcal{A}}$ is not necessarily a *-homomorphism (see Proposition 1.14 in [13]).

Proposition 3.59. The assignment

$$\mathbf{cC}^* - \mathbf{Alg}_0^{\mathrm{op}} \xrightarrow{\Gamma} \mathbf{cC}^* - \mathbf{Alg}_1^{\mathrm{op}}$$

$$\mathcal{A} \mapsto \left(\mathcal{A} \xrightarrow{\Gamma_{\mathcal{A}}} C(\sigma(\mathcal{A})) \right)$$
(3.60)

from Definition 3.55 defines a natural isomorphism $\Gamma: \mathrm{id}_{\mathbf{c}\mathbf{C}^*\text{-}\mathbf{Alg}} \Rightarrow C \circ \sigma$.

Proof. Associated to any *-homomorphism $\mathcal{B} \xrightarrow{f} \mathcal{A}$ of C^* -algebras is the diagram

$$\mathcal{B} \xrightarrow{\Gamma_{\mathcal{B}}} C(\sigma(\mathcal{B}))$$

$$f \downarrow \qquad \qquad \downarrow_{C^{\sigma(f)}}.$$

$$\mathcal{A} \xrightarrow{\Gamma_{\mathcal{A}}} C(\sigma(\mathcal{A}))$$

$$(3.61)$$

To see that it commutes, let $b \in \mathcal{B}$. Going along the top and right arrows gives $C^{\sigma(f)}(\Gamma_{\mathcal{A}}(b)) = \Gamma_{\mathcal{A}}(b) \circ \sigma^f$ by definition of C as a functor (see Proposition 3.7). Going along the left and bottom arrows gives $\Gamma_{\mathcal{A}}(f(b))$. To see that these two elements of $C(\sigma(\mathcal{A}))$ are equal, let $\chi \in \sigma(\mathcal{A})$. Then

$$\Gamma_{\mathcal{B}}(b)\left(\sigma^{f}(\chi)\right) \qquad \Gamma_{\mathcal{A}}(f(b))\left(\chi\right)$$

$$(3.44) \qquad \qquad (3.56)$$

$$\Gamma_{\mathcal{B}}(b)\left(\chi \circ f\right) \qquad \chi(f(b))$$

$$(3.56) \qquad \qquad (3.62)$$

which proves that Γ is a natural transformation. It is a natural *iso* morphism by ii. of Proposition 3.57.

Proposition 3.63. Let X be a compact Hausdorff space. The assignment

$$h_X: X \to \sigma(C(X))$$

$$x \mapsto (C(X) \ni f \mapsto (h_X(x))(f) := f(x))$$
(3.64)

is a well-defined homeomorphism.

Proof. See Theorem 1.16 of [13]. and Example 2 after Theorem 1.30 in [13].

Remark 3.65. If we had only assumed that X was compact and not Hausdorff, then h_X would not be one-to-one. To see this, let $x, x' \in X$ be two distinct points that cannot be separated by open sets. For any continuous function $\varphi : X \to \mathbb{C}$, the value of φ on these two points is forced to be the same, $\varphi(x) = \varphi(x')$, i.e. $h_X(x) = h_X(x')$.

Proposition 3.66. The assignment

$$\mathbf{cHaus}_0 \xrightarrow{h} \mathbf{cHaus}_1$$

$$X \mapsto \left(X \xrightarrow{h_X} \sigma(C(X)) \right)$$
(3.67)

from Proposition 3.63 defines a natural isomorphism $h: id_{\mathbf{cHaus}} \Rightarrow \sigma \circ C$.

Proof. Associated to any morphism $X \xrightarrow{f} Y$ of compact Hausdorff spaces is the diagram

$$X \xrightarrow{h_X} \sigma(C(X))$$

$$f \downarrow \qquad \qquad \downarrow \sigma^{C(f)} .$$

$$Y \xrightarrow{h_Y} \sigma(C(Y))$$

$$(3.68)$$

To see that this diagram commutes, let $x \in X$. Going along the top and right arrows gives $\sigma^{C(f)}(h_X(x)) = h_X(x) \circ C^f$ by definition of σ as a functor (see Proposition 3.43). Going along the left and bottom arrows gives $h_Y(f(x))$. To see that these two elements of $\sigma(C(Y))$, i.e. non-zero characters $C(Y) \to \mathbb{C}$, are equal, let $\varphi : Y \to \mathbb{C}$ be a continuous function on Y. Then

$$(h_{X}(x))(C^{f}(\varphi)) \qquad (h_{Y}(f(x)))(\varphi)$$

$$(3.4) \qquad \qquad (3.64)$$

$$(h_{X}(x))(\varphi \circ f) \qquad \varphi(f(x))$$

$$(3.64) \qquad \qquad (3.69)$$

$$(3.64) \qquad \qquad (\varphi \circ f)(x)$$

which proves that h is a natural transformation. It is a natural *iso* morphism by Proposition 3.63.

3.5 The commutative Gelfand-Naimark equivalence

The following theorem is a categorical phrasing of a theorem due to Gelfand and Naimark (Lemma 1 in [18]). It describes in what sense the functors C and σ are inverses of each other. The natural transformations Γ and h also satisfy a universal property that indicates in what sense they are inverses of each other.

Theorem 3.70. Using the notation from above,

$$\left(\mathbf{cC}^*\text{-}\mathbf{Alg}^{\mathrm{op}} \xrightarrow{\sigma} \mathbf{cHaus}, \mathbf{cHaus} \xrightarrow{C} \mathbf{cC}^*\text{-}\mathbf{Alg}^{\mathrm{op}}, \mathrm{id} \xrightarrow{\Gamma} C \circ \sigma, \sigma \circ C \xrightarrow{h^{-1}} \mathrm{id}\right)$$
(3.71)

is an adjoint equivalence of categories.

Proof. Because of Propositions 3.59 and 3.66, it suffices to check the zig-zag identities. The first one is given by

$$\mathbf{cHaus} \xrightarrow{C} \mathbf{cC^*-Alg^{op}} \xrightarrow{\mathrm{id}_{\mathbf{cC^*-Alg^{op}}}} \mathbf{c}$$

$$\mathbf{cHaus} \xrightarrow{C} \mathbf{cC^*-Alg^{op}} \xrightarrow{\sigma} \mathbf{cHaus} \xrightarrow{C} \mathbf{cC^*-Alg^{op}} = \mathbf{cHaus} \xrightarrow{\mathrm{id}_{C}} \mathbf{cC^*-Alg^{op}}$$

$$\downarrow \mathsf{id}_{C} \mathsf{id}_$$

which translates to commutativity of the diagram¹⁴

$$C\left(\sigma(C(X))\right)$$

$$C(h_X^{-1}) \nearrow \Gamma_{C(X)}$$

$$C(X) \xleftarrow{\operatorname{id}_{C(X)}} C(X)$$

$$(3.73)$$

for every compact Hausdorff space X. Because all morphisms here are invertible, it is equivalent to show that the diagram

$$C\left(\sigma(C(X))\right)$$

$$C(h_X) \qquad \Gamma_{C(X)}$$

$$C(X) \stackrel{\text{id}_{C(X)}}{\longleftarrow} C(X)$$

$$(3.74)$$

commutes, which would provide a sense in which h is the left inverse of Γ . Thus, let $\varphi \in C(X)$. Applying the composition along the top two arrows to this element gives

$$C^{h_X}(\Gamma_{C(X)}(\varphi)) = \Gamma_{C(X)}(\varphi) \circ h_X \tag{3.75}$$

by Proposition 3.3. This is a map $X \to \sigma(C(X)) \to \mathbb{C}$ and is therefore determined pointwise so let $x \in X$ and apply this map to it. The result is

$$\left(\Gamma_{C(X)}(\varphi)\right)\left(h_X(x)\right) \stackrel{\text{(3.56)}}{=} h_X(x)\left(\varphi\right) \stackrel{\text{(3.64)}}{=} \varphi(x),\tag{3.76}$$

which proves the first zig-zag identity. The other zig-zag identity

$$\mathbf{cC^*-Alg} \xrightarrow{\mathrm{id}_{\mathbf{cC^*-Alg}}} \mathbf{cHaus^{\mathrm{op}}} \xrightarrow{-C \to \mathbf{cC^*-Alg}} \mathbf{cHaus^{\mathrm{op}}} = \mathbf{cC^*-Alg} \xrightarrow{\sigma} \mathbf{cHaus^{\mathrm{op}}}$$

$$\downarrow^{\mathrm{id}_{\sigma}} \qquad \downarrow^{\mathrm{id}_{\sigma}} \qquad \downarrow^{\mathrm{id}_{\sigma}$$

follows by a similar calculation. For a commutative C^* -algebra \mathcal{A} , this identity says that the diagram

$$\sigma\left(C(\sigma(\mathcal{A}))\right) \\
\sigma(\Gamma_{\mathcal{A}}) \qquad h_{\sigma(\mathcal{A})} \\
\sigma(\mathcal{A}) \stackrel{h_{\sigma(\mathcal{A})}}{\longleftarrow} \sigma(\mathcal{A}) \tag{3.78}$$

commutes, which provides a sense in which h is the right inverse of Γ .

The upshot of the Gelfand-Naimark Theorem that we emphasize here is that it provides one with the perspective that the study of topology (compact Hausdorff spaces) and their continuous functions is equivalent to the study of *commutative* C^* -algebras and their *-homomorphisms.

¹⁴The arrows are in the direction required by contravariance of the functors.

Therefore, non-commutative C^* -algebras can be interpreted as non-commutative topology though a more satisfying relationship to topological concepts is still an area of active research [21]. Furthermore, the *-homomorphisms between these non-commutative spaces can be interpreted as deterministic processes. In the next section, we will describe how to fit in stochastic maps (non-deterministic processes) and motivate non-commutative probability theory.

4 Abstract probability theory

The table below collects several of the categories that have been used along with a few new ones.

Category name	Objects	Morphisms	
FinSetFun	finite sets	functions	
FinSetStoch	finite sets	stochastic maps	
cHaus	compact Hausdorff spaces	continuous functions	
cHausStoch	compact Hausdorff spaces	(continuous) stochastic maps	
$\mathrm{fdcC}^* ext{-}\mathrm{Alg}$	finite-dimensional commutative C^* -algebras	*-homomorphisms	
fdcC*-AlgPos	finite-dimensional commutative C^* -algebras	positive maps	
$\mathrm{cC}^* ext{-Alg}$	commutative C^* -algebras	*-homomorphisms	
cC*-AlgPos	commutative C^* -algebras	positive maps	

In this section, we define the category **cHausStoch** and generalize the (commutative) Gelfand-Naimark Theorem to prove an adjoint equivalence of categories

cHausStoch^{op}
$$\xrightarrow{C}$$
 cC*-AlgPos. (4.1)

During this procedure, we provide an explicit formula for the composition in **cHausStoch** in Proposition 4.79 by showing that a stochastic map induces a canonical Markov kernel in Lemma 4.76. This formula was shown to be valid in the category of stochastic maps on Polish spaces in [19] but seems to have not been verified in the case of compact Hausdorff spaces. One can nevertheless define the composition by passing to the algebra of continuous functions and using the Riesz-Markov-Kakutani Representation Theorem [12] or states [15] though these constructions are a bit formal. In addition, we construct a stochastic version of the Gelfand spectrum functor that does not use the Gelfand transform nor the Riesz-Markov-Kakutani Theorem in Theorem 4.145. Instead, Choquet theory [35] is used to send states onto probability measures on the spectrum. Several of the key ingredients in these constructions are standard [23], [9], [14], [16] though we illustrate how these results fit into a broader context by showing how all of these categories fit into the cube from (1.1) in the introduction.

Sections 4.1 and 4.2 review some measure-theoretic preliminaries. The main results begin in Section 4.3 with the construction of the category **cHausStoch** and end in Section 4.6 with the stochastic Gelfand-Naimark Theorem. The reader with the analytic background may skip ahead directly to these sections. We discuss the relationship between our work and that of others in more detail in Section 5.2.

4.1 Compact Hausdorff spaces and complex measures

Given a measurable space (Y, \mathcal{M}_Y) , where Y is a set and \mathcal{M}_Y is a set of measurable subsets, known as a σ -algebra, one can define the notion of a positive measure and a complex measure. Our measurable spaces will come from compact Hausdorff topological spaces equipped with the Borel subsets for their σ -algebra. As a result, with the additional assumption of finite total measure, the set of positive measures will be a subset of the set of complex measures. We review the definitions, which will be important in specifying a particular topology on the set of all such measures (Proposition 4.29), following Chapters 1 and 6 of [38]. We will obtain from this procedure, a topology on the set of all probability measures on a compact Hausdorff space. For technical reasons, we will then look at a particular subset of all measures satisfying a condition compatible with respect to the topology of the underlying space—such measures are called regular measures (also Radon measures). We will explain the technicalities and why they are needed in remarks. Henceforth, all topological spaces will be assumed compact and Hausdorff and the set of measurable subsets will always be taken to be the set of Borel subsets. A discussion of the pros and cons of using Baire sets is deferred to Section 5.3. Most of this background material does not require the measurable space to be compact Hausdorff though we phrase everything in this setting since all our applications are for such spaces.

Definition 4.2. Let Y be a compact Hausdorff space with associated Borel measurable subsets \mathcal{M}_Y . A <u>complex measure</u> on (Y, \mathcal{M}_Y) is a function $\mu : \mathcal{M}_Y \to \mathbb{C}$ that is <u>countably additive</u> in the sense that for every measurable set $E \in \mathcal{M}_Y$,

$$\mu(E) = \sum_{i \in I} \mu(E_i) \tag{4.3}$$

for every at most countable collection $\{E_i\}_{i\in I\subseteq\mathbb{N}}$ of measurable sets satisfying¹⁵

$$E_i \cap E_j = \varnothing \tag{4.4}$$

whenever $i \neq j$ and

$$\bigcup_{i \in I} E_i = E. \tag{4.5}$$

Such a collection $\{E_i\}_{i\in I}$ is called a <u>measurable partition</u> of E. A <u>positive measure</u> is a complex measure whose value on all measurable sets is non-negative. A <u>probability measure</u> is a positive measure satisfying $\mu(Y) = 1$. The set of all complex measures on Y will be denoted by $\mathbb{C}\text{Meas}(Y)$.

Occasionally, we may write Y instead of (Y, \mathcal{M}_Y) since our set of measurable sets will always be Borel. Note that every positive measure can be used to define an integral using simple functions (see Chapter 1 of [38]). Since the construction is used frequently for positive measurable functions, we state it here while also setting some notation.

Definition 4.6. Let Y be a compact Hausdorff space and let $E \in \mathcal{M}_Y$. The function $\chi_E : Y \to \mathbb{R}$ defined by

$$Y \ni y \mapsto \chi_E(y) := \begin{cases} 1 & \text{if } y \in E \\ 0 & \text{otherwise} \end{cases}$$
 (4.7)

 $^{^{15}\}varnothing$ denotes the empty set.

is called the <u>characteristic function</u> on E. A <u>simple function</u> is a function $s: Y \to \mathbb{R}_{\geq 0}$ of the form

$$s = \sum_{i}^{\text{finite}} s_i \chi_{E_i} \tag{4.8}$$

for some finite set of non-negative numbers $\{s_i\}$ and measurable subsets $\{E_i\}$. Let μ be a positive measure on Y. For a measurable function $f: Y \to \mathbb{R}_{\geq 0}$, the <u>integral of f with respect to μ </u> is the number

$$\int_{Y} f \, d\mu := \sup_{0 \le s \le f} \left\{ \sum_{i} s_{i} \mu(E_{i}) : s = \sum_{i} s_{i} \chi_{E_{i}} \text{ is simple} \right\}. \tag{4.9}$$

The integral of a complex measurable function can be defined by splitting the function up into a sum of its real and imaginary parts each of which can be decomposed into a sum of positive and negative parts. Similarly, every complex measure can be uniquely decomposed via Jordan decomposition into an appropriate sum and difference of positive measures so as to define the integral of a positive function with respect to a complex measure and then finally the integral of any complex measurable function with respect to a complex measure (see Section 6.18 of [38]—we will review a necessary fact to make sense of this definition in Theorem 4.23 below).

The following Lemma is useful for several of the facts that will follow. It is unfortunately long, but is used so often that we felt it was important to state it in its full form.

Lemma 4.10. Let Z be a compact topological space and let $\varphi: Z \to \mathbb{R}_{\geq 0}$ be a bounded measurable function. Then there exists a sequence $s: \mathbb{N} \to \mathbb{R}^Z$ of simple functions satisfying

$$0 \le s_1 \le s_2 \le s_3 \le \dots \le \varphi \qquad \& \qquad \lim_{n \to \infty} s_n(z) = \varphi(z) \quad \forall \ z \in Z. \tag{4.11}$$

Furthermore, the coefficients $\{s_{n,i_n}\}$ and measurable sets $\{E_{n,i_n}\}$ in the expression

$$s_n = \sum_{i_n} s_{n,i_n} \chi_{E_{n,i_n}} \tag{4.12}$$

can be chosen so that $i_n \in \{0, 1, \dots, 2^n - 1\},\$

$$E_{n,i_n} = E_{n+1,2i_n} \cup E_{n+1,2i_n+1} \qquad \& \qquad E_{n+1,2i_n} \cap E_{n+1,2i_n+1} = \varnothing,$$
 (4.13)

and

$$s_{n,i_n} \le \min \left\{ s_{n+1,2i_n}, s_{n+1,2i_n+1} \right\} \tag{4.14}$$

for all $i_n \in \{0, 1, \dots, 2^n - 1\}$ and $n \in \mathbb{N}$.

Proof. Since φ is bounded, there exists an $M \geq 0$ such that $\varphi(Z) \subseteq [0, M]$. For each $n \in \mathbb{N}$, set

$$F_{n,i_n} := \begin{cases} \left[\frac{i_n M}{2^n}, \frac{(i_n+1)M}{2^n} \right) & \text{for } i_n \in [0, 2^n - 1) \cap \mathbb{Z} \\ \left[\frac{i_n M}{2^n}, \frac{(i_n+1)M}{2^n} \right] & \text{for } i_n = 2^n - 1, \end{cases}$$
(4.15)

where $i_n \in \{0, 1, ..., 2^n - 1\}$. Since φ is Borel measurable, the preimages $E_{n,i_n} := \varphi^{-1}(F_{n,i_n})$ form a measurable partition of Z for each n. Set $s_n : Z \to \mathbb{R}_{\geq 0}$ to be the simple function

$$s_n := \sum_{i_n=0}^{2^n-1} \frac{i_n M}{2^n} \chi_{E_{n,i_n}}.$$
 (4.16)




Figure 2: A non-negative bounded measurable function is an increasing sequence of simple functions satisfying several convenient properties as described in Lemma 4.10.

An example of such a simple function together with the next function in the sequence is depicted in Figure 2. The sequence $\mathbb{N} \ni n \mapsto s_n$ of simple functions satisfies the required properties because for every n, the pointwise difference between s_n and φ is at most $\frac{M}{2^n}$. Furthermore, $s_n \leq s_{n+1}$ for each $n \in \mathbb{N}$ because $\{F_{n+1,i_{n+1}}\}$ is a refinement of $\{F_{n,i_n}\}$ (this was the reason for the choice of 2^n as opposed to merely n in the above formulas). The other properties are immediate from the construction.

Theorem 4.17 (Monotone Convergence Theorem). Let Y be a compact Hausdorff space and let $\varphi : \mathbb{N} \to [0,\infty]^Y$ be an increasing sequence of Borel measurable functions, i.e. $\varphi_{n+1}(y) \ge \varphi_n(y)$ for all $y \in Y$ and $n \in \mathbb{N}$. Then $\lim \varphi$ is Borel measurable and

$$\lim_{n \to \infty} \int_{Y} \varphi_n \ d\mu = \int_{Y} \lim \varphi \ d\mu \tag{4.18}$$

for every Borel measure μ on Y.

Proof. See Theorem 1.26 in [38].

Definition 4.19. Let Y be a compact Hausdorff space. Given a complex measure $\mu \in \mathbb{C}\mathrm{Meas}(Y)$, the total variation of μ is the function $|\mu| : \mathcal{M}_Y \to \mathbb{R}$ defined by

$$\mathcal{M}_Y \ni E \stackrel{|\mu|}{\longmapsto} \sup \left\{ \sum_{i \in I} \left| \mu(E_i) \right| \right\},$$
 (4.20)

where the supremum is over all measurable partitions $\{E_i\}_{i\in I}$ of E.

Proposition 4.21. Let Y be a compact Hausdorff space and $\mu \in \mathbb{C}\text{Meas}(Y)$. Then, the total variation of μ is a positive measure on Y. Furthermore, the set of all complex measures on Y has the structure of a normed complex vector space with norm given by

$$\mathbb{C}\mathrm{Meas}(Y) \ni \mu \mapsto |\mu|(Y). \tag{4.22}$$

Proof. See Chapter 6 of [38].

Theorem 4.23 (Polar decomposition for complex measures). Let Y be a compact Hausdorff space and $\mu \in \mathbb{C}\mathrm{Meas}(Y)$. Then there exists a measurable function $h: Y \to \mathbb{C}$ such that |h(y)| = 1 for all $y \in Y$ and

$$\mu(E) = \int_{E} h \ d|\mu| \qquad \forall \ E \in \mathcal{M}_{Y}. \tag{4.24}$$

This is often written in shorthand form as $d\mu = h d|\mu|$.

Proof. See Theorem 6.12 in [38].

This is an important consequence of the Lebesgue-Radon-Nikodym Theorem. It allows one to define the integral of complex-valued measurable functions with respect to complex measures, namely using the notation of Theorem 4.23, the integral of a complex-valued measurable function $\varphi: Y \to \mathbb{C}$ with respect to μ is

$$\int_{Y} \varphi \, d\mu := \int_{Y} \varphi h \, d|\mu|. \tag{4.25}$$

This definition satisfies the usual properties of the integral (see Section 6.18 in [38]). This theorem can also be used to prove the following result, which we will need.

Lemma 4.26. Let Y be a compact Hausdorff space and $\mu \in \mathbb{C}Meas(Y)$. Then

$$\left| \int_{Y} \varphi \ d\mu \right| \le \int_{Y} |\varphi| \ d|\mu| \tag{4.27}$$

for all $\varphi \in C(Y)$.

Proof. Using a polar decomposition $d\mu = h d|\mu|$,

$$\left| \int_{Y} \varphi \ d\mu \right| = \left| \int_{Y} \varphi h \ d|\mu| \right| \le \int_{Y} |\varphi| |h| \ d|\mu| = \int_{Y} |\varphi| \ d|\mu| \tag{4.28}$$

for all $\varphi \in C(Y)$.

Being a normed space, $\mathbb{C}Meas(Y)$ has a natural topology on it. However, as for many infinite-dimensional vector spaces, too few sequences have convergent subsequences with respect to this topology. There are several ways to implement topologies on $\mathbb{C}Meas(Y)$. Rather than using all measurable functions, we will use continuous functions to define a collection of seminorms (recall the discussion in Section 3.2).

Proposition 4.29. Let Y be a compact Hausdorff space. For every continuous function $\varphi : Y \to \mathbb{C}$, set $\|\cdot\|_{\varphi} : \mathbb{C}\mathrm{Meas}(Y) \to \mathbb{R}$ to be the function given by

$$\mathbb{C}\mathrm{Meas}(Y) \ni \mu \mapsto \int_{Y} |\varphi| \ d|\mu|. \tag{4.30}$$

Then $\|\cdot\|_{\varphi}$ is a seminorm on $\mathbb{C}\mathrm{Meas}(Y)$.

Proof. This follows immediately from the definition of the integral and the fact that the total variation $|\cdot|$ defines a norm on $\mathbb{C}\text{Meas}(Y)$. Note that the function $|\cdot|_{\varphi}$ makes sense even when φ is just a measurable function.

- i. $\|\mu\|_{\varphi} = \int_{Y} |\varphi| \, d|\mu| \ge 0$ for all complex measures $\mu \in \mathbb{C}\text{Meas}(Y)$ follows immediately from the definition of the integral of a positive function with respect to a positive measure.
- ii. Fix $\lambda \in \mathbb{C}$ and $\mu \in \mathbb{C}\mathrm{Meas}(Y)$. By Lemma 4.10, there exists a sequence $s : \mathbb{N} \to \mathbb{R}^Y$ of simple functions on Y of the form

$$s_n := \sum_{i_n}^{\text{finite}} s_{n,i_n} \chi_{E_{n,i_n}}, \tag{4.31}$$

satisfying

$$0 \le s_1 \le s_2 \le s_3 \le \dots \le |\varphi| \tag{4.32}$$

and

$$\lim_{n \to \infty} s_n(y) = |\varphi(y)| \qquad \forall \ y \in Y. \tag{4.33}$$

By the definition of the integral

$$\lim_{n \to \infty} \int_{V} s_n \ d|\lambda \mu| = \int_{V} |\varphi| \ d|\lambda \mu| = ||\lambda \mu||_{\varphi}. \tag{4.34}$$

Meanwhile, by definition of simple functions, the left-hand-side of this expression equals

$$\lim_{n \to \infty} \int_{Y} s_{n} \ d|\lambda \mu| = \lim_{n \to \infty} \sum_{i_{n}=1}^{\text{finite}} |\lambda \mu| (E_{n,i_{n}}) s_{n,i_{n}}$$

$$= \lim_{n \to \infty} \sum_{i_{n}=1}^{\text{finite}} |\lambda| |\mu| (E_{n,i_{n}}) s_{n,i_{n}}$$

$$= |\lambda| \lim_{n \to \infty} \sum_{i_{n}=1}^{\text{finite}} |\mu| (E_{n,i_{n}}) s_{n,i_{n}}$$

$$= |\lambda| \lim_{n \to \infty} \int_{Y} s_{n} \ d|\mu|$$

$$= |\lambda| \|\mu\|_{\varphi}.$$

$$(4.35)$$

Hence, $\|\lambda\mu\|_{\varphi} = |\lambda| \|\mu\|_{\varphi}$ as needed.

iii. Let $\mu, \nu \in \mathbb{C}\text{Meas}(Y)$ and let s be a sequence as in the previous part. A similar argument as above shows that

$$\|\mu + \nu\|_{\varphi} = \int_{Y} |\varphi| \ d|\mu + \nu| \le \int_{Y} |\varphi| \ d|\mu| + \int_{Y} |\varphi| \ d|\nu| = \|\mu\|_{\varphi} + \|\nu\|_{\varphi}. \tag{4.36}$$

All of the conditions of a seminorm have therefore been checked.

Henceforth, we equip $\mathbb{C}\text{Meas}(Y)$ with the topology generated by the family of seminorms $\{\|\cdot\|_{\varphi}\}_{\varphi\in C(Y)}$. This topology is known as the <u>vague topology</u> on $\mathbb{C}\text{Meas}(Y)$. It is helpful to know when nets converge with respect to this topology so that we may use nets to prove whether or not certain functions are continuous.

Proposition 4.37. Let Y be a compact Hausdorff space. A net $\mu : \Theta \to \mathbb{C}\text{Meas}(Y)$ converges to $\lim \mu$ with respect to the vague topology if and only if

$$\lim_{\theta \in \Theta} \int_{Y} \varphi \ d\mu_{\theta} = \int_{Y} \varphi \ d \lim \mu \tag{4.38}$$

for all $\varphi \in C(Y)$.

Proof.

 (\Rightarrow) Suppose that $\mu:\Theta\to\mathbb{C}\mathrm{Meas}(Y)$ converges to $\lim \mu$. Fix $\epsilon>0$. By assumption, for any continuous function $\varphi\in C(Y)$, there exists a $\theta_0\in\Theta$ such that

$$\|\lim \mu - \mu_{\theta}\|_{\varphi} < \epsilon \qquad \forall \ \theta \ge \theta_0. \tag{4.39}$$

Then, by Lemma 4.26,

$$\left| \int_{Y} \varphi \ d \lim \mu - \int_{Y} \varphi \ d\mu_{\theta} \right| \leq \left| \int_{Y} \varphi \ d(\lim \mu - \mu_{\theta}) \right|$$

$$\leq \int_{Y} |\varphi| \ d|\lim \mu - \mu_{\theta}|$$

$$= \|\lim \mu - \mu_{\theta}\|_{\varphi}$$

$$< \epsilon \qquad \forall \ \theta \geq \theta_{0}.$$

$$(4.40)$$

(\Leftarrow) Suppose that $\lim_{\theta \in \Theta} \int_Y \varphi \ d\mu_\theta = \int_Y \varphi \ d\lim \mu$ for all $\varphi \in C(Y)$. Fix $\epsilon > 0$, $n \in \mathbb{N}$, and $\alpha : \overline{n} \to C(Y)$ whose value at $i \in \overline{n}$ will be denoted by α_i (we will show that μ is eventually in a neighborhood base of $\lim \mu$). By assumption, for each $i \in \overline{n}$, there exist $\theta_i \in \Theta$ such that

$$\left| \int_{Y} \varphi \ d \lim \mu - \int_{Y} \varphi \ d\mu_{\theta} \right| < \epsilon \qquad \forall \ \theta \ge \theta_{i}. \tag{4.41}$$

Set

$$\theta_0 := \max_{i \in \overline{\eta}} \{\theta_i\}. \tag{4.42}$$

Then, $\mu_{\theta} \in B_{\epsilon}^{\alpha}(\lim \mu)$ for all $\theta \geq \theta_0$. Hence μ converges to $\lim \mu$ with respect to the vague topology.

4.2 Probability measures on compact Hausdorff spaces

Naively, one might use the set of *all* probability measures on Y and define suitably continuous stochastic maps between compact Hausdorff spaces using this definition. However, in order to obtain an equivalence of categories between such spaces and the category of commutative C^* -algebras and positive maps, an additional restriction needs to be made on the set of probability measures to ensure that the equivalence holds. This restriction is solely due to the fact that not all compact Hausdorff spaces are separable, or equivalently, metrizable (for instance, see counterexamples 24, 43, 107, and 111 from [45]). The measures we must restrict our attention to are regular (Radon) measures. Since we will mainly be concerned with probability measures, we will immediately assume throughout that all measures are finite.

Definition 4.43. Let Y be a compact Hausdorff space. A positive Borel measure $\mu: \mathcal{M}_Y \to \mathbb{R}_{\geq 0}$ is regular iff

$$\mu(E) = \inf \{ \mu(U) : E \subseteq U \text{ with } U \text{ open in } Y \}$$
(4.44)

and

$$\mu(E) = \sup \{ \mu(K) : K \subseteq E \text{ with } K \text{ compact in } Y \}$$
(4.45)

for all $E \in \mathcal{M}_Y$. A complex Borel measure $\mu : \mathcal{M}_Y \to \mathbb{C}$ is <u>regular</u> iff the positive and negative parts of the real and complex parts of μ are all regular.

Because being regular is a property of a measure, the set of regular measures on Y is a subset $\mathbb{C}\text{Reg}(Y) \subseteq \mathbb{C}\text{Meas}(Y)$ of all complex measures. Since $\mathbb{C}\text{Meas}(Y)$ has a normed vector space structure, one could ask if $\mathbb{C}\text{Reg}(Y)$ inherits the properties of this structure as well.

Theorem 4.46. $\mathbb{C}\text{Reg}(Y)$ with the total variation norm is a Banach space for all compact Hausdorff spaces Y.

Proof. See Exercise 3 in Chapter 6 of [38].

Notation 4.47. Because $\mathbb{C}\text{Reg}(Y) \subseteq \mathbb{C}\text{Meas}(Y)$, it also inherits the vague topology described after Proposition 4.29. Let ProbMeas(Y) denote the subset of $\mathbb{C}\text{Reg}(Y)$ of regular measures that are also probability measures. Equip ProbMeas(Y) with the subspace topology coming from the vague topology on $\mathbb{C}\text{Meas}(Y)$.

Notice that when Y is a finite set with the discrete topology, ProbMeas(Y) = Pr(Y).

Remark 4.48. When Y is just a locally compact Hausdorff space, one must use Radon probability measures instead (see Theorem 7.2 in [14]). When Y is a compact Hausdorff space, it is a fact that every Radon probability measure is regular (see Corollary 7.6 in [14]). When Y is, in addition, second countable, every Borel measure on Y is regular (see Theorem 7.8 in [14]). Hence, many of the complications that follow can be ignored provided that Y is second countable (and therefore separable). In fact, many treatments of probability theory on topological spaces assume the spaces are Polish. We do not need this requirement, which is one of the interesting features of our findings.

We mention several facts regarding regular measures on compact Hausdorff spaces that will be needed for many of our proofs.

Lemma 4.49 (Urysohn's Lemma). Let Y be a compact Hausdorff space and let $K \subseteq U \subseteq Y$ with K closed (and hence compact) and U open. Then there exists a continuous function $\varphi \in C(Y, [0, 1])$ such that $\varphi = 1$ on K and $\varphi = 0$ outside a compact subset of U.

Proof. See Theorem 4.32 in [14].

Definition 4.50. Let Y be a topological space. A function $\varphi: Y \to \mathbb{R}$ is <u>lower semi-continuous</u> iff $f^{-1}((a,\infty))$ is open for all $a \in \mathbb{R}$ and is <u>upper semi-continuous</u> iff $f^{-1}((-\infty,a))$ is open for all $a \in \mathbb{R}$.

It is immediate from the definition that semi-continuous functions are measurable. This is because sets of the form (a, ∞) also generate the Borel σ -algebra on \mathbb{R} and because a function is measurable if and only if the preimages of generating sets are measurable.

Proposition 4.51. Let Y be a topological space.

- (a) If $U \subseteq Y$ is open, then χ_U , the characteristic function on U, is lower semi-continuous.
- (b) If $K \subseteq Y$ is closed, then χ_K is upper semi-continuous.
- (c) If Y is compact and Hausdorff and $\chi: Y \to \mathbb{R}_{\geq 0}$ is lower semi-continuous, then

$$\chi(y) = \sup_{\varphi \in C(Y)} \left\{ \varphi(y) : 0 \le \varphi \le \chi \right\} \qquad \forall \ y \in Y.$$
 (4.52)

(d) If Y is compact and Hausdorff and $\chi: Y \to \mathbb{R}_{\geq 0}$ is upper semi-continuous, then

$$\chi(y) = \inf_{\varphi \in C(Y)} \left\{ \varphi(y) : 0 \le \chi \le \varphi \right\} \qquad \forall \ y \in Y.$$
 (4.53)

Proof. See Proposition 7.11 in [14] (though the reader may also check that this follows from the definitions and Urysohn's Lemma).

Corollary 4.54. Let μ be a positive regular measure on a compact Hausdorff space Y and let $\chi: Y \to \mathbb{R}_{>0}$ be a lower semi-continuous function on Y. Then

$$\int_{Y} \chi \ d\mu = \sup_{\varphi \in C(Y)} \left\{ \int_{Y} \varphi \ d\mu : 0 \le \varphi \le \chi \right\}. \tag{4.55}$$

Proof. See Corollary 7.13 in [14].

Remark 4.56. Corollary 4.54 follows from Lusin's theorem, which is a more quantitative version of Urysohn's Lemma for compact Hausdorff spaces *equipped with* a (finite) regular measure [14].

All regular probability measures are regular positive measures, and therefore all of these results apply to regular probability measures on a compact Hausdorff space.

4.3 Continuous stochastic maps

We are finally ready to define the category **cHausStoch**, the generalization of the category **FinSetStoch** to compact Hausdorff spaces and (continuous) stochastic maps. In the process of doing so, we prove several facts of independent interest (see Lemma 4.64 and 4.76 for instance).

Definition 4.57. Let X and Y be two compact Hausdorff spaces. A (continuous) <u>stochastic map</u> from X to Y is a continuous function $\mu: X \to \operatorname{ProbMeas}(Y)$ with respect to the vague topology and is denoted by $\mu: X \leadsto Y$. The value of μ at $x \in X$ is denoted by μ_x and is a regular probability measure on Y.

Example 4.58. Let $f: X \to Y$ be a continuous function of compact Hausdorff spaces. Then

$$X \xrightarrow{\delta_f} \text{ProbMeas}(Y)$$

$$x \mapsto \delta_{f(x)}, \tag{4.59}$$

where $\delta_{f(x)}$ is the measure defined by

$$\mathcal{M}_Y \ni E \mapsto \delta_{f(x)}(E) := \begin{cases} 1 & \text{if } f(x) \in E \\ 0 & \text{if } f(x) \notin E, \end{cases}$$

$$(4.60)$$

is a stochastic map from X to Y. $\delta_{f(x)}$ is called the Dirac delta measure on Y concentrated at the point f(x). The fact that $\delta_{f(x)}$ is a regular probability measure follows immediately from the definition. Before proving that this is a stochastic map, notice that for any continuous function $\varphi \in C(Y)$ and for any $x \in X$,

$$\int_{Y} \varphi \ d\delta_{f(x)} = \varphi(f(x)). \tag{4.61}$$

To see that $X \ni x \mapsto \delta_{f(x)}$ is continuous, let $x : \Theta \to X$ be a net converging to $\lim x \in X$. Then, for any continuous function $\varphi \in C(Y)$,

$$\lim_{\theta \in \Theta} \int_{Y} \varphi \ d\delta_{f(x_{\theta})} = \lim_{\theta \in \Theta} \varphi \big(f(x_{\theta}) \big) = \varphi \big(f(\lim x) \big) = \int_{Y} \varphi \ d\delta_{f(\lim x)}. \tag{4.62}$$

Since φ was arbitrary, this shows that $\lim_{\theta \in \Theta} \delta_{f(x_{\theta})} = \delta_{f(\lim x)}$ in the vague topology.

In particular, the identity function from a space to itself induces a stochastic map known as the <u>identity stochastic map</u>. Why it is called the identity will be justified after the definition of composition of stochastic maps is provided.

The composition of stochastic maps is quite subtle for compact Hausdorff spaces due to the possibility that the underlying spaces need not be metrizable. Before going into the details, we isolate the issue that occurs by what we expect the composition to be heuristically based on our understanding of the finite set case. In this regard, let X, Y, and Z be compact Hausdorff spaces and let $\mu: X \leadsto Y$ and $\nu: Y \leadsto Z$ be two stochastic maps. The composition of μ followed by ν should be some stochastic map $X \leadsto Z$ which when applied to a Borel set $E \subseteq Z$ and a point $x \in X$ should give the probability of x evolving through the intermediate possibilities in Y that end up at the "window" E as sketched in Figure 3.




Figure 3: $x \in X$ evolves to the probability measure μ_x and all the points in Y that have neighborhoods with non-vanishing measure themselves get sent to probability measures on Z. To calculate the overlap with a measurable set $E \subseteq Z$, the measures are evaluated on E and the results are integrated over all of Y. Only ν_y and $\nu_{y'}$ for two points $y, y' \in Y$ have been sketched on the left for ease of visualization.

In other words, we sum up $\nu_y(E)$ over all y using the measure μ_x on Y to obtain

$$\int_{Y} \nu_y(E) \ d\mu_x(y). \tag{4.63}$$

However, this formula assumes that the function $Y \ni y \mapsto \nu_y(E)$ is measurable for all Borel sets $E \in \mathcal{M}_Z$. The validity of this formula is well known if all the spaces in question are Polish [19] but does not seem to be known for arbitrary compact Hausdorff spaces. We will prove this result soon, but first we prove a crucial result that is interesting in its own right.

Lemma 4.64. Let Y be a compact Hausdorff space and let $E \in \mathcal{M}_Y$ be a Borel set. Then the evaluation function

$$\operatorname{ProbMeas}(Y) \xrightarrow{\operatorname{ev}_E} \mathbb{R}$$

$$\mu \mapsto \mu(E) \tag{4.65}$$

is Borel measurable with respect to the vague topology on ProbMeas(Y). In fact, ev_E is lower semi-continuous when E is open and upper semi-continuous when E is closed.

To prove this lemma, we recall a useful theorem, which we learned through Sengupta's lucid notes [43].

Theorem 4.66 (Dynkin's π - λ Theorem). Let X be a set and let P and L be collections of subsets of X satisfying the following conditions.

- i. P is closed under finite intersections (such a collection is called a $\underline{\pi}$ -system).
- ii. L contains the empty set, is closed under taking complements (in X), and is closed under countable unions of pairwise disjoint elements (such a collection is called a λ -system).

iii. $P \subseteq L$.

Then the σ -algebra generated by P is contained in L.

Proof. See Theorem 3.2 in [6] or the notes [43].

We are indebted to Iddo Ben-Ari for the crucial insight that Dynkin's π - λ theorem can be used to give a nice proof of Lemma 5.8, which lead to the proof of Lemma 4.64.

Proof of Lemma 4.64. The proof will be broken up into several steps. First, it will be shown that the collection, P, of all open subsets of Y forms a π -system. Then it will be shown that the set

$$L := \left\{ E \subseteq Y : \operatorname{ProbMeas}(Y) \xrightarrow{\operatorname{ev}_E} [0, 1] \text{ is Borel measurable} \right\}$$
 (4.67)

is a λ -system. Finally, it will be shown that ev_U is lower semi-continuous (and hence Borel measurable), for all open sets $U \subseteq Y$ showing that $P \subseteq L$.

- i. The fact that open sets are closed under finite intersections is part of the definition of a topology.
- ii. For each $\mu \in \text{ProbMeas}(Y)$,

$$\operatorname{ev}_{\varnothing}(\mu) = \mu(\varnothing) = 0 \tag{4.68}$$

so that ev_{\emptyset} is constant and hence measurable. For $E \in L$,

$$\operatorname{ev}_{E^c}(\mu) = \mu(E^c) = \mu(Y) - \mu(E) = (1 - \operatorname{ev}_E)(\mu)$$
 (4.69)

is the difference of two measurable functions with respect to μ and is therefore measurable with respect to μ . For $\{E_n\}$ a countable collection of pairwise disjoint measurable sets in L with $E := \bigcup_n E_n$,

$$\operatorname{ev}_{E}(\mu) = \mu\left(\bigcup_{n=1}^{\infty} E_{n}\right) = \sum_{n=1}^{\infty} \mu(E_{n}) = \left(\sum_{n=1}^{\infty} \operatorname{ev}_{E_{n}}\right)(\mu)$$
(4.70)

by additivity of the measure μ . Hence, since this is a bounded countable sum of measurable functions, the sum is measurable.

iii. Let $U \subseteq Y$ be open and non-empty (if U is empty, the result was already proved in part ii.). By Proposition 4.51, χ_U is lower semi-continuous. Hence, for any $a \in \mathbb{R}$, the preimage of (a, ∞) under ev_U is

$$\operatorname{ev}_{U}^{-1}((a, \infty)) = \left\{ \mu \in \operatorname{ProbMeas}(Y) : \mu(U) > a \right\}$$

$$= \left\{ \mu \in \operatorname{ProbMeas}(Y) : \int_{Y} \chi_{U} d\mu > a \right\}$$

$$= \left\{ \mu \in \operatorname{ProbMeas}(Y) : \sup_{\substack{\varphi \in C(Y) \\ 0 \leq \varphi \leq \chi_{U}}} \left\{ \int_{Y} \varphi d\mu \right\} > a \right\} \text{ by Corollary 4.54} \quad (4.71)$$

$$= \bigcup_{\substack{\varphi \in C(Y) \\ 0 < \varphi \leq \chi_{U}}} \left\{ \mu \in \operatorname{ProbMeas}(Y) : \int_{Y} \varphi d\mu > a \right\}.$$

By definition of the vague topology, the set in curly brackets in the last equality is open. Hence, this is a union of open sets and is therefore an open subset of ProbMeas(Y). Therefore, this shows that ev_U is lower semi-continuous for all open sets U in Y. An analogous argument using infima shows that ev_K is upper semi-continuous for all closed sets K in Y.

Thus, since all the conditions of Dynkin's π - λ theorem are satisfied, the σ -algebra generated by P which is the Borel σ -algebra, is contained in L.

Remark 4.72. Although ev_U was shown to be lower semi-continuous for all open sets U, it is in general *not* continuous. For an explicit example, enumerate the rationals in [0,1] by some bijective function $q: \mathbb{N} \to \mathbb{Q} \cap [0,1]$ and set

$$U := \left(\bigcup_{n=1}^{\infty} \left(q_n - \frac{\epsilon}{2^{n+1}}, q_n + \frac{\epsilon}{2^{n+1}}\right)\right) \cap [0, 1]$$

$$(4.73)$$

for some $\epsilon \in (0,1)$. Then, U is open, contains all of the rationals in [0,1], and has Lebesgue measure $\epsilon < 1$ so that it does not contain all of [0,1]. Fix $x_{\infty} \in [0,1] \setminus U$. Since $\mathbb{Q} \cap [0,1]$ is dense in [0,1], there exists a sequence $x : \mathbb{N} \to \mathbb{Q} \cap [0,1]$ converging to x_{∞} in [0,1]. Then, the corresponding sequence of Dirac measures $n \mapsto \delta_{x_n}$ converges to $\delta_{x_{\infty}}$ in the vague topology but

$$\delta_{x_n}(U) = 1$$
 & $\delta_{x_n}([0,1] \setminus U) = 0$ $\forall n \in \mathbb{N}$ (4.74)

while

$$\delta_{x_{\infty}}(U) = 0 \qquad \& \qquad \delta_{x_{\infty}}([0,1] \setminus U) = 1 \tag{4.75}$$

showing that ev_U is not continuous at δ_{x_∞} . In fact, checking that ev_U is lower semi-continuous at δ_{x_∞} directly follows from the definition since $\operatorname{ev}_U(\mu) \geq \operatorname{ev}_U(\delta_{x_\infty}) - \epsilon$ reads $\mu(U) \geq 0 - \epsilon = -\epsilon$, which is true for all $\epsilon > 0$ and probability measures μ . However, one runs into trouble checking if ev_U is upper semi-continuous for the reason illustrated above.

Lemma 4.76. Let $\mu: X \leadsto Y$ be a stochastic map as in Definition 4.57. For every Borel set $E \in \mathcal{M}_Y$, the function

$$X \ni x \mapsto \mu_x(E) \tag{4.77}$$

is Borel measurable. In fact, if E is open (closed), then this function is lower (upper) semi-continuous.

Proof. This follows from Lemma 4.64 and the fact that this function is the composition

$$X \xrightarrow{\mu} \operatorname{ProbMeas}(Y) \xrightarrow{\operatorname{ev}_E} \mathbb{R}$$
 (4.78)

of Borel measurable functions. If E is open (closed), then this function is the composition of a continuous function followed by a lower (upper) semi-continuous function, which is still lower (upper) semi-continuous (this is immediate from the definitions).

Note that this shows, in particular, that a stochastic map gives rise to a Markov kernel. The converse is not true as an arbitrary Markov kernel takes no account of continuity. Hence, although the composition of Markov kernels is a Markov kernel, the requirement of being a stochastic map imposes additional restrictions.

Proposition 4.79. Let X, Y, and Z be compact Hausdorff spaces and let $\mu : X \rightsquigarrow Y$ and $\nu : Y \rightsquigarrow Z$ be two stochastic maps. For each $x \in X$, the assignment

$$\mathcal{M}_Z \ni E \mapsto (\nu \circ \mu)_x(E) := \int_Y \nu_y(E) \ d\mu_x(y),$$
 (4.80)

is a regular probability measure on Z. Furthermore, the assignment

$$X \to \text{ProbMeas}(Z)$$

$$x \mapsto (\nu \circ \mu)_x,$$
(4.81)

is a stochastic map. This stochastic map is called the <u>composition</u> of μ followed by ν and is denoted by $\nu \circ \mu$.

Proof. The proof will be split into three parts to prove the following claims: i. $(\nu \circ \mu)_x$ is a Borel probability measure on Z, ii. it is regular, and iii. the assignment $X \ni x \mapsto (\nu \circ \mu)_x$ is a stochastic map $X \leadsto Z$.

i. First,

$$(\nu \circ \mu)_x(Z) = \int_Y \underbrace{\nu_y(Z)}_{1} d\mu_x(y) = \int_Y d\mu_x(y) = \mu_x(Y) = 1. \tag{4.82}$$

Now, fix a Borel set $E \in \mathcal{M}_Z$. Since $Y \ni y \mapsto \nu_y(E)$ is Borel measurable (by Lemma 4.76) and since $\nu_y(E)$ is non-negative for all Borel sets E in Z, its integral with respect to a positive measure is non-negative. We now check that $(\nu \circ \mu)_x$ is additive on (at most) countable measurable partitions. Let $\{E_i\}_{i\in I}$ be a countable measurable partition of E. Then

$$(\nu \circ \mu)_x(E) = \int_Y \nu_y(E) \, d\mu_x(y) \quad \text{by definition of } (\nu \circ \mu)_x$$

$$= \int_Y \sum_{i \in I} \nu_y(E_i) \, d\mu_x(y) \quad \text{since } \nu_y \text{ is a measure}$$

$$= \sum_{i \in I} \int_Y \nu_y(E_i) \, d\mu_x(y) \quad \text{by the Monotone Convergence Theorem}$$

$$= \sum_{i \in I} (\nu \circ \mu)_x(E_i) \quad \text{by definition of } (\nu \circ \mu)_x,$$

$$(4.83)$$

In the third equality, $\nu_y(E_i)$ is non-negative and measurable for all $y \in Y$ so the partial sums form an increasing sequence of measurable functions. Hence, the Monotone Convergence Theorem allows one to exchange the sum operation with the integral. Therefore, $(\nu \circ \mu)_x$ is a Borel probability measure.

ii. To prove regularity of the measure $(\nu \circ \mu)_x$, fix $E \in \mathcal{M}_Z$ and $\epsilon > 0$. The goal is to find an open set $U \subseteq Z$ and a compact set $K \subseteq Z$ such that $K \subseteq E \subseteq U$ and

$$(\nu \circ \mu)_x(U) - (\nu \circ \mu)_x(E) < \epsilon \qquad \& \qquad (\nu \circ \mu)_x(E) - (\nu \circ \mu)_x(K) < \epsilon. \tag{4.84}$$

Note, that it suffices to show that (by a single application of the triangle inequality)

$$(\nu \circ \mu)_x(U) - (\nu \circ \mu)_x(K) < \epsilon. \tag{4.85}$$

Fix $y \in Y$. Since ν_y is regular, there exist an open set $U_y \subseteq Z$ and a closed set $K_y \subseteq Z$ such that $K_y \subseteq E \subseteq U_y$ and

$$\nu_y(U_y) - \nu_y(K_y) < \frac{\epsilon}{3}. \tag{4.86}$$

Since $Y \ni y' \mapsto \nu_{y'}(U_y)$ is lower semi-continuous and $Y \ni y' \mapsto \nu_{y'}(K_y)$ is upper semi-continuous by Lemma 4.76, there exists an open neighborhood $V_y \subseteq Y$ containing y such that

$$\nu_y(U_y) - \nu_{y'}(U_y) \le \frac{\epsilon}{3} \qquad \& \qquad \nu_{y'}(K_y) - \nu_y(K_y) \le \frac{\epsilon}{3} \qquad \forall \ y' \in V_y. \tag{4.87}$$

The collection of such open and closed sets can be obtained for every $y \in Y$. Hence, $\{V_y\}_{y \in Y}$ forms an open cover of Y. By compactness of Y, there exists a finite collection $\{y_1, \ldots, y_n\}$ of points in Y for which $\{V_{y_1}, \ldots, V_{y_n}\}$ covers Y. Set

$$U := \bigcap_{i=1}^{n} U_{y_i} \qquad \& \qquad K := \bigcup_{i=1}^{n} K_{y_i}. \tag{4.88}$$

Since these intersections and unions are finite, U is open and K is closed in Z. Furthermore, they satisfy $K \subseteq E \subseteq U$. Now, fix $y \in Y$. There exists some $i \in \overline{n}$ such that $y \in V_{y_i}$ so that

$$\begin{aligned} \left| \nu_{y}(U) - \nu_{y}(K) \right| &\leq \left| \nu_{y}(U_{y_{i}}) - \nu_{y}(K_{y_{i}}) \right| & \text{by (4.88) and sub-additivity of } \nu_{y} \\ &= \left| \nu_{y}(U_{y_{i}}) - \nu_{y_{i}}(U_{y_{i}}) + \nu_{y_{i}}(U_{y_{i}}) - \nu_{y_{i}}(K_{y_{i}}) + \nu_{y_{i}}(K_{y_{i}}) - \nu_{y}(K_{y_{i}}) \right| \\ &\leq \left| \nu_{y}(U_{y_{i}}) - \nu_{y_{i}}(U_{y_{i}}) \right| + \left| \nu_{y_{i}}(U_{y_{i}}) - \nu_{y_{i}}(K_{y_{i}}) \right| + \left| \nu_{y_{i}}(K_{y_{i}}) - \nu_{y}(K_{y_{i}}) \right| \\ &< \epsilon & \text{by (4.86) and (4.87).} \end{aligned}$$

$$(4.89)$$

Since this inequality is true for every $y \in Y$,

$$(\nu \circ \mu)_x(U) - (\nu \circ \mu)_x(K) = \int_Y \left(\nu_y(U) - \nu_y(K)\right) d\mu_x(y) < \int_Y \epsilon d\mu_x = \epsilon$$
 (4.90)

because μ_x is a probability measure. This proves that $(\nu \circ \mu)_x$ is regular for arbitrary $x \in X$.

iii. Now, we check that the function $X \ni x \mapsto (\nu \circ \mu)_x$ is continuous so that $\nu \circ \mu : X \leadsto Z$ is a stochastic map. In this regard, fix an arbitrary $\varphi \in C(Z)$ and a net $x : \Theta \to X$ converging to $\lim x$. The goal is to show that

$$\lim_{\theta \in \Theta} \int_{Z} \varphi \ d(\nu \circ \mu)_{x_{\theta}} = \int_{Z} \varphi \ d(\nu \circ \mu)_{\lim x}. \tag{4.91}$$

By the Jordan decomposition theorem, it suffices to assume that φ is positive. Since φ is continuous, by Lemma 4.10 there exists a sequence of simple functions $0 \le s_1 \le s_2 \le \cdots \le \varphi$ of the form

$$s_n = \sum_{i_n} s_{n,i_n} \chi_{E_{n,i_n}} \tag{4.92}$$

defined on Borel sets $\{E_{n,i_n}\}$ in Z satisfying all of the conclusions of that Lemma. Hence

$$\lim_{\theta \in \Theta} \int_{Z} \varphi \ d(\nu \circ \mu)_{x_{\theta}} = \lim_{\theta \in \Theta} \lim_{n \to \infty} \sum_{i_{n}} s_{n,i_{n}} (\nu \circ \mu)_{x_{\theta}} (E_{n,i_{n}}) \quad \text{by definition of the integral}$$

$$= \lim_{\theta \in \Theta} \lim_{n \to \infty} \sum_{i_{n}} s_{n,i_{n}} \int_{Y} \nu_{y}(E_{n,i_{n}}) \ d\mu_{x_{\theta}} \quad \text{by definition of } \nu \circ \mu$$

$$= \lim_{\theta \in \Theta} \lim_{n \to \infty} \int_{Y} \sum_{i} s_{n,i_{n}} \nu_{y}(E_{n,i_{n}}) \ d\mu_{x_{\theta}} \quad \text{by linearity of the integral.}$$

$$(4.93)$$

Now, to verify that the limit can be interchanged with the integral, note that by the properties of the simple functions constructed above and since ν_y is a measure,

$$\sum_{i_n} s_{n,i_n} \nu_y(E_{n,i_n}) = \sum_{i_n} s_{n,i_n} \nu_y(E_{n+1,2i_n}) + \sum_{i_n} s_{n,i_n} \nu_y(E_{n+1,2i_n+1}) \quad \text{by (4.13)}$$

$$= \sum_{i_{n+1}} s_{n,\frac{i_{n+1}}{2}} \nu_y(E_{n+1,i_{n+1}}) + \sum_{i_{n+1}} s_{n,\frac{i_{n+1}-1}{2}} \nu_y(E_{n+1,i_{n+1}})$$

$$\leq \sum_{i_{n+1}} s_{n+1,i_{n+1}} \nu_y(E_{n+1,i_{n+1}}) + \sum_{i_{n+1}} s_{n,i_{n+1}-1} \nu_y(E_{n+1,i_{n+1}}) \quad \text{by (4.14)}$$

$$= \sum_{i_{n+1}} s_{n+1,i_{n+1}} \nu_y(E_{n+1,i_{n+1}})$$

for all $y \in Y$. Therefore, the sequence of functions

$$\mathbb{N} \times Y \ni (n, y) \mapsto \sum_{i_n} s_{n, i_n} \nu_y(E_{n, i_n}) \tag{4.95}$$

on Y is a monotonic increasing sequence of Borel measurable functions. Hence, continuing on from equation (4.93),

$$\lim_{\theta \in \Theta} \int_{Z} \varphi \ d(\nu \circ \mu)_{x_{\theta}} = \lim_{\theta \in \Theta} \int_{Y} \lim_{n \to \infty} \sum_{i_{n}} s_{n,i_{n}} \nu_{y}(E_{n,i_{n}}) \ d\mu_{x_{\theta}}$$
by the Monotone Convergence Theorem
$$= \lim_{\theta \in \Theta} \int_{Y} \left(\int_{Z} \varphi \ d\nu_{y} \right) \ d\mu_{x_{\theta}} \quad \text{by definition of the integral.}$$
(4.96)

To continue with this calculation, note that by assumption, the assignment $y \mapsto \nu_y$ is continuous with respect to the vague topology on ProbMeas(Z). This means that for any continuous function such as $\varphi \in C(Z)$ above, $y \mapsto \int_Z \varphi \ d\nu_y$ is a continuous real-valued function on Y. Hence, following from (4.96),

$$\lim_{\theta \in \Theta} \int_{Z} \varphi \ d(\nu \circ \mu)_{x_{\theta}} = \int_{Y} \left(\int_{Z} \varphi \ d\nu_{y} \right) \ d\mu_{\lim x}$$
 (4.97)

since μ is continuous with respect to the vague topology on ProbMeas(Y). By applying an analogous procedure (except without taking any limit in Θ), one similarly concludes that

$$\int_{Z} \varphi \ d(\nu \circ \mu)_{\lim x} = \int_{Y} \left(\int_{Z} \varphi \ d\nu_{y} \right) \ d\mu_{\lim x}. \tag{4.98}$$

Since these two results are equal, continuity of $\nu \circ \mu$ has been exhibited.

Therefore, $\nu \circ \mu : X \leadsto Z$ is a stochastic map.

From this fact, it immediately follows that the identity stochastic map is indeed an identity for the composition of stochastic maps.

Proposition 4.99. The composition of stochastic maps is associative.

Proof. Let $\mu: X \leadsto Y, \nu: Y \leadsto Z$, and $\zeta: Z \leadsto W$ be stochastic maps of compact Hausdorff spaces. The goal is to show that $(\zeta \circ (\nu \circ \mu))_x(E) = ((\zeta \circ \nu) \circ \mu)_x(E)$ for all Borel sets $E \in \mathcal{M}_W$ and points $x \in X$. By expanding out the definitions, it must be shown that

$$\int_{Z} \zeta_{z}(E) \ d(\nu \circ \mu)_{x}(z) = \int_{Y} \left(\int_{Z} \zeta_{z}(E) \ d\nu_{y}(z) \right) d\mu_{x}(y). \tag{4.100}$$

Contrary to what one might naively think, Fubini's Theorem does not apply here. Instead, since $Z \ni z \mapsto \zeta_z(E)$ is Borel measurable and bounded, by Lemma 4.10, there exists a sequence $s: \mathbb{N} \to \mathbb{R}^Z$ of simple functions satisfying

$$0 \le s_1(z) \le s_2(z) \le s_3(z) \le \dots \le \zeta_z(E) \qquad \& \qquad \lim_{n \to \infty} s_n(z) = \zeta_z(E) \qquad \forall \ z \in Z. \tag{4.101}$$

and all the other properties of that Lemma. Hence, by a completely analogous calculation as in (4.93)–(4.96) but without a net, the equality in (4.100) is obtained. This proves that composition of stochastic maps is associative.

In conclusion, we have proved the first part of the following theorem.

Theorem 4.102. The collection of compact Hausdorff spaces together with morphisms as stochastic maps forms a category, denoted by **cHausStoch**. In addition, the Dirac measure furnishes a faithful (but not full) functor δ : **cHaus** \rightarrow **cHausStoch** sending a space X to X and a continuous function $f: X \rightarrow Y$ to $\delta_f: X \leadsto Y$ as in Example 4.58.

Proof. The faithfulness of δ follows immediately from the Hausdorff condition. More precisely, seeking to prove the contrapositive, suppose that $f, g: X \to Y$ were two different functions. Then there exists an $x \in X$ such that $f(x) \neq g(x)$. Since Y is Hausdorff, there exist open sets U and V in Y such that $x \in U, y \in V$, and $U \cap V = \emptyset$. Then $\delta_{f(x)}(U) = 1$ while $\delta_{g(x)}(U) = 0$. Hence $\delta_f \neq \delta_g$.

4.4 From continuous probability theory to algebra

We have already associated to every compact Hausdorff space a commutative C^* -algebra by taking continuous functions on that space. To every stochastic map, we can also associate a canonical positive map of C^* -algebras.

Proposition 4.103. Let $\mu: X \leadsto Y$ be a stochastic map between compact Hausdorff spaces. For every continuous function $\varphi: Y \to \mathbb{C}$, the function $C^{\mu}(\varphi): X \to \mathbb{C}$ defined by

$$X \ni x \mapsto \int_{Y} \varphi \ d\mu_{x} \tag{4.104}$$

is a continuous function on X. Furthermore, $C^{\mu}: C(Y) \to C(X)$ is a (unital) positive map.

Proof. Linearity and positivity of C^{μ} follow immediately from the definitions and properties of a probability measure. To prove that $C^{\mu}(\varphi)$ is a continuous function for any continuous function $\varphi \in C(Y)$, let $x : \Theta \to X$ be a net converging to $\lim x \in X$. By definition of the vague topology on ProbMeas(Y) and because $\mu : X \to \text{ProbMeas}(Y)$ is continuous,

$$\lim_{\theta \in \Theta} \left(C^{\mu}(\varphi)(x_{\theta}) \right) = \lim_{\theta \in \Theta} \int_{Y} \varphi \ d\mu_{x_{\theta}} = \int_{Y} \varphi \ d\mu_{\lim x} = C^{\mu}(\varphi)(\lim x)$$
 (4.105)

showing that $C^{\mu}(\varphi)$ is continuous.

Theorem 4.106. The assignment

cHausStoch^{op}
$$\xrightarrow{C}$$
 cC *-AlgPos
 $X \mapsto C(X)$ (4.107)
 $(\mu : X \leadsto Y) \mapsto (C^{\mu} : C(Y) \leadsto C(X))$

is a functor. Moreover, the diagram

$$\begin{array}{ccc}
\mathbf{cHaus}^{\mathrm{op}} & \xrightarrow{C} \mathbf{cC} *-\mathbf{Alg} \\
\delta & & \downarrow \\
\mathbf{cHausStoch}^{\mathrm{op}} & \xrightarrow{C} \mathbf{cC} *-\mathbf{AlgPos}
\end{array} (4.108)$$

commutes (the functor on the right is the inclusion since every *-homomorphism is positive).

Proof. The fact that the codomain of C is as claimed follows from Proposition 4.103.

If $id_X: X \leadsto X$ is the identity stochastic map, then

$$\left(C^{\mathrm{id}_X}(\varphi)\right)(x) = \int_X \varphi \, d\delta_x = \varphi(x) \tag{4.109}$$

for all $x \in X$ and for all $\varphi \in C(X)$. Hence $C^{\mathrm{id}_X}(\varphi) = \varphi$ for all $\varphi \in C(X)$. Thus C preserves identities.

To see that C preserves the composition, let $\mu: X \leadsto Y$ and $\nu: Y \leadsto Z$ be two stochastic maps. The goal is to show that $C^{\nu \circ \mu} = C^{\mu} \circ C^{\nu}$. The proof will be similar to the proof that composition in **cHausStoch** is associative. To do this, fix $\varphi \in C(Z)$. By the Jordan decomposition theorem, it suffices to assume that $\varphi(Z) \subseteq \mathbb{R}_{\geq 0}$ since any measurable complex-valued function can be decomposed into its real and imaginary parts and each of those into positive and negative parts. Since φ is continuous and Z is compact, let s be a sequence of simple functions as in Lemma 4.10.

Then,

$$(C^{\nu\circ\mu}(\varphi))(x) = \int_{Z} \varphi \ d(\nu \circ \mu)_{x} \quad \text{by definition of } C^{\nu\circ\mu}$$

$$= \lim_{n \to \infty} \sum_{i_{n}} s_{n,i_{n}} (\nu \circ \mu)_{x} (E_{n,i_{n}}) \quad \text{by definition of the Lebesgue integral}$$

$$= \lim_{n \to \infty} \sum_{i_{n}} s_{n,i_{n}} \int_{Y} \nu_{y}(E_{n,i_{n}}) \ d\mu_{x}(y) \quad \text{by definition of } (\nu \circ \mu)_{x}$$

$$= \lim_{n \to \infty} \int_{Y} \sum_{i_{n}} s_{n,i_{n}} \nu_{y}(E_{n,i_{n}}) \ d\mu_{x}(y) \quad \text{by (finite) linearity of the integral}$$

$$= \int_{Y} \lim_{n \to \infty} \sum_{i_{n}} s_{n,i_{n}} \nu_{y}(E_{n,i_{n}}) \ d\mu_{x}(y) \quad \text{by Monotone Convergence Theorem}$$

$$= \int_{Y} \left(\int_{Z} \varphi \ d\nu_{y} \right) d\mu_{x}(y) \quad \text{by definition of the Lebesgue integral}$$

$$= \int_{Y} \left(C^{\nu}(\varphi) \right)(y) \ d\mu_{x}(y) \quad \text{by definition of } C^{\nu}$$

$$= \left(C^{\mu}(C^{\nu}(\varphi)) \right)(x) \quad \text{by definition of } C^{\mu}$$

for all $x \in X$. The Monotone Convergence Theorem can be applied in this calculation due to Lemma 4.10 and the way the simple functions were constructed in that Lemma (recall step iii. in the proof of Proposition 4.79). Since φ was arbitrary, this shows that $C^{\nu \circ \mu} = C^{\mu} \circ C^{\nu}$.

Finally, if $f: X \to Y$ is a continuous function, let $\varphi \in C(Y)$ and $x \in X$. Then

$$(C^f(\varphi))(x) = \int_Y \varphi \ d\delta_{f(x)} = \varphi(f(x))$$
 (4.111)

showing that the diagram in the statement of the theorem commutes.

4.5 The stochastic Gelfand spectrum functor

In this section, we prove that the functor $C: \mathbf{cHausStoch^{op}} \to \mathbf{cC^*-AlgPos}$ constructed in Section 4.4 is fully faithful. To do this, we review the Riesz-Markov-Kakutani Representation Theorem. Then we use these results to construct an analogue of the spectrum functor for positive maps on arbitrary commutative C^* -algebras.

Theorem 4.112 (Riesz-Markov-Kakutani (RMK) Representation Theorem). Let Y be a compact Hausdorff space and let $I: C(Y) \leadsto \mathbb{C}$ be a state on C(Y). Then, there exists a unique regular probability measure $\mu \in \operatorname{ProbMeas}(Y)$ such that

$$I(\varphi) = \int_{Y} \varphi \, d\mu \qquad \forall \, \varphi \in C(Y). \tag{4.113}$$

Furthermore, this measure μ satisfies

$$\mu(U) = \sup \{ I(\varphi) : \varphi \in C(Y) \text{ and } 0 \le \varphi \le \chi_U \}$$
(4.114)

for all open sets $U \subseteq Y$ and

$$\mu(K) = \inf \left\{ I(\varphi) : \varphi \in C(Y) \text{ and } \varphi \ge \chi_K \right\}$$
(4.115)

for all closed (and hence compact) sets $K \subseteq Y$.

Note that by our definition, a state is a positive linear functional that is unit-preserving. This is why the measure μ in the statement of the theorem is a probability measure. We will not prove this theorem in its entirety, but an explicit construction of the measure is useful and provides a close comparison to the finite-dimensional case. Namely, when X is finite, $C(X) = \mathbb{C}^X$ and it makes sense to assign $\mu(E) := I(\chi_E)$ for any finite subset $E \subseteq X$ as we have done in Section 2. This is because the characteristic function χ_E is continuous on a finite set. Hence, heuristically, given a measurable set $E \in \mathcal{M}_Y$, one would expect $\mu(E)$ to be given by $I(\chi_E)$. However, χ_E is not continuous and so this expression simply does not make sense because χ_E is not in the domain of I. Nevertheless, we know by regularity that E can be approximated using open and closed sets. One can use this fact to first extend I to be a positive linear functional on the larger space of lower semi-continuous functions as is done in [48].

Sketch of proof. First, let $U \subseteq Y$ be open and let $K \subseteq Y$ be closed. Set

$$\mu(U) := \sup_{\substack{\varphi \in C(Y) \\ 0 \le \varphi \le \chi_U}} \left\{ I(\varphi) \right\} \qquad \& \qquad \mu(K) := \inf_{\substack{\varphi \in C(Y) \\ 0 \le \chi_K \le \varphi}} \left\{ I(\varphi) \right\}. \tag{4.116}$$

For an arbitrary subset E of Y, set

$$\mu^{+}(E) = \inf_{\substack{U \text{ open} \\ E \subseteq U}} \{\mu(U)\} \qquad \& \qquad \mu^{-}(E) = \sup_{\substack{K \text{ closed} \\ K \subseteq E}} \{\mu(K)\}. \tag{4.117}$$

It is a fact that for all Borel subsets $E \in \mathcal{M}_Y$,

$$\mu^{+}(E) = \mu^{-}(E) \tag{4.118}$$

and we therefore denote this assignment by μ . It is also a fact that μ is the unique regular probability measure on Y satisfying the condition in the statement of the theorem. See Theorem 7.2 in [14], Theorem 8 in [48], or Theorem 2.14 in [38] for proofs of these claims.

Remark 4.119. In general, the assignments μ^{\pm} on arbitrary subsets E of Y given by

$$\mu^{+}(E) := \inf_{\substack{U \text{ open} \\ E \subseteq U}} \left\{ \sup_{\substack{\varphi \in C(Y) \\ 0 \le \varphi \le \chi_{U}}} \left\{ I(\varphi) \right\} \right\}$$
(4.120)

and

$$\mu^{-}(E) := \sup_{\substack{K \text{ closed} \\ K \subseteq E}} \left\{ \inf_{\substack{\varphi \in C(Y) \\ 0 \le \chi_K \le \varphi}} \left\{ I(\varphi) \right\} \right\}$$
(4.121)

are not the same. These are known as *outer* and *inner* measures associated to I. The collection of subsets E for which $\mu^+(E) = \mu^-(E)$ contains the Borel sets of Y. In general, however, the resulting σ -algebra is larger than the Borel σ -algebra (see Theorem 2.14 in [38]). Nevertheless, μ restricted to the Borel σ -algebra associated to Y provides Y with a regular probability measure. This is the measure constructed in the proof of Theorem 4.112.

Proposition 4.122. cHausStoch^{op} \xrightarrow{C} cC*-AlgPos is fully faithful. More explicitly, let X and Y be compact Hausdorff spaces and let $f: C(Y) \leadsto C(X)$ be a positive (unital) map. Then there exists a unique stochastic map $\mu: X \leadsto Y$ such that

$$(f(\varphi))(x) = \int_{Y} \varphi \ d\mu_x \qquad \forall \ x \in X \qquad \forall \ \varphi \in C(Y).$$
 (4.123)

Proof. First note that for every $\varphi \in C(Y)$ and $x \in X$,

$$(f(\varphi))(x) = (\operatorname{ev}_x \circ f)(\varphi),$$
 (4.124)

where $\operatorname{ev}_x:C(X)\to\mathbb{C}$ is the evaluation at x function (recall Example 2.48), which is a *-homomorphism and therefore a positive map. Hence, the composition $C(Y)\stackrel{f}{\leadsto}C(X)\stackrel{\operatorname{ev}_x}{\Longrightarrow}\mathbb{C}$ is a state on C(Y). By the RMK Representation Theorem, there exists a unique regular probability measure $\mu_x\in\operatorname{ProbMeas}(Y)$ satisfying (4.123). The only thing left to check is that the assignment $X\ni x\mapsto \mu_x\in\operatorname{ProbMeas}(Y)$ is continuous. However, this immediately follows from the fact that $\varphi(f)$ is continuous since if $x:\Theta\to X$ is a net converging to $\lim x$, then

$$\lim_{\theta \in \Theta} \int_{Y} \varphi \ d\mu_{x_{\theta}} = \lim_{\theta \in \Theta} (f(\varphi))(x_{\theta}) = (f(\varphi))(\lim x) = \int_{Y} \varphi \ d\mu_{\lim x}. \tag{4.125}$$

Hence, C is fully faithful.

Thus, the RMK Representation Theorem precisely encodes the fact that the functor C: $\mathbf{cHausStoch^{op}} \to \mathbf{cC^*-AlgPos}$ is fully faithful while the Gelfand-Naimark Theorem shows that this functor is essentially surjective. Together, these results hold if and only if C is part of adjoint equivalence of categories. However, this result is somewhat formal. It is useful to have an explicit functor $\sigma: \mathbf{cC^*-AlgPos} \to \mathbf{cHausStoch^{op}}$ that (1) requires no reference to any underlying space structure and (2) agrees with the usual spectrum functor from the ordinary (commutative) Gelfand-Naimark correspondence when restricted to *-homomorphisms of C^* -algebras.

We have already constructed the functor σ at the level of objects, namely, given any commutative C^* -algebra \mathcal{A} , take $\sigma(\mathcal{A})$ to be the spectrum of \mathcal{A} (see Definition 3.38). Given a *homomorphism of commutative C^* -algebras $f: \mathcal{B} \to \mathcal{A}$, the associated map $\sigma(f)$ assigns to each character $\chi \in \sigma(\mathcal{A})$ the linear functional $\chi \circ f: \mathcal{B} \to \mathbb{C}$. This linear functional is a character precisely because f and χ are both (non-vanishing) *-homomorphisms. Hence, when f is simply a positive map, $\chi \circ f$ is no longer a character in general. Nevertheless, from these data, namely a positive map $f: \mathcal{B} \leadsto \mathcal{A}$ and a character $\chi \in \sigma(\mathcal{A})$, we must construct a canonical regular probability measure σ_{χ}^f on $\sigma(\mathcal{B})$ in such a way so that this assignment $\sigma^f: \sigma(A) \to \operatorname{ProbMeas}(\sigma(\mathcal{B}))$ is a stochastic map. The idea should be that we smear $\chi \circ f$ onto $\sigma(\mathcal{B})$ in some way that captures the information of the state $\chi \circ f$ as depicted in Figure 4.

This can be done formally using the Gelfand transform and the fact that C is fully faithful. Namely, we can fill in the diagram

$$\begin{array}{c|c}
\mathcal{B} & & \mathcal{A} \\
\Gamma_{\mathcal{B}}^{-1} & & & & \downarrow \Gamma_{\mathcal{A}} \\
C(\sigma(\mathcal{B})) \sim & & & \sim C(\sigma(\mathcal{A}))
\end{array} (4.126)$$




Figure 4: A character provides a Dirac point measure on the spectrum. A state is "smeared" onto the spectrum in terms of a probability measure representing the associated "weight" of that state on the characters in the spectrum.

on the bottom by using the inverse of the Gelfand transform, the positive map f, and then the Gelfand transform once more. This defines a positive map $C^{\sigma(f)}: C(\sigma(\mathcal{B})) \leadsto C(\sigma(\mathcal{A}))$ which we know is uniquely characterized by a stochastic map $\sigma^f: \sigma(\mathcal{A}) \leadsto \sigma(\mathcal{B})$ by the RMK theorem. We could simply define $\sigma(f)$ in this manner and move on. However, a more geometrically appealing formula for $\sigma(f)$ is available in terms of the convex structure on the set of states with reference to neither the Gelfand transform nor the RMK theorem. As the reader may have guessed from the suggestive drawing above, this merits a review of convex sets, their extreme points, the Krein-Milman theorem, and Choquet theory [44], [35].

Definition 4.127. Let V be a topological vector space. A subset $C \subseteq V$ is said to be *convex* iff

$$\lambda v + (1 - \lambda)w \in C \qquad \forall \ \lambda \in [0, 1] \text{ whenever } v, w \in C.$$
 (4.128)

An <u>extreme point</u> of C is a point $u \in C$ such that if $u = \lambda v + (1 - \lambda)w$ for some $v, w \in C$ and $\lambda \in [0, 1]$, then v = u and/or w = u. The set of extreme points of C is denoted by ex(C).

Example 4.129. Let \mathcal{A} be a commutative C^* -algebra. Then, $\mathcal{S}(\mathcal{A})$, the set of all states, is a compact convex subset of \mathcal{A}^* in the weak* topology. $\sigma(\mathcal{A})$, the spectrum of \mathcal{A} , is the set of extreme points of $\mathcal{S}(\mathcal{A})$. This follows, for instance, from Theorem 11.33 in Rudin [39].

Definition 4.130. Let V be a topological vector space and let $X \subseteq V$. The <u>convex hull</u> of X is given by

$$\operatorname{ch}(X) := \left\{ \sum_{i=1}^{\text{finite}} \lambda_{i} v_{i} : v_{i} \in X, \ \lambda_{i} \in [0, 1], \ \sum_{i=1}^{n} \lambda_{i} = 1 \right\}, \tag{4.131}$$

i.e. the set of all finite convex combinations of elements of X.

The following is an application of the Krein-Milman theorem suitable for our purposes [44].

Theorem 4.132. Let A be a commutative C^* -algebra. Then the set of all states on A is the closure of the convex hull of the set of characters on A, i.e.

$$S(A) = \overline{\operatorname{ch}(\sigma(A))}. \tag{4.133}$$

Proof. This follows from the usual Krein-Milman theorem, the conclusions of Example 4.129, and the fact that \mathcal{A}^* is a locally convex topological vector space (it is locally convex due to the separating family of seminorms generating its topology).

This important result implies the existence of a probability measure associated to any state.

Theorem 4.134. Let \mathcal{A} be a commutative C^* -algebra and let $\omega \in \mathcal{S}(\mathcal{A})$ be a state on \mathcal{A} . Then there exists a unique regular probability measure μ on $\sigma(\mathcal{A})$ such that

$$\int_{\sigma(\mathcal{A})} \operatorname{ev}_a d\mu = \omega(a) \qquad \forall \ a \in \mathcal{A}. \tag{4.135}$$

Here $\operatorname{ev}_a : \sigma(\mathcal{A}) \to \mathbb{C}$ denotes the evaluation at the point a sending a character $\chi \in \sigma(\mathcal{A})$ to $\chi(a)$.

The statement of the theorem can be obtained from the RMK Representation Theorem combined with the Gelfand-Naimark Theorem. However, we prefer to prove this theorem directly using Choquet theory, which provides a concrete and geometric construction of the measure. We thank Benjamin Russo for discussions leading to this construction.

Proof. We first establish existence and then uniqueness.

By the Krein-Milman Theorem, $\omega \in \overline{\operatorname{ch}(\sigma(\mathcal{A}))}$. Hence, there exists a net $\xi : \Theta \to \operatorname{ch}(\sigma(\mathcal{A}))$ such that $\lim \xi = \omega$ in the weak* topology on $\mathcal{S}(\mathcal{A})$. Since each ξ_{θ} is in the convex hull of $\sigma(\mathcal{A})$, there exists an $N_{\theta} \in \mathbb{N}$ and a convex decomposition of ξ_{θ} into characters, i.e.

$$\xi_{\theta} = \sum_{i_{\theta}=1}^{N_{\theta}} \lambda_{i_{\theta}} \chi_{i_{\theta}}, \tag{4.136}$$

where $\chi_{i_{\theta}} \in \sigma(\mathcal{A})$ for all $i_{\theta} \in \{1, \dots, N_{\theta}\}$. Set

$$\mu_{\theta} := \sum_{i_{\theta}=1}^{N_{\theta}} \lambda_{i_{\theta}} \delta_{\chi_{i_{\theta}}} \tag{4.137}$$

to be the convex combination of Dirac measures on these characters. This defines a net of regular probability measures on $\sigma(\mathcal{A})$. Since ProbMeas $(\sigma(\mathcal{A}))$ is compact in the vague topology, there exists a subnet $(\mu_{\theta'})_{\theta' \in \Theta' \subseteq \Theta}$ of these measures converging in the vague topology to some regular probability measure, which we denote by μ . This probability measure satisfies (4.135) since

$$\int_{\sigma(\mathcal{A})} \operatorname{ev}_a d\mu = \lim_{\theta' \in \Theta'} \int_{\sigma(\mathcal{A})} \operatorname{ev}_a d\mu_{\theta'} \quad \text{by continuity of ev}_a \text{ and vague convergence}$$

$$= \lim_{\theta' \in \Theta'} \sum_{i_{\theta'}=1}^{N_{\theta'}} \lambda_{i_{\theta'}} \int_{\sigma(\mathcal{A})} \operatorname{ev}_a d\delta_{\chi_{i_{\theta'}}} \quad \text{by linearity of the measure and integral}$$

$$= \lim_{\theta' \in \Theta'} \sum_{i_{\theta'}=1}^{N_{\theta'}} \lambda_{i_{\theta'}} \chi_{i_{\theta'}}(a) \quad \text{by definition of the Dirac measure}$$

$$= \lim_{\theta' \in \Theta'} \sum_{i_{\theta'}=1}^{N_{\theta'}} \xi_{\theta'}(a) \quad \text{by definition of } \xi$$

$$= \omega(a) \quad \text{by weak* convergence.}$$

This proves existence of such a measure.

Uniqueness is more subtle and follows, for instance, from the Choquet-Meyer Theorem. More specifically, it follows from Corollary 10.9 in [35], whose notation we follow. We do not introduce all the necessary definitions to explain this in full detail, but we outline the idea using the notation and terminology from [35]. \mathcal{A}^* , the topological dual space of \mathcal{A} with the weak* topology, is a locally convex vector space (this follows, for instance, from Proposition 3.29 above and Theorem 1.37 in [39]). The set of all states $\mathcal{S}(\mathcal{A})$ on a commutative (unital) C^* -algebra is a compact convex subset of \mathcal{A}^* and is a Choquet simplex (see Theorem 3.1.18 in [40] and note that states on a commutative C^* -algebra are automatically tracial). Because $\exp(\mathcal{S}(\mathcal{A})) = \sigma(\mathcal{A})$ is a closed set (since it is compact), every maximal measure is supported on $\sigma(\mathcal{A})$ (see the comments preceding the Bishop-de Leeuw Theorem in Section 4 of [35]). To see that

$$\int_{\mathcal{S}(\mathcal{A})} \varphi \ d\mu = \int_{\mathcal{S}(\mathcal{A})} \varphi \ d\delta_{\omega} \tag{4.139}$$

 $(\mu \sim \delta_{\omega} \text{ in the notation of [35]})$ for all continuous real-valued affine functions φ on $\mathcal{S}(\mathcal{A})$, notice that all functions of the form $\mathrm{ev}_a : \mathcal{A}^* \to \mathbb{C}$ are continuous and linear and hence remain continuous when restricted to $\mathcal{S}(\mathcal{A}) \subseteq \mathcal{A}^*$. By construction of μ from the subnet $(\mu_{\theta'})$ and the definition of the Dirac measure,

$$\int_{\mathcal{S}(\mathcal{A})} \operatorname{ev}_a d\mu = \int_{\sigma(\mathcal{A})} \operatorname{ev}_a d\mu = \omega(a) = \int_{\mathcal{S}(\mathcal{A})} \operatorname{ev}_a d\delta_\omega \qquad \forall \ a \in \mathcal{A}.$$
(4.140)

Now, suppose that φ is an arbitrary continuous real-valued affine function on $\mathcal{S}(\mathcal{A})$. Then, φ restricted to $\sigma(\mathcal{A})$ is a continuous function and therefore equals ev_a for some $a \in \mathcal{A}$ by Proposition 3.57. If we show that $\varphi = \operatorname{ev}_a$ on all of $\mathcal{S}(\mathcal{A})$, then we will be done by using the previous calculation. Note that because $\mathcal{S}(\mathcal{A})$ is a compact convex subset of a locally convex space, the supremum and infimum of φ are achieved on a face of $\mathcal{S}(\mathcal{A})$. This is a consequence, for instance, of Proposition 8.26 in [44] by noting that the image of a non-empty compact convex set under such a continuous map is an interval in \mathbb{R} and the supremum and infimum of an interval are at most two points. These endpoints are faces of the interval and so their pre-images are faces of the simplex $\mathcal{S}(\mathcal{A})$. Furthermore, the extreme points of a face are extreme points of the original convex subset (a simple calculation from the definitions shows this). Hence, $|\varphi|$ achieves its supremum on $\sigma(\mathcal{A})$ (and possibly on a larger domain as well). Hence, by the Hahn-Banach Theorem (see Section 5.22 of [38]), there exists a unique extension of $\varphi|_{\sigma(\mathcal{A})} = \operatorname{ev}_a$ to $\mathcal{S}(\mathcal{A})$. But we already know that ev_a is such an extension. Hence $\varphi = \operatorname{ev}_a$. Since the result was already proved for such functions, $\mu \sim \delta_\omega$ so uniqueness of the regular probability measure follows from the Choquet-Meyer theorem.

Remark 4.141. Uniqueness also follows from a theorem due to Bauer since $\mathcal{S}(\mathcal{A})$ is a Bauer simplex for \mathcal{A} a commutative C^* -algebra (see Theorem II.4.1 in [1]).

Remark 4.142. Note that because the evaluation map ev_E : $\operatorname{ProbMeas}(\sigma(\mathcal{A})) \to \mathbb{R}$ is not necessarily continuous for every Borel subset $E \subseteq \sigma(\mathcal{A})$, in general

$$\mu(E) \neq \lim_{\theta \in \Theta} \sum_{i_{\theta} \in I_{\theta}} \lambda_{i_{\theta}}(E), \tag{4.143}$$

where I_{θ} is the subset of $\{1,\ldots,N_{\theta}\}$ for which $\chi_{i_{\theta}} \in E$. A counterexample is given by the sequence

$$\mathbb{N} \ni n \mapsto \mu_n := \frac{1}{n} \sum_{k=1}^n \delta_{\frac{k}{n}} \tag{4.144}$$

of a uniform sum of Dirac measures on [0,1]. This sequence converges to the standard Lebesgue measure μ on [0,1] in the vague topology. However, $\mu_n(\mathbb{Q} \cap [0,1]) = 1$ while $\mu(\mathbb{Q} \cap [0,1]) = 0$.

In the finite-dimensional setting, this does not occur, nets are not required, and ω is uniquely expressed as a convex decomposition of extremal states. The measure of E is the sum of the weights corresponding to restricting this sum to the contributions coming only from E. This is because the definition of a Choquet simplex is one that agrees with the definition of a simplex in finite dimensional vector spaces (see Proposition 10.10 in [35]).

Theorem 4.145. Let \mathcal{B} and \mathcal{A} be commutative C^* -algebras and let $f: \mathcal{B} \leadsto \mathcal{A}$ be a positive (unital) map. For each character $\chi \in \sigma(\mathcal{A})$, there exists a unique regular probability measure $\sigma_{\chi}^f \in \operatorname{ProbMeas}(\sigma(\mathcal{B}))$ such that

$$\int_{\sigma(\mathcal{B})} \operatorname{ev}_b \, d\sigma_{\chi}^f = \chi(f(b)) \qquad \forall \, b \in \mathcal{B}. \tag{4.146}$$

Furthermore, the assignment

$$\sigma(\mathcal{A}) \xrightarrow{\sigma(f) \equiv \sigma^f} \text{ProbMeas}(\sigma(\mathcal{B}))$$

$$\chi \longmapsto \sigma_{\chi}^f, \tag{4.147}$$

where σ_{χ}^{f} is the unique regular probability measure on $\sigma(\mathcal{B})$ associated to the state $\chi \circ f \in \mathcal{S}(\mathcal{B})$ via Theorem 4.134, defines a stochastic map $\sigma(\mathcal{A}) \leadsto \sigma(\mathcal{B})$.

Proof. Since $\chi \circ f$ is a state on \mathcal{A} , Theorem 4.134 guarantees there exists a unique regular probability measure σ_{χ}^f satisfying (4.146). The only thing left to check is that this defines a stochastic map, i.e. that the function $\sigma^f : \sigma(\mathcal{A}) \to \operatorname{ProbMeas}(\sigma(\mathcal{B}))$ is continuous with respect to the vague topology. Let $\chi : \Theta \to \sigma(\mathcal{A})$ be a net of characters converging to $\lim \chi$. The goal is to show that

$$\lim_{\theta \in \Theta} \int_{\sigma(\mathcal{B})} \varphi \, d\sigma_{\chi_{\theta}}^{f} = \int_{\sigma(\mathcal{B})} \varphi \, d\sigma_{\lim \chi}^{f} \qquad \forall \, \varphi \in C(\sigma(\mathcal{B})). \tag{4.148}$$

The Gelfand transform $\Gamma_{\mathcal{B}}: \mathcal{B} \to C(\sigma(\mathcal{B}))$ sending $b \in \mathcal{B}$ to the evaluation function ev_b is a C^* -algebra isomorphism by Proposition 3.57. Hence, every continuous function on the spectrum of \mathcal{B} is of this form. This fact simplifies the above goal to simply showing that

$$\lim_{\theta \in \Theta} \int_{\sigma(\mathcal{B})} \operatorname{ev}_b \, d\sigma_{\chi_\theta}^f = \int_{\sigma(\mathcal{B})} \operatorname{ev}_b \, d\sigma_{\lim \chi}^f \qquad \forall \, b \in \mathcal{B}. \tag{4.149}$$

This follows from the previous results since

$$\lim_{\theta \in \Theta} \int_{\sigma(\mathcal{B})} \operatorname{ev}_b \, d\sigma_{\chi_\theta}^f = \lim_{\theta \in \Theta} \chi_\theta \big(f(b) \big) \quad \text{by (4.146)}$$

$$= (\lim \chi) \big(f(b) \big) \quad \text{by weak* convergence of the net } \chi \qquad (4.150)$$

$$= \int_{\sigma(\mathcal{B})} \operatorname{ev}_b \, d\sigma_{\lim \chi}^f \quad \text{by (4.146)}$$

for all $b \in \mathcal{B}$. Therefore σ^f is a stochastic map.

Theorem 4.151. The assignment

$$\mathbf{cC}^*\text{-}\mathbf{AlgPos}^{\mathrm{op}} \xrightarrow{\sigma} \mathbf{cHausStoch}$$

$$\mathcal{A} \mapsto \sigma(\mathcal{A}) \qquad (4.152)$$

$$\left(\mathcal{B} \xrightarrow{f} \mathcal{A}\right) \mapsto \left(\sigma(\mathcal{A}) \xrightarrow{\sigma(f) \equiv \sigma^f} \sigma(\mathcal{B})\right)$$

from Definition 3.38 and Theorem 4.145 defines a functor, henceforth referred to as the stochastic spectrum functor.

Proof. The identity positive map gets sent to the identity stochastic map by uniqueness of the measure and the fact that the Dirac measure is a regular measure satisfying the required conditions. Let \mathcal{C}, \mathcal{B} , and \mathcal{A} be commutative C^* -algebras and let $f: \mathcal{B} \leadsto \mathcal{A}$ and $g: \mathcal{C} \leadsto \mathcal{B}$ be positive maps. Let $\sigma^f: \sigma(\mathcal{A}) \to \operatorname{ProbMeas}(\sigma(\mathcal{B}))$, $\sigma^g: \sigma(\mathcal{B}) \to \operatorname{ProbMeas}(\sigma(\mathcal{C}))$, and $\sigma^{f \circ g}: \sigma(\mathcal{A}) \to \operatorname{ProbMeas}(\sigma(\mathcal{C}))$ denote the images of f, g, and $f \circ g$, under σ , respectively. Let $\sigma^g \circ \sigma^f: \sigma(\mathcal{A}) \to \operatorname{ProbMeas}(\sigma(\mathcal{C}))$ denote the composition $\sigma(\mathcal{A}) \leadsto \sigma(\mathcal{B}) \leadsto \sigma(\mathcal{C})$ of the stochastic map σ^f followed by σ^g . The goal is to show $\sigma^{f \circ g} = \sigma^g \circ \sigma^f$. Since $\sigma_\chi^{f \circ g}$ and $(\sigma^g \circ \sigma^f)_\chi$ are regular probability measures for each $\chi \in \sigma(\mathcal{A})$, by uniqueness in Theorem 4.145, it suffices to show

$$\int_{\sigma(\mathcal{C})} \operatorname{ev}_c d(\sigma^g \circ \sigma^f)_{\chi} = \chi \Big(f(g(c)) \Big)$$
(4.153)

for all $c \in \mathcal{C}$ and for all $\chi \in \sigma(\mathcal{A})$. Therefore, first let $c \in \mathcal{C}$ be positive so that $\operatorname{ev}_c \geq 0$, and fix $\chi \in \sigma(\mathcal{A})$, and let

$$0 \le s_1 \le s_2 \le \dots \le \operatorname{ev}_c \tag{4.154}$$

be a sequence $s: \mathbb{N} \to \mathbb{R}^{\sigma(\mathcal{C})}$ of simple functions converging to ev_c and satisfying all the conditions

as in Lemma 4.10. Then,

$$\chi\Big(f\big(g(c)\big)\Big) = \int\limits_{\sigma(\mathcal{B})} \operatorname{ev}_{g(c)}(\psi) \ d\sigma_{\chi}^{f}(\psi) \quad \text{by definition of } \sigma_{\chi}^{f}$$

$$= \int\limits_{\sigma(\mathcal{B})} \psi\big(g(c)\big) \ d\sigma_{\chi}^{f}(\psi) \quad \text{by definition of } \operatorname{ev}_{g(c)}$$

$$= \int\limits_{\sigma(\mathcal{B})} \left(\int\limits_{\sigma(\mathcal{C})} \operatorname{ev}_{c} \ d\sigma_{\psi}^{g}\right) \ d\sigma_{\chi}^{f}(\psi) \quad \text{by definition of } \sigma_{\psi}^{g}$$

$$= \int\limits_{\sigma(\mathcal{B})} \lim\limits_{n \to \infty} \sum_{i_{n}} s_{n,i_{n}} \sigma_{\psi}^{g}(E_{n,i_{n}}) \ d\sigma_{\chi}^{f}(\psi) \quad \text{by definition of Lebesgue integral}$$

$$= \lim\limits_{n \to \infty} \sum_{i_{n}} s_{n,i_{n}} \int\limits_{\sigma(\mathcal{B})} \sigma_{\psi}^{g}(E_{n,i_{n}}) \ d\sigma_{\chi}^{f}(\psi) \quad \text{by Monotone Convergence Theorem}$$

$$= \lim\limits_{n \to \infty} \sum_{i_{n}} s_{n,i_{n}} (\sigma^{g} \circ \sigma^{f})_{\chi}(E_{n,i_{n}}) \quad \text{by definition of } (\sigma^{g} \circ \sigma^{f})_{\chi}$$

$$= \int\limits_{\sigma(\mathcal{C})} \lim\limits_{n \to \infty} \sum_{i_{n}} s_{n,i_{n}} \chi_{E_{n,i_{n}}} \ d(\sigma^{g} \circ \sigma^{f})_{\chi} \quad \text{by definition of Lebesgue integral}$$

$$= \int\limits_{\sigma(\mathcal{C})} \operatorname{ev}_{c} \ d(\sigma^{g} \circ \sigma^{f})_{\chi} \quad \text{by definition of } s.$$

Again, the Monotone Convergence Theorem applies by similar arguments as in step iii. in the proof of Proposition 4.79. The case of more general $c \in \mathcal{C}$ for which ev_c need not be positive is handled by splitting up ev_c using a Jordan decomposition and applying a similar procedure.

Example 4.156. Let $\rho: [-1,1] \to \mathbb{R}$ be the restriction of the Gaussian used to describe the distribution of heat after some time with initial condition a source of heat at the origin, namely

$$[-1,1] \ni x \mapsto \rho(x) := \frac{\exp(-100x^2)}{\int_{-1}^1 \exp(-100y^2) \, dy},\tag{4.157}$$

which has been properly normalized so that the integral of ρ with respect to the Lebesgue measure is 1. Consider the sequence of probability measures given by

$$\mathbb{N} \ni n \mapsto \mu_n := \sum_{k=1}^n \left(\int_{-1 + \frac{2(k-1)}{n}}^{-1 + \frac{2k}{n}} \rho(x) \, dx \right) \delta_{-1 + \frac{2k-1}{n}}. \tag{4.158}$$

The distribution and resulting convex sum approximation is depicted in Figure 5 for several values of $n \in \mathbb{N}$. Although the sequence of measures $d\mu_n$ does not converge to $\rho d\mu_{\text{Lebesgue}}$ in the total variation norm topology, the convergence holds in the vague topology, i.e.

$$\lim_{n \to \infty} \int_{-1}^{1} \varphi \ d\mu_n = \int_{-1}^{1} \varphi(x)\rho(x) \ dx \qquad \forall \ \varphi \in C([-1,1]). \tag{4.159}$$




Figure 5: The curve drawn in red is the graph of ρ in (4.157) (cut off from above to fit on the page) while the (larger) blue bullets represent the heights of the Dirac point distributions. The heights for the point measures decrease because the area over which they are used to approximate functions decreases as n increases.

This gives an example of a probability measure on [-1,1] that can be approximated by a sequence of convex sums of Dirac point measures with respect to the vague topology. Again, it is not true that $\lim_{n\to\infty} \mu_n(E) = \int_E \rho(x) \ dx$ for all Borel measurable sets E. For example, take $E := [-1,1] \cap (\mathbb{R} \setminus \mathbb{Q})$, all irrationals in [-1,1]. The term on the left is 0 while the term on the right is 1.

4.6 The stochastic commutative Gelfand-Naimark Theorem

Notice that when $\mathcal{A} = C(X)$ with X a compact Hausdorff space and $\omega \in \mathcal{S}(C(X))$, Theorem 4.134 says that there exists a unique regular probability measure μ on $\sigma(C(X))$ such that

$$\int_{\sigma(C(X))} \operatorname{ev}_{\varphi} d\mu = \omega(\varphi) \qquad \forall \ \varphi \in C(X). \tag{4.160}$$

Meanwhile, by the RMK Representation Theorem, there exists a unique regular probability measure ν on X such that

$$\omega(\varphi) = \int_X \varphi \ d\nu \qquad \forall \ \varphi \in C(X). \tag{4.161}$$

By the Gelfand-Naimark Theorem, we know that the spaces X and $\sigma(C(X))$ are homeomorphic and one expects these two measures to be related. In fact, the function $h_X: X \to \sigma(C(X))$ pushes forward the measure ν to μ and the inverse $h_X^{-1}: \sigma(C(X)) \to X$ pushes forward the measure μ to ν . This is explained in more detail in the following theorem.

Theorem 4.162. The stochastic spectrum functor together with the natural isomorphisms

cHausStoch
$$\leftarrow$$
 id cHausStoch \leftarrow cC*-AlgPos \leftarrow id cC*-AlgPos \leftarrow cC*-AlgPos \leftarrow cC*-AlgPos \leftarrow cHausStoch \leftarrow cC*-AlgPos \leftarrow cHausStoch \leftarrow

from the commutative Gelfand-Naimark Theorem form an adjoint equivalence of categories.

Proof. The only thing that has not been shown is that the transformations Γ and h still satisfy naturality. Note that each is an isomorphism in its respective category on objects due to the usual Gelfand-Naimark Theorem.

i. Proof that h is natural. Let $\mu: X \leadsto Y$ be a stochastic map and let $\sigma^{C(\mu)}: \sigma(C(X)) \leadsto \sigma(C(Y))$ denote the stochastic map obtained from applying the functor C and then σ to μ . The goal is to prove that the diagram

commutes. Note that for any Borel set $E \in \mathcal{M}_{\sigma(C(Y))}$ and for any $x \in X$, the lower left composition $\delta_{h_Y} \circ \mu$ applied to x evaluated at E gives

$$(\delta_{h_Y} \circ \mu)_x(E) = \int_Y \delta_{h_Y(y)}(E) \ d\mu_x(y) = \int_Y \chi_{h_Y^{-1}(E)}(y) \ d\mu_x(y) = \mu_x (h_Y^{-1}(E)). \tag{4.165}$$

In other words, $(\delta_{h_Y} \circ \mu)_x$ is the pushforward of the measure μ_x along the function h_Y . Now we check that this equals $(\sigma^{C(\mu)} \circ h_X)_x$ for every $x \in X$ by integrating ev_E with respect to this pushforward measure and $\varphi \in C(Y)$. By the change of variables formula for integrating measurable functions with respect to pushforward measures (see Theorem 3.6.1 in [7]),

$$\int_{\sigma(C(Y))} \operatorname{ev}_{\varphi} d(\delta_{h_Y} \circ \mu)_x = \int_{\sigma(C(Y))} \operatorname{ev}_{\varphi} d(\mu_x \circ h_Y^{-1}) = \int_Y (\operatorname{ev}_{\varphi} \circ h_Y) d\mu_x$$
(4.166)

since ev_{φ} is continuous (and hence measurable). Therefore,

$$\int_{Y} \operatorname{ev}_{\varphi} d(\delta_{h_{Y}} \circ \mu)_{x} = \int_{Y} \operatorname{ev}_{\varphi}(h_{Y}(y)) d\mu_{x}(y) \quad \text{by (4.166)}$$

$$= \int_{Y} h_{Y}(y)(\varphi) d\mu_{x}(y) \quad \text{by definition of } \operatorname{ev}_{\varphi}$$

$$= \int_{Y} \varphi(y) d\mu_{x}(y) \quad \text{by definition of } h_{Y} \text{ (3.64)}$$

$$= (C^{\mu}(\varphi))(x) \quad \text{by definition of } C^{\mu} \text{ (4.104)}$$

$$= h_{X}(x)(C^{\mu}(\varphi)) \quad \text{by definition of } h_{X} \text{ (3.64)}$$

$$= \int_{\varphi(C(Y))} \operatorname{ev}_{\varphi} d\sigma_{h_{X}(x)}^{C(\mu)} \quad \text{by definition of } \sigma_{h_{X}(x)}^{C(\mu)}.$$

Since this is true for all $\varphi \in C(Y)$, the uniqueness of measures on $\sigma(C(Y))$ from Theorem 4.134 implies $\sigma_{h_X(x)}^{C(\mu)} = (\delta_{h_Y} \circ \mu)_x$ for all $x \in X$. Therefore, the diagram above commutes and $h : \mathrm{id}_{\mathbf{cHausStoch}} \Rightarrow \sigma \circ C$ is a natural isomorphism.

ii. Proof that Γ is natural. Let $f: \mathcal{B} \leadsto \mathcal{A}$ be a positive map. The goal is to show that the diagram

$$\begin{array}{ccc}
\mathcal{B} & \xrightarrow{\Gamma_{\mathcal{B}}} & C(\sigma(\mathcal{B})) \\
\downarrow^{f} & & & \downarrow^{C^{\sigma(f)}} \\
\mathcal{A} & \xrightarrow{\Gamma_{\mathcal{A}}} & C(\sigma(\mathcal{A}))
\end{array} (4.168)$$

commutes, i.e. for arbitrary $b \in \mathcal{B}$, the two functions $C^{\sigma(f)}(\Gamma_B(b))$ and $\Gamma_A(f(a))$ are equal on $\sigma(A)$. To check equality of these functions, evaluating them on an arbitrary $\chi \in \sigma(A)$ gives

$$\left(C^{\sigma(f)}(\Gamma_B(b))\right)(\chi) = \int_{\sigma(\mathcal{B})} \operatorname{ev}_b d\sigma_{\chi}^f = \chi(f(b)) = \left(\Gamma_{\mathcal{A}}(f(a))\right)(\chi), \tag{4.169}$$

which is the desired result.

We should also check that this equivalence agrees with the usual Gelfand-Naimark equivalence for deterministic processes, i.e. for *-homomorphisms on the C^* -algebra side and measure-preserving functions on the analytic side.

Theorem 4.170. Let \mathcal{B} and \mathcal{A} be commutative C^* -algebras, let $f: \mathcal{B} \leadsto \mathcal{A}$ be a *-homomorphism, and let $\chi \in \sigma(\mathcal{A})$ be a character on \mathcal{A} . Then the corresponding regular probability measure σ_{χ}^f is the Dirac measure at $\chi \circ f$, i.e. $\sigma_{\chi}^f = \delta_{\chi \circ f}$. Therefore, $\sigma(f)$ uniquely determines a continuous function $\sigma(\mathcal{A}) \to \sigma(\mathcal{B})$ agreeing with the usual spectrum functor on \mathbf{cC}^* -Alg, i.e. the diagram

cHaus
$$\leftarrow \frac{\sigma}{cC} cC^*$$
-Alg^{op}

$$\downarrow \qquad \qquad \downarrow \qquad \qquad (4.171)$$
cHausStoch $\leftarrow \frac{\sigma}{\sigma} cC^*$ -AlgPos^{op}

commutes.

Proof. This follows from the fact that $\chi \circ f$ is a character on $\sigma(\mathcal{B})$ and that it immediately satisfies the conditions of Theorem 4.145 so that the measure is uniquely specified.


Remark 4.172. The full power of Theorem 4.145 is not needed to prove this last result. It follows from a simpler result due to Bauer (see Proposition 1.4 in [35]).

5 Closing discussion

We briefly discuss the broad physical scope of our results in the context of quantum mechanics and then the relationship to similar conclusions by other authors.


5.1 Categories of classical and quantum dynamics

In this paper, we have collected functors



between a variety of categories. The left face of the cube consists of dynamics on classical systems (phase space). The right face describes the algebraic counterparts in terms of observables of those classical systems (functions on phase space) and how those evolve under the dynamics. The top face describes deterministic dynamics in both space and algebraic settings. The bottom face describes stochastic dynamics in both space and algebraic settings. The front face describes finite classical systems (systems whose phase space is finite). The back face describes classical systems whose phase space is possibly infinite. The functors between the left face and the right face are all equivalences. The equivalences on the top face are due to Gelfand and Naimark while the back edge of the bottom face has been described in this work.

This cube has a natural extension to the right for arbitrary C^* -algebras. *-homomorphisms are therefore non-commutative analogues of deterministic processes while completely positive maps are non-commutative analogues of non-deterministic processes.



The theorems presented in this work offer more motivation for thinking of completely positive maps and the dynamics associated with them in quantum theory as a natural extension of stochastic dynamics in classical theory. This can be seen more concretely if one adds more data to these categories. The categories above only describe dynamics of the observables of a system, but not of individual states. In other words, we have so far described the Heisenberg picture of quantum mechanics. However, this is a red herring. Evolution of states is also described in this setting because a state is a completely positive map. If a completely positive map $f: \mathcal{B} \leadsto \mathcal{A}$ describes the evolution of observables described by \mathcal{A} to observables described by \mathcal{B} , and $\omega: \mathcal{A} \leadsto \mathbb{C}$ is a state on \mathcal{A} , then ω evolves to $\omega \circ f: \mathcal{B} \leadsto \mathbb{C}$ under the evolution described by f. Therefore, one can include both evolution of observables and states by taking the slice category of \mathbf{C}^* -AlgCP over \mathbb{C} , denoted by \mathbf{C}^* -AlgCP/ \mathbb{C} . The objects of this category are C^* -algebras equipped with states $\omega: \mathcal{A} \leadsto \mathbb{C}$. A morphism from $\xi: \mathcal{B} \leadsto \mathbb{C}$ to $\omega: \mathcal{A} \leadsto \mathbb{C}$ consists of a completely positive map $f: \mathcal{B} \leadsto \mathcal{A}$ such that $\xi = \omega \circ f$. This is a non-commutative generalization of spaces equipped with probability measures. This situation was described in Example 2.93 though more general dynamics is encompassed in Lindblad's formalism [27].

5.2 Categorical probability theory

As mentioned earlier, Sections 2 and 3 were entirely a review though phrased in a categorical setting that was used to state and prove various results in Section 4. Many of the results in Section 4 relied on well known facts in analysis though our categorical framework brought some questions to surface which seem to have not been addressed earlier. Similar results were obtained by Furber and Jacobs [15], who proved an equivalence of categories between **cHausStoch**^{op} and **cC*-AlgPos** by realizing the category **cHausStoch** as the Kleisli category of a particular monad, a slight variant of the Giry monad, on the category **cHaus**. There are three main differences between our findings.

i. Rather than constructing the Giry monad, we worked directly with the objects and morphisms of **cHausStoch** bypassing the monad entirely. This saves us quite a bit of space from explaining background material on monads, Kleisli categories, and how these are related to adjunctions, which is all beautifully explained in [42]. Furthermore, it makes our presentation more accessible to those with a minimal categorical background. However, as a result, some proofs become a bit more involved compared to [15]. For instance, given two stochastic maps $\mu: X \leadsto Y$ and $\nu: Y \leadsto Z$, proving that the composition $\nu \circ \mu$ is well-defined, continuous, regular, and a Markov kernel was quite involved (see Proposition 4.79 and the preceding lemmas). From the Giry monad perspective, ProbMeas can be viewed as a functor from the category **cHaus** to itself (one must prove this). In addition, there are two natural transformations that provide ProbMeas with the structure of a monad. The inclusion of a space X into ProbMeas(X) via the Dirac measure defines a natural transformation $\delta: \mathrm{id}_{\mathbf{cHaus}} \Rightarrow \mathrm{ProbMeas}$. In addition, there is a natural transformation $b: \mathrm{ProbMeas} \circ \mathrm{ProbMeas} \Rightarrow \mathrm{ProbMeas}$ given on a space X by the assignment

$$\operatorname{ProbMeas}(\operatorname{ProbMeas}(X)) \xrightarrow{b_X} \operatorname{ProbMeas}(X)$$

$$\Omega \mapsto \left(\mathcal{M}_X \ni E \xrightarrow{b_X(\Omega)} \int_{\operatorname{ProbMeas}(X)} \operatorname{ev}_E d\Omega \right). \tag{5.3}$$

Notice that one must still show that ev_E is measurable to define this map. b_X is surprisingly easy to interpret and has significant physical meaning. ProbMeas(X) is a convex subset of the vector space of all finite measures. A probability measure on it therefore has a barycenter, its center of mass. The map b_X is precisely the assignment of that probability measure to its barycenter. This follows from the fact that for any Borel set $E \in \mathcal{M}_X$,

$$\int_{\text{ProbMeas}(X)} \text{ev}_E d\Omega = \int_{\text{ProbMeas}(X)} \text{ev}_E(\mu) d\Omega(\mu) = \int_{\text{ProbMeas}(X)} \mu(E) d\Omega(\mu). \tag{5.4}$$

A physically relevant instance where the barycenter naturally occurs will be discussed after point iii. Going back to the Giry monad and **cHausStoch**, using the barycenter map, the definition of the composition $\nu \circ \mu$ can then be taken to be the composition of the functions

$$X \xrightarrow{\mu} \operatorname{ProbMeas}(Y) \xrightarrow{\operatorname{ProbMeas}(\nu)} \operatorname{ProbMeas}(\operatorname{ProbMeas}(Z)) \xrightarrow{bZ} \operatorname{ProbMeas}(Z)$$
 (5.5)

from which it immediately follows that the composition is continuous, since each function in this composition is continuous. The relationship between our formula for the composition and this composition of functions follows from the change of variables formula since applying the above composition to a point gives

$$X \ni x \mapsto \mu_x \mapsto \nu_* \mu_x \mapsto b_Z(\nu_* \mu_x), \tag{5.6}$$

where $\nu_*(\mu_x)$ denotes the pushforward measure of μ_x along the (measurable) function $\nu: Y \to \text{ProbMeas}(Z)$. This evaluates on measurable sets $E \in \mathcal{M}_Z$ to

$$(b_Z(\nu_*\mu_x))(E) = \int_{\text{ProbMeas}(Z)} \text{ev}_E \ d(\nu_*\mu_x) = \int_Y (\text{ev}_E \circ \nu) \ d\mu_x, \tag{5.7}$$

which agrees with our definition of composition (4.80). By using an intermediate amount of category theory, our work offers a bridge towards more categorical approaches to probability theory [19].

ii. In [15], Furber and Jacobs use the RMK Representation Theorem immediately so that a stochastic map $X \leadsto Y$ is defined to be a continuous function $X \to \mathcal{S}(C(Y))$ into states on the C^* -algebra of continuous functions on Y instead of a continuous function $X \to \operatorname{ProbMeas}(Y)$ into regular probability measures on Y (this is how the Kleisli category associated to their version of the Giry monad was defined). There is no significant difference between these perspectives. However, we prefer to think of the left side of our cube in Section 5.1 as purely analytical, topological, and measure-theoretic. In particular, we provided an explicit definition of the composition using Markov kernels and without any reference to the RMK theorem. The right face of this cube is more algebraic and utilizes algebras of function spaces, states, positive maps, etc. Of course, the point of the equivalence between these two faces is that this distinction is just psychological. Nevertheless, it is helpful to distinguish the different tools used in both subjects. As a result, many of the proofs in our paper are completely different in nature from those of Furber and Jacobs.

iii. Finally, Furber and Jacobs showed that $C: \mathbf{cHausStoch^{op}} \to \mathbf{cC^*-AlgPos}$ is fully faithful and essentially surjective [15]. However, any abstract nonsense construction of an explicit inverse will necessarily relate any commutative C^* -algebra to one of continuous functions on some space and will therefore use the RMK theorem in its construction. We have therefore provided an explicit construction of an inverse without using the Gelfand transform nor the RMK theorem. Furthermore, we showed that the usual Gelfand transform is part of the adjoint equivalence for the stochastic Gelfand-Naimark Theorem. Our construction of the stochastic spectrum functor applies to positive maps and take characters to states. Choquet theory was used to "spread" this state out onto the spectrum in terms of a probability measure which was obtained as a limit of successive approximations via decompositions of the state into convex combinations of Dirac measures supported on the space of characters. This perspective may be useful for explorations in the category of non-commutative C^* -algebras.

One instance where the barycenter map is important in physics occurs in preparing systems on which to perform experiments. In an experiment, one has a protocol for preparing a state, which will then be probed in some way, say, by measuring some observable. Because the probe may alter the state, one needs the protocol to be as accurate as possible so that the same state can be constructed so that it can be probed in subsequent experiments. Indeed, this is how a state is defined mathematically—via its expectation values. However, in practice, protocols, as well as instruments, are rarely perfect. As a result, the prepared states may differ slightly from one another so that they are more accurately represented by a probability measure on the set of states. Hence, an experimentalist is not always obtaining expectation values for the same state. Nevertheless, after viewing the set of states as probability measures via the RMK Representation Theorem, the barycenter map provides one with a canonical state that represents the average state prepared by the protocol. This is the state very often referred to in most textbooks, but one should keep in mind that the barycenter represents only one part of all the data.

5.3 The Baire approach

One may have circumvented many of the difficulties we encountered by using Baire sets instead of Borel sets. This perspective is emphasized in works such as [12]. In this case, one has the following analogue of Lemma 4.64.

Lemma 5.8. Let Y be a compact Hausdorff space, let $E \subseteq Y$ be a Baire set, and let Pr(Y) denote all probability measures on Y. Then the evaluation function

$$\Pr(Y) \xrightarrow{\operatorname{ev}_E} \mathbb{R}$$

$$\mu \mapsto \mu(E) \tag{5.9}$$

is Borel measurable.

Proof. Let P be the collection of all compact G_{δ} subsets of Y and let

$$L := \left\{ E \subseteq Y : \Pr(Y) \xrightarrow{\text{ev}_E} [0, 1] \text{ is Borel measurable} \right\}$$
 (5.10)

- i. The intersection of compact sets is compact and the finite intersection of a countable collection of open sets is still a countable intersection of open sets. Hence P is a π -system.
- ii. This is the same as in the proof of Lemma 4.64 so L is a λ -system.
- iii. Let $K = U_1 \cap U_2 \cap U_3 \cap \cdots$ be a compact G_δ set in Y with $\{U_n\}$ an at most countable collection of open sets in Y. It suffices to assume that $U_1 \supseteq U_2 \supseteq U_3 \supseteq \cdots$ since if this were not the case, the sequence $U_1, U_1 \cap U_2, U_1 \cap U_2 \cap U_3, \ldots$ would also have intersection equal to K. For each n, by Urysohn's Lemma, there exists a continuous function $f_n : Y \to [0,1]$ such that $f_n(y) = 1$ for $y \in K$ and $f_n(y) = 0$ for $y \in U_n^c$. Similarly, it suffices to assume that $f_1 \ge f_2 \ge f_3 \ge \cdots$ since if this were not the case, the sequence $\mathbb{N} \times Y \ni (n,y) \mapsto \min \{f_1(y), f_2(y), \ldots, f_n(y)\}$ would also have the same pointwise limit. From these assumptions, it follows that

$$\lim_{n \to \infty} f_n(y) = \chi_K(y) \qquad \forall \ y \in Y. \tag{5.11}$$

Hence, for any $a \in \mathbb{R}$, the preimage of (a, ∞) under ev_K is

$$\operatorname{ev}_{K}^{-1}((a,\infty)) = \left\{ \mu \in \Pr(Y) : \mu(K) > a \right\}$$

$$= \left\{ \mu \in \Pr(Y) : \lim_{n \to \infty} \int_{Y} f_{n} \, d\mu > a \right\}$$
by the Monotone Convergence Theorem
$$= \bigcap_{n=1}^{\infty} \left\{ \mu \in \Pr(Y) : \int_{Y} f_{n} \, d\mu > a \right\} \quad \text{since } \dots \geq f_{n} \geq f_{n+1} \geq \dots.$$

$$(5.12)$$

By definition of the vague topology, the set in curly brackets in the last equality is open. Hence, this is a countable intersection of open sets and is therefore a Borel subset of Pr(Y). This shows that ev_K is Borel measurable for all compact G_{δ} sets K.

Since all the conditions of Dynkin's π - λ theorem are satisfied, the sigma algebra generated by P, which is the Baire σ -algebra on Y, is contained L.

This allows one to define the composition of stochastic maps on Baire sets. One can then use the theorem that every Baire measure extends uniquely to a regular measure under these assumptions (see Corollary 7.3.4. in [8]). However, this would not guarantee that formula (4.80) for the composition of stochastic maps is still a valid mathematical expression. The construction of the regular Borel extension of this measure is somewhat formal and does not provide one with the explicit formula we have produced.

Index of notation

Notation	Name and/or description	Location	Page
FinSetFun	category of finite sets and functions	Notation 2.12	7
FinProb	category of probability measures on finite sets and measure-preserving functions	Notation 2.12	7
Pr(Y)	probability measures on a finite set Y	Definition 2.13	8
$X \leadsto Y$	stochastic map from X to Y	Definition 2.13	8
f_{yx}	yx -component of a stochastic map $f: X \leadsto Y$	Definition 2.13	8
$\delta_{yy'}$	the value of the Kronecker delta function on $(y, y') \in Y \times Y$	Equation 2.19	8
FinSetStoch	category of finite sets and stochastic maps	Notation 2.27	10
$\mathcal{A},\mathcal{B},\mathcal{C}$	an algebra (often a unital C^* -algebra)	Definition 2.34	11
$1_{\mathcal{A}}$	multiplicative unit in an algebra \mathcal{A}	Definition 2.34	11
$\ \cdot\ $	a norm or seminorm	Definition 2.35	12
iff	"if and only if" (used only in definitions)	Definition 2.35	12
\mathbb{C}^X	set of all functions from X to \mathbb{C}	Example 2.48	13
$M_n(\mathbb{C})$	set of all complex $n \times n$ matrices	Example 2.50	13
C*-Alg	category of C^* -algebras and *-homomorphisms	Definition 2.52	13
cC*-Alg	the subcategory of \mathbf{C}^* -Alg of commutative C^* -algebras and *-homomorphisms	Definition 2.52	13
fdcC*-Alg	the subcategory of \mathbf{cC}^* -Alg of finite-dimensional C^* -algebras and *-homomorphisms	Definition 2.52	13
CP map	completely positive (unital) map	Definition 2.63	15
$\mathcal{B} o \mathcal{A}$	*-homomorphism of C^* -algebras	Notation 2.66	15
$\mathcal{B} \rightsquigarrow \mathcal{A}$	completely positive map of C^* -algebras	Notation 2.66	15
C*-AlgCP	category of C^* -algebras and completely positive (CP) maps	Notation 2.67	15
cC*-AlgPos	the subcategory of C^* -AlgCP of commutative C^* -algebras and CP maps	Notation 2.67	15
fdcC*-AlgPos	the subcategory of \mathbf{cC}^* -AlgPos of finite-dimensional C^* -algebras and CP maps	Notation 2.67	15
C	functor $C : \mathbf{FinSetStoch}^{\mathrm{op}} \to \mathbf{fdcC^*-AlgPos}$	Theorem 2.72	16
	and its restriction to FinSetFun	Theorem 2.78	17
C(X)	continuous functions on X	Example 3.1	24
$C^f \equiv C(f)$	pullback $C(Y) \to C(X)$ associated to $f: X \to Y$	Proposition 3.3	24
cHaus	category of compact Hausdorff spaces and continuous maps	Notation 3.6	24
C	functor $\mathbf{cHaus}^{\mathrm{op}} \to \mathbf{cC^*} ext{-}\mathbf{Alg}$	Proposition 3.7	24
\mathfrak{B}	base on a set (for a topology)	Definition 3.13	25

\overline{n}	the set $\{1, 2, \dots, n\}$ with $n \in \mathbb{N}$	Notation 3.18	26
$\ \cdot\ _{lpha}$	seminorm indexed by α in some set	Proposition 3.26	27
$\overline{V^{\vee}}$	algebraic (linear) dual of vector space V	Proposition 3.23	27
V^*	topological dual of normed space V	Proposition 3.23	27
Θ	ordered index set for nets	Proposition 3.32	28
$\sigma(\mathcal{A})$	spectrum of commutative Banach algebra ${\cal A}$	Definition 3.38	29
$\sigma(f) \equiv \sigma^f$	spectrum of a homomorphism f	Proposition 3.43	29
σ	spectrum functor $\mathbf{cC^*} ext{-}\mathbf{Alg}^{\mathrm{op}} \to \mathbf{cHaus}$	Proposition 3.50	30
Γ	Gelfand transform $\Gamma : \mathrm{id}_{\mathbf{c}\mathbf{C}^*\text{-}\mathbf{Alg}} \Rightarrow C \circ \sigma$	Proposition 3.59	32
h	natural transformation $h: \mathrm{id}_{\mathbf{cHaus}} \Rightarrow \sigma \circ C$	Proposition 3.66	32
(Y, \mathcal{M}_Y)	compact Hausdorff space Y with its Borel σ -algebra \mathcal{M}_Y	Section 4.1	36
Ø	the empty set	Definition 4.2	36
μ, ν	measure (positive, real, or complex)	Definition 4.2	36
$\mathbb{C}\mathrm{Meas}(Y)$	set of complex measures on Y	Definition 4.2	36
χ_E	characteristic/indicator function on E	Definition 4.6	36
s	simple function	Definition 4.6	36
$\int \cdot d\mu$	integral with respect to a measure μ	Definition 4.6	36
$\overline{\text{ProbMeas}(Y)}$	set of regular probability measures on Y (equipped with the vague topology)	Notation 4.47	42
$\mu: X \leadsto Y$	stochastic map described by a continuous function $\mu: X \to \operatorname{ProbMeas}(Y)$	Definition 4.57	44
δ_f	Dirac delta stochastic map associated to a continuous function f	Example 4.58	44
$\nu \circ \mu$	composition of $\mu: X \leadsto Y$ followed by $\nu: Y \leadsto Z$	Proposition 4.79	48
cHausStoch	category of compact Hausdorff spaces and stochastic maps	Theorem 4.102	51
δ	Dirac measure functor $\delta : \mathbf{cHaus} \to \mathbf{cHausStoch}$	Theorem 4.102	51
C	$\text{functor } \mathbf{cHausStoch}^{\text{op}} \to \mathbf{cC^*\text{-}AlgPos}$	Theorem 4.106	52
RMK	short for "Riesz-Markov-Kakutani"	Theorem 4.112	53
ex(C)	extreme points of a convex set C	Definition 4.127	56
$\mathcal{S}(\mathcal{A})$	states on commutative C^* -algebra $\mathcal A$	Example 4.129	56
$\operatorname{ch}(X)$	convex hull of X	Definition 4.130	56
σ	${f stochastic\ spectrum\ functor} \ {f cC^*-AlgPos}^{op} ightarrow {f cHausStoch}$	Theorem 4.151	60

References

- [1] Erik M. Alfsen, Compact convex sets and boundary integrals, Ergebnisse der Mathematik und ihrer Grenzgebiete. 2. Folge, Springer-Verlag Berlin Heidelberg, 1971. ↑58
- [2] John C. Baez and Tobias Fritz, A bayesian characterization of relative entropy, Theory and Applications of Categories 29 (2014), no. 16, 422–456, available at 1402.3067. ↑5, 8
- [3] John C. Baez, Tobias Fritz, and Tom Leinster, A characterization of entropy in terms of information loss, Entropy 13 (2011), 1945–1957, available at 1106.1791. ↑8
- [4] John C. Baez and Mike Stay, *Physics, topology, logic and computation: a Rosetta Stone*, New structures for physics, 2011, pp. 95–172. ↑2
- [5] Patrick Billingsley, The invariance principle for dependent random variables, Trans. Amer. Math. Soc. 83 (1956), 250–268. ↑4
- [6] ______, Probability and measure, 3rd ed., New York: John Wiley & Sons, Inc., 1995. \dagger46
- [7] Vladimir I. Bogachev, Measure theory, Vol. 1, Springer-Verlag Berlin Heidelberg, 2007. †63
- [8] ______, Measure theory, Vol. 2, Springer-Verlag Berlin Heidelberg, 2007. \dagger69
- [9] Fabio Cipriani, Positive maps on c*-algebras, Encyclopedia of Mathematical Physics 3 (2006), 88–94. ↑2, 35
- [10] Kenneth R. Davidson, C^* -algebras by example, Fields Institute Monographs, vol. 6, American Mathematical Society, Providence, RI, 1996. \uparrow 13
- [11] Jacques Dixmier, C*-algebras, North-Holland Publishing Co., Amsterdam-New York-Oxford, 1977. Translated from the French by Francis Jellett, North-Holland Mathematical Library, Vol. 15. ↑11, 12, 13
- [12] Tanja Eisner, Bálint Farkas, Markus Haase, and Rainer Nagel, Operator theoretic aspects of ergodic theory, Graduate Texts in Mathematics, vol. 272, Springer, Cham, 2015. ↑2, 4, 35, 68
- [13] Gerald B. Folland, A course in abstract harmonic analysis, Textbooks in Mathematics, CRC Press, 1994. ↑23, 29, 31, 32
- [14] _____, Real analysis: Modern techniques and their applications, 2nd ed., Wiley, 2007. \dagger26, 27, 28, 35, 42, 43, 54
- [15] Robert Furber and Bart Jacobs, From kleisli categories to commutative C*-algebras: probabilistic gelfand duality, Logical Methods in Computer Science 11 (2015), 1–28. ↑3, 4, 35, 66, 67, 68
- [16] Alex Furman, What is ... a stationary measure?, Notices Amer. Math. Soc. 58 (2011), no. 9, 1276–1277. \\$\dagger35\$
- [17] Nicolas Gagné and Prakash Panangaden, A categorical characterization of relative entropy on polish spaces (2017), available at 1703.08853. ↑8
- [18] Israel Gelfand and Mark Neumark, On the imbedding of normed rings into the ring of operators in Hilbert space, Rec. Math. [Mat. Sbornik] N.S. 12(54) (1943), 197–213. †2, 33
- [19] Michèle Giry, A categorical approach to probability theory, Categorical aspects of topology and analysis (Ottawa, Ont., 1980), 1982, pp. 68–85. †3, 4, 11, 35, 45, 67
- [20] Brian C. Hall, Quantum theory for mathematicians, Graduate Texts in Mathematics, vol. 267, Springer, New York, 2013. ↑14
- [21] Chris Heunen, The many classical faces of quantum structures, Entropy 19 (2017), no. 4, 144, available at 1412.2177. ↑5, 35
- [22] Richard V. Kadison, A representation theory for commutative topological algebra, Mem. Amer. Math. Soc., No. 7 (1951), 39. ↑23

- [23] Shizuo Kakutani, Concrete representation of abstract (M)-spaces. (A characterization of the space of continuous functions.), Ann. of Math. (2) 42 (1941), 994–1024. †2, 23, 35
- [24] Dexter Kozen, Semantics of probabilistic programs, J. Comput. System Sci. 22 (1981), no. 3, 328–350. Special issue dedicated to Michael Machtey. ↑3
- [25] Karl Kraus, States, effects, and operations: Fundamental notions of quantum theory, Lecture Notes in Physics Vol. 190, Springer Berlin Heidelberg, 1983. †14, 15
- [26] William F. Lawvere, The category of probabilistic mappings, 1962. preprint. ↑3
- [27] Göran Lindblad, On the generators of quantum dynamical semigroups, Comm. Math. Phys. 48 (1976), no. 2, 119–130. ↑2, 15, 23, 66
- [28] Fred E. J. Linton, *The functorial foundations of measure theory*, Columbia University, 1963. dissertation thesis. ↑3
- [29] James R. Munkres, Topology: a first course, 2nd ed., Prentice-Hall, Inc., 2000. ↑24
- [30] Michael A. Nielsen and Isaac L. Chuang, Quantum computation and quantum information, 10th Anniversary, Cambridge University Press, Cambridge, 2010. ↑15, 22, 23
- [31] Prakash Panangaden, *The category of Markov kernels*, PROBMIV'98: First International Workshop on Probabilistic Methods in Verification (Indianapolis, IN), 1999, pp. 17. ↑3
- [32] Arthur J. Parzygnat, From observables and states to hilbert space and back: a 2-categorical adjunction (2016), available at 1609.08975. ↑14
- [33] Vern Paulsen, Completely bounded maps and operator algebras, Cambridge Studies in Advanced Mathematics, Cambridge University Press, 2003. ↑15
- [34] Gert K. Pedersen, Analysis now, Graduate Texts in Mathematics, vol. 118, Springer, 1989. ²⁷
- [35] Robert R. Phelps, *Lectures on choquet's theorem*, 2nd ed., Lecture Notes in Mathematics, vol. 1757, Springer-Verlag Berlin Heidelberg, 2001. ↑35, 56, 58, 59, 64
- [36] Miklós Rédei and Stephen J. Summers, *Quantum probability theory*, Studies in History and Philosophy of Modern Physics **38** (2007), 390–417, available at quant−ph/0601158. ↑14
- [37] Walter Rudin, *Principles of mathematical analysis*, Third, McGraw-Hill Book Co., New York-Auckland-Düsseldorf, 1976. International Series in Pure and Applied Mathematics. ↑24
- [38] ______, Real and complex analysis, 3rd ed., McGraw-Hill Inc., 1987. †36, 37, 38, 39, 42, 54, 58
- [39] _____, Functional analysis, 2nd ed., International Series in Pure and Applied Mathematics, McGraw-Hill Inc., 1991. ↑23, 28, 29, 56, 58
- [40] Shoichiro Sakai, c^* -algebras and w^* -algebras, 1st ed., Ergebnisse der Mathematik und ihrer Grenzgebiete, Springer-Verlag Berlin Heidelberg, 1971. $\uparrow 58$
- [41] Jun John Sakurai, Modern Quantum Mechanics; revised ed., Addison-Wesley, 1993. †21
- [42] Zbigniew Semadeni, Monads and their eilenberg-moore algebras in functional analysis, Queen's Papers in Pure and Applied Mathematics, Issue 33, Queen's University, 1973. ↑3, 66
- [43] Ambar Sengupta, Lecture 6. the dynkin π-λ theorem, Louisiana State University, 2009. Available as: https://www.math.lsu.edu/~sengupta/7360f09/DynkinPiLambda.pdf. ↑45, 46
- [44] Barry Simon, Convexity: An analytic viewpoint, 1st ed., Cambridge Tracts in Mathematics (Book 187), Cambridge University Press, 2011. ↑56, 58
- [45] Lynn Arthur Steen and J. Arthur Seebach Jr., Counterexamples in topology, Dover Publications, Inc., Mineola, NY, 1995. Reprint of the second (1978) edition. ↑42

- [46] Tadeusz Świrszcz, Monadic functors and convexity, Bull. Acad. Polon. Sci. Sér. Sci. Math. Astronom. Phys. 22 (1974), 39–42. ↑3
- [47] Terence Tao, 245b, notes 11: The strong and weak topologies, 2009. Available as: https://terrytao.wordpress.com/2009/02/21/245b-notes-11-the-strong-and-weak-topologies/. ↑23, 25, 26, 27, 28
- [48] ______, 245b, notes 12: Continuous functions on locally compact hausdorff spaces, 2009. Available as: https://terrytao.wordpress.com/2009/03/02/245b-notes-12-continuous-functions-on-locally-compact-hausd ↑54
- [49] Garth Warner, C^* -algebras, 2010. $\uparrow 15$
- [50] Stephen Willard, General topology, Dover Books on Mathematics, Dover Publications, 2004. †26, 28

Mathematics Department, University of Connecticut, 341 Mansfield Road U1009, Storrs, CT 06269, USA

Email: arthur.parzygnat@uconn.edu