Non-leptonic two-body weak decays of $\Lambda_c(2286)$

C.Q. Geng,^{1,2} Y.K. Hsiao,^{1,2} Yu-Heng Lin,² and Liang-Liang Liu^1

¹School of Physics and Information Engineering, Shanxi Normal University, Linfen 041004, China

²Department of Physics, National Tsing Hua University, Hsinchu, Taiwan 300

(Dated: July 29, 2018)

Abstract

We study the non-leptonic two-body weak decays of $\Lambda_c^+(2286) \to \mathbf{B}_n M$ with $\mathbf{B}_n(M)$ representing as the baryon (meson) states. Based on the SU(3) flavor symmetry, we can describe most of the data reexamined by the BESIII Collaboration with higher precisions. However, our result of $\mathcal{B}(\Lambda_c^+ \to p\pi^0) = (5.6 \pm 1.5) \times 10^{-4}$ is larger than the current experimental limit of 3×10^{-4} (90% C.L.) by BESIII. In addition, we find that $\mathcal{B}(\Lambda_c^+ \to \Sigma^+ K^0) = (8.0 \pm 1.6) \times 10^{-4}$, $\mathcal{B}(\Lambda_c^+ \to \Sigma^+ \eta') = (1.0^{+1.6}_{-0.8}) \times 10^{-2}$, and $\mathcal{B}(\Lambda_c^+ \to p\eta') = (12.2^{+14.3}_{-8.7}) \times 10^{-4}$, which are accessible to the BESIII experiments.

I. INTRODUCTION

Recently, the BESIII Collaboration has reanalyzed the two-body weak decays of $\Lambda_c^+(2286)$ with the final states to be the combinations of baryon (\mathbf{B}_n) and pseudoscalar meson (M) particles, where $\Lambda_c^+ \equiv \Lambda_c^+(2286)$ along with $\Xi_c^{+,0}(2470)$ belongs to the lowest-lying antitriplet charmed baryon (\mathbf{B}_c) state. In particular, the decay branching ratios of $\Lambda_c^+ \rightarrow p\bar{K}^0, \Lambda\pi^+, \Sigma^+\pi^0$ and $\Sigma^0\pi^+$ have been measured at the level of 10^{-2} with high precisions [1]. In addition, the Cabibbo-suppressed $\Lambda_c^+ \rightarrow p\eta$ decay has been observed for the first time [2]. According to the measurements of the two-body $\Lambda_c^+ \rightarrow \mathbf{B}_n M$ decays since 2016 [1], there have been 4 measured branching fractions listed in PDG [3], given as

$$\mathcal{B}(\Lambda_c^+ \to p\bar{K}^0) = (3.16 \pm 0.16)\%,$$

$$\mathcal{B}(\Lambda_c^+ \to \Lambda \pi^+) = (1.30 \pm 0.07)\%,$$

$$\mathcal{B}(\Lambda_c^+ \to \Sigma^+ \pi^0) = (1.24 \pm 0.10)\%,$$

$$\mathcal{B}(\Lambda_c^+ \to \Sigma^0 \pi^+) = (1.29 \pm 0.07)\%,$$

(1)

together with the new data [2], given by

$$\mathcal{B}(\Lambda_c^+ \to p\eta) = (1.24 \pm 0.28 \pm 0.10) \times 10^{-3},$$

$$\mathcal{B}(\Lambda_c^+ \to p\pi^0) < 3 \times 10^{-4} (90\% \text{ C.L.}).$$
(2)

Note that the limit of $\mathcal{B}(\Lambda_c^+ \to p\pi^0)$ in Eq. (2) comes from the original data of $\mathcal{B}(\Lambda_c^+ \to p\pi^0) = (7.95 \pm 13.61) \times 10^{-5}$ [4] by BESIII, while the $\Lambda_c^+ \to \Sigma^+ K^0, p\eta'$ and $\Sigma^+ \eta'$ decays, along with the neutron modes, have not been seen yet. It is interesting to see if these current data can be understood.

Theoretically, the factorization approach is demonstrated to well explain the B and bbaryon decays [5–7], such that it is also applied to the two-body $\Lambda_c^+ \to \mathbf{B}_n M$ decays [8], of which the amplitudes are derived as the combination of the two computable matrix elements for the $\Lambda_c^+ \to \mathbf{B}_n$ transition and the meson (M) production. However, the factorization approach does not work for most of the two-body $\Lambda_c^+ \to \mathbf{B}_n M$ ones. For example, the decays of $\Lambda_c^+ \to \Sigma^+ \pi^0$ and $\Xi^0 K^+$ are forbidden in the factorization approach [9], but their branching ratios turn out to be measured. As a result, several theoretical attempts to improve the factorization by taking into account the nonfactorizable effects have been made [10–14]. In contrast with the QCD-based models, the SU(3) symmetry approach is independent of the



FIG. 1. Feynman diagrams for the c quark decays, where (a,b,c) are the tree-level processes with the W-boson emissions to directly connect to quark pairs, where q = (d, s), while (d) corresponds to the penguin-level ones with the W-boson in the loop.

detailed dynamics, which has been widely used in the *B* meson [15–17], *b*-baryon [18, 19] and Λ_c^+ (Ξ_c) [9, 20–23] decays. With this advantage, the two-body $\Lambda_c^+ \to \mathbf{B}_n M$ decays can be related by the *SU*(3) parameters, which receive possible non-perturbative and nonfactorizable contributions [9, 20–24], despite of the unknown sources. The minimum χ^2 fit with the p-value estimation [3] can statistically test if the *SU*(3) flavor symmetry agrees with the data. Being determined from the fitting also, the *SU*(3) parameters are taken to predict the not-yet-measured modes for the future experimental tests. However, the global fit was once unachievable without the sufficient data and the use of the symmetry for $\Lambda_c^+ \to \mathbf{B}_n M$. Clearly, the reexamination with the global fit to match the currently more accurate data is needed. Note that, to study the $\Lambda_c^+ \to \mathbf{B}_n \eta^{(\prime)}$ decays, the singlet state of η_1 should be included [16, 17]. In this report, we will extract the *SU*(3) parameters in the global fit, and predict the branching fractions to be compared with the future BESIII experimental measurements.

II. FORMALISM

From Fig. 1, there are four types of diagrams for the non-leptonic charm quark decays, where Figs. 1a-1c with the W-boson emissions directly connected to quark pairs are the so-called tree-level processes, while Fig. 1d with the W-boson in the loop corresponds to the penguin-level ones. In Fig. 1c, the $c \rightarrow du\bar{s}$ transition that proceeds through $|V_{cd}V_{us}| \simeq$ $\sin^2 \theta_c$ is the doubly Cabibbo-suppressed one, with θ_c being the well-known Cabibbo angle. Meanwhile, the $c \rightarrow uq\bar{q}(s\bar{s})$ transitions in Fig. 1d have the higher-order contributions from the quark loops, with the effective Wilson coefficients [25] calculated to be smaller than the tree-level ones by one order of magnitude. As a result, the decay processes in Figs. 1c and 1d are both excluded in the present study. Accordingly, the effective Hamiltonian for the $c \rightarrow su\bar{d}$ and $c \rightarrow uq\bar{q}$ transitions with q = (d, s) in Figs. 1a and 1b, respectively, is given by [26]

$$\mathcal{H}_{eff} = \frac{G_F}{\sqrt{2}} \Big\{ V_{cs} V_{ud} [c_1 O_1 + c_2 O_2] + \sum_{q=d,s} V_{cq} V_{uq} [c_1 O_1^q + c_2 O_2^q] \Big\},$$
(3)

with the current-current operators $O_{1,2}^{(q)}$, written as

$$O_{1} = (\bar{u}d)_{V-A}(\bar{s}c)_{V-A}, \quad O_{2} = (\bar{s}d)_{V-A}(\bar{u}c)_{V-A},$$
$$O_{1}^{q} = (\bar{u}q)_{V-A}(\bar{q}c)_{V-A}, \quad O_{2}^{q} = (\bar{q}q)_{V-A}(\bar{u}c)_{V-A}, \quad (4)$$

where G_F is the Fermi constant, V_{ij} are the Cabibbo-Kobayashi-Maskawa (CKM) matrix elements, and $(\bar{q}_1q_2)_{V-A}$ stands for $\bar{q}_1\gamma_{\mu}(1-\gamma_5)q_2$. The operators $O_{1,2}$ and $O_{1,2}^q$ in Eq. (3) lead to the so-called Cabibbo-allowed and Cabibbo-suppressed decay modes due to the factor of $|(V_{cq}V_{uq})/(V_{cs}V_{ud})| = \sin\theta_c$. The Wilson coefficients $c_{1,2}$ in Eq. (3) are scale-dependent. In the NDR scheme [25, 27], one has that $(c_1, c_2) = (1.27, -0.51)$ at the scale $\mu = 1$ GeV. Note that one is able to recombine $V_{cs}V_{ud}[c_1O_1 + c_2O_2]$ and $\sum_{q=d,s}V_{cq}V_{uq}[c_1O_1^q + c_2O_2^q]$ in Eq. (3) into $V_{cs}V_{ud}[c_+O_+ + c_-O_-]$ and $V_{cd}V_{ud}[c_+\hat{O}_+ + c_-\hat{O}_-]$ with $\hat{O}_{\pm} \equiv O_{\pm}^d - O_{\pm}^s$, respectively, where $c_{\pm} = c_1 \pm c_2, O_{\pm}^{(q)} = (O_1^{(q)} \pm O_2^{(q)})/2$ and $V_{cs}V_{us} = -V_{cd}V_{ud}$.

For the four-quark operator $(\bar{q}^i q_k)(\bar{q}^j c)$ from the effective Hamiltonian in Eq. (4), $\bar{q}^i q_k \bar{q}^j$ that belongs to the SU(3) triplet of $q_i = (u, d, s)$ can be decomposed as the irreducible forms of $\bar{3} \times 3 \times \bar{3} = \bar{3} + \bar{3}' + 6 + \bar{15}$, which are in terms of the SU(3) flavor symmetry with the Lorentz-Dirac structures being disregarded. Consequently, the Cabibbo-allowed operators O_- and O_+ fall into 6 and $\bar{15}$, respectively, instead of $\bar{3} + \bar{3}'$ that actually appear in the penguin operators. Therefore, in the SU(3) picture the Cabibbo-allowed operators O_- and O_+ are presented as [20, 21]

$$\mathcal{O}_{6} = \frac{1}{2} [(\bar{u}d)(\bar{s}c) - (\bar{s}d)(\bar{u}c)],$$

$$\mathcal{O}_{\overline{15}} = \frac{1}{2} [(\bar{u}d)(\bar{s}c) + (\bar{s}d)(\bar{u}c)],$$
 (5)

which are formed as the tensor notations of $H(6)^{ij}$ and $H(\overline{15})^i_{jk}$, respectively. Note that the Cabibbo-suppressed operators \hat{O}_- and \hat{O}_+ have similar irreducible forms, leading to their own $\hat{H}(6)^{ij}$ and $\hat{H}(\overline{15})^i_{jk}$ [20, 21]. As a result, the effective Hamiltonian in Eq. (3) under the SU(3) representation becomes

$$\mathcal{H}_{eff} = \frac{G_F}{\sqrt{2}} \Big\{ V_{cs} V_{ud} [c_- H(6) + c_+ H(\overline{15})] + V_{cd} V_{ud} [c_- \hat{H}(6) + c_+ \hat{H}(\overline{15})] \Big\}, \tag{6}$$

where the non-zero entries are

$$H^{22}(6) = 2, H^{2}_{13}(\overline{15}) = H^{2}_{31}(\overline{15}) = 1,$$

$$\hat{H}^{23}(6) = \hat{H}^{32}(6) = -2,$$

$$\hat{H}^{2}_{12}(\overline{15}) = \hat{H}^{2}_{21}(\overline{15}) = -\hat{H}^{3}_{13}(\overline{15}) = -\hat{H}^{3}_{31}(\overline{15}) = 1.$$
(7)

To proceed, we take the amplitudes of $\mathbf{B}_c \to \mathbf{B}_n M$ under the SU(3) representations. First, the \mathbf{B}_c state acts as $\bar{3}$ under the SU(3) flavor symmetry, written as

$$(\mathbf{B}_c)^i = (\Xi_c^0, -\Xi_c^+, \Lambda_c^+), \qquad (8)$$

by which one defines $T_{ij} = \epsilon_{ijk} (\mathbf{B}_c)^k$. Second, \mathbf{B}_n is the baryon octet, given by

$$(\mathbf{B}_{n})_{j}^{i} = \begin{pmatrix} \frac{1}{\sqrt{6}}\Lambda + \frac{1}{\sqrt{2}}\Sigma^{0} & \Sigma^{+} & p \\ \Sigma^{-} & \frac{1}{\sqrt{6}}\Lambda - \frac{1}{\sqrt{2}}\Sigma^{0} & n \\ \Xi^{-} & \Xi^{0} & -\sqrt{\frac{2}{3}}\Lambda \end{pmatrix} .$$
 (9)

To include the octet (π, K, η_8) and singlet η_1, M is presented as the nonet, given by

$$(M)_{j}^{i} = \begin{pmatrix} \frac{1}{\sqrt{2}}(\pi^{0} + \cos\phi\eta + \sin\phi\eta') & \pi^{-} & K^{-} \\ \pi^{+} & \frac{-1}{\sqrt{2}}(\pi^{0} - \cos\phi\eta - \sin\phi\eta') & \bar{K}^{0} \\ K^{+} & K^{0} & -\sin\phi\eta + \cos\phi\eta' \end{pmatrix}, (10)$$

where (η, η') are the mixtures of (η_1, η_8) , decomposed as $\eta_1 = \sqrt{2/3}\eta_q + \sqrt{1/3}\eta_s$ and $\eta_8 = \sqrt{1/3}\eta_q - \sqrt{2/3}\eta_s$ with $\eta_q = \sqrt{1/2}(u\bar{u} + d\bar{d})$ and $\eta_s = s\bar{s}$. Explicitly, the $\eta - \eta'$ mixing matrix is given by [28]

$$\begin{pmatrix} \eta \\ \eta' \end{pmatrix} = \begin{pmatrix} \cos \phi & -\sin \phi \\ \sin \phi & \cos \phi \end{pmatrix} \begin{pmatrix} \eta_q \\ \eta_s \end{pmatrix}, \tag{11}$$

with the mixing angle $\phi = (39.3 \pm 1.0)^{\circ}$. Subsequently, the amplitude of $\mathbf{B}_c \to \mathbf{B}_n M$ is derived as

$$\mathcal{A}(\mathbf{B}_{c} \to \mathbf{B}_{n}M) = \langle \mathbf{B}_{n}M | H_{eff} | \mathbf{B}_{c} \rangle$$
$$= \frac{G_{F}}{\sqrt{2}} [V_{cs}V_{ud}T(\mathbf{B}_{c} \to \mathbf{B}_{n}M) + V_{cd}V_{ud}\hat{T}(\mathbf{B}_{c} \to \mathbf{B}_{n}M)], \qquad (12)$$

where $T(\mathbf{B}_c \to \mathbf{B}_n M) = T(\mathcal{O}_{\overline{15}}) + T(\mathcal{O}_6)$ are given by [9]

$$T(\mathcal{O}_{\overline{15}}) = aH^i_{jk}(\overline{15})(\mathbf{B}_c)^j(\mathbf{B}_n)^k_l(M)^l_i + bH^i_{jk}(\overline{15})(\mathbf{B}_c)^j(M)^k_l(\mathbf{B}_n)^l_i$$

$$+ cH_{jk}^{i}(\overline{15})(\mathbf{B}_{n})_{l}^{j}(M)_{i}^{k}(\mathbf{B}_{c})^{l} + dH_{jk}^{i}(\overline{15})(M)_{l}^{j}(\mathbf{B}_{n})_{i}^{k}(\mathbf{B}_{c})^{l} + h_{1}H_{jk}^{i}(\overline{15})(\mathbf{B}_{n})_{i}^{k}(M)_{l}^{l}(\mathbf{B}_{c})^{j}, T(\mathcal{O}_{6}) = eH^{ij}(6)T_{ik}(\mathbf{B}_{n})_{l}^{k}(M)_{j}^{l} + fH^{ij}(6)T_{ik}(M)_{l}^{k}(\mathbf{B}_{n})_{j}^{l} + gH^{ij}(6)(\mathbf{B}_{n})_{i}^{k}(M)_{j}^{l}T_{kl} + h_{2}H^{ij}(6)T_{ik}(\mathbf{B}_{n})_{j}^{k}(M)_{l}^{l},$$
(13)

with (c_+, c_-) absorbed in the SU(3) parameters of (a, b, c, d, h_1) and (e, f, g, h_2) , respectively, while $\hat{T}(\mathbf{B}_c \to \mathbf{B}_n M)$ is given by replacing $H(6, \overline{15})$ in $T(\mathbf{B}_c \to \mathbf{B}_n M)$ with $\hat{H}(6, \overline{15})$, respectively. Since the amplitudes are derived from the effective Hamiltonian in Eq. (3), where the $c \to su\bar{d}$ and $c \to uq\bar{q}$ transitions are the tree-level processes, $T(\mathbf{B}_c \to \mathbf{B}_n M)$ and $\hat{T}(\mathbf{B}_c \to \mathbf{B}_n M)$ are named as the tree-level (T) amplitudes. In Eq. (13), the expansions of $T(\mathbf{B}_c \to \mathbf{B}_n M)$ and $\hat{T}(\mathbf{B}_c \to \mathbf{B}_n M)$ are shown in Table I. Note that, although we

$\frac{1}{1} \frac{1}{1} \frac{1}$								
Decay modes	$T(\mathcal{O}_{\overline{15}})$	$T(\mathcal{O}_6)$						
$T(\Lambda_c^+ \to p\bar{K}^0)$	a+c	-2e						
$T(\Lambda_c^+ \to \Lambda \pi^+)$	$\sqrt{\frac{1}{6}}(a+b-2c)$	$-\sqrt{\frac{2}{3}}(e+f+g)$						
$T(\Lambda_c^+\to \Sigma^+\pi^0)$	$-\sqrt{\frac{1}{2}}(a-b)$	$\sqrt{2}(e-f-g)$						
$T(\Lambda_c^+\to \Sigma^0\pi^+)$	$\sqrt{\frac{1}{2}}(a-b)$	$-\sqrt{2}(e-f-g)$						
$T(\Lambda_c^+\to \Xi^0 K^+)$	b+d	-2f						
$\hat{T}(\Lambda_c^+ \to p\pi^0)$	$-\sqrt{\frac{1}{2}}(b+c)$	$\sqrt{2}(f+g)$						
$\hat{T}(\Lambda_c^+\to\Lambda K^+)$	$\sqrt{\frac{1}{6}}(-a+2b+2c+3d)$	$\sqrt{\tfrac{2}{3}}(e-2f+g)$						
$\hat{T}(\Lambda_c^+\to \Sigma^0 K^+)$	$-\sqrt{\frac{1}{2}}(a+d)$	$\sqrt{2}(e-g)$						
$\hat{T}(\Lambda_c^+\to \Sigma^+ K^0)$	(-a+d)	2(e-g)						
$T(\Lambda_c^+\to \Sigma^+\eta)$	$-d\sin\phi + \sqrt{\frac{1}{2}}(a+b)\cos\phi$	$-\sqrt{2}[e+(f-g)]\cos\phi$						
	$+h_1(\sqrt{2}\cos\phi-\sin\phi)$	$-2h_2(\sqrt{2}\cos\phi-\sin\phi)$						
$T(\Lambda_c^+\to \Sigma^+\eta')$	$d\cos\phi + \sqrt{\frac{1}{2}}(a+b)\sin\phi$	$-\sqrt{2}[e+(f-g)]\sin\phi$						
	$+h_1(\cos\phi+\sqrt{2}\sin\phi)$	$-2h_2(\sqrt{2}\sin\phi + \cos\phi)$						
$\hat{T}(\Lambda_c^+ \to p\eta)$	$-(a+c+d)\sin\phi + \sqrt{\frac{1}{2}}(b-c)\cos\phi$	$-\sqrt{2}[\sqrt{2}e\sin\phi - (f-g)\cos\phi]$						
	$+h_1(\sqrt{2}\cos\phi-\sin\phi)$	$+2h_2(\sqrt{2}\cos\phi-\sin\phi)$						
$\hat{T}(\Lambda_c^+ \to p\eta')$	$(a+c+d)\sin\phi + \sqrt{\frac{1}{2}}(b-c)\cos\phi$	$\sqrt{2}[\sqrt{2}e\cos\phi + (f-g)\sin\phi]$						
	$+h_1(\cos\phi+\sqrt{2}\sin\phi)$	$+2h_2(\sqrt{2}\sin\phi+\cos\phi)$						

TABLE I. The tree-level amplitudes for the $\Lambda_c^+ \to \mathbf{B}_n M$ decays

follow the approach in Ref. [21], the $h_{1,2}$ terms are newly added for the singlet η_1 . Due to $c_-/c_+ \simeq 2.4$, the contribution of $\mathcal{O}_6(\hat{\mathcal{O}}_6)$ to the decay branching ratio can be 5.5 times larger than that of $\mathcal{O}_{\overline{15}}(\hat{\mathcal{O}}_{\overline{15}})$, such that $\mathcal{O}_{\overline{15}}(\hat{\mathcal{O}}_{\overline{15}})$ is negligible. However, we will examine if the reduction is reasonable, in case the interferences between $\mathcal{O}_6(\hat{\mathcal{O}}_6)$ and $\mathcal{O}_{\overline{15}}(\hat{\mathcal{O}}_{\overline{15}})$ can be sizable. Subsequently, we only keep the SU(3) parameters e, f, g and h_2 from \mathcal{O}_6 to simplify the amplitudes. Since e, f, g and h_2 are complex numbers, we have 7 real independent parameters to be determined by the data, given by

$$e, fe^{i\delta_f}, ge^{i\delta_g}, h_2 e^{i\delta_{h_2}}, \tag{14}$$

where e is set to be real, while an overall phase can be removed without losing generality. To calculate the decay widths, we use [3]:

$$\Gamma(\Lambda_c \to \mathbf{B}_n M) = \frac{|\vec{p}_{cm}|}{8\pi m_{\Lambda_c}^2} |\mathcal{A}(\Lambda_c \to \mathbf{B}_n M)|^2, \qquad (15)$$

where $|\vec{p}_{cm}| = \sqrt{[(m_{\Lambda_c}^2 - (m_{\mathbf{B}_n} + m_M)^2][(m_{\Lambda_c}^2 - (m_{\mathbf{B}_n} - m_M)^2]/(2m_{\Lambda_c}))}$, with the integratedover variables of the phase spaces in the two-body decays.

III. NUMERICAL RESULTS AND DISCUSSIONS

For the numerical analysis, we use the minimum χ^2 fit to find the SU(3) parameters in Eq. (14). The theoretical inputs for the CKM matrix elements are given by [3]

$$(V_{cs}, V_{ud}, V_{us}, V_{cd}) = (1 - \lambda^2/2, 1 - \lambda^2/2, \lambda, -\lambda), \qquad (16)$$

with $\lambda = 0.225$ in the Wolfenstein parameterization. There are 9 branching ratios of $\Lambda_c \rightarrow B_n M$, which are the data inputs, given in the last column of Table II. The equation of the χ^2 fit is given by

$$\chi^2 = \sum_{i=1}^{9} \left(\frac{\mathcal{B}_{th}^i - \mathcal{B}_{ex}^i}{\sigma_{ex}^i} \right)^2, \tag{17}$$

where \mathcal{B}_{th}^{i} and \mathcal{B}_{ex}^{i} stand for the branching ratios from the theoretical SU(3) amplitudes in Table I and experimental data inputs in Table II, with σ_{ex}^{i} as the 1σ experimental errors, while i = 1, 2, ..., 9 denote the 9 observed decay modes involved in the global fit, respectively. Consequently, we obtain

$$(e, f, g, h_2) = (0.257 \pm 0.006, 0.121 \pm 0.015, 0.092 \pm 0.021, 0.111 \pm 0.081) \,\text{GeV}^3,$$
$$(\delta_f, \delta_g, \delta_{h_2}) = (79.0 \pm 6.8, 35.2 \pm 8.8, 102.4 \pm 29.8)^\circ,$$
$$\chi^2/d.o.f = 2.4,$$
(18)

where d.o.f stands for the degrees of freedom. The statistical p-value to be smaller than 0.05 will show the inconsistency between the theory and data [3], which is equivalent to $\chi^2/d.o.f > 3$ here. In our case, the value of $\chi^2/d.o.f = 2.4$ indicates a tolerable result to accommodate the current data of $\mathcal{B}(\Lambda_c \to \mathbf{B}_n M)$ under the SU(3) flavor symmetry, where the contributions of $\mathcal{O}_{\overline{15}}(\hat{\mathcal{O}}_{\overline{15}})$ and the broken effects of SU(3) are both neglected. Explicitly, it is found that the largest contributions to χ^2 are from $\mathcal{B}(\Lambda_c^+ \to \Lambda K^+, \Sigma^0 K^+)$, whereas the individual χ^2 values from the other seven data show no apparent violation of SU(3) or the sextet dominating assumption. The value of $\chi^2/d.o.f.$ being as large as 2.4 could suggest that the decays of $\Lambda_c^+ \to (\Lambda K^+, \Sigma^0 K^+)$ should be reexamined by BESIII with more precisions. Due to the lack of sufficient data, it leaves the room for more precise examinations by the future experimental measurements on the SU(3) flavor symmetry with or without $\mathcal{O}_{\overline{15}}(\hat{\mathcal{O}}_{\overline{15}})$. With the parameters in Eq. (18) we can reproduce the branching ratios of the measured two-body Λ_c decays as shown in Table II, where the results based on the heavy quark effective theory (HQET) [24], Sharma and Verma (SV) in Ref. [23], pole model (PM) [11] and current algebra (CA) [11] are also listed. In our fit, the SU(3) amplitudes in Eq. (18) have considerable imaginary parts, being included in δ_{f,g,h_2} . Nonetheless, the studies in Refs. [12, 24] depend on real ones. For a test, we turn off δ_{f,g,h_2} , which causes an unsatisfactory fit to the data with $\chi^2/d.o.f \approx 14 \gg 2.4$ in Eq. (18), suggesting that the imaginary parts are necessary to fit the nine data well. It is similar that, in the $D \to MM$ decays, the imaginary parts with the SU(3) flavor symmetry are also considerable, which correspond to the strong phases calculated from the on-shell quark loops in the next-leadingorder QCD models [29]. We hence conclude that the $\Lambda_c^+ \to \mathbf{B}_n M$ decays are like the $D \to MM$ ones, where the phases are in accordance with the higher order contributions in the QCD models, which have not been well developed yet.

From Table I, by keeping both \mathcal{O}_6 and $\mathcal{O}_{\overline{15}}$, we obtain that

$$\mathcal{A}(\Lambda_c^+ \to \Sigma^0 \pi^+) = -\mathcal{A}(\Lambda_c^+ \to \Sigma^+ \pi^0),$$

$$\mathcal{A}(\Lambda_c^+ \to \Sigma^+ K^0) = \sqrt{2}\mathcal{A}(\Lambda_c^+ \to \Sigma^0 K^+),$$
 (19)

where $\Lambda_c^+ \to \Sigma^0 \pi^+$ being identical to $\Lambda_c^+ \to \Sigma^+ \pi^0$ represents the conservation of the isospin (SU(2)) symmetry. By neglecting $\mathcal{O}_{\overline{15}}$, the relations in Eq. (19) can be extended to

$$\mathcal{A}(\Lambda_c^+ \to \Sigma^0 \pi^+) = -\mathcal{A}(\Lambda_c^+ \to \Sigma^+ \pi^0) ,$$

$$\mathcal{A}(\Lambda_c^+ \to \Sigma^+ K^0) = \sqrt{2} \mathcal{A}(\Lambda_c^+ \to \Sigma^0 K^+) ,$$

$$\sqrt{6}\mathcal{A}(\Lambda_c^+ \to \Lambda \pi^+) + \sqrt{2}\mathcal{A}(\Lambda_c^+ \to \Sigma^0 \pi^+) = 2\mathcal{A}(\Lambda_c^+ \to p\bar{K}^0),$$

$$\sqrt{6}\mathcal{A}(\Lambda_c^+ \to \Lambda \pi^+) - \sqrt{2}\mathcal{A}(\Lambda_c^+ \to \Sigma^0 \pi^+) = \frac{2\sqrt{2}}{\lambda}\mathcal{A}(\Lambda_c^+ \to p\pi^0),$$
 (20)

resulting in

$$\mathcal{B}(\Lambda_c^+ \to \Sigma^0 \pi^+) = \mathcal{B}(\Lambda_c^+ \to \Sigma^+ \pi^0),$$

$$\mathcal{B}(\Lambda_c^+ \to \Sigma^+ K^0) = 2\mathcal{B}(\Lambda_c^+ \to \Sigma^0 K^+),$$

$$\mathcal{B}(\Lambda_c^+ \to p\pi^0) \simeq \frac{\lambda^2}{2} [3\mathcal{B}(\Lambda_c^+ \to \Lambda \pi^+) + \mathcal{B}(\Lambda_c^+ \to \Sigma^0 \pi^+) - \mathcal{B}(\Lambda_c^+ \to p\bar{K}^0)].$$
(21)

With the inputs of the SU(3) parameters in Eq. (18), we show $\mathcal{B}(\Lambda_c^+ \to \Sigma^+ K^0) = (8.0 \pm 1.6) \times 10^{-4}$ and $\mathcal{B}(\Lambda_c^+ \to \Sigma^0 K^+) = (4.0 \pm 0.8) \times 10^{-4}$ to agree with the second relation in Eq. (21), which can be used to test the assumption of the dominant \mathcal{O}_6 contributions in comparison with the future measurements. We remark that the factorization approach predicts $\mathcal{B}(\Lambda_c^+ \to \Sigma^0 K^+)$

TABLE II. The branching ratios of the $\Lambda_b \to \mathbf{B}_n M$ decays, where the 2nd column is for our results, where the errors come from the parameters in Eq. (18), while 3, 4, ..., 7 ones correspond to the studies by the heavy quark effective theory (HQET) [24], Sharma and Verma (SV) in Ref. [23], pole model (PM) [11], current algbra (CA) [11] and data [1–3], respectively.

Branching ratios	Our results	HQET $[24]$	SV [23]	PM [11]	CA [11]	Data [1–3]
$10^2 \mathcal{B}(\Lambda_c^+ \to p\bar{K}^0)$	3.3 ± 0.2	1.23	2.67 ± 0.74	1.20	3.46	3.16 ± 0.16
$10^2 \mathcal{B}(\Lambda_c^+ \to \Lambda \pi^+)$	1.3 ± 0.2	1.17		0.84	1.39	1.30 ± 0.07
$10^2 \mathcal{B}(\Lambda_c^+ \to \Sigma^+ \pi^0)$	1.3 ± 0.2	0.69		0.68	1.67	1.24 ± 0.10
$10^2 \mathcal{B}(\Lambda_c^+ \to \Sigma^0 \pi^+)$	1.3 ± 0.2	0.69	0.87 ± 0.20	0.68	1.67	1.29 ± 0.07
$10^2 \mathcal{B}(\Lambda_c^+ \to \Xi^0 K^+)$	0.5 ± 0.1	0.07				0.50 ± 0.12
$10^4 \mathcal{B}(\Lambda_c^+ \to p\pi^0)$	5.6 ± 1.5		2			
$10^4 \mathcal{B}(\Lambda_c^+ \to \Lambda K^+)$	4.6 ± 0.9		14			6.1 ± 1.2
$10^4 \mathcal{B}(\Lambda_c^+ \to \Sigma^0 K^+)$	4.0 ± 0.8		4			5.2 ± 0.8
$10^4 \mathcal{B}(\Lambda_c^+ \to \Sigma^+ K^0)$	8.0 ± 1.6		9			
$10^2 \mathcal{B}(\Lambda_c^+ \to \Sigma^+ \eta)$	0.7 ± 0.4	0.25	0.50 ± 0.17			0.70 ± 0.23
$10^2 \mathcal{B}(\Lambda_c^+ \to \Sigma^+ \eta')$	$1.0^{+1.6}_{-0.8}$	0.08	0.20 ± 0.08			
$10^4 \mathcal{B}(\Lambda_c^+ \to p\eta)$	12.4 ± 4.1		21			12.4 ± 3.0
$10^4 \mathcal{B}(\Lambda_c^+ \to p\eta')$	$12.2^{+14.3}_{-8.7}$		4			

 $\Sigma^0 \pi^+, \Sigma^0 K^+) \simeq 0$, which contradicts the relations from the SU(3) symmetry. This is due to the fact that, when the decay proceeds with the $\Lambda_c^+ \to \Sigma^0$ transition, together with the recoiled meson π^+ or K^+ , the $c \to s$ transition currents transform $\Lambda_c^+ \to \Lambda = (ud - du)s$, which is unable to correlate to $\Sigma^0 = (ud + du)s$ [30], leading to $\mathcal{B} = 0$.

In Eq. (21), the simple estimation based on the data inputs gives that $\mathcal{B}(\Lambda_c^+ \to p\pi^0) = (5.1 \pm 0.7) \times 10^{-4}$, which agrees with our numerical fitting result of $\mathcal{B}(\Lambda_c^+ \to p\pi^0) = (5.6 \pm 1.5) \times 10^{-4}$, but is larger than the experimental upper bound of 3×10^{-4} (90%C.L.) in Eq. (2) by BESIII. To check if there is a discrepancy here, we have taken the original data of $\mathcal{B}(\Lambda_c^+ \to p\pi^0) = (7.95 \pm 13.61) \times 10^{-5}$ [4] by BESIII as the input. In this case, we get $\chi^2/d.o.f = 4.7$, which is two times larger than the value in Eq. (18), showing that the fitting cannot accommodate the present data of $\mathcal{B}(\Lambda_c^+ \to p\pi^0)$. Apart from the SU(3) flavor symmetry, we estimate that $\mathcal{B}(\Lambda_c^+ \to p\pi^0) \simeq 5 \times 10^{-4}$ in the approach of the factorization, which is also larger than the experimental upper bound. It is clear that a dedicated search for this mode with a more precise measurement should be done. An improved sensitivity to measure $\Lambda_c^+ \to p\pi^0$ will clarify if the currently unmovable discrepancy exists or not.

It is also interesting to see that $\mathcal{B}(\Lambda_c^+ \to \Sigma^+ \eta')$ and $\mathcal{B}(\Lambda_c^+ \to p\eta')$ fitted to be $(1.0^{+1.6}_{-0.8}) \times 10^{-2}$ and $(12.2^{+14.3}_{-8.7}) \times 10^{-4}$ are as large as their η counterparts, respectively, while $\mathcal{B}(\Lambda_b \to \Lambda \eta) \simeq \mathcal{B}(\Lambda_b \to \Lambda \eta')$ [31]. We note that there is a similar term in Ref. [32] as the $h_{1,2}$ terms, which relates $\Lambda_c^+ \to \Sigma^+ \eta'$ to $\Xi_c^0 \to \Xi^0 \eta'$. In contrast, the theoretical approach in Ref. [23] is based on the SU(3) flavor symmetry also, but without the $h_{1,2}$ terms to include the singlet η_1 , such that it leads to $\mathcal{B}(\Lambda_c^+ \to \Sigma^+(p)\eta') < \mathcal{B}(\Lambda_c^+ \to \Sigma^+(p)\eta)$ [23]. Finally, we remark that, with the SU(3) symmetry, we can extend our study to the two-body $\Xi_b^{+,0}$ decays, which are also accessible to the current experiments. Since the two-body $\Lambda_c^+ \to \mathbf{B}_n V$ with V the vector meson and three-body Λ_c^+ decays are observed, which require the interpretations, the approach of the SU(3) symmetry can be useful.

IV. CONCLUSIONS

We have studied the two-body $\Lambda_c^+ \to \mathbf{B}_n M$ decays, which have been recently reanalyzed or newly measured by BESIII. With the SU(3) flavor symmetry, we can describe the data except that for $\Lambda_c^+ \to p\pi^0$. We have found that $\mathcal{B}(\Lambda_c^+ \to p\pi^0) = (5.6 \pm 1.5) \times 10^{-4}$, which is almost 2σ above the experimental upper bound of 3×10^{-4} . We hope that the future experimental measurement of $\mathcal{B}(\Lambda_c^+ \to p\pi^0)$ can resolve this discrepancy. Unlike the previous results, we have predicted that $\mathcal{B}(\Lambda_c^+ \to \Sigma^+ \eta') = (1.0^{+1.6}_{-0.8}) \times 10^{-2}$ and $\mathcal{B}(\Lambda_c^+ \to p\eta') = (12.2^{+14.3}_{-8.7}) \times 10^{-4}$ which are as large as their η counterparts, due to the newly added $h_{1,2}$ terms with the singlet η_1 in the SU(3) flavor symmetry. With the SU(3) symmetry, one is able to study $\Lambda_c^+ \to \mathbf{B}_n V$ and the three-body \mathbf{B}_c decays, which have been observed but barely interpreted. Moreover, the extensions to study the $\Xi_b^{+,0}$ decays are possible, which are also accessible to the current experiments.

ACKNOWLEDGMENTS

We would like to thank X.G. He for useful discussions. This work was supported in part by National Center for Theoretical Sciences, MoST (MoST-104-2112-M-007-003-MY3), and National Science Foundation of China (11675030).

- [1] M. Ablikim et al. [BESIII Collaboration], Phys. Rev. Lett. 116, 052001 (2016).
- [2] M. Ablikim et al. [BESIII Collaboration], arXiv:1702.05279 [hep-ex].
- [3] C. Patrignani et al. [Particle Data Group], Chin. Phys. C 40, 100001 (2016).
- [4] The private communication with Dr. Peilian Li.
- [5] A. Ali, G. Kramer and C.D. Lu, Phys. Rev. D58, 094009 (1998).
- [6] C.Q. Geng, Y.K. Hsiao and J.N. Ng, Phys. Rev. Lett. 98, 011801 (2007).
- [7] Y.K. Hsiao and C.Q. Geng, Phys. Rev. D 91, 116007 (2015).
- [8] J.D. Bjorken, Phys. Rev. D 40, 1513 (1989).
- [9] C.D. Lu, W. Wang and F.S. Yu, Phys. Rev. D 93, 056008 (2016).
- [10] H.Y. Cheng and B. Tseng, Phys. Rev. D 46, 1042 (1992); 55, 1697(E) (1997).
- [11] H.Y. Cheng and B. Tseng, Phys. Rev. D 48, 4188 (1993).
- [12] P. Zenczykowski, Phys. Rev. D 50, 402 (1994).
- [13] T. Uppal, R.C. Verma and M.P. Khanna, Phys. Rev. D 49, 3417 (1994).
- [14] Fayyazuddin and Riazuddin, Phys. Rev. D 55, 255; 56, 531(E) (1997).
- [15] X.G. He, Y.K. Hsiao, J.Q. Shi, Y.L. Wu and Y.F. Zhou, Phys. Rev. D 64, 034002 (2001).
- [16] H.K. Fu, X.G. He and Y.K. Hsiao, Phys. Rev. D 69, 074002 (2004).

- [17] Y.K. Hsiao, C.F. Chang and X.G. He, Phys. Rev. D 93, 114002 (2016).
- [18] X.G. He and G.N. Li, Phys. Lett. B **750**, 82 (2015).
- [19] M. He, X.G. He and G.N. Li, Phys. Rev. D 92, 036010 (2015).
- [20] M.J. Savage and R.P. Springer, Phys. Rev. D 42, 1527 (1990).
- [21] M.J. Savage, Phys. Lett. B **257**, 414 (1991).
- [22] R.C. Verma and M.P. Khanna, Phys. Rev. D 53, 3723 (1996).
- [23] K.K. Sharma and R.C. Verma, Phys. Rev. D 55, 7067 (1997).
- [24] K.K. Sharma and R.C. Verma, Eur. Phys. J. C 7, 217 (1999).
- [25] H.n. Li, C.D. Lu and F.S. Yu, Phys. Rev. D 86, 036012 (2012).
- [26] A.J. Buras, hep-ph/9806471.
- [27] S. Fajfer, P. Singer and J. Zupan, Eur. Phys. J. C 27, 201 (2003).
- [28] T. Feldmann, P. Kroll and B. Stech, Phys. Rev. D 58, 114006 (1998); Phys. Lett. B 449, 339 (1999).
- [29] H.Y. Cheng and C.W. Chiang, Phys. Rev. D 86, 014014 (2012).
- [30] Y.K. Hsiao, Y. Yao and C.Q. Geng, Phys. Rev. D 95, 093001 (2017).
- [31] C.Q. Geng, Y.K. Hsiao, Y.H. Lin and Y. Yu, Eur. Phys. J. C 76, 399 (2016).
- [32] G. Altarelli, N. Cabibbo and L. Maiani, Phys. Lett. **57B**, 277 (1975).