

Sufficient condition for root reconstruction by parsimony on binary trees with general weights *

Sebastien Roch[†]

Kun-Chieh Wang

January 1, 2021

Abstract

We consider the problem of inferring an ancestral state from observations at the leaves of a tree, assuming the state evolves along the tree according to a two-state symmetric Markov process. We establish a general branching rate condition under which maximum parsimony, a common reconstruction method requiring only the knowledge of the tree, succeeds better than random guessing uniformly in the depth of the tree. We thereby generalize previous results of [GS10, ZSYL10]. Our results apply to both deterministic and i.i.d. edge weights.

*Keywords: Markov model on a tree, reconstruction problem, branching number, parsimony principle

[†]Department of Mathematics at the University of Wisconsin–Madison. Supported by NSF grants DMS-1248176, DMS-1149312 (CAREER), DMS-1614242, CCF-1740707 (TRIPODS), DMS-1902892, DMS-2023239, as well as a Simons Fellowship and a Vilas Associates Award. roch@math.wisc.edu

1 Introduction

Ancestral reconstruction In biology, the inferred evolutionary history of organisms is depicted by a phylogenetic tree, that is, a rooted tree whose branchings indicate past speciation events with the leaves representing living species. The evolution of features, such as the nucleotide at a given position in the genome of a species, the presence or absence of a protein or the number of horns in a lizard, is commonly assumed to follow Markovian dynamics along the tree [Ste16]. That is, on each edge, the state of the feature changes according to a Markov process; at bifurcations, two independent copies of the feature evolve along the outgoing edges starting from the state at the branching point.

Here we consider the problem of inferring an ancestral state from observations of a feature at the leaves of a known phylogenetic tree. We refer to this problem, which has important applications in biology [Mos04, DMR11, RS17], as the ancestral reconstruction problem. Many rigorous results have been obtained in previous work; see, e.g., [KS66, BRZ95, Iof96, TS97, Mos98, EKPS00, Mos01, MP03, BCMR06, GS10, ZSYL10, Sly11, MRS13, MS14, BST16, LJV18, FR18, MMP20, FR20, GRP20, ABDLV20] for a partial list. Typically, one seeks an estimator of the root state which is strictly superior to random guessing—uniformly in the depth of the tree—under a uniform prior on the root [Mos01]. Whether such an estimator exists has been shown to depend on a trade-off between the mixing rate of the Markov process and the growth rate of the tree. In some cases, for instance in two-state symmetric Markov chains on binary trees [KS66, Iof96] as well as on more general trees [EKPS00], sharp thresholds have been established. We focus on that special case here.

The threshold of maximum parsimony Maximum parsimony is an ancestral reconstruction method with a long history in evolutionary biology. See e.g. [Ste16]. Its underlying principle is simple and intuitive: it calls for assigning a state to *each* internal vertex in such a way as to *minimize the total number of changes* along the edges; the resulting state at the *root* (possibly not unique) is the desired reconstructed state. An advantage of maximum parsimony, which accounts partly for its occasional use in practice, is that it only requires knowledge of the tree—not of the substitution probabilities along the edges, which can be difficult to estimate accurately from data and can affect the accuracy of reconstruction [CR14]. In fact maximum parsimony is equivalent to maximum likelihood under those circumstances [TS97]. Another root state estimator with that property is recursive majority, studied in [Mos98, Mos04].

The theoretical properties of maximum parsimony for ancestral state reconstruction have been widely studied [Ste89, TS97, FT09, ZSYL10, GS10, CH11, GS14, HF17, HF18, HLS19]. From its very definition, one might expect maximum parsimony to perform well when the probabilities of substitution along the edges are “sufficiently small.” This statement was confirmed rigorously for the two-state symmetric model on complete binary trees with constant mutation probability in [ZSYL10] (as first conjectured in [Ste89]). Below a critical probability, maximum parsimony does indeed beat random guessing. A related result was obtained in [GS10] under a common random tree model known as the Yule tree. Here we substantially generalize both of these results. We give a general sufficient condition on the branching rate under which maximum parsimony succeeds. Our condition is related to the branching number, which plays an important role in many processes on trees. See e.g. [LP16]. Other generalized notions of degree have been used in related contexts [SStY17].

Definitions In order to state our results formally, we begin with some definitions. Let $\mathcal{T} = (\mathcal{V}, \mathcal{E})$ be an infinite complete binary tree rooted at ρ . That is, all vertices of \mathcal{T} have exactly two children; in particular, \mathcal{T} has no leaf. Every edge $e \in \mathcal{E}$ is assigned a weight $\theta_e \in [0, 1]$. If $e = (x, y)$ where x is the parent of y , we also write $\theta_y = \theta_e$. We use the notation $x \leq y$ to indicate that x is an ancestor of y and we write $x < y$ for $x \leq y$ and $x \neq y$. We also let $s(x)$ be the sibling of $x \neq \rho$.

Under the *Cavender-Farris (CF) model* over \mathcal{T} and $\theta = (\theta_y)_{y \in \mathcal{V}}$, also known as Neyman 2-state model, we associate to each vertex $x \in \mathcal{V}$ a state $\sigma_x \in \{0, 1\}$ as follows. The state at the root, σ_ρ , is picked uniformly at random in $\{0, 1\}$. Recursively, if y has parent x , state σ_y is equal to σ_x with probability θ_y , otherwise it is picked uniformly at random in $\{0, 1\}$. We let $p_y = \mathbb{P}_{\mathcal{T}, \theta}[\sigma_x \neq \sigma_y] = \frac{1-\theta_y}{2}$ be the *probability of a substitution* on edge $e = (x, y)$. Here $\mathbb{P}_{\mathcal{T}, \theta}$ stands for the probability operator under the CF model over \mathcal{T} and θ . The CF model is equivalent to the Fortuin-Kastelyn random cluster representation of the ferromagnetic Ising model on \mathcal{T} with a free boundary. See e.g. [EKPS00] and references therein.

Informally, a root state estimator is a map returning a (possibly randomized) guess for the state of the root, based on the knowledge of the states on an observed cutset.

Definition 1 (Cutset). *A cutset is a minimal, finite set of vertices $\pi \subseteq \mathcal{V}$ such that all infinite self-avoiding paths starting at ρ must visit π . We let $\mathcal{C}(\mathcal{T})$ be the collection of all cutsets of \mathcal{T} . We denote by $\mathcal{T}^\pi = (\mathcal{V}^\pi, \mathcal{E}^\pi)$ the finite tree obtained from \mathcal{T} after removing all descendants of the vertices in π .*

As mentioned above, our focus in this work is on a root state estimator known in phylogenetics as *maximum parsimony*. The parsimony principle dictates that one assigns to each vertex x (ancestor to the observed cutset π) a state $\hat{\sigma}_x$ such that the overall number of changes along the edges of \mathcal{T}^π , namely, $\sum_{(x,y) \in \mathcal{E}^\pi} \mathbf{1}\{\hat{\sigma}_x \neq \hat{\sigma}_y\}$, is minimized, where we let by default $\hat{\sigma}_z = \sigma_z$ for all $z \in \pi$. In case both 0 and 1 can be obtained in this way as root state, a uniformly random value in $\{0, 1\}$ is returned. We let $\text{RA}_{\mathcal{T}, \theta}^\pi$ be the *reconstruction accuracy* of parsimony, i.e. the probability that it correctly reconstructs the root state.

Main result: deterministic weights In our main result, we give conditions under which the reconstruction accuracy of maximum parsimony is uniformly bounded away from $1/2$.

Theorem 1 (Reconstruction accuracy of parsimony: deterministic weights). *Let $\mathcal{T} = (\mathcal{V}, \mathcal{E})$ be an infinite complete binary tree with edge weights $\theta = (\theta_z)_{z \in \mathcal{V}}$ satisfying $\theta_* := \inf_{z \in \mathcal{V}} \theta_z > 0$ and*

$$\sup \left\{ \kappa > 0 : \inf_{\pi \in \mathcal{C}(\mathcal{T})} \sum_{x \in \pi} \prod_{\rho \neq z \leq x} \kappa^{-1} \theta_z > 0 \right\} > \frac{3}{2}. \quad (1)$$

Then $\inf_{\pi \in \mathcal{C}(\mathcal{T})} \text{RA}_{\mathcal{T}, \theta}^\pi > \frac{1}{2}$, i.e., the reconstruction accuracy of parsimony on \mathcal{T} is away from $1/2$.

Condition (1) involves the *branching number*, a generalized notion of branching rate which plays a key role in the analysis of many stochastic processes on trees and tree-like graphs. See e.g. [LP16]. The following example provides some intuition in a special case.

Example 1 (Fixed edge weights). *As a simple illustration, observe that, when all weights are equal to $\theta_* \in (0, 1]$, the supremum in (1) is attained for $\kappa = 2\theta_*$. Indeed, the sum in (1) when $\kappa = 2\theta_*$ simplifies to*

$$\sum_{x \in \pi} \prod_{\rho \neq z \leq x} (2\theta_*)^{-1} \theta_z = \sum_{x \in \pi} \prod_{\rho \neq z \leq x} \frac{1}{2} = 1 > 0$$

for any cutset π , where the equality can be proved by induction on the graph distance from the root to the furthest vertex in π . On the other hand, letting π_n be the cutset of all vertices at graph distance n from ρ , for any $\varepsilon > 0$ it holds that

$$\sum_{x \in \pi_n} \prod_{\rho \neq z \leq x} ((2 + \varepsilon)\theta_*)^{-1} \theta_z = (2 + \varepsilon)^{-n} \cdot 2^n \rightarrow 0,$$

as $n \rightarrow +\infty$. Hence, in this case, condition (1) reduces to $2\theta_ > 3/2$, that is, $\theta_* > 3/4$. In terms of substitution probability, this is $p_* = \frac{1-\theta_*}{2} < 1/8$.*

The argument in the example above leads to the following corollary.

Corollary 1 (All substitution probabilities below the threshold). *Let the substitution probabilities $(p_z)_{z \in \mathcal{V}}$ satisfy $p_z \in [0, p_*]$ for all $z \in \mathcal{V}$, for some $p_* < 1/8$. Then the reconstruction accuracy of maximum parsimony on \mathcal{T} is bounded away from $1/2$.*

In general, Theorem 1 cannot be improved in the following sense. It was shown in [ZSYL10, Theorem 4.1] that when $p_z = p_* > 1/8$, that is, $\theta_z = 1 - 2p_z = \theta_* < 3/4$ for all z then $\inf_n \text{RA}_{\mathcal{T}, \theta}^{\pi_n} = \frac{1}{2}$, where π_n is defined in Example 1. On the other hand, it is not known whether the reconstruction accuracy necessarily converges to $1/2$ if (1) is not satisfied. We leave this as an open problem.

Zhang et al. [ZSYL10] also established the special case of Theorem 1 when $p_z = p_* < 1/8$ for all z and $\pi = \pi_n$. Their proof proceeds through a careful analysis of the limit of a recurrence for $\text{RA}_{\mathcal{T}, \theta}^{\pi_n}$ first derived in [Ste89]. Our more general result follows from a softer argument which relies on the instability of a fixed point of this recurrence corresponding to asymptotic reconstruction accuracy $1/2$. A more detailed proof sketch is given in Section 2 following some preliminaries. We note that our proof method may be of more general interest, e.g., to extend the results beyond the two-state case where the higher dimensionality of the system may complicate significantly the derivation of an explicit limit even when edge weights are constant.

Main result: i.i.d. weights We also obtain a related result in the case of edge weights that are i.i.d. No lower bound on the weights is needed in this case, unlike Theorem 1.

Theorem 2 (Reconstruction accuracy of parsimony: i.i.d. weights). *Let $\mathcal{T} = (\mathcal{V}, \mathcal{E})$ be an infinite complete binary tree with edge weights $\theta = (\theta_z)_{z \in \mathcal{V}}$ drawn i.i.d. from a distribution Θ over $(0, 1]$. Let μ be the mean of Θ and assume that $\mu > 3/4$. Then, for any $\delta > 0$, there is $\varepsilon > 0$ such that $\inf_{\pi \in \mathcal{C}(\mathcal{T})} \text{RA}_{\mathcal{T}, \theta}^{\pi} > \frac{1}{2}(1 + \varepsilon)$, with probability at least $1 - \delta$.*

The previous theorem covers in particular the case of the pure birth process, or Yule tree, which is a popular random model of phylogenetic trees. See e.g. [Ste16]. In that case, $\theta_z = e^{-2T_z}$, where T_z is an exponential with rate λ . To derive the corresponding threshold, we note that

$$\mu = \mathbb{E}[\theta_z] = \int_0^\infty e^{-2t} \lambda e^{-\lambda t} dt = \frac{\lambda}{\lambda + 2}.$$

Then $\mu > \frac{3}{4} \iff \lambda > 6$, which is consistent with the results of [GS10].

2 Preliminaries

Computing parsimony Our proofs are based on a recurrence for the reconstruction accuracy. Maximum parsimony can be computed efficiently by dynamic programming, which is referred to as the *Fitch method*.

Definition 2 (Parsimony recursion). *The Fitch method recursively constructs a set \widehat{S}_z^π of possible states for each vertex $z \in \mathcal{V}^\pi$, starting from π , as follows. If $z \in \pi$, $\widehat{S}_z^\pi = \{\sigma_z\}$. If $z \notin \pi$ and has children x and y ,*

$$\widehat{S}_z^\pi = \begin{cases} \widehat{S}_x^\pi \cap \widehat{S}_y^\pi, & \text{if } \widehat{S}_x^\pi \cap \widehat{S}_y^\pi \neq \emptyset \\ \widehat{S}_x^\pi \cup \widehat{S}_y^\pi, & \text{o.w.} \end{cases}$$

The method returns the maximum parsimony estimator $\hat{\sigma}_\rho$ which is equal to the unique state in \widehat{S}_ρ^π if $|\widehat{S}_\rho^\pi| = 1$, and otherwise returns a uniformly random value in $\{0, 1\}$.

What is described above is the bottom-up phase of the Fitch method. (A top-down phase, which we will not require here, then assigns a state to each vertex in \mathcal{V}^π consistent with a maximum parsimony solution; e.g. [Ste16].) Let π be an arbitrary cutset on \mathcal{T} with states σ_u , $u \in \pi$, and let \widehat{S}_z , $z \in \mathcal{V}^\pi$, be the corresponding reconstructed sets under the Fitch method. We define

$$\alpha_z^\pi = \mathbb{P}_{\mathcal{T},\theta} \left[\widehat{S}_z^\pi = \{\sigma_z\} \right], \quad \beta_z^\pi = \mathbb{P}_{\mathcal{T},\theta} \left[\widehat{S}_z^\pi = \{1 - \sigma_z\} \right].$$

Under our randomization rule, the reconstruction accuracy of maximum parsimony $\text{RA}_{\mathcal{T},\theta}^\pi$ is given by

$$\mathbb{P}_{\mathcal{T},\theta} \left[\widehat{S}_\rho^\pi = \{\sigma_\rho\} \right] + \frac{1}{2} \mathbb{P}_{\mathcal{T},\theta} \left[\widehat{S}_\rho^\pi = \{0, 1\} \right] = \alpha_\rho^\pi + \frac{1}{2}(1 - \alpha_\rho^\pi - \beta_\rho^\pi) = \frac{1}{2} + \frac{1}{2}(\alpha_\rho^\pi - \beta_\rho^\pi). \quad (2)$$

Proof sketch Fix θ satisfying the assumptions of Theorem 1 and fix a cutset π . To analyze $(\alpha_z^\pi, \beta_z^\pi)$, it is natural to take advantage of the recursive nature of \mathcal{T} . Let x and y be the children of z . The event $\widehat{S}_z^\pi = \{\sigma_z\}$ occurs when either (i) $\widehat{S}_x^\pi = \{\sigma_z\}$ and $\widehat{S}_y^\pi = \{\sigma_z\}$, or (ii) $\widehat{S}_x^\pi = \{\sigma_z\}$ and $\widehat{S}_y^\pi = \{0, 1\}$ or vice versa. By the Markov property of the CF model, the random variables \widehat{S}_x^π and \widehat{S}_y^π , which are functions only of the states of π under x and y respectively, are conditionally independent given σ_z . Hence, letting $q_u = 1 - p_u$ for $u = x, y$ and taking into account the possibility of a mutation along the edges (z, x) and (z, y) , it follows as first derived in [Ste89, Lemma 7.20] that

$$\begin{aligned} \alpha_z^\pi &= (q_x \alpha_x^\pi + p_x \beta_x^\pi)(q_y \alpha_y^\pi + p_y \beta_y^\pi) \\ &\quad + (q_x \alpha_x^\pi + p_x \beta_x^\pi)(1 - \alpha_y^\pi - \beta_y^\pi) + (1 - \alpha_x^\pi - \beta_x^\pi)(q_y \alpha_y^\pi + p_y \beta_y^\pi), \end{aligned} \quad (3)$$

where the first and second lines on the r.h.s. correspond respectively to cases (i) and (ii) above. Similarly,

$$\begin{aligned} \beta_z^\pi &= (p_x \alpha_x^\pi + q_x \beta_x^\pi)(p_y \alpha_y^\pi + q_y \beta_y^\pi) \\ &\quad + (p_x \alpha_x^\pi + q_x \beta_x^\pi)(1 - \alpha_y^\pi - \beta_y^\pi) + (1 - \alpha_x^\pi - \beta_x^\pi)(p_y \alpha_y^\pi + q_y \beta_y^\pi). \end{aligned} \quad (4)$$

In the case that $p_u = p$ for all u , a fixed point analysis was performed in [Ste89, Theorem 7.22]. It was found that, if $p \geq 1/8$, there is a single fixed point $(1/3, 1/3)$ which corresponds informally to ‘‘having no information about the root.’’ While if $p < 1/8$, there is an additional fixed point $(\alpha_p^\infty, \beta_p^\infty)$ with $\alpha_p^\infty > \beta_p^\infty$. Convergence to $(1/3, 1/3)$ in the first case and $(\alpha_p^\infty, \beta_p^\infty)$ in the second case was established rigorously in [ZSYL10]. One step in [ZSYL10] involved the derivation of a new recurrence in terms of the quantities $\alpha_z^{\pi_n} - \beta_z^{\pi_n}$ and $1 - (\alpha_z^{\pi_n} + \beta_z^{\pi_n})$, which facilitates the analysis of the limit in the fixed edge weight case.

Going back to binary trees with general weights, as our starting point we further modify the recurrence of [ZSYL10]. For all $z \in \mathcal{V}$, we define

$$\mathbf{d}_z^\pi = \alpha_z^\pi - \beta_z^\pi, \quad \text{and} \quad \mathbf{u}_z^\pi = 3(\alpha_z^\pi + \beta_z^\pi) - 2. \quad (5)$$

We show in Proposition 1 below that $(\mathbf{d}_z^\pi, \mathbf{u}_z^\pi)$ satisfies the following recurrence

$$\mathbf{d}_z^\pi = \left(\frac{4 - \mathbf{u}_y^\pi}{6} \right) \theta_x \mathbf{d}_x^\pi + \left(\frac{4 - \mathbf{u}_x^\pi}{6} \right) \theta_y \mathbf{d}_y^\pi, \quad (6)$$

$$\mathbf{u}_z^\pi = \frac{3}{2} \theta_x \theta_y \mathbf{d}_x^\pi \mathbf{d}_y^\pi - \frac{1}{2} \mathbf{u}_x^\pi \mathbf{u}_y^\pi, \quad (7)$$

for $z \in \mathcal{V}^\pi - \pi$ with children x, y , as well as the inequalities $0 \leq \mathbf{d}_z^\pi \leq 1$ and $-1/2 \leq \mathbf{u}_z^\pi \leq 1$ for all $z \in \mathcal{V}^\pi$ and the boundary conditions $\mathbf{d}_z^\pi = \mathbf{u}_z^\pi = 1$ for all $z \in \pi$. Our choice of parametrization is motivated

in part by the fact that the “no information” fixed point is now at $(0, 0)$ and that $|\mathbf{d}_z^\pi|, |\mathbf{u}_z^\pi| \leq 1$. At a high level we show that, under the branching rate condition (1), the fixed point $(0, 0)$ is “unstable” and that \mathbf{d}_z^π in particular stays bounded away from 0. That in terms implies a lower bound on the reconstruction accuracy as, by (2), we have

$$\text{RA}_{\mathcal{T}, \theta}^\pi = \frac{1}{2} + \frac{1}{2} \mathbf{d}_\rho^\pi.$$

The link between stability and the weighted branching rate in (1) is easily seen from (6). First, when all weights are equal to θ_* and $\pi = \pi_n$, we get by symmetry that close to $(0, 0)$ to the first order $\mathbf{d}_z^\pi \approx \frac{4}{3} \theta_* \mathbf{d}_x^\pi$. Hence, in that case, the solution can be expected to grow when $\frac{4}{3} \theta_* > 1$, corresponding to the condition derived in Example 1. More generally, we use (6) to relate the \mathbf{d} -value at the root to the \mathbf{d} -values on a cutset (see Lemma 1). To deal with the nonlinear nature of (6) and (7), we control the \mathbf{u} -values thanks to the quadratic form of (7) which implies a quick decay towards 0 (see Lemma 3).

Recurrence Before proceeding to the proof of our main results, we first establish a basic recurrence which follows from the work of [ZSYL10]. We give a short proof for completeness.

Proposition 1 (Recurrence and basic properties). *The following hold:*

- [Boundary conditions] For all $z \in \pi$, $\mathbf{d}_z^\pi = 1$ and $\mathbf{u}_z^\pi = 1$.
- [Recurrence] For all $z \in \mathcal{V}^\pi - \pi$, if x, y are the children of z , the system (6) and (7) holds.
- [Bounds] For all $z \in \mathcal{V}^\pi$, we have $0 \leq \mathbf{d}_z^\pi \leq 1$ and $-1/2 \leq \mathbf{u}_z^\pi \leq 1$.

Proof. We start with the boundary conditions. By Definition 2(a) and the definitions of α_z^π and β_z^π , we have for all $z \in \pi$ that $\alpha_z^\pi = \mathbb{P}_{\mathcal{T}, \theta}[\widehat{S}_z^\pi = \{\sigma_z\}] = 1$ and $\beta_z^\pi = \mathbb{P}_{\mathcal{T}, \theta}[\widehat{S}_z^\pi = \{1 - \sigma_z\}] = 0$. So $\mathbf{u}_z^\pi = 3(\alpha_z^\pi + \beta_z^\pi) - 2 = 1$ and $\mathbf{d}_z^\pi = \alpha_z^\pi - \beta_z^\pi = 1$.

The second statement is merely a change of variables. We briefly expand on the first equation (the other one being similar). Let $z \in \mathcal{V}^\pi - \pi$ with children x, y . We define $\Sigma_u = \alpha_u^\pi + \beta_u^\pi$ and $\Delta_u = \alpha_u^\pi - \beta_u^\pi$. By the definitions of p_x and q_x , note that

$$q_x \alpha_x^\pi + p_x \beta_x^\pi = \left(\frac{1 + \theta_x}{2} \right) \alpha_x^\pi + \left(\frac{1 - \theta_x}{2} \right) \beta_x^\pi = \frac{1}{2} \Sigma_x + \frac{1}{2} \theta_x \Delta_x,$$

and, similarly, $p_x \alpha_x^\pi + q_x \beta_x^\pi = \frac{1}{2} \Sigma_x - \frac{1}{2} \theta_x \Delta_x$. Hence, by (3) and (4),

$$\alpha_z^\pi = \frac{1}{4} (\Sigma_x + \theta_x \Delta_x) (\Sigma_y + \theta_y \Delta_y) + \frac{1}{2} (\Sigma_x + \theta_x \Delta_x) (1 - \Sigma_y) + \frac{1}{2} (1 - \Sigma_x) (\Sigma_y + \theta_y \Delta_y),$$

and

$$\beta_z^\pi = \frac{1}{4} (\Sigma_x - \theta_x \Delta_x) (\Sigma_y - \theta_y \Delta_y) + \frac{1}{2} (\Sigma_x - \theta_x \Delta_x) (1 - \Sigma_y) + \frac{1}{2} (1 - \Sigma_x) (\Sigma_y - \theta_y \Delta_y).$$

Subtracting the above two equations, we get

$$\begin{aligned} \Delta_z &= \frac{1}{2} \theta_x \Delta_x \Sigma_y + \frac{1}{2} \theta_y \Delta_y \Sigma_x + \theta_x \Delta_x (1 - \Sigma_y) + \theta_y \Delta_y (1 - \Sigma_x) \\ &= \left(1 - \frac{1}{2} \Sigma_y \right) \theta_x \Delta_x + \left(1 - \frac{1}{2} \Sigma_x \right) \theta_y \Delta_y, \end{aligned}$$

which after plugging in (5) gives (6).

Because α_z^π and β_z^π are probabilities and further $\alpha_z^\pi + \beta_z^\pi \leq 1$, we have that $\mathbf{d}_z^\pi \leq 1$ and $\mathbf{u}_z^\pi \leq 1$, for all z . Moreover, that together with the boundary conditions and (6) implies that $\mathbf{d}_z^\pi \geq 0$ for all z by induction. In turn, that together with the boundary conditions and (7) implies that $\mathbf{u}_z^\pi \geq -1/2$ for all z . \square

3 Deterministic weights

Before proceeding with the proof of Theorem 1, we first prove some lemmas.

3.1 Controlling d- and u-values

In the first lemma, we express the d-value at the root as a function of the d- and u-values above an arbitrary cutset. Recall that $s(z)$ is the sibling of z .

Lemma 1 (Controlling the root with a cutset). *For any cutset π' in \mathcal{T}^π , it holds that*

$$\mathbf{d}_\rho^\pi = \sum_{x \in \pi'} \mathbf{d}_x^\pi \prod_{\rho \neq z \leq x} \left\{ \frac{4 - \mathbf{u}_{s(z)}^\pi}{6} \right\} \theta_z. \quad (8)$$

Proof. The result follows by recursively applying (6) from the root down to π' . We implicitly use the fact that, by definition, a cutset is minimal. \square

Our second lemma shows that d-values cannot grow too fast down the tree. This fact will be useful to proving the next key lemma. We will need the lower bound $\theta_* = \inf_{z \in \mathcal{V}} \theta_z > 0$, on the θ -values. For $v, w \in \mathcal{V}^\pi$, we let $\gamma(v, w)$ be the graph distance between v and w in \mathcal{T}^π . Recall that $\theta_* \leq 1 < 2$.

Lemma 2 (Growth of d-values). *Fix any $v \in \mathcal{V}^\pi$. Then $\forall \varepsilon' > 0$, for all descendants w of v in \mathcal{V}^π*

$$\mathbf{d}_v^\pi \leq \varepsilon' \implies \mathbf{d}_w^\pi \leq \varepsilon' \left(\frac{2}{\theta_*} \right)^{\gamma(v, w)}.$$

Proof. Let $z \in \mathcal{V}^\pi$, not on π , have children x and y . (Note that, in the case where z is the parent of a vertex on the cutset π , z itself cannot be on the cutset by minimality and therefore both its children are in \mathcal{V}^π .) By Proposition 1, we have $\mathbf{u}_x^\pi, \mathbf{u}_y^\pi \leq 1$ and $\mathbf{d}_x^\pi, \mathbf{d}_y^\pi \geq 0$, which implies that both terms on the r.h.s. of (6) are non-negative. Hence, using $\theta_x \geq \theta_*$, (6) gives $\mathbf{d}_z^\pi \geq \frac{1}{2} \theta_* \mathbf{d}_x^\pi$. In particular, $\mathbf{d}_z^\pi \leq \varepsilon'$ implies that $\mathbf{d}_x^\pi \leq \varepsilon' (2/\theta_*)$. Recursing gives the claim. \square

Our final lemma controls u-values at the root of a subtree where d-values are uniformly small.

Lemma 3 (Controlling u when d is small). *Fix any $0 < \phi' \leq 1/9$ and $v \in \mathcal{V}^\pi$. Then there exists $\varepsilon' > 0$ depending only on θ_* and ϕ' such that: $\mathbf{d}_v^\pi \leq \varepsilon' (2/\theta_*)$ implies $|\mathbf{u}_v^\pi| \leq 4\phi'$.*

Proof. Let H be the smallest non-negative integer such that

$$\left(\frac{1}{2} \right)^{-1+2^H} \leq \phi'. \quad (9)$$

Define $\varepsilon' > 0$ to be the largest positive real such that

$$\varepsilon' \left(\frac{2}{\theta_*} \right)^{H+1} \leq 0.99 \quad \text{and} \quad \frac{3}{2} \left[\varepsilon' \left(\frac{2}{\theta_*} \right)^{H+1} \right]^2 \leq \phi'. \quad (10)$$

The rest of the proof proceeds in two steps: we derive a simplified recurrence for u-values and solve it.

1. *Simplified recurrence:* Assume that $\mathbf{d}_v^\pi \leq \varepsilon'(2/\theta_*)$. Let w be a descendant of v with graph distance $\gamma(v, w) \leq H$. Then, by Lemma 2, $\mathbf{d}_w^\pi \leq \varepsilon'(2/\theta_*)^{H+1} < 1$, where we used (10). This shows, in particular, that all descendants of v in \mathcal{V} within graph distance H are in fact strictly above π , because \mathbf{d} -values are 1 on the cutset π . Moreover, by the recurrence (7) and the inequality (10), for any descendant w_0 of v with children w_1, w_2 in \mathcal{V}^π that are within graph distance H of v , we have

$$|\mathbf{u}_{w_0}^\pi| \leq \frac{3}{2}\theta_{w_1}\theta_{w_2}\mathbf{d}_{w_1}^\pi\mathbf{d}_{w_2}^\pi + \frac{1}{2}|\mathbf{u}_{w_1}^\pi||\mathbf{u}_{w_2}^\pi| \leq \phi' + \frac{1}{2}|\mathbf{u}_{w_1}^\pi||\mathbf{u}_{w_2}^\pi|, \quad (11)$$

where we used that θ -values are ≤ 1 and (10). Define $\mathbf{U}_h = \sup\{|\mathbf{u}_w^\pi| : w \in \mathcal{V}^\pi, v \leq w \text{ and } \gamma(v, w) = h\}$, and $\mathbf{U}_0 = |\mathbf{u}_v^\pi|$. By the remark above, the set in the previous display is non-empty for all $h = 0, \dots, H-1$. Taking a supremum on both sides of (11) gives the recurrence

$$\mathbf{U}_h \leq \phi' + \frac{1}{2}\mathbf{U}_{h+1}^2, \quad 0 \leq h \leq H-1. \quad (12)$$

2. *Solution:* We show by induction on h (backwards from $H-1$) that

$$\mathbf{U}_h \leq 3\phi' + \left(\frac{1}{2}\right)^{-1+2^{H-h}}. \quad (13)$$

For the base of the induction $h = H-1$, we have indeed that

$$\mathbf{U}_{H-1} \leq \phi' + \frac{1}{2} \leq 3\phi' + \left(\frac{1}{2}\right)^{-1+2^{H-(H-1)}},$$

where we used (12) and the fact that \mathbf{u} -values are ≤ 1 in absolute value. Assume the induction claim (13) holds for all $h'+1 \leq h \leq H-1$. We show it then holds for $h = h'$. Indeed, by (12) again,

$$\begin{aligned} \mathbf{U}_{h'} &\leq \phi' + \frac{1}{2}\mathbf{U}_{h'+1}^2 \leq \phi' + \frac{1}{2} \left[3\phi' + \left(\frac{1}{2}\right)^{-1+2^{H-(h'+1)}} \right]^2 \\ &\leq \phi' \left[1 + \frac{9\phi'}{2} + 3 \left(\frac{1}{2}\right)^{-1+2^{H-(H-1)}} \right] + \left(\frac{1}{2}\right)^{2\{-1+2^{H-(h'+1)}\}+1} \\ &\leq \phi' \left[\frac{5}{2} + \frac{9\phi'}{2} \right] + \left(\frac{1}{2}\right)^{-1+2^{H-h'}}. \end{aligned}$$

Because by assumption $\phi' \leq 1/9$, the square bracket above is ≤ 3 . That concludes the induction.

By our choice of H in (9), that implies $|\mathbf{u}_v^\pi| = \mathbf{U}_0 \leq 3\phi' + (1/2)^{-1+2^H} \leq 4\phi'$. \square

3.2 Proof of main theorem

Proof of Theorem 1. Fix $\pi \in \mathcal{C}(\mathcal{T})$ and assume that $\theta_* > 0$ and that (1) holds. Then there is $0 < \phi' \leq 1/9$ and $0 < \zeta < 1$ such that

$$\sum_{x \in \pi'} \prod_{\rho \neq z \leq x} \left\{ \frac{2}{3}(1 - \phi') \right\} \theta_z \geq \zeta, \quad (14)$$

for all cutsets $\pi' \in \mathcal{C}(\mathcal{T})$. For this value of ϕ' , let ε' be as in Lemma 3 and define $\varepsilon = \varepsilon'\zeta < \varepsilon'$. The proof proceeds by contradiction. Assume that $\mathbf{d}_\rho^\pi \leq \varepsilon$. Let π' be the cutset of those nodes closest to the root where the \mathbf{d} -values first cross above ε' , i.e., formally $\pi' = \{x \in \mathcal{V}^\pi : \mathbf{d}_x^\pi > \varepsilon' \text{ and } \mathbf{d}_z^\pi \leq \varepsilon', \forall z \leq x\}$. Such a cutset (which is necessarily minimal) exists because $\mathbf{d}_v^\pi = 1$ for all $v \in \pi$ and $\varepsilon' > \varepsilon$. By Lemma 2, for all z on or above π' , i.e. such that $z \leq x$ for some $x \in \pi'$, we have $\mathbf{d}_z^\pi \leq \varepsilon' \frac{2}{\theta_*}$ and $\mathbf{d}_{s(z)}^\pi \leq \varepsilon' \frac{2}{\theta_*}$, since the immediate parent of z (and $s(z)$) has \mathbf{d} -value $\leq \varepsilon'$ by definition of π' . By Lemma 3, we then have

$$|\mathbf{u}_z^\pi| \leq 4\phi' \quad \text{and} \quad \left| \mathbf{u}_{s(z)}^\pi \right| \leq 4\phi'. \quad (15)$$

By Lemma 1, (14) and (15), summing over π'

$$\mathbf{d}_\rho^\pi = \sum_{x \in \pi'} \mathbf{d}_x^\pi \prod_{\rho \neq z \leq x} \left\{ \frac{4 - \mathbf{u}_{s(z)}^\pi}{6} \right\} \theta_z > \sum_{x \in \pi'} \varepsilon' \prod_{\rho \neq z \leq x} \left\{ \frac{2}{3}(1 - \phi') \right\} \theta_z \geq \varepsilon'\zeta = \varepsilon,$$

which is a contradiction. \square

4 I.i.d. weights

In this section, we prove our main result in the i.i.d. weight case. Because there is no lower bound on the weights, Theorem 1 cannot be applied directly to this case. In particular, the absence of a lower bound makes controlling the \mathbf{u} -values more challenging. Here we identify a subtree of \mathcal{T} where \mathbf{u} -values are well-behaved. The existence of such a subtree is established with a coupling to a percolation process, where open edges roughly indicate that weights are uniformly bounded in a properly defined neighborhood.

Proof of Theorem 2. First, we need the following percolation result. To each edge $e = (x, y)$ of \mathcal{T} , where x is the parent of y , we assign an independent random weight $\tilde{\theta}_y$ drawn from a distribution $\tilde{\Theta}$ over $(0, 1]$. We also pick an independent indicator variable \tilde{J}_y , which is 1 with probability $\tilde{q} \in [0, 1]$ and 0 otherwise. Let $\tilde{\mathcal{T}} = (\tilde{\mathcal{V}}, \tilde{\mathcal{E}})$ be the subtree of \mathcal{T} whose vertices x satisfy $\prod_{\rho \neq z \leq x} \tilde{J}_z = 1$. We let \mathcal{N}_{ext} be the event of non-extinction, i.e., the event that $\tilde{\mathcal{T}}$ is infinite. The following result can be proved along the lines of Proposition 3.2, Proposition 5.1 and Corollary 5.2 in [Per99].

Lemma 4 (Branching condition: random edge weights on open cluster). *Fix $\tilde{q} \in [0, 1]$ and assume that $\tilde{\Theta}$ has mean $\tilde{\mu} \in (0, 1)$. Then, conditioned on \mathcal{N}_{ext} , almost surely*

$$\sup \left\{ \kappa > 0 : \inf_{\pi \in \mathcal{C}(\tilde{\mathcal{T}})} \sum_{x \in \pi} \prod_{\rho \neq z \leq x} \kappa^{-1} \tilde{\theta}_z > 0 \right\} = 2\tilde{q}\tilde{\mu}. \quad (16)$$

By standard branching process arguments [AN72], the extinction probability $\tilde{\varphi}$ satisfies $\tilde{\varphi} = \tilde{q}^2 \tilde{\varphi}^2 + 2(1 - \tilde{q})\tilde{q}\tilde{\varphi} + (1 - \tilde{q})^2$, i.e.,

$$\mathbb{P}[\mathcal{N}_{\text{ext}}] = \left(\frac{1 - \tilde{q}}{\tilde{q}} \right)^2. \quad (17)$$

As discussed briefly above, we use a coupling argument. In order to describe the coupling, we first need to define some constants (not depending on θ). Recall that μ is the mean of the edge weight distribution Θ and that δ is the desired failure probability. Let $q \in (0, 1)$ be close enough to 1 that

$$\left(\frac{1 - q}{q} \right)^2 \leq \frac{\delta}{3}, \quad (18)$$

and $2q\mu > 3/2$. Let then $0 < \phi' \leq 1/9$ be such that

$$2q\mu > \frac{3}{2(1-\phi')}. \quad (19)$$

Let H be the smallest non-negative integer such that $(1/2)^{-1+2^H} \leq \phi'$. Because $\mathbb{P}_{\theta \sim \Theta}[\theta \leq \theta_*] \rightarrow 0$ as $\theta_* \rightarrow 0$, we can take $\theta_* \in (0, 1)$ small enough that $\tau = \mathbb{P}[\theta \leq \theta_*]$, satisfies

$$(1-\tau)^{2^{H+1}} > q, \quad \text{and} \quad 1 - (1-\tau)^{2^{H+2}} \leq \frac{\delta}{3}. \quad (20)$$

We are now ready to define the coupled process $(\tilde{\mathcal{T}}, \tilde{\theta})$. We use the following notation: $\text{Ber}(\psi)$ is a Bernoulli random variable with success probability ψ and $Z|\mathcal{A}$ is the random variable Z conditioned on the event \mathcal{A} . For each z , we generate $\theta_z = (1 - I_z)\theta_{z,0} + I_z\theta_{z,1}$, where $I_z \sim \text{Ber}(\mathbb{P}[\theta_z > \theta_*])$, $\theta_{z,0} \sim \theta_z|\{\theta_z \leq \theta_*\}$, and $\theta_{z,1} \sim \theta_z|\{\theta_z > \theta_*\}$ are independent. For $z \in \mathcal{V}$, let $\mathcal{D}_h(z)$ be the descendants of z lying exactly h levels below it, i.e., $\mathcal{D}_h(z) = \{w \in \mathcal{V} : z \leq w \text{ and } \gamma(z, w) = h\}$. Define $\tilde{J}_z = \prod_{w \in \mathcal{D}_{H+1}(z)} I_w$, and $\tilde{\theta}_z = \theta_{z,1}$. Note that by construction the random variables $\tilde{J}_z \sim \text{Ber}((1-\tau)^{2^{H+1}})$ and $\tilde{\theta}_z, z \in \mathcal{V}$, are independent. Let $\tilde{\mathcal{T}}$ be defined as in Lemma 4, let $\tilde{q} = (1-\tau)^{2^{H+1}}$ and let $\tilde{\mu}$ be the mean of Θ conditioned on being larger than θ_* . By (17), (18), and (20), $\mathbb{P}[\mathcal{N}_{\text{ext}}] \leq \frac{\delta}{3}$.

We apply Lemma 4 to $(\tilde{\mathcal{T}}, \tilde{\theta})$ to obtain a branching rate condition similar to (14).

Lemma 5 (Towards controlling \mathbf{d} at the root). *There is a deterministic (i.e., not depending on θ) $0 < \zeta < 1$ such that, on the event of non-extinction, with probability at least $1 - \delta/3$*

$$\inf_{\tilde{\pi}' \in \mathcal{C}(\tilde{\mathcal{T}})} \sum_{x \in \tilde{\pi}'} \prod_{\rho \neq z \leq x} \left\{ \frac{2}{3}(1-\phi') \right\} \tilde{\theta}_z \geq \zeta. \quad (21)$$

Proof. On the event of non-extinction, we have almost surely that

$$\sup \left\{ \kappa > 0 : \inf_{\pi \in \mathcal{C}(\tilde{\mathcal{T}})} \sum_{x \in \pi} \prod_{\rho \neq z \leq x} \kappa^{-1} \tilde{\theta}_z > 0 \right\} = 2\tilde{q}\tilde{\mu} > 2q\mu > \frac{3}{2(1-\phi')},$$

where we used the fact that $\tilde{\mu} > \mu$ as well as (19) and (20). In particular, we can choose a deterministic (i.e., not depending on θ) $0 < \zeta < 1$ such that, on the event of non-extinction, with probability at least $1 - \delta/3$ the inequality in (21) holds. \square

The purpose of the coupling is to show that the argument used in Lemma 3 to control the \mathbf{u} -values can be applied to the vertices in $\tilde{\mathcal{T}}$. This is stated in the next lemma. For the choice of ϕ' , H and θ_* above, let ε' be chosen as in the proof of Lemma 3, i.e., the largest positive real such that

$$\frac{3}{2} \left[\varepsilon' \left(\frac{2}{\theta_*} \right)^{H+1} \right]^2 \leq \phi', \quad \text{and} \quad \varepsilon' \left(\frac{2}{\theta_*} \right)^{H+1} \leq 0.99.$$

Lemma 6 (Controlling \mathbf{u} on $\tilde{\mathcal{T}}$). *For all $z \in \mathcal{V} \cap \tilde{\mathcal{V}}$*

$$\mathbf{d}_z^\pi \leq \varepsilon' \frac{2}{\theta_*} \implies |\mathbf{u}_z^\pi| \leq 4\phi' \text{ and } \mathbf{d}_{s(z)}^\pi \leq \varepsilon' \frac{2}{\theta_*} \implies |\mathbf{u}_{s(z)}^\pi| \leq 4\phi'. \quad (22)$$

Proof. First, observe that for any $z \in \mathcal{V} \cap \tilde{\mathcal{V}}$ it must be that $\theta_w > \theta_*$, $\forall w \in \cup_{h \leq H+1} \mathcal{D}_h(z)$. Indeed, note that the unique v such that $v \leq z$ and $\gamma(v, z) = h$ satisfies $v \in \tilde{\mathcal{T}}$ and $\mathcal{D}_{H+1-h}(z) \subseteq \mathcal{D}_{H+1}(v)$.

As a result, repeating the proof of Lemma 3, it follows that for all $z \in \mathcal{V} \cap \tilde{\mathcal{V}}$

$$\mathbf{d}_z^\pi \leq \varepsilon' \frac{2}{\theta_*} \implies |\mathbf{u}_z^\pi| \leq 4\phi' \quad \text{and} \quad \mathbf{d}_{s(z)}^\pi \leq \varepsilon' \frac{2}{\theta_*} \implies |\mathbf{u}_{s(z)}^\pi| \leq 4\phi',$$

where we used the fact that the parent of z is in $\tilde{\mathcal{T}}$ and therefore has all its descendants within $H + 1$ level with θ -values above θ_* . The latter in turn implies that $s(z)$ itself has all its descendants within H level with θ -values above θ_* . That concludes the proof of the lemma. \square

The rest of the proof of Theorem 2 is similar to that of Theorem 1, except that we restrict the argument to $\tilde{\mathcal{T}}$. Define $\varepsilon = \varepsilon' \zeta < \varepsilon'$. Let \mathcal{H} be the event that $\theta_w > \theta_*$ for all w such that $\gamma(\rho, w) \leq H$. By (20), $\mathbb{P}[\mathcal{H}] \geq 1 - \delta/3$. Condition on \mathcal{H} , \mathcal{N}_{ext} and (21), which jointly occur with probability at least $1 - \delta$. Assume by contradiction that $\mathbf{d}_\rho^\pi \leq \varepsilon$ and let $\pi' = \{x \in \mathcal{V}^\pi : \mathbf{d}_x^\pi > \varepsilon' \text{ and } \mathbf{d}_z^\pi \leq \varepsilon', \forall z \leq x\}$. By Lemma 2, for all $z \in \mathcal{V} \cap \tilde{\mathcal{V}}$ such that $z \leq x$ for some $x \in \pi'$, we have $\mathbf{d}_z^\pi \leq \varepsilon' \frac{2}{\theta_*}$ and $\mathbf{d}_{s(z)}^\pi \leq \varepsilon' \frac{2}{\theta_*}$, since the immediate parent of z (and $s(z)$) has \mathbf{d} -value $\leq \varepsilon'$ by definition of π' and θ -values on $\tilde{\mathcal{T}}$ are above θ_* . By (22), we then have

$$|\mathbf{u}_z^\pi| \leq 4\phi' \quad \text{and} \quad |\mathbf{u}_{s(z)}^\pi| \leq 4\phi', \quad (23)$$

for all $z \in \mathcal{V} \cap \tilde{\mathcal{V}}$ such that $z \leq x$ for some $x \in \pi'$. Similarly to Theorem 1, by Lemma 1, (21) and (23),

$$\mathbf{d}_\rho^\pi = \sum_{x \in \pi'} \mathbf{d}_x^\pi \prod_{\rho \neq z \leq x} \left\{ \frac{4 - \mathbf{u}_{s(z)}^\pi}{6} \right\} \theta_z \geq \sum_{x \in \pi' \cap \tilde{\mathcal{V}}} \mathbf{d}_x^\pi \prod_{\rho \neq z \leq x} \left\{ \frac{4 - \mathbf{u}_{s(z)}^\pi}{6} \right\} \tilde{\theta}_z \geq \varepsilon' \zeta = \varepsilon,$$

where we used again that θ -values on $\tilde{\mathcal{T}}$ are above θ_* as well as the fact that all terms in the sum are non-negative by the general bounds on the \mathbf{d} - and \mathbf{u} -values. This is a contradiction. \square

Acknowledgments

We thank Mike Steel for helpful discussions and an anonymous reviewer for suggested improvements to a previous version of the manuscript.

References

- [ABDLV20] Louigi Addario-Berry, Luc Devroye, Gabor Lugosi, and Vasiliki Velona. Broadcasting on random recursive trees. *arXiv:2006.11787 [math]*, June 2020. arXiv: 2006.11787.
- [AN72] K. B. Athreya and P. E. Ney. *Branching processes*. Springer-Verlag, New York, 1972. Die Grundlehren der mathematischen Wissenschaften, Band 196.
- [BCMR06] Christian Borgs, Jennifer Chayes, Elchanan Mossel, and Sébastien Roch. The Kesten-Stigum reconstruction bound is tight for roughly symmetric binary channels. In *FOCS*, pages 518–530, 2006.
- [BRZ95] P. M. Bleher, J. Ruiz, and V. A. Zagrebnov. On the purity of the limiting Gibbs state for the Ising model on the Bethe lattice. *J. Statist. Phys.*, 79(1-2):473–482, 1995.

- [BST16] Nayantara Bhatnagar, Allan Sly, and Prasad Tetali. Decay of correlations for the hardcore model on the d -regular random graph. *Electronic Journal of Probability*, 21, 2016. Publisher: The Institute of Mathematical Statistics and the Bernoulli Society.
- [CH11] Juanjuan Chai and Elizabeth Ann Housworth. On the Number of Binary Characters Needed to Recover a Phylogeny Using Maximum Parsimony. *Bulletin of Mathematical Biology*, 73(6):1398–1411, June 2011.
- [CR14] Natalie Cusimano and Susanne S. Renner. Ultrametric trees or phylograms for ancestral state reconstruction: Does it matter? *Taxon*, 63(4):721–726, 2014. Publisher: International Association for Plant Taxonomy (IAPT).
- [DMR11] Constantinos Daskalakis, Elchanan Mossel, and Sébastien Roch. Evolutionary trees and the Ising model on the Bethe lattice: a proof of Steel’s conjecture. *Probability Theory and Related Fields*, 149:149–189, 2011. 10.1007/s00440-009-0246-2.
- [EKPS00] W. S. Evans, C. Kenyon, Y. Peres, and L. J. Schulman. Broadcasting on trees and the Ising model. *Ann. Appl. Probab.*, 10(2):410–433, 2000.
- [FR18] Wai-Tong (Louis) Fan and Sébastien Roch. Necessary and sufficient conditions for consistent root reconstruction in Markov models on trees. *Electronic Journal of Probability*, 23, 2018. Publisher: The Institute of Mathematical Statistics and the Bernoulli Society.
- [FR20] Wai-Tong Fan and Sébastien Roch. Statistically consistent and computationally efficient inference of ancestral DNA sequences in the TKF91 model under dense taxon sampling. *Bulletin of Mathematical Biology*, 82(2):21, January 2020.
- [FT09] Mareike Fischer and Bhalchandra D. Thatte. Maximum parsimony on subsets of taxa. *Journal of Theoretical Biology*, 260(2):290–293, September 2009.
- [GRP20] Yuzhou Gu, Hajir Roozbehani, and Yury Polyanskiy. Broadcasting on trees near criticality. *arXiv:2005.07801 [cs, math]*, May 2020. arXiv: 2005.07801.
- [GS10] Olivier Gascuel and Mike Steel. Inferring ancestral sequences in taxon-rich phylogenies. *Math. Biosci.*, 227(2):125–135, 2010.
- [GS14] Olivier Gascuel and Mike Steel. Predicting the Ancestral Character Changes in a Tree is Typically Easier than Predicting the Root State. *Systematic Biology*, 63(3):421–435, May 2014.
- [HF17] Lina Herbst and Mareike Fischer. Ancestral Sequence Reconstruction with Maximum Parsimony. *Bulletin of Mathematical Biology*, 79(12):2865–2886, December 2017.
- [HF18] Lina Herbst and Mareike Fischer. On the Accuracy of Ancestral Sequence Reconstruction for Ultrametric Trees with Parsimony. *Bulletin of Mathematical Biology*, 80(4):864–879, April 2018.
- [HLS19] Lina Herbst, Heyang Li, and Mike Steel. Quantifying the accuracy of ancestral state prediction in a phylogenetic tree under maximum parsimony. *Journal of Mathematical Biology*, 78(6):1953–1979, May 2019.

- [Iof96] D. Ioffe. On the extremality of the disordered state for the Ising model on the Bethe lattice. *Lett. Math. Phys.*, 37(2):137–143, 1996.
- [KS66] H. Kesten and B. P. Stigum. Additional limit theorems for indecomposable multidimensional Galton-Watson processes. *Ann. Math. Statist.*, 37:1463–1481, 1966.
- [LJN18] Wenjian Liu, Sreenivasa Rao Jammalamadaka, and Ning Ning. The Tightness of the Kesten-Stigum Reconstruction Bound of Symmetric Model with Multiple Mutations. *Journal of Statistical Physics*, 170(3):617–641, February 2018.
- [LP16] Russell Lyons and Yuval Peres. *Probability on Trees and Networks*, volume 42 of *Cambridge Series in Statistical and Probabilistic Mathematics*. Cambridge University Press, New York, 2016.
- [MMP20] A. Makur, E. Mossel, and Y. Polyanskiy. Broadcasting on Random Directed Acyclic Graphs. *IEEE Transactions on Information Theory*, 66(2):780–812, February 2020. Conference Name: IEEE Transactions on Information Theory.
- [Mos98] E. Mossel. Recursive reconstruction on periodic trees. *Random Struct. Algor.*, 13(1):81–97, 1998.
- [Mos01] E. Mossel. Reconstruction on trees: beating the second eigenvalue. *Ann. Appl. Probab.*, 11(1):285–300, 2001.
- [Mos04] E. Mossel. Phase transitions in phylogeny. *Trans. Amer. Math. Soc.*, 356(6):2379–2404, 2004.
- [MP03] E. Mossel and Y. Peres. Information flow on trees. *Ann. Appl. Probab.*, 13(3):817–844, 2003.
- [MRS13] E. Mossel, S. Roch, and A. Sly. Robust Estimation of Latent Tree Graphical Models: Inferring Hidden States With Inexact Parameters. *IEEE Transactions on Information Theory*, 59(7):4357–4373, July 2013. Conference Name: IEEE Transactions on Information Theory.
- [MS14] Elchanan Mossel and Mike Steel. Majority rule has transition ratio 4 on Yule trees under a 2-state symmetric model. *Journal of Theoretical Biology*, 360:315–318, November 2014.
- [Per99] Yuval Peres. Probability on trees: an introductory climb. In *Lectures on probability theory and statistics (Saint-Flour, 1997)*, volume 1717 of *Lecture Notes in Math.*, pages 193–280. Springer, Berlin, 1999.
- [RS17] Sebastien Roch and Allan Sly. Phase transition in the sample complexity of likelihood-based phylogeny inference. *Probability Theory and Related Fields*, 169(1):3–62, October 2017.
- [Sly11] Allan Sly. Reconstruction for the Potts model. *Ann. Probab.*, 39(4):1365–1406, 07 2011.
- [SStY17] Alistair Sinclair, Piyush Srivastava, Daniel ?tefankovi?, and Yitong Yin. Spatial mixing and the connective constant: optimal bounds. *Probability Theory and Related Fields*, 168(1):153–197, June 2017.
- [Ste89] M. Steel. *Distributions on bicoloured evolutionary trees*. PhD thesis, Massey University, 1989.

- [Ste16] Mike Steel. *Phylogeny—discrete and random processes in evolution*, volume 89 of *CBMS-NSF Regional Conference Series in Applied Mathematics*. Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA, 2016.
- [TS97] C. Tuffley and M. Steel. Links between maximum likelihood and maximum parsimony under a simple model of site substitution. *Bull. Math. Biol.*, 59(3):581–607, 1997.
- [ZSYL10] Louxin Zhang, Jian Shen, Jialiang Yang, and Guoliang Li. Analyzing the Fitch method for reconstructing ancestral states of ultrametric phylogenetic trees. *Bull. Math. Biol.*, 72(7):1760–1782, 2010.