BOUNDED CONTRACTIONS FOR AFFINE BUILDINGS

MLADEN BESTVINA AND GORDAN SAVIN

ABSTRACT. We consider affine buildings with refined chamber structure. For each vertex x we construct a contraction, based at x, that is used to prove exactness of Schneider-Stuhler resolutions of arbitrary depth.

1. INTRODUCTION

Let k be a local non-archimedean field and G the group of k-points of an algebraic reductive group defined over k. Let X be the Bruhat-Tits building attached to G, which we view as a CAT(0) metric space. In [MP2], to every $x \in X$ and non-negative real number r, Moy and Prasad attach a subgroup $G_{x,r}$ such that $G_x := G_{x,0}$ is the parahoric subgroup of G attached to x, $G_{x,s} \subseteq G_{x,r}$ whenever $r \leq s$, and $G_{x,r}$ are normal subgroups of G_x . For every $r \geq 0$ we let $G_{x,r,+} = \bigcup_{s>r} G_{x,s}$. The building X has a (standard) structure of a chamber complex such that, for every facet σ and r an integer, the function $x \mapsto G_{x,r,+}$ is constant for x in the interior of σ . Let r be a rational number. Then the chamber structure can be refined so that $x \mapsto G_{x,r,+}$ is constant on the interior points of each facet [BKV]. For example, if $G = SL_2(k)$, the building X is a tree and if we divide each edge into two, of equal lengths, then $x \mapsto G_{x,r,+}$ is constant on the interior points of each edge if r is half-integral. Fix r and a refined chamber structure on X. Let $C_i(X)$ be the free abelian group with the basis consisting of the *i*-dimensional facets of X. Let $C_{-1}(X) = \mathbb{Z}$. Let $x \in X$ be a vertex. Recall that a contraction c of $C_*(X)$ based at x is a sequence of homomorphisms $c_i: C_i(X) \to C_{i+1}(X), i = -1, 0, 1, \cdots$, such that $c_{-1}(1) = x$ and $c_{i-1}\partial + \partial c_i = 1$. Our main technical result is the construction of a contraction of $\sigma \to c(\sigma) = \sum c(\sigma, \tau)\tau$ with the following properties:

- (1) the contraction c is G_x -equivariant.
- (2) if $c(\sigma, \tau) \neq 0$ then τ is contained in the smallest subcomplex of X containing the cone with the vertex x and the base σ .
- (3) the coefficients $c(\sigma, \tau)$ are uniformly bounded.

A more detailed meaning of (2) is the following. The cone is the union of all geodesic segments connecting x to a point in σ . If $c(\sigma, \tau) \neq 0$ then there exists a point $y \in \sigma$ and a point z on the geodesic [x, y] such that z is either an interior point of τ or τ is in the boundary of a facet containing z as an interior point.

Let V be a smooth representation of G of depth r. Following Schneider-Stuhler [SS] we define a projective resolution of V using the chain complex $C_*(X)$. Since we use a refined

M.B. partially supported by NSF grant DMS-1607236.

G.S. partially supported by NSF grant DMS-1359774.

chamber structure on X the projective modules in the resolution have the same depth r. The exactness of resolution is a simple consequence of the existence of the contraction c and niceness of Moy-Prasad groups $G_{x,r}$ if r > 0. An open compact subgroup K of G is nice if, for any smooth representation V of G generated by its K-fixed vectors, any subquotient of V is generated by its K-fixed vectors. It is recorded and used in the literature that $G_{x,r}$ is nice if x is a special point in the building. Niceness of all positive depth Moy-Prasad groups is possibly known to experts, however, we have not found a precise reference in the literature and hence have included a proof. We thank C. Bushnell and G. Henniart for a discussion on this matter. The property (3) is not used in this paper, however, it is critical to prove that the resolution stays exact after passing to a Schwartz completion, see [OS] for details.

2. LIPSCHITZ SIMPLICIAL APPROXIMATION

We fix a cell structure A on \mathbb{R}^n so that each (closed) cell σ is a convex polyhedron, and when $\tau \subset \sigma$ are cells in A then τ is a polyhedral face of σ . For simplicity, we will assume that A is invariant under a discrete group of translations of rank n, although the arguments below apply when A has bounded combinatorics and contains only finitely many isometry types of cells. We also fix an orientation of each cell. By A^i we denote the *i*-skeleton of A. Consider the augmented cellular chain complex $C_*(A)$ of A where $C_i(A)$ is the free abelian group with the basis consisting of the *i*-cells of A, and $C_{-1}(A) = \mathbb{Z}$. The boundary morphisms are defined by

$$\partial \sigma = \sum_{\tau} \epsilon(\sigma, \tau) \tau$$

where the sum runs over codimension 1 faces τ of σ and $\epsilon(\sigma, \tau) = 1$ if the orientation of σ is obtained from the orientation of τ by appending an outward normal to τ , and otherwise $\epsilon(\sigma, \tau) = -1$.

We fix a vertex v_0 of A^0 . Without loss of generality we can assume that $v_0 = 0$. Recall that a contraction c of $C_*(A)$ based at v_0 is a sequence of homomorphisms $c_i : C_i(A) \to C_{i+1}(A)$, $i = -1, 0, 1, \cdots$, where $c_{-1}(1) = v_0$, such that $c_{i-1}\partial + \partial c_i = 1$. The goal of this section is to construct $\sigma \to c(\sigma) = \sum c(\sigma, \tau)\tau$ with the coefficients $c(\sigma, \tau)$ uniformly bounded and $c(\sigma, \tau) \neq 0$ only if τ is contained in every convex subcomplex of A containing v_0 and σ .

The standard method of deforming a map $f : \mathbb{R}^k \to \mathbb{R}^n$ so it misses a given cell σ of dimension > k is to radially project to $\partial \sigma$ from a point x_{σ} in the interior of σ and in the complement of the image of f. For the purposes of controlling the coefficients $c(\sigma, \tau)$ in the contraction we need to control the Lipschitz constant, which will dramatically increase if x_{σ} is close to the image of f.

Proposition 2.1. For every N, L > 0 there is L' > 0 with the following properties. Let $D \subset \mathbb{R}^k$ be a polyhedral cell and $f: D \to \mathbb{R}^n$ a map such that

- f is L-Lipschitz,
- $f(\partial D) \subset A^{k-1}$,
- $f(D) \cap \sigma$ is contained in the union of N k-planes for every cell σ of A.

Then there is a map $g: D \to \mathbb{R}^n$ such that

- (i) g = f on ∂D , and on $f^{-1}(A^{k-1})$,
- (ii) g is L'-Lipschitz,
- (iii) for every $x \in D$, g(x) belongs to the closure of the open cell of A that contains f(x), (iv) $g(D) \subset A^k$.

Proof. We inductively construct maps $f = g_n, g_{n-1}, \dots, g_k = g : D \to \mathbb{R}^n$ satisfying (i), (iv), (modified) (ii)-(iii): $g_i(D) \subset A^i$, g_i is L_i -Lipschitz, and in addition $g_i(D) \cap \sigma$ is contained in the union of N_i *i*-planes. The construction from g_i to g_{i-1} is as follows.

By volume considerations, there is $\epsilon > 0$ that depends only on N_i and A, so that for every *i*-cell σ there is a point $x_{\sigma} \in \overset{\circ}{\sigma}$ at distance $> \epsilon$ from $g_i(D) \cap \sigma$ and also distance $> \epsilon$ from $\partial \sigma$. Compose g_i with the map which is the radial projection from x_{σ} to $\partial \sigma$ on σ for each *i*-simplex σ . This composition is g_{i-1} . The Lipschitz constant L_{i-1} of g_{i-1} is uniformly bounded, more precisely,

$$L_{i-1} \le \left(\frac{d}{\epsilon}\right)^2 L_i$$

where d is a bound on the diameter of all simplices, and $N_{i-1} \leq N_i M_i P_i$, where M_i is the maximal number of codimension 1 faces of a cell in A and P_i is the maximal number of cells intersecting a given cell.

Proposition 2.2. Let $v_0 = 0$ be a vertex of A. For every cell σ of A there is a homotopy $H_{\sigma}: \sigma \times [0, D_{\sigma}] \to \mathbb{R}^n$, with $D_{\sigma} = \text{diam}(\sigma \cup \{v_0\})$, so that the following holds.

(1) $H_{\sigma}(x,0) = v_0$ for every $x \in \sigma$,

- (2) $H_{\sigma}(\cdot, 1)$ is the inclusion,
- (3) the image of H_{σ} is contained in $A^{\dim(\sigma)+1}$, and it is also contained in every convex subcomplex of A that contains v_0 and σ ,
- (4) the restriction of H_{σ} to $\tau \times [0, D_{\sigma}]$ for a face $\tau < \sigma$ is the linear reparametrization of H_{τ} ,
- (5) H_{σ} is uniformly Lipschitz,
- (6) H_{σ} is uniformly bounded distance away from the straight line homotopy $(x, t) \mapsto tx/D_{\sigma}$.

In the proof we will need the following lemma.

Lemma 2.3. For every $L_0, \epsilon, D, \alpha_0 > 0$ there exists L' > 0 such that the following holds. Let $\sigma \subset \mathbb{R}^k$ be a convex polyhedral cell, $v \in \mathbb{R}^k$, $f : \sigma \to \mathbb{R}^n$, $w \in \mathbb{R}^n$ such that:

(a) f is L-Lipschitz,

(b) $|v - x| > \epsilon$ for every $x \in \sigma$,

(c) $|w - y| \le D$ for every $y \in f(\sigma)$,

(d) $\angle_y(x,v) > \alpha_0$ for every $x, y \in \sigma, x \neq y$.

Then the map $F: v * \sigma \to \mathbb{R}^n$ defined on the cone by

$$F((1-t)v + tx) = (1-t)w + tf(x)$$

is L'-Lipschitz.

Proof. We may assume v = w = 0. The extension F is clearly L-Lipschitz when restricted to each slice $t\sigma$ by (a), and by (b) and (c) F is also uniformly Lipschitz when restricted to each radial line $\{tx \mid t \in [0, 1]\}$ for any $x \in \sigma$. Any two points in the cone $v * \sigma$ can be connected by a segment in a slice followed by a segment in a radial line. By (d) the angle between these segments is bounded below, so the total length is bounded by a fixed multiple of the distance between the two points, implying the result.

Proof of Proposition 2.2. The construction is by induction on dim(σ). When σ is a vertex apply Proposition 2.1 to the geodesic f joining v_0 to σ . This gives $g = H_{\sigma} : [0, D_{\sigma}] \to \mathbb{R}^n$.

Assume now that $\dim(\sigma) = k$ and H_{τ} has been constructed for all faces $\tau < \sigma$. Define $f: \partial(\sigma \times [0, D_{\sigma}]) \to \mathbb{R}^n$ to be constant $v_0 = 0$ on $\sigma \times \{0\}$, inclusion on $\sigma \times \{D_{\sigma}\}$, and linearly reparametrized H_{τ} on $\tau \times [0, D_{\sigma}]$, for every face $\tau < \sigma$. We now wish to extend f to all of $\sigma \times [0, D_{\sigma}]$ and apply Proposition 2.1 to this extension. If we extend by coning off from a point in the interior of $\sigma \times [0, D_{\sigma}]$, the Lipschitz constant might blow up since the angles as in Lemma 2.3(d) will be small when D_{σ} is large. Instead, we first subdivide $[0, D_{\sigma}]$ into $0 = t_0 < t_1 < \cdots < t_{\sigma} = D_{\sigma}$ so that the length $t_i - t_{i-1}$ of each segment belongs to a fixed interval $[B_1, B_2]$ with $B_1 > 0$. Then we use Lemma 2.3 to extend f to $\sigma \times \{t_i\}$ for each *i*. For this we use $v_i = (x_{\sigma}, t_i)$ for a point x_{σ} in the interior of σ , and we use the same point for this isometry type. The image $f(\partial \sigma) \times \{t_i\}$ has uniformly bounded diameter, and we can set $w_i \in \mathbb{R}^n$ to be any point in this image. Because of (5) and (6), the assumptions of Lemma 2.3 are satisfied and we get uniformly Lipschitz extensions of f. It remains to extend f to each $\sigma \times [t_{i-1}, t_i]$ this is done in exactly the same way, by coming from $(x_{\sigma}, \frac{t_{i-1}, t_i}{2})$ with respect to a point in the image of $\partial(\sigma \times [t_{i-1}, t_i])$. Thus the extension $\sigma \times [0, D_{\sigma}] \to \mathbb{R}^n$ is uniformly Lipschitz. The third bullet in Proposition 2.1 holds since only a bounded number of images of $\sigma \times [t_{i-1}, t_i]$ intersect a given cell. Thus we can apply Proposition 2.1 to this extension to get the desired map H_{σ} .

When σ is a k-cell and τ a (k + 1)-cell (both oriented) we denote by $c(\sigma, \tau)$ the degree of the map

$$(\sigma \times [0, D_{\sigma}], \partial (\sigma \times [0, D_{\sigma}])) \xrightarrow{H_{\sigma}} (A^{k+1}, A^{k+1} \smallsetminus \overset{\circ}{\tau})$$

(the latter set is by excision equivalent to $(\tau, \partial \tau)$).

Proposition 2.4. The numbers $c(\sigma, \tau) \in \mathbb{Z}$ are uniformly bounded.

Proof. A warmup is the fact that any map $S^n \to S^n$ of large degree must have a large Lipschitz constant. This follows from Arzela-Ascoli. The set of maps $S^n \to S^n$ with Lipschitz constants bounded above is compact and hence represents finitely many homotopy classes.

The proof in our relative case is similar. Suppose there is a sequence of pairs (σ_i, τ_i) with $|c(\sigma_i, \tau_i)| \to \infty$. We may assume all σ_i are isometric and have dimension k, and all τ_i are isometric. By (5) the preimage of τ_i under H_{σ_i} is contained in $\sigma_i \times [u_i, u_i + C]$ for some fixed C independent of i. The image of $\sigma_i \times [u_i, u_i + C]$ is contained in a subcomplex $Y_i \subset A^{k+1}$ of τ_i and up to isomorphism there are only finitely many possibilities.

After translating the interval and identifying all σ_i and all τ_i and the Y_i as above we have a sequence of maps $\sigma \times [0, C] \to Y$, and these all have uniformly bounded Lipschitz

constants. By Arzela-Ascoli after a subsequence they will be close to each other and will determine the same degree. Contradiction. $\hfill \Box$

Now $\sigma \to c(\sigma) = -\sum c(\sigma, \tau)\tau$ defines a contraction.

Assume that X is a building corresponding to a reductive group G over a p-adic field k, with a refined chamber structure i.e. we have divided chambers into smaller ones in a G-equivariant fashion. Let x be a vertex in G and G_x the (largest) parahoric group fixing x. For every facet $\sigma \subset X$ the cone with the vertex x and the base σ is contained in any apartment A containing x and σ . Thus the contraction $c(\sigma)$ can be defined working in any such apartment. If $g \in G_x$ stabilizes σ then g fixes σ point-wise, hence also the cone and $c(\sigma)$. Hence c can be extended to whole G_x -orbit of σ in an G_x -equivariant fashion, and we have proved:

Proposition 2.5. For every vertex x in X there exists a G_x -invariant contraction c satisfying the conditions (1)-(3) in Introduction.

3. Moy-Prasad groups

Let k be a non-archimedean local field and G the group of k points of a reductive algebraic group defined over k. Let S be a maximal k-split torus in G and $\Phi = \Phi(G, S)$ the corresponding restricted root system. Let T be the centralizer of S in G. A decomposition $\Phi = \Phi^+ \cup \Phi^-$ of the root system into positive and negative roots defines a pair of maximal unipotent subgroups U and \overline{U} of G. Let A(S) be the apartment in the building X of G stabilized by S. Let $x \in A(S)$ and r > 0. The Moy-Prasad group $G_{x,r}$ has an Iwahori decomposition ([MP2] Theorem 4.2)

$$G_{x,r} = \bar{U}_{x,r}T_r U_{x,r}$$

where the three factors are the intersection of $G_{x,r}$ with \overline{U} , T and U, respectively. The factor T_r is independent of x as indicated.

Lemma 3.1. Let $x, y \in X$ and z be a point on the geodesic connecting x and y. If r > 0 then

$$G_{z,r} \subseteq G_{x,r}G_{y,r}.$$

Proof. Without loss of generality we can assume that $x, y \in A(S)$. The apartment A(S) is an affine space so v = x - y is a well defined vector in the space of translations of A(S). Elements of Φ (roots) are functionals on the space of translations of A(S). Thus $\alpha(v)$ is a well defined real number for every $\alpha \in \Phi$. Pick a decomposition $\Phi = \Phi^+ \cup \Phi^-$ such that $\alpha(v) \ge 0$ for all $\alpha \in \Phi^+$. Then

$$\overline{U}_{x,r} \supseteq \overline{U}_{z,r}$$
 and $U_{y,r} \supseteq U_{z,r}$.

This inclusions are a direct consequence of the definition of $G_{x,r}$ [MP1] if G is quasi-split. The general case is then deduced by checking it over a finite unramified extension of k, over which G is quasi-split, and then taking fixed points for the Galois action. Lemma follows from the Iwahori decomposition of $G_{z,r}$. The Iwahori decomposition and Lemma 3.1 hold for the Groups $G_{x,r,+}$ (for $r \ge 0$) since $G_{x,r,+} = G_{x,s}$ for all s > r and sufficiently close to r. Now fix a non-negative rational number r. Refine the chamber decomposition of X so that the function $x \mapsto G_{x,r,+}$ is constant on interiors of facets. Thus, for any facet σ we define $K_{\sigma} = G_{x,r,+}$ where x is any interior point of σ . If τ is a facet in the boundary of σ then $K_{\tau} \subseteq K_{\sigma}$. The refinement of X and a check of these inclusions is done explicitly for quasi-split groups. We shall give details in the split case below. The general case follows by unramified Galois descent.

Lemma 3.2. Let x be a vertex of X and σ a facet for a refined decomposition of X. Assume that a facet τ is contained in the smallest subcomplex of X containing the cone with the vertex x and the base σ . Then

$$K_{\tau} \subseteq K_x K_{\sigma}$$

Proof. By the assumption, there exists a point $y \in \sigma$ and a point z on the geodesic [x, y] such that the facet containing z, as an interior point, contains τ . Then $K_{\tau} \subseteq G_{z,r,+}$ and $G_{y,r,+} \subseteq K_{\sigma}$. Lemma follows from Lemma 3.1.

We shall work out the details in the case G is simple and split. (A similar treatment for quasi split groups can be easily given based on computations in [PR].) Since G is split, S is a maximal torus, hence T = S. Let O be the ring of integers in k and π and a uniformizing element. The group k^{\times} has a natural filtration $k^{\times} \supset O^{\times} \supset 1 + \pi O \supset 1 + \pi^2 O \supset \ldots$ which gives rise to a filtration

$$T \supset T_0 \supset T_1 \supset \ldots$$

where T_0 is the maximal compact subgroup, which is the same as the set of all $t \in T$ such that $\chi(t) \in O^{\times}$ for all algebraic characters χ of T, and T_r is the set of all $t \in T$ such that $\chi(t) \in 1 + \pi^r O$ for all algebraic characters χ . The apartment A = A(S) can be identified with $\operatorname{Hom}(\mathbb{G}_m, S) \otimes \mathbb{R}$, so 0 is a special vertex of A. The apartment is a Coxeter complex defined by an affine root system $\Psi = \bigcup_{n \in \mathbb{Z}} \Phi + n$.

For every root $\alpha \in \Phi$ we have a subgroup U_{α} of G isomorphic to k, and for every affine root a we have a subgroup isomorphic to O. More precisely, If a is an affine root whose gradient is α , then U_a is defined as the subgroup of U_{α} consisting of elements fixing the half-plane $a \geq 0$. The filtration

$$\ldots \supset U_{a-1} \supset U_a \supset U_{a+1} \supset \ldots$$

of U_{α} corresponds to the filtration $\ldots \supset \pi^{-1}O \supset O \supset \pi O \supset \ldots$ of k. For example, if $G = \operatorname{SL}_2(k)$ and T the torus of diagonal matrices, then one can identify A with \mathbb{R} and the set of affine roots with the set of affine functions $x \mapsto \pm x + n$, where $n \in \mathbb{Z}$, so that for a(x) = x + n

$$U_a = \left\{ \begin{bmatrix} 1 & x \\ 0 & 1 \end{bmatrix} \mid x \in \pi^n O \right\} \text{ and } U_{-a} = \left\{ \begin{bmatrix} 1 & 0 \\ x & 1 \end{bmatrix} \mid x \in \pi^{-n} O \right\}$$

It is convenient to introduce a notion of imaginary roots. The imaginary roots are constant, integer-valued functions on A. Let

$$\Psi = \Psi \cup \mathbb{Z}.$$

If $a \in \Psi$ is an imaginary non-negative root, we define $U_a = T_a \subseteq T$. Let $x \in A$ and $r \geq 0$, a real number. Then $G_{x,r}$ and $G_{x,r,+}$ are generated by U_a for all $a \in \tilde{\Psi}$ such that $a(x) \geq r$ and a(x) > r, respectively. If we fix a decomposition $\Phi = \Phi^+ \cup \Phi^-$ then the groups $U_{x,r}$ and $\bar{U}_{x,r}$, appearing in the Iwahori decomposition, are generated by U_a such that $a(x) \geq r$ and the gradient of a is in Φ^+ and Φ^- , respectively. Now the inclusions in the proof of Lemma 3.1 are clear.

If r is an integer and a(x) > r for one interior point of a facet σ then the same is true for any point in the interior of σ . Thus the groups $G_{x,r,+}$ are constant on interiors of facets. Moreover, if τ is contained in the boundary of σ , and a > r on the interior of τ then it is so on the interior of σ . Hence $K_{\tau} \subseteq K_{\sigma}$. Finally, assume $r = \frac{n}{m}$ is a non-negative rational number. We can refine the Coxeter complex by replacing Ψ with $\Psi_m = \bigcup_{n \in \mathbb{Z}} \Phi + \frac{n}{m}$. Now the groups $G_{x,r,+}$ are constant on interiors of facets.

4. Schneider-Stuhler complex

We fix a non-negative rational number r throughout this section. We refine the chamber decomposition of X so that the groups $G_{x,r,+}$ are constant on the interior of facetes and we set $K_{\sigma} := G_{x,r,+}$ where x is any interior point of a facet σ . Let V be a smooth representation of G, this means that any vector $v \in V$ is fixed by an open compact subgroup of G, depending on v. For every facet σ let V_{σ} be the subspace of all vectors in V fixed by K_{σ} . The complex $C_*(X) \otimes_{\mathbb{Z}} V$ admits a natural representation of G defined by $g(\sigma \otimes v) = g(\sigma) \otimes g(v)$, for all $g \in G$. Let $C_*(X, V)$ be the subcomplex spanned by $\tau \otimes v$ where $v \in V_{\tau}$. The boundary ∂ preserves $C_*(X, V)$ because $V_{\sigma} \subseteq V_{\rho}$ any time ρ is in the boundary of σ . The action of G on $C_*(X) \otimes V$ preserves the sub complex $C_*(X, V)$.

Theorem 4.1. Let x be a vertex in X and c an x-based contraction of $C_*(X)$ satisfying the conditions (1) and (2) in the Introduction. Then the complex $C_*(X, V)^{K_x}$ is exact.

Proof. The contraction c defines a contraction on $C_*(X) \otimes V$ by $c(\sigma \otimes v) = c(\sigma) \otimes v$. To prove the theorem, it suffices to show that the contraction preserves the subcomplex $C_*(X,V)^{K_x}$. Let $e_x : V \to V_x$ be the projection given by averaging the action of K_x on V with respect to a Haar measure on K_x of volume one. The subcomplex $C_*(X,V)^{K_x}$ is spanned by elements $e_x(\sigma \otimes v)$ where $v \in V_{\sigma}$. Since c is K_x -invariant,

$$c(e_x(\sigma \otimes v)) = e_x(c(\sigma) \otimes v) = \sum_{\tau} c(\sigma, \tau) e_x(\tau \otimes v).$$

By the property (2) and Lemma 3.2, for every τ appearing in this sum, there exist $g_1, \ldots, g_n \in K_x \cap K_\tau$ such that

$$K_{\tau} = \bigcup_{i=1}^{n} g_i (K_{\sigma} \cap K_{\tau})$$

(a disjoint sum). Since $g_i \in K_x$, it follows that $e_x \cdot g_i = e_x$, as operators on any representation of G. Hence

$$e_x(\tau \otimes v) = \frac{1}{n} \sum_{i=1}^n e_x g_i(\tau \otimes v).$$

Since $g_i \in K_{\tau}$, these elements fix τ . Hence

$$e_x(\tau \otimes v) = e_x(\tau \otimes \frac{1}{n} \sum_{i=1}^n g_i v).$$

Since v is fixed by $K_{\sigma} \cap K_{\tau}$ and g_i are representatives of all $K_{\sigma} \cap K_{\tau}$ -cosets in K_{τ} , it follows that

$$\frac{1}{n}\sum_{i=1}^{n}g_iv = e_\tau(v) \in V_\tau$$

where $e_{\tau}: V \to V_{\tau}$ is the projection given by averaging the action of K_{τ} on V with respect to a Haar measure on K_{τ} of volume one. Hence $e_x(\tau \otimes v) \in C_*(X, V)^{K_x}$ as desired. \Box

5. NICE OPEN COMPACT SUBGROUPS

Let S be a maximal split torus in G. Let P be a parabolic subgroup of G. Without loss of generality we shall assume that P contais S. In particular, we have a "standard" choice of the Levi L and the opposite \overline{U} of the radical U of P, both normalized by S. Let K be a an open compact subgroup of G and K^G the set of all G-conjugates of K. The parabolic group P acts on K^G with finitely many orbits. We say that K is nice with respect to P if in any P-conjugacy class in K^G there is K' such that the Iwahori decomposition holds:

$$K' = (K' \cap U)(K' \cap L)(K' \cap \overline{U}).$$

If that is the case, then $K'_L = K' \cap L$ is isomorphic to $K' \cap P/K' \cap U$. This observation implies the first of the following properties, for the second see Proposition 3.5.2. in [BD].

- (1) If W is an L-module and $(\operatorname{Ind}_P^G W)^K \neq 0$ then $W^{K'_L} \neq 0$ for some K' in K^G with the Iwahori decomposition.
- (2) For any *G*-module V, and K' in K^G with the Iwahori decomposition, the map $V^{K'} \to V_U^{K'_L}$ is surjective.

We say that K is nice if it is nice with respect to any parabolic P containing S.

Proposition 5.1. Assume that K is nice and V is a G-module generated by V^K . If V' is a non-trivial subquotient of V then $(V')^K \neq 0$.

Proof. It suffices to prove the statement for irreducible subquotients. Without loss of generality we can assume that V is contained in a single Bernstein component corresponding to a pair (L, W) where L is a Levi group containing S and W a quasi-cuspidal representation of L. Let V' be an irreducible subquotient of V. Then there exists an unramified twist W' of W such that V' is a submodule of $\operatorname{Ind}_P^G W'$ where P is a parabolic subgroup containing L. Since $V^K \neq 0$ there exists at least one irreducible subquotient V_0 such that $(V_0)^K \neq 0$. Let W_0 be an unramified twist of W such that V_0 is contained in $\operatorname{Ind}_P^G W_0$. It follows that $(\operatorname{Ind}_P^G W_0)^K \neq 0$. By (1) above, there exists $K' \in K^G$ with the Iwahori decomposition such that $(W_0)^{K'_L} \neq 0$. Hence $(W')^{K'_L} \neq 0$ for any unramified twist W' of W. Now let V' be any irreducible quotient of V. From the Frobenius reciprocity

$$\operatorname{Hom}_{G}(V', \operatorname{Ind}_{P}^{G}(W')) = \operatorname{Hom}_{L}(V'_{U}, W')$$

it follows that W' is a quotient of V'_U . Hence $(V'_U)^{K'_L} \neq 0$ and $(V')^{K'} \neq 0$ by (2) above. Hence $(V')^K \neq 0$ since K' is conjugate to K.

Proposition 5.2. For every $x \in X$ and r > 0 the Moy-Prasad group $G_{x,r}$ is nice.

Proof. Without loss of generality we can assume that $x \in A(S)$. Let P be a parabolic subgroup containing S. Let σ be a chamber in A(S) containing x. Then the point-wise stabilizer G_{σ} of σ is an Iwahori subgroup of G. Let N be the normalizer of S in G. It acts naturally on A(S). We have an Iwasawa decomposition ([BT1], Proposition 7.3.1)

$$G = PNG_{\sigma}$$

Since $G_{\sigma} \subset G_x$, the Iwahori group G_{σ} normalizes $G_{x,r}$, hence representatives of *P*-orbits in the *G*-conjugacy class of $K = G_{x,r}$ can be taken to be $K' = G_{x',r}$ where $x' = n(x) \in A(S)$ for some $n \in N$. And these groups have the Iwahori decomposition with respect to *P*, since *P* contains *S* by Theorem 4.2 in [MP2].

Corollary 5.3. Assume that, for every vertex $x \in X$ there exists a contraction of $C_*(X)$ satisfying the properties (1) and (2). Then the Schneider-Stuhler complex is exact for every smooth representation V.

Proof. It suffices to prove that the complex is exact in every Bernstein component. The complex $C_*(X, V)$ is a direct sum of G-modules isomorphic to $\operatorname{ind}_{S_\tau}^G V_\tau$ where τ is a facet in the refined chamber complex and S_τ is the stabilizer of τ . This module is generated by $V^{K_\tau} = V_\tau$. Let x be a vertex of τ . Since $K_x \subseteq K_\tau$, it follows that $\operatorname{ind}_{G_\tau}^G V_\tau$ is generated K_x fixed vectors. Thus any Bernstein summand of $C_*(X, V)$ is generated by K_x -fixed vectors, for some vertex x and exactness can be checked by passing to K_x -fixed vectors. \Box

References

- [BKV] R. Bezrukavnikov, D. Kazhdan and Y. Varshavsky, On the depth r Bernstein projector. Selecta Math. (N.S.) 22 (2016), no. 4, 2271-2311.
- [BT1] F. Bruhat and J. Tits, Groupes réductif sur un corps local I. Publ. Math. I.H.E.S. 41 (1972).
- [BT2] F. Bruhat and J. Tits, Groupes réductif sur un corps local —I. Publ. Math. I.H.E.S. 60 (1984).
- [BD] J. N. Bernstein, rédigé par P. Deligne, *Le centre de Bernstein*. Travaux en Cours, Herman, Paris,1984.
- [MP1] A. Moy and G. Prasad, Unramified minimal K-types for p-adic groups. Invent. Math. **116** (1994), 393-408.
- [MP2] A. Moy and G. Prasad, Jacquet functors and unramified minimal K-types. Comment. Math. Helvetici 71 (1996), 98-121.
- [OS] E. Opdam and M. Solleveld Resolutions of tempered representations of reductive p-adic groups. J. Funct. Anal. 265 (2013), no. 1, 108134.
- [PR] G. Prasad and M. S. Ragunathan, Topological extensions of semi-simple groups over local fields. Annals of Math. 119 (1984), 143-201.
- [SS] P. Schneider and U. Stuhler, Representation theory and sheaves on the Bruhat-Tits building. Inst. Hautes tudes Sci. Publ. Math. 85 (1997), 97191.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF UTAH, SALT LAKE CITY, UT 84112 E-mail address: bestvina@math.utah.edu E-mail address: savin@math.utah.edu