

A fix-point characterization of Herbrand equivalence of expressions in data flow frameworks

Jasine Babu¹, K. Murali Krishnan², and Vineeth Paleri²

- 1 Indian Institute of Technology Palakkad, India
jasine@iitpkd.ac.in
- 2 National Institute of Technology Calicut, India
{kmurali,vpaleri}@nitc.ac.in

Abstract

The problem of determining Herbrand equivalence of terms at each program point in a data flow framework is a central problem in program analysis and compiler optimizations. Two expressions e and e' are said to be equivalent at a program point p in a data flow framework if e and e' are equivalent on every path in the flow graph from the entry point to the point p . Whereas the algorithmic computation of Herbrand equivalence in data flow frameworks proceeds via iterative fix-point computation on an abstract lattice framework, a fix-point characterization for Herbrand equivalence for data flow frameworks does not seem to appear so far in the literature. In this paper, we develop a set theoretic fix-point characterization of Herbrand equivalences at each program point in a data flow framework and prove the equivalence of this characterization with a meet-over-all-paths formulation.

1998 ACM Subject Classification F.3.2 – please refer to <http://www.acm.org/about/class/ccs98-html>

Keywords and phrases Herbrand Equivalence, Data Flow Framework, Fix Point, Program Analysis

1 Introduction

A data flow framework is an abstract representation of a program, used in program analysis and compiler optimizations [1]. One central problem in program analysis is that of determining the semantic equivalence of expressions at each point in a program. However this is unsolvable [6]. The alternative is to define a weaker, syntactic notion of expression equivalence over the set of all possible expressions and detect equivalence under the syntactic notion. Such a syntactic notion of equivalence is called *Herbrand equivalence*. Stated informally, Herbrand equivalence treats operators as uninterpreted functions, and two expressions are considered equivalent if they are obtained by applying the same operator on equivalent operands [4, 8, 10]. Two expressions e and e' are said to be equivalent at a program point p if e and e' are equivalent on every path in the flow graph from the entry point to the point p .

The pioneering work of Kildall[7], which essentially is an abstract interpretation [3] of terms, showed that at each program point, Herbrand equivalence of expressions that occur in a program could be computed using an iterative refinement algorithm. The algorithm models each iteration as the application of a monotone function over a meet semi-lattice, and terminates at a fix-point of the function[5, 6]. Subsequently, several problems in program analysis have been shown to be solvable using iterative fix-point computation on

lattice frameworks. (see [1] for examples). Several algorithms for computation of Herbrand equivalence of program expressions also were proposed in the literature [2, 4, 9, 10].

Although algorithmic computation to detect Herbrand equivalence among expressions that appear in a program proceeds via iterative fix-point computation on an abstract lattice framework, a precise set theoretic fix-point characterization of the Herbrand equivalences at program points in a data flow framework does not seem to appear so far in the literature. The main difficulty here is that, even as algorithms work with finite *working sets*, a mathematically precise characterization of Herbrand equivalence requires consideration of all program paths and equivalence of all expressions - including expressions not appearing in the program. Consequently, an iterative characterization of Herbrand equivalence cannot be achieved without resorting to set theoretic machinery.

In this paper, we develop a set theoretic fix-point characterization of Herbrand equivalences at each program point in a data flow framework. We define the notion of a congruence relation on the set of all expressions, and show that the set of all congruences form a complete lattice. Given a data flow framework with n program points, we show how to define a continuous function over the n fold product of the above lattice, such that the maximum fix-point of the function yields the set of Herbrand equivalence classes at various program points. This characterization is then shown to be equivalent to a meet over all paths formulation of expression equivalence over the same lattice framework.

Section 2 introduces the basic notation. Sections 3 to 7 develop the basic theory of congruences and transfer functions, including non-deterministic assignment functions. Section 8 and Section 9 deal with the application of the formalism of congruences to derive a fix-point characterization of Herbrand equivalence at each program point. Section 10 describes a meet-over-all-paths formulation for expression equivalence and establishes the equivalence between the fix-point characterization and the meet-over-all-paths formulation.

2 Terminology

Let C be a countable set of constants and X be a countable set of variables. For simplicity, we assume that the set of operators $Op = \{+\}$. (More operators can be added without any difficulty). The set of all terms over $C \cup X$, $\mathcal{T} = \mathcal{T}(X, C)$ is defined by $t ::= c \mid x \mid (t + t)$, with $c \in C$ and $x \in X$. (Parenthesis is avoided when there is no confusion.) Let \mathcal{P} be a partition of \mathcal{T} . Let $[t]_{\mathcal{P}}$ (or simply $[t]$ when there is no confusion) denote the equivalence class containing the term $t \in \mathcal{T}$. If $t' \in [t]_{\mathcal{P}}$, we write $t \cong_{\mathcal{P}} t'$ (or simply $t \cong t'$). Note that \cong is reflexive, symmetric and transitive. For any $A \subseteq \mathcal{T}$, $A(x)$ denotes the set of all terms in A in which the variable x appears and $\bar{A}(x)$ denotes the set of all terms in A in which x does not appear. In particular, for any $x \in X$, $\mathcal{T}(x)$ is the set of all terms containing the variable x and $\bar{\mathcal{T}}(x)$ denotes the set of terms in which x does not appear.

► **Definition 2.1** (Substitution). For $t, \alpha \in \mathcal{T}$, $x \in X$, substitution of x with α in t , denoted by $t[x \leftarrow \alpha]$ is defined as follows:

1. If $t = x$, then $t[x \leftarrow \alpha] = \alpha$.
2. If $t \notin \mathcal{T}(x)$, $t[x \leftarrow \alpha] = t$
3. If $t = t_1 + t_2$ then $t[x \leftarrow \alpha] = t_1[x \leftarrow \alpha] + t_2[x \leftarrow \alpha]$.

► **Remark.** In the rest of the paper, complete proofs for statements that are left unproven in the main text are provided in the appendix. Proofs for some elementary properties of lattices, which are used in the main text, are also given in the appendix, to make the presentation self contained.

3 Congruences of Terms

We define the notion of congruence (of terms). The notion of congruence will be useful later to model equivalence of terms at various program points in a data flow framework.

► **Definition 3.1** (Congruence of Terms). Let \mathcal{P} be a partition of \mathcal{T} . \mathcal{P} is a Congruence (of terms) if the following conditions hold:

1. For each $c, c' \in C$, if $c \neq c'$ then $c \not\cong c'$. (No two distinct constants are congruent).
2. For $t, t', s, s' \in \mathcal{T}$, $t' \cong t$ and $s' \cong s$ if and only if $t' + s' \cong t + s$. (Congruences respect operators).
3. For any $c \in C$, $t \in \mathcal{T}$, if $t \cong c$ then either $t = c$ or $t \in X$. (The only non-constant terms that are allowed to be congruent to a constant are variables).

The motivation for the definition of congruence is the following. Given the representation of a program in a data flow framework (to be defined later), we will associate a congruence to each program point at each iteration. Each iteration refines the present congruence at each program point. We will see later that this process of refinement leads to a well defined “fix point congruence” at each program point. We will see that this fix point congruence captures Herbrand equivalence at that program point.

► **Definition 3.2.** The set of all congruences over \mathcal{T} is denoted by $\mathcal{G}(\mathcal{T})$. We first note the substitution property of congruences.

► **Observation 3.3** (Substitution Property). Let \mathcal{P} be a congruence over \mathcal{T} . Then, for each $\alpha, \beta \in \mathcal{T}$, $\alpha \cong \beta$ if and only if for all $x \in X$ and $t \in \mathcal{T}$, $t[x \leftarrow \alpha] \cong t[x \leftarrow \beta]$.

Proof. One direction is easy. If for all $x \in X$ and $t \in \mathcal{T}$, $t[x \leftarrow \alpha] \cong t[x \leftarrow \beta]$, then setting $t = x$ we get $\alpha = t[x \leftarrow \alpha] \cong t[x \leftarrow \beta] = \beta$. Conversely, suppose $\alpha \cong \beta$. Let $x \in X$ and $t \in \mathcal{T}$ be chosen arbitrarily. To prove $t[x \leftarrow \alpha] \cong t[x \leftarrow \beta]$, we use induction. If $t \notin \mathcal{T}(x)$, then $t[x \leftarrow \alpha] = t \cong t = t[x \leftarrow \beta]$. If $t = x$, then $t[x \leftarrow \alpha] = \alpha \cong \beta = t[x \leftarrow \beta]$. Otherwise, if $t = t_1 + t_2$, then $t[x \leftarrow \alpha] = t_1[x \leftarrow \alpha] + t_2[x \leftarrow \alpha] \cong t_1[x \leftarrow \beta] + t_2[x \leftarrow \beta]$ (by induction hypothesis) $= t[x \leftarrow \beta]$. ◀

The following observation is a direct consequence of condition (3) of the definition of congruence.

► **Observation 3.4.** Let $\mathcal{P} \in \mathcal{G}(\mathcal{T})$, $c \in C$ and let $t \in \mathcal{T}(y)$, with $t \neq y$. Then $c \not\cong_{\mathcal{P}} t[y \leftarrow c]$.

We define a binary *confluence operation* on the set of congruences, $\mathcal{G}(\mathcal{T})$. A confluence operation transforms a pair of congruences into a congruence.

► **Definition 3.5** (Confluence). Let $\mathcal{P}_1 = \{A_i\}_{i \in I}$ and $\mathcal{P}_2 = \{B_j\}_{j \in J}$ be two congruences. For all $i \in I$ and $j \in J$, define $C_{i,j} = A_i \cap B_j$. The confluence of \mathcal{P}_1 and \mathcal{P}_2 is defined by: $\mathcal{P}_1 \wedge \mathcal{P}_2 = \{C_{i,j} : i \in I, j \in J, C_{i,j} \neq \emptyset\}$.

► **Theorem 3.6.** If \mathcal{P}_1 and \mathcal{P}_2 are congruences, then $\mathcal{P}_1 \wedge \mathcal{P}_2$ is a congruence.

4 Structure of Congruences

In this section we will define an ordering on the set $\mathcal{G}(\mathcal{T})$ and then extend it to a complete lattice.

► **Definition 4.1** (Refinement of a Congruence). Let $\mathcal{P}, \mathcal{P}'$ be congruences over \mathcal{T} . We say $\mathcal{P} \preceq \mathcal{P}'$ (read \mathcal{P} is a refinement of \mathcal{P}') if for each equivalence class $A \in \mathcal{P}$, there exists an equivalence class $A' \in \mathcal{P}'$ such that $A \subseteq A'$.

► **Definition 4.2.** The partition in which each term in \mathcal{T} belongs to a different class is defined as: $\perp = \{\{t\} : t \in \mathcal{T}\}$.

The following observation is a direct consequence of the definition of \perp .

► **Observation 4.3.** \perp is a congruence. Moreover for any $\mathcal{P} \in \mathcal{G}(\mathcal{T})$, $\perp \wedge \mathcal{P} = \perp$.

► **Definition 4.4.** A partially ordered set (P, \leq) is a meet semi-lattice if every pair of elements $a, b \in P$ has a greatest lower bound (called the meet of a and b).

► **Lemma 4.5.** $(\mathcal{G}(\mathcal{T}), \preceq)$ is a meet semi-lattice with meet operation \wedge and bottom element \perp .

The following lemma extends the meet operation to arbitrary non-empty collections of congruences. The proof relies on the axiom of choice.

► **Lemma 4.6.** Every non-empty subset of $(\mathcal{G}(\mathcal{T}), \preceq)$ has a greatest lower bound.

Next, we extend the meet semi-lattice $(\mathcal{G}(\mathcal{T}), \preceq)$ by artificially adding a top element \top , so that the greatest lower bound of the empty set is also well defined.

► **Definition 4.7.** The lattice $(\overline{\mathcal{G}(\mathcal{T})}, \preceq, \perp, \top)$ is defined over the set $\overline{\mathcal{G}(\mathcal{T})} = \mathcal{G}(\mathcal{T}) \cup \{\top\}$ with $\mathcal{P} \preceq \top$ for each $\mathcal{P} \in \mathcal{G}(\mathcal{T})$. In particular, \top is the greatest lower bound of \emptyset and $\top \wedge \mathcal{P} = \mathcal{P}$ for every $\mathcal{P} \in \overline{\mathcal{G}(\mathcal{T})}$.

Hereafter, we will be referring the element \top as a congruence. It follows from Lemma 4.6 and the above definition that every subset of $\overline{\mathcal{G}(\mathcal{T})}$ has a greatest lower bound. Since a meet semi-lattice in which every subset has a greatest lower bound is a complete lattice (Theorem B.1), we have:

► **Theorem 4.8.** $(\overline{\mathcal{G}(\mathcal{T})}, \preceq, \perp, \top)$ is a complete lattice.

► **Definition 4.9 (Infimum).** Let $S = \{\mathcal{P}_i\}_{i \in I}$ be an arbitrary collection of congruences in $\overline{\mathcal{G}(\mathcal{T})}$ (S may be empty or may contain \top). The infimum of the set S , denoted by $\bigwedge S$ or $\bigwedge_{i \in I} \mathcal{P}_i$, is defined as the greatest lower bound of the set $\{\mathcal{P}_i\}_{i \in I}$.

5 Transfer Functions

We now define a class of unary operators on $\mathcal{G}(\mathcal{T})$ called *transfer functions*. A transfer function specifies how the assignment of a term to a variable transforms the congruence before the assignment into a new one.

► **Definition 5.1 (Transfer Functions).** Let $y \in X$ and $\beta \in \overline{\mathcal{T}}(y)$. (Note that y does not appear in β). Let $\mathcal{P} = \{A_i\}_{i \in I}$ be an arbitrary congruence. The transfer function $f_{y=\beta}(\mathcal{P}) : \mathcal{G}(\mathcal{T}) \rightarrow \mathcal{G}(\mathcal{T})$ transforms \mathcal{P} to a congruence $f_{y=\beta}(\mathcal{P})$ given by the following:


- For each $i \in I$, let $B_i = \{t \in \mathcal{T} : t[y \leftarrow \beta] \in A_i\}$.
- $f_{y=\beta}(\mathcal{P}) = \{B_i : i \in I, B_i \neq \emptyset\}$.

► **Remark.** It follows from the above definition that $\overline{A_i}(y) = (A_i \setminus A_i(y)) \subseteq B_i$. That is, B_i will contain all terms in A_i in which y does not appear. See Figure 1 for an example.

In the following, we write $f(\mathcal{P})$ instead of $f_{y=\beta}(\mathcal{P})$ to avoid cumbersome notation. The following is a direct consequence of Definition 5.1.

► **Observation 5.2.** For any $t, t' \in \mathcal{T}$, $t \cong_{f(\mathcal{P})} t'$ if and only if $t[y \leftarrow \beta] \cong_{\mathcal{P}} t'[y \leftarrow \beta]$.

To make Definition 5.1 well founded, we need to establish the following:



■ **Figure 1** Application of Transfer Functions

► **Theorem 5.3.** *If \mathcal{P} is a congruence, then for any $y \in X$, $\beta \in \overline{\mathcal{T}}(y)$, $f_{y=\beta}(\mathcal{P})$ is a congruence.*

Next, we extend the definition of transfer functions to $(\overline{\mathcal{G}}(\mathcal{T}), \preceq, \perp, \top)$.

► **Definition 5.4.** Let $y \in X$ and $\beta \in \overline{\mathcal{T}}(y)$. Let $\mathcal{P} \in \overline{\mathcal{G}}(\mathcal{T})$. The extended transfer function $\overline{f}_{y=\beta}(\mathcal{P}) : \overline{\mathcal{G}}(\mathcal{T}) \rightarrow \overline{\mathcal{G}}(\mathcal{T})$ transforms \mathcal{P} to $\overline{f}_{y=\beta}(\mathcal{P}) \in \overline{\mathcal{G}}(\mathcal{T})$ given by the following:

- If $\mathcal{P} \in \mathcal{G}(\mathcal{T})$, $\overline{f}_{y=\beta}(\mathcal{P}) = f_{y=\beta}(\mathcal{P})$.
- $\overline{f}_{y=\beta}(\top) = \top$.

To simplify the notation, we often write $f_{y=\beta}(\mathcal{P})$ (or even simply $f(\mathcal{P})$) instead of $\overline{f}_{y=\beta}(\mathcal{P})$, and refer to extended transfer functions as simply transfer functions, when the underlying assignment operation is clear from the context.

6 Properties of Transfer Functions

In this section we show that transfer functions are continuous over the complete lattice $(\overline{\mathcal{G}}(\mathcal{T}), \preceq, \perp, \top)$.

Consider the (extended) transfer function $f = \overline{f}_{y=\beta}$, where $y \in X$, $\beta \in \overline{\mathcal{T}}(y)$. Let \mathcal{P}_1 and \mathcal{P}_2 be congruences in $\overline{\mathcal{G}}(\mathcal{T})$, not necessarily distinct.

► **Lemma 6.1** (Distributivity). $f(\mathcal{P}_1) \wedge f(\mathcal{P}_2) = f(\mathcal{P}_1 \wedge \mathcal{P}_2)$.

Since distributive functions are monotone, we have:

► **Corollary 6.2** (Monotonicity). *If $\mathcal{P}_1 \preceq \mathcal{P}_2$, then $f(\mathcal{P}_1) \preceq f(\mathcal{P}_2)$.*

We next show that distributivity extends to arbitrary collections of congruences.

► **Definition 6.3.** Let (L, \leq, \perp, \top) and $(L', \leq', \perp', \top')$ be complete lattices. A function $f : L \rightarrow L'$ is continuous if for each $\emptyset \neq S \subseteq L$, $f(\bigwedge S) = \bigwedge' f(S)$, where $f(S) = \{f(s) : s \in S\}$ and \bigwedge, \bigwedge' denote the infimum operations in the lattices L and L' respectively.

► **Remark.** The definition of continuity given above is more stringent than the standard definition in the literature, which requires continuity only for subsets that are chains. Moreover, the definition above exempts the continuity condition to hold for the empty set, because otherwise even constant maps will fail to be continuous.

The proof of the next theorem uses the axiom of choice. Let $f = f_{y=\beta}$, where $y \in X$, $\beta \in \overline{\mathcal{T}}(y)$. For arbitrary collections of congruences $S \subseteq \overline{\mathcal{G}(\mathcal{T})}$, The notation $f(S)$ denotes the set $\{f(s) : s \in S\}$.

► **Theorem 6.4** (Continuity). *For any $\emptyset \neq S \subseteq \overline{\mathcal{G}(\mathcal{T})}$, $f(\bigwedge S) = \bigwedge f(S)$.*

7 Non-deterministic assignment

Next we define a special kind of transfer functions corresponding to input statements in the program. This kind of transfer functions are called non-deterministic assignments.

► **Definition 7.1.** Let $y \in X$ and let $\mathcal{P} \in \mathcal{G}(\mathcal{T})$. The transfer function $f_{y=*}(\mathcal{P}) : \mathcal{G}(\mathcal{T}) \mapsto \mathcal{G}(\mathcal{T})$ transforms \mathcal{P} to a congruence $f(\mathcal{P}) = f_{y=*}(\mathcal{P})$, given by: for every $t, t' \in \mathcal{T}$, $t \cong_{f(\mathcal{P})} t'$ if and only if both the following conditions are satisfied:

1. $t \cong_{\mathcal{P}} t'$
2. For every $\beta \in \mathcal{T} \setminus \mathcal{T}(y)$, $t[y \leftarrow \beta] \cong_{\mathcal{P}} t'[y \leftarrow \beta]$.

Since for every pair of terms $t, t' \in \mathcal{T}$ the above definition precisely decides whether $t \cong_{f(\mathcal{P})} t'$ or not, $f_{y=*}(\mathcal{P})$ is the unique relation containing exactly those pairs $t, t' \in \mathcal{T}$ satisfying both the conditions in Definition 7.1. The definition asserts that two terms that were equivalent before a non-deterministic assignment, will remain equivalent after the non-deterministic assignment to y if and only if the equivalence between the two terms is preserved under all possible substitutions to y .

To make the above definition well founded, we need to prove that $f_{y=*}(\mathcal{P})$ is a congruence.

► **Theorem 7.2.** *If \mathcal{P} is a congruence, then for any $y \in X$, $f_{y=*}(\mathcal{P})$ is a congruence.*

We write $\bigwedge_{\beta \in \overline{\mathcal{T}}(y)} f_{y=\beta}(\mathcal{P})$ to denote the set $\bigwedge \{f_{y=\beta}(\mathcal{P}) : \beta \in \overline{\mathcal{T}}(y)\}$. The next theorem shows that each non-deterministic assignment may be expressed as a confluence of (an infinite collection of) transfer function operations.

► **Theorem 7.3.** *If \mathcal{P} is a congruence, then for any $y \in X$,*

$$f_{y=*}(\mathcal{P}) = \mathcal{P} \wedge \left(\bigwedge_{\beta \in \overline{\mathcal{T}}(y)} f_{y=\beta}(\mathcal{P}) \right).$$

Next, we extend the definition of non-deterministic assignment transfer functions to the complete lattice $(\overline{\mathcal{G}(\mathcal{T})}, \preceq, \perp, \top)$.

► **Definition 7.4.** Let $y \in X$ and $\mathcal{P} \in \overline{\mathcal{G}(\mathcal{T})}$. The extended transfer function $\overline{f}_{y=*}(\mathcal{P}) : \overline{\mathcal{G}(\mathcal{T})} \mapsto \overline{\mathcal{G}(\mathcal{T})}$ transforms \mathcal{P} to $\overline{f}_{y=*}(\mathcal{P}) \in \overline{\mathcal{G}(\mathcal{T})}$ given by the following:

- If $\mathcal{P} \in \mathcal{G}(\mathcal{T})$, $\overline{f}_{y=*}(\mathcal{P}) = f_{y=*}(\mathcal{P})$.
- $\overline{f}_{y=*}(\top) = \top$.

The following theorem involves use of the axiom of choice. We will write $f_{y=*}(\mathcal{P})$ instead of $\overline{f}_{y=*}(\mathcal{P})$ to simplify notation.

► **Theorem 7.5** (Continuity). *For any $\emptyset \neq S \subseteq \overline{\mathcal{G}(\mathcal{T})}$, $f_{y=*}(\bigwedge S) = \bigwedge f_{y=*}(S)$, where $f_{y=*}(S) = \{f_{y=*}(s) : s \in S\}$.*

In the following, we derive a characterization for non-deterministic assignment that does not depend on the axiom of choice. Condition (3) of the definition of congruence (Definition 3.1) is necessary to derive this characterization. We first note a lemma which states that if the equivalence between two terms t, t' is preserved under substitution of y with any two distinct constants chosen arbitrarily, then the equivalence between the two terms will be preserved under substitution of y with any other term β in which y does not appear.

► **Lemma 7.6.** *Let $\mathcal{P} \in \mathcal{G}(\mathcal{T})$. Let $t, t' \in \mathcal{T}$ and $c_1, c_2 \in C$ with $c_1 \neq c_2$. Then $t[y \leftarrow c_1] \cong_{\mathcal{P}} t'[y \leftarrow c_1]$ and $t[y \leftarrow c_2] \cong_{\mathcal{P}} t'[y \leftarrow c_2]$ if and only if $t[y \leftarrow \beta] \cong_{\mathcal{P}} t'[y \leftarrow \beta]$ for all $\beta \in \overline{\mathcal{T}}(y)$.*

The above observation leads to a characterization of non-deterministic assignment that does not involve the axiom of choice.

► **Theorem 7.7.** *Let $\mathcal{P} \in \mathcal{G}(\mathcal{T})$ and let $c_1, c_2 \in C$, $c_1 \neq c_2$. Then, for any $y \in X$,*
 $f_{y=*}(\mathcal{P}) = \mathcal{P} \wedge f_{y \leftarrow c_1}(\mathcal{P}) \wedge f_{y \leftarrow c_2}(\mathcal{P})$

It follows from the above theorem that non-deterministic assignments can be characterized in terms of just three congruences (instead of dealing with infinitely many as in Theorem 7.3).

8 Data Flow Analysis Frameworks

We next formalize the notion of a data flow framework and apply the formalism developed above to characterize Herbrand equivalence at each point in a program.

► **Definition 8.1.** A control flow graph $G(V, E)$ is a directed graph over the vertex set $V = \{1, 2, \dots, n\}$ for some $n \geq 1$ satisfying the following properties:

- $1 \in V$ is called the entry point and has no predecessors.
- Every vertex $i \in V$, $i \neq 1$ is reachable from 1 and has at least one predecessor and at most two predecessors.
- Vertices with two predecessors are called confluence points.
- Vertices with a single predecessor are called (transfer) function points.

► **Definition 8.2.** A data flow framework over \mathcal{T} is a pair $D = (G, F)$, where $G(V, E)$ is a control flow graph on the vertex set $V = \{1, 2, \dots, n\}$ and F is a collection of transfer functions over $\mathcal{G}(\mathcal{T})$ such that for each function point $i \in V$, there is an associated transfer function $h_i \in F$, and $F = \{h_i : i \in V \text{ is a transfer function point}\}$.

Data flow frameworks can be used to represent programs. An example is given in Figure 2.

► **Remark.** In the sections that follow, for any $h_i \in F$, we will simply write h_i to actually denote the extended transfer function \bar{h}_i (see Definition 5.4 and Definition 7.4) without further explanation.

9 Herbrand Equivalence

Let $D = (G, F)$ be a data flow framework over \mathcal{T} . In the following, we will define the Herbrand Congruence function $H_D : V(G) \mapsto \overline{\mathcal{G}(\mathcal{T})}$. For each vertex $i \in V(G)$, the congruence $H_D(i)$ will be called the *Herbrand Congruence* associated with the vertex i of the data flow framework D . The function H_D will be defined as the maximum fix-point of a continuous function $f_D : \overline{\mathcal{G}(\mathcal{T})}^n \mapsto \overline{\mathcal{G}(\mathcal{T})}^n$. The function f_D will be called the *composite transfer function* associated with the data flow framework D .

► **Definition 9.1 (Product Lattice).** Let n a positive integer. The product lattice, $(\overline{\mathcal{G}(\mathcal{T})}^n, \preceq_n, \perp_n, \top_n)$ is defined as follows: for $\overline{\mathcal{P}} = (\mathcal{P}_1, \mathcal{P}_2, \dots, \mathcal{P}_n)$, $\overline{\mathcal{Q}} = (\mathcal{Q}_1, \mathcal{Q}_2, \dots, \mathcal{Q}_n) \in \overline{\mathcal{G}(\mathcal{T})}^n$, $\overline{\mathcal{P}} \preceq_n \overline{\mathcal{Q}}$ if $\mathcal{P}_i \preceq \mathcal{Q}_i$ for each $1 \leq i \leq n$, $\perp_n = (\perp, \perp, \dots, \perp)$ and $\top_n = (\top, \top, \dots, \top)$.

For $S \subset \overline{\mathcal{G}(\mathcal{T})}^n$, the notation $\bigwedge_n S$ will be used to denote the least upper bound of S in the product lattice.

By Theorem 4.8, Theorem B.4 and Corollary B.5, we have:

► **Theorem 9.2.** *The product lattice satisfies the following properties:*

1. $(\overline{\mathcal{G}(\mathcal{T})}^n, \preceq_n, \perp_n, \top_n)$ is a complete lattice.
2. If $\tilde{S} \subseteq \overline{\mathcal{G}(\mathcal{T})}^n$ is non-empty, with $\tilde{S} = S_1 \times S_2 \times \dots \times S_n$, where $S_i \subseteq \overline{\mathcal{G}(\mathcal{T})}$ for $1 \leq i \leq n$. Then $\bigwedge_n \tilde{S} = (\bigwedge S_1, \bigwedge S_2, \dots, \bigwedge S_n)$.

As preparation for defining the composite transfer function, we introduce the following functions:

- **Definition 9.3** (Projection Maps). Let n be a positive integer. For each $i \in \{1, 2, \dots, n\}$,
 - The projection map to the i^{th} co-ordinate, $\pi_i : \overline{\mathcal{G}(\mathcal{T})}^n \mapsto \overline{\mathcal{G}(\mathcal{T})}$ is defined by $\pi_i(\mathcal{P}_1, \mathcal{P}_2, \dots, \mathcal{P}_n) = \mathcal{P}_i$ for any $(\mathcal{P}_1, \mathcal{P}_2, \dots, \mathcal{P}_n) \in \overline{\mathcal{G}(\mathcal{T})}^n$.
 - The confluence map $\pi_{i,j} : \overline{\mathcal{G}(\mathcal{T})}^n \mapsto \overline{\mathcal{G}(\mathcal{T})}$ is defined by $\pi_{i,j}(\mathcal{P}_1, \mathcal{P}_2, \dots, \mathcal{P}_n) = \mathcal{P}_i \wedge \mathcal{P}_j$ for any $(\mathcal{P}_1, \mathcal{P}_2, \dots, \mathcal{P}_n) \in \overline{\mathcal{G}(\mathcal{T})}^n$.

In addition to the above functions, we will also use the constant map which maps each element in $\overline{\mathcal{G}(\mathcal{T})}^n$ to \perp . The following observation is a consequence of Theorem B.10.

- **Observation 9.4.** Constant maps, projection maps and confluence maps are continuous.

For each $k \in V(G)$, $\text{Pred}(k)$ denotes the set of predecessors of the vertex k in the control flow graph G .

- **Definition 9.5** (Composite Transfer Function). Let $D = (G, F)$ be a data flow framework over \mathcal{T} . For each $k \in V(G)$, define the component map $f_k : \overline{\mathcal{G}(\mathcal{T})}^n \mapsto \overline{\mathcal{G}(\mathcal{T})}$ as follows:

1. If $k = 1$, the entry point, then $f_k = \perp$. (f_1 is the constant function that always returns the value \perp).
2. If k is a function point with $\text{Pred}(k) = \{j\}$, then $f_k = h_k \circ \pi_j$, where h_k is the (extended) transfer function corresponding to the function point k , and π_k the projection map to the k^{th} coordinate as defined in Definition 9.3.
3. If k is a confluence point with $\text{Pred}(k) = \{i, j\}$, then $f_k = \pi_{i,j}$, where $\pi_{i,j}$ is the confluence map as defined in Definition 9.3.

The composite transfer function of D is defined to be the unique function (Observation B.13) f_D satisfying $\pi_k \circ f_D = f_k$ for each $k \in V(G)$.

The purpose of defining f_D is the following. Suppose we have associated a congruence with each program point in a data flow framework. Then f_D specifies how a simultaneous and synchronous application of all the transfer functions/confluence operations at the respective program points modifies the congruences at each program point. The definition of f_D conservatively sets the confluence at the entry point to \perp , treating terms in $\mathcal{G}(\mathcal{T})$ to be inequivalent to each other at the entry point. See Figure 2 for an example. The following observation is a direct consequence of the above definition.

- **Observation 9.6.** The composite transfer function f_D (Definition 9.5) satisfies the following properties:

1. If $k = 1$, the entry point, then $\pi_k \circ f_D = \perp$.
2. If k is a function point with $\text{Pred}(k) = \{j\}$, then $f_k = \pi_k \circ f_D = h_k \circ \pi_j$, where h_k is the (extended) transfer function corresponding to the function point k .
3. If k is a confluence point with $\text{Pred}(k) = \{i, j\}$, then $f_k = \pi_k \circ f_D = \pi_{i,j}$.

The following lemma is a consequence of Observation 9.6.

- **Lemma 9.7.** Let $D = (G, F)$ be a data flow framework over \mathcal{T} and $k \in V(G)$. Let $S = \{f_D(\top_n), f_D^2(\top_n), \dots\}$, where f_D is the composite transfer function of D .

1. If $k = 1$, the entry point, then $\pi_k \circ f_D^l(\top_n) = \perp$ for all $l \geq 1$, hence $\pi_k(\bigwedge_n S) = \top$.
2. If k is a function point with $\text{Pred}(k) = \{j\}$, then for all $l \geq 1$,

$$\begin{aligned} (\pi_k \circ f_D^l)(\top_n) &= (\pi_k \circ f_D)(f_D^{l-1}(\top_n)) \\ &= (h_k \circ \pi_j \circ f_D^{l-1})(\top_n) \end{aligned}$$


3. If k is a confluence point with $\text{Pred}(k) = \{i, j\}$, then for all $l \geq 1$,

$$\begin{aligned} (\pi_k \circ f_D^l)(\top_n) &= (\pi_k \circ f_D)(f_D^{l-1}(\top_n)) \\ &= (\pi_{i,j})(f_D^{l-1}(\top_n)) \\ &= ((\pi_i \circ f_D^{l-1})(\top_n)) \wedge ((\pi_j \circ f_D^{l-1})(\top_n)) \end{aligned}$$

By Theorem B.14, Observation B.7 and Corollary B.17, we have:

► **Theorem 9.8.** *The following properties hold for the composite transfer function f_D (Definition 9.5):*

1. f_D is monotone, distributive and continuous.
2. The component maps $f_k = \pi_k \circ f_D$ are continuous for all $k \in \{1, 2, \dots, n\}$.
3. f_D has a maximum fix-point.
4. If $S = \{\top, f_D(\top_n), f_D^2(\top_n), \dots\}$, then $\bigwedge_n S$ is the maximum fix-point of f_D .



■ **Figure 2** Component Maps of the Composite Transfer Function

The objective of defining Herbrand Congruence as the maximum fix point of the composite transfer function is possible now.

► **Definition 9.9** (Herbrand Congruence). Given a data flow framework $D = (G, F)$ over \mathcal{T} , the Herbrand Congruence function $H_D : V(G) \mapsto \overline{\mathcal{G}(\mathcal{T})}$ is defined as the maximum fix point of the composite transfer function f_D . For each $k \in V(G)$, the value $H_D(k) \in \overline{\mathcal{G}(\mathcal{T})}$ is referred to as the Herbrand Congruence at program point k .

The following is a consequence of Theorem 9.8 and the definition of Herbrand Congruence.

► **Observation 9.10.** For each $k \in V(G)$, $H_D(k) = \bigwedge \{(\pi_k \circ f_D^l)(\top_n) : l \geq 0\}$.

Proof.

$$\begin{aligned} H_D(k) &= \pi_k \left(\bigwedge_n \{f_D^l(\top_n) : l \geq 0\} \right) \text{ (by Theorem 9.8)} \\ &= \bigwedge \{ \pi_k(f_D^l(\top_n)) : l \geq 0 \} \text{ (by continuity of } \pi_k \text{)} \\ &= \bigwedge \{ (\pi_k \circ f_D^l)(\top_n) : l \geq 0 \} \end{aligned}$$

◀

The definition of Herbrand congruence must be shown to be consistent with the traditional meet-over-all-paths description of Herbrand equivalence of terms in a data flow framework. The next section addresses this issue.

10 MOP characterization

In this section, we give a meet over all paths characterization for the Herbrand Congruence at each program point. In the following, assume that we are given a data flow framework $D = (G, F)$ over \mathcal{T} , with $V(G) = \{1, 2, \dots, n\}$.

► **Definition 10.1** (Path). For any non-negative integer l , a program path (or simply a path) of length l to a vertex $k \in V(G)$ is a sequence $\alpha = (v_0, v_2, \dots, v_l)$ satisfying $v_0 = 1$, $v_l = k$ and $(v_{i-1}, v_i) \in E(G)$ for each $i \in \{1, 2, \dots, l\}$. For each $i \in \{0, 1, \dots, l\}$, α_i denotes the initial segment of α up to the i^{th} vertex, given by (v_0, v_1, \dots, v_i) .

► **Remark.** Note that the vertices in a path need not be distinct under this definition.

The next definition associates a congruence in $\overline{\mathcal{G}(\mathcal{T})}$ with each path in D . The path function captures the effect of application of transfer functions along the path on the initial congruence \perp , in the order in which the transfer functions appear along the path.

► **Definition 10.2** (Path Congruence). Let $\alpha = (v_0, v_1, \dots, v_l)$ be a path of length l to vertex $k \in V(G)$ for some $l \geq 0$. We define:

1. When $i = 0$, $m_{\alpha_i} = \perp$.
2. If $i > 0$ and $v_i = j$, where $j \in V(G)$ is a function point, then $m_{\alpha_i} = h_j(m_{\alpha_{i-1}})$, where $h_j \in F$ is the extended transfer function associated with the function point j .
3. If $i > 0$ and v_i is a confluence point, then $m_{\alpha_i} = m_{\alpha_{i-1}}$.
4. $m_\alpha = m_{\alpha_l}$.

The congruence m_α is defined as the path congruence associated with the path α .

For $k \in V(G)$ and $l \geq 0$, let $\Phi_l(k)$ denote the set of all paths of length *less than* l from the entry point 1 to the vertex k . In particular, $\Phi_0(k) = \emptyset$, for all $k \in V(G)$. The following observation is a consequence of the definition of $\Phi_l(k)$.

► **Observation 10.3.** If $k \in V(G)$ and $l \geq 1$,

1. If k is the entry point, then $\Phi_l(k) = \{(1)\}$, the set containing only the path of length zero, starting and ending at vertex 1.
2. If k is a function point with $\text{Pred}(k) = \{j\}$, then $\{\alpha_{l-1} : \alpha \in \Phi_l(k)\} = \{\alpha' : \alpha' \in \Phi_{l-1}(j)\} = \Phi_{l-1}(j)$.
3. If k is a confluence point with $\text{Pred}(k) = \{i, j\}$, then $\{\alpha_{l-1} : \alpha \in \Phi_l(k)\} = \{\alpha' : \alpha' \in \Phi_{l-1}(i)\} \cup \{\alpha' : \alpha' \in \Phi_{l-1}(j)\} = \Phi_{l-1}(i) \cup \Phi_{l-1}(j)$.

For $l \geq 0$, we define the congruence $M_l(k)$ to be the meet of all path congruences associated with paths of length less than l from the entry point to vertex k in G . Stated formally,

$$M_l(k) = \bigwedge \{m_\alpha : \alpha \in \Phi_l(k)\}, \text{ for } l \geq 0$$

► **Observation 10.4.** If $l = 0$, $\Phi_l(k) = \emptyset$ and hence $M_0(k) = \top$, for all $k \in V(G)$. Further, $M_1(1) = \perp$ and $M_1(k) = \top$, for $k \neq 1$. In general, $M_l(k) = \top$ if there are no paths of length less than l from 1 to k in G .

We define Φ_k to be the set of all paths from vertex 1 to vertex k in G , i.e., $\Phi_k = \bigcup_{l \geq 1} \Phi_l(k)$ and $MOP(k) = \bigwedge \{m_\alpha : \alpha \in \Phi(k)\} = \bigwedge \{M_l(k) : l \geq 0\}$. (The second equality follows from Lemma B.2 and Observation 10.4.) The congruence $MOP(k)$ is the meet of all path congruences associated with paths in Φ_k .

Our objective is to prove $MOP(k) = H_D(k)$ for each $k \in \{1, 2, \dots, n\}$ so that H_D captures the meet over all paths information about equivalence of expressions in \mathcal{T} . We begin with the following observations.

► **Lemma 10.5.** For each $k \in V(G)$ and $l \geq 1$

1. If $k = 1$, the entry point, then $M_l(k) = \perp$.
2. If k is a function point with $\text{Pred}(k) = \{j\}$, then $M_l(k) = h_k(M_{l-1}(j))$, where h_k is the (extended) transfer function corresponding to the function point k .
3. If k is a confluence point with $\text{Pred}(k) = \{i, j\}$, then $M_l(k) = M_{l-1}(i) \wedge M_{l-1}(j)$.

Proof. If $k = 1$, $\Phi_l(k) = \{(1)\}$, by Observation 10.3. Hence $M_l(k) = M_1(1) = \perp$, by Observation 10.4.

If k is a function point with $\text{Pred}(k) = \{j\}$, then

$$\begin{aligned} M_l(k) &= \bigwedge \{m_\alpha : \alpha \in \Phi_l(k)\} \\ &= \bigwedge \{h_k(m_{\alpha_{l-1}}) : \alpha \in \Phi_l(k)\} \text{ (by Definition 10.2)} \\ &= \bigwedge \{h_k(m_{\alpha'}) : \alpha' \in \Phi_{l-1}(j)\} \text{ (by Observation 10.3)} \\ &= h_k \left(\bigwedge \{m_{\alpha'} : \alpha' \in \Phi_{l-1}(j)\} \right) \text{ (by continuity of } h_k) \\ &= h_k(M_{l-1}(j)) \text{ (by definition of } M_{l-1}(j)) \end{aligned}$$

If k is a confluence point with $\text{Pred}(k) = \{i, j\}$, then

$$\begin{aligned} M_l(k) &= \bigwedge \{m_\alpha : \alpha \in \Phi_l(k)\} \\ &= \bigwedge (\{m_{\alpha_{l-1}} : \alpha \in \Phi_l(k)\}) \text{ (by Definition 10.2)} \\ &= \bigwedge (\{m_{\alpha'} : \alpha' \in \Phi_{l-1}(i)\} \cup \{m_{\alpha'} : \alpha' \in \Phi_{l-1}(j)\}) \text{ (by Observation 10.3)} \\ &= \left(\bigwedge \{m_{\alpha'} : \alpha' \in \Phi_{l-1}(i)\} \right) \wedge \left(\bigwedge \{m_{\alpha'} : \alpha' \in \Phi_{l-1}(j)\} \right) \text{ (by Lemma B.2)} \\ &= M_{l-1}(i) \wedge M_{l-1}(j) \text{ (by definition of } M_{l-1}(i) \text{ and } M_{l-1}(j)) \end{aligned}$$

◀

► **Lemma 10.6.** For each $k \in V(G)$ and $l \geq 0$, $M_l(k) = (\pi_k \circ f_D^l)(\top_n)$.

Proof. Let $k \in V(G)$ be chosen arbitrarily. We prove the lemma by induction on l . When $l = 0$, $M_l(k) = \top$, by Observation 10.4, as required. Otherwise,

1. If $k = 1$, the entry point, then by Lemma 9.7 and Lemma 10.5,
 $M_l(k) = \perp = (\pi_k \circ f_D^l)(\top_n)$.
2. If k is a function point with $\text{Pred}(k) = \{j\}$ and h_k is the (extended) transfer function corresponding to the function point k , then

$$\begin{aligned} M_l(k) &= h_k(M_{l-1}(j)) \text{ (By Lemma 10.5)} \\ &= h_k((\pi_j \circ f_D^{l-1})(\top_n)) \text{ (By induction hypothesis)} \\ &= (h_k \circ \pi_j \circ f_D^{l-1})(\top_n) \\ &= (\pi_k \circ f_D^l)(\top_n) \text{ (By Lemma 9.7)} \end{aligned}$$

3. If k is a confluence point with $\text{Pred}(k) = \{i, j\}$, then

$$\begin{aligned} M_l(k) &= M_{l-1}(i) \wedge M_{l-1}(j) \text{ (By Lemma 10.5)} \\ &= \left((\pi_i \circ f_D^{l-1})(\top_n) \right) \wedge \left((\pi_j \circ f_D^{l-1})(\top_n) \right) \text{ (By induction hypothesis)} \\ &= (\pi_k \circ f_D^l)(\top_n) \text{ (By Lemma 9.7)} \end{aligned}$$

◀

Finally, we show that the iterative fix-point characterization of Herbrand equivalence and the meet over all paths characterization coincide.

► **Theorem 10.7.** Let $D = (G, F)$ be a data flow framework. Then, for each $k \in V(G)$, $MOP(k) = H_D(k)$.

Proof.

$$\begin{aligned} MOP(k) &= \bigwedge \{m_\alpha : \alpha \in \Phi(k)\} \\ &= \bigwedge \{M_l(k) : l \geq 0\} \text{ (by Lemma B.2 and Observation 10.4)} \\ &= \bigwedge \{(\pi_k \circ f_D^l)(\top_n) : l \geq 0\} \text{ (by Lemma 10.6)} \\ &= H_D(k) \text{ (by Observation 9.10)} \end{aligned}$$

◀

11 Conclusion

We have shown that Herbrand equivalence of terms in a data flow framework admits a fix point characterization. Though not the concern addressed here, we note that this characterization naturally leads to algorithms that work on the restriction of congruences to terms that actually appear in given data flow framework and iteratively compute the maximum fix-point (see, for example, the algorithm presented in Appendix C). This allows the detection Herbrand equivalence between the expressions that actually appear in the given program.

References

- 1 Alfred V. Aho, Monica S. Lam, Ravi Sethi, and Jeffrey D. Ullman. *Compilers: principles, techniques, and tools*, volume 2. Addison-wesley Reading, 2006.
- 2 B. Alpern, M. N. Wegman, and F. K. Zadeck. Detecting equality of variables in programs. In *Proceedings of the 15th ACM SIGPLAN-SIGACT Symposium on Principles of Programming Languages*, POPL '88, pages 1–11, New York, NY, USA, 1988. ACM. Available from: <http://doi.acm.org/10.1145/73560.73561>, doi:10.1145/73560.73561.
- 3 Patrick Cousot and Radhia Cousot. Abstract interpretation: A unified lattice model for static analysis of programs by construction or approximation of fixpoints. In *Proceedings of the 4th ACM SIGACT-SIGPLAN Symposium on Principles of Programming Languages*, POPL '77, pages 238–252, New York, NY, USA, 1977. ACM. Available from: <http://doi.acm.org/10.1145/512950.512973>, doi:10.1145/512950.512973.
- 4 Sumit Gulwani and George C. Necula. A polynomial-time algorithm for global value numbering. *Science of Computer Programming*, 64(1):97–114, 2007. Special issue on the 11th Static Analysis Symposium - SAS 2004.
- 5 John B. Kam and Jeffrey D. Ullman. Global data flow analysis and iterative algorithms. *J. ACM*, 23(1):158–171, Jan 1976. Available from: <http://doi.acm.org/10.1145/321921.321938>, doi:10.1145/321921.321938.
- 6 John B. Kam and Jeffrey D. Ullman. Monotone data flow analysis frameworks. *Acta Informatica*, 7(3):305–317, Sep 1977. Available from: <https://doi.org/10.1007/BF00290339>, doi:10.1007/BF00290339.
- 7 Gary A. Kildall. A unified approach to global program optimization. In *Proceedings of the 1st Annual ACM SIGACT-SIGPLAN Symposium on Principles of Programming Languages*, POPL '73, pages 194–206, New York, NY, USA, 1973. ACM. Available from: <http://doi.acm.org/10.1145/512927.512945>, doi:10.1145/512927.512945.
- 8 Markus Müller-Olm, Oliver Rüthing, and Helmut Seidl. *Checking Herbrand Equalities and Beyond*, pages 79–96. Springer Berlin Heidelberg, Berlin, Heidelberg, 2005. Available from: https://doi.org/10.1007/978-3-540-30579-8_6, doi:10.1007/978-3-540-30579-8_6.
- 9 B. K. Rosen, M. N. Wegman, and F. K. Zadeck. Global value numbers and redundant computations. In *Proceedings of the 15th ACM SIGPLAN-SIGACT Symposium on Principles of Programming Languages*, POPL '88, pages 12–27, New York, NY, USA, 1988. ACM. Available from: <http://doi.acm.org/10.1145/73560.73562>, doi:10.1145/73560.73562.
- 10 Oliver Rüthing, Jens Knoop, and Bernhard Steffen. *Detecting Equalities of Variables: Combining Efficiency with Precision*, pages 232–247. Springer Berlin Heidelberg, Berlin, Heidelberg, 1999. Available from: https://doi.org/10.1007/3-540-48294-6_15, doi:10.1007/3-540-48294-6_15.
- 11 Alfred Tarski. A lattice-theoretical fixpoint theorem and its applications. *Pacific J. Math.*, 5(2):285–309, 1955.

A Proofs of Theorems

Theorem 3.6. If \mathcal{P}_1 and \mathcal{P}_2 are congruences, then $\mathcal{P}_1 \wedge \mathcal{P}_2$ is a congruence.

Proof. Let $\mathcal{P}_1 = \{A_i\}_{i \in I}$ and $\mathcal{P}_2 = \{B_j\}_{j \in J}$. Let $C_{i,j} = A_i \cap B_j$ for all $i \in I$ and $j \in J$. Clearly $\mathcal{P}_1 \wedge \mathcal{P}_2 = \{C_{i,j} : i \in I, j \in J, C_{i,j} \neq \emptyset\}$ is a partition of \mathcal{T} . It suffices to prove that $\mathcal{P}_1 \wedge \mathcal{P}_2$ satisfies properties (1) to (3) of Definition 3.1.

- (1) For any $c, c' \in C$, $c \cong_{\mathcal{P}_1 \wedge \mathcal{P}_2} c' \iff c \cong_{\mathcal{P}_1} c' \text{ and } c \cong_{\mathcal{P}_2} c' \iff c = c'$.
- (2) For $t, t', s, s' \in \mathcal{T}$, $t + s \cong_{\mathcal{P}_1 \wedge \mathcal{P}_2} t' + s'$ if and only if $t + s \cong_{\mathcal{P}_1} t' + s'$ and $t + s \cong_{\mathcal{P}_2} t' + s'$
 $\iff t' \cong_{\mathcal{P}_1} t, t' \cong_{\mathcal{P}_2} t, s' \cong_{\mathcal{P}_1} s \text{ and } s' \cong_{\mathcal{P}_2} s \iff t' \cong_{\mathcal{P}_1 \wedge \mathcal{P}_2} t \text{ and } s' \cong_{\mathcal{P}_1 \wedge \mathcal{P}_2} s$.
- (3) For any $c \in C, t \in \mathcal{T}$, $c \cong_{\mathcal{P}_1 \wedge \mathcal{P}_2} t$ if and only if $c \cong_{\mathcal{P}_1} t$ and $c \cong_{\mathcal{P}_2} t$, only if either $t = c$ or $t \in X$ (by condition (3) of the definition of congruence). ◀

Lemma 4.5. $(\mathcal{G}(\mathcal{T}), \preceq)$ is a meet semi-lattice with meet operation \wedge and bottom element \perp .

Proof. It is evident from the definition of \preceq that $(\mathcal{G}(\mathcal{T}), \preceq)$ is a partial order. Let $\mathcal{P}_1 = \{A_i\}_{i \in I}$ and $\mathcal{P}_2 = \{B_j\}_{j \in J}$ be congruences in $\mathcal{G}(\mathcal{T})$. We next show that $\mathcal{P}_1 \wedge \mathcal{P}_2$ is the meet of \mathcal{P}_1 and \mathcal{P}_2 with respect to \preceq . By Definition 3.5, the relations $\mathcal{P}_1 \wedge \mathcal{P}_2 \preceq \mathcal{P}_1$ and $\mathcal{P}_1 \wedge \mathcal{P}_2 \preceq \mathcal{P}_2$ must hold true. Suppose that a congruence $\mathcal{P} = \{C_k\}_{k \in K}$ satisfies $\mathcal{P} \preceq \mathcal{P}_1$ and $\mathcal{P} \preceq \mathcal{P}_2$. Then by definition of \preceq , for each $k \in K$, there must exist $i \in I$ and $j \in J$ such that $C_k \subseteq A_i$ and $C_k \subseteq B_j$. Consequently, $\emptyset \neq C_k \subseteq (A_i \cap B_j)$. By Definition 3.5 we have $A_i \cap B_j \in \mathcal{P}_1 \wedge \mathcal{P}_2$ and hence $\mathcal{P} \preceq \mathcal{P}_1 \wedge \mathcal{P}_2$. Thus, $\mathcal{P}_1 \wedge \mathcal{P}_2$ is the meet of \mathcal{P}_1 and \mathcal{P}_2 . Finally, \perp is the bottom element of $\mathcal{G}(\mathcal{T})$ by Observation 4.3. ◀

Lemma 4.6. Every non-empty subset of $(\mathcal{G}(\mathcal{T}), \preceq)$ has a greatest lower bound.

Proof. We will show that any arbitrary non-empty family of congruences $\{\mathcal{P}_r\}_{r \in R}$ of \mathcal{T} has a greatest lower bound. For each $r \in R$, let $\mathcal{P}_r = \{A_{r,i_r}\}_{i_r \in I_r}$, where $I_r \neq \emptyset$. Let \mathcal{I} be the Cartesian product of the index sets of the congruences; i.e., $\mathcal{I} = \prod_{r \in R} I_r$. Note that, each element in \mathcal{I} is a sequence $(i_r)_{r \in R}$. For each $(i_r)_{r \in R} \in \mathcal{I}$, define $B_{(i_r)} = \bigcap_{r \in R} A_{r,i_r}$. (Note that the index element $i_r \in I_r$ and A_{r,i_r} is the set in partition $\{\mathcal{P}_r\}_{r \in R}$ having index i_r). Let $\mathcal{P} = \{B_{(i_r)} : (i_r) \in \mathcal{I}, B_{(i_r)} \neq \emptyset\}$.

We claim that \mathcal{P} is a congruence and it is the greatest lower bound of $\{\mathcal{P}_r\}_{r \in R}$. To show that \mathcal{P} is a partition, first assume $t \in \mathcal{T}$. Then, for each $r \in R$, there exists some $i_r \in I_r$ such that $t \in A_{r,i_r}$. Hence $t \in B_{(i_r)}$. Thus every element appears in at least one set $B_{(i_r)}$ in \mathcal{P} . Next, if $(i_r), (j_r) \in \mathcal{I}$ such that $(i_r) \neq (j_r)$, we show that $B_{(i_r)} \cap B_{(j_r)} = \emptyset$. To this end, assume that some $t \in \mathcal{T}$ satisfies $t \in B_{(i_r)} \cap B_{(j_r)}$. Since $(i_r) \neq (j_r)$, there exists some index r_0 in which the sequences (i_r) and (j_r) differ; i.e., $i_{r_0} \neq j_{r_0}$. As $t \in B_{(i_r)} \cap B_{(j_r)}$, it must be the case that $t \in A_{r_0,i_{r_0}} \cap A_{r_0,j_{r_0}}$, which contradicts our assumption that \mathcal{P}_{r_0} is a congruence.

Next, we prove that \mathcal{P} satisfies properties (1) to (3) of Definition 3.1.

- (1) Let $c, c' \in C$ be constants such that $c \neq c'$. Suppose $c, c' \in B_{(i_r)}$ for some $(i_r) \in \mathcal{I}$. Then, for each $r \in R$, $c, c' \in A_{r,i_r}$. However, this is impossible as $\{A_{r,i_r}\}_{i_r \in I_r}$ is a congruence for each r . This proves (1).
- (2) Let $t, t', s, s' \in \mathcal{T}$. We have $t, t' \in B_{(i_r)}$ and $s, s' \in B_{(j_r)}$ for some $(i_r), (j_r) \in \mathcal{I}$ if and only if $t, t' \in A_{r,i_r}$ and $s, s' \in A_{r,j_r}$ for each $r \in R$ if and only if there exists $(k_r) \in \mathcal{I}$ such that $t + s, t' + s' \in A_{r,k_r}$ for each $r \in R$ (because each \mathcal{P}_r is a congruence) if and only if $t + s, t' + s' \in B_{(k_r)}$ (by definition of $B_{(k_r)}$), proving (3).

- (3) For any $c \in C$ and $t \in \mathcal{T}$, suppose $c, t \in B_{(i_r)}$. Then, $c, t \in A_{r, i_r}$ for each $r \in R$. Hence either $t = c$ or $t \in X$ because \mathcal{P}_r is a congruence for each $r \in R$.

Next we show that $\mathcal{P} \preceq \mathcal{P}_r$ for each $r \in R$. Let $B_{(i_r)} \in \mathcal{P}$. By definition, $B_{(i_r)} = \bigcap_{r \in R} A_{r, i_r}$. Thus $B_{(i_r)} \subseteq A_{r, i_r}$ for each $r \in R$. As $A_{r, i_r} \in \mathcal{P}_r$ for each $r \in R$, we have $\mathcal{P} \preceq \mathcal{P}_r$ for each $r \in R$.

To prove that \mathcal{P} is the greatest lower bound of $\{\mathcal{P}_r\}_{r \in R}$, assume $\mathcal{Q} \preceq \mathcal{P}_r$ for each $r \in R$. Let $\mathcal{Q} = \{C_k\}_{k \in K}$. Then, for each $k \in K$ and each $r \in R$, there exists $i_r \in I_r$ such that $C_k \subseteq A_{r, i_r}$. Hence $C_k \subseteq B_{(i_r)} = \bigcap_{r \in R} A_{r, i_r}$. Since $B_{(i_r)} \in \mathcal{P}$, we have $\mathcal{Q} \preceq \mathcal{P}$.

The proof of the lemma is complete. Note that axiom of choice was used in assuming that the set \mathcal{I} is non-empty. ◀

Theorem 5.3. If \mathcal{P} is a congruence, then for any $y \in X$, $\beta \in \overline{\mathcal{T}}(y)$, $f_{y=\beta}(\mathcal{P})$ is a congruence.

Proof. Let $\mathcal{P} = \{A_i\}_{i \in I}$. Let $f(\mathcal{P}) = f_{y=\beta}(\mathcal{P})$ and $\{B_i\}_{i \in I}$ be defined as in Definition 5.1. Since \mathcal{P} is a congruence, for each $t \in \mathcal{T}$, there exists classes $A_i, A_j \in \mathcal{P}$ such that $t \in A_i$ and $t[y \leftarrow \beta] \in A_j$. (Note that $i = j$ is possible, but A_i and A_j are uniquely determined by the terms t and $t[y \leftarrow \beta]$). First we show that $f(\mathcal{P})$ has t in exactly one class.

- If $t \notin A_i(y)$ (i.e., y does not appear in t), then clearly $t \in B_i$ and no other class of $f(\mathcal{P})$ by definition. In particular, when $t = c$ for any constant $c \in C$, $c \in B_i$ if and only if $c \in A_i$, establishing condition (1) of Definition 3.1.
- If $t \in A_i(y)$, then by definition of B_j , $t \in B_j$ and no other class in $f(\mathcal{P})$ contains t .

We have shown that $f(\mathcal{P})$ is a partition that satisfies condition (1) of Definition 3.1. Next, we prove that $f(\mathcal{P})$ satisfies the remaining conditions of the definition of congruence.

Condition (2): we need to prove that for all $t, t', s, s' \in \mathcal{T}$, $t' \cong_{f(\mathcal{P})} t$ and $s' \cong_{f(\mathcal{P})} s$ if and only if $t' + s' \cong_{f(\mathcal{P})} t + s$. We have:

$$\begin{aligned}
& t' + s' \cong_{f(\mathcal{P})} t + s \\
\iff & (t' + s')[y \leftarrow \beta] \cong_{\mathcal{P}} (t + s)[y \leftarrow \beta] \text{ (by definition of } f_{y=\beta}(\mathcal{P})) \\
& = t'[y \leftarrow \beta] + s'[y \leftarrow \beta] \cong_{\mathcal{P}} t[y \leftarrow \beta] + s[y \leftarrow \beta] \text{ (by Definition 2.1)} \\
\iff & t'[y \leftarrow \beta] \cong_{\mathcal{P}} t[y \leftarrow \beta] \text{ and } s'[y \leftarrow \beta] \cong_{\mathcal{P}} s[y \leftarrow \beta] \text{ (by definition of } \cong_{\mathcal{P}}) \\
\iff & t' \cong_{f(\mathcal{P})} t \text{ and } s' \cong_{f(\mathcal{P})} s \text{ (by definition of } f_{y=\beta}(\mathcal{P}))
\end{aligned}$$

Condition (3): For any $c \in C$, let $t \in \mathcal{T}$ such that $t \neq c$. Suppose $c \cong_{f(\mathcal{P})} t$, then by Observation 5.2 $t[y \leftarrow \beta] \in X \cup \{c\}$ (by condition (3) of the definition of congruence), which is possible only if one among the following cases are true: (1) $t = x$ for some $x \in X$, $x \neq y$ such that $x \cong_{\mathcal{P}} c$ or (2) $t = y$ and $\beta = x$ for some $x \in X$, $x \cong_{\mathcal{P}} c$ or (3) $t = y$ and $\beta = c$. In any case $t \in X$. ◀

Lemma 6.1. $f(\mathcal{P}_1) \wedge f(\mathcal{P}_2) = f(\mathcal{P}_1 \wedge \mathcal{P}_2)$.

Proof. If $\mathcal{P}_1 = \top$, then $f(\mathcal{P}_1) \wedge f(\mathcal{P}_2) = \top \wedge f(\mathcal{P}_2) = f(\mathcal{P}_2) = f(\top \wedge \mathcal{P}_2)$ and the lemma holds true. Similarly, if $\mathcal{P}_2 = \top$, the lemma holds true. Otherwise, let $\mathcal{P}_1 = \{A_i\}_{i \in I}$ and $\mathcal{P}_2 = \{B_j\}_{j \in J}$. By Definition 5.1, we have:

$$\begin{aligned}
f(\mathcal{P}_1) &= \{A'_i : i \in I, A'_i \neq \emptyset\}, \text{ where } A'_i = \{t \in \mathcal{T} : t[y \leftarrow \beta] \in A_i\} \\
f(\mathcal{P}_2) &= \{B'_j : j \in J, B'_j \neq \emptyset\}, \text{ where } B'_j = \{t \in \mathcal{T} : t[y \leftarrow \beta] \in B_j\}
\end{aligned}$$

By the definition of meet, we have:

$$f(\mathcal{P}_1) \wedge f(\mathcal{P}_2) = \{A'_i \cap B'_j : i \in I, j \in J, A'_i \cap B'_j \neq \emptyset\} \quad (1)$$

$$\mathcal{P}_1 \wedge \mathcal{P}_2 = \{A_i \cap B_j : i \in I, j \in J, A_i \cap B_j \neq \emptyset\} \quad (2)$$

Moreover,

$$f(\mathcal{P}_1 \wedge \mathcal{P}_2) = \{D_{i,j} : i \in I, j \in J, D_{i,j} \neq \emptyset\} \quad (3)$$

where,


$$\begin{aligned} D_{i,j} &= \{t \in \mathcal{T} : t[y \leftarrow \beta] \in A_i \cap B_j\} \\ &= \{t \in \mathcal{T} : t[y \leftarrow \beta] \in A_i \text{ and } t[y \leftarrow \beta] \in B_j\} \\ &= \{t \in \mathcal{T} : t \in A'_i \text{ and } t \in B'_j\} = A'_i \cap B'_j \end{aligned}$$

Thus, Equation (3) becomes,

$$f(\mathcal{P}_1 \wedge \mathcal{P}_2) = \{A'_i \cap B'_j : i \in I, j \in J, A'_i \cap B'_j \neq \emptyset\} \quad (4)$$

The result follows by comparing equations (1) and (4). \blacktriangleleft

Theorem 6.4. For any $\emptyset \neq S \subseteq \overline{\mathcal{G}(\mathcal{T})}$, $f(\bigwedge S) = \bigwedge f(S)$.



■ **Figure 3** Commutativity diagram for Theorem 6.4

Proof. If $S = \{\top\}$, $f(\bigwedge S) = \top = \bigwedge f(S)$. Otherwise, we may assume without loss of generality that $\top \notin S$ (else consider $S \setminus \{\top\}$).

Let $S = \{\mathcal{P}_r\}_{r \in R}$ be a non-empty collection of congruences, not containing \top . For each $r \in R$, let $\mathcal{P}_r = \{A_{r,i_r}\}_{i_r \in I_r}$ for some index set $I_r \neq \emptyset$. By Definition 5.1 of transfer function, for each $r \in R$ we have:

$$f(\mathcal{P}_r) = \{A'_{r,i_r} : i_r \in I_r, A'_{r,i_r} \neq \emptyset\}, \text{ where } A'_{r,i_r} = \{t \in \mathcal{T} : t[y \leftarrow \beta] \in A_{r,i_r}\} \quad (5)$$

Hence $f(S) = \{f(\mathcal{P}_r) : r \in R\}$.

Let \mathcal{I} be the Cartesian product of the index sets of the congruences; i.e., $\mathcal{I} = \prod_{r \in R} I_r$. Each element in \mathcal{I} is a sequence $(i_r)_{r \in R}$. For each $(i_r)_{r \in R} \in \mathcal{I}$, define $B_{(i_r)} = \bigcap_{r \in R} A_{r,i_r}$ and $B'_{(i_r)} = \bigcap_{r \in R} A'_{r,i_r}$. As shown in the proof of Lemma 4.6, we have:

$$\bigwedge S = \{B_{(i_r)} : (i_r) \in \mathcal{I}, B_{(i_r)} \neq \emptyset\} \quad (6)$$

$$\bigwedge f(S) = \{B'_{(i_r)} : (i_r) \in \mathcal{I}, B'_{(i_r)} \neq \emptyset\} \quad (7)$$

From Equation 6 and Definition 5.1 of transfer function we write:

$$f(\bigwedge S) = \{D_{(i_r)} : (i_r) \in \mathcal{I}, D_{(i_r)} \neq \emptyset\} \text{ where, } D_{(i_r)} = \{t \in \mathcal{T} : t[y \leftarrow \beta] \in B_{(i_r)}\} \quad (8)$$

Comparing Equation 7 and Equation 8, we see that to complete the proof, it is enough to prove that $D_{(i_r)} = B'_{(i_r)}$ for each $(i_r) \in I$ (see Figure 3).

$$\begin{aligned} D_{(i_r)} &= \{t \in \mathcal{T} : t[y \leftarrow \beta] \in B_{(i_r)}\} \\ &= \{t \in \mathcal{T} : t[y \leftarrow \beta] \in \bigcap_{r \in R} A_{r, i_r}\} \text{ (by definition of } B_{(i_r)}) \\ &= \{t \in \mathcal{T} : t[y \leftarrow \beta] \in A_{r, i_r} \text{ for each } r \in R\} \\ &= \{t \in \mathcal{T} : t \in A'_{r, i_r} \text{ for each } r \in R\} \text{ (by Equation 5)} \\ &= \{t \in \mathcal{T} : t \in \bigcap_{r \in R} A'_{r, i_r}\} \\ &= \{t \in \mathcal{T} : t \in B'_{(i_r)}\} \text{ (by definition of } B'_{(i_r)}) \\ &= B'_{(i_r)} \end{aligned}$$

The proof is now complete. Note that the axiom of choice was used in the definition of the set \mathcal{I} . \blacktriangleleft

Theorem 7.2. If \mathcal{P} is a congruence, then for any $y \in X$, $f_{y=*}(\mathcal{P})$ is a congruence.

Proof. We write $f(\mathcal{P})$ instead of $f_{y=*}(\mathcal{P})$ to avoid cumbersome notation. First we show that $\cong_{f(\mathcal{P})}$ is an equivalence relation, thereby establishing that $f(\mathcal{P})$ is a partition of \mathcal{T} . Reflexivity, symmetry and transitivity of $\cong_{f(\mathcal{P})}$ is clear from Definition 7.1. Moreover, each equivalence class in $f_{y=*}(\mathcal{P})$ is a subset of some equivalence class in the congruence \mathcal{P} . Consequently, *Condition (1)* and *Condition (3)* in the definition of congruence (Definition 3.1) will be satisfied by $f(\mathcal{P})$.

It suffices to show that $f_{y=*}(\mathcal{P})$ satisfies Condition (2) of Definition 3.1.

Condition (2): we need to prove that for all $t, t', s, s' \in \mathcal{T}$, $t' \cong_{f(\mathcal{P})} t$ and $s' \cong_{f(\mathcal{P})} s$ if and only if $t' + s' \cong_{f(\mathcal{P})} t + s$. We have:

$$\begin{aligned} t' + s' &\cong_{f(\mathcal{P})} t + s \\ \iff &(t' + s') \cong_{\mathcal{P}} (t + s) \text{ and, for every } \beta \in \mathcal{T} \setminus \mathcal{T}(y), \\ &(t + s)[y \leftarrow \beta] \cong_{\mathcal{P}} (t' + s')[y \leftarrow \beta] \text{ (by definition of } f_{y=*}(\mathcal{P})) \\ \iff &(t' + s') \cong_{\mathcal{P}} (t + s) \text{ and, for every } \beta \in \mathcal{T} \setminus \mathcal{T}(y), \\ &(t[y \leftarrow \beta] + s[y \leftarrow \beta]) \cong_{\mathcal{P}} (t'[y \leftarrow \beta] + s'[y \leftarrow \beta]) \\ \iff &(t' \cong_{\mathcal{P}} t) \text{ and } (s' \cong_{\mathcal{P}} s) \text{ and, for every } \beta \in \mathcal{T} \setminus \mathcal{T}(y), \\ &(t[y \leftarrow \beta] \cong_{\mathcal{P}} t'[y \leftarrow \beta]) \text{ and } s[y \leftarrow \beta] \cong_{\mathcal{P}} s'[y \leftarrow \beta]) \text{ (by Definition 3.1)} \\ \iff &(t' \cong_{f(\mathcal{P})} t) \text{ and } (s' \cong_{f(\mathcal{P})} s) \text{ (by definition of } f_{y=*}(\mathcal{P})) \end{aligned}$$

The proof of the theorem is complete. \blacktriangleleft

Theorem 7.3. If \mathcal{P} is a congruence, then for any $y \in X$,

$$f_{y=*}(\mathcal{P}) = \mathcal{P} \wedge \left(\bigwedge_{\beta \in \overline{\mathcal{T}}(y)} f_{y=\beta}(\mathcal{P}) \right).$$

Proof. Let $Q = \mathcal{P} \wedge \left(\bigwedge_{\beta \in \overline{\mathcal{T}}(y)} f_{y=\beta}(\mathcal{P}) \right)$. It follows from condition (1) of Definition 7.1 and Theorem 7.2 that $f_{y=*}(\mathcal{P})$ is a congruence, and a refinement of \mathcal{P} . That is, $f_{y=*}(\mathcal{P}) \preceq \mathcal{P}$. From condition (2) of Definition 7.1, we have,

$$\begin{aligned} t \cong_{f_{y=*}(\mathcal{P})} t' &\implies \text{for every } \beta \in \mathcal{T} \setminus \mathcal{T}(y), t[y \leftarrow \beta] \cong_{\mathcal{P}} t'[y \leftarrow \beta] \\ &\iff \text{for every } \beta \in \mathcal{T} \setminus \mathcal{T}(y), t \cong_{f_{y=\beta}(\mathcal{P})} t' \text{ (by Observation 5.2)} \end{aligned}$$

Hence for each $\beta \in \mathcal{T} \setminus \mathcal{T}(y)$, $f_{y=*}(\mathcal{P}) \preceq f_{y=\beta}(\mathcal{P})$. It follows from the definition of \bigwedge that $f_{y=*}(\mathcal{P}) \preceq \bigwedge_{\beta \in \overline{\mathcal{T}}(y)} f_{y=\beta}(\mathcal{P})$. Combining this with the fact that $f_{y=*}(\mathcal{P}) \preceq \mathcal{P}$ and using the definition of \bigwedge , we get $f_{y=*}(\mathcal{P}) \preceq Q$.

Conversely let $t, t' \in \mathcal{T}$ such that $t \cong_Q t'$. Then by definition of Q , we have $t \cong_{\mathcal{P}} t'$ and, for all $\beta \in \mathcal{T} \setminus \mathcal{T}(y)$ $t \cong_{f_{y=\beta}(\mathcal{P})} t'$. By the definition of $f_{y=*}(\mathcal{P})$ (Definition 7.1), we have $t \cong_{f_{y=*}(\mathcal{P})} t'$. Consequently $Q \preceq f_{y=*}(\mathcal{P})$.

Hence, $f_{y=*}(\mathcal{P}) = Q$. Note that the proof involves use of the axiom of choice in assuming existence of \bigwedge . ◀

Theorem 7.5. For any $\emptyset \neq S \subseteq \overline{\mathcal{G}(\mathcal{T})}$, $f_{y=*}(\bigwedge S) = \bigwedge f_{y=*}(S)$, where $f_{y=*}(S) = \{f_{y=*}(s) : s \in S\}$.

Proof. To avoid cumbersome notation, we will write f for $f_{y=*}$. If $S = \{\top\}$, $f(\bigwedge S) = \top = \bigwedge f(S)$. Otherwise, we may assume without loss of generality that $\top \notin S$ (else consider $S \setminus \{\top\}$).

Let $S = \{\mathcal{P}_r\}_{r \in R}$ be a non-empty collection of congruences, not containing \top . We denote by $\bigwedge_{r \in R} \mathcal{P}_r$ the congruence $\bigwedge\{\mathcal{P}_r : r \in R\}$ and $\bigwedge_{r \in R} f(\mathcal{P}_r)$ for the congruence $\bigwedge\{f(\mathcal{P}_r) : r \in R\}$.

$$\begin{aligned} f(\bigwedge S) &= f\left(\bigwedge_{r \in R} \mathcal{P}_r\right) \\ &= \left(\bigwedge_{r \in R} \mathcal{P}_r\right) \wedge \left(\bigwedge_{\beta \in \overline{\mathcal{T}}(y)} f\left(\bigwedge_{r \in R} \mathcal{P}_r\right)\right) \text{ (by Theorem 7.3)} \\ &= \left(\bigwedge_{r \in R} \mathcal{P}_r\right) \wedge \left(\bigwedge_{\beta \in \overline{\mathcal{T}}(y)} \bigwedge_{r \in R} f_{y=\beta}(\mathcal{P}_r)\right) \text{ (by continuity of } f_{y=\beta}) \\ &= \left(\bigwedge_{r \in R} \mathcal{P}_r\right) \wedge \left(\bigwedge_{r \in R} \bigwedge_{\beta \in \overline{\mathcal{T}}(y)} f_{y=\beta}(\mathcal{P}_r)\right) \text{ (by properties of } \bigwedge) \\ &= \bigwedge_{r \in R} \left(\mathcal{P}_r \wedge \bigwedge_{\beta \in \overline{\mathcal{T}}(y)} f_{y=\beta}(\mathcal{P}_r)\right) \text{ (by properties of } \bigwedge) \\ &= \bigwedge_{r \in R} f(\mathcal{P}_r) \text{ (by Theorem 7.3)} \\ &= \bigwedge f(S) \end{aligned}$$

Note that the axiom of choice was implicitly used in the proof. ◀

Lemma 7.6. Let $\mathcal{P} \in \mathcal{G}(\mathcal{T})$. Let $t, t' \in \mathcal{T}$ and $c_1, c_2 \in C$ with $c_1 \neq c_2$. Then $t[y \leftarrow c_1] \cong_{\mathcal{P}} t'[y \leftarrow c_1]$ and $t[y \leftarrow c_2] \cong_{\mathcal{P}} t'[y \leftarrow c_2]$ if and only if for every $\beta \in \overline{\mathcal{T}}(y)$, $t[y \leftarrow \beta] \cong_{\mathcal{P}} t'[y \leftarrow \beta]$.

Proof. One direction is trivial. For the other, assume that $\mathcal{P} \in \mathcal{G}(\mathcal{T})$ and $c_1, c_2 \in C$, $c_1 \neq c_2$. If $t = t'$ or $t, t' \notin \mathcal{T}(y)$, the lemma holds true trivially. Hence we assume that $t \in \mathcal{T}(y)$ and $t' \neq t$. If $t' \notin \mathcal{T}(y)$, Then our assumption leads to $t[y \leftarrow c_1] \cong_{\mathcal{P}} t'[y \leftarrow c_1] = t' = t'[y \leftarrow c_2] \cong_{\mathcal{P}} t[y \leftarrow c_2]$. This is impossible by Lemma A.1. Hence we may assume that $t, t' \in \mathcal{T}(y)$.

If $t = y$, then by Observation 3.4, $t[y \leftarrow c_1] = c_1 \not\cong_{\mathcal{P}} t'[y \leftarrow c_1]$ for any $t' \in \mathcal{T}(y)$, $t' \neq y$. Hence we may assume that $t \neq y$. Similarly, we may assume that $t' \neq y$ as well. Hence the lemma holds true whenever at least one among t, t' is in $X \cup C$. We now proceed by induction.

Suppose $t = t_1 + t_2$ and $t' = t'_1 + t'_2$. We have:

$$\begin{aligned} t[y \leftarrow c_1] \cong_{\mathcal{P}} t'[y \leftarrow c_1] &\iff t_1[y \leftarrow c_1] + t_2[y \leftarrow c_1] \cong_{\mathcal{P}} t'_1[y \leftarrow c_1] + t'_2[y \leftarrow c_1] \\ &\iff t_1[y \leftarrow c_1] \cong_{\mathcal{P}} t'_1[y \leftarrow c_1] \text{ and } t_2[y \leftarrow c_1] \cong_{\mathcal{P}} t'_2[y \leftarrow c_1] \\ &\quad \text{(By condition (2) of Definition 3.1)} \end{aligned}$$

Similarly,

$$t[y \leftarrow c_2] \cong_{\mathcal{P}} t'[y \leftarrow c_2] \iff t_1[y \leftarrow c_2] \cong_{\mathcal{P}} t'_1[y \leftarrow c_2] \text{ and } t_2[y \leftarrow c_2] \cong_{\mathcal{P}} t'_2[y \leftarrow c_2]$$

We may assume as induction hypothesis that:

$$\begin{aligned} t_1[y \leftarrow c_1] \cong_{\mathcal{P}} t'_1[y \leftarrow c_1] \quad \text{and} \quad t_1[y \leftarrow c_2] \cong_{\mathcal{P}} t'_1[y \leftarrow c_2] \\ \iff t_1[y \leftarrow \beta] \cong_{\mathcal{P}} t'_1[y \leftarrow \beta], \text{ for every } \beta \in \overline{\mathcal{T}}(y) \end{aligned}$$

$$\begin{aligned} t_2[y \leftarrow c_1] \cong_{\mathcal{P}} t'_2[y \leftarrow c_1] \quad \text{and} \quad t_2[y \leftarrow c_2] \cong_{\mathcal{P}} t'_2[y \leftarrow c_2] \\ \iff t_2[y \leftarrow \beta] \cong_{\mathcal{P}} t'_2[y \leftarrow \beta], \text{ for every } \beta \in \overline{\mathcal{T}}(y) \end{aligned}$$

Since the left sides of the above equivalences hold by assumption, we have:

$$t_1[y \leftarrow \beta] \cong_{\mathcal{P}} t'_1[y \leftarrow \beta] \text{ and } t_2[y \leftarrow \beta] \cong_{\mathcal{P}} t'_2[y \leftarrow \beta], \text{ for every } \beta \in \overline{\mathcal{T}}(y)$$

Hence, by condition (3) of Definition 3.1,

$$t[y \leftarrow \beta] \cong_{\mathcal{P}} t'[y \leftarrow \beta], \text{ for every } \beta \in \overline{\mathcal{T}}(y)$$

◀

► **Lemma A.1.** *Let $t \in \mathcal{T}(y)$ and $c_1, c_2 \in C$, $c_1 \neq c_2$. Let \mathcal{P} be a congruence. Then, $t[y \leftarrow c_1] \not\cong_{\mathcal{P}} t[y \leftarrow c_2]$.*

Proof. If $t = y$, the lemma holds by condition (1) of the definition of congruence (Definition 3.1). Otherwise, we proceed by induction. Let $t = t_1 + t_2$. Since $t \in \mathcal{T}(y)$, either $t_1 \in \mathcal{T}(y)$ or $t_2 \in \mathcal{T}(y)$. Without loss of generality, we assume that $t_1 \in \mathcal{T}(y)$. We have by induction hypothesis:

$$t_1[y \leftarrow c_1] \not\cong_{\mathcal{P}} t_1[y \leftarrow c_2] \tag{9}$$

Hence we have:

$$\begin{aligned} t[y \leftarrow c_1] &= t_1[y \leftarrow c_1] + t_2[y \leftarrow c_1] \\ &\not\cong_{\mathcal{P}} t_1[y \leftarrow c_2] + t_2[y \leftarrow c_2] \text{ (by condition (2) of Definition 3.1)} \\ &= t[y \leftarrow c_2] \end{aligned}$$

◀

Theorem 7.7. Let $\mathcal{P} \in \mathcal{G}(\mathcal{T})$ and let $c_1, c_2 \in C$, $c_1 \neq c_2$. Then, for any $y \in X$,
 $f_{y=*}(\mathcal{P}) = \mathcal{P} \wedge f_{y \leftarrow c_1}(\mathcal{P}) \wedge f_{y \leftarrow c_2}(\mathcal{P})$

Proof. Let $\mathcal{Q} = \mathcal{P} \wedge f_{y \leftarrow c_1}(\mathcal{P}) \wedge f_{y \leftarrow c_2}(\mathcal{P})$ and let $\mathcal{Q}' = \mathcal{P} \wedge \left(\bigwedge_{\beta \in \overline{\mathcal{T}}(y)} f_{y=\beta}(\mathcal{P}) \right)$ We have:

$$\begin{aligned}
 t \cong_{\mathcal{Q}} t' &\iff t \cong_{\mathcal{P}} t' \text{ and } t \cong_{f_{y=c_1}(\mathcal{P})} t' \text{ and } t \cong_{f_{y=c_2}(\mathcal{P})} t' \text{ (by definition of confluence)} \\
 &\iff t \cong_{\mathcal{P}} t' \text{ and } t[y \leftarrow c_1] \cong_{\mathcal{P}} t'[y \leftarrow c_1] \text{ and } t[y \leftarrow c_2] \cong_{\mathcal{P}} t'[y \leftarrow c_2] \\
 &\quad \text{(by Observation 5.2)} \\
 &\iff t \cong_{\mathcal{P}} t' \text{ and } \forall \beta \in \overline{\mathcal{T}}(y) \ t[y \leftarrow \beta] \cong_{\mathcal{P}} t'[y \leftarrow \beta] \text{ (by Lemma 7.6)} \\
 &\iff t \cong_{\mathcal{P}} t' \text{ and } \forall \beta \in \overline{\mathcal{T}}(y) \ t \cong_{f_{y=\beta}(\mathcal{P})} t' \text{ (by Observation 5.2)} \\
 &\iff t \cong_{\mathcal{Q}'} t' \text{ (by definition of confluence)}
 \end{aligned}$$

◀

B Lattice Properties

Certain standard facts from lattice theory used in the paper are proved here for easy reference.

In the following, (L, \leq, \perp, \top) denotes a lattice with greatest element \top and least element \perp . (We will refer to the lattice simply as L when there is not scope for confusion). For $a, b \in L$, $a \wedge b$ (respectively $a \vee b$) denotes the meet (respectively join) of a and b . For $S \subseteq L$, $\bigwedge S$ (respectively $\bigvee S$) denotes the greatest lower bound (respectively least upper bound) of the set S whenever it exists. The lattice L is meet complete if $\bigwedge S$ exists for every subset S of L . In particular, $\bigwedge \emptyset = \top$ and $\bigwedge L = \perp$. A lattice L is a complete lattice if $\bigwedge S$ and $\bigvee S$ exists for every $S \subseteq L$.

► **Theorem B.1.** *If (L, \leq, \perp, \top) is meet complete, then it is a complete lattice.*

Proof. It suffices to prove that for each $S \subseteq L$, $\bigvee S$ exists. Let S be a subset of L . Let $T = \{t \in L : s \leq t \text{ for each } s \in S\}$. Since L is meet complete there exists $t_0 \in L$ such that $t_0 = \bigwedge T$. We claim that t_0 is the least upper bound of S . By definitions of the set T , for each $t \in T$ and $s \in S$, $s \leq t$. Hence, by the definition of meet, $s \leq t_0$ for each $s \in S$. Consequently t_0 is an upper bound to S . Next, if any $t \in L$ satisfies $s \leq t$ for each $s \in S$, then by the definition of T , $t \in T$. Hence $t_0 \leq t$ (by the definition of t_0). Thus t_0 is the least upper bound of S . ◀

The following property of the meet operation will be frequently used.

► **Lemma B.2.** *If (L, \leq, \perp, \top) is a complete lattice. Let $\{S_i\}_{i \in I}$ be subsets of L . Then, $\bigwedge (\bigcup_{i \in I} S_i) = \bigwedge \{\bigwedge S_i : i \in I\}$.*

Proof. For each $i \in I$, let $\alpha_i = \bigwedge S_i$ and $\alpha = \bigwedge \{\alpha_i : i \in I\}$. Let $S = \bigcup_{i \in I} S_i$. We have to prove that $\alpha = \bigwedge S$. Since $\alpha \leq \alpha_i$ for each $i \in I$, $\alpha \leq s$ for each $s \in S_i$ for each $i \in I$. Hence α is a lower bound to S . If $\beta \in L$ satisfies $\beta \leq s$ for each $s \in S$, then $\beta \leq s_i$ for each $s_i \in S_i$ for all $i \in I$. Hence $\beta \leq \alpha_i$ for each $i \in I$. Consequently, $\beta \leq \alpha$. ◀

We next define the Cartesian product of lattices.

► **Definition B.3 (Product Lattice).** Let (L, \leq, \perp, \top) be a lattice and n a positive integer. The product lattice, $(L^n, \leq_n, \perp_n, \top_n)$ is defined as follows: for $\bar{a} = (a_1, a_2, \dots, a_n)$, $\bar{b} = (b_1, b_2, \dots, b_n) \in L^n$, $\bar{a} \leq_n \bar{b}$ if $a_i \leq b_i$ for each $1 \leq i \leq n$, $\perp_n = (\perp, \perp, \dots, \perp)$ and $\top_n = (\top, \top, \dots, \top)$.

It is easy to see that L^n is a lattice with the meet of \bar{a} and \bar{b} given by $(a_1 \wedge b_1, a_2 \wedge b_2, \dots, a_n \wedge b_n)$. The meet of \bar{a} and \bar{b} will be denoted by $\bar{a} \wedge_n \bar{b}$. Similarly, the join of \bar{a} and \bar{b} is $\bar{a} \vee_n \bar{b} = (a_1 \vee b_1, a_2 \vee b_2, \dots, a_n \vee b_n)$. For $\tilde{S} \subseteq L^n$, the notation $\bigwedge_n \tilde{S}$ (respectively $\bigvee_n \tilde{S}$) denotes the greatest lower bound (respectively least upper bound) of \tilde{S} in L^n whenever it exists.

► **Theorem B.4.** *If (L, \leq, \perp, \top) is a complete lattice, then $(L^n, \leq_n, \perp_n, \top_n)$ is a complete lattice.*

Proof. In view of Theorem B.1, it suffices to prove that for each $\tilde{S} \subseteq L^n$, $\bigwedge_n \tilde{S}$ exists in L^n . If $\tilde{S} = \emptyset$, the proof is trivial. Let $\tilde{S} = S_1 \times S_2 \times \dots \times S_n$, where $S_i \subseteq L$ for $1 \leq i \leq n$. We will show that $\bigwedge_n \tilde{S} = (\bigwedge S_1, \bigwedge S_2, \dots, \bigwedge S_n)$. Let $\alpha_i = \bigwedge S_i$ for $1 \leq i \leq n$. Let $\bar{\alpha} = (\alpha_1, \alpha_2, \dots, \alpha_n)$. Since $\alpha_i \leq s_i$ for each $s_i \in S_i$, by the definition of \leq_n , we have $\bar{\alpha} \leq_n \bar{s}$ for each $\bar{s} \in \tilde{S}$. Suppose $\bar{\beta} = (\beta_1, \beta_2, \dots, \beta_n) \in L^n$ satisfies $\bar{\beta} \leq_n \bar{s}$ for each $\bar{s} \in \tilde{S}$. Then, by the definition of \leq_n , we have $\beta_i \leq s_i$ for each $s_i \in S_i$, $1 \leq i \leq n$. It follows from the definition of α_i that $\beta_i \leq \alpha_i$ for all $1 \leq i \leq n$. Consequently, $\bar{\beta} \leq_n \bar{\alpha}$. The proof is complete. ◀

The proof of Theorem B.4 yields the following corollary.

► **Corollary B.5.** *Let $\tilde{S} \subseteq L^n$ be non-empty with $\tilde{S} = S_1 \times S_2 \times \dots \times S_n$, where $S_i \subseteq L$ for $1 \leq i \leq n$. Then $\bigwedge_n \tilde{S} = (\bigwedge S_1, \bigwedge S_2, \dots, \bigwedge S_n)$.*

Next we define continuous maps between lattices and show that continuous maps are distributive and monotone. Let (L, \leq, \perp, \top) and $(L', \leq', \perp', \top')$ be complete lattices. For arbitrary subsets $S \subseteq L$ and $S' \subseteq L'$, we use the notation $\bigwedge S$ and $\bigwedge' S'$ to denote the greatest lower bounds of S and S' in the respective lattices L and L' .

► **Definition B.6.** Let (L, \leq, \perp, \top) and $(L', \leq', \perp', \top')$ be complete lattices. A function $f : L \rightarrow L'$ is continuous if for each $\emptyset \neq S \subseteq L$, $f(\bigwedge S) = \bigwedge' f(S)$, where $f(S) = \{f(s) : s \in S\}$.

► **Observation B.7.** Let (L, \leq, \perp, \top) and $(L', \leq', \perp', \top')$ be complete lattices. A function $f : L \rightarrow L'$ be continuous. Let $s, s' \in L$ be chosen arbitrarily. Then f satisfies:

1. Distributivity: $f(s \wedge s') = f(s) \wedge' f(s')$.
2. Monotonicity: if $s \leq s'$ then $f(s) \leq' f(s')$.

The first property is immediate from the definition of continuity. For the second, assume that $s \leq s'$. Then $s = s \wedge s'$ and we have $f(s) = f(s \wedge s') = f(s) \wedge' f(s') \leq' f(s')$.

We next show that two particular families of maps from L^n to L called projection maps and confluence maps are continuous.

► **Definition B.8 (Projection Maps).** Let (L, \leq, \perp, \top) be a lattice. For each $i \in \{1, 2, \dots, n\}$, the projection map to the i^{th} co-ordinate, $\pi_i : L^n \rightarrow L$ is defined by $\pi_i(s_1, s_2, \dots, s_n) = s_i$ for any $(s_1, s_2, \dots, s_n) \in L^n$.

► **Definition B.9 (Confluence Maps).** Let (L, \leq, \perp, \top) be a lattice. For each $i, j \in \{1, 2, \dots, n\}$, the confluence map $\pi_{i,j} : L^n \rightarrow L$ is defined by $\pi_{i,j}(s_1, s_2, \dots, s_n) = s_i \wedge s_j$ for any $(s_1, s_2, \dots, s_n) \in L^n$.

Note that $\pi_{i,j}(\bar{s}) = \pi_i(\bar{s}) \wedge \pi_j(\bar{s})$ and $\pi_{i,i}(\bar{s}) = \pi_i(\bar{s})$ for all $\bar{s} \in L^n$ and $i, j \in \{1, 2, \dots, n\}$. Thus projection maps are special instances of confluence maps.

► **Theorem B.10.** *Projection maps and confluence maps over a complete lattice (L, \leq, \perp, \top) are continuous.*

Proof. Let $i, j \in \{1, 2, \dots, n\}$. Let $\emptyset \neq \tilde{S} \subseteq L^n$. We need to prove that $\pi_{i,j}(\bigwedge_n \tilde{S}) = \bigwedge \pi_{i,j}(\tilde{S})$. Let $\tilde{S} = S_1 \times S_2 \times \dots \times S_n$, $S_i \neq \emptyset$. Let $\alpha = \bigwedge S_i$ and $\beta = \bigwedge S_j$. By Corollary B.5 and the definition of $\pi_{i,j}$ we have:

$$\pi_{i,j}(\bigwedge_n \tilde{S}) = \pi_{i,j}(\bigwedge S_1, \bigwedge S_2, \dots, \bigwedge S_n) = \alpha \wedge \beta \quad (10)$$

$$\bigwedge \pi_{i,j}(\tilde{S}) = \bigwedge \pi_{i,j}(S_1 \times S_2 \times \dots \times S_n) = \bigwedge \{s_i \wedge s_j : s_i \in S_i, s_j \in S_j\} \quad (11)$$

The equality between the left sides of the two equations above follows from Lemma B.11. ◀

► **Lemma B.11.** *Let (L, \leq, \perp, \top) be a complete lattice. Let S_1 and S_2 be non empty subsets of L with $\alpha = \bigwedge S_1$ and $\beta = \bigwedge S_2$. Let $S' = \{s_1 \wedge s_2 : s_1 \in S_1, s_2 \in S_2\}$. Then $\bigwedge S' = \alpha \wedge \beta$.*

Proof. Since α and β are lower bounds to S_1 and S_2 , we have $\alpha \wedge \beta \leq \alpha \leq s_1$ and $\alpha \wedge \beta \leq \beta \leq s_2$, for each $s_1 \in S_1$ and $s_2 \in S_2$. Consequently, from the definition of meet, we have $\alpha \wedge \beta \leq s_1 \wedge s_2$ for any $s_1 \in S_1, s_2 \in S_2$ and thus $\alpha \wedge \beta$ is a lower bound to the set S' . Now, if $\gamma \leq s_1 \wedge s_2$ for all $s_1 \in S_1$ and $s_2 \in S_2$, then clearly $\gamma \leq s_1$ for all $s_1 \in S_1$ and $\gamma \leq s_2$ for all $s_2 \in S_2$. Since $\alpha = \bigwedge S_1$ and $\beta = \bigwedge S_2$, we have $\gamma \leq \alpha$ and $\gamma \leq \beta$. Consequently, by the definition of meet, $\gamma \leq \alpha \wedge \beta$. This shows that $\alpha \wedge \beta$ is the greatest lower bound of the set S' . ◀

Next we show that continuous maps are closed under composition.

► **Theorem B.12 (Composition).** *Let (L, \leq, \perp, \top) , $(L', \leq', \perp', \top')$ and $(L'', \leq'', \perp'', \top'')$ be complete lattices. Let $f : L \mapsto L'$ and $g : L' \mapsto L''$ be continuous maps. Then the composition map, $g \circ f : L \mapsto L''$, defined by $(g \circ f)(s) = g(f(s))$ for each $s \in L$, is continuous.*

Proof. Let $\emptyset \neq S \subseteq L$. We need to prove that $(g \circ f)(\bigwedge S) = \bigwedge (g \circ f)(S)$. Let $f(S) = \{f(s) : s \in S\}$ and $(g \circ f)(S) = g(f(S)) = \{(g \circ f)(s) : s \in S\}$.

$$\begin{aligned} (g \circ f)(\bigwedge S) &= g(f(\bigwedge S)) \\ &= g(\bigwedge f(S)) \text{ (By the continuity of } f) \\ &= \bigwedge g(f(S)) \text{ (By the continuity of } g) \\ &= \bigwedge (g \circ f)(S). \end{aligned}$$

◀

The next theorem shows that f is continuous if and only if each of its component maps are continuous. We first note the following:

► **Observation B.13.** *Let (L, \leq, \perp, \top) be a complete lattice. For each $i \in \{1, 2, \dots, n\}$, let $f_i : L^n \mapsto L$ be arbitrary functions. Then, the function $f : L^n \mapsto L^n$ defined by $f(\bar{s}) = (f_1(\bar{s}), f_2(\bar{s}), \dots, f_n(\bar{s}))$ satisfies $\pi_i \circ f = f_i$ for each $1 \leq i \leq n$. Conversely, for any function $f : L^n \mapsto L^n$, the i^{th} component map $f_i : L^n \mapsto L$ defined by $f_i = \pi_i \circ f$ for each $i \in \{1, 2, \dots, n\}$ satisfies $f_i(\bar{s}) = \pi_i(f(\bar{s}))$ for all $\bar{s} \in L^n$.*

► **Theorem B.14.** *Let (L, \leq, \perp, \top) be a complete lattice. The map $f : L^n \mapsto L^n$ is continuous if and only if $f_i = \pi_i \circ f$ is continuous for each $1 \leq i \leq n$.*

Proof. Suppose f is continuous. Since π_i is continuous (by Theorem B.10) for each $1 \leq i \leq n$, by Theorem B.12, $\pi_i \circ f$ is continuous. This establishes one direction of the theorem.

Conversely, suppose f_i is continuous for each $i \in \{1, 2, \dots, n\}$. Let $\emptyset \neq \tilde{S} \subseteq L^n$ be chosen arbitrarily. We need to prove that $f(\bigwedge_n \tilde{S}) = \bigwedge_n f(\tilde{S})$. Let $\tilde{S} = S_1 \times S_2 \times \dots \times S_n$, where

$S_i \subseteq L$ for each $1 \leq i \leq n$. Let $\alpha_i = \bigwedge S_i$ for each $1 \leq i \leq n$. Let $\bar{\alpha} = (\alpha_1, \alpha_2, \dots, \alpha_n)$. By Corollary B.5 we have:

$$\bigwedge_n \tilde{S} = (\bigwedge S_1, \bigwedge S_2, \dots, \bigwedge S_n) = (\alpha_1, \alpha_2, \dots, \alpha_n) = \bar{\alpha} \quad (12)$$

Hence,

$$f(\bigwedge_n \tilde{S}) = f(\bar{\alpha}) \quad (13)$$

Since for each $\bar{s} \in \tilde{S}$, $f(\bar{s}) = (f_1(\bar{s}), f_2(\bar{s}), \dots, f_n(\bar{s}))$, the above equation becomes:

$$f(\bigwedge_n \tilde{S}) = (f_1(\bar{\alpha}), f_2(\bar{\alpha}), \dots, f_n(\bar{\alpha})) \quad (14)$$

Now,

$$\begin{aligned} \bigwedge_n f(\tilde{S}) &= \bigwedge_n \{(f_1(\bar{s}), f_2(\bar{s}), \dots, f_n(\bar{s})) : \bar{s} \in \tilde{S}\} \\ &= \bigwedge_n f_1(\tilde{S}) \times f_2(\tilde{S}) \cdots \times f_n(\tilde{S}) \text{ (by definition of Cartesian product)} \\ &= (\bigwedge_n f_1(\tilde{S}), \bigwedge_n f_2(\tilde{S}), \dots, \bigwedge_n f_n(\tilde{S})) \text{ (by Corollary B.5)} \\ &= (f_1(\bigwedge_n \tilde{S}), f_2(\bigwedge_n \tilde{S}), \dots, f_n(\bigwedge_n \tilde{S})) \text{ (by continuity of } f_1, f_2, \dots, f_n) \\ &= (f_1(\bar{\alpha}), f_2(\bar{\alpha}), \dots, f_n(\bar{\alpha})) \text{ (by Equation 12)} \end{aligned}$$

The theorem follows by comparing the above equation with Equation 14. \blacktriangleleft

Next we turn to the computation of the maximum fix-point of a continuous map on a complete lattice. The well known Knaster-Tarski Theorem [11] asserts the existence of a maximum fix-point for every monotone function defined over a complete lattice. When the function is continuous, the maximum fix-point can be defined as the greatest lower bound of a specific iteratively defined subset of the lattice, as described below. In the following we use the notation $f^2 = f \circ f$, $f^3 = f \circ f^2$, etc.

► **Definition B.15 (Maximum Fix-Point).** Let (L, \leq, \perp, \top) be a complete lattice. Let $f : L \mapsto L$ be any function. $s \in L$ is a maximum fix point of f if:

- s is a fix point of f . That is, $f(s) = s$.
- for any $s' \in L$, if $f(s') = s'$, then $s' \leq s$.

► **Theorem B.16.** Let (L, \leq, \perp, \top) be a complete lattice. Let $f : L \mapsto L$ be continuous. Let $S = \{\top, f(\top), f^2(\top), f^3(\top), \dots\}$. Then $\bigwedge S$ is the maximum fix point of f .

Proof. Let $s_0 = \bigwedge S$. We need to prove that s_0 is the maximum fix point of f . If $f(\top) = \top$, then $S = \{\top\}$ and the theorem holds. Otherwise, as $f(\top) \leq \top$, it follows from the monotonicity of f (Observation B.7) that, S is a descending chain with $f^{i+1}(\top) \leq f^i(\top)$ for each $i \geq 1$; and we have $f(S) = S \setminus \{\top\} \neq \emptyset$. By continuity of f , we get:

$$f(s_0) = f(\bigwedge S) = \bigwedge f(S) = \bigwedge (S \setminus \{\top\}) = \bigwedge S = s_0$$

This shows s_0 is a fix-point. Suppose $s' \in L$ satisfies $f(s') = s'$. Since $s' \leq \top$, $s' = f(s') \leq f(\top)$. Extending the argument, we see that $s' \leq f^i(\top)$ for each $i \geq 1$. Hence by the definition of \bigwedge , $s' \leq \bigwedge S = s_0$. This proves that s_0 is the maximum fix-point of f . \blacktriangleleft

The following is a consequence of the above proof.

► **Corollary B.17.** *Let (L, \leq, \perp, \top) be a complete lattice. Let $f : L \mapsto L$ be continuous. Then the set $S' = \{\top, f(\top), f^2(\top), f^3(\top), \dots\}$ is a decreasing sequence of lattice values with $f^{i+1}(\top) \leq f^i(\top)$ for each $i \geq 0$. Moreover, $f(\bigwedge S')$ is the maximum fix point of f .*

C An Algorithm for Computing Program Expression Equivalence

A program expression in a program P is a term over $(X \cup C)$ that actually appear in the program P . Two program expressions e and e' are Herbrand equivalent at a program point if they belong to the same partition class of the Herbrand Congruence at that program point. The *Herbrand equivalence classes of program expressions* at a program point are the partition classes obtained by restricting the Herbrand Congruence of that point to only the set of program expressions. In this section we describe an algorithm that calculates the Herbrand equivalence classes of program expressions at each program point in any input program, represented as a data flow framework. We restrict our attention to only intra procedural analysis.

C.1 Description of Data Structures

An *ID* (which stands for a value identifier) is a composite data type with four fields, namely *fctype*, *valueNum*, *idOperand1* and *idOperand2*. If *fctype* is 0, we call it an atomic ID and it will have an associated value number called *valueNum*, which is a positive integer and the other two fields *idOperand1*, *idOperand2* are set to *NIL*. For a non-atomic ID, its *fctype* will be an operator and its *valueNum* field will be set to -1 . The field *idOperand1* will point to the *ID* of the first operand and *idOperand2* will point to the *ID* of the second operand. An *IdArray* type represents an array of *ID*s that holds one index for each element t in $(X \cup C) \cup [(X \cup C) \times (X \cup C) \times Op]$ in an order arbitrarily fixed in the beginning. Each array element indicates the value identifier of the corresponding element t .

Associated with each program point p , there is an *IdArray* and this together forms an array *Partitions* which has one index corresponding to each program point. This way, for each program point i , *Partitions*[i] will be an array of value identifiers, one index corresponding to each program expression and two program expressions t and t' are considered equivalent at program point i if and only if *Partitions*[i][t] = *Partitions*[i][t']. The notation *numClasses* represents the cardinality of the set $(X \cup C) \cup ((X \cup C) \times (X \cup C) \times Op)$ and *numProgPoints* represents the number of program points in the input program.

Algorithm 1 Data Structures

```

1: typedef struct {
    int fctype;                ▷ 0 if atomic and the operator number otherwise.
    int valueNum;             ▷ value number of the element if fctype is 0 and -1 otherwise.
    struct ID* idOperand1;    ▷ NIL if fctype is 0
    struct ID* idOperand2;    ▷ NIL if fctype is 0
2: } ID;
3: typedef IdArray ID[numClasses];
4: IdArray Partitions[numProgPoints];

```

C.2 Description of the Algorithm

Algorithm 2 Main Program

```

1: procedure MAIN
2:   StartCounter();
3:   for each term  $t \in X \cup C$  do
4:      $Partitions[1][t] \leftarrow CreateAtomicId()$ ;
5:   end for
6:   for each term  $t$  of the form  $t = t_1 + t_2$ , where  $t_1, t_2 \in X \cup C$  and  $+$  in Op do
7:      $Partitions[1][t] \leftarrow AssignCompoundId(\&Partitions[1][t_1], \&Partitions[1][t_2], +)$ ;
8:   end for
9:   for each program point  $k$  from 2 to numProgPoints do
10:     $Partitions[k] = \top$ ;
11:  end for
12:  ConvergeFlag  $\leftarrow 0$ ;
13:  while ConvergeFlag is 0 do
14:    ConvergeFlag  $\leftarrow 1$ ;
15:    for each program point  $k$  from 2 to numProgPoints do
16:       $OldPartition \leftarrow Partitions[k]$ ;
17:      if  $k$  is a function point with  $Pred(k) = \{j\}$  and assignment  $y \leftarrow \beta$  then
18:         $Partitions[k] \leftarrow AssignStmt(Partitions[j], y, \beta)$ ;
19:      else if  $k$  is a function point with  $Pred(k) = \{j\}$  and assignment  $y \leftarrow *$  then
20:         $Partitions[k] \leftarrow NonDetAssign(Partitions[j], y)$ ;
21:      else if  $k$  is a confluence point with  $Pred(k) = \{i, j\}$  then
22:         $Partitions[k] \leftarrow Confluence(Partitions[i], Partitions[j])$ ;
23:      end if
24:      if  $IsSame(OldPartition, Partitions[k])$  is 0 then ConvergeFlag  $\leftarrow 0$ ;
25:      end if
26:    end for
27:  end while
28: end procedure

```

Algorithm 3 Handling Assignment Statement $y \leftarrow \beta$

```

1: procedure ASSIGNSTMT(IdArray  $\mathcal{P}$ , variable  $y$ , term  $\beta$ )  $\triangleright$  Assume that  $y$  does not
   appear in  $\beta$ .
2:   IdArray  $\mathcal{Q}$ ;
3:   Initialize  $\mathcal{Q} = \mathcal{P}$ 
4:    $\mathcal{Q}[y] \leftarrow \mathcal{P}[\beta]$ 
5:   for each term  $t$  of the form  $t = y + t'$ , where  $t' \in X \cup C$  do
6:      $\mathcal{Q}[t] \leftarrow \text{AssignCompoundId}(\&\mathcal{Q}[y], \&\mathcal{Q}[t'], +)$ ;
7:   end for
8:   for each term  $t$  of the form  $t = t' + y$ , where  $t' \in X \cup C$  do
9:      $\mathcal{Q}[t] \leftarrow \text{AssignCompoundId}(\&\mathcal{Q}[t'], \&\mathcal{Q}[y], +)$ ;
10:  end for
11:  return  $\mathcal{Q}$ ;
12: end procedure

```

Algorithm 4 Handling Non-Deterministic Assignment Statement $y \leftarrow *$

```

1: procedure NONDETASSIGN(IdArray  $\mathcal{P}$ , variable  $y$ )  $\triangleright$ 
2:   IdArray  $\mathcal{Q}_1, \mathcal{Q}_2, \mathcal{Q}$ ;
3:   Initialize  $\mathcal{Q}_1 = \mathcal{P}; \mathcal{Q}_2 = \mathcal{P}$ ;
4:   Let  $c_1, c_2 \in C$ .
5:    $\mathcal{Q}_1[y] = \mathcal{P}[c_1]; \mathcal{Q}_2[y] = \mathcal{P}[c_2]$ ;
6:   for each term  $t$  of the form  $t = y + t'$ , where  $t' \in X \cup C$  do
7:      $\mathcal{Q}_1[t] = \text{AssignCompoundId}(\&\mathcal{Q}_1[y], \&\mathcal{Q}_1[t'], +)$ ;
8:      $\mathcal{Q}_2[t] = \text{AssignCompoundId}(\&\mathcal{Q}_2[y], \&\mathcal{Q}_2[t'], +)$ ;
9:   end for
10:  for each term  $t$  of the form  $t = t' + y$ , where  $t' \in X \cup C$  do
11:     $\mathcal{Q}_1[t] = \text{AssignCompoundId}(\&\mathcal{Q}_1[t'], \&\mathcal{Q}_1[y], +)$ ;
12:     $\mathcal{Q}_2[t] = \text{AssignCompoundId}(\&\mathcal{Q}_2[t'], \&\mathcal{Q}_2[y], +)$ ;
13:  end for
14:   $\mathcal{Q} = \text{Confluence}(\mathcal{Q}_1, \mathcal{Q}_2)$ ;
15:  return  $\text{Confluence}(\mathcal{Q}, \mathcal{P})$ ;
16: end procedure

```

Algorithm 5 Handling Confluence

```

1: procedure CONFLUENCE(IdArray  $\mathcal{P}_1$ , IdArray  $\mathcal{P}_2$ ) ▷
2:   IdArray  $\mathcal{Q}$ ; int AccessFlag[numClasses];
3:   for each term  $t \in X \cup C$  do
4:     AccessFlag[ $t$ ] = 0;
5:   end for
6:   for each term  $t \in X \cup C$  do
7:     if AccessFlag[ $t$ ] is equal to 0 then
8:       AccessFlag[ $t$ ] = 1;
9:       if  $\mathcal{P}_1[t]$  is equal to  $\mathcal{P}_2[t]$  then
10:         $\mathcal{Q}[t] = \mathcal{P}_1[t]$ ;
11:      else
12:         $S_1 = \text{getClass}(t, \mathcal{P}_1)$ ;
13:         $S_2 = \text{getClass}(t, \mathcal{P}_2)$ ;
14:         $\text{newId} = \text{CreateAtomicId}()$ ;
15:        for each term  $t'$  in  $S_1 \cap S_2 \cap (X \cup C)$  do
16:          AccessFlag[ $t'$ ] = 1;
17:           $\mathcal{Q}[t'] = \text{newId}$ ;
18:        end for
19:      end if
20:    end if
21:  end for
22:  for each term  $t$  of the form  $t = t_1 + t_2$ , where  $t_1, t_2 \in X \cup C$  do
23:     $\mathcal{Q}[t] = \text{AssignCompoundId}(\&\mathcal{Q}[t_1], \&\mathcal{Q}[t_2], +)$ ;
24:  end for
25:  return  $\mathcal{Q}$ ;
26: end procedure

```

Algorithm 6 Compare Congruences

```

1: procedure ISSAME(IdArray  $\mathcal{P}_1$ , IdArray  $\mathcal{P}_2$ ) ▷ See if two congruences are the same,
   considering class value-id to be abstract
2:   Set  $S_1, S_2$ ;
3:   for each term  $t$  in  $(X \cup C) \cup ((X \cup C) \times (X \cup C) \times Op)$  do
4:      $S_1 = \text{GetClass}(t, \mathcal{P}_1)$ ;
5:      $S_2 = \text{GetClass}(t, \mathcal{P}_2)$ ;
6:     if  $S_1$  is not equal to  $S_2$  then
7:       return 0;
8:     end if
9:   end for
10:  return 1;
11: end procedure

```

Algorithm 7 Get the class of t in the partition \mathcal{P}

```

1: procedure GETCLASS(term  $t$ ,  $IdArray \mathcal{P}$ )  $\triangleright$  Get the partition class of terms congruent
   to  $t$  in the partition  $\mathcal{P}$ 
2:   Set  $S = \emptyset$ ;
3:   for each term  $t'$  in  $(X \cup C) \cup ((X \cup C) \times (X \cup C) \times Op)$  do
4:     if  $\mathcal{P}[t']$  is equal to  $\mathcal{P}[t]$  then
5:        $S = S \cup \{t'\}$ ;
6:     end if
7:   end for
8:   return  $S$ ;
9: end procedure

```

Algorithm 8 Create a New Atomic Value Id

```

1: procedure CREATEATOMICID  $\triangleright$  Creates a new atomic value class and assigns it to
   term  $t$ 
2:   struct ID  $idClass$ ;
3:    $idClass.valueNum \leftarrow IncCounter()$ ;
4:    $idClass.fctype \leftarrow 0$ ;
5:    $idClass.idOperand1 \leftarrow NIL$ ;
6:    $idClass.idOperand2 \leftarrow NIL$ ;
7:   return  $idClass$ ;
8: end procedure

```

Algorithm 9 Assign a Compound Value Id to a Term

```

1: procedure ASSIGNCOMPOUNDID(struct ID*  $id1$ , struct ID*  $id2$ , int  $optype$ )  $\triangleright$  Create a
   compound value class
2:   struct ID  $idClass$ ;
3:    $idClass.valueNum \leftarrow -1$ 
4:    $idClass.fctype \leftarrow optype$ ;
5:    $idClass.idOperand1 \leftarrow id1$ ;
6:    $idClass.idOperand2 \leftarrow id2$ ;
7:   return  $idClass$ ;
8: end procedure

```

Algorithm 10 Initialize Global Counter

```

1: procedure STARTCOUNTER  $\triangleright$  A global counter is initialized to 0
2:    $counter \leftarrow 0$ 
3: end procedure

```

Algorithm 11 Increment Global Counter

```

1: procedure INCCOUNTER  $\triangleright$  A global counter is incremented on each function invocation
2:    $counter \leftarrow counter + 1$ 
3:   return  $counter$ 
4: end procedure

```
