

LINEAR POLYNOMIAL FOR THE REGULARITY OF POWERS OF EDGE IDEALS OF VERY WELL-COVERED GRAPHS

A. V. JAYANTHAN AND S. SELVARAJA

ABSTRACT. Let G be a finite simple graph and $I(G)$ denote the corresponding edge ideal. In this paper we prove that if G is a very well-covered graph then for all $s \geq 1$ the regularity of $I(G)^s$ is exactly $2s + \nu(G) - 1$, where $\nu(G)$ denotes the induced matching number of G .

1. INTRODUCTION

Let $R = K[x_1, \dots, x_n]$ be a polynomial ring over a field K with the standard grading (i.e., $\deg(x_i) = 1$ for $i = 1, \dots, n$). The Castelnuovo-Mumford regularity (or simply, regularity) of a finitely generated non-zero graded R -module M , denoted by $\text{reg}(M)$, is defined to be the least integer m for which we have for every j , the j^{th} syzygy of M is generated in degrees $\leq m + j$. For a homogeneous ideal I of R , the behavior of I^s for $s \geq 2$ is studied in various contexts. It was proved by Cutkosky, Herzog and Trung, [9], and independently by Kodiyalam [23], that for a homogeneous ideal I in a polynomial ring, $\text{reg}(I^s)$ is given by a linear function for $s \gg 0$, i.e., there exist non-negative integers a, b, s_0 such that

$$\text{reg}(I^s) = as + b \text{ for all } s \geq s_0.$$

They also proved that $a \leq \deg(I)$, where $\deg(I)$ denotes the maximal degree of a minimal generator. Finding exact values of b and s_0 are non-trivial tasks, even for monomial ideals, (see, for example, [7], [17]). There have been some attempts on computing the exact form of this linear function and the stabilization index s_0 for several classes of ideals, see for example [4], [6], [10], [11], [16]. In this paper, we obtain the linear polynomial corresponding to the regularity of powers of edge ideals of very well-covered graphs.

Let G be a finite simple (no loops, no multiple edges) undirected graph on the vertex set $V(G) = \{x_1, \dots, x_n\}$. Let $K[x_1, \dots, x_n]$ be the polynomial ring in n variables, where K is a field. Then the ideal $I(G)$ generated by $\{x_i x_j \mid \{x_i, x_j\} \in E(G)\}$ is called the *edge ideal* of G . For a graph G , there exist integers b and s_0 such that $\text{reg}(I(G)^s) = 2s + b$ for all $s \geq s_0$. Our objective in this paper is to find b and s_0 , for certain class of graphs, in terms of combinatorial invariants of the graph G . Regularity of edge ideals and their powers have been studied by several authors and bounds on regularity have been computed, (see, for example, [1], [2], [3], [5], [19], [21], [25], [27], [28], [29]).

In [13], Gitler and Valencia proved that if G is a well-covered graph without isolated vertices, then $\text{ht}(I(G)) \geq \frac{|V(G)|}{2}$. In this paper, we consider the class of graphs for

1991 *Mathematics Subject Classification.* AMS Classification 2010: 13D02, 13F20, 05C70, 05E40.
Key words and phrases. Castelnuovo-Mumford regularity, Edge ideals, Very well-covered graphs.

which the above inequality is an equality, namely *very well-covered graphs*. The Cohen-Macaulayness, regularity and projective dimension of very well-covered graphs have already been looked into by several authors, [8], [22], [24], [28], [30]. Since the class of very well-covered graphs contains unmixed bipartite graphs, whiskered graphs and grafted graphs (see [8], [12]), it is interesting in the algebraic sense as well.

The regularity of powers of edge ideals of unmixed bipartite graphs have been studied by Jayanthan et al., [21]. They showed that if G is an unmixed bipartite graphs, then $\text{reg}(I(G)^s) = 2s + \nu(G) - 1$ for all $s \geq 1$. Mahmoudi et al., [24], showed that for a very well-covered graph G , $\text{reg}(I(G)) = \nu(G) + 1$, where $\nu(G)$ denotes the induced matching number of G . Since unmixed bipartite graphs are very well-covered graphs, it is natural to ask if the same result generalizes to very well-covered graphs. Recently, Norouzi et al. showed that if G is a very well-covered graph with $\text{odd-girth}(G) \geq 2k + 1$, then $\text{reg}(I(G)^s) = 2s + \nu(G) - 1$, for $1 \leq s \leq k - 2$, [28].

The main result of this paper is the following:

Theorem 1.1. (*Theorem 4.8*) *Let G be a very well-covered graph. Then for all $s \geq 1$,*

$$\text{reg}(I(G)^s) = 2s + \nu(G) - 1.$$

Therefore, for this class of graphs, we have $b = \nu(G) - 1$ and $s_0 = 1$. As an immediate consequence, we get that the above equality holds for an unmixed bipartite graphs and whiskered graphs.

Our paper is organized as follows. In Section 2, we collect the necessary notation, terminology and some results that are used in the rest of the paper. The main tool in obtaining $\text{reg}(I(G)^{s+1})$ is a result of Banerjee which gives an upper bound on $\text{reg}(I(G)^{s+1})$ in terms of $\text{reg}((I(G)^{s+1} : M))$ and $\text{reg}(I(G)^s)$, where M is minimal generator of $I(G)^s$. In Section 3, we prove that the regularity of $(I(G)^{s+1} : M)$ is bounded above by $\nu(G) + 1$, when $(I(G)^{s+1} : M)$ is squarefree. We study the case when $(I(G)^{s+1} : M)$ has square monomial generators in Section 4 and show that in this case also, the regularity is bounded above by $\nu(G) + 1$. Using these upper bounds, we prove our main result.

2. PRELIMINARIES

Throughout this paper, G denotes a finite simple graph without isolated vertices. For a graph G , $V(G)$ and $E(G)$ denote the set of all vertices and the set of all edges of G respectively. A subgraph $H \subseteq G$ is called *induced* if for $u, v \in V(H)$, $\{u, v\} \in E(H)$ if and only if $\{u, v\} \in E(G)$. For $\{u_1, \dots, u_r\} \subseteq V(G)$, let $N_G(u_1, \dots, u_r) = \{v \in V(G) \mid \{u_i, v\} \in E(G) \text{ for some } 1 \leq i \leq r\}$ and $N_G[u_1, \dots, u_r] = N_G(u_1, \dots, u_r) \cup \{u_1, \dots, u_r\}$. For $U \subseteq V(G)$, define $G \setminus U$ to be the induced subgraph of G on the vertex set $V(G) \setminus U$.

A *matching* in a graph G is a collection of pairwise disjoint edges. A matching M of a graph G is called an *induced matching* if no two edges of M are joined by an edge of G . The largest size of an induced matching in G is called its *induced matching number* and denoted by $\nu(G)$. A subset X of $V(G)$ is called an *independent set* if $\{x, y\} \notin E(G)$ for $x, y \in X$. An independent set is said to be a *maximal independent set* if it is maximal, with respect to inclusion, among the independent sets.

A subset $M \subseteq V(G)$ is a *vertex cover* of G if for each $e \in E(G)$, $e \cap M \neq \emptyset$. If M is minimal with respect to inclusion, then M is called a *minimal vertex cover* of G . A graph

G is called *unmixed* (also called *well-covered*) if all minimal vertex covers of G have the same number of elements.

A graph G is called *very well-covered* if it is unmixed without isolated vertices and with $\text{ht}(I(G)) = \frac{|V(G)|}{2}$. The following is a useful result on very well-covered graphs that allow us to assume certain order on their vertices and edges.

Lemma 2.1. [14, Corollary 3.2] *Let G be a very well-covered graph with $2h$ vertices. Then there is a relabeling of vertices $V(G) = \{x_1, \dots, x_h, y_1, \dots, y_h\}$ such that the following two conditions hold:*

- (1) $X = \{x_1, \dots, x_h\}$ is a minimal vertex cover of G and $Y = \{y_1, \dots, y_h\}$ is a maximal independent set of G ;
- (2) For all $1 \leq i \leq h$, $\{x_i, y_i\} \in E(G)$.

The concept of *even-connectedness* was introduced by Banerjee in [3]. This has emerged as a fine tool in the inductive process of computation of asymptotic regularity. We recall the definition and some of its important properties from [3].

Definition 2.2. *Let G be a graph. Two vertices u and v (u may be same as v) are said to be even-connected with respect to an s -fold products $e_1 \cdots e_s$, where e_i 's are edges of G , not necessarily distinct, if there is a path $p_0 p_1 \cdots p_{2k+1}$, $k \geq 1$ in G such that:*

- (1) $p_0 = u, p_{2k+1} = v$.
- (2) For all $0 \leq \ell \leq k - 1$, $p_{2\ell+1} p_{2\ell+2} = e_i$ for some i .
- (3) For all i , $|\{\ell \geq 0 \mid p_{2\ell+1} p_{2\ell+2} = e_i\}| \leq |\{j \mid e_j = e_i\}|$.
- (4) For all $0 \leq r \leq 2k$, $p_r p_{r+1}$ is an edge in G .

The next theorem describes the minimal generators of the ideal $(I(G)^{s+1} : M)$, where M is minimal generator of $I(G)^s$ for $s \geq 1$.

Theorem 2.3. [3, Theorem 6.1 and Theorem 6.7] *Let G be a graph with edge ideal $I = I(G)$, and let $s \geq 1$ be an integer. Let M be a minimal generator of I^s . Then $(I^{s+1} : M)$ is minimally generated by monomials of degree 2, and uv (u and v may be the same) is a minimal generator of $(I^{s+1} : M)$ if and only if either $\{u, v\} \in E(G)$ or u and v are even-connected with respect to M .*

Polarization is a process to obtain a squarefree monomial ideal from a given monomial ideal.

Definition 2.4. *Let $M = x_1^{a_1} \cdots x_n^{a_n}$ be a monomial in $R = K[x_1, \dots, x_n]$. Then we define the squarefree monomial $P(M)$ (polarization of M) as*

$$P(M) = x_{11} \cdots x_{1a_1} x_{21} \cdots x_{2a_2} \cdots x_{n1} \cdots x_{na_n}$$

in the polynomial ring $S = K[x_{ij} \mid 1 \leq i \leq n, 1 \leq j \leq a_i]$. If $I = (M_1, \dots, M_q)$ is an ideal in R , then the polarization of I , denoted by \tilde{I} , is defined as $\tilde{I} = (P(M_1), \dots, P(M_q))$.

For various properties of polarization, we refer the reader to [18]. In this paper, we repeatedly use one of the important properties of the polarization, namely:

Corollary 2.5. [18, Corollary 1.6.3(a)] *Let I be a monomial ideal in $K[x_1, \dots, x_n]$. Then*

$$\text{reg}(I) = \text{reg}(\tilde{I}).$$

3. BOUNDING THE REGULARITY: THE SQUAREFREE MONOMIAL CASE

We obtain the asymptotic expression for the regularity by using induction and [3, Theorem 5.2] which says that $\text{reg}(I(G)^{s+1}) \leq \max_M \{\text{reg}(I(G)^{s+1} : M) + 2s, \text{reg}(I(G)^s)\}$, where M is a minimal monomial generator of $I(G)^s$. For this purpose, one needs to compute $\text{reg}(I(G)^{s+1} : M)$. In this section, we obtain an upper bound for the regularity of $(I(G)^{s+1} : M)$, when $(I(G)^{s+1} : M)$ is squarefree. We first fix certain set-up for the class of graphs that we consider throughout this paper.

Set-up 3.1. *Let G be a graph with $2h$ vertices, none of which are isolated and $V(G) = X \cup Y$, where $X = \{x_1, \dots, x_h\}$ is a minimal vertex cover of G and $Y = \{y_1, \dots, y_h\}$ is a maximal independent set of G such that $\{x_i, y_i\} \in E(G)$, for all $1 \leq i \leq h$.*

The following result is being used repeatedly in this paper:

Lemma 3.2. [8, Proposition 2.3] *Let G be a graph as in Set-up 3.1. Then G is a very well-covered if and only if the following conditions hold:*

- (1) *if $\{z_i, x_j\}, \{y_j, x_k\} \in E(G)$, then $\{z_i, x_k\} \in E(G)$ for distinct i, j, k and for $z_i \in \{x_i, y_i\}$;*
- (2) *if $\{x_i, y_j\} \in E(G)$, then $\{x_i, x_j\} \notin E(G)$.*

We make an observation which follows directly follows from the Lemma 3.2.

Observation 3.3. *If G is a very well-covered graph as in Set-up 3.1, then for any $1 \leq i \leq h$, $G \setminus N_G[x_i, y_i]$, $G \setminus N_G[x_i]$ and $G \setminus \{x_i, y_i\}$ are very well-covered.*

We begin by showing that if we start with a very well-covered graph, then we may make certain relabelling of the vertices with the hypotheses of Set-up 3.1 being preserved.

Lemma 3.4. *Let G be a very well-covered graph satisfying Set-up 3.1. For an $i \in \{1, \dots, h\}$, let $N_G(x_i) \setminus X = \{y_{i_1}, \dots, y_{i_t}\}$, for some $1 \leq i_1, \dots, i_t \leq h$. Let*

$$X' = \left\{ x'_j \mid j \in \{1, \dots, h\}, x'_j = \begin{cases} y_j & \text{if } j \in \{i_1, \dots, i_t\} \\ x_j & \text{otherwise} \end{cases} \right\}$$


and $Y' = V(G) \setminus X'$. Let G_1 denote the graph with the above relabelling. Then G_1 with $V(G_1) = X' \cup Y'$ satisfies Set-up 3.1 and the properties (1) and (2) of Lemma 3.2.

Proof. Let $X' = \{a_1, \dots, a_h\}$ and $Y' = \{b_1, \dots, b_h\}$. First we claim that Y' is a maximal independent set of G_1 . Suppose not, then there exists an edge $\{b_p, b_q\} \in E(G_1)$. Since G is a very well-covered, at least one of b_p and b_q is in $\{x_{j_1}, \dots, x_{j_t}\}$. Suppose $b_p, b_q \in \{x_{j_1}, \dots, x_{j_t}\}$. Let $b_p = x_{j_r}$ and $b_q = x_{j_{r'}}$. We have $\{x_i, y_{j_r}\}, \{x_i, y_{j_{r'}}\}, \{x_{j_r}, x_{j_{r'}}\} \in E(G)$. Since G is a very well-covered graph, there is an edge $\{x_i, x_{j_{r'}}\}$ in G . This contradicts Lemma 3.2(2). Suppose $b_q = x_{j_r} \in \{x_{j_1}, \dots, x_{j_t}\}$ for some $1 \leq r \leq t$ and $b_p \notin \{x_{j_1}, \dots, x_{j_t}\}$. Therefore, $\{x_i, y_{j_r}\} \in E(G)$. Since G is a very well-covered and $\{x_i, y_{j_r}\}, \{x_{j_r}, b_p\} \in E(G)$, we have $\{x_i, b_p\} \in E(G)$, i.e., $b_p \in \{x_{j_1}, \dots, x_{j_t}\}$, which is a contradiction to the assumption that $b_p \notin \{x_{j_1}, \dots, x_{j_t}\}$. Therefore Y' is a maximal independent set in G_1 . Hence, X' is a minimal vertex cover of G_1 . \square

In the Lemma 3.4, we have shown that we may conveniently swap some of the x_i 's and y_i 's preserving the hypotheses of Set-up 3.1 and the properties (1) and (2) of Lemma 3.2.

However, arbitrary swapping of x_i 's and y_i 's may not preserve the hypotheses of Set-up 3.1 as can be seen from the following example.

Example 3.5. Let G be the very well-covered graph on $\{x_1, x_2, x_3, y_1, y_2, y_3\}$ as given in the figure below.



Let G_1 be the graph obtained from G by swapping the vertices x_2 and y_2 , i.e., $V(G_1) = X' \cup Y'$, where $X' = \{x_1, x'_2, x_3\}$ and $Y' = \{y_1, y'_2, y_3\}$, where $x'_2 = y_2$ and $y'_2 = x_2$.

We can see that Y' is not an independent set of G_1 and hence G_1 does not satisfy the Set-up 3.1. The third graph, G_2 , is obtained by swapping the vertices x_3 and y_3 . Note that $N_G(x_2) \setminus X = \{y_3\}$. In this case, by taking $X' = \{x_1, x_2, x'_3\}$ and $Y' = \{y_1, y_2, y'_3\}$, it can be seen that G_2 satisfies the Set-up 3.1.

We now show that adding edges between even-connected vertices preserves the very well-covered property of a graph, provided there are no vertices which are even-connected to itself.

Theorem 3.6. Let G be a very well-covered graph with $2h$ vertices. If for some $e \in E(G)$, the ideal $(I(G)^2 : e)$ is squarefree, then G' is a very well-covered graph, where G' is the graph associated to $(I(G)^2 : e)$.

Proof. By Lemma 2.1, there is a relabeling of vertices $V(G) = X \cup Y$ such that G satisfies the Set-up 3.1, where $X = \{x_1, \dots, x_h\}$ and $Y = \{y_1, \dots, y_h\}$. Suppose $e = \{x_n, y_n\}$, for some $1 \leq n \leq h$. If u and v are even-connected with respect to $\{x_n, y_n\}$, then by Lemma 3.2, $\{u, v\} \in E(G)$. Therefore by Theorem 2.3 $(I(G)^2 : e) = I(G)$ so that G' is a very well-covered graph.

Suppose $e = \{x_n, y_m\}$, for some $1 \leq n, m \leq h$. Note that G' also satisfies Set-up 3.1. We need to show that G' satisfies Lemma 3.2(1)-(2). If $\{z_i, x_j\}, \{y_j, x_k\} \in E(G)$, then by Lemma 3.2, $\{z_i, x_k\} \in E(G)$. Therefore $\{z_i, x_k\} \in E(G')$. Suppose $\{z_i, x_j\} \in E(G)$ and $\{y_j, x_k\} \in E(G') \setminus E(G)$. Let $y_j p_1 p_2 x_k$ be an even-connection in G with respect to $\{p_1, p_2\} = e$. Since $\{z_i, x_j\}$ and $\{y_j, p_1\}$ are in $E(G)$ and G is very well-covered, $\{z_i, p_1\} \in E(G)$. Hence $z_i p_1 p_2 x_k$ is an even-connection in G with respect to e so that $\{z_i, x_k\} \in E(G')$. Similarly we can prove that, if $\{z_i, x_j\} \in E(G') \setminus E(G)$ and $\{y_j, x_k\} \in E(G)$, then $\{z_i, x_k\} \in E(G')$. Suppose $\{z_i, x_j\}, \{y_j, x_k\} \in E(G') \setminus E(G)$. Let $z_i p_1 p_2 x_j$ and $y_j q_1 q_2 x_k$ be an even-connection in G with respect to $e = \{p_1, p_2\} = \{q_1, q_2\}$. If $p_1 = q_1$, then there is an even-connection $z_i(p_1 = q_1)q_2 x_k$ in G with respect to e . Suppose $p_1 = q_2$. Then $\{p_2, x_j\} \in E(G)$ and $\{y_j, p_2\} \in E(G)$. This contradicts the fact that G is a very well-covered graph. Therefore $\{z_i, x_k\} \in E(G')$.

Now we show that if $\{x_i, y_j\} \in E(G')$, then $\{x_i, x_j\} \notin E(G')$. Suppose $\{x_i, y_j\} \in E(G)$ and $\{x_i, x_j\} \in E(G')$. Note that $\{x_i, x_j\} \notin E(G)$. Let $x_i p_1 p_2 x_j$ be an even-connection in G with respect to $e = \{p_1, p_2\}$. Since $\{x_i, y_j\}, \{x_j, p_2\} \in E(G)$, $\{x_i, p_2\} \in E(G)$. Then there is an even-connection $x_i p_1 p_2 x_i$ in G with respect to e . Therefore $x_i^2 \in (I(G)^2 : e)$, which is a contradiction. Therefore, $\{x_i, x_j\} \notin E(G')$. Suppose $\{x_i, x_j\}, \{x_i, y_j\} \in E(G') \setminus E(G)$. Therefore, there exist even-connections, $x_i p_1 p_2 x_j$ and $x_i q_1 q_2 y_j$. If $p_1 = q_1$, then there exist edges $\{p_2, x_j\}$ and $\{p_2, y_j\}$ in $E(G)$ which contradicts the assumption that G is very

well-covered. If $p_1 = q_2$, then there exists an even-connection $x_i p_2 p_1 x_i$. Therefore, $x_i^2 \in (I(G)^2 : e)$ which contradicts the assumption that $(I(G)^2 : e)$ is a squarefree monomial ideal. Hence G' is a very well-covered graph.

Suppose $e = \{x_n, x_m\}$, for some $1 \leq n, m \leq h$ and $N_G(x_m) \setminus X = \{y_{j_1}, \dots, y_{j_t}\}$. Let X' and Y' be as in Lemma 3.4. Let G_1 denote the graph with the above relabelling. By Lemma 3.4, $V(G_1) = X' \cup Y'$ satisfies Set-up 3.1 and Lemma 3.2. Let e' denote the edge e after relabelling. Then $e' = \{x_n, y_m\}$. Since $I(G_1)$ is obtained from $I(G)$ by relabelling certain variables, it follows that $(I(G_1)^2 : e')$ is also a squarefree monomial ideal. Let G'_1 be the graph associated to $(I(G_1)^2 : e')$. By previous case, G'_1 is a very well-covered graph. Since G'_1 is also obtained by relabelling of certain vertices of G' , it follows that G' is a very well-covered graph. \square

The below example shows that if $(I(G)^2 : e)$ is not squarefree, then the assertion of the Theorem 3.6 need not necessarily be true.

Example 3.7. *Let*

$$I = (x_1 y_1, x_2 y_2, x_3 y_3, x_4 y_4, x_1 x_2, x_1 x_4, x_1 y_3, x_2 y_3, x_2 x_4, x_3 x_4) \subset R = K[x_1, \dots, x_4, y_1, \dots, y_4]$$

and G be the associated graph. By Lemma 2.1, G is a very well-covered graph. It can be seen that $x_4^2, y_3^2 \in (I^2 : x_1 x_2)$ and that

$$I(G') = (\widetilde{I^2 : x_1 x_2}) = I + (y_1 y_2, y_1 x_4, y_1 y_3, y_2 x_4, y_2 y_3, y_3 x_4, x_4 z_2, y_3 z_1) \subseteq R[z_1, z_2].$$

Since $\mathfrak{p} = (x_1, y_2, y_3, x_4)$ and $\mathfrak{q} = (x_1, x_2, x_3, y_1, y_2, y_3, y_4, z_2)$ are minimal prime ideals of $I(G')$, G' is not a very well-covered graph.

Theorem 3.8. *Let G be a graph and e_1, \dots, e_s , $s \geq 1$ be some edges of G which are not necessarily distinct. Suppose $(I(G)^{s+1} : e_1 \cdots e_s)$ is squarefree ideal. For $1 \leq i \leq s$,*

$$(I(G)^{s+1} : e_1 \cdots e_s) = ((I(G)^2 : e_i)^s : \prod_{i \neq j} e_j).$$

Proof. This result has been proved for bipartite graphs in [1, Lemma 3.7]. In our case, the assumption that $(I(G)^{s+1} : e_1 \cdots e_s)$ is squarefree implies that there are no odd cycles in the even-connections. Using this property, one can see that their proof goes through in our case as well. \square

In [21, Theorem 4.1], it was proved that if G is an unmixed bipartite graph, then so is G' , the graph associated to $(I(G)^{s+1} : e_1 \cdots e_s)$. As a consequence of the above results, we generalize this to the case of very well-covered graphs G with $(I(G)^{s+1} : e_1 \cdots e_s)$ squarefree.

Corollary 3.9. *(with hypothesis as in Theorem 3.8). If G is a very well-covered graph, then so is the graph G' associated to $(I(G)^{s+1} : e_1 \cdots e_s)$, for every s -fold product $e_1 \cdots e_s$ and $s \geq 1$.*

Proof. We prove that G' is very well-covered by induction on s . If $s = 1$, then the assertion follows from Theorem 3.6. Assume by induction that for any very well-covered graph H

and edges f_1, \dots, f_{s-1} with $(I(H)^s : f_1 \cdots f_{s-1})$ a squarefree monomial ideal, the graph associated to $(I(H)^s : f_1 \cdots f_{s-1})$ is very well-covered. By Theorem 3.8 we have

$$(I(G)^{s+1} : e_1 \cdots e_s) = ((I(G)^2 : e_i)^s : \prod_{i \neq j} e_j).$$

Note that $(I(G)^2 : e_i)$ is squarefree monomial ideal. Let H be the graph associated to $(I(G)^2 : e_i)$. By the case $s = 1$, H is a very well-covered graph. Therefore, by induction, the graph associated to $((I(G)^2 : e_i)^s : \prod_{i \neq j} e_j)$ is a very well-covered graph. \square

We finally obtain an upper bound for the regularity in the squarefree case:

Corollary 3.10. *(with hypothesis as in Theorem 3.8). Let G be a very well-covered graph. Then*

$$\text{reg}((I(G)^{s+1} : e_1 \cdots e_s)) \leq \nu(G) + 1.$$

Proof. Let G' be the graph associated to $(I(G)^{s+1} : e_1 \cdots e_s)$. By Corollary 3.9, G' is a very well-covered graph. Therefore,

$$\begin{aligned} \text{reg}((I(G)^{s+1} : e_1 \cdots e_s)) &= \nu(G') + 1 \quad (\text{by [24, Theorem 4.12]}) \\ &\leq \nu(G) + 1. \quad (\text{by [21, Proposition 4.4]}) \end{aligned}$$

\square

4. REGULARITY OF POWERS OF EDGE IDEALS OF VERY WELL-COVERED GRAPHS

In this section, we obtain an upper bound for the regularity of $(I(G)^{s+1} : e_1 \cdots e_s)$ when this is not a squarefree monomial ideal. Using these results we prove the main theorem, namely, the asymptotic expression for the regularity of powers of edge ideals of very well-covered graphs.

Set-up 4.1. *Let G be a very well-covered graph with $2h$ vertices and $V(G) = \{x_1, \dots, x_h, y_1, \dots, y_h\}$ satisfying Lemma 2.1(1 - 2). By Theorem 2.3 and Definition 2.4, $(I(G)^{s+1} : e_1 \cdots e_s)$ is a quadratic squarefree monomial ideal in an appropriate polynomial ring. Let G' be the graph associated to $(I(G)^{s+1} : e_1 \cdots e_s)$.*

If $(I(G)^{s+1} : e_1 \cdots e_s)$ is not squarefree, then there will be new vertices along with new edges in G' and hence it need not necessarily be a very well-covered graph, see Example 3.7. Our aim in this section is to get an upper bound for $\text{reg}(I(G'))$. For this purpose, we need to get more details about the structure of the graph G' . With this aim in mind, in the next three Lemmas, we describe some of the edges that are in G' which are possibly not in G .

Lemma 4.2. *Let the notation be as in Set-up 4.1. For $t_i \in \{x_i, y_i\}$, if $t_i x_j$ and $y_j x_k \in I(G')$, then either $t_i x_k \in I(G')$ or $t_i y_j \in I(G')$ for distinct i, j, k .*

Proof. Suppose $\{t_i, x_j\} \in E(G)$. If $\{y_j, x_k\} \in E(G)$, then by Lemma 3.2, $\{t_i, x_k\} \in E(G)$. Suppose y_j and x_k are even-connected with respect to $e_1 \cdots e_s$ in G . For some $r' \geq 1$, let $(y_j = p_0)p_1 \cdots p_{2r'}(p_{2r'+1} = x_k)$ be an even-connection in G . Since $\{t_i, x_j\}, \{y_j, p_1\} \in E(G)$, by Lemma 3.2, $\{t_i, p_1\} \in E(G)$. Then there is an even-connection $t_i p_1 \cdots (p_{2r'+1} = x_k)$ with respect to $e_1 \cdots e_s$ in G . Therefore, $\{t_i, x_k\} \in E(G')$. Similarly we can prove that, if $\{t_i, x_j\} \in E(G') \setminus E(G)$ and $\{y_j, x_k\} \in E(G)$, then $\{t_i, x_k\} \in E(G')$.

Suppose $\{t_i, x_j\}, \{y_j, x_k\} \in E(G') \setminus E(G)$. For some $r_1 \geq 1$ and $r_2 \geq 1$, let

$$(t_i = q_0)q_1 \cdots q_{2r_1}(q_{2r_1+1} = x_j) \text{ and } (y_j = s_0)s_1 \cdots s_{2r_2}(s_{2r_2+1} = x_k)$$

be even-connections with respect to $e_1 \cdots e_s$ in G . Suppose $\{s_{2\alpha+1}, s_{2\alpha+2}\}$ and $\{q_{2\beta+1}, q_{2\beta+2}\}$ do not have common vertices, for all $0 \leq \alpha \leq r_2 - 1$ and $0 \leq \beta \leq r_1 - 1$. Since $\{q_{2r_1}, x_j\}, \{y_j, s_1\} \in E(G)$, $\{q_{2r_1}, s_1\} \in E(G)$. Then there is an even-connection $(t_i = q_0)q_1 \cdots q_{2r_1}s_1 \cdots s_{2r_2}(s_{2r_2+1} = x_k)$ with respect to $e_1 \cdots e_s$ in G . If for some $0 \leq \alpha \leq r_2 - 1$, $0 \leq \beta \leq r_1 - 1$, $\{s_{2\alpha+1}, s_{2\alpha+2}\}$ and $\{q_{2\beta+1}, q_{2\beta+2}\}$ have a common vertex, then by [3, Lemma 6.13], t_i is even-connected to either y_j or x_k with respect to $e_1 \cdots e_s$ in G . Therefore either $\{t_i, y_j\} \in E(G')$ or $\{t_i, x_k\} \in E(G')$. \square

Lemma 4.3. *Let the notation be as in Set-up 4.1. Suppose $(u = p_0)p_1 \cdots p_{2k}(p_{2k+1} = v)$ is an even-connection in G with respect to $e_1 \cdots e_s$, for some $k \geq 1$. If $\{w, p_i\} \in E(G')$, for some $0 \leq i \leq 2k + 1$, then either $\{u, w\} \in E(G')$ or $\{v, w\} \in E(G')$.*

Proof. If $i = 0, 2k + 1$, then we are done. Assume that $i = 2j + 1$, for some $j \geq 0$. For some $j \geq 0$, let $(w = q_0)q_1 \cdots (q_{2j+1} = p_i)$ be an even-connection with respect to $e_1 \cdots e_s$ in G . If $\{q_{2\alpha+1}, q_{2\alpha+2}\}$ and $\{p_{2\beta+1}, p_{2\beta+2}\}$ do not have a common vertex, for all $0 \leq \alpha \leq j - 1$, $j \leq \beta \leq k - 1$, then $(w = q_0)q_1 \cdots (q_{2j+1} = p_i)p_{i+1} \cdots (p_{2k+1} = v)$ is an even-connection with respect to $e_1 \cdots e_s$ in G . Therefore, $wv \in I(G')$. If $\{q_{2\alpha+1}, q_{2\alpha+2}\}$ and $\{p_{2\beta+1}, p_{2\beta+2}\}$ have a common vertex, for some $0 \leq \alpha \leq j - 1$, $j \leq \beta \leq k - 1$, then by [3, Lemma 6.13], w is even-connected either to u or to v in G . Therefore either $wu \in I(G')$ or $wv \in I(G')$. If $i = 2j + 2$, then proof is similar. \square

In the next lemma, we further obtain more even-connected edges in G' . Let G be a very well-covered graph as in Set-up 3.1. For $u = x_i$ or y_i , set $[u] = \{x_i, y_i\}$ and $N_G[[u]] = N_G[x_i, y_i]$.

Lemma 4.4. *Let the notation be as in Set-up 4.1. Let $u^2 \in (I(G)^{s+1} : e_1 \cdots e_s)$. If $a \in (N_{G'}([u] \setminus u) \cap V(G))$ and $b \in N_G[[u]]$, then $\{a, b\} \in E(G')$.*

Proof. Since $u^2 \in (I(G)^{s+1} : e_1 \cdots e_s)$, we have an even-connection $(p_0 = u)p_1 \cdots p_{2k}(p_{2k+1} = u)$ with respect to $e_1 \cdots e_s$ in G , for some $k \geq 1$. Note that, since $\{p_1, u\}, \{p_{2k}, u\} \in E(G)$, if $b \in N_G([u] \setminus u)$, then $p_1 \neq b$ and $p_{2k} \neq b$. Therefore, $(u = p_0)p_1 \cdots p_{2k}b$ and $(u = p_{2k+1})p_{2k} \cdots p_1b$ are an even-connections in G so that $\{u, b\} \in E(G')$. Hence, if $a = u$, then $\{a, b\} \in E(G')$.

We now assume that $a \neq u$. Suppose $\{a, [u] \setminus u\} \in E(G)$. If either $b = u$ or $b = [u] \setminus u$, then we are done. If $b \in N_G(u)$, then $\{a, b\} \in E(G)$. Suppose $b \in N_G([u] \setminus u)$. Since we have $\{u, b\} \in E(G')$, by the proof of Lemma 4.2, $\{a, b\} \in E(G')$.

Suppose $\{a, [u] \setminus u\} \in E(G') \setminus E(G)$. For some $t \geq 1$, let $(q_0 = a)q_1 \cdots (q_{2t+1} = [u] \setminus u)$ be an even-connection with respect to $e_1 \cdots e_s$ in G . If $\{u, b\} \in E(G)$, then by the proof of Lemma 4.2, $\{a, b\} \in E(G')$. Suppose $\{[u] \setminus u, b\} \in E(G)$. Note that u is even-connected to b with even-connections $(u = p_0)p_1 \cdots p_{2k}b$ and $(u = p_{2k+1})p_{2k} \cdots p_1b$ with respect to $e_1 \cdots e_s$ in G . Suppose $\{p_{2\lambda+1}, p_{2\lambda+2}\} \neq \{q_{2\lambda'+1}, q_{2\lambda'+2}\}$, for all $0 \leq \lambda \leq k - 1$, $0 \leq \lambda' \leq t - 1$. Then either $\{p_{2\lambda+1}, p_{2\lambda+2}\} \cap \{q_{2\lambda'+1}, q_{2\lambda'+2}\} = \emptyset$ or $\{p_{2\lambda+1}, p_{2\lambda+2}\} \cap \{q_{2\lambda'+1}, q_{2\lambda'+2}\}$ is a vertex. In either case, it follows from the proof of Lemma 4.2 that $\{a, b\} \in E(G')$. Suppose $\{p_{2\lambda+1}, p_{2\lambda+2}\} = \{q_{2\lambda'+1}, q_{2\lambda'+2}\}$, for some $0 \leq \lambda \leq k - 1$, $0 \leq \lambda' \leq t - 1$. Choose

the smallest λ' such that $\{p_{2\lambda+1}, p_{2\lambda+2}\} = \{q_{2\lambda'+1}, q_{2\lambda'+2}\}$ for some $0 \leq \lambda \leq k-1$. If $p_{2\lambda+1} = q_{2\lambda'+1}$ and $p_{2\lambda+2} = q_{2\lambda'+2}$, then there is an even-connection

$$(a = q_0)q_1 \cdots (q_{2\lambda'+1} = p_{2\lambda+1})(q_{2\lambda'+2} = p_{2\lambda+2})p_{2\lambda+3} \cdots p_{2k}b$$

with respect to $e_1 \cdots e_s$ in G . If $p_{2\lambda+1} = q_{2\lambda'+2}$ and $p_{2\lambda+2} = q_{2\lambda'+1}$, then there is an even-connection

$$(a = q_0)q_1 \cdots (q_{2\lambda'+1} = p_{2\lambda+2})(q_{2\lambda'+2} = p_{2\lambda+1})p_{2\lambda} \cdots p_1b$$

with respect to $e_1 \cdots e_s$ in G . Therefore $\{a, b\} \in E(G')$.

If $b = u$ and $a \in (N_{G'}([u] \setminus u) \cap V(G))$, then proceeding as in the previous case of the proof, one can show that $\{a, b\} \in E(G')$. \square

To get an upper bound for the regularity of $I(G')$, we need to bound the regularity of certain induced subgraphs of G' . In the next two lemmas, we understand more closely the structure of some of the induced subgraphs of G' . This, in turn, helps us during the induction process.

Lemma 4.5. *Let the notation be as in Set-up 4.1. Let $y \in V(G)$ and $H = G \setminus N_G[y]$. If $\{e_1, \dots, e_s\} \cap E(H) = \{e_{i_1}, \dots, e_{i_t}\}$ and H' is the graph associated to $(I(H)^{t+1} : e_{i_1} \cdots e_{i_t})$, then $G' \setminus N_{G'}[y]$ is an induced subgraph of H' . In particular, $\text{reg}(I(G' \setminus N_{G'}[y])) \leq \text{reg}(I(H'))$.*

Proof. Let $\{u, v\} \in E(G' \setminus N_{G'}[y])$. By Theorem 2.3, either $\{u, v\} \in E(G)$ or u is an even-connected to v in G with respect to $e_1 \cdots e_s$. If $\{u, v\} \in E(G)$, then $\{u, v\} \in E(H)$. Let $(u = p_0)p_1 \cdots (p_{2k+1} = v)$ be an even-connection in G with respect to $e_1 \cdots e_s$ for some $k \geq 0$. If $p_i \in N_{G'}[y]$, for some $0 \leq i \leq 2k+1$, then by Lemma 4.3, y is even-connected either to u or to v . This contradicts the assumption that $\{u, v\} \in G' \setminus N_{G'}[y]$. Therefore, for each $0 \leq i \leq 2k+1$, $p_i \notin N_{G'}[y]$. Hence $\{u, v\} \in E(H')$, which proves $G' \setminus N_{G'}[y]$ is a subgraph of H' . If $a, b \in V(G' \setminus N_{G'}[y])$ is such that $\{a, b\} \in E(H)$, then $\{a, b\} \in E(G' \setminus N_{G'}[y])$. Hence $G' \setminus N_{G'}[y]$ is an induced subgraph of H . The assertion on the regularity follows from [20, Proposition 4.1.1]. \square

It may be noted that, in the above proof, we did not really use the very well-covered property of G . The result holds true for an arbitrary graph.

Lemma 4.6. *Let the notation be as in Set-up 4.1. Let $u^2 \in (I(G)^{s+1} : e_1 \cdots e_s)$, $t \in (N_{G'}([u] \setminus u) \cap V(G))$ and $H = G \setminus N_G[[u]]$. Then $G' \setminus N_{G'}[t]$ is an induced subgraph of H' , where $\{e_1, \dots, e_s\} \cap E(H) = \{e_{i_1}, \dots, e_{i_k}\}$ and H' is the graph associated to $(I(H)^{k+1} : e_{i_1} \cdots e_{i_k})$. In particular, $\text{reg}(I(G' \setminus N_{G'}[t])) \leq \text{reg}(I(H'))$.*

Proof. Let $\{a, b\} \in E(G' \setminus N_{G'}[t])$. By Theorem 2.3, either $\{a, b\} \in E(G)$ or a is an even-connected to b in G with respect to $e_1 \cdots e_s$. Suppose $\{a, b\} \in E(G)$. If $\{a, b\} \cap N_G[[u]] = \emptyset$, then $\{a, b\} \in E(H)$. If $\{a, b\} \cap N_G[[u]] \neq \emptyset$, then by Lemma 4.4, either $\{a, t\} \in E(G')$ or $\{b, t\} \in E(G')$. This is a contradiction to $\{a, b\} \in E(G' \setminus N_{G'}[t])$. Therefore, if $\{a, b\} \in E(G)$, then $\{a, b\} \cap N_G[[u]] = \emptyset$ and hence $\{a, b\} \in E(H)$.

Suppose $\{a, b\} \in E(G') \setminus E(G)$. For $r \geq 1$, let $(a = q_0)q_1 \cdots q_{2r}(q_{2r+1} = b)$ be an even-connection in G with respect to $e_1 \cdots e_s$. If $q_i \in N_G[[u]]$, for some i , then by Lemma 4.4, $\{t, q_i\} \in E(G')$. Therefore, by Lemma 4.3, either $\{t, a\} \in E(G')$ or $\{t, b\} \in E(G')$. This

is a contradiction to the assumption that $\{a, b\} \in E(G' \setminus N_{G'}[t])$. Therefore $q_i \notin N_G[[u]]$, for all $0 \leq i \leq 2r + 1$ which implies that a is an even-connected to b in H with respect to $e_{i_1} \cdots e_{i_k}$, i.e., $\{a, b\} \in E(H')$. Hence $G' \setminus N_{G'}[t]$ is an induced subgraph of H' . The assertion on the regularity follows from [20, Proposition 4.1.1]. \square

Now we prove that the regularity of $I(G')$ is bounded above by $\nu(G) + 1$.

Theorem 4.7. *Let G be a very well-covered graph and e_1, \dots, e_s be edges of G , for some $s \geq 1$. Then,*

$$\text{reg}((I(G)^{s+1} : e_1 \cdots e_s)) \leq \nu(G) + 1.$$

Proof. For any graph K , let

$$W_K(e_1 \cdots e_s) = \left\{ u \in V(K) \mid u \text{ is an even-connected to itself in } K \text{ with respect to } e_1 \cdots e_s \right\}.$$

Let G' be the graph associated to $(I(G)^{s+1} : e_1 \cdots e_s)$ contained in an appropriate polynomial ring R_1 and $|W_G(e_1 \cdots e_s)| = r$. We prove the assertion by induction on r .

If $r = 0$, then for any $e_1, \dots, e_s \in E(G)$, $s \geq 1$, $(I(G)^{s+1} : e_1 \cdots e_s)$ is a squarefree monomial ideal. Therefore, by Corollary 3.10,

$$\text{reg}((I(G)^{s+1} : e_1 \cdots e_s)) \leq \nu(G) + 1.$$

By induction, assume that if L is a very well-covered graph with $|W_L(f_1 \cdots f_s)| < r$ for $f_1, \dots, f_s \in E(L)$, then $\text{reg}((I(L)^{s+1} : f_1 \cdots f_s)) \leq \nu(L) + 1$.

Let G be a very well-covered graph with $|W_G(e_1 \cdots e_s)| = r$ for $e_1, \dots, e_s \in E(G)$. Let G' be the graph associated to $(I(G)^{s+1} : e_1 \cdots e_s)$, for some $e_1, \dots, e_s \in E(G)$. Set $W_G(e_1 \cdots e_s) = \{u_1, \dots, u_r\}$, $[u_r] \setminus u_r = u'_r$, $U = (N_{G'}(u'_r) \cap V(G)) = \{t_1, \dots, t_l\}$ and $J = I(G')$. It follows from the exact sequences

$$\begin{array}{ccccccc} 0 & \longrightarrow & \frac{R_1}{(J : t_1)}(-1) & \xrightarrow{\cdot t_1} & \frac{R_1}{J} & \longrightarrow & \frac{R_1}{(J, t_1)} \longrightarrow 0; \\ 0 & \longrightarrow & \frac{R_1}{((J, t_1) : t_2)}(-1) & \xrightarrow{\cdot t_2} & \frac{R_1}{(J, t_1)} & \longrightarrow & \frac{R_1}{(J, t_1, t_2)} \longrightarrow 0; \\ & & \vdots & & \vdots & & \vdots \\ 0 & \longrightarrow & \frac{R_1}{((J, t_1, \dots, t_{l-1}) : t_l)}(-1) & \xrightarrow{\cdot t_l} & \frac{R_1}{(J, t_1, \dots, t_{l-1})} & \longrightarrow & \frac{R_1}{(J, U)} \longrightarrow 0 \end{array}$$

that

$$\text{reg}(R_1/J) \leq \max \left\{ \begin{array}{l} \text{reg} \left(\frac{R_1}{(J:t_1)} \right) + 1, \text{reg} \left(\frac{R_1}{((J,t_1):t_2)} \right) + 1, \\ \dots \dots \dots \\ \text{reg} \left(\frac{R_1}{(J,t_1,\dots,t_{l-1}):t_l} \right) + 1, \text{reg} \left(\frac{R_1}{(J,U)} \right). \end{array} \right.$$

We now prove that each of the regularities appearing on the right hand side of the above inequality is bounded above by $\nu(G)$.

Let $H = G \setminus N_G[u'_r]$ and $\{e_1, \dots, e_s\} \cap E(H) = \{e_{i_1}, \dots, e_{i_k}\}$. We have

$$\begin{aligned} \text{reg}(J, U) &= \text{reg}(I(G' \setminus N_{G'}[u'_r])) && \text{(by [5, Remark 2.5])} \\ &\leq \text{reg}((I(H)^{k+1} : e_{i_1} \cdots e_{i_k})) && \text{(by Lemma 4.5)} \end{aligned}$$

Since H is a very well-covered graph and $u_r^2 \notin (I(H)^{k+1} : e_{i_1} \cdots e_{i_k})$, $|W_H(e_{i_1} \cdots e_{i_k})| < r$. Hence, by induction, we get

$$\text{reg}(J, U) \leq \text{reg}(I(H)^{k+1} : e_{i_1} \cdots e_{i_k}) \leq \nu(H) + 1 \leq \nu(G) + 1.$$

Let $H = G \setminus N_G[[u_r]]$ and $E(H) \cap \{e_1, \dots, e_s\} = \{e_{i_1}, \dots, e_{i_\ell}\}$. Since H is a very well-covered graph and $u_r^2 \notin (I(H)^{\ell+1} : e_{i_1} \cdots e_{i_\ell})$, we have

$$\begin{aligned} \text{reg}(J : t_i) &= \text{reg}(I(G' \setminus N_{G'}[t_i]) \leq \text{reg}(I(H)^{\ell+1} : e_{i_1} \cdots e_{i_\ell}) && \text{(By Lemma 4.6)} \\ &\leq \nu(H) + 1 && \text{(By induction hypothesis)} \\ &\leq \nu(G), \end{aligned}$$

where the last inequality follows since $\{f_1, \dots, f_t, [u_r]\}$ is an induced matching in G if $\{f_1, \dots, f_t\}$ is an induced matching in H .

Since $((J, t_1, \dots, t_{i-1}) : t_i)$ corresponds to an induced subgraph of $(J : t_i)$, it follows that

$$\text{reg}\left(\frac{R_1}{((J, t_1, \dots, t_{i-1}) : t_i)}\right) + 1 \leq \text{reg}\left(\frac{R_1}{(J : t_i)}\right) + 1 \leq \nu(G).$$

Therefore, $\text{reg}\left(\frac{R_1}{J}\right) \leq \nu(G)$. \square

Now the main theorem can be derived as a consequence of the above results:

Theorem 4.8. *Let G be a very well-covered graph. Then for all $s \geq 1$,*

$$\text{reg}(I(G)^s) = 2s + \nu(G) - 1.$$

Proof. For any $s \geq 1$, by [5, Theorem 4.5], we have $2s + \nu(G) - 1 \leq \text{reg}(I(G)^s)$. We need to prove that $\text{reg}(I(G)^s) \leq 2s + \nu(G) - 1$, for all $s \geq 1$. We prove this by induction on s . If $s = 1$, then the assertion follows from [24, Theorem 4.12]. Assume that $s > 1$. By applying [3, Theorem 5.2] and using induction, it is enough to prove that for edges e_1, \dots, e_s of G , $\text{reg}(I(G)^{s+1} : e_1 \cdots e_s) \leq \nu(G) + 1$ for all $s \geq 1$. This follows from Theorem 4.7. \square

Since unmixed bipartite graphs are very well-covered graphs, we obtain

Corollary 4.9. [21, Corollary 5.1(1)] *If G is an unmixed bipartite graph, then for all $s \geq 1$,*

$$\text{reg}(I(G)^s) = 2s + \nu(G) - 1.$$

For a graph G on n vertices, let $W(G)$ be the whiskered graph on $2n$ vertices obtained by adding a pendent vertex (an edge to a new vertex of degree 1) to every vertex of G .

Moghimian et al., [25, Theorem 2.5] proved that $\text{reg}(I(W(C_n)^s)) = 2s + \nu(W(G)) - 1$ for all $s \geq 1$ and Jayanthan et al. [21, Corollary 5.1(2)] proved that if G is a bipartite graph, then $\text{reg}(I(W(G))^s) = 2s + \nu(W(G)) - 1$ for all $s \geq 1$. Since whiskered graphs are very well-covered graphs, we obtain asymptotic regularity expression for this class of graphs as well:

Corollary 4.10. *If G is a graph, then for all $s \geq 1$,*

$$\text{reg}(I(W(G))^s) = 2s + \nu(W(G)) - 1.$$

Next, we study the regularity of powers of edge ideals of join of very well-covered graphs.

Definition 4.11. Let $G_1 = (V(G_1), E(G_1))$ and $G_2 = (V(G_2), E(G_2))$ be graphs with disjoint vertex sets. The join of G_1 and G_2 , denoted by $G_1 * G_2$, is the graph on the vertex set $V(G_1) \cup V(G_2)$ whose edge set is $E(G_1 * G_2) = E(G_1) \cup E(G_2) \cup \left\{ \{x, y\} \mid x \in V(G_1) \text{ and } y \in V(G_2) \right\}$.

It may be noted that for very well-covered graphs G_1, \dots, G_k , the product, $G_1 * \dots * G_k$ is not necessarily a very well-covered graph. However, we obtain the linear expression for $\text{reg}(I(G_1 * \dots * G_k)^s)$, for all $s \geq 1$.

Corollary 4.12. Let G_1, \dots, G_k be very well-covered graphs with $V(G_i) \cap V(G_j) = \emptyset$, for all $1 \leq i \neq j \leq k$. Then for all $s \geq 1$,

$$\text{reg}(I(G_1 * \dots * G_k)^s) = 2s + \max\{\nu(G_1), \dots, \nu(G_k)\} - 1$$

Proof. By [5, Theorem 4.5] and [26, Lemma 3.14], for all $s \geq 1$

$$2s + \max\{\nu(G_1), \dots, \nu(G_k)\} - 1 \leq \text{reg}(I(G_1 * \dots * G_k)^s).$$

Let $\mathcal{A} = \{G \mid \text{reg}((I(G)^{s+1} : e_1 \dots e_s)) \leq \text{reg}(I(G)), \text{ for any } s\text{-fold product } e_1 \dots e_s, s \geq 1\}$. By Theorem 4.7, it follows that $G_1, \dots, G_k \in \mathcal{A}$. Hence by [29, Theorem 4.4], we get $G_1 * \dots * G_k \in \mathcal{A}$. Therefore, it follows from [29, Theorem 4.5] and [26, Lemma 3.14], that for all $s \geq 1$, $\text{reg}(I(G_1 * \dots * G_k)^s) \leq 2s + \max\{\nu(G_1), \dots, \nu(G_k)\} - 1$. Hence the assertion follows. \square

It follows from Theorem 4.8 that if G is a very well-covered graph, then $\text{reg}(I(G)^s) = 2s + \nu(G) - 1$ for all $s \geq 2$. As a natural extension of this result, one tend to think that the same expression may hold true for well-covered graphs. This is not the case. For example, let $I = (x_1x_2, x_2x_3, x_3x_4, x_4x_5, x_5x_1, x_1x_6, x_6x_9, x_6x_7, x_7x_8)$ and G be the associated graph. It can be easily verified that G is a well-covered graph with $\nu(G) = 2$, but not a very well-covered graph. By [2, Corollary 3.8 and Theorem 5.3], for $s \geq 1$, $2s + \nu(G) - 1 < \text{reg}(I^s) = 2s + \nu(G) = 2s + 2$. Also, there exist well-covered graphs G such that $\text{reg}(I(G)^s) = 2s + \nu(G) - 1$ for all $s \geq 2$. For example, if $G = C_5$, then by [5, Theorem 5.2], $\text{reg}(I(G)^s) < 2s + \nu(G) - 1$ for all $s \geq 2$. Beyarslan et al. raised the question for which classes of graphs the equality $\text{reg}(I(G)^s) = 2s + \nu(G) - 1$ holds for $s \gg 0$, [5, Question 5.4]. This seems to be a rather tough question to answer. Therefore, we would like to ask:

Question 4.13. Characterize well-covered graphs G for which $\text{reg}(I(G)^s) = 2s + \nu(G) - 1$ for all $s \gg 0$?

Acknowledgement: The computational commutative algebra package Macaulay 2 [15] was heavily used to compute several examples. We also would like to thank Selvi Beyarslan for going through the manuscript and making some useful suggestions. The second author would like to thank the National Board for Higher Mathematics, India for the financial support. We also thank the referee for carefully reading the manuscript and making several suggestions that improved the exposition.

REFERENCES

- [1] A. Alilooee and A. Banerjee. Powers of edge ideals of regularity three bipartite graphs. *J. Commut. Algebra*, 9(4):441–454, 2017.
- [2] A. Alilooee, S. Beyarslan, and S. Selvaraja. Regularity of powers of edge ideals of unicyclic graphs. *ArXiv e-prints*, Feb. 2017.
- [3] A. Banerjee. The regularity of powers of edge ideals. *J. Algebraic Combin.*, 41(2):303–321, 2015.
- [4] D. Berlekamp. Regularity defect stabilization of powers of an ideal. *Math. Res. Lett.*, 19(1):109–119, 2012.
- [5] S. Beyarslan, H. T. Hà, and T. N. Trung. Regularity of powers of forests and cycles. *J. Algebraic Combin.*, 42(4):1077–1095, 2015.
- [6] M. Chardin. Powers of ideals and the cohomology of stalks and fibers of morphisms. *Algebra Number Theory*, 7(1):1–18, 2013.
- [7] A. Conca. Regularity jumps for powers of ideals. In *Commutative algebra*, volume 244 of *Lect. Notes Pure Appl. Math.*, pages 21–32. Chapman & Hall/CRC, Boca Raton, FL, 2006.
- [8] M. Crupi, G. Rinaldo, and N. Terai. Cohen-Macaulay edge ideal whose height is half of the number of vertices. *Nagoya Math. J.*, 201:117–131, 2011.
- [9] S. D. Cutkosky, J. Herzog, and N. V. Trung. Asymptotic behaviour of the Castelnuovo-Mumford regularity. *Compositio Math.*, 118(3):243–261, 1999.
- [10] D. Eisenbud and J. Harris. Powers of ideals and fibers of morphisms. *Math. Res. Lett.*, 17(2):267–273, 2010.
- [11] D. Eisenbud and B. Ulrich. Notes on regularity stabilization. *Proc. Amer. Math. Soc.*, 140(4):1221–1232, 2012.
- [12] S. Faridi. Cohen-Macaulay properties of square-free monomial ideals. *J. Combin. Theory Ser. A*, 109(2):299–329, 2005.
- [13] I. Gitler and C. E. Valencia. Bounds for invariants of edge-rings. *Comm. Algebra*, 33(5):1603–1616, 2005.
- [14] I. Gitler and C. E. Valencia. On bounds for some graph invariants. *ArXiv Mathematics e-prints*, Oct. 2005.
- [15] D. R. Grayson and M. E. Stillman. Macaulay2, a software system for research in algebraic geometry. Available at <http://www.math.uiuc.edu/Macaulay2/>.
- [16] H. T. Hà. Asymptotic linearity of regularity and a^* -invariant of powers of ideals. *Math. Res. Lett.*, 18(1):1–9, 2011.
- [17] H. T. Hà. Regularity of squarefree monomial ideals. In *Connections between algebra, combinatorics, and geometry*, volume 76 of *Springer Proc. Math. Stat.*, pages 251–276. Springer, New York, 2014.
- [18] J. Herzog and T. Hibi. *Monomial ideals*, volume 260 of *Graduate Texts in Mathematics*. Springer-Verlag London, Ltd., London, 2011.
- [19] J. Herzog, T. Hibi, and X. Zheng. Monomial ideals whose powers have a linear resolution. *Math. Scand.*, 95(1):23–32, 2004.
- [20] S. Jacques. *Betti numbers of graph ideals*. PhD thesis, University of Sheffield, 2004.
- [21] A. V. Jayanthan, N. Narayanan, and S. Selvaraja. Regularity of powers of bipartite graphs. *J. Algebraic Combin.*, 47(1):17–38, 2018.
- [22] K. Kimura, N. Terai, and S. Yassemi. The projective dimension of the edge ideal of a very well-covered graph. *Nagoya Math. J.*, (to appear).
- [23] V. Kodiyalam. Asymptotic behaviour of Castelnuovo-Mumford regularity. *Proc. Amer. Math. Soc.*, 128(2):407–411, 2000.
- [24] M. Mahmoudi, A. Mousivand, M. Crupi, G. Rinaldo, N. Terai, and S. Yassemi. Vertex decomposability and regularity of very well-covered graphs. *J. Pure Appl. Algebra*, 215(10):2473–2480, 2011.
- [25] M. Moghimian, S. A. S. Fakhari, and S. Yassemi. Regularity of powers of edge ideal of whiskered cycles. *Comm. Algebra*, 45(3):1246–1259, 2017.
- [26] A. Mousivand. Algebraic properties of product of graphs. *Comm. Algebra*, 40(11):4177–4194, 2012.
- [27] E. Nevo and I. Peeva. C_4 -free edge ideals. *J. Algebraic Combin.*, 37(2):243–248, 2013.
- [28] P. Norouzi, S. A. Seyed Fakhari, and S. Yassemi. Regularity of powers of edge ideal of very well-covered graphs. *ArXiv e-prints*, July 2017.

- [29] S. Selvaraja. Regularity of powers of edge ideals of product of graphs. *Journal of Algebra and Its Applications*, 17(6):1850128, 2018.
- [30] S. A. Seyed Fakhari. Symbolic powers of cover ideal of very well-covered and bipartite graphs. *Proc. Amer. Math. Soc.*, 146(1):97–110, 2018.

DEPARTMENT OF MATHEMATICS, INDIAN INSTITUTE OF TECHNOLOGY MADRAS, CHENNAI, INDIA
- 600036

E-mail address: `jayanav@iitm.ac.in`

DEPARTMENT OF MATHEMATICS, INDIAN INSTITUTE OF TECHNOLOGY MADRAS, CHENNAI, INDIA
- 600036

E-mail address: `selva.y2s@gmail.com`