# ON HIRANO INVERSES IN RINGS

HUANYIN CHEN AND MARJAN SHEIBANI

ABSTRACT. We introduce and study a new class of Drazin inverses. An element a in a ring has Hirano inverse b if  $a^2-ab \in N(R)$ , ab = ba and b = bab. Every Hirao inverse of an element is its Drazin inverse. We derive several characterizations for this generalized inverse. An element  $a \in R$  has Hirano inverse if and only if  $a^2$  has strongly Drazin inverse, if and only if  $a - a^3 \in N(R)$ . If  $\frac{1}{2} \in R$ , we prove that  $a \in R$  has Hirano inverse if and only if there exists  $p^3 = p \in comm^2(a)$  such that  $a - p \in N(R)$ , if and only if there exist two idempotents  $e, f \in comm^2(a)$  such that  $a + e - f \in N(R)$ . Clines formula and additive results for this generalized inverse are thereby obtained.

#### 1. INTRODUCTION

Let R be an associative ring with an identity. An element a in R is said to be Drazin invertible provided that there is a common solution to the equations  $a^k = a^{k+1}x, x = xax, ax = xa$  for some  $k \in \mathbb{N}$ . As is well known, an element  $a \in R$  has Drazin inverse if there exists  $b \in R$  such that

$$a - a^2 b \in N(R), ab = ba$$
 and  $b = bab$ .

The preceding b is unique, if such element exists. As usual, it will be denoted by  $a^{D}$ , and called the Drazin inverse of a.

The Drazin inverse is an important tool in ring theory and Banach algebra. It is very useful in matrix theory and in various applications in matrices (see [8, 10, 9]). The purpose of this paper is to introduce and study a new class of Drazin inverses. An

<sup>2010</sup> Mathematics Subject Classification. 15A09, 16E50, 15A23.

*Key words and phrases.* Drazin inverse, nilpotent; tripotent; Clines formula; matrix ring.

element  $a \in R$  has Hirano inverse if there exists  $b \in R$  such that

$$a^2 - ab \in N(R), ab = ba$$
 and  $b = bab$ .

The preceding b shall be unique, if such element exists, we denote it by  $a^H$ , and call the Hirano inverse of a. We observed these inverses form a class of Drazin inverses which is related to tripotents. Here,  $p \in R$  is a tripotent if  $p^3 = p$  (see [1]).

Following [8], An element  $a \in R$  has strongly Drazin inverse b if

$$a - ab \in N(R), ab = ba$$
 and  $b = bab$ .

In Section 2, We prove that every Hirano inverse of an element is its Drazin inverse. An element  $a \in R$  has Hirano inverse if and only if  $a^2$  has strongly Drazin inverse,

In Section 3 we characterize Hirano inverses via tripotents and nilpotents. It is shown that an element  $a \in R$  has Hirano inverse if and only if  $a - a^3 \in N(R)$ . If  $\frac{1}{2} \in R$ , we prove that  $a \in R$ has Hirano inverse if and only if there exists  $p^3 = p \in comm^2(a)$ such that  $a - p \in N(R)$ , if and only if there exist two idempotents  $e, f \in comm^2(a)$  such that  $a + e - f \in N(R)$ .

Let R be a ring, and let  $a, b, c \in R$  with aba = aca. In Section 4, we establish Clines formula for Hirano inverses. We prove that ac has Hirano inverse if and only if ba has Hirano inverse. In the last section, we are concern on generalized Jacobson Lemma. We prove that 1+ac has Hirano inverse if and only if 1+ba has Hirano inverse. Related multiplicative and additive results are also obtained.

Throughout, all rings are associative with an identity. We use N(R) to denote the set of all nilpotent elements in R. The double commutant of  $a \in R$  is defined by  $comm^2(a) = \{x \in R \mid xy = yx \text{ if } ay = ya \text{ for } y \in R\}$ .  $\mathbb{N}$  stands for the set of all natural numbers.

#### 2. Drazin inverse

The aim of this section is to investigate the relations among Drazin inverse, strongly Drazin inverse and Hirano inverse. We have **Theorem 2.1.** Let R be a ring and  $a \in R$ . If a has Hirano inverse, then a has Drazin inverse.

*Proof.* Suppose that a has Hirano inverse b. Then  $a^2 - ab \in N(R)$ , ab = ba and bab = b. Hence,  $a^2 - a^2b^2 = a^2 - a(bab) = a^2 - ab \in N(R)$ , and so

$$a^{2}(1 - a^{2}b^{2}) = (a^{2} - a^{2}b^{2})(1 - a^{2}b^{2}) \in N(R).$$

It follows that  $a(a-a^2b) = a^2(1-a^2b^2) \in N(R)$ ; hence,  $(a-a^2b)^2 = a(a-a^2b)(1-ab) \in N(R)$ . Thus,  $a-a^2b \in N(R)$ . Moreover, we see that ab = ba and bab = b. Therefore, a has Drazin inverse b, as desired.

**Corollary 2.2.** Let R be a ring and  $a \in R$ . Then a has at most one Hirano inverse in R, and if the Hirano inverse of a exists, it is exactly the Drazin inverse of a, i.e.,  $a^H = a^D$ .

*Proof.* Let a has Hirano inverse x. In view of Theorem 2.1, a has Drazin inverse. Furthermore, we see that  $a^D = a^H$ . By virtue of [8, Theorem 2.4], the Drazin inverse of a is unique. Thus, a has at most one Hirano inverse in R, as desired.

**Example 2.3.** Every  $2 \times 2$  matrix over  $\mathbb{Z}_2$  has Drazin inverse, but  $a = \begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix} \in M_2(\mathbb{Z}_2)$  has no Hirano inverse.

*Proof.* Since  $M_2(\mathbb{Z}_2)$  is a finite ring, every element in  $M_2(\mathbb{Z}_2)$  has Drazin inverse. If  $a = \begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix} \in M_2(\mathbb{Z}_2)$  has Hirano inverse, it follows by Corollary 2.2 that  $a^H = a^D = \begin{pmatrix} -1 & 1 \\ 1 & 0 \end{pmatrix}$ . But  $a^2 - (a - 1)$ 

 $aa^{H} = \begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix}$  is not nilpotent. This gives a contradiction, and we are through.

**Theorem 2.4.** Let R be a ring and  $a \in R$ . Then a has Hirano inverse if and only if  $a^2$  has strongly Drazin inverse. In this case,  $(a^2)^{sD} = (a^H)^2$  and  $a^H = a(a^2)^{sD}$ .

*Proof.*  $\implies$  Since a has Hirano inverse, we have  $b \in R$  such that  $a^2-ab \in N(R)$ , ab = ba and b = bab. Thus,  $a^2-a^2b^2 = a^2-a(bab) =$ 

 $a^2 - ab \in N(R), a^2b^2 = b^2a^2$  and  $b^2a^2b^2 = b^2$ . Therefore  $a^2 \in R$  has strongly Drazin inverse. Moreover,  $(a^2)^{sD} = (a^H)^2$ .

 $\Leftarrow$  Since  $a^2$  has strongly Drazin inverse, we have  $x \in R$  such that  $a^2 - a^2 x \in N(R)$ , ax = xa and  $x = xa^2 x$ . Set b = ax. Then  $a^2 - ab = a^2 - a^2 x \in N(R), ab = ba$  and bab = (ax)a(ax) = $a(xa^2x) = ax = b$ . Hence, a has Hirano inverse b. Furthermore,  $a^H = a(a^2)^{sD}.$ 

**Example 2.5.** Every element in  $\mathbb{Z}_3$  has Hirano inverse, but  $2 \in \mathbb{Z}_3$ has no strongly Drazin inverse.

*Proof.* One directly checks that every element in  $\mathbb{Z}_3$  has Hirano inverse. If  $2 \in \mathbb{Z}_3$  has strongly Drazin inverse, then  $2 = 2(2^D)$ and  $2^{D} = 2(2^{D})^{2}$ , whence  $-1 = -2^{D}$ , and so  $2^{D} = 1$ . This gives a contradiction. This shows that  $2 \in \mathbb{Z}_3$  has no strongly Drazin inverse. 

**Example 2.6.** Let  $a = \begin{pmatrix} -2 & 3 & 2 \\ -2 & 3 & 2 \\ 1 & -1 & -1 \end{pmatrix} \in M_3(\mathbb{Z})$ . Then a has Hirano inverse, but it has no strongly Drazin inverse.

*Proof.* Clearly,  $a = a^3$ , and so  $a^2$  has strongly Drazin inverse  $a^2$ , thus a has Hirano inverse, by Theorem 2.1. On the other hand, the characteristic polynomial  $\chi(a - a^2) = t^2(t + 2) \not\equiv t^3 \pmod{N(R)}$ . In light of [3, Proposition 4.2],  $a - a^2$  is not nilpotent. Therefore A does not have strongly Drazin inverse by [8, Lemma 2.2].  $\square$ 

Thus, we have the relations of various type of inverses in a ring by the following:

 $\{\text{strongly Drazin inverses}\} \subseteq \{\text{Hirano inverses}\} \subseteq \{\text{Drazin inverses}\}.$ 

**Example 2.7.** Let  $\mathbb{C}$  be the field of complex numbers.

(1)  $2 \in \mathbb{C}$  is invertible, but 2 has no Hirano inverse in  $\mathbb{C}$ .

(2)  $A \in M_n(\mathbb{C})$  has Hirano inverse if and only if there exists a decomposition  $\mathbb{C}^n \oplus \mathbb{C}^n = U \oplus V$  such that U, V are A-invariant and  $A^2 \mid_U, (I_n - A^2) \mid_V$  are both nilpotent if and only if eigenvalues of  $A^2$  are only 0 or 1.

*Proof.* (1) Straightforward.

(2) In view of Theorem 2.4,  $A \in M_n(\mathbb{C})$  has Hirano inverse if and only if  $A^2 \in M_n(\mathbb{C})$  has strongly Drazin inverse if and only if  $A^2$  is the sum of an idempotent and an nilpotent matrices that commute. Therefore we are done by [8, Lemma 2.2].

## 3. CHARACTERIZATIONS

The purpose of this section is to characterize Hirano inverses by means of tripotents and nilpotents in a ring. The following result is crucial.

**Theorem 3.1.** Let R be a ring and  $a \in R$ . Then a has Hirano inverse if and only if  $a - a^3 \in N(R)$ .

*Proof.*  $\implies$  Since a has Hirano inverse b, we have

$$w := a^2 - ab \in N(R), ab = ba$$
 and  $b = bab$ .

Hence,  $a^2 = ab + w$  and (ab)w = w(ab). Thus,  $a^2 - a^4 = (ab + w) - (ab + w)^2 = (w - 2ab - w)w \in N(R)$ . Hence,  $a(a - a^3) \in N(R)$ , and so  $(a - a^3)^2 = a(a - a^3)(1 - a^2) \in N(R)$ . Accordingly,  $a - a^3 \in N(R)$ .  $\iff$  Suppose that  $a - a^3 \in N(R)$ . Then  $a^2 - a^4 = a(a - a^3) \in N(R)$ .

Suppose that  $a - a^{\circ} \in N(R)$ . Then  $a^{\circ} - a^{\circ} \equiv a(a - a^{\circ}) \in N(R)$ . In view of [2, Lemma 2.1], there exists an idempotent  $e \in \mathbb{Z}[a^2]$  such that  $w := a^2 - e \in N(R)$  and ae = ea. Set  $c = (1+w)^{-1}e$ . Then ac = ca and

$$\begin{aligned} a^2 c &= a^2 (1+w)^{-1} e \\ &= (1+w)^{-1} e (a^2+1-e) \\ &= (1+w)^{-1} e (1+w) \\ &= e. \end{aligned}$$

Hence,  $a^2 - a^2c = a^2 - e = w \in N(R)$ . Moreover,  $ca^2c = ce = c$ . Set b = ac. Then  $a^2 - ab \in N(R)$ , ab = ba and  $bab = (ac)a(ac) = (ca^2c)a = ca = b$ . Therefore a has Hirano inverse b, as required.  $\Box$ 

**Corollary 3.2.** Let R be a ring and  $a \in R$ . If a has Hirano inverse, then  $a^H$  has Hirano inverse, and that  $(a^H)^H = a^2 a^H$ .

*Proof.* Write  $b = a^H$ . Then  $w := a^2 - ab \in N(R)$ , ab = ba and b = bab. In view of Theorem 3.1,  $a - a^3 \in N(R)$ . One easily checks

that

$$\begin{array}{rcl} b-b^3 &=& b^2a-b^2(bab)\\ &=& b^2a-b^3(a^2-w)=b^2a-b(b^2a)a-b^3w\\ &=& b^2a-b^2(ba)a-b^3w\\ &=& b^2a-b(b^2a)a-b^3w\\ &=& b^2a-b^2(a^2-w)a-b^3w=b^2(a-a^3)+(b-b^3)w\\ &\in& N(R). \end{array}$$

By using Theorem 3.1 again, b has Hirano inverse. In light of Corollary 2.2,  $b = a^D$  and  $b^H = b^D$ . Therefore  $(a^H)^H = (a^D)^D = a^2 a^D = a^2 a^H$ , in terms of [5, Theorem 5.4]. This completes the proof.

We are now ready to prove the following.

**Theorem 3.3.** Let R be a ring with  $\frac{1}{2} \in R$ , and let  $a \in R$ . Then the following are equivalent:

- (1)  $a \in R$  has Hirano inverse.
- (2) There exists  $p^3 = p \in comm^2(a)$  such that  $a p \in N(R)$ .
- (3) There exist two idempotents  $e, f \in comm^2(a)$  such that  $a + e f \in N(R)$ .

*Proof.* (1)  $\Rightarrow$  (3) In view of Theorem 3.1,  $a - a^3 \in N(R)$ . Set  $b = \frac{a^3 + a}{2}$  and  $c = \frac{a^3 - a}{2}$ . Then

$$b^{2} - b = \frac{1}{4}(a^{4} + 2a^{3} - a^{2} - 2a) = \frac{1}{4}(a + 2)(a^{3} - a),$$
  

$$c^{2} - c = \frac{1}{4}(a^{4} - 2a^{3} - a^{2} + 2a) = \frac{1}{4}(a - 2)(a^{3} - a).$$

Thus,  $b - b^2$ ,  $c - c^2 \in N(R)$ . In view of [2, Lemma 2.1], there exist two idempotents  $e \in \mathbb{Z}[b]$  and  $f \in \mathbb{Z}[c]$  such that  $b - e, c - f \in N(R)$ . One easily checks that a = b - c and  $b, c \in \mathbb{Z}[a]$ . Therefore  $a - f + e = (b - f) - (c - e) \in N(R)$ . We see that  $e, f \in \mathbb{Z}[a] \subseteq comm^2(a)$ , as desired.

 $(3) \Rightarrow (2)$  Set p = f - e. Then  $a - p \in N(R)$  and  $p^3 = (e - f)^3 = e - f = p$ , as ef = fe.

(2)  $\Rightarrow$  (1) By hypothesis, there exists  $p^3 = p \in comm^2(a)$  such that  $w := a - p \in N(R)$ . Thus,  $a - a^3 = (p + w) - (p + w)^3 = w(1 - 3pw - 3p^2 - w^2) \in N(R)$ . This completes the proof, by Theorem 3.1.

6

**Corollary 3.4.** Let R be a ring with  $\frac{1}{2} \in R$ . Then  $a \in R$  has Hirano inverse if and only if there exist commuting  $b, c \in R$  such that a = b - c, where b, c have strongly Drazin inverses.

*Proof.*  $\implies$  In light of Theorem 3.3, there exist two idempotents  $e, f \in comm^2(a)$  such that  $w := a + e - f \in N(R)$ . Hence, a = f - (e + w) with  $w \in \mathbb{Z}[a]$ . Clearly,  $e \in R$  has strongly Drazin inverse. Furthermore, we see that  $(e + w) - (e + w)e \in N(R), e(e + w) = (e + w)e$  and e = e(e + w)e. Thus,  $e + w \in R$  has strongly Drazin inverse. Set b = f and c = e + w. Then a = b - c and bc = ca, as desired.

As 2 is invertible in any Banach algebra, we now derive

**Corollary 3.5.** Let A be a Banach algebra and let  $a \in A$ . Then the following are equivalent:

- (1) a has Hirano inverse.
- (2) There exists  $p^3 = p \in comm^2(a)$  such that  $a p \in N(A)$ .
- (3) There exist two idempotents  $e, f \in comm^2(a)$  such that  $a + e f \in N(A)$ .
- (4) There exist commuting  $b, c \in R$  such that a = b c, where b, c have strongly Drazin inverses.

Recall that a ring R is strongly 2-nil-clean if every element in R is the sum of a tripotent and a nilpotent that commute (see [2]). We characterize strongly 2-nil-clean rings by using Hirano inverses.

**Proposition 3.6.** A ring R is strongly 2-nil-clean if and only if every element in R has Hirano inverse.

*Proof.*  $\implies$  Let  $a \in R$ . In view of [2, Theorem 2.3],  $a - a^3 \in N(R)$ . Thus, a has Hirano inverse by Theorem 3.1.

 $\leftarrow$  Let  $a \in R$ . In light of Theorem 3.1,  $a-a^3 \in N(R)$ . Therefore we complete the proof, by [2, Theorem 2.3].

**Corollary 3.7.** A Banach algebra A is strongly 2-nil-clean if and only if every element in A is the difference of two commuting elements that have strongly Drazin inverses.

*Proof.* Combining Proposition 3.6 and Corollary 3.5, we obtain the result.  $\Box$ 

# 4. CLINES FORMULA

Let  $a, b \in R$ . Then ab has Drazin inverse if and only if ba has Drazin inverse. In this case,  $(ba)^D = b((ab)^D)^2 a$ . This was known as Clines formula for Drazin inverses. It plays an important role in revealing the relationship between the Drazin inverse of a sum of two elements and the Drazin inverse of a block triangular matrix. In [14] extended this result and proved that if aba = aca then 1+achas Drazin inverse if and only if ba has Drazin inverse. The goal of this section is to generalize Clines formula from Drazin inverses to Hirano inverses. We derive

**Theorem 4.1.** Let R be a ring, and let  $a, b, c \in R$ . If aba = aca, Then ac has Hirano inverse if and only if ba has Hirano inverse. In this case,  $(ba)^H = b((ac)^H)^2 a$ .

*Proof.*  $\implies$  By induction on *n* we shall prove

$$(ba - (ba)^3)^{n+1} = b(ac - (ac)^3)^n(a - ababa).$$

For n = 1,

 $((ba - (ba)^3))^2$ = (ba - bababa)(ba - bababa) = b(a - ababa)ba(1 - baba) = b(aba - abababa)(1 - baba) = b(aca - acacaca)(1 - baba) = b(ac - acacaca)(a - ababa) = b(ac - (ac)^3)(a - ababa).

Suppose that the assertion is true for any  $n \leq k(k \geq 2)$ . Then,

8

$$(ba - (ba)^3)^{k+1} = ((ba - (ba)^3))^k (ba - (ba)^3)$$
  
=  $b(ac - (ac)^3)^{k-1} (a - ababa)(ba - bababa)$   
=  $b(ac - (ac)^3)^{k-1} (aba - abababa)(1 - baba)$   
=  $b(ac - (ac)^3)^{k-1} (aca - acacaca)(1 - baba)$   
=  $b(ac - (ac)^3)^k (a - ababa).$ 

As ac has Hirano inverse, then by Theorem 3.1,  $(ac-(ac)^3)^n = 0$  for some  $n \in \mathbb{N}$ , which implies that  $ba-(ba)^3 \in N(R)$ ) by the preceding discussion. In light of Theorem 3.1, ba has Hirano inverse.

Now let  $e = (ac)^{H}$ , by Corollary 2.2, e is Drazin inverse of ac. As we proved, ba has Hirano inverse, and its Hirano inverse is its drazin inverse, then by [14, Theorem 2.7],  $(ba)^{D} = (b(ac)^{2})^{D}a$  and so  $(ba)^{H} = b(b(ac)^{2})^{H}a$ .

 $\Leftarrow$  This is symmetric.

**Corollary 4.2.** Let R be a ring and  $a, b \in R$ . If ab has Hirano inverse, then so has ba, and  $(ba)^H = b((ab)^H)^2 a$ .

*Proof.* It follows directly from Theorem 4.1, as aba = aba.

**Theorem 4.3.** Let R be a ring, and let  $a, b \in R$ . If  $(ab)^k$  has Hirano inverse, then so is  $(ba)^k$ .

*Proof.*  $\Longrightarrow$  Let k = 1, then by Corollary 4.2, ab has Hirano inverse if and only if ba has Hirano inverse. Now let  $k \ge 2$ , as  $(ab)^k = a(b(ab)^{k-1})$ , then by Corollary 4.2,  $(b(ab)^{k-1})a$  has Hirano inverse, so  $(ba)^k$  has Hirano inverse.

**Theorem 4.4.** Let R be a ring, and let  $a, b \in R$ . If a, b have Hirano inverses and ab = ba. Then ab has Hirano inverse and

$$(ab)^H = a^H b^H.$$

*Proof.* Since a, b have Hirano inverses, by Theorem 3.1, we have  $a - a^3, b - b^3 \in N(R)$ . Note that

$$ab - (ab)^3 = (a - a^3)(b - b^3) + ab^3 + a^3b - a^3b^3 - a^3b^3 = (a - a^3)(b - b^3) + a^3(b - b^3) + b^3(a - a^3) \in N(R),$$

as ab = ba. By using Theorem 3.1 again, ab has Hirano inverse. By Theorem 2.1, ab has Drazin inverse and by [13, Lemma 2],  $(ab)^D = b^D a^D$ . According to Corollary 2.2,  $(ab)^H = (ab)^D = b^D a^D = b^H a^H$ , as asserted.

**Corollary 4.5.** Let R be a ring. If  $a \in R$  has Hirano inverse, then  $a^n$  has Hirano inverse and  $(a^n)^H = (a^H)^n$  for all  $n \in \mathbb{N}$ .

*Proof.* By Theorem 4.4 and induction, we easily obtain the result.  $\Box$ 

We note that the converse of the preceding corollary is not true, as the following shows.

**Example 4.6.** Let  $\mathbb{Z}_5 = \mathbb{Z}/5\mathbb{Z}$  be the ring of integers modulo 5. Then  $-2 \in \mathbb{Z}_5$  has no Hirano inverse, as  $(-2) - (-2)^3 = 1 \in \mathbb{Z}_5$  is not nilpotent, but  $(-2)^2 = -1 \in \mathbb{Z}$  has Hirano inverse, as  $(-1) - (-1)^3 \in \mathbb{Z}_5$  is nilpotent.

### 5. Additive properties

Let  $a, b \in R$ . Jacobson Lemma asserted that  $1 + ab \in R$  has Drazin inverse if and only if  $1 + ba \in R$  has Drazin inverse. More recently, Yan and Fang studied the link between basic operator properties of 1 + ac and 1 + ba when aba = aca (see [11]). Though we do not know whether the Drazin invertibility of 1 + ac and 1 + bacoincide under aba = aca, we could generalize Jacobson Lemma for Hirano inverses.

**Theorem 5.1.** Let R be a ring, and let  $a, b, c \in R$ . If aba = aca, Then 1 + ac has Hirano inverse if and only if 1 + ba has Hirano inverse.

*Proof.*  $\implies$  First by induction on *n* we prove that

$$((1+ba)-(1+ba)^3)^{n+1} = b((1+ac)-(1+ac)^3)^n(-2a-3aba-ababa).$$

For n = 1,

 $\begin{array}{l} ((1+ba)-(1+ba)^3)^2 \\ = ((1+ba)-(1+ba)^3)((1+ba)-(1+ba)^3) \\ = (-2ba-3baba-bababa)(1+ba-1-3ba-3baba-bababa) \\ = (-2ba-3baba-bababa)ba(-2-3ba-baba) \\ = (-2baba-3bababa-babababa)(-2-3ba-baba) \\ = b(-2aba-3ababa-abababa)(-2-3ba-baba) \\ = b(-2ac-3acac-acacac)(-2a-3aba-ababa) \\ = b((1+ac)-(1+ac)^3)(-2a-3aba-ababa). \end{array}$ 

Let  $k \geq 2$  and the assertion works for any  $n \leq k$ . Then

 $\begin{array}{l} ((1+ba)-(1+ba)^3)^{k+1} \\ = ((1+ba)-(1+ba)^3)^k((1+ba)-(1+ba)^3) \\ = b((1+ac)-(1+ac)^3)^{k-1}(-2a-3ba-baba)((1+ba)-(1+ba)^3) \\ = b((1+ac)-(1+ac)^3)^{k-1}(-2a-3ba-baba)(ba-3ba-3baba-(ba)^3) \\ = b((1+ac)-(1+ac)^3)^{k-1}(-2aba-3baba-bababa)(-2-3ba-baba) \\ = b((1+ac)-(1+ac)^3)^k(-2a-3aba-ababa). \end{array}$ 

As 1 + ac has Hirano inverse, then by Theorem 3.1,  $(1 + ac) - (1 + ac)^3 \in N(R)$ . By the above equality,  $(1 + ba) - (1 + ba)^3 \in N(R)$ . In light of Theorem 3.1, (1 + ba) has Hirano inverse.

 $\Leftarrow$  This is symmetric.

**Corollary 5.2.** Let R be a ring, and let  $a, b \in R$ . Then 1 + ab has Hirano inverse if and only if 1 + ba has Hirano inverse.

*Proof.* The proof follows directly from the equality aba = aba and Theorem 5.1.

As is well known,  $a \in R$  has strongly Drazin inverse if and only if so does  $1 - a \in R$  (see [8, Lemma 3.3]). But Hirano inveses in a ring are not the case.

**Example 5.3.** Let  $a = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \in M_2(\mathbb{Z}_3)$ . Then a has Hirano inverse, but 1 - a does not have.

Proof. It is easy to verify that  $a - a^3 = \begin{pmatrix} 1 & 1 \\ -1 & 2 \end{pmatrix} \in N(M_2(\mathbb{Z}_4)),$ but  $(1-a) - (1-a)^3 = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \notin N(M_2(\mathbb{Z}_4)).$  Therefore we are done by Theorem 3.1. **Proposition 5.4.** Let R be a ring and  $a, b \in R$ . If a, b have Hirano inverses and ab = ba = 0, then a + b has Hirano inverse and

$$(a+b)^H = a^H + b^H$$

*Proof.* Clearly,  $ab^H = ab(b^H)^2 = 0$  and  $ba^H = ba(a^H)^2 = 0$ . Likewise,  $a^Hb = b^Ha = 0$ . Thus, we see that  $a, a^H, b, b^H$  commute. Thus,  $(a^2 - aa^H)(b^2 - bb^H) = (b^2 - bb^H)(a^2 - aa^H)$ . Also we have

$$(a+b)(a^{H}+b^{H})^{2} = (aa^{H}+bb^{H})(a^{H}+b^{H}) = a^{H}+b^{H}$$

and

$$(a+b)^2 - (a+b)(a^H + b^H) = (a^2 - aa^H) + (b^2 - bb^H) \in N(R).$$
  
Therefore  $(a+b)^H = a^H + b^H$ , as asserted,.

It is convenient at this stage to include the following.

**Theorem 5.5.** Let R be a ring, and let  $a, b \in R$ . If ab has strongly inverse and  $a^2 = b^2 = 0$ , then a + b has Hirano inverse and  $(a + b)^H = a(ba)^H + b(ab)^H$ .

*Proof.* Suppose that ab has strongly inverse and  $a^2 = b^2 = 0$ . Then ba has strongly inverse by [8, Theorem 3.1]. It is obvious that ab and ba have Drazin inverses. Let  $x = a(ba)^D + b(ab)(ab)^D$ . Then we easily check that (a + b)x = x(a + b) and  $(a + b)x^2 = x$ . We easily check that

$$(a+b)^2 - (a+b)x = ab + ba - (ab)(ab)^D - (ba)(ba)^D = (ab - (ab)(ab)^D) + (ba - (ba)(ba)^D) \in N(R),$$

as  $(ab-(ab)(ab)^D)(ba-(ba)(ba)^D) = (ba-(ba)(ba)^D)(ab-(ab)(ab)^D)$ = 0. Therefore  $a + b \in R$  has Hirano inverse, and that  $(a + b)^H = a(ba)^D + b(ab)^D = a(ba)^H + b(ab)^H$ .

**Example 5.6.** Let  $a = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$ ,  $b = \begin{pmatrix} 0 & 0 \\ 2 & 0 \end{pmatrix} \in M_2(\mathbb{Z}_3)$ . Then  $ab = \begin{pmatrix} 2 & 0 \\ 0 & 0 \end{pmatrix} \in M_2(\mathbb{Z})$  has Hirano inverse,  $a^2 = b^2 = 0$ . But  $a + b = \begin{pmatrix} 0 & 1 \\ 2 & 0 \end{pmatrix} \in M_2(\mathbb{Z}_3)$  has no Hirano inverse. **Example 5.7.** Let  $a = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$ ,  $b = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix} \in M_2(\mathbb{Z})$ . Then  $ab = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \in M_2(\mathbb{Z})$  has strongly Drazin inverse,  $a^2 = b^2 = 0$ . But  $a + b = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$  has no strongly Drazin inverse. In this case,  $(a + b)^H = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$ .

### References

- M.S. Abdolyousefi and H. Chen, Rings in which elements are sums of tripotents and nilpotents, J. Algebra App., https://doi.org/10.1142/S02194988 18500421.
- H. Chen and M. Sheibani, Strongly 2-nil-clean rings, J. Algebra Appl., 16(2017) 1750178 (12 pages), DOI: 10.1142/S021949881750178X.
- [3] A.J. Diesl, Nil clean rings, J. Algebra, **383**(2013), 197–211.
- [4] Y. Hirano and H. Tominaga, Rings in which every element is a sum of two idempotents, Bull. Austral. Math. Soc., 37(1988), 161–164.
- [5] J. J. Koliha, A generalized Drazin inverse, Glasgow. Math. J., 38(1996), 367–381.
- [6] T.Y. Lam and P. Nielsen, Jacobson's lemma for Drazin inverses, Ring theory and its applications, 185–195, Contemp. Math. 609, Amer. Math. Soc., Providence, RI, 2014.
- [7] P. Patricio and R.E. Hartwig, Some additive results on Drazin inverses, Appl. Math. Comput., 215(2009), 530–538.
- [8] Z. Wang, A class of Drazin inverses in rings, *Filomat*, **31**(2017), 1781-C1789.
- [9] Y.M. Wei and C.Y. Deng, A note on additive results for the Drazin inverse, Linear Multilinear Algebra, 59(2011), 1319–1329.
- [10] H. Yang; X.F. Liu, The Drazin inverse of the sum of two matrices and its applications, J. Comput. Appl. Math., 235(2011), 1412–1417.
- [11] K. Yang; X.C. Fang, Common properties of the operator products in local spectral theorey, Acta Math. Sinica, English Series, 31(2015), 1715–1724.
- [12] Y. Zhou, Rings in which elements are sum of nilpotents, idempotents and tripotents, J. Algebra App., 16(2017), DOI: 10.1142/S0219498818500093.

- [13] G.F. Zhuang; J.L. Chen; D.S. Cvetkovic-Ilic and Y.M. Wei, Additive property of Drazin invertibility of elements in a ring, *Linear Multilinear Algebra*, **60**(2012), 903–910.
- [14] QP. Zeng and HJ. Zhong, New results on common properties of the products AC and BA, J. Math. Annal. Appl., 427(2015), 830–840.

Department of Mathematics, Hangzhou Normal University, Hang -zhou, China

*E-mail address*: <huanyinchen@aliyun.com>

WOMEN'S UNIVERSITY OF SEMNAN (FARZANEGAN), SEMNAN, IRAN *E-mail address*: <sheibani@fgusem.ac.ir>