

***CP* violation in charmed hadron decays into neutral kaons**

Di Wang

School of Nuclear Science and Technology, Lanzhou University, Lanzhou 730000, People's Republic of China

E-mail: dwang15@lzu.edu.cn

Fu-Sheng Yu

School of Nuclear Science and Technology, Lanzhou University, Lanzhou 730000, People's Republic of China

E-mail: yufsh@lzu.edu.cn

Hsiang-nan Li

Institute of Physics, Academia Sinica, Taipei, Taiwan 115, Republic of China

E-mail: hnli@phys.sinica.edu.tw

We find a new *CP* violating effect in charmed hadron decays into neutral kaons, which is induced by the interference between the Cabibbo-favored and doubly Cabibbo-suppressed amplitudes with the $K^0 - \bar{K}^0$ mixing [1]. It is estimated to be of order of $\mathcal{O}(10^{-3})$, much larger than the direct *CP* asymmetry, but missed in the literature. To reveal this new *CP* violation effect, we propose a new observable, the difference of the *CP* asymmetries in the $D^+ \rightarrow \pi^+ K_S^0$ and $D_s^+ \rightarrow K^+ K_S^0$ modes. Once the new effect is determined by experiments, the direct *CP* asymmetry then can be extracted and used to search for new physics.

The 15th International Conference on Flavor Physics & CP Violation
5-9 June 2017
Prague, Czech Republic

CP asymmetry plays a unique role in understanding the matter-antimatter asymmetry and searching for new physics. It has been well established in the kaon and B meson systems [2, 3, 4, 5], but not yet in the charm sector. In the past decade, many efforts have been devoted to study the CP violation in the singly Cabibbo-suppressed (SCS) D meson decays. The most precise experimental result of CP violation in SCS decays is [6]

$$\Delta A_{CP} \equiv A_{CP}(D^0 \rightarrow K^+ K^-) - A_{CP}(D^0 \rightarrow \pi^+ \pi^-) = (-0.10 \pm 0.08 \pm 0.03)\%. \quad (1)$$

With the precision lower than 10^{-3} , the CP violation in charm decays has not been observed.

CP asymmetry can also occur in $D \rightarrow f K_S^0$ decays, where f is a final-state particle. For example, the CP violation in $D^+ \rightarrow \pi^+ K_S^0$ has been measured by Belle collaboration with 3.2σ from zero [7]. In this work, We point out a new CP -violation effect, which is induced by the interference between the Cabibbo-favored (CF) and doubly Cabibbo-suppressed (DCS) amplitudes with the mixing of final-state mesons [1]. It is estimated to be of order of 10^{-3} , much larger than the direct CP asymmetry, however, missed in the literature [7, 8, 9, 10]. We propose a new observable, the difference between the CP asymmetries in the $D^+ \rightarrow \pi^+ K_S^0$ and $D_s^+ \rightarrow K^+ K_S^0$ decays, to measure the new CP violation effect. Once the new effect is obtained, the direct CP asymmetry in charm decays can be extracted correctly and used to search for new physics.

In experiments, the K_S^0 state is reconstructed by $\pi^+ \pi^-$ final state. The time-dependent CP violation in D meson decays into neutral kaons is defined by

$$A_{CP}(t) \equiv \frac{\Gamma(D \rightarrow K(t)(\rightarrow \pi^+ \pi^-)f) - \Gamma(\bar{D} \rightarrow K(t)(\rightarrow \pi^+ \pi^-)\bar{f})}{\Gamma(D \rightarrow K(t)(\rightarrow \pi^+ \pi^-)f) + \Gamma(\bar{D} \rightarrow K(t)(\rightarrow \pi^+ \pi^-)\bar{f})}, \quad (2)$$

where $K(t)$ donates the immediate state of neutral kaons. The mass eigenstates of neutral kaons, $|K_S^0\rangle$ of mass m_S and width Γ_S and $|K_L^0\rangle$ of mass m_L and width Γ_L , are linear combinations of the flavor eigenstates $|K^0\rangle$ and $|\bar{K}^0\rangle$, $|K_{S,L}^0\rangle = p_K |K^0\rangle \mp q_K |\bar{K}^0\rangle$, with $p_K = (1 + \varepsilon)/\sqrt{2(1 + |\varepsilon|^2)}$ and $q_K = (1 - \varepsilon)/\sqrt{2(1 + |\varepsilon|^2)}$, and ε is a small parameter characterizing the indirect CP violation in neutral kaon mixing [5]. For convenience, the ratio between DCS and CF amplitudes is set as

$$\mathcal{A}(D \rightarrow K^0 f) / \mathcal{A}(D \rightarrow \bar{K}^0 f) = r_f e^{i(\phi + \delta_f)}, \quad (3)$$

where r_f is the size of the ratio, ϕ and δ_f are relative weak and strong phases respectively. In the SM, $r_f \sim |V_{cd}^* V_{us} / V_{cs}^* V_{ud}| \sim \mathcal{O}(10^{-2})$ and $\phi \equiv \text{Arg}[-V_{cd}^* V_{us} / V_{cs}^* V_{ud}] = (-6.17 \pm 0.43) \times 10^{-4}$ [5]. With the small parameters ε , r_f and ϕ , we obtain the time-dependent CP violation as

$$A_{CP}(t) \simeq [A_{CP}^{\bar{K}^0}(t) + A_{CP}^{\text{dir}}(t) + A_{CP}^{\text{int}}(t)] / D(t), \quad (4)$$

in which $D(t) = e^{-\Gamma_S t} (1 - 2r_f \cos \delta_f \cos \phi) + e^{-\Gamma_L t} |\varepsilon|^2$. The $A_{CP}^{\bar{K}^0}(t)$ term is the indirect CP violation in $K^0 - \bar{K}^0$ mixing,

$$A_{CP}^{\bar{K}^0}(t) = -2e^{-\Gamma t} (\text{Re}(\varepsilon) \cos(\Delta m t) + \text{Im}(\varepsilon) \sin(\Delta m t)) + 2\text{Re}(\varepsilon) e^{-\Gamma_S t}. \quad (5)$$

The $A_{CP}^{\text{dir}}(t)$ term is the direct CP violation in charm decay induced by the interference between the CF and DCS amplitudes,

$$A_{CP}^{\text{dir}}(t) = 2e^{-\Gamma_S t} r_f \sin \delta_f \sin \phi. \quad (6)$$




Figure 1: The plot of time-dependent CP asymmetries in the $D^+ \rightarrow \pi^+ K(t) (\rightarrow \pi^+ \pi^-)$ given by [1], where the right figure is the zoomed plot for the small t region of the left one, and the gray bands are the theoretical uncertainties.

The $A_{CP}^{\text{int}}(t)$ term is the interference effect between the CF and DCS amplitudes with $K^0 - \bar{K}^0$ mixing,

$$A_{CP}^{\text{int}}(t) = -4r_f \cos \phi \sin \delta_f (Im(\epsilon) e^{-\Gamma_S t} - e^{-\Gamma t} (Im(\epsilon) \cos(\Delta m t) - Re(\epsilon) \sin(\Delta m t))). \quad (7)$$

The parameters r_f and δ_f for the $D^+ \rightarrow \pi^+ K_S^0$ and $D_s^+ \rightarrow K^+ K_S^0$ decays have been estimated in the factorization-assisted topological-amplitude (FAT) approach [11, 12]. The dependences of the CP asymmetry in the $D^+ \rightarrow \pi^+ K(t) (\rightarrow \pi^+ \pi^-)$ decay on t/τ_S are displayed in Fig. 1. It is found that the total CP violation dominated by $A_{CP}^{\bar{K}^0}(t)$, and the deviation from $A_{CP}^{\bar{K}^0}(t)$ mainly comes from $A_{CP}^{\text{int}}(t)$. The direct CP asymmetries are too small to be seen in Fig. 1, being of order of $\mathcal{O}(10^{-5})$. According to Eqs. (5) and (7), $A_{CP}^{\bar{K}^0}(t=0) = A_{CP}^{\text{int}}(t=0) = 0$, resulting in $A_{CP}(t=0) = A_{CP}^{\text{dir}}(t=0)$. Both the forthcoming experiments cannot measure the direct CP asymmetries, unless the large weak phase differences are provided by new physics. Thereby, an observation with nonvanishing $A_{CP}(t=0)$ indicates new physics. Compared to the SCS processes, in which the CP asymmetry cannot discriminate new physics due to the ambiguities in estimating the penguin amplitudes, the direct CP asymmetry in neutral kaon modes would give a more unambiguous new physics signal.

The time-integrated CP asymmetry is

$$A_{CP} = \frac{\int_0^\infty F(t) [A_{CP}^{\bar{K}^0}(t) + A_{CP}^{\text{dir}}(t) + A_{CP}^{\text{int}}(t)] dt}{\int_0^\infty F(t) D(t) dt}, \quad (8)$$

where $F(t)$ is a function to take into account relevant experimental effects. With the approximation of $F(t) = 1$ in the interval $[t_1, t_2]$ and $F(t) = 0$ elsewhere [9], equation (8) yields

$$A_{CP}(t_1, t_2) = \frac{2Re(\epsilon) - 4Im(\epsilon)r_f \cos \phi \sin \delta_f}{1 - 2r_f \cos \delta_f \cos \phi} \left[1 - \frac{[c(t_1) - c(t_2)] + \frac{Im(\epsilon) + 2Re(\epsilon)r_f \cos \phi \sin \delta_f}{Re(\epsilon) - 2Im(\epsilon)r_f \cos \phi \sin \delta_f} [s(t_1) - s(t_2)]}{\tau_S \Gamma (1+x^2) (e^{-\Gamma_S t_1} - e^{-\Gamma_S t_2})} \right] + 2r_f \sin \delta_f \sin \phi, \quad (9)$$

where $x = \Delta m/\Gamma$, $c(t) = e^{-t\Gamma} [\cos(\Delta m t) - x \sin(\Delta m t)]$, and $s(t) = e^{-t\Gamma} [x \cos(\Delta m t) + \sin(\Delta m t)]$. In the first line, those terms proportional to r_f represent the new effect $A_{CP}^{\text{int}}(t_1, t_2)$, and those without




Figure 2: The plot of time-integrated CP asymmetries in the $D^+ \rightarrow \pi^+ K_S^0$ given in [1], where the left plot is the total CP asymmetry and the right one is the new CP -violation effect. The dashed lines is the theoretical uncertainties of our predictions.

r_f are the CP violation in the neutral kaon mixing. The second line, which is independent of $t_{1,2}$, corresponds to the direct CP asymmetry in charm decays. The time-integrated CP asymmetries in the $D^+ \rightarrow \pi^+ K_S^0$ and the new CP violating effect are exhibited in Fig. 2. In some ranges of t_1 and t_2 , these two quantities are relatively larger than other ranges. The experimental investigations could choose the favorable time intervals. In some experiments, including Belle and LHCb, the new CP violation effect is in absence [7, 13, 14, 15, 16]. However, since this new effect is of the same order as the direct CP asymmetries in the SCS processes, it cannot be neglected in these measurements.

In order to measure the new CP -violation effect in experiments, we propose an observable

$$\Delta A_{CP}^{\pi^+, K^+} \equiv A_{CP}^{D^+ \rightarrow \pi^+ K_S^0}(t_1, t_2) - A_{CP}^{D_s^+ \rightarrow K^+ K_S^0}(t_1, t_2). \quad (10)$$

The CP violation in the kaon mixing cancels in the above difference, and the direct CP violation is negligible. Our global-fit analysis indicates that the $2r_f \cos \phi \cos \delta_f$ term in denominator of Eq. (9) matters little due to the large strong phases δ_f [12], which is consistent with those derived in the literature [17, 18, 19, 20] and supported by experiment [21]. Then we have

$$\Delta A_{CP}^{\pi^+, K^+} \simeq A_{CP}^{\text{int}, D^+ \rightarrow \pi^+ K_S^0}(t_1, t_2) - A_{CP}^{\text{int}, D_s^+ \rightarrow K^+ K_S^0}(t_1, t_2). \quad (11)$$

The model-independent $SU(3)$ symmetry analysis shows the new effects in two modes are constructive in $\Delta A_{CP}^{\pi^+, K^+}$. The dependencies of $\Delta A_{CP}^{\pi^+, K^+}$ on t_1 and t_2 are plotted in Fig. 3. $\Delta A_{CP}^{\pi^+, K^+}$ is of order of 10^{-3} in most of time intervals, which is accessible at Belle II and LHCb upgrade experiments [6, 15, 22, 23].

In summary, we investigated the time-dependent and time-integrated CP violation in charm decays into neutral kaons. We first pointed out a new measurable CP -violating effect, the interference between charm decays and kaon mixing, exists in these modes. It could be revealed by measuring the difference of CP asymmetries in the $D^+ \rightarrow \pi^+ K_S^0$ and $D_s^+ \rightarrow K^+ K_S^0$ modes on Belle




Figure 3: The dependences of $\Delta A_{CP}^{\pi^+, K^+}$ on t_1 and t_2 given in [1].

II and LHCb upgrade. In addition, an observation with non-zero CP violation at $t = 0$ would signal new physics.

Acknowledgements

This work was supported in part by the National Natural Science Foundation of China under Grants No. 11347027, 11505083, by the Ministry of Science and Technology of R.O.C. under Grant No. MOST-104-2112-M-001-037-MY3, and by the Fundamental Research Funds for the Central Universities under Grant No. lzujbky-2015-241 and lzujbky-2017-97.

References

- [1] F. S. Yu, D. Wang and H. n. Li, arXiv:1707.09297 [hep-ph].
- [2] J. H. Christenson, J. W. Cronin, V. L. Fitch and R. Turlay, Phys. Rev. Lett. **13**, 138 (1964).
- [3] B. Aubert *et al.* [BaBar Collaboration], Phys. Rev. Lett. **87**, 091801 (2001).
- [4] K. Abe *et al.* [Belle Collaboration], Phys. Rev. Lett. **87**, 091802 (2001).
- [5] C. Patrignani *et al.* [Particle Data Group Collaboration], Chin. Phys. C **40**, 100001 (2016).
- [6] R. Aaij *et al.* [LHCb Collaboration], Phys. Rev. Lett. **116**, 191601 (2016).
- [7] B. R. Ko *et al.* [Belle Collaboration], Phys. Rev. Lett. **109**, 021601 (2012), Erratum: [Phys. Rev. Lett. **109**, 119903 (2012)].
- [8] H. J. Lipkin and Z. z. Xing, Phys. Lett. B **450**, 405 (1999); G. Dmbrosio and D.-N. Gao, Phys. Lett. B **513**, 123 (2001).
- [9] Y. Grossman and Y. Nir, JHEP **1204**, 002 (2012).

- [10] S. Bianco, F. L. Fabbri, D. Benson and I. Bigi, Riv. Nuovo Cim. **26N7**, 1 (2003).
- [11] H. n. Li, C. D. Lu and F. S. Yu, Phys. Rev. D **86**, 036012 (2012).
- [12] D. Wang, F. S. Yu, P. F. Guo and H. Y. Jiang, Phys. Rev. D **95**, 073007 (2017).
- [13] R. Aaij *et al.* [LHCb Collaboration], JHEP **1306**, 112 (2013).
- [14] R. Aaij *et al.* [LHCb Collaboration], JHEP **1407**, 041 (2014).
- [15] R. Aaij *et al.* [LHCb Collaboration], JHEP **1410**, 25 (2014).
- [16] R. Aaij *et al.* [LHCb Collaboration], Phys. Lett. B **767**, 177 (2017).
- [17] S. Müller, U. Nierste and S. Schacht, Phys. Rev. D **92**, 014004 (2015).
- [18] B. Bhattacharya and J. L. Rosner, Phys. Rev. D **81**, 014026 (2010).
- [19] H. Y. Cheng and C. W. Chiang, Phys. Rev. D **81**, 074021 (2010).
- [20] D. N. Gao, Phys. Rev. D **91**, 014019 (2015).
- [21] Q. He *et al.* [CLEO Collaboration], Phys. Rev. Lett. **100**, 091801 (2008).
- [22] A. J. Schwartz, [arXiv:1701.07159 [hep-ex]].
- [23] R. Aaij *et al.* [LHCb Collaboration], Eur. Phys. J. C **73**, 2373 (2013).