

# Drawing bobbin lace graphs, or, fundamental cycles for a subclass of periodic graphs

Therese Biedl and Veronika Irvine \*

David R. Cheriton School of Computer Science, University of Waterloo

**Abstract.** In this paper, we study a class of graph drawings that arise from bobbin lace patterns. The drawings are periodic and require a combinatorial embedding with specific properties which we outline and demonstrate can be verified in linear time. In addition, a lace graph drawing has a topological requirement: it contains a set of non-contractible directed cycles which must be homotopic to  $(1, 0)$ , that is, when drawn on a torus, each cycle wraps once around the minor meridian axis and zero times around the major longitude axis. We provide an algorithm for finding the two fundamental cycles of a canonical rectangular schema in a supergraph that enforces this topological constraint. The polygonal schema is then used to produce a straight-line drawing of the lace graph inside a rectangular frame. We argue that such a polygonal schema always exists for combinatorial embeddings satisfying the conditions of bobbin lace patterns, and that we can therefore create a pattern, given a graph with a fixed combinatorial embedding of genus one.


## 1 Introduction

Bobbin lace is a 500-year-old fibre art-form created by braiding threads together in complex patterns. See Figure 1. Bobbin lace can depict landscapes, figures, flowers, as well geometric and abstract designs. Common to all bobbin lace compositions is the use of doubly periodic patterns to fill regions of any shape or size. It is the study of these periodic patterns that we pursue here. To create bobbin lace, threads, which are wound around wooden bobbins to facilitate handling, are arranged left to right in linear order  $t_1, t_2, \dots, t_{2n-1}, t_{2n}$ . The lacemaker selects four consecutive threads, starting at an odd index, and crosses the four threads over and under each other to form an alternating braid. After one or several crossings are made in this manner, the four threads are set aside and another set of four (not the same four, but possibly using a subset of the original four) is selected, again starting at an odd index, and braided. Since the selected threads are consecutive starting at an odd index, we can describe the pattern by tracking the movement of pairs of threads rather than individual strands.

The application of mathematics to the study of fibre arts dates back to the beginnings of computer science. A survey of various areas, including knitting and weaving, is presented by Belcastro and Yackel [3]. Grishanov et al. take a deep

---

\* Research supported by NSERC. Thank you to Anna Lubiw for helpful input.



**Fig. 1.** Bobbin lace. (a) Pattern, (b) Lace with two different  $\zeta(v)$  mappings, (c) Periodic graph drawing. Four osculating circuits distinguished using black and gray.

look into knot theory and its relevance to the topology of textiles [7]. The second author and Ruskey [8] were the first to develop a mathematical model for bobbin lace and express its patterns using graph drawings. Specifically, a lace pattern can be represented as  $(\Gamma(G), \zeta(v))$  where  $G$  is a combinatorial embedding that captures the flow of pairs of threads from one grouping of four to another,  $\Gamma(G)$  gives a specific drawing of  $G$  which assigns a geometry to the position of the braids, and  $\zeta(v)$  is a mapping from each node  $v \in V(G)$  to a mathematical braid word which specifies the over and under crossings performed on the subset of four threads that meet at  $v$ . A systematic exploration of different  $\zeta(v)$  mappings is straight-forward, although time consuming, and has been undertaken by lacemakers for several traditional patterns; see [8] for more details. Discovering which pairs of threads can be successfully combined, as represented by  $\Gamma(G)$ , is a much harder task and is therefore the focus of this paper.

Lace patterns are doubly periodic. As a result, the graphs are drawn on the surface of a torus and can be lifted to their universal cover of infinite periodic planar graphs. The main question investigated in this paper is to decide, for a directed graph with a fixed rotation system, whether we can construct a straight-line drawing that is a lace pattern. We argue that recognizing such graphs can be done in linear time and depends only on their combinatorial structure (some of these results were reported earlier [8]). A toroidal embedding can be drawn as a straight-line planar graph drawing with a rectangular outer face by choosing a canonical rectangular schema [1,5]. The main challenge is to find two fundamental cycles to serve as the borders of the rectangle under the constraints of the topological requirements of the lace pattern and restrictions imposed by the drawing algorithm itself. We show that a suitable polygonal schema can always be found.

## 2 Background

We assume familiarity with graphs and combinatorial embeddings; see for example [9]. Throughout this paper,  $G = (V, E)$  is a directed graph that comes with a rotation system, i.e., a clockwise order of edges around each vertex. We will assume that the rotation system describes a cellular combinatorial embedding on an orientable surface (i.e., the complement of  $G$  on the orientable surface is a collection of open topological disks). The genus of the cellular embedding can be

determined from the rotation system by computing the *facial walk* consisting of edges and vertices incident to each face in order while walking around the face. All embeddings of interest here have a genus of one.

A circuit is a closed path in which a vertex may be visited more than once but no edges are repeated. A cycle is a closed path in which each vertex and edge is only visited once. A *separating cycle* of a graph drawn on a torus  $\mathcal{T}$  is a cycle  $C$  that splits the surface of  $\mathcal{T}$  into two disjoint pieces when removed. A *contractible cycle* is a cycle that can be continuously retracted to a point. On the torus, a contractible cycle bounds a topological disk and is therefore the same as a separating cycle. From now on, we will use the term contractible cycle. A canonical rectangular representation of a torus is bounded by a pair of *non-contractible* cycles, also referred to as fundamental cycles, that have only one vertex in common, their point of intersection.

## 2.1 Conditions on lace pattern graphs

As should be obvious from the description, in a lace pattern graph every vertex  $v$  must have exactly two incoming and two outgoing edges, corresponding to the two pairs of threads that meet at  $v$ , are being braided together, and then separate again. We call such a graph a 2-2-regular digraph.

Lace patterns are doubly periodic, i.e., they can tile the plane by translation in two non-parallel directions. One “tile” of the repeat can be drawn using a canonical rectangular schema in which the position of a thread intersecting the horizontal boundaries of the schema has the same abscissa top and bottom and a thread intersecting the vertical boundaries of the schema has the same ordinate value left and right. In graph drawing this is called periodic.

We do not want the lace created from the pattern to “fall apart”, i.e., the graph needs to be suitably connected. In graph-theoretical terms, this means  $G$  must be a cellular embedding.

For a lace pattern to be executable, there must exist a partial ordering of the vertices such that, for any directed edge, the braid mapped to the tail vertex is worked before that of the head. The pattern, when repeated over the infinite plane, cannot contain any directed cycles, or equivalently, the associated graph on the torus must be free from contractible directed cycles.

The graph may have loops but they must be non-contractible. Parallel edges with the same orientation do not represent a change in the four threads being braided and therefore do not appear in a lace graph. Parallel edges with opposite orientation must form a non-contractible directed cycle. Summarizing, a lace graph may be a multigraph but no loop or parallel edge can be a face.

In summary, the following three conditions are required of any lace pattern graph  $G$  with a fixed combinatorial embedding:

- C1.  $G$  is a directed 2-2-regular digraph.
- C2. The rotation system of  $G$  describes a toroidal cellular embedding in which all facial walks contain at least 3 edges.
- C3. All directed cycles of  $G$  are non-contractible.

It is easy to check in linear time whether (C1) holds. To test (C2), we first compute the facial walks to ascertain the number of faces, and can then determine whether the embedding is toroidal since, by Euler’s formula, such an embedding with  $n$  vertices and  $2n$  edges must have exactly  $n$  faces.

We are not aware of any results regarding how to test (C3), i.e., to check whether some directed cycle is contractible. It is possible to check whether a given graph has a contractible cycle [4], but this does not take edge direction into account. We will test whether (C3) holds via a detour: we use another condition (C3’) that is a necessary condition for (C3), and then argue that if (C3’) holds then (C3) can be tested in linear time.

Assume we have a 2-2-regular digraph with a fixed combinatorial embedding. There are two ways in which edges can be arranged at a vertex  $v$  with  $\text{indeg}(v) = \text{outdeg}(v) = 2$ : Either *rotationally alternating* in which edges alternate between incoming and outgoing directions or *rotationally consecutive* with edges in the order incoming, incoming, outgoing, outgoing. We define the following condition:


C3’. At all vertices, the outgoing arcs are rotationally consecutive.

Clearly (C3’) can be tested in linear time. Irvine and Ruskey showed [8] that if (C3) holds, then (C3’) must hold as well. We will argue in the next section that if (C3’) holds, then one can test in linear time whether (C3) holds.

There is one more important restriction for a workable bobbin lace pattern. The threads in the periodic pattern are continuous; threads are neither removed (by cutting) nor added (by knotting or weaving in). In other words, to fill a rectangle of fixed width and undetermined height, a fixed set of threads starts at the top of the rectangle and the same set of threads terminates (not necessarily in the same order) at the bottom of the rectangle. To achieve this, the threads in one repeat of the pattern must not have a net drift to the right or left. See [8] for a more detailed discussion of this *thread conservation* property.

The thread conservation property is impossible to formulate as a condition of the combinatorial embedding because it is affected by how the graph is drawn on the torus. In particular, consider Figure 2 which shows two drawings of the same graph on the torus. Both drawings have the same combinatorial embedding, yet on the left side of Figure 2 the red (thick) circuit returns to its starting point while on the right side of the figure the circuit “wraps around” or drifts to the right. Thread conservation demands that we fix more than the combinatorial embedding of the graph; it imposes an additional topological constraint. Creating a drawing consistent with this topological constraint is the main problem addressed in this paper.

To formulate the thread conservation property in a mathematically precise way, we need to introduce more terminology. Fix a digraph  $G$  with a combinatorial embedding such that (C1,C2,C3’) holds. Thus every vertex  $v$  has two incoming and two outgoing edges, arranged rotationally consecutive. Define the *left/right incoming/outgoing* edge of  $v$  as follows: Going in clockwise order around  $v$ , we encounter first the left outgoing edge, then the right outgoing edge, then the right incoming edge and then the left incoming edge. The *left face* of a directed edge  $v \rightarrow w$  is the face to the left of it while walking from  $v$  to  $w$ .



**Fig. 2.** A Dehn twist changes a valid lace graph on left into an invalid one on right.

Consider two edge-disjoint directed circuits  $C_1, C_2$  that have a vertex  $v$  in common. There are two possible ways in which  $C_1$  and  $C_2$  meet at  $v$ . In an *osculating intersection* these circuits only touch (“kiss”), i.e., both incoming and outgoing edges of  $C_1$  are on the left side of  $v$  and the edges of  $C_2$  are on the right side of  $v$  (or vice versa). In contrast, at a *transverse intersection* the circuits truly cross, i.e.,  $C_1$  enters from one side of  $v$  and exits from the other and  $C_2$  does the same but in the opposite direction. If  $C_1$  uses the right incoming (and the left outgoing) edge of  $v$  we say that  $C_1$  *crosses  $C_2$  from right to left* at  $v$ . Summing over all shared vertices, we define the *algebraic crossing number* of  $C_1$  and  $C_2$  to be:

$$\hat{i}(C_1, C_2) = \#\{C_1 \text{ crosses } C_2 \text{ left-to-right}\} - \#\{C_1 \text{ crosses } C_2 \text{ right-to-left}\}$$

A (*canonical*) *polygonal schema* for a toroidal graph consists of two fundamental cycles (called the *meridian  $M$*  and the *longitude  $L$* ) such that  $M$  and  $L$  intersect in exactly one point. Cutting  $G$  along the edges of  $M \cup L$  will result in a topological disk. A circuit  $C$  belongs to *homotopy class  $(m, \ell)$*  (with respect to a fixed polygonal schema) if  $\hat{i}(C, L) = m$  and  $\hat{i}(C, M) = \ell$ . The following is known:

**Lemma 1.** [10, p. 209] *Two closed simple curves  $C, C'$  have  $\hat{i}(C, C') = 0$  if and only if they belong to the same homotopy class.*

With these terms in place, we can now imply the thread conservation property via the following constraint:

- C4. There exists a meridian  $M$ , a longitude  $L$  and a partition  $\mathcal{P}(G)$  of edges into osculating directed circuits such that each circuit in the partition is in the  $(1, 0)$ -homotopy class.

### 3 Osculating circuits and some properties

**Lemma 2.** *Presume  $(C1, C2, C3')$  holds. The edges of  $G$  can be partitioned into a set  $\mathcal{P}(G)$  of disjoint directed circuits such that no two circuits in  $\mathcal{P}(G)$  have a transverse intersection. Furthermore, this partition is unique and can be found in linear time.*

*Proof.* Arbitrarily select an edge  $e$  of  $G$  as the start of a circuit  $P_e$ . If this edge is left incoming at its head  $v$ , then let  $e_1$  be the left outgoing edge at  $v$ , else let

$e_1$  be the right outgoing edge at  $v$ . Put differently,  $e$  and  $e_1$  are on the “same side” of  $v$ . Append  $e_1$  to  $P_e$  and repeat the operation at the head of  $e_1$ . Since the graph is finite, we eventually must close a circuit  $P_e$ ; this is the first element of the partition  $\mathcal{P}(G)$ . In fact, circuit  $P_e$  must exactly finish at edge  $e$ , because for any edge the rule of “stay on the same side” uniquely determines the edge before and after in the circuit.

Now select edge  $e'$  of  $G$  that was not in  $P_e$ , and repeat the process starting at  $e'$ . The new circuit will not contain an edge of  $P_e$  by the same “stay on the same side” rule. Thus we obtain the next circuit in the partition. Repeat until all edges belong to some circuit. Since there is never a choice about which next edge to take, the partition is unique. Each edge is visited exactly once resulting in a linear runtime.  $\square$

We call the circuits in  $\mathcal{P}(G)$  the *osculating circuits* of  $G$ , see also Figure 1(c). The following observation allows us to test whether (C3) holds:

**Lemma 3.** *Presume (C1,C2,C3') holds. If  $G$  has a simple contractible directed cycle  $C$ , then  $C$  bounds a face of  $G$ .*

*Proof.* (Sketch) Arbitrarily declare one face to contain the origin so that “inside” and “outside” is well-defined for any contractible directed cycle  $C$ . Let  $O_C$  and  $I_C$  be the number of faces outside and inside  $C$ , and consider a directed contractible cycle  $C$  that maximizes  $|O_C - I_C|$ . Prove, by contradiction, that a directed cycle containing an edge in its interior cannot maximize  $|O_C - I_C|$  which means that the inside of  $C$  is a face as required. Details are in the appendix.  $\square$

It follows that to test condition (C3), we first test for condition (C1,C2,C3'). If these hold, simply test whether any facial walk is a directed cycle. Clearly this takes linear time.

From now on, we will assume that (C1-C3) hold and focus our attention on finding a meridian and a longitude to satisfy (C4). We distinguish two major cases based on whether or not an osculating circuit is simple. It turns out that when a circuit visits a vertex twice,  $\mathcal{P}(G)$  has a trivial structure.

**Lemma 4.** *Presume (C1-C3) hold. If some osculating circuit  $C \in \mathcal{P}(G)$  visits a vertex twice, then  $C$  is the only element in  $\mathcal{P}(G)$ .*

*Proof.* Follow  $C$ , a non-simple directed circuit, until the first time a vertex  $v \in C$  is reached for the second time.  $C$  can be partitioned into a simple directed cycle  $C'$  (by taking the part from  $v$  to  $v$  that we just followed) and a directed circuit  $C''$  (the rest of  $C$ ). Both  $C'$  and  $C''$  are incident to  $v$ .

Fix an arbitrary drawing of  $G$  on the torus  $\mathcal{T}$ . By (C3)  $C'$  is non-contractible, so cutting  $\mathcal{T}$  along  $C'$  produces a cylinder. The cut will split  $v$  into two vertices,  $v'$  and  $v''$ , one on each boundary of the cylinder.

Because of the rotationally consecutive construction of  $C$ ,  $C'$  and  $C''$  must intersect transversely at  $v$  (otherwise,  $C$  would have terminated the first time it returned to  $v$ ). Thus  $C''$  contains one incident edge each at the two copies  $v'$  and  $v''$  of  $v$ . Taking the subpath  $D$  of  $C''$  between  $v'$  and  $v''$ , we obtain a path

that travels on the cylinder from one boundary to the other. Cutting along  $D$  will cut the cylinder into one or more disks.

Now consider some other circuit  $P \in \mathcal{P}(G)$ . It cannot have a transverse crossing with either  $C'$  or  $C''$ , because  $C$  and  $P$  are osculating circuits. Therefore, it intersects neither  $C'$  nor  $D$  and we can conclude that  $P$  resides entirely within (or on the boundary) of one of the disks of  $\mathcal{T} - C' - D$ . But then  $P$  is a contractible directed circuit, a contradiction. So  $\mathcal{P}(G)$  contains no osculating circuits other than  $C$ .  $\square$

The osculating partition can contain more than one directed circuit, see Figure 1, but then all circuits in  $\mathcal{P}(G)$  must be simple directed cycles. Further, we can say:

**Lemma 5.** *If (C1-C3) hold, then all directed cycles  $P_i \in \mathcal{P}(G)$  belong to the same homotopy class.*

*Proof.* If  $\mathcal{P}(G)$  contains a non-simple circuit then, by the previous lemma, it is the only member of  $\mathcal{P}(G)$  and the claim holds trivially. Otherwise, the osculating circuits of  $\mathcal{P}(G)$  are all simple closed curves. None of them intersect transversally, which means that  $\hat{i}(P_i, P_j) = 0$  for all pairs of osculating circuits  $i \neq j$ . This proves the result by Lemma 1.  $\square$

## 4 Finding a polygonal schema

For the rest of this paper, we assume that the input graph embedding satisfies (C1-C3), and work on finding a meridian  $M$  and longitude  $L$  such that (C4) is satisfied. Additionally we require that  $M$  and  $L$  intersect each edge of the graph at most once, a necessary constraint for applying existing drawing algorithms.

In general, finding a polygonal schema with vertex-disjoint interiors is NP-hard [5]. However, for the purpose of drawing the lace-pattern, we do not need to find a polygonal schema within the given graph; it suffices (and in fact, is preferable) to add vertices and edges to the graph and find a polygonal schema within the additions. In this manner, the original edges of  $G$  are not on the schema boundary giving more freedom to where they can be placed. We explain how to find such a supergraph with  $O(n)$  nodes which, in combination with existing drawing algorithms, gives:

**Theorem 1.** *Given a digraph  $G$  that satisfies (C1-C3), we can draw a lace pattern in linear time. The drawing resides in an  $O(n^4) \times O(n^4)$ -grid.*


Proving the existence of a meridian and longitude that satisfy (C4) is straightforward if the graph can be partitioned into simple, non-contractible, osculating cycles. By the Dehn-Lickorish theorem (see e.g. [6]) there exists a homeomorphism that maps any simple, non-contractible cycle to the  $(1, 0)$  homotopy class of the torus. However, finding a homeomorphism that will convert a given drawing of the graph into a lace pattern is not trivial. Further, it is not clear how to handle the case in which the (unique) osculating circuit is not simple. For

this reason, we present a five step algorithm for finding a suitable meridian  $M$  and longitude  $L$  in a supergraph of  $G$  and use this rectangular representation to enforce the desired homotopy class.

<b>DrawLacePattern Algorithm</b>
<p>A. Partition <math>G</math> into a set <math>\mathcal{P}(G)</math> of osculating circuits and select one directed circuit <math>P^*</math> from the set.</p> <p>B. Create the offset graph <math>\mathcal{O}(G)</math>.</p> <p>C. Find a simple cycle <math>M</math> in <math>\mathcal{O}(G)</math> such that <math>\hat{i}(M, P^*) = 0</math> and <math>M</math> intersects every edge of <math>G</math> at most once.</p> <p>D. Find a simple cycle <math>L</math> in <math>\mathcal{O}(G)</math> such that <math>\hat{i}(L, P^*) = 1</math>, <math>M</math> and <math>L</math> intersect exactly once, and <math>L</math> intersects every edge of <math>G</math> at most once.</p> <p>E. Use existing torus-drawing techniques to draw <math>G</math> on a rectangle with meridian <math>M</math> and longitude <math>L</math>.</p>

All steps are linear or constant time, and step E. will create a drawing of the required size, proving the theorem. Step A. was explained in Section 3 already; all other steps are explained below.

**Creating the offset graph:** We first create an offset graph  $\mathcal{O}(G)$  in which we will choose a suitable meridian  $M$  and longitude  $L$ . Roughly speaking,  $\mathcal{O}(G)$  is obtained by creating two copies of every osculating circuit  $P$  in  $\mathcal{P}(G)$ , which we will represent as  $\ddot{P}$  (used to find  $M$ ) and  $\mathbb{L}P$  (used to find  $L$ ). In each copy, the circuits are separated (they do not share vertices as they do in  $G$ ) and are simple cycles. The copies lie on top of  $G$  introducing crossings which we remove by inserting dummy vertices. Finally, for the simple cycles of  $\mathbb{L}P$ , we connect the two halves of each vertex, split in the process of separating osculating paths, by introducing “crossover-edges”.



**Fig. 3.** Create offset graph. (a) Close-up near a vertex showing  $\ddot{P}$  (thick, dashed, square, orange),  $\mathbb{L}P$  (thick, solid, triangle, green), crossover-edges (thin, solid, purple) and shortcut-edges (thin, dotted, blue). (b) Offset graph of non-simple osculating circuit. (c) Offset graph with multiple simple osculating cycles.

Figure 3 illustrates such a toroidal graph embedding  $\mathcal{O}(G)$ , with  $O(n)$  vertices and edges, which we will now define formally. For each vertex in  $\mathcal{O}(G)$  we define  $f_\ell(v)$  to be the face to the left of the left incoming and left outgoing edges of  $v$ ,



$f_r(v)$  to be the face to the right of the right incoming and right outgoing edges of  $v$ , and  $f_c(v)$  to be the face incident to the left incoming and right incoming edges of  $v$ .

- Initially,  $\mathcal{O}(G)$  contains all vertices and edges of  $G$ , embedded as in  $G$ .
- For every vertex  $v \in V(G)$ , add four new vertices  $\ddot{v}_\ell, \mathbb{L}v_\ell, \ddot{v}_r, \mathbb{L}v_r$ . Place  $\ddot{v}_\ell, \mathbb{L}v_\ell$  in  $f_\ell(v)$  such that  $\mathbb{L}v_\ell$  is closer to the left incoming edge than  $\ddot{v}_\ell$ . Place  $\ddot{v}_r, \mathbb{L}v_r$  in  $f_r(v)$  such that  $\ddot{v}_r$  is closer to the right incoming edge than  $\mathbb{L}v_r$ .
- For every edge  $e = v \rightarrow w \in E(G)$ , we add two new edges  $\ddot{e}$  and  $\mathbb{L}e$ . If  $e$  is left outgoing at  $v$ , then  $\ddot{e}$  starts at  $\ddot{v}_\ell$  and  $\mathbb{L}e$  starts at  $\mathbb{L}v_\ell$ . else  $\ddot{e}$  starts at  $\ddot{v}_r$  and  $\mathbb{L}e$  starts at  $\mathbb{L}v_r$ . If  $e$  is left incoming at  $w$ , then  $\ddot{e}$  ends at  $\ddot{w}_\ell$  and  $\mathbb{L}e$  ends at  $\mathbb{L}w_\ell$ , else  $\ddot{e}$  ends at  $\ddot{w}_r$  and  $\mathbb{L}e$  ends at  $\mathbb{L}w_r$ .
- Note that edge  $e \in E(G)$  may be intersected by its copies  $\ddot{e}$  and  $\mathbb{L}e$ . This occurs, for example, when  $e$  is right outgoing at its tail and left incoming at its head. We remove these crossings (so that we again have a toroidal embedding) in the standard way by inserting dummy vertices that subdivide  $e, \ddot{e}, \mathbb{L}e$ . (In the following descriptions, we will ignore these dummy-vertices, and speak of an edge  $e$ , even though it has become a path with 3 edges.)
- For every osculating circuit  $P$  there are now two circuits  $\ddot{P}$  and  $\mathbb{L}P$ , using for each edge  $e \in P$  the corresponding edges  $\ddot{e}$  and  $\mathbb{L}e$ . Note that  $\ddot{P}$  and  $\mathbb{L}P$  are simple, even if  $P$  is not. When  $P$  visits a vertex  $v$  twice, once via the incoming left and outgoing left edges of  $v$  and once via the incoming right and outgoing right edges of  $v$ , the corresponding  $\ddot{P}$  visits  $\ddot{v}_\ell$  and  $\ddot{v}_r$  respectively and similarly for  $\mathbb{L}P$ . Due to the order of the copies near  $v$ ,  $\ddot{P}$  and  $\mathbb{L}P$  do not cross.
- Next add the *crossover-edge*  $\mathbb{L}e_v = (\mathbb{L}v_\ell, \mathbb{L}v_r)$  for each vertex  $v \in V$ . To obtain a toroidal embedding, route this edge so that it crosses both incoming edges of  $G$  at  $v$ , and also the edge  $\ddot{e}$  that is incoming to  $\ddot{v}_r$ . These crossings are again replaced by dummy-vertices. In the crossover-edge insert a vertex  $v_c$  in the face  $f_c(v)$ .
- For a straight-line drawing, an edge  $e$  in  $E(G)$  must not cross the rectangular frame twice. Consider an edge  $\mathbb{L}e$  that crosses  $e = v \rightarrow w$  where  $\mathbb{L}e$  originates inside the face  $f_c(w)$ , say  $\mathbb{L}e = \mathbb{L}v_r \rightarrow \mathbb{L}w_\ell$ . It may happen that the chosen longitude  $L$  contains  $\mathbb{L}e$  followed by the crossover-edge  $\mathbb{L}e_w = (\mathbb{L}w_\ell, \mathbb{L}w_r)$  resulting in a double crossing of  $e$  by  $L$ . To avoid this situation, we add a *shortcut-edge* to  $\mathcal{O}(G)$  that connects a point on  $\mathbb{L}e$  inside  $f_c(w)$  to  $w_c$ . Note that any circuit using  $\mathbb{L}e\mathbb{L}e_w$  acts the same (with respect to algebraic crossing numbers) as a circuit shortened via the shortcut-edge, since all other crossings are unaffected.

**Finding the meridian:** In step A. we selected an osculating circuit  $P^*$ . Define  $M$  to be the copy  $\ddot{P}^*$  of circuit  $P^*$  in the offset graph.  $M$  is a simple cycle. It may cross  $P^*$  repeatedly but it does so only by switching back and forth between being left of  $P^*$  and right of  $P^*$ . In other words,  $\hat{i}(P^*, M) = 0$  as desired. Finally  $M$  intersects every edge of  $G$  at most once by construction of  $\ddot{P}$ .

**Finding the longitude:** Define  $\mathcal{L}(G)$  to be the subgraph of  $\mathcal{O}(G)$  taking only the copies  $LP$  of osculating circuits, the crossover-edges, and the shortcut-edges. We claim that we can find a suitable longitude in  $\mathcal{L}(G)$ . The algorithm is quite simple (find a shortest path within  $\mathcal{L}(G)$ , with some edges removed), but arguing that it works is not.

**Case 1) Circuit  $P^*$ :** First consider the easier case in which  $P^*$  is not simple and is therefore the only element in  $\mathcal{P}(G)$ . Let  $v$  be a vertex visited twice by  $P^*$ , hence  $Lv_\ell$  and  $Lv_r$  both belong to  $LP^*$ . Define  $L^*$  to be a subpath of  $LP^*$  between  $Lv_\ell$  and  $Lv_r$  and  $\hat{L}$  is  $L^*$  plus the crossover-edge  $(Lv_r, Lv_\ell)$ . We have constructed  $LP^*$  such that it does not intersect  $\ddot{P}^*$ . So  $L^*$  does not intersect  $M$ . The crossover-edge  $(Lv_r, Lv_\ell)$  intersects  $M$  exactly once. Therefore,  $\hat{L}$  intersects  $M$  exactly once as desired.

$L^*$  and  $P^*$  may intersect numerous times between  $Lv_\ell$  and  $Lv_r$ . But  $Lv_\ell$  is to the left of  $P^*$  while  $Lv_r$  is to the right, so in total  $L^*$  has one more left-to-right intersection than right-to-left-intersection. The crossover-edge  $(Lv_r, Lv_\ell)$  intersects both incoming edges at  $Lv$ , and both of these edges belong to  $P^*$ . This adds two more right-to-left intersections, and so in total  $\hat{i}(\hat{L}, P^*) = 1$  as desired.

**Case 2) Simple  $P^*$ :** If  $P^*$  is simple, then we find  $L$  using a path that connects “both sides” of  $P^*$  without crossing  $P^*$ . The existence of such a path is non-trivial, and crucially needs condition (C2), i.e., that the embedding is cellular. Specifically, we show:

**Lemma 6.** *Presume (C1-C3) holds. Let  $P$  be a simple osculating cycle. Then there exists a directed walk  $W$  in  $G$  that starts at a vertex  $v \in P$  with a left outgoing edge  $e_1 \notin P$ , ends at a vertex  $w \in P$  with a right incoming edge  $e_k \notin P$ , and has no transverse intersection or shared edges with  $P$ .*

*Proof.* (Sketch) There must exist such an edge  $e_1$ , else  $P$  would bound a face, contradicting (C2) or (C3) since  $P$  is a directed cycle. One can now argue that starting from  $e_1$  and always taking the left outgoing edge, we must reach such an edge  $e_k$  or find a contradiction to (C2). Details are in the appendix.  $\square$

Let  $Q$  be such a walk in  $G$  from  $v \in P^*$  to  $w \in P^*$ , shortened by eliminating directed cycles (if any) so that it becomes a simple path. We obtain the longitude  $L$  by “translating”  $Q$  into  $\mathcal{L}(G)$  and adding a subpath of  $LP^*$ .


Note that the directed edges of  $Q$  need not be rotationally consecutive. For example,  $e_i \in E(Q)$  may be right incoming at its head vertex  $x$  while  $e_{i+1} \in E(Q)$  may be left outgoing at its tail vertex  $x$ . If  $x \notin P^*$  then we can add the crossover-edge  $(Lx_\ell, Lx_r)$  to connect  $Le_i$  and  $Le_{i+1}$  without crossing  $M = \ddot{P}^*$ . If  $x \in P^*$  but  $e_i \notin P^*$  and  $e_{i+1} \notin P^*$  then no crossover-edge is required because  $e_i$  and  $e_{i+1}$  are rotationally consecutive. It is not possible to have  $x \in P^*$  and only one of  $e_i \in P^*$  or  $e_{i+1} \in P^*$  because of the way in which  $Q$  is defined.

Formally, let  $Q = e_1, \dots, e_k$  be a simple path in  $G - E(P^*)$  from  $v \in P^*$  to  $w \in P^*$  and let  $L^- = Le_1, \dots, Le_k$  be a simple path using the corresponding edges in  $\mathcal{L}(G)$  and crossover-edges as needed.  $L^-$  begins at  $Lv_\ell$  since  $e_1$  is left outgoing and ends at  $Lw_r$  since  $e_k$  is right incoming. Define  $L^*$  to be the subpath

of  $\mathbb{L}P^*$  connecting  $\mathbb{L}w_\ell$  to  $\mathbb{L}v_r$ . Finally, define  $\hat{L}$  to be the simple cycle consisting of  $L^-$ , crossover-edge  $(\mathbb{L}w_r, \mathbb{L}w_\ell)$ ,  $L^*$  and crossover-edge  $(\mathbb{L}v_r, \mathbb{L}v_\ell)$ .

Note that  $L^-$  and  $L^*$  never cross the meridian  $M = \mathring{P}^*$ . The only place where  $\hat{L}$  can intersect  $M$  is at the two crossover-edges  $(\mathbb{L}v_r, \mathbb{L}v_\ell)$  and  $(\mathbb{L}w_r, \mathbb{L}w_\ell)$ . We claim that exactly one of them intersects  $\mathring{P}^*$ . To see this, recall that  $P^*$  uses the right incoming and outgoing edge at  $v$  and the left incoming and outgoing edge at  $w$ . The order of vertices in the vicinity of  $v$  is  $\mathring{v}_\ell, \mathbb{L}v_\ell, v, \mathring{v}_r, \mathbb{L}v_r$ , and  $\mathring{P}^*$  uses  $\mathring{v}_r$ , so  $(\mathbb{L}v_r, \mathbb{L}v_\ell)$  crosses  $\mathring{P}^*$ . On the other hand the order of vertices in the vicinity of  $w$  is  $\mathring{w}_\ell, \mathbb{L}w_\ell, w, \mathring{w}_r, \mathbb{L}w_r$ , and  $\mathring{P}^*$  uses  $\mathring{w}_\ell$ , so  $(\mathbb{L}w_r, \mathbb{L}w_\ell)$  does not cross  $\mathring{P}^*$ . Therefore,  $L$  and  $M$  cross exactly once as desired.

To see that  $\hat{i}(\hat{L}, P^*) = 1$ , observe that  $L^-$  has no transverse intersections with  $P^*$  and so contributes nothing.  $L^*$  may intersect  $P^*$  repeatedly, alternatingly from left to right and from right to left, however,  $L^*$  starts on the left side of  $P^*$  at  $\mathbb{L}w_\ell$  and ends on the right side of  $P^*$  at  $\mathbb{L}v_r$ , so it has a net of one left to right crossing. The crossover-edges  $(\mathbb{L}w_r, \mathbb{L}w_\ell)$  and  $(\mathbb{L}v_r, \mathbb{L}v_\ell)$  are both right to left crossings. Thus  $\hat{L}$  has one more crossing from right to left than it has crossings from left to right, yielding  $\hat{i}(\hat{L}, P^*) = 1$  as desired.



**Fig. 4.** (a)  $M$  is a shifted copy of  $P^*$ . (b)  $L$  exits from the left side of  $M$  at  $e_1$  and returns on the right side of  $M$  at  $e_k$ . Loops where shortcuts are required are shaded purple. (c) Connect  $e_1$  to  $e_k$  following  $P^*$ .

**Applying shortcuts:** As desired, the simple cycle  $\hat{L}$  intersects  $M$  once and has  $\hat{i}(\hat{L}, P^*) = 1$ . However,  $\hat{L}$  may intersect an edge  $e$  of  $G$  repeatedly. This happens only when  $\mathbb{L}e$  transversely crosses  $e$ , and a crossover-edge at the tail of  $\mathbb{L}e$  follows immediately after. Let  $L$  be the simple cycle obtained from  $\hat{L}$  by substituting the shortcut-edge at any such edge  $e$ . This has the same algebraic crossing number with  $P^*$ , still crosses  $M$  exactly once, and is a suitable longitude.

We finally remark that the simplest way to find  $L$  is as follows. Start with graph  $\mathcal{L}(G)$  and remove all dummy-vertices that lie on  $\mathring{P}^*$ . This effectively disallows any path in  $\mathcal{L}(G)$  that crosses  $M$ . Pick any vertex  $v \in P^*$  and, in what remains of  $\mathcal{L}(G)$ , find a shortest path  $L^-$  from  $\mathbb{L}v_\ell$  to  $\mathbb{L}v_r$  (the vertex within the crossover-edge  $(\mathbb{L}v_r, \mathbb{L}v_\ell)$ ). Finally add the edge  $(\mathbb{L}v_r, \mathbb{L}v_\ell)$  to close the cycle into a suitable longitude  $L$ .


**Drawing the graph:** Let  $\mathcal{T}(G)$  be the graph obtained from  $\mathcal{O}(G)$  by removing all edges that do not belong to  $G, M$  or  $L$  and smoothing any degree-2 vertices.  $\mathcal{T}(G)$  has a natural embedding on a torus with a well defined meridian and longitude. Cut  $\mathcal{T}(G)$  along  $L$  to form graph  $\mathcal{C}(G)$  which has a natural embedding on a cylinder with top and bottom boundaries  $L_t$  and  $L_b$ . Cut  $\mathcal{C}(G)$  along  $M$  to obtain a planar graph  $\mathcal{R}(G)$  with a natural embedding on a rectangle whose top/right/bottom/left sides are formed by the four boundary-paths  $L_t/M_r/L_b/M_\ell$ . Figure 8 in the appendix illustrates this process.

In our construction, we have been careful to ensure that boundary-cycles do not cross an edge of  $G$  more than once. As a consequence, we can now apply known algorithms for straight-line rectangular-frame drawings to  $\mathcal{R}(G)$ . The first such algorithm was given by Duncan et al. [5] and creates a drawing in an  $O(n) \times O(n^2)$ -grid in linear time. Unfortunately, the algorithm does not necessarily produce a periodic drawing. We therefore turn to the drawing algorithm of Aleardi et al. [2], a modification of the algorithm of Duncan et al. which ensures periodic drawings. This is still achieved in linear time but at the cost of increasing the grid-size to  $O(n^4) \times O(n^4)$ -grid. This proves Theorem 1.

## 5 Discussion and open problems

In this paper we provide an algorithm to generate a lace pattern given a directed graph and a fixed combinatorial embedding that satisfy the conditions (C1-C3). Our main challenge was to produce a graph drawing such that the threads in the pattern weave back and forth within a fixed width (follow a directed circuit that wraps once around the minor meridian axis and zero times around the major longitude axis). It turns out that if (C1-C3) are satisfied we can easily define a supergraph of linear size supporting a suitable rectangular schema and extract the schema in linear time. With care, we can sure that neither the meridian nor longitude of the schema intersects an edge of the graph twice, allowing us to reuse existing graph drawing techniques.

Our drawings are still somewhat unsatisfying for use as lace patterns. Consider an edge  $e \in E(G)$  that crosses the rectangular schema, say the horizontal border  $L$ . The edge is broken into two parts  $e_t$  and  $e_b$ , where  $e_t$  intersects the top of the rectangle along  $L_t$  and  $e_b$  intersects the bottom along  $L_b$ . The algorithm of Aleardi et al. [2] ensures that the end points of  $e_t$  and  $e_b$  on  $L$  have the same  $x$ -coordinate but for  $e$  to be truly seen as “one edge”,  $e_t$  and  $e_b$  should also have the same slope. The bend is clearly visible when viewing several tiled repeats of the pattern as shown in Figure 5. At the current time, we cannot see a way to simultaneously satisfy the topological  $(1, 0)$  homotopy constraint enforced by the rectangular representation and the smooth continuity of edges across this rectangular boundary.




**Fig. 5.** Unwanted bends along border. (a) Result of algorithm. (b) Desired result.

## References

1. Aleardi, L.C., Devillers, O., Fusy, É.: Canonical ordering for triangulations on the cylinder, with applications to periodic straight-line drawings. In: International Symposium on Graph Drawing. pp. 376–387. Springer (2012)
2. Aleardi, L.C., Fusy, É., Kostygin, A.: Periodic planar straight-frame drawings with polynomial resolution. In: Latin American Symposium on Theoretical Informatics. pp. 168–179. Springer (2014)
3. Belcastro, S.M., Yackel, C.: Making mathematics with needlework: ten papers and ten projects. AK Peters (2008)
4. Cabello, S., Colin de Verdière, É., Lazarus, F.: Finding cycles with topological properties in embedded graphs. *SIAM Journal on Discrete Mathematics* 25(4), 1600–1614 (2011)
5. Duncan, C.A., Goodrich, M.T., Kobourov, S.G.: Planar drawings of higher-genus graphs. *J. Graph Algorithms Appl.* 15(1), 7–32 (2011)
6. Farb, B., Margalit, D.: A Primer on Mapping Class Groups. Princeton University Press (2011)
7. Grishanov, S., Meshkov, V., Omelchenko, A.: A topological study of textile structures. part i: An introduction to topological methods. *Textile Research Journal* 79(8), 702–713 (2009)
8. Irvine, V., Ruskey, F.: Developing a mathematical model for bobbin lace. *Journal of Mathematics and the Arts* 8(3–4), 95–110 (2014)
9. Mohar, B., Thomassen, C.: Graphs on surfaces. John Hopkins University Press (2001)
10. Stillwell, J.: Classical topology and combinatorial group theory, Graduate Texts in Mathematics, vol. 72. Springer-Verlag (1980)

## A Introduction to bobbin lace



**Fig. 6.** Pattern in progress on lace pillow.

Perhaps the easiest way to understand bobbin lace is to look at the six step process by which it is made.

**Step 1) Prepare the threads.** To manage many long threads without creating a tangled mess, each end of a thread is wound onto one of a pair of bobbins. A bobbin, commonly made from wood, is about 10cm in length. One end is flanged to hold a length of thread and the other end, usually thicker and sometimes weighted with beads, is the handle.

**Step 2) Prepare the pattern.** The lace is worked on top of a firm pillow stuffed with material that can hold pins (such as styrofoam or sawdust). As threads are braided together, they are held in place by pins pushed into the pillow. To start a piece of lace, a pattern is copied onto stiff material such as card stock. Black dots in the pattern represent the position of pins. Before starting to make the lace, all of these dots are pricked through to make small holes. The pattern is then pinned to the pillow.

**Step 3) Hang the bobbins.** The middle of each thread is draped around an anchoring pin at the top of the pattern with the pair of bobbins hanging down on either side. Often the first row of the pattern is a simple weave to anchor the threads.

**Step 4) Braid and pin.** The lacemaker braids the threads working with four consecutive threads at a time. During a sequence of braiding actions, the lacemaker may insert a pin to hold the braid in place. The pin provides resistance so that the lacemaker can apply tension to an individual thread without distorting its neighbours.


**Step 5) Advance to next set.** Once four threads have been braided and pinned, the lace maker moves on to braid another set of four consecutive threads. This new set of threads may include two threads from the previous set but it may also be formed from four completely new threads. Which four threads comprise the next set depends on the pattern.

Steps 4 and 5 are repeated until the lace pattern is completed.

**Step 6) Finish.** The threads are secured (sometimes with a knot, sometimes by weaving them back into the lace) and trimmed off. The pins are removed and the lace may be lifted off the pillow. Once the pins are removed, the lace is held together by the over and under crossings of the threads and friction.

## B Proof of Lemma 3

**Lemma 3.** *Presume  $(C1, C2, C3')$  holds. If  $G$  has a simple contractible directed cycle  $C$ , then  $C$  bounds a face of  $G$ .*



**Fig. 7.** Two cases in proof of Lemma 3. Origin is represented by  $\times$ . (a) Origin inside  $C'$ . (b) Origin outside  $C'$ .

*Proof.* Arbitrarily declare one face to contain the origin so that “inside” and “outside” is well-defined for any contractible directed cycle  $C$ . Let  $O_C$  and  $I_C$  be the number of faces outside and inside  $C$ , and consider a directed contractible cycle  $C$  that maximizes  $|O_C - I_C|$ .

Assume that some vertex  $v \in C$  has an incident edge  $e'$  that is not on  $C$  and belongs to the inside of  $C$ . Follow the osculating circuit  $P'$  containing  $e'$  while staying inside  $C$  until, for the first time, we reach  $C$  again, say at vertex  $w$ . If  $v = w$ , then  $P$  has formed another directed cycle  $C'$  that stays on or inside  $C$  throughout and therefore also is contractible. If  $v \neq w$ , then we can find such a directed cycle  $C'$  by taking  $P$  from  $v$  to  $w$  and then  $C$  from  $w$  to  $v$ .

Consider the case where  $O_C \geq I_C$  (the other case is symmetric). Examine  $|O_{C'} - I_{C'}|$ . If the origin is inside  $C'$ , then  $I_{C'} < I_C$  and  $O_{C'} > O_C$ , therefore  $|O_{C'} - I_{C'}| \geq O_{C'} - I_{C'} > O_C - I_C = |O_C - I_C|$ , a contradiction. If the origin is outside  $C'$  then the outside of  $C'$  is strictly inside the inside of  $C$ , so  $O_{C'} < I_C$  and  $I_{C'} > O_C$ , which implies  $|O_{C'} - I_{C'}| = |I_{C'} - O_{C'}| \geq I_{C'} - O_{C'} > O_C - I_C = |O_C - I_C|$ , a contradiction to the choice of  $C$ .

We therefore conclude that no vertex on  $C$  has an incident edge strictly inside  $C$  and therefore the inside of  $C$  is a face as required.  $\square$

## C Proof of Lemma 6

Before we can prove Lemma 6, we need a few definitions and observations. We will call a simple directed circuit  $C$  a *leftmost circuit* if all edges of  $C$  are left outgoing at their respective tails. Observe that any leftmost circuit  $C$  in a 2-2-regular digraph is actually a cycle, because no vertex can have two left outgoing edges incident to it.

Consider two simple cycles  $P, Q$ . We already defined the algebraic crossing number  $\hat{i}(P, Q)$  if  $P$  and  $Q$  meet in a finite number of points. However, we need a similar concept even if  $P$  and  $Q$  share one or more edges. To define this, replace  $P$  and  $Q$  by cycles  $P'$  and  $Q'$  that are arbitrarily close to  $P$  and  $Q$  in such a way that  $P'$  and  $Q'$  intersect in a finite number of points, and define  $\hat{i}(P, Q) := \hat{i}(P', Q')$ . It is not hard to verify that this number is independent of the choice of  $P'$  and  $Q'$  since we use the algebraic crossing number (rather than the geometric one).

Now we can make the following observation of leftmost cycles:

**Lemma 7.** *Presume (C1-C3) holds. Let  $C$  be a leftmost cycle. Then  $C$  does not bound a face, and  $\hat{i}(C, P) \neq 0$  for all osculating circuits  $P \in \mathcal{P}(G)$ .*

*Proof.*  $C$  cannot possibly bound a face, because it is a directed circuit (hence non-contractible by (C3)) while facial circuits are contractible by (C2). Therefore, there must be some edge  $e_\ell$  on the left side of  $C$  that shares a vertex  $v$  with  $C$ , but does not belong to  $C$ . Since  $C$  uses only left outgoing edges,  $e_\ell$  is necessarily a left incoming edge. Let  $P_\ell$  be the osculating circuit containing  $e_\ell$ . This circuit continues from  $e_\ell$  along  $C$  for possibly some time, but eventually must leave  $C$  again to return back to  $e_\ell$ .  $P_\ell$  can leave  $C$  only at a right outgoing edge, because the left outgoing edges always belong to  $C$ . Therefore in this stretch  $P_\ell$  has crossed  $C$  going from left to right.

Note that  $P_\ell$  can never cross  $C$  from right to left, because all left outgoing edges incident to a vertex in  $C$  belong to  $C$ . Therefore  $\hat{i}(P_\ell, C) \neq 0$ .

Following from Lemma 1,  $P_\ell$  and  $C$  do not belong to the same homotopy class. From Lemma 5, we know that all  $P \in \mathcal{P}(G)$  belong to the same homotopy class so we can conclude  $\hat{i}(P, C) \neq 0$  for all  $P \in \mathcal{P}(G)$ .

Now we can prove our main theorem:

**Lemma 6.** *Presume (C1-C3) holds. Let  $P$  be a simple osculating cycle. Then there exists a directed walk  $W$  in  $G$  that starts at a vertex  $v \in P$  with a left outgoing edge  $e_1 \notin P$ , ends at a vertex  $w \in P$  with a right incoming edge  $e_k \notin P$ , and has no transverse intersection or shared edges with  $P$ .*

*Proof.* Since  $P$  is a directed cycle, it is non-contractible by (C3) and does not bound a face by (C2). Therefore, there must be some edge on the left side of  $P$  that shares a vertex  $v$  with  $P$ , but does not belong to  $P$ . If, say, the left incoming edge of  $v$  is not in  $P$ , then  $P$  uses the right incoming and right outgoing edge at  $v$  (because  $P$  follows rotationally consecutive edges), and so the left outgoing edge of  $v$  is also not in  $P$ . Therefore edge  $e_1$  exists.




Now create a leftmost walk  $W_\ell$  starting with  $e_1$ , then taking the left outgoing edge at its head, and continue taking left outgoing edges until one of the following happens:

1. We reach a vertex  $w \in P$  via a right incoming edge. In this case we have found the required path  $W$  and stop. Since we stop as soon as the head vertex  $w$  of a right incoming edge  $e_k$  of  $W_\ell$  is also a vertex of  $P$ , we can conclude that, before arriving at  $w$ , if we visited any other vertices in  $P$ , then these vertices were reached from a left incoming edge,  $e_i, 1 \leq i < k$ . Since  $W_\ell$  always follows the left outgoing edge and  $P$  always follows rotationally consecutive edges, we can conclude that the left incoming edge  $e_i$  and left outgoing edge  $e_{i+1}$  were not edges in  $P$  and the meeting of  $W_\ell$  and  $P$  was an osculating intersection rather than a transverse intersection.
2.  $W_\ell$  repeats a vertex, and the resulting directed cycle  $C$  does not contain vertices of  $P$ . Then  $\hat{i}(C, P) = 0$ , which contradicts Lemma 7. This case cannot happen.
3.  $W_\ell$  repeats a vertex, and the resulting directed cycle  $C$  contains vertices of  $P$ . Since we did not stop with Case 1, all such vertices of  $C$  reach  $P$  from the left. Following the same argument used in Case 1, all meetings of  $C$  with  $P$  are osculating resulting in  $\hat{i}(C, P) = 0$ . This again contradicts Lemma 7 so this case cannot happen.

The graph is finite and neither Case 2 nor 3 can happen, therefore Case 1 eventually must happen and we have found  $W$ .

## D An example of the meridian and longitude

Figure 8 illustrates how the graph  $G$  (of Figure 3(b)) is converted into a rectangular representation. We consider two different choices for  $L$ . Note that while both of them are perfectly fine longitudes, the left one is more visually appealing, principally because  $L$  and the (unique) osculating circuit  $P$  have significantly fewer crossings. One way to minimize the number of times  $L$  crosses  $G$  is to assign a weight to the edges of  $\mathcal{L}(G)$  according to the number of endpoints an edge has on  $G$  and finding a path with the least weight.



**Fig. 8.** Find rectangular representation. (a) Shortest path  $L$ . (b) Alternate  $L$ . (c) Draw graph with polygonal schema defined in (a). (d) Draw graph with polygonal schema defined in (b).