Leray's fundamental work on the Navier–Stokes equations: a modern review of "Sur le mouvement d'un liquide visqueux emplissant l'espace"

Wojciech S. Ożański, Benjamin C. Pooley

September 1, 2017

Abstract

This article offers a modern perspective which exposes the many contributions of Leray in his celebrated work on the Navier-Stokes equations from 1934. Although the importance of his work is widely acknowledged, the precise contents of his paper are perhaps less well known. The purpose of this article is to fill this gap. We follow Leray's results in detail: we prove local existence of strong solutions starting from divergence-free initial data that is either smooth, or belongs to $H^1, L^2 \cap L^p$ (with $p \in (3, \infty]$), as well as lower bounds on the norms $\|\nabla u(t)\|_2$, $\|u(t)\|_p$ $(p \in (3,\infty])$ as t approaches a putative blow-up time. We show global existence of a weak solution and weak-strong uniqueness. We present Leray's characterisation of the set of singular times for the weak solution, from which we deduce that its upper box-counting dimension is at most $\frac{1}{2}$. Throughout the text we provide additional details and clarifications for the modern reader and we expand on all ideas left implicit in the original work, some of which we have not found in the literature. We use some modern mathematical tools to bypass some technical details in Leray's work, and thus expose the elegance of his approach.

1 Introduction

The Navier–Stokes equations,

$$\partial_t u + (u \cdot \nabla)u - \nu \Delta u + \nabla p = 0,$$

 $\nabla \cdot u = 0,$

where u denotes the velocity of a fluid, p denotes the scalar pressure and $\nu > 0$ denotes viscosity of the fluid, comprise the fundamental model for the flow of an incompressible viscous fluid. They are named in recognition of Claude-Louis Navier (1822) and George Stokes (1845) who first formulated them, and they form the basis for many models in engineering and

mathematical fluid mechanics. The equations have been studied extensively and a number of excellent textbooks on the subject are available, see for instance Constantin (2008), Constantin & Foias (1988), Lemarié-Rieusset (2002), Robinson, Rodrigo & Sadowski (2016), Sohr (2001) and Temam (2001). However, the fundamental issue of the well-posedness of the equations in three dimensions remains unsolved.

In this article we focus solely on the work of Jean Leray (1934b), which to this day remains of fundamental importance in the study of the Navier– Stokes equations. Leray was the first to study the Navier–Stokes equations in the context of *weak solutions*. It is remarkable that such solutions are defined using a distributional form of the equations while the distribution theory was only formally introduced later by Schwartz (1950).

Many of the ideas in the modern treatment of these equations and a number of other systems originate from Leray's (1934) paper (which we shall often refer to simply as "Leray's work" or "Leray's paper"). The importance of that work is witnessed by the fact that it is one of the most cited works in mathematical fluid mechanics.

Leray studied the Navier–Stokes equations on the whole space (\mathbb{R}^3). Unlike later authors who have largely adopted Faedo-Galerkin techniques (see Hopf (1951) and Kiselev & Ladyzhenskaya (1957) for early examples), a characteristic of earlier works, including Leray's, is the use of explicit kernels. In this instance, Leray applied the Oseen kernel (herein denoted by \mathcal{T}), as derived by Oseen (1911), to obtain solutions of the *inhomogeneous* Stokes equations:

$$\partial_t u - \nu \Delta u + \nabla p = F, \qquad \nabla \cdot u = 0,$$

for a given forcing F. At the time these were also known as the equations for *infinitely slow motion*.

Oseen had previously applied this kernel iteratively to prove local wellposedness of the Navier–Stokes equations in $C^2(\mathbb{R}^3)$ (with bounded velocity, decay conditions on $\omega = \nabla \times u$, and polynomial growth estimates on $\nabla \omega$ and ∇u ; see Section 3.8 of Lemarié-Rieusset (2016) for a more complete discussion of Oseen's contributions). Leray applied a more elegant iteration scheme (a Picard iteration) to prove existence, uniqueness, and regularity results for local-in-time strong solutions for initial data $u_0 \in H^1 \cap L^{\infty} \cap C^1$ (Leray used the term *regular solutions*). We will see that in fact, $u_0 \in L^2 \cap$ L^{∞} suffices to construct strong solutions when his arguments are rewritten using a distributional form of equations.

Leray then derives lower bounds on various norms of the strong solution u(t) as t approaches the maximal time of existence T_0 if T_0 is finite, which indicate the rate of blow-up of a strong solution if such a blow-up occurs. He leaves open the issue of existence of blow-ups.

Next, Leray considers a generalised notion of solution of the Navier– Stokes equations, *weak solutions*. For this he studies the so-called *regularised* equations, which are obtained by replacing the nonlinear term $(u \cdot \nabla)u$ by $(J_{\varepsilon}u \cdot \nabla)u$ (where J_{ε} is the standard mollification operator). He shows that the regularised equations admit local well-posedness results similar to those for strong solutions of the Navier–Stokes equations, with an additional global-in-time estimate on the L^{∞} norm of the velocity. This extra property results in global-in-time existence and uniqueness of solutions u_{ε} for each $\varepsilon > 0$. By a careful compactness argument, he constructs a sequence of solutions $\{u_{\varepsilon_n}\}$ converging to a global-in-time weak solution to the Navier–Stokes equations. Such solutions, which he termed *turbulent solutions*, can be thought of as weak continuations of the strong solution beyond the blow-up time, a revolutionary idea at the time. He then shows that these weak solutions are strong locally-in-time except on a certain compact set of singular times with Lebesgue measure zero. To this end he uses a certain *weak-strong uniqueness* property.

We now briefly highlight a few important developments that proceeded from Leray's work. Eberhard Hopf (1951) performed a study of the Navier-Stokes equations on the bounded domain $\Omega \subset \mathbb{R}^3$, and proved global-in-time existence of weak solutions. Then Ladyzhenskaya (1959) proved existence and uniqueness of global-in-time strong solutions to the Navier–Stokes equations in bounded two-dimensional domains (Leray (1933) dealt with wellposedness in \mathbb{R}^2 in his thesis but was less successful in studying the case of bounded domains Leray (1934a)). Fujita & Kato (1964) used fractional powers of operators and the theory of semigroups to construct local-intime strong solutions of the (three-dimensional) Navier–Stokes equations on bounded domains (Kato (1984) used similar methods in the case of unbounded domains). As for the smoothness of weak solutions, it follows from the work of Serrin (1962), Prodi (1959), and Ladyženskaja (1967), that if a weak solution u belongs to $L^r((a,b); L^s)$ with $2/r+3/s \le 1$, s > 3 then u is a strong solution on the time interval (a, b); the critical case $r = \infty$, s = 3 was proved by Escauriaza et al. (2003). In addition, Beale et al. (1984) showed that if curl $u \in L^1((a, b); L^\infty)$ then u is a strong solution on the time interval (a,b].

Scheffer (1977) was the first to study the size of the singular set in both time and in space. Subsequently, Caffarelli, Kohn & Nirenberg (1982) proved that the one-dimensional Hausdorff measure of the singular set is zero. We refer the reader to the textbooks above for a wider description of the contributions to the theory of the Navier–Stokes equations in the last 80 years.

It is remarkable that despite many significant contributions, it is still not known whether the (unique) local-in-time strong solutions to the Navier– Stokes equations develop blow-ups or whether the global-in-time weak solutions are unique. This remains one of the most important open problems in mathematics, at the turn of the millennium, it was announced as one of seven Millennium Problems, see Fefferman (2006). A number of concepts and methods that found early use in Leray's work are now ubiquitous in the analysis of PDEs. These include: weak compactness of bounded sequences in L^2 , the concept of weak derivatives (called *quasi-derivatives* by Leray), the mollification operation and the fact that a weakly convergent sequence converges strongly if and only if the norm of the limit is the limit of the norms. He made an extensive use of the space of L^2 functions with weak derivatives in L^2 two years before the celebrated work of Sobolev (1936). This space would later be called Sobolev space H^1 . Furthermore, Leray was the first to introduce the compactness method of solving partial differential equations (see the proof of Theorem 4.7); in fact this, together with his work with Juliusz Schauder opened a new branch in mathematics, the use of topological methods in PDEs.

The terms Leray weak solution (or Leray-Hopf weak solution) of the Navier–Stokes equations (see Definition 4.5), Leray regularisation (see (4.1)) and Leray projection (for the projection of L^2 onto the space of weakly divergence-free vector fields) have become part of the mathematical lexicon in recognition of his seminal paper on the Navier–Stokes equations. We refer the interested reader to Lax (1998) for a broader description of Leray's work in the field of PDEs.

This article arose from series of lectures presented by the authors for a fluid mechanics reading group organised at the University of Warwick by James Robinson and José Rodrigo, and its purpose is to offer a modern exposition of Leray's work. We update the notation and we simplify some technical details by applying some modern methods; in particular we use the Fourier transform (see Theorem 2.2) and the distributional forms of the partial differential equations appearing in Leray's work. It is perhaps remarkable that these updates do not detract from the originality of Leray's arguments; rather they make them even more elegant.

We have also endeavoured to give a rigorous account of all non-trivial results that are left implicit in the original work, some of which we were not able to find in the literature. These include Leray's derivations of the blow-up rate of the norm $||u(t)||_p$ (with p > 3) of a strong solution u as t approaches the putative blow-up time (see Corollary 3.12), and a result on local existence of strong solutions for initial data $u_0 \in L^2 \cap L^p$ (with $p \in (3,\infty]$) that is weakly divergence free (see Corollary 3.18). In order to make the exposition self-contained we have also added appendices on relevant facts from the theory of the heat equation, integral inqualities, the Volterra equation and other topics.

For simplicity of notation we focus only on the case $\nu = 1$. The corresponding results for any $\nu > 0$ can be recovered using the following rescaling argument: if u, p is a solution of the Navier–Stokes equations with $\nu = 1$, then $\tilde{u}(x,t) := u(x\nu, t\nu)$, $\tilde{p} := p(x\nu, t\nu)$ is a solution for given $\nu > 0$.

The structure of the article is as follows. In the remainder of this section we describe some notation, we recall some preliminary results, and we introduce the Oseen kernel \mathcal{T} , which will be the main tool for solving the Stokes equations.

In Section 2 we study the Stokes equations. We first show that if the forcing F is sufficiently smooth (see (2.9)) the equations can be solved classically using the representation formulae

$$u(t) \coloneqq \Phi(t) * u_0 + \int_0^t \mathcal{T}(t-s) * F(s) \, \mathrm{d}s,$$

$$p(t) \coloneqq -(-\Delta)^{-1}(\operatorname{div} F(t)),$$

see (2.3), (2.4), where $\Phi(t)$ denotes the heat kernel (Theorem 2.2). We also show some further properties of the representation formula for u in the case of less regular forcing F (Lemma 2.1). We then focus on a special form of the forcing

$$F = -(Y \cdot \nabla)Y,$$

see (2.12), which is reminiscent of the nonlinear term in the Navier–Stokes equations. This special form of F gives rise to the modified representation formulae,

$$u(t) \coloneqq \Phi(t) * u_0 + \int_0^t \nabla \mathcal{T}(t-s) * [Y(s)Y(s)] \, \mathrm{d}s,$$
$$p(t) \coloneqq \partial_k \partial_i (-\Delta)^{-1} (Y_i(t)Y_k(t)),$$

see (2.14), (2.15). After deducing some properties of this modified representation formula (Lemma 2.3) we show that it gives a unique solution (in some wide sense) to the Stokes equations with the forcing F of the form above, in the sense of distributions (Theorem 2.5 and Theorem 2.6).

In Section 3 we study strong solutions of the Navier–Stokes equations. After defining strong solutions on an open time interval (0,T) (Definition 3.1) we use the theory of the Stokes equations developed in Section 2 to deduce the smoothness of such solutions (Corollary 3.3), as well as other interesting properties, such as the energy equality

$$||u(t_2)||^2 + 2 \int_{t_1}^{t_2} ||\nabla u(s)||^2 ds = ||u(t_1)||^2,$$

(Theorem 3.4) and the comparison of strong solutions

$$\|(u-v)(t_2)\|^2 \le \|(u-v)(t_1)\|^2 e^{\frac{1}{2}\int_{t_1}^{t_2} \|u(s)\|_{\infty}^2 \, \mathrm{d}s}, \quad t_1 < t_2$$

(Lemma 3.5). We then define strong solutions on a half-closed time interval [0,T) (Definition 3.7) and show local-in-time existence and uniqueness of such solutions with weakly divergence-free initial data $u_0 \in L^2 \cap L^\infty$ (Theorem 3.9). Next, we discuss the issue of the maximal time of existence T_0 of strong solutions (Lemma 3.10 and Corollary 3.11), from which we deduce

the rates of blow-up of u(t) in various norms as t approaches T_0 (if T_0 is finite):

$$||u(t)||_{\infty} \ge \frac{C}{\sqrt{T_0 - t}}, \quad ||\nabla u(t)|| \ge \frac{C}{(T_0 - t)^{1/4}},$$

and

$$||u(t)||_p \ge \frac{C^{(1-3/p)/2}(1-3/p)}{(T_0-t)^{(1-3/p)/2}}$$

(Corollary 3.12). This study of strong solutions is concluded with an observation that less regular initial data u_0 also gives rise to a unique strong solution on the time interval (0,T) for some T > 0. This motivates the definition of *semi-strong solutions* (Definition 3.15); we show that if $u_0 \in L^2$ is weakly divergence free, and either

$$\nabla u_0 \in L^2$$
 or $u_0 \in L^p$ (with $p > 3$),

then there exists a unique local-in-time semi-strong solution starting from initial data u_0 (Theorem 3.17 and Corollary 3.18).

In Section 4 we study weak solutions of the Navier–Stokes equations. To this end, for each $\varepsilon > 0$ we consider regularised equations, where a mollification operator is applied in the nonlinear term,

$$\partial_t u - \Delta u + ((J_\varepsilon u) \cdot \nabla)u + \nabla p = 0$$

(Definition 4.1). We show that for each $\varepsilon > 0$ the regularised equations can be analysed in a similar way to strong solutions of the Navier–Stokes equations in Section 3, the difference being that the maximal time of existence of the solution u_{ε} is infinite (Theorem 4.2). In order to take the limit $\varepsilon \to 0^+$ we first show that the kinetic energy of $u_{\varepsilon}(t)$ outside a ball can be estimated independently of ε ,

$$\int_{|x|>R_2} |u_{\varepsilon}(t)|^2 \, \mathrm{d}x \le \int_{|x|>R_1} |u_0|^2 \, \mathrm{d}x + \frac{C(u_0, t)}{R_2 - R_1}, \qquad R_2 > R_2 > 0,$$

(Lemma 4.3). Thanks to this separation of energy result we can let $\varepsilon \rightarrow 0^+$ (along a carefully chosen subsequence) to obtain a global-in-time weak solution of the Navier–Stokes equations (Theorem 4.7). We then show the so-called weak-strong uniqueness result (Lemma 4.9) and we deduce that the weak solution admits a particular structure, namely that it is (locally) a strong solution at times

$$t \in \bigcup_i (a_i, b_i)$$

where the intervals $(a_i, b_i) \subset (0, \infty)$ are pairwise disjoint (Theorem 4.11). Finally we show that the complement of their union

$$\Sigma \coloneqq (0,\infty) \setminus \bigcup_i (a_i, b_i),$$

(the set of putative singular times) is bounded, its box-counting dimension is bounded above by 1/2 and that the weak solution admits certain decay for sufficiently large times (Theorem 4.12 and Corollary 4.13).

Unless specified otherwise, each proof follows Leray (1934b), possibly with minor modifications. We also comment on Leray's methodology throughout the text, in footnotes and in the "Notes" at the end of each section. Equation numbers marked in italics correspond to expressions in Leray's paper.

1.1 Preliminaries

The letter C denotes a numerical constant, whose value may change at each occurrence. Occasionally we write C' (or C'') to denote a constant that has the same value wherever it appears within a given section. Also, C_m (and c_m) denotes a numerical constant for each m. Throughout the article (unless specified otherwise) we consider function spaces on \mathbb{R}^3 , for example $L^p := L^p(\mathbb{R}^3), H^m := H^m(\mathbb{R}^3), C_0^\infty := C_0^\infty(\mathbb{R}^3)$. We also define $\int := \int_{\mathbb{R}^3}, \|\cdot\|_p := \|\cdot\|_{L^p}$ and we reserve the notation $\|\cdot\|$ for the L^2 norm. We say that a velocity field v is *weakly divergence free* if div v = 0 in the sense of distributions, that is

$$\int v \cdot \nabla g = 0 \tag{1.1}$$

for all $g \in C_0^{\infty}$, and we set

$$H \coloneqq \{ f \in L^2 : f \text{ is weakly divergence free} \}.$$

$$V \coloneqq \{ f \in H^1 : \operatorname{div} f = 0 \}.$$
(1.2)

We understand \mathbb{Q}^+ as the nonnegative rational numbers and we define $\partial_j := \partial/\partial x_j$ and $\nabla^m := D^m$, where we understand all derivatives in the weak sense. We use the convention of summing over repeated indices. For example, we write $v_j \partial_j u_i$ to denote the vector $(v \cdot \nabla)u$ (here i = 1, 2, 3). For an interval I define

$$\mathcal{H}^{1/2}(I) \coloneqq \{f : \mathbb{R}^3 \times I : \exists C(t), \text{ a continuous function on } I, \text{ such that} \\ |f(x,t) - f(y,t)| \le C(t)|x - y|^{1/2} \text{ for all } t \in I, x, y \in \mathbb{R}^3\},$$

that is $\mathcal{H}^{1/2}(I)$ is the space of functions such that $||f(t)||_{C^{0,1/2}} \leq C(t)$ for some continuous C(t) (where $|| \cdot ||_{C^{0,1/2}}$ denotes 1/2-Hölder seminorm). Note that $\mathcal{H}^{1/2}(I)$ is defined in the same way for vector-valued functions.

We recall the integral Minkowski inequality

$$\|f(t)\|_{p} \leq \int_{0}^{t} \|g(s)\|_{p} \,\mathrm{d}s.$$
(1.3)

whenever $p \in [1, \infty]$ and f(x, t) is of the form $f(x, t) = \int_0^t g(x, s) ds$. More generally, for integrable and nonnegative ξ , η , if

$$F(x,t) = \int_{\mathbb{R}} \xi(s) \int \eta(y) f(x,y,t,s) \, \mathrm{d}y \, \mathrm{d}s$$

then

$$|F(t)||_p \le \int_{\mathbb{R}} \xi(s) \int \eta(y) ||f(\cdot, y, t, s)||_p \,\mathrm{d}y \,\mathrm{d}s.$$
(1.4)

Now let p, q, r > 0 satisfy 1/q = 1/p + 1/r - 1 and let $f \in L^p, g \in L^r$. Then

$$||f * g||_q \le ||f||_p ||g||_r.$$
(1.5)

This is Young's inequality for convolutions (see e.g. Stein & Weiss (1971), p. 178, for the proof). Here "*" denotes the convolution, that is

$$(f * g)(x) \coloneqq \int f(x - y)g(y) \, \mathrm{d}y.$$

If f and g are also functions of time t we omit x and simply write u(t) = f(t)*g(t). We apply this notation in the statement of the following extension of Young's inequality to the case of space-time convolutions, whose proof we give in Appendix A.2 (see Lemma A.3).

Lemma 1.1. If $p, q, r \ge 1$ satisfy

$$\frac{1}{q} = \frac{1}{p} + \frac{1}{r} - 1,$$

 $A \in L^1_{\text{loc}}([0,T);L^p)$ and $B \in C((0,T);L^r)$ with $||B(t)||_r$ bounded as $t \to 0^+$ then u defined by

$$u(t) \coloneqq \int_0^t A(t-s) * B(s) \,\mathrm{d}s$$

belongs to $C([0,T);L^q)$ and

$$||u(t)||_q \le \int_0^t ||A(t-s)||_p ||B(s)||_r \,\mathrm{d}s.$$

Let J_{ε} denote the standard *mollification* operator, that is

$$J_{\varepsilon}v \coloneqq \eta_{\varepsilon} * v,$$

where η_{ε} is a standard mollifier, e.g. $\eta_{\varepsilon}(x) \coloneqq \varepsilon^{-3}\eta(x/\varepsilon)$, where $\eta(x) \coloneqq C \exp((|x|^2 - 1)^{-1})$ for |x| < 1 and $\eta(x) \coloneqq 0$ for $|x| \ge 1$ with the constant C > 0 chosen such that $\int \eta = 1$.

Lemma 1.2 (properties of mollification). The mollification operator J_{ε} (on \mathbb{R}^3) enjoys the following properties:

- (i) $||J_{\varepsilon}v||_p \le ||v||_p$ for all $p \in [1, \infty]$, $\varepsilon > 0$,
- (ii) $\partial_k J_{\varepsilon} v = J_{\varepsilon}(\partial_k v)$ for every k = 1, 2, 3 whenever $\nabla v \in L^1_{\text{loc}}$,
- (iii) $||J_{\varepsilon}v||_{\infty} \leq C\varepsilon^{-3/2}||v||,$
- (iv) if $v \in L^2$ then $J_{\varepsilon}v \in H^m$ for all m with $\|J_{\varepsilon}v\|_{H^m} \leq C_m \varepsilon^{-m} \|v\|$,
- (v) if $v \in L^1_{loc}$ then $J_{\varepsilon}v \to v$ almost everywhere as $\varepsilon \to 0$,
- (vi) if $v \in L^p$, where $p \in [1, \infty)$, then $J_{\varepsilon}v \to v$ in L^p as $\varepsilon \to 0$.

The proofs of the above properties are elementary (and can be found in Appendix C in Evans (2010), Section 3.5.2 in Majda & Bertozzi (2002) or Appendix A.3 in Robinson et al. (2016)).

For $f \in L^2$ define

$$(-\Delta)^{-1}f(x) \coloneqq \int \frac{f(y)}{4\pi |x-y|} \,\mathrm{d}y. \tag{1.6}$$

The symbol $(-\Delta)^{-1}$ relates to the fact that $g := (-\Delta)^{-1}f$ satisfies the Poisson equation $-\Delta g = f$ in \mathbb{R}^3 in the sense of distributions (which follows by an application of Fubini's theorem). Since we will often estimate terms similar to the right-hand side of (1.6), we formulate the following lemma (which is Leray's (1.14)).

Lemma 1.3. If $f \in H^1$ then for any $y \in \mathbb{R}^3$

$$\int \frac{|f(x)|^2}{|x-y|^2} \,\mathrm{d}x \le 4 \|\nabla f\|^2.$$
(1.7)

Proof. It is enough to prove the claim when y = 0 and when f is a scalar function. If $f \in C_0^{\infty}$ then integration by parts, the Cauchy–Schwarz inequality and Young's inequality give

$$\int \frac{(f(x))^2}{|x|^2} \, \mathrm{d}x = -\int \frac{x}{|x|^2} \cdot \nabla (f(x))^2 \, \mathrm{d}x = -2\int \frac{x}{|x|^2} \cdot \nabla f(x) f(x) \, \mathrm{d}x$$
$$\leq 2\|\nabla f\| \sqrt{\int \frac{(f(x))^2}{|x|^2} \, \mathrm{d}x} \leq 2\|\nabla f\|^2 + \frac{1}{2}\int \frac{(f(x))^2}{|x|^2} \, \mathrm{d}x.$$

For $f \in H^1$ the claim follows from the density of C_0^{∞} functions in H^1 and Fatou's lemma.

Lemma 1.4 (The Plancherel Lemma). The operator

$$f \mapsto \partial_i \partial_k (-\Delta)^{-1} f$$

is a bounded operator from L^2 to L^2 for every *i*, *k*. Consequently an application of Fubini's theorem gives that $\partial_i \partial_k (-\Delta)^{-1}$ is a bounded operator from H^m to H^m for every $m \ge 0$. *Proof.* This follows by considering the Fourier transform and using the Plancherel property:

$$\|\partial_i \partial_k (-\Delta)^{-1} f\|^2 = \int \left| \frac{\xi_i \xi_k}{|\xi|^2} \widehat{f}(\xi) \right|^2 \mathrm{d}\xi \le \int \left| \widehat{f}(\xi) \right|^2 \mathrm{d}\xi = \|f\|^2,$$

where \hat{f} denotes the Fourier transform of f.

Consider the heat equation in $\mathbb{R}^3 \times (0, T)$:

$$v_t - \Delta v = 0$$

with initial condition $v(0) = v_0$ (understood in the sense of L^2 limit as $t \to 0^+$) for some $v_0 \in L^2$. Then a classical solution v of the heat equation is given by the convolution

$$v(t) = \Phi(t) * v_0,$$

where

$$\Phi(x,t) := \frac{1}{(4\pi t)^{3/2}} e^{-|x|^2/4t}$$
(1.8)

is the *heat kernel*. In what follows, we will rely on some well-known properties of the heat equation and the heat kernel, which we discuss in Appendix A.1.

Finally, we will often use an integral version of an elementary fact from the theory of ordinary differential equations: if $f, \phi \colon [0,T) \to \mathbb{R}^+$ are C^1 functions such that

$$\begin{cases} f' \le g f^k + a, \\ \phi' \ge g \phi^k + b \end{cases} \quad \text{on } [0, T) \qquad \text{with} \quad f(0) < \phi(0), \end{cases}$$

where k > 0 and g, a, b are continuous functions such that g > 0 and $a \le b$, then $f < \phi$ on [0, T). In particular, we will use the following result, which corresponds to the case k = 2.

Lemma 1.5 (Integral inequalities). Suppose g > 0 is a continuous function on (0,T) that is locally integrable [0,T), that functions $f, \phi : (0,T) \to \mathbb{R}^+$ satisfy

$$f(t) \leq \int_0^t g(t-s)f(s)^2 \,\mathrm{d}s + a(t),$$

$$\phi(t) \geq \int_0^t g(t-s)\phi(s)^2 \,\mathrm{d}s + b(t)$$

for all $t \in (0,T)$, where a, b are continuous functions satisfying $a \leq b$, ϕ is continuous, and that f^2 and ϕ^2 are integrable near 0. Then $f \leq \phi$ on (0,T).

Note that no assumption on the continuity of f is needed. We prove this lemma, along with a few related results, in Appendix A.5 (see Lemma A.5).

1.2 The Oseen kernel \mathcal{T}

The Oseen kernel is the main tool in solving the Stokes equations in \mathbb{R}^3 (which we discuss in the next section; see also the comment following Theorem 2.2). It is a 3×3 matrix-valued function $\mathcal{T} = [\mathcal{T}_{ij}]$ defined by

$$\mathcal{T}_{ij}(x,t) \coloneqq \delta_{ij} \Phi(x,t) + \partial_i \partial_j P(x,t), \qquad x \in \mathbb{R}^3, t > 0, \tag{1.9}$$

where δ_{ij} denotes the Kronecker delta and

$$P(x,t) \coloneqq \frac{1}{4\pi^{3/2}t^{1/2}|x|} \int_0^{|x|} e^{-\xi^2/4t} \,\mathrm{d}\xi.$$
(1.10)

It was first introduced by Oseen (1911) (see pages 3, 19 and 41 therein)

Note that $P(\cdot, t)$ is a smooth function for each t > 0. Indeed, for fixed t the function $\tilde{P} : \mathbb{R} \to \mathbb{R}$ defined by $\tilde{P}(0) := 1$,

$$\tilde{P}(s) := \frac{1}{4\pi^{3/2}s} \int_0^s \frac{\mathrm{e}^{-r^2/4t}}{t^{1/2}} \,\mathrm{d}r$$

is even and smooth (one can verify that \tilde{P} , $\frac{d\tilde{P}}{ds}$ are continuous and that $\frac{d^2\tilde{P}}{ds^2} = e^{-s^2/4t}/(4\pi t)^{3/2}$, a smooth function). Therefore, since $P(x,t) = \tilde{P}(|x|)$, $P(\cdot,t)$ is smooth as well (for each t > 0). A direct computation shows that

$$-\Delta P = \Phi. \tag{1.11}$$

This yields an equivalent definition of \mathcal{T}_{ij} :

$$\begin{aligned} \mathcal{T}_{1,1} &= -(\partial_2^2 + \partial_3^2)P, & \mathcal{T}_{1,2} &= \partial_1 \partial_2 P, & \mathcal{T}_{1,3} &= \partial_1 \partial_3 P, \\ \mathcal{T}_{2,1} &= \partial_1 \partial_2 P, & \mathcal{T}_{2,2} &= -(\partial_1^2 + \partial_3^2)P, & \mathcal{T}_{2,3} &= \partial_2 \partial_3 P, \\ \mathcal{T}_{3,1} &= \partial_1 \partial_3 P, & \mathcal{T}_{3,2} &= \partial_2 \partial_3 P, & \mathcal{T}_{3,3} &= -(\partial_1^2 + \partial_2^2)P. \end{aligned}$$

$$\end{aligned}$$

$$(1.12)$$

Since the derivatives $\nabla^m P$ satisfy the pointwise estimate

$$|\nabla^m P(x,t)| \le \frac{C_m}{(|x|^2 + t)^{(m+1)/2}}, \qquad m \ge 0$$

(see Theorem A.4), we obtain the following pointwise estimates on the Oseen kernel

$$|\nabla^m \mathcal{T}(x,t)| \le \frac{C_m}{(|x|^2 + t)^{(m+3)/2}} \qquad \text{for } x \in \mathbb{R}^3, t > 0, m \ge 0.$$
(1.13)

Using these bounds we can easily deduce the integral estimates

$$\|\mathcal{T}(t)\| \le C t^{-3/4}$$
 and $\|\nabla \mathcal{T}(t)\|_1 \le C t^{-1/2}$ (1.14)

for t > 0, where we used the facts $\int (|x|^2 + t)^{-3} dx = C/t^{3/2}$ and $\int (|x|^2 + t)^{-2} dx = C/t^{1/2}$. Similarly, by (1.13) and an application of the Dominated Convergence Theorem we obtain

$$\mathcal{T} \in C((0,\infty); L^2)$$
 and $\nabla \mathcal{T} \in C((0,\infty); L^1).$ (1.15)

Finally \mathcal{T} enjoys a certain integral continuity property, which we will use later to show Hölder continuity of the solution to Stokes equations (see Lemma 2.3 (ii)).

Lemma 1.6 (1/2-Hölder continuity of $\nabla \mathcal{T}$ in an L^1 sense). For $x, y \in \mathbb{R}^3$, t > 0

$$\int |\nabla \mathcal{T}(x-z,t) - \nabla \mathcal{T}(y-z,t)| \,\mathrm{d}z \le C|x-y|^{1/2}t^{-3/4}.$$

Leray mentions this inequality on page 213, and he frequently uses it in his arguments (in (2.14), (2.18), the inequality following (3.3), and the first inequality on page 219) to show Hölder continuity of functions given by representation formulae involving ∇T . We provide a proof for the sake of completeness.

Proof. Let $R \coloneqq |x - y|$ and

$$\Omega \coloneqq B(x, 2R) \cup B(y, 2R).$$

Since $\Omega \subset B(x, 3R)$ we can use (1.13) to write

$$\int_{\Omega} |\nabla \mathcal{T}(x-z,t)| \, \mathrm{d}z$$

$$\leq C \int_{B(x,3R)} \frac{\mathrm{d}z}{(|x-z|^2+t)^2} = C \int_{0}^{3R} \frac{r^2}{(r^2+t)^2} \, \mathrm{d}r \qquad (1.16)$$

$$\leq C \int_{0}^{3R} \frac{r^2+t}{(r^2+t)^2} \, \mathrm{d}r = \frac{C}{t^{1/2}} \tan^{-1}\left(\frac{3R}{t^{1/2}}\right) \leq C \frac{R^{1/2}}{t^{3/4}},$$

since $\tan^{-1} \alpha \leq \alpha^{1/2}$ for $\alpha > 0$. Analogously

$$\int_{\Omega} |\nabla \mathcal{T}(y - z, t)| \, \mathrm{d}z \le C R^{1/2} t^{-3/4}.$$
(1.17)

As for $z \in \mathbb{R}^3 \setminus \Omega$ note that $|z - x| \ge 2R$. Hence for any point ξ from the line segment [x, y]

$$|z - x| \le |z - \xi| + |\xi - x| \le |z - \xi| + R \le |z - \xi| + |z - x|/2,$$

and so $|z - \xi| \ge |z - x|/2$. Thus, using the Mean Value Theorem and the bound on $\nabla^2 \mathcal{T}$ (see (1.13)) we obtain

where $\xi(z) \in [x, y]$ for each z. This together with (1.16) and (1.17) proves the claim.

2 The Stokes equations

In this section we consider the Stokes equations,

$$\partial_t u - \Delta u + \nabla p = F, \tag{2.1}$$

$$\operatorname{div} u = 0 \qquad \text{in } \mathbb{R}^3 \times (0, T), \qquad (2.2)$$

where T > 0 and F(x, t) is a vector-valued forcing. Leray calls these equations the *infinitely slow motion*. The Stokes equations model a drift-diffusion flow of a incompressible velocity field u. Here p denotes the pressure function. One can think of the appearance of the pressure function as providing the extra freedom necessary to impose the incompressibility constraint (2.2) for an arbitrary F, see the comment after Theorem 2.2. As usual, we denote the initial condition for (2.1), (2.2) by $u_0 \in H$, which is understood in the sense of the L^2 limit, that is $||u(t) - u_0|| \to 0$ as $t \to 0^+$.

In his paper, Leray includes an essentially complete analysis of the Stokes initial value problem in \mathbb{R}^3 . The results for this problem are fundamental in the analysis of the Navier–Stokes equations that follows; while the arguments are at times somewhat technical, we therefore present them in full, but with some details in Appendix A.4.

The Stokes equations with the general form of the forcing F can be solved using the representation formulae¹,

$$u(t) = u_1(t) + u_2(t) \coloneqq \Phi(t) * u_0 + \int_0^t \mathcal{T}(t-s) * F(s) \, \mathrm{d}s, \quad (2.3)$$

$$p(t) := -(-\Delta)^{-1}(\operatorname{div} X(t)), \qquad (2.4)$$

¹These formulae are stated by Leray in (2.2) and (2.9).

see Theorem 2.2 below (in which we focus only on the case of regular F). (See (1.6) for the definition of $(-\Delta)^{-1}$, and recall that $\Phi(t)$ denotes the heat kernel (1.8).) Here the convolution of the matrix function $\mathcal{T}(t-s)$ and a vector function F(s) is understood as a matrix-vector operation, that is

$$u_{2,i}(x,t) = \int_0^t \int \mathcal{T}_{ij}(x-y,t-s)F_j(y,s)\,\mathrm{d}y\,\mathrm{d}s.$$

In this section we study the representation formulae (2.3), (2.4) and certain modified representation formulae (that is (2.14), (2.15)) in the case when F is of the special form $F = -(Y \cdot \nabla)Y$ for some vector field Y. Of the two cases

F of general form and
$$F = -(Y \cdot \nabla)Y$$
 for some Y

Leray considers² mainly the former case; studying the formula (2.3) given appropriate regularity of F, and only mentioning briefly the latter case³. Here we treat the two cases separately. We treat the former case briefly, and we focus more on the latter case. An advantage of this approach is that it makes our results for each of the two cases directly applicable in the analysis of the Navier–Stokes equations. Moreover, in this slight refinement of Leray's approach, we construct the solution using the representation formulae, rather than deducing the representation as a property of the solution. As a result, we obtain a simple existence and uniqueness theorem for the Stokes equations (Theorem 2.6).

2.1 A general forcing F

Consider a forcing $F \in C([0,T); L^2)$ and let u, p be given by the representation formulae (2.3), (2.4) above.

Lemma 2.1. If $F \in C([0,T); L^2)$ then the function u defined above satisfies

(i) $u \in C((0,T); L^{\infty})$ with⁴,

$$\|u(t)\|_{\infty} \le C \int_0^t \frac{\|F(s)\|}{(t-s)^{3/4}} \,\mathrm{d}s + C \|u_0\|t^{-3/4}.$$
 (2.5)

(ii) $\nabla u \in C((0,T);L^2)$ with⁵

$$\|\nabla u(t)\| \le C \int_0^t \frac{\|F(s)\|}{(t-s)^{1/2}} \,\mathrm{d}s + C \|u_0\| t^{-1/2}.$$
 (2.6)

 $^{^{2}}$ This corresponds to Sections 11–13.

 $^{^{3}}$ See Lemma 8 in his work.

⁴Leray does not state this bound (we state it as a tool for proving (iii)).

⁵This is Leray's (2.8), (2.12) and (2.19).

More generally, if $F, \ldots, \nabla^m F \in C([0,T); L^2)$ then

$$\nabla^{m+1} u \in C((0,T);L^2)$$

with

$$\|\nabla^{m+1}u(t)\| \le C \int_0^t \frac{\|\nabla^m F(s)\|}{(t-s)^{1/2}} \,\mathrm{d}s + C_{m+1} \|u_0\| t^{-(m+1)/2}.$$

(iii) $u \in C([0,T); L^2)$ with⁶

$$||u(t)|| \le \int_0^t ||F(s)|| \,\mathrm{d}s + ||u_0|| \qquad \text{for all } t \in (0,T).$$
(2.7)

Moreover u satisfies the energy dissipation equality

$$||u(t)||^{2} - ||u_{0}||^{2} + 2\int_{0}^{t} ||\nabla u(s)||^{2} \,\mathrm{d}s = 2\int_{0}^{t} \int u \cdot F \,\mathrm{d}x \,\mathrm{d}s \qquad (2.8)$$

for all $t \in (0,T)$.

The properties (i), (ii) of the lemma follow from Lemma 1.1, the integral bounds on the Oseen kernel (see (1.14)) and the corresponding property of the heat kernel (see (ii), (iii) in Appendix A.1). As for (iii), assuming first that F is smooth, the functions u, p constitute a classical solution to the Stokes equations (2.1), (2.2) (which is proved in the following theorem). The required estimates are straightforward for classical solutions. If F is not smooth, one obtains (iii) by a density argument. See Appendix A.4.2 for the detailed proof of (iii).

Theorem 2.2 (Classical solution for smooth forcing F). Suppose that for some R > 0

$$F \in C^{\infty}(\mathbb{R}^3 \times [0,T);\mathbb{R}^3) \quad and \quad \operatorname{supp} F(t) \subset B(0,R) \text{ for } t \in [0,T).$$
(2.9)

Then the pair of functions u, p given above is a classical solution of the Stokes equations (2.1), (2.2) with $u(0) = u_0$. Moreover $u \in C([0,T); L^2)$.

In fact the functions u, p constitute a unique solution in a much wider class, namely the class of distributional solutions u, p such that $u \in C([0, T); L^2)$ and $p \in L^1_{\text{loc}}(\mathbb{R}^n \times [0, T))$, see Theorem 2.5 in Section 2.2.

Theorem 2.2 follows by showing that u_1 , u_2 satisfy the equations⁷

$$\begin{cases} \partial_t u_1 - \Delta u_1 = 0, \\ \operatorname{div} u_1 = 0, \\ u_1(0) = u_0, \end{cases} \qquad \begin{cases} \partial_t u_2 - \Delta u_2 + \nabla p = F, \\ \operatorname{div} u_2 = 0, \\ u_2(0) = 0, \end{cases}$$
(2.10)

⁷The study of u_1 and u_2 corresponds to Leray's Sections 11 and 12, respectively.

⁶This is Section 13 in Leray (1934b).

The part of the claim for u_1 follows directly from the analysis of the heat equation, see Appendix A.1. As for u_2 , using the Fourier transform one can see that it is enough to prove the claim in Fourier space. It therefore suffices to use the Fourier transform of the Oseen kernel,

$$\mathcal{F}[\mathcal{T}(t)] = \left(I - \frac{\xi \otimes \xi}{|\xi|^2}\right) e^{-4\pi^2 t |\xi|^2} \quad t > 0,$$
(2.11)

obtained from (1.9) and (1.11), where I denotes the identity matrix and $\xi \otimes \xi$ denotes the 3×3 matrix with components $\xi_i \xi_j$. An interested reader is referred to Appendix A.4.1 for the detailed proof.

At this point it is interesting to note that the Stokes equations are in fact a nonhomogeneous heat equation for u under the incompressibility constraint div u = 0. Since $\Delta p = \text{div } F$ we see that p appearing in the Stokes equations acts as a modification of the forcing F to make it divergence free (that is div $(F - \nabla p) = 0$). Since any solution of a nonhomogeneous heat equation with divergence-free forcing and initial data remains divergence free for positive times, we see that the role of p in the Stokes equations is to guarantee that u(t) remains divergence free for t > 0. One can also think of it as the projection of X(t) onto the space of weakly divergence-free vector fields (which is often called the *Leray projection*).

Moreover, from (2.4) we see that the Fourier transform of the modified forcing $F - \nabla p$ is

$$\left(I - \frac{\xi \otimes \xi}{|\xi|^2}\right) \widehat{F}(\xi, t).$$

Thus we see from (2.11) that the Oseen kernel is precisely the modification of the heat kernel that accounts for this modification of the forcing. This is particularly clear from a calculation in the Fourier space in Appendix A.4.1.

2.2 A forcing of the form $F = -(Y \cdot \nabla)Y$

Here we assume that F is of a particular form, namely

$$F = -(Y \cdot \nabla)Y \tag{2.12}$$

(in components $F_k = Y_i \partial_i Y_k$) for some weakly divergence-free $Y \in C((0, T), L^{\infty})$ such that $||Y(t)||_{\infty}$ remains bounded as $t \to 0^+$. Note that since the derivatives $\partial_i Y_k$ are not well-defined we understand (2.12) in a formal sense and we will consider the Stokes equations (2.1), (2.2) in the sense of distributions. More precisely, we say that u, p is a *distributional solution* of (2.1), (2.2) with F of the form (2.12) if u(t) is weakly divergence free for $t \in (0, T)$ and

$$\int u_0 \cdot \phi(0) \,\mathrm{d}x + \int_0^T \int \left(u \cdot (\phi_t + \Delta \phi) + p \operatorname{div} \phi \right) = \int_0^T \int Y \cdot (Y \cdot \nabla) \phi \quad (2.13)$$

for $\phi \in C_0^{\infty}(\mathbb{R}^3 \times [0,T);\mathbb{R}^3)$. We will also consider a modified form of the representation formulae (2.3), (2.4) that accounts for this special form of the forcing,

$$u(t) = u_1(t) + u_2(t) \coloneqq \Phi(t) * u_0 + \int_0^t \nabla \mathcal{T}(t-s) * [Y(s)Y(s)] \, \mathrm{d}s, \quad (2.14)$$

$$p(t) \coloneqq \partial_k \partial_i (-\Delta)^{-1} (Y_i(t) Y_k(t)), \qquad (2.15)$$

where we write

$$\left(\nabla \mathcal{T}(t-s) * [Y(s)Y(s)]\right)_j(x) \coloneqq \int \partial_i \mathcal{T}_{jk}(x-y,t-s)Y_i(y,s)Y_k(y,s)\,\mathrm{d}y.$$
(2.16)

Clearly, such u, p are well-defined since no derivatives fall on Y in these modified representation formulae. If Y is regular (in the sense of (2.9)) then the above definition of u, p is equivalent to (2.3), (2.4), and so Theorem 2.2 implies that such u, p constitute a classical solution of the Stokes equations (and hence also a distributional solution). In this section we show that u, p constructed above constitute the unique distributional solution in a wide class if we have $Y \in C([0,T); L^2)$ in addition to the assumptions on Ymentioned in (2.12), see Theorem 2.6 below.⁸

For this purpose we derive several properties of such u, p. In view of Lemma 2.1, we now prove refined bounds on $||u(t)||_{\infty}$ and ||u(t)||, and show that $u \in \mathcal{H}^{1/2}((0,T))$ and $\nabla u \in C((0,T); L^{\infty})$.

Lemma 2.3 (Properties of u, p given by (2.14-2.15)). Let $u_0 \in L^2$ and u, p given by (2.14) and (2.15) for some Y with the properties described following (2.12).

(i) If u_0 is bounded then $u \in C((0,T), L^{\infty})$ with⁹

$$||u(t)||_{\infty} \le C \int_0^t \frac{||Y(s)||_{\infty}^2}{\sqrt{t-s}} \,\mathrm{d}s + ||u_0||_{\infty}.$$

Moreover $u \in C([0,T), L^{\infty})$ if $u_0 \in L^{\infty}$ is uniformly continuous.

(ii) $u \in \mathcal{H}^{1/2}((0,T))$ and the corresponding Hölder constant $C_0(t)$ satisfies¹⁰

$$C_0(t) \le c_0 \int_0^t \frac{\|Y(s)\|_{\infty}^2}{(t-s)^{3/4}} \,\mathrm{d}s + c_0 \frac{\|u_0\|}{t}$$

⁸This corresponds to Leray's Lemma 8, in which he states that the representation formula (2.14) is a property of the solution.

⁹Leray does not state this bound explicitly, but he uses it in later sections during the study of the Navier–Stokes equations (for instance in (3.5), at the bottom of p. 222, and at the top of p. 232).

¹⁰Leray shows a similar property of ∇u in the case of F of general form (which he obtains in (2.18) and as a consequence of (2.7) and (2.8)). We translate this result to the case of F of the form $F = -(Y \cdot \nabla)Y$.

for some $c_0 > 0$.

More generally, if $Y, \nabla Y, \ldots, \nabla^m Y \in C((0,T), L^{\infty})$ with the respective L^{∞} norms bounded as $t \to 0^+$ then $\nabla^m u \in \mathcal{H}^{1/2}((0,T))$ and the corresponding constant $C_m(t)$ satisfies

$$C_m(t) \le c_m \sum_{\alpha+\beta=m} \int_0^t \frac{\|\nabla^{\alpha} Y(s)\|_{\infty} \|\nabla^{\beta} Y(s)\|_{\infty}}{(t-s)^{3/4}} \,\mathrm{d}s + c_m \frac{\|u_0\|}{t^{(m+2)/2}}$$

(iii) If additionally $Y \in C([0,T); L^2)$ then $p \in C([0,T); L^2)$ and $u \in C([0,T); L^2)$ with¹¹

$$\|u(t)\| \le C \int_0^t \frac{\|Y(s)\|_{\infty} \|Y(s)\|}{\sqrt{t-s}} \,\mathrm{d}s + \|u_0\|.$$
(2.17)

Moreover if T' < T and $\{Y^{(n)}\}$ is a sequence such that $Y^{(n)} \to Y$ in $C([0,T'];L^2)$ and $\max_{t \in [0,T']} \|Y^{(n)}(t)\|_{\infty} \le \max_{t \in [0,T']} \|Y(t)\|_{\infty}$ then

$$u^{(n)} \to u$$
 and $p^{(n)} \to p$ in $C([0,T'];L^2)$.

(iv) If additionally $Y \in \mathcal{H}^{1/2}((0,T))$ with the corresponding constant $C_0(t)$ bounded as $t \to 0^+$ then $\nabla u \in C((0,T); L^{\infty})$ with¹²

$$\|\nabla u(t)\|_{\infty} \le C \int_0^t \frac{\|Y(s)\|_{\infty} C_0(s)}{(t-s)^{3/4}} \,\mathrm{d}s + C \frac{\|u_0\|}{t^{5/4}}.$$
 (2.18)

More generally if for every multi-index α with $|\alpha| \leq m-1$, $D^{\alpha}Y \in \mathcal{H}^{1/2}((0,T)) \cap C((0,T);L^{\infty})$ with the corresponding Hölder constant $C_{\alpha}(t)$ and $\|D^{\alpha}Y(t)\|_{\infty}$ bounded as $t \to 0^+$ then $\nabla^m u \in C((0,T);L^{\infty})$ with

$$\|\nabla^m u(t)\|_{\infty} \le C_m \int_0^t \frac{\sum_{\alpha+\beta=m-1} \|\nabla^\alpha Y(s)\|_{\infty} C_{\beta}(s)}{(t-s)^{3/4}} \,\mathrm{d}s + C_m t^{-\frac{m}{2}-\frac{3}{4}} \|u_0\|$$

Proof. Since $Y \in C((0,T); L^{\infty})$ with $||Y(t)||_{\infty}$ bounded as $t \to 0^+$ the same is true of $Y_i Y_k$ for each pair i, k and so claim (i) follows from Lemma 1.1 and from the properties of the heat kernel (see Appendix A.1; note also that Lemma A.1 verifies the comment in (i)). In a similar way one obtains (iii), where the claim for p follows directly from the Plancherel Lemma (Lemma 1.4) and the limiting property follows from (2.17) and the Plancherel Lemma.

¹¹Leray does not state this property, but he uses it in showing existence of strong solutions to the Navier–Stokes equations (in the inequality he states at the bottom of page 223). We will apply it in a similar way (see Theorem 3.9) and also it in the existence and uniqueness theorem for the Stokes equations (Theorem 2.6).

¹²This corresponds to Leray's property of ∇u (which he obtains in (2.7) and (2.20)).

Property (ii) is a consequence of the Hölder continuity of the heat kernel (see (v) in Appendix A.1) and the Hölder continuity of $\nabla \mathcal{T}$ in L^1 (see Lemma 1.6). Indeed we have

$$\begin{aligned} u_2(x,t) &- u_2(y,t) |\\ &\leq \int_0^t \int |\nabla \mathcal{T}(x-z,t-s) - \nabla \mathcal{T}(y-z,t-s)| \, \mathrm{d}z \|Y(s)\|_\infty^2 \, \mathrm{d}s \\ &\leq C |x-y|^{1/2} \int_0^t \frac{\|Y(s)\|_\infty^2}{(t-s)^{3/4}} \, \mathrm{d}s. \end{aligned}$$

As for property (iv) note that the bound on ∇u_1 in (2.18) (that is $\|\nabla u_1(t)\|_{\infty} \leq C \|u_0\| t^{-5/4}$) and the continuity $\nabla u_1 \in C((0,T); L^{\infty})$ follow from properties of the heat kernel (see (iv) in Appendix A.1). As for u_2 , the bound on ∇u_2 in (2.18) can be shown using the following trick. Recalling that Y is weakly divergence free, we obtain

$$\int \partial_{li} \mathcal{T}_{kj}(x-y,t) Y_i(y,s) \,\mathrm{d}y = 0$$

for all j, k, l and $s, t \in (0, T), x \in \mathbb{R}^3$, where the integral exists due to (1.13). Hence, for each j, l and $t \in (0, T)$

$$\partial_l u_{2,j}(x,t) = -\int_0^t \int \partial_{li} \mathcal{T}_{jk}(x-y,t-s) Y_i(y,s) Y_k(y,s) \,\mathrm{d}y \,\mathrm{d}s$$
$$= -\int_0^t \int \partial_{li} \mathcal{T}_{jk}(x-y,t-s) Y_i(y,s) \left[Y_k(y,s) - Y_k(x,s) \right] \,\mathrm{d}y \,\mathrm{d}s, \quad (2.19)$$

and so using the bound $|\nabla^2 \mathcal{T}(x,t)| \leq C(|x|^2 + t)^{-5/2}$ (see (1.13)) and the assumption $Y \in \mathcal{H}^{1/2}((0,T))$ we obtain

$$\begin{aligned} |\nabla u_2(x,t)| &\leq C \int_0^t \int \frac{C_0(s) ||Y(s)||_\infty |x-y|^{1/2}}{(|x-y|^2 + (t-s))^{5/2}} \, \mathrm{d}y \, \mathrm{d}s \\ &= C \int_0^t \frac{||Y(s)||_\infty C_0(s)}{(t-s)^{3/4}} \, \mathrm{d}s, \end{aligned}$$

where we used the fact $\int |y|^{1/2}/(|y|^2+t)^{5/2} dy = Ct^{-3/4}$. Thus (2.18) follows. One can also employ this trick to show the continuity $\nabla u_2 \in C((0,T); L^{\infty})$, see Appendix A.4.3 for the details.

Finally, claims (ii) and (iv) for higher derivatives $\nabla^m u$ follow in a similar way. Indeed, the claims corresponding to u_1 follow from the properties of the heat kernel (see (iv) in Appendix A.1) and, as for u_2 , we write any $D^{\gamma}u_{2,j}$ with $|\gamma| = m$ as the sum of the integrals

$$\int_0^t \int \partial_l \partial_i \mathcal{T}_{jk}(x-y,t-s) Y_{\alpha,i}(y,s) Y_{\beta,k}(y,s) \,\mathrm{d}y \,\mathrm{d}s$$

where Y_{α} , Y_{β} denote appropriate derivatives of Y of orders α , β , respectively, where $\alpha + \beta = m - 1$, l, j = 1, 2, 3, and we repeat the reasoning above. \Box

Corollary 2.4. The results of the above lemma extend to the case $F_i = -\partial_k(Y_iZ_k)$ for some weakly divergence-free $Y, Z \in C((0,T), L^{\infty})$ with the L^{∞} norms bounded as $t \to 0^+$. In particular, if such Y, Z satisfy also $Y, Z \in C([0,T), L^2)$ and v is given by

$$v(t) \coloneqq \Phi(t) * u_0 + \int_0^t \nabla \mathcal{T}(t-s) * [Y(s)Z(s)] \,\mathrm{d}s \tag{2.20}$$

then $v \in C((0,T), L^{\infty}) \cap C([0,T), L^2)$ with

$$\|v(t)\|_{\infty} \le C \int_0^t \frac{\|Y(s)\|_{\infty} \|Z(s)\|_{\infty}}{\sqrt{t-s}} \,\mathrm{d}s + \|u_0\|_{\infty},$$
$$\|v(t)\| \le C \int_0^t \frac{\|Y(s)\|_{\infty} \|Z(s)\|}{\sqrt{t-s}} \,\mathrm{d}s + \|u_0\|.$$

The existence and uniqueness theorem for distributional solutions to the Stokes equations is based on the following uniqueness result.

Theorem 2.5 (Uniqueness of distributional solutions to the Stokes equations¹³). If u, p are such that $u \in C([0,T); L^2)$ is weakly divergence-free, $p \in L^1_{loc}(\mathbb{R}^3 \times [0,T))$, and

$$\int_0^T \int \left((\phi_t + \Delta \phi) \cdot u + p \operatorname{div} \phi \right) \, \mathrm{d}x \, \mathrm{d}t = 0 \tag{2.21}$$

for all $\phi \in C_0^{\infty}(\mathbb{R}^3 \times [0,T);\mathbb{R}^3)$, then $u \equiv 0$.

It follows that $\int_0^T \int p \operatorname{div} \phi \, dx \, dt = 0$, and so integration by parts and the Fundamental Lemma of Calculus of Variations give $\nabla p \equiv 0$; that is pis a function of t only. Since both the Stokes equations and the Navier– Stokes equations are invariant under addition to the pressure function any function of time, we will identify two solutions u_1, p_1 and u_2, p_2 of the Stokes equations (or the Navier–Stokes equations) if $u_1 = u_2$ and p_1 differs from p_2 by a function of time.

Proof (sketch; see Appendix A.4.4 for details). The proof of the theorem is based on considering the regularisations of u, p,

$$v(x,t) \coloneqq \int_0^t (J_{\varepsilon}u)(x,s) \,\mathrm{d}s, \qquad q(x,t) \coloneqq \int_0^t (J_{\varepsilon}p)(x,s) \,\mathrm{d}s, \quad \varepsilon > 0.$$

Such a regularisation is still a solution of (2.21) and one can show that $\Delta q = 0$ in a distributional sense. Thus Δv satisfies the homogeneous heat equation in a distributional sense and the uniqueness of the solution to the heat equation gives $\Delta v = 0$. An application of Liouville's theorem and the assumption $||u(t)|| < \infty$ for all t then gives $v \equiv 0$ for all $\varepsilon > 0$, and consequently $u \equiv 0$.

¹³This is a version of the argument from Section 14 of Leray (1934b).

We are now ready to prove the existence and uniqueness of distributional solutions, the central result of the study of the Stokes equations.

Theorem 2.6 (Distributional solution for F of the form (2.12)). Let $Y \in C([0,T), L^2) \cap C((0,T), L^{\infty})$ be weakly divergence free such that $||Y(t)||_{\infty}$ remains bounded as $t \to 0^+$. Then u, p given by (2.14), (2.15) comprise a distributional solution of (2.1), (2.2) with initial data u_0 and $F = -(Y \cdot \nabla)Y$. Moreover this solution is unique in the class $u \in C([0,T), L^2)$, $p \in L^1_{\text{loc}}(\mathbb{R}^3 \times [0,T))$.

Proof. Uniqueness follows from the theorem above. The fact that $u \in C([0,T); L^2)$ and the L^2 continuity at t = 0, $||u(t) - u_0|| \to 0$ as $t \to 0$, is clear from Lemma 2.3, (iii). Fix $\phi \in C_0^{\infty}(\mathbb{R}^3 \times [0,T), \mathbb{R}^3)$ and let $T' \in (0,T)$ be such that $\phi = 0$ for $t \ge T'$. Let $\{Y^{(n)}\}$ be a sequence of functions $Y^{(n)} \in C^{\infty}(\mathbb{R}^3 \times [0,T))$ such that $\operatorname{supp} Y^{(n)}(t) \subset B(0,R_n)$ for some $R_n > 0$,

$$||Y - Y^{(n)}||_{C([0,T'],L^2)} \to 0 \quad \text{as } n \to \infty$$

and $\max_{t\in[0,T']} \|Y^{(n)}(t)\|_{\infty} \leq \max_{t\in[0,T']} \|Y(t)\|_{\infty}$. Note that the above convergence means that also

$$||Y_iY_k - Y_i^{(n)}Y_k^{(n)}||_{C([0,T'],L^2)} \to 0 \quad \text{as } n \to \infty$$

for all i, k. The existence of such $Y^{(n)}$'s is guaranteed by Lemma A.12. Let (u_n, p_n) be given by (2.3), (2.4) with F replaced by $F^{(n)}$, where $F_k^{(n)} := -\partial_i(Y_i^{(n)}Y_k^{(n)})$. By Theorem 2.2 (u_n, p_n) satisfies the equations (2.1), (2.2) with F replaced by $F^{(n)}$ in the classical sense, and so also in the sense of distributions, that is u_n is weakly divergence free and

$$\int u_0 \cdot \phi \,\mathrm{d}x + \int_0^T \int \left(u_n \cdot (\phi_t + \Delta \phi) + p_n \,\mathrm{div}\,\phi \right) = \int_0^T \int Y^{(n)} \cdot (Y^{(n)} \cdot \nabla)\phi.$$
(2.22)

By Lemma 2.3, (iii), we have

$$||u_n - u||_{C([0,T'],L^2)} \to 0, \quad ||p_n - p||_{C([0,T'],L^2)} \to 0 \quad \text{as } n \to \infty$$

and so we can take the limit $n \to \infty$ to obtain that u is weakly divergence free and, from (2.22),

$$\int u_0 \cdot \phi \, \mathrm{d}x + \int_0^T \int \left(u \cdot (\phi_t + \Delta \phi) + p \operatorname{div} \phi \right) = \int_0^T \int Y \cdot (Y \cdot \nabla) \phi,$$

that is u, p is indeed the distributional solution.

Corollary 2.7. The conclusion of Theorem 2.6 also holds if F is of the form $F_i = -\partial_i(Y_iZ_k)$, where $Y, Z \in C([0,T), L^2) \cap C((0,T), L^\infty)$ are weakly divergence free with $||Y(t)||_{\infty}$, $||Z(t)||_{\infty}$ bounded as $t \to 0^+$, and the representation formula (2.14) is replaced by (2.20).

Notes

As remarked in the beginning of the section, we focused on the forcing of the form $F = -(Y \cdot \nabla)Y$ more directly than Leray. In particular Lemma 2.3 is not stated by explicitly by Leray. Thanks to the use of the distributional form of the Stokes equations (2.13) and the limiting property of the representation formulae (2.14), (2.15) (that is Lemma 2.3 (iii)) the main results of the section can be encapsulated in Theorem 2.6.

In the next section we will follow Leray in applying the results for the Stokes equations to study the Navier–Stokes equations. In particular we will employ the other properties of the modified representation formulae (2.14), (2.15), that is Lemma 2.3 (i),(ii),(iv). These properties were presented by Leray either implicitly during the study of the Navier–Stokes equations or by showing a related result for a general of the forcing F, as we pointed out in the footnotes in Lemma 2.3.

3 Strong solutions of the Navier–Stokes equations

We now consider the Navier–Stokes equations

$$\partial_t u - \Delta u + \nabla p = -(u \cdot \nabla)u, \qquad (3.1)$$

$$\operatorname{div} u = 0 \tag{3.2}$$

in $\mathbb{R}^3 \times (0, T)$. We will consider a weak form of these equations,

$$\int_0^T \int (u \cdot (f_t + \Delta f) + p \operatorname{div} f) = \int_0^T \int u \cdot (u \cdot \nabla) f$$
(3.3)

for $f \in C_0^{\infty}(\mathbb{R}^3 \times (0,T))$, where u(t) is weakly divergence free. We first define solutions on the open time interval (0,T) (see below) and we study their properties in Section 3.1. In Section 3.2 we equip the problem (3.1)-(3.2) with initial data, and for this reason we extend the definition of strong solutions to the half-closed time interval [0,T) (Definition 3.7). We then show existence and uniqueness of local-in-time strong solutions (Theorem 3.9). In Section 3.3 we study the maximal time of existence for strong solutions and the rate of blow-up of strong solutions if the maximal time is finite. In Section 3.4 we study local existence and uniqueness of strong solutions with less regular initial data.

Definition 3.1. A function u is a strong solution of the Navier–Stokes equations on the time interval (0,T) if it satisfies the weak form of the equations with some $p \in L^1_{loc}(\mathbb{R}^3 \times (0,T))$ and if

$$u \in C((0,T); L^2) \cap C((0,T); L^\infty).$$

This is how Leray defines a strong solution, except that he requires the continuity of all terms appearing in the Navier–Stokes equations (3.1) (see p. 217). Here we make use of the weak form of equations and thus we avoid specifying any conditions on derivatives of u. However, smoothness of strong solutions (Corollary 3.3) implies that the two definitions are equivalent to each other. Note also that if u is a strong solution of the Navier–Stokes equations then (3.3) is equivalent to requiring that

$$\int u(t_1) \cdot f(t_1) - \int u(t_2) \cdot f(t_2) + \int_{t_1}^{t_2} \int (u \cdot (f_t + \Delta f) + p \operatorname{div} f) = \int_{t_1}^{t_2} \int u \cdot (u \cdot \nabla) f \quad (3.4)$$

for all $f \in C_0^{\infty}(\mathbb{R}^3 \times (0,T))$ and $t_1, t_2 \in (0,T)$ with $t_1 < t_2$. While the " \Leftarrow " part this equivalence is trivial, the " \Rightarrow " part is not immediate but can be obtained by a simple cut-off procedure, which we now explain.

For h > 0 let $F_h(x, s) := f(x, s)\theta_h(s)$, where $\theta_h \in C^{\infty}(\mathbb{R})$ is a nonincreasing function such that $\theta_h(s) = 1$ for $s \le t_2$, $\theta_h(s) = 0$ for $s \ge t_2 + h$. Using F_h as a test function in (3.3) we obtain

$$\int_0^{t_2+h} \int \left(u \cdot (f_t + \Delta f) + p \operatorname{div} f \right) \theta_h + \int_{t_2}^{t_2+h} \int u \cdot f \theta'_h$$
$$= \int_0^{t_2+h} \int u \cdot (u \cdot \nabla) f \theta_h.$$

Since $u, f \in C((0,T); L^2)$ the function $s \mapsto \int u(s) \cdot f(s)$ is continuous. Thus, since θ_h is nonincreasing and $\int_{\mathbb{R}} \theta'_h = -1$ we obtain

$$\int_{t_2}^{t_2+h} \int u \cdot f \,\theta'_h \to -\int u(t_2) \cdot f(t_2) \quad \text{as } h \to 0^+$$

Thus taking the limit $h \to 0^+$ in the last equation (via the Dominated Convergence Theorem) gives

$$-\int u(t_2) \cdot f(t_2) + \int_0^{t_2} \int (u \cdot (f_t + \Delta f) + p \operatorname{div} f) = \int_0^{t_2} \int u \cdot (u \cdot \nabla) f.$$

Applying a similar cut-off procedure at time t_1 gives (3.4).

From (3.4) and the theorem about the existence and uniqueness of distributional solutions to Stokes equations (Theorem 2.6) we see that a strong solution of the Navier–Stokes equations admits representation formulae¹⁴,

$$u(t_2) = \Phi(t_2 - t_1) * u(t_1) + \int_{t_1}^{t_2} \int \nabla \mathcal{T}(t_2 - t_1 - s) * [u(s)u(s)] \, \mathrm{d}s,$$

$$p(t_2) = \partial_i \partial_k (-\Delta)^{-1} (u_i(t_2)u_k(t_2)).$$
(3.5)

¹⁴These are (3.2) and (3.3) in Leray (1934b).

for all $t_1, t_2 \in (0, T)$ with $t_1 < t_2$. Recall we employ the notation

$$\left(\nabla \mathcal{T}(t-s) * [Y(s)Z(s)]\right)_j(x) \coloneqq \int \partial_i \mathcal{T}_{jk}(x-y,t-s)Y_i(y,s)Z_k(y,s)\,\mathrm{d}y.$$

Note this representation formula also determines uniquely the pressure function p.

3.1 Properties of strong solutions

In this section we study the properties of strong solutions of the Navier– Stokes equations on the open time interval (0, T). We will show that if u is a strong solution and p is the corresponding pressure function then u and pare smooth and u satisfies an energy equality, along with some other useful results. The theorem below as well as the following corollary show that u, p are smooth.

Theorem 3.2. ¹⁵ If u is a strong solution of the Navier–Stokes equations on (0,T) then

$$\nabla^m u \in C((0,T); L^2) \cap C((0,T); L^\infty)$$
 for all $m \ge 0$.

The proof of the theorem (as well as the corollary that follows) is a simplification¹⁶ of Leray's arguments (which he presents on pages 218-219).

Proof. The proof proceeds by a double use of induction. First, we show that

$$\nabla^m u \in C((0,T); L^{\infty}) \cap \mathcal{H}^{1/2}((0,T)) \quad \text{for } m \ge 0.$$

Here the base case follows from the definition of a strong solution and from Lemma 2.3, (ii), and the induction step follows from the same lemma, properties (iv) and (ii).

Second, we show that

$$\nabla^m u \in C((0,T); L^2), \ \nabla^m [(u \cdot \nabla)u] \in C((0,T); L^2) \text{ for } m \ge 0.$$

Here the base case follows from the definition of a strong solution and by deducing that $(u \cdot \nabla)u \in C((0,T); L^2)$ by using Hölder's inequality,

$$\begin{aligned} \|(u(t) \cdot \nabla)u(t) - (u(s) \cdot \nabla)u(s)\| \\ &\leq \|u(t)\| \, \|\nabla u(t) - \nabla u(s)\|_{\infty} + \|\nabla u(s)\|_{\infty} \|u(t) - u(s)\|. \end{aligned}$$

The induction step follows from Lemma 2.1 (ii) and from a similar use of Hölder's inequality. $\hfill \Box$

¹⁵Theorem 3.2 and Corollary 3.3 correspond to pp. 218-219 in Leray (1934b).

 $^{^{16}}$ This simplification is due to our choice to organise the properties of the representation formulae (2.3), (2.4) and (2.14), (2.15) into Lemmas 2.1 and 2.3, see Notes at the end of the section

Note that in fact for each $s \in (0, T)$ we have bounded the norms $\|\nabla^m u(s)\|$, $\|\nabla^m u(s)\|_{\infty}$, $m \ge 0$ using only the norms $\|u(t)\|$, $\|u(t)\|_{\infty}$, $t \in (0,T)$.

Corollary 3.3 (Smoothness of strong solutions). If u is a strong solution of the Navier–Stokes equations on (0,T) and p is the corresponding pressure then

$$\partial_t^k \nabla^m u, \partial_t^k \nabla^m p \in C((0,T); L^2) \cap C((0,T); L^\infty) \quad \text{for all } m, k \ge 0.$$

In particular $u, p \in C^{\infty}(\mathbb{R}^3 \times (0,T))$ and u, p constitute a classical solution of the Navier-Stokes equations on $\mathbb{R}^3 \times (0,T)$.

Proof. From the representation of p, (3.5), and the Plancherel Lemma (Lemma 1.4), we obtain that $\nabla^m p \in C((0,T); L^2)$ for all m. Moreover, using the Cauchy-Schwarz inequality and Lemma 1.3 we obtain for any $x \in \mathbb{R}^3$

$$\begin{aligned} |p(x,t) - p(x,s)| &\leq \int \frac{|\partial_i u_j(y,t)(\partial_j u_i(y,t) - \partial_j u_i(y,s))|}{4\pi |x-y|} \,\mathrm{d}y \\ &+ \int \frac{|\partial_j u_i(y,s)(\partial_i u_j(y,t) - \partial_i u_j(y,s))|}{4\pi |x-y|} \,\mathrm{d}y \\ &\leq C \|\nabla u(t) - \nabla u(s)\| \left(\sqrt{\int \frac{|\nabla u(y,t)|^2}{|x-y|^2} \,\mathrm{d}y} + \sqrt{\int \frac{|\nabla u(y,s)|^2}{|x-y|^2} \,\mathrm{d}y} \right) \\ &\leq C \|\nabla u(t) - \nabla u(s)\| \left(\|\nabla^2 u(t)\| + \|\nabla^2 u(s)\| \right). \end{aligned}$$

Thus $p \in C((0,T); L^{\infty})$ and performing a similar calculation for each of the spatial derivatives of p shows that $\nabla^m p \in C((0,T);L^\infty)$ for all m. Thus the distributional form of the Navier–Stokes equations (3.3) gives $\nabla^m u_t \in$ $C((0,T); L^2) \cap C((0,T); L^{\infty})$ for all m.

The regularity of higher derivatives in time follows by induction: regularity of $\partial_t^k u$ follows from the regularity of $u, \partial_t u, \ldots, \partial_t^{k-1} u$ and $\partial_t^{k-1} p$, and by taking (k-1)-th time derivative (in a weak sense) of the Navier–Stokes equations, and the regularity of $\partial_t^k p$ follows by taking k time derivatives of the representation formula of p, (3.5).

Theorem 3.4 (Energy equality for strong solutions¹⁷). A strong solution uof the Navier-Stokes equations on (0,T) satisfies

$$\|u(t_2)\|^2 + 2\int_{t_1}^{t_2} \|\nabla u(s)\|^2 \,\mathrm{d}s = \|u(t_1)\|^2 \tag{3.6}$$

 $[\]frac{\text{for all } t_1, t_2 \in (0, T).}{^{17}\text{This is } (3.4) \text{ in Leray (1934b).}}$

Proof. Since Theorem 3.2 gives in particular $(u \cdot \nabla)u \in C((0,T); L^2)$, Lemma 2.1, (iii) gives

$$\|u(t_2)\|^2 - \|u(t_1)\|^2 + 2\int_{t_1}^{t_2} \|\nabla u(s)\|^2 \,\mathrm{d}s = 2\int_{t_1}^{t_2} \int u \cdot (u \cdot \nabla) u \,\mathrm{d}x \,\mathrm{d}s.$$

The theorem follows by noting that the right-hand side vanishes: integration by parts and the incompressibility constraint, $\partial_k u_k = 0$, give

$$\int u \cdot (u \cdot \nabla) u = \int u_i u_k \,\partial_k u_i = -\int \partial_k u_i \,u_k \,u_i, \qquad (3.7)$$

that is $\int u \cdot (u \cdot \nabla) u = 0.$

We now show that we can control the separation of two strong solutions.

Lemma 3.5 (Comparison of two strong solutions¹⁸). Suppose that u, v are strong solutions to the Navier–Stokes equations on (0,T) and let

$$w \coloneqq u - v$$

Then

$$||w(t_2)||^2 \le ||w(t_1)||^2 e^{\frac{1}{2} \int_{t_1}^{t_2} ||u(s)||_{\infty}^2 ds}$$
(3.8)

for $t_1, t_2 \in (0, T)$ with $t_1 < t_2$.

In particular, if u, v coincide at time t_1 then they continue to coincide for the later times. We will extend this uniqueness property to account for the initial data in Section 3.2 (Lemma 3.8).

Proof. Since both u and v satisfy the Navier–Stokes equations pointwise, subtracting them gives

$$\partial_t w - \Delta w + \nabla q = -(u \cdot \nabla)u + (v \cdot \nabla)v = -(v \cdot \nabla)w - (w \cdot \nabla)u.$$

As in (3.7) we have $\int w \cdot (v \cdot \nabla) w = 0$ and hence multiplying the above equality by w, integrating by parts in spatial variables, using the incompressibility constraint, $\partial_k w_k = 0$, we obtain for $t \in (0, T)$

$$\begin{aligned} \frac{1}{2} \frac{\mathrm{d}}{\mathrm{d}t} \|w(t)\|^2 + \|\nabla w(t)\|^2 &= -\int w \cdot ((v \cdot \nabla)w + (w \cdot \nabla)u) \\ &= \int w_i w_k \partial_k u_i = -\int \partial_k w_i w_k u_i \\ &\leq \|\nabla w(t)\| \|w(t)\| \|u(t)\|_{\infty} \\ &\leq \|\nabla w(t)\|^2 + \frac{1}{4} \|w(t)\|^2 \|u(t)\|_{\infty}^2, \end{aligned}$$

¹⁸This is Section 18 of Leray (1934b).

where we also used the Cauchy-Schwarz and Young inequalities (and we omitted the argument "t" under the integrals). Hence

$$\frac{\mathrm{d}}{\mathrm{d}t} \|w(t)\|^2 \le \frac{1}{2} \|w(t)\|^2 \|u(t)\|_{\infty}^2,$$

and the claim follows by applying Gronwall's inequality.

Finally, the remark after the proof of Theorem 3.2 suggests the following convergence property of a family of strong solutions of the Navier–Stokes equations.

Lemma 3.6 (Convergence lemma¹⁹). Suppose $\{u_{\varepsilon}\}_{\varepsilon>0}$ is a family of strong solutions of the Navier-Stokes equations such that

$$\|u_{\varepsilon}(t)\|_{\infty}, \|u_{\varepsilon}(t)\| \le f(t) \qquad \text{for } t \in (0,T),$$

where f is a continuous function on (0,T). Then there exists a sequence $\varepsilon_k \to 0^+$ and a function u, such that $u_{\varepsilon_k} \to u, \nabla u_{\varepsilon_k} \to \nabla u$ uniformly on compact sets in $\mathbb{R}^3 \times (0,T)$ as $\varepsilon_k \to 0^+$.

Moreover u is a strong solution of the Navier-Stokes equations in (0,T)and satisfies $||u(t)||_{\infty}, ||u(t)|| \leq f(t)$ for all $t \in (0,T)$.

Proof. Let $p_{\varepsilon}(t)$ denote the pressure function corresponding to u_{ε} (which is determined uniquely by the representation formula (3.5) with u replaced by u_{ε}). Fix a bounded domain $\Omega \subset \mathbb{R}^3$ and $\delta > 0$. We see from Corollary 3.3 that for a multi-index α and m such that $m, |\alpha| \leq 3$

$$|\partial_{t^m} D^{\alpha} u_{\varepsilon}|, |\partial_{t^m} D^{\alpha} p_{\varepsilon}| \le C_{\Omega, \delta} \quad \text{on } \Omega_{\delta} \times (\delta/2, T - \delta/2),$$

where $\Omega_{\delta} \coloneqq \Omega + B(0, \delta)$. Thus an application of the Arzelà–Ascoli theorem and a simple diagonalization argument produces a sequence $\{\varepsilon_k\}$ such that all derivatives $\partial_{t^m} D^{\alpha} u_{\varepsilon_k}$, $\partial_{t^m} D^{\alpha} p_{\varepsilon_k}$ with $m, |\alpha| \leq 2$ converge to the respective derivatives of u and p uniformly on $\Omega \times (\delta, T - \delta)$, for some functions u, p. In particular the Navier–Stokes equations

$$\partial_t u_{\varepsilon_k} - \Delta u_{\varepsilon_k} + \nabla p_{\varepsilon_k} = -(u_{\varepsilon_k} \cdot \nabla) u_{\varepsilon_k},$$

div $u_{\varepsilon_k} = 0$

converge uniformly on $\Omega \times (\delta, T - \delta)$ to Navier–Stokes equations for u (in the sense that all terms converge). Now consider a sequence of bounded sets $\Omega_n \nearrow \mathbb{R}^3$, a sequence $\delta_n \to 0^+$ and apply another diagonal argument to obtain a subsequence $\{\varepsilon_k\}$ (which we relabel) such that for m, α with $m, |\alpha| \leq 2$

$$\partial_{t^m} D^{\alpha} u_{\varepsilon_k} \to \partial_{t^m} D^{\alpha} u, \quad \partial_{t^m} D^{\alpha} p_{\varepsilon_k} \to \partial_{t^m} D^{\alpha} p$$

¹⁹This is Lemma 9 in Leray (1934b).

uniformly on compact sets in $\mathbb{R}^3 \times (0,T)$. In particular u, p satisfy the Navier–Stokes equations pointwise, and thus also in the sense of distributions (3.3). That $||u(t)||_{\infty} \leq f(t)$ holds for all t is clear, and the inequality $||u(t)|| \leq f(t)$ follows by an application of Fatou's lemma. According to Definition 3.1 it remains to verify that $u \in C((0,T); L^2) \cap C((0,T); L^{\infty})$. For this let I be a closed interval in (0,T) and note that there exists M > 0 such that $||\partial_t u_{\varepsilon_k}(t)||_{\infty}, ||\partial_t u_{\varepsilon_k}(t)|| \leq M$ for $t \in I, k \geq 0$ (see Corollary 3.3). Thus the mean value theorem gives

$$\|u_{\varepsilon_k}(t) - u_{\varepsilon_k}(s)\|_{\infty}, \|u_{\varepsilon_k}(t) - u_{\varepsilon_k}(s)\| \le M |t - s| \quad \text{for } s, t \in I, \ k \ge 0.$$

Thus taking the limit in k (and applying Fatou's lemma) gives the required continuity.

3.2 Local existence and uniqueness of strong solutions

In this section we study the Navier–Stokes initial value problem, that is we consider the equations (3.1), (3.2) with initial data. For this reason we extend the definition of strong solutions (Definition 3.1) to the half-closed time interval [0, T).

Definition 3.7. A function u is a strong solution of the Navier–Stokes equations on [0,T) if for some $p \in L^1_{loc}(\mathbb{R}^3 \times [0,T))$

$$\int u(0) \cdot f(0) + \int_0^T \int (u \cdot (f_t + \Delta f) + p \operatorname{div} f) = \int_0^T \int u \cdot (u \cdot \nabla) f \quad (3.9)$$

for all $f \in C_0^{\infty}(\mathbb{R}^3 \times [0,T);\mathbb{R}^3)$, u(t) is weakly divergence free for $t \in (0,T)$, and

$$u \in C([0,T); L^2) \cap C((0,T); L^{\infty})$$
(3.10)

with $||u(t)||_{\infty}$ bounded as $t \to 0^+$.

The regularity (3.10) is a part of Leray's definition of solutions on the time interval [0, T), but he also requires $\nabla u \in C([0, T); L^2)$ and that u and its spacial derivatives are continuous at t = 0 (see pp. 220-221 of his paper). It is remarkable that by use of the weak formulation (3.9), these additional assumptions are not necessary for showing local well-posedness (see Theorem 3.9 below). Moreover, the above definition is much less restrictive than the commonly used definition of strong solutions, which usually requires

$$u \in L^{\infty}_{\text{loc}}([0,T); H^1) \cap L^2_{\text{loc}}([0,T); H^2),$$

and so consequently $u \in C([0,T); H^1)$ see, for example, Definition 6.1 and the following discussion in Robinson et al. (2016). In particular, Definition 3.7 makes no assumption on the regularity of $\nabla u(t)$ for times t near 0. Since Definition 3.7 is an extension of the definition of the strong solution on the open time interval (0,T) (Definition 3.1), we see that u, p admit the representation formulae (3.5). Moreover, now the representation formula also holds for $t_1 = 0$,

$$u(t) = \Phi(t) * u(0) + \int_0^t \nabla \mathcal{T}(t-s) * [u(s) u(s)] \, \mathrm{d}s$$
 (3.11)

for $t \in (0, T)$, a consequence of the definition above and Theorem 2.6.

We also see that (3.9) is equivalent to

$$\int u(0) \cdot f(0) - \int u(t) \cdot f(t) + \int_0^t \int (u \cdot (f_t + \Delta f) + p \operatorname{div} f) = \int_0^t \int u \cdot (u \cdot \nabla) f$$
(3.12)

being satisfied for all $t \in (0,T)$, $f \in C_0^{\infty}(\mathbb{R}^3 \times [0,T); \mathbb{R}^3)$ (see the discussion following Definition 3.1). A consequence of this fact is that a strong solution on the time interval [0,T) is also a strong solution on the time interval $[\tau,T)$ for any $\tau \in (0,T)$.

Given the definition of strong solutions on the half-closed time interval [0, T), we immediately obtain the energy equality (3.6) with $t_1 = 0$ and the uniqueness of strong solutions.

Lemma 3.8 (Uniqueness and the energy equality for local strong solution). A strong solution u to the Navier–Stokes equations on [0,T) satisfies the energy equality

$$||u(t)||^{2} + 2\int_{0}^{t} ||\nabla u(s)||^{2} \,\mathrm{d}s = ||u_{0}||^{2}$$
(3.13)

for $t \in [0, T)$.

Moreover, if v is another strong solution to the Navier–Stokes equations on [0,T) with v(0) = u(0) then $u \equiv v$.

Proof. The claim follows by taking the limit $t_1 \to 0^+$ in (3.6) and (3.8) and applying the Monotone Convergence Theorem.

From the lemma we obtain the semigroup property: if $\tau \in (0, T_1)$ and u, \tilde{u} are strong solutions on the intervals $[0, T_1)$, $[\tau, T_2)$ respectively with $\tilde{u}(\tau) = u(\tau)$, then $\tilde{u} = u$ on the time interval $[\tau, \min\{T_1, T_2\})$. We now state the central theorem regarding strong solutions to the Navier–Stokes equations.

Theorem 3.9 (Local existence of strong solutions). If $u_0 \in H \cap L^{\infty}$ (that is $u_0 \in L^2 \cap L^{\infty}$ is weakly divergence free, see (1.2)) then there exists a unique strong solution u of the Navier–Stokes equations on [0,T) with $u(0) = u_0$, where $T > C/||u_0||_{\infty}^2$.

This theorem is proved by Leray in Section 19 in the case when $u_0 \in C^1 \cap H^1 \cap L^\infty$. Here, we use the distributional form of equations to relax this regularity requirement to only $u_0 \in H \cap L^\infty$, and to demonstrate how Leray's original proof can be simplified while exposing his main ideas.

Proof. Uniqueness is guaranteed by Lemma 3.8. As for existence, we consider the following iterative definition of $u^{(n)}$:

$$u^{(0)}(t) \coloneqq \Phi(t) * u_0$$

and

$$u^{(n+1)}(t) \coloneqq \int_0^t \nabla \mathcal{T}(t-s) * \left[u^{(n)}(s) \, u^{(n)}(s) \right] \, \mathrm{d}s + u^{(0)}(t),$$

using the notation from (2.16). From properties of the heat kernel (see (iii) in Appendix A.1) we have

$$u^{(0)} \in C([0,\infty); L^2) \cap C((0,\infty); L^\infty),$$

 $||u^{(0)}(t)||_{\infty} \leq ||u_0||_{\infty}$ and $||u^{(0)}(t)|| \leq ||u_0||$ for $t \geq 0$. Moreover, using induction we deduce from Lemma 2.3, (i) and (iii) (applied with $Y \coloneqq u^{(n)}$) that for all $n \geq 0, t \geq 0$,

$$u^{(n+1)} \in C([0,\infty); L^2) \cap C((0,\infty); L^\infty),$$

$$\|u^{(n+1)}(t)\|_{\infty} \leq C' \int_0^t \frac{\|u^{(n)}(s)\|_{\infty}^2}{\sqrt{t-s}} \,\mathrm{d}s + \|u_0\|_{\infty}, \tag{3.14}$$

and

$$\|u^{(n+1)}(t)\| \le C' \int_0^t \frac{\|u^{(n)}(s)\|_{\infty} \|u^{(n)}(s)\|}{\sqrt{t-s}} \,\mathrm{d}s + \|u_0\|, \qquad (3.15)$$

Theorem 2.6 guarantees that if

$$p^{(n+1)}(t) \coloneqq \partial_k \partial_i (-\Delta)^{-1} \left(u_i^{(n)}(t) \, u_k^{(n)}(t) \right)$$

(recall (1.6) for the notation) then for each $n \ge 0$ the pair $u^{(n+1)}, p^{(n+1)}$ is a distributional solution of the problem

$$\partial_t u^{(n+1)} - \Delta u^{(n+1)} + \nabla p^{(n+1)} = (u^{(n)} \cdot \nabla) u^{(n)}, \quad \text{div} \, u^{(n+1)} = 0$$

with initial condition $u^{(n+1)}(0) = u_0$, that is $u^{(n+1)}$ is weakly divergence free and

$$\int u_0 \cdot f(0) \,\mathrm{d}x + \int_0^\infty \int \left(u^{(n+1)} \cdot (f_t + \Delta f) + p^{(n+1)} \,\mathrm{div}\, f \right)$$
$$= \int_0^\infty \int u^{(n)} \cdot (u^{(n)} \cdot \nabla) f \quad (3.16)$$

for all $f \in C_0^{\infty}(\mathbb{R}^3 \times [0, \infty), n \ge 0$. In order to take the limit in n we will find a uniform bound on $||u^{(n+1)}||_{\infty}$ on the finite time interval [0, T], where

$$T := \frac{1}{32(1+C')^4 \|u_0\|_{\infty}^2}$$

For such choice of T > 0 the constant function $\phi(t) \coloneqq (1+C') ||u_0||_{\infty}$ satisfies the integral inequality

$$\phi(t) \ge C' \int_0^t \frac{\phi^2(s)}{\sqrt{t-s}} \,\mathrm{d}s + \|u_0\|_\infty$$
 (3.17)

for all $t \in [0, 8T)$. A use of induction and (3.14) thus gives

$$||u^{(n)}(t)||_{\infty} \le \phi(t).$$
 (3.18)

for all such t's and all $n \ge 0$. Now noting that for all $j, k = 1, 2, 3, n \ge 1$

$$u_k^{(n)}u_j^{(n)} - u_k^{(n-1)}u_j^{(n-1)} = u_k^{(n)}\left(u_j^{(n)} - u_j^{(n-1)}\right) + u_j^{(n-1)}\left(u_k^{(n)} - u_k^{(n-1)}\right)$$

we use Corollary 2.4 twice (first with $Y \coloneqq u^{(n)}, Z \coloneqq u^{(n)} - u^{(n-1)}$ and then with $Y \coloneqq u^{(n-1)}, Z \coloneqq u^{(n)} - u^{(n-1)}$) to obtain

$$\|u^{(n+1)}(t) - u^{(n)}(t)\|_{\infty}$$

 $\leq C' \int_0^t \frac{\left(\|u^{(n)}(s)\|_{\infty} + \|u^{(n-1)}(s)\|_{\infty}\right) \|u^{(n)}(s) - u^{(n-1)}(s)\|_{\infty}}{\sqrt{t-s}} \,\mathrm{d}s, \quad (3.19a)$

$$\begin{aligned} \|u^{(n+1)}(t) - u^{(n)}(t)\| \\ &\leq C' \int_0^t \frac{\left(\|u^{(n)}(s)\|_{\infty} + \|u^{(n-1)}(s)\|_{\infty}\right) \|u^{(n)}(s) - u^{(n-1)}(s)\|}{\sqrt{t-s}} \,\mathrm{d}s. \end{aligned} (3.19b)$$

Applying (3.18) to the second of the above inequalities gives for $t \in [0, T]$

$$\begin{aligned} \|u^{(n+1)}(t) - u^{(n)}(t)\| \\ &\leq 2C'(1+C')\|u_0\|_{\infty}\|u^{(n)} - u^{(n-1)}\|_{C([0,T],L^2)} \int_0^t \frac{1}{\sqrt{t-s}} \,\mathrm{d}s \\ &= 4C'(1+C')\|u_0\|_{\infty}\sqrt{t}\|u^{(n)} - u^{(n-1)}\|_{C([0,T],L^2)} \\ &\leq \lambda \|u^{(n)} - u^{(n-1)}\|_{C([0,T],L^2)}, \end{aligned}$$

where $\lambda \coloneqq 4C'(1+C') \|u_0\|_{\infty} \sqrt{T}$. Hence

$$\|u^{(n+1)} - u^{(n)}\|_{C([0,T],L^2)} \le \lambda \|u^{(n)} - u^{(n-1)}\|_{C([0,T],L^2)}$$

Because the definition of T implies $\lambda \in (0, 1)$ we see that $\{u^{(n)}\}\$ is a Cauchy sequence in $C([0, T]; L^2)$ and so

$$u^{(n)} \to u$$
 in $C([0,T]; L^2)$

for some $u \in C([0,T]; L^2)$ such that $u(0) = u_0$ and u(t) is weakly divergence free for each $t \in [0,T]$ (since $u^{(n)}(0) = u_0$ and $u^{(n)}(t)$ is weakly divergence free for each n). Similarly, applying (3.18) to the first inequality in (3.19) gives

$$\|u^{(n+1)} - u^{(n)}\|_{C([0,T];L^{\infty})} \le \lambda \|u^{(n)} - u^{(n-1)}\|_{C([0,T];L^{\infty})}.$$

Although this does not imply that $\{u^{(n)}\}$ is Cauchy in $C([0,T]; L^{\infty})$ (recall each $u^{(n)}$ need not belong to this space; it is continuous into L^{∞} only on the open time interval $(0,\infty)$), it does follow that $\{u^{(n)}\}$ is a Cauchy sequence in $C([\delta,T]; L^{\infty})$ for any $\delta \in (0,T)$, and therefore

$$u^{(n)} \to u \quad \text{in } C([\delta, T]; L^{\infty})$$

$$(3.20)$$

for each δ . Note that the limit function is u since L^2 convergence implies convergence almost everywhere on a subsequence. Therefore

$$u \in C([0,T];L^2) \cap C((0,T];L^\infty)$$

and (3.18) gives $||u(t)||_{\infty} \leq (1+C')||u_0||_{\infty}$ for all $t \in [0,T]$. Letting

$$p(t) \coloneqq \partial_i \partial_k (-\Delta)^{-1} \left(u_i(t) \, u_k(t) \right)$$

we see that the Plancherel Lemma (Lemma 1.4) gives that

$$p^{(n)} \to p$$
 in $C([0,T];L^2)$.

Therefore, taking the limit $n \to \infty$ in (3.16) we obtain

$$\int u_0 \cdot f \, \mathrm{d}x + \int_0^T \int \left(u \cdot (f_t + \Delta f) + p \operatorname{div} f \right) = \int_0^T \int u \cdot (u \cdot \nabla) f$$

for all $f \in C_0^{\infty}(\mathbb{R}^3 \times [0, T))$. Thus u is a strong solution of the Navier–Stokes equations on [0, T).

Note that if u_0 is more regular then the continuity as $t \to 0^+$ of the corresponding strong solution u on [0,T) may be stronger. This is, in essence, the issue of continuity as $t \to 0^+$ of the solution of the heat equation (cf. Lemma A.1). If u_0 is uniformly continuous then the representation formula (3.11) gives that $u(t) \to u_0$ in L^{∞} as $t \to 0^+$ (cf. the proof of Lemma 2.3 (i)). Furthermore, for such u_0 the proof above simplifies since each $u^{(n)}$ belong to $C([0,\infty); L^{\infty})$ (cf. the same lemma) and so they converge in $C([0,T]; L^{\infty})$ rather than in $C([\delta,T]; L^{\infty})$ for all δ 's as in (3.20).

3.3 Characterisation of singularities

Here we investigate the maximal time of existence of strong solutions and derive the rates of blow-up of u in various norms at an approach to a (putative) blow-up time.

Let $u_0 \in V \cap L^{\infty}$ (that is $u_0 \in H^1 \cap L^{\infty}$ is divergence free, see (1.2)), let u be the strong solution of the Navier–Stokes equations starting from u_0 and let T_0 be its maximal time of existence, that is u cannot be extended to a solution on [0, T') for any $T' > T_0$. Note that Theorem 3.9 gives that $T_0 \geq C/||u_0||_{\infty}^2$ and, if T_0 is finite,

$$||u(t)||_{\infty}$$
 blows up as $t \to T_0^-$,

as otherwise we could extend u beyond T_0 and hence obtain a contradiction.

In this section we will apply the theory of integral inequalities (see Lemma 1.5) to bound the L^{∞} norm of a strong solution u on some time interval starting from 0 and thus obtain lower bounds on T_0 as well as lower bounds on $||u(t)||_{\infty}$, $||\nabla u(t)||$ and $||u(t)||_p$ with p > 3 when $t \to T_0^-$ (if T_0 is finite).

Since (3.11) holds for $t \in (0, T_0)$, Lemma 2.3 (i) gives that for all $t \in [0, T_0), p > 3$

$$\|u(t)\|_{\infty} \le C' \int_0^t \frac{\|u(s)\|_{\infty}^2}{\sqrt{t-s}} \,\mathrm{d}s + \min\left(\|u_0\|_{\infty}, C'\frac{\|\nabla u_0\|}{t^{1/4}}, C'\frac{\|u_0\|_p}{t^{3/2p}}\right).$$
(3.21)

Here, the minimum on the right-hand side is obtained by applying Young's inequality for convolutions (1.5) to $u_1(t) = \Phi(t) * u_0$ (with exponents $(1, \infty)$, (6/5, 6) and (p/(p-1), p) respectively), the fact that $\|\Phi(t)\|_p \leq C/t^{-3(p-1)/2p}$, and the Gagliardo-Nirenberg-Sobolev²⁰ inequality $\|u_0\|_6 \leq C \|\nabla u_0\|$. Similarly, the convolution under the time integral in the representation formula (3.11) can be bounded in terms of $\|\nabla \mathcal{T}(t-s)\|_{\infty} \|u(s)\|^2$ (rather than by $\|\nabla \mathcal{T}(t-s)\|_1 \|u(s)\|_{\infty}^2$ as in Lemma 2.3, (i)), and this gives for $t \in [0, T_0)$,

$$\|u(t)\|_{\infty} \le C'' \int_0^t \min\left(\frac{\|u(s)\|_{\infty}^2}{\sqrt{t-s}}, \frac{\|u_0\|^2}{(t-s)^2}\right) \,\mathrm{d}s + \|u_0\|_{\infty},\tag{3.22}$$

where we also used the facts $\|\nabla \mathcal{T}(t)\|_{\infty} \leq Ct^{-2}$ (see (1.13)) and $\|u(t)\| \leq \|u_0\|$ (see the energy equality (3.13)). The two inequalities above²¹ allow us to obtain bounds on $\|u(t)\|_{\infty}$ in terms of various norms of the initial data.

$$\left| \int \Phi(x-y,t)u_0(y) \, \mathrm{d}y \right| \le 2 \left| \int |\Phi(x-y,t)|^2 |x-y|^2 \right|^{1/2} \|\nabla u_0\| \le C \|\nabla u_0\| t^{-1/4}.$$

 $^{^{20}\}mathrm{Note}$ that this inequality were not available in the 1930's. Instead, Leray used Lemma 1.3 to obtain

²¹These comprise (3.5) in Leray (1934b).

Lemma 3.10. ²² If u is the strong solution with initial data $u_0 \in V \cap L^{\infty}$ then

- (i) $||u(t)||_{\infty} \leq C ||u_0||_{\infty}$ for $t \leq C/||u_0||_{\infty}^2$,
- (ii) $||u(t)||_{\infty} \leq C ||\nabla u_0|| t^{-1/4}$ for $t \leq C/||\nabla u_0||^4$,
- (iii) $||u(t)||_{\infty} \leq C ||u_0||_p t^{-3/2p}$ for $t \leq (C(1-3/p)/||u_0||_p)^{2p/(p-3)}, p > 3.$

Moreover, there exists $\varepsilon > 0$ such that $||u(t)||_{\infty} \leq C ||u_0||_{\infty}$ for all $t \geq 0$ if $||u_0||^2 ||u_0||_{\infty} < \varepsilon$.

Proof. As in the proof of Theorem 3.9, we note that the constant function $\phi(t) = C ||u_0||_{\infty}$ satisfies the integral inequality (3.17) for $t \in [0, C/||u_0||_{\infty}^2]$ and (i) follows from this, (3.21) and from the theory of integral inequalities (see Lemma 1.5). Similarly, a direct calculation shows that the function $\phi(t) = C ||\nabla u_0|| t^{-1/4}$ satisfies the integral inequality

$$\phi(t) \ge C' \int_0^t \frac{\phi(s)^2}{\sqrt{t-s}} \,\mathrm{d}s + C' \|\nabla u_0\| t^{-1/4}$$

for $t \in (0, C/||\nabla u_0||^4]$, and so (ii) follows. One can also check that the function $\phi(t) = C||u_0||_p t^{-3/2p}$ satisfies the integral inequality

$$\phi(t) \ge C' \int_0^t \frac{\phi(s)^2}{\sqrt{t-s}} \,\mathrm{d}s + C' \|u_0\|_p t^{-3/2p}$$

for $t \in (0, (C(1-3/p)/||u_0||_p)^{2p/(p-3)}]$, and (iii) follows.

The last claim follows from (3.22) and the fact that the constant function $\phi(t) = C ||u_0||_{\infty}$ satisfies the integral inequality

$$\phi(t) > C'' \int_0^t \min\left(\frac{\phi(s)^2}{\sqrt{t-s}}, \frac{\|u_0\|^2}{(t-s)^2}\right) \,\mathrm{d}s + \|u_0\|_{\infty},\tag{3.23}$$

for all t > 0 if and only if

$$||u_0||_{\infty} > C \int_0^\infty \min\left(\frac{||u_0||_{\infty}^2}{\sqrt{s}}, \frac{||u_0||^2}{s^2}\right) \mathrm{d}s.$$

One can check that this last condition is equivalent to the smallness condition $||u_0||^2 ||u_0||_{\infty} < \varepsilon$ for some $\varepsilon > 0$. Therefore, the integral inequalities (3.23) and (3.22) show that, given the smallness condition, $||u(t)||_{\infty} \le \phi(t)$ for all $t \ge 0$ (where we apply another fact from the theory of integral inequalities, see Corollary A.7).

 $^{^{22}\}mathrm{This}$ lemma and the two following corollaries correspond to Sections 21 and 22 in Leray (1934b).

From the lemma we immediately obtain lower bounds on the maximal time of existence T_0 , the rates of blow-up of norms $||u(t)||_{\infty}$, $||\nabla u(t)||$ and $||u(t)||_p$, where p > 3, as $t \to T_0^-$, as well as global existence result for small data, which we formulate in the following three corollaries.

Corollary 3.11 (Lower bounds on the existence time T_0). If T_0 is the maximal time of existence of the strong solution u with initial data $u_0 \in V \cap L^{\infty}$ then

- (i) $T_0 > C/||u_0||_{\infty}^2$,
- (ii) $T_0 > C / \| \nabla u_0 \|^4$,

(iii)
$$T_0 > \left(C\left(1-\frac{3}{p}\right)/\|u_0\|_p\right)^{2p/(p-3)}$$
 for all $p > 3$.

Proof. It is a direct consequence of Lemma 3.10 (i)-(iii).

Corollary 3.12 (Blow-up rates). If u is a strong solution of the Navier– Stokes equations on the time interval (T, T_0) , where $T_0 < \infty$ is the maximal existence time, then for $t \in (T, T_0)$,

$$||u(t)||_{\infty} \ge \frac{C}{\sqrt{T_0 - t}}, \quad ||\nabla u(t)|| \ge \frac{C}{(T_0 - t)^{1/4}},$$

and, for p > 3,

$$||u(t)||_p \ge \frac{C^{(1-3/p)/2}(1-3/p)}{(T_0-t)^{(1-3/p)/2}}$$

Proof. Let $t \in (T, T_0)$. Since $u(t) \in H \cap L^{\infty}$, the local existence and uniqueness theorem (Theorem 3.9) gives that $(T_0 - t) \geq C/||u(t)||_{\infty}^2$, which gives the first bound. The other two follow in a similar way using Corollary 3.11, (ii) and (iii).

Corollary 3.13 (Global existence for small initial data). There exists $\varepsilon > 0$ such that if either $||u_0||^2 ||u_0||_{\infty} < \varepsilon$, $||u_0|| ||\nabla u_0|| < \varepsilon$ or

$$(C||u_0||)^{2(p-3)} ||u_0||_p^p < C\left(1 - \frac{3}{p}\right)\varepsilon^{p-3} \qquad \text{for any } p > 3 \qquad (3.24)$$

then $T_0 = \infty$, that is the strong solution with initial data u_0 exists for all times.

Proof. The first claim follows directly from the last claim of Lemma 3.10. As for the smallness condition on $||u_0|| ||\nabla u_0||$, let $t_0 \coloneqq C/||\nabla u_0||^4$, the endpoint time in Lemma 3.10 (ii). Then

$$||u(t_0)||_{\infty} \le C ||\nabla u_0|| t_0^{-1/4} = C ||\nabla u_0||^2,$$

and so, using the energy equality (3.13), we obtain

$$||u(t_0)||^2 ||u(t_0)||_{\infty} \le C ||u_0||^2 ||\nabla u_0||^2.$$

In other words, the condition $||u(t_0)||^2 ||u(t_0)||_{\infty} < \varepsilon$ holds if $||u_0|| ||\nabla u_0||$ is sufficiently small, as required.

As for the smallness condition on $(C||u_0||)^{2(p-3)}||u_0||_p^p$, take $t_0 := (C(1-3/p)/||u_0||_p)^{2p/(p-3)}$, the endpoint time in Lemma 3.10 (iii). Then

$$\|u(t_0)\|_{\infty} \le C \|u_0\|_p t_0^{-3/2p} = C \|u_0\|_p^{p/(p-3)} \left(C\left(1-\frac{3}{p}\right)\right)^{-3/(p-3)}$$

Thus the energy equality (3.13) and (3.24) gives

$$||u(t_0)||^2 ||u(t_0)||_{\infty} \le ||u_0||^2 ||u(t_0)||_{\infty} < \varepsilon,$$

as required.

Finally, we deduce the following result²³, which we will only use later in analysing the structure of a weak solution (Theorem 4.12).

Corollary 3.14. Let u be a strong solution of the Navier–Stokes equations on the time interval (T,T_0) , let $t_1 \in (T,T_0)$ and $t_2 > t_1$. If $t_2 - t_1 \leq C \|\nabla u(t_1)\|^{-4}$ then

$$\|u(t_2)\|_{\infty} \le C \frac{\|\nabla u(t_1)\|}{(t_2 - t_1)^{1/4}}, \quad and \quad \|\nabla u(t_2)\| \le C \|\nabla u(t_1)\|.$$

Proof. The first inequality follows directly from Lemma 3.10 (ii). For the second one, note that from the smoothness of strong solutions (see Corollary 3.3) we have $(u \cdot \nabla)u \in C([t_1, T_0); L^2)$, so by the representation (3.5) and Lemma 2.1 (ii), we obtain²⁴

$$\begin{aligned} \|\nabla u(t)\| &\leq C \int_{t_1}^t \frac{\|\nabla u(s)\| \|u(s)\|_{\infty}}{\sqrt{t-s}} \,\mathrm{d}s + \|\nabla u(t_1)\| \\ &\leq C''' \|\nabla u(t_1)\| \int_{t_1}^t \frac{\|\nabla u(s)\|}{\sqrt{t-s(s-t_1)^{1/4}}} \,\mathrm{d}s + \|\nabla u(t_1)\| \end{aligned}$$

for all $t \in [t_1, T_0)$. Now, a direct calculation shows that the constant function $\phi(t) \coloneqq C \|\nabla u(t_1)\|$ satisfies the integral inequality

$$\phi(t) \ge C''' \|\nabla u(t_1)\| \int_{t_1}^t \frac{\phi(s)}{\sqrt{t-s}(s-t_1)^{1/4}} \,\mathrm{d}s + \|\nabla u(t_1)\|$$

for $t \in [t_1, t_1 + C/||\nabla u(t_1)||^4]$. Therefore,

 $\|\nabla u(t)\| \le \phi(t)$ for $t \in [t_1, t_1 + C/\|\nabla u(t_1)\|^4]$,

where we also used a fact from the theory of integral inequalities, see Corollary A.6. Thus we obtain the second of the required inequalities. \Box

²³This corollary is a consequence of Leray's (3.19).

 $^{^{24}}$ This is (3.6) in Leray (1934b).
3.4 Semi-strong solutions

In this section we focus on the regularity required from u_0 in order to generate a unique strong solution. In Section 3.2 we have shown that $u_0 \in H \cap L^{\infty}$ generates such a solution that is strong on [0, T) (see Definition 3.7) for some T > 0 (see Theorem 3.9). We also observed that the high regularity of u_0 guarantees some further properties of such solutions; in particular the representation formula (3.11).

It turns out that relaxing the regularity of u_0 still gives a unique strong solution for (sufficiently small) positive times. This motivates the following definition.²⁵

Definition 3.15. A function u is a semi-strong solution of the Navier– Stokes equations (3.1), (3.2) on the time interval [0,T) if it is a strong solution on the open time interval (0,T) (see Definition 3.1) such that $u \in C([0,T); L^2)$ and

$$\int_{0}^{t} \|u(s)\|_{\infty}^{2} \,\mathrm{d}s < \infty \quad \text{for all } t \in (0,T).$$
(3.25)

Note this definition is less restrictive than the definition of strong solutions on the time interval [0,T) (Definition 3.7). Namely, we replace the weak form of the equations (3.9) by the weak form (3.3), which does not include the initial data u_0 , and we replace the boundedness of $||u(t)||_{\infty}$ as $t \to 0^+$ by the integral condition (3.25). Note that the initial condition $u(0) = u_0$ is now incorporated in the assumption $u \in C([0,T); L^2)$.

Lemma 3.16. Semi-strong solutions to the Navier–Stokes equations on [0,T) (that satisfy a given initial condition) are unique and satisfy the energy equality

$$||u(t)||^{2} + 2 \int_{0}^{t} ||\nabla u(s)||^{2} ds = ||u(0)||^{2}$$

for all $t \in [0,T)$.

Proof. As in Lemma 3.8, the claim follows by taking the limit $t_1 \to 0^+$ in (3.6) and (3.8) and applying Monotone Convergence Theorem (note that the integral condition (3.25) guarantees that the exponent in (3.8) remains finite as $t_1 \to 0^+$).

We now use the notion of semi-strong solutions to obtain a local-in-time well-posedness for initial data $u_0 \in V$ (rather than $u_0 \in H \cap L^{\infty}$ as in Theorem 3.9).²⁶

 $^{^{25}\}mathrm{Definition}$ 3.15 and the uniqueness result in Lemma 3.16 are stated in Section 23 in Leray's paper.

²⁶This is Section 24 in Leray (1934b).

Theorem 3.17. If $u_0 \in V$ (that is $u_0 \in H^1$ is divergence free) then there exists a unique semi-strong solution u on time interval [0,T), where $T \geq C/||\nabla u_0||^4$, such that $u(0) = u_0$.

Proof. Note that $J_{\varepsilon}u_0 \in H \cap L^{\infty}$ and $\operatorname{div}(J_{\varepsilon}u_0) = 0$ (see Section 1.1). Hence Theorem 3.9 implies the existence of a unique strong solution $u_{\varepsilon}(t)$ to the Navier–Stokes equations on some time interval $[0, T_{\varepsilon})$ such that $u_{\varepsilon}(0) = J_{\varepsilon}u_0$. The energy equality (3.13) and the properties of mollification (see Lemma 1.2) give

$$||u_{\varepsilon}(t)|| \le ||u_{\varepsilon}(0)|| = ||J_{\varepsilon}u_{0}|| \le ||u_{0}||,$$
 (3.26)

$$\|\nabla u_{\varepsilon}(0)\| = \|\nabla (J_{\varepsilon}u_0)\| = \|J_{\varepsilon}(\nabla u_0)\| \le \|\nabla u_0\|.$$
(3.27)

The last bound and Corollary 3.11 let us bound the existence time T_{ε} from below independently of ε ,

$$T_{\varepsilon} \ge C / \|\nabla u_0\|^4 =: T.$$

Moreover, Lemma 3.10 (ii) gives

$$||u_{\varepsilon}(t)||_{\infty} \le C ||\nabla u_{\varepsilon}(0)||t^{-1/4} \le C ||\nabla u_{0}||t^{-1/4}$$
(3.28)

for $t \in [0, T)$. By (3.26) and (3.28) we may apply the convergence lemma (Lemma 3.6) to extract a sequence $\{\varepsilon_k\}$ such that $u_{\varepsilon_k} \to u$ almost everywhere, where u is a strong solution of the Navier–Stokes equations on $\mathbb{R}^3 \times (0, T)$ with

$$\|u(t)\| \le \|u_0\| \tag{3.29}$$

and

$$|u(t)||_{\infty} \le C ||\nabla u_0|| t^{-1/4} \tag{3.30}$$

for $t \in (0, T)$. It follows from the last inequality that $\int_0^t ||u(s)||_{\infty}^2 ds$ is finite for all $t \in (0, T)$. It remains to verify that $u(t) \to u_0$ in L^2 as $t \to 0$. Since u_{ε_k} is a strong solution to the Navier–Stokes equations on [0, T) (3.12) gives

$$0 = \int u_{\varepsilon_k}(t) \cdot \phi - \int (J_{\varepsilon_k} u_0) \cdot \phi - \int_0^t \int u_{\varepsilon_k} \cdot \Delta \phi - \int_0^t \int u_{\varepsilon_k} \cdot (u_{\varepsilon_k} \cdot \nabla) \phi$$

for $t \in [0,T)$ and $\phi \in C_0^{\infty}$. By the fact that $J_{\varepsilon}u_0 \to u_0$ in L^2 as $\varepsilon \to 0$, (3.26) and the Dominated Convergence Theorem (applied to the time integrals) we can pass to the limit in the above equation to obtain

$$0 = \int u(t) \cdot \phi - \int u_0 \cdot \phi - \int_0^t \int u \cdot \Delta \phi - \int_0^t \int u \cdot (u \cdot \nabla) \phi,$$

which gives that

$$\int u(t) \cdot \phi \to \int u_0 \cdot \phi$$
 as $t \to 0^+$

for all $\phi \in C_0^{\infty}$, which, by L^2 boundedness (3.29), gives that $u(t) \rightarrow u_0$ weakly in L^2 as $t \rightarrow 0^+$. In order to show that $u(t) \rightarrow u_0$ strongly in L^2 it is enough to show the convergence of the norms, $||u(t)|| \rightarrow ||u_0||$ as $t \rightarrow 0^+$. This last claim follows from properties of weak limits and (3.29) by writing

$$||u_0|| \le \liminf_{t \to 0} ||u(t)|| \le \limsup_{t \to 0} ||u(t)|| \le ||u_0||.$$

Similarly, we obtain that the notion of semi-strong solutions gives localin-time well-posedness for $u_0 \in H \cap L^p$, where $p > 3.^{27}$

Corollary 3.18. Given $u_0 \in H \cap L^p$ with $p \in (3, \infty)$ there exists a semistrong solution u of the Navier–Stokes equations on [0,T) with $u(0) = u_0$, where $T \ge (C(1-3/p)/||u_0||_p)^{2p/(p-3)}$.

Proof. Copy the proof above making the following replacements. Replace (3.27) by $||u_{\varepsilon}(0)||_p \leq ||u_0||_p$, (3.28) by $||u_{\varepsilon}(t)||_{\infty} \leq C||u_0||_p t^{-3/2p}$, (3.30) by $||u(t)||_{\infty} \leq C||u_0||_p t^{-3/2p}$, and T by $(C(1-3/p)/||u_0||_p)^{2p/(p-3)}$.

Notes

This section corresponds to Chapters III and IV of Leray (1934b).

In Section 20 Leray (1934b) considers the issue of the existence of solutions that blow up. He points out that such a solution exists if

$$\begin{cases} \Delta U(x) - \alpha U(x) - \alpha (x \cdot \nabla) U(x) - \nabla P(x) = (U(x) \cdot \nabla) U(x), \\ \operatorname{div} U(x) = 0, \end{cases}$$

has a nontrivial solution in \mathbb{R}^3 for some $\alpha > 0$. In that case there would exist a strong solution of the Navier–Stokes equations on the time interval $(-\infty, T)$ of the following self-similar form

$$u(x,t) \coloneqq \frac{1}{\sqrt{2\alpha(T-t)}} U\left(\frac{x}{\sqrt{2\alpha(T-t)}}\right),$$

which would blow up at time T. However, Nečas, Růžička & Šverák (1996) have shown that this system of equations has no nontrivial L^3 solutions.

In fact, it is rather remarkable that the issue of the existence of solutions that blow up is one of the most important open problems in mathematics to this day, one of seven Millennium Problems (see Fefferman (2006)).

Leray showed the smoothness of strong solutions via similar bounds as in the analysis of the Stokes equations (pp. 218-219). Since in our presentation the properties of the representation formulae (2.3), (2.4) and (2.14), (2.15)

²⁷This is Section 25 of Leray (1934b). Note that the case $u_0 \in H \cap L^{\infty}$, for which Leray states well-posedness of semi-strong solutions, was covered in this article in the well-posedness result for strong solutions, see Theorem 3.9.

are organised in Lemmas 2.1 and 2.3, we were able to prove the smoothness of strong solutions by induction.

Furthermore, Section 3.3 shows that the analysis of the maximal time of existence and the blow-up rates can be done without the use of Leray's (3.6),

$$\|\nabla u(t)\| \le C \int_0^t \frac{\|\nabla u(s)\| \, \|u(s)\|_{\infty}}{\sqrt{t-s}} \mathrm{d}s + \|\nabla u_0\|,$$

which we use only in the proof of Corollary 3.14.

4 Weak solutions of the Navier–Stokes equations

In this section we show the global existence of a weak solution (which Leray termed a *turbulent solution*) of the Navier-Stokes equations. His approach is characterised by considering the following modified system, for $\varepsilon > 0$:

$$\partial_t u - \Delta u + ((J_\varepsilon u) \cdot \nabla)u + \nabla p = 0,$$

div $u(t) = 0,$ (4.1)

which is often called the *Leray regularisation*. We will see that this regularisation of the nonlinear term gives for each $\varepsilon > 0$ a unique, global in time, strong solution. We then study the limit $\varepsilon \to 0$ of the solutions of the above equations.

4.1 Well-posedness for the regularised equations

Definition 4.1. A function u_{ε} is a strong solution of the regularised equations (4.1) on the interval [0,T) if for some $p_{\varepsilon} \in L^1_{loc}(\mathbb{R}^3 \times [0,T))$

$$\int u_{\varepsilon}(0) \cdot f(0) + \int_{0}^{T} \int (u_{\varepsilon} \cdot (f_{t} + \Delta f) + p_{\varepsilon} \operatorname{div} f) = \int_{0}^{T} \int (J_{\varepsilon} u_{\varepsilon}) \cdot (u_{\varepsilon} \cdot \nabla) f \quad (4.2)$$

for all $f \in C_0^{\infty}(\mathbb{R}^3 \times [0,T);\mathbb{R}^3)$, $u_{\varepsilon}(t)$ is weakly divergence free for $t \in (0,T)$, and

$$u_{\varepsilon} \in C([0,T);L^2) \cap C((0,T);L^{\infty})$$

with $||u_{\varepsilon}(t)||_{\infty}$ bounded as $t \to 0^+$.

Note this definition follows the lines of the definition of a strong solution of the Navier–Stokes equations on time interval [0,T) (Definition 3.7), the difference appearing only in the form of the distributional equations (4.2). Moreover, we see that a solution u_{ε} and the corresponding pressure p_{ε} are given by

$$u_{\varepsilon}(t) = \Phi(t) * u_{\varepsilon}(t_1) + \int_{t_1}^t \int \nabla T(t-s) * \left[(J_{\varepsilon}u_{\varepsilon})(s) u_{\varepsilon}(s) \right] ds, \qquad (4.3)$$
$$p_{\varepsilon}(t) = \partial_i \partial_k (-\Delta)^{-1} \left((J_{\varepsilon}u_{\varepsilon,i})(t) u_{\varepsilon,k}(t) \right)$$

for all $0 \le t_1 < t < T$, cf. (3.5) and (3.11), see also Corollary 2.7.

Now let $u_0 \in H^1 \cap L^\infty$ be divergence free; we will show existence and uniqueness of global-in-time strong solution of the regularised equations with initial data u_0 .

Theorem 4.2 (Global well-posedness of the regularised equations²⁸). For each $\varepsilon > 0$ there exists a unique strong solution u_{ε} of the regularised equations (4.1) on the time interval $[0,\infty)$ such that $u_{\varepsilon}(0) = u_0$, u_{ε} is smooth on the time interval $(0,\infty)$ (in the sense of Corollary 3.3) and the energy equality

$$||u_{\varepsilon}(t)||^{2} + 2\int_{0}^{t} ||\nabla u_{\varepsilon}(s)||^{2} \,\mathrm{d}s = ||u_{0}||^{2}$$
(4.4)

holds for all $t \geq 0$.

Proof. We note that the analysis from Section 3 can be applied to the regularised equations (4.1). In particular, by noting that $||J_{\varepsilon}v|| \leq ||v||$ and $||J_{\varepsilon}v||_{\infty} \leq ||v||_{\infty}$ for any $v \in L^2 \cap L^{\infty}$ (see Lemma 1.2, (i)), we can prove a local existence and uniqueness theorem following Theorem 3.9, to obtain a unique strong solution u_{ε} of the system (4.1) on the time interval [0,T) for some $T \geq C/||u_0||_{\infty}^2$. Now following the arguments in Section 3.1 we note that (using the representation formulae (4.3) instead of (3.5)) u_{ε} is smooth on the time interval (0,T), and so following Theorem 3.4 and Lemma 3.8 we obtain the energy equality

$$||u_{\varepsilon}(t)||^{2} + 2 \int_{0}^{t} ||\nabla u_{\varepsilon}(s)||^{2} ds = ||u_{0}||^{2}$$

for $t \in [0, T)$. It remains to show that T, the maximal time of existence, is infinite.

As in Section 3.3 we see that $||u_{\varepsilon}(t)||_{\infty}$ must blow-up as $t \to T^{-}$ if $T < \infty$. We also obtain

$$\|u(t)\|_{\infty} \le C \int_0^t \frac{\|(J_{\varepsilon}u_{\varepsilon})(s)\|_{\infty}\|u(s)\|_{\infty}}{\sqrt{t-s}} \,\mathrm{d}s + \|u_0\|_{\infty}$$

for $t \in [0, T)$, in a similar way to the derivation of (3.21). This inequality, however, is fundamentally different from (3.21) in the sense that we can now apply the bound

$$\|(J_{\varepsilon}u_{\varepsilon})(s)\|_{\infty} \le C\varepsilon^{3/2} \|u_{\varepsilon}(s)\| \le C\varepsilon^{3/2} \|u_0\|,$$

(see Lemma 1.2 (iii) and the energy equality (4.4)). Moving this bound outside of the integral, we obtain a linear integral inequality,

$$\|u_{\varepsilon}(t)\|_{\infty} \leq C'\varepsilon^{-3/2}\|u_0\| \int_0^t \frac{\|u_{\varepsilon}(s)\|_{\infty}}{\sqrt{t-s}} \,\mathrm{d}s + \|u_0\|_{\infty}.$$

²⁸This is Section 26 of Leray (1934b).

Letting $\phi_{\varepsilon} \in C([0,\infty))$ be the unique solution to the corresponding linear integral equation²⁹,

$$\phi_{\varepsilon}(t) = C' \varepsilon^{-3/2} \|u_0\| \int_0^t \frac{\phi_{\varepsilon}(s)}{\sqrt{t-s}} \,\mathrm{d}s + \|u_0\|_{\infty}, \quad t \ge 0,$$
(4.5)

we see that $||u_{\varepsilon}(t)||_{\infty} \leq \phi_{\varepsilon}(t)$ for all $t \geq 0$ (see Corollary A.6). Therefore $||u_{\varepsilon}(t)||_{\infty}$ remains bounded on every finite interval and hence $T = \infty$, as required.

In order to study the limit as $\varepsilon \to 0$ of the solutions u_{ε} to the system (4.1), we first show that the kinetic energy of $u_{\varepsilon}(t)$ outside of a ball can be bounded independently of ε .

Lemma 4.3 (Separation of energy³⁰). Let $\varepsilon > 0$, $0 < R_1 < R_2$ and let u_{ε} be the solution of (4.1) with initial condition u_0 . Then for $t \ge 0$

$$\int_{|x|>R_2} |u_{\varepsilon}(t)|^2 \, \mathrm{d}x \le \int_{|x|>R_1} |u_0|^2 \, \mathrm{d}x + \frac{C(u_0, t)}{R_2 - R_1},$$

where $C(u_0, t) := C ||u_0||^2 \sqrt{t} + C ||u_0||^3 t^{1/4}$.

Proof. For brevity we will write u in place of u_{ε} . Let

$$f(x) := \begin{cases} 0 & |x| < R_1, \\ \frac{|x| - R_1}{R_2 - R_1} & R_1 \le |x| \le R_2, \\ 1 & |x| > R_2. \end{cases}$$

Taking the scalar product of the regularised equations (4.1) against $-2f(x)u_i(x,t)$, integrating in space and time and using $\operatorname{div} u = 0$ yields

$$2\int_{0}^{t} \int f|\nabla u|^{2} + \int f|u|^{2}$$

= $\int f|u_{0}|^{2} - \int_{0}^{t} \int \left(2\partial_{k}f \, u_{i} \, \partial_{k}u_{i} + 2\partial_{i}f \, p \, u_{i} + \partial_{k}f(J_{\varepsilon}u_{k})|u|^{2}\right) \quad (4.6)$

Bounding below the second term on the left-hand side by $\int_{|x|>R_2} |u(t)|^2$ and

 $^{^{29}(4.5)}$ is an example of the Volterra equation; see Appendix A.6 for the proof of existence and uniqueness of solutions. 30 This is Section 27 of Leray (1934b).

using the nonnegativity of the first term yields

$$\begin{split} &\int_{|x|>R_2} |u(t)|^2 \\ &\leq \int f |u_0|^2 - \int_0^t \int \left(2\partial_k f u_i \partial_k u_i + 2\partial_i f \, p \, u_i + \partial_k f (J_\varepsilon u_k) |u|^2 \right) \\ &\leq \int_{|x|>R_1} |u_0|^2 + \frac{1}{R_2 - R_1} \int_0^t \left(2 ||u|| ||\nabla u|| + 2 ||u|| ||p|| + ||J_\varepsilon u|| ||u||_4^2 \right) \\ &\leq \int_{|x|>R_1} |u_0|^2 + \frac{||u_0||}{R_2 - R_1} \left(2 \int_0^t ||\nabla u|| + 2 \int_0^t ||p|| + \int_0^t ||u||_4^2 \right). \end{split}$$

Let us denote the last three integrals on the right hand side by I_1 , I_2 and I_3 respectively. We have

$$I_1 \le \sqrt{t} \left(\int_0^t \|\nabla u(s)\|^2 \, \mathrm{d}s \right)^{1/2} \le \frac{\sqrt{t}}{2} \|u_0\|$$
(4.7)

by the energy equality (4.4). As for I_2 , I_3 note that since $|u|^2$ solves (trivially) the Poisson equation $\Delta |u|^2 = \Delta |u|^2$ we can integrate by parts to obtain

$$|u|^{2} = \frac{-1}{4\pi} \int \frac{1}{|x-y|} \Delta |u(y)|^{2} \,\mathrm{d}y = \frac{1}{4\pi} \int \frac{x-y}{|x-y|^{3}} \cdot \nabla |u(y)|^{2} \,\mathrm{d}y.$$

Thus

$$\begin{aligned} \|u\|_{4}^{4} &= \frac{1}{4\pi} \int \int |u(x)|^{2} \frac{x-y}{|x-y|^{3}} \cdot \nabla |u(y)|^{2} \, \mathrm{d}x \, \mathrm{d}y \\ &\leq \frac{1}{2\pi} \int \int \frac{|u(x)|^{2}}{|x-y|^{2}} \, |u(y)| \, |\nabla u(y)| \, \mathrm{d}x \, \mathrm{d}y \\ &\leq C \|\nabla u\|^{2} \int |u(y)| \, |\nabla u(y)| \, \mathrm{d}y \leq C \|\nabla u\|^{3} \|u\|, \end{aligned}$$

where we used Lemma 1.3 and the Cauchy-Schwarz inequality. Note that this estimate is independent of t (which we omitted in the notation). Moreover, the representation formula (4.3) for p together with the Plancherel Lemma (Lemma 1.4) and the bound $||J_{\varepsilon}u||_4 \leq ||u||_4$ (see Lemma 1.2 (i)) give

$$||p|| \le C ||J_{\varepsilon}u| |u||| \le C ||u||_4^2 \le C ||\nabla u||^{3/2} ||u||^{1/2}.$$
(4.8)

Therefore ||p|| and $||u||_4^2$ enjoy the same bound $C||\nabla u||^{3/2}||u||^{1/2}$. Thus, by the energy equality (4.4) and Hölder's inequality, we obtain

$$I_{2}, I_{3} \leq C \int_{0}^{t} \|\nabla u(s)\|^{3/2} \|u(s)\|^{1/2} \, \mathrm{d}s \leq C \|u_{0}\|^{1/2} \int_{0}^{t} \|\nabla u(s)\|^{3/2} \, \mathrm{d}s$$
$$\leq C \|u_{0}\|^{1/2} t^{1/4} \left(\int_{0}^{t} \|\nabla u(s)\|^{2} \, \mathrm{d}s\right)^{3/4} \leq C \|u_{0}\|^{2} t^{1/4}.$$

Hence, finally

$$\int_{|x|>R_2} |u(t)|^2 \leq \int_{|x|>R_1} |u_0|^2 + \frac{||u_0||}{R_2 - R_1} (2I_1 + 2I_2 + I_3)$$
$$\leq \int_{|x|>R_1} |u_0|^2 + \frac{C}{R_2 - R_1} \left(t^{1/2} ||u_0||^2 + t^{1/4} ||u_0||^3 \right). \quad \Box$$

Remark 4.4. It is interesting to note that Leray (1934b) presented a way of deriving the bound (4.8) that does not use the Plancherel Lemma (in fact Leray does not mention the use of Fourier transform). See Appendix A.7 for details.

4.2 Global existence of a weak solution

Here we study the limit as $\varepsilon \to 0$ of the solutions u_{ε} of the regularised equations (4.1) to obtain a global-in-time weak solution of the Navier–Stokes equations, as defined below.

Definition 4.5. ³¹ A function u is a weak solution of the Navier–Stokes equations if there exists a set $S \subset (0, \infty)$ of measure zero such that u(t) is weakly divergence free for all $t \in (0, \infty) \setminus S$,

$$\int u(t) \cdot f(t) = \int u_0 \cdot f(0) + \int_0^t \int u \cdot (\partial_t f + \Delta f) + \int_0^t \int u \cdot (u \cdot \nabla) f \quad (4.9)$$

for all t > 0 and all divergence-free test functions f such that $\partial_t^m \nabla^k f \in C([0,\infty); L^2) \cap C([0,\infty); L^\infty)$ for all $k, m \ge 0$, and

$$\|u(t)\|^{2} + 2\int_{s}^{t} \|\nabla u(r)\|^{2} \,\mathrm{d}r \le \|u(s)\|^{2} \tag{4.10}$$

for every $s \in [0, \infty) \setminus S$ and every $t \ge s$.

Such a solution is often called a *Leray-Hopf weak solution* (Eberhard Hopf (1951) considered weak solutions in a similar sense on a bounded domain), while weak solutions are functions (belonging to $L^{\infty}(0,T;H) \cap L^2(0,T;V)$) satisfying the first part but not necessarily the energy inequality, see, for example, Definitions 4.9 and 3.3 in Robinson et al. (2016) (see also Lemma 6.6 therein for the equivalence of the spaces of test functions). The set S is often called the set of singular times.

Corollary 4.6. A weak solution u satisfies

$$u \in L^{\infty}((0,\infty); L^2) \cap L^2((0,\infty); H^1).$$

Moreover, it is L^2 weakly continuous in time and for $s \in [0, \infty) \setminus S$, $u(t) \to u(s)$ in L^2 as $t \to s^+$.

³¹This is the definition in Section 31 of Leray (1934b).

Proof. The first property is a consequence of the energy inequality (4.10), the L^2 weakly continuity is a consequence of (4.9), and the last property is a consequence of the weak continuity and the convergence of the norms $||u(t)|| \rightarrow ||u(s)||$ for $s \in [0, \infty) \setminus S$ (which follows from (4.10)).

Theorem 4.7 (Global existence of a weak solution³²). If $u_0 \in H$ then there exists a weak solution u of the Navier–Stokes equations such that $||u(t) - u_0|| \to 0$ as $t \to 0^+$.

Proof. For each $\varepsilon > 0$ let u_{ε} be the unique strong solution of (4.1) with $u_{\varepsilon}(0) = J_{\varepsilon}u_0$. The existence of such u_{ε} is guaranteed by Theorem 4.2, which also gives

$$\|u_{\varepsilon}(t)\|^{2} + 2\int_{0}^{t} \|\nabla u_{\varepsilon}(s)\|^{2} \,\mathrm{d}s = \|J_{\varepsilon}u_{0}\|^{2} \le \|u_{0}\|^{2}$$
(4.11)

for $t \ge 0$, $\varepsilon > 0$ (see (4.4); the last inequality is a property of the mollification operator, see Lemma 1.2 (i)). The weak solution is constructed in the following four steps.

Step 1. Construct the sequence $\{\varepsilon_n\}$.

Multiplying the equation in (4.1) by f and integrating by parts we obtain, for $t \ge 0$,

$$\int u_{\varepsilon}(t) \cdot f(t) = \int J_{\varepsilon} u_0 \cdot f(0) + \int_0^t \int u_{\varepsilon} \cdot (\partial_t f + \Delta f) + \int_0^t \int u_{\varepsilon} \cdot (J_{\varepsilon} u_{\varepsilon} \cdot \nabla) f.$$
(4.12)

Inequality (4.11) implies that for each t the numbers $||u_{\varepsilon}(t)||$ are bounded independently of ε , and so using a diagonal argument we extract a subsequence $\{\varepsilon_n\}$ such that for all $t \in \mathbb{Q}^+$, $||u_{\varepsilon_n}(t)|| \to W(t)$ as $k \to \infty$, for some function $W \colon \mathbb{Q}^+ \to [0, \infty)$. Extend W to the whole of \mathbb{R}_+ by letting $W(t) \coloneqq \liminf_{s \to t^-} W(s)$ for $t \in \mathbb{R}_+ \setminus \mathbb{Q}^+$. Using the following fact, we see that $||u_{\varepsilon_n}(t)|| \to W(t)$ as $n \to \infty$ for times t at which W is continuous.

Fact 4.8 (Helly's theorem³³). If $g_n \in C([0,1])$ is nonincreasing for each n and $g_n(t) \to g(t)$ for all $t \in \mathbb{Q} \cap [0,1]$ then $g_n(t) \to g(t)$ at each continuity point t of g.

Since $||u_{\varepsilon_n}(t)||$ is a nonincreasing function of t for each n (see the energy equality (4.4)), the same is true of the limit function W(t). Therefore, since any non-increasing non-negative function has at most countably many points of discontinuity, we can apply the diagonal argument to account for such points and obtain

$$||u_{\varepsilon_n}(t)|| \to W(t) \qquad \text{for } t \ge 0, \tag{4.13}$$

 $^{^{32}}$ This theorem corresponds to Sections 28-31 of Leray (1934b).

³³This result is due to Helly (1912), see also Lemma 13.15 in Carothers (2000).

where we have also relabelled the sequence $\{\varepsilon_n\}$ and redefined W at its points of discontinuity. Note that since

$$u_{\varepsilon_n}(0) = J_{\varepsilon_n} u_0 \to u_0 \quad \text{in } L^2 \tag{4.14}$$

(see Lemma 1.2 (vi)) we obtain

$$W(0) = \lim_{n \to \infty} \|u_{\varepsilon_n}(0)\| = \lim_{n \to \infty} \|J_{\varepsilon_n} u_0\| = \|u_0\|.$$

We now want to take the limit of the functions themselves (rather than the norms). Since

$$\|J_{\varepsilon_n} u_{\varepsilon_n}(t)\| \le \|u_{\varepsilon_n}(t)\| \le \|u_0\|, \qquad \varepsilon_n > 0, t \ge 0$$

$$(4.15)$$

(see Lemma 1.2 (i) and (4.11)) we can apply the diagonal argument once more (and relabel the sequence $\{\varepsilon_n\}$) to deduce that the numbers

$$\int_{t_1}^{t_2} \int_{\Omega} (u_{\varepsilon_n})_k \quad \text{and} \quad \int_{t_1}^{t_2} \int_{\Omega} (u_{\varepsilon_n})_k (J_{\varepsilon_n} u_{\varepsilon_n})_l \quad \text{converge as } n \to \infty$$
(4.16)

for all $t_1, t_2 \in \mathbb{Q}^+$, k, l = 1, 2, 3 and all cubes $\Omega \subset \mathbb{R}^3$ with rational vertices, that is vertices whose coordinates are rational numbers. Moreover, from the timewise uniform continuity of the above integrals (that is from the bound $C(T)|t_2 - t_1|$ of the integrals whenever $t_1, t_2 \in (0, T)$ for some T) we obtain that in fact they converge as $n \to \infty$ for every pair $t_1, t_2 \ge 0$, every k, l = 1, 2, 3 and every cube Ω with rational vertices.

From the convergence in (4.16) we see that

$$\int_{t_1}^{t_2} \int u_{\varepsilon_n} \cdot (\partial_t f + \Delta f) \quad \text{and} \quad \int_{t_1}^{t_2} \int u_{\varepsilon_n} \cdot (J_{\varepsilon_n} u_{\varepsilon_n} \cdot \nabla) f \qquad (4.17)$$

converge as $n \to \infty$ for all $t_1, t_2 \ge 0$ and all test functions f (see Definition 4.5). Indeed, fix $t_1, t_2 \ge 0$, a test function f, and $\epsilon > 0$. Without loss of generality we can assume $t_1 < t_2$. There exists $N \in \mathbb{N}$, $\{g_k\}_{k=1}^N \subset \mathbb{R}^3$, $\{\Omega_k\}_{k=1}^N$ (cubes with rational coordinates), and a family of intervals

$$\{(p_k, q_k): 0 \le p_k < q_k \le \infty, \ k = 1, \dots, N\}$$

such that the (vector-valued) function

$$G_N(x,t) \coloneqq \sum_{k=1}^N g_k I_{\Omega_k}(x) I_{(p_k,q_k)}(t),$$

satisfies

$$\|(\partial_t f + \Delta f) - G_N\|_{L^{\infty}((t_1, t_2); L^2)} \le \frac{\epsilon}{4\|u_0\|(t_2 - t_1)}.$$
(4.18)

Moreover, the first part of (4.16) gives that for sufficiently large n, m

$$\left| \int_{t_1}^{t_2} \int (u_{\varepsilon_n} - u_{\varepsilon_m}) \cdot G_N \right| \le \frac{\epsilon}{2}$$

Thus

$$\begin{split} \left| \int_{t_1}^{t_2} \int (u_{\varepsilon_n} - u_{\varepsilon_m}) \cdot (\partial_t f + \Delta f) \right| \\ &\leq \left| \int_{t_1}^{t_2} \int (u_{\varepsilon_n} - u_{\varepsilon_m}) \cdot G_N \right| + \int_{t_1}^{t_2} \int (|u_{\varepsilon_n}| + |u_{\varepsilon_m}|) \left| (\partial_t f + \Delta f) - G_N \right| \\ &\leq \frac{\epsilon}{2} + \frac{\epsilon}{4 \|u_0\| (t_2 - t_1)} \int_{t_1}^{t_2} (\|u_{\varepsilon_n}(s)\| + \|u_{\varepsilon_m}(s)\|) \, \mathrm{d}s \\ &\leq \frac{\epsilon}{2} + \frac{\epsilon}{2 \|u_0\| (t_2 - t_1)} \int_{t_1}^{t_2} \|u_0\| \, \mathrm{d}s = \epsilon, \end{split}$$

where we also used the energy inequality (4.11). Thus we obtain the first part of (4.17). The second part follows in a similar way, by choosing a simple (matrix) function G_N such that

$$\|\nabla f - G_N\|_{L^{\infty}((t_1, t_2); L^{\infty})} \le \frac{\epsilon}{2\|u_0\|^2(t_2 - t_1)},$$

rather than (4.18). Thus we obtain (4.17).

On the other hand, the convergence $J_{\varepsilon_n}u_0 \to u_0$ in L^2 (see (4.14)) implies, in particular, the convergence of numbers

$$\int J_{\varepsilon_n} u_0 \cdot f(0) \to \int u_0 \cdot f(0) \qquad \text{for all test functions } f.$$

Combining this with (4.17) we can use the weak form of the regularised equations (4.12) to obtain that the numbers

$$\int u_{\varepsilon_n}(t) \cdot f(t) \quad \text{converge for all test functions } f \text{ and all } t \ge 0.$$

Thus letting $f(x,t) := \phi(x)$ for some $\phi \in L^2$ such that $\phi \in H^m$ for all $m \ge 1$ and div $\phi = 0$, we obtain that the numbers

$$\int u_{\varepsilon_n}(t) \cdot \phi \quad \text{converge for all } t \ge 0, \phi.$$
(4.19)

This together with the fact that $||u_{\varepsilon_n}(t)|| \leq ||u_0||$ (see (4.11)) implies that³⁴

$$u_{\varepsilon_n}(t) \to u(t) \qquad \text{as } n \to \infty \text{ in } L^2, \quad t \ge 0, \tag{4.20}$$

³⁴This functional analytical fact is one of Leray's remarkable contributions, which he discusses on page 209. To see it, note that if (4.20) does not hold then (using the boundedness in L^2) one could extract subsequences $\{n_k\}, \{m_k\}$ such that $u_{\varepsilon_{n_k}}(t) \rightarrow v(t), u_{\varepsilon_{m_k}}(t) \rightarrow w(t)$ for some $v(t), w(t) \in L^2, v(t) \neq w(t)$. In that case (4.19) gives $\int (u_{\varepsilon_{n_k}}(t) - u_{\varepsilon_{m_k}}(t))\phi \to 0$ for all ϕ , and taking $\phi \coloneqq J_{\delta}(v(t) - w(t))$ gives $\int (v(t) - w(t))J_{\delta}(v(t) - w(t)) = 0$, which in the limit $\delta \to 0^+$ gives ||v(t) - w(t)|| = 0, a contradiction.

for some $u(t) \in L^2$ (note $u(0) = u_0$ by (4.14) and u(t) is weakly divergence free for each t). We have thus constructed the sequence $\{\varepsilon_n\}$ and we obtained u as the weak limit of u_{ε_n} 's. In order to show that u is the required weak solution (Step 4), we first show that in fact $u_{\varepsilon_n}(t) \to u(t)$ strongly in L^2 for almost all t (Step 3).

Step 2. Define the set of singular times S.

Fatou's lemma and (4.11) give

$$\int_0^\infty \liminf_{\varepsilon_n \to 0} \|\nabla u_{\varepsilon_n}(t)\|^2 \, \mathrm{d}t \le \liminf_{\varepsilon_n \to 0} \int_0^\infty \|\nabla u_{\varepsilon_n}(t)\|^2 \, \mathrm{d}t \le \frac{1}{2} \|u_0\|^2.$$

Hence $\liminf_{\varepsilon_n\to 0} \|\nabla u_{\varepsilon_n}(t)\|^2 < \infty$ for almost every t > 0, that is |S| = 0, where

$$S \coloneqq \{t > 0 : \|\nabla u_{\varepsilon_n}(t)\| \to \infty \text{ as } n \to \infty\}$$

$$(4.21)$$

is the set of singular times.

Step 3. Show that $u_{\varepsilon_n}(t) \to u(t)$ strongly in L^2 for $t \in (0,\infty) \setminus S$.

Recalling that $W(t) = \lim_{\varepsilon_n \to 0} ||u_{\varepsilon_n}(t)||$ (see (4.13)) we see that it is enough to show that

$$||u(t)|| = W(t), \quad t \in (0,\infty) \setminus S, \tag{4.22}$$

since weak convergence (4.20) together with the convergence of the norms is equivalent to strong convergence. The estimate $||u(t)|| \leq W(t)$ holds for all $t \geq 0$ by the property of weak limits,

$$\|u(t)\| \le \liminf_{\varepsilon_n \to 0} \|u_{\varepsilon_n}(t)\| = W(t).$$
(4.23)

In order to prove the converse inequality, fix $t \in (0, \infty) \setminus S$. For such t let $\{\varepsilon_{n_k}\}$ be a subsequence along which $\liminf_{\varepsilon_n \to 0} \|\nabla u_{\varepsilon_n}(t)\|^2$ is attained, that is

$$\lim_{k \to \infty} \|\nabla u_{\varepsilon_{n_k}}(t)\| = \liminf_{\varepsilon_n \to 0} \|\nabla u_{\varepsilon_n}(t)\|.$$
(4.24)

It follows that the sequence $\{\nabla u_{\varepsilon_{n_k}}(t)\}$ is bounded in L^2 and therefore

$$\nabla u_{\varepsilon_{n_k}}(t) \rightharpoonup \nabla u(t) \text{ in } L^2$$

$$(4.25)$$

on a subsequence (which we relabel back to ε_{n_k} ; note on such a subsequence (4.24) still holds). The limit function is $\nabla u(t)$ by the definition of weak derivatives, and

$$\|\nabla u(t)\| \le \liminf_{\varepsilon_n \to 0} \|\nabla u_{\varepsilon_n}(t)\| \quad \text{for } t \in (0,\infty) \setminus S.$$
(4.26)

At this point we need to apply the separation of energy result (Lemma 4.3). We fix $\eta > 0$, and we let $R_1(\eta) > 0$ be large enough that

$$\int_{|x|>R_1(\eta)} |u_0|^2 \le \frac{\eta}{2},$$

and set

$$R_2(\eta, t) \coloneqq R_1(\eta) + \frac{4}{\eta}C(u_0, t),$$

where $C(u_0, t)$ is the constant from the lemma. The lemma now implies that

$$\int_{|x|>R_2(\eta,t)} |u_{\varepsilon}(t)|^2 \le \eta, \qquad \varepsilon > 0.$$
(4.27)

Since $u_{\varepsilon_{n_k}}(t) \rightharpoonup u(t)$ in H^1 (see (4.20) and (4.25)) we have in particular

$$u_{\varepsilon_{n_k}}(t) \rightharpoonup u(t) \quad \text{in } H^1(B(R_2(\eta, t))),$$

and so the compact embedding ^35 $H^1(B(R_2(\eta,t))) \subset \subset L^2(B(R_2(\eta,t)))$ implies

$$u_{\varepsilon_{n_k}}(t) \to u(t)$$
 in $L^2(B(R_2(\eta, t)))$.

Therefore, from (4.27), we obtain

$$\begin{split} \limsup_{\varepsilon_{n_k} \to 0} \|u_{\varepsilon_{n_k}}(t)\|^2 \\ &\leq \limsup_{\varepsilon_{n_k} \to 0} \int_{|x| \leq R_2(\eta, t)} |u_{\varepsilon_{n_k}}(t)|^2 + \limsup_{\varepsilon_{n_k} \to 0} \int_{|x| > R_2(\eta, t)} |u_{\varepsilon_{n_k}}(t)|^2 \\ &\leq \int_{|x| \leq R_2(\eta, t)} |u(t)|^2 + \eta \leq \|u(t)\|^2 + \eta \qquad \eta > 0, \end{split}$$

And hence, taking $\eta \to 0$,

$$W(t) = \limsup_{\varepsilon_{n_k} \to 0} \|u_{\varepsilon_{n_k}}(t)\| \le \|u(t)\|.$$

Thus we obtained the \geq inequality in (4.22), as required.

Step 4. Verify that u is a weak solution.

We already established (after (4.20)) that u(t) is weakly divergence free. Step 3 gives

$$\int u_{\varepsilon_n}(t) \cdot (\partial_t f(t) + \Delta f(t)) \to \int u(t) \cdot (\partial_t f(t) + \Delta f(t)),$$
$$\int u_{\varepsilon_n}(t) \cdot (J_{\varepsilon_n} u_{\varepsilon_n}(t) \cdot \nabla) f(t) \to \int u(t) \cdot (u(t) \cdot \nabla) f(t)$$

for almost every t > 0. Thus, using (4.15), we obtain (4.9) by taking the limit $\varepsilon_n \to 0^+$ in (4.12) and applying the Dominated Convergence Theorem to the time integrals. As for the energy inequality (4.10), let $s \in (0, \infty) \setminus S$

³⁵Leray's Lemma 2 provides a similar compactness result.

and t > s. Since $\|\nabla u(\tau)\| \le \liminf_{\varepsilon_n \to 0} \|\nabla u_{\varepsilon_n}(\tau)\|$ for almost every $\tau \ge 0$ (see (4.26)), we can use (4.23), Fatou's lemma, the identity

$$\|u_{\varepsilon}(t)\|^{2} + 2\int_{s}^{t} \|\nabla u_{\varepsilon}(s)\|^{2} \,\mathrm{d}s = \|u_{\varepsilon}(s)\|^{2}, \qquad \varepsilon > 0$$

(see (4.11)) and the fact that $\lim_{\varepsilon_n\to 0} ||u_{\varepsilon_n}(s)|| = ||u(s)||$ (see (4.22)) to obtain

$$\begin{aligned} \|u(t)\|^{2} + 2\int_{s}^{t} \|\nabla u(\tau)\|^{2} \,\mathrm{d}\tau \\ &\leq \liminf_{\varepsilon_{n} \to 0} \|u_{\varepsilon_{n}}(t)\|^{2} + 2\int_{s}^{t}\liminf_{\varepsilon_{n} \to 0} \|\nabla u_{\varepsilon_{n}}(\tau)\|^{2} \,\mathrm{d}\tau \\ &\leq \liminf_{\varepsilon_{n} \to 0} \left(\|u_{\varepsilon_{n}}(t)\|^{2} + 2\int_{0}^{t} \|\nabla u_{\varepsilon_{n}}(\tau)\|^{2} \,\mathrm{d}\tau \right) \\ &= \liminf_{\varepsilon_{n} \to 0} \|u_{\varepsilon_{n}}(s)\|^{2} = \|u(s)\|^{2} \end{aligned}$$

The case s = 0 follows similarly by using (4.14) (rather than (4.22)) in the last equality (recall also $u(0) = u_0$). Finally $||u(t) - u_0|| \to 0$ as $t \to 0$ follows from the weak convergence $u(t) \to u_0$ in L^2 (a consequence of (4.9)) and the convergence of norms $||u(t)|| \to ||u_0||$ (a consequence of Fatou's lemma and energy inequality with s = 0).

4.3 Structure of the weak solution

Let $u_0 \in H$ and consider the weak solution u given by Theorem 4.7. We first show that such solution enjoys the following *weak-strong uniqueness* property³⁶.

Lemma 4.9 (Weak-strong uniqueness). If $\|\nabla u(t_0)\| < \infty$ for some $t_0 \ge 0$ then u = v on the time interval $[t_0, T)$ for some $T > t_0$, where v is the semi-strong solution corresponding to the initial data $u(t_0)$ (recall Definition 3.15).

Proof. The assumption gives that $u(t_0) \in V$ (that is $u(t_0) \in H^1$ and $u(t_0)$ is divergence free, recall (1.2)) and so Theorem 3.17 gives the existence of a unique semi-strong solution v(t) on $[t_0, T)$, for some $T > t_0$. We will show that ${}^{37} u = v$ on time interval $[t_0, T)$.

 $^{^{36}}$ Leray calls this Comparison of a regular solution and a turbulent solution, see pp. 242-244.

 $^{^{37}}$ This is essentially the so-called *weak-strong uniqueness* property (see, for instance, Section 6.3 in Robinson et al. (2016)).

From the energy equality for v (see Lemma 3.16) and the energy inequality for u (see (4.10)), we obtain for a.e. $t_1 \in (t_0, T)$ and every $t \in (t_1, T)$

$$\|v(t)\|^{2} + 2\int_{t_{1}}^{t} \|\nabla v(s)\|^{2} \,\mathrm{d}s = \|v(t_{1})\|^{2},$$

$$\|u(t)\|^{2} + 2\int_{t_{1}}^{t} \|\nabla u(s)\|^{2} \,\mathrm{d}s \le \|u(t_{1})\|^{2}.$$

Adding these together and letting w := u - v gives

$$\|u(t_1)\|^2 + \|v(t_1)\|^2 \ge \|w(t)\|^2 + 2\int_{t_1}^t \|\nabla w(s)\|^2 \,\mathrm{d}s + 2\int u(t) \cdot v(t) + 4\int_{t_1}^t \int \nabla u : \nabla v \,\mathrm{d}s + 2\int_{t_1}^t \int \nabla u : \nabla v \,\mathrm{d}s + 2\int_{t_1}^t \int \nabla u : \nabla v \,\mathrm{d}s + 2\int_{t_1}^t \int \nabla u : \nabla v \,\mathrm{d}s + 2\int_{t_1}^t \int \nabla u : \nabla v \,\mathrm{d}s + 2\int_{t_1}^t \int \nabla u : \nabla v \,\mathrm{d}s + 2\int_{t_1}^t \int \nabla u : \nabla v \,\mathrm{d}s + 2\int_{t_1}^t \int \nabla u : \nabla v \,\mathrm{d}s + 2\int_{t_1}^t \int \nabla u : \nabla v \,\mathrm{d}s + 2\int_{t_1}^t \int \nabla u \,\mathrm{d}s + 2\int_{t_1$$

where $A: B \coloneqq A_{ij}B_{ij}$ denotes the inner product of matrices. Since v is strong on (t_0, T) (see Definition 3.15), it satisfies $\partial_t v - \Delta v + (v \cdot \nabla)v + \nabla p = 0$ in $(t_0, T) \times \mathbb{R}^3$ and it can be used as a test function for u on time interval (t_1, t) (see (4.9)) to write

$$\int u(t_1) \cdot v(t_1) = \int u(t) \cdot v(t) - \int_{t_1}^t \int u \cdot (\partial_t v + \Delta v + (u \cdot \nabla)v)$$
$$= \int u(t) \cdot v(t) - \int_{t_1}^t \int u \cdot (2\Delta v + (w \cdot \nabla)v - \nabla p)$$
$$= \int u(t) \cdot v(t) + \int_{t_1}^t \int v \cdot (w \cdot \nabla)w + 2\int_{t_1}^t \int \nabla u : \nabla v,$$

where in the last step we integrated by parts all three terms under the last integral and used the facts that u(t) and v(t) are weakly divergence free and that $\int v \cdot (w \cdot \nabla) v = 0$ (cf. (3.7)). Hence, using the inequality above and the Cauchy-Schwarz inequality we obtain

$$\begin{split} \|w(t_1)\|^2 &= \|u(t_1)\|^2 + \|v(t_1)\|^2 - 2\int u(t_1) \cdot v(t_1) \\ &\geq \|w(t)\|^2 + 2\int_{t_0}^t \|\nabla w(s)\|^2 \,\mathrm{d}s - 2\int_{t_1}^t \int v \cdot (w \cdot \nabla) \,w \\ &\geq \|w(t)\|^2 + 2\int_{t_1}^t \|\nabla w(s)\|^2 \,\mathrm{d}s - 2\int_{t_1}^t \|v(s)\|_\infty \|w(s)\| \|\nabla w(s)\| \,\mathrm{d}s \\ &\geq \|w(t)\|^2 - \frac{1}{2}\int_{t_1}^t \|v(s)\|_\infty^2 \|w(s)\|^2, \end{split}$$

where in the last step we used Young's inequality $ab \leq a^2 + b^2/4$. Hence Gronwall's inequality implies

$$||w(t)||^2 \le ||w(t_1)||^2 \exp\left(\frac{1}{2}\int_{t_1}^t ||v(s)||_{\infty}^2 \,\mathrm{d}s\right).$$

Since $\int_{t_0}^t ||v(s)||_{\infty} < \infty$ (see Definition 3.15) and since $t_0 \notin S$ both u and v are continuous in L^2 as $t_1 \to t_0^-$ (see Corollary 4.6) and we can take the limit $t_1 \to t_0^-$ to obtain

$$||w(t)||^2 \le ||w(t_0)||^2 \exp\left(\frac{1}{2}\int_{t_1}^t ||v(s)||_{\infty}^2 \,\mathrm{d}s\right) = 0 \qquad t \in (t_0, T).$$

Thus u(t) = v(t) for $t \in [t_0, T)$, as required.

We can use the weak-strong uniqueness to obtain that u, the weak solution given by Theorem 4.7, is regular in certain time intervals.

Definition 4.10. We call an open interval $(a, b) \subset (0, \infty)$ a maximal interval of regularity if u is a strong solution on (a, b) (see Definition 3.1) and u is not a strong solution on any open interval I strictly containing (a, b).

Theorem 4.11 (The structure of the weak solution³⁸). If u is a weak solution given by Theorem 4.7 then there exists a family of pairwise disjoint maximal intervals of regularity $(a_i, b_i) \subset (0, \infty)$ of u such that the set

$$\Sigma \coloneqq (0,\infty) \setminus \bigcup_i (a_i, b_i)$$

has measure zero.

Clearly $S \subset (0, \infty) \setminus \bigcup_i (a_i, b_i)$ (where S is the set of singular times of u, see Definition 4.5), since a strong solution is divergence free for all times and satisfies the energy equality (3.6). Therefore this theorems asserts that any $t_0 \in (0, \infty) \setminus S$ is an initial point of an interval in the interior of which u coincides with a strong solution, and the coincidence continues as long as the strong solution exists. Moreover, the energy inequality (4.10) shows that ||u(t)|| is strictly decreasing on every interval of regularity (a_i, b_i) . Note also that the family $\{(a_i, b_i)\}_i$ is at most countable (as a family of pairwise disjoint open intervals on \mathbb{R}).

Proof. Since for the weak solution u constructed in the last theorem the set S is given by (4.21), we see that if $t_0 \notin S$ then $\|\nabla u(t_0)\| < \infty$ and so Lemma 4.9 gives that t_0 either belongs to a maximal interval of regularity (see Definition 4.10) or is a left endpoint of one such interval. That is $t_0 \in \bigcup_i [a_i, b_i)$, and so

$$\Sigma \subset S \cup \bigcup_{i} \{a_i\}.$$

Since there are at most countably many maximal intervals of regularity we obtain $|\Sigma| = 0$, as required.

³⁸This theorem corresponds to Sections 32 and 33 in Leray (1934b).

Furthermore, it turns out that one of the intervals of regularity (a_i, b_i) of the weak solution u contains $(C||u_0||^4, \infty)$ and on this interval u enjoys a certain decay, which we make precise in the following theorem.

Theorem 4.12 (Supplementary information³⁹). Under the assumptions of the last theorem the set of singular times Σ is bounded above by $C||u_0||^4$, and for $t > C||u_0||^4$

$$\|\nabla u(t)\| < C \|u_0\| t^{-1/2},$$
 (4.28)

$$||u(t)||_{\infty} < C||u_0||t^{-3/4}.$$
(4.29)

Moreover

$$\sum_{i: b_i < \infty} \sqrt{b_i - a_i} \le C \|u_0\|^2.$$
(4.30)

Proof. If the union $\bigcup_i (a_i, b_i)$ has only one element $(0, \infty)$ then the first claim follows trivially. If not then let *i* be such that $b_i < \infty$ and let $t \in (a_j, b_j)$, where *j* is such that $b_j \leq b_i$ (that is (a_i, b_i) does not preceded (a_j, b_j) on the line). Since *u* becomes singular at b_j Corollary 3.12 gives

$$\|\nabla u(t)\| \ge C(b_j - t)^{-1/4} \ge C(b_i - t)^{-1/4}.$$
(4.31)

Applying this lower bound in the energy inequality (4.10) gives

$$||u_0||^2 \ge 2\int_0^{b_i} ||\nabla u(t)||^2 \, \mathrm{d}t \ge C\int_0^{b_i} (b_i - t)^{-1/2} \, \mathrm{d}s = C(b_i)^{1/2}.$$

Thus all finite b_i 's are bounded by $C||u_0||^4$ and the first claim follows. As for the last claim of the lemma apply the first inequality of (4.31) to the energy inequality (4.10) to obtain

$$||u_0||^2 \ge 2 \sum_{j: b_j < \infty} \int_{a_j}^{b_j} ||\nabla u(t)||^2 dt$$
$$\ge C \sum_{j: b_j < \infty} \int_{a_j}^{b_j} (b_j - t)^{-1/2} dt = C \sum_{j: b_j < \infty} \sqrt{b_j - a_j},$$

as required. It remains to show the decay estimates (4.28), (4.29). Let $s \in \bigcup_i (a_i, b_i)$ and t > s. Since u is strong on an interval containing s, we can use Corollary 3.14 to write that if $t - s \leq C \|\nabla u(s)\|^{-4}$ then

$$\|\nabla u(t)\| \le C \|\nabla u(s)\|,\tag{4.32}$$

$$||u(t)||_{\infty} \le C ||\nabla u(s)|| (t-s)^{-1/4}.$$
(4.33)

³⁹This is Section 34 in Leray (1934b).

From the first of these two inequalities we see that either $t-s \ge C \|\nabla u(s)\|^{-4}$ or $\|\nabla u(t)\| \le C \|\nabla u(s)\|$, and so

$$\|\nabla u(s)\| \ge C \min\left((t-s)^{-1/4}, \|\nabla u(t)\|\right).$$

Using this lower bound in the energy inequality (4.10) we obtain

$$||u_0||^2 \ge 2C \int_0^t \min\left((t-s)^{-1/2}, ||\nabla u(t)||^2\right) ds$$
$$\ge C \int_0^t \min\left(t^{-1/2}, ||\nabla u(t)||^2\right) ds$$
$$= C \min\left(t^{1/2}, t ||\nabla u(t)||^2\right).$$

Therefore, since the first argument of the minimum tends to ∞ as $t \to \infty$, we see that for $t^{1/2} > ||u_0||^2/C'$ we must have min $(t^{1/2}, t||\nabla u(t)||^2) = t||\nabla u(t)||^2$ and hence $||u_0||^2 \ge Ct||\nabla u(t)||^2$, from which (4.28) follows.

To obtain (4.29), let $s \ge ||u_0||^4/2(C')^2$ and t := 2s. Then $t^{1/2} \ge ||u_0||^2/C'$ and so using the above result gives

$$t - s = t/2 = \frac{1}{2} (t^{-1/4} t^{1/2})^4 \le C(||u_0||^{-1} t^{1/2})^4 \le C ||\nabla u(t)||^{-4}$$

Therefore we may apply (4.33) to obtain $||u(t)||_{\infty} \leq C ||\nabla u(s)||t^{-1/4}$. Now (4.29) follows by an application of (4.28),

$$\|u(t)\|_{\infty} \le C \|\nabla u(s)\| t^{-1/4} \le C \|u_0\| t^{-3/4}.$$

Finally, we remark on a consequence of (4.30).

Corollary 4.13. The set of singular times Σ satisfies

$$d_H(\Sigma) \le d_B(\Sigma) \le 1/2.$$

Here d_H denotes the Hausdorff dimension and d_B denotes the upper boxcounting dimension (see Sections 2.1 and 3.2 in Falconer (2014) for the definitions). This corollary does not appear in Leray's paper. Here we show it following the proof of a more general result by Lapidus & van Frankenhuijsen (2006) (see Theorem 1.10 therein); see also Besicovitch & Taylor (1954). An alternative approach can be found in the proof of Theorem 8.13 of Robinson et al. (2016).

Proof. Note that due to the general inequality $d_H(K) \leq d_B(K)$ for any compact $K \subset \mathbb{R}$ (see Lemma 3.7 in Falconer (2014)) it is enough to show the bound $d_B(\Sigma) \leq 1/2$. In this context the upper box-counting dimension is equivalent to the Minkowski-Bouligand dimension, which (for subsets of \mathbb{R}) is given by

$$d_B(\Sigma) = 1 - \liminf_{\delta \to 0^+} \frac{\log |\Sigma_\delta|}{\log \delta}, \qquad (4.34)$$

where Σ_{δ} denotes the δ -neighbourhood of Σ (see, for instance, Proposition 2.4 in Falconer (2014)). Since Σ is bounded there exists a unique index i_0 such that $b_{i_0} = \infty$, and so the set $\bigcup_{i \neq i_0} (a_i, b_i)$ is bounded by a_{i_0} . Thus

$$\Sigma = (0, a_{i_0}] \setminus \bigcup_i (a_i, b_i),$$

where now each b_i is finite. Thus we can renumber the intervals (a_i, b_i) such that the length of (a_i, b_i) does not increase with i, that is $b_i - a_i \ge b_{i+1} - a_{i+1}$ for all i.

For $\delta > 0$ let N_{δ} be such that

$$\begin{cases} b_i - a_i < 2\delta & i > N_\delta, \\ b_i - a_i \ge 2\delta & i \le N_\delta. \end{cases}$$

Observe that

$$\Sigma_{\delta} = \Sigma \cup \left(\bigcup_{i=1}^{N_{\delta}} (a_i, a_i + \delta) \cup (b_i - \delta, b_i)\right) \cup \bigcup_{i > N_{\delta}} (a_i, b_i).$$

Thus, since $|\Sigma| = 0$,

$$|\Sigma_{\delta}| \le 2\delta N_{\delta} + \sum_{i > N_{\delta}} (b_i - a_i) \le 2\delta N_{\delta} + \sqrt{2\delta} \sum_{i > N_{\delta}} \sqrt{b_i - a_i} \le 2\delta N_{\delta} + C\sqrt{\delta},$$
(4.35)

where we used the assumption $\sum_{i} \sqrt{b_i - a_i} \leq C$ (see the last inequality in Theorem 4.12). Now since for each k

$$k\sqrt{b_k - a_k} = \sum_{j=1}^k \sqrt{b_k - a_k} \le \sum_{j=1}^k \sqrt{b_j - a_j} \le C,$$

we see that $b_k - a_k < 2\delta$ for $k > C/\sqrt{2\delta}$. In other words $N_{\delta} \leq C/\sqrt{\delta}$. Hence (4.35) gives $|\Sigma_{\delta}| \leq C\sqrt{\delta}$ and so (since log $\delta < 0$ for small $\delta > 0$)

$$\frac{\log |\Sigma_{\delta}|}{\log \delta} \ge \frac{\log C + \frac{1}{2}\log \delta}{\log \delta} \to \frac{1}{2}$$

as $\delta \to 0^+$. Therefore $d_B(\Sigma) \leq 1/2$, as required.

It is interesting to note that apart from the bound $d_H(\Sigma) \leq 1/2$ a stronger property $\mathcal{H}^{1/2}(\Sigma) = 0$ holds. This can be shown by an (earlier) argument due to Scheffer (1976), which is similar to the above and is also presented by Robinson (2006), see Proposition 8 therein.

Notes

This section corresponds to Chapters V and VI of Leray (1934b), except that Leray did not discuss the dimension of the set of singular times. In fact the notion of the box-counting dimension was not properly unified in the 1930's, although some variants were already being studied at the time (see Bouligand (1928) and Pontrjagin & Schnirelmann (1932) for instance). On the other hand, the notion of the Hausdorff dimension was already quite well developed (see Hausdorff (1918), Besicovitch (1935)), but it is not apparent whether or not Leray was aware of these developments. In any case, it was Scheffer (1976) who was the first to study the dimension of the set of singular times for the Navier–Stokes equations.

Acknowledgements

We would like to thank Robert Terrell for his English translation of Leray's original work, and Simon Baker, Tobias Barker, Antoine Choffrut and Kenneth Falconer for their helpful comments. We are particularly grateful to James Robinson for his enthusiasm, encouragement and interest in this work.

WSO is supported by EPSRC as part of the MASDOC DTC at the University of Warwick, Grant No. EP/HO23364/1.

BCP was partially supported by an EPSRC Doctoral Training Award and partially by postdoctoral funding from ERC 616797.

A Appendix

A.1 The heat equation and the heat kernel

Let $v_0 \in L^2$ and $v(t) := \Phi(t) * v_0$, where $\Phi = (4\pi t)^{-3/2} e^{-|x|^2/4t}$ is the heat kernel.

If v_0 is a vector-valued function which is weakly divergence free then div v(t) = 0 for each t > 0 (which can be shown directly by approximating $\nabla \Phi(x - y, t)$ in L^2 by the gradient of a smooth and compactly supported gat each t). Furthermore for $m \ge 1$, v satisfies

- (i) $v \in C([0,T); L^2)$ with $v(0) = v_0$, and $||v(t)|| \le ||v_0||$,
- (ii) $\nabla^m v \in C((0,T);L^2)$ and

$$\|\nabla^m v(t)\| \le C_m \|v_0\| t^{-m/2}.$$

Moreover if $\nabla^m v_0 \in L^2$ then $\nabla^m v \in C([0,T); L^2)$ with $\|\nabla^m v(t)\| \leq C_m \min_{k \leq m} \{\|\nabla^k v_0\| t^{-(m-k)/2}\}.$

(iii) $v \in C((0,T); L^{\infty})$ and $||v(t)||_{\infty} \leq C \min\{||v_0||t^{-3/4}, ||v_0||_{\infty}\},\$

(iv) $\nabla^m v \in C((0,T); L^\infty)$ and

$$\|\nabla^m v\|_{\infty} \le C_m \min\{\|v_0\|t^{-m/2-3/4}, \|v_0\|_{\infty}t^{-m/2}\}$$

(v) $v, \nabla^m v \in \mathcal{H}^{1/2}((0,T))$ with the corresponding constants $C_0(t) \leq c_0 ||v_0||/t$, $C_m(t) \leq c_m ||v_0|| t^{-1-m/2}$.

The inequalities in (i)-(iv) follow directly from Young's inequality (1.5) and the bounds $\|\nabla^m \Phi(t)\|_1 \leq C_m t^{-m/2}$, $\|\nabla^m \Phi(t)\| \leq C_m t^{-m/2-3/4}$, $m \geq 0$ (which can be verified by a direct calculation). The claims regarding continuity in (0,T) follow from the fact that Φ and all its spacial derivatives $\nabla^m \Phi$, $m \geq 1$, belong to $C((0,T); L^1) \cap C((0,T); L^2)$ (a consequence of the Dominated Convergence Theorem). The claims regarding continuity at t = 0 in (i) and (ii) are known as the *approximation of identity*, see e.g. Theorem 1.18 in Chapter 1 of Stein & Weiss (1971) for a proof. Finally, property (v) follows from Morrey's inequality (see, for example, Section 4.5.3 in Evans & Gariepy (2015)) and Young's inequality for convolutions (1.5) by writing

$$\|\Phi(t) * v_0\|_{C^{0,1/2}} \le C \|\nabla\Phi(t) * v_0\|_6 \le C \|\nabla\Phi(t)\|_{3/2} \|v_0\| = C \|v_0\|/t.$$
(A.1)

The claim for the derivatives follows similarly by noting that

$$\|\nabla^{m+1}\Phi(t)\|_{3/2} = C_m t^{-1-m/2}.$$

As for the pointwise convergence as $t \to 0^+$ we have

Lemma A.1 (pointwise convergence as $t \to 0^+$ of the heat flow). If $x_0 \in \mathbb{R}^3$ is a continuity point of v_0 then

$$v(x_0, t) \to v_0(x_0)$$
 as $t \to 0^+$.

If v_0 is bounded and uniformly continuous then

$$\|v(t) - v_0\|_{\infty} \to 0 \qquad \text{as } t \to 0^+.$$

See Section 4.2 in Giga et al. (2010) for a proof.

One can verify that v(t) is a solution of the heat equation $v_t = \Delta v$ in $\mathbb{R}^3 \times (0, \infty)$ by a direct calculation. It is a unique such solution in the class of functions $C([0, \infty); L^2)$ satisfying $v(0) = v_0$ for some $v_0 \in L^2$, which we make precise in the following lemma.

Lemma A.2. Let $w \in C([0,T); L^2)$ be a weak solution of the heat equation with w(0) = 0, that is

$$\int_0^T \int (\phi_t + \Delta \phi) w \, \mathrm{d}x \, \mathrm{d}t = 0 \tag{A.2}$$

for all $\phi \in C_0^{\infty}(\mathbb{R}^3 \times [0,T))$. Then $w \equiv 0$.

Proof. We modify the argument from Section 4.4.2 of Giga et al. (2010). We focus on the case $T < \infty$ (the case $T = \infty$ follows trivially by applying the result for all T > 0).

We first show that assumption $w \in C([0,T); L^2)$ implies that (A.2) holds also for all $\phi \in C^{\infty}(\mathbb{R}^3 \times [0,T))$ such that

$$\begin{cases} \sup_{t \in [0,T)} \left(\|\partial_t \phi(t)\| + \|D^{\alpha} \phi(t)\| \right) \le C & \text{ for some } C > 0, \\ \sup \phi \subset \mathbb{R}^3 \times [0,T') & \text{ for some } T' < T. \end{cases}$$
(*)

Indeed, let $\theta \in C^{\infty}(\mathbb{R}; [0, 1])$ be such that $\theta(\tau) = 1$ for $\tau \leq 1$ and $\theta(\tau) = 0$ for $\tau \geq 2$ (take for instance $\theta(\tau) \coloneqq q(2-\tau)/(q(2-\tau)+q(1-\tau))$, where $q(s) \coloneqq e^{-1/s}$ for s > 0 and $q(s) \coloneqq 0$ otherwise). For $x \in \mathbb{R}^3$ Let $\theta_j(x) \coloneqq \theta(|x|/j)$. Then $\theta_j \in C_0^{\infty}$, for some $M > 0 \|\nabla \theta_j\|_{\infty} \leq M/j$ and $\|D^2 \theta_j\|_{\infty} \leq M/j^2$, and

$$\theta_j(x) = \begin{cases} 1 & |x| < j, \\ 0 & |x| > 2j \end{cases}$$

Now for $\phi \in C^{\infty}(\mathbb{R}^3 \times [0,T))$ satisfying (*) let T' < T be such that $\phi(t) \equiv 0$ for $t \geq T'$ and let

$$\phi_j(x,t) \coloneqq \theta_j(x)\phi(x,t).$$

Then $\phi_j \in C_0^{\infty}(\mathbb{R}^3 \times [0,T))$ and so (A.2) gives

$$\int_0^T \int (\partial_t \phi_j + \Delta \phi_j) w \, \mathrm{d}x \, \mathrm{d}t = 0,$$

or equivalently

$$\int_0^T \int \theta_j (\partial_t \phi + \Delta \phi) w \, \mathrm{d}x \, \mathrm{d}t + \int_0^T \int w \left(2\nabla \theta_j \cdot \nabla \phi + \phi \Delta \theta_j \right) \, \mathrm{d}x \, \mathrm{d}t = 0,$$
(A.3)

Since θ_j converges pointwise to 1 and

$$\begin{split} \int_0^T \int |(\partial_t \phi + \Delta \phi) w| \, \mathrm{d}x \, \mathrm{d}t &\leq \int_0^T \|\partial_t \phi(t) + \Delta \phi(t)\| \| \|w(t)\| \, \mathrm{d}t \\ &\leq TC \sup_{t \in [0, T']} \|w(t)\| < \infty \end{split}$$

the Dominated Convergence Theorem gives that the first integral on the left-hand side of (A.3) converges to $\int_0^T \int (\partial_t \phi + \Delta \phi) w \, dx \, dt$. The second integral is bounded by

$$\begin{split} \int_0^T \int |w| \, |2\nabla\theta_j \cdot \nabla\phi + \phi \Delta\theta_j| \, \mathrm{d}x \, \mathrm{d}t \\ &\leq T \sup_{t \in [0,T']} \|w(t)\| (2CM/j + CM/j^2) \xrightarrow{j \to \infty} 0 \end{split}$$

Therefore taking the limit $j \to \infty$ in (A.3) gives

$$\int_0^T \int (\partial_t \phi + \Delta \phi) w \, \mathrm{d}x \, \mathrm{d}t = 0$$

as required.

We will show that

$$\int_0^T \int \Psi w \, \mathrm{d}x \, \mathrm{d}t = 0$$

for all $\Psi \in C_0^{\infty}(\mathbb{R}^3 \times (0,T))$. This finishes the proof as C_0^{∞} is dense in L^2 and so $\int_0^T \int vw \, dx \, dt = 0$ for all $v \in L^2(\mathbb{R}^3 \times (0,T))$. Hence $w \equiv 0$. Given $\Psi \in C_0^{\infty}(\mathbb{R}^3 \times (0,T))$ let T' < T be such that $\Psi(t) \equiv 0$ for $t \geq T'$.

Given $\Psi \in C_0^{\infty}(\mathbb{R}^3 \times (0, T))$ let T' < T be such that $\Psi(t) \equiv 0$ for $t \geq T'$. Extend Ψ by zero for $t \leq 0$ and $t \geq T$. There exists a classical solution ϕ to the problem

$$\begin{cases} \phi_t + \Delta \phi = \Psi & \text{in } \mathbb{R}^3 \times (-\infty, T), \\ \phi(x, T) = 0 & \text{for } x \in \mathbb{R}^3. \end{cases}$$
(A.4)

Indeed, denoting $\tilde{\phi}(x,t) = \phi(x,T-t)$ and $\tilde{\Psi}(x,t) = \Psi(x,T-t)$ the above problem becomes

$$\begin{cases} \widetilde{\phi}_t - \Delta \widetilde{\phi} = -\widetilde{\Psi} & \text{ in } \mathbb{R}^3 \times (0, \infty), \\ \widetilde{\phi}(x, 0) = 0 & \text{ for } x \in \mathbb{R}^3. \end{cases}$$

A classical solution ϕ is obtained by an application of the Duhamel principle,

$$\widetilde{\phi}(t) \coloneqq -\int_0^t \Phi(t-s) * \widetilde{\Psi}(s) \,\mathrm{d}s,$$

and such ϕ is smooth (see for example Section 4.3.2 of Giga et al. (2010)). Observe that, since $\Psi(s) \equiv 0$ for $s \geq T'$, we have that $\phi(t) \equiv 0$ for $t \in [0, T - T']$. Moreover, since Ψ is smooth and compactly supported in $\mathbb{R}^3 \times \mathbb{R}$ we have for $|\alpha| \leq 2$

$$D^{\alpha}\widetilde{\phi}(t) = -\int_{0}^{t} \Phi(t-s) * D^{\alpha}\widetilde{\Psi}(s) \,\mathrm{d}s,$$

and it follows from Lemma 1.1 that $D^{\alpha} \widetilde{\phi} \in C([0,T], L^2)$ (note that $\Phi \in L^1_{\text{loc}}([0,\infty); L^1)$ due to the bound $\|\Phi(t)\|_1 \leq C_0$). Therefore, we also obtain $\widetilde{\phi}_t = \Delta \widetilde{\phi} - \widetilde{\Psi} \in C([0,T]; L^2)$. Hence $\phi(x,t) = \widetilde{\phi}(x,T-t)$ is smooth, $\phi(t) \equiv 0$ for $t \in [T',T]$, and

$$\sup_{t \in [0,T)} \left(\|\partial_t \phi(t)\| + \|D^{\alpha} \phi(t)\| \right) = \sup_{t \in [0,T]} \left(\|\partial_t \widetilde{\phi}(t)\| + \|D^{\alpha} \widetilde{\phi}(t)\| \right),$$

that is ϕ satisfies (*). We can therefore use the first part and (A.4) to write

$$0 = \int_0^T \int (\phi_t + \Delta \phi) w \, \mathrm{d}x \, \mathrm{d}t = \int_0^T \int \Psi w \, \mathrm{d}x \, \mathrm{d}t.$$

A.2 The extension of Young's inequality for convolutions

We will prove the following lemma (Lemma 1.1).

Lemma A.3. If $p,q,r \ge 1$ are such that 1/q = 1/p + 1/r - 1, $A \in L^1_{loc}([0,T); L^p)$ and $B \in C((0,T); L^r)$ with $||B(t)||_r$ bounded as $t \to 0^+$ then u defined by

$$u(t) := \int_0^t A(t-s) * B(s) \,\mathrm{d}s$$

belongs to $C([0,T); L^q)$ and

$$\|u(t)\|_{q} \le \int_{0}^{t} \|A(t-s)\|_{p} \|B(s)\|_{r} \,\mathrm{d}s.$$
(A.5)

Proof. The bound (A.5) is clear from the integral version of Minkowski inequality (1.3) and from Young's inequality (1.5). This bound also implies the continuity $u(t) \to 0$ in L^q as $t \to 0^+$. It remains to show that $u \in C((0,T); L^q)$. For this reason let T' < T and $\varepsilon > 0$. Let M > 0 be such that

$$\max_{s \in [0,T']} \|B(s)\|_r \le M, \quad \int_0^{T'} \|A(s)\|_p \, \mathrm{d}s \le M.$$

From the assumption on A we see that there exist $\eta > 0$ such that

$$\int_0^\eta \|A(s)\|_p \,\mathrm{d} s \le \varepsilon.$$

Since B is uniformly continuous into L^r on $[\eta/2,T']$ there exists $\delta\in(0,\eta/2)$ such that

$$||B(s) - B(t)||_r < \varepsilon$$

whenever $s, t \in [\eta/2, T']$, t > s and $t - s < \delta$. Letting $t_1, t_2 \in (0, T']$ be such that $t_1 < t_2, t_2 - t_1 < \delta$ we obtain

$$u(t_{2}) - u(t_{1}) = \int_{0}^{t_{2}} A(t_{2} - s) * B(s) ds - \int_{0}^{t_{1}} A(t_{1} - s) * B(s) ds$$

$$= \int_{0}^{t_{2}} A(t_{2} - s) * B(s) ds - \int_{t_{2} - t_{1}}^{t_{2}} A(t_{2} - s) * B(s - (t_{2} - t_{1})) ds$$

$$= \int_{0}^{t_{2} - t_{1}} A(t_{2} - s) * B(s) ds$$

$$+ \int_{t_{2} - t_{1}}^{t_{2}} A(t_{2} - s) * (B(s) - B(s - (t_{2} - t_{1}))) ds.$$
(A.6)

Thus, using the integral Minkowski inequality (1.3) and Young's inequality for convolutions (1.5) we obtain

$$\|u(t_2) - u(t_1)\|_q \le \int_0^{t_2 - t_1} \|A(t_2 - s)\|_p \|B(s)\|_r \,\mathrm{d}s$$

+ $\int_{t_2 - t_1}^{t_2} \|A(t_2 - s)\|_p \|B(s) - B(s - (t_2 - t_1))\|_r \,\mathrm{d}s$

Since $t_2 - t_1 < \delta < \eta$ the first term on the right-hand side is bounded by $M\varepsilon$. As for the second term, if $t_2 < \eta$ it can be similarly bounded by $2M\varepsilon$. If not, then we decompose it into $\int_{t_2-t_1}^{\eta} + \int_{\eta}^{t_2}$, we bound the first of the resulting integrals by $2M\varepsilon$ and, as for the second one, observe that $s, s - (t_2 - t_2) \in [\eta/2, T']$ whenever $s \in [\eta, T']$ to write

$$\int_{\eta}^{t_2} \|A(t_2 - s)\|_p \|B(s) - B(s - (t_2 - t_1))\|_r \, \mathrm{d}s$$
$$\leq \varepsilon \int_{\eta}^{t_2} \|A(t_2 - s)\|_p \, \mathrm{d}s \leq \varepsilon M.$$

Therefore $||u(t_2) - u(t_1)||_q \le 4M\varepsilon$.

A.3 Decay estimates of P(x,t)

Let $P(x,t) := \frac{1}{|x|} \int_0^{|x|} \frac{e^{-\xi^2/4t}}{t^{1/2}} d\xi$ (see (1.10)). We have the following decay estimates.

Theorem A.4.

$$|\nabla^m P(x,t)| \le \frac{C_m}{(|x|^2 + t)^{(m+1)/2}}$$
(A.7)

for all $x \in \mathbb{R}^3$, t > 0.

Proof. Let us first assume that t = 1 (the general case will follow from the rescaling $P(x,t) = \frac{1}{\sqrt{t}}P(x/\sqrt{t},1)$). We claim that any partial derivative $D^{\alpha}P(x,1)$ of order $|\alpha| = m$ is of the form

$$D^{\alpha}P(x,1) = \frac{Q_{\alpha}(x)}{|x|^{2m}}P(x,1) + E_{\alpha}(x)$$
(A.8)

for $m \ge 0$ and |x| > 1, where $Q_{\alpha}(x)$ denotes a polynomial of degree less than or equal to $|\alpha| = m$ and $E_{\alpha}(x)$ denotes a function that is smooth in |x| > 1and whose derivatives of all orders decay exponentially when $|x| \to \infty$. The case m = 0 follows trivially with $E_0 \equiv 0$. For the inductive step, assume (A.8) holds for all partial derivatives $D^{\alpha}P(x,t)$ with $|\alpha| = m \ge 0$ and write

$$\begin{split} \partial_{x_i} D^{\alpha} P(x,1) &= \partial_{x_i} \left(\frac{Q_{\alpha}(x)}{|x|^{2m}} P(x,1) + E_{\alpha}(x) \right) \\ &= \frac{\tilde{Q}_{m-1}(x)}{|x|^{2m}} P(x,1) - \frac{2mQ_{\alpha}(x)x_i}{|x|^{2m+2}} P(x,1) \\ &+ \frac{Q_{\alpha}(x)}{|x|^{2m}} \left(\frac{x_i}{|x|^2} P(x,1) + \frac{x_i}{|x|^2} e^{-|x|^2/4} \right) + \partial_{x_i} E_{\alpha}(x) \\ &= \frac{P(x,1)}{|x|^{2m+2}} \left(|x|^2 \tilde{Q}_{m-1}(x) - (2m-1)Q_{\alpha}(x)x_i \right) \\ &+ \frac{Q_{\alpha}(x)x_i}{|x|^{2m+2}} e^{-|x|^2/4} + \partial_{x_i} E_{\alpha}(x) \end{split}$$

for i = 1, 2, 3, where $\tilde{Q}_{m-1}(x)$ denotes some polynomial of degree less than or equal to m-1. Clearly, the last bracket is some polynomial of degree less than or equal to m+1 and the remaining two terms are smooth in |x| > 1and decay exponentially as $|x| \to \infty$. Hence the induction follows. Because P(x, 1) decays like $|x|^{-1}$ as $|x| \to \infty$ we see from (A.8) that

$$|\nabla^m P(x,1)| \le \frac{C_m}{|x|^{m+1}} \le \frac{C_m}{(|x|^2+1)^{(m+1)/2}}$$

holds for $|x| \ge 2$. As $P(\cdot, 1)$ is a smooth function (see Section 1.2) we have $|\nabla^m P(x, 1)| \le C_m \le C_m (|x|^2 + 1)^{-(m+1)/2}$ for all $x \in B(0, 2)$. Hence (A.7) follows in the case t = 1. Finally, the rescaling $P(x, t) = \frac{1}{\sqrt{t}} P(x/\sqrt{t}, 1)$ yields

$$\begin{aligned} |\nabla^m P(x,t)| &= \left| t^{-(m+1)/2} \left[\nabla^m P(y,1) \right]_{y=x/\sqrt{t}} \right| \\ &\leq t^{-(m+1)/2} \frac{C_m}{\left(\left| x/\sqrt{t} \right|^2 + 1 \right)^{(m+1)/2}} = \frac{C_m}{\left(\left| x \right|^2 + t \right)^{(m+1)/2}}. \quad \Box \end{aligned}$$

A.4 Properties of the Stokes equations

Here we present proofs of some results from Section 2. Namely we complete the proof of Theorem 2.2, show property (iii) of the representation (2.3) and the continuity $\nabla u \in C((0,T); L^{\infty})$ given representation (2.14) with $Y \in \mathcal{H}^{1/2}((0,T))$, and we show uniqueness of distributional solutions of the Stokes equations (Theorem 2.5).

A.4.1 Proof of Theorem 2.2

Recall that it remains to verify that if for some R > 0

$$F \in C^{\infty}(\mathbb{R}^3 \times [0,T);\mathbb{R}^3)$$
 and $\operatorname{supp} F(t) \subset B(0,R)$ for $t \in [0,T)$

then the pair of functions u_2 , p given by (2.3), (2.4) is a classical solution of the problem

$$\begin{cases} \partial_t u_2 - \Delta u_2 + \nabla p &= F, \\ \operatorname{div} u_2 &= 0, \\ u_2(0) &= 0, \end{cases}$$

and $u_2 \in C([0,T); L^2)$. Indeed, as remarked after the statement of Theorem 2.2, then $u = u_1 + u_2$ is a classical solution of the Stokes equations (2.1), (2.2) with $u(0) = u_0$ and $u \in C([0,T); L^2)$.

Note that, since F is smooth and compactly supported in space and since $\mathcal{T} \in L^1_{\text{loc}}([0,T); L^2)$ (see (1.14)) we can use Lemma 1.1 to obtain $u_2 \in C([0,T); L^2)$ and $||u_2(t)|| \to 0$ as $t \to 0^+$ (which means that u_2 satisfies the initial condition $u_2(0) = 0$). Moreover, since $\mathcal{T} \in C((0,\infty); L^2)$ (see (1.15)), we deduce that the functions ∇u_2 , Δu_2 , $\partial_t u_2$ are continuous (an application of the Dominated Convergence Theorem) and belong to $C((0,T); L^2)$ (a consequence of Lemma 1.1). Similarly ∇p is continuous and $\nabla p \in C([0,T); L^2)$ (by an application of the Plancherel Lemma (Lemma 1.4)). Therefore, since the Fourier transform is an isometry from L^2 into L^2 , we see that u_2 , p satisfy the claim above if and only if

$$\partial_t \widehat{u_2}(\xi, t) + 4\pi^2 |\xi|^2 \widehat{u_2}(\xi, t) + 2\pi i \,\xi \, \widehat{p}(\xi, t) = \widehat{F}(\xi, t) \tag{A.9}$$

and

$$\xi \cdot \widehat{u_2}(\xi, t) = 0 \tag{A.10}$$

hold for t > 0 and almost every $\xi \in \mathbb{R}^3$. Here $\widehat{u_2}$, \widehat{p} , \widehat{F} denote the Fourier transform \mathcal{F} of u_2 , p, F, respectively. Since p satisfies $\Delta p = \operatorname{div} F$ we obtain

$$2\pi \mathrm{i}\,\xi\,\widehat{p}(\xi,t) = \frac{\xi\otimes\xi}{|\xi|^2}\widehat{F}(\xi,t),$$

where $\xi \otimes \xi$ denotes the 3 × 3 matrix with components $\xi_i \xi_j$. Therefore (A.9) is equivalent to

$$\left(\partial_t + 4\pi^2 |\xi|^2\right) \widehat{u_2}(\xi, t) = \left(I - \frac{\xi \otimes \xi}{|\xi|^2}\right) \widehat{F}(\xi, t).$$
(A.11)

Now since

$$\mathcal{F}[\Phi(\cdot,t)] = e^{-4\pi^2 t |\xi|^2}$$
 (A.12)

(see e.g. Theorem 1.13 in Chapter 1 of Stein & Weiss (1971) for a proof of this fact) and $-\Delta P = \Phi$ (see (1.11)), we obtain

$$\mathcal{F}[P(\cdot,t)] = \frac{1}{4\pi^2 |\xi|^2} e^{4\pi^2 t |\xi|^2},$$

and so

$$\mathcal{F}\left[\partial_i \partial_j P(\cdot, t)\right] = \frac{-\xi_i \xi_j}{|\xi|^2} e^{-4\pi^2 t |\xi|^2}.$$

Hence for all t > 0

$$\mathcal{F}[\mathcal{T}(t)] = \left(I - \frac{\xi \otimes \xi}{|\xi|^2}\right) e^{-4\pi^2 t |\xi|^2}.$$

Since $u_2(t) = \int_0^t \mathcal{T}(t-s) * F(s) \, \mathrm{d}s$,

$$\widehat{u_2}(\xi,t) = \int_0^t \left(I - \frac{\xi \otimes \xi}{|\xi|^2}\right) \widehat{F}(\xi,s) \mathrm{e}^{-4\pi^2(t-s)|\xi|^2} \,\mathrm{d}s, \qquad \xi \neq 0, \qquad (A.13)$$

and so (A.10) holds for $\xi \neq 0$ and

$$(\partial_t + 4\pi^2 |\xi|^2) \,\widehat{u_2}(\xi, t)$$

$$= \left(\partial_t + 4\pi^2 |\xi|^2\right) \int_0^t \left(I - \frac{\xi \otimes \xi}{|\xi|^2}\right) e^{-4\pi^2 (t-s)|\xi|^2} \widehat{F}(\xi, s) \,\mathrm{d}s$$

$$= \left(I - \frac{\xi \otimes \xi}{|\xi|^2}\right) \widehat{F}(\xi, t)$$

for $\xi \neq 0$, that is (A.9), as claimed.

A.4.2 Property (iii) of the representation (2.3)

We need to show that if $F \in C([0,T); L^2)$, $u_0 \in L^2$ and u is given by (2.3),

$$u(t) = u_1(t) + u_2(t) = \Phi(t) * u_0 + \int_0^t \mathcal{T}(t-s) * F(s) \, \mathrm{d}s,$$

then $u \in C([0,T);L^2)$ with

$$||u(t)|| \le \int_0^t ||F(s)|| \, \mathrm{d}s + ||u_0|| \quad \text{for } t \in (0,T).$$
 (A.14)

Moreover u satisfies the energy dissipation equality

$$\|u(t)\|^{2} - \|u_{0}\|^{2} + 2\int_{0}^{t} \|\nabla u(s)\|^{2} \,\mathrm{d}s = 2\int_{0}^{t} \int u \cdot F \,\mathrm{d}x \,\mathrm{d}s \tag{A.15}$$

for $t \in (0, T)$.

We prove these claim by considering two cases.

Case 1. F is regular, that is satisfies (2.9) for some R > 0. For such F, due to Theorem 2.2 (or rather to the proof above), $u = u_1 + u_2$ and p satisfy $u(0) = u_0$,

$$\partial_t u_1 - \Delta u_1 = 0, \quad \partial_t u_2 - \Delta u_2 + \nabla p = F \qquad \text{in } \mathbb{R}^3 \times (0, T) \qquad (A.16)$$

and $u_1, u_2 \in C([0,T); L^2)$, where p is given by (2.4).

Note that for $y \in B(0, R)$ and |x| > 2R

$$|x|/2 < |x| - R < |x - y| < |x| + R$$

and using (1.13) we can write for such x and for all t

$$\begin{aligned} |u_2(x,t)| &\leq \int_0^t \int_{B(0,R)} \frac{C_0}{(|x-y|^2+t-s)^{3/2}} |F(y,s)| \, \mathrm{d}y \, \mathrm{d}s \\ &\leq \frac{C|B(0,R)|^{1/2}t}{|x|^3} \max_{s \in [0,t]} \|F(s)\|, \end{aligned}$$

which gives a decay $\sim |x|^{-3}$ of $|u_2(x,t)|$ as $|x| \to \infty$ that is uniform on any compact time interval [0,T'], where T' < T. Similarly one can derive a decay $\sim |x|^{-5}$ of $|\Delta u_2(x,t)|$ and decay $\sim |x|^{-3}$ of $|\nabla p(x,t)|$. Thus (A.16) gives the decay $\sim |x|^{-3}$ of $\partial_t u_2(x,t)$. Letting $\delta, t \in (0,T), t > \delta$ and using this decay as well as the fact that u_1 and all its derivatives belong to $C((0,T); L^2)$ (see (ii) in Appendix A.1) we can integrate the equality

$$u \cdot \partial_t u - u \cdot \Delta u + u \cdot \nabla p = u \cdot F$$

on $\mathbb{R}^3 \times (\delta, t)$ to obtain

$$||u(t)||^{2} - ||u(\delta)||^{2} + 2\int_{\delta}^{t} ||\nabla u(s)||^{2} \,\mathrm{d}s = 2\int_{\delta}^{t} \int u \cdot F \,\mathrm{d}x \,\mathrm{d}s.$$
(A.17)

Since $u \in C([0, T); L^2)$ we can take the limit $\delta \to 0^+$ (and apply the Monotone Convergence Theorem) to obtain (A.15). As for the bound (A.14) note that the above equality and the Cauchy-Schwarz inequality give

$$\frac{\mathrm{d}}{\mathrm{d}t} \|u(t)\|^2 \le 2 \int u(t) \cdot F(t) \, dx \le 2 \|u(t)\| \, \|F(t)\|, \qquad t \in (0,T),$$

that is

$$\frac{\mathrm{d}}{\mathrm{d}t} \|u(t)\| \le \|F(t)\|, \qquad t \in (0,T).$$

Integrating this inequality in t gives (A.14).

Case 2. $F \in C([0,T); L^2)$. Let u be the velocity field corresponding to F and the initial velocity field u_0 , and consider a compact time interval $[0,T'] \subset [0,T)$. For such F consider a sequence $\{F_R\} \subset C([0,T']; L^2)$ such that $F_R \in C^{\infty}(\mathbb{R}^3 \times \mathbb{R})$, supp $F_R(t) \subset B(0,R)$ for all t and

$$F_R \to F$$
 in $C([0, T']; L^2)$ as $R \to \infty$.

(See Lemma A.12 for a proof of the existence of such a sequence.) Let u_R denote the velocity field corresponding to F_R and to the initial velocity u_0 . Since the representation formula (2.3) is linear we can apply (A.14)

to the difference $u_{R_1} - u_{R_2}$ for $R_1, R_2 > 0$ to see that $\{u_R\}$ is Cauchy in $C([0, T']; L^2)$ as $R \to \infty$. Thus $u_R \to u'$ in $C([0, T']; L^2)$ as $R \to \infty$ for some $u' \in C([0, T']; L^2)$. However, Lemma 2.1 (i) applied to the difference $u_R - u$ gives $||u(t) - u_R(t)||_{\infty} \to 0$ for every $t \in (0, T']$, and thus u = u' (since convergence in L^2 gives convergence almost everywhere on a subsequence). Hence $u \in C([0, T']; L^2)$ and (A.14) follows for $t \in [0, T']$ by taking $R \to \infty$ in the corresponding inequality for u_R . As for the energy dissipation equality (A.15), let $\delta \in (0, T')$ and note that property (ii) of the representation (see Section 2.1) applied to the difference $u_R - u$ gives

$$\nabla u_R \to \nabla u$$
 in $C([\delta, T']; L^2)$ as $R \to \infty$.

(Note that the convergence is not in $C([0, T']; L^2)$ since each ∇u_R need not belong to this space; see (ii).) Since Case 1 gives for each $R > 0, t \in [0, T']$

$$\frac{1}{2} \|u_R(t)\|^2 - \frac{1}{2} \|u_R(\delta)\|^2 + \int_{\delta}^t \|\nabla u_R(s)\|^2 \,\mathrm{d}s = \int_{\delta}^t \int u_R \cdot F_R \,\mathrm{d}x \,\mathrm{d}s,$$

(see (A.17)) we can take the limit $R \to \infty$ and then take the limit $\delta \to 0^+$ (and apply the Monotone Convergence Theorem again) to obtain (A.15) for all $t \in [0, T']$.

A.4.3 The continuity $\nabla u \in C((0,T);L^{\infty})$ for u given by formula (2.14)

We need to show that if $Y \in C((0,T), L^{\infty})$ is weakly divergence free, $||Y(t)||_{\infty}$ remains bounded as $t \to 0^+$ and $Y \in \mathcal{H}^{1/2}((0,T))$ and if u_2 is given by

$$u_2(t) \coloneqq \int_0^t \nabla \mathcal{T}(t-s) * [Y(s)Y(s)] \,\mathrm{d}s$$

(recall the notation (2.16)) then $\nabla u \in C((0,T); L^{\infty})$.

In order to prove it, fix T' < T and let M > 0 be such that

$$||Y(t)||_{\infty}, C_0(t) \le M$$
 for $t \in [0, T']$.

Fix $\varepsilon > 0$. Let $\eta > 0$ be such that

$$\int_0^t \frac{1}{(t-s)^{3/4}} \, \mathrm{d}s \le \frac{\varepsilon}{M^2} \quad \text{ for } t \le 2\eta.$$

Since for each pair i, k we have $Y_i Y_k \in C((0,T); L^{\infty}), Y_i Y_k$ is uniformly continuous into L^{∞} on time interval $[\eta/2, T']$. Thus there exists $\delta \in (0, \eta/2)$ such that for all i, k

$$||Y_i(t)Y_k(t) - Y_i(s)Y_k(s)||_{\infty} \le \frac{\varepsilon \eta}{T'}$$

whenever $s, t \in [\eta/2, T']$ and $|t - s| < \delta$. Now let $t_1, t_2 \in [0, T']$ be such that $t_1 < t_2$ and $t_2 - t_1 < \delta$, and calculate

$$\partial_l u_{2,j}(t_1) - \partial_l u_{2,j}(t_2) = \int_0^{t_2 - t_1} \partial_{li} \mathcal{T}_{jk}(t_2 - s) * [Y_i(s)Y_k(s)] \, \mathrm{d}s$$

+
$$\int_{t_2 - t_1}^{t_2} \partial_{li} \mathcal{T}_{jk}(t_2 - s) * [Y_i(s)Y_k(s)$$

-
$$Y_i(s - (t_2 - t_1))Y_k(s - (t_2 - t_1))] \, \mathrm{d}s$$

(see calculation (A.6)) and denote the two integrals on the right-hand side by I_1 , I_2 respectively. As for I_1 , using the same trick of employing the Hölder continuity of $Y_k(s)$ as indicated in (2.19), we obtain

$$\begin{aligned} |I_1| &\leq \int_0^{t_2 - t_1} \int |\partial_{li} \mathcal{T}_{jk}(x - y, t_2 - s) Y_i(y, s) \left[Y_k(y, s) - Y_k(x, s) \right] | \, \mathrm{d}y \, \mathrm{d}s \\ &\leq M^2 \int_0^{t_2 - t_1} \int \frac{C |x - y|^{1/2}}{\left(|x - y|^2 + (t_2 - s) \right)^{5/2}} \, \mathrm{d}y \, \mathrm{d}s \\ &= M^2 \int_0^{t_2 - t_1} \frac{C}{(t_2 - s)^{3/4}} \, \mathrm{d}s \leq C\varepsilon. \end{aligned}$$

If $t_2 \leq 2\eta$ then one can bound $|I_2|$ in a similar way to obtain

 $|I_2| \le 2C\varepsilon.$

Otherwise, we write $\int_{t_2-t_1}^{t_2} = \int_{t_2-t_1}^{\eta} + \int_{\eta}^{t_2-\eta} + \int_{t_2-\eta}^{t_2}$ and denote the resulting three integrals by $I_{2,1}$, $I_{2,2}$, $I_{2,3}$, respectively. Since the length of the intervals of integration in $I_{2,1}$, $I_{2,3}$ is less than 2η we obtain, as above,

$$\left|I_{2,1}\right|, \left|I_{2,3}\right| \le 2C\varepsilon.$$

For $I_{2,2}$ note that $s, s - (t_2 - t_1) \in [\eta/2, T']$ for each s from the interval of integration to write

$$\begin{aligned} |I_{2,2}| &\leq \int_{\eta}^{t_2 - \eta} \|\partial_{li} \mathcal{T}_{jk}(t_2 - s)\|_1 \|Y_i(s)Y_k(s) \\ &- Y_i(s - (t_2 - t_1))Y_k(s - (t_2 - t_1))\|_{\infty} \,\mathrm{d}s \\ &\leq \frac{\varepsilon \,\eta}{T'} \int_{\eta}^{t_2 - \eta} \frac{C}{t_2 - s} \,\mathrm{d}s \leq C\varepsilon, \end{aligned}$$

where we used the Minkowski inequality (1.3), Young's inequality (1.5) and the bound $\|\nabla^2 \mathcal{T}(t)\|_1 \leq C/t$ (see (1.13)). Thus altogether

$$|\partial_l u_{2,j}(t_1) - \partial_l u_{2,j}(t_2)| \le 5C\varepsilon \qquad l, j = 1, 2, 3$$

and the continuity $\nabla u_2 \in C((0,T); L^{\infty})$ follows.

A.4.4 Uniqueness of distributional solutions of the Stokes equations

We will show that if u, p are such that $u \in C([0,T); L^2)$ is weakly divergence free, $p \in L^1_{loc}(\mathbb{R}^3 \times [0,T))$, and

$$\int_0^T \int \left((\phi_t + \Delta \phi) \cdot u + p \operatorname{div} \phi \right) \, \mathrm{d}x \, \mathrm{d}t = 0 \tag{A.18}$$

for all $\phi \in C_0^{\infty}(\mathbb{R}^3 \times [0,T);\mathbb{R}^3)$, then $u \equiv 0$.

Proof. Fix $\psi \in C_0^{\infty}(\mathbb{R}^n \times [0,T))$ let $\phi \coloneqq \nabla \psi$. Then div $\phi = \Delta \psi$ and so (A.18) gives

$$\int_0^T \int p \,\Delta\psi \,\mathrm{d}x \,\mathrm{d}t = \int_0^T \int \nabla(\psi_t + \Delta\psi) \cdot u \,\mathrm{d}x \,\mathrm{d}t = 0 \tag{A.19}$$

since u is weakly divergence free.

For $\varepsilon > 0$ let

$$v(x,t) \coloneqq \int_0^t (J_{\varepsilon}u)(x,s) \,\mathrm{d}s, \qquad q(x,t) \coloneqq \int_0^t (J_{\varepsilon}p)(x,s) \,\mathrm{d}s.$$

We first show that v, q also satisfy (A.18),

$$\int_0^T \int \left((\phi_t + \Delta \phi) \cdot v + q \operatorname{div} \phi \right) \, \mathrm{d}x \, \mathrm{d}t = 0 \tag{A.20}$$

for all $\phi \in C_0^{\infty}(\mathbb{R}^3 \times [0,T);\mathbb{R}^3)$, or equivalently

$$0 = \int_0^T \int \int_0^t \int (\eta_{\varepsilon}(y)u(x-y,s) \cdot (\phi_t(x,t) + \Delta\phi(x,t)) + \eta_{\varepsilon}(y)p(x-y,s) \operatorname{div} \phi(x,t)) \, \mathrm{d}y \, \mathrm{d}s \, \mathrm{d}x \, \mathrm{d}t = \int \eta_{\varepsilon}(y) \int \int_0^T \left(\int_0^t u(x,s) \, \mathrm{d}s \cdot (\phi_t(x+y,t) + \Delta\phi(x+y,t)) + \int_0^t p(x,s) \, \mathrm{d}s \, \operatorname{div} \phi(x+y,t) \right) \, \mathrm{d}t \, \mathrm{d}x \, \mathrm{d}y.$$

We will show that the expression under the y integral vanishes. In fact, for fixed $y \in \mathbb{R}^3$ let

$$\Psi(x,t) \coloneqq -\int_t^{T'} \phi(x+y,s) \,\mathrm{d}s,$$

where T' < T is such that $\phi(t) \equiv 0$ for $t \geq T'$. Clearly $\Psi \in C_0^{\infty}(\mathbb{R}^3 \times [0,T);\mathbb{R}^3)$ and so (A.18) gives

$$0 = \int \int_0^T u(x,t) \cdot (\Psi_t(x,t) + \Delta \Psi(x,t)) \, \mathrm{d}t \, \mathrm{d}x + \int \int_0^T q(x,t) \, \mathrm{div} \, \Psi(x,t) \, \mathrm{d}t \, \mathrm{d}x.$$

Integration by parts in t and the identity

$$\partial_t(\Psi_t(x,t) + \Delta\Psi(x,t)) = \phi_t(x+y,t) + \Delta\phi(x+y,t)$$

give

$$\int \int_0^T \left(\int_0^t u(x,s) \, \mathrm{d}s \cdot (\phi_t(x+y,t) + \Delta \phi(x+y,t)) + \int_0^t p(x,s) \, \mathrm{d}s \, \operatorname{div} \phi(x+y,t) \right) \, \mathrm{d}t \, \mathrm{d}x = 0$$

as required. Therefore v, q indeed satisfy (A.20). Moreover, letting

$$\Psi(x,t) \coloneqq -\int_t^{T'} \psi(x+y,s) \,\mathrm{d}s,$$

this time for a scalar test function ψ we obtain from (A.19) that for each $y\in \mathbb{R}^3$

$$0 = \int_0^T \int p\Delta \Psi \, \mathrm{d}x \, \mathrm{d}t = -\int_0^T \int p(x,t) \int_t^{T'} \Delta \psi(x+y,s) \, \mathrm{d}s \, \mathrm{d}x \, \mathrm{d}t$$
$$= -\int_0^T \int \left(\int_0^t p(x,s) \, \mathrm{d}s\right) \Delta \psi(x+y,t) \, \mathrm{d}x \, \mathrm{d}t.$$

Hence Fubini's theorem gives

$$\int_0^T \int q\Delta\psi \, \mathrm{d}x \, \mathrm{d}t = \int_0^T \int \int_0^t \int \eta_\varepsilon(y) p(x-y,s) \Delta\psi(x,t) \, \mathrm{d}y \, \mathrm{d}s \, \mathrm{d}x \, \mathrm{d}t$$
$$= \int \eta_\varepsilon(y) \int_0^T \int \left(\int_0^t p(x,s) \, \mathrm{d}s\right) \Delta\psi(x+y,t) \, \mathrm{d}x \, \mathrm{d}t \, \mathrm{d}y = 0,$$

that is, like p, q satisfies (A.19). Therefore (A.20) applied with $\Delta \phi$ in place of ϕ gives

$$0 = \int_0^T \int \Delta(\phi_t + \Delta\phi) \cdot v \, \mathrm{d}x \, \mathrm{d}t = \int_0^T \int (\phi_t + \Delta\phi) \cdot \Delta v \, \mathrm{d}x \, \mathrm{d}t.$$

The uniqueness of weak solutions to the heat equation (see Lemma A.2) now implies $\Delta v \equiv 0$ almost everywhere in $\mathbb{R}^3 \times [0, T)$, hence $\Delta v \equiv 0$ everywhere by continuity of Δv . Hence Liouville's theorem implies that v(t) is constant for each t. Therefore $v(t) \equiv 0$ at each t due to the fact $v \in C([0,T); L^2)$. Hence, differentiating the definition of v (in t), we see that $J_{\varepsilon}u(t) \equiv 0$ for each t and ε . The almost everywhere convergence of the mollification (see Lemma 1.2, (v)) gives $u(t) \equiv 0$ for each t, as required.

A.5 Integral inequalities

Lemma A.5. Suppose g > 0 is a continuous function on (0,T) that is locally integrable [0,T), that functions $f, \phi : (0,T) \to \mathbb{R}^+$ satisfy

$$f(t) \leq \int_{0}^{t} g(t-s)f(s)^2 \,\mathrm{d}s + a(t),$$
 (A.21)

$$\phi(t) \geq \int_0^t g(t-s)\phi(s)^2 \,\mathrm{d}s + b(t) \tag{A.22}$$

for all $t \in (0,T)$, where a, b are continuous functions satisfying $a \leq b$, ϕ is continuous, and that f^2 and ϕ^2 are integrable near 0. Then $f \leq \phi$ on (0,T).

Proof. The proof proceeds in three steps.

Step 1. The case $a(t) \leq b(t) - \delta$ for $t \in (0, \tau)$ for some $\delta, \tau > 0$. Let

$$I := \left\{ t' : \int_0^t g(t-s)f(s)^2 \, \mathrm{d}s + a(t) < \phi(t) \text{ for all } t \in (0,t'] \right\}.$$

Note that (A.21) gives $f(t) < \phi(t)$ for $t \in I$. Let $t_0 \in (0, \tau)$ be such that

$$\left|\int_0^t g(t-s)f(s)^2 \,\mathrm{d}s\right| < \frac{\delta}{2}, \quad \left|\int_0^t g(t-s)\phi(s)^2 \,\mathrm{d}s\right| < \frac{\delta}{2}$$

for $t \in (0, t_0]$. Then for $t \in (0, t_0]$ (A.22) gives

$$\int_{0}^{t} g(t-s)f(s)^{2} ds + a(t) < \delta/2 + b(t) - \delta$$

$$\leq \phi(t) - \int_{0}^{t} g(t-s)\phi(s)^{2} ds - \delta/2 < \phi(t),$$

that is $t_0 \in I$. Now let $T' := \sup I$. We need to show that T' = T. Suppose otherwise that T' < T. Then

$$\int_0^{T'} g(T'-s)f(s)^2 \,\mathrm{d}s + a(T') < \int_0^{T'} g(T'-s)\phi(s)^2 \,\mathrm{d}s + b(T') \le \phi(T')$$

by (A.22). By continuity we obtain $\int_0^t g(t-s)f(s)^2 ds + a(t) < \phi(t)$ for $t \in [T', T'']$ for some T'' > T'. Hence $T'' \in I$, which contradicts the definition of T'. Therefore indeed T' = T and the lemma follows in this case.

Step 2. The case $\liminf_{t\to 0^+} (b(t) - a(t)) = 0$: there exists $t_0 > 0$ such that $f(t) \leq \phi(t)$ for $t \in (0, t_0)$.

Let $t_0 > 0$ be small enough such that

$$\left| \int_{0}^{t} g(t-s) \, \mathrm{d}s \right| < \frac{1}{4}, \quad \left| \int_{0}^{t} g(t-s) f(s) \, \mathrm{d}s \right| < \frac{1}{4} \qquad \text{for } t \in [0, t_0].$$

Let $\varepsilon \in (0, 1/3)$ and consider $f_{\varepsilon}(t) := f(t) - \varepsilon$. Then $\varepsilon/2 + 3\varepsilon^2/4 - \varepsilon \le -\varepsilon/4$ and so f_{ε} satisfies the inequality

$$\begin{split} f_{\varepsilon}(t) &\leq \int_{0}^{t} g(t-s)(f_{\varepsilon}(s)+\varepsilon)^{2} \,\mathrm{d}s + a(t) - \varepsilon \\ &\leq \int_{0}^{t} g(t-s)f_{\varepsilon}(s)^{2} \,\mathrm{d}s + 2\varepsilon \int_{0}^{t} g(t-s)(f(s)-\varepsilon) \,\mathrm{d}s \\ &+ \varepsilon^{2} \int_{0}^{t} g(t-s) \,\mathrm{d}s + a(t) - \varepsilon \\ &\leq \int_{0}^{t} g(t-s)f_{\varepsilon}(s)^{2} \,\mathrm{d}s + \varepsilon/2 + \varepsilon^{2}/2 + \varepsilon^{2}/4 + a(t) - \varepsilon \\ &\leq \int_{0}^{t} g(t-s)f_{\varepsilon}(s)^{2} \,\mathrm{d}s + \varepsilon/2 + \varepsilon^{2}/4 + a(t) - \varepsilon \end{split}$$

Because $a(t) - \varepsilon/4 \le b(t) - \varepsilon/4$ on $(0, t_0)$, similarly as in Step 1 we obtain $f_{\varepsilon} \le \phi$ on $(0, t_0)$. The claim follows by taking the limit $\varepsilon \to 0^+$.

Step 3. The case $\liminf_{t\to 0^+} (b(t) - a(t)) = 0$: $f(t) \le \phi(t)$ for all $t \in (0, T)$. Let

$$I_1 := \{ t \in (0,T) : f(s) \le \phi(s) \text{ for } s \in (0,t) \}.$$

Let $t_1 := \sup I_1$. Note that Step 2 gives $t_1 \ge t_0 > 0$. Suppose that $t_1 < T$. Let $F(t) := f(t_1 + t), \ \Phi(t) := \phi(t_1 + t)$. Then $F, \ \Phi$ satisfy

$$F(t) \leq \int_0^t g(t-s)F(s)^2 \,\mathrm{d}s + A(t),$$

$$\Phi(t) \geq \int_0^t g(t-s)\Phi(s)^2 \,\mathrm{d}s + B(t),$$

where

$$A(t) \coloneqq a(t_1 + t) + \int_0^{t_1} g(t_1 + t - s) f(s)^2 \, \mathrm{d}s,$$

$$B(t) \coloneqq b(t_1 + t) + \int_0^{t_1} g(t_1 + t - s)\phi(s)^2 \,\mathrm{d}s$$

$$\geq A(t) + b(t_1 + t) - a(t_1 + t).$$

Noting that A, B are continuous (by the Dominated Convergence Theorem) and that $A(t) \leq B(t)$ for all $t \in (0, T - t_1)$, we can apply Step 1 and Step 2 to the functions F, Φ , to conclude that $F(t) \leq \Phi(t)$ for all $t \in (0, t_2]$ for some $t_2 > 0$. Thus $f(t) \leq \phi(t)$ for all $t \in [0, t_1 + t_2)$, which contradicts the definition of t_1 .

The above lemma can be modified to fit several other settings.

Corollary A.6. Let g, a, b be as in Lemma 1.5 and let h satisfy the same conditions as g. Let functions $f, \phi: (0,T) \to \mathbb{R}$ be such that f and ϕ are bounded near as $t \to 0^+$ and ϕ is continuous and

$$\begin{cases} f(t) &\leq \int_0^t g(t-s)h(s)f(s)\,\mathrm{d}s + a(t), \\ \phi(t) &\geq \int_0^t g(t-s)h(s)\phi(s)\,\mathrm{d}s + b(t) \end{cases}$$

for all $t \in (0,T)$. Then $f \leq \phi$ on (0,T).

Proof. Similar to the proof above.

Corollary A.7. Let a, b > 0 be such that $a \leq b$ and let $h, g: (0, \infty) \to \mathbb{R}^+$ be continuous functions such that g is locally integrable on $[0, \infty)$ and h is integrable on $(1, \infty)$. Suppose also that there exist $\tau > 0$, C' > 0 such that

$$\begin{split} h(s) &\geq C'^2 g(s) \quad for \ s \in (0,\tau), \\ h(s) &\leq C'^2 g(s) \quad for \ s \in [\tau,\infty), \end{split}$$

and for all t > 0

$$C' > \int_0^t \min\left(C'^2 g(t-s), h(t-s)\right) \, \mathrm{d}s + b.$$

If T > 0 and f is a positive function on (0,T) that is bounded near 0 and

$$f(t) \le \int_0^t \min\left(g(t-s)f(s)^2, h(t-s)\right) \,\mathrm{d}s + a$$

for $t \in (0,T)$, then $f \leq C'$ on (0,T).

Proof. If $t < \tau$ then the minimum under the second last integral is $(C')^2 g(t-s)$ and so Lemma A.5 gives $f(t) \leq C'$ for such t's. Thus letting

$$t_0 \coloneqq \sup\{t' > 0 \colon f(t) \le C' \text{ for } t < t'\}$$

we see that $t_0 \ge \tau > 0$. If $t_0 < T$ we obtain

$$\int_0^{t_0} \min\left(g(t_0 - s)f(s)^2, h(t_0 - s)\right) \, \mathrm{d}s + a$$

$$\leq \int_0^{t_0} \min\left(g(t_0 - s)(C')^2, h(t_0 - s)\right) \, \mathrm{d}s + b < C'.$$

Thus, by continuity

$$\int_0^t \min(g(t-s)f(s)^2, h(t-s)) \, \mathrm{d}s + a < C'$$

for $t \in [t_0, t_0 + \delta)$ for some $\delta > 0$, which contradicts the definition of t_0 . Thus $t_0 = T$, as required.
A.6 The Volterra equation

In this section we show that the equation

$$\phi(x) = C \int_0^x \frac{\phi(y)}{\sqrt{x-y}} \,\mathrm{d}y + D, \tag{A.23}$$

where C, D > 0, has a unique solution $\phi \in C[0, \infty)$. This is equivalent to showing that (A.23) has a unique solution $\phi \in C[0, T]$ for every T > 0. We can rewrite (A.23) in the form

$$\phi - A\phi = D, \tag{A.24}$$

where

$$A\phi(x) := \int_0^T K(x, y)\phi(y) \,\mathrm{d}y \tag{A.25}$$

with $K(x,y) := C \chi_{\{y < x\}} (x - y)^{-1/2}$. This is an example of the Volterra integral equation of the 2nd kind with with a weakly singular kernel K, that is any K such that for all $x, y \in [0,T]$ with $x \neq y$ K is continuous at (x,y)and $|K(x,y)| \leq M|x - y|^{\alpha-1}$ for some $\alpha \in (0,1]$, M > 0. In what follows we apply the theory of such equations to (A.24). We consider only the case $\alpha = 1/2$; other cases follow similarly. We follow the arguments from Kress (2014).

First note that the set of compact operators is closed in the operator norm.

Lemma A.8. Let X be a Banach space and $A_n \in L(X)$ be a sequence of compact operators such that $||A_n - A|| \to 0$ as $n \to \infty$ for some $A \in L(X)$. Then A is compact.

This is elementary (see e.g. Kress (2014), p. 26, for the proof). We now show that A (defined by (A.25)) is a compact operator on X := C[0,T] by cutting off the singularity and using the above lemma.

Lemma A.9. The operator $A: X \to X$ is compact.

Proof. We see that A is continuous by writing

$$|A\phi(t)| = C \left| \int_0^t \frac{\phi(s)}{\sqrt{t-s}} \, \mathrm{d}s \right|$$

$$\leq C \|\phi\|_{\sup} \int_0^t (t-s)^{-1/2} \, \mathrm{d}s = 2C \|\phi\|_{\sup} t^{1/2} \leq 2CT^{1/2} \|\phi\|_{\sup}$$

For $n \in \mathbb{N}$ we define a cut-off K_n of the kernel K by

$$K_n(t,s) := \begin{cases} h(n|t-s|)K(t,s) & t \neq s \\ 0 & t = s, \end{cases}$$

where $h: [0, \infty) \to [0, 1]$ is a continuous function such that h(t) = 0 for $t \in [0, 1/2]$ and h(t) = 1 for $t \ge 1$. Because $K_n \in C([0, T]^2)$ for every n, the corresponding integral operators A_n are compact by the Arzelà-Ascoli theorem. Moreover

$$|A\phi(t) - A_n\phi(t)| = \left| \int_{t-1/n}^t (1 - h(n|t-s|))K(t,s)\phi(s) \,\mathrm{d}s \right|$$

$$\leq \|\phi\|_{\sup} \int_{t-1/n}^t \frac{C}{\sqrt{t-s}} \,\mathrm{d}s = C \|\phi\|_{\sup} \ n^{-1/2} \to 0$$

as $n \to \infty$ uniformly in t. Hence $||A_n - A|| \to 0$ and Lemma A.8 gives compactness of A.

We now show the claim by applying Fredholm Alternative.

Theorem A.10. The equation (A.24) has a unique solution $\phi \in C[0,T]$.

Proof. Because X is a Banach space and $A : X \to X$ is compact we can apply Fredholm Alternative to conclude that (A.24) has a unique solution if the equation

$$\phi - A\phi = 0 \tag{A.26}$$

has no non-zero solution. We will use induction to show that a solution ϕ to this homogeneous problem satisfies

$$|\phi(t)| \le \|\phi\|_{\sup} \frac{M^k t^k}{k!} \tag{A.27}$$

for some M > 0 and all $t \in [0, T]$, $k = 0, 1, \ldots$ The base case k = 0 is trivial. For the inductive step we first note that that for any $t, s \in [0, T]$ with $0 \le s < t \le T$ we have

$$\int_{s}^{t} \frac{\mathrm{d}\tau}{\sqrt{t-\tau}\sqrt{\tau-s}} = \int_{0}^{1} \frac{\mathrm{d}z}{\sqrt{z(1-z)}} =: I$$

by the change of variable $z := \frac{\tau - s}{t - s}$. Now assume that a solution ϕ to (A.26) satisfies $|\phi(t)| \leq M \|\phi\|_{\sup} \frac{C^k t^k}{k!}$ for some k. Then, because $\phi = A\phi = A^2\phi$, we have for all $t \in (0, T]$

$$\begin{aligned} |\phi(t)| &= |A^2 \phi(t)| = C^2 \left| \int_0^t \int_0^\tau \frac{\phi(s)}{\sqrt{\tau - s}\sqrt{t - \tau}} \, \mathrm{d}s \, \mathrm{d}\tau \right| \\ &\leq C^2 \int_0^t \int_0^\tau \frac{|\phi(s)|}{\sqrt{\tau - s}\sqrt{t - \tau}} \, \mathrm{d}s \, \mathrm{d}\tau = C^2 \int_0^t \int_s^t \frac{|\phi(s)|}{\sqrt{\tau - s}\sqrt{t - \tau}} \, \mathrm{d}\tau \, \mathrm{d}s \\ &= IC^2 \int_0^t |\phi(s)| \, \mathrm{d}s \leq \|\phi\|_{\sup} \frac{IC^2 M^k}{k!} \int_0^t s^k \, \mathrm{d}s = \|\phi\|_{\sup} \frac{IC^2 M^k t^{k+1}}{(k+1)!} \, . \end{aligned}$$

where we used Fubini's theorem. The bound (A.27) now follows with $M := IC^2$. Taking the limit $k \to \infty$ in (A.27) gives $\phi \equiv 0$.

A.7 A proof of (4.8) without the use of the Plancherel Lemma

Here we give an elementary proof of (4.8),

$$||p||^2 \le C ||\nabla u||^3 ||u||.$$

First note that the representation formula for p (see (4.3)) can be rewritten in the form

$$p = \frac{1}{4\pi} \int \nabla \left(\frac{1}{|x-y|} \right) \cdot g(y) \, \mathrm{d}y,$$

where $g \coloneqq (J_{\varepsilon} u \cdot \nabla) u$. Therefore

$$\begin{split} \|p\|^2 &= \frac{1}{(4\pi)^2} \iiint \left[\nabla \frac{1}{|x-y|} \cdot g(y) \right] \left[\nabla \frac{1}{|x-z|} \cdot g(z) \right] \, \mathrm{d}x \, \mathrm{d}y \, \mathrm{d}z \\ &= \frac{-1}{(4\pi)^2} \iiint \left[\frac{1}{|x-y|} \mathrm{div} \, g(y) \right] \left[\nabla \frac{1}{|x-z|} \cdot g(z) \right] \, \mathrm{d}x \, \mathrm{d}y \, \mathrm{d}z \end{split}$$

Since $\nabla_x |x - y|^{-1} = \nabla_y |x - y|^{-1}$, integration by parts in x and then in y gives

$$\|p\|^2 = \frac{-1}{(4\pi)^2} \iiint \left[\frac{1}{|x-y|} \nabla(\operatorname{div} g(y)) \right] \cdot \frac{g(z)}{|x-z|} \, \mathrm{d}x \, \mathrm{d}y \, \mathrm{d}z.$$

Now the calculus identity

$$\nabla(\operatorname{div} g) = \Delta g + \operatorname{curl}(\operatorname{curl} g)$$

gives

$$\|p\|^{2} = \frac{1}{4\pi} \iint \frac{g(x) \cdot g(z)}{|x - z|} \, \mathrm{d}x \, \mathrm{d}z - \frac{1}{(4\pi)^{2}} \iiint \left[\frac{1}{|x - y|} \operatorname{curl}(\operatorname{curl} g(y)) \right] \cdot \frac{g(z)}{|x - z|} \, \mathrm{d}x \, \mathrm{d}y \, \mathrm{d}z.$$
(A.28)

Since the *i*th component of curl g can be expressed in the form $(\operatorname{curl} g)_i = \epsilon_{ijk} \partial_j g_k$, where the coefficients

$$\epsilon_{ijk} \coloneqq \begin{cases} 1 & \text{if } ijk \text{ is an even permutation of } 123, \\ -1 & \text{if } ijk \text{ is an odd permutation of } 123, \\ 0 & \text{otherwise} \end{cases}$$

satisfy $\epsilon_{ijk} = -\epsilon_{kji}$, we can write the last triple integral as

$$\iiint \frac{\epsilon_{ijk}\partial_j \left(\operatorname{curl} g(y)\right)_k g_i(z)}{|x-y| |x-z|} \,\mathrm{d}x \,\mathrm{d}y \,\mathrm{d}z,$$

which integrated by parts three times (first in y then in x and z) gives

$$-\iiint \frac{\epsilon_{ijk} \left(\operatorname{curl} g(y)\right)_k \partial_j g_i(z)}{|x-y| |x-z|} \, \mathrm{d}x \, \mathrm{d}y \, \mathrm{d}z$$

$$= \iiint \frac{\left(\operatorname{curl} g(y)\right)_k \epsilon_{kji} \partial_j g_i(z)}{|x-y| |x-z|} \, \mathrm{d}x \, \mathrm{d}y \, \mathrm{d}z$$

$$= \iiint \frac{\left(\operatorname{curl} g(y)\right) \cdot \left(\operatorname{curl} g(z)\right)}{|x-y| |x-z|} \, \mathrm{d}x \, \mathrm{d}y \, \mathrm{d}z = \int |F(x)|^2 \, \mathrm{d}x \ge 0,$$

where $F(x) \coloneqq \int \operatorname{curl} g(y) / |x - y| \, dy$. Therefore (A.28) gives

$$||p||^{2} \leq \frac{1}{4\pi} \iint \frac{g(x) \cdot g(z)}{|x-z|} \, \mathrm{d}x \, \mathrm{d}z$$
$$= \frac{1}{4\pi} \iint \frac{(J_{\varepsilon} u_{k}(x))\partial_{k} u_{i}(x)(J_{\varepsilon} u_{j}(z))\partial_{j} u_{i}(z)}{|x-z|} \, \mathrm{d}x \, \mathrm{d}z$$

(this inequality appears in Leray (1934b) on p. 233 with " \leq " wrongly replaced by "="), from where the Cauchy-Schwarz inequality and Lemma 1.3 give

$$\begin{aligned} \|p\|^2 &\leq \frac{1}{4\pi} \int \|\nabla u\| \left(\int \frac{|J_{\varepsilon}u_k(x)|^2}{|x-z|^2} \,\mathrm{d}x \right)^{1/2} \,|J_{\varepsilon}u_j(z)| \,|\nabla u(z)| \,\mathrm{d}z \\ &\leq C \|\nabla u\|^2 \int |J_{\varepsilon}u_j(z)| \,|\nabla u(z)| \,\mathrm{d}z \leq C \|\nabla u\|^3 \|u\|, \end{aligned}$$

where we also used $||J_{\varepsilon}(\nabla u)|| \le ||\nabla u||, ||J_{\varepsilon}u|| \le ||u||$ (see Lemma 1.2 (i)).

A.8 Smooth approximation of the forcing

Lemma A.11 (Dini's lemma). Let I be a compact interval and let $f_n, f \in C(I; \mathbb{R})$ be continuous functions such that $f_n(t) \to f(t)$ as $n \to \infty$ and $f_{n+1}(t) \leq f_n(t)$ for each $t \in I$. Then $||f_n - f||_{C(I)} \to 0$.

Proof. This is elementary.

Lemma A.12. Let $p \in [1, \infty)$ and $F \in C([0, T), L^p)$. For any $T' \in (0, T)$ and any $\varepsilon > 0$ there exists $\widetilde{F} \in C^{\infty}(\mathbb{R}^3 \times \mathbb{R})$ and R > 0 such that supp $\widetilde{F}(t) \subset B(0, R)$ for all $t \in \mathbb{R}$ and

$$\|F - \widetilde{F}\|_{C([0,T'],L^2)} < \varepsilon.$$
(A.29)

Moreover $\max_{t \in [0,T']} \|\widetilde{F}(t)\|_{\infty} \le \max_{t \in [0,T']} \|F(t)\|_{\infty}.$

Proof. It suffices to consider $T < \infty$. First extend F in time from [0, T'] to the whole line by taking $F(\cdot, t) \coloneqq F(\cdot, 0)$ for t < 0 and $F(\cdot, t) \coloneqq F(\cdot, T')$ for t > T'. For R > 0 let

$$F_R(x,t) \coloneqq \chi_{B(0,R)}(x)F(x,t).$$

Clearly $F_R \in C(\mathbb{R}; L^p)$ (as a product of two such functions) and so $||F_R(t) - F(t)||_p$ is continuous in t for each R. Hence, noting that $||F_R(t) - F(t)||_p$ is a nonincreasing function of R converging to zero as $R \to \infty$ for each $t \in [0, T']$, we can use Dini's Lemma (Lemma A.11) to fix R > 0 such that

$$||F_R - F||_{C([0,T'];L^p)} < \varepsilon/3.$$
 (A.30)

We will now mollify F_R to obtain \widetilde{F} . Let η_{δ} , ξ_{δ} be mollifiers in \mathbb{R}^3 and \mathbb{R} respectively, that is let $\xi(t) \coloneqq C \exp(1/(|t|^2 - 1))$ for $t \in (0, 1)$ and $\xi(t) \coloneqq 0$ if $t \notin (0, 1)$, where C is such that $\int_{\mathbb{R}} \xi = 1$, and let $\xi_{\delta}(t) \coloneqq \xi(t/\delta)/\delta$, $\eta(x) \coloneqq \xi(|x|), \eta_{\delta}(x) \coloneqq \eta(x/\delta)/\delta^3$. Define the mollification F_R^{δ} of F_R by

$$\begin{split} F_R^{\delta}(x,t) &\coloneqq \int_{\mathbb{R}} \xi_{\delta}(s) \int \eta_{\delta}(y) F_R(x-y,t-s) \, \mathrm{d}y \, \mathrm{d}s \\ &= \int_{\mathbb{R}} \xi(s) \int \eta(y) F_R(x-\delta y,t-\delta s) \, \mathrm{d}y \, \mathrm{d}s. \end{split}$$

Clearly $\widetilde{F} := F_R^{\delta}(x,t)$ has the required regularity for each $\delta > 0$ (in particular $\max_{t \in [0,T']} \|\widetilde{F}(t)\|_{\infty} \leq \max_{t \in [0,T']} \|F(t)\|_{\infty}$ holds by the property of mollifiers (see (i) in Lemma 1.2)). Therefore the proof will be complete if we show the approximation property (A.29) for some $\delta > 0$. Because $F_R(x,t) - F_R^{\delta}(x,t) = \int_{\mathbb{R}} \xi(s) \int \eta(y) \left(F_R(x,t) - F_R(x - \delta y, t - \delta s)\right) \, dy \, ds$, we can use the Minkowski inequality (see 1.4) and the triangle inequality

$$||F_R(x - \delta y, t - \delta s) - F_R(x, t)||_p \le ||F_R(x - \delta y, t - \delta s) - F_R(x - \delta y, t)||_p + ||F_R(x - \delta y, t) - F_R(x, t)||_p$$

to write

$$\begin{aligned} \|F_R(t) - F_R^{\delta}(t)\|_p &\leq \int_{\mathbb{R}} \xi(s) \int \eta(y) \|F_R(\cdot - \delta y, t - \delta s) - F_R(\cdot, t)\|_p \, \mathrm{d}y \, \mathrm{d}s \\ &\leq \int_{\mathbb{R}} \xi(s) \|F_R(t - \delta s) - F_R(t)\|_p \, \mathrm{d}s \\ &+ \int \eta(y) \|F_R(\cdot - \delta y, t) - F_R(\cdot, t)\|_p \, \mathrm{d}y. \end{aligned}$$
(A.31)

Since F_R is uniformly continuous on [-1, T' + 1] into L^p (recall we extended F in time to the whole line) there exists sufficiently small $\delta_1 > 0$ such that for $\delta \in (0, \delta_1)$

$$||F_R(t-\delta s) - F_R(t)||_p < \varepsilon/3$$

for all $t \in [0, T']$, and so the first term on the right-hand side of (A.31) is less than $\varepsilon/3$ for all $t \in [0, T']$. As for the second term there exists $\delta > 0$ such that $\delta < \delta_1$ and

$$||F_R(\cdot - z, t) - F_R(\cdot, t)||_p < \varepsilon/3$$

whenever $|z| < \delta$ and $t \in [0, T']$. Indeed, by the continuity of translation in space of L^p functions for each $t_0 \in [0, T']$ there exists a δ_{t_0} such that $||F_R(\cdot - z, t_0) - F_R(\cdot, t_0)||_p < \varepsilon/3$ whenever $|z| < \delta_{t_0}$. Moreover, the continuity of F_R in time into L^p and triangle inequality gives that $||F_R(\cdot - z, t) - X_R(\cdot, t)||_p < \varepsilon/3$ whenever $|z| < \delta_{t_0}$ and t belongs to some open set J_{t_0} containing t_0 . By compactness of [0, T'] we obtain a finite cover $\{J_{t_i}\}_{i=1,...,m}$ of [0, T']consisting of such open sets and δ is obtained by taking the minimum of the corresponding δ_{t_i} , i = 1, ..., m. This means in particular that

$$||F_R(\cdot - \delta y, t) - F_R(\cdot, t)||_p < \varepsilon/3$$

whenever |y| < 1, that is for all $y \in \text{supp }\eta$. Therefore, for such δ the second term on the right-hand side of (A.31) is bounded by $\varepsilon/3$ uniformly in $t \in [0, T']$. The approximation property (A.29) therefore follows directly from (A.30) and (A.31).

References

- Beale, J. T., Kato, T. & Majda, A. (1984), 'Remarks on the breakdown of smooth solutions for the 3-D Euler equation', *Comm. Math. Phys.* 94, 61– 66.
- Besicovitch, A. S. (1935), 'On the sum of digits of real numbers represented in the dyadic system', *Math. Ann.* **110**(1), 321–330.
- Besicovitch, A. S. & Taylor, S. J. (1954), 'On the complementary intervals of a linear closed set of zero Lebesgue measure', J. London Math. Soc. 29, 449–459.
- Bouligand, G. (1928), 'Ensembles impropres et nombre dimensionnel', Bull. Sciences Mathématiques 52, 320–334, 361–376.
- Caffarelli, L., Kohn, R. & Nirenberg, L. (1982), 'Partial regularity of suitable weak solutions of the Navier-Stokes equations', *Comm. Pure Appl. Math.* 35(6), 771–831.
- Carothers, N. L. (2000), *Real analysis*, Cambridge University Press, Cambridge.
- Constantin, P. (2008), 'Euler and Navier-Stokes Equations', *Publ. Mat* 52(2), 235–265.
- Constantin, P. & Foias, C. (1988), Navier-Stokes Equations, The University of Chicago Press, Chicago.
- Escauriaza, L., Seregin, G. & Šverák, V. (2003), $L_{3,\infty}$ -solutions of Navier–Stokes equations and backwards uniqueness', *Russian Math. Surveys* **58**, 211–250.

- Evans, L. (2010), *Partial differential equations: second edition*, American Mathematical Society, Providence R.I.
- Evans, L. C. & Gariepy, R. F. (2015), *Measure theory and fine properties* of functions, Textbooks in Mathematics, revised edn, CRC Press, Boca Raton, FL.
- Falconer, K. (2014), Fractal geometry, Mathematical foundations and applications, Third Edition, John Wiley & Sons, Ltd., Chichester.
- Fefferman, C. (2006), Existence and Smoothness of the Navier-Stokes Equation, in J. Carlson, A. Jaffe & A. Wiles, eds, 'The Millenium Prize Problems', AMS.
- Fujita, H. & Kato, T. (1964), 'On the Navier–Stokes initial value problem. I', Arch. Ration. Mech. Anal. 16, 269–315.
- Giga, M.-H., Giga, Y. & Saal, J. (2010), Nonlinear partial differential equations, Asymptotic behavior of solutions and self-similar solutions, Progress in Nonlinear Differential Equations and their Applications, 79, Birkhäuser Boston, Inc., Boston, MA.
- Hausdorff, F. (1918), 'Dimension und äußeres Maß', Math. Ann. **79**(1-2), 157–179.
- Helly, E. (1912), 'Uber lineare Funktionaloperationen', Sitzungsberichte der Kaiserlichen Akademie der Wissenschaften zu Wien. Mathematisch-Naturwissenschaftlichen Klasse, 121 II A1, 265–297.
- Hopf, E. (1951), 'Über die Anfangswertaufgabe für die hydrodynamischen Grundgleichungen', Math. Nachr. 4, 213–231. (An English translation due to Andreas Klöckner is available at http://www.dam.brown.edu/people/menon/publications/notes/hopf-NS.pdf.).
- Kato, T. (1984), 'Strong l^p -solutions of the Navier–Stokes equations in \mathbb{R}^m with applications to weak solutions', *Math. Zeit.* **187**, 471–480.
- Kiselev, A. & Ladyzhenskaya, O. (1957), 'On the existence and uniqueness of the solution of the nonstationary problem for a viscous, incompressible fluid.', *Izv. Akad. Nauk SSSR. Ser. Mat.* **21**, 655–680. in Russian.
- Kress, R. (2014), Linear integral equations, Vol. 82 of Applied Mathematical Sciences, third edn, Springer, New York.
- Ladyženskaja, O. A. (1967), 'Uniqueness and smoothness of generalized solutions of Navier-Stokes equations', Zap. Naučn. Sem. Leningrad. Otdel. Mat. Inst. Steklov. (LOMI) 5, 169–185.

- Ladyzhenskaya, O. A. (1959), 'Solution "in the large" of the nonstationary boundary value problem for the Navier–Stokes system in two space variables', Comm. Pure Appl. Math. 12, 427–433.
- Lapidus, M. L. & van Frankenhuijsen, M. (2006), Fractal geometry, complex dimensions and zeta functions, Springer Monographs in Mathematics, Springer, New York. Geometry and spectra of fractal strings.
- Lax, P. (1998), Jean Leray and Partial Differential Equations, Springer-Verlag, Berlin; Société Mathématique de France, Paris. Introduction to Jean Leray, Selected papers. Œuvres scientifiques. Vol. II: Fluid dynamics and real partial differential equations/Équations aux dérivées partielles réelles et mécanique des fluides, Springer.
- Lemarié-Rieusset, P. G. (2002), Recent developments in the Navier-Stokes problem, Vol. 431 of Chapman & Hall/CRC Research Notes in Mathematics, Chapman & Hall/CRC, Boca Raton, FL.
- Lemarié-Rieusset, P. G. (2016), *The Navier-Stokes problem in the 21st century*, CRC Press, Boca Raton, FL.
- Leray, J. (1933), 'Étude de diverses équations intégrales non linéaires et de quelques problèmes que pose l'hydrodynamique', J. Math. Pures Appl. (9) 12, 1–82. (thesis).
- Leray, J. (1934a), 'Essai sur les mouvements plans d'un liquide visquex que limitent des parois', J. Math. Pures Appl. (9) 13, 331–418. (Available in Jean Leray, Selected papers. Œuvres scientifiques. Vol. II: Fluid dynamics and real partial differential equations/Équations aux dérivées partielles réelles et mécanique des fluides, Springer, 1998.).
- Leray, J. (1934b), 'Sur le mouvement d'un liquide visqueux emplissant l'espace', Acta Math. 63, 193–248. (An English translation due to Robert Terrell is available at http://www.math.cornell.edu/~bterrell/leray.pdf and https://arxiv.org/abs/1604.02484.).
- Majda, A. & Bertozzi, A. (2002), Vorticity and incompressible flow, CUP, Cambridge.
- Navier, C. L. M. H. (1822), 'Mémoire sur les lois du mouvement des fluides', Mém. Ac. R. Sc. de l'Institut de France 6, 389–440.
- Nečas, J., Růžička, M. & Šverák, V. (1996), 'On Leray's self-similar solutions of the Navier-Stokes equations', Acta Math. 176(2), 283–294.
- Oseen, C. (1911), 'Sur les formules de green généralisées qui se présentent dans l'hydrodynamique et sur quelquesunes de leurs applications', Acta. Math. 34, 205–288.

- Pontrjagin, L. & Schnirelmann, L. (1932), 'Sur une propriété métrique de la dimension', Ann. of Math. (2) 33(1), 156–162.
- Prodi, G. (1959), 'Un teorema di unicità per le equazioni di Navier-Stokes', Ann. Mat. Pura Appl. (4) 48, 173–182.
- Robinson, J. C. (2006), 'Regularity and singularity in the three-dimensional Navier-Stokes equations', Bol. Soc. Esp. Mat. Apl. SeMA 35, 43–71.
- Robinson, J., Rodrigo, J. & Sadowski, W. (2016), The three-dimensional Navier-Stokes equations. Classical Theory, Cambridge University Press, Cambridge.
- Scheffer, V. (1976), 'Turbulence and Hausdorff dimension'. In *Turbulence and Navier-Stokes equations* (Proc. Conf., Univ. Paris-Sud, Orsay, 1975), Springer LNM 565: 174–183, Springer Verlag, Berlin.
- Scheffer, V. (1977), 'Hausdorff measure and the Navier–Stokes equations', Comm. Math. Phys. 55, 97–112.
- Schwartz, L. (1950), Théorie des distributions. Tome I, Actualités Sci. Ind., no. 1091, Publ. Inst. Math. Univ. Strasbourg 9, Hermann & Cie., Paris.
- Serrin, J. (1962), 'On the interior regularity of weak solutions of the Navier-Stokes equations', Arch. Rational Mech. Anal. 9, 187–195.
- Sobolev, S. (1936), 'Méthode nouvelle à résoudre le problème de cauchy pour les équations linéaires hyperboliques normales', *Rec. Math. [Mat. Sbornik] N.S.* 1(43)(1), 39–72.
- Sohr, H. (2001), *The Navier-Stokes equations, An elementary functional analytic approach*, Birkhäuser Advanced Texts: Basel Textbooks, Birkhäuser Verlag, Basel.
- Stein, E. M. & Weiss, G. (1971), Introduction to Fourier analysis on Euclidean spaces, Princeton University Press, Princeton, N.J. Princeton Mathematical Series, No. 32.
- Stokes, G. (1845), 'On the theories of the internal friction of fluids in motion, and of the equilibrium and motion of elastic solids', *Trans. Cam. Phil. Soc.* 8, 287–319.
- Temam, R. (2001), Navier-Stokes equations, AMS Chelsea Publishing, Providence, RI. Theory and numerical analysis, Reprint of the 1984 edition.