

# Seifert surgery on knots via Reidemeister torsion and Casson-Walker-Lescop invariant III

Teruhisa Kadokami, Noriko Maruyama and Tsuyoshi Sakai

March 16th, 2018

## Abstract

For a knot  $K$  in a homology 3-sphere  $\Sigma$ , let  $M$  be the result of  $2/q$ -surgery on  $K$ , and let  $X$  be the universal abelian covering of  $M$ . Our first theorem is that if the first homology of  $X$  is finite cyclic and  $M$  is a Seifert fibered space with  $N \geq 3$  singular fibers, then  $N \geq 4$  if and only if the first homology of the universal abelian covering of  $X$  is infinite. Our second theorem is that under an appropriate assumption on the Alexander polynomial of  $K$ , if  $M$  is a Seifert fibered space, then  $q = \pm 1$  (i.e. integral surgery).

## 1 Introduction

It is conjectured that Seifert surgeries on non-trivial knots are integral (except some cases), and that the resultant Seifert fibered spaces have three singular fibers (for example, see [DMM]). We [KMS1, KMS2] have studied the  $2/q$ -Seifert surgeries. One reason why the coefficients are  $2/q$  is that we started from the following shown in [Kd1] :

**Theorem 1.1** ([Kd1, Theorem 1.4]) *Let  $K$  be a knot in a homology 3-sphere  $\Sigma$  such that the Alexander polynomial of  $K$  is  $t^2 - 3t + 1$ . The only surgeries on  $K$  that may produce a Seifert fibered space with base  $S^2$  and with  $H_1 \neq \{0\}, \mathbb{Z}$  have coefficients  $2/q$  and  $3/q$ , and produce Seifert fibered space with three singular fibers. Moreover (1) if the coefficient is  $2/q$ , then the set of multiplicities is  $\{2\alpha, 2\beta, 5\}$  where  $\gcd(\alpha, \beta) = \gcd(\alpha, 5) = \gcd(\beta, 5) = 1$ , and (2) if the coefficient is  $3/q$ , then the set of multiplicities is  $\{3\alpha, 3\beta, 4\}$  where  $\gcd(\alpha, \beta) = \gcd(\alpha, 2) = \gcd(\beta, 2) = 1$ .*

In this paper, we continue to study the  $2/q$ -Seifert surgeries, where we replace the condition on the Alexander polynomial with a condition on the first homology of the double branched covering. We note that the results below are related to the conjectures mentioned above.

(1) Let  $\Sigma$  be a homology 3-sphere, and let  $K$  be a knot in  $\Sigma$ . Then  $\Delta_K(t)$  denotes the Alexander polynomial of  $K$ ,  $a_2(K)$  denotes the Conway's  $a_2$ -term of  $K$ , and  $\Sigma(K; p/r)$  denotes the result of  $p/r$ -surgery on  $K$ .

(2) The first author [Kd2] introduced the norm of polynomials and homology lens spaces: Let  $\zeta_d$  be a primitive  $d$ -th root of unity. For an element  $\alpha$  of  $\mathbb{Q}(\zeta_d)$ ,  $N_d(\alpha)$  denotes the norm of  $\alpha$

---

<sup>0</sup>2010 *Mathematics Subject Classification*: 11R04, 11R27, 57M25, 57M27.

*Keywords*: Reidemeister torsion, Casson-Walker-Lescop invariant, Seifert fibered space.

associated to the algebraic extension  $\mathbb{Q}(\zeta_d)$  over  $\mathbb{Q}$ . Let  $f(t)$  be a Laurent polynomial over  $\mathbb{Z}$ . We define  $|f(t)|_d$  by

$$|f(t)|_d = |N_d(f(\zeta_d))| = \left| \prod_{i \in (\mathbb{Z}/d\mathbb{Z})^\times} f(\zeta_d^i) \right|.$$

Let  $X$  be a homology lens space with  $H_1(X) \cong \mathbb{Z}/p\mathbb{Z}$ . Then there exists a knot  $K$  in a homology 3-sphere  $\Sigma$  such that  $X = \Sigma(K; p/r)$  ([BL, Lemma 2.1]). We define  $|X|_d$  and  $\|X\|_d$  by

$$|X|_d = |\Delta_K(t)|_d \quad \text{and} \quad \|X\|_d = \prod_{d'|d, d' \geq 1} |X|_{d'},$$

where  $d$  is a divisor of  $p$ . Then both  $|X|_d$  and  $\|X\|_d$  are topological invariants of  $X$  (Refer to [Kd2] for details).

(3) Let  $X$  be a closed oriented 3-manifold. Then  $\lambda(X)$  denotes the Lescop invariant of  $X$  ([Ls]). Note that  $\lambda(S^3) = 0$ .

## 2 Results

Let  $K$  be a knot in a homology 3-sphere  $\Sigma$ . For an odd integer  $q$ , let  $M$  be the result of  $2/q$ -surgery on  $K$ :  $M = \Sigma(K; 2/q)$ . Let  $\pi : X \rightarrow M$  be the universal abelian covering of  $M$  (i.e. the covering associated to  $\text{Ker}(\pi_1(M) \rightarrow H_1(M))$ ). Since  $H_1(M) \cong \mathbb{Z}/2\mathbb{Z}$ ,  $\pi$  is the 2-fold unbranched covering.

Let  $\Sigma_2$  be the double branched covering space of  $\Sigma$  branched along  $K$ , and  $\overline{K}$  the lifted knot of  $K$  in  $\Sigma_2$ . Since  $\overline{K}$  is null-homologous in  $\Sigma_2$ , and  $X$  is the result of  $1/q$ -surgery on  $\overline{K}$ , we have  $H_1(X) \cong H_1(\Sigma_2)$ .

$$\begin{array}{ccccc} \overline{K} \subset \Sigma_2 & \rightsquigarrow & X & = & \Sigma_2(\overline{K}; 1/q) \\ & \downarrow & \downarrow & & \\ K \subset \Sigma & \rightsquigarrow & M & = & \Sigma(K; 2/q) \end{array}$$

Assume that  $H_1(\Sigma_2) \cong \mathbb{Z}/m\mathbb{Z}$  with  $m \geq 3$ . Then  $H_1(X) \cong \mathbb{Z}/m\mathbb{Z}$ . Hence  $|X|_d$  is defined for each divisor  $d$  of  $m$ , and so is  $\|X\|_d$ .

We then have the following.

**Theorem 2.1** *Let  $K$  be a knot in a homology 3-sphere  $\Sigma$ . With the notation above, we assume that  $H_1(\Sigma_2) \cong \mathbb{Z}/m\mathbb{Z}$  with  $m \geq 3$ , and  $M = \Sigma(K; 2/q)$  is a Seifert fibered space with  $N$  singular fibers where  $N \geq 3$ . Then the following hold.*

(1)  $N \geq 4$  if and only if  $\|X\|_m = 0$ .

(2) If  $m$  is a power of a prime, then  $N = 3$  and the set of multiplicities is  $\{2\alpha, 2\beta, m\}$ , where  $\alpha, \beta$  and  $m$  are pairwise coprime.

**Theorem 2.2** *Let  $K$  be a knot in a homology 3-sphere  $\Sigma$ . We assume the following.*

(2.1)  $\lambda(\Sigma) = 0$ .

(2.2)  $|\Delta_K(-1)| = 5$ .

(2.3)  $|q| \geq 3$ ,  $\sqrt{|X|_5} > 4\{qa_2(K)\}^2$ .

*Then  $M = \Sigma(K; 2/q)$  is not a Seifert fibered space.*

### 3 Proof of Theorem 2.1

Since  $H_1(M) \cong \mathbb{Z}/2\mathbb{Z}$ , the base surface of  $M$  is  $S^2$ . As noted in Section 2,  $H_1(X) \cong H_1(\Sigma_2) \cong \mathbb{Z}/m\mathbb{Z}$ . Hence  $\|X\|_m$  is defined, and also we have

$$\|M\|_2 = |M|_2 = |\Delta_K(-1)| = |H_1(\Sigma_2)| = m \geq 3.$$

Hence, by [Kd1, Theorem 1.3 (2)], we may assume that  $M$  has a framed link presentation as in Figure 1, where  $\gcd(p_1, p_2) \geq 2$ ,  $p_k \geq 2$  for  $k = 1, \dots, N$ , and  $\gcd(p_i, p_j) = 1$  for  $1 \leq i < j \leq N$  and  $\{i, j\} \neq \{1, 2\}$ .




Figure 1: A framed link presentation of  $M = \Sigma(K; 2/q)$

Since  $H_1(M) \cong \mathbb{Z}/2\mathbb{Z}$ , we have  $\gcd(p_1, p_2) = 2$  by [Kd1, Theorem 1.2 (1)]. Hence we may assume that  $M$  has a framed link presentation as in Figure 2, where  $\alpha, \beta, p_3, \dots, p_N$  are pairwise coprime and  $p_3, \dots, p_N$  are all odd.




Figure 2: A framed link presentation of  $M = \Sigma(K; 2/q)$

Hence, as in [KMS1], we see that  $X$  has a framed link presentation as in Figure 3.




Figure 3: A framed link presentation of  $X$

Hence, by [Kd1, Theorem 1.3], we have (1).

Moreover, by [Kd1, Theorem 1.2 (3)], we have  $|M|_2 = p_3 \cdots p_N$ , and hence  $p_3 \cdots p_N = m$ . Therefore if  $m$  is a power of a prime, we have  $N = 3$  and  $p_3 = m$ . This completes the proof.  $\square$

## 4 Proof of Theorem 2.2

Suppose that  $M = \Sigma(K; 2/q)$  is a Seifert fibered space. Recall that  $|H_1(\Sigma_2)| = |\Delta_K(-1)| = 5$ . Then, by Theorem 2.1 (2), we may assume that  $M$  has a framed link presentation as in Figure 4, where  $1 \leq \alpha < \beta$  and  $\gcd(\alpha, \beta) = 1$ .




Figure 4: A framed link presentation of  $M = \Sigma(K; 2/q)$

Since  $\Delta_K(-1) \equiv \pm 3 \pmod{8}$  according to **(2.2)**, [Lc, Theorem 10.7, p108] implies that  $a_2(K) \neq 0$ .

As shown in [Kd1, Theorem 1.2 (3)],  $\sqrt{|X|_5} = (\alpha\beta)^2$ . By **(2.1)** and [Ls, 1.5 T2], we have  $\lambda(M) = qa_2(K)$ . By substituting in  $\sqrt{|X|_5} > 4\{qa_2(K)\}^2$ , we have  $(\alpha\beta)^2 > 4\{\lambda(M)\}^2$ . Hence we have  $|\lambda(M)| < (\alpha\beta)/2$ . The remaining part of the proof goes parallel to the proof of [KMS2, Theorem 2.1]. This completes the proof.  $\square$

**Acknowledgement** The first author is supported by Grant-in-Aid for Scientific Research(C) Number 17K05246, and the second author is supported by Grant-in-Aid for Scientific Research(C) Number 16K05158.

## References

- [BL] S. Boyer and D. Lines, *Surgery formulae for Casson's invariant and extensions to homology lens spaces*, J. Reine Angew. Math., **45** (1990), 181–220.
- [DMM] A. Deruelle, K. Miyazaki and K. Motegi, *Networking Seifert surgeries on knots*, Mem. Amer. Math. Soc. **217** (2012), no. 1021.
- [Kd1] T. Kadokami, *Reidemeister torsion of Seifert fibered homology lens spaces and Dehn surgery*, Algebr. Geom. Topol., **7** (2007), 1509–1529.
- [Kd2] T. Kadokami, *Reidemeister torsion and lens surgeries on knots in homology 3-spheres II*, Top. Appl., **155**, no.15 (2008), 1699–1707.
- [KMS1] T. Kadokami, N. Maruyama and T. Sakai, *Seifert surgery on knots via Reidemeister torsion and Casson-Walker-Lescop invariant*, Top. Appl., **188** (2015), 64–73.
- [KMS2] T. Kadokami, N. Maruyama and T. Sakai, *Seifert surgery on knots via Reidemeister torsion and Casson-Walker-Lescop invariant II*, Osaka J. Math., **53** (2016), 767–773.
- [Ls] C. Lescop, *Global surgery formula for the Casson-Walker invariant*, Ann. of Math. Studies, Princeton Univ. Press., **140** (1996).
- [Lc] W. B. R. Lickorish, *An introduction to knot theory*, Graduate texts in mathematics, **175** (1997).

Teruhisa Kadokami  
School of Mechanical Engineering,  
College of Science and Engineering, Kanazawa University,  
Kakuma-machi, Kanazawa, Ishikawa, 920-1192, Japan  
[kadokami@se.kanazawa-u.ac.jp](mailto:kadokami@se.kanazawa-u.ac.jp)

Noriko Maruyama  
Musashino Art University,  
Ogawa 1-736, Kodaira, Tokyo 187-8505, Japan  
[maruyama@musabi.ac.jp](mailto:maruyama@musabi.ac.jp)

Tsuyoshi Sakai  
Department of Mathematics, Nihon University,  
3-25-40, Sakurajosui, Setagaya-ku, Tokyo 156-8550, Japan