# Inference of topology and the nature of synapses, and the flow of information in neuronal networks

F. S. Borges<sup>1</sup>, E. L. Lameu<sup>2</sup>, K. C. Iarosz<sup>1,3</sup>, P. R. Protachevicz<sup>4</sup>, I. L.

Caldas<sup>1</sup>, R. L. Viana<sup>5</sup>, E. E. N. Macau<sup>2</sup>, A. M. Batista<sup>1,3,4,5</sup>, M. S. Baptista<sup>3</sup>

<sup>1</sup>Physics Institute, University of São Paulo, São Paulo, SP, Brazil.

<sup>2</sup>National Institute for Space Research, São José dos Campos, SP, Brazil.

<sup>3</sup>Institute for Complex Systems and Mathematical Biology, University of Aberdeen, SUPA, UK.

<sup>4</sup>Post-Graduation in Science, State University of Ponta Grossa, Ponta Grossa, PR, Brazil.

<sup>5</sup>Physics Department, Federal University of Paraná, Curitiba, PR, Brazil.

<sup>6</sup>Mathematics and Statistics Department, State University of Ponta Grossa, Ponta Grossa, PR, Brazil.

(Dated: July 19, 2018)

The characterisation of neuronal connectivity is one of the most important matters in neuroscience. In this work, we show that a recently proposed informational quantity, the causal mutual information, employed with an appropriate methodology, can be used not only to correctly infer the direction of the underlying physical synapses, but also to identify their excitatory or inhibitory nature, considering easy to handle and measure bivariate time-series. The success of our approach relies on a surprising property found in neuronal networks by which non-adjacent neurons do "understand" each other (positive mutual information), however this exchange of information is not capable of causing effect (zero transfer entropy). Remarkably, inhibitory connections, responsible for enhancing synchronisation, transfer more information than excitatory connections, known to enhance entropy in the network. We also demonstrate that our methodology can be used to correctly infer directionality of synapses even in the presence of dynamic and observational Gaussian noise, and is also successful in providing the effective directionality of inter modular connectivity, when only mean fields can be measured.

PACS numbers: 87.10Hk, 87.19.lj, 87.19.lw

Many real systems have been modelled by complex networks with different topological characteristics. Network theory has been applied in a large number of examples and different research fields, such as biology [1], economics [2] and physics [3]. In neuroscience, the application of network theory provides a way to analyse the structure and the functional behaviour of neuronal systems [4]. A fundamental research topic in neuroscience is the determination of the structure of the brain, to better understand its functioning. Some neuronal networks had their structure directly mapped by means of diffusion tensor imaging tractography [5].

One of the most challenging problems in neuronal networks is the inference of its topology, that is, the determination of the underlying synaptic connectivity by indirect means, based on functional measurements of time-series of the membrane potential [6, 7]. There are works that infer the topology based on functional measures such as correlation [8, 9] and synchronisation [10], or functional magnetic resonance imaging [11]. And there are those based on informational quantities [12-14, 17, 18]. Inference based on functional measures requires a threshold analysis that establishes a link between the measurement and the physical connection [12, 19, 20]. Rubido et al. [17] showed that a threshold can be calculated whenever a functional measure between nodes (cross correlation (CC) or mutual information) in a network is dissimilar. Higher functional values correspond to a pair of adjacent nodes, lower functional values to non-adjacent nodes. Bianco-Martinez et al. [18] used the mutual information rate (MIR) to successfully infer the connectivity

of a network composed of Hindmarsh-Rose (HR) neurons [21] connected by electrical synapses. Both works in Refs. [17, 18] have shown that the threshold technique could surprisingly provide an inferred network that matched exactly with the real network. These works have considered undirected networks, where nodes were connected bidirectionally with the same intensity.

This work considers HR networks with chemical synapses. Unlike electrical synapses that are undirected, chemical synapses are directed [22]. Whereas undirected networks can have their topologies properly inferred by CC and MIR, directed networks require methodologies capable of detecting the directionality of the physical influence [12, 19, 23]. Granger causality [12] is a concept construct on the idea that one can obtain optimal fittings of mathematical models about the measured time-series that provide the structure and direction of the connectivity. These models are statistically optimised to improve the predictability of events in one time-series based on observations of other time-series and have been shown to be a powerful tool to infer [24]. Informational quantities have also been demonstrated to provide a framework that is at the frontier to infer. In Ref. [17] it was shown that inference based on mutual information is more reliable than those based on correlational measurements. In Ref. [23] it was shown that directed information had advantages over Granger causality for quantifying effective connectivity in the brain. One question that remains open is whether information measures can reliably infer the connectivity of complex neuronal networks for all existing synapses by only accessing bivariate measurements, in

contrast to more complex and computational demanding techniques such as multivariate analysis based on informational analysis [14, 15], a technique that takes into consideration time-series from more than two neurons at each time, or modelled-based multivariate approaches such as those that employ compressive sensing [16].

In this work, we use the recently defined causal mutual information (CaMI) [25, 26] calculated using an appropriate methodology to infer the direction of chemical synapses in complex neuronal networks without any mistake, by only considering easy to handle and to measure bivariate time-series. Moreover, we show that inhibitory connections are responsible for a considerably larger amount of information transfer than that compared to neurons connected by excitatory synapses. This allows one to infer also the nature of the connection (excitatory and inhibitory), and not only its existence as previous techniques. Furthermore, we will also show that non-adjacent neurons transmit roughly null amount of directed information, indicating that indeed causal information has a direct relationship with the existence of a synapse.

The CaMI was constructed from the idea that if there is a flow of information from a system A to a system B, then longer time-series (or measurements with higher precision) from B should have a positive mutual information to short time-series (or to observations with lower precision) in A. This quantity, measuring the influence from A to B, was shown to be equal to the transfer entropy (TE) [13] from A to B plus the mutual information between A and B when both systems are being measured with the same resolution. The advantage of CaMI however is that it allows one to calculate TE, and therefore the directionality of the flow of information, by using measurements with arbitrary resolution. Which in turn also allows for the correct calculation of the TE using binary partitions of the phase space, i.e., appropriated when measurements have the lowest possible resolution. Moreover, CaMI can be calculated in lower-dimensional space of only 2 dimensions, without the need to consider conditional probabilities, but only marginal and joint probabilities, and finally, it is a quantity that fully express not only the exchange of information (MIR), but also its causal directionality (TE).

We consider the random neuronal network (RNN) [27, 28] introduced by Gelenbe [29] and the neuronal network of the nematode worm C. elegans [30] whose structure was completely mapped at a cellular level [31]. The node dynamics in the network is expressed by the Hidmarsh-Rose (HR) neuron model. Hindmarsh and Rose [21] proposed a phenomenological neuron model that is a simplification of the Hodgkin-Huxley model [32]. The HR is described by  $\dot{p} = q - ap^3 + bp^2 - n + I_{\text{ext}}$ ,  $\dot{q} = c - dp^2 - q$ ,  $\dot{n} = r[s(p - p_0) - n]$ , where p(t) is the action potential of the membrane, q(t) is related to the fast current,  $K^+$  or  $Na^+$ , and n(t) is associated with the slow current, for instance,  $Ca^{2+}$ . We use the parameters  $a = 1, b = 3, c = 1, d = 5, s = 4, r = 0.005, p_0 = -1.60$ 

and  $3.24 \leq I_{\text{ext}} \leq 3.25$ , so that the HR neuron exhibits a chaotic burst behaviour. Pre-synaptic neurons with an action potential  $p_j$  coupled by chemical synapses to neurons *i* modifying its action potential  $p_i$  according to  $\dot{p}_i = q_i - ap_i^3 + bp_i^2 - n_i + I_{\text{ext}} + g_c(V_{\text{syn}} - p_i) \sum_{j=1}^N \varepsilon_{ij} \Gamma(p_j)$ , where  $(i, j) = 1, \dots, N$  and *N* is the neurons number. The chemical synapse function is modelled by the sigmoidal function  $\Gamma(p_j) = \frac{1}{1 + \exp[-\lambda(p_j - \Theta_{\text{syn}})]}$ , with  $\Theta_{\text{syn}} =$  $1.0, \lambda = 10, V_{\text{syn}} = 2.0$  for excitatory and  $V_{\text{syn}} = -1.5$  for inhibitory synapses. The adjacency matrix  $\varepsilon_{ij}$  describes the neurons chemically connected. To do our analysis, we normalise  $p_i$  through the equation  $x_i = \frac{p_i^{\max} - p_i}{p_i^{\max} - p_i^{\min}}$ , where  $p_i^{\max}$  and  $p_i^{\min}$  are the maximum and minimum values, respectively, of the time-series of  $p_i(t)$ .

In order to be able to describe most of the information content of the time-series by a short-length binary symbolic representation, we make a time-Poincaré map of the time-series. Ideally, in the case one wants short symbolic sequences to fully express the amount of information of infinitely long sequences, points in the mapping should be spaced by a time step such that the symbolic representation of the time-series behaves as a random process, i.e., the next symbolic sequence is decorrelated with the previous. We are interested in obtaining a good estimation of CaMI to correctly infer the network's topology, its synaptic nature and to obtain a sufficiently accurate value for the magnitude of the flow of information (e.g., CaMI, MIR and TE). Given a time step  $\Delta t$ , a mapping for neuron  $i X_i$  is constructed by collecting a point of the membrane potential at times  $t = n\Delta t$  producing the discrete time-series described  $x_i^n = x_i$   $(t = n\Delta t)$ . In this way, we obtain the mapping  $X_i = x_i^0, x_i^1, x_i^2, ..., x_i^{T-1}$  for neuron i, where T is the number of points in the mapping. In the following, we will study coupled neurons to determine a time step for which CaMI is maximised, aiming with this maximisation to construct a time Poincaré map that tends to behave as a Markov process, allowing CaMI, MI and TE to express a good approximant of their real values. Figure 1 shows the normalised membrane potential for two chemical coupled HR neurons with connection from  $x_2$  (red line) to  $x_1$  (black line). The black and red circles correspond to  $X_1$  and  $X_2$ , respectively, where the mapping step time  $\Delta t$  is equal to 1ms. With the forward-time trajectory  $X_i^L(n) = x_i^n, \dots, x_i^{n+L-1}$ , where L is the length of the time-series X(n) and n is the discrete time, we generate a symbolic sequence  $S_i^L(n) = s_i^n, \cdots, s_i^{n+L-1}$ (Supplementary Material), where we consider  $s_i^n = 0$  if  $x_i^n \leq 0.5 \text{ and } s_i^n = 1 \text{ if } x_i^n > 0.$ 

Bianco-Martinez and Baptista [25, 26] defined a new quantity named CaMI from  $X_i$  to  $X_j$  (CaMI<sub> $X_i \to X_j$ </sub>) as the MI between joint events in  $X_i^{-L}$  and the set composed by the joint events of  $X_j^{-L}$  and  $X_j^L$  as CaMI<sub> $X_i \to X_j$ </sub> = MI( $X_i^{-L}; (X_j^{-L}, X_j^L)$ ) = MI( $X_i^L; W_j^{2L}$ ), where the mutual Information MI( $X_i^L; X_j^L$ ) is given by MI( $X_i^L; X_j^L$ ) =  $H(X_i^L) + H(X_j^L) - H(X_i^L, X_j^L)$ , and  $H(X_i^L)$  is the Shannon entropy of length-L tra-



FIG. 1. (Colour online) Normalised membrane potential of two chemical coupled HR neurons with connection from  $x_2$ (red line) to  $x_1$  (black line). We consider the coupling strength  $g_c = 1$  and the mapping time step  $\Delta t = 1$ ms. The black and red circles correspond to  $X_1$  and  $X_2$ , respectively.

jectory points of the discrete mapping. It is also true that  $\operatorname{CaMI}_{X_i \to X_j} = \operatorname{MI}(X_i^L; X_j L) + \operatorname{TE}_{X_i \to X_j}$ . Probabilities to calculate CaMI are constructed considering the probabilities of the encoded binary symbolic sequences. CaMI is thus calculated by  $CaMI_{X_i \to X_i} =$  $\sum_{S_i} \sum_{S_j} P(S_i^L, S_j^{2L}) \log \frac{P(S_i^L, S_j^{2L})}{P(S_i^L)P(S_j^{2L})}, \text{ where the summation indexes } S_i \text{ and } S_j \text{ represent the space of possible}$ length-L symbolic sequences coming from neuron i and  $S_i$  the space of possible joint events of finding a length-L symbolic sequence coming from neuron j at time n - Lfollowed by a length-L symbolic sequence in this same neuron at time n, or in other words, of finding a length-2L symbolic sequence in neuron j starting at the time n-L.  $P(S_i^L)$  is the probability of finding symbolic sequences  $S_i^L = \{s_i, \dots, s_i^{L-1}\}$  in  $X_i$ ,  $P(S_j^{2L})$  is the probability of finding a particular length-*L* symbolic sequences  $S_i^{2L} = \{s_i, \dots, s_i^{2L-1}\}$  in  $X_j$ , and  $P(S_i^L, S_j^{2L})$ is the joint probability between length-L symbolic sequences in neuron i and length-2L symbolic sequences in neuron j. The directionality index defined in Ref. [13] in terms of the TE can be calculated by  $DI_{X_i \to X_i} =$  $\operatorname{CaMI}_{X_i \to X_j} - \operatorname{CaMI}_{X_j \to X_i}$ . For simplicity in notation we consider that  $DI_{X_i \to X_j} \equiv DI_{ij}$ . This index measures the net amount of directed information flowing from  $X_i$ to  $X_i$ . Thus, if  $DI_{ij}$  is positive (negative), there is a net amount of information flowing from neuron i to neuron j (from neuron j to neuron i). Our hypothesis, also sustained by the works of [17, 18] and others is that if there is a directed adjacent connection from neuron i to j, thus  $DI_{ij}$  will be considerably larger than the directionality index of neurons that are not adjacently connected. So, the connection is  $X_i \to X_j$  if  $DI_{ij} > h$ , the connection is  $X_j \to X_i$  if  $\mathrm{DI}_{ij} < -h$ , and there is no connection if  $DI_{ij} \cong 0$ . In the latter case, the directionality index will be close to zero because the transfer entropy

will be roughly zero for non-adjacent nodes. The mutual information is a symmetric quantity and therefore  $MI(X_j, X_i) = MI(X_i, X_j).$ 



FIG. 2. (Colour online) Directional index (DI) as a function of the coupling strength  $(g_c)$  for the mapping step time  $\Delta t =$ 0.25ms. We consider L = 1 (black line), L = 2 (red line), L = 4 (green line), and L = 8 (blue line).

In Fig. 2 we calculate DI as a function of  $g_c$  for two coupled neurons with one directional connection from  $x_1$ to  $x_2$ . We observe that DI = 0 when the neurons are uncoupled ( $g_c = 0$ ), and DI > 0 for  $g_c > 0$ . The information is transmitted from  $x_1$  to  $x_2$ , in accordance with the direction of the connection. We compute DI for L = 1(black line), L = 2 (red line), and L = 8 (blue line). For the following analysis, we fix  $\Delta t = 0.25$ ms, and L = 8that maximises DI values.

Next, we build a directed network where the connections among the neurons are randomly chosen. We consider a random neuronal network with 64 HR neurons and average degree of connectivities K equal to 4. As a consequence, the network has 256 of a total of 4096 directed connections (ij). Figure 3 shows the normalised directional index, ranked from larger to smaller values, for 3 different neuronal connectivity configurations: 256 excitatory synapses (black line), 256 inhibitory synapses (red line), and 128 excitatory and 128 inhibitory synapses (blue line). In Fig. 3(a) there are 2 regions with  $DI_{ij} \neq 0$ , that represent the connections from *i* to *j*, while  $DI_{ij} \approx 0$ corresponds to the situation in that there is no connection between i and j. The magnification (Fig. 3(b)) exhibits two abrupt transitions. The transition to  $DI_{ij} \approx h$  allows the detection of directed connections in the neuronal network. The transition that occurs for  $DI_{ij} > h$  allows to infer the excitatory and inhibitory synapses, as shown by the blue line, where we observe the existence of 128 excitatory and 128 inhibitory synapses.

Notice that the DI values between adjacent and nonadjacent neurons are notably dissimilar, meaning that a small threshold h can be chosen such that  $DI_{ij} > h$ implies a directed connection from neuron i to neuron j. For the network whose neurons are connected by both inhibitory and excitatory synapses, we notice in the blue line of Fig. 3 two ranges of DI dissimilar values. For  $h < DI_{ij} < 0.4$ , the connection is excitatory and for  $DI_{ij} > 0.4$  the connection is inhibitory. In Fig. 3(c) we see the adjacency matrix, where the coloured elements of the matrix indicate if the pairs of neurons are connected.



FIG. 3. (Colour online) (a) Normalised directional index, ranked from larger to smaller values, for a random neuronal network with N = 64 HR neurons, K = 4,  $\Delta t = 0.25$ , L = 8,  $T = 4 \ 10^6$ , and  $g_c = 0.1$ . We consider 3 cases for the connectivity: 256 excitatory synapses (black line), 256 inhibitory synapses (red line), and 128 excitatory and 128 inhibitory synapses (blue line). (b) Magnification of (a). (c) Matrix of the normalised directional index (DI<sub>ii</sub>) of latter case.

The uncoupled pairs of neurons are indicated in black, while the coupled pairs are in colour scale according to the normalised directional index. We consider the same parameters used to calculate the blue line in Fig. 3. For  $DI_{ij} < 0.4$  the colour scale shows the excitatory synapses and for  $DI_{ij} \geq 0.4$  the synapses are inhibitory.



FIG. 4. (Colour online) Normalised directional index for  $\sigma_d = 3$  (black line) and  $\sigma_d = 4$  (green line). We consider  $\Delta t = 0.25$ , L = 8, and  $g_c = 0.15$ .

We analyse the noise effect in the inference of the connections. Neuronal noise can be related to several sources, such as synaptic noise [33] and ion conductance noise [34]. In the action potential equation, we add a Gaussian noise with zero mean and variance  $\sigma_d$ . We calculate the DI<sub>ij</sub> values for the neuronal network with  $\sigma_d = 3$  (black line) and  $\sigma_d = 4$  (green line), as shown

in Fig. 4. We verify that the inference for the existence of a synapse is robust to dynamic noise in the membrane potential. However, for  $\sigma_d \gtrsim 3.5$  it is not possible to infer whether the synapse is excitatory or inhibitory. Therefore, the inference of the connectivities is more robust than the inference of its nature of the synapses. CaMI-based inference is also robust to additive noise of moderate amplitude (Supplementary Material).



FIG. 5. Normalised directional index, ordered from larger to smaller values for N = 277 HR neurons. We consider  $\sigma_d = 1$ ,  $g_c = 0.035$ ,  $\Delta t = 0.25$ , and L = 8.

In the literature, there are many works that consider C. elegans neuronal network to study nervous system [35, 36]. The C. elegans is a soil worm with body size about 1mm and a simple nervous system [37]. We consider in our study the connectome of the large somatic nervous system according to Ref. [38] that consists of 277 neurons. To test our inference approach, we consider approximately 50% of excitatory and 50% of inhibitory synapses in the C. elegans network with 1731 directed connections. The directed adjacency matrix  $(\varepsilon_{ii})$  is obtained from the brain connectivity of the C. elegans. Figure 5 exhibits the DI values, where the two discontinuities transitions in the DI values correspond to the excitatory and inhibitory synapses. In Fig. 5 is possible to identify the connected neurons of the C. elegans, where from i = 1 to i = 138 and from i = 139 to i = 277 there are 850 inhibitory synapses and 881 excitatory synapses, respectively.

In conclusion, we propose a successful methodology based on CaMI to infer, characterise and investigate the transmission of information in neuronal networks with chemical synapses. Through the CaMI, we show not only how to infer the existence of synapses, but also to identify the nature of the synapse. Our technique can be applied to time-series generated with Gaussian dynamical noise inbuilt in the neuron equations, or to time-series contaminated by observational noise (Supplementary Material). Moreover, we also showed that when access to the neuron potential is not possible, but rather only local mean fields can be measured, such as those coming from EEG signals, our CaMI-based technique can correctly determine the effective net directed connectivity between different neuronal clusters. This work also shows that excitatory connections are not so efficient to transfer information as inhibitory connections, and that non-adjacent neurons transfer roughly zero amount of information. This latter

observation suggests that a pre-synaptic neuron (a neuron that has an adjacent connection to the post-synaptic one) not only exchange information (positive mutual information), but is also capable of using information to cause an effect in a post-synaptic neuron (positive transfer entropy). Non-adjacent neurons only exchange information. This one-to-one relationship between structure and information transmission remains valid for a wide range of the coupling strength  $g_c$ , constraint within an interval with not so small (to prevent full decorrelation)

- A.-L. Barabási and Z. N. Oltvai, Nat. Rev. Genet. 5, 101 (2004).
- [2] M. P. Niemira and T. L. Saaty, Int. J. Forecasting 20, 573 (2004).
- [3] A. Arenas, A. Díaz-Guilera, J. Kurths, Y. Moreno, and C. Zhow, Phys. Rep. 469, 93 (2008).
- [4] E. Bullmore and O. Sporns, Nat. Rev. Neurosci. 10, 186 (2009).
- [5] G. Gong, Y. He, L. Concha, C. Lebel, D. W. Gross, A. C. Evans, and C. Beaulieu, Cerb. Cortex **19**, 524 (2009).
- [6] H. X. Ta, C. N. Yoon, L. Holm, and S. K. Han, BMC Syst. Biol. 4, 70 (2010).
- [7] A. M. Bastos and J.-M. Schoffelen, Front. Syst. Neurosci. 9, 1 (2016).
- [8] M. Takigawa, G. Wang, H. Kawasaki, and H. Fukuzako, Int. J. Psychophysiol. 21, 65 (1996).
- [9] A. Baccalá and K. Sameshima, Biol. Cybern. 84, 463 (2001).
- [10] X.-M. Cui, W. S. Kim, D.-U. Hwang, and S. K. Han, Europhys. Lett. **110**, 38001 (2015); Y. Chen, G. Rangarajan, J. Feng, and M. Ding, Phys. Lett. A 324, 26 (2004); N. Ancona, D. Marinazzo, and S. Stramaglia, Phys. Rev. E **70**, 056221 (2004).
- [11] M. P. van den Heuvel and H. E. H. Pol, Eur. Neuropsychopharmacol. 20, 519 (2010).
- [12] P.-O. Amblard and O. J. J. Michel, J. Comput. Neurosci. 30, 7 (2011).
- [13] T. Schreiber, Phys. Rev. Lett. 85, 461 (2000).
- [14] J. Runge, Phys. Rev. E **92**, 062829 (2015).
- [15] J. Sun, D. Taylor, and E. M. Bolt, SIAM J. Appl. Dyn. Syst. 14, 73 (2015).
- [16] W.-X. Wang, Y.-C. Lai, and C. Grebogi, Phys. Rep. 644, 1 (2016).
- [17] N. Rubido, A. C. Martí, E. Bianco-Martínez, C. Grebogi, M. S. Baptista, and C. Masoller, New J. Phys. 16, 093010 (2014).
- [18] E. Bianco-Martinez, N. Rubido, C. G. Antonopoulos, and M. S. Baptista, Chaos 26, 043102 (2016).
- [19] J. Massey, Causality, feedback and directed information (In Proc. Intl. Symp. on Info. Theory and Its Applications, Waikiki, Hawai, USA, 27, 1990).
- [20] I. Vlachos and D. Kugiumtzis, Phys. Rev. E 82, 016207 (2010).
- [21] J. L. Hindmarsh and R. M. Rose, Proc. R. Soc. London B 221, 87 (1984).
- [22] A. Pereda, Nat. Rev. Neurosci. 15, 250 (2014).
- [23] Y. Liu and S. Aviyente, Comp. Math. Methods Med. 2012, 635103 (2012).

and not so large (to prevent full synchronization) bounds. For an inference with no mistakes, time-series should be sufficiently long, more specifically, their size in seconds (i.e., t) should scale with the size of the network, and it is a function of the coupling strength, a relationship that was studied in much detail in Ref. [17].

Acknowledgement: CAPES, DFG-IRTG 1740/2, Fundação Araucária, Newton Fund, CNPq (154705/2016-0, 311467/2014-8), FAPESP (2011/19296-1, 2015/07311-7, 2016/16148-5, 2016/23398-8, 2015/50122-0), EPSRC-EP/I032606.

- [24] B. Schelter, M. Mader, W. Mader, L. Sommerlade, B. Platt, Y.-C. Lai, C. Grebogi, and M. Thiel, Europhys. Lett. 105, 30004 (2014).
- [25] E. J. Bianco-Martinez, PhD Thesis, University of Aberdeen, 2015.
- [26] E. Bianco-Martinez and M. S. Baptista, arXiv:1612 05023v1.
- [27] A.-L. Barabási and R. Albert, Science 286, 509 (1999).
- [28] S. Timotheou, Comput. J. **53**, 251 (2010).
- [29] E. Gelenbe, Neural Comput. 1, 502 (1989).
- [30] M. Kaiser and C. C. Hilgetag, PLoS Comput. Biol. 2, e95 (2006).
- [31] J. G. White, E. Southgate, J. N. Thomson, and S. Brenner, Phil. Trans. R. Soc. Lond. B 314, 1 (1986).
- [32] A. L. Hodgkin and A. F. Huxley, J. Physiol. 117, 500 (1952).
- [33] G. Györgyi, Phys. Rev. Lett 64, 2957 (1990).
- [34] X. J. Cao and D. Oertel, J. Neurophysiol. 94, 821 (2005).
- [35] K. S. Liu and P. W. Sternberg, Neuron 14, 79 (1995).
- [36] L. R. Varshney, B. L. Chen, E. Paniagua, D. H. Hall, and D. B. Chklovskii, PLoS Comput. Biol. 7, e1001066 (2011).
- [37] C. Gally, J. L. Bessereau JL, Med. Sci. 19, 725 (2003).
- [38] Connectome File Format-Datasets (Version 2.0). Available: http://cmtk.org/viewer/datasets/.

# INFERENCE OF TOPOLOGY AND THE NATURE OF SYNAPSES, AND THE FLOW OF INFORMATION IN NEURONAL NETWORKS: SUPPLEMENTARY MATERIAL

# ENCODING THE TRAJECTORY INTO SYMBOLIC SEQUENCES

Figure 1 (in the paper) shows the normalised membrane potential for two neurons coupled from  $x_2$  (red line) to  $x_1$  (black line). In table I, we show some mapped values of  $x_1^n$  and  $x_2^n$  with their respective length-1 symbolic values  $s_1^n$  and  $s_2^n$ , and also the length-2 and length-4 symbolic sequence  $S_1^{L=2}(n)$  and  $S_2^{L=4}(n)$ , respectively.

n	$x_1^n$	$s_1^n$	$S_1^{L=2}(n)$	$x_2^n$	$s_2^n$	$S_2^{L=4}(n)$
285	0.163043	0	00	0.374431	0	0110
286	0.161350	0	00	0.500448	1	1100
287	0.274266	0	00	0.886694	1	1000
288	0.265589	0	00	0.213396	0	0000
289	0.279589	0	00	0.174788	0	0000
290	0.306991	0	00	0.174349	0	0000
291	0.349396	0	00	0.173966	0	0000
292	0.427650	0	01	0.173642	0	0000
293	0.645130	1	11	0.173384	0	
294	0.725724	1	10	0.173200	0	
295	0.180110	0		0.173100	0	

TABLE I. Mappings for  $\Delta t = 1$ ms and L = 2.

Considering the symbolic sequences  $S_i^L$  and  $S_j^{2L}$  is possible to find the probabilities  $P(S_i^L)$ ,  $P(S_j^{2L})$  and  $P(S_i^L, S_j^{2L})$ . These probabilities are used to calculate the Casual Mutual Information

$$CaMI_{X_i \to X_j} = \sum_{S_i} \sum_{S_j} P(S_i^L, S_j^{2L}) \log \frac{P(S_i^L, S_j^{2L})}{P(S_i^L)P(S_j^{2L})}$$

and Directionality Index

$$DI_{X_i \to X_j} = CaMI_{X_i \to X_j} - CaMI_{X_j \to X_i}.$$
 (2)

## INFORMATIONAL QUANTITIES FOR ADJACENT NEURONS

The CaMI<sub>ij</sub>, measuring the influence from *i* to *j*, was shown to be equal to CaMI<sub>ij</sub> = MI<sub>ij</sub> + TE<sub>ij</sub>, where MI<sub>ij</sub> = MI<sub>ji</sub> is the mutual information, and TE<sub>ij</sub> is the transfer entropy from *i* to *j*. In Fig. 6, we compare the values of DI<sub>ij</sub>, CaMI<sub>ij</sub>, CaMI<sub>ji</sub>, and MI<sub>ij</sub>, as a function of  $g_c$  for two coupled neurons with one directional connection from  $x_i$  to  $x_j$ . We observe that CaMI<sub>ji</sub>  $\approx$  MI<sub>ij</sub>, therefore TE<sub>ji</sub>  $\approx$  0 and TE<sub>ij</sub>  $\approx$  DI<sub>ij</sub>.

In Fig. 7, we show the values of  $DI_{ij}$ ,  $CaMI_{ij}$ ,  $CaMI_{ji}$ ,  $CaMI_{ji}$ , and  $MI_{ij}$ , for o neuron i = 1 in a network with N = 64 neurons. There are conections from i = 1 to j = 29, 46, 54, and 65. In these case, we observe that  $CaMI_{ji} \approx MI_{ij}$ , therefore  $TE_{ij} \approx DI_{ij} > 0$ . Moreover,



FIG. 6. Directional index (DI), causal mutual information (CaMI), and mutual information (MI), as a function of the coupling strength  $(g_c)$  for the mapping step time  $\Delta t = 0.5$ ms and L = 8.

there are conections from j = 3, 5, and 19 to i = 1, where  $\mathrm{DI}_{ij} < 0$ . Finally, we can observe  $\mathrm{DI}_{ij} \approx 0$  when there are no connections.



FIG. 7. Directional index (DI), causal mutual information (CaMI), and mutual information (MI), for o neuron i = 1 in a network with N = 64 neurons, for the coupling strength  $g_c = 0.1$ , mapping step time  $\Delta t = 0.5$ ms, and L = 8.

### DELAY ANALYSIS

In Table 1, we show the case where no mapping delay is considered. We can insert a time delay by generating the symbol  $s_2^n$  using the values of  $x_2^{n+\text{delay}}$ . In Fig. 1 we show the Directionality Index (DI) as a function of mapping time step  $\Delta t$  and delay in this mapping for L = 2 and L = 6. For the dynamical model used, we find that the DI is higher if L = 6 than if L = 2. In addition, the DI values are higher in the region where  $\Delta t < 1.0$ and delay< 6ms. This shows that a possible delay in connections can be neglected in this case.



FIG. 8. Directionality index (DI) as a function of mapping step time  $\Delta t$  and delay in this mapping. (a) L = 2 and (b) L = 6.

#### ADDITIVE NOISE

The additive noise is related to the imprecision of the equipment responsible for capturing the electrical signals in the neural membrane, so in our simulations we add to the values of p(t) a noise with zero mean and standard deviation  $\sigma_a$ . In Figs. 9(a) and 9(b) we observe the change in the dynamics of the membrane potential of a network neuron under the application of additive noise with  $\sigma_a = 0.1$  and  $\sigma_a = 0.35$ , respectively. The difference between the minimum and maximum values reached by the membrane potential of the HR model is approximately 3.5, so  $\sigma_a = 0.35$  corresponds to 10% of this value. For  $\sigma_a = 0.1$  the observed dynamics remains very similar to the case with no noise observed in Fig. 1, however, when  $\sigma_a = 0.35$  the noise intensity can change the values of the symbolic sequence  $S_i^L(n)$ . In Fig. 9(c) we see that the DI calculation does not present significant changes when considering the additive noise with  $\sigma_a = 0.1$  (black line). For  $\sigma_a = 0.35$  (green line) it is no longer possible to distinguish excitatory connections from inhibitory ones, but all 256 connections are detected.

#### INFORMATION FLOW BETWEEN NETWORKS

In many experimental cases it is not possible to directly measure the membrane potential of each neuron, but only an average field of a group of them, or a brain region. Through the analysis of the mean field between two neural networks, we show that is possible to infer if distinct networks are connected to each other, and identify the direction of the effective connectivity by the direction of the flow of information.

In order to do this analysis, we considered two random networks with N = 64 neurons each, with average degree of intra connections within the networks  $K_{\text{intra}} = 24$ and average degree of inter connections between networks  $K_{\text{inter}} = 12$ . To study the flow of information between the two networks, we consider that there are only directed connections from neurons of network 1 to neurons of network 2. In each of the networks we calculated the mean



FIG. 9. Membrane potential for the neuron i = 32 for additive noise standard deviation (a)  $\sigma_a = 0.1$  and (b)  $\sigma_a = 0.35$ . (c) Normalised Directionality index for  $\sigma_a = 0.1$  (black line) and  $\sigma_a = 0.35$  (green line). We consider  $\Delta t = 0.5$ , L = 4, and  $g_c = 0.1$ .

field of the membrane potential and made the symbolic sequence using this time series. The process of calculating DI was performed in the same way as in the case of isolated neurons.



FIG. 10. (a) Directionality Index (DI) as a function of coupling strength  $(g_c)$ . For  $\Delta t = 0.25$  ms, where L = 1 (black line), L = 2 (red line), L = 4 (green line), and L = 8 (blue line). Time evolution of the Normalised Mean Field (NMF)for network 1 (black line) and network 2 (red line), both with (b)  $g_c = 0.025$ , (c)  $g_c = 0.175$ , and (d)  $g_c = 0.275$ .

In Fig. 10(a) we show the values obtained from the DI as a function of the intensity of the coupling  $g_c$ , where we set  $\Delta t = 0.25$ ms and we evaluate different sizes for

the symbolic trajectory: L = 1 (black line), L = 2 (red line), L = 4 (green line) and L = 8 (blue line). We find that, as in the case of two neurons, the highest DIvalues are observed when using symbolic trajectories of size L = 8. In this case, we observed that when the coupling is low, the DI values are small, since the influence of the network dynamics 1 on the network 2 is smaller. For a coupling around  $g_c = 0.175$  we have the highest calculated value of DI and for  $g_c > 0.275$  the value of DI decreases, tending to a constant value. This happens when the neurons of both networks are roughly completely synchronous. The neurons had been completely synchronous, thus the Transfer Entropy would be zero, resulting in a DI of zero. To understand more about the dynamical behaviour leading to the curve presented by the blue line in Fig. 10(a), we analyse the temporal evolution of the normalised mean field (NMF) for three values of the coupling. In Fig. 10(b) we have  $g_c = 0.025$ and we observe that the NMF of network 1 (black line) and network 2 (red line) show that the neurons of these

networks present the behaviour of bursting synchronisation, when neurons start the bursting of firing activities roughly simultaneously. Firings are asynchronous. In Fig. 10(c) we have  $g_c = 0.175$  and the NMF of network 1 and 2 show that not only intra but also inter neurons are roughly synchronous. Firing spikes in the NMF indicates intra synchronisation. Inter synchronisation is evidenced by the fact that the curves are roughly identical. These both factors are responsible for the high DI values. Finally, in Fig. 10(d) we have  $g_c = 0.275$  which is intense enough to make the networks to almost fully synchronise.

Therefore, even in the case when we have only the data of the average field of networks, we show that it is possible to infer the effective directionality of the connections in a similar way to the case between two neurons only. This method may be thus suitable to be considered for information flow studies in different regions of the brain, analysing data obtained from several experimental sources such as structural and functional MRI, diffusion tensor imaging, magnetoencephalography, and electroencephalography.