Robust Estimation in High Dimensional Generalized Linear Models

M. Valdora^{*1}, C. Agostinelli², and V.J. Yohai³

¹Departamento de Matematicas, Facultad de Ciencias Exactas y Naturales, University of Buenos Aires

²Department of Mathematics, University of Trento, Trento, Italy

³Departamento de Matematicas, Facultad de Ciencias Exactas y Naturales, University of Buenos Aires and CONICET

October 2, 2017

Abstract

Generalized Linear Models are routinely used in data analysis. The classical procedures for estimation are based on Maximum Likelihood and it is well known that the presence of outliers can have a large impact on this estimator. Robust procedures are presented in the literature but they need a robust initial estimate in order to be computed. This is especially important for robust procedures with non convex loss function such as redescending M-estimators. Subsampling techniques are often used to determine a robust initial estimate; however when the number of unknown parameters is large the number of subsamples needed in order to have a high probability of having one subsample free of outliers become infeasible. Furthermore the subsampling procedure provides a non deterministic starting point. Based on ideas in Peña and Yohai [1999], we introduce a deterministic robust initial estimate for M-estimators based on transformations [Valdora and Yohai, 2014] for which we also develop an iteratively reweighted least squares algorithm. The new methods are studied by Monte Carlo experiments.

^{*}Corresponding author mvaldora@gmail.com

Keywords: Initial estimates, Outliers, Transformed Least Squares estimators, Transformed M-estimators,

1 Introduction

We consider high dimensional generalized linear models and we study a robust method for estimating its parameters. Robust estimators for generalized linear models (GLM) have been studied by Künsch et al. [1989], Cantoni and Ronchetti [2001], Bergesio and Yohai [2011], Bianco et al. [2013], Valdora and Yohai [2014] and Algallaf and Agostinelli [2016]. However, these proposals either lack robustness or require a robust initial estimator. We propose a method for computing an initial estimator which can be used to begin an iterative algorithm, as needed by redescending estimators. We apply this method in the computation of M-estimators based on transformations (MT) proposed by Valdora and Yohai [2014]. MT-estimators are a family of Mestimators based on variance stabilizing transformations which are shown to be highly robust and efficient by means of a Monte Carlo study. These estimators are redescending M-estimators applied after transforming the responses by means of a variance stabilizing function. Stabilizing the variance allows the correct scaling of the loss function used in the definition of the M-estimator.

Consider a GLM in which y is the response and \mathbf{x} is a p-dimensional vector of explanatory variables. We assume that

$$g(\mu) = \boldsymbol{\beta}_0^{\top} \mathbf{x},\tag{1}$$

where $\beta_0 \in \mathbb{R}^p$ is an unknown vector of parameters and $g : \mathbb{R} \to \mathbb{R}$ is a known link function. We further assume that

$$y|\mathbf{x} \sim F_{\mu},$$
 (2)

where F_{μ} is a discrete or continuous model in the exponential family of distributions in \mathbb{R} , that is to say, it has a density of the form

$$f_{\mu}(y) = \exp(((y\mu - b(\mu))/a(\phi) + c(y,\phi)), \qquad (3)$$

for given functions a, b and c. We assume ϕ is known. MT-estimators are

defined as follows

$$L(\boldsymbol{\beta}) = \sum_{i=1}^{n} \rho \left(t(y_i) - m \left(g^{-1} \left(\mathbf{x}_i^{\top} \boldsymbol{\beta} \right) \right) \right)$$
$$\hat{\boldsymbol{\beta}} = \arg \min_{\boldsymbol{\beta}} L(\hat{\boldsymbol{\beta}})$$
(4)

where $\rho(u)$ is a symmetric, bounded, continuous and non decreasing on |t| function, t is a variance stabilizing transformation and m is the function defined by

$$m(\mu) = \operatorname{argmin}_{\gamma} \mathbb{E}_{\mu} \left(\rho \left(t(y) - \gamma \right) \right), \tag{5}$$

where $\mathbb{E}_{\mu}(y)$ denotes the expectation of y when y has distribution F_{μ} . It is assumed that m is univocally defined, therefore (5) implies the Fisher consistency of $\hat{\beta}$. Other assumptions necessary to have consistency and asymptotic normality of this estimators are listed in Valdora and Yohai [2014]. The solution to (4) can be found by iterative methods which typically solve the corresponding system of estimating equations

$$\sum_{i=1}^{n} \psi(\mathbf{x}_i, y_i, \boldsymbol{\beta}) = 0.$$
(6)

where $\psi(\mathbf{x}_i, y_i, \boldsymbol{\beta})$ is the derivative with respect to $\boldsymbol{\beta}$ of $\rho(t(y_i) - m(g^{-1}(\mathbf{x}_i^{\top}\boldsymbol{\beta}))))$. In the Appendix we provide an iteratively reweighted least squares (IRWLS) algorithm to find a solution to equation (6). The difficulty in the case of redescending M-estimators is that the goal function $L(\beta)$ might have several local minima. As a consequence, it might happen that the iterative procedure converges to a solution of equation (6) that is not a solution of the optimization problem (4). To avoid this, one must begin the iterative algorithm at an initial estimator which is a very good approximation of the absolute minimum of L, i.e. the solution of (4). If p is small, this approximate solution may be obtained by the subsampling method Valdora and Yohai [2014, see]. Based on the algorithm described in Rousseeuw and Leroy [1987] for linear models, this method consists in computing a finite set A of candidate solutions to (4) and then replace the minimization over \mathbb{R}^p by a minimization over A. The set A is obtained by randomly drawing subsamples of size p and computing the maximum likelihood (ML) estimator based on the subsample. If the original sample contains a proportion ϵ of outliers, then the probability that a given subsample is free of outliers is $(1 - \epsilon)^p$ and the probability of

having at least one subsample free of outliers is $1 - (1 - (1 - \epsilon)^p)^N$, where N is the number of subsamples drawn. If we want this probability to be greater than a given α , we must draw a number of subsamples such that

$$1 - (1 - (1 - \epsilon)^p)^N > \alpha,$$

that is to say,

$$N > \frac{\log(\alpha)}{\log(1 - (1 - \varepsilon)^p)} \simeq \left| \frac{\log(\alpha)}{(1 - \varepsilon)^p} \right|.$$

This makes the algorithm infeasible for large p.

Peña and Yohai [1999] studied this problem in the case of linear models, introducing an alternative method to compute the set of candidate solutions A. Their proposal succeeds in obtaining a set A which contains very good approximations of the actual solution and, on the other hand, requires the computation of a small number of subsamples, namely 3p + 1. This makes the algorithm much faster and feasible even for very large values of p.

We modify the method introduced by Peña and Yohai [1999] in order to apply it to generalized linear models. We study its application to MTestimators by means of an extensive Monte-Carlo study, which shows that the method is very fast and robust for large values of p.

As a particular case of the MT-estimator we define the Least Squares estimator based on Transformations (LST), which corresponds to $\rho(u) = u^2$, in the following way

$$\hat{\boldsymbol{\beta}} = \operatorname{argmin}_{\boldsymbol{\beta}} \sum_{i=1}^{n} \left(t(y_i) - \mathbb{E}_{\left(g^{-1}\left(\mathbf{x}_i^{\top} \boldsymbol{\beta}\right)\right)}\left(t(y_i)\right) \right)^2.$$
(7)

This estimator can be seen as a natural generalization of the Least Squares estimator (LS) for linear models to the case of GLM. LST estimators are Fisher consistent, however since ρ is not bounded, they are, in general, non robust. In the Appendix we provide an iteratively reweighted least squares algorithm to find the solution to the optimization problem (7).

2 Detecting Outliers Using Principal Sensitivity Components

The classical statistic used to measure the influence of an observation is the Cook statistic introduced by Cook [1977] for linear models, which can be adapted for generalized linear models (see Chapter 12 of McCullagh and Nelder [1989]). This statistic is a measure of the distance between $\hat{\boldsymbol{\beta}}$, the maximum likelihood estimator and $\hat{\boldsymbol{\beta}}_{(i)}$, the maximum likelihood estimator computed without observation *i*. However, these measure is non-robust and therefore, when there are several ouliers, it may be completely unreliable. In these cases, some outliers (y_i, \mathbf{x}_i) with high influence may have a small Cook statistic if there are other similar outliers that still influence $\hat{\boldsymbol{\beta}}_{(i)}$. This is known as masking effect. To make things worse, high leverage outliers may have small residuals making their detection difficult. This situation usually arises when there are several similar, or highly correlated outliers.

The proposal of Peña and Yohai (1999) follows the same idea as the subsampling method but it computes the set of candidate solutions A in a different way. The candidates are obtained, as before, by computing the least squares estimates on subsamples. However, the subsamples are not chosen at random. Instead, they are chosen by deleting from the sample, groups of similar or highly correlated outliers, which can potentially cause a masking effect. The set A will, in this way, contain candidates which are already quite robust estimates and therefore it will not need to have a large number of candidates in the set A is only 3p + 1.

Let $(\mathbf{x}_1, y_1), \ldots, (\mathbf{x}_n, y_n)$ be random vectors which follow a generalized linear model as defined by (2) and (1). Let $\hat{\boldsymbol{\beta}}$ be the LST estimator and

$$\hat{\boldsymbol{\mu}} = (\hat{\mu}_1, \dots, \hat{\mu}_n)^\top = g^{-1}(\mathbf{X}\hat{\boldsymbol{\beta}})$$

be the vector of fitted values. Let $\hat{\mu}_{i(j)}$ be the fitted value for observation i computed without using observation j, that is $\hat{\mu}_{i(j)} = g^{-1}(\mathbf{x}_i^{\top} \hat{\boldsymbol{\beta}}_{(j)})$, where $\hat{\boldsymbol{\beta}}_{(j)}$ is the LST estimate based on the original sample without observation j. We define the *i*-th residual e_i as the difference between $t_i = t(y_i)$ and its predicted value $\hat{t}_i = m(g^{-1}(\mathbf{x}_i^{\top} \hat{\boldsymbol{\beta}}))$, that is $e_i = t_i - \hat{t}_i$. Following the ideas introduced by Peña and Yohai (1999) for linear models, we define the sensitivity vectors as the vectors \mathbf{r}_i with entries

$$r_{ij} = \hat{t}_i - \hat{t}_{i(j)}$$

where $\hat{t}_{i(j)} = m(\hat{\mu}_{i(j)})$ is the predicted value of t_i computed without using observation j. Then, r_{ij} is the sensitivity of the forecast of the t_i to the deletion of observation j and the sensitivity vectors are defined by

$$\mathbf{r}_i = (r_{i1}, \ldots, r_{in}) \quad , \qquad 1 \le i \le n.$$

The sensitivity matrix \mathbf{R} is defined as the matrix whose rows are the vectors \mathbf{r}_i . Let

$$\mathbf{v}_{1} = \operatorname{argmax}_{||\mathbf{v}||=1} \sum_{i=1}^{n} \left(\mathbf{v}^{\top} \mathbf{r}_{i} \right)^{2}.$$
(8)

 \mathbf{v}_1 is the direction in which the projections of the sensitivity vectors is largest. Let

$$\mathbf{z}_1 = \mathbf{R}\mathbf{v}_1; \tag{9}$$

then \mathbf{z}_1 is the vector whose entries are the terms of the sum in (8). Therefore, the largest entries in \mathbf{z}_1 correspond to the largest terms in the sum in (8), which in turn correspond to the observations that have the largest projected sensitivity in the direction \mathbf{v}_1 .

In the same way, we can define recursively \mathbf{v}_i , $2 \leq i \leq n$ as the solution of

$$\mathbf{v}_{i} = \operatorname{argmax}_{||\mathbf{v}||=1} \sum_{i=1}^{n} \left(\mathbf{v}^{\top} \mathbf{r}_{i} \right)^{2}.$$
(10)

subject to
$$\mathbf{v}_i \mathbf{v}_j = 0$$
 for all $1 \le j < i$ (11)

The vectors $\mathbf{v}_1 \dots \mathbf{v}_p$ are the directions in which the projected sensitivity of the observations are the largest. The corresponding projections

$$\mathbf{z}_i = \mathbf{R}\mathbf{v}_i \tag{12}$$

are called the principal sensitivity components. The entries of \mathbf{z}_i are the projections of the sensitivity vectors on the direction \mathbf{v}_i . Large entries correspond to observations whose projected sensitivity in the direction \mathbf{v}_i is large. Therefore, large entries are considered potential outliers.

High leverage observations typically have large sensitivity because a small change in the estimated slopes will cause a large change in the fitted values. Peña and Yohai [1999] prove that, in the case of linear models, if the sample is contaminated with less than (n-p+1)/(2n-p+1) high leverage outliers, then, at least for one eigenvector, the coordinates corresponding to the outliers have absolute value larger than the median.

3 Procedure for obtaining a robust initial estimate in generalized linear models

Consider a random sample following a generalized linear model as defined by (1), (2) and (3). The following procedure computes an approximation of β_0 which will be used as an initial estimator in the IRWLS algorithm for the estimating equation (6). The procedure has two stages. Stage 1 aims at finding a highly robust but possibly inefficient estimate and stage 2 aims at increasing its efficiency.

Stage 1. In this stage, the idea is to find a robust, but possibly inefficient, estimate of β by an iterative procedure. In each iteration $k \ge 1$ we get

$$\hat{\boldsymbol{\beta}}^{(k)} = \arg\min_{\boldsymbol{\beta} \in A_k} L(\boldsymbol{\beta}).$$
(13)

In the first iteration (k = 1) the set A_1 is constructed as follows. We begin by computing the LST estimate with the complete sample and the principal sensitivity components. For each principal sensitivity component \mathbf{z}_i we compute three estimates by the LST method. The first eliminating the half of the observations corresponding to the smallest entries in \mathbf{z}_i , the second eliminating the half corresponding to the largest entries in \mathbf{z}_i and the third eliminating the half corresponding to the largest absolute values. To these 3p initial candidates we add the LST estimate computed using the complete sample, obtaining a set of 3p + 1 elements. Once we have A_1 we obtain $\hat{\boldsymbol{\beta}}^{(1)}$ by minimizing $L(\boldsymbol{\beta})$ over the elements of A_1 .

Suppose now that we are on stage k. Let $0 < \alpha < 0.5$ be a trimming constant; in all our applications we set $\alpha = 0.05$. Then, for k > 1, we first delete the observations $(i = 1, \dots, n)$ such that $y_i > F_{\hat{\mu}_i}^{-1}(1 - \alpha/2)$ or $y_i < F_{\hat{\mu}_i}^{-1}(\alpha/2)$ where $\hat{\mu}_i = g^{-1}\left(\mathbf{x}_i^{\top}\hat{\boldsymbol{\beta}}^{(k-1)}\right)$ and, with the remaining observations, we re-compute the LST estimator $\hat{\boldsymbol{\beta}}_{\text{LTS}}^{(k)}$ and the principal sensitivity components. Let us remark that, for the computation of $\hat{\boldsymbol{\beta}}_{\text{LTS}}^{(k)}$ we have deleted the observations that have large residuals, since $\hat{\mu}_i$ is the fitted value obtained using $\hat{\boldsymbol{\beta}}^{(k-1)}$. In this way, while candidates on the first step of the iteration are protected from high leverage outliers, candidate $\hat{\boldsymbol{\beta}}_{\text{LTS}}^{(k)}$ is protected from low leverage outliers, which may not be extreme entries of the \mathbf{z}_i .

Now the set A_k will contain $\hat{\boldsymbol{\beta}}_{\text{LST}}^{(k)}$, $\hat{\boldsymbol{\beta}}^{(k-1)}$ and the 3p LST estimates computed deleting extreme values according to the principal sensitivity components as in the first iteration. $\hat{\boldsymbol{\beta}}^{(k)}$ is the element of A_k minimizing $L(\boldsymbol{\beta})$. The iterations will continue until $\hat{\boldsymbol{\beta}}^{(k)} \approx \hat{\boldsymbol{\beta}}^{(k-1)}$. Let $\hat{\boldsymbol{\beta}}_1$ be the final

estimate obtained at this stage.

Stage 2. We first delete the observations y_i $(i = 1, \dots, n)$ such that $y_i > 1$ $F_{\hat{\boldsymbol{\mu}}_i}^{-1}(1-\alpha/2)$ or $y_i < F_{\hat{\boldsymbol{\mu}}_i}^{-1}(\alpha/2)$, where $\hat{\boldsymbol{\mu}}_i = g^{-1}\left(\mathbf{x}_i^{\top}\hat{\boldsymbol{\beta}}_1\right)$ and compute the LST estimate $\hat{\boldsymbol{\beta}}^{(*)}$ with the reduced sample. Then for each of the deleted observations we check whether $y_i > F_{\hat{\boldsymbol{\mu}}_i}^{-1}(1 - \alpha/2)$ or $y_i < F_{\hat{\boldsymbol{\mu}}_i}^{-1}(\alpha/2)$, where $\hat{\boldsymbol{\mu}}_i = g^{-1} \left(\mathbf{x}_i^{\top} \hat{\boldsymbol{\beta}}^{(*)} \right)$. Observations which are not within these bounds are finally eliminated and those which are, are restored to the sample. With the resulting set of observations we compute the LST estimate β_2 which is our proposal as a starting value for solving the estimating equations of the MT-estimates.

Monte Carlo Study 4

In this section we report the results of a Monte Carlo study in which we compare the MT-estimator computed with the proposed initial estimate (FMT), to the robust quasi likelihood estimator (RQL) proposed by Cantoni and Ronchetti [2001], the Conditionally Unbiased Bounded Influence (CUBIF) estimator proposed by Künsch et al. [1989], and the MT-estimate beginning at an initial estimator computing by subsampling (SMT). For computing the RQL estimator, we used function glmrob from the R package robustbase with method "Mqle" and argument weights.on.x set to "robCov" so that weights based on robust Mahalanobis distance of the design matrix (intercept excluded) are used to downweight potential outliers in x-space. The Conditionally Unbiased Bounded Influence (CUBIF) estimators proposed by Künsch et al. [1989] was computed using an implementation kindly provided by Prof. Alfio Marazzi (personal communication); the implementation available in function glmrob from the R package robustbase with method "cubif" perform substantially bad and it is not reported here. For the computation of the SMT estimator, the number of subsamples was set to 2500. Both FMT and SMT are computed using an iteratively reweighted least squares method described in the appendix. They only differ in the starting point. We study

the case of Poisson regression and log link.

Let $\mathbf{x} = (1, \mathbf{x}^*)$ be a random vector in \mathbb{R}^p such that \mathbf{x}^* has distribution $\mathcal{N}_{p-1}(\mathbf{0},\mathbf{I})$ and let y be a random variable such that $y|\mathbf{x} \sim \mathcal{P}\left(\exp(\boldsymbol{\beta}_0^{\top}\mathbf{x})\right)$. We consider p = 100 and three different models. In model 1 data are generated with $\boldsymbol{\beta}_0 = \mathbf{e}_2$, in model 2 $\boldsymbol{\beta}_0 = 2\mathbf{e}_1 + \mathbf{e}_2$ and in model 3 $\boldsymbol{\beta}_0 = 2\mathbf{e}_1 + 1.5\mathbf{e}_2$, where \mathbf{e}_i is the vector of \mathbb{R}^p with all entries equal to zero except for the *i*-th entry which is equal to one. For each of these models we simulate the case in which the samples do not contain outliers and the case in which the samples have 10 per cent of identical outliers at point (\mathbf{x}_0, y_0) . The outliers are located at $\mathbf{x}_0 = \mathbf{e}_1 + 3\mathbf{e}_2$. The values of y_0 are taken in a grid ranging from $\boldsymbol{\mu}_0 - K_1$ to $\boldsymbol{\mu}_0 + K_2$ where $\boldsymbol{\mu}_0 = \exp(\boldsymbol{\beta}_0^{\top} \mathbf{x}) = \mathbb{E}_{\boldsymbol{\beta}_0}(y | \mathbf{x} = \mathbf{x}_0)$. The values K_1 and K_2 and the grid step are chosen so that the maximum mean squared error of our proposed estimator can be identified. For model 1 we also consider high leverage outliers with $\mathbf{x}_{0,2} = \mathbf{e}_1 + 3\mathbf{e}_2 + 4\mathbf{e}_3$. We have considered samples of size n = 400 an n = 1000, but since the behaviour is similar we report the results only for the larger sample size. Given an estimator β , we denote by MSE, the mean squared error defined by $\mathbb{E}_{\beta_0}(||\hat{\boldsymbol{\beta}} - \boldsymbol{\beta}_0||^2)$, where $||\cdot||$ denotes the L_2 norm. We estimate the MSE by

$$\widehat{\mathrm{MSE}} = \frac{1}{N} \sum_{j=1}^{N} ||\hat{\boldsymbol{\beta}}_j - \boldsymbol{\beta}_0||^2,$$

where $\hat{\boldsymbol{\beta}}_{j}$ is the value of the estimator at the *j*-th replication and N is the number of replications which was chosen equal to 1000.

Our simulations show that the proposed estimator has smaller MSE than all other proposals for almost all the contaminations considered. CUBIF estimator has a smaller MSE for some values of y_0 but, since it is based on a monotone score function, its MSE increases as y_0 increases. On the other hand, the MSE of FMT estimator is bounded; we observe that it decreases as y_0 increases beyond a certain value. To see this we study the MSE as a function of y_0 and consider, as a measure of robustness, the maximum MSE for $y_0 \in \mathbb{Z}_{\geq 0}$. The proposed estimator has the smallest maximum MSE for all the models considered. In Figures 1 to 3 we plot the MSE as a function of y_0 for samples of size n = 1000 with 10% contamination level.

In Figure 5 we report the execution time for the different methods. This figure shows that our proposed method is a great improvement over the subsampling method, as far as computational time is concerned.



Figure 1: MSE for model 1, p = 100, n = 1000 with 10% outliers at $\mathbf{x}_0 = 3\mathbf{e}_1 + \mathbf{e}_2$.



Figure 2: MSE for model 2, p = 100, n = 1000 with 10% outliers at $\mathbf{x}_0 = 3\mathbf{e}_1 + \mathbf{e}_2$.



Figure 3: MSE for model 3, p = 100, n = 1000 with 10% outliers at $\mathbf{x}_0 = 3\mathbf{e}_1 + \mathbf{e}_2$.



Figure 4: MSE for model 4, p = 100, n = 1000 with 10% outliers at $\mathbf{x}_0 = 3\mathbf{e}_1 + \mathbf{e}_2 + 4\mathbf{e}_4$.



Figure 5: Execution time in seconds for the four models, first row model 1 and 2, second row model 3 and 4, p = 100, n = 1000 with 10% outliers.

5 Conclusion

We introduce a deterministic robust initial estimate for generalized linear models. This initial estimate is used in an iteratively reweighted least squares algorithm to obtain a solution of a transformed M-estimator. We illustrate the procedures for the Poisson model. Monte Carlo experiments show that the performance of MT-estimators computed with the proposed initial estimator have a small bounded mean squared error exhibiting a redescending behavior. This is not the case for other proposals such RQL, CUBIF and MT with initial estimator based on subsampling.

Computational details and algorithms

In this Appendix we describe the iteratively reweighted least squares algorithms that were used to compute the LST and the MT estimators.

Suppose that we have an initial estimator $\boldsymbol{\beta}_0$ and call $s(t) = m(g^{-1}(t))$, then using a Taylor expansion of order one we can approximate $m\left(g^{-1}\left(\mathbf{x}_i^{\top}\boldsymbol{\beta}\right)\right) = s\left(\mathbf{x}_i^{\top}\boldsymbol{\beta}\right)$ by

$$s(\mathbf{x}_i^{\top}\boldsymbol{\beta}_0) + s'(\mathbf{x}_i^{\top}\boldsymbol{\beta}_0)\mathbf{x}_i^{\top}(\boldsymbol{\beta} - \boldsymbol{\beta}_0).$$
(14)

Then an approximate value to the LST estimator can be found by the value β_1 that minimizes

$$\sum_{i=1}^{n} \left(t(y_i) - s(\mathbf{x}_i^{\top} \boldsymbol{\beta}_0) - s'(\mathbf{x}_i^{\top} \boldsymbol{\beta}_0) \mathbf{x}_i^{\top} (\boldsymbol{\beta} - \boldsymbol{\beta}_0) \right)^2$$

Therefore $\beta_1 - \beta_0$ is the LS estimator for a linear model with responses $t(y_1), \ldots, t(y_n)$ and regressor vectors $s'(\mathbf{x}_1^\top \boldsymbol{\beta}_0) \mathbf{x}_1, \ldots, s'(\mathbf{x}_n^\top \boldsymbol{\beta}_0) \mathbf{x}_n$ and consequently

$$\boldsymbol{\beta}_{1} = \boldsymbol{\beta}_{0} + \left(\mathbf{X}^{\top} \mathbf{W} (\boldsymbol{X} \boldsymbol{\beta}_{0})^{2} \mathbf{X} \right)^{-1} \mathbf{X}^{\top} \mathbf{W} (\mathbf{X} \boldsymbol{\beta}_{0}) (\mathbf{T} - \mathbf{s} (\mathbf{X} \boldsymbol{\beta}_{0})), \quad (15)$$

where **X** is the $n \times p$ matrix whose *i*-th row is \mathbf{x}_i^{\top} , $\mathbf{s}(\mathbf{X}\boldsymbol{\beta}) = (s(\mathbf{x}_1^{\top}\boldsymbol{\beta}), \dots, s(\mathbf{x}_n^{\top}\boldsymbol{\beta}))^{\top}$, $\mathbf{W}(\mathbf{X}\boldsymbol{\beta})$ is the diagonal matrix with diagonal elements $s'(\mathbf{x}_1^{\top}\boldsymbol{\beta}), \dots, s'(\mathbf{x}_n^{\top}\boldsymbol{\beta})$ and $\mathbf{T} = (t(y_1), \dots, t(y_n))^{\top}$.

An iterative procedure to compute the LST estimator can be obtained putting

$$\boldsymbol{\beta}_{k+1} = \boldsymbol{\beta}_k + \left(\mathbf{X}^{\top} \mathbf{W} (\boldsymbol{X} \boldsymbol{\beta}_k)^2 \mathbf{X} \right)^{-1} \mathbf{X}^{\top} \mathbf{W} (\mathbf{X} \boldsymbol{\beta}_k) (\mathbf{T} - \mathbf{s} (\mathbf{X} \boldsymbol{\beta}_k)), \quad (16)$$

and stopping when $\|\boldsymbol{\beta}_{k+1} - \boldsymbol{\beta}_{k+1}\| / \|\boldsymbol{\beta}_k\| \le \delta$, where δ is the error tolerance.

Suppose that β_k converges to β^* , then this value should satisfy the LST estimating equation. In fact, taking limit in both sides of (16) we get

$$\left(\mathbf{X}^{\top}\mathbf{W}(\boldsymbol{X}\boldsymbol{\beta}^{*})^{2}\mathbf{X}\right)^{-1}\mathbf{X}^{\top}\mathbf{W}(\mathbf{X}\boldsymbol{\beta}^{*})(\mathbf{T}-\mathbf{s}(\boldsymbol{\beta}^{*}))=\mathbf{0},$$

which is equivalent to

$$\mathbf{X}^{\top}\mathbf{W}(\mathbf{X}\boldsymbol{\beta}^{*})^{\top}(\mathbf{T}-\mathbf{s}(\boldsymbol{\beta}^{*}))=\mathbf{0},$$

and then β^* satisfies the estimating equation of the LST estimator.

To start the algorithm, it will be convenient to write equation (15) in the following slightly different way

$$\boldsymbol{\beta}_{1} = \left(\mathbf{X}^{\top}\mathbf{W}(\boldsymbol{X}\boldsymbol{\beta}_{0})^{2}\mathbf{X}\right)^{-1}\left(\mathbf{X}^{\top}\mathbf{W}(\boldsymbol{X}\boldsymbol{\beta}_{0})^{2}\mathbf{X}\boldsymbol{\beta}_{0} + \mathbf{X}^{\top}\mathbf{W}(\mathbf{X}\boldsymbol{\beta}_{0})(\mathbf{T} - \mathbf{s}(\mathbf{X}\boldsymbol{\beta}_{0}))\right).$$
(17)

Observe that according to (17) to compute β_1 we only need to give $\eta_0 = \mathbf{X}\boldsymbol{\beta}_0$. Then, since for Poisson regression and log link it holds $\mathbf{x}_i^{\mathsf{T}}\boldsymbol{\beta} = \log(E(y_i))$, it seems reasonable to take $\eta_0 = (\log(y_1+0.1), \ldots, \log(y_n+0.1))^{\mathsf{T}}$. The value 0.1 is added to avoid numerical problem when $y_i = 0$. To compute the estimators $\hat{\boldsymbol{\beta}}_{(j)}$ only one iteration is performed. The reason is that for these auxiliary estimators the accuracy is not as important as the speed at which they can be computed. Our experiments show that there is no noticeable loss in the precision of the final estimate by doing this but, on the other hand, the computation times decrease significantly.

We describe now an analogous iterative algorithm for computing the MT estimator. Suppose that we have an initial robust estimator $\boldsymbol{\beta}_0$. We compute a new value using two approximations. As in the case of the LST estimator, replacing, in (4), $m\left(g^{-1}\left(\mathbf{x}_i^{\top}\boldsymbol{\beta}\right)\right)$ by (14) we consider the approximate loss function

$$\sum_{i=1}^{n} \rho\left(t(y_i) - s(\mathbf{x}_i^{\top} \boldsymbol{\beta}_0) - s'(\mathbf{x}_i^{\top} \boldsymbol{\beta}_0) \mathbf{x}_i^{\top} (\boldsymbol{\beta} - \boldsymbol{\beta}_0)\right)$$

Differentiating with respect to β we obtain the estimating equation

$$\sum_{i=1}^{n} \psi \left(t(y_i) - s(\mathbf{x}_i^{\mathsf{T}} \boldsymbol{\beta}_0) - s'(\mathbf{x}_i^{\mathsf{T}} \boldsymbol{\beta}_0) \mathbf{x}_i^{\mathsf{T}} (\boldsymbol{\beta} - \boldsymbol{\beta}_0) \right) s'(\mathbf{x}_i^{\mathsf{T}} \boldsymbol{\beta}_0) \mathbf{x}_i = \mathbf{0}, \quad (18)$$

where $\psi = \rho'$. Note that this equation can be written as

$$\sum_{i=1}^{n} \left(t(y_i) - s(\mathbf{x}_i^{\top} \boldsymbol{\beta}_0) - s'(\mathbf{x}_i^{\top} \boldsymbol{\beta}_0) \mathbf{x}_i^{\top} (\boldsymbol{\beta} - \boldsymbol{\beta}_0) \right) w(\mathbf{x}_i^{\top} \boldsymbol{\beta}, \mathbf{x}_i^{\top} \boldsymbol{\beta}_0) s'(\mathbf{x}_i^{\top} \boldsymbol{\beta}_0) \mathbf{x}_i,$$
(19)

where

$$w(u,v) = \frac{\psi(t(y_i) - s(v) - s'(v)(u - v))}{t(y_i) - s(v) - s'(v)(u - v)}.$$

Since $\boldsymbol{\beta}$ should be close to $\boldsymbol{\beta}_0$, the second approximation is to replace, in (19), $w(\mathbf{x}_i^{\top}\boldsymbol{\beta}, \mathbf{x}_i^{\top}\boldsymbol{\beta}_0)$ by $w^*(\mathbf{x}_i^{\top}\boldsymbol{\beta}_0) = w(\mathbf{x}_i^{\top}\boldsymbol{\beta}_0, \mathbf{x}_i^{\top}\boldsymbol{\beta}_0)$. Then $\boldsymbol{\beta}_1$ is defined as the solution of the approximate estimating equation

$$\sum_{i=1}^{n} \left(t(y_i) - s(\mathbf{x}_i^{\top} \boldsymbol{\beta}_0) - s'(\mathbf{x}_i^{\top} \boldsymbol{\beta}_0) \mathbf{x}_i^{\top} (\boldsymbol{\beta} - \boldsymbol{\beta}_0) \right) w^*(\mathbf{x}_i^{\top} \boldsymbol{\beta}_0) s'(\mathbf{x}_i^{\top} \boldsymbol{\beta}_0) \mathbf{x}_i$$

and is given by

$$\boldsymbol{\beta}_1 = \boldsymbol{\beta}_0 + \left(\mathbf{X}^\top \mathbf{W}^2 (\boldsymbol{X} \boldsymbol{\beta}_0)^\top \mathbf{W}^* (\boldsymbol{X} \boldsymbol{\beta}_0) \mathbf{X} \right)^{-1} \mathbf{X}^\top \mathbf{W} (\mathbf{X} \boldsymbol{\beta}_0) \mathbf{W}^* (\boldsymbol{X} \boldsymbol{\beta}_0) (\mathbf{T} - \mathbf{s} (\mathbf{X} \boldsymbol{\beta}_0)),$$

where $\mathbf{W}^*(\mathbf{X}\boldsymbol{\beta})$ is the $n \times n$ diagonal matrix with diagonal elements $w^*(\mathbf{x}_1^\top \boldsymbol{\beta}), \ldots, w^*(\mathbf{x}_n^\top \boldsymbol{\beta})$. Then, the iterative procedure to compute the MT estimator is given by

$$\boldsymbol{\beta}_{k+1} = \boldsymbol{\beta}_k + \left(\mathbf{X}^\top \mathbf{W}^2 (\boldsymbol{X} \boldsymbol{\beta}_k)^\top \mathbf{W}^* (\boldsymbol{X} \boldsymbol{\beta}_k) \mathbf{X} \right)^{-1} \mathbf{X}^\top \mathbf{W} (\mathbf{X} \boldsymbol{\beta}_k) \mathbf{W}^* (\boldsymbol{X} \boldsymbol{\beta}_k) (\mathbf{T} - \mathbf{s} (\mathbf{X} \boldsymbol{\beta}_k))$$
(20)

Suppose that $\beta_k \to \beta^*$, then taking limits in both sides of (20), we get

$$\mathbf{X}^{\top}\mathbf{W}(\mathbf{X}\boldsymbol{\beta}^{*})\mathbf{W}^{*}(\boldsymbol{X}\boldsymbol{\beta}^{*})(\mathbf{T}-\mathbf{s}(\mathbf{X}\boldsymbol{\beta}^{*}))=\mathbf{0},$$

and this is equivalent to

$$\mathbf{X}^{\top}\mathbf{W}(\mathbf{X}\boldsymbol{\beta}^{*})\boldsymbol{\Psi}\ (\boldsymbol{X}\boldsymbol{\beta}^{*}) = \mathbf{0},$$
(21)

where $\Psi(\boldsymbol{X}\boldsymbol{\beta}) = (\psi(t(y_1) - s(\mathbf{x}_i^{\top}\boldsymbol{\beta})), \dots, \psi(t(y_n) - s(\mathbf{x}_i^{\top}\boldsymbol{\beta}))^{\top}$. Then $\boldsymbol{\beta}^*$ satisfies the estimating equation of the MT estimator.

References

F. Alqallaf and C. Agostinelli. Robust inference in generalized linear models. Communications in Statistics - Simulation and Computation, 45(9):3053– 3073, 2016. doi: 10.1080/03610918.2014.911896.

- A. Bergesio and V.J. Yohai. Projection estimators for generalized linear models. Journal of the American Statistical Association, 106:661–671, 2011.
- A.M. Bianco, G. Boente, and I.M. Rodrigues. Resistant estimators in poisson and gamma models with missing responses and an application to outlier detection. *Journal of Multivariate Analysis*, 114:209–226, 2013.
- E. Cantoni and E. Ronchetti. Robust inference for generalized linear models. Journal of the American Statistical Association, 96:1022–1030, 2001.
- R.D. Cook. Detection of influential observation in linear regression. *Techno*metrics, 19(1):15–18, 1977. doi: 10.2307/1268249.
- H. Künsch, L. Stefanski, and R. Carroll. Conditionally unbiased boundedinfluence estimation in general regression models, with applications to generalized linear models. *Journal of the American Statistical Association*, 84: 460–466, 1989.
- P. McCullagh and J.A. Nelder. *Generalized Linear Models*. Chapman and Hall/CRC, second edition, 1989.
- D. Peña and V.J. Yohai. A fast procedure for outlier diagnostics in large regression problems. *Journal of the American Statistical Association*, 94: 434–445, 1999.
- P.J. Rousseeuw and A.M. Leroy. Robust regression and outlier detection. Wiley and Sons, 1987.
- M. Valdora and V.J. Yohai. Robust estimators for generalized linear models. Journal of Statistical Planning and Inference, 146:31–48, 2014.