

Continuum Mechanics and Thermodynamics in the Hamilton and the Godunov-type Formulations

ILYA PESHKOV,¹ MICHAL PAVELKA,² EVGENIY ROMENSKI,³ MIROSLAV GRMELA⁴

Spoiler

Continuum mechanics with dislocations, with the Cattaneo type heat conduction, with mass transfer, and with electromagnetic fields is put into the Hamiltonian form and into the form of the Godunov type system of the first order, symmetric hyperbolic partial differential equations (SHTC equations). The compatibility with thermodynamics of the time reversible part of the governing equations is mathematically expressed in the former formulation as degeneracy of the Hamiltonian structure and in the latter formulation as the existence of a companion conservation law. In both formulations the time irreversible part represents gradient dynamics. The Godunov type formulation brings the mathematical rigor (the well-posedness of the Cauchy initial value problem) and the possibility to discretize while keeping the physical content of the governing equations (the Godunov finite volume discretization).

CONTENTS

1	Introduction	3
2	Preliminaries	4
2.1	SHTC framework, Symmetric Hyperbolic Equations, Well-Posedness, Causality	4
2.2	GENERIC	6
3	SHTC formulation of continuum mechanics	8
3.1	Lagrangian frame	8
3.1.1	Energy formulation	9
3.1.2	Conjugate formulation and symmetric form	10
3.1.3	Complimentary structure and odd-even parity	11
3.1.4	Involution constraints	11
3.1.5	Variational nature of the SHTC equations	11
3.2	Eulerian frame	13
3.2.1	Energy formulation	14
3.2.2	Involution constraints	15
3.2.3	Weakly non-local conservative non-symmetrizable formulation	17
3.2.4	Conjugate formulation	18
3.2.5	Symmetric form of conjugate formulation for curl-free and divergence-free involution constraints	20
3.2.6	Symmetric form of conjugate weakly non-local formulation in general case	21

¹Institut de Mathématiques de Toulouse, Toulouse, France and Sobolev Institute of Mathematics, Novosibirsk, Russia, [e-mail](#)

²Mathematical Institute, Faculty of Mathematics and Physics, Charles University, Sokolovská 83, 186 75 Prague, Czech Republic, [e-mail](#)

³Sobolev Institute of Mathematics and Novosibirsk State University, Novosibirsk, Russia, [e-mail](#)

⁴École Polytechnique de Montréal, C.P.6079 suc. Centre-ville, Montréal, H3C 3A7, Québec, Canada, [e-mail](#)

4	GENERIC form of SHTC equations	23
4.1	Lagrangian SHTC system as Hamiltonian evolution	23
4.2	Hydrodynamics of ideal fluids	24
4.3	Distortion matrix	25
4.3.1	Dynamics of labels	25
4.3.2	Differential geometric derivation	27
4.3.3	Dynamics of general distortion matrix	28
4.3.4	Dynamics of the left Cauchy-Green (or Finger) tensor	29
4.4	Electrodynamics of slowly moving medium	30
4.5	Ballistic transport	32
4.5.1	The SHTC-compatible bracket	32
4.5.2	Alternative bracket, Cattaneo hydrodynamics	34
4.6	Complete Poisson bracket	35
5	Dissipation	35
5.1	Irreversible dynamics	36
5.1.1	Energy representation	36
5.1.2	Entropy representation and gradient dynamics	38
5.2	Involution constraints in case of dissipation	39
5.3	Dispersive dynamics	40
6	Examples	40
6.1	MaxEnt reduction	40
6.2	Navier-Stokes dynamics	41
6.3	Elastic and elastoplastic solids	42
6.4	Non-Newtonian fluid dynamics	43
6.5	Heat and mass transfer	44
6.5.1	Heat conduction	44
6.5.2	Mass transfer	45
6.6	Electrodynamics of slowly moving medium	47
7	Concluding remarks	47
A	Lagrange-to-Euler transformation	48
A.1	Auxiliary relations	48
A.2	(w_i, σ) -pair	49
A.2.1	Transformation (38f)→(18f)	49
A.2.2	Transformation (38e)→(18e):	50
A.3	(e, h) -pair, transformation of the nonlinear Maxwell equations	51
A.4	(m, F) -pair	53
A.4.1	Distortion matrix	53
A.4.2	Transformation of the momentum conservation	53
B	Derivation of the energy conservation law	54
C	Derivation of the time evolution for $\nabla \times w$ and $\nabla \times A$	55

1. INTRODUCTION

Results of experimental observations are seen in mathematical models as properties of solutions of their governing equations. Universality of some results translates in the models into the universality of the mathematical structure of the equations. For instance one of the universal experimental observations is the approach of externally unforced macroscopic systems to the thermodynamic equilibrium states at which the behavior is found to be well described by the classical equilibrium thermodynamics. The structure responsible for this observation has been identified to be the following. The irreversible part of the time evolution is of gradient type generated by a potential called entropy. This potential is then preserved in the reversible part of the time evolution that is Hamiltonian, since it is the part of the time evolution directly inherited from the underlying microscopic mechanics. The time-evolution equations involving both the Hamiltonian and the gradient part have appeared first in [20], in [44] (that was presented at the AMS-IMS-SIAM Joint Summer Research Conference in the Mathematical Sciences on Fluids and Plasmas “Geometry and Dynamics”, held at the University of Colorado, Boulder, CO, USA, 17–23 July 1983) and in [59, 67, 45, 7, 50, 72, 71]. In [50, 72], the abstract equation (14) has been called GENERIC.

In the smaller pool of governing equations that arise in continuum mechanics we may expect to identify even stronger common structure. In particular, what is missing in GENERIC is the guarantee that the initial value problem is well posed. This is indeed one of the basic validations of a model. The time evolution is seen to exist and thus the time evolution seen in the model should also exist. Another aspect that is not included in GENERIC is the adaptability of the governing equation to searching their solutions numerically.

The structure addressing both of these aspects have been identified by Godunov [35, 36] for a system of the first order partial differential equations having the form of local conservation laws (i.e. equations having the form of the time derivative of a field equals divergence of a flux). The compatibility with thermodynamics of these equations (representing the time reversible part of the time evolution) is expressed by the existence of a companion local conservation law for a new field that is an appropriately regular and convex function of the fields playing the role of state variables. It is very interesting that this requirement of the agreement with experimental observations implies then directly the mathematical well-posedness (the companion local conservation law enables to symmetrize the system of the first order hyperbolic equations which then implies their well posedness). Godunov has also introduced a discretization (the modification of the governing equations needed when computers are brought to assistance in the process of finding details of their solutions) preserving the physical content of the original partial differential equations [34].

In this paper we introduce a large family of partial differential equations that:

1. Govern the time evolution of fluids, solids without and with dislocations, the Cattaneo type heat conduction and the electromagnetic fields, all coupled together.
2. Possess the GENERIC structure.
3. Possess the SHTC structure, that extends to Godunov structure to a larger class of the first order partial differential equations.

The GENERIC route begins with the well known Hamiltonian structure of the Euler hydrodynamic equations [16, 2]. By adopting the field of labels into the set of state variables, we pass to the Lagrangian formulation and to the solid mechanics without dislocations [46] and (by using the extension developed in this paper) with dislocations. This Hamiltonian formulation of the reversible part of the continuum mechanics is then coupled to the Hamiltonian formulation of the Cattaneo-like heat conduction and to the Hamiltonian formulation of the time evolution of electromagnetic fields.

The Godunov route begins with the Lagrangian formulation in which the time reversible governing equations have the form of the local conservation laws and the original Godunov analysis thus applies to them. By passing to the Eulerian formulation (the Lagrange \rightarrow Euler passage is worked out in detail in this paper) the governing equations do not remain to be, at least in general, local conservation laws. Their weakly nonlocal extension is

however shown to possess a new structure (called SHTC structure) that also allows symmetrization, and thus the mathematical well posedness, as well as the applicability of the Godunov discretization, is guaranteed for them. The weakly nonlocal extension (having the physical meaning, for example, of including dislocations into the consideration) consists of:

1. Constructing new local conservation laws for new fields that are appropriate spatial derivatives of the original fields included in the set of the state variables (e.g. the vorticity) by appropriately combining the original system of equations with equations obtained by applying appropriate spatial derivatives on the original system of equations.
2. Adopting the new fields (that are appropriate spatial derivatives of the fields included in the original set of the state variables) as new independent fields.

Both the GENERIC and the Godunov routes lead to the same equations. Having proven that the equations possess both the GENERIC and the SHTC structures, all the results and techniques that are available for these structures (in particular the physical, the mathematical, and the numerical well posedness) are now available for the governing equations investigated in this paper.

2. PRELIMINARIES

2.1. SHTC framework, Symmetric Hyperbolic Equations, Well-Posedness, Causality

The origin of the SHTC formulation of continuum mechanics should be attributed to the work of Godunov [35] where he considered “an interesting class” of nonlinear conservation laws. The key motivation (from the words of S.K. Godunov) for this seminal paper was the will to understand what physical principles may guarantee the *well-posedness* of the initial value problem (IVP) for a *nonlinear* system of time dependent PDEs. Indeed, when one deals with nonlinear dynamical phenomena, and in particular with nonlinear time-dependent partial differential equations (PDEs), perhaps, the first examination a new model has to pass is to verify if the IVP is well-posed, at least locally, i.e. whether the solution to the system of PDEs with given initial data exists, is unique and stable (depends continuously on the initial data). We emphasize that the well-posedness of the IVP should not be considered as a purely mathematical requirement but as a *fundamental physical observation* about the time evolution of physical systems, i.e. exactly as we consider causality, conservation and thermodynamic principles, Galilean or Lorentz invariance to be essential features of macroscopic time evolution. In other words, a model describing the time evolution of a physical system and having an ill-posed IVP should be regarded as wrong. Moreover, the well-posedness of the IVP is a fundamental property of time-dependent PDEs in order for them to be numerically solvable.

Not all nonlinear systems of PDEs, of course, are well posed. This even cannot be guaranteed for a first order quasi-linear system [5, 19, 69], or for models which were believed to be consistently derived from microscopic theories as, for example, the Burnett equations derived from the gas kinetic theory [8, 90, 91]. However, there is a class of nonlinear PDEs for which the IVP is locally well-posed in time, by definition, which is the class of hyperbolic PDEs. Unfortunately, it is difficult to prove that a given non-linear first order system is hyperbolic in the domain of interest (domain of state parameters) because this would require to prove the existence of the full basis of eigenvectors for a matrix whose entries depends nonlinearly on the state variables. So how can one guaranty that the IVP for a new nonlinear continuum mechanics model is well-posed, i.e. that the model is hyperbolic?

The only possibility we see is to deliberately develop the model within a very important *subclass* of first order hyperbolic systems

$$A(\mathbf{p}) \frac{\partial \mathbf{p}}{\partial t} + B_k(\mathbf{p}) \frac{\partial \mathbf{p}}{\partial x_k} = 0, \quad (1)$$

where $A^T = A > 0$ and $B_k^T = B_k$, for which (local) well-posedness is known to hold true [58, 5, 69, 88]. This subclass is called *symmetric hyperbolic systems of PDEs* which is a generalization of Friedrichs-symmetrizable linear systems [27] and is known to be globally hyperbolic by definition. However, one should naturally question how restrictive it is for a model to be symmetric hyperbolic and, at the same time, represent diverse physical phenomena admitting continuum mechanics description? As it was shown by Godunov [35, 36, 37] and later by others [28, 9, 89] there is an intimate connection between the symmetric hyperbolicity and thermodynamics for a specific subclass of *system of conservation laws*.

In particular, if a first order system of conservation laws for conservative state variables $\mathbf{q} = (q_1, q_2, \dots, q_n)$

$$\frac{\partial \mathbf{q}}{\partial t} + \frac{\partial \mathbf{F}^k(\mathbf{q})}{\partial x_k} = 0 \quad (2)$$

admits an extra conservation law

$$\frac{\partial E(\mathbf{q})}{\partial t} + \frac{\partial G^k(\mathbf{q})}{\partial x_k} = 0 \quad (3)$$

for a *strictly convex potential* $E(\mathbf{q})$, which plays the role of the total energy for the system, then such a system can be parametrized with new state variables $\mathbf{p} = (p_1, p_2, \dots, p_n)$ and a potential $L(\mathbf{p})$ as follows (which shall be referred to as the *Godunov form* of conservation laws)

$$\frac{\partial L_{\mathbf{p}}}{\partial t} + \frac{\partial L_{\mathbf{p}}^k}{\partial x_k} = 0, \quad (4)$$

where, as through out this paper, $L_{\mathbf{p}}$ and $L_{\mathbf{p}}^k$ are the vectors with components $L_{p_i} = \partial L / \partial p_i$, $L_{p_i}^k = \partial L^k / \partial p_i$, see below for the concrete form of these vectors for the conservative system. It is obvious that (4) can be rewritten in a symmetric quasilinear form

$$L_{p_i p_j} \frac{\partial p_j}{\partial t} + L_{p_i p_j}^k \frac{\partial p_j}{\partial x_k} = 0. \quad (5)$$

Moreover, the conservative state variables \mathbf{q} and the potential $E(\mathbf{q})$ are related to the new variables \mathbf{p} and potential $L(\mathbf{p})$ as

$$\mathbf{q} = L_{\mathbf{p}}, \quad E(\mathbf{q}) = \mathbf{p} \cdot L_{\mathbf{p}} - L(\mathbf{p}) = \mathbf{p} \cdot \mathbf{q} - L(\mathbf{p}) \quad (6)$$

and vice versa,

$$\mathbf{p} = E_{\mathbf{q}}, \quad L(\mathbf{p}) = \mathbf{q} \cdot E_{\mathbf{q}} - E(\mathbf{q}) = \mathbf{q} \cdot \mathbf{p} - E(\mathbf{q}). \quad (7)$$

The relations $\mathbf{p} = E_{\mathbf{q}}$ and $\mathbf{q} = L_{\mathbf{p}}$ are one-to-one relations because we assume that $E(\mathbf{q})$ is convex and hence, the Hessian $\frac{\partial \mathbf{p}}{\partial \mathbf{q}} = E_{\mathbf{q}\mathbf{q}} = (L_{\mathbf{p}\mathbf{p}})^{-1} > 0$. Thus, the convexity of $E(\mathbf{q})$, in fact, means that (5) is symmetric hyperbolic. This result is referred to as the Godunov-Friedrichs-Lax (GFL) theorem [29].

The key point of the Godunov observation is that the system (2) and conservation law (3) together forms an *overdetermined* system of PDEs, i.e. the number of PDEs is $n + 1$ while the number of state variables is n (E is not an unknown but the potential $E(\mathbf{q})$), and thus, in order such a system has a solution, one of the conservation laws should be a consequence of the others. Therefore, the simultaneous validity of (2) and (3) implies that

$$E_{\mathbf{q}} \cdot \left(\frac{\partial \mathbf{q}}{\partial t} + \frac{\partial \mathbf{F}^k(\mathbf{q})}{\partial x_k} \right) \equiv \frac{\partial E(\mathbf{q})}{\partial t} + \frac{\partial G^k(\mathbf{q})}{\partial x_k}, \quad (8)$$

where the dot denotes the scalar product. After introducing the potentials $L^k(\mathbf{p}) = E_{\mathbf{q}} \cdot \mathbf{F}^k(\mathbf{q}) - G^k(\mathbf{q}) = \mathbf{p} \cdot \mathbf{F}^k(\mathbf{q}) - G^k(\mathbf{q})$ and noting that $L_{p_i}^k = \mathbf{F}^k$, the energy conservation (3) can be rewritten as

$$\frac{\partial (p_i L_{p_i} - L)}{\partial t} + \frac{\partial (p_i L_{p_i}^k - L^k)}{\partial x_k} = 0. \quad (9)$$

In other words, L is the Legendre-conjugate energy and L^k is Legendre conjugation of energy flux. Therefore, returning back to our last question, such a subclass of symmetric hyperbolic PDEs can be associated with the *thermodynamically compatible* systems of first order nonlinear conservation laws.

However, it turns out that there are some important continuum mechanics models, like ideal magnetohydrodynamics (MHD) equations for example, which *seems* to be represented by systems of conservation laws (2) with an additional conservation law (3) but still can not be symmetrizable in such a way. A more attentive analysis by Godunov of this inconsistency of the MHD equations in [38] shows that, in fact, the MHD equations written in a classical conservative form are not compatible with the energy conservation, i.e. the energy conservation law is not the consequence of the mass, momentum, entropy and magnetic field conservation laws, like in (8). More precisely, the energy flux of the MHD equation is inconsistent with the other fluxes unless the *non-conservative* product $B_i v_i \frac{\partial B_k}{\partial x_k} \equiv 0$ is added to the left-hand side of (8), where v_i and B_i are the velocity and magnetic fields accordingly. We discuss this in more details in Section 3.2.3. Godunov concludes that the ideal MHD equations do not admit the original Godunov structure (4) but generalize it in the following way

$$\frac{\partial L_{\mathbf{p}}}{\partial t} + \frac{\partial (v_k L)_{\mathbf{p}}}{\partial x_k} + C_k \frac{\partial \mathbf{p}}{\partial x_k} = 0, \quad (10)$$

where $C_k^T = C_k$ are symmetric matrices and v_k being the velocity field.

The latter analysis by Godunov and Romenski in nineties in a series of papers [32, 43, 30, 42, 42, 87, 21, 41] of various equations (ideal MHD equations, nonlinear elasticity, electrodynamics of slowly moving medium, superfluid flow, superconductivity) studied in the books of Landau and Lifshitz [62, 61] reveals that for a proper understanding of the *structure* of the continuum mechanics models it is important to distinguish the Lagrangian and Eulerian frames of reference, and that the original Godunov structure (4) is, in fact, rather inherent to the conservation laws written in the Lagrangian frame, while their Eulerian counterparts are *inherently non-conservative* time evolutions and have more complicated structure (10).

The departure point of the SHTC framework is the invariant under arbitrary rotations conservation laws in the Lagrangian frame, which were proposed in [30] and can be derived from Hamilton's principle of stationary action, see Section 3.1.5. Here, we also demonstrate for the first time the the Lagrangian SHTC equations can be generated by *canonical Poisson brackets*. Then, Eulerian non-conservative SHTC equations (10) are the direct consequence of the Lagrange-to-Euler transformation applied to the Lagrangian conservation laws. It is the primary goal of this paper to demonstrate that the Eulerian non-conservative SHTC equations (10) are fully compatible with the GENERIC framework presented in the subsequent section. Another goal of the paper is to provide the reader with some non-trivial details of the Lagrange-to-Euler transformation between the conservative Lagrangian SHTC equations and their non-conservative Eulerian counterparts which have never been published before. It is thus one of the goal of our paper to provide these important, though quite technical, details. Moreover, we review all properties of the SHTC formulation which were previously published rather separately in a series of paper by Godunov and Romenski [32, 43, 30, 42, 42, 87, 21, 41].

Eventually, we also recall that apart of the well-posedness of the IVP the hyperbolicity also naturally accounts for another fundamental observation about time evolution of physical systems, namely the finite velocity for any perturbation propagation, i.e. *causality*.

2.2. GENERIC

General Equation for Non-Equilibrium Reversible-Irreversible Coupling (GENERIC) was first introduced in [50, 72]. The fundamental idea behind GENERIC is that evolution equations for chosen state variables are generated from four building blocks: a Poisson bracket $\{\bullet, \bullet\}$, Hamiltonian $\mathcal{E} = \int E dr$, dissipation potential Ξ and entropy functional $\mathcal{S} = \int s dr$, where the potentials E and s are the total energy and entropy densities. Once these building blocks are at hand, the evolution equations follow by direct calculation.

The building blocks are required to fulfill certain degeneracies and conditions:

1. Entropy should be a Casimir of the Poisson bracket, which means that $\{\mathcal{A}, \mathcal{S}\} = 0$ for all functionals \mathcal{A} of the state variables.
2. Energy is not changed by the dissipation potential¹, i.e. $\langle \mathcal{E}_x, \Xi_{\mathcal{S}_x} \rangle = 0$, and
3. the dissipation potential is a monotonous operator, i.e. $\langle \Xi_{\mathcal{S}_x}, \mathcal{S}_x \rangle = 0$ for all entropies.²

Apart from these conditions, the Poisson bracket has the following properties:

1. antisymmetry, $\{\mathcal{A}, \mathcal{B}\} = -\{\mathcal{B}, \mathcal{A}\}$,
2. Leibniz rule, $\{\mathcal{A}\mathcal{B}, \mathcal{C}\} = \{\mathcal{A}, \mathcal{C}\}\mathcal{B} + \mathcal{A}\{\mathcal{B}, \mathcal{C}\}$,
3. Jacobi identity, $\{\{\mathcal{A}, \mathcal{B}\}, \mathcal{C}\} + \{\{\mathcal{B}, \mathcal{C}\}, \mathcal{A}\} + \{\{\mathcal{C}, \mathcal{A}\}, \mathcal{B}\} = 0$.

Moreover, the Poisson bracket is constructed from a Poisson bivector L^{ij} (antisymmetric twice contravariant tensor field) as

$$\{\mathcal{A}, \mathcal{B}\} = \mathcal{A}_{x^i} L^{ij} \mathcal{B}_{x^j}. \quad (11)$$

Note that summation over an index can be interpreted as integral over fields. Finally, Hamiltonian evolution of functionals is given by

$$\frac{\partial \mathcal{A}}{\partial t} = \{\mathcal{A}, \mathcal{E}\}. \quad (12)$$

GENERIC then gives the following evolution equations for functionals of state variables x :

$$\frac{\partial \mathcal{A}}{\partial t} = \{\mathcal{A}, \mathcal{E}\} + \langle \mathcal{A}_x, \frac{\delta \Xi}{\delta \mathcal{S}_x} \rangle. \quad (13)$$

From antisymmetry of the Poisson bracket and from the degeneracy of the dissipation potential it follows that energy is conserved, $\partial \mathcal{E} / \partial t = 0$. From the degeneracy of the Poisson bracket and from monotonicity of the dissipation potential it follows that entropy grows, $\partial \mathcal{S} / \partial t \geq 0$. Moreover, when energy, entropy and the dissipation potential are taken as even functions with respect to time reversal transformation, the Hamiltonian part generates reversible evolution while the dissipative part generates irreversible evolution [74]. Finally, Onsager-Casimir reciprocal relations are satisfied automatically [74, 71]. These results manifest compatibility of GENERIC with thermodynamics.

Instead of working with evolution of functionals, which can be interpreted as a sort of weak formulation, one can recover the evolution equations of the state variables easily. Taking $\mathcal{A} = x^i$, we obtain

$$\frac{\partial x^i}{\partial t} = L^{ij} \mathcal{E}_{x^j} + \Xi_{\mathcal{S}_x^i}. \quad (14)$$

This last equation shows how to write down the evolution equations (usually partial differential equations) implied by GENERIC.

Since Poisson brackets are a cornerstone of GENERIC, a question often arises how the Poisson brackets can be derived from first principles. For example the canonical Poisson bracket, that generates Hamilton canonical equations in classical mechanics, is generated by the canonical two-form present on the cotangent bundle of classical mechanics (positions and momenta of all particles). The canonical Poisson bracket can be thus interpreted as a geometric result of construction of the cotangent bundle.

The hydrodynamic Poisson bracket, that generates compressible Euler equations, is different from the canonical Poisson bracket. Of course, it generates evolution equations for the hydrodynamic fields: density, momentum

¹ $\langle \bullet, \bullet \rangle$ denoting integration (L^2 scalar product)

²Note that subscripts denote functional derivatives $\mathcal{S}_x = \frac{\delta \mathcal{S}}{\delta x}$. In the special case when the functional is a spatial integral over a density which is a real-valued function of the state variables, the functional derivative becomes the usual derivative of the density with respect to the state variable. For example $\mathcal{E} = \int E(x(r)) dr$ and $\mathcal{E}_x = \frac{\partial E}{\partial x}$ if E is a real-valued function of real variables.

density and entropy density. It can be derived in several ways. One can start for example from dynamics of particles in the Lagrangian frame and perform transformation to the Eulerian frame [1]. The calculation is, however, rather complicated. A shorter way is to construct dual of the Lie algebra of the group of diffeomorphisms of a volume [64]. Perhaps the shortest way is to use projection from the Boltzmann Poisson bracket to the hydrodynamic fields, see e.g. [73].

It is, in fact, often more difficult to derive the dissipation potential. In the simplest cases dissipation potentials can be quadratic functions of derivatives of entropy. That is the setting of classical irreversible thermodynamics [51]. The matrix of second derivatives of the dissipation potential is then referred to as the dissipative matrix M^{ij} , and the evolution equations then become

$$\frac{\partial x^i}{\partial t} = L^{ij} \mathcal{E}_{x^j} + M^{ij} \mathcal{S}_{x^j} \quad \text{for} \quad \Xi = \frac{1}{2} \mathcal{S}_{x^i} M^{ij}(\mathbf{x}) \mathcal{S}_{x^j}. \quad (15)$$

Apart from quadratic dissipation potentials, exponential dissipation potentials (like hyperbolic cosine) are used in case of chemical reactions and Boltzmann equation [50, 47], and such dissipation potentials have also statistical reasoning [66].

In summary, the GENERIC framework is an alternative approach to non-equilibrium thermodynamics, which does not rely on classical balance laws and their extensions as usual non-equilibrium thermodynamics. The reversible (Hamiltonian) part of the evolution can be constructed by various methods, for example by means of differential geometry (Lie groups, Lie algebras, semidirect products) or projections. The irreversible part is constructed as gradient dynamics satisfying the first and second laws of thermodynamics. Applicability of GENERIC is very wide, covering for example particle mechanics, kinetic theory, motions of fluids and solids, complex fluids, electromagnetism and dynamics of interfaces.

3. SHTC FORMULATION OF CONTINUUM MECHANICS

In this Section, we present the SHTC formulation of continuum mechanics and discuss its properties in both Lagrangian and Eulerian frames. We first, present the Lagrangian SHTC equations. Exactly these equations are fully compatible with original Godunov's structure (4). The Lagrangian SHTC equations can be also seen as Hamiltonian dynamics generated by a canonical Poisson bracket of the cotangent bundle of two vector fields, see Section 4.1. We then proceed with presenting the Eulerian SHTC equations which are the equations of principle interest in this paper. The derivation of Poisson brackets for the Eulerian SHTC equations and thus demonstrating their fully compatibility with GENERIC in Section 4 and for which we show that they have a structure fully compatible with GENERIC in Section 4.

3.1. Lagrangian frame

In the series of papers [32, 43, 30, 42, 42, 87, 21, 41], Godunov and Romenski investigated the question of whether time evolution equations of various continuum mechanics models share some common elements of a general structure. More specifically, in [30], they examined the following important question. Whether it is possible to conclude something meaningful about the structure of a PDE system representing a continuum mechanics model without specifying the physical settings but being based merely on *universally accepted fundamental physical principles* such as invariance principles, conservation principles, causality principle, the laws of thermodynamics and well-posedness of the IVP. In [30], this question was considered in the Lagrangian coordinates and the following requirements to PDEs were imposed:

- PDEs are invariant under any rotations of the space
- PDEs are compatible with an extra conservation law

- PDEs are *generated* by only one thermodynamic potential
- PDEs are nonlinear but have well-posed IVP (i.e. they are first-order symmetric hyperbolic)
- PDEs are conservative and the fluxes are generated by invariant differential operators only, such as div, grad and curl.

Then, based on the analysis of various continuum mechanics models and the group representation theory, a quite general system (system (25) in [30]) of Lagrangian conservation laws satisfying all the above requirements was proposed. Moreover, it has appeared that even a smaller system of PDEs is sufficient for many practical applications and it was proposed to consider a subsystem of the general one which we shall refer to as the *master system* and is discussed in the following section.

3.1.1 Energy formulation

As will be shown thorough this paper, one of the apparent common things between the SHTC and GENERIC frameworks is the use of the state variables and the generating potentials and their conjugate ones with respect to the Legendre transformation, see also a discussion in [76]. In both frameworks the primary state variables are the density like fields and the primary potential is the total energy conservation. The conjugate state variables and potential have the meaning of the fluxes and a generalized pressure accordingly. In this section, we present the Energy formulation of the SHTC Lagrangian equations, i.e. they are written in terms of the primary density fields and the total energy potential. The conjugate formulation is discussed in Section 3.1.2.

The master system is formulated with respect to the following state variables

$$\mathbf{q} = (\mathbf{m}, \mathbf{F}, \mathbf{h}, e, \mathbf{w}, \sigma) \quad (16)$$

and the thermodynamic potential

$$U = U(\mathbf{q}), \quad (17)$$

where $\mathbf{m} = [m_i]$ is a vector field with the meaning of the generalized momentum, e.g. see [18], $\mathbf{F} = [F_{ij}]$ is the deformation gradient, $e = [e_i]$ and $\mathbf{h} = [h_i]$ are vector fields which can be interpreted as electromagnetic fields [18], σ is a scalar field which is transported according to the vector field $\mathbf{w} = [w_i]$. The later two are used for modeling of the mass and heat transfer where σ has the meaning of mass concentration or the entropy, accordingly.

In fact, we do not need to give a precise meaning to the fields (16) and the further analysis is independent of specific physical interpretations. The meanings of the fields (16) depend on the specification of the generating potential U in the SHTC framework. For instance, in the example Section 6, we show that the same equations can describe the heat and mass transfer and the represented physics is fully determined by the way we define the generating potential.

All over the paper, we use the summation over the repeated indexes and we denote the partial derivative of the potentials with respect to the state variables as $U_{q_i} = \partial U / \partial q_i$, $U_{\mathbf{q}} = [U_{q_i}]$, $U_{m_i} = \partial U / \partial m_i$, etc.

The master system that fulfills all of the requirements listed in Section 2.1 was proposed in [30]. A more general master system however can be easily proposed, see [30], but the minimal system then was selected after analysis of

a large number of continuum mechanics models. Such a minimal master system reads as

$$\frac{dm_i}{dt} - \frac{\partial U_{F_{ij}}}{\partial y_j} = 0, \quad (18a)$$

$$\frac{dF_{ij}}{dt} - \frac{\partial U_{m_i}}{\partial y_j} = 0, \quad (18b)$$

$$\frac{dh_i}{dt} + \varepsilon_{ijk} \frac{\partial U_{e_k}}{\partial y_j} = 0, \quad (18c)$$

$$\frac{de_i}{dt} - \varepsilon_{ijk} \frac{\partial U_{h_k}}{\partial y_j} = 0, \quad (18d)$$

$$\frac{dw_j}{dt} + \frac{\partial U_\sigma}{\partial y_j} = 0, \quad (18e)$$

$$\frac{d\sigma}{dt} + \frac{\partial U_{w_j}}{\partial y_j} = 0. \quad (18f)$$

In addition, an extra conservation law is the consequence of the above PDEs

$$\frac{dU}{dt} - \frac{\partial}{\partial y_j} \left(U_{v_i} U_{F_{ij}} + \varepsilon_{ijk} U_{e_i} U_{h_k} - U_\sigma U_{w_j} \right) = 0. \quad (19)$$

It can be obtained as the sum of the governing PDEs multiplied by the corresponding fluxes:

$$(19) \equiv U_{m_i} \cdot (18a) + U_{F_{ij}} \cdot (18b) + U_{e_i} \cdot (18d) + U_{h_i} \cdot (18c) + U_\sigma \cdot (18f) + U_{w_j} \cdot (18e). \quad (20)$$

Usually, in such a formulation, the potential U plays the role of the *total energy density* (per unit mass), i.e. it relates to the total specific energy \mathcal{U} (per unit volume) as $U = \rho_0 \mathcal{U}$, where ρ_0 is the reference mass density.

The first pair of equations usually represents the equations of motion, while the second pair, (18d), (18c), may represent a non-linear generalization of the Maxwell equations. However, as we already mentioned, they may have a drastically different interpretations. For example, in [76], the equations with the structure of (18d), (18c) were used to describe the dynamics of flow defects, while in Section 3.2.6, the Eulerian counterpart of (18c) is used to describe a weakly non-local dynamics.

3.1.2 Conjugate formulation and symmetric form

System (18) is a system of conservation laws and it is compatible with the accompanying conservation law (19) and hence, according to the Godunov-Friedrichs-Lax theorem discussed in Section 2, it can be written in Godunov form (4) and hence it is symmetric hyperbolic. In order to see this explicitly, it is necessary to introduce fluxes of (18) as new state variables $\mathbf{p} = [p_i]$ and a potential $M(\mathbf{p})$ (it has the meaning close to the pressure).

The new variables and the potential (\mathbf{p}, M) are *thermodynamically conjugate* to the pair (\mathbf{q}, U) in the following sense

$$\mathbf{p} = U_{\mathbf{q}}, \quad M(\mathbf{p}) = q_i U_{q_i} - U = q_i p_i - U, \quad (21)$$

i.e. $M(\mathbf{p})$ is the Legendre transformation of $U(\mathbf{q})$. Of course, the vice versa is also true

$$\mathbf{q} = M_{\mathbf{p}}, \quad U(\mathbf{q}) = p_i M_{p_i} - M = q_i p_i - M. \quad (22)$$

In order the relation (21) and (22) be a one-to-one relation, it is necessary that $U(\mathbf{q})$ would be a convex function

($M(\mathbf{p})$ would be then also convex because of the properties of the Legendre transformation), i.e.

$$U_{\mathbf{q}\mathbf{q}} = (M_{\mathbf{p}\mathbf{p}})^{-1} > 0. \quad (23)$$

3.1.3 Complimentary structure and odd-even parity

One may clearly note a *complimentary* structure of the equations (18), i.e. when a field under the time derivative, say m_i in (18a), then appears in the flux of the complimentary equation (18b) in the partial derivative U_{m_i} of the generating potential U . Thus equations (18) are split into complimentary pairs for (m_i, F_{ij}) , (h_i, e_i) and (w_i, σ) . Therefore, when one constructs a specific model in the SHTC framework, it is impossible to take only one PDE in a pair because its flux will be undetermined. However, an arbitrary number of ordinary differential equations

$$\frac{\partial q_i}{\partial t} = f_i(\mathbf{q}) \quad (24)$$

can be added to system (18). Here, q_i might be scalars or entries of a vector or tensor field, $f_i(\mathbf{q})$ are sufficiently smooth functions of the entire vector of state variables. For example, equations (18a), (18a) and ordinary differential equations of type (24) constitute the nonlinear elastoplasticity model studied in [31].

The complimentary structure of the SHTC equations can be also seen from the point of view of the *odd-even parity* with respect to *time-reversal transformation* (TRT) [71, 74] which has been playing a central role in thermodynamics for very long time. If a field changes its sign under TRT, i.e. when the velocities of all microscopic particles (and magnetic field intensity) are inverted, the quantity is odd. If the quantity is not altered by the inversion of velocities, it is even. In applications we dealt with so far, in each complimentary pair (m_i, F_{ij}) , (h_i, e_i) and (w_i, σ) one of the fields has even parity while the second has odd parity. For example, if m_i is treated as the momentum density and F_{ij} is treated as the deformation gradient then m_i is odd while F_{ij} is even; if h_i is treated as the magnetic field and e_i is treated as the electric field then h_i is odd while e_i is even; if w_i is treated as the momentum of heat carriers and σ is treated as the entropy density then w_i is odd while σ is even. The odd-even parity of the state variables is also evident from the geometric form of the Lagrangian SHTC equations presented in Section 4.1.

3.1.4 Involution constraints

Solutions to the master system satisfy some *stationary laws* that are conditioned by the structure of the fluxes:

$$\frac{\partial F_{ij}}{\partial y_k} - \frac{\partial F_{ik}}{\partial y_j} = \text{const}, \quad \frac{\partial h_i}{\partial y_i} = \text{const}, \quad \frac{\partial e_i}{\partial y_i} = \text{const}, \quad \frac{\partial w_j}{\partial y_k} - \frac{\partial w_k}{\partial y_j} = \text{const}, \quad (25)$$

In fact, these stationary laws impose constraints on the initial data and hold for every $t > 0$ if they were satisfied at $t = 0$. Indeed, applying the divergence operator, for instance, to equation (18d) we obtain

$$\frac{d}{dt} \left(\frac{\partial e_i}{\partial y_i} \right) = 0. \quad (26)$$

Thus, the stationary laws (25) should be considered as *involution constraints*. Moreover, as we shall see in Section 3.1.5 where the master system is derived from the variational principle, these involution constraints play the role of *integrability conditions* to the equations of motion if, however, the constants on the right-hand sides of (25) equal to zero.

3.1.5 Variational nature of the SHTC equations

The master system was proposed in [30] based merely on the requirements listed at the beginning of Section 3.1. In this section, we show that the master system can be also obtained from Hamilton's principle of stationary action if

the Lagrangian is taken as a partial Legendre transformation (see details below) of the total energy potential and stationary laws (25) have zeros on the right-hand sides. Partly, these results were published in Chapter 23 of the Russian edition [42] of the book [41] as well as in the recent paper [18].

We start by introducing two vector potentials and two scalar potentials:

$$x_i(t, \mathbf{y}), \quad a_i(t, \mathbf{y}), \quad \varphi(t, \mathbf{y}), \quad \chi(t, \mathbf{y}) \quad (27)$$

and denote their derivatives as follows

$$\hat{v}_i = \frac{\partial x_i}{\partial t}, \quad \hat{F}_{ij} = \frac{\partial x_i}{\partial y_j}, \quad (28a)$$

$$\hat{e}_i = -\frac{\partial a_i}{\partial t} - \frac{\partial \varphi}{\partial y_i}, \quad \hat{h}_i = \varepsilon_{ijk} \frac{\partial a_k}{\partial y_j}, \quad (28b)$$

$$\hat{\sigma} = \frac{\partial \chi}{\partial t}, \quad \hat{w}_i = \frac{\partial \chi}{\partial y_i}. \quad (28c)$$

The independent variables t , $\mathbf{y} = [y_i]$ and $\mathbf{x} = [x_i]$ can be seen as the time, the Lagrangian and Eulerian spatial coordinates respectively, a_i and φ can be considered as the conventional electromagnetic potentials, while at this stage it is difficult to give a certain meaning to χ .

Then, we define the action integral

$$\mathcal{L} = \int \Lambda \, d\mathbf{y} dt, \quad (29)$$

where $\Lambda = \Lambda(\hat{v}_i, \hat{F}_{ij}, \hat{h}_i, \hat{e}_i, \hat{w}_i, \hat{\sigma})$ is the Lagrangian.

The first variation of \mathcal{L}

$$\delta \mathcal{L} = \int \left[\left(\frac{\partial \Lambda}{\partial t} + \frac{\partial \Lambda}{\partial y_j} \hat{F}_{ij} \right) \delta x_i + \left(\frac{\partial \Lambda}{\partial t} + \varepsilon_{ijk} \frac{\partial \Lambda}{\partial y_j} \hat{h}_k \right) \delta a_i + \frac{\partial \Lambda}{\partial y_j} \delta \varphi + \left(\frac{\partial \Lambda}{\partial t} + \frac{\partial \Lambda}{\partial y_j} \hat{w}_j \right) \delta \chi \right] d\mathbf{y} dt \quad (30)$$

gives us the Euler-Lagrange equations

$$\frac{\partial \Lambda}{\partial t} + \frac{\partial \Lambda}{\partial y_j} \hat{F}_{ij} = 0, \quad (31a)$$

$$\frac{\partial \Lambda}{\partial t} + \varepsilon_{ijk} \frac{\partial \Lambda}{\partial y_j} \hat{h}_k = 0, \quad \frac{\partial \Lambda}{\partial y_j} = 0, \quad (31b)$$

$$\frac{\partial \Lambda}{\partial t} + \frac{\partial \Lambda}{\partial y_j} \hat{w}_j = 0 \quad (31c)$$

To this system, the following integrability conditions should be added (they are trivial consequences of the definitions (28))

$$\frac{\partial \hat{F}_{ij}}{\partial t} - \frac{\partial \hat{v}_i}{\partial y_j} = 0, \quad \frac{\partial \hat{F}_{ij}}{\partial y_k} - \frac{\partial \hat{F}_{ik}}{\partial y_j} = 0, \quad (32a)$$

$$\frac{\partial \hat{h}_i}{\partial t} + \varepsilon_{ijk} \frac{\partial \hat{e}_k}{\partial y_j} = 0, \quad \frac{\partial \hat{h}_j}{\partial y_i} = 0. \quad (32b)$$

$$\frac{\partial \hat{w}_j}{\partial t} - \frac{\partial \hat{\sigma}}{\partial y_j} = 0, \quad \frac{\partial \hat{w}_i}{\partial y_j} - \frac{\partial \hat{w}_j}{\partial y_i} = 0. \quad (32c)$$

In order to rewrite equations (31)–(32) in the form of system (18), let us introduce a potential U as a partial Legendre transformation of the Lagrangian Λ

$$\begin{aligned} dU &= d(\hat{v}_i \Lambda_{\hat{v}_i} + \hat{e}_i \Lambda_{\hat{e}_i} + \hat{\sigma} \Lambda_{\hat{\sigma}} - \Lambda) = \hat{v}_i d\Lambda_{\hat{v}_i} + \hat{e}_i d\Lambda_{\hat{e}_i} + \hat{\sigma} d\Lambda_{\hat{\sigma}} - \Lambda_{\hat{F}_{ij}} d\hat{F}_{ij} - \Lambda_{\hat{h}_i} d\hat{h}_i - \Lambda_{\hat{w}_i} d\hat{w}_i = \\ &\hat{v}_i d\Lambda_{\hat{v}_i} + \hat{e}_i d\Lambda_{\hat{e}_i} + \Lambda_{\hat{F}_{ij}} d(-\hat{F}_{ij}) + \Lambda_{\hat{h}_i} d(-\hat{h}_i) + \Lambda_{\hat{w}_i} d(-\hat{w}_i). \end{aligned} \quad (33)$$

Hence, denoting $m_i = \Lambda_{\hat{v}_i}$, $e_i = \Lambda_{\hat{e}_i}$, $\sigma = \Lambda_{\hat{\sigma}}$, $F_{ij} = -\hat{F}_{ij}$, $h_i = -\hat{h}_i$ and $w_i = -\hat{w}_i$, we get the thermodynamic identity

$$dU = U_{m_i} dm_i + U_{F_{ij}} dF_{ij} + U_{h_i} dh_i + U_{e_i} de_i + U_{w_i} dw_i + U_{\sigma} d\sigma.$$

Eventually, in terms of the variables

$$\mathbf{q} = (m_i, F_{ij}, h_i, e_i, w_i, \sigma) \quad (34)$$

and the potential $U = U(\mathbf{q})$, equations (31a), (31b)₁, (31c), (32a)₁, (32b)₁ and (32c)₁ transform exactly into the equations (18) which should be supplemented by stationary constraints (32a)₂, (31b)₂, (32b)₂ and (32c)₂ which now read as

$$\frac{\partial F_{ij}}{\partial y_k} - \frac{\partial F_{ik}}{\partial y_j} = 0, \quad \frac{\partial e_i}{\partial y_i} = 0, \quad \frac{\partial h_i}{\partial y_i} = 0, \quad \frac{\partial w_i}{\partial y_j} - \frac{\partial w_j}{\partial y_i} = 0. \quad (35)$$

3.2. Eulerian frame

We now present the Eulerian SHTC equations which later will be shown to be compatible with GENERIC in Section 4. We first present the equations which are directly obtained from the Lagrangian master system (18) by means of the Lagrange-to-Euler change of the spatial variables which results in the change of the time and spatial derivatives

$$\frac{d}{dt} = \frac{\partial}{\partial t} + v_k \frac{\partial}{\partial x_k}, \quad \frac{\partial}{\partial y_j} = F_{kj} \frac{\partial}{\partial x_k}. \quad (36)$$

The resulting Eulerian master system have a more complicated structure of PDEs than its Lagrangian counterpart (18). The main difference is that, in contrast to the fully conservative Lagrangian master system, the Eulerian equations are inherently non-conservative ones and do not admit the original Godunov structure (4) but generalized it in the following way

$$\frac{\partial L_{\mathbf{p}}}{\partial t} + \frac{\partial (v_k L)_{\mathbf{p}}}{\partial x_k} + C_k \frac{\partial \mathbf{p}}{\partial x_k} = 0, \quad (37)$$

where one can recognize the Godunov structure (4) in the first two terms, while $C_k^T = C_k$ are some symmetric matrices. The reason for the insufficiency of the original conservative Godunov structure (4) is the fact that the involution constraints in the Eulerian frame are not just auxiliary equations as in the Lagrangian frame but they are an intrinsic part of the structure of the Eulerian SHTC equations and contribute into matrices C_k .

As in the previous section, we start the discussion with the Eulerian SHTC equations in terms of the total energy potential and density-type state variables. We then discuss the role of the involution constraints and only for pedagogical reasons we present a fully conservative reformulation of the SHTC equations in Section 3.2.2. This reformulation results in the lose of Galilean invariance property of the Eulerian equations and it is should be avoided to thoughtlessly use this conservative formulation in practice. We conclude the section by presenting the conjugate formulation and demonstrating how the involution constraints essentially affect the symmetrization procedure of the SHTC equations in the Eulerian frame.

3.2.1 Energy formulation

Transformation from the Lagrangian SHTC master system (18) to the Eulerian equations has been done in [30] while omitting non-trivial details of the calculation, which can be now found in Appendix A. The resulting non-conservative Eulerian system of equations is

$$\frac{\partial m_i}{\partial t} + \frac{\partial}{\partial x_k} \left(m_i v_k + \delta_{ik} (\rho E_\rho + \sigma E_\sigma + m_l E_{m_l} + e_l E_{e_l} + h_l E_{h_l} - E) - e_k E_{e_i} - h_k E_{h_i} + w_i E_{w_k} + A_{jk} E_{A_{ji}} \right) = 0, \quad (38a)$$

$$\frac{\partial A_{ik}}{\partial t} + \frac{\partial (A_{il} v_l)}{\partial x_k} + v_j \left(\frac{\partial A_{ik}}{\partial x_j} - \frac{\partial A_{ij}}{\partial x_k} \right) = 0, \quad (38b)$$

$$\frac{\partial h_i}{\partial t} + \frac{\partial (h_i v_k - v_i h_k + \varepsilon_{ikl} E_{e_l})}{\partial x_k} + v_i \frac{\partial h_k}{\partial x_k} = 0, \quad (38c)$$

$$\frac{\partial e_i}{\partial t} + \frac{\partial (e_i v_k - v_i e_k - \varepsilon_{ikl} E_{h_l})}{\partial x_k} + v_i \frac{\partial e_k}{\partial x_k} = 0, \quad (38d)$$

$$\frac{\partial w_k}{\partial t} + \frac{\partial (v_l w_l + E_\sigma)}{\partial x_k} + v_j \left(\frac{\partial w_k}{\partial x_j} - \frac{\partial w_j}{\partial x_k} \right) = 0. \quad (38e)$$

$$\frac{\partial \sigma}{\partial t} + \frac{\partial (\sigma v_k + E_{w_k})}{\partial x_k} = 0, \quad (38f)$$

$$\frac{\partial \rho}{\partial t} + \frac{\partial (\rho v_k)}{\partial x_k} = 0. \quad (38g)$$

We note that it is always implied in the SHTC framework (as well as in the GENERIC) that the velocity field v_i and the total momentum field are conjugate fields, i.e. $v_i = E_{m_i}$. Thus, in all the equations of system (38), v_i can be substituted by E_{m_i} but we prefer to keep v_i explicitly written for historical reasons. Later, in Section 4 we shall always use E_{m_i} instead of v_i .

Also, note that we use the same notations for the state variables in system (18) and (38), however they are different fields which relate to each other by the formulas

$$\mathbf{m}' = w \mathbf{m}, \quad \mathbf{F} = \mathbf{A}^{-1}, \quad \rho_0 = w \rho, \quad w = \det(\mathbf{F}), \quad (39a)$$

$$\mathbf{e}' = w \mathbf{A} \mathbf{e}, \quad \mathbf{h}' = w \mathbf{A} \mathbf{h}, \quad (39b)$$

$$\sigma' = w \sigma, \quad \mathbf{w}' = \mathbf{F}^T \mathbf{w}, \quad (39c)$$

while the Lagrangian total energy density U is related to Eulerian total energy density E as

$$U = w E. \quad (39d)$$

In these relations, \mathbf{m}' , \mathbf{F} , \mathbf{h}' , \mathbf{e}' , \mathbf{w}' and σ' are the Lagrangian fields, i.e. exactly those fields in equations (18), while \mathbf{m} , \mathbf{A} , \mathbf{h} , \mathbf{e} , \mathbf{w} and σ are the fields in the Eulerian equations (38). Also, ρ_0 is the reference (Lagrangian) mass density.

Exactly as in the Lagrangian framework (see summation rule (20)), the conservation of the total energy density

E ,

$$\frac{\partial E}{\partial t} + \frac{\partial}{\partial x_k} \left(v_k E + v_i \left[(\rho E_\rho + \sigma E_\sigma + m_l E_{m_l} + e_l E_{e_l} + h_l E_{h_l} - E) \delta_{ik} - e_k E_{e_i} - h_k E_{h_i} + w_i E_{w_k} + A_{jk} E_{A_{ji}} \right] \right. \\ \left. + \varepsilon_{ijk} E_{e_i} E_{h_j} + E_\sigma E_{w_k} \right) = 0, \quad (40)$$

is just a consequence of the governing PDEs (38). Namely, it is the sum of PDEs (38) multiplied by the corresponding factors, the conjugate state variables:

$$(40) \equiv E_{m_i} \cdot (38a) + E_{A_{ik}} \cdot (38b) + E_{h_i} \cdot (38c) + E_{e_i} \cdot (38d) + E_{w_k} \cdot (38e) + E_\sigma \cdot (38f) + E_\rho \cdot (38g). \quad (41)$$

As discussed in Section 5, this summation rule also plays a key role in the introduction of the dissipative terms in the SHTC framework. The calculation can be found in the Russian edition [42] of the book [41]. However, for completeness, we shall also demonstrate it in this paper after we present the conjugate formulation in Section 3.2.4.

We note that there is no need to use the mass conservation equation in the Lagrangian frame because the mass density is $\rho = \rho_0 w^{-1}$. Moreover, as it has been shown several times, e.g. [42, 76], the mass conservation (38g) is the consequence of the time evolution (38b) for the distortion matrix \mathbf{A} and hence, can be dropped out. But in fact, in the Eulerian frame, for many reasons, it is easier to treat ρ as an independent state variable governing by its own time evolution. In this sense, the complimentary pair of Lagrangian equations (18a) and (18b) corresponds to three equations (38a), (38b) and (38g) of the Eulerian SHTC master system. Also, remark that an arbitrary number of equations with the structure of (38g) can be added to system (38). In practical cases, the entropy conservation law for the ideal fluids or time evolution for the volume fraction in multi-phase models have the same structure as (38g), e.g. see Section 6.5.2. Thus, (38g) symbolizes various equations with such a structure.

Eventually, we mention a few examples in which equations (38) were used, see also more details in Section 6. Thus, equations (38a), (38b), were used in [38, 85, 41, 4, 23, 3, 14] for modeling of elastic and elastoplastic deformations in metals, while the same equations constitute the unified formulation of continuum mechanics proposed in [77, 17, 78, 18] and can be used for modeling of fluid flows. Equations (38a)–(38c) were used in [86, 18] to describe the electrodynamics of moving medium, equations (38a), (38e) and (38f) can be used to model multiphase flows, heat conduction, superfluid helium flows, see [86, 83, 82, 81].

3.2.2 Involution constraints

As we already mentioned, the involution constraints play an exceptional role in the Eulerian SHTC equations. This role will show up in all the subsequent sections.

We start the discussion by demonstrating that the following *stationary constraints*³

$$\nabla \times \mathbf{A} = \mathbf{B}, \quad \nabla \cdot \mathbf{h} = Q, \quad \nabla \cdot \mathbf{e} = R, \quad \nabla \times \mathbf{w} = \mathbf{\Omega}, \quad (42)$$

or, in the component-wise notations,

$$\varepsilon_{ajk} \frac{\partial A_{ik}}{\partial x_j} = B_{ia}, \quad \frac{\partial h_k}{\partial x_k} = Q, \quad \frac{\partial e_k}{\partial x_k} = R, \quad \varepsilon_{ajk} \frac{\partial w_k}{\partial x_j} = \Omega_a, \quad (43)$$

³The curl operator is applied to matrix \mathbf{A} in the column-wise manner.

are compatible with system (38), where the quantities \mathbf{B} , Q , R and $\mathbf{\Omega}$ satisfy the following conservation laws

$$\frac{\partial B_{ij}}{\partial t} + \frac{\partial(B_{ij}v_k - v_j B_{ik})}{\partial x_k} + v_j \frac{\partial B_{ik}}{\partial x_k} = 0, \quad (44a)$$

$$\frac{\partial R}{\partial t} + \frac{\partial(Rv_k)}{\partial x_k} = 0, \quad (44b)$$

$$\frac{\partial Q}{\partial t} + \frac{\partial(Qv_k)}{\partial x_k} = 0, \quad (44c)$$

$$\frac{\partial \Omega_j}{\partial t} + \frac{\partial(\Omega_j v_k - v_j \Omega_k)}{\partial x_k} + v_j \frac{\partial \Omega_k}{\partial x_k} = 0. \quad (44d)$$

Indeed, we are going to demonstrate that the following involution constraints

$$\frac{\partial}{\partial t} (\nabla \times \mathbf{A} - \mathbf{B}) \equiv 0, \quad \frac{\partial}{\partial t} (\nabla \cdot \mathbf{e} - R) \equiv 0, \quad \frac{\partial}{\partial t} (\nabla \cdot \mathbf{h} - Q) \equiv 0, \quad \frac{\partial}{\partial t} (\nabla \times \mathbf{w} - \mathbf{\Omega}) \equiv 0, \quad (45)$$

are satisfied on the solutions to (38) if they hold true at the initial moment of time. Indeed, $\nabla \times \mathbf{A}$, $\nabla \cdot \mathbf{e}$, $\nabla \cdot \mathbf{h}$ and $\nabla \times \mathbf{w}$ satisfy the following evolution equations⁴

$$\frac{\partial(\nabla \times \mathbf{A})_{ij}}{\partial t} + \frac{\partial((\nabla \times \mathbf{A})_{ij}v_k - v_j(\nabla \times \mathbf{A})_{ik})}{\partial x_k} + v_j \frac{\partial(\nabla \times \mathbf{A})_{ik}}{\partial x_k} = 0, \quad (46a)$$

$$\frac{\partial(\nabla \cdot \mathbf{h})}{\partial t} + \frac{\partial(v_k(\nabla \cdot \mathbf{h}))}{\partial x_k} = 0, \quad (46b)$$

$$\frac{\partial(\nabla \cdot \mathbf{e})}{\partial t} + \frac{\partial(v_k(\nabla \cdot \mathbf{e}))}{\partial x_k} = 0, \quad (46c)$$

$$\frac{\partial(\nabla \times \mathbf{w})_j}{\partial t} + \frac{\partial((\nabla \times \mathbf{w})_j v_k - v_j(\nabla \times \mathbf{w})_k)}{\partial x_k} + v_j \frac{\partial(\nabla \times \mathbf{w})_k}{\partial x_k} = 0. \quad (46d)$$

By comparing (44) and (46), one may conclude that (45) is satisfied if it was true at $t = 0$. We emphasize that even though the non-conservative terms in (46a) and (46d) are zeros they can not be omitted because this would violate the Galilean invariance property of these equations and also change the characteristic structure.

In some cases, at $t = 0$, one may assume that $\mathbf{B} = 0$, $\mathbf{h} = 0$, $\mathbf{e} = 0$ and $\mathbf{\Omega} = 0$ and hence, (44) has trivial solution, and (45) becomes

$$\nabla \times \mathbf{A} \equiv 0, \quad \nabla \cdot \mathbf{h} \equiv 0, \quad \nabla \cdot \mathbf{e} \equiv 0, \quad \nabla \times \mathbf{w} \equiv 0 \quad (47)$$

for any $t > 0$. However, in some applications we have to set $\mathbf{B} \neq 0$, $\mathbf{h} \neq 0$, $\mathbf{e} \neq 0$ or $\mathbf{\Omega} \neq 0$ in the initial conditions and hence, system (44) has non-trivial time-dependent solution. Thus, it is necessary to note that quantities \mathbf{B} , \mathbf{h} , \mathbf{e} and $\mathbf{\Omega}$ may have a certain physical meaning. For example, $\nabla \times \mathbf{A}$ has the meaning of the density of microscopic defects (dislocation density tensor) [42, 41, 76] in the elatoplasticity theory. Thus, if the material has suffered from plastic deformations in the past ($t < 0$) then at $t = 0$ we have $\nabla \times \mathbf{A} \neq 0$. For example, the impact of the dislocation density on the dispersive properties of the elastic waves was studied in [84] where it was assumed that the material has non-zero initial concentration of defects.

As we shall see later, the involution constraints (45) and (47) have strong impact on the symmetrization of system (38). Thus, if constraints (47) hold then the symmetrization is much simpler than in the case of (45). If only

⁴For the derivation of (46a) and (46d), see Appendix C, while equations (46c) and (46b) can be easily obtained by applying $\partial/\partial x_i$ to the SHTC equations (38d) and (38c) accordingly.

(45) holds, then system (38) can be symmetrized in an extended sense. Namely, we shall demonstrate that system (38) extended by adding to it constraints (44) can be symmetrized. This is, in fact, a very interesting feature of the SHTC formulation because one may note that the extended system (38), (44) is *weakly nonlocal* because the quantities $\nabla \times \mathbf{A}$, $\nabla \cdot \mathbf{e}$, etc, represent a coarser scale than the state variables \mathbf{A} , \mathbf{e} , etc. This will be even more pronounced when we shall discuss the irreversible dynamics within the SHTC framework in Section 5 because in the presence of dissipative processes the quantities $\nabla \times \mathbf{A}$, $\nabla \cdot \mathbf{e}$, $\nabla \times \mathbf{w}$ are not zeros even if they were zero at $t = 0$.

3.2.3 Weakly non-local conservative non-symmetrizable formulation

In contrast to the Lagrangian master system (18), the Eulerian system (38) is not a system of *conservation laws* due to the presence of the non-conservative differential terms,

$$v_j(\partial_j A_{ik} - \partial_k A_{ij}), \quad v_i \partial_k h_k, \quad v_i \partial_k e_k, \quad v_j(\partial_j w_k - \partial_k w_j), \quad (48)$$

in equations (38b), (38c), (38d) and (38e) accordingly. Therefore, there is a temptation to rewrite Eulerian SHTC equations in a fully conservative form using the results of Section 3.2.2 and to symmetrize it in a way discussed in Section 2.1. Unfortunately, such an idea is not viable, and the pedagogical goal of this section is to demonstrate this.

For the sake of simplicity, we consider the ideal MHD equations in this section. However, all the analysis can be applied to the entire system (38) without any changes. Thus, let us consider the MHD equations which using the results of Section 3.2.2 can be written as

$$\partial_t m_i + \partial_k (m_i v_k + \delta_{ik} (\rho E_\rho + \sigma E_\sigma + m_l E_{m_l} + B_l E_{B_l} - E) - B_k E_{B_i}) = 0, \quad (49a)$$

$$\partial_t B_i + \partial_k (B_i v_k - v_i B_k) = -v_i Q, \quad (49b)$$

$$\partial_t Q + \partial_k (Q v_k) = 0, \quad (49c)$$

$$\partial_t \rho + \partial_k (\rho v_k) = 0, \quad (49d)$$

$$\partial_t \sigma + \partial_k (\sigma v_k) = 0, \quad (49e)$$

$$\partial_t E + \partial_k (v_k E + v_i [(\rho E_\rho + \sigma E_\sigma + m_l E_{m_l} + B_l E_{B_l} - E) \delta_{ik} - B_k E_{w_i}]) = 0, \quad (50)$$

which look like the system of conservation laws (i.e. all the differential terms are written in a conservative manner) with the algebraic source term $-v_i Q$ in the PDE for B_i which can be ignored in the symmetrization process as a low order term. Here, instead of the field h_i used in system (38) we use a convectional notation B_i for the magnetic field, ρ is the mass density, σ is treated as the entropy density. As discussed in Section 3.2.2, on the solution of the system, the involution constraint

$$\frac{\partial}{\partial t} (\nabla \cdot \mathbf{B} - Q) = 0 \quad (51)$$

is fulfilled. This involution constraint can be reduced to $\partial_t Q = 0$ because $\nabla \cdot \mathbf{B} = 0$ which has the meaning of the absence of magnetic monopoles. Therefore, one may even suggest to consider a simpler model

$$\partial_t m_i + \partial_k (m_i v_k + \delta_{ik} (\rho E_\rho + \sigma E_\sigma + m_l E_{m_l} + B_l E_{B_l} - E) - B_k E_{B_i}) = 0, \quad (52a)$$

$$\partial_t B_i + \partial_k (B_i v_k - v_i B_k) = 0, \quad (52b)$$

$$\partial_t \rho + \partial_k (\rho v_k) = 0, \quad (52c)$$

$$\partial_t \sigma + \partial_k (\sigma v_k) = 0, \quad (52d)$$

as it is done in many works, e.g. see discussion in [79]. However, neither (49) nor (52) is compatible with the energy conservation law (50). Indeed, equations (49) and (50) form an overdetermined system of PDEs. Hence, the energy

conservation should be a combination of equations (49) or (52). However, as it was shown by Godunov in [38], the energy flux can not be obtained as a combination of the conservative fluxes (49) or (52) because the energy flux is exactly (see details in Appendix B)

$$\text{flux}(50) \equiv E_{m_i} \cdot \text{flux}(52a) + E_{B_i} \cdot \text{flux}(52b) + E_\rho \cdot \text{flux}(52c) + E_{B_i} v_i \partial_k B_k. \quad (53)$$

Therefore, the energy conservation is *not a combination* of either (49) or (52) and hence the symmetrization is impossible by means of the GFL theorem discussed in Section 2.1. Other ways of symmetrization of the MHD equations are unknown. Eventually, both formulations (49) and (52) are not Galilean invariant, e.g. see [79], and should be avoided in practical use.

The MHD formulation which is compatible with the energy conservation (50) is the following *non-conservative* system

$$\partial_t m_i + \partial_k (m_i v_k + \delta_{ik} (\rho E_\rho + \sigma E_\sigma + m_l E_{m_l} + B_l E_{B_l} - E) - B_k E_{B_i}) = 0, \quad (54a)$$

$$\partial_t B_i + \partial_k (B_i v_k - v_i B_k) + v_i \partial_k B_k = 0, \quad (54b)$$

$$\partial_t \rho + \partial_k (\rho v_k) = 0, \quad (54c)$$

$$\partial_t \sigma + \partial_k (\sigma v_k) = 0. \quad (54d)$$

This formulation is non-conservative due to the presence of the non-conservative differential term $v_i \partial_k B_k$ in the PDE for B_i . However, exactly this formulation is compatible with the energy law (50), it is symmetrizable as was shown in [38], it has the structure of the equation (38c) of the SHTC master system (38), it is Galilean invariant, and exactly equation (54b) can be obtained from the Lagrangian equation for the magnetic field

$$\frac{dB_i}{dt} = 0, \quad (55)$$

see details in Section A.3.

In summary, it is important not only to pay attention to casting the equations into the form of conservation laws, but also to the companion conservation law, which has to be compatible with the rest of the evolution equations. Otherwise, the GFL theorem can not be applied.

3.2.4 Conjugate formulation

In the previous section, the governing equations were formulated in terms of the state variables \mathbf{q} and the total energy density $E(\mathbf{q})$. In this section, we also provide another, *thermodynamically conjugate*, formulation in terms of the conjugate state variables \mathbf{p} (flux fields) and a potential $L(\mathbf{p})$ which is the Legendre transform of $E(\mathbf{q})$ and has the physical meaning of a generalized pressure. This formulation allows to emphasize an exceptional role of the generating potentials L and E and shall be used to prove the symmetric hyperbolicity of the governing PDEs in Sections 3.2.6 and 3.2.5 within the SHTC formalism.

As in the Lagrangian framework, we introduce the conjugate variables

$$\mathbf{p} = (r, v_i, \alpha_{ik}, b_i, d_i, \eta_k, \theta) \quad (56)$$

as the partial derivatives of the total energy density $E(\mathbf{q})$ with respect to the conservative state variables

$$v_i = E_{m_i}, \quad \alpha_{ik} = E_{A_{ik}}, \quad b_i = E_{h_i}, \quad d_i = E_{e_i}, \quad \eta_k = E_{w_k}, \quad \theta = E_\sigma, \quad r = E_\rho,$$

and the new potential

$$L(\mathbf{p}) = \mathbf{q} \cdot \mathbf{p} - E = r\rho + v_i m_i + \alpha_{ij} A_{ij} + d_i e_i + b_i h_i + \eta_i w_i + \theta\sigma - E \quad (57)$$

as the Legendre transformation of E . In terms of the new variables \mathbf{p} and potential $L(\mathbf{p})$, system (38) can be rewritten as

$$\frac{\partial L_{v_i}}{\partial t} + \frac{\partial}{\partial x_k} [(v_k L)_{v_i} + \alpha_{mk} L_{\alpha_{mi}} - \delta_{ik} \alpha_{mn} L_{\alpha_{mn}} - d_i L_{d_k} - b_i L_{b_k} + \eta_k L_{\eta_i} - \delta_{ik} \eta_n L_{\eta_n}] = 0, \quad (58a)$$

$$\frac{\partial L_{\alpha_{ik}}}{\partial t} + \frac{\partial [(v_m L)_{\alpha_{im}}]}{\partial x_k} + v_j \left(\frac{\partial L_{\alpha_{ik}}}{\partial x_j} - \frac{\partial L_{\alpha_{ij}}}{\partial x_k} \right) = 0, \quad (58b)$$

$$\frac{\partial L_{b_i}}{\partial t} + \frac{\partial [(v_k L)_{b_i} - v_i L_{b_k} + \varepsilon_{ikl} d_l]}{\partial x_k} + v_i \frac{\partial L_{b_k}}{\partial x_k} = 0, \quad (58c)$$

$$\frac{\partial L_{d_i}}{\partial t} + \frac{\partial [(v_k L)_{d_i} - v_i L_{d_k} - \varepsilon_{ikl} b_l]}{\partial x_k} + v_i \frac{\partial L_{d_k}}{\partial x_k} = 0, \quad (58d)$$

$$\frac{\partial L_{\eta_k}}{\partial t} + \frac{\partial [(v_i L)_{\eta_i} + \theta]}{\partial x_k} + v_j \left(\frac{\partial L_{\eta_k}}{\partial x_j} - \frac{\partial L_{\eta_j}}{\partial x_k} \right) = 0, \quad (58e)$$

$$\frac{\partial L_{\theta}}{\partial t} + \frac{\partial [(v_k L)_{\theta} + \eta_k]}{\partial x_k} = 0, \quad (58f)$$

$$\frac{\partial L_r}{\partial t} + \frac{\partial (v_k L)_r}{\partial x_k} = 0, \quad (58g)$$

while the energy conservation (40) now reads (the prove that it is the consequence of the equations (58) is given in the Appendix B)

$$\begin{aligned} & \frac{\partial}{\partial t} (rL_r + v_i L_{v_i} + \alpha_{ij} L_{\alpha_{ij}} + d_i L_{d_i} + b_i L_{b_i} + \theta L_{\theta} + \eta_k L_{\eta_k} - L) + \\ & \frac{\partial}{\partial x_k} \left(v_k \left(rL_r + v_i L_{v_i} + \alpha_{ij} L_{\alpha_{ij}} + d_i L_{d_i} + b_i L_{b_i} + \theta L_{\theta} + \eta_k L_{\eta_k} - L \right) + \right. \\ & \left. v_i [(L - \alpha_{ab} L_{\alpha_{ab}} - \eta_a L_{\eta_a}) \delta_{ik} + \alpha_{nk} L_{\alpha_{ni}} - d_i L_{d_k} - b_i L_{b_k} + \eta_k L_{\eta_i}] + \varepsilon_{kij} d_i b_j + \theta \eta_k \right) = 0. \quad (59) \end{aligned}$$

It is still not clear though whether (58) can be written as a symmetric hyperbolic system because of the non-trivial structure of the fluxes and non-conservative differential terms. Apparently, the Godunov-Friedrichs-Lax theorem discussed in Section 2.1 can not be applied because (58) is not a system of *conservation laws*. Moreover, it can not be treated as a conservative system in the extended sense as discussed in Section 3.2.3 because energy conservation would be *incompatible* with such conservative equations⁵ and the GFL theorem would still be inapplicable.

The source of the problem is in the following. Unlike the Lagrangian frame, in the Eulerian frame, it has appeared [38, 33, 86, 87] that the involution constraints (45) play an important role in the symmetrization of system (58). If the initial conditions were chosen in such a way that (47) holds true, then the symmetrization is much easier and is discussed in Section 3.2.5 while the general case for non curl-free and divergence-free involution constraints (45) is discussed in Section 3.2.6.

⁵See details in Appendix B, where we prove that if the non-conservative terms are ignored then the energy flux is not a consequence of the other fluxes.

3.2.5 Symmetric form of conjugate formulation for curl-free and divergence-free involution constraints

As in the Lagrangian framework, the parametrization of the governing equations in terms of the conjugate variables \mathbf{p} and generating potential $L(\mathbf{p})$ allows to rewrite the system in a symmetric quasilinear form. However, this is not that straightforward as it is in the Lagrangian frame. In order, to show that (58) is equivalent to a quasilinear symmetric system it is necessary to add the following linear combination of the involution constraints (47)

$$\alpha_{jk} \left(\frac{\partial L_{\alpha_{jk}}}{\partial x_i} - \frac{\partial L_{\alpha_{ji}}}{\partial x_k} \right) + d_i \frac{\partial L_{d_k}}{\partial x_k} + b_i \frac{\partial L_{b_k}}{\partial x_k} + \eta_k \left(\frac{\partial L_{\eta_k}}{\partial x_i} - \frac{\partial L_{\eta_i}}{\partial x_k} \right) \equiv 0 \quad (60)$$

to equation (58a) and *formally* brake its conservative form. After that and some trivial term rearrangements in the rest of the equations, (58) can be rewritten as

$$\frac{\partial L_{v_i}}{\partial t} + \frac{\partial (v_k L)_{v_i}}{\partial x_k} + L_{\alpha_{ij}} \frac{\partial \alpha_{kj}}{\partial x_k} - L_{\alpha_{jk}} \frac{\partial \alpha_{ji}}{\partial x_i} - L_{d_k} \frac{\partial d_i}{\partial x_k} - L_{b_k} \frac{\partial b_i}{\partial x_k} + L_{\eta_i} \frac{\partial \eta_k}{\partial x_k} - L_{\eta_j} \frac{\partial \eta_j}{\partial x_i} = 0, \quad (61a)$$

$$\frac{\partial L_{\alpha_{ij}}}{\partial t} + \frac{\partial (v_k L)_{\alpha_{ij}}}{\partial x_k} + L_{\alpha_{mj}} \frac{\partial v_m}{\partial x_i} - L_{\alpha_{ij}} \frac{\partial v_k}{\partial x_k} = 0, \quad (61b)$$

$$\frac{\partial L_{b_i}}{\partial t} + \frac{\partial (v_k L)_{b_i}}{\partial x_k} - L_{b_k} \frac{\partial v_i}{\partial x_k} + \varepsilon_{ikj} \frac{\partial d_j}{\partial x_k} = 0, \quad (61c)$$

$$\frac{\partial L_{d_i}}{\partial t} + \frac{\partial (v_k L)_{d_i}}{\partial x_k} - L_{d_k} \frac{\partial v_i}{\partial x_k} - \varepsilon_{ikj} \frac{\partial b_j}{\partial x_k} = 0, \quad (61d)$$

$$\frac{\partial L_{\eta_i}}{\partial t} + \frac{\partial (v_k L)_{\eta_i}}{\partial x_k} + L_{\eta_j} \frac{\partial v_j}{\partial x_i} - L_{\eta_i} \frac{\partial v_k}{\partial x_k} + \frac{\partial \theta}{\partial x_i} = 0, \quad (61e)$$

$$\frac{\partial L_{\theta}}{\partial t} + \frac{\partial (v_k L)_{\theta}}{\partial x_k} + \frac{\partial \eta_k}{\partial x_k} = 0, \quad (61f)$$

$$\frac{\partial L_r}{\partial t} + \frac{\partial (v_k L)_r}{\partial x_k} = 0. \quad (61g)$$

Now, these modified equations can be rewritten as a quasilinear system

$$A \frac{\partial \mathbf{p}}{\partial t} + B_k \frac{\partial \mathbf{p}}{\partial x_k} + C_k \frac{\partial \mathbf{p}}{\partial x_k} = 0 \quad (62)$$

which is apparently symmetric because the first two terms in each equation in (61) form symmetric matrices $A(\mathbf{p}) = L_{\mathbf{p}\mathbf{p}}$ and $B_k(\mathbf{p}) = (v_k L)_{\mathbf{p}\mathbf{p}}$ as second order derivatives of the potentials L and $v_k L$ while the rest terms contribute to these matrices only in a symmetric way via symmetric matrices $C_k(\mathbf{p})$. Moreover, if L is a convex potential, then system (58) is *symmetric hyperbolic* because $A = L_{\mathbf{p}\mathbf{p}} > 0$. Recall that due the properties of the Legendre transformation, the convexity of $L(\mathbf{p})$ in \mathbf{p} is equivalent to convexity of $E(\mathbf{q})$ in \mathbf{q} .

To conclude this section, we demonstrate the symmetric structure of the matrix C_k . For example, for $k = 1$ the

submatrix of C_1 corresponding to the variables v_i , η_i and θ is

$$\begin{pmatrix} 0 & 0 & 0 & 0 & -L_{\eta_2} & -L_{\eta_3} & 0 \\ 0 & 0 & 0 & L_{\eta_2} & 0 & 0 & 0 \\ 0 & 0 & 0 & L_{\eta_3} & 0 & 0 & 0 \\ 0 & L_{\eta_2} & L_{\eta_3} & 0 & 0 & 0 & 1 \\ -L_{\eta_2} & 0 & 0 & 0 & 0 & 0 & 0 \\ -L_{\eta_3} & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 \end{pmatrix}. \quad (63)$$

3.2.6 Symmetric form of conjugate weakly non-local formulation in general case

In the previous section, we have demonstrated that conjugate form (58) of master equations (38) can be written as a symmetric quasilinear system. The key step was a modification of the momentum equation (58a) by *formally* breaking its conservative form via adding to it the linear combination of involution constraints (60). In the case of non curl-free and non divergence-free involution constraints (45) such a step is not valid. However, the symmetrization is still possible but in an extended sense when the PDEs (44) for the new state variables B , Q , R and Ω are added to the master system. Because these quantities coincide with $\nabla \times \mathbf{A}$, $\nabla \cdot \mathbf{h}$, $\nabla \cdot \mathbf{e}$ and $\nabla \times \mathbf{w}$ accordingly at each time instant, the extended system should be treated as a *weakly non-local* extension system of the original Eulerian SHTC equations (38). Here, we follow the idea presented in [21] and for the sake of simplicity, we demonstrate the symmetrization algorithm for the complimentary couple (w_i, σ) , or in fact, for their conjugate couple (η_i, θ) . As will be shown in the example section 6, these equations constitute an important class of equations based on which heat and mass transfer hyperbolic models are developed within the both frameworks, the SHTC and GENERIC formalism. All the steps of the algorithm then can be applied for symmetrizing the PDEs for the couples (m_i, A_{ij}) and (e_i, h_i) and eventually the entire system (38). Thus, we consider the following system for the (w_i, σ) -pair which is coupled with the evolution PDE for the vector field Ω_i from which we expect to be equal to $\Omega_i = \varepsilon_{ijk} \partial_j w_k$

$$\partial_t m_i + \partial_k (m_i v_k + \delta_{ik} (\rho E_\rho + \sigma E_\sigma + m_l E_{m_l} + \Omega_l E_{\Omega_l} - E) - \Omega_k E_{\Omega_i} + w_i E_{w_k}) = 0, \quad (64a)$$

$$\partial_t w_k + \partial_k (w_a v_a + E_\sigma) + v_j (\partial_j w_k - \partial_k w_j) = 0, \quad (64b)$$

$$\partial_t \sigma + \partial_k (\sigma v_k + E_{w_k}) = 0, \quad (64c)$$

$$\partial_t \Omega_i + \partial_k (\Omega_i v_k - \Omega_k v_i) + v_i \partial_k \Omega_k = 0, \quad (64d)$$

$$\partial_t \rho + \partial_k (\rho v_k) = 0. \quad (64e)$$

Here, it is implied that we have extended the set of state variables by adding to it the vector field Ω . However, the addition of a new state variable and its evolution PDE can not be done arbitrary in the SHTC framework because this might be incompatible with the energy conservation law. In order to make it compatible, we have to add the extra stress $-\Omega_k E_{\Omega_i}$ in the momentum flux, otherwise the summation rule (41) does not give the energy conservation. Thus, the energy density and the conjugate potential

$$E = E(m_i, w_i, \sigma, \Omega_i, \rho), \quad L = L(v_i, \eta_i, \theta, \omega_i, r) \quad (65)$$

now also depend on Ω_i and its conjugate $\omega_i = E_{\Omega_i}$ accordingly. The entire system (64) including new equation for Ω_i forms a sub-system of the master system (38). For example, one may clearly notice that the new PDE for Ω_i has the same structure as the PDE (38c) for the field h_i which, in turn, also has the corresponding extra stress $-h_k E_{h_i}$. The following involution constraints are implied as well

$$\frac{\partial}{\partial t} (\varepsilon_{ijk} \partial_j w_k - \Omega_i) = 0, \quad \frac{\partial}{\partial t} (\partial_k \Omega_k) = 0. \quad (66)$$

Therefore, the extended system (67) is still formulated within the SHTC framework.

The conjugate formulation of (64) is

$$\partial_t L_{v_i} + \partial_k ((v_k L)_{v_i} + \eta_k L_{\eta_i} - \delta_{ik} \eta_n L_{\eta_n} - \omega_i L_{\omega_k}) = 0, \quad (67a)$$

$$\partial_t L_{\eta_k} + \partial_k (L_{\eta_a} v_a + \theta) + v_j (\partial_j L_{\eta_k} - \partial_k L_{\eta_j}) = 0, \quad (67b)$$

$$\partial_t L_\theta + \partial_k ((v_k L)_\theta + \eta_k) = 0, \quad (67c)$$

$$\partial_t L_{\omega_i} + \partial_j (L_{\omega_i} v_j - L_{\omega_j} v_i) + v_i \partial_k L_{\omega_k} = 0, \quad (67d)$$

$$\partial_t L_r + \partial_k (v_k L)_r = 0, \quad (67e)$$

while the involution constraints (66) are

$$\frac{\partial}{\partial t} (\nabla \times L_\eta - L_\omega) \equiv 0, \quad \frac{\partial}{\partial t} (\nabla \cdot L_\omega) \equiv 0. \quad (68)$$

The energy conservation law for such an extended formulation now reads as

$$\begin{aligned} \frac{\partial}{\partial t} (r L_r + v_i L_{v_i} + \omega_i L_{\omega_i} + \theta L_\theta + \eta_k L_{\eta_k} - L) + \frac{\partial}{\partial x_k} (v_k (r L_r + v_i L_{v_i} + \omega_i L_{\omega_i} + \theta L_\theta + \eta_k L_{\eta_k} - L) + \\ v_i [(L - \eta_a L_{\eta_a}) \delta_{ik} - \omega_i L_{\omega_k}] + \theta \eta_k) = 0, \end{aligned} \quad (69)$$

which, we emphasize, has the fluxes fully compatible with the spatial differential terms of the governing equations (67), i.e. it can be obtained by the summation rule (41).

Now, we are in position to demonstrate that the conjugate formulation (67) is indeed symmetrizable. At the first step of the symmetrization algorithm we open the brackets in the PDE for L_{η_k} , which, after that, becomes

$$\partial_t L_{\eta_k} + L_{\eta_a} \partial_k v_a + v_j \partial_j L_{\eta_k} + \partial_k \theta = 0, \quad (70)$$

At the second step, we modify this PDE by adding to it $L_{\eta_k} \partial_j v_j - L_{\eta_k} \partial_j v_j \equiv 0$. We have

$$\partial_t L_{\eta_i} + \partial_k (v_k L)_{\eta_i} - L_{\eta_i} \partial_k v_k + L_{\eta_a} \partial_i v_a + \partial_i \theta = 0. \quad (71)$$

At the last step of the algorithm, we formally ‘‘destroy’’ the conservative form of the momentum equation by adding to it the linear combination of the constraints (68)

$$0 \equiv \eta_k (\partial_i L_{\eta_k} - \partial_k L_{\eta_i}) - \varepsilon_{ijk} \eta_j L_{\omega_k} + \omega_i \partial_k L_{\omega_k}. \quad (72)$$

Eventually, we have

$$\partial_t L_{v_i} + \partial_k (v_k L)_{v_i} + L_{\eta_i} \partial_k \eta_k - L_{\eta_a} \partial_i \eta_a - L_{\omega_k} \partial_k \omega_i = \varepsilon_{ijk} \eta_j L_{\omega_k}, \quad (73a)$$

$$\partial_t L_{\eta_i} + \partial_k (v_k L)_{\eta_i} - L_{\eta_i} \partial_k v_k + L_{\eta_a} \partial_i v_a + \partial_i \theta = 0, \quad (73b)$$

$$\partial_t L_\theta + \partial_k (v_k L)_\theta + \partial_k \eta_k = 0, \quad (73c)$$

$$\partial_t L_{\omega_i} + \partial_k (v_k L)_{\omega_i} - L_{\omega_k} \partial_k v_i = 0, \quad (73d)$$

$$\partial_t L_r + \partial_k (v_k L)_r = 0, \quad (73e)$$

which can be written as a symmetric quasilinear system (62) but with an algebraic source term $S = (\varepsilon_{ijk}\eta_j L_{\omega_k}, 0, 0, 0, 0)$:

$$A \frac{\partial \mathbf{p}}{\partial t} + B_k \frac{\partial \mathbf{p}}{\partial x_k} + C_k \frac{\partial \mathbf{p}}{\partial x_k} = S.$$

Indeed, because the non-conservative differential terms have the same structure as in (61), they contribute in a symmetric way into the matrices C_k . The key idea of such a symmetrization is that the source term $\varepsilon_{ijk}\eta_j L_{\omega_k}$ in the momentum equation (73a) is an algebraic term due to the fact that we treat $\nabla \times \mathbf{w}$ as the new independent state variable $\Omega = L_{\omega}$. This source term is therefore treated as a low order term and does not affect the type of the PDEs (i.e. it does not contribute into the structure of the coefficient matrices A , B_k and C_k).

Let us demonstrate the symmetric structure of the matrix C_k . For example, for $k = 1$ the submatrix of C_1 corresponding to the variables v_i , η_i , θ and ω_i is

$$\begin{pmatrix} 0 & 0 & 0 & 0 & -L_{\eta_2} & -L_{\eta_3} & -L_{\omega_1} & 0 & 0 & 0 \\ 0 & 0 & 0 & L_{\eta_2} & 0 & 0 & 0 & -L_{\omega_1} & 0 & 0 \\ 0 & 0 & 0 & L_{\eta_3} & 0 & 0 & 0 & 0 & -L_{\omega_1} & 0 \\ 0 & L_{\eta_2} & L_{\eta_3} & 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ -L_{\eta_2} & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ -L_{\eta_3} & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ -L_{\omega_1} & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & -L_{\omega_1} & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & -L_{\omega_1} & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}. \quad (74)$$

The equations for the pairs (\mathbf{m}, \mathbf{A}) and (\mathbf{h}, \mathbf{e}) are symmetrizable exactly in the same way. For example, in order to symmetrize the equations for the (\mathbf{m}, \mathbf{A}) -pair, one has to add the time evolution (44a) for $\mathbf{B} = \nabla \times \mathbf{A}$ to the system, extend the set of state variables by considering the total energy as $E = E(\mathbf{m}, \mathbf{A}, \mathbf{B}, \rho)$ and repeat the above steps.

In summary, extending the idea from [21], we have demonstrated that in case the constraints $\nabla \times \mathbf{A} = 0$, $\nabla \cdot \mathbf{h} = 0$, $\nabla \cdot \mathbf{e} = 0$ and $\nabla \times \mathbf{w} = 0$ are not met, the Eulerian SHTC equations (38) can be symmetrized only after a weakly non-local extension when the quantities $\nabla \times \mathbf{A}$, $\nabla \cdot \mathbf{h}$, $\nabla \cdot \mathbf{e}$ and $\nabla \times \mathbf{w}$ starts to play the role of the state variables governed by their own time evolutions (44).

4. GENERIC FORM OF SHTC EQUATIONS

After having recalled the SHTC equations, let us turn to the question whether equations (38) can be seen as a Hamiltonian evolution. The answer to that question is affirmative, and the equation are thus fully compatible with GENERIC framework. Moreover, focusing on the Poisson bracket provides an alternative way of derivation and generalization of the SHTC equations.

We, first, demonstrate in sub-section 4.1 that the Lagrangian SHTC equations can be generated by *canonical* Poisson brackets, while the rest of the section is dedicated to the central and more challenging question of whether the Eulerian SHTC system (38) can be seen as a Hamiltonian evolution.

4.1. Lagrangian SHTC system as Hamiltonian evolution

Consider the cotangent bundle of two vector fields $\mathbf{q}(\mathbf{r})$ and $\mathbf{p}(\mathbf{r})$, where the Poisson bracket is canonical, i.e.

$$\{\mathcal{A}, \mathcal{B}\} = \int (\mathcal{A}_{\mathbf{q}} \cdot \mathcal{B}_{\mathbf{p}} - \mathcal{B}_{\mathbf{q}} \cdot \mathcal{A}_{\mathbf{p}}) d\mathbf{r}. \quad (75)$$

Let us first construct a Poisson bracket for the pair (18a), (18b). Let us denote \mathbf{q} as \mathbf{m} and assume that the functionals depend only on gradients of \mathbf{p} , namely on $F_{ij} = -\partial_i p_j$. Poisson bracket (75) then becomes

$$\{\mathcal{A}, \mathcal{B}\} = \int (\mathcal{A}_{m_i} \partial_j \mathcal{B}_{F_{ij}} - \mathcal{B}_{m_i} \partial_j \mathcal{A}_{F_{ij}}) d\mathbf{r}, \quad (76)$$

which generates the first couple of evolution equations (18).

Let us now denote \mathbf{q} as \mathbf{e} and assume that the functionals depend only on curl of \mathbf{p} , namely on $h_i = -\varepsilon_{ijk} \partial_j p_k$. Poisson bracket (75) then becomes

$$\{\mathcal{A}, \mathcal{B}\} = \int (\mathcal{A}_{e_i} \varepsilon_{ijk} \partial_j \mathcal{B}_{h_k} - \mathcal{B}_{e_i} \varepsilon_{ijk} \partial_j \mathcal{A}_{h_k}) d\mathbf{r}, \quad (77)$$

which generates the second couple of evolution equations (18).

Finally, let us denote \mathbf{q} as \mathbf{w} and assume that the functionals depend only on divergence of \mathbf{p} , namely on $s = \partial_j p_j$. Poisson bracket (75) then becomes

$$\{\mathcal{A}, \mathcal{B}\} = \int (\mathcal{A}_{w_i} (-\partial_i \mathcal{B}_s) - \mathcal{B}_{w_i} (-\partial_i \mathcal{A}_s)) d\mathbf{r}, \quad (78)$$

which generates the last couple of evolution equations (18).

The Lagrangian STHC evolution equations (18) can be thus seen as canonical Hamiltonian evolution of a triplet of cotangent bundles, Poisson bracket of which is the sum of brackets (76), (77) and (78). The triplet of operations, namely the rotationally invariant differential operators gradient, curl and divergence, forms a sort of complete set of operations in three-dimensional space, see p. 181 of book [25] and [30].

4.2. Hydrodynamics of ideal fluids

Let us start with the Hamiltonian formulation of the hydrodynamics of ideal fluids. The Poisson bracket for Euler equations of ideal fluids, i.e. for fields (\mathbf{m}, ρ, s) , is (e.g. see [16, 2, 64, 45, 68, 73])

$$\{\mathcal{A}, \mathcal{B}\}^{(Euler)} = \int \left[m_i (\mathcal{B}_{m_i} \partial_j \mathcal{A}_{m_i} - \mathcal{A}_{m_i} \partial_j \mathcal{B}_{m_i}) + \rho (\mathcal{B}_{m_i} \partial_i \mathcal{A}_\rho - \mathcal{A}_{m_i} \partial_i \mathcal{B}_\rho) + s (\mathcal{B}_{m_i} \partial_i \mathcal{A}_s - \mathcal{A}_{m_i} \partial_i \mathcal{B}_s) \right] d\mathbf{r}, \quad (79)$$

where $\mathcal{A}(\mathbf{m}, \rho, s)$ and $\mathcal{B}(\mathbf{m}, \rho, s)$ are two arbitrary functionals and \mathcal{A}_ρ , \mathcal{B}_ρ , \mathcal{A}_{m_i} , \mathcal{B}_{m_i} , etc. are the functional derivatives, i.e. $\mathcal{A}_\rho = \frac{\delta \mathcal{A}}{\delta \rho}$, $\mathcal{A}_{m_i} = \frac{\delta \mathcal{A}}{\delta m_i}$, etc.

The Poisson bracket expresses kinematics of the state variables, which are the fields $\mathbf{x} = (\mathbf{m}, \rho, s)$. Taking an arbitrary functional $\mathcal{A}(\mathbf{m}, \rho, s)$, evolution of that functional is given by the Poisson bracket

$$\frac{\partial \mathcal{A}}{\partial t} = \{\mathcal{A}, \mathcal{H}\}^{(Euler)}, \quad (80)$$

together with the Hamiltonian

$$\mathcal{H}(\mathbf{m}, \rho, s) = \int E(\mathbf{m}, \rho, s) d\mathbf{r}, \quad (81)$$

where the integrand $E(\mathbf{m}, \rho, s)$ is the total energy potential. Poisson bracket and the Hamiltonian thus generate reversible evolution of functionals of state variables, and the evolution equations for the fields (\mathbf{m}, ρ, s) are

$$\frac{\partial m_i}{\partial t} = \{m_i, \mathcal{H}\}^{(Euler)}, \quad \frac{\partial \rho}{\partial t} = \{\rho, \mathcal{H}\}^{(Euler)}, \quad \frac{\partial s}{\partial t} = \{s, \mathcal{H}\}^{(Euler)}. \quad (82)$$

Alternatively, evolution of a functional $\mathcal{A}(\mathbf{x}) = \int A(\mathbf{x})d\mathbf{r}$ can be given by

$$\frac{\partial \mathcal{A}}{\partial t} = \int \frac{\delta \mathcal{A}}{\delta x^i} \frac{\partial x^i}{\partial t} d\mathbf{r}, \quad (83)$$

and hence, by collecting terms multiplying derivatives of the functional \mathcal{A} in the Poisson bracket, i.e. by rewriting the bracket as

$$\{\mathcal{A}, \mathcal{H}\}^{(Euler)} = \int [\mathcal{A}_{m_i} \cdot (\dots) + \mathcal{A}_\rho \cdot (\dots) + \mathcal{A}_s \cdot (\dots)] d\mathbf{r}, \quad (84)$$

one can directly read the evolution equations of the state variables denoted as “...” in (84).

Eventually, by or using (84), one can obtain the Euler equations of compressible hydrodynamics

$$\frac{\partial m_i}{\partial t} = -\partial_j(m_i E_{m_j}) - \rho \partial_i E_\rho - m_j \partial_i E_{m_j} - s \partial_i E_s = -\partial_j(m_i E_{m_j}) - \partial_i p \quad (85a)$$

$$\frac{\partial \rho}{\partial t} = -\partial_i(\rho E_{m_i}) \quad (85b)$$

$$\frac{\partial s}{\partial t} = -\partial_i(s E_{m_i}), \quad (85c)$$

where the pressure is identified as

$$p = \rho E_\rho + s E_s + m_i E_{m_i} - E. \quad (85d)$$

This construction of pressure is general within the context of Hamiltonian dynamics and SHTC formalism. Compressible Euler equations can be thus seen as Hamiltonian evolution.

4.3. Distortion matrix

The goal of this section is to formulate the Poisson brackets generating the time evolutions (38a), (38b) and (38g) for the momentum, distortion and the mass density fields of the SHTC formulation, and hence to demonstrate that these SHTC equations are compatible with GENERIC. The distortion field is a very important concept, which is used to describe elastic and inelastic deformation of continuum (viscous flows, elastoplastic deformations) in the SHTC theory. Before discussing the dynamics of distortion matrix, let us first consider dynamics of labels advected by the fluid particles.

4.3.1 Dynamics of labels

When describing dynamics of solids, the usual approach is to formulate the evolution equations in the Lagrangian coordinates, since dynamics of the material points is then simply related to classical Newtonian mechanics. Since the balance equations (of energy, momentum, mass and angular momentum) are formulated in the Eulerian coordinates, one needs to use the mapping from the Lagrangian coordinates to the Eulerian coordinates. The classical treatment of elasticity, e.g. [52], is formulated this way. After the transformation to the Eulerian frame, the standard equations consist of balance laws (conservation of mass, momentum and entropy (or energy)) equipped with an equation expressing advection of the Lagrangian coordinates of the particles, i.e. *labels* of the particles.

However, when the material undergoes plastic deformations or even chemical reactions or phase transitions, where the material points disappear and are created, the mapping from the Lagrangian coordinates to the Eulerian coordinates loses its physical significance. The reference configuration itself loses its physical significance and it plays merely the role of an initial condition.

Because the balance equations still keep their physical meaning even when the reference configuration does not, it seems to be advantageous to formulate the evolution equations of solids purely within the Eulerian coordinates.

GENERIC provides a way for such a construction [46], and the result turns out to be compatible with the Godunov-Romenski approach [40, 39, 31, 4, 23, 77, 17, 78]. Let us now develop the GENERIC way.

Note that dynamics of solids has already been formulated within GENERIC. For example in [55] evolution equation for the deformation tensor is generated by a Poisson bracket which is constructed by using the requirements of antisymmetry and momentum conservation.

The aim here is to find evolution equations for solids capturing balance of mass, momentum and entropy (or energy) as well as dynamics of individual points of the continuum (particle labels). Therefore, the state variables will be density ρ , momentum density \mathbf{u} , entropy density s and volume density of the labels \mathbf{b} . The volume density of labels is a vector field that expresses where the material at given Eulerian position belongs.

The reversible evolution of the state variables is required to be Hamiltonian, i.e. generated by a Poisson bracket. Poisson bracket governing evolution of variables ρ , \mathbf{m} and s is the standard hydrodynamic Poisson bracket (79). The bracket expresses that mass density, momentum density and entropy density are passively advected by the momentum density. Therefore, it is clear how to add an another passively advected field – the volume density of labels \mathbf{b} . The overall bracket becomes

$$\{\mathcal{A}, \mathcal{B}\}^{(Euler+Labels)} = \int \left[m_i (\partial_j \mathcal{A}_{m_i} \mathcal{B}_{m_j} - \partial_j \mathcal{B}_{m_i} \mathcal{A}_{m_j}) + \rho (\partial_i \mathcal{A}_\rho \mathcal{B}_{m_i} - \partial_i \mathcal{B}_\rho \mathcal{A}_{m_i}) + s (\partial_i \mathcal{A}_s \mathcal{B}_{m_i} - \partial_i \mathcal{B}_s \mathcal{A}_{m_i}) + b_i (\partial_j \mathcal{A}_{b_i} \mathcal{B}_{m_j} - \partial_j \mathcal{B}_{b_i} \mathcal{A}_{m_j}) \right] d\mathbf{r} \quad (86)$$

for any two functionals \mathcal{A} and \mathcal{B} of the state variables. This is the Poisson bracket expressing kinematics of hydrodynamic fields and volume density of labels.

However, each label should correspond to a concrete particle of the material rather than to a volume of the material, and thus we perform transformation from \mathbf{b} to

$$\mathbf{a} \stackrel{def}{=} \frac{\mathbf{b}}{\rho}, \quad (87)$$

which is the mass-density of labels. Bracket (86) then transforms (by plugging in functionals of type $\mathcal{A}(\mathbf{m}, \mathbf{b}(\rho, \mathbf{a}), \rho, s)$) to

$$\{\mathcal{A}, \mathcal{B}\}^{(Lin)} = \{\mathcal{A}, \mathcal{B}\}^{(Euler)} + \int \partial_j a_i (\mathcal{A}_{m_j} \mathcal{B}_{a_i} - \mathcal{B}_{m_j} \mathcal{A}_{a_i}) d\mathbf{r}. \quad (88)$$

This Poisson bracket was first introduced in [46], where it was referred to as the Lin Poisson bracket. The evolution equations implied by bracket (88) are

$$\frac{\partial m_i}{\partial t} = -\partial_j (m_i E_{m_j}) - \rho \partial_i E_\rho - m_j \partial_i E_{m_j} - s \partial_i E_s + E_{a_j} \partial_j a_i \quad (89a)$$

$$\frac{\partial \rho}{\partial t} = -\partial_i (\rho E_{m_i}) \quad (89b)$$

$$\frac{\partial s}{\partial t} = -\partial_i (s E_{m_i}) \quad (89c)$$

$$\frac{\partial a_i}{\partial t} = -E_{m_j} \partial_j a_i. \quad (89d)$$

When the field \mathbf{a} is interpreted as the Lagrangian coordinate of a particle, the evolution equations are the same as the standard equations of continuum mechanics transformed into the Eulerian frame.

In order to write down the equations explicitly, we have to choose an energy potential $E(\mathbf{m}, \mathbf{a}, \rho, s)$. Besides the kinetic energy, which is of course $m^2/(2\rho)$, and internal energy, which is a function of ρ and s , the deformation energy should be taken into account. This energy can not depend on the field \mathbf{a} itself but only on gradient of the

field due to the invariance with respect to translations of the whole body. Thus, the energy can be taken as

$$E = E(\mathbf{m}, \nabla \mathbf{a}, \rho, s). \quad (90)$$

Derivative of energy with respect to the field of labels \mathbf{a} is then

$$E_{a_i} = -\partial_j \frac{\partial E}{\partial \partial_j a_i} = -\partial_j E_{A_{ij}}, \quad A_{ij} = \partial_j a_i, \quad (91)$$

and the evolution equation for the field of labels becomes

$$\frac{\partial \mathbf{a}}{\partial t} + \mathbf{v} \cdot \nabla \mathbf{a} = 0, \quad (92)$$

where velocity \mathbf{v} is identified with $E_{\mathbf{m}}$. The evolution equation for momentum density now reads

$$\frac{\partial m_i}{\partial t} = -\partial_j (m_i E_{m_j}) - \rho \partial_i E_\rho - m_j \partial E_{m_j} - s \partial_i E_s + \partial_j \sigma_{ij}, \quad (93)$$

where the stress tensor generated by the labels is

$$\sigma_{ij} = E \delta_{ij} - \partial_i a_l \frac{\partial E}{\partial \partial_j a_l} = E \delta_{ij} - A_{li} E_{A_{lj}}. \quad (94)$$

The rest of evolution equations (89) is as in classical hydrodynamics. We have thus explicitly written down the evolution equations for the hydrodynamic variables with an advected field of particle labels. These equations can be regarded as standard equations for continuum mechanics in the Eulerian frame.

4.3.2 Differential geometric derivation

In this section we provide an alternative derivation of Poisson bracket (88) by means of differential geometry. More specifically, we employ the formulas for matched pairs and semidirect products derived in [22], in particular formula (43) from that paper,

$$\begin{aligned} \{\mathcal{A}, \mathcal{B}\}^{(\ell^* \times T^*V)} &= \{\mathcal{A}, \mathcal{B}\}^{(\ell^*)} + \{\mathcal{A}, \mathcal{B}\}^{(T^*V)} \\ &\quad + \langle \mathcal{A}_v, \mathcal{B}_\zeta \triangleright v \rangle - \langle \mathcal{B}_v, \mathcal{A}_\zeta \triangleright v \rangle \\ &\quad + \langle \alpha, \mathcal{A}_\zeta \triangleright \mathcal{B}_\alpha \rangle - \langle \alpha, \mathcal{B}_\zeta \triangleright \mathcal{A}_\alpha \rangle. \end{aligned} \quad (95)$$

The bracket expresses kinematics of a Lie algebra dual ℓ^* , $\zeta \in \ell^*$, coupled with a cotangent bundle T^*V , $(v, \alpha) \in T^*V$. The coupling is just one-sided, the Lie algebra dual acts on the cotangent bundle. The $\langle \bullet, \bullet \rangle$ brackets denote a scalar product, here standard L^2 scalar product, i.e. simple integration. Finally, for example action $\mathcal{B}_\zeta \triangleright v$ stands for $-\mathcal{L}_{\mathcal{B}_\zeta} v$, \mathcal{L} being the Lie derivative, and the other actions are interpreted analogically. This is the meaning of bracket (95).

Let us suppose that the functionals do not depend on α , i.e. elements of the dual to the vector space V . Bracket (95) then reduces to

$$\{\mathcal{A}, \mathcal{B}\}^{(\ell^* \times V)} = \{\mathcal{A}, \mathcal{B}\}^{(\ell^*)} + \langle \mathcal{A}_v, \mathcal{B}_\zeta \triangleright v \rangle - \langle \mathcal{B}_v, \mathcal{A}_\zeta \triangleright v \rangle. \quad (96)$$

The hydrodynamic fields (ρ, \mathbf{m}) can be seen as elements of a Lie algebra dual ℓ^* , see e.g. [64] or [73]. Consider now

the vector field of labels interpreted as an element of a vector space V . Bracket (96) then becomes

$$\begin{aligned} \{\mathcal{A}, \mathcal{B}\}^{(\ell^* \times V)} &= \int \rho (\partial_i \mathcal{A}_\rho \mathcal{B}_{m_i} - \partial_i \mathcal{B}_\rho \mathcal{A}_{m_i}) d\mathbf{r} + \int m_i (\partial_j \mathcal{A}_{m_i} \mathcal{B}_{m_j} - \partial_j \mathcal{B}_{m_i} \mathcal{A}_{m_j}) d\mathbf{r} \\ &\quad + \int (\mathcal{B}_{a_i} \mathcal{A}_{m_j} \partial_j a_i - \mathcal{A}_{a_i} \mathcal{B}_{m_j} \partial_j a_i) d\mathbf{r}, \end{aligned} \quad (97)$$

which is the same Poisson bracket as bracket (88). We have thus interpreted the Lin bracket in terms of differential geometry.

4.3.3 Dynamics of general distortion matrix

Evolution equations (89) for hydrodynamics with labels are required to be invariant with respect to shift of the field of labels by a constant. To fulfill such a requirement, it was necessary to let the energy depend only on the spatial gradient of the field of labels. Let us now reformulate the dynamics solely in terms of the gradient $\nabla \mathbf{a}$, which is referred to as the distortion matrix (following [39, 42, 41, 77, 17]),

$$A_{ij} = \partial_j a_i, \quad (98)$$

and expresses local variations of the field of labels. This definition of the distortion matrix implies the integrability conditions

$$\partial_k A_{ij} = \partial_j A_{ik} \quad \forall i, j, k, \quad \text{that is} \quad \varepsilon_{ljk} \partial_j A_{ik} = 0 \quad \forall l \quad \iff \quad \nabla \times \mathbf{A} = 0. \quad (99)$$

We first propose a Poisson bracket for the time evolution of the compatible distortion, i.e. for which the integrability conditions are fulfilled and which is thus the gradient of the field of labels. Our main goal, however, is to provide a Poisson bracket for generally incompatible distortion fields, for which $\nabla \times \mathbf{A} \neq 0$. The incompatible distortion field is the most important case because such distortions are used to describe irreversible deformations of viscous fluids and elastoplastic solids [41, 77, 17, 78], see also examples in Section 6, and non-Newtonian fluids, see Sec. 4.3.4.

Transforming Poisson bracket (86) into variables $(\mathbf{m}, \mathbf{A}, \rho, s)$, which also means that the two functionals depend only on the new variables, leads to the Poisson bracket of hydrodynamics with the distortion matrix,

$$\{\mathcal{A}, \mathcal{B}\}^{(Euler+A)} = \{\mathcal{A}, \mathcal{B}\}^{(Euler)} + \int A_{ij} (\mathcal{B}_{m_j} \partial_k \mathcal{A}_{A_{ik}} - \mathcal{A}_{m_j} \partial_k \mathcal{B}_{A_{ik}}) d\mathbf{r}. \quad (100)$$

This bracket is a Poisson bracket, in particular it fulfills Jacobi identity (checked with program [60]), if and only if the integrability conditions (98) hold. However, it can be expected that irreversible evolution violates these conditions and hence the distortion matrix loses the connection (99) with the field of labels. Indeed, having $\nabla \times \mathbf{A} \neq 0$ also means that integral of \mathbf{A} over a closed loop does not necessarily give zero, which means that there might be a discontinuity in the field of labels – the material undergoes irreversible deformation and identity of the particles (the labels) are altered. We need to extend the Poisson bracket so that it fulfills the Jacobi identity also for the incompatible distortion fields.

The extension should be done in such a way that for curl-free distortion matrices bracket (100) is recovered, which means that the extra terms (to be added to the bracket) should thus be multiplied by $\nabla \times \mathbf{A}$. Moreover, in order to generate reversible evolution, the extra terms should provide additional coupling between \mathbf{A} (even with respect to time reversal) and momentum \mathbf{m} (odd) because \mathbf{m} is the only variable with opposite parity than \mathbf{A} , see Section 3.1.3 and [75]. The new distortion matrix bracket then reads

$$\{\mathcal{A}, \mathcal{B}\}^{(DM)} = \{\mathcal{A}, \mathcal{B}\}^{(Euler+A)} + \int (\partial_k A_{ij} - \partial_j A_{ik}) (\mathcal{A}_{A_{ik}} \mathcal{B}_{m_j} - \mathcal{B}_{A_{ik}} \mathcal{A}_{m_j}) d\mathbf{r}, \quad (101)$$

which fulfills the Jacobi identity unconditionally (checked with program [60]). Note that the extra terms can not be multiplied by any number or function without violating the Jacobi identity, which makes the choice of the extension

unique. This Poisson bracket leads to evolution equations

$$\frac{\partial m_i}{\partial t} = -\partial_j(m_i E_{m_j}) - \rho \partial_i E_\rho - m_j \partial_i E_{m_j} - s \partial_i E_s - A_{ji} \partial_k E_{A_{jk}} - (\partial_k A_{ji} - \partial_i A_{jk}) E_{A_{jk}} \quad (102a)$$

$$= -\partial_i p - \partial_j(m_i E_{m_j}) + \partial_j \sigma_{ij}$$

$$\frac{\partial A_{ik}}{\partial t} = -\partial_k(A_{ij} E_{m_j}) + (\partial_k A_{ij} - \partial_j A_{ik}) E_{m_i}, \quad (102b)$$

$$\frac{\partial \rho}{\partial t} = -\partial_i(\rho E_{m_i}) \quad (102c)$$

$$\frac{\partial s}{\partial t} = -\partial_i(s E_{m_i}). \quad (102d)$$

where the generalized pressure p and extra stress tensor σ_{ij} were identified as (taken $\mathcal{H} = \int E(\mathbf{m}, \mathbf{A}, \rho, s) d\mathbf{r}$)

$$p = \rho E_\rho + s E_s + m_i E_{m_i} + A_{ij} E_{A_{ij}} - E, \quad \sigma_{ij} = -A_{ki} E_{A_{kj}} + A_{kl} E_{A_{kl}} \delta_{ij}. \quad (103)$$

Equation (102) are the same as the Eulerian SHTC equation (38a), (38b) (recall that equations (102c) and (102d) have the same structure and thus are represented by one equation (38g) in system (38)). It should be borne in mind that the distortion matrix is no longer required to have zero curl. Also, evolution equations (102) are Galilean invariant, as they can be rewritten in terms of material derivatives $\partial_t + \mathbf{v} \cdot \nabla$ and gradients of velocity.

Kinematics of the field of labels is the same as kinematics of Lagrangian coordinates of the fluid particles in the Eulerian frame. Therefore, kinematics of the distortion matrix expressed by Poisson bracket (100), where only matrices with zero curl are allowed, is the same as kinematics of inverse of the deformation tensor, which is standard in continuum mechanics. The extended Poisson bracket (101), however, allows for distortion matrices with non-zero curl, and the resulting kinematics thus lies outside of the scope of classical continuum mechanics because the Lagrangian coordinates can not be reconstructed uniquely.

In summary, the hydrodynamic Poisson bracket was first enriched with the field of labels, kinematics of which is pure advection. That is the setting of classical continuum mechanics in the Eulerian frame. Due to the required invariance with respect to spatial shifts, only spatial gradients of the field of labels (the distortion matrix) were kept among the state variables, and a new Poisson bracket followed. The bracket fulfilled Jacobi identity only for curl-free distortion matrices, and to satisfy the identity also for distortion matrices with non-zero curl, it was necessary to add (in a unique way) extra terms to the Poisson bracket. The Poisson bracket expressing kinematics of non curl-free distortion matrices has then been found, which generates the same evolution equations (38a), (38b) and (38g) of the SHTC framework. To our best knowledge, these results are new.

4.3.4 Dynamics of the left Cauchy-Green (or Finger) tensor

One of the main motivations for developing GENERIC framework was modeling of rheological properties of complex fluids [50, 72], where it is popular to include a symmetric positive definite tensor into the set of state variables, e.g. the conformation tensor. Here we choose the left Cauchy-Green tensor (or Finger tensor), which has been shown useful when describing complex fluids [80, 63, 54].

The goal of this section is to discuss the relation of a tensor \mathcal{B} , which becomes the left Cauchy-Green tensor (or Finger tensor) in the case of curl-free distortion field, e.g. in the absence of dissipation. Indeed, the distortion matrix is identical to the inverse deformation gradient if dissipation is absent [41]. In particular, we demonstrate that a Poisson bracket for the Left Cauchy-Green tensor can be derived from the Poisson bracket (101), which generates reversible evolution of the hydrodynamic fields and the general distortion matrix.

The left Cauchy-Green tensor is defined as

$$\mathcal{B}_{ij} = A_{ik}^{-1} A_{jk}^{-1}. \quad (104)$$

Derivative of a functional $\mathcal{A}(\mathbf{m}, \mathcal{B}, \rho, s)$ with respect to the distortion matrix then becomes

$$\mathcal{A}_{A_{mn}} = \frac{\delta \mathcal{A}}{\delta A_{mn}} = \frac{\delta \mathcal{A}}{\delta \mathcal{B}_{ij}} \frac{\partial \mathcal{B}_{ij}}{\partial A_{mn}}. \quad (105)$$

By differentiating definition (104), we can obtain that

$$\frac{\partial \mathcal{B}_{ij}}{\partial A_{mn}} = -A_{im}^{-1} \mathcal{B}_{nj} - A_{jm}^{-1} \mathcal{B}_{in}, \quad (106)$$

from which it follows that

$$\mathcal{A}_{A_{ik}} = -\mathcal{A}_{\mathcal{B}_{mn}} A_{mi}^{-1} \mathcal{B}_{kn} - \mathcal{A}_{\mathcal{B}_{mn}} A_{ni}^{-1} \mathcal{B}_{mk}. \quad (107)$$

Substituting this last relation into equation (101) leads to the following Poisson bracket

$$\begin{aligned} \{\mathcal{A}, \mathcal{B}\}^{(Finger)} &= \{\mathcal{A}, \mathcal{B}\}^{(Euler)} + \int \mathcal{B}_{ik} \left(\partial_j \mathcal{A}_{\mathcal{B}_{ik}} \mathcal{B}_{mj} - \partial_j \mathcal{B}_{\mathcal{B}_{ik}} \mathcal{A}_{mj} \right) d\mathbf{r} \\ &+ \int \mathcal{B}_{km} \left(\left(\mathcal{A}_{\mathcal{B}_{mj}} + \mathcal{A}_{\mathcal{B}_{jm}} \right) \partial_k \mathcal{B}_{mj} - \left(\mathcal{B}_{\mathcal{B}_{mj}} + \mathcal{B}_{\mathcal{B}_{jm}} \right) \partial_k \mathcal{A}_{mj} \right) d\mathbf{r} \\ &+ \int \mathcal{B}_{ik} \left(\mathcal{A}_{\mathcal{B}_{ik}} \partial_j \mathcal{B}_{mj} - \mathcal{B}_{\mathcal{B}_{ik}} \partial_j \mathcal{A}_{mj} \right) d\mathbf{r}, \end{aligned} \quad (108)$$

which generates the following reversible evolution equations:

$$\frac{\partial m_i}{\partial t} = -\partial_j (m_i E_{mj}) - \rho \partial_i E_\rho - m_j \partial_i E_{m_j} - s \partial_i E_s - \mathcal{B}_{jk} \partial_i E_{\mathcal{B}_{jk}} + \partial_k (\mathcal{B}_{km} (E_{\mathcal{B}_{mi}} + E_{\mathcal{B}_{im}})) + \partial_i (\mathcal{B}_{jk} E_{\mathcal{B}_{jk}}) \quad (109a)$$

$$= -\partial_j (m_i E_{mj}) - \rho \partial_i E_\rho - m_j \partial_i E_{m_j} - s \partial_i E_s + E_{\mathcal{B}_{jk}} \partial_i \mathcal{B}_{jk} + \partial_k (\mathcal{B}_{km} (E_{\mathcal{B}_{mi}} + E_{\mathcal{B}_{im}})) \quad (109b)$$

$$\frac{\partial \mathcal{B}_{ij}}{\partial t} = -E_{u_k} \partial_k \mathcal{B}_{ij} + \mathcal{B}_{ki} \partial_k E_{u_j} + \mathcal{B}_{kj} \partial_k E_{u_i}, \quad (109c)$$

$$\frac{\partial \rho}{\partial t} = -\partial_i (\rho E_{m_i}) \quad (109d)$$

$$\frac{\partial s}{\partial t} = -\partial_i (s E_{m_i}). \quad (109e)$$

This means that the left Cauchy-Green tensor has zero upper-convected derivative unless dissipation is introduced.

Note also that kinematics of the tensor \mathcal{B} can be thus seen as a consequence of kinematics of the distortion matrix (101), which in turn lies outside of the scope of classical continuum solid mechanics due to the possibility of non-zero curl of the matrix.

4.4. Electrodynamics of slowly moving medium

The goal of this section is to demonstrate that SHTC equations (38a), (38c), (38d) and (38g) governing the dynamics of the fields $(\mathbf{m}, \mathbf{h}, \mathbf{e}, \rho, s)$ has the structure which is compatible with GENERIC, i.e. these equations can be generated by a Poisson bracket. However, as usually in GENERIC, while constructing the Poisson brackets, we have to be more specific and thus the fields $(\mathbf{m}, \mathbf{h}, \mathbf{e}, \rho, s)$ have to be endowed with certain physical meanings. For example, the electrodynamics of slowly moving continuous media is formulated in [18] within the SHTC framework. In this

paper, the fields (\mathbf{e}, \mathbf{h}) were recognized as

$$\mathbf{e} = \epsilon(\mathbf{E} + \mathbf{v} \times (\mathbf{v} \times \mathbf{E})), \quad \mathbf{h} = \mathbf{B} + \mathbf{v} \times (\mathbf{v} \times \mathbf{B}), \quad (110)$$

while the conjugate fields are

$$E_e = \mathbf{E} + \mathbf{v} \times \mathbf{B}, \quad E_h = \mathbf{B} - \epsilon \mathbf{v} \times \mathbf{E}, \quad (111)$$

where ϵ is the electric permittivity, \mathbf{E} and \mathbf{B} are the electric and magnetic fields in the laboratory frame and the speed of light is assumed to be one.

Thus, if the motion is absent ($\mathbf{v} = 0$) then the standard electric displacement field \mathbf{D} and magnetic field are recovered

$$\mathbf{e} = \epsilon \mathbf{E} = \mathbf{D}, \quad \mathbf{h} = \mathbf{B}. \quad (112)$$

It has appeared that a Poisson bracket governing evolution of exactly the fields (\mathbf{D}, \mathbf{B}) and fields \mathbf{m} , ρ and s has been referred to as the *electromagnetohydrodynamic* Poisson bracket (EMHD) in [53] or in [22] (see equation (66) there), and it reads⁶

$$\begin{aligned} \{\mathcal{A}, \mathcal{B}\}^{(EMHD)} &= \{\mathcal{A}, \mathcal{B}\}^{(Euler)} + \{\mathcal{A}, \mathcal{B}\}^{(EM)} \\ &+ \int D_i (\mathcal{B}_{m_j} \partial_j \mathcal{A}_{D_i} - \mathcal{A}_{m_j} \partial_j \mathcal{B}_{D_i}) d\mathbf{r} + \int \partial_j D_j (\mathcal{A}_{m_i} \mathcal{B}_{D_i} - \mathcal{B}_{m_i} \mathcal{A}_{D_i}) d\mathbf{r} + \int (\mathcal{A}_{m_i} D_j \partial_j \mathcal{B}_{D_i} - \mathcal{B}_{m_i} D_j \partial_j \mathcal{A}_{D_i}) d\mathbf{r} \\ &+ \int B_i (\mathcal{B}_{m_j} \partial_j \mathcal{A}_{B_i} - \mathcal{A}_{m_j} \partial_j \mathcal{B}_{B_i}) d\mathbf{r} + \int \partial_j B_j (\mathcal{A}_{m_i} \mathcal{B}_{B_i} - \mathcal{B}_{m_i} \mathcal{A}_{B_i}) d\mathbf{r} + \int (\mathcal{A}_{m_i} B_j \partial_j \mathcal{B}_{B_i} - \mathcal{B}_{m_i} B_j \partial_j \mathcal{A}_{B_i}) d\mathbf{r}. \end{aligned} \quad (113)$$

Note that the momentum density \mathbf{m} is the coupled matter-field momentum in [22] and [18]. This Poisson bracket contains also the electromagnetic Poisson bracket,

$$\{\mathcal{A}, \mathcal{B}\}^{(EM)} = \int (\mathcal{A}_{D_i} \varepsilon_{ijk} \partial_j \mathcal{B}_{B_k} - \mathcal{B}_{D_i} \varepsilon_{ijk} \partial_j \mathcal{A}_{B_k}) d\mathbf{r}, \quad (114)$$

which generates the Maxwell equations.

By taking a Hamiltonian $\mathcal{H} = \int E(\mathbf{m}, \mathbf{B}, \mathbf{D}, \rho, s) d\mathbf{r}$ and rewriting bracket (113) in a form similar to (84), one directly obtains the following evolution equations

$$\frac{\partial m_i}{\partial t} = -\partial_j (m_i E_{m_j}) - \rho \partial_i E_\rho - m_j \partial_i E_{m_j} - s \partial_i E_s - D_j \partial_i E_{D_j} - B_j \partial_i E_{B_j} + \partial_j (D_j E_{D_i} + B_j E_{B_i}), \quad (115a)$$

$$\frac{\partial B_i}{\partial t} = -\partial_j (B_i E_{m_j} - E_{m_i} B_j + \varepsilon_{ijk} \partial_j E_{D_k}) - E_{m_i} \partial_j B_j, \quad (115b)$$

$$\frac{\partial D_i}{\partial t} = -\partial_j (D_i E_{m_j} - E_{m_i} D_j - \varepsilon_{ijk} \partial_j E_{B_k}) - E_{m_i} \partial_j D_j, \quad (115c)$$

$$\frac{\partial \rho}{\partial t} = -\partial_i (\rho E_{m_i}), \quad (115d)$$

$$\frac{\partial s}{\partial t} = -\partial_i (s E_{m_i}), \quad (115e)$$

where pressure p is defined as

$$p = \rho E_\rho + s E_s + m_i E_{m_i} + D_i E_{D_i} + B_i E_{B_i} - E, \quad (115f)$$

and $E(\mathbf{m}, \mathbf{B}, \mathbf{D}, \rho, s)$ being the total energy density. These evolution equations are also equipped with two con-

⁶ Bracket (113) can be derived from Eq. (65) in paper [22] by rewriting action \triangleright as minus Lie derivative when interpreting fields \mathbf{D} and \mathbf{B} as vector fields, i.e. H_D and H_B as covector fields (in notations of [22]). By applying constraints (115g), bracket (113) becomes bracket (66) from paper [22]. Constraints (115g), which are two of four Maxwell equations, can be seen either as results of gauge invariance of the theory of electromagnetism coupled with matter [64] or as conditions under which Poisson bracket (66) from paper [22] becomes a projection from kinetic theory coupled with matter, equation (58) in [22].

straints imposed on the Poisson bracket

$$\operatorname{div}(\mathbf{D}) = \frac{ze}{\epsilon}\rho, \quad \operatorname{div}(\mathbf{B}) = 0, \quad (115g)$$

where z is number of elementary charges per particle and e is the elementary charge.

By subtracting and adding $\partial_i E$ to the momentum equation, it can be transformed exactly into the SHTC momentum conservation law (38a), while the equations for \mathbf{B} and \mathbf{D} already has exactly the structure of equations (38c) and (38d) accordingly. Thus, the entire GENERIC formulation and the SHTC formulation for electrodynamics of moving medium are fully compatible.

4.5. Ballistic transport

The goal of this section is to present a Poisson bracket which generates equations (38a), (38f) and (38e), and thus to prove their compatibility with GENERIC. In applications, these equations can be used for modeling of heat and mass transfer if equipped with appropriate dissipative algebraic source terms (see Section 5 and 6 for details). However, in this section we shall use the terminology for heat conduction and we shall call the scalar field s the entropy density while the vector field \mathbf{w} as the conjugate entropy flux. Absence of the dissipative terms, i.e. absence of the interaction between the mass or heat carriers and the medium, corresponds to the most non-equilibrium state, *ballistic transport*, which explains the title of this Section.

4.5.1 The SHTC-compatible bracket

Consider first a rigid heat-conducting body. We wish to prescribe a field of entropy density s expressing local thermodynamic state of the body. Let us now address dynamics of the field. Taking a Lagrangian $\Lambda(s, \dot{s})$, the Legendre transformation introduces the corresponding Hamiltonian,

$$\frac{\delta}{\delta \dot{s}} \left(-\Lambda(s, \dot{s}) + \int \psi(\mathbf{r}) \dot{s}(\mathbf{r}) \right) d\mathbf{r} = 0 \quad \Leftrightarrow \quad \Lambda_{\dot{s}} = \psi, \quad (116)$$

and the conjugate scalar field $\psi(\mathbf{r}, t)$. The couple (ψ, s) is equipped with the canonical Poisson bracket

$$\{\mathcal{A}, \mathcal{B}\}^{(\psi, s)} = \int (\mathcal{A}_{\psi} \mathcal{B}_s - \mathcal{B}_{\psi} \mathcal{A}_s) d\mathbf{r}. \quad (117)$$

However, the aim is to work with a vector (covariant) field that plays the role of conjugate entropy flux. Therefore, taking functionals dependent only on $\mathbf{w} = -\nabla\psi$, the canonical bracket becomes

$$\{\mathcal{A}, \mathcal{B}\}^{(\mathbf{w}, s)} = \int (\partial_k \mathcal{A}_s \mathcal{B}_{w_k} - \partial_k \mathcal{B}_s \mathcal{A}_{w_k}) d\mathbf{r}, \quad (118)$$

evolution equations generated by bracket (118) are

$$\frac{\partial s}{\partial t} = -\partial_k E_{w_k}, \quad (119a)$$

$$\frac{\partial w_k}{\partial t} = -\partial_k E_s, \quad (119b)$$

equipped with the condition that \mathbf{w} is a potential vector field, or that

$$\partial_i w_j = \partial_j w_i. \quad (119c)$$

From (119a) it is clear why w is referred to as the conjugate entropy flux. Equations (119) express evolution of entropy within rigid heat conductors.

How can one couple the dynamics of heat with hydrodynamics, i.e. with transport of matter? As in section 4.3.2, the hydrodynamic bracket for (m, ρ) can be extended to describe also action of hydrodynamics on elements of a vector space V , $\psi \in V$, and the resulting Poisson bracket would be analogical to bracket (97) with a_i replaced by ψ . Here, however, we will consider coupling between the hydrodynamic Lie algebra and a cotangent bundle $(\psi, s) \in T^*V$, kinematics of which is expressed by bracket (95),

$$\{\mathcal{A}, \mathcal{B}\}^{(m, \rho) \times (\psi, s)} = \{\mathcal{A}, \mathcal{B}\}^{(Euler)} + \int (\mathcal{A}_\psi \mathcal{B}_s - \mathcal{A}_s \mathcal{B}_\psi) d\mathbf{r} + \int (\mathcal{B}_\psi \mathcal{A}_{m_j} \partial_j \psi - \mathcal{A}_\psi \mathcal{B}_{m_j} \partial_j \psi) d\mathbf{r}.$$

Taking functionals dependent only on gradients of ψ , denoted by $w_i = -\partial_i \psi$, this last bracket becomes

$$\{\mathcal{A}, \mathcal{B}\}^{(m, \rho) \times (w, s)} = \{\mathcal{A}, \mathcal{B}\}^{(Euler)} + \int (\mathcal{B}_{w_i} \partial_i \mathcal{A}_s - \mathcal{A}_{w_i} \partial_i \mathcal{B}_s) d\mathbf{r} + \int w_j (\partial_i \mathcal{A}_{w_i} \mathcal{B}_{m_j} - \partial_i \mathcal{B}_{w_i} \mathcal{A}_{m_j}) d\mathbf{r}. \quad (120)$$

This Poisson bracket expresses kinematics of variables (m, ρ, w, s) provided w is a potential vector field, i.e. $\nabla \times w = 0$. Indeed, Jacobi identity is fulfilled only when this involution constraint is met (as in the case of the transition from kinematics of labels to kinematics of distortion matrix).

Validity of Jacobi identity for bracket (120) even for fields w with non-zero curl can be achieved by extending the Poisson bracket as in the case of distortion matrix in Section 4.3.3. A new term has to be added to the bracket proportional to $\partial_i w_j - \partial_j w_i$ (the term can not be multiplied by any constant). Moreover, the term has to provide coupling between w , which is an odd variable, and an another odd field, for example with the field m . The only way to extend the bracket (120) is then as follows (the Jacobi identity for this bracket has been checked by program [60])

$$\{\mathcal{A}, \mathcal{B}\}^{(Heat)} = \{\mathcal{A}, \mathcal{B}\}^{(m, \rho) \times (w, s)} + \int (\partial_i w_j - \partial_j w_i) (\mathcal{A}_{w_i} \mathcal{B}_{m_j} - \mathcal{B}_{w_i} \mathcal{A}_{m_j}) d\mathbf{r}, \quad (121)$$

which⁷ after rearranging the terms as in (84), leads to the time evolutions equations

$$\frac{\partial m_i}{\partial t} = -\partial_k (m_i E_{m_k}) - \rho \partial_i E_\rho - s \partial_i E_s - m_k \partial_i E_{m_k} - w_i \partial_k E_{w_k} - E_{w_k} (\partial_a w_i - \partial_i w_k), \quad (122a)$$

$$\frac{\partial w_k}{\partial t} = -\partial_k (w_j E_{m_j} + E_s) - (\partial_j w_k - \partial_k w_j) E_{m_j}, \quad (122b)$$

$$\frac{\partial s}{\partial t} = -\partial_k (s E_{m_k} + E_{w_k}), \quad (122c)$$

$$\frac{\partial \rho}{\partial t} = -\partial_k (\rho E_{m_k}) \quad (122d)$$

which coincide with the Eulerian SHTC equations (38a), (38e), (38f) and (38g).

In summary, Poisson bracket (120) has been derived as a result of coupling between the Lie algebra dual of classical hydrodynamics and the cotangent bundle of dynamics of entropy. The Jacobi identity is valid only for potential vector fields w . Validity of Jacobi identity can be extended to fields w with non-zero curl by extending the bracket by a term coupling fields w and m , and such an extension can be carried out in only one way. The final extended Poisson bracket is bracket (121), which generates the SHTC evolution equations (122) for ballistic transport of the scalar field s .

⁷This bracket was first proposed in [70].

4.5.2 Alternative bracket, Cattaneo hydrodynamics

A Poisson bracket expressing the Cattaneo-like dynamics of heat was proposed in [49],

$$\{\mathcal{A}, \mathcal{B}\}^{(Cat)} = \int \pi_i \left(\partial_j \mathcal{A} \pi_i \mathcal{B} \pi_j - \partial_j \mathcal{B} \pi_i \mathcal{A} \pi_j \right) d\mathbf{r} + \int s \left(\partial_i \mathcal{A} s \mathcal{B} \pi_i - \partial_i \mathcal{A} s \mathcal{B} \pi_i \right) d\mathbf{r}, \quad (123)$$

where π is a momentum flux associated with transport of entropy. There are at least two reasons why such a bracket plays a role. Firstly, it can be obtained by projection from the Boltzmann Poisson bracket expressing kinematics of phonons. Secondly, it can be seen as the Poisson bracket of the Lie algebra dual of the semidirect product where entropy is advected by its momentum.

It is a matter of straightforward calculation that by transformation

$$\pi = \mathbf{w} s \quad (124)$$

and by using condition (119c), bracket (123) becomes bracket (118). The two brackets are thus equivalent and related by change of variables provided the vector field \mathbf{w} is curl-free.

However, bracket (123) is not constrained to the case of curl-free \mathbf{w} fields. In fact, it can be derived by projection to fields s (a scalar function of the phonon distribution function) and π (first moment of the distribution function) from the Boltzmann Poisson bracket governing dynamics of phonons. Bracket (123) is the hydrodynamic bracket expressing kinematics of phonons.

When bracket (123) has to be coupled with hydrodynamics, one can proceed simply by summing the hydrodynamic bracket for density and momentum of matter, (ρ, \mathbf{u}) , given by (79), and bracket (123),

$$\{\mathcal{A}, \mathcal{B}\}^{(\mathbf{u}, \rho, \pi, s)} = \{\mathcal{A}, \mathcal{B}\}^{(\mathbf{u}, \rho)} + \{\mathcal{A}, \mathcal{B}\}^{(Cat)}. \quad (125)$$

Coupling between those brackets is introduced by transformation from matter and phonon momentum onto total momentum and phonon momentum, $(\mathbf{u}, \rho, \pi, s) \rightarrow (\mathbf{m} = \mathbf{u} + \pi, \rho, \pi, s)$. After this transformation, Poisson bracket (125) becomes

$$\begin{aligned} \{\mathcal{A}, \mathcal{B}\}^{(\mathbf{m}, \rho, \pi, s)} &= \{\mathcal{A}, \mathcal{B}\}^{(Euler)} + \int s \left(\partial_i \mathcal{A} s \mathcal{B} \pi_i - \partial_i \mathcal{B} s \mathcal{A} \pi_i \right) d\mathbf{r} \\ &+ \int \pi_i \left[\left(\partial_j \mathcal{A} m_i \mathcal{B} \pi_j - \partial_j \mathcal{B} m_i \mathcal{A} \pi_j \right) + \left(\partial_j \mathcal{A} \pi_i \mathcal{B} m_j - \partial_j \mathcal{B} \pi_i \mathcal{A} m_j \right) + \left(\partial_j \mathcal{A} \pi_i \mathcal{B} \pi_j - \partial_j \mathcal{B} \pi_i \mathcal{A} \pi_j \right) \right] d\mathbf{r}. \end{aligned} \quad (126)$$

Further transformation from $(\rho, \mathbf{m}, s, \pi)$ to $(\rho, \mathbf{m}, s, \mathbf{w} = \pi/s)$ leads to bracket

$$\{\mathcal{A}, \mathcal{B}\}^{(CatHydAlt)} = \{\mathcal{A}, \mathcal{B}\}^{(Heat)} + \int d\mathbf{r} \frac{1}{s} (\partial_i \mathcal{A} w_j - \partial_j \mathcal{A} w_i) A_{w_i} B_{w_j}, \quad (127)$$

where the first term on the right hand side is given by Eq. (120) and the second term disappears if condition (119c) is satisfied. The condition, however, does not need to be satisfied in general, especially in the case when dissipation (usually proportional to E_w) is taken into account. Poisson bracket (127) is new.

In summary, Poisson bracket (127) has been derived in a way alternative to the way bracket (121) was derived. These brackets, however, are nearly the same. The difference lies in how the Jacobi identity is ensured for fields \mathbf{w} with non-zero curl. In the case of bracket (121) Jacobi identity was satisfied by adding a term coupling field \mathbf{w} with momentum \mathbf{m} whereas in the case of bracket (127) a yet another term coupling field \mathbf{w} with itself appeared. We have thus arrived at a SHTC-compatible ballistic heat conduction expressed in Poisson bracket (121) and an alternative bracket, (127), that contains advection of field \mathbf{w} by itself. It will be a matter of future research to compare the alternatives with results on non-Fourier heat transport, e.g. [92] or [6].

Poisson bracket (127) generates evolution equations

$$\frac{\partial m_i}{\partial t} = -\partial_j(m_i E_{m_j}) - \partial_j(w_i E_{w_j}) - \rho \partial_i E_\rho - m_j \partial_i E_{m_j} - s \partial_i E_s - w_k \partial_i E_{w_k} + \partial_i(E_{w_k} w_k) \quad (128a)$$

$$\frac{\partial w_k}{\partial t} = -\partial_k E_s - \partial_k(w_j E_{m_j}) + (\partial_k w_j - \partial_j w_k) \left(E_{m_j} + \frac{1}{s} E_{w_j} \right), \quad (128b)$$

$$\frac{\partial s}{\partial t} = -\partial_k (s E_{m_k} + E_{w_k}), \quad (128c)$$

$$\frac{\partial \rho}{\partial t} = -\partial_k (\rho E_{m_k}). \quad (128d)$$

The difference between the SHTC Cattaneo hydrodynamics and this alternative Cattaneo hydrodynamics lies in the last term of (128b) with prefactor $\frac{1}{s}$.

4.6. Complete Poisson bracket

The complete Poisson bracket generating SHTC Eulerian equations (38) is then the combination of brackets (79), (101), (113) and (121),

$$\{\mathcal{A}, \mathcal{B}\}^{(SHTC)} = -2\{\mathcal{A}, \mathcal{B}\}^{(Euler)} + \{\mathcal{A}, \mathcal{B}\}^{(DM)} + \{\mathcal{A}, \mathcal{B}\}^{(EMHD)} + \{\mathcal{A}, \mathcal{B}\}^{(Heat)}. \quad (129)$$

Bracket $\{\mathcal{A}, \mathcal{B}\}^{(Euler)}$ is contained in the last three terms and therefore we have to subtract it twice so that it is contained only once in the final Poisson bracket. Validity of the Jacobi identity for this bracket was checked by the program developed in [60].

5. DISSIPATION

Dissipative evolution is usually meant as the evolution that raises entropy. In the sense of the hyperbolic conservation laws (SHTC) entropy density is among state variables and entropy is thus conserved unless dissipative terms are added to equations (38). In the sense of GENERIC, entropy is required to be a Casimir of the Poisson bracket, i.e. $\{\mathcal{S}, \mathcal{E}\} = 0$ for all possible energies \mathcal{E} , and it is thus also conserved unless some dissipative terms are added. It is the goal of this section to compare the way dissipation is added in the SHTC and GENERIC formalisms.

Before introducing concrete forms of dissipative terms, let us make a comment on reversibility and irreversibility. Reversible evolution equations is such that it is intact by the *time-reversal transformation* (TRT), where velocities of all particles and magnetic field are inverted as well as the sign of the time increment in time derivatives. Evolution equations in non-equilibrium thermodynamics usually contain both reversible and irreversible parts. Assuming that the left hand sides of the evolution equations are constituted by only partial time derivatives of the state variables, the reversible parts of the right hand sides transform under TRT exactly as the left hand sides of the corresponding equations (partial time derivatives of the state variable) while the irreversible parts of the right hand sides gain opposite signs than the corresponding left hand sides. Reversible evolution is invariant with respect to TRT while the irreversible changes its sign.

In order to be able to apply TRT to an equation, however, one has to first introduce parities with respect to TRT. A quantity is called *even with respect to TRT* (parity equal to 1) if TRT does not alter the quantity, and it is called *odd* (parity equal to -1) if TRT changes sign of the quantity. How can we determine whether a quantity is even or odd? The odd quantities are those that change its sign when velocities of particles and magnetic field are inverted. For example, density is even, momentum is odd, entropy and energy are even. Electric displacement field is even and magnetic field is odd. There are, however, quantities with no definite parity, which means that TRT does neither leave them intact nor changes their sign, e.g. the one-particle distribution function in kinetic theory. In that case one has to consider a more geometric definition of TRT (a push-forward on vector fields), see [75]. Finally, we require (as

is usual in non-equilibrium thermodynamics) that the reversible parts of evolution equation do not change entropy (i.e. are non-dissipative) while the irreversible terms raise entropy (are dissipative), and the adjectives irreversibility and dissipativity will be then considered equivalent.

In general, dissipative (or irreversible) evolution within the SHTC framework is constructed by means of a quadratic dissipation potential, i.e. the irreversible terms are linear in derivatives of energy (conjugate state variables within SHTC). Within GENERIC we also prefer using a dissipation potential although some authors prefer using a possibly non-symmetric dissipative bracket [56, 71]. The geometric character and statistical interpretation of dissipation potentials [66], discussed also in [57], convince us to prefer dissipation potentials. The dissipation potential is, however, used in a slightly different way in SHTC than within GENERIC, which differ by the way conjugate variables are constructed from the state variables. In the SHTC formalism conjugate variables are constructed as derivatives of energy while in GENERIC they are derivatives of entropy. Let us now discuss all the mentioned possibilities.

5.1. Irreversible dynamics

5.1.1 Energy representation

In the SHTC framework, irreversible dynamics is modeled via only algebraic (independent of spatial gradients) dissipative terms which do not harm the hyperbolic type of the PDEs. In the light of our recent results [77, 17, 18] on modeling of dissipative processes such as viscous momentum, heat and charge transfer, the modeling of dissipative phenomena with algebraic terms of relaxation type does not seem to be something restrictive.

In order to simplify the notations, in this sections, we also denote the conjugate variables as

$$A_{ij}^* = E_{A_{ij}}, \quad e_i^* = E_{e_i}, \quad h_i^* = E_{h_i}, \quad s^* = E_s, \quad w_i^* = E_{w_i}. \quad (130)$$

In what follows we provide a recipe on how to introduce dissipative terms into system (38). It is strongly motivated by the key fact around the SHTC equations, the summation rule (41). In this consideration we do not consider the momentum and mass conservation as they are pure reversible time evolutions. Thus, we add, so far, arbitrary functions of the state variables into the right hand side of each of the remaining equations:

$$\frac{\partial A_{ik}}{\partial t} + \partial_k(A_{il}v_l) + v_j(\partial_j A_{ik} - \partial_k A_{ij}) = S_{ik}^A, \quad (131a)$$

$$\frac{\partial h_i}{\partial t} + \partial(h_i v_k - v_i h_k + \varepsilon_{ikl} E_{e_l}) + v_i \partial_k h_k = S_i^h, \quad (131b)$$

$$\frac{\partial e_i}{\partial t} + \partial_k(e_i v_k - v_i e_k - \varepsilon_{ikl} E_{h_l}) + v_i \partial_k e_k = S_i^e, \quad (131c)$$

$$\frac{\partial w_k}{\partial t} + \partial_k(v_i w_l + E_s) + v_j(\partial_j w_k - \partial_k w_j) = S_k^w, \quad (131d)$$

$$\frac{\partial s}{\partial t} + \partial_k(s v_k + E_{w_k}) = S^s. \quad (131e)$$

Because the functions S_{ij}^A , S_i^h , S_i^e , S_i^w and S^s are quite arbitrary, we can not guaranty that

$$A_{ij}^* S_{ij}^A + h_i^* S_i^h + e_i^* S_i^e + w_i^* S_i^w + s^* S^s \equiv 0, \quad (132)$$

which is required by the summation rule (41) and energy conservation (40). Therefore, the energy conservation, the first law of thermodynamics, might be, in general, violated. Thus, the central question is how can we introduce the dissipative sources in (131) which fulfills the first law (40)? It seems that the only degree of freedom we have is to assign to one of the variables the role of the entropy. Bearing in mind the Hamiltonian nature of the reversible part

of the SHTC equations, only a Casimir of the corresponding Poisson bracket (129) can be chosen as the entropy. The field s indeed fulfills this criterion. We emphasize that the introduction of the entropy in the SHTC framework is not a spontaneous step but it is fully motivated by the necessity to respect the first law of thermodynamics.

We now show how the entropy field can be used to fix the energy conservation, i.e. to guarantee the zero on the right-hand side of (132). It is evident that if we define the entropy source term as

$$S^s = -\frac{1}{s^*}(A_{ij}^* S_{ij}^A + h_i^* S_i^h + e_i^* S_i^e + w_i^* S_i^w) \quad (133)$$

then (132) holds automatically (recall that $s^* = E_s$ is usually interpreted as the temperature). However, there is also the second law of thermodynamics which postulates the positivity of the entropy production, $S^s \geq 0$. How can we guarantee the second law is fulfilled? Perhaps, the simplest (but not exhaustive) solution is as follows. Let us introduce the vectors of state variables and conjugate variables

$$\mathbf{Q} = (\mathbf{q}, s), \quad \mathbf{Q}^* = (\mathbf{q}^*, s^*), \quad (134)$$

where $\mathbf{q} = (A_{ij}, h_i, e_i, w_i)$ and $\mathbf{q}^* = (A_{ij}^*, h_i^*, e_i^*, w_i^*)$, i.e. they do not include s and s^* accordingly. Then, we define the dissipative sources as the partial derivatives

$$S_{ij}^A = -\frac{\partial \Psi}{\partial A_{ij}^*}, \quad S_i^h = -\frac{\partial \Psi}{\partial h_i^*}, \quad S_i^e = -\frac{\partial \Psi}{\partial e_i^*}, \quad S_i^w = -\frac{\partial \Psi}{\partial w_i^*}, \quad (135)$$

of a potential $\Psi(\mathbf{q}^*)$ which does not *explicitly* depend on s^* and is taken as

$$\Psi(\mathbf{q}^*) = \frac{1}{2} \mathbf{q}^* \Lambda \mathbf{q}^*, \quad (136)$$

where $\Lambda = \Lambda(\mathbf{Q})$ being a symmetric positive semidefinite matrix, $\Lambda^\top = \Lambda \geq 0$, whose entries may depend on the primary state variables \mathbf{Q} (as well as on other state variables whose time evolutions are reversible and are omitted in this section, e.g. ρ) and will be associated with the inverse of the characteristic dissipation times in Section 6. Therefore, with the choice (135), (136), the second law, i.e. the inequality

$$S^s \geq 0, \quad (137)$$

is automatically fulfilled.

We note that the main nonlinearity of the dissipative processes is not defined by the nonlinearity of the dissipation potential (136) or nonlinearity of the conjugate variables \mathbf{q}^* but it is, in fact, hidden in the entries of $\Lambda(\mathbf{q})$. For example, when dealing with irreversible deformations in solids, the entries of Λ are associated with the inverse strain relaxation times [39, 85, 42, 41, 31, 4, 23, 14] which may vary over several orders of magnitude.

Therefore, we have shown that the SHTC irreversible time evolution can be expressed in terms of the quadratic dissipation potential $\Psi(\mathbf{q}^*)$ (136) as

$$\left(\frac{\partial q_i}{\partial t} \right)_{\text{irr}} = -\Psi_{q_i^*}, \quad (138a)$$

except for the right hand side of the evolution equation for entropy density, which is constructed as

$$\left(\frac{\partial s}{\partial t} \right)_{\text{irr}} = \frac{1}{s^*} q_i^* \Psi_{q_i^*} \geq 0. \quad (138b)$$

This particular choice of the dissipative source terms respects the both laws of thermodynamics.

5.1.2 Entropy representation and gradient dynamics

In the previous Section, we have shown that the irreversible part of the SHTC time evolution of all the state variables can be generated by the potential $\Psi(\mathbf{q}^*)$ which does not depend explicitly on s^* , and \mathbf{q}^* were defined as the conjugate variables with respect to the energy, $\mathbf{q}^* = E_{\mathbf{q}}$. Whereas, within GENERIC, conjugate variables in the irreversible part are usually identified as derivatives of entropy with respect to the state variables and irreversible time evolution can be formulated as *nonlinear gradient dynamics* with a prefactor playing the role of temperature

$$\left(\frac{\partial \mathbf{q}}{\partial t}\right)_{irr} = \frac{1}{s_E} \frac{\delta \Xi}{\delta s_{\mathbf{q}}} \quad \text{and} \quad \left(\frac{\partial E}{\partial t}\right)_{irr} = \frac{1}{s_E} \frac{\delta \Xi}{\delta s_E}, \quad (139)$$

where δ/δ stands for a functional derivative and E is the total energy density. The prefactor was introduced for example in [75]. Is the SHTC irreversible evolution compatible with this GENERIC irreversible evolution?

The answer is affirmative, but in order to demonstrate it we have to first recall transformations between the energetic representation, where state variables are (\mathbf{q}, s) , and entropic representation, where state variables are (\mathbf{q}, E) . In the energetic representation conjugate variables are identified with derivatives of energy with respect to the state variables, $(\mathbf{q}^*, s^*) = (E_{\mathbf{q}}, E_s)$, while in the entropic representation the conjugate variables are derivatives of entropy, $(\mathbf{p}^*, E^*) = (s_{\mathbf{q}}, s_E)$. The two representations thus have different conjugate variables.

In order to show the compatibility of both formalisms, we need to recall the following relations between conjugate variables in the two representations.

$$\left(\frac{\partial E}{\partial q_i}\right)_s = -\left(\frac{\partial E}{\partial s}\right)_{\mathbf{q}} \cdot \left(\frac{\partial s}{\partial q_i}\right)_E \quad \text{and} \quad \frac{\partial E}{\partial s} = \frac{1}{\frac{\partial s}{\partial E}}, \quad (140)$$

or

$$q_i^* = -s^* p_i^* \quad \text{and} \quad s^* = \frac{1}{E^*}. \quad (141)$$

From (140) it follows that

$$\left(\frac{\partial q_i^*}{\partial p_j^*}\right)_{E^*} = -s^* \delta_{ij}, \quad \left(\frac{\partial q_i^*}{\partial E^*}\right)_{\mathbf{p}^*} = \frac{p_i^*}{(E^*)^2} \quad (142a)$$

$$\left(\frac{\partial s^*}{\partial p_i^*}\right)_{E^*} = 0, \quad \left(\frac{\partial s^*}{\partial E^*}\right)_{\mathbf{p}^*} = -\frac{1}{(E^*)^2}. \quad (142b)$$

Derivatives of the dissipation potential then transform as

$$\frac{\partial \Xi}{\partial p_i^*} = \frac{\partial \Xi}{\partial q_j^*} \frac{\partial q_j^*}{\partial p_i^*} + \frac{\partial \Xi}{\partial s^*} \frac{\partial s^*}{\partial p_i^*} = -s^* \frac{\partial \Xi}{\partial q_i^*} \quad (143a)$$

$$\frac{\partial \Xi}{\partial E^*} = \frac{\partial \Xi}{\partial q_j^*} \frac{\partial q_j^*}{\partial E^*} + \frac{\partial \Xi}{\partial s^*} \frac{\partial s^*}{\partial E^*} = -s^* \left(q_j^* \frac{\partial \Xi}{\partial q_j^*} + s^* \frac{\partial \Xi}{\partial s^*} \right). \quad (143b)$$

Irreversible GENERIC evolution equations (139) can be then reformulated as

$$\left(\frac{\partial q^i}{\partial t}\right)_{irr} = -(s^*)^2 \Xi_{q_i^*} \quad (144a)$$

$$\left(\frac{\partial E}{\partial t}\right)_{irr} = -(s^*)^2 \left(q_j^* \frac{\partial \Xi}{\partial q_j^*} + s^* \frac{\partial \Xi}{\partial s^*} \right). \quad (144b)$$

Since the SHTC dissipation potential Ψ is always assumed algebraic, i.e. does not contain spatial gradients, the sought corresponding GENERIC dissipation potential Ξ should be also algebraic. Energy conservation then requires that the right hand side of the evolution equation for total energy density disappears, i.e.

$$\Xi_{E^*} = 0 = q_j^* \frac{\partial \Xi}{\partial q_j^*} + s^* \frac{\partial \Xi}{\partial s^*}, \quad (144c)$$

which means that the dissipation potential must be a zero-homogeneous function in the energetic representation. These evolution equations imply the irreversible evolution of entropy density,

$$\left(\frac{\partial s}{\partial t} \right)_{irr} = \frac{p_j^*}{E^*} \Xi_{p_j^*} = q_j^* s^* \Xi_{q_j^*}. \quad (144d)$$

We have arrived at the algebraic energy-conserving irreversible GENERIC evolution equations in the energetic representation.

Let us now go back to the irreversible evolution formulated within the SHTC framework, Eqs. (138), which are generated by the dissipation potential Ψ , which is independent of s^* , and by the requirement of energy conservation. In order to find the gradient dynamics in the entropic representation that is compatible with the SHTC irreversible evolution, we need to find the dissipation potential Ξ such that Eqs. (144) are the same as Eqs. (138). This is achieved by choosing

$$\Xi(q^*, s^*) = \Psi(q^*)|_{q^* := q^*/s^*}. \quad (145)$$

Such a dissipation potential clearly conserves energy because it can be written as a function of p^* only (not E^*). Derivatives of this dissipation potential are

$$\Xi_{q^*} = \frac{1}{s^*} \Psi_{q^*}|_{q^* := q^*/s^*} \quad \text{and} \quad \Xi_{s^*} = -\frac{q_i^*}{(s^*)^2} \Psi_{q_i^*}|_{q^* := q^*/s^*}, \quad (146)$$

and evolution equations (144) become

$$\left(\frac{\partial q^i}{\partial t} \right)_{irr} = -s^* \Psi_{q_i^*}|_{q^* := q^*/s^*} \quad (147a)$$

$$\left(\frac{\partial s}{\partial t} \right)_{irr} = q_i^* \Psi_{q_i^*}|_{q^* := q^*/s^*}, \quad (147b)$$

which are exactly the same as Eqs. (138) for quadratic potentials Ψ .

In summary, energy-conserving algebraic irreversible GENERIC gradient dynamics with the temperature prefactor thus completely compatible with the irreversible SHTC equations provided the SHTC dissipation potential is quadratic. If the potential is not quadratic, extra dependence on temperature pops out, which does not play any crucial role because of the possible dependence of the phenomenological equations on temperature. One can even start with purely gradient dynamics without any prefactor, as is usual within GENERIC, and the resulting evolution equations in the energetic representation would be compatible with the SHTC evolution up to a prefactor given by a power of temperature.

5.2. Involution constraints in case of dissipation

It is important to remark that there is an intimate connection between the dissipation and involution constraints discussed in Section 3.2.2. The presence of the dissipative algebraic source terms S_{ik}^A , S_i^h , S_i^e and S_i^{wv} in (131) also

affects the time evolutions of $\mathbf{B} = \nabla \times \mathbf{A}$, $Q = \nabla \cdot \mathbf{h}$, $R = \nabla \cdot \mathbf{e}$ and $\mathbf{\Omega} = \nabla \times \mathbf{w}$ in the following way

$$\frac{\partial B_{ij}}{\partial t} + \partial_k (B_{ij} v_k - v_j B_{ik} + \varepsilon_{jkl} S_{il}^A) + v_j \partial_k B_{ik} = 0, \quad (148a)$$

$$\frac{\partial Q}{\partial t} + \partial_k (Q v_k + S_k^h) = 0, \quad (148b)$$

$$\frac{\partial R}{\partial t} + \partial_k (R v_k + S_k^e) = 0, \quad (148c)$$

$$\frac{\partial \Omega_j}{\partial t} + \partial_k (\Omega_j v_k - v_j \Omega_k + \varepsilon_{jkl} S_l^w) + v_j \partial_k \Omega_k = 0, \quad (148d)$$

i.e. the dissipative sources emerge as constitutive fluxes in the time evolutions for the fields \mathbf{B} , Q , R and $\mathbf{\Omega}$. Therefore, in contrast to the reversible dynamics, these fields deviate from zero values even if initially ($t = 0$) $\mathbf{B} = 0$, $Q = 0$, $R = 0$ and $\mathbf{\Omega} = 0$ and hence, may serve as indicators of irreversible dynamics. See also Section 6 with some examples for more discussions.

5.3. Dispersive dynamics

It is not obligatory in the SHTC theory that addition of the algebraic source terms to the master equations results in that the dynamics becomes irreversible, i.e. rise the entropy. In fact, the source terms can be added in such a way that both, the entropy and the total energy, are *conserved*. This can be achieved by introducing new state variables representing the dynamics at a microscale and designing the source terms which models the energy conversion between the macro- and micro- scales without any lose. For example, the impact of the dislocation density on the dispersive properties of the elastic waves was studied in [84] where it was assumed that the material has non-zero initial concentration of defects and the energy converts reversibly between the kinetic energy carried by the elastic waves and the energy stored in the microscopic defects.

We recall that in the classical continuum mechanics, the dispersive dynamics, as well as the dissipative dynamics, is modeled by means of high-order PDEs. However, as we have demonstrated in [77, 17, 18] that the classical dissipative models can be successfully modeled with first-order PDEs, the dispersive properties of the continuous media can be also modeled with merely first-order PDEs, e.g. see [84, 24, 65].

6. EXAMPLES

6.1. MaxEnt reduction

In non-equilibrium thermodynamics one often deals with the problem of how to reduce some detailed evolution to a less detailed evolution. In particular, one often has a fast state variable which is to be eliminated from the evolution equations. The standard procedure for elimination of the fast variable is the Chapman-Enskog reduction, see e.g. [51]. However, the reduction often relies on demanding calculations and rigorous origin of the method can also be questioned.

An alternative method to the Chapman-Enskog reduction was proposed in [75]. The method relies on identification of state and conjugate variables in the evolution equations. Note that the Chapman-Enskog reduction does not use this extra information. It is becoming fashionable to formulate non-equilibrium thermodynamics within contact geometry, where the conjugate variables have their *independent* meaning and own evolution equations, see e.g. [48]. In this sense, it is reasonable to let conjugate and state variables relax with different speeds. For example, in the reduction proposed in [75], let us refer to it as to the MaxEnt reduction, the fast variable is set to the corresponding value given by maximization of entropy while the conjugate fast variable is set to a value solving the stationary fast

evolution equation. This way the conjugate fast variable becomes a function of the other state variables, and due to the coupling between the fast and slow variables, evolution equations for the slow variables obtain irreversible terms caused by relaxation of the fast variable. Let us demonstrate this reduction on the following examples.

6.2. Navier-Stokes dynamics

Compressible Navier-Stokes (NS) equations consist of a reversible and irreversible part. The reversible part, the compressible Euler equations, is of course implied by the SHTC and GENERIC formalisms. Taking for example Poisson bracket (79), the implied evolution equations are equations (85), which represent the Euler equations.

The irreversible part of NS equations is more difficult to obtain. The NS equations can not be incorporated into the SHTC framework because of the parabolic character of the irreversible terms (Laplacian of velocity). Within GENERIC, the NS equations can be recovered by taking a quadratic dissipation potential, or a dissipative bracket, which is composed of second derivatives of the potential, as in [71],

$$\Xi(\mathbf{x}^*) = \frac{1}{2} S_{x^i} M^{ij} S_{x^j}, \quad (149)$$

where state variables are $\mathbf{x} = (\mathbf{m}, \rho, E)$ and \mathbf{x}^* are the corresponding conjugate (with respect to entropy) variables.

Our goal, however, is to see the Navier-Stokes equation as an approximation of the SHTC equations. This, in particular, was demonstrated in [77, 17], but this time we would like also to demonstrate the work of the MaxEnt reduction. For this purpose, let us consider dynamics of the Finger (or left Cauchy-Green) tensor from Sec. 4.3.4, which is a consequence of kinematics of the generalized distortion matrix. Let the energy potential depend on the fields through

$$E = \varepsilon(\rho, s) + \frac{1}{2} \alpha^2 (\text{dev} \mathbf{B})^2 + \frac{1}{2} \beta^2 (\text{tr} \mathbf{B} - 3)^2 + \frac{\mathbf{m}^2}{2\rho}, \quad (150)$$

derivative of which with respect to the Finger tensor is

$$E_{\mathbf{B}} = \alpha^2 \text{dev} \mathbf{B} + \beta^2 (\text{tr} \mathbf{B} - 3) \mathbf{I}. \quad (151)$$

Note that $\text{dev} \mathbf{B}$ is the deviatoric (traceless) part of tensor \mathbf{B} and α and β are assumed to be material-dependent constants (typically sound speeds, e.g. see [17]). The derivative $E_{\mathbf{B}}$ is zero for $\mathbf{B} = \mathbf{I}$. Similarly, one can express entropy as function of $(\mathbf{m}, \mathbf{B}, \rho, E)$ and derivative of entropy with respect to \mathbf{B} will be zero only for Finger tensor equal to the unit matrix. The MaxEnt value of \mathbf{B} is thus the unit matrix \mathbf{I} .

Putting the MaxEnt value of \mathbf{B} into the evolution equations (109), and adding the shear and volume dissipative terms⁸ $-\tau_S^{-1} \text{dev} \mathbf{B}^*$ and $-\tau_V^{-1} \text{tr}(\mathbf{B}^*) \mathbf{I}$ to the right hand side of the equation for \mathbf{B} , leads to

$$\frac{\partial m_i}{\partial t} = -\partial_j (m_i m_j^*) - \rho \partial_i \rho^* - m_j \partial_i m_j^* - s \partial_i s^* + \partial_k (\mathbf{B}_{ki}^* + \mathbf{B}_{ik}^*) \quad (152a)$$

$$0 = \partial_i m_j^* + \partial_j m_i^* - \frac{1}{\tau_S} \text{dev} \mathbf{B}_{ij}^* - \frac{1}{\tau_V} \text{tr}(\mathbf{B}^*) \delta_{ij}, \quad (152b)$$

$$\frac{\partial \rho}{\partial t} = -\partial_i (\rho m_i^*) \quad (152c)$$

$$\frac{\partial s}{\partial t} = -\partial_i (s m_i^*). \quad (152d)$$

⁸The dissipative terms can be seen as derivatives of a quadratic dissipation potential with respect to \mathbf{B}^* , see Section 5. It should be also borne in mind that \mathbf{B}^* is interpreted as an element of the dual space to the space where \mathbf{B} lives, but in the precise variant of the evolution, Eqs. (109), it can be identified with derivative of energy with respect to \mathbf{B} .

The last equation means that

$$\text{dev}\mathcal{B}_{ij}^* = \tau_S \left(\partial_i m_j^* + \partial_j m_i^* - \frac{2}{3} \partial_k m_k^* \delta_{ij} \right) \quad \text{and} \quad \text{tr}(\mathcal{B}^*) = \frac{2}{3} \tau_V \partial_k m_k^*, \quad (153)$$

which, after substitution into the equation for m and evaluating the conjugate variables as the corresponding derivatives of energy, leads to the Navier-Stokes equations

$$\begin{aligned} \frac{\partial m_i}{\partial t} = & -\partial_j (m_i E_{m_j}) - \rho \partial_i E_\rho - m_j \partial_i E_{m_j} - s \partial_i E_s \\ & + \partial_k \left(2\tau_S \left(\partial_i E_{m_k} + \partial_k E_{m_i} - \frac{2}{3} \partial_l E_{m_l} \delta_{ik} \right) \right) + \partial_i \left(\frac{4}{3} \tau_V \partial_k E_{m_k} \right), \end{aligned} \quad (154)$$

where τ_S and τ_V are related to shear and volume viscosities.

Note that if the volumetric relaxation time, τ_V , goes to infinity, it follows from Eq. (152b) that the spherical part of the \mathcal{B} tensor remains undetermined and the condition of incompressibility, $\nabla \cdot E_m = 0$, is enforced after the MaxEnt reduction. This way the incompressible Navier-Stokes equation can be recovered, and the undetermined spherical part of \mathcal{B} plays the role of pressure in incompressible fluids.

In summary, compressible (and also incompressible) Navier-Stokes equation is an approximation of the evolution of the Finger tensor, which is implied by evolution of the generalized distortion matrix, compatible with the SHTC equations.

6.3. Elastic and elastoplastic solids

Classical evolution of elastic solids can be formulated by means of the distortion matrix easily [39, 42, 41]. The distortion matrix is then interpreted as inverse of the deformation gradient, and evolution equations (102) or (38a), (38b), (38g) become equivalent with the balance of mass, balance of momentum, and evolution equation for the inverse deformation gradient in the Eulerian frame. Recall that the fact that the distortion matrix is the inverse deformation gradient is expressed by the integrability condition $\nabla \times \mathbf{A} = 0$.

Equations (102) or (38b) are however much more general than the mentioned Eulerian equations of classical continuum theory because matrix \mathbf{A} is allowed to have non-zero curl, in which case the interpretation as a *global* deformation field becomes invalid. The case of distortion matrix with non-zero curl corresponds to irreversible deformations, where indeed one can not reconstruct the original Lagrangian frame precisely, e.g. see [39, 85, 42, 41]. Non-zero curl of \mathbf{A} is caused by dissipative term $\frac{1}{\tau} E_{\mathbf{A}}$ (see (148a)) in the evolution equation for the distortion matrix because the time evolution for the Burgers tensor $\mathbf{B} = \nabla \times \mathbf{A}$ (dislocation density tensor) is, e.g. see Appendix C or [39, 42, 41, 76],

$$\frac{\partial B_{ij}}{\partial t} + \partial_k \left(B_{ij} v_k - v_j B_{ik} + \varepsilon_{jkm} \frac{1}{\tau} E_{A_{im}} \right) + v_j \partial_k B_{ik} = 0, \quad (155)$$

from which it follows that even if $B_{ij} = 0$ initially it will, in general, distinct from zero at later times. One can clearly see that this equation has the SHTC structure of equation (38c). However, to have the full SHTC and GENERIC structure, this equation should be accompanied by a time evolution for a complimentary tensor field \mathbf{D} , exactly as \mathbf{h} is the complimentary field to \mathbf{e} , see Section 3.1.3. In the theory of dislocations, the complimentary field \mathbf{D} has the meaning of the flux dislocation density, e.g. see a discussion in [76].

It is interesting to note that, in case $\mathbf{B} \neq 0$, i.e. in case of irreversible deformations, the SHTC Eulerian equations (38a), (38b) can be symmetrized only in the extended weakly non-local sense, see Section 3.2.6, i.e. when the Burgers tensor \mathbf{B} and the flux dislocation density tensor \mathbf{D} are considered as new independent state variables and the total energy potential $E = E(m, \mathbf{A}, \mathbf{B}, \mathbf{D}, \rho, s)$ depends on the extended set of state variables. The later leads to the

appearing of extra stress in the momentum flux (38a) due to dislocation motion:

$$\frac{\partial m_i}{\partial t} + \frac{\partial}{\partial x_k} \left(m_i E_{m_k} + \delta_{ik} \left(\rho E_\rho + \sigma E_\sigma + m_l E_{m_l} + B_{jl} E_{B_{jl}} + D_{jl} E_{D_{jl}} - E \right) - B_{jk} E_{B_{ji}} - D_{jk} E_{D_{ji}} + A_{jk} E_{A_{ji}} \right) = 0. \quad (156)$$

Thus, evolution of the distortion matrix \mathbf{A} gives, in fact, exact evolution of the Burgers tensor, defining dynamics of dislocations [39, 42, 41]. The evolution equation of the distortion matrix coupled with mass, momentum and energy conservation, thus represents an Eulerian framework covering elastic and elasto-plastic solids as well as fluids, see also [77, 17, 78, 85, 4, 14, 23]. In practice, the transition between the elastic and plastic response of solids is governed by the dependence of the strain relaxation time on the state parameters, $\tau = \tau(\mathbf{A}, \mathbf{B}, \mathbf{D}, \rho, s)$, see details in [78, 85, 4, 14, 23].

6.4. Non-Newtonian fluid dynamics

Dynamics of viscoelastic fluids (or solids) can be also formulated within the SHTC and GENERIC formalisms by the proper choice of the dissipative source term in the time evolution for the distortion matrix:

$$\frac{\partial A_{ik}}{\partial t} + \partial_k (A_{im} v_m) + v_j (\partial_j A_{ik} - \partial_k A_{ij}) = -\frac{1}{\tau'} E_{A_{ik}} \quad (157)$$

or by using the Finger tensor as in Sec. 4.3.4. Here, the relaxation function $\tau' = \tau'(\tau)$ depends on the real relaxation time τ and typically is defined as $\tau' \sim \tau b^2$, while its further specification depends on the specific choice of the total energy, see [77, 17]. Here, b is the shear sound speed (the speed of propagation of small transversal perturbations). For the further discussion, let us recall the physical interpretation of the strain dissipation time τ . In the unified flow theory [77, 17, 78], τ is understood as a continuum interpretation of the seminal idea of the so-called *particle settled life time* of Frenkel [26], who applied it to describe the ability of liquids to flow, see also recent promising experimental and theoretical advances [15, 10, 12, 11, 13] confirming and further developing Frenkel's idea. Thus, in our continuum approach, the time τ is the time taken by a given continuum particle (finite volume) to “escape” from the cage composed of its neighbor particles, i.e. the time taken to rearrange with one of its neighbors. The more viscous a fluid is, the larger the time τ , i.e. the longer the continuum particles stay in contact with each other.

Such a concept of the strain dissipation time, in contrast to the phenomenological viscosity concept, allows to cover the entire spectrum of material responses, from ideal fluids ($\tau = 0$) to elastic solids ($\tau = \infty$) through the viscous fluids, either Newtonian or non-Newtonian ($0 < \tau < \infty$). Note that the viscosity concept does not allow to model deformable solids. Moreover, in the proposed framework, the division on the Newtonian and non-Newtonian fluids becomes artificial because it does not rely on any constitutive laws designed specifically for either Newtonian or non-Newtonian flows. For example, to apply our model to modeling Newtonian flows one may even do not know that Newton's viscous law exists. All flows are treated as being non-equilibrium and no assumptions on the closeness to the global equilibrium was assumed in both formalisms. Thus, according to our approach, simple fluids as water for example exhibit elastic response at time scales $T < \tau$, viscoelastic at time scales $T \sim \tau$ and Newtonian response at time scales $T \gg \tau$, where T is the characteristic flow time scale. We emphasize that the viscoelastic and even solid like properties of simple liquids at small time scales is an experimental fact, see e.g. [15, 10, 12, 11, 13] and was predicted by Frenkel [26].

In practice, as in the modeling of elastoplastic solids, the user have to provide a suitable function $\tau = \tau(\mathbf{A}, \rho, s)$ to cover diverse non-Newtonian responses. We emphasize that, in contrast to the viscosity approach, there is no need to use the dependence of τ on the strain rate because our relaxation model is rate dependent by the construction, e.g. see Fig. 1 in [77]. Moreover, in contrast to the nonlinear viscosity, the switching from $\tau = \text{const}$ to nonlinear $\tau(\mathbf{A}, \rho, s)$ does not increase the complexity of the model. Thus, in classical phenomenological approach the substitution of the constant viscosity onto a nonlinear viscosity transforms a linear parabolic PDE system into a nonlinear parabolic system which is a way more difficult problem than the linear theory from both, the theoretical

as well as numerical viewpoint. In our case, $\tau(\mathbf{A}, \rho, s)$ does not affect the hyperbolic differential part (reversible part of the time evolution) of the model at all.

6.5. Heat and mass transfer

In this section, we demonstrate the viewpoint discussed in the Introduction section on the *universality of the mathematical structure of the equations*. We shall demonstrate that equations (38e), (38f) equipped with proper dissipative source terms and coupled with momentum and mass conservation, i.e.

$$\frac{\partial m_i}{\partial t} + \partial_k(m_i v_k + [\rho E_\rho + \sigma E_\sigma + m_l E_{m_l} - E]\delta_{ik} + w_i E_{w_k}) = 0, \quad (158a)$$

$$\frac{\partial w_k}{\partial t} + \partial_k(v_l w_l + E_\sigma) + v_j(\partial_j w_k - \partial_k w_j) = -\frac{1}{\tau} E_{w_k}, \quad (158b)$$

$$\frac{\partial \sigma}{\partial t} + \partial_k(\sigma v_k + E_{w_k}) = \frac{1}{E_\sigma \tau} E_{w_i} E_{w_i}, \quad (158c)$$

$$\frac{\partial \rho}{\partial t} + \partial_k(\rho v_k) = 0, \quad (158d)$$

possess a sort of universal structure because they can be applied to quite different physical settings. Namely, to the transport of heat and transport of mass. In particular, we shall demonstrate that the solution to this system satisfies the equilibrium laws of heat and mass transfer of Fourier and Fick accordingly, if the energy potential E is properly designed. The non-Fourier heat conduction and non-Fickian diffusion corresponds to $\tau \sim T$ and are beyond the scope of this paper. Here, τ is the relaxation parameter in (158b), and T is the characteristic macroscopic time scale. Let us start with the discussion of the heat transfer.

6.5.1 Heat conduction

Energy conservation law (40) for system (158) reduces to

$$\frac{\partial E}{\partial t} + \partial_k(v_k E + v_i([\rho E_\rho + \sigma E_\sigma + m_l E_{m_l} - E]\delta_{ik} + w_i E_{w_k}) + E_\sigma E_{w_k}) = 0, \quad (159)$$

from which one can recognize the heat flux

$$q_i = E_\sigma E_{w_i}. \quad (160)$$

By taken the energy potential in the most simple form

$$E = E^{hydro}(\rho, s) + \frac{\alpha^2}{2} \mathbf{w}^2 + \frac{1}{2\rho} m^2 \quad (161)$$

and expanding the solution in a series in τ , e.g. $\mathbf{w} = \mathbf{w}_0 + \tau \mathbf{w}_1 + \tau^2 \mathbf{w} + \dots$, one can show (exactly as in [17]) that in the leading terms, the Fourier law of heat conduction is recovered as

$$q_i = -\rho \theta \tau \alpha^2 \frac{\partial \theta}{\partial x_i}, \quad \theta = E_\sigma, \quad (162)$$

where θ is the temperature. From this relation, it is easy to recognize the heat conductivity κ as $\kappa = \rho \theta \tau \alpha^2$.

Let us now demonstrate that similar results can be obtained with the help of MaxEnt approach (see Section 6.1) which we have already applied to the Navier-Stokes equations in Section 6.2. Again, for this purpose, it is more convenient to use equivalent form (122) of (158). Thus, the goal is to reduce the Cattaneo-type hydrodynamics,

represented by the equations

$$\frac{\partial m_i}{\partial t} = -\partial_k(m_i E_{m_k}) - \rho \partial_i E_\rho - s \partial_i E_s - m_k \partial_i E_{m_k} - w_i \partial_k w_k^* - w_k^* (\partial_a w_i - \partial_i w_k), \quad (163a)$$

$$\frac{\partial w_k}{\partial t} = -\partial_k E_s - w_j \partial_k E_{m_j} - E_{m_j} \partial_j w_k + \frac{1}{s} (\partial_k w_j - \partial_j w_k) w_j^* - \frac{1}{\tau} w_k^*, \quad (163b)$$

$$\frac{\partial s}{\partial t} = -\partial_k (s E_{m_k} + w_k^*) \quad (163c)$$

into hydrodynamics with Fourier heat conduction. Here, we use the notation $w_i^* = E_{w_i}$ for the w conjugate field.

As always, the energy is assumed to be a convex function of w and thus the entropy is a concave function of w . Derivative of entropy with respect to w is zero for $w = 0$, which is the value to which field w is set in the MaxEnt reduction. This leads to equations

$$\frac{\partial m_i}{\partial t} = -\partial_j (m_i E_{m_j}) - \rho \partial_i E_\rho - m_j \partial_i E_{m_j} - s \partial_i E_s \quad (164a)$$

$$0 = -\partial_k E_s - \frac{1}{\tau} w_k^*, \quad (164b)$$

$$\frac{\partial s}{\partial t} = -\partial_k (s E_{m_k} + w_k^*). \quad (164c)$$

The second equation gives

$$w_k^* = -\tau \partial_k \theta, \quad \theta = E_s, \quad (164d)$$

where θ being the temperature.

Plugging the relaxed value of w^* into the equation for entropy yields

$$\frac{\partial m_i}{\partial t} = -\partial_j (m_i E_{m_j}) - \rho \partial_i E_\rho - m_j \partial_i E_{m_j} - s \partial_i E_s \quad (165a)$$

$$\frac{\partial s}{\partial t} = -\partial_k (s E_{m_k} - \tau E_s), \quad (165b)$$

which represent Euler equations with Fourier heat conduction.

Eventually we note that, by combining the Cattaneo-type hydrodynamics with the dynamics of the Finger tensor, Sec. 4.3.4, one can obtain the full Navier-Stokes-Fourier (NSF) system of equations by the MaxEnt reduction from the SHTC framework although the NSF system is not part of the framework itself.

6.5.2 Mass transfer

In this section we continue exploiting equations (158) and, this time, show that the classical Fickian diffusion usually described by parabolic PDE can be successfully modeled by equations (158). However, for this purpose we shall prescribe different meanings to the variables (w, σ) .

Let us consider a mixture of two inviscid fluids. We characterize the mixture morphology by three independent scalars, the mass density of the mixture ρ , and the mass and volume fractions of one of the phases (denoted by the index "1") c and α correspondingly. These three scalars relate to each other through the phase mass densities ρ_1 and ρ_2 as

$$\rho = \alpha \rho_1 + (1 - \alpha) \rho_2, \quad c = \frac{\alpha \rho_1}{\rho}. \quad (166)$$

Obviously that the mass and volume fraction of the second phase are $1 - c$ and $1 - \alpha$ accordingly.

The kinematics of the mixture is characterized by two velocity fields

$$\mathbf{v} = c\mathbf{v}^1 + (1-c)\mathbf{v}^2, \quad \mathbf{w} = \mathbf{v}^1 - \mathbf{v}^2, \quad (167)$$

which are the mixture velocity and the relative velocity of the phases.

Thus, if one identifies the scalar σ in (158) with ρc and introduces the specific total energy $E = E/\rho$ then system (158) can be written as

$$\frac{\partial \rho v_i}{\partial t} + \partial_k (\rho v_i v_k + \rho^2 E_\rho \delta_{ik} + \rho w_i E_{w_k}) = 0, \quad (168a)$$

$$\frac{\partial w_k}{\partial t} + \partial_k (w_j v_j + E_c) + v_j (\partial_j w_k - \partial_k w_j) = -\frac{1}{\tau^w} E_{w_k}, \quad (168b)$$

$$\frac{\partial \rho c}{\partial t} + \partial_k (\rho c v_k + \rho E_{w_k}) = 0, \quad (168c)$$

$$\frac{\partial \rho}{\partial t} + \partial_k (\rho v_k) = 0, \quad (168d)$$

$$\frac{\partial \rho \alpha}{\partial t} + \partial_k (\rho \alpha v_k) = -\frac{1}{\tau^\alpha} \rho E_\alpha, \quad (168e)$$

$$\frac{\partial \rho s}{\partial t} + \partial_k (\rho s v_k) = \frac{1}{E_s} \left(\frac{1}{\tau^A} E_{A_{ij}} E_{A_{ij}} + \frac{\rho^2}{\tau^\alpha} E_\alpha E_\alpha + \frac{1}{\tau^w} E_{w_i} E_{w_i} \right) \geq 0, \quad (168f)$$

where we also add the time evolution of $\rho \alpha$, while the entropy time evolution now has the structure of fluxes not as in the heat conduction case in the previous section, i.e. as in the time evolution for the scalar σ in (158), but as the mass conservation. Recall that PDEs with such a simple structure of fluxes as in (168d), (168e) and (168f) can be added in an arbitrary amount, see Section 3.1.3.

To close the system (168), one has to provide the specific total energy potential E . For example, it can be taken as

$$E = c e^1(\rho_1, s) + (1-c) e^2(\rho_2, s) + \frac{1}{2} c(1-c) \mathbf{w}^2 + E^{\text{mix}}(\rho, s, \alpha, c) + \frac{1}{2} \mathbf{v}^2, \quad (169)$$

where e^1 and e^2 are the internal energies of the phases, E^{mix} defines the chemical interaction of the phase molecules and can be left unspecified for our purposes. Then, the mixture chemical potential is defined as the derivative of the energy with respect to the mass fraction

$$E_c = e^1 + \rho_1 e_{\rho_1}^1 - e^2 - \rho_2 e_{\rho_2}^2 + \frac{1}{2} (1-2c) \mathbf{w}^2 + E_c^{\text{mix}} = \mu_1 - \mu_2 + \frac{1}{2} (1-2c) \mathbf{w}^2 + E_c^{\text{mix}}, \quad (170)$$

where $\mu_i = e^i + \frac{p_i}{\rho_i}$, $i = 1, 2$ are the chemical potentials of the phases.

Taking the derivatives of the total energy with respect to α we obtain the pressure relaxation term in the volume fraction time evolution

$$E_\alpha = \frac{1}{\rho} (p_2 - p_1), \quad (171)$$

where $p_i = \rho_i^2 \frac{\partial e^i}{\partial \rho_i}$ are the phase pressures.

Now, we are in position to show that the Fick law of diffusion is an inherent property of the model (168). As in the case of the Fourier law of heat conduction, it can be shown that the Fick law is recovered in our hyperbolic model in the leading terms for small relaxation time τ^w when the mass fraction time evolution reduces to

$$\rho \frac{dc}{dt} = \nabla \cdot [\rho \tau^w \nabla (\mu_1 - \mu_2 + E_c^{\text{mix}})] \quad (172)$$

from which one can recognize the Fick law

$$\rho \frac{dc}{dt} = \nabla \cdot \mathbf{J}, \quad \mathbf{J} = \eta \nabla E_c \quad (173)$$

or

$$\rho \frac{dc}{dt} = \nabla \cdot (D \nabla c), \quad D = \eta E_{cc} \quad (174)$$

if to require

$$\eta = \rho \tau^w, \quad (175)$$

and hence the effective diffusion coefficient is $D = \rho \tau^w E_{cc}$.

Eventually, we recall that the viscous momentum transfer is modeled by the distortion field in the SHTC formulation [77, 17, 18]. Thus, if one wants to generalize the above equations to model the viscous properties of mixtures then it is sufficient to add only one distortion field representing the distortion of the mixture elements. Then, the strain relaxation time, see [17], should be composed of the phase relaxation times. We hope to discuss this in a subsequent paper dedicated to modeling of multi-phase flows.

6.6. Electrodynamics of slowly moving medium

The electrodynamics of slowly moving medium ($|v| \ll c$, where c is the speed of light) can be modeled by equations (38a), (38d), (38c) and (38g), see [18, 22]. By means of adding the dissipative source term $-\frac{1}{\eta} E_{e_i}$ to the right hand side of the equation (38d) which represents the dynamics of the electric field, one can describe dielectrics ($\eta \rightarrow \infty$), ideal conductors ($\eta \rightarrow 0$), and resistive conductors ($0 < \eta < \infty$) as particular cases, where η is the resistivity, see details in [18, 22].

7. CONCLUDING REMARKS

Continuum mechanics of fluids and solids with dislocations, with Cattaneo type heat conduction, with mass transfer, and with electromagnetic fields is formulated in the Hamiltonian and the Godunov type form. In order to emphasize that the Godunov type structure that we consider is an extension of the original Godunov structure of local conservation laws we also call it SHTC structure. The Hamiltonian structure guarantees the physical well-posedness and the Godunov like structure the mathematical and the numerical well-posedness.

From the microscopic point of view, the physical systems under consideration are mechanical systems, their microscopic governing equations thus possess the Hamiltonian structure. This structure then passes also to the time reversible part of the dynamics in the continuum formulation. However, due to the ignorance of microscopic details in the continuum formulation, a new dynamics, that is time irreversible and of gradient type, emerges. The governing equations of continuum mechanics are thus an appropriate combination of the Hamiltonian and the gradient dynamics (called GENERIC).

The governing equations in the SHTC form are first order symmetric hyperbolic partial differential equations admitting a companion local conservation law. We have seen that the Hamiltonian continuum equations can be cast into the SHTC form. The companion local conservation law in the SHTC formulation appears in the Hamiltonian formulation as a statement about the degeneracy of the Hamiltonian structure. The time irreversible part of the continuum dynamics is in both the GENERIC and the SHTC formulations of gradient type. In the SHTC formulation the time irreversible part is moreover required to be free of spatial derivatives. The advantage of the SHTC formulation is the mathematical well-posedness in the sense that the partial differential equations can be symmetrized (due to the existence of the companion local conservation law) and as such the Cauchy initial value problem for them is well posed. The numerical well-posedness means that the Godunov finite volume discretization method, that keeps the physical content of the governing equations also in their discrete form, can be applied.

ACKNOWLEDGMENTS

I.P. acknowledges a financial support from ANR-11-LABX-0040-CIMI within the program ANR-11-IDEX-0002-02. M.P. and M.G. were supported by Czech Science Foundation, project no. 17-15498Y, by Natural Sciences and Engineering Research Council of Canada (NSERC). E.R. acknowledges a partial support by the Program N15 of the Presidium of RAS (project 121) and the Russian Foundation for Basic Research (grant No 16-29-15131).

A. LAGRANGE-TO-EULER TRANSFORMATION

In this section we recover all the technical details omitted in [30] of the transformation of the Lagrangian master system (18) into its Eulerian counterpart (38).

In what follows, we use the notations $\mathbf{A} = \mathbf{F}^{-1}$, $w = \det(\mathbf{F})$, and we use primes to denote the Lagrangian fields, e.g. m' is the field appearing in equation (18a), etc. Yet the reference (Lagrangian) density will be denoted not as ρ' but ρ_0 .

The Lagrangian and Eulerian fields relate to each other by the following formulas

$$\mathbf{m}' = w \mathbf{m}, \quad \mathbf{F} = \mathbf{A}^{-1}, \quad \rho_0 = w \rho, \quad (176a)$$

$$\mathbf{e}' = w \mathbf{A} \mathbf{e}, \quad \mathbf{h}' = w \mathbf{A} \mathbf{h}, \quad (176b)$$

$$\mathbf{w}' = \mathbf{F}^T \mathbf{w}, \quad \sigma' = w \sigma, \quad (176c)$$

while the Lagrangian total energy density U is related to Eulerian total energy density E as

$$U = w E. \quad (176d)$$

For the Lagrange-Euler transformation, we also need the following relations between the derivatives of E and U with respect to the state variables

$$E_{h_i} = A_{ji} U_{h'_j}, \quad E_{e_i} = A_{ji} U_{e'_j}, \quad (177a)$$

$$E_{\sigma} = U_{\sigma'}, \quad E_{w_k} = \frac{1}{\rho_0} F_{kj} U_{w'_j}. \quad (177b)$$

A.1. Auxiliary relations

Here, we summarize the definitions and formulas used in the Lagrange-to-Euler transformation. The total deformation gradient $\mathbf{F} = [F_{ij}]$, the distortion matrix $\mathbf{A} = [A_{ij}]$ and the velocity are defined as

$$F_{ij} = \frac{\partial x_i}{\partial y_j}, \quad \mathbf{A} = \mathbf{F}^{-1}, \quad w = \det(\mathbf{F}) = \frac{\rho_0}{\rho}, \quad v_i = \frac{\partial x_i}{\partial t}, \quad \frac{d}{dt} = \frac{\partial}{\partial t} + v_k \frac{\partial}{\partial x_k} \quad (178)$$

where y_j are the Lagrangian coordinates and x_i are the Eulerian ones, ρ and ρ_0 are the actual and the reference mass densities, respectively. The time evolution equation for F_{ij} in the Lagrangian coordinates

$$\frac{dF_{ij}}{dt} - \frac{\partial v_i}{\partial y_j} = 0 \quad (179)$$

is a trivial consequence of definitions (178)₁ and (178)₄.

The following standard definitions and formulas are also introduced

$$\mathbf{C} = \text{cof}(\mathbf{F}) = w\mathbf{A}^T = [C_{ij}], \text{ or } A_{ij} = w^{-1}C_{ji}, \quad (180)$$

$$\frac{\partial C_{ij}}{\partial y_j} = 0, \quad \varepsilon_{ijk} \frac{\partial F_{ik}}{\partial y_j} = 0, \quad (181)$$

$$\frac{dw}{dt} - \frac{\partial w A_{jk} v_k}{\partial y_j} = 0, \text{ or using (180) and (181): } \frac{dw}{dt} - w A_{jk} \frac{\partial v_k}{\partial y_j} = 0, \text{ or } \frac{d\rho}{dt} + \rho A_{jk} \frac{\partial v_k}{\partial y_j} = 0, \quad (182)$$

$$\frac{\partial w^{-1} F_{kj}}{\partial x_k} = 0, \quad \frac{\partial}{\partial t} \left(\frac{1}{w} \right) + \frac{\partial}{\partial x_k} \left(\frac{v_k}{w} \right) = 0, \quad (183)$$

$$\varepsilon_{ikl} A_{mi} A_{jk} A_{al} = \varepsilon_{mja} w^{-1}, \quad (184)$$

$$C_{km} = \frac{1}{2} \varepsilon_{lnk} \varepsilon_{pqm} F_{lp} F_{nq}, \quad (185)$$

$$\varepsilon_{imn} \varepsilon_{jmn} = 2\delta_{ij}. \quad (186)$$

Here, ε_{ijk} is the Levi-Civita symbol and δ_{ik} is the Kronecker delta.

A.2. (w_i, σ) -pair

Here, we work with the complimentary pair

$$\begin{aligned} \frac{dw'_j}{dt} + \frac{\partial U'_\sigma}{\partial y_j} &= 0, \\ \frac{d\sigma'}{dt} + \frac{\partial U_{w'_j}}{\partial y_j} &= 0. \end{aligned}$$

This pair of equation can be used (if proper energy potential and dissipation source terms are introduced) to model heat transfer (σ then should be treated as the entropy while w_j as some heat vector) or multi-phase flows (σ should be treated as the mass fraction while w_j as the relative velocity vector).

Moreover, we also assume that $\sigma = \rho s$ and $E = \rho E$. Note that the specific energy E is related to the Lagrangian energy density U as $E = \rho_0 U$.

A.2.1 Transformation (38f)→(18f)

$$\begin{aligned} \frac{\partial \sigma}{\partial t} + \frac{\partial (\sigma v_k + E_{w_k})}{\partial x_k} &= \frac{\partial \rho s}{\partial t} + \frac{\partial (\rho s v_k + \rho E_{w_k})}{\partial x_k} = s \frac{\partial \rho}{\partial t} + \rho \frac{\partial s}{\partial t} + \rho s \frac{\partial v_k}{\partial x_k} + v_k \rho \frac{\partial s}{\partial x_k} + v_k s \frac{\partial \rho}{\partial x_k} + \frac{\partial \rho E_{w_k}}{\partial x_k} = \\ &= s \frac{d\rho}{dt} + \rho \frac{ds}{dt} + \rho s \frac{\partial v_k}{\partial x_k} + \frac{\partial \rho E_{w_k}}{\partial x_k} = s \frac{d\rho}{dt} + \rho \frac{ds}{dt} + \rho s A_{jk} \frac{\partial v_k}{\partial y_j} + A_{jk} \frac{\partial \rho E_{w_k}}{\partial y_j} \end{aligned}$$

We now use (182)₃ to substitute time derivative of the density:

$$-s \rho A_{jk} \frac{\partial v_k}{\partial y_j} + \rho \frac{ds}{dt} + \rho s A_{jk} \frac{\partial v_k}{\partial y_j} + A_{jk} \frac{\partial \rho E_{w_k}}{\partial y_j} = \rho \frac{ds}{dt} + A_{jk} \frac{\partial \rho E_{w_k}}{\partial y_j} = \rho_0 w^{-1} \frac{ds}{dt} + w^{-1} C_{kj} \frac{\partial \rho E_{w_k}}{\partial y_j} = 0$$

Using constraint (181)₁ for the cofactor matrix, the last equality can be rewritten as

$$\frac{d\rho_0^s}{dt} + \frac{\partial \rho C_{kj} E_{w_k}}{\partial y_j} = 0,$$

and after using (176c) and $w^{-1}C_{kj} = A_{jk}$, as

$$\frac{d\sigma'}{dt} + \frac{\partial A_{jk} \rho_0 E_{w_k}}{\partial y_j} = 0.$$

Eventually, after noting that $E_{w_k} = \left(\frac{1}{\rho_0} U(F_{ba} w_a) \right)_{w_k} = \frac{1}{\rho_0} F_{kj} U_{w'_j}$ the latter equation becomes

$$\frac{d\sigma'}{dt} + \frac{\partial U_{w'_j}}{\partial y_j} = 0, \quad (188)$$

which is the sought Lagrangian equation (18f).

A.2.2 Transformation (38e)→(18e):

Denoting $\omega_i = \varepsilon_{ijk} \partial_j w_k$, equation (38e) reads as

$$\begin{aligned} \frac{\partial w_k}{\partial t} + \frac{\partial (v_\alpha w_\alpha + E_\sigma)}{\partial x_k} + \varepsilon_{klj} \omega_l v_j &= \frac{\partial w_k}{\partial t} + v_\alpha \frac{\partial w_\alpha}{\partial x_k} + w_\alpha \frac{\partial v_\alpha}{\partial x_k} + \frac{\partial E_\sigma}{\partial x_k} + \varepsilon_{klj} \omega_l v_j = \frac{\partial w_k}{\partial t} + v_\alpha \frac{\partial w_\alpha}{\partial x_k} + w_\alpha \frac{\partial v_\alpha}{\partial x_k} + \frac{\partial E_\sigma}{\partial x_k} + \\ + v_\alpha \left(\frac{\partial w_k}{\partial x_\alpha} - \frac{\partial w_\alpha}{\partial x_k} \right) &= \frac{\partial w_k}{\partial t} + v_\alpha \left(\frac{\partial w_k}{\partial x_\alpha} \right) + w_\alpha \frac{\partial v_\alpha}{\partial x_k} + \frac{\partial E_\sigma}{\partial x_k} = \frac{dw_k}{dt} + w_\alpha \frac{\partial v_\alpha}{\partial x_k} + \frac{\partial E_\sigma}{\partial x_k} = \frac{dw_k}{dt} + A_{jk} \left(w_\alpha \frac{\partial v_\alpha}{\partial y_j} + \frac{\partial E_\sigma}{\partial y_j} \right) \end{aligned}$$

The intermediate result is

$$\frac{dw_k}{dt} + A_{jk} \left(w_\alpha \frac{\partial v_\alpha}{\partial y_j} + \frac{\partial E_\sigma}{\partial y_j} \right) = 0.$$

By multiplying the last equation by F^T , we obtain

$$F_{ki} \frac{dw_k}{dt} + F_{ki} A_{jk} \left(w_\alpha \frac{\partial v_\alpha}{\partial y_j} + \frac{\partial E_\sigma}{\partial y_j} \right) = 0.$$

Further, using the identity $A_{jk} F_{ki} = \delta_{ji}$ we can rewrite the last equations as

$$F_{ki} \frac{dw_k}{dt} + F_{ki} A_{jk} \left(w_\alpha \frac{\partial v_\alpha}{\partial y_j} + \frac{\partial E_\sigma}{\partial y_j} \right) = F_{ki} \frac{dw_k}{dt} + \left(w_\alpha \frac{\partial v_\alpha}{\partial y_i} + \frac{\partial E_\sigma}{\partial y_i} \right) = 0.$$

Then using (179), we get

$$F_{ki} \frac{dw_k}{dt} + \left(w_\alpha \frac{\partial v_\alpha}{\partial y_i} + \frac{\partial E_\sigma}{\partial y_i} \right) = F_{ki} \frac{dw_k}{dt} + \left(w_\alpha \frac{dF_{\alpha i}}{dt} + \frac{\partial E_\sigma}{\partial y_i} \right) = \frac{d(F_{ki} w_k)}{dt} + \frac{\partial E_\sigma}{\partial y_i} = 0$$

Then using that $U_{\sigma'} = (wE)_{w\sigma'} = E_{\sigma'}$, we get the final result, i.e. sought Lagrangian equation (18e),

$$\frac{dw'_k}{dt} + \frac{\partial U_{\sigma'}}{\partial y_k} = 0. \quad (189)$$

Therefore, equations (188) and (189) suggest that the Lagrangian fields w' , σ' and Eulerian w , σ are related by (176c).

A.3. (e, h) -pair, transformation of the nonlinear Maxwell equations

We shall transform (38d) into (18d) while (38c) transforms into (18c) analogously. Thus, (38d) is equivalent to

$$\frac{\partial e_i}{\partial t} + v_k \frac{\partial e_i}{\partial x_k} + \frac{\partial v_k}{\partial x_k} e_i - \frac{\partial v_i}{\partial x_k} e_k - \varepsilon_{ikl} \frac{\partial E_{h_l}}{\partial x_k} = 0. \quad (190)$$

Using $d/dt = \partial/\partial t + v_k/\partial x_k$ we have

$$\frac{de_i}{dt} + \frac{\partial v_k}{\partial x_k} e_i - \frac{\partial v_i}{\partial x_k} e_k - \varepsilon_{ikl} \frac{\partial E_{h_l}}{\partial x_k} = 0. \quad (191)$$

From (178) it follows that $\frac{\partial}{\partial x_k} = A_{jk} \frac{\partial}{\partial y_j}$ and thus we change the variables x_k on y_j

$$\frac{de_i}{dt} + A_{jk} \frac{\partial v_k}{\partial y_j} e_i - A_{jk} \frac{\partial v_i}{\partial y_j} e_k - \varepsilon_{ikl} A_{jk} \frac{\partial E_{h_l}}{\partial y_j} = 0. \quad (192)$$

Applying (182)₂ to the second term and (179) to the third term of the last equation, we have

$$\frac{de_i}{dt} + \frac{1}{w} \frac{dw}{dt} e_i - A_{jk} \frac{dF_{ij}}{dt} e_k - \varepsilon_{ikl} A_{jk} \frac{\partial E_{h_l}}{\partial y_j} = 0, \quad (193)$$

and then

$$\frac{dwe_i}{dt} - we_k A_{jk} \frac{dF_{ij}}{dt} - w\varepsilon_{ikl} A_{jk} \frac{\partial E_{h_l}}{\partial y_j} = 0. \quad (194)$$

Now, we add $0 \equiv we_k \frac{d\delta_{ik}}{dt}$ to the left hand side and then using $\delta_{ik} = F_{ij} A_{jk}$ one can obtain that

$$\frac{dwe_i}{dt} + we_k \frac{d\delta_{ik}}{dt} - we_k A_{jk} \frac{dF_{ij}}{dt} - w\varepsilon_{ikl} A_{jk} \frac{\partial E_{h_l}}{\partial y_j} = 0, \quad (195)$$

$$\frac{dwe_i}{dt} + we_k \frac{dF_{ij} A_{jk}}{dt} - we_k A_{jk} \frac{dF_{ij}}{dt} - w\varepsilon_{ikl} A_{jk} \frac{\partial E_{h_l}}{\partial y_j} = 0, \quad (196)$$

$$\frac{dwe_i}{dt} + we_k F_{ij} \frac{dA_{jk}}{dt} + we_k \frac{dF_{ij}}{dt} A_{jk} - we_k A_{jk} \frac{dF_{ij}}{dt} - w\varepsilon_{ikl} A_{jk} \frac{\partial E_{h_l}}{\partial y_j} = 0. \quad (197)$$

After multiplying the last equation by A_{mi}

$$A_{mi} \left(\frac{dwe_i}{dt} + we_k F_{ij} \frac{dA_{jk}}{dt} - w\varepsilon_{ikl} A_{jk} \frac{\partial E_{h_l}}{\partial y_j} \right) = 0, \quad (198)$$

$$A_{mi} \frac{dwe_i}{dt} + we_k \frac{dA_{mk}}{dt} - w\varepsilon_{ikl} A_{mi} A_{jk} \frac{\partial E_{h_l}}{\partial y_j} = 0, \quad (199)$$

we have an intermediate result:

$$\frac{dwA_{mke_k}}{dt} - w\varepsilon_{ikl} A_{mi} A_{jk} \frac{\partial E_{h_l}}{\partial y_j} = 0. \quad (200)$$

Now, we introduce the change of unknowns (176b): $wA_{mke_k} = e'_m$, $wA_{mk}h_k = h'_m$, and we also change the energy potential $E(e_i, h_i) = E(w^{-1}F_{ij}e'_j, w^{-1}F_{ij}h'_j) = w^{-1}U(e'_j, h'_j)$. Hence, $E_{h_i} = A_{ji}U_{h'_j}$. After this, the intermediate result (200) reads as

$$\frac{de'_m}{dt} - w\varepsilon_{ikl} A_{mi} A_{jk} \frac{\partial A_{al}U_{h'_a}}{\partial y_j} = 0, \quad (201)$$

$$\frac{de'_m}{dt} - w\varepsilon_{ikl} A_{mi} A_{jk} A_{al} \frac{\partial U_{h'_a}}{\partial y_j} - w\varepsilon_{ikl} A_{mi} A_{jk} \frac{\partial A_{al}}{\partial y_j} U_{h'_a} = 0. \quad (202)$$

Applying (184) and (180)₃ to the second term, we get

$$\frac{de'_m}{dt} - \varepsilon_{mja} \frac{\partial U_{h'_a}}{\partial y_j} - w\varepsilon_{ikl} A_{mi} A_{jk} \frac{\partial A_{al}}{\partial y_j} U_{h'_a} = 0, \quad (203)$$

Now using the cofactor definition (180) and then applying (185) to the third term, we have

$$\frac{de'_m}{dt} - \varepsilon_{mja} \frac{\partial U_{h'_a}}{\partial y_j} - \varepsilon_{ikl} A_{mi} C_{kj} \frac{\partial A_{al}}{\partial y_j} U_{h'_a} = 0, \quad (204)$$

$$\frac{de'_m}{dt} - \varepsilon_{mja} \frac{\partial U_{h'_a}}{\partial y_j} - \frac{1}{2} \varepsilon_{ikl} \varepsilon_{lnk} \varepsilon_{pqj} F_{lp} F_{nq} A_{mi} \frac{\partial A_{al}}{\partial y_j} U_{h'_a} = 0. \quad (205)$$

Using (186) in the third term gives us

$$\frac{de'_m}{dt} - \varepsilon_{mja} \frac{\partial U_{h'_a}}{\partial y_j} - \delta_{in} \varepsilon_{pqj} F_{lp} F_{nq} A_{mi} \frac{\partial A_{al}}{\partial y_j} U_{h'_a} = 0, \quad (206)$$

and subsequently,

$$\frac{de'_m}{dt} - \varepsilon_{mja} \frac{\partial U_{h'_a}}{\partial y_j} - \varepsilon_{pqj} F_{lp} F_{iq} A_{mi} \frac{\partial A_{al}}{\partial y_j} U_{h'_a} = 0, \quad (207)$$

$$\frac{de'_m}{dt} - \varepsilon_{mja} \frac{\partial U_{h'_a}}{\partial y_j} - \varepsilon_{mj p} F_{lp} \frac{\partial A_{al}}{\partial y_j} U_{h'_a} = 0. \quad (208)$$

Now, adding $0 \equiv \varepsilon_{mj p} \frac{\partial F_{lp}}{\partial y_j} A_{al} U_{h'_a}$ (see (181)₂), we get

$$\frac{de'_m}{dt} - \varepsilon_{mja} \frac{\partial U_{h'_a}}{\partial y_j} - \varepsilon_{mj p} F_{lp} \frac{\partial A_{al}}{\partial y_j} U_{h'_a} - \varepsilon_{mj p} \frac{\partial F_{lp}}{\partial y_j} A_{al} U_{h'_a} = 0, \quad (209)$$

$$\frac{de'_m}{dt} - \varepsilon_{mja} \frac{\partial U_{h'_a}}{\partial y_j} - U_{h'_a} \left(\varepsilon_{mj p} \frac{\partial A_{al} F_{lp}}{\partial y_j} \right) = 0, \quad (210)$$

$$\frac{de'_m}{dt} - \varepsilon_{mja} \frac{\partial U_{h'_a}}{\partial y_j} - U_{h'_a} \left(\varepsilon_{mj p} \frac{\partial \delta_{ap}}{\partial y_j} \right) = 0. \quad (211)$$

Eventually, we have

$$\frac{de'_m}{dt} - \varepsilon_{mja} \frac{\partial U_{h'_a}}{\partial y_j} = 0, \quad (212)$$

which is identical to (18d).

A.4. (m, F) -pair

A.4.1 Distortion matrix

Here, we derive Eulerian equation (38b) from its Lagrangian form (18a).

Recall that the velocity field v_i is conjugate to the momentum field, i.e. $v_i = U_{m_i}$. Thus, (18b) reads as

$$\frac{dF_{ij}}{dt} - \frac{\partial v_i}{\partial y_j} = 0. \quad (213)$$

Then, applying (178)₁ and (178)₅ we arrive at

$$\frac{\partial F_{ij}}{\partial t} + v_k \frac{\partial F_{ij}}{\partial x_k} - F_{kj} \frac{\partial v_i}{\partial x_k} = 0. \quad (214)$$

Using the identity

$$\mathbf{0} \equiv d\mathbf{I} \equiv d(\mathbf{F}\mathbf{A}) \equiv \mathbf{F}d\mathbf{A} + (d\mathbf{F})\mathbf{A}, \quad (215)$$

(214) can be rewritten as

$$\frac{\partial A_{ij}}{\partial t} + v_k \frac{\partial A_{ij}}{\partial x_k} + A_{ik} \frac{\partial v_k}{\partial x_j} = 0, \quad (216)$$

and eventually as

$$\frac{\partial A_{ik}}{\partial t} + \frac{\partial(A_{im}v_m)}{\partial x_k} + v_j \left(\frac{\partial A_{ik}}{\partial x_j} - \frac{\partial A_{ij}}{\partial x_k} \right) = 0, \quad (217)$$

which is the sought equation (38b).

A.4.2 Transformation of the momentum conservation

In this section, we demonstrate that Eulerian momentum conservation law (38a) can be derived from its Lagrangian counterpart (18a). As earlier, we use prime to denote Lagrangian fields. Now, (18a) reads as

$$\frac{dm'_i}{dt} - \frac{\partial U_{F_{ij}}}{\partial y_j} = 0. \quad (218)$$

Using that $d/dt = \partial/\partial t + v_k \partial/\partial x_k$ and $F_{ij} = \partial x_i / \partial y_j$, equation (218) can be rewritten as

$$\frac{\partial m'_i}{\partial t} + v_k \frac{\partial m'_i}{\partial x_k} - F_{kj} \frac{\partial U_{F_{ij}}}{\partial x_k} = 0. \quad (219)$$

Subsequently, using the Eulerian stationary constraint for F_{ij} and time evolution of $w = \det(\mathbf{F})$ (183):

$$\frac{\partial w^{-1} F_{kj}}{\partial x_k} = 0, \quad \frac{\partial}{\partial t} \left(\frac{1}{w} \right) + \frac{\partial}{\partial x_k} \left(\frac{v_k}{w} \right) = 0, \quad (220)$$

(219) can be rewritten as

$$\frac{\partial}{\partial t} \left(\frac{m'_i}{w} \right) + \frac{\partial}{\partial x_k} \left(\frac{v_k m'_i - F_{kj} U_{F_{ij}}}{w} \right) = 0. \quad (221)$$

We now introduce change of the potential

$$U = w \mathcal{U},$$

and change of the momentum (176a), $m_i = w^{-1} m'_i$. Note that \mathcal{U} equals to E but depends on the Lagrangian fields, i.e.

$$\mathcal{U}(F_{ij}, m'_i, e'_i, h'_i, \sigma', w'_i) = E(\rho, F_{ij}, m_i, e_i, h_i, \sigma, w_i). \quad (222)$$

After that, we arrive at

$$\frac{\partial m_i}{\partial t} + \frac{\partial}{\partial x_k} \left(v_k m_i - \mathcal{U} \delta_{ki} - F_{kj} \mathcal{U}_{F_{ij}} \right) = 0. \quad (223)$$

Finally, it remains to transform the term $F_{kj} \mathcal{U}_{F_{ij}}$ in the flux of equation (223). Thus, according to (222) and change of the variables (176a), (176b) and (176c): $m'_a = w m_a$, $e'_b = w A_{ba} e_a$, $h'_b = w A_{ba} h_a$, $\sigma' = w \sigma$ and $w'_b = F_{ab} w_a$, we have

$$\mathcal{U}_{F_{ij}} = \frac{\partial}{\partial F_{ij}} \left(E(\rho, F_{ab}, w^{-1} m'_a, w^{-1} F_{ab} e'_b, w^{-1} F_{ab} h'_b, w^{-1} \sigma', A_{ba} w'_b) \right), \quad (224)$$

and hence

$$F_{kj} \mathcal{U}_{F_{ij}} = F_{kj} \left(-\rho A_{ji} E_\rho + E_{F_{ij}} + E_{m_a} (w^{-1} m'_a)_{F_{ij}} + E_{e_a} (w^{-1} F_{ab} e'_b)_{F_{ij}} + E_\sigma (w^{-1} \sigma')_{F_{ij}} + E_{w_a} (A_{ba} w'_b)_{F_{ij}} \right). \quad (225)$$

After using the formulas

$$\frac{\partial w}{\partial F_{ij}} = w A_{ji}, \quad \frac{\partial A_{ab}}{\partial F_{cd}} = -A_{ac} A_{db}, \quad (226)$$

and doing some trivial algebra, we arrive at

$$F_{kj} \mathcal{U}_{F_{ij}} = -(\rho E_\rho + m_l E_{m_l} + \sigma E_\sigma + e_l E_{e_l} + h_l E_{h_l}) \delta_{ki} + e_k E_{e_i} + h_k E_{h_i} - w_i E_{w_k} + F_{kj} E_{F_{ij}}. \quad (227)$$

Eventually, using that $F_{kj} E_{F_{ij}} = -A_{jk} E_{A_{ji}}$, the momentum equation becomes

$$\frac{\partial m_i}{\partial t} + \frac{\partial}{\partial x_k} \left(m_i E_{m_k} + (\rho E_\rho + m_l E_{m_l} + e_l E_{e_l} + h_l E_{h_l} + \sigma E_\sigma - E) \delta_{ki} - e_k E_{e_i} - h_k E_{h_i} + w_i E_{w_k} + A_{jk} E_{A_{ji}} \right) = 0. \quad (228)$$

B. DERIVATION OF THE ENERGY CONSERVATION LAW

In this section, we prove that the energy conservation law (59)

$$\begin{aligned} & \frac{\partial}{\partial t} (r L_r + v_i L_{v_i} + \alpha_{ij} L_{\alpha_{ij}} + d_i L_{d_i} + b_i L_{b_i} + \theta L_\theta + \eta_k L_{\eta_k} - L) + \\ & \frac{\partial}{\partial x_k} \left(r L_r + v_i L_{v_i} + \alpha_{ij} L_{\alpha_{ij}} + d_i L_{d_i} + b_i L_{b_i} + \theta L_\theta + \eta_k L_{\eta_k} - L \right) + \\ & v_i \left[(L - \alpha_{ab} L_{\alpha_{ab}} - \eta_a L_{\eta_a}) \delta_{ik} + \alpha_{nk} L_{\alpha_{ni}} - d_i L_{d_k} - b_i L_{b_k} + \eta_k L_{\eta_i} \right] + \varepsilon_{kij} d_i b_j + \theta \eta_k \end{aligned} = 0.$$

is compatible with the governing equations (58), that is it can be obtained as the consequence of all the equations (58) multiplied by the corresponding conjugate state variables (56). From this prove, it will be clear that the energy flux, in fact, is not the consequence of only the fluxes in (58), but the non-conservative terms in equations (58b), (58d),

(58c) and (58e) contribute into the energy flux as well.

It is obvious that the time derivative of the energy conservation can be obtained as the sum of the time derivatives of the equations (58) multiplied by the corresponding conjugate fields, i.e.

$$\frac{\partial(p_i L_{p_i} - L)}{\partial t} = p_i \frac{\partial L_{p_i}}{\partial t}, \quad (229)$$

while the main challenge is to show that the energy flux is

$$\begin{aligned} \text{flux}(59) = & r \cdot \text{flux}(58g) + v_i \cdot \text{flux}(58a) + \alpha_{ik} \cdot \left(\text{flux}(58b) + v_j (\partial_j L_{\alpha_{ik}} - \partial_k L_{\alpha_{ij}}) \right) + \\ & d_i \cdot \left(\text{flux}(58d) + v_i \partial_k L_{d_k} \right) + b_i \cdot \left(\text{flux}(58c) + v_i \partial_k L_{b_k} \right) + \\ & \theta \cdot \text{flux}(58f) + \eta_k \cdot \left(\text{flux}(58e) + v_j (\partial_j L_{\eta_k} - \partial_k L_{\eta_j}) \right). \end{aligned} \quad (230)$$

The right-hand side of (230) is

$$\begin{aligned} & r \partial_k (v_k L_r) + v_i \partial_k (\delta_{ik} L + v_k L_{v_i} + \alpha_{mk} L_{\alpha_{mi}} - \delta_{ik} \alpha_{mn} L_{\alpha_{mn}} - d_i L_{d_k} - b_i L_{b_k} + \eta_k L_{\eta_i} - \delta_{ik} \eta_n L_{\eta_n}) + \\ & \alpha_{ik} \left(\partial_k (v_m L_{\alpha_{im}}) + v_j (\partial_j L_{\alpha_{ik}} - \partial_k L_{\alpha_{ij}}) \right) + \\ & d_i (\partial_k (v_k L_{d_i} - v_i L_{d_k} - \varepsilon_{ikl} b_l) + v_i \partial_k L_{d_k}) + b_i (\partial_k (v_k L_{b_i} - v_i L_{b_k} + \varepsilon_{ikl} d_l) + v_i \partial_k L_{b_k}) + \\ & \theta \partial_k (v_k L_\theta + \eta_k) + \eta_k (\partial_k (v_i L_{\eta_i} + \theta) + v_j (\partial_j L_{\eta_k} - \partial_k L_{\eta_j})) = \end{aligned} \quad (231)$$

$$\begin{aligned} & r \partial_k (v_k L_r) + v_i \partial_k (v_k L_{v_i}) + \alpha_{mi} \partial_k (v_k L_{\alpha_{mi}}) + d_i \partial_k (v_k L_{d_i}) + b_i \partial_k (v_k L_{b_i}) + \theta \partial_k (v_k L_\theta) + \eta_i \partial_k (v_k L_{\eta_i}) - \\ & v_i \partial_k (\delta_{ik} \alpha_{mn} L_{\alpha_{mn}} - \alpha_{mk} L_{\alpha_{mi}} + d_i L_{d_k} + b_i L_{b_k} + \delta_{ik} \eta_n L_{\eta_n} - \eta_k L_{\eta_i}) - \\ & (\delta_{ik} \alpha_{mn} L_{\alpha_{mn}} - \alpha_{mk} L_{\alpha_{mi}} + d_i L_{d_k} + b_i L_{b_k} + \delta_{ik} \eta_n L_{\eta_n} - \eta_k L_{\eta_i}) \partial_k v_i + \\ & v_k \partial_k L - \varepsilon_{ikl} d_i \partial_k b_l + \varepsilon_{ikl} b_i \partial_k d_l + \theta \partial_k \eta_k + \eta_k \partial_k \theta = \end{aligned} \quad (232)$$

$$\begin{aligned} & \partial_k (v_k (r L_r + v_i L_{v_i} + \alpha_{mi} L_{\alpha_{mi}} + d_i L_{d_i} + b_i L_{b_i} + \theta L_\theta + \eta_k L_{\eta_k})) - \\ & v_k (L_r \partial_k r + L_{v_i} \partial_k v_i + L_{\alpha_{mi}} \partial_k \alpha_{mi} + L_{d_i} \partial_k d_i + L_{b_i} \partial_k b_i + L_\theta \partial_k \theta + L_{\eta_k} \partial_k \eta_k) + v_k \partial_k L - \\ & \partial_k (v_i (\delta_{ik} \alpha_{mn} L_{\alpha_{mn}} - \alpha_{mk} L_{\alpha_{mi}} + d_i L_{d_k} + b_i L_{b_k} + \delta_{ik} \eta_n L_{\eta_n} - \eta_k L_{\eta_i})) + \partial_k (\varepsilon_{kij} d_i b_j + \theta \eta_k), \end{aligned} \quad (233)$$

which eventually transforms exactly into the energy flux (59).

C. DERIVATION OF THE TIME EVOLUTION FOR $\nabla \times w$ AND $\nabla \times A$

Here, we derive the time evolutions (46a) and (46d) for $\nabla \times A$ and $\nabla \times w$ accordingly. In fact, because the time evolution for each column A_{ik} , $k = 1, 2, 3$ has the same structure as the time evolution for w_i (apart of the constitutive flux E_σ) we can perform the derivation only for w_i .

We start by rewriting (38e) (we also assume the presence of an algebraic source term $-f_k$ in the right-hand side)

$$\frac{\partial w_k}{\partial t} + \partial_k (v_a w_a + E_\sigma) + v_j (\partial_j w_k - \partial_k w_j) = -f_k \quad (234)$$

in an equivalent form

$$\frac{\partial w_k}{\partial t} + w_a \partial_k v_a + \partial_k E_\sigma + v_a \partial_a w_k = -f_k.$$

Then, applying ∂_m to the k -th equation and ∂_k to the m -th equation and subtracting the later from the former we get

$$\frac{\partial}{\partial t} (\partial_m w_k - \partial_k w_m) + v_a \partial_a (\partial_m w_k - \partial_k w_m) + \partial_m w_a \partial_k v_a - \partial_k w_a \partial_m v_a - \partial_a w_m \partial_k v_a + \partial_a w_k \partial_m v_a = -(\partial_m f_k - \partial_k f_m). \quad (235)$$

This equation can be written in an elegant form if all the derivatives of the components of w_i are arranged in such a way that the resulting equation contains only the components of the vector $\Omega = \nabla \times \mathbf{w}$. Indeed, it suffices to explain how to handle with last four terms in the left-hand side of (235) for some m and k . We consider only the case $m = 1, k = 2$ and write explicitly the summation with respect to a :

$$\begin{aligned} & \partial_1 w_a \partial_2 v_a + \partial_a w_2 \partial_1 v_a - \partial_2 w_a \partial_1 v_a - \partial_a w_1 \partial_2 v_a = \\ & \partial_1 w_1 \partial_2 v_1 + \partial_1 w_2 \partial_1 v_1 - \partial_2 w_1 \partial_1 v_1 - \partial_1 w_1 \partial_2 v_1 + \partial_1 w_2 \partial_2 v_2 + \partial_2 w_2 \partial_1 v_2 - \\ & \partial_2 w_2 \partial_1 v_2 - \partial_2 w_1 \partial_2 v_2 + \partial_1 w_3 \partial_2 v_3 + \partial_3 w_2 \partial_1 v_3 - \partial_2 w_3 \partial_1 v_3 - \partial_3 w_1 \partial_2 v_3 = \\ & (\partial_1 w_2 - \partial_2 w_1) \partial_1 v_1 + (\partial_3 w_2 - \partial_2 w_3) \partial_1 v_3 + (\partial_1 w_3 - \partial_3 w_1) \partial_2 v_3 + (\partial_1 w_2 - \partial_2 w_1) \partial_2 v_2 = \\ & (\partial_1 w_2 - \partial_2 w_1) (\partial_1 v_1 + \partial_2 v_2) + (\partial_3 w_2 - \partial_2 w_3) \partial_1 v_3 + (\partial_1 w_3 - \partial_3 w_1) \partial_2 v_3 = \\ & \Omega_3 (\partial_1 v_1 + \partial_2 v_2) - \Omega_1 \partial_1 v_3 - \Omega_2 \partial_2 v_3 = \Omega_3 (\partial_1 v_1 + \partial_2 v_2 + \partial_3 v_3) - \Omega_1 \partial_1 v_3 - \Omega_2 \partial_2 v_3 - \Omega_3 \partial_3 v_3 = \\ & \Omega_3 \partial_a v_a - \Omega_a \partial_a v_3. \end{aligned} \quad (236)$$

Finally, we conclude that

$$\frac{\partial \Omega_i}{\partial t} + v_k \partial_k \Omega_i + \Omega_i \partial_k v_k - \Omega_k \partial_k v_i + \varepsilon_{ikl} \partial_k f_l = 0 \quad (237)$$

or equivalently

$$\frac{\partial \Omega_i}{\partial t} + \partial_k (\Omega_i v_k - \Omega_k v_i + \varepsilon_{ikl} f_l) + v_i \partial_k \Omega_k = 0, \quad (238)$$

which apparently has the same structure as equation (38c) of the Eulerian SHTC system and the sought equation (44d) but with an additional term in the flux conditioned by the presence of the source term f_k in (234).

Eventually we make an important remark. By construction, $\partial_k \omega_k = 0$ because $\Omega = \nabla \times \mathbf{w}$ and hence, one may want to write (238) in a conservative form by eliminating the non-conservative term $v_i \partial_k \Omega_k = 0$. This however should be avoided because the obtained conservative equation

$$\frac{\partial \Omega_i}{\partial t} + \partial_k (v_k \Omega_i - \Omega_k v_i + \varepsilon_{ikl} f_l) = 0 \quad (239)$$

has different characteristic speeds, e.g. see [79], and thus is not equivalent to the original equations (237) and (238). Moreover, in contrast to (237) and (238), conservative equation (239) is not Galilean invariant.

REFERENCES

- [1] Abarbanel, H.D.I., Brown, R., Yang, Y.M.: Hamiltonian formulation of inviscid flows with free boundaries. *Physics of Fluids* **31**, 2802 (1988) (Cited on page 8.)
- [2] Arnold, V.I.: Sur la géométrie différentielle des groupes de Lie de dimension infini et ses applications dans l'hydrodynamique des fluides parfaits. *Ann. Inst. Fourier* **16**, 319 (1966) (Cited on pages 3 and 24.)

- [3] Barton, P.T., Deiterding, R., Meiron, D., Pullin, D.: Eulerian adaptive finite-difference method for high-velocity impact and penetration problems. *Journal of Computational Physics* **240**, 76–99 (2013) (Cited on page 15.)
- [4] Barton, P.T., Drikakis, D., Romenski, E.I.: An Eulerian finite-volume scheme for large elastoplastic deformations in solids. *International journal for numerical methods in engineering* **81(4)**, 453–484 (2010) (Cited on pages 15, 26, 37, and 43.)
- [5] Benzoni-Gavage, S., Serre, D.: *Multidimensional Hyperbolic Partial Differential Equations*. Oxford University Press, Oxford (2007) (Cited on pages 4 and 5.)
- [6] Berezovski, A., Ván, P.: *Internal Variables in Thermoelasticity*. Solid Mechanics and Its Applications. Springer International Publishing (2017). URL <https://books.google.de/books?id=vv3MDgAAQBAJ> (Cited on page 34.)
- [7] Beris, A.N., Edwards, B.J.: *Thermodynamics of Flowing Systems: With Internal Microstructure*. Oxford University Press, USA (1994) (Cited on page 3.)
- [8] Bobylev, A.: The Chapman-Enskog and Grad methods for solving the Boltzmann equation. *Akademiia Nauk SSSR Doklady* **262**, 71–75 (1982) (Cited on page 4.)
- [9] Boillat, G.: Sur l'existence et la recherche d'équations de conservation supplémentaires pour les systèmes hyperboliques. *C. R. Acad. Sc. Paris, Sér A* **278** (1974) (Cited on page 5.)
- [10] Bolmatov, D., Brazhkin, V.V., Trachenko, K.: Thermodynamic behaviour of supercritical matter. *Nature communications* **4** (2013) (Cited on page 43.)
- [11] Bolmatov, D., Zav'yalov, D., Zhernenkov, M., Musaev, E.T., Cai, Y.Q.: Unified phonon-based approach to the thermodynamics of solid, liquid and gas states. *Annals of Physics* **363**, 221–242 (2015). DOI 10.1016/j.aop.2015.09.018. URL <http://dx.doi.org/10.1016/j.aop.2015.09.018> (Cited on page 43.)
- [12] Bolmatov, D., Zhernenkov, M., Zav'yalov, D., Stoupin, S., Cai, Y.Q., Cunsolo, A.: Revealing the Mechanism of the Viscous-to-Elastic Crossover in Liquids. *The journal of physical chemistry letters* **6(15)**, 3048–3053 (2015) (Cited on page 43.)
- [13] Bolmatov, D., Zhernenkov, M., Zav'yalov, D., Stoupin, S., Cunsolo, A., Cai, Y.Q.: Thermally triggered phononic gaps in liquids at THz scale. *Scientific Reports* **6(November 2015)**, 19,469 (2016). DOI 10.1038/srep19469. URL <http://www.nature.com/articles/srep19469> (Cited on page 43.)
- [14] Boscheri, W., Dumbser, M., Loubère, R.: Cell centered direct Arbitrary-Lagrangian-Eulerian ADER-WENO finite volume schemes for nonlinear hyperelasticity. *Computers & Fluids* **134-135**, 111–129 (2016). DOI 10.1016/j.compfluid.2016.05.004. URL <http://linkinghub.elsevier.com/retrieve/pii/S004579301630144X> (Cited on pages 15, 37, and 43.)
- [15] Brazhkin, V.V., Fomin, Y.D., Lyapin, A.G., Ryzhov, V.N., Trachenko, K.: Two liquid states of matter: A dynamic line on a phase diagram. *Physical Review E* **85(3)**, 31,203 (2012) (Cited on page 43.)
- [16] Clebsch, A.: Über die Integration der hydrodynamische Gleichungen. *J. Reine Angew. Math.* **56(1)** (1859) (Cited on pages 3 and 24.)
- [17] Dumbser, M., Peshkov, I., Romenski, E., Zanotti, O.: High order ADER schemes for a unified first order hyperbolic formulation of continuum mechanics: Viscous heat-conducting fluids and elastic solids. *Journal of Computational Physics* **314**, 824–862 (2016). DOI 10.1016/j.jcp.2016.02.015. URL <http://www.sciencedirect.com/science/article/pii/S0021999116000693> (Cited on pages 15, 26, 28, 36, 40, 41, 43, 44, and 47.)

- [18] Dumbser, M., Peshkov, I., Romenski, E., Zanotti, O.: High order ADER schemes for a unified first order hyperbolic formulation of Newtonian continuum mechanics coupled with electro-dynamics. *Journal of Computational Physics* **348**, 298–342 (2017). DOI 10.1016/j.jcp.2017.07.020. URL <http://dx.doi.org/10.1016/j.jcp.2017.07.020><http://www.sciencedirect.com/science/article/pii/S0021999117> (Cited on pages 9, 12, 15, 30, 31, 36, 40, and 47.)
- [19] Dupret, F., Marchal, J.: Loss of evolution in the flow of viscoelastic fluids. *Journal of Non-Newtonian Fluid Mechanics* **20**, 143–171 (1986) (Cited on page 4.)
- [20] Dzyaloshinskii, I.E., Volovick, G.E.: Poisson brackets in condensed matter physics. *Annals of Physics* **125**(1), 67–97 (1980). DOI 10.1016/0003-4916(80)90119-0 (Cited on page 3.)
- [21] E. I. Romenski: Thermodynamics and Balance Laws for Processes of Inelastic Deformations. In: Proceedings "WASCOM 2001" 11th Conference on Waves and Stability in Continuous Media, pp. 484–495. World Scientific (2002) (Cited on pages 6, 8, 21, and 23.)
- [22] Esen, O., Pavelka, M., Grmela, M.: Hamiltonian coupling of electromagnetic field and matter. *International Journal of Advances in Engineering Sciences and Applied Mathematics* (2017). DOI 10.1007/s12572-017-0179-4. URL <http://link.springer.com/10.1007/s12572-017-0179-4> (Cited on pages 27, 31, and 47.)
- [23] Favrie, N., Gavriluyuk, S.: Dynamics of shock waves in elastic-plastic solids. In: ESAIM proceedings, vol. 30, pp. 50–67 (2011). DOI 10.1051/proc/201133005. URL <https://www.esaim-proc.org/component/makeref/?task=show{&}type=html{&}doi=10.1051/proc/201133005> (Cited on pages 15, 26, 37, and 43.)
- [24] Favrie, N., Gavriluyuk, S.: A rapid numerical method for solving Serreâ'sGreenâ'sNaghdi equations describing long free surface gravity waves. *Nonlinearity* **30**(7), 2718–2736 (2017). DOI 10.1088/1361-6544/aa712d. URL <http://stacks.iop.org/0951-7715/30/i=7/a=2718?key=crossref.139c98587b84970534e28823dcd579eb> (Cited on page 40.)
- [25] Fecko, M.: *Differential Geometry and Lie Groups for Physicists*. Cambridge University Press (2006). URL <https://books.google.de/books?id=vQR0mN1dgUEC> (Cited on page 24.)
- [26] Frenkel, J.: *Kinetic theory of liquids*. Dover (1955) (Cited on page 43.)
- [27] Friedrichs, K.O.: Symmetric positive linear differential equations. *Communications on Pure and Applied Mathematics* **11**(3), 333–418 (1958) (Cited on page 5.)
- [28] Friedrichs, K.O., Lax, P.D.: Systems of conservation equations with a convex extension. *Proceedings of the National Academy of Sciences* **68**(8), 1686–1688 (1971) (Cited on page 5.)
- [29] Gavriluyuk, S.L., Makarenko, N.I., Sukhinin, S.V.: *Waves in Continuous Media*. Lecture Notes in Geosystems Mathematics and Computing. Springer International Publishing (2017) (Cited on page 5.)
- [30] Godunov, S., Mikhailova, T., Romenskii, E.: Systems of thermodynamically coordinated laws of conservation invariant under rotations. *Siberian Mathematical Journal* **37**(4), 690–705 (1996) (Cited on pages 6, 8, 9, 11, 14, 24, and 48.)
- [31] Godunov, S., Peshkov, I.: Thermodynamically Consistent Nonlinear Model of Elastoplastic Maxwell Medium. *Computational Mathematics and Mathematical Physics* **50**(8), 1409–1426 (2010). DOI 10.1134/S0965542510080117 (Cited on pages 11, 26, and 37.)

- [32] Godunov, S., Romenskiy, E.: Thermodynamics, conservation laws and symmetric forms of differential equations in mechanics of continuous media. In: *Computational Fluid Dynamics Review 1995*, vol. 95, pp. 19–31. John Wiley, NY (1995). DOI 10.1142/7799 (Cited on pages 6 and 8.)
- [33] Godunov, S., Yu Mikhailova, T., Romenskii, E.: Systems of thermodynamically coordinated laws of conservation invariant under rotations. *Siberian Mathematical Journal* **37**(4), 790–806 (1996) (Cited on page 19.)
- [34] Godunov, S.K.: A difference method for numerical calculation of discontinuous solutions of the equations of hydrodynamics. *Matematicheskii Sbornik* **89**(3), 271–306 (1959) (Cited on page 3.)
- [35] Godunov, S.K.: An interesting class of quasilinear systems. *Dokl. Akad. Nauk SSSR* **139**(3), 521–523 (1961) (Cited on pages 3, 4, and 5.)
- [36] Godunov, S.K.: The problem of a generalized solution in the theory of quasilinear equations and in gas dynamics. *Russian Mathematical Surveys* **17**(3), 145–156 (1962) (Cited on pages 3 and 5.)
- [37] Godunov, S.K.: Symmetric form of the magnetohydrodynamic equation. *Numerical Methods for Mechanics of Continuum Medium* **3**(1), 26–34 (1972). URL <http://citeseerx.ist.psu.edu/viewdoc/download?doi=10.1.1.55.9645{%&}rep=rep1{%&}type=pdf> (Cited on page 5.)
- [38] Godunov, S.K.: Symmetric form of the magnetohydrodynamic equation. *Numerical Methods for Mechanics of Continuum Medium* **3**(1), 26–34 (1972). URL <http://citeseerx.ist.psu.edu/viewdoc/download?doi=10.1.1.55.9645{%&}rep=rep1{%&}type=pdf> (Cited on pages 6, 15, 18, and 19.)
- [39] Godunov, S.K.: *Elements of mechanics of continuous media*, 1st russia edn. Nauka (1978) (Cited on pages 26, 28, 37, 42, and 43.)
- [40] Godunov, S.K., Romenskii, E.I.: Nonstationary equations of nonlinear elasticity theory in Eulerian coordinates. *Journal of Applied Mechanics and Technical Physics* **13**(6), 868–884 (1972) (Cited on page 26.)
- [41] Godunov, S.K., Romenskii, E.I.: *Elements of continuum mechanics and conservation laws*. Kluwer Academic/Plenum Publishers (2003) (Cited on pages 6, 8, 12, 15, 16, 28, 29, 37, 42, and 43.)
- [42] Godunov, S.K., Romenskii, E.I.: *Elements of mechanics of continuous media*. Kluwer Academic/Plenum Publishers, New York (2003) (Cited on pages 6, 8, 12, 15, 16, 28, 37, 42, and 43.)
- [43] Godunov, S.K., Romenskiy, E.I.: Symmetric forms of thermodynamically compatible systems of conservation laws in continuum mechanics. In: *ECCOMAS Conference on numerical methods in engineering*, pp. 54–57 (1996) (Cited on pages 6 and 8.)
- [44] Grmela, M.: Particle and Bracket Formulations of Kinetic Equations. *Contemporary Mathematics* **28**, 125–132 (1984) (Cited on page 3.)
- [45] Grmela, M.: Bracket formulation of diffusion-convection equations. *Phys. D* **21**, 179–212 (1986) (Cited on pages 3 and 24.)
- [46] Grmela, M.: A framework for elasto-plastic hydrodynamics. *Physics letters A* **312**, 134–146 (2003) (Cited on pages 3 and 26.)
- [47] Grmela, M.: Fluctuations in extended mass-action-law dynamics. *Physica D: Nonlinear Phenomena* **241**(10), 976–986 (2012) (Cited on page 8.)

- [48] Grmela, M.: Contact geometry of mesoscopic thermodynamics and dynamics. *Entropy* **16**(3), 1652–1686 (2014). DOI 10.3390/e16031652 (Cited on page 40.)
- [49] Grmela, M., Lebon, G., Dubois, C.: Multiscale thermodynamics and mechanics of heat. *Physical Review E - Statistical, Nonlinear, and Soft Matter Physics* **83**(6), 1–15 (2011). DOI 10.1103/PhysRevE.83.061134 (Cited on page 34.)
- [50] Grmela, M., Öttinger, H.C.: Dynamics and thermodynamics of complex fluids. I. Development of a general formalism. *Physical Review E* **56**(6), 6620–6632 (1997). DOI 10.1103/PhysRevE.56.6620 (Cited on pages 3, 6, 8, and 29.)
- [51] de Groot, S.R., Mazur, P.: *Non-equilibrium Thermodynamics*. Dover Books on Physics. Dover Publications (1984) (Cited on pages 8 and 40.)
- [52] Gurtin, M.E.: *An Introduction to Continuum Mechanics*. Mathematics in Science and Engineering. Elsevier Science (1982) (Cited on page 25.)
- [53] Holm, D.D.: Hamiltonian dynamics of a charged fluid, including electro-and magnetohydrodynamics. *Phys. Lett. A* **114**(3), 137–141 (1986) (Cited on page 31.)
- [54] Hron, J., Miloš, V., Průruša, V., Souček, O., Třřuma, K.: On thermodynamics of viscoelastic rate type fluids with temperature dependent material coefficients. *Int. J. Non-Linear Mech.* **95**, 193–208 (2017). DOI 10.1016/j.ijnonlinmec.2017.06.011 (Cited on page 29.)
- [55] Hřřtter, M., Svendsen, B.: Thermodynamic model formulation for viscoplastic solids as general equations for non-equilibrium reversible–irreversible coupling. *Continuum Mechanics and Thermodynamics* **24**(3), 211–227 (2012). DOI 10.1007/s00161-011-0232-7. URL <http://link.springer.com/10.1007/s00161-011-0232-7> (Cited on page 26.)
- [56] Hřřtter, M., Svendsen, B.: Quasi-linear versus potential-based formulations of force–flux relations and the GENERIC for irreversible processes: comparisons and examples. *Continuum Mechanics and Thermodynamics* **25**(6), 803–816 (2013). DOI 10.1007/s00161-012-0289-y (Cited on page 36.)
- [57] Janečka, A., Pavelka, M.: Gradient dynamics and entropy production maximization. *Journal of Non-Equilibrium Thermodynamics* (2017) (Cited on page 36.)
- [58] Kato, T.: The Cauchy problem for quasi-linear symmetric hyperbolic systems. *Archive for Rational Mechanics and Analysis* **58**(3), 181–205 (1975). DOI 10.1007/BF00280740. URL <http://link.springer.com/10.1007/BF00280740> (Cited on page 5.)
- [59] Kaufman, A.N.: Dissipative Hamiltonian systems: A unifying principle. *Physics Letters A* **100**(8), 419–422 (1984) (Cited on page 3.)
- [60] Kroeger, M., Huetter, M.: Automated symbolic calculations in nonequilibrium thermodynamics. *Comput. Phys. Commun.* **181**, 2149–2157 (2010) (Cited on pages 28, 33, and 35.)
- [61] Landau, L.D., Lifshitz, E.M.: *Fluid Mechanics, Course of Theoretical Physics, Volume 6*. Elsevier Butterworth-Heinemann, Oxford (2004) (Cited on page 6.)
- [62] Landau L.D. Lifshitz, E.M.: *Electrodynamics of continuous media, vol. 8, 2 edn.* elsevier (1984) (Cited on page 6.)
- [63] Mřřlek, J., Průruša, V.: Derivation of equations for continuum mechanics and thermodynamics of fluids. In: Y. Giga, A. Novotný (eds.) *Handbook of Mathematical Analysis in Mechanics of Viscous Fluids*, pp. 1–70. Springer (2017). DOI 10.1007/978-3-319-10151-4_1-1 (Cited on page 29.)

- [64] Marsden, J.E., Weinstein, A.: Coadjoint orbits, vortices and Clebsch variables for incompressible fluids. *Physica D* **7**, 305–323 (1983) (Cited on pages [8](#), [24](#), [27](#), and [31](#).)
- [65] Mazaheri, A., Ricchiuto, M., Nishikawa, H.: A first-order hyperbolic system approach for dispersion. *Journal of Computational Physics* **321**, 593–605 (2016). DOI 10.1016/j.jcp.2016.06.001. URL <http://www.sciencedirect.com/science/article/pii/S0021999116302261> (Cited on page [40](#).)
- [66] Mielke, A., Peletier, M.A., Renger, D.R.M.: On the Relation between Gradient Flows and the Large-Deviation Principle, with Applications to Markov Chains and Diffusion. *Potential Analysis* **41**(4), 1293–1327 (2014) (Cited on pages [8](#) and [36](#).)
- [67] Morrison, P.J.: Bracket formulation for irreversible classical fields. *Physics Letters A* **100**(8), 423–427 (1984) (Cited on page [3](#).)
- [68] Morrison, P.J.: Hamiltonian description of the ideal fluid. *Reviews of Modern Physics* **70**(2), 467–521 (1998). DOI 10.1103/RevModPhys.70.467. URL <http://link.aps.org/doi/10.1103/RevModPhys.70.467> (Cited on page [24](#).)
- [69] Muller, I., Ruggeri, T.: *Rational Extended Thermodynamics*, vol. 16. Springer (1998) (Cited on pages [4](#) and [5](#).)
- [70] Öttinger, H.C.: On the structural compatibility of a general formalism for nonequilibrium dynamics with special relativity. *Physica A: Statistical Mechanics and its Applications* **259**(1-2), 24–42 (1998). DOI 10.1016/S0378-4371(98)00298-2. URL <http://linkinghub.elsevier.com/retrieve/pii/S0378437198002982> (Cited on page [33](#).)
- [71] Öttinger, H.C.: *Beyond Equilibrium Thermodynamics*. Wiley (2005) (Cited on pages [3](#), [7](#), [11](#), [36](#), and [41](#).)
- [72] Öttinger, H.C., Grmela, M.: Dynamics and thermodynamics of complex fluids. II. Illustrations of a general formalism. *Physical Review E* **56**(6), 6633–6655 (1997). DOI 10.1103/PhysRevE.56.6633. URL <http://link.aps.org/doi/10.1103/PhysRevE.56.6633> (Cited on pages [3](#), [6](#), and [29](#).)
- [73] Pavelka, M., Klika, V., Esen, O., Grmela, M.: A hierarchy of Poisson brackets in non-equilibrium thermodynamics. *Physica D: Nonlinear Phenomena* **335**, 54–69 (2016). DOI 10.1016/j.physd.2016.06.011. URL <http://linkinghub.elsevier.com/retrieve/pii/S0167278915301019> (Cited on pages [8](#), [24](#), and [27](#).)
- [74] Pavelka, M., Klika, V., Grmela, M.: Time reversal in nonequilibrium thermodynamics. *Physical Review E* **90**(6), 1–19 (2014). DOI 10.1103/PhysRevE.90.062131. URL <http://link.aps.org/doi/10.1103/PhysRevE.90.062131> (Cited on pages [7](#) and [11](#).)
- [75] Pavelka, M., Klika, V., Grmela, M.: Time reversal in nonequilibrium thermodynamics. *Phys. Rev. E* **90**(062131) (2014) (Cited on pages [28](#), [35](#), [38](#), and [40](#).)
- [76] Peshkov, I., Grmela, M., Romenski, E.: Irreversible mechanics and thermodynamics of two-phase continua experiencing stress-induced solid-fluid transitions. *Continuum Mechanics and Thermodynamics* **27**(6), 905–940 (2015). DOI 10.1007/s00161-014-0386-1 (Cited on pages [9](#), [10](#), [15](#), [16](#), and [42](#).)
- [77] Peshkov, I., Romenski, E.: A hyperbolic model for viscous Newtonian flows. *Continuum Mechanics and Thermodynamics* **28**(1-2), 85–104 (2016). DOI 10.1007/s00161-014-0401-6 (Cited on pages [15](#), [26](#), [28](#), [36](#), [40](#), [41](#), [43](#), and [47](#).)
- [78] Peshkov, I., Romenski, E., Dumbser, M.: A unified hyperbolic formulation for viscous fluids and elastoplastic solids. *ArXiv e-prints (Accepted for Springer Proceedings in Mathematics and Statistics, XVI International Conference on Hyperbolic Problems)* (2017). URL <http://arxiv.org/abs/1705.02151> (Cited on pages [15](#), [26](#), [28](#), and [43](#).)

- [79] Powell, K.G., Roe, P.L., Linde, T.J., Gombosi, T.I., De Zeeuw, D.L., Keck, W.M.: A Solution-Adaptive Upwind Scheme for Ideal Magnetohydrodynamics. *Journal of Computational Physics* **154**, 284–309 (1999). URL <http://www.idealibrary.com> (Cited on pages 17, 18, and 56.)
- [80] Rajagopal, K.R., Srinivasa, A.R.: A thermodynamic frame work for rate type fluid models. *J. Non-Newton. Fluid Mech.* **88**(3), 207–227 (2000). DOI 10.1016/S0377-0257(99)00023-3 (Cited on page 29.)
- [81] Romenski, E., Belozerov, A., Peshkov, I.: Conservative formulation for compressible multiphase flows. *Quarterly of Applied Mathematics* **74**(1), 1–24 (2016). DOI 10.1090/qam/1409 (Cited on page 15.)
- [82] Romenski, E., Drikakis, D., Toro, E.: Conservative Models and Numerical Methods for Compressible Two-Phase Flow. *Journal of Scientific Computing* **42**(1), 68–95 (2010) (Cited on page 15.)
- [83] Romenski, E., Resnyansky, A.D., Toro, E.F.: Conservative hyperbolic model for compressible two-phase flow with different phase pressures and temperatures. *Quarterly of applied mathematics* **65**(2), 259–279 (2007) (Cited on page 15.)
- [84] Romenski, E.I., Sadykov, A.D.: On Modeling the Frequency Transformation Effect in Elastic Waves. *Journal of Applied and Industrial Mathematics* **5**(2), 282–289 (2011). DOI 10.1134/S1990478911020153 (Cited on pages 16 and 40.)
- [85] Romenskii, E.I.: Hyperbolic equations of Maxwell’s nonlinear model of elastoplastic heat-conducting media. *Siberian Mathematical Journal* **30**(4), 606–625 (1989) (Cited on pages 15, 37, 42, and 43.)
- [86] Romenskiy, E.I.: Hyperbolic systems of thermodynamically compatible conservation laws in continuum mechanics. *Mathematical and computer modelling* **28**(10), 115–130 (1998) (Cited on pages 15 and 19.)
- [87] Romenskiy, E.I.: Thermodynamics and hyperbolic systems of balance laws in continuum mechanics. In: *Godunov Methods: Theory and Applications*, pp. 745–761 (2001) (Cited on pages 6, 8, and 19.)
- [88] Ruggeri, T.: Global existence of smooth solutions and stability of the constant state for dissipative hyperbolic systems with applications to extended thermodynamics. In: S. Rionero, G. Romano (eds.) *Trends and Applications of Mathematics to Mechanics*, pp. 215–224. Springer, Milano (2005). DOI 10.1007/88-470-0354-7_17 (Cited on page 5.)
- [89] Ruggeri, T., Strumia, A.: Main field and convex covariant density for quasi-linear hyperbolic systems. *Relativistic Fluid Dynamics. Ann. Inst. Henri Poincaré* **34**(1), 65–84 (1981) (Cited on page 5.)
- [90] Struchtrup, H., Torrilhon, M.: Regularization of Grad’s 13 moment equations: Derivation and linear analysis. *Physics of Fluids* **15**(9), 2668–2680 (2003). DOI 10.1063/1.1597472 (Cited on page 4.)
- [91] Torrilhon, M.: Modeling Nonequilibrium Gas Flow Based on Moment Equations. *Annual Review of Fluid Mechanics* **48**(1), 429–458 (2016). DOI 10.1146/annurev-fluid-122414-034259. URL <http://dx.doi.org/10.1146/annurev-fluid-122414-034259> (Cited on page 4.)
- [92] Ván, P., Berezovski, A., Fülöp, T., Gróf, G., Kovács, R., Lovas, Á., Verhás, J.: Guyer-Krumhansl’s type heat conduction at room temperature. *EPL (Europhysics Letters)* **118**(5), 50,005 (2017). DOI 10.1209/0295-5075/118/50005. URL <http://stacks.iop.org/0295-5075/118/i=5/a=50005> (Cited on page 34.)