Variations of q-Garnier system

By

Hidehito Nagao¹ and Yasuhiko Yamada² $(Akashi College¹ and Kobe University², Japan)$ November 5, 2018

Abstract. We study several variants of q-Garnier system corresponding to various directions of discrete time evolutions. We also investigate a relation between the q-Garnier system and Suzuki's higher order q -Painlevé system by using a duality of the q -KP system.

Key Words and Phrases. q -Garnier system, q -Painlevé equation, Lax pair, q -KP system, Padé method.

2010 MSC Numbers. 14H70, 34M55, 37K20, 39A13, 41A21.

1 Introduction

The q-Garnier system is a multivariable extension of the q-Painlevé VI equation [\[8\]](#page-17-0). It was first formulated in [\[28\]](#page-18-0) and has been studied in [\[22,](#page-18-1) [25,](#page-18-2) [29\]](#page-18-3) from differing points of view.^{[1](#page-0-0)} For continuous/discrete Painlevé and Garnier systems in general, one can consider various directions of deformations (evolutions). In this paper, we consider such variations for q -Garnier system.^{[2](#page-0-1)} Our motivations and results are as follows.

- In general, we need to choose dependent variables appropriately according to their directions in order to make evolution equations in tractable form. We will derive simple equations for fundamental directions of q -Garnier systems by using the contiguity type scalar Lax pairs.
- In [\[22\]](#page-18-1) a connection between Suzuki's higher order q -Painlevé system [\[31,](#page-18-4) [32\]](#page-18-5) and q-Garnier system is suggested by comparing their special solutions. We clarify the connection by using a duality of the q -KP system [\[9,](#page-17-1) [10\]](#page-17-2).

This paper is organized as follows. In Section [2,](#page-1-0) based on our previous work [\[22\]](#page-18-1), we rewrite Sakai's 2×2 matrix Lax form into a scalar Lax form. In Section [3,](#page-3-0) we exemplify the scalar Lax pairs and the evolution equations for the various q -Garnier systems, by choosing the proper dependent variables. In Section [4,](#page-7-0) as an application of the results in Section [3,](#page-3-0) we give two reductions from case $N = 3$ of the various q-Garnier systems to q-Painlevé systems of type $E_7^{(1)}$ $q⁽¹⁾$. In Section [5,](#page-10-0) extending the duality for reduced q -KP system [\[9,](#page-17-1) [10\]](#page-17-2), we show the equivalence between Suzuki's system and a certain variation of q-Garnier system.

¹Other discrete analogues such as elliptic and additive Garnier systems have been studied in [\[23,](#page-18-6) [26,](#page-18-7) [40\]](#page-19-0) and [\[2,](#page-17-3) [3\]](#page-17-4).

²For differential Garnier system, the "continuous time evolutions" and the "Schlesinger transformations" are usually considered as different kind of equations. However, in difference case, we call both of them "Garnier system", since we can treat them in similar manner.

2 q-Garnier system

In this section we recall Sakai's 2×2 matrix Lax from [\[28\]](#page-18-0) for the q-Garnier system and its corresponding scalar Lax form [\[22\]](#page-18-1).

2.1 Sakai's matrix Lax form

In [\[28\]](#page-18-0), the q-Garnier system has been formulated by H.Sakai. For an unknown function $Y(z) = {}^t[y_1(z), y_2(z)]$, the 2×2 matrix Lax form for the q-Garnier system is given by

$$
Y(qz) = \mathcal{A}(z)Y(z),\tag{2.1}
$$

$$
\overline{Y}(z) = \mathcal{B}(z)Y(z). \tag{2.2}
$$

Here q is a (complex) parameter, and $\mathcal A$ and $\mathcal B$ are 2×2 matrices depending on z and other parameters. The matrix $\mathcal A$ is a polynomial in z (see properties [\(2.4\)](#page-1-1) below), on the other hand the matrix B is rational in z. We denote the time evolution by $T(*) = \overline{*}$. The equation [\(2.1\)](#page-1-2) gives the linear q-difference equation and the equation [\(2.2\)](#page-1-3) gives its deformation equation^{[3](#page-1-4)}. The compatibility condition for the matrix systems (2.1) and (2.2) reads

$$
\overline{\mathcal{A}}(z)\mathcal{B}(z) = \mathcal{B}(qz)\mathcal{A}(z),\tag{2.3}
$$

which gives the q-Garnier system. Let $\alpha_1, \ldots \alpha_{2N+2}, \kappa_1, \kappa_2, \theta_1$ and θ_2 be (complex) parameters. Then the matrix $\mathcal A$ is characterized by the following properties :

(i)
$$
\mathcal{A}(z) = A_0 + A_1 z + \dots + A_{N+1} z^{N+1}
$$
,
\n(ii) $A_{N+1} = \begin{bmatrix} \kappa_1 & 0 \\ 0 & \kappa_2 \end{bmatrix}$, A_0 has eigenvalues θ_1 and θ_2 ,
\n(iii) $|\mathcal{A}(z)| = \kappa_1 \kappa_2 \prod_{i=1}^{2N+2} (z - \alpha_i)$, $(\kappa_1 \kappa_2 \prod_{i=1}^{2N+2} \alpha_i = \theta_1 \theta_2)$. (2.4)

The matrix A satisfying these properties (i)–(iii) has $2N + 1$ arbitrary parameters except parameters $\{\alpha_i, \kappa_i, \theta_i\}$. The 2N parameters of them can be interpreted as variables for the q-Garnier system and the remaining parameter is a "gauge" freedom. The deformation direction in [\[28\]](#page-18-0) is given by $\overline{\alpha}_i = q\alpha_i \ \overline{\alpha}_j = \alpha_j$, $\overline{\kappa}_i = \kappa_i$ and $\theta_i = q\theta_i$ $(i = 1, 2, j = 3, \ldots, 2N + 2).$

We must choose proper coordinates of variables according to the deformation directions, in order to obtain simple forms of the q-Garnier systems. This content will be shown in Section [3.](#page-3-0)

2.2 Scalar Lax form

We recall the scalar Lax form and derive the simple form of the time evolution equation from its compatibility condition.

³The matrix β can be chosen according to various deformation directions.

Lemma 2.2.1 The matrix system (2.1) and (2.2) can be rewritten into the following linear q-difference equations:

$$
L_1: F(\frac{z}{q})y(qz) - \left\{ F(\frac{z}{q})\mathcal{A}_{11}(z) + F(z)\mathcal{A}_{22}(\frac{z}{q}) \right\} y(z) + F(z)|\mathcal{A}(\frac{z}{q})|y(\frac{z}{q}) = 0,
$$

\n
$$
L_2: F(z)\overline{y}(z) - H(z)y(qz) + G(z)y(z) = 0,
$$

\n
$$
L_3: \overline{F}(z)|\mathcal{B}(z)|y(qz) - H(qz)|\mathcal{A}(z)|\overline{y}(z) + G(z)\overline{y}(qz) = 0.
$$
\n(2.5)

Here

$$
y(z) = y_1(z), \ F(z) = \mathcal{A}_{12}(z), \ G(z) = \mathcal{A}_{11}(z)\mathcal{B}_{12}(z) - \mathcal{A}_{12}(z)\mathcal{B}_{11}(z), \ H(z) = \mathcal{B}_{12}(z). \tag{2.6}
$$

Proof. We write the matrix system (2.1) as

$$
\begin{cases}\ny_1(qz) = \mathcal{A}_{11}(z)y_1(z) + \mathcal{A}_{12}(z)y_2(z), \\
y_2(qz) = \mathcal{A}_{21}(z)y_1(z) + \mathcal{A}_{22}(z)y_2(z).\n\end{cases}
$$
\n(2.7)

Eliminating the component $y_2(z)$ from the equation [\(2.7\)](#page-2-0), we obtain the linear equation L_1 [\(2.5\)](#page-2-1). The matrix system [\(2.2\)](#page-1-3) is also rewritten as

$$
\begin{cases} \overline{y}_1(z) = \mathcal{B}_{11}(z)y_1(z) + \mathcal{B}_{12}(z)y_2(z), \\ \overline{y}_2(z) = \mathcal{B}_{21}(z)y_1(z) + \mathcal{B}_{22}(z)y_2(z). \end{cases}
$$
\n(2.8)

Eliminating the component $y_2(z)$ from the first equations in [\(2.7\)](#page-2-0) and [\(2.8\)](#page-2-2), we obtain the deformation equation L_2 [\(2.5\)](#page-2-1).

On the other hand, rewriting [\(2.1\)](#page-1-2) and [\(2.2\)](#page-1-3) into

$$
\overline{Y}(qz) = \overline{\mathcal{A}}(z)\overline{Y}(z), \quad \overline{Y}(qz) = \mathcal{B}(qz)Y(qz), \tag{2.9}
$$

and eliminating $y_2(qz)$, $\overline{y}_2(z)$ and $\overline{y}_2(qz)$ from the system [\(2.9\)](#page-2-3), then we have

$$
\overline{\mathcal{A}}_{12}(z)|\mathcal{B}(qz)|y(qz)-H(qz)|\overline{\mathcal{A}}(z)|\overline{y}(x)-\left\{\mathcal{A}_{12}(z)\mathcal{B}_{22}(qz)-\overline{\mathcal{A}}_{22}(z)\mathcal{B}_{12}(z)\right\}\overline{y}(qz)=0.
$$
\n(2.10)

Moreover applying the compatibility condition (2.3) to (2.10) , we we can derive the deformation equation L_3 [\(2.5\)](#page-2-1).

We remark that unknown variables of q -Garnier systems appear as the coefficients in z of the polynomials F, G and H . The polynomial F is of degree N in z and the N points such that $F(z) = 0$ are not singularities but "*apparent singularities*" (i.e. the solutions are regular there). This means that the two relations

$$
\frac{y(qz)}{y(z)} = \mathcal{A}_{11}(z), \quad \text{for} \quad F(z) = 0,
$$
\n
$$
\frac{y(z)}{y(\frac{z}{q})} = \frac{|\mathcal{A}(\frac{z}{q})|}{\mathcal{A}_{22}(\frac{z}{q})}, \quad \text{for} \quad F(\frac{z}{q}) = 0,
$$
\n
$$
(2.11)
$$

derived from the L_1 equation [\(2.5\)](#page-2-1) are consistent with each other.

A pair of the deformation equations $\{L_2, L_3\}$ is equivalent to the scalar Lax pair ${L_1, L_2}$, so we call the pair ${L_2, L_3}$ the "scalar Lax pair (of contiguous relation type)". We conveniently deal with the Lax pair $\{L_2, L_3\}$ [\[24\]](#page-18-8), since the explicit form of the L_1 equation is rather complicated.

Lemma 2.2.2 The compatibility of the scalar Lax pair $\{L_2, L_3\}$ [\(2.5\)](#page-2-1) gives the following relation:

$$
G(z)\underline{G}(z) = H(z)\underline{H}(qz)|\underline{A}(z)|, \quad \text{for} \quad F(z) = 0,
$$
\n
$$
|\mathcal{B}(z)|F(z)\overline{F}(z) = H(z)H(qz)|\underline{A}(z)|, \quad \text{for} \quad G(z) = 0.
$$
\n
$$
\Box
$$

Proof. Under the condition $F(z) = 0$, eliminating $y(z)$ and $y(qz)$ from $L_2(z) = L_3(qz) =$ 0, we obtain the first relation of [\(2.12\)](#page-3-1). Similarly, for $G(z) = 0$, eliminating $y(qx)$ and $\overline{y}(x)$ from $L_2(z) = L_3(qz) = 0$, we have the second relation of [\(2.12\)](#page-3-1).

The relation [\(2.12\)](#page-3-1) may be insufficient for the compatibility of the Lax pair $\{L_2, L_3\}$. Additional conditions for the sufficiency will be considered case by case in Section [3.](#page-3-0)

Though we have derived the scalar Lax pair from matrix Lax form, the scalar Lax pair $\{L_2, L_3\}$ can be also obtained from a certain method using Padé interpolation (see for example [\[22\]](#page-18-1)). In the following sections, we will discuss based on the Lax pair obtained by Padé method^{[4](#page-3-2)}.

3 Variations of the q-Garnier system

In this section, we consider Lax pairs and evolution equations for several directions of q-Garnier systems.

3.1 Notation

Fix a positive integer N and a parameter q. Let $a_1, \ldots, a_{N+1}, b_1, \ldots, b_{N+1}, c_1, c_2, d_1, d_2$ be parameters with a constraint $\prod_{i=1}^{N+1} \frac{a_i}{b_i}$ $\frac{a_i}{b_i} = q \prod_{i=1}^2 \frac{c_i}{d_i}$ $\frac{c_i}{d_i}$ and $T_a: a \mapsto qa$ be the q-shift operator of parameter a.

We consider the following four directions T_1, \ldots, T_4 defined by

$$
T_1 = T_{a_1}^{-1} T_{b_1}^{-1}, \quad T_2 = T_{a_1}^{-1} T_{a_2}^{-1} T_{b_1}^{-1} T_{b_2}^{-1}, \quad T_3 = T_{c_1} T_{d_1}, \quad T_4 = T_{a_{N+1}}^{-1} T_{c_1}^{-1}, \tag{3.1}
$$

and the corresponding shifts are denoted as $\overline{X} := T_i(X)$ and $\underline{X} := T_i^{-1}$ $i^{-1}(X)$. The operators T_i play the role of time evolutions of the q -Garnier system.

The directions T_1 , T_3 and T_4 (together with related ones obtained by obvious symmetry: $T_{a_i}^{-1}T_{b_j}^{-1}$ $L_{b_j}^{-1}$, $T_{c_i}T_{d_j}$, $T_{a_i}^{-1}T_{c_j}^{-1}$ are fundamental ones, and others (e.g. T_2) are given by compositions of them (and their inverses).

⁴For the proof of the correspondence between (2.5) and (3.11) independent of Padé method, see Appendix [A.1](#page-13-0) as an example

In this subsection, we use the following notations:

$$
A(z) = \prod_{i=1}^{N+1} (z - a_i), \quad B(z) = \prod_{i=1}^{N+1} (z - b_i), \quad A_i(z) = \frac{A(z)}{z - a_i}, \quad B_i(z) = \frac{B(z)}{z - b_i},
$$

$$
A_{i,j}(z) = \frac{A(z)}{(z - a_i)(z - a_j)}, \quad B_{i,j}(z) = \frac{B(z)}{(z - b_i)(z - b_j)}, \quad F(z) = \sum_{i=0}^{N} f_i z^i,
$$
 (3.2)

where f_0, \ldots, f_N are variables depending on parameters a_i, b_i, c_i and d_i .

The following data will be described by the polynomials $F(z)$, $G(z)$ and $H(z)$. The polynomial F is common for all cases as given above, while one should take the polynomials G and H differently case by case as given below.

3.2 Results for each direction

In the following subsubsections, we show the three items^{[5](#page-4-0)}: (a) Scalar Lax pair, (b) Time evolution equation, (c) The L_1 equation.

In item (b), we give the time evolution equation as necessary and sufficient condition for compatibility of the scalar Lax pair (For the proof of the sufficiency, see Appendix [A.2\)](#page-14-0). In item (c), the L_1 equations in each subsections are expressed in different forms, however, we will show that they are equivalent with each other (Theorem [3.2.1\)](#page-7-1).

3.2.1 Direction $T_1 = T_{a_1}^{-1} T_{b_1}^{-1}$ b_1

This case is considered in [\[22\]](#page-18-1) and the direction T_1 is (an inverse of) the original direction in [\[28\]](#page-18-0).

(a) Scalar Lax pair

$$
L_2(z) = F(z)\overline{y}(z) - A_1(z)y(qz) + (z - b_1)G(z)y(z),
$$

\n
$$
L_3(z) = \overline{F}(\frac{z}{q})y(z) + (z - a_1)G(\frac{z}{q})\overline{y}(z) - qc_1c_2B_1(\frac{z}{q})\overline{y}(\frac{z}{q}),
$$
\n(3.3)

where $A_1(z)$, $B_1(z)$, $F(z)$ are as in [\(3.2\)](#page-4-1) and $G(z) = \sum_{i=0}^{N-1} g_i z^i$. (b) Time evolution equation

$$
G(z)\underline{G}(z) = c_1 c_2 \frac{A_1(z)B_1(z)}{(z - a_1)(z - b_1)}, \quad \text{for} \quad F(z) = 0,
$$

\n
$$
F(z)\overline{F}(z) = qc_1 c_2 A_1(z)B_1(z), \quad \text{for} \quad G(z) = 0,
$$

\n
$$
f_N \overline{f}_N = q(g_{N-1} - c_1)(g_{N-1} - c_2),
$$

\n
$$
f_0 \overline{f}_0 = a_1 b_1 \Big(g_0 - \frac{d_1}{a_1 b_1} A(0)\Big) \Big(g_0 - \frac{d_2}{a_1 b_1} A(0)\Big),
$$
\n(3.4)

where $2N$ variables $\frac{f_1}{f_0}, \ldots, \frac{f_N}{f_0}$ $\frac{f_N}{f_0}, g_0, \ldots, g_{N-1}$ are the dependent variables.

⁵The data for each items is obtained by considering a same Padé problem with various deformations (see Appendix [B\)](#page-15-0).

(c) The L_1 equation

$$
L_1(z) = A(z)F(\frac{z}{q})y(qz) + qc_1c_2B(\frac{z}{q})F(z)y(\frac{z}{q})
$$

$$
- \left\{ (z-a_1)(z-b_1)F(\frac{z}{q})G(z) + \frac{F(z)}{G(\frac{z}{q})}V(\frac{z}{q}) \right\}y(z),
$$
 (3.5)

where $V(z) = qc_1c_2A_1(z)B_1(z) - F(z)\overline{F}(z)$.

3.2.2 Direction $T_2 = T_{a_1}^{-1} T_{a_2}^{-1} T_{b_1}^{-1}$ $T_{b_1}^{-1}T_{b_2}^{-1}$ $b₂$

(a) Scalar Lax pair

$$
L_2(z) = F(z)\overline{y}(z) - A_{1,2}(z)(1 + hz)y(qz) + (z - b_1)(z - b_2)G(z)y(z),
$$

\n
$$
L_3(z) = \overline{F}(\frac{z}{q})y(z) + (z - a_1)(z - a_2)G(\frac{z}{q})\overline{y}(z) - qc_1c_2B_{1,2}(\frac{z}{q})(1 + hz)\overline{y}(\frac{z}{q}),
$$
\n(3.6)

where $A_{1,2}(z)$, $B_{1,2}(z)$, $F(z)$ are as in [\(3.2\)](#page-4-1) and $G(z) = \sum_{i=0}^{N-2} g_i z^i$. (b) Time evolution equation

$$
\frac{G(z)\underline{G}(z)}{(1+hz)(1+q\underline{h}z)} = \frac{c_1c_2A_{1,2}(z)B_{1,2}(z)}{q(z-a_1)(z-a_2)(z-b_1)(z-b_2)}, \text{ for } F(z) = 0,
$$

\n
$$
F(z)\overline{F}(z) = qc_1c_2A_{1,2}(z)B_{1,2}(z)(1+hz)(1+qhz), \text{ for } G(z) = 0,
$$

\n
$$
F(z)\overline{F}(\frac{z}{q}) = (z-a_1)(z-a_2)(z-b_1)(z-b_2)G(z)G(\frac{z}{q}), \text{ for } 1+hz = 0,
$$

\n
$$
f_N\overline{f}_N = q^2(g_{N-2} - c_1h)(g_{N-2} - c_2h),
$$

\n
$$
f_0\overline{f}_0 = a_1a_2b_1b_1(g_0 - \frac{d_1A(0)}{a_1a_2b_1b_2})(g_0 - \frac{d_2A(0)}{a_1a_2b_1b_2}),
$$
\n(3.7)

where $2N$ variables $\frac{f_1}{f_0}, \ldots, \frac{f_N}{f_0}$ $\frac{f_N}{f_0}, g_0, \ldots, g_{N-2}$ and h are the dependent variables. (c) The L_1 equation

$$
L_1(z) = A(z)F(\frac{z}{q})y(qz) + qc_1c_2B(\frac{z}{q})F(z)y(\frac{z}{q})
$$

$$
-\frac{1}{1+hz}\Big\{(z-a_1)(z-b_1)(z-a_2)(z-b_2)F(\frac{z}{q})G(z) + \frac{F(z)}{G(\frac{z}{q})}V(\frac{z}{q})\Big\}y(z),
$$
 (3.8)

where $V(z) = qc_1c_2(1 + hz)(1 + qhz)A_{1,2}(z)B_{1,2}(z) - F(z)\overline{F}(z).$

3.2.3 Direction $T_3 = T_{c_1} T_{d_1}$

The deformation direction T_3 is the same as that of Suzuki's system [\[31\]](#page-18-4) (see §[5.3\)](#page-11-0). (a) Scalar Lax pair

$$
L_2(z) = F(z)\overline{y}(z) - A(z)y(qz) + G(z)y(z),
$$

\n
$$
L_3(z) = \frac{z}{q}\overline{F}(\frac{z}{q})y(z) + G(\frac{z}{q})\overline{y}(x) - qc_1c_2B(\frac{z}{q})\overline{y}(\frac{z}{q}),
$$
\n(3.9)

where $A(z)$, $B(z)$, $F(z)$ are as in [\(3.2\)](#page-4-1) and $G(z) = \sum_{i=0}^{N+1} g_i z^i$, $g_{N+1} = \frac{c_2 g_0}{d_1 A(0)}$. (b) Time evolution equation

$$
G(z)\underline{G}(z) = c_1c_2A(z)B(z), \quad \text{for} \quad F(z) = 0,
$$

\n
$$
zF(z)\overline{F}(z) = qc_1c_2A(z)B(z), \quad \text{for} \quad G(z) = 0,
$$
\n(3.10)

where $2N$ variables $\frac{f_1}{f_0}, \ldots, \frac{f_N}{f_0}$ $\frac{f_N}{f_0}, \frac{g_1}{g_0}$ $\frac{g_1}{g_0}, \ldots, \frac{g_N}{g_0}$ $\frac{g_N}{g_0}$ are the dependent variables. (c) The L_1 equation

$$
L_1(z) = A(z)F(\frac{z}{q})y(qz) + qc_1c_2B(\frac{z}{q})F(z)y(\frac{z}{q})
$$

$$
- \left\{ F(\frac{z}{q})G(z) + \frac{F(z)}{G(\frac{z}{q})}V(\frac{z}{q}) \right\}y(z),
$$
 (3.11)

where $V(z) = qc_1c_2A(z)B(z) - zF(z)\overline{F}(z)$.

3.2.4 Direction $T_4 = T_{a_{N+1}}^{-1} T_{c_1}^{-1}$

(a) Scalar Lax pair

$$
L_2(z) = F(z)\overline{y}(z) - A_{N+1}(z)y(qz) + G(z)y(z),
$$

\n
$$
L_3(z) = \overline{F}(\frac{z}{q})y(z) + (z - a_{N+1})G(\frac{z}{q})\overline{y}(x) - qc_1c_2B(\frac{z}{q})\overline{y}(\frac{z}{q}),
$$
\n(3.12)

where $A_{N+1}(z)$, $B(z)$, $F(z)$ are as in [\(3.2\)](#page-4-1) and $G(z) = \sum_{i=0}^{N} g_i z^i$, $g_N = c_1$. (b) Time evolution equation

$$
G(z)\underline{G}(z) = \frac{qc_1c_2A_{N+1}(z)B(z)}{z - a_{N+1}}, \quad \text{for} \quad F(z) = 0,
$$

\n
$$
F(z)\overline{F}(z) = qc_1c_2A_{N+1}(z)B(z), \quad \text{for} \quad G(z) = 0,
$$

\n
$$
f_0\overline{f}_0 = \left(g_0 + \frac{d_1A(0)}{a_{N+1}}\right)\left(g_0 + \frac{d_2A(0)}{a_{N+1}}\right),
$$
\n(3.13)

where $2N$ variables $\frac{f_1}{f_0}, \ldots, \frac{f_N}{f_0}$ $\frac{f_N}{f_0}, g_0, \ldots, g_{N-1}$ are the dependent variables. (c) The L_1 equation

$$
L_1(z) = A(z)F(\frac{z}{q})y(qz) + qc_1c_2B(\frac{z}{q})F(z)y(\frac{z}{q})
$$

$$
-\left\{(z-a_{N+1})F(\frac{z}{q})G(z) + \frac{F(z)}{G(\frac{z}{q})}V(\frac{z}{q})\right\}y(z),
$$
 (3.14)

where $V(z) = qc_1c_2A_{N+1}(z)B(z) - F(z)\overline{F}(z)$.

3.2.5 Relations among variables g_i and h for each direction

Theorem 3.2.1 The L_1 equations for each direction T_1, \ldots, T_4 : [\(3.5\)](#page-5-0), [\(3.8\)](#page-5-1), [\(3.11\)](#page-6-0) and (3.14) are equivalent with each other if the coefficients g_i in $G(z)$ and h are related as

$$
\frac{(z - b_1)G(z)^{\S^{3.2.1}}}{A_1(z)} = \frac{(z - b_1)(z - b_2)}{A_{1,2}(z)} \left(\frac{G(z)}{1 + hz}\right)^{\S^{3.2.2}}
$$
\n
$$
= \frac{G(z)^{\S^{3.2.3}}}{A(z)} = \frac{G(z)^{\S^{3.2.4}}}{A_{N+1}(z)}, \quad \text{for } F(z) = 0.
$$
\n
$$
\Box
$$

Proof. The relation [\(3.15\)](#page-7-2) is obtained by comparing ratios $\frac{y(qz)}{y(z)}$ in L_1 under the condition $F(z) = 0$. Then the equivalence of the L_1 equations can be checked using their characteristic properties by a similar argument as the Appendix [A.1.](#page-13-0)

4 Reduction to q - $E_7^{(1)}$ $\frac{1}{7}^{(1)}$ system

In this section we consider some q-Painlevé equations of type $E_7^{(1)}$ $q_7^{(1)}$ as a reduction of q-Garnier systems.

4.1 Notation

Throughout this section, we consider the case $N = 3$ and specialize parameters as

$$
c_1 = c_2, \quad d_1 = d_2. \tag{4.1}
$$

Among the directions T_i considered in previous section, the directions consistent with this specialization are the following three:

$$
T_1 = T_{a_1}^{-1} T_{b_1}^{-1}, \quad T_2 = T_{a_1}^{-1} T_{a_2}^{-1} T_{b_1}^{-1} T_{b_2}^{-1}, \tag{4.2}
$$

and inconsistent ones T_3 and T_4 will be omitted. Under the specialization [\(4.1\)](#page-7-3), we can and will impose constraints on dependent variables as

$$
f_0 = f_3 = 0, \quad f_1 = w, \quad f_2 = -fw,
$$
\n(4.3)

where f is one of the unknown variables for the reduced system and w is a gauge freedom. In the followings, we also impose additional constraints [\(4.4\)](#page-8-0) or [\(4.10\)](#page-9-0) in order to reduce the variables g_i and h of q-Garnier systems to a variable q and a parameter e. Note that the meaning of the parameter e and the reduced variable g are different depending on directions. The results of §[4.2.1](#page-8-1) are known in [\[22,](#page-18-1) [20\]](#page-18-9) and those of §[4.2.2](#page-9-1) are new.

4.2 Results for each direction

4.2.1 Reduction from q-Garnier system in §[3.2.1](#page-4-2)

The direction is $T_1 = T_{a_1}^{-1} T_{b_1}^{-1}$ b_1^{-1} . In this case we can and will impose an additional condition:

$$
g_0 = e, \quad g_1 = eg, \quad g_2 = c_1,
$$
 (4.4)

where $e = d_1 a_2 a_3 a_4 b_1^{-1}$. (a) Scalar Lax pair

Under the conditions (4.1) , (4.3) and (4.4) , we can reduce the Lax pair (3.3) to the following Lax pair:

$$
L_2(z) = wz(1 - fz)\overline{y}(z) - \prod_{i=2}^4 (z - a_i)y(qz) + (z - b_1)(e + egz + c_1z^2)y(z),
$$

\n
$$
L_3(z) = \overline{w}_q^2(1 - \overline{f}_q^2)y(z) + (z - a_1)\{e + eg_q^2 + c_1(\frac{z}{q})^2\}\overline{y}(z) - qc_1^2\prod_{i=2}^4(\frac{z}{q} - b_i)\overline{y}(\frac{z}{q}).
$$
\n(4.5)

The Lax pair [\(4.5\)](#page-8-2) has been given in [\[20\]](#page-18-9).

(b) Time evolution equation and equation for w

$$
\left(\frac{e}{c_1}f^2 + \frac{e}{c_1}gf + 1\right)\left(\frac{e}{qc_1}f^2 + \frac{e}{qc_1}gf + 1\right) = \frac{\prod_{i=2}^4 (1 - a_i f)(1 - b_i f)}{(1 - a_1 f)(1 - b_1 f)},
$$
\n
$$
\frac{z_1^2 (1 - fz_1)(1 - \overline{f}z_1)}{z_2^2 (1 - fz_2)(1 - \overline{f}z_2)} = \prod_{i=2}^4 \frac{(z_1 - a_i)(z_1 - b_i)}{(z_2 - a_i)(z_2 - b_i)},
$$
\n(4.6)

and

$$
w\overline{w} = \frac{\prod_{j=2}^{4} (z_i - a_j)(z_i - b_j)}{z_i^2 (1 - fz_i)(1 - \overline{f}z_i)}, \quad (i = 1, 2),
$$
\n(4.7)

where $z = z_1, z_2$ are solutions of the equation $e + egz + c_1z^2 = 0$. The bi-rational equation [\(4.6\)](#page-8-3) is equivalent to the variation of q-Painlevé equation of type $E_7^{(1)}$ $\frac{1}{7}$ ⁽¹⁾ in [\[20,](#page-18-9) [22\]](#page-18-1).

We remark that the configuration of $8 \text{ singular points}^6$ $8 \text{ singular points}^6$ is given by 2 points on a line $g = \infty$ and 6 points on a parabolic curve $ef^2 + egf + c_1 = 0$ as follows:

$$
(f,g) = (\frac{1}{a_1}, \infty), (\frac{1}{b_1}, \infty), (\frac{1}{a_i}, -\frac{1}{a_i} - \frac{a_i c_1}{e}), (\frac{1}{b_i}, -\frac{1}{b_i} - \frac{b_i c_1}{e}), \quad (i = 2, 3, 4). \tag{4.8}
$$

(c) The L_1 equation

$$
L_1(z) = \prod_{i=1}^4 (z - a_i)(1 - \frac{fz}{q})y(qz) + q^2 c_1^2 \prod_{i=1}^4 (\frac{z}{q} - b_i)(1 - fz)y(\frac{z}{q})
$$

$$
- \left\{ (z - a_1)(z - b_1)(1 - \frac{fz}{q})\varphi(z) + \frac{q(1 - fz)}{\varphi(\frac{z}{q})}V(\frac{z}{q}) \right\}y(z),
$$
 (4.9)

where $V(z) = qc_1^2 \prod_{i=2}^4 (z - a_i)(z - b_i) - w\overline{w}z^2(1 - fz)(1 - \overline{f}z)$ and $\varphi(z) = e + egz + c_1z^2$. The L_1 equation [\(4.9\)](#page-8-5) has been given in [\[20\]](#page-18-9).

 6 For the theory of the configuration of 8 singular points, see [\[11,](#page-17-5) [27\]](#page-18-10).

4.2.2 Reduction from q-Garnier system in §[3.2.2](#page-5-2)

The direction is $T_2 = T_{a_1}^{-1} T_{a_2}^{-1} T_{b_1}^{-1}$ $T_{b_1}^{-1}T_{b_1}^{-1}$ b_1^{-1} . We impose an additional condition:

$$
g_0 = e, \quad g_1 = -\frac{e}{g}, \quad h = -\frac{e}{c_1 g},
$$
 (4.10)

where $e = d_1 a_3 a_4 (b_1 b_2)^{-1}$.

(a) Scalar Lax pair

Under the conditions (4.1) , (4.3) and (4.10) , one can reduce the Lax pair (3.6) to the following Lax pair:

$$
L_2(z) = wz(1 - fz)\overline{y}(z) - \prod_{i=3}^4 (z - a_i)(1 - \frac{ez}{c_1g})y(qz) + e \prod_{i=1}^2 (z - b_i)(1 - \frac{z}{g})y(z),
$$

\n
$$
L_3(z) = \overline{w}_q^{\frac{z}{2}}(1 - \overline{f}_q^{\frac{z}{2}})y(z) + e \prod_{i=1}^2 (z - a_i)(1 - \frac{z}{qg})\overline{y}(z) - qc_1^2 \prod_{i=3}^4 (\frac{z}{q} - b_i)(1 - \frac{ez}{c_1g})\overline{y}(\frac{z}{q}).
$$
\n(4.11)

The scalar Lax pair [\(4.11\)](#page-9-2) is equivalent to the one in [\[11,](#page-17-5) [38\]](#page-19-1).

(b) Time evolution equation and equation for w

$$
\frac{\left(\frac{c_1}{e}fg - 1\right)\left(\frac{qc_1}{e}fg - 1\right)}{(fg - 1)(fg - 1)} = \frac{\prod_{i=1}^{2} (1 - a_i f)(1 - b_i f)}{\prod_{i=3}^{4} (1 - a_i f)(1 - b_i f)},
$$
\n
$$
\frac{\left(fg - \frac{e}{c_1}\right)\left(\overline{fg} - \frac{qe}{c_1}\right)}{\left(fg - 1\right)\left(\overline{fg} - 1\right)} = \frac{\prod_{i=1}^{2} (g - a_i \frac{e}{c_1})(g - b_i \frac{e}{c_1})}{\prod_{i=3}^{4} (g - a_i)(g - b_i)},
$$
\n(4.12)

and

$$
w\overline{w} = \frac{\prod_{i=3}^{4} (g - a_i)(g - b_i)}{qg^2(c_1 - e)(c_1 - qe)(fg - 1)(\overline{f}g - 1)}.
$$
\n(4.13)

The bi-rational equation [\(4.12\)](#page-9-3) is equivalent to the *q*-Painlevé equation of type $E_7^{(1)}$ $\frac{1}{7}^{\left(1\right)}$ in [\[5,](#page-17-6) [11,](#page-17-5) [27,](#page-18-10) [38\]](#page-19-1), and the configuration of 8 singular points is the well-known standard one. (c) The L_1 equation

$$
L_1(z) = \prod_{i=1}^4 (z - a_i)(1 - \frac{fz}{q})y(qz) + q^2 c_1^2 \prod_{i=1}^4 (\frac{z}{q} - b_i)(1 - fz)y(\frac{z}{q})
$$

$$
- \frac{1}{1 + \frac{ez}{c_1 g}} \Big\{ e \prod_{i=1}^2 (z - a_i)(z - b_i)(1 - \frac{fz}{q})(1 - \frac{z}{g}) + \frac{q(1 - fz)}{e(1 - \frac{z}{qg})}V(\frac{z}{q}) \Big\}y(z),
$$
\n(4.14)

where $V(z) = qc_1^2(1 - \frac{ez}{c_1}$ $\frac{ez}{c_1g}$) $\left(1 - \frac{qez}{c_1g}\right)$ $\frac{qez}{q_1g}$) $\prod_{i=3}^4(z-a_i)(z-b_i) - w\overline{w}z^2(1-fz)(1-\overline{f}z).$

Remark 4.2.1 The L_1 equations for each directions T_1 and T_2 are equivalent under the relation [\[20\]](#page-18-9)

$$
\frac{(1-a_2f)(1-b_2f)}{\left(\frac{e}{c_1}f^2 + \frac{e}{c_1}gf + 1\right)^{\S 4.2.1}} \left(\frac{1-fg}{1 - \frac{c_1}{e}fg}\right)^{\S 4.2.2} = 1.
$$
\n(4.15)

This result follows as a reduction of Theorem [3.2.1.](#page-7-1) A direct proof is given in [\[20\]](#page-18-9). \Box

Remark 4.2.2 If we include more directions such as $T_{a_1}^{-1}T_{a_2}^{-1}T_{c_1}^{-1}$ and $T_{a_1}^{-1}T_{a_2}^{-1}T_{d_1}$ in addition to T_1 as fundamental ones, then all the other directions can be obtained by compositions of them. Thanks to the symmetry of the parameters, the equations for such additional fundamental directions are similar to that of T_1 . Therefore the results of this section are enough to obtain all the directions of q-Painlevé equation of type $E_7^{(1)}$ 7 \Box

5 Relation to Suzuki's system

5.1 q -KP and its duality

We first recall a Lax formalism of a certain isomonodromic system formulated as a reduction of the q -KP hierarchy $[10]^7$ $[10]^7$ $[10]^7$.

Let $x_{i,j}, r_i, t_j \in \mathbb{C}$ and $m, n \in \mathbb{N}$ be parameters. For an unknown *n*-vector $Y(z)$, we consider a linear q-difference equation of $n \times n$ matrix system

$$
Y(qz) = \mathcal{A}(z)Y(z),\tag{5.1}
$$

and its deformation system

$$
\overline{Y}(z) = \mathcal{B}(z)Y(z). \tag{5.2}
$$

Here^{[8](#page-10-2)}

$$
\mathcal{A}(z) = dX_m \cdots X_2 X_1,
$$
\n
$$
X_i(z) = \begin{bmatrix} x_{i,1} & 1 & & & \\ & x_{i,2} & 1 & & \\ & & \ddots & \ddots & \\ & & & x_{i,n-1} & 1 \\ & & & & x_{i,n} \end{bmatrix}, \quad d = \text{diag}[t_1, \dots, t_n].
$$
\n(5.3)

In this section, we focus consideration on the linear equation [\(5.1\)](#page-10-3).

Extending the result of [\[9,](#page-17-1) [10\]](#page-17-2), one obtains the following:

Theorem 5.1.1 Two cases (m, n) and (n, m) of the linear equation [\(5.1\)](#page-10-3) are equivalent under the following transformation:

$$
x_{i,j} \leftrightarrow -x_{j,i}, \quad r_k \leftrightarrow t_k, \quad z \leftrightarrow T_z. \tag{5.4}
$$

 \Box

Proof. Setting $Y_1 = Y, Y_{i+1} = X_i Y_i$ ($1 \leq i \leq m$) and defining components of Y by $Y_{i,j} = (Y_i)_j$, then we have relations

$$
Y_{i+1,j} = x_{i,j} Y_{i,j} + Y_{i,j+1}, \quad Y_{m+1,j} = t_j^{-1} T_z Y_{1,j}, \quad Y_{i,n+1} = r_i^{-1} z Y_{i,1}.
$$
 (5.5)

⁷For the related works on q -KP, see [\[9\]](#page-17-1) (affine Weyl group symmetry), [\[15,](#page-18-11) [33\]](#page-19-2) (the case (m, n) = $(2, 3), (2, 4), [12, 36]$ $(2, 3), (2, 4), [12, 36]$ $(2, 3), (2, 4), [12, 36]$ (Yang-Baxter maps), [\[34\]](#page-19-4) $(q$ -UC hierarchy), [\[1,](#page-17-8) [6\]](#page-17-9) (non-commutative version).

⁸For convenience, extra parameters r_i and t_j are introduced though there are redundancy.

Applying a transformation [\(5.4\)](#page-10-4) and

$$
m \leftrightarrow n, \quad Y_{i,j} \leftrightarrow Y_{j,i}, \tag{5.6}
$$

into the relations [\(5.5\)](#page-10-5), then the cases (m, n) and (n, m) of the linear equation [\(5.1\)](#page-10-3) can be transformed with each other.

We remark that the transformation $z \leftrightarrow T_z$ in [\(5.4\)](#page-10-4) is a kind of q-Laplace transformation, and it transforms $T_z z = q z T_z$ into $z T_z = q T_z z$ i.e. $T_z^{-1} z = q z T_z^{-1}$.

5.2 q-Garnier system as the case $(m, n) = (2N + 2, 2)$

Consider the case $(m, n) = (2N + 2, 2)$ of the matrix A given by

$$
\mathcal{A}(z) = dX_{2N+2} \cdots X_2 X_1,
$$

\n
$$
X_i = \begin{bmatrix} x_{i,1} & 1 \\ r_i^{-1} z & x_{i,2} \end{bmatrix}, \quad d = \text{diag}[t_1, t_2].
$$
\n(5.7)

Proposition 5.2.1 The matrix \mathcal{A} [\(5.7\)](#page-11-1) is equivalent to the Sakai's matrix \mathcal{A} [\(2.4\)](#page-1-1).

Proof. The matrix $\mathcal{A}(5.7)$ $\mathcal{A}(5.7)$ takes the form

(i)
$$
\mathcal{A}(z) = \sum_{i=0}^{N+1} A_i z^i
$$
,
\n(ii) $A_{N+1} = \begin{bmatrix} t_1 \prod_{i=1}^{N+1} r_{2i-1}^{-1} & 0 \\ * & t_2 \prod_{i=1}^{N+1} r_{2i}^{-1} \end{bmatrix}$, $A_0 = \begin{bmatrix} \theta_1 & * \\ 0 & \theta_2 \end{bmatrix}$, (5.8)
\n(iii) $\det \mathcal{A}(z) = t_1 t_2 \prod_{i=1}^{2N+2} r_i^{-1} (z - \alpha_i)$,

where $\theta_j = t_j \prod_{i=1}^{2N+2} x_{i,j}, \ \alpha_i = r_i x_{i,1} x_{i,2}.$ These properties [\(5.8\)](#page-11-2) corresponds exactly to the properties (2.4) for Sakai's matrix A, by a gauge transformation with lower triangular matrix $\begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix}$ ∗ 1 1 .

Conversely, any matrix $\mathcal A$ with properties [\(5.8\)](#page-11-2) can be factorized into the form $\mathcal A$ [\(5.7\)](#page-11-1). Note that the variables $x_{i,j}$ can be obtained by considering the KerA at $z = \alpha_i$ due to the property (iii).

Remark 5.2.2 In the case of differential Garnier system, its relation to the KP hierarchy is not so clear. However, it is known that the Garnier system can been derived through a similarity reduction of the UC hierarchy [\[35\]](#page-19-5).

5.3 Suzuki's system as the case $(m, n) = (2, 2N + 2)$

Consider the case $(m, n) = (2, 2N + 2)$ of the matrix A [\(5.3\)](#page-10-6):

 $\mathcal{A}(z) = dX_2X_1,$

$$
X_{i} = \begin{bmatrix} x_{i,1} & 1 & & & \\ & x_{i,2} & 1 & & \\ & & \ddots & \ddots & \\ & & & x_{i,2N+1} & 1 \\ r_{i}^{-1}z & & & x_{i,2N+2} \end{bmatrix}, \quad d = \text{diag}[t_{1}, \ldots, t_{2N+2}].
$$
 (5.9)

Proposition 5.3.1 The matrix $A(5.9)$ $A(5.9)$ has the following form:

(i)
$$
\mathcal{A}(z) = \begin{bmatrix} \alpha_1 & \varphi_1 & t_1 \\ \alpha_2 & \varphi_2 & t_2 \\ \vdots & \vdots & \ddots \\ \alpha_{2N} & \varphi_{2N} & t_{2N} \\ \frac{t_{2N+1}}{r_1}z & \alpha_{2N+1} & \varphi_{2N+1} \\ \varphi_{2N+2}z & \frac{t_{2N+2}}{r_2}z & \alpha_{2N+2} \end{bmatrix}
$$
, (5.10)
(ii) $\det \mathcal{A}(z) = r_1^{-1} r_2^{-1} \prod_{i=1}^{2N+2} t_i (z - \theta_1)(z - \theta_2)$.

Conversely, any matrix A of the form (5.10) can be factorized as (5.9) .

Proof. It is easy to check that the matrix $\mathcal{A}(5.9)$ $\mathcal{A}(5.9)$ takes the form (5.10) under substitutions $\alpha_i = t_i x_{1,i} x_{2,i}, \ \theta_i = r_i \prod_{j=1}^{2N+2} x_{i,j}, \ \varphi_k = t_k (x_{1,1+k} + x_{2,k}) \quad (k \neq 2N+2) \text{ and } \varphi_{2N+2} =$ $t_{2N+2}(r_2^{-1}x_{1,1}+r_1^{-1}x_{2,2N+2}).$

Conversely, the matrix \mathcal{A} [\(5.10\)](#page-12-0) can be factorized as \mathcal{A} [\(5.9\)](#page-11-3); consider the Ker \mathcal{A} at $z = \theta_i$ using the property (ii).

Remark 5.3.2 In [\[32\]](#page-18-5) T.Suzuki considered the matrix \mathcal{A} [\(5.10\)](#page-12-0), with redundant parameters fixed as $r_1 = \frac{1}{t}$ $\frac{1}{t}$, $r_2 = 1$, $t_1 = \ldots = t_{2N+2} = 1$. Note that two variables among φ_1 , \ldots , φ_{2N+2} can be reduced by a gauge transformation and the constraint coming from the property (ii). He also gave the matrix B corresponding with a time evolution $\overline{r_1} = qr_1$ (*i.e.* $\overline{t} = \frac{t}{a}$ $\frac{t}{q}$), $\overline{\theta_1} = q\theta_1$.

Thanks to Theorem [5.1.1](#page-10-7) and Proposition [5.2.1,](#page-11-4) Proposition [5.3.1,](#page-11-5) we have the conclusion that the q -Garnier system and Suzuki's higher order Painlevé system are related with each other in the sense that they are the deformations of the linear equations equivalent under a q-Laplace transformation.

By the correspondence of parameters $(n = N + 1)$

the Suzuki's direction $\bar{r}_1 = qr_1$, $\theta_1 = q\theta_1$ for [\(5.10\)](#page-12-0) corresponds to $T_3 = T_{c_1}T_{d_1}$ in §[3.2.3.](#page-5-3)

6 Summary

In this paper, extending the previous work [\[22\]](#page-18-1), we obtained the following two main results.

• Choosing the proper variables according to several deformation directions, we obtained simple expression of the scalar Lax pairs and the evolution equations for the variations of the q-Garnier system. Consequently, we gave relations among these q -Garnier systems. As a byproduct, we obtained the standard q -Painlevé equations of type $E_7^{(1)}$ $q^{-1/1}$ as a reduction of a certain q-Garnier system.

• We clarified the relation between the q -Garnier system [\[22,](#page-18-1) [28\]](#page-18-0) and the Suzuki's system [\[32\]](#page-18-5) by formulating the duality of the q -KP system.

A Lax equations

A.1 Correspondence to Sakai's Lax form

The L_1 equations in [\(2.5\)](#page-2-1) and those in §[3.2](#page-4-4) are all equivalent with each other. In this appendix, we give the correspondence^{[9](#page-13-1)} in case of (2.5) and (3.11) as an example.

We call q^{ρ_0} (resp. q^{ρ_∞}) characteristic exponents at $x = 0$ (resp. $x = \infty$) when solutions $y(x)$ have the form

$$
y(x) = k_{\rho_0} x^{\rho_0} (1 + O(x)), \text{ at } x = 0,
$$
 (A.1)

$$
y(x) = k_{\rho_{\infty}} x^{\rho_{\infty}} (1 + O\left(\frac{1}{x}\right)), \quad \text{at } x = \infty.
$$
 (A.2)

Eliminating $\overline{y}(z)$ and $\overline{y}(\frac{z}{a})$ $\frac{z}{q}$) from $L_2(z) = L_2(\frac{z}{q})$ $\frac{z}{q}$) = $L_3(z) = 0$ [\(3.9\)](#page-5-5), we obtain $L_1(z) = 0$ [\(3.11\)](#page-6-0). Then we have

Lemma A.1.1 The L_1 equation [\(3.11\)](#page-6-0) takes the following properties:

(i) It is a linear three term equation between $y(qz)$, $y(z)$ and $y(\frac{z}{q})$ $\frac{z}{q}$), and the coefficients of $y(qz)$, $y(z)$ and $y(\frac{z}{q})$ $\frac{z}{q}$) are polynomials of degree $2N+1$ in z,

(ii) The coefficient of $y(qz)$ (resp. $y(\frac{z}{q})$ $\left(\frac{z}{q}\right)$) has zeros at $z = a_1, \cdots, a_{N+1}$ (resp. $z =$ $q b_1, \cdots, q b_{N+1}$,

(iii) The characteristic exponents of solutions $y(z)$ are d_1, d_2 (at $z = 0$) and c_1, c_2 (at $z = \infty$) generically^{[10](#page-13-2)}.

(iv) The N points z such that $F(z) = 0$ are apparent singularities (i.e., the solutions are regular there), where

$$
\frac{y(qz)}{y(z)} = \frac{G(z)}{A(z)}, \quad \text{for} \quad F(z) = 0,
$$
\n(A.3)

holds (c.f. [\(2.11\)](#page-2-5)).

Conversely, the L_1 equation [\(3.11\)](#page-6-0) is uniquely characterized by these properties (i)–(iv). \Box

On the other hand, owing to Lemma [2.2.1,](#page-1-6) Sakai's Lax equation [\(2.1\)](#page-1-2) is rewritten into the L_1 equation [\(2.5\)](#page-2-1). Furthermore we have

⁹Other cases are similar. The case of (2.5) and (3.5) is given in $[22, §2.2]$ $[22, §2.2]$.

¹⁰Generic case means $f_0 f_N \neq 0$. We remark that the reduced cases to q - $E_7^{(1)}$ system are nongeneric, where their L_1 equations [\(4.9\)](#page-8-5) and [\(4.14\)](#page-9-4) have exponents d_1 , qd_1 (at $z = 0$) and c_1 , $\frac{c_1}{q}$ ($z = \infty$).

Lemma A.1.2 The L_1 equation [\(2.5\)](#page-2-1) can be expressed as a form

$$
L_1: \prod_{i=1}^{N+1} (z - \alpha_i) F(\frac{z}{q}) y(qz) - \left\{ F(\frac{z}{q}) \mathcal{A}_{11}(z) + F(z) \mathcal{A}_{22}(z) \right\} y(z) + \kappa_1 \kappa_2 \prod_{i=N+2}^{2N+2} (\frac{x}{q} - \alpha_i) F(z) y(\frac{z}{q}) = 0.
$$
 (A.4)

 \Box

 \blacksquare

 \Box

Proof. The proof is given by using a gauge transformation:

$$
y(z) \mapsto H(z)y(z)
$$
, with $\frac{H(qz)}{H(z)} = \prod_{i=1}^{N+1} (z - \alpha_i)$. (A.5)

Similarly to Lemma [A.1.1,](#page-13-3) we have

Lemma A.1.3 Then the L_1 equation $(A.4)$ has the following properties:

(i) It is a linear three term equation between $y(qz)$, $y(z)$ and $y_1(\frac{z}{q})$ $\frac{z}{q}$), and the coefficients of $y(qz)$, $y(z)$ and $y(\frac{z}{q})$ $\frac{z}{q}$) are polynomials of degree $2N+1$ in z,

(ii) The coefficient of $y(qz)$ (resp. $y(\frac{z}{q})$ $(\frac{z}{q})$) has zeros at $z = \alpha_1, \cdots, \alpha_{N+1}$ (resp. $z =$ $q\alpha_{N+2}, \cdots, q\alpha_{2N+2}$.

(iii) The characteristic exponents of the solutions $y(z)$ are $\frac{\theta_1}{\prod_{i=1}^{N+1}(-\alpha_i)}, \frac{\theta_2}{\prod_{i=1}^{N+1}(-\alpha_i)}$ (at $z = 0$) and $\kappa_1, q^{-1}\kappa_2$ (at $z = \infty$),

(iv) N points $z = \lambda_i$ such that $F(z) = 0$ are apparent singularities, where

$$
\frac{y(qz)}{y(z)} = \frac{A_{11}(z)}{\prod_{i=1}^{N+1} (z - \alpha_i)}, \quad \text{for} \quad F(z) = 0,
$$
\n(A.6)

holds.

Conversely, the L_1 equation $(A.4)$ is uniquely characterized by these properties $(i)–(iv)$.

Hence we obtain

Proposition A.1.4 The L_1 equations [\(3.11\)](#page-6-0) and [\(2.5\)](#page-2-1) are equivalent up to the gauge transformation $(A.5)$ and changes of variables and parameters.

Proof. Thanks to Lemmas [A.1.1,](#page-13-3) [A.1.2](#page-13-4) and [A.1.3.](#page-14-3)

A.2 Sufficiency for the compatibility

In §[3.2,](#page-4-4) we stated that the equations in item (b) are sufficient for the compatibility of the equations in item (a). Here we will prove this fact in the case of $\S 3.2.3$ $\S 3.2.3$ as an example^{[11](#page-14-4)}.

¹¹Other cases can be proved in the similar way and the case of $\S 3.2.1$ $\S 3.2.1$ has been done in [\[22,](#page-18-1) $\S 2.3$].

Eliminating $y(z)$ and $y(qz)$ from $L_2(z) = L_3(z) = L_3(qz) = 0$ [\(3.9\)](#page-5-5), we obtain the following expression:

$$
L_4(z) = \overline{A}(z)\overline{F}(\frac{z}{q})\overline{y}(qz) + q^2 c_1 c_2 \overline{B}(\frac{z}{q})\overline{F}(z)\overline{y}(\frac{z}{q})
$$

$$
-\left\{q\overline{F}(z)G(\frac{z}{q}) + \frac{\overline{F}(\frac{z}{q})}{G(z)}V(z)\right\}\overline{y}(x),
$$
 (A.7)

where V is given in (3.11) .

Lemma A.2.1 The L_4 equation $(A.7)$ has the following properties:

(i) It is a linear three term equation between $\overline{y}(qz)$, $\overline{y}(z)$ and $\overline{y}(\frac{z}{q})$ $\frac{z}{q}$), and the coefficients of $\overline{y}(qz)$, $\overline{y}(z)$ and $\overline{y}(\frac{z}{q})$ $\frac{z}{q}$) are polynomials of degree $2N+1$ in z,

(ii) The coefficient of $\overline{y}(qz)$ (resp. $\overline{y}(\frac{z}{qz})$ $(\frac{z}{q})$) has zeros at $z = a_1, a_2, \cdots, a_{N+1}$ (resp. $z = qb_1, qb_2, \cdots, qb_{N+1}),$

(iii) The characteristic exponents of solutions $\bar{y}(z)$ are qd_1, d_2 (at $z = 0$) and qc_1, c_2 $(at z = \infty),$

(iv) The N points z such that $\overline{F}(z) = 0$ are apparent singularities, where

$$
\frac{\overline{y}(qz)}{\overline{y}(z)} = \frac{qc_1c_2B(z)}{G(z)}, \quad \text{for} \quad \overline{F}(z) = 0,
$$
\n(A.8)

holds.

Conversely, the equation $L_4 = 0$ [\(A.7\)](#page-15-1) is uniquely characterized by these properties $(i)–(iv).$

Proposition A.2.2 The linear q-difference equations L_1 [\(3.11\)](#page-6-0) and L_2 [\(3.9\)](#page-5-5) are compatible if and only if the bi-rational equation (3.10) .

Proof. The compatibility of L_1 and L_2 means that $T(L_1) = L_4$. This can be checked by the characterizations of the equations L_1 (resp. L_4) in Lemma [A.1.1](#page-13-3) (resp. [A.2.1\)](#page-15-2).

B From Padé interpolation to the q -Garnier system

There is a convenient method to approach the continuous/discrete Painlevé equations by using certain problem of Padé approximation $[19, 37]$ $[19, 37]$ (see also $[13, 14]$ $[13, 14]$) or interpolation $[7, 17, 18, 20, 21, 24, 39, 40]$ $[7, 17, 18, 20, 21, 24, 39, 40]$ $[7, 17, 18, 20, 21, 24, 39, 40]$ $[7, 17, 18, 20, 21, 24, 39, 40]$ $[7, 17, 18, 20, 21, 24, 39, 40]$ $[7, 17, 18, 20, 21, 24, 39, 40]$ $[7, 17, 18, 20, 21, 24, 39, 40]$ $[7, 17, 18, 20, 21, 24, 39, 40]$. In $[22]$, this method is applied to the q-Garnier system in the case of §[3.2.1.](#page-4-2) Here, we illustrate the derivation of the Lax pair [\(3.9\)](#page-5-5) through the Padé method in case of q-Garnier system in $\S 3.2.3$ $\S 3.2.3$ as an example.

B.1 Derivation of scalar Lax pair

Fix a positive integer $N \in \mathbb{N}$ and a parameter q $(0 \lt |q| \lt 1)$. For parameters $a_1, \ldots, a_N, b_1, \ldots, b_N$ and c, we consider a function

$$
\psi(z) = c^{\log_q z} \prod_{i=1}^N \frac{(a_i z, b_i)_{\infty}}{(a_i, b_i z)_{\infty}}.
$$
\n(B.1)

Let $P(z)$ and $Q(z)$ be polynomials of degree m and $n \in \mathbb{Z}_{\geq 0}$ in z determined by the following Padé interpolation condition:

$$
\psi(z_s) = \frac{P(z_s)}{Q(z_s)}, \quad (z_s = q^s, s = 0, 1, \dots m + n). \tag{B.2}
$$

The common normalizations of the polynomials P and Q in z are fixed as $P(0) = 1$. We consider the shift operation: $\overline{x} = T(x)$ where $T = T_c$.

We construct two linear q-difference relations: $L_2 = 0$ among $y(z), y(qz), \overline{y}(z)$ and $L_3 = 0$ among $y(z), \overline{y}(z), \overline{y}(\frac{z}{a})$ $\frac{z}{q}$) satisfied by the functions $y = P$ and $y = \psi Q$.

Proposition B.1.1 The linear relations L_2 and L_3 can be expressed as

$$
L_2(z) = F(z)\overline{y}(z) + A(z)\left(\frac{z}{q^{m+n}}\right)_1 y(qz) + \frac{c}{g_0}G(z)y(z) = 0,
$$

\n
$$
L_3(z) = \frac{z}{q}\overline{F}(\frac{z}{q})y(x) + \frac{1}{g_0}G(\frac{z}{q})\overline{y}(z) - (z)_1 B(\frac{z}{q})\overline{y}(\frac{z}{q}) = 0,
$$
\n(B.3)

where $A(z) = \prod_{i=1}^{N} (a_i z)_1$, $B(z) = \prod_{i=1}^{N} (b_i z)_1$ and $F(z)$, $G(z)$ are as in [\(3.9\)](#page-5-5) and $f_0, \ldots, f_N, g_0, \ldots, g_N$ are constants depending on parameters a_i, b_j, c and $m, n \in \mathbb{Z}_{\geq 0}$. \Box

Proof. By the definition above, the linear relations L_2 and L_3 can be written as

$$
L_2(z) \propto \begin{vmatrix} y(z) & y(qz) & \overline{y}(z) \\ \mathbf{y}(z) & \mathbf{y}(qz) & \overline{\mathbf{y}}(z) \end{vmatrix} = D_1(z)\overline{y}(z) - D_2(z)y(qz) + D_3(z)y(z) = 0,
$$

\n
$$
L_3(z) \propto \begin{vmatrix} y(z) & \overline{y}(z) & \overline{y}(\frac{z}{q}) \\ \mathbf{y}(z) & \overline{\mathbf{y}}(z) & \overline{\mathbf{y}}(\frac{z}{q}) \end{vmatrix} = \overline{D}_1(\frac{z}{q})y(z) + D_3(\frac{z}{q})\overline{y}(z) - D_2(z)\overline{y}(\frac{z}{q}) = 0,
$$
\n(B.4)

where $\mathbf{y}(z) = \begin{bmatrix} P(z) \\ P(z) \end{bmatrix}$ $\psi(z)Q(z)$ 1 and Casorati determinants $D_1(z) = |\mathbf{y}(z), \mathbf{y}(qz)|, \quad D_2(z) = |\mathbf{y}(z), \mathbf{\bar{y}}(z)|, \quad D_3(z) = |\mathbf{y}(qz), \mathbf{\bar{y}}(z)|.$ (B.5)

Using the relations

$$
\frac{\psi(qz)}{\psi(z)} = c \frac{B(z)}{A(z)}, \quad \frac{\overline{\psi}(z)}{\psi(z)} = z,
$$
\n(B.6)

we can rewrite the Casorati determinants [\(B.5\)](#page-16-0) into the following determinants:

$$
D_1(z) = \frac{\psi(z)}{A(z)} \{cB(z)P(z)Q(qz) - A(z)P(qz)Q(z)\} =: \frac{\psi(z)}{A(z)} \prod_{i=0}^{m+n-1} (\frac{z}{q^i})_1 c_0 F(z),
$$

\n
$$
D_2(z) = \psi(z) \{zP(z)\overline{Q}(z) - \overline{P}(z)Q(z)\} =: \psi(z) \prod_{i=0}^{m+n} (\frac{z}{q^i})_1 c_0,
$$

\n
$$
D_3(z) = \frac{\psi(z)}{A(z)} \{zA(z)P(qz)\overline{Q}(z) - cB(z)\overline{P}(z)Q(qz)\} =: \frac{\psi(z)}{A(z)} \prod_{i=0}^{m+n-1} (\frac{z}{q^i})_1 G(z).
$$
\n(B.7)

Then the constants $c_0 =$ \overline{g}_0 $\frac{g_0}{c},\ g_{N+1}=-\frac{g_0\prod_{i=1}^N(-a_i)}{cq^n}$ $\frac{a_{i-1}(\cdots a_i)}{cq^n}$ are determined through the expansions around $x = 0$ and $x = \infty$. As a result, we obtain the desired equations.

Acknowledgment. The authors shall thank Professors Kenji Kajiwara, Tetsu Masuda, Masatoshi Noumi, Hidetaka Sakai, Takao Suzuki and Teruhisa Tsuda for stimulating comments. This work is partially supported by JSPS KAKENHI (26287018) and Expenses Revitalizing Education and Research of Akashi College (0217030).

References

- [1] Doliwa A., Non-cmmutative rational Yang-Baxter maps, Lett. Math. Phys., 104, $(2014), 299-309.$
- [2] Dzhamay A., Sakai H., and Takenawa T., Discrete Hamiltonian Structure of Schlesinger Transformations, [arXiv:1302.2972](http://arxiv.org/abs/1302.2972) [math-ph].
- [3] Dzhamay A., and Takenawa T., Geometric Analysis of Reductions from Schlesinger transformations to difference Painlevé equations, [arXiv:1408.3778](http://arxiv.org/abs/1408.3778) [math-ph].
- [4] Gasper G., and Rahman M., Basic Hypergeometric Series. With a foreword by Richard Askey. Second edition. Encyclopedia of Mathematics and its Applications, 96. Cambridge University Press Cambridge, (2004).
- [5] Grammaticos B., Ramani A., On a novel q -discrete analogue of the Painlevé VI equation, Phys. Lett. A, 257 (1999), 288–292.
- [6] Hasegawa K., Quantizing the discrete Painlevé VI equation: the Lax formalism, Lett. Math. Phys., 103 (2013), 865–879.
- [7] Ikawa Y., *Hypergeometric Solutions for the q-Painlevé Equation of Type* $E_6^{(1)}$ $b_6^{(1)}$ by the Padé method, Lett. Math. Phys., 103 (2013), 743-763.
- [8] Jimbo M. and Sakai H., A q-analog of the sixth Painlevé equation, Lett. Math. Phys., 38 (1996), 145–154.
- [9] Kajiwara K., Noumi M., and Yamada Y., *Discrete dynamical systems with* $W(A_{m-1}^{(1)} \times$ $A_{n-}^{(1)}$ $_{n-1}^{(1)}$) symmetry, Lett. Math. Phys., 60 (2002), 211–219.
- [10] Kajiwara K., Noumi M., and Yamada Y., q -Painlevé Systems Arising from q -KP Hierarchy, Lett. Math. Phys., 62 (2002), 259–268.
- [11] Kajiwara K., Noumi M., and Yamada Y., *Geometric aspects of Painlevé equations*, J. Phys. A: Math. Theor., 50 (2017), 073001(164pp) (Topical Review).
- [12] Kuniba A., Okado M., Takagi T., and Yamada Y., Geometric crystal and tropical R for $D_n^{(1)}$, Int. Math. Res. Not., 5 (2003), 2565–2620.
- [13] Mano T., Determinant formula for solutions of the Garnier system and Padé approximation. J. Phys. A: Math. Theor., 45 (2012), 135206–135219.
- [14] Mano T., and Tsuda T., *Hermite-Padé approximation, isomonodromic deformation* and hypergeometric integral. Math. Z., 285 (2017), no. 1-2, 397–431.
- [15] Masuda T., On the rational solutions of q-Painlevé V equation, Nagoya Math. J., 169 (2003), 119–143.
- [16] Masuda T., *Hypergeometric* τ -functions of the q-Painlevé system of type $E_7^{(1)}$,(1)
7, SIGMA, 5 (2009), 035 (30pp).
- [17] Nagao H., The Padé interpolation method applied to q -Painlevé equations. Lett. Math. Phys., 105 (2015), 503–521.
- [18] Nagao H., Lax pairs for additive difference Painlevé equations, $arXiv:1604.02530$ [nlin.SI].
- [19] Nagao H., The Padé interpolation method applied to q-Painlevé equations II (differential grid version), Lett. Math. Phys., 107 (2017), 107–127.
- [20] Nagao H., A variation of the q-Painlevé system with affine Weyl group symmetry of type $E_7^{(1)}$ $T_7^{(1)}$, [arXiv:1706.10087](http://arxiv.org/abs/1706.10087) [nlin.SI].
- [21] Nagao H., *Hypergeometric special solutions for d-Painlevé equations*, [arXiv:1706.10101](http://arxiv.org/abs/1706.10101) [nlin.SI].
- [22] Nagao H., and Yamada Y., *Study of q-Garnier system by Padé method*, Funkcialaj Ekvacioj, to appear, [arXiv:1601.01099](http://arxiv.org/abs/1601.01099) [nlin.SI].
- [23] Nijhoff F., and Delice N., On elliptic Lax pairs and isomonodromic deformation systems for elliptic lattice equations, [arXiv:1605.00829](http://arxiv.org/abs/1605.00829) [nlin.SI].
- [24] Noumi M., Tsujimoto S., and Yamada Y., *Padé Interpolation for Elliptic Painlevé* Equation, Springer 40 (2013), 463–482.
- [25] Ormerod, C.M., and Rains E.M., Commutation Relations and Discrete Garnier Systems, SIGMA, 12 (2016), 110, 50 pages.
- [26] Ormerod, C.M., and Rains E.M., An elliptic Garnier system, [arXiv:1607.07831](http://arxiv.org/abs/1607.07831) [nlin.SI].
- [27] Sakai H., Rational surfaces with affine root systems and geometry of the Painlevé equations, Commun. Math. Phys. 220 (2001), 165–221.
- [28] Sakai H., A q-analog of the Garnier system, Funkcialaj Ekvacioj, 48 (2005), 273–297.
- [29] Sakai H., Hypergeometric Solution of q-Schlesinger System of Rank Two, Lett. Math. Phys., 73 (2005), 237–247.
- [30] Sakai H., Lax form of the q-Painlevé equation associated with the $A_2^{(1)}$ $_2^{(1)}$ surface, J. Phys. A: Math. Gen., 39 (2006), 12203–12210.
- [31] Suzuki T., A q-analogue of the Drinfeld-Sokolov hierarchy of type A and q -Painlevé system, AMS Contemp. Math., 651 (2015), 25–38.
- [32] Suzuki T., A reformulation of the generalized q-Painlevé VI system with $W(A_{2n+1}^{(1)})$ symmetry, J. Integrable Syst., 2 (2017), 1–18.
- [33] Takenawa T., Weyl group symmetry of type $D_5^{(1)}$ $_5^{(1)}$ in the q-Painlevé V equation, Funkcialaj Ekvacoj, 46 (2003), 173–186.
- [34] Tsuda T., On an integrable system of q-difference equations satisfied by the universal $characters:$ its Lax formalism and an application to q -Painlevé equations, Comm. Math. Phys., 293 (2010), 347–359.
- [35] Tsuda T., UC hierarchy and monodromy preserving deformation, J. reine angew. Math., **690** (2014), 1–34.
- [36] Yamada Y., A birational representation of Weyl group, combinatorial R-matrix and discrete Toda equation, in "Physics and combinatorics, 2000", World Sci. Publ., (2001), 305–319.
- [37] Yamada Y., *Padé method to Painlevé equations*, Funkcialaj Ekvacoj, 52 (2009), 83– 92.
- [38] Yamada Y., Lax formalism for q -Painlevé equations with affine Weyl group symmetry of type $E_n^{(1)}$, Int. Math. Res. Not., 17 (2011), 3823-3838.
- [39] Yamada Y., A simple expression for discrete Painlevé equations, RIMS Kokyuroku Bessatsu, B47 (2014), 087–095.
- [40] Yamada Y., An elliptic Garnier system from interpolation, [arXiv:1706.05155](http://arxiv.org/abs/1706.05155) [mathph].