## The Quantum Anomalous Hall Majorana Platform

Yongxin Zeng,<sup>1</sup> Chao Lei,<sup>2</sup> Gaurav Chaudhary,<sup>2</sup> and Allan H. MacDonald<sup>2</sup>

<sup>1</sup>CAS Key Laboratory of Strongly-Coupled Quantum Matter Physics, and Department of Physics,

<sup>2</sup>Department of Physics, The University of Texas at Austin, Austin, Texas 78712, USA

(Dated: June 12, 2022)

We show that quasi-one-dimensional (1D) quantum wires can be written onto the surface of magnetic topological insulator (MTI) thin films by gate arrays. When the MTI is in a quantum anomalous Hall (QAH) state, MTI/superconductor quantum wires have especially broad stability regions for both topological and non-topological states, facilitating creation and manipulation of Majorana particles on the MTI surface.

PACS numbers: 71.70.Ej, 71.10.Pm, 74.45.+c,

Introduction:- Non-Abelian anyons that can be localized at controllable positions provide an attractive platform for fault-tolerant quantum computation[1–7]. Candidate non-Abelian particles that have already been realized in solid state systems include fractionally charged excitations of the  $\nu = 5/2$  fractional quantum Hall liquid [8], Abrikosov vortices in two-dimensional topological superconductors [9], and Majorana zero mode (MZM) end states of *p*-wave superconducting quantum wires[10–13]. In this article we propose a new strategy for creation of MZMs and manipulations of MZM positions on a twodimensional surface.

MZMs appear at boundaries between normal and topological quasi-1D superconductors [10, 12–14]. The simplest 1D topological superconductor (TSC) consists of spinless electrons with near-neighbor attractive interactions [10]. Topological superconducting states can occur, however, in any quasi-1D system with superconductivity and broken time-reversal symmetry [11]. Much progress has been achieved both theoretically [12, 13, 15] and experimentally [16–21] by studying strongly spin-orbitcoupled proximitized quantum wires with large q-factors that are perturbed by an external magnetic field, and also by placing magnetic-atom arrays on superconducting substrates [22]. Here we suggest an alternate approach, based on the QAH state [23–25] of MTIs, that has potentially valuable advantages. We show that ribbons formed from MTIs are often topological when proximity-coupled to a superconductor. MZMs also appear when quasi-1D topological regions are written onto a portion of the surface of a MTI by external gates, as illustrated schematically in Fig. 1. Furthermore, the MTI/SC phase diagram has particularly broad regions of stability for both normal and topological phases when the isolated MTI is close to a phase boundary between normal insulator and QAH states. [23, 24] This property provides a practical route toward electrical control of MZM positions.

Theoretical model of a proximized MTI:- We focus on Cr-doped Bi<sub>2</sub>Se<sub>3</sub> thin films in which proximitized superconductivity has already been demonstrated [24, 25] experimentally, and which are close to the MTI's QAH in-



FIG. 1: Majorana zero modes can be created locally by separating the surface of a magnetic topological insulator in a quantum Anomalous Quantum Hall state into alternating normal and topological regions using remote gates.

sulator/normal insulator phase boundary when 4-6 quintile layers thick. In the  $\psi_{\mathbf{k}} = (c_{\mathbf{k}\uparrow}^t, c_{\mathbf{k}\downarrow}^t, c_{\mathbf{k}\uparrow}^b, c_{\mathbf{k}\downarrow}^b)^T$  basis, where  $c_{\mathbf{k}\sigma}^l$  annihilates an electron with momentum  $\mathbf{k}$  spin  $\sigma = \uparrow, \downarrow$  and surface s = t/b (top/bottom), the single particle Hamiltonian of a MTI thin film at energies below the bulk gap is

$$H_0(\mathbf{k}) = h_{\rm D}(\mathbf{k})\tau_z + m_k\tau_x + \lambda\sigma_z + \lambda'\tau_z \tag{1}$$

where  $\tau_i$  acts on surface and  $\sigma_i$  on spin. In Eq. 1  $h_D(\mathbf{k}) = v(k_y\sigma_x - k_x\sigma_y)$  is the 2D Dirac isolated surface Hamiltonian with Fermi velocity v;  $m_k = m_0 + m_1(k_x^2 + k_y^2)$  accounts for hybridization between top and bottom surfacess,  $\lambda$  is the exchange field produced by the ferromagnetically ordered magnetic dopants; and  $\lambda'$  accounts for the energetic displacement between Dirac cones on top on bottom surfaces produced by vertical electric fields in the bulk of the TI. When placed in proximity to an *s*-wave SC, the system is described by the Bogoliubov-de Gennes (BdG) Hamiltonian  $H(\mathbf{k}) = \sum_{\mathbf{k}} \psi_{\mathbf{k}}^{\dagger} H_{\text{BdG}}(\mathbf{k}) \psi_{\mathbf{k}}/2$ , where

$$H_{\rm BdG}(\boldsymbol{k}) = \begin{pmatrix} H_0(\boldsymbol{k}) - \mu & \Delta_{\rm sc} \\ \Delta_{\rm sc}^{\dagger} & -H_0^*(-\boldsymbol{k}) + \mu \end{pmatrix}$$
(2)

University of Science and Technology of China, Hefei, Anhui 230026, China.

with

$$\Delta_{\rm sc} = \begin{pmatrix} i\Delta_t \sigma_y & 0\\ 0 & i\Delta_b \sigma_y \end{pmatrix}.$$
 (3)

A regularized square-lattice version of Eq. 2 with nearest neighbor hopping<sup>[26]</sup> accurately and conveniently captures the physics of topological phase transitions (TPT) at the surface of MTI. In the lattice model  $k_{x/y}$  is replaced by  $\sin k_{x/y}$  and  $k^2$  by  $2(2 - \cos k_x - \cos k_y)$ . We absorb the lattice constant into the wavevectors in Eq. 1 to make them dimensionless, so that  $m_0, m_1$  and v all have dimensions of energy. The inter-surface hybridization parameter  $m_1$  plays an essential role by preventing the appearance of unphysical states at low energies away from  $\mathbf{k} = 0$ . For  $m_0 \ll m_1$  only the  $\Gamma$ -point avoided crossing is relevant at low energies. We can obtain realistic values of v,  $m_0$  and  $m_1$  by comparing the model spectrum with DFT band structures of  $Bi_2Se_3$ , like the one illustrated Fig. S1 of the Supplemental Material. For a five layer film we find that the gap at  $\Gamma$  is about 12 meV, giving  $m_0 = 6 \text{ meV}$ . Rough estimates for  $m_1$ and v can be obtained by fitting to the DFT Dirac velocity  $v ~(\sim 4 \times 10^5 \text{ms}^{-1})$  and the DFT gaps at M and K. We find that  $m_1 \sim 0.2 \,\mathrm{eV}$  and  $v \sim 0.7 \,\mathrm{eV}$  for a lattice model with lattice constant  $a \approx 4$ Å.

Topological classification of 2D bulk and 1D ribbon states:- The 2D classification of MTI/SC states presented in Ref. 27 demonstrates the possibility of BdG bands with odd Chern numbers. Motivated by the simplifications that occur when  $\Delta_t = -\Delta_b = \Delta$  recognized in that paper, we first address that special case and set  $\lambda' = \mu = 0$ to obtain simple analytical expressions for phase boundaries. When the basis transformation used in Ref. 27 is employed the  $8 \times 8$  BdG Hamiltonian block diagonalizes into four  $2 \times 2$  matrices, each of which has the form of a spinless  $p \pm ip$  superconductor. The gaps close at  $\Gamma$ for  $\pm \lambda \pm \Delta = m_0$ , and at X, Y and M Brillouin-zone points when the other gap-closing conditions are satisfied. Because  $m_0$ ,  $\lambda$  and  $\Delta$  are all comparable in size in typical samples, the  $\Gamma$  gap-closing condition is readily satisfied. The other gap closing conditions cannot be satisfied, however, when realistic values are used for  $m_1$ . For  $|\lambda| > m_0$ , *i.e.* when the quantum anomalous Hall effect occurs, weak pairing gives rise to superconducting states with Chern numbers  $\pm 2$ , corresponding to Nambu-doubled quantum Hall edge states. As explained below, we propose using these states as a resource for MZM formation. (The Chern number phase diagram of our proximitized MTI model is described in detail in the Supplementary Material.)

Our main interest is in the possibility of gate-defined quasi-1D topological regions embedded within the 2D film surface. To set the stage, however, we first address the closely related case of narrow ribbons.[10, 28, 29] For width W = 1, where W is the number of lattice model rows, the  $\Delta_t = -\Delta_b$  BdG Hamiltonian reduces



FIG. 2:  $Z_2$  ribbon phase diagram in  $(\lambda, \mu)$  space for  $\lambda' = 0, m_0 = 6$  meV,  $m_1 = 200$  meV,  $\Delta_b = 1$  meV,  $\Delta_t = 0$ , ribbon width W = 300, and velocity v = 45 meV. The orange regions are topologically non-trivial and support MZMs.

to four effective Kitaev chains with hopping terms that have different functional dependences on  $m_0, m_1, \lambda$ , and  $\Delta$  and different chemical potentials. Each yields MZMs localized near the chain ends when topologically nontrivial. The Supplementary material contains a detailed summary of how the the  $Z_2$  (1D) topological phase diagram of the  $\Delta_t = -\Delta_b$  model depends on W. In Fig. 2 we plot the  $Z_2$  phase diagram in  $(\lambda, \mu)$  space for a ribbon with proximity coupling only to the bottom MTI surface. The  $Z_2$  invariant is evaluated using  $\nu = \text{sgn}[\text{Pf}\tilde{H}(0)]\text{sgn}[\text{Pf}\tilde{H}(\pi)] = -1$ , where  $\tilde{H}$  is the skewsymmetrized Hamiltonian, Pf denotes the Pfaffian number and sgn[x] is the sign of x. For weak pairing  $\nu = \pm 1$ depending on whether the number of normal state bands that cross the Fermi level is even or odd.

The broad region of topologically nontrivial behavior at large  $\lambda$  and small  $\mu$  in Fig. 2, *i.e.* when the unproximitized MTI is in a QAH state, reflects the property that only a single ribbon band is present in this energy range at any value of W. The  $Z_2$  invariant is much more sensitive to  $\lambda$ ,  $\mu$ , W and other ribbon model parameters at larger values of  $\mu$  that lie within the gapped surface state bands. For a lattice model of a normal 2D p-wave superconductor, for example, there are 4W phase boundaries in the quasi-1D  $Z_2$  classification of ribbon states at width W. This sensitivity is not favorable for reliable realization of either trivial or non-trivial states. In the QAH state there is at most a single band, but the MZMs present for finite ribbon length are protected only by ex-



FIG. 3: Phase diagram of a MTI on a superconducting substrate in  $\lambda' - \mu$  space. Here  $\Delta_b = 0.2m_0$ ,  $\Delta_t = 0$ , and from left to right  $\lambda = 0.5m_0$ ,  $m_0$ , and  $2m_0$ .

ponentially small superconducting gaps (~  $\Delta e^{-W/\xi_P}$ ), where  $\xi \sim_P v/(\lambda - m_0)$  is the 2D edge state localization length in lattice constant units, making the  $Z_2$  classification academic. (For typical parameters  $\xi \sim 10$ nm in physical units.) (The phase diagram in Fig. 2 was calculated using a Dirac velocity that is unrealistically small for this reason.) In order to obtain MZMs that are reasonably localized near ribbon edges it is necessary to have ribbon widths that are not too large compared to  $\xi$ , and also to be able to conveniently tune between topologically trivial and non-trivial states. Since the exchange coupling parameter  $\lambda$  in Fig. 2 is fixed for a given MTI sample and a given operating temperature, a different tuning parameter must be identified.

Quantum Anomalous Hall Ribbon Majoranas:- We propose controlling MZMs in ribbons by placing a MTI film that supports a QAH state on a superconducting substrate and fabricating a top gate. Varying the gate field will alter the carrier density (and hence the chemical potential  $\mu$ ) of the MTI, and also shift the energy of the top surface Dirac cone relative to the bottom surface. The latter effect is due to the unscreened portion of the gate electric field that survives in the interior of the TI and is represented in our model by the parameter  $\lambda'$ . Applying a gate voltage moves the system along a line in  $(\mu, \lambda')$  space that depends on the effective bulk dielectric constant of the TI. As we now show a gate voltage can therefore tune the proximitized MTI between  $Z_2 = 0$  and  $Z_2 = 1$  ribbon states.

Fig. 3 illustrates 2D Chern number phase diagrams in  $(\lambda',\mu)$  space for MTIs on superconducting substrates with  $\Delta_b = 0.2m_0$  and  $\Delta_t = 0$ , at three different values of exchange field  $\lambda$ . For  $\lambda < m_0$  (left panel), the unproximitzed MTI is in a normal insulator state, but superconductivity that is sufficiently strong can still induce odd Chern number BdG states. Our main interest is in the case in the right panel where  $\lambda > m_0$  so that the unproximitized MTI is in a quantum anomalous Hall state at  $\lambda' = 0$ . Proximitized superconductivity of quantum anomalous Hall states is now routinely achieved experimentally.[24, 25] For  $\sqrt{\lambda'^2 + m_0^2} > |\lambda|$  The QAH state occurs at small  $|\mu$  only when  $\sqrt{\lambda'^2 + m_0^2} > |\lambda|$ ; the gate field  $\lambda'$  efficiently converts a QAH insulator with edge states into a normal insulator with no edge states in the gap, and no opportunity for  $Z_2$  states in ribbons.

For weak pairing, ribbons have  $Z_2 = 1$  states when an odd number of subbands cross the Fermi level. In a ribbon formed from a semiconductor quantum well with strong Zeeman and spin-orbit coupling, or from a MTI with  $m_0 \gg \lambda$ , a series of closely spaced non-degenerate 1D sub-bands appear close to the extrema of the bulk 2D bands, as illustrated schematically in Fig. 4a. When the ribbon width increases, the spacing between subbands becomes smaller and the  $Z_2$  phase diagrams will have more closely spaced boundaries between  $Z_2 = 0$ and  $Z_2 = 1$  phases. In the  $\lambda \gg m_0$  QAH case on the other hand, illustrated schematically in Fig. 4a, a single pair of bands crosses the Fermi level at all energies inside the bulk 2D gap, except for a narrow gapped region due to the avoided crossing between QAH edge states localized on opposite sides of the ribbon. The system is therefore almost always nontrivial in the broad range  $\mu \in (-m_0, m_0)$ , independent of the ribbon width.

In Fig. 4c we plot the  $Z_2$  phase diagram  $(\mu, \lambda')$  plane of  $W = 300, m_0 = 6 \text{ meV}, m_1 = 0.2 \text{ eV}, v = 0.7 \text{ eV}, \lambda =$  $10 \text{ meV}, \Delta_t = 1 \text{ meV}, \Delta_b = 0 \text{ ribbon, which exhibits large}$ adjacent trivial and nontrivial regions near  $\mu = 0$ . For these realistic parameters  $|\lambda|$  is larger than  $m_0$ , but not much larger. (Larger QAH regions can be achieved by going to thicker films with smaller  $m_0$  but only at the expense of reducing all relevant energy scales.) QAH edge states overlap strongly even at W = 300, so strongly in fact that the normal state gap produced by avoided crossing of edge states is comparable to  $m_0$ , creating a substantial  $Z_2 = 0$  region near  $\mu = 0$ . This region is however still bordered at  $\lambda' = 0$  by a large  $Z_2 = 1$  region that can be identified with pairing of QAH edge states. The gate field  $\lambda'$  sweeps this state into a large adjacent  $Z_2 = 0$  region, which can be identified with a proximity coupled ordinary insulator.



FIG. 4: (a) Bandstructure of a nanoribbon. When the Fermi level (the red dotted line) coincides with the bottom of a subband, a TPT occurs; (b) Bandstructure in the QAH regime. Grey regions are conduction and valence bands, blue curves are edge states with a small hybridization gap, and the red dotted line shows the Fermi level. When the chemical potential is inside the bulk gap and outside the hybridization gap, with SC the system should always be nontrivial. (c) Phase diagram in  $\lambda'$ - $\mu$  space for a quasi-1D chain with  $W = 300, m_0 = 6 \text{ meV}, m_1 = 0.2 \text{ eV}, v = 0.7 \text{ eV}, \lambda = 12 \text{ meV}, \Delta_t = 1 \text{ meV}, \Delta_b = 0$ ; (d) Relation between gap and width, with  $\mu = 5 \text{ meV}, \lambda' = 0$  and other parameters the same as in (c).

Fig. 4d plots the ribbon width dependence of the gap at  $k_x = 0$ ,  $\lambda' = 0$ , and  $\mu = 5 \text{ meV}$  out to  $W \sim 1000$ in lattice constant units. Here we notice that there is only one gap closing as a function of W which signals a phase transition between a small  $W Z_2 = 0$  state associated with a large avoided crossing gap between QAH edge states, and a large  $W Z_2 = 1$  state associated with pairing of QAH edge states. The gaps remain large out to  $\sim 1000$  lattice constants, corresponding to a physical length  $\sim 400 \,\mathrm{nm}$ , partly because the edge state localization length is enhanced by the relatively high velocity of quantum Hall edge states compared to the velocity of states present near the bottom of a bulk 2D band. As the system becomes wider, the QAH edge states have less overlap and the gap is eventually reduced to very small values.

Because of the bulk-edge correspondence of topological states we expect Majorana zero modes to appear not only in ribbons, but also in wide films in which quasione-dimensional regions are formed that have local model parameters in the topologically non-trivial range. In order to define a quasi-1D region, we can form a gate array above a proximitized MTI thin film whose chemical potential is in the QAH gap, as illustrated schematically in Fig. 1. Just as in the ribbon case, the gate electric field  $\lambda'$  will form local regions that have  $Z_2 = 0$ , isolating the  $Z_2 = 1$  regions along the less strongly disturbed portions of the MTI thin film. As discussed in the Supplementary material, we have explicitly verified that MZMs appear near the ends of quasi-1D topological superconductors that are written onto the MTI surface in this way.

MZMs localized at quantum wire ends are likely to prove more amenable to electrical control than MZMs localized at the vortices of a bulk 2D effective pwave superconductor. [15] With this motivation, we have demonstrated the possibility of using gate arrays to write MZMs onto the surface of a 2D MTI placed on a superconducting substrate. This strategy for generating MZM resources for quantum information processing has obvious advantages compared to arrays of separately grown one-dimensional quantum wires, which may be difficult to align. It has some similarity with systems[30] in which a gate array writes quantum wires onto a quantum well by periodically depleting all carriers or varying the number of locally occupied subbands between even and odd values but has advantages in this case as well because i: it is not necessary to apply a magnetic field to break degeneracies at time-reversal invariant points in one-dimensional momentum space and because ii: there can be a large energy separation between the quasi-1D bands formed by quantum Hall edge states and higher energy subbands, providing a large target for efforts to tune to  $Z_2 = 1$  superconductors.

This work was supported by the Army Research Office (ARO) under contract W911NF-15-1-0561:P00001, and by the Welch Foundation under grant TBF1473.

- D. A. Ivanov, Physical Review Letters 86, 268 (2001), pRL.
- [2] C. Nayak, S. H. Simon, A. Stern, M. Freedman, and S. Das Sarma, Reviews of Modern Physics 80, 1083 (2008), rMP.
- [3] A. Y. Kitaev, Annals of Physics **303**, 2 (2003).
- [4] A. Kitaev, Annals of Physics **321**, 2 (2006).
- [5] S. D. Sarma, M. Freedman, and C. Nayak, 1, 15001 (2015).
- [6] D. Aasen, M. Hell, R. V. Mishmash, A. Higginbotham, J. Danon, M. Leijnse, T. S. Jespersen, J. A. Folk, C. M. Marcus, K. Flensberg, and J. Alicea, Physical Review X 6, 031016 (2016), pRX.
- [7] T. Karzig, C. Knapp, R. M. Lutchyn, P. Bonderson, M. B. Hastings, C. Nayak, J. Alicea, K. Flensberg, S. Plugge, Y. Oreg, C. M. Marcus, and M. H. Freedman, Physical Review B 95, 235305 (2017), pRB.
- [8] G. Moore and N. Read, Nuclear Physics B 360, 362

(1991).

- [9] H.-H. Sun and J.-F. Jia, npj Quantum Materials 2, 34 (2017).
- [10] A. Y. Kitaev, Physics-Uspekhi 44, 131 (2001).
- [11] A. Jason, Reports on Progress in Physics **75**, 076501 (2012).
- [12] R. M. Lutchyn, J. D. Sau, and S. Das Sarma, Physical Review Letters 105, 077001 (2010), pRL.
- [13] J. D. Sau, R. M. Lutchyn, S. Tewari, and S. Das Sarma, Physical Review Letters 104, 040502 (2010), pRL.
- [14] C. Beenakker, Annual Review of Condensed Matter Physics 4, 113 (2013).
- [15] L. Fu and C. L. Kane, Physical Review Letters 100, 096407 (2008), pRL.
- [16] V. Mourik, K. Zuo, S. M. Frolov, S. R. Plissard, E. P. A. M. Bakkers, and L. P. Kouwenhoven, Science **336**, 1003 (2012).
- [17] M. T. Deng, C. L. Yu, G. Y. Huang, M. Larsson, P. Caroff, and H. Q. Xu, Nano Letters **12**, 6414 (2012).
- [18] A. Das, Y. Ronen, Y. Most, Y. Oreg, M. Heiblum, and H. Shtrikman, Nat Phys 8, 887 (2012), 10.1038/nphys2479.
- [19] M. T. Deng, S. Vaitieknas, E. B. Hansen, J. Danon, M. Leijnse, K. Flensberg, J. Nygrd, P. Krogstrup, and C. M. Marcus, Science **354**, 1557 (2016).
- [20] L. P. Rokhinson, X. Liu, and J. K. Furdyna, Nat Phys 8, 795 (2012), 10.1038/nphys2429.
- [21] S. M. Albrecht, A. P. Higginbotham, M. Madsen, F. Kuemmeth, T. S. Jespersen, J. Nygrd, P. Krogstrup,

and C. M. Marcus, Nature 531, 206 (2016).

- [22] S. Nadj-Perge, I. K. Drozdov, J. Li, H. Chen, S. Jeon, J. Seo, A. H. MacDonald, B. A. Bernevig, and A. Yazdani, Science **346**, 602 (2014).
- [23] R. Yu, W. Zhang, H.-J. Zhang, S.-C. Zhang, X. Dai, and Z. Fang, Science **329**, 61 (2010).
- [24] C.-Z. Chang, J. Zhang, X. Feng, J. Shen, Z. Zhang, M. Guo, K. Li, Y. Ou, P. Wei, L.-L. Wang, Z.-Q. Ji, Y. Feng, S. Ji, X. Chen, J. Jia, X. Dai, Z. Fang, S.-C. Zhang, K. He, Y. Wang, L. Lu, X.-C. Ma, and Q.-K. Xue, Science **340**, 167 (2013).
- [25] Q. L. He, L. Pan, A. L. Stern, E. C. Burks, X. Che, G. Yin, J. Wang, B. Lian, Q. Zhou, E. S. Choi, K. Murata, X. Kou, Z. Chen, T. Nie, Q. Shao, Y. Fan, S.-C. Zhang, K. Liu, J. Xia, and K. L. Wang, Science **357**, 294 (2017).
- [26] D. J. J. Marchand and M. Franz, Physical Review B 86, 155146 (2012), pRB.
- [27] J. Wang, Q. Zhou, B. Lian, and S.-C. Zhang, Physical Review B 92, 064520 (2015).
- [28] A. C. Potter and P. A. Lee, Physical Review Letters 105, 227003 (2010), pRL.
- [29] S. Tewari and J. D. Sau, Physical Review Letters 109, 150408 (2012), pRL.
- [30] F. Pientka, A. Keselman, E. Berg, A. Yacoby, A. Stern, and B. I. Halperin, Physical Review X 7, 021032 (2017), pRX.