

Global stability of the multi-strain Kermack-McKendrick (renewal) epidemic model

Michael T. Meehan^a, Daniel G. Cocks^b, Emma S. McBryde^a

^a*Australian Institute of Tropical Health and Medicine, James Cook University, Townsville, Australia*

^b*Research School of Science and Engineering, Australian National University, Canberra, Australia*

Abstract

We extend a recent investigation by Meehan et al. (2019) [20] regarding the global stability properties of the general Kermack-McKendrick (renewal) model to the multi-strain case. We demonstrate that the basic reproduction number of each strain R_{0j} represents a sharp threshold parameter such that when $R_{0j} \leq 1$ for all j each strain dies out and the infection-free equilibrium is globally asymptotically stable, whereas for $R_{01} \equiv \max_j R_{0j} > 1$ the endemic equilibrium point \bar{P}^1 , at which only the fittest strain (i.e. strain 1) remains in circulation, becomes globally asymptotically stable.

Keywords: multi-strain global stability Lyapunov competitive exclusion

1. Introduction

The combination of genetic evolution and primary transmission has driven an explosion in the number of phenotypically distinct lineages of infectious diseases (i.e. strains) circulating in the global population. To simulate the dynamics of several co-circulating pathogen strains, several authors have developed multi-strain extensions of canonical single-strain epidemic models [3, 5, 8, 4, 17, 1, 6]. Often, these models are constructed by making n copies of the various infectious states considered for a single strain, with the additional constraint that each of the n strains dips from a common susceptible pool. In this case, many authors (e.g. [3, 5, 8]) have rediscovered the well-known competitive exclusion principle (first appearing in the ecological literature [25, 15]) which asserts that when several species are competing over a shared resource only one of them can survive indefinitely — namely, the one with the greatest reproductive capacity. Although this result can often be deduced by investigating the asymptotic dynamics of each system, the global stability of the various equilibria has often proven to be more difficult to establish (see for instance [8]).

Many of the multi-strain models investigated thus far have been of compartmental type for which infected individuals have a fixed infectiousness for the duration of their infectious period (which may or may not follow an exponentially distributed latency period). In this article we extend these approaches by adapting the general Kermack-McKendrick renewal model to the multi-strain context (section 2) and apply the direct Lyapunov method to establish the necessary and sufficient conditions for the global asymptotic stability of the infection-free and endemic equilibrium points of our system (section 3). Importantly, since the general Kermack-McKendrick model incorporates many of the familiar transmission dynamic models as limiting cases (e.g. the SIR and SEIR models) [9, 21, 7], the analysis and results presented in this article generalize a number of results derived in previous investigations [6]. (For applications of the direct Lyapunov method to single-strain compartmental epidemic models see e.g. [14, 16, 12] and for renewal-type models see e.g. [18, 19].)

2. Model description

The model we investigate in this article is a multi-strain extension of the general Kermack-McKendrick model outlined in detail in [20] (see also [13, 18, 7]). In particular, we develop a model with n distinct,

uncoupled pathogen strains which each dip from a common susceptible pool, S . We assume perfect cross-immunity such that, once infected with strain $j \in [1, n]$, individuals are immune to further infection with an alternate strain. In the absence of cross-immunity it is possible that chaotic solutions may arise [2, 23].

Firstly, we introduce the force of infection of each strain j , $F_j(t)$, which, by definition, is the per-capita rate at which susceptibles are infected with strain j at time t . It follows then, that the incidence of strain j at time t , which we denote $v_j(t)$, is given by

$$v_j(t) = F_j(t)S(t)$$

where $S(t)$ denotes the number of susceptibles. Here, we assume that the force of infection depends linearly on the size of each infectious population, such that $F_j(t)$ can be expressed in terms of a renewal equation:

$$F_j(t) = \int_0^{\bar{\tau}_j} A_j(\tau)F_j(t-\tau)S(t-\tau) d\tau. \quad (1)$$

The kernel $A_j(\tau)$ gives the expected contribution to the force of infection for an individual who has been infected with strain j for τ units of time and $\bar{\tau}_j$ is the maximum infection-age at which an individual infected with strain j contributes to the force of infection $F_j(t)$:

$$\bar{\tau}_j = \sup \{ \tau \geq 0 : A_j(\tau) > 0 \} < \infty.$$

Note that in our analysis we assume $\bar{\tau}_j < \infty$ for all j because the lack of compactness in the infinite case makes the problem much more difficult [11, 10].

We also define

$$\bar{\tau} = \max_j \bar{\tau}_j.$$

Next, we assume that individuals are recruited (i.e. born) directly into the susceptible class at a constant rate λ and that all individuals experience a constant per-capita natural death rate, μ . Therefore, if we combine the demographic influences with the loss of susceptible individuals due to all types of infection, we find that the susceptible population varies according to

$$\frac{dS(t)}{dt} = \lambda - \mu S(t) - \sum_{j=1}^n F_j(t)S(t). \quad (2)$$

Finally, if we integrate the expected contribution to the force of infection $A_j(\tau)$ over all possible infection ages we find that the basic reproduction number for each strain, R_{0j} , is given by

$$R_{0j} = S^0 \int_0^{\bar{\tau}_j} A_j(\tau) d\tau \quad (3)$$

where $S^0 = \lambda/\mu$ is the steady-state susceptible population in the absence of infection. In the analysis that follows, we assume that

$$R_{0i} \neq R_{0j} \quad \text{for } i \neq j,$$

and, without loss of generality, we label the strain with the maximum reproduction number strain 1, such that

$$R_{01} \equiv \max_j R_{0j}.$$

Supplementing our model with appropriate initial conditions, $\mathcal{S}_0 \in C_+^0([-\bar{\tau}, 0])$ and $\mathcal{F}_{j,0} \in L_+^1(-\bar{\tau}_j, 0)$, where \mathcal{S}_0 and $\mathcal{F}_{j,0}$ are the initial histories of the susceptible population and the force of infection of each strain j respectively, we see that the phase-space of our system is given by the product topology

$$\Omega = C_+^0([-\bar{\tau}, 0]) \times \prod_{j=1}^n L_+^1(-\bar{\tau}_j, 0)$$

which is a Banach space with the natural norm

$$\|(\mathcal{S}, \mathcal{F}_1, \mathcal{F}_2, \dots, \mathcal{F}_n)\| = \sup_{s \in [-\bar{\tau}, 0]} |\mathcal{S}(s)| + \sum_{j=1}^n \int_{-\bar{\tau}_j}^0 |\mathcal{F}_j(s)| ds.$$

With this choice of state space standard arguments show that the model (1)-(2) is well defined and induces a continuous semi-flow $\Phi_t : \Omega \rightarrow \Omega$. Importantly, by Lemma 1 of [20], which invokes the smoothing properties of convolution integrals detailed in [22], it is straightforward to show that when the infectivity kernels A_j are of bounded variation, i.e. $A_j \in BV([0, \bar{\tau}_j])$, system trajectories generated by the continuous semiflow Φ_t originating in Ω enter a bounded subset $\Omega^c \subset \Omega$ that is relatively compact. In this case, which we shall assume holds forthwith, the ω -limit set of the system (1)-(2) is non-empty such that the infinite-dimensional form of LaSalle's extension to Lyapunov's global asymptotic stability theorem [24, Theorem 5.17] can be applied.

As a useful shorthand, we introduce the notation $\mathcal{F} = (\mathcal{F}_1, \mathcal{F}_2, \dots)$ to denote the set of force of infection states and observe that the system trajectory $(\mathcal{S}_t(\cdot), \mathcal{F}_t(\cdot)) \in \Omega$ with

$$\mathcal{S}_t(s) = S(t+s), \quad \mathcal{F}_t(s) = (\mathcal{F}_{1,t}(s), \mathcal{F}_{2,t}(s), \dots) = (F_1(t+s), F_2(t+s), \dots), \quad s \in [-\bar{\tau}, 0].$$

Next, we discuss the equilibrium states of the system. We have already observed from equations (1)-(2) that the trivial infection-free equilibrium solution P^0 is given by $\mathcal{F}^0 = 0$ and $\mathcal{S}^0 = S^0 = \lambda/\mu$, where, in general, P^0 belongs to the collection of infection-free states $\partial\Omega$:

$$P^0 \in \partial\Omega = \{(\mathcal{S}, \mathcal{F}) \in \Omega \mid \mathcal{F} = 0\}.$$

Here we use the notation $\partial\Omega$ not in the strict topological sense, but rather in view of our application: $\partial\Omega$ is the set of states for which the force of infection is vanishing and which therefore lead to trivial dynamics.

To determine the remaining (endemic) solutions, $\bar{P}^j = (\bar{\mathcal{S}}, \bar{\mathcal{F}}^j)$, we solve equation (1) to obtain

$$\mathcal{K}[\bar{\mathcal{F}}] = \frac{\bar{\mathcal{S}}}{S^0} \bar{\mathcal{F}}$$

where we introduce the next-generation operator \mathcal{K} , defined as

$$\mathcal{K} = \text{diag} \left(S^0 \int_0^{\bar{\tau}_j} A_j(\tau) d\tau \right) = \text{diag} (R_{0j}).$$

Hence, the endemic solutions, \bar{P}^j , are determined by the spectrum of the (diagonal) operator \mathcal{K} :

$$\bar{S}^j = \frac{S^0}{R_{0j}} \quad \text{and} \quad \bar{F}_i^j = \begin{cases} \mu(R_{0j} - 1), & i = j \\ 0, & \text{otherwise} \end{cases} \quad (4)$$

where the $\bar{\mathcal{F}}^j$ have been calculated by substituting the solution for \bar{S}^j into (2) and rearranging.

From (4), we see that at each of the endemic equilibrium points \bar{P}^j only strain j survives, i.e. we have competitive exclusion. Moreover, $\bar{P}^j \in \Omega$ if, and only if $R_{0j} \geq 1$.

In the following, we would like to consider the trajectories of the system (1)-(2) originating from a space for which all strains $j \in [1, n]$ are active — that is, the set of states, $\hat{\Omega}$, for which all strains have a non-zero force of infection at some point over the interval $[0, \bar{\tau}]$. More formally, if we define the set of states for which a subset $\mathcal{A} \subseteq \{1, \dots, n\}$ of strains are active as

$$\hat{\Omega}^{\mathcal{A}} = \left\{ (\mathcal{S}, \mathcal{F}) \in \Omega \mid \int_0^{\bar{\tau}_j} \int_0^{\bar{\tau}_j} A_j(\tau+a) \mathcal{F}_j(-\tau) \mathcal{S}(-\tau) d\tau da > 0 \iff j \in \mathcal{A} \right\},$$

we then have

$$\hat{\Omega} = \hat{\Omega}^{\{1, \dots, n\}}.$$

Moreover, we see that each of the endemic equilibrium points $\bar{P}^j \in \widehat{\Omega}^j$ whilst, conversely, $\partial\Omega \subset \widehat{\Omega}^\emptyset$.

Given the parallel structure of the strains in our model, it is clear that once a strain becomes inactive (i.e. $j \notin \mathcal{A}$), it remains so thereafter. Consequently, the number of active strains can only diminish as the system evolves: $\Phi_{t>0}(\widehat{\Omega}^{\mathcal{A}}) \subseteq \widehat{\Omega}^{\mathcal{A}}$. Indeed, as we will show below, in the case of endemic infection, the system evolves towards a state in which only the fittest strain (strain 1) survives, i.e. $\Phi_{t \rightarrow \infty}(\widehat{\Omega}^{\{1, \dots, n\}}) \subseteq \widehat{\Omega}^1$.

3. Global stability analysis

We now establish the stability properties of the equilibrium solutions of the system (1)-(2), for which we recall the definition $R_{01} \equiv \max_j R_{0j}$ to identify strain 1.

3.1. Infection-free equilibrium

Theorem 1. *The infection-free equilibrium point P^0 is globally asymptotically stable in Ω if $R_{01} \leq 1$. However, if $R_{01} > 1$, solutions of (1)-(2) starting sufficiently close to P^0 in $\widehat{\Omega} = \widehat{\Omega}^{\{1, \dots, n\}}$ move away from P^0 .¹ In all cases, solutions starting in $\partial\Omega \subset \widehat{\Omega}^\emptyset$ approach P^0 .*

Proof. Recall that $\bar{\tau} = \max_j \bar{\tau}_j$ and consider the forward-invariant and attracting region $D = \Phi_{\bar{\tau}}(\Omega)$. From (2) we observe $\mathcal{S}(0) > 0$ for all $(\mathcal{S}, \mathcal{F}) \in D$. Next we define $U : D \rightarrow \mathbb{R}_+$ (c.f. [20]):

$$U(\mathcal{S}, \mathcal{F}) = g\left(\frac{\mathcal{S}(0)}{S^0}\right) + \sum_{j=1}^n \int_0^{\bar{\tau}_j} \eta_j(\tau) \mathcal{F}_j(-\tau) \mathcal{S}(-\tau) d\tau \quad (5)$$

where

$$g(x) = x - 1 - \log x \quad \text{and} \quad \eta_j(\tau) = \int_\tau^{\bar{\tau}_j} A_j(s) ds. \quad (6)$$

In particular we have $\eta_j(\bar{\tau}_j) = 0$,

$$\eta_j(0) = \frac{R_{0j}}{S^0} \quad \text{and} \quad \eta'_j(\tau) = -A_j(\tau) \quad (7)$$

where a $'$ denotes differentiation with respect to τ . Note, the functional U is positive, continuous and well defined in D , and has a global minimum in Ω at P^0 .

Using $\mathcal{S}_t(s) = \mathcal{S}(t+s)$ and $\mathcal{F}_{j,t}(s) = F_j(t+s)$ and rewriting the integral in (5) we find that U , evaluated along system trajectories, is given by

$$\begin{aligned} U(\mathcal{S}_t(\cdot), \mathcal{F}_t(\cdot)) &= g\left(\frac{\mathcal{S}_t(0)}{S^0}\right) + \sum_{j=1}^n \int_0^{\bar{\tau}_j} \eta_j(\tau) \mathcal{F}_{j,t}(-\tau) \mathcal{S}_t(-\tau) d\tau, \\ &= g\left(\frac{S(t)}{S^0}\right) + \sum_{j=1}^n \int_{t-\bar{\tau}_j}^t \eta_j(t-s) F_j(s) S(s) ds. \end{aligned} \quad (8)$$

Taking the time derivative of each term in (8) separately, and substituting in the model equations (1) and (2),

¹The extension to systems for which only a subset of strains are active initially, i.e. $\widehat{\Omega}^{\mathcal{A}}$ where $\mathcal{A} \subseteq \{1, \dots, n\}$, is trivial, provided R_{01} is redefined appropriately: $R_{01} = \max_{j \in \mathcal{A}} R_{0j}$.

we get

$$\begin{aligned}
\frac{d}{dt} g\left(\frac{S(t)}{S^0}\right) &= \left(\frac{1}{S^0} - \frac{1}{S(t)}\right) \frac{dS(t)}{dt}, \\
&= \frac{\lambda}{S^0} - \mu \frac{S(t)}{S^0} - \sum_{j=1}^n F_j(t) \frac{S(t)}{S^0} - \frac{\lambda}{S(t)} + \mu + \sum_{j=1}^n F_j(t), \\
&= \mu \left(2 - \frac{S(t)}{S^0} - \frac{S^0}{S(t)}\right) - \sum_{j=1}^n F_j(t) \left(\frac{S(t)}{S^0} - 1\right), \\
&= -\mu \frac{S(t)}{S^0} \left(1 - \frac{S^0}{S(t)}\right)^2 - \sum_{j=1}^n F_j(t) \left(\frac{S(t)}{S^0} - 1\right)
\end{aligned}$$

where in the second line we have substituted in the identity $\lambda = \mu S^0$, and

$$\begin{aligned}
\frac{d}{dt} \left[\sum_{j=1}^n \int_{t-\bar{\tau}_j}^t \eta_j(t-s) F_j(s) S(s) ds \right] &= \sum_{j=1}^n \left[\eta_j(0) F_j(t) S(t) - \eta_j(\bar{\tau}_j) F_j(t - \bar{\tau}_j) S(t - \bar{\tau}_j) \right. \\
&\quad \left. + \int_{t-\bar{\tau}_j}^t \eta_j'(t-s) F_j(s) S(s) ds \right], \\
&= \sum_{j=1}^n \left[R_{0j} F_j(t) \frac{S(t)}{S^0} - \int_{t-\bar{\tau}_j}^t A_j(t-s) F_j(s) S(s) ds \right], \\
&= \sum_{j=1}^n F_j(t) \left(R_{0j} \frac{S(t)}{S^0} - 1 \right).
\end{aligned}$$

Combining these results we then have

$$\frac{d}{dt} U(\mathcal{S}_t, \mathcal{F}_t) = -\mu \frac{S_t(0)}{S^0} \left(1 - \frac{S^0}{S_t(0)}\right)^2 - \sum_{j=1}^n (1 - R_{0j}) \mathcal{F}_{j,t}(0) \frac{S_t(0)}{S^0} \leq 0. \quad (9)$$

For system trajectories $(\mathcal{S}_t, \mathcal{F}_t) \in D \subset \Omega$ equations (1) and (2) imply that $\mathcal{F}_{j,t} \in C^0([-\bar{\tau}_j, 0])$ for $t > \bar{\tau}$ (see [20]) such that (9) is well defined and U is a proper Lyapunov functional on the domain D .

We observe that the derivative $\dot{U} = 0$ if and only if $S_t(0) = S^0$ and either (a) $R_{0j} = 1$ or (b) $\mathcal{F}_{j,t}(0) = 0$ for all j . Therefore, the largest invariant subset in Ω for which $\dot{U} = 0$ is the singleton $\{P^0\}$. Since by Lemma 1 of [20] the orbit is eventually precompact, by the infinite-dimensional form of LaSalle's extension of Lyapunov's global asymptotic stability theorem [24, Theorem 5.17], the infection-free equilibrium point P^0 is globally asymptotically stable in Ω if $R_{01} \leq 1$.

Conversely, if $R_{0j} > 1$ for any j , the derivative $\dot{U} > 0$ for $S_t(0)$ sufficiently close to S^0 , provided $\mathcal{F}_{j,t}(0) > 0$. Therefore, solutions starting sufficiently close to the infection-free equilibrium point P^0 in $\widehat{\Omega}^j$ leave a neighbourhood of P^0 . Since $\dot{U} \leq 0$ for solutions starting in $\partial\Omega$ these solutions approach P^0 . \square

3.2. Endemic equilibrium

For convenience, in this section we adopt the shorthand notation that an overbar refers to the value of a state variable at the endemic equilibrium point \bar{P}^1 such that, for example, $\bar{S} \equiv S^1$ and $\bar{\mathcal{F}} \equiv \mathcal{F}^1$.

Theorem 2. *If $R_{01} > 1$ the endemic equilibrium point \bar{P}^1 is globally asymptotically stable in $\widehat{\Omega} = \widehat{\Omega}^{\{1, \dots, n\}}$.²*

²As before, the extension to systems for which only a subset of strains are active initially, i.e. $\widehat{\Omega}^{\mathcal{A}}$ where $\mathcal{A} \subseteq \{1, \dots, n\}$, is trivial, provided R_{01} is redefined appropriately: $R_{01} = \max_{j \in \mathcal{A}} R_{0j}$.

Proof. From theorem 1 we have that $F_1(t)$ is bounded away from zero for $t > 0$ when $R_{01} > 1$, such that $\Phi_t : \widehat{\Omega} \rightarrow \widehat{\Omega}^A \supseteq \widehat{\Omega}^1$.³ Recall that $\bar{\tau} = \max_j \bar{\tau}_j$ and define $\widehat{D} = \Phi_{\bar{\tau}}(\widehat{\Omega})$ which is forward invariant and attracting for $R_{01} > 1$. Moreover, $\mathcal{S}, \mathcal{F}_1 > 0$ for $(\mathcal{S}, \mathcal{F}) \in \widehat{D}$. Consider the Lyapunov functional $W : \widehat{D} \rightarrow \mathbb{R}_+$ (c.f. [20]):

$$W(\mathcal{S}, \mathcal{F}) = g\left(\frac{\mathcal{S}(0)}{\mathcal{S}}\right) + \bar{F}_1 \bar{S} \int_0^{\bar{\tau}_1} \chi_1(\tau) g\left(\frac{\mathcal{F}_1(-\tau) \mathcal{S}(-\tau)}{\bar{F}_1 \bar{S}}\right) d\tau + \sum_{j=2}^n \int_0^{\bar{\tau}_j} \chi_j(\tau) \mathcal{F}_j(-\tau) \mathcal{S}(-\tau) d\tau \quad (10)$$

where $g(x)$ has been defined previously in (6) and

$$\chi_j(\tau) = \int_{\tau}^{\bar{\tau}_j} A_j(s) ds.$$

In particular, we have

$$\chi_j(0) = \frac{R_{0j}}{S^0} = \frac{R_{0j}}{R_{01}} \frac{1}{\bar{S}} \quad \text{and} \quad \chi_j'(\tau) = -A_j(\tau) \quad (11)$$

so that $\chi_1(0) = 1/\bar{S}$ and $\chi_j(\bar{\tau}_j) = 0$. Note the functional W is positive, continuous, and has a global minimum in $\widehat{\Omega}$ at \bar{P}^1 .

Rewriting the integral terms as in theorem 1 and evaluating W along system trajectories gives

$$W(\mathcal{S}_t, \mathcal{F}_t) = g\left(\frac{\mathcal{S}(t)}{\bar{S}}\right) + \bar{F}_1 \bar{S} \int_{t-\bar{\tau}_1}^t \chi_1(t-s) g\left(\frac{\mathcal{F}_1(s) \mathcal{S}(s)}{\bar{F}_1 \bar{S}}\right) ds + \sum_{j=2}^n \int_{t-\bar{\tau}_j}^t \chi_j(t-s) \mathcal{F}_j(s) \mathcal{S}(s) ds. \quad (12)$$

Next we differentiate the first term in (12) to get

$$\begin{aligned} \frac{d}{dt} g\left(\frac{\mathcal{S}(t)}{\bar{S}}\right) &= \left(\frac{1}{\bar{S}} - \frac{1}{\mathcal{S}(t)}\right) \frac{d\mathcal{S}(t)}{dt}, \\ &= \frac{\lambda}{\bar{S}} - \mu \frac{\mathcal{S}(t)}{\bar{S}} - \sum_{j=1}^n F_j(t) \frac{\mathcal{S}(t)}{\bar{S}} - \frac{\lambda}{\mathcal{S}(t)} + \mu + \sum_{j=1}^n F_j(t), \\ &= \mu \left(2 - \frac{\mathcal{S}(t)}{\bar{S}} - \frac{\bar{S}}{\mathcal{S}(t)}\right) + \sum_{j=1}^n \bar{F}_j \left(1 - \frac{\bar{S}}{\mathcal{S}(t)}\right) + \sum_{j=1}^n F_j(t) \left(1 - \frac{\mathcal{S}(t)}{\bar{S}}\right), \\ &= -\mu \frac{\mathcal{S}(t)}{\bar{S}} \left(1 - \frac{\bar{S}}{\mathcal{S}(t)}\right)^2 + \bar{F}_1 \left(1 - \frac{\bar{S}}{\mathcal{S}(t)}\right) + \sum_{j=1}^n F_j(t) \left(1 - \frac{\mathcal{S}(t)}{\bar{S}}\right), \end{aligned} \quad (13)$$

where in the second line we have used the identity $\lambda = \mu \bar{S} + \sum_{j=1}^n \bar{F}_j \bar{S} = \mu \bar{S} + \bar{F}_1 \bar{S}$. Similarly, we differentiate the final term in (12) and substitute in the definition of $\chi_j(\tau)$ to get

$$\begin{aligned} \frac{d}{dt} \left[\sum_{j=2}^n \int_{t-\bar{\tau}_j}^t \chi_j(t-s) \mathcal{F}_j(s) \mathcal{S}(s) ds \right] &= \sum_{j=2}^n \left[\chi_j(0) \mathcal{F}_j(t) \mathcal{S}(t) - \chi_j(\bar{\tau}_j) \mathcal{F}_j(t-\bar{\tau}_j) \mathcal{S}(t-\bar{\tau}_j) \right. \\ &\quad \left. + \int_{t-\bar{\tau}_j}^t \chi_j'(t-s) \mathcal{F}_j(s) \mathcal{S}(s) ds \right], \\ &= \sum_{j=2}^n \left[\mathcal{F}_j(t) \frac{R_{0j}}{R_{01}} \frac{\mathcal{S}(t)}{\bar{S}} - \int_{t-\bar{\tau}_j}^t A_j(t-s) \mathcal{F}_j(s) \mathcal{S}(s) ds \right], \\ &= \sum_{j=2}^n \mathcal{F}_j(t) \left(\frac{R_{0j}}{R_{01}} \frac{\mathcal{S}(t)}{\bar{S}} - 1 \right). \end{aligned} \quad (14)$$

³Although $F_1(t)$ is not uniformly bounded away from zero, it is sufficiently so to ensure that our Lyapunov function W remains well defined.

Differentiating the second term in (12) and using (11), yields

$$\begin{aligned}
\frac{d}{dt} \left[\bar{F}_1 \bar{S} \int_{t-\bar{\tau}_1}^t \chi_1(t-s) g \left(\frac{F_1(s)S(s)}{\bar{F}_1 \bar{S}} \right) ds \right] &= \bar{F}_1 \bar{S} \left[\chi_1(0) g \left(\frac{F_1(t)S(t)}{\bar{F}_1 \bar{S}} \right) - \chi_1(\bar{\tau}_1) g \left(\frac{F_1(t-\bar{\tau}_1)S(t-\bar{\tau}_1)}{\bar{F}_1 \bar{S}} \right) \right. \\
&\quad \left. + \int_{t-\bar{\tau}_1}^t \chi_1'(t-s) g \left(\frac{F_1(s)S(s)}{\bar{F}_1 \bar{S}} \right) ds \right], \\
&= \bar{F}_1 g \left(\frac{F_1(t)S(t)}{\bar{F}_1 \bar{S}} \right) - \bar{F}_1 \bar{S} \int_{t-\bar{\tau}_1}^t A_1(t-s) g \left(\frac{F_1(s)S(s)}{\bar{F}_1 \bar{S}} \right) ds, \\
&= F_1(t) \left(\frac{S(t)}{\bar{S}} - 1 \right) - \bar{F}_1 \log \left(\frac{F_1(t)S(t)}{\bar{F}_1 \bar{S}} \right) \\
&\quad + \bar{F}_1 \bar{S} \int_{t-\bar{\tau}_1}^t A_1(t-s) \log \left(\frac{F_1(s)S(s)}{\bar{F}_1 \bar{S}} \right) ds
\end{aligned}$$

where in the last line we have invoked the identity $\bar{S} \int_0^{\bar{\tau}_1} A_1(\tau) d\tau = 1$. As in [20], we can bound this expression using Jensen's inequality:⁴

$$\begin{aligned}
\frac{d}{dt} \left[\bar{F}_1 \bar{S} \int_{t-\bar{\tau}_1}^t \chi_1(t-s) g \left(\frac{F_1(s)S(s)}{\bar{F}_1 \bar{S}} \right) ds \right] &\leq F_1(t) \left(\frac{S(t)}{\bar{S}} - 1 \right) - \bar{F}_1 \log \left(\frac{F_1(t)S(t)}{\bar{F}_1 \bar{S}} \right) \\
&\quad + \bar{F}_1 \log \left[\bar{S} \int_{t-\bar{\tau}_1}^t A_1(t-s) \frac{F_1(s)S(s)}{\bar{F}_1 \bar{S}} ds \right], \\
&\leq F_1(t) \left(\frac{S(t)}{\bar{S}} - 1 \right) - \bar{F}_1 \log \left(\frac{F_1(t)S(t)}{\bar{F}_1 \bar{S}} \right) + \bar{F}_1 \log \left(\frac{F_1(t)}{\bar{F}_1} \right), \\
&\leq F_1(t) \left(\frac{S(t)}{\bar{S}} - 1 \right) - \bar{F}_1 \log \left(\frac{S(t)}{\bar{S}} \right), \\
&\leq F_1(t) \left(\frac{S(t)}{\bar{S}} - 1 \right) - \bar{F}_1 \left(1 - \frac{\bar{S}}{S(t)} \right)
\end{aligned} \tag{15}$$

where in the last line we have also used $\log x \geq 1 - \frac{1}{x}$.

Finally, recalling that $R_{01} = \max_j R_{0j}$, and combining (13), (14) and (15) yields

$$\frac{d}{dt} W(\mathcal{S}_t, \mathcal{F}_t) \leq -\mu \frac{\mathcal{S}_t(0)}{\bar{S}} \left(1 - \frac{\bar{S}}{\mathcal{S}_t(0)} \right)^2 - \sum_{j=2}^n \left(1 - \frac{R_{0j}}{R_{01}} \right) \mathcal{F}_{j,t}(0) \frac{\mathcal{S}_t(0)}{\bar{S}} \leq 0. \tag{16}$$

From equation (16) we see that the largest invariant subset in $\widehat{\Omega}$ for which $\dot{W} = 0$ is the endemic equilibrium point \bar{P}^1 . Since by Lemma 1 of [20] the orbit is eventually precompact, by LaSalle's extension of Lyapunov's asymptotic stability theorem [24, Theorem 5.17], the endemic equilibrium point \bar{P}^1 is globally asymptotically stable in $\widehat{\Omega}$ for $R_{01} > 1$. \square

4. Conclusions

In this article we investigated the global stability properties of the multi-strain Kermack-McKendrick model. We found that when the basic reproduction number $R_{0j} \leq 1$ for all strains j the infection-free equilibrium P^0 is unique in Ω and is globally asymptotically stable. We also discovered a set of n endemic

⁴For a concave function $\varphi(\cdot)$, and probability distribution $h(t)$, the following inequality holds:

$$\varphi \left(\int_0^\infty h(t) f(t) dt \right) \geq \int_0^\infty h(t) \varphi(f(t)) dt.$$

equilibrium solutions, $\{\bar{P}^j\}$, at which only strain j survives with a positive infected population, à la competitive exclusion. Moreover, we found that \bar{P}^j only exists in the positive cone $\widehat{\Omega}$ if $R_{0j} > 1$. Our main result, which was derived using the direct Lyapunov method, was to show that of this set, \bar{P}^1 — at which the fittest strain, defined by $R_{01} \equiv \max_j R_{0j}$, survives indefinitely — is globally asymptotically stable when it exists.

Acknowledgements

The authors would like to express their sincere gratitude to Prof. Johannes Müller for his invaluable advice and feedback in the preparation of this manuscript.

References

- [1] S. Ackleh and J.S. Allen. Competitive exclusion and coexistence for pathogens in an epidemic model with variable population size. *Journal of Mathematical Biology*, 47(2):153–168, 2003. ISSN 1432-1416. doi: 10.1007/s00285-003-0207-9.
- [2] M. Aguiar, B. Kooi, and N. Stollenwerk. Epidemiology of dengue fever: a model with temporary cross-immunity and possible secondary infection shows bifurcation and chaotic behaviour in wide parameter regions. *Math. Model. Nat. Pheno.*, 3(4):48–70, January 2008.
- [3] R.M. Anderson and R.M. May. Coevolution of hosts and parasites. *Parasitology*, 85(02):411–426, 1982.
- [4] V. Andreasen and A. Pugliese. Pathogen coexistence induced by density-dependent host mortality. *Journal of Theoretical Biology*, 177(2):159–166, 1995.
- [5] K. Beck. Coevolution: Mathematical analysis of host-parasite interactions. *Journal of Mathematical Biology*, 19(1):63–77, 1984. ISSN 1432-1416. doi: 10.1007/BF00275931.
- [6] D. Bichara, A. Iggidr, and G. Sallet. Global analysis of multi-strains SIS, SIR and MSIR epidemic models. *Journal of Applied Mathematics and Computing*, 44(1):273–292, 2013.
- [7] D. Breda, O. Diekmann, W.F. De Graaf, A. Pugliese, and R. Vermiglio. On the formulation of epidemic models (an appraisal of Kermack and McKendrick). *Journal of Biological Dynamics*, 6(sup2):103–117, 2012.
- [8] H.J. Bremermann and H.R. Thieme. A competitive exclusion principle for pathogen virulence. *Journal of Mathematical Biology*, 27(2):179–190, 1989. ISSN 1432-1416. doi: 10.1007/BF00276102.
- [9] O. Diekmann. Limiting behaviour in an epidemic model. *Nonlinear Analysis: Theory, Methods & Applications*, 1(5):459–470, 1977. ISSN 0362-546X.
- [10] O. Diekmann and M. Gyllenberg. Equations with infinite delay: Blending the abstract and the concrete. *Journal of Differential Equations*, 252(2):819–851, 2012. ISSN 0022-0396.
- [11] O. Diekmann, P. Getto, and M. Gyllenberg. Stability and bifurcation analysis of Volterra functional equations in the light of Suns and Stars. *SIAM Journal on Mathematical Analysis*, 39(4):1023–1069, 2008.
- [12] M. Fan, M.Y. Li, and K. Wang. Global stability of an SEIS epidemic model with recruitment and a varying total population size. *Mathematical Biosciences*, 170(2):199–208, 2001.
- [13] W.O. Kermack and A.G. McKendrick. A contribution to the mathematical theory of epidemics. *Proceedings of the Royal Society of London A: Mathematical, Physical and Engineering Sciences*, 115(772):700–721, 1927.
- [14] A. Korobeinikov and G.C. Wake. Lyapunov functions and global stability for SIR, SIRS, and SIS epidemiological models. *Applied Mathematics Letters*, 15(8):955–960, 2002.
- [15] S.A. Levin. Community equilibria and stability, and an extension of the competitive exclusion principle. *American Naturalist*, pages 413–423, 1970.
- [16] M.Y. Li and J.S. Muldowney. Global stability for the SEIR model in epidemiology. *Mathematical Biosciences*, 125(2):155–164, 1995.
- [17] M. Lipsitch and E.R. Moxon. Virulence and transmissibility of pathogens: what is the relationship? *Trends in Microbiology*, 5(1):31–37, 1997. ISSN 0966-842X. doi: http://dx.doi.org/10.1016/S0966-842X(97)81772-6.
- [18] P. Magal, C.C. McCluskey, and G.F. Webb. Lyapunov functional and global asymptotic stability for an infection-age model. *Applicable Analysis*, 89(7):1109–1140, 2010.
- [19] C.C. McCluskey. Complete global stability for an SIR epidemic model with delay: distributed or discrete. *Nonlinear Analysis: Real World Applications*, 11(1):55–59, 2010.
- [20] M.T. Meehan, D.G. Cocks, J. Müller, and E.S. McBryde. Global stability properties of a class of renewal epidemic models. *Journal of Mathematical Biology*, 78:1713–1725, 2019.
- [21] J. A. J. Metz and O. Diekmann. *The Dynamics of Physiologically Structured Populations*. Lecture Notes in Biomathematics 68. Springer-Verlag, Berlin, Heidelberg, New York, London, Paris, Tokyo, 1986.
- [22] J. Mikusiński and Cz. Ryll-Nardzewski. Sur le produit de composition. *Studia Mathematica*, 12:51–57, 1951.
- [23] P. Minayev and N. Ferguson. Improving the realism of deterministic multi-strain models: implications for modelling influenza a. *J. R. Soc. Interface*, 6(35):509–518, September 2008.
- [24] H. Smith. *An introduction to delay differential equations with applications to the life sciences*. Springer, 2010.
- [25] V. Volterra. Variations and fluctuations of the number of individuals in animal species living together. *J. Cons. Int. Explor. Mer*, 3(1):3–51, 1928.