HYPOELLIPTICITY WITHOUT LOSS OF DERIVATIVES FOR FEDII'S TYPE OPERATORS

TIMUR AKHUNOV, LYUDMILA KOROBENKO, AND CRISTIAN RIOS

ABSTRACT. We prove that second order linear operators on \mathbb{R}^{n+m} of the form $L(x, y, D_x, D_y) = L_1(x, D_x) + g(x)L_2(y, D_y)$, where L_1 and L_2 satisfy Morimoto's super-logarithmic estimates and g is smooth, nonnegative, and vanishes only at the origin in \mathbb{R}^n (but to any arbitrary order) are hypoelliptic without loss of derivarives. We also show examples in which our hypotheses are necessary for hypoellipticity.

1. INTRODUCTION

The regularity question, or whether singular solutions are possible, is central to the study of PDEs. More precisely, a differential operator $P = \sum_{\alpha} a_{\alpha}(x) \partial_x^{\alpha}$ is (locally) hypoelliptic, if

(1) u is smooth near x_0 , whenever Lu is smooth near x_0 .

Equivalently, the formal inverse operator L^{-1} preserves smoothness. The question of which operators are hypoelliptic goes back Laurent Schwartz in the 50s [Sch50, Sch63]. Lars Hörmander classified constant coefficient operators in [Hör55]. For variable coefficients the classification is incomplete even for the operators of the form

(2)
$$Lu = -\sum_{i,j=1}^{n} a_{ij}(x)\partial_{x_i}\partial_{x_j} + \sum_{i=1}^{n} a_i(x)\partial_{x_i} + a_0(x), \text{ for matrix } (a_{ij}) \ge 0$$

Such differential operators are called (degenerately) elliptic and generalize the Laplace operator $L = \Delta$. These operators are among the most important and intensely studied ones. The non-degenerate case of (2), or $0 < \lambda \leq (a_{ij}(x)) \leq \Lambda < \infty$ closely resembles the Laplacian Δ , and in particular every solution of Lu = 0 must be analytic. The degenerate case of (2) may fail to be hypoelliptic, i.e. there may exist non-smooth functions u such that Lu is smooth.

Our goal is to prove hypoellipticity for a wide class of linear degenerate elliptic operators. Starting with the famous Hörmander bracket condition [Hör67], there has

Date: November 12, 2021.

¹⁹⁹¹ Mathematics Subject Classification. 35H10, 35H20, 35S05, 35G05, 35B65, 35A18.

Key words and phrases. hypoellipticity, infinite vanishing, loss of derivatives.

The third author is supported by the Natural Sciences and Engineering Research Council of Canada.

been a lot of theory built trying to classify hypoelliptic operators. It is known that Hörmander's bracket condition is equivalent to the following estimate

(3)
$$||u||_{H^{\varepsilon}} := ||\langle \xi \rangle^{\varepsilon} \hat{u}(\xi)||_{L^{2}} \le (Lu, u) + C||u||_{L^{2}}$$

called subellipticity. Here $\hat{u}(\xi)$ denotes the Fourier transform of the function u(x), and $\langle \xi \rangle := (1 + |\xi|^2)^{1/2}$. It is known that subelliptic operators are hypoelliptic, however the latter class is much wider. For example, it includes operators of the form $\partial_x^2 + g(x)\partial_y^2$, where the function g(x) is allowed to vanish at a point together with all its derivatives. It has been shown by Fediĭ [Fed71] that such operators are hypoelliptic independently of the order of vanishing of g(x). Moreover, this result has been generalized by Kohn in [Koh98] (see also [KR14]) to include operators of the form

(4)
$$L(x, y, D_x, D_y) = L_1(x, D_x) + g(x)L_2(y, D_y)$$

where $x \in \mathbb{R}^m$, $y \in \mathbb{R}^n$, and the operators L_1 and L_2 are subelliptic in their variables. Again, the function g(x) was allowed to vanish at isolated points together with all its derivatives. Using Hörmander's bracket condition it is easy to check that the operator L defined by (4) is not subelliptic for infinitely vanishing g(x). In particular, estimate (3) does not hold for all test functions u. On the other hand in [Mor87] (see also [Chr01]) Morimoto established the following sufficient condition for the hypoellipticity of an operator L of the form (2), which we call the superlogarithmic estimate: for all $\varepsilon > 0$ and $K \Subset R^m$

(5)
$$||w(\xi)\hat{u}(\xi)||^2 \le \varepsilon \operatorname{Re}(Lu, u) + C_{\varepsilon, K}||u||^2, \quad u \in C_0^{\infty}(K),$$

where $w(\xi)$ is a function satisfying $w(\xi) \ge \log (1 + |\xi|^2)$. Note that the subelliptic estimate (3) can be thought of as (5) with $w(\xi) = \langle \xi \rangle^{\varepsilon}$. Thus, roughly speaking, more degenerate operators satisfy estimates of the form (5) with smaller weights $w(\xi)$. Inspired by these results we were interested in constructing the widest collection of linear operators of the form (4) that are hypoelliptic, i.e. $u \in C^{\infty}$ whenever $Lu \in C^{\infty}$ and L has smooth coefficients. It is known that simply requiring L_1 and L_2 to be hypoelliptic in their variables is not sufficient, examples can be found in [KS84] and [Mor87]; see also Example 1 below.

We consider two linear (degenerately) elliptic operators

(6)
$$L_1 = -\sum_{j,k=1}^n a_{jk}(x)\partial_{x_j}\partial_{x_k} + \sum_{j=1}^n a_j(x)\partial_{x_j} + a_0(x)$$

where $x \in \mathbb{R}^n$, a_{jk} , a_j are smooth real functions and a_{jk} is a non-negative matrix:

$$\sum_{j,k=1}^{n} a_{j,k}(y)\eta_j\eta_k \ge 0$$

and similarly

(7)
$$L_2 = -\sum_{j,k=1}^m b_{jk}(y)\partial_{y_j}\partial_{y_k} + \sum_{j=1}^m b_j(y)\partial_{y_j} + b_0(y)$$

where $y \in \mathbb{R}^m$, b_{jk} , b_j are smooth real functions and b_{jk} is a non-negative matrix:

$$\sum_{j,k=1}^m b_{j,k}(y)\eta_j\eta_k \ge 0$$

We were looking for conditions on each L_i , i = 1, 2 which are close to necessary for hypoellipticity, but are still strong enough to yield the hypoellipticity of $L_1 + g(x)L_2$. The (degenerate) function g(x) is assumed to be smooth, non-negative, and it does not vanish for $x \neq 0$:

(8)
$$g \in C^{\infty}(\mathbb{R}^n), \ g(x) > 0 \text{ for } x \neq 0$$

At x = 0 we allow vanishing, possibly of an infinite type, i.e. $\partial_x^{\alpha} g(0) = 0$ possibly for all multiindices α . Therefore, we consider the operator

(9)
$$L(x, y, D_x, D_y) = L_1(x, D_x) + g(x)L_2(y, D_y).$$

The main result of this paper is the following.

Theorem 1 (Main theorem). Let L be defined by (9) with assumptions on L_1 , L_2 and g as above, and let L_1 , L_2 satisfy the superlogarithmic estimate, that is for each $\varepsilon > 0$ and any compact sets $K_1 \subset \mathbb{R}^m$ and $K_2 \subset \mathbb{R}^n$, there exist constants C_{ε,K_i} , i = 1, 2, such that

(10)
$$\begin{aligned} \|\log\langle\xi\rangle^2 \,\hat{u}(\xi)\|^2 &\leq \varepsilon(L_1 u, u) + C_{\varepsilon, K_1} \|u\|^2, \quad \forall u \in C_0^\infty(K_1);\\ \|\log\langle\eta\rangle^2 \,\hat{v}(\eta)\|^2 &\leq \varepsilon(L_2 v, v) + C_{\varepsilon, K_2} \|v\|^2 \quad \forall v \in C_0^\infty(K_2); \end{aligned}$$

Then L is hypoelliptic.

For g(x) sufficiently degenerate, the operator L will violate the Morimoto's superlogarithmic estimate (10), yet still be hypoelliptic. The techniques of the proof rely mainly on pseudodifferential calculus and clever use of the operator Λ introduced by Michael Christ in [Chr01]. The general idea of the proof is similar to the previous results [Koh98] and [Chr01], however, the analysis is more delicate due to the presence of infinite degeneracy and the absence of subelliptic estimates at the same time. Moreover, the operator we consider is not of the form of sums of squares of vector fields, c.f. [Hör67, Chr01].

1.1. Necessity. As mentioned above, Morimoto in [Mor87], motivated by probabilistic techniques of [KS84], found that a symmetric operator $\partial_x^2 + L_2$ has to satisfy estimate (10) in order to be hypoelliptic. In the case of our operator $L = L_1 + g(x)L_2$, the validity of superlog estimates for L_1 and L_2 is also necessary for hypoellipticity as the following example due to Kusuoka and Stroock [KS84] demonstrates. **Example 1.** Let $L_1 := \partial_{x_1}^2 + a^2(x_1)\partial_{x_2}^2$, $L_2 := \partial_{y_1}^2$, and $g(x_1, x_2) \equiv 1$. Assume that the function $a(x_1)$ is smooth, nonnegative and satisfies a(0) = 0 and $a(x_1) \neq 0$ if $x_1 \neq 0$. L_1 is thus a Fedii type operator. It is has been shown by Kusuoka and Stroock, and later by Morimoto, that the operator $L = L_1 + L_2$ is hypoelliptic iff

(11)
$$\lim_{x \to 0} x \log a(x) = 0$$

It has been later shown by Christ [Chr01, Lemma 5.2] that (11) is equivalent to

(12)
$$\lim_{\tau \to \infty} \frac{\lambda_0(\tau)}{\log \tau} = \infty,$$

where $\lambda_0(\tau)$ is the principal eigenvalue of $L_{\tau} := -\partial_x^2 + \tau^2 a^2(x)$, i.e.

$$\lambda_0(\tau) = \inf \frac{\langle L_{\tau} f, f \rangle^{1/2}}{||f||_{L^2}}$$

where the infimum is taken over all $0 \neq f \in C_0^2(\mathbb{R})$. We will show that (12) implies the superlogarithmic estimate for L_1

 $\left| \left| \log \left(1 + |\xi_1|^2 + |\xi_2|^2 \right) \cdot \hat{u}(\xi_1, \xi_2) \right| \right|^2 \le \varepsilon(L_1 u, u) + C_{\varepsilon, K} ||u||^2, \quad u \in C_0^\infty(K), \quad K \subset \mathbb{R}^2.$

To estimate the left hand side we divide the integral into two

$$\int \int \log^2 \langle \xi_1, \xi_2 \rangle \hat{u}(\xi_1, \xi_2)^2 d\xi_1 d\xi_2 = \int \int_{|\xi_1| > |\xi_2|} \log^2 \langle \xi_1, \xi_2 \rangle \hat{u}(\xi_1, \xi_2)^2 d\xi_1 d\xi_2 + \int \int_{|\xi_1| \le |\xi_2|} \log^2 \langle \xi_1, \xi_2 \rangle \hat{u}(\xi_1, \xi_2)^2 d\xi_1 d\xi_2 =: I + II$$

For the first integral we have

$$I \lesssim \int \int_{|\xi_1| > |\xi_2|} \log^2 \langle \xi_1 \rangle \hat{u}(\xi_1, \xi_2)^2 d\xi_1 d\xi_2 \lesssim \int \int (\varepsilon \langle \xi_1 \rangle^2 + \frac{1}{\varepsilon}) \hat{u}(\xi_1, \xi_2)^2 d\xi_1 d\xi_2$$

= $\varepsilon \int \int (\partial_{x_1} u)^2 dx_1 dx_2 + \frac{1}{\varepsilon} \int \int u(x, y)^2 dx_1 dx_2 \leq \varepsilon (L_1 u, u) + \frac{1}{\varepsilon} ||u||^2$

where we used Plancherel and the definition of L_1 . For the second integral we have

(14)
$$II = \int \int_{|\xi_1| \le |\xi_2|} \log^2 \langle \xi_1, \xi_2 \rangle \hat{u}(\xi_1, \xi_2)^2 d\xi_1 d\xi_2 \le C \int \int \log^2 \langle \xi_2 \rangle \hat{u}(\xi_1, \xi_2)^2 d\xi_1 d\xi_2$$

(15)
$$= C \int \int \log^2 \langle \xi_2 \rangle \mathcal{F}_{\xi_2}(u)(x_1, \xi_2)^2 dx_1 d\xi_2$$

where \mathcal{F}_{ξ_2} stands for the partial Fourier transform in the second variable. Next, by definition of λ_0 we have for $f(x_1, \xi_2) := \mathcal{F}_{\xi_2}(u)(x_1, \xi_2)$

$$\int f(x_1,\xi_2)^2 dx_1 \leq \frac{1}{\lambda_0(\xi_2)^2} \langle L_{\xi_2} f(x_1,\xi_2), f(x_1,\xi_2) \rangle_{x_1}$$
$$= -\frac{1}{\lambda_0(\xi_2)^2} \langle \mathcal{F}_{\xi_2}(L_1 u)(x_1,\xi_2), \mathcal{F}_{\xi_2}(u)(x_1,\xi_2) \rangle_{x_1}$$

Combining with (14) we obtain by Plancherel and (12)

$$II \le C \int \int \frac{\log^2 \langle \xi_2 \rangle}{\lambda_0(\xi_2)^2} (\mathcal{F}_{\xi_2}(L_1 u)(x_1, \xi_2), \mathcal{F}_{\xi_2}(u)(x_1, \xi_2)) dx_1 d\xi_2 \le \varepsilon \langle L_1 u, u \rangle + C_\varepsilon ||u||^2$$

1.2. Outline of the paper. The paper is organized as follows. Section 2 is dedicated to the proof of the main sufficiency result Theorem 1. First, in Section 2.2 we reduce Theorem 1 to Theorem 10 which makes use of a special pseudodifferential operator Λ . Then, in Section 2.4 we introduce relevant symbol classes and derive some of their properties relevant to our estimates. Section 2.3 outlines the proof of Theorem 10, and sections 2.6 and 3 are dedicated to the proofs of the main technical tools, Poincaré inequality and commutator estimates. Finally, in the Appendix we provide some auxiliary results from calculus of pseudodifferential operators, and basic PDE theory, that are used throughout the paper.

2. Proof of sufficiency

2.1. Compactly supported distributions. We first claim that to show hypoellipticity it is sufficient to consider compactly supported distributions $u \in \mathscr{E}'$. In fact it is sufficient to consider functions supported near the degenacy of g(x). The following definition and two lemmas make the satement precise.

Definition 2. We say that the operator L is **locally** H^s **hypoelliptic** near $(x, y) \in \mathbb{R}^{n+m}$ if for any $u \in D'$ and $\phi \in C_0^{\infty}$, $\phi \equiv 1$ in a neighborhood of (x, y) satisfying $\phi Lu \in H^s$, there exists $\tilde{\phi} \in C_0^{\infty}$, $\tilde{\phi} \equiv 1$ in a (possibly smaller) neighborhood of (x, y), such that $\tilde{\phi}u \in H^s$.

Lemma 3. Let the operator L satisfy Definition 2 in a neighborhood of $(x_0, y_0) \in \mathbb{R}^{n+m}$ but only for compactly supported distributions. Then L is locally H^s hypoelliptic near $(x_0, y_0) \in \mathbb{R}^{n+m}$.

As $H^{\infty} = \bigcap_{s} H^{s}$, Lemma 3 for all s implies local hypoellipticity in the sense of (1).

Proof. Let $v \in D'$, not necessarily compactly supported, and suppose that $\phi Lv \in H^s$ for some $\phi \in C_0^{\infty}$. Let $\phi^* \in C_0^{\infty}$ be such that $\phi \subset \subset \phi^*$. Since $\phi^* = 1$ on the support of ϕ we then have $\phi L(\phi^*v) = \phi Lv \in H^s$, and obviously $\phi^*v \in \mathscr{E}'$. Therefore, since L is locally H^s hypoelliptic with compactly supported distributions, there exists $\tilde{\phi} \subset \subset \phi$ and $\tilde{\phi} = 1$ near (x_0, y_0) such that $\tilde{\phi}v = \tilde{\phi}\phi^*v \in H^s$. This concludes that Lis locally H^s hypoelliptic near (x_0, y_0) . **Lemma 4.** Let the operator L be defined as in (9) and let the assumptions of Theorem 1 hold. Assume that $v \in \mathscr{E}'$ and there exists a constant a > 0 such that for each $(x, y) \in \text{supp } v$ we have |x| > a, i.e. v is supported away from the degeneracy set of g(x). If $\phi L v \in H^s$ for some $\phi \in C_0^{\infty}$, $\phi \equiv 1$ in a neighborhood of $(0, y_0)$, then there exists $\tilde{\phi} \in C_0^{\infty}$, $\tilde{\phi} \equiv 1$ in a neighborhood of $(0, y_0)$, such that $\tilde{\phi} u \in H^s$.

Proof. Let $\zeta \in C_0^{\infty}$ and $\zeta = 1$ on $\operatorname{supp} v =: V$, and without loss of generality we may assume that $\zeta(x, y) = 0$ for |x| < a/2. We then have $\zeta Lv = Lv$, and we denote $\tilde{L} := \zeta L$. We now claim that operator \tilde{L} satisfies the superlogarithmic estimate, i.e.

(16)
$$||\log \langle \xi, \eta \rangle \cdot \hat{u}(\xi, \eta)||^2 \le \varepsilon(\tilde{L}u, u) + C_{\varepsilon, K} ||u||^2, \quad u \in C_0^{\infty}(K),$$

for each $\varepsilon > 0$ and any compact set $K \subset V$. To show that let $\inf_{|x|>a/2} g(x) = g_a > 0$, and let $K_1 \subset \mathbb{R}^m$ and $K_2 \subset \mathbb{R}^n$ satisfy $K \subset K_1 \times K_2 \subset V$. Then $u \in C_0^{\infty}(K)$ restricted to each variable, belongs to $C_0^{\infty}(K_i)$ and moreover $\tilde{L}u = L_1u + gL_2u$. We now apply the superlog estimate to each L_1 and L_2

$$\begin{aligned} ||\log(1+|\xi|^{2}) \cdot \mathcal{F}_{\xi}(u)(\xi,y)||^{2} &\leq \frac{\varepsilon}{2} (L_{1}u,u)_{x} + C_{\varepsilon,K_{1}} ||u(\cdot,y)||^{2}, \\ ||\log(1+|\eta|^{2}) \cdot \mathcal{F}_{\eta}(u)(x,\eta)||^{2} &\leq \frac{\varepsilon \cdot g_{a}}{2} (L_{2}u,u)_{y} + C_{\varepsilon,a,K_{2}} ||u(\cdot,y)||^{2} \\ &\leq \frac{\varepsilon}{2} (g(x)L_{2}u,u)_{y} + C_{\varepsilon,a,K_{2}} ||u(\cdot,y)||^{2}. \end{aligned}$$

Integrating each inequality in the second variable and adding, we obtain (16). We therefore can apply Theorem 1 of [Mor87] to conclude that \tilde{L} is hypoelliptic in V. Recalling the definition of \tilde{L} concludes the proof.

Remark 5. Lemmas 3 and 4 effectively say that that the operator L from Theorem 1 is hypoelliptic away from x = 0 based on earlier work of Morimoto [Mor87] (and Christ [Chr01], in the case of sum of squares). This is done for convenience, as many arguments are more delicate in the case of g(0) = 0. However, if desired, one can proceed with our argument, even if $g(0) \neq 0$. In particular, shifting coordinates makes our argument self-contained and not relying on Lemma 4.

Remark 6. Lemma 3 reduces the question of local hypoellipticity to a neighborhood of a point. Thus, when needed we may change the operator L away from a specified neighborhood. In particular, by changing coefficients of L away from a large compact set, we may assume that all the coefficient functions are bounded with bounded derivatives. I.e. $\|g\|_{W^{\infty,\infty}} < \infty$, $\|a_{ij}\|_{W^{\infty,\infty}} < \infty$, etc.

2.2. Reduction to Λ .

2.2.1. Definition of Λ . To recast the main Theorem 1 in terms of the special operator Λ we first introduce a relevant cutoff function

Definition 7 (Fixing cutoffs). Let $\chi_x(x) = \begin{cases} 0, |x| \ge 2\\ 1, |x| \le 1 \end{cases}$ be a compactly supported non-negative function. Let $y_0 \in \mathbb{R}^m$ and $\chi_y \in C_0^{\infty}(\mathbb{R}^m)$ be a bump function that is 1




FIGURE 1. A sketch of ϕ

in some neighborhood of y_0 .

Given $u \in \mathscr{E}'$ and $(0, y_0) \in \mathbb{R}^n \times \mathbb{R}^m$ let $R \gg 1$ be large enough, so that $\operatorname{supp} u \subset B_R(0) \times B_R(y_0)$. Let $0 < \delta_2 \leq \frac{1}{2R}$ be a small parameter and let

(17)
$$\phi(x,y) = \chi_x\left(\frac{x}{R}\right) \cdot \chi_y\left(\frac{y-y_0}{R}\right) - \chi_x\left(\frac{x}{\delta_2}\right)\chi_y(R(y-y_0)) \in C_0^\infty$$

be a localization near $(0, y_0)$ (see also Figure 1 on p.7). Then $\phi(x, y) = 1$, whenever (x, y) is away from $B_{\delta_2}(0) \times B_{\frac{1}{R}}(y_0)$ on supp u. In particular, except for a small neighborhood of $(0, y_0)$, $\phi(x, y) = 1$ on supp u.

Moreover, the key property that drives our argument, is that

(18)
$$\partial_{\mu}^{\beta}\partial_{x}^{\alpha}\phi(x,y) \equiv 0 \text{ for } |x| \le \delta_{2}, \text{ for } |\alpha| > 0$$

(19)
$$|\partial_y^\beta \phi(x,y)| \le C(\beta,R), \text{ for } |\beta| \ge 0$$

Define a PDO Λ by the following symbol

(20)
$$\lambda(x, y, \xi, \eta) = \langle \xi, \eta \rangle^s \exp(-(N + s + 3)\phi(x, y) \log\langle \xi, \eta \rangle) S_{\delta_5}$$

where $S_{\delta_5} = \langle \delta_5 \cdot (\xi, \eta) \rangle^{-(N+s+3)}$ is regularization operator of order -(N+s+3), defined in Lemma 45. In particular, its seminorms of order 0 are uniformly bounded in δ_5 , see Appendix.

Remark 8. Note, that S_{δ_5} is dependent on the parameter δ_5 , but its S^0 seminorms are S_{δ_5} independent. Other than a statement that if $u \in H^{-N}$, then $S_{\delta_5}u \in H^{s+3}$, all estimates from now on are uniform in δ_5 as $\delta_5 \to 0$.

2.2.2. Cutoff functions. The operator and symbol λ defined in (20) is a crucial ingredient of the paper. In particular, the proof of the Theorem 1 demands close attention to a parameter $\delta_2 > 0$. This parameter, δ_2 is closely related to the size of the localization in the degenerate regime from ϕ in (17). From now on we allow and keep track of dependence of all functions and parameters on δ_2 , chief among them λ .

Motivated by (18), we want to emphasize two classes of families of bump functions adapted to our degenerate operator (4). Roughly, those functions that behave like $\partial_x^{\alpha} \phi(x, y)$ for $|\alpha| > 0$, and those that do not have x derivatives. More precisely,

(21)
$$\tilde{\mathcal{O}}_{\delta_2} := \left\{ \tilde{\psi}_{\delta_2}(x, y) \in C_0^\infty : \tilde{\psi}_{\delta_2}(x, y) \equiv 0 \text{ for } |x| < \frac{\delta_2}{2} \right\}$$

As mentioned above, our motivating example is $\partial_x^{\alpha} \phi$ with ϕ from (17). With this notation $\partial_x^{\alpha} \phi \in \tilde{\mathcal{O}}_{\delta_2}$ for $|\alpha| > 1$.

We now introduce a second class, of functions, whose support may include the most degenerate region.

$$\mathcal{O} := \left\{ \psi_{\delta_2}(x,y) \in C^\infty : \partial_x^\alpha \psi_{\delta_2}(x,y) \in \tilde{\mathcal{O}}_{\delta_2} \text{ for } |\alpha| > 0 \text{ and } \limsup_{\delta_2 \to 0} |\partial_y^\beta \psi_{\delta_2}(x,y)| < \infty \right\}$$

The motivating example for \mathcal{O} is ϕ from (17) with bounds from (18). The key difference between $\tilde{\mathcal{O}}$ and \mathcal{O} classes are the bounds on the functions and their derivatives, which motivates the following remark.

Remark 9. All the symbols and functions below, unless explicitly stated otherwise, are dependent on the parameter $\delta_2 > 0$. For this reason, we suppress the subscript δ_2 below in the notation for functions and symbols. In particular, we will speak of $\tilde{\mathcal{O}}$ implying $\tilde{\mathcal{O}}_{\delta_2}$.

We reduce the proof of Theorem 1 to the following

Theorem 10. Let the operator L be defined by (9). There exists $\delta_2 > 0$ small enough such that with ϕ defined by (17) and the PDO Λ defined by (20), we have for any

(22)

 $u \in \mathscr{E}'$ and N so that $u \in H^{-N}$, with $Lu \in H^s_{loc}$ the following estimate $\|\Lambda u\| \lesssim \|\Lambda Lu\| + \|u\|_{-N}$

Proposition 11. Theorem 10 implies Theorem 1

We reduce the proof of the Proposition to the following two Lemmas that allow us to "remove" the operator Λ once the essential work of Theorem 10 is achieved.

Lemma 12. Let $\psi \in C_0^{\infty}$ be given, such that $\psi \equiv 1$ on a neighborhood of $(0, y_0)$. Let δ_2 be small enough, so that

(23)
$$B_{2\delta_2}(0) \times B_{2\delta_2}(y_0) \Subset \{(x,y) : \psi(x,y) = 1\}$$

Further, let $u \in \mathscr{E}'$ with $\operatorname{supp} u \subset B_R(0) \times B_R(y_0)$ and N be such that $u \in H^{-N}$. Then there exists a constant $C = C(\psi, s, R, N, \delta_2) > 0$, so that

(24)
$$||\Lambda Lu|| \le C(\psi, s, R, N, \delta_2) \left(||\psi Lu||_{H^s} + ||u||_{H^{-N}} \right)$$

Lemma 13. Let $\delta_2 > 0$ be given and consider $\phi_0 \in C_0^{\infty}(\mathbb{R}^n \times \mathbb{R}^m)$, so that

$$\operatorname{supp} \phi_0 \subset B_{\frac{1}{2}\delta_2}(0) \times B_{\frac{1}{2}\delta_2}(y_0)$$

Let u and N be as in Lemma 12. Then there exists a constant C independent of the parameter δ_5 , more precisely $C = C(\phi_0, s, R, N, \delta_2) > 0$, so that

(25)
$$||S_{\delta_5}\phi_0 u||_{H^s} \le C(\phi_0, s, R, N, \delta_2)||\Lambda u|| + ||u||_{H^{-N}}$$

Proof of Proposition 11. Let $u \in \mathscr{E}'$, such that $Lu \in H^s_{loc}$. Thus there exists $\psi \in C_0^{\infty}$ with $\psi \equiv 1$ on the relevant neighborhood, so that $\psi Lu \in H^s$. From $u \in \mathscr{E}'$, there exist an $N \in \mathbb{R}$, so that $u \in H^{-N}$.

Let $\delta_2 > 0$ be small enough, so that Theorem 10 holds. Choose a cutoff $\phi_0 \in C_0^\infty$ as in Lemma 13

Then

$$\begin{split} \|S_{\delta_5}\phi_0 u\|_{H^s} &\lesssim \|\Lambda u\| + \|u\|_{-N} \text{ by Lemma 13} \\ &\lesssim \|\Lambda L u\| + \|u\|_{-N} \text{ by Theorem 10} \\ &\lesssim \|\psi L u\|_{H^s} + \|u\|_{-N} \text{ by Lemma 12} \end{split}$$

where none of the constants depend on δ_5 .

We now take $\delta_5 \to 0$ to conclude $\phi_0 u \in H^s$. We proceed as follows.

First, take a sequence $\delta_5 \to 0$. By Banach-Alaouglu, since $S_{\delta_5}\phi_0 u$ is a bounded sequence in the Hilbert space H^s , it converges weakly in H^s up to a subsequence. Let $v = w - \lim_{\delta_5 \to 0} S_{\delta_5}\phi_0 u$ along this subsequence. Note that $v \in H^s$, and since weak limits in H^s are also distributional limits we have that $\lim_{\delta_5 \to 0} S_{\delta_5}\phi_0 u = v$ in the sense of distributions.

We now turn to strong limits. Recall that we have $u \in H^{-N}$, and thus $\lim_{\delta_5 \to 0} S_{\delta_5} \phi_0 u = \phi_0 u$ in H^{-N} as $S_{\delta_5} \in S^0$ uniformly in δ_5 . This in turn implies $\lim_{\delta_5 \to 0} S_{\delta_5} \phi_0 u = \phi_0 u$

in the sense of distributions. Since distributional limits are unique, we conclude that $\phi_0 u = v$. Hence $\phi_0 u \in H^s$ and $u \in H^s_{loc}$.

Proof of Lemma 12. Let ϕ_3 be a cut-off function, so that

$$B_{2\delta_2}(0) \times B_{2\delta_2}(y_0) \Subset \phi_3 \Subset \psi$$

I.e. $\phi_3 \equiv 1$ on $B_{2\delta_2}(0) \times B_{2\delta_2}(y_0)$, while the latter members of the family nest, i.e. $\psi \equiv 1$ on support of ϕ_3 .

We now begin the estimates.

$$\Lambda Lu = \phi_3 \Lambda Lu + (1 - \phi_3) \Lambda Lu$$
$$= \phi_3 \Lambda \psi Lu + \phi_3 \Lambda (1 - \psi) Lu + (1 - \phi_3) \Lambda Lu$$

Using triangle inequality, we separate the terms.

$$\|\Lambda Lu\| \le \|\phi_3 \Lambda \psi Lu\| + \|\phi_3 \Lambda (1-\psi) Lu\| + \|(1-\phi_3) \Lambda Lu\|$$

= I + II + III

Since $\phi_3 \Lambda \in S_{1,\varepsilon}^s$, by the boundedness of the ΨDO [Kg81]

$$I \le C(\delta_2, s, N) ||\psi Lu||_{H^s}$$

From the supports of ϕ_3 and ψ , $\phi_3 \Lambda(1-\psi) \in S^{-\infty}$, therefore

$$II \le C(\delta_2, s, N) ||u||_{H^{-N}}$$

Finally, for the third term, observe that on $\operatorname{supp}(1 - \phi_3) \cap \operatorname{supp} u$, $\phi \equiv 1$. Hence in this region Λ is smoothing of order N + 3:

$$||(1 - \phi_3)\Lambda Lu|| \le C(\delta_2, s, N)||u||_{H^{-N}}$$

Combining the estimates I - III we complete the proof.

Proof of Lemma 13. Define the following sequence of cutoffs, that nest inside the "degenerate" region, where ϕ from (17) is designed to vanish. I.e. let $\phi_0 \Subset \phi_1 \Subset \phi_2 \Subset \chi_{B_{\delta_2}(0) \times B_{\delta_2}(y_0)}$.

With cutoffs fixed to be used later, we want to commute the localization ϕ_0 with the derivative operators. By the calculus of Ψ DO we can express:

(26)
$$\langle (\xi,\eta) \rangle^s S_{\delta_5} \circ \phi_0 = \phi_1 a_0 \langle (\xi,\eta) \rangle^s S_{\delta_5} \mod S^{-N}$$

where we defined the symbol a_0 as

$$a_0(x, y, \xi, \eta) = \sum_{|\alpha| < s+N} \partial_{x,y}^{\alpha} \phi_0(x, y) \frac{\partial_{\xi, \eta}^{\alpha} [\langle (\xi, \eta) \rangle^s S_{\delta_5}]}{\langle (\xi, \eta) \rangle^s S_{\delta_5}}$$

As $\phi_1 \equiv 1$ on the support of ϕ_0 , it is harmless to add it in (26).

Note, that the bounds on the semi-norms of a_0 depend on (δ_2, s, N) through the support of ϕ_0 . However, a_0 seminorms can be bounded independely of δ_5 , as all the

 S^0 semi-norms of S_{δ_5} are bounded uniformly in δ_5 .

Next, recall that $\phi_2 \equiv 1$ on the support of ϕ_1 . Therefore, $\phi_1 a_0 \circ (1 - \phi_2) \in S^{-\infty}$. We thus have

$$\langle (\xi,\eta) \rangle^s S_{\delta_5} \circ \phi_0 = \phi_1 a_0 \phi_2 \langle (\xi,\eta) \rangle^s S_{\delta_5} \mod S^{-N}$$

We are now ready to complete the proof of (25). From the computations above

$$||S_{\delta_{5}}\phi_{0}u||_{H^{s}} \leq ||\phi_{1}a_{0}\phi_{2}\langle(\xi,\eta)\rangle^{s} S_{\delta_{5}}u|| + ||R_{-N}u||$$

$$\leq C(s,N,\delta_{2})||\phi_{2}\langle(\xi,\eta)\rangle^{s} S_{\delta_{5}}u|| + C||u||_{H^{-N}}$$

Where we applied the L^2 boundedness of $a_0 \in S^0$ and the $H^{-N} \to L^2$ boundedness of the, previously unnamed, remainder $R_{-N} \in S^{-N}$.

Finally, as on supp ϕ_2 , $\langle (\xi, \eta) \rangle^s S_{\delta_5} = \Lambda$, and ϕ_2 is bounded, the proof is complete.

2.3. **Proof of Theorem 10.** By Proposition 11, the proof of the hypoellipticity of L reduces to an estimate of the form

$$\|\Lambda u\| \lesssim \|\Lambda Lu\| + \|u\|_{-N}$$

In this section we reduce the argument to a series of the Lemmas and Propositions, whose proof is deferred to the later sections. The argument is inspired by [Fed71] and [Koh98].

(1) First, using a Poincaré type inequality we can bound the norm of a function with small support by the norm of its derivative. More precisely,

Proposition 14. Given $\delta_2 > 0$, set $\delta_1 = \frac{2(2\pi)^n}{\log^2 2\pi\delta_2}$. Then for Λ defined in (20) we get:

(28)
$$\|\Lambda u\|^2 \le \delta_1 \|\log\langle\xi\rangle \,\widehat{\Lambda u}(\xi, y)\|^2 + C_{\delta_1} \|u\|_{H^{-N}}$$

where we take the Fourier transform with respect to the variable x only.

We defer the proof of this result to section 2.6.

(2) Next, to make the operator appear, we use the superlog estimate for L_1 with $\varepsilon = 1$:

(29)
$$\|\log\langle\xi\rangle\,\widehat{\Lambda u}(\xi,y)\|^2 \le Re(L_1\Lambda u,\Lambda u) + C\|\Lambda u\|^2 \le Re(L\Lambda u,\Lambda u) + C\|\Lambda u\|^2$$

where the last inequality is due to positivity of the operators (Lemma 41 in the appendix applied to the operator $g(x)L_2(y,\partial_y)$). Note that we need to take real parts of expressions like $(L\Lambda u, \Lambda u)$ since even though u is a real function, Λu does not have to be.

(3) Combining estimates (29) and (28) gives

(30)
$$\|\Lambda u\|^2 \le \delta_1 \left[Re(L\Lambda u, \Lambda u) + C \|\Lambda u\|^2 \right] + C_{\delta_1} \|u\|_{H^{-N}}$$

To prove the main estimate (27), we want to commute L and Λ . I.e. the main task of the argument is the analysis of the commutator term

$$[L,\Lambda] = [L_1,\Lambda] + g[L_2,\Lambda] + [g,\Lambda]L_2 \equiv Q_1 + gQ_2 + Q_3$$

The commutators can be estimated at a price of extra logs, which appear in the non-degenerate region away from x = 0 or with a weight g(x):

Proposition 15 (Commutator estimate). For any $\delta_2 > 0$, there exists a function $\tilde{\phi} \in \tilde{\mathcal{O}}_{\delta_2}$, so that the following estimates hold.

$$Re(Q_1u,\Lambda u) \leq \frac{1}{2}Re(L\Lambda u,\Lambda u) + C_{\delta_2} \|\log\langle (\xi,\eta)\rangle \,\tilde{\phi}\Lambda u\|^2 + C \|\Lambda u\|^2 + C_{\delta_2} \|u\|_{H^{-N}}$$

$$Re(Q_1u,\Lambda u) \leq C \|\langle \overline{\chi}|_{\delta_2} |\langle \overline{\chi}|_{\delta_2} |\langle$$

 $Re(gQ_2u,\Lambda u) + Re(Q_3u,\Lambda u) \le C \|\sqrt{g}\log\langle(\xi,\eta)\rangle \Lambda u\|^2 + C\|\Lambda u\|^2 + C_{\delta_2}\|u\|_{H^{-N}}$

We defer the proof of this key proposition to a series of Lemmas in Section 3 and continue the argument below.

(4) Based on the superlogarithmic estimates (10) for L_1 and L_2 , we estimate the logarithmic terms in the commutators above.

Lemma 16. Let $\delta_0 > 0$ be given. Then there exists a decreasing function C_{δ_0} , so that

(32)
$$\|\sqrt{g}\log\langle(\xi,\eta)\rangle\Lambda u\|^2 \le \delta_0 Re(L\Lambda u,\Lambda u) + C_{\delta_0}\|\Lambda u\|^2$$

 $(L\Lambda u, \Lambda u) = (\Lambda L u, \Lambda u) + ([L, \Lambda]u, \Lambda u)$

Lemma 17 (Interpolation in nondegenerate region). Let $\varepsilon > 0$ and $\phi \in \tilde{\mathcal{O}}$. Then there exist constants \tilde{C}_{δ_2} and C_{ε} (increasing in the reciprocals of parameters ε and δ_2) such that the following estimate holds

(33)
$$\|\log\langle(\xi,\eta)\rangle\,\tilde{\phi}\Lambda u\| \le \varepsilon(L\Lambda u,\Lambda u) + \varepsilon\tilde{C}_{\delta_2}\|\Lambda u\|^2 + C_{\varepsilon}\|u\|_{H^{-N}}$$

We defer the proofs of Lemmas 16 and 17 to the section 2.7

(5) Combining Proposition 15 and Lemmas 16 and 17 we obtain

$$Re([L,\Lambda]u,\Lambda u) \leq (C + \varepsilon \cdot C_{\delta_2}C_{\delta_2} + C_{\delta_0}) \|\Lambda u\|^2 + (\frac{1}{2} + \varepsilon \cdot C_{\delta_2} + \delta_0 \cdot C)Re(L\Lambda u,\Lambda u) + C_{\varepsilon,\delta_2} \|u\|_{H^{-N}}$$

(6) We substitute this estimate into (31) with a choice of $\delta_0 = 1/(8C)$ and $\varepsilon = \min\{1/(8C_{\delta_2}), 1/(C_{\delta_2}\tilde{C}_{\delta_2})\}$ for δ_2 as in (17)

$$Re(L\Lambda u, \Lambda u) \le Re(\Lambda Lu, \Lambda u) + \frac{3}{4}Re(L\Lambda u, \Lambda u) + C\|\Lambda u\|^2 + C_{\delta_2}\|u\|_{H^{-N}}$$

and thus

(34)
$$Re(L\Lambda u, \Lambda u) \le 4Re(\Lambda Lu, \Lambda u) + C \|\Lambda u\|^2 + C_{\delta_2} \|u\|_{H^{-N}}$$

(7) Note that δ_1 is uniquely determined by δ_2 , and the connection is given in Proposition 14. Therefore, the constants depending on δ_1 can be thought of as constants depending on δ_2 and vice versa. Keeping that in mind, we use the estimate (34) in (30) to obtain:

$$\|\Lambda u\|^{2} \leq \delta_{1} \left[4Re(\Lambda Lu, \Lambda u) + C \|\Lambda u\|^{2} \right] + C_{\delta_{1}} \|u\|_{H^{-N}}$$

(8) Using Cauchy-Schwarz in the estimate above gives

$$\|\Lambda u\|^{2} \leq \|\Lambda Lu\|^{2} + \delta_{1} \cdot C \|\Lambda u\|^{2} + C_{\delta_{1}} \|u\|_{H^{-N}}$$

Finally, choosing δ_1 small enough (or equivalently choosing δ_2 small enough) so that

$$\delta_1 \cdot C = \frac{1}{2},$$

the estimate above implies (27) and completes the proof.

2.4. Symbol classes. We introduce the following bump function and symbol classes to simplify recurring calculations. Simply put, the proof of Theorem 1 fails with simple $L^2 \to H^s$ boundedness of Ψ DO in the class S^s . The key ingredient in the proof of the main theorem (Theorem 1) is a careful analysis of the commutator $[L, \Lambda]$ performed in the Proposition 15. In turn, the heart of the commutator argument are the support properties of the symbols and/or bounds on their semi-norms. The commutator $[L, \Lambda]$ has over 15 different terms on which calculus of Ψ DO is performed repeatedly causing "migration" of properties. After a lot of trial and error the following classes of symbols proved helpful. We first define three symbol classes, $\tilde{\mathcal{A}}$, $\tilde{\mathcal{B}}$ and \mathcal{D} , and prove their boundedness properties to serve as motivation.

2.4.1. Symbol classes. First, we introduce classes of families of symbols depending on parameter δ_2 . All the functions, symbols, and constants in the definitions below depend on δ_2 , unless explicitly stated otherwise, but we will suppress the dependence in our notation later on. These are standard symbols with an additional requirement that their semi-norms are δ_2 independent. More precisely,

(35)

$$\mathcal{A}_{j} \equiv \mathcal{A}_{j}^{\delta_{2}} := \{ a_{j}^{\delta_{2}} \in S^{j} : |\partial_{\xi,\eta}^{\alpha} \partial_{x,y}^{\beta} a_{j}^{\delta_{2}}(x,y,\xi,\eta)| \le C_{\alpha,\beta}^{\delta_{2}} \langle \xi \rangle^{j-|\alpha|} \text{ and } \limsup_{\delta_{2} \to 0} C_{\alpha,\beta}^{\delta_{2}} < \infty \}$$

What is the difference between a class $\tilde{\mathcal{A}}_j$ and a standard Ψ DO class S^j ? The interested reader may recall the Remark 9. That is, when talking about $a_j \in \mathcal{A}_j$, we really think of a δ_2 dependent function $a_{j,\delta_2}(x, y, \xi, \eta)$ with the property

(36)
$$\sup_{(x,y,\xi,\eta);\delta_2\to 0} \left|\partial^{\alpha}_{\xi,\eta}\partial^{\beta}_{x,y}a_{j,\delta_2}(x,y,\xi,\eta)\right| < \infty$$

From now on, whenever we say that a symbol (or semi-norm) is δ_2 independent, we imply that the symbol satisfies (36).

Ideally, all the terms arising from $[L, \Lambda]$ would be in the \mathcal{A}_j classes for some j. Since this is not the case, we need to introduce the negative order symbols.

(37)
$$S^{-} := \{ b \in \bigcup_{0 < \rho \le 1} S_{1,1/2}^{-\rho} \}$$

The essential properties of S^- class will be established in Lemma 21.

Third, we need symbols supported away from degeneracy.

(38)
$$\tilde{\mathcal{B}}_j \equiv \tilde{\mathcal{B}}_j^{\delta_2} := \left\{ \tilde{b}_j^{\delta_2} \in S^j : \exists b_j \in S^j \text{ and } \tilde{\psi}_{\delta_2} \in \tilde{\mathcal{O}}_{\delta_2}, \text{ with } \tilde{b}_j = \tilde{\psi}_{\delta_2}(x, y) b_j \right\}$$

As above, $\tilde{\mathcal{B}}_j$ is a subset or subclass of S^j , with a condition on the support of the symbol.

Fourth, we need symbols with \mathcal{O} factor, i.e. symbols whose derivatives vanish in certain directions:

(39)

$$\tilde{\mathcal{A}}_{j} \equiv \tilde{\mathcal{A}}_{j}^{\delta_{2}} := \left\{ \tilde{a}_{j}^{\delta_{2}} \in S^{j} : \exists m, a_{j,1}^{\delta_{2}}, \dots, a_{j,m}^{\delta_{2}} \in \mathcal{A}^{j} \text{ and } \psi_{1}^{\delta_{2}}, \dots, \psi_{m}^{\delta_{2}} \in \mathcal{O} \\
\text{with } \tilde{a}_{j}^{\delta_{2}} = \sum_{k=1}^{m} \psi_{k}^{\delta_{2}}(x, y) a_{j,k}^{\delta_{2}} \right\}$$

We need to allow for more than one $\tilde{\mathcal{O}}$ bump function, because we demand to factor them from the \mathcal{A} . E.g. $\partial_{y_k}\phi + \frac{\xi}{\langle \xi \rangle}\phi$ for $\phi \in \tilde{\mathcal{O}}$ cannot be written as $\psi(x, y) \cdot a_0(x, y, \xi, \eta)$ for $\psi \in \tilde{\mathcal{O}}$.

Fifth, we need a class of symbols $\tilde{\mathcal{A}}$ -like symbols, that are, in addition, independent of parameter ξ .

(40)
$$\mathcal{D}_j \equiv \mathcal{D}_j^{\delta_2} := \{ d_j^{\delta_2} \in \tilde{\mathcal{A}}_j^{\delta_2} : d_j^{\delta_2}(x, y, \xi, \eta) = d_j^{\delta_2}(x, y, \eta) \}$$

This implies that in addition to the boundedness properties of $\tilde{\mathcal{A}}_j$, the symbols $d_j \in \mathcal{D}$ class commute with all functions of x. In particular,

(41)
$$d_j(x, y, \eta)\sqrt{g(x)} = \sqrt{g(x)}d_j(x, y, \eta).$$

2.4.2. Boundedness of symbol classes. The main justification for introducing the language of $\tilde{\mathcal{A}}$ (and $\tilde{\mathcal{B}}$) classes is parameter-management of constants, that would be too tedious for the number of terms we have when we work with the commutator $[L, \Lambda]$. Namely, if we treat $\tilde{\mathcal{A}}_0$ terms as generic S^0 symbols, their $L^2 \to L^2$ boundedness constant would depend on δ_2 . Such a crude analysis would not allow our argument to proceed. Instead, we prove the following simple lemma that motivates the definition of $\tilde{\mathcal{A}}$:

Lemma 18. Let $\tilde{a}_j \in \tilde{\mathcal{A}}_j$. Then there exist a constant C, independent of δ_2 , so that $\|\tilde{a}_j v\|_{L^2} \leq C \|v\|_{H^j}$

Note that if a_j was a member of \mathcal{A}_j instead of $\tilde{\mathcal{A}}_j$, no work would have been needed. Indeed, by (35), the seminorms of such symbol are independent of the parameter δ_2 . Hence their $H^j \to L^2$ norms are independent.

Proof. From (39)

$$\tilde{a}_j = \sum_{k=1}^m \psi_k(x, y) a_{j,k}(x, y, \xi, \eta) \text{ for } a_{j,k} \in \mathcal{A}_j$$

and thus

$$\|\tilde{a}_{j}v\|_{L^{2}} \leq \sum_{k} \|\psi_{k}(x,y)\|_{L^{\infty}} \|a_{j,k}v\|_{L^{2}}$$
$$\leq C \|v\|_{H^{j}}$$

since $\|\psi_k(x,y)\|_{L^{\infty}}$ is bounded independently of δ_2 , which follows from the definition (22).

The following Lemma demonstrates the motivation for the class \mathcal{B} . The actual estimate needed has a few more technical details but the spirit of the argument is still captured.

Lemma 19. Let $\tilde{b} \in \tilde{\mathcal{B}}_0$, and $\tilde{b} = \tilde{\psi}b_0$ with $\tilde{\psi} \in \mathcal{O}$ as in (38). Then there exists a constant C_{δ_2} , that depends on δ_2 , such that

(42)
$$\|\tilde{b}v\|_{L^2} \le C_{\delta_2} \|\tilde{\psi}v\|_{L^2} + \|v\|_{H^{-1}}$$

Proof. From (38) we have $\tilde{b} = \tilde{\psi}b_0$, where $b_0 \in S^0$. Note that the semi-norms of the symbol b_0 may depend on δ_2 .

Now doing the composition calculus of the Ψ DO backwards

$$\tilde{b} = b_0 \circ \tilde{\psi} \mod S^{-1}$$

Now use L^2 boundedness to complete the proof.

We also motivate the use of class S^- , which is simply the symbols of order less than 0, whose support properties we do not track and whose semi-norms may depend on the parameter δ_2 . We also state a simple interpolation lemma, that we use repeatedly.

Lemma 20. Let $a(\xi)$ be a positive increasing function, with $\lim_{|\xi|\to\infty} a(\xi) = \infty$. Then for any $\varepsilon > 0$ and N > 0, there exists a constant $C_{\varepsilon,N}$, so that for any $v \in \mathscr{S}(\mathbb{R}^n)$

(43)
$$||v||^{2} \leq \varepsilon ||a(\xi)\hat{v}(\xi)||^{2} + C_{\varepsilon}||v||_{H^{-N}}^{2}$$

Proof. Let R > 1. By Plancherel

$$\|v\|^{2} = \frac{1}{(2\pi)^{n}} \left(\int_{|\xi| \ge R} \frac{a(\xi)}{a(\xi)} |\hat{v}^{2}(\xi)| d\xi + \int_{|\xi| \le R} \frac{\langle R \rangle^{N}}{\langle R \rangle^{N}} |\hat{v}^{2}(\xi)| d\xi \right)$$

Since $a(\xi)$ and $\langle \xi \rangle$ are monotone functions we can estimate the integrals above as

$$\|v\|^{2} \leq \frac{1}{(2\pi)^{n}} \left(\frac{1}{a(R)} \int_{|\xi| \geq R} a(\xi) |\hat{v}^{2}(\xi)| d\xi + \langle R \rangle^{N} \int_{|\xi| \leq R} \langle \xi \rangle^{-N} |\hat{v}^{2}(\xi)| d\xi \right)$$

Choosing R large enough, so that $\frac{1}{a(R)(2\pi)^n} \leq \varepsilon$ completes the proof.

Lemma 21. Let $\varepsilon > 0$, N, s and δ_2 be given. Then given a symbol $b^- \in S^-$ there exists a constant $C_{\varepsilon,\delta_2,N,s}$, such that for any $v \in \mathscr{S}$

$$||b^{-}v|| \leq \varepsilon ||v|| + C_{\varepsilon,\delta_2,N,s} ||v||_{H^{-N-s}}$$

Proof. By the definition of S^- in (37) $b^- \in S_{1,\frac{1}{2}}^{-\rho}$ for some $\rho > 0$. Applying boundedness of Ψ DO gives

$$||b^{-}v|| \le C_{\delta_2} ||v||_{H^{-\rho}}$$

Apply Lemma 20 with $a(\xi) = \langle \xi \rangle^{\rho}$. In order to eliminate the constant C_{δ_2} in front of v, replace ε in Lemma 20 with $\frac{\varepsilon}{C_{\delta_2}}$.

2.5. Derivatives of Operators and symbols. The following lemma provides useful estimates on the derivatives of λ .

Lemma 22. Let $\lambda(x, y, \xi, \eta)$ be defined by (20). Then we have

(44)
$$\frac{\partial_x^{\alpha}\lambda}{\left[\log\langle(\xi,\eta)\rangle\right]^{|\alpha|}\cdot\lambda} \in \tilde{\mathcal{B}}_0 \text{ for } |\alpha| > 0$$

(45)
$$\frac{\partial_y^{\alpha}\lambda}{\left[\log\langle(\xi,\eta)\rangle\right]^{|\alpha|}\cdot\lambda} \in \tilde{\mathcal{A}}_0 \text{ for } |\alpha| > 0$$

(46)
$$\frac{\partial_{\xi,\eta}^{\beta}\lambda}{\lambda} \in \tilde{\mathcal{A}}_{-|\beta|}, \text{ for } |\beta| > 0$$

Proof. First, we can calculate

$$\partial_{x_k} \lambda = -\phi_{x_k} (N + s + 3) \log\langle \xi, \eta \rangle \lambda$$
$$\partial_{y_k} \lambda = -\phi_{y_k} (N + s + 3) \log\langle \xi, \eta \rangle \lambda$$

Therefore, for $|\alpha| = 1$ (44) and (45) follow immediately from the definitions of ϕ and $\tilde{\mathcal{B}}_0$ and $\tilde{\mathcal{A}}_0$ classes. Differentiating again and iterating gives the result for $|\alpha| \geq 2$. To show (46) we calculate

$$\partial_{\xi_k} \lambda = \left(\left[s \frac{\xi_k}{\langle \xi, \eta \rangle^2} + \frac{\partial_{\xi_k} S_\delta}{S_\delta} \right] \cdot 1 - (N + s + 3) \frac{\xi_k}{\langle \xi, \eta \rangle^2} \phi \right) \lambda$$

and therefore we obtain

$$\tilde{a}_{-\alpha} := \frac{\partial_{\xi}^{\alpha} \lambda}{\lambda} \in \tilde{\mathcal{A}}_{-|\alpha|}$$

for $|\alpha| = 1$. With $\tilde{a}_{\alpha'}$ defined as above for any $|\alpha'| > 0$, we use the product rule for $|\alpha| = 1$ to obtain

(47)
$$\frac{\partial_{\xi}^{\alpha+\alpha'}\lambda}{\lambda} = \tilde{a}_{-\alpha} \cdot \tilde{a}_{-\alpha'} + \partial_{\xi}^{\alpha}\tilde{a}_{-\alpha'}$$

From the definition of $\tilde{\mathcal{A}}_j$, $\partial^{\alpha}_{\xi,\eta}$ derivatives lower the order into $\tilde{\mathcal{A}}_{j-|\alpha|}$ and by moving all $\tilde{\mathcal{O}}$ functions we preserve class membership as well. Thus (47) implies that

$$\frac{\partial_{\xi}^{\alpha}\lambda}{\lambda} \in \tilde{\mathcal{A}}_{-|\alpha|}$$

A similar computation is valid, for $\partial_{\eta}^{\alpha}\lambda$.

HYPOELLIPTICITY WITHOUT LOSS OF DERIVATIVES FOR FEDII'S TYPE OPERATORS 17

We now derive some useful properties of the following symbols

(48)
$$l_1(x,\xi) = \sum_{j,k=1}^n \alpha_{jk}(x)\xi_j\xi_k + \sum_{j=1}^n i\alpha_j(x)\xi_j + \alpha_0(x)$$

(49)
$$l_2(y,\eta) = \sum_{j,k=1}^m \beta_{jk}(y)\eta_j\eta_k + \sum_{j=1}^m i\beta_j(y)\eta_j + \beta_0(y)$$

Where we denote by l_1 and l_2 the symbols of the operators L_1 and L_2 respectively.

Lemma 23. Let l_1 and l_2 be defined by (48) and (49). The following properties hold

- $l_1, l_2 \in \mathcal{A}_2$
- l_1 and l_2 are self-adjoint principal symbols, i.e. $\overline{l_i} l_i \in \mathcal{A}_1$
- for $|\alpha| = 2$, $\partial_{\xi}^{\alpha} l_1$ and $\partial_{\eta}^{\alpha} l_2$ depend only on x and y respectively.

Proof. The first two properties follow immediately from (48) and (49), as the operators are independent of δ_2 and related localization. To show the last property we differentiate

$$\partial_{\xi_k \xi_j} l_1 = \alpha_{jk}(x)$$
$$\partial_{\eta_k \eta_j} l_2 = \beta_{jk}(y)$$

which concludes the proof.

2.6. **Poincaré inequality.** We first prove a preliminary Lemma that is only valid for compactly supported functions, and then deduce Proposition 14 as a simple corollary.

Lemma 24 (Poincare). Suppose $v \in C_0^1(B_{\delta_2}(0))$, where $B_{\delta_2}(0) \subset \mathbb{R}^n$ is a ball of radius δ_2 centered at the origin. Then

$$\|v\|^2 \le \delta_1 \|\log\langle\xi\rangle \,\hat{v}(\xi)\|^2$$

where $\delta_1 = \frac{2(2\pi)^n}{\log^2 4\pi\delta_2}$. In particular, by choosing δ_2 small enough, we can make δ_1 appropriately small.

Proof. From the definition of the Fourier transform and Cauchy-Schwartz inequality

(50)
$$|\hat{v}(\xi)| = \left| \int_{|x| \le \delta_2} e^{-ix \cdot \xi} v(x) dx \right| \le \omega_n^{\frac{1}{2}} \delta_2^{\frac{n}{2}} ||v||$$

where ω_n is the volume of an *n*-dimensional unit ball. Next by Plancherel

$$\|v\|^{2} \leq (2\pi)^{n} \int_{|\xi| \leq \frac{1}{C\delta_{2}}} |\hat{v}(\xi)|^{2} d\xi + (2\pi)^{n} \int_{|\xi| \geq \frac{1}{C\delta_{2}}} |\hat{v}(\xi)|^{2} d\xi := II_{1} + II_{2}$$

for any C > 0. Using (50) for II_1 and monotonicity of the logarithm for II_2 gives

$$\|v\|^{2} \leq \left(\frac{2\pi}{C}\right)^{n} \|v\|^{2} + \frac{(2\pi)^{n}}{\log^{2}\frac{1}{C\delta_{2}}} \int_{|\xi| \geq \frac{1}{C\delta_{2}}} \log^{2}|\xi| |\hat{v}(\xi)|^{2} d\xi$$

Thus for $C = 4\pi$,

$$||v||^2 \le \frac{2(2\pi)^n}{\log^2 4\pi\delta_2} \int_{|\xi| \ge \frac{1}{C\delta_2}} \log^2 \langle \xi \rangle \, |\hat{v}(\xi)|^2 d\xi$$

Proof of Proposition 14. Let $v = \psi \Lambda u$ for $\phi \in \psi$ with ϕ from (20) and use the fact that $(1 - \psi)\Lambda \in S^{-N}$

2.7. Superlog estimates. In this section we prove Lemmas 16 and 17

Proof of Lemma 16. First, an elementary calculation shows:

$$\begin{aligned} \|\sqrt{g}\log\langle(\xi,\eta)\rangle v\| &\leq \|\sqrt{g}(x)\log\langle\eta\rangle v\| + \|\sqrt{g}(x)\left[\log\langle(\xi,\eta)\rangle - \log\langle\eta\rangle\right]v\| \\ &\leq \|\sqrt{g}(x)\log\langle\eta\rangle v\| + C\|\left[\log\langle(\xi,\eta)\rangle - \log\langle\eta\rangle\right]v\| \end{aligned}$$

where we used $||g||_{L^{\infty}} \leq C$ in the second line. Next, it is easy to see that

$$0 \le \log \langle (\xi, \eta) \rangle - \log \langle \eta \rangle \le \log \langle \xi \rangle$$

Thus we have $\|[\log\langle(\xi,\eta)\rangle - \log\langle\eta\rangle]v\| \le \|\log\langle\xi\rangle v\|$ and hence

(51)
$$\|\sqrt{g}\log\langle(\xi,\eta)\rangle v\| \le \|\sqrt{g}(x)\log\langle\eta\rangle v\| + C\|\log\langle\xi\rangle v\|$$

We now apply this estimate to our operator. From estimate (10) for L_2 with $\varepsilon = \frac{\delta_0}{2}$

$$\|\sqrt{g(x)}\log\langle\eta\rangle\Lambda u\|^2 \le \frac{\delta_0}{2}(\sqrt{g}L_2\sqrt{g}\Lambda u,\Lambda u) + C_{\delta_0}\|\sqrt{g}\Lambda u\|^2$$

Using positivity of the operator L_1 (Lemma 41) and the estimate $||g||_{L^{\infty}} \leq C$ (see Remark 6) we obtain from the estimate above

$$\|\sqrt{g(x)}\log\langle\eta\rangle\Lambda u\|^2 \le \frac{\delta_0}{2}Re(L\Lambda u,\Lambda u) + C_{\delta_0}\|\Lambda u\|^2$$

Now let $\varepsilon = \frac{\delta_0}{2C}$ for C > 0 from (51). Apply (10) for L_1 with this choice of ε and use the positivity of gL_2 , Lemma 41, as above to obtain

$$\|\log\langle\xi\rangle\Lambda u\|^2 \le \frac{\delta_0}{2C}Re(L\Lambda u,\Lambda u) + C_{\delta_0}\|\Lambda u\|^2$$

Combining the last two estimates with (51) completes the proof.

Proof of Lemma 17. Recall that we are aiming to prove the following estimate:

$$\|\log\langle(\xi,\eta)\rangle\,\tilde{\phi}\Lambda u\|^2 \le \varepsilon Re(L\Lambda u,\Lambda u) + \varepsilon C_{\delta_2}\|\Lambda u\|^2 + C_{\varepsilon}\|u\|_{H^{-N}}^2$$

The proof will proceed in five steps:

- (1) Split the logarithm similar to Lemma 16.
- (2) Use estimate (10) for L_1 and L_2 .
- (3) Insert g(x) using the support of $\tilde{\phi}\Lambda u$.
- (4) Remove ϕ from the operator.
- (5) Interpolate the remainders.

The first three steps are essentially identical to the Proof of Lemma 4, with more detail.

(1) From
$$\langle (\xi, \eta) \rangle \leq \langle \xi \rangle \langle \eta \rangle$$
 we get

$$\log\langle(\xi,\eta)\rangle \le \log\langle\eta\rangle + \log\langle\xi\rangle$$

Thus

(52)
$$\|\log\langle(\xi,\eta)\rangle\,\hat{\phi}\Lambda u\| \le \|\log\langle\eta\rangle\,\hat{\phi}\Lambda u\| + \|\log\langle\xi\rangle\,\hat{\phi}\Lambda u\|$$

(2) Using (10) for L_1 we get for any $\varepsilon_1 > 0$

$$|\log\langle\xi\rangle\,\tilde{\phi}\Lambda u\|^2 \le \varepsilon_1 Re(L_1\tilde{\phi}\Lambda u,\tilde{\phi}\Lambda u) + C_{\varepsilon_1}\|\tilde{\phi}\Lambda u\|^2$$

Meanwhile, (10) for L_2 gives for any $\varepsilon_2 > 0$

$$|\log\langle\eta\rangle\,\tilde{\phi}\Lambda u\|^2 \leq \varepsilon_2 Re(L_2\tilde{\phi}\Lambda u,\tilde{\phi}\Lambda u) + C_{\varepsilon_2}\|\tilde{\phi}\Lambda u\|^2$$

where we have integrated in the complementary variables to recover the full norms. Adding to (52) gives

(53)

$$\|\log\langle(\xi,\eta)\rangle\,\tilde{\phi}\Lambda u\|^2 \le 2\varepsilon_1 Re(L_1\tilde{\phi}\Lambda u,\tilde{\phi}\Lambda u) + 2\varepsilon_2 Re(L_2\tilde{\phi}\Lambda u,\tilde{\phi}\Lambda u) + C_{\varepsilon_1,\varepsilon_2}\|\tilde{\phi}\Lambda u\|^2$$

(3) Define a decreasing function C_{δ_2} by

$$\frac{1}{C_{\delta_2}} = \inf_{|x| \ge \delta_2} \{g(x)\}$$

In particular, we have $|x| \ge \delta_2$ on support of $\tilde{\phi}$, and thus $g(x) - \frac{1}{C_{\delta_2}} \ge 0$ there. Therefore, the principal symbol of the operator $(g(x) - \frac{1}{C_{\delta_2}})L_2$ is nonnegative on the support of $\tilde{\phi}$. We now use this positivity together with Lemma 41 from Appendix to get an operator estimate:

$$Re\left((g(x) - \frac{1}{C_{\delta_2}})L_2\tilde{\phi}\Lambda u, \tilde{\phi}\Lambda u\right) \ge -\tilde{C}_{\delta_2}\|\tilde{\phi}\Lambda u\|^2$$

Moving all the terms to one side and multiplying by $2\varepsilon_2 C_{\delta_2} > 0$ we obtain

(54)
$$0 \le 2\varepsilon_2 C_{\delta_2} Re\left((g(x)L_2\tilde{\phi}\Lambda u, \tilde{\phi}\Lambda u) - 2\varepsilon_2 Re(L_2\tilde{\phi}\Lambda u, \tilde{\phi}\Lambda u) + 2\varepsilon_2\tilde{C}_{\delta_2}C_{\delta_2} \|\tilde{\phi}\Lambda u\|^2 \right)$$

Finally, we add (53) to (54) and set $\varepsilon_1 = \varepsilon_2 C_{\delta_2} = \frac{\varepsilon}{2}$ for $\varepsilon > 0$ to obtain

$$\|\log\langle(\xi,\eta)\rangle\,\tilde{\phi}\Lambda u\|^2 \leq \varepsilon Re(L\tilde{\phi}\Lambda u,\tilde{\phi}\Lambda u) + C_\varepsilon \|\tilde{\phi}\Lambda u\|^2$$

(4) We now proceed to eliminate $\tilde{\phi}$ from the operator. The process is similar to the previous step. First, observe that

$$\tilde{\phi}L\tilde{\phi} = \tilde{\phi}^2L + \tilde{\phi}[L,\tilde{\phi}] = I + II$$

We next compute the symbol of the operator $II = \tilde{\phi}[L, \tilde{\phi}] \in \tilde{S}^1$, whose principal symbol is anti-self adjoint. Therefore, we can estimate it

$$Re(\tilde{\phi}[L,\tilde{\phi}]v,v) \le C_{\tilde{\phi}} \|v\|^2.$$

To estimate the *I* term we add non-negative expression to recreate operator *L*. More precisely, $(1 - \tilde{\phi}^2)L \ge 0$ is a non-negative operator and satisfies

$$Re((1 - \tilde{\phi}^2)Lv, v) \ge -C \|v\|^2$$

Hence

$$Re(\tilde{\phi}^2 Lv, v) \le Re(Lv, v) + C \|v\|^2$$

Combining the estimates for I and II we obtain

$$Re(L\phi\Lambda u,\phi\Lambda u) \leq Re(L\Lambda u,\Lambda u) + C_{\phi} \|\Lambda u\|^2.$$

Returning to the end of the previous step we conclude with

(55)
$$\|\log\langle(\xi,\eta)\rangle\,\tilde{\phi}\Lambda u\|^2 \le \varepsilon Re(L\Lambda u,\Lambda u) + \varepsilon C_{\tilde{\phi}}\|\Lambda u\|^2 + C_{\varepsilon}\|\tilde{\phi}\Lambda u\|^2$$

(5) To complete the proof it suffices to eliminate the last term in the estimate. We do so by interpolation, Lemma 20. Namely, we apply Lemma 20 with $a(\langle (\xi, \eta) \rangle) = \log \langle (\xi, \eta) \rangle$ and N + s + 3 in place of N to obtain

$$\|v\| \le \varepsilon' \|\log\langle (\xi,\eta)\rangle \,\hat{v}\| + C_{\varepsilon'} \|v\|_{H^{-N-s-3}}$$

Now let $v = \tilde{\phi} \Lambda u$, it is easy to check that $v \in H^3$ so we can substitute it in the formula above. Using $\Lambda \in S^s$, we obtain for $\varepsilon' = \frac{C_{\varepsilon}}{2}$ small enough

$$C_{\varepsilon} \| \tilde{\phi} \Lambda u \| \le \frac{1}{2} \| \log \langle (\xi, \eta) \rangle \, \tilde{\phi} \Lambda u \| + C_{\varepsilon} \| u \|_{H^{-N}}$$

The last estimate provides together with (55) gives (33).

3. Commutators

We now estimate the commutators Q_1 , Q_2 , and Q_3 separately. The outline of all three commutators goes through the same four steps, which we outline here.

- (0) When working with commutators we distinguish symbols of several classes that were defined in section 2.4:
 - Classes \mathcal{B} supported away from the degenerate region, which allows them to absorb logarithms.
 - \bullet Classes $\mathcal A$ with norms independent of the localization to degenerate region
 - Classes \mathcal{A} symbols that are flat in the degenerate region. These operators give good constants without the logs or when the weight g(x) is added.
 - Classes $\tilde{\mathcal{D}}$ symbols in $\tilde{\mathcal{A}}$ class that in addition do not depend on ξ .
 - Classes S^- symbols of negative order
 - Symbols that we denote c_0 that do not belong to any of the above classes, and are dealt with separately in Lemma 29.
- (1) Express the symbol of the commutator $q_j = p_j \cdot \lambda \mod S^{-N}$ using calculus of PDO and accounting for symbol classes from above

- (2) Convert the product of symbols into the composition: $Q_j = \tilde{P}_j \circ \Lambda \mod S^{-N}$, again accounting for supports as above
- (3) Find the adjoint part of the operator P_i to use the cancellation
- (4) Use the structure of the classes to obtain the estimates needed for Proposition 15

3.1. Commutator Q_1 . Calculus. We consider $Q_1 = [L_1, \Lambda]$. From the calculus of PDO (treating $\lambda \in S_{1,\varepsilon}^s$ for arbitrary small $\varepsilon > 0$) we get

(56)
$$q_1 = \sum_{|\alpha|=1}^{2} \frac{i^{|\alpha|}}{\alpha!} \left(\partial_{\xi}^{\alpha} l_1 \partial_x^{\alpha} \lambda - \partial_x^{\alpha} l_1 \partial_{\xi}^{\alpha} \lambda \right) - \sum_{|\alpha|=3}^{(N+s+1)} \frac{i^{|\alpha|}}{\alpha!} \partial_x^{\alpha} l_1 \partial_{\xi}^{\alpha} \lambda \mod S_{1,\varepsilon}^{-N}$$

This computation used *y*-independence of l_1 .

We now begin following the outline by decomposing the q_1 term into the classes of symbols $\tilde{\mathcal{A}}, \tilde{\mathcal{B}}, \ldots$ See section 2.4 as technical details arise.

Lemma 25. There exists an operator

(57)
$$p_1 = b_1 \log \langle (\xi, \eta) \rangle + b_0 \log^2 \langle (\xi, \eta) \rangle + a_1 + a_0 + r^-$$
$$so \ that \ q_1 = p_1 \cdot \lambda \mod S_{1,\varepsilon}^{-N}$$

where every term is explicitly defined in terms of given operators in the proof, but in particular,

(58)
$$b_i \in \tilde{\mathcal{B}}_i; \ a_i \in \tilde{\mathcal{A}}_i, \ for \ i = 0, 1; \ r^- \in S^-$$

Furthermore, the principal part of symbols b_1 and a_1 are purely imaginary.

Proof. From the calculus of (56) we obtain the following expressions by separating terms by first order, zero-th order and negative order and factoring λ :

$$q_1 = \lambda \left([b_1 \log \langle (\xi, \eta) \rangle + a_1] + [b_0 \log^2 \langle (\xi, \eta) \rangle + a_0] + r^- \right) \mod S^{-N}$$

Recall, from Lemma 22, $\partial_x^{\alpha} \lambda$ terms lead to logarithms, with a redeeming $\tilde{\mathcal{B}}$ class symbol. We group them with derivatives of l_1 into b_i s. $\partial_{\xi}^{\alpha} \lambda$ terms are logarithm free and have symbols in $\tilde{\mathcal{A}}$ classes. By Lemma 23 the factors of l_1 are independent of the degeneracy and do not influence class membership. To confirm the reasoning above we explicitly write down all the terms of non-negative order, where we need the explicit for of the a_1 for later calculations.

$$(59) \quad b_{1} := i \sum_{|\alpha|=1} \partial_{\xi}^{\alpha} l_{1} \frac{\partial_{x}^{\alpha} \lambda}{\lambda \cdot \log\langle(\xi,\eta)\rangle} \in \tilde{\mathcal{B}}_{1} \quad b_{0} := -\sum_{|\alpha|=2} \frac{1}{\alpha!} \partial_{\xi}^{\alpha} l_{1} \frac{\partial_{x}^{\alpha} \lambda}{\lambda \cdot \log^{|\alpha|} \langle(\xi,\eta)\rangle} \in \tilde{\mathcal{B}}_{0};$$

$$a_{1} := i \sum_{|\alpha|=1} \frac{1}{\alpha!} \partial_{x}^{\alpha} l_{1} \frac{\partial_{\xi}^{\alpha} \lambda}{\lambda} \in \tilde{\mathcal{A}}_{1} \qquad a_{0} := -\sum_{|\alpha|=2} \frac{1}{\alpha!} \partial_{x}^{\alpha} l_{1} \frac{\partial_{\xi}^{\alpha} \lambda}{\lambda} \in \tilde{\mathcal{A}}_{0};$$

$$(60) \quad r^{-} := \sum_{j=3}^{N+s+1} \sum_{|\alpha|=j} \frac{i^{|\alpha|}}{\alpha!} \partial_{x}^{\alpha} l_{1} \frac{\partial_{\xi}^{\alpha} \lambda}{\lambda} \in S^{-}$$

Finally, the principal parts of symbols b_1 and a_1 are purely imaginary, from their definition and because corresponding parts of λ and l_1 are real by the definition of λ in (20) and (48).

We now convert the product in Lemma 25 into a composition.

Lemma 26. q_1 can be rewritten as $q_1 = \tilde{p}_1 \circ \lambda \mod S^{-N}$ with (61) $\tilde{p}_1 = b_1 \log\langle (\xi, \eta) \rangle + a_1 + b'_0 \log^2 \langle (\xi, \eta) \rangle + a_0 + c_0 \log \langle (\xi, \eta) \rangle + \tilde{r}^-;$

where b_1 , a_1 and a_0 are from Lemma 25; $b'_0 \in \tilde{\mathcal{B}}_0$ and $\tilde{r}^- \in S^-$ are explicitly defined in terms of given operators in the proof. We also have $c_0 \in \tilde{\mathcal{A}}$, where c_0 is defined explicitly by the following formula:

(62)
$$c_0 = \sum_{|\alpha| = |\beta| = 1} \partial_x^{\alpha} l_1 \partial_{\eta}^{\beta} \left[\frac{\partial_{\xi}^{\alpha} \lambda}{\lambda} \right] \frac{\partial_y^{\beta} \lambda}{\lambda \log\langle (\xi, \eta) \rangle}$$

Proof. We need to show that $q_1 = \tilde{p}_1 \circ \lambda$ with \tilde{p}_1 defined in the Lemma. We express the composition using the calculus of Ψ DO and compare $\tilde{p}_1 \circ \lambda$ against $p_1 \cdot \lambda$ from Lemma 25 for terms of order 1, 0 and below.

In particular, the principal symbol of $\tilde{p}_1 \circ \lambda$ is $(b_1 \log \langle (\xi, \eta) \rangle + a_1) \cdot \lambda$, which coincides with $p_1 \cdot \lambda$. For terms of order 0 (with logarithms), we get

(63)
$$\left[b_0' \log^2 \langle (\xi, \eta) \rangle + a_0 + \mathbf{c_0} \log \langle (\xi, \eta) \rangle \right] \cdot \lambda + i \sum_{|\beta|=1} \partial_{\xi,\eta}^{\beta} \left[b_1 \log \langle (\xi, \eta) \rangle + a_1 \right] \partial_{x,y}^{\beta} \lambda$$

we match these terms with $[b_0 \log^2 \langle (\xi, \eta) \rangle + a_0] \cdot \lambda$ from (57). Note, that a_0 remains unchanged and the symbol b'_0 gathers all terms, where a derivative $\partial_x^{\alpha} \lambda$ is present for $|\alpha| > 0$. However, the expression (63), contains the term without $\partial_y^{\alpha} \lambda$, which is not in the $\tilde{\mathcal{B}}$ class, yet has a logarithm. We include such terms into a new symbol c_0 . More precisely, we define c_0 by terms we need to absorb:

(64)
$$c_0 = -i \sum_{|\alpha|=1} \partial_{\eta}^{\alpha} a_1 \frac{\partial_y^{\alpha} \lambda}{\lambda \log\langle (\xi, \eta) \rangle}$$

Substitution of a_1 from (59) gives the formula (62).

HYPOELLIPTICITY WITHOUT LOSS OF DERIVATIVES FOR FEDIJ'S TYPE OPERATORS 23

Terms of order 0 in $\tilde{p}_1 \circ \lambda$ are chosen to match with $p_1 \cdot \lambda$ by making an appropriate choice of b'_0 . I.e.

$$b_0' = b_0 - i \sum_{|\beta|=1} \partial_{\xi,\eta}^{\beta} \left(b_1 \log \langle (\xi,\eta) \rangle \right) \frac{\partial_{x,y}^{\beta} \lambda}{\lambda \log^2 \langle (\xi,\eta) \rangle} - i \sum_{|\beta|=1} \partial_{\xi}^{\beta} a_1 \frac{\partial_x^{\beta} \lambda}{\lambda \log^2 \langle (\xi,\eta) \rangle} \in \tilde{\mathcal{B}}^0$$

where b_j , a_j are from (59) in Lemma 25. Terms of order less than 0: \tilde{r}^- are treated similarly. We do it explicitly defining inductively as follows:

(66)
$$\tilde{r}^{-} = \sum_{k=1}^{N+s+1} r_{k}^{-} \in S^{-}$$

with the first term r_1^- defined as follows

(67)
$$r_1^- = r^- - \sum_{j=2}^{N+s+1} \sum_{|\beta|=j} \frac{i^{|\beta|}}{\beta!} \partial_{\xi,\eta}^\beta [b_1 \log\langle (\xi,\eta) \rangle + a_1] \frac{\partial_{x,y}^\beta \lambda}{\lambda}$$

(68)
$$-\sum_{j=1}^{N+s+1}\sum_{|\beta|=j}\frac{i^{|\beta|}}{\beta!}\partial_{\xi,\eta}^{\beta}[c_0\log\langle(\xi,\eta)\rangle+b_0'\log^2\langle(\xi,\eta)\rangle+a_0]\frac{\partial_{x,y}^{\beta}\lambda}{\lambda}$$

and the lower order terms defined inductively for $k\geq 2$

$$r_{k}^{-} = -\sum_{|\gamma|=1}^{k-1} \partial_{\xi,\eta}^{\gamma} r_{k-1}^{-} \cdot \frac{\partial_{x,y}^{\gamma} \lambda}{\lambda} \in S^{-}$$

Lemma 27. The operator \tilde{p}_1 from Lemma 26 is anti-self-adjoint to the top order. *I.e.*

(69)
$$\tilde{p}_1 + \tilde{p}_1^* = b_0'' \log^2 \langle (\xi, \eta) \rangle + a_0' + \mathbf{c}_0' \log \langle (\xi, \eta) \rangle + R^-$$

where we have explicit expression for every term:

$$\begin{aligned} a_0' &= a_0 + \sum_{|\beta| = |\alpha| = 1} \partial_{\xi}^{\beta} \left[\partial_x^{\alpha + \beta} l_1 \frac{\partial_{\xi}^{\alpha} \lambda}{\lambda} \right] \in \tilde{\mathcal{A}}_0; \\ c_0' &= c_0 + \sum_{|\beta| = |\alpha| = 1} \frac{\partial_x^{\alpha} l_1}{\log \langle (\xi, \eta) \rangle} \partial_{\eta}^{\beta} \partial_y^{\beta} \left[\frac{\partial_{\xi}^{\alpha} \lambda}{\lambda} \right]; \\ b_0'' &= b_0' + \sum_{|\beta| = |\alpha| = 1} \partial_{\xi}^{\beta} \left[\partial_x^{\alpha} l_1 \partial_x^{\beta} \frac{\partial_{\xi}^{\alpha} \lambda}{\lambda} \right] - i \sum_{|\beta| = 1} \overline{\partial_{\xi, \eta}^{\beta} \partial_{\xi, \eta}^{\beta} b_1} \in \tilde{\mathcal{B}}_0; \end{aligned}$$

and the remainder $R^- \in S^-$.

Proof. By the Lemma 26, b_1 and a_1 are purely imaginary to the top order. Hence from the calculus of Ψ DO the operator \tilde{p}_1 is anti-self adjoint at the top order. More explicitly,

$$\tilde{p}_1^* = \sum_{|\beta|=0}^{N+1} \frac{i^{|\beta|}}{\beta!} \overline{\partial_{x,y}^\beta \partial_{\xi,\eta}^\beta \tilde{p}_1};$$

All terms of negative order are absorbed into R^- . Terms with symbols of the form $\log^2\langle (\xi,\eta)\rangle \tilde{\mathcal{B}}_0$ are absorbed into b''_0 . Symbols in $\tilde{\mathcal{A}}_0$ are absorbed into a'_0 . Finally, terms that belong to neither $\tilde{\mathcal{B}}_0$ nor $\tilde{\mathcal{A}}_0$ are explicitly added to c'_0 .

3.2. Bounds on Q1. The Lemmas in the previous subsections allow us to rewrite the commutator Q_1 into a sum of a, b and c terms, up to a remainder. The a terms, at least for Q_1 , do not have logarithmic derivatives; b terms have additional logs, but this is redeemed by the support of the symbol. Finally, c terms have all the flaws, but their redeeming feature is that they preserve a substantial part of the original operator. More precisely from Lemma 26

(70)
$$Q_1 = \tilde{P}_1 \Lambda + R_{-N}$$

Whereas (69) in Lemma 27 together with Lemma 44 from the Appendix imply

(71)
$$Re(\tilde{P}_1v, v) \le Re\left((b_0''\log^2\langle(\xi, \eta)\rangle + a_0' + c_0'\log\langle(\xi, \eta)\rangle)v, v\right) + ||R^-v||||v||$$

Combining these two estimates for $v = \Lambda u$ reduces the first half of the Proposition 15 to bounds on symbols a_0-c_0 and the remainder. Half of them were already done in Lemmas 18 and 21. The term $c_0 \log\langle (\xi, \eta) \rangle$ was not treated in section 2.4.1 and requires a different analysis. Meanwhile $b''_0 \log^2 \langle (\xi, \eta) \rangle$ follows the ideas of Lemma 19 with a bit more of Ψ DO calculus. We establish these facts here, starting with an argument for $c_0 \log \langle (\xi, \eta) \rangle$ term. This argument, requires the following Lemma.

Lemma 28 (Oleinik-Radkevich). Let K be a compact set and L_1 a non-negative elliptic operator with smooth coefficients of the form (2) with symbol l_1 . Then there exists a constant C_K , such that for all $v \in C_0^{\infty}(K)$ the following estimate holds

$$\|(\partial_x^\beta l_1)v\|_{H^{-1}}^2 \le C_K Re\left((L_1v, v) + \|v\|^2\right)$$

Proof. This estimate is stated as equation (1.2) on p.3 of [Mor87], with reference to the proof in [OR73].

Lemma 29. Let c'_0 be from (69). Then the following estimate holds

(72)

$$Re(c_{0}'\log\langle(\xi,\eta)\rangle\Lambda u,\Lambda u) \leq \frac{1}{2}Re(L\Lambda u,\Lambda u) + C\|\Lambda u\|^{2} + C_{\delta_{2}}\|\Lambda u\|^{2}_{H^{-1/2}} + C\|u\|^{2}_{H^{-N}}$$

HYPOELLIPTICITY WITHOUT LOSS OF DERIVATIVES FOR FEDII'S TYPE OPERATORS 25

Proof. From (69) and definition of c'_0 in the Lemma 27, we can express this symbol as

$$c_{0}' = \sum_{|\beta|=|\alpha|=1} \frac{\partial_{x}^{\alpha} l_{1}}{\log\langle(\xi,\eta)\rangle} \left(\partial_{\eta}^{\beta} \partial_{y}^{\beta} \left[\frac{\partial_{\xi}^{\alpha} \lambda}{\lambda}\right] + \partial_{\eta}^{\beta} \left[\frac{\partial_{\xi}^{\alpha} \lambda}{\lambda}\right] \frac{\partial_{y}^{\beta} \lambda}{\lambda}\right);$$
$$= \sum_{|\alpha|=1} \partial_{x}^{\alpha} (l_{1}) \cdot a_{-2,\alpha}(x, y, \xi, \eta)$$

for $a_{-2,\alpha} \in \tilde{\mathcal{A}}_{-2}$. Since $l_1 \in \mathcal{A}_2$, the composition is of order 0 (with good constant). The obstacle is an extra logarithm that has to be treated differently.

First, observe that by the calculus of ΨDO

$$c_0' \log \langle (\xi, \eta) \rangle = \sum_{|\alpha|=1} a_{-2,\alpha} \log \langle (\xi, \eta) \rangle \circ \partial_x^{\alpha} l_1 + r_{-\frac{1}{2}}, \quad \text{where } r_{-\frac{1}{2}} \in S^{-\frac{1}{2}}$$

Second, we apply the boundedness of Ψ DO for the second term above and Lemma 18 for the first to obtain

(73)
$$\begin{aligned} \|c_{0}' \log\langle(\xi,\eta)\rangle v\| &\leq C \sum_{|\alpha|=1} \|\log\langle(\xi,\eta)\rangle \,\partial_{x}^{\alpha} l_{1} \cdot v\|_{H^{-2}} + C_{\delta_{2}} \|v\|_{H^{-\frac{1}{2}}} \\ &\leq C \sum_{|\alpha|=1} \|\partial_{x}^{\alpha} l_{1} \cdot v\|_{H^{-1}} + C_{\delta_{2}} \|v\|_{H^{-\frac{1}{2}}}. \end{aligned}$$

Third, let $\psi_1 \in \psi_2 \in C_0^{\infty}$ with $\psi_1 \equiv 1$ on supp u. We split

$$v = \Lambda u = \psi_2 \Lambda u + (1 - \psi_2) \Lambda u$$

Note, that by pseudo-locality of Ψ DO $(1-\psi_2)\Lambda\psi_1 \in S^{-\infty}$ and similarly for $\partial_x^{\beta}\psi_2\Lambda\psi_1 \in S^{-\infty}$ for $|\beta| > 0$.

We now apply Lemma 28 to $\psi_2 \Lambda u \in H^3$ (with δ_5 dependent bounds) and compact support by the ψ_2 localization. That Lemma implies

(74)
$$\|\partial_x^{\alpha} l_1 \psi_2 \Lambda u\|_{H^{-1}} \le C \left(Re(L_1 \psi_2 \Lambda u, \psi_2 \Lambda u) + \|\psi_2 \Lambda u\|_{L^2} \right)$$

We now commute L_1 and ψ_2 observing that $[L_1, \psi_2] \Lambda \psi_1 \in S^{-\infty}$:

$$(L_1\psi_2\Lambda u, \psi_2\Lambda u) = (\psi_2^2 L_1\Lambda u, \Lambda u) + (\psi_2[L_1, \psi_2]L\Lambda u, \Lambda u)$$
$$Re(L_1\psi_2\Lambda u, \psi_2\Lambda u) \le Re(\psi_2^2 L_1\Lambda u, \Lambda u) + C \|u\|_{H^{-N}}^2$$

Similarly, by $S^{-\infty}$ property,

$$\|\partial_x^{\alpha} l_1 (1 - \psi_2) \Lambda u\|_{H^{-1}} \le C \|u\|_{H^{-N}}^2$$

Combining the last three estimates with (73), we conclude

(75)

$$\|c_0'\log\langle(\xi,\eta)\rangle v\| \le C \ Re(\psi_2^2 L_1 \Lambda u, \Lambda u) + C \|u\|_{H^{-N}}^2 + C \|\psi_2 \Lambda u\|_{L^2} + C_{\delta_2} \|\Lambda u\|_{H^{-\frac{1}{2}}}$$

Fourth, as $(1 - \psi_2)L_1 + g(x)L_2$ is a non-negative operator, we can use Lemma 41 to replace $\psi_2^2 L_1$ with L as follows

$$Re(\psi_2^2 L_1 v, v) \le Re(Lv, v) + C \|v\|^2$$

We now combine this positivity estimate, with (75) and Cauchy-Schwartz inequality to obtain:

$$Re(c_0' \log\langle (\xi, \eta) \rangle \circ \Lambda u, \Lambda u) \le \varepsilon \|c_0' \log\langle (\xi, \eta) \rangle \Lambda u\|^2 + C_\varepsilon \|\Lambda u\|^2$$
$$\le \varepsilon CRe(Lu, u) + C_\varepsilon \|\Lambda u\|^2 + (\varepsilon C_{\delta_2} + C_\varepsilon) \|\Lambda u\|_{H^{-\frac{1}{2}}} + C \|u\|_{H^{-N}}^2$$

Now a choice of $\varepsilon = \frac{1}{2C}$ completes the proof.

The next lemma gives estimates for the term $b_0'' \log^2 \langle (\xi, \eta) \rangle + a_0'$.

Lemma 30. Let a'_0 be as in (69). Then

(76)
$$Re(a'_0\Lambda u, \Lambda u) \le C \|\Lambda u\|^2$$

Proof. The result follows immediately from Lemma 18 by taking j = 0.

Lemma 31. Let b_0'' be as in (69), and let $\tilde{\psi} \in \tilde{O}$ with $\tilde{\psi} \equiv 1$ on supp b_0'' as in Lemma 12. Then

$$Re(b_0''\log^2\langle(\xi,\eta)\rangle\Lambda u,\Lambda u) \le C_{\delta_2} \|\log\langle(\xi,\eta)\rangle\,\tilde{\psi}\Lambda u\|^2 + C_{\delta_2} \|\Lambda u\|_{H^{-1/2}}^2.$$

The argument is essentially a slight modification of (42). We present the full justification below.

Proof. Note, that from the definition of λ , (20) and symbol classes \tilde{B} in (38), b''_0 is compactly supported in the physical space (x, y). Therefore, there exists $\tilde{\psi} \equiv 1$ on its support. Moreover, from the definition of \tilde{B} , we may ensure that $\tilde{\psi} \in \tilde{\mathcal{O}}$.

We claim, that b_0'' can be rewritten as follows with the help of the Ψ DO calculus (77) $b_0'' \log^2 \langle (\xi, \eta) \rangle = \tilde{\psi}^2 b_0'' \log^2 \langle (\xi, \eta) \rangle = \tilde{\psi} \log \langle (\xi, \eta) \rangle \circ b_0'' \circ \log \langle (\xi, \eta) \rangle \tilde{\psi} \mod S^{-\frac{1}{2}}$ Indeed, on the symbol level, the two sides agree perfectly, because of the support properties of b_0'' .

Applying (77) we get

(78)
$$Re(\tilde{b}_{0}'' \log^{2}\langle(\xi,\eta)\rangle \Lambda u, \Lambda u) = Re(\tilde{b}_{0}'' \circ \log\langle(\xi,\eta)\rangle \tilde{\psi}\Lambda u, \log\langle(\xi,\eta)\rangle \tilde{\psi}\Lambda u)$$

(79)
$$\leq C_{\delta_{2}} \|\log\langle(\xi,\eta)\rangle \tilde{\psi}\Lambda u\|^{2}$$

which completes the proof

We now return to (70) and (71) to conclude the Q_1 estimates.

Corollary 32. The first estimate in the Proposition 15 holds. I.e. (80)

$$Re(Q_1u,\Lambda u) \le C \|\Lambda u\|^2 + \frac{1}{2} Re(L\Lambda u,\Lambda u) + C_{\delta_2} \|\log\langle (\xi,\eta)\rangle \,\tilde{\phi}\Lambda u\|^2 + C_{\delta_2} \|u\|_{H^{-N}}$$

Proof. Combining (70) and (71) for $v = \Lambda u$ it remains to estimate 4 terms. Lemma 30 estimates a'_0 term. Lemma 29 estimates c_0 term. Lemma 21 estimates $R^- \in S^-$ term.

Meanwhile, Lemma 31 estimates $b_0^{''}$ term, where we interpolate the $H^{-\frac{1}{2}}$ norm between H^{-N-s} and L^2 norms using Lemma 21.

3.3. Commutator Q_2 . Calculus. We consider $Q_2 = [L_2, \Lambda]$. From the calculus of PDO (treating $\lambda \in S_{1,\varepsilon}^s$ for arbitrary small $\varepsilon > 0$) we get

$$q_2 = \sum_{|\alpha|=1}^2 \frac{i^{|\alpha|}}{\alpha!} \left(\partial_{\eta}^{\alpha} l_2 \partial_y^{\alpha} \lambda - \partial_y^{\alpha} l_2 \partial_{\eta}^{\alpha} \lambda \right) - \sum_{|\alpha|=3}^{N+s+1} \frac{i^{|\alpha|}}{\alpha!} \partial_y^{\alpha} l_2 \partial_{\eta}^{\alpha} \lambda \mod S_{1,\varepsilon}^{-N}$$

which using (45) and (46) gives

Proposition 33. There exist operators $d_j = d_j(x, y, \eta) \in \mathcal{D}_j$ and $a_j = a_j(x, y, \eta, \xi) \in \tilde{\mathcal{A}}_j$ for j = 0, 1 and $r^- \in S^-$ satisfying $\overline{a_1} - a_1 \in \tilde{\mathcal{A}}_0$, such that

 $q_2 = p_2 \lambda := \left(d_1 \log \langle (\xi, \eta) \rangle + a_1 + d_0 \log^2 \langle (\xi, \eta) \rangle + a_0 + r^- \right) \lambda + R_{-N}$

where $R_{-N} \in S_{1,\varepsilon}^{-N}$.

Proof. For $|\alpha| = 1$ we have $\partial_{\eta}^{\alpha} l_2 \in \mathcal{A}_1$ and $\frac{\partial_y^{\alpha} \lambda}{\log^{|\alpha|} \langle (\xi,\eta) \rangle \cdot \lambda} \in \mathcal{O}$ both independent of ξ , which gives the first term $d_1 \log \langle (\xi,\eta) \rangle \lambda$. We also have $\partial_y^{\alpha} l_2 \in \mathcal{A}_2$, $\frac{\partial_{\eta}^{\alpha} \lambda}{\lambda} \in \tilde{\mathcal{A}}_{-1}$ for $|\alpha| = 1$ which gives the second term $a_1 \lambda$. Similarly, we obtain the third and forth terms by taking $|\alpha| = 2$, and $r^- \lambda$ accommodates $|\alpha| \geq 3$ terms.

Lemma 34. $p_2\lambda$ can be rewritten as $p_2\lambda = \tilde{p}_2 \circ \lambda \mod S_{1,\varepsilon}^{-N}$ with

$$\tilde{p}_2 = d_1 \log \langle (\xi, \eta) \rangle + a_1 + \tilde{d}_0 \log^2 \langle (\xi, \eta) \rangle + (\tilde{a}_0 + \tilde{b}_0) \log \langle (\xi, \eta) \rangle + a_0 + \tilde{r}^-$$

for \tilde{a}_j , \tilde{d}_j with the same properties as above and $\tilde{b}_j \in \tilde{\mathcal{B}}_j$ for j = 0, 1, and $\tilde{r}^- \in S^-$. *Proof.* We can write

$$(\tilde{p}_2 - \tilde{r}^-) \circ \lambda = (\tilde{p}_2 - \tilde{r}^-) \lambda + \sum_{\alpha} \frac{i^{|\alpha|}}{\alpha!} \left(\partial_{\xi}^{\alpha} \left(\tilde{p}_2 - \tilde{r}^- \right) \partial_x^{\alpha} \lambda + \partial_{\eta}^{\alpha} \left(\tilde{p}_2 - \tilde{r}^- \right) \partial_y^{\alpha} \lambda \right)$$
(81)

$$= \left(d_1 \log \langle (\xi, \eta) \rangle + a_1 + \left(\tilde{d}_0 + \sum_{|\alpha|=1} i \partial_\eta^{\alpha} d_1 \frac{\partial_y^{\alpha} \lambda}{\lambda \log \langle (\xi, \eta) \rangle} \right) \log^2 \langle (\xi, \eta) \rangle \right) \lambda$$

$$+ \left(\left(\left(\tilde{a}_0 + \tilde{b}_0 + \sum_{|\alpha|=1} i \left(\partial_{\xi,\eta}^{\alpha} a_1 + d_1 \partial_{\xi,\eta}^{\alpha} \log \langle (\xi, \eta) \rangle \right) \frac{\partial_{x,y}^{\alpha} \lambda}{\lambda \log \langle (\xi, \eta) \rangle} \right) \log \langle (\xi, \eta) \rangle + a_0 \right) \lambda$$

$$+ r_{def}^- \lambda \mod S_{1,\varepsilon}^{-N}$$

where we defined

$$\begin{aligned} r_{def}^{-} &:= \sum_{|\alpha|=1} \partial_{\xi,\eta}^{\alpha} \left(\tilde{d}_0 \log^2 \langle (\xi,\eta) \rangle + (\tilde{a}_0 + \tilde{b}_0) \log \langle (\xi,\eta) \rangle \right) \frac{\partial_{x,y}^{\alpha} \lambda}{\lambda} \\ &+ \sum_{|\alpha|=2}^{N+s+1} \partial_{\xi,\eta}^{\alpha} \left(\tilde{p}_2 - \tilde{r}^- \right) \frac{\partial_{x,y}^{\alpha} \lambda}{\lambda} \in S^-. \end{aligned}$$

Now we choose

(82)
$$\tilde{d}_0 = d_0 - \sum_{|\alpha|=1} i \partial_\eta^\alpha d_1 \frac{\partial_y^\alpha \lambda}{\lambda \log\langle (\xi, \eta) \rangle}$$

(83)
$$\tilde{a}_0 = -\sum_{|\alpha|=1} i \left(\partial_\eta^\alpha a_1 + d_1 \partial_\eta^\alpha \log\langle (\xi, \eta) \rangle \right) \frac{\partial_y^\alpha \lambda}{\lambda \log\langle (\xi, \eta) \rangle}$$

(84)
$$\tilde{b}_0 = -\sum_{|\alpha|=1} i \left(\partial_{\xi}^{\alpha} a_1 + d_1 \partial_{\xi}^{\alpha} \log \langle (\xi, \eta) \rangle \right) \frac{\partial_x^{\alpha} \lambda}{\lambda \log \langle (\xi, \eta) \rangle}$$

It follows from (44)-(46) that these symbols belong to the corresponding classes. Here the coefficients from $\tilde{\mathcal{B}}$ class appear when differentiating λ with respect to x. We define the remainder term $\tilde{r}^- := \sum_{k=1}^{N+s+1} r_k^-$, where each r_k^- is defined inductively as follows

$$\begin{aligned} r_1^- &= r^- - r_{def}^- \\ r_k^- &= -\sum_{|\alpha|=1}^{N+s+1} \partial_{\xi,\eta}^{\alpha} r_{k-1}^- \frac{\partial_{x,y}^{\alpha} \lambda}{\lambda}, \quad k \geq 2. \end{aligned}$$

One can check that this gives

$$\tilde{p}_2 \circ \lambda = p_2 \lambda \mod S_{1,\varepsilon}^{-N}.$$

We now use the fact that the top order terms are purely imaginary to obtain a better estimate for the real part of $(gQ_2u, \Lambda u)$.

Lemma 35. Let $(g\tilde{p}_2)^*$ be the operator adjoint of $(g\tilde{p}_2)$. Then

$$(g\tilde{p}_2)^* + (g\tilde{p}_2) = g\left(\tilde{d}'_0\log^2\langle(\xi,\eta)\rangle + (\tilde{a}'_0 + \tilde{b}'_0)\log\langle(\xi,\eta)\rangle + a_0\right) \\ + \sum_{|\alpha_1|=1} \partial_x^{\alpha_1} g\left(\tilde{a}'_0 + \tilde{a}''_0\log\langle(\xi,\eta)\rangle\right) + r^-$$

where the symbols are different from before but belong to the same symbol classes. Proof. First, using Lemma 34 we can write

$$g\tilde{p}_2 = gd_1 \log\langle (\xi,\eta) \rangle + ga_1 + g\tilde{d}_0 \log^2 \langle (\xi,\eta) \rangle + g(\tilde{a}_0 + \tilde{b}_0) \log \langle (\xi,\eta) \rangle + ga_0 + \tilde{r}^-,$$

HYPOELLIPTICITY WITHOUT LOSS OF DERIVATIVES FOR FEDII'S TYPE OPERATORS 29

where r^{-} is a symbol of negative order. Using this we have

$$(g\tilde{p}_2)^* = \sum_{|\alpha|=0}^{1} \frac{i^{|\alpha|}}{\alpha!} \partial^{\alpha}_{\xi,\eta} \partial^{\alpha}_{x,y} \overline{(g\tilde{p}_2)} + r_1^-$$

= $\overline{(g\tilde{p}_2)} + ga'_0 \log\langle(\xi,\eta)\rangle + ga''_0 + \sum_{|\alpha_1|=1} (\partial^{\alpha_1}_x g) \left(\tilde{a}_0 + \tilde{a}'_0 \log\langle(\xi,\eta)\rangle\right) + r_2^-$

so the anti self-adjoint terms cancel and we obtain the desired result.

3.4. Commutator Q_3 . Calculus. Let $Q_3 \equiv [g(x), \Lambda]L_2$, then

$$q_3 = \sum_{|\alpha|=1}^{2} \frac{i^{|\alpha|}}{\alpha!} \left(-\partial_x^{\alpha} g \partial_{\xi}^{\alpha} \lambda \right) \circ l_2 - \sum_{|\alpha|=3}^{N+s+1} \frac{i^{|\alpha|}}{\alpha!} \partial_x^{\alpha} g \partial_{\xi}^{\alpha} \lambda \circ l_2 \quad mod \quad S_{1,\varepsilon}^{-N}$$

Proposition 36. There exist operators $a_j \in \tilde{\mathcal{A}}_j$ for j = 0, 1, -1, and $r^- \in S^-$ satisfying

 $\overline{a_1} - a_1 \in \tilde{\mathcal{A}}_0$, such that

$$q_3 = p_3 \lambda := \left(\sum_{|\alpha|=1} \partial_x^{\alpha} g a_1 + a_0 + a_{-1} \right) \lambda + r^- \quad mod \quad S_{1,\varepsilon}^{-N}.$$

Proof. First, we compute

$$\sum_{|\alpha|=1}^{2} \frac{i^{|\alpha|}}{\alpha!} \left(-\partial_x^{\alpha} g \partial_{\xi}^{\alpha} \lambda \right) - \sum_{|\alpha|=3}^{N+s+1} \frac{i^{|\alpha|}}{\alpha!} \partial_x^{\alpha} g \partial_{\xi}^{\alpha} \lambda = \sum_{|\alpha|=1} \partial_x^{\alpha} g a_{-1} \lambda + a_{-2} \lambda + a_{-3} \lambda + R_{-N} \lambda + R_{-$$

where $a_{-j} \in \tilde{\mathcal{A}}_{-j}$ and $R_{-N} \in S_{1,\varepsilon}^{-N}$. Next, using the PDO calculus for compositions

$$\begin{pmatrix} \sum_{|\alpha|=1} \partial_x^{\alpha} g a_{-1} \lambda + a_{-2} \lambda + a_{-3} \lambda \end{pmatrix} \circ l_2 = \left(\sum_{|\alpha|=1} \partial_x^{\alpha} g a_{-1} \lambda + a_{-2} \lambda + a_{-3} \lambda \right) l_2 + \sum_{|\beta| \ge 1} \partial_{\eta}^{\beta} (a_{-1}' \lambda) \partial_y^{\beta} l_2 \\ = \sum_{|\alpha|=1} \partial_x^{\alpha} g a_1 \lambda + a_0 \lambda + a_{-1} \lambda$$

where we used the notation

$$a'_{-1} := \sum_{|\alpha|=1} \partial_x^{\alpha} g a_{-1} + a_{-2} + a_{-3}$$

This concludes the proof.

Lemma 37. There exist operators $a_j \in \tilde{\mathcal{A}}_j$ and $b_j \in \tilde{\mathcal{B}}_j$ for j = 0, 1 and $r^- \in S^-$, such that $p_3\lambda$ can be rewritten as $p_3\lambda = \tilde{p}_3 \circ \lambda \mod S_{1,\varepsilon}^{-N}$ with

$$\tilde{p}_3 = \sum_{|\alpha_1|=1} \partial_x^{\alpha_1} g a_1 + \partial_x^{\alpha_1} g (a'_0 + b'_0) \log \langle (\xi, \eta) \rangle + \tilde{a}_0 + r^-.$$

Proof. Proceeding as in the proof of Lemma 34 we compute

$$(\tilde{p}_{3} - r^{-}) \circ \lambda = (\tilde{p}_{3} - r^{-}) \lambda + \sum_{\alpha} \frac{i^{|\alpha|}}{\alpha!} \left(\partial_{\xi}^{\alpha} \left(\tilde{p}_{3} - r^{-} \right) \partial_{x}^{\alpha} \lambda + \partial_{\eta}^{\alpha} \left(\tilde{p}_{3} - r^{-} \right) \partial_{y}^{\alpha} \lambda \right) (85) \qquad = \sum_{|\alpha_{1}|=1} \partial_{x}^{\alpha_{1}} g a_{1} \lambda + \partial_{x}^{\alpha_{1}} g \left((a_{0}' + b_{0}) + \sum_{|\alpha|=1} i \partial_{\xi,\eta}^{\alpha} a_{1} \frac{\partial_{x,y}^{\alpha} \lambda}{\lambda \log\langle(\xi,\eta)\rangle} \right) \log\langle(\xi,\eta)\rangle \lambda + r_{def}^{-} \lambda + \tilde{a}_{0} \lambda \mod S_{1,\varepsilon}^{-N}$$

where we defined

$$r_{def}^{-} := \sum_{|\alpha|=1} \partial_{\xi,\eta}^{\alpha} \left(\partial_{x}^{\alpha_{1}} g a_{1} (a_{0}' + b_{0}') \log \langle (\xi, \eta) \rangle + \tilde{a}_{0} \right) \frac{\partial_{x,y}^{\alpha} \lambda}{\lambda} + \sum_{|\alpha|=2}^{N+s+1} \partial_{\xi,\eta}^{\alpha} \left(\tilde{p}_{3} - r^{-} \right) \frac{\partial_{x,y}^{\alpha} \lambda}{\lambda} \in S^{-}.$$

Now choose

(86)
$$a_0' = -\sum_{|\alpha|=1} i \partial_\eta^\alpha a_1 \frac{\partial_y^\alpha \lambda}{\lambda \log\langle (\xi, \eta) \rangle}$$

(87)
$$b_0' = -\sum_{|\alpha|=1} i \partial_{\xi}^{\alpha} a_1 \frac{\partial_x^{\alpha} \lambda}{\lambda \log\langle (\xi, \eta) \rangle}$$

Finally, let $\tilde{r}^- := \sum_{k=1}^{N+s+1} r_k^-$, and define inductively r_k^- , $k = 1, \ldots N + s + 1$ the same way as in Lemma 34 to obtain

$$\tilde{p}_3 \circ \lambda = p_3 \lambda \mod S_{1,\varepsilon}^{-N}.$$

Lemma 38. Let \tilde{p}_3^* be the operator adjoint to \tilde{p}_3 . Then

$$\tilde{p}_3 + \tilde{p}_3^* = \sum_{|\alpha|=1} \partial_x^{\alpha} g(a_0'' + b_0'') \log\langle (\xi, \eta) \rangle + \tilde{a}_0 + r^-.$$

where the symbols are different from before but belong to the same symbol classes.

Proof. The proof proceeds the same way as the proof of Lemma 35 so we omit it. \Box

3.5. Bounds on Q_2 and Q_3 . Gathering the estimates from the previous two sections together and using Lemma 44 from the Appendix we obtain

$$\begin{aligned} Re(gQ_{2}u,\Lambda u) + Re(Q_{3}u,\Lambda u) \\ = & Re\left(g(\tilde{d}_{0}\log^{2}\langle(\xi,\eta)\rangle + (\tilde{a}_{0} + \tilde{b}_{0})\log\langle(\xi,\eta)\rangle) \circ \Lambda u,\Lambda u\right) \\ & + \sum_{|\alpha|=1} \frac{1}{2}Re\left((\partial_{x}^{\alpha}g(a_{0}'' + b_{0}'')\log\langle(\xi,\eta)\rangle) \circ \Lambda u,\Lambda u\right) \\ & + Re(a_{0}\Lambda u,\Lambda u) + Re(r^{-}\Lambda u,\Lambda u) + Re(R_{-N}u,\Lambda u) \end{aligned}$$

We will now obtain bounds for the terms on the right to finish the proof of the second part of Proposition 15. First, using Lemmas 21 and 18 we obtain the estimates

$$|(r^{-}\Lambda u, \Lambda u)| \leq C ||\Lambda u||^2 + C_{\delta_2} ||u||^2_{-N},$$

$$|(a_0\Lambda u, \Lambda u)| \leq C ||\Lambda u||^2,$$

and by Cauchy-Schwartz

$$|(R_{-N}u, \Lambda u)| \le C_{\delta_2} ||u||_{-N}^2 + C||\Lambda u||^2.$$

The next Lemma provides a bound for $((\partial_x^{\alpha}g(a_0''+b_0'')\log\langle(\xi,\eta)\rangle) \circ \Lambda u, \Lambda u).$

Lemma 39. Let $|\alpha| = 1$, then $Re\left(\left(\partial_x^{\alpha}g(a_0''+b_0'')\log\langle(\xi,\eta)\rangle\right)\circ\Lambda u,\Lambda u\right) \leq C||\sqrt{g}\log\langle(\xi,\eta)\rangle\circ\Lambda u||^2 + C||\Lambda u||^2 + C_{\delta_2}||u||_{H^{-N}}$

Proof. First note that $a_0'' + b_0''$ is of the same class as a_0'' , $\tilde{\mathcal{A}}_0$, so we can denote it as \tilde{a}_0 . Now we write

$$\begin{aligned} Re\left(\partial_x^{\alpha} g \tilde{a}_0 \log\langle(\xi,\eta)\rangle \circ \Lambda u, \Lambda u\right) &= Re(\left[\partial_x^{\alpha} g, \tilde{a}_0\right] \log\langle(\xi,\eta)\rangle \circ \Lambda u, \Lambda u) \\ &+ Re(\tilde{a}_0 \partial_x^{\alpha} g \log\langle(\xi,\eta)\rangle \circ \Lambda u, \Lambda u) \\ &\leq C||\Lambda u||_{-\varepsilon}^2 + ||\tilde{a}_0 \partial_x^{\alpha} g \log\langle(\xi,\eta)\rangle \circ \Lambda u||^2 + ||\Lambda u||^2 \\ &\leq C||\partial_x^{\alpha} g \log\langle(\xi,\eta)\rangle \circ \Lambda u||^2 + C||\Lambda u||^2 + C||u||_{H^{-N}}^2 \end{aligned}$$

where we used Cauchy-Scwartz, the estimate $||\tilde{a}_0|| \leq C$, and interpolation

$$||\Lambda u||_{-\varepsilon}^2 \le C ||\Lambda u||^2 + C ||u||_{H^{-N}}^2.$$

We now use the following Wirtinger-type inequality (see e.g. [MK86]): If $\phi \in C^2(U)$ with U open in \mathbb{R}^n , ϕ nonnegative, then for any compact subset $F \subset U$ there exists a constant C depending on $\|D^2\phi\|_{L^{\infty}(V)}$, with V open and $F \subset V \subseteq U$, and dist $(F, \partial V) > 0$ such that

$$\left|D\phi\left(x\right)\right|^{2} \le C\phi\left(x\right).$$

Applied to the function g on the support of u this gives

$$\left|\partial_x^{\alpha}g\right| \le C\sqrt{g}$$

for $|\alpha| = 1$, and therefore, since g = g(x), the estimate

$$||\partial_x^{\alpha} g \log \langle (\xi, \eta) \rangle \circ \Lambda u||^2 \le C ||\sqrt{g} \log \langle (\xi, \eta) \rangle \circ \Lambda u||^2.$$

Lemma 40. Let $\tilde{d}_0 = \tilde{d}_0(x, y, \eta) \in \mathcal{D}_0$. There exists $r^- \in S^-$ such that

$$\operatorname{Re}(g\tilde{d}_0\log^2\langle(\xi,\eta)\rangle u, u) \le C \|\sqrt{g}\log\langle(\xi,\eta)\rangle u\|^2 + (r^-u, u)$$

Proof. We can write

$$(g\tilde{d}_0 \log^2 \langle (\xi, \eta) \rangle \, u, u) = (g\tilde{d}_0 \log \langle (\xi, \eta) \rangle \, u, \log \langle (\xi, \eta) \rangle \, u) + ([g\tilde{d}_0, \log \langle (\xi, \eta) \rangle] \log \langle (\xi, \eta) \rangle \, u, u)$$

Since $\tilde{d}_0 = \tilde{d}_0(x, y, \eta)$ and $g = g(x)$ for the first term we have
$$|(g\tilde{d}_0 \log \langle (\xi, \eta) \rangle \, u, \log \langle (\xi, \eta) \rangle \, u)| = |(\tilde{d}_0 \sqrt{g} \log \langle (\xi, \eta) \rangle \, u, \sqrt{g} \log \langle (\xi, \eta) \rangle \, u)| \le C \|\sqrt{g} \log \langle (\xi, \eta) \rangle \, u\|^2$$

While for the second term it is clear that

le for the second term it is clear that

$$[g\tilde{d}_0, \log\langle(\xi, \eta)\rangle] = r^{-1}$$

where $r^- \in S^-$ which concludes the proof.

From the lemmas above we thus get

$$Re(gQ_2u,\Lambda u) + Re(Q_3u,\Lambda u) \le C \|\sqrt{g}\log\langle(\xi,\eta)\rangle\Lambda u\|^2 + C\|\Lambda u\|^2 + C_{\delta_2}\|u\|_{H^{-N}},$$

which finishes the proof of the second part of Proposition 15.

APPENDIX

Notation. The following multipliers will be frequently used, where as usual multipliers are defined via the Fourier transform $\mathcal{F}(m(D)f)(\xi) := m(\xi)\hat{f}(\xi)$. In particular for any $s \in \mathbb{R}$

$$\langle \xi \rangle^s := (e^2 + \xi^2)^{\frac{s}{2}}$$

Then $\langle (\xi, \eta) \rangle^s = \langle D \rangle^s \in S^s$.

For cut off functions we will use the following notation,

(88)
$$\phi \subset \subset \psi \in C_0^\infty(K)$$

to mean that $\psi \equiv 1$ on the support of ϕ , and $\phi \equiv 1$ on a typically predefined compact set.

Norms without subscript will be L^2 norms, i.e. $||f|| = ||f||_{L^2}$ and subscripts would mean H^s norms, i.e. $||f||_s := ||f||_{H^s} := \left(\int \langle \xi \rangle^{2s} |\hat{f}(\xi)|^2 d\xi \right)^{\frac{1}{2}}$.

Positivity of elliptic operator. First we establish positivity of a second order elliptic operator, which is used throughout the paper.

Lemma 41. Let L be degenerately elliptic, i.e. of the form (2). Then for any compact set $K \in \mathbb{R}^n$ there is a constant C_K , such that for all $u \in C_0^{\infty}(K)$

(89)
$$Re(Lu, u) \ge -C_K \|u\|^2$$

Proof. We consider terms one by one from the second order down to the second.

Let I_2 denote all second order terms and integrate by parts:

$$I_{2} := -\sum_{j,k} \int a_{jk}(x) \partial_{x_{j}} \partial_{x_{k}} u \cdot u dx$$
$$= \sum_{j,k} \int a_{jk}(x) \partial_{x_{j}} u \cdot \partial_{x_{k}} u dx + \sum_{j,k} \int \partial_{x_{k}} a_{jk}(x) \partial_{x_{j}} u \cdot u dx$$

Observe that $\partial_{x_k} u \cdot u = \frac{1}{2} \partial_{x_k} (u^2)$. With this knowledge in mind integrate the second term by parts

$$I_2 = \sum_{j,k} \int a_{jk}(x) \partial_{x_j} u \cdot \partial_{x_k} u dx - \frac{1}{2} \sum_{j,k} \int \partial_{x_j} \partial_{x_k} a_{jk}(x) u \cdot u dx$$

The non-negative definite property of $a_{jk} \ge 0$ implies the first term above is non-negative. While for the second term we apply the Hölder inequality:

$$I_2 \ge 0 - \frac{1}{2} \sum_{j,k} \|\partial_{x_j} \partial_{x_k} a_{jk}(x)\|_{L^{\infty}_x} \cdot \|u\|^2$$

By Holder inequality

(90)
$$|(a_0u, u)| \le C_K ||u||^2$$

The first and zero-th terms are treated similarly to $\int \partial_{x_k} a_{jk}(x) \partial_{x_j} u \cdot u dx$ and $\int \partial_{x_k} a_{jk}(x) \partial_{x_j} u \cdot u dx$ respectively.

Properties of pseudodifferential operators.

Proposition 42 (Adjoint Ψ DO). Let p be the symbol of the Ψ DO P. If P^* , i.e. $(Pu, v) = (u, P^*v), \forall u, v \in \mathscr{S}$, then the symbol p^* has the following asymptotic expansion

$$p^*(x,\xi) \sim \sum_{\alpha} \frac{i^{|\alpha|}}{\alpha!} \overline{\partial_{\xi}^{\alpha} \partial_{x}^{\alpha} p(x,\xi)}.$$

Proof. See e.g. [Kg81] Theorem 2.1.7.

Proposition 43 (Composition of Ψ DO). Let p be the symbol of the Ψ DO P, and q — the symbol of the Ψ DO Q. Then the symbol r of the composition $P \circ Q$ has the following asymptotic expansion

$$r(x,\xi) \sim \sum_{\alpha} \frac{(-i)^{|\alpha|}}{\alpha!} \partial_{\xi}^{\alpha} p(x,\xi) \partial_{x}^{\alpha} q(x,\xi).$$

Proof. See e.g. [Kg81] Theorem 2.1.7.

Lemma 44. Let P be a pseudodifferential operator of order m such that $P + P^*$ is of order m - l. Then for any $v \in H^m$ we have

(91)
$$Re(Pv,v) = \left(\frac{P+P^*}{2}v,v\right) \le C||v||_{\frac{m-l}{2}}^2$$

Proof. We can write

$$Re(Pv,v) = \frac{1}{2}\left((Pv,v) + \overline{(Pv,v)}\right) = \frac{1}{2}\left((Pv,v) + \overline{(v,P^*v)}\right) = \frac{1}{2}\left((Pv,v) + (P^*v,v)\right),$$

which establishes the first equality in (91). The inequality in (91) then follows from the boundedness of Ψ DO, see [Kg81].

Regularization S_{δ_5} .

Lemma 45. Let S_{δ_5} be defined by the symbol $s_{\delta_5} = \langle \delta_5 \cdot (\xi, \eta) \rangle^{-(N+s+3)}$. We then have

- (1) S_{δ_5} is an operator of order -N s 3 whose seminorms depend on δ_5 ;
- (2) S_{δ_5} is an operator of order 0 uniformly in δ_5 , i.e. $\limsup_{\delta_5 \to 0} |s_{\delta_5}| < \infty$;
- (3) the operator with the symbol $\frac{\partial_{\xi,\eta}^{\alpha} s_{\delta_5}}{s_{\delta_5}}$ for $|\alpha| = 1$ is uniformly of order -1. Inductively, same argument can be repeated for $|\alpha| > 1$.

Proof. (1) This follows immediately from the definition.

- (2) It is easy to check that $|s_{\delta_5}| \leq 1 \ \forall \xi, \eta$.
- (3) Differentiating we get

$$\frac{\partial_{\xi_k} s_{\delta_5}}{s_{\delta_5}} = \frac{\partial_{\xi_k} (1 + \delta_5^2 |\xi|^2 + \delta_5^2 |\eta|^2)^{\frac{-(N+s+3)}{2}}}{(1 + \delta_5^2 |\xi|^2 + \delta_5^2 |\eta|^2)^{\frac{-(N+s+3)}{2}}} = -(N+s+3) \frac{\delta_5^2 \xi_k}{(1 + \delta_5^2 |\xi|^2 + \delta_5^2 |\eta|^2)}$$

which is easily seen to be an operator of order -1 uniformly in δ_5 .

References

- [Chr01] Michael Christ, Hypoellipticity in the infinitely degenerate regime, Complex analysis and geometry (Columbus, OH, 1999), Ohio State Univ. Math. Res. Inst. Publ., vol. 9, de Gruyter, Berlin, 2001, pp. 59–84. MR 1912731 (2003e:35044)
- [Fed71] V. S. Fediĭ, A certain criterion for hypoellipticity, Mat. Sb. (N.S.) 85 (127) (1971), 18–48. MR 0287160 (44 #4367)
- [Hör55] Lars Hörmander, On the theory of general partial differential operators, Acta Math. 94 (1955), 161–248. MR 0076151 (17,853d)
- [Hör67] _____, Hypoelliptic second order differential equations, Acta Math. **119** (1967), 147–171. MR 0222474 (36 #5526)
- [Kg81] Hitoshi Kumano-go, Pseudodifferential operators, MIT Press, Cambridge, Mass., 1981, Translated from the Japanese by the author, Rémi Vaillancourt and Michihiro Nagase. MR MR666870 (84c:35113)
- [Koh98] Joseph J. Kohn, Hypoellipticity of some degenerate subelliptic operators, J. Funct. Anal. 159 (1998), no. 1, 203–216. MR 1654190 (99m:35035)

HYPOELLIPTICITY WITHOUT LOSS OF DERIVATIVES FOR FEDII'S TYPE OPERATORS 35

- [KR14] Lyudmila Korobenko and Cristian Rios, Hypoellipticity of certain infinitely degenerate second order operators, J. Math. Anal. Appl. 409 (2014), no. 1, 41–55. MR 3095016
- [KS84] Shigeo Kusuoka and Daniel Stroock, Applications of the Malliavin calculus. I, Stochastic analysis (Katata/Kyoto, 1982), North-Holland Math. Library, vol. 32, North-Holland, Amsterdam, 1984, pp. 271–306. MR 780762 (86k:60100a)
- [MK86] Vladimir G. Maz'ja and Alois Kufner, Variations on the theme of the inequality $(f')^2 \leq 2f \sup |f''|$, Manuscripta Math. 56 (1986), no. 1, 89–104. MR 846988
- [Mor87] Yoshinori Morimoto, A criterion for hypoellipticity of second order differential operators, Osaka J. Math. 24 (1987), no. 3, 651–675. MR 923880 (89b:35023)
- [OR73] O. A. Oleĭnik and E. V. Radkevič, Second order equations with nonnegative characteristic form, Plenum Press, New York-London, 1973, Translated from the Russian by Paul C. Fife. MR 0457908 (56 #16112)
- [Sch50] L. Schwartz, Théorie des distributions. Tome I, Actualités Sci. Ind., no. 1091 = Publ. Inst. Math. Univ. Strasbourg 9, Hermann & Cie., Paris, 1950. MR 0035918
- [Sch63] Laurent Schwartz, Some applications of the theory of distributions, Lectures on Modern Mathematics, Vol. I, Wiley, New York, 1963, pp. 23–58. MR 0178353

DEPARTMENT OF MATHEMATICAL SCIENCES, BINGHAMPTON UNIVERSITY, PO Box 6000, BINGHAMTON, NY13902-6000, USA

MATHEMATICS DEPARTMENT, REED COLLEGE, 3203 SOUTHEAST WOODSTOCK BOULEVARD, PORTLAND, OR 97202-8199, USA

UNIVERSITY OF CALGARY, DEPARTMENT OF MATHEMATICS AND STATISTICS, 2500 UNIVER-SITY DR. NW, CALGARY, AB, T2X 3B5, CANADA