

submitted to Uzbek Mathematical Journal

Direct and inverse source problems for two-term time-fractional diffusion equation with Hilfer derivative
Salakhitdinov M.S., Karimov E.T.

Abstract

In this paper, we investigate direct and inverse source problems for the diffusion equation with two-term generalized fractional derivative (Hilfer derivative) in a rectangular domain. Using spectral expansion method, we derive two-term fractional differential equation together with appropriate initial condition (Cauchy problem). Based on solution of that Cauchy problem, we represent solution of formulated problems as a combination of sinus and multinomial Mittag-Leffler function of two variables. Imposing certain conditions to the given data, we prove uniform convergence of certain infinite series.

MSC 2010: 35R11, 26A33

Keywords: Hilfer derivative, multi-term fractional diffusion equation, inverse source problem, multivariate Mittag-Leffler function.

1 Inverse source problem.

Consider the following diffusion equation with Hilfer derivatives

$$D_{0t}^{\alpha_1, \beta_1} u(t, x) + \mu D_{0t}^{\alpha_2, \beta_2} u(t, x) - u_{xx}(t, x) = g(x) \tag{1}$$

in a rectangular domain $\Omega = \{(t, x) : 0 < t < T, 0 < x < 1\}$. Here α_i, β_i ($i = 1, 2$), μ, T are given real numbers such that $T > 0, 0 < \alpha_2 < \alpha_1 < 1, 0 \leq \beta_i \leq 1$,

$$D_{0t}^{\alpha, \beta} f(t) = \left(I_{0t}^{\beta(n-\alpha)} \frac{d^n}{dt^n} \left(I_{0t}^{(1-\beta)(n-\alpha)} f \right) \right) (t), t > 0 \tag{2}$$

is a Hilfer fractional derivative of the order α and type β with $n-1 < \alpha \leq n \in \mathbb{N}$ (see [1]),

$$I_{0t}^\alpha f(t) = \frac{1}{\Gamma(\alpha)} \int_0^t (t-z)^{\alpha-1} f(z) dz, t > 0 \tag{3}$$

is the Riemann-Liouville fractional integral of order $\alpha > 0$ such that $I_{0t}^0 f(t) = f(t)$ (see, for example [2]).

We note that in case of $\beta = 0$, Hilfer derivative (2) coincides with the Riemann-Liouville derivative and in case of $\beta = 1$ with the Caputo derivative [1].

Problem 1 is to find a pair of functions $\{u(t, x), g(x)\}$ from the class of functions

$$W_1 = \{u(t, x) : u \in C_{-1}[0, T], I_{0t}^{(1-\beta_1)(1-\alpha_1)} u \in C_{-1}^1[0, T], u \in C[0, 1], u_{xx} \in C(0, 1); g(x) \in C[0, 1]\},$$

which satisfies equation (1) in Ω together with boundary conditions

$$u(t, 0) = u(t, 1) = 0, 0 \leq t \leq T \tag{4}$$

and initial conditions

$$\lim_{t \rightarrow +0} I_{0t}^{(1-\beta_i)(1-\alpha_i)} u(t, x) = 0, \quad 0 \leq x \leq 1 \quad (i = 1, 2), \quad (5)$$

and also over-determining condition

$$u(T, x) = \varphi(x), \quad 0 \leq x \leq 1. \quad (6)$$

Here $\varphi(x)$ is a given function such that $\varphi(0) = \varphi(1) = 0$.

We remind the definition of the space C_α^m .

Definition. A function $f(x)$, $x > 0$, is said to be in the space C_α^m , $m \in \mathbb{N}_0 = \mathbb{N} \cup \{0\}$, $\alpha \in \mathbb{R}$ if there exists a real number p , $p > \alpha$, such that $f^{(m)}(x) = x^p f_1(x)$ with a function $f_1 \in C[0, \infty)$, where $C_\alpha^0 = C_\alpha$.

Hilfer derivative has appeared in the theoretical modeling of a broadband dielectric relaxation spectroscopy for glasses [3]. Some properties and applications of this derivative were studied in [4-7]. We will use in this work result of the work [8], where explicit solution of the modified Cauchy problem for multi-term fractional differential equation with Hilfer derivatives found by operational method.

Separation of variables lead us to the following spectral problem

$$X''(x) - \lambda X(x) = 0, \quad X(0) = X(1) = 0,$$

whose eigenvalues are $\lambda_k = (k\pi)^2$, $k \in \mathbb{Z}$ and corresponding eigenfunctions $\{X_k = \sin k\pi x\}$. Since this system forms complete orthogonal basis, we can expand solution of the problem 1 by the following series:

$$u(t, x) = \sum_{k=1}^{\infty} U_k(t) \sin k\pi x, \quad 0 \leq t \leq T \quad (7)$$

$$g(x) = \sum_{k=1}^{\infty} g_k \sin k\pi x, \quad (8)$$

where

$$U_k(t) = 2 \int_0^1 u(t, x) \sin k\pi x dx, \quad k = 1, 2, \dots \quad (9)$$

$$g_k = 2 \int_0^1 g(x) \sin k\pi x dx, \quad k = 1, 2, \dots \quad (10)$$

Substituting (7)-(8) into (1), we get

$$D_{0t}^{\alpha_1, \beta_1} U_k(t) + \mu D_{0t}^{\alpha_2, \beta_2} U_k(t) + (k\pi)^2 U_k(t) = g_k. \quad (11)$$

Initial condition (5) gives us

$$\lim_{t \rightarrow +0} I_{0t}^{(1-\beta_i)(1-\alpha_i)} U_k(t) = 0 \quad (i = 1, 2). \quad (12)$$

According to [8], solution of (11) together with (12) has a form

$$U_k(t) = g_k \int_0^t z^{\alpha_1-1} E_{(\alpha_1-\alpha_2, \alpha_1), \alpha_1}(-\mu z^{\alpha_1-\alpha_2}, -(k\pi)^2 z^{\alpha_1}) dz, \quad (13)$$

where

$$E_{(\alpha-\beta,\alpha),\alpha}(x,y) = \sum_{n=0}^{\infty} \sum_{i=0}^n \frac{n!}{i!(n-i)!} \frac{x^i y^{n-i}}{\Gamma(\rho + \alpha n - \beta i)} \quad (14)$$

is a particular case (in two variables) of multi-variate Mittag-Leffler function [9] with $\alpha, \beta, \rho > 0$.

Applying the following formula (see, for example [10])

$$\int_0^t z^{\alpha-1} E_{(\alpha-\beta,\alpha),\alpha}(m_1 z^{\alpha-\beta}, m_2 z^{\alpha}) dz = t^{\alpha} E_{(\alpha-\beta,\alpha),\alpha+1}(m_1 t^{\alpha-\beta}, m_2 t^{\alpha})$$

from (13) we obtain

$$U_k(t) = g_k \cdot t^{\alpha_1-1} E_{(\alpha_1-\alpha_2,\alpha_1),\alpha_1+1}(-\mu t^{\alpha_1-\alpha_2}, -(k\pi)^2 t^{\alpha_1}). \quad (15)$$

In order to define g_k we use over-determined condition (6), which will take a form

$$U_k(T) = \varphi_k, \quad (16)$$

where

$$\varphi_k = 2 \int_0^1 \varphi(x) \sin k\pi x dx, \quad k = 1, 2, \dots$$

From (15)-(16) we deduce

$$g_k = \frac{\varphi_k}{T^{\alpha_1-1} E_{(\alpha_1-\alpha_2,\alpha_1),\alpha_1+1}(-\mu T^{\alpha_1-\alpha_2}, -(k\pi)^2 T^{\alpha_1})}$$

with

$$T^{\alpha_1-1} E_{(\alpha_1-\alpha_2,\alpha_1),\alpha_1+1}(-\mu T^{\alpha_1-\alpha_2}, -(k\pi)^2 T^{\alpha_1}) \neq 0. \quad (17)$$

Finally, $U_k(t)$ will have a form

$$U_k(t) = \frac{t^{\alpha_1-1} E_{(\alpha_1-\alpha_2,\alpha_1),\alpha_1+1}(-\mu t^{\alpha_1-\alpha_2}, -(k\pi)^2 t^{\alpha_1})}{T^{\alpha_1-1} E_{(\alpha_1-\alpha_2,\alpha_1),\alpha_1+1}(-\mu T^{\alpha_1-\alpha_2}, -(k\pi)^2 T^{\alpha_1})} \varphi_k. \quad (18)$$

Now, let us estimate $U_k(t)$. Since

$$\frac{t^{\alpha_1-1} E_{(\alpha_1-\alpha_2,\alpha_1),\alpha_1+1}(-\mu t^{\alpha_1-\alpha_2}, -(k\pi)^2 t^{\alpha_1})}{T^{\alpha_1-1} E_{(\alpha_1-\alpha_2,\alpha_1),\alpha_1+1}(-\mu T^{\alpha_1-\alpha_2}, -(k\pi)^2 T^{\alpha_1})} \leq C_1 \quad (C_1 = const > 0),$$

we have

$$|U_k(t)| \leq \frac{C_1}{(k\pi)^2} |\varphi_k^{(2)}|, \quad (19)$$

where

$$\varphi_k^{(2)} = -2 \int_0^1 \varphi''(x) \sin k\pi x dx.$$

We need more "strong estimate" for $U_k(t)$ in order to provide convergence of infinite series corresponding for $u_{xx}(t, x)$. Precisely,

$$|U_k(t)| \leq \frac{C_2}{(k\pi)^3} |\varphi_k^{(3)}|, \quad (20)$$

where C_2 is a positive constant and

$$\varphi_k^{(3)} = -2 \int_0^1 \varphi'''(x) \cos k\pi x dx.$$

We have to impose more condition to the given function $\varphi(x)$ in order to guarantee uniform convergence of the following series

$$u_{xx}(t, x) = \sum_{k=1}^{\infty} U_k(t) (k\pi)^2 \sin k\pi x. \quad (21)$$

In fact, considering (20), we have

$$|u_{xx}(t, x)| \leq \sum_{k=1}^{\infty} |U_k(t)| (k\pi)^2 \leq \sum_{k=1}^{\infty} \frac{C_2}{k\pi} |\varphi_k^{(3)}|.$$

If we use $2ab \leq a^2 + b^2$, we have

$$|u_{xx}(t, x)| \leq \sum_{k=1}^{\infty} \left(\frac{C_2^2}{4(k\pi)^2} + |\varphi_k^{(3)}|^2 \right).$$

Due to $\sum_{k=1}^{\infty} \frac{1}{(k\pi)^2} = 1/6$ and $\sum_{k=1}^{\infty} |\varphi_k^{(3)}|^2 \leq \|\varphi\|_{L_2(0,1)}^2$, we assume that

$$\varphi(x) \in C^2[0, 1], \varphi'''(x) \in L_2(0, 1), \varphi(0) = \varphi(1) = \varphi''(0) = \varphi''(1) = 0, \quad (22)$$

then by Weierstrass M-test theorem, we can conclude that series (21) uniformly converges.

Proof of the uniform convergence of series corresponding to the functions $u(t, x)$, $D_{0t}^{\alpha_i, \beta_i} u(t, x)$ ($i = 1, 2$) and $g(x)$ can be done by similar way considering (2), (3), (17), (19), but with less conditions to the given function $\varphi(x)$.

The uniqueness of the solution of the problem 1, can be obtained based on the completeness of the system $\{\sin k\pi x, k = 1, 2, \dots\}$ in L_2 . In fact, if we consider corresponding homogeneous problem, i.e. $\varphi(x) = 0$, from (18) we will get $U_k(t) \equiv 0$, which implies

$$\int_0^1 u(t, x) \sin k\pi x dx = 0, \quad 0 \leq t \leq T.$$

Due to the completeness of the system $\{\sin k\pi x, k = 1, 2, \dots\}$ in L_2 , we will get $u(t, x) \equiv 0$ in $\bar{\Omega}$.

We proved the following theorem:

Theorem 1. If conditions (17) and (22) are valid, then the problem 1 is uniquely solvable and solution is represented by

$$u(t, x) = \sum_{k=1}^{\infty} \frac{t^{\alpha_1-1} E_{(\alpha_1-\alpha_2, \alpha_1), \alpha_1+1}(-\mu t^{\alpha_1-\alpha_2}, -(k\pi)^2 t^{\alpha_1})}{T^{\alpha_1-1} E_{(\alpha_1-\alpha_2, \alpha_1), \alpha_1+1}(-\mu T^{\alpha_1-\alpha_2}, -(k\pi)^2 T^{\alpha_1})} \varphi_k \sin k\pi x,$$

$$g(x) = \sum_{k=1}^{\infty} \frac{1}{T^{\alpha_1-1} E_{(\alpha_1-\alpha_2, \alpha_1), \alpha_1+1}(-\mu T^{\alpha_1-\alpha_2}, -(k\pi)^2 T^{\alpha_1})} \varphi_k \sin k\pi x.$$

2 Direct problem.

Now let us consider the following direct problem.

Problem 2. To find a solution of the equation

$$D_{0t}^{\alpha_1, \beta_1} u(t, x) + \mu D_{0t}^{\alpha_2, \beta_2} u(t, x) - u_{xx}(t, x) = \bar{g}(t, x) \quad (23)$$

from the class of functions

$$W_2 = \{u(t, x) : u \in C_{-1}[0, T], I_{0t}^{(1-\beta_1)(1-\alpha_1)} u \in C_{-1}^1[0, T], \\ u \in C[0, 1], u_{xx} \in C(0, 1)\},$$

satisfying conditions (4) and (5).

Here $\bar{g}(t, x)$ is a given function.

Similarly as in the case of problem 1, we search solution in the form of

$$u(t, x) = \sum_{k=1}^{\infty} \bar{U}_k(t) \sin k\pi x, \quad 0 \leq t \leq T. \quad (24)$$

Substituting (24) into (23) we get

$$D_{0t}^{\alpha_1, \beta_1} \bar{U}_k(t) + \mu D_{0t}^{\alpha_2, \beta_2} \bar{U}_k(t) + (k\pi)^2 \bar{U}_k(t) = \bar{g}_k(t), \quad (25)$$

where

$$\bar{g}_k(t) = 2 \int_0^1 \bar{g}(t, x) \sin k\pi x dx.$$

Solution of (25) satisfying initial condition

$$\lim_{t \rightarrow +0} I_{0t}^{(1-\beta_1)(1-\alpha_1)} \bar{U}_k(t) = 0$$

has a form

$$\bar{U}_k(t) = \int_0^t z^{\alpha_1-1} E_{(\alpha_1-\alpha_2, \alpha_1), \alpha_1}(-\mu z^{\alpha_1-\alpha_2}, -(k\pi)^2 z^{\alpha_1}) \bar{g}_k(t-z) dz. \quad (26)$$

For the estimation of $\bar{U}_k(t)$ we use two different estimation of the function (14). Precisely, first of them is

$$|E_{(\alpha-\beta, \alpha), \alpha}(x, y)| \leq \frac{C_3}{1+|x|} \quad (27)$$

with $C_3 = \text{const} > 0$, which is proved in [9]. Another one is

$$|E_{(\alpha-\beta, \alpha), \alpha}(x, y)| \leq \frac{C_4}{1+|x+y|}, \quad (28)$$

with $C_4 = \text{constant} > 0$, which has the following additional condition to the fractional orders (see [10], lemma 1.3)

$$\Gamma(\rho + n(\alpha - \beta) + k\beta) > \Gamma(\rho + n(\alpha - \beta)), \quad n, k \in \mathbb{N}, \quad n \geq k. \quad (29)$$

If we use estimation (27), we get

$$|\bar{U}_k(t)| \leq \frac{C_5}{(k\pi)^2} \left| \bar{g}_k^{(2)}(t) \right|,$$

$$\bar{g}_k^{(2)}(t) = -2 \int_0^1 \frac{\partial^2 \bar{g}(t, x)}{\partial x^2} \sin k\pi x dx.$$

As we mentioned in previous case, we need another estimation for the $\bar{U}_k(t)$ in order to guarantee uniform convergence of infinite series corresponding to the function $u_{xx}(t, x)$, namely

$$|\bar{U}_k(t)| \leq \frac{C_6}{(k\pi)^3} \left| \bar{g}_k^{(3)}(t) \right|, \quad (30)$$

$$\bar{g}_k^{(3)}(t) = -2 \int_0^1 \frac{\partial^3 \bar{g}(t, x)}{\partial x^3} \cos k\pi x dx.$$

We impose the following conditions to the $\bar{g}(t, x)$:

$$\frac{\partial^2 \bar{g}(t, x)}{\partial x^2} \in C[0, 1], \quad \frac{\partial^3 \bar{g}(t, x)}{\partial x^3} \in L_2(0, 1),$$

$$\bar{g}(t, 0) = \bar{g}(t, 1) = 0, \quad \frac{\partial^2 \bar{g}(t, x)}{\partial x^2} \Big|_{x=0} = \frac{\partial^2 \bar{g}(t, x)}{\partial x^2} \Big|_{x=1} = 0 \quad (31)$$

in order to get

$$|u_{xx}(t, x)| \leq C_7 + \left\| \bar{g}_k^{(3)}(t) \right\|, \quad (32)$$

where C_7 is a positive constant.

Now, if we use estimation (28), we obtain

$$|\bar{U}_k(t)| \leq \frac{C_8}{(k\pi)^2} \left| \bar{g}_k^{(1)}(t) \right|,$$

$$\bar{g}_k^{(1)}(t) = 2 \int_0^1 \frac{\partial \bar{g}(t, x)}{\partial x} \cos k\pi x dx.$$

Consequently, in order to provide the uniform convergence of series

$$u_{xx}(t, x) = \sum_{k=1}^{\infty} \bar{U}_k(t) (k\pi)^2 \sin k\pi x \quad (33)$$

we impose the following condition to the given function $\bar{g}(t, x)$:

$$\frac{\partial \bar{g}(t, x)}{\partial x} \in C[0, 1], \quad \frac{\partial^2 \bar{g}(t, x)}{\partial x^2} \in L_2(0, 1), \quad \bar{g}(t, 0) = \bar{g}(t, 1) = 0, \quad (34)$$

which yields

$$|u_{xx}(t, x)| \leq C_9 + \left\| \bar{g}_k^{(2)}(t) \right\|.$$

Using Weierstrass M-test theorem, one can easily prove the uniform convergence of (33). The uniqueness of the solution for the problem 2 can be proved similarly to the proof of the problem 1.

Hence, we proved the following theorems:

Theorem 2. If condition (31) is valid, then problem 2 is uniquely solvable and solution is represented by

$$u(t, x) = \sum_{k=1}^{\infty} \sin k\pi x \int_0^t z^{\alpha_1-1} E_{(\alpha_1-\alpha_2, \alpha_1), \alpha_1}(-\mu z^{\alpha_1-\alpha_2}, -(k\pi)^2 z^{\alpha_1}) \bar{g}_k(t-z) dz. \quad (35)$$

Theorem 3. If condition (29) and (34) are valid, then problem 2 is uniquely solvable and solution is represented by (35).

References

1. R. Hilfer, Applications of Fractional Calculus in Physics. World Scientific, Singapore (2000).
2. I. Podlubny, Fractional Differential Equations, in: Mathematics in Science and Engineering, vol. 198, Acad. Press, 1999.
3. R. Hilfer, Experimental evidence for fractional time evolution in glass forming materials. Chem. Phys. 284 (2002), 399-408.
4. R. Hilfer, Y. Luchko, Z. Tomovski, Operational method for the solution of fractional differential equations with generalized Riemann-Liouville fractional derivatives. Fract. Calc. Appl. Anal. 12, No 3 (2009), 299-318.
5. K.M. Furati, M.D. Kassim, N.e.-Tatar, Existence and uniqueness for a problem involving Hilfer fractional derivative. Comp. Math. Appl. 64, No 6 (2012), 16161626.
6. K.M. Furati, O.S. Iyiola, M. Kirane, An inverse problem for a generalised fractional diffusion, Appl. Math. Comput. 249 (2014) 2431.
7. S.A.Malik, S.Aziz. An inverse source problem for a two parameter anomalous diffusion equation with nonlocal boundary conditions. Computer and Mathematics with Applications, 2017, <http://dx.doi.org/10.1016/j.camwa.2017.03.019>
8. M.-Ha. Kim, G.Chol-Ri, Chol O.H. Operational method for solving multi-term fractional differential equations with the generalized fractional derivatives. Fractional Calculus and Applied Analysis. 2014. vol.17. No 1, pp.79-95.
9. Luchko Y. and Gorenflo R. An Operational Method for Solving Fractional Differential Equations with the Caputo Derivatives, Acta Math.Vietnamica, N 2, 1999, 207-233.
10. E.T. Karimov, S. Kerbal, N. Al-Salti. Inverse source problem for multi-term fractional mixed type equation. M.Ruzhansky et al. (eds.), Advances in Real and Complex Analysis with Applications, Trends in Mathematics. 2017, pp. 289-301

Institute of Mathematics
named after V.I.Romanovsky