

# UNIQUENESS OF CONVEX ANCIENT SOLUTIONS TO MEAN CURVATURE FLOW IN $\mathbb{R}^3$

SIMON BRENDLE AND KYEONGSU CHOI

ABSTRACT. A well-known question of Perelman concerns the classification of noncompact ancient solutions to the Ricci flow in dimension 3 which have positive sectional curvature and are  $\kappa$ -noncollapsed. In this paper, we solve the analogous problem for mean curvature flow in  $\mathbb{R}^3$ , and prove that the rotationally symmetric bowl soliton is the only noncompact ancient solution of mean curvature flow in  $\mathbb{R}^3$  which is strictly convex and noncollapsed.

## 1. INTRODUCTION

This paper is concerned with the classification of ancient solutions to mean curvature flow. By definition, an ancient solution is a solution which is defined for  $t \in (-\infty, T]$  for some  $T$ . Ancient solutions play an important role in understanding singularity formation in geometric flows. For example, Perelman's famous Canonical Neighborhood Theorem [22] states that, for a solution to the Ricci flow in dimension 3, the high curvature regions are modeled on ancient solutions, which have nonnegative curvature and are  $\kappa$ -noncollapsed. Moreover, Perelman [22] proved that every noncompact ancient  $\kappa$ -solution in dimension 3 has the structure of a tube with a positively curved cap attached on one side.

Results of a similar nature hold for mean curvature flow, assuming that the initial hypersurface is mean convex and embedded (see [6],[12],[13],[16],[17],[25],[26]). In particular, Huisken-Sinestrari [16],[17] proved that, for any mean convex solution to mean curvature flow, the high curvature regions are almost convex. Under the stronger assumption of two-convexity, one can show that the mean curvature flow will only form neck-pinch singularities. Moreover, the flow can be continued beyond singularities by a surgery procedure similar in spirit to the one devised by Hamilton and Perelman for the Ricci flow (see [6],[13],[18]).

Our goal in this paper is to classify all convex ancient solutions to mean curvature flow in  $\mathbb{R}^3$  which are  $\alpha$ -noncollapsed in the sense of Sheng and Wang [23]. Recall that a mean convex surface is called  $\alpha$ -noncollapsed if, for each point  $x$  on the surface, there exists a ball of radius  $\frac{\alpha}{H}$  in ambient

---

The first author was supported by the National Science Foundation under grant DMS-1649174 and by the Simons Foundation. The second author was supported by the National Science Foundation under grant DMS-1811267. The authors gratefully acknowledge the hospitality of Tübingen University, where part of this work was carried out.

space, which lies inside the surface and which touches the surface at the given point  $x$ . Examples of noncollapsed ancient solutions include the shrinking cylinders and the rotationally symmetric bowl soliton (cf. [1]). In this paper, we show that these are the only ancient solutions which are noncompact, convex, and noncollapsed:

**Theorem 1.1.** *Let  $M_t$ ,  $t \in (-\infty, 0]$ , be a noncompact ancient solution of mean curvature flow in  $\mathbb{R}^3$  which is strictly convex and noncollapsed. Then  $M_t$  agrees with the bowl soliton, up to scaling and ambient isometries.*

By combining Theorem 1.1 with known results in the literature (see [12], Theorem 1.10, or [25],[26]), we can draw the following conclusion:

**Corollary 1.2.** *Consider an arbitrary closed, embedded, mean convex surface in  $\mathbb{R}^3$ , and evolve it by mean curvature flow. At the first singular time, the only possible blow-up limits are shrinking round spheres; shrinking round cylinders; and the translating bowl soliton.*

Let us indicate how Corollary 1.2 follows from Theorem 1.1. Suppose we evolve a closed, embedded, mean convex surface in  $\mathbb{R}^3$  by mean curvature flow. At the first singular time, every blow-up limit is an ancient solution which is weakly convex and noncollapsed (cf. [25], [26], or [12]). Indeed, every blow-up limit must be 1-noncollapsed by [4]. If a blow-up limit is compact and strictly convex, then the original flow eventually becomes convex, and therefore converges to a family of shrinking round spheres (cf. [14]). If a blow-up limit is noncompact and strictly convex, then that blow-up limit is the bowl soliton by Theorem 1.1. Finally, if a blow-up limit is not strictly convex, then it splits off a line, and is a family of shrinking round cylinders.

We next discuss the background of Theorem 1.1. Note that an ancient solution which is mean convex and two-sided noncollapsed is necessarily convex (cf. [12]), but it is not uniformly convex unless it is a family of shrinking spheres (see [19]). Theorem 1.1 can be viewed as a parabolic analogue of the classical Bernstein theorem, which classifies entire solutions to the minimal surface equation. Theorem 1.1 can be generalized to higher dimensions, if we assume that the solution is uniformly two-convex (see [5]).

Daskalopoulos, Hamilton, and Šešum obtained a complete classification of all compact ancient solutions to the Ricci flow in dimension 2 (cf. [9]). Moreover, they were able to classify all compact, convex ancient solutions to curve shortening flow in the plane (cf. [8]). Remarkably, these results do not require any noncollapsing assumptions. In a very important recent paper [2], Angenent, Daskalopoulos, and Šešum studied compact, convex ancient solutions to the mean curvature flow. Under suitable symmetry assumptions, they obtained precise asymptotic estimates for the solution as  $t \rightarrow -\infty$ .

A special case of ancient solutions are solitons; these are solutions that move in a self-similar fashion under the evolution. In a recent paper [3], the first author proved that every noncollapsed steady Ricci soliton in dimension

3 is rotationally symmetric, and hence is isometric to the Bryant soliton up to scaling. Using similar techniques, Haslhofer [11] subsequently proved that every noncollapsed, convex translating soliton for the mean curvature flow in  $\mathbb{R}^3$  is rotationally symmetric, and hence coincides with the bowl soliton up to scaling and ambient isometries. A related uniqueness result for the bowl soliton was proved in an important paper by Wang [24]; this relies on a completely different approach.

In Section 2, we study the asymptotic behavior of the flow as  $t \rightarrow -\infty$ . To that end, we write  $M_t = (-t)^{\frac{1}{2}} \bar{M}_{-\log(-t)}$ . As  $\tau \rightarrow -\infty$ , the rescaled surfaces  $\bar{M}_\tau$  converge in  $C_{\text{loc}}^\infty$  to a cylinder of radius  $\sqrt{2}$  with axis passing through the origin. More precisely, we show that  $\bar{M}_\tau$  can be approximated by a cylinder up to error terms of order  $O(e^{\frac{\tau}{2}})$ . As in [7], a major difficulty is the presence of a non-trivial eigenfunction for the linearized problem with eigenvalue 0. This eigenfunction corresponds to the second Hermite polynomial. Using the convexity of  $\bar{M}_\tau$  and the Brunn-Minkowski inequality, we show that this eigenfunction cannot become dominant as  $\tau \rightarrow -\infty$ .

In Section 3, we show that  $\liminf_{t \rightarrow -\infty} H_{\max}(t) > 0$ , where  $H_{\max}(t)$  denotes the supremum of the mean curvature at time  $t$ . To do that, we consider the complement  $M_t \setminus B_{8(-t)^{\frac{1}{2}}}(0)$ . This set has two connected components, one of which is compact and one of which is noncompact. By combining the asymptotic analysis in Section 2 with a barrier argument, we prove that the bounded connected component of  $M_t \setminus B_{8(-t)^{\frac{1}{2}}}(0)$  has diameter at least  $\sim (-t)$ . This implies that  $H_{\max}(t)$  is uniformly bounded from below as  $t \rightarrow -\infty$ .

In Section 4, we establish the Neck Improvement Theorem, which asserts that a neck becomes more symmetric under the evolution. This result does not require that the solution is ancient; it can be applied whenever we have a solution of mean curvature flow which is close to a cylinder on a sufficiently large parabolic neighborhood.

In Section 5, we iterate the Neck Improvement Theorem to conclude that  $M_t$  is rotationally symmetric.

Finally, in Section 6, we analyze ancient solutions with rotational symmetry, and complete the proof of Theorem 1.1.

## 2. ASYMPTOTIC ANALYSIS AS $t \rightarrow -\infty$

Let  $M_t$ ,  $t \in (-\infty, 0]$ , be a noncompact ancient solution of mean curvature flow in  $\mathbb{R}^3$  which is strictly convex and noncollapsed. We first recall some known results concerning the blowdown limit as  $t \rightarrow -\infty$ . Given any sequence  $t_j \rightarrow -\infty$ , we can find a subsequence with the property that the rescaled surfaces  $(-t_j)^{-\frac{1}{2}} M_{t_j}$  converge in  $C_{\text{loc}}^\infty$  to a smooth limit, which is either a plane, or a round sphere, or a cylinder of radius  $\sqrt{2}$  with axis passing through the origin (see [12], Theorem 1.11). Since the original flow  $M_t$  is noncompact, the limit cannot be a sphere. Moreover, it follows from

Huisken's monotonicity formula [15] that the limit cannot be a plane. Therefore, the limit must be a cylinder.

In the following, we consider the rescaled flow  $\bar{M}_\tau = e^{\frac{\tau}{2}} M_{-e^{-\tau}}$ . The surfaces  $\bar{M}_\tau$  move with velocity  $-(H - \frac{1}{2} \langle x, \nu \rangle) \nu$ . Given any sequence  $\tau_j \rightarrow -\infty$ , there exists a subsequence with the property that the surfaces  $\bar{M}_{\tau_j}$  converges in  $C_{\text{loc}}^\infty$  to a cylinder of radius  $\sqrt{2}$  with axis passing through the origin. To fix notation, we denote by  $\Sigma = \{x_1^2 + x_2^2 = 2\}$  the cylinder of radius  $\sqrt{2}$  whose axis coincides with the  $x_3$ -axis.

**Proposition 2.1.** *For each  $\tau$ , we have*

$$\int_{\bar{M}_\tau} e^{-\frac{|x|^2}{4}} \leq \int_{\Sigma} e^{-\frac{|x|^2}{4}}.$$

**Proof.** Every convex surface is star-shaped, hence outward-minimizing by a standard calibration argument. This implies  $\text{area}(\bar{M}_\tau \cap B_r(p)) \leq Cr^2$  for all  $p \in \mathbb{R}^3$  and all  $r > 0$ . We next consider an arbitrary sequence  $\tau_j \rightarrow -\infty$ . After passing to a subsequence, the surfaces  $\bar{M}_{\tau_j}$  converge in  $C_{\text{loc}}^\infty$  to a cylinder of radius  $\sqrt{2}$  with axis passing through the origin. This gives

$$\int_{\bar{M}_{\tau_j}} e^{-\frac{|x|^2}{4}} \rightarrow \int_{\Sigma} e^{-\frac{|x|^2}{4}}$$

as  $j \rightarrow \infty$ . On the other hand, Huisken's monotonicity formula [15] implies that the function

$$\tau \mapsto \int_{\bar{M}_\tau} e^{-\frac{|x|^2}{4}}$$

is monotone decreasing in  $\tau$ . From this, the assertion follows.

In view of the discussion above, there exists a smooth function  $S(\tau)$  taking values in  $SO(3)$  such that the rotated surfaces  $\tilde{M}_\tau = S(\tau)\bar{M}_\tau$  converge to the cylinder  $\Sigma$  in  $C_{\text{loc}}^\infty$ . Hence, we can find a function  $\rho(\tau)$  with the following properties:

- $\lim_{\tau \rightarrow -\infty} \rho(\tau) = \infty$ .
- $-\rho(\tau) \leq \rho'(\tau) \leq 0$ .
- We may write the surface  $\tilde{M}_\tau$  as a graph of some function  $u(\cdot, \tau)$  over  $\Sigma \cap B_{2\rho(\tau)}(0)$ , so that

$$\{x + u(x, \tau)\nu_\Sigma(x) : x \in \Sigma \cap B_{2\rho(\tau)}(0)\} \subset \tilde{M}_\tau,$$

where  $\nu_\Sigma$  denotes the unit normal to  $\Sigma$  and  $\|u(\cdot, \tau)\|_{C^4(\Sigma \cap B_{2\rho(\tau)}(0))} \leq \rho(\tau)^{-4}$ .

In the next step, we fine-tune the choice of  $S(\tau)$ . To that end, we fix a smooth cutoff function  $\varphi \geq 0$  satisfying  $\varphi = 1$  on  $[-\frac{1}{2}, \frac{1}{2}]$  and  $\varphi = 0$  outside  $[-\frac{2}{3}, \frac{2}{3}]$ . By the implicit function theorem, we can choose  $S(\tau) \in SO(3)$  such

that  $u(x, \tau)$  satisfies the orthogonality relations

$$\int_{\Sigma \cap B_{\rho(\tau)}(0)} e^{-\frac{|x|^2}{4}} \langle Ax, \nu_\Sigma \rangle u(x, \tau) \varphi\left(\frac{x_3}{\rho(\tau)}\right) = 0$$

for every matrix  $A \in so(3)$ . Finally, we can arrange that the matrix  $A(\tau) := S'(\tau)S(\tau)^{-1} \in so(3)$  satisfies  $A(\tau)_{12} = 0$ . (Otherwise, we replace  $S(\tau)$  by  $R(\tau)S(\tau)$ , where  $R(\tau)$  is a rotation in the  $x_1x_2$ -plane. This does not affect the orthogonality relations above.)

Our next two results are straightforward adaptations of powerful estimates in [2]. These estimates will play a key role in the subsequent arguments. Recall the foliation by self-shrinkers  $\Sigma_a$  and  $\tilde{\Sigma}_b$  given in [2] (see also [20]). Let  $\nu_{\text{fol}}$  denote the unit normal vector to this foliation. As explained in [2], the union of all the leaves in this foliation contains a neighborhood of the cylinder  $\Sigma$ , and opens up like a cone at infinity. In particular, the union of all the leaves in the foliation contains the cylinder  $\Sigma$  and the surface  $\tilde{M}_\tau$  (for  $-\tau$  sufficiently large), as well as the region  $\Delta_\tau$  which lies in between  $\Sigma$  and  $\tilde{M}_\tau$ .

**Proposition 2.2** (cf. [2], Lemma 4.10). *There exists a constant  $L_0$  such that for all  $L \in [L_0, \rho(\tau)]$*

$$\int_{\tilde{M}_\tau \cap \{|x_3| \geq L\}} e^{-\frac{|x|^2}{4}} - \int_{\Sigma \cap \{|x_3| \geq L\}} e^{-\frac{|x|^2}{4}} \geq - \int_{\Delta_\tau \cap \{|x_3| = L\}} e^{-\frac{x^2}{4}} |\langle \omega, \nu_{\text{fol}} \rangle|,$$

where  $\omega = (0, 0, 1)$  denotes the vertical unit vector in  $\mathbb{R}^3$ .

**Proof.** Since each leaf of the foliation is a self-similar shrinker, the normal vector  $\nu_{\text{fol}}$  satisfies  $\text{div}(e^{-\frac{|x|^2}{4}} \nu_{\text{fol}}) = 0$ . Note that  $\nu_{\text{fol}} = \nu_\Sigma$  at each point on the cylinder  $\Sigma$ . Using the divergence theorem, we obtain

$$\begin{aligned} & \int_{\tilde{M}_\tau \cap \{L \leq |x_3| \leq z\}} e^{-\frac{|x|^2}{4}} \langle \nu, \nu_{\text{fol}} \rangle - \int_{\Sigma \cap \{L \leq |x_3| \leq z\}} e^{-\frac{|x|^2}{4}} \\ & \geq - \int_{\Delta_\tau \cap \{|x_3| = L\}} e^{-\frac{x^2}{4}} |\langle \omega, \nu_{\text{fol}} \rangle| - \int_{\Delta_\tau \cap \{|x_3| = z\}} e^{-\frac{x^2}{4}} |\langle \omega, \nu_{\text{fol}} \rangle|. \end{aligned}$$

We know  $\langle \nu, \nu_{\text{fol}} \rangle \leq 1$  on  $\tilde{M}_\tau$ . The convexity of  $\tilde{M}_\tau$  implies that the area of  $\Delta_\tau \cap \{|x_3| = z\}$  is at most  $Cz^2$ . Hence, passing to the limit as  $z \rightarrow \infty$  gives

$$\int_{\tilde{M}_\tau \cap \{|x_3| \geq L\}} e^{-\frac{|x|^2}{4}} - \int_{\Sigma \cap \{|x_3| \geq L\}} e^{-\frac{|x|^2}{4}} \geq - \int_{\Delta_\tau \cap \{|x_3| = L\}} e^{-\frac{x^2}{4}} |\langle \omega, \nu_{\text{fol}} \rangle|,$$

as desired.

**Proposition 2.3** (cf. [2], Lemma 4.7). *There exists a constant  $L_0$  such that*

$$\int_{\Sigma \cap \{|x_3| \leq L\}} e^{-\frac{|x|^2}{4}} |\nabla^\Sigma u(x, \tau)|^2 \leq C \int_{\Sigma \cap \{|x_3| \leq \frac{L}{2}\}} e^{-\frac{|x|^2}{4}} u(x, \tau)^2$$

and

$$\int_{\Sigma \cap \{\frac{L}{2} \leq |x_3| \leq L\}} e^{-\frac{|x|^2}{4}} u(x, \tau)^2 \leq CL^{-2} \int_{\Sigma \cap \{|x_3| \leq \frac{L}{2}\}} e^{-\frac{|x|^2}{4}} u(x, \tau)^2$$

for all  $L \in [L_0, \rho(\tau)]$ .

**Proof.** Lemma 4.11 in [2] implies that  $|\langle \omega, \nu_{\text{fol}} \rangle| \leq CL^{-1} |x_1^2 + x_2^2 - 2|$  for each point  $x \in \Delta_\tau \cap \{|x_3| = L\}$ . This gives

$$\begin{aligned} \int_{\Delta_\tau \cap \{|x_3| = L\}} e^{-\frac{x^2}{4}} |\langle \omega, \nu_{\text{fol}} \rangle| &\leq CL^{-1} \int_{\Delta_\tau \cap \{|x_3| = L\}} e^{-\frac{x^2}{4}} |x_1^2 + x_2^2 - 2| \\ &\leq CL^{-1} \int_{\Sigma \cap \{|x_3| = L\}} e^{-\frac{|x|^2}{4}} u^2. \end{aligned}$$

Using Proposition 2.2, we obtain

$$\int_{\tilde{M}_\tau \cap \{|x_3| \geq L\}} e^{-\frac{|x|^2}{4}} - \int_{\Sigma \cap \{|x_3| \geq L\}} e^{-\frac{|x|^2}{4}} \geq -CL^{-1} \int_{\Sigma \cap \{|x_3| = L\}} e^{-\frac{|x|^2}{4}} u^2$$

(compare [2], equation (4.33)). On the other hand,

$$\begin{aligned} &\int_{\tilde{M}_\tau \cap \{|x_3| \leq L\}} e^{-\frac{|x|^2}{4}} - \int_{\Sigma \cap \{|x_3| \leq L\}} e^{-\frac{|x|^2}{4}} \\ &= \int_{-L}^L \int_0^{2\pi} e^{-\frac{z^2}{4}} \left[ e^{-\frac{(\sqrt{2}+u)^2}{4}} \sqrt{(\sqrt{2}+u)^2 \left(1 + \left(\frac{\partial u}{\partial z}\right)^2\right) + \left(\frac{\partial u}{\partial \theta}\right)^2} - e^{-\frac{1}{2}} \sqrt{2} \right] d\theta dz. \end{aligned}$$

Since  $L \leq \rho(\tau)$ , we have  $|u| + \left|\frac{\partial u}{\partial z}\right| + \left|\frac{\partial u}{\partial \theta}\right| \leq o(1)$  for  $|x_3| \leq L$ . This gives

$$\begin{aligned} &\int_{\tilde{M}_\tau \cap \{|x_3| \leq L\}} e^{-\frac{|x|^2}{4}} - \int_{\Sigma \cap \{|x_3| \leq L\}} e^{-\frac{|x|^2}{4}} \\ &\geq \int_{-L}^L \int_0^{2\pi} e^{-\frac{z^2}{4}} \left[ e^{-\frac{(\sqrt{2}+u)^2}{4}} (\sqrt{2}+u) - e^{-\frac{1}{2}} \sqrt{2} + \frac{1}{C} |\nabla^\Sigma u|^2 \right] d\theta dz \\ &\geq \int_{-L}^L \int_0^{2\pi} e^{-\frac{z^2}{4}} \left[ -Cu^2 + \frac{1}{C} |\nabla^\Sigma u|^2 \right] d\theta dz \end{aligned}$$

where  $C > 0$  is a large numerical constant. Putting these facts together, we obtain

$$\begin{aligned} \int_{\tilde{M}_\tau} e^{-\frac{|x|^2}{4}} - \int_{\Sigma} e^{-\frac{|x|^2}{4}} &\geq \int_{\Sigma \cap \{|x_3| \leq L\}} e^{-\frac{z^2}{4}} \left[ -Cu^2 + \frac{1}{C} |\nabla^\Sigma u|^2 \right] \\ &\quad - CL^{-1} \int_{\Sigma \cap \{|x_3| = L\}} e^{-\frac{|x|^2}{4}} u^2. \end{aligned}$$

Using Proposition 2.1, we conclude that

$$\begin{aligned} &\int_{\Sigma \cap \{|x_3| \leq L\}} e^{-\frac{|x|^2}{4}} |\nabla^\Sigma u|^2 \\ &\leq C \int_{\Sigma \cap \{|x_3| \leq L\}} e^{-\frac{|x|^2}{4}} u^2 + CL^{-1} \int_{\Sigma \cap \{|x_3| = L\}} e^{-\frac{|x|^2}{4}} u^2. \end{aligned}$$

The divergence theorem gives

$$\begin{aligned} L \int_{\Sigma \cap \{|x_3|=L\}} e^{-\frac{|x|^2}{4}} u^2 &= \int_{\Sigma \cap \{|x_3| \leq L\}} \operatorname{div}_\Sigma (e^{-\frac{|x|^2}{4}} u^2 x^{\tan}) \\ &= \int_{\Sigma \cap \{|x_3| \leq L\}} e^{-\frac{|x|^2}{4}} \left( u^2 - \frac{1}{2} x_3^2 u^2 + 2u \langle x^{\tan}, \nabla^\Sigma u \rangle \right) \\ &\leq \int_{\Sigma \cap \{|x_3| \leq L\}} e^{-\frac{|x|^2}{4}} \left( u^2 - \frac{1}{4} x_3^2 u^2 + 4 |\nabla^\Sigma u|^2 \right), \end{aligned}$$

hence

$$\begin{aligned} &L^2 \int_{\Sigma \cap \{|x_3| \leq L\}} e^{-\frac{|x|^2}{4}} u^2 + L \int_{\Sigma \cap \{|x_3|=L\}} e^{-\frac{|x|^2}{4}} u^2 \\ &\leq C \int_{\Sigma \cap \{|x_3| \leq L\}} e^{-\frac{|x|^2}{4}} |\nabla^\Sigma u|^2 + CL^2 \int_{\Sigma \cap \{|x_3| \leq \frac{L}{2}\}} e^{-\frac{|x|^2}{4}} u^2. \end{aligned}$$

Putting these facts together, we conclude that

$$\begin{aligned} &\int_{\Sigma \cap \{|x_3| \leq L\}} e^{-\frac{|x|^2}{4}} |\nabla^\Sigma u|^2 \\ &\leq CL^{-2} \int_{\Sigma \cap \{|x_3| \leq L\}} e^{-\frac{|x|^2}{4}} |\nabla^\Sigma u|^2 + C \int_{\Sigma \cap \{|x_3| \leq \frac{L}{2}\}} e^{-\frac{|x|^2}{4}} u^2. \end{aligned}$$

If  $L$  is sufficiently large, the first term on the right hand side can be absorbed into the left hand side. This gives

$$\int_{\Sigma \cap \{|x_3| \leq L\}} e^{-\frac{|x|^2}{4}} |\nabla^\Sigma u|^2 \leq C \int_{\Sigma \cap \{|x_3| \leq \frac{L}{2}\}} e^{-\frac{|x|^2}{4}} u^2.$$

This proves the first statement. Using the inequality

$$0 \leq \int_{\Sigma \cap \{|x_3| \leq L\}} e^{-\frac{|x|^2}{4}} \left( u^2 - \frac{1}{4} x_3^2 u^2 + 4 |\nabla^\Sigma u|^2 \right),$$

the second statement follows.

Let us denote by  $\mathcal{H}$  the space of all functions  $f$  on  $\Sigma$  such that

$$\|f\|_{\mathcal{H}}^2 = \int_{\Sigma} e^{-\frac{|x|^2}{4}} f^2 < \infty.$$

We define an operator  $\mathcal{L}$  on the cylinder  $\Sigma$  by

$$\mathcal{L}f = \Delta_\Sigma f - \frac{1}{2} \langle x^{\tan}, \nabla^\Sigma f \rangle + f.$$

In coordinates,  $\mathcal{L}$  takes the form

$$\mathcal{L}f = \frac{\partial^2}{\partial z^2} f + \frac{1}{2} \frac{\partial^2}{\partial \theta^2} f - \frac{1}{2} z \frac{\partial}{\partial z} f + f.$$

The eigenfunctions of  $\mathcal{L}$  are of the form  $H_n\left(\frac{z}{2}\right) \cos(m\theta)$  and  $H_n\left(\frac{z}{2}\right) \sin(m\theta)$ , where  $m$  and  $n$  are nonnegative integers and  $H_n$  denotes the Hermite polynomial of degree  $n$ . The corresponding eigenvalues are given by  $1 - \frac{n+m^2}{2}$ .

Thus, there are four eigenfunctions that correspond to positive eigenvalues of  $\mathcal{L}$ , and these are given by  $1, z, \cos \theta, \sin \theta$  up to scaling. The span of these eigenfunctions will be denoted by  $\mathcal{H}_+$ . Moreover, there are three eigenfunctions of  $\mathcal{L}$  with eigenvalue 0, and these are given by  $z^2 - 2, z \cos \theta, z \sin \theta$  up to scaling. The span of these eigenfunctions will be denoted by  $\mathcal{H}_0$ . The span of all other eigenfunctions will be denoted by  $\mathcal{H}_-$ . Clearly,

$$\begin{aligned} \langle \mathcal{L}f, f \rangle_{\mathcal{H}} &\geq \frac{1}{2} \|f\|_{\mathcal{H}}^2 && \text{for } f \in \mathcal{H}_+, \\ \langle \mathcal{L}f, f \rangle_{\mathcal{H}} &= 0 && \text{for } f \in \mathcal{H}_0, \\ \langle \mathcal{L}f, f \rangle_{\mathcal{H}} &\leq -\frac{1}{2} \|f\|_{\mathcal{H}}^2 && \text{for } f \in \mathcal{H}_-. \end{aligned}$$

**Lemma 2.4.** *The function  $u(x, \tau)$  satisfies*

$$\frac{\partial}{\partial \tau} u = \mathcal{L}u + E + \langle A(\tau)x, \nu_{\Sigma} \rangle,$$

where  $E$  satisfies the pointwise estimate  $|E| \leq O(\rho(\tau)^{-1})(|u| + |\nabla^{\Sigma}u| + |A(\tau)|)$ .

**Proof.** Recall that the rescaled surfaces  $\bar{M}_{\tau}$  move with velocity  $-(H - \frac{1}{2}\langle x, \nu \rangle)\nu$ . Hence, the rotated surfaces  $\tilde{M}_{\tau} = S(\tau)\bar{M}_{\tau}$  move with velocity  $-(H - \frac{1}{2}\langle x, \nu \rangle - \langle A(\tau)x, \nu \rangle)\nu$ , where  $x \in \tilde{M}_{\tau}$ . Therefore, the function  $u(x, \tau)$  satisfies the equation

$$\begin{aligned} \frac{\partial}{\partial \tau} u &= -\frac{1}{\langle \nu_{\Sigma}, \nu(x + u\nu_{\Sigma}) \rangle} H(x + u\nu_{\Sigma}) \\ &\quad + \frac{1}{2\langle \nu_{\Sigma}, \nu(x + u\nu_{\Sigma}) \rangle} \langle x + u\nu_{\Sigma}, \nu(x + u\nu_{\Sigma}) \rangle \\ &\quad + \frac{1}{\langle \nu_{\Sigma}, \nu(x + u\nu_{\Sigma}) \rangle} \langle A(\tau)(x + u\nu_{\Sigma}), \nu(x + u\nu_{\Sigma}) \rangle \end{aligned}$$

for  $x \in \Sigma$ . By assumption,  $\|u\|_{C^4(\Sigma \cap B_{2\rho(\tau)}(0))} \leq O(\rho(\tau)^{-2})$ . This gives

$$|\nu(x + u\nu_{\Sigma}) - \nu_{\Sigma} + \nabla^{\Sigma}u| \leq O(\rho(\tau)^{-2})(|u| + |\nabla^{\Sigma}u|)$$

and

$$\left| H(x + u\nu_{\Sigma}) + \Delta_{\Sigma}u + \frac{1}{2}u \right| \leq O(\rho(\tau)^{-2})(|u| + |\nabla^{\Sigma}u|).$$

Putting these facts together, we obtain

$$\frac{\partial}{\partial \tau} u = \mathcal{L}u + E + \langle A(\tau)x, \nu_{\Sigma} \rangle,$$

where  $E$  satisfies the pointwise estimate  $|E| \leq O(\rho(\tau)^{-1})(|u| + |\nabla^{\Sigma}u| + |A(\tau)|)$ .

**Lemma 2.5.** *The function  $\hat{u}(x, \tau) = u(x, \tau) \varphi\left(\frac{x_3}{\rho(\tau)}\right)$  satisfies*

$$\frac{\partial}{\partial \tau} \hat{u} = \mathcal{L}\hat{u} + \hat{E} + \langle A(\tau)x, \nu_{\Sigma} \rangle \varphi\left(\frac{x_3}{\rho(\tau)}\right),$$

where  $\hat{E}$  satisfies  $\|\hat{E}\|_{\mathcal{H}} \leq O(\rho(\tau)^{-1}) (\|\hat{u}\|_{\mathcal{H}} + |A(\tau)|)$ .

**Proof.** We compute

$$\frac{\partial}{\partial \tau} \hat{u} = \mathcal{L}\hat{u} + \hat{E} + \langle A(\tau)x, \nu_{\Sigma} \rangle \varphi\left(\frac{x_3}{\rho(\tau)}\right)$$

where

$$\begin{aligned} \hat{E} &= E \varphi\left(\frac{x_3}{\rho(\tau)}\right) - \frac{2}{\rho(\tau)} \frac{\partial u}{\partial z} \varphi'\left(\frac{x_3}{\rho(\tau)}\right) - \frac{1}{\rho(\tau)^2} u \varphi''\left(\frac{x_3}{\rho(\tau)}\right) \\ &\quad + \frac{x_3}{2\rho(\tau)} u \varphi'\left(\frac{x_3}{\rho(\tau)}\right) - \frac{x_3 \rho'(\tau)}{\rho(\tau)^2} u \varphi'\left(\frac{x_3}{\rho(\tau)}\right). \end{aligned}$$

Using Lemma 2.4, we deduce that

$$|\hat{E}| \leq O(\rho(\tau)^{-1}) (|u| + |\nabla^{\Sigma} u| + |A(\tau)|)$$

for  $|x_3| \leq \frac{\rho(\tau)}{2}$ . Moreover, since  $|\rho'(\tau)| \leq \rho(\tau)$ , we obtain

$$|\hat{E}| \leq O(1) |u| + O(\rho(\tau)^{-1}) (|\nabla^{\Sigma} u| + |A(\tau)|)$$

for  $\frac{\rho(\tau)}{2} \leq |x_3| \leq \rho(\tau)$ . Using Proposition 2.3, we conclude that

$$\begin{aligned} \int_{\Sigma} e^{-\frac{|x|^2}{4}} |\hat{E}|^2 &\leq O(\rho(\tau)^{-2}) \int_{\Sigma \cap \{|x_3| \leq \frac{\rho(\tau)}{2}\}} e^{-\frac{|x|^2}{4}} u^2 \\ &\quad + O(1) \int_{\Sigma \cap \{\frac{\rho(\tau)}{2} \leq |x_3| \leq \rho(\tau)\}} e^{-\frac{|x|^2}{4}} u^2 \\ &\quad + O(\rho(\tau)^{-2}) \int_{\Sigma \cap \{|x_3| \leq \rho(\tau)\}} e^{-\frac{|x|^2}{4}} |\nabla^{\Sigma} u|^2 \\ &\quad + O(\rho(\tau)^{-2}) |A(\tau)|^2 \\ &\leq O(\rho(\tau)^{-2}) \int_{\Sigma \cap \{|x_3| \leq \frac{\rho(\tau)}{2}\}} e^{-\frac{|x|^2}{4}} u^2 \\ &\quad + O(\rho(\tau)^{-2}) |A(\tau)|^2 \\ &\leq O(\rho(\tau)^{-2}) \int_{\Sigma} e^{-\frac{|x|^2}{4}} \hat{u}^2 \\ &\quad + O(\rho(\tau)^{-2}) |A(\tau)|^2. \end{aligned}$$

Thus,  $\|\hat{E}\|_{\mathcal{H}} \leq O(\rho(\tau)^{-1}) \|\hat{u}\|_{\mathcal{H}} + O(\rho(\tau)^{-1}) |A(\tau)|$ , as claimed.

**Lemma 2.6.** *We have  $|A(\tau)| \leq O(\rho(\tau)^{-1}) \|u\|_{\mathcal{H}}$  and*

$$\left\| \frac{\partial}{\partial \tau} \hat{u} - \mathcal{L}\hat{u} \right\|_{\mathcal{H}} \leq O(\rho(\tau)^{-1}) \|\hat{u}\|_{\mathcal{H}}.$$

**Proof.** The orthogonality relations imply that  $\hat{u}$  is orthogonal to  $\langle Ax, \nu_{\Sigma} \rangle$  for every  $A \in so(3)$ . Since this is true for each  $\tau$ , it follows that  $\frac{\partial}{\partial \tau} \hat{u}$  is orthogonal to  $\langle Ax, \nu_{\Sigma} \rangle$  for every  $A \in so(3)$ . Moreover, since the function  $\langle Ax, \nu_{\Sigma} \rangle$  is an eigenfunction of  $\mathcal{L}$  with eigenvalue 0, we deduce that  $\mathcal{L}\hat{u}$  is

orthogonal to  $\langle Ax, \nu_\Sigma \rangle$  for every  $A \in so(3)$ . Consequently,  $\frac{\partial}{\partial \tau} \hat{u} - \mathcal{L} \hat{u}$  is orthogonal to  $\langle Ax, \nu_\Sigma \rangle$  for every  $A \in so(3)$ . Therefore,  $\hat{E} + \langle A(\tau)x, \nu_\Sigma \rangle \varphi\left(\frac{x_3}{\rho(\tau)}\right)$  is orthogonal to  $\langle Ax, \nu_\Sigma \rangle$  for every  $A \in so(3)$ . In particular,

$$\int_{\Sigma} e^{-\frac{|x|^2}{4}} \left( \hat{E} + \langle A(\tau)x, \nu_\Sigma \rangle \varphi\left(\frac{x_3}{\rho(\tau)}\right) \right) \langle A(\tau)x, \nu_\Sigma \rangle = 0.$$

Using the fact that  $A(\tau)_{12} = 0$ , we obtain

$$\begin{aligned} |A(\tau)|^2 &\leq O(1) \int_{\Sigma} e^{-\frac{|x|^2}{4}} \langle A(\tau)x, \nu_\Sigma \rangle^2 \varphi\left(\frac{x_3}{\rho(\tau)}\right) \\ &\leq O(1) \int_{\Sigma} e^{-\frac{|x|^2}{4}} |\hat{E}| |\langle A(\tau)x, \nu_\Sigma \rangle| \\ &\leq O(1) \|\hat{E}\|_{\mathcal{H}} |A(\tau)| \\ &\leq O(\rho(\tau)^{-1}) (\|\hat{u}\|_{\mathcal{H}} + |A(\tau)|) |A(\tau)|, \end{aligned}$$

where in the last step we have used Lemma 2.5. Consequently,  $|A(\tau)| \leq O(\rho(\tau)^{-1}) \|\hat{u}\|_{\mathcal{H}}$ . Using Lemma 2.5, we obtain

$$\left\| \frac{\partial}{\partial \tau} \hat{u} - \mathcal{L} \hat{u} \right\|_{\mathcal{H}} \leq \|\hat{E}\|_{\mathcal{H}} + O(1) |A(\tau)| \leq O(\rho(\tau)^{-1}) \|\hat{u}\|_{\mathcal{H}},$$

as claimed.

We now define

$$\begin{aligned} U_+(\tau) &:= \|P_+ \hat{u}(\cdot, \tau)\|_{\mathcal{H}}^2, \\ U_0(\tau) &:= \|P_0 \hat{u}(\cdot, \tau)\|_{\mathcal{H}}^2, \\ U_-(\tau) &:= \|P_- \hat{u}(\cdot, \tau)\|_{\mathcal{H}}^2, \end{aligned}$$

where  $P_+, P_0, P_-$  denote the orthogonal projections to  $\mathcal{H}_+, \mathcal{H}_0, \mathcal{H}_-$ , respectively. Using Lemma 2.6, we obtain

$$\begin{aligned} \frac{d}{d\tau} U_+(\tau) &\geq U_+(\tau) - O(\rho(\tau)^{-1}) (U_+(\tau) + U_0(\tau) + U_-(\tau)), \\ \left| \frac{d}{d\tau} U_0(\tau) \right| &\leq O(\rho(\tau)^{-1}) (U_+(\tau) + U_0(\tau) + U_-(\tau)), \\ \frac{d}{d\tau} U_-(\tau) &\leq -U_-(\tau) + O(\rho(\tau)^{-1}) (U_+(\tau) + U_0(\tau) + U_-(\tau)). \end{aligned}$$

Clearly,  $U_+(\tau) + U_0(\tau) + U_-(\tau) = \|\hat{u}\|_{\mathcal{H}}^2 \rightarrow 0$  as  $\tau \rightarrow -\infty$ . Moreover,  $U_+(\tau) + U_0(\tau) + U_-(\tau) = \|\hat{u}\|_{\mathcal{H}}^2 > 0$  since  $\tilde{M}_\tau$  is strictly convex.

**Lemma 2.7.** *We have  $U_0(\tau) + U_-(\tau) \leq o(1)U_+(\tau)$ .*

**Proof.** Applying an ODE lemma of Merle and Zaag (cf. Lemma 5.4 in [2] or Lemma A.1 in [21]), we conclude that either  $U_0(\tau) + U_-(\tau) \leq o(1)U_+(\tau)$  or  $U_+(\tau) + U_-(\tau) \leq o(1)U_0(\tau)$ .

The second case can be ruled out as follows: Suppose  $U_+(\tau) + U_-(\tau) \leq o(1)U_0(\tau)$ . Then  $\frac{\hat{u}(\cdot, \tau)}{\|\hat{u}(\cdot, \tau)\|_{\mathcal{H}}}$  converges with respect to  $\|\cdot\|_{\mathcal{H}}$  to the subspace

$\mathcal{H}_0 = \text{span}\{z^2 - 2, z \cos \theta, z \sin \theta\}$ . On the other hand, the orthogonality relations used to define  $S(\tau)$  imply that the function  $\hat{u}(\cdot, \tau)$  is orthogonal to the function  $\langle Ax, \nu_\Sigma \rangle$  for each  $A \in so(3)$ . In other words, the function  $\hat{u}(\cdot, \tau)$  is orthogonal to the functions  $z \cos \theta$  and  $z \sin \theta$ . Consequently,  $\frac{\hat{u}(\cdot, \tau)}{\|\hat{u}(\cdot, \tau)\|_{\mathcal{H}}}$  converges with respect to  $\|\cdot\|_{\mathcal{H}}$  to a non-zero multiple of  $z^2 - 2$ .

Let  $\Omega_\tau$  denote the region enclosed by  $\tilde{M}_\tau$ . Moreover, we denote by  $\mathcal{A}(z, \tau)$  the area of the intersection  $\Omega_\tau \cap \{x_3 = z\}$ . By the Brunn-Minkowski inequality, the function  $z \mapsto \sqrt{\mathcal{A}(z, \tau)}$  is concave. Since  $\tilde{M}_\tau$  is noncompact, it follows that the function  $z \mapsto \sqrt{\mathcal{A}(z, \tau)}$  is monotone.

For  $|z| \leq \rho(\tau)$ , we have the exact identity  $\mathcal{A}(z, \tau) = \frac{1}{2} \int_0^{2\pi} (\sqrt{2} + u(\theta, z, \tau))^2 d\theta$ . Thus, the function  $z \mapsto \int_0^{2\pi} (2\sqrt{2} u(\theta, z, \tau) + u(\theta, z, \tau)^2) d\theta$  is monotone. In particular, we either have

$$\begin{aligned} & \int_{-3}^{-1} \int_0^{2\pi} (2\sqrt{2} u(\theta, z, \tau) + u(\theta, z, \tau)^2) d\theta dz \\ & \leq \int_{-1}^1 \int_0^{2\pi} (2\sqrt{2} u(\theta, z, \tau) + u(\theta, z, \tau)^2) d\theta dz \\ & \leq \int_1^3 \int_0^{2\pi} (2\sqrt{2} u(\theta, z, \tau) + u(\theta, z, \tau)^2) d\theta dz \end{aligned}$$

or

$$\begin{aligned} & \int_{-3}^{-1} \int_0^{2\pi} (2\sqrt{2} u(\theta, z, \tau) + u(\theta, z, \tau)^2) d\theta dz \\ & \geq \int_{-1}^1 \int_0^{2\pi} (2\sqrt{2} u(\theta, z, \tau) + u(\theta, z, \tau)^2) d\theta dz \\ & \geq \int_1^3 \int_0^{2\pi} (2\sqrt{2} u(\theta, z, \tau) + u(\theta, z, \tau)^2) d\theta dz. \end{aligned}$$

However, neither of these possibilities is consistent with the fact that the limit of  $\frac{\hat{u}(\cdot, \tau)}{\|\hat{u}(\cdot, \tau)\|_{\mathcal{H}}}$  is a non-zero multiple of  $z^2 - 2$ . This is a contradiction. This completes the proof of Lemma 2.7.

**Lemma 2.8.** *For each  $\varepsilon > 0$ , we have  $\|u(\cdot, \tau)\|_{C^4([0, 2\pi] \times [-100, 100])} \leq o(e^{\frac{(1-\varepsilon)\tau}{2}})$  and  $|A(\tau)| \leq o(e^{\frac{(1-\varepsilon)\tau}{2}})$ .*

**Proof.** Recall that  $U_0(\tau) + U_-(\tau) \leq o(1)U_+(\tau)$  by Lemma 2.7. This directly implies

$$\frac{d}{d\tau} U_+(\tau) \geq U_+(\tau) - o(1)U_+(\tau).$$

Consequently, for every  $\varepsilon > 0$ , we have  $U_+(\tau) \leq o(e^{(1-\varepsilon)\tau})$ . This gives  $U_0(\tau) + U_-(\tau) \leq o(1)U_+(\tau) \leq o(e^{(1-\varepsilon)\tau})$ , hence

$$\|\hat{u}\|_{\mathcal{H}}^2 = U_+(\tau) + U_0(\tau) + U_-(\tau) \leq o(e^{(1-\varepsilon)\tau}).$$

Using Lemma 2.6, we obtain  $|A(\tau)| \leq o(1) \|\hat{u}\|_{\mathcal{H}} \leq o(e^{\frac{(1-\varepsilon)\tau}{2}})$ . Finally, the inequality  $\|u(\cdot, \tau)\|_{C^4([0, 2\pi] \times [-100, 100])} \leq o(e^{\frac{(1-\varepsilon)\tau}{2}})$  follows from standard interpolation inequalities. This completes the proof of Lemma 2.8.

Recall that  $A(\tau) = S'(\tau)S(\tau)^{-1}$ . Since  $|A(\tau)| \leq o(e^{\frac{(1-\varepsilon)\tau}{2}})$  by Lemma 2.8, the limit  $\lim_{\tau \rightarrow -\infty} S(\tau)$  exists. Without loss of generality, we may assume that  $\lim_{\tau \rightarrow -\infty} S(\tau) = \text{id}$ . Clearly,  $|S(\tau) - \text{id}| \leq o(e^{\frac{(1-\varepsilon)\tau}{2}})$ .

**Lemma 2.9.** *We have*

$$\sup_{\bar{M}_\tau \cap \{|x_3| \leq e^{-\frac{\tau}{10}}\}} |x_1^2 + x_2^2 - 2| \leq e^{\frac{\tau}{10}}$$

if  $-\tau$  is sufficiently large.

**Proof.** Using Lemma 2.8 and the estimate  $|S(\tau) - \text{id}| \leq o(e^{\frac{(1-\varepsilon)\tau}{2}})$ , we obtain

$$\sup_{x \in \bar{M}_\tau \cap B_{10}(0)} |x_1^2 + x_2^2 - 2| \leq o(e^{\frac{(1-\varepsilon)\tau}{2}}).$$

The convexity of  $\bar{M}_\tau$  implies

$$\sup_{\bar{M}_\tau \cap \{|x_3| \leq e^{-\frac{\tau}{10}}\}} (x_1^2 + x_2^2) \leq 2 + e^{\frac{\tau}{10}}$$

if  $-\tau$  is sufficiently large. Let

$$\Sigma_a = \{(x_1, x_2, x_3) : x_1^2 + x_2^2 = u_a(-x_3)^2, -a \leq x_3 \leq 0\}$$

denote the self-similar shrinker constructed in [2]. By Lemma 4.4 in [2],  $u_a(2) \leq \sqrt{2} - a^{-2}$ . Since  $\bar{M}_\tau$  converges to  $\Sigma$  in  $C_{\text{loc}}^\infty$ , the surface  $\bar{M}_\tau \cap \{x_3 \leq -2\}$  encloses the surface  $\Sigma_a \cap \{x_3 \leq -2\}$  if  $-\tau$  is sufficiently large (depending on  $a$ ). On the other hand, since  $\inf_{x \in \bar{M}_\tau \cap B_{10}(0)} (x_1^2 + x_2^2) \geq 2 - o(e^{\frac{(1-\varepsilon)\tau}{2}})$ , the boundary  $\bar{M}_\tau \cap \{x_3 = -2\}$  encloses the boundary  $\Sigma_a \cap \{x_3 = -2\}$  provided that  $-\tau$  is sufficiently large and  $a \leq e^{-\frac{(1-\varepsilon)\tau}{4}}$ . By the maximum principle, the surface  $\bar{M}_\tau \cap \{x_3 \leq -2\}$  encloses  $\Sigma_a \cap \{x_3 \leq -2\}$  whenever  $-\tau$  is sufficiently large and  $a \leq e^{-\frac{(1-\varepsilon)\tau}{4}}$ . Using Theorem 8.2 in [2], we obtain  $u_a(y) \geq \sqrt{2(1 - a^{-2}y^2)}$  for all  $y \in [0, a]$ , provided that  $a$  is sufficiently large. Putting these facts together, we obtain

$$\inf_{\bar{M}_\tau \cap \{-e^{-\frac{\tau}{10}} \leq x_3 \leq -2\}} (x_1^2 + x_2^2) \geq 2 - e^{\frac{\tau}{10}}$$

if  $-\tau$  is sufficiently large. An analogous argument gives

$$\inf_{\bar{M}_\tau \cap \{2 \leq x_3 \leq e^{-\frac{\tau}{10}}\}} (x_1^2 + x_2^2) \geq 2 - e^{\frac{\tau}{10}}$$

if  $-\tau$  is sufficiently large. Putting these facts together, we obtain

$$\inf_{\bar{M}_\tau \cap \{|x_3| \leq e^{-\frac{\tau}{10}}\}} (x_1^2 + x_2^2) \geq 2 - e^{\frac{\tau}{10}}$$

if  $-\tau$  is sufficiently large. This completes the proof of Lemma 2.9.

**Lemma 2.10.** *Let  $\varepsilon_0 > 0$  be given. If  $-\tau$  is sufficiently large (depending on  $\varepsilon_0$ ), then every point in  $\bar{M}_\tau \cap \{|x_3| \leq \frac{1}{2} e^{-\frac{\tau}{10}}\}$  lies at the center of an  $\varepsilon_0$ -neck.*

**Proof.** Suppose that there exists a sequence of times  $\tau_j \rightarrow -\infty$  and a sequence of points  $q_j \in \bar{M}_{\tau_j} \cap \{|x_3| \leq \frac{1}{2} e^{-\frac{\tau_j}{10}}\}$  such that  $q_j$  does not lie on an  $\varepsilon_0$ -neck. Using Lemma 2.9 and the noncollapsing property, we conclude that the mean curvature at  $q_j$  is bounded from below by a positive constant. We now consider the triangle in  $\mathbb{R}^3$  with vertices  $q_j$ ,  $(0, 0, e^{-\frac{\tau_j}{10}})$ , and  $(0, 0, -e^{-\frac{\tau_j}{10}})$ . Using Lemma 2.9 and the convexity of  $\bar{M}_{\tau_j}$ , we conclude that this triangle lies inside  $\bar{M}_{\tau_j}$ . Moreover, the angle at  $q_j$  converges to  $\pi$  as  $j \rightarrow -\infty$ . We now dilate the surface  $\bar{M}_{\tau_j}$  to make the mean curvature at  $q_j$  equal to  $\frac{1}{\sqrt{2}}$ . Passing to the limit as  $j \rightarrow \infty$ , we obtain a noncollapsed ancient solution of mean curvature flow which is weakly, but not strictly convex. By the strong maximum principle, the limit splits off a line. By Lemma 3.6 in [12], the limit is a round cylinder. Therefore, the point  $q_j$  lies on an  $\varepsilon_0$ -neck if  $j$  is sufficiently large. This is a contradiction.

After these preparations, we now state the main result of this section:

**Proposition 2.11.** *We have*

$$\sup_{x \in \bar{M}_\tau \cap B_{10}(0)} |x_1^2 + x_2^2 - 2| \leq O(e^{\frac{\tau}{2}}).$$

**Proof.** In view of Lemma 2.9, Lemma 2.10, and standard interpolation inequalities, we may write  $\bar{M}_\tau$  as a graph over the cylinder  $\Sigma \cap B_{e^{-\frac{\tau}{100}}}(0)$ , and the  $C^4$ -norm of the height function is bounded by  $O(e^{\frac{\tau}{100}})$ . We now repeat the argument above, this time with  $\rho(\tau) = e^{-\frac{\tau}{1000}}$ . As above, we consider the rotated surfaces  $\tilde{M}_\tau = S(\tau)\bar{M}_\tau$ , where  $S(\tau)$  is a function taking values in  $SO(3)$ . We write each surface  $\tilde{M}_\tau$  as a graph over the cylinder, so that

$$\{x + u(x, \tau)\nu_\Sigma(x) : x \in \Sigma \cap B_{2e^{-\frac{\tau}{1000}}}(0)\} \subset \tilde{M}_\tau,$$

where  $\|u(\cdot, \tau)\|_{C^4(\Sigma \cap B_{2e^{-\frac{\tau}{1000}}}(0))} \leq O(e^{\frac{\tau}{200}})$ . We choose the matrices  $S(\tau)$  in such a way that the orthogonality relations

$$\int_{\Sigma \cap B_{e^{-\frac{\tau}{1000}}}(0)} e^{-\frac{|x|^2}{4}} \langle Ax, \nu_\Sigma \rangle u(x, \tau) \varphi(e^{\frac{\tau}{1000}} x_3) = 0$$

are satisfied for all  $A \in so(3)$ . As above, the function  $\hat{u}(x, \tau) = u(x, \tau) \varphi(e^{\frac{\tau}{1000}} x_3)$  satisfies

$$\left\| \frac{\partial}{\partial \tau} \hat{u} - \mathcal{L} \hat{u} \right\|_{\mathcal{H}} \leq O(e^{\frac{\tau}{1000}}) \|\hat{u}\|_{\mathcal{H}}.$$

Hence, if we define

$$\begin{aligned} U_+(\tau) &:= \|P_+\hat{u}(\cdot, \tau)\|_{\mathcal{H}}^2, \\ U_0(\tau) &:= \|P_0\hat{u}(\cdot, \tau)\|_{\mathcal{H}}^2, \\ U_-(\tau) &:= \|P_-\hat{u}(\cdot, \tau)\|_{\mathcal{H}}^2, \end{aligned}$$

then

$$\begin{aligned} \frac{d}{d\tau}U_+(\tau) &\geq U_+(\tau) - O(e^{\frac{\tau}{1000}})(U_+(\tau) + U_0(\tau) + U_-(\tau)), \\ \left| \frac{d}{d\tau}U_0(\tau) \right| &\leq O(e^{\frac{\tau}{1000}})(U_+(\tau) + U_0(\tau) + U_-(\tau)), \\ \frac{d}{d\tau}U_-(\tau) &\leq -U_-(\tau) + O(e^{\frac{\tau}{1000}})(U_+(\tau) + U_0(\tau) + U_-(\tau)). \end{aligned}$$

As above, the ODE lemma of Merle and Zaag implies that either  $U_0(\tau) + U_-(\tau) \leq o(1)U_+(\tau)$  or  $U_+(\tau) + U_-(\tau) \leq o(1)U_0(\tau)$ , and the latter case can be ruled out as above using the orthogonality relations and the Brunn-Minkowski inequality. Thus,  $U_0(\tau) + U_-(\tau) \leq o(1)U_+(\tau)$ . This gives

$$\frac{d}{d\tau}U_+(\tau) \geq U_+(\tau) - O(e^{\frac{\tau}{1000}})U_+(\tau),$$

hence  $U_+(\tau) \leq O(e^\tau)$ . This implies  $U_0(\tau) + U_-(\tau) \leq o(1)U_+(\tau) \leq O(e^\tau)$ . From this, we deduce that  $\|\hat{u}\|_{\mathcal{H}} \leq O(e^{\frac{\tau}{2}})$ . Using Lemma 2.6, we obtain  $|A(\tau)| \leq O(e^{\frac{\tau}{2}})$ . Since  $\lim_{\tau \rightarrow -\infty} S(\tau) = \text{id}$ , it follows that  $|S(\tau) - \text{id}| \leq O(e^{\frac{\tau}{2}})$ . Finally, we observe that  $u$  satisfies an equation of the form  $\frac{\partial}{\partial \tau}u = \tilde{\mathcal{L}}u + \langle A(\tau)x, \nu_\Sigma \rangle$ , where  $\tilde{\mathcal{L}}$  is an elliptic operator of second order whose coefficients depend on  $u$ ,  $\nabla u$ ,  $\nabla^2 u$ , and  $A(\tau)$ . As  $\tau \rightarrow -\infty$ , the coefficients of  $\tilde{\mathcal{L}}$  converge smoothly to the corresponding coefficients of  $\mathcal{L}$ . Using standard interior estimates for parabolic equations, we obtain  $\|u(\cdot, \tau)\|_{C^4([0, 2\pi] \times [-100, 100])} \leq O(e^{\frac{\tau}{2}})$ . Since  $|S(\tau) - \text{id}| \leq O(e^{\frac{\tau}{2}})$ , we conclude that

$$\sup_{x \in \bar{M}_\tau \cap B_{10}(0)} |x_1^2 + x_2^2 - 2| \leq O(e^{\frac{\tau}{2}}).$$

This completes the proof of Proposition 2.11.

### 3. LOWER BOUND FOR $H_{\max}(t)$ AS $t \rightarrow -\infty$

Let  $M_t$ ,  $t \in (-\infty, 0]$ , be a noncompact ancient solution of mean curvature flow in  $\mathbb{R}^3$  which is strictly convex and noncollapsed. For each  $t$ , we denote by  $H_{\max}(t)$  the supremum of the mean curvature of  $M_t$ .

**Proposition 3.1.** *For each  $t$ ,  $H_{\max}(t) < \infty$ .*

**Proof.** Let us fix a time  $t$  and a small number  $\varepsilon$ . By Proposition 3.1 in [13], we can find a compact subset of  $M_t$  with the property that every point in the complement of that set lies at the center of an  $\varepsilon$ -neck. Hence, if  $H_{\max}(t) = \infty$ , then the surface  $M_t$  contains a sequence of  $\varepsilon$ -necks with radii

converging to 0, but this is impossible in a convex surface.

**Corollary 3.2.** *The function  $H_{\max}(t)$  is continuous and monotone increasing in  $t$ .*

**Proof.** The pointwise curvature derivative estimate of Haslhofer and Kleiner [12],[13] gives  $|\frac{\partial}{\partial t}H| \leq CH^3$  for some uniform constant  $C$ . Consequently,  $H_{\max}(t)$  is continuous in  $t$ . In particular,  $H_{\max}(t)$  is uniformly bounded from above on every compact time interval. Hence, we can apply Hamilton's Harnack inequality [10] to conclude that  $H_{\max}(t)$  is monotone increasing in  $t$ .

**Proposition 3.3.** *We have  $\liminf_{t \rightarrow -\infty} H_{\max}(t) > 0$ .*

**Proof.** Proposition 2.11 gives

$$\sup_{x \in (-t)^{-\frac{1}{2}}(M_t \cap B_{10(-t)^{\frac{1}{2}}}(0))} |x_1^2 + x_2^2 - 2| \leq O((-t)^{-\frac{1}{2}}).$$

By assumption,  $M_t$  is noncompact and strictly convex. Hence,  $M_t$  has exactly one end. Without loss of generality, we may assume that  $M_t \cap \{x_3 \leq 0\}$  is compact and  $M_t \cap \{x_3 \geq 0\}$  is noncompact. We can find a large constant  $K$  such that the curve

$$(-t)^{-\frac{1}{2}}(M_t \cap \{x_3 = -2(-t)^{\frac{1}{2}}\})$$

lies outside the circle

$$\{x_1^2 + x_2^2 = (\sqrt{2} - K(-t)^{-\frac{1}{2}})^2, x_3 = -2\}$$

if  $-t$  is sufficiently large. Let us consider the self-similar solutions constructed in [2]. For  $a > 0$  large, there exists a surface

$$\Sigma_a = \{(x_1, x_2, x_3) : x_1^2 + x_2^2 = u_a(-x_3)^2, -a \leq x_3 \leq 0\}$$

which satisfies the shrinker equation  $H = \frac{1}{2}\langle x, \nu \rangle$ . Hence, the surfaces

$$\begin{aligned} \Sigma_{a,t} &:= (-t)^{\frac{1}{2}}\Sigma_a + (0, 0, Ka^2) \\ &= \{(x_1, x_2, x_3) : x_1^2 + x_2^2 = (-t)u_a((-x_3 + Ka^2)(-t)^{-\frac{1}{2}})^2, \\ &\quad Ka^2 - a(-t)^{\frac{1}{2}} \leq x_3 \leq Ka^2\} \end{aligned}$$

evolve by mean curvature flow.

We will use the surfaces  $\Sigma_{a,t} \cap \{x_3 \leq -2(-t)^{\frac{1}{2}}\}$  as barriers for the flow  $M_t \cap \{x_3 \leq -2(-t)^{\frac{1}{2}}\}$ . As  $t \rightarrow -\infty$ , the rescaled surfaces  $(-t)^{-\frac{1}{2}}M_t$  converge in  $C_{\text{loc}}^\infty$  to the cylinder  $\{x_1^2 + x_2^2 = 2\}$ . Moreover, as  $t \rightarrow -\infty$ , the rescaled surfaces  $(-t)^{-\frac{1}{2}}(\Sigma_{a,t} \cap \{x_3 \leq -2(-t)^{\frac{1}{2}}\})$  converge to  $\Sigma_a \cap \{x_3 \leq -2\}$ , which is a compact subset of  $\{x_1^2 + x_2^2 < 2\}$ . Therefore,  $\Sigma_{a,t} \cap \{x_3 \leq -2(-t)^{\frac{1}{2}}\}$  lies inside  $M_t \cap \{x_3 \leq -2(-t)^{\frac{1}{2}}\}$  if  $-t$  is sufficiently large (depending on  $a$ ).

We next examine the boundary curves  $M_t \cap \{x_3 = -2(-t)^{\frac{1}{2}}\}$  and  $\Sigma_{a,t} \cap \{x_3 = -2(-t)^{\frac{1}{2}}\}$ . By our choice of  $K$ , the curve

$$(-t)^{-\frac{1}{2}} (M_t \cap \{x_3 = -2(-t)^{\frac{1}{2}}\})$$

lies outside the circle

$$\{x_1^2 + x_2^2 = (\sqrt{2} - K(-t)^{-\frac{1}{2}})^2, x_3 = -2\}.$$

Moreover, the curve

$$(-t)^{-\frac{1}{2}} (\Sigma_{a,t} \cap \{x_3 = -2(-t)^{\frac{1}{2}}\})$$

is a circle

$$\{x_1^2 + x_2^2 = u_a(2 + Ka^2(-t)^{-\frac{1}{2}})^2, x_3 = -2\}.$$

Using Lemma 4.4 in [2], we obtain  $u_a(2) \leq \sqrt{2}$  and  $u_a(2) - u_a(1) \leq -a^{-2}$  if  $a$  is sufficiently large. Moreover, by Lemma 4.2 in [2], the function  $u_a : [0, a] \rightarrow \mathbb{R}$  is concave. Hence, we obtain

$$\begin{aligned} u_a(2 + Ka^2(-t)^{-\frac{1}{2}}) &\leq u_a(2) + Ka^2(-t)^{-\frac{1}{2}} (u_a(2) - u_a(1)) \\ &\leq \sqrt{2} - K(-t)^{-\frac{1}{2}} \end{aligned}$$

for  $-t \geq 4K^2a^2$ . Therefore, the curve  $\Sigma_{a,t} \cap \{x_3 = -2(-t)^{\frac{1}{2}}\}$  lies inside the curve  $M_t \cap \{x_3 = -2(-t)^{\frac{1}{2}}\}$  whenever  $-t \geq 4K^2a^2$  and  $a$  is sufficiently large. Using the maximum principle, we conclude that the surface  $\Sigma_{a,t} \cap \{x_3 \leq -2(-t)^{\frac{1}{2}}\}$  lies inside the surface  $M_t \cap \{x_3 \leq -2(-t)^{\frac{1}{2}}\}$  whenever  $-t \geq 4K^2a^2$  and  $a$  is sufficiently large. For  $-t = 4K^2a^2$ , the tip of  $\Sigma_{a,t}$  has distance  $a(-t)^{\frac{1}{2}} - Ka^2 = Ka^2 = -\frac{t}{4K}$  from the origin. Consequently, the intersection of  $M_t$  with the halfline  $\{x_1 = x_2 = 0, x_3 \leq \frac{t}{4K}\}$  is non-empty if  $-t$  is sufficiently large. From this, we deduce that  $\limsup_{t \rightarrow -\infty} H_{\max}(t) > 0$ . Since  $H_{\max}(t)$  is monotone increasing in  $t$ , we conclude that  $\liminf_{t \rightarrow -\infty} H_{\max}(t) > 0$ .

#### 4. THE NECK IMPROVEMENT THEOREM

**Definition 4.1.** Let  $K$  be a vector field in  $\mathbb{R}^3$ . We say that  $K$  is a normalized rotation vector field if there exists a matrix  $S \in O(3)$  and a point  $q \in \mathbb{R}^3$  such that  $K(x) = SJS^{-1}(x - q)$ , where

$$J = \begin{bmatrix} 0 & 1 & 0 \\ -1 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}.$$

Note that we do not require that the origin lies on the axis of rotation.

**Lemma 4.2.** *There exists a large constant  $C$  and small constant  $\varepsilon_0 > 0$  with the following property. Suppose that  $M$  is a surface in  $\mathbb{R}^3$  and let  $\bar{x}$  be a point on  $M$ . We assume that, after rescaling, the surface  $M$  is  $\varepsilon_0$ -close (in the  $C^4$ -norm) to a cylinder  $S^1 \times [-5, 5]$  of radius 1. Suppose that  $K^{(1)}$  and  $K^{(2)}$  are normalized rotation vector fields with the following properties:*

- $|K^{(1)}|H \leq 10$  and  $|K^{(2)}|H \leq 10$  at the point  $\bar{x}$ .
- $|\langle K^{(1)}, \nu \rangle|H \leq \varepsilon$  and  $|\langle K^{(2)}, \nu \rangle|H \leq \varepsilon$  in a geodesic ball around  $\bar{x}$  in  $M$  of radius  $H(\bar{x})^{-1}$ .

Then

$$\min \left\{ \sup_{B_{100H(\bar{x})^{-1}}(\bar{x})} |K^{(1)} - K^{(2)}|, \sup_{B_{100H(\bar{x})^{-1}}(\bar{x})} |K^{(1)} + K^{(2)}| \right\} H(\bar{x}) \leq C\varepsilon.$$

**Proof.** By scaling, we may assume that  $H(\bar{x}) = 1$ . We argue by contradiction. If the assertion is false, then there exist a sequence of surfaces  $M^{(j)}$ , a sequence of points  $\bar{x}_j \in M^{(j)}$  satisfying  $H(\bar{x}_j) = 1$ , sequences of normalized rotation vector fields  $K^{(1,j)}$  and  $K^{(2,j)}$ , and a sequences of real number  $\varepsilon_j \rightarrow 0$  with the following properties:

- The surfaces  $M^{(j)}$  are  $\frac{1}{j}$ -close (in the  $C^4$ -norm) to a cylinder  $M = S^1 \times [-5, 5]$  of radius 1. Moreover, we may assume that the axis of the cylinder is the  $x_3$ -axis.
- $|K^{(1,j)}| \leq 10$  and  $|K^{(2,j)}| \leq 10$  at the point  $\bar{x}_j$ .
- $|\langle K^{(1,j)}, \nu \rangle|H \leq \varepsilon_j$  and  $|\langle K^{(2,j)}, \nu \rangle|H \leq \varepsilon_j$  in a geodesic ball around  $\bar{x}_j$  in  $M$  of radius 1.
- $\sup_{B_{100}(\bar{x}_j)} |K^{(1,j)} - K^{(2,j)}| \geq j\varepsilon_j$ .
- $\sup_{B_{100}(\bar{x}_j)} |K^{(1,j)} + K^{(2,j)}| \geq j\varepsilon_j$ .

Note that the distance of  $\bar{x}_j$  from the axis of rotation of  $K^{(1,j)}$  is at most 10. Hence, after passing to a subsequence if necessary, the vector fields  $K^{(1,j)}$  converge to a normalized vector field  $K^{(1)}$ . Similarly, the vector fields  $K^{(2,j)}$  converge to a vector field  $K^{(2)}$ . Clearly,  $K^{(1)}$  and  $K^{(2)}$  are tangential to the cylinder  $S^1 \times [-5, 5]$ . Consequently, we have  $K^{(1)}(x) = \pm Jx$  and  $K^{(2)}(x) = \pm Jx$ , where  $J$  is defined as above. Without loss of generality, we assume that  $K^{(1)}(x) = K^{(2)}(x) = Jx$ . For each  $j$ , we define  $\delta_j := \sup_{B_{100}(\bar{x}_j)} |K^{(1,j)} - K^{(2,j)}| \geq j\varepsilon_j$ . Clearly,  $\delta_j \rightarrow 0$ . We next consider the Killing vector field  $V^{(j)} := \delta_j^{-1} (K^{(1,j)} - K^{(2,j)})$ . Then  $\sup_{B_{100}(\bar{x}_j)} |V^{(j)}| = 1$ , and  $|\langle V^{(j)}, \nu \rangle|H \leq 2\delta_j^{-1}\varepsilon_j \leq 2j^{-1}$  in a geodesic ball around  $\bar{x}_j$  in  $M$  of radius 1. Hence, after passing to a subsequence, the vector fields  $V^{(j)}$  converge to a non-trivial Killing vector field  $V$  on  $\mathbb{R}^3$  which is tangential to the cylinder  $S^1 \times [-5, 5]$ . Since  $K^{(1,j)}$  and  $K^{(2,j)}$  are normalized rotation vector fields, the limit vector field  $V$  must be of the form  $V(x) = [A, J]x - Jb$  for some matrix  $A \in so(3)$  and some vector  $b \in \mathbb{R}^3$ . However, such a vector field cannot be tangential to the cylinder  $S^1 \times [-5, 5]$  unless  $V$  vanishes identically. This is a contradiction.

As in [18], pp. 189–190, we denote by  $\mathcal{P}(\bar{x}, \bar{t}, r, \tau)$  the set of all points  $(x, t)$  in space-time such that  $x \in B_{g(\bar{t})}(\bar{x}, r)$  and  $t \in [\bar{t} - \tau, \bar{t}]$ . Moreover, we put  $\hat{\mathcal{P}}(\bar{x}, \bar{t}, L, \theta) = \mathcal{P}(\bar{x}, \bar{t}, L H(\bar{x}, \bar{t})^{-1}, \theta H(\bar{x}, \bar{t})^{-2})$ . We say that  $(\bar{x}, \bar{t})$  lies on

an  $\varepsilon$ -neck if the parabolic neighborhood  $\hat{\mathcal{P}}(\bar{x}, \bar{t}, 100, 100)$  is, after rescaling,  $\varepsilon$ -close (in the  $C^{10}$ -norm), to a family of shrinking cylinders.

**Definition 4.3.** Let  $M_t$  be a solution of mean curvature flow with positive mean curvature. We say that a point  $(\bar{x}, \bar{t})$  is  $\varepsilon$ -symmetric if there exists a normalized rotation vector field  $K$  on  $\mathbb{R}^3$  such that  $|K|H \leq 10$  at the point  $(\bar{x}, \bar{t})$  and  $|\langle K, \nu \rangle|H \leq \varepsilon$  in the parabolic neighborhood  $\hat{\mathcal{P}}(\bar{x}, \bar{t}, 10, 100)$ .

Note that the condition that  $|K|H \leq 10$  at the point  $(\bar{x}, \bar{t})$  is equivalent to the condition that the distance of the point  $\bar{x}$  from the axis of rotation of  $K$  is at most  $10H(\bar{x}, \bar{t})^{-1}$ .

**Theorem 4.4** (Neck Improvement Theorem). *There exists a large constant  $L$  and a small constant  $\varepsilon_1$  with the following property. Let  $M_t$  be a solution of mean curvature flow, and let  $(\bar{x}, \bar{t})$  be a point in space-time. Suppose that every point in the parabolic neighborhood  $\hat{\mathcal{P}}(\bar{x}, \bar{t}, L, L^2)$  lies on an  $\varepsilon_1$ -neck. Moreover, suppose that every point in  $\hat{\mathcal{P}}(\bar{x}, \bar{t}, L, L^2)$  is  $\varepsilon$ -symmetric, where  $\varepsilon \leq \varepsilon_1$ . Then  $(\bar{x}, \bar{t})$  is  $\frac{\varepsilon}{2}$ -symmetric.*

**Proof.** Without loss of generality, we assume  $\bar{t} = -1$  and  $H(\bar{x}, -1) = \frac{1}{\sqrt{2}}$ . Throughout the proof, we assume that  $L$  is sufficiently large, and  $\varepsilon_1$  is sufficiently small depending on  $L$ . In the parabolic neighborhood  $\hat{\mathcal{P}}(\bar{x}, \bar{t}, L, L^2)$ , we can approximate  $M_t$  by a cylinder  $S^1((-2t)^{\frac{1}{2}}) \times \mathbb{R}$ , up to errors which are bounded by  $C(L)\varepsilon_1$  in the  $C^{100}$ -norm.

*Step 1:* By assumption, for every point  $(x_0, t_0) \in \hat{\mathcal{P}}(\bar{x}, -1, L, L^2)$ , there exists a normalized vector field  $K^{(x_0, t_0)}$  such that  $|K^{(x_0, t_0)}|H \leq 10$  at the point  $(x_0, t_0)$ , and  $|\langle K^{(x_0, t_0)}, \nu \rangle|H \leq \varepsilon$  on the parabolic neighborhood  $\hat{\mathcal{P}}(x_0, t_0, 10, 100)$ . A repeated application of Lemma 4.2 gives

$$\min \left\{ \sup_{B_{10L}(0)} |K^{(\bar{x}, -1)} - K^{(x_0, t_0)}|, \sup_{B_{10L}(0)} |K^{(\bar{x}, -1)} + K^{(x_0, t_0)}| \right\} \leq C(L)\varepsilon$$

for all  $(x_0, t_0) \in \hat{\mathcal{P}}(\bar{x}, -1, L, L^2)$ . Without loss of generality, we may assume that

$$\sup_{B_{10L}(0)} |K^{(\bar{x}, -1)} - K^{(x_0, t_0)}| \leq C(L)\varepsilon$$

for all  $(x_0, t_0) \in \hat{\mathcal{P}}(\bar{x}, -1, L, L^2)$ . Moreover, we may assume without loss of generality that  $\bar{K} = K^{(\bar{x}, -1)}$  is an infinitesimal rotation around the  $x_3$ -axis, so that  $\bar{K}(y) = Jy$ , where  $J$  is defined as in Definition 4.1. Finally, we may assume that the point  $\bar{x}$  lies in the plane  $\{x_3 = 0\}$ .

Let us write  $M_t$  as a graph over the  $x_3$ -axis, so that

$$\left\{ (r(\theta, z, t) \cos \theta, r(\theta, z, t) \sin \theta, z) : \theta \in [0, 2\pi], z \in \left[ -\frac{L}{4}, \frac{L}{4} \right] \right\} \subset M_t.$$

By assumption, the difference  $r(\theta, z, t) - (-2t)^{\frac{1}{2}}$  is bounded by  $C(L)\varepsilon_1$  in the  $C^{100}$ -norm. The unit normal vector to  $M_t$  is given by

$$\nu = \frac{1}{\sqrt{1 + r^{-2}\left(\frac{\partial r}{\partial \theta}\right)^2 + \left(\frac{\partial r}{\partial z}\right)^2}} \cdot \left[ (\cos \theta, \sin \theta, 0) - r^{-1} \frac{\partial r}{\partial \theta} (-\sin \theta, \cos \theta, 0) - \frac{\partial r}{\partial z} (0, 0, 1) \right].$$

We define

$$u = \langle \bar{K}, \nu \rangle = \frac{1}{\sqrt{1 + r^{-2}\left(\frac{\partial r}{\partial \theta}\right)^2 + \left(\frac{\partial r}{\partial z}\right)^2}} \frac{\partial r}{\partial \theta}.$$

*Step 2:* For each point  $(x_0, t_0) \in \hat{\mathcal{P}}(\bar{x}, -1, L, L^2)$ , we know that

$$|\langle K^{(x_0, t_0)}, \nu \rangle| \leq C\varepsilon (-t_0)^{\frac{1}{2}}$$

on the parabolic neighborhood  $\hat{\mathcal{P}}(x_0, t_0, 10, 100)$ . Moreover, we can find a matrix  $S \in O(3)$  and a vector  $q \in \mathbb{R}^3$  (depending on  $(x_0, t_0)$ ) such that  $\bar{K}(y) - K^{(x_0, t_0)}(y) = Jy - SJS^{-1}(y - q)$  and  $|S - \text{id}| + |q| \leq C(L)\varepsilon$ . Consequently, there exist real numbers  $a_0, a_1, b_0, b_1$  (depending on  $(x_0, t_0)$ ) such that

$$|a_0| + |a_1| + |b_0| + |b_1| \leq C(L)\varepsilon$$

and

$$|\langle \bar{K} - K^{(x_0, t_0)}, \nu \rangle - (a_0 + a_1 z) \cos \theta - (b_0 + b_1 z) \sin \theta| \leq C(L)\varepsilon_1\varepsilon$$

on the parabolic neighborhood  $\hat{\mathcal{P}}(x_0, t_0, 10, 100)$ . Note that the numbers  $a_0, a_1, b_0, b_1$  account for the fact that the rotation vector fields  $K^{(x_0, t_0)}$  and  $\bar{K}$  may have different axes of rotation.

Putting these facts together, we obtain

$$|\langle \bar{K}, \nu \rangle - (a_0 + a_1 z) \cos \theta - (b_0 + b_1 z) \sin \theta| \leq C\varepsilon (-t_0)^{\frac{1}{2}} + C(L)\varepsilon_1\varepsilon$$

on the parabolic neighborhood  $\hat{\mathcal{P}}(x_0, t_0, 10, 100)$ . To summarize, given any point  $(z_0, t_0) \in [-\frac{L}{2}, \frac{L}{2}] \times [-\frac{L^2}{4}, -1]$ , there exist real numbers  $a_0, a_1, b_0, b_1$  (depending on  $(z_0, t_0)$ ) such that

$$|a_0| + |a_1| + |b_0| + |b_1| \leq C(L)\varepsilon$$

and

$$|u(\theta, z, t) - (a_0 + a_1 z) \cos \theta - (b_0 + b_1 z) \sin \theta| \leq C\varepsilon (-t_0)^{\frac{1}{2}} + C(L)\varepsilon_1\varepsilon$$

for  $z \in [z_0 - (-t_0)^{\frac{1}{2}}, z_0 + (-t_0)^{\frac{1}{2}}]$  and  $t \in [2t_0, t_0]$ .

*Step 3:* The function  $u = \langle \bar{K}, \nu \rangle$  satisfies the evolution equation

$$\frac{\partial}{\partial t} u = \Delta_{M_t} u + |A|^2 u.$$

Since  $|u| \leq C(L)\varepsilon$ , it follows from standard interior estimates for parabolic equations that  $|\nabla u| + |\nabla^2 u| \leq C(L)\varepsilon$  for  $z \in [-\frac{L}{4}, \frac{L}{4}]$  and  $t \in [-\frac{L^2}{16}, -1]$ . Hence, we obtain

$$\left| \frac{\partial}{\partial t} u - \frac{\partial^2}{\partial z^2} u - \frac{1}{(-2t)} \frac{\partial^2}{\partial \theta^2} u - \frac{1}{(-2t)} u \right| \leq C(L)\varepsilon_1\varepsilon$$

for  $z \in [-\frac{L}{4}, \frac{L}{4}]$  and  $t \in [-\frac{L^2}{16}, -1]$ .

Let  $\tilde{u}$  be the solution of the linear equation

$$\frac{\partial}{\partial t} \tilde{u} = \frac{\partial^2}{\partial z^2} \tilde{u} + \frac{1}{(-2t)} \frac{\partial^2}{\partial \theta^2} \tilde{u} + \frac{1}{(-2t)} \tilde{u}$$

in the parabolic cylinder  $\{z \in [-\frac{L}{4}, \frac{L}{4}], t \in [-\frac{L^2}{16}, -1]\}$  such that  $\tilde{u} = u$  on the parabolic boundary  $\{|z| = \frac{L}{4}\} \cup \{t = -\frac{L^2}{16}\}$ . Using the maximum principle, we obtain

$$|u - \tilde{u}| \leq C(L)\varepsilon_1\varepsilon$$

in the parabolic cylinder  $\{z \in [-\frac{L}{4}, \frac{L}{4}], t \in [-\frac{L^2}{16}, -1]\}$ .

*Step 4:* We now analyze the function  $\tilde{u}$  using separation of variables. For  $m \geq 1$ , we put

$$v_m(z, t) = \frac{1}{\pi} \int_0^{2\pi} \tilde{u}(\theta, z, t) \cos(m\theta) d\theta$$

and

$$w_m(z, t) = \frac{1}{\pi} \int_0^{2\pi} \tilde{u}(\theta, z, t) \sin(m\theta) d\theta.$$

The functions  $v_m$  and  $w_m$  satisfy the evolution equations

$$\frac{\partial}{\partial t} v_m = \frac{\partial^2}{\partial z^2} v_m + \frac{1-m^2}{(-2t)} v_m, \quad \frac{\partial}{\partial t} w_m = \frac{\partial^2}{\partial z^2} w_m + \frac{1-m^2}{(-2t)} w_m.$$

Consequently, the functions  $\hat{v}_m = (-t)^{\frac{1-m^2}{2}} v_m$  and  $\hat{w}_m = (-t)^{\frac{1-m^2}{2}} w_m$  satisfy the linear heat equation

$$\frac{\partial}{\partial t} \hat{v}_m = \frac{\partial^2}{\partial z^2} \hat{v}_m, \quad \frac{\partial}{\partial t} \hat{w}_m = \frac{\partial^2}{\partial z^2} \hat{w}_m$$

for  $m \geq 1$ .

*Step 5:* We first consider the modes with  $m \geq 2$ . For  $m \geq 2$ , we have

$$|v_m| + |w_m| \leq (C\varepsilon + C(L)\varepsilon_1\varepsilon) (-t)^{\frac{1}{2}}$$

in the parabolic cylinder  $[-\frac{L}{4}, \frac{L}{4}] \times [-\frac{L^2}{16}, -1]$ . This implies

$$|\hat{v}_m| + |\hat{w}_m| \leq (C\varepsilon + C(L)\varepsilon_1\varepsilon) (-t)^{1-\frac{m^2}{2}}$$

in the parabolic cylinder  $[-\frac{L}{4}, \frac{L}{4}] \times [-\frac{L^2}{16}, -1]$ . Using the solution formula for the one-dimensional heat equation with Dirichlet boundary condition on

the rectangle  $[-\frac{L}{4}, \frac{L}{4}] \times [-\frac{L^2}{16}, -1]$ , we may express  $\hat{v}_m(z, t)$  and  $\hat{w}_m(z, t)$  as integrals of initial and boundary data. This gives

$$\begin{aligned} |\hat{v}_m(z, t)| + |\hat{w}_m(z, t)| &\leq (C\varepsilon + C(L)\varepsilon_1\varepsilon) \left(\frac{L}{4}\right)^{2-m^2} \\ &\quad + (C\varepsilon + C(L)\varepsilon_1\varepsilon)L \int_{-\frac{L^2}{16}}^t e^{-\frac{L^2}{100(t-s)}} (t-s)^{-\frac{3}{2}} (-s)^{1-\frac{m^2}{2}} ds \\ &\leq (C\varepsilon + C(L)\varepsilon_1\varepsilon) \left(\frac{L}{4}\right)^{2-m^2} \\ &\quad + (C\varepsilon + C(L)\varepsilon_1\varepsilon)L^{-1} \int_{-\frac{L^2}{16}}^t (-s)^{\frac{1-m^2}{2}} ds \\ &\leq (C\varepsilon + C(L)\varepsilon_1\varepsilon) \left(\frac{L}{4}\right)^{2-m^2} \\ &\quad + (C\varepsilon + C(L)\varepsilon_1\varepsilon)L^{-1}m^{-2}(-t)^{\frac{3-m^2}{2}} \end{aligned}$$

for  $z \in [-20, 20]$  and  $t \in [-400, -1]$ . Therefore, we obtain

$$\begin{aligned} |v_m(z, t)| + |w_m(z, t)| &\leq (C\varepsilon + C(L)\varepsilon_1\varepsilon) \left(\frac{L^2}{16(-t)}\right)^{\frac{2-m^2}{2}} \\ &\quad + (C\varepsilon + C(L)\varepsilon_1\varepsilon)L^{-1}m^{-2} \end{aligned}$$

for  $z \in [-20, 20]$  and  $t \in [-400, -1]$ . Summation over  $m \geq 2$  yields

$$\left| \sum_{m=2}^{\infty} v_m(z, t) \right| + \left| \sum_{m=2}^{\infty} w_m(z, t) \right| \leq CL^{-1}\varepsilon + C(L)\varepsilon_1\varepsilon$$

for  $z \in [-20, 20]$  and  $t \in [-400, -1]$ .

*Step 6:* We next consider the modes with  $m = 1$ . We have

$$\frac{\partial}{\partial t} v_1 = \frac{\partial^2}{\partial z^2} v_1, \quad \frac{\partial}{\partial t} w_1 = \frac{\partial^2}{\partial z^2} w_1.$$

Moreover, given any point  $(z_0, t_0) \in [-\frac{L}{4}, \frac{L}{4}] \times [-\frac{L^2}{16}, -1]$ , there exist constants  $a_0, a_1, b_0, b_1$  (depending on  $z_0$  and  $t_0$ ) such that

$$|a_0|, |a_1|, |b_0|, |b_1| \leq C(L)\varepsilon$$

and

$$|v_1(z, t) - a_0 - a_1 z| + |w_1(z, t) - b_0 - b_1 z| \leq C\varepsilon(-t_0)^{\frac{1}{2}} + C(L)\varepsilon_1\varepsilon$$

for  $z \in [z_0 - (-t_0)^{\frac{1}{2}}, z_0 + (-t_0)^{\frac{1}{2}}]$  and  $t \in [2t_0, t_0]$ . Using standard interior estimates for the linear heat equation, we obtain

$$\left| \frac{\partial^2 v_1}{\partial z^2} \right| + \left| \frac{\partial^2 w_1}{\partial z^2} \right| \leq (C\varepsilon + C(L)\varepsilon_1\varepsilon) (-t)^{-\frac{1}{2}}$$

in the parabolic cylinder  $[-\frac{L}{4}, \frac{L}{4}] \times [-\frac{L^2}{16}, -1]$ . As above, we can use the solution formula for the one-dimensional heat equation with Dirichlet boundary

condition on the rectangle  $[-\frac{L}{4}, \frac{L}{4}] \times [-\frac{L^2}{16}, -1]$  to express  $\frac{\partial^2 v_1}{\partial z^2}$  and  $\frac{\partial^2 w_1}{\partial z^2}$  as integrals of initial and boundary data. This gives

$$\begin{aligned} & \left| \frac{\partial^2 v_1}{\partial z^2}(z, t) \right| + \left| \frac{\partial^2 w_1}{\partial z^2}(z, t) \right| \\ & \leq (C\varepsilon + C(L)\varepsilon_1\varepsilon) \left( \frac{L}{4} \right)^{-1} \\ & + (C\varepsilon + C(L)\varepsilon_1\varepsilon)L \int_{-\frac{L^2}{16}}^t e^{-\frac{L^2}{100(t-s)}} (t-s)^{-\frac{3}{2}} (-s)^{-\frac{1}{2}} ds \\ & \leq (C\varepsilon + C(L)\varepsilon_1\varepsilon) \left( \frac{L}{4} \right)^{-1} \\ & + (C\varepsilon + C(L)\varepsilon_1\varepsilon)L^{-2} \int_{-\frac{L^2}{16}}^t (-s)^{-\frac{1}{2}} ds \\ & \leq (C\varepsilon + C(L)\varepsilon_1\varepsilon) L^{-1} \end{aligned}$$

for  $z \in [-20, 20]$  and  $t \in [-400, -1]$ . Consequently, we can find real numbers  $A_0, A_1, B_0, B_1$  such that

$$|v_1(z, t) - A_0 - A_1 z| + |w_1(z, t) - B_0 - B_1 z| \leq CL^{-1}\varepsilon + C(L)\varepsilon_1\varepsilon$$

for  $z \in [-20, 20]$  and  $t \in [-400, -1]$ .

*Step 7:* Finally, we consider the mode with  $m = 0$ . Using identity

$$\int_0^{2\pi} u \sqrt{1 + r^{-2} \left( \frac{\partial r}{\partial \theta} \right)^2 + \left( \frac{\partial r}{\partial z} \right)^2} d\theta = \int_0^{2\pi} \frac{\partial r}{\partial \theta} d\theta = 0$$

together with the estimates  $|u| \leq C(L)\varepsilon$ ,  $|\frac{\partial r}{\partial \theta}| + |\frac{\partial r}{\partial z}| \leq C(L)\varepsilon_1$ , we obtain

$$\left| \int_0^{2\pi} u(\theta, z, t) d\theta \right| \leq C(L)\varepsilon_1\varepsilon,$$

hence

$$\left| \int_0^{2\pi} \tilde{u}(\theta, z, t) d\theta \right| \leq C(L)\varepsilon_1\varepsilon$$

for  $z \in [-20, 20]$  and  $t \in [-400, -1]$ .

*Step 8:* To summarize, we have shown that there exist  $A_0, A_1, B_0, B_1$  such that

$$|\tilde{u} - (A_0 + A_1 z) \cos \theta - (B_0 + B_1 z) \sin \theta| \leq CL^{-1}\varepsilon + C(L)\varepsilon_1\varepsilon$$

for  $z \in [-20, 20]$  and  $t \in [-400, -1]$ . This directly implies

$$|u - (A_0 + A_1 z) \cos \theta - (B_0 + B_1 z) \sin \theta| \leq CL^{-1}\varepsilon + C(L)\varepsilon_1\varepsilon$$

for  $z \in [-20, 20]$  and  $t \in [-400, -1]$ . In particular,  $|A_0| + |A_1| + |B_0| + |B_1| \leq C(L)\varepsilon$ . Hence, there exists a normalized rotation vector field  $\tilde{K}$  such that

$$|\langle \tilde{K}, \nu \rangle| \leq CL^{-1}\varepsilon + C(L)\varepsilon_1\varepsilon$$

in the parabolic neighborhood  $\hat{\mathcal{P}}(\bar{x}, -1, 10, 100)$ . Therefore,  $(\bar{x}, -1)$  is  $(CL^{-1}\varepsilon + C(L)\varepsilon_1\varepsilon)$ -symmetric. In particular, if we choose  $L$  sufficiently large and  $\varepsilon_1$  sufficiently small (depending on  $L$ ), then  $(\bar{x}, -1)$  is  $\frac{\varepsilon}{2}$ -symmetric.

5. PROOF OF ROTATIONAL SYMMETRY

Let  $M_t, t \in (-\infty, 0]$ , be a noncompact ancient solution of mean curvature flow in  $\mathbb{R}^3$  which is strictly convex and noncollapsed.

**Lemma 5.1.** *If  $-t$  is sufficiently large, then there exists a unique point  $p_t \in M_t$ , where the mean curvature attains its maximum. Moreover, the Hessian of the mean curvature at  $p_t$  is negative definite. In particular,  $p_t$  varies smoothly in  $t$ .*

**Proof.** We know that  $M_t \cap B_{8(-t)^{\frac{1}{2}}}(0)$  is a neck with radius  $(-2t)^{\frac{1}{2}}$ . The complement  $M_t \setminus B_{8(-t)^{\frac{1}{2}}}(0)$  has two connected components, one of which is compact and one of which is noncompact. On the noncompact connected component, the mean curvature is bounded from above by  $C(-t)^{-\frac{1}{2}}$ . On the other hand, we have shown in Proposition 3.3 that  $H_{\max}(t)$  is bounded away from 0. Consequently, if  $-t$  is sufficiently large, then the maximum of the mean curvature is attained at some point  $p_t \in M_t$ .

We next consider an arbitrary sequence  $t_j \rightarrow -\infty$ , and define  $M_t^{(j)} := M_{t+t_j} - p_{t_j}$ , where  $p_{t_j}$  is the point on  $M_{t_j}$  where the mean curvature attains its maximum. After passing to a subsequence if necessary, the sequence  $M_t^{(j)}$  converges in  $C_{\text{loc}}^\infty$  to a smooth eternal solution. Moreover, there exists a point on the limit solution where the mean curvature attains its space-time maximum. By work of Hamilton [10], the limit solution must be a translating soliton. By [11], the limit is the bowl soliton. Hence, if  $-t$  is sufficiently large, then  $p_t$  is the only point on  $M_t$  where the maximum of the mean curvature is attained, and the Hessian of the mean curvature at  $p_t$  is negative definite. This completes the proof of Lemma 5.1.

Let  $\varepsilon_1$  and  $L$  be the constants in the Neck Improvement Theorem. Since  $H_{\max}(t)$  is uniformly bounded from below, Proposition 3.1 in [13] implies that there exists a large constant  $\Lambda$  with the property that every point  $x \in M_t$  with  $|x - p_t| \geq \Lambda$  lies at the center of an  $\varepsilon_1$ -neck and satisfies  $H(x, t) |x - p_t| \geq 10^6 L$ .

**Lemma 5.2.** *There exists a time  $T < 0$  with the following property: Suppose that  $\bar{t} \leq T$ , and  $\bar{x}$  is a point on  $M_{\bar{t}}$  satisfying  $|\bar{x} - p_{\bar{t}}| \geq \Lambda$ . Then  $|\bar{x} - p_t| \geq |\bar{x} - p_{\bar{t}}|$  for all  $t \leq \bar{t}$ .*

**Proof.** If  $-t$  is sufficiently large, then  $M_t$  looks like the bowl soliton near the point  $p_t$ . Hence, if  $-t$  is sufficiently large, then the vector  $\frac{d}{dt}p_t$  is almost parallel to  $-\nu(p_t, t)$ . Consequently, we can find a time  $T < 0$  with the property that  $\langle x - p_t, \frac{d}{dt}p_t \rangle > 0$  whenever  $t \leq T$  and  $|x - p_t| \geq \Lambda$ . This implies  $\frac{d}{dt}|x - p_t| = -\langle \frac{x - p_t}{|x - p_t|}, \frac{d}{dt}p_t \rangle < 0$  whenever  $t \leq T$  and  $|x - p_t| \geq \Lambda$ .

We will show that  $T$  has the desired property. To prove this, we consider a time  $\bar{t} \leq T$  and a point  $\bar{x} \in M_{\bar{t}}$  such that  $|\bar{x} - p_{\bar{t}}| \geq \Lambda$ . We claim that  $|\bar{x} - p_t| \geq |\bar{x} - p_{\bar{t}}|$  for all  $t \leq \bar{t}$ . Indeed, if this is false, then we define

$\tilde{t} := \sup\{t \leq \bar{t} : |\bar{x} - p_t| < |\bar{x} - p_{\bar{t}}|\}$ . Clearly,  $\tilde{t} < \bar{t}$ , and  $|\bar{x} - p_t| \geq |\bar{x} - p_{\bar{t}}| \geq \Lambda$  for all  $t \in [\tilde{t}, \bar{t}]$ . In view of our choice of  $T$ , we obtain  $\frac{d}{dt}|\bar{x} - p_t| < 0$  for all  $t \in [\tilde{t}, \bar{t}]$ . Consequently,  $|\bar{x} - p_{\tilde{t}}| > |\bar{x} - p_{\bar{t}}|$ , which contradicts the definition of  $\tilde{t}$ . This completes the proof of Lemma 5.2.

**Proposition 5.3.** *If  $t \leq T$ ,  $x \in M_t$  and  $|x - p_t| \geq 2^{\frac{j}{400}} \Lambda$ , then  $(x, t)$  is  $2^{-j}\varepsilon_1$ -symmetric.*

**Proof.** We argue by induction on  $j$ . For  $j = 0$ , the assertion is true. Suppose now that  $j \geq 1$  and the assertion holds for  $j - 1$ . We claim that the assertion holds for  $j$ . Suppose this is false. Then there exists a time  $\bar{t} \leq T$  and a point  $\bar{x} \in M_{\bar{t}}$  such that  $|\bar{x} - p_{\bar{t}}| \geq 2^{\frac{j}{400}} \Lambda$  and  $(\bar{x}, \bar{t})$  is not  $2^{-j}\varepsilon_1$ -symmetric. By the Neck Improvement Theorem, there exists a point  $(x, t) \in \hat{\mathcal{P}}(\bar{x}, \bar{t}, L, L^2)$  such that either  $(x, t)$  is not  $2^{-j+1}\varepsilon_1$ -symmetric or  $(x, t)$  does not lie at the center of an  $\varepsilon_1$ -neck. In view of the induction hypothesis, we conclude that  $|x - p_t| \leq 2^{\frac{j-1}{400}} \Lambda$ . Since  $t \leq \bar{t} \leq T$ , Lemma 5.2 gives  $|\bar{x} - p_{\bar{t}}| \leq |\bar{x} - p_t|$ . Putting these facts together, we obtain

$$\begin{aligned} |\bar{x} - p_{\bar{t}}| &\leq |\bar{x} - p_t| \\ &\leq |x - p_t| + |x - \bar{x}| \\ &\leq 2^{\frac{j-1}{400}} \Lambda + 10 L H(\bar{x}, \bar{t})^{-1} \\ &\leq 2^{-\frac{1}{400}} |\bar{x} - p_{\bar{t}}| + \frac{1}{1000} |\bar{x} - p_{\bar{t}}| \\ &< |\bar{x} - p_{\bar{t}}|. \end{aligned}$$

This is a contradiction.

**Theorem 5.4.** *The surface  $M_t$  is rotationally symmetric for each  $t \leq T$ .*

**Proof.** Let us fix a time  $\bar{t} \leq T$ . For  $j$  sufficiently large, we denote by  $\Omega^{(j)}$  the set of all points  $(x, t)$  in space-time such that  $t \leq \bar{t}$  and  $|x - p_t| \leq 2^{\frac{j}{400}} \Lambda$ . If  $j$  is sufficiently large, then  $H(x, t) \geq 2 \cdot 2^{-\frac{j}{400}}$  for each point  $(x, t) \in \Omega^{(j)}$ . By Proposition 5.3, every point  $(x, t) \in \partial\Omega^{(j)}$  is  $2^{-j}\varepsilon_1$ -symmetric. In other words, for each point  $(x, t) \in \partial\Omega^{(j)}$ , there exists a normalized rotation vector field  $K^{(x,t)}$  such that  $|\langle K^{(x,t)}, \nu \rangle| H \leq 2^{-j}\varepsilon_1$  on  $\hat{\mathcal{P}}(x, t, 10, 100)$ . Using Lemma 4.2, we can control how the axis of rotation of  $K^{(x,t)}$  varies as we vary the point  $(x, t)$ . More precisely, if  $(x_1, t_1)$  and  $(x_2, t_2)$  are two points on  $\partial\Omega^{(j)}$  satisfying  $(x_2, t_2) \in \hat{\mathcal{P}}(x_1, t_1, 1, 1)$ , then

$$\begin{aligned} &\min \left\{ \sup_{B_{10H(x_2, t_2)^{-1}}(x_2)} |K^{(x_1, t_1)} - K^{(x_2, t_2)}|, \sup_{B_{10H(x_2, t_2)^{-1}}(x_2)} |K^{(x_1, t_1)} + K^{(x_2, t_2)}| \right\} \\ &\leq C 2^{-j} H(x_2, t_2)^{-1}. \end{aligned}$$

Hence, there exists a single normalized rotation vector field  $K^{(j)}$  with the following property: if  $(x, t)$  is a point in  $\partial\Omega^{(j)}$  satisfying  $\bar{t} - 2\frac{j}{100} \leq t \leq \bar{t}$ , then

$$\min\{|K^{(x,t)} - K^{(j)}|, |K^{(x,t)} + K^{(j)}|\} \leq C 2^{-\frac{j}{2}}$$

at the point  $(x, t)$ . This implies  $|\langle K^{(j)}, \nu \rangle| \leq C 2^{-\frac{j}{2}}$  for all points  $(x, t) \in \partial\Omega^{(j)}$  satisfying  $\bar{t} - 2\frac{j}{100} \leq t \leq \bar{t}$ . Moreover, we clearly have  $|\langle K^{(j)}, \nu \rangle| \leq C 2\frac{j}{100}$  for all points  $(x, t) \in \Omega^{(j)}$  with  $t = \bar{t} - 2\frac{j}{100}$ .

We now define a function  $f^{(j)} : \Omega^{(j)} \rightarrow \mathbb{R}$  by

$$f^{(j)} := \exp(-2^{-\frac{j}{200}}(\bar{t} - t)) \frac{\langle K^{(j)}, \nu \rangle}{H - 2^{-\frac{j}{400}}}.$$

Using the estimate for  $\langle K^{(j)}, \nu \rangle$ , we obtain

$$|f^{(j)}(x, t)| \leq \frac{C 2^{-\frac{j}{2}}}{2 \cdot 2^{-\frac{j}{400}} - 2^{-\frac{j}{400}}} \leq C 2^{-\frac{j}{4}}$$

for all points  $(x, t) \in \partial\Omega^{(j)}$  satisfying  $\bar{t} - 2\frac{j}{100} \leq t \leq \bar{t}$ , and

$$|f^{(j)}(x, t)| \leq \exp(-2^{-\frac{j}{200} + \frac{j}{100}}) \cdot \frac{C 2\frac{j}{100}}{2 \cdot 2^{-\frac{j}{400}} - 2^{-\frac{j}{400}}} \leq C 2^{-\frac{j}{4}}$$

for all points  $(x, t) \in \Omega^{(j)}$  with  $t = \bar{t} - 2\frac{j}{100}$ . Using the identities

$$\frac{\partial}{\partial t} H = \Delta H + |A|^2 H$$

and

$$\frac{\partial}{\partial t} \langle K^{(j)}, \nu \rangle = \Delta \langle K^{(j)}, \nu \rangle + |A|^2 \langle K^{(j)}, \nu \rangle,$$

we compute

$$\frac{\partial}{\partial t} f^{(j)} = \Delta f^{(j)} + \frac{2}{H - 2^{-\frac{j}{400}}} \langle \nabla H, \nabla f^{(j)} \rangle - 2^{-\frac{j}{400}} \left( \frac{|A|^2}{H - 2^{-\frac{j}{400}}} - 2^{-\frac{j}{400}} \right) f^{(j)}.$$

On the set  $\Omega^{(j)}$ , we have

$$\frac{|A|^2}{H - 2^{-\frac{j}{400}}} - 2^{-\frac{j}{400}} \geq \frac{1}{2} \frac{H^2}{H - 2^{-\frac{j}{400}}} - 2^{-\frac{j}{400}} \geq \frac{1}{2} H - 2^{-\frac{j}{400}} \geq 0.$$

Using the maximum principle, we conclude that

$$\begin{aligned} & \sup_{(x,t) \in \Omega^{(j)}, \bar{t} - 2\frac{j}{100} \leq t \leq \bar{t}} |f^{(j)}(x, t)| \\ & \leq \max \left\{ \sup_{(x,t) \in \partial\Omega^{(j)}, \bar{t} - 2\frac{j}{100} \leq t \leq \bar{t}} |f^{(j)}(x, t)|, \sup_{(x,t) \in \Omega^{(j)}, t = \bar{t} - 2\frac{j}{100}} |f^{(j)}(x, t)| \right\} \\ & \leq C 2^{-\frac{j}{4}}. \end{aligned}$$

This gives  $|\langle K^{(j)}, \nu \rangle| \leq C 2^{-\frac{j}{4}}$  for all points  $(x, t) \in \Omega^{(j)}$  with  $t = \bar{t}$ . From this, we deduce that the distance of the point  $p_{\bar{t}}$  from the axis of rotation of  $K^{(j)}$  is bounded from above by a uniform constant which is independent of  $j$ . Hence, if we send  $j \rightarrow \infty$ , the vector fields  $K^{(j)}$  converge to a normalized rotation vector field in  $\mathbb{R}^3$  which is tangential along  $M_{\bar{t}}$ . This completes the proof of Theorem 5.4.

Once we know that  $M_t$  is rotationally symmetric for  $-t$  sufficiently large, it follows from standard arguments that  $M_t$  is rotationally symmetric for all  $t$ :

**Proposition 5.5.** *Suppose that  $M_{\bar{t}}$  is rotationally symmetric for some  $\bar{t} \in (-\infty, 0]$ . Then, for each  $t \in [\bar{t}, 0]$ ,  $M_t$  is rotationally symmetric with the same axis.*

**Proof.** By Proposition 3.1 and Corollary 3.2, the flow  $M_t$ ,  $t \in (-\infty, 0]$ , has bounded curvature. Without loss of generality, we may assume that  $\sup_{M_t} |A|^2 \leq 2$  for each  $t \in (-\infty, 0]$ . If  $K$  is a rotation vector field in  $\mathbb{R}^3$ , then

$$\frac{\partial}{\partial t} \langle K, \nu \rangle = \Delta \langle K, \nu \rangle + |A|^2 \langle K, \nu \rangle.$$

Moreover, since  $|A|^2 \leq 2$ , the function  $\rho(x, t) := e^{8t}(|x|^2 + 1)$  satisfies

$$\frac{\partial}{\partial t} \rho > \Delta \rho + |A|^2 \rho$$

for  $t \in (-\infty, 0]$ . By the maximum principle, the quantity  $\sup_{M_t} \frac{|\langle K, \nu \rangle|}{\rho}$  is monotone decreasing for  $t \in (-\infty, 0]$ . In particular, if  $\langle K, \nu \rangle = 0$  at each point on  $M_{\bar{t}}$ , then  $\langle K, \nu \rangle = 0$  on  $M_t$  for all  $t \in [\bar{t}, 0]$ .

## 6. UNIQUENESS OF ANCIENT SOLUTIONS WITH ROTATIONAL SYMMETRY

Let  $M_t$  be an ancient solution satisfying the assumptions of Theorem 1.1. By the results in Section 5,  $M_t$  is rotationally symmetric. Without loss of generality, we may assume that  $M_t$  is symmetric with respect to the  $x_3$ -axis. Thus, there exists a function  $f(r, t)$  such that the solution  $M_t$  consists of the points  $(r \cos \theta, r \sin \theta, f(r, t)) \in \mathbb{R}^3$ . Moreover, the function  $f(r, t)$  satisfies the following evolution equation:

$$f_t = \frac{f_{rr}}{1 + f_r^2} + \frac{1}{r} f_r.$$

Note that  $f(r, t)$  may not be defined for all  $r$ .

Next, we can consider the radius  $r$  as a function of  $(z, t)$ . Namely, the radius function  $r(z, t)$  is defined by

$$f(r(z, t), t) = z.$$

Then  $r(z, t)$  satisfies the following equation (see also [2]):

$$r_t = \frac{r_{zz}}{1 + r_z^2} - \frac{1}{r}.$$

Note that the convexity of  $M_t$  yields

$$r > 0, \quad r_z > 0, \quad r_t < 0, \quad r_{zz} < 0.$$

Without loss of generality, we assume that the tip of  $M_0$  is at the origin. In other words,  $f(0, 0) = 0$  and  $r(0, 0) = 0$ .

Let  $q_t = (0, 0, f(0, t))$  denote the tip of  $M_t$ , and let  $H_{\text{tip}}(t)$  denote the mean curvature of  $M_t$  at the tip  $q_t$ . Using the Harnack inequality, we conclude that  $H_{\text{tip}}(t)$  is monotone increasing. In particular, the limit  $\mathcal{H} := \lim_{t \rightarrow -\infty} H_{\text{tip}}(t)$  exists. Using Proposition 3.3, we obtain  $\mathcal{H} > 0$ .

We first prove that  $f_t(r, t)$  is monotone increasing in  $t$ .

**Proposition 6.1.** *We have  $f_{tt}(r, t) \geq 0$  everywhere.*

**Proof.** Hamilton's Harnack inequality [10] implies that

$$\frac{\partial}{\partial t} H + 2V^i \nabla_i H + h_{ij} V^i V^j \geq 0$$

for every vector field  $V$ .

Let  $\omega = (0, 0, -1)$  denote the vertical vector field in  $\mathbb{R}^3$ , and let  $V = -H \langle \omega, \nu \rangle^{-1} \omega^{\text{tan}}$ . For this choice of  $V$ , the Harnack inequality takes the form

$$\left( \frac{\partial}{\partial t} + V^i \nabla_i \right) (H \langle \omega, \nu \rangle^{-1}) \geq 0.$$

A straightforward calculation gives

$$f_t(r, t) = H \langle \omega, \nu \rangle^{-1}$$

and

$$f_{tt}(r, t) = \left( \frac{\partial}{\partial t} + V^i \nabla_i \right) (H \langle \omega, \nu \rangle^{-1}).$$

Putting these facts together, the assertion follows.

We next show that  $f_t(r, t)$  is bounded from below.

**Proposition 6.2.** *We have  $f_t(r, t) \geq \mathcal{H}$  at each point in space-time. Moreover, for each  $r_0 > 0$ ,*

$$\lim_{t \rightarrow -\infty} \sup_{r \leq r_0} f_t(r, t) = \mathcal{H}.$$

**Proof.** We consider an arbitrary sequence  $t_j \rightarrow -\infty$ , and define  $M_t^{(j)} := M_{t+t_j} - q_{t_j}$ . We apply the compactness theorem for ancient solutions (cf. [12], Theorem 1.10) to the sequence  $M_t^{(j)}$ . Hence, after passing to a subsequence if necessary, the sequence  $M_t^{(j)}$  converges in  $C_{\text{loc}}^\infty$  to a smooth eternal solution, which is rotationally symmetric. Moreover, on the limit solution, the mean curvature at the tip equals  $\mathcal{H}$  at all times. Hence, equality holds

in the Harnack inequality. By work of Hamilton [10], the limit solution must be a soliton which is translating with speed  $\mathcal{H}$ . This directly implies

$$\lim_{j \rightarrow \infty} \sup_{r \leq r_0} |f_t(r, t_j) - \mathcal{H}| = 0$$

for every  $r_0 > 0$ . Since  $f_{tt}(r, t) \geq 0$  by Proposition 6.1, we conclude that  $f_t(r, t) \geq \mathcal{H}$  for all  $r$  and  $t$ .

We next prove that  $f_t(r, t)$  is monotone increasing in  $r$ .

**Proposition 6.3.** *We have  $f_{tr}(r, t) \geq 0$  everywhere.*

**Proof.** Let us fix a time  $t_0$  and a radius  $r_0$  such that  $f(r_0, t_0)$  is defined. For each  $T < t_0$ , we consider the parabolic region  $Q_T = \{x_1^2 + x_2^2 \leq r_0^2, t \in [T, t_0]\}$ . Using the equations

$$\frac{\partial}{\partial t} H = \Delta H + |A|^2 H$$

and

$$\frac{\partial}{\partial t} \langle \omega, \nu \rangle = \Delta \langle \omega, \nu \rangle + |A|^2 \langle \omega, \nu \rangle,$$

we conclude that the maximum  $\sup_{Q_T} H \langle \omega, \nu \rangle^{-1}$  must be attained on the parabolic boundary of  $Q_T$ . This gives

$$\begin{aligned} & \sup_{x_1^2 + x_2^2 \leq r_0^2, t = t_0} H \langle \omega, \nu \rangle^{-1} \\ & \leq \max \left\{ \sup_{x_1^2 + x_2^2 = r_0^2, T \leq t \leq t_0} H \langle \omega, \nu \rangle^{-1}, \sup_{x_1^2 + x_2^2 \leq r_0^2, t = T} H \langle \omega, \nu \rangle^{-1} \right\}. \end{aligned}$$

Since  $f_t(r, t) = H \langle \omega, \nu \rangle^{-1}$ , it follows that

$$\begin{aligned} \sup_{r \leq r_0} f_t(r, t_0) & \leq \max \left\{ \sup_{T \leq t \leq t_0} f_t(r_0, t), \sup_{r \leq r_0} f_t(r, T) \right\} \\ & = \max \left\{ f_t(r_0, t_0), \sup_{r \leq r_0} f_t(r, T) \right\}, \end{aligned}$$

where in the last step we have used Proposition 6.1. We now send  $T \rightarrow -\infty$ . Since  $\lim_{t \rightarrow -\infty} \sup_{r \leq r_0} f_t(r, t) = \mathcal{H}$ , we conclude that

$$\sup_{r \leq r_0} f_t(r, t_0) \leq \max\{f_t(r_0, t_0), \mathcal{H}\} = f_t(r_0, t_0).$$

This completes the proof of Proposition 6.3.

We recall that  $M_t$  is strictly convex and noncollapsed and  $H_{\text{tip}}(t)$  is bounded from below by  $\mathcal{H}$ . By Proposition 3.1 in [13], there exists a small constant  $\varepsilon_0 \in (0, \frac{1}{20})$  and a decreasing function  $\Lambda : (0, \varepsilon_0] \rightarrow \mathbb{R}$  such that given any  $\varepsilon \in (0, \varepsilon_0]$ , if  $|\bar{x} - q_t| \geq \Lambda(\varepsilon)$ , then  $(\bar{x}, \bar{t})$  is a center of  $\varepsilon$ -neck. (Alternatively, this can be deduced from Theorem 7.14 and Lemma 7.4 in [18].)

**Lemma 6.4.** *On every  $\varepsilon_0$ -neck,  $rr_z = \frac{r}{f_r} \leq (1 + 2\varepsilon_0)\mathcal{H}^{-1}$ .*

**Proof.** On an  $\varepsilon_0$ -neck, we have we have  $\frac{1}{f_r} = r_z \leq \varepsilon_0$ . Moreover, the principal curvature in radial direction is bounded by  $\frac{\varepsilon_0}{r}$ . This gives

$$\frac{f_{rr}}{(1 + f_r^2)^{\frac{3}{2}}} \leq \frac{\varepsilon_0}{r}.$$

Using Proposition 6.2, we obtain

$$\mathcal{H} \leq f_t = \frac{f_{rr}}{1 + f_r^2} + \frac{f_r}{r} \leq \frac{\varepsilon_0}{r} (1 + f_r^2)^{\frac{1}{2}} + \frac{f_r}{r} \leq (1 + 2\varepsilon_0) \frac{f_r}{r},$$

as claimed.

**Lemma 6.5.** *There exists a constant  $C_0 \geq 1$  such that  $r^m \left| \frac{\partial^m}{\partial z^m} r \right| \leq C_0$  holds for  $m = 1, 2, 3$  at center of  $\varepsilon_0$ -necks with  $r \geq 1$ .*

**Proof.** For  $m = 1$ , the assertion follows from Lemma 6.4. Let  $u = \langle \omega, \nu \rangle$ , where  $\omega = (0, 0, -1)$ . Then

$$\frac{\partial}{\partial t} u = \Delta u + |A|^2 u.$$

Moreover,  $u \leq \mathcal{H}^{-1} H$  by Proposition 6.2. Standard interior estimates imply that

$$|\nabla^m u|^2 \leq C H^{2m+2}$$

for  $m = 1, 2$  at the center of an  $\varepsilon_0$ -necks.

In the parametrization  $(z, \theta) \mapsto (r(z) \cos \theta, r(z) \sin \theta, z)$ , the induced metric is given by  $g_{zz} = 1 + r_z^2$ ,  $g_{z\theta} = 0$ ,  $g_{\theta\theta} = r^2$ . Moreover,  $u = r_z (1 + r_z^2)^{-\frac{1}{2}}$  and  $u_z = r_{zz} (1 + r_z^2)^{-\frac{3}{2}}$ . Hence,  $|\nabla u|^2 = g^{zz} u_z^2 = r_{zz}^2 (1 + r_z^2)^{-4}$ . In addition,  $r_z \leq \varepsilon_0$  and  $Hr \leq 1 + \varepsilon_0$  hold in every  $\varepsilon_0$ -neck. Therefore, the inequality  $|\nabla u|^2 \leq C H^4 \leq C r^{-4}$  implies  $r^4 r_{zz}^2 \leq C$ . Similarly,  $|\nabla^2 u|^2 \leq C H^6 \leq C r^{-6}$  gives  $r^6 r_{zz}^2 \leq C$ .

**Proposition 6.6.** *Let  $C_1 = 2 + 2\Lambda(\varepsilon_0) + 9\mathcal{H}^{-2}$ . If  $r \geq C_1$ , then  $0 \leq -r_{zz}(z, t) \leq C_2 r(z, t)^{-\frac{5}{2}}$ .*

**Proof.** Clearly,  $-r_{zz} \geq 0$  since  $M_t$  is convex. To prove the upper bound for  $-r_{zz}$ , let us fix a point  $(\bar{r}, \bar{t})$  satisfying  $\bar{r} \geq C_1 \geq 2$ , and let  $\bar{z} = f(\bar{r}, \bar{t})$ . Then we have  $\frac{1}{2}\bar{r} \geq \frac{1}{2}C_1 \geq \Lambda(\varepsilon_0)$  by definition of  $C_1$ . Hence, every point  $(x, t)$  with  $r = (x_1^2 + x_2^2)^{\frac{1}{2}} \geq \frac{1}{2}\bar{r}$  lies at the center of an  $\varepsilon_0$ -neck.

Using Lemma 6.4 and  $\varepsilon_0 \leq \frac{1}{20}$ , we obtain

$$\bar{z} - f\left(\frac{\bar{r}}{2}, \bar{t}\right) = \int_{\frac{\bar{r}}{2}}^{\bar{r}} f_r(r, \bar{t}) dr \geq \int_{\frac{\bar{r}}{2}}^{\bar{r}} \frac{\mathcal{H}}{1 + 2\varepsilon_0} r dr \geq \frac{1}{3} \mathcal{H} \bar{r}^2.$$

Since  $\bar{r} \geq C_1 \geq 9\mathcal{H}^{-2}$ , it follows that  $f\left(\frac{\bar{r}}{2}, \bar{t}\right) \leq \bar{z} - \bar{r}^{\frac{3}{2}}$ . In other words,  $r(z, \bar{t}) \geq \frac{\bar{r}}{2}$  for  $z \in [\bar{z} - \bar{r}^{\frac{3}{2}}, \bar{z} + \bar{r}^{\frac{3}{2}}]$ . Since  $r(z, t)$  is decreasing in  $t$ , it follows

that  $r(z, t) \geq \frac{\bar{r}}{2}$  for  $(z, t) \in Q = [\bar{z} - \bar{r}^{\frac{3}{2}}, \bar{z} + \bar{r}^{\frac{3}{2}}] \times [\bar{t} - \bar{r}^3, \bar{t}]$ . Hence, every point  $(x, t)$  with  $(x_3, t) \in Q$  lies at the center of an  $\varepsilon_0$ -neck.

We next consider the evolution equation of  $rr_z$ . We compute

$$(rr_z)_t = (rr_z)_{zz} - \frac{(2 + 3r_z^2)r_z r_{zz} + rr_z^2 r_{zzz}}{1 + r_z^2} - \frac{2rr_z r_{zz}^2}{(1 + r_z^2)^2}.$$

Using Lemma 6.4 and Lemma 6.5, we obtain

$$\sup_Q |rr_z| \leq C$$

and

$$\sup_Q |(rr_z)_t - (rr_z)_{zz}| \leq C\bar{r}^{-3}.$$

Standard interior estimates for the linear heat equation give

$$|(rr_z)_z| \leq C\bar{r}^{-\frac{3}{2}} \sup_Q |rr_z| + C\bar{r}^{\frac{3}{2}} \sup_Q |(rr_z)_t - (rr_z)_{zz}| \leq C\bar{r}^{-\frac{3}{2}}$$

at  $(\bar{z}, \bar{t})$ . Thus,  $|r_{zz}| \leq C\bar{r}^{-\frac{5}{2}}$  at  $(\bar{z}, \bar{t})$ . This completes the proof of Proposition 6.6.

For each  $z < 0$ , we define a real number  $\mathcal{T}(z)$  by

$$r(z, t) > 0 \quad \text{for } t < \mathcal{T}(z), \quad \lim_{t \rightarrow \mathcal{T}(z)} r(z, t) = 0.$$

The following result allows us to estimate  $r(z, t)$  in terms of  $\mathcal{T}(z) - t$ .

**Corollary 6.7.** *We have*

$$2[\mathcal{T}(z) - t] \leq r(z, t)^2 \leq 2[\mathcal{T}(z) - t] + 8C_2[\mathcal{T}(z) - t]^{\frac{1}{4}} + C_1^2$$

if  $z < 0$  and  $r(z, t)$  is sufficiently large.

**Proof.** We again fix a point  $(\bar{z}, \bar{t})$ . Since  $(r^2 + 2t)_t = \frac{2rr_{zz}}{1+r_z^2} < 0$ , we have

$$r(\bar{z}, \bar{t})^2 \geq 2[\mathcal{T}(\bar{z}) - \bar{t}].$$

Moreover, Proposition 6.6 implies that  $(r^2 + 2t)_t = \frac{2rr_{zz}}{1+r_z^2} \geq -2C_2r^{-\frac{3}{2}}$  whenever  $r \geq C_1$ . Let  $\tilde{t} \leq \mathcal{T}(\bar{z})$  denote the time when  $r(\bar{z}, \tilde{t}) = C_1$ . Since  $r(\bar{z}, t)$  is a decreasing function of  $t$ ,  $r(\bar{z}, t) \leq C_1$  for  $t \leq \tilde{t}$ . Therefore,

$$\begin{aligned} r(\bar{z}, \bar{t})^2 &= C_1^2 + 2(\tilde{t} - \bar{t}) - \int_{\tilde{t}}^{\bar{t}} (r(\bar{z}, t)^2 + 2t)_t dt \\ &\leq C_1^2 + 2(\tilde{t} - \bar{t}) + 2C_2 \int_{\tilde{t}}^{\bar{t}} r(\bar{z}, t)^{-\frac{3}{2}} dt \\ &\leq C_1^2 + 2(\tilde{t} - \bar{t}) + 2C_2 \int_{\tilde{t}}^{\bar{t}} [\mathcal{T}(\bar{z}) - t]^{-\frac{3}{4}} dt \\ &\leq C_1^2 + 2(\tilde{t} - \bar{t}) - 8C_2[\mathcal{T}(\bar{z}) - \tilde{t}]^{\frac{1}{4}} + 8C_2[\mathcal{T}(\bar{z}) - \bar{t}]^{\frac{1}{4}} \\ &\leq C_1^2 + 2[\mathcal{T}(\bar{z}) - \bar{t}] + 8C_2[\mathcal{T}(\bar{z}) - \bar{t}]^{\frac{1}{4}}, \end{aligned}$$

as claimed.

Lemma 6.4 gives a sharp upper bound for the limit of  $rr_z$ . More precisely,  $\limsup_{z \rightarrow \infty} r(z, t)r_z(z, t) \leq \mathcal{H}^{-1}$  for each  $t$ . We next establish a lower bound for  $\liminf_{z \rightarrow \infty} r(z, t)r_z(z, t)$ . To derive this estimate, we need a lower bound for  $r(0, t)r_z(0, t)$ .

**Lemma 6.8.** *Let  $\delta > 0$  be arbitrary. Then*

$$r(0, t)r_z(0, t) \geq \mathcal{H}^{-1} - \delta$$

*whenever  $-t$  is sufficiently large.*

**Proof.** In the following, we assume that  $-t$  is sufficiently large so that  $R = r(0, t) \geq C_1$ . Consequently, every point  $(x, t)$  with  $x_3 = 0$  lies at the center of an  $\varepsilon_0$ -neck. This implies  $|r(z, t) - R| \leq \varepsilon_0 R$  for  $|z| \leq 2R$ . Recall that  $rr_z \leq (1 + 2\varepsilon_0)\mathcal{H}^{-1}$  by Lemma 6.4, and  $|(rr_z)_z| = |rr_{zz} + r_z^2| \leq C_3 R^{-\frac{3}{2}}$  for some constant  $C_3$  by Proposition 6.6. Hence, if we choose  $-t$  sufficiently large so that  $R^{\frac{1}{2}} \geq 4C_3\delta^{-1}$ , then we obtain

$$|r(z, t)r_z(z, t) - r(0, t)r_z(0, t)| \leq 2C_3 R^{-\frac{1}{2}} \leq \frac{\delta}{2}$$

for all  $z \in [-2R, 2R]$ .

It follows from Corollary 6.7 that

$$\begin{aligned} r(-R, t)^2 &\geq 2[\mathcal{T}(-R) - t], \\ r(-2R, t)^2 &\geq 2[\mathcal{T}(-2R) - t], \end{aligned}$$

and

$$\begin{aligned} r(-2R, t)^2 &\leq 2[\mathcal{T}(-2R) - t] + 8C_2[\mathcal{T}(-2R) - t]^{\frac{1}{4}} + C_1^2 \\ &\leq 2[\mathcal{T}(-2R) - t] + 8C_2 r(-2R, t)^{\frac{1}{2}} + C_1^2 \\ &\leq 2[\mathcal{T}(-2R) - t] + 8C_2 R^{\frac{1}{2}} + C_1^2. \end{aligned}$$

This implies

$$r(-R, t)^2 - r(-2R, t)^2 \geq 2[\mathcal{T}(-R) - \mathcal{T}(-2R)] - 8C_2 R^{\frac{1}{2}} - C_1^2.$$

Moreover, if  $R$  is sufficiently large, then

$$\mathcal{T}(-R) - \mathcal{T}(-2R) \geq \left(\mathcal{H}^{-1} - \frac{\delta}{2}\right) R.$$

This gives

$$r(-R, t)^2 - r(-2R, t)^2 \geq 2\left(\mathcal{H}^{-1} - \frac{\delta}{2}\right) R$$

if  $-t$  is sufficiently large. Hence, if  $-t$  is sufficiently large, then

$$\sup_{z \in [-2R, R]} r(z, t)r_z(z, t) \geq \mathcal{H}^{-1} - \frac{\delta}{2}.$$

Putting these facts together, we conclude that

$$r(0, t)r_z(0, t) \geq \mathcal{H}^{-1} - \delta$$

whenever  $-t$  is sufficiently large.

We next recall a solution  $\psi(z, t)$  to the heat equation satisfying Dirichlet boundary condition on the half line.

**Proposition 6.9.** *We define a smooth function  $\psi : (0, \infty) \times (0, \infty) \rightarrow \mathbb{R}$  by*

$$\psi(z, t) = \frac{1}{\sqrt{4\pi t}} \int_0^\infty (e^{-\frac{(z-y)^2}{4t}} - e^{-\frac{(z+y)^2}{4t}}) dy.$$

*Then  $\psi$  is a solution to the heat equation  $\psi_t = \psi_{zz}$ . Moreover, for each  $z > 0$  and  $t > 0$  we have  $\psi_{zz}(z, t) < 0$  and*

$$\lim_{z \rightarrow 0} \psi(z, t) = 0, \quad \lim_{z \rightarrow \infty} \psi(z, t) = 1, \quad \lim_{t \rightarrow 0} \psi(z, t) = 1, \quad \lim_{t \rightarrow \infty} \psi(z, t) = 0.$$

**Proof.** We only need to show  $\psi_{zz} < 0$ . Direct computations yield

$$\begin{aligned} \psi_{zz} &= \frac{1}{\sqrt{4\pi t}} \left[ - \int_0^\infty \left( \frac{1}{2t} - \frac{(z-y)^2}{4t^2} \right) e^{-\frac{(z-y)^2}{4t}} dy + \int_0^\infty \left( \frac{1}{2t} - \frac{(z+y)^2}{4t^2} \right) e^{-\frac{(z+y)^2}{4t}} dy \right] \\ &= \frac{1}{\sqrt{8\pi t^2}} \left[ - \int_{-\frac{z}{\sqrt{2t}}}^\infty (1 - \xi^2) e^{-\frac{\xi^2}{2}} d\xi + \int_{\frac{z}{\sqrt{2t}}}^\infty (1 - \xi^2) e^{-\frac{\xi^2}{2}} d\xi \right] \\ &= - \frac{1}{\sqrt{8\pi t^2}} \int_{-\frac{z}{\sqrt{2t}}}^{\frac{z}{\sqrt{2t}}} (1 - \xi^2) e^{-\frac{\xi^2}{2}} d\xi. \end{aligned}$$

Clearly,  $\psi_{zz} < 0$  for  $0 < z \leq \sqrt{2t}$ . Moreover,  $\psi_{zzz} > 0$  for  $z \geq \sqrt{2t}$ , and

$$\lim_{z \rightarrow \infty} \psi_{zz}(z, t) = - \frac{1}{\sqrt{8\pi t^2}} \int_{-\infty}^\infty (1 - \xi^2) e^{-\frac{\xi^2}{2}} d\xi = 0.$$

Therefore,  $\psi_{zz} < 0$  also holds for  $z \geq \sqrt{2t}$ . This completes the proof of Proposition 6.9.

**Proposition 6.10.** *Given  $\delta > 0$ , there exists a time  $\bar{t} \in (-\infty, 0]$  (depending on  $\delta$ ) such that*

$$r(z, t)r_z(z, t) \geq \mathcal{H}^{-1} - 2\delta,$$

*holds for all  $z \geq 0$  and  $t \leq \bar{t}$ .*

**Proof.** By Proposition 6.6, we have  $1 + rr_{zz} \geq 0$  for  $r \geq C_1 + C_2$ . This implies

$$(rr_z)_t = \frac{(rr_z)_{zz}}{1 + r_z^2} - \frac{2r_z r_{zz}(1 + r_z^2 + rr_{zz})}{(1 + r_z^2)^2} \geq \frac{(rr_z)_{zz}}{1 + r_z^2}$$

for  $r \geq C_1 + C_2$ . By Lemma 6.8, we can choose  $\bar{t}$  large enough so that  $r(0, t)r_z(0, t) \geq \mathcal{H}^{-1} - \delta$  for  $t \leq \bar{t}$ . Moreover, by a suitable choice of  $\bar{t}$  we can arrange that  $r(z, t) \geq C_1 + C_2$  for all  $z \geq 0$  and  $t \leq \bar{t}$ . For each  $s < \bar{t}$ , we define a barrier function  $\psi^{\delta, s}(z, t)$  by

$$\psi^{\delta, s}(z, t) = \mathcal{H}^{-1} - 2\delta - \mathcal{H}^{-1} \psi(2z, t - s)$$

for  $t \in (s, \bar{t}]$ . We claim that  $rr_z > \psi^{\delta, s}$  for all  $z \geq 0$  and all  $t \in (s, \bar{t}]$ .

By our choice of  $\bar{t}$ ,  $r(0, t)r_z(0, t) \geq \mathcal{H}^{-1} - \delta > \limsup_{z \rightarrow 0} \psi^{\delta, s}(z, t)$  for each  $t \in (s, \bar{t}]$ . Moreover,  $\liminf_{z \rightarrow \infty} r(z, t)r_z(z, t) \geq 0 > \limsup_{z \rightarrow \infty} \psi^{\delta, s}(z, t)$  for each  $t \in (s, \bar{t}]$ . Finally, Proposition 6.9 gives  $r(z, s)r_z(z, s) \geq 0 > \limsup_{t \rightarrow s} \psi^{\delta, s}(z, t)$  for each  $z > 0$ .

Thus, if the inequality  $rr_z > \psi^{\delta, s}$  fails, there exists some point  $(z_0, t_0) \in (0, \infty) \times (s, \bar{t}]$  such that  $r(z_0, t_0)r_z(z_0, t_0) = \psi^{\delta, s}(z_0, t_0)$  and  $r(z, t)r_z(z, t) \geq \psi^{\delta, s}(z, t)$  for all  $z \geq 0$  and all  $t \in (s, t_0]$ . Then, at the point  $(z_0, t_0)$  we have

$$\frac{(\psi^{\delta, s})_{zz}}{1 + r_z^2} \leq \frac{(rr_z)_{zz}}{1 + r_z^2} \leq (rr_z)_t \leq (\psi^{\delta, s})_t = \frac{1}{4}(\psi^{\delta, s})_{zz}.$$

This contradicts the fact that  $r_z \leq \varepsilon_0$  and  $(\psi^{\delta, s})_{zz} > 0$ .

Thus, we conclude that  $rr_z > \psi^{\delta, s}$  for all  $z \geq 0$  and all  $t \in (s, \bar{t}]$ . Sending  $s \rightarrow -\infty$ , we obtain  $rr_z \geq \mathcal{H}^{-1} - 2\delta$  for all  $z \geq 0$  and all  $t \leq \bar{t}$ .

**Corollary 6.11.** *We can find a time  $T \in (-\infty, 0]$  such that  $r(z, t)^2 \geq \mathcal{H}^{-1}z$  for all  $z \geq 0$  and  $t \leq T$ . In particular, if  $t \leq T$ , then the function  $f(r, t)$  is defined for all  $r \in [0, \infty)$ .*

**Proof.** By Proposition 6.10, we can find a time  $T \in (-\infty, 0]$  such that  $r(z, t)r_z(z, t) \geq \frac{1}{2}\mathcal{H}^{-1}$  for all  $z \geq 0$  and all  $t \leq T$ . From this, the assertion follows easily.

After these preparations, we now compute the limit  $\lim_{z \rightarrow \infty} r(z, t)r_z(z, t)$ .

**Proposition 6.12.** *For each  $t \leq T$ , we have  $\lim_{z \rightarrow \infty} r(z, t)r_z(z, t) = \mathcal{H}^{-1}$ .*

**Proof.** Lemma 6.4 gives  $\limsup_{z \rightarrow \infty} r(z, t)r_z(z, t) \leq \mathcal{H}^{-1}$ . So, it is enough to show that  $\liminf_{z \rightarrow \infty} r(z, t)r_z(z, t) \geq \mathcal{H}^{-1}$  for each  $t \leq T$ . Given any  $\delta > 0$ , Proposition 6.10 implies that we can find a number  $\bar{t} \leq T$  such that  $\liminf_{z \rightarrow \infty} r(z, \bar{t})r_z(z, \bar{t}) \geq \mathcal{H}^{-1} - 2\delta$ . Moreover, Lemma 6.5 guarantees that

$$|(rr_z)_t| = |r_t r_z + rr_{zt}| = \left| \frac{r_z r_{zz}}{1 + r_z^2} + \frac{rr_{zzz}}{1 + r_z^2} - \frac{2rr_z r_{zz}^2}{(1 + r_z^2)^2} \right| \leq \frac{4C_0^3}{r^2}$$

for  $r \geq C_1$ . Using Corollary 6.11, we obtain

$$\liminf_{z \rightarrow \infty} r(z, t)r_z(z, t) = \liminf_{z \rightarrow \infty} r(z, \bar{t})r_z(z, \bar{t}) \geq \mathcal{H}^{-1} - 2\delta$$

for each  $t \leq T$ . Since  $\delta > 0$  is arbitrary, we conclude that  $\liminf_{z \rightarrow \infty} r(z, t)r_z(z, t) \geq \mathcal{H}^{-1}$  for each  $t \leq T$ . This completes the proof of Proposition 6.12.

**Theorem 6.13.** *For each  $t \leq T$ ,  $M_t$  is a translating soliton.*

**Proof.** Since  $rr_z = \frac{r}{f_r}$ , Proposition 6.12 implies

$$\lim_{r \rightarrow \infty} \frac{f_r(r, t)}{r} = \mathcal{H}$$

for each  $t \leq T$ . Using the evolution equation for  $f(r, t)$ , we obtain

$$\lim_{r \rightarrow \infty} f_t(r, t) = \lim_{r \rightarrow \infty} \frac{f_r(r, t)}{r} = \mathcal{H}$$

for each  $t \leq T$ . Using Proposition 6.3, we conclude that  $f_t(r, t) \leq \mathcal{H}$  for all  $r \geq 0$  and all  $t \leq T$ . Therefore, Proposition 6.2 gives  $f_t(r, t) = \mathcal{H}$  for all  $r \geq 0$  and all  $t \leq T$ . Consequently,  $M_t$  is a translating soliton for each  $t \leq T$ .

Once we know that  $M_t$  is a translating soliton for  $-t$  sufficiently large, it follows from standard arguments that  $M_t$  is a translating soliton for all  $t$ :

**Proposition 6.14.** *Suppose that  $M_{\bar{t}}$  is a translating soliton for some  $\bar{t} \in (-\infty, 0]$ . Then  $M_t$  is a translating soliton for all  $t \in [\bar{t}, 0]$ .*

**Proof.** By Proposition 3.1 and Corollary 3.2, the flow  $M_t$ ,  $t \in (-\infty, 0]$ , has bounded curvature. Without loss of generality, we may assume that  $\sup_{M_t} |A|^2 \leq 2$  for each  $t \in (-\infty, 0]$ . If  $\omega$  is a fixed vector in  $\mathbb{R}^3$ , then

$$\frac{\partial}{\partial t}(H - \langle \omega, \nu \rangle) = \Delta(H - \langle \omega, \nu \rangle) + |A|^2(H - \langle \omega, \nu \rangle).$$

Moreover, since  $|A|^2 \leq 2$ , the function  $\rho(x, t) := e^{8t}(|x|^2 + 1)$  satisfies

$$\frac{\partial}{\partial t}\rho > \Delta\rho + |A|^2\rho$$

for  $t \in (-\infty, 0]$ . By the maximum principle, the quantity  $\sup_{M_t} \frac{|H - \langle \omega, \nu \rangle|}{\rho}$  is monotone decreasing for  $t \in (-\infty, 0]$ . In particular, if  $H = \langle \omega, \nu \rangle$  at each point on  $M_{\bar{t}}$ , then  $H = \langle \omega, \nu \rangle$  on  $M_t$  for all  $t \in [\bar{t}, 0]$ .

## REFERENCES

- [1] S. Altschuler and L.F. Wu, *Translating surfaces of the non-parametric mean curvature flow with prescribed contact angle*, Calc. Var. 2, 101–111 (1994)
- [2] S. Angenent, P. Daskalopoulos, and N. Šešum, *Unique asymptotics of ancient convex mean curvature flow solutions*, arXiv:1503.01178v3
- [3] S. Brendle, *Rotational symmetry of self-similar solutions to the Ricci flow*, Invent. Math. 194, 731–764 (2013)
- [4] S. Brendle, *A sharp bound for the inscribed radius under mean curvature flow*, Invent. Math. 202, 217–237 (2015)
- [5] S. Brendle and K. Choi, *Uniqueness of convex ancient solutions to mean curvature flow in higher dimensions*, arXiv:1804.00018
- [6] S. Brendle and G. Huisken, *Mean curvature flow with surgery of mean convex surfaces in  $\mathbb{R}^3$* , Invent. Math. 203, 615–654 (2016)
- [7] T.H. Colding and W.P. Minicozzi, *Uniqueness of blowups and Lojasiewicz inequalities*, Ann. of Math. 182, 221–285 (2015)
- [8] P. Daskalopoulos, R. Hamilton, and N. Šešum, *Classification of compact ancient solutions to the curve shortening flow*, J. Diff. Geom. 84, 455–464 (2010)
- [9] P. Daskalopoulos, R. Hamilton, and N. Šešum, *Classification of ancient compact solutions to the Ricci flow on surfaces*, J. Diff. Geom. 91, 171–214 (2012)
- [10] R. Hamilton, *Harnack estimate for the mean curvature flow*, J. Diff. Geom. 41, 215–226 (1995)

- [11] R. Haslhofer, *Uniqueness of the bowl soliton*, *Geom. Topol.* 19, 2393–2406 (2015)
- [12] R. Haslhofer and B. Kleiner, *Mean curvature flow of mean convex hypersurfaces*, *Comm. Pure Appl. Math.* 70, 511–546 (2017)
- [13] R. Haslhofer and B. Kleiner, *Mean curvature flow with surgery*, *Duke Math. J.* 166, 1591–1626 (2017)
- [14] G. Huisken, *Flow by mean curvature of convex surfaces into spheres*, *J. Diff. Geom.* 20, 237–266 (1984)
- [15] G. Huisken, *Asymptotic behavior for singularities of mean curvature flow*, *J. Diff. Geom.* 31, 285–299 (1990)
- [16] G. Huisken and C. Sinestrari, *Mean curvature flow singularities for mean convex surfaces*, *Calc. Var.* 8, 1–14 (1999)
- [17] G. Huisken and C. Sinestrari, *Convexity estimates for mean curvature flow and singularities of mean convex surfaces*, *Acta Math.* 183, 45–70 (1999)
- [18] G. Huisken and C. Sinestrari, *Mean curvature flow with surgeries of two-convex hypersurfaces*, *Invent. Math.* 175, 137–221 (2009)
- [19] G. Huisken and C. Sinestrari, *Convex ancient solutions of the mean curvature flow*, *J. Diff. Geom.* 101, 267–287 (2015)
- [20] S. Kleene and N.M. Møller, *Self-shrinkers with a rotational symmetry*, *Trans. Amer. Math. Soc.* 366, 3943–3963 (2014)
- [21] F. Merle and H. Zaag, *Optimal estimates for blowup rate and behavior for nonlinear heat equations*, *Comm. Pure Appl. Math.* 51, 139–196 (1998)
- [22] G. Perelman, *The entropy formula for the Ricci flow and its geometric applications*, arxiv:0211159
- [23] W. Sheng and X.J. Wang, *Singularity profile in the mean curvature flow*, *Methods Appl. Anal.* 16, 139–155 (2009)
- [24] X.J. Wang, *Convex solutions to the mean curvature flow*, *Ann. of Math.* 173, 1185–1239 (2011)
- [25] B. White, *The size of the singular set in mean curvature flow of mean convex sets*, *J. Amer. Math. Soc.* 13, 665–695 (2000)
- [26] B. White, *The nature of singularities in mean curvature flow of mean convex sets*, *J. Amer. Math. Soc.* 16, 123–138 (2003)

DEPARTMENT OF MATHEMATICS, COLUMBIA UNIVERSITY, 2990 BROADWAY, NEW YORK, NY 10027, USA.

DEPARTMENT OF MATHEMATICS, MASSACHUSETTS INSTITUTE OF TECHNOLOGY, 77 MASSACHUSETTS AVENUE, CAMBRIDGE MA 02138, USA.