

An alternative approach for stability analysis of discrete time nonlinear dynamical systems

R. Bouyekhf^{a**} and L.T. Gruyitch^b

^aNanomedicine Lab, Université Bourgogne Franche-Comté,
UTBM, F-90010 Belfort, France

^bJohn Kennedy 31/15, Belgrade 11070, Serbia

Abstract

The paper endeavours to solve the problem of the necessary and sufficient conditions for testing asymptotic stability of the equilibrium state without using a positive definite or semi-definite Lyapunov function for time-invariant nonlinear discrete-time dynamical systems. The solution is based on the concept of the \mathbf{G} -functions introduced in this paper. As a result, new necessary and sufficient conditions for asymptotic stability of such systems and an estimation or the exact determination of the asymptotic stability domain of the state $x = 0$ are established. Examples are worked out to illustrate the results.

keywords Discrete-time nonlinear systems, asymptotic stability, stability domain, \mathbf{G} -functions.

1 Introduction

So far, the study of the stability of discrete-time nonlinear systems has been elaborated by the Lyapunov method by following and accepting its basis established for continuous-time systems, which is the concept of definite [8] and semi-definite functions [7], [10]. It is well known that two of the main properties of such functions are their continuity and their global minimum (usually zero) value at the equilibrium state. Hence, the Lyapunov method reduces the stability analysis to a search for a Lyapunov function which is decreasing along trajectories of the system as long as $x \neq 0$. Unfortunately, even in some simple cases, it can be very difficult to find such an appropriate function.

With the best of our knowledge, the efforts that have been made up to the present time in the framework of discrete-time systems are limited only to

**CONTACT R. Bouyekhf. Email: rachid.bouyekhf@utbm.fr

some special classes of systems (linear, polynomial systems [2], [13], homogeneous [15], exponential stable systems [3], [4]). Generally speaking, for a given discrete time nonlinear system one has little hope to find the Lyapunov function unless the system belongs to one of the special classes. Hence, two questions arise naturally: When the method of the first approximation fails, why the stability test must be restricted in general to only Lyapunov functions? Is it possible to broaden Lyapunov's method to certain functions without requiring their positive definiteness or semi-definiteness in the framework of discrete-time nonlinear systems? Such a possibility will enable us to enlarge the scope of the stability to other functions which have no mathematical reason to be excluded.

This paper endeavours to give positive answer to the above mentioned question. More precisely, we present analytical tests for the asymptotic stability and an estimation or the exact determination of the asymptotic stability domain of the equilibrium state for discrete-time nonlinear systems. The tests have to be carried out by using the concept of the \mathbf{G} -functions rather than that of the positive definite or semi-definite Lyapunov functions.

The paper is organized as follows. Section 2 is devoted to the terminology and notations and summarizes some results that are needed in the subsequent development. Section 3 defines the concept of \mathbf{G} -functions on which our approach is based and discusses their properties. The final results are given in Section 4 with illustrative examples.

2 System description and definitions

We consider discrete-time nonlinear systems of the form

$$x(k+1) = f(x(k)), \quad x(0) = x_0, \quad (1)$$

where $x(k) \in \mathbb{R}^n$ is the state vector at discrete time $k \in \mathbb{N}$ defined on an open neighbourhood $S \subseteq \mathbb{R}^n$ of $x = 0$ and $f : S \rightarrow S$ is a nonlinear vector function such that $f(0) = 0$. The solution of the system (1), which starts at $x_0 \in S$ is defined by $x(k; x_0) \equiv x(k) = f^k(x_0)$, where f^k is the k -th multiple composition of the mapping f . Throughout the paper, we shall use the following notations: $\mathbb{R}_{\geq 0} = [0, +\infty)$ and $\mathbb{R}_{> 0} = (0, +\infty)$. The set $B_\varepsilon = \{x : \|x\| < \varepsilon\}$ is an open ball with the radius ε and center at the origin. A neighbourhood of $x = 0$ is a set N , which contains an open ball B_ε , i.e.; $0 \in B_\varepsilon \subset N$. We denote by $\text{Cl}A$, ∂A the closure and the boundary of the set A , respectively. Let $V : \mathbb{R}^n \rightarrow \mathbb{R}$ be a real function. $\Delta V(x) = V(f(x)) - V(x)$ denotes the variation of the function V along the solution of the system. For the sake of clarity we state the following:

Definition 2.1 The equilibrium state $x = 0$ of system (1) is:

- (i) stable if, and only if, for every $\varepsilon > 0$ there exists $\delta(\varepsilon) > 0$ such that $\|x_0\| < \delta(\varepsilon)$ implies $\|x(k; x_0)\| < \varepsilon, \forall k \in \mathbb{N}$.
- (ii) attractive if, and only if, there exists $\eta > 0$ and for every $\rho > 0$ there exists $K(x_0, \rho) > 0$ such that $\|x_0\| < \eta$ implies $\|x(k; x_0)\| < \rho, \quad \forall k > K(x_0, \rho)$, or equivalently: $\lim_{k \rightarrow +\infty} \|x(k, x_0)\| = 0$.

(iii) asymptotically stable if, and only if, it is both stable and attractive.

The notions of the domain of stability, the domain of attraction and the domain of asymptotic stability used in this paper preserve the same meaning as proposed and used in [5], [6], [12].

Definition 2.2 The equilibrium state $x = 0$ of system (1) has the domain of:

- 1) stability $D_s \subseteq S$ if, and only if, both:
 - (i) for every $\varepsilon > 0$ there is a neighbourhood $D_s(\varepsilon) \subseteq S$ of $x = 0$ depending on ε such that $\|x(k; x_0)\| < \varepsilon$ for all $k \in \mathbb{N}$, provided only that $x_0 \in D_s(\varepsilon)$,
 - (ii) $D_s = \bigcup_{\varepsilon > 0} D_s(\varepsilon)$.
- 2) attraction $D_a \subseteq S$ if, and only if, the set D_a is the largest neighbourhood of $x = 0$ such that for every $x_0 \in D_a$ and $\rho > 0$ there exists $K(x_0, \rho) > 0$ guaranteeing that $\|x(k; x_0)\| < \rho$ for every $k > K(x_0, \rho)$.
- 3) asymptotic stability $D_{as} \subseteq S$ if, and only if, both:
 - (i) $D_{as} = D_s \cap D_a$,
 - (ii) the set D_{as} is a neighbourhood of $x = 0$.

Remark 1: It is important to note that, the domain of stability D_s introduced in Definition 2.2 is defined as the union of all $D_s(\varepsilon)$, where $D_s(\varepsilon)$ is a neighbourhood of $x = 0$ relative to each $\varepsilon > 0$. Besides, the domain of attraction is defined as the largest neighbourhood of $x = 0$ that satisfies condition 2).

Notice that, the invariance of the asymptotic stability domain is an essential element of the stability concept. Three propositions given here will be employed in the next sections. Although the results presented seem to be classical in nature (see our earlier work on the subject [1]), proofs are included here in order to render the paper self-contained.

Proposition 2.3 *The stability domain D_s of the state $x = 0$ of the system (1) is positively invariant with respect to system motions, that is, $x_0 \in D_s$ implies $x(k; x_0) \in D_s$ for every $k \in \mathbb{N}$.*

Proof: Let $\varepsilon > 0$ and $x_0 \in D_s(\varepsilon)$. If the proposition were not true, there would exist an instant $k_b \in \mathbb{N}$ for which $x(k_b) = x(k_b; x_0) \notin D_s(\varepsilon)$. Hence, $\exists k_c \geq k_b$ such that $\|x(k_c; x(k_b))\| \geq \varepsilon$. But $x(k_c; x(k_b)) = x(k_c; x_0)$. Therefore, $\|x(k_c; x_0)\| \geq \varepsilon$, which implies $x_0 \notin D_s(\varepsilon)$ that contradicts $x_0 \in D_s(\varepsilon)$ and proves the proposition. ■

Proposition 2.4 *The domain of attraction D_a of the state $x = 0$ of the system (1) is positively invariant with respect to system motions.*

Proof: Let $x_0 \in D_a$, then $\lim_{k \rightarrow +\infty} \|x(k, x_0)\| = 0$. Since $x(k+j; x_0) = f^{k+j}(x_0) = f^k(f^j(x_0)) = f^k(x(j)) = x(k; x(j; x_0))$ for all $j < k$, it follows $\lim_{k \rightarrow +\infty} \|x(k, x(j; x_0))\| = 0$. Hence, $x(j; x_0) \in D_a$ for all $j < +\infty$, which proves the proposition. \blacksquare

Proposition 2.5 *The domain of asymptotic stability D_{as} of the state $x = 0$ of the system (1) is positively invariant with respect to system motions.*

Proposition 2.5 is a direct consequence of Definition 2.2, Proposition 2.3 and Proposition 2.4.

Definition 2.6 A set E is an estimate of the asymptotic stability domain D_{as} if, and only if:

- (i) E is a neighbourhood of $x = 0$,
- (ii) $E \subseteq D_{as}$ and,
- (iii) E is positively invariant set of the system.

3 Definition of the G-functions

We introduce here the class of **G**-functions (not to be confused with the Meijer or Siegel **G**-functions [11]) that will serve to stability analysis without requiring their positive definiteness or semi-definiteness. Let the extended real valued function $g : \mathbb{R}^n \rightarrow \widehat{\mathbb{R}} = \mathbb{R} \cup \{-\infty\}$ be given. Needless to say that, such a function may be also real valued, but the extended real valued function used here simply allows the limit operation to take $-\infty$ into account. Now, denote by $\mathcal{R}(g)$ the range of the function g and for $\zeta \in \mathcal{R}(g)$, denote $G_\zeta = \{x \in \mathbb{R}^n : g(x) < \zeta\}$ the ζ -sub-level set of g .

Definition 3.1 We say that, the extended real valued function $g : \mathbb{R}^n \rightarrow \widehat{\mathbb{R}}$ is a **G**-function if, and only if, it obeys:

- 1) Either, $\min_{x \in \mathbb{R}^n} g(x) = g(0)$ if $g(0)$ is defined or $\lim_{x \rightarrow 0} g(x) = -\infty$,
- 2) There is no point $x^* \in \mathbb{R}^n$ such that $\lim_{x \rightarrow x^*} g(x) = +\infty$,
- 3) $g(x) \leq \zeta$ if, and only if, $x \in ClG_\zeta$ for all $\zeta > \zeta_m \in \mathcal{R}(g)$, where $\zeta_m = g(0)$ if $g(0)$ is defined or $\zeta_m = \lim_{x \rightarrow 0} g(x) = -\infty$,
- 4) $g(x) = \zeta$ if, and only if, $x \in \partial G_\zeta$ for all $\zeta > \zeta_m \in \mathcal{R}(g)$.

Remark 2: It is important to note that, no continuity, positive definiteness or semi-definiteness conditions are required for the \mathbf{G} -functions. This explains that the stability conditions expressed in terms of a \mathbf{G} -function are substantially different than the existing Lyapunov stability criteria for discrete-time nonlinear systems.

Now, in order to show the meaning of the \mathbf{G} -functions, let us present some examples. Let the following extended real valued functions $g_1, g_2 : \mathbb{R}^n \rightarrow \widehat{\mathbb{R}}$ be in the form

$$g_1(x) = \begin{cases} \frac{\ln(\|x\|)}{\|x\|} & \text{if } x \neq 0 \in \mathbb{R}^n, \\ -\infty & \text{if } x = 0. \end{cases}, \quad g_2(x) = \begin{cases} \ln(\|x\|) & \text{if } x \neq 0 \in \mathbb{R}^n, \\ -\infty & \text{if } x = 0. \end{cases}$$

They are discontinuous at $x = 0$ and neither positive definite nor semi-definite. These functions are the \mathbf{G} -functions because in both cases $\zeta_m = -\infty$. Obviously, conditions 2), 3) and 4) are verified.

Another examples where the \mathbf{G} -function is continuous everywhere are

$$g_1(x) = \begin{cases} -\frac{\sin(\|x\|)}{\|x\|} & \text{if } x \neq 0 \in \mathbb{R}^n, \\ -1 & \text{if } x = 0. \end{cases}, \quad g_2(x) = -\cos(\sum_i x_i) \exp(-\|x\|^2).$$

These functions are not positive definite or semi-definite. Both have a global minimum at 0, so that $\zeta_m = g(0) = -1$. Clearly, the other conditions are satisfied.

Finally, let us present an example where the \mathbf{G} -function is discontinuous at some x . Indeed, let the following extended real valued function $g : \mathbb{R}^n \rightarrow \widehat{\mathbb{R}}$

$$g(x) = \begin{cases} \ln(\|x\|) & \text{if } \|x\| \neq 0 < 1 \\ -\infty & \text{if } x = 0, \\ \|x\| & \text{if } \|x\| \geq 1. \end{cases}$$

Clearly, this function is not continuous at the set $A = \{x \in \mathbb{R}^n : \|x\| = 1\} \cup \{0\}$. Of course, it is a \mathbf{G} -function because all conditions of Definition 3.1 are fulfilled.

In summary, we note that the class of \mathbf{G} -functions requires only either a global minimum at zero (not necessary equal to 0) or $\lim_{x \rightarrow 0} g(x) = -\infty$. Conditions 3) and 4) imply simply that $G_{\zeta_1} \subset G_{\zeta_2}$ for all $\zeta_1 < \zeta_2$ in $\mathcal{R}(g)$. On the other hand, condition 2) ensures, among other things, the connectedness of the set G_ζ because the connectedness is an essential element for the proof of the stability. It is not restrictive because there is a large range of functions that satisfy this condition.

4 Main results

We begin this section by stating some technical lemmas necessary for the proofs of the main results of this paper.

Lemma 4.1 *Let g be a \mathbf{G} -function. In order that the ζ -sub-level set G_ζ to be positively invariant with respect to motions of the system (1) it is sufficient that the function g obeys*

$$\Delta g(x) = g(f(x)) - g(x) \leq 0, \quad \forall x \neq 0 \in G_\zeta. \quad (2)$$

Proof: Let $x_0 \in G_\zeta$ be arbitrary. Suppose that $\exists k_i \in \mathbb{N}$ such that $x_i = x(k_i; x_0) \in G_\zeta$ and $x_{i+1} = x(k_i + 1; x(k_i)) = f(x_i) \notin G_\zeta$. We then have $g(x_i) < \zeta$ and $g(f(x_i)) \geq \zeta$. Hence, $\Delta g(x_i) = g(f(x_i)) - g(x_i) > 0$. This result contradicts condition (2) of the Lemma and completes the proof. ■

Lemma 4.2 *Let g be a \mathbf{G} -function and let $G_\zeta^c = \mathbb{R}^n \setminus G_\zeta$ be the complement in \mathbb{R}^n of G_ζ . If the function g obeys*

$$\Delta g(x) \geq 0, \quad \forall x \in G_\zeta^c, \quad (3)$$

then G_ζ^c is positively invariant set with respect to motions of system (1).

Proof: The proof follows the analogous arguments to those of the proof of Lemma 4.1. ■

We are now ready to state our stability criterion. To this end, recall that ζ_m introduced in Definition 3.1 is defined as $\zeta_m = g(0)$ if $g(0)$ is defined or $\zeta_m = \lim_{x \rightarrow 0} g(x) = -\infty$.

Theorem 4.3 *The existence of \mathbf{G} -function g and a number $\Lambda \in (\zeta_m, +\infty)$ such that $G_\Lambda \subseteq S$ is an open connected neighbourhood of $x = 0$ and*

$$\Delta g(x) = g(f(x)) - g(x) < 0, \quad \forall x \neq 0 \in G_\Lambda, \quad (4)$$

are necessary and sufficient conditions for the state $x = 0$ of the system (1) to be asymptotically stable and G_Λ to be an estimate of its domain of the asymptotic stability D_{as} , i.e., $G_\Lambda \subseteq D_{as}$.

Proof: Necessity: Let $x = 0$ of system (1) be asymptotically stable. From converse theorem of Lyapunov (see [9]), we know that there exists a positive definite function V defined on a neighbourhood \mathbb{U} of $x = 0$ such that along the trajectories of the system we have $\Delta V(x) = V(f(x)) - V(x) < 0$, $\forall x \neq 0 \in D_{as}$. Besides, the α -sub-level set of V , $\mathbb{V}_\alpha = \{x \in \mathbb{U} : V(x) < \alpha\}$ is a subset of D_{as} and positively invariant for some $\alpha > 0$. From this information, we have to show the existence of a number Λ such that the \mathbf{G} -function to be proposed gives rise to an open connected set G_Λ . Indeed, note first that the set \mathbb{V}_α might not be connected in general and the connectedness of \mathbb{V}_α depends on the nature of V . Let σ be such that $0 < \sigma \leq \alpha$ and the σ -sub-level set $\mathbb{V}_\sigma = \{x \in \mathbb{U} : V(x) < \sigma\}$ is the largest open connected subset of \mathbb{U} containing $x = 0$. Because V is positive definite, such a number σ exists since V is continuous and $x = 0$ is the global minimum of V . It is clear that $\mathbb{V}_\sigma \subseteq \mathbb{V}_\alpha \subseteq D_{as}$. Furthermore, \mathbb{V}_σ is positively

invariant with respect to motions of the system. With this in mind, let now the extended real valued function $g : \mathbb{R}^n \rightarrow \widehat{\mathbb{R}}$ be in the form

$$g(x) = \begin{cases} \ln(V(x)) & \text{if } x \neq 0 \in \mathbb{V}_\sigma, \\ -\infty & \text{if } x = 0, \\ \ln(\sigma) & \text{if } x \notin \mathbb{V}_\sigma. \end{cases}$$

It is neither continuous at $x = 0$ nor positive definite or semi-definite. We shall first prove that this function is a **G**-function. Indeed, observe that $V(0) = 0$ implies $\zeta_m = \lim_{x \rightarrow 0} g(x) = -\infty$, so that the condition 1) of Definition 3.1 is verified. On the other hand, since for $x \in \mathbb{V}_\sigma$ we have $\ln(V(x)) < \ln(\sigma) < \sigma < +\infty$, then the condition 2) of Definition 3.1 is also verified. Finally, it is clear from the definition of g that conditions 3) and 4) of Definition 3.1 are trivially fulfilled. This proves the existence of the **G**-function g .

Next, observe that if we take $\Lambda = \ln(\sigma)$, then $G_\Lambda = \{x \in \mathbb{R}^n : \ln(V(x)) < \Lambda\} = \mathbb{V}_\sigma$. Hence, G_Λ is an open connected neighbourhood of $x = 0$. This proves the existence of $G_\Lambda \subseteq D_{as}$ and $G_\Lambda \subseteq S$ due to $D_{as} \subseteq S$ in view of Definition 2.2. Finally, since $\Delta V(x) < 0$ for all $x \neq 0 \in \mathbb{V}_\sigma$ along the trajectories of the system and $\mathbb{V}_\sigma = G_\Lambda$ is positively invariant, it follows

$$V(f(x)) < V(x), \quad \forall x \neq 0 \in G_\Lambda.$$

Thus, for all $x \neq 0 \in G_\Lambda$ we have

$$\Delta g(x) = \ln(V(f(x))) - \ln(V(x)) < 0,$$

which proves the necessity.

Sufficiency: The hypotheses of the theorem prove that for all $\zeta \in (\zeta_m, \Lambda]$ the set G_ζ is positively invariant with respect to the motions of the system (1) (Lemma 4.1), i.e.,

$$x_0 \in G_\zeta \Rightarrow x(k; x_0) \in G_\zeta, \quad \forall k \in \mathbb{N}, \forall \zeta \in (\zeta_m, \Lambda]. \quad (5)$$

Let $\varepsilon > 0$ be arbitrarily chosen and let $\zeta(\varepsilon) \in (\zeta_m, \Lambda]$ be such that $C\ell G_{\zeta(\varepsilon)} \subset B_\varepsilon$. The existence of the number $\zeta(\varepsilon)$ obeying the last condition is assured by the definitions of g and $G_{\zeta(\varepsilon)}$. Now, let $\delta(\varepsilon) > 0$ be such that

$$B_{\delta(\varepsilon)} \subset C\ell G_{\zeta(\varepsilon)} \subset B_\varepsilon. \quad (6)$$

Such $\delta(\varepsilon)$ exists because G_Λ is an open connected neighbourhood of $x = 0$. From (5) and (6) we have

$$x_0 \in B_{\delta(\varepsilon)} \Rightarrow x(k; x_0) \in G_{\zeta(\varepsilon)} \subset B_\varepsilon, \quad \forall k \in \mathbb{N}. \quad (7)$$

which proves the stability of $x = 0$ and $G_{\zeta(\varepsilon)} \subseteq D_s$ for all $\zeta(\varepsilon) \in (\zeta_m, \Lambda]$. Moreover, (7) and $\zeta(\varepsilon) \in (\zeta_m, \Lambda]$ show that G_Λ is positively invariant with respect to the system motions. Hence, G_Λ is also an estimate of D_s in view of Definition 2.6, i.e., $G_\Lambda \subseteq D_s$.

In order to prove attraction we suppose opposite, i.e., that there exist $x_0^* \neq 0 \in G_\Lambda$ and $\alpha > 0$ such that $\lim_{k \rightarrow +\infty} \|x(k; x_0^*)\| = \alpha$. Let $\lim_{k \rightarrow +\infty} g(x(k; x_0^*)) = \gamma$. Since $x(k; x_0^*) \neq 0$ and $\Delta g(x) < 0$ for all $x \neq 0 \in G_\Lambda$, it follows from 1) of Definition 3.1 that $\gamma > \zeta_m$ (otherwise $\gamma \leq \zeta_m$ would imply $x(+\infty; x_0^*) = 0$ in view of 1) of Definition 3.1, which is a contradiction since $\|x(+\infty; x_0^*)\| \neq 0$ by assumption). Now, observe that since g is decreasing and $\lim_{k \rightarrow +\infty} g(x(k; x_0^*)) = \gamma$, then the trajectory $x(k; x_0^*)$ lies outside G_γ . It follows for all $k \in \mathbb{N}$

$$x(k, x_0^*) \in G(x_0^*, \gamma) = \{x \in \mathbb{R}^n : \gamma \leq g(x) \leq g(x_0^*)\} \subset G_\Lambda.$$

Let $-\mu = \sup_{x \in G(x_0^*, \gamma)} \Delta g(x)$. The supremum clearly exists due to stability of $x = 0$ of system (1) and positive invariance of G_Λ . Thus, $-\mu < 0$ by (4). Hence, $\Delta g(x) \leq -\mu < 0$ for all $x \in G(x_0^*, \gamma)$. This infers that for all $x(k) \in G(x_0^*, \gamma)$

$$g(x(k; x_0^*)) \leq g(x_0^*) - \sum_{i=0}^{k-1} \mu = g(x_0^*) - k\mu.$$

Therefore, for $k \rightarrow +\infty$ we have $g(x(k; x_0^*)) \rightarrow -\infty$ due to $\mu > 0$ and $g(x_0^*) \neq +\infty$ in view of condition 2) of Definition 3.1. But $\lim_{k \rightarrow +\infty} g(x(k; x_0^*)) = \gamma \neq -\infty$, a contradiction with the definition of γ . Consequently $\lim_{k \rightarrow +\infty} \|x(k; x_0)\| = 0$ for all $x_0 \in G_\Lambda$, so that $G_\Lambda \subseteq D_a$ and hence $G_\Lambda \subseteq D_s \cap D_a = D_{as}$, which completes the proof of the theorem. \blacksquare

Remark 3: As in the Lyapunov approach, the concept of \mathbf{G} -functions used in theorem 4.3 has an intuitive geometric meaning. If a solution $x(k, x_0)$ starts at $x_0 \in G_\Lambda$, then the positive invariance of G_Λ with respect to the system ensures that the solution remains in G_Λ for all $k \in \mathbb{N}$. Moreover, the positive invariance of G_ζ for all $\zeta \leq \Lambda$ and the strict inequality $\Delta g(x) < 0$ will ensure also that the system solution will progress by entering a decreasing sequence of ζ -sub-level-sets $G_{\zeta_1} \supset G_{\zeta_2} \cdots \supset G_{\zeta_i} \supset \cdots$ with $\zeta_1 = g(x(0)) > \zeta_2 = g(x(1)) > \cdots > \zeta_i = g(x(i-1)) > \cdots > \zeta_m$. Consequently, in the long run the system solution will asymptotically converge to $C\ell G_{\zeta_m} = \{x \in \mathbb{R}^n : g(x) \leq \zeta_m\} = \{0\}$.

Remark 4: It is important to note that, if $g(0)$ is finite, one might be tempted to consider $V(x) = g(x) - g(0)$ and thus obtain a Lyapunov function. However, observe in this case that $\Delta V(x) = g(f(x)) - g(x)$. Hence, by doing so, we bring back to the fundamental problem of the analysis of the sign of $\Delta V(x)$ with respect to g (which is sign indefinite) rather than with respect to the positive definite function V . The routine proof of attraction of $x = 0$ using $\Delta V(x) < 0$ in the existing sufficient conditions given by the Lyapunov method is based on the fact that V is positive for $x \neq 0$. Therefore, taking $V(x) = g(x) - g(0)$ does not recapture the technique of Lyapunov function.

Next, to illustrate the notion of the \mathbf{G} -functions, let us look at the following simple but illustrative examples. First of all, it is important to note that, the

asymptotic stability of the presented examples can be tested by several tools, but the aim here is to show that the \mathbf{G} -functions are also applicable.

Example 1: Consider the following system defined on $S = (-1; 1)$

$$x(k+1) = \sin(x(k))(\tanh(x(k)) - x(k)) = f(x(k)).$$

The system has $x = 0$ as a unique equilibrium state on S . As shown in Section 3, let the following extended function

$$g(x) = \begin{cases} \frac{\ln(|x|)}{|x|} & \text{if } x \neq 0 \in \mathbb{R}, \\ -\infty & \text{if } x = 0. \end{cases}$$

be candidate \mathbf{G} -function for such system. Since, the function f maps S into itself, we take $\Lambda = 0$ so that $G_\Lambda = (-1; 1) = S$ is an open connected neighbourhood of $x = 0$. Now, we test $\Delta g(x)$ on $G_\Lambda/\{0\}$. Indeed, for all $x \neq 0 \in G_\Lambda$ we have

$$\Delta g(x) = \frac{\ln(|f(x)|)}{|f(x)|} - \frac{\ln(|x|)}{|x|} = \frac{1}{|f(x)||x|} \left(|x| \ln(|f(x)|) - |f(x)| \ln(|x|) \right).$$

Let $p(x) = |x| \ln(|f(x)|) - |f(x)| \ln(|x|)$. Since $p(-x) = p(x)$, it is enough to study $p(x)$ on $(0, 1)$. Indeed, for all $x \in (0, 1)$, we have

$$p(x) = x \ln(\sin(x)) + x \ln(|\tanh(x) - x|) - \sin(x) |\tanh(x) - x| \ln(x).$$

But, $0 < \sin(x) < 1$ and $\ln(x) < 0$ on $(0, 1)$, it follows

$$p(x) < x \ln(|\tanh(x) - x|) - |\tanh(x) - x| \ln(x).$$

Since $x \neq |\tanh(x) - x|$ on $(0, 1)$, then by applying the fact that $(a-b)(\ln(b) - \ln(a)) < 0$ for all $a \neq b \in \mathbb{R}_{>0}$, we get after some elementary transformations

$$p(x) < [x - |\tanh(x) - x|] [\ln(|\tanh(x) - x|) - \ln(x)] + \ln(x) [x - 2|\tanh(x) - x| + |\tanh(x) - x| \ln(|\tanh(x) - x|)].$$

Now, remark that for $x \in (0, 1)$ we have $\ln(|\tanh(x) - x|) < 0$ and $x - 2|\tanh(x) - x| > 0$. We conclude that $p(x) < 0$ on $(-1, 1)$ because $p(x)$ is even. Therefore, $\Delta g(x) < 0$ for all $x \neq 0 \in G_\Lambda$. All conditions of Theorem 4.3 are satisfied, the state $x = 0$ of the system is asymptotically stable and the set G_Λ is an estimate of its domain of asymptotic stability D_{as} .

Example 2: Consider the following system defined on $S = (-\pi; \pi)$

$$x(k+1) = \frac{\sin(x(k))}{2 \cos^3(\frac{x(k)}{3})} = f(x(k)). \quad (8)$$

The system has $x = 0$ as a unique equilibrium state on S . As shown in Section 3, let the following extended function

$$g(x) = \begin{cases} -\frac{\sin(x)}{x} & \text{if } x \neq 0, \\ -1 & \text{if } x = 0. \end{cases}$$

be candidate \mathbf{G} -function for such system with $\zeta_m = -1$. It is well known that the function g enjoys the following properties (see [14], [16]).

- Jordan's inequality

$$-1 \leq -\frac{\sin(x)}{x} \leq -\frac{2}{\pi}, \quad \text{for } |x| \leq \frac{\pi}{2}, \quad (9)$$

- Sándor's inequality

$$-\cos\left(\frac{x}{2}\right) < -\frac{\sin(x)}{x} < -\frac{1 + \cos(x)}{2}, \quad \text{for } |x| < \frac{\pi}{2}. \quad (10)$$

From the Jordan's inequality (9), if we take $\Lambda = -\frac{2}{\pi}$, then $G_\Lambda = (-\frac{\pi}{2}; \frac{\pi}{2}) \subset S$ is an open connected neighbourhood of $x = 0$. In order to test $\Delta g(x)$ on $G_\Lambda/\{0\}$, observe that for all $x \neq 0 \in G_\Lambda$ we have $\Delta g(x) = \frac{\sin(x)}{x} - \frac{\sin(f(x))}{f(x)}$. To exploit inequality (10) we have to show that $f(x) \in (-\frac{\pi}{2}; \frac{\pi}{2})$. To this end, since $f(x)$ is odd, it suffices to study f on $(0, \frac{\pi}{2})$. Indeed, elementary computation yields $f'(x) = \frac{1}{2} \cos(\frac{2x}{3}) \cos^{-4}(\frac{x}{3}) > 0$ on $(0, \frac{\pi}{2})$. Therefore, $0 = f(0) < f(x) < f(\frac{\pi}{2}) = \frac{4}{3\sqrt{3}} < \frac{\pi}{2}$ so that $f(x) \in (-\frac{\pi}{2}; \frac{\pi}{2})$. Consequently, we can use inequalities (10) to obtain

$$\Delta g(x) < \cos\left(\frac{x}{2}\right) - \frac{1 + \cos(f(x))}{2}.$$

But, $\frac{1 + \cos(f(x))}{2} = \cos^2\left(\frac{f(x)}{2}\right)$ and $\cos^2\left(\frac{x}{4}\right) = \frac{1 + \cos(\frac{x}{2})}{2} \geq \cos\left(\frac{x}{2}\right)$. This implies

$$\Delta g(x) < \cos^2\left(\frac{x}{4}\right) - \cos^2\left(\frac{f(x)}{2}\right) = \left[\cos\left(\frac{x}{4}\right) - \cos\left(\frac{f(x)}{2}\right)\right] \left[\cos\left(\frac{x}{4}\right) + \cos\left(\frac{f(x)}{2}\right)\right].$$

Let now $h(x) = \cos\left(\frac{x}{4}\right) - \cos\left(\frac{f(x)}{2}\right)$. Since, $\cos\left(\frac{x}{4}\right) + \cos\left(\frac{f(x)}{2}\right) > 0$ on $(-\frac{\pi}{2}; \frac{\pi}{2})$ and $h(-x) = h(x)$, it is sufficient to study $h(x)$ on $(0, \frac{\pi}{2})$. For, define $p(x) = \frac{x}{4} - \frac{f(x)}{2}$. After some elementary transformations we get $p'(x) = \frac{1}{4} - \frac{1}{2}f'(x) = \frac{1}{4} \tan^4(x) > 0$. Therefore, $p(x) > p(0) = 0$ so that $\frac{f(x)}{2} < \frac{x}{4}$ on $(0, \frac{\pi}{2})$. Since $\cos(x)$ is decreasing on $(0, \frac{\pi}{2})$ we get $\cos\left(\frac{x}{4}\right) < \cos\left(\frac{f(x)}{2}\right)$ and thus $h(x) < 0$ for all $x \in (-\frac{\pi}{2}; \frac{\pi}{2})$. Consequently, $\Delta g(x) < 0$ for all $x \neq 0 \in G_\Lambda$. The state $x = 0$ of the system is asymptotically stable and the set G_Λ is an estimate of its domain of asymptotic stability D_{as} .

Next, recall that a function $\phi(\cdot) : \mathbb{R}_{\geq 0} \rightarrow \mathbb{R}_{\geq 0}$ is a comparison function of the class \mathcal{K} defined by Hahn [6] if, and only if, it is continuous, strictly increasing and $\phi(0) = 0$.

Classically, in the existing Lyapunov stability theory, it is customary to require also the existence of a function ϕ of class \mathcal{K} such that $\Delta V(x) \leq -\phi(\|x\|)$ instead of requiring only $\Delta V(x) < 0$, where V is a Lyapunov function. It is worthwhile to show how this requirement can be also extended to \mathbf{G} -functions. In this framework, we have the next result.

Theorem 4.4 *The existence of a \mathbf{G} -function g , a function ϕ of class \mathcal{K} and a number $\Lambda \in (\zeta_m, +\infty)$ such that $G_\Lambda \subseteq S$ is an open connected neighbourhood of $x = 0$ and*

$$\Delta g(x) \leq -\phi(\|x\|), \quad \forall x \neq 0 \in G_\Lambda, \quad (11)$$

are necessary and sufficient conditions for the state $x = 0$ of the system (1) to be asymptotically stable and G_Λ to be an estimate of its domain of asymptotic stability D_{as} , i.e., $G_\Lambda \subseteq D_{as}$.

Proof: By Theorem 4.3, it is clear that we need only to show here the existence of a function ϕ of class \mathcal{K} such that the inequality (11) is verified. Indeed, from converse theorem of Lyapunov we know also that there exist a positive definite function V and a function ψ of class \mathcal{K} such that along the trajectories of the system (1) we have for all $x \neq 0 \in D_{as}$ (see [9], Theorem 4.10.2)

$$\Delta V(x) = V(f(x)) - V(x) \leq -\psi(\|x\|).$$

This in turn implies that $V(x) > \psi(\|x\|)$, $\forall x \neq 0 \in D_{as}$ in view of positive definiteness of V . With this in mind, let g and $G_\Lambda = \mathbb{V}_\sigma$ as defined in the proof of the necessity part of Theorem 4.3. Then we have

$$V(f(x)) \leq V(x) - \psi(\|x\|), \quad \forall x \neq 0 \in G_\Lambda,$$

which implies

$$\frac{V(f(x))}{V(x)} \leq 1 - \frac{\psi(\|x\|)}{V(x)}, \quad \forall x \neq 0 \in G_\Lambda.$$

Now, observe that since $V(x) > \psi(\|x\|)$ for all $x \neq 0 \in G_\Lambda$, then $\ln\left(1 - \frac{\psi(\|x\|)}{V(x)}\right)$ is well defined. Thus, we get

$$\Delta g(x) = \ln\left(\frac{V(f(x))}{V(x)}\right) \leq \ln\left(1 - \frac{\psi(\|x\|)}{V(x)}\right) \leq -\frac{\psi(\|x\|)}{V(x)},$$

where we have used that fact that $\ln(1 - x) \leq -x$ for $0 \leq x < 1$. Let $\gamma = \sup_{x \in G_\Lambda} V(x) > 0$. It follows

$$\Delta g(x) \leq -\frac{1}{\gamma}\psi(\|x\|), \quad \forall x \neq 0 \in G_\Lambda.$$

Clearly $\phi(\|x\|) = \frac{1}{\gamma}\psi(\|x\|)$ belongs to class \mathcal{K} . This completes the proof. \blacksquare

Now, for the global asymptotic stability, we have the following.

Corollary 4.5 *Under the conditions of Theorem 4.3 or Theorem 4.4, if $S = \mathbb{R}^n$ and Λ can be arbitrarily large, then the state $x = 0$ is globally asymptotically stable, i.e., $D_{as} = \mathbb{R}^n$.*

Example 3: Consider the following two dimensional system defined on \mathbb{R}^2

$$x(k+1) = x(k)y(k) \exp(-y^2(k)) \quad (12)$$

$$y(k+1) = \alpha x(k) \quad (13)$$

where $\alpha \neq 0 \in \mathbb{R}$. We are interested in discovering a condition on α under which the global asymptotic stability of the fixed point $(0, 0)$ is guaranteed. Let the extended real valued function $g : \mathbb{R}^n \rightarrow \widehat{\mathbb{R}}$ be in the form

$$g(x) = \begin{cases} \ln(|x| + |y|) & \text{if } (x, y) \neq (0, 0) \in \mathbb{R}^n, \\ -\infty & \text{if } (x, y) = (0, 0). \end{cases}$$

This function is the **G**-function which evidently obeys Definition 3.1 with $\zeta_m = -\infty$. Now, Let $\Lambda \in \mathbb{R}$ be arbitrarily chosen. Clearly, $G_\Lambda = \{x \in \mathbb{R}^n : \ln(|x| + |y|) < \Lambda\}$, is on an open connected neighbourhood of $(x, y) = (0, 0)$. We test now $\Delta g(x)$ on $G_\Lambda / \{(0, 0)\}$. Indeed,

$$\Delta g(x) = \ln\left(\frac{|x|}{|x| + |y|}\right) + \ln\left(|y| \exp(-y^2) + |\alpha|\right).$$

Since, $\frac{|x|}{|x| + |y|} \leq 1$ and $|y| \exp(-y^2) \leq \frac{1}{\sqrt{2e}}$, $\forall y \in \mathbb{R}$, it follows $\Delta g(x) \leq \ln\left(\frac{1}{\sqrt{2e}} + |\alpha|\right)$. If $|\alpha| < 1 - \frac{1}{\sqrt{2e}}$, then $\Delta g(x) < 0$ for all $(x, y) \in G_\Lambda / \{(0, 0)\}$. We conclude that all conditions of Theorem 4.3 are satisfied for arbitrary $\Lambda \in \mathbb{R}$. The state $(x, y) = (0, 0)$ is then globally asymptotically stable.

So far, we have investigated the asymptotic stability problem of discrete-time nonlinear systems by means of sign indefinite **G**-functions. Now, a fundamental question arises: How can the domain of asymptotic stability be completely determined by the **G**-functions? By enhancing the sufficient conditions for asymptotic stability of the state $x = 0$, the following theorem establishes a complete answer to this question.

Theorem 4.6 *If all conditions of Theorem 4.3 or Theorem 4.4 hold and if in addition*

$$\Delta g(x) \geq 0, \quad \forall x \in \mathbb{R}^n / G_\Lambda, \quad (14)$$

then the set G_Λ is the domain of the asymptotic stability of the state $x = 0$ of the system (1), i.e., $G_\Lambda \equiv D_{as}$.

Proof: By Theorem 4.3 we have

$$G_\Lambda \subseteq D_a \cap D_s = D_{as}. \quad (15)$$

Let $x_0 \in \mathbb{R}^n / G_\Lambda$. Lemma 4.2 shows that \mathbb{R}^n / G_Λ is positively invariant, i.e., $x_0 \in \mathbb{R}^n / G_\Lambda$ implies $x(k; x_0) \in \mathbb{R}^n / G_\Lambda$ for all $k \in \mathbb{N}$. Since \mathbb{R}^n / G_Λ does not contain the origin, it follows $\lim_{k \rightarrow +\infty} \|x(k; x_0)\| \neq 0$, which implies $D_a \subseteq G_\Lambda$. This and (15) yield $D_a \cap D_s = G_\Lambda$ so that $D_{as} = G_\Lambda$, which completes the proof. ■

Example 4: Let the following family of systems

$$x(k+1) = \frac{f(x(k))}{1 + \|\|x(k)\| - e^\alpha} x(k), \quad (16)$$

where $x(k) \in \mathbb{R}^n$, $\alpha \in \mathbb{R}$ and $f : \mathbb{R}^n \rightarrow \mathbb{R}$. We are interested in discovering functions family $f(x)$ under which the set $S(\alpha) = \{x \in \mathbb{R}^n : \|x\| < e^\alpha\}$ is the domain of the asymptotic stability of the fixed point $x = 0$. As shown in Section 3, let the following **G**-function

$$g(x) = \begin{cases} \ln(\|x\|) & \text{if } x \neq 0 \in \mathbb{R}^n, \\ -\infty & \text{if } x = 0. \end{cases}$$

be candidate for such system. Let $\Lambda = \alpha$, then $G_\Lambda = \{x \in \mathbb{R}^n : \ln(\|x\|) < \Lambda\} = S(\alpha)$ is an open connected neighbourhood of $x = 0$. Now, on $G_\Lambda/\{0\}$ we have

$$\Delta g(x) = \ln(|f(x)|) + \ln\left(\frac{1}{1 + \|\|x\| - e^\alpha}\right)$$


According to Theorem 4.6, the set $S(\alpha)$ is the domain of asymptotic stability if:

- on $S(\alpha)/\{0\}$, $|f(x)| \leq 1$,
- on $x \in \partial S(\alpha)$, $f(x) = 1$,
- on $\mathbb{R}^n/S(\alpha)$, $f(x) \geq 1 + \|x\| - e^\alpha$.


For the simulation purpose, consider the case of one dimensional systems. For $\alpha = 0$, it is clear that $f(x) = x^2$ or $f(x) = e^{x^2} + 1 - e$ satisfy all the above conditions. In Both cases, system motions are shown in Figure 1 for $x(0) = 0.999999999 \in S(0)$ and in Figure 2 for $x(0) = 1.00000001$ out of $S(0)$. Both initial states are close to the boundary of $S(0)$. In the former case the system motion converges to $x = 0$, while in the latter case it does not. Notice also a very small difference between initial values in the two cases. The set $S(0)$ is therefore the domain of the asymptotic stability of $x = 0$.

5 Conclusion

The theoretical study of the stability problems for the discrete-time nonlinear systems has been developed by using properties of a class of functions called the **G**-functions in this paper. It has resulted in the new necessary and sufficient conditions for asymptotic stability of such systems and an estimation or the exact determination of the asymptotic stability domain of the zero state $x = 0$. Hence, the paper establishes essentially an alternative approach for the asymptotic stability analysis of time-invariant nonlinear discrete-time systems. The essence is the possibility to use henceforth the sign indefinite and discontinuous functions for testing asymptotic stability of such systems.




(a) The graph for $f(x) = x^2$




(b) The graph for $f(x) = e^{x^2} - e + 1$

Figure 1: Stable trajectories



(a) The graph for $f(x) = x^2$



(b) The graph for $f(x) = e^{x^2} - e + 1$

Figure 2: Unstable trajectories

Finally, we would like to point out that the **G**-functions approach presented here simply offers an additional tool for the study of the stability of discrete-time nonlinear systems. It can be further developed and possibly combined with the Lyapunov function approach to generate more interesting and advanced results in the future, specifically for stabilization and control of discrete-time nonlinear systems.

References

- [1] R. Bouyekh and L.T. Grujic, *Lyapunov-like solutions to stability problems for discrete-time systems on asymptotically contractive sets*, Nonlinear Dynamics 18 (1999), pp. 107–127.
- [2] P. Giesl, *Construction of a local and global Lyapunov function using radial basis functions*, IMA J. Appl. Math 75 (2008), pp. 782–802.
- [3] P. Giesl and S. Hafsteinb, *Computation of Lyapunov functions for nonlinear discrete time systems by linear programming*, Journal of Difference Equations and Applications 20 (2014), pp. 610–640.

- [4] P. Giesl and S. Hafsteinb, *Review on computational methods for Lyapunov functions*, Discrete and Continuous Dynamical Systems Series B 20 (2015), pp. 2291–2331.
- [5] L.T. Gruyitch, J.P. Richard, P. Borne, and J.C. Gentina, *Stability Domains*, Chapman and Hall/ CRC press, London, 2003.
- [6] W. Hahn, *Stability of Motion*, Springer-Verlag, Berlin, 1967.
- [7] A. Iggidr and M. Bensoubaya, *New results on the stability of discrete-time systems and applications to control problems*, Journal of Mathematical Analysis and Applications 219 (1998), pp. 392–414.
- [8] R.E. Kalman and J.E. Bertram, *Control system analysis and design via the second method of Lyapunov, part II, discrete systems*, Transactions of the ASME Journal of Basic Engineering 82 (1960), pp. 394–400.
- [9] V. Lakshmikantham and D. Trigiante, *Theory of Difference Equations with Applications to Numerical Analysis*, Academic Press, San Diego, 1988.
- [10] J.P. LaSalle, *The Stability of Dynamical Systems*, SIAM, Philadelphia, Pennsylvania, 1976.
- [11] Y.L. Luke, *The Special Functions and their Approximations*, Academic Press, New York, 1969.
- [12] R. Miller and A. Michel, *Ordinary Differential Equation*, Academic Press, New York, 1982.
- [13] R. O'schea, *The extension of the Zobov's method to sampled data control systems described by nonlinear autonomous difference equations*, IEEE Transactions on Automatic Control January (1964), pp. 62–70.
- [14] B.N.G. Qi. Feng Qi D-W. Niu, *Refinements, generalizations, and applications of Jordan's inequality and related problems*, Journal of Inequalities and Applications 2009, Article ID 271923 (2009), p. 52 pages.
- [15] L. Rosier, *Homogeneous Lyapunov function for homogeneous continuous vector fields*, Systems and Control Letters 19 (1992), pp. 467–473.
- [16] J. Sándor, *Certain trigonometric inequalities*, Octagon Mathematical Magazine 9 (2001), pp. 331–336.