Existence of a heteroclinic solution for a double well potential equation in an infinite cylinder of \mathbb{R}^N

Claudianor O. Alves*

Universidade Federal de Campina Grande, Unidade Acadêmica de Matemática CEP: 58429-900 - Campina Grande-PB, Brazil e-mail: coalves@mat.ufcg.edu

Abstract

This paper concerns with the existence of a heteroclinic solution for the following class of elliptic equations

 $-\Delta u + A(\epsilon x, y)V'(u) = 0, \quad \text{in} \quad \Omega,$

where $\epsilon > 0$, $\Omega = \mathbb{R} \times \mathcal{D}$ is an infinite cylinder of \mathbb{R}^N with $N \ge 2$. Here, we have considered a large class of potential V that includes the Ginzburg-Landau potential $V(t) = (t^2 - 1)^2$ and two geometric conditions on the function A. In the first condition we assume that A is asymptotic at infinity to a periodic function, while in the second one A satisfies

$$0 < A_0 = A(0, y) = \inf_{(x, y) \in \Omega} A(x, y) < \liminf_{|(x, y)| \to +\infty} A(x, y) = A_\infty < \infty, \quad \forall y \in \mathcal{D}.$$

2000 Mathematics Subject Classifications: 26A33, 34C37, 35A15, 35B38. **Key words.** Heteroclinic solutions, Variational methods, Double potential, Critical points

1 Introduction

This paper concerns with the existence of a heteroclinic solution for the following class of elliptic equations

$$-\Delta u + A(\epsilon x, y)V'(u) = 0, \quad \text{in} \quad \Omega, \tag{PDE}$$

together with the Neumann boundary condition

$$\frac{\partial u}{\partial \nu}(x,y) = 0, \ x \in \mathbb{R}, \ y \in \partial \mathcal{D},$$
 (NC)

^{*}Research of C. O. Alves partially supported by CNPq 304036/2013-7 and INCTMAT/CNPq/Brazil

where $N \geq 2$, $\epsilon > 0$, Ω is an infinite cylinder of the type $\Omega = \mathbb{R} \times \mathcal{D}$ with $\mathcal{D} \subset \mathbb{R}^{N-1}$ being a smooth bounded domain and $\nu = \nu(y)$ is the normal vector outward pointing to $\partial \mathcal{D}$. Related to the functions $A : \overline{\Omega} \to \mathbb{R}$ and $V : \mathbb{R} \to \mathbb{R}$, we are assuming the following conditions:

Conditions on V:

 $(V_1) V \in C^1(\mathbb{R}, \mathbb{R}).$

$$(V_2)$$
 $V(-1) = V(1) = 0$ and $V(t) \ge 0$ for all $t \in \mathbb{R}$,

and

 (V_3) V(t) > 0 for all $t \neq -1, 1$.

An example of V satisfying $(V_1) - (V_3)$ is the Ginzburg-Landau potential $V(t) = (t^2 - 1)^2$.

Conditions on A:

In whole this paper A is a C^1 -function that belongs to one of the following classes:

Class 1: A is asymptotic at infinity to a periodic function.

In this class, we assume that there exists a C^1 -function $A_p: \overline{\Omega} \to \mathbb{R}$, which is 1- periodic in x, such that

(A₁)
$$|A(x,y) - A_p(x,y)| \to 0 \text{ as } |(x,y)| \to +\infty$$

and

$$(A_2) \qquad \qquad 0 < A_0 = \inf_{(x,y) \in \Omega} A(x,y) \le A(x,y) < A_p(x,y), \ \forall (x,y) \in \Omega.$$

This type of condition is well known when we are working with periodic asymptotically problem of the type

$$-\Delta u + A(x)u = f(u), \text{ in } \mathbb{R}^N,$$

see for example Alves, Carrião and Miyagaki [7], Jianfu and Xiping [9] and their references.

Class 2: A satisfies the Rabinowitz's condition.

In this class of functions, we suppose that

$$(A_3) \qquad 0 < A_0 = A(0, y) = \inf_{(x,y) \in \Omega} A(x, y) < \liminf_{|(x,y)| \to +\infty} A(x, y) = A_\infty < \infty, \quad \forall y \in \mathcal{D}.$$

A condition like above has been introduced by Rabinowitz [11, Theorem 4.33] to study the existence of solution for a P.D.E. of the type

$$-\epsilon^2 \Delta u + A(x)u = f(u)$$
 in \mathbb{R}^N ,

where $\epsilon > 0, f : \mathbb{R} \to \mathbb{R}$ is a continuous function with subcritical growth and $A : \mathbb{R}^N \to \mathbb{R}$ is a continuous function satisfying

$$0 < \inf_{x \in \mathbb{R}^N} A(x) < \liminf_{|x| \to \infty} A(x).$$

By using variational methods, more precisely the mountain pass theorem, Rabinowitz has established the existence of solution for ϵ small enough. For this reason, throughout this article, we will call (A_3) of Rabinowitz's condition.

By $(V_1) - (V_3)$, V is a double well potential and we are interested in the existence of solutions for (PDE) and (NC) that are heteroclinic in x from 1 to -1. A heteroclinic solution from 1 to -1 is a function $u \in C^2(\overline{\Omega}, \mathbb{R})$ verifying (PDE)-(NC) with

$$u(x,y) \to 1$$
 as $x \to -\infty$ and $u(x,y) \to -1$ as $x \to +\infty$, uniformly in $y \in \mathcal{D}$.

In [12], Rabinowitz has proved the existence of a heteroclinic solution for elliptic equations of the type

$$-\Delta u = g(x, y, u), \quad \text{in} \quad \Omega,$$

together with the boundary condition (NC) and also with the Dirichlet boundary condition, that is,

$$u(x,y) = 0, \quad x \in \mathbb{R}, \ y \in \partial \mathcal{D}.$$
 (DC)

In order to prove the existence of heteroclinic solution, in Section 2, Rabinowitz has used variational methods by supposing on g the conditions below:

 $(q_1) \quad g \in C^1(\overline{\Omega} \times \mathbb{R}, \mathbb{R}).$

 (g_2) g(x, y, t) is even and 1-periodic in x.

In Section 3, Rabinowitz has considered some conditions on g that permit to study other classes of nonlinearity. From these comments, we see that if

$$g(x, y, t) = A(x, y)V'(t), \qquad (g)$$

Rabinowitz has studied the case when A(x, y) is 1-periodic in x, see Section 2 of the paper above mentioned. Here, we continue this study, because we will work with two new classes of function A that were not considered in that paper, more precisely the Classes 1 and 2.

After Byeon, Montecchiari and Rabinowitz [8] have established the existence of heteroclinic solution $u: \Omega \to \mathbb{R}^m$ for a large class of elliptic system like

$$-\Delta u + V_u(x, u) = 0, \quad \text{in} \quad \Omega,$$

together with the boundary condition (NC) by supposing the following conditions on potential V:

 (H_1) $V \in C^1(\overline{\Omega} \times \mathbb{R}^m, \mathbb{R})$ and $V(x_1 + 1, x_2, ..., x_N, y) = V(x, y)$, i.e., V is 1-periodic in x_1 . (H_2) There are points $a^- \neq a^+$ such that $V(x, a^{\pm}) = 0$ for all $x \in \Omega$ and V(x, y) > 0 otherwise.

(H₃) There is a constant $\underline{V} > 0$ such that $\liminf V(x, t) \ge \underline{V}$ uniformly in $x \in \Omega$.

 (H_4) For $N \ge 2$, there exist constants $c_1; C_1 > 0$ such that

$$|V_u(x,t)| \le c_1 + C_1 |t|^p,$$

where $1 for <math>N \ge 3$ and there is no upper growth restriction on p if N = 2. In the present paper, we are working with the potential V(x, y, u) = A(x, y)V(u), with A belonging to Classes 1 or 2 and V satisfying $(H_1) - (H_4)$. Our paper also continues the study made in [8] for m = 1, because we are working with other classes of function A. Here, it is very important to mention that the study of elliptic system as above is very subtle because some arguments used for the scalar case m = 1 cannot be used for general case m > 1 as for example maximum principle.

In the literature we also find interesting papers that study the existence of heteroclinic solution for elliptic equations in whole \mathbb{R}^N like

$$-\Delta u(x,y) + A(x,y)V'(u(x,y)) = 0, \quad (x,y) \in \mathbb{R}^N,$$

by supposing different conditions on A and V, see for example, Alessio and Montecchiari [2], Alessio, Jeanjean and Montecchiari [5], Alessio, Gui and Montecchiari [6], Rabinowitz [13], Rabinowitz and Stredulinsky [14, 15, 16] and their references. The reader can find versions for elliptic systems of the above equation in Alama, Bronsard and Gui [1], Alessio, Jeanjean and Montecchiari [5], Montecchiari and Rabinowitz [10] and references therein.

Motived by papers [8] and [12], we intend to establish the existence of a heteroclinic solution for the equation (PDE) under the Neumann boundary conditions by working with the Classes 1 and 2. As in the above papers, we have used variational method, more precisely minimization technical on a special set, however new ideas have been introduced in the study of the problem, see for example, Proposition 3.1 in Section 2. The regularity and behavior of the heteroclinic are obtained by using the same arguments found in [8].

Our main results are the following

Theorem 1.1. Assume $(V_1) - (V_3)$, $\epsilon = 1$ and that A belongs to Class 1. Then problem (PDE)-(NC) has a heteroclinic solution from 1 to -1.

Theorem 1.2. Assume $(V_1) - (V_3)$ and that A belongs to Class 2. Then, there is $\epsilon_0 > 0$ such that problem (PDE)-(NC) possesses a heteroclinic solution from 1 to -1 for all $\epsilon \in (0, \epsilon_0)$.

The plan of the paper is as follows: In Section 2, we prove some technical results, which will be useful to prove the above theorems. In Section 3 we prove the Theorem 1.1, while in Section 4 we prove the Theorem 1.2.

2 Preliminary Results

Consider the problem (PDE)-(NC) with $\epsilon = 1$, more precisely,

$$\begin{cases} -\Delta u + A(x,y)V'(u) = 0, \quad \forall \ (x,y) \in \Omega = \mathbb{R} \times \mathcal{D}, \\ \frac{\partial u}{\partial \nu}(x,y) = 0, \quad \forall x \in \mathbb{R}, \ y \in \partial \mathcal{D}. \end{cases}$$

In the sequel, we define the set

$$\Gamma = \{ U \in W_{loc}^{1,2}(\Omega) : |\nabla U| \in L^{2}(\Omega), \|P_{k}U - 1\|_{L^{2}(\Omega_{1})} \to 0 \text{ as } (2.1) \\ k \to -\infty \text{ and } \|P_{k}U + 1\|_{L^{2}(\Omega_{1})} \to 0 \text{ as } k \to +\infty \},$$

where $\Omega_1 = (0, 1) \times \mathcal{D}$ and

$$P_k U(x,y) = U(x+k,y), \text{ for } (x,y) \in \Omega \text{ and } k \in \mathbb{Z}.$$

It is very important to observe that $\Gamma \neq \emptyset$, because the function Φ given by

$$\Phi(x,y) = \begin{cases}
1, & \text{if} \quad x \le j, y \in \mathcal{D}, \\
2j+1-2x, & \text{if} \quad j < x \le j+1, y \in \mathcal{D}, \\
-1, & \text{if} \quad j+1 < x, y \in \mathcal{D},
\end{cases}$$
(2.2)

belongs to Γ . Furthermore, we also fix

$$\mathcal{L}(u) = \frac{1}{2} |\nabla u|^2 + A(x, y)V(u)$$

and the functionals $J: \Gamma \to \mathbb{R} \cup \{+\infty\}$ given by

$$J(U) = \sum_{k \in \mathbb{Z}} I_k(U) \tag{2.3}$$

and $I_k: W^{1,2}((k, k+1) \times \mathcal{D}) \to \mathbb{R}$ defined by

$$I_k(U) = \int_k^{k+1} \int_{\mathcal{D}} \mathcal{L}(U) dx dy.$$

Associated with functional J we have the number

$$\Theta^* = \inf\{J(U) : U \in \Gamma\}.$$
(2.4)

By (2.2), $\Phi \in \Gamma$, then $\Theta^* < +\infty$. From definition of Θ^* , there exists a minimizing sequence $(U_n) \subset \Gamma$ for J, that is,

$$J(U_n) \to \Theta^* \text{ as } n \to \infty.$$
 (2.5)

Without loss of generality, we can assume that (U_n) verifies

$$-1 \le U_n(x,y) \le 1, \quad \forall (x,y) \in \Omega.$$
(2.6)

Indeed, for each $n \in \mathbb{N}$ let us consider

$$\tilde{U}_n(x,y) = \begin{cases} -1, & \text{if } U_n(x,y) \le -1, \\ U_n(x,y), & \text{if } -1 \le U_n(x,y) \le 1, \\ 1, & \text{if } U_n(x,y) \ge 1. \end{cases}$$

It is easy to check that $\tilde{U}_n \in W^{1,2}_{loc}(\Omega)$ with

$$|\tilde{U}_n(x,y) - 1| \le |U_n(x,y) - 1|, \quad \forall (x,y) \in \Omega$$

and

$$|\tilde{U}_n(x,y)+1| \le |U_n(x,y)+1|, \quad \forall (x,y) \in \Omega.$$

Hence $(\tilde{U}_n) \subset \Gamma$, and so,

$$\Theta^* \le J(U_n), \quad \forall n \in \mathbb{N}.$$

Since

$$J(U_n) \le J(U_n), \quad \forall n \in \mathbb{N},$$

it follows that

$$\Theta^* \le J(\tilde{U}_n) \le J(U_n) = \Theta^* + o_n(1),$$

thereby showing that (\tilde{U}_n) is also a minimizing sequence for J on Γ with

$$-1 \le \tilde{U}_n(x,y) \le 1, \quad \forall (x,y) \in \Omega.$$

From (2.5)-(2.6), there is M > 0 independent of k and m such that

$$\||\nabla U_m|\|_{L^2((k,k+1)\times\mathcal{D})} + \|U_m\|_{L^2((k,k+1)\times\mathcal{D})} \le M, \quad \forall m \in \mathbb{N} \quad \text{and} \quad k \in \mathbb{Z}.$$

Consequently, (U_n) is bounded in $E_k = W^{1,2}((k, k+1) \times \mathcal{D})$, endowed with the usual norm, for all $k \in \mathbb{Z}$. Then for some subsequence, there is $U \in W^{1,2}_{loc}(\Omega)$ such that

$$U_n \rightharpoonup U$$
 in $E_k, \quad \forall k \in \mathbb{Z},$ (2.7)

$$U_n \to U$$
 in $L^2((k, k+1) \times \mathcal{D}), \quad \forall k \in \mathbb{Z},$ (2.8)

and

$$U_n(x,y) \to U(x,y), \quad \text{a.e. in} \quad \Omega.$$
 (2.9)

Therefore, from (2.5)-(2.9),

$$J(U) \le \Theta^*$$
 and $-1 \le U(x, y) \le 1$, a.e. in Ω . (2.10)

In the next section, our main goal is proving that U is the desired heteroclinic solution, and in this point, the conditions on function A play their role. However, before doing that we need to say that if A is 1- periodic in x, the same arguments explored in [8] guarantee the existence of a heteroclinic solution W^* from 1 to -1.

3 Proof of Theorem 1.1: *A* is asymptotic at infinity to a periodic function

By hypothesis,

$$A(x,y) < A_p(x,y), \quad \forall (x,y) \in \Omega.$$

Then, if $W^* \in \Gamma$ is a heteroclinic solution for the periodic case, we must have

$$\Theta^* \le J(W^*) < J_p(W^*) = \Theta_p^*,$$

that is,

$$\Theta^* < \Theta_p^*. \tag{3.1}$$

The last inequality will be a key point in our approach. In what follows, $(U_m) \subset \Gamma$ is a minimizing sequence for J with

$$-1 \le U_m(x,y) \le 1, \quad \forall (x,y) \in \Omega.$$

By using the fact that $(U_n) \subset \Gamma$, given $\tau \in (0, \sqrt{|\Omega_1|})$, for each $m \in \mathbb{N}$, there are $k_1(m), k_2(m) \in \mathbb{Z}$ such that

$$||P_{-j}Q_m - 1||_{L^2(\Omega_1)} < \tau, \quad ||Q_m - 1||_{L^2(\Omega_1)} \ge \tau$$
 (3.2)

and

$$||P_j R_m + 1||_{L^2(\Omega_1)} < \tau, \quad ||R_m + 1||_{L^2(\Omega_1)} \ge \tau$$
 (3.3)

for all $j \in \mathbb{N}$, where

$$Q_m(x,y) = U_m(x - k_1(m), y)$$
 and $R_m(x,y) = U_m(x + k_2(m), y), \quad \forall (x,y) \in \Omega.$

The reader is invited to observe that only one of the following conditions holds:

(I) $k_1(m), k_2(m) < 0$ for some subsequence. (II) $k_1(m) > 0$ for all $m \in \mathbb{N}$, and $k_2(m) < 0$ for some subsequence. (III) $k_2(m) > 0$ for all $m \in \mathbb{N}$, and $k_1(m) < 0$ for some subsequence. (IV) $k_1(m), k_2(m) > 0$ for all $m \in \mathbb{N}$.

The above conditions are crucial to prove the following proposition

Proposition 3.1. (Main proposition) There is $j_0 \in \mathbb{N}$ such that

$$\|U - 1\|_{L^2((-j, -j+1) \times \mathcal{D})} \le \tau \quad and \quad \|U + 1\|_{L^2((j, j+1) \times \mathcal{D})} \le \tau, \quad \forall j \ge j_0.$$
(3.4)

We will assume for a moment that Proposition 3.1 is proved and show Theorem 1.1.

Proof of Theorem 1.1 From the limit $J(U_n) \to \Theta^*$, we get

$$\sum_{j\in\mathbb{Z}} I_j(U) = J(U) \le \Theta^*, \tag{3.5}$$

from where it follows that

$$I_j(U) \to 0 \quad \text{as} \quad j \to -\infty,$$

or equivalently

$$\int_0^1 \int_{\mathcal{D}} |\nabla P_{-j}U|^2 \, dx \, dy + \int_0^1 \int_{\mathcal{D}} A(x, y) V(P_{-j}U) \, dx \, dy \to 0 \quad \text{as} \quad j \to -\infty.$$

As $U \in L^{\infty}(\Omega)$, we have that $(P_{-j}U)$ is a bounded sequence in $W^{1,2}(\Omega_1)$. Thus, there is a subsequence $(P_{-jk}U)$ of $(P_{-j}U)$ and $\hat{U} \in W^{1,2}(\Omega_1)$ such that

$$P_{-j_k}U \rightarrow \hat{U}$$
 in $W^{1,2}(\Omega_1)$ as $j_k \rightarrow +\infty$

$$P_{-j_k}U \to \hat{U}$$
 in $L^2(\Omega_1)$ as $j_k \to +\infty$

and

$$P_{-j_k}U(x,y) \to U(x,y)$$
 a.e. in Ω_1 , as $j_k \to +\infty$.

From this,

$$\int_0^1 \int_{\mathcal{D}} V(\hat{U}) \, dx dy = 0,$$

then

$$\hat{U} = 1$$
 or $\hat{U} = -1$,

and so,

$$P_{-j_k}U \to 1$$
 or $P_{-j_k}U \to -1$ in $L^2(\Omega_1)$ as $j_k \to +\infty$.

Since $\tau \in (0, \sqrt{|\Omega_1|})$, these limits combine with (3.4) to give

$$P_{-j_k}U \to 1$$
 in $L^2(\Omega_1)$ as $j_k \to +\infty$.

The above argument also yields

$$P_{-j}U \to 1$$
 in $L^2(\Omega_1)$ as $j \to +\infty$.

Similar reasoning proves

$$P_j U \to -1$$
 in $L^2(\Omega_1)$ as $j \to +\infty$.

Consequently, $U \in \Gamma$ and $-1 \leq U(x, y) \leq 1$ for all $(x, y) \in \overline{\Omega}$. Moreover, by (3.5),

 $J(U) = \Theta^*.$

Now, we claim that for each $\phi \in C_0^{\infty}(\overline{\Omega})$, we have $\frac{\partial J}{\partial \phi}(U) = 0$, where $\frac{\partial J}{\partial \phi}(U)$ denotes the directional derivative of J at U in the direction of $\phi \in C_0^{\infty}(\overline{\Omega})$, where

$$C_0^{\infty}(\overline{\Omega}) = \{ \phi : \overline{\Omega} \to \mathbb{R} : \exists \psi \in C_0^{\infty}(\mathbb{R}^N, \mathbb{R}) \text{ such that } \psi(x) = \phi(x), \ \forall x \in \overline{\Omega} \}.$$

Indeed, taking $w = U + t\phi$ with $\phi \in C_0^{\infty}(\overline{\Omega})$ and $t \in \mathbb{R}$, we derive that for k large enough, let's say, $|k| > \ell_0$, we have

$$I_k(U+t\phi) = I_k(U), \ \forall |k| > \ell_0.$$

Thereby

$$\frac{J(U+t\phi) - J(U)}{t} = \frac{1}{t} \left(\sum_{k \in \mathbb{Z}} (I_k(U+t\phi) - I_k(U)) = \sum_{k=-\ell_0}^{\ell_0} \left(\frac{I_k(U+t\phi) - I_k(U)}{t} \right),$$

and so,

$$\frac{\partial J}{\partial \phi}(U) = \lim_{t \to 0} \frac{J(U + t\phi) - J(U)}{t} = \sum_{k=-\ell_0}^{\ell_0} I'_k(U)\phi.$$

As $w \in \Gamma$ and $J(U) \leq J(w)$, a standard argument ensures that $\frac{\partial J}{\partial \phi}(U) = 0$, for all $\phi \in C_0^{\infty}(\overline{\Omega})$. Therefore,

$$\int_{\Omega} \nabla U \nabla \phi \, dx + \int_{\Omega} A(x, y) V(U) \phi \, dx = 0, \quad \forall \phi \in C_0^{\infty}(\overline{\Omega}).$$

From this, U is a weak solution of (PDE). A regularity argument from [8, Section 6] implies that $U \in C^2(\overline{\Omega}, \mathbb{R})$, and that U is a classical solution of

$$-\Delta U + A(x,y)V'(U) = 0$$
, in Ω and $\frac{\partial U}{\partial \nu} = 0$, $x \in \mathbb{R}$, $y \in \partial \mathcal{D}$,

with

 $U(x,y) \to 1$ as $x \to -\infty$ and $U(x,y) \to -1$ as $x \to +\infty$, uniformly in $y \in \mathcal{D}$.

From this, U is a heteroclinic solution from 1 to -1, which finishes the proof of Theorem 1.1. \Box

Proof of Proposition 3.1 Note that if (I) holds, then

$$\|U_m - 1\|_{L^2((-j-1,-j)\times\mathcal{D})} \le \tau \quad \text{and} \quad \|U_m + 1\|_{L^2((j,j+1)\times\mathcal{D})} \le \tau, \quad \forall j, m \in \mathbb{N}.$$

This together with (2.8) yield

$$\|U-1\|_{L^2((-j-1,-j)\times\mathcal{D})} \le \tau \quad \text{and} \quad \|U+1\|_{L^2((j,j+1)\times\mathcal{D})} \le \tau, \quad \forall j \in \mathbb{N},$$

and the proposition follows with $j_0 = 0$.

Now, we will prove the proposition by supposing that (II) holds. To begin with, we make the following claim

Claim 3.1. $(k_1(m))$ is bounded.

In what follows, let us denote $(k_1(m))$ by (k(m)). Assume by contradiction that there is a subsequence of (k(m)), still denoted by itself, with $k(m) \to +\infty$. The boundedness of (U_m) in E_k , implies that (Q_m) is also bounded in E_k . Thus, for some subsequence, there is $W \in W_{loc}^{1,2}(\Omega)$ such that

$$Q_m \rightharpoonup W \quad \text{in} \quad E_k, \quad \forall k \in \mathbb{Z},$$

$$(3.6)$$

$$Q_m \to W$$
in $L^2((k, k+1) \times \mathcal{D}), \quad \forall k \in \mathbb{Z},$ (3.7)

$$Q_m(x,y) \to W(x,y), \quad \text{a.e. in} \quad \Omega,$$
(3.8)

and

$$-1 \le W(x, y) \le 1, \quad \text{a.e. in} \quad \Omega. \tag{3.9}$$

A simples change of variables gives us

$$\sum_{k \in \mathbb{Z}} \tilde{I}_k(Q_m) = J(U_m) = \Theta^* + o_m(1) \le \Theta^* + 1$$
(3.10)

where

$$\tilde{I}_k(U) = \int_k^{k+1} \int_{\mathcal{D}} \tilde{\mathbf{L}}_m(U) dx dy$$

and

$$\tilde{\mathbf{L}}_m(u) = \frac{1}{2} |\nabla u|^2 + A(x - k(m), y)V(u).$$

Consequently, the Fatou's Lemma together with (A_1) and (3.6)-(3.10) provides

$$J_p(W) \le \Theta^*, \tag{3.11}$$

which gives

$$I_{p,-j}(W) \to 0 \quad \text{as} \quad j \to +\infty.$$
 (3.12)

Setting for each $j \in \mathbb{N}$ the function $\widetilde{W}_j = P_{-j}W$, the fact that $W \in L^{\infty}(\Omega)$ together with the Sobolev embeddings guarantee the existence of $W_0 \in L^2(\Omega_1)$, and a subsequence of (\widetilde{W}_j) , still denoted by itself, such that

$$\widetilde{W}_j \to W_0 \quad \text{in} \quad L^2(\Omega_1),$$

 $\|\widetilde{W}_j - W_0\|_{L^2(\Omega_1)} \to 0.$ (3.13)

that is,

This limit and (3.2) lead to

$$||W_0 - 1||_{L^2(\Omega_1)} \le \tau.$$

On the other hand, by (3.12),

$$I_{p,0}(W_0) = 0$$

from where it follows that $W_0 = 1$ or $W_0 = -1$. As $\tau \in (0, \sqrt{|\Omega_1|})$, we must have $W_0 = 1$. Thereby,

$$\|\widetilde{W}_j - 1\|_{L^2(\Omega_1)} \to 0 \quad \text{as} \quad j \to +\infty.$$
(3.14)

Now, fixing $W_j = P_j W$ for $j \in \mathbb{N}$, the same reasoning works to show that there exists $\widehat{W}_0 \in L^2(\Omega_1)$ and a subsequence of (W_j) , still denoted by itself, such that

$$W_j \to \widehat{W}_0 \text{ in } L^2(\Omega_1) \quad \text{as} \quad j \to \infty,$$
(3.15)

or equivalently,

$$\|W_j - \widetilde{W}_0\|_{L^2(\Omega_1)} \to 0.$$
(3.16)

This information gathering with the limit

$$I_{p,j}(W) \to 0$$
 as $j \to +\infty$

leads to $\widehat{W}_0 = 1$ or $\widehat{W}_0 = -1$. Next we are going to show that $\widehat{W}_0 = -1$. To see why, assume by contradiction that $\widehat{W}_0 = 1$. From (3.2) and (3.15), there is $j_1 \in \mathbb{N}$ such that

$$||W - 1||_{L^2((j_1 - 1, j_1) \times D)} \ge \tau$$
 and $||W - 1||_{L^2((j_1, j_1 + 1) \times D)} \le \tau$

As $Q_m \to W$ in $L^2((j_1 - 1, j_1 + 1) \times \mathcal{D})$, there is $m_0 \in \mathbb{N}$ satisfying

$$||Q_m - 1||_{L^2((j_1 - 1, j_1) \times \mathcal{D})} \ge \tau/2$$
 and $||Q_m - 1||_{L^2((j_1, j_1 + 1) \times \mathcal{D})} \le 2\tau$, $\forall m \ge m_0$.

In what follows, we denoted by $\beta = \beta(\tau)$ the real number given by

$$\frac{\beta}{\tilde{A}_0} = \inf_{u \in \mathcal{N}_\tau} I_{*,\tau}(u),$$

where $\tilde{A}_0 = \min\{1, A_0\},\$

 $\mathcal{N}_{\tau} = \{ u \in W^{1,2}((-1,1) \times \mathcal{D}) : \|u-1\|_{L^{2}((-1,0) \times \mathcal{D})} \ge \tau/2 \text{ and } \|u-1\|_{L^{2}((0,1) \times \mathcal{D})} \le 2\tau \}$ and $I_{*,\tau} : W^{1,2}((-1,1) \times \mathcal{D}) \to \mathbb{R}$ is defined by

$$I_{*,\tau}(u) = \int_{-1}^{1} \int_{\mathcal{D}} (|\nabla u|^2 + V(u)) \, dx dy.$$

Hence, by a simple change of variable

$$\int_{j_1-1}^{j_1+1} \int_{\mathcal{D}} (|\nabla Q_m|^2 + V(Q_m)) \, dx dy \ge \frac{\beta}{\tilde{A}_0}, \quad \forall m \ge m_0. \tag{3.17}$$

Here we would like point out that the same arguments found in [8, Proposition 2.14] work to show that $\beta > 0$. Having this in mind, we can assume without loss of generality that

$$J(U_m) \le \Theta^* + \beta/4, \quad \forall m \ge m_0. \tag{3.18}$$

In the sequel, for each $j \ge j_1 + 2$ and $m \ge m_0$, let us consider the function

$$Z_{j,m}(x,y) = \begin{cases} 1, & \text{if } x \leq j, y \in \mathcal{D}, \\ ((j+1)-x) + (x-j)Q_m(x,y), & \text{if } j < x \leq j+1, y \in \mathcal{D}, \\ Q_m(x,y), & \text{if } j+1 < x, y \in \mathcal{D}. \end{cases}$$

By a direct computation, we see that $Z_{j,m} \in \Gamma$ and

$$J_p(Z_{j,m}) = I_{p,j}(Z_{j,m}) + \sum_{k=j+1}^{\infty} I_{p,k}(Q_m) = I_{p,j}(Z_{j,m}) + \sum_{k=j+1+k(m)}^{\infty} I_{p,k}(U_m),$$

and so,

$$\Theta_p^* \le J_p(Z_{j,m}) = I_{p,j}(Z_{j,m}) + \sum_{k=j+1+k(m)}^{\infty} I_{p,k}(U_m).$$

As A verifies $(A_1) - (A_2)$ and $(J(U_n))$ is bounded, increasing m_0 if necessary, we have

$$\sum_{k=j+1+k(m)}^{\infty} I_{p,k}(U_m) \le \sum_{k=j+1+k(m)}^{\infty} I_k(U_m) + \beta/4, \quad \forall m \ge m_0.$$
(3.19)

Now, as $j \ge j_1 + 2$, (3.19) implies in the inequality

$$\Theta_p^* \le I_{p,j}(Z_{j,m}) + J(U_m) - \tilde{A}_0 \int_{j_{1}-1}^{j_{1}+1} \int_{\mathcal{D}} (|\nabla Q_m|^2 + V(Q_m)) \, dxdy + \beta/4,$$

which combine with (3.17)-(3.18) to give

$$\Theta_p^* \le I_{p,j}(Z_{j,m}) + \Theta^* + \frac{\beta}{4} - \tilde{A}_0 \frac{\beta}{\tilde{A}_0} + \frac{\beta}{4} = I_{p,j}(Z_{j,m}) + \Theta^* - \frac{\beta}{2}.$$
 (3.20)

Since

$$-1 \le W_j(x,y) \le 1$$
 and $W_j \to 1$ in $W^{1,2}(\Omega_1)$ as $j \to +\infty$,

it is easy to check that

$$\lim_{j \to +\infty} \int_0^1 \int_{\mathcal{D}} A(x+j,y) V((-x+1+xW_j) \, dx \, dy = 0$$

and

$$\lim_{j \to +\infty} \int_0^1 \int_{\mathcal{D}} |1 - W_j|^2 \, dx \, dy = 0.$$

Thus, given $\delta > 0$, there is $j_0 = j_0(\delta) > j_1 + 2$, which is independent of m, such that

$$\int_0^1 \int_{\mathcal{D}} A(x+j,y) V(-x+1+xW_j) \, dx \, dy < \delta, \quad \forall j \ge j_0 \tag{3.21}$$

and

$$\int_0^1 \int_{\mathcal{D}} |1 - W_j|^2 \, dx \, dy < \delta, \quad \forall j \ge j_0. \tag{3.22}$$

To continue, we further claim there is $j = j(m) \ge j_0$ and $m \ge m_0$ such that

$$I_{p,j}(Z_{j,m}) = \int_{j}^{j+1} \int_{\mathcal{D}} \mathcal{L}_{p}(Z_{j,m}) dx dy < \beta/2.$$
(3.23)

If the claim does not hold, for each $j \ge j_0$, there exists $m_1 = m_1(j) \ge m_0$ verifying

$$\int_{j}^{j+1} \int_{\mathcal{D}} \mathcal{L}_{p}(Z_{j,m}) dx dy \ge \beta/2, \quad \forall m \ge m_{1}.$$

From definition of $Z_{j,m}$ and (A_2) ,

$$\int_{j}^{j+1} |\nabla Z_{j,m}|^2 \, dx \, dx \ge \beta/2 - \int_{j}^{j+1} \int_{\mathcal{D}} A(x,y) V((j+1) - x + (x-j)Q_m) \, dx \, dy.$$

Recalling that

$$\lim_{m \to +\infty} \int_{j}^{j+1} \int_{\mathcal{D}} A(x,y) V((j+1) - x + (x-j)Q_m) \, dx dy = \int_{0}^{1} \int_{\mathcal{D}} A(x+j,y) V((-x+1+xW_j) \, dx dy < \delta,$$

for $j \ge j_0$ and $\delta < \beta/4$, there exists $m_2 = m_2(j) \ge m_1(j)$ such that

$$\int_{j}^{j+1} \int_{\mathcal{D}} |\nabla Z_{j,m}|^2 \, dx \, dx \ge \beta/4, \quad \forall m \ge m_2.$$

Using again the definition of $Z_{i,m}$, there is a constant C > 0 such that

$$\int_{j}^{j+1} \int_{\mathcal{D}} |\nabla Z_{j,m}|^2 \, dx dy \le C \left(\int_0^1 \int_{\mathcal{D}} |1 - P_j(Q_m)|^2 \, dx dy + \int_j^{j+1} \int_{\mathcal{D}} |\nabla Q_m|^2 \, dx dy \right).$$

Now, fixing $\delta < \frac{\beta}{8C}$ in (3.22), we obtain

$$\int_{j}^{j+1} \int_{\mathcal{D}} |\nabla Q_m|^2 \, dx \, dy \ge \beta/8, \quad \forall m \ge m_2(j).$$

Let $l \in \mathbb{N}$ such that

$$(l+1)\beta/8 > \Theta^* + 1$$

and fix $m > \max\{m_2(j) : j_0 \le j \le j_0 + l\}$. Then,

$$\sum_{k\in\mathbb{Z}}\tilde{I}_k(Q_m)\geq\Theta^*+1,$$

which contradicts (3.10), thereby showing (3.23). Thus, by (3.20) and (3.23),

 $\Theta_p^* < \Theta^*,$

contrary to (3.1), and this ensures that $\widehat{W}_0 = -1$. From the above study, we deduce that $W \in \Gamma$, then by (3.11),

$$\Theta_p^* \le J(W) \le \Theta^*,$$

which is absurd. This proves the Claim 3.1. Hence, there is a subsequence $(k_1(m))$, still denoted by itself, and $k_* \in \mathbb{N}$ such that $k_1(m) = k_*$ for all m in \mathbb{N} . Hence, the inequality below

$$||P_{-j}Q_m - 1||_{L^2(\Omega_1)} < \tau, \quad \forall m \in \mathbb{N} \quad \text{and} \quad j \in \mathbb{N}$$

is equivalent to

$$||U_m - 1||_{L^2((j-1,j) \times \mathcal{D})} < \tau, \quad \forall m \in \mathbb{N} \text{ and } \forall j \le -k_*$$

This inequality combined with (2.8) gives

$$||U - 1||_{L^2((j-1,j) \times \mathcal{D})} \le \tau, \quad \forall j \le -k_*.$$

Therefore, when (II) occurs, the Proposition 3.1 holds with $j_0 = k_*$. The cases (III) and (IV) can be analyzed of the same way, then we omit their proofs, and the proposition is proved. \Box

4 Proof of Theorem 1.2: A verifies the Rabinowitz's condition

In this section we establish the existence of a heteroclinic solution for the Class 2. In what follows, we are considering the equation

$$-\Delta u + A(\epsilon x, y)V'(u) = 0, \quad \text{in} \quad \Omega, \tag{P_{\epsilon}}$$

together with the Neumann boundary condition

$$\frac{\partial u}{\partial \nu}(x,y) = 0, \ x \in \mathbb{R}, y \in \partial \mathcal{D},$$
 (NC)

where ϵ is a positive parameter and A satisfies

$$(A_3) \qquad 0 < A_0 = A(0, y) = \inf_{(x,y) \in \Omega} A(x, y) \le \liminf_{|(x,y)| \to +\infty} A(x, y) = A_\infty < \infty, \quad \forall y \in \mathcal{D}.$$

From now on, we are denoting by $J_{\epsilon}, J_{\infty}: \Gamma \to \mathbb{R} \cup \{+\infty\}$ the functionals

$$J_{\epsilon}(U) = \sum_{k \in \mathbb{Z}} I_{\epsilon,k}(U)$$

and

$$J_{\infty}(U) = \sum_{k \in \mathbb{Z}} I_{\infty,k}(U),$$

where $I_{\epsilon,k}, I_{\infty,k} : E_k \to \mathbb{R}$ are given by

$$I_{\epsilon,k}(U) = \int_{k}^{k+1} \int_{\mathcal{D}} \left(|\nabla U|^2 + A(\epsilon x, y) V(U) \right) \, dx dy$$

and

$$I_{\infty,k}(U) = \int_{k}^{k+1} \int_{\mathcal{D}} \left(|\nabla U|^2 + A_{\infty} V(U) \right) \, dx \, dy.$$

Moreover, we denote by Θ_{ϵ} and Θ_{∞} the following numbers

$$\Theta_{\epsilon} = \inf\{J_{\epsilon}(U) : U \in \Gamma\}$$

and

$$\Theta_{\infty} = \inf\{J_{\infty}(U) : U \in \Gamma\}$$

By Section 2, we know that there are $W_0, W_\infty \in \Gamma$ verifying $J_0(W_0) = \Theta_0$ and $J_\infty(W_\infty) = \Theta_\infty$. This fact permit us to prove the following lemma

Lemma 4.1. $\limsup_{\epsilon \to 0} \Theta_{\epsilon} \leq \Theta_0$ and $\Theta_0 < \Theta_{\infty}$.

Proof. For each $\epsilon > 0$,

 $\Theta_{\epsilon} \leq J_{\epsilon}(W_0).$

Since

$$\lim_{\epsilon \to 0} J_{\epsilon}(W_0) = J_0(W_0) = \Theta_0$$

it follows that

$$\limsup_{\epsilon \to 0} \Theta_{\epsilon} \le \Theta_0.$$

On the other hand, by (A_3) ,

$$\Theta_0 \le J_0(W_0) < J_\infty(W_\infty) = \Theta_\infty,$$

which shows the lemma.

In the sequel, we fix $\epsilon_0 > 0$ small enough a such way that

$$\Theta_{\epsilon} < \Theta_{\infty}, \quad \forall \epsilon \in (0, \epsilon_0).$$
 (4.1)

4.1 Proof of Theorem 1.2

Arguing as in Section 2, for each $\epsilon > 0$ there is a minimizing sequence $(U_n) \subset \Gamma$ with $-1 \leq U_n(x, y) \leq 1$ for all $(x, y) \in \Omega$ and $U \in W^{1,2}_{loc}(\Omega)$ such that

$$J_{\epsilon}(U_n) \to \Theta_{\epsilon},$$

$$U_n \rightharpoonup U \quad \text{in} \quad E_k, \quad \forall k \in \mathbb{Z},$$

$$U_n \to U \quad \text{in} \quad L^2((k, k+1) \times \mathcal{D}), \quad \forall k \in \mathbb{Z},$$

$$U_n(x, y) \to U(x, y), \quad \text{a.e. in} \quad \Omega,$$

$$-1 \le U(x, y) \le 1, \quad \forall (x, y) \in \Omega,$$

and

$$J_{\epsilon}(U) \le \Theta_{\epsilon}.\tag{4.2}$$

In the sequel, we will use the same approach explored in Section 3. As $(U_n) \subset \Gamma$ and $\tau \in (0, \sqrt{|\Omega_1|})$, for each $m \in \mathbb{N}$ there are $k_1(m), k_2(m) \in \mathbb{Z}$ such that

$$||P_{-j}Q_m - 1||_{L^2(\Omega_1)} < \tau, \quad ||Q_m - 1||_{L^2(\Omega_1)} \ge \tau$$
 (4.3)

and

$$||P_j R_m + 1||_{L^2(\Omega_1)} < \tau, \quad ||R_m + 1||_{L^2(\Omega_1)} \ge \tau$$
(4.4)

for all $j \in \mathbb{N}$, where

$$Q_m(x,y) = U_m(x - k_1(m), y)$$
 and $R_m(x,y) = U_m(x + k_2(m), y), \quad \forall (x,y) \in \Omega.$

Proposition 4.1. If $\epsilon \in (0, \epsilon_0)$, then there is $j_0 \in \mathbb{N}$ such that

$$||U - 1||_{L^2((-j, -j+1) \times \mathcal{D})} \le \tau \quad and \quad ||U + 1||_{L^2((j, j+1) \times \mathcal{D})} \le \tau, \quad \forall j \ge j_0.$$
(4.5)

Proof. As in the proof of Proposition 3.1, we must study the cases (I) - (IV). The case (I) follows of the same way, however for the other cases we need to do some modifications. Next, we will consider the case (II). As in the last section, we begin by showing the claim below

Claim 4.1. $(k_1(m))$ is bounded.

In what follows we denote $k_1(m)$ by k(m). Assume by contradiction that there is a $\epsilon \in (0, \epsilon_0)$ such that (k(m)) is unbounded and $k(m) \to +\infty$. The boundedness of (U_m) in E_k implies that (Q_m) is also bounded in E_k for all $k \in \mathbb{N}$. Thus, for some subsequence, there is $W \in W^{1,2}_{loc}(\Omega)$ such that

$$Q_m \to W \quad \text{in} \quad E_k, \quad \forall k \in \mathbb{N},$$

$$(4.6)$$

$$Q_m(x,y) \to W(x,y), \quad \text{a.e. in} \quad \Omega,$$

$$(4.7)$$

and

$$-1 \le W(x, y) \le 1$$
, a.e. in Ω . (4.8)

By a simple change variable,

$$\sum_{k\in\mathbb{Z}} \tilde{I}_k(W_m) \le J_{\epsilon}(U_m) = \Theta_{\epsilon} + o_m(1)$$
(4.9)

where

$$\tilde{I}_{\epsilon,k}(U) = \int_{k}^{k+1} \int_{\mathcal{D}} \tilde{\mathbf{L}}_{\epsilon,m}(U) dx dy$$

with

$$\tilde{\mathcal{L}}_{\epsilon,m}(u) = \frac{1}{2} |\nabla u|^2 + A(\epsilon x - \epsilon k(m), y) V(u).$$

Now, the Fatou's Lemma combined with (4.9) leads to

$$J_{\infty}(W) \le \Theta_{\epsilon}.\tag{4.10}$$

Then,

$$I_{\infty,j}(W) \to 0 \quad \text{as} \quad j \to +\infty.$$
 (4.11)

Setting for each $j \in \mathbb{N}$ the function $\widetilde{W}_j = P_{-j}W$, the fact that $W \in L^{\infty}(\Omega)$ implies that there are $W_0 \in W^{1,2}_{loc}(\Omega, \mathbb{R})$ and a subsequence of (\widetilde{W}_j) , still denoted by itself, such that

 $\widetilde{W}_j \to W_0 \text{ in } W^{1,2}(\Omega_1) \quad \text{as} \quad j \to +\infty,$

and so,

$$\|\widetilde{W}_j - W_0\|_{L^2(\Omega_1)} \to 0.$$
 (4.12)

This limit combined (4.3) yields

 $||W_0 - 1||_{L^2(\Omega_1)} \le \tau.$

On the other hand, by (4.11),

$$I_{\infty,0}(W_0) = 0,$$

which gives $W_0 = 1$ or $W_0 = -1$. As $\tau \in (0, \sqrt{|\Omega_1|})$, we must have $W_0 = 1$. Then,

$$||W_j - 1||_{L^2(\Omega_1)} \to 0 \text{ as } j \to +\infty.$$
 (4.13)

By using the same type of argument, fixing $W_j = P_j W$ for $j \in \mathbb{N}$, it is possible to prove that there exist $\widehat{W}_0 \in W^{1,2}_{loc}(\overline{\Omega})$ and a subsequence of (W_j) , still denoted by itself, such that

$$W_j \to \widehat{W}_0 \text{ in } W^{1,2}(\Omega_1) \text{ as } j \to -\infty,$$

and so,

$$\|W_j - \widehat{W}_0\|_{L^2(\Omega_1)} \to 0.$$
 (4.14)

Thereby, $\widehat{W}_0 = 1$ or $\widehat{W}_0 = -1$. Here, as in the previous section, we have that $\widehat{W}_0 = -1$. Indeed, assuming by contradiction that $\widehat{W}_0 = 1$, we set the function

$$H_{j}(x,y) = \begin{cases} 1, & x \leq j, y \in \mathcal{D}, \\ ((j+1)-x) + (x-j)Q_{m}(x,y), & j < x \leq j+1, y \in \mathcal{D}, \\ Q_{m}(x,y), & j+1 < x, y \in \mathcal{D}. \end{cases}$$

Arguing as Proposition 3.1, we will find

$$\Theta_{\infty} \leq \Theta_{\epsilon}$$

which contradicts (4.1), and then $\widehat{W}_0 = -1$. Now we follow the same idea explored in Proposition 3.1 to conclude the proof.

Now, we are ready to prove Theorem 1.2.

Proof of Theorem 1.2 As an immediate consequence of the last proposition, for each $\epsilon \in (0, \epsilon_0)$, there is $j_0 \in \mathbb{N}$ such that

$$||U-1||_{L^2((-j,-j+1)\times\mathcal{D})} < \tau$$
 and $||U+1||_{L^2((j,j+1)\times\mathcal{D})} < \tau$, $\forall j \ge j_0$.

Now, arguing as in the proof of Theorem 1.1, it follows that $U \in C^2(\overline{\Omega}, \mathbb{R})$. Moreover, U is a classical solution of

$$-\Delta U + A(\epsilon x, y)V'(U) = 0$$
, in Ω and $\frac{\partial U}{\partial \nu} = 0, x \in \mathbb{R}, y \in \partial \mathcal{D}$

with

$$U(x,y) \to 1$$
 as $x \to -\infty$ and $U(x,y) \to -1$ as $x \to +\infty$, uniformly in $y \in \mathcal{D}$.

From this, U is a heteroclinic solution from 1 to -1, which finishes the proof of Theorem 1.2. \Box

Acknowledgment. The author would like to warmly thank Prof. Olimpio Hiroshi Miyagaki for several discussions about this subject, and also to Professor Rabinowitz by his comments that were very important to improve this manuscript. Moreover, the author would like to thank to the referee for his/her very nice remarks and suggestions, which were very important to improve this manuscript.

References

- S. Alama, L. Bronsard and C. Gui, Stationary layered solution in ℝ² for an Allen-Cahn system with multiple well potential, Calc. Var. Partial Differential Equations 5 (1997), 359–390.
- [2] F. Alessio and P. Montecchiari, *Entire solutions in* \mathbb{R}^2 for a class of Allen-Cahn equations, ESAIM: COCV 11 (2005), 633-672.
- [3] F. Alessio and P. Montecchiari, Layered solutions with multiple asymptotes for non autonomous Allen-Cahn equations in ℝ³, Calc. Var. Partial Differential Equations 46 (2013), 591-622.
- [4] F. Alessio, L. Jeanjean and P. Montecchiari, Stationary layered solutions in ℝ² for a class of non autonomous Allen-Cahn equations, Calc. Var. Partial Differential Equations 11 (2000), 177-202.
- [5] F. Alessio, L. Jeanjean and P. Montecchiari, Existence of infinitely many stationary layered solutions in ℝ² for a class of periodic Allen-Chan equations, Comm. Partial Differential Equations 27 (2002), 1537-1574.
- [6] F. Alessio, C. Gui and P. Montecchiari, Saddle Solutions to Allen-Cahn Equations in Doubly Periodic Media, Indiana Univ. Math. J. 65 No. 1 (2016), 199-221
- [7] C.O. Alves, P.C. Carrião and O.H. Miyagaki, Nonlinear perturbations of a periodic elliptic problem with critical growth, J. Math. Anal. Appl. 260 (2001), 133-146.
- [8] J. Byeon, P. Montecchiari and P. H. Rabinowitz, A Double well potential system, Analysis and PDE 9(7) (2016), 1737-1772.
- [9] Y. Jianfu and Z. Xiping, On the existence of nontrivial solution of quasilinear elliptic boundary value problem for unbounded domains, Acta Math. Sci. 7(3) 1987, 341-359.
- [10] P. Montecchiari and P. H. Rabinowitz, On the existence of multi-transition solutions for a class of elliptic systems, Ann. Inst. H. Poincaré Anal. Non Linéaire 33(1) (2016), 199219.
- [11] P.H. Rabinowitz, On a class of nonlinear Schrödinger equations, Z. Angew Math. Phys. 43 (1992), 270-291.
- [12] P. H. Rabinowitz, Solutions of heteroclinic type for some classes of semilinear elliptic partial differential equations, J. Math. Sci. Univ. Tokyo 1(3) (1994), 525550.
- [13] P.H. Rabinowitz, A new variational characterization of spatially heteroclinic solutions of a semilinear elliptic EDP, Discrete Contin. Dyn. Syst 10 (2004), 507-515.
- [14] P. H. Rabinowitz and E. Stredulinsky, On a class of infinite transition solutions for an Allen-Cahn model equation, Discrete Contin. Dynam. Systems 21 (2008), 207–20.

- [15] P. H. Rabinowitz and E. Stredulinsky, Mixed states for an Allen-Cahn type equation, Commun. Pure Appl. Anal. Vol. LVI (2003), 1078-2003
- [16] P. H. Rabinowitz and E. Stredulinsky, Mixed states for an Allen-Cahn equation, II, Cal. Var. 21 (2004), 157-207.