THE PRIMITIVE SPECTRUM OF A SEMIGROUP OF MARKOV OPERATORS

HENRIK KREIDLER

ABSTRACT. For a semigroup S of Markov operators on a space of continuous functions, we use S-invariant ideals to describe qualitative properties of S such as mean ergodicity and the structure of its fixed space. For this purpose we focus on *primitive S-ideals* and endow the space of those ideals with an appropriate topology. This approach is inspired by the representation theory of C*-algebras and can be adapted to our dynamical setting.

In the particularly important case of Koopman semigroups, we characterize the centers of attraction of the underlying dynamical system in terms of the invariant ideal structure of S.

Mathematics Subject Classification (2010). Primary 47A35, 47D03; Secondary 37B05, 37B25, 47B65.

1. INTRODUCTION

The primitive spectrum is a useful tool in the study of C*-algebras (see, e.g., Chapter IV of [Dix77], Section 4.3 of [Ped79] or Section II.6.5 of [Bla06]) and plays a crucial role in representation theory (cf. [Hof11]). Given a C*-algebra A it is defined as

 $Prim(A) := \{ \ker \pi \mid 0 \neq \pi \text{ irreducible representation of A} \}.$

Equipped with the *hull-kernel topology* (also called *Jacobson topology*) it becomes a quasi-compact T_0 -space. A nice application is the so called Dauns-Hofmann Theorem asserting that—in the unital case—the center of A is canonically isomorphic to C(Prim(A)).

In this note we study a dynamical version of the primitive spectrum in the commutative and unital case. Starting from a right amenable semigroup S of Markov operators on the space of continuous functions C(K) on some compact space K we introduce the primitive spectrum Prim(S) of S as the set of absolute kernels of ergodic measures. Again we equip the primitive spectrum with a hull-kernel topology and obtain a quasi-compact T_0 -space. We then describe the space C(Prim(S)) and give applications to topological dynamics and ergodic theory.

We now give a more detailed description of the results.

Based on two papers of H. H. Schaefer (see [Sch67] and [Sch68]) as well as Paragraph III.8 of [Sch74] we consider S-invariant ideals and measures in

The author wants to thank Roland Derndinger, Nikolai Edeko, Ulrich Groh and Rainer Nagel for ideas, suggestions and inspiring discussions. The author is also very grateful to the referee for their suggestions. Their thorough reports have led to major improvements of the article. In particular, their valuable advice vastly helped to improve the abstract, the introduction and the readability of the article overall.

HENRIK KREIDLER

Section 2 recalling some basic definitions and facts.

In the subsequent sections we introduce and study radical S-ideals. In Section 3 we give the definition and prove an equivalent characterization in the metric case (see Proposition 3.7). In the fourth section we then establish a close connection between radical S-ideals, centers of attraction appearing in topological dynamics and stability conditions of the semigroup (see Theorem 4.2 and Theorem 4.9).

The last three sections are devoted to the primitive spectrum of S as a topological space and its applications. In Section 5 we define the topology (cf. Proposition 5.3), state its basic properties and give some examples.

In Section 6 we then prove a Dauns-Hofmann-type theorem showing that if S is radical free (see Definition 3.1) the space of continuous functions on the primitive spectrum C(Prim(S)) is canonically isomorphic to the fixed space fix(S) of the semigroup S (see Theorem 6.3). We then extend this result to the general case of not necessarily radical free S (see Theorem 6.5).

As an application we obtain in Section 7 a new description of mean ergodicity of semigroups of Markov operators (see Theorem 7.1) which generalizes Schaefer's Theorem 2 of [Sch67] in two different ways. On one hand we consider the more general setting of right amenable semigroups instead of single operators. But more importantly, we obtain—in contrast to Schaefer's work—a full characterization of mean ergodicity . We finally look at some examples illustrating these results (cf. Examples 7.5).

It should be pointed out that while maximal invariant ideals (which have been the central objects in [Sch67] and [Sch68]) are enough to describe mean ergodic Markov operators and semigroups, the results of our paper show that primitive ideals are the natural algebraic structure to describe dynamical properties of general Markov semigroups, see also Remark 7.2 below.

In the following we always assume K to be a compact (Hausdorff) space. We denote the Banach lattice of continuous complex-valued functions on Kby C(K) and identify the dual space C(K)' of C(K) with the Banach lattice of complex regular Borel masures on K. Moreover, we refer to [Sch74] and [MN91] for Banach lattices and their ideal structure and remind the reader that the closed lattice ideals of C(K) coincide with the closed algebra ideals and are precisely the sets

$$I_L := \{ f \in \mathcal{C}(K) \mid f|_L = 0 \}$$

with $L \subseteq K$ closed. Recall also that a positive operator $T \in \mathscr{L}(\mathcal{C}(K))$ is called *Markov* if $T\mathbb{1} = \mathbb{1}$.

We now fix a semigroup $S \subseteq \mathscr{L}(C(K))$ of Markov operators which is *right* amenable (cf. Section 2.3 of [BJM89]) if endowed with the strong operator topology, i.e., there is a positive element $m \in C_b(S)'$ (called *right invariant* mean) such that m(1) = 1 and $m(R_S f) = m(f)$ for every $f \in C_b(S)$ where $R_S(f)(T) := f(TS)$ for all $T, S \in S$. All abelian topological semigroups and compact topological groups are amenable and, in particular, right amenable. For more examples and counterexamples we refer to [Day61] and Chapter 1 of [Pat88].

Note that the important cases of semigroups generated by a single operator and one-parameter semigroups are contained in our results since these are always abelian.

Many examples of Markov operators and semigroups arise from topological dynamical systems on K. In fact, if $\varphi \colon K \longrightarrow K$ is a continuous mapping, then the associated *Koopman operator* $T_{\varphi} \in \mathscr{L}(\mathcal{C}(K))$ defined by $f := f \circ \varphi$ for $f \in \mathcal{C}(K)$ is a Markov lattice operator. We write \mathcal{S}_{φ} for the semigroup $\{T_{\varphi}^{n} \mid n \in \mathbb{N}_{0}\}$.

2. Ergodic Measures and Primitive Ideals

In this section we introduce primitive S-ideals adapting concepts from the theory of C*-algebras and start with the following definition going back to H. H. Schaefer (see [Sch67]). Recall that an ideal I of C(K) is called *proper* if $I \neq C(K)$.

Definition 2.1. A closed proper ideal $I \subseteq C(K)$ is an S-*ideal* if it is S-invariant, i.e., $SI \subseteq I$ for each $S \in S$. It is called *maximal* if it is maximal among all S-ideals with respect to inclusion.

Remark 2.2. If I is an S-ideal, then a standard application of Zorn's lemma shows that I is contained in a maximal proper S-invariant ideal. Since there are no dense proper ideals in C(K), this ideal is already closed and therefore each S-ideal is contained in a maximal S-ideal (cf. Proposition 1 in [Sch67]).

In [Sin68] R. Sine used the concept of a self-supporting set of a Markov operator. Generalizing this to our setting, a non-empty closed set $L \subseteq K$ is called *self-supporting* if the measure $S'\delta_x \in C(K)'$ has support in L for each $x \in L$ and $S \in S$. Recall that here the *support* supp μ of a probability measure $\mu \in C(K)'$ is the smallest closed subset $A \subseteq K$ with $\mu(A) = 1$. Each self-supporting set L defines an S-ideal

$$I_L := \{ f \in \mathcal{C}(K) \mid f|_L = 0 \}.$$

Conversely, each S-ideal is an I_L for some self-supporting set L and the mapping $L \mapsto I_L$ is bijective. Moreover, each maximal S-ideal corresponds to a minimal self-supporting set.

Given an S-ideal I we call the unique self-supporting set L with $I_L = I$ the support of I and write L = supp I.

Remark 2.3. For each S-ideal I, the semigroup S induces a semigroup S_I of Markov operators on C(supp I) given by

$$\mathcal{S}_I := \{ S_I \mid S \in \mathcal{S} \}$$

with $S_I f := SF|_{\text{supp}I}$ for $S \in S$ and $f \in C(\text{supp}I)$ where $F \in C(K)$ is any extension of f to K. It is readily checked that I is maximal if and only if S_I is irreducible, i.e., there are no non-trivial S_I -ideals (see the corollary to Proposition III.8.2 in [Sch74]).

We are primarily interested in S-ideals defined by measures. The *absolute* kernel of a measure $0 \leq \mu \in C(K)'$ is

$$I_{\mu} := \{ f \in \mathcal{C}(K) \mid \langle |f|, \mu \rangle = 0 \}.$$

If μ is invariant, i.e., $S'\mu = \mu$ for each $S \in S$, this is an S-ideal.

We write $P_{\mathcal{S}}(K) \subseteq C(K)'$ for the space of invariant probability measures on K equipped with the weak* topology. By right amenability of S this is always a nonempty compact convex set (this is a simple consequence of Day's fixed point theorem, see Theorem 3 of [Day61]).

We recall that for each $\mu \in P_{\mathcal{S}}(K)$ the space $L^1(K,\mu)$ is the completion of $C(K)/I_{\mu}$ with respect to the L¹-norm. Since I_{μ} is S-invariant, every $S\in \mathbb{S}$ induces an operator on $\mathcal{C}(K)/I_{\mu}$ which then uniquely extends to a bi-Markov operator S_{μ} on $L^{1}(K, \mu)$, i.e., S_{μ} is a positive operator on $L^{1}(K, \mu)$ with $S_{\mu}\mathbb{1} = \mathbb{1}$ and $S'_{\mu}\mathbb{1} = \mathbb{1}$. We write $S_{\mu} := \{S_{\mu} \mid S \in S\}$ for the semigroup on $L^1(K, \mu)$ induced by S.

Definition 2.4. A measure $\mu \in P_{\mathcal{S}}(K)$ is called *ergodic* if the fixed space $fix(\mathcal{S}_{\mu})$ in $L^{1}(K,\mu)$ is one-dimensional.

The following characterization of ergodicity generalizes a result of M. Rosenblatt (cf. [Ros76]) and is well-known for single operators. We give a short proof in case of semigroup actions inspired by the proof of Proposition 10.4 of [EFHN15]. Here and in the following we write ex M for the set of extreme points of a convex subset M of a vector space.

Proposition 2.5. A measure $\mu \in P_{\mathcal{S}}(K)$ is ergodic if and only if $\mu \in$ $\exp P_{\mathcal{S}}(K).$

Proof. Assume that $fix(S_{\mu})$ is not one-dimensional. Since $fix(S_{\mu})$ is an ALsublattice of $L^1(K,\mu)$ with weak order unit 1, the set

 $B = \{ f \in \text{fix}(\mathcal{S}_{\mu}) \mid f \ge 0 \text{ and } \sup(f, \mathbb{1} - f) = 0 \}$

is total in fix(S_{μ}) (cf. page 115 of [Sch74]). But B is just the set of characteristic functions in fix (S_{μ}) . Thus there is a measurable set $A \subseteq K$ with $S_{\mu}\mathbb{1}_A = \mathbb{1}_A$ and $0 < \mu(A) < 1$. Now consider the measures μ_1, μ_2 defined by

$$\mu_1(g) := \frac{1}{\mu(A)} \int_A g \,\mathrm{d}\mu \text{ and } \mu_2(g) := \frac{1}{1 - \mu(A)} \int_{K \setminus A} g \,\mathrm{d}\mu$$

for $g \in C(K)$. For every $g \in C(K)$ with $0 \leq g \leq 1$ and each $S \in S$ we obtain

$$\int_{A} g \, \mathrm{d}\mu = \int g \wedge \mathbb{1}_{A} \, \mathrm{d}\mu = \int S_{\mu}(g \wedge \mathbb{1}_{A}) \, \mathrm{d}\mu \leqslant \int Sg \wedge \mathbb{1}_{A} \, \mathrm{d}\mu = \int_{A} Sg \, \mathrm{d}\mu$$

id, similarly

and 1,

$$\int_{A^c} g \,\mathrm{d}\mu \leqslant \int_{A^c} Sg \,\mathrm{d}\mu,$$

which implies $\mu_i \in P_{\mathcal{S}}(K)$ for i = 1, 2. Moreover,

$$\mu = \mu(A)\mu_1 + (1 - \mu(A))\mu_2,$$

so $\mu \notin \exp P_{\mathcal{S}}(K)$.

Conversely, take an ergodic measure $\mu \in P_{\delta}(K)$ and suppose that $\mu =$ $\frac{1}{2}(\mu_1 + \mu_2)$ for some $\mu_1, \mu_2 \in P_{\mathcal{S}}(K)$. Since

$$|\langle f, \mu_1 \rangle| \leqslant 2 \langle |f|, \mu \rangle \leqslant 2 ||f||_{\mathrm{L}^1(K,\mu)}$$

for each $f \in C(K)$ and C(K) is dense in $L^1(K,\mu)$, we conclude that μ_1 extends uniquely to a continuous functional $\tilde{\mu}_1 \in L^{\infty}(K,\mu) = L^1(K,\mu)'$. The semigroup S_{μ} is mean ergodic (in the sense of Definition 8.31 of [EFHN15]) on $L^{1}(K,\mu)$ (see Example 13.24 of [EFHN15]) and therefore fix(S_{μ}) separates $fix(\mathfrak{S}'_{\mu})$ by Theorem 8.33 of [EFHN15]. Since $fix(\mathfrak{S}_{\mu})$ is one-dimensional,

fix (S'_{μ}) is also one-dimensional. Consequently we obtain $\tilde{\mu}_1 = \mathbb{1} \in L^{\infty}(K, \mu)$ which implies $\mu_1 = \mu$.

We are now ready to introduce primitive S-ideals.

Definition 2.6. An S-ideal p is called *primitive* if there is an ergodic measure $\mu \in P_{\mathcal{S}}(K)$ with $p = I_{\mu}$.

The set of all primitive S-ideals is called the *primitive spectrum of* S, denoted by Prim(S).

Remark 2.7. The supports of primitive S-ideals are precisely the supports of ergodic measures. Instead of looking at the ideal space it is therefore justified (and sometimes helpful) to see the primitive spectrum as a subset of the power set of K.

Remark 2.8. If $S \subseteq \mathscr{L}(C(K))$ is irreducible, i.e., there are no non-trivial S-ideals, then Prim(S) is a singleton. Other examples are given below (cf. Examples 5.7).

We need the following result which relates invariant measures for quotient systems to invariant measures on K.

Proposition 2.9. Let $L \subseteq K$ be the support of an S-ideal and consider the semigroup S_L of Markov operators on C(L) induced by S. The canonical continuous embedding

$$i: \mathcal{C}(L)' \longrightarrow \mathcal{C}(K)'$$

with $i(\mu)(f) := \langle f|_L, \mu \rangle$ for each $f \in C(K)$ and $\mu \in C(L)'$ restricts to continuous embeddings

$$\begin{split} &i\colon \mathrm{P}_{\mathbb{S}_L}(L) \longrightarrow \mathrm{P}_{\mathbb{S}}(K), \\ &i\colon \exp \mathrm{P}_{\mathbb{S}_L}(L) \longrightarrow \exp \mathrm{P}_{\mathbb{S}}(K) \end{split}$$

with

$$i(\mathcal{P}_{\mathcal{S}_L}(L)) = \{ \tilde{\mu} \in \mathcal{P}_{\mathcal{S}}(K) \mid \operatorname{supp} \tilde{\mu} \subseteq L \} and$$

$$i(\operatorname{ex} \mathcal{P}_{\mathcal{S}_L}(L)) = \{ \tilde{\mu} \in \operatorname{ex} \mathcal{P}_{\mathcal{S}}(K) \mid \operatorname{supp} \tilde{\mu} \subseteq L \}.$$

Proof. It is obvious that images of invariant measures remain invariant. Now assume that $\mu \in \exp_{\mathcal{S}}(L)$ and suppose that $i(\mu) = \frac{1}{2}(\tilde{\mu}_1 + \tilde{\mu}_2)$ for measures $\tilde{\mu}_1, \tilde{\mu}_2 \in P_{\mathcal{S}}(K)$. Then $\operatorname{supp} \tilde{\mu}_i \subseteq \operatorname{supp} \mu$ for i = 1, 2 and therefore $\tilde{\mu}_1$ and $\tilde{\mu}_2$ restrict to measures $\mu_1, \mu_2 \in P_{\mathcal{S}_L}(L)$ with $\mu = \frac{1}{2}(\mu_1 + \mu_2)$, so $\mu_1 = \mu = \mu_2$ since μ is ergodic.

Corollary 2.10. Each maximal S-ideal is primitive.

Proof. Take a maximal S-ideal $I = I_L$. Then the induced semigroup S_I on C(L) is irreducible and consequently every S_I -invariant measure μ is strictly positive, i.e., supp $\mu = L$.

We give two simple examples showing that the converse of Corollary 2.10 does not hold.

HENRIK KREIDLER

- **Example 2.11.** (i) If $K = \mathbb{T} := \{z \in \mathbb{C} \mid |z| = 1\}$ and $\varphi(z) := z^2$ for $z \in \mathbb{T}$, then the Haar measure of \mathbb{T} is ergodic by Proposition 2.17 of [EW11]. However, $1 \in \mathbb{T}$ is a fixed point of φ and therefore $I_{\{1\}}$ is a non-trivial S_{φ} -ideal. Therefore, the zero ideal is a primitive, but not maximal S_{φ} -ideal.
 - (ii) Consider $K = \{0,1\}^{\mathbb{N}}$ and $\varphi((x_n)_{n \in \mathbb{N}}) := (x_{n+1})_{n \in \mathbb{N}}$ for $(x_n)_{n \in \mathbb{N}} \in K$. Clearly, φ has fixed points whence the zero ideal is not a maximal \mathcal{S}_{φ} -ideal. Let $\nu := \frac{1}{2}(\delta_0 + \delta_1) \in C(\{0,1\})'$. Then the product measure $\mu := \prod_{n \in \mathbb{N}} \nu \in C(K)'$ on K is ergodic by Proposition 6.20 of [EFHN15] and has full support. Therefore $I_{\mu} = \{0\}$ is a primitive \mathcal{S}_{φ} -ideal.

Remark 2.12. In view of Examples 2.11 considering all primitive ideals instead of maximal ideals yields more information on the semigroup action.

3. RADICAL IDEALS

The Jacobson topology on the primitive spectrum of C*-algebras can be defined using the notions of *hull* and *kernel* (see Section 4.3 of [Ped79] or Section II.6.5 of [Bla06]). In our context they also yield a natural correspondence between closed subsets of Prim(S) and so-called *radical S-ideals*.

Definition 3.1. For subsets $A \subseteq Prim(S)$ and $I \subseteq C(K)$ we set

$$\ker(A) := \bigcap_{p \in A} p,$$
$$\operatorname{hull}(I) := \left\{ p \in \operatorname{Prim}(\mathbb{S}) \mid I \subseteq p \right\}.$$

(i) For a subset $I \subseteq C(K)$ the S-radical of I is

$$\operatorname{rad}_{\mathbb{S}}(I) := \operatorname{ker}(\operatorname{hull}(I))) = \bigcap_{\substack{p \in \operatorname{Prim}(\mathbb{S})\\I \subseteq p}} p.$$

- (ii) An S-ideal I is a radical S-ideal if $I = \operatorname{rad}_{S}(I)$.
- (iii) The semigroup S is *radical free* if the zero ideal is a radical S-ideal, i.e., if $rad_{S}(0) = 0$.

We denote the set of all radical S-ideals by Rad(S).

Remark 3.2. We point out that our definition of a radical free semigroup does not coincide with the one of Schaefer (using maximal S-ideals, see [Sch68]). Every radical free semigroup in the sense of Schaefer is also radical free in our terminology. However, the converse does not hold (see Examples 2.11).

Remark 3.3. By the Krein-Milman theorem $P_{\mathcal{S}}(K)$ is the closed convex hull of ex $P_{\mathcal{S}}(K)$ with respect to the weak* topology and therefore

$$\operatorname{rad}_{\mathbb{S}}(0) = \bigcap_{\mu \in \mathcal{P}_{\mathbb{S}}(K)} I_{\mu}.$$

Remark 3.4. The S-radical of a subset $I \subseteq C(K)$ is either C(K) or a radical S-ideal. Moreover, we always have $hull(I) = hull(rad_{S}(I))$.

Remark 3.5. Just as primitive ideals correspond to the supports of ergodic measures, radical ideals correspond to the closures of unions of supports of ergodic measures. Therefore S is radical free if and only if the union of all supports of invariant ergodic measures is dense in K. Note that the latter set is not closed in general (see Example 5.7 (iii) below). For the Markov semigroup induced by the right shift on $K = \beta \mathbb{N} \setminus \mathbb{N}$ this set is nowhere dense (cf. Corollary 1.5 in [Cho67]).

We need the following result which relates radical and primitive ideals of quotient systems to the corresponding S-ideals of C(K).

Proposition 3.6. Let $I = I_L \subseteq C(K)$ be an S-ideal and S_I the semigroup of Markov operators on C(L) induced by S. Then the mappings

$$\{p \in \operatorname{Prim}(\mathbb{S}) \mid I \subseteq p\} \longrightarrow \operatorname{Prim}(\mathbb{S}_I), \quad p \mapsto p|_L, \\ \{J \in \operatorname{Rad}(\mathbb{S}) \mid I \subseteq J\} \longrightarrow \operatorname{Rad}(\mathbb{S}_I), \quad J \mapsto J|_L, \end{cases}$$

where $J|_L := \{f|_L \mid f \in J\}$ for $J \subseteq C(K)$, are inclusion preserving bijections with inclusion preserving inverses. Moreover, $\operatorname{rad}_{S_I}(0) = \operatorname{rad}_S(I)|_L$.

Proof. We first recall that the natural projection $P: C(K) \longrightarrow C(L)$ is a surjetive Banach lattice homomorphism. Thus, if $\tilde{J} \subseteq C(L)$ is a closed ideal, then $J := P^{-1}(\tilde{J})$ is a closed ideal containing I with $\tilde{J} = P(P^{-1}(\tilde{J})) = J|_L$. It is readily checked that J is the unique closed ideal H containing I with $H|_L = \tilde{J}$. Clearly \tilde{J} is S_I -invariant if and only if J is S-invariant.

We therefore obtain mutually inverse and inclusion preserving mappings

$$\{J \subseteq \mathcal{C}(K) \mid J \text{ S-ideal with } I \subseteq J\} \leftrightarrow \{J \subseteq \mathcal{C}(L) \mid J \text{ S}_I\text{-ideal}\}, \\ J \mapsto J|_L \\ P^{-1}(\tilde{J}) \leftrightarrow \tilde{J}.$$

We now prove that

$$\{p \in \operatorname{Prim}(\mathfrak{S}) \mid I \subseteq p\} \longrightarrow \operatorname{Prim}(\mathfrak{S}_I), \quad p \mapsto p|_L$$

is a bijective map. Assume that $I \subseteq J = I_{\mu}$ for some $\mu \in \exp_{\mathbb{S}}(K)$. Then supp $\mu \subseteq L$ and we thus find $\nu \in \exp_{\mathbb{S}_{I}}(L)$ with $i(\nu) = \mu$ (see Proposition 2.9). Moreover we obtain for every $f \in C(L)$ that

(1)
$$\langle |f|, \nu \rangle = \int_L |F| \, \mathrm{d}\mu,$$

for each extension $F \in C(K)$ of f to K. Thus $f \in I_{\nu}$ if and only if $f \in I_{\mu}|_{L}$. If, on the other hand, $\tilde{J} = I_{\nu}$ for some $\nu \in \exp_{\mathfrak{S}_{I}}(L)$, then Equation (1) holds for $\mu = i(\nu)$ and thus $\tilde{J} = I_{\mu}|_{L}$.

Before proceeding with the remaining assertions, we make the following two observations.

• For a family $(J_{\alpha})_{\alpha \in A}$ of S-ideals with $I \subseteq J_{\alpha}$ for every $\alpha \in A$

$$\left(\bigcap_{\alpha\in A} J_{\alpha}\right)|_{L} = \bigcap_{\alpha\in A} (J_{\alpha}|_{L}).$$

• For two S-ideals J_1, J_2 with $I \subseteq J_1, J_2$ the inclusion $J_1|_L \subseteq J_2|_L$ implies $J_1 \subseteq J_2$.

We use these facts to show that

$$\{J \in \operatorname{Rad}(\mathfrak{S}) \mid I \subseteq J\} \longrightarrow \operatorname{Rad}(\mathfrak{S}_I), \quad J \mapsto J|_L$$

is a well-defined bijection. Take an S-ideal $J \subseteq C(K)$ with $I \subseteq J$. Then J is radical if and only if

$$J = \bigcap_{\substack{p \in \operatorname{Prim}(\mathbb{S})\\J \subseteq p}} p$$

which is—by the observations above—equivalent to

$$J|_{L} = \bigcap_{\substack{p \in \operatorname{Prim}(\mathbb{S})\\ J \subseteq p}} p|_{L} = \bigcap_{\substack{p \in \operatorname{Prim}(\mathbb{S})\\ J|_{L} \subseteq p|_{L}}} p|_{L} = \bigcap_{\substack{p \in \operatorname{Prim}(\mathbb{S}_{I})\\ J|_{L} \subseteq p}} p,$$

i.e., $J|_L$ being a radical S_I -ideal.

Finally, the identity $\operatorname{rad}_{S_I}(0) = \operatorname{rad}_{S}(I)|_L$ follows from the fact that $\operatorname{rad}_{S_I}(0)$ is the smallest radical S_I -ideal and $\operatorname{rad}_{S}(I)$ is the smallest radical S-ideal containing I.

Our main class of examples for radical ideals are the absolute kernels of (possibly non-ergodic) invariant measures. The following result generalizes Proposition 12 of [Sch68] using similar arguments.

Proposition 3.7. The following assertions are valid.

- (i) For each $\mu \in P_{\mathcal{S}}(K)$ the S-ideal I_{μ} is a radical S-ideal.
- (ii) If K is metrizable, then for each radical S-ideal I there is $\mu \in P_{S}(K)$ with $I = I_{\mu}$.

Proof. For (i) let $\mu \in P_{\mathcal{S}}(K)$. By Lemma 3.6 we may assume that $K = \operatorname{supp} \mu$ and it then suffices to show that \mathcal{S} is radical free. But this directly follows from Remark 3.3.

We now prove (ii) and assume that K is metrizable and I is a radical S-ideal. We may assume that I = 0 (otherwise we pass to C(supp I), cf. Lemma 3.6). Take a countable base of the topology consisting of nonempty open sets U_n , $n \in \mathbb{N}$. Since the supports of ergodic measures are dense in K we find $\mu_n \in \exp(K)$ with $\operatorname{supp} \mu_n \cap U_n \neq \emptyset$ for each $n \in \mathbb{N}$. For

$$\mu := \sum_{n=1}^{\infty} 2^{-n} \mu_n \in \mathcal{P}_{\mathcal{S}}(K)$$

we obtain $\mu(U_n) > 0$ for each $n \in \mathbb{N}$, hence $\mu(U) > 0$ for each non-empty open set $U \subseteq K$.

Remark 3.8. Taking $S = {Id}$ in Proposition 3.7 (ii) yields the probably wellknown fact that every compact metric space has a fully supported regular Borel probability measure.¹

The following examples show that part (ii) of Proposition 3.7 is wrong in the non-metric case.

- **Example 3.9.** (i) If $K = \Omega \cup \{\infty\}$ is the one-point compactification of an uncountable discrete space Ω , then C(K)' can be identified with $\ell^1(K)$. Thus there is no fully supported probability measure $\mu \in C(K)'$.¹
 - (ii) If $K = \beta \mathbb{N} \setminus \mathbb{N}$ and S_{φ} is the Markov semigroup induced by the right shift φ , then

$$\bigcap_{n\in\mathbb{N}}I_{\mu_n}\not\subseteq\operatorname{rad}_{\mathcal{S}_{\varphi}}(0)$$

for every sequence of probability measures $(\mu_n)_{n \in \mathbb{N}} \subseteq C(K)'$ (see Corollary 1.10 of [Cho67]).

4. Centers of Attraction

Radical S-ideals can also be described via an ergodic stability condition. To formulate our theorem we write $\overline{\operatorname{co}} S$ for the closed convex hull of S with respect to the strong operator topology and recall that a net $(T_{\alpha})_{\alpha \in A} \subseteq \overline{\operatorname{co}} S \subseteq \mathscr{L}(\mathcal{C}(K))$ of operators is *right ergodic* if

$$\lim_{\alpha} T_{\alpha}(\mathrm{Id} - S) = 0$$

for each $S \in S$ with respect to the strong operator topology. We note that there always are right ergodic operator nets for S (see Corollary 1.5 of [Sch13]). We give some examples (see Examples 1.2 of [Sch13]).

Example 4.1. (i) If $S = \{S^n \mid n \in \mathbb{N}_0\}$ for some Markov operator $S \in \mathscr{L}(\mathbb{C}(K))$, then the *Cesàro means*

$$C_N := \frac{1}{N} \sum_{n=0}^{N-1} S^n \text{ for } N \in \mathbb{N}$$

define a right ergodic operator sequence $(C_N)_{N \in \mathbb{N}}$ for S. Likewise, the net of Abel means $(A_r)_{r \in (0,1)}$ defined by

$$A_r := (1-r) \sum_{n=0}^{\infty} (rS)^n$$
 for $r \in (0,1)$

is right ergodic for S.

(ii) If $S = \{S(t) \mid t \ge 0\}$ is a strongly continuous one-parameter semigroup of Markov operators on C(K), then the *Cesàro means*

$$C_T f := \frac{1}{T} \int_0^T S(t) f \, \mathrm{d}t \text{ for } f \in \mathcal{C}(K) \text{ and } T > 0$$

define a right ergodic operator net $(C_T)_{T>0}$ for S.

¹Remark 3.8 and Example 3.9 (i) were kindly suggested by the referee.

The following result generalizes Theorem 4 of [Sch68].

Theorem 4.2. For each support $L \subseteq K$ of an S-ideal

$$\operatorname{rad}_{\mathbb{S}}(I_L) = \left\{ f \in \mathcal{C}(K) \left| \lim_{\alpha} \int_L T_{\alpha} |f| \, \mathrm{d}\mu = 0 \text{ for each } \mu \in \mathcal{C}(L)' \right\} \right.$$
$$= \left\{ f \in \mathcal{C}(K) \left| \lim_{\alpha} (T_{\alpha} |f|) \right|_L = 0 \text{ in the norm of } \mathcal{C}(L) \right\}$$

where $(T_{\alpha})_{\alpha \in A}$ is any right ergodic operator net for S. In particular, if $(T_{\alpha})_{\alpha \in A}$ is any right ergodic operator net for S, then an S-ideal I_L is a radical S-ideal if and only if every $f \in C(K)$ satisfying

$$\lim_{\alpha} (T_{\alpha}|f|)|_{L} = 0$$

vanishes on L.

Proof. By Lemma 3.6 we may assume L = K. Take $f \in \operatorname{rad}_{\mathbb{S}}(0)$ and any right ergodic operator net $(T_{\alpha})_{\alpha \in A}$ for S.

Let $\mu \in C(K)'$ and observe that each subnet of $(T'_{\alpha}\mu)_{\alpha\in A}$ has a subnet converging to some $\nu \in P_{\mathcal{S}}(K)$. Since $\langle |f|, \nu \rangle = 0$ (see Remark 3.3), we obtain that each subnet of $(\langle T_{\alpha}|f|, \mu \rangle)_{\alpha\in A}$ has a subnet converging to zero which implies

$$\lim_{\alpha} \langle T_{\alpha} | f |, \mu \rangle = 0.$$

Now let $f \in C(K)$ with $\lim_{\alpha} T_{\alpha}|f| = 0$ weakly for some right ergodic operator net $(T_{\alpha})_{\alpha \in A}$ for S. Then for $\mu \in P_{S}(K)$

$$0 = \lim_{\alpha} \langle T_{\alpha} | f |, \mu \rangle = \langle | f |, \mu \rangle$$

which proves $f \in \operatorname{rad}_{\mathbb{S}}(0)$ and thus the first equation. By Theorem 1.7 of [Sch13] the semigroup \mathbb{S} is mean ergodic on $\operatorname{rad}_{\mathbb{S}}(0)$ with mean ergodic projection P = 0 and therefore

$$\operatorname{rad}_{\mathfrak{S}}(0) \subseteq \left\{ f \in \operatorname{C}(K) \, \middle| \, \lim_{\alpha} T_{\alpha} |f| = 0 \text{ in the norm of } \operatorname{C}(K) \right\}.$$

The converse inclusion is obvious.

If S has a right ergodic operator sequence (for example if it has a Følner sequence as defined in Assumption 4.5 below), then Lebesgue's Theorem yields the following result.

Corollary 4.3. Suppose that $(T_n)_{n \in \mathbb{N}}$ is a right ergodic operator sequence for S. Then

$$\operatorname{rad}_{\mathbb{S}}(I) = \left\{ f \in \mathcal{C}(K) \, \middle| \, \lim_{n \to \infty} T_n |f|(x) = 0 \text{ for each } x \in \operatorname{supp} I \right\}$$

for each S-ideal I.

The next corollary shows that if S is the semigroup generated by a Markov lattice homomorphism $T \in \mathscr{L}(C(K))$ (i.e., a Koopman operator), the radical rad_S(0) of the zero ideal coincides with the almost weakly stable part of C(K) with respect to T as defined in (9.4) on page 176 of [EFHN15].

Corollary 4.4. Assume that $\varphi \colon K \longrightarrow K$ is a continuous mapping and $S = S_{\varphi}$. Then

$$\operatorname{rad}_{\mathbb{S}}(0) = \left\{ f \in \mathcal{C}(K) \left| \lim_{N \to \infty} \frac{1}{N} \sum_{n=0}^{N-1} |\langle T_{\varphi}^{n} f, \mu \rangle| = 0 \text{ for each } \mu \in \mathcal{C}(K)' \right\}.\right.$$

Proof. If $f \in \operatorname{rad}_{\mathbb{S}}(0)$ and $\mu \in C(K)'$, we obtain

$$\frac{1}{N}\sum_{n=0}^{N-1}|\langle T_{\varphi}^{n}f,\mu\rangle|\leqslant \frac{1}{N}\sum_{n=0}^{N-1}\langle T_{\varphi}^{n}|f|,|\mu|\rangle$$

for every $N \in \mathbb{N}$ and therefore $\lim_{N \to \infty} \frac{1}{N} \sum_{n=0}^{N-1} |\langle T_{\varphi}^n f, \mu \rangle| = 0$ by Theorem 4.2. The converse inclusion follows directly from Corollary 4.3.

For Koopman semigroups S we also obtain a further dynamical characterization of $\operatorname{rad}_{S}(0)$. For the rest of this section we make the following assumption (cf. Examples 1.2 (e) of [Sch13]).

Assumption 4.5. Let S be a closed subsemigroup of a locally compact group G with left-invariant Haar measure λ acting on K such that

$$\mathcal{S} \times K \longrightarrow K, \quad (s, x) \mapsto sx$$

is continuous. Let S be the associated Koopman semigroup, i.e.,

$$\mathbb{S} = \{T_s \mid s \in \mathcal{S}\}$$

with $T_s f(x) := f(sx)$ for $f \in C(K)$, $s \in S$ and $x \in K$, which is strongly continuous by Theorem 4.17 of [EFHN15]. Moreover, we assume that $(F_n)_{n \in \mathbb{N}}$ is a *Følner sequence* for S, i.e., each F_n is a compact subset of S with positive measure satisfying

$$\lim_{n \to \infty} \frac{\lambda(F_n \Delta s F_n)}{\lambda(F_n)} = 0$$

for each $s \in \mathcal{S}$.

- **Example 4.6.** (i) If S is the additive semigroup \mathbb{N}_0 , then the sequence $(F_n)_{n\in\mathbb{N}}$ defined by $F_n := \{0, ..., n-1\}$ for $n \in \mathbb{N}$ is a Følner sequence for S.
 - (ii) If S is the additive semigroup $\mathbb{R}_{\geq 0}$ and $(t_n)_{n \in \mathbb{N}}$ is any sequence in $(0, \infty)$ with $\lim_{n \to \infty} t_n = \infty$ then $(F_n)_{n \in \mathbb{N}}$ defined by $F_n := [0, t_n]$ for $n \in \mathbb{N}$ is a Følner sequence for S.

Lemma 4.7. Under Assumption 4.5 S is left amenable and thus S is right amenable. Moreover we obtain an ergodic operator sequence $(\mathcal{F}_n)_{n \in \mathbb{N}}$ for S by setting

$$\mathfrak{F}_n f := \frac{1}{\lambda(F_n)} \int_{F_n} T_s f \, \mathrm{d}\lambda(s)$$

for $f \in C(K)$ and $n \in \mathbb{N}$.

Proof. For each $n \in \mathbb{N}$ set

$$m_n(f) := \frac{1}{\lambda(F_n)} \int_{F_n} f(s) \, \mathrm{d}\lambda(s)$$

for $f \in C_b(\mathcal{S})$. Then $m_n \in C_b(\mathcal{S})'$ with $m_n(\mathbb{1}) = 1$ and $m_n \ge 0$ for each $n \in \mathbb{N}$. Let m be any weak^{*} limit point of $(m_n)_{n \in \mathbb{N}}$. Since

$$\left|\frac{1}{\lambda(F_n)}\int_{F_n} f(s) \,\mathrm{d}\lambda(s) - \frac{1}{\lambda(F_n)}\int_{F_n} f(ts) \,\mathrm{d}\lambda(s)\right| \leq \frac{\lambda(F_n\Delta tF_n)}{\lambda(F_n)} \cdot \|f\|$$

for each $f \in C_b(S)$, $n \in \mathbb{N}$ and $t \in S$, *m* is an invariant mean. The second assertion is obvious.

We now introduce certain "attractors" of the dynamical system (K; S) with respect to the Følner sequence $(F_n)_{n \in \mathbb{N}}$.

Definition 4.8. A closed non-empty set $L \subseteq K$ is a *(global) center of attraction* if for each open set $U \supseteq L$ we have

$$\lim_{n \to \infty} \frac{1}{\lambda(F_n)} \lambda(\{s \in F_n \mid sx \in U\}) = 1$$

for every $x \in K$.

This type of attraction is quite weak. Loosely speaking, orbits of points may move arbitarily far away from a center of attraction as long as they come back "often enough" with respect to the Følner sequence.

Global as well as point-dependent centers of attraction for \mathbb{N}_{0} - and $\mathbb{R}_{\geq 0}$ actions have been examined by several authors (see, e.g., [Hil36], [Ber51], [JR72], [Sig77], Exercise I.8.3 in [Man87] and [Dai16]). In a recent paper Z. Chen and X. Dai study the chaotic behavior of minimal centers of attraction with respect to a point for discrete amenable group actions (see [CD17]).

It is known that in case of \mathbb{N}_0 -actions on metric compact spaces there always is a unique minimal (global) center of attraction given by the closure of the union of the supports of ergodic measures (see Exercises I.8.3 and II.1.5 in [Man87]). The following result shows that this still holds in our more general situation.

Theorem 4.9. Under Assumption 4.5 the definition of a center of attraction does not depend on the Følner sequence. Moreover, for a closed non-empty set $L \subseteq K$ the following assertions are equivalent.

- (a) L is a center of attraction.
- (b) $I_L := \{ f \in C(K) \mid f \mid_L = 0 \} \subseteq rad_{\mathcal{S}}(0).$

In particular there is a unique minimal center of attraction M(S) given by the closure of the union of the supports of ergodic measures, i.e.,

 $M(S) = \operatorname{supp} \operatorname{rad}_{S}(0).$

Proof. Take a non-empty and closed set $L \subseteq K$. The mapping

$$I_L \longrightarrow \mathcal{C}_0(K \setminus L), \quad f \mapsto f|_{K \setminus L}$$

is an isomorphism of Banach lattices. Now L is a center of attraction if and only if

$$\lim_{n \to \infty} \frac{1}{\lambda(F_n)} \lambda(\{s \in F_n \mid sx \in A\}) = 0,$$

i.e.,

(2)
$$\lim_{n \to \infty} \frac{1}{\lambda(F_n)} \int_{F_n} \mathbb{1}_A(sx) \, \mathrm{d}\lambda(s) = 0$$

for each compact set $A \subseteq K \setminus L$ and each $x \in K$. Since the mapping

 $\mathcal{S} \times K \longrightarrow K, \quad (s, x) \mapsto sx$

is continuous, the function $f: \mathcal{S} \times K \longrightarrow \mathbb{C}, (s, x) \mapsto \mathbb{1}_A(sx)$ is Borel measurable. By Lebesgue's and Fubini's theorems Equation (2) is thus equivalent to

$$\lim_{n \to \infty} \int_K \int_{F_n} \mathbb{1}_A(sx) \, \mathrm{d}\lambda(s) \, \mathrm{d}\mu(x) = \lim_{n \to \infty} \frac{1}{\lambda(F_n)} \int_{F_n} T'_s \mu(A) \, \mathrm{d}\lambda(s) = 0$$

for each $\mu \in C_0(K \setminus L)'$ and each compact set $A \subseteq K \setminus L$. Since the space of compactly supported continuous functions $C_c(K \setminus L)$ is dense in $C_0(K \setminus L)$, this is the case if and only if

$$\lim_{n \to \infty} \frac{1}{\lambda(F_n)} \int_{F_n} \langle T_s | f |, \mu \rangle \mathrm{d}\lambda(s) = 0$$

for each $f \in I_L$ and every $\mu \in I'_L$. This means

$$\lim_{n \to \infty} \mathcal{F}_n |f| = 0$$

with respect to the weak topology for each $f \in I_L$, i.e., $I_L \subseteq \operatorname{rad}_{\mathbb{S}}(0)$. \Box

5. The Primitive Spectrum as a Topological Space

In this section we return to a general right amenable Markov semigroup $S \subseteq \mathscr{L}(C(K))$ and analyze the topology of Prim(S). It turns out that it basically has the same properties as the (non-dynamical) primitive spectrum of C^{*}algebras and the topology of affine schemes of algebraic geometry (cf. Section (2.2) of [GW10]). We employ methods as in Chapter IV of [Dix77] and Section 4.3 of [Ped79] and first prove two technical lemmas before introducing a topology on Prim(S). Recall that given $\mu \in P_S(K)$ we write S_{μ} for the induced semigroup on $L^1(K, \mu)$.

Lemma 5.1. If $\mu \in P_{\mathcal{S}}(K)$ and $L \subseteq K$ is the support of an S-ideal, then $\mathbb{1}_L \in fix(\mathfrak{S}_{\mu})$.

Proof. We fix $S \in S$. For each open set $O \supseteq L$ take a continuous function f_O with $f_O(K) \subseteq [0, 1]$, $f|_L = \mathbb{1}$ and $f_O|_{(K \setminus O)} = 0$. The set Λ of open sets containing L is directed with respect to converse set inclusion and thus we obtain a net $(f_O)_{O \in \Lambda}$ with

 $\|\mathbb{1}_L - f_O\|_{\mathrm{L}^1(K,\mu)} \leqslant \mu(O \setminus L) \to 0$

by regularity of μ . By definition of S_{μ}

$$S_{\mu}\mathbb{1}_{L} = \lim_{O} Sf_{O}.$$

in $L^1(K,\mu)$. Moreover,

$$Sf_O(x) = \langle Sf_O, \delta_x \rangle = \langle f_O, S'\delta_x \rangle = 1$$

for each $x \in L$ since L is the support of an S-ideal. This implies

$$0 = \lim_{O} ((\mathbb{1} - Sf_O) \cdot \mathbb{1}_L) = \mathbb{1}_L - S_\mu \mathbb{1}_L \cdot \mathbb{1}_L,$$

where the limit is taken in $L^1(K, \mu)$. Thus $\mathbb{1}_L = S_\mu \mathbb{1}_L \cdot \mathbb{1}_L$ which shows $\mathbb{1}_L \leq S_\mu \mathbb{1}_L$ and consequently $\mathbb{1}_L \in \text{fix}(\mathbb{S}_\mu)$ by Theorem 13.2 (d) of [EFHN15]. \Box

Lemma 5.2. Consider two S-ideals I_1, I_2 . If p is a primitive S-ideal with $I_1 \cap I_2 \subseteq p$, then $I_1 \subseteq p$ or $I_2 \subseteq p$.

Proof. Let $p = I_{\mu}$ for some $\mu \in \exp P_{\mathbb{S}}(K)$ and let $L_j := \operatorname{supp} I_j$ for j = 1, 2. By Lemma 5.1, $\mathbb{1}_{L_j} \in \operatorname{fix}(\mathbb{S}_{\mu})$ for j = 1, 2 and therefore $\mu(L_j) \in \{0, 1\}$ since μ is ergodic. Now $\operatorname{supp} \mu \subseteq L_1 \cup L_2$ implies $\mu(L_1 \cup L_2) = 1$, so there is $j \in \{1, 2\}$ with $\mu(L_j) = 1$. But this means $\operatorname{supp} \mu \subseteq L_j$ and consequently $I_j \subseteq p$. \Box

We are now ready to equip Prim(S) with a topology by defining a Kuratowski closure operator (see page 43 of [Kel75] for this notion). Recall that the concepts of *hull* and *kernel* have been introduced in Definition 3.1.

Proposition 5.3. The mapping

$$\overline{}: \mathscr{P}(\operatorname{Prim}(\mathbb{S})) \longrightarrow \mathscr{P}(\operatorname{Prim}(\mathbb{S})), \quad A \mapsto \overline{A} := \operatorname{hull}(\ker(A))$$

defines a Kuratowski closure operator.

Proof. It is readily checked that

$$\overline{\emptyset} = \emptyset, A \subseteq \overline{A} \text{ and } \overline{A} = \overline{A}$$

for each $A \subseteq \text{Prim}(S)$. It remains to show that $\overline{A_1 \cup A_2} = \overline{A_1} \cup \overline{A_2}$ for all $A_1, A_2 \subseteq \text{Prim}(S)$. Applying Lemma 5.2 to the ideals $I_j := \text{ker}(A_j)$ for j = 1, 2 yields

$$A_1 \cup A_2 = \operatorname{hull}(\operatorname{ker}(A_1 \cup A_2)) = \operatorname{hull}(\operatorname{ker}(A_1) \cap \operatorname{ker}(A_2))$$
$$= \{p \in \operatorname{Prim}(\mathbb{S}) \mid \operatorname{ker}(A_1) \cap \operatorname{ker}(A_2) \subseteq p\}$$
$$= \{p \in \operatorname{Prim}(\mathbb{S}) \mid \operatorname{ker}(A_1) \subseteq p \text{ or } \operatorname{ker}(A_2) \subseteq p\} = \overline{A_1} \cup \overline{A_2},$$

which shows the claim.

Definition 5.4. The topology on Prim(S) induced by the closure operator of Proposition 5.3 is called the *hull-kernel-topology*.

We from now on equip Prim(S) with the hull-kernel-topology.

Proposition 5.5. The following assertions are valid.

(i) The mappings

$$\{ \emptyset \neq A \subseteq \operatorname{Prim}(\mathbb{S}) \operatorname{closed} \} \leftrightarrow \operatorname{Rad}(\mathbb{S}) \\ A \mapsto \ker(A) \\ \operatorname{hull}(I) \leftarrow I$$

are mutually inverse bijections.

(ii) The sets

$$U_f := \{ p \in \operatorname{Prim}(\mathcal{S}) \mid f \notin p \}$$

for $f \in C(K)$ define a base for the hull-kernel-topology of Prim(S).

- (iii) If K is metrizable, then Prim(S) has a countable base.
- (iv) The space Prim(S) is T_0 . Given $p \in Prim(S)$, the set $\{p\}$ is closed if and only if p is a maximal S-ideal.
- (v) The space Prim(S) is quasi-compact.
- (vi) The mapping

$$\pi : \exp \mathcal{P}_{\mathcal{S}}(K) \longrightarrow \operatorname{Prim}(\mathcal{S}), \quad \mu \mapsto I_{\mu}$$

is continuous and surjective.

Proof. Assertion (i) is obvious. For (ii) observe that $Prim(S) = U_{\mathbb{1}}$. Now take a closed set $\emptyset \neq A \subseteq Prim(S)$. Then $A = \ker(I)$ for some S-ideal I and we obtain

$$\operatorname{Prim}(\mathfrak{S}) \setminus A = \bigcup_{f \in I} \{ p \in \operatorname{Prim}(\mathfrak{S}) \mid f \notin p \} = \bigcup_{f \in I} U_f.$$

Moreover, each U_f is open since $Prim(\mathfrak{S}) \setminus U_f = hull(\{f\})$. This proves (ii). Assertion (iii) is a direct consequence of (ii).

We proceed with part (iv) and prove that Prim(S) is a T₀-space. If $p_1, p_2 \in Prim(S)$ with $p_1 \neq p_2$, then $M_1 \neq M_2$ for the supports $M_i := \operatorname{supp} p_i$, i = 1, 2. We may assume that there is $x \in M_2 \setminus M_1$ and find $f \in C(K)$ with $f|_{M_1} = 0$ and f(x) = 1. Then $p_1 \notin U_f$ and $p_2 \in U_f$.

For the second part of (iv) take a maximal S-ideal and assume that $p \in \overline{\{m\}}$. Then $m \subseteq p$ and thus m = p by maximality of m.

Conversely, suppose that $\{m\}$ is closed and take a maximal S-ideal p with $m \subseteq p$. Then ker $(\{m\}) = m \subseteq p$ and thus

$$p \in \operatorname{hull}(\ker(\{m\})) = \overline{\{m\}},$$

i.e., p = m.

For the proof of (v) take closed subsets $A_j \subseteq Prim(S)$ for $j \in J$ with

$$\bigcap_{j\in J} A_j = \emptyset$$

and let $I_j := \ker(A_j)$ be the corresponding radical ideals for $j \in J$. We show that

$$\sum_{j \in J} I_j = \mathcal{C}(K).$$

Denote the ideal on the left side by I and assume that it is a proper invariant ideal. Since there are no dense ideals in C(K), the closure \overline{I} is contained in a maximal S-ideal p. But then $p \in A_j$ for each $j \in J$ since the sets A_j are closed, a contradiction.

Take $j_1, ..., j_k$ with $1 \in I_{j_1} + ... + I_{j_k}$ for some $k \in \mathbb{N}$. Then

$$\sum_{m=1}^{k} I_{j_m} = \mathcal{C}(K)$$

and consequently

$$\bigcap_{m=1}^{k} A_{j_m} = \emptyset.$$

Finally, assertion (vi) follows from the fact that the set

$$\pi^{-1}(U_f) = \{ \mu \in \exp \mathsf{P}_{\mathcal{S}}(K) \mid \langle |f|, \mu \rangle \neq 0 \}$$

is open in $\exp P_{\mathcal{S}}(K)$ for each $f \in C(K)$.

Corollary 5.6. A net $(p_{\alpha})_{\alpha \in A}$ in Prim(S) converges to $p \in Prim(S)$ if and only if for each open set $U \subseteq K$ with $\operatorname{supp} p \cap U \neq \emptyset$ there is $\alpha_0 \in A$ with $\operatorname{supp} p_{\alpha} \cap U \neq \emptyset$ for every $\alpha \ge \alpha_0$.

Proof. Consider the sets $V_f := \{x \in K \mid f(x) \neq 0\}$ for $f \in C(K)$. Proposition 5.5 (ii) shows that a net $(p_\alpha)_{\alpha \in A}$ in Prim(S) converges to $p \in Prim(S)$ if and only if for each $f \in C(K)$ with $f|_{\operatorname{supp}(p_\alpha)} \neq 0$ there is $\alpha_0 \in A$ with $f|_{\operatorname{supp}(p_\alpha)} \neq 0$ for every $\alpha \ge \alpha_0$, i.e., for each $f \in C(K)$ with $V_f \cap \operatorname{supp} p \neq \emptyset$ there is α_0 with $V_f \cap \operatorname{supp} p_\alpha \neq \emptyset$ for every $\alpha \ge \alpha_0$. Since the sets V_f are a base of the topology of K, this shows the claim. \Box

- **Example 5.7.** (i) For the trivial semigroup $S = {\text{Id}}$ every ideal is invariant and Prim(S) coincides with the maximal ideal space of the commutative C*-algebra C(K), i.e., it is homeomorphic to K.
 - (ii) Consider the torus $K = \mathbb{T} = \{z \in \mathbb{C} \mid |z| = 1\}$ and the rotation $\varphi_a(z) := az$ for $z \in \mathbb{T}$ and some fixed $a \in \mathbb{T}$ with $a^k = 1$ for some $k \in \mathbb{N}$. Denote the group of kth roots of unity by G_k . The ergodic measures are then precisely the measures $\mu_b \in P_{\mathcal{S}_{\varphi}}(K)$ with $\mu_b := \frac{1}{k} \sum_{j=0}^{k-1} \delta_{a^j b}$ for $b \in \mathbb{T}$. Their supports are clearly the sets

$$M_b := bG_k = \{ bz \mid z \in \mathbb{T}, \, z^k = 1 \}$$

for $b \in \mathbb{T}$. Using Corollary 5.6 a moment's thought reveals that

$$\mathbb{T}/G_k \longrightarrow \operatorname{Prim}(\mathfrak{S}_{\varphi}), \quad bG_k \mapsto I_{M_b}$$

is a homeomorphism if we endow the factor group \mathbb{T}/G_k with the quotient topology.

(iii) Consider the space $L := \{0, 1\}^{\mathbb{N}}$ and the shift $\varphi : L \longrightarrow L$ given by $\varphi((x_n)_{n \in \mathbb{N}}) := (x_{n+1})_{n \in \mathbb{N}}$ for each $(x_n)_{n \in \mathbb{N}} \in L$. For each $k \in \mathbb{N}$ consider the minimal non-empty closed invariant set

$$M_k := \{\varphi^n(x^k) \mid n \in \{0, ..., 2k - 1\}\}$$

with $x^k = (x_m^k)_{m \in \mathbb{N}}$ defined by

$$x_m^k := \begin{cases} 0 & \text{if } m \in \{1, \dots, k\} + 2k\mathbb{N}_0 \\ 1 & \text{else.} \end{cases}$$

16

Now if

$$K := \overline{\bigcup_{k \in \mathbb{N}} M_k},$$

then it is readily seen that K is the invariant set

 $\bigcup_{k \in \mathbb{N}} M_k \cup \{ (x_m)_{m \in \mathbb{N}} \in L \mid (x_m)_{m \in \mathbb{N}} \text{ increasing or decreasing} \}.$

We restrict φ to K and claim that $\operatorname{Prim}(\mathbb{S}_{\varphi})$ is not Hausdorff. It suffices to show that the sequence $(m_n)_{n \in \mathbb{N}}$ in $\operatorname{Prim}(\mathbb{S}_{\varphi})$ with $m_n := I_{M_n}$ converges to two different points.

To this end, consider $k \in \mathbb{N}$ and the open subset

$$U := \left(\prod_{i=1}^{k} \{1\} \times \prod_{i=k+1}^{\infty} \{0,1\}\right) \cap K.$$

Then $M_l \cap U \neq \emptyset$ for each $l \ge k$. By Remark 5.6 this implies $m_l \to I_{\{(1)_{n \in \mathbb{N}}\}}$ and a similar argument shows $m_l \to I_{\{(0)_{n \in \mathbb{N}}\}}$.

6. CONTINUOUS FUNCTIONS ON THE PRIMITIVE SPECTRUM

It is our goal to describe the continuous functions on Prim(S). As above we write M(S) for the support of $rad_S(0)$, i.e., the closure of the union of all supports of invariant ergodic measures, and recall that the semigroup on C(M(S)) induced by S is denoted by $S_{rad_S(0)}$. Now consider the following functions.

Definition 6.1. For a function $f \in fix(S_{rad_{S}(0)})$ we define

$$\hat{f} \colon \operatorname{Prim}(\mathfrak{S}) \longrightarrow \mathbb{C}, \quad I_{\mu} \mapsto \int_{\mathcal{M}(\mathfrak{S})} f \, \mathrm{d}\mu.$$

Note that each $f \in \text{fix}(S_{\text{rad}_{\mathbb{S}}(0)})$ is constant on supports of ergodic measures and therefore $\int_{\mathcal{M}(\mathbb{S})} f \, d\mu$ only depends on I_{μ} and not on μ itself. Thus \hat{f} is in fact well-defined for each $f \in \text{fix}(S_{\text{rad}_{\mathbb{S}}(0)})$ and the next lemma shows that it is even continuous.

Lemma 6.2. If $f \in \text{fix}(\mathcal{S}_{\text{rad}_{\mathcal{S}}(0)})$, then $\hat{f} \in C(\text{Prim}(\mathcal{S}))$.

Proof. Let $p = I_{\mu} \in \text{Prim}(S)$ and $\varepsilon > 0$. We set

$$f_{\varepsilon} := \sup\left(\varepsilon \cdot \mathbb{1} - \left| f - \int_{\mathcal{M}(\mathcal{S})} f \, \mathrm{d}\mu \cdot \mathbb{1} \right|, 0 \right) \Big|_{\mathcal{M}(\mathcal{S})} \in \mathcal{C}(\mathcal{M}(\mathcal{S})).$$

Then $U := U_{f_{\varepsilon}}$ is an open neighborhood of p. Moreover, for each $q = I_{\nu} \in U$

$$\varepsilon - \left| \int_{\mathcal{M}(\mathcal{S})} f \, \mathrm{d}\nu - \int_{\mathcal{M}(\mathcal{S})} f \, \mathrm{d}\mu \right| > 0$$

which means $|\hat{f}(p) - \hat{f}(q)| < \varepsilon$.

In the radical free case (i.e., M(S) = K) we now obtain a linear mapping from the fixed space fix(S) to C(Prim(S)). It turns out that this is actually an isomorphism.

Theorem 6.3. If S is radical free, then the fixed space fix(S) is a Banach sublattice of C(K) and the mapping

$$\hat{}: \operatorname{fix}(\mathfrak{S}) \longrightarrow \operatorname{C}(\operatorname{Prim}(\mathfrak{S})), \quad f \mapsto \hat{f}$$

is an isometric Markov lattice isomorphism.

Proof. We first show that fix(S) is a sublattice of C(K). Take $f \in fix(S)$ and $S \in S$. Since M(S) = K, it suffices to prove that $S_p|f||_{supp p} = |f||_{supp p}$ for each $p \in Prim(S)$. However, this is true since the fixed space $fix(S_p)$ consists only of constant functions for every $p \in Prim(S)$.

The mapping $\hat{}$ is clearly linear and $\hat{1} = 1$. Next we show that $\hat{}$ is an isometric Markov lattice homomorphism. For $f \in \text{fix}(\mathbb{S})$

$$\|\hat{f}\| = \sup_{p \in \operatorname{Prim}(\mathbb{S})} |\hat{f}(p)| = \sup_{\mu \in \operatorname{ex} \operatorname{P}_{\mathbb{S}}(K)} |\langle f, \mu \rangle| \leqslant \|f\|.$$

The set $\{x \in K \mid |f(x)| = ||f||\}$ is the support of an S-ideal (see Theorem 1.2 in [Sin68]) and thus contains a minimal support of an S-ideal M which in turn supports an ergodic measure μ . This implies

$$|f(I_M)| = |\langle f, \mu \rangle| = |f(x)| = ||f||$$

for each $x \in M$ and consequently $\|\hat{f}\| = \|f\|$. Now take $f \in \text{fix}(S)$ and $p = I_{\mu} \in \text{Prim}(S)$. For each $x \in \text{supp}(\mu)$

$$|\hat{f}(p)| = |f(x)| = |f|(x) = \widehat{|f|}(p).$$

It remains to show that $\hat{}$ is surjective. Take $f \in C(Prim(S))$ with $0 \leq f \leq 1$. We fix $n \in \mathbb{N}$ and consider the open sets

$$U_{k,n} := \left\{ p \in \operatorname{Prim}(\mathbb{S}) \, \middle| \, \frac{k-1}{n} < f(p) < \frac{k+1}{n} \right\}$$

for $k \in \{0, ..., n\}$. Then $U_{k,n}^c = \text{hull}(I_{k,n})$ for invariant ideals $I_{k,n} \subseteq C(K)$ and $k \in \{0, ..., n\}$. Assume

$$I := \sum_{k=0}^{n} I_{k,n} \neq \mathcal{C}(K).$$

Then I is contained in a maximal S-ideal p. Since $I_{k,n} \subseteq p$ for all $k \in \{0, ..., n\}$,

$$p \in \bigcap_{k=0}^{n} \operatorname{hull}(I_{k,n}) = \left(\bigcup_{k=0}^{n} U_{k,n}\right)^{c} = \emptyset,$$

a contradiction.

We thus find $0 \leq f_{k,n} \in I_{n,k}$ for $k \in \{0, ..., n\}$ with $\mathbb{1} = \sum_{k=0}^{n} f_{k,n}$ (see II.5.1.4 in [Bla06]). Now set $g_n := \sum_{k=1}^{n} \frac{k}{n} f_{k,n}$. Take $p \in \text{Prim}(\mathcal{S})$. If $k \in \{0, ..., n\}$ with $p \notin U_{k,n}$, then $I_{k,n} \subseteq p$ and therefore

ī

 $f_{k,n} \in p$, i.e., $f_{k,n}|_{\operatorname{supp} p} = 0$. This implies

$$|g_n(x) - f(p)| = \left| \sum_{k: p \in U_{k,n}} \frac{k}{n} f_{k,n}(x) - \sum_{k: p \in U_{k,n}} f_{k,n}(x) f(p) \right|$$
$$\leqslant \sum_{k: p \in U_{k,n}} \left| \frac{k}{n} - f(p) \right| |f_{k,n}(x)| \leqslant \frac{1}{n}$$

for $x \in \operatorname{supp} p$. In particular we obtain

$$|g_n(x) - g_m(x)| \leq \frac{1}{n} + \frac{1}{m}$$

for all $x \in M := \bigcup_{p \in \operatorname{Prim}(S)} \operatorname{supp} p$ and all $n, m \in \mathbb{N}$. Since S is radical free, M is dense whence $(g_n)_{n \in \mathbb{N}}$ is a Cauchy sequence in C(K). Denoting its limit by g we obtain g(x) = f(p) for $x \in \operatorname{supp} p, p \in \operatorname{Prim}(S)$ and, since M is dense, $g \in \operatorname{fix}(S)$. Moreover, we clearly have $\hat{g} = f$.

The first part of the proof of Theorem 6.3 is based on the proof of Theorem 5 in [Sch68], while the second part uses arguments from the Dauns-Hofmann Theorem II.6.5.10 in [Bla06].

We now focus on the general case, i.e., S not being radical free.

Lemma 6.4. The mapping

$$\vartheta \colon \operatorname{Prim}(\mathbb{S}) \longrightarrow \operatorname{Prim}(\mathbb{S}_{\operatorname{rad}_{\mathbb{S}}(0)}), \quad p \mapsto p|_{\mathcal{M}(\mathbb{S})}$$

is a homeomorphism.

Proof. Note first that ϑ is well-defined and bijective by Lemma 3.6 since $\operatorname{rad}_{\mathbb{S}}(0) \subseteq p$ for each $p \in \operatorname{Prim}(\mathbb{S})$. Lemma 3.6 also implies that

$$\operatorname{Rad}(S) \longrightarrow \operatorname{Rad}(S_{\operatorname{rad}_{S}(0)}), \quad I \mapsto I|_{\operatorname{M}(S)}$$

is bijective. By Proposition 5.5 (i) we obtain that $A \subseteq Prim(S)$ is closed if and only if A = hull(I) for some $I \in Rad(S)$. If A is closed, we therefore obtain

$$\begin{split} \vartheta(A) &= \vartheta(\{p \in \operatorname{Prim}(\mathbb{S}) \mid I \subseteq p\}) \\ &= \{q \in \operatorname{Prim}(\mathbb{S}_{\operatorname{rad}_{\mathbb{S}}(0)}) \mid I \subseteq \vartheta^{-1}(q)\} \\ &= \{q \in \operatorname{Prim}(\mathbb{S}_{\operatorname{rad}_{\mathbb{S}}(0)}) \mid I|_{\operatorname{M}(\mathbb{S})} \subseteq q\} \\ &= \operatorname{hull}(I|_{\operatorname{M}(\mathbb{S})}), \end{split}$$

and therefore $\vartheta(A)$ is closed. Conversely, if $\vartheta(A)$ is closed, then by Lemma 3.6 and Proposition 5.5 (i) there is $I \in \operatorname{Rad}(S)$ with $\vartheta(A) = \operatorname{hull}(I|_{\operatorname{M}(S)})$ and, by the above, we obtain $A = \operatorname{hull}(I)$.

By combining Lemma 6.4 with Theorem 6.3 we obtain the main result of this section.

Theorem 6.5. The fixed space $fix(S_{rad_{S}(0)})$ is a Banach sublattice of C(M(S)) and the mapping

 $\hat{}: \ \mathrm{fix}(\mathbb{S}_{\mathrm{rad}_{\mathbb{S}}(0)}) \longrightarrow \mathrm{C}(\mathrm{Prim}(\mathbb{S})), \quad f \mapsto \widehat{f}$

is an isometric Markov lattice isomorphism.

HENRIK KREIDLER

7. Mean Ergodic Semigroups of Markov Operators

Using the space C(Prim(S)) we can now analyze mean ergodicity of Markov semigroups and extend Theorem 2 of [Sch67]. Recall that $\overline{co} S$ denotes the closed convex hull of S with respect to the strong operator topology. The right amenable semigroup S is *mean ergodic* if there is $P \in \overline{co} S$ with PS =SP = P for each $S \in S$ (see [Nag73] or [Sch13] for this concept). In this case P is unique with these properties and a projection onto the fixed space fix(S) of S, called the *mean ergodic projection*.

Theorem 7.1. The following assertions are equivalent.

- (a) S is mean ergodic.
- (b) The following two conditions are satisfied.
 - (i) The mapping

$$\exp P_{\mathcal{S}}(K) \longrightarrow \Pr(\mathcal{S}), \quad \mu \mapsto I_{\mu}$$

is a homeomorphism.

- (ii) For each $f \in \text{fix}(S_{\text{rad}_{S}(0)})$ there is $F \in \text{fix}(S)$ with $f = F|_{M(S)}$.
- (c) The following three conditions are satisfied.
 - (i) The primitive spectrum Prim(S) is a Hausdorff space.
 - (ii) For each $\mu \in \exp P_{S}(K)$ the support $\operatorname{supp} \mu$ is uniquely ergodic, i.e., μ is the only invariant measure having its support in $\operatorname{supp} \mu$.
 - (iii) For each $f \in \text{fix}(S_{\text{rad}_{S}(0)})$ there is $F \in \text{fix}(S)$ with $f = F|_{M(S)}$.

Remark 7.2. Note that assertion (c) (i) of Theorem 7.1 implies that each primitive ideal is maximal (see Proposition 5.5 (iv)). This shows that the concept of maximal S-ideals is sufficient for mean ergodic semigroups. In particular, combining Theorem 7.1 with Proposition 3.7 (i) implies Theorem 2 of [Sch67].

Proof (of Theorem 7.1). "(a) \Rightarrow (c)": Assume that S is mean ergodic with mean ergodic projection $P \in \mathscr{L}(C(K))$. We first show that Prim(S) is Hausdorff.

Consider I_{μ_1} , $I_{\mu_2} \in \text{Prim}(S)$ with $\mu_1 \neq \mu_2$. Since S is mean ergodic, we find $f \in \text{fix}(S)$ with

$$c_1 \coloneqq \langle f, \mu_1 \rangle < \langle f, \mu_2 \rangle \eqqcolon c_2$$

by Theorem 1.7 of [Sch13]. Choose $c \in (c_1, c_2)$ and set $U_1 := f^{-1}((-\infty, c))$ and $U_2 := f^{-1}((c, \infty))$. The sets

 $V_i := \{ p \in \operatorname{Prim}(\mathcal{S}) \mid \operatorname{supp} p \cap U_i \neq \emptyset \} \subseteq \operatorname{Prim}(\mathcal{S})$

are open by Proposition 5.5 (ii) and $I_{\mu_i} \in V_i$ for i = 1, 2. Assume there is $p \in V_1 \cap V_2$. Then there are $x_i \in U_i \cap \operatorname{supp} p$ for i = 1, 2and thus $f(x_1) < c < f(x_2)$. Since f is constant on supports of ergodic measures, this is a contradiction.

Given $\mu \in \exp P_{\mathcal{S}}(K)$ we know that $\mathcal{S}_{I_{\mu}} \subseteq \mathscr{L}(C(\operatorname{supp} \mu))$ is also mean ergodic. Since fix $(\mathcal{S}_{I_{\mu}})$ is one-dimensional, we obtain that fix $(\mathcal{S}'_{I_{\mu}})$ is one dimensional, too. Thus, the supports of ergodic measures are uniquely ergodic. Next, take $f \in \text{fix}(S_{\text{rad}_{S}(0)})$ and let G be any continuous extension of f to K. Then $F := PG \in \text{fix}(S)$ with $F|_{M(S)} = PG|_{M(S)} = f$. Thus (a) implies (c).

"(c) \Rightarrow (b)": Now suppose that (i), (ii) and (iii) of (c) are valid. We first show that ex $P_{\mathcal{S}}(K)$ is compact. Take a net $(\mu_{\alpha})_{\alpha \in A}$ in ex $P_{\mathcal{S}}(K)$ with $\lim_{\alpha} \mu_{\alpha} = \mu \in P_{\mathcal{S}}(K)$. Since I_{μ} is a radical ideal by Proposition 3.7 (i) we obtain

$$I_{\mu} = \bigcap_{\substack{p \in \operatorname{Prim}(S)\\ I_{\mu} \subseteq p}} p.$$

Now take $p \in Prim(S)$ with $p \supseteq I_{\mu}$ and any $f \in C(K)$ with $p \in U_f$. Then $f \notin p$ and consequently $f \notin I_{\mu}$. This implies $\langle |f|, \mu \rangle \neq 0$ and thus there is $\alpha_0 \in A$ with $\langle |f|, \mu_{\alpha} \rangle \neq 0$ for all $\alpha \ge \alpha_0$. But then $I_{\mu_{\alpha}} \to p$ and, since Prim(S) is Hausdorff, this implies that there is only one such p, hence I_{μ} is primitive. Applying (ii) shows that μ is ergodic.

We now obtain that

$$\pi \colon \exp \mathcal{P}_{\mathcal{S}}(K) \longrightarrow \operatorname{Prim}(\mathcal{S}), \quad \mu \mapsto I_{\mu}$$

is a homeomorphism since the mapping is injective by (ii), $\exp P_{\mathcal{S}}(K)$ is compact and $\operatorname{Prim}(\mathcal{S})$ is Hausdorff by (i).

"(b) \Rightarrow (a)": Finally assume that (b) is valid. The mapping

$$\Phi_1 \colon \mathcal{C}(\operatorname{Prim}(\mathfrak{S})) \longrightarrow \mathcal{C}(\exp_{\mathfrak{S}}(K)), \quad f \mapsto f \circ \pi$$

is then an isometric Markov lattice isomorphism and by Theorem 6.5 the mapping

$$: \operatorname{fix}(\mathcal{S}_{\operatorname{rad}_{\mathfrak{S}}(0)}) \longrightarrow \operatorname{C}(\operatorname{Prim}(\mathfrak{S}))$$

is so, too. Now consider the map

$$\Phi_2\colon \operatorname{fix}(\mathfrak{S}) \longrightarrow \operatorname{fix}(\mathfrak{S}_{\operatorname{rad}_{\mathfrak{S}}(0)}), \quad f \mapsto f|_{\mathcal{M}(\mathfrak{S})}.$$

This is an isometric Banach space embedding (isometry follows with the same arguments as in the proof of Theorem 6.5) and by (ii) it is surjective. Thus

$$\Phi := \Phi_1 \circ \circ \Phi_2 \colon \operatorname{fix}(\mathfrak{S}) \longrightarrow \operatorname{C}(\operatorname{ex} \operatorname{P}_{\mathfrak{S}}(K)), \quad f \mapsto \langle f, \cdot \rangle$$

is an isometric isomorphism of Banach spaces.

Now take $\mu_1, \mu_2 \in P_{\mathbb{S}}(K)$ with $\mu_1 \neq \mu_2$. The space $\exp P_{\mathbb{S}}(K)$ is compact since it is homeomorphic to $\operatorname{Prim}(\mathbb{S})$. By Choquet theory (see Proposition 1.2 in [Phe01]) we thus find measures $\tilde{\mu}_1, \tilde{\mu}_2 \in C(\exp P_{\mathbb{S}}(K))'$ with $\tilde{\mu}_1 \neq \tilde{\mu}_2$ such that

$$\langle f, \mu_i \rangle = \int_{\exp \mathsf{P}_{\mathsf{S}}(K)} \langle f, \nu \rangle \, \mathrm{d}\tilde{\mu}_i(\nu)$$

for each $f \in C(K)$ and i = 1, 2. We then obtain

$$\langle f, \mu_i \rangle = \int_{\mathrm{ex}\,\mathrm{P}_{\mathbb{S}}(K)} \Phi(f)(\nu) \,\mathrm{d}\tilde{\mu}_i(\nu) = \langle \Phi(f), \tilde{\mu}_i \rangle$$

for each $f \in \text{fix}(S)$ and i = 1, 2. Since $C(\exp P_S(K))$ separates $C(\exp P_S(K))'$, this proves that fix(S) separates $P_S(K)$. Now S consists of Markov operators

and therefore fix(S) separates fix(S'). Thus S is mean ergodic by Theorem 1.7 of [Sch13].

Corollary 7.3. If S is radical free, then S is mean ergodic if and only if $\exp P_{S}(K) \longrightarrow \operatorname{Prim}(S), \quad \mu \mapsto I_{\mu}$

is a homeomorphism.

The next corollary follows from Proposition 2.9 and Lemma 6.4.

Corollary 7.4. The semigroup S is mean ergodic if and only if $S_{rad_S(0)}$ is mean ergodic and for each $f \in fix(S_{rad_S(0)})$ there is $F \in fix(S)$ with $f = F|_{M(S)}$.

Finally we discuss some examples showing that the conditions of Theorem 7.1 (c) are independent of each other.

Example 7.5. Consider the following continuous mappings $\varphi \colon K \longrightarrow K$ and the induced semigroups $\mathcal{S} = \mathcal{S}_{\varphi} \subseteq \mathscr{L}(\mathcal{C}(K))$.

- (i) If K = [0,1] and $\varphi(x) = x^2$ for $x \in K$, then $M(S) = \{0,1\}$ and the primitive spectrum is the two point discrete space. Clearly, both fixed points define uniquely ergodic sets, so conditions (i) and (ii) of Theorem 7.1 (c) are fulfilled. However, S is not mean ergodic since the function $f: \{0,1\} \longrightarrow \mathbb{C}$ defined by f(0) := 0 and f(1) := 1 has no invariant continuous extension to K.
- (ii) If K = T then there is a homeomorphism φ: T → T such that S is not uniquely ergodic, but minimal (see Theorem 5.8 of [Par81]). In particular, the support of every ergodic measure is K. Thus the primitive spectrum is trivial and S is radical free whence Theorem 7.1 (c) (i) and (iii) are valid. S is not mean ergodic since supports of ergodic measures are not uniquely ergodic.
- (iii) If K and φ are defined as in Example 5.7 (iii), then supports of ergodic measures are uniquely ergodic and S is radical free. Thus Theorem 7.1 (c) (ii) and (iii) are fulfilled. The semigroup S is not mean ergodic since the primitive spectrum is not Hausdorff.

References

- [Ber51] R. R. Bernard. Probability in dynamical transformation groups. Duke Math. J., 18:307–319, 1951.
- [BJM89] J. F. Berglund, H. Junghenn, and P. Milnes. Analysis on Semigroups. Function Spaces, Compactifications, Representations. Wiley, 1989.
- [Bla06] B. Blackadar. Operator Algebras. Theory of C*-Algebras and von Neumann Algebras. Springer, 2006.
- [CD17] Z. Chen and X. Dai. Chaotic dynamics of minimal center of attraction for a flow with discrete amenable phase group. J. Math. Anal. Appl., 456:1397–1414, 2017.
- [Cho67] C. Chou. Minimal sets and ergodic measures for $\beta N \setminus N$. Illinois J. Math., 13:777–788, 1967.
- [Dai16] X. Dai. On chaotic minimal center of attraction of a Lagrange stable motion for topological semi flows. J. Differential Equations, 260:4393–4409, 2016.

22

- [Day61] M. M. Day. Fixed-point theorems for compact convex sets. Illinois J. Math., 5:585–590, 1961.
- [Dix77] J. Dixmier. C*-Algebras. North Holland, 1977.
- [EFHN15] T. Eisner, B. Farkas, M. Haase, and R. Nagel. Operator Theoretic Aspects of Ergodic Theory. Springer, 2015.
- [EW11] M. Einsiedler and T. Ward. Ergodic Theory with a view towards Number Theory. 2011.
- [GW10] U. Görtz and T. Wedhorn. Algebraic Geometry I. Vieweg+Teubner, 2010.
- [Hil36] H. Hilmy. Sur les centres d'attraction minimaux des systèmes dynamiques. Compos. Math., 3:227–238, 1936.
- [Hof11] K. H. Hofmann. The Dauns–Hofmann theorem revisited. J. Algebra Appl., 10:29–37, 2011.
- [JR72] K. Jacobs and J. Rosenmüller. Selecta Mathematica IV. Springer, 1972.
- [Kel75] J. L. Kelley. General Topology. Springer, 1975.
- [Man87] R. Mané. Ergodic Theory and Differentiable Dynamics. Springer, 1987.
- [MN91] P. Meyer-Nieberg. Banach Lattices. Springer, 1991.
- [Nag73] R. Nagel. Mittelergodische Halbgruppen linearer Operatoren. Ann. Inst. Fourier, 23:75–87, 1973.
- [Par81] W. Parry. Topics in Ergodic Theory. Cambridge University Press, 1981.
- [Pat88] A. L. T. Paterson. Amenability. American Mathematical Society, 1988.
- [Ped79] G. K. Pedersen. C*-Algebras and Their Automorphism Groups. Academic Press, 1979.
- [Phe01] R. R. Phelps. Lectures on Choquet's Theorem. Springer, 2nd edition, 2001.
- [Ros76] M. Rosenblatt. Transition probability operators. In Proc. Fifth Berkeley Symp. on Math. Statist. and Prob., volume 2 of Part 2, pages 473–483. Unif. of Calif. Press, 1976.
- [Sch67] H. H. Schaefer. Invariant ideals of positive operators in C(X). I. Illinois J. Math., 11:703–715, 1967.
- [Sch68] H. H. Schaefer. Invariant ideals of positive operators in C(X). II. Illinois J. Math., 12:525–538, 1968.
- [Sch74] H. H. Schaefer. Banach Lattices and Positive Operators. Springer, 1974.
- [Sch13] M. Schreiber. Uniform families of ergodic operator nets. Semigroup Forum, 86:321–336, 2013.
- [Sig77] K. Sigmund. On minimal centers of attraction and generic points. J. Reine Angew. Math., 295:72–79, 1977.
- [Sin68] R. Sine. Geometric theory of a single Markov operator. Pacific J. Math., 27:155–166, 1968.

Henrik Kreidler, Mathematisches Institut, Universität Tübingen, Auf der Morgenstelle 10, D-72076 Tübingen, Germany

E-mail address: hekr@fa.uni-tuebingen.de