### **TRIANGLE-ROUNDEDNESS IN MATROIDS**

## JOÃO PAULO COSTALONGA<sup>1</sup> AND XIANGQIAN ZHOU<sup>2</sup>

ABSTRACT. A matroid N is said to be triangle-rounded in a class of matroids  $\mathcal{M}$  if each 3connected matroid  $M \in \mathcal{M}$  with a triangle T and an N-minor has an N-minor with T as triangle. Reid gave a result useful to identify such matroids as stated next: suppose that M is a binary 3-connected matroid with a 3-connected minor N, T is a triangle of M and  $e \in T \cap E(N)$ ; then M has a 3-connected minor M' with an N-minor such that T is a triangle of M' and  $|E(M')| \leq |E(N)| + 2$ . We strengthen this result by dropping the condition that such element e exists

and proving that there is a 3-connected minor M' of M with an N-minor N' such that T is a triangle of M' and  $E(M') - E(N') \subseteq T$ . This result is extended to the non-binary case and, as an application, we prove that  $M(K_5)$  is triangle-rounded in the class of the regular matroids.

Key words: matroid minors; roundedness; matroid connectivity.

## 1. INTRODUCTION

Let  $\mathcal{M}$  be a class of matroids closed for minors and isomorphisms and let  $\mathcal{F}$  be a family of matroids. An  $\mathcal{F}$ -minor of a matroid M is a minor of M isomorphic to a member of  $\mathcal{F}$ . A matroid M uses a set T if  $T \subseteq E(M)$ . We say that  $\mathcal{F}$  is (k, t)-rounded in  $\mathcal{M}$  if each element of  $\mathcal{F}$  is k-connected and, for each k-connected matroid  $M \in \mathcal{M}$  with an  $\mathcal{F}$ -minor and each t-subset  $T \subseteq E(M)$ , M has an  $\mathcal{F}$ -minor using T. We define  $\mathcal{F}$  to be t-rounded in  $\mathcal{M}$  if it is (t+1, t)-rounded in  $\mathcal{M}$ . A matroid N is said (k, t)-rounded (resp. t-rounded) in  $\mathcal{M}$  if so is  $\{N\}$ . When we simply say that a matroid or family of matroids is (k, t)-rounded or t-rounded with no mention to a specific class of matroids, we are referring to the class of all matroids.

Bixby [1] proved that  $U_{2,4}$  is 1-rounded. Seymour [14] established a method to find a minimal 1-rounded family containing a given family of matroids; in that work it is established that  $\{U_{2,4}, M(K_4)\}$ ,  $\{U_{2,4}, F_7, F_7^*\}$ ,  $\{U_{2,4}, F_7, F_7^*, M^*(K_{3,3}), M^*(K_5), M^*(K'_{3,3})\}$  and  $\{U_{2,5}, U_{3,5}, F_7, F_7^*\}$  are 1-rounded.

Seymour [16] proved that  $U_{2,4}$  is also 2-rounded and, later, in [17], established a method to find a minimal 2-rounded family containing a given family of matroids.

Khan [6] and Coullard [5] proved independently that  $U_{2,4}$  is not 3-rounded. To the best of our knowledge, there is no known criterion to check (k, t)-roundedness for  $k \ge 4$ . For  $t \ge 3$ , Oxley [9] proved that  $\{U_{2,4}, \mathcal{W}^3\}$  is (3,3)-rounded. Moss [7] proved that  $\{\mathcal{W}^2, \mathcal{W}^3, \mathcal{W}^4, M(\mathcal{W}_3), M(\mathcal{W}_4), Q_6\}$ 

<sup>&</sup>lt;sup>1</sup>Departamento de Matemática, Universidade Federal do Espírito Santo. Av. Fernando Ferrari, 514; Campus de Goiabeiras, 29075-910, Vitória, ES, Brazil. e-mail: joaocostalonga@gmail.com (corresponding author).

<sup>&</sup>lt;sup>2</sup>Department of Mathematics and Statistics, Wright State University, Dayton, OH, 45435, USA and School of Mathematical Sciences, Huaqiao University, Fujian, China. e-mail: xiangqian.zhou@wright.edu.

is (3,4)-rounded and  $\{M(\mathcal{W}_3), M(\mathcal{W}_4), M(\mathcal{W}_5), M(K_5 \setminus e), M^*(K_5 \setminus e), M(K_{1,2,3}), M^*(K_{1,2,3}), S_8\}$  is (3,5)-rounded in the class of the binary matroids.

There are results on classification of small *t*-rounded families of matroids for t = 1, 2. Oxley [8] proved that for  $|E(N)| \ge 4$ , *N* is 1-rounded if and only if  $N \cong U_{2,4}$ ,  $P(U_{1,3}, U_{1,1})$  or  $Q_6$  and 2-rounded if and only if  $N \cong U_{2,4}$ . Reid and Oxley [12] proved that, up to isomorphisms, the unique 2-rounded matroids with more than three members in the class of GF(q)-representable matroids are  $M(\mathcal{W}_3)$  and  $\mathcal{M}(\mathcal{W}_4)$  for q = 2,  $U_{2,4}$  and  $\mathcal{W}^3$  for q = 3 and  $U_{2,4}$  for  $q \ge 4$ .

In this work, we focus on a different type of roundedness. A family of matroids  $\mathscr{F}$  is said to be **triangle-rounded** in  $\mathscr{M}$  if all members of  $\mathscr{F}$  are 3-connected and, for each matroid  $M \in \mathscr{M}$ with an  $\mathscr{F}$ -minor and each triangle T of M, there is an  $\mathscr{F}$ -minor of M with T as triangle. We say that a matroid N is **triangle-rounded** in  $\mathscr{M}$  if so is  $\{N\}$ . Some examples of trianglerounded matroids and families are  $U_{2,4}$  in the class of all matroids,  $F_7$  in the class of binary matroids and  $M^*(K_{3,3})$  in the class of regular matroids (Asano, Nishizeki and Seymour [2]),  $M(K_5 \setminus e)$  in the class of regular matroids and  $\{S_8, J_{10}\}$  in the class of binary matroids (Reid [13]). The proofs for the triangle-roundedness of the later two rely on the following criterion:

**Theorem 1.** (*Reid* [13, Theorem 1.1]) Let  $\{e, f, g\}$  be a triangle of a 3-connected binary matroid M and N be a 3-connected minor of M with  $e \in E(N)$ . Then, there exists a 3-connected minor M' of M using  $\{e, f, g\}$  such that M' has a minor which is isomorphic to N and E(M') has at most |E(N)| + 2 elements.

Here we establish a stronger result for binary matroids:

**Theorem 2.** If M is a 3-connected binary matroid with a 3-connected minor N and T is a triangle of M, then M has a 3-connected minor M' using T with an N-minor N' such that  $E(M') - E(N') \subseteq T$ .

Theorem 2 will be proved using a stronger result:

**Theorem 3.** Let M be a matroid with a 3-connected minor N satisfying  $|E(N)| \ge 4$ . Suppose that T is a triangle of M and M is minor-minimal with the property that M is 3-connected and has an N-minor using T. Then  $r(M) - r(N) \le 2$  and for some N-minor N' of M,  $|E(M) - (E(N') \cup T)| \le 1$ . Moreover, if  $E(M) - E(N') \nsubseteq T$ , then one of the following assertions holds:

- (a)  $r(M)-r(N) = 1, r^*(M)-r^*(N) \in \{2,3\}$  and M has an element x such that  $E(M)-E(N') \subseteq T \cup x, T \cup x$  is a 4-point line of M,  $M \setminus x$  has no N-minor and x is the unique element of M such that si(M/x) is 3-connected with an N-minor; or
- (b)  $r^*(M) r^*(N) = 2$ ,  $r(M) r(N) \in \{1, 2\}$  and M has an element y such that  $E(M) E(N') \subseteq T \cup y$ ,  $T \cup y$  is a 4-cocircuit of M, M/y has no N-minor and  $M \setminus A$  has no N-minor for each subset A of T with  $|A| \ge 2$ .

All the possible cases described in this theorem indeed occur, we give examples in Section 3. We say that a graph *G* is **triangle-rounded** if so is M(G) in the class of graphic matroids. Using Theorem 2, we establish that  $K_5$  is triangle-rounded, in other words:

**Theorem 4.** If G is a 3-connected graph with a triangle T and a  $K_5$ -minor, then G has a  $K_5$ -minor with E(T) as edge-set of a triangle.

**Remark:**  $K_3$  and  $K_4$  are triangle-rounded, but no larger complete graph than  $K_5$  is trianglerounded. Indeed, consider, for disjoint sets X, Y and  $\{z\}$  satisfying  $|X|, |Y| \ge 2$ , a complete graph K with n vertices with  $X \cup Y \cup \{z\}$  as vertex set. Consider also a graph G extending K by two vertices x and y with  $E(G) - E(K) = \{xy, xz, yz\} \cup \{xx' : x' \in X\} \cup \{yy' : y' \in Y\}$ . Note that  $G \setminus xz/xy \cong K_n$ . But no  $K_n$ -minor of G uses  $\{xy, xz, yz\}$  because contracting any other edge than xy in G results in a graph with more than one parallel pair of edges.

The next result allows us to derive triangle-roundedness in the class of regular matroids from triangle-roundedness in the classes of graphic and cographic matroids.

**Theorem 5.** If a family  $\mathscr{F}$  of internally 4-connected matroids with no triads is triangle-rounded both in the class of graphic and cographic matroids, then  $\mathscr{F}$  is triangle-rounded in the class of regular matroids not isomorphic to  $R_{10}$ .

As  $R_{10}$  has no  $M(K_5)$ -minor and  $M(K_5)$  is internally 4-connected with no triads and trivially triangle-rounded in the class of cographic matroids, it follows from Theorems 4 and 5 that:

**Corollary 6.**  $M(K_5)$  is triangle-rounded in the class of regular matroids.

All proofs are in the next section.

# 2. Proofs

In this section we prove the theorems. For matroids *M* and *N* we write N < M to say that *M* has a proper minor isomorphic to *N*. The notation  $N \le M$  means that N < M or N = M. Next we state some results used in the proofs.

**Lemma 7.** (Whittle, [18, Lemma 3.6]) Let M be a 3-connected matroid with elements x and p such that si(M/x) and si(M/x, p) are 3-connected, but si(M/p) is not 3-connected. Then,  $r(M) \ge 4$  and there is a rank-3 cocircuit  $C^*$  of M containing x such that  $p \in cl_M(C^*) - C^*$ .

**Lemma 8.** (Whittle [18, Lemma 3.7]) Let  $C^*$  be a rank-3 cocircuit of a 3-connected matroid M such that  $p \in cl_M(C^*) - C^*$ .

- (a) If  $z_1, z_2 \in C^*$ , then  $si(M/p, z_1) \cong si(M/p, z_2)$ .
- (b) If N is a matroid and for some  $x \in C^*$ , si(N/x, p) is 3-connected with an N-minor, then si(N/z, p) is 3-connected with an N-minor for each  $z \in C^*$ .

**Lemma 9.** (Whittle [18, Lemma 3.8]) Let  $C^*$  be a rank-3 cocircuit of a 3-connected matroid M. If  $x \in C^*$  has the property that  $cl_M(C^*) - x$  contains a triangle of M/x, then si(M/x) is 3-connected.

From Lemma 9, we may conclude:

**Corollary 10.** Let *M* be a 3-connecteed matroid with a triangle *T* and a triad  $T^*$  such that  $T^* - T = \{x\}$  and  $T - T^* = \{y\}$ . Then si(M/x) and  $co(M \setminus y)$  are 3-connecteed.

**Lemma 11.** (*Wu*, [19, Lemma 3.15]) If  $I^*$  is a coindependent set in a matroid M and  $M \setminus I^*$  is vertically 3-connected, then so is M.

Using Seymour's Splitter Theorem (as stated in [11, Corollary 12.2.1]) and proceeding by induction on *i* using Lemma 11, we may conclude:

**Corollary 12.** Let N < M be 3-connected matroids such that M has no larger wheel or whirlminor than N in case N is a wheel or whirl respectively. Then, there is a chain of 3-connected matroids  $N \cong M_n < \cdots < M_1 < M_0 = M$  such that for each  $i = 1, \ldots, n$  there is  $x_i \in E(M_i)$  satisfying  $M_i = M_{i-1}/x_i$  or  $M_i = M_{i-1} \setminus x_i$ . Moreover, for  $I := \{x_i : M_{i-1} = M_i/x_i\}$  and  $I^* := \{x_i : M_{i-1} = M_i/x_i\}$ ,

- (a) I is an independent set and  $I^*$  is a coindependent set of M.
- (b) for each  $1 \le i \le n$ ,  $M/(I \cap \{x_1, \dots, x_i\})$  and  $(M \setminus (I^* \cap \{x_1, \dots, x_i\}))^*$  are vertically 3-connected.

**Theorem 13.** (Whittle, [18, Corollary 3.3]) Let N be a 3-connected minor of the 3-connected matroid M. If  $r(M) \ge r(N) + 3$ , then for each element x such that si(M/x) is 3-connected with an N-minor, there exists  $y \in E(M)$  such that si(M/y) and si(M/x, y) are 3-connected with N-minors.

**Theorem 14.** (*Whittle* [18, Lemma 3.4 and Theorem 3.1] and Costalonga [3, Theorem 1.3]) Let  $k \in \{1,2,3\}$  and let M be a 3-connected matroid with a 3-connected minor N such that  $r(M) - r(N) \ge k$ . Then M has a independent k-set J such that si(M/x) is 3-connected with an N-minor for all  $x \in J$ .

**Lemma 15.** (*Costalonga* [4, Corollary 4]) Suppose that N < M are 3-connected matroids with  $r^*(M) - r^*(N) \ge 4$  and N is cosimple. Then:

- (a) *M* has a coindependent set *S* of size 4 such that  $co(M \setminus e)$  is 3-connected with an *N*-minor for all  $e \in S$ ; or
- (b) *M* has distinct elements  $a_1, a_2, b_1, b_2, b_3$  such that,  $T_s := \{a_s, b_t, b_3\}$  is a triangle for  $\{s, t\} = \{1, 2\}, T^* := \{b_1, b_2, b_3\}$  is a triad of *M* and  $co(M \setminus T^*)$  is 3-connected with an *N*-minor.

Next, we prove Theorem 3 and, after, using Theorem 3, we prove Theorem 2. For a non-negative integer *k*, a *k*-segment of a matroid is a *k*-subset of a line of this matroid.

*Proof of Theorem 3:* Suppose that the result does not hold. This is, for each *N*-minor *N'* of *M*,  $E(M) - E(N') \nsubseteq T$  and items (a) and (b) of the theorem do not hold. It is already known that  $U_{2,4}$  is triangle-rounded [2], so, we may assume that  $|E(N)| \ge 5$ . The proof will be based on a series of assertions. First, note that it follows from the minimality of *M* that:

(I). If, for  $x \in E(M)$ , si(M/x) is 3-connected with an N-minor, then  $x \in cl_M(T)$ .

(II). If  $T^*$  is a triad and T is a triangle of M such that  $T^* - T = \{x\}$ , then  $M \setminus x$  has no N-minor.

*Subproof:* Suppose the contrary. Let  $T^* \cap T = \{a, b\}$ . As *N* is simple and cosimple,  $M \setminus x/a \setminus b$  has an *N*-minor. But  $M \setminus x/a \setminus b \cong M \setminus a, b/x$  and, therefore, M/x has an *N*-minor. By Corollary 10, si(M/x) is 3-connected. By (I),  $x \in cl_M(T)$ . As  $x \notin T$ , then  $T^*$  meets a 4-segment of *M*. This implies that  $M \cong U_{2,4}$ , a contradiction.

**(III).** *If, for*  $x \in E(M)$ *,*  $co(M \setminus x)$  *is* 3*-connected with an* N*-minor, then*  $x \in T$ *.* 

*Subproof:* Suppose the contrary. Then  $T \nsubseteq E(co(M \setminus x))$  and therefore, there is a triad  $T^*$  meeting *x* and *T*. This contradicts (II).

**(IV).** If si(M/x) and si(M/x, y) are 3-connected with N-minors, then so is si(M/y) and  $x, y \in T$ .

*Subproof:* First we prove that si(M/y) is 3-connected. Suppose the contrary. By Lemma 7, there is a rank-3 cocircuit  $C^*$  such that  $x \in C^*$  and  $y \in cl(C^*) - C^*$ . By (I),  $x \in cl(T)$ . By orthogonality [11, Proposition 2.1.11],  $T \subseteq cl(C^*)$ . As  $r(C^*) = 3$ , there is  $z \in C^* - cl(T)$  and *T* is a triangle of M/z contained in  $cl_{M/z}(C^*)$ . By Lemma 9, si(M/z) is 3-connected and, by Lemma 8, M/z has an *N*-minor. This contradicts (I). So, si(M/y) is 3-connected.

By (I),  $x, y \in cl(T)$ . If, for some  $\{a, b\} = \{x, y\}$ ,  $a \notin T$ , then, as M/b has an N-minor and a is in a parallel pair of M/b, it follows that  $M \setminus a$  has an N-minor. Moreover, in this case,  $T \cup a$  is a 4-segment of M and  $M \setminus a$  is 3-connected with an N-minor, contradicting the minimality of M. Thus,  $x, y \in T$ .

(V).  $r(M) - r(N) \le 2$  and  $r^*(M) - r^*(N) \le 3$ .

*Subproof:* If  $r(M) - r(N) \ge 3$ , then, by Theorem 14, there is an independent set *J* of size 3 such that si(M/x) is 3-connected with an *N*-minor for all  $x \in J$ . So, there is  $x \in J - cl_M(T)$ , a contradiction to (I). Thus,  $r(M) - r(N) \le 2$ .

If  $r^*(M) - r^*(N) \ge 4$ , then Lemma 15 applies. If item (a) of that Lemma holds, then we have an element  $x \in E(M) - T$  such that  $co(M \setminus x)$  is 3-connected with an *N*-minor, contradicting (III). So, consider the elements given by item (b) of Lemma 15. Let  $s \in \{1,2\}$ . Note that  $M \setminus T^*$ is isomorphic to a minor of  $M \setminus a_s$ . By Corollary 10,  $co(M \setminus a_s)$  is 3-connected with an *N*-minor and, therefore,  $a_s \in T$  by (III). So,  $a_1, a_2 \in T$ . By orthogonality between *T* and  $T^*$ ,  $b_3 \notin T$ . As  $M \setminus b_3$  has an *N*-minor, then, by (III),  $co(M \setminus b_3)$  is not 3-connected. By Bixby's Lemma [11, Lemma 8.7.3],  $si(M/b_3)$  is 3-connected. As  $M \setminus T^*$  has an *N*-minor, then, so has  $M/b_3$  and, therefore,  $si(M/b_3)$ . By (I),  $b_3 \in cl_M(T)$ . But  $b_3$  is in a triad and, therefore, in no 4-segment of M. So,  $b_3 \in T$ , a contradiction.

(VI). The theorem holds if there is a wheel or whirl W such that  $N < W \le M$ .

*Subproof:* Let  $W \cong M(W_n)$  or  $W^n$ . Since *N* is a minor of *W*, then *N* is isomorphic to a wheel or whirl. As  $|E(N)| \ge 5$ , then  $|E(N)| \ge 6$  and  $n \ge 4$ . As  $r(M) - r(N) \le 2$  and  $r(W) - r(N) \ge 1$ , hence  $r(M) - r(W) \le 1$ .

Suppose for a contradiction that r(M) = r(W). So, for some coindependet set  $J^*$  of M,  $W = M \setminus J^*$ . If y is a non-spoke of W, then W/y is vertically 3-connected by Corollary 10. Moreover, si(W/y) has an N-minor and so does si(M/y). So, by Lemma 11, si(M/y) is 3-connected with an N-minor for each non-spoke y of W. By (I), all non-spokes of W are in  $cl_M(T)$ , contradicting the fact that they are the elements of a set with rank at least 3 in W.

Thus, r(M) = r(W) + 1. Now, by Theorem 14, there is an element x such that si(M/x) is 3-connected with an W-minor W'. As r(M/x) = r(W'), there is coindependent set  $I^*$  of M/x such that  $W' = M/x \setminus I^*$ . If y is a non-spoke of W', W'/y = M/x,  $y \setminus I^*$  is vertically 3-connected and, by Lemma 11, so is M/x, y. By (IV), all non-spokes of W' are in T, a contradiction again.  $\Diamond$ 

As the theorem fails for *M*, *N* and *T*, by (VI), there is no wheel or whirl *W* such that  $N < W \le M$  and the hypotheses of Seymour's Splitter Theorem hold for *M* and *N*.

**(VII).** If  $x \in E(M)$  and si(M/x) is 3-connected with an N-minor, then  $x \in T$ .

Subproof: Suppose the contrary. By (I),  $x \in cl_M(T)$ , which is a line with more than 3 points. As  $M \setminus z$  is 3-connected for all  $z \in cl_M(T) - T$ , then  $M \setminus z$  has no *N*-minor if  $z \in cl_M(T) - T$ . As M/x has an *N*-minor,  $cl_M(T) = T \cup x$  and  $M \setminus x$  has no *N*-minor. This implies that  $r^*(M) - r^*(N) \ge 2$  as *T* is in a parallel class of M/x.

Let us check that for each  $z \in E(M) - x$ , si(M/z) is not 3-connected with an *N*-minor. Suppose the contrary, by (I),  $z \in cl_M(T)$ . So, *x* is in the non-trivial parallel class  $cl_M(T)$  of M/z. Since *N* is simple and M/z has an *N*-minor, then  $M/z \setminus x$ , and, therefore,  $M \setminus x$  have an *N*-minor. This is a contradiction to what we proved before. So, *x* is the unique element of *M* such that si(M/x) is 3-connected with an *N*-minor. By Theorem 14, r(M) - r(N) = 1.

Consider the structures defined as in Corollary 12. By what we proved, for all choices of  $M_1, \ldots, M_n$ , we have  $I = \{x\}$  and n = 3 or 4. As M/x has a parallel class with 3 elements, then  $x = x_3$  or  $x = x_4$ , so, we have two cases to consider:

*Case 1. We may pick*  $M_1, ..., M_n$  *with*  $x = x_n$ : For all  $y \in E(M) - x$  such that  $M \setminus y$  is 3connected with an *N*-minor, we have  $y \in T$  by (III). In particular this holds for each  $y \in I^* \cup$  $\operatorname{cl}_M(T) - x$ . So,  $I^* \cup (\operatorname{cl}_M(T) - x) \subseteq T$ . This implies the validity of the theorem and, in particular, of item (a). Contradicting the assumption that the Theorem fails for M, N and T.

*Case 2. Otherwise:* Now, necessarily, n = 4 and  $x = x_3$ . If  $M' := M \setminus x_1, x_2, x_4$  is 3-connected, then, as M'/x is 3-connected, we could choose  $x_4 = x$ , which does not hold in Case 2. Note, that x is in a cocircuit with size at most two in M'. As  $M_2 = M \setminus x_1, x_2$  is 3-connected, then x is in a serial pair of M' with an element z. This implies that  $M \setminus x_1, x_2, x_4/z \cong N$ . By Lemma 11, si(M/z) is 3-connected with an *N*-minor. This contradicts the uniqueness of x established before.

**(VIII).** If  $co(M \setminus x)$  and  $co(M \setminus x, y)$  are 3-connected with N-minors, then  $co(M \setminus y)$  is 3-connected and  $x, y \in T$ .

*Subproof:* Suppose the contrary. By (III),  $co(M \setminus y)$  is not 3-connected. By the dual of Lemma 7, there is a corank-3 circuit *C* containing *x* with  $y \in cl^*(C) - C$ .

First assume that  $C \neq T$ . If *M* has a 4-cocircuit  $D^*$  contained in  $C \cup y$ , then, as  $|D^* \cap C| \ge 3$ and  $T \nsubseteq C$ , there is  $z \in (D^* \cap C) - T$ . So,  $D^* - z$  is a triad of  $M \setminus z$  contained in  $cl_{M \setminus z}^*(C - z)$  and, by the dual of Lemmas 8 and 9,  $co(M \setminus z)$  is 3-connected with an *N*-minor. But this contradicts (III). Thus,  $C \cup y$  contains no 4-cocircuit of *M*. But  $r^*(C \cup y) = 3$  and  $y \in cl^*(C) - C$ , so  $C \cup y$  is the disjoint union of a singleton set  $\{e\}$  and a non-trivial coline  $L^*$  containing *y*. By the dual of Lemmas 8 and 9 again,  $co(M \setminus e)$  is 3-connected with an *N*-minor. By (III),  $e \in T$ . By the dual of Lemma 8, for some  $f \in L^*$ ,  $co(M \setminus x, y) \cong co(M \setminus f, y) \cong co(M \setminus L^*)$  has an *N*-minor. Thus, M/y has an *N*-minor and, by Bixby's Lemma, si(M/y) is 3-connected with an *N*-minor. By (VII),  $y \in T$ . Since *T* meets  $L^*$ , it follows that  $L^*$  is a triad and, as a consequence, |C| = 3. By orthogonality, there is  $g \in (L^* \cap T) - y \subseteq C$ . Since  $e \in (C \cap T) - L^*$ , then  $C \cup T$  is a 4-segment of *M* meeting a triad, a contradiction since *M* is 3-connected and not isomorphic to  $U_{2,4}$ . Therefore, C = T. Let  $C^*$  be a cocircuit such that  $y \in C^* \subseteq T \cup y$ . If  $C^*$  is a triad, we have a contradiction to (II) since  $y \in C^* - T$  and  $M \setminus y$  has an *N*-minor. So,  $C^*$  is a 4-cocircuit and  $C^* = T \cup y$ . By Bixby's Lemma, si(M/y) is 3-connected. Since  $y \notin T$ , then M/y has no *N*-minor by (VII). If r(M) = r(N), then  $N \cong M \setminus x$ , y and  $M \setminus y$  is 3-connected, therefore  $r(M) - r(N) \in \{1, 2\}$ .

For all 2-subsets *A* of *T*, *M*\*A* has no *N*-minor because, otherwise, *y* would be in the serial pair  $C^* - A$  of *M*\*A* and *M*/*y* would have an *N*-minor.

As  $co(M \setminus x, y)$  has an N-minor, hence  $r^*(M) - r^*(N) \ge 2$ . If  $r^*(M) - r^*(N) \ge 3$ , then, by Theorem 13, there is  $z \in E(M)$  such that  $co(M \setminus z)$  and  $co(M \setminus x, z)$  are 3-connected with *N*minors. By (III),  $z \in T$ . So, for  $A := \{x, z\} \subseteq T$ ,  $M \setminus A$  has an *N*-minor, a contradiction. Therefore,  $r^*(M) - r^*(N) = 2$ . To prove the theorem and item (b), we have to find an N-minor N' of M with  $E(M) - E(N') \subseteq T \cup y$ . Consider a chain of matroids, sets and elements as in Corollary 12. Let *a* and *b*, in this order, be the elements deleted from *M* in order to get  $M_n$  from *M* as in the chain (recall that  $r^*(M) - r^*(N) = 2$ ). By Lemma 11,  $co(M \setminus a)$  is 3-connected with an *N*-minor, hence, by (III),  $a \in T$ . It follows from Lemma 11, (IV), (V) and (VII) that  $I \subseteq T$ . We just have to prove now that b = y. Suppose the contrary. If  $b \in T$ , then, for  $A := \{a, b\} \subseteq T$ ,  $M \setminus A$ has an *N*-minor, a contradiction, as we saw before. Thus,  $b \notin T$  and, by (III),  $co(M \setminus b)$  is not 3-connected. So, a and b play similar roles as x and y and applying the same steps for a and b as we did for x and y, we conclude that  $D^* := T \cup b$  is a cocircuit of M. By circuit elimination on  $C^* := T \cup y$ ,  $D^*$  and any element e of T, it follows from the cosimplicity of M and from the orthogonality with T that  $(T - e) \cup \{y, b\}$  is a cocircuit of M. Therefore, y is in a series class of  $M \setminus a, b$ , which has an N-minor. But this implies that M/y has an N-minor, a contradiction. So, b = y and (b) holds.  $\Diamond$ 

Now, consider the structures as given by Corollary 12. It follows from (IV), (V) and (VII) that  $I \subseteq T$ . If  $r^*(M) - r^*(N) \leq 2$ , it follows from (III) and (VIII) that  $I^* \subseteq T$ . This implies that  $T \subseteq E(M) - E(N')$  for  $N' = M_n$  and the theorem holds in this case. So, we may assume that  $r^*(M) - r^*(N) = 3$ . If |I| = 0, then  $N' = M \setminus I^*$  and  $M \setminus e$  is 3-connected with an *N*-minor for all  $e \in I^*$  and by (III),  $I^* \subseteq T$ . So,  $|I| \geq 1$ . By Lemma 11 and (VIII), the elements of  $I^*$  with the two least indices in  $x_1, \ldots, x_n$  are in *T*. So,  $|T \cap I| \leq 1$  and as  $I \subseteq T$ , |I| = 1. Therefore, n = 4. Since  $I \subseteq T$ ,  $x_1 \notin I$  by the simplicity of  $M_1$ . If  $I = \{x_4\}$ , then  $M \setminus e$  is 3-connected for all  $e \in I^*$ , a contradiction, as before. Therefore,  $I = \{x_3\}$  or  $\{x_2\}$ . This implies that  $T = \{x_1, x_2, x_3\}$ .

By Theorem 14, there is a coindependent 3-set  $J^*$  of M such that  $co(M \setminus e)$  is 3-connected with an N-minor for all  $e \in J^*$ . By (III),  $J^* = T = \{x_1, x_2, x_3\}$ . If T meets a triad  $T^*$ , then, for  $f \in T^* - T$  and  $e \in T \cap T^*$  we have that  $M \setminus e$  and, therefore,  $M \setminus e/f$ , have an N-minor. But in this case, by Lemma 9, si(M/f) is 3-connected with an N-minor, a contradiction to (IV). Thus, T meets no triads of M. Next, we check:

## **(IX).** We may not pick $M_1, M_2, M_3, M_4$ in such a way that $I = \{x_3\}$ .

*Subproof:* Suppose the contrary. Then, we may pick the chain of matroids in such a way that  $M_4 = M \setminus x_1, x_2/x_3 \setminus x_4$  with  $I = \{x_3\}$ . As  $x_4 \notin T$ , by (III),  $\operatorname{co}(M \setminus x_4)$  is not 3-connected and  $M^*$  has a vertical 3-separation  $(A, x_4, B)$ . This is, both *A* and *B* are 3-separating sets of *M*,  $x_4 \in \operatorname{cl}^*(A) \cap \operatorname{cl}^*(B)$  and  $r^*(A), r^*(B) \ge 3$ . So, (A, B) is a 2-separation of  $M \setminus x_4$ , but  $M \setminus x_4, x_1, x_2/x_3$  is 3-connected and we may assume, therefore, that  $|A - \{x_1, x_2, x_3\}| = |A - T| \le 1$ . As  $|A| \ge 3$ , then

 $|A \cap T| \ge 2$  and *A* spans *T*. This implies that  $Y = A \cup T$  is a 3-separating set of *M*. Moreover, Y = T or  $Y = T \cup y$  for some  $y \in E(M) - T$ .

Next we prove that  $T \cup x_4$  is a cocircuit of M. Note that  $x_4 \in cl^*(Y)$ . If Y = T, then, as T meets no triads, it follows that  $T \cup x_4$  is a 4-cocircuit. So, we may assume that  $Y = T \cup y$  for some  $y \in E(M) - T$ . If  $r_M(Y) = 2$ , Y is a 4-segment of M. As  $M/x_3$  has an N-minor, then so has  $M \setminus y$ . In this case,  $M \setminus y$  is 3-connected, contradicting (III) since  $y \notin T$ . Thus,  $r_M(Y) = 3$ . Since |Y| = 4 and Y is 3-separating, it follows that  $r^*(Y) = 3 = r^*(T)$ . Now, T cospans y, and as  $T \cup y$  cospans  $x_4$ , it follows that T cospans  $x_4$  and  $T \cup x_4$  is a cocircuit of M since T meets no triads.

As  $T \cup x_4$  is a cocircuit of M, hence  $T - \{x_3, x_4\}$  is a serial pair of  $M_2 = M \setminus x_1, x_2$  which is 3-connected with at least 4 elements, a contradiction. Thus, (IX) holds.

Now, By (IX),  $I = \{x_2\}$  for all choices of chains. This implies that there are no pair of elements  $\{a, b\} \subseteq E(M)$  such that  $M \setminus a$  and  $M \setminus a, b$  are 3-connected with an *N*-minor. In particular,  $M \setminus x_1, x_3$  is not 3-connected. But  $M_3 = M \setminus x_1, x_3/x_2$  is 3-connected. As  $M \setminus x_1$  is 3-connected, then  $x_2$  is in a serial pair  $\{x_2, z\}$  of  $M \setminus x_1, x_3$ . Hence, M/z has an *N*-minor. Since  $M \setminus x_1$  is 3-connected, it follows that  $\{z, x_2, x_3\}$  is a triad of  $M \setminus x_1$ . But *T* meets no triads of *M* and, therefore,  $C^* := \{z, x_1, x_2, x_3\} = T \cup z$  is a 4-cocircuit of *M*. If  $z \in cl(T)$ , then  $r^*(C^*) = 2$  and  $C^*$  is a 2-separating set of *M*. This implies that  $r(M) = r(C^*) = 2$ , contradicting the fact that  $r(M) = r(N) + 1 \ge 3$ . So,  $z \notin cl(T)$  and *T* is a triangle of M/z contained in  $C^*$ . By Lemma 9, si(M/z) is 3-connected. But M/z has an *N*-minor. By (VII),  $z \in T$ , a contradiction. This proves the theorem.

*P*roof of Theorem 2: If  $|E(N)| \ge 4$ , then, since binary matroids cannot contains 4-point lines nor circuits meeting cocircuits in an odd number of elements, then items (a) and (b) of Theorem 3 does not hold. In this case, Theorem 3 implies the theorem. So we may assume that  $|E(N)| \le 3$ . If  $N \not\cong U_{1,3}$ , then, since *N* is 3-connected, if follows that *N* is isomorphic to a minor of  $M' := M|T \cong U_{2,3}$ . So, we may assume that  $N \cong U_{1,3}$ . In this case *M* has a circuit *C* meeting *T* but different from *T*. Now, for any *e* ∈ *C*−*T*, the theorem holds for  $M' = (M|T \cup C)/(C - (T \cup e))$ . □

We define  $K_{3,3}^{1,1}$  as the graph in Figure 1. The following lemma is a well-known result and is a straightforward consequence of Seymour's Splitter Theorem.

**Lemma 16.** If G is a 3-connected graph with a  $K_5$ -minor then, either  $G \cong K_5$  or G has a  $K_{3,3}^{1,1}$ -minor.



FIGURE 1.  $K \cong K_{3,3}^{1,1}$ 

First suppose that |V(G)| = 7. Now, *G* is obtained from  $K_5$  by two vertex splittings by edges  $e_1$  and  $e_2$  and, possibly, by the addition of an edge  $e_3$  with  $e_1, e_2 \in E(T)$  and  $e_3 \in E(T)$  in case it is added. So,  $e_1$  and  $e_2$  are adjacent in *G*. When obtaining *G* in this process, we split a vertex by  $e_1$ , then we split an endvertex of  $e_1$  by  $e_2$ . If x, y, and z are the three vertices incident to  $e_1$  or  $e_2$  in after these two splittings, then the sum of their degrees is 8. So, in order to get *G* with all vertices with degree at least 3, the addition of  $e_3$  is necessary. Note that this process describes that *G* is obtained from  $K_5$  by expanding a vertex into the triangle *T* with vertices x, y and z. Observe that there are two vertices  $u, v \in V(T)$  with degree 3. It is clear that for the edges  $e, f \notin E(T)$  incident to u and v respectively, we have  $si(G/e, f) \cong K_5$ .

So, we may assume that |V(G)| = 6. By Lemma 16, up to labels, *G* is obtained from  $K \cong K_{3,3}^{1,1}$  (the graph in Figure 1), by adding the edges of E(T) - E(K). Since  $K/uv \cong K_5$ , then  $uv \in T$  and we may assume without losing generality that  $V(T) = \{u, v, a\}$ , so G = K + va. Now, it is clear that  $G \setminus ba/ub$  is a  $K_5$ -minor of *G* using *T*. This proves the theorem.

The following lemma has a slightly stronger conclusion than [11, Proposition 9.3.5] (beyond the conclusions of [11, Proposition 9.3.5], it states beyond that *R* has a *K*-minor and describes the way it is obtained), but the proof for [11, Proposition 9.3.5] also holds for the following lemma.

**Lemma 17.** Let  $R = K \oplus_3 L$  be a 3-sum of binary matroids, where K and L are 3-connected and  $E(K) \cap E(L) = S$ . Then there are  $X, Y \subseteq E(L) - S$  such that  $K \cong R/X \setminus Y$ , where K is obtained from  $R/X \setminus Y$  by relabeling the elements  $s_1$ ,  $s_2$  and  $s_3$  of S in K by respective elements  $l_1$ ,  $l_2$  and  $l_3$  of L.

*Proof of Theorem 5:* Suppose that *R* is a matroid contradicting the theorem minimizing |E(R)|. So, *R* has a triangle *T* and an *M*-minor for some  $M \in \mathscr{F}$  but *M* has no  $\mathscr{F}$ -minor using *T*. If *R* is graphic or cographic, the theorem holds for *R*, so assume the contrary. By Seymour's Decomposition Theorem for Regular Matroids [11, Theorem 13.1.1], there are matroids *K* and *L* with at least 7 elements each, intersecting in a common triangle *S* such that  $R = K \oplus_3 L$  with *L* being 3-connected and *K* being 3-connected up to parallel classes of size two meeting *S*. Under these circunstances, we may assume that  $|E(K) \cap E(M)| \ge |E(L) \cap E(M)|$ .

If *C* is a cycle of *R* meeting both E(K) and E(L), then there is  $s \in S$  such that  $(C \cap E(N)) \cup s$  is a cycle of *N* for N = K, L. As we picked *L* with no parallel pairs, it follows that  $cl_R(E(K) - S) \cap E(L) = \emptyset$ .

Let us first check that *K* has an *M*-minor. Let  $M = R/I \setminus I^*$  for some independent set *I* and coindependent set  $I^*$  of *R*. Since  $\lambda_M(E(K) \cap E(M)) \leq \lambda_R(E(K) \cap E(M)) = 2$ , then as *M* is internally 4-connected, it follows that  $|E(M) \cap E(L)| \leq 3$ , and, moreover,  $E(M) \cap E(L)$  is not a triad of *M* because *M* has no triads. This implies that  $E(M) \cap E(L) \subseteq cl_M(E(M) - E(L))$ . By the format of the family of circuits of *R*, it follows that  $E(M) \cap E(L) \subseteq cl_R(E(K) - S)$ , which is empty. So,  $E(M) \subseteq E(K)$ . By Lemma 17, there is a minor *K'* of *R* obtained by relabeling the elements  $s_1$ ,  $s_2$  and  $s_3$  of *S* in *K* by respective elements  $l_1$ ,  $l_2$  and  $l_3$  of *L*. Consider the matroid *K''* 

obtained from K' by contracting each  $l_i$  for those indices  $i \in \{1, 2, 3\}$  such that  $s_i \in cl_L(I \cap E(L))$ . Now K'' is obtained from  $R/(E(L) \cap I) \setminus (E(L) \cap I^*)$  by relabeling the remaining elements of S. This implies that K'' and, therefore, K have M-minors.

If  $T \subseteq E(K)$ , then *K* has an  $\mathscr{F}$ -minor using *T* by the minimality of *R*. But *R* has an *K*-minor using *T* by Lemma 17 and this implies the theorem. So, *T* meets E(L). As  $cl_R(E(K) - S) \cap E(L) = \emptyset$ , it follows that  $X := T \cap E(L)$  has at least two elements. As *L* is 3-connected and  $\lambda_L(S) = \lambda_L(X) = 2$ , then  $\kappa_L(S, X) = 2$ . By Tutte's Linking Theorem [11, Theorem 8.5.2], there is a minor *N* of *L* with  $E(N) = S \cup X$  such that  $\lambda_N(S) = 2$ . Hence:

$$2 = \lambda_N(S) = r_N(S) + r_N(X) - r(N) \le 4 - r(N).$$

So,  $r(N) \le 2$ . But  $r(N) \ge r_N(S) \ge \lambda_N(S) = 2$ . Also  $r_N(X) \ge \lambda_N(X) = 2$ . This implies that *S* spans *N* and *X* contains no parallel pairs of *N*. Now, each element of *X* is in parallel with an element of *S* in *N*. Therefore, for  $N = L/A \setminus B$ , we have that  $R/A \setminus B$  is obtained from *K* by relabeling the elements of *S* by elements of *T*. So,  $R/A \setminus B$  is 3-connected with *T* as triangle and has an *M*-minor. By the minimality of *R*,  $R/A \setminus B$  has an  $\mathscr{F}$ -minor using *T* and this proves the lemma.

#### **3.** Sharpness

In this section we establish that Theorem 3 is sharp in the sense that all described cases may occur indeed.

First we construct an example for Theorem 3 with  $E(M) \subseteq E(N) \cup T$ . Consider a complete graph K on  $n \ge 14$  vertices. Let  $X := \{v_{i,j} : i = 1,2,3 \text{ and } j = 1,2,3,4\}$  be a 12-subset of V(K). Consider a triangle T on vertices  $u_1$ ,  $u_2$  and  $u_3$ , disjoint from K. Let  $G = K \cup T + \{u_i v_{i,j} : i = 1,2,3 \text{ and } j = 1,2,3,4\}$ . Define, for disjoint subsets A and B of E(T),  $H := G \setminus A/B$ , M := M(G) and N := M(H) provided H is a simple graph. For each  $e \in E(G) - E(T)$ , G/e has at least 3 parallel pairs; so,  $|E(si(M/e))| \le |E(G)| - 4 < |E(H)|$  and, therefore M/e has no N-minor. For e incident to  $v \in V(K)$ , in order to get a minor of  $G \setminus e$  with |V(K)| - 12 vertices with degree  $|V(K)| - 1 \ge 13$  and 12 vertices with degree  $|V(K)| \ge 14$  as in H, it is necessary to contract some edge out of T, thus  $G \setminus e$  has no H-minor either. Note that we may pick A with any size from 0 to 3, We always may pick  $B = \emptyset$ , and provided  $|A| \ge 1$  we may pick B with size from 1 to 3 - |A|.

Now, let us construct an example satisfying item (a) of Theorem 3. Let us pick M as a restriction of the affine space  $\mathbb{R}^3$ . Consider a 4-subset  $L := \{a, b, c, x\}$  of an line R. Let T := L - x. Now consider for each  $y \in T$  a line  $R_y$  meeting L in y in such a way that no three lines among R,  $R_a$ ,  $R_b$  and  $R_c$  lay in a same plane. Let  $m \ge 6$ . For each  $y \in T$ , pick a m-subset  $L_y$  of  $R_y$  containing y. Let M be the restriction of the affine space to  $L \cup L_a \cup L_b \cup L_c$ . Let  $N = M/x \setminus a, b$  or  $N = M/x \setminus T$ . Note that it is not possible to get a rank-3 minor of M with 3 disjoint (m - 1)-segments by contracting an element other than x. So all N-minors of M are minors of M/x and therefore, deleting at least two elements of T is necessary to get an N-minor of M. Moreover, deleting the element in the intersection of the three m-lines is the unique way to get and  $M/x \setminus T$  from an  $M/x \setminus a, b$ -minor of M.

Next, we construct an example satisfying item (b) of Theorem 3. We denote by M + e the matroid obtained by adding e freely to M. Start with a projective geometry P with  $r(P) \ge 6$ . Let F be a flat of P with  $4 \le r(F) \le r(P) - 2$ . Consider a copy U of  $U_{2,4}$  on ground set  $T \cup x := \{x, x_1, x_2, x_3\}$  with  $(T \cup x) \cap E(P) = \emptyset$ . Let y be an element out of  $E(P) \cup T \cup x$ . Let M be the matroid obtained by adding y freely to the flat  $F \cup T$  of  $(P + x) \oplus_2 U$ . Note that E(P) is a hyperplane of M and, therefore  $T \cup y$  is a 4-cocircuit of M. Define  $N_1 = M \setminus y/x_2 \setminus x_3$  and  $N_2 := N_1/x_1$ . Note that  $N_1 = P + x_1$  and  $N_2$  is the truncation of P with rank r(P) - 1 (the matroid on E(P) whose independent sets are those independent sets of P with size at most r(P) - 1).

Let  $i \in \{1,2\}$  and  $N = M/X \setminus Y$  be an  $N_i$ -minor of M. Note that  $r^*(M) - r^*(N) = 2$  and  $r(M) - r(N) \in \{1,2\}$ . For each  $p \in E(P)$ ,  $|E(si(M/p))| < |E(N_i)|$ . Thus, no element  $p \in P$  may be contracted in M in order to get an  $N_i$ -minor. So,  $X \subseteq T \cup Y$ .

Let us check that  $T \cup y$  meets no circuit of M with less than six elements other than T. Indeed,  $M \setminus y$  is a two sum of a 4-point line on  $T \cup x$  and a matroid with rank greater than five with x as free element. Thus all circuits of  $M \setminus y$  meeting T, except for T itself, have more than five elements. Moreover, M is obtained from  $M \setminus y$  adding y as a free element to a flat with rank greater than 4 and, therefore, all circuits of M containing y also have more than five elements. Hence, the triangles of M/X are precisely the triangles of P. Moreover, those must be the same triangles of N since all triangles of N are triangles of M/X and they occur in the same number. As deleting an element of P from M/X would result in a matroid with less triangles than  $N_i$ , it follows that  $Y \subseteq T \cup y$ . Hence,  $E(P) \subseteq E(N)$  for each minor N of Misomorphic to  $N_1$  or  $N_2$ .

Let us check that M/y has no  $N_2$ -minor and, therefore, no  $N_1$ -minor too. Suppose for a contradiction that N is an  $N_2$ -minor of M/y. We may assume that  $N = M/y, x_1 \setminus x_2, x_3$ . Note that  $x_1$  is a free element of the rank- $r_P(F)$  flat  $F \cup x_1$  of  $M/y \setminus x_2, x_3$ . This implies that N|F is a truncation of rank  $r_P(F) - 1$  of the rank- $r_P(F)$  projective geometry F. But, as  $N_2$  is the rank-(r(P)-1) truncation of P and  $r_P(F) \le r(P)-2$ , then all rank- $(r_P(F)-1)$  flats of  $N_1$  are projective geometries and so is F, a contradiction.

Now, for i = 1, 2, each  $N_i$ -minor of M is in the form  $M \setminus y, x_i / A$  with  $1 \le i \le 3$  and A being an i-subset of  $T - x_i$ . Let A be a 2-subset of T. As we proved for  $A = \{x_2, x_3\}$ , it follows that  $M \setminus A$  has no  $N_2$ -minor. Moreover, for  $x_k \in T - A$  it is clear that y is not a free element of  $M \setminus A / x_k$ , which, therefore, is not isomorphic to  $N_1$ . Thus,  $M \setminus A$  has no minor isomorphic to  $N_1$ . This implies that  $M \setminus A$  has no minor isomorphic to  $N_2$  neither.

# 4. ACKNOWLEDGEMENTS

The authors thank the Visiting Scholar Fund of the Department of Mathematics and Statistics at Wright State University for the support.

#### REFERENCES

- [1] R. E. Bixby,  $\ell$ -matrices and a characterization of binary matroids, Discr. Math. 8 (1974), 139-145.
- [2] T. Asano, T. Nishizeki and P. D. Seymour, A note on non-graphic matroids, J. Combin. Theory Ser. B 37 (1984) 290–293.
- [3] J. P. Costalonga, On 3-connected minors of 3-connected matroids and graphs, European J. Combin. **33** (2012), 72–81.

#### JOÃO PAULO COSTALONGA AND XIANGQIAN ZHOU

- [4] J. P. Costalonga, A splitter theorem on 3-connected matroids, European J. Combin. 69 (2018) 7-18.
- [5] C. R. Coullard, Counterexamples to conjectures on 4-connected matroids, Combinatorica, 6 (1986), 315–320.
- [6] J. Kahn, A problem of P. Seymour on nonbinary matroids, Combinatorica, 5 (1985), 319–323.
- [7] T. Moss, A minor-based characterization of matroid 3-connectivity, Adv. Appl. Math 50 (2013) 132-141.
- [8] J. G. Oxley, On the intersections of circuits and cocircuits in matroids. Combinatorica 4 (1984) 187–195.
- [9] J. G. Oxley, On nonbinary 3-connected matroids. Trans. Amer. Math. Soc. 300 (1987) 663-679.
- [10] J. G. Oxley, A characterization of certain excluded-minor classes of matroids. European J. Combin. **10** (1989) 275–279.
- [11] J. G. Oxley, *Matroid theory, Second Edition*, Oxford graduate texts in mathematics, Oxford University Press, 2006.
- [12] J. G. Oxley and T. J. Reid, The smallest rounded sets of binary matroids, European J. Combin. 11 (1990) 47-56.
- [13] T. J. Reid, Triangles in 3-connected matroids, Discrete Mathematics 90 (1991) 281–296.
- [14] P. D. Seymour, A note on the production of matroid minors. J. Combin. Theory Ser. B 22 (1977) 289–295.
- [15] P. D. Seymour, Decomposition of regular matroids, J. Combin. Theory Ser. B 28 (1980), 305–359.
- [16] P. D. Seymour, On minors of non-binary matroids, Combinatorica 1 (1981), 387–394.
- [17] P. D. Seymour, Minors of 3-connected matroids, European J. Combin. 6 (1985), 375–382.
- [18] G. Whittle, Stabilizers of classes of representable matroids, J. Combin. Theory Ser. B 77 (1999), 39–72.
- [19] H. Wu, On contractible and vertically contractible elements in 3-connected matroids and graphs, Discrete Math. **179** (1998), 185–203.