LINEAR INVISCID DAMPING AND ENHANCED DISSIPATION FOR THE KOLMOGOROV FLOW

DONGYI WEI, ZHIFEI ZHANG, AND WEIREN ZHAO

ABSTRACT. In this paper, we prove the linear inviscid damping and voticity depletion phenomena for the linearized Euler equations around the Kolmogorov flow. These results confirm Bouchet and Morita's predictions based on numerical analysis. By using the wave operator method introduced by Li, Wei and Zhang, we solve Beck and Wayne's conjecture on the optimal enhanced dissipation rate for the 2-D linearized Navier-Stokes equations around the bar state called Kolmogorov flow. The same dissipation rate is proved for the Navier-Stokes equations if the initial velocity is included in a basin of attraction of the Kolmogorov flow with the size of $\nu^{\frac{2}{3}+}$, here ν is the viscosity coefficient.

1. INTRODUCTION

In this paper, we study the incompressible Navier-Stokes equations on the torus $\mathbb{T}^2_{\delta} = \{(x,y) : x \in \mathbb{T}_{2\pi\delta}, y \in \mathbb{T}_{2\pi}\}$ with $0 < \delta \leq 1$:

(1.1)
$$\begin{cases} \partial_t V - \nu \Delta V + V \cdot \nabla V + \nabla P = 0, \\ \nabla \cdot V = 0, \\ V|_{t=0} = V_0(x, y). \end{cases}$$

Here $V = (V^1(t, x, y), V^2(t, x, y)), P(t, x, y)$ denote the velocity and pressure of the fluid respectively, and $\nu \ge 0$ is the viscosity coefficient. When $\nu = 0$, (1.1) is reduced to the Euler equations.

It is easy to see that $V_{NS} = (-a_0 e^{-\nu t} \cos y, 0)$ for any constant a_0 is a special solution of (1.1) called the bar state or Kolmogorov flow. The bar states are physically relevant due to the fact that they can be viewed as maximum entropy solutions of the Euler equations, as well as the fact that they dominate the flow when the aspect ratio of the torus $\delta < 1$ [11].

The first part of this paper is devoted to studying the linear inviscid damping for the linearized Euler equations around the Kolmogorov flow $(-\cos y, 0)$, which take in the vorticity formulation:

(1.2)
$$\begin{cases} \partial_t \omega + \mathcal{L}\omega = 0, \\ \omega|_{t=0} = \omega_0(x, y), \end{cases}$$

where $\omega = \partial_x V^2 - \partial_y V^1$ is the vorticity and

(1.3)
$$\mathcal{L} = u(y)\partial_x + u''(y)\partial_x(-\Delta)^{-1}, \quad u(y) = -\cos y.$$

Since the breakthrough work on Landau damping by Mouhot and Villani [28], the inviscid damping has been a very active field as an analogues of Landau damping in hydrodynamics. Bedrossian and Masmoudi [7] proved nonlinear inviscid damping for the 2-D Euler equations around the Couette flow (y, 0) for the perturbation in Gevrey class. On the other hand, Lin and Zeng [23] proved that nonlinear inviscid damping is not true for the perturbation in H^s for $s < \frac{3}{2}$. For general shear flows, the linear inviscid damping is also a difficult problem

Date: November 7, 2017.

due to the presence of nonlocal part $u''(y)\partial_x(-\Delta)^{-1}$. In such case, the linear dynamics is associated with the singularities at the critical layers u = c of the solution for the Rayleigh equation

$$(u-c)(\phi''-\alpha^2\phi) - u''\phi = f.$$

Case [12] gave a first prediction of linear damping for monotone shear flow. His prediction was confirmed by a series of works [29, 30, 36, 37], and finally by our work [34]. Roughly speaking, if $u(y) \in C^4$ is monotone and \mathcal{L} has no embedding eigenvalues, then

$$\|V(t)\|_{L^{2}} \leq \frac{C}{\langle t \rangle} \|\omega_{0}\|_{H^{-1}_{x}H^{1}_{y}}, \quad \|V^{2}(t)\|_{L^{2}} \leq \frac{C}{\langle t \rangle^{2}} \|\omega_{0}\|_{H^{-1}_{x}H^{2}_{y}},$$

for the initial vorticity ω_0 satisfying $\int_{\mathbb{T}} \omega_0(x, y) dx = 0$ and $P_{\mathcal{L}} \omega_0 = 0$, where $P_{\mathcal{L}}$ is the spectral projection to $\sigma_d(\mathcal{L})$.

For non-monotone flows such as Poiseuille flow $u(y) = y^2$ and Kolmogorov flow $u(y) = \cos y$, it is even difficult to predict the inviscid damping due to strong degeneration of the Rayleigh equation at critical points. Based on Laplace tools and numerical computations, Bouchet and Morita [10] gave a first prediction of the inviscid damping for non-monotone flows, especially Kolmogorov flow. Instead of vorticity mixing mechanism in the monotone case, they found that the depletion phenomena of the vorticity at the stationary streamlines plays the key mechanism leading to the inviscid damping for non-monotone flows. In our work [35], we confirmed the predictions of Bouchet and Morita about the linear damping and the vorticity depletion phenomena for a class of shear flows (u(y), 0) in a finite channel denoted by \mathcal{K} , which consists of the function u(y) satisfying $u(y) \in H^3(-1, 1)$, and $u''(y) \neq 0$ for critical points(i.e., u'(y) = 0) and $u'(\pm 1) \neq 0$. Moreover, for a class of symmetric flows including Poiseuille flow, we obtained the same decay rate of the velocity as in the monotone flow.

In this paper, we extend the linear inviscid damping and the vorticity depletion phenomena to the Kolmogorov flow on the torus. These results confirm Bouchet and Morita's predictions based on numerical computations. The first result is the linear inviscid damping.

Theorem 1.1. Let $\delta \in (0,1)$ and $\omega(t,x,y)$ be the solution of (1.2) with $\int_{\mathbb{T}_{2\pi\delta}} \omega_0(x,y) dx = 0$. Then it holds that

1. if $\omega_0(x,y) \in H_x^{-\frac{1}{2}} H_y^1$, then

$$\|V(t)\|_{L^2} \le \frac{C}{\langle t \rangle} \|\omega_0\|_{H_x^{-\frac{1}{2}}H_y^1};$$

2. if $\omega_0(x,y) \in H_x^{\frac{1}{2}}H_y^2$, then

$$\|V^{2}(t)\|_{L^{2}} \leq \frac{C}{\langle t \rangle^{2}} \|\omega_{0}\|_{H^{\frac{1}{2}}_{x}H^{2}_{y}};$$

3. if $\omega_0(x,y) \in H_x^{\frac{1}{2}} H_y^k$ for k = 0, 1, there exists $\omega_\infty(x,y) \in H_x^{\frac{1}{2}} H_y^k$ such that $\|\omega(t, x + tu(y), y) - \omega_\infty\|_{L^2} \longrightarrow 0$ as $t \to +\infty$.

Remark 1.2. When $\delta = 1$, Proposition 4.3, 4.5, 4.7 still hold for $|\alpha| \ge 2$. Thus, we can obtain the following decay result:

1. if
$$\omega_0(x,y) \in H_x^{-\frac{1}{2}} H_y^1$$
, then
 $\|P_{\geq 2}V(t)\|_{L^2} \leq \frac{C}{\langle t \rangle} \|\omega_0\|_{H_x^{-\frac{1}{2}} H_y^1};$

2. if $\omega_0(x,y) \in H_x^{\frac{1}{2}}H_y^2$, then

$$\|P_{\geq 2}V^{2}(t)\|_{L^{2}} \leq \frac{C}{\langle t \rangle^{2}} \|\omega_{0}\|_{H^{\frac{1}{2}}_{x}H^{2}_{y}}$$

Here and in what follows, $P_{\geq 2}$ denotes the orthogonal projection of $L^2(\mathbb{T}^2)$ to the subspace $\{f \in L^2(\mathbb{T}^2) : f(x,y) = \sum_{|\alpha| \geq 2} \widehat{f}(\alpha, y)e^{i\alpha x}\}.$

 α

The second result is the behavior of the solution at critical points $y = k\pi$ for $k \in \mathbb{Z}$, which in particular confirms the vorticity depletion phenomena found by Bouchet and Morita [10].

Theorem 1.3. Let $\delta \in (0,1)$ and $\omega(t,x,y)$ be the solution of (1.2) with $\int_{\mathbb{T}_{2\pi\delta}} \omega_0(x,y) dx = 0$. Then for $s > \frac{1}{2}$, it holds that for all $k \in \mathbb{Z}$,

1. if $\omega_e(x,y) \in H^{1+s}_x H^3_y$, then for $x \in \mathbb{T}_{2\pi\delta}$, |t| > 1,

$$|\omega(t, x, k\pi)| \le \frac{C}{|t|} \|\omega_e\|_{H^{1+s}_x H^3_y};$$

2. if $\omega_e(x,y) \in H_x^{1+s}H_y^3$, then for $x \in \mathbb{T}_{2\pi\delta}$, |t| > 1, $|V^2(t,x,k\pi)| \le \frac{C}{|t|^2} \|\omega_e\|_{H_x^{1+s}H_y^3}$;

3. if
$$\omega_o(x,y) \in H_x^{s-\frac{1}{2}} H_y^3$$
, then for $x \in \mathbb{T}_{2\pi\delta}$, $|t| > 1$,
 $|V^1(t,x,k\pi)| \le \frac{C}{|t|^{\frac{3}{2}}} \|\omega_o\|_{H_x^{s-\frac{1}{2}} H_y^3}$

Here $\omega_e(x,y) = \frac{1}{2}(\omega_0(x,y) + \omega_0(x,-y))$ and $\omega_o(x,y) = \frac{1}{2}(\omega_0(x,y) - \omega_0(x,-y)).$

Recently, Lin and Xu [22] proved the linear inviscid damping for shear flows including Kolmogorov flow in the following sense:

$$\lim_{T \to \infty} \frac{1}{T} \int_0^T \|V(t)\|_{L^2}^2 dt = 0.$$

Their proof is based on the Hamiltonian structure of the linearized Euler equation and an instability index theory recently developed by Lin and Zeng [24].

The second part of this paper is devoted to studying the 2-D incompressible Navier-Stokes equation with small viscosity on the torus. The vorticity formulation takes

(1.4)
$$\begin{cases} \omega_t - \nu \Delta \omega + V \cdot \nabla \omega = 0\\ \omega|_{t=0} = \omega_0(x, y). \end{cases}$$

Experiments and numerical studies [14, 26] have shown that the solutions of the nearly inviscid 2-D incompressible Navier-Stokes equations will approach to some quasi-stationary(metastable) states such as the bar stats and dipole states in a much shorter time than the diffusive time $O(\frac{1}{\nu})$. Then they dominate the dynamics for very long time intervals. Afterwards they decay on the diffusive time scale. See [1] for a similar phenomena for Burgers equation with small viscosity.

To explain the metastability phenomena, Beck and Wayne [2] considered the linearized Navier-Stokes equations around the bar states, which are relatively easy to analyse from a mathematical view of point. Moreover, the bar states are more physically relevant in the case when the aspect ratio of the torus $\delta < 1$ [11]. The linearized Navier-Stokes equations around the Kolmogorov flow $V_{NS} = (-a_0 e^{-\nu t} \cos y, 0)$ take as follows

(1.5)
$$\begin{cases} \partial_t \omega + \mathcal{L}_{\nu}(t)\omega = 0\\ \omega|_{t=0} = \omega_0(x, y), \end{cases}$$

where

(1.6)
$$\mathcal{L}_{\nu}(t) = -\nu\Delta - a_0 e^{-\nu t} \cos y \partial_x (1 + \Delta^{-1}).$$

To study the dynamic of (1.5), the main difficulty is that the linearized operator $\mathcal{L}_{\nu}(t)$ is nonself-adjoint, nonlocal and time-dependent. In [2], the authors dropped nonlocal part of $\mathcal{L}_{\nu}(t)$ and considered the following model equation

$$\partial_t \omega - \nu \Delta - a_0 e^{-\nu t} \cos y \partial_x w = 0.$$

Using the hypocoercive method introduced by Villani [33], they proved the enhanced dissipation rate of the solution in some Banach space X(see (3.7) in [2]):

$$\|\omega(t)\|_X \le Ce^{-M\sqrt{\nu t}} \|w_0\|_X$$

for any $t \in [0, \tau/\nu]$. This decay rate is much faster than the diffusive decay of $e^{-\nu t}$. The mechanism leading to the enhanced dissipation is due to mixing. See [3, 13, 20, 8, 9] and references therein for more relevant works. Let us mention recent important progress [4, 5, 6] on the enhanced dissipation for Couette flow (y, 0) in $\mathbb{T} \times \mathbb{R}$ applied to the subcritical transition.

For the full linearized operator $\mathcal{L}_{\nu}(t)$, Beck and Wayne numerically computed the eigenvalues of $-\mathcal{L}_{\nu}(t)$ for a fixed time. The real parts of the eigenvalues are negative and scale like $\sqrt{\nu}$. Based on these analysis, Beck and Wayne conjectured that the same result should hold for the full linearized equation. The following theorem gives an affirmative answer to Beck and Wayne's conjecture.

Theorem 1.4. Given $\delta \in (0,1)$ and $\tau > 0$, there exist constants $c_1 > 0$, C > 0, such that if ω satisfies (1.5) with $\omega_0 \in L^2$ and $\int_{\mathbb{T}_{2\pi\delta}} \omega_0(x, y) dx = 0$, then it holds that for $0 \le t \le \tau/\nu$,

$$\|\omega(t)\|_{L^{2}} \leq Ce^{-c_{1}\sqrt{\nu}t}\|\omega_{0}\|_{L^{2}},$$
$$\|V(t)\|_{\dot{H}^{1}_{x}L^{2}_{y}} \leq \frac{Ce^{-c_{1}\sqrt{\nu}t}}{\sqrt{1+\nu t^{3}}}\|\omega_{0}\|_{L^{2}}.$$

When $\delta = 1$, it holds that for $0 \le t \le \tau/\nu$,

$$\begin{aligned} \|P_{\geq 2}\omega(t)\|_{L^{2}} &\leq Ce^{-c_{1}\sqrt{\nu}t}\|P_{\geq 2}\omega_{0}\|_{L^{2}}, \\ \|P_{\geq 2}V(t)\|_{\dot{H}^{1}_{x}L^{2}_{y}} &\leq \frac{Ce^{-c_{1}\sqrt{\nu}t}}{\sqrt{1+\nu t^{3}}}\|P_{\geq 2}\omega_{0}\|_{L^{2}} \end{aligned}$$

Remark 1.5. When $\delta = 1$, the operator $1 + \Delta^{-1}$ has two additional kernels $\{\sin x, \cos x\}$, which yield two family of exact solutions $\{e^{-\nu t} \sin x, e^{-\nu t} \cos x\}$ of the linearized Navier-Stokes equations. Thus, we have to make the projection $P_{\geq 2}$ for the solution in order to obtain the enhanced dissipation.

In [22], Lin and Xu proved the enhanced dissipation in the sense: if $\int_{\mathbb{T}_{2\pi\delta}} \omega_0(x, y) dx = 0$, then for any $\tau > 0$ and $\epsilon > 0$, if ν is small enough,

$$\begin{aligned} \|P_{\geq 2}\omega(\tau/\nu)\|_{L^2} &\leq \epsilon \|P_{\geq 2}\omega_0\|_{L^2} \quad \text{for} \quad \delta = 1, \\ \|\omega(\tau/\nu)\|_{L^2} &\leq \epsilon \|\omega_0\|_{L^2} \quad \text{for} \quad \delta < 1. \end{aligned}$$

The proof is based on the Hamiltonian structure of the linearized Euler equation and RAGE theorem as in [13].

Very recently, Ibrahim, Maekawa and Masmoudi [19] proved the same decay rate of the vorticity for the following equation:

$$\partial_t \omega + \mathcal{L}_{\nu} w = 0, \quad \mathcal{L}_{\nu} = -\nu \Delta - a \cos y \partial_x (1 + \Delta^{-1}),$$

which is the linearized equation of the 2-D Navier-Stokes equation with a force $(-a\nu \cos y, 0)$ around the Kolmogorov flow $(-a \cos y, 0)$. Their proof is based on the pseudospectral bound of \mathcal{L}_{ν} . Pseudospectra is an important concept in understanding the hydrodynamic stability [31, 32] due to the non-normality of the linearized operators. Gallay et al. applied the pseudospectra method to study the stability of the Lamb-Oseen vortex in the regime of high circulation Reynolds number [18, 15, 25, 16, 21, 17]. Recently, McQuighan and Wayne [27] applied a similar method to revisit the metastability problem of the viscous Burgers equation. However, it seems difficult to apply the pseudospectra method to the time-dependent operators such as $\mathcal{L}_{\nu}(t)$.

As announced in a work by Li and the first two authors [21], the wave operator method could be used to solve Beck and Wayne's conjecture. In [21], we solved Gallay's conjecture on pseudospectral and spectral bounds on the Oseen vortices operator. One of the key ideas is to transform a nonlocal operator into a local one by constructing a wave operator. In this work, the analysis on linear inviscid damping naturally gives rise to our desired wave operator. Moveover, we also use the completely different method to handle the local operator.

Let us point out that the extra decay factor $\frac{1}{\sqrt{1+\nu t^3}}$ of the velocity in Theorem 1.4 comes from the inviscid damping mechanism of Kolmogorov flow. Even for the time-independent operator \mathcal{L}_{ν} , it seems difficult to deduce this decay estimate from the resolvent estimate. This decay estimate of velocity plays an important role in the following nonlinear enhanced dissipation result.

Theorem 1.6. Given $\delta \in (0,1), \tau > 0, \gamma > \frac{2}{3}$ and $C_1 > 1$, there exist constants $c_2, c_3 \in (0,1)$, such that if $0 < \nu < c_2, \omega_0 \in L^2$, $||(I - P_2)\omega_0||_{L^2} \le \nu^{\gamma}$, and $C_1^{-1} \le ||P_2\omega_0||_{L^2} \le C_1$, then the solution to (1.4) satisfies

$$\|P_{\neq 0}\omega(t)\|_{L^2} \le Ce^{-c_3\sqrt{\nu t}}\|P_{\neq 0}\omega_0\|_{L^2}$$

for all $0 < t < \tau/\nu$. Here P_2 denotes the orthogonal projection of $L^2(\mathbb{T}^2_{\delta})$ to the subspace $W_2 = span\{\sin y, \cos y\}.$

Remark 1.7. The stability threshold ν^{γ} for $\gamma > \frac{2}{3}$ may be not optimal. We can only achieve the stability threshold $\nu^{\frac{3}{4}}$ if we do not use the enhanced dissipation with an extra decay factor of the velocity.

Remark 1.8. The case of $\delta = 1$ is a challenging problem. In this case, we need to consider the linearized Navier-Stokes equations around the dipole states such as

$$e^{-\nu t}(-\sin y, \sin x), \quad e^{-\nu t}(-\cos y, \cos x).$$

2. Sketch of the proof

In this section, we present a sketch of the proof of main theorems.

2.1. Linear inviscid damping. The basic ideas are similar to [34, 35]. We introduce the stream function ψ so that $V = (\partial_y \psi, -\partial_x \psi)$ and $-\Delta \psi = \omega$. In terms of ψ , the linearized Euler equation (1.2) takes

$$\partial_t \Delta \psi + u(y) \partial_x \Delta \psi - u''(y) \partial_x \psi = 0,$$

here $u(y) = -\cos y$. Taking the Fourier transform in x, we get

$$(\partial_y^2 - \alpha^2)\partial_t\widehat{\psi} = i\alpha \big(u''(y) - u(y)(\partial_y^2 - \alpha^2)\big)\widehat{\psi}.$$

Inverting the operator $(\partial_y^2 - \alpha^2)$, we obtain

(2.1)
$$-\frac{1}{i\alpha}\partial_t\widehat{\psi} = \mathcal{R}_\alpha\widehat{\psi},$$

where

(2.2)
$$\mathcal{R}_{\alpha}\widehat{\psi} = -(\partial_y^2 - \alpha^2)^{-1} \big(u''(y) - u(\partial_y^2 - \alpha^2) \big) \widehat{\psi}.$$

Let Ω be a simple connected domain including the spectrum $\sigma(\mathcal{R}_{\alpha})$ of \mathcal{R}_{α} . We have the following representation formula of the solution to (2.1):

(2.3)
$$\widehat{\psi}(t,\alpha,y) = \frac{1}{2\pi i} \int_{\partial\Omega} e^{-i\alpha tc} (c - \mathcal{R}_{\alpha})^{-1} \widehat{\psi}(0,\alpha,y) dc.$$

In this way, the large time behaviour of the solution $\widehat{\psi}(t, \alpha, y)$ is reduced to the study of the resolvent $(c - \mathcal{R}_{\alpha})^{-1}$.

Let Φ be a solution of the inhomogeneous Rayleigh equation:

(2.4)
$$\begin{cases} \partial_y^2 \Phi - \alpha^2 \Phi - \frac{u''}{u-c} \Phi = f, \\ \partial_y \Phi(-\pi) = \partial_y \Phi(\pi), \quad \Phi(-\pi) = \Phi(\pi) \end{cases}$$

with $f(\alpha, y, c) = \frac{\widehat{\omega}_0(\alpha, y)}{i\alpha(u(y)-c)}$. Then we find that

(2.5)
$$(c - \mathcal{R}_{\alpha})^{-1}\widehat{\psi}(0, \alpha, y) = i\alpha\Phi.$$

To study the large time behaviour of $\hat{\psi}$, one of key ingredients is to establish the limiting absorption principle:

$$\Phi(\alpha, y, c \pm i\varepsilon) \to \Phi_{\pm}(\alpha, y, c) \quad \text{for } c \in \operatorname{Ran} u,$$

as $\varepsilon \to 0$. In [35], we established the limiting absorption principle for shear flows in the class \mathcal{K} , and proved that Φ_{\pm} is bounded in H_y^1 . These information is enough to show that the velocity decays to 0 as the time tends to infinity in L^2 sense. These results can be easily extended to the Kolmogorov flow.

To obtain the explicit decay estimates of the velocity, we need to know more precise behaviour of the limit function Φ_{\pm} . As in the case of symmetric flows, we decompose the solution of (2.4) into the odd part and even part. Each part can be handled as in the monotonic flow. Main difference is that $\frac{1}{u(y)-c}$ is more singular for c = u(0) and $c = u(\pi)$. For the symmetric flow considered in [35], $\frac{1}{u(y)-c}$ is singular only at c = u(0). Due to the singularity of $\frac{1}{u(y)-c}$, we can only establish the uniform estimates of the solution of the homogeneous Rayleigh equation in the weighted space:

(2.6)
$$(u-c)(\phi''-\alpha^2\phi) - u''\phi = 0.$$

The details will be presented in section 3.

In section 4, we solve the inhomogeneous Rayleigh equation. In section 5, we prove the limiting absorption principle and give the solution formula of the linearized Euler equations. Again, the decay estimates of the velocity will be proved by using the dual method. More precisely, for $f = (\partial_y^2 - \alpha^2)g$ with $g \in H^2(0,\pi) \cap H^1_0(0,\pi)$,

$$\int_0^\pi \widehat{\psi}_o(t,\alpha,y) f(y) dy = -\int_{u(0)}^{u(1)} K_o(c,\alpha) e^{-i\alpha ct} dc$$

and for $f = (\partial_y^2 - \alpha^2)g$ with $g \in H^2(0, \pi)$ and $g'(0) = g'(\pi) = 0$,

$$\int_0^\pi \widehat{\psi}_e(t,\alpha,y) f(y) dy = -\int_{u(0)}^{u(1)} K_e(c,\alpha) e^{-i\alpha ct} dc,$$

where $\hat{\psi}_o$ and $\hat{\psi}_o$ are the odd part and even part of $\hat{\psi}$ respectively. Thus, the decay estimates are reduced to the regularity of the kernels K_o and K_e . In section 6, we prove the decay estimates of the velocity under the regularity assumptions on the kernels. Section 7 and section 8 are devoted to the regularity of the kernels.

In section 9, we prove the decay estimates of the vorticity and velocity at critical points.

2.2. Enhanced dissipation. Motivated by [21], we will use the wave operator method to study the enhanced dissipation. We will construct a wave operator \mathbb{D} , which satisfies the following basic properties:

1.
$$\mathbb{D}(\cos y(1 + (\partial_y^2 - \alpha^2)^{-1})\omega) = \cos y\mathbb{D}(\omega);$$

2. $(1 - \alpha^{-2})\|\omega\|_{L^2}^2 \le \|\mathbb{D}(\omega)\|_{L^2}^2 \le \|\omega\|_{L^2}^2;$
3. $\|\sin y(\mathbb{D}(\partial_y^2\omega) - \partial_y^2\mathbb{D}(\omega))\|_{L^2} \le C(|\alpha|\|\omega\|_{L^2} + \|\partial_y\omega\|_{L^2})$

See section 10 for more properties.

In section 11, we establish the linear enhanced dissipation. First of all, we study the decay estimates for the model without nonlocal term in a short time scale $\nu^{-\frac{1}{2}}$. Our method is based on the change of unknown, which is very different from the hypocoercive method used by Beck and Wayne. Then we use the wave operator method to establish the decay estimates for the model with nonlocal term. Finally, the linear enhanced dissipation was obtained by using some kind of iteration argument.

In section 12, we establish the nonlinear enhanced dissipation. The proof is based on the linear enhanced dissipation, especially the decay estimate of the velocity. Another key observation is that the basic energy dissipation ensures that if $C_1^{-1} \leq ||P_2\omega(0)||_{L^2} \leq C_1$, then

$$C^{-1} \le \|P_2\omega(t)\|_{L^2} \le C \quad \text{for } 0 < t < \tau/\nu.$$

The proof used the iteration argument again.

3. The homogeneous Rayleigh equation

To solve (2.4), we first construct a smooth solution of the homogeneous Rayleigh equation on $[-\pi, \pi]$ with $u(y) = -\cos y$:

$$(u-c)(\phi'' - \alpha^2 \phi) - u'' \phi = 0,$$

where the complex constant c will be taken in four kinds of domains given by

$$D_{0} \triangleq \left\{ c \in (-1,1) \right\},$$

$$D_{\epsilon_{0}} \triangleq \left\{ c = c_{r} + i\epsilon, \ c_{r} \in [-1,1], 0 < |\epsilon| < \epsilon_{0} \right\},$$

$$B_{\epsilon_{0}}^{l} \triangleq \left\{ c = -1 + \epsilon e^{i\theta}, \ 0 < \epsilon < \epsilon_{0}, \ \frac{\pi}{2} \le \theta \le \frac{3\pi}{2} \right\},$$

$$B_{\epsilon_{0}}^{r} \triangleq \left\{ c = 1 - \epsilon e^{i\theta}, \ 0 < \epsilon < \epsilon_{0}, \ \frac{\pi}{2} \le \theta \le \frac{3\pi}{2} \right\},$$

for some $\epsilon_0 \in (0, 1)$. We denote

(3.1)
$$\Omega_{\epsilon_0} \triangleq \overline{D}_0 \cup D_{\epsilon_0} \cup B^l_{\epsilon_0} \cup B^r_{\epsilon_0}.$$

3.1. Rayleigh integral operator. Given $|\alpha| \ge 1$, let A be a constant larger than $C|\alpha|$ with $C \ge 1$ independent of α .

Definition 3.1. For a function f(y,c) defined on $[0,\pi] \times \Omega_{\epsilon_0}$, for k = 0,1,2 and j = 0,1, we define

$$\begin{split} \|f\|_{X_0^k} &\stackrel{def}{=} \sup_{(y,c) \in [0,\pi] \times D_0} \left| \frac{u'(y_c)^k f(y,c)}{\cosh(A(y-y_c))} \right|, \\ \|f\|_{X_j} &\stackrel{def}{=} \sup_{(y,c) \in [0,\pi] \times \Omega_{\epsilon_0}} \left| \frac{u'(y_c)^j f(y,c)}{\cosh(A(y-y_c))} \right|, \end{split}$$

where $y_c \in [0, \pi]$ satisfies $u(y_c) = c_r$ with

(3.2) $c_r = \operatorname{Re} c \quad \text{for } c \in D_{\epsilon_0} \cup \overline{D}_0, \quad c_r = -1 \quad \text{for } c \in B^l_{\epsilon_0}, \quad c_r = 1 \quad \text{for } c \in B^r_{\epsilon_0}.$

Definition 3.2. For a function f(y,c) defined on $[0,\pi] \times \Omega_{\epsilon_0}$, we define

$$\|f\|_{Y_C} \stackrel{def}{=} \sum_{k=0}^2 \sum_{\beta+\gamma=k} A^{-k} \|\partial_y^{\beta} \partial_c^{\gamma} f\|_{X_0^{\gamma}} + A^{-3} \|\partial_c^2 \partial_y f\|_{X_1^{3}}.$$

where

$$\|f\|_{X_1^3} \stackrel{\text{def}}{=} \sup_{\substack{(y,c) \in [0,\pi] \times D_0}} \left| \frac{Au'(y_c)^2 f(y,c)}{\cosh(A(y-y_c))(A+u'(y_c)/u_1(y,c)^2)} \right|$$

with $u_1(y,c) = \frac{\cos y_c - \cos y}{y - y_c}$.

Definition 3.3. Let y_c be as in Definition 3.1 with c_r defined by (3.2). The Rayleigh integral operator T is defined by

$$T(f) \stackrel{def}{=} T_0 \circ T_{2,2}(f) = \int_{y_c}^y \frac{1}{(u(y') - c)^2} \int_{y_c}^{y'} f(z,c)(u(z) - c)^2 dz dy',$$

where

$$T_0 f(y,c) \stackrel{def}{=} \int_{y_c}^{y} f(z,c) dz,$$

$$T_{k,j} f(y,c) \stackrel{def}{=} \frac{1}{(u(y)-c)^j} \int_{y_c}^{y} f(z,c) (u(z)-c)^k dz, \quad j \le k+1$$

Let us first list some basic properties of $\cos y$ and $\sin y$, which will be used frequently. The proof is easy and we omit it.

Lemma 3.4. There exists a constant C so that for any $y, y_c \in [0, \pi]$,

$$C^{-1} \max_{\substack{|y'-y_c|+|y'-y|=|y-y_c| \\ y'-y_c|+|y'-y|=|y-y_c| \\ z' = \frac{\cos y_c - \cos y}{y - y_c} \le C \max_{\substack{|y'-y_c|+|y'-y|=|y-y_c| \\ y'-y_c|+|y'-y|=|y-y_c| \\ z' = \frac{\cos y_c - \cos y}{y - y_c} \le C \sin \frac{y + y_c}{2}, \quad \frac{1}{\pi} y(\pi - y) \le \sin y \le \frac{4}{\pi^2} y(\pi - y).$$

In particular, we have

$$\left|\frac{(y-y_c)(\sin y + \sin y_c)}{\cos y_c - \cos y}\right| \le C,$$
$$\left|\int_y^{y_c} \frac{(y-y_c)^2}{(\cos y_c - \cos y)^2} dy'\right| \le \frac{C}{\sin y_c}$$

Throughout this paper, we denote

(3.3)
$$u_1(y,c) = \frac{\cos y_c - \cos y}{y - y_c} = \int_0^1 \sin(y_c + t(y - y_c))dt$$

Lemma 3.5. There exists a constant C independent of A such that

$$||Tf||_{Y_C} \le \frac{C}{A^2} ||f||_{Y_C}, \quad ||Tf||_{X_0} \le \frac{C}{A^2} ||f||_{X_0}.$$

Moreover, if $f \in C([0,\pi] \times \Omega_{\epsilon_0})$, then

$$T_0f, T_{2,2}f, Tf \in C([0,\pi] \times \Omega_{\epsilon_0}).$$

Proof. Let us prove the first inequality. A direct calculation shows that

$$\begin{aligned} \|T_0 f\|_{X_0^k} &= \sup_{(y,c) \in [0,\pi] \times D_0} \left| \frac{1}{\cosh A(y-y_c)} \int_{y_c}^y \frac{u'(y_c)^k f(z,c)}{\cosh A(z-y_c)} \cosh A(z-y_c) dz \right| \\ &\leq \sup_{(y,c) \in [0,\pi] \times D_0} \left| \frac{1}{\cosh A(y-y_c)} \int_{y_c}^y \cosh A(z-y_c) dz \right| \|f\|_{X_0^k} \\ &\leq \frac{1}{A} \|f\|_{X_0^k}. \end{aligned}$$

Using the fact that for any $0 < y < y' < y_c$ or $y_c < y' < y < \pi$,

(3.5)
$$|u(y') - u(y_c)| \le |u(y) - u(y_c)|,$$

we deduce that

(3.4)

(3.6)
$$\begin{aligned} \|T_{2,2}f\|_{X_0^k} &\leq \sup_{(y,c)\in[0,\pi]\times D_0} \left|\frac{y-y_c}{\cosh A(y-y_c)} \int_0^1 \cosh tA(y-y_c)dt\right| \|f\|_{X_0^k} \\ &\leq \frac{C}{A} \|f\|_{X_0^k}, \end{aligned}$$

which along with (3.4) shows that for $k \ge 0$,

(3.7)
$$\|Tf\|_{X_0^k} \le \frac{C}{A^2} \|f\|_{X_0^k}.$$

Using the fact that for any $0 < y < y' < y_c$ or $y_c < y' < y < \pi$, $|u(y') - u(y_c)| \le |u(y) - u(y_c)|$, thus $|u(y') - c| \le |u(y) - c|$, we get by Lemma 3.4 that

$$||T_{k,k+1}f||_{X_0^{j+1}} + ||u'(y)T_{k,k+1}f||_{X_0^j}$$

(3.8)
$$\leq C \sup_{(y,c)\in[0,\pi]\times D_0} \left| \frac{1}{\cosh A(y-y_c)} \int_0^1 \cosh t A(y-y_c) dt \right| \|f\|_{X_0^j} \leq C \|f\|_{X_0^j}.$$

It is easy to see that

$$\begin{split} \partial_y Tf(y,c) &= T_{2,2}f(y,c), \\ \partial_c Tf(y,c) &= 2T_0 \circ T_{2,3}f(y,c) - 2T_0 \circ T_{1,2}f(y,c) + T\partial_c f(y,c), \\ \partial_y^2 Tf(y,c) &= -2u'(y)T_{2,3}f(y,c) + f(y,c). \end{split}$$

Then it follows from (3.4), (3.6) and (3.8) that

(3.9)
$$\|\partial_y Tf\|_{X_0^0} + \frac{1}{A} \|\partial_y^2 Tf\|_{X_0^0} + \|\partial_c Tf\|_{X_0^1} \le \frac{C}{A} \|f\|_{X_0^0} + \frac{C}{A^2} \|\partial_c f\|_{X_0^1}.$$
 It is easy to see that

$$T_{k,k+1}f(y,c) = \int_0^1 f(y_c + t(y - y_c), c) \frac{t^k u_1(y_c + t(y - y_c), c)^k}{u_1(y, c)^{k+1}} dt,$$

from which and the fact that $|\partial_c u_1(y,c)| \leq \frac{C}{u'(y_c)}$, we can deduce that

(3.10)
$$\left|\frac{u'(y_c)^j \partial_c T_{k,k+1} f(y,c)}{\cosh A(y-y_c)}\right| \le C \left(\frac{\|f\|_{X_0^{j-1}}}{u_1(y,c)^2} + \frac{\|\partial_y f\|_{X_0^{j-1}} + \|\partial_c f\|_{X_0^j}}{u_1(y,c)}\right).$$

Direct calculations show that

$$\begin{split} \partial_y \partial_c T f(y,c) &= 2T_{2,3} f(y,c) - 2T_{1,2} f(y,c) + T_{2,2} \partial_c f(y,c), \\ \partial_y \partial_c^2 T f(y,c) &= 2 \partial_c T_{2,3} f(y,c) - 2 \partial_c T_{1,2} f(y,c) \\ &+ 2T_{2,3} \partial_c f(y,c) - 2T_{1,2} \partial_c f(y,c) + T_{2,2} \partial_c^2 f(y,c), \end{split}$$

and

$$\begin{split} \partial_c^2 Tf(y,c) =& 2\partial_c T_0 T_{2,3} f(y,c) - 2\partial_c T_0 T_{1,2} f(y,c) + \partial_c T \partial_c f(y,c) \\ =& 2T_0 \partial_c T_{2,3} f(y,c) - 2T_0 \partial_c T_{1,2} f(y,c) + \frac{f(y_c,c)}{3u'(y_c)^2} \\ &+ 2T_0 T_{2,3} \partial_c f(y,c) - 2T_0 T_{1,2} \partial_c f(y,c) + T \partial_c^2 f(y,c). \end{split}$$

•

It follows from (3.8) and (3.6) that

(3.11)
$$\|\partial_y \partial_c Tf\|_{X_0^1} \le C \|f\|_{X_0^0} + \frac{C}{A} \|\partial_c f\|_{X_0^1}$$

By (3.10), we have

$$\begin{aligned} \left| \frac{u'(y_c)^2 T_0 \partial_c T_{k,k+1} f(y,c)}{\cosh A(y-y_c)} \right| &\leq \frac{u'(y_c)}{\cosh A(y-y_c)} \int_{y_c}^{y} \cosh A(y'-y_c) \left| \frac{u'(y_c) \partial_c T_{k,k+1} f(y',c)}{\cosh A(y'-y_c)} \right| dy' \\ &\leq C u'(y_c) \left| \int_{y_c}^{y} \frac{1}{u_1(y',c)^2} dy' \right| \|f\|_{X_0^0} + \frac{C}{A} \left(\|\partial_y f\|_{X_0^0} + \|\partial_c f\|_{X_0^1} \right) \\ &\leq C \|f\|_{X_0^0} + \frac{C}{A} \left(\|\partial_y f\|_{X_0^0} + \|\partial_c f\|_{X_0^1} \right), \end{aligned}$$

from which, (3.4) and (3.8), we infer that

$$(3.12) \|\partial_c^2 Tf\|_{X_0^2} \le C \|f\|_{X_0^0} + \frac{C}{A} \|\partial_y f\|_{X_0^0} + \frac{C}{A} \|\partial_c f\|_{X_0^1} + \frac{C}{A^2} \|\partial_c^2 f\|_{X_0^2},$$

By (3.8) and (3.10), we get

$$\left|\frac{u'(y_c)\partial_y\partial_c^2 Tf(y,c)}{\cosh A(y-y_c)}\right| \le \frac{C\|f\|_{X_0^0}}{u_1(y,c)^2} + \frac{C\|\partial_y f\|_{X_0^0}}{u'(y_c)} + \frac{C\|\partial_c f\|_{X_0^1}}{u'(y_c)} + \frac{C\|\partial_c^2 f\|_{X_0^2}}{Au'(y_c)}$$

which implies that

$$(3.13) \|\partial_y \partial_c^2 Tf\|_{X_1^3} \le CA \|f\|_{X_0^0} + C \|\partial_y f\|_{X_0^0} + C \|\partial_c f\|_{X_0^1} + \frac{C}{A} \|\partial_c^2 f\|_{X_0^2}.$$

Putting (3.7), (3.9) and (3.11)-(3.13) together, we conclude the first inequality. The estimate of T in X_0 norm is similar to (3.7).

Now we check the continuity. We have

$$T_{2,2}(f) = \int_0^1 \frac{(y - y_c)(u(y_c + t(y - y_c)) - c)^2}{(u(y) - c)^2} f(y_c + t(y - y_c), c) dt.$$

Using the fact that for any $0 < y < y' < y_c$ or $y_c < y' < y < \pi$, $\left|\frac{(u(y')-c)^2}{(u(y)-c)^2}\right| \leq 1$ and the continuity of $\frac{(y-y_c)(u(y_c+t(y-y_c))-c)^2}{(u(y)-c)^2}$, and Lebesgue's dominated convergence theorem, we conclude the continuity of $T_{2,2}f$. The continuity of Tf follows from $Tf = T_0 \circ T_{2,2}f$.

3.2. Existence of the solution. We solve the homogeneous Rayleigh equation on $[0, \pi]$:

(3.14)
$$\begin{cases} \phi'' - \alpha^2 \phi - \frac{u''}{u-c} \phi = 0, \\ \phi(y_c, c) = u(y_c) - c, \quad \phi'(y, c) \Big|_{y=y_c} = u'(y_c). \end{cases}$$

Proposition 3.6. There exists $\phi_1(y,c) \in C([0,\pi] \times \Omega_{\epsilon_0})$ such that $\phi(y,c) = (u(y)-c)\phi_1(y,c)$ is a solution of the Rayleigh equation (3.14). Moreover, $\partial_y \phi_1(y,c) \in C([0,\pi] \times \Omega_{\epsilon_0})$, and there exists $\epsilon_1 > 0, C > 0$ such that for any $\epsilon_0 \in [0,\epsilon_1)$ and $(y,c) \in [0,\pi] \times \Omega_{\epsilon_0}$,

$$|\phi_1(y,c)| \ge \frac{1}{2}, \quad |\phi_1(y,c)-1| \le C|y-y_c|^2,$$

where the constants ϵ_1, C may depend on α .

Proof. By Lemma 3.5, the operator $1 - \alpha^2 T$ is invertible in the spaces Y_C and $X_0 \cap C([0, \pi] \times \Omega_{\epsilon_0})$. Therefore, we may take

$$\phi_1(y,c) = (1 - \alpha^2 T)^{-1} = \sum_{k=0}^{+\infty} \alpha^{2k} T^k 1$$

which converges in both Y_C and $X_0 \cap C([0,\pi] \times \Omega_{\epsilon_0})$. Then $\phi_1(y,c) \in C([0,\pi] \times \Omega_{\epsilon_0})$. Moreover, for $c \in D_0$, we have

$$T^{k}1(y,c) \ge 0, \quad \phi_{1}(y,c) \ge 1,$$

which ensures that there exists $\epsilon_1 > 0$ so that for any $\epsilon_0 \in [0, \epsilon_1)$ and $(y, c) \in [0, \pi] \times \Omega_{\epsilon_0}$,

$$|\phi_1(y,c)| \ge \frac{1}{2}, \ |\phi_1(y,c)| \le C$$

Thanks to $\phi_1 = 1 + \alpha^2 T \phi_1$, we have

(3.15)
$$\phi_1(y,c) = 1 + \int_{y_c}^y \frac{\alpha^2}{(u(y') - c)^2} \int_{y_c}^{y'} \phi_1(z,c)(u(z) - c)^2 dz dy',$$

from which, it follows that

$$((u-c)^2 \phi_1')' = \alpha^2 \phi_1 (u-c)^2, \quad \phi_1(y_c,c) = 1.$$

Let $\phi(y,c) = (u(y) - c)\phi_1(y,c)$. Then $\phi(y,c)$ satisfies the Rayleigh equation (3.14). It follows from (3.15) that

$$|\phi_1(y,c) - 1| \le \alpha^2 \int_{y_c}^y \int_{y_c}^z |\phi_1(y',c)| \left| \frac{u(y') - c}{u(z) - c} \right|^2 dy' dz \le C|y - y_c|^2.$$

As $\partial_y \phi_1(y,c) = \alpha^2 T_{2,2} \phi_1(y,c)$, by Lemma 3.5, we have $\partial_y \phi_1(y,c) \in C([0,1] \times \Omega_{\epsilon_0})$.

We need the following extra properties of $\phi_1(y, c)$.

Lemma 3.7. For $c \in D_0$, it holds that

- 1. $\phi_1(y,c) \ge 1;$
- 2. $\partial_y \phi_1(y,c) > 0$ for $y \in (y_c,\pi]$ and $\partial_y \phi_1(y,c) < 0$ for $y \in [0,y_c)$;
- 3. for any given $M_0 > 0$, there exists a constant C only depending on M_0 such that for $\beta + \gamma \leq 2$, $|y y_c| \leq M_0/|\alpha|$

$$|u'(y_c)^{\gamma}\partial_y^{\beta}\partial_c^{\gamma}\phi_1(y,c)| \le C|\alpha|^{\beta+\gamma},$$

$$|u'(y_c)^2\partial_y\partial_c^2\phi_1(y,c)| \le C\left(|\alpha|^3 + \frac{\alpha^2 u'(y_c)}{u_1(y,c)^2}\right)$$

Proof. The first two properties are a direct consequence of (3.15). By Lemma 3.5, we have $\|\phi_1\|_{Y_C} \leq 2$ if we take $A = C|\alpha|$ for some constant C independent of α . The third property follows from the definition of Y_C norm.

3.3. Uniform estimates of the solution. The goal of this subsection is to establish some uniform estimates in wave number α for the solution $\phi(y, c)$ for $c \in D_0$ of the Rayleigh equation given by Proposition 3.6.

Without loss of generality, we always assume $\alpha \geq 1$ in the sequel. We introduce

$$\mathcal{F}(y,c) = \frac{\partial_y \phi_1(y,c)}{\phi_1(y,c)}, \quad \mathcal{G}(y,c) = \frac{\partial_c \phi_1(y,c)}{\phi_1(y,c)},$$
$$\mathcal{G}_1(y,c) = \frac{\mathcal{F}}{u'(y_c)} + \mathcal{G} = \frac{1}{\phi_1} \Big(\frac{\partial_y}{u'(y_c)} + \partial_c \Big) \phi_1,$$

where $\phi_1(y,c) = \frac{\phi(y,c)}{u(y)-c}$ and $y_c \in [0,\pi]$ satisfying $c = u(y_c)$ for $c \in D_0$.

Proposition 3.8. There exists a constant C independent of α such that

$$\begin{split} \phi_1(y,c) &-1 \le C \min\{\alpha^2 | y - y_c |^2, 1\} \phi_1(y,c), \\ C^{-1} \alpha \min\{\alpha | y - y_c |, 1\} \le |\mathcal{F}(y,c)| \le C \alpha \min\{\alpha | y - y_c |, 1\}, \\ C^{-1} e^{C^{-1} \alpha | y - y_c |} \le \phi_1(y,c) \le e^{C \alpha | y - y_c |}, \\ |\partial_y^\beta \partial_c^\gamma \phi_1(y,c)| \le C \alpha^{\beta + \gamma} \phi_1(y,c) / u'(y_c)^\gamma, \quad \beta + \gamma \le 2, \\ |\partial_c \phi_1(y,c)| \le C \phi_1(y,c) \frac{\alpha \min\{1, \alpha | y - y_c |\}}{u'(y_c)}, \end{split}$$

and

$$\left| \left(\frac{\partial_y}{u'(y_c)} + \partial_c \right) \phi_1(y,c) \right| \le C \frac{\min\{1,\alpha^2 | y - y_c |^2\} \phi_1}{u'(y_c)^2},$$
$$\left| \left(\frac{\partial_y}{u'(y_c)} + \partial_c \right)^2 \phi_1(y,c) \right| \le C \frac{\min\{1,\alpha^2 | y - y_c |^2\} \phi_1}{u'(y_c)^4}.$$

Proof. Step 1. Estimates of \mathcal{F} and $\partial_u^k \phi_1$ for k = 0, 1, 2.

Recall that $\phi_1 = 1 + \alpha^2 T(\phi_1)$, which implies that

(3.16)
$$\phi_1(y,c) - 1 \le C \min\{\alpha^2 | y - y_c |^2, 1\} \phi_1(y,c)$$

It is easy to check that \mathcal{F} satisfies

(3.17)
$$\mathcal{F}' + \mathcal{F}^2 + \frac{2u'}{u-c}\mathcal{F} = \alpha^2, \quad \mathcal{F}(y_c, c) = 0.$$

Thanks to $\mathcal{F}(y,c) \geq 0$ for $y \geq y_c$ and $\mathcal{F}(y,c) \leq 0$ for $y \leq y_c$, we have $\frac{2u'}{u-c}\mathcal{F} \geq 0$. Due to $\phi'_1(y_c,c) = 0$, we get

$$\lim_{y \to y_c} \frac{u'(y)\mathcal{F}(y,c)}{u-c} = \lim_{y \to y_c} \frac{u'(y_c)\phi_1'(y,c)}{u-c} = \phi_1''(y_c,c) = \mathcal{F}'(y_c,c),$$

which along with (3.17) implies that $\mathcal{F}'(y_c, c) = \frac{\alpha^2}{3} > 0$. Take $y_0 \in [y_c, \pi]$ such that $\mathcal{F}(y_0, c) = \sup_{y \in [y_c, \pi]} \mathcal{F}(y, c)$. Then $y_0 > y_c$, $\mathcal{F}'(y_0, c) \ge 0$ and $\alpha^2 \ge (\mathcal{F}' + \mathcal{F}^2)(y_0, c) \ge \mathcal{F}^2(y_0, c)$, which implies that $\mathcal{F}(y, c) \le \mathcal{F}(y_0, c) \le \alpha$ for $y \in [y_c, \pi]$. For $y \in [0, y_c]$, we consider the minimal point of $\mathcal{F}(y, c)$ and obtain $\alpha \le \mathcal{F}(y_c, c)$. This shows that $|\mathcal{F}(y, c)| \le \alpha$, which in

point of $\mathcal{F}(y,c)$, and obtain $-\alpha \leq \mathcal{F}(y,c) \leq 0$. This shows that $|\mathcal{F}(y,c)| \leq \alpha$, which in particular gives

(3.18)
$$e^{-\alpha|y-y'|} \le \frac{\phi_1(y',c)}{\phi_1(y,c)} \le e^{\alpha|y-y'|}.$$

Using the fact that

$$\partial_y \phi_1(y,c) = \frac{\alpha^2}{(u(y)-c)^2} \int_{y_c}^y \phi_1(y',c)(u(y')-c)^2 \, dy',$$

and $(u(y')-c)^2 \leq (u(y)-c)^2$ and $\phi_1(y',c) \leq \phi_1(y,c)$ for $y_c \leq y' \leq y$ or $y \leq y' \leq y_c$, we infer that

$$|\mathcal{F}(y,c)| \le C\alpha^2 |y - y_c|.$$

On the other hand, we deduce from (3.18) and Lemma 3.4 that for $0 \le y_c \le y \le \min\{\frac{1}{\alpha} + y_c, \pi\}$,

$$\begin{aligned} \mathcal{F}(y,c) &= \frac{\alpha^2}{(u(y)-c)^2} \int_{y_c}^y \frac{\phi_1(y',c)}{\phi_1(y,c)} (u(y')-c)^2 \, dy' \\ &\geq \frac{\alpha^2}{(u(y)-c)^2} \int_{y_c}^y e^{-\alpha|y-y'|} (u(y')-c)^2 \, dy' \\ &\geq C^{-1} \frac{\alpha^2}{\sin^2 \frac{y+y_c}{2} (y-y_c)^2} \int_{\frac{y_c+y}{2}}^y \sin^2 \frac{y'+y_c}{2} (y'-y_c)^2 \, dy' \\ &\geq C^{-1} \alpha^2 |y-y_c|. \end{aligned}$$

Similarly, for $\max\{0, y_c - \frac{1}{\alpha}\} \le y \le y_c \le \pi$, we have

$$\begin{aligned} -\mathcal{F}(y,c) &\geq \frac{\alpha^2}{(u(y)-c)^2} \int_y^{y_c} e^{-\alpha|y-y'|} (u(y')-c)^2 \, dy' \\ &\geq C^{-1} \frac{\alpha^2}{\sin^2 \frac{y+y_c}{2} (y-y_c)^2} \int_y^{\frac{y_c+y}{2}} \sin^2 \frac{y'+y_c}{2} (y'-y_c)^2 \, dy' \\ &\geq C^{-1} \alpha^2 |y-y_c|. \end{aligned}$$

For $0 \leq \frac{1}{\alpha} + y_c \leq y \leq \pi$, we have

$$\begin{aligned} \mathcal{F}(y,c) &= \frac{\alpha^2}{(u(y)-c)^2} \int_{y_c}^y \frac{\phi_1(y',c)}{\phi_1(y,c)} (u(y')-c)^2 \, dy' \\ &\geq \frac{\alpha^2}{(u(y)-c)^2} \int_{y_c}^y e^{-\alpha|y-y'|} (u(y')-c)^2 \, dy' \\ &\geq C^{-1} \frac{\alpha^2}{\sin^2 \frac{y+y_c}{2} (y-y_c)^2} \int_{\max\left\{\frac{y_c+y}{2}, y-\frac{1}{\alpha}\right\}}^y e^{-\alpha|y-y'|} \sin^2 \frac{y'+y_c}{2} (y'-y_c)^2 \, dy' \\ &\geq C^{-1} \alpha. \end{aligned}$$

For $0 \le y \le y_c - \frac{1}{\alpha} \le \pi$, we have

$$\begin{aligned} -\mathcal{F}(y,c) &= \frac{\alpha^2}{(u(y)-c)^2} \int_y^{y_c} \frac{\phi_1(y',c)}{\phi_1(y,c)} (u(y')-c)^2 \, dy' \\ &\ge C^{-1} \frac{\alpha^2}{(u(y)-c)^2} \int_y^{y_c} e^{-\alpha|y-y'|} (u(y')-c)^2 \, dy' \\ &\ge C^{-1} \frac{\alpha^2}{\sin^2 \frac{y+y_c}{2} (y-y_c)^2} \int_y^{\min\left\{\frac{y_c+y}{2}, y+\frac{1}{\alpha}\right\}} e^{-\alpha|y-y'|} \sin^2 \frac{y'+y_c}{2} (y'-y_c)^2 \, dy' \\ &\ge C^{-1} \alpha. \end{aligned}$$

This shows that

 $(3.19) C^{-1}\alpha\min\{\alpha|y-y_c|,1\} \le |\mathcal{F}(y,c)| \le C\alpha\min\{\alpha|y-y_c|,1\},$ which along with $\phi_1(y_c,c) = 1$ implies that

(3.20)
$$C^{-1}e^{C^{-1}\alpha|y-y_c|} \le \phi_1(y,c) \le e^{C\alpha|y-y_c|},$$

(3.21)
$$|\partial_y \phi_1| \le C\alpha \min\{\alpha|y-y_c|,1\}\phi_1.$$

Using $\phi_1'' + \frac{2u'}{u-c}\phi_1' = \alpha^2\phi_1$ and $(u'(y) + u'(y_c))|y - y_c| \le C|u - c|$ (by Lemma 3.4), we obtain $|\partial_y^2\phi_1| \le C\alpha^2\phi_1$.

Step 2. Estimates of $\partial_c \phi_1$, $\partial_y \partial_c \phi_1$ and $\left(\frac{\partial_y}{u'(y_c)} + \partial_c\right) \phi_1$. It is easy to check that

$$\partial_c \mathcal{F} = \partial_y \mathcal{G}, \quad \left(\frac{\partial_y}{u'(y_c)} + \partial_c\right) \mathcal{F} = \partial_y \mathcal{G}_1, (\phi^2 \partial_c \mathcal{F})' + 2\phi_1^2 u' \mathcal{F} = 0, \quad (\phi^2 \partial_y \mathcal{F})' + 2\phi_1^2 (u''(u-c) - u'^2) \mathcal{F} = 0,$$

and

$$\mathcal{F}(y_c, c) = \mathcal{G}(y_c, c) = \mathcal{G}_1(y_c, c) = 0,$$
$$\left(\frac{\partial_y}{u'(y_c)} + \partial_c\right)\mathcal{G}_1(y_c, c) = 0.$$

Therefore, we obtain

(3.22)
$$\partial_c \mathcal{F}(y,c) = \frac{-2}{\phi(y,c)^2} \int_{y_c}^{y} \phi_1(y',c)^2 u'(y') \mathcal{F}(y',c) dy' = \frac{-2}{\phi(y,c)^2} T_0(\phi_1^2 u' \mathcal{F}),$$

(3.23)
$$\partial_y \mathcal{F}(y,c) = \frac{-2}{\phi(y,c)^2} T_0(\phi_1^2(u''(u-c) - u'^2)\mathcal{F}),$$

which also show that

(3.24)
$$\left(\frac{\partial_y}{u'(y_c)} + \partial_c\right)\mathcal{F} = \partial_y \mathcal{G}_1(y,c) = \frac{-2}{\phi(y,c)^2} T_0\left(\phi_1^2 \frac{a_1(y,c)}{u'(y_c)}\mathcal{F}\right),$$

where

$$a_1(y,c) = u'(y)u'(y_c) + u''(y)(u(y) - c) - u'(y)^2.$$

It is easy to see that

$$a_1(y_c, c) = 0, \quad \partial_y a_1(y, c) = u''(y)(u'(y_c) - u'(y)) + u'''(y)(u(y) - c),$$

$$\partial_c a_1(y, c) = u'(y)\frac{u''(y_c)}{u'(y_c)} - u''(y),$$

therefore,

(3.25)
$$|\partial_y a_1(y,c)| \le C|y-y_c|, \quad |a_1(y,c)| \le C|y-y_c|^2.$$

By Lemma 3.7, we know that $\partial_y \phi_1(y,c)$ has the same sign as $y - y_c$. For $y_c \leq y' \leq y$ or $y \leq y' \leq y_c$, $u'(y') \leq C \sin \frac{y+y_c}{2}$. So, we infer from (3.22) and (3.16) that

$$\begin{aligned} |\partial_{y}\mathcal{G}(y,c)| &= |\partial_{c}\mathcal{F}(y,c)| \\ &\leq \frac{C\sin\frac{y+y_{c}}{2}}{\phi(y,c)^{2}} \int_{y_{c}}^{y} \phi_{1}(y',c)\partial_{y'}\phi_{1}(y',c)dy' \\ &\leq C\frac{\phi_{1}(y,c)^{2}-1}{\phi(y,c)^{2}}\sin\frac{y+y_{c}}{2} \\ (3.26) &\leq C\frac{\phi_{1}(y,c)-1}{\phi_{1}(y,c)}\frac{1}{|y-y_{c}|^{2}\sin\frac{y+y_{c}}{2}} \leq \frac{C\min\{1,\alpha^{2}|y-y_{c}|^{2}\}}{|y-y_{c}|^{2}\sin y_{c}}, \end{aligned}$$

and by (3.24) and (3.25),

$$\begin{aligned} \left| \left(\frac{\partial_y}{u'(y_c)} + \partial_c \right) \mathcal{F} \right| &= |\partial_y \mathcal{G}_1(y, c)| \\ &\leq \frac{C|y - y_c|^2}{u'(y_c)\phi(y, c)^2} \int_{y_c}^y \phi_1(y', c) \partial_{y'} \phi_1(y', c) dy' \\ &\leq C \frac{\phi_1(y, c)^2 - 1}{u'(y_c)\phi(y, c)^2} |y - y_c|^2 \\ &\leq \frac{C \min\{1, \alpha^2 | y - y_c|^2\}}{u'(y_c)} \frac{|y - y_c|^2}{(u(y) - c)^2}, \end{aligned}$$

$$(3.27)$$

from which and $\mathcal{G}(y_c, c) = \mathcal{G}_1(y_c, c) = 0$, it follows that

(3.28)
$$|\mathcal{G}(y,c)| \le C \left| \int_{y_c}^{y} \frac{\min\{1,\alpha^2 | y' - y_c|^2\}}{|y' - y_c|^2 u'(y_c)} dy' \right| \le \frac{C\alpha \min\{1,\alpha | y - y_c|\}}{u'(y_c)},$$

and by Lemma 3.4,

$$|\mathcal{G}_1(y,c)| \le C \frac{\min\{1,\alpha^2|y-y_c|^2\}}{u'(y_c)} \left| \int_{y_c}^y \frac{|y'-y_c|^2}{(u(y')-c)^2} dy' \right| \le \frac{C\min\{1,\alpha^2|y-y_c|^2\}}{u'(y_c)^2}.$$

Thus, we deduce that

(3.29)
$$|\partial_c \phi_1| \le \frac{C\alpha\phi_1}{u'(y_c)} \min(1, \alpha|y - y_c|), \quad |\partial_y \partial_c \phi_1| \le \frac{C\alpha^2\phi_1}{u'(y_c)},$$

and

(3.30)
$$\left| \left(\frac{\partial_y}{u'(y_c)} + \partial_c \right) \phi_1(y,c) \right| \le C \frac{\min\{1, \alpha^2 | y - y_c |^2\} \phi_1}{u'(y_c)^2}$$

Step 3. Estimates of $\partial_c^2 \phi_1$ and $\left(\frac{\partial_y}{u'(y_c)} + \partial_c\right)^2 \phi_1$. By Lemma 3.7, we get for $|y - y_c| \le \frac{M_0}{\alpha}$,

$$\begin{aligned} |\partial_c^2 \mathcal{F}(y,c)| &\leq \frac{C|\partial_c^2 \partial_y \phi_1|}{\phi_1} + \frac{C|\partial_c \partial_y \phi_1 \partial_c \phi_1|}{\phi_1^2} + \frac{C|\partial_c^2 \phi_1 \partial_y \phi_1|}{\phi_1^2} + \frac{C|\partial_c \phi_1|^2 |\partial_y \phi_1|}{\phi_1^3} \\ &\leq \frac{C\alpha^3}{u'(y_c)^2} + \frac{C\alpha^2}{u'(y_c)\sin^2\frac{y+y_c}{2}}. \end{aligned}$$

It follows from (3.22) that

$$\partial_c^2 \mathcal{F}(y,c) = \partial_y \partial_c \mathcal{G}(y,c) = -2 \int_{y_c}^y \partial_c \Big(\frac{\phi_1(z,c)^2}{\phi(y,c)^2} \mathcal{F}(z,c) \Big) u'(z) dz.$$

Then by (3.29), we have

$$\begin{aligned} |\partial_c^2 \mathcal{F}(y,c)| &\leq 4 \frac{|\partial_c \phi(y,c)|}{|\phi(y,c)|^3} \sin(\frac{y+y_c}{2}) \int_{y_c}^y \phi_1(z,c) \partial_z \phi_1(z,c) dz \\ &+ \frac{C\alpha}{|\phi(y,c)|^2 u'(y_c)} \sin(\frac{y+y_c}{2}) \int_{y_c}^y \phi_1(z,c) \partial_z \phi_1(z,c) dz \\ &+ \frac{C\alpha}{(u(y)-c)^2} \sin(\frac{y+y_c}{2}) \Big| \int_{y_c}^y |\partial_c \mathcal{F}(z,c)| dz \Big| \end{aligned}$$

which along with the fact $\int_{y_c}^{y} 2\phi_1(z,c)\partial_z \phi_1(z,c)dz = \phi_1(y,c)^2 - 1$, (3.16) and (3.26) implies that for $|y - y_c| \ge \frac{M_0}{\alpha}$,

$$|\partial_c^2 \mathcal{F}(y,c)| \le \frac{C\alpha}{|y-y_c|^2 u'(y_c)^2}$$

Thanks to $\partial_c \mathcal{G}(y_c,c) = \partial_c^2 \phi_1(y_c,c) = \alpha^2 \partial_c^2 T \phi_1 = \frac{\alpha^2}{3u'(y_c)^2}$, we get

$$\left|\partial_{c}\mathcal{G}(y,c)\right| \leq \left|\partial_{c}\mathcal{G}(y_{c},c)\right| + \left|\int_{y_{c}}^{y} \partial_{g}\partial_{c}\mathcal{G}(z,c)dz\right| \leq \frac{C\alpha^{2}}{u'(y_{c})^{2}}$$

which implies that $|\partial_c^2 \phi_1| \leq \frac{C\alpha^2 \phi_1}{u'(y_c)^2}$. Using (3.24) and the formula

$$\left(\frac{\partial_y}{u'(y_c)} + \partial_c\right)(T_0 a) = \frac{a(y,c)}{u'(y_c)} - \frac{a(y_c,c)}{u'(y_c)} + T_0(\partial_c a) = T_0\left(\left(\frac{\partial_y}{u'(y_c)} + \partial_c\right)a\right),$$

we deduce that

(3.31)
$$\left(\frac{\partial_y}{u'(y_c)} + \partial_c \right) \partial_y \mathcal{G}_1(y,c) = \left(\left(\frac{\partial_y}{u'(y_c)} + \partial_c \right) \frac{-2}{\phi(y,c)^2} \right) T_0 \left(\phi_1^2 \frac{a_1(y,c)}{u'(y_c)} \mathcal{F} \right) + \frac{-2}{\phi(y,c)^2} T_0 \left(\left(\frac{\partial_y}{u'(y_c)} + \partial_c \right) \left(\phi_1^2 \frac{a_1(y,c)}{u'(y_c)} \mathcal{F} \right) \right).$$

Using (3.30) and the fact that

(3.32)
$$\left| \left(\frac{\partial_y}{u'(y_c)} + \partial_c \right) (u(y) - c) \right| = \left| \frac{u'(y)}{u'(y_c)} - 1 \right| \le C \frac{|y - y_c|}{u'(y_c)} \le C \frac{|u(y) - c|}{u'(y_c)^2},$$

we infer that

(3.33)
$$\left| \left(\frac{\partial_y}{u'(y_c)} + \partial_c \right) \frac{-2}{\phi(y,c)^2} \right| \le \frac{C}{\phi(y,c)^2 u'(y_c)^2}.$$

On the other hand, we have

$$\begin{pmatrix} \frac{\partial_y}{u'(y_c)} + \partial_c \end{pmatrix} \left(\phi_1^2 \frac{a_1(y,c)}{u'(y_c)} \mathcal{F} \right)$$

= $2\phi_1 \frac{a_1(y,c)}{u'(y_c)} \mathcal{F} \left(\frac{\partial_y}{u'(y_c)} + \partial_c \right) \phi_1 + \phi_1^2 \frac{a_2(y,c)}{u'(y_c)^2} \mathcal{F}$
 $- \phi_1^2 \frac{a_1(y,c)}{u'(y_c)^3} u''(y_c) \mathcal{F} + \phi_1^2 \frac{a_1(y,c)}{u'(y_c)} \left(\frac{\partial_y}{u'(y_c)} + \partial_c \right) \mathcal{F},$

where

$$a_2(y,c) = u'(y)(u''(y_c) - u''(y)) + u'''(y)(u(y) - c) = 0.$$

Then we get by (3.25), (3.30) and (3.27) that

$$\begin{split} & \left| \left(\frac{\partial_y}{u'(y_c)} + \partial_c \right) \left(\phi_1^2 \frac{a_1(y,c)}{u'(y_c)} \mathcal{F} \right) \right| \\ & \leq C \phi_1^2 \frac{|y - y_c|^2}{u'(y_c)^3} |\mathcal{F}| + C \phi_1^2 \frac{|y - y_c|^2}{u'(y_c)} \frac{\phi_1(y,c)^2 - 1}{u'(y_c)\phi_1(y,c)^2} \frac{|y - y_c|^2}{(u(y) - c)^2} \\ & \leq C \phi_1^2 \frac{|y - y_c|^2}{u'(y_c)^3} |\mathcal{F}| + C |y - y_c| \frac{\phi_1(y,c)^2 - 1}{u'(y_c)^3}, \end{split}$$

which gives

(3.34)
$$\begin{aligned} \left| T_0 \left(\left(\frac{\partial_y}{u'(y_c)} + \partial_c \right) \left(\phi_1^2 \frac{a_1(y,c)}{u'(y_c)} \mathcal{F} \right) \right) \right| \\ &\leq C \frac{|y - y_c|^2}{u'(y_c)^3} |T_0(\phi_1^2|\mathcal{F}|)| + C \left| T_0 \left(|y - y_c| \frac{\phi_1(y,c)^2 - 1}{u'(y_c)^3} \right) \right| \\ &\leq C |y - y_c|^2 \frac{\phi_1(y,c)^2 - 1}{u'(y_c)^3}. \end{aligned}$$

It follows from (3.31), (3.33) and (3.34) that

$$\left| \left(\frac{\partial_y}{u'(y_c)} + \partial_c \right) \partial_y \mathcal{G}_1(y, c) \right|$$

 $\leq C \frac{|\partial_y \mathcal{G}_1(y, c)|}{u'(y_c)^2} + C \frac{|y - y_c|^2}{\phi(y, c)^2} \frac{\phi_1(y, c)^2 - 1}{u'(y_c)^3}$
 $\leq C \frac{\phi_1(y, c)^2 - 1}{u'(y_c)^3 \phi_1(y, c)^2} \frac{|y - y_c|^2}{(u(y) - c)^2}.$

This together with $\left(\frac{\partial_y}{u'(y_c)} + \partial_c\right) \mathcal{G}_1(y_c, c) = 0$ gives

$$\left| \left(\frac{\partial_y}{u'(y_c)} + \partial_c \right) \mathcal{G}_1(y,c) \right| \le C \frac{\phi_1(y,c)^2 - 1}{u'(y_c)^3 \phi_1(y,c)^2} \left| \int_{y_c}^y \frac{|y' - y_c|^2}{(u(y') - c)^2} dy' \right| \le \frac{C(\phi_1(y,c)^2 - 1)}{u'(y_c)^4 \phi_1(y,c)^2},$$

from which, (3.30) and (3.16), we infer that

$$\left| \left(\frac{\partial_y}{u'(y_c)} + \partial_c \right)^2 \phi_1(y,c) \right| \le C \frac{\min\{1, \alpha^2 | y - y_c |^2\} \phi_1}{u'(y_c)^4}.$$

The proposition follows by collecting the estimates in Step 1-Step 3.

It is easy to see from the proof of Proposition 3.8 that

Lemma 3.9. We have following estimates for \mathcal{F} and \mathcal{G} at $y = 0, \pi$:

$$\begin{aligned} C^{-1} \alpha \min\{\alpha y_{c}, 1\} &\leq |\mathcal{F}(0, c)| \leq C \alpha \min\{\alpha y_{c}, 1\}, \\ C^{-1} \alpha \min\{\alpha(\pi - y_{c}), 1\} \leq |\mathcal{F}(\pi, c)| \leq C \alpha \min\{\alpha(\pi - y_{c}), 1\}, \\ |\partial_{c} \mathcal{F}(0, c)| &\leq \frac{C|\mathcal{F}(0, c)|^{2}}{\alpha^{2} y_{c}^{2} \sin y_{c}}, \quad |\partial_{c}^{2} \mathcal{F}(0, c)| \leq \frac{C \alpha \min\{\alpha y_{c}, 1\}}{y_{c}^{2} \sin^{2} y_{c}}, \\ |\partial_{c} \mathcal{F}(\pi, c)| &\leq \frac{C|\mathcal{F}(\pi, c)|^{2}}{\alpha^{2} (\pi - y_{c})^{2} \sin y_{c}}, \quad |\partial_{c}^{2} \mathcal{F}(\pi, c)| \leq \frac{C \alpha \min\{\alpha(\pi - y_{c}), 1\}}{(\pi - y_{c})^{2} \sin^{2} y_{c}} \\ |\mathcal{G}(0, c)| &\leq \frac{C \alpha \min\{1, \alpha y_{c}\}}{\sin y_{c}}, \quad |\partial_{c} \mathcal{G}(0, c)| \leq \frac{C \alpha^{2}}{\sin^{2} y_{c}}, \\ |\mathcal{G}(\pi, c)| &\leq \frac{C \alpha \min\{1, \alpha(\pi - y_{c})\}}{\sin y_{c}}, \quad |\partial_{c} \mathcal{G}(\pi, c)| \leq \frac{C \alpha^{2}}{\sin^{2} y_{c}}. \end{aligned}$$

To obtain better estimates near critical points, we introduce the following functions: for $0 \le y_c < \frac{1}{\alpha}$ and $0 \le y \le 1$, let

$$\widetilde{\phi_1}(y,c) = \phi_1(yy_c,c),$$

and for $\pi - \frac{1}{\alpha} < y_c \le \pi$ and $0 \le y \le 1$, let

$$\widetilde{\phi_1}(y,c) = \phi_1((1-y)\pi + yy_c,c).$$

Proposition 3.10. It holds that for any $y \in [0,1]$ and $0 \le y_c < \frac{1}{\alpha}$,

$$1 \leq \widetilde{\phi_1} \leq 2, \quad |\partial_c \widetilde{\phi_1}| \leq C\alpha^2, \quad |\partial_c^2 \widetilde{\phi_1}| \leq C\alpha^4, \\ \left|\partial_c^k \left(\frac{\partial_y \widetilde{\phi_1}(y,c)}{y_c^2}\right)\right| \leq C\alpha^{2+2k}, \quad k = 0, 1, 2,$$

and for any $y \in [0,1]$ and $\pi - \frac{1}{\alpha} < y_c \leq \pi$, it holds that

$$1 \le \widetilde{\widetilde{\phi_1}} \le 2, \quad |\partial_c \widetilde{\widetilde{\phi_1}}| \le C\alpha^2, \quad |\partial_c^2 \widetilde{\widetilde{\phi_1}}| \le C\alpha^4, \\ \left|\partial_c^k \left(\frac{\partial_y \widetilde{\widetilde{\phi_1}}(y,c)}{(\pi - y_c)^2}\right)\right| \le C\alpha^{2+2k}, \quad k = 0, 1, 2.$$

Proof. For $0 \le y_c < \frac{1}{\alpha}$ and $0 \le y \le 1$, by (3.15), we have $\widetilde{\phi_1} = 1 + \alpha^2 T_1 \widetilde{\phi_1}$, where

$$(T_1f)(y,c) = \int_1^y \frac{y_c^2}{(u(y'y_c) - c)^2} \int_1^{y'} f(z,c)(u(zy_c) - c)^2 dz dy'$$
$$= \int_1^y \int_1^{y'} f(z,c) \frac{F(z,y_c)^2}{F(y',y_c)^2} y_c^2 dz dy'$$

with

$$F(z,c) = \frac{u(zy_c) - c}{y_c^2} = \frac{\cos y_c - \cos zy_c}{y_c^2}$$

$$=\frac{1-z^2}{2}m_1\left(\frac{(1-z)y_c}{2}\right)m_1\left(\frac{(1+z)y_c}{2}\right),$$

and $m_1(y) = \sin y/y \in C^4([-1,1])$. One can check that $1/C \leq m_1(-y) = m_1(y) \leq C$, and $|\partial_c^k m_1(ay_c)| \leq C$ for $y_c < \frac{1}{\alpha}, k = 1, 2$. Then we can deduce that for $y_c < \frac{1}{\alpha}$,

$$\frac{|z^2 - 1|}{C} \le |F(z, c)| \le C|z^2 - 1|, \ |\partial_c F(z, c)| \le C|z^4 - 1|, \ |\partial_c^2 F(z, c)| \le C|z^4 - 1|,$$

and for $y_c < \frac{1}{\alpha}$ and 0 < y' < z < 1,

$$0 \le \frac{F(z, y_c)^2}{F(y', y_c)^2} \le C, \ \left| \partial_c \left(\frac{F(z, y_c)^2}{F(y', y_c)^2} \right) \right| \le C, \ \left| \partial_c^2 \left(\frac{F(z, y_c)^2}{F(y', y_c)^2} \right) \right| \le C.$$

Let $\Omega_{\alpha} = \{(y,c): y \in [0,1], c \in [u(0), u(1/\alpha)]\}$. Then we can infer that

(3.35)
$$||T_1f||_{L^{\infty}(\Omega_{\alpha})} \leq \frac{1}{2\alpha^2} ||f||_{L^{\infty}(\Omega_{\alpha})},$$

(3.36)
$$\|\partial_c T_1 f - T_1 \partial_c f\|_{L^{\infty}(\Omega_{\alpha})} \le C \|f\|_{L^{\infty}(\Omega_{\alpha})},$$

(3.37)
$$\|\partial_c^2 T_1 f - T_1 \partial_c^2 f\|_{L^{\infty}(\Omega_{\alpha})} \le C(\|f\|_{L^{\infty}(\Omega_{\alpha})} + \|\partial_c f\|_{L^{\infty}(\Omega_{\alpha})})$$

We infer from (3.35) that $1 \leq \widetilde{\phi_1} \leq 2$ for $(y,c) \in \Omega_{\alpha}$. By Proposition 3.8, we have for $(y,c) \in \Omega_{\alpha}$,

$$|\partial_c \widetilde{\phi_1}| = \left| (\partial_c \phi_1)(yy_c, c) + \frac{y}{u'(y_c)} (\partial_y \phi_1)(yy_c, c) \right| \le \frac{C \alpha \widetilde{\phi_1}}{u'(y_c)} \le \frac{C \alpha}{y_c}.$$

Thus, we obtain

$$||T_1(\partial_c \phi_1)||_{L^{\infty}(\Omega_{\alpha})} \le C,$$

which along with (3.36) shows that for $(y, c) \in \Omega_{\alpha}$,

$$(3.38) |\partial_c \widetilde{\phi_1}| \le C\alpha^2.$$

By Proposition 3.8, we have for $(y, c) \in \Omega_{\alpha}$,

$$\partial_c^2 \widetilde{\phi_1}| \le \frac{C \alpha^2 \widetilde{\phi_1}}{u'(y_c)^2} \le \frac{C \alpha^2}{y_c^2}.$$

Thus, we have

$$||T_1(\partial_c^2 \widetilde{\phi_1})||_{L^{\infty}(\Omega_{\alpha})} \le C\alpha^2,$$

which along with (3.36) shows that for $(y, c) \in \Omega_{\alpha}$, (3.39) $|\partial_c^2 \widetilde{\phi_1}| \le C \alpha^4$.

Using the fact that
$$\frac{\partial_y \widetilde{\phi_1}(y,c)}{y_c^2} = \alpha^2 \int_1^y \widetilde{\phi_1}(z,c) \frac{F(z,y_c)^2}{F(y,y_c)^2} dz$$
, we can deduce that for $(y,c) \in \Omega_{\alpha}$,
 $\left| \partial_c^k \left(\frac{\partial_y \widetilde{\phi_1}(y,c)}{y_c^2} \right) \right| \le C \alpha^{2+2k}, \quad k = 0, 1, 2.$

Due to $\phi_1(0,c) = \widetilde{\phi_1}(0,c)$, we have by Proposition 3.8,

(3.40)
$$-\frac{\partial_y \phi_1(0,c)}{y_c^2} \ge C^{-1} \alpha^2.$$

For $\pi - \frac{1}{\alpha} < y_c < \pi$ and $0 \le y \le 1$, we have $\widetilde{\widetilde{\phi_1}} = 1 + \alpha^2 T_2 \widetilde{\widetilde{\phi_1}}$, where

$$(T_2f)(y,c) = \int_1^y \frac{(\pi - y_c)^2}{(u((1 - y')\pi + y'y_c) - c)^2} \int_1^{y'} f(z,c)(u((1 - z)\pi + zy_c) - c)^2 dz dy',$$

and we also have

$$\frac{u((1-y')\pi + y'y_c) - c}{(\pi - y_c)^2} = \cos(\pi - y_c) - \cos(y'(\pi - y_c))$$
$$= \frac{1 - y'^2}{2} m_1 \Big(\frac{(1-y')\pi - y_c}{2}\Big) m_1 \Big(\frac{(1+y')\pi - y_c}{2}\Big).$$

Using similar arguments as above, we can deduce that for $y \in [0, 1]$ and $c \in [u(\pi - 1/\alpha), u(\pi)]$,

$$1 \le \widetilde{\widetilde{\phi_1}} \le 2, \quad |\partial_c \widetilde{\widetilde{\phi_1}}| \le C\alpha^2, \quad |\partial_c^2 \widetilde{\widetilde{\phi_1}}| \le C\alpha^4, \\ \left|\partial_c^k \left(\frac{\partial_y \widetilde{\widetilde{\phi_1}}(y,c)}{(\pi - y_c)^2}\right)\right| \le C\alpha^{2+2k}, \quad k = 0, 1, 2,$$

and

(3.41)
$$\frac{\partial_y \widetilde{\phi_1}(0,c)}{(\pi - y_c)^2} \ge C^{-1} \alpha^2.$$

4. The inhomogeneous Rayleigh equation

In this section, we solve the inhomogeneous Rayleigh equation:

(4.1)
$$\begin{cases} \Phi'' - \alpha^2 \Phi - \frac{u''}{u - c} \Phi = f, \\ \Phi'(-\pi) = \Phi'(\pi), \quad \Phi(-\pi) = \Phi(\pi), \end{cases}$$

where $u(y) = -\cos y$. We split the equation (4.1) into two parts:

(4.2)
$$\begin{cases} \Phi_o'' - \alpha^2 \Phi_o - \frac{u''}{u - c} \Phi_o = f_o, \\ \Phi_o(0) = \Phi_o(\pi) = 0, \end{cases}$$

and

(4.3)
$$\begin{cases} \Phi_e'' - \alpha^2 \Phi_e - \frac{u''}{u-c} \Phi_e = f_e, \\ \Phi_e'(0) = \Phi_e'(\pi) = 0, \end{cases}$$

where $f_o(y,c) = \frac{f(y,c)-f(-y,c)}{2}$ and $f_e(y,c) = \frac{f(y,c)+f(-y,c)}{2}$. Thus, $\Phi = \Phi_o + \Phi_e$ is a solution of (4.1) with Φ_o being an odd function and Φ_e being an even function.

Throughout this section, we denote by $\phi(y,c)$ a solution of (2.6) obtained by Proposition 3.6 and let $\phi_1(y,c) = \frac{\phi(y,c)}{u(y)-c}$ for $c \in \Omega_{\epsilon}$. Here and in what follows, we denote $\phi' = \partial_y \phi$ and $\phi'_1 = \partial_y \phi_1$.

4.1. The Wronskian. Before we present the solution formulation, let us first calculate the Wronskian of (4.2) and (4.3).

Lemma 4.1. For any $c \in \Omega_{\epsilon_0} \setminus \overline{D}_0$, the Wronskian of (4.2) has the form

$$W_o(c) = \phi(0, c)\phi(\pi, c) \int_0^{\pi} \frac{1}{\phi(y, c)^2} dy,$$

and the Wronskian of (4.3) has the form

$$W_e(c) = \frac{\partial_y \phi(\pi, c)}{\phi(0, c)} - \frac{\partial_y \phi(0, c)}{\phi(\pi, c)} - \phi'(\pi, c) \phi'(0, c) \int_0^\pi \frac{1}{\phi(y', c)^2} dy'.$$

Proof. For $c \notin \overline{D}_0$, the integration $\int_0^y \frac{1}{\phi(y',c)^2} dy'$ is well defined. It is easy to check that

$$\varphi_o(y,c) = \phi(0,c)\phi(y,c) \int_0^y \frac{1}{\phi(y',c)^2} dy',$$

is a solution to (2.6) with boundary conditions $\varphi(0,c) = 0$ and $\varphi'(0,c) = 1$. So, $W_o(c) = \varphi_o(\pi,c)$.

One can also check that

$$\varphi_e(y,c) = \frac{\phi(y,c)}{\phi(0,c)} - \phi(y,c)\partial_y\phi(0,c)\int_0^y \frac{1}{\phi(y',c)^2}dy'$$

is a solution of (2.6) with boundary conditions $\varphi(0,c) = 1$ and $\partial_y \varphi(0,c) = 0$. So, $W_e(c) = \partial_y \varphi_e(\pi,c)$.

By Proposition 3.6, $\phi(y, c)$ and $\partial_y \phi(y, c)$ are continuous. Thus, both $W_o(c)$ and $W_e(c)$ are continuous for $c \in \Omega_{\epsilon_0} \setminus \overline{D}_0$. The following lemma will show that $\sin y_c W_o(c)$ is continuous up to the boundary. To this end, we introduce some notations. For $c = u(y_c) \in D_0$ and j = 0, 1, k = 1, 2, let

(4.4)
$$\operatorname{II}(c) = \int_0^{\pi} \frac{1}{(u(y) - c)^2} \Big(\frac{1}{\phi_1(y, c)^2} - 1 \Big) dy,$$

(4.5)
$$A(c) = \sin^3 y_c II(c), \quad B(c) = \pi \cos y_c,$$

(4.6)
$$J_j^k(c) = \frac{-u'(y_c)(u(j\pi) - c)^k}{\phi_1((1-j)\pi, c)^{k-1}\phi_1'((1-j)\pi, c)}, \quad J_j(c) = J_j^2(c),$$

(4.7)
$$A_1(c) = J_1(c) - J_0(c) + \sin^2 y_c A(c), \quad B_1(c) = \sin^2 y_c B_{,c}$$

(4.8)
$$\rho(c) = (u(\pi) - c)(c - u(0)) = 1 - c^2$$

Lemma 4.2. For $c_{\epsilon} = c + i\epsilon = u(y_c) + i\epsilon \in D_{\epsilon_0}$, $c \in (-1, 1)$, we have

$$\lim_{\epsilon \to 0+} \rho(c_{\epsilon})^{1/2} W_o(c_{\epsilon}) = -\phi_1(0, c) \phi_1(\pi, c) (A(c) - iB(c)),$$

$$\lim_{\epsilon \to 0-} \rho(c_{\epsilon})^{1/2} W_o(c_{\epsilon}) = -\phi_1(0, c) \phi_1(\pi, c) (A(c) + iB(c)),$$

$$\lim_{\epsilon \to 0+} \phi(0, c_{\epsilon}) \phi(\pi, c_{\epsilon}) W_e(c_{\epsilon}) = -\frac{(\phi_1 \phi_1')(\pi, c)(\phi_1 \phi_1')(0, c)}{u'(y_c)} (A_1(c) - iB_1(c)),$$

$$\lim_{\epsilon \to 0-} \phi(0, c_{\epsilon}) \phi(\pi, c_{\epsilon}) W_e(c_{\epsilon}) = -\frac{(\phi_1 \phi_1')(\pi, c)(\phi_1 \phi_1')(0, c)}{u'(y_c)} (A_1(c) + iB_1(c)).$$

Here $\rho(c_{\epsilon}) = 1 - c_{\epsilon}^2$ and $\rho(c_{\epsilon})^{1/2}$ is taken so that $\operatorname{Re} \rho(c_{\epsilon})^{1/2} > 0$.

Proof. Let us prove the case of $\epsilon > 0$. Let $c_{\epsilon} = u(y_c) + i\epsilon \in D_{\epsilon_0}$. Then

$$\begin{split} \rho(c_{\epsilon})^{3/2} \int_0^{\pi} \frac{1}{(u(y) - c_{\epsilon})^2} dy &= \rho(c_{\epsilon})^{3/2} \partial_c \left(\int_0^{\pi} \frac{1}{u(y) - c_{\epsilon}} dy \right) \\ &= -\rho(c_{\epsilon})^{3/2} \partial_c \left(\int_0^{\pi} \frac{1}{\cos y + c_{\epsilon}} dy \right) \\ &= -\rho(c_{\epsilon})^{3/2} \partial_c \left(\int_0^{\infty} \frac{2}{1 - t^2 + c_{\epsilon}(1 + t^2)} dt \right) \\ &= -\rho(c_{\epsilon})^{3/2} \partial_c \left(\frac{1}{(1 + c_{\epsilon}) d_{\epsilon}} \ln \left(\frac{1 + d_{\epsilon}t}{1 - d_{\epsilon}t} \right) \Big|_{t=0}^{\infty} \right), \end{split}$$

here d_{ϵ} satisfies $d_{\epsilon}^2 = \frac{1-c_{\epsilon}}{1+c_{\epsilon}}$ with $\operatorname{Re} d_{\epsilon} > 0$. Then $\operatorname{Im} d_{\epsilon} < 0$, $\rho(c_{\epsilon})^{1/2} = (1+c_{\epsilon})d_{\epsilon}$ and

$$\rho(c_{\epsilon})^{3/2} \int_0^{\pi} \frac{1}{(u(y) - c_{\epsilon})^2} dy = -\rho(c_{\epsilon})^{3/2} \partial_c \left(\frac{-1}{(1 + c_{\epsilon})d_{\epsilon}} \pi i\right)$$
$$= -\partial_c \rho(c_{\epsilon})\pi i/2 \to -\cos y_c \pi i, \text{ as } \epsilon \to 0 + .$$

By Proposition 3.6 and Lemma 3.4, we get

$$\left|\frac{\rho(c_{\epsilon})}{(u(y)-c_{\epsilon})^{2}}\left(\frac{1}{\phi_{1}(y,c_{\epsilon})^{2}}-1\right)\right| \leq C \left|\frac{(\rho(c)+\epsilon)|y-y_{c}|^{2}}{(u(y)-c)^{2}+\epsilon^{2}}\right| \leq C,$$

with C a constant independent of ϵ , which ensures that as $\epsilon \to 0+$,

$$\rho(c_{\epsilon}) \int_{0}^{\pi} \frac{1}{(u(y) - c_{\epsilon})^{2}} \Big(\frac{1}{\phi_{1}(y, c_{\epsilon})^{2}} - 1 \Big) dy \to \sin^{2} y_{c} \int_{0}^{\pi} \frac{1}{(u(y) - c)^{2}} \Big(\frac{1}{\phi_{1}(y, c)^{2}} - 1 \Big) dy.$$

Thus, we deduce that as $\epsilon \to 0+,$

$$\rho(c_{\epsilon})^{3/2} \int_0^\pi \frac{1}{\phi(y, c_{\epsilon})^2} dy \to \sin^3 y_c \mathrm{II}(c) - \cos y_c \pi i.$$

This together with the continuity of $\phi_1(y,c)$ gives

$$\lim_{\epsilon \to 0+} \rho(c_{\epsilon})^{1/2} W_o(c_{\epsilon}) = -\phi_1(0, c) \phi_1(\pi, c) \big(\mathbf{A}(c) - i \mathbf{B}(c) \big).$$

Using the fact that $u'(0) = u'(\pi) = 0$, we get $\phi'(j\pi, c) = (u(j\pi) - c)\phi'_1(j\pi, c)$ for j = 0, 1. Then we infer that

$$\begin{split} \phi(0,c_{\epsilon})\phi(\pi,c_{\epsilon})W_{e}(c_{\epsilon}) &= (\phi\phi')(\pi,c_{\epsilon}) - (\phi\phi')(0,c_{\epsilon}) - (\phi\phi')(\pi,c_{\epsilon})(\phi\phi')(0,c_{\epsilon}) \int_{0}^{\pi} \frac{1}{\phi(y',c_{\epsilon})^{2}} dy' \\ &\to (\phi\phi')(\pi,c) - (\phi\phi')(0,c) - \frac{(\phi\phi')(\pi,c)(\phi\phi')(0,c)}{\sin^{3}y_{c}} (\mathbf{A}(c) - i\mathbf{B}(c)) \\ &= (1 + \cos y_{c})^{2}(\phi_{1}\phi'_{1})(\pi,c) - (1 - \cos y_{c})^{2}(\phi_{1}\phi'_{1})(0,c) \\ &- \frac{(\phi_{1}\phi'_{1})(\pi,c)(\phi_{1}\phi'_{1})(0,c)}{u'(y_{c})} \sin^{2}y_{c}(\mathbf{A}(c) - i\mathbf{B}(c)) \\ &= -\frac{(\phi_{1}\phi'_{1})(\pi,c)(\phi_{1}\phi'_{1})(0,c)}{u'(y_{c})} (\mathbf{A}_{1}(c) - i\mathbf{B}_{1}(c)). \end{split}$$

The case of $\epsilon < 0$ is similar, here we omit the details.

4.2. Uniform estimates of A(c), B(c).

Proposition 4.3. Let A(c), B(c) be defined by (4.5). Then there exists a constant C independent of α , such that

$$C^{-1}(1 + \alpha \sin y_c)^2 \le A(c)^2 + B(c)^2 \le C(1 + \alpha \sin y_c)^2,$$

$$\partial_c \left(\frac{1}{A(c)^2 + B(c)^2}\right) \le \frac{C}{(1 + \alpha \sin y_c)^2 \sin^2 y_c},$$

$$\partial_c^2 \left(\frac{1}{A(c)^2 + B(c)^2}\right) \le \frac{C}{(1 + \alpha \sin y_c)^2 \sin^4 y_c}.$$

We need the following lemma.

Lemma 4.4. Let II(c) be defined by (4.4). Then there exists a constant C independent of α , such that

$$C^{-1}\min\left\{\frac{\alpha^2}{\sin y_c}, \frac{|\alpha|}{\sin^2 y_c}\right\} \le -\mathrm{II}(c) \le C\min\left\{\frac{\alpha^2}{\sin y_c}, \frac{|\alpha|}{\sin^2 y_c}\right\},\\left|\rho(c)^k \partial_c^k \mathrm{II}(c)\right| \le C\min\left\{\frac{\alpha^2}{\sin y_c}, \frac{|\alpha|}{\sin^2 y_c}\right\}, \quad k = 1, 2.$$

Proof. Let us first prove the first inequality of the lemma. Due to $\phi_1(y,c) \ge 1$, we get by Proposition 3.8 and Lemma 3.4 that

$$\begin{split} -\mathrm{II}(c) &= \int_{0}^{\pi} \frac{1}{(u(y) - c)^{2}} \Big(1 - \frac{1}{\phi_{1}(y, c)^{2}} \Big) dy \\ &\geq C^{-1} \int_{0}^{\pi} \frac{\alpha^{2} |y - y_{c}|^{2}}{(u(y) - c)^{2}} \chi_{\{|y - y_{c}| \leq \frac{1}{\alpha}\}} dy \\ &= \begin{cases} C^{-1} \int_{0}^{y_{c}} \frac{\alpha^{2} |y - y_{c}|^{2}}{\sin^{2} \frac{y + y_{c}}{2} |y - y_{c}|^{2}} dy \geq \frac{\alpha^{2}}{Cu'(y_{c})}, \ (0 < y_{c} \leq \frac{1}{\alpha}), \\ \\ C^{-1} \int_{y_{c} - \frac{1}{\alpha}}^{y_{c}} \frac{\alpha^{2} |y - y_{c}|^{2}}{u'(y_{c})^{2} |y - y_{c}|^{2}} dy \geq \frac{\alpha}{Cu'(y_{c})^{2}}, \ (\frac{1}{\alpha} \leq y_{c} < \pi - \frac{1}{\alpha}), \\ \\ C^{-1} \int_{y_{c}}^{\pi} \frac{\alpha^{2} |y - y_{c}|^{2}}{\sin^{2} \frac{y + y_{c}}{2} |y - y_{c}|^{2}} dy \geq \frac{\alpha^{2}}{Cu'(y_{c})}, \ (\pi - \frac{1}{\alpha} < y_{c} \leq \pi). \end{split}$$

On the other hand, by Proposition 3.8 and Lemma 3.4 again, we have

$$-\mathrm{II}(c) = \int_0^\pi \frac{1}{(u(y) - c)^2} \left(1 - \frac{1}{\phi_1(y, c)^2}\right) dy$$
$$\leq C \int_0^\pi \frac{\alpha^2 |y - y_c|^2}{(u(y) - c)^2} dy \leq C \frac{\alpha^2}{u'(y_c)},$$

or

$$-\mathrm{II}(c) = \int_0^\pi \frac{1}{(u(y) - c)^2} \left(1 - \frac{1}{\phi_1(y, c)^2}\right) dy$$
$$\leq C \int_0^\pi \frac{\min\{\alpha^2 | y - y_c|^2, 1\}}{u'(y_c)^2 | y - y_c|^2} dy \leq C \frac{\alpha}{u'(y_c)^2} dy$$

Next we prove the derivative estimates of II. Direct calculations show that

$$\partial_c II(c) = \partial_c \int_0^{\pi} \frac{1}{(u(y) - c)^2} \Big(\frac{1}{\phi_1(y, c)^2} - 1 \Big) dy$$

$$= \int_0^{\pi} \left(\frac{\partial_y}{u'(y_c)} + \partial_c \right) \left(\frac{1}{(u(y) - c)^2} \left(\frac{1}{\phi_1(y, c)^2} - 1 \right) \right) dy \\ - \frac{1}{u'(y_c)(u(y) - c)^2} \left(\frac{1}{\phi_1(y, c)^2} - 1 \right) \Big|_{y=0}^{\pi}.$$

and

$$\begin{aligned} \partial_c^2 \mathrm{II}(c) &= \int_0^\pi \left(\frac{\partial_y}{u'(y_c)} + \partial_c \right)^2 \left(\frac{1}{(u(y) - c)^2} \left(\frac{1}{\phi_1(y, c)^2} - 1 \right) \right) dy \\ &- \frac{1}{u'(y_c)} \left(\frac{\partial_y}{u'(y_c)} + \partial_c \right) \left(\frac{1}{(u(y) - c)^2} \left(\frac{1}{\phi_1(y, c)^2} - 1 \right) \right) \Big|_{y=0}^\pi, \\ &- \partial_c \left(\frac{1}{u'(y_c)(u(y) - c)^2} \left(\frac{1}{\phi_1(y, c)^2} - 1 \right) \Big|_{y=0}^\pi \right). \end{aligned}$$

By (3.32) and Proposition 3.8, we get

$$\left| \left(\frac{\partial_y}{u'(y_c)} + \partial_c \right) \left(\frac{1}{(u(y) - c)^2} \left(\frac{1}{\phi_1(y, c)^2} - 1 \right) \right) \right| \le \frac{C \min\{\alpha^2 | y - y_c |^2, 1\}}{u'(y_c)^2 (u(y) - c)^2},$$

and

$$\left| \left(\frac{\partial_y}{u'(y_c)} + \partial_c \right)^2 \left(\frac{1}{(u(y) - c)^2} \left(\frac{1}{\phi_1(y, c)^2} - 1 \right) \right) \right| \le \frac{C \min\{\alpha^2 | y - y_c |^2, 1\}}{u'(y_c)^4 (u(y) - c)^2}.$$

Here we used the fact that

$$\left| \left(\frac{\partial_y}{u'(y_c)} + \partial_c \right)^2 (u(y) - c) \right| = \frac{|u''(y)u'(y_c) - u''(y_c)u'(y)|}{u'(y_c)^3} = \frac{\sin|y - y_c|}{\sin^3 y_c} \le \frac{C|u(y) - c|}{\sin^4 y_c}.$$

Using Proposition 3.6 and the facts that $\rho(c) \leq |u(y) - c|$ and $|u(y) - c| \sim |y - y_c|^2 \geq C^{-1}u'(y_c)^2$ for $y = 0, \pi$, we deduce that for $y = 0, \pi$:

$$\left|\frac{1}{u'(y_c)(u(y)-c)^2} \left(\frac{1}{\phi_1(y,c)^2} - 1\right)\right| \le C \frac{\min\{\alpha^2 | y - y_c|^2, 1\}}{u'(y_c)(u(y)-c)^2} \le C \frac{\min\{\alpha^2, \alpha/u'(y_c)\}}{u'(y_c)\rho(c)},$$

and

$$\begin{split} & \left| \partial_c \left(\frac{1}{u'(y_c)(u(y) - c)^2} \left(\frac{1}{\phi_1(y, c)^2} - 1 \right) \right) \right| \\ & \leq C \left(\frac{1}{u'(y_c)^3(u(y) - c)^2} + \frac{1}{u'(y_c)|u(y) - c|^3} \right) \left| \frac{1}{\phi_1(y, c)^2} - 1 \right| \\ & + 2 \frac{1}{u'(y_c)(u(y) - c)^2} \frac{|\mathcal{G}(y, c)|}{\phi_1(y, c)^2} \\ & \leq C \left(\frac{\min\{\alpha^2 | y - y_c|^2, 1\}}{u'(y_c)^3(u(y) - c)^2} + \frac{\min\{\alpha^2 | y - y_c|^2, 1\}}{u'(y_c)|u(y) - c|^3} + \frac{\alpha \min\{\alpha | y - y_c|, 1\}}{u'(y_c)^2(u(y) - c)^2\phi_1(y, c)^2} \right) \\ & \leq C \left(\frac{\min\{\alpha^2 | y - y_c|^2, 1\}}{u'(y_c)|u(y) - c|\rho(c)^2} + \frac{\alpha \min\{\alpha | y - y_c|, 1\}}{u'(y_c)^2(u(y) - c)^2\phi_1(y, c)^2} \right) \\ & \leq \frac{C}{\rho(c)^2} \min\left\{ \frac{\alpha^2}{u'(y_c)}, \frac{\alpha}{u'(y_c)^2} \right\}, \end{split}$$

here $\mathcal{G}(y,c) = \frac{\partial_c \phi_1}{\phi_1}(y,c)$. With the estimates above and Lemma 3.4, we can deduce the high order derivative estimates of II(c).

Let us begin with the proof of Proposition 4.3.

Proof. It follow from Lemma 4.4 that

$$A(c)^{2} + B(c)^{2} \ge C^{-1} (2\min\{\alpha^{2}\sin^{2}y_{c}, \alpha^{4}\sin^{4}y_{c}\} + \cos^{2}y_{c}) \ge C^{-1} (1 + \alpha\sin y_{c})^{2},$$

and the upper bound of $A(c)^2 + B(c)^2$.

By Lemma 4.4 again, we have

$$\begin{aligned} |\mathbf{A}(c)| &= |\sin^3 y_c \mathbf{II}(c)| \le C\alpha \sin y_c, \\ |\partial_c \mathbf{A}(c)| &= |\partial_c (\sin^3 y_c \mathbf{II}(c))| \le C \min\{\alpha^2, \alpha/\sin y_c\}, \\ |\partial_c^2 \mathbf{A}(c)| &= |\partial_c^2 (u'(y_c)\rho \mathbf{II}(c))| \le C \min\{\alpha^2/\sin^2 y_c, \alpha/\sin^3 y_c\}, \end{aligned}$$

which together with the fact $\partial_c B(c) = -\pi$ and $\partial_c^2 B(c) = 0$ show that

$$\begin{aligned} \left| \partial_c \left(\frac{1}{\mathbf{A}(c)^2 + \mathbf{B}(c)^2} \right) \right| &\leq \frac{C}{(1 + \alpha \sin y_c)^2 \sin^2 y_c}, \\ \left| \partial_c^2 \left(\frac{1}{\mathbf{A}(c)^2 + \mathbf{B}(c)^2} \right) \right| &\leq \frac{C}{(\mathbf{A}(c)^2 + \mathbf{B}(c)^2) \sin^4 y_c} + \frac{C |\partial_c \mathbf{A}(c)|^2}{(\mathbf{A}(c)^2 + \mathbf{B}(c)^2)^2} + \frac{C |\mathbf{A}(c)\partial_c^2 \mathbf{A}(c)|}{(\mathbf{A}(c)^2 + \mathbf{B}(c)^2)^2} \\ &\leq \frac{C}{(1 + \alpha \sin y_c)^2 \sin^4 y_c} + \frac{C (|\alpha/\sin y_c|^2)}{(1 + \alpha \sin y_c)^4} + \frac{C \alpha^2 / \sin^2 y_c}{(1 + \alpha \sin y_c)^4} \\ &\leq \frac{C}{(1 + \alpha \sin y_c)^2 \sin^4 y_c}. \end{aligned}$$

4.3. Uniform estimates of $A_1(c), B_1(c)$.

Proposition 4.5. Let $A_1(c), B_1(c)$ be defined by (4.7). Then there exists a constant C independent of α , such that

$$\frac{(1+\alpha\sin y_c)^6}{C\alpha^4} \le A_1(c)^2 + B_1(c)^2 \le \frac{C(1+\alpha\sin y_c)^6}{\alpha^4},\\ \left|\partial_c \left(\frac{1}{A_1(c)^2 + B_1(c)^2}\right)\right| \le \frac{C\alpha^6}{(1+\alpha\sin y_c)^8},\\ \left|\partial_c^2 \left(\frac{1}{A_1(c)^2 + B_1(c)^2}\right)\right| \le \frac{C\alpha^8}{(1+\alpha\sin y_c)^{10}}.$$

To prove Proposition 4.5, we need a criterion on embedding eigenvalues.

Definition 4.6. We say that $c \in Ran u$ is an embedding eigenvalue of \mathcal{R}_{α} if there exists $0 \neq \psi \in H^1(\mathbb{T})$ such that for all $\varphi \in H^1(\mathbb{T})$,

(4.9)
$$\int_{\mathbb{T}} (\psi'\varphi' + \alpha^2\psi\varphi)dy + p.v. \int_{\mathbb{T}} \frac{u''\psi\varphi}{u-c}dy + i\pi \sum_{y \in u^{-1}\{c\}, u'(y) \neq 0} \frac{(u''\psi\varphi)(y)}{|u'(y)|} = 0.$$

For $u(y) = -\cos y$, one can check that if (4.9) holds, then u''(y) = 0 for some point where c = u(y) (see Lemma 5.2 in [35]), which means that c = 0. Thus, $\varphi \in H^2(\mathbb{T})$ is a classical solution to

(4.10)
$$-\psi'' + \alpha^2 \psi + \frac{u'' \psi}{u - c} = 0 \Leftrightarrow -\psi'' + (\alpha^2 - 1)\psi = 0,$$

which implies that $|\alpha| = 0$ or 1. On the torus \mathbb{T}^2_{δ} with $0 < \delta < 1$, we have $\alpha \delta \in \mathbb{Z}$. If $\alpha \neq 0$, then $|\alpha| > 1$, which implies that \mathcal{R}_{α} has no embedding eigenvalues.

Proposition 4.7. If $A_1(c)^2 + B_1(c)^2 = 0$, then $c \in D_0$ is an embedding eigenvalue of \mathcal{R}_{α} . Thus, if $|\alpha| > 1$, then $A_1(c)^2 + B_1(c)^2 > 0$.

Proof. If $A_1(c)^2 + B_1(c)^2 = 0$, then $B_1(c) = 0$. As $B_1(c) = \sin^2 y_c B(c) = (1-c^2)(-\pi c)$, we have c = 0 or ± 1 . Due to $\sin y_c^2 A(c) = 0$ at $c = \pm 1$, using $|J_j(c)| = \frac{-u'(y_c)(u(j\pi) - c)^2}{\phi_1((1-j)\pi, c)^2 \mathcal{F}((1-j)\pi, c)}$ and Lemma 3.9, we find that

$$J_1(c) \neq 0, \ J_0(c) = 0 \quad \text{for } c = -1,$$

 $J_1(c) = 0, \ J_0(c) \neq 0 \quad \text{for } c = 1,$

here we define a function at $c = \pm 1$ as its limit as $c \to \pm 1$. Therefore, $A_1(c) \neq 0$ for $c = \pm 1$, and c = 0.

Now we may assume that c = 0, $A_1(c) = 0$, then $y_c = \pi/2$. Let

$$\phi_e(y,c) = \phi(\pi,c)\phi'_1(\pi,c)\widetilde{\varphi}(y,c) - \phi(y,c),$$

where $\widetilde{\varphi}(y,c)$ is given by

$$\begin{split} \widetilde{\varphi}(y,c) &= \frac{\phi_1(y,c)}{u'(y_c)} (u(y) - u(\pi)) \\ &+ \frac{\phi_1(y,c)}{u'(y_c)} (u(y) - c) (u(\pi) - c) \int_{\pi}^{y} \frac{u'(y_c) - u'(z)}{(u(z) - c)^2} dz \\ &+ \phi_1(y,c) (u(y) - c) (u(\pi) - c) \int_{\pi}^{y} \frac{1}{(u(z) - c)^2} \Big(\frac{1}{\phi_1(z,c)^2} - 1 \Big) dz. \end{split}$$

Then $\phi_e(y,c)$ satisfies the homogeneous Rayleigh equation for $y \in (0,\pi)$. Moreover,

$$\begin{aligned} \partial_y \phi_e(0,c) &= \frac{\phi(\pi,c)\phi_1'(\pi,c)\phi_1'(0,c)}{u'(y_c)} (u(0) - u(\pi)) \\ &+ \frac{\phi(\pi,c)\phi_1'(\pi,c)\phi_1'(0,c)}{u'(y_c)} (u(0) - c)(u(\pi) - c) \int_{\pi}^0 \frac{u'(y_c) - u'(z)}{(u(z) - c)^2} dz \\ &+ \phi(\pi,c)\phi_1'(\pi,c)\phi_1'(0,c)(u(0) - c)(u(\pi) - c) \int_{\pi}^0 \frac{1}{(u(z) - c)^2} \Big(\frac{1}{\phi_1(z,c)^2} - 1\Big) dz \\ &- \frac{\phi(\pi,c)\phi_1'(\pi,c)}{\phi_1(0,c)} - \phi'(0,c). \end{aligned}$$

Using the facts that $u(0) - u(\pi) = -2$, $(u(0) - c)(u(\pi) - c) = -1$, and

$$\int_{\pi}^{0} \frac{u'(y_c) - u'(z)}{(u(z) - c)^2} dz = \int_{\pi}^{0} \frac{1 - \sin z}{(\cos z)^2} dz = \int_{\pi}^{0} \frac{dz}{1 + \sin z} = \tan\left(\frac{z}{2} - \frac{\pi}{4}\right)\Big|_{\pi}^{0} = -2,$$

we deduce that

$$\partial_y \phi_e(0,c) = \phi(\pi,c) \phi_1'(\pi,c) \phi_1'(0,c) \operatorname{II}(c) - \frac{\phi(\pi,c) \phi_1'(\pi,c)}{\phi_1(0,c)} - \phi'(0,c)$$

Using the facts that u(0) - c = -1, $u(\pi) - c = 1$, $u'(0) = u'(\pi) = 0$, $u'(y_c) = \sin y_c = 1$, we infer that $\phi_1(\pi, c) = \phi(\pi, c)$, $-\phi'(0, c) = \phi'_1(0, c)$ and

$$\partial_y \phi_e(0,c) = \phi(\pi,c) \phi'_1(\pi,c) \phi'_1(0,c) (\mathrm{II}(c) + J_1(c) - J_0(c))$$

= $\phi(\pi,c) \phi'_1(\pi,c) \phi'_1(0,c) \mathrm{A}_1(c) = 0.$

It is easy to verify that $\partial_y \phi_e(\pi, c) = 0$.

Now we extend ϕ_e to be an even function with periodic 2π . Then $\phi_e \in H^2(\mathbb{T})$ satisfies (4.9). This means that c = 0 is an embedding eigenvalue of \mathcal{R}_{α} .

Lemma 4.8. Let j = 0, 1, and $J_j^k(c)$ be defined by (4.6). Then there exists a constant C independent of α , such that for k = 1, 2, m = 0, 1, 2,

$$|\partial_c^m J_{1-j}^k(c)| \le C \min \Big\{ \frac{|(1-j)\pi - y_c|^{2k+1}}{\alpha^2 |\sin y_c|^{2m} \phi_1(j\pi, c)^{k-1}}, \frac{\alpha^{2m-2}}{|(1-j)\pi - y_c|^s} \Big\},$$

where s = 1 for k = 1, m = 2, and s = 0 otherwise.

Proof. Recall that $\mathcal{F}(y,c) = \frac{\partial_y \phi_1}{\phi_1}(y,c)$ and $\mathcal{G}(y,c) = \frac{\partial_c \phi_1}{\phi_1}(y,c)$. A direct calculation gives

$$\begin{split} \frac{1}{\phi_1^{k-1}\partial_y\phi_1}(0,c) &= \frac{1}{\phi_1(0,c)^k\mathcal{F}(0,c)},\\ \partial_c\Big(\frac{1}{\phi_1^{k-1}\partial_y\phi_1}\Big)(0,c) &= -\frac{\partial_c\mathcal{F}}{\phi_1^k\mathcal{F}^2}(0,c) - \frac{k\mathcal{G}}{\phi_1^k\mathcal{F}}(0,c),\\ \partial_c^2\Big(\frac{1}{\phi_1^{k-1}\partial_y\phi_1}\Big)(0,c) &= \frac{-k\partial_c\mathcal{G} + k^2\mathcal{G}^2}{\phi_1^k\mathcal{F}}(0,c) + \frac{2k\mathcal{G}\partial_c\mathcal{F}}{\phi_1^k\mathcal{F}^2}(0,c) - \frac{\partial_c^2\mathcal{F}}{\phi_1^k\mathcal{F}^2}(0,c) + \frac{2|\partial_c\mathcal{F}|^2}{\phi_1^k\mathcal{F}^3}(0,c), \end{split}$$

from which and Lemma 3.9, we infer that

$$\left| \frac{1}{\phi_1(0,c)^{k-1}\phi_1'(0,c)} \right| \sim \frac{1 + \alpha y_c}{\alpha^2 \phi_1(0,c)^2 y_c},$$
$$\left| \partial_c \left(\frac{1}{\phi_1^{k-1}\phi_1'} \right)(0,c) \right| \leq \frac{C(1 + \alpha y_c)^2}{\alpha^2 \phi_1(0,c)^k y_c^2 \sin y_c},$$
$$\left| \partial_c^2 \left(\frac{1}{\phi_1^{k-1}\phi_1'} \right)(0,c) \right| \leq \frac{C(1 + \alpha y_c)^3}{\alpha^2 \phi_1(0,c)^k y_c^3 \sin^2 y_c}.$$

Recall that

$$J_1^k(c) = \frac{-u'(y_c)(u(\pi) - c)^k}{\phi_1(0, c)^{k-1}\phi_1'(0, c)}.$$

Thus, we have

$$\left|\frac{J_1^k}{(\pi - y_c)^{2k}}\right| \le \frac{Cu'(y_c)}{\phi_1(0,c)^k |\mathcal{F}(0,c)|} \le \frac{C\sin y_c(1 + \alpha y_c)}{\phi_1(0,c)^k \alpha^2 y_c} \le \frac{C|\pi - y_c|}{\alpha^2 \phi_1(0,c)^{k-1}},$$

here we used $\sin y_c \sim y_c(\pi - y_c)$ and $\phi_1(0, c) \geq C^{-1} e^{C^{-1} \alpha y_c}$. For m = 1, 2, we have

$$\begin{aligned} \left| \frac{\partial_c^m J_1^k}{\left(u(\pi) - c \right)^k} \right| &\leq \sum_{n=0}^m C u'(y_c)^{2n-2m+1} \left| \partial_c^n \frac{1}{\phi_1^{k-1} \phi_1'(0,c)} \right| \\ &\leq \frac{C(1 + \alpha y_c)^{m+1}}{\alpha^2 \phi_1(0,c)^k y_c u'(y_c)^{2m-1}} \leq \frac{C(\pi - y_c)}{\alpha^2 (\sin y_c)^{2m} \phi_1(0,c)^{k-1}}. \end{aligned}$$

To complete the proof, it suffices to improve the estimate for $y_c < \frac{1}{\alpha}$. In this case, thanks to $y_c \partial_y \phi_1(0, c) = \partial_y \widetilde{\phi_1}(0, c)$, we get

$$J_1^k = \frac{-u'(y_c)(u(\pi) - c)^2}{\phi_1(0, c)^{k-1}\phi_1'(0, c)} = \frac{-m_1(y_c)(u(\pi) - c)^k y_c^2}{\widetilde{\phi_1}(0, c)^{k-1}\widetilde{\phi_1}'(0, c)},$$

where $m_1(y) = \frac{\sin y}{y}$. Since $m_1 \in C^{\infty}([-1,1])$ is even, we have $|\partial_c^m m_1(y_c)| \leq C$ for $y_c < 1$, m = 0, 1, 2. Thus, for $0 < y_c < \frac{1}{\alpha}$, by Proposition 3.10 and (3.40),

$$\left|J_{1}^{k}\right| \leq C\alpha^{-2}, \quad \left|\partial_{c}J_{1}^{k}\right| \leq C, \quad \left|\partial_{c}^{2}J_{1}^{k}\right| \leq C\alpha^{2}.$$

This completes the proof of the case of j = 0. The case of j = 1 can be proved similarly. \Box

Now we are in a position to prove Proposition 4.5.

Proof. By Lemma 4.8 and Lemma 4.4, we get

$$|\mathbf{A}_{1}(c)| + |\mathbf{B}_{1}(c)| \le |J_{1}| + |J_{0}| + C\alpha \sin^{3} y_{c} + \pi |\sin^{2} y_{c} \cos y_{c}|$$
$$\le C(\frac{1}{\alpha} + \alpha \sin^{3} y_{c} + \sin^{2} y_{c}) \le C\frac{(1 + \alpha \sin y_{c})^{3}}{\alpha^{2}}.$$

Due to $\phi'_1(y,c) \leq 0$ for $y \leq y_c$, we get $J_1(c) \geq 0$. Similarly, we have $J_0(c) \leq 0$. Proposition 4.7 ensures that $|A_1(c)| + |B_1(c)| > C_{\alpha} > 0$. Thus, we only need to consider the case of the large α . Let M be a large number determined later. Then for $y_c \leq \frac{1}{M\alpha}$, we choose M large enough so that $1 \leq \phi_1(0,c) \leq 2$. Then

$$|J_1(c)| \ge \frac{u'(y_c)}{C|\mathcal{F}(0,c)|} \ge \frac{y_c}{C\alpha\min\{\alpha y_c,1\}} \ge \frac{1}{C\alpha^2}$$

here C is a constant independent of M and α . Thus, for $y_c \leq \frac{1}{M\alpha}$,

$$|\mathbf{A}_1(c)| \ge |J_1(c)| + |J_0(c)| - |\sin^2 y_c \mathbf{A}(c)| \ge \frac{1}{C\alpha^2} - C\alpha \sin^3 y_c \ge \frac{(1 + \alpha \sin y_c)^3}{C\alpha^2}.$$

For $\frac{1}{M\alpha} \le y_c \le \frac{M}{\alpha} < 1$, we have $|\mathbf{B}(c)| \ge |\mathbf{B}(0)/2| \ge 1$, then

$$|\mathbf{B}_1(c)| = |\rho(c)\mathbf{B}(c)| \ge \rho(c) \ge \frac{(1 + \alpha \sin y_c)^3}{C\alpha^2}$$

This shows that for $y_c \leq \frac{M}{\alpha} < 1$, we have

$$|\mathbf{A}_1(c)| + |\mathbf{B}_1(c)| \ge \frac{(1 + \alpha \sin y_c)^3}{C\alpha^2}.$$

Similarly, for $\pi - y_c \leq \frac{M}{\alpha} < 1$, we have

$$|\mathbf{A}_1(c)| + |\mathbf{B}_1(c)| \ge \frac{(1 + \alpha \sin y_c)^3}{C\alpha^2}$$

While, for $y_c \in [\frac{M}{\alpha}, \pi - \frac{M}{\alpha}], \ \alpha > M$, and M large enough, we have

$$|A_1(c)| \ge |\sin^2 y_c A(c)| - |J_1(c)| - |J_0(c)| \ge \frac{|\alpha \sin^3 y_c|}{C} - \frac{C}{\alpha^2} \ge \frac{1 + \alpha^3 \sin^3 y_c}{C\alpha^2}.$$

Therefore, we obtain

$$\frac{(1 + \alpha \sin y_c)^6}{C\alpha^4} \le A_1(c)^2 + B_1(c)^2 \le \frac{C(1 + \alpha \sin y_c)^6}{\alpha^4}$$

By Lemma 4.6 and Lemma 4.4, we have

$$\begin{aligned} |\mathbf{A}_1(c)| &\leq C(1+\alpha \sin y_c)^3/\alpha^2, \quad |\partial_c \mathbf{A}_1(c)| \leq C(1+\alpha \sin y_c), \\ |\partial_c^2 \mathbf{A}_1(c)| &\leq C \min\{\alpha^2, \frac{\alpha}{\sin y_c}\}, \\ |\mathbf{B}_1(c)| &\leq C \sin^2 y_c, \ |\partial_c \mathbf{B}_1(c)| \leq C, \ |\partial_c^2 \mathbf{B}_1(c)| \leq C. \end{aligned}$$

Then we can deduce that

$$\begin{aligned} \left| \partial_c \left(\frac{1}{\mathbf{A}_1(c)^2 + \mathbf{B}_1(c)^2} \right) \right| &\leq \frac{C\alpha^6}{(1 + \alpha \sin y_c)^8}, \\ \left| \partial_c^2 \left(\frac{1}{\mathbf{A}_1(c)^2 + \mathbf{B}_1(c)^2} \right) \right| \\ &\leq \frac{C}{(\mathbf{A}_1(c)^2 + \mathbf{B}_1(c)^2)^2} + \frac{C |\partial_c \mathbf{A}_1(c)|^2}{(\mathbf{A}_1(c)^2 + \mathbf{B}_1(c)^2)^2} + \frac{C |\mathbf{A}_1(c)\partial_c^2 \mathbf{A}_1(c)|}{(\mathbf{A}_1(c)^2 + \mathbf{B}_1(c)^2)^2} \\ &\leq \frac{C\alpha^8}{(1 + \alpha \sin y_c)^{12}} + \frac{C\alpha^8}{(1 + \alpha \sin y_c)^{10}} + \frac{C\alpha^8}{(1 + \alpha \sin y_c)^{10}} \leq \frac{C\alpha^8}{(1 + \alpha \sin y_c)^{10}}. \end{aligned}$$

Remark 4.9. Since the functions A, B, A_1, B_1 are continuous with respect to α , Proposition 4.3 and 4.5 are true for every $\alpha \ge c_0 > 1$ with the constant C depending only on c_0 .

4.4. Solution formula of the inhomogeneous Rayleigh equation. The fact that the Kolmogorov flow is stable for the case of $\alpha > 1$, implies that $W_o(c) \neq 0$ and $W_e(c) \neq 0$ for all c with $\text{Im } c \neq 0$.

Now we define for $y \in [0, \pi]$ and $c \notin [-1, 1]$,

$$\Phi_{o}(y,c) = \phi(y,c) \int_{0}^{y} \frac{1}{\phi(y',c)^{2}} \int_{y_{c}}^{y'} f_{o}\phi(y'',c)dy''dy' + \mu^{o}(c)\phi(y,c) \int_{0}^{y} \frac{1}{\phi(y',c)^{2}}dy'$$

$$(4.11) \qquad = \phi(y,c) \int_{\pi}^{y} \frac{1}{\phi(y',c)^{2}} \int_{y_{c}}^{y'} f_{o}\phi(y'',c)dy''dy' + \mu^{o}(c)\phi(y,c) \int_{\pi}^{y} \frac{1}{\phi(y',c)^{2}}dy',$$

where

(4.12)
$$\mu^{o}(c) = \frac{-\phi(0,c)\phi(\pi,c)\int_{0}^{\pi} \frac{1}{\phi(y',c)^{2}}\int_{y_{c}}^{y'} f_{o}\phi(y'',c)dy''dy'}{W_{o}(c)}$$

For $y \in [-\pi, 0]$, let $\Phi_o(y, c) = -\Phi_o(-y, c)$.

We define for $y \in [0, \pi]$ and $c \notin [-1, 1]$,

$$\Phi_{e}(y,c) = \phi(y,c) \int_{0}^{y} \frac{1}{\phi(y',c)^{2}} \int_{y_{c}}^{y'} f_{e}\phi(y'',c)dy''dy' + \mu_{*}^{e}(c)\phi(y,c) \int_{0}^{y} \frac{1}{\phi(y',c)^{2}}dy' + \nu_{0}^{e}(c)\phi(y,c) = \phi(y,c) \int_{\pi}^{y} \frac{1}{\phi(y',c)^{2}} \int_{y_{c}}^{y'} f_{e}\phi(y'',c)dy''dy' + \mu_{*}^{e}(c)\phi(y,c) \int_{\pi}^{y} \frac{1}{\phi(y',c)^{2}}dy' + \nu_{1}^{e}(c)\phi(y,c),$$

$$(4.13)$$

where μ^e_* and ν^e_0 , ν^e_1 are determined by

$$\int_{0}^{\pi} \frac{1}{\phi(y',c)^{2}} \int_{y_{c}}^{y'} f_{e}\phi(y'',c)dy''dy' + \mu_{*}^{e}(c) \int_{0}^{\pi} \frac{1}{\phi(y',c)^{2}}dy' + \nu_{0}^{e}(c) - \nu_{1}^{e}(c) = 0,$$

$$\nu_{j}^{e}(c)\phi(j\pi,c)\phi'(j\pi,c) + \int_{y_{c}}^{j\pi} f_{e}\phi(y'',c)dy'' + \mu_{*}^{e}(c) = 0, \quad j = 0, 1.$$

For $y \in [-\pi, 0]$, let $\Phi_e(y, c) = \Phi_e(-y, c)$. We have (4.14)

$$\nu_1^e(c) = \frac{\int_0^{\pi} f_e \phi(y',c) dy' + (\phi\phi') (0,c) \left(\int_0^{\pi} \frac{\int_{y_c}^{y_c} f_e \phi(y'',c) dy''}{\phi(y',c)^2} dy' - \int_0^{\pi} \frac{\int_{y_c}^{\pi} f_e \phi(y'',c) dy''}{\phi(y',c)^2} dy' \right)}{-\phi(0,c)\phi(\pi,c)W_e(c)}$$

$$\nu_0^e(c) = \frac{\int_0^{\pi} f_e \phi(y',c) dy' + (\phi\phi') \left(\pi,c\right) \left(\int_0^{\pi} \frac{\int_{y_c}^{y'_c} f_e \phi(y'',c) dy''}{\phi(y',c)^2} dy' - \int_0^{\pi} \frac{\int_{y_c}^0 f_e \phi(y'',c) dy''}{\phi(y',c)^2} dy'\right)}{-\phi(0,c)\phi(\pi,c)W_e(c)}$$

$$\mu_*^e(c) = \frac{(\phi\phi')(\pi,c)(\phi\phi')(0,c)\int_0^\pi \frac{\int_{y_c}^{y'} f_e\phi(y'',c)dy'}{\phi(y',c)^2}dy' - (\phi\phi')(j\pi,c)\int_{y_c}^{(1-j)\pi} f_e\phi(y'',c)dy''\big|_{j=0}^1}{\phi(0,c)\phi(\pi,c)W_e(c)}$$

Proposition 4.10. Let $c \in D_{\epsilon_0}$. Then the solution of (4.1) takes as follows

$$\Phi(y,c) = \Phi_o(y,c) + \Phi_e(y,c),$$

where Φ_o and Φ_e are defined by (4.11) and (4.13) respectively.

Proof. It is easy to check that the inhomogeneous Rayleigh equations (4.2) and (4.3) are equivalent to

(4.17)
$$\begin{cases} \left(\phi^2 \left(\frac{\Phi_o}{\phi}\right)'\right)' = f_o \phi, \\ \Phi_o(0) = \Phi_o(\pi) = 0 \end{cases}$$

and

(4.18)
$$\begin{cases} \left(\phi^2 \left(\frac{\Phi_e}{\phi}\right)'\right)' = f_e \phi, \\ \Phi'_e(0) = \Phi'_e(\pi) = 0. \end{cases}$$

As $\phi(y,c) \neq 0$ for $c \in D_{\epsilon_0}$, we conclude the proposition by integration twice, and matching the boundary conditions by using the fact that $\phi \int^y \frac{1}{\phi^2} dy'$ and ϕ are two independent solutions of the homogeneous Rayleigh equation.

For $c \in B^l_{\epsilon_0} \cup B^r_{\epsilon_0}$, Howard's semi-circle theorem tells us that $c \notin \sigma(\mathcal{R}_{\alpha})$. Thus, there exists a unique solution of (4.1).

5. The linearized Euler equations

5.1. The limiting absorption principle. In this subsection, we establish the limiting absorption principle for the inhomogeneous Rayleigh equation

(5.1)
$$(u-c)(\Phi''-\alpha^2\Phi)-u''\Phi=\omega,$$

when $c \in \left\{ z \in \mathbb{C} : 0 < \inf_{y \in \mathbb{T}} |u(y) - z| < \epsilon_0 \right\}$ for $\epsilon_0 > 0$ small enough.

In [35], we established the limiting absorption principle for a class of shear flows in \mathcal{K} by using blow-up analysis and compactness argument. This result can be easily extended to the following periodic shear flows (u(y), 0):

- (a) Periodic: $u(y + 2\pi) = u(y);$
- (b) Regularity: $u \in H^3(\mathbb{T})$, where $\mathbb{T} = \mathbb{T}_{2\pi}$;

- (c) Spectrum: \mathcal{R}_{α} has no embedding eigenvalues for $\alpha \neq 0$;
- (d) Curvature: $u''(y) \neq 0$ at critical points (i.e., u'(y) = 0).

We denote by \mathcal{P} the set of periodic flows satisfying (a)-(d). When $\alpha > 1$, $u(y) = -\cos y \in \mathcal{P}$.

Proposition 5.1. Assume that $u \in \mathcal{P}$. There exists ϵ_0 such that for $c \in \{z \in \mathbb{C} : 0 < \inf_{y \in \mathbb{T}} |u(y) - z| < \epsilon_0\}$, the solution to (5.1) has the following uniform bound

$$\|\Phi\|_{H^1(\mathbb{T})} \le C \|\omega\|_{H^1(\mathbb{T})}.$$

Here C is a constant independent of ϵ_0 . Moreover, there exists $\Phi_{\pm}(\alpha, y, c) \in H^1_0(\mathbb{T})$ for $c \in \operatorname{Ran} u$, such that $\Phi(\alpha, \cdot, c \pm i\epsilon) \to \Phi_{\pm}(\alpha, \cdot, c)$ in $C(\mathbb{T})$ as $\epsilon \to 0+$ and

$$\|\Phi_{\pm}(\alpha,\cdot,c)\|_{H^1(\mathbb{T})} \le C \|\omega\|_{H^1(\mathbb{T})}.$$

One can refer to section 6 in [35] for more details.

5.2. Explicit formulas of the limits. In this subsection, we give the explicit formulas of the limits Φ_{\pm} , which are important to obtain the explicit decay estimates of the velocity. In what follows, we denote by $\phi(y,c)$ a solution of (2.6) obtained by Proposition 3.6 and let $\phi_1(y,c) = \frac{\phi(y,c)}{u(y)-c}$ for $c \in \Omega_{\epsilon}$.

We will use the notations A, B, A₁, B₁ and II defined in section 4.1. Let us also introduce some new notations. We define $Int(\varphi)$ to be a 2π periodic function so that

(5.2)
$$\operatorname{Int}(\varphi)(y) = \int_0^y \varphi(y') \, dy' \quad \text{for } y \in [0,\pi], \quad \operatorname{Int}(\varphi)(y) = \operatorname{Int}(\varphi)(-y) \quad \text{for } y \in [-\pi,0].$$

We introduce

(5.3)
$$II_{1,1}(\varphi)(y_c) = p.v. \int_0^\pi \frac{\operatorname{Int}(\varphi)(y) - \operatorname{Int}(\varphi)(y_c)}{(u(y) - u(y_c))^2} \, dy.$$

(5.4)
$$II_1(\varphi)(y_c) = p.v. \int_0^\pi \frac{\int_{y_c}^y \varphi(y')\phi_1(y',c)dy'}{\phi(y,c)^2} dy,$$

(5.5)
$$\mathbf{E}_{j}(\varphi)(y_{c}) = \int_{y_{c}}^{j\pi} \varphi(y)\phi_{1}(y,c)dy \quad \text{for } j = 0, 1,$$

and for k = 0, 1, 2,

(5.6)
$$\mathcal{L}_k(\varphi)(y_c) = \int_0^\pi \int_{y_c}^z \varphi(y) \left(\frac{\partial_z + \partial_y}{u'(y_c)} + \partial_c\right)^k \left(\frac{1}{(u(z) - c)^2} \left(\frac{\phi_1(y, c)}{\phi_1(z, c)^2} - 1\right)\right) dy dz.$$

It is easy to see that

$$II_1(\varphi) = II_{1,1}(\varphi) + \mathcal{L}_0(\varphi).$$

Note. Since the map from $c \in [-1, 1]$ to $y_c \in [0, \pi]$ is one-to-one, we do not make a difference between c and y_c in some places.

Remark 5.2. We have the following formulation:

(5.7)
$$\operatorname{II}_{1,1}(\varphi)(y_c) = -\partial_c \Big(\frac{H(\operatorname{Int}(\varphi))(y_c)}{2\sin y_c}\Big),$$

where H is the Hilbert transform on \mathbb{T} defined by

$$H(\varphi)(y_c) = p.v. \int_{-\pi}^{\pi} \cot \frac{y_c - y}{2} \varphi(y) dy.$$

Indeed, we have

$$\begin{aligned} \mathrm{II}_{1,1}(\varphi) &= p.v. \int_0^\pi \frac{\mathrm{Int}(\varphi)(y) - \mathrm{Int}(\varphi)(y_c)}{(u(y) - u(y_c))^2} \, dy \\ &= \partial_c \Big(\int_0^\pi \frac{\mathrm{Int}(\varphi)(y) - \mathrm{Int}(\varphi)(y_c)}{u(y) - c} \, dy \Big) + \frac{\varphi(y_c)}{u'(y_c)} p.v. \int_0^\pi \frac{1}{u(y) - c} \, dy \\ &= \partial_c \Big(p.v. \int_0^\pi \frac{\mathrm{Int}(\varphi)(y)}{u(y) - c} \, dy \Big) - \mathrm{Int}(\varphi)(y_c) \partial_c \Big(p.v. \int_0^\pi \frac{dy}{u(y) - c} \Big). \end{aligned}$$

For an even function f, we have

$$p.v. \int_0^\pi \frac{f(y)}{u(y) - c} \, dy = \frac{1}{2} p.v. \int_{-\pi}^\pi \frac{f(y)}{u(y) - c} \, dy$$
$$= \frac{1}{2\sin y_c} p.v. \int_{-\pi}^\pi \frac{(\sin y_c + \sin y)f(y)}{\cos y_c - \cos y} \, dy = -\frac{(Hf)(y_c)}{2\sin y_c},$$

which along with the facts that H1 = 0, $p.v. \int_0^{\pi} \frac{dy}{u(y)-c} = 0$, shows (5.7).

We denote

$$\begin{split} \widehat{\omega}_o(y) &= \frac{\widehat{\omega}_0(\alpha, y) - \widehat{\omega}_0(\alpha, -y)}{2}, \quad \widehat{\omega}_e(y) = \frac{\widehat{\omega}_0(\alpha, y) + \widehat{\omega}_0(\alpha, -y)}{2}, \\ f_o(y, c) &= \frac{\widehat{\omega}_o(\alpha, y)}{i\alpha(u(y) - c)}, \quad f_e(y, c) = \frac{\widehat{\omega}_e(\alpha, y)}{i\alpha(u(y) - c)}. \end{split}$$

For $c \in (-1, 1)$, we define

$$\Phi^{o}_{\pm}(y,c) \stackrel{\text{def}}{=} \begin{cases} \phi(y,c) \int_{0}^{y} \frac{1}{\phi(z,c)^{2}} \int_{y_{c}}^{z} \phi f_{o}(y',c) dy' dz \\ + \mu^{o}_{\pm}(\widehat{\omega}_{o})(c)\phi(y,c) \int_{0}^{y} \frac{1}{\phi(y',c)^{2}} dy' & 0 \leq y < y_{c}, \\ \phi(y,c) \int_{\pi}^{y} \frac{1}{\phi(z,c)^{2}} \int_{y_{c}}^{z} \phi f_{o}(y',c) dy' dz \\ + \mu^{o}_{\pm}(\widehat{\omega}_{o})(c)\phi(y,c) \int_{\pi}^{y} \frac{1}{\phi(y',c)^{2}} dy' & y_{c} < y \leq \pi, \end{cases}$$

where

$$(5.8) \quad \mu_{+}^{o}(\widehat{\omega}_{o})(c) = \frac{1}{\alpha} \frac{i \sin^{3} y_{c} \mathrm{II}_{1}(\widehat{\omega}_{o})(y_{c}) - \sin y_{c} \widehat{\omega}_{o}(\alpha, y_{c})\pi}{-i\pi \cos y_{c} + \sin^{3} y_{c} \mathrm{II}(c)} = \frac{1}{\alpha} \frac{-\mathrm{C}_{o}(\widehat{\omega}_{o})(c) + i\mathrm{D}_{o}(\widehat{\omega}_{o})(c)}{\mathrm{A}(c) - i\mathrm{B}(c)},$$

$$(5.9) \quad \mu_{-}^{o}(\widehat{\omega}_{o})(c) = \frac{1}{\alpha} \frac{i \sin^{3} y_{c} \mathrm{II}_{1}(\widehat{\omega}_{o})(y_{c}) + \sin y_{c} \widehat{\omega}_{o}(\alpha, y_{c})\pi}{i\pi \cos y_{c} + \sin^{3} y_{c} \mathrm{II}(c)} = \frac{1}{\alpha} \frac{\mathrm{C}_{o}(\widehat{\omega}_{o})(c) + i\mathrm{D}_{o}(\widehat{\omega}_{o})(c)}{\mathrm{A}(c) + i\mathrm{B}(c)},$$

with

(5.10)
$$C_o(\varphi)(c) = \pi \sin y_c \varphi(y_c), \quad D_o(\varphi)(c) = u'(y_c)\rho(c) II_1(\varphi)(y_c).$$

For $y \in [-\pi, 0]$, let $\Phi^o_{\pm}(y, c) = -\Phi^o_{\pm}(-y, c)$.

For $c \in (-1, 1)$, we define

$$\Phi_{\pm}^{e}(y,c) \stackrel{\text{def}}{=} \begin{cases} \phi(y,c) \int_{0}^{y} \frac{\int_{y_{c}}^{y'} \phi f_{e}(y'',c)dy''}{\phi(y',c)^{2}} dy' \\ + \mu_{*\pm}^{e}(\widehat{\omega}_{e})(c)\phi(y,c) \int_{0}^{y} \frac{1}{\phi(y',c)^{2}} dy' + \nu_{0\pm}^{e}(\widehat{\omega}_{e})(c)\phi(y,c), \ 0 \leq y < y_{c}, \\ \phi(y,c) \int_{\pi}^{y} \frac{\int_{y_{c}}^{y'} \phi f_{e}(y'',c)dy''}{\phi(y',c)^{2}} dy' \\ + \mu_{*\pm}^{e}(\widehat{\omega}_{e})(c)\phi(y,c) \int_{\pi}^{y} \frac{1}{\phi(y',c)^{2}} dy' + \nu_{1\pm}^{e}(\widehat{\omega}_{e})(c)\phi(y,c), \ y_{c} < y \leq \pi, \end{cases}$$

where

(5.11)
$$\mu_{*+}^{e}(\varphi)(c) = \frac{1}{\alpha} \frac{I(c) (i \mathbf{D}_{o}(\varphi)(c) - \mathbf{C}_{o}(\varphi)(c)) - i(\phi \phi')(j\pi, c) \mathbf{E}_{1-j}(\varphi)(c) \big|_{j=0}^{1}}{I(c) (\mathbf{A}_{1}(c) - i \mathbf{B}_{1}(c)) / \rho(c)},$$

(5.12)
$$\mu_{*-}^{e}(\varphi)(c) = \frac{1}{\alpha} \frac{I(c) (i \mathcal{D}_{o}(\varphi)(c) + \mathcal{C}_{o}(\varphi)(c)) - i(\phi \phi')(j\pi, c) \mathcal{E}_{1-j}(\varphi)(c)|_{j=0}^{1}}{I(c) (\mathcal{A}_{1}(c) + i \mathcal{B}_{1}(c)) / \rho(c)},$$

(5.13)
$$\nu_{k+}^{e}(\varphi)(c) = \frac{1}{\alpha} \frac{i \mathrm{E}(\varphi) + (\phi \phi') ((1-k)\pi, c) (\mathrm{D}_{+}(\varphi) - i \mathrm{E}_{k}(\varphi)(\mathrm{A} - i\mathrm{B})) / \sin^{3} y_{c}}{I(c)(\mathrm{A}_{1}(c) - i \mathrm{B}_{1}(c)) / \rho(c)},$$

(5.14)
$$\nu_{k-}^{e}(\varphi)(c) = \frac{1}{\alpha} \frac{i \mathbf{E}(\varphi) + (\phi \phi') ((1-k)\pi, c) (\mathbf{D}_{-}(\varphi) - i \mathbf{E}_{k}(\varphi)(\mathbf{A}+i\mathbf{B})) / \sin^{3} y_{c}}{I(c)(\mathbf{A}_{1}(c) + i \mathbf{B}_{1}(c)) / \rho(c)},$$

with

$$E(\varphi) = E_1(\varphi)(y_c) - E_0(\varphi)(y_c), \quad D_{\pm}(\varphi) = iD_o(\varphi)(c) \mp C_o(\varphi)(c),$$

and

$$I(c) = \frac{\phi_1(\pi, c)\phi'(\pi, c)\phi_1(0, c)\phi'(0, c)}{u'(y_c)}$$

= $-\sin y_c(\phi_1\phi_1')(\pi, c)(\phi_1\phi_1')(0, c) = -\frac{(\phi\phi')(\pi, c)(\phi\phi')(0, c)}{u'(y_c)\rho(c)}.$

Proposition 5.3. Let $\Phi(y,c)$ be a solution of (4.1) given by Proposition 4.10 with $f = \frac{\widehat{\omega_0}(\alpha,y)}{i\alpha(u-c)}$ for $\widehat{\omega}_0 \in H^1(\mathbb{T})$. Then it holds that for any $y \in [-\pi,\pi]$, $y_c \in (-\pi,\pi) \setminus \{0\}$ and $y \neq \pm y_c$,

$$\lim_{\epsilon \to 0+} \Phi(y, c_{\epsilon}) = \Phi_+(y, u(y_c)), \quad \lim_{\epsilon \to 0-} \Phi(y, c_{\epsilon}) = \Phi_-(y, u(y_c)),$$

where $\Phi_{\pm} = \Phi^o_{\pm} + \Phi^e_{\pm}$, and $c_{\epsilon} = c + i\epsilon \in D_{\epsilon_0}$ and $c = u(y_c)$.

The proposition is an immediate consequence of Lemma 5.5 and Lemma 5.6.

Lemma 5.4. Let $c_{\epsilon} = c + i\epsilon \in D_{\epsilon_0}$, $c \in (-1, 1)$. Then for any $\varphi \in H^1(0, \pi)$,

$$\lim_{\epsilon \to 0+} \rho(c_{\epsilon}) \int_0^{\pi} \frac{\int_{y_c}^y \varphi(y') dy'}{(u(y) - c_{\epsilon})^2} dy = \rho(c) \operatorname{II}_{1,1}(\varphi)(c) + i\pi\varphi(y_c),$$
$$\lim_{\epsilon \to 0-} \rho(c_{\epsilon}) \int_0^{\pi} \frac{\int_{y_c}^y \varphi(y') dy'}{(u(y) - c_{\epsilon})^2} dy = \rho(c) \operatorname{II}_{1,1}(\varphi)(c) - i\pi\varphi(y_c).$$

Proof. Set $g(y) = \int_{y_c}^y \varphi(y') dy' - \frac{\varphi(y_c)}{u'(y_c)} (u(y) - c)$. Then we have $\begin{aligned} \rho(c_\epsilon) \int_0^\pi \frac{\int_{y_c}^y \varphi(y') dy'}{(u(y) - c_\epsilon)^2} dy \\ &= \rho(c_\epsilon) \int_0^\pi \frac{g(y)}{(u(y) - c_\epsilon)^2} dy + \rho(c_\epsilon) \frac{\varphi(y_c)}{u'(y_c)} \int_0^\pi \frac{(u(y) - c)}{(u(y) - c_\epsilon)^2} dy. \end{aligned}$

As $\varphi \in H^1(0,\pi)$, we get

$$|g'(y)| = |\varphi(y) - u'(y)\varphi(y_c)/u'(y_c)| \le C|y - y_c|^{\frac{1}{2}}/u'(y_c),$$

which along with $g(y_c) = 0$ ensures that

$$\int_0^\pi \frac{g(y)}{(u(y) - c_\epsilon)^2} dy \longrightarrow p.v. \int_0^\pi \frac{g(y)}{(u(y) - c)^2} dy.$$

From the proof of Lemma 4.2 and $p.v. \int_0^{\pi} \frac{dy}{u(y)-c} = 0$, we deduce that as $\epsilon \to 0+$,

$$\begin{split} \rho(c_{\epsilon}) \int_{0}^{\pi} \frac{(u(y)-c)}{(u(y)-c_{\epsilon})^{2}} dy &= \rho(c_{\epsilon}) \int_{0}^{\pi} \frac{dy}{(u(y)-c_{\epsilon})} dy + i\epsilon\rho(c_{\epsilon}) \int_{0}^{\pi} \frac{dy}{(u(y)-c_{\epsilon})^{2}} \\ &= \rho(c_{\epsilon}) \frac{\pi i}{\rho(c_{\epsilon})^{\frac{1}{2}}} - i\epsilon \frac{\partial_{c}\rho(c_{\epsilon})\pi i}{2\rho(c_{\epsilon})^{\frac{1}{2}}} \\ &\longrightarrow \pi i\rho(c)^{\frac{1}{2}} = p.v. \int_{0}^{\pi} \frac{(u(y)-c)}{(u(y)-c)^{2}} dy + i\pi u'(y_{c}). \end{split}$$

This shows that as $\epsilon \to 0+$,

$$\rho(c_{\epsilon}) \int_{0}^{\pi} \frac{\int_{y_{c}}^{y} \varphi(y') dy'}{(u(y) - c_{\epsilon})^{2}} dy \longrightarrow \rho(c) p.v. \int_{0}^{\pi} \frac{g(y) + \frac{\varphi(y_{c})}{u'(y_{c})} (u(y) - c)}{(u(y) - c)^{2}} dy + i\pi u'(y_{c}) \frac{\varphi(y_{c})}{u'(y_{c})}$$
$$= \rho(c) \mathrm{II}_{1,1}(\varphi)(c) + i\pi \varphi(y_{c}).$$

The case of $\epsilon \to 0-$ can be proved similarly.

Lemma 5.5. Let $k \in \{0,1\}$ and $\widehat{\omega}_0(\alpha, y) \in H^1(0, \pi)$. Let μ^o , ν^e_k and μ^e_* be defined by (4.12), (4.14), (4.15) and (4.16). Let μ^o_{\pm} , $\mu^e_{\pm\pm}$ and $\nu^e_{k\pm}$ be defined by (5.8), (5.9) and (5.11)-(5.14). Then it holds that for any $y_c \in (0, \pi)$,

$$\begin{split} &\lim_{\epsilon \to 0+} \mu^o(c_\epsilon) = \mu^o_+(c), \quad \lim_{\epsilon \to 0-} \mu^o(c_\epsilon) = \mu^o_-(c), \\ &\lim_{\epsilon \to 0+} \mu^e_*(c_\epsilon) = \mu^e_{*+}(c), \quad \lim_{\epsilon \to 0-} \mu^e_*(c_\epsilon) = \mu^e_{*-}(c), \\ &\lim_{\epsilon \to 0+} \nu^e_k(c_\epsilon) = \nu^e_{k+}(c), \quad \lim_{\epsilon \to 0-} \nu^e_k(c_\epsilon) = \nu^e_{k-}(c). \end{split}$$

Here $c_{\epsilon} = c + i\epsilon = u(y_c) + i\epsilon \in D_{\epsilon_0}$.

Proof. By Proposition 3.6, $\phi_1(y,c)$ is continuous for $(y,c) \in [0,\pi] \times \Omega_{\epsilon_0}$ and for ϵ_0 small enough,

$$|\phi_1(y,c)| > \frac{1}{2}, \quad |\phi_1(y,c)-1| \le C|y-y_c|^2.$$

Thus, for $(y,c) \in [0,\pi] \times \Omega_{\epsilon_0}$ with $(z-y_c)(z-y) \leq 0$, there exists a constant C so that

$$\left|\frac{\rho(c)}{(u(y)-c)^2} \left(\frac{\phi_1(z,c)}{\phi_1(y,c)^2} - 1\right)\right| \le C.$$

This implies that as $c_{\epsilon} \to u(y_c) = c$,

$$\rho(c_{\epsilon}) \int_0^{\pi} \int_{y_c}^y \frac{\varphi(z)}{(u(y) - c_{\epsilon})^2} \Big(\frac{\phi_1(z, c_{\epsilon})}{\phi_1(y, c_{\epsilon})^2} - 1 \Big) dz dy \longrightarrow \rho(c) \mathcal{L}_0(\varphi)(c),$$

from which and Lemma 5.4, it follows that as $\epsilon \to 0\pm$,

$$i\alpha u'(y_c)\rho(c_{\epsilon})\int_0^{\pi} \frac{1}{\phi(y',c_{\epsilon})^2} \int_{y_c}^{y'} f_o\phi(y'',c_{\epsilon})dy''dy'$$
$$\longrightarrow \rho(c)u'(y_c)\mathcal{L}_0(\widehat{\omega})(c) + \rho(c)u'(y_c)\mathrm{II}_{1,1}(\widehat{\omega})(c) \pm i\pi\widehat{\omega}(y_c)u'(y_c)$$

Then by Lemma 4.2, we infer that

$$\lim_{\epsilon \to 0\pm} \mu^o(c_\epsilon) = \mu^o_\pm(c).$$

Notice that for $c_{\epsilon} = u(y_c) + i\epsilon$ and $k \in \{0, 1\}$,

$$\int_{y_c}^{k\pi} f_e \phi(y, c_\epsilon) dy \longrightarrow \frac{1}{i\alpha} \mathcal{E}_k(\widehat{\omega}_e), \quad \text{as } \epsilon \to 0.$$

The other limits can be proved similarly.

Lemma 5.6. Let $\varphi(y) \in H^1(0, \pi)$. Then it holds that for $0 \le y < y_c \le \pi$,

$$\lim_{\epsilon \to 0} \phi(y, c_{\epsilon}) \int_{0}^{y} \frac{\int_{y_{c}}^{y'} \varphi(y'')\phi_{1}(y'', c_{\epsilon})dy''}{\phi(y', c_{\epsilon})^{2}} dy' = \phi(y, u(y_{c})) \int_{0}^{y} \frac{\int_{y_{c}}^{y'} \varphi(y'')\phi_{1}(y'', u(y_{c}))dy''}{\phi(y', u(y_{c}))^{2}} dy',$$
$$\lim_{\epsilon \to 0} \phi(y, c_{\epsilon}) \int_{0}^{y} \frac{1}{\phi(y', c_{\epsilon})^{2}} dy' = \phi(y, u(y_{c})) \int_{0}^{y} \frac{1}{\phi(y', u(y_{c}))^{2}} dy',$$

and for $0 \le y_c < y \le \pi$,

$$\begin{split} \lim_{\epsilon \to 0} \phi(y, c_{\epsilon}) \int_{\pi}^{y} \frac{\int_{y_{c}}^{y'} \varphi(y'') \phi_{1}(y'', c_{\epsilon}) dy''}{\phi(y', c_{\epsilon})^{2}} dy' &= \phi(y, u(y_{c})) \int_{\pi}^{y} \frac{\int_{y_{c}}^{y'} \varphi(y'') \phi_{1}(y'', u(y_{c})) dy''}{\phi(y', u(y_{c}))^{2}} dy', \\ \lim_{\epsilon \to 0} \phi(y, c_{\epsilon}) \int_{\pi}^{y} \frac{1}{\phi(y', c_{\epsilon})^{2}} dy' &= \phi(y, u(y_{c})) \int_{\pi}^{y} \frac{1}{\phi(y', u(y_{c}))^{2}} dy'. \\ Here \ c_{\epsilon} &= c + i\epsilon = u(y_{c}) + i\epsilon \in D_{\epsilon_{0}}. \end{split}$$

Proof. We consider the case of $0 \le y < y_c \le \pi$. Another case is similar. Notice that for $0 \le y' \le y$, $y' \le y'' \le y_c$,

(5.15)
$$\left|\frac{\phi(y,c_{\epsilon})}{\phi(y',c_{\epsilon})^{2}}\right| + \left|\frac{\phi(y,c_{\epsilon})\varphi(y'')\phi_{1}(y'',c_{\epsilon})}{\phi(y',c_{\epsilon})^{2}}\right|$$
$$\leq C\frac{|u(y)-c_{\epsilon}|}{|u(y')-c_{\epsilon}|^{2}} \leq \frac{C}{|u(y')-c_{\epsilon}|} \leq \frac{C}{|u(y)-u(y_{c})|}$$

Then the result follows from Lebesgue's dominated convergence theorem and the continuity of ϕ_1 , ϕ .

5.3. Solution formula of the linearized Euler equations. Let $\widehat{\psi}(t, \alpha, y)$ be a solution of (2.1) with initial data $\widehat{\psi}(0, \alpha, y) = -(\partial_y^2 - \alpha^2)^{-1}\widehat{\omega}_0(\alpha, y)$. Then we have

$$\widehat{\psi}(t,\alpha,y) = \frac{1}{2\pi i} \oint_{\partial\Omega} e^{-i\alpha tc} (c - \mathcal{R}_{\alpha})^{-1} \widehat{\psi}(0,\alpha,y) dc,$$

where Ω is a simply connected domain including the spectrum $\sigma(\mathcal{R}_{\alpha}) = [-1, 1]$ of \mathcal{R}_{α} .

Let $\Phi(\alpha, y, c)$ be a solution of (4.1) with $f(\alpha, y, c) = \frac{\widehat{\omega}_0(\alpha, y)}{i\alpha(u(y)-c)}$. We have $(c - \mathcal{R}_{\alpha})^{-1}\widehat{\psi}(0, \alpha, y) = i\alpha\Phi(\alpha, y, c).$

Therefore, we have

(5.16)
$$\widehat{\psi}(t,\alpha,y) = \frac{1}{2\pi} \oint_{\partial\Omega_{\epsilon_0}} e^{-i\alpha tc} \alpha \Phi(\alpha,y,c) dc.$$

In particular,

(5.17)
$$\widehat{\psi}(0,\alpha,y) = \frac{1}{2\pi} \oint_{\partial \Omega_{\epsilon_0}} \alpha \Phi(\alpha,y,c) dc.$$

Lemma 5.7. Let $\widehat{\psi}(t, \alpha, y)$ be as in (5.16). Then we have

$$\begin{split} \widehat{\psi}(t,\alpha,y) = & \frac{1}{2\pi} \int_{-1}^{1} e^{-i\alpha tc} \alpha \widetilde{\Phi}_{o}(\alpha,y,c) dc + \frac{1}{2\pi} \int_{-1}^{1} e^{-i\alpha tc} \alpha \widetilde{\Phi}_{e}(\alpha,y,c) dc \\ & \triangleq & \widehat{\psi}_{o}(t,\alpha,y) + \widehat{\psi}_{e}(t,\alpha,y). \end{split}$$

In particular,

$$\widehat{\psi}(0,\alpha,y) = \frac{1}{2\pi} \int_{-1}^{1} \alpha \widetilde{\Phi}_o(\alpha,y,c) dc + \frac{1}{2\pi} \int_{-1}^{1} \alpha \widetilde{\Phi}_e(\alpha,y,c) dc$$

where

$$\widetilde{\Phi}_{o}(y,c) = \begin{cases} (\mu_{-}^{o}(c) - \mu_{+}^{o}(c))\phi(y,c) \int_{0}^{y} \frac{1}{\phi(z,c)^{2}} dz, \ 0 \le y \le y_{c}, \\ (\mu_{-}^{o}(c) - \mu_{+}^{o}(c))\phi(y,c) \int_{\pi}^{y} \frac{1}{\phi(z,c)^{2}} dz, \ y_{c} \le y \le \pi, \end{cases}$$

and

$$\widetilde{\Phi}_{e}(y,c) = \begin{cases} \left(\mu_{*-}^{e}(c) - \mu_{*+}^{e}(c)\right) \int_{0}^{y} \frac{\phi(y,c)}{\phi(z,c)^{2}} dz + \left(\nu_{0-}^{e}(c) - \nu_{0+}^{e}(c)\right)\phi(y,c), \ 0 \le y < y_{c}, \\ \left(\mu_{*-}^{e}(c) - \mu_{*+}^{e}(c)\right) \int_{\pi}^{y} \frac{\phi(y,c)}{\phi(z,c)^{2}} dz + \left(\nu_{1-}^{e}(c) - \nu_{1+}^{e}(c)\right)\phi(y,c), \ y_{c} < y \le \pi, \\ and \ \widetilde{\Phi}_{o}(y,c) = -\widetilde{\Phi}_{o}(-y,c), \ \widetilde{\Phi}_{e}(y,c) = \widetilde{\Phi}_{e}(-y,c) \ for \ y \in [-\pi,0]. \end{cases}$$

Proof. For any $0 < \epsilon \leq \epsilon_0$, we have

$$\widehat{\psi}(t,\alpha,y) = \frac{1}{2\pi} \oint_{\partial\Omega_{\epsilon}} e^{-i\alpha tc} \alpha \Phi(\alpha,y,c) dc = I_1 + I_2 + I_3 + I_4,$$

where

$$\begin{split} I_1 &= \frac{1}{2\pi} \int_{-1}^{1} e^{-i\alpha t(c-i\epsilon)} \alpha \Phi(\alpha, y, c-i\epsilon) dc, \\ I_2 &= \frac{1}{2\pi} \int_{1}^{-1} e^{-i\alpha t(c+i\epsilon)} \alpha \Phi(\alpha, y, c+i\epsilon) dc, \\ I_3 &= \frac{1}{2\pi} \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} e^{-i\alpha t(1+\epsilon e^{i\theta})} \alpha \Phi(\alpha, y, 1+\epsilon e^{i\theta}) i\epsilon e^{i\theta} d\theta, \\ I_4 &= \frac{1}{2\pi} \int_{\frac{\pi}{2}}^{\frac{3\pi}{2}} e^{-i\alpha t(-1+\epsilon e^{i\theta})} \alpha \Phi(\alpha, y, -1+\epsilon e^{i\theta}) i\epsilon e^{i\theta} d\theta. \end{split}$$
Proposition 5.3, Proposition 5.1 and Lebesgue's dominated convergence theorem ensure that

$$\lim_{\epsilon \to 0+} I_1 = \frac{1}{2\pi} \int_{-1}^{1} \alpha e^{-i\alpha tc} (\Phi^o_-(\alpha, y, c) + \Phi^e_-(\alpha, y, c)) dc,$$
$$\lim_{\epsilon \to 0+} I_2 = -\frac{1}{2\pi} \int_{-1}^{1} \alpha e^{-i\alpha tc} (\Phi^o_+(\alpha, y, c) + \Phi^e_+(\alpha, y, c)) dc,$$

and

$$\lim_{\epsilon \to 0+} I_3 = 0, \quad \lim_{\epsilon \to 0+} I_4 = 0$$

Now, the lemma follows from the facts that $\widetilde{\Phi}_o = \Phi_-^o - \Phi_+^o, \ \widetilde{\Phi}_e = \Phi_-^e - \Phi_+^e.$

5.4. Dual formulation. As in [35], we will use the dual method to derive the decay estimates of the velocity. So, we introduce the dual formulation of the stream function. Let

$$\mu_{-}^{o}(c) - \mu_{+}^{o}(c) = \frac{2}{\alpha} \frac{\operatorname{AC}_{o}(\widehat{\omega}_{o}) + \operatorname{BD}_{o}(\widehat{\omega}_{o})}{\operatorname{A}^{2} + \operatorname{B}^{2}} \stackrel{\text{def}}{=} \frac{2}{\alpha} \rho(c) \mu_{1}(c),$$

$$\mu_{-}^{e}(c) - \mu_{+}^{e}(c) = \frac{2}{\alpha} \frac{I(c)^{2}(\operatorname{AC}_{e} + \operatorname{BD}_{e}) + I(c)(\phi\phi')(j\pi, c)(\operatorname{C}_{e} + E_{1-j}\operatorname{B})|_{j=0}^{1}}{I(c)^{2}(\operatorname{A}_{1}^{2} + \operatorname{B}_{1}^{2})/\rho(c)^{2}} \stackrel{\text{def}}{=} \frac{2}{\alpha} \mu_{2}(c),$$

$$\nu_{j-}^{e}(c) - \nu_{j+}^{e}(c) = -\frac{2}{\alpha} \frac{\mu_{2}}{\phi(j\pi, c)\phi'(j\pi, c)} \stackrel{\text{def}}{=} \frac{2}{\alpha} \nu_{j}(c).$$

Here $C_e = C_o(\widehat{\omega}_e)(c)$, $D_e = D_o(\widehat{\omega}_e)(c)$ and $E_{1-j} = E_{1-j}(\widehat{\omega}_e)(y_c)$. We denote

(5.18)
$$\Lambda_1(\varphi)(y_c) = \Lambda_{1,1}(\varphi)(y_c) + \Lambda_{1,2}(\varphi)(y_c),$$

(5.19)
$$\Lambda_2(\varphi)(y_c) = \Lambda_{2,1}(\varphi)(y_c) + \Lambda_{2,2}(\varphi)(y_c),$$

where

(5.20)
$$\Lambda_{1,1}(\varphi)(y_c) = \rho(c)u''(y_c) \Pi_{1,1}(\varphi)(c),$$

(5.21)
$$\Lambda_{1,2}(\varphi)(y_c) = \rho(c)u''(y_c)\mathcal{L}_0(\varphi)(c) + u'(y_c)\rho(c)\mathrm{II}(c)\varphi(y_c),$$

(5.22)
$$\Lambda_{2,1}(\varphi)(y_c) = \rho(c) \operatorname{II}_{1,1}(u''\varphi)(c)$$

 $\Lambda_{2,1}(\varphi)(y_c) = \rho(c)\Pi_{1,1}(u''\varphi)(c),$ $\Lambda_{2,2}(\varphi)(y_c) = \rho(c)\mathcal{L}_0(u''\varphi)(c) + u'(y_c)\rho(c)\Pi(c)\varphi(y_c),$ (5.23)

We denote

(5.24)
$$\Lambda_3(\widehat{\omega}_e)(y_c) = \rho(c)\Lambda_1(\widehat{\omega}_e)(y_c) + \Lambda_{3,1}(\widehat{\omega}_e)(y_c),$$

(5.25)
$$\Lambda_4(g)(y_c) = \rho(c)\Lambda_2(g)(y_c) + \Lambda_{4,1}(g)(y_c),$$

where

(5.26)
$$\Lambda_{3,1}(\widehat{\omega}_e)(y_c) = J_j \left(\frac{u''(y_c)}{u'(y_c)} \mathbf{E}_{1-j}(\widehat{\omega}_e) + \widehat{\omega}_e(y_c) \right) \Big|_{j=0}^1,$$

(5.27)
$$\Lambda_{4,1}(g)(y_c) = J_j \left(\frac{\mathrm{E}_{1-j}(gu'')}{u'(y_c)} + g(y_c) \right) \Big|_{j=0}^1,$$

with

$$J_j(c) = \frac{(\phi\phi')(j\pi, c)\rho(c)}{I(c)} = \frac{-u'(y_c)\rho(c)^2}{(\phi\phi')((1-j)\pi, c)}$$

(5.28)
$$= \frac{-u'(y_c)(u(j\pi) - c)^2}{(\phi_1 \phi'_1)((1-j)\pi, c)} \quad \text{for } j = 0, 1.$$

Lemma 5.8. Let $f(\alpha, y) = (\partial_y^2 - \alpha^2)g(\alpha, y)$ with $g \in H^2(0, \pi) \cap H^1_0(0, \pi)$. Then we have

$$\int_0^{\pi} \widehat{\psi}_o(t,\alpha,y) f(\alpha,y) dy = -\int_{-1}^1 K_o(c,\alpha) e^{-i\alpha ct} dc,$$

where

(5.29)
$$K_o(c,\alpha) = \frac{\Lambda_1(\widehat{\omega}_o)(y_c)\Lambda_2(g)(y_c)}{(\mathbf{A}(c)^2 + \mathbf{B}(c)^2)u'(y_c)}.$$

Proof. By Lemma 5.7, we get

$$\begin{split} &\int_{0}^{\pi} \widehat{\psi}_{o}(t,\alpha,y) f(\alpha,y) dy \\ &= \frac{1}{\pi} \int_{0}^{\pi} f(\alpha,y) \int_{-1}^{u(y)} \rho(c) \mu_{1}(c) \phi(y,c) \int_{\pi}^{y} \frac{1}{\phi(z,c)^{2}} dz e^{-i\alpha c t} dc dy \\ &+ \frac{1}{\pi} \int_{0}^{\pi} f(\alpha,y) \int_{u(y)}^{1} \rho(c) \mu_{1}(c) \phi(y,c) \int_{0}^{y} \frac{1}{\phi(z,c)^{2}} dz e^{-i\alpha c t} dc dy \\ &= -\frac{1}{\pi} \int_{-1}^{1} \rho(c) \mu_{1}(c) e^{-i\alpha c t} \int_{0}^{\pi} \frac{\int_{y_{c}}^{z} f(\alpha,y) \phi(y,c) dy}{\phi(z,c)^{2}} dz dc. \end{split}$$

First of all, we have

$$\rho(c)\mu_1(c) = \pi \frac{\mathcal{A}(c)\rho(c)\frac{\widehat{\omega}_o(y_c)}{u'(y_c)} + \rho(c)^2 \frac{u''(y_c)}{u'(y_c)} \mathrm{II}_1(\widehat{\omega}_o)(c)}{\mathcal{A}(c)^2 + \mathcal{B}(c)^2} = \pi \frac{\sin y_c \Lambda_1(\widehat{\omega}_o)}{\mathcal{A}(c)^2 + \mathcal{B}(c)^2}.$$

We write

$$\begin{split} &\int_{0}^{\pi} \frac{\int_{y_{c}}^{z} f(\alpha, y) \phi(y, c) dy}{\phi(z, c)^{2}} dz \\ &= \int_{0}^{\pi} \frac{\int_{y_{c}}^{z} (g''(y) - \alpha^{2}g(y)) \phi(y, c) dy}{\phi(z, c)^{2}} dz \\ &= \int_{0}^{\pi} \frac{\int_{y_{c}}^{z} g(y) (\phi'' - \alpha^{2}\phi)(y, c) dy + g'(z) \phi(z, c) - g(z) \phi'(z, c) + g(y_{c}) \phi'(y_{c}, c)}{\phi(z, c)^{2}} dz \\ &= \int_{0}^{\pi} \left[\frac{\int_{y_{c}}^{z} g(y) u''(y) \phi_{1}(y, c) dy}{\phi(z, c)^{2}} + \left(\frac{g(z)}{\phi(z, c)} \right)' + \frac{g(y_{c}) u'(y_{c})}{\phi(z, c)^{2}} \right] dz \\ &= \Pi_{1}(gu'') + p.v. \int_{0}^{\pi} \left[\left(\frac{g(z)}{\phi(z, c)} \right)' + \frac{g(y_{c}) u'(y_{c})}{\phi(z, c)^{2}} \right] dz. \end{split}$$

Notice that for $c \in (-1, 1)$, we have

$$\frac{g(z)}{\phi(z,c)} - \frac{g(y_c)}{u(z) - c} = \frac{g(z)(1 - \phi_1(z,c))}{(u(z) - c)\phi_1(z,c)} + \frac{g(z) - g(y_c)}{u(z) - c} \in C([0,\pi]),$$

then we get

$$p.v. \int_0^\pi \left[\left(\frac{g(z)}{\phi(z,c)} \right)' + \frac{g(y_c)u'(y_c)}{\phi(z,c)^2} \right] dz$$

$$= \left(\frac{g(z)}{\phi(z,c)} - \frac{g(y_c)}{u(z) - c}\right) \Big|_0^\pi + p.v. \int_0^\pi \left[\left(\frac{g(y_c)}{u(z) - c}\right)' + \frac{g(y_c)u'(y_c)}{\phi(z,c)^2} \right] dz \\ = \frac{g(y_c)}{u(0) - c} - \frac{g(y_c)}{u(\pi) - c} + p.v. \int_0^\pi \left[-\frac{g(y_c)(u'(z) - u'(y_c))}{(u(z) - c)^2} + \frac{g(y_c)u'(y_c)}{(u(z) - c)^2} \left(\frac{1}{\phi(z,c)^2} - 1\right) \right] dz \\ = \frac{g(y_c)(u(0) - u(\pi))}{\rho(c)} - g(y_c)p.v. \int_0^\pi \frac{(u'(z) - u'(y_c))}{(u(z) - c)^2} dz + g(y_c)u'(y_c) \mathrm{II}(c).$$

From Remark 5.2 and $p.v. \int_0^{\pi} \frac{1}{u(z)-c} dz = 0$, we know that

$$p.v. \int_0^\pi \frac{u'(z) - u'(y_c)}{(u(z) - c)^2} dz = II_{1,1}(u'')(c)$$
$$= \partial_c \left(p.v. \int_0^\pi \frac{u'(y) - u'(0)}{u(y) - c} dy \right) = \partial_c \left(\ln \frac{u(\pi) - c}{c - u(0)} \right) = \frac{u(0) - u(\pi)}{\rho(c)}.$$

This gives

$$p.v. \int_0^\pi \left[\left(\frac{g(z)}{\phi(z,c)} \right)' + \frac{g(y_c)u'(y_c)}{\phi(z,c)^2} \right] dz = u'(y_c)g(y_c)\mathrm{II}(c),$$

which implies

$$\int_0^{\pi} \frac{\int_{y_c}^z f(\alpha, y)\phi(y, c)dy}{\phi(z, c)^2} dz = \mathrm{II}_1(gu'') + u'(y_c)g(y_c)\mathrm{II}(c) = \Lambda_2(g)(y_c)/\rho(c).$$

Therefore,

$$\int_0^\pi \widehat{\psi}_o(t,\alpha,y) f(y) dy = -\int_{-1}^1 \frac{\Lambda_1(\widehat{\omega}_o)(y_c)\Lambda_2(g)(y_c)}{(\mathbf{A}(c)^2 + \mathbf{B}(c)^2)u'(y_c)} e^{-i\alpha ct} dc,$$

which proves the lemma.

Remark 5.9. From the above proof, we know that if $g \in H^2(0,\pi)$, then we have

$$p.v. \int_0^\pi \left[\left(\frac{g(z)}{\phi(z,c)} \right)' + \frac{g(y_c)u'(y_c)}{\phi(z,c)^2} \right] dz = \frac{g(z)}{\phi(z,c)} \Big|_{z=0}^\pi + u'(y_c)g(y_c)\mathrm{II}(c).$$

Lemma 5.10. Let $f(\alpha, y) = (\partial_y^2 - \alpha^2)g(\alpha, y)$ with $g \in H^2(0, \pi)$ and $g'(0) = g'(\pi) = 0$. Then we have

$$\int_0^\pi \widehat{\psi}_e(t,\alpha,y) f(\alpha,y) dy = -\int_{-1}^1 K_e(c,\alpha) e^{-i\alpha ct} dc,$$

where

(5.30)
$$K_e(c,\alpha) = \frac{\Lambda_3(\widehat{\omega}_e)(y_c)\Lambda_4(g)(y_c)}{u'(y_c)(\Lambda_1(c)^2 + B_1(c)^2)}.$$

Proof. By Lemma 5.7, we have

$$\begin{split} &\int_{0}^{\pi} \widehat{\psi}_{e}(t,\alpha,y) f(\alpha,y) dy \\ &= \frac{1}{\pi} \int_{0}^{\pi} f(\alpha,y) \int_{-1}^{u(y)} \mu_{2}(c) \phi(y,c) \int_{\pi}^{y} \frac{1}{\phi(z,c)^{2}} dz e^{-i\alpha c t} dc dy \\ &+ \frac{1}{\pi} \int_{0}^{\pi} f(\alpha,y) \int_{u(y)}^{1} \mu_{2}(c) \phi(y,c) \int_{0}^{y} \frac{1}{\phi(z,c)^{2}} dz e^{-i\alpha c t} dc dy \end{split}$$

$$\begin{aligned} &+ \frac{1}{\pi} \int_0^{\pi} f(\alpha, y) \int_{u(y)}^1 \nu_0(c) \phi(y, c) e^{-i\alpha ct} dc dy + \frac{1}{\pi} \int_0^{\pi} f(\alpha, y) \int_{-1}^{u(y)} \nu_1(c) \phi(y, c) e^{-i\alpha ct} dc dy \\ &= -\frac{1}{\pi} \int_{-1}^1 \mu_2(c) e^{-i\alpha ct} \int_0^{\pi} \frac{\int_{y_c}^z f(y) \phi(y, c) dy}{\phi(z, c)^2} dz dc \\ &+ \frac{1}{\pi} \int_{-1}^1 \nu_0(c) e^{-i\alpha ct} \int_0^{y_c} f(y) \phi(y, c) dy dc + \frac{1}{\pi} \int_{-1}^1 \nu_1(c) e^{-i\alpha ct} \int_{y_c}^{\pi} f(y) \phi(y, c) dy dc. \end{aligned}$$

We have

$$\int_0^{\pi} \frac{\int_{y_c}^z f(y)\phi(y,c)dy}{\phi(z,c)^2} dz = \mathrm{II}_1(gu'') + p.v. \int_0^{\pi} \left[\left(\frac{g(z)}{\phi(z,c)} \right)' + \frac{g(y_c)u'(y_c)}{\phi(z,c)^2} \right] dz,$$

and by Remark 5.9,

$$p.v. \int_0^{\pi} \left[\left(\frac{g(z)}{\phi(z,c)} \right)' + \frac{g(y_c)u'(y_c)}{\phi(z,c)^2} \right] dz = \frac{g(y_c)}{\rho(c)} \mathcal{A}(c) - \frac{g(0)}{\phi(0,c)} + \frac{g(\pi)}{\phi(\pi,c)}.$$

On the other hand, we have

$$\int_{0}^{y_c} f(y)\phi(y,c)dy = -\mathbf{E}_0(gu'') - g(y_c)u'(y_c) + g(0)\phi'(0,c),$$

$$\int_{y_c}^{\pi} f(y)\phi(y,c)dy = \mathbf{E}_1(gu'') + g(y_c)u'(y_c) - g(\pi)\phi'(\pi,c).$$

Then we conclude that

$$\begin{split} &\int_{0}^{\pi} \widehat{\psi}_{e}(t,\alpha,y) f(y) dy \\ &= -\frac{1}{\pi} \int_{-1}^{1} \Big[\mu_{2} \Big(\frac{g(y_{c}) \mathbf{A}(c)}{\rho(c)} + \mathrm{II}_{1}(gu'') \Big) + \nu_{0} \big(\mathbf{E}_{0}(gu'') + g(y_{c})u'(y_{c}) \big) \\ &- \nu_{1} \big(\mathbf{E}_{1}(gu'') + g(y_{c})u'(y_{c}) \big) \Big] e^{-i\alpha ct} dc, \end{split}$$

Here we used

$$\nu_0(c) = -\frac{\mu_2(c)}{\phi(0,c)\phi'(0,c)}, \quad \nu_1(c) = -\frac{\mu_2(c)}{\phi(\pi,c)\phi'(\pi,c)}.$$

Recall that

$$\begin{split} \mu_{2}(c) &= \frac{I(c)^{2}(\mathrm{AC}_{e} + \mathrm{BD}_{e}) + I(c)\phi(j\pi,c)\phi'(j\pi,c)(\mathrm{C}_{e} + E_{1-j}B)|_{j=0}^{1}}{I(c)^{2}(\mathrm{A}_{1}(c)^{2} + \mathrm{B}_{1}(c)^{2})/\rho(c)^{2}} \\ &= \frac{\rho(c)^{2}(\mathrm{AC}_{e} + \mathrm{BD}_{e}) + \rho(c)J_{j}(c)(\mathrm{C}_{e} + E_{1-j}B)|_{j=0}^{1}}{\mathrm{A}_{1}(c)^{2} + \mathrm{B}_{1}(c)^{2}} \\ &= \pi\rho^{2}\frac{\left(\mathrm{A}\rho(c)\frac{\widehat{\omega}_{e}(y_{c})}{u'(y_{c})} + \rho(c)\frac{u''(y_{c})}{u'(y_{c})^{2}}u'(y_{c})\rho(c)\mathrm{II}_{1}(\widehat{\omega}_{e})\right) + J_{j}\left(\frac{u''(y_{c})}{u'(y_{c})^{2}}\mathrm{E}_{1-j} + \frac{\widehat{\omega}_{e}(y_{c})}{u'(y_{c})}\right)|_{j=0}^{1}}{\mathrm{A}_{1}(c)^{2} + \mathrm{B}_{1}(c)^{2}} \\ &= \frac{\pi\rho(c)^{2}}{u'(y_{c})}\frac{\rho(c)\Lambda_{1}(\widehat{\omega}_{e})(y_{c}) + \Lambda_{3,1}(\widehat{\omega}_{e})(y_{c})}{\mathrm{A}_{1}(c)^{2} + \mathrm{B}_{1}(c)^{2}} = \frac{\pi\rho(c)^{2}}{u'(y_{c})}\frac{\Lambda_{3}(\widehat{\omega}_{e})(y_{c})}{\mathrm{A}_{1}(c)^{2} + \mathrm{B}_{1}(c)^{2}}. \end{split}$$

Then we have

$$\mu_2 \Big(\frac{g(y_c) \mathbf{A}(c)}{\rho(c)} + \mathrm{II}_1(gu'') \Big) + \nu_0 \big(\mathbf{E}_0(gu'') + g(y_c)u'(y_c) \big) - \nu_1 \big(\mathbf{E}_1(gu'') + g(y_c)u'(y_c) \big)$$

LINEAR INVISCID DAMPING AND ENHANCED DISSIPATION

$$= \mu_2(c) \frac{\Lambda_2(g)(y_c)}{\rho(c)} - \frac{\mu_2(c) \left(\mathcal{E}_0(gu'') + g(y_c)u'(y_c) \right)}{\phi(0, c)\phi'(0, c)} + \frac{\mu_2(c) \left(\mathcal{E}_1(gu'') + g(y_c)u'(y_c) \right)}{\phi(\pi, c)\phi'(\pi, c)}$$

$$= \mu_2 \frac{\Lambda_2(g)(y_c)}{\rho(c)} + \frac{\mu_2 J_1 \left(\mathcal{E}_0(gu'') + g(y_c)u'(y_c) \right)}{u'(y_c)\rho(c)^2} - \frac{\mu_2 J_0 \left(\mathcal{E}_1(gu'') + g(y_c)u'(y_c) \right)}{u'(y_c)\rho(c)^2}$$

$$= \mu_2(c) \frac{\Lambda_2(g)(y_c)}{\rho(c)} + \frac{\mu_2(c)\Lambda_{4,1}(g)(y_c)}{\rho(c)^2} = \frac{\mu_2(c)\Lambda_4(g)(y_c)}{\rho(c)^2}$$

$$= \frac{\pi\rho(c)^2}{u'(y_c)} \frac{\Lambda_3(\widehat{\omega}_e)(y_c)}{\Lambda_1(c)^2 + \mathcal{B}_1(c)^2} \frac{\Lambda_4(g)(y_c)}{\rho(c)^2} = \pi \frac{\Lambda_3(\widehat{\omega}_e)(y_c)\Lambda_4(g)(y_c)}{u'(y_c)(\Lambda_1(c)^2 + \mathcal{B}_1(c)^2)} = \pi K_e(c, \alpha),$$

which gives the lemma.

6. LINEAR INVISCID DAMPING

The decay estimates of the velocity rely on the following uniform $W^{1,2}$ estimates of the kernels K_o and K_e .

Proposition 6.1. Let K_o be defined by (5.29) with $g \in H^2(0,\pi)$, $g(0) = g(\pi) = 0$ and $\widehat{\omega}_o = \frac{1}{2}(\widehat{\omega}_0(\alpha, y) - \widehat{\omega}_0(\alpha, -y)) \in H^2$. Then $K_o(-1, \alpha) = K_o(1, \alpha) = 0$, and there exists a constant C independent of α such that,

$$\begin{aligned} \|K_o(\cdot,\alpha)\|_{L^1_c(-1,1)} &\leq C \|\widehat{\omega}_o\|_{L^2} \|g\|_{L^2}, \\ \|(\partial_c K_o)(\cdot,\alpha)\|_{L^1_c(-1,1)} &\leq C \|\widehat{\omega}_o\|_{H^1} \|g\|_{H^1}, \\ \|(\partial_c^2 K_o)(\cdot,\alpha)\|_{L^1_c(-1,1)} &\leq C \alpha^{\frac{1}{2}} \|\widehat{\omega}_o\|_{H^2} \|f\|_{L^2}. \end{aligned}$$

Proposition 6.2. Let K_e be defined by (5.30) with $g \in H^2(0,\pi)$, $g'(0) = g'(\pi) = 0$ and $\widehat{\omega}_e = \frac{1}{2}(\widehat{\omega}_0(\alpha, y) + \widehat{\omega}_0(\alpha, -y)) \in H^2$. Then $K_e(-1, \alpha) = K_e(1, \alpha) = 0$, and there exists a constant C independent of α such that,

$$\begin{split} \|K_e(\cdot,\alpha)\|_{L^1_c(-1,1)} &\leq C \|\widehat{\omega}_e\|_{L^2} \|g\|_{L^2},\\ \|(\partial_c K_e)(\cdot,\alpha)\|_{L^1_c(-1,1)} &\leq C |\alpha|^{\frac{1}{2}} \|\widehat{\omega}_e\|_{H^1} (\|g'\|_{L^2} + \alpha \|g\|_{L^2}),\\ \|(\partial_c^2 K_e)(\cdot,\alpha)\|_{L^1_c(-1,1)} &\leq C |\alpha|^{\frac{3}{2}} \|\widehat{\omega}_e\|_{H^2} \|f\|_{L^2}. \end{split}$$

For the moment, let us admit these two propositions and complete the proof of Theorem 1.1.

We have $\hat{\psi}(t, \alpha, y) = \hat{\psi}_o(t, \alpha, y) + \hat{\psi}_e(t, \alpha, y)$. Using integration by parts, Proposition 6.1 and Proposition 6.2 ensure that

$$\begin{split} \|\widehat{\psi}_{o}(t,\alpha,\cdot)\|_{L^{2}} &= 2 \sup_{\|f\|_{L^{2}} \leq 1} \left| \int_{0}^{\pi} \widehat{\psi}_{o}(t,\alpha,y) f(y) dy \right| \\ &\leq 2 \sup_{\|f\|_{L^{2}} \leq 1} \left| \int_{-1}^{1} K_{o}(c,\alpha) e^{-i\alpha ct} dc \right| \leq C \frac{1}{|\alpha|^{\frac{3}{2}} t^{2}} \|\widehat{\omega}_{o}\|_{H^{2}}, \end{split}$$

and

$$\begin{split} \|\widehat{\psi}_{e}(t,\alpha,\cdot)\|_{L^{2}} &= 2 \sup_{\|f\|_{L^{2}} \leq 1} \left| \int_{0}^{\pi} \widehat{\psi}_{e}(t,\alpha,y) f(y) dy \right| \\ &\leq 2 \sup_{\|f\|_{L^{2}} \leq 1} \left| \int_{-1}^{1} K_{e}(c,\alpha) e^{-i\alpha ct} dc \right| \leq C \frac{1}{|\alpha|^{\frac{1}{2}} t^{2}} \|\widehat{\omega}_{e}\|_{H^{2}}. \end{split}$$

Proposition 6.1 and Proposition 6.2 also imply that

$$\begin{aligned} \alpha^2 \|\widehat{\psi_o}(t,\alpha,\cdot)\|_{L^2}^2 + \|\partial_y \widehat{\psi_o}(t,\alpha,\cdot)\|_{L^2}^2 \\ &= -2\int_0^\pi \widehat{\psi_o}(t,\alpha,y)(\overline{\psi_o}'' - \alpha^2 \overline{\psi_o})(t,\alpha,y)dy \le \begin{cases} C\|\widehat{\omega}_o\|_{L^2}\|\widehat{\psi_o}(t,\alpha,\cdot)\|_{L^2}, \\ C\frac{1}{|\alpha t|}\|\widehat{\omega}_o\|_{H^1}\|\widehat{\psi_o}(t,\alpha,\cdot)\|_{H^1}, \end{cases} \end{aligned}$$

and

$$\begin{split} &\alpha^2 \|\widehat{\psi_e}(t,\alpha,\cdot)\|_{L^2}^2 + \|\partial_y \widehat{\psi_e}(t,\alpha,\cdot)\|_{L^2}^2 \\ &= -2\int_0^\pi \widehat{\psi_e}(t,\alpha,y)(\overline{\psi_e}'' - \alpha^2 \overline{\psi_e})(t,\alpha,y)dy \\ &\leq \begin{cases} C\|\widehat{\omega}_e(\alpha,\cdot)\|_{L^2_y}\|\widehat{\psi}_e(t,\alpha,\cdot)\|_{L^2_y}, \\ C\frac{1}{|\alpha|^{\frac{1}{2}}|t|}\|\widehat{\omega}_e(\alpha,\cdot)\|_{H^1_y}(|\alpha|\|\widehat{\psi}_e(t,\alpha,\cdot)\|_{L^2_y} + \|\partial_y \widehat{\psi}_e(t,\alpha,\cdot)\|_{L^2_y}). \end{cases} \end{split}$$

Thus, we conclude that for $|t| \leq 1$,

$$\begin{aligned} \|\widehat{V_o^1}\|_{L^2} + \|\widehat{V_o^2}\|_{L^2} &\leq |\alpha| \|\widehat{\psi_o}(t,\alpha,\cdot)\|_{L^2} + \|\partial_y \widehat{\psi_o}(t,\alpha,\cdot)\|_{L^2} \leq C|\alpha|^{-1} \|\widehat{\omega}_o\|_{L^2}, \\ \|\widehat{V_e^1}\|_{L^2} + \|\widehat{V_e^2}\|_{L^2} &\leq |\alpha| \|\widehat{\psi_e}(t,\alpha,\cdot)\|_{L^2} + \|\partial_y \widehat{\psi_e}(t,\alpha,\cdot)\|_{L^2} \leq C|\alpha|^{-1} \|\widehat{\omega}_e\|_{L^2}, \end{aligned}$$

and for $|t| \ge 1$,

$$\begin{aligned} \|\widehat{V_o^1}\|_{L^2} + \|\widehat{V_o^2}\|_{L^2} &\leq |\alpha| \|\widehat{\psi_o}(t,\alpha,\cdot)\|_{L^2} + \|\partial_y \widehat{\psi_o}(t,\alpha,\cdot)\|_{L^2} \leq C \frac{1}{|\alpha t|} \|\widehat{\omega}_o\|_{H^1}, \\ \|\widehat{V_e^1}\|_{L^2} + \|\widehat{V_e^2}\|_{L^2} &\leq |\alpha| \|\widehat{\psi_e}(t,\alpha,\cdot)\|_{L^2} + \|\partial_y \widehat{\psi_e}(t,\alpha,\cdot)\|_{L^2} \leq C \frac{1}{|\alpha|^{\frac{1}{2}} |t|} \|\widehat{\omega}_e\|_{H^1}, \end{aligned}$$

Due to $V^1 = V_o^1 + V_e^1$ and $V^2 = V_o^2 + V_e^2$, we obtain

$$\|V^1\|_{L^2_{x,y}} + \|V^2\|_{L^2_{x,y}} \le \frac{C}{\langle t \rangle} \|\omega_0\|_{H^{-\frac{1}{2}}_x H^1_y}.$$

As $\|\widehat{V^2}(t, \alpha, \cdot)\|_{L^2} \leq C|\alpha| \|\widehat{\psi}(t, \alpha, \cdot)\|_{L^2} \leq C \frac{|\alpha|^{\frac{1}{2}}}{t^2} \|\widehat{\omega}_0(t, \alpha, \cdot)\|_{H^2}$, we infer that $\|V^2\|_{L^2_{x,y}} \leq \frac{C}{\langle t^2 \rangle} \|\omega_0\|_{H^{\frac{1}{2}}_x H^2_y}.$

This completes the proof of Theorem 1.1.

Let us remark that if $\omega_0(x, y)$ is odd in y, then we have

$$\|V\|_{L^{2}_{x,y}} \leq \frac{C}{\langle t \rangle} \|\omega_{0}\|_{H^{-1}_{x}H^{1}_{y}}, \quad \|V^{2}\|_{L^{2}_{x,y}} \leq \frac{C}{\langle t^{2} \rangle} \|\omega_{0}\|_{H^{-\frac{1}{2}}_{x}H^{2}_{y}}.$$

7. Estimates of some integral operators

In this section, we present the estimates of some integral operators, which will be used to establish the $W^{2,1}$ estimates of K_o and K_e .

7.1. Basic properties of Hilbert transform. Recall that the Hilbert transform H on \mathbb{T} is defined by

$$H(f)(x) = p.v. \int_{-\pi}^{\pi} \cot \frac{x-y}{2} f(y) dy.$$

Lemma 7.1. It holds that

$$\frac{d}{dx}(H(\varphi)(x)) = H(\varphi')(x),$$

$$\frac{d}{dx}(\sin x(H\varphi)'(x) - \cos xH\varphi(x)) = \sin x(H\varphi)''(x) + \sin xH(\varphi)(x).$$

If φ is even, then

$$\cot x H(\varphi)(x) - H(\cot y\varphi)(x) = -\int_{-\pi}^{\pi} \varphi(z) \, dz,$$

$$\sin x (H\varphi)(x) - H(\sin y\varphi)(x) = \int_{-\pi}^{\pi} (\cos x + \cos y)\varphi(y) \, dy,$$

$$H(\cos y\varphi) - \cos x H(\varphi) = \sin x \int_{-\pi}^{\pi} \varphi(y) dy,$$

$$H(\varphi)(x) = -2p.v. \int_{0}^{\pi} \frac{1}{\sin y} \ln \left| \frac{\sin \frac{x+y}{2}}{\sin \frac{x-y}{2}} \right| \sin y\varphi'(y) \, dy,$$

and if φ is odd, then

$$\sin x(H\varphi)(x) - H(\sin y\varphi)(x) = 0.$$

Proof. The first two equalities are obvious. Notice that

$$\sin x (H\varphi)(x) - H(\sin y\varphi)(x) = p.v. \int_{-\pi}^{\pi} (\sin x - \sin y) \cot \frac{x - y}{2} \varphi(y) dy$$
$$= \int_{-\pi}^{\pi} (\cos x + \cos y) \varphi(y) dy,$$

in particular, if φ is odd, then

$$\sin x(H\varphi)(x) - H(\sin x\varphi)(x) = 0.$$

If φ is even, then we have

$$H(\varphi)(x) = p.v. \int_{-\pi}^{\pi} \cot \frac{x-y}{2} \varphi(y) dy$$

= $p.v. \int_{0}^{\pi} \left(\cot \frac{x-y}{2} + \cot \frac{x+y}{2} \right) \varphi(y) dy$
= $-2p.v. \int_{0}^{\pi} \frac{1}{\sin y} \ln \left| \frac{\sin \frac{x+y}{2}}{\sin \frac{x-y}{2}} \right| \sin y \varphi'(y) dy,$

and

$$\cot x (H\varphi)(x) - H(\cot y\varphi)(x)$$

= $p.v. \int_{-\pi}^{\pi} (\cot x - \cot y) \cot \frac{x-y}{2} \varphi(y) dy$
= $-p.v. \int_{-\pi}^{\pi} \frac{1 + \cos(x-y)}{\sin x \sin y} \varphi(y) dy = -\int_{-\pi}^{\pi} \varphi(y) dy,$

and

$$H(\cos y\varphi) - \cos x H(\varphi) = p.v. \int_{-\pi}^{\pi} (\cos y - \cos x) \cot \frac{x - y}{2} \varphi(y) dy$$
$$= \int_{-\pi}^{\pi} (\sin x + \sin y) \varphi(y) dy = \sin x \int_{-\pi}^{\pi} \varphi(y) dy.$$

This proves the lemma.

The following lemma can be easily proved by Hardy's inequality.

Lemma 7.2. If $\varphi \in H^2(\mathbb{T})$ is even, then $\varphi'(0) = \varphi'(\pi) = 0$ and

$$\left\|\frac{\partial_y \varphi}{\sin y}\right\|_{L^2(0,\pi)} \le C \|\varphi\|_{H^2(0,\pi)}$$

If φ is even and $\varphi(0) = \varphi(\pi) = 0$, then

$$\begin{aligned} \left\|\frac{\varphi}{\sin^2 y}\right\|_{L^2(0,\pi)} + \left\|\frac{\varphi}{\sin y}\right\|_{L^\infty(0,\pi)} &\leq C \|\varphi\|_{H^2(0,\pi)}, \\ \left\|\frac{\varphi}{\sin y}\right\|_{L^2(0,\pi)} &\leq C \|\varphi\|_{H^1(0,\pi)}. \end{aligned}$$

Lemma 7.3. Let $Z(f)(y) = f'(y) - \cot y f(y)$. If $f \in H^m(0,\pi)$ is an even function for m = 1, 2, 3, then $Z(f) \in H^{m-1}(0,\pi)$ can be extended to a periodic odd function. Moreover, we have

$$||Z(f)||_{H^{m-1}(\mathbb{T})} \le C ||f||_{H^m(0,\pi)}.$$

If we $f \in H^2(0,\pi) \cap H^1_0(0,\pi)$ is an even function, then

$$||Z(f)/\sin y||_{L^2(\mathbb{T})} \le ||f||_{H^2(0,\pi)}.$$

Proof. For $0 < y \leq \frac{\pi}{4}$, we may write

$$Z(f)(y) = m(y)y\left(\frac{f}{ym(y)}\right)' = f' - \frac{\cos yf}{ym(y)}$$

where $m(y) = \frac{\sin y}{y} \in C^{\infty}(\mathbf{R})$ and $\frac{1}{m} \in C^{\infty}([0, \frac{3\pi}{4}])$. On the other hand, for $\pi - y \leq \frac{3\pi}{4}$, we write

$$Z(f)(y) = m(\pi - y)(\pi - y) \left(\frac{f}{(\pi - y)m(\pi - y)}\right)' = f' - \frac{\cos yf}{(\pi - y)m(\pi - y)}.$$

Thus, we can conclude that

$$||Z(f)||_{H^{m-1}(0,\pi)} \le C ||f||_{H^m(0,\pi)}.$$

To prove that Z(f) can be extended to an even π -periodic function in $H^{m-1}(\mathbb{T})$, we only need to show that $Z(f)(0) = Z(f)(\pi) = 0$ and $\partial_y Z(f)$ is an even function. This is because $\lim_{y\to 0+} Z(f)(y) = \lim_{y\to 0+} f'(y) - \lim_{y\to 0+} \cot y f(y) = f'(0+) - f'(0+) = 0$, $\lim_{y\to \pi-} Z(f)(y) = f'(\pi-) - f'(\pi-) = 0$, and Z(f) is an odd function. So, $\|Z(f)\|_{H^{m-1}(\mathbb{T})} \leq C \|f\|_{H^m(0,\pi)}$. \Box

7.2. Estimates of $II_{1,1}$. Recall that $II_{1,1}$ is defined by

$$\operatorname{II}_{1,1}(\varphi)(y_c) = p.v. \int_0^\pi \frac{\operatorname{Int}(\varphi)(y) - \operatorname{Int}(\varphi)(y_c)}{(u(y) - u(y_c))^2} \, dy$$

where $Int(\varphi)$ is defined by

$$\operatorname{Int}(\varphi)(y) = \int_0^y \varphi(y') \, dy' \quad \text{for } y \in [0,\pi], \quad \operatorname{Int}(\varphi)(y) = \operatorname{Int}(\varphi)(-y) \quad \text{for } y \in [-\pi,0].$$

In what follows, we denote

$$\|f\|_{H^k_{y_c}} = \sum_{n=0}^k \|\partial_{y_c}^n f\|_{L^2_{y_c}(\mathbb{T})}, \quad \|\cdot\|_{H^k} = \|\cdot\|_{H^k(0,\pi)}.$$

Lemma 7.4. It holds that for k = 0, 1, 2,

$$\begin{aligned} \|\rho \mathrm{II}_{1,1}(\varphi)\|_{H^k_{y_c}} &\leq C \|\varphi\|_{H^k}, \\ \|\partial_{y_c}(\rho \mathrm{II}_{1,1}(\varphi))/\sin y_c\|_{L^2_{y_c}} &\leq C \|\varphi\|_{H^2}. \end{aligned}$$

Proof. By Lemma 7.1, we have for φ even,

$$\frac{\sin y_c}{2} \partial_{y_c} \left(\frac{H(\varphi)(y_c)}{\sin y_c} \right) = \frac{1}{2} ((H\varphi)'(y_c) - \cot y_c H\varphi(y_c))$$
$$= \frac{1}{2} H(\varphi' - \cot y_c \varphi)(y_c) + \frac{1}{2} \int_{-\pi}^{\pi} \varphi(y) dy.$$

Let $\operatorname{In}(\varphi)(y) = \operatorname{Int}(\varphi)(y) - \frac{\operatorname{Int}(\varphi)(\pi)}{2}(1 - \cos y) = \int_0^y \varphi(y') \, dy' - \sin^2 \frac{y}{2} \int_0^\pi \varphi(y') \, dy'$ for $y \in [0, \pi]$. We extend $\operatorname{In}(\varphi)$ to be an even π -periodic function. Then we get by Remark 5.2 that

(7.1)
$$\begin{aligned} \mathrm{II}_{1,1}(\varphi)(y_c) &= -\frac{1}{2\sin y_c} \partial_{y_c} \Big(\frac{H(\mathrm{In}(\varphi))(y_c)}{\sin y_c} \Big) \\ &= -\frac{1}{2\sin^2 y_c} \mathcal{H}(\varphi)(y_c) - \frac{1}{2\sin^2 y_c} \int_{-\pi}^{\pi} \mathrm{In}(\varphi)(y') dy'. \end{aligned}$$

Here $\mathcal{H} = H \circ Z \circ \text{In.}$ Then by Lemma 7.3, we get

$$\begin{aligned} \|\rho \mathrm{II}_{1,1}(\varphi)\|_{H^k_{y_c}} &\leq C \left(\|\mathcal{H}(\varphi)\|_{H^k(\mathbb{T})} + \|\varphi\|_{L^2} \right) \\ &\leq C \left(\|Z \circ \mathrm{In}(\varphi)\|_{H^k(\mathbb{T})} + \|\varphi\|_{L^2} \right) \\ &\leq C \left(\|\mathrm{In}(\varphi)\|_{H^{k+1}(0,\pi)} + \|\varphi\|_{L^2} \right) \leq C \|\varphi\|_{H^k} \end{aligned}$$

As $\partial_{y_c}(\rho \operatorname{II}_{1,1}(\varphi)) = H\Big(\partial_y(Z \circ \operatorname{In}(\varphi))\Big), H\Big(\partial_y(Z \circ \operatorname{In}(\varphi))\Big) \in H^1(\mathbb{T})$ is odd, and by Lemma 7.3, we have

$$\left\|\partial_{y_c}(\rho \mathrm{II}_{1,1})/\sin y_c\right\|_{L^2_{y_c}} \le C \left\|H\left(\partial_y(Z \circ \mathrm{In}(\varphi))\right)\right\|_{H^1} \le C \|\mathrm{In}(\varphi)\|_{H^3} \le C \|\varphi\|_{H^2}.$$

This proves the lemma.

The following lemma shows that for φ vanishing at 0 and π , we have better estimates, which will be used in the odd case.

Lemma 7.5. If $\varphi \in H_0^1(0,\pi)$, then

$$\|\sin y_c \mathrm{II}_{1,1}(\varphi)\|_{L^2_{y_c}} \le C \|\varphi\|_{H^1}.$$

If $\varphi \in H^2(0,\pi) \cap H^1_0(0,\pi)$, then

$$\|\sin y_{c}\mathrm{II}_{1,1}(\varphi)\|_{L^{\infty}} + \|\mathrm{II}_{1,1}(\varphi)\|_{L^{2}_{y_{c}}} \leq C \|\varphi\|_{H^{2}}.$$

$$If \varphi \in H^{2}(0,\pi) \cap H^{1}_{0}(0,\pi) \text{ and } \varphi'' / \sin y_{c} \in L^{2}(0,\pi), \text{ then} \\ \|\partial_{y_{c}} \mathrm{II}_{1,1}(\varphi)\|_{L^{2}_{y_{c}}} + \|\mathrm{II}_{1,1}(\varphi)\|_{L^{\infty}} + \|\partial^{2}_{y_{c}}(\sin y_{c}\mathrm{II}_{1,1}(\varphi))\|_{L^{2}_{y_{c}}} \\ + \|\sin y_{c}\partial_{y_{c}}\mathrm{II}_{1,1}(\varphi)\|_{L^{\infty}} \leq C (\|\varphi\|_{H^{2}} + \|\varphi'' / \sin y_{c}\|_{L^{2}_{y_{c}}}).$$

Proof. For $\varphi \in H^1_0(0,\pi)$, we get by Hardy's inequality that

$$\left\|\frac{\operatorname{In}(\varphi)}{\sin^2 y}\right\|_{L^2} + \left\|\frac{\partial_y \operatorname{In}(\varphi)}{\sin y}\right\|_{L^2} + \left\|\partial_y \left(\frac{\operatorname{In}(\varphi)}{\sin y}\right)\right\|_{L^2} \le C \|\varphi\|_{H^1}.$$

Recall that

(7.2)
$$\operatorname{II}_{1,1}(\varphi)(y_c) = -\frac{1}{2\sin y_c} \partial_{y_c} \Big(\frac{H(\operatorname{In}(\varphi))}{\sin y_c} \Big),$$

and $\frac{\ln(\varphi)}{\sin y}$ is odd, by Lemma 7.1, we have

$$\sin y_c \operatorname{II}_{1,1}(\varphi)(y_c) = -\frac{1}{2} \partial_{y_c} H\Big(\frac{\operatorname{In}(\varphi)}{\sin y}\Big) = -\frac{1}{2} H\Big(\partial_y \Big(\frac{\operatorname{In}(\varphi)}{\sin y}\Big)\Big),$$

which implies that for $\varphi \in H_0^1(0,\pi)$,

$$\|\sin y_c \mathrm{II}_{1,1}(\varphi)\|_{L^2_{y_c}} \le C \|\varphi\|_{H^1}.$$

For
$$\varphi \in H^2(0,\pi) \cap H^1_0(0,\pi)$$
, $\partial_y^2\left(\frac{\ln(\varphi)}{\sin y}\right) \in L^2(\mathbb{T})$ and

$$\partial_{y_c} \big(\sin y_c \mathrm{II}_{1,1}(\varphi)(y_c) \big) = -\frac{1}{2} H \Big(\partial_y^2 \big(\frac{\mathrm{In}(\varphi)}{\sin y} \big) \Big).$$

which shows that

$$\left\|\partial_{y_c}\left(\sin y_c \operatorname{II}_{1,1}(\varphi)(y_c)\right)\right\|_{L^2_{y_c}} \le C \left\|\partial_y^2\left(\frac{\operatorname{In}(\varphi)}{\sin y}\right)\right\|_{L^2} \le C \|\varphi\|_{H^2},$$

which implies that

$$\|\sin y_c \mathrm{II}_{1,1}(\varphi)\|_{L^{\infty}} + \|\mathrm{II}_{1,1}(\varphi)\|_{L^2_{y_c}} \le C \|\varphi\|_{H^2}.$$

For f(0) = f'(0) = f''(0) = 0 and $f'''/y \in L^2$, we have $\left\|\frac{f}{y^4}\right\|_{L^2} + \left\|\frac{f'}{y^3}\right\|_{L^2} + \left\|\frac{f''}{y^3}\right\|_{L^2} \le C \|f'''/y\|_{L^2}.$

For $\varphi \in H^2(0,\pi) \cap H^1_0(0,\pi)$ and $\varphi''/u' \in L^2(0,\pi)$, let

 $\operatorname{In}_2(\varphi) = \operatorname{In}(\varphi) - (\sin y)^2 (\varphi'(0)(1 + \cos y) + \varphi'(\pi)(1 - \cos y) + \operatorname{Int}(\varphi)(\pi) \cos y) / 4,$ then $\operatorname{In}_2(\varphi) = \operatorname{In}_2(\varphi)' = \operatorname{In}_2(\varphi)'' = 0 \text{ for } y = 0, \pi.$ Thus,

$$\left\|\frac{\mathrm{In}_{2}(\varphi)}{\sin^{4} y}\right\|_{L^{2}} + \left\|\frac{\mathrm{In}_{2}(\varphi)'}{\sin^{3} y}\right\|_{L^{2}} + \left\|\frac{\mathrm{In}_{2}(\varphi)''}{\sin^{2} y}\right\|_{L^{2}} \le C\left(\|\mathrm{In}_{2}(\varphi)''/\sin y\|_{L^{2}} + \|\mathrm{In}_{2}(\varphi)\|_{H^{2}}\right),$$

which implies that

$$\left\| \partial_y^3 \left(\frac{\ln(\varphi)}{\sin y} \right) \right\|_{L^2} \le \left\| \partial_y^3 \left(\frac{\ln_2(\varphi)}{\sin y} \right) \right\|_{L^2} + C \|\varphi\|_{H^2} \le C \left(\|\varphi'' / \sin y\|_{L^2} + \|\varphi\|_{H^2} \right).$$

Therefore, by Lemma 7.3,

$$\begin{aligned} \|\partial_{y_c} \mathrm{II}_{1,1}(\varphi)\|_{L^2_{y_c}} &\leq C \Big\| \frac{Z \circ H\big(\partial_y (\mathrm{In}(\varphi)/\sin y)\big)}{\sin y} \Big\|_{L^2} \\ &\leq C \|H\big(\partial_y (\mathrm{In}(\varphi)/\sin y)\big)\|_{H^2} \\ &\leq C \|\mathrm{In}(\varphi)/\sin y\|_{H^3} \leq C\big(\|\varphi''/\sin y\|_{L^2} + \|\varphi\|_{H^2}\big), \end{aligned}$$

as well as

$$\|\partial_{y_c}^2(\sin y_c \mathrm{II}_{1,1}(\varphi))\|_{L^2_{y_c}} \le \|H(\partial_y^3(\mathrm{In}(\varphi/\sin y)))\|_{L^2} \le C(\|\varphi''/\sin y\|_{L^2} + \|\varphi\|_{H^2}),$$

which implies that $\|\sin y_c \partial_{y_c} \mathrm{II}_{1,1}(\varphi)\|_{L^{\infty}} \le C(\|\varphi\|_{H^2} + \|\varphi''/u'\|_{L^2}).$

7.3. Estimates of \mathcal{L}_k . In this subsection, we denote by $\phi(y,c)$ a solution of (2.6) obtained by Proposition 3.6 and let $\phi_1(y,c) = \frac{\phi(y,c)}{u(y)-c}$ for $c \in D_0$. Here $u(y) = -\cos y$. We denote $\|\cdot\|_{L^p} = \|\cdot\|_{L^p(0,\pi)}$. Recall that

$$\mathcal{L}_k(\varphi)(y_c) = \int_0^\pi \int_{y_c}^z \varphi(y) \left(\frac{\partial_z + \partial_y}{u'(y_c)} + \partial_c\right)^k \left(\frac{1}{(u(z) - c)^2} \left(\frac{\phi_1(y, c)}{\phi_1(z, c)^2} - 1\right)\right) dy dz$$

Lemma 7.6. For any $p \in [1, \infty)$, there exists a constant C independent of α such that for k = 0, 1, 2,

$$\begin{aligned} \|u'(y_c)^{2k+2}\mathcal{L}_k(\varphi)\|_{L^{\infty}} &\leq C |\alpha|^{\frac{1}{p}} \|\varphi\|_{L^p} ,\\ \|u'(y_c)^{2k+1}\mathcal{L}_k(\varphi)\|_{L^{\infty}} &\leq C |\alpha|^{1+\frac{1}{p}} \|\varphi\|_{L^p} ,\\ \|u'(y_c)^{2k+1}\mathcal{L}_k(\varphi)\|_{L^{\infty}} &\leq C |\alpha| \|\varphi\|_{L^{\infty}}. \end{aligned}$$

Proof. By Proposition 3.8, we know that for k = 0, 1, 2, and $0 \le z \le y \le y_c \le \pi$ or $0 \le y_c \le y \le z \le \pi$,

(7.3)
$$\left| \left(\frac{\partial_z + \partial_y}{u'(y_c)} + \partial_c \right)^k \left(\frac{1}{(u(z) - c)^2} \left(\frac{\phi_1(y, c)}{\phi_1(z, c)^2} - 1 \right) \right) \right| \le C \frac{\min\{\alpha^2 | z - y_c|^2, 1\}}{(u(z) - c)^2 u'(y_c)^{2k}}.$$

Recall that $u_1(y,c) = \frac{\cos y_c - \cos y}{y - y_c}$ and

$$C^{-1}\sin\frac{y+y_c}{2} \le u_1(y,c) \le C\sin\frac{y+y_c}{2}.$$

Thus,

$$\begin{split} &\int_{0}^{\pi} \frac{\left|z - y_{c}\right|^{1 - \frac{1}{p}} \min\{\alpha^{2} | z - y_{c}|^{2}, 1\}}{\left|u(z) - c\right|^{2}} \, dz \\ &\leq C \int_{0}^{\pi} \frac{\min\{\alpha^{2} | z - y_{c}|^{2}, 1\}}{\sin^{2} \frac{z + y_{c}}{2} | z - y_{c}|^{1 + \frac{1}{p}}} \, dz \\ &\leq C \int_{|z - y_{c}| \leq \frac{1}{\alpha}, 0 \leq z \leq \pi} \frac{\alpha^{2} |z - y_{c}|^{1 - \frac{1}{p}}}{\sin^{2} \frac{z + y_{c}}{2}} \, dz + C \int_{|z - y_{c}| > \frac{1}{\alpha}, 0 \leq z \leq \pi} \frac{|z - y_{c}|^{-1 - \frac{1}{p}}}{\sin^{2} \frac{z + y_{c}}{2}} \, dz. \end{split}$$

Using the fact that

$$\int_{|z-y_c| \le \frac{1}{\alpha}, 0 \le z \le \pi} \frac{\alpha^2 |z-y_c|^{1-\frac{1}{p}}}{\sin^2 \frac{z+y_c}{2}} dz \le C \min \Big\{ \frac{\alpha^{1+\frac{1}{p}}}{\sin y_c}, \frac{\alpha^{\frac{1}{p}}}{\sin^2 y_c} \Big\},$$

$$\int_{|z-y_c| > \frac{1}{\alpha}, 0 \le z \le \pi} \frac{|z-y_c|^{-1-\frac{1}{p}}}{\sin^2 \frac{z+y_c}{2}} dz \le C \min\left\{\frac{\alpha^{1+\frac{1}{p}}}{\sin y_c}, \frac{\alpha^{\frac{1}{p}}}{\sin^2 y_c}\right\},$$

and for $p = \infty$,

$$\begin{split} &\int_{0}^{\pi} \frac{|z - y_{c}| \min\{\alpha^{2}|z - y_{c}|^{2}, 1\}}{|u(z) - c|^{2}} dz \\ &\leq C \int_{|z - y_{c}| \leq \frac{1}{\alpha}, 0 \leq z \leq \pi} \frac{\alpha^{2}|z - y_{c}|}{\sin^{2} \frac{z + y_{c}}{2}} dz + C \int_{|z - y_{c}| > \frac{1}{\alpha}, 0 \leq z \leq \pi} \frac{|z - y_{c}|^{-1}}{\sin^{2} \frac{z + y_{c}}{2}} dz \\ &\leq C \frac{|\alpha|}{\sin y_{c}}, \end{split}$$

we deduce that for k = 0, 1, 2, and $p \in [1, \infty)$,

$$\begin{aligned} \left| u'(y_c)^{2k} \mathcal{L}_k(\varphi)(c) \right| &\leq C \int_0^\pi \frac{\left| z - y_c \right|^{1 - \frac{1}{p}} \min\{\alpha^2 | z - y_c |^2, 1\}}{|u(z) - c|^2} dz \, \|\varphi\|_{L^p} \\ &\leq C \min\left\{ \frac{\alpha^{1 + \frac{1}{p}}}{u'(y_c)}, \frac{\alpha^{\frac{1}{p}}}{u'(y_c)^2} \right\} \|\varphi\|_{L^p} \,, \end{aligned}$$

and

(7.4)
$$|u'(y_c)^{2k} \mathcal{L}_k(\varphi)| \le C \frac{|\alpha|}{\sin y_c} \|\varphi\|_{L^{\infty}}.$$

This proves the lemma.

To estimate the derivatives of \mathcal{L}_0 , we introduce for k, j = 0, 1,

(7.5)
$$I_{k,j}(\varphi)(c) = \int_{y_c}^{z} \varphi(y) \left(\frac{\partial_z + \partial_y}{u'(y_c)} + \partial_c\right)^k \left(\frac{1}{(u(z) - c)^2} \left(\frac{\phi_1(y, c)}{\phi_1(z, c)^2} - 1\right)\right) dy \Big|_{z=j\pi}.$$

Thus, for k = 0, 1,

(7.6)
$$\partial_c \mathcal{L}_k(\varphi) = \frac{1}{u'(y_c)} \left(\mathcal{L}_k(\varphi') - I_{k,1}(\varphi) + I_{k,0}(\varphi) \right) + \mathcal{L}_{k+1}(\varphi).$$

Lemma 7.7. It holds that

$$\begin{aligned} \left\| u'(y_c)^3 \partial_c \mathcal{L}_0(\varphi) \right\|_{L^{\infty}} &\leq C \alpha \|\varphi\|_{L^{\infty}} + C \alpha^{\frac{1}{2}} \|\varphi'\|_{L^2}, \\ \|u'(y_c)^3 \partial_c \mathcal{L}_0(\varphi)\|_{L^{\infty}} &\leq C \alpha \|\varphi\|_{L^{\infty}} + C \alpha u'(y_c) \|\varphi'\|_{L^{\infty}}. \end{aligned}$$

1

If $\varphi/u' \in L^p(0,\pi)$ for $p \in (1,\infty)$, then

$$\begin{aligned} \|\sin y_c \mathcal{L}_0(\varphi)\|_{L^{\infty}} &\leq C\alpha^{\frac{1}{p}} \|\varphi\|_{W^{1,p}}, \\ \|\mathcal{L}_0(\varphi)\|_{L^{\infty}} &\leq C\alpha \|\varphi\|_{W^{1,\infty}}, \\ \|\sin^3 y_c \partial_c \mathcal{L}_0(\varphi)\|_{L^{\infty}} &\leq C\alpha^{\frac{1}{2}} \|\varphi\|_{H^1}. \end{aligned}$$

Proof. By (7.3), we have for k = 0, 1,

$$|I_{k,j}(\varphi)(c)| \le C \left| \int_{y_c}^{j\pi} \varphi(y) dy \right| \frac{\min\{\alpha^2 | j\pi - y_c|^2, 1\}}{(\cos j\pi - \cos y_c)^2 u'(y_c)^{2k}} \\ \le C \frac{|j\pi - y_c| \min\{\alpha^2 | j\pi - y_c|^2, 1\}}{(\cos j\pi - \cos y_c)^2 u'(y_c)^{2k}} \|\varphi\|_{L^{\infty}}$$

(7.7)
$$\leq C \frac{\min\{\alpha^2 | j\pi - y_c|^2, 1\}}{| j\pi - y_c|^3 u'(y_c)^{2k}} \|\varphi\|_{L^{\infty}},$$

thus by the fact that $\sin y_c \leq C |j\pi - y_c|$, we deduce that for j = 0, 1,

(7.8)
$$|u(y_c)^{2k+3}I_{k,j}(\varphi)(c)| \le C \|\varphi\|_{L^{\infty}},$$

(7.9)
$$|u(y_c)^{2k+2}I_{k,j}(\varphi)(c)| \le C\alpha \|\varphi\|_{L^{\infty}}.$$

Notice that

(7.10)
$$\partial_c \mathcal{L}_0(\varphi) = \frac{1}{u'(y_c)} \left(\mathcal{L}_0(\varphi') - I_{0,1}(\varphi) + I_{0,0}(\varphi) \right) + \mathcal{L}_1(\varphi).$$

We get by Lemma 7.6 that

$$\begin{aligned} \|u'(y_c)^2 \mathcal{L}_0(\varphi')\|_{L^{\infty}} &\leq C \alpha^{\frac{1}{2}} \|\varphi'\|_{L^2}, \\ \|u'(y_c)^3 \mathcal{L}_1(\varphi)\|_{L^{\infty}} &\leq C \alpha \|\varphi\|_{L^{\infty}}, \end{aligned}$$

which along with the estimates of ${\cal I}_{0,0}$ and ${\cal I}_{0,1}$ imply that

$$\|u'(y_c)^3 \partial_c \mathcal{L}_0(\varphi)\|_{L^{\infty}} \le C \alpha \|\varphi\|_{L^{\infty}} + C \alpha^{\frac{1}{2}} \|\varphi'\|_{L^2}.$$

By (7.4), we get

$$\|u'(y_c)\mathcal{L}_0(\varphi')\|_{L^{\infty}} \le C|\alpha| \|\varphi'\|_{L^{\infty}}.$$

This along with the estimates of $I_{0,0}$ and $I_{0,1}$ gives

$$\|u'(y_c)^3 \partial_c \mathcal{L}_0(\varphi)\|_{L^{\infty}} \le C \alpha \|\varphi\|_{L^{\infty}} + C \alpha u'(y_c) \|\varphi'\|_{L^{\infty}}.$$

If $\varphi/u' \in L^p$, then $\frac{\varphi(y)}{u'(y)} = \frac{1}{u'(y)} \int_{j\pi}^y \varphi'(z) dz$, thus, $\left\| \frac{\varphi(y)}{u'(y)} \right\|_{L^p(0,\pi)} \leq C \|\varphi'\|_{L^p(0,\pi)}$ for any $p \in (1,\infty]$. Due to $u(y) - c = u_1(y,c)(y-y_c)$, we have

$$\left| \int_{y_c}^{z} u'(y)^{p'} dy \right|^{\frac{1}{p'}} \le C \max_{|2y-y_c-z| \le |z-y_c|} u'(y) |z-y_c|^{\frac{1}{p'}} \le C u_1(z,c) |z-y_c|^{\frac{1}{p'}}.$$

Then we have

$$\begin{aligned} |\mathcal{L}_{k}(\varphi)(c)| &\leq C \int_{0}^{\pi} \frac{\min\{\alpha^{2}|z-y_{c}|^{2},1\} \left| \int_{y_{c}}^{z} u'(y)^{p'} dy \right|^{\frac{1}{p'}}}{u_{1}(z,c)^{2}(z-y_{c})^{2} u'(y_{c})^{2k}} dz \left\| \frac{\varphi(y)}{u'(y)} \right\|_{L^{p}} \\ &\leq C \int_{0}^{\pi} \frac{\min\{\alpha^{2}|z-y_{c}|^{2},1\}}{u'(y_{c})^{2k+1}|z-y_{c}|^{1+\frac{1}{p}}} \|\varphi'\|_{L^{p}} \leq C \frac{\alpha^{\frac{1}{p}}}{u'(y_{c})^{2k+1}} \|\varphi'\|_{L^{p}}, \end{aligned}$$

and similarly,

(7.11)
$$\|u'(y_c)^{2k+1}\mathcal{L}_k(\varphi)\|_{L^{\infty}} \le C\alpha \|\varphi\|_{L^{\infty}}.$$

In particular, we get

$$|\sin y_{c}\mathcal{L}_{0}(\varphi)||_{L^{\infty}} \leq C\alpha^{\frac{1}{p}} \|\varphi\|_{W^{1,p}}.$$
As $|u(z) - c| = \left|\sin\frac{y_{c}+z}{2}\sin\frac{y_{c}-z}{2}\right| \geq \left|\sin\frac{y_{c}-z}{2}\right|^{2} \geq C^{-1}|z - y_{c}|^{2}, \text{ we have}$

$$|u'(y_{c})^{2k}\mathcal{L}_{k}(\varphi)(c)| \leq C\int_{0}^{\pi}\frac{\min\{\alpha^{2}|z - y_{c}|^{2}, 1\}\int_{y_{c}}^{z}u'(y)dy}{(u(z) - c)^{2}}dz \left\|\frac{\varphi(y)}{u'(y)}\right\|_{L^{\infty}}$$

$$(7.12) \leq C\int_{0}^{\pi}\frac{\min\{\alpha^{2}|z - y_{c}|^{2}, 1\}}{|u(z) - c|}dz \|\varphi'\|_{L^{\infty}}$$

$$\leq C \int_0^{\pi} \frac{\min\{\alpha^2 | z - y_c|^2, 1\}}{|z - y_c|^2} dz \|\varphi'\|_{L^{\infty}} \leq C |\alpha| \|\varphi'\|_{L^{\infty}},$$

In particular, we get

$$\|\mathcal{L}_0(\varphi)\|_{L^{\infty}} \le C\alpha \|\varphi\|_{W^{1,\infty}}.$$

On the other hand, we have

$$|u'(y_c)^{2k+2}I_{k,0}(\varphi)(c)| \le C \frac{u'(y_c)^{2k+2}y_c^{2-\frac{1}{p}}\min\{\alpha^2 y_c^2, 1\}}{u'(y_c)^{2k+2}y_c^2} \left\|\frac{\varphi}{u'}\right\|_{L^p} \le C\alpha^{\frac{1}{p}} \|\varphi\|_{W^{1,p}(0,\pi)}.$$

The bound of $I_{k,1}$ is similar. Then we can conclude the last inequality of the lemma by using (7.10).

Lemma 7.8. It holds that

 $\left|u'(y_c)^5 \partial_c^2 \mathcal{L}_0(\varphi)(c)\right| \le C \left(\alpha^{\frac{1}{2}} u'(y_c) \|\varphi''\|_{L^2} + \alpha \|\varphi\|_{L^{\infty}} + \alpha u'(y_c) \|\varphi'\|_{L^{\infty}} + \alpha^{\frac{1}{2}} \|\varphi'\|_{L^2}\right).$ If $\varphi/u' \in L^{\infty}$, then for k = 0, 1, 2,

$$\|u'(y_c)^{2k}\partial_c^k \mathcal{L}_0(\varphi)\|_{L^{\infty}} \le C\alpha \|\varphi'\|_{L^{\infty}} + C\alpha^{\frac{1}{2}} \|\varphi''\|_{L^2}.$$

Proof. Since we have

$$\begin{aligned} \partial_c I_{0,j}(\varphi)(c) &= -\frac{\varphi(y_c)}{u'(y_c)} \frac{1}{(u(z) - c)^2} \left(\frac{1}{\phi_1(z,c)^2} - 1 \right) \Big|_{z=j\pi} \\ &+ \int_{y_c}^{j\pi} \partial_c \left(\frac{\varphi(y)}{(u(z) - c)^2} \left(\frac{\phi_1(y,c)}{\phi_1(z,c)^2} - 1 \right) \right) \, dy \Big|_{z=j\pi}, \end{aligned}$$

and by Proposition 3.8, we get for $|2y - y_c - z| \le |z - y_c|$,

$$\begin{aligned} \left| \partial_c \left(\frac{1}{(u(z) - c)^2} \left(\frac{\phi_1(y, c)}{\phi_1(z, c)^2} - 1 \right) \right) \right| \\ &\leq C \left| \left(\frac{1}{|u(z) - c|^3} \left(\frac{\phi_1(y, c)}{\phi_1(z, c)^2} - 1 \right) \right) \right| + C \left| \frac{1}{|u(z) - c|^2} \partial_c \left(\frac{\phi_1(y, c)}{\phi_1(z, c)^2} \right) \right| \\ &\leq C \frac{\min\{\alpha^2 | z - y_c|^2, 1\}}{|u(z) - c|^3} + C \frac{\min\{\alpha^2 | z - y_c|, \alpha\}}{u'(y_c)|u(z) - c|^2}. \end{aligned}$$

Thus, we obtain

$$\begin{aligned} |\partial_c I_{0,1}(\varphi)(c)| &\leq C \|\varphi\|_{L^{\infty}} \frac{\min\{\alpha^2(\pi - y_c)^2, 1\}}{(\pi - y_c)^2 u'(y_c)^3} \\ &+ C \left| \partial_c \left(\frac{1}{(u(\pi) - c)^2} \left(\frac{\phi_1(y, c)}{\phi_1(\pi, c)^2} - 1 \right) \right) \right| |\pi - y_c| \|\varphi\|_{L^{\infty}} \\ &\leq C \|\varphi\|_{L^{\infty}} \frac{\min\{\alpha^2(\pi - y_c)^2, 1\}}{(\pi - y_c)^2 u'(y_c)^3} + C \|\varphi\|_{L^{\infty}} \frac{\min\{\alpha^2(\pi - y_c), \alpha\}}{|\pi - y_c| u'(y_c)^3} \end{aligned}$$

and

$$|\partial_c I_{0,0}(\varphi)(c)| \le C \|\varphi\|_{L^{\infty}} \Big(\frac{\min\{\alpha^2 y_c^2, 1\}}{y_c^2 u'(y_c)^3} + \frac{\min\{\alpha^2 y_c, \alpha\}}{y_c u'(y_c)^3} \Big).$$

This gives

$$|\partial_c I_{0,1}(\varphi)(c)| + |\partial_c I_{0,0}(\varphi)(c)| \le C \|\varphi\|_{L^{\infty}} \Big(\frac{\min\{\alpha^2 \sin^2 y_c, 1\}}{\sin^5 y_c} + \frac{\min\{\alpha^2 \sin^2 y_c, \alpha \sin y_c\}}{\sin^5 y_c} \Big).$$

Therefore, for any $\gamma \in [0, 1]$,

(7.13)
$$|u'(y_c)^{3+\gamma}\partial_c I_{0,0}(\varphi)(c)| + |u'(y_c)^{3+\gamma}\partial_c I_{0,1}(\varphi)(c)| \le C\alpha^{2-\gamma} \|\varphi\|_{L^{\infty}}.$$

Notice that

$$\begin{aligned} \partial_c^2 \mathcal{L}_0(\varphi) &= \partial_c \left(\frac{1}{u'(y_c)} \left(\mathcal{L}_0(\varphi') - I_{0,1}(\varphi) + I_{0,0}(\varphi) \right) + \mathcal{L}_1(\varphi) \right) \\ &= \frac{1}{u'(y_c)^2} \left(\mathcal{L}_0(\varphi'') - I_{0,1}(\varphi') + I_{0,0}(\varphi') \right) + \frac{1}{u'(y_c)} \mathcal{L}_1(\varphi') \\ &- \frac{1}{u'(y_c)} \partial_c (I_{0,1}(\varphi)(c) - I_{0,0}(\varphi)(c)) \\ &- \frac{u''(y_c)}{u'(y_c)^3} \left(\mathcal{L}_0(\varphi') - I_{0,1}(\varphi) + I_{0,0}(\varphi) \right) \\ &+ \frac{1}{u'(y_c)} \left(\mathcal{L}_1(\varphi') - I_{1,1}(\varphi) + I_{1,0}(\varphi) \right) + \mathcal{L}_2(\varphi). \end{aligned}$$

Thus, by Lemma 7.6, (7.8), (7.9), (7.11) and (7.13), we obtain

$$|\sin^5 y_c \partial_c^2 \mathcal{L}_0(\varphi)(c)| \le C\alpha^{\frac{1}{2}} u'(y_c) \|\varphi''\|_{L^2} + C\alpha \|\varphi\|_{L^{\infty}} + C\alpha u'(y_c) \|\varphi'\|_{L^{\infty}} + C\alpha^{\frac{1}{2}} \|\varphi'\|_{L^2}.$$

If $\varphi(0) = \varphi(\pi) = 0$, then

$$\left|\partial_c I_{0,0}(\varphi)(c)\right| \le C \left\|\frac{\varphi}{u'}\right\|_{L^{\infty}} \left(\frac{\min\{\alpha^2 y_c^2, 1\}}{y_c^2 u'(y_c)^2} + \frac{\min\{\alpha^2 y_c, \alpha\}}{y_c u'(y_c)^2}\right)$$

Therefore,

$$|u'(y_c)^3 \partial_c I_{0,0}(\varphi)(c)| \le C \alpha \left\| \frac{\varphi}{u'} \right\|_{L^{\infty}}.$$

Similarly, we have

$$|u'(y_c)^3 \partial_c I_{0,1}(\varphi)(c)| \le C \alpha \left\| \frac{\varphi}{u'} \right\|_{L^{\infty}}.$$

We also have

$$\|u'(y_c)^{2k+1}I_{k,j}(\varphi)\|_{L^{\infty}} \le C\alpha \left\|\frac{\varphi}{u'}\right\|_{L^{\infty}}.$$

With the estimates above, we deduce the second inequality of the lemma by using (7.12), (7.9) and Lemma 7.6.

8. $W^{2,1}$ estimates of K_o and K_e

This section is devoted to the proof of Proposition 6.1 and Proposition 6.2. For the sake of simplicity, we introduce some notations:

- We use \mathcal{L}^p to denote a function f which satisfies $||f||_{L^p_{yc}} \leq C$.
- We use $\mathcal{L}^p \cap \mathcal{L}^q$ to denote a function f which satisfies $\|f\|_{L^p_{y_c}} + \|f\|_{L^q_{y_c}} \leq C$.
- We use $\rho \mathcal{L}^p$ denote a function f which satisfies $\left\| \frac{f}{\rho} \right\|_{L^p_{y_c}} \leq C$.
- We use $\mathcal{L}^p + \mathcal{L}^q$ to denote a function f which can be divided into two parts $f = f_1 + f_2$ with f_1, f_2 satisfing $||f_1||_{L^p_{y_c}} \leq C, ||f_2||_{L^q_{y_c}} \leq C$.

8.1. $W^{2,1}$ estimate of K_o . In this subsection, we prove Proposition 6.1. Recall that

$$K_o(c,\alpha) = \frac{\Lambda_1(\widehat{\omega}_o)(y_c)\Lambda_2(g)(y_c)}{(\mathbf{A}(c)^2 + \mathbf{B}(c)^2)u'(y_c)},$$

where A, B are defined by (4.5) and Λ_1, Λ_2 are defined by (5.18) and (5.19).

Step 1. L^1 estimate.

We normalize $\widehat{\omega}_0, g$ so that $\|\widehat{\omega}_o\|_{L^2} \leq 1$, $\|g\|_{L^2} \leq 1$. By (5.20), (5.22) and Lemma 7.4, we get $\Lambda_{1,1}(\widehat{\omega}_o)(y_c) = \mathcal{L}^2$ and $\Lambda_{2,1}(g)(y_c) = \mathcal{L}^2$. By (5.21), (5.23) Lemma 4.4 and Lemma 7.6, we get $\Lambda_{1,2}(\widehat{\omega}_o)(y_c) = |\alpha|^{\frac{1}{2}}\mathcal{L}^{\infty} + |\alpha| \sin y_c \mathcal{L}^2$ and $\Lambda_{2,2}(g)(y_c) = |\alpha|^{\frac{1}{2}}\mathcal{L}^{\infty} + |\alpha| \sin y_c \mathcal{L}^2$. Thus, we have

$$\Lambda_1(\widehat{\omega}_o)(y_c) = (1 + |\alpha| \sin y_c)\mathcal{L}^2 + |\alpha|^{\frac{1}{2}}\mathcal{L}^{\infty}, \ \Lambda_2(g)(y_c) = (1 + |\alpha| \sin y_c)\mathcal{L}^2 + |\alpha|^{\frac{1}{2}}\mathcal{L}^{\infty}.$$

Then by Proposition 4.3, we infer that

$$u'(y_c)K_o(c,\alpha) = \mathcal{L}^1 + \frac{|\alpha|^{\frac{1}{2}}\mathcal{L}^2}{1+|\alpha|\sin y_c} + \frac{\alpha\mathcal{L}^{\infty}}{(1+|\alpha|\sin y_c)^2},$$

which implies that

$$||K_o(\cdot, \alpha)||_{L^1_c(-1,1)} \le C ||\widehat{\omega}_o||_{L^2} ||g||_{L^2}.$$

Step 2. $W^{1,1}$ estimate.

We normalize $\widehat{\omega}_0, g$ so that $\|\widehat{\omega}_o\|_{H^1} \leq 1$, $\|g\|_{H^1} \leq 1$. By the definition of $\widehat{\omega}_o$, we know that $\widehat{\omega}_o(\alpha, 0) = \widehat{\omega}_o(\alpha, \pi) = g(0) = g(\pi) = 0$, then $\|\frac{\widehat{\omega}_o}{u'}\|_{L^2} + \|\frac{g}{u'}\|_{L^2} \leq C$. By (5.20), (5.22) and Lemma 7.5, we get

$$\Lambda_{1,1}(\widehat{\omega}_o)(y_c) = \sin y_c \mathcal{L}^2, \quad \Lambda_{2,1}(g)(y_c) = \sin y_c \mathcal{L}^2,$$

and by Lemma 7.4, we get

$$\partial_{y_c} \Lambda_{1,1}(\widehat{\omega}_o)(y_c) = \mathcal{L}^2, \quad \partial_{y_c} \Lambda_{2,1}(g)(y_c) = \mathcal{L}^2.$$

By (5.21), (5.23), Lemma 4.4 and Lemma 7.7, we get

$$\begin{split} \Lambda_{1,2}(\widehat{\omega}_o)(y_c) &= |\alpha|^{\frac{1}{2}} \sin y_c \mathcal{L}^{\infty} + |\alpha| \sin^2 y_c \mathcal{L}^2, \\ \Lambda_{2,2}(g)(y_c) &= |\alpha|^{\frac{1}{2}} \sin y_c \mathcal{L}^{\infty} + |\alpha| \sin^2 y_c \mathcal{L}^2, \\ \partial_c \Lambda_{1,2}(\widehat{\omega}_o)(y_c) &= \frac{\alpha^{\frac{1}{2}} \mathcal{L}^{\infty}}{\sin y_c} + |\alpha| \mathcal{L}^2, \\ \partial_c \Lambda_{2,2}(g)(y_c) &= \frac{\alpha^{\frac{1}{2}} \mathcal{L}^{\infty}}{\sin y_c} + |\alpha| \mathcal{L}^2. \end{split}$$

Thus, we obtain

$$\Lambda_1(\widehat{\omega}_o)(y_c) = \sin y_c((1+|\alpha|\sin y_c)\mathcal{L}^2 + |\alpha|^{\frac{1}{2}}\mathcal{L}^{\infty}),$$

$$\Lambda_2(g)(y_c) = \sin y_c((1+|\alpha|\sin y_c)\mathcal{L}^2 + |\alpha|^{\frac{1}{2}}\mathcal{L}^{\infty}),$$

$$\partial_c\Lambda_1(\widehat{\omega}_o)(y_c) = \frac{1}{\sin y_c} \big((1+|\alpha|\sin y_c)\mathcal{L}^2 + \alpha^{\frac{1}{2}}\mathcal{L}^{\infty}\big),$$

$$\partial_c\Lambda_2(g)(y_c) = \frac{1}{\sin y_c} \big((1+|\alpha|\sin y_c)\mathcal{L}^2 + \alpha^{\frac{1}{2}}\mathcal{L}^{\infty}\big).$$

Therefore,

$$\Lambda_1(\widehat{\omega}_o)\Lambda_2(g) = \sin^2 y_c \left((1+|\alpha|\sin y_c)^2 \mathcal{L}^1 + \alpha^{\frac{1}{2}} (1+|\alpha|\sin y_c) \mathcal{L}^2 + \alpha \mathcal{L}^\infty \right),$$

$$\partial_c(\Lambda_1(\widehat{\omega}_o)\Lambda_2(g)) = (1+|\alpha|\sin y_c)^2 \mathcal{L}^1 + \alpha^{\frac{1}{2}}(1+|\alpha|\sin y_c)\mathcal{L}^2 + \alpha \mathcal{L}^{\infty},$$

which along with Proposition 4.3 give

$$\|\partial_c K_o(\cdot, \alpha)\|_{L^1_c(-1,1)} \le C \|\widehat{\omega}_o\|_{H^1} \|g\|_{H^1}.$$

Step 3. $W^{2,1}$ estimate.

We normalize $\widehat{\omega}_0, g$ so that $\|\widehat{\omega}_0\|_{H^2} \leq 1$, $\|f\|_{L^2} \leq 1$. Then we have

$$\begin{aligned} \|g''\|_{L^2}^2 + 2\alpha^2 \|g'\|_{L^2}^2 + \alpha^4 \|g\|_{L^2}^2 &\leq 1, \\ \|\frac{u''g}{u'}\|_{L^{\infty}} &\leq C \|(u''g)'\|_{L^{\infty}} \leq C/\alpha^{\frac{1}{2}}, \quad \|(u''g)''\|_{L^2} \leq C \end{aligned}$$

By (5.20), (5.22) and Lemma 7.5, we get

 $\Lambda_{1,1}(\widehat{\omega}_o)(y_c) = \sin^2 y_c \mathcal{L}^2 \cap \sin y_c \mathcal{L}^\infty, \quad \Lambda_{2,1}(g)(y_c) = \sin^2 y_c \mathcal{L}^2 \cap \sin y_c \mathcal{L}^\infty.$ By Lemma 7.4, we get

by Lemma 7.4, we get

$$\partial_c \Lambda_{1,1}(\widehat{\omega}_o)(y_c) = \mathcal{L}^2, \quad \partial_c \Lambda_{2,1}(g)(y_c) = \mathcal{L}^2, \\ \partial_c^2 \Lambda_{1,1}(\widehat{\omega}_o)(y_c) = \frac{\mathcal{L}^2}{\sin^2 y_c}, \quad \partial_c^2 \Lambda_{2,1}(g)(y_c) = \frac{\mathcal{L}^2}{\sin^2 y_c}$$

By (5.21), (5.23), Lemma 4.4 and Lemma 7.8, we get

$$\begin{split} \Lambda_{1,2}(\widehat{\omega}_o)(y_c) &= |\alpha| \sin^2 y_c \mathcal{L}^{\infty}, \quad \Lambda_{1,2}(g)(y_c) = |\alpha|^{\frac{1}{2}} \sin^2 y_c \mathcal{L}^{\infty}, \\ \partial_c \Lambda_{1,2}(\widehat{\omega}_o)(y_c) &= |\alpha| \mathcal{L}^{\infty}, \quad \partial_c \Lambda_{1,2}(g)(y_c) = |\alpha|^{\frac{1}{2}} \mathcal{L}^{\infty}, \\ \partial_c^2 \Lambda_{1,2}(\widehat{\omega}_o)(y_c) &= \frac{|\alpha| \mathcal{L}^{\infty}}{\sin^2 y_c} + \frac{|\alpha| \mathcal{L}^2}{\sin y_c}, \quad \partial_c^2 \Lambda_{1,2}(g)(y_c) = \frac{|\alpha|^{\frac{1}{2}} \mathcal{L}^{\infty}}{\sin^2 y_c} + \frac{|\alpha| \mathcal{L}^2}{\sin y_c}. \end{split}$$

Therefore, we obtain

$$\begin{split} \Lambda_1(\widehat{\omega}_o) &= \sin^2 y_c \mathcal{L}^2 \cap \sin y_c \mathcal{L}^\infty + |\alpha| \sin^2 y_c \mathcal{L}^\infty, \\ \partial_c(\Lambda_1(\widehat{\omega}_o)) &= \mathcal{L}^2 + \alpha \mathcal{L}^\infty, \\ \partial_c^2(\Lambda_1(\widehat{\omega}_o)) &= \frac{(1+|\alpha| \sin y_c) \mathcal{L}^2 + |\alpha| \mathcal{L}^\infty}{\sin^2 y_c}, \\ \Lambda_2(g) &= \sin^2 y_c (\mathcal{L}^2 + |\alpha|^{\frac{1}{2}} \mathcal{L}^\infty), \\ \partial_c(\Lambda_2(g)) &= \mathcal{L}^2 + |\alpha|^{\frac{1}{2}} \mathcal{L}^\infty, \\ \partial_c^2(\Lambda_2(g)) &= \frac{(1+|\alpha| \sin y_c) \mathcal{L}^2 + |\alpha|^{\frac{1}{2}} \mathcal{L}^\infty}{\sin^2 y_c}. \end{split}$$

Thus,

$$\Lambda_1(\widehat{\omega}_o)\Lambda_2(g) = \sin^4 y_c \left(\mathcal{L}^1 + \alpha \mathcal{L}^2 + \alpha^{\frac{3}{2}} \mathcal{L}^\infty\right),\\ \partial_c \left(\Lambda_1(\widehat{\omega}_o)\Lambda_2(g)\right) = \sin^2 y_c \left(\mathcal{L}^1 + \alpha \mathcal{L}^2 + \alpha^{\frac{3}{2}} \mathcal{L}^\infty\right),\\ \partial_c^2 \left(\Lambda_1(\widehat{\omega}_o)\Lambda_2(g)\right) = (1 + \alpha \sin y_c) \left(\mathcal{L}^1 + \alpha \mathcal{L}^2 + \alpha^2 \mathcal{L}^\infty\right),$$

from which and Proposition 4.3, we infer that

$$\|\partial_c^2 K_o(\cdot, \alpha)\|_{L^1_c(-1,1)} \le C \alpha^{\frac{1}{2}} \|\widehat{\omega}_o\|_{H^2} \|f\|_{L^2},$$

and

$$K_o(c,\alpha) = C(\alpha)\sin^2 y_c \mathcal{L}^2,$$

which implies that $K_o(\pm 1, \alpha) = 0$.

8.2. $W^{2,1}$ estimate of K_e . In this subsection, we prove Proposition 6.2. Recall that

$$K_e(c,\alpha) = \frac{\Lambda_3(\widehat{\omega}_e)(y_c)\Lambda_4(g)(y_c)}{u'(y_c)(\Lambda_1(c)^2 + B_1(c)^2)},$$

where A_1, B_2 are defined by (4.7), and Λ_3, Λ_4 are defined by (5.24) and (5.25).

We need the following lemma.

Lemma 8.1. Let $\phi(y,c)$ be as in Proposition 3.6 with $(y,c) \in [0,\pi] \times [-1,1]$, and $\Lambda_{3,1}(\widehat{\omega}_e)$ and $\Lambda_{4,1}(g)$ be defined by (5.26) and (5.27) with \mathbf{E}_j defined by (5.5) for j = 0, 1. It holds that (1) if $\|\widehat{\omega}_e\|_{L^2} \leq 1$ and $\|g\|_{L^2} \leq 1$, then

$$\Lambda_{3,1}(\widehat{\omega}_e) = rac{\mathcal{L}^2}{lpha^2}, \quad \Lambda_{4,1}(g) = rac{\mathcal{L}^2}{lpha^2};$$

(2) if $\|\widehat{\omega}_e\|_{H^1} \leq 1$ and $|\alpha| \|g\|_{L^2} + \|g'\|_{L^2} \leq 1$, then

$$\Lambda_{3,1}(\widehat{\omega}_e) = \frac{\sin y_c(\mathcal{L}^2 + \alpha \mathcal{L}^\infty)}{\alpha^2}, \quad \Lambda_{4,1}(g) = \frac{\sin y_c(\mathcal{L}^2 + \alpha^{\frac{1}{2}}\mathcal{L}^\infty)}{\alpha^2}, \\ \partial_c(\Lambda_{3,1}(\widehat{\omega}_e)) = \frac{\mathcal{L}^2 + \alpha \mathcal{L}^\infty}{\alpha^2 \sin y_c}, \quad \partial_c(\Lambda_{4,1}(g)) = \frac{\mathcal{L}^2 + \alpha^{\frac{1}{2}}\mathcal{L}^\infty}{\alpha^2 \sin y_c};$$

(3) if
$$\|\widehat{\omega}_e\|_{H^2} \leq 1$$
 and $\|g'' - \alpha^2 g\|_{L^2} \leq 1$ and $\widehat{\omega}'_e(0) = \widehat{\omega}'_e(\pi) = g'(0) = g'(\pi) = 0$, then
 $\Lambda_{3,1}(\widehat{\omega}_e) = \alpha^{-2}(\rho \mathcal{L}^2 \cap u' \mathcal{L}^\infty) + \rho \mathcal{L}^\infty$, $\Lambda_{4,1}(g) = \alpha^{-2}(\rho \mathcal{L}^2 \cap u' \mathcal{L}^\infty) + \alpha^{-\frac{3}{2}}\rho \mathcal{L}^\infty$,
 $\partial_c (\Lambda_{3,1}(\widehat{\omega}_e)) = \alpha^{-2}\mathcal{L}^2 + \mathcal{L}^\infty$, $\partial_c (\Lambda_{4,1}(g)) = \alpha^{-2}(\mathcal{L}^2 + \alpha^{\frac{1}{2}}\mathcal{L}^\infty)$,
 $\partial_c^2 (\Lambda_{3,1}(\widehat{\omega}_e)) = \frac{\alpha^{-2}\mathcal{L}^2 + \mathcal{L}^\infty}{\sin^2 y_c}$, $\partial_c^2 (\Lambda_{4,1}(g)) = \frac{\mathcal{L}^2 + \alpha^{\frac{1}{2}}\mathcal{L}^\infty}{\alpha^2 \sin^2 y_c}$;

(4) if $\|\widehat{\omega}_e\|_{H^2} \le 1$ and $\|\partial_y \widehat{\omega}_e / u'\|_{H^1} \le 1$, then

$$\partial_c (\Lambda_{3,1}(\widehat{\omega}_e)) = \mathcal{L}^{\infty}, \quad \partial_c^2 (\Lambda_{3,1}(\widehat{\omega}_e)) = \frac{\alpha^2 \mathcal{L}^{\infty}}{(1 + \alpha \sin y_c)^2} + \frac{\mathcal{L}^2}{\alpha \sin y_c}.$$

Remark 8.2. Following the proof of Case 2, we can show that if $|\alpha| ||\omega||_{L^2} + ||\partial_y \omega||_{L^2} \le 1$, then

$$\Lambda_{3,1}(\omega) = \frac{\sin y_c(\mathcal{L}^2 + \alpha^{\frac{1}{2}}\mathcal{L}^\infty)}{\alpha^2}, \quad \partial_c(\Lambda_{3,1}(\omega)) = \frac{\mathcal{L}^2 + \alpha^{\frac{1}{2}}\mathcal{L}^\infty}{\alpha^2 \sin y_c}.$$

This will be used in the proof of Proposition 10.3.

Proof. Case 1. $\|\widehat{\omega}_e\|_{L^2} \leq 1$ and $\|g\|_{L^2} \leq 1$.

Using the fact that $\phi_1(y,c) \le \phi_1(0,c)$ for $0 < y < y_c$ and $\phi_1(y,c) \le \phi_1(\pi,c)$ for $y_c < y < \pi$, we get

(8.1)
$$\left\| \frac{\mathcal{E}_j(\varphi)}{\phi_1(j\pi, c)(j\pi - y_c)} \right\|_{L^p} \le C \|\varphi\|_{L^p}, \quad 1$$

which implies that

$$u''(y_c) \mathcal{E}_0(\widehat{\omega}_e) + u'(y_c) \widehat{\omega}_e(y_c) = \phi_1(0, c) y_c \mathcal{L}^2,$$

$$u''(y_c) \mathcal{E}_1(\widehat{\omega}_e) + u'(y_c) \widehat{\omega}_e(y_c) = \phi_1(\pi, c) (\pi - y_c) \mathcal{L}^2,$$

and

$$E_0(gu'') + u'(y_c)g(y_c) = \phi_1(0,c)y_c\mathcal{L}^2, E_1(gu'') + u'(y_c)g(y_c) = \phi_1(\pi,c)(\pi - y_c)\mathcal{L}^2,$$

from which and Lemma 4.8, we deduce (1).

...

Case 2. $\|\widehat{\omega}_e\|_{H^1} \le 1$ and $|\alpha| \|g\|_{L^2} + \|g'\|_{L^2} \le 1$ (thus, $\|g\|_{L^{\infty}} \le C |\alpha|^{-\frac{1}{2}}$.) We write

$$u''(y_c) \mathbf{E}_j(\widehat{\omega}_e) + u'(y_c) \widehat{\omega}_e(y_c)$$

$$= \int_{y_c}^{j\pi} \left(u''(y_c) \phi_1(y, c) \widehat{\omega}_e(y) - u''(y) \widehat{\omega}_e(y_c) \right) dy$$

$$= \int_{y_c}^{j\pi} \left(u''(y_c) - u''(y) \right) \phi_1(y, c) \widehat{\omega}_e(y) dy$$

$$+ \int_{y_c}^{j\pi} \left(\phi_1(y, c) - 1 \right) u''(y) \widehat{\omega}_e(y) dy + \int_{y_c}^{j\pi} u''(y) \left(\widehat{\omega}_e(y) - \widehat{\omega}_e(y_c) \right) dy$$

$$(8.2) \qquad = I_1 + I_2 + I_3.$$

Using the fact that $\phi_1(y,c) - 1 \le C \min\{\alpha^2 | y - y_c|^2, 1\} \phi_1(y,c) \le C \min\{\alpha^2 | y - y_c|^2, |\alpha| | y - y_c|\} \phi_1(y,c)$ and $|u''(y) - u''(y_c)| \le \sin \frac{y + y_c}{2} | y - y_c|$, we obtain

(8.3)
$$|I_1| \le C |j\pi - y_c|^3 \phi_1(j\pi, c) \|\widehat{\omega}_e\|_{L^{\infty}},$$

 $|I_2| \le C |\alpha| |j\pi - y_c|^2 \min\{1, |\alpha| |j\pi - y_c|\} \phi_1(j\pi, c) \|\widehat{\omega}_e\|_{L^{\infty}},$ (8.4)

(8.5)
$$\left\|\frac{I_3}{|j\pi - y_c|^2}\right\|_{L^2} \le C \|\widehat{\omega}'_e\|_{L^2}.$$

Thus, we infer that

$$u''(y_c) \mathcal{E}_0(\widehat{\omega}_e) + u'(y_c) \widehat{\omega}_e(y_c) = \phi_1(0, c) y_c^2 \alpha \mathcal{L}^{\infty} + y_c^2 \mathcal{L}^2, u''(y_c) \mathcal{E}_1(\widehat{\omega}_e) + u'(y_c) \widehat{\omega}_e(y_c) = \phi_1(\pi, c) (\pi - y_c)^2 \alpha \mathcal{L}^{\infty} + (\pi - y_c)^2 \mathcal{L}^2,$$

which along with Lemma 4.8 show that

$$\Lambda_{3,1}(\widehat{\omega}_e) = \frac{\sin y_c(\mathcal{L}^2 + \alpha \mathcal{L}^\infty)}{\alpha^2}$$

We write

$$\begin{split} & \mathcal{E}_{j}(u''g) + u'(y_{c})g(y_{c}) \\ &= \int_{y_{c}}^{j\pi} \left(u''(y)\phi_{1}(y,c)g(y) - u''(y)g(y_{c}) \right) dy \\ &= \int_{y_{c}}^{j\pi} \left(\phi_{1}(y,c) - 1 \right) u''(y)g(y) dy + \int_{y_{c}}^{j\pi} u''(y) \left(g(y) - g(y_{c}) \right) dy \\ &= I_{2}' + I_{3}'. \end{split}$$

We have

(8.6)

(8.7)
$$|I'_{2}| \leq C |\alpha| |j\pi - y_{c}|^{2} \min \left\{ 1, |\alpha| |j\pi - y_{c}| \right\} \phi_{1}(j\pi, c) ||g||_{L^{\infty}}, \\ \left\| \frac{I'_{3}}{|j\pi - y_{c}|^{2}} \right\|_{L^{2}} \leq C ||g'||_{L^{2}}.$$

Thus, we obtain

$$E_0(gu'') + u'(y_c)g(y_c) = \phi_1(0,c)y_c^2 \alpha^{\frac{1}{2}} \mathcal{L}^{\infty} + y_c^2 \mathcal{L}^2,$$

$$E_1(gu'') + u'(y_c)g(y_c) = \phi_1(\pi,c)(\pi - y_c)^2 \alpha^{\frac{1}{2}} \mathcal{L}^{\infty} + (\pi - y_c)^2 \mathcal{L}^2,$$

which along with Lemma 4.8 show that

$$\Lambda_{4,1}(g) = \frac{\sin y_c(\mathcal{L}^2 + \alpha^{\frac{1}{2}}\mathcal{L}^\infty)}{\alpha^2}.$$

A direct calculation gives

(8.8)
$$\partial_c \left(u''(y_c) \mathcal{E}_j(\widehat{\omega}_e) + u'(y_c) \widehat{\omega}_e(y_c) \right) \\ = \frac{u'''(y_c)}{u'(y_c)} \mathcal{E}_j(\widehat{\omega}_e) - \frac{u''(y_c) \widehat{\omega}_e(y_c)}{u'(y_c)} + u''(y_c) \int_{y_c}^{j\pi} \widehat{\omega}_e \partial_c \phi_1(y,c) dy + \frac{(u'\widehat{\omega}_e)'(y_c)}{u'(y_c)} \\ = \frac{u'''(y_c)}{u'(y_c)} \mathcal{E}_j(\widehat{\omega}_e) + \widehat{\omega}'_e(y_c) + u''(y_c) \int_{y_c}^{j\pi} \widehat{\omega}_e \partial_c \phi_1(y,c) dy,$$

and

$$\partial_c \left(\mathcal{E}_j(gu'') + u'(y_c)g(y_c) \right) = -\frac{gu''(y_c)}{u'(y_c)} + \int_{y_c}^{j\pi} gu'' \partial_c \phi_1(y,c) dy + \frac{(u'g)'(y_c)}{u'(y_c)}$$
$$= g'(y_c) + \int_{y_c}^{j\pi} gu'' \partial_c \phi_1(y,c) dy.$$

By Proposition 3.8, we get for $0 < y < y_c$, j = 0 or $y_c < y < \pi$, j = 1,

$$\frac{\partial_c \phi_1(y,c)}{\phi_1(j\pi,c)} \le \frac{C|\alpha|}{u'(y_c)} \min\{|\alpha||y-y_c|,1\} \le \frac{C|\alpha|}{u'(y_c)} \min\{|\alpha||j\pi-y_c|,1\}.$$

Then we have for j = 0

(8.9)
$$\left\|\frac{(\pi - y_c)}{\min\{\alpha y_c, 1\}} \int_{y_c}^0 \varphi(y) \frac{\partial_c \phi_1(y, c)}{\phi_1(0, c)} dy\right\|_{L^p} \le C\alpha \|\varphi\|_{L^p}, \quad 1$$

Similarly, we get for j = 1

(8.10)
$$\left\|\frac{y_c}{\min\{\alpha(\pi - y_c), 1\}} \int_{y_c}^{\pi} \varphi(y) \frac{\partial_c \phi_1(y, c)}{\phi_1(\pi, c)} dy\right\|_{L^p} \le C\alpha \|\varphi\|_{L^p}, \quad 1$$

Recall that $\|\widehat{\omega}_e\|_{H^1} \leq 1$ and $\alpha \|g\|_{L^2} + \|g'\|_{L^2} \leq 1$, then $\|g\|_{L^{\infty}} \leq C\alpha^{-\frac{1}{2}}$. Thus, we obtain

$$\partial_c \left(u''(y_c) \mathcal{E}_0(\widehat{\omega}_e) + u'(y_c) \widehat{\omega}_e(y_c) \right) = \phi_1(0, c) \frac{\alpha \mathcal{L}^{\infty}}{\pi - y_c} + \mathcal{L}^2,$$

$$\partial_c \left(u''(y_c) \mathcal{E}_1(\widehat{\omega}_e) + u'(y_c) \widehat{\omega}_e(y_c) \right) = \phi_1(\pi, c) \frac{\alpha \mathcal{L}^{\infty}}{y_c} + \mathcal{L}^2,$$

$$\partial_c \left(\mathcal{E}_0(gu'') + u'(y_c)g(y_c) \right) = \frac{\phi_1(0, c)\alpha^{\frac{1}{2}}\mathcal{L}^{\infty}}{\pi - y_c} + \mathcal{L}^2,$$

$$\partial_c \left(\mathcal{E}_1(gu'') + u'(y_c)g(y_c) \right) = \frac{\phi_1(\pi, c)\alpha^{\frac{1}{2}}\mathcal{L}^{\infty}}{y_c} + \mathcal{L}^2,$$

from which and Lemma 4.8, we infer that

$$\partial_c \big(\Lambda_{3,1}(\widehat{\omega}_e) \big) = \frac{\mathcal{L}^2 + \alpha \mathcal{L}^\infty}{\alpha^2 \sin y_c}, \quad \partial_c \big(\Lambda_{4,1}(g) \big) = \frac{\mathcal{L}^2 + \alpha^{\frac{1}{2}} \mathcal{L}^\infty}{\alpha^2 \sin y_c}.$$

Case 3. $\|\widehat{\omega}_e\|_{H^2} \leq 1$ and $\|g'' - \alpha^2 g\|_{L^2} \leq 1$ and $\widehat{\omega}'_e(0) = \widehat{\omega}'_e(\pi) = g'(0) = g'(\pi) = 0$. Thus, we have ~ 1

$$\|g''\|_{L^{2}}^{2} + 2\alpha^{2} \|g'\|_{L^{2}}^{2} + \alpha^{4} \|g\|_{L^{2}}^{2} \leq 1, \quad \left\|\frac{\widehat{\omega}'_{e}}{u'}\right\|_{L^{2}} \leq C \|\widehat{\omega}''_{e}\|_{L^{2}},$$
$$\|g\|_{L^{\infty}} \leq C\alpha^{-\frac{3}{2}}, \quad \left\|\frac{g'}{u'}\right\|_{L^{2}} \leq C \|g''\|_{L^{2}} \leq C, \quad \|g'\|_{L^{\infty}} \leq C\alpha^{-\frac{1}{2}}$$

First of all, we have for I_3 in (8.2) and I'_3 in (8.6),

$$|I_3| \le C \Big| \int_{y_c}^{j\pi} \int_y^{y_c} \widehat{\omega}'_e(y') dy' dy \Big| \le C \Big| \int_{y_c}^{j\pi} |j\pi - y'| \widehat{\omega}'_e(y') dy' \Big| \le C |j\pi - y_c|^2 \Big| \int_{y_c}^{j\pi} \frac{\widehat{\omega}'_e(y')}{j\pi - y'} dy' \Big|,$$
which implies that

which implies that

$$\left\|\frac{I_3}{|j\pi - y_c|^3}\right\|_{L^2} \le C \left\|\frac{\widehat{\omega}'_e}{u'}\right\|_{L^2} \le C \|\widehat{\omega}''_e\|_{L^2}, \quad \left\|\frac{I_3}{|j\pi - y_c|^2}\right\|_{L^{\infty}} \le C \|\widehat{\omega}'_e\|_{L^{\infty}}.$$

Similarly, for I'_3 , we have

$$\left\|\frac{I_3'}{|j\pi - y_c|^3}\right\|_{L^2} \le C \left\|\frac{g'}{u'}\right\|_{L^2} \le C \|g''\|_{L^2}, \quad \left\|\frac{I_3'}{|j\pi - y_c|^2}\right\|_{L^\infty} \le C \|g'\|_{L^\infty}.$$

Thus, by (8.3), (8.4) and (8.7), we obtain

$$u''(y_c) \mathcal{E}_0(\widehat{\omega}_e) + u'(y_c) \widehat{\omega}_e(y_c) = \phi_1(0, c) y_c^3 \alpha^2 \mathcal{L}^{\infty} + (y_c^3 \mathcal{L}^2 \cap y_c^2 \mathcal{L}^{\infty}),$$

$$u''(y_c) \mathcal{E}_1(\widehat{\omega}_e) + u'(y_c) \widehat{\omega}_e(y_c) = \phi_1(\pi, c) (\pi - y_c)^3 \alpha^2 \mathcal{L}^{\infty} + (\pi - y_c)^2 ((\pi - y_c) \mathcal{L}^2 \cap \mathcal{L}^{\infty}),$$

$$\mathcal{E}_0(u''g) + u'(y_c) g(y_c) = \phi_1(0, c) y_c^3 \alpha^{\frac{1}{2}} \mathcal{L}^{\infty} + (y_c^3 \mathcal{L}^2 \cap y_c^2 \mathcal{L}^{\infty}),$$

$$\mathcal{E}_1(u''g) + u'(y_c) g(y_c) = \phi_1(\pi, c) (\pi - y_c)^3 \alpha^{\frac{1}{2}} \mathcal{L}^{\infty} + (\pi - y_c)^2 ((\pi - y_c) \mathcal{L}^2 \cap \mathcal{L}^{\infty}),$$

which along with Lemma 4.8 show that

$$\Lambda_{3,1}(\widehat{\omega}_e) = \alpha^{-2}(\rho \mathcal{L}^2 \cap u' \mathcal{L}^\infty) + \rho \mathcal{L}^\infty, \quad \Lambda_{4,1}(g) = \alpha^{-2}(\rho \mathcal{L}^2 \cap u' \mathcal{L}^\infty) + \alpha^{-\frac{3}{2}}\rho \mathcal{L}^\infty.$$

We have

$$\begin{aligned} \partial_c \left(u''(y_c) E_j(\widehat{\omega}_e) + u'(y_c) \widehat{\omega}_e(y_c) \right) \\ &= \int_{y_c}^{j\pi} u''(y_c) \partial_c \phi_1(y,c) \widehat{\omega}_e(y) dy + \widehat{\omega}'_e(y_c) - \int_{y_c}^{j\pi} \phi_1(y,c) \widehat{\omega}_e(y) dy \\ &= I_4 + I_5 + I_6. \end{aligned}$$

By Proposition 3.8, we have

$$|I_4| \leq \frac{C|\alpha||j\pi - y_c|\phi_1(j\pi, c)}{u'(y_c)} \min\{|\alpha||j\pi - y_c|, 1\} \|\widehat{\omega}_e\|_{L^{\infty}}, \\ \left\|\frac{I_5}{u'}\right\|_{L^2} \leq C \left\|\frac{\widehat{\omega}'_e}{u'}\right\|_{L^2}, \ |I_6| \leq C |\phi_1(j\pi, c)|j\pi - y_c| \|\widehat{\omega}_e\|_{L^{\infty}},$$

which implies that

$$\partial_c \left(u''(y_c) E_j(\widehat{\omega}_e) + u'(y_c) \widehat{\omega}_e(y_c) \right) = \frac{\phi_1(j\pi, c) \alpha^2 |j\pi - y_c| \mathcal{L}^{\infty}}{((1-j)\pi - y_c)} + u'(y_c) \mathcal{L}^2.$$

Similarly, due to

$$\partial_c \left(E_j(u''g) + u'(y_c)g(y_c) \right) = \int_{y_c}^{j\pi} u''(y)\partial_c \phi_1(y,c)g(y)dy + g'(y_c),$$

we get

$$\partial_c \left(\mathcal{E}_j(u''g) + u'(y_c)g(y_c) \right) = \frac{\phi_1(j\pi, c)\alpha^{\frac{1}{2}}|j\pi - y_c|\mathcal{L}^{\infty}}{((1-j)\pi - y_c)} + u'(y_c)\mathcal{L}^2,$$

from which and Lemma 4.8, we infer that

$$\partial_c \big(\Lambda_{3,1}(\widehat{\omega}_e) \big) = \alpha^{-2} \mathcal{L}^2 + \mathcal{L}^{\infty}, \quad \partial_c \big(\Lambda_{4,1}(g) \big) = \alpha^{-2} (\mathcal{L}^2 + \alpha^{\frac{1}{2}} \mathcal{L}^{\infty}).$$

We have

$$\partial_c^2 \left(\mathcal{E}_j(gu'') + u'(y_c)g(y_c) \right) = \frac{g''(y_c)}{u'(y_c)} + \int_{y_c}^{j\pi} gu'' \partial_c^2 \phi_1(y,c) dy,$$

and

$$(8.11) \qquad \partial_{c}^{2} \left(u''(y_{c}) \mathrm{E}_{j}(\widehat{\omega}_{e}) + u'(y_{c})\widehat{\omega}_{e}(y_{c}) \right) \\ = \left(\frac{u'''(y_{c})}{u'(y_{c})^{2}} - \frac{(u'''u'')(y_{c})}{u'(y_{c})^{3}} \right) \mathrm{E}_{j}(\widehat{\omega}_{e}) + \frac{\widehat{\omega}''_{e}(y_{c})}{u'(y_{c})} + 2\frac{u'''(y_{c})}{u'(y_{c})} \int_{y_{c}}^{j\pi} \widehat{\omega}_{e} \partial_{c} \phi_{1}(y, c) dy \\ + u''(y_{c}) \int_{y_{c}}^{j\pi} \widehat{\omega}_{e} \partial_{c}^{2} \phi_{1}(y, c) dy - \frac{u'''(y_{c})}{u'(y_{c})^{2}} \widehat{\omega}_{e}(y_{c}) \\ = \frac{\widehat{\omega}''_{e}(y_{c})}{u'(y_{c})} - 2 \int_{y_{c}}^{j\pi} \widehat{\omega}_{e} \partial_{c} \phi_{1}(y, c) dy + u''(y_{c}) \int_{y_{c}}^{j\pi} \widehat{\omega}_{e} \partial_{c}^{2} \phi_{1}(y, c) dy - \frac{u'''(y_{c})}{u'(y_{c})^{2}} \widehat{\omega}_{e}(y_{c}) \\ = \frac{\widehat{\omega}''_{e}(y_{c})}{u'(y_{c})} - 2 \int_{y_{c}}^{j\pi} \widehat{\omega}_{e} \partial_{c} \phi_{1}(y, c) dy + u''(y_{c}) \int_{y_{c}}^{j\pi} \widehat{\omega}_{e} \partial_{c}^{2} \phi_{1}(y, c) dy - \frac{u'''(y_{c})}{u'(y_{c})^{2}} \widehat{\omega}_{e}(y_{c}) \\ = \frac{\widehat{\omega}''_{e}(y_{c})}{u'(y_{c})} - 2 \int_{y_{c}}^{j\pi} \widehat{\omega}_{e} \partial_{c} \phi_{1}(y, c) dy + u''(y_{c}) \int_{y_{c}}^{j\pi} \widehat{\omega}_{e} \partial_{c}^{2} \phi_{1}(y, c) dy - \frac{u'''(y_{c})}{u'(y_{c})^{2}} \widehat{\omega}_{e}(y_{c}) \\ = \frac{\widehat{\omega}''_{e}(y_{c})}{u'(y_{c})} - 2 \int_{y_{c}}^{j\pi} \widehat{\omega}_{e} \partial_{c} \phi_{1}(y, c) dy + u''(y_{c}) \int_{y_{c}}^{j\pi} \widehat{\omega}_{e} \partial_{c}^{2} \phi_{1}(y, c) dy - \frac{u'''(y_{c})}{u'(y_{c})^{2}} \widehat{\omega}_{e}(y_{c}) \\ = \frac{\widehat{\omega}''_{e}(y_{c})}{u'(y_{c})} - 2 \int_{y_{c}}^{j\pi} \widehat{\omega}_{e} \partial_{c} \phi_{1}(y, c) dy + u''(y_{c}) \int_{y_{c}}^{j\pi} \widehat{\omega}_{e} \partial_{c}^{2} \phi_{1}(y, c) dy - \frac{u''(y_{c})}{u'(y_{c})^{2}} \widehat{\omega}_{e}(y_{c}) \\ = \frac{\widehat{\omega}''_{e}(y_{c})}{u'(y_{c})} - 2 \int_{y_{c}}^{j\pi} \widehat{\omega}_{e} \partial_{c} \phi_{1}(y, c) dy + u''(y_{c}) \int_{y_{c}}^{j\pi} \widehat{\omega}_{e} \partial_{c}^{2} \phi_{1}(y, c) dy \\ = \frac{\widehat{\omega}_{e}^{j}(y_{c})}{u'(y_{c})} - 2 \int_{y_{c}}^{j\pi} \widehat{\omega}_{e} \partial_{c} \phi_{1}(y, c) dy + u''(y_{c}) \int_{y_{c}}^{j\pi} \widehat{\omega}_{e} \partial_{c}^{j\pi} \phi_{1}(y, c) dy \\ = \frac{\widehat{\omega}_{e}^{j}(y_{c})}{u'(y_{c})} - 2 \int_{y_{c}}^{j\pi} \widehat{\omega}_{e} \partial_{c} \phi_{1}(y, c) dy \\ = \frac{\widehat{\omega}_{e}^{j}(y_{c})}{u'(y_{c})} - 2 \int_{y_{c}}^{j\pi} \widehat{\omega}_{e} \partial_{c} \phi_{1}(y, c) dy \\ = \frac{\widehat{\omega}_{e}^{j}(y_{c})}{u'(y_{c})} - 2 \int_{y_{c}}^{j\pi} \widehat{\omega}_{e} \partial_{c} \phi_{1}(y, c) dy \\ = \frac{\widehat{\omega}_{e}^{j}(y_{c})}{u'(y_{c})} - 2 \int_{y_{c}}^{j\pi} \widehat{\omega}_{e} \partial_{c} \phi_{1}(y, c) dy \\ = \frac{\widehat{\omega}_{e}^{j}(y_{c})}{$$

By Proposition 3.8, we deduce that for $0 < y < y_c$, j = 0 or $y_c < y < \pi$, j = 1, $\left|\frac{\partial_c^2 \phi_1(y,c)}{\phi_1(j\pi,c)}\right| \leq \frac{C\alpha^2}{u'(y_c)^2}$, and then

(8.12)
$$\left\| y_c (\pi - y_c)^2 \int_{y_c}^0 \varphi(y) \frac{\partial_c^2 \phi_1(y,c)}{\phi_1(0,c)} dy \right\|_{L^p} \le C \alpha^2 \|\varphi\|_{L^p}, \quad 1$$

(8.13)
$$\left\| y_c^2(\pi - y_c) \int_{y_c}^{\pi} \varphi(y) \frac{\partial_c^2 \phi_1(y,c)}{\phi_1(\pi,c)} dy \right\|_{L^p} \le C\alpha^2 \|\varphi\|_{L^p}, \quad 1$$

Then we can deduce that

$$y_c \partial_c^2 \left(u''(y_c) \mathcal{E}_0(\widehat{\omega}_e) + u'(y_c) \widehat{\omega}_e(y_c) \right) = \frac{\phi_1(0,c) \alpha^2 \mathcal{L}^\infty}{(\pi - y_c)^2} + \mathcal{L}^2,$$
$$(\pi - y_c) \partial_c^2 \left(u''(y_c) \mathcal{E}_1(\widehat{\omega}_e) + u'(y_c) \widehat{\omega}_e(y_c) \right) = \frac{\phi_1(\pi,c) \alpha^2 \mathcal{L}^\infty}{y_c^2} + \mathcal{L}^2,$$

from which and Lemma 4.8, we infer that

(8.14)
$$\partial_c^2 \left(\frac{J_{1-j}}{u'(y_c)} \left(u''(y_c) \mathcal{E}_j(\widehat{\omega}_e) + u'(y_c) \widehat{\omega}_e(y_c) \right) \right) = \frac{\mathcal{L}^{\infty}}{(j\pi - y_c)^2} + \frac{\mathcal{L}^2}{\alpha^2 (j\pi - y_c)^2},$$

which implies that

$$\partial_c^2 (\Lambda_{3,1}(\widehat{\omega}_e)) = \frac{\alpha^{-2} \mathcal{L}^2 + \mathcal{L}^\infty}{\sin^2 y_c}.$$

Similarly, we have

$$y_c \partial_c^2 \left(\mathcal{E}_0(u''g) + u'(y_c)g(y_c) \right) = \frac{\phi_1(0,c)\alpha^{\frac{1}{2}}\mathcal{L}^{\infty}}{(\pi - y_c)^2} + \mathcal{L}^2,$$

$$(\pi - y_c)\partial_c^2 \left(\mathbf{E}_1(u''g) + u'(y_c)g(y_c) \right) = \frac{\phi_1(\pi, c)\alpha^{\frac{1}{2}}\mathcal{L}^{\infty}}{y_c^2} + \mathcal{L}^2,$$

from which and Lemma 4.8, we infer that

$$\partial_c^2 \big(\Lambda_{4,1}(g) \big) = \frac{\mathcal{L}^2 + \alpha^{\frac{1}{2}} \mathcal{L}^\infty}{\alpha^2 \sin^2 y_c}.$$

Case 4. $\|\widehat{\omega}_e\|_{H^2} \leq 1$ and $\|\partial_y \widehat{\omega}_e / u'\|_{H^1} \leq 1$. Recall that

$$\partial_c \left(u''(y_c) E_j(\widehat{\omega}_e) + u'(y_c) \widehat{\omega}_e(y_c) \right) \\= \int_{y_c}^{j\pi} u''(y_c) \partial_c \phi_1(y,c) \widehat{\omega}_e(y) dy + \widehat{\omega}'_e(y_c) - \int_{y_c}^{j\pi} \phi_1(y,c) \widehat{\omega}_e(y) dy$$

Now since $\partial_y \widehat{\omega}_e / u' \in H^1 \subset L^\infty$, we get

$$\partial_c \left(u''(y_c) E_j(\widehat{\omega}_e) + u'(y_c) \widehat{\omega}_e(y_c) \right) = \frac{\phi_1(j\pi, c) \alpha^2 |j\pi - y_c| \mathcal{L}^{\infty}}{((1-j)\pi - y_c)} + u'(y_c) \mathcal{L}^{\infty},$$

from which and Lemma 4.8, we infer that $\partial_c \Lambda_{3,1} = \mathcal{L}^{\infty}$.

For the estimates of higher derivative, we only need to improve the estimate of $\frac{J_{1-j}}{u'(y_c)} (u''(y_c) \mathbf{E}_j(\widehat{\omega}_e) + u'(y_c)\widehat{\omega}_e(y_c))$ for $|y_c - j\pi| \leq \frac{1}{\alpha}$. We show the proof of the case j = 0, the other case can be proved similarly.

Let $\omega_1 = \frac{\partial_y \hat{\omega}_e}{y} \in H^1(0, 1)$, and then

$$E_0(\widehat{\omega}_e) = \int_{y_c}^0 \widehat{\omega}_e(y)\phi_1(y,c)dy = y_c \int_1^0 \widehat{\omega}_e(yy_c)\widetilde{\phi_1}(y,c)dy,$$

here we recall the definition of $\phi_1(y,c) = \phi_1(yy_c,c)$ and its estimate in the proof of Lemma 4.8.

Using the fact that

$$\partial_c \widehat{\omega}_e(yy_c) = \frac{y\widehat{\omega}'_e(yy_c)}{u'(y_c)} = \frac{y^2\omega_1(yy_c)}{m_1(y_c)},$$

with $m_1(y) = \frac{\sin y}{y}$ and

$$\partial_c^2 \widehat{\omega}_e(yy_c) = \frac{y^3 \omega_1'(yy_c)}{m_1(y_c)u'(y_c)} + \frac{y^2 \omega_1(yy_c)m_1'(y_c)}{m_1(y_c)^2 u'(y_c)},$$

and that $1 \leq |\widetilde{\phi_1}(y,c)| \leq 2$ and (3.38), (3.39) hold for $y \in [0,1]$ and $c \in [u(0), u(1/\alpha)]$, we deduce that for $0 < y_c < \frac{1}{\alpha}$,

$$\begin{aligned} |\partial_c(E_0(\widehat{\omega}_e)/y_c)| &\leq C(\|\omega_1\|_{L^{\infty}} + \alpha^2 \|\widehat{\omega}_e\|_{L^{\infty}}), \\ |\partial_c^2(E_0(\widehat{\omega}_e)/y_c)| &\leq C(\alpha^2 \|\omega_1\|_{L^{\infty}} + \alpha^4 \|\widehat{\omega}_e\|_{L^{\infty}}) + \frac{C}{y_c} \int_0^1 y^3 |\omega_1'(yy_c)| dy. \end{aligned}$$

Now using the fact that

$$J_1\left(\frac{u''(y_c)}{u'(y_c)}\mathcal{E}_0(\widehat{\omega}_e) + \widehat{\omega}_e(y_c)\right) = \frac{-(u(\pi) - c)^2 y_c^2 u''(y_c)}{\widetilde{\phi_1}(0, c)\widetilde{\phi_1}'(0, c)} \frac{\mathcal{E}_0(\widehat{\omega}_e)}{y_c} + J_1\widehat{\omega}_e(y_c),$$

and by Proposition 3.10 and Lemma 4.8, we deduce that for $0 < y_c < \frac{1}{\alpha}$,

$$\partial_c^2 \left(J_1 \left(\frac{u''(y_c)}{u'(y_c)} \mathcal{E}_0(\widehat{\omega}_e) + \widehat{\omega}_e(y_c) \right) \right) \chi_{[u(0), u(1/\alpha)]} = \alpha^2 \mathcal{L}^\infty + \frac{\mathcal{L}^2}{\alpha^2 y_c},$$

which along with (8.14) gives

$$\partial_c^2 \left(J_1 \left(\frac{u''(y_c)}{u'(y_c)} \mathcal{E}_0(\widehat{\omega}_e) + \widehat{\omega}_e(y_c) \right) \right) = \frac{\alpha^2 \mathcal{L}^\infty}{(1 + \alpha \sin y_c)^2} + \frac{\mathcal{L}^2}{\alpha y_c}$$

Similarly, for j = 1, we have

$$E_1(\widehat{\omega}_e) = \int_{y_c}^{\pi} \widehat{\omega}_e(y)\phi_1(y,c)dy = (y_c - \pi) \int_1^0 \widehat{\omega}_e((1-z)\pi + zy_c)\widetilde{\phi_1}(z,c)dz.$$

Then by the same argument, we obtain

$$\partial_c^2 \left(J_0 \left(\frac{u''(y_c)}{u'(y_c)} \mathcal{E}_1(\widehat{\omega}_e) + \widehat{\omega}_e(y_c) \right) \right) = \frac{\alpha^2 \mathcal{L}^\infty}{(1 + \alpha \sin y_c)^2} + \frac{\mathcal{L}^2}{\alpha (\pi - y_c)}$$

from which, we deduce (4) of the lemma.

Now we are in a position to prove Proposition 6.2.

Proof. Step1. L^1 estimate.

We normalize $\widehat{\omega}_e, g$ so that $\|\widehat{\omega}_e\|_{L^2} \leq 1$ and $\|g\|_{L^2} \leq 1$. By Lemma 7.4, we get $\Lambda_{1,1}(\widehat{\omega}_e) = \mathcal{L}^2$ and $\Lambda_{2,1}(g) = \mathcal{L}^2$. By Lemma 7.6 and Lemma 4.4, we get $\Lambda_{1,2}(\widehat{\omega}_e) = |\alpha|^{\frac{1}{2}}\mathcal{L}^{\infty} + |\alpha| \sin y_c \mathcal{L}^2$ and $\Lambda_{2,2}(g) = |\alpha|^{\frac{1}{2}}\mathcal{L}^{\infty} + |\alpha| \sin y_c \mathcal{L}^2$. Then by Lemma 8.1, we get

$$\Lambda_3(\widehat{\omega}_e) = \rho(1+|\alpha|\sin y_c)\mathcal{L}^2 + |\alpha|^{\frac{1}{2}}\sin^2 y_c\mathcal{L}^\infty + \frac{\mathcal{L}^2}{\alpha^2}$$
$$\Lambda_4(g) = \rho(1+|\alpha|\sin y_c)\mathcal{L}^2 + |\alpha|^{\frac{1}{2}}\sin^2 y_c\mathcal{L}^\infty + \frac{\mathcal{L}^2}{\alpha^2},$$

from which and Proposition 4.5, we deduce that

$$||K_e(\cdot, \alpha)||_{L^1_c(-1,1)} \le C ||\widehat{\omega}_e||_{L^2} ||g||_{L^2}.$$

Step 2. $W^{1,1}$ estimate.

We normalize $\widehat{\omega}_e, g$ so that $\|\widehat{\omega}_e\|_{H^1} \leq 1$ and $\|\alpha\|\|g\|_{L^2} + \|g'\|_{L^2} \leq 1$, then $\|g\|_{L^{\infty}} \leq C|\alpha|^{-\frac{1}{2}}$. By Lemma 7.4, we get $\Lambda_{1,1}(\widehat{\omega}_e) = \mathcal{L}^{\infty}$ and $\Lambda_{2,1}(g) = |\alpha|^{-\frac{1}{2}}\mathcal{L}^{\infty}$. By Lemma 7.6 and Lemma 4.4, we get $\Lambda_{1,2}(\widehat{\omega}_e) = |\alpha| \sin y_c \mathcal{L}^{\infty}$ and $\Lambda_{2,2}(g) = |\alpha|^{\frac{1}{2}} \sin y_c \mathcal{L}^{\infty}$. Then by Lemma 8.1, we get

$$\begin{split} \Lambda_3(\widehat{\omega}_e) &= \sin^2 y_c (1+|\alpha|\sin y_c) \mathcal{L}^\infty + \frac{C\sin y_c (\mathcal{L}^2 + \alpha \mathcal{L}^\infty)}{\alpha^2} \\ &= \frac{\sin y_c}{\alpha} (1+|\alpha|\sin y_c)^2 \mathcal{L}^\infty + \frac{\sin y_c}{\alpha^2} \mathcal{L}^2 = |\alpha|^{-\frac{3}{2}} \sin y_c (1+|\alpha|\sin y_c)^3 \mathcal{L}^2 \\ \Lambda_4(g) &= |\alpha|^{-\frac{1}{2}} \sin^2 y_c (1+|\alpha|\sin y_c) \mathcal{L}^\infty + \frac{\sin y_c (\mathcal{L}^2 + |\alpha|^{\frac{1}{2}} \mathcal{L}^\infty)}{\alpha^2} \\ &= \frac{\sin y_c}{|\alpha|^{-\frac{3}{2}}} (1+|\alpha|\sin y_c)^2 \mathcal{L}^\infty + \frac{\sin y_c}{\alpha^2} \mathcal{L}^2 = \frac{\sin y_c}{\alpha^2} (1+|\alpha|\sin y_c)^3 \mathcal{L}^2. \end{split}$$

By Lemma 7.4, we get $\partial_c \Lambda_{1,1}(\widehat{\omega}_e) = \frac{\mathcal{L}^2}{\sin y_c}$ and $\partial_c \Lambda_{2,1}(g) = \frac{\mathcal{L}^2}{\sin y_c}$. By Lemma 7.7 and Lemma 4.4, we get $\partial_c \Lambda_{1,2}(\widehat{\omega}_e) = \frac{\alpha \mathcal{L}^{\infty}}{\sin y_c} + \alpha \mathcal{L}^2$ and $\partial_c \Lambda_{2,2}(g) = \frac{\alpha^{\frac{1}{2}} \mathcal{L}^{\infty}}{\sin y_c} + \alpha \mathcal{L}^2$. Then by Lemma 8.1, we obtain

$$\partial_c \left(\Lambda_3(\widehat{\omega}_e) \right) = (1 + \alpha \sin y_c) (\sin y_c \mathcal{L}^2 + \mathcal{L}^\infty) + \frac{(\mathcal{L}^2 + \alpha \mathcal{L}^\infty)}{\alpha^2 \sin y_c} \\ = \frac{(1 + |\alpha| \sin y_c)^2}{\alpha \sin y_c} \mathcal{L}^\infty + \frac{(1 + |\alpha| \sin y_c)^3}{\alpha^2 \sin y_c} \mathcal{L}^2 = \frac{(1 + |\alpha| \sin y_c)^3}{|\alpha|^{\frac{3}{2}} \sin y_c} \mathcal{L}^2, \\ \partial_c \left(\Lambda_4(g) \right) = \alpha^{\frac{1}{2}} \sin y_c \mathcal{L}^\infty + \sin y_c (1 + \alpha \sin y_c) \mathcal{L}^2 + \frac{(\mathcal{L}^2 + \alpha^{\frac{1}{2}} \mathcal{L}^\infty)}{\alpha^2 \sin y_c} \\ = \frac{(1 + |\alpha| \sin y_c)^2}{|\alpha|^{\frac{3}{2}} \sin y_c} \mathcal{L}^\infty + \frac{(1 + |\alpha| \sin y_c)^3}{\alpha^2 \sin y_c} \mathcal{L}^2 = \frac{(1 + |\alpha| \sin y_c)^3}{|\alpha|^2 \sin y_c} \mathcal{L}^2,$$

from which and Proposition 4.5, we infer that

$$\|\partial_c K_e(\cdot, \alpha)\|_{L^1_c(-1,1)} \le C\alpha^{\frac{1}{2}} \|\widehat{\omega}_e\|_{H^1} \|g\|_{H^1}.$$

Step 3. $W^{2,1}$ estimate.

We normalize $\widehat{\omega}_e, g$ so that $\|\widehat{\omega}_e\|_{H^2} \leq 1$ with $\widehat{\omega}'_e(0) = \widehat{\omega}'_e(1) = 0$ and $\|g'' - \alpha^2 g\|_{L^2} \leq 1$ with g'(0) = g'(1) = 0. Then we have

$$\|g''\|_{L^{2}}^{2} + \alpha^{2} \|g'\|_{L^{2}}^{2} + \alpha^{4} \|g\|_{L^{2}} \leq 1, \quad \|\frac{\widehat{\omega}'_{e}}{u'}\|_{L^{2}} \leq C \|\widehat{\omega}''_{e}\|_{L^{2}} \leq C, \\ \|g\|_{L^{\infty}} \leq C\alpha^{-\frac{3}{2}}, \quad \|\frac{g'}{u'}\|_{L^{2}} \leq C \|g''\|_{L^{2}} \leq C, \quad \|g'\|_{L^{\infty}} \leq C\alpha^{-\frac{1}{2}}, \quad \|u''g\|_{H^{k}} \leq C \|g\|_{H^{k}}$$

for k = 0, 1, 2.

By Lemma 7.4, we get $\Lambda_{1,1}(\widehat{\omega}_e) = \mathcal{L}^{\infty}$ and $\Lambda_{2,1}(g) = |\alpha|^{-\frac{3}{2}}\mathcal{L}^{\infty}$. By Lemma 7.6 and Lemma 4.4, we get $\Lambda_{1,2}(\widehat{\omega}_e) = |\alpha| \sin y_c \mathcal{L}^{\infty}$ and $\Lambda_{2,2}(g) = |\alpha|^{-\frac{1}{2}} \sin y_c \mathcal{L}^{\infty}$. Then by Lemma 8.1, we get

$$\Lambda_3(\widehat{\omega}_e) = \alpha^{-2} (\sin^2 y_c \mathcal{L}^2 \cap \sin y_c \mathcal{L}^\infty) + \sin^2 y_c (1 + \alpha \sin y_c) \mathcal{L}^\infty = \alpha^{-\frac{1}{2}} \sin^2 y_c (1 + \alpha \sin y_c)^2 \mathcal{L}^2,$$

$$\Lambda_4(g) = \alpha^{-2} (\rho \mathcal{L}^2 \cap u' \mathcal{L}^\infty) + \alpha^{-\frac{3}{2}} \rho (1 + \alpha \sin y_c) \mathcal{L}^\infty = \alpha^{-2} \sin^2 y_c (1 + \alpha \sin y_c)^2 \mathcal{L}^2.$$

By Lemma 7.4, we get $\partial_c \Lambda_{1,1}(\widehat{\omega}_e) = \mathcal{L}^2$ and $\partial_c \Lambda_{2,1}(g) = \mathcal{L}^2$. By Lemma 7.7 and Lemma 4.4, we get $\partial_c \Lambda_{1,2}(\widehat{\omega}_e) = \frac{\alpha \mathcal{L}^{\infty}}{\sin y_c}$ and $\partial_c \Lambda_{2,2}(g) = \frac{\alpha^{-\frac{1}{2}}\mathcal{L}^{\infty} + \alpha^{\frac{1}{2}} \sin y_c \mathcal{L}^{\infty}}{\sin y_c}$. Then by Lemma 8.1, we obtain

$$\partial_c \left(\Lambda_3(\widehat{\omega}_e) \right) = \sin^2 y_c \mathcal{L}^2 + \alpha^{-2} \mathcal{L}^2 + (1 + \alpha \sin y_c) \mathcal{L}^\infty = \alpha^{-\frac{1}{2}} (1 + \alpha \sin y_c)^2 \mathcal{L}^2,$$

$$\partial_c \left(\Lambda_4(g) \right) = \alpha^{-2} (\mathcal{L}^2 + \alpha^{\frac{1}{2}} (1 + \alpha \sin y_c) \mathcal{L}^\infty) + \sin^2 y_c (\mathcal{L}^2 + \alpha^{\frac{1}{2}} \mathcal{L}^\infty) = \alpha^{-2} (1 + \alpha \sin y_c)^3 \mathcal{L}^2$$

By Lemma 7.4, we get $\partial_c^2 \Lambda_{1,1}(\widehat{\omega}_e) = \frac{\mathcal{L}^2}{\sin^2 y_c}$ and $\partial_c^2 \Lambda_{2,1}(g) = \frac{\mathcal{L}^2}{\sin^2 y_c}$. By Lemma 7.8 and Lemma 4.4, we get $\partial_c^2 \Lambda_{1,2}(\widehat{\omega}_e) = \frac{\alpha \mathcal{L}^\infty}{\sin^3 y_c} + \frac{\alpha \mathcal{L}^2}{\sin y_c}$ and $\partial_c^2 \Lambda_{2,2}(g) = \frac{(\alpha^{\frac{1}{2}} \sin y_c + \alpha^{-\frac{1}{2}})}{\sin^3 y_c} \mathcal{L}^\infty + \frac{\alpha \mathcal{L}^2}{\sin y_c}$. Then by Lemma 8.1, we obtain

$$\partial_c^2 \left(\Lambda_3(\widehat{\omega}_e) \right) = (1 + \alpha \sin y_c) (\mathcal{L}^2 + \rho^{-1} \mathcal{L}^\infty) + \rho^{-1} (\alpha^{-2} \mathcal{L}^2 + \mathcal{L}^\infty) = \alpha^{-\frac{1}{2}} \rho^{-1} (1 + \alpha \sin y_c)^3 \mathcal{L}^2,$$

$$\partial_c^2 \left(\Lambda_4(g) \right) = (\alpha^{-2} \rho^{-1} + 1) (\mathcal{L}^2 + \alpha^{\frac{1}{2}} \mathcal{L}^\infty) + (1 + \alpha \sin y_c) \mathcal{L}^2 = \alpha^{-2} \rho^{-1} (1 + \alpha \sin y_c)^3 \mathcal{L}^2,$$

from which and Proposition 4.5, we infer that

$$\|\partial_c^2 K_e(\cdot, \alpha)\|_{L^1_c(-1,1)} \le C\alpha^{\frac{3}{2}} \|\widehat{\omega}_e\|_{H^2} \|f\|_{L^2}$$

and

$$K_e(c,\alpha) = C(\alpha)\sin^2 y_c \mathcal{L}^2,$$

which implies that $K_e(\pm 1, \alpha) = 0$.

9. Decay estimates at critical points

9.1. Vorticity depletion phenomena. In this subsection, we give a formula of the vorticity at a critical point and confirm the vorticity depletion phenomena found by Bouchet and Morita [10].

Lemma 9.1. Let $\omega(t, x, y)$ be a solution to (1.2). Then we have

$$\widehat{w}(t,\alpha,0) = \int_{-1}^{1} \frac{(1+\cos y_c)^2 \Lambda_3(\widehat{\omega}_e) e^{-i\alpha ct}}{u'(y_c) \phi_1'(0,c) (\mathbf{A}_1^2 + \mathbf{B}_1^2)} dc.$$

and $\frac{(1+\cos y_c)^2 \Lambda_3(\widehat{\omega}_e)}{u'(y_c)\phi'_1(0,c)(\mathbf{A}_1^2+\mathbf{B}_1^2)} \in L^1_c(-1,1)$. In particular, for any $\alpha \neq 0$,

$$\lim_{t \to \infty} \widehat{w}(t, \alpha, 0) = 0.$$

Proof. Let $\eta \in C^{\infty}(\mathbf{R})$ so that $\eta(y) = 1$ for $|y| < \frac{1}{2}$ and $\eta(y) = 0$ for |y| > 1 and $\eta'(y) \le 0$ for y > 0. We denote $\eta_n(y) = \eta(ny)$, $g_n(y) = \eta'_n(y) = n\eta'(ny)$ and $f_n(y) = g''_n(y) - \alpha^2 g_n(y)$. Assume that $\omega_0(x, y) \in H^2$, then for any $t \ge 0$, $\omega(t, x, y) \in H^2$. Then $\widehat{\omega}_o(t, \alpha, 0) = 0$, and

by Lemma 5.10,

$$\widehat{w}(t,\alpha,0) = -\lim_{n \to +\infty} \int_0^{\pi} \widehat{\omega}_e(t,\alpha,y) g_n(y) dy$$
$$= \lim_{n \to +\infty} \int_0^{\pi} \widehat{\psi}_e(t,\alpha,y) f_n(y) dy$$
$$= -\lim_{n \to +\infty} \int_{u(0)}^{u(\pi)} \frac{\Lambda_3(\widehat{\omega}_e) \Lambda_4(g_n)}{u'(y_c)(\Lambda_1^2 + B_1^2)} e^{-i\alpha ct} dc.$$

Pointwise limit of $\Lambda_4(g_n)$.

Since $\lim_{n \to +\infty} \eta_n(y_c) = 0$ and $\lim_{n \to +\infty} g_n(y_c) = 0$ for $y_c \in (0, \pi]$, we get

$$\lim_{n \to +\infty} \mathcal{E}_0(g_n u'')(c) = \lim_{n \to +\infty} -\int_0^{y_c} u''(y)\phi_1(y,c)d\eta_n(y) = u''(0)\phi_1(0,c),$$
$$\lim_{n \to +\infty} \mathcal{E}_1(g_n u'')(c) = \lim_{n \to +\infty} \int_{y_c}^{\pi} g_n(y)u''(y)\phi_1(y,c)dy = 0.$$

By the proof of Lemma 5.5, we get $II_{1,1}(\varphi) = p.v. \int_0^{\pi} \frac{\int_{y_c}^{y} \varphi(y') dy'}{(u(y)-c)^2} dy$. Then for $1/n < y_c$, as $n \to \infty$,

$$|\mathrm{II}_{1,1}(g_n u'')(c)| \leq \int_0^{1/n} \frac{\int_y^{y_c} |g_n u''|(z)dz}{(u(y) - c)^2} dy$$
$$\leq ||g_n u''||_{L^1} \int_0^{1/n} \frac{dy}{(u(y) - c)^2} \to 0.$$

62

By (7.3), we also have for $1/n < y_c$, as $n \to \infty$,

$$(9.1) |\mathcal{L}_{0}(g_{n}u'')| \leq \left| \int_{0}^{\frac{1}{n}} \int_{y_{c}}^{z} g_{n}u''(y) \left(\frac{1}{(u(z)-c)^{2}} \left(\frac{\phi_{1}(y,c)}{\phi_{1}(z,c)^{2}} - 1 \right) \right) dydz \right| \\ \leq C ||g_{n}u''||_{L^{1}} \int_{0}^{\frac{1}{n}} \sup_{y \in [z,1/n]} \left| \frac{1}{(u(z)-c)^{2}} \left(\frac{\phi_{1}(y,c)}{\phi_{1}(z,c)^{2}} - 1 \right) \right| dz \\ \leq \frac{C|\alpha|}{ny_{c}^{3}} ||g_{n}u''||_{L^{1}} \to 0$$

Thus, for $y_c \in (0, \pi]$, we obtain

$$\lim_{n \to +\infty} \Lambda_4(g_n)(c) = \frac{J_1(c)u''(0)\phi_1(0,c)}{u'(y_c)}$$
$$= -\frac{u'(y_c)\rho(c)^2 u''(0)\phi_1(0,c)}{\phi(0,c)\phi'(0,c)u'(y_c)} = -\frac{(u(\pi)-c)^2}{\phi_1'(0,c)}.$$

Uniform bound of $\Lambda_4(g_n)$.

For $c \in (-1, 1]$, by Lemma 4.8 we have for j = 0, 1,

$$\|\mathbf{E}_{j}(g_{n}u'')(c)\| \le \phi_{1}(j\pi, c) \|g_{n}u''\|_{L^{1}} \le C\phi_{1}(j\pi, c), \quad |J_{j}| \le \frac{C}{\alpha^{2}\phi_{1}((1-j)\pi, c)}.$$

By Lemma 4.4, we get $|\rho II(c)| \leq C \alpha \sin y_c$. By Lemma 7.6, we have

$$|\sin y_c^2 \mathcal{L}_0(g_n u'')| \le C |\alpha| ||g_n u''||_{L^1}.$$

Thus, $|\Lambda_{2,2}(g_n u'')| \leq C|\alpha|$. By (7.1), we get

(9.2)
$$\sin^3 y_c \operatorname{II}_{1,1}(\varphi)(y_c) = -\frac{\sin y_c}{2} \mathcal{H}(\varphi) - \frac{\sin y_c}{2} \int_{-\pi}^{\pi} \operatorname{In}(\varphi)(y') dy'.$$

As $Z \circ In(u''g_n)$ is odd, we get by Lemma 7.1 that

$$in y_c H(Z \circ In(u''g_n)) = H(sin yZ \circ In(u''g_n)).$$

Using the fact that for $y \in [0, \pi]$,

$$\sin yZ \circ \ln(u''g_n)(y) = \sin yu''g_n(y) - \sin y \int_0^\pi u''g_n(y)dy - \cos y \ln(u''g_n)(y),$$

we deduce that

$$\sin y_c \mathcal{H}(u''g_n) = H(\sin yu''g_n(y)) - \int_0^\pi u''g_n(y)dyH(\sin y) - H(\cos y \ln(u''g_n)(y)).$$

Thanks to $\|\sin y u'' g_n\|_{H^1} \le C\sqrt{n}$ and $\|\sin y u'' g_n\|_{L^2} \le \frac{C}{\sqrt{n}}$, we get

$$\|H(\sin yu''g_n(y))\|_{L^{\infty}} \le C \|H(\sin yu''g_n)\|_{H^1}^{\frac{1}{2}} \|H(\sin yu''g_n)\|_{L^2}^{\frac{1}{2}} \le C.$$

Thanks to $\|\cos y \operatorname{In}(u''g_n)(y)\|_{H^1} \le C\sqrt{n}$ and $\|\cos y \operatorname{In}(u''g_n)(y)\|_{L^2} \le \frac{C}{\sqrt{n}}$, we get

 $\|H(\cos y \ln(u''g_n))\|_{L^{\infty}} \leq C \|H(\cos y \ln(u''g_n))\|_{H^1}^{\frac{1}{2}} \|H(\cos y \ln(u''g_n))\|_{L^2}^{\frac{1}{2}} \leq C.$ With the estimates above, we conclude $\|u'(y_c)\Lambda_4(g_n)\|_{L^{\infty}} \leq C|\alpha|.$

By the proof of Proposition 6.2, we get for $\widehat{\omega}_e \in H^2$,

$$\Lambda_3(\widehat{\omega}_e) = C\alpha^{-2}(\sin^2 y_c \mathcal{L}^2 \cap \sin y_c \mathcal{L}^\infty) + C\sin^2 y_c(1 + \alpha \sin y_c) \mathcal{L}^\infty.$$

Thus, by Lebesgue's dominated convergence theorem, we obtain

$$\begin{aligned} \widehat{\omega}(t,\alpha,0) &= \lim_{n \to +\infty} -\int_{u(0)}^{u(\pi)} \frac{\Lambda_3(\widehat{\omega}_e)\Lambda_4(g_n)}{u'(y_c)(\mathbf{A}_1^2 + \mathbf{B}_1^2)} e^{-i\alpha ct} dc \\ &= \int_{u(0)}^{u(\pi)} \frac{(u(\pi) - c)^2 \Lambda_3(\widehat{\omega}_e) e^{-i\alpha ct}}{u'(y_c) \phi_1'(0, c)(\mathbf{A}_1^2 + \mathbf{B}_1^2)} dc. \end{aligned}$$
By Proposition 4.5, we get $\frac{(u(\pi) - c)^2 \Lambda_3(\widehat{\omega}_e) e^{-i\alpha ct}}{u'(y_c) \phi_1'(0, c)(\mathbf{A}_1^2 + \mathbf{B}_1^2)} \in L^2_{y_c} \subset L^1_{y_c} \subset L^1_c.$

9.2. Decay estimates at critical points. First of all, we need the following formula of the stream function at a critical point.

Lemma 9.2. Let $\omega(t, x, y)$ be the solution to (1.2), and $-\Delta \psi = \omega$. Then it holds that

$$\begin{aligned} \widehat{\psi}(t,\alpha,0) &= \int_{-1}^{1} \frac{(1+\cos y_c)\sin y_c \Lambda_3(\widehat{\omega}_e) e^{-i\alpha ct}}{\phi_1'(0,c)(\mathbf{A}_1^2+\mathbf{B}_1^2)} dc, \\ \partial_y \widehat{\psi}(t,\alpha,0) &= \int_{-1}^{1} \frac{\sin y_c \Lambda_1(\widehat{\omega}_o)}{(\mathbf{A}^2+\mathbf{B}^2)\phi(0,c)} e^{-i\alpha ct} dc. \end{aligned}$$

Proof. By (2.1), we get

$$u''(y)i\alpha\widehat{\psi} = -\partial_t\widehat{\omega} - i\alpha u(y)\widehat{\omega},$$

which gives

$$\widehat{\psi}(t,\alpha,0) = \frac{-1}{i\alpha u''(0)} \left(\partial_t \widehat{\omega}_e(t,\alpha,0) + i\alpha u(0)\widehat{\omega}_e(t,\alpha,0)\right).$$

By Lemma 9.1, we get

$$\partial_t \widehat{\omega}(t, \alpha, 0) = \int_{u(0)}^{u(\pi)} -i\alpha c \frac{(u(\pi) - c)^2 \Lambda_3(\widehat{\omega}_e) e^{-i\alpha ct}}{u'(y_c) \phi_1'(0, c) (A_1^2 + B_1^2)} dc,$$

which implies the first identity.

Thanks to $\partial_y \psi(t, x, 0) = \partial_y \psi_o(t, x, 0)$, we get by Lemma 5.7 that

$$\begin{aligned} \partial_y \psi(t, x, 0) &= \frac{1}{2\pi} \int_{-1}^1 e^{-i\alpha tc} \alpha \partial_y \widetilde{\Phi}_o(\alpha, 0, c) dc \\ &= \frac{1}{2\pi} \int_{-1}^1 e^{-i\alpha tc} \frac{\alpha(\mu_-^o(c) - \mu_+^o(c))}{\phi(0, c)} dc, \\ &= \int_{-1}^1 \frac{\sin y_c \Lambda_1(\widehat{\omega}_o)}{(\mathbf{A}^2 + \mathbf{B}^2)\phi(0, c)} e^{-i\alpha tc} dc. \end{aligned}$$

This finishes the proof.

We denote

$$\begin{split} K_o^0(c,\alpha) &= \frac{\sin y_c \Lambda_1(\widehat{\omega}_o)(y_c)}{(\mathbf{A}^2 + \mathbf{B}^2)\phi(0,c)}, \\ K_e^0(c,\alpha) &= \frac{(1 + \cos y_c) \sin y_c \Lambda_3(\widehat{\omega}_e)}{\phi_1'(0,c)(\mathbf{A}_1^2 + \mathbf{B}_1^2)} = -\frac{J_1^1(c)\Lambda_3(\widehat{\omega}_e)(y_c)}{(\mathbf{A}_1^2 + \mathbf{B}_1^2)} \end{split}$$

Lemma 9.3. If
$$\|\widehat{\omega}_o\|_{H^2} + \|\widehat{\omega}''_o/u'\|_{L^2} \leq 1$$
, then we have
 $\partial_c K_o^0 = \alpha y_c^{-2} \sin y_c \mathcal{L}^\infty$, $\partial_c^2 K_o^0 = \alpha y_c^{-2} ((\sin y_c)^{-1} \mathcal{L}^\infty + \mathcal{L}^2).$

Proof. By the identity $\frac{\rho(c)}{\phi(0,c)} = -\frac{1-c}{\phi_1(0,c)}$ and Remark 3.9, we get (9.3) $\left|\frac{\rho(c)}{\phi(0,c)}\right| \leq C(1-c), \left|\partial_c \frac{\rho(c)}{\phi(0,c)}\right| \leq \left|\frac{1}{\phi_1(0,c)}\right| + \left|\frac{(c-1)\mathcal{G}}{\phi_1(0,c)^2}\right| \leq \frac{C\alpha(\pi-y_c)}{y_c} + C,$ and

(9.4)
$$\left| \partial_c^2 \frac{\rho(c)}{\phi(0,c)} \right| \le \left| \frac{2\mathcal{G} + (c-1)\partial_c \mathcal{G}}{\phi_1(0,c)^2} \right| + 2 \left| \frac{(c-1)\mathcal{G}^2}{\phi_1(0,c)^3} \right| \le \frac{C\alpha^2}{y_c^2} + \frac{C\alpha}{\sin y_c} \right|$$

Then by (5.20) and Lemma 7.5, we have

$$\partial_c \Lambda_{1,1}(\widehat{\omega}_o)(y_c) = \mathcal{L}^{\infty}, \quad \Lambda_{1,1}(\widehat{\omega}_o)(y_c) = \sin^2 y_c \mathcal{L}^{\infty}, \quad \partial_c^2 \Lambda_{1,1}(\widehat{\omega}_o)(y_c) = \frac{\mathcal{L}^2}{\sin y_c}.$$

By (5.21), Lemma 4.4 and Lemma 7.8, we obtain

$$\Lambda_{1,2}(\widehat{\omega}_o)(y_c) = |\alpha| \sin^2 y_c \mathcal{L}^{\infty}, \quad \partial_c \Lambda_{1,2}(\widehat{\omega}_o)(y_c) = |\alpha| \mathcal{L}^{\infty}, \quad \partial_c^2 \Lambda_{1,2}(\widehat{\omega}_o)(y_c) = \frac{|\alpha| \mathcal{L}^{\infty}}{\sin^2 y_c} + \frac{|\alpha| \mathcal{L}^2}{\sin y_c}.$$

Therefore, we obtain

(9.5) $\Lambda_1(\widehat{\omega}_o) = \alpha \rho \mathcal{L}^{\infty}, \quad \partial_c \Lambda_1(\widehat{\omega}_o) = \alpha \mathcal{L}^{\infty}, \quad \partial_c^2 \Lambda_1(\widehat{\omega}_o) = \rho^{-1} \alpha (\sin y_c \mathcal{L}^2 + \mathcal{L}^{\infty}).$ Notice that $K_o^0(c, \alpha) = \frac{\rho(c)}{\phi(0,c)} \frac{\Lambda_1(\widehat{\omega}_o)}{\sin y_c(A^2 + B^2)}.$ By (9.3), (9.4), (9.5) and Proposition 4.3, we deduce the result.

Lemma 9.4. Assume that $\widehat{\omega}_e \in H^2$ and $\frac{\partial_y \widehat{\omega}_e}{u'} \in H^1$. Then $K_e^0|_{c=\pm 1} = 0$ and $\|\partial_c^2 K_e^0\|_{L_c^1} \leq C |\alpha|^2 (\|\widehat{\omega}_e\|_{H^2} + \|\widehat{\omega}'_e/u'\|_{H^1}).$

Proof. We may assume $\|\widehat{\omega}_e\|_{H^2} + \|\widehat{\omega}'_e/u'\|_{H^1} \leq 1$ by normalization.

By Lemma 4.8, we obtain

$$\left|J_{1}^{1}(c)\right| \leq \frac{C}{\alpha^{2}}, \ \left|\partial_{c}J_{1}^{1}(c)\right| \leq \frac{C}{(1+\alpha\sin y_{c})^{2}}, \ \left|\partial_{c}^{2}J_{1}^{1}(c)\right| \leq \frac{C\alpha}{(1+\alpha\sin y_{c})^{3}\sin y_{c}}$$

By the proof of Proposition 6.2, we have

$$\Lambda_3(\widehat{\omega}_e) = \alpha^{-2} (\sin^2 y_c \mathcal{L}^2 \cap \sin y_c \mathcal{L}^\infty) + \sin^2 y_c (1 + \alpha \sin y_c) \mathcal{L}^\infty,$$

and by Lemma 7.4, we get $\partial_c \Lambda_{1,1}(\widehat{\omega}_e) = \mathcal{L}^2$; by Lemma 7.7 and Lemma 4.4, we get $\partial_c \Lambda_{1,2}(\widehat{\omega}_e) = \frac{\alpha \mathcal{L}^{\infty}}{\sin y_c}$. We also have $\Lambda_1(\widehat{\omega}_e) = (1 + \alpha \sin y_c)\mathcal{L}^{\infty}$. Then by Lemma 8.1, we get

$$\partial_c (\Lambda_3(\widehat{\omega}_e)) = \sin^2 y_c \mathcal{L}^2 + (1 + \alpha \sin y_c) \mathcal{L}^\infty$$

By Lemma 7.4, we get $\partial_c^2 \Lambda_{1,1}(\widehat{\omega}_e) = \frac{\mathcal{L}^2}{\sin^2 y_c}$; and by Lemma 7.8 and Lemma 4.4, we get $\partial_c^2 \Lambda_{1,2}(\widehat{\omega}_e) = \frac{\alpha \mathcal{L}^\infty}{\sin^3 y_c} + \frac{\alpha \mathcal{L}^2}{\sin y_c}$. Then by Lemma 8.1, we get

$$\partial_c^2 \left(\Lambda_3(\widehat{\omega}_e) \right) = \left(1 + \alpha \sin y_c + \frac{1}{\alpha \sin y_c} \right) \mathcal{L}^2 + \frac{\alpha^2}{(1 + \alpha \sin y_c)^2} \mathcal{L}^\infty + \frac{\alpha}{\sin y_c} \mathcal{L}^\infty.$$

Then by Proposition 4.5, we obtain

$$u'(y_c)\partial_c^2 K_e^0 = \frac{\alpha^3 \mathcal{L}^\infty}{(1+\alpha \sin y_c)^6} + \frac{\alpha \mathcal{L}^2}{(1+\alpha \sin y_c)^4} = \alpha^2 \mathcal{L}^1,$$

and

$$K_e^0 = C(\alpha) \sin y_c \mathcal{L}^\infty$$

which implies that $K_e^0|_{c=\pm 1} = 0$ and $\partial_c K_e^0 = \alpha^2 \mathcal{L}^{\infty}$.

Now we are in a position to prove Theorem 1.3.

Proof. Step 1. Let $c' \in (u(0), u(1))$, which will be determined later. Without loss of generality, let us consider the case of $|\alpha t| > 1$. Then by Lemma 9.2, we have

$$\begin{split} \partial_y \widehat{\psi}(t,\alpha,0) &= \int_{u(0)}^{u(\pi)} K_o^0(c,\alpha) e^{-i\alpha ct} dc \\ &= \frac{1}{i\alpha t} \int_{u(0)}^{u(\pi)} \partial_c K_o^0(c,\alpha) e^{-i\alpha ct} dc \\ &= \frac{1}{i\alpha t} \int_{u(0)}^{c'} \partial_c K_o^0(c,\alpha) e^{-i\alpha ct} dc \\ &- \frac{1}{(\alpha t)^2} \left(\partial_c K_o^0(c',\alpha) e^{-i\alpha c't} + \int_{c'}^{u(\pi)} \partial_c^2 K_o^0(c,\alpha) e^{-i\alpha ct} dc \right). \end{split}$$

By Lemma 9.3, we have

$$\begin{aligned} \|\partial_c^2 K_o^0\|_{L^1_c(c',u(\pi))} &= \|u'(y_c)\partial_c^2 K_o^0\|_{L^1_{y_c}(y_{c'},1)} \\ &\leq C\alpha(\|y_c^{-2}\|_{L^1_{y_c}(y_{c'},1)} + \|y_c^{-1}\|_{L^2_{y_c}(y_{c'},1)}) \leq C\alpha y_{c'}^{-1}. \end{aligned}$$

Therefore, we deduce that for any $y_{c'} \in (0, 1)$,

$$|\partial_y \widehat{\psi}(t,\alpha,0)| \le \frac{C\alpha}{|\alpha t|} \int_{u(0)}^{c'} \frac{dc}{y_c} + C \frac{1}{(\alpha t)^2} \frac{\alpha}{y_{c'}} \le C \left(\frac{y_{c'}}{|t|} + \frac{1}{\alpha y_{c'} t^2}\right)$$

Taking $y_{c'} = |\alpha t|^{-\frac{1}{2}}$, we deduce that

$$|\partial_y \widehat{\psi}(t, \alpha, 0)| \le C \alpha^{-\frac{1}{2}} |t|^{-\frac{3}{2}}.$$

Step 2. Using the fact that $\widehat{\psi}(t, \alpha, 0) = \int_{-1}^{1} K_{e}^{0}(c, \alpha) e^{-i\alpha ct} dc$, and Lemma 9.4, we obtain

$$\begin{aligned} |\widehat{\psi}(t,\alpha,0)| &= \left| \frac{1}{i\alpha t} \int_{-1}^{1} \partial_{c} K_{e}^{0}(c,\alpha) e^{-i\alpha c t} dc \right| \\ &\leq \left| \frac{1}{\alpha^{2} t^{2}} \int_{-1}^{1} \partial_{c}^{2} K_{e}^{0}(c,\alpha) e^{-i\alpha c t} dc \right| + \frac{1}{\alpha^{2} t^{2}} \|\partial_{c} K_{e}^{0}(c,\alpha)\|_{L^{\infty}} \\ &\leq C t^{-2} (\|\widehat{\omega}_{e}\|_{H^{2}} + \|\widehat{\omega}_{e}'/u'\|_{H^{1}}). \end{aligned}$$

Let $W(t, x, y) = \omega(t, x + u(y)t, y)$. Then $\widehat{W}(t, \alpha, y) = e^{i\alpha t u(y)}\widehat{\omega}(t, \alpha, y)$ and $\partial_t \widehat{W} = -i\alpha e^{i\alpha u(y)t} u''\widehat{\psi}.$

By Lemma 9.1, $\widehat{\omega}(t, \alpha, 0) \to 0$ as $t \to \infty$, so does $\widehat{W}(t, \alpha, 0)$. Thus,

$$\widehat{W}(t,\alpha,0) = \int_{t}^{\infty} i\alpha e^{i\alpha u(0)s} u''(0)\widehat{\psi}(s,\alpha,0)ds$$

Thus, if t > 1 then

$$\begin{aligned} |\widehat{\omega}(t,\alpha,0)| &= |\widehat{W}(t,\alpha,0)| \le C|\alpha| \int_{t}^{+\infty} |\widehat{\psi}(s,\alpha,0)| ds \\ &\le C|\alpha| \int_{t}^{+\infty} s^{-2} (\|\widehat{\omega}_{e}\|_{H^{2}} + \|\widehat{\omega}_{e}'/u'\|_{H^{1}}) ds \\ &\le C|\alpha||t|^{-1} (\|\widehat{\omega}_{e}\|_{H^{2}} + \|\widehat{\omega}_{e}'/u'\|_{H^{1}}), \end{aligned}$$

and if t < -1 then

$$\begin{aligned} |\widehat{\omega}(t,\alpha,0)| &= |\widehat{W}(t,\alpha,0)| \le C|\alpha| \int_{-\infty}^{t} |\widehat{\psi}(s,\alpha,0)| ds \\ &\le C|\alpha| |t|^{-1} (\|\widehat{\omega}_e\|_{H^2} + \|\widehat{\omega}'_e/u'\|_{H^1}). \end{aligned}$$

Step 3. By Step 1 and Step 2, we deduce that

$$\begin{split} |\omega(t,x,0)| &= \left| \sum_{\alpha \neq 0} \widehat{\omega}(t,\alpha,0) e^{i\alpha x} \right| \leq \sum_{\alpha \neq 0} |\widehat{\omega}(t,\alpha,0)| \leq \sum_{\alpha \neq 0} C|\alpha| |t|^{-1} \|\widehat{\omega}_{e}(\alpha,\cdot)\|_{H^{3}} \\ &\leq C|t|^{-1} \Big(\sum_{\alpha \neq 0} |\alpha|^{-2s} \Big)^{\frac{1}{2}} \Big(\sum_{\alpha \neq 0} |\alpha|^{2+2s} \|\widehat{\omega}_{e}(\alpha,\cdot)\|_{H^{3}}^{2} \Big)^{\frac{1}{2}} \leq C|t|^{-1} \|\omega_{e}\|_{H^{s+1}_{x}H^{3}_{y}}, \\ |V^{2}(t,x,0)| &= \left| \sum_{\alpha \neq 0} i\alpha \widehat{\psi}(t,\alpha,0) e^{i\alpha x} \right| \leq \sum_{\alpha \neq 0} |\alpha| |\widehat{\psi}(t,\alpha,0)| \leq \sum_{\alpha \neq 0} C|\alpha| |t|^{-2} \|\widehat{\omega}_{e}(\alpha,\cdot)\|_{H^{3}} \\ &\leq C|t|^{-2} \Big(\sum_{\alpha \neq 0} |\alpha|^{-2s} \Big)^{\frac{1}{2}} \Big(\sum_{\alpha \neq 0} |\alpha|^{2+2s} \|\widehat{\omega}_{e}(\alpha,\cdot)\|_{H^{3}}^{2} \Big)^{\frac{1}{2}} \leq C|t|^{-2} \|\omega_{e}\|_{H^{s+1}_{x}H^{3}_{y}}, \\ |V^{1}(t,x,0)| &= \left| \sum_{\alpha \neq 0} \partial_{y} \widehat{\psi}(t,\alpha,0) e^{i\alpha x} \right| \leq \sum_{\alpha \neq 0} |\partial_{y} \widehat{\psi}(t,\alpha,0)| \leq \sum_{\alpha \neq 0} C|\alpha|^{-\frac{1}{2}} |t|^{-\frac{3}{2}} \|\widehat{\omega}_{o}(\alpha,\cdot)\|_{H^{3}} \\ &\leq C|t|^{-\frac{3}{2}} \Big(\sum_{\alpha \neq 0} |\alpha|^{-2s} \Big)^{\frac{1}{2}} \Big(\sum_{\alpha \neq 0} |\alpha|^{2s-1} \|\widehat{\omega}_{o}(\alpha,\cdot)\|_{H^{3}}^{2} \Big)^{\frac{1}{2}} \leq C|t|^{-\frac{3}{2}} \|\omega_{o}\|_{H^{s-\frac{1}{2}}_{x}H^{3}_{y}}. \end{split}$$

The above results also hold with 0 replaced by π , thus all $k\pi(k \in \mathbb{Z})$.

10. WAVE OPERATOR

To prove the enhanced dissipation, we use the wave operator method introduced in [21].

10.1. Basic properties of wave operator.

Definition 10.1. Let $|\alpha| > 1$. Let $\phi(y,c)$ be defined in Proposition 3.6 with $\phi_1(y,c) = \frac{\phi(y,c)}{u(y)-c}$ for $(y,c) \in [0,\pi] \times [-1,1]$. For any function $\varphi(y)$ defined on $[-\pi,\pi]$, we define

$$-\mathbb{D}(\varphi_o)(-y_c) = \mathbb{D}(\varphi_o)(y_c) = \frac{\Lambda_1(\varphi_o)(y_c)}{(\mathbf{A}^2 + \mathbf{B}^2)^{\frac{1}{2}}},$$
$$\mathbb{D}(\varphi_e)(-y_c) = \mathbb{D}(\varphi_e)(y_c) = \frac{\Lambda_3(\varphi_e)(y_c)}{(\mathbf{A}_1^2 + \mathbf{B}_1^2)^{\frac{1}{2}}},$$

where $\varphi_o = \frac{\varphi(y) - \varphi(-y)}{2}$ and $\varphi_e = \frac{\varphi(y) + \varphi(-y)}{2}$.

The operator \mathbb{D} is called the wave operator, which has the following basic properties.

Proposition 10.2. It holds that

$$\mathbb{D}\big(\cos y(1+(\partial_y^2-\alpha^2)^{-1})\omega\big)=\cos y\mathbb{D}(\omega),\\\|\mathbb{D}(\omega)\|_{L^2}^2=\langle\omega,\omega+(\partial_y^2-\alpha^2)^{-1}\omega\rangle.$$

Moreover, there exists a constant C independent of α so that

$$(1 - \alpha^{-2}) \|\omega\|_{L^2}^2 \le \|\mathbb{D}(\omega)\|_{L^2}^2 \le \|\omega\|_{L^2}^2, \|\sin y \mathbb{D}(\omega)\|_{L^2}^2 \ge \|\partial_y \psi\|_{L^2}^2 + (\alpha^2 - 1) \|\psi\|_{L^2}^2,$$

$$\|\partial_y \mathbb{D}(\omega)\|_{L^2} \le C |\alpha|^{\frac{1}{2}} \|\omega\|_{H^1}, \quad \|\partial_y^2 \mathbb{D}(\omega)\|_{L^2} \le C |\alpha|^{\frac{3}{2}} \|\omega\|_{H^2}.$$

where $-(\partial_y^2 - \alpha^2)\psi = \omega$.

Proof. Step 1. If ω is odd, then ψ is odd and $\psi(0) = \psi(\pi) = 0$. Then we have

$$\begin{split} & \Pi_{1}(u''\psi) \\ &= p.v. \int_{0}^{\pi} \frac{\int_{y_{c}}^{y} \psi(y')u''(y')\phi_{1}(y',c)dy'}{\phi(y,c)^{2}}dy \\ &= p.v. \int_{0}^{\pi} \frac{\int_{y_{c}}^{y} \psi(y')(\partial_{y}^{2} - \alpha^{2})\phi(y',c)dy'}{\phi(y,c)^{2}}dy \\ &= -\int_{0}^{\pi} \frac{\int_{y_{c}}^{y} \omega(y')\phi(y',c)dy'}{\phi(y,c)^{2}}dy + p.v. \int_{0}^{\pi} \frac{\psi(y)\partial_{y}\phi(y,c) - u'(y_{c})\psi(y_{c})}{\phi(y,c)^{2}}dy \\ &= -\int_{0}^{\pi} \frac{\partial_{y}\psi(y)\phi(y,c)}{\phi(y,c)^{2}}dy \\ &= -\int_{0}^{\pi} \frac{\int_{y_{c}}^{y} \omega(y')\phi(y',c)dy'}{\phi(y,c)^{2}}dy - \frac{\psi(y)}{\phi(y,c)}\Big|_{y=0}^{\pi} - \frac{\psi(y_{c})}{\rho(c)}A \\ &= u(y_{c})\Pi_{1}(\omega) - \frac{\psi(y_{c})}{\rho(c)}A - p.v. \int_{0}^{\pi} \frac{\int_{y_{c}}^{y} \omega(y')u(y')\phi_{1}(y',c)dy'}{\phi(y,c)^{2}}dy. \end{split}$$

Here we used Remark 5.9. Thus, we get

$$\rho \mathrm{II}_1 (u'' \psi + u\omega) = \rho u \mathrm{II}_1(\omega) - \psi(y_c) \mathrm{A}.$$

Recall that $\Lambda_1(\varphi) = A\varphi + u'' \rho II_1(\varphi)$. This shows that for ω odd,

$$\mathbb{D}(u''\psi + u\omega) = u\mathbb{D}(\omega).$$

If ω is even, then ψ is also even and $\partial_y \psi(0) = \partial_y \psi(\pi) = 0$. Then we have

$$\begin{split} & \Pi_{1}(u''(y)\psi(\alpha,y)) \\ &= p.v. \int_{0}^{\pi} \frac{\int_{y_{c}}^{y} \psi(y')u''(y')\phi_{1}(y',c)dy'}{\phi(y,c)^{2}}dy \\ &= p.v. \int_{0}^{\pi} \frac{\int_{y_{c}}^{y} \psi(y')(\partial_{y}^{2} - \alpha^{2})\phi(y',c)dy'}{\phi(y,c)^{2}}dy \\ &= -\int_{0}^{\pi} \frac{\int_{y_{c}}^{y} \omega(y')\phi(y',c)dy'}{\phi(y,c)^{2}}dy + p.v. \int_{0}^{\pi} \frac{\psi(y)\partial_{y}\phi(y,c) - u'(y_{c})\psi(y_{c})}{\phi(y,c)^{2}}dy \\ &- p.v. \int_{0}^{\pi} \frac{\partial_{y}\psi(y)\phi(y,c)}{\phi(y,c)^{2}}dy \\ &= -\int_{0}^{\pi} \frac{\int_{y_{c}}^{y} \omega(y')\phi(y',c)dy'}{\phi(y,c)^{2}}dy - \frac{\psi(y)}{\phi(y,c)}\Big|_{y=0}^{\pi} - \frac{\psi(y_{c})}{\rho(c)}A \\ &= u(y_{c})\Pi_{1}(\omega) - \frac{\psi(y_{c})}{\rho(c)}A - \Pi_{1}(u\omega) - \frac{\psi(\pi)}{\phi(\pi,c)} + \frac{\psi(0)}{\phi(0,c)}, \end{split}$$

and for j = 0, 1,

$$j = 0, 1,$$

$$E_j(u''\psi) = \int_{y_c}^{j\pi} u''\psi\phi_1(y,c) \, dy = \int_{y_c}^{j\pi} \psi(\partial_y^2 - \alpha^2)\phi(y',c) \, dy$$

LINEAR INVISCID DAMPING AND ENHANCED DISSIPATION

$$= -\int_{y_c}^{j\pi} \omega \phi(y',c) \, dy + \psi \partial_y \phi(y',c) \Big|_{y'=y_c}^{j\pi} - \partial_y \psi \phi(y',c) \Big|_{y'=y_c}^{j\pi}$$
$$= u(y_c) \mathcal{E}_j(\omega) - \mathcal{E}_j(u\omega) + \psi(j\pi)(u(j\pi)-c) \partial_y \phi_1(j\pi,c) - \psi(y_c)u'(y_c)$$

Thus, we obtain

$$\begin{split} \Lambda_{3}(u''\psi + u\omega) \\ &= \rho(c)\Lambda(u''\psi + u\omega) + u''\rho^{2}\Pi_{1}(u''\psi + u\omega) \\ &- \frac{u'(y_{c})(u(\pi) - c)^{2}}{\phi_{1}(0, c)\phi_{1}'(0, c)} \left(\frac{u''(y_{c})}{u'(y_{c})} E_{0}(u''\psi + u\omega) + u''\psi + u\omega\right) \\ &+ \frac{u'(y_{c})(u(0) - c)^{2}}{\phi_{1}(\pi, c)\phi_{1}'(\pi, c)} \left(\frac{u''(y_{c})}{u'(y_{c})} E_{1}(u''\psi + u\omega) + u''\psi + u\omega\right) \\ &= \rho(c)\Lambda(u''\psi + u\omega) + u''\rho^{2} \left(u(y_{c})\Pi_{1}(\omega) - \frac{\Lambda\psi}{\rho(c)} - \frac{\psi(\pi)}{\phi(\pi, c)} + \frac{\psi(0)}{\phi(0, c)}\right) \\ &- \frac{u'(y_{c})(u(\pi) - c)^{2}}{\phi_{1}(0, c)\phi_{1}'(0, c)} \left(\frac{u''(y_{c})}{u'(y_{c})} (uE_{0}(\omega) + \psi(0)(u(0) - c)\phi_{1}'(0, c) - u'\psi) + u''\psi + u\omega\right) \\ &+ \frac{u'(y_{c})(u(0) - c)^{2}}{\phi_{1}(\pi, c)\phi_{1}'(\pi, c)} \left(\frac{u''(y_{c})}{u'(y_{c})} (uE_{1}(\omega) + \psi(\pi)(u(\pi) - c)\phi_{1}'(\pi, c) - u'\psi) + u''\psi + u\omega\right) \\ &= u(y_{c})\Lambda_{3}(\omega). \end{split}$$

This shows that for ω even,

$$\mathbb{D}(u''\psi + u\omega) = u\mathbb{D}(\omega).$$

Step 2. Let $\omega = \omega_o + \omega_e$ with $\omega_o = \frac{\omega(y) - \omega(-y)}{2}$ and $\omega_e = \frac{\omega(y) + \omega(-y)}{2}$. Let $(\partial_y^2 - \alpha^2)g_o = \overline{\omega_o} + (\partial_y^2 - \alpha^2)\overline{\omega_o}$, where we denote by \overline{f} the complex conjugate of f. Then by Lemma 5.8, we get

$$\langle \psi_o, \omega_o + (\partial_y^2 - \alpha^2) \omega_o \rangle = -2 \int_{u(0)}^{u(\pi)} \frac{\Lambda_1(\omega_o) \Lambda_2(g_o)}{(\mathbf{A}^2 + \mathbf{B}^2) u'(y_c)} dc.$$

As $g_o = \overline{\omega_o} - \overline{\psi_o}$, we have

$$\Lambda_2(g_o) = \rho II_1(u''g_o) + Ag_o = -\rho II_1(u''\overline{\psi_o} + u\overline{\omega_o}) + Ag_o$$
$$= -\rho u II_1(\overline{\omega_o}) + A\overline{\psi_o} + Ag_o = \rho u'' II_1(\overline{\omega_o}) + A\overline{\omega_o} = \Lambda_1(\overline{\omega_o})$$

This shows that

$$\begin{aligned} \langle \omega_o, \omega_o - \psi_o \rangle &= -\langle \psi_o, \omega_o + (\partial_y^2 - \alpha^2) \omega_o \rangle \\ &= 2 \int_{u(0)}^{u(\pi)} \frac{\Lambda_1(\omega_o) \overline{\Lambda_1(\omega_o)}}{(\mathbf{A}^2 + \mathbf{B}^2) u'(y_c)} dc = \| \mathbb{D}(\omega_o) \|_{L^2}^2. \end{aligned}$$

Let $(\partial_y^2 - \alpha^2)g_e = \overline{\omega_e} + (\partial_y^2 - \alpha^2)\overline{\omega_e}$. Then we have

$$\langle \psi_e, \omega_e + (\partial_y^2 - \alpha^2) \omega_e \rangle = -2 \int_{u(0)}^{u(\pi)} \frac{\Lambda_3(\omega_e) \Lambda_4(g_e)}{(\Lambda_1^2 + B_1^2) u'(y_c)} dc.$$

Due to $g_e = \overline{\omega_e} - \overline{\psi_e}$, we have

$$u''(y_c)\Lambda_4(g_e) = \Lambda_3(u''g_e) = -\Lambda_3(u''\overline{\psi_e} + u\overline{\omega_e})$$

= $-u(y_c)\Lambda_3(\overline{\omega_e}) = u''(y_c)\Lambda_3(\overline{\omega_e}),$

thus, $\Lambda_4(g_e) = \Lambda_3(\overline{\omega_e})$ a.e. and

$$\begin{aligned} \langle \omega_e, \omega_e - \psi_e \rangle &= -\langle \psi_e, \omega_e + (\partial_y^2 - \alpha^2)\omega_e \rangle \\ &= 2 \int_{u(0)}^{u(\pi)} \frac{\Lambda_3(\omega_e)\overline{\Lambda_3(\omega_e)}}{(A_1^2 + B_1^2)u'(y_c)} dc = \|\mathbb{D}(\omega_e)\|_{L^2}^2. \end{aligned}$$

Thus we get $\|\mathbb{D}(\omega)\|_{L^2}^2 = \langle \omega, \omega + (\partial_y^2 - \alpha^2)^{-1}\omega \rangle.$

Step 3. Let $-(\partial_y^2 - \alpha^2)\psi = \omega$. Then we have

$$\|\omega\|_{L^2}^2 = \|\partial_y^2\psi\|_{L^2}^2 + 2\alpha^2 \|\partial_y\psi\|_{L^2}^2 + \alpha^4 \|\psi\|_{L^2}^2 \ge \alpha^2 (\|\partial_y\psi\|_{L^2}^2 + \alpha^2 \|\psi\|_{L^2}^2),$$

which gives

$$\|\mathbb{D}(\omega)\|_{L^{2}}^{2} = \langle \omega, \omega - \psi \rangle = \|\omega\|_{L^{2}}^{2} - (\|\partial_{y}\psi\|_{L^{2}}^{2} + \alpha^{2}\|\psi\|_{L^{2}}^{2}) \le \|\omega\|_{L^{2}}^{2},$$

$$\|\mathbb{D}(\omega)\|_{L^{2}}^{2} \ge (1 - \alpha^{-2})\|\omega\|_{L^{2}}^{2}.$$

$$\begin{split} \|\sin y \mathbb{D}(\omega)\|_{L^{2}}^{2} &= \|\mathbb{D}(\omega)\|_{L^{2}}^{2} - \|\cos y \mathbb{D}(\omega)\|_{L^{2}}^{2} \\ &= \|\mathbb{D}(\omega)\|_{L^{2}}^{2} - \|\mathbb{D}(\cos y(\omega - \psi))\|_{L^{2}}^{2} \\ &\geq \|\mathbb{D}(\omega)\|_{L^{2}}^{2} - \|\cos y(\omega - \psi)\|_{L^{2}}^{2} \geq \|\mathbb{D}(\omega)\|_{L^{2}}^{2} - \|\omega - \psi\|_{L^{2}}^{2} \\ &= \langle \omega, \omega - \psi \rangle - \|\omega - \psi\|_{L^{2}}^{2} = \langle \psi, \omega - \psi \rangle \\ &= \|\partial_{y}\psi\|_{L^{2}}^{2} + (\alpha^{2} - 1)\|\psi\|_{L^{2}}^{2} \geq 0. \end{split}$$

Step 4. By Proposition 4.3, we obtain

$$\left| \partial_{y_c} \left(\frac{1}{(\mathbf{A}^2 + \mathbf{B}^2)^{1/2}} \right) \right| \le C \left| (\mathbf{A}^2 + \mathbf{B}^2)^{1/2} \sin y_c \partial_c \left(\frac{1}{\mathbf{A}^2 + \mathbf{B}^2} \right) \right|$$
$$\le \frac{C}{(1 + \alpha \sin y_c) \sin y_c},$$

and by the proof of Proposition 6.1, we get for $\omega \in H^1(\mathbb{T})$ odd, $\|\omega\|_{H^1} = 1$,

$$\Lambda_1(\omega)(y_c) = \sin y_c ((1+|\alpha|\sin y_c)\mathcal{L}^2 + |\alpha|^{\frac{1}{2}}\mathcal{L}^{\infty}),$$

$$\partial_{y_c}\Lambda_1(\omega)(y_c) = \left((1+|\alpha|\sin y_c)\mathcal{L}^2 + \alpha^{\frac{1}{2}}\mathcal{L}^{\infty}\right).$$

Thus, for $\omega \in H^1(\mathbb{T})$ odd, $\|\omega\|_{H^1} = 1$,

$$\partial_{y_c} \mathbb{D}(\omega) = \Lambda_1(\omega)(y_c) \partial_{y_c} \left(\frac{1}{(\mathbf{A}^2 + \mathbf{B}^2)^{1/2}}\right) + \frac{\partial_{y_c} \Lambda_1(\omega)(y_c)}{(\mathbf{A}^2 + \mathbf{B}^2)^{1/2}}$$
$$= \mathcal{L}^2 + \frac{|\alpha|^{\frac{1}{2}} \mathcal{L}^{\infty}}{1 + \alpha \sin y_c} = \mathcal{L}^2.$$

Thus, for $\omega \in H^1(\mathbb{T})$ odd,

 $\|\mathbb{D}(\omega)\|_{H^1} \le C \|\omega\|_{H^1}.$

By Proposition 4.3, we get

$$\left|\partial_{y_c}^2 \left(\frac{1}{(\mathbf{A}^2 + \mathbf{B}^2)^{1/2}}\right)\right| \le \frac{C}{(1 + \alpha \sin y_c) \sin^2 y_c}$$

and by the proof of Proposition 6.1, for $\omega \in H^2(\mathbb{T})$ odd, $\|\omega\|_{H^2} = 1$, we get $\Lambda_1(\omega) = (\sin^2 y_c \mathcal{L}^2 \cap \sin y_c \mathcal{L}^\infty) + |\alpha| \sin^2 y_c \mathcal{L}^\infty$,

$$\partial_{y_c}(\Lambda_1(\omega)) = \sin y_c(\mathcal{L}^2 + \alpha \mathcal{L}^\infty), \partial_{y_c}^2(\Lambda_1(\omega)) = ((1 + |\alpha| \sin y_c)\mathcal{L}^2 + |\alpha|\mathcal{L}^\infty)$$

Thus for $\omega \in H^2(\mathbb{T})$ odd, $\|\omega\|_{H^2} = 1$, we get

$$\partial_{y_c}^2 \mathbb{D}(\omega) = \Lambda_1(\omega) \partial_{y_c}^2 \left(\frac{1}{(\mathbf{A}^2 + \mathbf{B}^2)^{1/2}} \right) + \frac{\partial_{y_c}^2 \Lambda_1(\omega)}{(\mathbf{A}^2 + \mathbf{B}^2)^{1/2}} + 2\partial_{y_c} \Lambda_1(\omega) \partial_{y_c} \left(\frac{1}{(\mathbf{A}^2 + \mathbf{B}^2)^{1/2}} \right)$$
$$= \mathcal{L}^2 + \frac{|\alpha|^{\frac{1}{2}} \mathcal{L}^{\infty}}{1 + \alpha \sin y_c} + \frac{|\alpha| \mathcal{L}^{\infty}}{1 + \alpha \sin y_c} = \alpha^{\frac{1}{2}} \mathcal{L}^2.$$

Thus, for $\omega \in H^2(\mathbb{T})$ odd,

$$\|\mathbb{D}(\omega)\|_{H^2} \le C\alpha^{\frac{1}{2}} \|\omega\|_{H^2}.$$

By Proposition 4.5, we get

$$\frac{1}{(\mathbf{A}_1^2 + \mathbf{B}_1^2)^{\frac{1}{2}}} \le \frac{C\alpha^2}{(1 + \alpha \sin y_c)^3},\\ \partial_{y_c} \left(\frac{1}{(\mathbf{A}_1^2 + \mathbf{B}_1^2)^{\frac{1}{2}}}\right) \le \frac{C\alpha^2}{(1 + \alpha \sin y_c)^3 \sin y_c},\\ \partial_{y_c}^2 \left(\frac{1}{(\mathbf{A}_1^2 + \mathbf{B}_1^2)^{\frac{1}{2}}}\right) \le \frac{C\alpha^2}{(1 + \alpha \sin y_c)^3 \sin^2 y_c}$$

For $\omega \in H^1$ even, $\|\omega\|_{H^1} = 1$, by the proof of Proposition 6.2, we get

$$\Lambda_3(\omega) = |\alpha|^{-\frac{3}{2}} \sin y_c (1+|\alpha| \sin y_c)^3 \mathcal{L}^2,$$

$$\partial_{y_c} (\Lambda_3(\omega)) = |\alpha|^{-\frac{3}{2}} (1+|\alpha| \sin y_c)^3 \mathcal{L}^2.$$

Thus, for $\omega \in H^1$ even, $\|\omega\|_{H^1} = 1$,

$$\partial_{y_c} \mathbb{D}(\omega) = \Lambda_3(\omega)(y_c) \partial_{y_c} \left(\frac{1}{(A_1^2 + B_1^2)^{1/2}} \right) + \frac{\partial_{y_c} \Lambda_3(\omega)(y_c)}{(A_1^2 + B_1^2)^{1/2}} = \alpha^{\frac{1}{2}} \mathcal{L}^2.$$

This shows that for $\omega \in H^1$ even,

$$\|\mathbb{D}(\omega)\|_{H^1} \le C |\alpha|^{\frac{1}{2}} \|\omega\|_{H^1}.$$

For $\omega \in H^2$ even, $\|\omega\|_{H^2} = 1$, by the proof of Proposition 6.2, we can get

$$\partial_{y_c}^m \Lambda_3(\omega) = \alpha^{-1/2} (\sin y_c)^{2-m} (1 + \alpha \sin y_c)^3 \mathcal{L}^2, \quad m = 0, 1, 2.$$

Thus, for $\omega \in H^2$ even, $\|\omega\|_{H^2} = 1$,

$$\begin{aligned} \partial_{y_c}^2 \mathbb{D}(\omega) &= \Lambda_3(\omega) \partial_{y_c}^2 \left(\frac{1}{(\mathbf{A}_1^2 + \mathbf{B}_1^2)^{1/2}} \right) + \frac{\partial_{y_c}^2 \Lambda_3(\omega)}{(\mathbf{A}_1^2 + \mathbf{B}_1^2)^{1/2}} + 2\partial_{y_c} \Lambda_3(\omega) \partial_{y_c} \left(\frac{1}{(\mathbf{A}_1^2 + \mathbf{B}_1^2)^{1/2}} \right) \\ &= \alpha^{\frac{3}{2}} \mathcal{L}^2. \end{aligned}$$

This shows that for $\omega \in H^2$ even,

$$\|\mathbb{D}(\omega)\|_{H^2} \le C |\alpha|^{\frac{3}{2}} \|\omega\|_{H^2}.$$

This completes the proof of the proposition.

10.2. Commutator estimate.

Proposition 10.3. It holds that

$$\|\sin y(\mathbb{D}(\partial_y^2\omega) - \partial_y^2\mathbb{D}(\omega))\|_{L^2} \le C(|\alpha|\|\omega\|_{L^2} + \|\partial_y\omega\|_{L^2}).$$

We need following lemmas.

Lemma 10.4. It holds that

$$\|\sin y_c[\partial_y^2, \rho \Pi_{1,1}]\omega\|_{L^2} \le C \|\omega\|_{H^1}.$$

Proof. By (7.1), we get

$$\rho \operatorname{II}_{1,1}(\omega) = -\frac{1}{2}H(Z \circ \operatorname{In}(\omega)) - \frac{1}{2}\int_{-\pi}^{\pi} \operatorname{In}(\omega)(y')dy'.$$

As $Z \circ \operatorname{In}(\omega)$ is odd, so does $\partial_{y_c}^2 Z \circ \operatorname{In}(\omega)$. Thus,

$$\sin y_c \partial_{y_c}^2 \rho \mathrm{II}_{1,1}(\omega) = -\frac{1}{2} H(\sin y_c \partial_{y_c}^2 Z \circ \mathrm{In}(\omega))$$

Then we get

$$\sin y_c[\partial_y^2, \rho \mathrm{II}_{1,1}]\omega = -\frac{1}{2}H(\sin y_c[\partial_{y_c}^2, Z \circ \mathrm{In}](\omega)) + \frac{\sin y_c}{2}\int_{-\pi}^{\pi} \mathrm{In}(\partial_y^2\omega)(y')dy'.$$

On one hand, we have

$$\frac{1}{2} \int_{-\pi}^{\pi} \ln(\partial_y^2 \omega)(y) dy = \int_0^{\pi} \ln(\partial_y^2 \omega)(y) dy = \int_0^{\pi} \omega'(y) - \omega'(0) - \sin^2 \frac{y}{2} (\omega'(\pi) - \omega'(0)) dy$$
$$= \omega(\pi) - \omega(0) - \pi(\omega'(0) + \omega'(\pi))/2.$$

On the other hand,

$$\sin y_c \left(Z \circ \operatorname{In}(\partial_y^2 \omega) - \partial_y^2 Z \circ \operatorname{In}(\omega) \right) \\ = \frac{\omega'(0) + \omega'(\pi)}{2} \cos y + \frac{\omega'(0) - \omega'(\pi)}{2} - \frac{2\omega}{\sin y} + \frac{2\cos y}{\sin^2 y} \operatorname{In}(\omega)(y) - \frac{\operatorname{Int}(\omega)(\pi)}{2} + \frac{2\cos y}{\cos^2 y} + \frac{2\cos y}{\cos^2 y} \operatorname{In}(\omega)(y) - \frac{\operatorname{Int}(\omega)(\pi)}{2} + \frac{2\cos y}{\cos^2 y} +$$

and $H(\cos y) = 2\pi \sin y$. Then we deduce that

$$\sin y_c [\partial_y^2, \rho \mathrm{II}_{1,1}] \omega = \frac{1}{2} H \Big(-\frac{2\omega}{\sin y} + \frac{2\cos y}{\sin^2 y} \mathrm{In}(\omega)(y) \Big) + \sin y_c(\omega(\pi) - \omega(0)),$$

from which, it follows that

$$\begin{split} \left\| \sin y_c [\partial_y^2, \rho \mathrm{II}_{1,1}] \omega \right\|_{L^2} &\leq C \left\| H \Big(-\frac{2\omega}{\sin y} + \frac{2\cos y}{\sin^2 y} \mathrm{In}(\omega)(y) \Big) \right\|_{L^2} + C \|\omega\|_{L^\infty} \\ &\leq C \left\| -\frac{2\omega}{\sin y} + \frac{2\cos y}{\sin^2 y} \mathrm{In}(\omega)(y) \right\|_{L^2} + C \|\omega\|_{L^\infty} \\ &\leq C \left\| -\frac{\omega}{\sin y} + \frac{\cos y}{\sin^2 y} \int_0^y \omega(y') dy' \right\|_{L^2(0,\frac{\pi}{2})} \\ &+ C \left\| -\frac{\omega}{\sin y} + \frac{\cos y}{\sin^2 y} \int_\pi^y \omega(y') dy' \right\|_{L^2[\frac{\pi}{2},\pi]} + C \|\omega\|_{L^\infty}. \end{split}$$

Using the fact that

$$-\frac{\omega}{\sin y} + \frac{\cos y}{\sin^2 y} \int_0^y \omega(y') dy' = \frac{-1}{\sin^2 y} \Big(\int_0^y \sin y' \omega'(y') dy' + \int_0^y \omega(y') (\cos y' - \cos y) dy' \Big),$$
$$-\frac{\omega}{\sin y} + \frac{\cos y}{\sin^2 y} \int_{\pi}^{y} \omega(y') dy' = \frac{-1}{\sin^2 y} \Big(\int_{\pi}^{y} \sin y' \omega'(y') dy' + \int_{\pi}^{y} \omega(y') (\cos y' - \cos y) dy' \Big),$$

and Hardy's inequality, we infer that

$$\left\| -\frac{\omega}{\sin y} + \frac{\cos y}{\sin^2 y} \int_0^y \omega(y') dy' \right\|_{L^2(0,\frac{\pi}{2})} + \left\| -\frac{\omega}{\sin y} + \frac{\cos y}{\sin^2 y} \int_{\pi}^y \omega(y') dy' \right\|_{L^2(\frac{\pi}{2},\pi)} \le C \|\omega\|_{H^1}.$$

Summing up, we conclude the lemma.

Lemma 10.5. It holds that

$$|\sin^3 y_c[\partial_{y_c}^2, \mathcal{L}_0]\varphi| \le C\alpha^{\frac{1}{2}} (\|\varphi\|_{H^1} + \alpha \|\varphi\|_{L^2}).$$

Proof. By (7.6), we get

$$\begin{aligned} \partial_{y_c}^2 \mathcal{L}_0(\varphi) &= u'(y_c) \partial_c \left(\mathcal{L}_0(\varphi') - I_{0,1}(\varphi) + I_{0,0}(\varphi) \right) + u'(y_c) \partial_c (u'(y_c) \mathcal{L}_1(\varphi)) \\ &= \mathcal{L}_0(\varphi'') - I_{0,1}(\varphi') + I_{0,0}(\varphi') + u'(y_c) \mathcal{L}_1(\varphi') \\ &- u'(y_c) \partial_c I_{0,1}(\varphi) + u'(y_c) \partial_c I_{0,0}(\varphi) + u''(y_c) \mathcal{L}_1(\varphi) \\ &+ u'(y_c) \left(\mathcal{L}_1(\varphi') - I_{1,1}(\varphi) + I_{1,0}(\varphi) \right) + u'(y_c)^2 \mathcal{L}_2(\varphi), \end{aligned}$$

which gives

$$\begin{aligned} [\partial_{y_c}^2, \mathcal{L}_0](\varphi) &= -I_{0,1}(\varphi') + I_{0,0}(\varphi') + u'(y_c) \big(-I_{1,1}(\varphi) + I_{1,0}(\varphi) \big) \\ &- u'(y_c) \partial_c I_{0,1}(\varphi) + u'(y_c) \partial_c I_{0,0}(\varphi) \\ &+ u''(y_c) \mathcal{L}_1(\varphi) + 2u'(y_c) \mathcal{L}_1(\varphi') + u'(y_c)^2 \mathcal{L}_2(\varphi). \end{aligned}$$

By (7.3), we have for j = 0, 1

$$\begin{aligned} |I_{0,j}(\varphi)(c)| &\leq C \left| \int_{y_c}^{j\pi} \varphi(y) dy \right| \frac{\min\{\alpha^2 | j\pi - y_c|^2, 1\}}{(\cos j\pi - \cos y_c)^2 u'(y_c)} \\ &\leq C \frac{|j\pi - y_c|^{\frac{1}{2}} \min\{\alpha^2 | j\pi - y_c|^2, 1\}}{(\cos j\pi - \cos y_c)^2 u'(y_c)} \|\varphi\|_{L^2} \\ &\leq C \frac{\min\{\alpha^2 | j\pi - y_c|^2, 1\}}{|j\pi - y_c|^{7/2} u'(y_c)} \|\varphi\|_{L^2}, \end{aligned}$$

which gives

$$|\sin^3 y_c I_{0,j}(\varphi')| \le C\alpha^{\frac{1}{2}} \|\varphi'\|_{L^2}.$$

We get by (7.7) that

$$|\sin^4 y_c I_{1,j}(\varphi)| \le C\alpha \|\varphi\|_{L^{\infty}} \le C\alpha^{\frac{1}{2}} (\|\varphi'\|_{L^2} + \alpha \|\varphi\|_{L^2}).$$

Using (7.13) with $\gamma = 1$, we get

$$|\sin^4 y_c \partial_c I_{0,j}(\varphi)| \le C\alpha \|\varphi\|_{L^{\infty}} \le C\alpha^{\frac{1}{2}} (\|\varphi'\|_{L^2} + \alpha \|\varphi\|_{L^2}).$$

By (7.4), we get

$$|\sin^3 y_c \mathcal{L}_1(\varphi)| + |\sin^5 y_c \mathcal{L}_2(\varphi)| \le C\alpha \|\varphi\|_{L^{\infty}} \le C\alpha^{\frac{1}{2}} (\|\varphi'\|_{L^2} + \alpha \|\varphi\|_{L^2}),$$

and by Lemma 7.6, we get

$$|u'(y_c)^4 \mathcal{L}_1(\varphi')| \le C\alpha^{\frac{1}{2}} \|\varphi'\|_{L^2}.$$

Summing up, we conclude the lemma.

Lemma 10.6. For $\omega \in H^2[0,\pi]$, $\omega'(0) = \omega'(\pi) = 0$, we have

$$\sin y_c[\partial_{y_c}^2, \Lambda_{3,1}](\omega) = C(\alpha^{-2} + \rho)(\|\omega\|_{H^1} + \alpha\|\omega\|_{L^2})(\alpha^{\frac{1}{2}}\mathcal{L}^\infty + \mathcal{L}^2).$$

Proof. We have

$$-\mathbf{E}_{j}(\partial_{y}^{2}\omega) = \omega'(y_{c}) + \int_{y_{c}}^{j\pi} \partial_{y}\phi_{1}(y,c)\partial_{y}\omega dy.$$

Then by (8.8) and (8.11), we get $\partial^2 \Lambda_{0,1}(\omega) = \Lambda_{0,1}(\partial^2 \omega)$

$$\begin{split} \partial_{y_c}^2 \Lambda_{3,1}(\omega) &- \Lambda_{3,1}(\partial_y^2 \omega) \\ &= \partial_{y_c}^2 \left(\frac{J_j}{u'}\right) (u'' \mathcal{E}_{1-j}(\omega) + u'\omega) + 2\partial_{y_c} \left(\frac{J_j}{u'}\right) \partial_{y_c} (u'' \mathcal{E}_{1-j}(\omega) + u'\omega) \\ &+ \frac{J_j}{u'} \left(\partial_{y_c}^2 (u'' \mathcal{E}_{1-j}(\omega) + u'\omega) - (u'' \mathcal{E}_{1-j}(\partial_y^2 \omega) + u' \partial_y^2 \omega)\right) \Big|_{j=0}^1 \\ &= \partial_{y_c}^2 \left(\frac{J_j}{u'}\right) (u'' \mathcal{E}_{1-j}(\omega) + u'\omega) + 2\partial_{y_c} \left(\frac{J_j}{u'}\right) \left(u''' \mathcal{E}_{1-j}(\omega) + u'\omega' + u''u' \int_{y_c}^{(1-j)\pi} \partial_c \phi_1(y,c) \omega(y) dy\right) \\ &+ \frac{J_j}{u'} \left(u \mathcal{E}_{1-j}(\omega) + 2u'' \partial_y \omega + u'' \int_{y_c}^{(1-j)\pi} \partial_y \phi_1(y,c) \partial_y \omega dy - u'''\omega \\ &+ (u''^2 - 2u'^2) \int_{y_c}^{(1-j)\pi} \partial_c \phi_1(y,c) \omega(y) dy + u'^2 u'' \int_{y_c}^{(1-j)\pi} \partial_c^2 \phi_1(y,c) \omega(y) dy \right) \Big|_{j=0}^1 \\ &= T_1 + T_2 + T_3. \end{split}$$

By (8.2)-(8.5) and Lemma 4.8, we get

$$\sin y_c T_1 = \alpha^{-2} (\|\omega\|_{H^1} + \alpha \|\omega\|_{L^2}) (\alpha^{\frac{1}{2}} \mathcal{L}^{\infty} + \mathcal{L}^2).$$

Using $u'''(y) = -\sin y$, (8.1), (8.9), (8.10) and Lemma 4.8, we deduce that

$$|\sin y_c T_2| \le C\alpha^{-2} (\|\omega\|_{H^1} + \alpha\|\omega\|_{L^2}) (\alpha^{\frac{1}{2}} \mathcal{L}^{\infty} + \mathcal{L}^2),$$

By (8.1) and Lemma 4.8, we get

$$\begin{aligned} \left| J_j \mathcal{E}_{1-j}(\omega) \right| + \left| J_j \omega \right| &\leq C \alpha^{-2} \| \omega \|_{L^{\infty}} \leq C \alpha^{-2} (\| \omega \|_{H^1} + \alpha \| \omega \|_{L^2}), \\ \left\| J_j \partial_y \omega \right\|_{L^2} &\leq C \alpha^{-2} \| \omega \|_{H^1}, \end{aligned}$$

and

$$\begin{aligned} \left| J_j \int_{y_c}^{(1-j)\pi} \partial_y \phi_1(y,c) \partial_y \omega dy \right| &\leq C |J_j| \int_{y_c}^{(1-j)\pi} \alpha \alpha^{\frac{1}{2}} |y-y_c|^{\frac{1}{2}} |\partial_y \omega| dy \phi_1((1-j)\pi,c) \\ &\leq C \frac{\sin y_c}{\alpha^{\frac{1}{2}}} \|\partial_y \omega\|_{L^2}. \end{aligned}$$

By (8.9), (8.10) and Lemma 4.8, we get

$$\left| J_j \int_{y_c}^{(1-j)\pi} \partial_c \phi_1(y,c) \omega(y) dy \right| \le C |J_j| \alpha ||\omega||_{L^{\infty}} \frac{\phi_1((1-j)\pi,c)}{|j\pi - y_c|} \le C \alpha^{-1} ||\omega||_{L^{\infty}}.$$

By (8.12), (8.13) and Lemma 4.8, we get

$$\left| J_{j} u'^{2} \int_{y_{c}}^{(1-j)\pi} \partial_{c}^{2} \phi_{1}(y,c) \omega(y) dy \right| \leq C |J_{j}| \alpha^{2} ||\omega||_{L^{\infty}} \phi_{1}((1-j)\pi,c) |(1-j)\pi - y_{c}|$$
$$\leq C \sin y_{c} ||\omega||_{L^{\infty}}.$$

Notice that

$$\|\omega\|_{L^{\infty}} \le C\alpha^{-\frac{1}{2}} (\|\omega\|_{H^{1}} + \alpha\|\omega\|_{L^{2}}), \qquad \sin y_{c} \le \alpha^{-1} + \alpha\rho.$$

Then we conclude that

$$\sin y_c T_3 = (\alpha^{-2} + \rho)(\|\omega\|_{H^1} + \alpha\|\omega\|_{L^2})(\alpha^{\frac{1}{2}}\mathcal{L}^{\infty} + \mathcal{L}^2).$$

This proves the lemma.

Now we are in a position to prove the proposition.

Proof. Step 1. the odd case. Let $\omega \in H^2$ be odd and $\omega(0) = \omega(\pi) = 0$. Direct calculation gives

$$\begin{aligned} \partial_{y_c}^2 \mathbb{D}(\omega) &- \mathbb{D}(\partial_y^2 \omega) \\ &= \rho \mathrm{II}_{1,1}(\omega) \partial_{y_c}^2 \Big(\frac{u''}{(\mathrm{A}^2 + \mathrm{B}^2)^{\frac{1}{2}}} \Big) + 2\partial_{y_c} \Big(\rho \mathrm{II}_{1,1}(\omega) \Big) \partial_{y_c} \Big(\frac{u''}{(\mathrm{A}^2 + \mathrm{B}^2)^{\frac{1}{2}}} \Big) + \frac{u'' [\partial_{y_c}^2, \rho \mathrm{II}_{1,1}] \omega}{(\mathrm{A}^2 + \mathrm{B}^2)^{\frac{1}{2}}} \\ &+ \mathcal{L}_0(\omega) \partial_{y_c}^2 \Big(\frac{\rho u''}{(\mathrm{A}^2 + \mathrm{B}^2)^{\frac{1}{2}}} \Big) + 2\partial_{y_c} \mathcal{L}_0(\omega) \partial_{y_c} \Big(\frac{\rho u''}{(\mathrm{A}^2 + \mathrm{B}^2)^{\frac{1}{2}}} \Big) + \frac{\rho u'' [\partial_{y_c}^2, \mathcal{L}_0] \omega}{(\mathrm{A}^2 + \mathrm{B}^2)^{\frac{1}{2}}} \\ &+ \omega \partial_{y_c}^2 \Big(\frac{\rho u' \mathrm{II}}{(\mathrm{A}^2 + \mathrm{B}^2)^{\frac{1}{2}}} \Big) + 2\partial_{y_c} \omega \partial_{y_c} \Big(\frac{\rho u' \mathrm{II}}{(\mathrm{A}^2 + \mathrm{B}^2)^{\frac{1}{2}}} \Big) \\ &= I_1 + I_2 + I_3 + I_4 + I_5 + I_6 + I_7 + I_8. \end{aligned}$$

By Lemma 7.5 and Proposition 4.3, we get

$$\|\sin y_c I_1\|_{L^2} \le C \|\sin y_c \operatorname{II}_{1,1}\|_{L^2} \left\| \rho \partial_{y_c}^2 \left(\frac{u''}{(\mathbf{A}^2 + \mathbf{B}^2)^{\frac{1}{2}}} \right) \right\|_{L^{\infty}} \le C \|\omega\|_{H^1},$$

and by Lemma 7.4 and Proposition 4.3,

$$\|\sin y_c I_2\|_{L^2} \le C \|\rho \Pi_{1,1}\|_{H^1} \|\sin y_c \partial_{y_c} \left(\frac{u''}{(A^2 + B^2)^{\frac{1}{2}}}\right) \|_{L^{\infty}} \le C \|\omega\|_{H^1},$$

and by Lemma 10.4 and Proposition 4.3,

$$\|\sin y_c I_3\|_{L^2} \le C \|\sin y_c[\partial_y^2, \rho II_{1,1}]\omega\|_{L^2} \le C \|\omega\|_{H^1},$$

and by Lemma 7.7 and Proposition 4.3,

$$\begin{aligned} \|\sin y_{c}I_{4}\|_{L^{2}} &\leq C\alpha^{-\frac{1}{2}} \|\sin y_{c}\mathcal{L}_{0}(\omega)\|_{L^{\infty}} \left\|\alpha^{\frac{1}{2}}\partial_{y_{c}}^{2}\left(\frac{\rho u''}{(\mathbf{A}^{2}+\mathbf{B}^{2})^{\frac{1}{2}}}\right)\right\|_{L^{2}} \\ &\leq C\|\omega\|_{H^{1}} \left\|\frac{\alpha^{\frac{1}{2}}}{(1+\alpha\sin y_{c})}\right\|_{L^{2}} \leq C\|\omega\|_{H^{1}}, \\ \|\sin y_{c}I_{5}\|_{L^{2}} &\leq C\alpha^{-\frac{1}{2}} \|\sin^{2} y_{c}\partial_{y_{c}}\mathcal{L}_{0}(\omega)\|_{L^{\infty}} \left\|\frac{\alpha^{\frac{1}{2}}}{\sin y_{c}}\partial_{y_{c}}\left(\frac{\rho u''}{(\mathbf{A}^{2}+\mathbf{B}^{2})^{\frac{1}{2}}}\right)\right\|_{L^{2}} \\ &\leq C\|\omega\|_{H^{1}} \left\|\frac{\alpha^{\frac{1}{2}}}{(1+\alpha\sin y_{c})}\right\|_{L^{2}} \leq C\|\omega\|_{H^{1}}. \end{aligned}$$

By Lemma 10.5, we get

$$\|\sin y_c I_6\|_{L^2} \le C\alpha^{-\frac{1}{2}} \left\| \frac{\alpha^{\frac{1}{2}}}{(1+\alpha\sin y_c)} \right\|_{L^2} \|\sin^3 y_c[\partial_{y_c}^2, \mathcal{L}_0]\omega\|_{L^\infty} \le C(\|\omega\|_{H^1} + \alpha\|\omega\|_{L^2})$$

By Lemma 4.4 and Proposition 4.3, we get

$$\|\sin y_{c}I_{7}\|_{L^{2}} \leq C \|\omega/u'\|_{L^{2}} \left\|\rho\partial_{y_{c}}^{2}\left(\frac{\rho u'\mathrm{II}}{(\mathrm{A}^{2}+\mathrm{B}^{2})^{\frac{1}{2}}}\right)\right\|_{L^{\infty}} \leq C \|\omega\|_{H^{1}},$$

$$\|\sin y_{c}I_{8}\|_{L^{2}} \leq C \|\partial_{y}\omega\|_{L^{2}} \left\|\sin y_{c}\partial_{y_{c}}\left(\frac{\rho u'\mathrm{II}}{(\mathrm{A}^{2}+\mathrm{B}^{2})^{\frac{1}{2}}}\right)\right\|_{L^{\infty}} \leq C \|\partial_{y}\omega\|_{L^{2}}.$$

Summing up, we conclude the odd case.

Step 2. the even case. Let $\omega \in H^2$ be even, $\omega'(0) = \omega'(\pi) = 0$. Direct calculation gives $\partial^2_{u_c} \mathbb{D}(\omega) - \mathbb{D}(\partial^2_u \omega)$

$$\begin{split} &= \frac{\rho u''}{(\mathbf{A}_{1}^{2} + \mathbf{B}_{1}^{2})^{\frac{1}{2}}} [\partial_{y_{c}}^{2}, \rho \mathbf{I} \mathbf{I}_{1,1}] \omega + 2\partial_{y_{c}} \Big(\frac{\rho u''}{(\mathbf{A}_{1}^{2} + \mathbf{B}_{1}^{2})^{\frac{1}{2}}} \Big) \partial_{y_{c}} \big(\rho \mathbf{I} \mathbf{I}_{1,1}(\omega) \big) + \partial_{y_{c}}^{2} \Big(\frac{\rho u''}{(\mathbf{A}_{1}^{2} + \mathbf{B}_{1}^{2})^{\frac{1}{2}}} \Big) \rho \mathbf{I} \mathbf{I}_{1,1}(\omega) \\ &+ \frac{\rho^{2} u''}{(\mathbf{A}_{1}^{2} + \mathbf{B}_{1}^{2})^{\frac{1}{2}}} [\partial_{y_{c}}^{2}, \mathcal{L}_{0}] \omega + 2\partial_{y_{c}} \Big(\frac{\rho^{2} u''}{(\mathbf{A}_{1}^{2} + \mathbf{B}_{1}^{2})^{\frac{1}{2}}} \Big) \partial_{y_{c}} \big(\mathcal{L}_{0}(\omega) \big) + \partial_{y_{c}}^{2} \Big(\frac{\rho^{2} u''}{(\mathbf{A}_{1}^{2} + \mathbf{B}_{1}^{2})^{\frac{1}{2}}} \Big) \mathcal{L}_{0}(\omega) \\ &+ \omega \partial_{y_{c}}^{2} \Big(\frac{\rho^{2} u' \mathbf{I} \mathbf{I}}{(\mathbf{A}_{1}^{2} + \mathbf{B}_{1}^{2})^{\frac{1}{2}}} \Big) + 2\partial_{y_{c}} \Big(\frac{\rho^{2} u' \mathbf{I} \mathbf{I}}{(\mathbf{A}_{1}^{2} + \mathbf{B}_{1}^{2})^{\frac{1}{2}}} \Big) \partial_{y_{c}} \omega \\ &+ \partial_{y_{c}}^{2} \Big(\frac{1}{(\mathbf{A}_{1}^{2} + \mathbf{B}_{1}^{2})^{\frac{1}{2}}} \Big) \Lambda_{3,1}(\omega) + \partial_{y_{c}} \Big(\frac{1}{(\mathbf{A}_{1}^{2} + \mathbf{B}_{1}^{2})^{\frac{1}{2}}} \Big) \partial_{y_{c}} \Lambda_{3,1}(\omega) + \frac{[\partial_{y_{c}}^{2}, \Lambda_{3,1}] \omega}{(\mathbf{A}_{1}^{2} + \mathbf{B}_{1}^{2})^{\frac{1}{2}}} \\ &= I_{1}' + I_{2}' + I_{3}' + I_{4}' + I_{5}' + I_{6}' + I_{7}' + I_{8}' + I_{9}' + I_{10}' + I_{11}'. \end{split}$$

By Proposition 4.5 and Lemma 10.4, we get

$$\|\sin y_c I_1'\|_{L^2} \le C \left\| \frac{\rho u''}{(A_1^2 + B_1^2)^{\frac{1}{2}}} \right\|_{L^{\infty}} \|\sin y_c [\partial_{y_c}^2, \rho II_{1,1}] \omega \|_{L^2} \le C(\alpha \|\omega\|_{L^2} + \|\partial_y \omega\|_{L^2}),$$

and by Proposition 4.5 and Lemma 7.4,

$$\begin{aligned} \|\sin y_c I_2'\|_{L^2} &\leq C \left\|\sin y_c \partial_{y_c} \left(\frac{\rho u''}{(\mathbf{A}_1^2 + \mathbf{B}_1^2)^{\frac{1}{2}}}\right)\right\|_{L^{\infty}} \left\|\partial_{y_c} \left(\rho \mathbf{II}_{1,1}(\omega)\right)\right\|_{L^2} &\leq C \|\omega\|_{H^1}, \\ \|\sin y_c I_3'\|_{L^2} &\leq C \left\|\sin y_c \partial_{y_c}^2 \left(\frac{\rho u''}{(\mathbf{A}_1^2 + \mathbf{B}_1^2)^{\frac{1}{2}}}\right)\right\|_{L^{\infty}} \left\|\rho \mathbf{II}_{1,1}(\omega)\right\|_{L^2} &\leq C \alpha \|\omega\|_{L^2}. \end{aligned}$$

By Proposition 4.5 and Lemma 10.5, we get

$$\|\sin y_c I_4'\|_{L^2} \le C \left\| \frac{\rho u''}{(A_1^2 + B_1^2)^{\frac{1}{2}}} \right\|_{L^2} \|\sin^3 y_c[\partial_{y_c}^2, \mathcal{L}_0]\omega\|_{L^\infty} \le C(\alpha \|\omega\|_{L^2} + \|\partial_y \omega\|_{L^2}),$$

and by Proposition 4.5 and Lemma 7.7,

$$\begin{aligned} \|\sin y_{c}I_{5}'\|_{L^{2}} &\leq C \left\| \frac{1}{\sin y_{c}} \partial_{y_{c}} \left(\frac{\rho^{2}u''}{(\mathbf{A}_{1}^{2} + \mathbf{B}_{1}^{2})^{\frac{1}{2}}} \right) \right\|_{L^{2}} \|\sin^{2} y_{c} \partial_{y_{c}} \left(\mathcal{L}_{0}(\omega) \right) \|_{L^{\infty}} \\ &\leq C \left(\alpha^{\frac{1}{2}} \|\omega\|_{L^{\infty}} + \|\omega'\|_{L^{2}} \right) \leq C \left(\alpha \|\omega\|_{L^{2}} + \|\omega'\|_{L^{2}} \right), \end{aligned}$$

and by Proposition 4.5 and Lemma 7.6,

$$\|\sin y_c I_6'\|_{L^2} \le C \left\| \frac{1}{\sin y_c} \partial_{y_c}^2 \left(\frac{\rho^2 u''}{(A_1^2 + B_1^2)^{\frac{1}{2}}} \right) \right\|_{L^2} \left\| \rho \mathcal{L}_0(\omega) \right\|_{L^\infty}$$

$$\leq C\alpha \|\omega\|_{L^2}.$$

By Proposition 4.5 and Lemma 4.4, we get

$$\begin{aligned} \|\sin y_c I_7'\|_{L^2} &\leq C \|\omega\|_{L^2} \left\|\sin y_c \partial_{y_c}^2 \left(\frac{\rho^2 u' \mathrm{II}}{(\mathrm{A}_1^2 + \mathrm{B}_1^2)^{\frac{1}{2}}}\right)\right\|_{L^{\infty}} \leq C \alpha \|\omega\|_{L^2}, \\ \|\sin y_c I_8'\|_{L^2} &\leq C \|\partial_y \omega\|_{L^2} \left\|\sin y_c \partial_{y_c} \left(\frac{\rho^2 u' \mathrm{II}}{(\mathrm{A}_1^2 + \mathrm{B}_1^2)^{\frac{1}{2}}}\right)\right\|_{L^{\infty}} \leq C \|\partial_y \omega\|_{L^2}. \end{aligned}$$

By Proposition 4.5 and Remark 8.2, we get

$$\sin y_c I'_9 = \frac{(\mathcal{L}^2 + \alpha^{\frac{1}{2}} \mathcal{L}^\infty)}{(1 + \alpha \sin y_c)^3} (\alpha \|\omega\|_{L^2} + \|\partial_y \omega\|_{L^2}) = \mathcal{L}^2,$$

$$\sin y_c I'_{10} = \frac{(\mathcal{L}^2 + \alpha^{\frac{1}{2}} \mathcal{L}^\infty)}{(1 + \alpha \sin y_c)^3} (\alpha \|\omega\|_{L^2} + \|\partial_y \omega\|_{L^2}) = \mathcal{L}^2.$$

By Proposition 4.5 and Lemma 10.6, we get

$$\sin y_c I'_{11} = \frac{(1+\alpha^2 \rho)(\mathcal{L}^2 + \alpha^{\frac{1}{2}} \mathcal{L}^\infty)}{(1+\alpha \sin y_c)^3} (\alpha \|\omega\|_{L^2} + \|\partial_y \omega\|_{L^2}) = \mathcal{L}^2.$$

Summing up, we conclude the even case.

11. LINEAR ENHANCED DISSIPATION

11.1. Decay estimates on the model without nonlocal term. In this subsection, we consider a toy model without nonlocal term:

(11.1)
$$\partial_t \omega - \nu \partial_y^2 \omega - iae^{-\nu t} (\cos y)\omega = 0.$$

Lemma 11.1. Let $0 < \nu < |a|$, $a \in \mathbf{R}$, and $\omega(t, y)$ solve (11.1) for $t \ge 0$, $y \in \mathbb{T}$. Then $\|\omega(t)\|_{L^2} \le \|\omega(0)\|_{L^2}$ and

$$\|\sin y\omega(t)\|_{L^2} \le C(\nu t^3 a^2)^{-\frac{1}{2}} \|\omega(0)\|_{L^2}$$

for $0 < t \le (\nu |a|)^{-\frac{1}{2}}$.

Proof. First of all, we have

$$\partial_t \|\omega\|_{L^2}^2 = 2\operatorname{Re}\langle\partial_t\omega,\omega\rangle = 2\operatorname{Re}\langle\nu\partial_y^2\omega + iae^{-\nu t}\cos y\omega,\omega\rangle = -2\nu\|\partial_y\omega\|_{L^2}^2 \le 0,$$

which gives $\|\omega(t)\|_{L^2} \le \|\omega(0)\|_{L^2}$.

Let
$$u(y) = -\cos y$$
, $\gamma(t,s) = a \frac{e^{-\nu s} - e^{-\nu t}}{\nu}$. Then $\gamma(s,s) = 0$, $\partial_t \gamma = a e^{-\nu t}$, and
 $\partial_t (e^{i\gamma u}\omega) = e^{i\gamma u} (\omega_t + iu\gamma_t\omega) = \nu e^{i\gamma u} \partial_u^2 \omega$,

from which, we infer that

$$\begin{aligned} \partial_t \|\partial_y (e^{i\gamma u}\omega)\|_{L^2}^2 &= 2\operatorname{Re}\langle \partial_t \partial_y (e^{i\gamma u}\omega), \partial_y (e^{i\gamma u}\omega)\rangle \\ &= -2\operatorname{Re}\langle \partial_t (e^{i\gamma u}\omega), \partial_y^2 (e^{i\gamma u}\omega)\rangle \\ &= -2\operatorname{Re}\langle \nu e^{i\gamma u} \partial_y^2 \omega, \partial_y^2 (e^{i\gamma u}\omega)\rangle \\ &= -2\nu\operatorname{Re}\langle \partial_y^2 \omega, e^{-i\gamma u} \partial_y^2 (e^{i\gamma u}\omega)\rangle \\ &= -2\nu\operatorname{Re}\langle \partial_y^2 \omega, \partial_y^2 \omega + 2i\gamma u' \partial_y \omega + (i\gamma u'' - \gamma^2 u'^2)\omega\rangle \\ &= -2\nu\operatorname{Re}\langle \partial_y^2 \omega + i\gamma u' \partial_y \omega + i\gamma u'' \omega/2\|_{L^2}^2 + 2\nu\gamma^2 \|u' \partial_y \omega + u'' \omega/2\|_{L^2}^2 \end{aligned}$$

$$+ 2\nu\gamma^2 \operatorname{Re}\langle \partial_y^2 \omega, u'^2 \omega \rangle.$$

For the last term, we get by integration by parts that

$$\operatorname{Re}\langle\partial_{y}^{2}\omega, u'^{2}\omega\rangle = -\operatorname{Re}\langle\partial_{y}\omega, \partial_{y}(u'^{2}\omega)\rangle$$

$$= -\operatorname{Re}\langle\partial_{y}\omega, u'^{2}\partial_{y}\omega + 2u'u''\omega\rangle$$

$$= -\operatorname{Re}\langle u'\partial_{y}\omega, u'\partial_{y}\omega + 2u''\omega\rangle$$

$$= -\|u'\partial_{y}\omega + u''\omega/2\|_{L^{2}}^{2} + \|u''\omega/2\|_{L^{2}}^{2} - \operatorname{Re}\langle u'\partial_{y}\omega, u''\omega\rangle,$$

where

$$-2\operatorname{Re}\langle u'\partial_y\omega, u''\omega\rangle = -\int_{-\pi}^{\pi} u'u''(\partial_y\omega\overline{\omega} + \partial_y\overline{\omega}\omega)dy$$
$$= -\int_{-\pi}^{\pi} u'u''\partial_y|\omega|^2dy = \int_{-\pi}^{\pi} (u'u'')'|\omega|^2dy.$$

Putting these identities above together and using the fact that

$$(u'')^2/2 + (u'u'')' = 3(u'')^2/2 + u'u''' = 3(\cos y)^2/2 - (\sin y)^2 \le 3/2,$$

we deduce that

$$\begin{aligned} \partial_t \|\partial_y (e^{i\gamma u}\omega)\|_{L^2}^2 &\leq 2\nu\gamma^2 \left(\|u''\omega/2\|_{L^2}^2 - \operatorname{Re}\langle u'\partial_y\omega, u''\omega\rangle\right) \\ &= \nu\gamma^2 \int_{-\pi}^{\pi} \left((u'')^2/2 + (u'u'')'\right)|\omega|^2 dy \leq \frac{3}{2}\nu\gamma^2 \|\omega\|_{L^2}^2. \end{aligned}$$

Therefore, for 0 < s < t,

(11.2)
$$\|\partial_y(e^{i\gamma u}\omega)(t,s)\|_{L^2}^2 \le \|\partial_y(e^{i\gamma u}\omega)(s,s)\|_{L^2}^2 + \frac{3}{2}\nu \int_s^t \gamma(\tau,s)^2 \|\omega(\tau)\|_{L^2}^2 d\tau \\ \le \|\partial_y\omega(s)\|_{L^2}^2 + \frac{3}{2}\nu\|\omega(0)\|_{L^2}^2 \int_s^t \gamma(\tau,s)^2 d\tau.$$

Thanks to $\partial_y(e^{i\gamma u}\omega) = e^{i\gamma u}(i\gamma u'\omega + \partial_y\omega), \|\partial_y(e^{i\gamma u}\omega)\|_{L^2}^2 = \|i\gamma u'\omega + \partial_y\omega\|_{L^2}^2$. Then by (11.2), we get

$$\int_{0}^{t} \|i\gamma(t,s)u'\omega(t) + \partial_{y}\omega(t)\|_{L^{2}}^{2} ds = \int_{0}^{t} \|\partial_{y}(e^{i\gamma u}\omega)(t,s)\|_{L^{2}}^{2} ds$$
$$\leq \int_{0}^{t} \|\partial_{y}\omega(s)\|_{L^{2}}^{2} ds + \frac{3}{2}\nu\|\omega(0)\|_{L^{2}}^{2} \int_{0}^{t} \int_{s}^{t} \gamma(\tau,s)^{2} d\tau ds.$$

On the other hand, let

$$\gamma_1(t) = \int_0^t \gamma(t,s) ds, \quad \gamma_2(t) = \int_0^t \gamma(t,s)^2 ds, \quad \gamma_0(t) = \gamma_2(t) - \gamma_1(t)^2 / t.$$

We find that

$$\begin{split} &\int_0^t \|i\gamma(t,s)u'\omega(t) + \partial_y\omega(t)\|_{L^2}^2 ds \\ &= \gamma_2(t)\|u'\omega(t)\|_{L^2}^2 + 2\gamma_1(t)\operatorname{Re}\langle iu'\omega(t),\partial_y\omega(t)\rangle + t\|\partial_y\omega(t)\|_{L^2}^2 \\ &= \gamma_0(t)\|u'\omega(t)\|_{L^2}^2 + t\|i\gamma_1(t)u'\omega(t)/t + \partial_y\omega(t)\|_{L^2}^2. \end{split}$$

Moreover,

$$t\gamma_0(t) = t\gamma_2(t) - \gamma_1(t)^2$$

$$\begin{split} &= \int_0^t \int_0^t \gamma(t,s)^2 ds d\tau - \int_0^t \int_0^t \gamma(t,s)\gamma(t,\tau) ds d\tau \\ &= \int_0^t \int_0^t (\gamma(t,s)^2/2 + \gamma(t,\tau)^2/2 - \gamma(t,s)\gamma(t,\tau)) ds d\tau \\ &= \frac{1}{2} \int_0^t \int_0^t (\gamma(t,s) - \gamma(t,\tau))^2 ds d\tau \\ &= \frac{1}{2} \int_0^t \int_0^t \gamma(\tau,s)^2 ds d\tau = \int_0^t \int_s^t \gamma(\tau,s)^2 d\tau ds, \end{split}$$

and

$$\int_0^t \|\partial_y \omega(s)\|_{L^2}^2 ds = \frac{1}{2\nu} \big(\|\omega(0)\|_{L^2}^2 - \|\omega(t)\|_{L^2}^2\big).$$

Therefore, we obtain

$$\gamma_0(t) \| u'\omega(t) \|_{L^2}^2 \le \frac{1}{2\nu} (\| \omega(0) \|_{L^2}^2 - \| \omega(t) \|_{L^2}^2) + \frac{3}{2}\nu t\gamma_0(t) \| \omega(0) \|_{L^2}^2,$$

which gives

$$\|u'\omega(t)\|_{L^2}^2 \le (1/(2\nu\gamma_0(t)) + 3\nu t/2) \|\omega(0)\|_{L^2}^2.$$

Now, we give a lower bound of $\gamma_0(t)$. As $\gamma(\tau, s) = \int_s^{\tau} a e^{-\nu t'} dt' \ge a(\tau - s) e^{-\nu t}$ for $0 < s < \tau < t$, we have

$$t\gamma_0(t) = \int_0^t \int_s^t \gamma(\tau, s)^2 d\tau ds \ge \int_0^t \int_s^t a^2(\tau - s)^2 e^{-2\nu t} d\tau ds = a^2 e^{-2\nu t} \frac{t^4}{12},$$

which gives

$$\frac{1}{2\nu\gamma_0(t)} \le \frac{1}{2\nu a^2} e^{2\nu t} \frac{12}{t^3} = \frac{6e^{2\nu t}}{\nu a^2 t^3}$$

For $0 < t \le (\nu |a|)^{-\frac{1}{2}}$, $0 < \nu < |a|$, we have $\nu t \le 1$, $\nu t \le 1/(\nu a^2 t^3)$. Then we conclude that $\|u'\omega(t)\|_{L^2}^2 \le (6e^2 + 2)\|\omega(0)\|_{L^2}^2/(\nu a^2 t^3)$.

This completes the proof.

Now, we deal with the inhomogeneous equation.

Lemma 11.2. Let $0 < \nu < |a|, a \in \mathbf{R}, and \omega(t, y)$ solve (11.3) $\partial_t \omega - \nu \partial_y^2 \omega - iae^{-\nu t} (\cos y)\omega = f.$

Then we have

$$\|\sin y\omega(t)\|_{L^2} \le \frac{C}{\sqrt{\nu t^3 a^2}} \sup_{0 \le s \le t} \|\omega(s)\|_{L^2} + \int_0^t \|\sin yf(s)\|_{L^2} ds$$

for $0 < t \le (\nu |a|)^{-\frac{1}{2}}$.

Proof. We decompose $\omega = \omega_1 + \omega_2$, where

$$\partial_t \omega_1 - \nu \partial_y^2 \omega_1 - iae^{-\nu t} (\cos y) \omega_1 = 0, \quad \omega_1(0) = \omega(0),$$

$$\partial_t \omega_2 - \nu \partial_y^2 \omega_2 - iae^{-\nu t} (\cos y) \omega_2 = f, \quad \omega_2(0) = 0.$$

 $u_t \omega_2 - \nu u_y \omega_2 - iae^{-i\pi} (\cos y) \omega_2 = f, \quad \omega_2(0) = 0$ By Lemma 11.1, we have $\|\omega_1(t)\|_{L^2} \le \|\omega_1(0)\|_{L^2} = \|\omega(0)\|_{L^2}$, and

$$\|\sin y\omega_1(t)\|_{L^2} \le C(\nu t^3 a^2)^{-\frac{1}{2}} \|\omega_1(0)\|_{L^2} = C(\nu t^3 a^2)^{-\frac{1}{2}} \|\omega(0)\|_{L^2}$$

for
$$0 < t \le (\nu |a|)^{-\frac{1}{2}}$$
. Let $A(t) = \sup_{0 \le s \le t} \|\omega(s)\|_{L^2}$. Then
 $\|\omega_2(t)\|_{L^2} \le \|\omega_1(t)\|_{L^2} + \|\omega(t)\|_{L^2} \le 2A(t).$

For any b > 0, we have

$$\begin{aligned} \frac{d}{dt} \|\sin y\omega_2\|_{L^2}^2 &= 2\operatorname{Re}\langle\sin y\partial_t\omega_2,\sin y\omega_2\rangle \\ &= 2\operatorname{Re}\langle\sin y(\nu\partial_y^2\omega_2 + iae^{-\nu t}\cos y\omega_2 + f),\sin y \,\omega_2\rangle \\ &= -2\nu\|\sin y \,\partial_y\omega_2 + \cos y \,\omega_2\|_{L^2}^2 + 2\nu\|\cos y \,\omega_2\|_{L^2}^2 + 2\operatorname{Re}\langle\sin yf,\sin y \,\omega_2\rangle \\ &\leq 2\nu\|\omega_2\|_{L^2}^2 + 2\|\sin yf\|_{L^2}\|\sin y\omega_2\|_{L^2} \\ &\leq 2\left(\|\sin yf\|_{L^2} + 4\nu A(t)^2/b\right)\left(\|\sin y\omega_2\|_{L^2}^2 + b^2\right)^{\frac{1}{2}}, \end{aligned}$$

therefore,

7

$$\frac{d}{dt} \left(\|\sin y\omega_2\|_{L^2}^2 + b^2 \right)^{\frac{1}{2}} \le \|\sin yf\|_{L^2} + 4\nu A(t)^2/b,$$

which implies that

$$\begin{aligned} |\sin y\omega_2(t)||_{L^2} &\leq (\|\sin y\omega_2(t)\|_{L^2}^2 + b^2)^{\frac{1}{2}} \\ &\leq b + \int_0^t (\|\sin yf(s)\|_{L^2} + 4\nu A(s)^2/b) ds \\ &\leq b + 4\nu t A(t)^2/b + \int_0^t \|\sin yf(s)\|_{L^2} ds. \end{aligned}$$

As it is true for any b > 0, we minimize the right hand side to obtain

$$\|\sin y\omega_2(t)\|_{L^2} \le 4\sqrt{\nu t}A(t) + \int_0^t \|\sin yf(s)\|_{L^2} ds$$

For $0 < t \le (\nu |a|)^{-\frac{1}{2}}$, we have $\sqrt{\nu t} \le (\nu t^3 a^2)^{-\frac{1}{2}}$. Then

$$\begin{aligned} \|\sin y\omega(t)\|_{L^{2}} &\leq \|\sin y\omega_{1}(t)\|_{L^{2}} + \|\sin y\omega_{2}(t)\|_{L^{2}} \\ &\leq C(\nu t^{3}a^{2})^{-\frac{1}{2}}\|\omega(0)\|_{L^{2}} + 4\sqrt{\nu t}A(t) + \int_{0}^{t} \|\sin yf(s)\|_{L^{2}}ds \\ &\leq C(\nu t^{3}a^{2})^{-\frac{1}{2}}A(t) + \int_{0}^{t} \|\sin yf(s)\|_{L^{2}}ds. \end{aligned}$$

This gives our result.

11.2. Decay estimates on the model with nonlocal term.

Lemma 11.3. Let $0 < \nu < |a|, |\alpha| > 1, a, \alpha \in \mathbf{R}, and \omega(t, y)$ solve

(11.4)
$$\partial_t \omega - \nu \partial_y^2 \omega - iae^{-\nu t} \cos y(\omega - \psi) = 0, \quad -(\partial_y^2 - \alpha^2)\psi = \omega.$$

Then for $0 < t \le (\nu |a|)^{-\frac{1}{2}}, \ \nu \alpha^2 t < 1,$

$$-\langle \psi(t), \omega(t) - \psi(t) \rangle \le \frac{C}{\nu t^3 a^2} \langle \omega(0), \omega(0) - \psi(0) \rangle$$

Proof. By (11.4), we have

$$\begin{aligned} \partial_t \langle \omega, \omega - \psi \rangle &= \langle \omega_t, \omega - \psi \rangle + \langle \omega, \omega_t \rangle + \langle (\partial_y^2 - \alpha^2) \psi, \psi_t \rangle \\ &= \langle \omega_t, \omega - \psi \rangle + \langle \omega, \omega_t \rangle - \langle \psi, \omega_t \rangle \\ &= -2\nu \langle \partial_y \omega, \partial_y \omega - \partial_y \psi \rangle. \end{aligned}$$

By Proposition 10.2, we have $\|\mathbb{D}(\omega)\|_{L^2}^2 = \langle \omega, \omega - \psi \rangle$. Using the facts that

$$\begin{split} \|\partial_{y}\omega\|_{L^{2}}^{2} &= \|\partial_{y}^{3}\psi\|_{L^{2}}^{2} + \alpha^{4}\|\partial_{y}\psi\|_{L^{2}}^{2} + 2\alpha^{2}\|\partial_{y}^{2}\psi\|_{L^{2}}^{2} \ge \alpha^{4}\|\partial_{y}\psi\|_{L^{2}}^{2} + \alpha^{2}\|\partial_{y}^{2}\psi\|_{L^{2}}^{2}, \\ &-\langle\partial_{y}\omega,\partial_{y}\psi\rangle = -\|\partial_{y}^{2}\psi\|_{L^{2}}^{2} - \alpha^{2}\|\partial_{y}\psi\|_{L^{2}}^{2}, \end{split}$$

we deduce that

(11.5)
$$\langle \partial_y \omega, \partial_y \omega - \partial_y \psi \rangle = \| \partial_y \omega \|_{L^2}^2 - \langle \partial_y \omega, \partial_y \psi \rangle \ge (1 - \alpha^{-2}) \| \partial_y \omega \|_{L^2}^2.$$

Then we obtain $\partial_t \|\mathbb{D}(\omega)(t)\|_{L^2}^2 \leq 0$, thus $\|\mathbb{D}(\omega)(t)\|_{L^2} \leq \|\mathbb{D}(\omega)(0)\|_{L^2}$ and

$$2\nu \int_0^t \|\partial_y \omega(s)\|_{L^2}^2 \le \frac{\alpha^2}{\alpha^2 - 1} \|\mathbb{D}(\omega)(0)\|_{L^2}^2.$$

By Proposition 10.2, $\mathbb{D}(\omega)$ satisfies (11.3) with $f(s, y) = \nu[\mathbb{D}, \partial_y^2]\omega$. Thus by Lemma 11.2, we know that for $0 < t \leq (\nu|a|)^{-\frac{1}{2}}$,

$$\|\sin y\mathbb{D}(\omega)(t)\|_{L^{2}} \le C(\nu t^{3}a^{2})^{-\frac{1}{2}}\|\mathbb{D}(\omega)(0)\|_{L^{2}} + \int_{0}^{t}\nu\|\sin y[\mathbb{D},\partial_{y}^{2}]\omega(s)\|_{L^{2}}ds.$$

Again by Proposition 10.2, we have

$$\begin{split} \int_{0}^{t} \nu \|\sin y \ [\mathbb{D}, \partial_{y}^{2}] \omega(s) \|_{L^{2}} ds &\leq \int_{0}^{t} \nu C(|\alpha| \|\omega(s)\|_{L^{2}} + \|\partial_{y}\omega(s)\|_{L^{2}}) ds \\ &\leq \int_{0}^{t} \nu C |\alpha| \sqrt{\frac{\alpha^{2}}{\alpha^{2} - 1}} \|\mathbb{D}(\omega)(s)\|_{L^{2}} ds + C\nu t^{\frac{1}{2}} \|\partial_{y}\omega\|_{L^{2}_{t}(L^{2})} \\ &\leq C\nu t |\alpha| \sqrt{\frac{\alpha^{2}}{\alpha^{2} - 1}} \|\mathbb{D}(\omega)(0)\|_{L^{2}} + C\nu t^{\frac{1}{2}} \sqrt{\frac{\alpha^{2}/(2\nu)}{\alpha^{2} - 1}} \|\mathbb{D}(\omega)(0)\|_{L^{2}}. \end{split}$$

Therefore, for $0 < t \le (\nu |a|)^{-\frac{1}{2}}$,

$$\|\sin y\mathbb{D}(\omega)(t)\|_{L^{2}} \leq C\left((\nu t^{3}a^{2})^{-\frac{1}{2}} + \nu t|\alpha| + (\nu t)^{\frac{1}{2}}\right)\|\mathbb{D}(\omega)(0)\|_{L^{2}}.$$

Thus, for $0 < t \leq (\nu |a|)^{-\frac{1}{2}}$, $\nu \alpha^2 t < 1$, we have $\nu t |\alpha| \leq (\nu t)^{\frac{1}{2}} \leq (\nu t^3 a^2)^{-\frac{1}{2}}$. As $-\langle \psi, \omega \rangle = -\|\partial_y \psi\|_{L^2}^2 - \alpha^2 \|\psi\|_{L^2}^2$, then by Proposition 10.2,

(11.6)
$$\langle \psi, \omega - \psi \rangle(t) = (\alpha^2 - 1) \|\psi\|_{L^2}^2 + \|\partial_y \psi\|_{L^2}^2$$

$$\leq \|\sin y \ \mathbb{D}(\omega)(t)\|_{L^2}^2$$

$$\leq C \|\mathbb{D}(\omega)(0)\|_{L^2}^2 / (\nu t^3 a^2) = C \langle \omega, \omega - \psi \rangle(0) / (\nu t^3 a^2).$$

The proof is completed.

Let $-(\partial_y^2 - \alpha^2)\widehat{\psi} = \widehat{\omega}$. We introduce

$$K_1(t,\alpha) = \langle \widehat{\omega}, \widehat{\omega} - \psi \rangle (t,\alpha),$$

$$K_2(t,\alpha) = -\langle (\partial_y^2 - \alpha^2) \widehat{\omega}, \widehat{\omega} - \widehat{\psi} \rangle (t,\alpha),$$

$$K_3(t,\alpha) = \langle \widehat{\psi}, \widehat{\omega} - \widehat{\psi} \rangle (t,\alpha).$$

Lemma 11.4. It holds that

$$\begin{split} K_1 &= \|\mathbb{D}(\widehat{\omega})\|_{L^2}^2 \ge \alpha^2 K_3, \\ K_2 &= (\alpha^2 - 1) \|\widehat{\omega}\|_{L^2}^2 + \|\partial_y \widehat{\omega}\|_{L^2}^2 \ge \alpha^2 K_1, \\ K_3 &= (\alpha^2 - 1) \|\widehat{\psi}\|_{L^2}^2 + \|\partial_y \widehat{\psi}\|_{L^2}^2 \ge 0, \\ K_2 K_3 \ge K_1^2. \end{split}$$

Proof. By (11.6) and Proposition 10.2, we get

$$\begin{split} \|\mathbb{D}(\widehat{\omega})\|_{L^{2}}^{2} &= K_{1} \geq (1 - \alpha^{-2}) \|\widehat{\omega}\|_{L^{2}}^{2} \\ &= (1 - \alpha^{-2}) (\|\partial_{y}^{2}\widehat{\psi}\|_{L^{2}}^{2} + \alpha^{2} \|\partial_{y}\widehat{\psi}\|_{L^{2}}^{2} + \alpha^{4} \|\widehat{\psi}\|_{L^{2}}^{2}) \\ &\geq \alpha^{2} ((\alpha^{2} - 1) \|\widehat{\psi}\|_{L^{2}}^{2} + \|\partial_{y}\widehat{\psi}\|_{L^{2}}^{2}) = \alpha^{2} K_{3} \geq 0. \end{split}$$

It is easy to check that

$$K_2 = -\langle (\partial_y^2 - \alpha^2)\widehat{\omega}, \widehat{\omega} - \widehat{\psi} \rangle = \langle \partial_y \widehat{\omega}, \partial_y \widehat{\omega} - \partial_y \widehat{\psi} \rangle + \alpha^2 \langle \widehat{\omega}, \widehat{\omega} - \widehat{\psi} \rangle,$$

which along with (11.5) gives $K_2 \ge \alpha^2 K_1 \ge 0$.

Using the facts that

$$\begin{split} \|\widehat{\omega}\|_{L^{2}}^{2} &= \|\partial_{y}^{2}\widehat{\psi}\|_{L^{2}}^{2} + \alpha^{2}\|\partial_{y}\widehat{\psi}\|_{L^{2}}^{2} + \alpha^{4}\|\widehat{\psi}\|_{L^{2}}^{2}, \\ \|\partial_{y}\widehat{\omega}\|_{L^{2}}^{2} &= \|\partial_{y}^{3}\widehat{\psi}\|_{L^{2}}^{2} + \alpha^{2}\|\partial_{y}^{2}\widehat{\psi}\|_{L^{2}}^{2} + \alpha^{4}\|\partial_{y}\widehat{\psi}\|_{L^{2}}^{2}, \end{split}$$

we can deduce that

$$K_{2}K_{3} = \left((\alpha^{2} - 1) \|\widehat{\omega}\|_{L^{2}}^{2} + \|\partial_{y}\widehat{\omega}\|_{L^{2}}^{2} \right) \left((\alpha^{2} - 1) \|\widehat{\psi}\|_{L^{2}}^{2} + \|\partial_{y}\widehat{\psi}\|_{L^{2}}^{2} \right)$$

$$\geq \left((\alpha^{2} - 1) \langle \widehat{\omega}, \widehat{\psi} \rangle + \langle \partial_{y}\widehat{\omega}, \partial_{y}\widehat{\psi} \rangle \right)^{2} = \left(\langle \widehat{\omega}, (\alpha^{2} - 1)\widehat{\psi} - \partial_{y}^{2}\widehat{\psi} \rangle \right)^{2}$$

$$= \langle \widehat{\omega}, \widehat{\omega} - \widehat{\psi} \rangle^{2} = K_{1}^{2}.$$

Г		
L		
L		
L		

11.3. Proof of Theorem 1.4.

Proof. We first consider the case of $\delta < 1$. Taking Fourier transform in x to (1.5), we obtain $\partial_t \hat{\omega} + \mathcal{L}_{\nu}(\alpha, t) \hat{\omega} = 0,$

where

$$\mathcal{L}_{\nu}(\alpha,t) = \nu(-\partial_y^2 + \alpha^2) - i\alpha a_0 e^{-\nu t} \cos y \left(1 + (\partial_y^2 - \alpha^2)^{-1}\right).$$

Let $-(\partial_y^2 - \alpha^2)\widehat{\psi} = \widehat{\omega}$. Then we have

$$\begin{aligned} \partial_t \langle \widehat{\omega}, \widehat{\omega} - \widehat{\psi} \rangle &= \langle \widehat{\omega}_t, \widehat{\omega} \rangle + \langle \widehat{\omega}, \widehat{\omega}_t \rangle - \langle \widehat{\psi}, \widehat{\omega}_t \rangle - \langle \widehat{\omega}_t, \widehat{\psi} \rangle \\ &= -2\nu \langle \partial_y \widehat{\omega}, \partial_y \widehat{\omega} - \partial_y \widehat{\psi} \rangle - 2\nu \alpha^2 \langle \widehat{\omega}, \widehat{\omega} - \widehat{\psi} \rangle. \end{aligned}$$

This gives by Lemma 11.4 that

$$\partial_t K_1 = -2\nu K_2 \le -2\nu\alpha^2 K_1,$$

which implies that

(11.7)
$$K_1(t,\alpha) \le e^{-2\nu\alpha^2 t} K_1(0,\alpha), \quad \int_0^T 2\nu K_2(s,\alpha) ds = K_1(0,\alpha) - K_1(T,\alpha).$$

Now fix α , consider the following cases.

Case 1. $\nu \ge a_0 |\alpha| e^{-\tau}$ or $\nu |\alpha|^3 \ge 1$.

Due to $|\alpha| > 1$, we have $\nu^{\frac{1}{3}} + \nu^{\frac{1}{2}} \leq C\nu\alpha^2$. Then by Lemma 11.4 and (11.7), we get

$$\alpha^{2}K_{3}(t,\alpha) \leq K_{1}(t,\alpha) \leq e^{-2\nu\alpha^{2}t}K_{1}(0,\alpha) \leq \frac{Ce^{-\nu\alpha^{2}t}}{1+\nu t^{3}}K_{1}(0,\alpha) \leq \frac{Ce^{-c\sqrt{\nu}t}}{1+\nu t^{3}}K_{1}(0,\alpha).$$

Case 2: $\nu < a_0 |\alpha| e^{-\tau}$ and $\nu |\alpha|^3 < 1$.

Take $t_1 = (\nu |\alpha|)^{-\frac{1}{2}}$, then $\nu t_1 \alpha^2 < 1$. Fix $0 \le t \le \tau / \nu$, then $(e^{\nu \alpha^2 s} \widehat{\omega}(t+s, \alpha, y), e^{\nu \alpha^2 s} \widehat{\psi}(t+s, \alpha, y))$ solves (11.4) with $a = a_0 \alpha e^{-\nu t}$. By our assumption, $0 < \nu < |a|$. Then by Lemma 11.3, we obtain

$$e^{2\nu\alpha^{2}t_{1}}K_{3}(t+t_{1},\alpha)$$

$$= \langle e^{\nu\alpha^{2}t_{1}}\widehat{\psi}(t+t_{1},\alpha), e^{\nu\alpha^{2}t_{1}}\widehat{\omega}(t+t_{1},\alpha) - e^{\nu\alpha^{2}s}\widehat{\psi}(t+t_{1},\alpha)\rangle$$

$$\leq \frac{C}{\nu t_{1}^{3}a^{2}} \langle e^{\nu\alpha^{2}t_{1}}\widehat{\omega}(t,\alpha), e^{\nu\alpha^{2}t_{1}}\widehat{\omega}(t,\alpha) - e^{\nu\alpha^{2}t_{1}}\widehat{\psi}(t,\alpha)\rangle$$

$$\leq \frac{C}{\nu t_{1}^{3}a^{2}}K_{1}(t,\alpha) \leq \frac{C}{\nu t_{1}^{3}\alpha^{2}}K_{1}(t,\alpha) = CK_{1}(t,\alpha)t_{1}\nu.$$

Thus, there exists an absolute constant $c_0 \in (0, 1)$ so that

(11.8)
$$c_0 e^{c_0} K_3(t+t_1,\alpha) \le K_1(t,\alpha) t_1 \nu_1$$

Set $M = \max_{0 \le t \le \tau/\nu} e^{c_0 t/t_1} K_1(t, \alpha)$. Then there exists $t_2 \in [0, \tau/\nu]$ so that $e^{c_0 t_2/t_1} K_1(t_2, \alpha) = M$. Next we will show that $t_2 \le t_1$, otherwise $K_1 = 0$. If $t_2 > t_1$, then $\partial_t (e^{c_0 t/t_1} K_1(t, \alpha))|_{t=t_2} \ge 0$. Since

$$\partial_t (e^{c_0 t/t_1} K_1(t, \alpha)) = e^{c_0 t/t_1} (c_0/t_1 K_1(t, \alpha) + \partial_t K_1(t, \alpha))$$

= $e^{c_0 t/t_1} ((c_0/t_1) K_1(t, \alpha) - 2\nu K_2(t, \alpha)),$

we obtain

(11.9)
$$K_2(t_2, \alpha) \le \frac{c_0}{2\nu t_1} K_1(t_2, \alpha).$$

By (11.8), we get

$$c_0 e^{c_0} K_3(t_2, \alpha) \le K_1(t_2 - t_1, \alpha) t_1 \nu$$

$$\le e^{-c_0(t_2 - t_1)/t_1} M t_1 \nu$$

$$= e^{-c_0(t_2 - t_1)/t_1} e^{c_0 t_2/t_1} K_1(t_2, \alpha) t_1 \nu = e^{c_0} K_1(t_2, \alpha) t_1 \nu,$$

which gives

(11.10)
$$K_3(t_2, \alpha) \le K_1(t_2, \alpha) t_1 \nu / c_0.$$

It follows from (11.9) and (11.10) that

$$K_2(t_2, \alpha) K_3(t_2, \alpha) \le \frac{1}{2} K_1(t_2, \alpha)^2$$

On the other hand, $K_2K_3 \ge K_1^2$ by Lemma 11.4. Then we conclude that $K_1(t_2, \alpha) = 0$, which implies M = 0.

Thus, we only need to deal with the case $t_2 \leq t_1$. Then

$$M = e^{c_0 t_2/t_1} K_1(t_2, \alpha) \le e^{c_0} K_1(t_2, \alpha) \le e^{c_0} K_1(0, \alpha).$$

Therefore, for $0 \le t \le \tau/\nu$,

(11.11)
$$K_1(t,\alpha) \le e^{-c_0 t/t_1} M \le e^{c_0 - c_0 t/t_1} K_1(0,\alpha) = e^{c_0 - c_0 \sqrt{\nu |\alpha|} t} K_1(0,\alpha).$$

This inequality is also true for **Case 1** by (11.7), since we have $\nu |\alpha|^3 \ge C^{-1}$ or $\nu \alpha^2 \ge C^{-1} \sqrt{\nu |\alpha|}$ in **Case 1**.

As $\left(e^{\nu\alpha^2 t}\widehat{\omega}(t,\alpha,y), e^{\nu\alpha^2 t}\widehat{\psi}(t,\alpha,y)\right)$ solves (11.4) with $a = a_0\alpha$ and $|\alpha| > 1$, we deduce from Lemma 11.3 that $0 < t \le (\nu|\alpha|)^{-\frac{1}{2}}$ and $\nu\alpha^2 t < 1$,

$$e^{2\nu\alpha^2 t} K_3(t,\alpha) \le \frac{C}{\nu t^3 a^2} K_1(0,\alpha) \le \frac{C e^{-c_0 t/(2t_1)}}{\nu t^3 \alpha^2} K_1(0,\alpha)$$

for $0 \le t \le t_1$. For $t_1 \le t \le \tau/\nu$, by (11.8), we have

$$c_0 e^{c_0} K_3(t,\alpha) \le K_1(t-t_1,\alpha) t_1 \nu \le e^{c_0 - c_0(t-t_1)/t_1} K_1(0,\alpha) t_1 \nu,$$

which gives

$$\begin{aligned} \alpha^2 K_3(t,\alpha) &\leq c_0^{-1} e^{c_0 - c_0 t/t_1} K_1(0,\alpha) t_1 \nu \alpha^2 \\ &= c_0^{-1} e^{c_0 - c_0 t/t_1} K_1(0,\alpha) t_1 \nu / (t_1^2 \nu)^2 \\ &= c_0^{-1} e^{c_0 - c_0 t/t_1} K_1(0,\alpha) / (t_1^3 \nu) \leq C e^{-c_0 t/(2t_1)} K_1(0,\alpha) / (t^3 \nu). \end{aligned}$$

Therefore, for $0 \le t \le \tau/\nu$,

$$\alpha^2 K_3(t,\alpha) \le C e^{-c_0 t/(2t_1)} K_1(0,\alpha)/(\nu t^3).$$

While by (11.11),

$$\alpha^2 K_3(t,\alpha) \le K_1(t,\alpha) \le e^{c_0 - c_0 t/t_1} K_1(0,\alpha).$$

This shows that for $0 \le t \le \tau/\nu$,

$$\alpha^2 K_3(t,\alpha) \le C e^{-c_0 t/(2t_1)} K_1(0,\alpha) / (1+\nu t^3) = \frac{C e^{-c_0} \sqrt{\nu |\alpha| t/2}}{1+\nu t^3} K_1(0,\alpha).$$

Combining two cases, we deduce that there exists an absolute constant $c_1 \in (0, 1)$ so that for $0 \le t \le \tau/\nu$,

(11.12)
$$K_1(t,\alpha) \le Ce^{-2c_1\sqrt{\nu}t}K_1(0,\alpha),$$

(11.13)
$$\alpha^2 K_3(t,\alpha) \le \frac{Ce^{-2c_1\sqrt{\nu t}}}{1+\nu t^3} K_1(0,\alpha).$$

By Proposition 10.2, we have for $0 \le t \le \tau/\nu$,

$$\begin{split} \|\omega(t)\|_{L^{2}}^{2} &= \sum_{\alpha \neq 0} \|\widehat{\omega}(t,\alpha)\|_{L^{2}}^{2} \leq \sum_{\alpha \neq 0} \frac{\alpha^{2}}{\alpha^{2} - 1} \|\mathbb{D}(\widehat{\omega})(t,\alpha)\|_{L^{2}}^{2} \\ &= \sum_{\alpha \neq 0} \frac{\alpha^{2}}{\alpha^{2} - 1} K_{1}(t,\alpha) \leq C \sum_{\alpha \neq 0} e^{-2c_{1}\sqrt{\nu}t} K_{1}(0,\alpha) \\ &\leq C \sum_{\alpha \neq 0} e^{-2c_{1}\sqrt{\nu}t} \|\widehat{\omega}(0,\alpha)\|_{L^{2}}^{2} \\ &= C e^{-2c_{1}\sqrt{\nu}t} \|\omega_{0}\|_{L^{2}}^{2}. \end{split}$$

Using (11.6), we have for $0 \le t \le \tau/\nu$,

$$\|V(t)\|_{\dot{H}^{1}_{x}L^{2}_{y}}^{2} = \sum_{\alpha \neq 0} \alpha^{2} (\alpha^{2} \|\widehat{\psi}(t,\alpha)\|_{L^{2}}^{2} + \|\partial_{y}\widehat{\psi}(t,\alpha)\|_{L^{2}}^{2})$$

$$\leq \sum_{\alpha \neq 0} \frac{\alpha^4}{\alpha^2 - 1} K_3(t, \alpha)$$

$$\leq C \sum_{\alpha \neq 0} \frac{e^{-2c_1 \sqrt{\nu t}}}{1 + \nu t^3} K_1(0, \alpha) \leq \frac{C e^{-2c_1 \sqrt{\nu t}}}{1 + \nu t^3} \|\omega_0\|_{L^2}^2$$

Therefore for $0 \le t \le \tau/\nu$,

$$\|\omega(t)\|_{L^2} \le Ce^{-c_1\sqrt{\nu}t} \|\omega_0\|_{L^2}, \quad \|V(t)\|_{\dot{H}^1_x L^2_y} \le \frac{Ce^{-c_1\sqrt{\nu}t}}{\sqrt{1+\nu t^3}} \|\omega_0\|_{L^2}$$

For the case of $\delta = 1$, the proof is almost the same by making the orthogonal projection $P_{|\alpha|\geq 2}$, since \mathcal{R}_{α} has no eigenvalue and embedding eigenvalues for $|\alpha|\geq 2$ and Lemma 11.3-Lemma 11.4 hold for $|\alpha|\geq 2$.

Let us conclude this section by introducing some space-time estimates of the solution for the linearized Navier-Stokes equations, which will be used in the proof of nonlinear enhanced dissipation.

Lemma 11.5. Let $\delta < 1$. Under the same assumptions as in Theorem 1.4, we have

$$\int_{0}^{\tau/\nu} \|\nabla\omega(t)\|_{L^{2}}^{2} dt \leq C\nu^{-1} \|\omega_{0}\|_{L^{2}}^{2},$$
$$\int_{0}^{\tau/\nu} \|\partial_{x}\omega(t)\|_{L^{2}} dt \leq C\nu^{-\frac{2}{3}} \|\omega_{0}\|_{L^{2}},$$
$$\int_{0}^{\tau/\nu} \|V(t)\|_{L^{\infty}}^{2} dt \leq C(|\ln\nu|+1)\nu^{-\frac{1}{3}} \|\omega_{0}\|_{L^{2}}^{2}.$$

Proof. By (11.7), we have

$$\begin{split} \int_{0}^{\tau/\nu} \|\nabla\omega(t)\|_{L^{2}}^{2} dt &= \int_{0}^{\tau/\nu} \sum_{\alpha \neq 0} \left(|\alpha|^{2} \|\widehat{\omega}(t,\alpha)\|_{L^{2}}^{2} + \|\partial_{y}\widehat{\omega}(t,\alpha)\|_{L^{2}}^{2} \right) dt \\ &\leq \int_{0}^{\tau/\nu} \sum_{\alpha \neq 0} \frac{\alpha^{2}}{\alpha^{2} - 1} K_{2}(t,\alpha) dt \\ &= \frac{1}{2\nu} \sum_{\alpha \neq 0} \frac{\alpha^{2}}{\alpha^{2} - 1} (K_{1}(0,\alpha) - K_{1}(\tau/\nu,\alpha)) \\ &\leq \frac{C}{\nu} \sum_{\alpha \neq 0} K_{1}(0,\alpha) \leq \frac{C}{\nu} \sum_{\alpha \neq 0} \|\widehat{\omega}(0,\alpha)\|_{L^{2}}^{2} = \frac{C}{\nu} \|\omega_{0}\|_{L^{2}}^{2}. \end{split}$$

Using the fact that $\partial_t K_1 \leq -2\nu \alpha^2 K_1$, we have

$$|\alpha|^{2} K_{1}(t,\alpha) \leq |\alpha|^{2} e^{-2\nu\alpha^{2}t} K_{1}(0,\alpha) \leq C K_{1}(0,\alpha)/(\nu t)$$

and (11.11) implies that

$$|\alpha|^2 K_1(t,\alpha) \le |\alpha|^2 C e^{-2c_1 \sqrt{\nu |\alpha|} t} K_1(0,\alpha) \le C K_1(0,\alpha) / (\nu^2 t^4),$$

from which, we infer that

$$\|\partial_x \omega(t)\|_{L^2}^2 = \sum_{\alpha \neq 0} |\alpha|^2 \|\widehat{\omega}(t, \alpha)\|_{L^2}^2 \le \sum_{\alpha \neq 0} |\alpha|^2 \frac{\alpha^2}{\alpha^2 - 1} K_1(t, \alpha)$$

$$\leq \sum_{\alpha \neq 0} CK_1(0, \alpha) \min\{(\nu t)^{-1}, (\nu^2 t^4)^{-1}\}$$

$$\leq C \|\omega_0\|_{L^2}^2 \min\{(\nu t)^{-1}, (\nu^2 t^4)^{-1}\}.$$

This gives

$$\int_0^{\tau/\nu} \|\partial_x \omega(t)\|_{L^2} dt \le \int_0^{\tau/\nu} C \|\omega_0\|_{L^2} \min\{(\nu t)^{-\frac{1}{2}}, (\nu t^2)^{-1}\} dt \le C\nu^{-\frac{2}{3}} \|\omega_0\|_{L^2}.$$

By Minkowski inequality, we get

$$\int_0^{\tau/\nu} \|V(t)\|_{L^\infty}^2 dt \le \int_0^{\tau/\nu} \left(\sum_{\alpha \ne 0} \|\widehat{V}(t,\alpha,\cdot)\|_{L^\infty}\right)^2 dt \le \left(\sum_{\alpha \ne 0} \left(\int_0^{\tau/\nu} \|\widehat{V}(t,\alpha,\cdot)\|_{L^\infty}^2 dt\right)^{\frac{1}{2}}\right)^2,$$

and by the interpolation,

$$\begin{split} \|\widehat{V}(t,\alpha,\cdot)\|_{L^{\infty}}^{2} &\leq C \|\widehat{V}(t,\alpha,\cdot)\|_{L^{2}} \|\widehat{V}(t,\alpha,\cdot)\|_{H^{1}} \\ &\leq C \left(\alpha^{2} \|\widehat{\psi}(t,\alpha,\cdot)\|_{L^{2}}^{2} + \|\partial_{y}\widehat{\psi}(t,\alpha,\cdot)\|_{L^{2}}^{2}\right)^{\frac{1}{2}} \\ & \times \left(\alpha^{4} \|\widehat{\psi}(t,\alpha,\cdot)\|_{L^{2}}^{2} + 2\alpha^{2} \|\partial_{y}\widehat{\psi}(t,\alpha,\cdot)\|_{L^{2}}^{2} + \|\partial_{y}^{2}\widehat{\psi}(t,\alpha,\cdot)\|_{L^{2}}^{2}\right)^{\frac{1}{2}} \\ &\leq C K_{3}(t,\alpha)^{\frac{1}{2}} K_{1}(t,\alpha)^{\frac{1}{2}} \leq C K_{3}(t,\alpha)^{\frac{3}{4}} K_{2}(t,\alpha)^{\frac{1}{4}}, \end{split}$$

from which and (11.13), (11.7), we infer that

$$\int_{0}^{\tau/\nu} \|\widehat{V}(t,\alpha)\|_{L^{\infty}}^{2} dt \leq \int_{0}^{\tau/\nu} CK_{3}(t,\alpha)^{\frac{1}{2}} K_{1}(t,\alpha)^{\frac{1}{2}} dt$$
$$\leq \int_{0}^{\tau/\nu} \frac{CK_{1}(0,\alpha)}{|\alpha|\sqrt{1+\nu t^{3}}} dt \leq \frac{CK_{1}(0,\alpha)}{|\alpha|\nu^{\frac{1}{3}}},$$

and

$$\begin{split} \int_{0}^{\tau/\nu} \|\widehat{V}(t,\alpha)\|_{L^{\infty}}^{2} dt &\leq \int_{0}^{\tau/\nu} CK_{3}(t,\alpha)^{\frac{3}{4}} K_{2}(t,\alpha)^{\frac{1}{4}} dt \\ &\leq C \left(\int_{0}^{\tau/\nu} K_{3}(t,\alpha) dt \right)^{\frac{3}{4}} \left(\int_{0}^{\tau/\nu} K_{2}(t,\alpha) dt \right)^{\frac{1}{4}} \\ &\leq C \left(\int_{0}^{\tau/\nu} \frac{K_{1}(0,\alpha)}{\alpha^{2}(1+\nu t^{3})} dt \right)^{\frac{3}{4}} \left(\frac{K_{1}(0,\alpha) - K_{1}(\tau/\nu,\alpha)}{2\nu} \right)^{\frac{1}{4}} \\ &\leq C \left(\frac{K_{1}(0,\alpha)}{\alpha^{2}\nu^{\frac{1}{3}}} \right)^{\frac{3}{4}} \left(\frac{K_{1}(0,\alpha)}{2\nu} \right)^{\frac{1}{4}} \leq \frac{CK_{1}(0,\alpha)}{|\alpha|^{\frac{3}{2}}\nu^{\frac{1}{2}}}. \end{split}$$

Therefore,

$$\int_{0}^{\tau/\nu} \|V(t)\|_{L^{\infty}}^{2} dt \leq \left(\sum_{\alpha \neq 0} \left(\int_{0}^{\tau/\nu} \|\widehat{V}(t,\alpha)\|_{L^{\infty}}^{2} dt \right)^{\frac{1}{2}} \right)^{2} \\ \leq \left(\sum_{0 < |\alpha| \le \nu^{-\frac{1}{3}}} \left(\frac{CK_{1}(0,\alpha)}{|\alpha|\nu^{\frac{1}{3}}} \right)^{\frac{1}{2}} + \sum_{|\alpha| > \nu^{-\frac{1}{3}}} \left(\frac{CK_{1}(0,\alpha)}{|\alpha|^{\frac{3}{2}}\nu^{\frac{1}{2}}} \right)^{\frac{1}{2}} \right)^{2}$$

$$\leq C\left(\sum_{\alpha\neq0}K_{1}(0,\alpha)\right)\left(\sum_{0<|\alpha|\leq\nu^{-\frac{1}{3}}}\frac{1}{|\alpha|^{\frac{1}{3}}} + \sum_{|\alpha|>\nu^{-\frac{1}{3}}}\frac{1}{|\alpha|^{\frac{3}{2}}\nu^{\frac{1}{2}}}\right)$$
$$\leq C\left(\sum_{\alpha\neq0}\|\widehat{\omega}(0,\alpha)\|_{L^{2}}^{2}\right)\left(\frac{1+|\ln\nu|}{\nu^{\frac{1}{3}}} + \frac{\nu^{\frac{1}{6}}}{\nu^{\frac{1}{2}}}\right)$$
$$\leq C\|\omega_{0}\|_{L^{2}}^{2}(1+|\ln\nu|)/\nu^{\frac{1}{3}},$$

which gives the third inequality.

12. Nonlinear enhanced dissipation

In this section, we prove Theorem 1.6. We define $W_2 = \operatorname{span}\{\sin y, \cos y\}$ and the non-shear space

$$X = \Big\{ \omega \in L^2(\mathbb{T}^2_{\delta}) : \omega(x,y) = \sum_{\alpha \neq 0} \widehat{\omega}(\alpha,y) e^{i\alpha x} \Big\}.$$

We denote by P_2 the orthogonal projection to $W_2 = \operatorname{span}\{\sin y, \cos y\}$, and $P_{\neq 0}$ the orthogonal projection to X, and $P_0 = I - P_{\neq 0}$ the orthogonal projection to the shear space $X_0 = \{\omega \in L^2(\mathbb{T}) : \partial_x \omega = 0\}$.

12.1. Semigroup estimates. Let $S(t, s) = S(t, s, a_0)$ be the solution operator of linearized equation (1.5). That is, $\omega(t, y) = S(t, s)f(y)$ solves (1.5) with $\omega(s, y) = f(y)$, here we assume $t \ge s$. Now Theorem 1.4 and Lemma 11.5 can be restated as follows

$$\begin{split} \|S(t,0)f\|_{L^{2}} &\leq Ce^{-c_{1}\sqrt{\nu t}} \|f\|_{L^{2}} \text{ for } 0 < t < \tau/\nu, \\ \int_{0}^{\tau/\nu} \|\nabla S(t,0)f\|_{L^{2}}^{2} dt \leq C\nu^{-1} \|f\|_{L^{2}}^{2}, \\ \int_{0}^{\tau/\nu} \|\partial_{x}S(t,0)f\|_{L^{2}} dt \leq C\nu^{-\frac{2}{3}} \|f\|_{L^{2}}, \\ \int_{0}^{\tau/\nu} \|\nabla \Delta^{-1}S(t,0)f\|_{L^{\infty}}^{2} dt \leq C(|\ln\nu|+1)\nu^{-\frac{1}{3}} \|f\|_{L^{2}}^{2}. \end{split}$$

Notice that $S(t, s, a_0) = S(t - s, 0, a_0 e^{-\nu s})$ and $a_0 e^{-\nu s}$ has uniform positive upper and lower bounds. Then we also have for $0 \le s < t < \tau/\nu$,

(12.1)
$$\|S(t,s)f\|_{L^2} \le Ce^{-c_1\sqrt{\nu}(t-s)}\|f(s)\|_{L^2},$$

(12.2)
$$\int_{s}^{\tau/\nu} \|\nabla S(t,s)f\|_{L^{2}}^{2} dt \leq C\nu^{-1} \|f(s)\|_{L^{2}}^{2},$$

(12.3)
$$\int_{s}^{\tau/\nu} \|\partial_x S(t,s)f\|_{L^2} dt \le C\nu^{-\frac{2}{3}} \|f(s)\|_{L^2},$$

(12.4)
$$\int_{s}^{\tau/\nu} \|\nabla \Delta^{-1} S(t,s) f\|_{L^{\infty}}^{2} dt \leq C \nu^{-\gamma_{1}} \|f(s)\|_{L^{2}}^{2},$$

where γ_1 is a constant such that $\frac{1}{3} < \gamma_1 < 2\gamma - 1$.

Let f be the solution of the inhomogeneous linearized Navier-Stokes equations:

(12.5)
$$\partial_t f + \mathcal{L}_{\nu}(t)f = g.$$

Then we have

$$f(t) = S(t,0)f(0) + \int_0^t S(t,s)g(s)ds.$$

Using (12.1)-(12.4) and the Minkowski inequality, we deduce that for $0 < t < \tau/\nu$,

$$(12.6) ||f(t)||_{L^{2}} \leq Ce^{-c_{1}\sqrt{\nu}t} ||f(0)||_{L^{2}} + C \int_{0}^{t} ||g(s)||_{L^{2}} ds, \nu^{\frac{1}{2}} \left(\int_{0}^{t} ||\nabla f(s)||_{L^{2}}^{2} ds \right)^{\frac{1}{2}} + \nu^{\frac{2}{3}} \int_{0}^{t} ||\partial_{x}f(s)||_{L^{2}} ds + \nu^{\frac{\gamma_{1}}{2}} \left(\int_{0}^{t} ||\nabla \Delta^{-1}f(s)||_{L^{\infty}}^{2} ds \right)^{\frac{1}{2}} (12.7) \leq C ||f(0)||_{L^{2}} + C \int_{0}^{t} ||g(s)||_{L^{2}} ds,$$

where C, c_1 are constants independent of ν, t .

12.2. **Proof of Theorem 1.6.** First of all, we have the following energy dissipation law: $\partial_t \|V\|_{L^2}^2 = -2\nu \|\nabla V\|_{L^2}^2, \quad \partial_t \|\omega\|_{L^2}^2 = -2\nu \|\nabla \omega\|_{L^2}^2,$

which gives

(12.8)
$$\partial_t (\|\omega\|_{L^2}^2 - \|\nabla\psi\|_{L^2}^2) = -2\nu (\|\nabla\omega\|_{L^2}^2 - \|\omega\|_{L^2}^2),$$

where $\psi = -\Delta^{-1}\omega$. The key point is that if $\delta < 1$, it holds that for $\omega \in \{1\}^{\perp}$,

(12.9)
$$\|(I-P_2)\omega\|_{L^2}^2 \ge \|\omega\|_{L^2}^2 - \|\nabla\psi\|_{L^2}^2 \ge C_0 \|(I-P_2)\omega\|_{L^2}^2,$$

(12.10)
$$\|\nabla \omega\|_{L^2}^2 - \|\omega\|_{L^2}^2 \ge C_0 \|\nabla (I - P_2)\omega\|_{L^2}^2$$

which can be easily proved by using Fourier transform.

By (12.8)-(12.10), we have

(12.11)
$$\| (I - P_2)\omega(t) \|_{L^2}^2 \le C(\|\omega(t)\|_{L^2}^2 - \|\nabla\psi(t)\|_{L^2}^2)$$

$$\le C(\|\omega(0)\|_{L^2}^2 - \|\nabla\psi(0)\|_{L^2}^2) \le C\|(I - P_2)\omega(0)\|_{L^2}^2 \le C\nu^{2\gamma},$$

for $0 < t < \tau/\nu$, and

(12.12)
$$\int_{0}^{\tau/\nu} \|\nabla (I - P_2)\omega(t)\|_{L^2}^2 dt \leq C \int_{0}^{\tau/\nu} (\|\nabla \omega(t)\|_{L^2}^2 - \|\omega(t)\|_{L^2}^2) dt$$
$$\leq C(\|\omega(0)\|_{L^2}^2 - \|\nabla \psi(0)\|_{L^2}^2)/(2\nu)$$
$$\leq C\|(I - P_2)\omega(0)\|_{L^2}^2/(2\nu) \leq C\nu^{2\gamma-1}.$$

Now, we decompose $\omega = \omega_s + \omega_n$, where $\omega_s = P_0 \omega$ is the shear part and $\omega_n = P_{\neq 0} \omega$. Correspondingly, $V = V_s + V_n$, $\psi = \psi_s + \psi_n$. We denote $V_s = (V^1(t, y), 0)$, $V_n = (V_n^1, V_n^2)$. Then the equation (1.4) can be written as

(12.13)
$$\partial_t \omega_s = \nu \Delta \omega_s - P_0(V_n \cdot \nabla \omega_n) = \nu \partial_y^2 \omega_s - \partial_y P_0(V_n^2 \omega_n),$$

(12.14)
$$\partial_t \omega_n = \nu \Delta \omega_n - V_s^1 \partial_x \omega_n - V_n \cdot \nabla \omega_s - P_{\neq 0}(V_n \cdot \nabla \omega_n).$$

For $P_2\omega = P_2\omega_s$, we have

$$\partial_t P_2 \omega = \nu \Delta P_2 \omega - \partial_y P_2(V_n^2 \omega_n) = -\nu P_2 \omega - \partial_y P_2(V_n^2 \omega_n),$$

therefore,

$$P_{2}\omega(t) = e^{-\nu t} P_{2}\omega(0) - \int_{0}^{t} e^{-\nu(t-s)} \partial_{y} P_{2}(V_{n}^{2}\omega_{n})(s) ds,$$

from which and (12.11), we deduce that for $0 < t < \tau/\nu$,

$$(12.15) \quad \|P_{2}\omega(t) - e^{-\nu t}P_{2}\omega(0)\|_{L^{2}} \leq \int_{0}^{t} \|\partial_{y}P_{2}(V_{n}^{2}\omega_{n})(s)\|_{L^{2}}ds$$
$$\leq \int_{0}^{t} C\|(V_{n}^{2}\omega_{n})(s)\|_{L^{1}}ds \leq \int_{0}^{t} C\|V_{n}^{2}(s)\|_{L^{2}}\|\omega_{n}(s)\|_{L^{2}}ds$$
$$\leq \int_{0}^{t} C\|\omega_{n}(s)\|_{L^{2}}^{2}ds \leq \int_{0}^{t} C\|(I - P_{2})\omega(s)\|_{L^{2}}^{2}ds$$
$$\leq \int_{0}^{t} C\nu^{2\gamma}ds = Ct\nu^{2\gamma}.$$

In particular, we have

$$||P_2\omega(t) - e^{-\nu t}P_2\omega(0)||_{L^2} \le C\tau\nu^{2\gamma-1},$$

Thanks to $C_1^{-1} \le ||P_2\omega(0)||_{L^2} \le C_1$, there exist $c_2 \in (0,1), C > 1$ so that if $0 < \nu < c_2$, then (12.16) $C^{-1} \le ||P_2\omega(t)||_{L^2} \le C$ for $0 < t < \tau/\nu$.

Now we choose $t_0 \sim \nu^{-\frac{1}{2}}$ so that $Ce^{-c_1\sqrt{\nu}t_0} = \frac{1}{4}$. Then it suffices to show that for $0 \leq t' < t < \tau/\nu$,

(12.17)
$$\begin{aligned} \|\omega_n(t)\|_{L^2} &\leq C \|\omega_n(t')\|_{L^2} &\text{if } t-t' < t_0, \\ \|\omega_n(t)\|_{L^2} &\leq \frac{1}{2} \|\omega_n(t')\|_{L^2} &\text{if } t-t' = t_0. \end{aligned}$$

Indeed, Theorem 1.6 follows from (12.17) by using the following iteration

$$\|\omega_n(t)\|_{L^2} \le C \|\omega_n(mt_0)\|_{L^2} \le C \frac{1}{2^m} \|\omega_n(0)\|_{L^2} \le C e^{-c\sqrt{\nu}t} \|\omega_n(0)\|_{L^2},$$

with $m = [t/t_0]$ and $t_0 \le C\nu^{-\frac{1}{2}}$.

By time translation and (12.16), we only need to prove that (12.17) for t' = 0. Let $P_2\omega(0) = -a_0 \sin(y+\theta)$, where $a_0 > 0$, $\theta \in [0, 2\pi)$. Then $||P_2\omega(0)||_{L^2}/||\sin y||_{L^2} = a_0$. By translation, we may assume $\theta = 0$. Thus,

$$P_2\omega(0) = -a_0 \sin y, \quad P_2V(0) = (-a_0 \cos y, 0).$$

Let

$$\omega(t) = e^{-\nu t} P_2 \omega(0) + \widetilde{\omega}(t), \quad V(t) = e^{-\nu t} P_2 V(0) + \widetilde{V}(t).$$

We decompose $\tilde{\omega} = \tilde{\omega}_s + \omega_n$ and $\tilde{V} = \tilde{V}_s + V_n$, where $\tilde{\omega}_s = P_0 \tilde{\omega}$ and $\tilde{V}_s = P_0 \tilde{V}$. Then (12.14) can be rewritten as

(12.18)
$$\partial_t \omega_n + \mathcal{L}_{\nu}(t)\omega_n = -\left(\widetilde{V}_s^1 \partial_x \omega_n + V_n \cdot \nabla \widetilde{\omega}_s + P_{\neq 0}(V_n \cdot \nabla \omega_n)\right).$$

For $0 < t \le \min\{t_0, \tau/\nu\}$, we infer from (12.15) that

(12.19)
$$\|P_2\widetilde{\omega}_s(t)\|_{L^2} = \|P_2\omega(t) - e^{-\nu t}P_2\omega(0)\|_{L^2} \le Ct\nu^{2\gamma} \le Ct_0\nu^{2\gamma} \le C\nu^{2\gamma-\frac{1}{2}}.$$

Using (12.11), (12.19) and the fact that $(I - P_2)\widetilde{\omega}_s(t) = P_0(I - P_2)\omega(t)$, we deduce that

(12.20)
$$\|V_s^1(t)\|_{L^{\infty}} \leq C \|\widetilde{\omega}_s(t)\|_{L^2} \leq C \|P_2\widetilde{\omega}_s(t)\|_{L^2} + C \|(I-P_2)\omega(t)\|_{L^2}$$
$$\leq C\nu^{2\gamma-\frac{1}{2}} + C\nu^{\gamma} \leq C\nu^{\gamma}.$$

Using (12.12) and (12.19), we deduce that

(12.21)
$$\int_{0}^{t} \|\nabla\widetilde{\omega}(s)\|_{L^{2}}^{2} ds = \int_{0}^{t} \|\nabla P_{2}\widetilde{\omega}_{s}(s)\|_{L^{2}}^{2} ds + \int_{0}^{t} \|\nabla (I - P_{2})\widetilde{\omega}_{s}(s)\|_{L^{2}}^{2} ds$$
$$\leq C \int_{0}^{t} \|P_{2}\widetilde{\omega}_{s}(s)\|_{L^{2}}^{2} ds + \int_{0}^{t} \|\nabla (I - P_{2})\omega(s)\|_{L^{2}}^{2} ds$$
$$\leq C \int_{0}^{\tau/\nu} \nu^{4\gamma - 1} ds + C\nu^{2\gamma - 1}$$
$$= C\tau\nu^{4\gamma - 2} + C\nu^{2\gamma - 1} \leq C\nu^{2\gamma - 1}.$$

Let $f = \omega_n$ and

$$g = -\left(\widetilde{V}_s^1 \partial_x \omega_n + V_n \cdot \nabla \widetilde{\omega}_s + P_{\neq 0}(V_n \cdot \nabla \omega_n)\right) = g_1 + g_2 + g_3,$$

Then (f,g) solves (12.5). We denote

$$\begin{aligned} A_1(t) &= \nu^{\frac{1}{2}} \left(\int_0^t \|\nabla \omega_n(s)\|_{L^2}^2 ds \right)^{\frac{1}{2}} + \nu^{\frac{2}{3}} \int_0^t \|\partial_x \omega_n(s)\|_{L^2} ds + \nu^{\frac{\gamma_1}{2}} \left(\int_0^t \|V_n(s)\|_{L^{\infty}}^2 ds \right)^{\frac{1}{2}}, \\ A_2(t) &= \int_0^t \|g(s)\|_{L^2} ds \le \sum_{j=1}^3 \int_0^t \|g_j(s)\|_{L^2} ds. \end{aligned}$$

Then (12.7) implies that

(12.22) $A_1(t) \le C \|\omega_n(0)\|_{L^2} + CA_2(t).$

By (12.20), we have

(12.23)
$$\int_{0}^{t} \|g_{1}(s)\|_{L^{2}} ds = \int_{0}^{t} \|\widetilde{V}_{s}^{1}(s)\partial_{x}\omega_{n}(s)\|_{L^{2}} ds \leq \int_{0}^{t} \|\widetilde{V}_{s}^{1}(s)\|_{L^{\infty}} \|\partial_{x}\omega_{n}(s)\|_{L^{2}} ds \\ \leq C\nu^{\gamma} \int_{0}^{t} \|\partial_{x}\omega_{n}(s)\|_{L^{2}} ds \leq C\nu^{\gamma-\frac{2}{3}}A_{1}(t),$$

and by (12.21),

(12.24)
$$\int_{0}^{t} \|g_{2}(s)\|_{L^{2}} ds = \int_{0}^{t} \|V_{n}(s) \cdot \nabla \widetilde{\omega}_{s}(s)\|_{L^{2}} ds \leq \int_{0}^{t} \|V_{n}(s)\|_{L^{\infty}} \|\nabla \widetilde{\omega}_{s}(s)\|_{L^{2}} ds$$
$$\leq \left(\int_{0}^{t} \|V_{n}(s)\|_{L^{\infty}}^{2} ds\right)^{\frac{1}{2}} \left(\int_{0}^{t} \|\nabla \widetilde{\omega}_{s}(s)\|_{L^{2}}^{2} ds\right)^{\frac{1}{2}}$$
$$\leq \nu^{-\frac{\gamma_{1}}{2}} A_{1}(t) (C\nu^{2\gamma-1})^{\frac{1}{2}} \leq C\nu^{\gamma-\frac{\gamma_{1}+1}{2}} A_{1}(t).$$

Using (12.12) and the fact that $\nabla \omega_n(t) = P_{\neq 0} \nabla (I - P_2) \omega(t)$, we infer that

(12.25)
$$\int_{0}^{t} \|g_{3}(s)\|_{L^{2}} ds = \int_{0}^{t} \|P_{\neq 0}(V_{n} \cdot \nabla \omega_{n})\|_{L^{2}} ds \leq \int_{0}^{t} \|V_{n}(s)\|_{L^{\infty}} \|\nabla \omega_{n}(s)\|_{L^{2}} ds$$
$$\leq \left(\int_{0}^{t} \|V_{n}(s)\|_{L^{\infty}}^{2} ds\right)^{\frac{1}{2}} \left(\int_{0}^{t} \|\nabla (I - P_{2})\omega(s)\|_{L^{2}}^{2} ds\right)^{\frac{1}{2}}$$
$$\leq \nu^{-\frac{\gamma_{1}}{2}} A_{1}(t) (C\nu^{2\gamma-1})^{\frac{1}{2}} \leq C\nu^{\gamma-\frac{\gamma_{1}+1}{2}} A_{1}(t).$$

It follows from (12.22)-(12.25) that

$$A_1(t) \le C \|\omega_n(0)\|_{L^2} + C(\nu^{\gamma - \frac{2}{3}} + \nu^{\gamma - \frac{\gamma_1 + 1}{2}})A_1(t).$$

As $\gamma > \frac{2}{3}$, $\gamma_1 < 2\gamma - 1$, we can find $c_2 \in (0,1)$ so that $C(\nu^{\gamma-\frac{2}{3}} + \nu^{\gamma-\frac{\gamma_1+1}{2}}) \leq \frac{1}{2}$ for $0 < \nu < c_2$. Then we obtain

$$A_1(t) \le C \|\omega_n(0)\|_{L^2}, \quad A_2(t) \le C (\nu^{\gamma - \frac{2}{3}} + \nu^{\gamma - \frac{\gamma_1 + 1}{2}}) \|\omega_n(0)\|_{L^2}.$$

While, by (12.6), we have

(12.26)
$$\|\omega_n(t)\|_{L^2} \le K e^{-c_1 \sqrt{\nu} t} \|\omega_n(0)\|_{L^2} + C A_2(t).$$

Therefore, $\|\omega_n(t)\|_{L^2} \leq C \|\omega_n(0)\|_{L^2}$ for $0 < t < \min\{t_0, \tau/\nu\}$. If $t = t_0 < \tau/\nu$, then we have

$$\begin{aligned} \|\omega_n(t)\|_{L^2} &\leq C e^{-c_1 \sqrt{\nu t_0}} \|\omega_n(0)\|_{L^2} + C A_2(t) \\ &\leq \frac{1}{4} \|\omega_n(0)\|_{L^2} + C_{(\nu^{\gamma - \frac{2}{3}} + \nu^{\gamma - \frac{\gamma_1 + 1}{2}})} \|\omega_n(0)\|_{L^2} \end{aligned}$$

Take c_2 small enough (if necessary) so that $C(\nu^{\gamma-\frac{2}{3}} + \nu^{\gamma-\frac{\gamma_1+1}{2}}) \leq \frac{1}{4}$ for $0 < \nu < c_2$. Thus, $\|\omega_n(t)\|_{L^2} \leq \frac{1}{2} \|\omega_n(0)\|_{L^2}$ for $t = t_0 < \tau/\nu$.

This completes the proof of Theorem 1.6.

Acknowledgement

The authors thank Zhiwu Lin for profitable discussions. Z. Zhang is partially supported by NSF of China under Grant 11425103. W. Zhao is partially supported by China Postdoctoral Science Foundation under Grant 2017M610007.

References

- M. Beck and C. E. Wayne, Using global invariant manifolds to understand metastability in the Burgers equation with small viscosity, SIAM Rev., 53(2011), 129-153.
- [2] M. Beck and C. E. Wayne, Metastability and rapid convergence to quasi-stationary bar states for the two-dimensional Navier-Stokes equations, Proc. Roy. Soc. Edinburgh Sect. A, 143 (2013), 905-927.
- [3] J. Bedrossian, M. Coti Zelati, Enhanced dissipation, hypoellipticity, and anomalous small noise inviscid limits in shear flows, Arch. Ration. Mech. Anal., 224(2017), 1161-1204.
- [4] J. Bedrossian, P. Germain and N. Masmoudi, Dynamics near the subcritical transition of the 3D Couette flow I: Below threshold case, arXiv 1506.03720.
- [5] J. Bedrossian, P. Germain and N. Masmoudi, Dynamics near the subcritical transition of the 3D Couette flow II: Above threshold case, arXiv 1506.03721.
- [6] J. Bedrossian, P. Germain and N. Masmoudi, On the stability threshold for the 3D Couette flow in Sobolev regularity, Annals of Math., 185(2017), 541-608.
- [7] J. Bedrossian and N. Masmoudi, Inviscid damping and the asymptotic stability of planar shear flows in the 2D Euler equations, Publ. Math. Inst. Hautes Études Sci., 122(2015), 195-300.
- [8] J. Bedrossian, N. Masmoudi and V. Vicol, Enhanced dissipation and inviscid damping in the inviscid limit of the Navier-Stokes equations near the two dimensional Couette flow, Arch. Ration. Mech. Anal., 219(2016), 1087-1159.
- [9] J. Bedrossian, F. Wang and V. Vicol, The Sobolev stability threshold for 2D shear flows near Couette, arXiv:1604.01831.
- [10] F. Bouchet and H. Morita, Large time behavior and asymptotic stability of the 2D Euler and linearized Euler equations, Physica D, 239(2010), 948-966.
- [11] F. Bouchet and E. Simonnet, Random changes of flow topology in twodimensional and geophysical turbulence, Phys. Rev. Lett., 102 (2009), 094504.
- [12] K. M. Case, Stability of inviscid plane Couette flow, Phys. Fluids, 3(1960), 143-148.
- [13] P. Constantin, A. Kiselev, L. Ryzhik and A. Zlatos, Diffusion and mixing in fluid flow, Ann. of Math. (2), 168(2008), 643-674.
- [14] Y. Couder, Observation expérimental de la turbulence bidimensionnelle dans un film liquide mince, C. R. Acad. Sci. Paris Sèr. II, 297(1983), 641-645.
- [15] W. Deng, Pseudospectrum for Oseen vortices operators, Int. Math. Res. Not., IMRN 2013, 1935-1999.
- [16] T. Gallay and Y. Maekawa, Existence and stability of viscous vortices, arxiv:1610.08384.

- [17] T. Gallay, Enhanced dissipation and axisymmetrization of two-dimensional viscous votices, arXiv:1707.05525.
- [18] I. Gallagher, Th. Gallay and F. Nier, Spectral asymptotics for large skew-symmetric perturbations of the harmonic oscillator, Int. Math. Res. Notices, IMRN 2009, 2147-2199.
- [19] S. Ibrahim, Y. Maekawa and N. Masmoudi, On pseudospectral bound for non-selfadjoint operators and its application to stability of Kolmogorov flows, arXiv:1710.05132.
- [20] A. Kiselev and X. Xu, Suppression of chemotactic explosion by mixing, Arch. Ration. Mech. Anal., 222(2016), 1077-1112.
- [21] T. Li, D. Wei and Z. Zhang, *Pseudospectral and spectral bounds for the Oseen vortices operator*, arXiv:1701.06269.
- [22] Z. Lin and M. Xu, Metastability of Kolmogorov flows and inviscid damping of shear flows, arXiv:1707.00278.
- [23] Z. Lin and C. Zeng, Inviscid dynamic structures near Couette flow, Arch. Rat. Mech. Anal., 200(2011), 1075-1097.
- [24] Z. Lin and C. Zeng, Instability, index theorem, and exponential trichotomy for Linear Hamiltonian PDEs, arXiv:1703.04016.
- [25] Y. Maekawa, Spectral properties of the linearization at the Burgers vortex in the high rotation limit, J. Math. Fluid Mech., 13 (2011), 515-532.
- [26] W. H. Matthaeus, W. T. Stribling, D. Martinez, S. Oughton and D. Montgomery, Decaying, twodimensional, Navier-Stokes turbulence at very long times, Physica D, 51(1991), 531-538.
- [27] K. McQuighan, C. E. Wayne, An explanation of metastability in the viscous Burgers equation with periodic boundary conditions via a spectral analysis, SIAM J. Appl. Dyn. Syst., 15(2016), 1916-1961.
- [28] C. Mouhot and C. Villani, On Landau damping, Acta Math., 207(2011), 29-201.
- [29] S. I. Rosencrans and D. H. Sattinger, On the spectrum of an operator occurring in the theory of Hydrodynamics stability, J. Math. Phys., 45(1966), 289-300.
- [30] S. A. Stepin, Nonself-adjoint Friedrichs models in Hydrodynamics stability, Functional Analysis and its Applications, 29(1995), 91-101.
- [31] L. N. Trefethen, Pseudospectra of linear operators, SIAM Review, 39(1997), 383-406.
- [32] L. N. Trefethen, A. E. Trefethe, S. C. Reddy and T. A. Driscoll, Hydrodynamic stability without eigenvalues, Science, 261(5121), 578-584 (1993).
- [33] C. Villani, Hypocoercivity, Mem. Amer. Math. Soc., 202 (2009), no. 950.
- [34] D. Wei, Z. Zhang and W. Zhao, Linear inviscid damping for a class of momotone shear flow in Sobolev spaces, Comm. Pure Appl. Math., online.
- [35] D. Wei, Z. Zhang and W. Zhao, *Linear inviscid damping and vorticity depletion for shear flows*, arXiv:1704.00428.
- [36] C. Zillinger, Linear inviscid damping for monotone shear flows, Trans. Amer. Math. Soc., 369 (2017), 8799-8855.
- [37] C. Zillinger, Linear inviscid damping for monotone shear flows in a finite periodic channel, boundary effects, blow-up and critical Sobolev regularity, Arch. Ration. Mech. Anal., 221(2016), 1449-1509.

SCHOOL OF MATHEMATICAL SCIENCE, PEKING UNIVERSITY, 100871, BEIJING, P. R. CHINA *E-mail address*: jnwdyi@163.com

SCHOOL OF MATHEMATICAL SCIENCE, PEKING UNIVERSITY, 100871, BEIJING, P. R. CHINA *E-mail address:* zfzhang@math.pku.edu.cn

SCHOOL OF MATHEMATICAL SCIENCE, PEKING UNIVERSITY, 100871, BEIJING, P. R. CHINA *E-mail address:* zjzjzwr@126.com