

Reply to “Comment on ‘Strong Measurements Give a Better Direct Measurement of the Quantum Wave Function’ ”

Giuseppe Vallone¹ and Daniele Dequal²

¹Department of Information Engineering, University of Padova, I-35131 Padova, Italy and

²Italian Space Agency, ASI, Rome, Italy

In Ref. [1], we generalized the method introduced in Ref. [2] to obtain a direct measurement of a quantum wave-function. The method of Ref. [2] is based on weak measurements [3] to determine the wave-function. In Ref. [1] we showed that weak measurements are not necessary and indeed a better direct measurement (in term of precision and accuracy) of the quantum wave-function can be obtained by using strong measurements.

Recently, our work received a comment by V. R. Pande, S. Kanjilal and D. Das (PKD) [4] about a flaw our method. The argument of PKD can be summarized in one of their statements:

“we show that certain entities in Eq. (S6) of their supplemental materials on which the central claim of their result is based have no operational existence and are therefore not experimentally measurable.”

In this reply, we show that their claim is not correct and that the quantities appearing in our method can be indeed measured.

As correctly stated in [4], the probabilities needed for our method (namely $P_+^{(x)}$, $P_-^{(x)}$, $P_1^{(x)}$, $P_R^{(x)}$ and $P_L^{(x)}$, defined in the supplementary material of [1]) are calculated by using the unnormalized pointer wave-function $|\varphi\rangle_P$ (Eq. (S2)) such as

$$P_j^{(x)} = |\langle j|\varphi\rangle|^2, \quad (1)$$

with $j = +, -, 1, R, L$ and the pointer states defined as follows

$$\begin{aligned} |+\rangle_P &= \frac{1}{\sqrt{2}}(|0\rangle + |1\rangle), & |-\rangle_P &= \frac{1}{\sqrt{2}}(|0\rangle - |1\rangle) \\ |L\rangle_P &= \frac{1}{\sqrt{2}}(|0\rangle + i|1\rangle), & |R\rangle_P &= \frac{1}{\sqrt{2}}(|0\rangle - i|1\rangle). \end{aligned} \quad (2)$$

In the above equations, $\{|0\rangle, |1\rangle\}$ is a basis for the pointer Hilbert space. The unnormalized pointer state is defined as

$$|\varphi\rangle_P = {}_X\langle p_0|U_x(\theta)|\psi\rangle_X \otimes |0\rangle_P, \quad (3)$$

with $|0\rangle_P$ the initial state of the pointer, $|\psi\rangle_X$ the unknown state whose wave-function will be determined, $U_x(\theta)$ the interaction unitary and $|p_0\rangle_X = \frac{1}{\sqrt{d}}\sum_x |x\rangle$ the zero transverse momentum state. Since $|\varphi\rangle$ is not normalized, clearly $P_+^{(x)} + P_-^{(x)} = \langle\varphi|\varphi\rangle \neq 1$.

The mistake of PKD is to consider that the quantities $P_j^{(x)}$ should be measured on the post-selected quantum

ensemble and that they correspond to conditional probabilities (as in the original proposal [2]).

Indeed, as shown below, the quantities $P_j^{(x)}$ are not *conditional probabilities* but *joint probabilities*. By defining $|\Psi'\rangle = U_x(\theta)|\psi\rangle_X \otimes |0\rangle_P$ as the joint system-pointer state after the interaction, it is easy to show that the probabilities in eq. (1) can be written as

$$P_j^{(x)} = |\langle j|\otimes\langle p_0|\Psi'\rangle|^2 \equiv \text{Prob}_{\Psi'}(j, p_0). \quad (4)$$

The above equation shows that the quantities $P_j^{(x)}$ represent the *joint probabilities* of measuring the system in the state $|p_0\rangle$ and the pointer in the state $|j\rangle$ after the interaction. Such probabilities are well defined, can be measured and thus have “operational existence”.

On the other hand, probabilities measured on the post-selected quantum ensemble (after the post-selection on the state $|p_0\rangle$) are *conditional probabilities*, and indeed they can be obtained by Bayes’ rule as

$$\text{Prob}_{\Psi'}(j|p_0) = \frac{\text{Prob}_{\Psi'}(j, p_0)}{P(p_0)} = \frac{P_j^{(x)}}{\langle\varphi|\varphi\rangle}. \quad (5)$$

We note that the normalization factor $\langle\varphi|\varphi\rangle$ corresponds to the probability of post-selection, since

$$\langle\varphi|\varphi\rangle = \langle p_0|\text{Tr}_P[|\Psi'\rangle\langle\Psi'|]|p_0\rangle = P_{\text{post-selection}}. \quad (6)$$

The mistake of PKD is claiming that only the conditional probabilities $\text{Prob}_{|\Psi'\rangle}(j|p_0)$ can be measured while the probabilities that are “calculated from an unnormalized pointer state [...] cannot be measured experimentally”. As we have shown in eq. (4), the probabilities $P_j^{(x)}$ obtained from the unnormalized state $|\varphi\rangle_P$ are *joint probabilities* that can be easily measured in experiments. By using joint probabilities of eq. (4) it is possible to show that the wavefunction $\psi_x = \langle x|\psi\rangle$ can be expressed as

$$\psi_x = \frac{d}{2\tilde{\psi}\sin\theta} \left[P_+^{(x)} - P_-^{(x)} + 2P_1^{(x)} \tan\frac{\theta}{2} + i(P_L^{(x)} - P_R^{(x)}) \right] \quad (7)$$

with $\tilde{\psi} = \sum_x \psi_x$. In the above equation, the normalization factor $\frac{d}{2\tilde{\psi}\sin\theta}$ does not depend on x , and therefore it can be determined at the end of the procedure by imposing the overall normalization of the wave-function.

It is worth noticing that any experiment designed to measure the conditional probabilities is also able to measure joint conditional probabilities.

-
- [1] G. Vallone and D. Dequal, Physical Review Letters **116**, 040502 (2016).
- [2] J. S. Lundeen, B. Sutherland, A. Patel, C. Stewart, and C. Bamber, Nature **474**, 188 (2011).
- [3] Y. Aharonov, D. Z. Albert, and L. Vaidman, Phys. Rev. Lett. **60**, 1351 (1988).
- [4] V. R. Pande, S. Kanjilal, and D. Das, (2017), arXiv:1711.00764.