

RIEMANNIAN CURVATURE MEASURES

JOSEPH H.G. FU AND THOMAS WANNERER

ABSTRACT. A famous theorem of Weyl states that if M is a compact submanifold of euclidean space, then the volumes of small tubes about M are given by a polynomial in the radius r , with coefficients that are expressible as integrals of certain scalar invariants of the curvature tensor of M with respect to the induced metric. It is natural to interpret this phenomenon in terms of curvature measures and smooth valuations, in the sense of Alesker, canonically associated to the Riemannian structure of M . This perspective yields a fundamental new structure in Riemannian geometry, in the form of a certain abstract module over the polynomial algebra $\mathbb{R}[t]$ that reflects the behavior of Alesker multiplication. This module encodes a key piece of the array of kinematic formulas of any Riemannian manifold on which a group of isometries acts transitively on the sphere bundle. We illustrate this principle in precise terms in the case where M is a complex space form.

1. INTRODUCTION

The so-called Weyl tube formula states that if $M \hookrightarrow \mathbb{R}^N$ is a smooth isometric embedding of a compact smooth Riemannian manifold, then the volume of a tube around M of sufficiently small radius $r > 0$ is given by a polynomial of degree $\leq N$ in r , whose coefficients may be expressed as integrals of scalar invariants (Lipschitz-Killing curvatures) of the curvature tensor of M . Up to scale, the coefficient in degree N is the Euler characteristic $\chi(M)$. That this invariant admits such an expression is the Chern-Gauss-Bonnet theorem, proved by Chern [15, 16] by integrating certain canonical differential forms on M and its tangent sphere bundle SM , derived from the Cartan apparatus of curvature and connection forms. The coefficients of the lower degree terms arise similarly [14]. The tube coefficients may be localized by integrating these forms over subsets, and extended to subsets M more general than smooth submanifolds, yielding Federer's theory of curvature measures, formulated in terms of normal cycles in [28].

From a different perspective, the tube coefficients coincide up to scale with natural extensions of the *intrinsic volumes* μ_k of Hadwiger [23]. These are the principal examples of the concept of *convex valuation*. Alesker's recent work [1–7] introduces for any smooth manifold M the space $\mathcal{V}(M)$ of *smooth valuations on M* , equipped with a natural commutative product. With respect to this product, the intrinsic volumes then appear (up to scale) as powers of μ_1 . A smooth immersion $M \hookrightarrow N$ induces a restriction homomorphism $r_{MN} : \mathcal{V}(N) \rightarrow \mathcal{V}(M)$. Combining these facts with the Nash embedding theorem, and taking N to be euclidean space, Alesker observed that Weyl's theorem yields a canonical embedding $i_M : \mathbb{R}[t]/(t^{\dim M+1}) \hookrightarrow \mathcal{V}(M)$ associated to any smooth Riemannian M , with generator t identified with $\mu_1|_M$. This system of embeddings of algebras is obviously *reproductive* in the sense that if $P \hookrightarrow M$ is again an isometric immersion, then $i_P = r_{PM} \circ i_M$. The image of i_M is the *Lipschitz-Killing algebra* $\mathcal{LK}(M)$.

The fundamental impetus for the present paper is to understand these phenomena in purely valuation-theoretic terms, and without reference to existence of isometric immersions in euclidean spaces. We accomplish this by introducing a fundamental new structure in

Date: December 14, 2024.

Key words and phrases. Curvature measures, smooth valuations, Lipschitz-Killing curvatures.

JHGF was supported by NSF grant DMS-1406252. He wishes also to thank the Courant Institute at NYU, and the Technical University of Vienna, for their kind hospitality during the preparation of this paper. TW was supported by DFG grant WA 3510/1-1. The authors also wish to thank A. Bernig and G. Solanes for their many helpful comments.

Riemannian geometry, a further refinement of the general picture described above. In order to describe it we recall that any smooth valuation may be localized, albeit non-uniquely. The resulting space $\mathcal{C}(M)$ of *curvature measures on M* carries the structure of a module over the algebra $\mathcal{V}(M)$. This language is described in Section 2.1 below. We define the space \mathcal{R} of *Riemannian curvature measures*, abstracted from the space of summands of the tube coefficients described above. An element $\Psi \in \mathcal{R}$ is an object that assigns to any smooth Riemannian manifold M a concrete curvature measure Ψ^M on M , giving rise to a canonical subspace $\mathcal{R}(M) \subset \mathcal{C}(M)$. Our main results, given in Section 3, may be stated in general terms as follows. We describe the universal behavior of the elements of \mathcal{R} under isometric immersion of one Riemannian manifold into another (Theorem 3.7), and show that the Lipschitz-Killing curvatures are precisely the elements invariant under all such immersions (Theorem 3.11). This is accomplished via a natural identification (18) of \mathcal{R} with the space $\mathbb{R}[[\xi, \eta]]$ of bivariate formal power series. We then give an explicit description of an action of $\mathbb{R}[t]$ in these terms, reflecting the universal action of $\mathcal{LK}(M)$ on $\mathcal{R}(M)$ (Theorem 3.13).

The arguments in Section 3 correspond to the soft part of the *template method*, a familiar procedure in integral geometry: one proves the existence of a formula of a certain type (typically the hard part), then evaluates the constants by examining enough special cases, or templates (the soft part). Surprisingly, the cases of spheres of varying radius are enough. The technical heart of the paper is Section 4, giving the proofs of Lemmas 3.8 and 3.12 (the hard part). There we adapt the classical method of moving frames to display the geometric processes of Alesker multiplication by μ_1 , and of isometric immersion, in terms of formal models based on the Cartan apparatus. We lay the foundations for these models in Section 2.2.

Key to recent progress in integral geometry is the fact that if M admits a group G of isometries acting transitively on SM , then Alesker multiplication on the spaces $\mathcal{V}^G(M), \mathcal{C}^G(M)$ of G -invariant valuations and curvature measures is intimately related to the array of kinematic formulas for (M, G) ([10], Section 2). Since necessarily $\mathcal{R}(M) \subset \mathcal{C}^G(M)$, and $\mathcal{LK}(M) \subset \mathcal{V}^G(M)$, the structure studied here represents a universal component of any such array. The final Section 5 applies this observation to the case of complex projective (or hyperbolic) spaces $\mathbb{C}P^n$. Stabilizing by taking the dimension $n \rightarrow \infty$, it turns out that $\mathcal{R}(\mathbb{C}P^\infty)$ is a faithful copy of \mathcal{R} , and may be characterized as the space of all invariant curvature measures that enjoy the fundamental geometric property of *angularity*. This space admits a natural basis Δ_{kp} adapted to the complex structure, distinct from the natural basis for \mathcal{R} . In Theorem 5.4 we translate the formulas of Theorem 3.13 in terms of the Δ basis, by means of certain simple yet remarkable transforms \mathcal{O}, \mathcal{P} on the space of bivariate power series. These transforms already played a role in Section 3.3 of [10], although the relation between that appearance and this one remains mysterious.

2. PRELIMINARIES

2.1. Curvature measures and valuations. A more detailed account of the notions in the present section appears in Section 2.2 of [10].

Although the theory of smooth valuations on a manifold M is entirely independent of orientation or orientability, it will be convenient to assume that M is oriented. With this assumption we can frame the discussion in terms of integration, over the normal cycles associated to sufficiently regular subsets of M , of smooth differential forms on the cosphere bundle of M . If the orientation is switched to its opposite, then both the differential forms that we consider and the normal cycles change sign, so that the resulting set function is unchanged. If M is not oriented (even unorientable) it is possible to recast the whole theory in terms of differential forms twisted by the orientation line bundle of M , but, since the theory is essentially local in nature, one may alternatively avoid any loss of generality by working on the orientation double cover of M if needed.

Another simplification available in the present Riemannian context is to replace the cosphere bundle by the sphere bundle SM . Let $\dim M = m + 1$. In this language, we recall

that any closed submanifold with corners (which we refer to henceforth as a *simple smooth polyhedron*) $P \subset M$, admits a *normal cycle* $\text{nc}(P)$, a closed integral current of dimension m in the tangent sphere bundle SM . The current $\text{nc}(P)$ is *Legendrian*, in the sense that it annihilates any multiple of the contact (canonical) 1-form (and hence also any multiple of its exterior derivative). If M is a submanifold of N , we distinguish between the normal cycles with respect to the two different ambient spaces by the notations nc_M, nc_N . One may assign to each pair $(\psi, \phi) \in \Omega^m(SM) \oplus \Omega^{m+1}(M)$ the **curvature measure** Ψ that associates to such P the signed measure

$$(1) \quad \Psi(P, \cdot) := \pi_*(\text{nc}(P) \lrcorner \psi) + \int_{\cdot \cap P} \phi.$$

We denote this element Ψ by $[\psi, \phi]$. The space of all such curvature measures on M is denoted $\mathcal{C}(M)$. Such a pair (ψ, ϕ) determines also a **valuation** given by $\mu(P) := \int_{\text{nc}(P)} \psi + \int_P \phi$ for *compact* submanifolds with corners P . The space of all such set functions is denoted $\mathcal{V}(M)$, and the assignment $\Psi \mapsto \mu$ is the **globalization map**

$$\text{glob} : \mathcal{C}(M) \rightarrow \mathcal{V}(M).$$

The classical examples are the Federer curvature measures $\Phi_0, \dots, \Phi_{m+1} \in \mathcal{C}(\mathbb{R}^{m+1})$ [17, 28] and the corresponding intrinsic volumes $\mu_i := \text{glob} \Phi_i$. The domain of geometric shapes P subject to these set functions may be enlarged to the class of sets with positive reach, or still larger classes [21]. However, these extensions are irrelevant to the present paper, in which we emphasize the set functions Ψ, μ , and the P play the role of test objects.

If $\Psi \in \mathcal{C}(M)$, and P_0, P_1, P_2, \dots are closed submanifolds with corners such that $\text{nc}(P_i) \rightarrow c \text{nc}(P_0)$ in the flat metric topology, then the associated signed measures on M converge weakly, which we write as

$$(2) \quad \Psi(P_i, \cdot) \rightarrow c\Psi(P_0, \cdot)$$

It is natural to think of this process in terms of the specialization of a family of constructible functions [22]. In the simplest instance, the P_i are k -dimensional spheres of radii $r_i \rightarrow 0$ and common center x , and $P_0 = \{x\}$, in which case $c = 1 + (-1)^k$.

If $f : M \rightarrow N$ is a smooth immersion then there are restriction (or pullback) maps $\mathcal{C}(N) \rightarrow \mathcal{C}(M)$ and $\mathcal{V}(N) \rightarrow \mathcal{V}(M)$, both of which we denote by f^* , given by

$$(f^*\Psi)(P, E) := \Psi(f(P), f(E)), \quad (f^*\mu)(P) := \mu(f(P)).$$

Clearly these pullbacks maps commute with globalization. The point is that the pulled back objects may again be represented by differential forms on the domain manifolds M, SM . We carry this out explicitly in the special case of an isometric immersion of Riemannian manifolds in Proposition 4.7 below. We will also use the standard notation

$$(f_*m)(E) := m(f^{-1}(E))$$

for the pushforward of a (signed) measure m by a map f . Thus in the circumstances above

$$f_*((f^*\Psi)(A, \cdot))(E) = \Psi(f(A), E).$$

The main facts that we use in this paper from the theory of valuations are summarized in the following.

Theorem 2.1 ([1, 20]).

- (1) *The space $\mathcal{V}(M)$ admits a natural commutative multiplication (Alesker product), with the Euler characteristic χ acting as the multiplicative identity. Furthermore $\mathcal{V}(M)$ acts on $\mathcal{C}(M)$ in a natural way, compatible with the product of valuations, i.e. if $\mu \in \mathcal{V}(M), \Psi \in \mathcal{C}(M)$ then $\text{glob}(\mu \cdot \Psi) = \mu \cdot \text{glob}(\Psi)$. If f is a smooth immersion as above then f^* is an algebra and module homomorphism, i.e. if $\mu, \nu \in \mathcal{V}(N), \Psi \in \mathcal{C}(N)$ then*

$$(f^*\mu) \cdot (f^*\nu) = f^*(\mu \cdot \nu), \quad (f^*\mu) \cdot (f^*\Psi) = f^*(\mu \cdot \Psi).$$

- (2) Suppose $X \subset M$ is a compact simple smooth polyhedron, and $P \times M \rightarrow M$ is a smooth proper family of diffeomorphisms $\varphi_p : M \rightarrow M$, $p \in P$, equipped with a smooth measure dp . Suppose further that the map $P \times S^*M \rightarrow S^*M$, induced by the derivative maps $\varphi_{p*} : S^*M \rightarrow S^*M$, is a submersion. Then $\nu(A) := \int_P \chi(\varphi_p(X) \cap A) dp$ defines a smooth valuation $\nu \in \mathcal{V}(M)$. Given $\mu \in \mathcal{V}(M)$, $\Psi \in \mathcal{C}(M)$ we have

$$\begin{aligned} (\nu \cdot \mu)(A) &= \int_P \mu(\varphi_p(X) \cap A) dp, \\ (\nu \cdot \Psi)(A, E) &= \int_P \Psi(\varphi_p(X) \cap A, E) dp. \quad \square \end{aligned}$$

We will rely on an explicit construction of the Alesker product in terms of the underlying differential forms, originally due to Alesker and Bernig, and stated in Theorem 4.1 below.

2.1.1. *Angular curvature measures.* We recall briefly this concept from Section 2.5 of [10].

If V is an n -dimensional real affine space of dimension n , we denote by $\text{Curv}(V) \subset \mathcal{C}(V)$ the space of translation-invariant curvature measures on V . Any element of $\text{Curv}(V)$ may be expressed as $[\psi, \phi]$ where both ψ, ϕ are translation-invariant. The space $\text{Curv}(V)$ is graded by degree $k \in \{0, \dots, n\}$.

Let us now suppose further that V is a euclidean space. A curvature measure $\Psi \in \text{Curv}(V)$ is *angular* if there exists a function c_Ψ on the k -Grassmannian of V such that for any convex polytope P

$$(3) \quad \Psi(P, \cdot) = \sum_{k=0}^n \sum_{F \in \mathfrak{F}_k(P)} c_\Psi(\vec{F}) \angle(F, P) \mathcal{H}_k|_F$$

where $\mathfrak{F}_k(P)$ is the set of all k -faces of P , $\angle(F, P)$ is the exterior solid angle of P along F , and \vec{F} is the element of the k -Grassmannian parallel to F . Any such Ψ is clearly translation-invariant. Any element of $\text{Curv}(V)$ of degree $\geq n-1$ is vacuously angular.

Passing now to a Riemannian manifold, Section 2.2.2 of [10] describes a canonical isomorphism $\mathcal{C}(M) \leftrightarrow \Gamma(\text{Curv}(TM))$, the space of smooth sections of the bundle over M whose fiber over x is the space of translation-invariant curvature measures in $T_x M$. Thus $\mathcal{C}(M)$ inherits a natural grading. A curvature measure $\Psi \in \mathcal{C}(M)$ is said to be angular iff each value of the associated section has this property.

2.2. **The formal Cartan apparatus.** We give a brief but explicit account of a formal algebraic model for the calculus of moving frames on an oriented Riemannian manifold, implicitly used in classical Riemannian geometry, notably the work of Chern [15, 16]. In these and related calculations we adopt the usual summation convention.

2.2.1. $\mathcal{A}_m, \bar{\mathcal{A}}_m, \bar{\mathcal{A}}_m^+, \mathcal{B}_m, \mathcal{B}_m^+$, and their realizations. Let \mathcal{A}_m denote the anticommutative graded algebra with generators

$$\theta_i, \quad \omega_{ij} = -\omega_{ji}, \quad \Omega_{ij} = -\Omega_{ji}, \quad 0 \leq i, j \leq m,$$

of respective degrees 1, 1, and 2, equipped with the left action of the group $O(m+1) \ni g$ given by

$$(4) \quad \begin{aligned} L'_g \theta_i &:= g_{ji} \theta_j \\ L'_g \Omega_{ij} &:= g_{ki} g_{lj} \Omega_{kl} \\ L'_g \omega_{ij} &:= g_{ki} g_{lj} \omega_{kl} \equiv g_{0i} g_{lj} \omega_{0l} + g_{ki} g_{0j} \omega_{k0} \pmod{\langle \omega_{ij} : 0 \notin \{i, j\} \rangle}. \end{aligned}$$

This algebra admits the formal differential

$$(5) \quad \begin{aligned} d' \theta_i &:= -\omega_{ij} \theta_j \\ d' \omega_{ij} &:= -\omega_{ik} \omega_{kj} + \Omega_{ij} \\ d' \Omega_{ij} &:= \Omega_{ik} \omega_{kj} - \omega_{ik} \Omega_{kj}, \end{aligned}$$

modeled on the Cartan structure equations, where we follow the conventions of [13], p. 100. Note that $d'^2 \neq 0$, since we have not imposed relations corresponding to the Bianchi identity, but this property is unnecessary for our purposes.

Put $\bar{\mathcal{A}}_m$ for the quotient obtained by setting all $\omega_{ij} = 0$ unless $0 \in \{i, j\}$. There is an obvious inclusion map $\bar{\mathcal{A}}_m \hookrightarrow \mathcal{A}_m$. The restriction to $\bar{\mathcal{A}}_m$ of the action of $h \in O(m)$ by fixing the 0 coordinate, denoted L_h , descends to $\bar{\mathcal{A}}_m$ and intertwines the inclusion and quotient maps. We may define the differential d on $\bar{\mathcal{A}}_m$, and also endomorphisms $\tilde{L}_g, g \in O(m+1)$, by precomposing d', L' with the inclusion and postcomposing with the quotient map. Clearly

$$(6) \quad \tilde{L}_{gh} = \tilde{L}_g \circ L_h, \quad \tilde{L}_{hg} = L_h \circ \tilde{L}_g, \quad d \circ L_h = L_h \circ d$$

for any $h \in O(m)$. Put $\bar{\mathcal{A}}_m^+ \subset \bar{\mathcal{A}}_m$ for the subspace of all $\phi \in \bar{\mathcal{A}}_m$ such that $L_g \phi = (\det g)\phi$ for $g \in O(m)$. For $0 \leq k \leq m, 0 \leq 2p \leq k$, we define the elements

$$(7) \quad \bar{\mathcal{A}}_m^+ \ni \phi_{kp} := \sum_{\pi} \operatorname{sgn}(\pi) \Omega_{\pi_1 \pi_2} \cdots \Omega_{\pi_{2p-1} \pi_{2p}} \theta_{\pi_{2p+1}} \cdots \theta_{\pi_k} \omega_{\pi_{k+1}, 0} \cdots \omega_{\pi_m, 0}$$

where the sum extends over all permutations of $\{1, \dots, m\}$.

Put \mathcal{B}_m for the further quotient of $\bar{\mathcal{A}}_m$ obtained by setting all remaining $\omega_{ij} = 0$. Now the action of $O(m+1)$ descends. Put \mathcal{B}_m^+ for the subspace of all $\psi \in \mathcal{B}_m$ such that $L_g \psi = (\det g)\psi, g \in O(m+1)$. Define

$$(8) \quad \mathcal{B}_m^+ \ni \psi_p := \sum_{\pi} \operatorname{sgn}(\pi) \Omega_{\pi_0 \pi_1} \cdots \Omega_{\pi_{2p-2} \pi_{2p-1}} \theta_{\pi_{2p}} \cdots \theta_{\pi_m}$$

where the sum extends over all permutations of $\{0, \dots, m\}$.

Proposition 2.2. *Modulo θ_0 , the space of elements of formal degree m in $\bar{\mathcal{A}}_m^+$ is spanned by the ϕ_{kp} . The space of elements of formal degree $m+1$ in \mathcal{B}_m^+ is spanned by the ψ_p .*

Proof. We prove the first assertion. The proof of the second is similar and simpler. Given an $O(m)$ module M , we refer to the submodule on which the group acts by multiplication by the determinant as the *determinant submodule*.

The subspace of $\bar{\mathcal{A}}_m^+$ of elements of degree m decomposes by the degree p in the Ω and the degree $k-2p$ in the θ . We wish to show that, modulo θ_0 , the subspace \mathcal{C}_{kp} thus described is spanned by ϕ_{kp} . Put for simplicity $V := \mathbb{R}^m$. Let $O(m)$ act on V in the standard way, and on $\mathfrak{so}(m), \mathfrak{gl}(m)$ by conjugation. Clearly $\mathcal{C}_{kp}/(\theta_0)$ is isomorphic to the $O(m)$ -submodule of the determinant submodule of $(\mathfrak{so}(m)^{\otimes p} \otimes V^{\otimes(k-2p)} \otimes V^{\otimes(m-k)})^*$ that consists of elements that are symmetric in the $\mathfrak{so}(m)$ factors and antisymmetric in each of the two clusters of V factors. Pulling back via the projection $\mathfrak{gl}(m) \rightarrow \mathfrak{so}(m), M \mapsto (M-M^t)/2$, in fact \mathcal{C}_{kp} is isomorphic to a submodule of the corresponding submodule of $(\mathfrak{gl}(m)^{\otimes p} \otimes V^{\otimes(k-2p)} \otimes V^{\otimes(m-k)})^* \simeq V^{\otimes m^*}$.

The First Fundamental Theorem of invariant theory for $SO(m)$ ([25], Chapter 11.2) states that the algebra of such invariants in variables $y_i \in V$ is generated by functions of the form $(y_1, \dots, y_m) \mapsto \det[y_1 \dots y_m]$ and $\langle y_i, y_j \rangle$. In our case the degree is m , so if the latter appear in any term then all other factors must have the same form. Since these expressions are $O(m)$ -invariant, this cannot happen. Thus the determinant submodule is spanned by $y_1 \otimes \cdots \otimes y_m \mapsto \det[y_1 \dots y_m]$. This corresponds to $\phi_{kp} \in \mathcal{C}_{kp}$ under the identifications above. \square

2.2.2. Realization maps. Now let M be a smooth oriented Riemannian manifold of dimension $m+1$, with frame bundle $\hat{\pi} : FM \rightarrow M$ and tangent sphere bundle $\pi : SM \rightarrow M$. Thus the total space FM consists of all orthonormal frames $b = (b_0, \dots, b_m)$ for $T_x M, x = \hat{\pi}(b) \in M$. We may regard FM also as a bundle $z : FM \rightarrow SM$ by taking $z(b_0, \dots, b_m) := b_0$. The bundles $\hat{\pi}, z$ are principal bundles, with groups $O(m+1), O(m)$ respectively. We denote the right action in each case by R .

There is a well-defined *realization map* $\rho : \mathcal{A}_m \rightarrow \Omega^*(FM)$ given by taking the generators of \mathcal{A}_m to the coframe forms, connection forms, and curvature forms associated to a given $b \in FM$ [13, 15]. The Cartan structure equations state that this map intertwines the formal

differential above and the exterior differential on $\Omega^*(FM)$. It also intertwines the formal left action L' with the action on $\Omega^*(FM)$ induced by pullback under the right action of the structure group.

If P is a smooth manifold, with $\bar{b} : P \rightarrow FM$ a smooth map (typically a section of a bundle with total space FM), and $g \in O(m+1)$, then

$$(9) \quad (R_g \circ \bar{b})^* \circ \rho = \bar{b}^* \circ \rho \circ L'_g.$$

Definition 2.3. Given $b = (b_0 = \xi_0, b_1, \dots, b_m) \in FM$ we define $\bar{\rho}_b : \bar{\mathcal{A}}_m \rightarrow \Lambda^* T_{\xi_0}(SM)$ as follows. Let \bar{b} be a local section of the bundle z such that $\bar{b}(\xi_0) = b$. Then $\bar{\rho}_b$ is the restriction to $\bar{\mathcal{A}}_m$ of the evaluation at ξ_0 of $\bar{b}^* \circ \rho$.

Define $\bar{\rho} = \bar{\rho}_M : \bar{\mathcal{A}}_m^+ \rightarrow \Omega^*(SM)$ by setting its value at any given $\xi_0 \in SM$ by restricting $\bar{\rho}_b$ to $\bar{\mathcal{A}}_m^+$, where $b = (b_0 = \xi_0, b_1, \dots, b_m)$ is any positive adapted frame.

Abusing notation, we use the same symbol $\bar{\rho}$ to denote the corresponding maps $\mathcal{B}, \mathcal{B}^+ \rightarrow \Omega^*(M)$.

Lemma 2.4. *The maps $\bar{\rho}_b, \bar{\rho}$ are well-defined. If $g \in O(m)$ then $\bar{\rho}_{R_g b} = \bar{\rho}_b \circ L_g$.*

Proof. We must show that the map $\bar{\rho}_b$ is independent of the choice of the extension \bar{b} . This is clearly true of the values $\bar{\rho}_b(\theta_i), \bar{\rho}_b(\Omega_{ij})$. As for $\rho_b(\omega_{0i})$, note first that $\bar{\rho}_b(\theta_0)$ is the canonical contact 1-form α of SM , and hence independent even of the other vectors b_1, \dots, b_m of the frame. Now if $\tilde{b}_i \in T_{\xi_0}SM$ is the horizontal lift of $b_i \in T_{\pi(\xi_0)}M$ then $\bar{\rho}_b(\omega_{0i})$ is the interior product of \tilde{b}_i with $d\alpha$.

The final conclusion follows from (9), and implies in turn that $\bar{\rho}$ is well defined. \square

2.2.3. *The infinitesimally parallel extension of a frame and the canonical connection of z .* It is well known that any frame $b, z(b) = \xi_0$, may be extended to a local section \bar{b} of z such that $\bar{b}^* \circ \rho(\omega_{ij})|_{\xi_0} = 0$ if $0 \notin \{i, j\}$. Let us introduce an explicit terminology for such an extension for later use. Put $x_0 := \pi(\xi_0) \in M$, and let $V \subset M$ be a normal neighborhood of x_0 . Put $E : V \times S_{x_0}M \rightarrow SM$ to be the map induced by parallel translation along geodesics from x_0 . The **infinitesimally parallel extension** of b is the adapted moving frame \bar{b} defined on the neighborhood $\tilde{V}_{\xi_0} := E(V \times (S_{x_0}M \setminus \{-\xi_0\}))$ of ξ_0 , constructed as follows. For $\xi \in S_{x_0}M \setminus \{-\xi_0\}$, define

$$\text{rot}_{\xi}^{\xi_0} \in SO(T_{x_0}M) \quad \text{by } \text{rot}_{\xi}^{\xi_0}(\xi_0) = \xi, \quad \text{rot}_{\xi}^{\xi_0}|_{\xi^\perp \cap \xi_0^\perp} = \text{identity}.$$

Now set $\bar{b}(\xi) := \text{rot}_{\xi}^{\xi_0}(b) := (\text{rot}_{\xi}^{\xi_0}(b_0), \dots, \text{rot}_{\xi}^{\xi_0}(b_m))$ for $\xi \in S_{x_0}M \setminus \{-\xi_0\}$. In other words, we extend the frame initially to $S_{x_0} \setminus \{-\xi_0\}$ by parallel translation along great circles from ξ_0 . Now put

$$\bar{b}(E(x, \xi)) := E(x, \bar{b}(\xi)) := (E(x, \bar{b}_0(\xi)), \dots, E(x, \bar{b}_m(\xi))).$$

The map $b \mapsto \bar{b}$ is clearly compatible with the right $O(m)$ action on $z : FM \rightarrow SM$, i.e.

$$(10) \quad \overline{(R_h b)} = R_h \bar{b}, \quad h \in O(m).$$

Therefore the images of the derivatives $\bar{b}_* : T_{\xi}SM \rightarrow T_b FM$ corresponding to the infinitesimally parallel extensions of all adapted frames b at all points $\xi \in SM$ constitute the family of horizontal subspaces of a **canonical connection** on the principal $O(m)$ bundle z .

Consider the standard sphere $S^m \subset \mathbb{R}^{m+1}$, with standard coordinates u_0, \dots, u_m and standard basis vectors e_0, \dots, e_m . For $-e_0 \neq u \in S^m$, put $\overline{\text{rot}}_u \in SO(m+1)$ for the element that acts as the identity on vectors perpendicular to e_0 and u , and such that $\overline{\text{rot}}_u(e_0) = u$. Define $\psi_b : S_{x_0}M \rightarrow S^m$ by $\psi_b(u_i b_i) := (u_0, \dots, u_m)$. We will need the following obvious facts.

Lemma 2.5. *For all $\xi \neq -\xi_0$,*

$$(11) \quad \overline{\text{rot}}_{\psi_b(\xi)} = \psi_b \circ \text{rot}_{\xi}^{\xi_0} \circ \psi_b^{-1}.$$

If $u \not\perp e_0$, then the map $v \mapsto \overline{\text{rot}}_v(u)$ yields a diffeomorphism between small neighborhoods of e_0, u .

2.2.4. *The virtual antipodal map.* Let $a : SM \rightarrow SM$ denote the fiberwise antipodal map. Clearly

$$(12) \quad \bar{\rho} \circ L_{-1} = a^* \circ \bar{\rho}.$$

2.2.5. *The immersed version.* We will also need a version of this apparatus that describes in formal terms an isometric immersion of one Riemannian manifold into another. Let $n > m$ be given. The coordinates in the range $0, \dots, m$ will be treated in an essentially different way from the range $m+1, \dots, n$; we denote the former by Roman letters i, j, \dots from the first half of the alphabet and the latter by Greek letters α, β, \dots . Roman letters r, s, t, \dots from the second half of the alphabet will take values from the entire range $0, 1, \dots, n$. Put $\mathcal{A}_{m,n} := \mathcal{A}_n / (\theta_\alpha = 0)$. Introduce new symbols

$$\bar{\Omega}_{ij} := \omega_{i\alpha} \omega_{j\alpha}, \quad 0 \leq i, j \leq m.$$

Consider the standardly embedded subgroup $O(m+1) \times O(n-m) \subset O(n+1)$, where the two factors act respectively on the first $m+1$ and last $n-m$ coordinates. The restriction $L_{g,h}$ of the left action of $O(n+1)$ descends to this quotient.

Define the quotient $\bar{\mathcal{A}}_{m,n}$ by setting

$$(13) \quad \begin{aligned} \omega_{ij} &= 0 \quad \text{unless } 0 \in \{i, j\}, \\ \omega_{\alpha\beta} &= 0. \end{aligned}$$

There is a natural inclusion into $\mathcal{A}_{m,n}$. The group $O(m) \times O(n-m)$ acts by fixing the 0 coordinate. Put

$$\bar{\mathcal{A}}_{m,n}^{++} \text{ (resp. } \bar{\mathcal{A}}_{m,n}^+) := \{\phi : L_{g,h}\phi = (\det g)(\det h)\phi \text{ (resp. } (\det g)\phi) \text{ for } (g, h) \in O(m) \times O(n-m)\},$$

so that the quotient map takes $\bar{\mathcal{A}}_n^+ \rightarrow \bar{\mathcal{A}}_{m,n}^{++}$.

Put $\bar{\bar{\mathcal{A}}}_{m,n}$ to be the subspace $\Omega_{\alpha\beta} = \Omega_{\alpha i} = 0$. The $O(m) \times O(n-m)$ action descends to this space. Put $\bar{\bar{\mathcal{A}}}_{m,n}^+ = \{\phi \in \bar{\bar{\mathcal{A}}}_{m,n} : L_{g,h}\phi = (\det g)\phi, (g, h) \in O(m) \times O(n-m)\}$. The elements of degree m

$$\phi_{kpl} := \sum_{\pi} \text{sgn}(\pi) \Omega_{\pi_1 \pi_2} \cdots \Omega_{\pi_{2p-1} \pi_{2p}} \bar{\Omega}_{\pi_{2p+1} \pi_{2p+2}} \cdots \bar{\Omega}_{\pi_{2p+2l-1} \pi_{2p+2l}} \theta_{\pi_{2p+2l+1}} \cdots \theta_{\pi_k} \omega_{\pi_{k+1}, 0} \cdots \omega_{\pi_m, 0}$$

belong to this space.

Define the quotient $\mathcal{B}_{m,n}$ by setting all the remaining ω_{ij} , and the $\omega_{0\alpha}$, to zero as well. Thus the only remaining formal connection forms are the $\omega_{i\alpha}, i \neq 0$. The group $O(m+1) \times O(n-m)$ acts on this space. Put $\mathcal{B}_{m,n}^+$ for the subspace of all elements ϕ such that $L_{(g,h)}\phi = (\det g)\phi$, in particular the elements of degree $m+1$

$$\psi_{pl} = \sum_{\pi} \text{sgn}(\pi) \theta_{\pi_0} \Omega_{\pi_1 \pi_2} \cdots \Omega_{\pi_{2p-1} \pi_{2p}} \bar{\Omega}_{\pi_{2p+1} \pi_{2p+2}} \cdots \bar{\Omega}_{\pi_{2p+2l-1} \pi_{2p+2l}} \theta_{\pi_{2p+2l+1}} \cdots \theta_{\pi_m}$$

are of this type.

Proposition 2.6. *Modulo θ_0 , the space of elements of formal degree m in $\bar{\bar{\mathcal{A}}}_{m,n}^+$ is spanned by the ϕ_{kpl} . The space of elements of formal degree $m+1$ in $\mathcal{B}_{m,n}^+$ is spanned by the ψ_{pl} .*

Proof. Put $\mathcal{N} \subset \bar{\bar{\mathcal{A}}}_{m,n}$ for the subalgebra generated by the $\omega_{i\alpha}$. Clearly $\bar{\bar{\mathcal{A}}}_{m,n} \simeq \bar{\mathcal{A}}_m \otimes \mathcal{N}$, under the convention that $a \otimes b \cdot c \otimes d = (-1)^{(\deg b)(\deg c)} ac \otimes bd$. The groups $SO(m)$ and $O(n-m)$ act on the two factors separately. Thus any $\omega \in \bar{\bar{\mathcal{A}}}_{m,n}^+$ may be expressed in the form $\omega = \sum \mu \otimes \nu$, where the $\nu \in \mathcal{N}$ and are $O(n-m)$ -invariant. Abbreviating $W := \mathbb{R}^{n-m}$, with coordinates indexed by $\alpha = m+1, \dots, n$, the space of all such ν is isomorphic to the space S of $O(n-m)$ -invariant elements of $\Lambda^*(W^{m+1})^*$, where the W factors are indexed by $i = 0, \dots, m$. The First Fundamental Theorem implies that S is generated by the $\bar{\Omega}_{ij} = \omega_{i\alpha} \omega_{j\alpha}$.

Thus, as in the proof of Proposition 2.2, the degree m subspace of $\bar{\mathcal{A}}_{m,n}^+(\theta_0)$ is isomorphic as an $O(m)$ -module to the direct sum over k, p, l of submodules of the determinant submodules of $(\mathfrak{so}(m)^{\otimes p} \otimes \mathfrak{so}(m)^{\otimes l} \otimes V^{\otimes k-2p-2l} \otimes V^{\otimes m-k})^*$, where the additional $\mathfrak{so}(m)$ factors correspond to the $\bar{\Omega}_{ij}$. The proof now concludes in similar fashion to that of Proposition 2.2, where the invariant element that emerges now corresponds to ϕ_{kpl} . \square

These algebras may be realized on an isometric immersion $e : M^{m+1} \hookrightarrow N^{n+1}$. Put $F(M, N) \subset FN|_M$ for the subbundle of adapted frames $b = (b_0, \dots, b_n)$, i.e. frames such that $b_0, \dots, b_m \in T_{\hat{\pi}(b)}M$. This is a principal $O(m+1) \times O(n-m)$ bundle over M . The realization map $\rho : \mathcal{A}_n \rightarrow \Omega^*(FN)$ restricts to give an $O(m+1) \times O(n-m)$ -equivariant map $\rho_{rel} : \mathcal{A}_{m,n} \rightarrow \Omega^*(F(M, N))$. The Gauss equation becomes

$$(14) \quad \rho_{rel}(\bar{\Omega}_{ij}) = \Omega_{ij}^M - \Omega_{ij}^N,$$

where these two terms refer to the curvature forms of M, N respectively.

Given a fixed frame b , we likewise have a map $\bar{\rho}_{b,rel} : \bar{\mathcal{A}}_{m,n} \rightarrow \wedge^* T_{z(b)}(SM)$ since the second fundamental form is well defined. The restriction of z makes $F(M, N)$ into a principal bundle over SM with group $O(m) \times O(n-m)$. By restricting to frames b such that b_0, \dots, b_m is a positive frame for $T_{\hat{\pi}(b)}M$, we obtain maps $\bar{\rho}_{rel} : \bar{\mathcal{A}}_{m,n}^+ \rightarrow \Omega^*(SM), \mathcal{B}_{m,n}^+ \rightarrow \Omega^*(M)$ as above. Taking b_{m+1}, \dots, b_n also to be positive we obtain a map $\bar{\mathcal{A}}_{m,n}^{++} \rightarrow \Omega^*(SM)$, again denoted $\bar{\rho}_{rel}$.

2.3. Fiber integration in a fiber bundle with a connection. Let $q : P \rightarrow X$ be a smooth fiber bundle with model fiber F and group G . Suppose further that q is equipped with a G -connection, with horizontal subspaces $H_p \subset T_pP$. Thus the derivative q_* induces isomorphisms $H_p \rightarrow T_{q(p)}X$. For $x \in X$, denote by $F_x \subset P$ the fiber over x . Given $p \in F_x$, we define the evaluation map

$$\text{ev}_p : \Omega^*(P) \rightarrow \wedge^* T_p F_x \otimes \wedge^* T_x X$$

by identifying $T_pP = H_p \oplus T_p F_x \simeq T_x X \oplus T_p F_x$ via the connection. For given $x \in X$, consider the pullback of the bundle of exterior algebras $\wedge^* TP$ over P under the inclusion map $F_x \hookrightarrow P$. This gives rise to a *restriction map*

$$r_x : \Omega^*(P) \rightarrow \Omega^*(F_x, \wedge^* T_x X)$$

by taking the value of $r_x(\phi)|_p$ at $p \in F_x$ to be $\text{ev}_p(\phi)$.

Lemma 2.7. *Let C be a G -invariant integral current of dimension k in F , and for each $x \in X$ let C_x be the corresponding current in the fiber F_x . Then the natural fiber integration operator $p_{C*} : \Omega^*(P) \rightarrow \Omega^*(X)$ of degree $-k$ is given by*

$$(p_{C*}\phi)|_x = \int_{C_x} r_x(\phi) \in \wedge^* T_x X.$$

Proof. Given $x \in X$, by parallel translation we may find a neighborhood $U \subset X$ of x and a local trivialization $\tau : U \times F \rightarrow P$ such that $D\tau_{x,f}(T_x X \oplus \{0\}) = H_{\tau(x,f)}$ for all $f \in F$. Thus, given $\phi \in \Omega^*(P)$ we have $\tau^*(\phi)|_{x,f} = \text{ev}_{\tau(x,f)} \phi$. The Lemma now follows directly from the definition of fiber integration. \square

2.4. Generating functions. We will make extensive use of the Maclaurin expansions

$$(15) \quad (1-x)^{-(k+1)} = \sum_{j=0}^{\infty} \binom{k+j}{j} x^j.$$

The special cases $k = \pm\frac{1}{2}$ admit the alternate expressions

$$(16) \quad (1-x)^{-\frac{1}{2}} = \sum_{j=0}^{\infty} \binom{2j}{j} \left(\frac{x}{4}\right)^j,$$

$$(17) \quad (1-x)^{-\frac{3}{2}} = \sum_{j=0}^{\infty} (2j+1) \binom{2j}{j} \left(\frac{x}{4}\right)^j.$$

3. RIEMANNIAN CURVATURE MEASURES

In this section we introduce the space \mathcal{R} of *Riemannian curvature measures*, given as the vector space of all possibly infinite linear combinations of certain objects $\{C_{kp}\}_{0 \leq k, p \in \mathbb{Z}, 2p \leq k}$. We will describe this situation briefly by saying that \mathcal{R} is the ω -span of the C_{kp} , and refer similarly to ω -linear maps, etc. Formally, an element of \mathcal{R} is a universal object Ψ that associates to every smooth Riemannian manifold M a curvature measure $\Psi^M \in \mathcal{C}(M)$ —the *realization* of Ψ on M —that is expressible in a universal way in terms of the Cartan apparatus. Taking M to be a euclidean space, the Federer curvature measures Φ_k are included. If $k > \dim M$ then C_{kp}^M is the zero curvature measure. We normalize the C_{kp} so as to minimize the role played by the dimension of M . In particular, they are invariant under totally geodesic isometric embeddings (cf. Corollary 3.9 below). The *Lipschitz-Killing curvature measures* Λ_k , defined in Section 3.4 below, are distinguished elements of \mathcal{R} . We denote their ω -span by $\widetilde{\mathcal{L}\mathcal{K}}$.

3.1. Definition and basic properties.

Definition 3.1. For $2p \leq k$, define C_{kp} to be the object that assigns to any smooth Riemannian manifold M the curvature measure

$$\iota_M(C_{kp}) := C_{kp}^M := \begin{cases} \frac{\omega_k}{\pi^{k(m-k+1)! \omega_{n-k+1}}} [\bar{\rho}_M(\phi_{kp}), 0] & \text{if } k < \dim M \\ \frac{\omega_k}{\pi^k} [0, \bar{\rho}_M(\psi_p)] & \text{if } k = \dim M \\ 0 & \text{if } k > \dim M. \end{cases}$$

Here ω_k denotes the volume of the unit ball in \mathbb{R}^k . The globalization of C_{21}^M coincides, up to scale, with the valuation first described in Theorem 2 of [12].

The ω -span of the C_{kp} is denoted by \mathcal{R} . The space \mathcal{R} is ω -graded, by setting $\deg(C_{kp}) = k$. Given any smooth Riemannian M , the realization map $\iota_M : \mathcal{R} \rightarrow \mathcal{C}(M)$ is an ω -graded map of degree zero. We denote the image of the realization map by $\mathcal{R}(M) := \iota_M(\mathcal{R}) \subset \mathcal{C}(M)$.

The normalizing constants in the definition of the C_{kp} are chosen so that

- (1) the C_{kp} are invariant under totally geodesic embeddings (Corollary 3.9);
- (2) $C_{kp}^M = \lambda^p C_{k0}^M$ if M is a sphere of curvature λ ;
- (3) if V is a euclidean space then $C_{k,0}^V = \frac{k! \omega_k}{\pi^k} \Phi_k$, where Φ_k is the k th Federer curvature measure [17].

The constants reflect also the behavior of Alesker multiplication, as will be evident. From this perspective it is convenient to introduce the ω -linear isomorphism

$$(18) \quad \alpha : \mathcal{R} \rightarrow \mathbb{R}[[\xi, \eta]], \quad C_{kp} \mapsto \xi^{k-2p} \eta^p.$$

Proposition 3.2. *For every k, p, M , the curvature measure $C_{kp}^M \in \mathcal{C}(M)$ is angular.*

Proof. Given a point $x \in M$, let $\Psi \in \text{Curv}(T_x M)$ denote the value at x of the section of $\Gamma(\text{Curv}(TM))$ that corresponds to C_{kp}^M under the transfer map described in Section 2.2.2 of [10]. Let $P \subset T_x M$ be a convex polytope and $F \subset P$ a face of dimension k . Choosing moving frames so as to include some fixed orthonormal frame e_1, e_2, \dots, e_k for \vec{F} , the expression (7) implies that the coefficient $c_\Psi(\vec{F})$ from (3) is equal to a constant multiple of $\sum_{\pi} \int_{B_F} \text{sgn } \pi \Omega_{\pi_1 \pi_2} \dots \Omega_{\pi_{2p-1} \pi_{2p}} \theta_{\pi_{2p+1}} \dots \theta_{\pi_k}$, where π ranges over all permutations of $1, \dots, k$ and B_F is the unit ball in \vec{F} . \square

3.2. Some facts from spherical integral geometry. Our main calculations all follow ultimately from a few simple yet fundamental observations about the integral geometry of spheres. This discussion also represents the simplest instance of the apparatus just described. We denote the standard $(n+1)$ -dimensional sphere of curvature λ (or radius $\lambda^{-\frac{1}{2}}$) by S_λ^{n+1} , $\lambda > 0$.

Proposition 3.3 ([18], 0.4.3). *Modulo the canonical 1-form and its differential, the space of all differential forms on the sphere bundle of S_λ^{n+1} that are invariant under the usual action of $SO(n+2)$ is spanned by the $\bar{\rho}(\phi_{k,0}), k = 0, \dots, n$.*

Proof. Cf. pp. 184–5 of [18]. □

Up to scale, these forms correspond to the elements $C_{k,0}^{S_\lambda^{n+1}} \in \mathcal{R}(S_\lambda^{n+1}) \subset \mathcal{C}(S_\lambda^{n+1})$, $k \leq n$, to which we append the rescaled volume curvature measure $C_{n+1,0}^{S_\lambda^{n+1}}$. Since $\Omega_{ij} = \lambda \theta_i \wedge \theta_j$ for $i, j = 0, \dots, n$, we have

$$(19) \quad C_{kp}^{S_\lambda^{n+1}} = \lambda^p C_{k,0}^{S_\lambda^{n+1}}$$

so that

$$\mathcal{R}(S_\lambda^{n+1}) = \text{span}\{C_{k,0}^{S_\lambda^{n+1}} : 0 \leq k \leq n+1\}.$$

The associated valuations

$$(20) \quad \tau_k := \text{glob } C_{k,0}^{S_\lambda^{n+1}} \in \mathcal{V}(S_\lambda^{n+1})$$

constitute a natural basis for the space $\mathcal{V}^{SO(n+2)}$ of all invariant valuations. Using the identity $\pi(k+1)!\omega_k\omega_{k+1} = (2\pi)^{k+1}$ we obtain

$$(21) \quad C_{k0}^{S_\lambda^{n+1}}(S_\lambda^j, \cdot) = \delta_j^k \frac{k! \omega_k}{\pi^k} \mathcal{H}^j \Big|_{S_\lambda^j}$$

$$(22) \quad \tau_k(S_\lambda^j) = \delta_j^k 2 \left(\frac{2}{\sqrt{\lambda}} \right)^k$$

where \mathcal{H}^j denotes j -dimensional Hausdorff measure. Unfortunately this basis is not multiplicative, i.e. in general $\tau_j \cdot \tau_k \neq \tau_{j+k}$, even up to scale. It is helpful to define two other natural bases that *are* multiplicative. For the first, let dg denote the Haar probability measure on $SO(n+2)$ and define the valuation

$$\phi := \frac{2}{\sqrt{\lambda}} \int_{SO(n+2)} \chi(\cdot \cap gS_\lambda^n) dg$$

Then by Theorem 2.1 (2),

$$(23) \quad \phi^k = \left(\frac{2}{\sqrt{\lambda}} \right)^k \int_{SO(n+2)} \chi(\cdot \cap gS_\lambda^{n-k+1}) dg$$

and we again arrive at a basis $\phi^0, \dots, \phi^{n+1}$.

For the second, let $t = t_{\mathbb{R}^{n+2}} := \text{glob}(C_{10}^{\mathbb{R}^{n+2}})$. By Theorem 2.3.18 of part 2 of [7], the Alesker powers of t satisfy

$$(24) \quad \frac{\pi^k t^k}{k!} = \omega_k \mu_k,$$

where $\mu_k \in \mathcal{V}(\mathbb{R}^{n+2})$ is the k th intrinsic volume. In particular, $t = \frac{2}{\pi} \mu_1$. We abuse notation and denote again by t the restriction of this valuation to the sphere. Thus t^0, \dots, t^{n+1} constitute another basis for $\mathcal{V}^{SO(n+2)}$.

We describe the relations among these three bases. Clearly

$$(25) \quad \phi^k(S_\lambda^j) = \begin{cases} 2 \left(\frac{2}{\sqrt{\lambda}} \right)^k & j - k \geq 0 \text{ and even} \\ 0 & \text{otherwise} \end{cases}$$

which with (22) implies that

$$\phi^k = \sum_{j=0}^{\infty} \left(\frac{\lambda}{4}\right)^j \tau_{k+2j}.$$

Thus the generating function for the coefficients c_i of τ_i in the expansion of ϕ^k is $x^k \left(1 - \frac{\lambda x^2}{4}\right)^{-1}$, i.e.

$$(26) \quad \phi^k = \sum_j [x^j] \left(x^k \left(1 - \frac{\lambda x^2}{4}\right)^{-1} \right) \tau_j$$

where we use the standard generating functions notation

$$[x^j] \left(\sum_i c_i x^i \right) := c_j.$$

Lemma 3.4.

$$(27) \quad t^k = \phi^k \left(1 - \frac{\lambda \phi^2}{4}\right)^{-\frac{k}{2}}$$

$$(28) \quad = \sum_{j=0}^{\infty} \binom{\frac{k}{2} + j - 1}{j} \left(\frac{\lambda}{4}\right)^j \phi^{k+2j}$$

$$(29) \quad = \sum_{j=0}^{\infty} \binom{\frac{k}{2} + j}{j} \left(\frac{\lambda}{4}\right)^j \tau_{k+2j}$$

Proof. The relation (27) is Proposition 2.4.20 in [7]. In fact it is easy to prove directly by the template method: simply evaluate both sides on the great spheres S_λ^j , using (24) and the known values of the intrinsic volumes of spheres (e.g. [24]) (the case $k = 2$ is especially convenient, from which the other cases follow, since t is the unique square root of t^2 in $\mathbb{R}[[\phi]]$ that assigns positive values to curves). The explicit expansion (28) then follows from (15).

To obtain (29), note that by (26), (27),

$$(30) \quad t^k = \sum_j [x^j] \left(x^k \left(1 - \frac{\lambda x^2}{4}\right)^{-\frac{k}{2}} \left(1 - \frac{\lambda x^2}{4}\right)^{-1} \right) \tau_j.$$

□

Corollary 3.5.

$$t \cdot \tau_k = \sum_{j \geq 0} \binom{2j}{j} \left(\frac{\lambda}{16}\right)^j \tau_{k+2j+1}.$$

Proof. By (30), for any formal power series p we have

$$p(t) = \sum_j [x^j] \left(\left(1 - \frac{\lambda x^2}{4}\right)^{-1} p \left(x \left(1 - \frac{\lambda x^2}{4}\right)^{-\frac{1}{2}} \right) \right) \tau_j.$$

It follows that for any formal power series q

$$t \cdot \sum_j [x^j] (q(x)) \tau_j = \sum_j [x^j] \left(x \left(1 - \frac{\lambda x^2}{4}\right)^{-\frac{1}{2}} q(x) \right) \tau_j.$$

In particular, this is true of $q(x) = x^k$.

□

3.3. Behavior of the C_{kp} under isometric immersions.

Definition 3.6. For $2p + 2l \leq k$, define C_{kpl} to be the object that assigns to any isometric immersion $e : M \hookrightarrow N$ of smooth Riemannian manifolds the curvature measure

$$\mathcal{C}(M) \ni \iota_e(C_{kpl}) := C_{kpl}^e := \begin{cases} \frac{\omega_k}{\pi^{k(m+1-k)} \omega_{m+1-k}} [\bar{\rho}_{rel}(\phi_{kpl}), 0] & \text{if } k < \dim M \\ \frac{\omega_k}{\pi^k} [0, \bar{\rho}_{rel}(\psi_{pl})] & \text{if } k = \dim M \\ 0 & \text{if } k > \dim M. \end{cases}$$

We denote the ω -span of the C_{kpl} , the space of *relative Riemannian curvature measures*, by \mathcal{R}_{rel} .

We introduce the ω -linear isomorphism $\mathcal{R}_{rel} \rightarrow \mathbb{R}[[x, y, z]]$ by

$$(31) \quad \beta : C_{kpj} \mapsto x^{k-2p-2j} y^p z^j.$$

The normalizing constants in the definition of the C_{kpl} are chosen such that

- (1) $C_{k,0,l}^e = C_{k,l}^M$ if $N = \mathbb{R}^{n+1}$
- (2) $C_{k,p,l}^e = \lambda^p C_{k,0,l}^e$ if N is a sphere of curvature λ .

Theorem 3.7. *Let $e : M \hookrightarrow N$ be an isometric immersion of one smooth Riemannian manifold into another. Then for all k, p*

$$(32) \quad e^* C_{k,p}^N = \sum_{j=0}^{\infty} \binom{\frac{k}{2} + j}{j} \left(\frac{1}{4}\right)^j C_{k+2j,p,j}^e$$

In other words, if $\delta : \mathbb{R}[[\xi, \eta]] \rightarrow \mathbb{R}[[x, y, z]]$ is the ω -linear map

$$(33) \quad \delta : \xi^k \eta^p \mapsto \left(1 - \frac{z}{4}\right)^{-1} \left(x \left(1 - \frac{z}{4}\right)^{-\frac{1}{2}}\right)^k \left(y \left(1 - \frac{z}{4}\right)^{-1}\right)^p$$

then

$$\iota_e \circ \beta^{-1} \circ \delta = e^* \circ \iota_N \circ \alpha^{-1}.$$

The proof is based on the following, to be proved in Section 4.2.

Lemma 3.8. *There exist universal constants $c_{k,p,j}^{m,n}$ such that*

$$(34) \quad e^* C_{kp}^N = \sum_j c_{k,p,j}^{m,n} C_{k+2j,p,j}^e.$$

Proof of Theorem 3.7. Assuming Lemma 3.8, we evaluate the constants in (34) via the template method. First, choose $N = S_\lambda^{n+1}$ and let $M = S_\mu^{m+1}$ be a subsphere of curvature $\mu \geq \lambda$. Then

$$e^* C_{kp}^N = \sum_{j=0}^{\infty} c_{k,p,j}^{m,n} C_{k+2j,p,j}^e = \lambda^p \sum_{j=0}^{\infty} c_{k,p,j}^{m,n} C_{k+2j,0,j}^e.$$

Since $C_{kp}^N = \lambda^p C_{k0}^N$ we also have

$$e^* C_{kp}^N = \lambda^p e^* C_{k0}^N = \lambda^p \sum_{j=0}^{\infty} c_{k,0,j}^{m,n} C_{k+2j,0,j}^e.$$

The Gauss equation (14) now yields $C_{k+2j,0,j}^e = (\mu - \lambda)^j C_{k+2j,0,0}^e$, so we conclude that $c_{k,p,j}^{m,n} = c_{k,0,j}^{m,n} =: c_{k,j}^{m,n}$.

In order to determine these constants we consider a standard embedding of $M = S_\lambda^{m+1}$ into $N = \mathbb{R}^{n+1}$. Then

$$e^* C_{k0}^N = \sum_{j=0}^{\infty} c_{k,j}^{m,n} C_{k+2j,0,j}^e = \sum_{j=0}^{\infty} c_{k,j}^{m,n} C_{k+2j,j}^M = \sum_{j=0}^{\infty} c_{k,j}^{m,n} \lambda^j C_{k+2j,0}^M$$

Globalizing this relation yields

$$t^k = \sum_{j=0}^{\infty} C_{k,j}^{m,n} \lambda^j \tau_{k+2j}.$$

Now (32) follows from Lemma 3.4. \square

Corollary 3.9. *If the embedding $e : M \rightarrow N$ is totally geodesic, then $e^* C_{kp}^N = C_{kp}^M$ for all k, p .*

3.4. Lipschitz-Killing curvature measures.

Definition 3.10. Define the subspace $\widetilde{\mathcal{L}\mathcal{K}} \subset \mathcal{R}$ as the ω -span of the elements

$$(35) \quad \bar{\Lambda}_k := \sum_{j=0}^{\infty} \binom{\frac{k}{2} + j}{j} \left(\frac{1}{4}\right)^j C_{k+2j,j}, \quad k = 0, 1, \dots$$

The alternative normalization $\Lambda_k := \frac{\pi^k}{k! \omega_k} \bar{\Lambda}_k$ agrees with the corresponding definition in [10]. If $k = \dim M$ then $\Lambda_k^M(A, \cdot)$ equals the Riemannian volume measure restricted to $A \subset M$. However, the $\bar{\Lambda}_k$ are more natural algebraically, since

$$(36) \quad \alpha(\bar{\Lambda}_k) = \xi^k \left(1 - \frac{\eta}{4}\right)^{-\frac{k}{2}-1}.$$

Theorem 3.11. *The subspace $\widetilde{\mathcal{L}\mathcal{K}}$ consists precisely of all elements $\Psi \in \mathcal{R}$ such that*

$$(37) \quad e^* \Psi^N = \Psi^M$$

for every smooth isometric immersion $e : M \hookrightarrow N$ of smooth Riemannian manifolds.

Proof. We introduce an alternate basis for the space of relative Riemannian curvature measures as follows. Put

$$\Xi_{kpl} = \sum_{\pi} \operatorname{sgn}(\pi) \Omega_{\pi_1 \pi_2}^N \cdots \Omega_{\pi_{2p-1} \pi_{2p}}^N \Omega_{\pi_{2p+1} \pi_{2p+2}}^M \cdots \Omega_{\pi_{2p+2l-1} \pi_{2p+2l}}^M \theta_{\pi_{2p+2l+1}} \cdots \theta_{\pi_k} \omega_{\pi_{k+1}, 0} \cdots \omega_{\pi_m, 0}$$

for $2p + 2l \leq k$, and

$$\Theta_{pl} = \sum_{\pi} \operatorname{sgn}(\pi) \theta_{\pi_0} \Omega_{\pi_1 \pi_2}^N \cdots \Omega_{\pi_{2p-1} \pi_{2p}}^N \Omega_{\pi_{2p+1} \pi_{2p+2}}^M \cdots \Omega_{\pi_{2p+2l-1} \pi_{2p+2l}}^M \theta_{\pi_{2p+2l+1}} \cdots \theta_{\pi_m}$$

For $2p + 2l \leq k$, define $\Gamma_{kpl} \in \mathcal{R}_{rel}$ to be the object that assigns to any smooth isometric embedding $e : M^{m+1} \rightarrow N^{n+1}$ the curvature measure

$$\Gamma_{kpl}^e := \begin{cases} \frac{\omega_k}{\pi^k (m+1-k)! \omega_{m+1-k}} [\Xi_{kpl}, 0] & \text{if } k < m+1 \\ \frac{\omega_k}{\pi^k} [0, \Theta_{pl}] & \text{if } k = m+1 \\ 0 & \text{if } k > m+1. \end{cases}$$

on M . Thus

$$(38) \quad \Gamma_{k,0,p}^e = C_{k,p}^M$$

and

$$(39) \quad C_{kpl} = \sum_{j=0}^l (-1)^j \binom{l}{j} \Gamma_{k,p+j,l-j}.$$

In other words, if we define the ω -linear map

$$\gamma : \Gamma_{kpl} \mapsto x^{k-2p-2j} y^p z^j$$

then $\gamma \circ \beta^{-1}(x^a y^b z^c) = x^a y^b (z - y)^c$.

We first give a proof, using this formalism, of the well known fact (cf. the Introduction) that $e^* \bar{\Lambda}_k^N = \bar{\Lambda}_k^M$ for any smooth isometric immersion $e : M \hookrightarrow N$. By (36) and (38) this is encoded in the relation

$$(40) \quad e^* \circ \iota_N \circ \alpha^{-1} \left(\xi^k \left(1 - \frac{\eta}{4} \right)^{-\frac{k}{2}-1} \right) = \iota_e \circ \gamma^{-1} \left(x^k \left(1 - \frac{z}{4} \right)^{-\frac{k}{2}-1} \right).$$

By Theorem 3.7, this relation holds, since

$$\begin{aligned} \xi^k \left(1 - \frac{\eta}{4} \right)^{-\frac{k}{2}-1} &\xrightarrow{\delta} \left(1 - \frac{z}{4} \right)^{-1} \left(x \left(1 - \frac{z}{4} \right)^{-\frac{1}{2}} \right)^k \left(1 - \frac{y}{4} \left(1 - \frac{z}{4} \right)^{-1} \right)^{-\frac{k}{2}-1} \\ &= x^k \left(1 - \frac{z}{4} - \frac{y}{4} \right)^{-\frac{k}{2}-1} \\ &\xrightarrow{\gamma \circ \beta^{-1}} x^k \left(1 - \frac{z}{4} \right)^{-\frac{k}{2}-1}. \end{aligned}$$

Conversely, suppose $\Psi = \sum_{kp} a_{kp} C_{kp} \in \mathcal{R}$ satisfies (37) for all isometric immersions $e : M \hookrightarrow N$. We wish to show that for each $k \geq 0$ the sequence of coefficients $(a_{k+2j,j})_j$ is proportional to $\left(\binom{k+j}{j} \left(\frac{1}{4} \right)^j \right)_j$. We know this to be the case if $\Psi \in \widetilde{\mathcal{L}\mathcal{K}}$. Fixing k , it is therefore enough merely to show that the $a_{k+2j,j}, j > 0$, are determined by $A := a_{k,0} \frac{\omega_k k!}{\pi^k}$. By induction we may assume that the statement is true for $0, \dots, k-1$; subtracting the terms that correspond to these values, we may assume that all $a_{jp} = 0, j \leq k-1$.

Given $j > 0$, consider the standard embedding $e : \mathbb{R}^k \rightarrow \mathbb{R}^{k+2j+1}$. Thus $e^*(\Psi^{\mathbb{R}^{k+2j+1}}) = \Psi^{\mathbb{R}^k} = A \Phi_k^{\mathbb{R}^k}$. Taking $M_\lambda := \mathbb{R}^k \times S_\lambda^{2j}$, with standard isometric embeddings $e_\lambda : M_\lambda \hookrightarrow \mathbb{R}^{k+2j+1}$, as subsets of \mathbb{R}^{k+2j+1} the normal cycles $\text{nc}(e_\lambda(M_\lambda)) \rightarrow \chi(S^{2j})N(\mathbb{R}^k) = 2N(\mathbb{R}^k)$ in the flat metric topology as $\lambda \rightarrow \infty$. This is an elementary case of Theorem 3.7 of [22]; alternatively it follows the product formula for normal cycles and the fact that if $S_r^{2k} \subset \mathbb{R}^{2k+1}$ is the standard sphere of radius r then $\lim_{r \downarrow 0} \text{nc}(S_r^{2k}) = 2 \text{nc}(\{0\})$. Therefore by (2)

$$\Psi^{\mathbb{R}^{k+2j+1}}(e_\lambda(M_\lambda), \cdot) \rightarrow 2\Psi^{\mathbb{R}^{k+2j+1}}(\mathbb{R}^k, \cdot) = 2A\Phi_k(\mathbb{R}^k, \cdot)$$

as measures on \mathbb{R}^{k+2j+1} . On the other hand, the hypothesis of isometric invariance implies that

$$\Psi^{\mathbb{R}^{k+2j+1}}(e_\lambda(M_\lambda), \cdot) = e_{\lambda*} \left[\Psi^{M_\lambda}(M_\lambda, \cdot) \right]$$

But it is clear that the signed measures $e_{\lambda*} \left[C_{lm}^{M_\lambda}(M_\lambda, \cdot) \right] \rightarrow 0$ for $(l, m) \neq (k+2j, j)$, and that $C_{k+2j,j}^{M_\lambda}(M_\lambda, [0, 1]^k \times S_\lambda^{2j}) =: c \neq 0$, independent of λ . It follows that $a_{k+2j,j} = \frac{2A}{c}$. \square

3.5. Lipschitz-Killing valuations. The Lipschitz-Killing curvature measures are distinguished also by the fact that their globalizations behave in a universal fashion with respect to Alesker multiplication.

3.5.1. The universal module. Given a smooth Riemannian manifold M , we set $\mathcal{L}\mathcal{K}(M) \subset \mathcal{V}(M)$ for the **Lipschitz-Killing algebra** of M , generated by

$$(41) \quad t_M := \text{glob}(\bar{\Lambda}_1^M) \in \mathcal{V}(M).$$

Lemma 3.12. *Given k, p, m , there exists $\Psi_{k,p,m} \in \mathcal{R}$ such that, if M is any smooth oriented Riemannian manifold of dimension $m+1$, then*

$$t_M \cdot C_{kp}^M = \Psi_{k,p,m}^M.$$

In particular, $t_M \cdot \mathcal{R}(M) \subset \mathcal{R}(M)$ for every oriented Riemannian manifold M .

We postpone the proof of Lemma 3.12 to Section 4.1 below.

Assuming Lemma 3.12, we now determine the $\Psi_{kpm} \in \mathcal{R}$. Using the identification (18) of \mathcal{R} with $\mathbb{R}[[\xi, \eta]]$, we define an action of $\mathbb{R}[t]$ on \mathcal{R} by

$$(42) \quad \alpha(t \cdot \Psi) := \frac{\xi}{\sqrt{1 - \frac{\eta}{4}}} \alpha(\Psi).$$

Thus by (36)

$$(43) \quad t \cdot \bar{\Lambda}_k = \bar{\Lambda}_{k+1}.$$

Writing out (42) explicitly:

$$(44) \quad t \cdot C_{kp} = \sum_{j=0}^{\infty} \binom{2j}{j} \frac{1}{16^j} C_{k+2j+1, p+j},$$

$$(45) \quad t^i \cdot C_{kp} = \sum_{j=0}^{\infty} \binom{\frac{i}{2} + j - 1}{j} \frac{1}{4^j} C_{k+i+2j, p+j}.$$

Theorem 3.13. *The Alesker powers of t_M are given by*

$$(46) \quad t_M^k = \text{glob}(\bar{\Lambda}_k^M).$$

The algebra homomorphism $\mathbb{R}[t] \rightarrow \mathcal{LK}(M) \subset \mathcal{V}(M)$ determined by $t \mapsto t_M$ intertwines the actions of $\mathbb{R}[t]$ on \mathcal{R} and of $\mathcal{LK}(M)$ on $\mathcal{R}(M)$, i.e.

$$(47) \quad (t \cdot \Phi)^M = t_M \cdot \Phi^M.$$

Proof. We must show that

$$(48) \quad \Psi_{k,p,m} = \sum_{j=0}^{\infty} \binom{2j}{j} \left(\frac{1}{16}\right)^j C_{k+2j+1, p+j}.$$

This formula is determined by the cases $M = S_\lambda^{n+1}$ for varying n, λ , as follows. By Theorem 3.11, the valuation $t \in \mathcal{V}(S_\lambda^{n+1})$, described in Section 3.2, is identical to $t_{S_\lambda^{n+1}}$ as defined in (41). Globalizing $t \cdot C_{kp}$ in S_λ^n and using (19), (20), and Corollary 3.5, we obtain

$$\text{glob}(t \cdot C_{kp}^M) = \lambda^p t \cdot \tau_k = \lambda^p \sum_{j=0}^{\infty} \binom{2j}{j} \left(\frac{\lambda}{16}\right)^j \tau_{k+2j+1} = \text{glob} \left(\sum_{j=0}^{\infty} \binom{2j}{j} \left(\frac{1}{16}\right)^j C_{k+2j+1, p+j}^M \right)$$

as claimed. \square

3.5.2. There are no further universal Riemannian valuations. Our next result expresses the fact that $\widetilde{\mathcal{LK}}$ comprises all elements of \mathcal{R} whose globalizations act on \mathcal{R} in a universal way.

Theorem 3.14. *Let $\Phi \in \mathcal{R}$, and suppose that there exists a map $L = L_\Phi : \mathcal{R} \rightarrow \mathcal{R}$ such that for every Riemannian manifold M and every $\Psi \in \mathcal{R}$*

$$\text{glob}(\Phi^M) \cdot \Psi^M = L(\Psi)^M.$$

Then $\Phi \in \widetilde{\mathcal{LK}}$, and L is described by (42).

In order to prove this we examine the particular cases $M = \mathbb{C}P_\lambda^n$, the complex space form of constant holomorphic sectional curvature 4λ , and exploit the detailed understanding of the integral geometry of such spaces achieved in [10, 11]. The needed facts from hermitian integral geometry are given in the next Proposition. For simplicity we denote the realizations $C_{kp}^{\mathbb{C}P_\lambda^n} \in \mathcal{R}(\mathbb{C}P_\lambda^n)$ simply by C_{kp}^λ . Put \mathcal{V}_λ^n for the algebra of isometry-invariant valuations on $\mathbb{C}P_\lambda^n$. Following [10], we denote the canonical generator of the Lipschitz-Killing algebra by $t_\lambda := t_{\mathbb{C}P_\lambda^n} \in \mathcal{V}_\lambda^n$, and take $s \in \mathcal{V}_\lambda^n$ to be the valuation

$$s(Q) := \int_{Gr_{n-1}} \chi(Q \cap P) dP$$

where \overline{Gr}_{n-1} is the homogeneous space of all totally geodesic subspaces of $\mathbb{C}P_\lambda^n$ of complex codimension 1, and dP is the (appropriately normalized) Haar measure (cf. [10], (3.1)).

Proposition 3.15.

- (1) *The restriction of the globalization map to the space of invariant angular curvature measures gives a linear isomorphism with \mathcal{V}_λ^n .*
- (2) *If $\lambda \neq 0$ and $0 \leq k \leq n$, then the curvature measures*

$$C_{k,0}^\lambda, C_{k,1}^\lambda, \dots, C_{k,[k/2]}^\lambda$$

are linearly independent.

- (3) *$\mathcal{R}(\mathbb{C}P_\lambda^n)$ coincides with the space of all invariant angular curvature measures on $\mathbb{C}P_\lambda^n$.*
- (4) *Let $p(\tau, \sigma)$ be a polynomial in the formal variables τ, σ , where $\deg \tau = 1, \deg \sigma = 2$. If multiplication by $p(t_\lambda, s) \in \mathcal{V}_\lambda^n$ stabilizes the space of invariant angular curvature measures on $\mathbb{C}P_\lambda^n$ then the terms of $p(\tau, \sigma)$ of weighted degree less than $n - 2$ are independent of σ .*

Proof. Conclusion (1) follows from Lemma 3.9 of [10]. Corollary 5.3, proved below, yields (2). Therefore the span of the C_{kp} coincides with that of the Δ_{kp} (defined in Section 5.1.2 below) for k lying in this range. Examining the standard embedding of $\mathbb{C}P_\lambda^n$ into $\mathbb{C}P_\lambda^N, N \gg n$, conclusion (3) now follows from Lemma 3.6 of [10] and Corollary 3.9 above.

To prove (4), recall that the algebra $\text{Val}^{U(n)}$ is the quotient $\mathbb{R}[s, t]/(f_{n+1}, f_{n+2})$, where each $f_i(s, t)$ is a certain polynomial of weighted degree i , where $\deg s = 2, \deg t = 1$. We apply Theorem 2 of [11], which states that the algebra of isometry invariant valuations on $\mathbb{C}P_\lambda^n$ whose action preserves the space of angular invariant curvature measures is given by all $p(t_\lambda, s)$ such that

$$(49) \quad ts(4s - t^2) \frac{\partial p}{\partial \sigma} \left(\frac{t}{\sqrt{1 - \lambda s}}, s \right) = 0 \quad \text{in } \text{Val}^{U(n)}.$$

Clearly we may assume that $p(\tau, \sigma)$ is divisible by σ . Let $\tau^i \sigma^j$ be the term of smallest degree of $p(\tau, \sigma)$ (we may assume that the coefficient is unity). It follows that the term of lowest degree of the left hand expression in (49) is $ts(4s - t^2)t^i s^{j-1}$. The ideal of relations between s, t in $\text{Val}^{U(n)}$ is generated by homogeneous polynomials in degrees $n + 1, n + 2$, so the presence of this last monomial implies that $i + 2j \geq n - 2$. \square

Proof of Theorem 3.14. Suppose $\Phi \in \mathcal{R} \setminus \widetilde{\mathcal{L}\mathcal{K}}$. Take N large enough that this is still true modulo the ω -span of the $C_{kp}, k > N$. By Proposition 3.15, (1) and (2), for sufficiently large $n \geq N$ the globalization $\text{glob}(\Phi^{\mathbb{C}P_\lambda^n}) \notin \mathcal{L}\mathcal{K}(\mathbb{C}P_\lambda^n)$. Choose a polynomial $p(\tau, \sigma)$ such that $p(t_\lambda, s) = \text{glob}(\Phi^{\mathbb{C}P_\lambda^n})$ in \mathcal{V}_λ^n . By Proposition 3.15, items (4) and (2), it follows that $\Phi \in \widetilde{\mathcal{L}\mathcal{K}}$ modulo the ω -span of the $C_{kp}, k \geq n - 2$. This is a contradiction for $n \geq N$ sufficiently large. \square

4. PROOFS OF LEMMAS 3.8 AND 3.12

4.1. Proof of Lemma 3.12.

4.1.1. *The Alesker-Bernig formula.* We use the description provided in [5, 20] of the Alesker product in terms of differential forms. Given a smooth oriented Riemannian manifold M^{m+1} , denote by Σ the oriented $(3m + 2)$ -dimensional manifold

$$(50) \quad \Sigma = \left\{ (\xi, \zeta, \eta) \in SM \times_M SM \times_M SM : \xi \neq \pm\eta, \zeta \in \overline{\xi, \eta} \right\}$$

where $\overline{\xi, \eta}$ denotes the open geodesic segment in $S_{\pi\xi}M$ joining ξ, η . We have three associated projections

$$\xi, \zeta, \eta : \Sigma \rightarrow SM$$

to the first, second and third factors respectively.

We regard the projection $\xi : SM \times_M SM \times_M SM \rightarrow SM$ as a bundle associated to the principal $O(m)$ bundle $z : FM \rightarrow SM$. Here $O(m)$ acts diagonally on the model fiber $S^m \times S^m \subset \mathbb{R}^{m+1} \times \mathbb{R}^{m+1}$ of ξ via the following action on \mathbb{R}^{m+1} : let u_0, \dots, u_m be standard coordinates for \mathbb{R}^{m+1} , and let $O(m)$ fix the 0 coordinate and act as usual on (u_1, \dots, u_m) . Thus the fibers Σ_ξ correspond to the $O(m)$ -invariant subset

$$C := \{(v, u) : u \neq \pm e_0, v \in \overline{e_0, u}\} \subset S^m \times S^m.$$

We endow the bundle ξ with the connection induced by the canonical connection on the principal bundle z , described in Section 2.2.3.

Theorem 4.1 ([5, 20]). *If $\mu \in \mathcal{V}^\infty(M)$, $\beta \in \Omega^m(SM)$, and $\gamma \in \Omega^{m+1}(M)$, then*

$$\mu \cdot [\beta, \gamma] = [\theta, \psi]$$

with

$$(51) \quad \theta = (-1)^{m+1} \xi_* (\zeta^* \beta \wedge \eta^* a^* \Delta_\mu) + \pi^* \mathcal{F}_\mu \cdot \beta,$$

$$(52) \quad \psi = \pi_* (\beta \wedge a^* \Delta_\mu) + \mathcal{F}_\mu \cdot \gamma.$$

Here

$$\mathcal{F} : \mathcal{V}^\infty(M) \rightarrow C^\infty(M), \quad \Delta : \mathcal{V}^\infty(M) \rightarrow \Omega^n(SM)$$

are defined by $\mathcal{F}_\mu(x) = \mu(\{x\})$ and $\Delta_\mu = D\rho + \pi^* \tau$, provided $\mu = [[\rho, \tau]]$.

The sign in front of the first term of (51) corrects the corresponding formula in [20]. Here the operator $D\rho$ is the *Rumin differential* of ρ [8, 26], defined as the unique $(m+1)$ -form $D\rho = d(\rho + \theta_0 \wedge \xi)$ such that

$$\theta_0 \wedge d(\rho + \theta_0 \wedge \xi) = 0$$

for some $(m-1)$ -form ξ , where $\theta_0 \in \Omega^1(SM)$ is the contact form. Recall that $a : SM \rightarrow SM$, $a(\xi) = -\xi$ denotes the fiberwise antipodal map. The fibers Σ_ξ may be compactified so as to make clear the convergence of the fiber integrals. The orientations of the fibers in these integrals are canonical; we need not make them precise here since we only wish to confirm the existence of the universal product formula. Actually the term $\pi^* \tau$ may be ignored, for trivial dimensional reasons: the integration is of forms of degree $2m+1$ over fibers of dimension $m+1$, so that any terms including a horizontal factor of degree $m+1$ must yield zero. By the same token Δ_μ may be perturbed by any horizontal form of degree $m+1$ without affecting the result.

Define

$$(53) \quad \begin{aligned} \bar{\mathcal{A}}_m^+ \ni \lambda_k &:= \sum_{j=0}^{\lfloor \frac{m-k}{2} \rfloor} \binom{\frac{k}{2} + j}{j} \left(\frac{1}{4}\right)^j \frac{\omega_{k+2j}}{\pi^{k+2j} (m-k-2j+1)! \omega_{m-k-2j+1}} \phi_{k+2j,j} \\ &= \frac{\omega_k}{\pi^k (m-k+1)! \omega_{m-k+1}} \sum_{j=0}^{\lfloor \frac{m-k}{2} \rfloor} \frac{1}{2^j j! (m-k-2j)!} \phi_{k+2j,j}. \end{aligned}$$

Thus $\bar{\rho}(\lambda_k)$ is the boundary term of $\bar{\Lambda}_k \in \mathcal{R}$, as in Definition 3.1. Here we use the relation

$$k\omega_k = 2\pi\omega_{k-2}, \quad k = 1, 2, \dots$$

For nonnegative integers k, p satisfying $2p \leq k \leq m-1$ we define elements of degree $m+1$ by

$$\bar{\mathcal{A}}_m^+ \ni D_{kp} := (-1)^k \sum_{\pi} \operatorname{sgn}(\pi) \Omega_{\pi_1 \pi_2} \cdots \Omega_{\pi_{2p-1} \pi_{2p}} \theta_{\pi_{2p+1}} \cdots \theta_{\pi_k} \Omega_{\pi_{k+1}, 0} \omega_{\pi_{k+2}, 0} \cdots \omega_{\pi_m, 0}.$$

The sum ranges over all permutations of $1, \dots, m$.

The proof of Lemma 3.12 using the Alesker-Bernig formula is based on the following observation.

Lemma 4.2. *The formal exterior derivative $d\lambda_1$ is the sum of a multiple of θ_0 and an element of \mathcal{B}_m .*

Proof. More precisely and generally, we compute

$$(54) \quad d\lambda_k = \frac{\omega_k}{\pi^k (m-k-1)!! \omega_{m-k+1}} \times \left[\theta_0 \wedge \sum_{j=0}^{\lfloor \frac{m-k}{2} \rfloor} \frac{k}{2^j j! (m-k-2j)!!} \phi_{k+2j-1, j} + \delta_{m-1} \right]$$

where

$$\delta_{m-1} := \frac{1}{2^{\frac{m-k-1}{2}} \left(\frac{m-k-1}{2}\right)!} D_{m-1, \frac{m-k-1}{2}}$$

for $m-k$ odd and $\delta_{m-1} := 0$ otherwise. This follows by telescoping from the formula

$$d\phi_{kp} - (k-2p)\theta_0 \wedge \phi_{k-1, p} + 2pD_{k-2, p-1} - (m-k)D_{kp} = 0$$

since $\omega_{ij} = 0$ whenever both $i, j \geq 0$, which in turn follows from the structure equations (5). \square

Proposition 4.3. *For any smooth oriented manifold M of dimension $m+1$, we have $D(\bar{\rho}(\lambda_k)) \equiv \bar{\rho}(d\lambda_k)$ modulo the space of horizontal forms.*

Proof. Recall that $\bar{\rho} \circ d = d \circ \bar{\rho}$. Since δ_{m-1} has degree $m+1$ and no ω_0 factors, it follows that for any Riemannian M^{m+1} its realization is a functional multiple of $\bar{\rho}(\theta_0)$. \square

4.1.2. *The formal model.* Since normal cycles are Legendrian, and the realization of θ_0 is the contact form of any SM , with Proposition 2.2 the following implies Lemma 3.12.

Proposition 4.4. *Given k, p, m , there exist $\varphi_1 \in \bar{\mathcal{A}}_m^+$, $\varphi_2 \in \mathcal{B}_m^+$ with the following property. Given an oriented Riemannian manifold M^{m+1} , let θ, ψ be as in Theorem 4.1 for $\beta = \bar{\rho}(\phi_{kp}), \mu = t_M$. Then*

$$(55) \quad \theta = \bar{\rho}(\varphi_1), \quad \psi = \bar{\rho}(\varphi_2).$$

We will only give explicitly the construction of φ_1 , corresponding to the term of the Alesker-Bernig formula that involves fiber integration over Σ ; that of φ_2 is similar but significantly simpler.

The gist of the construction is contained in Lemmas 4.5 and 4.6. The idea is to carry out the fiber integration in the first term of the right hand side of (51) in purely symbolic terms (the second term vanishes since $\mathcal{F}_\mu = 0$, which is to say that t_M clearly vanishes on singletons). The underlying observation is Lemma 4.2, which implies that in the present setting the forms β, Δ_μ from (51) are realizations of elements of $\bar{\mathcal{A}}_m^+$. Lemma 4.6 shows that the process of pulling them back under ζ, η may be viewed as the realization of a parallel symbolic process. With the aid of Lemma 2.7, Lemma 4.5 shows that the fiber integration step in the Alesker-Bernig formula may also be carried out symbolically, and that the symbolic result belongs to $\bar{\mathcal{A}}_m^+$. Proposition 2.2 then concludes the proof.

Recalling the left actions of $O(m)$ on S^m and on $\bar{\mathcal{A}}_m$, we define the left action Λ of $O(m)$ on $\Omega^*(S^m, \bar{\mathcal{A}}_m)$ by

$$(56) \quad \Lambda_g \phi := L_g \circ (g^{-1*} \phi) = g^{-1*} (L_g \circ \phi).$$

Define similarly an action $\hat{\Lambda}$ of the same group on $\Omega^*(S^m \times S^m, \bar{\mathcal{A}}_m)$ via the diagonal action on $S^m \times S^m$. Let $\tilde{\zeta}, \tilde{\eta} : S^m \times S^m \rightarrow S^m$ denote the projections to the respective factors; these maps clearly intertwine the $O(m)$ actions. Denote by $\Omega^*(S^m, \bar{\mathcal{A}}_m)^+$ the subspace on which $O(m)$ acts by multiplication by the determinant, and note that, if $\alpha, \omega \in \Omega^*(S^m, \bar{\mathcal{A}}_m)^+$, then $\tilde{\eta}^* \alpha \wedge \tilde{\zeta}^* \omega \in \Omega^*(S^m \times S^m, \bar{\mathcal{A}}_m)^{O(m)}$.

Given $\xi_0 \in SM$ with $x_0 := \pi(\xi_0) \in M$, let b be an adapted frame at ξ_0 . Define the diffeomorphism $\psi_b : S_{x_0}M \rightarrow S^m$, $\psi_b^{-1}(u) := u_i b_i$, and define $\hat{\rho}_b : \Omega^*(S^m, \bar{\mathcal{A}}_m) \rightarrow \Omega^*(S_{x_0}M, \wedge^* T_{\xi_0}^* SM)$ by

$$(57) \quad \hat{\rho}_b(\phi) := \psi_b^*(\bar{\rho}_b \circ \phi) = \bar{\rho}_b \circ (\psi_b^* \phi).$$

If $b, b' = R_g b$ are adapted frames at $\xi_0 \in SM$, where $g \in O(m)$, then by Lemma 2.4

$$\hat{\rho}_{b'} \phi = \bar{\rho}_{R_g b} \circ \psi_{R_g b}^* \phi = \bar{\rho}_b \circ L_g \circ \psi_b^* ((g^{-1})^* \phi) = \bar{\rho}_b \circ \psi_b^* (L_g \circ (g^{-1})^* \phi) = \hat{\rho}_b (\Lambda_g \phi).$$

It follows that this value is independent of the choice of positively oriented frame b if ϕ is $SO(m)$ -invariant.

Consider the bundle $SM \times_M SM \rightarrow SM$ under projection to the first factor as an $O(m)$ bundle associated to z , endowed with the connection induced by the canonical connection on z . Here the action of $O(m)$ on the model fiber $S^m \subset \mathbb{R}^{m+1}$ is again induced by the standard action on the $1, \dots, m$ coordinates, and fixes the 0 coordinate; thus the projections from the bundle ξ , obtained by omitting the second or third coordinate, are both bundle maps that intertwine the connections. We then obtain a map $\hat{\rho} : \Omega^*(S^m, \bar{\mathcal{A}}_m)^+ \rightarrow \Omega^*(SM \times_M SM)$ as the restriction of $\hat{\rho}_b$ for any choice of positive adapted frame b , via the constructions of Section 2.3. In the same manner we have $\hat{\hat{\rho}} : \Omega^*(S^m \times S^m, \bar{\mathcal{A}}_m)^{O(m)} \rightarrow \Omega^*(SM \times_M SM \times_M SM)$. Factor the maps $\zeta = \pi_2 \circ \zeta_1, \eta = \pi_2 \circ \eta_1$, where the $\zeta_1, \eta_1 : SM \times_M SM \times_M SM \rightarrow SM \times_M SM$ respectively omit the third and second coordinates.

Lemma 4.5. *Let I denote integration over the model fiber $C := \{(z, y) : y \neq \pm e_0, z \in \bar{e}_0, \bar{y}\} \subset S^m \times S^m$, with an appropriate choice of orientation. Then the following diagram commutes for all $(m+1)$ -dimensional oriented Riemannian manifolds M :*

$$(58) \quad \begin{array}{ccc} (\Omega^*(S^m, \bar{\mathcal{A}}_m)^+)^{\otimes 2} & \xrightarrow{\hat{\rho}^{\otimes 2}} & \Omega^*(SM \times_M SM)^{\otimes 2} \\ \bar{\zeta}^* \wedge \bar{\eta}^* \downarrow & & \zeta_1^* \wedge \eta_1^* \downarrow \\ \Omega^*(S^m \times S^m, \bar{\mathcal{A}}_m)^{O(m)} & \xrightarrow{\hat{\hat{\rho}}} & \Omega^*(SM \times_M SM \times_M SM) \\ I \downarrow & & \xi_*^\Sigma \downarrow \\ \bar{\mathcal{A}}_m^+ & \xrightarrow{\bar{\rho}} & \Omega^*(SM) \end{array}$$

Proof. That the top square commutes follows straightforwardly from the definitions.

Lemma 2.7 implies that a variant of the bottom square commutes, in which the invariance conditions on the spaces on the left are relaxed, and the maps $\bar{\rho}, \hat{\hat{\rho}}$ are replaced by $\bar{\rho}_b, \hat{\rho}_b$ for some frame b at ξ_0 . Since the orientation of C is reversed under the action of $g \in O(m)$ if $\det g = -1$, we find that

$$I(\hat{\Lambda}_g \phi) = (\det g) L_g I(\phi), \quad g \in O(m).$$

Thus the restriction of I to the subspace of invariant elements indeed takes values in $\bar{\mathcal{A}}_m^+$, and the given diagram commutes. \square

The next step is to use $\hat{\rho}$ to construct a universal model for the pullback of $\bar{\rho}(\bar{\mathcal{A}}_m^+)$ under the projection π_2 of $SM \times_M SM$ to the second factor.

Lemma 4.6. *There exists a canonical map σ such that the following diagram commutes:*

$$(59) \quad \begin{array}{ccc} \bar{\mathcal{A}}_m^+ & \xrightarrow{\bar{\rho}} & \Omega^*(SM) \\ \sigma \downarrow & & \pi_2^* \downarrow \\ \Omega^*(S^m, \bar{\mathcal{A}}_m)^+ & \xrightarrow{\hat{\rho}} & \Omega^*(SM \times_M SM) \end{array}$$

Proof. Given $g \in O(m+1)$ we define an algebra homomorphism $s_g : \bar{\mathcal{A}}_m \rightarrow \Omega^*(S^m, \bar{\mathcal{A}}_m)$ by taking $s_g(\theta_i) = \tilde{L}_g(\theta_i), s_g(\Omega_{ij}) = \tilde{L}_g(\Omega_{ij})$ and

$$(60) \quad s_g(\omega_{i0}) = \tilde{L}_g(\omega_{i0}) + g_{ji} du_j,$$

where we recall the definition of \tilde{L}_g from the discussion around (4), (6). One sees readily that if $h \in O(m)$ then

$$s_{gh} = s_g \circ L_h, \quad s_{hg} = \Lambda_h \circ s_g.$$

From the first relation it follows that if $\phi \in \bar{\mathcal{A}}_m^+$ then $s_g(\phi)$ depends only on $g \cdot e_0 = u$ and $\det g$. Thus we may define $\sigma : \bar{\mathcal{A}}_m^+ \rightarrow \Omega^*(S^m, \bar{\mathcal{A}}_m)$ by $\sigma(\phi)|_u := s_g(\phi)|_u$, where $g \cdot e_0 = u$ and $\det g = 1$. The second relation now implies that $\sigma(\bar{\mathcal{A}}_m^+) \subset \Omega^*(S^m, \bar{\mathcal{A}}_m)^+$.

It remains to show that $\hat{\rho}, \sigma$ fit into (59). Selecting a point ξ_0 , a positive frame b at ξ_0 , and $g \in SO(m+1)$, we accomplish this by proving the corresponding fact for the contingently defined maps $\hat{\rho}_b, s_g$ above. This amounts to the following assertion. Let $u := g \cdot e_0 \in S^m$, $\eta_0 := \psi_b^{-1}(u) \in S_{x_0}M$, and $\beta := R_g b$. Thus β is a positive adapted frame at η_0 . The proof of the Lemma will be completed by showing that the diagram

$$(61) \quad \begin{array}{ccc} \bar{\mathcal{A}}_m & \xrightarrow{\pi_2^* \circ \bar{\rho}_\beta|_{\xi_0, \eta_0}} & \Lambda^* T_{\xi_0, \eta_0}(SM \times_M SM) \\ s_g|_u \downarrow & & r_{\xi_0} \downarrow \\ \Lambda^* T_u S^m \otimes \bar{\mathcal{A}}_m & \xrightarrow{\hat{\rho}_b|_u} & \Lambda^* T_{\eta_0} S_{x_0}M \otimes \Lambda^* T_{\xi_0} SM \end{array}$$

commutes, where the map on the right is defined as usual via the connection.

Observe first that if $\bar{g} : S^m \supset U \rightarrow SO(m+1)$ is a local section extending g , i.e.

$$\bar{g}(v) \cdot e_0 = (\bar{g}_{00}(v), \bar{g}_{10}(v), \dots, \bar{g}_{m0}(v)) = v, \quad \bar{g}(u) = g,$$

then (60) may be rewritten as

$$(62) \quad s_g(\omega_{i_0})|_u = \bar{g}_{00}(u) \bar{g}_{ji}(u) \omega_{j_0} + \bar{g}_{0i}(u) \bar{g}_{j_0}(u) \omega_{0j} + \bar{g}_{ji}(u) d\bar{g}_{j_0}|_u.$$

We construct a convenient choice of such an extension as follows. By continuity we may assume that $\xi_0 \not\perp \eta_0$, or equivalently that $u \not\perp e_0$. Thus by Lemma 2.5, we may take

$$(63) \quad \bar{g}(\overline{\text{rot}}_w(u)) := \overline{\text{rot}}_w \cdot g$$

for w in some small neighborhood of e_0 .

Let \bar{b} be the infinitesimally parallel extension of b , and define an extension $\bar{\beta}$ of β to an adapted moving frame on a neighborhood of η_0 in SM as follows: for v in a neighborhood of u in $S_{x_0}M$, take $\bar{\beta}(v) := R_{\bar{g}\psi_b(v)} b$; then extend by parallel translation along geodesics from x_0 . For $(\xi, \eta) \in SM \times_M SM$ lying in the appropriate domain we define $\bar{g}(\xi, \eta) \in SO(m+1)$ by the condition $R_{\bar{g}(\xi, \eta)} \bar{b}(\xi) = \bar{\beta}(\eta)$. Thus, by Lemma 2.5 and the definition of the infinitesimally parallel extension \bar{b} ,

$$(64) \quad \bar{g}(\xi, \eta) = \overline{\text{rot}}_{\psi_b(\xi)}^{-1} \cdot (\bar{g} \circ \psi_b)(\eta)$$

for $\xi \in S_{x_0} \setminus \{-\xi_0\}, \eta \in \psi_b^{-1}(U)$, so that by Lemma 2.5 and (63)

$$(65) \quad \bar{g}(\xi, \overline{\text{rot}}_{\xi}^{\xi_0}(\eta_0)) = \overline{\text{rot}}_{\psi_b(\xi)}^{-1} \cdot (\bar{g} \circ \psi_b(\overline{\text{rot}}_{\xi}^{\xi_0}(\eta_0))) = \overline{\text{rot}}_{\psi_b(\xi)}^{-1} \cdot (\bar{g}(\overline{\text{rot}}_{\psi_b(\xi)}(\psi_b(\eta_0)))) \equiv g.$$

Since the group action commutes with parallel translation, the map \bar{g} is determined by these restricted values: if $\xi', \eta' \in S_x M$ are the parallel translates of ξ, η along the geodesic from x_0 to x , then $\bar{g}(\xi', \eta') = \bar{g}(\xi, \eta)$.

Let $R : FM \times O(m+1) \rightarrow FM$ be the right action. We have the locally defined commutative diagram

$$(66) \quad \begin{array}{ccccccc} \mathcal{A}_m & \xrightarrow{\rho} & \Omega^*(FM) & \xrightarrow{R^*} & \Omega^*(FM \times O(m+1)) & \xrightarrow{\text{ev}_{b,g}} & \Lambda^* T_b FM \otimes \Lambda^* T_g O(m+1) \\ \uparrow & & \downarrow \bar{\beta}^* & & \downarrow (\bar{b}, \bar{g})^* & & \downarrow (\bar{b}, \bar{g})^*|_{\xi_0, \eta_0} \\ \bar{\mathcal{A}}_m & \xrightarrow{\bar{\rho}_{\bar{\beta}}} & \Omega^*(SM) & \xrightarrow{\pi_2^*} & \Omega^*(SM \times_M SM) & \xrightarrow{\text{ev}_{\xi_0, \eta_0}} & \Lambda^* T_{\xi_0} SM \otimes \Lambda^* T_{\eta_0} S_{x_0} M \end{array}$$

where, as in (61), the space on the lower right is identified with $\Lambda^* T_{\xi_0, \eta_0}(SM \times_M SM)$ via the connection. We claim that the composition of maps from the lower left corner across the top line to the lower right coincides with $\hat{\rho}_b \circ s_g|_u$, which will imply commutativity of (61).

Note first that the derivatives \bar{b}_*, \bar{g}_* at (ξ_0, η_0) vanish on $T_{\eta_0} S_{x_0} M, T_{\xi_0} SM$ respectively. The first of these statements is obvious, while the second follows from (65) and the definition of the connection. By the definitions of \bar{b} and of the realization maps we have

$$\begin{aligned} \bar{b}^*(\rho(\theta_i))\Big|_{\xi_0} &= \bar{\rho}_b(\theta_i), & \bar{b}^*(\rho(\Omega_{ij}))\Big|_{\xi_0} &= \bar{\rho}_b(\Omega_{ij}), \\ \bar{b}^*(\rho(\omega_{i0}))\Big|_{\xi_0} &= \bar{\rho}_b(\omega_{i0}), \\ \bar{b}^*(\rho(\omega_{ij}))\Big|_{\xi_0} &= 0, & 0 &\notin \{i, j\}. \end{aligned}$$

Meanwhile, the restrictions of \bar{g}_* and $(\bar{g} \circ \psi_b)_*$ to $T_{\eta_0}^* S_{x_0} M$ clearly coincide. Thus the basic theory of connections on principal bundles ([13], Lemma 1 of Section 5.1, and Lemma 1 of Section 6.1.1) gives the values at $(b, g) \in FM \times O(m+1)$ as

$$\begin{aligned} R^* \rho(\theta_i)\Big|_{b,g} &= g_{ji} \rho(\theta_j)\Big|_b \\ R^* \rho(\Omega_{ij})\Big|_{b,g} &= g_{ki} g_{lj} \rho(\Omega_{kl})\Big|_b \\ R^* \rho(\omega_{i0})\Big|_{b,g} &= g_{ki} g_{l0} \rho(\omega_{kl})\Big|_b + g_{ji} dg_{j0}. \end{aligned}$$

Applying the adjoints of the derivatives described above and comparing with the definition of s_g in (60) and (62) completes the proof. \square

Proof of Proposition 4.4. Recalling (12), we construct φ_1 by feeding $L_{(-1)} d\lambda_1 \otimes \phi_{kp}$ into the tensor square of (59), then applying (58). Note that, in applying Theorem 4.1 in our present circumstances, we have $\mathcal{F}_\mu = 0$, and we may ignore the $\pi^* \tau$ term. \square

4.2. Proof of Lemma 3.8. The proof is similar to that of Lemma 3.12, with the Alesker-Bernig formula replaced by the more elementary construction giving the restriction of smooth curvature measures on the level of differential forms, which we now describe.

4.2.1. Restrictions of curvature measures. Let $e : M^{m+1} \hookrightarrow N^{n+1}$ be an isometric immersion, $m < n$. Let $S_M^\perp N \subset SN|_M$ denote the bundle of spheres normal to M , with projection $q : S_M^\perp N \rightarrow M$. Consider the open submanifold $SN|_M^+ := SN|_M \setminus S_M^\perp N$. Put

$$i : SN|_M^+ \rightarrow SN, \quad j : S_M^\perp N \rightarrow SN.$$

for the inclusion maps. Normalized projection to the tangent spaces of M gives a projection $p : SN|_M^+ \rightarrow SM$, whose fibers are open hemispheres of dimension $n - m$.

Proposition 4.7. *Let $\omega \in \Omega^n(SN), \gamma \in \Omega^{n+1}(N)$. Then the fibers of the bundles p, q may be canonically oriented so that*

$$(67) \quad e^*[\omega, \gamma] = e^*[\omega, 0] = [p_* i^* \omega, q_* j^* \omega]$$

Although the fibers of p are not compact, they may be easily compactified by a blowing up process, so that the fiber integral operator p_* that occurs here is well defined.

Proof. Let a smooth polyhedron $A \subset M$ be given. The proof of Theorem 4.5 of [19] shows how to construct the normal cycle $\text{nc}_N(A)$ of A , considered as a subset of N , from its normal cycle $\text{nc}_M(A)$ as a subset of M : under a suitable choice of orientations of the fibers S_+^{n-m} of p , we have

$$\text{nc}_N(A) = i_*(\text{nc}_M(A) \times_p S_+^{n-m}) + \text{nc}_N(M)|_A.$$

The conclusion now follows from the fiber integration definition of products of currents.

Actually this construction is valid without regard to the Riemannian structure via passage to the conormal cycle. From that perspective, that the fibers are oriented canonically follows from the fact that the pair (M, N) is locally diffeomorphic to the model case $(\mathbb{R}^{m+1}, \mathbb{R}^{n+1})$. \square

The bundle p admits a natural connection, as follows. Take the subspace $SM \subset SN|_M^+$ to be parallel. Using the map $(\xi, \eta, \theta) \mapsto \cos \theta \xi + \sin \theta \eta$, the deleted bundle $SN|_M^+ \setminus SM \rightarrow SM$ may be identified with the pullback to SM of $S_M^\perp N \times (0, \frac{\pi}{2}) \rightarrow M$. The normal connection on $S_M^\perp N \rightarrow M$ coming from the Levi-Civita connection on N induces a connection on the latter space as a bundle over M . Now pull this connection back to SM via the projection to M . These two descriptions clearly fit together to give the desired connection on p . This connection now yields for each $\xi \in SM$ a canonical restriction map $r_\xi : \Omega^*(SN|_M^+) \rightarrow \Omega^*(S_\xi^+ N, \wedge^* T_\xi SM)$, such that any ϕ is determined uniquely by the family of $r_\xi(\phi)$, $\xi \in SM$.

4.2.2. *The formal model.* With Proposition 2.6, the following will imply Lemma 3.8.

Proposition 4.8. *Let k, p, m, n be given. Then there exist elements $\psi_1 \in \bar{\mathcal{A}}_{m,n}^+$, $\psi_2 \in \mathcal{B}_{m,n}^+$ such that for any smooth isometric immersion $e : M^{m+1} \hookrightarrow N^{n+1}$*

$$(68) \quad p_* i^* \bar{\rho}_N(\phi_{kp}) = \bar{\rho}_{rel}(\psi_1), \quad \pi_* j^* \bar{\rho}_N(\phi_{kp}) = \bar{\rho}_{rel}(\psi_2).$$

The proof of the first assertion is parallel to the proof of the first assertion of Proposition 4.4. Consider the sphere $S^{n-m} \subset \mathbb{R} \oplus \mathbb{R}^{n-m}$, with standard basis e_0, e_{m+1}, \dots, e_n and associated coordinates u_0, u_{m+1}, \dots, u_n . The group $O(n-m)$ acts on S_+^{n-m} by fixing the 0 coordinate. Put S_+^{n-m} for the open hemisphere $\{u_0 > 0\}$. The group $O(m) \times O(n-m)$ admits a left action $\bar{\Lambda}$ on $\Omega^*(S_+^{n-m}, \bar{\mathcal{A}}_{m,n})$ by

$$\bar{\Lambda}_{g,h} \phi := L_{g,h} \circ (h^{-1})^* \phi.$$

Denote by $\Omega^*(S_+^{n-m}, \bar{\mathcal{A}}_{m,n})^{++}$ the subspace $\{\phi : \bar{\Lambda}_{g,h} \phi = \det g \cdot \det h \cdot \phi\}$.

Lemma 4.9. *There exist canonical maps $\sigma_{rel}, \hat{\rho}_{rel}$ such that the following diagram commutes, where $J = \int_{S_+^{n-m}}$:*

$$\begin{array}{ccc} & \bar{\mathcal{A}}_n^+ & \\ & \swarrow & \searrow \bar{\rho} \\ \bar{\mathcal{A}}_{m,n}^{++} & \xrightarrow{\quad} & \Omega^*(SN) \\ \sigma_{rel} \downarrow & & \downarrow i^* \\ \Omega^*(S_+^{n-m}, \bar{\mathcal{A}}_{m,n})^{++} & \xrightarrow{\hat{\rho}_{rel}} & \Omega^*(SN|_M^+) \\ J \downarrow & & \downarrow p_* \\ \bar{\mathcal{A}}_{m,n}^+ & \xrightarrow{\bar{\rho}_{rel}} & \Omega^*(SM) \end{array}$$

Proof of Lemma 4.9. Put Q for the index set $\{0, m+1, \dots, n\}$. Let $O(n-m+1)$ act on $\mathbb{R}^Q = \mathbb{R}^{n-m+1}$ in the expected way, so that the action of the subgroup $O(n-m)$ is as described above, fixing e_0 . Let $\bar{g} : S_+^{n-m} \rightarrow SO(m) \times SO(n-m+1) \subset O(n+1)$ be a section, and define $\sigma_{rel} : \bar{\mathcal{A}}_{m,n} \rightarrow \Omega^*(S_+^{n-m}, \bar{\mathcal{A}}_{m,n})$ by putting, for $u \in S_+^{n-m}$,

$$(69) \quad \begin{aligned} \sigma_{rel}(\theta_i)|_u &:= \bar{g}_{ji}(u) \theta_j \\ \sigma_{rel}(\Omega_{ij})|_u &:= \bar{g}_{ki}(u) \bar{g}_{lj}(u) \Omega_{kl} \\ \sigma_{rel}(\omega_{ij})|_u &:= \bar{g}_{ki}(u) \bar{g}_{lj}(u) \omega_{kl} + \bar{g}_{ji}(u) du_j \\ &= \bar{g}_{ki}(u) \bar{g}_{lj}(u) \omega_{kl} + \bar{g}_{ji}(u) d\bar{g}_{j0}, \end{aligned}$$

where we take into account the relations (13). As in the remarks following the statement of Proposition 4.4, one readily sees that the restriction to $\bar{\mathcal{A}}_{m,n}^{++}$ of the resulting algebra

homomorphism is independent of the choice of \bar{g} , and that the values of the restricted map enjoy the indicated symmetries.

Applying the constructions of Section 2.3 to the connection on the bundle p described above, we may define the map $\hat{\rho}_{rel}$ by specifying a map into $\Omega^*(S_\xi^+ N, \wedge^* T_\xi SM)$ for each $\xi \in SM$. Let $b = (b_0, \dots, b_n)$ be a positive adapted frame at ξ , so that $b_0 = \xi$ and b_1, \dots, b_m are tangent to M at $x := \pi(\xi)$. Put $\psi_b : S_x^+ N \rightarrow S_+^{n-m}$ by $\psi_b^{-1}(u_0, u_{m+1}, \dots, u_n) := u_0 b_0 + u_\alpha b_\alpha$. Now put $\hat{\rho}_{b,rel}(\phi) := \bar{\rho}_{b,rel} \circ \psi_b^* \phi$. Restricting this map to $\phi \in \Omega^*(S_+^{n-m}, \bar{\mathcal{A}}_{m,n})^{++}$ yields $\hat{\rho}_{rel}$.

We now show that the top square commutes. Extend b to an adapted moving frame \bar{b} defined on a neighborhood of ξ in SM , such that $(\bar{b}_0, \dots, \bar{b}_m)$ is constructed as in the proof of Lemma 4.6 above, and $\bar{b}_{m+1}, \dots, \bar{b}_n$ depend only on the base point $\pi(\bar{b}_0)$ and all have covariant derivative zero at $\pi(\xi)$ with respect to the normal connection. Thus $\bar{\rho}_{\bar{b},rel}(\omega_{ij})|_\xi = \bar{\rho}_{\bar{b},rel}(\omega_{\alpha\beta})|_\xi = 0$, $0 \notin \{i, j\}$.

We may now use the section \bar{g} above to construct an adapted moving frame on $SN|_M^+$ in the neighborhood of $S_\xi^+ M$, by taking for $(u_0, u_{m+1}, \dots, u_n) \in S_+^{n-m}$, $\xi \in SM$,

$$\beta(u_0 \xi + u_\alpha b_\alpha) := R_{\bar{g}(u_0, u_{m+1}, \dots, u_n)} \bar{b}(\xi).$$

Let $\bar{\beta}$ be an extension to an adapted moving frame on an open subset of SN . Put $\tilde{g}(u_0 \xi + u_\alpha b_\alpha) := \bar{g}(u_0, u_{m+1}, \dots, u_n)$. We have the locally defined commutative diagram

$$(70) \quad \begin{array}{ccccccc} \mathcal{A}_n & \xrightarrow{\rho} & \Omega^*(FN) & \xrightarrow{R^*} & \Omega^*(F(M, N) \times O(n-m+1)) & \xrightarrow{r_b} & \Omega^*(O(n-m+1), \wedge^* T_b F(M, N)) \\ \uparrow & & \downarrow \bar{\beta}^* & & \downarrow (\bar{b} \circ p, \tilde{g})^* & & \downarrow \\ \bar{\mathcal{A}}_m & \xrightarrow{\bar{\rho}_{\bar{\beta}}} & \Omega^*(SN) & \xrightarrow{i^*} & \Omega^*(SN|_M^+) & \xrightarrow{r_{\xi_0}} & \Omega^*(S_{\xi_0}^+ N, \wedge^* T_{\xi_0} SM) \end{array}$$

The commutativity of the top square in the diagram in the statement of the Lemma now follows by comparing the values (69) with the results of the right action R , as in the proof of Lemma 4.5.

We now examine the bottom square. Removing the covariance conditions from the spaces on the left, and replacing the horizontal maps by $\hat{\rho}_{b,rel}, \bar{\rho}_{b,rel}$, commutativity follows from Lemma 2.7. It remains to show that J maps $\Omega^*(S_+^{n-m}, \bar{\mathcal{A}}_{m,n})^{++}$ to $\bar{\mathcal{A}}_{m,n}^+$. Taking into account the change of orientation of the fiber, we have for any $(g, h) \in O(m) \times O(n-m)$

$$(71) \quad J(\bar{\Lambda}_{g,h} \phi) = (\det h) \bar{\Lambda}_{g,h} J(\phi).$$

This implies that

$$(72) \quad (g, h) \cdot J(\bar{\phi}_{kp}) = (\det g) J(\bar{\phi}_{kp}), \quad (g, h) \in O(m) \times O(n-m),$$

which implies that J maps $\Omega^*(S_+^{n-m}, \bar{\mathcal{A}}_{m,n})^{++}$ to $\bar{\mathcal{A}}_{m,n}^+$. \square

Proof of Proposition 4.8. Insert $\phi_{kp} \in \bar{\mathcal{A}}_n^+$ into the top of the diagram in Lemma 4.9 and chase it down in both directions. Applying (69) to the terms of the expansion (7), it is clear that the only terms that yield $n-m$ factors du_β are those for which $\{m+1, \dots, n\} \subset \{\pi_{k+1}, \dots, \pi_n\}$. Thus these terms include no factors $\Omega_{\alpha t}$. Since only these terms contribute to the fiber integral J , it follows that $\psi_1 := J \circ \sigma_{rel}(\phi_{kp}) \in \bar{\mathcal{A}}_{m,n}^+$. That the bottom square of the diagram commutes is now immediate.

We omit the simpler proof of the second assertion. \square

5. AN APPLICATION TO HERMITIAN INTEGRAL GEOMETRY

We apply our results to study the space \mathcal{C}_λ^n of invariant curvature measures, and the corresponding algebra \mathcal{V}_λ^n , on the complex space forms $\mathbb{C}P_\lambda^n$.

5.1. Review of hermitian integral geometry. Let us recall the main notions of this subject, developed mostly in [10].

5.1.1. *General theory.* A *Riemannian isotropic space* is a pair (M, G) , where M is a Riemannian manifold and G is a group of isometries of M that acts transitively on the tangent sphere bundle SM . For such a space there exist kinematic formulas both for G -invariant curvature measures and for G -invariant valuations, the latter being the image of the former under the globalization map:

$$\begin{array}{ccc} \mathcal{C}^G(M) & \xrightarrow{K} & \mathcal{C}^G(M) \otimes \mathcal{C}^G(M) \\ \text{glob} \downarrow & & \downarrow \text{glob} \otimes \text{glob} \\ \mathcal{V}^G(M) & \xrightarrow{k} & \mathcal{V}^G(M) \otimes \mathcal{V}^G(M) \end{array}$$

where the coproducts K, k are cocommutative and coassociative. Picking a representative point $o \in M$ and setting $G_o \subset G$ to be the subgroup $\{g : go = o\}$, the pair $(T_oM, \overline{G_o})$ is again a Riemannian isotropic space, where $\overline{G_o}$ is the group generated by the group of translations and the derivative of the action of G_o . Howard's transfer principle ([10], Theorem 2.23) implies that the natural map $\mathcal{C}^G(M) \rightarrow \mathcal{C}^{\overline{G_o}}(T_oM)$ is an isomorphism of graded coalgebras.

By the fundamental theorem of algebraic integral geometry ([10], Theorem 2.21), the kinematic operator k is adjoint to the multiplication of $\mathcal{V}^G(M)$ via a natural Poincaré duality map $p : \mathcal{V}^G(M) \rightarrow \mathcal{V}^G(M)^*$. By the same token, the action of $\mathcal{V}^G(M)$ on $\mathcal{C}^G(M)$ encodes information about the operator K ([10], Corollary 2.20). In view of the obvious fact that the Riemannian curvature measures are invariant under isometries, we now observe:

Lemma 5.1. *If (M, G) is a Riemannian isotropic space then $\mathcal{R}(M) \subset \mathcal{C}^G(M)$ and $\mathcal{LK}(M) \subset \mathcal{V}^G(M)$.*

It follows that the realization at M of the action of $\mathbb{R}[t]$ on \mathcal{R} carries kinematic information. We now illustrate this principle by examining the case of the integral geometry of complex space forms.

5.1.2. *The case of the complex space forms.* Put $\mathbb{C}P_\lambda^n$ for the complex space form of complex dimension n and holomorphic sectional curvature 4λ , equipped with its isometry groups G_λ^n . For $\lambda = 0$, we take $G_0^n := \overline{U(n)}$, and the resulting isotropic euclidean space is canonically isomorphic to each tangent space $(T_o\mathbb{C}P_\lambda^n, (G_\lambda^n)_o)$ described in the general case above. This implies that for each n there exists a single model \mathcal{C}^n canonically isomorphic as a coalgebra to each $\mathcal{C}^{G_\lambda^n}(\mathbb{C}P_\lambda^n)$, $\lambda \in \mathbb{R}$.

By definition, an element of $\mathcal{C}^{G_\lambda^n}(\mathbb{C}P_\lambda^n)$ is angular iff the corresponding element of \mathcal{C}^n is. The space of all such elements is graded, and denoted by $\text{Ang}^n = \bigoplus_{k=0}^{2n} \text{Ang}_k^n$. The space \mathcal{C}^n decomposes as

$$\mathcal{C}^n = \text{Ang}^n \oplus \text{Null}_\lambda^n$$

where $\text{Null}_\lambda^n := \ker(\text{glob}_\lambda^n)$ (cf. [10]). The space Ang^n admits a canonical graded basis

$$\Delta_{kq} \in \text{Ang}_k^n, \quad \max(0, k-n) \leq q \leq \frac{k}{2} \leq n,$$

characterized by the property that $\Delta_{kq}(\mathbb{C}^p \oplus \mathbb{R}^{l-2p}, \cdot) = \delta_{l,p}^{k,q} \mathcal{H}^k \Big|_{\mathbb{C}^q \oplus \mathbb{R}^{k-2q}}$, where \mathbb{R}^{l-2p} denotes a subspace of \mathbb{C}^n of the given real dimension that is orthogonal both to \mathbb{C}^p and to $\sqrt{-1}$ times itself.

By Lemma 3.6 of [10] and the remarks preceding it, there are natural restriction maps $r_n : \mathcal{C}^{n+1} \rightarrow \mathcal{C}^n$ that act formally as the identity on the span of the Δ_{kp} , $k \leq n$, and that respect the decompositions above. Take $\mathcal{C}^\infty = \text{Ang}^\infty \oplus \text{Null}_\lambda^\infty$ to be the inverse limit of this system. The inclusions also induce surjective algebra homomorphisms $\mathcal{V}_\lambda^{n+1} \rightarrow \mathcal{V}_\lambda^n$, intertwining the actions on $\mathcal{C}^{n+1}, \mathcal{C}^n$ and the restriction maps r_n . The inverse limit algebra $\mathcal{V}_\lambda^\infty$ thus acts on \mathcal{C}^∞ , and includes a canonical copy of $\mathbb{R}[[t]]$ obtained as the inverse limit of the $\mathcal{LK}(\mathbb{C}P_\lambda^n)$. The generator thus gives rise to an operator on \mathcal{C}^∞ that we denote by t_λ , partially described in Theorem 6.7 of [10]. Clearly $t_\lambda(\text{Null}_\lambda^\infty) \subset \text{Null}_\lambda^\infty$.

By Proposition 3.2 and Lemma 5.1, the realization maps $\mathcal{R} \rightarrow \mathcal{C}(\mathbb{C}P_\lambda^n)$ may be represented as maps

$$\rho_\lambda^n : \mathcal{R} \rightarrow \text{Ang}^n \subset \mathcal{C}^n.$$

Applying Corollary 3.9 to the standard totally geodesic inclusions $\mathbb{C}P_\lambda^n \hookrightarrow \mathbb{C}P_\lambda^{n+1}$, we deduce that

$$(73) \quad r_n \circ \rho_\lambda^{n+1} = \rho_\lambda^n.$$

Thus there are maps $\rho_\lambda : \mathcal{R} \rightarrow \text{Ang}^\infty, r_n : \mathcal{C}^\infty \rightarrow \mathcal{C}^n$, such that $\rho_\lambda^n = r_n \circ \rho_\lambda$.

In the next section we show first that if $\lambda \neq 0$ then ρ_λ is a linear isomorphism. Therefore t_λ stabilizes Ang^∞ , by Theorem 3.13. We then give explicitly the relation between the $\rho_\lambda(C_{kp}), \Delta_{lq}$, and express the action of t_λ in terms of the latter.

5.2. The correspondence $\mathcal{R} \leftrightarrow \mathcal{C}^\infty$.

5.2.1. *The exponential generating functions for $\widetilde{\mathcal{L}\mathcal{K}}(\mathbb{C}P_\lambda^n)$.* We abbreviate

$$C_{kp}^\lambda := \rho_\lambda(C_{kp}), \quad \bar{\Lambda}_k^\lambda := \rho_\lambda(\bar{\Lambda}_k).$$

We introduce the alternative graded basis

$$(74) \quad \tilde{\Delta}_{kl} := \sum_{j \geq 0} \binom{l+j}{l} \Delta_{k,l+j}$$

so that

$$(75) \quad \Delta_{kj} := \sum_{l \geq 0} (-1)^j \binom{l+j}{l} \tilde{\Delta}_{k,l+j}.$$

Thus, in the case of complex euclidean space (viz. $\lambda = 0$), the globalizations of the $\tilde{\Delta}$ coincide with the Tasaki valuations introduced in [9].

We may express the Lipschitz-Killing curvature measures of the $\mathbb{C}P_\lambda^n$ in terms of the $\tilde{\Delta}_{kp}$ by means of exponential generating functions. Define

$$(76) \quad g_k(z, y) := \binom{2k}{k} z^k (1-4z)^{-k-\frac{1}{2}} (1-4y)^{-\frac{3}{2}}$$

$$(77) \quad h_k(z, y) := z^k (1-4z)^{-k-\frac{3}{2}} (1-4y)^{-\frac{3}{2}}$$

Proposition 5.2. *Define the graded ω -linear maps $L, M : \mathbb{R}[[z, y]] \rightarrow \text{Ang}^\infty$ by*

$$L(z^m y^p) := m!p! \tilde{\Delta}_{2m+2p,p},$$

$$M(z^m y^p) := m!p! \tilde{\Delta}_{2m+2p+1,p}$$

of degrees 0, 1 respectively. Then for any $\lambda \in \mathbb{R}$

$$(78) \quad \left(\frac{4}{\lambda}\right)^k L\left(g_k\left(\frac{\lambda z}{4\pi}, \frac{\lambda y}{4\pi}\right)\right) = \bar{\Lambda}_{2k}^\lambda,$$

$$(79) \quad \frac{2}{\pi} \left(\frac{16}{\lambda}\right)^k M\left(h_k\left(\frac{\lambda z}{4\pi}, \frac{\lambda y}{4\pi}\right)\right) = \bar{\Lambda}_{2k+1}^\lambda.$$

In other words, the $\bar{\Lambda}_j^\lambda$ may be expressed as sums of the $\tilde{\Delta}_{kp}$, with coefficients given by the values at $(0, 0)$ of appropriate partial derivatives of the functions on the left of (78), (79).

Proof. This is a straightforward modification of Lemma 3.12 of [10]. \square

The following now implies Proposition 3.15 (1).

Corollary 5.3. *Suppose $\lambda \neq 0$. Then the map $\rho_\lambda : \mathcal{R} \rightarrow \text{Ang}^\infty$ is surjective. Furthermore, if $m \leq n$, then $\rho_\lambda^n|_{\mathcal{R}_m}$ is an isomorphism onto Ang_m^n .*

Proof. By (73), it is enough to prove the second assertion. Suppose $2l := m \leq n$. Clearly $\tilde{\Delta}_{2l,0}, \dots, \tilde{\Delta}_{2l,l}$ are linearly independent. On the other hand, for $j \leq l$ the component of degree $2l$ in Λ_{2j} is $C_{2l,j-l}$, up to a nonzero factor. Thus the matrix expressing the $C_{2l,\cdot}^\lambda$ in terms of the $\tilde{\Delta}_{2l,\cdot}$ is given by the values at $(0,0)$ of $\partial_z^m \partial_y^p L'$ with $m+p=l, k \leq l$, where L' is the function on the left of (78). The entries vanish if $m < k$, and are nonzero if $m = k$. Thus the matrix is the reflection of a triangular matrix with nonzero diagonal entries. The case where m is odd is similar. \square

5.2.2. *Expressing the $\tilde{\Delta}_{kp}$ in terms of the C_{lq}^λ .* Fix $\lambda \neq 0$. Recalling the identification $\alpha : \mathcal{R} \leftrightarrow \mathbb{R}[[\xi, \eta]]$, we introduce new coordinates of degree 2 by

$$\bar{\xi} := \xi^2, \quad \bar{\eta} := \frac{\eta}{\lambda} - \bar{\xi}.$$

Take z, y both have degree 2, and define the graded ω -linear maps $\mathcal{O}, \mathcal{P} : \mathbb{R}[[z, y]] \rightarrow \mathbb{R}[[\xi, \eta]]$ by

$$\begin{aligned} \mathcal{O}(z^m y^p) &:= \pi^{m+p} \frac{\binom{m+p}{p}}{(2p+1) \binom{2p}{p} \binom{2m}{m}} \bar{\xi}^m \bar{\eta}^p, \\ \mathcal{P}(z^m y^p) &:= \left(\frac{\pi}{4}\right)^{m+p} \frac{(2m+2p+1) \binom{2m+2p}{m+p} \binom{m+p}{p}}{(2m+1)(2p+1) \binom{2m}{m} \binom{2p}{p}} \bar{\xi}^m \bar{\eta}^p. \end{aligned}$$

Clearly \mathcal{O}, \mathcal{P} intertwine the operators $z^k \partial_z^k, \bar{\xi}^k \partial_{\bar{\xi}}^k$, and by (16), (17),

$$\begin{aligned} \mathcal{O}(g_0(z, y)) &= \left(1 - \pi(\bar{\xi} + \bar{\eta})\right)^{-1}, \\ \mathcal{P}(h_0(z, y)) &= \left(1 - \pi(\bar{\xi} + \bar{\eta})\right)^{-\frac{3}{2}}. \end{aligned}$$

Since $z^k \partial_z^k g_0 = k! g_k$ and $z^k \partial_z^k h_0 = \frac{(2k+1)!}{k!} h_k$, we thus compute

$$\begin{aligned} \mathcal{O}(g_k(z, y)) &= \pi^k \bar{\xi}^k \left(1 - \pi(\bar{\xi} + \bar{\eta})\right)^{-k-1}, \\ \mathcal{P}(h_k(z, y)) &= \left(\frac{\pi}{4}\right)^k \bar{\xi}^k \left(1 - \pi(\bar{\xi} + \bar{\eta})\right)^{-k-\frac{3}{2}}, \end{aligned}$$

and

$$\begin{aligned} \mathcal{O}\left(\left(\frac{4}{\lambda}\right)^k g_k\left(\frac{\lambda z}{4\pi}, \frac{\lambda y}{4\pi}\right)\right) &= \xi^{2k} \left(1 - \frac{\eta}{4}\right)^{-k-1} = \alpha(\bar{\Lambda}_{2k}), \\ \xi \cdot \mathcal{P}\left(\left(\frac{16}{\lambda}\right)^k h_k\left(\frac{\lambda z}{4\pi}, \frac{\lambda y}{4\pi}\right)\right) &= \xi^{2k+1} \left(1 - \frac{\eta}{4}\right)^{-k-\frac{3}{2}} = \alpha(\bar{\Lambda}_{2k+1}). \end{aligned}$$

Proposition 5.2 now implies that $f := \rho_\lambda \circ \alpha^{-1} \circ \mathcal{O} \circ L^{-1}$ maps each $\bar{\Lambda}_{2k}^\lambda$ to itself, and that $g := \rho_\lambda \circ \alpha^{-1} \circ m_{\frac{\pi\xi}{2}} \circ \mathcal{P} \circ M^{-1}$ maps each $\bar{\Lambda}_{2k+1}^\lambda$ to itself, where $m_{\frac{\pi\xi}{2}}$ denotes multiplication by $\frac{\pi\xi}{2}$. As graded maps, f, g act also as the identity on the graded components of these elements. But by the proof of Corollary 5.3, these graded components constitute an ω -basis for $\text{Ang}_{\text{even}}^\infty$ and $\text{Ang}_{\text{odd}}^\infty$, respectively. Therefore f, g are the identity maps on these spaces. In other words, putting

$$c_{mp} := \frac{(m+p)!}{(2m)!(2p+1)!}, \quad d_{mp} := \frac{(2m+2p+1)!}{(m+p)!(2m+1)!(2p+1)!},$$

we obtain

$$\begin{aligned}
(80) \quad \tilde{\Delta}_{2m+2p,p} &= \pi^{m+p} c_{mp} \rho_\lambda \circ \alpha^{-1} (\bar{\xi}^m \bar{\eta}^p) \\
&= \pi^{m+p} c_{mp} \rho_\lambda \circ \alpha^{-1} \left(\xi^{2m} \left(\frac{\eta}{\lambda} - \xi^2 \right)^p \right) \\
&= \pi^{m+p} \lambda^{-p} c_{mp} \sum_{j=0}^p (-\lambda)^j \binom{p}{j} C_{2m+2p,p-j}^\lambda.
\end{aligned}$$

and similarly

$$\begin{aligned}
(81) \quad \tilde{\Delta}_{2m+2p+1,p} &= \frac{\pi}{2} \left(\frac{\pi}{4} \right)^{m+p} d_{mp} \rho_\lambda \circ \alpha^{-1} (\xi \bar{\xi}^m \bar{\eta}^p) \\
&= \frac{\pi}{2\lambda^p} \left(\frac{\pi}{4} \right)^{m+p} d_{mp} \sum_{j=0}^p (-\lambda)^j \binom{p}{j} C_{2m+2p+1,p-j}^\lambda
\end{aligned}$$

The relations (80), (81) admit the common form

$$(82) \quad \tilde{\Delta}_{kp} = \left(\frac{\pi}{2} \right)^{\lceil \frac{k}{2} \rceil} \frac{k!!}{(k-2p)!(2p+1)!} \sum_{j=0}^p (-1)^{p-j} \lambda^{-j} \binom{p}{j} C_{kj}^\lambda.$$

This expression is easily inverted:

$$(83) \quad C_{kj}^\lambda = \left(\frac{2}{\pi} \right)^{\lceil \frac{k}{2} \rceil} \frac{\lambda^j}{k!!} \sum_{p=0}^j \binom{j}{p} (k-2p)!(2p+1)! \tilde{\Delta}_{kp}.$$

5.2.3. *Action of t_λ .* We may now use (80) and Theorem 3.13 to compute

$$t_\lambda \cdot \tilde{\Delta}_{2m+2p,p} = \pi^{m+p} c_{mp} \rho_\lambda \circ \alpha^{-1} \left(\xi \left(1 - \frac{\eta}{4} \right)^{-\frac{1}{2}} \bar{\xi}^m \bar{\eta}^p \right)$$

where

$$\begin{aligned}
\left(1 - \frac{\eta}{4} \right)^{-\frac{1}{2}} \bar{\xi}^m \bar{\eta}^p &= \left(1 - \frac{\lambda}{4} (\bar{\eta} + \bar{\xi}) \right)^{-\frac{1}{2}} \bar{\xi}^m \bar{\eta}^p \\
&= \sum_{n \geq m, q \geq p} \left(\frac{\lambda}{16} \right)^{n+q-m-p} \binom{2n+2q-2m-2p}{n+q-m-p} \binom{n+q-m-p}{n-m} \bar{\xi}^n \bar{\eta}^q
\end{aligned}$$

so that, by (81),

$$\begin{aligned}
(84) \quad t_\lambda \cdot \tilde{\Delta}_{2m+2p,p} &= \\
&= \frac{2c_{mp}}{\pi} \left(\frac{16\pi}{\lambda} \right)^{m+p} \sum_{n \geq m, q \geq p} d_{nq}^{-1} \left(\frac{\lambda}{4\pi} \right)^{n+q} \binom{2n+2q-2m-2p}{n+q-m-p} \binom{n+q-m-p}{n-m} \tilde{\Delta}_{2n+2q+1,q}
\end{aligned}$$

Similarly,

$$\begin{aligned}
(85) \quad t_\lambda \cdot \tilde{\Delta}_{2m+2p+1,p} &= \\
&= \frac{d_{mp}}{2} \left(\frac{4\pi}{\lambda} \right)^{m+p} \sum_{n \geq m, q \geq p} c_{n+1,q}^{-1} \left(\frac{\lambda}{16\pi} \right)^{n+q} \binom{2n+2q-2m-2p}{n+q-m-p} \binom{n+q-m-p}{n-m} \tilde{\Delta}_{2n+2q+2,q}
\end{aligned}$$

Theorem 5.4. $t_\lambda \cdot \tilde{\Delta}_{kp} =$

$$\frac{2\omega_{k-1}}{\omega_k} \frac{(k-1)!!}{(k-2p)!(2p+1)!} \sum_{l \geq 0, q \geq p} \frac{(k+2l-2q+1)!(2q+1)!}{(k+2l+1)!} \left(\frac{\lambda}{8\pi} \right)^l \binom{2l}{l} \binom{l}{q-p} \tilde{\Delta}_{k+2l+1,q}$$

Proof. Substituting $n + q = m + p + l$ and using the fact that

$$\frac{2\omega_{k-1}(k-1)!!}{\omega_k k!!} = \begin{cases} \frac{2}{\pi} & \text{if } k \text{ is even} \\ 1 & \text{if } k \text{ is odd} \end{cases}$$

this expression for $t \cdot \tilde{\Delta}_{kp}$ follows immediately from (84), (85). □

REFERENCES

- [1] S. Alesker, *Algebraic structures on valuations, their properties and applications*, Higher Ed. Press, 2002, Proceedings of the International Congress of Mathematicians, Vol. II (Beijing, 2002).
- [2] ———, *Theory of valuations on manifolds. I. Linear spaces*, Israel J. Math. **156** (2006), 311–339.
- [3] ———, *Theory of valuations on manifolds. II*, Adv. Math. **207** (2006), 420–454.
- [4] ———, *Theory of valuations on manifolds: a survey.*, Geom. Funct. Anal. **17** (2007), 1321–1341.
- [5] S. Alesker and A. Bernig, *The product on smooth and generalized valuations*, Amer. J. Math. **134** (2012), 507–560.
- [6] S. Alesker and J.H.G. Fu, *Theory of valuations on manifolds. III. Multiplicative structure in the general case*, Trans. Amer. Math. Soc. **360** (2008), 1951–1981.
- [7] S. Alesker and J.H.G. Fu, *Integral geometry and valuations* (E. Gallego and G. Solanes, eds.), Advanced Courses in Mathematics. CRM Barcelona, Birkhäuser/Springer, Basel, 2014, pp. viii+112.
- [8] A. Bernig and L. Bröcker, *Valuations on manifolds and Rumin cohomology*, J. Differential Geom. **75** (2003), 433–457.
- [9] A. Bernig and J.H.G. Fu, *Hermitian integral geometry*, Ann. of Math. (2) **173** (2011), 907–945.
- [10] A. Bernig, J.H.G. Fu, and G. Solanes, *Integral geometry of complex space forms*, Geom. Funct. Anal. **24** (2014), 403–492.
- [11] ———, *Dual curvature measures in hermitian integral geometry*, Analytic aspects of Convexity (G. Bianchi, A. Colesanti, and P. Gronchi, eds.), INdAM-Springer Series, Springer, to appear.
- [12] A. Bernig and F. Voide, *Spin(9)-invariant valuations on the octonionic plane*, Israel J. Math. **214** (2016), 831–855.
- [13] R. L. Bishop and R. J. Crittenden, *Geometry of manifolds*, Pure and Applied Mathematics, Vol. XV, Academic Press, New York-London, 1964.
- [14] J. Cheeger, R. Müller, and R. Schrader, *On the curvature of piecewise flat spaces*, Comm. Math. Phys. **92** (1984), 405–454.
- [15] S.S. Chern, *A simple intrinsic proof of the Gauss-Bonnet formula for closed Riemannian manifolds*, Ann. of Math. **45** (1944), 747–752.
- [16] ———, *On the curvatura integra in a Riemannian manifold*, Ann. of Math. **46** (1945), 674–684.
- [17] H. Federer, *Curvature measures*, Trans. Amer. Math. Soc. **93** (1959), 418–491.
- [18] J. H. G. Fu, *Some remarks on Legendrian rectifiable currents*, Manuscripta Math. **97** (1998), 175–187.
- [19] ———, *Curvature measures of subanalytic sets*, Amer. J. Math. **116** (1994), 819–880.
- [20] ———, *Intersection theory and the Alesker product*, Indiana Univ. Math. J. **65** (2016), 1347–1371.
- [21] ———, *Integral geometric regularity*, Tensor valuations and their Applications in Stochastic Geometry and Imaging (E.B.V. Jensen and M. Kiderlen, eds.), Lecture Notes in Mathematics, vol. 2177, 2017, pp. 266–290.
- [22] J. H. G. Fu and C. McCrory, *Stiefel-Whitney classes and the conormal cycle of a singular variety*, Trans. Amer. Math. Soc. **349** (1997), 809–835.
- [23] H. Hadwiger, *Vorlesungen über Inhalt, Oberfläche und Isoperimetrie*, Springer, Berlin, 1957.
- [24] D. Klain and G.-C. Rota, *Introduction to geometric probability*, Pure and Applied Mathematics, Vol. XV, Cambridge U. Press, 1994.
- [25] C. Procesi, *Lie groups*, Universitext, Springer, New York, 2007, An approach through invariants and representations.
- [26] M. Rumin, *Formes différentielles sur les variétés de contact*, J. Differential Geom. **39** (1994), 281–330.
- [27] H. Weyl, *On the volume of tubes*, Duke Math. J. **61** (1939), 461–472.
- [28] M. Zähle, *Curvature and currents for unions of sets with positive reach*, Geom. Dedicata **23** (1987), 155–171.

E-mail address: joefu@uga.edu

E-mail address: thomas.wannerer@uni-jena.de