LOCALLY CONFORMALLY SYMPLECTIC AND KÄHLER GEOMETRY

GIOVANNI BAZZONI

ABSTRACT. The goal of this note is to give an introduction to locally conformally symplectic and Kähler geometry. In particular, Sections 1 and 3 aim to provide the reader with enough mathematical background to appreciate this kind of geometry. The reference book for locally conformally Kähler geometry is [36] by Sorin Dragomir and Liviu Ornea. Many progresses in this field, however, were accomplished after the publication of this book, hence are not contained there – see the introduction of [97]. On the other hand, there is no book on locally conformally symplectic geometry and many recent advances lie scattered in the literature. Sections 2 and 4 would like to demonstrate how these geometries can be used to give precise mathematical formulations to ideas deeply rooted in classical and modern Physics.

1. SYMPLECTIC AND LOCALLY CONFORMALLY SYMPLECTIC GEOMETRY

A symplectic manifold is a smooth manifold M^{2n} with a 2-form $\omega \in \Omega^2(M)$ which is nondegenerate, i. e. $\omega_p^n \neq 0$ for every $p \in M$, and closed, i. e. $d\omega = 0$. The non-degeneracy condition can be rephrased by saying that ω provides an isomorphism of vector bundles $b^{\flat}: TM \to T^*M, X \mapsto X^{\flat} = \iota_X \omega$.

The word *symplectic* was coined by Hermann Weyl in 1939: he replaced the old terminology *complex group* with *symplectic group* to indicate the Lie group of matrices preserving the bilinear skew-symmetric form $\omega_0 = \sum_{i=1}^n dx_i \wedge dy_i$ on \mathbb{R}^{2n} , see [136, Page 165]. The etymology is from the Greek $\sigma \upsilon \mu \pi \lambda \varepsilon \kappa \tau \iota \kappa \delta \varsigma$, which actually means *complex*.

In the process of getting acquainted with symplectic geometry, something that one experiences quite early is, paraphrasing Gromov, a *curious mixture of "hard" and "soft"*, see [58] as well as [90, Page 81]. This applies both to the mathematical aspects and to the techniques employed in symplectic geometry. An indication of the soft side of symplectic geometry is certainly *Darboux theorem*, asserting that, locally, two symplectic manifolds can not be distinguished from one another¹ – see for instance [9, Section 8.43] or [90, Theorem 3.15] for a modern proof. Thus, symplectic geometry is somehow a global thing. Of the two conditions ensuring that a 2-form on an even-dimensional manifold is symplectic, however, only one is of global nature, namely closedness. Closedness imposes strong cohomological restrictions on the existence of a symplectic structure on an even-dimensional compact manifold²: for instance, all Betti numbers of even-degree must be non-zero. The general problem of determining which compact manifolds admit a symplectic structure is far from being solved, see [112].

¹This is very different from the Riemannian case, where curvature provides a local invariant.

 $^{^{2}}$ The same is not true for open manifolds: as proved by Gromov [56, 57], any open manifold with a non-degenerate 2-form admits a symplectic structure.

The first true mathematical exposition of what a symplectic manifold is appeared in a paper of Hwa-Chung Lee in 1941, see [77]. Lee considers the general setting of an even-dimensional manifold M^{2n} endowed with a non-degenerate 2-form ω . He studies first the *flat* case, in which $d\omega = 0$, that is, what is nowadays known as symplectic. Then, he discusses the problem of two 2-forms ω and ω' which are *conformal* to one another: on an open set $U \subset M$ with local coordinates (x_1, \ldots, x_{2n}) , write

$$\boldsymbol{\omega} = \sum_{i < j} \boldsymbol{\omega}_{ij}(x) dx_i \wedge dx_j$$
 and $\boldsymbol{\omega}' = \sum_{i < j} \boldsymbol{\omega}'_{ij}(x) dx_i \wedge dx_j;$

 ω and ω' are (locally) conformal if there exists $\varphi \in \mathscr{C}^{\infty}(U)$, nowhere vanishing, with $\omega'_{ij} = \varphi \omega_{ij}$. Lee then finds necessary and sufficient conditions for a given $\omega \in \Omega^2(M)$ to be (locally) conformal to a flat, i. e. closed, one: for $n \ge 3$ this happens³ if and only if there exists a 1-form ϑ such that $d\omega = \vartheta \wedge \omega$. It is interesting to notice that the mathematical birthplace of *both* symplectic and locally conformally symplectic geometry is the very same paper of Lee!

The development of symplectic geometry since 1941 was dramatic, kept up first by the French school (Charles Ehresmann, Paulette Libermann, André Lichnerowicz, Georges Reeb) in the 1950's, then by the Russian school, with the central figure of Vladimir Arnol'd, and by the American school (Dusa McDuff, Victor Guillemin, Alan Weinstein); another special place is occupied by Mikhaïl Gromov⁴. This is however not the right place to extol the ubiquity of symplectic geometry in modern Mathematics – I refer the reader to the nice surveys [10, 50, 89].

The fate of locally conformally symplectic geometry, on the contrary, was very different. Except for works of Libermann in 1955 [80] and Jean Lefebvre in 1966 and 1969 [78, 79], the subject remained in hibernation until the seminal papers of Izu Vaisman: *On locally conformal almost Kähler manifolds*, published in 1976 – see [123], and *Locally conformal symplectic manifolds*, published in 1984 – see [128].

In [123], Vaisman defines a *locally conformally symplectic manifold*⁵ as a manifold M^{2n} , $n \ge 1$, endowed with a non-degenerate 2-form $\omega \in \Omega^2(M)$ such that every point $p \in M$ has an open neighborhood U such that

(1)
$$d\left(e^{\sigma}\omega\right|_{U}\right) = 0,$$

where $\sigma: U \to \mathbb{R}$ is a smooth function. If (1) holds for U = M, then (M, ω) is globally conformally symplectic; if it holds for σ a constant function, (M, ω) is clearly a symplectic manifold. The work of Lee shows that the above definition is equivalent to the following one: a manifold M^{2n} , $n \ge 1$, endowed with a non-degenerate 2-form $\omega \in \Omega^2(M)$, is locally conformally symplectic manifold if there exists a globally defined 1-form $\vartheta \in \Omega^1(M)$ such that

(2)
$$d\omega = \vartheta \wedge \omega \text{ and } d\vartheta = 0.$$

³The case n = 1 is trivial: as remarked by Lee, every ω is in this case conformal to a flat one, due to dimension reasons. The case n = 2 is only slightly different – see the discussion below.

⁴The list of quoted mathematicians is of course far from being complete.

⁵Vaisman uses locally *conformal* symplectic, while I stick with the terminology locally *conformally* symplectic in this note. Some recent papers use *conformal symplectic*, see [30, 37].

The 1-form ϑ was baptized the *Lee form* by Vaisman. If n = 1 one has $d\omega = 0 = \vartheta \land \omega$ for any choice of ϑ . For $n \ge 2$, ϑ is completely determined by ω ; moreover, as remarked by Libermann in [80], the second condition in (2) follows from the first one if $n \ge 3$. (ω, ϑ) is called a locally conformally symplectic structure on M. According to this alternative definition, a locally conformally symplectic manifold is globally conformally symplectic if ϑ is exact and symplectic if $\vartheta = 0$.

Given a locally conformally symplectic manifold (M, ω) , the *conformal class* of ω is

$$\{\omega' \in \Omega^2(M) \mid \exists f \in C^{\infty}(M) \mid \omega' = e^f \omega\}.$$

If ϑ is the Lee form of (M, ω) and $\omega' = e^f \omega$, then the Lee form of (M, ω') is $\vartheta' = \vartheta + df$, hence the cohomology class of ϑ in $H^1_{dR}(M)$ is an invariant of the conformal class.

Formula (1) implies, in particular, that at a local scale a symplectic manifold can not be distinguished from a locally conformally symplectic manifold. Thus not only all symplectic manifolds locally look alike, in view of Darboux theorem, but potentially there may exist manifolds which locally look like symplectic manifolds and however fail to do so globally! Locally conformally symplectic structures exist on open manifolds, as proved by Fernandes and Frejlich using an *h*-principle – see [39], in particular the Acknowledgements. It was proved very recently by Eliashberg and Murphy using again *h*-principle that a *closed* almost complex manifold (M,J) with a non zero cohomology class $\mu \in H^1(M; \mathbb{R})$ admits a locally conformally symplectic structure – see [37, Theorem 1.8] for the precise statement. In [4, 20] explicit examples of compact locally conformally symplectic manifolds which do not admit any symplectic structure are provided.

For this reason, I prefer to consider locally conformally symplectic manifolds as something different from symplectic manifolds. Concretely, this means that our locally conformally symplectic structures will always be assumed to have a Lee form ϑ which is not exact.

In his 1976 paper Vaisman proves a few results about locally conformally symplectic manifolds but turns quickly his attention to the metric case, in the wake of Gray's work on almost Hermitian structures. It is in his 1984 article that he extensively studies the non-metric case. Motivated by the metric case, which I will discuss in Section 3, Vaisman distinguishes between locally conformally symplectic structures of the first kind and of the second kind. A locally conformally symplectic structure (ω, ϑ) on M is of the first kind if there exists a vector field $U \in \mathfrak{X}(M)$ such that

$$\mathscr{L}_U \boldsymbol{\omega} = 0$$
 and $\vartheta(U) = 1$.

Otherwise, it is of the second kind. The above conditions characterize U uniquely; it is the *Lee field* of the locally conformally symplectic structure. A sophisticated way to rephrase this goes as follows: define

(3)
$$\mathfrak{X}(M,\omega) = \{X \in \mathfrak{X}(M) \mid \mathscr{L}_X \omega = 0\};$$

then $\mathfrak{X}(M, \omega) \subset \mathfrak{X}(M)$ is a subalgebra. If $X \in \mathfrak{X}(M, \omega)$ then $\mathscr{L}_X \vartheta = 0$, hence $\vartheta(X)$ is a constant function on M. The *Lee morphism* is $\ell \colon \mathfrak{X}(M, \omega) \to \mathbb{R}$, $\ell(X) = \vartheta(X)$ and is a morphism of Lie algebras. Thus (ω, ϑ) is of the first kind if and only if the Lee morphism is non zero, hence surjective; of the second kind otherwise. In particular, the Lee form of a locally conformally symplectic structure of the first kind is nowhere zero. I should remark here that in the conformal class of a locally conformally symplectic structure of the second kind. To see this, it is enough

to choose a function f such that $df_p = -\vartheta_p$ for some $p \in M$; then the Lee form of $e^f \omega$ will have a zero. Notice that (3) defines an automorphism of a given element in the conformal class of a locally conformally symplectic structure. If one wants to deal with the whole conformal class, then the object to be considered is the subalgebra

$$\mathfrak{X}(M,\omega) = \{ X \in \mathfrak{X}(M) \mid \exists f_X \in \mathscr{C}^{\infty}(M) \mid \mathscr{L}_X \omega = f_X \omega \};$$

here f_X should be nowhere 0. In this case as well one sees that the *extended Lee morphism* $\hat{\ell}: \hat{\mathfrak{X}}(M, \omega) \to \mathbb{R}, \hat{\ell}(X) = \vartheta(X) + f_X$ is a morphism of Lie algebras (see [14]). The Lee morphism and its extended version have been investigated extensively, see for instance [14, 62, 128].

Another way to tell locally conformally symplectic structures apart is according to the Morse-Novikov class of the 2-form ω . Given a 1-form ϑ on a manifold M, one can define a differential operator $d_{\vartheta} \colon \Omega^k(M) \to \Omega^{k+1}(M)$ by setting $d_{\vartheta}\sigma = d\sigma - \vartheta \wedge \sigma$. If ϑ is closed, then $d_{\vartheta}^2 = 0$ and the *Morse-Novikov*⁶ cohomology of $(\Omega^*(M), d_{\vartheta})$ is

$$H^k_{\vartheta}(M) = \frac{\ker\{d_{\vartheta} \colon \Omega^k(M) \to \Omega^{k+1}(M)\}}{d_{\vartheta}(\Omega^{k-1}(M))}$$

If *M* is compact, these cohomology spaces are always finite-dimensional, and $H^*_{\vartheta}(M) \cong H^*_{dR}(M)$ if ϑ is exact. Further, as noticed in [12], the Euler-Poincaré characteristic of the Morse-Novikov cohomology of a compact, orientable manifold equals that of the de Rham cohomology, hence it is topological. In general, however, Morse-Novikov cohomology behaves very differently from de Rham cohomology: indeed, if ϑ is not exact and *M* is connected then $H^0_{\vartheta}(M) = 0$, see [59]; if, in addition, M^n is compact and orientable, then a Poincaré duality holds, that is, $H^i_{\vartheta}(M) \cong H^{n-i}_{\vartheta}(M)^*$, hence $H^n_{\vartheta}(M) = 0$, see [61]. In [126] Vaisman proved that $H^*_{\vartheta}(M)$ is isomorphic to the cohomology of *M* with coefficients in the sheaf of smooth functions $f \in \mathscr{C}^{\infty}(M)$ which satisfy $d_{\vartheta}f = 0$. It was proved in [34] that if *M* carries a Riemannian metric for which ϑ is parallel, then $H^*_{\vartheta}(M) = 0$. Aside from these general results, the computation of Morse-Novikov cohomology is in general very difficult. For a nilmanifold or a completely solvable solvmanifold⁷ the computation of the Morse-Novikov cohomology can be performed algebraically – see [3, 5, 91, 93]. For more details on the Morse-Novikov cohomology cohomology, I refer the reader to [17, 34, 38, 61, 62].

The significance of Morse-Novikov cohomology in the context of locally conformally symplectic geometry stems from (2): if (M, ω, ϑ) is a locally conformally symplectic manifold then $d\vartheta = 0$ and $d_{\vartheta}\omega = d\omega - \vartheta \wedge \omega = 0$, hence the 2-form ω defines a cohomology class $[\omega]_{\vartheta} \in H^2_{\vartheta}(M)$. The locally conformally symplectic structure is *exact* if $[\omega]_{\vartheta} = 0$, *non exact* otherwise. It is easy to see that a locally conformally symplectic structure of the first kind is exact: by defining $\eta = -\iota_U \omega$, where U is the Lee field, one has $\omega = d\eta - \vartheta \wedge \eta$. The converse

⁶The Morse-Novikov cohomology has more than two fathers. In the context of locally conformally symplectic geometry, for instance, it was first considered by Guédira and Lichnerowicz in [59]. It was also considered by Witten in his celebrated paper [137].

⁷A *nilmanifold* is the quotient of a connected, simply connected nilpotent Lie group by a discrete and cocompact subgroup. More generally, a *solvmanifold* is a compact quotient of a connected, simply connected solvable Lie group. A solvmanifold is *completely solvable* if the adjoint representation on the corresponding Lie algebra has only real eigenvalues.

does not hold: in fact, being exact is an invariant of the conformal class of a locally conformally symplectic structure, while being of the first kind is not, as it was shown above. The locally conformally symplectic structures constructed by Eliashberg and Murphy in [37] are exact. The importance of Morse-Novikov cohomology in the context of locally conformally symplectic geometry is highlighted, for instance, by the recent research papers [5, 76, 107].

Locally conformally symplectic structures of the first kind are strictly related to contact structures. A (*co-orientable*) contact structure on an odd-dimensional manifold P^{2n+1} ($n \ge 1$) consists of a 1-form α such that $\alpha \wedge (d\alpha)^n \neq 0$ at every point, see [46]. Frobenius' integrability theorem shows that the distribution $\xi = \ker \alpha$ is then maximally non-integrable. Let (P, α) be a contact manifold and consider a *strict contactomorphism*, that is, a diffeomorphism $\varphi : P \rightarrow P$ satisfying $\varphi^* \alpha = \alpha$. Then, as observed for instance by Banyaga in [16], the mapping torus⁸ P_{φ} admits a locally conformally symplectic structure of the first kind. In the same paper, Banyaga proves a sort of converse to this result: namely, if a compact manifold M is endowed with a locally conformally symplectic structure, then there exist a compact contact manifold (P, α) and a strict contactomorphism $\varphi : P \rightarrow P$ such that M is diffeomorphic to the mapping torus P_{φ} . Banyaga's result, however, does not claim that the original locally conformally symplectic structure is preserved, is proved in [21].

It is interesting to notice that contact and locally conformally symplectic structures come together also in the context of Jacobi structures. According to [59], indeed, a *transitive* Jacobi manifold is a contact manifold if the dimension is odd and a locally conformally symplectic manifold if it is even.

Locally conformally symplectic structures of the second kind are much less understood. Concerning, in particular, non exact structures, Banyaga [17] proved that there exist two families of locally conformally symplectic structures on the 4-dimensional solvmanifold constructed in [33] and that they are non exact. These are the first acknowledged examples of this type of locally conformally symplectic structures. In [87, Appendix A] it was shown that the locally conformally symplectic structure of the Oeljeklaus-Toma manifolds constructed in [95] is not exact. In [4] the properties of non exact locally conformally symplectic structures extensively are investigated, producing many new examples.

I conclude this section with a collection of results in locally conformally symplectic geometry.

The problem of reduction in locally conformally symplectic geometry was tackled in [62, 87, 94]. In [87] the authors also produce *universal models* for exact locally conformally symplectic manifolds, on the line of Tischler's result on universal models for symplectic manifolds, see [119]. A *Moser trick* for locally conformally symplectic forms was proved in [12]. The blow-up of a locally conformally symplectic manifold at a point or along a compact symplectic submanifold, i. e. a submanifold such that the locally conformally symplectic form restricts to a *closed* form, was constructed in [31, 139]. The notion of *Lagrangian submanifold*

⁸Given a topological space X and a homeomorphism $\varphi : X \to X$, the mapping torus or suspension X_{φ} is the quotient space of $X \times \mathbb{R}$ by the \mathbb{Z} -action generated by $1 \cdot (x,t) = (\varphi(x), t+1)$. The projection $\pi : X_{\varphi} \to S^1$, $[(x,t)] \mapsto [t]$ is a fiber bundle with fiber X. If M is a smooth manifold and φ is a diffeomorphism, then M_{φ} is a smooth manifold and $M \to M_{\varphi} \to S^1$ is a smooth fiber bundle.

makes perfectly sense in the locally conformally symplectic setting. A result on neighbourhoods of Lagrangian submanifolds in locally conformally symplectic manifolds was obtained in [109], analogous to the known result of Weinstein in the symplectic case, [134]. The problem of displacing a Lagrangian submanifold in a locally conformally symplectic manifold is tackled in [30]. The paper also contains some interesting observations on the issues that appear when one tries to apply Floer's machinery or results such as Gromov compactness to the locally conformally symplectic situation. Such issues depend, essentially, on the fact that ω is not closed, hence no bound à *la Gromov* on the energy of a *J*-holomorphic map is possible. The paper [113] suggests some ideas on how to control the failure of Gromov compactness. The properties of the group of diffeomorphisms preserving the conformal class of a locally conformally symplectic structure are studied in [62] – see also [14, 15, 79]. Finally, for a description of locally conformally symplectic structures in the language of Lie algebroids as well as some generalizations I refer the reader to the papers [67, 68].

2. CLASSICAL MECHANICS

Now those Quantities which I consider as gradually and indefinitely increasing, I shall hereafter call *Fluents*, or *Flowing Quantities*, [...] And the Velocities by which every Fluent is increased by its generating Motion, (which I may call *Fluxions*, or simply Velocities or Celerities,) [...]

The Relation of the Flowing Quantities to one another being given, to determine the Relation of their Fluxions.

A relation being proposed, including the Fluxions of Quantities, to find the Relations of those Quantities to one another.

Sir Isaac Newton, "De methodis serierum et fluxionum", 1671.⁹

The three sentences of Newton define the objects of interest and summarize the goals of the study of dynamical systems. It was Newton who gave a mathematically precise definition of the three laws that govern classical mechanics, that is, the study of the movement of a body as a response to being exposed to a force. He developed a theory, called in his honor *Newtonian mechanics*, to state and solve the problems posed by classical mechanics, notably arising from planetary motions. In this formalism, the equations of motion of a physical system with n degrees of freedom are given as solutions of n differential equations involving velocities and their derivatives (that is, differential equations of order 2).

Analytical techniques in the study of the problems of classical mechanics, especially celestial mechanics, were brought in by Lagrange at the beginning of the 19th century, founding what is nowadays known as *Lagrangian formalism*; an important role in this formalism is played by the *principle of minimal action*. In particular, as recalled by Weinstein in [135], in his 1808 book *Mémoire sur la théorie des variations des éléments des planètes*, Lagrange uses explicitly a certain skew-symmetric 6 by 6 matrix. The appearence of geometric techniques

⁹I am grateful to Prof. Antonio Giorgilli for having written amazing lecture notes for the Mathematical Physics courses he taught at University of Milano-Bicocca. As an undergrad I was lucky enough to attend a few of them, a very fruitful experience. His lecture notes contained, among other things, this reference to Newton's original work – see http://www.mat.unimi.it/users/antonio/meccanica/meccanica.html.

in classical mechanics is due to Hamilton, who rewrote Newton's equations as a set of 2*n* differential equations of order 1. In terms of position coordinates (q_1, \ldots, q_n) and corresponding momenta (p^1, \ldots, p^n) , the motions are governed by a function $H = H(q_1, \ldots, q_n, p^1, \ldots, p^n)$, the *Hamiltonian* of the system, through the equations

(4)
$$\begin{cases} \dot{q}_i = \frac{\partial H}{\partial p^i} \\ \dot{p}^i = -\frac{\partial H}{\partial q_i} \end{cases}$$

Nowadays¹⁰ it is known that, in the simplest case, the phase space of a Hamiltonian system is the cotangent bundle T^*Q of a manifold Q which parametrizes the positions q of the physical system; the corresponding momenta p live on the fibers of the cotangent bundle over a point $q \in Q$ and the Hamiltonian of the system is $H \in C^{\infty}(T^*Q)$. T^*Q is in a natural way a symplectic manifold; the symplectic form on T^*Q is very easy to describe: if $\pi: T^*Q \to Q$ is the canonical projection, define the *Liouville* or *tautological* 1-form $\lambda_{can} \in \Omega^1(T^*Q)$ by $\lambda_{(q,p)}(v) = p(d\pi_{(q,p)}(v))$ for a tangent vector v at $T_{(q,p)}T^*Q$. Then $\omega_{can} = -d\lambda_{can}$ is a symplectic form on T^*Q ; in local coordinates, one has $\omega_{can} = \sum_{i=1}^n dq_i \wedge dp^i$. In the Hamiltonian formalism, the equations of motions (4) are given as integral curves of the Hamiltonian vector field X_H ; if $H: T^*Q \to \mathbb{R}$ is the Hamiltonian function of the system, then X_H is uniquely determined by the condition $dH = \omega_{can}(X_H, \cdot)$.

From the point of view of Hamiltonian formalism, the fact that non-degeneracy is a local condition implies that the definition of the Hamiltonian vector field is local. Following the illuminating introduction of Vaisman's paper [123], I propose to show that locally conformally symplectic manifolds provide an adequate and more general context for Hamiltonian mechanics. One can make the *Ansatz* that the dynamics on the phase space consists of the orbits of a globally defined vector field X. Consider an open set $U_{\alpha} \subset T^*Q$ with local coordinates $(q_1^{\alpha}, \ldots, q_n^{\alpha}, p_{\alpha}^1, \ldots, p_{\alpha}^n)$. Then one obtains a local function $H_{\alpha}: U_{\alpha} \to \mathbb{R}$ such that the orbits of X are defined by a local version of Hamilton's equations,

(5)
$$\begin{cases} \dot{q}_{i}^{\alpha} = \frac{\partial H_{\alpha}}{\partial p_{\alpha}^{i}} \\ \dot{p}_{\alpha}^{i} = -\frac{\partial H_{\alpha}}{\partial q_{\alpha}^{i}} \end{cases}$$

Of course, X is the Hamiltonian vector field of the local Hamiltonian function H_{α} with respect to the local symplectic form $\omega_{can}^{\alpha} = \sum_{i=1}^{n} dq_{i}^{\alpha} \wedge dp_{\alpha}^{i}$. Suppose $\{U_{\alpha}\}_{\alpha}$ is an open covering of $T^{*}Q$. One the usually requires $\{\omega_{can}^{\alpha}\}$ and $\{H_{\alpha}\}$ to piece together to a global symplectic form ω_{can} and a global Hamiltonian H. However, following our *Ansatz*, in order to globalize this local assertion one only needs to prescribe the fact that the transition functions

(6)
$$q_i^\beta = q_i^\beta (q_j^\alpha, p_\alpha^k) \text{ and } p_\beta^i = p_\beta^i (q_j^\alpha, p_\alpha^k)$$

on $U_{\alpha} \cap U_{\beta}$ preserve (5). Of course, if (6) are *canonical* transformations of the phase space, then $\omega_{can}^{\alpha} = \omega_{can}^{\beta}$ and one is back to the symplectic context. However, allowing a *homothetical* change of coordinates, i. e. taking $H_{\beta} = \mu_{\beta\alpha} H_{\alpha}$ for a constant $\mu_{\beta\alpha} \neq 0$, then $\omega_{can}^{\alpha} = \mu_{\beta\alpha} \omega_{can}^{\beta}$.

¹⁰One could name here many other scientists who contributed to elaborate thorough foundations for classical mechanics – I prefer to direct the reader to the much more complete references [1, 9, 32, 50] for further historical and mathematical background.

Thus our phase space consists of T^*Q with an open covering $\{U_\alpha\}$ and a symplectic form ω_{can}^{α} on each U_{α} such that, on $U_{\alpha} \cap U_{\beta} \neq \emptyset$,

(7)
$$\omega_{can}^{\alpha} = \mu_{\beta\alpha} \omega_{can}^{\beta}.$$

Equation (7) implies that the collection $\{\mu_{\alpha\beta}\}\$ satisfies the *cocycle condition* $\mu_{\beta\gamma} = \mu_{\beta\alpha}\mu_{\alpha\gamma}$, hence one obtains a real line bundle $L \to T^*Q$ with transition functions $\{\mu_{\alpha\beta}\}\$. The *global* Hamiltonian is not anymore a smooth function on T^*Q but rather a smooth section of L. The cocycle condition can be rephrased by saying that

$$\mu_{\beta\alpha} = \frac{e^{\sigma_{\alpha}}}{e^{\sigma_{\beta}}}$$

for functions $\sigma_{\alpha} : U_{\alpha} \to \mathbb{R}$ (resp. $\sigma_{\beta} : U_{\beta} \to \mathbb{R}$). Now equation (7) shows that the collection of local 2-forms $\{e^{\sigma_{\alpha}}\omega_{can}^{\alpha}\}$ piece together to a global, non-degenerate 2-form ω on T^*Q . Clearly, the 1-forms $\{d\sigma_{\alpha}\}$ piece together to a 1-form ϑ and $d\omega = \vartheta \land \omega$. Thus ϑ is the Lee form of the locally conformally symplectic structure (ω, ϑ) and $(T^*Q, \omega, \vartheta)$ is a locally conformally symplectic manifold.

As pointed out in [62], given any manifold Q with a closed 1-form $\bar{\vartheta}$, the cotangent bundle T^*Q admits a canonical exact locally conformally symplectic structure

$$(\boldsymbol{\omega},\boldsymbol{\vartheta}) = (d_{\boldsymbol{\vartheta}}(-\boldsymbol{\lambda}_{can}),\boldsymbol{\vartheta}),$$

where $\pi: T^*Q \to Q$ is the canonical projection and $\vartheta = \pi^* \overline{\vartheta}$.

I conclude this jaunt into classical mechanics by mentioning a couple of more papers where ideas of conformally symplectic geometry find applications to physical problems.

Let (M, ω) be a symplectic manifold, let $H \in C^{\infty}(M)$ be a Hamiltonian function and X_H be the corresponding Hamiltonian vector field. If $f \in C^{\infty}(M)$ is a function, then the vector field $e^f X_H$ is *conformally Hamiltonian* with Hamiltonian H and conformal factor e^f . It clearly satisfies $e^f dH = \omega(e^f X_H, \cdot)$. Moreover, $e^f X_H$ is the Hamiltonian vector field of H for the 2-form $\omega' = e^{-f} \omega$, which is not closed anymore, but conformally closed with Lee form -df.

In [82] Maciejewski, Przybylska and Tsiganov consider conformally Hamiltonian vector fields in the theory of bi-Hamiltonian systems, in order to produce examples of completely integrable systems.

In [86] Marle used conformally Hamiltonian vector fields to study, in a new perspective, a certain diffeomorphism between the phase space of the Kepler problem and an open subset of the cotangent bundle of S^3 (resp. of a 2-sheeted hyperboloid, according to the energy of the motion).

In [138], Wojtkowski and Liverani apply the formalism of conformally symplectic geometry in order to model concrete physical situations such as the Gaussian isokinetic dynamics, also with collisions, and the Nosé-Hoover dynamics. More precisely, the authors show that such systems fall under the formalism of conformal Hamiltonian dynamics and explain how to easily deduce results about the symmetric of the Lyapunov spectrum.

LOCALLY CONFORMALLY SYMPLECTIC AND KÄHLER GEOMETRY

3. KÄHLER AND LOCALLY CONFORMALLY KÄHLER GEOMETRY

A Kähler manifold is a complex manifold with a compatible Riemannian metric such that the induced complex structure is parallel with respect to the Levi-Civita connection. The Riemannian metric and the complex structure provide a non-degenerate 2-form which is also parallel, in particular closed. Thus Kähler geometry lies at the intersection between *complex*, *Riemannian* and *symplectic* geometry. The combination of three geometries produces a class of manifolds which possess distinctive properties within each of the three geometries.

As complex manifolds, Kähler manifolds can, to a certain extent, be studied with methods of complex algebraic geometry; indeed, the main source of examples of compact Kähler manifolds is provided by projective varieties, i. e. zero loci of homogeneous polynomials in $\mathbb{C}P^N$. I should point out, however, that the "generic" Kähler manifold is not projective. Informally, this question is the content of the *Kodaira problem*:

Can every compact Kähler manifold be deformed to a projective manifold?

The answer to this question is, perhaps surprisingly, no, as proved by Claire Voisin in [131, 132]; see also the survey [64] by Daniel Huybrechts. A certain class of compact Kähler manifolds, namely *Hodge manifolds*, can be holomorphically embedded into a complex projective space: this is the content of Kodaira's embedding theorem, see [130, Theorem 7.11]. In this case the Kähler class is the pullback of the Fubini-Study class but the embedding is, in general, not isometric.

From the perspective of Riemannian geometry, the reduced holonomy of a compact Kähler manifold is contained in the unitary group U(n), where *n* is half the dimension of the manifold. Manifolds with special holonomy turn out to have many applications in Physics, see for instance [66].

From the point of view of symplectic geometry, compact Kähler manifolds satisfy the *Hard Lefschetz property*, see [63], while symplectic manifolds need not, see [19]. The Lefschetz property implies the well-known fact that the Betti numbers of odd degree are even on a compact Kähler manifold (this follows also directly from Hodge theory). In a very actual research area such as homological mirror symmetry, the fact that a symplectic structure is part of a Kähler structure on a compact manifold sheds a great deal of light in the study of such duality – see by way of example [115]. I should also mention here that it was originally believed, and to some extent even erroneously proved, see [60], that every compact symplectic manifold admitted a Kähler metric. It was only in 1976 that Thurston provided the first example of a compact symplectic manifold with first Betti number equal to 3, hence no Kähler metric, see [118]. Since then, the quest for compact symplectic manifolds with no Kähler metrics has inspired beautiful Mathematics – see for instance the papers [41, 49, 81, 88] and the book [120].

Finally, concerning the topology of compact Kähler manifolds, I should point out that they are formal in the sense of Sullivan, see [35].

For many purposes¹¹ it can be convenient to relax the strong integrability properties characterizing the three geometries that come together in a Kähler structure. The right framework to do this is that of *almost Hermitian structures*. An almost Hermitian structure on a manifold

¹¹I will come back to this point in Section 4.

consists of a triple (g, J, ω) , where g is a Riemannian metric, J is an almost complex structure and ω is a 2-form, called the Kähler form, such that J is an isometry for g. Actually two of the three structures determine the third one through the equation

$$\omega(X,Y) = g(X,JY) \, .$$

In their celebrated 1980 paper *The sixteen classes of almost Hermitian manifolds and their linear invariants* [55], Alfred Gray and Luis Hervella classified almost Hermitian structures in terms of the covariante derivative, with respect to the Levi-Civita connection, of the Kähler form. Kähler structures are recovered as those almost Hermitian structures whose Kähler form is parallel with respect to the Levi-Civita connection. This opens the doors to a whole series of almost Hermitian structures in which some of the integrability properties are not satisfied. Starting with this paper, the study of these structures was undertaken in a systematic way. However, some of them had already appeared before in the literature. For instance, nearly Kähler structures were considered by Fukami and Ishihara in 1955 ([42], on the six sphere) and then studied extensively by Gray [52, 53, 54]. Locally conformally almost Kähler structures were discussed in Vaisman's 1976 paper [123]. To an almost Hermitian structure (g, J, ω) on M^{2n} with $n \ge 2$ one can associate the *Lee form* $\vartheta \in \Omega^1(M)$, defined as

$$\vartheta = -\frac{1}{n-1}J(d^*\omega).$$

An almost Hermitian structure (g, J, ω) is *almost Kähler* if $d\omega = 0$, *locally conformally almost Kähler* if $d\omega = \vartheta \wedge \omega$ and $d\vartheta = 0$. If *J* is integrable, than the almost Kähler structure is Kähler and the locally conformally almost Kähler is *locally conformally Kähler*. For n = 2, the Lee form of a Hermitian structure is uniquely determined by the condition $d\omega = \vartheta \wedge \omega$. If $\vartheta = 0$ (resp. $d\vartheta = 0$) then the Hermitian structure is Kähler (resp. locally conformally Kähkler). If $n \ge 3$ the closedness of the Lee form follows from the equation $d\omega = \vartheta \wedge \omega^{12}$.

Thus, similarly to what happened for Kähler manifolds, locally conformally Kähler manifolds can be considered simultaneously as complex, Riemannian and locally conformally symplectic manifolds. As I mention above, a Kähler manifold M^{2n} can be characterized as a Riemannian manifold whose holonomy lies in U(n). One could think of a conformal version of manifolds with special holonomy. For instance, locally conformally hyperkähler manifolds are studied in [36, Chapter 11]. References for locally conformally G₂ and Spin(7) structures are [40, 65].

As it happens in the locally conformally symplectic case, a locally conformally Kähler manifold is actually Kähler, in case $\vartheta = 0$, or *globally* conformal to a Kähler manifold if ϑ is exact. In general, one can only argue that this conformal property holds locally. I prefer to consider locally conformally Kähler manifolds as a class which is distinct from that of Kähler manifolds (soemtimes one uses the terminology *strictly* locally conformally Kähler). The reference for locally conformally Kähler geometry is the monograph [36] by Dragomir and Ornea; see also [97, 103].

Of particular importance within locally conformally Kähler manifolds are *Vaisman manifolds*¹³; these are characterized by the property that the Lee form is parallel with respect to the

¹²This apparent discrepancy between the cases n = 2 and $n \ge 3$ is due to the fact that, in complex dimension 2, there exist only two "pure" classes in the Gray-Hervella classification.

¹³Vaisman manifolds were first called *generalized Hopf manifolds* by Vaisman, see [127].

Levi-Civita connection. I will implicitly assume that $\|\vartheta\| \neq 0$ on a compact Vaisman manifold, hence ϑ is nowhere zero. An interesting example of a Vaisman *surface* is the Hopf surface. This is defined as a compact complex surface whose universal cover is $\mathbb{C}^2 \setminus \{0\}$. As shown in [45], each primary¹⁴ Hopf surface admits a locally conformally Kähler metric and some Hopf surfaces (those of class 1) admit Vaisman metrics (see also [110]). Since every primary Hopf surface is diffeomorphic to $S^3 \times S^1$, no Hopf surface admits Kähler metrics.

As complex manifolds, locally conformally Kähler manifolds are different from Kähler manifolds¹⁵. For instance, a small deformation of the complex structure of a Kähler manifold remains Kähler (see [130, Theorem 9.23]). On the other hand, it was shown by Belgun in [24] that this is not the case for complex structures neither on locally conformally Kähler nor on Vaisman manifolds. In the same paper, Belgun carried out a systematical analysis of locally conformally Kähler metrics on compact complex surfaces. His paper was in some sense groundbreaking since it was conjectured that all non-Kähler compact complex surfaces admitted locally conformally Kähler metrics. Locally conformally Kähler, albeit non Vaisman, metrics on some Inoue surfaces had been previously constructed by Tricerri in [121]. Most non-Kähler compact complex surfaces admit locally conformally Kähler metrics (see [24, 27, 45]); in fact, only one of the three Inoue surfaces is known not to admit any. The *spherical shell conjecture* predicts that every class VII₀ surface with $b_2 > 0$ is a Kato surface, i. e. it contains a spherical shell¹⁶, see [105]. If the spherical shell conjecture holds, the remaining non Kähler compact complex surfaces admit locally conformally Kähler metrics.

A locally conformally Kähler manifold can be equivalently defined as a manifold admitting a Kähler covering whose deck group acts by conformal transformations (see [127]). As proved by Verbitsky in [129], the Kähler metric on the universal covering of a Vaisman manifold admits a global Kähler potential. Since this property is stable under small deformations, a Vaisman structure deforms to a locally conformally Kähler one, not necessarily a Vaisman one. Motivated by this observations, Ornea and Verbitsky defined a class of locally conformally Kähler manifolds, which strictly contains Vaisman manifolds, namely locally conformally Kähler manifolds with (proper) potential, see [101, 104]. Nice results for such manifolds are available. For instance, it was proved in [101] that they admit an embedding into a Hopf manifold, provided the complex dimension is at least 3 (see also [100]). Hopf manifolds are generalizations to arbitrary complex dimensions of Hopf surfaces: they are defined as quotients of $\mathbb{C}^n \setminus \{0\}$ by a discrete subgroup of linear holomorphisms. A *primary* Hopf manifold is the quotient of $\mathbb{C}^n \setminus \{0\}$ by the action of the abelian group generated by complex numbers $\lambda_1, \ldots, \lambda_n$, with $0 < |\lambda_0| \le \ldots \le |\lambda_n| < 1$, where the action sends x_i to $\lambda_i x_i$, for $i = 1, \ldots, n$ (see [69]). Compact Vaisman manifolds can be embedded into primary Hopf manifolds. In this sense, Vaisman manifolds and, more generally, locally conformally Kähler manifolds with proper potential are the analogue of Hodge manifolds in Kähler geometry. In locally conformally Kähler geometry, the statement corresponding the the Kodaira problem in Kähler geometry would be the following:

¹⁴A Hopf surface is called *primary* if its fundamental group is isomorphic to \mathbb{Z} . Every Hopf surface is finitely covered by a primary one.

¹⁵In the short note [11], Aubin erroneously claimed that a compact locally conformally Kähler manifold is actually Kähler.

¹⁶A spherical shell S in a compact complex surface M is a real submanifold diffeomorphic to S^3 , such that $M \setminus S$ is connected and S has a neighbourhood which is biholomorphic to an annulus in \mathbb{C}^2 .

Can every compact locally conformally Kähler manifold be deformed to a Vaisman manifold?

In fact, every compact locally conformally Kähler manifold with potential can be deformed to a Vaisman one, as shown in [102, Theorem 2.1]. A compact locally conformally Kähler manifold is globally conformally Kähler if and only if it admits a Kähler metric. Related to the above question, I mention the following two conjectures (see [124, 125]):

- A compact locally conformally Kähler manifold satisfying the topological conditions of a Kähler manifold admits some global Kähler metric.
- A compact locally but not globally conformally Kähler manifold has an odd odd-degree Betti number.

It is easy to see that the first Betti number of a compact Vaisman manifold is odd, hence the second conjecture holds for locally conformally Kähler manifolds with potential. A compact complex surface which admits a locally conformally Kähler but no Kähler metrics has odd first Betti number. In [95] Oeljeklaus and Toma disproved the second conjecture by constructing a compact complex 3-fold admitting locally conformally Kähler metrics with all odd-degree Betti numbers even. This also settles in the negative the Kodaira problem in the locally conformally Kähler context. The so-called *Oeljeklaus-Toma manifolds* are generalizations to arbitrary complex dimensions of Inoue surfaces. They can also be described as solvmanifolds, see [70].

The holonomy of locally conformally Kähler manifolds has been investigated in [83]. Although the complex structure is not parallel with respect to the Levi-Civita connection, it can be useful to have an auxiliary metric connection which does fulfill this property. To any Hermitian structure (g, J) on a manifold M one can associate a unique connection, called *Chern connection* ∇^C , which satisfies $\nabla^C g = 0 = \nabla^C J$ and whose torsion T is of type (2,0), that is,

$$T(JX,Y) = JT(X,Y) \quad \forall X,Y \in \mathfrak{X}(M).$$

The Chern connection coincides with the Levi-Civita connection if the Hermitian structure is Kähler. In [43] Gauduchon associated a 1-form $\tilde{\vartheta}$ to the Chern connection, the *torsion 1-form*, as follows:

$$\vartheta(X) = \operatorname{trace}(Y \mapsto T(X,Y)).$$

One can show that $\tilde{\vartheta} = (n-1)\vartheta$, hence the Lee form and the torsion 1-form are strictly related. Thus the Lee form of a locally conformally Kähler structure measures, in a certain sense, its lack of integrability, where integrability is the Kähler case.

A Weyl structure on a conformal manifold (M, c) is a torsion-free linear connection ∇^W , the Weyl connection, which preserves the conformal class c. This means that there exists a 1-form ϑ such that $\nabla^W g = g \otimes \vartheta$ for every $g \in c$. A conformal Hermitian manifold is a conformal manifold (M, c) with a complex structure J which is Hermitian for some, hence all, $g \in c$. If $\nabla^W J = 0$, then (M, c, J) is a Kähler-Weyl manifold. As pointed out by Kokarev in [71], locally conformally Kähler manifolds are examples of Kähler-Weyl manifolds; the Weyl connection is related to the Levi-Civita connection ∇ by the formula

$$\nabla_X^W Y = \nabla_X Y - \frac{1}{2}\vartheta(X)Y - \frac{1}{2}\vartheta(Y)X + \frac{1}{2}g(X,Y)U,$$

where $U = \vartheta^{\sharp}$ is the *Lee field*. This point of view on locally conformally Kähler geometry was adopted in [71, 72], with applications to the topology of compact Vaisman¹⁷ manifolds, in particular their fundamental group.

Since the Lee form of a Vaisman structure is parallel, the results of [34] imply that the underlying locally conformally symplectic structure is exact. But more is true: if (g,J) is Vaisman then, up to a homothety, one can assume that $\|\vartheta\| = 1$ and one can show that the underlying locally conformally symplectic structure is of the first kind, see [12, 36]; more precisely, one has $\mathscr{L}_U \omega = 0$ and $\omega = d\eta - \eta \wedge \vartheta$ for $\eta = -\iota_U \omega$ (see also [97, Section 9]).

In Section 1 we discussed the relation between locally conformally symplectic structures of the first kind and contact structures. A similar relation exists between Vaisman and Sasakian structures. A Sasakian structure is a normal contact metric structure, see [25, 26]. Indeed, the mapping torus of a Sasakian manifold and a Sasakian automorphism, that is, a diffeomorphism which respects the whole Sasakian structure, carries a natural Vaisman structure. In [99] the authors claimed that, in the compact case, the converse also holds; as explained in [104], however, the proof is flawed. Nevertheless, the result holds up to diffeomorphism: a compact Vaisman manifold is diffeomorphic to the mapping torus of a Sasakian manifold and a Sasakian automorphism. Morally, this discrepancy between the two directions in similar to what happens in the non-metric case. Based on this approach, a global splitting result for compact Vaisman manifolds was obtained in [22]. As in the non-metric case, let me notice the absence of structure results for compact locally conformally Kähler manifolds which are not Vaisman.

Analogous to the symplectic versus Kähler case, Ornea and Verbitsky formulated in [103] the following problem:

Construct a compact locally conformally symplectic manifolds which admits no locally conformally Kähler metrics.

A first answer to this question was provided by Bande and Kotschick in [13]. Different answers are contained in [20, 21].

Related to this problem is a conjecture of Ugarte which aims to give a complete characterization of locally conformally Kähler structures on nilmanifolds. In [122, Page 200], he conjectured the following:

A compact nilmanifold of dimension $2n \ge 4$ admitting a locally conformally Kähler structure is the product of *N* with S^1 , where *N* is a quotient of H(1,n).

¹⁷Kokarev defined in [71] *pluricanonical* locally conformally Kähler metrics (actually Kähler-Weyl structures) as those for which $(\nabla \vartheta)^{1,1} = 0$. In [102] it was erroneously claimed that a locally conformally Kähler metric is pluricanonical if and only if it admits a potential. The mistake was clarified in [92, 104], where it was proved that a compact pluricanonical locally conformally Kähler manifold is in fact Vaisman.

Here H(1,n) is the generalized Heisenberg group,

$$H(1,n) = \left\{ \begin{pmatrix} 1 & y_1 & y_2 & \dots & y_n & z \\ 0 & 1 & 0 & \dots & 0 & x_1 \\ \vdots & 0 & \ddots & \ddots & \vdots & x_2 \\ \vdots & \vdots & \ddots & \ddots & 0 & \vdots \\ \vdots & \vdots & \ddots & 1 & x_n \\ 0 & 0 & \dots & \dots & 0 & 1 \end{pmatrix} \mid x_i, y_i, z \in \mathbb{R}, \ i = 1, \dots, n \right\}.$$

The conjecture holds in full generality in dimension 4 ([20]). In higher dimension it holds if one assumes that the complex structure of the locally conformally Kähler structure is left-invariant¹⁸ (see [114]) or if the locally conformally Kähler structure is Vaisman (see [18]).

We mention here the fact that compact Vaisman manifolds satisfy a Hard Lefschetz property (see [28]); this result builds on the Hard Lefschetz property for compact Sasakian manifolds proved in [29]. Again, the lack of structure theorems for general locally conformally Kähler manifolds reflects on the absence of a Hard Lefschetz property in the most general setting.

Compact Vaisman manifolds are, in general, non formal. In 2001, Kotschick introduced the notion of *geometric formality*: a closed manifold is geometrically formal if it admits a Riemannian metric such that the product of two harmonic forms is harmonic (see [73]). Geometric formality implies formality in the sense of Sullivan, but the converse is not true, see for instance [74]. In [98], Ornea and Pilca showed that geometrically formal compact Vaisman manifolds obey to strong topological restrictions. It is not yet clear the extent to which a compact Vaisman manifold is non formal.

I end this section by quoting some other results about locally conformally Kähler manifolds.

Homogeneous locally conformally Kähler structures are in fact Vaisman, see [2, 44]. The papers [47, 48] consider the problem of reduction in locally conformally Kähler geometry. In [48] the authors introduce the notions of *presentation* and *rank* of a locally conformally Kähler manifold. The rank of a locally conformally Kähler structure and his relation with other properties such as the existence of a potential have been further investigated in [111]. Toric locally conformally Kähler manifolds, and in particular Vaisman, are considered in [84]. The blow-up of a locally conformally Kähler manifold was studied in [106, 121, 133]. An interesting contact point between locally conformally symplectic and Kähler geometry appears in the papers [7, 8]. The authors consider locally conformally symplectic structures (ω, ϑ) on compact complex surfaces (M,J) such that ω tames J, i. e. the (1,1)-part of ω is positive definite. The Morse-Novikov cohomology of locally conformally Kähler surfaces has been investigated in [108]. Results on the deformations of Lee classes of locally conformally Kähler structures, metrics which are locally conformal to notable ones, for instance to balanced ones, have been studied in [6].

¹⁸This means that it comes from a left-invariant complex structure on the corresponding Lie group.

LOCALLY CONFORMALLY SYMPLECTIC AND KÄHLER GEOMETRY

4. STRINGS, SUPERSYMMETRY AND M-THEORY

Passer de la mécanique de Newton à celle d'Einstein doit être un peu, pour le mathématicien, comme de passer du bon vieux dialecte provençal à l'argot parisien dernier cri. Par contre, passer à la mécanique quantique, j'imagine, c'est passer du français au chinois.

Alexandre Grothendieck, "Récoltes et Semailles", 1986.

Given the fact that I am not particularly well-versed in chinese, I will keep this section as low-key as possible. My only goal here is to show that, albeit at a different level from what I discussed in Section 2, Physics can motivate and foster research also in the case of locally conformally Kähler structures.

At the end of the thirties of the twentieth century, the physical community was able to catch a breath after the establishment of two very important theories, namely General Relativity by Einstein and Quantum Mechanics by Bohr, Heisenberg and Schrödinger among others¹⁹. By the end of the seventies the four foundamental forces of Physics, Gravitation, Electromagnetism, Weak and Strong interactions, had been completely described: while General Relativity took care of Gravitation, Quantum Mechanics was able to explain the other three. However, since General Relativity is formulated in the framework of classical physics, in order to fill the gap and elaborate a theory which subsumes the four forces, it is necessary to develop a quantized version of Gravity; this task includes, in particular, the search of a particle, called *graviton*, which carries Gravity. In the Standard Model atoms are broken down to particles, called *fermions* which are the very constituents of matter while the three interactions (electromagnetic, weak and strong) are described as trasmitted by another kind of particles, called *bosons*. These ultimate particles are thought of as punctiform. Among other things, fermions have half-integer spin, while bosons have integer spin. Particle physics in the formalisms of the Standard Model, however, presents some problems.

String theory emerged during the sixties and the seventies, with the goal to explain these incongruencies; roughly speaking, in string theory punctiform particles are interpreted as 1-dimensional manifolds with or without boundary. The vibration mode of the string determines the type of particle. In particular, one of the possible states of a string corresponds to the graviton; thus string theory has Quantum Gravity built in. One should keep in mind that, at a large scale, strings look punctiform; the standard analogy with everyday's life is that of a hanging cable or a garden hose: at a certain distance those look one-dimensional, but an ant moving on them would perceive the second dimension.

At a very naïf level, *Supersymmetry* is a theory in which to each fermion corresponds a boson under a supersymmetry operator. One of the peculiar features of supersymmetry is that it requires a universe with extra dimensions apart from the standard 4-dimensional space time²⁰. The number of such extra dimensions is constrained by supersymmetry and can be at most 11 and the universe is thought of as a product $M^n \times K^{11-n}$, where *M* is Minkowski space-time and

¹⁹Although Einstein is acknowledged as the founding father of General Relativity, his 1905 article "Über einen die Erzeugung und Verwandlung des Lichtes betreffenden heuristischen Gesichtspunkt", which won him the Nobel Prize, laid the foundations of the theory of quanta.

²⁰The idea of requiring extra dimensions in order to unify Gravity with Electromagnetism goes back to Theodor Kaluza and Oskar Klein in the twenties of last century.

K is the so-called *internal space* on which supersymmetry operators act. The extra dimensions encoded in *K*, however, escape our perception. In Physics one uses the term *compactification* to indicate that some extra dimensions "wrap up" around a lower dimensional, perceptible universe. This encompasses the idea of string theory that extra dimensions should be "small".

Supersymmetry is a further step toward the search for unification and nowadays string theory does include supersymmetry, so that one speaks of *supersymmetric string theory*. Physicists came up with five different supersymmetric string theories: type I, SO(32)- and $E_8 \times E_8$ heterotic, type IIA and type IIB. Over the years, however, they were able to prove that such theories were related to one another in a highly non-trivial way, through some *dualities*. This led Edward Witten, in 1995, to formulate *M*-theory, a theory which unifies all known supersymmetric string theories; in this sense, all of them are different incarnations of the same theory.

From the point of view of a mathematician, the bridge between M-theory and Geometry is provided by compactification²¹. Indeed, not every way of compactifying the extra dimensions predicted by M-theory is compatible with the properties of the observed space-time. Indeed, supersymmetry equations for the internal space constrain its geometry. In most models, the internal space is assumed to be a compact Calabi-Yau²² manifold. This requirement can be relaxed to include structures with torsion, see for instance [117].

An explicit compactification of M-theory in 8 dimensions (that is, a solution for which space-time is 3-dimensional and the internal manifold 8-dimensional) was constructed in [23]. The authors show that such 8-dimensional manifold is endowed with a Riemannian metric which, up to a global conformal factor called the warp factor, is Calabi-Yau.

In [116], Shahbazi makes an interesting remark. He asserts that the way in which physicists obtain solutions to their field equations, be they supersymmetry or general relativity, is by performing explicit computations on a local patch of the manifold they are looking for. A (local) solution consists then of an open set with distinguished tensors; the subsequent problem consists in determining which compact manifolds exhibit the particular set as an open set (this is the problem of the maximally analytic extension of a given local patch with a locally defined metric). In order to glue together different open sets on which a solution is known one can require that the change of coordinates respect the distinguished tensors. But more general transformations could be allowed, as we saw in Section 2, especially if our goal is to preserve the equations of motions (we refer the reader to [75] for an explanation of this principle in the setting of supergravity and supersymmetry). In the case we consider here, Shahbazi makes the Ansatz that the warp factor considered in [23] does not necessarily need to be of global nature. If one is able to construct a solution of the equations of motions under this Ansatz, then it would be impossible to distinguish, at a local scale, the two solutions. Shahbazi constructs an explicit compactification of M-theory on a compact 8-dimensional Riemannian manifold whose metric is conformal to a Kähler one (actually Calabi-Yau) only locally. More precisely, he constructs an internal 8-dimensional locally conformally Kähler manifold locally equipped with a preferred Calabi-Yau structure. It is a special type of Hopf manifold, diffeomorphic to $S^1 \times S^7$. It is remarkable that the global topology of this explicit compactification does not carry any Kähler (hence Calabi-Yau) metric. Shahbazi's solution can be seen as a principal

²¹See [96] for a nice explanation of how a string theorist sees geometry.

²²Recall that a Riemannian manifold (M^n, g) is Calabi-Yau if its holonomy is contained in $SU(\frac{n}{2})$.

torus bundle over a projective manifold; moreover, the solution is endowed with a codimension 1 foliation whose leaves carry nearly parallel G_2 -structures. Another nice outcome of Shahbazi's approach is that his solution evades the Maldacena-Nuñez *no-go theorem*, see [85]. This asserts, very roughly speaking, that every solution to the equations of supersymmetry which is compatible with a certain zeroth-order approximation of the theory (in a parameter, the Planck length, which corresponds to the tension of the string) must have, in particular, constant warp factor. In order to obtain non-trivial solutions, therefore, one has to allow at least a first-order approximation. In [23] a particular correction of order 6 in the Planck length was included. Shahbazi's solution evades the no-go theorem without needing any kind of correction. The trick lies in the topology of the solution, which is completely different from that of a Kähler manifold.

Again we see how a certain relaxation of the Kähler (or Calabi-Yau) condition leads to new geometries that can be of use in Physics. It is undeniable that these inputs from Physics are of paramount importance in motivating future research in the area of locally conformally Kähler geometry.

ACKNOWLEDGEMENTS

This note is part of my *Habilitationssarbeit* at the Philipps-Universität Marburg, and most of it was written while I was working there. I am grateful to the whole *Arbeitsgruppe* for the continuous support. The final editing was done in Madrid, where I am supported by a Juan de la Cierva Fellowship at Universidad Complutense de Madrid. I am grateful to Daniele Angella, Marcos Origlia, Alexandra Otiman, Maurizio Parton and Sönke Rollenske for their comments.

REFERENCES

- Ralph Abraham and Jerrold E. Marsden. *Foundations of mechanics*. Benjamin/Cummings Publishing Co., Inc., Advanced Book Program, Reading, Mass., 1978. Second edition, revised and enlarged, With the assistance of Tudor Raţiu and Richard Cushman. (cited on p. 7)
- [2] D. V. Alekseevsky, V. Cortés, K. Hasegawa, and Y. Kamishima. Homogeneous locally conformally Kähler and Sasaki manifolds. *Internat. J. Math.*, 26(6):1541001, 29, 2015. (cited on p. 14)
- [3] Adrián Andrada and Marcos Origlia. Lattices in almost abelian Lie groups with locally conformal Kähler or symplectic structures. *manuscripta math.*, 2017. (cited on p. 4)
- [4] Daniele Angella, Giovanni Bazzoni, and Maurizio Parton. Structure of locally conformally symplectic lie algebras and solvmanifolds. 2017. https://arxiv.org/abs/1704.01197. (cited on p. 3, 5)
- [5] Daniele Angella, Alexandra Otiman, and Nicoletta Tardini. Cohomologies of locally conformally symplectic manifolds and solvmanifolds. *Ann. Global Anal. Geom.*, 2017. (cited on p. 4, 5)
- [6] Daniele Angella and Luis Ugarte. Locally conformal Hermitian metrics on complex non-Kähler manifolds. *Mediterr. J. Math.*, 13(4):2105–2145, 2016. (cited on p. 14)
- [7] Vestislav Apostolov and Georges Dloussky. Locally conformally symplectic structures on compact non-Kähler complex surfaces. *Int. Math. Res. Not. IMRN*, (9):2717–2747, 2016. (cited on p. 14)
- [8] Vestislav Apostolov and Georges Dloussky. On the Lee classes of locally conformally symplectic complex surfaces. 2016. https://arxiv.org/abs/1611.00074. (cited on p. 14)
- [9] Vladimir I. Arnol'd. Mathematical methods of classical mechanics, volume 60 of Graduate Texts in Mathematics. Springer-Verlag, New York, second edition, 1989. Translated from the Russian by K. Vogtmann and A. Weinstein. (cited on p. 1, 7)
- [10] Vladimir I. Arnol'd. Symplectic geometry and topology. J. Math. Phys., 41(6):3307–3343, 2000. (cited on p. 2)

- [11] Thierry Aubin. Variétés hermitiennes compactes localement conformément kählériennes. C. R. Acad. Sci. Paris, 261:2427–2430, 1965. (cited on p. 11)
- [12] Gianluca Bande and Dieter Kotschick. Moser stability for locally conformally symplectic structures. *Proc. Amer. Math. Soc.*, 137(7):2419–2424, 2009. (cited on p. 4, 5, 13)
- [13] Gianluca Bande and Dieter Kotschick. Contact pairs and locally conformally symplectic structures. In *Harmonic maps and differential geometry*, volume 542 of *Contemp. Math.*, pages 85–98. Amer. Math. Soc., Providence, RI, 2011. (cited on p. 13)
- [14] Augustin Banyaga. A geometric integration of the extended Lee homomorphism. J. Geom. Phys., 39(1):30–44, 2001. (cited on p. 4, 6)
- [15] Augustin Banyaga. Quelques invariants des structures localement conformément symplectiques. C. R. Acad. Sci. Paris Sér. I Math., 332(1):29–32, 2001. (cited on p. 6)
- [16] Augustin Banyaga. On the geometry of locally conformal symplectic manifolds. In *Infinite dimensional Lie groups in geometry and representation theory (Washington, DC, 2000)*, pages 79–91. World Sci. Publ., River Edge, NJ, 2002. (cited on p. 5)
- [17] Augustin Banyaga. Examples of non d_{ω} -exact locally conformal symplectic forms. J. Geom., 87(1-2):1–13, 2007. (cited on p. 4, 5)
- [18] Giovanni Bazzoni. Vaisman Nilmanifolds. Bull. Lond. Math. Soc., 49(5):824-830, 2017. (cited on p. 14)
- [19] Giovanni Bazzoni, Marisa Fernández, and Vicente Muñoz. Formality and the Lefschetz property in symplectic and cosymplectic geometry. *Complex Manifolds*, 2:53–77, 2015. (cited on p. 9)
- [20] Giovanni Bazzoni and Juan C. Marrero. Locally conformal symplectic nilmanifolds with no locally conformal Kähler metrics. *Complex Manifolds*, 2017. (cited on p. 3, 13, 14)
- [21] Giovanni Bazzoni and Juan C. Marrero. On locally conformal symplectic manifolds of the first kind. *Bull. Sci. Math.*, 2017. (cited on p. 5, 13)
- [22] Giovanni Bazzoni, Juan C. Marrero, and John Oprea. A splitting theorem for compact Vaisman manifolds. *Rend. Semin. Mat. Univ. Politec. Torino*, 74(1):21–29, 2016. (cited on p. 13)
- [23] Katrin Becker and Melanie Becker. *M*-theory on eight-manifolds. *Nuclear Phys. B*, 477(1):155–167, 1996. (cited on p. 16, 17)
- [24] Florin A. Belgun. On the metric structure of non-Kähler complex surfaces. *Math. Ann.*, 317(1):1–40, 2000. (cited on p. 11)
- [25] David E. Blair. Riemannian geometry of contact and symplectic manifolds, volume 203 of Progress in Mathematics. Birkhäuser Boston, Inc., Boston, MA, second edition, 2010. (cited on p. 13)
- [26] Charles P. Boyer and Krzysztof Galicki. Sasakian geometry. Oxford Mathematical Monographs. Oxford University Press, Oxford, 2008. (cited on p. 13)
- [27] Marco Brunella. Locally conformally K\u00e4hler metrics on Kato surfaces. Nagoya Math. J., 202:77–81, 2011. (cited on p. 11)
- [28] Beniamino Cappelletti-Montano, Antonio De Nicola, Juan C. Marrero, and Ivan Yudin. Hard Lefschetz theorem for Vaisman manifolds. 2015. https://arxiv.org/abs/1510.04946. (cited on p. 14)
- [29] Beniamino Cappelletti-Montano, Antonio De Nicola, and Ivan Yudin. Hard Lefschetz theorem for Sasakian manifolds. *J. Differential Geom.*, 101(1):47–66, 2015. (cited on p. 14)
- [30] Baptiste Chantrain and Emmy Murphy. Conformal symplectic geometry of cotangent bundles. 2016. https://arxiv.org/abs/1606.00861. (cited on p. 2, 6)
- [31] Youming Chen and Song Yang. On the blow-up of points on locally conformally symplectic manifolds. *C. R. Math. Acad. Sci. Paris*, 354(4):411–414, 2016. (cited on p. 5)
- [32] Vicente Cortés and Alexander S. Haupt. *Mathematical Methods of Classical Physics*. SpringerBriefs in Physics. Springer International Publishing, first edition, 2017. (cited on p. 7)
- [33] Luis C. de Andrés, Luis A. Cordero, Marisa Fernández, and José J. Mencía. Examples of four-dimensional compact locally conformal Kähler solvmanifolds. *Geom. Dedicata*, 29(2):227–232, 1989. (cited on p. 5)
- [34] Manuel de León, Belén López, Juan C. Marrero, and Edith Padrón. On the computation of the Lichnerowicz-Jacobi cohomology. *J. Geom. Phys.*, 44(4):507–522, 2003. (cited on p. 4, 13)
- [35] Pierre Deligne, Phillip Griffiths, John Morgan, and Dennis Sullivan. Real homotopy theory of Kähler manifolds. *Invent. Math.*, 29(3):245–274, 1975. (cited on p. 9)
- [36] Sorin Dragomir and Liviu Ornea. Locally conformal Kähler geometry, volume 155 of Progress in Mathematics. Birkhäuser Boston, Inc., Boston, MA, 1998. (cited on p. 1, 10, 13)

- [37] Yakov Eliashberg and Emmy Murphy. Making cobordisms symplectic. 2015. https://arxiv.org/abs/1504.06312. (cited on p. 2, 3, 5)
- [38] Michael Farber. *Topology of closed one-forms*, volume 108 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 2004. (cited on p. 4)
- [39] Rui Loja Fernandes and Pedro Frejlich. An *h*-principle for symplectic foliations. *Int. Math. Res. Not. IMRN*, (7):1505–1518, 2012. (cited on p. 3)
- [40] Marisa Fernández, Anna Fino, and Alberto Raffero. Locally conformal calibrated G₂-manifolds. Ann. Mat. Pura Appl. (4), 195(5):1721–1736, 2016. (cited on p. 10)
- [41] Marisa Fernández and Vicente Muñoz. An 8-dimensional nonformal, simply connected, symplectic manifold. Ann. of Math. (2), 167(3):1045–1054, 2008. (cited on p. 9)
- [42] Tetsuzo Fukami and Shigeru Ishihara. Almost Hermitian structure on S⁶. Tôhoku Math. J. (2), 7:151–156, 1955. (cited on p. 10)
- [43] Paul Gauduchon. La 1-forme de torsion d'une variété hermitienne compacte. *Math. Ann.*, 267(4):495–518, 1984. (cited on p. 12)
- [44] Paul Gauduchon, Andrei Moroianu, and Liviu Ornea. Compact homogeneous lcK manifolds are Vaisman. Math. Ann., 361(3-4):1043–1048, 2015. (cited on p. 14)
- [45] Paul Gauduchon and Liviu Ornea. Locally conformally Kähler metrics on Hopf surfaces. *Ann. Inst. Fourier* (*Grenoble*), 48(4):1107–1127, 1998. (cited on p. 11)
- [46] Hansjörg Geiges. An introduction to contact topology, volume 109 of Cambridge Studies in Advanced Mathematics. Cambridge University Press, Cambridge, 2008. (cited on p. 5)
- [47] Rosa Gini, Liviu Ornea, and Maurizio Parton. Locally conformal Kähler reduction. J. Reine Angew. Math., 581:1–21, 2005. (cited on p. 14)
- [48] Rosa Gini, Liviu Ornea, Maurizio Parton, and Paolo Piccinni. Reduction of Vaisman structures in complex and quaternionic geometry. J. Geom. Phys., 56(12):2501–2522, 2006. (cited on p. 14)
- [49] Robert E. Gompf. A new construction of symplectic manifolds. *Ann. of Math.* (2), 142(3):527–595, 1995. (cited on p. 9)
- [50] Mark J. Gotay and James A. Isenberg. La symplectification de la science (La géométrie symplectique aux fondements de la physique et des mathématiques). *Gaz. Math.*, (54):59–79, 1992. (cited on p. 2, 7)
- [51] Ryushi Goto. On the stability of locally conformal Kähler structures. J. Math. Soc. Japan, 66(4):1375–1401, 2014. (cited on p. 14)
- [52] Alfred Gray. Some examples of almost Hermitian manifolds. *Illinois J. Math.*, 10:353–366, 1966. (cited on p. 10)
- [53] Alfred Gray. Nearly Kähler manifolds. J. Differential Geometry, 4:283–309, 1970. (cited on p. 10)
- [54] Alfred Gray. The structure of nearly Kähler manifolds. Math. Ann., 223(3):233–248, 1976. (cited on p. 10)
- [55] Alfred Gray and Luis M. Hervella. The sixteen classes of almost Hermitian manifolds and their linear invariants. *Ann. Mat. Pura Appl.* (4), 123:35–58, 1980. (cited on p. 10)
- [56] Mikhael Gromov. Stable mappings of foliations into manifolds. *Izv. Akad. Nauk SSSR Ser. Mat.*, 33:707–734, 1969. (cited on p. 1)
- [57] Mikhael Gromov. Partial differential relations, volume 9 of Ergebnisse der Mathematik und ihrer Grenzgebiete (3) [Results in Mathematics and Related Areas (3)]. Springer-Verlag, Berlin, 1986. (cited on p. 1)
- [58] Mikhael Gromov. Soft and hard symplectic geometry. In Proceedings of the International Congress of Mathematicians, Vol. 1, 2 (Berkeley, Calif., 1986), pages 81–98. Amer. Math. Soc., Providence, RI, 1987. (cited on p. 1)
- [59] Fouzia Guedira and André Lichnerowicz. Géométrie des algèbres de Lie locales de Kirillov. *J. Math. Pures Appl.* (9), 63(4):407–484, 1984. (cited on p. 4, 5)
- [60] Heinrich Guggenheimer. Sur les variétés qui possèdent une forme extérieure quadratique fermée. *C. R. Acad. Sci. Paris*, 232:470–472, 1951. (cited on p. 9)
- [61] Stefan Haller. *Perfectness and Simplicity of Certain Groups of Diffeomorphisms*. PhD thesis, Universität Wien, 1998. (cited on p. 4)
- [62] Stefan Haller and Tomasz Rybicki. On the group of diffeomorphisms preserving a locally conformal symplectic structure. Ann. Global Anal. Geom., 17(5):475–502, 1999. (cited on p. 4, 5, 6, 8)

- [63] Daniel Huybrechts. *Complex geometry. An introduction*. Universitext. Springer-Verlag, Berlin, 2005. (cited on p. 9)
- [64] Daniel Huybrechts. Projectivity of Kähler manifolds—Kodaira's problem (after C. Voisin). *Astérisque*, (311):Exp. No. 954, vii, 55–73, 2007. Séminaire Bourbaki. Vol. 2005/2006. (cited on p. 9)
- [65] Stefan Ivanov, Maurizio Parton, and Paolo Piccinni. Locally conformal parallel G₂ and Spin(7) manifolds. *Math. Res. Lett.*, 13(2-3):167–177, 2006. (cited on p. 10)
- [66] Dominic D. Joyce. *Compact manifolds with special holonomy*. Oxford Mathematical Monographs. Oxford University Press, Oxford, 2000. (cited on p. 9)
- [67] Roman Kadobianski and Jan Kubarski. Locally conformal symplectic structures and their generalizations from the point of view of Lie algebroids. *Ann. Acad. Pedagog. Crac. Stud. Math.*, 4:87–102, 2004. (cited on p. 6)
- [68] Roman Kadobianski, Jan Kubarski, Vitalij Kushnirevitch, and Robert Wolak. Transitive Lie algebroids of rank 1 and locally conformal symplectic structures. *J. Geom. Phys.*, 46(2):151–158, 2003. (cited on p. 6)
- [69] Yoshinobu Kamishima and Liviu Ornea. Geometric flow on compact locally conformally Kähler manifolds. *Tohoku Math. J.* (2), 57(2):201–221, 2005. (cited on p. 11)
- [70] Hisashi Kasuya. Vaisman metrics on solvmanifolds and Oeljeklaus-Toma manifolds. *Bull. Lond. Math. Soc.*, 45(1):15–26, 2013. (cited on p. 12)
- [71] Gerasim Kokarev. On pseudo-harmonic maps in conformal geometry. *Proc. Lond. Math. Soc.* (3), 99(1):168–194, 2009. (cited on p. 12, 13)
- [72] Gerasim Kokarev and Dieter Kotschick. Fibrations and fundamental groups of Kähler-Weyl manifolds. *Proc. Amer. Math. Soc.*, 138(3):997–1010, 2010. (cited on p. 13)
- [73] Dieter Kotschick. On products of harmonic forms. Duke Math. J., 107(3):521-531, 2001. (cited on p. 14)
- [74] Dieter Kotschick and Svjetlana Terzić. On formality of generalized symmetric spaces. Math. Proc. Cambridge Philos. Soc., 134(3):491–505, 2003. (cited on p. 14)
- [75] Calin Lazaroiu and Carlos Shahbazi. Geometric U-folds in four dimensions. J. Phys. A: Math. Theor., 2017. (cited on p. 16)
- [76] Hông Vân Lê and Jiři Vanžura. Cohomology theories on locally conformal symplectic manifolds. Asian J. Math., 19(1):45–82, 2015. (cited on p. 5)
- [77] Hwa-Chung Lee. A kind of even-dimensional differential geometry and its application to exterior calculus. *Amer. J. Math.*, 65:433–438, 1943. (cited on p. 2)
- [78] Jean Lefebvre. Transformations conformes et automorphismes de certaines structures presque symplectiques. C. R. Acad. Sci. Paris Sér. A-B, 262:A752–A754, 1966. (cited on p. 2)
- [79] Jean Lefebvre. Propriétés du groupe des transformations conformes et du groupe des automorphismes d'une variété localement conformément symplectique. C. R. Acad. Sci. Paris Sér. A-B, 268:A717–A719, 1969. (cited on p. 2, 6)
- [80] Paulette Libermann. Sur les structures presque complexes et autres structures infinitésimales régulières. Bull. Soc. Math. France, 83:195–224, 1955. (cited on p. 2, 3)
- [81] Gregory Lupton and John Oprea. Cohomologically symplectic spaces: toral actions and the Gottlieb group. *Trans. Amer. Math. Soc.*, 347(1):261–288, 1995. (cited on p. 9)
- [82] Andrzej J. Maciejewski, Maria Przybylska, and Andreĭ V. Tsiganov. On algebraic construction of certain integrable and super-integrable systems. *Phys. D*, 240(18):1426–1448, 2011. (cited on p. 8)
- [83] Farid Madani, Andrei Moroianu, and Mihaela Pilca. Conformally related K\u00e4hler metrics and the holonomy of lck manifolds. 2015. https://arxiv.org/abs/1511.09212. (cited on p. 12)
- [84] Farid Madani, Andrei Moroianu, and Mihaela Pilca. On toric locally conformally Kähler manifolds. 2016. https://arxiv.org/abs/1611.01707. (cited on p. 14)
- [85] Juan Maldacena and Carlos Nuñez. Supergravity description of field theories on curved manifolds and a no go theorem. In *Strings 2000. Proceedings of the International Superstrings Conference (Ann Arbor, MI*), volume 16, pages 822–855, 2001. (cited on p. 17)
- [86] Charles-Michel Marle. A property of conformally Hamiltonian vector fields; application to the Kepler problem. *J. Geom. Mech.*, 4(2):181–206, 2012. (cited on p. 8)
- [87] Juan C. Marrero, David Martínez Torres, and Edith Padrón. Universal models via embedding and reduction for locally conformal symplectic structures. Ann. Global Anal. Geom., 40(3):311–337, 2011. (cited on p. 5)

- [88] Dusa McDuff. Examples of simply-connected symplectic non-Kählerian manifolds. J. Differential Geom., 20(1):267–277, 1984. (cited on p. 9)
- [89] Dusa McDuff. Symplectic structures—a new approach to geometry. *Notices Amer. Math. Soc.*, 45(8):952–960, 1998. (cited on p. 2)
- [90] Dusa McDuff and Dietmar Salamon. *Introduction to symplectic topology*. Oxford Mathematical Monographs. The Clarendon Press, Oxford University Press, New York, second edition, 1998. (cited on p. 1)
- [91] Dmitri V. Millionschikov. Cohomology of solvmanifolds with local coefficients and problems in the Morse-Novikov theory. Uspekhi Mat. Nauk, 57(4(346)):183–184, 2002. (cited on p. 4)
- [92] Andrei Moroianu and Sergiu Moroianu. On Pluricanonical Locally Conformally Kähler manifolds. Int. Math. Res. Not. IMRN, 2017(14):4398–4405, 2017. (cited on p. 13)
- [93] G. D. Mostow. Cohomology of topological groups and solvmanifolds. *Ann. of Math.* (2), 73:20–48, 1961. (cited on p. 4)
- [94] Tomonori Noda. Reduction of locally conformal symplectic manifolds with examples of non-Kähler manifolds. *Tsukuba J. Math.*, 28(1):127–136, 2004. (cited on p. 5)
- [95] Karl Oeljeklaus and Matei Toma. Non-Kähler compact complex manifolds associated to number fields. *Ann. Inst. Fourier (Grenoble)*, 55(1):161–171, 2005. (cited on p. 5, 12)
- [96] Hirosi Ooguri. Geometry as seen by string theory. Jpn. J. Math., 4(2):95–120, 2009. (cited on p. 16)
- [97] Liviu Ornea. Locally conformally Kähler manifolds. A selection of results. volume 4 of *Lecture notes of Seminario Interdisciplinare di Matematica.*, pages 121–152. Semin. Interdiscip. Mat. (S.I.M.), Potenza, 2005. (cited on p. 1, 10, 13)
- [98] Liviu Ornea and Mihaela Pilca. Remarks on the product of harmonic forms. *Pacific J. Math.*, 250(2):353–363, 2011. (cited on p. 14)
- [99] Liviu Ornea and Misha Verbitsky. Structure theorem for compact Vaisman manifolds. *Math. Res. Lett.*, 10(5-6):799–805, 2003. (cited on p. 13)
- [100] Liviu Ornea and Misha Verbitsky. An immersion theorem for Vaisman manifolds. Math. Ann., 332(1):121– 143, 2005. (cited on p. 11)
- [101] Liviu Ornea and Misha Verbitsky. Locally conformal Kähler manifolds with potential. Math. Ann., 348(1):25–33, 2010. (cited on p. 11)
- [102] Liviu Ornea and Misha Verbitsky. Topology of locally conformally Kähler manifolds with potential. Int. Math. Res. Not. IMRN, (4):717–726, 2010. (cited on p. 12, 13)
- [103] Liviu Ornea and Misha Verbitsky. A report on locally conformally Kähler manifolds. In *Harmonic maps and differential geometry*, volume 542 of *Contemp. Math.*, pages 135–149. Amer. Math. Soc., Providence, RI, 2011. (cited on p. 10, 13)
- [104] Liviu Ornea and Misha Verbitsky. LCK rank of locally conformally Kähler manifolds with potential. J. Geom. Phys., 107:92–98, 2016. (cited on p. 11, 13)
- [105] Liviu Ornea and Misha Verbitsky. Embedding of LCK manifolds with potential into Hopf manifolds using Riesz-Schauder theorem. In *Complex and Symplectic Geometry*, pages 137–148. Springer, [Cham], 2017. (cited on p. 11)
- [106] Liviu Ornea, Misha Verbitsky, and Victor Vuletescu. Blow-ups of locally conformally Kähler manifolds. Int. Math. Res. Not. IMRN, (12):2809–2821, 2013. (cited on p. 14)
- [107] Alexandra Otiman. Morse-Novikov cohomology of closed one-forms of rank 1. 2016. https://arxiv.org/abs/1607.01663. (cited on p. 5)
- [108] Alexandra Otiman. Morse-Novikov cohomology of locally conformally Kähler surfaces. 2016. https://arxiv.org/abs/1609.07675. (cited on p. 14)
- [109] Alexandra Otiman and Miron Stanciu. Darboux-Weinstein theorem for locally conformally symplectic manifolds. J. Geom. Phys., 111:1–5, 2017. (cited on p. 6)
- [110] Maurizio Parton. Hopf surfaces: locally conformal Kähler metrics and foliations. *Ann. Mat. Pura Appl.* (4), 182(3):287–306, 2003. (cited on p. 11)
- [111] Maurizio Parton and Victor Vuletescu. Examples of non-trivial rank in locally conformal Kähler geometry. Math. Z., 270(1-2):179–187, 2012. (cited on p. 14)
- [112] Dietmar Salamon. Uniqueness of symplectic structures. Acta Math. Vietnam., 38(1):123–144, 2013. (cited on p. 1)

- [113] Yasha Savelyev. Gromov-Witten theory of a locally conformally symplectic manifold and the Fuller index. 2016. https://arxiv.org/abs/1609.08991. (cited on p. 6)
- [114] Hiroshi Sawai. Locally conformal Kähler structures on compact nilmanifolds with left-invariant complex structures. *Geom. Dedicata*, 125:93–101, 2007. (cited on p. 14)
- [115] Paul Seidel. Homological mirror symmetry for the quartic surface. *Mem. Amer. Math. Soc.*, 236(1116):vi+129, 2015. (cited on p. 9)
- [116] Carlos S. Shahbazi. M-theory on non-Kähler eight-manifolds. J. High Energy Phys., (9):178, front matter+30, 2015. (cited on p. 16)
- [117] Andrew Strominger. Superstrings with torsion. Nuclear Phys. B, 274(2):253-284, 1986. (cited on p. 16)
- [118] William P. Thurston. Some simple examples of symplectic manifolds. Proc. Amer. Math. Soc., 55(2):467–468, 1976. (cited on p. 9)
- [119] David Tischler. Closed 2-forms and an embedding theorem for symplectic manifolds. J. Differential Geometry, 12(2):229–235, 1977. (cited on p. 5)
- [120] Aleksy Tralle and John Oprea. *Symplectic manifolds with no Kähler structure*, volume 1661 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 1997. (cited on p. 9)
- [121] Franco Tricerri. Some examples of locally conformal Kähler manifolds. Rend. Sem. Mat. Univ. Politec. Torino, 40(1):81–92, 1982. (cited on p. 11, 14)
- [122] Luis Ugarte. Hermitian structures on six-dimensional nilmanifolds. *Transform. Groups*, 12(1):175–202, 2007. (cited on p. 13)
- [123] Izu Vaisman. On locally conformal almost K\"ahler manifolds. Israel J. Math., 24(3-4):338–351, 1976. (cited on p. 2, 7, 10)
- [124] Izu Vaisman. A theorem on compact locally conformal Kähler manifolds. Proc. Amer. Math. Soc., 75(2):279–283, 1979. (cited on p. 12)
- [125] Izu Vaisman. On locally and globally conformal Kähler manifolds. Trans. Amer. Math. Soc., 262(2):533– 542, 1980. (cited on p. 12)
- [126] Izu Vaisman. Remarkable operators and commutation formulas on locally conformal Kähler manifolds. *Compositio Math.*, 40(3):287–299, 1980. (cited on p. 4)
- [127] Izu Vaisman. Generalized Hopf manifolds. Geom. Dedicata, 13(3):231-255, 1982. (cited on p. 10, 11)
- [128] Izu Vaisman. Locally conformal symplectic manifolds. Internat. J. Math. Math. Sci., 8(3):521–536, 1985. (cited on p. 2, 4)
- [129] Misha Verbitsky. Theorems on the vanishing of cohomology for locally conformally hyper-Kähler manifolds. *Tr. Mat. Inst. Steklova*, 246(Algebr. Geom. Metody, Svyazi i Prilozh.):64–91, 2004. (cited on p. 11)
- [130] Claire Voisin. Hodge theory and complex algebraic geometry. I, volume 76 of Cambridge Studies in Advanced Mathematics. Cambridge University Press, Cambridge, 2002. (cited on p. 9, 11)
- [131] Claire Voisin. On the homotopy types of compact Kähler and complex projective manifolds. *Invent. Math.*, 157(2):329–343, 2004. (cited on p. 9)
- [132] Claire Voisin. On the homotopy types of Kähler manifolds and the birational Kodaira problem. J. Differential Geom., 72(1):43–71, 2006. (cited on p. 9)
- [133] Victor Vuletescu. Blowing-up points on l.c.K. manifolds. Bull. Math. Soc. Sci. Math. Roumanie (N.S.), 52(100)(3):387–390, 2009. (cited on p. 14)
- [134] Alan Weinstein. Symplectic manifolds and their Lagrangian submanifolds. Advances in Math., 6:329–346 (1971), 1971. (cited on p. 6)
- [135] Alan Weinstein. Lectures on symplectic manifolds. American Mathematical Society, Providence, R.I., 1977. Expository lectures from the CBMS Regional Conference held at the University of North Carolina, March 8–12, 1976, Regional Conference Series in Mathematics, No. 29. (cited on p. 6)
- [136] Hermann Weyl. *The classical groups*. Princeton Landmarks in Mathematics. Princeton University Press, Princeton, NJ, 1997. (cited on p. 1)
- [137] Edward Witten. Supersymmetry and Morse theory. J. Differential Geom., 17(4):661–692, 1982. (cited on p. 4)
- [138] Maciej P. Wojtkowski and Carlangelo Liverani. Conformally symplectic dynamics and symmetry of the Lyapunov spectrum. *Comm. Math. Phys.*, 194(1):47–60, 1998. (cited on p. 8)
- [139] Song Yang, Xiangdong Yang, and Guosong Zhao. Locally conformal symplectic blow-ups. *Differential Geom. Appl.*, 50:11–19, 2017. (cited on p. 5)

22