GLOBAL REGULARITY AND CONVERGENCE TO EQUILIBRIUM OF REACTION-DIFFUSION SYSTEMS WITH NONLINEAR DIFFUSION

KLEMENS FELLNER, EVANGELOS LATOS, BAO QUOC TANG

ABSTRACT. We study the boundedness and convergence to equilibrium of weak solutions to reaction-diffusion systems with nonlinear diffusion. The nonlinear diffusion is of porous medium type and the nonlinear reaction terms are assumed to grow polynomially and to dissipate (or conserve) the total mass. By utilising duality estimates, the dissipation of the total mass and the smoothing effect of the porous medium equation, we prove that if the exponents of the nonlinear diffusion terms are high enough, then weak solutions are bounded, locally Hölder continuous and their $L^{\infty}(\Omega)$ -norm grows in time at most polynomially.

In order to show convergence to equilibrium, we consider a specific class of nonlinear reaction-diffusion models, which describe a single reversible reaction with arbitrarily many chemical substances. By exploiting a generalised Logarithmic Sobolev Inequality, an indirect diffusion effect and the polynomial in time growth of the $L^{\infty}(\Omega)$ -norm, we show an entropy entropy-production inequality which implies exponential convergence to equilibrium in $L^{p}(\Omega)$ -norm, for any $1 \leq p < \infty$, with explicit rates and constants.

Contents

1.	Introduction and Main results	1
2.	Boundedness and local continuity of weak solutions	7
3.	Convergence to equilibrium	13
4.	Entropy-entropy production Inequality	21
5.	Proof Theorem 1.1: existence of global weak solution to (S)	25
References		27

1. INTRODUCTION AND MAIN RESULTS

In this article, we study the boundedness and convergence to equilibrium of weak solutions to reactiondiffusion systems with nonlinear diffusion

$$\begin{cases} \partial_t u_i - d_i \Delta(u_i^{m_i}) = f_i(u), & x \in \Omega, & t > 0, & i = 1, \dots, S, \\ d_i \nabla(u_i^{m_i}) \cdot \vec{n} = 0, & x \in \partial\Omega, & t > 0, & i = 1, \dots, S, \\ u_i(x, 0) = u_{i,0}(x), & x \in \Omega, & i = 1, \dots, S, \end{cases}$$
(S)

with the unknown functions $u = (u_1, \ldots, u_S)$ and $u_i : \Omega \times \mathbb{R}_+ \to \mathbb{R}$, the positive diffusion coefficients $d_i > 0$, the porous medium exponents $m_i > 1$ and where $\Omega \subset \mathbb{R}^d$ denotes a bounded domain with sufficiently smooth boundary $\partial \Omega$ (e.g. $\partial \Omega$ is of class $C^{2+\epsilon}$ for some $\epsilon > 0$) with outward unit normal \overrightarrow{n} on $\partial \Omega$. Moreover, the conditions imposed on the nonlinear reaction terms $f_i(u)$ and the nonnegative initial data $u_{i,0}$ will be specified later.

The first part of this paper considers weak solutions to system (S). Our aim is to provide sufficient conditions on the porous medium exponents m_i and on the nonlinearities $f_i(u)$, under which weak solutions are indeed bounded in L^{∞} (and thus locally Hölder-continuous) for all times and grow at most polynomially in time. More precisely, we assume the following conditions on the nonlinearities:

²⁰¹⁰ Mathematics Subject Classification. 35B35, 35B40, 35K57, 35Q92.

Key words and phrases. Reaction-Diffusion Systems; Nonlinear diffusion; Porous medium; Convergence to Equilibrium; Entropy Method.

(i) The nonlinearities $f_i : \mathbb{R}^S \to \mathbb{R}$ are locally Lipschitz functions and satisfy

$$|f_i(u)| \le C(1+|u|^{\nu}), \quad \forall u = (u_1, \dots, u_S) \in \mathbb{R}^S, \quad \forall i = 1, \dots, S,$$
(G)

where $\mathbb{R} \ni \nu \ge 1$ is the maximal growth exponent of the reaction terms.

(ii) There exist positive constants $\lambda_1, \ldots, \lambda_S > 0$ such that:

$$\sum_{i=1}^{S} \lambda_i f_i(u) \le 0, \qquad \forall u \in \mathbb{R}^S, \tag{M}$$

which formally implies the following mass dissipation law

$$\frac{d}{dt} \int_{\Omega} \sum_{i=1}^{S} \lambda_i u_i dx \le 0$$

(iii) The nonlinearities are assumed quasi-positive, that is for all i = 1, ..., S, holds

$$f(u_1, \dots, u_{i-1}, 0, u_{i+1}, \dots, u_S) \ge 0, \quad \forall u_1, \dots, u_S \ge 0.$$
 (P)

The quasi-positivity condition (P) ensures global nonnegativity of solutions subject to nonnegative initial data, see e.g. [Pie10, LP17].

The existence of global weak solutions to (S) subject to homogeneous Dirichlet boundary conditions and under the assumptions (G)-(M)-(P) was recently obtained in [LP17]. The proof of the following Theorem 1.1 on the existence of weak solutions to (S) subject to Neumann boundary conditions uses similar arguments to [LP17] and is postponed to Section 5.

Theorem 1.1. Assume the conditions (G), (M) and (P) and consider nonnegative initial data $(u_{i,0}) \in L^2(\Omega)^S$. If

$$m_i > \max\{\nu - 1; 1\} \quad for \ all \quad i = 1 \dots S,$$

then, there exists a global weak nonnegative solution to system (S) in the sense that, for all i = 1, ..., S, $u_i \in C([0, +\infty); L^1(\Omega)), u_i^{m_i} \in L^1(0, T; W^{1,1}(\Omega)), f_i(u) \in L^1(\Omega \times [0, T])$ and

$$-\int_{\Omega}\psi(0)u_{i,0}dx - \int_{0}^{T}\int_{\Omega}(u_{i}\partial_{t}\psi + d_{i}u_{i}^{m_{i}}\Delta\psi)dxdt = \int_{0}^{T}\int_{\Omega}\psi f_{i}(u)dxdt$$

for all test function $\psi \in C^{2,1}(\overline{\Omega} \times [0,T])$ with $\nabla \psi \cdot \overrightarrow{n} = 0$ on $\partial \Omega \times (0,T)$ and $\psi(\cdot,T) = 0$.

Moreover, a solution $u = (u_1, \ldots, u_S)$ to (S) with (M) and (P) satisfy

 $\|u_i\|_{L^{m_i+1}(Q_T)} \le C \quad for \ all \quad T > 0 \quad and \quad i = 1, \dots, S,$

where the constant C depends on the L^2 -norm of the initial data, the constants λ_i in (M), the diffusion coefficients $d_i > 0$ and the domain Ω .

Remark 1.1. With a more careful analysis, it seems possible to generalise Theorem 1.1 and consider initial data $u_{i,0} \in L^1(\Omega)$. We refer the interested reader to [PR16] for the case of systems with quadratic nonlinearities and L^1 initial data.

Given the weak solutions of Theorem 1.1, our aim is to establish their boundedness and a polynomially in time growing L^{∞} -estimate under stronger assumptions on the porous medium exponents m_i : First, we recall the a-priori estimate $u_i \in L^{m_i+1}(Q_T)$ of Theorem 1.1 and the growth condition (G) imply $f_i(u) \in L^{1+\epsilon}(Q_T)$ for some $\epsilon > 0$, which also justifies the definition of weak solutions in Theorem 1.1. In fact, the $L^{1+\epsilon}$ integrability guarantees uniform integrability of nonlinearities in a suitable approximating scheme (see the proof of Theorem 1.1 in the Section 5).

Intuitively, Theorem 1.1 states that larger exponents m_i yield higher integrability of the nonlinearities $f_i(u)$. Moreover, the functions u_i solve a porous medium equation with the right hand side having higher integrability. Thus, by quantifying the smoothing effect from the porous medium equation, this allows to start a bootstrap argument, which eventually leads to boundedness of u_i in L^{∞} . In particular, it is of importance that our argument allows to show that the growth in time of the L^{∞} -norms is at most polynomial. The first main result of this article is the following theorem.

Theorem 1.2 (Global bounded weak solutions).

Let $\Omega \subset \mathbb{R}^d$ be bounded with sufficiently smooth boundary. Let the initial data $0 \leq u_{i,0} \in L^{\infty}(\Omega)$, assume the conditions (G),(M) and (P) and $m_i > \max\{\nu - 1; 1\}$ for all $i = 1 \dots S$ as required by Theorem 1.1. Finally, in dimensions $d \geq 3$, we additionally assume

$$m_i > \nu - \frac{4}{d+2}, \qquad \forall i = 1 \dots S.$$
(1)

Then, any weak solution of (S) obtained in Theorem 1.1 is bounded in $L^{\infty}(\Omega)$ and grows in time at most polynomially in the sense that, for any T > 0,

$$||u_i||_{L^{\infty}(Q_T)} \le C_T, \quad \forall i = 1 \dots S$$

where C_T is a constant which depends at most polynomially on time. Consequently, these solutions are locally (in Q_T) Hölder continuous, see e.g. [Vaz07].

Remark 1.2 (Weakened assumptions on mass dissipation and initial data). If one is only interested in the boundedness of solutions but not in the polynomial growth of the L^{∞} -norm, then the mass dissipation condition (M) can in fact be weakened to

$$\sum_{i=1}^{S} \lambda_i f_i(u) \le C_1 \sum_{i=1}^{S} |u_i| + C_2 \quad \text{for all } u \in \mathbb{R}^S$$

for some positive constants C_1, C_2 .

Also the assumed initial regularity $u_{i,0} \in L^{\infty}(\Omega)$ is not optimal and could be relaxed to L^p integrability for sufficiently large p according to the details of the proof yet at the price of the readability of the Theorem.

Theorem 1.2 contributes to the large literature on global existence and boundedness of solutions to reaction-diffusion systems, which nevertheless poses still many open questions due to the lack of a unified approach (maximum principles do not hold for general systems). The largest part of the available literature, however, considers the case of linear diffusion, i.e. $m_i = 1$ in system (S). We refer the reader to the extensive review of Michel Pierre [Pie10] and the references therein, in particular [Ba94, BR10, BP00, CV09, DFPV07, HLV98, HMP87, KK00, Laa11, Mas83, Mor89, Pie03, PS97]

The case of nonlinear diffusion, on the other hand, is much less investigated. Most of the existing results considered special systems with special structures, see e.g. [Smo94, Leu09]. Up to the best of our knowledge, system (S) under the general structural assumptions (G)-(M)-(P) was only studied very recently in [LP17], where the authors showed the global existence of weak solutions. Therefore, the present paper serves as the first result to show the boundedness of weak solutions by assuming stronger conditions on porous medium exponents. Moreover, our proof allows to estimate explicitly the growth in time of the L^{∞} -norm, which turns out to be essential in studying the large time behaviour of solutions in the following second part of the paper.

The second main result of this paper proves exponential convergence to equilibrium for a class of reactiondiffusion systems with porous media diffusion of the form (S), where the nonlinearities model the following reversible reaction with arbitrarily many chemical substances

$$\alpha_1 \mathcal{A}_1 + \dots + \alpha_M \mathcal{A}_M \stackrel{k_b}{\underset{k_f}{\longleftarrow}} \beta_1 \mathcal{B}_1 + \dots + \beta_N \mathcal{B}_N.$$
⁽²⁾

Here $\alpha_i, \beta_i \in [1, +\infty)$ are the stoichiometric coefficients of the M + N involved substances $\mathcal{A}_1, \ldots, \mathcal{A}_M$, $\mathcal{B}_1, \ldots, \mathcal{B}_N$ and $k_f, k_b > 0$ are the forward and backward reaction rate constants. For simplicity, yet without loss of generality, we assume $k_f = k_b = 1$. By applying mass action kinetics to (2) and by using the short notation

$$a = (a_1, \dots, a_M), \quad b = (b_1, \dots, b_N), \quad \alpha = (\alpha_1, \dots, \alpha_M), \quad \beta = (\beta_1, \dots, \beta_N),$$
$$a^{\alpha} = \prod_{i=1}^M a_i^{\alpha_i}, \qquad b^{\beta} = \prod_{j=1}^N b_j^{\beta_j},$$

we study the following reaction-diffusion system:

$$\begin{cases} \partial_t a_i - d_i \Delta(a_i^{m_i}) = f_i(a,b) := -\alpha_i \left[a^\alpha - b^\beta \right], \ \forall i = 1, \dots, M & x \in \Omega, \quad t > 0, \\ \partial_t b_j - h_j \Delta(b_j^{p_j}) = g_j(a,b) := \beta_j \left[a^\alpha - b^\beta \right], \ \forall j = 1, \dots, N & x \in \Omega, \quad t > 0, \\ d_i \nabla(a_i^{m_i}) \cdot \overrightarrow{n} = 0, \quad \forall i = 1, \dots, M, & x \in \partial\Omega, \quad t > 0, \\ h_j \nabla(b_j^{p_j}) \cdot \overrightarrow{n} = 0, \quad \forall j = 1, \dots, N, & x \in \partial\Omega, \quad t > 0, \\ a_i(x,0) = a_{i,0}(x), \quad \forall i = 1, \dots, M, & x \in \Omega, \\ b_j(x,0) = b_{j,0}(x), \quad \forall j = 1, \dots, N, & x \in \Omega. \end{cases}$$
(R)

Here $d_i, h_j > 0$ are diffusion coefficients and $m_i, p_j > 1$ are nonlinear diffusion exponents. It is clear that (R) is a special case of (S). It is also straightforward to verify condition (P), while condition (G) is satisfied by choosing,

$$\nu = \max\left\{\sum_{i=1}^{M} \alpha_i, \sum_{j=1}^{N} \beta_j\right\}.$$

Finally condition (M) is a consequence from noting that

$$\frac{1}{M}\sum_{i=1}^{M}\frac{1}{\alpha_i}f_i(a,b) + \frac{1}{N}\sum_{j=1}^{N}\frac{1}{\beta_j}g_j(a,b) = 0.$$

After having the conditions (P), (G) and (M) verified, Theorem 1.1 implies the existence of global weak nonnegative solutions of system (R) provided

$$m_i, p_j > \max\{\nu - 1; 1\}$$
 for all $i = 1...M, j = 1...N$

Moreover by Theorem 1.2, these solutions are bounded in dimensions d = 1, 2, or in dimensions $d \ge 3$ when additionally assuming

$$m_i, p_j > \nu - \frac{4}{d+2}$$
 for all $i = 1...M, j = 1...N$

By multiplying the equations for a_i and b_j with β_j and α_i , respectively, and by adding the resulting terms, integration by parts with the homogeneous Neumann boundary conditions implies that these solutions satisfies the following mass conservation laws:

$$\beta_j \int_{\Omega} a_i(x,t) dx + \alpha_i \int_{\Omega} b_j(x,t) dx = \beta_j \int_{\Omega} a_{i,0}(x) dx + \alpha_i \int_{\Omega} b_{j,0}(x) dx =: M_{ij} > 0, \qquad \forall i,j, \qquad (3)$$

amongst which exactly M + N - 1 linearly independent conservation laws ought to be selected and only the corresponding M + N - 1 components of the initial mass vector M_{ij} need to be calculated from the initial data.

System (R) possesses for each fixed positive initial mass vector (M_{ij}) a unique positive detailed balanced equilibrium $(a_{\infty}, b_{\infty}) = (a_{1,\infty}, \ldots, a_{M,\infty}, b_{1,\infty}, \ldots, b_{N,\infty}) \in (0, \infty)^{M+N}$, which is the solutions of the following equilibrium equations:

$$\begin{cases} \prod_{i=1}^{M} a_{i\infty}^{\alpha_i} = \prod_{j=1}^{N} b_{j\infty}^{\beta_j}, \\ \beta_j a_{i\infty} + \alpha_i b_{j\infty} = M_{ij}, \quad \forall i, j, \end{cases}$$

where we recall that the second line constitutes of only M + N - 1 linearly independent conditions.

To study the convergence to equilibrium for (R), we will use the so-called entropy method, which recently proved a highly suitable tool in the analysis of the large-time-behaviour of dissipative PDE systems. With respect to reaction-diffusion systems with linear diffusion, we refer in particular to [DF06, DF07, DF08, MHM15, DFT16, FT17a, FT17].

The key *entropy functional* (or in this case the free energy functional) of system (\mathbf{R}) is defined by

$$E[a,b] = \sum_{i=1}^{M} \int_{\Omega} (a_i \ln a_i - a_i + 1) dx + \sum_{j=1}^{N} \int_{\Omega} (b_j \ln b_j - b_j + 1) dx$$

which dissipates according to the nonnegative *entropy production functional*, that is formally

$$-\frac{d}{dt}E[a,b] =: D[a,b] = \sum_{i=1}^{M} d_i \int_{\Omega} \frac{|\nabla a_i|^2}{a_i^{2-m_i}} dx + \sum_{j=1}^{N} h_j \int_{\Omega} \frac{|\nabla b_j|^2}{b_j^{2-p_j}} dx + \int_{\Omega} (a^{\alpha} - b^{\beta}) \ln \frac{a^{\alpha}}{b^{\beta}} dx \ge 0.$$

In the case of linear diffusion, i.e. $m_i = p_j = 1$ for all $i = 1 \dots M, j = 1 \dots N$, the convergence to equilibrium of solutions of (R) (or some special cases) was recently studied in e.g. [DF06, DF08, MHM15, FT17a, PSZ16].

Let us briefly review the entropy method used in the case of linear diffusion and then highlight the difficulties to be overcome in the current paper when dealing with nonlinear diffusion. In the case of linear diffusion, the entropy production writes as

$$D_{lin}[a,b] = \sum_{i=1}^{M} d_i \int_{\Omega} \frac{|\nabla a_i|^2}{a_i} dx + \sum_{j=1}^{N} h_j \int_{\Omega} \frac{|\nabla b_j|^2}{b_j} dx + \int_{\Omega} (a^{\alpha} - b^{\beta}) \ln \frac{a^{\alpha}}{b^{\beta}} dx \ge 0$$

and the entropy method consists in establishing a functional inequality of the form

$$D_{lin}[a,b] \ge \lambda(E[a,b] - E[a_{\infty},b_{\infty}]) \tag{4}$$

for all functions $a = (a_i)$, $b = (b_j)$ satisfying the conservation laws (3). In order to do that, one first uses an additivity property of the relative entropy to calculate

$$E[a,b] - E[a_{\infty}, b_{\infty}] = \left[\sum_{i=1}^{M} \int_{\Omega} a_i \log \frac{a_i}{\overline{a}_i} dx + \sum_{j=1}^{N} \int_{\Omega} b_j \log \frac{b_j}{\overline{b}_j} dx\right] \\ + \left[\sum_{i=1}^{M} (\overline{a}_i \log \frac{\overline{a}_i}{a_{i,\infty}} - \overline{a}_i + a_{i,\infty}) + \sum_{j=1}^{N} (\overline{b}_j \log \frac{\overline{b}_j}{b_{j,\infty}} - \overline{b}_j + b_{j,\infty})\right] \\ =: I_1 + I_2.$$

The term I_1 is controlled in terms of the entropy production $D_{lin}[a, b]$ thanks to the Logarithmic Sobolev Inequality (LSI)

$$\int_{\Omega} \frac{|\nabla f|^2}{f} dx \ge C_{\text{LSI}} \int_{\Omega} f \log \frac{f}{\overline{f}} dx \quad \text{for all} \quad 0 \le f \in H^1(\Omega).$$
(5)

The remain term I_2 only involves the averages of the concentrations $\overline{a}_i, \overline{b}_j$ and can be controlled by $D_{lin}[a, b]$ through lengthly, technical, but constructive estimates (see e.g. [FT17a, PSZ16] for more details). Note that this entropy approach applies successfully to more complex chemical reaction networks than (R), see [MHM15, DFT16, FT17, Mie]. We emphasised that the Logarithmic Sobolev Inequality (5) is not only used to control the term I_1 but also plays an important role in the estimates controlling the term I_2 .

In the case of nonlinear diffusion as here considered, we need a generalisation of the LSI (5) to exponents $m_i, p_j \ge 1$. In this paper, we utilise the following generalisation (see e.g. [MM17]): for any $m > (d-2)_+/d$ with $(d-2)_+ = \max\{d-2; 0\}$, there exists a constant $C(\Omega, m) > 0$ such that

$$\int_{\Omega} \frac{|\nabla f|^2}{f^{2-m}} dx \ge C(\Omega, m) \,\overline{f}^{m-1} \int_{\Omega} f \log \frac{f}{\overline{f}} dx.$$

When m = 1, this coincides with the classical Logarithmic Sobolev inequality (5). For system (R), we have in particular

$$\int_{\Omega} \frac{|\nabla a_i|^2}{a_i^{2-m_i}} dx \ge C(\Omega, m_i) \,\overline{a}_i^{m_i-1} \int_{\Omega} a_i \log \frac{a_i}{\overline{a}_i} dx \quad \text{and} \quad \int_{\Omega} \frac{|\nabla b_j|^2}{b_j^{2-p_j}} dx \ge C(\Omega, p_j) \,\overline{b}_j^{p_j-1} \int_{\Omega} b_j \log \frac{b_j}{\overline{b}_j} dx.$$
(6)

Note that if we assume the averages \overline{a}_i and \overline{b}_j to be bounded below by a positive constant, then one can apply the same strategy as for the linear diffusion case in order to obtain the convergence to equilibrium. However, there is no chemical/physical reason for such a lower bound to hold in the transient behaviour of system (**R**) subject to general initial data. There are even perfectly admissible initial conditions, where some averages are zero since the corresponding species have not yet been formed.

To overcome this difficulty, we first observe that the mass conservation laws (3) subject to a positive mass vector $M_{i,j} > 0$ implies that the averages \overline{a}_i and \overline{b}_j cannot be simultaneously small. Thus, at any fixed time at least one of the inequalities in (6) is useful, since either $\overline{a}_i \ge \varepsilon$ or $\overline{b}_j \ge \varepsilon$ for some suitably chosen $\varepsilon > 0$ depending on $M_{i,j} > 0$. Secondly, we are able to compensate the still lacking lower bounds in (6) by a phenomena which can be called "indirect diffusion effect" and which means in our context that the reversible reaction (2) transfers diffusion from a species a_i (with strictly positive diffusion bound in (6) due to $\overline{a_i} \ge \varepsilon$) to other species b_j (with lacking positive lower diffusion bound) in terms of a functional inequality, see Lemma 3.2 below.

Examples of indirect diffusion effect inequalities were already derived in e.g. [DF07, FLT17, FPT17], yet typically with a proof which requires uniform in time L^{∞} -bounds on the solutions, which is a severe technical restriction as L^{∞} -bounds for general reaction-diffusion systems are often unknown due to the lack of comparison principles. Note that also the L^{∞} -bounds of Theorem 1.1 would be insufficient since polynomially growing and not uniform in time.

In this work, we are able to prove an indirect diffusion functional inequality without using any L^{∞} bounds on solutions but instead by exploiting the special structure of (R), see Lemma 3.2. Nevertheless, in the remaining part of applying the entropy method, the polynomial growth in time of the L^{∞} -norm of Theorem 1.2 is still needed in one estimate concerning the relative entropy, yet the L^{∞} -norm appears only within a logarithm. While it is unclear to us whether this is essential or just technical necessary in our approach, it allows to derive a *time-dependent* entropy-entropy production inequality (as a generalisation of the functional inequality (4)) of the form

$$D[a(T), b(T)] \ge \Theta(T)(E[a(T), b(T)] - E[a_{\infty}, b_{\infty}]) \quad \text{for all} \quad T > 0,$$

$$\tag{7}$$

where the function $\Theta : \mathbb{R}_+ \to \mathbb{R}_+$ is of order $1/\ln(1+T)$ and satisfies $\int_0^{+\infty} \Theta(\tau) d\tau = +\infty$. Thus, a classical Gronwall argument implies explicit algebraic decay of $E[a(T), b(T)] - E[a_{\infty}, b_{\infty}]$ to zero and thus, algebraic convergence to equilibrium in relative entropy.

To obtain exponential from algebraic decay, we show that after some sufficiently large time $T_0 > 0$, the averages $\overline{a}_i(T)$ and $\overline{b}_j(T)$ are bounded below by a positive constant for all $T \ge T_0$ (since the equilibrium (a_{∞}, b_{∞}) consists of positive constants). Hence, for $T \ge T_0$, we can use the inequalities (6) like in the case for systems with linear diffusion and obtain accordingly exponential convergence to equilibrium. Finally, since T_0 can be explicitly estimated, one recovers global exponential convergence to equilibrium (i.e. for all $T \ge 0$) at the price of a smaller, yet explicit constant. Hence, the second main result of this paper is the following theorem.

Theorem 1.3. Let $\Omega \subset \mathbb{R}^d$ be bounded with sufficiently smooth boundary. Consider system (**R**) – which satisfies the conditions (**G**),(**M**) and (**P**) – subject to non-negative initial data $a_{i,0}, b_{j,0} \in L^{\infty}(\Omega)$. Assume for all $i = 1 \dots M, j = 1 \dots N$ that

$$m_i, p_j > \max\{\nu - 1; 1\}, \quad where \quad \nu = \max\left\{\sum_{i=1}^M \alpha_i, \sum_{j=1}^N \beta_j\right\}.$$

Moreover, in dimensions $d \geq 3$, we additionally assume

$$m_i, p_j > \nu - \frac{4}{d+2},$$
 for all $i = 1...M, j = 1...N.$

Finally, consider a positive initial mass vector $M_{ij} > 0$, which uniquely determines a positive equilibrium $(a_{i\infty}, b_{j\infty})$ of system (R).

Then, the bounded global weak solutions of Theorem 1.2 converge exponentially to (a_{∞}, b_{∞}) in all L^p -norms for $1 \leq p < \infty$, that is

$$\sum_{i=1}^{M} \|a_i(t) - a_{i\infty}\|_{L^p(\Omega)} + \sum_{j=1}^{N} \|b_j(t) - b_{j\infty}\|_{L^p(\Omega)} \le C e^{-\lambda_p t}$$

where the constant C > 0 and the convergence rate $\lambda_p > 0$ can be computed explicitly.

Notation:

• We denote by $\|\cdot\|$ the usual norm of $L^2(\Omega)$. For other $1 \le p < +\infty$, we write $\|\cdot\|_p$ as the norm of $L^p(\Omega)$.

• For any T > 0, $Q_T = \Omega \times (0,T)$ and $L^p(Q_T) =: L^p(0,T;L^p(\Omega))$. The space-time norm is defined as usual

$$\|f\|_{L^{p}(Q_{T})}^{p} = \int_{0}^{T} \int_{\Omega} |f(x,t)|^{p} dx dt.$$

• Throughout this work, we will denote by C_T a generic positive constant which depends on certain parameters, and more importantly C_T grows at most polynomially, i.e. there exists a polynomial P(x) such that $C_T \leq P(T)$ for all T > 0.

Organisation of the paper: Section 2 states the proof of Theorem 1.2. The proof of Theorem 1.3 is detailed in Section 3. This proof uses also a previously proven entropy-entropy production estimate for reaction-diffusion systems with linear diffusion, which is recalled in Section 4 for the sake of completeness. Finally, the existence of global weak solution is stated in Section 5.

2. Boundedness and local continuity of weak solutions

In this section, we prove for sufficiently large diffusion exponents m_i that the weak solutions obtained in Theorem 1.1 are actually bounded in L^{∞} and thus locally Hölder continuous. In Lemma 2.1, we device a bootstrap argument for the inhomogeneous porous media equation which proves that if the porous media exponents m_i and the initial integrability are high enough, then the weak solutions of Theorem 1.1 satisfy an improve integrability in a space $L^s(Q_T)$ and the L^s -norm grows at most polynomially in time T.

Lemma 2.1 (Smoothing effect of porous medium equation).

Suppose that $m \ge 1$. Assume $f \in L^{p_0}(Q_T)$ for some $p_0 > 1$ with $||f||_{L^{p_0}(Q_T)} \le C_T$. Let u be a solution to the inhomogeneous porous medium equation with positive diffusion coefficient $\delta > 0$

$$\begin{cases} \partial_t u - \delta \Delta(|u|^{m-1}u) = f, & x \in \Omega, \\ \delta \nabla(|u|^{m-1}u) \cdot \overrightarrow{n} = 0, & x \in \partial \Omega, \\ u(x,0) = u_0(x), & x \in \Omega, \end{cases}$$
(8)

and subject to initial data $u_0 \in L^{\infty}(\Omega)$. Then, u satisfies

$$\|u\|_{L^r(Q_T)} \le C_T, \quad \forall r \in [1, s)$$

where

$$s = \begin{cases} +\infty, & \text{if } p_0 \ge \frac{d+2}{2} \\ \frac{(md+2)p_0}{d+2-2p_0}, & \text{if } p_0 < \frac{d+2}{2} \end{cases}$$

and with a constant C_T , which only depends on q, d, m, Ω and at most polynomially on T.

Remark 2.1. In the linear case m = 1 Lemma 2.1 recovers the corresponding regularity estimates of the heat equation, see [CDF14]. While the smoothing effect stated in Lemma 2.1 is certainly well-known, our main contribution here lies in the polynomial growth in time of the norms, which will be crucial in Section 3.

Proof. The idea of the proof of this lemma follows [CDF14, Lemma 3.3] and is divided into several steps. Step 1. Let $\mu > 1$. By multiplying (8) by $\mu |u|^{\mu-1} \operatorname{sign}(u)$ then integrating over Ω , we obtain

$$\frac{d}{dt}\|u\|^{\mu}_{\mu} - \delta\mu \int_{\Omega} \Delta(|u|^{m-1}u)|u|^{\mu-1} \operatorname{sign}(u)dx = \mu \int_{\Omega} f|u|^{\mu-1} \operatorname{sign}(u)dx.$$
(9)

Integration by parts and the homogeneous Neumann boundary condition $\nabla(|u|^{m-1}u) \cdot \overrightarrow{n} = 0$ lead to

$$\begin{split} -\delta\mu \int_{\Omega} \Delta(|u|^{m-1})|u|^{\mu-1} \mathrm{sign}(u) dx &= m(\mu-1)\mu\delta \int_{\Omega} |u|^{m+\mu-3} |\nabla u|^2 dx + m\mu\delta \int_{\Omega} |u|^{m+p-2} |\nabla u|^2 \delta(u) dx \\ &\geq \underbrace{\frac{4m(\mu-1)\mu\delta}{(m+\mu-1)^2}}_{=:C(\mu)} \int_{\Omega} \left| \nabla \left(|u|^{\frac{m+\mu-1}{2}} \right) \right|^2 dx. \end{split}$$

By Young's inequality

$$\left| \mu \int_{\Omega} f|u|^{\mu-1} \operatorname{sign}(u) dx \right| \le \mu \|f\|_{p_0} \|u\|_{\frac{p_0(\mu-1)}{p_0-1}}^{\mu-1}$$

Therefore, it follows from (9) that

$$\frac{d}{dt} \|u\|_{\mu}^{\mu} + C(\mu) \int_{\Omega} \left| \nabla \left(|u|^{\frac{m+\mu-1}{2}} \right) \right|^2 dx \le \mu \|f\|_{p_0} \|u\|_{\frac{p_0(\mu-1)}{p_0-1}}^{\mu-1}.$$
(10)

Step 2. Choose $\mu = p_0 > 1$ in (10), we get

$$\frac{d}{dt} \|u\|_{p_0}^{p_0} + C(p_0) \int_{\Omega} \left| \nabla \left(|u|^{\frac{m+p_0-1}{2}} \right) \right|^2 dx \le p_0 \|f\|_{p_0} \|u\|_{p_0}^{p_0-1}.$$
(11)

By applying for r < 1 the elementary inequality

$$y' \le \alpha(t)y^{1-r} \implies y(T) \le \left[y(0)^r + r \int_0^T \alpha(t)dt\right]^{1/r},$$
(12)

to (11) with $r = 1/p_0$ and $y(t) = ||u(t)||_{p_0}^{p_0}$, we obtain

$$\|u(T)\|_{p_0}^{p_0} \le \left[\|u_0\|_{p_0} + \int_0^T \|f\|_{p_0} dt\right]^{p_0} \le \left[\|u_0\|_{p_0} + \|f\|_{L^{p_0}(Q_T)} T^{(p_0-1)/p_0}\right]^{p_0} =: C_{T,0}.$$
 (13)

That means

$$u \in L^{\infty}(0,T; L^{p_0}(\Omega))$$
 and $||u(T)||_{p_0}^{p_0} \le C_{T,0}$ (14)

with $C_{T,0}$ is defined in (13) grows at most polynomially in T. By integrating (11) with respect to t on (0,T)and by using Young's inequality and the convention $r_0 := m + p_0 - 1 > 1$, we get

$$C(p_0) \int_0^T \int_\Omega \left| \nabla \left(|u|^{\frac{r_0}{2}} \right) \right|^2 dx dt \le \|u_0\|_{p_0}^{p_0} + p_0 \int_0^T \|f\|_{p_0} \|u\|_{p_0}^{p_0-1} dt \le \|u_0\|_{p_0}^{p_0} + p_0 \|f\|_{L^{p_0}(Q_T)} \|u\|_{L^{p_0}(Q_T)}^{p_0-1}.$$

By adding $C(p_0) \int_0^T \!\!\int_\Omega \left| |u|^{\frac{r_0}{2}} \right|^2 dx dt$ to both sides, we have

$$C(p_{0})\int_{0}^{T}\left\|\left|u\right|^{\frac{r_{0}}{2}}\right\|_{H^{1}(\Omega)}^{2}dt = C(p_{0})\int_{0}^{T}\left[\int_{\Omega}\left|\nabla\left(\left|u\right|^{\frac{r_{0}}{2}}\right)\right|^{2}dx + \int_{\Omega}\left|\left|u\right|^{\frac{r_{0}}{2}}\right|^{2}dx\right]dt$$

$$\leq \|u_{0}\|_{p_{0}}^{p_{0}} + p_{0}\|f\|_{L^{p_{0}}(Q_{T})}\|u\|_{L^{p_{0}}(Q_{T})}^{p_{0}-1} + C(p_{0})\int_{0}^{T}\|u\|_{r_{0}}^{r_{0}}dt.$$
(15)

By the Sobolev's embedding, we have

$$C(p_0) \int_0^T \left\| |u|^{\frac{r_0}{2}} \right\|_{H^1(\Omega)}^2 \ge C(p_0) C_S^2 \int_0^T \|u\|_{s_0}^{r_0} dt \quad \text{with} \quad s_0 = \begin{cases} \frac{r_0 d}{d-2} & \text{if } d \ge 3, \\ r_0 < s_0 < \infty \text{ arbitrary} & \text{if } d = 1, 2. \end{cases}$$
(16)

On the other hand, by using the bound $||u(t)||_{p_0}^{p_0} \leq C_{T,0}$ in (14) and the interpolation inequality

$$\|u\|_{r_0} \le \|u\|_{p_0}^{\gamma} \|u\|_{s_0}^{1-\gamma} \le C_{T,0}^{\gamma/p_0} \|u\|_{s_0}^{1-\gamma} \quad \text{with} \quad \frac{1}{r_0} = \frac{\gamma}{p_0} + \frac{1-\gamma}{s_0} \quad \text{for} \quad \gamma = \frac{2p_0}{2p_0 + (m-1)d} \in (0,1],$$

we estimate in the cases m > 1 for which $\gamma < 1$

$$C(p_0) \int_0^T \|u\|_{r_0}^{r_0} dt \le C(p_0) \int_0^T C_{T,0}^{\gamma r_0/p_0} \|u\|_{s_0}^{(1-\gamma)r_0} dt \le \frac{C(p_0) C_S^2}{2} \int_0^T \|u\|_{s_0}^{r_0} dt + C C_{T,0}^{r_0/p_0} T,$$
(17)

where we have used Young's inequality (with the exponents $1 = (1 - \gamma) + \gamma$) in the last step. Note that if m = 1, the bound (17) holds still true yet without the first term and with $r_0/p_0 = 1$. Inserting (16) and (17) into (15) leads to

$$\int_{0}^{T} \|u\|_{s_{0}}^{r_{0}} dt \leq \frac{2}{C(p_{0}) C_{S}^{2}} \left[\|u_{0}\|_{p_{0}}^{p_{0}} + p_{0}\|f\|_{L^{p_{0}}(Q_{T})} \|u\|_{L^{p_{0}}(Q_{T})}^{p_{0}-1} + CC_{T,0}^{r_{0}/p_{0}}T \right] \\
\leq \frac{2}{C(p_{0}) C_{S}^{2}} \left[\|u_{0}\|_{p_{0}}^{p_{0}} + p_{0}\|f\|_{L^{p_{0}}(Q_{T})} \left(TC_{T,0}\right)^{\frac{p_{0}-1}{p_{0}}} + CC_{T,0}^{r_{0}/p_{0}}T \right] =: D_{T,0} \quad (\text{use (14)}).$$
(18)

It follows that

$$u \in L^{r_0}(0,T;L^{s_0}(\Omega)) \quad \text{with} \quad \begin{cases} s_0 = \frac{r_0 d}{d-2} & \text{if } d \ge 3, \\ r_0 < s_0 < \infty \text{ arbitrary} & \text{if } d = 1,2, \end{cases}$$
(19)

and

$$\int_0^T \|u\|_{s_0}^{r_0} dt \le D_{T,0}$$

with $D_{T,0}$ defined in (18).

Next, we construct a sequence $p_n \ge 1$ based on the estimate (14) and (19) such that

$$\|u(T)\|_{p_n}^{p_n} \le C_{T,n} \tag{20}$$

and

$$\int_{0}^{T} \|u\|_{s_{n}}^{r_{n}} dt \le D_{T,n} \quad \text{with} \quad r_{n} = m + p_{n} - 1 \quad \text{and} \quad \begin{cases} s_{n} = \frac{r_{n}d}{d-2} & \text{if } d \ge 3, \\ r_{n} < s_{n} < \infty \text{ arbitrary} & \text{if } d = 1, 2, \end{cases}$$
(21)

in which $C_{T,n}$ and $D_{T,n}$ are constants growing at most polynomially in T.

Step 3 (Iteration of (20)). In (10), we set $\mu = p_{n+1}$ for p_{n+1} to be chosen later. Thus, we have

$$\frac{d}{dt} \|u\|_{p_{n+1}}^{p_{n+1}} + C(p_{n+1}) \int_{\Omega} \left| \nabla \left(|u|^{\frac{r_{n+1}}{2}} \right) \right|^2 dx \le p_{n+1} \|f\|_{p_0} \|u\|_{\frac{p_0(p_{n+1}-1)}{p_0-1}}^{p_{n+1}-1},$$
(22)

where we recall that $r_{n+1} = m + p_{n+1} - 1$. By L^{p} - interpolation, we have

$$\|u\|_{\frac{p_0(p_{n+1}-1)}{p_0-1}} \le \|u\|_{p_{n+1}}^{1-\theta} \|u\|_{s_n}^{\theta}$$

and where $p_{n+1} > 1$ has to be chosen such that $\frac{p_0(p_{n+1}-1)}{p_0-1} \in (p_{n+1}, s_n)$ with $p_{n+1} < s_n$, which entails $\theta \in (0, 1)$ in

$$\frac{p_0 - 1}{p_0(p_{n+1} - 1)} = \frac{1 - \theta}{p_{n+1}} + \frac{\theta}{s_n}.$$
(23)

Note that $\frac{p_0(p_{n+1}-1)}{p_0-1} > p_{n+1}$ is always satisfied provided that $p_{n+1} > p_0$, i.e. that the sequence p_n is strictly monotone increasing.

It then follows from (22) (by neglecting the second term on the left hand side) that

$$\frac{d}{dt} \|u\|_{p_{n+1}}^{p_{n+1}} \le p_{n+1} \|f\|_{p_0} \|u\|_{s_n}^{\theta(p_{n+1}-1)} \left(\|u\|_{p_{n+1}}^{p_{n+1}}\right)^{1-\frac{1+\theta(p_{n+1}-1)}{p_{n+1}}}$$

By applying again the elementary inequality (12) with $y(t) = ||u(t)||_{p_{n+1}}^{p_{n+1}}$ and $r = \frac{1+\theta(p_{n+1}-1)}{p_{n+1}} < 1$, it yields

$$\begin{aligned} \|u(T)\|_{p_{n+1}}^{p_{n+1}} &\leq \left[\|u_0\|_{p_{n+1}}^{1+\theta(p_{n+1}-1)} + (1+\theta(p_{n+1}-1))\int_0^T \|f\|_{p_0} \|u\|_{s_n}^{\theta(p_{n+1}-1)} dt \right]^{\frac{p_{n+1}}{1+\theta(p_{n+1}-1)}} \\ &\leq \left[\|u_0\|_{p_{n+1}}^{1+\theta(p_{n+1}-1)} + (1+\theta(p_{n+1}-1))\|f\|_{L^{p_0}(Q_T)} \left(\int_0^T \|u\|_{s_n}^{\theta(p_{n+1}-1)\frac{p_0}{p_0-1}} dt \right)^{\frac{p_{0-1}}{p_0}} \right]^{\frac{p_{n+1}}{1+\theta(p_{n+1}-1)}}. \end{aligned}$$

$$(24)$$

In order to continue estimating by using (21), we choose p_{n+1} as

$$\theta(p_{n+1}-1)\frac{p_0}{p_0-1} = r_n.$$
(25)

Since $r_n = s_n \frac{d-2}{d}$, eq. (25) implies $\frac{\theta}{s_n} = (1 - \frac{2}{d}) \frac{p_0 - 1}{p_0(p_{n+1} - 1)}$ and thus with (23) $\theta = 1 - \frac{2}{d} \frac{p_0 - 1}{p_0} \frac{p_{n+1}}{p_{n+1} - 1} < 1.$

In order to verify that above choice of p_{n+1} satisfies $\frac{p_0(p_{n+1}-1)}{p_0-1} < s_n$, we insert (26) into (25) and calculate

$$(p_{n+1}-1)\frac{p_0}{p_0-1} - \frac{2}{d}p_{n+1} = s_n \frac{d-2}{d} \quad \Rightarrow \quad s_n - \frac{p_0(p_{n+1}-1)}{p_0-1} = \frac{2}{d}(s_n - p_{n+1}) > 0.$$

(26)

Similar, by recalling $s_n \frac{d-2}{d} = r_n = m - 1 + p_n$, we get the iteration

$$p_{n+1} = p_n \frac{d(p_0 - 1)}{p_0(d - 2) + 2} + \frac{d[(m - 1)(p_0 - 1) + p_0]}{p_0(d - 2) + 2}.$$
(27)

Altogether, by inserting (25) into (24), we obtain thanks to (21)

$$\|u(T)\|_{p_{n+1}}^{p_{n+1}} \leq \left[\|u_0\|_{p_{n+1}}^{1+\theta(p_{n+1}-1)} + (1+\theta(p_{n+1}-1))\|f\|_{L^{p_0}(Q_T)} \left(\int_0^T \|u\|_{s_n}^{r_n} dt \right)^{\frac{p_0-1}{p_0}} \right]^{\frac{r_{n+1}}{1+\theta(p_{n+1}-1)}} \leq \left[\|u_0\|_{p_{n+1}}^{1+\theta(p_{n+1}-1)} + (1+\theta(p_{n+1}-1))\|f\|_{L^{p_0}(Q_T)} D_{T,n}^{\frac{p_0-1}{p_0}} \right]^{\frac{p_{n+1}}{1+\theta(p_{n+1}-1)}} =: C_{T,n+1}$$

$$(28)$$

and thus

 $u \in L^{\infty}(0,T; L^{p_{n+1}}(\Omega))$ and $||u(T)||_{p_{n+1}}^{p_{n+1}} \le C_{T,n+1}.$ (29)

Step 4 (Iteration of (21)). We will use similar arguments to **Step 2**. Integrating (22) and adding $\int_0^T \int_{\Omega} \left| |u|^{\frac{r_{n+1}}{2}} \right|^2 dx dt$ to both sides yields in particular

$$C(p_{n+1}) \int_{0}^{T} \left\| |u|^{\frac{r_{n+1}}{2}} \right\|_{H^{1}(\Omega)}^{2} dt = C(p_{n+1}) \int_{0}^{T} \int_{\Omega} \left[\left| \nabla \left(|u|^{\frac{r_{n+1}}{2}} \right) \right|^{2} dx + \left| |u|^{\frac{r_{n+1}}{2}} \right|^{2} dx \right] dt$$

$$\leq \| u_{0} \|_{p_{n+1}}^{p_{n+1}} + p_{n+1} \int_{0}^{T} \| f \|_{p_{0}} \| u \|_{s_{n}}^{p_{0}(p_{n+1}-1)} dt + C(p_{n+1}) \int_{0}^{T} \| u \|_{r_{n+1}}^{r_{n+1}} dt$$

$$\leq \| u_{0} \|_{p_{n+1}}^{p_{n+1}} + p_{n+1} \int_{0}^{T} \| f \|_{p_{0}} \| u \|_{s_{n}}^{\theta(p_{n+1}-1)} \| u \|_{p_{n+1}}^{(1-\theta)(p_{n+1}-1)} dt + C(p_{n+1}) \int_{0}^{T} \| u \|_{r_{n+1}}^{r_{n+1}} dt \qquad (\theta \text{ in } (23))$$

$$\leq \| u_{0} \|_{p_{n+1}}^{p_{n+1}} + p_{n+1} C_{T,n+1}^{(1-\theta)(\frac{p_{n+1}-1)}{p_{n+1}}} \int_{0}^{T} \| f \|_{p_{0}} \| u \|_{s_{n}}^{\theta(p_{n+1}-1)} dt + C(p_{n+1}) \int_{0}^{T} \| u \|_{r_{n+1}}^{r_{n+1}} dt \qquad (\text{using } (29))$$

$$\leq \|u_0\|_{p_{n+1}}^{p_{n+1}} + p_{n+1}C_{T,n+1}^{(1-\theta)\frac{(p_{n+1}-1)}{p_{n+1}}} \|f\|_{L^{p_0}(Q_T)} \left(\int_0^T \|u\|_{s_n}^{r_n} dt\right)^{\frac{p_0-1}{p_0}} + C(p_{n+1})\int_0^T \|u\|_{r_{n+1}}^{r_{n+1}} dt \quad (\text{using (25)})$$

$$\leq \|u_0\|_{p_{n+1}}^{p_{n+1}} + p_{n+1}C_{T,n+1}^{(1-\theta)\frac{(p_{n+1}-1)}{p_{n+1}}}\|f\|_{L^{p_0}(Q_T)}D_{T,n}^{\frac{p_0-1}{p_0}} + C(p_{n+1})\int_0^T \|u\|_{r_{n+1}}^{r_{n+1}}dt \qquad (using (21)).$$
(30)

Now by Sobolev's embedding

$$C(p_{n+1}) \int_{0}^{T} \left\| |u|^{\frac{r_{n+1}}{2}} \right\|_{H^{1}(\Omega)}^{2} dt \ge C(p_{n+1}) C_{S}^{2} \int_{0}^{T} \|u\|_{s_{n+1}}^{r_{n+1}} dt$$
with $s_{n+1} = \begin{cases} \frac{r_{n+1}d}{d-2} & \text{if } d \ge 3, \\ r_{n+1} < s_{n+1} < \infty \text{ arbitrary} & \text{if } d = 1, 2. \end{cases}$
(31)

By the bound $||u(t)||_{p_{n+1}}^{p_{n+1}} \leq C_{T,n+1}$, the interpolation inequality

$$\|u\|_{r_{n+1}} \le \|u\|_{p_{n+1}}^{\gamma} \|u\|_{s_{n+1}}^{1-\gamma} \le C_{T,n+1}^{\gamma/p_{n+1}} \|u\|_{s_{n+1}}^{1-\gamma}$$
(32)

with
$$\frac{1}{r_{n+1}} = \frac{\gamma}{p_{n+1}} + \frac{1-\gamma}{s_{n+1}}$$
 for $\gamma = \frac{2p_{n+1}}{2p_{n+1} + (m-1)d} \in (0,1]$

Like in **Step 2** in case m > 1 and $\gamma < 1$, we have by Young's inequality,

$$C(p_{n+1}) \int_0^T \|u\|_{r_{n+1}}^{r_{n+1}} dt \le C(p_{n+1}) \int_0^T C_{T,n+1}^{\gamma r_{n+1}/p_{n+1}} \|u\|_{s_{n+1}}^{(1-\gamma)r_{n+1}} dt$$
$$\le \frac{C(p_{n+1}) C_S^2}{2} \int_0^T \|u\|_{s_{n+1}}^{r_{n+1}} dt + CTC_{T,n+1}^{r_{n+1}/p_{n+1}}$$

analog to (17) while the case m = 1 and $r_{n+1}/p_{n+1} = 1$ follows without interpolation and the first term on the right-hand-side above. Combining (30), (31) and (32) yields

$$\frac{C(p_{n+1})C_S^2}{2}\int_0^T \|u\|_{s_{n+1}}^{r_{n+1}}dt \le \|u_0\|_{p_{n+1}}^{p_{n+1}} + p_{n+1}C_{T,n+1}^{(1-\theta)\frac{(p_{n+1}-1)}{p_{n+1}}}\|f\|_{L^{p_0}(Q_T)}D_{T,n}^{\frac{p_0-1}{p_0}} + CTC_{T,n+1}^{r_{n+1}/p_{n+1}},$$

hence

$$\int_0^T \|u\|_{s_{n+1}}^{r_{n+1}} dt \le D_{T,n+1}$$

with

$$D_{T,n+1} := \frac{2}{C(p_{n+1})C_S^2} \left[\|u_0\|_{p_{n+1}}^{p_{n+1}} + p_{n+1}C_{T,n+1}^{(1-\theta)\frac{(p_{n+1}-1)}{p_{n+1}}} \|f\|_{L^{p_0}(Q_T)} D_{T,n}^{\frac{p_0-1}{p_0}} + CTC_{T,n+1}^{r_{n+1}/p_{n+1}} \right].$$
(33)

Step 5. Passing to the limit as $n \to \infty$. Considering the iteration (27), the only possible fixed point p_{∞} of the sequence p_n is

$$p_{\infty} = \frac{d[(m-1)(p_0-1)+p_0]}{2[\frac{d+2}{2}-p_0]}$$

Hence $p_{\infty} < 0$ if and only if $p_0 > \frac{d+2}{2}$. In particular, it is straightforward to check that the sequence p_n define by (27) is strictly monotone increasing if and only if either $p_n < p_{\infty}$ in the case $p_0 < \frac{d+2}{2}$ or $p_n > p_{\infty}$ in the case $p_0 > \frac{d+2}{2}$ when $p_{\infty} < 0$ holds or $p_0 = \frac{d+2}{2}$ where $p_{\infty} = +\infty$. Therefore, we have as $n \to \infty$

$$p_n \longrightarrow \begin{cases} p_{\infty} & \text{if} \quad p_0 < \frac{d+2}{2}, \\ +\infty & \text{if} \quad p_0 \ge \frac{d+2}{2} \end{cases}$$

Step 6 (Interpolation). From (20) and (21) and by using the interpolation

$$L^{\infty}(0,T;L^{p_n}(\Omega)) \cap L^{r_n}(0,T;L^{s_n}(\Omega)) \hookrightarrow L^{\frac{N+2}{N}p_n+m-1}(Q_T)$$

we get $u \in L^r(Q_T)$ for all $r < \infty$ in the case $p_0 \ge \frac{d+2}{2}$. In the case $p_0 < \frac{d+2}{2}$, we obtain $u \in L^s(Q_T)$ for all

$$s < \frac{d+2}{d}p_{\infty} + m - 1 = \frac{(md+2)p_0}{d+2 - 2p_0}.$$

This completes the proof of Lemma 2.1.

Lemma 2.2. Let u be a weak solution to (S) and

$$||u||_{L^{q_0}(Q_T)} \le C_T, \quad \forall i = 1, \dots, S, \quad with \quad q_0 > \frac{d(\nu - m) + 2(\nu - 1)}{2},$$

where $m = \min\{m_i : i = 1...S\}$ and ν is defined in (G).

Then, it follows that $||u_i||_{L^{\infty}(Q_T)} \leq C_T$ for all $i = 1 \dots S$.

Proof. From $u_i \in L^{q_0}(Q_T)$ for all $i = 1, \ldots, S$, we have $f_i(u) \in L^{q_0/\nu}(Q_T)$. Moreover note that the quasipositivity assumption (P) ensure non-negative solutions u for non-negative initial data $u_{i,0}$. Hence, the concentrations u_i satisfy the (non-sign-changing) porous media equation

$$\partial_t u_i - d_i \Delta(u_i^{m_i}) = f_i(u) \in L^{q_0/\nu}(Q_T).$$

Lemma 2.1 implies that if $q_0/\nu \geq \frac{d+2}{2}$, then $u_i \in L^r(Q_T)$ for all $r < \infty$, while if $q_0/\nu < \frac{d+2}{2}$, then

$$u_i \in L^s(Q_T)$$
 for all $s < q_1 := \frac{(md+2)q_0}{\nu(d+2) - 2q_0} \le \frac{(m_i d+2)q_0}{\nu(d+2) - 2q_0}$, for all $i = 1 \dots S$,

since $m \leq m_i$. We then construct a sequence q_n (equally for all $i = 1, \ldots, S$) such that

$$q_{n+1} = \frac{(md+2)q_n}{\nu(d+2) - 2q_n} \quad \text{for } n \ge 0.$$
(34)

It follows that

$$\frac{q_{n+1}}{q_n} = \frac{md+2}{\nu(d+2) - 2q_n}$$

Therefore, as long as $\nu(d+2) - 2q_n > 0 \iff q_n < \frac{(d+2)\nu}{2}$,

$$\frac{q_{n+1}}{q_n} > 1 \text{ for all } n \ge 0 \quad \iff \quad q_0 > \frac{d(\nu - m) + 2(\nu - 1)}{2}.$$

Hence with $q_0 > \frac{d(\nu-m)+2(\nu-1)}{2}$, after finitely many steps we arrive at $q_n > \frac{(d+2)\nu}{2}$. From $u_i \in L^s(Q_T)$ for all $s < q_n$, we have in particular $u_i \in L^{\frac{(d+2)\nu}{2}}(Q_T)$, which implies $f_i(u) \in L^{\frac{d+2}{2}}(Q_T)$ for $i = 1, \ldots, S$. By applying Lemma 2.1 once more we obtain $u_i \in L^r(Q_T) \cap L^\infty(0,T;L^q(\Omega))$ for all $r, q < \infty$. Thus,

$$\partial_t u_i - d_i \Delta(u_i^{m_i}) = f_i(u) \in L^{\infty}(0, T; L^s(\Omega)) \quad \text{ for all } s < \infty$$

with $||f_i(u)||_{L^{\infty}(0,T;L^s(\Omega))} \leq C_T$. By considering s large enough such that

$$\frac{d}{d(m_i - 1) + 2s} < 1$$

holds, we have

$$||u_i||_{L^{\infty}(Q_T)} \le C_T \quad \text{for all} \quad i = 1, \dots, S,$$

thanks to the following Lemma 2.3.

Lemma 2.3. Let $m \ge 1$. Let u be the solution to the no-flux porous media equation

$$u_t - \delta \Delta(|u|^{m-1}u) = f, \qquad \nu \cdot \nabla(|u|^{m-1}u) = 0, \qquad u(\cdot, 0) = u_0$$

with $u_0 \in L^{\infty}(\Omega)$. If $||f||_{L^{\infty}(0,T;L^p(\Omega))} \leq C_T$ with p large enough such that d/(d(m-1)+2p) < 1, then

$$\|u\|_{L^{\infty}(Q_T)} \le C_T,$$

where the constant C_T depends polynomially on T.

Remark 2.2. This regularity is well known for porous medium equation. However, since we were unable to find a reference, which includes the polynomial grow of the constant C_T , we provide in the following a proof for the sake of completeness.

Proof. Let S(t) be the nonlinear semigroup corresponding to the homogeneous equation $u_t - \delta \Delta(|u|^{m-1}u) = 0$ with homogeneous Neumann boundary condition $\nabla(|u|^{m-1}u) \cdot \vec{n} = 0$. Then, we have the following well known $L^p - L^\infty$ estimate (see e.g. [GM13, Theorem 3.2])

$$||S(t)u_0||_{L^{\infty}(\Omega)} \le C(||u_0||_{L^{p}(\Omega)} + ||u_0||_{L^{p}(\Omega)}^{\sigma_p} t^{-\alpha_p})$$

where

$$\sigma_p = \frac{2p}{d(m-1)+2p}$$
 and $\alpha_p = \frac{d}{d(m-1)+2p}$

Moreover, the semi-group propagates the L^{∞} -norm, i.e.

$$||S(t)u_0||_{L^{\infty}(\Omega)} \le ||u_0||_{L^{\infty}(\Omega)}.$$

Hence, we estimate the solution of the inhomogeneous equation by using Duhamel's formula for all $0 \le t \le T$,

$$\begin{aligned} \|u(t)\|_{L^{\infty}(\Omega)} &\leq \|S(t)u_{0}\|_{L^{\infty}(\Omega)} + \int_{0}^{t} \|S(t-s)f(s)\|_{L^{\infty}(\Omega)} ds \\ &\leq \|u_{0}\|_{L^{\infty}(\Omega)} + \int_{0}^{t} C\left(\|f(s)\|_{L^{p}(\Omega)} + \|f(s)\|_{L^{p}(\Omega)}^{\sigma_{p}}(t-s)^{-\alpha_{p}}\right) ds \\ &\leq \|u_{0}\|_{L^{\infty}(\Omega)} + CT\|f\|_{L^{\infty}(0,T;L^{p}(\Omega))} + C\|f\|_{L^{\infty}(0,T;L^{p}(\Omega))}^{\sigma_{p}} \frac{T^{1-\alpha_{p}}}{1-\alpha_{p}}, \end{aligned}$$

since $\alpha_p < 1$ by the assumption on p guarantees the convergence of the last integral on the right hand side. This finishes the proof.

Now we are ready to prove the boundedness of solutions to (S):

12

Proof of Theorem 1.2. Assuming $m_i > \nu - 1$, the existence of weak solutions follows similar to [LP17, PR16] and is proven in Section 5 in detail. By the duality estimates in Lemma 5.1, we have

$$u_i \in L^{m_i+1}(Q_T)$$
 for all $i = 1, \dots, S$.

Because $m_i > \nu - \frac{4}{2+d}$ it follows that

$$m_i + 1 > \frac{d(\nu - m_i) + 2(\nu - 1)}{2}$$

Therefore, Lemma 2.2 yields $u_i \in L^{\infty}(Q_T)$ and $||u_i||_{L^{\infty}(Q_T)} \leq C_T$ for arbitrary T > 0, which shows that the weak solutions are bounded and the $L^{\infty}(\Omega)$ norms grows at most polynomially in time.

The local Hölder continuity of the bounded weak solutions is a classical result, see e.g. [DF85] or [Vaz07, Theorem 7.17].

3. Convergence to equilibrium

In this section, we prove exponential convergence to equilibrium of solutions to (\mathbf{R}) by using the entropy method. We start by recalling the entropy (free energy) functional

$$E[a,b] = \sum_{i=1}^{M} \int_{\Omega} (a_i \ln a_i - a_i + 1) dx + \sum_{j=1}^{N} \int_{\Omega} (b_j \ln b_j - b_j + 1) dx$$

and its non-negative entropy production (free energy dissipation) functional $D[a, b] := -\frac{d}{dt}E[a, b]$, i.e.

$$D[a,b] = \sum_{i=1}^{M} d_i \int_{\Omega} \frac{|\nabla a_i|^2}{a_i^{2-m_i}} dx + \sum_{j=1}^{N} h_j \int_{\Omega} \frac{|\nabla b_j|^2}{b_j^{2-p_j}} dx + \int_{\Omega} (a^{\alpha} - b^{\beta}) \ln \frac{a^{\alpha}}{b^{\beta}} dx \ge 0,$$

where we have used the short hand notation

$$a^{\alpha} = \prod_{i=1}^{M} a_i^{\alpha_i}$$
 and $b^{\beta} = \prod_{j=1}^{N} b_j^{\beta_j}$

Moreover, the following additivity property of the relative entropy holds

$$E[a,b] - E[a_{\infty}, b_{\infty}] = \sum_{i=1}^{M} \int_{\Omega} \left(a_i \ln \frac{a_i}{a_{i\infty}} - a_i + a_{i\infty} \right) dx + \sum_{j=1}^{N} \int_{\Omega} \left(b_j \ln \frac{b_j}{b_{j\infty}} - b_j + b_{j\infty} \right) dx$$
$$= \sum_{i=1}^{M} \int_{\Omega} \left(a_i \ln \frac{a_i}{\overline{a_i}} \right) dx + \sum_{j=1}^{N} \int_{\Omega} \left(b_j \ln \frac{b_j}{\overline{b_j}} \right) dx$$
$$+ \sum_{i=1}^{M} \int_{\Omega} \left(\overline{a_i} \ln \frac{\overline{a_i}}{a_{i\infty}} - \overline{a_i} + a_{i\infty} \right) dx + \sum_{j=1}^{N} \int_{\Omega} \left(\overline{b_j} \ln \frac{\overline{b_j}}{b_{j\infty}} - \overline{b_j} + b_{j\infty} \right) dx.$$

The first Lemma 3.1 of this section states the generalisation of the Logarithmic Sobolev Inequality, which shall use in our approach.

Lemma 3.1 (A generalised Logarithmic Sobolev Inequalities, [MM17]).

Assume that $m \ge (d-2)_+/d$ where $(d-2)_+ = \max\{0, d-2\}$. Then, there exists a constant $C(\Omega, m) > 0$ such that

$$\int_{\Omega} \frac{|\nabla u|^2}{u^{2-m}} dx \ge C(\Omega, m) \,\overline{u}^{m-1} \int_{\Omega} u \ln \frac{u}{\overline{u}} dx \ge C(\Omega, m) \,\overline{u}^{m-1} \|\sqrt{u} - \overline{\sqrt{u}}\|^2$$

where $\overline{u} = \int_{\Omega} u dx$.

Proof. The first inequality follows from [MM17]. The second estimate follows from an elementary inequality:

$$\int_{\Omega} u \ln \frac{u}{\overline{u}} dx = \int_{\Omega} (u \ln \frac{u}{\overline{u}} - u + \overline{u}) dx \ge \int_{\Omega} (\sqrt{u} - \sqrt{\overline{u}})^2 dx.$$

The estimates in Lemma 3.1 constitute a generalisation of the Logarithmic Sobolev Inequality (5), which is recovered by setting m = 1 and for which the pre-factor \overline{u}^{m-1} vanishes. In the case of porous media diffusion m > 1, the pre-factor \overline{u}^{m-1} causes the lower bounds in Lemma 3.1 to degenerate for small spatial averages \overline{u} . In particular, we have by Lemma 3.1 the following lower bound for the entropy production

$$D[a,b] \geq \sum_{i=1}^{M} d_i C(\Omega,m_i) \overline{a_i}^{m_i-1} \int_{\Omega} a_i \ln \frac{a_i}{\overline{a_i}} dx + \sum_{j=1}^{N} h_j C(\Omega,p_j) \overline{b_j}^{p_j-1} \int_{\Omega} b_j \ln \frac{b_j}{\overline{b_j}} dx + \int_{\Omega} (a^{\alpha} - b^{\beta}) \ln \frac{a^{\alpha}}{b^{\beta}} dx$$

$$(35)$$

$$\geq C_0 \left[\sum_{i=1}^{M} \overline{a_i}^{m_i-1} \int_{\Omega} a_i \ln \frac{a_i}{\overline{a_i}} dx + \sum_{j=1}^{N} \overline{b_j}^{p_j-1} \int_{\Omega} b_j \ln \frac{b_j}{\overline{b_j}} dx + \int_{\Omega} (a^{\alpha} - b^{\beta}) \ln \frac{a^{\alpha}}{b^{\beta}} dx \right]$$

The problem of degeneracy appears when some averages $\overline{a_i}$ or $\overline{b_j}$ do not satisfy a positive lower bound. To overcome this problem, we first observe that due to the mass conservation laws (3) not all spatial averages can be small at the same time. If, for instance, a particular $\overline{a_i}$ is sufficiently small (w.r.t. M_{ij}) then another $\overline{b_j}$ can't be arbitrarily small because of a mass conservation law (3) connecting these two species, i.e.

$$\beta_j \overline{a_i} + \alpha_i \overline{b_j} = M_{ij} > 0, \tag{36}$$

The following crucial Lemma 3.2 shows functional inequalities, which quantity the so-called "indirect diffusion effect" and allows to compensate the lacking lower bounds for the species, whose spatial averages do not satisfy a lower bound.

We first introduce some convenient notations:

$$\begin{aligned} A_i &= \sqrt{a_i}, \ A_{i\infty} &= \sqrt{a_{i\infty}}, \\ \delta_i(x) &= A_i(x) - \overline{A_i}, \quad \forall x \in \Omega, \end{aligned} \qquad \qquad B_j &= \sqrt{b_j}, \ B_{j\infty} &= \sqrt{b_{j\infty}}, \\ \eta_j(x) &= B_j(x) - \overline{B_j}, \quad \forall x \in \Omega, \end{aligned}$$

where

$$\overline{A_i} = \int_{\Omega} A_i dx$$
 and $\overline{B_j} = \int_{\Omega} B_j dx$.

Moreover,

$$A^{\alpha} = \prod_{i=1}^{M} A_i^{\alpha_i}$$
 and $B^{\beta} = \prod_{j=1}^{N} B_j^{\beta_j}$

The conservation laws are now rewritten as

$$\beta_j \overline{A_i^2} + \alpha_i \overline{B_j^2} = M_{ij} > 0 \qquad \forall i = 1 \dots M, j = 1 \dots N.$$
(37)

Lemma 3.2 ("Indirect diffusion transfer" functional inequality).

Let $A_i, B_j : \Omega \to \mathbb{R}_+$ with i = 1...M and j = 1...N be nonnegative functions satisfying the conservation laws (37) and $\varepsilon > 0$ be a constant to be determined later. Assume that for some $J \in \{1, ..., N\}$,

$$\overline{B_j^2} \le \varepsilon \quad \text{for all} \quad j = 1 \dots J.$$

Then, there exists a constant K_1 which depends on ε such that:

$$\sum_{i=1}^{M} \|\delta_i\|^2 + \sum_{j=J+1}^{N} \|\eta_j\|^2 + \|A^{\alpha} - B^{\beta}\|^2 \ge K_1 \sum_{j=1}^{J} \|\eta_j\|^2$$
(38)

Remark 3.1. Note that when the last term on the left hand side $||A^{\alpha} - B^{\beta}||^2$ diverges, the inequality holds trivially. Therefore, in the proof we only consider the case when it is finite.

Proof. Due to the mass conservation laws (37), we have the following natural bounds,

$$\overline{A_i^2}, \overline{B_j^2} \le M_0^2, \qquad \forall i = 1, \dots, M, \ \forall j = 1, \dots, N$$

for some constant $M_0 > 0$. Therefore, by Jensen's inequality, recalling that $|\Omega| = 1$,

$$\overline{A_i} \le \sqrt{\overline{A_i^2}} \le M_0, \qquad \overline{B_j} \le \sqrt{\overline{B_j^2}} \le M_0, \quad \forall i, j.$$

From these bounds we get an upper bound for the right hand side of (38)

$$\sum_{j=1}^{J} \|\eta_j\|^2 = \sum_{j=1}^{J} (\overline{B_j^2} - \overline{B_j}^2) \le \sum_{j=1}^{J} \overline{B_j^2} \le M_0^2 J.$$

We consider the following two cases.

Case 1: If there exists $i \in \{1, ..., M\}$ such that $\|\delta_i\|^2 \ge \varepsilon$ or there exists a $j \in \{J + 1, ..., N\}$ such that $\|\eta_j\|^2 \ge \varepsilon$, we have:

$$\sum_{i=1}^{M} \|\delta_i\|^2 + \sum_{j=J+1}^{N} \|\eta_j\|^2 + \|A^{\alpha} - B^{\beta}\|^2 \ge \varepsilon \ge \frac{\varepsilon}{M_0^2 J} \sum_{j=1}^{J} \|\eta_j\|^2$$

hence, the desired inequality (38) holds with $K_1 = \frac{\varepsilon}{M_0^2 J}$. **Case 2**: Assume $\|\delta_i\|^2 \leq \varepsilon$ for all $i \in \{1, \ldots, M\}$ and $\|\eta_j\|^2 \leq \varepsilon$ for all $j \in \{J+1, \ldots, N\}$, which together with the above assumption $\overline{B_j^2} \leq \varepsilon$ and $\overline{\eta_j^2} \leq \overline{B_j^2}$ for all $j = 1 \ldots J$ implies $\|\eta_j\|^2 \leq \varepsilon$ for all $j \in \{1, \ldots, N\}$., Let $\lambda > 0$ and denote by

$$\Omega_{iA} = \{ x \in \Omega : |\delta_i(x)| \le \lambda \sqrt{\varepsilon} \} \quad \text{for } i = 1, \dots, M.$$

Then

$$\varepsilon \geq \int_{\Omega} |\delta_i(x)|^2 dx \geq \int_{\Omega \setminus \Omega_{iA}} |\delta_i(x)|^2 dx \geq \lambda^2 \varepsilon |\Omega \setminus \Omega_{iA}|$$

thus

$$|\Omega \setminus \Omega_{iA}| \le \frac{1}{\lambda^2}$$
 which implies $|\Omega_{iA}| \ge 1 - \frac{1}{\lambda^2}$

Similarly we get,

$$|\Omega_{jB}| \ge 1 - \frac{1}{\lambda^2}$$
 where $\Omega_{jB} = \{x \in \Omega : |\eta_j(x)| \le \lambda \sqrt{\varepsilon}\}$ $\forall j = 1, \dots, N.$

Now choose $\lambda^2 = 2(M+N)$ and consider $G = \bigcap_{i=1}^M \Omega_{iA} \cap_{j=1}^N \Omega_{jB}$. Then, we have $|G| \ge \frac{1}{2}$. Note that $|\delta_i(x)| \le \lambda \sqrt{\varepsilon}$ and $|\eta_j(x)| \le \lambda \sqrt{\varepsilon}$ for all $x \in G$ and for all i, j. Moreover, $\forall x \in G$

$$A_i(x) = \overline{A_i} + \delta_i(x) \le \overline{A_i} + |\delta_i(x)| \le M_0 + \lambda \sqrt{\varepsilon} \le 2M_0$$

and similarly $B_j(x) \leq 2M_0, \ \forall i, j \text{ if we choose } \varepsilon$ such that

$$\lambda \sqrt{\varepsilon} \leq M_0.$$

By Taylor's expansion, we have

$$A^{\alpha} = \prod_{i=1}^{M} A_{i}^{\alpha_{i}} = \prod_{i=1}^{M} (\overline{A_{i}} + \delta_{i})^{\alpha_{i}} = \prod_{i=1}^{M} \overline{A_{i}}^{\alpha_{i}} + R(\overline{A_{i}}, \delta_{i}) \sum_{i=1}^{M} \delta_{i}$$

where the remainder terms R depends polynomially on $\overline{A_i}$ and δ_i . Note that $|R(\overline{A_i}, \delta_i)| \leq C_0(M_0)$ on G, we estimate with $(x - y)^2 \geq \frac{1}{2}x^2 - y^2$

$$\begin{split} \|A^{\alpha} - B^{\beta}\|^{2} &= \int_{\Omega} \left(\prod_{i=1}^{M} A_{i}^{\alpha_{i}} - B^{\beta} \right)^{2} dx \\ &\geq \int_{G} \left(\prod_{i=1}^{M} \overline{A_{i}}^{\alpha_{i}} - B^{\beta} + R(\overline{A_{i}}, \delta_{i}) \sum_{i=1}^{M} \delta_{i} \right)^{2} dx \\ &\geq \frac{1}{2} \int_{G} \left(\prod_{i=1}^{M} \overline{A_{i}}^{\alpha_{i}} - B^{\beta} \right)^{2} dx - \int_{G} |R(\overline{A_{i}}, \delta_{i})|^{2} |\sum_{i=1}^{M} \delta_{i}|^{2} \\ &\geq \frac{1}{2} \int_{G} \left(\prod_{i=1}^{M} \overline{A_{i}}^{\alpha_{i}} - B^{\beta} \right)^{2} dx - C_{0}(M_{0})^{2} M \int_{G} \sum_{i=1}^{M} |\delta_{i}|^{2} \\ &\geq \frac{1}{2} \int_{G} \left(\prod_{i=1}^{M} \overline{A_{i}}^{\alpha_{i}} - B^{\beta} \right)^{2} dx - C_{0}(M_{0})^{2} M \int_{G} \sum_{i=1}^{M} |\delta_{i}|^{2} \\ &\geq \frac{1}{2} \int_{G} \left(\prod_{i=1}^{M} \overline{A_{i}}^{\alpha_{i}} - B^{\beta} \right)^{2} dx - C_{0}(M_{0})^{2} M^{2} \varepsilon \end{split}$$

where we used $\|\delta_i\|^2 \leq \varepsilon$ in the last inequality. In order to estimate further, we use again Taylor's expansion

$$B^{\beta} = \prod_{j=1}^{N} (\overline{B_j} + \eta_j)^{\beta_j} = \prod_{j=1}^{N} \overline{B_j}^{\beta_j} + Q(\overline{B_j}, \eta_j) \sum_{j=1}^{N} \eta_j$$

where again, Q depends polynomially on $\overline{B_j}, \eta_j$, which implies $|Q(\overline{B}_j, \eta_j)| \leq C_1(M_0)$ on G. Therefore,

$$\begin{split} \int_{G} \left(\prod_{i=1}^{M} \overline{A_{i}}^{\alpha_{i}} - B^{\beta} \right)^{2} dx &= \int_{G} \left(\prod_{i=1}^{M} \overline{A_{i}}^{\alpha_{i}} - \prod_{j=1}^{N} \overline{B_{j}}^{\beta_{j}} - Q(\overline{B_{j}}, \eta_{j}) \sum_{j=1}^{N} \eta_{j} \right)^{2} dx \\ &\geq \frac{1}{2} \int_{G} \left(\prod_{i=1}^{M} \overline{A_{i}}^{\alpha_{i}} - \prod_{j=1}^{N} \overline{B_{j}}^{\beta_{j}} \right)^{2} dx - \int_{G} |Q(\overline{B_{j}}, \eta_{j})|^{2} |\sum_{j=1}^{N} \eta_{j}|^{2} dx \\ &\geq \frac{1}{2} \int_{G} \left(\prod_{i=1}^{M} \overline{A_{i}}^{\alpha_{i}} - \prod_{j=1}^{N} \overline{B_{j}}^{\beta_{j}} \right)^{2} dx - C_{1}(M_{0})^{2} N^{2} \varepsilon \end{split}$$

where we used that $\|\eta_j\|^2 \leq \varepsilon$ for all j = 1, ..., N. Combining these two estimates, we arrive at

$$\|A^{\alpha} - B^{\beta}\|^{2} \ge \frac{1}{4} |G| \left(\prod_{i=1}^{M} \overline{A_{i}}^{\alpha_{i}} - \prod_{j=1}^{N} \overline{B_{j}}^{\beta_{j}} \right)^{2} - \varepsilon \left(\frac{1}{2} C_{1} (M_{0})^{2} N^{2} + C_{0} (M_{0})^{2} M^{2} \right).$$
(39)

By Jensen's inequality and the assumption of the Lemma, we have

$$\overline{B_j} \le \sqrt{\overline{B_j^2}} \le \sqrt{\varepsilon}, \quad \forall j = 1, \dots, J$$

On the other hand $\overline{B_j} \leq \sqrt{\overline{B_j^2}} \leq M_0, \ \forall j = J+1, \dots, N$. Thus, the conservation law (37) and $\|\delta_i\|^2 \leq \varepsilon$ yield

$$\overline{A}_i = \sqrt{\overline{A_i^2} - \|\delta_i\|^2} = \sqrt{\frac{1}{\beta_1}(M_{i1} - \alpha_i \overline{B_1^2}) - \|\delta_i\|^2} \ge \sqrt{\frac{M_{i1}}{\beta_1} - \frac{\alpha_i}{\beta_1}\varepsilon - \varepsilon} \quad \forall i = 1, \dots, M$$

Hence, by using $|G| \ge \frac{1}{2}$ we get from (39) that

$$\|A^{\alpha} - B^{\beta}\|^{2} \geq \frac{1}{8} \left[\prod_{i=1}^{M} \left(\frac{M_{i1}}{\beta_{1}} - \frac{\alpha_{i}}{\beta_{1}} \varepsilon - \varepsilon \right)^{\alpha_{i}/2} - \prod_{j=1}^{J} (\sqrt{\varepsilon})^{\beta_{j}} \prod_{j=J+1}^{N} M_{0}^{\beta_{j}} \right]^{2} - C_{2}\varepsilon.$$

Because the right hand side of the above inequality converges to $\frac{1}{8} \prod_{i=1}^{M} \left(\frac{M_{i1}}{\beta_1}\right)^{\alpha_i}$ as $\varepsilon \to 0$, we can choose $\varepsilon > 0$ small enough, but still explicit, such that

$$\|A^{\alpha} - B^{\beta}\|^{2} \ge \frac{1}{16} \prod_{i=1}^{M} \left(\frac{M_{i1}}{\beta_{1}}\right)^{\alpha_{i}} \ge \frac{1}{16M_{0}^{2}J} \prod_{i=1}^{M} \left(\frac{M_{i1}}{\beta_{1}}\right)^{\alpha_{i}} \sum_{j=1}^{J} \|\eta_{j}\|^{2},$$

which implies the desired inequality (38) with the constant

$$K_1 = \frac{1}{16M_0^2 J} \prod_{i=1}^M \left(\frac{M_{i1}}{\beta_1}\right)^{\alpha_i}.$$

Lemma 3.3 (An time-dependent entropy-entropy production estimate). Let $(a,b) = (a_1,\ldots,a_M,b_1,\ldots,b_N)$ with $a_i,b_j : Q_T \to \mathbb{R}_+$ be nonnegative functions, which satisfy the conservation laws (3). Moreover,

 $||a_i||_{L^{\infty}(Q_T)} \leq C_T$ and $||b_j||_{L^{\infty}(Q_T)} \leq C_T$ for all i, j.

Then, there exists a constant $K_2 > 0$ such that for all T > 0,

$$D[a(T), b(T)] \ge K_2 \frac{1}{1 + \ln(1+T)} (E[a(T), b(T)] - E[a_{\infty}, b_{\infty}]).$$

Proof. Let $\varepsilon > 0$ be a small constant chosen in Lemma 3.2. We will consider two cases and for convenience we will drop T in $a_i(T)$ and $b_i(T)$ when there is no confusion.

Case 1. Assume $\overline{a}_i \geq \varepsilon$ for all i = 1, ..., M and $\overline{b}_j \geq \varepsilon$ for all j = 1, ..., N. By applying Lemma 3.1, we have

$$D[a,b] \ge \sum_{i=1}^{M} d_i C(\Omega,m_i) \varepsilon^{m_i-1} \int_{\Omega} a_i \ln \frac{a_i}{\overline{a_i}} dx + \sum_{j=1}^{N} h_j C(\Omega,p_j) \varepsilon^{p_j-1} \int_{\Omega} b_j \ln \frac{b_j}{\overline{b_j}} dx + \int_{\Omega} (a^{\alpha} - b^{\beta}) \ln \frac{a^{\alpha}}{b^{\beta}} dx$$
$$\ge K_3 \left[\sum_{i=1}^{M} \int_{\Omega} a_i \ln \frac{a_i}{\overline{a_i}} dx + \sum_{j=1}^{N} \int_{\Omega} b_j \ln \frac{b_j}{\overline{b_j}} dx + \int_{\Omega} (a^{\alpha} - b^{\beta}) \ln \frac{a^{\alpha}}{b^{\beta}} dx \right]$$

with

$$K_3 = \min_{i=1...N} \{ d_i C(\Omega, m_i) \varepsilon^{m_i - 1}; h_j C(\Omega, p_j) \varepsilon^{p_j - 1}; 1 \}.$$

Using an entropy-entropy production inequality in case of system (\mathbf{R}) with linear diffusion, see Lemma 4.1 below, we know that

$$\sum_{i=1}^{M} \int_{\Omega} a_i \ln \frac{a_i}{\overline{a}_i} dx + \sum_{j=1}^{N} \int_{\Omega} b_j \ln \frac{b_j}{\overline{b}_j} dx + \int_{\Omega} (a^{\alpha} - b^{\beta}) \ln \frac{a^{\alpha}}{b^{\beta}} dx \ge K_4(E[a, b] - E[a_{\infty}, b_{\infty}])$$

for an explicit constant $K_4 > 0$. Therefore,

$$D[a,b] \ge K_3 K_4 (E[a,b] - E[a_\infty, b_\infty]).$$

Case 2. Suppose either $\overline{a}_i \leq \varepsilon$ for some $i \in \{1, \ldots, M\}$ or $\overline{b}_j \leq \varepsilon$ for some $j = 1, \ldots, N$. Due to the mass conservation laws $\beta_j \overline{a}_i + \alpha_i \overline{b}_j = M_{ij}$, it cannot happen that $\overline{a}_i \leq \varepsilon$ and $\overline{b}_j \leq \varepsilon$ simultaneously for a sufficiently small ε , e.g. $\varepsilon < \frac{M_{ij}}{2} \min\left\{\frac{1}{\beta_j}; \frac{1}{\alpha_i}\right\}$. Therefore, without loss of generality, we can assume that

$$\overline{b}_j \leq \varepsilon \quad \forall j = 1, \dots, J \qquad \text{and} \qquad \overline{b}_j \geq \varepsilon \quad \forall j = J+1, \dots, N$$

for some $J \in \{1, \ldots, N\}$. Moreover, by mass conservation laws

$$\overline{a}_i = \frac{1}{\beta_1} (M_{i1} - \alpha_i \overline{b}_1) \ge \frac{1}{\beta_1} (M_{i1} - \alpha_i \varepsilon), \quad \text{for all } i = 1, \dots, M.$$

Thus, we can apply Lemma 3.1 to D[a, b] and estimate

$$D[a,b] \ge \sum_{i=1}^{M} d_i C(\Omega, m_i) \left[\frac{1}{\beta_1} (M_{i1} - \alpha_i \varepsilon) \right]^{m_i - 1} \int_{\Omega} a_i \ln \frac{a_i}{a_i} dx + \sum_{j=J+1}^{N} h_j C(\Omega, p_j) \varepsilon^{p_j - 1} \int_{\Omega} b_j \ln \frac{b_j}{\overline{b}_j} dx + \int_{\Omega} (a^{\alpha} - b^{\beta}) \ln \frac{a^{\alpha}}{b^{\beta}} dx \ge K_5 \left[\sum_{i=1}^{M} \|\sqrt{a_i} - \overline{\sqrt{a_i}}\|^2 + \sum_{j=J+1}^{N} \|\sqrt{b_j} - \overline{\sqrt{b_j}}\|^2 + \|A^{\alpha} - B^{\beta}\|^2 \right] = K_5 \left[\sum_{i=1}^{M} \|\delta_i\|^2 + \sum_{j=J+1}^{N} \|\eta_j\|^2 + \|A^{\alpha} - B^{\beta}\|^2 \right],$$

where we have used $(x - y) \ln(x/y) \ge 4(\sqrt{x} - \sqrt{y})^2$ and

$$K_5 = \min_{i=1\dots M; j=J+1\dots N} \left\{ d_i C(\Omega, m_i) \left[\frac{1}{\beta_1} (M_{i1} - \alpha_i \varepsilon) \right]^{m_i - 1}; h_j C(\Omega, p_j) \varepsilon^{p_j - 1}; 4 \right\}.$$

Applying Lemma 3.2 yields

$$D[a,b] \ge K_6 \left[\sum_{i=1}^M \|\delta_i\|^2 + \sum_{j=1}^N \|\eta_j\|^2 + \|A^{\alpha} - B^{\beta}\|^2 \right]$$

where

$$K_6 = \frac{1}{2} \min\{K_5; K_5 K_1\}.$$

By using another functional inequality, which was already proven in the case of linear diffusion, see (48) in Section 4, we have

$$D[a,b] \ge K_7 \left[\sum_{i=1}^{M} (\|\delta_i\|^2 + |\sqrt{A_i^2} - A_{i,\infty}|^2) + \sum_{j=1}^{N} (\|\eta_j\|^2 + |\sqrt{B_j^2} - B_{j,\infty}|^2) \right].$$
(40)

Now, we estimate $E[a, b] - E[a_{\infty}, b_{\infty}]$ from above. Consider the two variables function

$$\Phi(x,y) = \frac{x\ln(x/y) - x + y}{(\sqrt{x} - \sqrt{y})^2}$$

which is continuous in $(0,\infty)^2$ and $\Phi(\cdot,y)$ is increasing for each fixed y > 0. It holds that

$$\begin{split} E[a,b] - E[a_{\infty},b_{\infty}] \\ &= \sum_{i=1}^{M} \int_{\Omega} \Phi(a_{i},a_{i,\infty}) (A_{i} - A_{i,\infty})^{2} dx + \sum_{j=1}^{N} \int_{\Omega} \Phi(b_{j},b_{j,\infty}) (B_{j} - B_{j,\infty})^{2} dx \\ &\leq \max_{i=1...M; j=1...N} \{ \Phi(\|a_{i}\|_{L^{\infty}(Q_{T})},a_{i,\infty}); \Phi(\|b_{j}\|_{L^{\infty}(Q_{T})},b_{j,\infty}) \} \left[\sum_{i=1}^{M} \|A_{i} - A_{i,\infty}\|^{2} + \sum_{j=1}^{N} \|B_{j} - B_{j,\infty}\|^{2} \right] \\ &\leq K_{8}(1 + \ln(1+T)) \left[\sum_{i=1}^{M} (\|\delta_{i}\|^{2} + |\overline{A}_{i} - A_{i,\infty}|^{2}) + \sum_{j=1}^{N} (\|\eta_{j}\|^{2} + |\overline{B}_{j} - B_{j,\infty}|^{2}) \right], \end{split}$$

$$(41)$$

where in the last inequality we have used the estimates $||a_i||_{L^{\infty}(Q_T)} \leq C_T$ and $||b_j||_{L^{\infty}(Q_T)} \leq C_T$ and that C_T is a constant growing at most polynomially w.r.t. T.

Next, from
$$\|\delta_i\|^2 = \overline{A_i^2} - \overline{A}_i^2 = (\sqrt{\overline{A_i^2}} - \overline{A}_i)(\sqrt{\overline{A_i^2}} + \overline{A}_i)$$
, we have
 $\overline{A}_i = \sqrt{\overline{A_i^2}} - \frac{\|\delta_i\|^2}{\sqrt{\overline{A_i^2}} + \overline{A}_i} = \sqrt{\overline{A_i^2}} - Q_i(A_i)\|\delta_i\|$ with $Q_i(A_i) = \frac{\|\delta_i\|}{\sqrt{\overline{A_i^2}} + \overline{A}_i}$.

It's obvious that $Q(A_i) \ge 0$ and moreover

$$Q_i(A_i)^2 = \frac{\overline{A_i^2} - \overline{A}_i^2}{(\sqrt{\overline{A_i^2}} + \overline{A}_i)^2} = \frac{\sqrt{\overline{A_i^2} - \overline{A}_i}}{\sqrt{\overline{A_i^2}} + \overline{A}_i} \le 1.$$

Therefore,

$$|\overline{A}_{i} - A_{i,\infty}|^{2} \leq 2 \left(|\sqrt{\overline{A_{i}^{2}}} - \overline{A}_{i}|^{2} + |\sqrt{\overline{A_{i}^{2}}} - A_{i,\infty}|^{2} \right)$$
$$= 2 \left(Q_{i}(A_{i})^{2} ||\delta_{i}||^{2} + |\sqrt{\overline{A_{i}^{2}}} - A_{i,\infty}|^{2} \right)$$
$$\leq 2 \left(||\delta_{i}||^{2} + |\sqrt{\overline{A_{i}^{2}}} - A_{i,\infty}|^{2} \right) \quad \text{for all } i = 1 \dots M$$

and similarly

$$|\overline{B}_j - B_{j,\infty}|^2 \le 2\left(\|\eta_i\|^2 + |\sqrt{\overline{B_j^2}} - B_{j,\infty}|^2\right) \quad \text{for all } j = 1\dots N.$$

Hence it follows from (41) that

$$E[a,b] - E[a_{\infty},b_{\infty}] \le 3K_8(1 + \ln(1+T)) \left[\sum_{i=1}^{M} (\|\delta_i\|^2 + |\sqrt{A_i^2} - A_{i,\infty}|^2) + \sum_{j=1}^{N} (\|\eta_j\|^2 + |\sqrt{B_j^2} - B_{j,\infty}|^2) \right].$$
(42)

A combination of (40) and (42) yields

$$D[a,b] \ge \frac{K_7}{3K_8(1+\ln(1+T))} (E[a,b] - E[a_{\infty},b_{\infty}]).$$

Finally, from Case 1 and Case 2, we can conclude the proof of Lemma 3.3 with

$$K_2 = \min\left\{K_3 K_4; \frac{K_7}{3K_8}\right\}.$$

Remark 3.2. The assumptions $||a_i||_{L^{\infty}(Q_T)} \leq C_T$ and $||b_j||_{L^{\infty}(Q_T)} \leq C_T$ in Lemma 3.3 are only needed to estimate $E[a,b] - E[a_{\infty},b_{\infty}]$ above as in (41). In the case of linear diffusion, it is possible to avoid these L^{∞} -bounds by using the additivity of the relative entropy (see also the proof of Lemma 4.1 in Section 4), i.e.

$$E[a,b] - E[a_{\infty},b_{\infty}] = (E[a,b] - E[\overline{a},\overline{b}]) + (E[\overline{a},\overline{b}] - E[a_{\infty},b_{\infty}])$$

However, while for linear diffusion, the Logarithmic Sobolev Inequality controls to first part $E[a, b] - E[\overline{a}, \overline{b}] \leq C(C_{\text{LSI}})D[a, b]$, such an estimate is unclear in the case of porous media diffusion, where the generalised Logarithmic Sobolev Inequality in Lemma 3.1 degenerates for states without lower bounds on the spatial averages.

We need also the following Csiszár-Kullback-Pinsker type inequality. The proof is standard and can be found in e.g. [DFT16, FT17a].

Lemma 3.4. There exists a constant $C_{CKP} > 0$ such that for any measurable nonnegative functions $a_i, b_j : \Omega \to \mathbb{R}_+$ satisfying the mass conservation (36), there holds

$$E[a,b] - E[a_{\infty},b_{\infty}] \ge C_{CKP} \left(\sum_{i=1}^{M} \|a_i - a_{i,\infty}\|_1^2 + \sum_{j=1}^{N} \|b_j - b_{j,\infty}\|_1^2 \right).$$

We are ready to prove Theorem 1.3.

Proof of Theorem 1.3. Due to the condition

$$m_i, p_j > \max\left\{\nu - \min\left\{\frac{4}{d+2}; 1\right\}; 1\right\} \qquad \forall i = 1 \dots M, j = 1 \dots N,$$

we can apply Theorem 1.2 to show boundedness of the weak solution (a, b) to (\mathbf{R}) , i.e.

$$||a_i||_{L^{\infty}(Q_T)} \le C_T, \quad ||b_j||_{L^{\infty}(Q_T)} \le C_T, \quad \forall i = 1...M, j = 1...N$$

By applying Lemma 3.3 this yields

$$D[a(T), b(T)] \ge K_2 \frac{1}{1 + \ln(1+T)} (E[a(T), b(T)] - E[a_{\infty}, b_{\infty}]).$$

Moreover, due to the boundedness of solutions, we have the entropy-entropy production relation

$$\frac{d}{dt}(E[a,b] - E[a_{\infty},b_{\infty}]) = \frac{d}{dt}E[a,b] = -D[a,b] \le -K_2 \frac{1}{1 + \ln(1+T)}(E[a,b] - E[a_{\infty},b_{\infty}]).$$

A classical Gronwall's inequality leads to

$$E[a(T), b(T)] - E[a_{\infty}, b_{\infty}] \le \exp\left(-K_2 \int_0^T \frac{d\tau}{1 + \ln(1 + \tau)}\right) (E[a_0, b_0] - E[a_{\infty}, b_{\infty}]).$$

By direct calculations

$$\exp\left(-K_2 \int_0^T \frac{d\tau}{1+\ln(1+\tau)}\right) \ge \exp\left(-K_2 \int_0^T \frac{d\tau}{1+\tau}\right) = (1+T)^{-K_2}.$$

Hence,

$$E[a(T), b(T)] - E[a_{\infty}, b_{\infty}] \le (1+T)^{-K_2} (E[a_0, b_0] - E[a_{\infty}, b_{\infty}]),$$
(43)

and therefore thanks to the Csiszár-Kullback-Pinsker inequality in Lemma 3.4

$$\sum_{i=1}^{M} \|a_i(T) - a_{i,\infty}\|_1^2 + \sum_{j=1}^{N} \|b_j(T) - b_{j,\infty}\|_1^2 \le C_{CKP}^{-1} (1+T)^{-K_2} (E[a_0, b_0] - E[a_\infty, b_\infty])$$
(44)

which implies algebraic convergence to equilibrium of solutions to (R).

We will now show that from this it is possible to recover exponential convergence. Since the right hand side of (44) tends to zero as $T \to \infty$, we can choose

$$T_{0} = \max\left\{1; \left[\frac{C_{CKP}^{-1}(E[a_{0}, b_{0}] - E[a_{\infty}, b_{\infty}])}{\frac{1}{2}\min_{i=1...N}\{a_{i,\infty}^{2}, b_{j,\infty}^{2}\}}\right]^{1/K_{2}} - 1\right\}$$
(45)

which implies for all $t \ge T_0$

$$||a_i(t) - a_{i,\infty}||_1 \le \frac{1}{2}a_{i,\infty}$$
 and $||b_j(t) - b_{j,\infty}||_1 \le \frac{1}{2}b_{j,\infty}$

and thus

$$\overline{a}_i(t) = \|a_i(t)\|_1 \ge \frac{1}{2}a_{i,\infty}$$
 and $\overline{b}_j(t) = \|b_j(t)\|_1 \ge \frac{1}{2}b_{j,\infty}$ for all $t \ge T_0$.

Therefore, for all $t \ge T_0$, we can apply these lower bounds on the spatial averages bounds and Lemma 3.1 to estimate the entropy-entropy production as follows

$$D[a(t), b(t)] \ge C_1 \left[\sum_{i=1}^M \int_{\Omega} a_i \ln \frac{a_i}{a_i} dx + \sum_{j=1}^N \int_{\Omega} b_j \ln \frac{b_j}{b_j} dx + \int_{\Omega} (a^{\alpha} - b^{\beta}) \ln \frac{a^{\alpha}}{b^{\beta}} dx \right] \quad \text{for all } t \ge T_0,$$

with

$$C_{1} = \min_{i=1...M; j=1...N} \left\{ d_{i}C(\Omega, m_{i}) \left(\frac{1}{2}a_{i,\infty}\right)^{m_{i}-1}; h_{j}C(\Omega, p_{j}) \left(\frac{1}{2}b_{j,\infty}\right)^{p_{j}-1}; 1 \right\}.$$

By applying again Lemma 4.1, we obtain

$$D[a(t), b(t)] \ge C_1 \lambda(E[a(t), b(t)] - E[a_{\infty}, b_{\infty}]) \quad \text{for all} \quad t \ge T_0,$$

which in a combination with the classical Gronwall's inequality yields for all $t \ge T_0$,

$$E[a(t), b(t)] - E[a_{\infty}, b_{\infty}] \le e^{-\lambda C_1 (t - T_0)} (E[a(T_0), b(T_0)] - E[a_{\infty}, b_{\infty}])$$

$$\le e^{-\lambda C_1 t} e^{\lambda C_1 T_0} (1 + T_0)^{-K_2} (E[a_0, b_0] - E[a_{\infty}, b_{\infty}])$$

$$\le e^{-\lambda C_1 t} e^{\lambda C_1 T_0} (E[a_0, b_0] - E[a_{\infty}, b_{\infty}])$$

where we used (43) for the second inequality. On the other hand, it follows from (43) that for all $0 \le t < T_0$,

$$E[a(t), b(t)] - E[a_{\infty}, b_{\infty}] \le (1+t)^{-K_2} (E[a_0, b_0] - E[a_{\infty}, b_{\infty}])$$
$$\le e^{-\lambda C_1 t} e^{\lambda C_1 T_0} (E[a_0, b_0] - E[a_{\infty}, b_{\infty}])$$

Due to the explicitness of T_0 in (45), we eventually get the exponential convergence

$$E[a(t), b(t)] - E[a_{\infty}, b_{\infty}] \le C_2 e^{-\hat{\lambda}t} (E[a_0, b_0] - E[a_{\infty}, b_{\infty}]) \text{ for all } t \ge 0,$$

with the constant $C_2 = e^{\lambda C_1 T_0}$ and the rate $\hat{\lambda} = \lambda C_1$. Note that C_2 is explicit since T_0 is explicit (see (45)). With another application of the Csiszár-Kullback-Pinsker inequality in Lemma 3.4, this yields

$$\sum_{i=1}^{M} \|a_i(t) - a_{i,\infty}\|_1^2 + \sum_{j=1}^{N} \|b_j(t) - b_{j,\infty}\|_1^2 \le C_2 C_{CKP}^{-1} e^{-\widehat{\lambda}t} (E[a_0, b_0] - E[a_\infty, b_\infty]) \le C_3 e^{-\widehat{\lambda}t}$$

with $C_3 = C_2 C_{CKP}^{-1}(E[a_0, b_0] - E[a_{\infty}, b_{\infty}])$. Finally, by combining the above exponential L^1 -convergence with the at most polynomial grow L^{∞} a-priori estimates $||a_i||_{L^{\infty}(Q_T)}, ||b_j||_{L^{\infty}(Q_T)} \leq C_T$, interpolation yields for any 1 ,

$$\|a_i(T) - a_{i,\infty}\|_p \le \|a_i(T) - a_{i,\infty}\|_{\infty}^{\theta} \|a_i(T) - a_{i,\infty}\|_1^{1-\theta} \le C_T^{\theta} C_3^{1-\theta} e^{-\widehat{\lambda}(1-\theta)T} \le C_4 e^{-\lambda_p T}$$

for some $0 < \lambda_p < \hat{\lambda}(1-\theta)$ since C_T grows at most polynomially in T, and similarly

$$\|b_j(T) - b_{j,\infty}\|_p \le \|b_j(T) - b_{j,\infty}\|_{\infty}^{\theta} \|b_j(T) - b_{j,\infty}\|_1^{1-\theta} \le C_5 e^{-\lambda_p T}.$$

This concludes the proof of Theorem 1.3.

4. Entropy-entropy production Inequality

Lemma 4.1 (Entropy-entropy production estimate). Let $a_{\infty} \in (0, \infty)^M$ and $b_{\infty} \in (0, \infty)^N$ satisfy

$$a_{\infty}^{\alpha} = b_{\infty}^{\beta}$$

where $\alpha \in [1,\infty)^M$ and $\beta \in [1,\infty)^N$.

Then, there exists an explicit constant $\lambda > 0$ depending on a_{∞} , b_{∞} , α , β and the domain Ω , such that for any nonnegative functions $a = (a_i) : \Omega \to \mathbb{R}^M_+$ and $b = (b_j) : \Omega \to \mathbb{R}^N_+$ satisfying

$$\beta_j \overline{a}_i + \alpha_i \overline{b}_j = \beta_j a_{i,\infty} + \alpha_i b_{j,\infty}$$
 for all $i = 1, \dots, M, \ j = 1, \dots, N,$

the following entropy-entropy production inequality holds

$$D[a,b] \ge \lambda(E[a,b] - E[a_{\infty},b_{\infty}])$$

where

$$\widetilde{D}[a,b] = \sum_{i=1}^{M} \int_{\Omega} a_i \ln \frac{a_i}{\overline{a}_i} dx + \sum_{j=1}^{N} \int_{\Omega} b_j \ln \frac{b_j}{\overline{b}_j} dx + \int_{\Omega} (a^{\alpha} - b^{\beta}) \ln \frac{a^{\alpha}}{b^{\beta}} dx$$

and

$$E[a,b] = \sum_{i=1}^{M} \int_{\Omega} (a_i \ln a_i - a_i + 1) dx + \sum_{j=1}^{N} \int_{\Omega} (b_j \ln b_j - b_j + 1) dx$$

Remark 4.1. The above entropy-entropy production inequality was first proved in [FT17a] in a constructive way with explicit bounds on the constant λ . The proof stated here follows the line of a significantly simplified version presented in [FT17].

Proof. First, by the additivity of the relative entropy, we have

$$\begin{split} E[a,b] - E[a_{\infty},b_{\infty}] &= (E[a,b] - E[\overline{a},\overline{b}]) + (E[\overline{a},\overline{b}] - E[a_{\infty},b_{\infty}]) \\ &= \left[\sum_{i=1}^{M} \int_{\Omega} a_{i} \ln \frac{a_{i}}{\overline{a}_{i}} dx + \sum_{j=1}^{N} \int_{\Omega} b_{j} \ln \frac{b_{j}}{\overline{b}_{j}} dx \right] \\ &+ \left[\sum_{i=1}^{M} \left(\overline{a}_{i} \ln \frac{\overline{a}_{i}}{a_{i,\infty}} - \overline{a}_{i} + a_{i,\infty} \right) + \sum_{j=1}^{N} \left(\overline{b}_{j} \ln \frac{\overline{b}_{j}}{b_{j,\infty}} - \overline{b}_{j} + b_{j,\infty} \right) \right] \\ &=: (I) + (II). \end{split}$$

It is straightforward that (I) can be controlled by $\widetilde{D}[a, b]$, i.e.

$$\frac{1}{2}\widetilde{D}[a,b] \ge \frac{1}{2} \times (I).$$

It remains to control (II). To do that, we first introduce the following useful notations and definitions

$$A_{i} = \sqrt{a_{i}}, \quad B_{j} = \sqrt{b_{j}}, \quad A_{i,\infty} = \sqrt{a_{i,\infty}}, \quad B_{j,\infty} = \sqrt{b_{j,\infty}}$$
$$\delta_{i}(x) = A_{i}(x) - \overline{A}_{i}, \qquad \eta_{j}(x) = B_{j}(x) - \overline{B}_{j},$$

and

$$A^{\alpha} = \prod_{i=1}^{M} A_i^{\alpha_i}, \quad B^{\beta} = \prod_{j=1}^{N} B_j^{\beta_j}.$$

By the elementary inequality $(x - y) \ln(x/y) \ge 4(\sqrt{x} - \sqrt{y})^2$, we have

$$\int_{\Omega} a_i \ln \frac{a_i}{\overline{a_i}} dx = \int_{\Omega} \left(a_i \ln \frac{a_i}{\overline{a_i}} - a_i + \overline{a_i} \right) dx \ge 4 \int_{\Omega} (\sqrt{a_i} - \sqrt{\overline{a_i}})^2 dx \ge 4 \|\delta_i\|^2$$

and similarly $\int_{\Omega} b_j \ln \frac{b_j}{b_j} dx \ge 4 \|\eta_j\|^2$. Moreover, $\int_{\Omega} (a^{\alpha} - b^{\beta}) \ln \frac{a^{\alpha}}{b^{\beta}} dx \ge 4 \|A^{\alpha} - B^{\beta}\|^2$. Therefore,

$$\frac{1}{2}\widetilde{D}[a,b] \ge 2\left[\sum_{i=1}^{M} \|\delta_i\|^2 + \sum_{j=1}^{N} \|\eta_j\|^2 + \|A^{\alpha} - B^{\beta}\|^2\right].$$
(46)

In order to bound to estimate the right-hand-side of (46) with an upper bound of (II), we first observe from the conservation laws

$$\beta_j \overline{a}_i + \alpha_i \overline{b}_j = \beta_j a_{i,\infty} + \alpha_i b_{j,\infty}, \qquad \text{for all } i, j$$

that there exists a constant $M_0 > 0$ such that

$$\overline{a}_i, \overline{b}_j \leq M_0^2, \quad \text{for all } i, j.$$

Next, we note that the two variables function

$$\Phi(x,y) = \frac{x \ln(x/y) - x + y}{(\sqrt{x} - \sqrt{y})^2}$$

is continuous on $(0,\infty)^2$ and $\Phi(\cdot,y)$ is increasing for each fixed y. Then, the term (II) is estimated as

$$(II) = \sum_{i=1}^{M} \Phi(\overline{a}_{i}, a_{i,\infty}) (\sqrt{\overline{a}_{i}} - \sqrt{a_{i,\infty}})^{2} + \sum_{j=1}^{N} \Phi(\overline{b}_{j}, b_{j,\infty}) (\sqrt{\overline{b}_{j}} - \sqrt{b_{j,\infty}})^{2}$$

$$\leq \max_{i,j} \{\Phi(M_{0}^{2}, a_{i,\infty}); \Phi(M_{0}^{2}, b_{j,\infty})\} \left(\sum_{i=1}^{M} (\sqrt{\overline{A_{i}^{2}}} - A_{i,\infty})^{2} + \sum_{j=1}^{N} (\sqrt{\overline{B_{j}^{2}}} - B_{j,\infty})^{2}\right).$$
(47)

From (46) and (47), it remains to show that

$$\sum_{i=1}^{M} \|\delta_i\|^2 + \sum_{j=1}^{N} \|\eta_j\|^2 + \|A^{\alpha} - B^{\beta}\|^2 \ge C_0 \left(\sum_{i=1}^{M} (\sqrt{\overline{A_i^2}} - A_{i,\infty})^2 + \sum_{j=1}^{N} (\sqrt{\overline{B_j^2}} - B_{j,\infty})^2 \right)$$
(48)

for some constant $C_0 > 0$. By using Lemma 4.2, we have with $\overline{A} = (\overline{A}_1, \ldots, \overline{A}_M)$ and $\overline{B} = (\overline{B}_1, \ldots, \overline{B}_N)$

$$\sum_{i=1}^{M} \|\delta_i\|^2 + \sum_{j=1}^{N} \|\eta_j\|^2 + \|A^{\alpha} - B^{\beta}\|^2 \ge C_1 \left(\sum_{i=1}^{M} \|\delta_i\|^2 + \sum_{j=1}^{N} \|\eta_j\|^2 + \left|\overline{A}^{\alpha} - \overline{B}^{\beta}\right|^2 \right)$$
(49)

for some constant $C_1 > 0$. Using the ansatz

$$\overline{A_i^2} = A_{i,\infty}^2 (1+\mu_i)^2 \quad \text{and} \quad \overline{B_j^2} = B_{j,\infty}^2 (1+\zeta_j)^2, \quad \text{where} \quad \mu_i, \zeta_j \in [-1,\infty),$$
(50)

the right hand side of (48) writes as

RHS of (48) =
$$C_0 \left(\sum_{i=1}^{M} \mu_i^2 + \sum_{j=1}^{N} \zeta_j^2 \right).$$
 (51)

Moreover, the bounds $\overline{a_i} = \overline{A_i^2} \le M_0^2$ and $\overline{b_j} = \overline{B_j^2} \le M_0^2$ imply

$$-1 \le \mu_i \le M_1$$
 and $-1 \le \zeta_j \le M_1$ (52)

for some constant $M_1 > 0$. From the ansatz (50) (and similar to the proof of Lemma 3.3), we have

$$\overline{A}_{i} = \sqrt{\overline{A}_{i}^{2}} - Q_{i}(A_{i}) \|\delta_{i}\| = A_{i,\infty}(1+\mu_{i}) - Q_{i}(A_{i}) \|\delta_{i}\|$$

$$\overline{B}_{j} = \sqrt{\overline{B}_{j}^{2}} - R_{j}(B_{j}) \|\eta_{j}\| = B_{j,\infty}(1+\zeta_{j}) - R_{j}(B_{j}) \|\eta_{j}\|$$

where

$$0 \le Q_i(A_i) := \frac{\|\delta_i\|}{\sqrt{A_i^2} + \overline{A_i}} \le 1 \quad \text{and} \quad 0 \le R_j(B_j) := \frac{\|\eta_j\|}{\sqrt{\overline{B_j^2}} + \overline{B_j}} \le 1.$$

Next, we use Taylor expansion to estimate

$$\overline{A_i}^{\alpha_i} = (A_{i,\infty}(1+\mu_i) - Q_i(A_i) \|\delta_i\|)^{\alpha_i} = A_{i,\infty}^{\alpha_i}(1+\mu_i)^{\alpha_i} + \widehat{Q}_i \|\delta_i\|$$

in which the Lagrange remainder term $\widehat{Q}_i = \widehat{Q}(\mu_i, \|\delta_i\|)$ is uniformly bounded above by a constant for all admissible values of μ_i and $\|\delta_i\|$ thanks to the boundedness of μ_i and $\|\delta_i\| \leq \sqrt{A_i^2} \leq M_0$. Similarly,

$$\overline{B_j}^{\beta_j} = B_{j,\infty}^{\beta_j} (1+\zeta_j)^{\beta_j} + \widehat{R}_j ||\eta_j|$$

with uniformly bounded remainder $\widehat{R}_j(\zeta_j, ||\eta_j||)$. Thus

$$\begin{aligned} \left|\overline{A}^{\alpha} - \overline{B}^{\beta}\right|^{2} &= \left|\prod_{i=1}^{M} \overline{A}_{i}^{\alpha_{i}} - \prod_{j=1}^{N} \overline{B}_{j}^{\beta_{j}}\right|^{2} \\ &= \left|\prod_{i=1}^{M} \left(A_{i,\infty}^{\alpha_{i}} (1+\mu_{i})^{\alpha_{i}} + \widehat{Q}_{i} \|\delta_{i}\|\right) - \prod_{j=1}^{N} \left(B_{j,\infty}^{\beta_{j}} (1+\zeta_{j})^{\beta_{j}} + \widehat{R}_{j} \|\eta_{j}\|\right)\right|^{2} \\ &= \left|A_{\infty}^{\alpha} \prod_{i=1}^{M} (1+\mu_{i})^{\alpha_{i}} - B_{\infty}^{\beta} \prod_{j=1}^{N} (1+\zeta_{j})^{\beta_{j}} + \Theta(\widehat{Q}_{i},\widehat{R}_{j}) \left(\sum_{i=1}^{M} \|\delta_{i}\| + \sum_{j=1}^{N} \|\eta_{j}\|\right)\right|^{2} \end{aligned}$$

with $\Theta(\hat{Q}_i, \hat{R}_j)$ is also uniformly bounded. Thus, by using $(x+y)^2 \ge \frac{1}{2}x^2 - y^2$ and $A_{\infty}^{\alpha} = \sqrt{a_{\infty}^{\alpha}} = \sqrt{b_{\infty}^{\beta}} = B_{\infty}^{\beta}$ and the Cauchy-Schwarz inequality,

$$\left|\overline{A}^{\alpha} - \overline{B}^{\beta}\right|^{2} \ge \frac{1}{2} A_{\infty}^{\alpha} \left|\prod_{i=1}^{M} (1+\mu_{i})^{\alpha_{i}} - \prod_{j=1}^{N} (1+\zeta_{j})^{\beta_{j}}\right|^{2} - |\Theta|^{2} (M+N)^{2} \left(\sum_{i=1}^{M} \|\delta_{i}\|^{2} + \sum_{j=1}^{N} \|\eta_{j}\|^{2}\right).$$
(53)

Hence, for any $\delta \in (0, 1)$ holds

$$\begin{split} \sum_{i=1}^{M} \|\delta_{i}\|^{2} + \sum_{j=1}^{N} \|\eta_{j}\|^{2} + \left|\overline{A}^{\alpha} - \overline{B}^{\beta}\right|^{2} \\ &\geq \sum_{i=1}^{M} \|\delta_{i}\|^{2} + \sum_{j=1}^{N} \|\eta_{j}\|^{2} \\ &+ \delta \left(\frac{1}{2}A_{\infty}^{\alpha} \left|\prod_{i=1}^{M} (1+\mu_{i})^{\alpha_{i}} - \prod_{j=1}^{N} (1+\zeta_{j})^{\beta_{j}}\right|^{2} - |\Theta|^{2} (M+N)^{2} \left(\sum_{i=1}^{M} \|\delta_{i}\|^{2} + \sum_{j=1}^{N} \|\eta_{j}\|^{2}\right)\right) \\ &\geq \frac{\delta}{2}A_{\infty}^{\alpha} \left|\prod_{i=1}^{M} (1+\mu_{i})^{\alpha_{i}} - \prod_{j=1}^{N} (1+\zeta_{j})^{\beta_{j}}\right|^{2} \end{split}$$

by choosing δ small enough such that $1 \ge \delta |\Theta|^2 (M+N)^2$ since Θ is uniformly bounded above. This leads in combination with (49) to a lower bound of the left hand side of (48)

LHS of (48)
$$\geq C_1 \frac{\delta}{2} A_{\infty}^{\alpha} \left| \prod_{i=1}^{M} (1+\mu_i)^{\alpha_i} - \prod_{j=1}^{N} (1+\zeta_j)^{\beta_j} \right|^2.$$
 (54)

From (51) and (54), it is sufficient to prove

$$\left|\prod_{i=1}^{M} (1+\mu_i)^{\alpha_i} - \prod_{j=1}^{N} (1+\zeta_j)^{\beta_j}\right|^2 \ge C_2 \left(\sum_{i=1}^{M} \mu_i^2 + \sum_{j=1}^{N} \zeta_j^2\right).$$
(55)

In order to do so, we note that the conservation laws

$$\beta_j \overline{a}_i + \alpha_i \overline{b}_j = \beta_j a_{i,\infty} + \alpha_i b_{j,\infty}$$

rewritten in terms of the ansatz (50), i.e.

$$\beta_j A_{i,\infty}^2(\mu_i^2 + 2\mu_i) + \alpha_i B_{j,\infty}^2(\zeta_j^2 + 2\zeta_j) = 0.$$

imply $\mu_i \zeta_j \leq 0$ thanks to $\mu_i, \zeta_j \geq -1$ for all i, j. Without loss of generality, we assume $\mu_i \geq 0$ and $\zeta_j \leq 0$ for all i, j. Then, for any $1 \leq i_0 \leq M$ and $1 \leq j_0 \leq N$,

$$\left| \prod_{i=1}^{M} (1+\mu_i)^{\alpha_i} - \prod_{j=1}^{N} (1+\zeta_j)^{\beta_j} \right| \ge \prod_{i=1}^{M} (1+\mu_i)^{\alpha_i} - \prod_{j=1}^{N} (1+\zeta_j)^{\beta_j} \ge (1+\mu_{i_0})^{\alpha_{i_0}} - (1+\zeta_{j_0})^{\beta_{j_0}} \ge (1+\mu_{i_0}) - (1+\zeta_{j_0}) \ge \mu_{i_0} - \zeta_{j_0} \ge 0.$$

Thus

$$\left|\prod_{i=1}^{M} (1+\mu_i)^{\alpha_i} - \prod_{j=1}^{N} (1+\zeta_j)^{\beta_j}\right|^2 \ge (\mu_{i_0} - \zeta_{j_0})^2 = \mu_{i_0}^2 - 2\mu_{i_0}\zeta_{j_0} + \zeta_{j_0}^2 \ge \mu_{i_0}^2 + \zeta_{j_0}^2.$$

Since $1 \le i_0 \le M$ and $1 \le j_0 \le N$ are arbitrary, we finally obtain (55) with $C_2 = 1/\max\{M; N\}$.

Lemma 4.2. Let a_i, b_j be functions defined in Lemma 4.1. Then, there exists a constant C such that

$$\sum_{i=1}^{M} \|\delta_i\|^2 + \sum_{j=1}^{N} \|\eta_j\|^2 + \|A^{\alpha} - B^{\beta}\|^2 \ge C \left|\overline{A}^{\alpha} - \overline{B}^{\beta}\right|^2.$$

Proof. Fix a constant L > 0. Denote by

$$S = \{x \in \Omega : |\delta_i(x)| \le L, |\eta_j(x)| \le L \text{ for all } i = 1, \dots, M, \ j = 1, \dots, N\} \text{ and } S^{\perp} = \Omega \setminus S^{\perp}$$

Recalling $\overline{A_i} \leq \sqrt{\overline{A_i^2}} \leq M_0$ and $\overline{B_j} \leq \sqrt{\overline{B_j^2}} \leq M_0$, we use Taylor expansion to estimate

$$|A^{\alpha} - B^{\beta}||^{2} \geq \int_{S} \left| \prod_{i=1}^{M} (\overline{A}_{i} + \delta_{i}(x))^{\alpha_{i}} - \prod_{j=1}^{N} (\overline{B}_{j} + \eta_{j}(x))^{\beta_{j}} \right|^{2} dx$$

$$\geq \frac{1}{2} \left| \overline{A}^{\alpha} - \overline{B}^{\beta} \right|^{2} |S| - \widetilde{R}(\overline{A}_{i}, \overline{B}_{j}, |\delta_{i}|, |\eta_{j}|) \left(\sum_{i=1}^{M} \|\delta_{i}\|^{2} + \sum_{j=1}^{N} \|\eta_{j}\|^{2} \right)$$
(56)

where $|\tilde{R}| \leq C(M_0, L)$ due to the boundedness of δ_i and η_j in S. In S^{\perp} , we have

$$\sum_{i=1}^{M} \|\delta_i\|^2 + \sum_{j=1}^{N} \|\eta_j\|^2 \ge \int_{S^{\perp}} \left(\sum_{i=1}^{M} |\delta_i(x)|^2 + \sum_{j=1}^{N} |\eta_j(x)|^2 \right) dx \ge L^2 |S^{\perp}|$$

Next, there clearly exists a constant $\Lambda > 0$ such that $\left|\overline{A}^{\alpha} - \overline{B}^{\beta}\right|^2 \leq \Lambda$ since $\overline{A}_i, \overline{B}_j \leq M_0$. Therefore,

$$\sum_{i=1}^{M} \|\delta_i\|^2 + \sum_{j=1}^{N} \|\eta_j\|^2 \ge L^2 |S^{\perp}| \ge \frac{L^2}{\Lambda} \left|\overline{A}^{\alpha} - \overline{B}^{\beta}\right|^2 |S^{\perp}|.$$
(57)

Combining (56) and (57) we find for any $\theta_1, \theta_2 \in (0, 1)$

$$\begin{split} \sum_{i=1}^{M} \|\delta_i\|^2 + \sum_{j=1}^{N} \|\eta_j\|^2 + \|A^{\alpha} - B^{\beta}\|^2 \ge \theta_1 \frac{L^2}{\Lambda} \left|\overline{A}^{\alpha} - \overline{B}^{\beta}\right|^2 |S^{\perp}| + (1 - \theta_1) \left(\sum_{i=1}^{M} \|\delta_i\|^2 + \sum_{j=1}^{N} \|\eta_j\|^2\right) \\ &+ \theta_2 \frac{1}{2} \left|\overline{A}^{\alpha} - \overline{B}^{\beta}\right|^2 |S| - \theta_2 |\widetilde{R}| \left(\sum_{i=1}^{M} \|\delta_i\|^2 + \sum_{j=1}^{N} \|\eta_j\|^2\right) \\ &\ge \min\left\{\theta_1 \frac{L^2}{\Lambda}; \theta_2 \frac{1}{2}\right\} \left|\overline{A}^{\alpha} - \overline{B}^{\beta}\right|^2 (|S| + |S^{\perp}|) \\ &= \min\left\{\theta_1 \frac{L^2}{\Lambda}; \theta_2 \frac{1}{2}\right\} \left|\overline{A}^{\alpha} - \overline{B}^{\beta}\right|^2 \end{split}$$

by choosing θ_1, θ_2 small enough such that $1 - \theta_1 - \theta_2 |\tilde{R}| \ge 0$ and using $|S| + |S^{\perp}| = |\Omega| = 1$. The proof of Lemma 4.2 is hence complete.

5. Proof Theorem 1.1: existence of global weak solution to (S)

In this section, we give a proof Theorem 1.1 about the global existence of weak solutions to (S) under the conditions (G)-(M)-(P). Consider the approximating system

$$\partial_t u_{i,\varepsilon} - d_i \Delta(u_{i,\varepsilon}^{m_i}) = f_{i,\varepsilon}(u_\varepsilon) := \frac{f_i(u_\varepsilon)}{1 + \varepsilon \sum_{i=1}^S |f_i(u_\varepsilon)|}, \quad \nabla(u_{i,\varepsilon}^{m_i}) \cdot \overrightarrow{n} = 0, \quad u_{i,\varepsilon}(x,0) = u_{i,0,\varepsilon}(x) \tag{58}$$

where $u_{\varepsilon} = (u_{1,\varepsilon}, \ldots, u_{S,\varepsilon})$ and the sequence of approximating nonnegative initial data $u_{i,0,\varepsilon} \in L^{\infty}(\Omega)$ converges to $u_{i,0}$ in $L^2(\Omega)$. By the construction of the approximative system, it directly follows that the nonlinearities $f_{i,\varepsilon}$ still satisfy the conditions (M) and (P). Moreover, for $\varepsilon > 0$

$$|f_{i,\varepsilon}(u_{\varepsilon})| \leq \frac{|f_i(u_{\varepsilon})|}{1 + \varepsilon \sum_{i=1}^S |f_i(u_{\varepsilon})|} \leq \frac{1}{\varepsilon} \quad \text{for all } u_{\varepsilon} \in \mathbb{R}^S.$$

Hence, by a classical result for the porous medium equation with L^{∞} data, there exists a strong nonnegative solution $u_{\varepsilon} = (u_{i,\varepsilon})_{i=1...S}$ (see e.g. [Vaz07, Section 8]) in the sense that

$$u_{i,\varepsilon}^{m_i} \in L^2_{loc}(0, +\infty; H^1(\Omega)), \quad \partial_t u_{i,\varepsilon} = d_i \Delta(u_{i,\varepsilon}^{m_i}) + f_{i,\varepsilon}(u_\varepsilon) \in L^1_{loc}(0, +\infty; L^1(\Omega)),$$

 $u_{i,\varepsilon} \in C([0,T); L^1(\Omega))$ and $u_{i,\varepsilon}(0) = u_{i,0,\varepsilon}$,

and the equation for $u_{i,\varepsilon}$ holds a.e. in Q_T for any T > 0. Therefore, it follows immediately that

$$-\int_{\Omega} u_{i,0,\varepsilon}\psi(0)dx - \int_{0}^{T}\int_{\Omega} (\partial_{t}\psi u_{i,\varepsilon} + u_{i,\varepsilon}^{m_{i}}\Delta\psi)dxdt = \int_{0}^{T}\int_{\Omega} f_{i,\varepsilon}(u_{\varepsilon})\psi dxdt$$
(59)

for any test function $\psi \in C^{2,1}(\overline{\Omega} \times [0,T])$ with $\psi(T) = 0$ and $\nabla \psi \cdot \overrightarrow{n} = 0$ on $\partial \Omega \times (0,T)$.

In order to pass to the limit as $\varepsilon \to 0$ in the weak formula (59), we use the following uniform a-priori estimates, which are a consequence of a duality argument in the spirit of e.g. [Pie10] and references therein.

Lemma 5.1 (Duality estimates and uniform a-priori estimates for the approximating solutions, cf. [LP17]). Let $u_{\varepsilon} = (u_{1,\varepsilon}, \ldots, u_{S,\varepsilon})$ be the nonnegative solutions to the approximating system (58). Then,

$$\|u_{i,\varepsilon}\|_{L^{m_i+1}(Q_T)} \le C \quad for \ all \quad T > 0 \quad and \quad i = 1, \dots, S,$$

where the ε -independent constant C only depends on the L^2 -norm of the initial data, the positive constants λ_i of assumption (M), the positive diffusion coefficients d_i and the domain Ω . Moreover, we have

$$||f_{i,\varepsilon}(u_{\varepsilon})||_{L^{1+\delta}(Q_T)} \le C$$

for some $\delta > 0$, where the constant C depends only on the L^2 -norm of $u_{i,0,\varepsilon}$, the positive constants λ_i of assumption (M), the diffusion coefficients d_i , the exponents m_i and the domain Ω .

Proof. The proof follows [LP17] with straightforward changes due to the considered Neumann (instead of Dirichlet) boundary conditions. By setting

$$Z = \sum_{i=1}^{S} \lambda_i u_{i,\varepsilon} \quad \text{and} \quad W = \sum_{i=1}^{S} d_i \lambda_i u_{i,\varepsilon}^{m_i}$$

and by summing up the equations of systems (S), the mass dissipation property (M) implies

 $\partial_t Z - \Delta W \leq 0 \quad \text{ and } \quad \nabla W \cdot \overrightarrow{n} = 0.$

Then, integration over (0, t) and multiplication with W(t) in $L^2(\Omega)$ (due to the regularity of the approximative solutions) leads after integration over Ω to

$$\int_{\Omega} \left(Z(t) - Z(0) \right) W(t) dx - \int_{\Omega} W(t) \Delta \int_{0}^{t} W(s) ds dx \le 0.$$
⁽⁶⁰⁾

Next, we integrate by parts with homogeneous Neumann boundary conditions the second term on the left hand side and calculate

$$-\int_{\Omega} W(t)\Delta \int_{0}^{t} W(s)ds \ dx = \int_{\Omega} \nabla W(t) \cdot \nabla \int_{0}^{t} W(s)ds \ dx = \frac{1}{2} \frac{d}{dt} \int_{\Omega} |\nabla \int_{0}^{t} W(s)ds|^{2} dx.$$

Therefore, by integrating (60) with respect to t on (0, T), we obtain

$$\int_0^T \int_\Omega Z(t)W(t)dxdt + \frac{1}{2} \int_\Omega |\nabla \int_0^T W(s)ds|^2 dx \le \int_0^T \int_\Omega Z(0)W(t)dxdt.$$
(61)

Moreover, we note that

$$\int_{0}^{T} \int_{\Omega} Z(t) W(t) dx dt = \int_{0}^{T} \int_{\Omega} \left(\sum_{i=1}^{S} \lambda_{i} u_{i} \right) \left(\sum_{i=1}^{S} d_{i} \lambda_{i} u_{i}^{m_{i}} \right) dx dt \ge \sum_{i=1}^{S} d_{i} \lambda_{i}^{2} \|u_{i}\|_{L^{m_{i}+1}(Q_{T})}^{m_{i}+1}$$
(62)

due to the nonnegativity of functions u_i and the constant λ_i . To estimate the right hand side of (61) in terms of the L^2 -norm of Z(0), we first notice from $\partial_t Z - \Delta W \leq 0$ that

$$Z(T) - \Delta \int_0^T W dt \le Z(0).$$

Multiplying this inequality with θ_0 in $L^2(\Omega)$, where $\theta_0 \ge 0$ solves $-\Delta \theta_0 = Z(0)$, $\theta_0|_{\partial\Omega} = 0$, and using integration by parts $-\int_{\Omega} \theta_0 \Delta \int_0^T W(t) dt dx = -\int_{\Omega} \Delta \theta_0 \int_0^T W(t) dt dx$, leads to

$$\int_{\Omega} Z(T)\theta_0 dx + \int_{\Omega} \left(Z(0) \int_0^T W(t) dt \right) dx \le \int_{\Omega} Z(0)\theta_0 dx = \|\nabla \theta_0\|^2 \le C \|Z(0)\|^2,$$

which, together with $\int_{\Omega} Z(T) \theta_0 dx \ge 0$, implies

$$\int_{0}^{T} \int_{\Omega} Z(0)W(t)dxdt \le C \|Z(0)\|^{2}.$$
(63)

By inserting (62) and (63) into (61), we obtain

$$\sum_{i=1}^{S} d_i \lambda_i^2 \| u_{i,\varepsilon} \|_{L^{m_i+1}(Q_T)}^{m_i+1} \le C \| Z(0) \|^2$$

which completes the proof of the first a-priori estimate of Lemma 5.1.

Concerning the second uniform a-priori estimate for the nonlinearities, we have

$$|f_{i,\varepsilon}(u_{\varepsilon})| \le |f_i(u_{\varepsilon})| \le C(1+|u_{\varepsilon}|^{\nu}),$$

where C does not depend on ε . By the assumption $m_i > \nu - 1$ and the estimate of $||u_{i,\varepsilon}||_{L^{m_i+1}(Q_T)}$, we obtain $||f_{i,\varepsilon}(u_{\varepsilon})||_{L^{1+\delta}(Q_T)} \leq C$.

The following compactness lemma allows to extract a converging subsequence from the approximating system.

Lemma 5.2. [Bar78] Let $m > (d-2)_+/d$ with $(d-2)_+ = \max\{0, d-2\}$. The mapping $L^1(\Omega) \times L^1(Q_T) \ni (u_0, f) \mapsto u \in L^1(Q_T)$ where $u \in C([0, T]; L^1(\Omega))$ is the weak solution to

$$\partial_t u - \delta \Delta(u^m) = f, \quad \nabla(u^m) \cdot \overrightarrow{n} = 0, \quad u(0) = u_0,$$

with $\delta > 0$, is compact.

Proof of Theorem 1.1. Thanks to the uniform bounds of the nonlinearities in Lemma 5.1 and the compactness Lemma 5.2, there exists a subsequence (not relabeled) $\{u_{i,\varepsilon}\}_{\varepsilon}$ which converges in $L^1(Q_T)$ to limit functions $u_i \in L^1(Q_T)$. From the L^{m_i+1} -bound in Lemma 5.1, it holds in fact that $u_{i,\varepsilon}$ (up to another subsequence) converges strongly to u_i in $L^{m_i}(Q_T)$. For the nonlinearities, we first notice from Lemma 5.1 that the sequence $\{f_{i,\varepsilon}(u_{\varepsilon})\}$ is uniformly integrable. Moreover, for another subsequence $u_{i,\varepsilon} \to u_i$ a.e. in Q_T it follows that

$$f_{i,\varepsilon}(u_{\varepsilon}) \to f_i(u_i)$$
 a.e. in Q_T .

Therefore, we can apply Vitali's Lemma, see e.g. [Sch05, Chapter 16], to obtain $f_{i,\varepsilon}(u_{\varepsilon}) \to f_i(u_i)$ strongly in $L^1(Q_T)$. All this allows to pass to the limit in the weak formulation (59) for any test function $\psi \in C^{2,1}(\overline{\Omega} \times [0,T])$ with $\psi(T) = 0$ and $\nabla \psi \cdot \overrightarrow{n} = 0$ on $\partial \Omega \times (0,T)$. Hence, we get

$$-\int_{\Omega}\psi(0)u_{i,0}dx - \int_{Q_T}(\partial_t\psi u_i + u_i^{m_i}\Delta\psi)dxdt = \int_{Q_T}f_i(u)\psi dxdt.$$

The additional regularity $u_i^{m_i} \in L^1(0,T; W^{1,1}(\Omega))$ follows immediately from [Luk10, Lemma 4.7], where

$$\int_0^T \int_\Omega |\nabla u_i^{m_i}|^\beta dx dt \le C(T, \|u_{i,0}\|_1, \|f_i(u)\|_{L^1(Q_T)}) \quad \text{for all } 1 \le \beta < 1 + \frac{1}{1 + m_i d}.$$

From the above estimate and $f_i(u) \in L^1(Q_T)$, we also have $\partial_t u_i \in L^1(0,T; (W^{1,1}(\Omega))^*)$ which implies in particular $u_i \in C([0,T]; L^1(\Omega))$. This completes the proof of existence of global weak solutions.

Acknowledgements. The second author was supported by the DFG Project CH 955/3-1. This work is partially supported by International Research Training Group IGDK 1754 and NAWI Graz.

References

- [Ba94] A. Barabanova, "On the global existence of solutions of a reaction-diffusion system with exponential nonlinearity", Proc. Am. Math. Soc. 122, (1994) 827–831.
- [Bar78] P. Baras. "Compacité de l'opérateur $f \mapsto u$ solution d'une équation non linéaire $\frac{du}{dt} + Au \ni f$." C. R. Acad. Sci., Sér. A 286 (1978) 1113–1116.

[BR10] S. Benachour, B. Rebiai, "Global classical solutions for reaction-diffusion systems with nonlinearities of exponential growth", J. Evol. Equ. 10 (2010) 511–527.

- [BP00] N. Boudiba and M. Pierre, "Global existence for Coupled Reaction-Diffusion Systems". J. Math. Anal. Appl. 250 (2000) 1–12.
- [CV09] C. Caputo and A. Vasseur, "Global Regularity of Solutions to Systems of Reaction-Diffusion with Sub-Quadratic Growth in Any Dimension". Comm. Partial Differential Equations, 34.10 (2009) 1228–1250.
- [CDF14] J.A. Cañizo, L. Desvillettes, K. Fellner. "Improved duality estimates and applications to reaction-diffusion equations." Comm. Partial Differential Equations 39.6 (2014), 1185–1204.
- [DF85] E. DiBenedetto, A. Friedman. "Hölder estimates for nonlinear degenerate parabolic systems." J. Reine Angew. Math., 357 (1985) 1–22.

- [DF06] L. Desvillettes, K. Fellner. "Exponential decay toward equilibrium via entropy methods for reaction-diffusion equations." J. Math. Anal. Appl. 319.1 (2006) 157–176.
- [DF07] L. Desvillettes, K. Fellner. "Entropy methods for reaction-diffusion systems." Discrete Contin. Dyn. Syst. (suppl). Proceedings of the 6th AIMS International Conference (2007) 304–312.
- [DF08] L. Desvillettes, K. Fellner. "Entropy methods for reaction-diffusion equations: slowly growing a-priori bounds." Rev. Mat. Iberoamericana 24.2 (2008) 407–431.
- [DFT16] L. Desvillettes, K. Fellner, B.Q. Tang. "Trend to equilibrium for reaction-diffusion systems arising from complex balanced chemical reaction networks." SIAM J. Math. Anal. 49 (2017) 2666–2709.
- [DFPV07] L. Desvillettes, K. Fellner, M. Pierre, J. Vovelle, "About Global Existence for Quadratic Systems of Reaction-Diffusion", Adv. Nonlinear Stud. 7 (2007) 491–511.
- [FLT17] K. Fellner, E. Latos, B.Q. Tang. "Well-posedness and exponential equilibration of a volume-surface reactiondiffusion system with nonlinear boundary coupling." to appear in Ann. Inst. H. Poincaré Anal. Non Linéaire. DOI: https://doi.org/10.1016/j.anihpc.2017.07.002
- [FPT17] K. Fellner, W. Prager, B.Q. Tang. "The entropy method for reaction-diffusion systems without detailed balance: first order chemical reaction networks.", Kinet. Relat. Models. 10 (4) (2017) 1055–1087.
- [FT17a] K. Fellner, B.Q. Tang. "Explicit exponential convergence to equilibrium for nonlinear reaction-diffusion systems with detailed balance condition." Nonlinear Anal. 159 (2017) 145–180.
- [FT17] K. Fellner, B.Q. Tang. "Convergence to equilibrium for renormalised solutions to nonlinear chemical reaction-diffusion systems", arXiv:1708.01427.
- [GM13] G. Grillo, M. Muratori. "Sharp short and long time L^{∞} bounds for solutions to porous media equations with homogeneous Neumann boundary conditions." J. Differential Equations, 254 (2013) 2261–2288.
- [HLV98] M. A. Herrero, A. A. Lacey and J. L.Velázquez, "Global Existence for Reaction-Diffusion Systems Modelling Ignition", Arch. Rational Mech. Anal. 142 (1998) 219–251.
- [HMP87] S.L. Hollis, R.H. Martin, M. Pierre. "Global existence and boundedness in reaction diffusion systems." SIAM J. Math. Anal. 18(3) (1987) 744–761.
- [KK00] J.I. Kanel, M. Kirane, "Global solutions of reaction-diffusion systems with a balance law and nonlinearities of exponential growth", J. Differential Equations 165 (2000) 24–41.
- [MP91] R.H. Martin and M. Pierre, "Nonlinear reaction-diffusion systems in Nonlinear Equations in the Applied Sciences", W.F. Ames and C. Rogers ed., Math. Sci. Eng. 185, Acad. Press, New York 1991.
- [Mas83] K. Masuda, "On the global existence and asymptotic behavior of reaction-diffusion equations", Hokkaido Math. J. 12 (1983) 360–370.
- [Laa11] E.-H. Laamri, "Global existence of classical solutions for a class of reaction-diffusion systems", Acta Appl. Math. 115(2) (2011) 153–165.
- [LP17] E.H. Laamri, M. Pierre. "Global existence for reaction-diffusion systems with nonlinear diffusion and control of mass." Ann. Inst. H. Poincaré Anal. Non Linéaire. 34.3 (2017) 571–591.
- [Leu09] A.W. Leung. "Nonlinear Systems of Partial Differential Equations: Applications to Life and Physical Sciences." World Scientific, 2009.
- [Luk10] T. Lukkari. "The porous medium equation with measure data." J. Evol. Equ. 10 (2010) 711–729.
- [Mie] A. Mielke. "Uniform exponential decay for reaction-diffusion systems with complex-balanced mass-action kinetics." to appear in Springer Proceedings in Mathematics & Statistics.
- [MHM15] A. Mielke, J. Haskovec, P. A. Markowich, "On uniform decay of the entropy for reaction-diffusion systems", J. Dynam. Differential Equations, 27 (2015) 897–928.
- [MM17] A. Mielke, M. Mittnenzweig. "Convergence to equilibrium in energy-reaction-diffusion systems using vector-valued functional inequalities." WIAS Preprint No. 2349, (2016).
- [Mor89] J. Morgan. "Global existence for semilinear parabolic systems." SIAM J. Math. Anal. 20(5) (1989) 1128-1144.
- [Pie10] M. Pierre. "Global existence in reaction-diffusion systems with control of mass: a survey." Milan J. Math. 78.2 (2010) 417–455.
- [Pie03] M. Pierre, "Weak solutions and supersolutions in L¹ for reaction-diffusion systems", J. Evol. Equ. 3 (2003) 153–168.
- [PR16] M. Pierre, G. Rolland. "Global existence for a class of quadratic reaction-diffusion systems with nonlinear diffusions and L^1 initial data." Nonlinear Anal. 138 (2016) 369–387.
- [PS97] M. Pierre and D. Schmitt, "Blow up in reaction-diffusion systems with dissipation of mass", SIAM J. Math. Anal. 28, (1997) 259–269.
- [PSZ16] M. Pierre, T. Suzuki and R. Zou. "Asymptotic behavior of solutions to chemical reaction-diffusion systems." J. Math. Anal. Appl., 450.1 (2017) 152–168.
- [Sch05] R.L. Schilling. "Measures, Integrals and Martingales." Cambridge University Press, 2005.
- [Smo94] J. Smoller. "Shock Waves and Reaction-Diffusion Equations." Springer, 1994.
- [Vaz07] J. Vázquez. "The porous medium equation: Mathematical Theory", Oxford Science Publications, 2007.

KLEMENS FELLNER

INSTITUTE OF MATHEMATICS AND SCIENTIFIC COMPUTING, UNIVERSITY OF GRAZ, HEINRICHSTRASSE 36, 8010 GRAZ, AUSTRIA E-mail address: klemens.fellner@uni-graz.at

Evangelos Latos

UNIVERSITY OF MANNHEIM, D-68131 MANNHEIM, GERMANY

E-mail address: evangelos.latos@math.uni-mannheim.de

Bao Quoc Tang Institute of Mathematics and Scientific Computing, University of Graz, Heinrichstrasse 36, 8010 Graz, Austria E-mail address: quoc.tang@uni-graz.at