INVASION MOVING BOUNDARY PROBLEM FOR A BIOFILM REACTOR MODEL

B. D'ACUNTO, L. FRUNZO, V. LUONGO, AND M. R. MATTEI

ABSTRACT. The work presents the analysis of the free boundary value problem related to the invasion model of new species in biofilm reactors. In the framework of continuum approach to mathematical modelling of biofilm growth, the problem consists of a system of nonlinear hyperbolic partial differential equations governing the microbial species growth and a system of semi-linear elliptic partial differential equations describing the substrate trends. The model is completed with a system of elliptic partial differential equations governing the diffusion and reaction of planktonic cells, which are able to switch their mode of growth from planktonic to sessile when specific environmental conditions are found. Two systems of nonlinear differential equations for the substrate and planktonic cells mass balance within the bulk liquid are also considered. The free boundary evolution is governed by a differential equation that accounts for detachment. The qualitative analysis is performed and a uniqueness and existence result is discussed. Furthermore, two special models of biological and engineering interest are discussed numerically. The invasion of Anammox bacteria in a constituted biofilm inhabiting the deammonification units of the wastewater treatment plants is simulated. Numerical simulations are run to evaluate the influence of the colonization process on biofilm structure and activity.

1. Introduction

The term biofilm is used nowadays to indicate the prevailing form of microbial lifestyle, which consists of dynamic complex microbial structures composed of various prokaryotic cells and other microorganisms, forming on solid or liquid surfaces and encased in a self-produced protective matrix of extracellular polymeric substances (EPS). The roles biofilms exert on both natural and human environments are disparate: they have proven detrimental to human health or undesirable in the open water environment but, on the other hand they can be used beneficially in resource recovery systems as well as water treatment [1]. With specific reference to the last point, biofilm reactors represent the primary means to harness the usefulness of biofilms for pollutant removal from wastewater by means of the synergistic interactions and biochemical transformations characterizing these microbial communities [2]. The biofilm structure results from the interplay of different interactions, such as mass transfer, conversion rates and detachment forces. The main biofilm expansion is due to bacterial growth and to extracellular polymer production. The soluble substrates necessary for bacterial growth are dissolved in the liquid flow and to reach the cells, first they pass through a boundary layer, characterized by a negligible flow over the biofilm/liquid interface, and then through the biofilm matrix. The external fluid flow regulates biofilm growth by establishing the concentration of substrates and products

Date: October 1, 2018.

Key words and phrases. Invasion Model, Biofilm reactor, Hyperbolic free boundary value problem, Numerical simulations, Anammox process.

at the solid-liquid interface and shearing the biofilm surface. Other biological phenomena are found to play significant roles in the establishment of mixed species biofilms, i.e. dispersal, bacteriophage, quorum sensing [3].

Among these phenomena, there is a growing interest in the study of microbial invasion and colonization of pre-existing biofilms as it can determine biofilm landscape and contribute to rapid alterations in biofilm populations. Recent advances in microbial ecology have identified motility as one of the main mediator of such process. Indeed, once a motile bacteria, supplied by the liquid phase or the biofilm itself (as a consequence of dispersal phenomenon), has successfully infiltrated the biofilm matrix, it can invade a resident community and establish where the environmental conditions are optimal for its growth. An accurate modeling of such a system has to take all of these factors into account. In a recent contribution [4], the authors have introduced a multispecies biofilm model which explicitly takes into account the invasion phenomenon pursued by planktonic cells. The core of the model lies on the introduction of new state variables which represent the concentrations of planktonic colonizing cells within the biofilm. These cells are supposed to be characterized by a diffusive movement within the biofilm and to be able to give up the ability to move in order to settle down in specific environmental niches.

In this work, we introduce the free boundary value problem for the invasion phenomenon in biofilm reactors which takes into account the dynamics of the bulk liquid phase in terms of both substrates and planktonic cells. The mathematical problem consists of a system of hyperbolic partial differential equations governing the biofilm growth, a system of elliptic partial differential equations for substrate dynamics within the biofilm and a system of elliptic partial differential equations, regulating the diffusion and reaction of planktonic cells. Mass balance equations for the dissolved substrates and planktonic cells within the bulk liquid phase of the biofilm reactor have been taken into account as well. The free boundary evolution is governed by a nonlinear ordinary differential equation.

The qualitative analysis of such a complex system is not an easy task as outlined in [5]. Due to the high non-linearity of the problem, the fixed point theorem seems the natural tool to be used for the existence and uniqueness of the solutions. However, we are considering a moving boundary problem where the domain is not fixed. To overcome this issue, we follow the methodology used in [6] for the analysis of the biofilm reactor model and in [4; 7] for the modeling of the planktonic cells dynamics both within the biofilm and the bulk phase. In particular, we use the method of characteristics to convert the differential problem to an integral one where the unknown functions are defined on a fixed domain and the existence and uniqueness of the solutions are proved in the class of continuous functions.

In addition, the work is completed with some numerical applications related to a real engineering/biological case which examines the invasion of specific microbial species in a constituted biofilm. More precisely, the case study reproduces the invasion of Anammox bacteria within a multispecies biofilm devoted to the concurrent oxidization of ammonium nitrogen and organic carbon occurring in the biological units of the wastewater treatment plants. Traditionally, ammonium oxidation leads to the formation of residual nitrogen compounds that need to be further removed by means of other treatment phases. The establishment of a biofilm community constituted by Anammox bacteria and Aerobic ammonium oxidizers may lead instead to the complete conversion of ammonium nitrogen to nitrogen gas within a single treatment unit. The establishment of this syntrophy is catalyzed by the formation of an anoxic zone, where the Anammox bacteria can effectively proliferate. The invasion model has been adopted to illustrate the trends related to the

establishment of such a multispecies community and to assess the effect of specific operational conditions on the biofilm colonization by Anammox bacteria. For all the cases analyzed real data from existing literature are used to feed numerical simulations, which produce results in nice agreement with experimental findings.

The paper is organized as follows. In Section 2 the invasion moving boundary problem for a biofilm reactor model is introduced: assumptions, boundary and initial conditions are discussed. Section 3 introduces the Volterra integral equations. Section 4 describes the experimental case to which the model is applied and presents the numerical results. Finally, in Section 5 we present the conclusions and the future recommendations of the work.

2. Invasion boundary problem for biofilm reactors

We analyze the free boundary value problem related to the invasion problem in biofilm reactors. In this model we consider the biofilm as constituted by various particulate components (i.e. bacteria, EPS, etc.) growing in a liquid environment, and planktonic cells belonging to various microbial species and able to move within the biofilm and the bulk liquid as well. The biofilm expansion depends on growth limiting nutrients which are dissolved in the liquid region or produced within the biofilm itself. The planktonic cells can diffuse from the bulk liquid to the biofilm, invade, and switch their mode of growth from suspended to sessile when appropriate environmental conditions are found.

The model is formulated for the variables concentration of microbial species in sessile form X_i , the concentration of planktonic invading cells Ψ_i , the concentration of dissolved substrates S_j , all expressed as functions of time t and z which denotes the spatial coordinate. The substratum is assumed to be placed at z=0. The one-dimensional form of the model writes

(2.1)
$$\frac{\partial X_i}{\partial t} + \frac{\partial}{\partial z}(uX_i) = \rho_i r_{M,i}(z, t, \mathbf{X}, \mathbf{S}) + \rho_i r_i(z, t, \mathbf{S}, \mathbf{\Psi}), \ i = 1, ..., n,$$

where ρ_i denotes the constant density of species i, u(z,t) the biomass velocity at which the microbial mass is displaced with respect to the film-support, and $\mathbf{X} = (\mathbf{X_1}, ..., \mathbf{X_n})$, $\mathbf{S} = (\mathbf{S_1}, ..., \mathbf{S_m})$, $\mathbf{\Psi} = (\Psi_1, ..., \Psi_n)$. The reaction terms $r_{M,i}$ describe the growth of sessile cells, which is controlled by the local availability of nutrients and usually modelled as standard Monod kinetics, and natural cell death. In most biological processes the function $r_{M,i}$ depends on z,t only trough the functions \mathbf{X} and \mathbf{S} . The explicit dependence has been considered mainly for mathematical generality. The variable t is positive and $0 \le z \le L(t)$, where L(t) denotes the biofilm thickness at time t. Equation (2.1) without the term r_i was first derived in [8] by mass balance principle. The initial conditions for (2.1) are provided by the initial concentrations $\varphi_i(z)$ of biofilm particulate components

$$(2.2) X_i(z,0) = \varphi_i(z), i = 1, ..., n, 0 \le z \le L(0).$$

The initial concentrations of the invading microbial species are set to zero. The equation in the form (2.1) was presented in [4]. The terms r_i represent the growth rates of the microbial species X_i due to the invasion process which induces the switch of planktonic cells to a sessile mode of growth. This phenotypic alteration is catalyzed by the formation within the biofilm matrix of specific environmental niches. The explicit dependence on z, t has been introduced only for mathematical generality.

Similarly to traditional continuum models of biofilm growth, equations (2.1) can be rewritten in terms of volume fractions $f_i = X_i/\rho_i$, which indicate the fraction of space at a particular location that is occupied by species i,

$$\frac{\partial f_i}{\partial t} + \frac{\partial}{\partial z}(uf_i) = r_{M,i} + r_i.$$

Of course, the sum of all volume fractions at each location and time must always sum to one $\sum_{i=1}^{n} f_i = 1$.

From the equations above it follows immediately that the function u(z,t) satisfies the following problem

(2.3)
$$\frac{\partial u}{\partial z} = \sum_{i=1}^{n} \left(r_{M,i}(z, t, \mathbf{X}, \mathbf{S}) + r_i(z, t, \mathbf{S}, \mathbf{\Psi}) \right), \quad 0 < z \leq L(t), \quad u(0, t) = 0,$$

where the initial condition u(0,t) = 0 comes from no flux condition on substratum.

The function L(t) is solution of the following problem

$$\dot{L}(t) = u(L(t), t) - \sigma_d(L(t)), \ L(0) = L_0.$$

Therefore, it is apparent that the evolution of the free boundary depends on the displacement velocity of microbial biomass u and detachment flux σ_d as well. Equation in (2.4) comes from global mass conservation principle.

The diffusion of planktonic cells within the biofilm matrix is governed by the following diffusion-reaction equations

(2.5)
$$\frac{\partial \Psi_i}{\partial t} - \frac{\partial}{\partial z} \left(D_{M,i} \frac{\partial \Psi_i}{\partial z} \right) = r_{\Psi,i}(z, t, \mathbf{S}, \boldsymbol{\Psi}), \ i = 1, ..., n, \ 0 < z < L(t),$$

where the reaction terms $r_{\Psi,i}$ represent loss terms for the invading species. Homogeneous Neumann conditions are adopted on the substratum at z=0 due to a no-flux conditions and Dirichlet boundary conditions are prescribed on the free boundary z=L(t)

(2.6)
$$\frac{\partial \Psi_i}{\partial z}(0,t) = 0, \ \Psi_i(L(t),t) = \psi_i^*(t), \ i = 1,...,n.$$

The initial conditions are set to zero if it is assumed that the invasion process starts at t = 0, but specific functions can also be considered.

The functions $\psi_i^*(t)$ denote the concentrations of planktonic cells within the bulk liquid and are governed by the following initial value problem for ordinary differential equations

(2.7)
$$V\dot{\psi}_{i}^{*} = -AD_{M,i}\frac{\partial\Psi_{i}}{\partial z}(L(t),t) + Q(\psi_{i}^{in} - \psi_{i}^{*}(t)), \ \psi_{i}^{*}(0) = \psi_{i}^{in}, \ i = 1,...,n.$$

Equations (2.7) come from a mass balance within the bulk liquid and account for the inlet and outlet flux to the biofilm reactor and the exchange fluxes to or from the biofilm as well. The bulk liquid is modelled as a completely mixed compartment of volume V and continuously fed and withdrawn at the same flow rate Q. The initial concentrations of planktonic cells within the bulk liquid are provided by the inlet concentrations ψ_i^{in} .

The substrate diffusion within the biofilm is governed by the following reaction-diffusion equations

(2.8)
$$\frac{\partial S_j}{\partial t} - \frac{\partial}{\partial z} \left(D_j \frac{\partial S_j}{\partial z} \right) = r_{S,j}(z, t, \mathbf{X}, \mathbf{S}), \ j = 1, ..., m, \ 0 < z < L(t),$$

where the terms $r_{S,j}$ represent the substrate production or consumption rates due to microbial metabolism and D_j denotes the diffusion coefficient of substrate j within the biofilm. As to the boundary conditions it is assumed that

(2.9)
$$\frac{\partial S_j}{\partial z}(0,t) = 0, \ h \frac{D_j}{D_j^*} \frac{\partial S_j}{\partial z}(L(t),t) + S_j(L(t),t) = S_j^*(t), \ j = 1, ..., m.$$

The first conditions is a no-flux boundary condition on the substratum placed at z=0. The second condition derives from the following reasonings. According to [12] we assume that at a certain distance from the substratum H(t) = L(t) + h, with h being a given positive constant, the substrate concentration $S_j(H(t),t)$ is the same as the bulk liquid concentration denoted by $S_j^*(t)$. This dissolved substrate diffuses from the bulk liquid to the biofilm $0 \le z \le L(t)$ where it is consumed according to equations (2.8). No biochemical reactions are supposed to occur for $L(t) \le z \le H(t)$ which leads to consider homogeneous parabolic equations for $S_j(z,t)$. Solving at steady-state leads to (2.9), where D_j^* represents the diffusion coefficient of substrate j within the bulk liquid. Note that, condition (2.9) reduces to $S_j(L(t),t) = S_j^*(t)$ for h = 0.

The functions $S_j^*(t)$ are governed by the following initial value problem for ordinary differential equations

$$(2.10) V\dot{S}_{j}^{*} = -AD_{j}\frac{\partial S_{j}}{\partial z}(L(t), t) + Q(S_{j}^{in} - S_{j}^{*}(t)), \ j = 1, ..., m, \ S_{j}^{*}(0) = S_{j}^{in}.$$

Equations above are derived from mass balance on the bulk liquid taking into account the inlet and outlet flux from the reactor and the exchange flux between the biofilm and the bulk liquid. The initial conditions for S_j^* are the same as the inlet concentrations.

Finally, due to the slow evolution of the system [6], $S_j(z,t)$ profiles can be considered to evolve quasi-statically and thus equations (2.8) are rewritten as

(2.11)
$$-D_j \frac{\partial^2 S_j}{\partial z^2} = r_{S,j}(z, \mathbf{X}, \mathbf{S}), \ j = 1, ..., m, \ 0 < z < L(t),$$

with boundary conditions (2.9). In addition, same arguments as before lead to replace equations (2.5) with the following

(2.12)
$$-D_{M,i} \frac{\partial^2 \Psi_i}{\partial z^2} = r_{\Psi,i}(z, \mathbf{S}, \mathbf{\Psi}), \ i = 1, ..., n, \ 0 < z < L(t),$$

with boundary conditions (2.6).

In conclusion the invasion free boundary problem for biofilm reactor is expressed by equations (2.1)-(2.12). In the next section, following [6; 7], an equivalent integral formulation of the problem will be provided. As it will be apparent at the end of the following section, the integral form of the free boundary problem presents the great advantage that the space variable is defined on a fixed domain whereas in the differential formulation (2.1)-(2.12) the space variable belongs to the moving domain $0 \le z \le L(t)$.

3. Volterra integral equations

The differential problem introduced in the previous section is herein converted to Volterra integral equations by using the method of characteristics. The characteristic-like lines of system (2.1) are defined by

(3.1)
$$\frac{\partial c}{\partial t}(z_0, t) = u(c(z_0, t), t), \ c(z_0, 0) = z_0, \ 0 \le z_0 \le L_0, \ t > 0,$$

Considering (3.1), equations (2.1) are converted to

$$\frac{d}{dt}X_i(c(z_0,t),t) =$$

(3.2)
$$F_i(c(z_0, t), t, \mathbf{X}(c(z_0, t), t), \mathbf{S}(c(z_0, t), t), \mathbf{\Psi}(c(z_0, t), t)), \ 0 \le z_0 \le L_0, \ t > 0,$$
 with

$$F_i = \rho_i r_{M,i}(c(z_0, t), t, \mathbf{X}(c(z_0, t), t), \mathbf{S}(c(z_0, t), t))$$

(3.3)
$$+ \rho_i r_i(c(z_0, t), t, \mathbf{S}(c(z_0, t), t), \boldsymbol{\Psi}(c(z_0, t), t)) - X_i(c(z_0, t), t) \sum_{i=1}^n (r_{M,i} + r_i),$$

and initial conditions

$$(3.4) X_i(c(z_0,0),0) = \varphi_i(z_0), \quad 0 \le z_0 \le L_0.$$

Integrating (3.2) and considering (3.4) yields

$$X_{i}(c(z_{0},t),t) = \int_{0}^{t} F_{i}(c(z_{0},\tau),\tau,\mathbf{X}(c(z_{0},\tau),\tau),\mathbf{S}(c(z_{0},\tau),\tau),\boldsymbol{\Psi}(c(z_{0},\tau),\tau))d\tau$$

$$(3.5) + \varphi_i(z_0), \quad i = 1, ..., n, \ 0 \le z_0 \le L_0, \ t > 0.$$

The following integral equation for $c(z_0,t)$ is derived from (3.1) and (2.3)

$$c(z_0,t) = z_0 + \int_0^t d\tau \int_0^{z_0} \sum_{i=1}^n ((r_{M,i}(c(\zeta_0,\tau),\tau,\mathbf{X}(c(\zeta_0,\tau),\tau),\mathbf{S}(c(\zeta_0,\tau),\tau))$$

$$(3.6) \qquad + r_i(c(\zeta_0,\tau),\tau,\mathbf{S}(c(\zeta_0,\tau),\tau),\boldsymbol{\Psi}(c(\zeta_0,\tau),\tau)))\frac{\partial c}{\partial \zeta_0}(\zeta_0,\tau) \ d\zeta_0, 0 \leqslant z_0 \leqslant L_0, t > 0.$$

From (3.6) it follows easily

$$\frac{\partial c}{\partial z_0}(z_0,t) = 1 + \int_0^t \sum_{i=1}^n ((r_{M,i}(c(z_0,\tau),\tau,\mathbf{X}(c(z_0,\tau),\tau),\mathbf{S}(c(z_0,\tau),\tau)))$$

$$(3.7) + r_i(c(z_0, \tau), \tau, \mathbf{S}(c(z_0, \tau), \tau), \mathbf{\Psi}(c(z_0, \tau), \tau))) \frac{\partial c}{\partial \zeta_0}(z_0, \tau) d\tau, 0 \leqslant z_0 \leqslant L_0, \ t > 0.$$

The integral equations for $S_j(z,t)$ are obtained by integrating (2.11) and considering the boundary conditions (2.9)

$$S_{j}(z,t) = S_{j}^{*}(t) + D_{j}^{-1} \int_{0}^{z} (L-z) r_{S,j}(\zeta, \mathbf{X}(\zeta,t), \mathbf{S}(\zeta,t)) d\zeta$$
$$+ D_{j}^{-1} \int_{z}^{L} (L-\zeta) r_{S,j}(\zeta, \mathbf{X}(\zeta,t), \mathbf{S}(\zeta,t)) d\zeta$$

(3.8)
$$+ \frac{h}{D_i^*} \int_0^L r_{S,j}(\zeta, \mathbf{X}(\zeta, t), \mathbf{S}(\zeta, t)) d\zeta, \ j = 1, ..., m, \ 0 \le z \le L(t), \ t > 0.$$

Similarly, the following integral equations for Ψ_i are obtained

$$\Psi_i(z,t) = \psi_i^*(t) + D_{M,i}^{-1} \int_0^z (L-z) r_i(\zeta, \mathbf{S}(\zeta,t), \Psi(\zeta,t)) d\zeta$$

(3.9)
$$+ D_{M,i}^{-1} \int_{z}^{L} (L - \zeta) r_{i}(\zeta, \mathbf{S}(\zeta, t), \mathbf{\Psi}(\zeta, t)) d\zeta \quad i = 1, ..., n, \quad 0 \le z \le L(t), \ t > 0.$$

From (3.8) it follows

(3.12)

(3.10)
$$\frac{\partial S_j}{\partial z}(L,t) = -D_j^{-1} \int_0^L r_{S,j}(\zeta, \mathbf{X}(\zeta, t), \mathbf{S}(\zeta, t)) d\zeta.$$

Considering (3.10) in (2.10), equation for $S_i^*(t)$ writes

$$\dot{S}_{j}^{*}(t) = (A/V) \int_{0}^{L} r_{S,j}(\zeta,t), \mathbf{X}(\zeta,t), \mathbf{S}(\zeta,t)) d\zeta + (Q/V) (S_{j}^{in} - S_{j}^{*}(t)).$$

Integrating the last equation over time leads to the following integral equation for $S_i^*(t)$

$$S_{j}^{*}(t) = \int_{0}^{t} \exp(-Q(t-\tau)/V) d\tau \int_{0}^{L} (A/V) r_{S,j}(\zeta, \mathbf{X}(\zeta, \tau), \mathbf{S}(\zeta, \tau)) d\zeta$$

$$+ S_{j}^{in}, \quad j = 1, ..., m, \ t > 0.$$
(3.11)

Following the same reasoning, a similar equation is obtained for $\psi_i^*(t)$

$$\psi_i^*(t) = \int_0^t \exp(-Q(t-\tau)/V) d\tau \int_0^L (A/V) r_{\psi,i}(\zeta, \mathbf{S}(\zeta, \tau), \mathbf{\Psi}(\zeta, \tau)) d\zeta$$
$$+ \psi_i^{in}, \quad i = 1, ..., n, \ t > 0.$$

The integral equation for L(t) is obtained from (2.4)

(3.13)
$$L(t) = L_0 + \int_0^t u(L(\tau), \tau) \ d\tau - \int_0^t \sigma_d(L(\tau)) \ d\tau, \ t > 0.$$

Let us note that, as outlined at the end of Section 2, the integral equations above depend on time and the space variable z_0 defined in the fixed domain $0 \le z \le L_0$. This result is essential to prove the existence and uniqueness of solutions. Indeed, following [6; 7], a suitable contractive map can be introduced in the space of continuous functions and the fixed point theorem can be applied. We neglect the calculations since they are a generalization of [6; 7] with small modifications.

4. Anammox invasion model

In the previous sections, we performed the qualitative analysis for the invasion free boundary value problem of a biofilm reactor model. In particular, a result on the existence and uniqueness of solutions was provided. However, it is apparent that when complex biological cases are discussed, only numerical simulations can provide satisfactory predictions. The previous qualitative analysis gives a solid base to calculations. For the numerical solution of the model we use an extension of the numerical method proposed in [9]. The code is implemented in MatLab platform and simulations are run for a set target simulation time T that will be specified later on.

The simulated biofilm system consists of bacterial cells accumulating on a surface surrounded by an aquatic region and reproduces a typical multi-culture and multi-substrate process which establishes in the deammonification units of the wastewater treatment plants. The deammonification process consists in the autotrophic nitrogen removal carried out by two microbial groups, the ammonium oxidizing bacteria AOB (X_1) which oxidize ammonium S_1 partially to nitrite S_2 aerobically and the anaerobic ammonium oxidizing bacteria AMX (X_2) , which subsequently convert the remaining ammonium and the newly formed nitrite into nitrogen gas and nitrate S_3 in trace concentrations. This process is also known as partial nitritation/anammox [10]. In multispecies biofilms, the AOB and AMX compete with other two major microbial groups: the nitrite oxidizing bacteria NOB (X_3) , which oxidize S_2 to S_3 in aerobic conditions and compete with X_1 for oxygen and X_2 for nitrite, and heterotrophic bacteria HB (X_4) . The latter can be further classified in ordinary heterotrophic organisms oxidizing the organic material and denitrifiers reducing nitrate to nitrite and nitrite to dinitrogen gas by consuming organic substrate S_4 . X_4 compete with X_1 and X_3 for oxygen S_5 and with X_2 for nitrite, the limiting substrate of X_2 in most instances. The establishment and proliferation of X_2 in such constituted biofilms strictly depends on the formation of an anoxic zone in the inner parts of the matrix where X_3 cannot grow, due to oxygen limitation.

The mathematical model takes into consideration the dynamics of the five microbial species $X_i(z,t)$, including inert material X_5 which derives from microbial decay, and the five reactive components $S_j(z,t)$ within the biofilm. The corresponding concentrations in the bulk liquid $S_j^*(t)$ are taken into account as well. Planktonic cells have been considered for both X_2 and X_4 species as the model is aimed at simulating the invasion of a constituted biofilm by heterotrophic and Anammox bacteria after the establishment of a favorable environmental niche. Two modelling scenarios have been considered: the case of X_2 as single invading species and the case of X_2 and X_4 invasion and establishment in an autotrophic biofilm. Hereafter, they will be referred as Model 1, considered in Section 4.1, and Model 2 that will be discussed in Section 4.2.

4.1. Model 1 – One invading species. Model 1 considers a single invading species: the anaerobic ammonium oxidizing bacteria AMX (X_2) . The mathematical formalization of the problem is provided below. The microbial species dynamics is governed by equations (2.1) rewritten in terms of f_i for convenience

(4.1)
$$\frac{\partial f_i}{\partial t} + \frac{\partial}{\partial z}(uf_i) = r_{M,i}(z, t, \mathbf{X}, \mathbf{S}) + r_i(z, t, \mathbf{S}, \mathbf{\Psi}), \ i = 1, ..., 5.$$

The following initial volume fractions are associated to equations (4.1)

$$(4.2) f_1(z,0) = 0.65, f_2(z,0) = 0.0, f_3(z,0) = 0.25, f_4(z,0) = 0.1, f_5(z,0) = 0.0.$$

The biofilm is assumed to be initially constituted only by the species X_1 , X_3 , X_4 . The invasion of the species X_2 is simulated. The initial biofilm thickness L_0 is given by

$$(4.3) L_0 = 0.1 mm.$$

A representation of the initial microbial distribution is reported in Fig. 1. The net specific biomass growth rates $r_{M,i}$, i = 1, ..., 4 are expressed as

$$r_{M,i} = (\mu_i(\mathbf{S}) - k_{d,i})f_i,$$

where the function $\mu_i(\mathbf{S})$ denotes the *ith* biomass specific growth rate and depends on the anabolic reactions performed by the *ith* microbial species. It is usually formulated as




FIGURE 1. Initial biofilm configuration for Model 1.

Monod kinetics as detailed below. The term $k_{d,i}$ accounts instead for the forms of biomass loss and energy requirements not associated with growth, including decay, maintenance, endogenous respiration, lysis, predation, death. The net specific growth rates associated to X_i , i = 1, ..., 4 are the following

(4.4)
$$r_{M,1} = (\mu_1(\mathbf{S}) - k_{d,1}) f_1 = \left(\mu_{\max,1} \frac{S_1}{K_{1,1} + S_1} \frac{S_5}{K_{1,5} + S_5} - k_{d,1}\right) f_1,$$

$$(4.5) r_{M,2} = (\mu_2(\mathbf{S}) - k_{d,2})f_2 = \left(\mu_{\max,2} \frac{K_{2,5}}{K_{2,5} + S_5} \frac{S_1}{K_{2,1} + S_1} \frac{S_2}{K_{2,2} + S_2} - k_{d,2}\right) f_2,$$

(4.6)
$$r_{M,3} = (\mu_3(\mathbf{S}) - k_{d,3})f_3 = \left(\mu_{\text{max},3} \frac{S_2}{K_{3,2} + S_2} \frac{S_5}{K_{3,5} + S_5} - k_{d,3}\right) f_3,$$

$$r_{M,4} = (\mu_{4,1}(\mathbf{S}) + \mu_{4,2}(\mathbf{S}) + \mu_{4,3}(\mathbf{S}) - k_{d,4})f_4$$

$$= \left(\mu_{\max,4} \frac{S_4}{K_{4,4} + S_4} \frac{S_5}{K_{4,5} + S_5} + \beta_1 \mu_{\max,4} \frac{K_{4,5}}{K_{4,5} + S_5} \frac{S_4}{K_{4,4} + S_4} \frac{S_3}{K_{4,3} + S_3} \frac{S_3}{S_2 + S_3} \right)$$

$$(4.7) +\beta_2 \mu_{\max,4} \frac{K_{4,5}}{K_{4,5} + S_5} \frac{S_4}{K_{4,4} + S_4} \frac{S_2}{K_{4,2} + S_2} \frac{S_2}{S_3 + S_2} - k_{d,4} f_4,$$

where $\mu_{\max,i}$ denotes the maximum net growth rate for biomass i, $K_{i,j}$ the affinity constant of substrate j for biomass i, β_1 and β_2 the reduction factor for denitrification nitrate to nitrite and nitrite to nitrogen gas respectively.

The autotrophic performance in the deammonification process relies on the activity of X_1 and X_2 and results in the S_1 conversion to dinitrogen gas via S_2 . In aerobic environments, S_1 represents the preferential substrate for X_1 growth (4.4). X_2 proliferate, in turn, on S_1 and S_2 and their metabolic activity is strongly affected by the oxygen concentration, the latter being inhibitory even at low concentrations (4.5). Moreover, they rely on the production of S_2 by X_1 , when that substrate is not provided from the bulk liquid. X_3 oxidize S_2 to S_3 under aerobic conditions and thus they compete with X_2 for S_2 (4.6). X_4 are considered facultative bacteria: they can aerobically oxidize the organic matter ($\mu_{4,1}$) or perform denitrification reactions over S_3 and S_2 ($\mu_{4,2}$ and $\mu_{4,3}$ respectively). Indeed, in presence of S_4 , S_2 and S_3 can be contextually consumed by X_4

according to equation (4.7). In particular, S_3 and S_2 are reduced to dinitrogen gas in a sequential process which first converts S_3 into S_2 , the latter being then reduced to N_2 . In addition, the ratios $S_3/(S_2 + S_3)$ and $S_2/(S_2 + S_3)$ varying between $0 \div 1$, indicate the percentage of biomass growing on nitrate and/or nitrite respectively. Inert has been treated as an additional microbial species whose growth rate depends on the decay of all the active species

$$(4.8) r_{M.5} = k_{d.1}f_1 + k_{d.2}f_2 + k_{d.3}f_3 + k_{d.4}f_4.$$

The specific growth rates r_i induced by the switch of the planktonic cells to the sessile mode of growth are defined as

$$(4.9) r_1 = r_3 = r_4 = r_5 = 0,$$

(4.10)
$$r_2 = k_{col,2} \frac{\Psi_2}{k_{\psi,2} + \Psi_2} \frac{K_{2,5}}{K_{2,5} + S_5} \frac{S_1}{K_{2,1} + S_1} \frac{S_2}{K_{2,2} + S_2}.$$

Note that the growth rate terms r_2 for X_2 is newly introduced as Monod kinetics and indicate that the transition of bacteria from planktonic state ψ_2 into the sessile state X_2 is controlled by the formation of a specific environmental niche which is strictly connected to the local concentration of dissolved substrates. The presence of planktonic species is fundamental for the occurrence of the invasion process, as better specified in the following

Remark 1. Consider the second equation in (2.8) with $r_{M,2}$ given by (4.5) and initial condition $f_2(z,0) = 0$. If it is supposed that $r_2 = 0$, then the mentioned equation admits the unique solution $f_2(z,t) = 0$ and the species X_2 cannot develop.

The diffusion of substrates is governed by

(4.11)
$$\frac{\partial S_j}{\partial t} - D_j \frac{\partial^2 S_j}{\partial z^2} = r_{S,j}(z, t, \mathbf{X}, \mathbf{S}), \ j = 1, ..., 5,$$

with the following initial-boundary conditions

(4.12)
$$S_j(z,0) = 0, \ \frac{\partial S_j}{\partial z}(0,t) = 0, \ j = 1,...,5,$$

(4.13)
$$S_j(L(t),t) = S_j^*(t), \ j = 1,...,4, \ S_5(L(t),t) = \overline{S}_5 = 1.5 \ mgO_2/L.$$

The last condition simulates a continuous aeration of the biofilm reactor, [5].

The net substrate conversion rates account for both the microbial production and consumption (positive and negative terms respectively) and can be formulated from the corresponding microbial growth rates through the specific microbial yield Y_i . They are usually expressed as double-Monod kinetics as presented below.

The ammonium conversion rate $r_{S,1}$ is expressed as

4.14)

$$r_{S,1} = (-\frac{1}{Y_1} - i_{N,B})\mu_1 X_1 + (-\frac{1}{Y_2} - i_{N,B})\mu_2 X_2 - i_{N,B}(\mu_3 X_3 + \mu_{4,1} X_4 + \mu_{4,2} X_4 + \mu_{4,3} X_4),$$

where Y_i denotes the yield for biomass i and $i_{N,B}$ is the nitrogen content in biomass. Ammonium can be directly consumed by AOB and AMX (first and second term in 4.14), and it is usually uptaken by other microbial species for anabolic reactions (third term in 4.14). The nitrite and nitrate conversion rates $r_{S,2}$ and $r_{S,3}$ can be written as

$$(4.15) \\ r_{S,2} = \frac{1}{Y_1} \mu_1 X_1 - (\frac{1}{Y_2} + \frac{1}{1.14}) \mu_2 X_2 - \frac{1}{Y_3} \mu_3 X_3 - (1 - \frac{1}{Y_4}) \frac{1}{1.14} \mu_{4,2} X_4 + (1 - \frac{1}{Y_4}) \frac{1}{1.72} \mu_{4,3} X_4,$$

$$(4.16) r_{S,3} = (\frac{1}{1.14})\mu_2 X_2 + + \frac{1}{Y_3}\mu_3 X_3 + (1 - \frac{1}{Y_4})\frac{1}{1.14}\mu_{4,2} X_4.$$

In aerobic environments, nitrite S_2 are produced through the ammonium conversion catalyzed by AOB and they are further oxidized to nitrate S_3 by NOB (first and third term in 4.15). Obviously, the latter represents a production rate for nitrate (second term in 4.16) Conversely, in anoxic conditions AMX bacteria convert nitrite and ammonium in dinitrogen gas (second term in 4.15), while HB consume both the oxidized nitrogen compounds by reducing nitrate to nitrite (fourth and third term in 4.15 and 4.16 respectively) and by using nitrite as oxygen source (fifth term in 4.15).

The organic carbon conversion rate $r_{S,4}$ is expressed by

(4.17)
$$r_{S,4} = -\frac{1}{Y_4}(\mu_{4,1}X_4 + \mu_{4,2}X_4 + \mu_{4,3}X_4)$$

and indicates the S_4 consumption due to X_4 metabolism in both aerobic and anoxic conditions.

Finally, $r_{S,5}$ describes the oxygen conversion rate within the multispecies biofilm

$$(4.18) r_{S,5} = \left(1 - \frac{3.43}{Y_1}\right)\mu_1 X_1 + \left(1 - \frac{1.14}{Y_3}\right)\mu_3 X_3 + \left(1 - \frac{1}{Y_4}\right)\mu_{4,1} X_4,$$

where the three terms in (4.18) are net consumption rates due to AOB, NOB and HB species using oxygen for their metabolisms.

The functions $S_j^*(t)$ are governed by the following initial value problem for ordinary differential equations

(4.19)
$$V\dot{S}_{j}^{*} = -AD_{j}\frac{\partial S_{j}}{\partial z}(L(t), t) + Q(S_{j}^{in} - S_{j}^{*}(t)), \ j = 1, ..., 4,$$

The initial conditions for S_i^* are the following

$$(4.20) S_1^{in} = 1200 \ mgN/L, \ S_2^{in} = S_3^{in} = 0, \ S_4^{in} = 120 \ mgCOD/L.$$

The inlet concentrations are non-zero only for S_1 and S_4 , reproducing the case of a biofilm reactor fed with a wastewater containing both ammonium nitrogen and organic carbon.

The diffusion and reaction of planktonic cells within the biofilm matrix is governed by the following equations

(4.21)
$$\frac{\partial \Psi_i}{\partial t} - D_{M,i} \frac{\partial^2 \Psi_i}{\partial z^2} = r_{\Psi,i}(z, t, \mathbf{S}, \boldsymbol{\Psi}), \ i = 1, ..., 5,$$

where $D_{M,i}$ denotes the diffusivity coefficient. The conversion rates of planktonic cells due to invasion process are expressed by

(4.22)
$$r_{\psi,i} = -\frac{1}{Y_{\psi,i}} r_i, \ i = 1, ..., 5,$$

Symbol	Definition	Value	Units
Y_1	X_1 yield on S_1	0.150	gCOD/gN
Y_2	X_2 yield on S_1	0.159	gCOD/gN
Y_3	X_3 yield on S_1	0.041	gCOD/gN
Y_4	X_4 yield on S_4	0.63	gCOD/gCOD
$\mu_{\mathrm{max},1}$	Maximum growth rate of X_1	2.05	d^{-1}
$\mu_{\mathrm{max},2}$	Maximum growth rate of X_2	0.08	d^{-1}
$\mu_{\mathrm{max},3}$	Maximum growth rate of X_3	1.45	d^{-1}
$\mu_{\mathrm{max,4}}$	Maximum growth rate of X_4	6.0	d^{-1}
$K_{1,1}$	S_1 affinity constant for X_1	2.4	mgN/L
$K_{1,5}$	S_5 affinity constant for X_1	0.6	mgO_2/L
$K_{2,1}$	S_1 affinity constant for X_2	0.07	mgN/L
$K_{2,2}$	S_2 affinity constant for X_2	0.05	mgN/L
$K_{2,5}$	S_5 inhibiting constant for X_2	0.01	mgO_2/L
$K_{3,2}$	S_2 affinity constant for X_3	5.5	mgN/L
$K_{3,5}$	S_5 affinity constant for X_3	2.2	mgO_2/L
$K_{4,4}$	S_4 affinity constant for X_4	4.0	mgCOD/L
$K_{4,5}$	S_5 affinity/inhibiting constant for X_4	0.2	mgO_2/L
$K_{4,2}$	S_2 affinity constant for X_4	0.5	mgN/L
$K_{4,3}$	S_3 affinity constant for X_4	0.5	mgN/L
$k_{d,1}$	Decay constant for X_1	0.0068	d^{-1}
$k_{d,2}$	Decay constant for X_2	0.0026	d^{-1}
$k_{d,3}$	Decay constant for X_3	0.04	d^{-1}
$k_{d,4}$	Decay constant for X_4	0.06	d^{-1}

Table 1. Kinetic and Stoichiometric Parameters used for Numerical Simulations

with $Y_{\psi,i}$ being the yield of sessile species on planktonic ones. They are assumed proportional to r_i , i.e. described by the same Monod kinetics [4]. The following initial-boundary conditions are associated to equations (4.21)

(4.23)
$$\Psi_i(z,0) = 0, \ \frac{\partial \Psi_i}{\partial z}(0,t) = 0, \Psi_i(L(t),t) = \psi_i^*(t), \ i = 1,..., 5.$$

The functions $\psi_i^*(t)$ satisfy the following initial value problem for ordinary differential equations

$$(4.24) V\dot{\psi}_{i}^{*} = -AD_{M,i}\frac{\partial\Psi_{i}}{\partial z}(L(t),t) + Q(\psi_{i}^{in} - \psi_{i}^{*}(t)), i = 1,...,5,$$

(4.25)
$$\psi_1^{in} = 0, \ \psi_2^{in} = 1.0 \ mgCOD/L, \ \psi_3^{in} = \psi_4^{in} = \psi_5^{in} = 0.$$

Note that for i=1 equation for Ψ_1 is homogeneous because of hypothesis (4.9) and equation for ψ_1^* do not contain the term ψ_1^{in} because of hypothesis (4.25). Therefore, the system of the two equations admits the unique solution $\Psi_1(z,t)=0$, $\psi_1^*(t)=0$. Same result holds for $\Psi_3(z,t)=0$, $\psi_3^*(t)=0$, $\Psi_4(z,t)=0$, $\psi_4^*(t)=0$, $\Psi_5(z,t)=0$, $\psi_5^*(t)=0$.

The biofilm reactor is characterized by the following operational parameters: the flow rate Q is set to 3.15L/d, the surface area available for biofilm attachment and proliferation A is equal to $1m^2$ and the reactor volume is of 3.15L, leading to a hydraulic retention time of 1 day.

The values of the stoichiometric and kinetic parameters used for numerical simulations are adopted from [2] and are reported for convenience in Tables 1-2.

The simulation results for the multispecies biofilm performance when the AMX invasion is considered are reported in Figs. 2-4. After one day of simulation time (Fig. 2-A), it is possible to notice that the microbial distribution into the biofilm is still affected by

Symbol	Definition	Value	Units
$i_{N,B}$	N content of biomass	0.07	gN/gCOD
eta_1	Reduction factor for denitrification $NO_3 - NO_2$	0.8	-
eta_2	Reduction factor for denitrification $NO_2 - N_2$	0.8	-
$k_{col,2}$	Maximum colonization rate of ψ_2	0.0001	d^{-1}
$k_{col,4}$	Maximum colonization rate of ψ_4	0.0001	d^{-1}
$Y_{\psi,2}$	Yield of X_2 on ψ_2	0.001	-
$Y_{\psi,4}$	Yield of X_4 on ψ_4	0.001	-
$k_{\psi,2}$	Kinetic constant for ψ_2	0.000001	mgCOD/L
$k_{\psi,4}$	Kinetic constant for ψ_4	0.000001	mgCOD/L

Table 2. Kinetic and Stoichiometric Parameters used for Numerical Simulations




FIGURE 2. Microbial species distribution of a multispecies biofilm undergoing ψ_2 colonization after 1(A) and 5(B) days simulation time.

the initial conditions and the colonization phenomenon has not occurred yet. After 5 days of simulation time (Fig. 2-B), the biofilm experiences oxygen limitation, due to the low concentration maintained within the bulk liquid. As a consequence, the NOB concentration significantly decreases with respect to the initial fraction, with the AOB and HB being the two species proliferating the most. The AOB activity is confirmed by the decrease in S_1 concentration within the bulk liquid with respect to the inlet concentration and a concurrent increase in S_2 concentration (Fig. 4-A,B). Note that the latter keeps higher than S_3 concentration as the metabolic activity of NOB is limited by the low oxygen concentration. The organic carbon is completely depleted within the biofilm and its concentration keeps lower than $1mgl^{-1}$.

Figure 3-A displays biofilm configuration after 20 days of system operation: nevertheless the concomitant formation of an anoxic zone in the inner part of the biofilm (Fig. 4-C) and a non-zero ψ_2 concentration all over the biofilm (data not shown), AMX have not yet established in sessile form. This might be due to the very slow growth rate of X_2 . The biofilm is dominated by X_1 while X_5 predominate in the inner layer. Substrate trends assume the following configuration: S_1 increases in the bulk liquid due to the lower AOB activity in the outer art of the biofilm where S_5 is totally consumed and its depletion determines the formation of an anoxic zone. Moreover, S_2 represents the main abundant product while S_3 and S_4 keep close to zero all over the biofilm (Fig. 4-C). At day 50 (Figs. 3-B and 4-D), AMX have colonized the environmental niche which formed at the bottom of the biofilm (Fig. 3-B). AOB still dominate the aerobic zone while NOB are confined to the internal layers. S_3 concentration is close to zero as the metabolism of NOB significantly




FIGURE 3. Microbial species distribution of a multispecies biofilm undergoing ψ_2 colonization after 20(A) and 50(B) days simulation time.

slows down. Note that the availability of S_5 within the biofilm is strictly connected to the penetration depth whose decrease leads to an increasing anoxic zone (Fig. 4-D). Furthermore, AMX grow only where favorable environmental conditions establish despite the biofilm results fully penetrated by the same bacteria in motile/colonizing form ψ_2 for all simulation times (data not shown).

AMX invasion is significantly influenced by many parameters such as environmental factors (i.e. pH and temperature) and operational conditions (i.e. dilution rate, C/N ratio, aeration pattern). The main goals for the further computational studies are to determine how the invasion phenomenon is affected by the oxygen and organic carbon availability. For this reason we vary the concentration of oxygen in the bulk liquid S_{5L} and the organic carbon concentration in the inlet S_4^{in} in the range [0.5-6] and [120-750] respectively. In the following, we will refer to Secs. 4.1.1 and 4.1.2 for the applications with the variable oxygen and organic carbon. We assumed the initial condition reported in Fig. 1 for all the simulation studies.

4.1.1. Model 1 - Assessment -A- Effect of oxygen concentration. Model outcomes for the simulation studies with variable oxygen are summarized in Figs. 5 and 6 in terms of biomass distribution and substrate concentrations within the bulk liquid. Four different oxygen levels $(0.5-1.5-3-6 mgO_2/L)$ have been tested and the simulations have been run for a target time of 50 days. AMX are strictly inhibited by the oxygen concentration and as expected, their total volume fraction is found to slightly increase when varying the oxygen level from 6 to 1.5. A lower oxygen concentration leads to a decrease in AMX fraction; indeed, under this condition HB reaches the highest fraction competing with AMX and NOB for S_2 . The optimal condition for AMX establishment and proliferation within the biofilm occurs at $3 mgO_2/L$, even if the relative total biofilm fraction is lower with respect to 1.5 mgO_2/L . Of course, NOB fraction is higher when the oxygen concentration is equal to 6 mgO_2/L . Regarding nitrogen removal, it is possible to note that the S_1 concentration progressively decreases and consequently S_2 increases going from 0.5 to 6 mgO_2/L (Fig. 4.6). These substrates show fully penetrated profiles (data not shown) and consequently the AMX can grow for all the cases with their maximum specific growth rate, but only where anoxic conditions are established.

4.1.2. Model 1 – Assessment -B- Effect of inlet organic carbon concentration. The second simulation studies investigated the effect of increasing C/N ratios on AMX performances. The oxygen concentration within the bulk liquid has been fixed to $3 \text{ } mgO_2/L$. As shown




FIGURE 4. Substrate trends within a multispecies biofilm undergoing ψ_2 colonization after 1(A), 5(B), 20(C) and 50(D) days simulation time. Dotted black line S_1 , continuous grey line S_2 , dashed grey line S_3 , continuous black line S_4 , dashed-dotted black line S_5 . S_1, S_2, S_3 concentrations are reduced by a factor of 0.002, 0.005 and 0.005 respectively.

in Figs. 7 and 8, four different concentrations of the inlet organic carbon S_4^{in} have been tested $(120-250-500-750\ mgCOD/L)$ and all the simulations have been run for 50 days. Figure 7 shows that the AOB volume fraction is prevalent when low organic carbon is available for HB, which compete for oxygen with all the other aerobic species in the external part of the biofilm. AMX invasion and proliferation is favored at $S_4^{in}=500mgCOD/L$ since the NOB significantly decrease when increasing the inlet organic carbon concentration. The highest carbon content leads to the highest inerts volume fraction as HB are strongly predominant and out-compete all the other species.

According to the volume fraction distribution, total nitrogen removal is higher when both AMX and AOB can easily perform their metabolisms while NOB activity is inhibited by HB. This particular condition is more evident when the inlet carbon concentration is $250 \ mgCODl^{-1}$ and a higher amount of dissolved oxygen is utilized by HB. Ammonium removal is not significant when both AMX and AOB are not prevalent within the biofilm and organic carbon removal starts to be incomplete when increasing S_4^{in} concentration to $750 \ mgCODl^{-1}$.




FIGURE 5. Total biofilm volume fractions at different O_2 concentrations after 50 days simulation time.




FIGURE 6. Substrate concentrations within the bulk liquid at different O_2 concentrations after 50 days simulation time.

4.2. Model 2 – Two invading species. In this section, the model was applied to the case of two species invasion, HB and AMX respectively. The microbial species growth is governed by equations (4.1) with the following initial volume fractions

$$(4.26) f_1(z,0) = 0.7, f_2(z,0) = 0, f_3(z,0) = 0.3, f_4(z,0) = 0, f_5(z,0) = 0.$$

Only the species X_1 and X_3 are supposed to inhabit the biofilm at t = 0. The invasion of the species X_2 and X_4 is simulated. The initial biofilm thickness L_0 is given by (4.3). A representation of the initial microbial distribution is reported in Fig. 9.




FIGURE 7. Total biofilm volume fractions at different S_4^{in} values after 50 days simulation time.




FIGURE 8. Substrate concentrations within the bulk liquid at different S_4^{in} values after 50 days simulation time.

The biomass growth rates $r_{M,i}$ are the same as Model 1, formulas (4.4)-(4.8). The specific growth rates r_i induced by the switch of the planktonic cells to the sessile mode of growth are defined as

$$(4.27) r_1 = r_3 = r_5 = 0,$$

(4.28)
$$r_2 = k_{col,2} \frac{\Psi_2}{k_{\psi,2} + \Psi_2} \frac{K_{2,5}}{K_{2,5} + S_5} \frac{S_1}{K_{2,1} + S_1} \frac{S_2}{K_{2,2} + S_2},$$




FIGURE 9. Initial biofilm configuration for Model 2.

$$r_{4} = k_{col,4} \frac{\Psi_{4}}{k_{\psi,4} + \Psi_{4}} \left(\frac{S_{4}}{K_{4,4} + S_{4}} \frac{S_{5}}{K_{4,5} + S_{5}} + \beta_{1} \frac{K_{4,5}}{K_{4,5} + S_{5}} \frac{S_{4}}{K_{4,4} + S_{4}} \frac{S_{3}}{K_{4,3} + S_{3}} \frac{S_{3}}{S_{2} + S_{3}} \right) + \beta_{2} \frac{K_{4,5}}{K_{4,5} + S_{5}} \frac{S_{4}}{K_{4,4} + S_{4}} \frac{S_{2}}{K_{4,2} + S_{2}} \frac{S_{2}}{S_{2} + S_{3}} \right).$$

The growth rate terms r_2 and r_4 for X_2 and X_4 , respectively, indicate that the transition of bacteria from planktonic state ψ_2 , ψ_4 into the sessile state X_2 , X_4 is controlled by the formation of specific environmental niches connected to the local concentration of dissolved substrates. As in Model 1, consider second and fourth equation in (2.8) with $r_{M,2}$ and $r_{M,4}$ given by (4.5) and (4.7), respectively. If it is supposed that $r_2 = r_4 = 0$, then the mentioned equations with initial condition $f_2(z,0) = f_4(z,0) = 0$ admit the unique solution $f_2(z,t) = f_4(z,t) = 0$ and the species X_2 and X_4 cannot develop.

The initial-boundary conditions for S_j and net conversion rates of substrates are the same as Model 1, formulas (4.12)-(4.13) and (4.14)-(4.18), respectively. The initial conditions for S_j^* are given by (4.20). The initial-boundary conditions for Ψ_i are same as Model 1, formula (4.23). The initial conditions for ψ_i^* are the following

$$(4.30) \psi_1^{in} = 0, \ \psi_2^{in} = 1.0 \ mgCOD/L, \ \psi_3^{in} = 0, \ \psi_4^{in} = 1.2 \ mgCOD/L, \ \psi_5^{in} = 0.$$

Note that, by using the same arguments as Model 1, it can be shown that $\Psi_1(z,t) = 0$, $\psi_1^*(t) = 0$, $\Psi_3(z,t) = 0$, $\psi_3^*(t) = 0$, $\Psi_5(z,t) = 0$, $\psi_5^*(t) = 0$.

The operational parameters of the biofilm reactor are the same as Model 1.

In Figs. 10 and 11 the simulation results for the multispecies biofilm system with two invading species are reported. Differently from Model 1, the X_4 invasion is very fast and it is already visible after 5 days of simulation time (Fig. 10-B). This is due to the different environmental conditions that trigger the invasion of the two microbial species. Indeed the establishment of X_2 is dependent on the formation of an anoxic zone within the biofilm while X_4 are facultative bacteria and can grow in both aerobic and anoxic environments. After 20 days the biofilm configuration is the same of the previous application (Figs. 3-A and 11-A) and as we can expect, the further evolution of the system is practically the same for the two cases studied. Simulation results confirm model capability of predicting the invasion phenomenon on time and space. Indeed, the model is able to predict the delays between the X_2 and X_4 colonizations and the location where the two planktonic species




FIGURE 10. Microbial species distribution of a multispecies biofilm undergoing ψ_2 and ψ_4 colonization after 2(A) and 5(B) days simulation time.




FIGURE 11. Microbial species distribution of a multispecies biofilm undergoing ψ_2 and ψ_4 colonization after 20(A) and 50(B) days simulation time.

establish. To the best of our knowledge, such results cannot be achieved by the existing continuum biofilm models but they might have a significant impact on the developing of new strategies for such biofilm reactors operation.

5. Conclusion

In this work, the qualitative analysis of the free boundary problem related to the invasion phenomenon in biofilm reactors has been discussed. The model takes into account the dynamics of sessile species, nutrients and microbial products, and planktonic cells, the latter diffusing from the bulk liquid within the biofilm matrix, where they might switch their status from motile to sessile and thus colonize the pre-existing biofilm. The dynamics of bulk liquid have been explicitly modeled by considering two systems of nonlinear ordinary differential equations which derive from mass conservation principles. An existence and uniqueness result has been provided for the related free boundary value problem by using the method of characteristics and the fixed point theorem. It is important to notice that the planktonic species are just provided by the bulk liquid; however, the reverse process which accounts for the switch from sessile to planktonic form of life might occur under specific conditions. This phenomenon could be explicitly taken into account by considering a direct dependence of the free planktonic cell reaction rates on the concentration of the sessile bacteria. The same methodology adopted in this work could be easily adapted to

address the existence and uniqueness questions of this new system. Numerical simulations related to a real biofilm system have been performed. Two specific model applications have been analyzed. Simulation results demonstrate the underlying conclusion that the invasion model can be effectively used as a predictive tool to develop specific reactor operation strategies. Further developments might be related to the definition of a calibration protocol through experimental data.

References

- [1] Boltz, J. P., Smets, B. F., Rittmann, B. E., van Loosdrecht, M. C., Morgenroth, E., Daigger, G. T. 2017. From biofilm ecology to reactors: a focused review. *Water Science and Technology* **75**, 1753: 1760.
- [2] Mattei, M.R., Frunzo, L., D'Acunto, B., Esposito, G., Pirozzi, F. 2015. Modelling microbial population dynamics in multispecies biofilms including anammox bacteria. *Ecological Modelling* **304**, 44: 58.
- [3] Ward, J. P., King, J. R. 2012. Thin-film modelling of biofilm growth and quorum sensing. *Journal of Engineering Mathematics* **73**, 71: 92.
- [4] D'Acunto, B., Frunzo, L., Klapper, I., Mattei, M.R. 2015. Modeling multispecies biofilms including new bacterial species invasion. *Mathematical biosciences* **259**, 20: 26.
- [5] Mašić, A., Eberl, H. J. 2014. A modeling and simulation study of the role of suspended microbial populations in nitrification in a biofilm reactor. *Bulletin of Mathematical Biology* **76**, 27: 58.
- [6] D'Acunto, B., Frunzo, L., Mattei, M. R. 2016. Qualitative analysis of the moving boundary problem for a biofilm reactor model. *Journal of Mathematical Analysis and Applications* 438(1), 474: 491.
- [7] Frunzo, L., Mattei, M. R. 2017. Qualitative analysis of the invasion free boundary problem in biofilms. *Ricerche di Matematica* **66**, 171: 188.
- [8] Wanner, O., Gujer, W. 1986. A multispecies biofilm model. *Biotechnology Bioengineering* 28, 314: 328.
- [9] D'Acunto, B., Esposito, G., Frunzo, L., Pirozzi, F. 2011. Dynamic modeling of sulfate reducing biofilms. *Computers Mathematics with Applications* **62**, 2601: 2608.
- [10] Cao, Y., van Loosdrecht, M. C., Daigger, G. T. 2017. Mainstream partial nitritation-anammox in municipal wastewater treatment: status, bottlenecks, and further studies. *Applied Microbiology and Biotechnology* **101**, 1365: 1383.
- [11] Emerenini, B. O., Hense, B. A., Kuttler, C., Eberl, H. J. 2015. A mathematical model of quorum sensing induced biofilm detachment. *PloS one* **10**(7), e0132385.
- [12] Dockery, J., Klapper, I. 2002. Finger formation in biofilm layers. SIAM Journal on Applied Mathematics 62, 853: 869.
- [13] Emerenini, B., Sonner, S., Eberl, H. 2017. Mathematical analysis of a quorum sensing induced biofilm dispersal model and numerical simulation of hollowing effects. *Mathematical Biosciences and Engineering* 14(3), 625: 653.

University of Naples "Federico II", Department of Mathematics and Applications, Complesso Monte Sant'Angelo, 80124, Naples, Italy dacunto@unina.it, Luigi.frunzo@unina.it, vincenzo.luongo@unina.it, mariarosaria.mattei@unina.it