

# Differential Virial Constraints on Orbital-free Kinetic Energy Density Functionals

Kai Luo<sup>1,\*</sup> and S.B. Trickey<sup>2,†</sup>

<sup>1</sup>Quantum Theory Project, Department of Physics, University of Florida, Gainesville, FL 32611

<sup>2</sup>Quantum Theory Project, Departments of Physics and of Chemistry, University of Florida, Gainesville, FL 32611

(Dated: 09 Nov. 2017)

Various constraints on the spatial derivative dependencies of approximate orbital-free kinetic energy density functionals have been based on application of differential virial theorems. We point out a defect in the core argument: the relationships are not true for an arbitrary density but hold only for the minimizing density and corresponding chemical potential associated with a specified external potential. Contrary to the claims therefore, the relationships are not constraints and provide no independent information about the spatial derivative dependencies of approximate functionals. A simple argument also shows that validity for arbitrary  $v$ -representable densities is not restored by appeal to the density-potential bijection that is the heart of the original Hohenberg-Kohn theorem.

## I. INTRODUCTION

Density functional theory in its Kohn-Sham (KS) form<sup>1</sup> decomposes the ground-state density functional for the total electronic energy of an  $N_e$  electron system with number density  $n(\mathbf{r})$  in an external field  $v_{\text{ext}}$  into contributions identifiable from the non-interacting KS system at the same density. The total energy functional in KS form is<sup>1</sup>

$$E[n] = T_s[n] + E_{\text{ext}}[n] + E_{\text{H}}[n] + E_{\text{xc}}[n], \quad (1)$$

with  $T_s[n]$  the KS kinetic energy functional,  $E_{\text{ext}}[n]$  the external field interaction energy,  $E_{\text{H}}[n]$  the Hartree energy (classical electron-electron repulsion), and  $E_{\text{xc}}[n]$  the exchange-correlation (XC) energy functional. (Remark: any external system configurational energy is omitted as irrelevant here.) Minimization formally gives a single Euler equation,

$$\frac{\delta T_s[n]}{\delta n(\mathbf{r})} + v_{\text{KS}}([n]; \mathbf{r}) = \mu. \quad (2)$$

Here  $v_{\text{KS}} = \delta(E_{\text{ext}} + E_{\text{H}} + E_{\text{xc}})/\delta n$  is the KS potential and  $\mu$  is the chemical potential associated with the minimizing density  $n_0$  that yields fixed  $N_e$ .

The familiar orbital-dependent KS problem follows from introducing

$$\begin{aligned} T_s[\{\varphi_i\}_{i=1}^{N_e}] &:= \frac{1}{2} \sum_{i=1}^{N_e} \int d\mathbf{r} |\nabla \varphi_i(\mathbf{r})|^2 \\ &\equiv \int d\mathbf{r} t_s^{\text{orb}}(\mathbf{r}) \end{aligned} \quad (3)$$

(in Hartree a.u.). In this form the integrand is manifestly positive definite. The more common Laplacian form is not. The difference is a surface integral which ordinarily is zero.

Minimization to find the ground-state density, hence the ground-state energy, gives the KS equation

$$\left\{-\frac{1}{2}\nabla^2 + v_{\text{KS}}([n]; \mathbf{r})\right\}\varphi_i(\mathbf{r}) = \epsilon_i\varphi_i(\mathbf{r}). \quad (4)$$

with

$$n_0(\mathbf{r}) = \sum_i^{N_e} \varphi_i^2(\mathbf{r}) \quad (5)$$

the ground-state density. For simplicity and without loss of generality, we have assumed singly occupied KS orbitals and no degeneracy at the Fermi level. Satisfaction of the Euler equation in either form, (2) or (4), is a critical element in the discussion below. Therefore that is explicit in the notation of Eq. (5).

Orbital-free DFT aims to provide useful approximations to  $T_s[n]$  without explicit use of the KS orbitals. If one restricts attention to single-point representations,

$$T_s[n] := \int d\mathbf{r} t_s[n(\mathbf{r})], \quad (6)$$

a basic issue is the maximum order of spatial derivative dependence to include in an approximate  $t_s$ . Generalized gradient approximations<sup>2</sup> (GGA) and Laplacian-level functionals<sup>3-6</sup> are the practical limits so far. Various dimensionless spatial derivative combinations (reduced density derivatives) have been proposed<sup>7</sup> but little is known about how to select from among them. An exception would seem to be papers by BalTin<sup>8</sup> and co-workers<sup>9</sup> and others<sup>10,11</sup>. Those use differential virial theorems to derive constraints on the order of spatial derivative that can appear. Here we show that those relationships are not constraints but trivial identities of complicated form satisfied only by the equilibrium density for a given  $v_{\text{ext}} = \delta E_{\text{ext}}/\delta n$ .

In the next section we rehearse the original arguments from Ref. [8] using the one-dimensional (1D) case presented there. (The three-dimensional case uses identical logic but is more cumbersome, so we do not treat it explicitly.) In the subsequent section, we discuss two related omissions in those arguments which significantly alter the claimed consequences to the point of triviality. We illustrate by reconsidering two cases originally treated in Ref. [8]. Brief consideration to show that a seemingly plausible Hohenberg-Kohn bijectivity argument does not alter the result concludes the presentation.

## II. DIFFERENTIAL VIRIAL CONSTRAINT- 1D

### A. Original Argument

The original argument of Ref. [8] follows in our notation. Consider a 1D system and its KS potential and states. For it the differential virial theorem (Eq. (13) of Ref. [12]) is

$$\frac{dt_s(x)}{dx} = \frac{1}{8} \frac{d^3 n(x)}{dx^3} - \frac{1}{2} n(x) \frac{dv_{KS}}{dx} \quad (7)$$

or in primed notation as used in Ref. [8],

$$t'_s(x) = \frac{n'''(x)}{8} - \frac{1}{2} n(x) v'_{KS}(x). \quad (8)$$

(Remark: To get to the Euler equation equivalent to our Eq. (2), Eq. (7) of Ref. [8] writes the supposed equivalent of our Eq. (1) in 1D as

$$E[n] = \int_{-\infty}^{\infty} dx t_s(x) + \int_{-\infty}^{\infty} dx n(x) v_{KS}(x). \quad (9)$$

This is incorrect since  $v_{KS}$  is not solely the external potential but the error is inconsequential for the discussion at hand.)

Ref. [8] then considers a one-point approximation for  $t_s$  that depends on spatial derivatives of  $n$  through  $g^{th}$  order:

$$t_s(x) := f(n, n', n'', n''', \dots, n^{(g)}). \quad (10)$$

Straightforwardly one gets

$$t'_s(x) = \sum_{\nu=0}^g \frac{\partial f}{\partial n^{(\nu)}} \frac{dn^{(\nu)}}{dx} = \sum_{\nu=0}^g \frac{\partial f}{\partial n^{(\nu)}} n^{(\nu+1)} \quad (11)$$

which is Eq. (11) in Ref. [8]. Alternatively, repeated integration by parts gives Eq. (6) of that reference,

$$\frac{\delta T_s}{\delta n} = \sum_{\nu=0}^g (-1)^\nu \frac{d^\nu}{dx^\nu} \left[ \frac{\partial f}{\partial n^{(\nu)}} \right]. \quad (12)$$

Ref. [8] then rewrites the Euler equation (2) with (12) and takes one spatial derivative to get

$$v'_{KS}(x) = \sum_{\nu=0}^g (-1)^{\nu+1} \frac{d^{\nu+1}}{dx^{\nu+1}} \left[ \frac{\partial f}{\partial n^{(\nu)}} \right]. \quad (13)$$

Substitution of both this result and the result from (11) in Eq. (8) then gives

$$\sum_{\nu=0}^g \left[ (-1)^\nu n \frac{d^{\nu+1}}{dx^{\nu+1}} \left( \frac{\partial f}{\partial n^{(\nu)}} \right) - 2n^{(\nu+1)} \frac{\partial f}{\partial n^{(\nu)}} \right] = -\frac{1}{4} n'''. \quad (14)$$

Ref. [8] then says that "...this equation has to be looked upon as a relation to be satisfied identically with respect to the variables  $n, n', \dots, n^{(\ell)}$  occurring in it and

that the equation "is a condition to be imposed on the dependence of  $f$  upon the variables  $n, n', \dots, n^{(\ell)}$ ". There follows an examination of functions  $f$  which depend on  $n^{(g)}$  through  $g = 2$  with the conclusion that the only allowable form consists of the full von Weizsäcker term<sup>13</sup> plus an arbitrarily scaled Thomas-Fermi term<sup>14,15</sup>.

### B. Difficulty

There are two consequential flaws in the foregoing argument that seem not to have been recognized heretofore. They have a common stem. First, the differential virial relation from which Eq. (8) is derived holds *only* for the exact eigenstates of the given Hamiltonian. In the KS case with fixed external potential, that differential virial relation therefore properly reads

$$t'_s(n_0(x)) = \frac{n_0'''(x)}{8} - \frac{1}{2} n_0(x) v'_{KS}(n_0(x)). \quad (15)$$

The same error occurs in use of the Euler equation to get the spatial derivative of the potential. The Euler equation is not a general functional relation for arbitrary density  $n$ . Rather it is a relationship between the minimizing density  $n_0$  and the unique (up to a constant) external potential which is paired with that  $n_0$ . Thus Eq. (13) must be replaced by

$$v'_{KS}(n_0(x)) = \sum_{\nu=0}^g (-1)^{\nu+1} \frac{d^{\nu+1}}{dx^{\nu+1}} \left[ \frac{\partial f}{\partial n^{(\nu)}} \right]_{n_0}. \quad (16)$$

As a consequence, the purported constraint on functional dependence becomes

$$\sum_{\nu=0}^g \left[ (-1)^\nu n_0 \frac{d^{\nu+1}}{dx^{\nu+1}} \left( \frac{\partial f}{\partial n^{(\nu)}} \right)_{n_0} - 2n_0^{(\nu+1)} \frac{\partial f}{\partial n^{(\nu)}} \Big|_{n_0} \right] = -\frac{1}{4} n_0'''. \quad (17)$$

This is a requirement on the behavior of  $f$  at a single point  $n_0$  in the space of one-body densities and paired with a specific  $v_{ext}$ . Contrary to Ref. [8], Eq. (17) is *not* a condition on the dependence of  $f$  upon the variables  $n, n', \dots, n^{(\ell)}$  for arbitrary density  $n$  given a  $v_{ext}$ . Rather, given a dependence through order  $n^{(\ell)}$ , and a particular  $v_{ext}$ , the requirement is to find the equilibrium density  $n_0$  that satisfies (17).

## III. 1D EXAMPLES

### A. 1D Homogeneous Electron Gas

For the 1D homogeneous electron gas (HEG), the Thomas-Fermi functional,  $T_{TF}$

$$T_{TF} = c_{TF} \int dx n^3(x) \quad (18)$$

is exact. But put aside that knowledge for a moment and simply consider  $T_{TF}$  as a candidate approximate KE functional. The associated kinetic energy density and partial derivative are

$$t_{TF} = c_{TF} n^3 := f(n) \quad (19)$$

$$\frac{\partial f}{\partial n} = 3c_{TF} n^2. \quad (20)$$

Then Eq. (17) becomes

$$n_0 \frac{d}{dx} (3c_{TF} n_0^2) - 2n_0^{(1)} (3c_{TF} n_0^2) = -\frac{1}{4} n_0^{(3)}. \quad (21)$$

Its solution is

$$n_0(x) = \frac{1}{2} a x^2 + b x + c, \quad (22)$$

with coefficients to be determined. An appropriate boundary condition is periodic

$$n_0(x+L) = n_0(x) \quad (23)$$

where  $L$  is a suitable length. As a result

$$a = b = 0. \quad (24)$$

The constant  $c$  is trivially set by imposition of the desired value of the density. The outcome of the supposed constraint is simply to demonstrate that  $t_{TF}$  is compatible with the HEG.

If, on the other hand, one imposes box boundary (BB) conditions

$$n_0(0) = n_0(L) = 0 \quad (25)$$

one has  $c = 0$  and

$$n_{0,BB}(x) = \frac{6N_e}{L^3} x(L-x). \quad (26)$$

This density, however is unacceptable, since it violates Lieb's condition<sup>16</sup> for the finitude of the KE:

$$\int_0^L dx \left( \frac{d\phi}{dx} \right)^2 < \infty \quad (27)$$

$$\phi(x) := \sqrt{n_{0,BB}(x)} \quad (28)$$

Alternatively, one may see the problem with  $n_{0,BB}$  by attempting direct inversion of the Schrödinger equation for  $\phi$  in the  $N_e = 1$  case to recover the one-body potential. Up to a constant, the purported potential is negative definite with poles at  $x = 0, L$ :  $(-L^2/[8x^2(L-x)^2])$ , i.e.,  $n_{0,BB}$  is not  $v$ -representable. There is nothing special about  $N_e = 1$  to rescue the case.

Thus, all that Eq. (17) yields in the  $T_{TF}$  case is confirmation that  $T_{TF}$  is indeed correct for the 1D HEG and that it also has solutions which upon detailed inspection do not correspond to any potential. For cases in which  $v_{\text{ext}}$  does exist, Eq. (17) has no information about it and provides no information on the accuracy of the approximation  $t_s \approx t_{TF}$ . Thus, contrary to Ref. [8], no general requirement on the dependence of  $T_s[n]$  upon spatial derivative order is obtained from Eq. (17) when  $T_{TF}$  is put to the test.

## B. One Electron in 1D

For a 1-electron system, the von Weizsäcker functional  $T_W$

$$T_s = T_W = \int dx t_W(x) \quad (29)$$

is exact. Its kinetic energy density is

$$t_W = \frac{[n'(x)]^2}{8n(x)} := f_W(n, n'). \quad (30)$$

For convenience, the relevant partial derivatives for use of Eq. (17) are

$$\begin{aligned} \frac{\partial f_W}{\partial n} &= -\frac{(n')^2}{8n^2} \\ \frac{\partial f_W}{\partial n'} &= \frac{n'}{4n} \end{aligned}$$

Substitute these results in the left-hand side of Eq. (17) to get

$$\begin{aligned} &\left( n_0 \frac{d}{dx} \frac{-(n'_0)^2}{8n_0^2} - 2n'_0 \frac{-(n'_0)^2}{8n_0^2} \right) \\ &+ \left( -n_0 \frac{d^2}{dx^2} \frac{n'_0}{4n_0} - 2n''_0 \frac{n'_0}{4n_0} \right) = \\ &\left( -\frac{n'_0 n''_0}{4n_0} + \frac{(n'_0)^3}{4n_0^2} + \frac{(n'_0)^3}{4n_0^2} \right) \\ &+ \left( -\frac{1}{4} n_0^{(3)} + \frac{3n'_0 n''_0}{4n_0} - \frac{(n'_0)^3}{2n_0^2} - \frac{n'_0 n''_0}{2n_0} \right) = \\ &\quad -\frac{1}{4} n_0^{(3)} \end{aligned}$$

This is the same as the right hand side of Eq. (17) so that equation reduces to a trivial identity for all equilibrium densities associated with the combination  $T_W$ , some  $E_{\text{xc}}$ , and some  $E_{\text{ext}}$ . Therefore, no information is provided by the differential virial constraint, Eq. (17), about the functional dependence of  $T_s$  upon spatial derivatives except that  $t_W$  is a valid form. Note also that unlike the 1D HEG or box-bounded  $t_{TF}$  cases considered above, there are infinitely many densities that lead to the trivial identity because there are infinitely many single-electron external potentials. Thus, there is no access to a unique solution  $n_0$  provided by the purported constraint.

## IV. DISCUSSION AND CONCLUSIONS

Examination of the 3D version of the differential virial constraint argument as summarized, for example, in Ref. [11], shows that the same critical mis-use of the Euler equation occurs in 3D as in 1D. The preceding analysis therefore holds unchanged.

It might seem that Hohenberg-Kohn bijectivity between  $v_{\text{ext}}$  and  $n_0$  could rescue the argument by making the Euler equation true for an arbitrary  $v$ -representable density. Note that even if that were the case, the differential virial theorem part of the argument itself still would hold only for the extremalizing density. But HK bijectivity does not remove the Euler equation restriction either. Bijectivity is true for an arbitrary density precisely and only in the case that the arbitrarily chosen density is the *minimizing* density  $n_0$  paired with its unique external potential. Bijectivity obviously is meaningless for the case of interest, namely a *fixed*  $v_{\text{ext}}$  but *arbitrary*  $n(\mathbf{r})$ . The required pairing of density and potential is missing. Thus, though the Euler equation holds for arbitrary  $n_0$  with the associated  $v_{\text{ext}}[n_0]$  and corresponding  $\mu[n_0]$ , the flaw identified above persists. Spatial differentiation of the Euler equation to replace  $v'_{\text{KS}}$  (recall Eq. (13)) in the differential virial relation still ties the result to equilibrium densities  $n_0$ , not arbitrary ones.

It is worth noting that the non-uniqueness of KE densities<sup>17</sup> (indeed, any energy density) should raise suspicions about the validity of any supposed constraint on the spatial derivative dependence of an approximation for  $t_s$ . The well-known vanishing of  $\nabla^2 n$  terms is an example. In fact, it is the counterexample to the Ref. [8] argument (just after Eq. (48) of that reference) that the approximate KE density must obey  $t_s^{\text{approx}} \geq 0$ . That constraint is highly valuable but it is a choice of gauge. The issue is discussed in detail in Refs. [18,19]. A related issue is that it is not self-evident that a function having up through  $g^{\text{th}}$  order derivatives, Eq. (11), necessarily is itself differentiable for arbitrary densities. Nor is it always true that one can do the repeated integration by parts, Eq. (12) with vanishing surface terms; see Ref.

[20] for counterexamples.

The present analysis resolves at least one other peculiar finding in Refs. [8] and [9], namely that through second-spatial derivatives the only KE density form consistent with the supposed differential virial constraint is  $t_s^{\text{approx}} = \lambda t_{TF} + t_W$  with  $\lambda$  an undetermined constant. This restriction is suspect on its face because of the Lieb conjecture<sup>21</sup> that  $T_s \leq T_{TF} + T_W$ . That conjecture is consistent with the  $N_e \rightarrow \infty$  limit of the bound found by Gázquez and Robles<sup>22</sup>. (See also Acharya et al.<sup>23</sup> for a heuristic formulation with number dependence that has the Lieb bound as the  $N_e \rightarrow \infty$  limit.) For finite systems, the only straightforward way to make that peculiar result consistent with the Gázquez-Robles expression would be for  $\lambda$  to be number-dependent, thereby raising an obvious problem of size-consistency. There is no obvious simple way to make the result consistent with the Lieb bound in the thermodynamic limit. The analysis presented here removes that problem by showing that all the supposed constraint really does is confirm that for a specific  $n_0$  one always can find a  $\lambda$  which makes the claim true. Just pick

$$\lambda[n_0] = \frac{T_s[n_0] - T_W[n_0]}{T_{TF}[n_0]}. \quad (31)$$

While true, it is essentially tautological, hence useless.

### Acknowledgments

We acknowledge informative conversations with Jim Dufty and Valentin Karasiev and helpful email exchange with Paul Ayers. This work was supported by the U.S. Dept. of Energy grant de-sc0002139.

\* Electronic address: kluo@ufl.edu

† Electronic address: trickey@qtp.ufl.edu

<sup>1</sup> W. Kohn and L.J. Sham, Phys. Rev. **140**, A1133 (1965).

<sup>2</sup> V.V. Karasiev, D. Chakraborty, O.A. Shukruto, and S.B. Trickey, Phys. Rev. B **88**, 161108(R) (2013) and refs. therein.

<sup>3</sup> J. P. Perdew and L.A. Constantin, Phys. Rev. B **75**, 155109 (2007).

<sup>4</sup> S. Laricchia, L.A. Constantin, E. Fabiano, and F. Della Sala, J. Chem. Th. Comput. **10**, 164 (2014).

<sup>5</sup> A.C. Cancio, D. Stewart and A. Kuna, J. Chem. Phys. **144**, 084107 (2016).

<sup>6</sup> A.C. Cancio and J.J. Redd, Mol. Phys. **15**, 618 (2017).

<sup>7</sup> V.V. Karasiev, R.S. Jones, S.B. Trickey, and F.E. Harris, Phys. Rev. B **80**, 245120 (2009).

<sup>8</sup> R. Baltin, J. Phys. A: Math. Gen. **20**, 111 (1987).

<sup>9</sup> J. Shao and R. Baltin, J. Phys. A: Math. Gen. **23** 5939 (1990).

<sup>10</sup> A. Holas and N.H. March, Int. J. Quantum Chem. **56**, 371 (1995).

<sup>11</sup> F.H. Alharbi and S. Kais, Int. J. Quantum Chem. **117**,

e25373 (2017).

<sup>12</sup> R. Baltin, Phys. Lett. **113A**, 121 (1985).

<sup>13</sup> C.F. von Weizsäcker, Z. Phys. **96**, 431 (1935).

<sup>14</sup> L.H. Thomas, Proc. Cambridge Phil. Soc. **23**, 542 (1927).

<sup>15</sup> E. Fermi, Atti Accad. Nazl. Lincei **6**, 602 (1927).

<sup>16</sup> E.H. Lieb, Int. J. Quantum Chem. **24**, 243 (1983)

<sup>17</sup> E. Sim, J. Larkin, K. Burke, and C.W. Bock, J. Chem. Phys. **118**, 8140 (2003) and refs. therein.

<sup>18</sup> P.W. Ayers, R.G. Parr, and Á. Nagy, Int. J. Quantum Chem. **90**, 309 (2002).

<sup>19</sup> S.B. Trickey, V.V. Karasiev, and R.S. Jones, Int. J. Quantum Chem. **109**, 2943 (2009).

<sup>20</sup> J.P. Perdew, V. Sahni, M.K. Harbola, and R.K. Pathak, Phys. Rev. B **34**, 686 (1986); erratum *ibid.* **37**, 4267 (1988).

<sup>21</sup> E.H. Lieb, Lecture Notes in Physics **116**, 91 (1980).

<sup>22</sup> J.L. Gázquez and J. Robles, J. Chem. Phys. **76**, 1467 (1982).

<sup>23</sup> P.K. Acharya, L.J. Bartolotti, S.B. Sears, and R.G. Parr Proc. Nat. Acad. Sci. (US) - Phys. Sci. **77**, 6978 (1980).